Electronic Version
Stylesheet Version v1.1.1

Description

BIOINFORMATICALLY DETECTABLE GROUP OF NOVEL REGULATORY GENES AND USES THEREOF

Cross Reference to Related Applications

[0001] The present invention is a continuation of application 10/293,338 incorporated herein, now abandoned.

Appendix Data

[0002] Computer Disk OBJECT_ID Sequence listing part

[0003] Computer Disk 2 OBJECT_ID #2 Sequence listing (copy 2)

Background of Invention

FIELD OF THE INVENTION

[0004] The present invention relates to a group of bioinformatically detectable novel genes, here identified as "genomic address messenger"or "GAM"genes, which are believed to be related to the micro RNA (miRNA) group of genes.

Description of Prior Art

[0005] Over 300 small noncoding RNA"s, termed Micro RNA"s (miRNA"s), have been identified to date in animals and plants (1-8). The MIR genes, code for ~70-120nt long non-protein-coding RNA sequences, that are the precursors of the ~22 nt miRNA sequences, which in some cases, specifically inhibit translation of target genes, by complementary binding to their untranslated regions (UTR) (9-13). The first two MIR genes discovered, Lin-4 and Let-7, were found in the nematode Ceanorhabditis elegans, and shown to be involved in developmental timing (14, 15), hence originally called Short Temporal RNA (stRNA) (16). The ~22 nt sequences inhibit translation of two respective target genes, Lin-14 and Lin-41, by complementary binding to their 3" untranslated regions (UTR) (14, 17). Later studies by three separate groups (1-3) identified several precursor and mature miRNA"s, some of them in clusters (1), expressed as sequences found in size fractionated (<80nt) total RNA, in a wide spectrum of species, including C. elegans, Drosophila melanogaster, Homo sapiens, as well as plants. More recently, additional miRNA"s from human (mir-91 to mir-121 (6)), mouse (mir-122 to mir-155 (4)) and plants (mir-156 to mir-171 (5)), have been reported. Furthermore, Llave et al identi-

fied in Arabidopsis thaliana 125 sequences, 21-24 nucleotides in length and presumably miRNAs (8). Each of the MIR genes is transcribed and processed to an ~80 nt-long hairpin shaped miRNA precursor, which is then processed by an enzyme called Dicer, to yield the mature ~22 nt-long single stranded miRNA. Such small RNAs will inhibit translation of target genes by complementary binding to sites in the 3" or 5" untranslated regions (18). Their structural similarity to the two well-studied miRNA"s in C. elegans, Lin-4 and Let-7, supports the belief that MIR genes code for specific translation inhibitors of target genes. However, determining the targets of the miRNAs is more complicated than just searching for antisense complementarity, because bulges and loops, disrupting perfect complementarity, are not only tolerated, but seem to be the rule in the postulated binding between the miRNA"s and their target UTR"s(19). Target-gene binding sites of most reported miRNA"s, except of Lin-4 and Let-7, have not been found, and therefore the specific functionality of these genes is still unknown. Using a computational approach, Rhoades et al (19), predicted targets for 14 A. thaliana miRNA"s by identifying their near complementarity to the predicted targets. However, using the

same approach was not helpful in identifying miRNA targets in C. elegans and D. melanogaster, raising the possibility that the near-perfect complementarity appears to be specific to plants (19).

[0006] The ability to detect novel RNA genes is limited by the methodologies used to detect such genes. All RNA genes identified so far either present a visibly discernable whole body phenotype, as do Lin-4 and Let-7 (Wightman et. al., Cell 75, 855 (1993); Reinhart et al., Nature 403, 901 (2000)), or produce significant enough quantities of RNA so as to be detected by the standard biochemical genomic techniques, as do the 93 recently detected miRNA genes. Since a limited number clones were sequenced by the researchers discovering these genes, 300 by Bartel and 100 by Tuschl (Bartel et. al., Science 294 ,858 (2001); Tuschl et. al., Science 294 ,853 (2001)), the RNA genes found can not be much rarer than 1% of all RNA genes. The recently detected miRNA genes therefore represent the more prevalent among the miRNA gene family.

[0007] Current methodology has therefore been unable to detect RNA genes which either do not present a visually discernable whole body phenotype, or are rare (e.g. rarer than 0.1% of all RNA genes), and therefore do not produce sig-

nificant enough quantities of RNA so as to be detected by standard biochemical technique.

SUMMARY OF INVENTION

[0008] The present invention relates to a novel group of regulatory, non-protein coding genes, which are functional in specifically inhibiting translation of other genes, some of which are known to be involved in various diseases. Each gene in this novel group of genes, here identified as "GAM"or "Genomic Address Messengers", specifically inhibits translation of one of more other "target" genes by means of complimentary hybridization of a segment of the RNA transcript encoded by GAM2, to an inhibitor site located in the 3"untranslated region of the mRNA of the one or more "target" genes.

[0009] In various preferred embodiments, the present invention seeks to provide improved method and system for specific modulation of expression of specific known "target" genes involved in significant human diseases, and improved method and system for detection of expression of these target genes.

[0010] Accordingly, the invention provides several substantially pure DNAs (e.g., genomic DNA, cDNA or synthetic DNA) each encoding a novel gene of the GAM group of gene,

vectors comprising the DNAs, probes comprising the DNAs, a method and system for selectively modulating translation of known "target" genes utilizing the vectors, and a method and system for detecting expression of known "target" genes utilizing the probe.

[0011] By "substantially pure DNA" is meant DNA that is free of the genes which, in the naturally-occurring genome of the organism from which the DNA of the invention is derived, flank the genes discovered and isolated by the present invention. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote at a site other than its natural site; or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

[0012] "Inhibiting translation" is defined as the ability to prevent synthesis of a specific protein encoded by a respective gene, by means of inhibiting the translation of the mRNA of this gene. "Translation inhibiter site" is defined as the

minimal DNA sequence sufficient to inhibit translation.

[0013] There is thus provided in accordance with a preferred embodiment of the present invention a bioinformatically detectable novel gene encoding substantially pure DNA wherein: RNA encoded by the bioinformatically detectable novel gene is about 18 to about 24 nucleotides in length, and originates from an RNA precursor, which RNA precursor is about 50 to about 120 nucleotides in length, a nucleotide sequence of a first half of the RNA precursor is a partial inversed-reversed sequence of a nucleotide sequence of a second half thereof, a nucleotide sequence of the RNA encoded by the novel gene is a partial inversed-reversed sequence of a nucleotide sequence of a binding site associated with at least one target gene, the novel gene cannot be detected by either of the following: a visually discernable whole body phenotype, and detection of 99.9% of RNA species shorter than 25 nucleotides expressed in a tissue sample, and a function of the novel gene is bioinformatically deducible.

[0014] There is further provided in accordance with another preferred embodiment of the present invention a bioinformatically detectable novel gene encoding substantially pure DNA wherein: RNA encoded by the bioninformatically

detectable novel gene includes a plurality of RNA sections, each of the RNA sections being about 50 to about 120 nucleotides in length, and including an RNA segment, which RNA segment is about 18 to about 24 nucleotides in length, a nucleotide sequence of a first half of each of the RNA sections encoded by the novel gene is a partial inversed-reversed sequence of nucleotide sequence of a second half thereof, a nucleotide sequence of each of the RNA segments encoded by the novel gene is a partial inversed-reversed sequence of the nucleotide sequence of a binding site associated with at least one target gene, and a function of the novel gene is bioinformatically deducible from the following data elements: the nucleotide sequence of the RNA encoded by the novel gene, a nucleotide sequence of the at least one target gene, and function of the at least one target gene.

[0015] There is still further provided in accordance with another preferred embodiment of the present invention a bioinformatically detectable novel gene encoding substantially pure DNA wherein: RNA encoded by the bioinformatically detectable novel gene is about 18 to about 24 nucleotides in length, and originates from an RNA precursor, which RNA precursor is about 50 to about 120 nucleotides in

length, a nucleotide sequence of a first half of the RNA precursor is a partial inversed-reversed sequence of a nucleotide sequence of a second half thereof, a nucleotide sequence of the RNA encoded by the novel gene is a partial inversed-reversed sequence of a nucleotide sequence of a binding site associated with at least one target gene, a function of the novel gene is modulation of expression of the at least one target gene, and the at least one target gene does not encode a protein.

[0016] There is additionally provided in accordance with another preferred embodiment of the present invention A bioinformatically detectable novel gene encoding substantially pure DNA wherein: the bioinformatically detectable novel gene does not encode a protein, RNA encoded by the bioinformatically detectable novel gene is maternally transferred by a cell to at least one daughter cell of the cell, a function of the novel gene includes modulation of a cell type of the daughter cell, and the modulation is bioinformatically deducible.

[0017] There is moreover provided in accordance with another preferred embodiment of the present invention a bioinformatically detectable novel gene encoding substantially pure DNA wherein: the bioinformatically detectable novel

gene does not encode a protein, a function of the novel gene is promotion of expression of the at lease one target gene, and the at least one target gene is bioinformatically deducible.

[0018] Further in accordance with a preferred embodiment of the present invention the function of the novel gene is bioinformatically deducible from the following data elements: the nucleotide sequence of the RNA encoded by the bioinformatically detectable novel gene, a nucleotide sequence of the at least one target gene, and a function of the at least one target gene.

[0019] Still further in accordance with a preferred embodiment of the present invention the RNA encoded by the novel gene complementarily binds the binding site associated with the at least one target gene, thereby modulating expression of the at least one target gene.

[0020] Additionally in accordance with a preferred embodiment of the present invention the binding site associated with at least one target gene is located in an untranslated region of RNA encoded by the at least one target gene.

[0021] Moreover in accordance with a preferred embodiment of the present invention the function of the novel gene is selective inhibition of translation of the at least one target

gene, which selective inhibition includes complementary hybridization of the RNA encoded by the novel gene to the binding site.

[0022] Further in accordance with a preferred embodiment of the present invention the invention includes a vector including the DNA.

[0023] Still further in accordance with a preferred embodiment of the present invention the invention includes a method of selectively inhibiting translation of at least one gene, including introducing the vector.

[0024] Moreover in accordance with a preferred embodiment of the present invention the introducing includes utilizing RNAi pathway.

[0025] Additionally in accordance with a preferred embodiment of the present invention the invention includes a gene expression inhibition system including: the vector, and a vector inserter, functional to insert the vector into a cell, thereby selectively inhibiting translation of at least one gene.

[0026] Further in accordance with a preferred embodiment of the present invention the invention includes a probe including the DNA.

[0027] Still further in accordance with a preferred embodiment of

the present invention the invention includes a method of selectively detecting expression of at least one gene, including using the probe.

[0028] Additionally in accordance with a preferred embodiment of the present invention the invention includes a gene expression detection system including: the probe, and a gene expression detector functional to selectively detect expression of at least one gene.

Brief Description of Drawings

[0029] Fig.1 is a simplified diagram illustrating the genomic differentiation enigma that the present invention addresses;

[0030] Figs. 2 through 4 are schematic diagrams which when taken together provide an analogy that illustrates a conceptual model of the present invention, addressing the genomic differentiation enigma;

[0031] Figs. 5A and 5B are schematic diagrams, which when taken together illustrate a "genomic records" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma;

[0032] Fig. 6 is a schematic diagram illustrating a "genomically programmed cell differentiation" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma;

[0033] Fig. 7 is a schematic diagram illustrating a "genomically programmed cell-specific protein expression modulation" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma;

[0034] Fig. 8 is a simplified diagram illustrating a mode by which genes of a novel group of genes of the present invention, modulate expression of known target genes;

[0035] Fig. 9 is a simplified block diagram illustrating a bioinformatic gene detection system capable of detecting genes of the novel group of genes of the present invention, which system is constructed and operative in accordance with a preferred embodiment of the present invention;

[0036] Fig. 10 is a simplified flowchart illustrating operation of a mechanism for training of a computer system to recognize the novel genes of the present invention, which mechanism is constructed and operative in accordance with a preferred embodiment of the present invention;

[0037] Fig. 11A is a simplified block diagram of a non-coding genomic sequence detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0038] Fig. 11B is a simplified flowchart illustrating operation of a non-coding genomic sequence detector constructed and

operative in accordance with a preferred embodiment of the present invention;

[0039] Fig. 12A is a simplified block diagram of a hairpin detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0040] Fig. 12B is a simplified flowchart illustrating operation of a hairpin detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0041] Fig. 13A is a simplified block diagram of a dicer-cut location detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0042] Fig. 13 B is a simplified flowchart illustrating training of a dicer-cut location detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0043] Fig. 14A is a simplified block diagram of a target-gene binding-site detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0044] Fig. 14B is a simplified flowchart illustrating operation of a target-gene binding-site detector constructed and operative in accordance with a preferred embodiment of the present invention;

[0045] Fig. 15 is a simplified flowchart illustrating operation of a function & utility analyzer constructed and operative in accordance with a preferred embodiment of the present invention;

[0046] Fig. 16 is an annotated sequence of an EST comprising a novel gene detected by the gene detection system of the present invention;

[0047] Figs. 17A and 17B are pictures of laboratory results, which when taken together demonstrate laboratory confirmation of expression of the bioinformatically detected novel gene of Fig. 16;

[0048] Fig. 18

[0049] Fig. 19

[0050] Fig. 20

[0051] Figs. 21A through 28020D are schematic diagrams illustrating sequences, functions and utilities of 28,000 specific genes of the novel group of genes of the present invention, detected using the bioinformatic gene detection system described hereinabove with reference to Figs. 8 through 15;

[0052] and Figs. 28021A through 3121B are schematic diagrams illustrating sequences, functions and utilities of 3,000

specific genes of a second group of novel genes of the present invention, detected using the bioinformatic gene detection system described hereinabove with reference to Figs. 8 through 15.

Brief Description of Sequences

[0053] A Sequence Listing of genomic sequences of the present invention is attached to this application, enclosed in computer readable form.

Detailed Description

[0054] Reference is now made to Fig. 1 which is a simplified diagram providing a conceptual explanation of a genomic differentiation enigma, which the present invention addresses.

[0055] Referring to Fig. 1, it is appreciated that different cell types in an organism, such as CARTILAGE CELL, LIVER CELL, FIBROBLAST CELL and BONE CELL all contain identical DNA, and derive from the initial FERTILIZED EGG CELL, and yet each of these cells expresses different proteins, and hence acquires different shape and function.

[0056] The present invention proposes that the inevitable conclusion from this constraint is, however, strikingly simple: the coding system used must be modular. It must com-

prise multiple modules, or records, one for each cell-type, and a mechanism whereby each cell at its inception is instructed which record to open, and behaves according to instructions in that record.

[0057] This modular code concept is somewhat difficult to grasp, since we are strongly habituated to viewing things from an external viewpoint. An architect, for example, looks at a blueprint of a building, which details exactly where each element (block, window, door, electrical switch, etc.) is to be placed relative to all other elements, and then instructs builders to place these elements in their designated places. This is an external viewpoint: the architect is external to the blueprint, which itself is external to the physical building, and its different elements. The architect may therefore act as an "external organizing agent": seeing the full picture and the relationships between all elements, and being able to instruct from the outside where to place each of them.

[0058] Genomics differentiation coding evidently works differently, without any such external organizing agent: It comprises only one smart block (the first cell), which is the architect and the blueprint, and which continuously duplicates itself, somehow knowing when to manifest itself as

a block and when as a window, door, or electrical switch.

[0059] Reference is now made to Figs. 2 through 4 which are schematic diagrams which when taken together provide an analogy that illustrates a conceptual model of the present invention, addressing the genomic differentiation enigma.

[0060] Referring to Fig. 2A, imagine a very talented chef, capable of preparing any meal provided he is given specific written cooking instructions. This chef is equipped with two items: (a) a thick recipe book, and (b) a small note with a number scribbled on it. The book comprises multiple pages, each page detailing how to prepare a specific meal. The small note indicates the page to be opened, and therefore the meal to be prepared. The chef looks at the page-number written on the note, opens the recipe book at the appropriate page, and prepares the meal according to the written instructions on this page. As an example, Fig. 2A depicts a CHEF holding a note with the number 12 written on it, he opens the book on page 12, and since that page contains the recipe for preparing BREAD, the CHEF prepares a loaf of BREAD.

[0061] Referring now to Fig. 2B, which depicts two identical chefs, CHEF A and CHEF B, holding an identical recipe

book. Despite their identity, and the identity of their recipe book, since CHEF A holds a note numbered 12, and therefore opens the book on page 12 and prepares BREAD, whereas CHEF B holds a note numbered 34 and therefore opens the book on page 34 and prepares a PIE.

[0062] Referring now to Fig. 3. Imagine the chef of the analogy is also capable of duplicating himself once he has finished preparing the specified meal. The format of the book is such that at the bottom of each page, two numbers are written. When he has finished preparing the meal specified on that page, the chef is trained to do the following: (i) divide himself into two identical duplicate chefs, (ii) duplicate the recipe book and hand a copy to each of his duplicate chefs, and (iii) write down the two numbers found at the bottom of the page of the meal he prepared, on two small notes, handing one note to each of his two duplicate chefs.

[0063] Each of the two resulting duplicate chefs are now equipped with the same book, and have the same talent to prepare any meal, but since each of them received a different note, they will now prepare different meals.

[0064] Fig. 3 depicts CHEF A holding a recipe book and receiving a note numbered 12. CHEF A therefore opens the book on

page 12 and prepares BREAD. When he is finished making bread, CHEF A performs the following actions: (i) divides himself into two duplicate chefs, designated CHEF B and CHEF C, (ii) duplicates his recipe book handing a copy to each of CHEF B and CHEF C, (iii) writes down the numbers found at the bottom of page 12, numbers 34 and 57, on two notes, handing note numbered 34 to CHEF B and note numbered 57 to CHEF C.

[0065] Accordingly, CHEF B receives a note numbered 34 and therefore opens the recipe book on page 34 and prepares PIE, whereas CHEF C receives a note numbered 57 and therefore opens the book on page 57 and therefore prepares RICE.

[0066] It is appreciated that while CHEF A, CHEF B & CHEF C are identical and hold identical recipe books, they each prepare a different meal. It is also appreciated that the meals prepared by CHEF B and CHEF C are determined CHEF A, and are mediated by the differently numbered notes passed on from CHEF A to CHEF B and CHEF C.

[0067] It is further appreciated that the mechanism illustrated by Fig. 3 enables an unlimited lineage of chefs to divide into duplicate, identical chefs and to determine the meals those duplicate chefs would prepare. For example, having

been directed to page 34, when CHEF B divides into duplicate chefs (not shown), he will instruct its two duplicate chefs to prepare meals specified on pages 14 and 93 respectively, according to the numbers at the bottom of page 34 to which he was directed. Similarly, CHEF C will instruct its duplicate chefs to prepare meals specified on pages 21 and 46 respectively, etc.

[0068] Referring now to Fig. 4. Imagine that the cooking instructions on each page of the recipe book are written in shorthand format: The main meal-page to which the chef was directed by the scribbled note, merely contains a list of numbers which direct him to multiple successive pages, each specifying how to prepare an ingredient of that meal.

[0069] As an example, Fig. 4 depicts CHEF A of FIGS 2 and 3, holding a recipe book and a note numbered 12. Accordingly, CHEF A opens the recipe book on page 12, which details the instructions for preparing BREAD. However, the "instructions" on making BREAD found on page 12 comprise only of 3 numbers, 18, 7 and 83, which "refer" CHEF A to pages detailing preparation of the ingredients of BREAD FLOUR, MILK and SALT, respectively.

[0070] As illustrated in Fig. 4, turning from the main "meal page" (e.g. 12) to respective "ingredients pages" (e.g. pages 18,

7 & 83) is mediated by scribbled notes with the page-numbers written on them. In this analogy, the scribbled notes are required for seeking the target pages to be turned to both when turning to main "meal pages" (e.g. page 12), as well as when turning to "ingredient pages" (e.g. pages 18, 7 & 83).

[0071] The chef in the given analogy, schematically depicted in FIGS 2 through 4, represents a cell; the thick recipe book represents the DNA; preparing a meal in the given analogy represents the cell manifesting itself as a specific cell-type; and ingredients of a meal represent proteins expressed by that cell-type. Like the chef equipped with the thick recipe book in the given analogy, all cells in an organism contain the same DNA and are therefore each potentially capable of manifesting itself as any cell-type, expressing proteins typical of that cell type.

[0072] Reference is now made to Figs. 5A and 5B which are schematic diagrams, which when taken together illustrate a "genomic records" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma.

[0073] The Genomic Records concept asserts that the DNA (the thick recipe book in the illustration) comprises a very

large number of Genomic Records (analogous to pages in the recipe book), each containing the instructions for differentiation of a different cell-type, or developmental process. Each Genomic Record is headed by a very short genomic sequence which functions as a "Genomic Address" of that Genomic Record (analogous to the page number in the recipe book). At its inception, in addition to the DNA, each cell also receives a short RNA segment (the scribbled note in the illustration). This short RNA segment binds complementarily to a "Genomic Address" sequence of one of the Genomic Records, thereby activating that Genomic Record, and accordingly determining the cell"s-fate (analogous to opening the book on the page corresponding to the number on the scribbled note, thereby determining the meal to be prepared).

[0074] Referring to Fig. 5A, a CELL is illustrated which comprises a GENOME. The GENOME comprises a plurality of GENOMIC RECORDS, each of which correlates to a specific cell type (for clarity only 6 sample genomic records are shown). Each genomic record comprises genomic instructions on differentiation into a specific cell-type, as further elaborated below with reference to Fig. 7. At cell inception, the CELL receives a maternal short RNA segment,

which activates one of the GENOMIC RECORDS, causing the cell to differentiate according to the instructions comprised in that genomic record. As an example, Fig. 5A illustrates reception of a maternal short RNA segment designated A" and outlined by a broken line, which activates the FIBRO genomic record, causing the cell to differentiate into a FIBROBLAST CELL.

[0075] Referring now to Fig. 5B, which is a simplified schematic diagram, illustrating cellular differentiation mediated by the "Genomic Records" concept. Fig. 5B depicts 2 cells in an organism, designated CELL A and CELL B, each having a GENOME. It is appreciated that since CELL A and CELL B are cells in the same organism, the GENOME of CELL A is identical to that of CELL B. Despite having an identical GENOME, CELL A differentiates differently from CELL B, due to activation of different genomic records in these two cells. In CELL A the FIBRO GENOMIC RECORD is activated, causing CELL A to differentiate into a FIBROBLAST CELL, whereas in CELL B the BONE GENOMIC RECORD is activated, causing the CELL B to differentiate into a BONE CELL. The cause for activation of different genomic records in these two cells is the different maternal short RNA which they both received: CELL A received a maternal

short RNA segment designated A" which activated genomic record FIBRO, whereas CELL B received a maternal short RNA segment designated B" which activated genomic record BONE.

[0076] Reference is now made to Fig. 6 which is a schematic diagram illustrating a "genomically programmed cell differentiation" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma.

[0077] A cell designated CELL A divides into 2 cells designated CELL B and CELL C. CELL A, CELL B and CELL C each comprise a GENOME, which GENOME comprises a plurality of GENOMIC RECORDS. It is appreciated that since CELL A, CELL B and CELL C are cells in the same organism, the GENOME of these cells, and the GENOMIC RECORDS comprised therein, are identical.

[0078] As described above with reference to Fig. 5B, at its inception, CELL A receives a maternal short RNA segment, designated A" and marked by a broken line, which activates the FIBRO genomic record, thereby causing CELL A to differentiate into a FIBROBLAST CELL. However, Fig. 6 shows further details of the genomic records: each cell genomic record also comprises two short genomic sequences, referred to here as Daughter Cell Genomic Addresses.

Blocks designated B and C are Daughter Cell Genomic Addresses of the FIBRO Genomic Record. At cell division, each parent cell transcribes two short RNA segments, corresponding to the two Daughter Cell Genomic Addresses of the Genomic Record of that parent cell, and transfers one to each of its two daughter cells. CELL A of Fig. 6 transcribes and transfers to its two respective daughter cells, two short RNA segments, outlined by a broken line and designated B" and C", corresponding to daughter cell genomic addresses designated B and C comprised in the FIBRO genomic record.

[0079] CELL B therefore receives the above mentioned maternal short RNA segment designated B", which binds complementarily to genomic address designated B of genomic record BONE, thereby activating this genomic record, which in turn causes CELL B to differentiate into a BONE CELL. Similarly, CELL C receives the above mentioned maternal short RNA segment designated C", which binds complementarily to genomic address designated C of genomic record CARTIL., thereby activating this genomic record, which in turn causes CELL C to differentiate into a CARTILAGE CELL.

[0080] It is appreciated that the mechanism illustrated by Fig. 6

enables an unlimited lineage of cells to divide into daughter cells containing the same DNA, and to determine the cell-fate of these daughter cells. For example, when CELL B and CELL C divide into their respective daughter cells (not shown), they will transfer short RNA segments designated D" & E", and F" & G" respectively, to their respective daughter cells. The cell fate of each of these daughter cells would be determined by the identity of the maternal short RNA segment they receive, which would determine the genomic record activated.

[0081] Reference is now made to Fig. 7 which is a schematic diagram illustrating a "genomically programmed cell-specific protein expression modulation" concept of the conceptual model of the present invention, addressing the genomic differentiation enigma.

[0082] Cell A receives a maternal short RNA segment designated A", which activates a genomic record designated FIBRO, by anti-sense binding to a binding site "header" of this genomic record, designated A. Genomic record FIBRO encodes 3 short RNA segments, designated 1, 2 and 4 respectively, which modulate expression of target genes designated GENE1, GENE2 and GENE4 respectively. Modulation of expression of these genes results in CELL A dif-

ferentiating into a FIBROBLAST CELL.

[0083] Reference is now made to Fig. 8 which is a simplified diagram illustrating a mode by which genes of a novel group of genes of the present invention, modulate expression of known target genes.

[0084] The novel genes of the present invention are micro RNA (miRNA)-like, regulatory RNA genes, modulating expression of known target genes. This mode of modulation is common to other known miRNA genes, as described hereinabove with reference to the background of the invention section.

[0085] GAM GENE and TARGET GENE are two human genes contained in the DNA of the human genome.

[0086] GAM GENE encodes a GAM PRECURSOR RNA. However, similar to other miRNA genes, and unlike most ordinary genes, its RNA, GAM PRECURSOR RNA, does not encode a protein.

[0087] GAM PRECURSOR RNA folds onto itself, forming GAM FOLDED PRECURSOR RNA. As Fig.8 illustrates, GAM FOLDED PRECURSOR RNA forms a "hairpin structure", folding onto itself. As is well known in the art, this "hairpin structure", is typical genes of the miRNA genes, and is due to the fact that nucleotide sequence of the first half of

the RNA of a gene in this group is an accurate or partial inversed-reversed sequence of the nucleotide sequence of its second half. By "inversed-reversed"is meant a sequence which is reversed and wherein each nucleotide is replaced by a complimentary nucleotide, as is well known in the art (e.g. ATGGC is the inversed-reversed sequence of GCCAT).

[0088] An enzyme complex, designated DICER COMPLEX, "dices" the GAM FOLDED PRECURSOR RNA into a single stranded RNA segment, about 22 nucleotides long, designated GAM RNA. As is known in the art, "dicing" of the hairpin structured RNA precursor into shorter RNA segments about 22 nucleotides long by a Dicer type enzyme is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins.

[0089] TARGET GENE encodes a corresponding messenger RNA, designated TARGET RNA. This TARGET RNA comprises 3 regions: a 5" untranslated region, a protein coding region and a 3" untranslated region, designated 5"UTR, PROTEIN CODING and 3"UTR respectively.

[0090] GAM RNA binds complimentarily a BINDING SITE, located on the 3"UTR segment of TARGET RNA. This complimentarily binding is due to the fact that the nucleotide se-

quence of GAM RNA is an accurate or partial inversed-reversed sequence of the nucleotide sequence of BINDING SITE.

[0091] The complimentary binding of GAM RNA to BINDING SITE inhibits translation of TARGET RNA into TARGET PROTEIN. TARGET PROTEIN is therefore outlined by a broken line.

[0092] It is appreciated by one skilled in the art that the mode of transcriptional inhibition illustrated by Fig. 1 with specific reference to GAM genes of the present invention, is in fact common to all other miRNA genes. A specific complimentary binding site has been demonstrated only for Lin-4 and Let-7. All the other 93 newly discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complimentary binding, although specific complimentary binding sites for these genes have not yet been found (Ruvkun G., "Perspective: Glimpses of a tiny RNA world", Science 294 ,779 (2001)). The present invention discloses a novel group of genes, the GAM genes, belonging to the miRNA genes group, and for which a specific an complimentary binding has been determined.

[0093] Reference is now made to Fig. 9 which is a simplified block diagram illustrating a bioinformatic gene detection

system capable of detecting genes of the novel group of genes of the present invention, which system is constructed and operative in accordance with a preferred embodiment of the present invention.

[0094] A bioinformatic gene detection engine 100, is a preferred implementation of a mechanism capable of detecting genes of the novel group of genes of the present invention.

[0095] The function of the bioinformatic gene detection engine 100 is as follows: it receives three types of input, expressed RNA data 102, sequenced DNA data 104, and protein function data 106, performs a complex process of analysis of this data as elaborated below, and based on this analysis produces output of a bioinformatically detected group of novel genes designated 108.

[0096] Expressed RNA data 102 comprises published expressed sequence tags (EST) data, , published mRNA data, as well as other sources of published RNA data. Sequenced DNA data 104 comprises alphanumeric data describing sequenced genomic data, which preferably includes annotation data such as location of known protein coding regions relative to the sequenced data. Protein function data 106 comprises scientific publications reporting studies

which elucidated physiological function known proteins, and their connection, involvement and possible utility in treatment and diagnosis of various diseases. Expressed RNA data 102, sequenced DNA data 104 may preferably be obtained from data published by the National Center for Bioinformatics (NCBI) at the National Institute of Health (NIH), as well as from various other published data sources. Protein function data 106 may preferably be obtained from any one of numerous relevant published data sources, such as the Online Mendelian Inherited Disease In Man (OMIM) database developed by John Hopkins University, and also published by NCBI.

[0097] Prior to actual detection of bioinformatically detected novel genes 108 by the bioinformatic gene detection engine 100, a process of bioinformatic gene detection engine training & validation designated 110 takes place. This process uses the known miRNA genes as a training set (some 200 such genes have been found to date using biological laboratory means), to train the bioinformatic gene detection engine 100 to bioinformatically recognize miRNA-like genes, and their respective potential target binding sites. Bioinformatic gene detection engine training & validation 110 is further describe hereinbelow with ref-

erence to Fig. 10.

[0098] The bioinformatic gene detection engine 100 comprises several modules which are preferably activated sequentially, and are described as follows:A non-coding genomic sequence detector 112 operative to screen for and identify non-protein coding genomic sequences. The non-coding genomic sequence detector 112 is further described hereinbelow with reference to Figs. 11A and 11B.

[0099] A hairpin detector 114 operative to locate genomic "hairpin-shaped" sequences, similar to GAM FOLDED PRECURSOR of Fig. 8. The hairpin detector 114 is further described hereinbelow with reference to Figs. 12A and 12B.

[0100] A dicer-cut location detector 116 operative to detect the location on a hairpin shaped sequence which is enzymatically cut by DICER COMPLEX of Fig. 8. The dicer-cut location detector 116 is further described hereinbelow with reference to Fig. 13A.

[0101] A target-gene binding-site detector 118 operative to detect In these ESTs, the system locates "hairpin-shaped" (i.e. hairpin) sequences, which resemble known miRNA sequences. RNA 2D folding is calculated based on free-energy, using the Zucker algorithm (used throughout the process) [reference]. The 2D structure is then evaluated to

locate hairpin structures, and meticulously compared to that of known miRNA genes.

[0102] A combined neural and Bayesian networks system is used to detect location in which the Dicer enzyme-complex is predicted to cut the final novel miRNA gene from the hairpin shaped gene-precursor. Nucleotide sequence, and "Bulges" (i.e. mismatches) on the hairpin shaped precursor, are represented as the input and output layers of several neural networks, and hidden layer/s are designed for these neural networks accordingly. The networks are trained, evaluated and validated based on the "training set" of 120 known miRNA genes, using accepted neural networks methodology; prediction is further optimized by integrating results of multiple networks [reference].

[0103] Potential target-gene binding-sites are detected, by comparison of the final-gene sequence, calculated above, to untranslated regions of known genes, followed by a comparison of the 2D structure and free-energy of the binding site to that of known miRNA binding sites.

[0104] Lastly, biological and clinical function of target genes, utilizing reference databases such as Johns Hopkins" OMIM [reference] is manually reviewed, and is compared to tissue-data of EST in which novel gene was detected.

[0105] Referring to Fig. 9, the following is a technical description of the computer hardware configuration which was used:Hardware and Software SetupComputing power presented a major challenge. Initial time estimates indicated that over 30 months (!) of computing time would be required to detect all miRNA genes in human EST data, and their respective binding sites, using a powerful 8-processor machine.

[0106] In order to address this challenge at reasonable cost, a system has been built which, comprises a cluster of one hundred PCs (Pentium IV, 1.7GHz, with 40GB storage each), connected by Ethernet to 4 servers (2-CPU, Xeon 2.2GHz, with 200GB storage each), combined with an 8-processor server (8-CPU, Xeon 550Mhz w/ 8GB RAM) connected via 2 HBA fiber-channels to an EMC Clariion 100-disks, 3.6 Terabyte storage device. Figure 2 provides an overview of this hardware configuration.

[0107] Using this configuration, a database comprising 30 billion records was created, using Microsoft"s SQL-Server database software, demonstrating remarkable performance results, including bulk insert of 1 million records in 0.8 seconds. To the best of our knowledge, this is the largest number-of-records database ever built on an In-

tel-machine, achieving unprecedented record-insert performance. The system has further been optimized such that all one hundred PCs run at 100% CPU usage continually. Using this configuration reduced the calculation time from 30 months to 20 days.

[0108] Reference is now made to Fig. 10 which is a simplified flowchart illustrating operation of a mechanism for training of a computer system to recognize the novel genes of the present invention. This mechanism is a preferred implementation of the bioinformatic gene detection engine training & validation 110 described hereinabove with reference to Fig. 9.

[0109] Reference is now made to Fig. 11A which is a simplified block diagram of a preferred implementation of the non-coding genomic sequence detector 112 described hereinabove with reference to Fig. 9.

[0110] *The system uses as input published genomic data, including expressed RNA data (EST data and mRNA data), sequenced DNA data, and protein function data, from NCBI. EST data is blast-compared [reference] to known protein coding sequences; only non-protein-coding ESTs are searched for novel RNA genes.* Reference is now made to Fig. 11B which is a simplified flowchart illustrating a preferred operation of the non-coding ge-

nomic sequence detector 112 of Fig. 9.

[0111] Reference is now made to Fig. 12A which is a simplified block diagram of a preferred implementation of the hairpin detector 114 described hereinabove with reference to Fig. 9.

[0112] Reference is now made to Fig. 12B which is a simplified flowchart illustrating a preferred operation of the hairpin detector 114 of Fig. 9.

[0113] Reference is now made to Fig. 13A which is a simplified block diagram of a preferred implementation of the dicer-cut location detector 116 described hereinabove with reference to Fig. 9.

[0114] Reference is now made to Fig. 13 B which is a simplified flowchart illustrating a preferred implementation of dicer-cut location detector training & validation 126 of Fig. 10.

[0115] Reference is now made to Fig. 14A which is a simplified block diagram of a preferred implementation of the target-gene binding-site detector 118 described hereinabove with reference to Fig. 9.

[0116] Reference is now made to Fig. 14B which is a simplified flowchart illustrating a preferred operation of the target-gene binding-site detector 118 of Fig. 9.

[0117] Reference is now made to Fig. 15 which is a simplified

flowchart illustrating a preferred operation of the function & utility analyzer 120 described hereinabove with reference to Fig. 9.

[0118] Reference is now made to Fig. 16 which is an annotated sequence of an EST comprising a novel gene detected by the gene detection system of the present invention. Fig. 16 shows the nucleotide sequence of a known human non-protein coding EST (Expressed Sequence Tag), identified as EST72223. It is appreciated that the sequence of this EST comprises sequences of one known miRNA gene, identified as MIR98, and of one novel GAM gene, referred to here as GAM-4, detected by the bioinformatic gene detection system of the present invention, described hereinabove with reference to Fig. 9.

[0119] Reference is now made to Figs. 17A and 17B that are pictures of laboratory results, which when taken together demonstrate laboratory confirmation of expression of the bioinformatically detected novel gene of Fig. 16.

[0120] Referring to Fig. 17A which is a Northern blot analysis of MIR-98 and EST72223 transcripts. MIR-98 and EST72223 were reacted with MIR-98 and GAM-4 probes as indicated in the figure. It is appreciated that the probes of both MIR-98 and GAM-4 reacted with EST72223, indicating

that EST72223 contains the sequences of MIR-98 and of GAM-4. It is further appreciated that the probe of GAM-4 did not cross-react with MIR-98.

[0121] With reference to Fig. 17B, a Northern blot analysis of EST72223 and MIR-98 transfections were performed, subsequently marking RNA by the MIR-98 and GAM-4 probes . Left, Northern reacted with MIR-98, Right, Northern reacted with GAM-4. The molecular Sizes of EST72223, MIR-98 and GAM-4 are indicated by arrows. Hela are control cells that have not been introduced to exogenous RNA. EST and MIR-98 Transfections are RNA obtained from Hela transfected with EST72223 and MIR-98, respectively. MIR-98 and EST are the transcripts used for the transfection experiment. The results indicate that EST72223, when transfected into Hela cells, is cut yielding known miRNA gene MIR-98 and novel miRNA gene GAM-4.

[0122] Referring to Figs. 17A and 17B, the following technical methods used are specified as follows: Transcript preparations: Digoxigenin (DIG) labeled transcripts were prepared from EST72223 (TIGER), MIR98 and predicted precursor hairpins by using a DIG RNA labeling kit (Roche Molecular Biochemicals) according to the manufacture"s

protocol. Briefly, PCR products with T7 promoter at the 5" end or T3 promoter at the 3"end were prepared from each DNA in order to use it as a template to prepare sense and antisense transcripts, respectively.

[0123] MIR-98 was amplified using EST72223 as a templet with T7miR98 forward primer: 5-"TAATACGACTCACTATAGGGTGAGGTAGTAAGTTGTATTGTT-3"and T3miR98 revse primer: 5"-AATTAACCCTCACTAAAGGGAAAGTAGTAAGTTGTATAGTT-3"EST72223 was amplified with T7-EST 72223 forward primer:5"-TAATACGACTCACTATAGGCCCTTATTAGAGGATTCTGCT-3"and T3-EST72223 reverse primer:5"-AATTAACCCTCACTAAAGGTTTTTTTTTCCTGAGACAGAGT-3" Bet-4 was amplified using EST72223 as a templet with Bet-4 forward primer: 5"-GAGGCAGGAGAATTGCTTGA- 3" and T3-EST72223 reverse primer:5"-AATTAACCCTCACTAAAGGCCTGAGACAGAGTCTTGCTC-3"The PCR products were cleaned and used for DIG-labeled or unlabeled transcription reactions with the appropriate polymerase. For transfection experiments, CAP reaction was performed by using a mMassage mMachine kit (Ambion).

[0124] Transfection procedure: Transfection of Hela cells was performed by using TransMessenger reagent (Qiagen) according to the manufacture"s protocol. Briefly, Hela cells were seeded to 1-2x 10^6 cells per plate a day before transfection. Two µg RNA transcripts were mixed with 8µl Enhancer in a final volume of 100µl, mixed and incubated at room temperature for 5 min. 16µl TransMessenger reagent was added to the RNA-Enhancer, mixed and incubated for additional 10 min. Cell plates were washed with sterile PBS twice and then incubated with the transfection mix diluted with 2.5 ml DMEM medium without serum. Cells were incubated with transfection mix for three hours under their normal growth condition (370C and 5% CO2) before the transfection mix was removed and a fresh DMEM medium containing serum was added to the cells. Cells were left to grow 48 hours before harvesting.

[0125] *Northern analysis:*RNAs were extracted from cells by using Tri-reagent according to the manufacture"s protocol. The RNAs were dissolved in water and heated to 650C to disrupt any association of the 25nt RNA with larger RNA molecules. RNA were placed on ice and incubated for 30 min with PEG (MW=8000) in a final concentration of 5% and NaCl in a final concentration of 0.5M to precipitate

high molecular weight nucleic acid. The RNAs were centrifuged at 10,000xg for 10 min to pellet the high molecular weight nucleic acid. The supernatant containing the low molecular weight RNAs was collected and three volumes of ethanol was added. The RNAs were placed at -200C for at least two hours and then centrifuged at 10,000xg for 10 min. The pellets were dissolved in Urea-TBE buffer (1Xtbe, 7M urea) for further analysis by a Northern blot.

[0126] RNA samples were boiled for 5 min before loading on 15%-8% polyacrylamide (19:1) gels containing 7M urea and 1xTBE. Gels were run in 1xTBE at a constant voltage of 300V and then transferred into a nylon membrane. The membrane was exposed to 3min ultraviolet light to cross link the RNAs to the membrane. Hybridization was performed overnight with DIG-labeled probes at 420C. Membranes were washed twice with SSCx2 and 0.2% SDS for 10 min. at 420C and then washed twice with SSCx0.5 for 5 min at room temperature. The membrane was then developed by using a DIG luminescent detection kit (Roche) using anti DIG and CSPD reaction, according to the manufacture"s protocol.

[0127] Reference is now made to Fig. 8, which is a simplified dia-

gram describing each of a plurality of novel bioinformatically detected genes of the present invention, referred to here as Genomic Address Messenger (GAM) genes, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0128] GAM is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM was detected is described hereinabove with reference to Figs. 2-8.

[0129] GAM GENE and GAM TARGET GENE are human genes contained in the human genome.

[0130] GAM GENE encodes a GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM PRECURSOR RNA does not encode a protein.

[0131] GAM PRECURSOR RNA folds onto itself, forming GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof. By inversed-reversed

is meant a sequence which is reversed and wherein each nucleotide is replaced by a complementary nucleotide, as is well known in the art (e.g. ATGGC is the inversed-reversed sequence of GCCAT).

[0132] An enzyme complex designated DICER COMPLEX, dices the GAM FOLDED PRECURSOR RNA into GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins.

[0133] GAM TARGET GENE encodes a corresponding messenger RNA, GAM TARGET RNA. GAM TARGET RNA comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0134] GAM RNA binds complementarily to one or more target binding sites located in untranslated regions of GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM RNA is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illus-

tration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM RNA may have a different number of target binding sites in untranslated regions of a GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0135] The complementary binding of GAM RNA to target binding sites on GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM TARGET RNA into GAM TARGET PROTEIN. GAM TARGET PROTEIN is therefore outlined by a broken line.

[0136] It is appreciated that GAM TARGET GENE in fact represents a plurality of GAM target genes. The mRNA of each one of this plurality of GAM target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM RNA, and which when bound by GAM RNA causes inhibition of translation of respective one or more GAM target proteins.

[0137] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM GENE on one or more GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0138] It is yet further appreciated that specific functions, and accordingly utilities, of GAM correlate with, and may be deduced from, the identity of the target genes which GAM binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0139] Nucleotide sequences of the GAM PRECURSOR RNA, and of the diced GAM RNA, and a schematic representation of the secondary folding of GAM FOLDED PRECURSOR RNA of

each of the plurality of GAM GENEs described by Fig. 8 are further described hereinbelow with reference to Table.

[0140] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM RNA are described hereinbelow with reference to Table.

[0141]

[0142] Fig. 8 further provides a conceptual description of a novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 24 (GAM24) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0143] GAM24 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM24 was detected is described hereinabove with reference to Figs. 2-8.

[0144] GAM24 gene, herein designated GAM GENE, and GAM24 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0145] GAM24 gene, herein designated GAM GENE, encodes a

GAM24 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM24 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM24 precursor RNA is designated SEQ ID:1, and is provided hereinbelow with reference to the sequence listing part.

[0146] GAM24 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM24 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0147] An enzyme complex designated DICER COMPLEX, dices the GAM24 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM24 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured

RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM24 RNA is designated SEQ ID:2, and is provided hereinbelow with reference to the sequence listing part.

[0148] GAM24 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM24 target RNA, herein designated GAM TARGET RNA. GAM24 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0149] GAM24 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM24 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM24 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM24 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM24 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0150] The complementary binding of GAM24 RNA, herein designated GAM RNA, to target binding sites on GAM24 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM24 target RNA, herein designated GAM TARGET RNA, into GAM24 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0151] It is appreciated that GAM24 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM24 target genes. The mRNA of each one of this plurality of GAM24 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM24 RNA, herein designated GAM RNA, and which when bound by GAM24 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM24 target proteins.

[0152] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM24 gene, herein designated GAM GENE, on one or more GAM24 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0153] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM24 correlate with, and may be deduced from, the identity of the target genes which GAM24 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0154] Nucleotide sequences of the GAM24 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM24 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM24 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM24 are further described hereinbelow with reference to Table Table1.

[0155] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM24 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0156]

[0157] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 25 (GAM25) gene, which modulates expression of respective target genes thereof, the function and utility of which

target genes is known in the art.

[0158] GAM25 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM25 was detected is described hereinabove with reference to Figs. 2-8.

[0159] GAM25 gene, herein designated GAM GENE, and GAM25 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0160] GAM25 gene, herein designated GAM GENE, encodes a GAM25 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM25 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM25 precursor RNA is designated SEQ ID:11, and is provided hereinbelow with reference to the sequence listing part.

[0161] GAM25 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM25 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0162] An enzyme complex designated DICER COMPLEX, dices the GAM25 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM25 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM25 RNA is designated SEQ ID:12, and is provided hereinbelow with reference to the sequence listing part.

[0163] GAM25 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM25 target RNA, herein designated GAM TARGET RNA. GAM25 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0164] GAM25 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM25 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM25 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM25 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM25 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0165] The complementary binding of GAM25 RNA, herein designated GAM RNA, to target binding sites on GAM25 target RNA, herein designated GAM TARGET RNA, such as BIND-

ING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM25 target RNA, herein designated GAM TARGET RNA, into GAM25 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0166] It is appreciated that GAM25 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM25 target genes. The mRNA of each one of this plurality of GAM25 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM25 RNA, herein designated GAM RNA, and which when bound by GAM25 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM25 target proteins.

[0167] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM25 gene, herein designated GAM GENE, on one or more GAM25 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site

has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0168] It is yet further appreciated that specific functions, and accordingly utilities, of GAM25 correlate with, and may be deduced from, the identity of the target genes which GAM25 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0169] Nucleotide sequences of the GAM25 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM25 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM25 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM25 are further described hereinbelow with reference to Table Table1.

[0170] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM25 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0171]

[0172] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 26 (GAM26) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0173] GAM26 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM26 was detected is described hereinabove with reference to Figs. 2-8.

[0174] GAM26 gene, herein designated GAM GENE, and GAM26 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0175] GAM26 gene, herein designated GAM GENE, encodes a GAM26 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM26 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM26 precursor RNA is designated SEQ ID:36, and is provided hereinbelow with reference to the sequence listing part.

[0176] GAM26 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM26 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0177] An enzyme complex designated DICER COMPLEX, dices the GAM26 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM26 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM26 RNA is designated SEQ ID:37, and is provided hereinbelow with

reference to the sequence listing part.

[0178] GAM26 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM26 target RNA, herein designated GAM TARGET RNA. GAM26 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0179] GAM26 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM26 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM26 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM26 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM26 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0180] The complementary binding of GAM26 RNA, herein designated GAM RNA, to target binding sites on GAM26 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM26 target RNA, herein designated GAM TARGET RNA, into GAM26 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0181] It is appreciated that GAM26 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM26 target genes. The mRNA of each one of this plurality of GAM26 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM26 RNA, herein designated GAM RNA, and which when bound by GAM26 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM26 tar-

get proteins.

[0182] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM26 gene, herein designated GAM GENE, on one or more GAM26 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0183] It is yet further appreciated that specific functions, and accordingly utilities, of GAM26 correlate with, and may be deduced from, the identity of the target genes which GAM26 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0184] Nucleotide sequences of the GAM26 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM26 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM26 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM26 are further described hereinbelow with reference to Table Table1.

[0185] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM26 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0186]

[0187] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 27 (GAM27) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0188] GAM27 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM27 was detected is described hereinabove with reference to Figs. 2-8.

[0189] GAM27 gene, herein designated GAM GENE, and GAM27 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0190] GAM27 gene, herein designated GAM GENE, encodes a GAM27 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM27 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM27 precursor RNA is designated SEQ ID:46, and is provided hereinbelow with reference to the sequence listing part.

[0191] GAM27 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM27 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0192] An enzyme complex designated DICER COMPLEX, dices

the GAM27 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM27 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM27 RNA is designated SEQ ID:47, and is provided hereinbelow with reference to the sequence listing part.

[0193] GAM27 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM27 target RNA, herein designated GAM TARGET RNA. GAM27 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0194] GAM27 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM27 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM27

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM27 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM27 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0195] The complementary binding of GAM27 RNA, herein designated GAM RNA, to target binding sites on GAM27 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM27 target RNA, herein designated GAM TARGET RNA, into GAM27 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0196] It is appreciated that GAM27 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM27 target genes. The mRNA of each one of this plurality of GAM27 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM27 RNA, herein designated GAM RNA, and which when bound by GAM27 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM27 target proteins.

[0197] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM27 gene, herein designated GAM GENE, on one or more GAM27 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complemen-

tary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0198] It is yet further appreciated that specific functions, and accordingly utilities, of GAM27 correlate with, and may be deduced from, the identity of the target genes which GAM27 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0199] Nucleotide sequences of the GAM27 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM27 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM27 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM27 are further described hereinbelow with reference to Table Table1.

[0200] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM27 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0201]

[0202] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 28 (GAM28) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0203] GAM28 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM28 was detected is described hereinabove with reference to Figs. 2-8.

[0204] GAM28 gene, herein designated GAM GENE, and GAM28 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0205] GAM28 gene, herein designated GAM GENE, encodes a GAM28 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM28 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM28 precursor RNA is designated SEQ ID:54, and is provided hereinbelow with reference to the sequence listing part.

[0206] GAM28 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM28 folded precur-

sor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0207] An enzyme complex designated DICER COMPLEX, dices the GAM28 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM28 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM28 RNA is designated SEQ ID:55, and is provided hereinbelow with reference to the sequence listing part.

[0208] GAM28 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM28 target RNA, herein designated GAM TARGET RNA. GAM28 target RNA, herein designated GAM TARGET RNA, com-

prises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0209] GAM28 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM28 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM28 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM28 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM28 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[0210] The complementary binding of GAM28 RNA, herein designated GAM RNA, to target binding sites on GAM28 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM28 target RNA, herein designated GAM TARGET RNA, into GAM28 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0211] It is appreciated that GAM28 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM28 target genes. The mRNA of each one of this plurality of GAM28 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM28 RNA, herein designated GAM RNA, and which when bound by GAM28 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM28 target proteins.

[0212] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM28 gene, herein designated GAM GENE, on one or

more GAM28 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0213] It is yet further appreciated that specific functions, and accordingly utilities, of GAM28 correlate with, and may be deduced from, the identity of the target genes which GAM28 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0214] Nucleotide sequences of the GAM28 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM28 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM28 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM28 are further described

hereinbelow with reference to Table Table1.

[0215] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM28 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0216]

[0217] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 29 (GAM29) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0218] GAM29 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM29 was detected is described hereinabove with reference to Figs. 2-8.

[0219] GAM29 gene, herein designated GAM GENE, and GAM29 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0220] GAM29 gene, herein designated GAM GENE, encodes a GAM29 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM29 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM29 precursor RNA is designated SEQ ID:67, and is provided hereinbelow with reference to the sequence listing part.

[0221] GAM29 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM29 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0222] An enzyme complex designated DICER COMPLEX, dices the GAM29 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM29 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is

catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 81%) nucleotide sequence of GAM29 RNA is designated SEQ ID:68, and is provided hereinbelow with reference to the sequence listing part.

[0223] GAM29 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM29 target RNA, herein designated GAM TARGET RNA. GAM29 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0224] GAM29 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM29 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM29 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM29 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM29 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0225] The complementary binding of GAM29 RNA, herein designated GAM RNA, to target binding sites on GAM29 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM29 target RNA, herein designated GAM TARGET RNA, into GAM29 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0226] It is appreciated that GAM29 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM29 target genes. The mRNA of each one of this plurality of GAM29 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM29 RNA, herein designated GAM RNA, and which when bound by GAM29 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM29 target proteins.

[0227] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM29 gene, herein designated GAM GENE, on one or more GAM29 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0228] It is yet further appreciated that specific functions, and accordingly utilities, of GAM29 correlate with, and may be

deduced from, the identity of the target genes which GAM29 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0229] Nucleotide sequences of the GAM29 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM29 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM29 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM29 are further described hereinbelow with reference to Table Table1.

[0230] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM29 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0231]

[0232] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 30 (GAM30) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0233] GAM30 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM30 was detected is described hereinabove with reference to Figs. 2-8.

[0234] GAM30 gene, herein designated GAM GENE, and GAM30 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0235] GAM30 gene, herein designated GAM GENE, encodes a GAM30 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM30 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM30 precursor RNA is designated SEQ ID:97, and is provided hereinbelow with reference to the sequence listing part.

[0236] GAM30 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM30 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0237] An enzyme complex designated DICER COMPLEX, dices the GAM30 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM30 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM30 RNA is designated SEQ ID:98, and is provided hereinbelow with reference to the sequence listing part.

[0238] GAM30 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM30 target RNA, herein designated GAM TARGET RNA. GAM30 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0239] GAM30 RNA, herein designated GAM RNA, binds comple-

mentarily to one or more target binding sites located in untranslated regions of GAM30 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM30 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM30 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM30 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0240] The complementary binding of GAM30 RNA, herein designated GAM RNA, to target binding sites on GAM30 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits

translation of GAM30 target RNA, herein designated GAM TARGET RNA, into GAM30 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0241] It is appreciated that GAM30 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM30 target genes. The mRNA of each one of this plurality of GAM30 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM30 RNA, herein designated GAM RNA, and which when bound by GAM30 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM30 target proteins.

[0242] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM30 gene, herein designated GAM GENE, on one or more GAM30 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known

miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0243] It is yet further appreciated that specific functions, and accordingly utilities, of GAM30 correlate with, and may be deduced from, the identity of the target genes which GAM30 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0244] Nucleotide sequences of the GAM30 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM30 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM30 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM30 are further described hereinbelow with reference to Table Table1.

[0245] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM30 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0246]

[0247] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 31 (GAM31) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0248] GAM31 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM31 was detected is described hereinabove with reference to Figs. 2-8.

[0249] GAM31 gene, herein designated GAM GENE, and GAM31 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0250] GAM31 gene, herein designated GAM GENE, encodes a GAM31 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM31 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM31 precursor RNA is designated

SEQ ID:119, and is provided hereinbelow with reference to the sequence listing part.

[0251] GAM31 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM31 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0252] An enzyme complex designated DICER COMPLEX, dices the GAM31 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM31 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM31 RNA is designated SEQ ID:120, and is provided hereinbelow with reference to the sequence listing part.

[0253] GAM31 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM31 target RNA, herein designated GAM TARGET RNA. GAM31 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0254] GAM31 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM31 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM31 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM31 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM31 tar-

get RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0255] The complementary binding of GAM31 RNA, herein designated GAM RNA, to target binding sites on GAM31 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM31 target RNA, herein designated GAM TARGET RNA, into GAM31 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0256] It is appreciated that GAM31 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM31 target genes. The mRNA of each one of this plurality of GAM31 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM31 RNA, herein designated GAM RNA, and which when bound by GAM31 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM31 target proteins.

[0257] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM31 gene, herein designated GAM GENE, on one or more GAM31 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0258] It is yet further appreciated that specific functions, and accordingly utilities, of GAM31 correlate with, and may be deduced from, the identity of the target genes which GAM31 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0259] Nucleotide sequences of the GAM31 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM31 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM31 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM31 are further described hereinbelow with reference to Table Table1.

[0260] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM31 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0261]

[0262] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 32 (GAM32) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0263] GAM32 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM32 was detected is described hereinabove with reference to Figs. 2-8.

[0264] GAM32 gene, herein designated GAM GENE, and GAM32

target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0265] GAM32 gene, herein designated GAM GENE, encodes a GAM32 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM32 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM32 precursor RNA is designated SEQ ID:134, and is provided hereinbelow with reference to the sequence listing part.

[0266] GAM32 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM32 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0267] An enzyme complex designated DICER COMPLEX, dices the GAM32 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, into GAM32 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM32 RNA is designated SEQ ID:135, and is provided hereinbelow with reference to the sequence listing part.

[0268] GAM32 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM32 target RNA, herein designated GAM TARGET RNA. GAM32 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0269] GAM32 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM32 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM32 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM32 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM32 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0270] The complementary binding of GAM32 RNA, herein designated GAM RNA, to target binding sites on GAM32 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM32 target RNA, herein designated GAM TARGET RNA, into GAM32 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0271] It is appreciated that GAM32 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM32 target genes. The mRNA of each one of this plurality of GAM32 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM32 RNA, herein designated GAM RNA, and which when bound by GAM32 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM32 target proteins.

[0272] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM32 gene, herein designated GAM GENE, on one or more GAM32 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not

yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0273] It is yet further appreciated that specific functions, and accordingly utilities, of GAM32 correlate with, and may be deduced from, the identity of the target genes which GAM32 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0274] Nucleotide sequences of the GAM32 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM32 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM32 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM32 are further described hereinbelow with reference to Table Table1.

[0275] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM32 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0276]

[0277] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 33 (GAM33) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0278] GAM33 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM33 was detected is described hereinabove with reference to Figs. 2-8.

[0279] GAM33 gene, herein designated GAM GENE, and GAM33 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0280] GAM33 gene, herein designated GAM GENE, encodes a GAM33 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM33 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM33 precursor RNA is designated SEQ ID:160, and is provided hereinbelow with reference to the sequence listing part.

[0281] GAM33 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM33 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0282] An enzyme complex designated DICER COMPLEX, dices the GAM33 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM33 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM33 RNA is designated SEQ ID:161, and is provided hereinbelow with reference to the sequence listing part.

[0283] GAM33 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM33 target RNA, herein designated GAM TARGET RNA. GAM33 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein

coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0284] GAM33 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM33 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM33 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM33 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM33 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0285] The complementary binding of GAM33 RNA, herein designated GAM RNA, to target binding sites on GAM33 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM33 target RNA, herein designated GAM TARGET RNA, into GAM33 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0286] It is appreciated that GAM33 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM33 target genes. The mRNA of each one of this plurality of GAM33 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM33 RNA, herein designated GAM RNA, and which when bound by GAM33 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM33 target proteins.

[0287] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM33 gene, herein designated GAM GENE, on one or more GAM33 target gene, herein designated GAM TARGET

GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0288] It is yet further appreciated that specific functions, and accordingly utilities, of GAM33 correlate with, and may be deduced from, the identity of the target genes which GAM33 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0289] Nucleotide sequences of the GAM33 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM33 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM33 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM33 are further described hereinbelow with reference to Table Table1.

[0290] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM33 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0291]

[0292] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 34 (GAM34) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0293] GAM34 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM34 was detected is described hereinabove with reference to Figs. 2-8.

[0294] GAM34 gene, herein designated GAM GENE, and GAM34 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0295] GAM34 gene, herein designated GAM GENE, encodes a GAM34 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM34 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM34 precursor RNA is designated SEQ ID:177, and is provided hereinbelow with reference to the sequence listing part.

[0296] GAM34 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM34 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0297] An enzyme complex designated DICER COMPLEX, dices the GAM34 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM34 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme

called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM34 RNA is designated SEQ ID:178, and is provided hereinbelow with reference to the sequence listing part.

[0298] GAM34 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM34 target RNA, herein designated GAM TARGET RNA. GAM34 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0299] GAM34 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM34 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM34 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM34 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM34 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0300] The complementary binding of GAM34 RNA, herein designated GAM RNA, to target binding sites on GAM34 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM34 target RNA, herein designated GAM TARGET RNA, into GAM34 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0301] It is appreciated that GAM34 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM34 target genes. The mRNA of each one of this plurality of GAM34 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM34 RNA,

herein designated GAM RNA, and which when bound by GAM34 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM34 target proteins.

[0302] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM34 gene, herein designated GAM GENE, on one or more GAM34 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0303] It is yet further appreciated that specific functions, and accordingly utilities, of GAM34 correlate with, and may be deduced from, the identity of the target genes which

GAM34 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0304] Nucleotide sequences of the GAM34 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM34 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM34 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM34 are further described hereinbelow with reference to Table Table1.

[0305] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM34 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0306]

[0307] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 35 (GAM35) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0308] GAM35 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM35 was detected is described hereinabove with reference to Figs. 2-8.

[0309] GAM35 gene, herein designated GAM GENE, and GAM35 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0310] GAM35 gene, herein designated GAM GENE, encodes a GAM35 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM35 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM35 precursor RNA is designated SEQ ID:188, and is provided hereinbelow with reference to the sequence listing part.

[0311] GAM35 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM35 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[0312] An enzyme complex designated DICER COMPLEX, dices the GAM35 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM35 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 27%) nucleotide sequence of GAM35 RNA is designated SEQ ID:189, and is provided hereinbelow with reference to the sequence listing part.

[0313] GAM35 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM35 target RNA, herein designated GAM TARGET RNA. GAM35 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0314] GAM35 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM35 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM35 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM35 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM35 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0315] The complementary binding of GAM35 RNA, herein designated GAM RNA, to target binding sites on GAM35 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM35 target RNA, herein designated GAM

TARGET RNA, into GAM35 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0316] It is appreciated that GAM35 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM35 target genes. The mRNA of each one of this plurality of GAM35 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM35 RNA, herein designated GAM RNA, and which when bound by GAM35 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM35 target proteins.

[0317] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM35 gene, herein designated GAM GENE, on one or more GAM35 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other re-

cently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0318] It is yet further appreciated that specific functions, and accordingly utilities, of GAM35 correlate with, and may be deduced from, the identity of the target genes which GAM35 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0319] Nucleotide sequences of the GAM35 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM35 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM35 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM35 are further described hereinbelow with reference to Table Table1.

[0320] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM35 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[0321]

[0322] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 36 (GAM36) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0323] GAM36 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM36 was detected is described hereinabove with reference to Figs. 2-8.

[0324] GAM36 gene, herein designated GAM GENE, and GAM36 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0325] GAM36 gene, herein designated GAM GENE, encodes a GAM36 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM36 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM36 precursor RNA is designated SEQ ID:208, and is provided hereinbelow with reference to

the sequence listing part.

[0326] GAM36 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM36 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0327] An enzyme complex designated DICER COMPLEX, dices the GAM36 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM36 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 39%) nucleotide sequence of GAM36 RNA is designated SEQ ID:209, and is provided hereinbelow with reference to the sequence listing part.

[0328] GAM36 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM36 target RNA, herein designated GAM TARGET RNA. GAM36 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0329] GAM36 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM36 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM36 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM36 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM36 target RNA, herein designated GAM TARGET RNA. It is further

appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0330] The complementary binding of GAM36 RNA, herein designated GAM RNA, to target binding sites on GAM36 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM36 target RNA, herein designated GAM TARGET RNA, into GAM36 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0331] It is appreciated that GAM36 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM36 target genes. The mRNA of each one of this plurality of GAM36 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM36 RNA, herein designated GAM RNA, and which when bound by GAM36 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM36 target proteins.

[0332] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM36 gene, herein designated GAM GENE, on one or more GAM36 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0333] It is yet further appreciated that specific functions, and accordingly utilities, of GAM36 correlate with, and may be deduced from, the identity of the target genes which GAM36 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0334] Nucleotide sequences of the GAM36 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM36 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM36 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM36 are further described hereinbelow with reference to Table Table1.

[0335] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM36 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0336]

[0337] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 37 (GAM37) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0338] GAM37 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM37 was detected is described hereinabove with reference to Figs. 2-8.

[0339] GAM37 gene, herein designated GAM GENE, and GAM37 target gene, herein designated GAM TARGET GENE, are

human genes contained in the human genome.

[0340] GAM37 gene, herein designated GAM GENE, encodes a GAM37 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM37 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM37 precursor RNA is designated SEQ ID:214, and is provided hereinbelow with reference to the sequence listing part.

[0341] GAM37 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM37 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0342] An enzyme complex designated DICER COMPLEX, dices the GAM37 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM37 RNA, herein desig-

nated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM37 RNA is designated SEQ ID:215, and is provided hereinbelow with reference to the sequence listing part.

[0343] GAM37 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM37 target RNA, herein designated GAM TARGET RNA. GAM37 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0344] GAM37 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM37 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM37 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM37 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM37 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0345] The complementary binding of GAM37 RNA, herein designated GAM RNA, to target binding sites on GAM37 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM37 target RNA, herein designated GAM TARGET RNA, into GAM37 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0346] It is appreciated that GAM37 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM37 target genes. The mRNA of each one of this plurality of GAM37 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM37 RNA, herein designated GAM RNA, and which when bound by GAM37 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM37 target proteins.

[0347] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM37 gene, herein designated GAM GENE, on one or more GAM37 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny

RNA world, Science 294,779 (2001)).

[0348] It is yet further appreciated that specific functions, and accordingly utilities, of GAM37 correlate with, and may be deduced from, the identity of the target genes which GAM37 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0349] Nucleotide sequences of the GAM37 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM37 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM37 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM37 are further described hereinbelow with reference to Table Table1.

[0350] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM37 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0351]

[0352] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

38 (GAM38) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0353] GAM38 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM38 was detected is described hereinabove with reference to Figs. 2-8.

[0354] GAM38 gene, herein designated GAM GENE, and GAM38 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0355] GAM38 gene, herein designated GAM GENE, encodes a GAM38 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM38 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM38 precursor RNA is designated SEQ ID:235, and is provided hereinbelow with reference to the sequence listing part.

[0356] GAM38 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM38 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0357] An enzyme complex designated DICER COMPLEX, dices the GAM38 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM38 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM38 RNA is designated SEQ ID:236, and is provided hereinbelow with reference to the sequence listing part.

[0358] GAM38 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM38 target RNA, herein designated GAM TARGET RNA. GAM38 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding re-

gion and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0359] GAM38 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM38 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM38 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM38 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM38 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0360] The complementary binding of GAM38 RNA, herein desig-

nated GAM RNA, to target binding sites on GAM38 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM38 target RNA, herein designated GAM TARGET RNA, into GAM38 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0361] It is appreciated that GAM38 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM38 target genes. The mRNA of each one of this plurality of GAM38 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM38 RNA, herein designated GAM RNA, and which when bound by GAM38 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM38 target proteins.

[0362] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM38 gene, herein designated GAM GENE, on one or more GAM38 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As

mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0363] It is yet further appreciated that specific functions, and accordingly utilities, of GAM38 correlate with, and may be deduced from, the identity of the target genes which GAM38 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0364] Nucleotide sequences of the GAM38 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM38 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM38 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM38 are further described hereinbelow with reference to Table Table1.

[0365] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM38 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0366]

[0367] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 39 (GAM39) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0368] GAM39 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM39 was detected is described hereinabove with reference to Figs. 2-8.

[0369] GAM39 gene, herein designated GAM GENE, and GAM39 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0370] GAM39 gene, herein designated GAM GENE, encodes a GAM39 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM39 precursor RNA, herein designated

GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM39 precursor RNA is designated SEQ ID:245, and is provided hereinbelow with reference to the sequence listing part.

[0371] GAM39 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM39 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0372] An enzyme complex designated DICER COMPLEX, dices the GAM39 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM39 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A

probable (over 64%) nucleotide sequence of GAM39 RNA is designated SEQ ID:246, and is provided hereinbelow with reference to the sequence listing part.

[0373] GAM39 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM39 target RNA, herein designated GAM TARGET RNA. GAM39 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0374] GAM39 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM39 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM39 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM39 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM39 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0375] The complementary binding of GAM39 RNA, herein designated GAM RNA, to target binding sites on GAM39 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM39 target RNA, herein designated GAM TARGET RNA, into GAM39 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0376] It is appreciated that GAM39 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM39 target genes. The mRNA of each one of this plurality of GAM39 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM39 RNA, herein designated GAM RNA, and which when bound by

GAM39 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM39 target proteins.

[0377] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM39 gene, herein designated GAM GENE, on one or more GAM39 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0378] It is yet further appreciated that specific functions, and accordingly utilities, of GAM39 correlate with, and may be deduced from, the identity of the target genes which GAM39 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[0379] Nucleotide sequences of the GAM39 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM39 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM39 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM39 are further described hereinbelow with reference to Table Table1.

[0380] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM39 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0381]

[0382] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 40 (GAM40) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0383] GAM40 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM40 was detected is described hereinabove with reference to Figs. 2-8.

[0384] GAM40 gene, herein designated GAM GENE, and GAM40 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0385] GAM40 gene, herein designated GAM GENE, encodes a GAM40 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM40 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM40 precursor RNA is designated SEQ ID:259, and is provided hereinbelow with reference to the sequence listing part.

[0386] GAM40 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM40 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[0387] An enzyme complex designated DICER COMPLEX, dices the GAM40 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM40 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM40 RNA is designated SEQ ID:260, and is provided hereinbelow with reference to the sequence listing part.

[0388] GAM40 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM40 target RNA, herein designated GAM TARGET RNA. GAM40 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0389] GAM40 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM40 target RNA, herein desig-

nated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM40 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM40 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM40 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0390] The complementary binding of GAM40 RNA, herein designated GAM RNA, to target binding sites on GAM40 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM40 target RNA, herein designated GAM TARGET RNA, into GAM40 target protein, herein desig-

nated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0391] It is appreciated that GAM40 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM40 target genes. The mRNA of each one of this plurality of GAM40 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM40 RNA, herein designated GAM RNA, and which when bound by GAM40 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM40 target proteins.

[0392] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM40 gene, herein designated GAM GENE, on one or more GAM40 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those

skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0393] It is yet further appreciated that specific functions, and accordingly utilities, of GAM40 correlate with, and may be deduced from, the identity of the target genes which GAM40 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0394] Nucleotide sequences of the GAM40 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM40 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM40 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM40 are further described hereinbelow with reference to Table Table1.

[0395] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM40 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0396]

[0397] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 41 (GAM41) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0398] GAM41 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM41 was detected is described hereinabove with reference to Figs. 2-8.

[0399] GAM41 gene, herein designated GAM GENE, and GAM41 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0400] GAM41 gene, herein designated GAM GENE, encodes a GAM41 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM41 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM41 precursor RNA is designated SEQ ID:263, and is provided hereinbelow with reference to the sequence listing part.

[0401] GAM41 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM41 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0402] An enzyme complex designated DICER COMPLEX, dices the GAM41 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM41 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM41 RNA is designated SEQ ID:264, and is provided hereinbelow with reference to the sequence listing part.

[0403] GAM41 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM41

target RNA, herein designated GAM TARGET RNA. GAM41 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0404] GAM41 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM41 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM41 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM41 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM41 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites

in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0405] The complementary binding of GAM41 RNA, herein designated GAM RNA, to target binding sites on GAM41 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM41 target RNA, herein designated GAM TARGET RNA, into GAM41 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0406] It is appreciated that GAM41 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM41 target genes. The mRNA of each one of this plurality of GAM41 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM41 RNA, herein designated GAM RNA, and which when bound by GAM41 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM41 target proteins.

[0407] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM41 gene, herein designated GAM GENE, on one or more GAM41 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0408] It is yet further appreciated that specific functions, and accordingly utilities, of GAM41 correlate with, and may be deduced from, the identity of the target genes which GAM41 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0409] Nucleotide sequences of the GAM41 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM41 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM41 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM41 are further described hereinbelow with reference to Table Table1.

[0410] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM41 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0411]

[0412] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 42 (GAM42) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0413] GAM42 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM42 was detected is described hereinabove with reference to Figs. 2-8.

[0414] GAM42 gene, herein designated GAM GENE, and GAM42 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0415] GAM42 gene, herein designated GAM GENE, encodes a GAM42 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM42 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM42 precursor RNA is designated SEQ ID:283, and is provided hereinbelow with reference to the sequence listing part.

[0416] GAM42 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM42 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0417] An enzyme complex designated DICER COMPLEX, dices the GAM42 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM42 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA seg-

ment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM42 RNA is designated SEQ ID:284, and is provided hereinbelow with reference to the sequence listing part.

[0418] GAM42 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM42 target RNA, herein designated GAM TARGET RNA. GAM42 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0419] GAM42 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM42 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM42 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM42 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM42 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0420] The complementary binding of GAM42 RNA, herein designated GAM RNA, to target binding sites on GAM42 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM42 target RNA, herein designated GAM TARGET RNA, into GAM42 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0421] It is appreciated that GAM42 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM42 target genes. The mRNA of each one of this plu-

rality of GAM42 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM42 RNA, herein designated GAM RNA, and which when bound by GAM42 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM42 target proteins.

[0422] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM42 gene, herein designated GAM GENE, on one or more GAM42 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0423] It is yet further appreciated that specific functions, and accordingly utilities, of GAM42 correlate with, and may be deduced from, the identity of the target genes which GAM42 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0424] Nucleotide sequences of the GAM42 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM42 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM42 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM42 are further described hereinbelow with reference to Table Table1.

[0425] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM42 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0426]

[0427] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 43 (GAM43) gene, which modulates expression of respec-

tive target genes thereof, the function and utility of which target genes is known in the art.

[0428] GAM43 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM43 was detected is described hereinabove with reference to Figs. 2-8.

[0429] GAM43 gene, herein designated GAM GENE, and GAM43 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0430] GAM43 gene, herein designated GAM GENE, encodes a GAM43 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM43 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM43 precursor RNA is designated SEQ ID:300, and is provided hereinbelow with reference to the sequence listing part.

[0431] GAM43 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM43 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0432] An enzyme complex designated DICER COMPLEX, dices the GAM43 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM43 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM43 RNA is designated SEQ ID:301, and is provided hereinbelow with reference to the sequence listing part.

[0433] GAM43 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM43 target RNA, herein designated GAM TARGET RNA. GAM43 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PRO-

TEIN CODING and 3UTR respectively.

[0434] GAM43 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM43 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM43 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM43 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM43 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0435] The complementary binding of GAM43 RNA, herein designated GAM RNA, to target binding sites on GAM43 target

RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM43 target RNA, herein designated GAM TARGET RNA, into GAM43 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0436] It is appreciated that GAM43 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM43 target genes. The mRNA of each one of this plurality of GAM43 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM43 RNA, herein designated GAM RNA, and which when bound by GAM43 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM43 target proteins.

[0437] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM43 gene, herein designated GAM GENE, on one or more GAM43 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background

section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0438] It is yet further appreciated that specific functions, and accordingly utilities, of GAM43 correlate with, and may be deduced from, the identity of the target genes which GAM43 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0439] Nucleotide sequences of the GAM43 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM43 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM43 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM43 are further described hereinbelow with reference to Table Table1.

[0440] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM43 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0441]

[0442] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 44 (GAM44) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0443] GAM44 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM44 was detected is described hereinabove with reference to Figs. 2-8.

[0444] GAM44 gene, herein designated GAM GENE, and GAM44 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0445] GAM44 gene, herein designated GAM GENE, encodes a GAM44 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM44 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nu-

cleotide sequence identical or highly similar to the nucleotide sequence of GAM44 precursor RNA is designated SEQ ID:309, and is provided hereinbelow with reference to the sequence listing part.

[0446] GAM44 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM44 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0447] An enzyme complex designated DICER COMPLEX, dices the GAM44 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM44 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM44 RNA is

designated SEQ ID:310, and is provided hereinbelow with reference to the sequence listing part.

[0448] GAM44 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM44 target RNA, herein designated GAM TARGET RNA. GAM44 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0449] GAM44 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM44 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM44 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM44 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM44 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0450] The complementary binding of GAM44 RNA, herein designated GAM RNA, to target binding sites on GAM44 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM44 target RNA, herein designated GAM TARGET RNA, into GAM44 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0451] It is appreciated that GAM44 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM44 target genes. The mRNA of each one of this plurality of GAM44 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM44 RNA, herein designated GAM RNA, and which when bound by GAM44 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM44 target proteins.

[0452] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM44 gene, herein designated GAM GENE, on one or more GAM44 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0453] It is yet further appreciated that specific functions, and accordingly utilities, of GAM44 correlate with, and may be deduced from, the identity of the target genes which GAM44 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0454] Nucleotide sequences of the GAM44 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM44 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM44 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM44 are further described hereinbelow with reference to Table Table1.

[0455] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM44 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0456]

[0457] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 45 (GAM45) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0458] GAM45 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM45 was detected is described hereinabove

with reference to Figs. 2-8.

[0459] GAM45 gene, herein designated GAM GENE, and GAM45 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0460] GAM45 gene, herein designated GAM GENE, encodes a GAM45 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM45 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM45 precursor RNA is designated SEQ ID:337, and is provided hereinbelow with reference to the sequence listing part.

[0461] GAM45 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM45 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0462] An enzyme complex designated DICER COMPLEX, dices the GAM45 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM45 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 17%) nucleotide sequence of GAM45 RNA is designated SEQ ID:338, and is provided hereinbelow with reference to the sequence listing part.

[0463] GAM45 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM45 target RNA, herein designated GAM TARGET RNA. GAM45 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0464] GAM45 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM45 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM45 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM45 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM45 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0465] The complementary binding of GAM45 RNA, herein designated GAM RNA, to target binding sites on GAM45 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM45 target RNA, herein designated GAM TARGET RNA, into GAM45 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is there-

fore outlined by a broken line.

[0466] It is appreciated that GAM45 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM45 target genes. The mRNA of each one of this plurality of GAM45 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM45 RNA, herein designated GAM RNA, and which when bound by GAM45 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM45 target proteins.

[0467] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM45 gene, herein designated GAM GENE, on one or more GAM45 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes

by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0468] It is yet further appreciated that specific functions, and accordingly utilities, of GAM45 correlate with, and may be deduced from, the identity of the target genes which GAM45 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0469] Nucleotide sequences of the GAM45 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM45 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM45 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM45 are further described hereinbelow with reference to Table Table1.

[0470] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM45 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0471]

[0472] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 46 (GAM46) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0473] GAM46 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM46 was detected is described hereinabove with reference to Figs. 2-8.

[0474] GAM46 gene, herein designated GAM GENE, and GAM46 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0475] GAM46 gene, herein designated GAM GENE, encodes a GAM46 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM46 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM46 precursor RNA is designated SEQ ID:365, and is provided hereinbelow with reference to the sequence listing part.

[0476] GAM46 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM46 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0477] An enzyme complex designated DICER COMPLEX, dices the GAM46 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM46 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 80%) nucleotide sequence of GAM46 RNA is designated SEQ ID:366, and is provided hereinbelow with reference to the sequence listing part.

[0478] GAM46 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM46 target RNA, herein designated GAM TARGET RNA. GAM46

target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0479] GAM46 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM46 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM46 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM46 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM46 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0480] The complementary binding of GAM46 RNA, herein designated GAM RNA, to target binding sites on GAM46 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM46 target RNA, herein designated GAM TARGET RNA, into GAM46 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0481] It is appreciated that GAM46 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM46 target genes. The mRNA of each one of this plurality of GAM46 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM46 RNA, herein designated GAM RNA, and which when bound by GAM46 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM46 target proteins.

[0482] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM46 gene, herein designated GAM GENE, on one or more GAM46 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0483] It is yet further appreciated that specific functions, and accordingly utilities, of GAM46 correlate with, and may be deduced from, the identity of the target genes which GAM46 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0484] Nucleotide sequences of the GAM46 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM46 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM46 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM46 are further described hereinbelow with reference to Table Table1.

[0485] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM46 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0486]

[0487] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 47 (GAM47) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0488] GAM47 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM47 was detected is described hereinabove with reference to Figs. 2-8.

[0489] GAM47 gene, herein designated GAM GENE, and GAM47 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0490] GAM47 gene, herein designated GAM GENE, encodes a

GAM47 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM47 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM47 precursor RNA is designated SEQ ID:387, and is provided hereinbelow with reference to the sequence listing part.

[0491] GAM47 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM47 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0492] An enzyme complex designated DICER COMPLEX, dices the GAM47 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM47 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured

RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 42%) nucleotide sequence of GAM47 RNA is designated SEQ ID:388, and is provided hereinbelow with reference to the sequence listing part.

[0493] GAM47 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM47 target RNA, herein designated GAM TARGET RNA. GAM47 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0494] GAM47 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM47 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM47 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM47 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM47 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0495] The complementary binding of GAM47 RNA, herein designated GAM RNA, to target binding sites on GAM47 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM47 target RNA, herein designated GAM TARGET RNA, into GAM47 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0496] It is appreciated that GAM47 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM47 target genes. The mRNA of each one of this plurality of GAM47 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM47 RNA, herein designated GAM RNA, and which when bound by GAM47 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM47 target proteins.

[0497] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM47 gene, herein designated GAM GENE, on one or more GAM47 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0498] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM47 correlate with, and may be deduced from, the identity of the target genes which GAM47 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0499] Nucleotide sequences of the GAM47 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM47 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM47 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM47 are further described hereinbelow with reference to Table Table1.

[0500] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM47 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0501]

[0502] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 48 (GAM48) gene, which modulates expression of respective target genes thereof, the function and utility of which

target genes is known in the art.

[0503] GAM48 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM48 was detected is described hereinabove with reference to Figs. 2-8.

[0504] GAM48 gene, herein designated GAM GENE, and GAM48 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0505] GAM48 gene, herein designated GAM GENE, encodes a GAM48 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM48 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM48 precursor RNA is designated SEQ ID:422, and is provided hereinbelow with reference to the sequence listing part.

[0506] GAM48 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM48 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0507] An enzyme complex designated DICER COMPLEX, dices the GAM48 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM48 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM48 RNA is designated SEQ ID:423, and is provided hereinbelow with reference to the sequence listing part.

[0508] GAM48 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM48 target RNA, herein designated GAM TARGET RNA. GAM48 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0509] GAM48 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM48 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM48 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM48 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM48 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0510] The complementary binding of GAM48 RNA, herein designated GAM RNA, to target binding sites on GAM48 target RNA, herein designated GAM TARGET RNA, such as BIND-

ING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM48 target RNA, herein designated GAM TARGET RNA, into GAM48 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0511] It is appreciated that GAM48 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM48 target genes. The mRNA of each one of this plurality of GAM48 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM48 RNA, herein designated GAM RNA, and which when bound by GAM48 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM48 target proteins.

[0512] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM48 gene, herein designated GAM GENE, on one or more GAM48 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site

has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0513] It is yet further appreciated that specific functions, and accordingly utilities, of GAM48 correlate with, and may be deduced from, the identity of the target genes which GAM48 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0514] Nucleotide sequences of the GAM48 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM48 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM48 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM48 are further described hereinbelow with reference to Table Table1.

[0515] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM48 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0516]

[0517] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 49 (GAM49) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0518] GAM49 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM49 was detected is described hereinabove with reference to Figs. 2-8.

[0519] GAM49 gene, herein designated GAM GENE, and GAM49 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0520] GAM49 gene, herein designated GAM GENE, encodes a GAM49 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM49 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM49 precursor RNA is designated SEQ ID:429, and is provided hereinbelow with reference to the sequence listing part.

[0521] GAM49 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM49 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0522] An enzyme complex designated DICER COMPLEX, dices the GAM49 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM49 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM49 RNA is designated SEQ ID:430, and is provided hereinbelow with

reference to the sequence listing part.

[0523] GAM49 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM49 target RNA, herein designated GAM TARGET RNA. GAM49 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0524] GAM49 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM49 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM49 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM49 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM49 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0525] The complementary binding of GAM49 RNA, herein designated GAM RNA, to target binding sites on GAM49 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM49 target RNA, herein designated GAM TARGET RNA, into GAM49 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0526] It is appreciated that GAM49 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM49 target genes. The mRNA of each one of this plurality of GAM49 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM49 RNA, herein designated GAM RNA, and which when bound by GAM49 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM49 tar-

get proteins.

[0527] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM49 gene, herein designated GAM GENE, on one or more GAM49 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0528] It is yet further appreciated that specific functions, and accordingly utilities, of GAM49 correlate with, and may be deduced from, the identity of the target genes which GAM49 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0529] Nucleotide sequences of the GAM49 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM49 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM49 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM49 are further described hereinbelow with reference to Table Table1.

[0530] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM49 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0531]

[0532] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 50 (GAM50) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0533] GAM50 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM50 was detected is described hereinabove with reference to Figs. 2-8.

[0534] GAM50 gene, herein designated GAM GENE, and GAM50 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0535] GAM50 gene, herein designated GAM GENE, encodes a GAM50 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM50 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM50 precursor RNA is designated SEQ ID:453, and is provided hereinbelow with reference to the sequence listing part.

[0536] GAM50 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM50 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0537] An enzyme complex designated DICER COMPLEX, dices

the GAM50 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM50 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM50 RNA is designated SEQ ID:454, and is provided hereinbelow with reference to the sequence listing part.

[0538] GAM50 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM50 target RNA, herein designated GAM TARGET RNA. GAM50 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0539] GAM50 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM50 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM50

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM50 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM50 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0540] The complementary binding of GAM50 RNA, herein designated GAM RNA, to target binding sites on GAM50 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM50 target RNA, herein designated GAM TARGET RNA, into GAM50 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0541] It is appreciated that GAM50 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM50 target genes. The mRNA of each one of this plurality of GAM50 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM50 RNA, herein designated GAM RNA, and which when bound by GAM50 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM50 target proteins.

[0542] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM50 gene, herein designated GAM GENE, on one or more GAM50 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complemen-

tary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0543] It is yet further appreciated that specific functions, and accordingly utilities, of GAM50 correlate with, and may be deduced from, the identity of the target genes which GAM50 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0544] Nucleotide sequences of the GAM50 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM50 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM50 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM50 are further described hereinbelow with reference to Table Table1.

[0545] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM50 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0546]

[0547] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 51 (GAM51) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0548] GAM51 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM51 was detected is described hereinabove with reference to Figs. 2-8.

[0549] GAM51 gene, herein designated GAM GENE, and GAM51 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0550] GAM51 gene, herein designated GAM GENE, encodes a GAM51 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM51 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM51 precursor RNA is designated SEQ ID:459, and is provided hereinbelow with reference to the sequence listing part.

[0551] GAM51 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM51 folded precur-

sor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0552] An enzyme complex designated DICER COMPLEX, dices the GAM51 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM51 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 26%) nucleotide sequence of GAM51 RNA is designated SEQ ID:460, and is provided hereinbelow with reference to the sequence listing part.

[0553] GAM51 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM51 target RNA, herein designated GAM TARGET RNA. GAM51 target RNA, herein designated GAM TARGET RNA, com-

prises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0554] GAM51 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM51 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM51 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM51 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM51 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[0555] The complementary binding of GAM51 RNA, herein designated GAM RNA, to target binding sites on GAM51 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM51 target RNA, herein designated GAM TARGET RNA, into GAM51 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0556] It is appreciated that GAM51 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM51 target genes. The mRNA of each one of this plurality of GAM51 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM51 RNA, herein designated GAM RNA, and which when bound by GAM51 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM51 target proteins.

[0557] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM51 gene, herein designated GAM GENE, on one or

more GAM51 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0558] It is yet further appreciated that specific functions, and accordingly utilities, of GAM51 correlate with, and may be deduced from, the identity of the target genes which GAM51 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0559] Nucleotide sequences of the GAM51 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM51 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM51 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM51 are further described

hereinbelow with reference to Table Table1.

[0560] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM51 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0561]

[0562] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 52 (GAM52) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0563] GAM52 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM52 was detected is described hereinabove with reference to Figs. 2-8.

[0564] GAM52 gene, herein designated GAM GENE, and GAM52 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0565] GAM52 gene, herein designated GAM GENE, encodes a GAM52 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM52 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM52 precursor RNA is designated SEQ ID:480, and is provided hereinbelow with reference to the sequence listing part.

[0566] GAM52 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM52 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0567] An enzyme complex designated DICER COMPLEX, dices the GAM52 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM52 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is

catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 41%) nucleotide sequence of GAM52 RNA is designated SEQ ID:481, and is provided hereinbelow with reference to the sequence listing part.

[0568] GAM52 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM52 target RNA, herein designated GAM TARGET RNA. GAM52 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0569] GAM52 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM52 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM52 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM52 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM52 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0570] The complementary binding of GAM52 RNA, herein designated GAM RNA, to target binding sites on GAM52 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM52 target RNA, herein designated GAM TARGET RNA, into GAM52 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0571] It is appreciated that GAM52 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM52 target genes. The mRNA of each one of this plurality of GAM52 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM52 RNA, herein designated GAM RNA, and which when bound by GAM52 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM52 target proteins.

[0572] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM52 gene, herein designated GAM GENE, on one or more GAM52 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0573] It is yet further appreciated that specific functions, and accordingly utilities, of GAM52 correlate with, and may be

deduced from, the identity of the target genes which GAM52 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0574] Nucleotide sequences of the GAM52 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM52 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM52 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM52 are further described hereinbelow with reference to Table Table1.

[0575] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM52 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0576]

[0577] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 53 (GAM53) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0578] GAM53 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM53 was detected is described hereinabove with reference to Figs. 2-8.

[0579] GAM53 gene, herein designated GAM GENE, and GAM53 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0580] GAM53 gene, herein designated GAM GENE, encodes a GAM53 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM53 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM53 precursor RNA is designated SEQ ID:522, and is provided hereinbelow with reference to the sequence listing part.

[0581] GAM53 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM53 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0582] An enzyme complex designated DICER COMPLEX, dices the GAM53 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM53 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM53 RNA is designated SEQ ID:523, and is provided hereinbelow with reference to the sequence listing part.

[0583] GAM53 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM53 target RNA, herein designated GAM TARGET RNA. GAM53 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0584] GAM53 RNA, herein designated GAM RNA, binds comple-

mentarily to one or more target binding sites located in untranslated regions of GAM53 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM53 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM53 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM53 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0585] The complementary binding of GAM53 RNA, herein designated GAM RNA, to target binding sites on GAM53 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits

translation of GAM53 target RNA, herein designated GAM TARGET RNA, into GAM53 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0586] It is appreciated that GAM53 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM53 target genes. The mRNA of each one of this plurality of GAM53 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM53 RNA, herein designated GAM RNA, and which when bound by GAM53 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM53 target proteins.

[0587] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM53 gene, herein designated GAM GENE, on one or more GAM53 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known

miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0588] It is yet further appreciated that specific functions, and accordingly utilities, of GAM53 correlate with, and may be deduced from, the identity of the target genes which GAM53 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0589] Nucleotide sequences of the GAM53 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM53 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM53 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM53 are further described hereinbelow with reference to Table Table1.

[0590] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM53 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0591]

[0592] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 54 (GAM54) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0593] GAM54 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM54 was detected is described hereinabove with reference to Figs. 2-8.

[0594] GAM54 gene, herein designated GAM GENE, and GAM54 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0595] GAM54 gene, herein designated GAM GENE, encodes a GAM54 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM54 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM54 precursor RNA is designated

SEQ ID:531, and is provided hereinbelow with reference to the sequence listing part.

[0596] GAM54 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM54 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0597] An enzyme complex designated DICER COMPLEX, dices the GAM54 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM54 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM54 RNA is designated SEQ ID:532, and is provided hereinbelow with reference to the sequence listing part.

[0598] GAM54 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM54 target RNA, herein designated GAM TARGET RNA. GAM54 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0599] GAM54 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM54 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM54 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM54 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM54 tar-

get RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0600] The complementary binding of GAM54 RNA, herein designated GAM RNA, to target binding sites on GAM54 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM54 target RNA, herein designated GAM TARGET RNA, into GAM54 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0601] It is appreciated that GAM54 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM54 target genes. The mRNA of each one of this plurality of GAM54 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM54 RNA, herein designated GAM RNA, and which when bound by GAM54 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM54 target proteins.

[0602] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM54 gene, herein designated GAM GENE, on one or more GAM54 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0603] It is yet further appreciated that specific functions, and accordingly utilities, of GAM54 correlate with, and may be deduced from, the identity of the target genes which GAM54 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0604] Nucleotide sequences of the GAM54 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM54 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM54 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM54 are further described hereinbelow with reference to Table Table1.

[0605] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM54 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0606]

[0607] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 55 (GAM55) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0608] GAM55 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM55 was detected is described hereinabove with reference to Figs. 2-8.

[0609] GAM55 gene, herein designated GAM GENE, and GAM55

target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0610] GAM55 gene, herein designated GAM GENE, encodes a GAM55 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM55 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM55 precursor RNA is designated SEQ ID:545, and is provided hereinbelow with reference to the sequence listing part.

[0611] GAM55 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM55 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0612] An enzyme complex designated DICER COMPLEX, dices the GAM55 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, into GAM55 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM55 RNA is designated SEQ ID:546, and is provided hereinbelow with reference to the sequence listing part.

[0613] GAM55 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM55 target RNA, herein designated GAM TARGET RNA. GAM55 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0614] GAM55 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM55 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM55 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM55 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM55 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0615] The complementary binding of GAM55 RNA, herein designated GAM RNA, to target binding sites on GAM55 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM55 target RNA, herein designated GAM TARGET RNA, into GAM55 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0616] It is appreciated that GAM55 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM55 target genes. The mRNA of each one of this plurality of GAM55 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM55 RNA, herein designated GAM RNA, and which when bound by GAM55 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM55 target proteins.

[0617] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM55 gene, herein designated GAM GENE, on one or more GAM55 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not

yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0618] It is yet further appreciated that specific functions, and accordingly utilities, of GAM55 correlate with, and may be deduced from, the identity of the target genes which GAM55 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0619] Nucleotide sequences of the GAM55 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM55 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM55 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM55 are further described hereinbelow with reference to Table Table1.

[0620] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM55 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0621]

[0622] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 56 (GAM56) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0623] GAM56 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM56 was detected is described hereinabove with reference to Figs. 2-8.

[0624] GAM56 gene, herein designated GAM GENE, and GAM56 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0625] GAM56 gene, herein designated GAM GENE, encodes a GAM56 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM56 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM56 precursor RNA is designated SEQ ID:556, and is provided hereinbelow with reference to the sequence listing part.

[0626] GAM56 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM56 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0627] An enzyme complex designated DICER COMPLEX, dices the GAM56 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM56 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM56 RNA is designated SEQ ID:557, and is provided hereinbelow with reference to the sequence listing part.

[0628] GAM56 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM56 target RNA, herein designated GAM TARGET RNA. GAM56 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein

coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0629] GAM56 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM56 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM56 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM56 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM56 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0630] The complementary binding of GAM56 RNA, herein designated GAM RNA, to target binding sites on GAM56 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM56 target RNA, herein designated GAM TARGET RNA, into GAM56 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0631] It is appreciated that GAM56 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM56 target genes. The mRNA of each one of this plurality of GAM56 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM56 RNA, herein designated GAM RNA, and which when bound by GAM56 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM56 target proteins.

[0632] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM56 gene, herein designated GAM GENE, on one or more GAM56 target gene, herein designated GAM TARGET

GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0633] It is yet further appreciated that specific functions, and accordingly utilities, of GAM56 correlate with, and may be deduced from, the identity of the target genes which GAM56 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0634] Nucleotide sequences of the GAM56 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM56 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM56 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM56 are further described hereinbelow with reference to Table Table1.

[0635] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM56 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0636]

[0637] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 57 (GAM57) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0638] GAM57 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM57 was detected is described hereinabove with reference to Figs. 2-8.

[0639] GAM57 gene, herein designated GAM GENE, and GAM57 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0640] GAM57 gene, herein designated GAM GENE, encodes a GAM57 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM57 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM57 precursor RNA is designated SEQ ID:561, and is provided hereinbelow with reference to the sequence listing part.

[0641] GAM57 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM57 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0642] An enzyme complex designated DICER COMPLEX, dices the GAM57 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM57 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme

called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM57 RNA is designated SEQ ID:562, and is provided hereinbelow with reference to the sequence listing part.

[0643] GAM57 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM57 target RNA, herein designated GAM TARGET RNA. GAM57 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0644] GAM57 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM57 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM57 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM57 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM57 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0645] The complementary binding of GAM57 RNA, herein designated GAM RNA, to target binding sites on GAM57 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM57 target RNA, herein designated GAM TARGET RNA, into GAM57 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0646] It is appreciated that GAM57 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM57 target genes. The mRNA of each one of this plurality of GAM57 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM57 RNA,

herein designated GAM RNA, and which when bound by GAM57 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM57 target proteins.

[0647] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM57 gene, herein designated GAM GENE, on one or more GAM57 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0648] It is yet further appreciated that specific functions, and accordingly utilities, of GAM57 correlate with, and may be deduced from, the identity of the target genes which

GAM57 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0649] Nucleotide sequences of the GAM57 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM57 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM57 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM57 are further described hereinbelow with reference to Table Table1.

[0650] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM57 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0651]

[0652] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 58 (GAM58) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0653] GAM58 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM58 was detected is described hereinabove with reference to Figs. 2-8.

[0654] GAM58 gene, herein designated GAM GENE, and GAM58 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0655] GAM58 gene, herein designated GAM GENE, encodes a GAM58 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM58 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM58 precursor RNA is designated SEQ ID:573, and is provided hereinbelow with reference to the sequence listing part.

[0656] GAM58 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM58 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[0657] An enzyme complex designated DICER COMPLEX, dices the GAM58 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM58 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM58 RNA is designated SEQ ID:574, and is provided hereinbelow with reference to the sequence listing part.

[0658] GAM58 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM58 target RNA, herein designated GAM TARGET RNA. GAM58 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0659] GAM58 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM58 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM58 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM58 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM58 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0660] The complementary binding of GAM58 RNA, herein designated GAM RNA, to target binding sites on GAM58 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM58 target RNA, herein designated GAM

TARGET RNA, into GAM58 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0661] It is appreciated that GAM58 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM58 target genes. The mRNA of each one of this plurality of GAM58 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM58 RNA, herein designated GAM RNA, and which when bound by GAM58 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM58 target proteins.

[0662] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM58 gene, herein designated GAM GENE, on one or more GAM58 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other re-

cently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0663] It is yet further appreciated that specific functions, and accordingly utilities, of GAM58 correlate with, and may be deduced from, the identity of the target genes which GAM58 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0664] Nucleotide sequences of the GAM58 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM58 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM58 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM58 are further described hereinbelow with reference to Table Table1.

[0665] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM58 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[0666]

[0667] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 59 (GAM59) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0668] GAM59 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM59 was detected is described hereinabove with reference to Figs. 2-8.

[0669] GAM59 gene, herein designated GAM GENE, and GAM59 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0670] GAM59 gene, herein designated GAM GENE, encodes a GAM59 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM59 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM59 precursor RNA is designated SEQ ID:588, and is provided hereinbelow with reference to

the sequence listing part.

[0671] GAM59 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM59 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0672] An enzyme complex designated DICER COMPLEX, dices the GAM59 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM59 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM59 RNA is designated SEQ ID:589, and is provided hereinbelow with reference to the sequence listing part.

[0673] GAM59 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM59 target RNA, herein designated GAM TARGET RNA. GAM59 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0674] GAM59 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM59 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM59 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM59 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM59 target RNA, herein designated GAM TARGET RNA. It is further

appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0675] The complementary binding of GAM59 RNA, herein designated GAM RNA, to target binding sites on GAM59 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM59 target RNA, herein designated GAM TARGET RNA, into GAM59 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0676] It is appreciated that GAM59 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM59 target genes. The mRNA of each one of this plurality of GAM59 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM59 RNA, herein designated GAM RNA, and which when bound by GAM59 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM59 target proteins.

[0677] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM59 gene, herein designated GAM GENE, on one or more GAM59 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0678] It is yet further appreciated that specific functions, and accordingly utilities, of GAM59 correlate with, and may be deduced from, the identity of the target genes which GAM59 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0679] Nucleotide sequences of the GAM59 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM59 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM59 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM59 are further described hereinbelow with reference to Table Table1.

[0680] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM59 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0681]

[0682] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 60 (GAM60) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0683] GAM60 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM60 was detected is described hereinabove with reference to Figs. 2-8.

[0684] GAM60 gene, herein designated GAM GENE, and GAM60 target gene, herein designated GAM TARGET GENE, are

human genes contained in the human genome.

[0685] GAM60 gene, herein designated GAM GENE, encodes a GAM60 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM60 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM60 precursor RNA is designated SEQ ID:604, and is provided hereinbelow with reference to the sequence listing part.

[0686] GAM60 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM60 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0687] An enzyme complex designated DICER COMPLEX, dices the GAM60 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM60 RNA, herein desig-

nated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM60 RNA is designated SEQ ID:605, and is provided hereinbelow with reference to the sequence listing part.

[0688] GAM60 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM60 target RNA, herein designated GAM TARGET RNA. GAM60 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0689] GAM60 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM60 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM60 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM60 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM60 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0690] The complementary binding of GAM60 RNA, herein designated GAM RNA, to target binding sites on GAM60 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM60 target RNA, herein designated GAM TARGET RNA, into GAM60 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0691] It is appreciated that GAM60 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM60 target genes. The mRNA of each one of this plurality of GAM60 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM60 RNA, herein designated GAM RNA, and which when bound by GAM60 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM60 target proteins.

[0692] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM60 gene, herein designated GAM GENE, on one or more GAM60 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny

RNA world, Science 294,779 (2001)).

[0693] It is yet further appreciated that specific functions, and accordingly utilities, of GAM60 correlate with, and may be deduced from, the identity of the target genes which GAM60 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0694] Nucleotide sequences of the GAM60 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM60 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM60 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM60 are further described hereinbelow with reference to Table Table1.

[0695] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM60 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0696]

[0697] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

61 (GAM61) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0698] GAM61 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM61 was detected is described hereinabove with reference to Figs. 2-8.

[0699] GAM61 gene, herein designated GAM GENE, and GAM61 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0700] GAM61 gene, herein designated GAM GENE, encodes a GAM61 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM61 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM61 precursor RNA is designated SEQ ID:617, and is provided hereinbelow with reference to the sequence listing part.

[0701] GAM61 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM61 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0702] An enzyme complex designated DICER COMPLEX, dices the GAM61 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM61 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM61 RNA is designated SEQ ID:618, and is provided hereinbelow with reference to the sequence listing part.

[0703] GAM61 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM61 target RNA, herein designated GAM TARGET RNA. GAM61 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding re-

gion and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0704] GAM61 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM61 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM61 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM61 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM61 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0705] The complementary binding of GAM61 RNA, herein desig-

nated GAM RNA, to target binding sites on GAM61 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM61 target RNA, herein designated GAM TARGET RNA, into GAM61 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0706] It is appreciated that GAM61 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM61 target genes. The mRNA of each one of this plurality of GAM61 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM61 RNA, herein designated GAM RNA, and which when bound by GAM61 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM61 target proteins.

[0707] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM61 gene, herein designated GAM GENE, on one or more GAM61 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As

mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0708] It is yet further appreciated that specific functions, and accordingly utilities, of GAM61 correlate with, and may be deduced from, the identity of the target genes which GAM61 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0709] Nucleotide sequences of the GAM61 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM61 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM61 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM61 are further described hereinbelow with reference to Table Table1.

[0710] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM61 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0711]

[0712] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 62 (GAM62) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0713] GAM62 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM62 was detected is described hereinabove with reference to Figs. 2-8.

[0714] GAM62 gene, herein designated GAM GENE, and GAM62 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0715] GAM62 gene, herein designated GAM GENE, encodes a GAM62 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM62 precursor RNA, herein designated

GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM62 precursor RNA is designated SEQ ID:630, and is provided hereinbelow with reference to the sequence listing part.

[0716] GAM62 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM62 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0717] An enzyme complex designated DICER COMPLEX, dices the GAM62 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM62 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A

probable (over 53%) nucleotide sequence of GAM62 RNA is designated SEQ ID:631, and is provided hereinbelow with reference to the sequence listing part.

[0718] GAM62 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM62 target RNA, herein designated GAM TARGET RNA. GAM62 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0719] GAM62 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM62 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM62 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM62 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM62 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0720] The complementary binding of GAM62 RNA, herein designated GAM RNA, to target binding sites on GAM62 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM62 target RNA, herein designated GAM TARGET RNA, into GAM62 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0721] It is appreciated that GAM62 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM62 target genes. The mRNA of each one of this plurality of GAM62 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM62 RNA, herein designated GAM RNA, and which when bound by

GAM62 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM62 target proteins.

[0722] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM62 gene, herein designated GAM GENE, on one or more GAM62 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0723] It is yet further appreciated that specific functions, and accordingly utilities, of GAM62 correlate with, and may be deduced from, the identity of the target genes which GAM62 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[0724] Nucleotide sequences of the GAM62 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM62 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM62 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM62 are further described hereinbelow with reference to Table Table1.

[0725] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM62 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0726]

[0727] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 63 (GAM63) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0728] GAM63 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM63 was detected is described hereinabove with reference to Figs. 2-8.

[0729] GAM63 gene, herein designated GAM GENE, and GAM63 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0730] GAM63 gene, herein designated GAM GENE, encodes a GAM63 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM63 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM63 precursor RNA is designated SEQ ID:652, and is provided hereinbelow with reference to the sequence listing part.

[0731] GAM63 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM63 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[0732] An enzyme complex designated DICER COMPLEX, dices the GAM63 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM63 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM63 RNA is designated SEQ ID:653, and is provided hereinbelow with reference to the sequence listing part.

[0733] GAM63 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM63 target RNA, herein designated GAM TARGET RNA. GAM63 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0734] GAM63 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM63 target RNA, herein desig-

nated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM63 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM63 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM63 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0735] The complementary binding of GAM63 RNA, herein designated GAM RNA, to target binding sites on GAM63 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM63 target RNA, herein designated GAM TARGET RNA, into GAM63 target protein, herein desig-

nated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0736] It is appreciated that GAM63 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM63 target genes. The mRNA of each one of this plurality of GAM63 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM63 RNA, herein designated GAM RNA, and which when bound by GAM63 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM63 target proteins.

[0737] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM63 gene, herein designated GAM GENE, on one or more GAM63 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those

skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0738] It is yet further appreciated that specific functions, and accordingly utilities, of GAM63 correlate with, and may be deduced from, the identity of the target genes which GAM63 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0739] Nucleotide sequences of the GAM63 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM63 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM63 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM63 are further described hereinbelow with reference to Table Table1.

[0740] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM63 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0741]

[0742] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 64 (GAM64) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0743] GAM64 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM64 was detected is described hereinabove with reference to Figs. 2-8.

[0744] GAM64 gene, herein designated GAM GENE, and GAM64 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0745] GAM64 gene, herein designated GAM GENE, encodes a GAM64 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM64 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM64 precursor RNA is designated SEQ ID:673, and is provided hereinbelow with reference to the sequence listing part.

[0746] GAM64 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM64 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0747] An enzyme complex designated DICER COMPLEX, dices the GAM64 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM64 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 38%) nucleotide sequence of GAM64 RNA is designated SEQ ID:674, and is provided hereinbelow with reference to the sequence listing part.

[0748] GAM64 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM64

target RNA, herein designated GAM TARGET RNA. GAM64 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0749] GAM64 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM64 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM64 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM64 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM64 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites

in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0750] The complementary binding of GAM64 RNA, herein designated GAM RNA, to target binding sites on GAM64 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM64 target RNA, herein designated GAM TARGET RNA, into GAM64 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0751] It is appreciated that GAM64 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM64 target genes. The mRNA of each one of this plurality of GAM64 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM64 RNA, herein designated GAM RNA, and which when bound by GAM64 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM64 target proteins.

[0752] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM64 gene, herein designated GAM GENE, on one or more GAM64 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0753] It is yet further appreciated that specific functions, and accordingly utilities, of GAM64 correlate with, and may be deduced from, the identity of the target genes which GAM64 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0754] Nucleotide sequences of the GAM64 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM64 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM64 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM64 are further described hereinbelow with reference to Table Table1.

[0755] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM64 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0756]

[0757] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 65 (GAM65) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0758] GAM65 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM65 was detected is described hereinabove with reference to Figs. 2-8.

[0759] GAM65 gene, herein designated GAM GENE, and GAM65 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0760] GAM65 gene, herein designated GAM GENE, encodes a GAM65 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM65 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM65 precursor RNA is designated SEQ ID:697, and is provided hereinbelow with reference to the sequence listing part.

[0761] GAM65 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM65 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0762] An enzyme complex designated DICER COMPLEX, dices the GAM65 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM65 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA seg-

ment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM65 RNA is designated SEQ ID:698, and is provided hereinbelow with reference to the sequence listing part.

[0763] GAM65 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM65 target RNA, herein designated GAM TARGET RNA. GAM65 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0764] GAM65 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM65 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM65 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM65 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM65 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0765] The complementary binding of GAM65 RNA, herein designated GAM RNA, to target binding sites on GAM65 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM65 target RNA, herein designated GAM TARGET RNA, into GAM65 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0766] It is appreciated that GAM65 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM65 target genes. The mRNA of each one of this plu-

rality of GAM65 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM65 RNA, herein designated GAM RNA, and which when bound by GAM65 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM65 target proteins.

[0767] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM65 gene, herein designated GAM GENE, on one or more GAM65 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0768] It is yet further appreciated that specific functions, and accordingly utilities, of GAM65 correlate with, and may be deduced from, the identity of the target genes which GAM65 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0769] Nucleotide sequences of the GAM65 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM65 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM65 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM65 are further described hereinbelow with reference to Table Table1.

[0770] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM65 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0771]

[0772] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 66 (GAM66) gene, which modulates expression of respec-

tive target genes thereof, the function and utility of which target genes is known in the art.

[0773] GAM66 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM66 was detected is described hereinabove with reference to Figs. 2-8.

[0774] GAM66 gene, herein designated GAM GENE, and GAM66 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0775] GAM66 gene, herein designated GAM GENE, encodes a GAM66 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM66 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM66 precursor RNA is designated SEQ ID:721, and is provided hereinbelow with reference to the sequence listing part.

[0776] GAM66 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM66 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0777] An enzyme complex designated DICER COMPLEX, dices the GAM66 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM66 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM66 RNA is designated SEQ ID:722, and is provided hereinbelow with reference to the sequence listing part.

[0778] GAM66 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM66 target RNA, herein designated GAM TARGET RNA. GAM66 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PRO-

TEIN CODING and 3UTR respectively.

[0779] GAM66 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM66 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM66 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM66 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM66 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0780] The complementary binding of GAM66 RNA, herein designated GAM RNA, to target binding sites on GAM66 target

RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM66 target RNA, herein designated GAM TARGET RNA, into GAM66 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0781] It is appreciated that GAM66 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM66 target genes. The mRNA of each one of this plurality of GAM66 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM66 RNA, herein designated GAM RNA, and which when bound by GAM66 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM66 target proteins.

[0782] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM66 gene, herein designated GAM GENE, on one or more GAM66 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background

section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0783] It is yet further appreciated that specific functions, and accordingly utilities, of GAM66 correlate with, and may be deduced from, the identity of the target genes which GAM66 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0784] Nucleotide sequences of the GAM66 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM66 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM66 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM66 are further described hereinbelow with reference to Table Table1.

[0785] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM66 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0786]

[0787] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 67 (GAM67) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0788] GAM67 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM67 was detected is described hereinabove with reference to Figs. 2-8.

[0789] GAM67 gene, herein designated GAM GENE, and GAM67 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0790] GAM67 gene, herein designated GAM GENE, encodes a GAM67 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM67 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nu-

cleotide sequence identical or highly similar to the nucleotide sequence of GAM67 precursor RNA is designated SEQ ID:737, and is provided hereinbelow with reference to the sequence listing part.

[0791] GAM67 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM67 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0792] An enzyme complex designated DICER COMPLEX, dices the GAM67 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM67 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 37%) nucleotide sequence of GAM67 RNA is

designated SEQ ID:738, and is provided hereinbelow with reference to the sequence listing part.

[0793] GAM67 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM67 target RNA, herein designated GAM TARGET RNA. GAM67 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0794] GAM67 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM67 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM67 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM67 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM67 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0795] The complementary binding of GAM67 RNA, herein designated GAM RNA, to target binding sites on GAM67 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM67 target RNA, herein designated GAM TARGET RNA, into GAM67 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0796] It is appreciated that GAM67 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM67 target genes. The mRNA of each one of this plurality of GAM67 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM67 RNA, herein designated GAM RNA, and which when bound by GAM67 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM67 target proteins.

[0797] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM67 gene, herein designated GAM GENE, on one or more GAM67 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0798] It is yet further appreciated that specific functions, and accordingly utilities, of GAM67 correlate with, and may be deduced from, the identity of the target genes which GAM67 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0799] Nucleotide sequences of the GAM67 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM67 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM67 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM67 are further described hereinbelow with reference to Table Table1.

[0800] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM67 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0801]

[0802] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 68 (GAM68) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0803] GAM68 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM68 was detected is described hereinabove

with reference to Figs. 2-8.

[0804] GAM68 gene, herein designated GAM GENE, and GAM68 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0805] GAM68 gene, herein designated GAM GENE, encodes a GAM68 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM68 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM68 precursor RNA is designated SEQ ID:748, and is provided hereinbelow with reference to the sequence listing part.

[0806] GAM68 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM68 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0807] An enzyme complex designated DICER COMPLEX, dices the GAM68 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM68 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 48%) nucleotide sequence of GAM68 RNA is designated SEQ ID:749, and is provided hereinbelow with reference to the sequence listing part.

[0808] GAM68 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM68 target RNA, herein designated GAM TARGET RNA. GAM68 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0809] GAM68 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM68 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM68 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM68 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM68 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0810] The complementary binding of GAM68 RNA, herein designated GAM RNA, to target binding sites on GAM68 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM68 target RNA, herein designated GAM TARGET RNA, into GAM68 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is there-

fore outlined by a broken line.

[0811] It is appreciated that GAM68 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM68 target genes. The mRNA of each one of this plurality of GAM68 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM68 RNA, herein designated GAM RNA, and which when bound by GAM68 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM68 target proteins.

[0812] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM68 gene, herein designated GAM GENE, on one or more GAM68 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes

by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0813] It is yet further appreciated that specific functions, and accordingly utilities, of GAM68 correlate with, and may be deduced from, the identity of the target genes which GAM68 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0814] Nucleotide sequences of the GAM68 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM68 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM68 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM68 are further described hereinbelow with reference to Table Table1.

[0815] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM68 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0816]

[0817] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 69 (GAM69) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0818] GAM69 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM69 was detected is described hereinabove with reference to Figs. 2-8.

[0819] GAM69 gene, herein designated GAM GENE, and GAM69 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0820] GAM69 gene, herein designated GAM GENE, encodes a GAM69 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM69 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM69 precursor RNA is designated SEQ ID:782, and is provided hereinbelow with reference to the sequence listing part.

[0821] GAM69 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM69 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0822] An enzyme complex designated DICER COMPLEX, dices the GAM69 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM69 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM69 RNA is designated SEQ ID:783, and is provided hereinbelow with reference to the sequence listing part.

[0823] GAM69 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM69 target RNA, herein designated GAM TARGET RNA. GAM69

target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0824] GAM69 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM69 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM69 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM69 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM69 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0825] The complementary binding of GAM69 RNA, herein designated GAM RNA, to target binding sites on GAM69 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM69 target RNA, herein designated GAM TARGET RNA, into GAM69 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0826] It is appreciated that GAM69 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM69 target genes. The mRNA of each one of this plurality of GAM69 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM69 RNA, herein designated GAM RNA, and which when bound by GAM69 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM69 target proteins.

[0827] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM69 gene, herein designated GAM GENE, on one or more GAM69 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0828] It is yet further appreciated that specific functions, and accordingly utilities, of GAM69 correlate with, and may be deduced from, the identity of the target genes which GAM69 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0829] Nucleotide sequences of the GAM69 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM69 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM69 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM69 are further described hereinbelow with reference to Table Table1.

[0830] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM69 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0831]

[0832] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 70 (GAM70) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0833] GAM70 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM70 was detected is described hereinabove with reference to Figs. 2-8.

[0834] GAM70 gene, herein designated GAM GENE, and GAM70 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0835] GAM70 gene, herein designated GAM GENE, encodes a

GAM70 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM70 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM70 precursor RNA is designated SEQ ID:792, and is provided hereinbelow with reference to the sequence listing part.

[0836] GAM70 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM70 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0837] An enzyme complex designated DICER COMPLEX, dices the GAM70 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM70 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured

RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM70 RNA is designated SEQ ID:793, and is provided hereinbelow with reference to the sequence listing part.

[0838] GAM70 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM70 target RNA, herein designated GAM TARGET RNA. GAM70 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0839] GAM70 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM70 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM70 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM70 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM70 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0840] The complementary binding of GAM70 RNA, herein designated GAM RNA, to target binding sites on GAM70 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM70 target RNA, herein designated GAM TARGET RNA, into GAM70 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0841] It is appreciated that GAM70 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM70 target genes. The mRNA of each one of this plurality of GAM70 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM70 RNA, herein designated GAM RNA, and which when bound by GAM70 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM70 target proteins.

[0842] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM70 gene, herein designated GAM GENE, on one or more GAM70 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0843] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM70 correlate with, and may be deduced from, the identity of the target genes which GAM70 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0844] Nucleotide sequences of the GAM70 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM70 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM70 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM70 are further described hereinbelow with reference to Table Table1.

[0845] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM70 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0846]

[0847] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 71 (GAM71) gene, which modulates expression of respective target genes thereof, the function and utility of which

target genes is known in the art.

[0848] GAM71 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM71 was detected is described hereinabove with reference to Figs. 2-8.

[0849] GAM71 gene, herein designated GAM GENE, and GAM71 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0850] GAM71 gene, herein designated GAM GENE, encodes a GAM71 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM71 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM71 precursor RNA is designated SEQ ID:818, and is provided hereinbelow with reference to the sequence listing part.

[0851] GAM71 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM71 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0852] An enzyme complex designated DICER COMPLEX, dices the GAM71 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM71 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM71 RNA is designated SEQ ID:819, and is provided hereinbelow with reference to the sequence listing part.

[0853] GAM71 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM71 target RNA, herein designated GAM TARGET RNA. GAM71 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0854] GAM71 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM71 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM71 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM71 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM71 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0855] The complementary binding of GAM71 RNA, herein designated GAM RNA, to target binding sites on GAM71 target RNA, herein designated GAM TARGET RNA, such as BIND-

ING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM71 target RNA, herein designated GAM TARGET RNA, into GAM71 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0856] It is appreciated that GAM71 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM71 target genes. The mRNA of each one of this plurality of GAM71 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM71 RNA, herein designated GAM RNA, and which when bound by GAM71 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM71 target proteins.

[0857] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM71 gene, herein designated GAM GENE, on one or more GAM71 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site

has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0858] It is yet further appreciated that specific functions, and accordingly utilities, of GAM71 correlate with, and may be deduced from, the identity of the target genes which GAM71 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0859] Nucleotide sequences of the GAM71 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM71 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM71 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM71 are further described hereinbelow with reference to Table Table1.

[0860] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM71 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0861]

[0862] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 72 (GAM72) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0863] GAM72 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM72 was detected is described hereinabove with reference to Figs. 2-8.

[0864] GAM72 gene, herein designated GAM GENE, and GAM72 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0865] GAM72 gene, herein designated GAM GENE, encodes a GAM72 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM72 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM72 precursor RNA is designated SEQ ID:835, and is provided hereinbelow with reference to the sequence listing part.

[0866] GAM72 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM72 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0867] An enzyme complex designated DICER COMPLEX, dices the GAM72 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM72 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM72 RNA is designated SEQ ID:836, and is provided hereinbelow with

reference to the sequence listing part.

[0868] GAM72 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM72 target RNA, herein designated GAM TARGET RNA. GAM72 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0869] GAM72 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM72 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM72 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM72 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM72 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0870] The complementary binding of GAM72 RNA, herein designated GAM RNA, to target binding sites on GAM72 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM72 target RNA, herein designated GAM TARGET RNA, into GAM72 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0871] It is appreciated that GAM72 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM72 target genes. The mRNA of each one of this plurality of GAM72 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM72 RNA, herein designated GAM RNA, and which when bound by GAM72 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM72 tar-

get proteins.

[0872] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM72 gene, herein designated GAM GENE, on one or more GAM72 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0873] It is yet further appreciated that specific functions, and accordingly utilities, of GAM72 correlate with, and may be deduced from, the identity of the target genes which GAM72 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0874] Nucleotide sequences of the GAM72 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM72 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM72 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM72 are further described hereinbelow with reference to Table Table1.

[0875] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM72 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0876]

[0877] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 73 (GAM73) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0878] GAM73 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM73 was detected is described hereinabove with reference to Figs. 2-8.

[0879] GAM73 gene, herein designated GAM GENE, and GAM73 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0880] GAM73 gene, herein designated GAM GENE, encodes a GAM73 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM73 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM73 precursor RNA is designated SEQ ID:845, and is provided hereinbelow with reference to the sequence listing part.

[0881] GAM73 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM73 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0882] An enzyme complex designated DICER COMPLEX, dices

the GAM73 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM73 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM73 RNA is designated SEQ ID:846, and is provided hereinbelow with reference to the sequence listing part.

[0883] GAM73 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM73 target RNA, herein designated GAM TARGET RNA. GAM73 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0884] GAM73 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM73 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM73

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM73 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM73 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0885] The complementary binding of GAM73 RNA, herein designated GAM RNA, to target binding sites on GAM73 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM73 target RNA, herein designated GAM TARGET RNA, into GAM73 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0886] It is appreciated that GAM73 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM73 target genes. The mRNA of each one of this plurality of GAM73 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM73 RNA, herein designated GAM RNA, and which when bound by GAM73 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM73 target proteins.

[0887] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM73 gene, herein designated GAM GENE, on one or more GAM73 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complemen-

tary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0888] It is yet further appreciated that specific functions, and accordingly utilities, of GAM73 correlate with, and may be deduced from, the identity of the target genes which GAM73 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0889] Nucleotide sequences of the GAM73 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM73 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM73 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM73 are further described hereinbelow with reference to Table Table1.

[0890] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM73 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0891]

[0892] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 74 (GAM74) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0893] GAM74 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM74 was detected is described hereinabove with reference to Figs. 2-8.

[0894] GAM74 gene, herein designated GAM GENE, and GAM74 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0895] GAM74 gene, herein designated GAM GENE, encodes a GAM74 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM74 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM74 precursor RNA is designated SEQ ID:862, and is provided hereinbelow with reference to the sequence listing part.

[0896] GAM74 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM74 folded precur-

sor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0897] An enzyme complex designated DICER COMPLEX, dices the GAM74 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM74 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM74 RNA is designated SEQ ID:863, and is provided hereinbelow with reference to the sequence listing part.

[0898] GAM74 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM74 target RNA, herein designated GAM TARGET RNA. GAM74 target RNA, herein designated GAM TARGET RNA, com-

prises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0899] GAM74 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM74 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM74 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM74 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM74 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[0900] The complementary binding of GAM74 RNA, herein designated GAM RNA, to target binding sites on GAM74 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM74 target RNA, herein designated GAM TARGET RNA, into GAM74 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0901] It is appreciated that GAM74 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM74 target genes. The mRNA of each one of this plurality of GAM74 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM74 RNA, herein designated GAM RNA, and which when bound by GAM74 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM74 target proteins.

[0902] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM74 gene, herein designated GAM GENE, on one or

more GAM74 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0903] It is yet further appreciated that specific functions, and accordingly utilities, of GAM74 correlate with, and may be deduced from, the identity of the target genes which GAM74 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0904] Nucleotide sequences of the GAM74 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM74 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM74 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM74 are further described

hereinbelow with reference to Table Table1.

[0905] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM74 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0906]

[0907] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 75 (GAM75) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0908] GAM75 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM75 was detected is described hereinabove with reference to Figs. 2-8.

[0909] GAM75 gene, herein designated GAM GENE, and GAM75 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0910] GAM75 gene, herein designated GAM GENE, encodes a GAM75 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM75 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM75 precursor RNA is designated SEQ ID:876, and is provided hereinbelow with reference to the sequence listing part.

[0911] GAM75 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM75 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0912] An enzyme complex designated DICER COMPLEX, dices the GAM75 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM75 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is

catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM75 RNA is designated SEQ ID:877, and is provided hereinbelow with reference to the sequence listing part.

[0913] GAM75 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM75 target RNA, herein designated GAM TARGET RNA. GAM75 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0914] GAM75 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM75 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM75 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM75 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM75 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0915] The complementary binding of GAM75 RNA, herein designated GAM RNA, to target binding sites on GAM75 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM75 target RNA, herein designated GAM TARGET RNA, into GAM75 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0916] It is appreciated that GAM75 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM75 target genes. The mRNA of each one of this plurality of GAM75 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM75 RNA, herein designated GAM RNA, and which when bound by GAM75 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM75 target proteins.

[0917] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM75 gene, herein designated GAM GENE, on one or more GAM75 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0918] It is yet further appreciated that specific functions, and accordingly utilities, of GAM75 correlate with, and may be

deduced from, the identity of the target genes which GAM75 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0919] Nucleotide sequences of the GAM75 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM75 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM75 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM75 are further described hereinbelow with reference to Table Table1.

[0920] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM75 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0921]

[0922] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 76 (GAM76) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0923] GAM76 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM76 was detected is described hereinabove with reference to Figs. 2-8.

[0924] GAM76 gene, herein designated GAM GENE, and GAM76 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0925] GAM76 gene, herein designated GAM GENE, encodes a GAM76 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM76 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM76 precursor RNA is designated SEQ ID:905, and is provided hereinbelow with reference to the sequence listing part.

[0926] GAM76 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM76 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0927] An enzyme complex designated DICER COMPLEX, dices the GAM76 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM76 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM76 RNA is designated SEQ ID:906, and is provided hereinbelow with reference to the sequence listing part.

[0928] GAM76 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM76 target RNA, herein designated GAM TARGET RNA. GAM76 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0929] GAM76 RNA, herein designated GAM RNA, binds comple-

mentarily to one or more target binding sites located in untranslated regions of GAM76 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM76 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM76 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM76 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0930] The complementary binding of GAM76 RNA, herein designated GAM RNA, to target binding sites on GAM76 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits

translation of GAM76 target RNA, herein designated GAM TARGET RNA, into GAM76 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0931] It is appreciated that GAM76 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM76 target genes. The mRNA of each one of this plurality of GAM76 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM76 RNA, herein designated GAM RNA, and which when bound by GAM76 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM76 target proteins.

[0932] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM76 gene, herein designated GAM GENE, on one or more GAM76 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known

miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0933] It is yet further appreciated that specific functions, and accordingly utilities, of GAM76 correlate with, and may be deduced from, the identity of the target genes which GAM76 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0934] Nucleotide sequences of the GAM76 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM76 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM76 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM76 are further described hereinbelow with reference to Table Table1.

[0935] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM76 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0936]

[0937] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 77 (GAM77) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0938] GAM77 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM77 was detected is described hereinabove with reference to Figs. 2-8.

[0939] GAM77 gene, herein designated GAM GENE, and GAM77 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0940] GAM77 gene, herein designated GAM GENE, encodes a GAM77 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM77 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM77 precursor RNA is designated

SEQ ID:919, and is provided hereinbelow with reference to the sequence listing part.

[0941] GAM77 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM77 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0942] An enzyme complex designated DICER COMPLEX, dices the GAM77 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM77 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 91%) nucleotide sequence of GAM77 RNA is designated SEQ ID:920, and is provided hereinbelow with reference to the sequence listing part.

[0943] GAM77 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM77 target RNA, herein designated GAM TARGET RNA. GAM77 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0944] GAM77 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM77 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM77 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM77 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM77 tar-

get RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0945] The complementary binding of GAM77 RNA, herein designated GAM RNA, to target binding sites on GAM77 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM77 target RNA, herein designated GAM TARGET RNA, into GAM77 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0946] It is appreciated that GAM77 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM77 target genes. The mRNA of each one of this plurality of GAM77 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM77 RNA, herein designated GAM RNA, and which when bound by GAM77 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM77 target proteins.

[0947] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM77 gene, herein designated GAM GENE, on one or more GAM77 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0948] It is yet further appreciated that specific functions, and accordingly utilities, of GAM77 correlate with, and may be deduced from, the identity of the target genes which GAM77 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0949] Nucleotide sequences of the GAM77 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM77 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM77 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM77 are further described hereinbelow with reference to Table Table1.

[0950] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM77 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0951]

[0952] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 78 (GAM78) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0953] GAM78 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM78 was detected is described hereinabove with reference to Figs. 2-8.

[0954] GAM78 gene, herein designated GAM GENE, and GAM78

target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0955] GAM78 gene, herein designated GAM GENE, encodes a GAM78 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM78 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM78 precursor RNA is designated SEQ ID:938, and is provided hereinbelow with reference to the sequence listing part.

[0956] GAM78 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM78 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0957] An enzyme complex designated DICER COMPLEX, dices the GAM78 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, into GAM78 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 37%) nucleotide sequence of GAM78 RNA is designated SEQ ID:939, and is provided hereinbelow with reference to the sequence listing part.

[0958] GAM78 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM78 target RNA, herein designated GAM TARGET RNA. GAM78 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0959] GAM78 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM78 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM78 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM78 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM78 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0960] The complementary binding of GAM78 RNA, herein designated GAM RNA, to target binding sites on GAM78 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM78 target RNA, herein designated GAM TARGET RNA, into GAM78 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0961] It is appreciated that GAM78 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM78 target genes. The mRNA of each one of this plurality of GAM78 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM78 RNA, herein designated GAM RNA, and which when bound by GAM78 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM78 target proteins.

[0962] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM78 gene, herein designated GAM GENE, on one or more GAM78 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not

yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0963] It is yet further appreciated that specific functions, and accordingly utilities, of GAM78 correlate with, and may be deduced from, the identity of the target genes which GAM78 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0964] Nucleotide sequences of the GAM78 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM78 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM78 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM78 are further described hereinbelow with reference to Table Table1.

[0965] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM78 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0966]

[0967] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 79 (GAM79) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0968] GAM79 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM79 was detected is described hereinabove with reference to Figs. 2-8.

[0969] GAM79 gene, herein designated GAM GENE, and GAM79 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0970] GAM79 gene, herein designated GAM GENE, encodes a GAM79 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM79 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM79 precursor RNA is designated SEQ ID:952, and is provided hereinbelow with reference to the sequence listing part.

[0971] GAM79 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM79 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0972] An enzyme complex designated DICER COMPLEX, dices the GAM79 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM79 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM79 RNA is designated SEQ ID:953, and is provided hereinbelow with reference to the sequence listing part.

[0973] GAM79 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM79 target RNA, herein designated GAM TARGET RNA. GAM79 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein

coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0974] GAM79 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM79 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM79 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM79 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM79 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0975] The complementary binding of GAM79 RNA, herein designated GAM RNA, to target binding sites on GAM79 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM79 target RNA, herein designated GAM TARGET RNA, into GAM79 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0976] It is appreciated that GAM79 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM79 target genes. The mRNA of each one of this plurality of GAM79 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM79 RNA, herein designated GAM RNA, and which when bound by GAM79 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM79 target proteins.

[0977] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM79 gene, herein designated GAM GENE, on one or more GAM79 target gene, herein designated GAM TARGET

GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0978] It is yet further appreciated that specific functions, and accordingly utilities, of GAM79 correlate with, and may be deduced from, the identity of the target genes which GAM79 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0979] Nucleotide sequences of the GAM79 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM79 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM79 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM79 are further described hereinbelow with reference to Table Table1.

[0980] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM79 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0981]

[0982] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 80 (GAM80) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0983] GAM80 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM80 was detected is described hereinabove with reference to Figs. 2-8.

[0984] GAM80 gene, herein designated GAM GENE, and GAM80 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[0985] GAM80 gene, herein designated GAM GENE, encodes a GAM80 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM80 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM80 precursor RNA is designated SEQ ID:971, and is provided hereinbelow with reference to the sequence listing part.

[0986] GAM80 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM80 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[0987] An enzyme complex designated DICER COMPLEX, dices the GAM80 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM80 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme

called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM80 RNA is designated SEQ ID:972, and is provided hereinbelow with reference to the sequence listing part.

[0988] GAM80 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM80 target RNA, herein designated GAM TARGET RNA. GAM80 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[0989] GAM80 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM80 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM80 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM80 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM80 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[0990] The complementary binding of GAM80 RNA, herein designated GAM RNA, to target binding sites on GAM80 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM80 target RNA, herein designated GAM TARGET RNA, into GAM80 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[0991] It is appreciated that GAM80 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM80 target genes. The mRNA of each one of this plurality of GAM80 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM80 RNA,

herein designated GAM RNA, and which when bound by GAM80 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM80 target proteins.

[0992] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM80 gene, herein designated GAM GENE, on one or more GAM80 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[0993] It is yet further appreciated that specific functions, and accordingly utilities, of GAM80 correlate with, and may be deduced from, the identity of the target genes which

GAM80 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[0994] Nucleotide sequences of the GAM80 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM80 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM80 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM80 are further described hereinbelow with reference to Table Table1.

[0995] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM80 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[0996]

[0997] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 81 (GAM81) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[0998] GAM81 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM81 was detected is described hereinabove with reference to Figs. 2-8.

[0999] GAM81 gene, herein designated GAM GENE, and GAM81 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1000] GAM81 gene, herein designated GAM GENE, encodes a GAM81 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM81 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM81 precursor RNA is designated SEQ ID:997, and is provided hereinbelow with reference to the sequence listing part.

[1001] GAM81 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM81 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[1002] An enzyme complex designated DICER COMPLEX, dices the GAM81 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM81 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 35%) nucleotide sequence of GAM81 RNA is designated SEQ ID:998, and is provided hereinbelow with reference to the sequence listing part.

[1003] GAM81 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM81 target RNA, herein designated GAM TARGET RNA. GAM81 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1004] GAM81 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM81 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM81 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM81 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM81 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1005] The complementary binding of GAM81 RNA, herein designated GAM RNA, to target binding sites on GAM81 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM81 target RNA, herein designated GAM

TARGET RNA, into GAM81 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1006] It is appreciated that GAM81 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM81 target genes. The mRNA of each one of this plurality of GAM81 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM81 RNA, herein designated GAM RNA, and which when bound by GAM81 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM81 target proteins.

[1007] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM81 gene, herein designated GAM GENE, on one or more GAM81 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other re-

cently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1008] It is yet further appreciated that specific functions, and accordingly utilities, of GAM81 correlate with, and may be deduced from, the identity of the target genes which GAM81 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1009] Nucleotide sequences of the GAM81 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM81 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM81 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM81 are further described hereinbelow with reference to Table Table1.

[1010] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM81 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[1011]

[1012] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 82 (GAM82) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1013] GAM82 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM82 was detected is described hereinabove with reference to Figs. 2-8.

[1014] GAM82 gene, herein designated GAM GENE, and GAM82 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1015] GAM82 gene, herein designated GAM GENE, encodes a GAM82 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM82 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM82 precursor RNA is designated SEQ ID:1017, and is provided hereinbelow with reference

to the sequence listing part.

[1016] GAM82 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM82 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1017] An enzyme complex designated DICER COMPLEX, dices the GAM82 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM82 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM82 RNA is designated SEQ ID:1018, and is provided hereinbelow with reference to the sequence listing part.

[1018] GAM82 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM82 target RNA, herein designated GAM TARGET RNA. GAM82 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1019] GAM82 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM82 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM82 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM82 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM82 target RNA, herein designated GAM TARGET RNA. It is further

appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1020] The complementary binding of GAM82 RNA, herein designated GAM RNA, to target binding sites on GAM82 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM82 target RNA, herein designated GAM TARGET RNA, into GAM82 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1021] It is appreciated that GAM82 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM82 target genes. The mRNA of each one of this plurality of GAM82 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM82 RNA, herein designated GAM RNA, and which when bound by GAM82 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM82 target proteins.

[1022] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM82 gene, herein designated GAM GENE, on one or more GAM82 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1023] It is yet further appreciated that specific functions, and accordingly utilities, of GAM82 correlate with, and may be deduced from, the identity of the target genes which GAM82 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1024] Nucleotide sequences of the GAM82 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM82 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM82 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM82 are further described hereinbelow with reference to Table Table1.

[1025] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM82 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1026]

[1027] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 83 (GAM83) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1028] GAM83 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM83 was detected is described hereinabove with reference to Figs. 2-8.

[1029] GAM83 gene, herein designated GAM GENE, and GAM83 target gene, herein designated GAM TARGET GENE, are

human genes contained in the human genome.

[1030] GAM83 gene, herein designated GAM GENE, encodes a GAM83 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM83 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM83 precursor RNA is designated SEQ ID:1037, and is provided hereinbelow with reference to the sequence listing part.

[1031] GAM83 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM83 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1032] An enzyme complex designated DICER COMPLEX, dices the GAM83 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM83 RNA, herein desig-

nated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM83 RNA is designated SEQ ID:1038, and is provided hereinbelow with reference to the sequence listing part.

[1033] GAM83 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM83 target RNA, herein designated GAM TARGET RNA. GAM83 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1034] GAM83 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM83 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM83 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM83 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM83 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1035] The complementary binding of GAM83 RNA, herein designated GAM RNA, to target binding sites on GAM83 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM83 target RNA, herein designated GAM TARGET RNA, into GAM83 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1036] It is appreciated that GAM83 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM83 target genes. The mRNA of each one of this plurality of GAM83 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM83 RNA, herein designated GAM RNA, and which when bound by GAM83 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM83 target proteins.

[1037] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM83 gene, herein designated GAM GENE, on one or more GAM83 target gene, herein designated GAM TARGET

GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1038] It is yet further appreciated that specific functions, and accordingly utilities, of GAM83 correlate with, and may be deduced from, the identity of the target genes which GAM83 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1039] Nucleotide sequences of the GAM83 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM83 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM83 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM83 are further described hereinbelow with reference to Table Table1.

[1040] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM83 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1041]

[1042] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 84 (GAM84) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1043] GAM84 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM84 was detected is described hereinabove with reference to Figs. 2-8.

[1044] GAM84 gene, herein designated GAM GENE, and GAM84 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1045] GAM84 gene, herein designated GAM GENE, encodes a GAM84 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM84 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM84 precursor RNA is designated SEQ ID:1061, and is provided hereinbelow with reference to the sequence listing part.

[1046] GAM84 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM84 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1047] An enzyme complex designated DICER COMPLEX, dices the GAM84 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM84 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme

called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM84 RNA is designated SEQ ID:1062, and is provided hereinbelow with reference to the sequence listing part.

[1048] GAM84 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM84 target RNA, herein designated GAM TARGET RNA. GAM84 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1049] GAM84 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM84 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM84 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM84 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM84 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1050] The complementary binding of GAM84 RNA, herein designated GAM RNA, to target binding sites on GAM84 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM84 target RNA, herein designated GAM TARGET RNA, into GAM84 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1051] It is appreciated that GAM84 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM84 target genes. The mRNA of each one of this plurality of GAM84 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM84 RNA,

herein designated GAM RNA, and which when bound by GAM84 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM84 target proteins.

[1052] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM84 gene, herein designated GAM GENE, on one or more GAM84 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1053] It is yet further appreciated that specific functions, and accordingly utilities, of GAM84 correlate with, and may be deduced from, the identity of the target genes which

GAM84 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1054] Nucleotide sequences of the GAM84 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM84 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM84 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM84 are further described hereinbelow with reference to Table Table1.

[1055] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM84 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1056]

[1057] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 85 (GAM85) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1058] GAM85 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM85 was detected is described hereinabove with reference to Figs. 2-8.

[1059] GAM85 gene, herein designated GAM GENE, and GAM85 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1060] GAM85 gene, herein designated GAM GENE, encodes a GAM85 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM85 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM85 precursor RNA is designated SEQ ID:1078, and is provided hereinbelow with reference to the sequence listing part.

[1061] GAM85 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM85 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[1062] An enzyme complex designated DICER COMPLEX, dices the GAM85 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM85 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 18%) nucleotide sequence of GAM85 RNA is designated SEQ ID:1079, and is provided hereinbelow with reference to the sequence listing part.

[1063] GAM85 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM85 target RNA, herein designated GAM TARGET RNA. GAM85 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1064] GAM85 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM85 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM85 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM85 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM85 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1065] The complementary binding of GAM85 RNA, herein designated GAM RNA, to target binding sites on GAM85 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM85 target RNA, herein designated GAM

TARGET RNA, into GAM85 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1066] It is appreciated that GAM85 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM85 target genes. The mRNA of each one of this plurality of GAM85 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM85 RNA, herein designated GAM RNA, and which when bound by GAM85 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM85 target proteins.

[1067] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM85 gene, herein designated GAM GENE, on one or more GAM85 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other re-

cently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1068] It is yet further appreciated that specific functions, and accordingly utilities, of GAM85 correlate with, and may be deduced from, the identity of the target genes which GAM85 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1069] Nucleotide sequences of the GAM85 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM85 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM85 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM85 are further described hereinbelow with reference to Table Table1.

[1070] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM85 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[1071]

[1072] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 86 (GAM86) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1073] GAM86 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM86 was detected is described hereinabove with reference to Figs. 2-8.

[1074] GAM86 gene, herein designated GAM GENE, and GAM86 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1075] GAM86 gene, herein designated GAM GENE, encodes a GAM86 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM86 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM86 precursor RNA is designated SEQ ID:1095, and is provided hereinbelow with reference

to the sequence listing part.

[1076] GAM86 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM86 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1077] An enzyme complex designated DICER COMPLEX, dices the GAM86 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM86 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM86 RNA is designated SEQ ID:1096, and is provided hereinbelow with reference to the sequence listing part.

[1078] GAM86 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM86 target RNA, herein designated GAM TARGET RNA. GAM86 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1079] GAM86 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM86 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM86 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM86 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM86 target RNA, herein designated GAM TARGET RNA. It is further

appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1080] The complementary binding of GAM86 RNA, herein designated GAM RNA, to target binding sites on GAM86 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM86 target RNA, herein designated GAM TARGET RNA, into GAM86 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1081] It is appreciated that GAM86 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM86 target genes. The mRNA of each one of this plurality of GAM86 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM86 RNA, herein designated GAM RNA, and which when bound by GAM86 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM86 target proteins.

[1082] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM86 gene, herein designated GAM GENE, on one or more GAM86 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1083] It is yet further appreciated that specific functions, and accordingly utilities, of GAM86 correlate with, and may be deduced from, the identity of the target genes which GAM86 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1084] Nucleotide sequences of the GAM86 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM86 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM86 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM86 are further described hereinbelow with reference to Table Table1.

[1085] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM86 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1086]

[1087] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 87 (GAM87) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1088] GAM87 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM87 was detected is described hereinabove with reference to Figs. 2-8.

[1089] GAM87 gene, herein designated GAM GENE, and GAM87 target gene, herein designated GAM TARGET GENE, are

human genes contained in the human genome.

[1090] GAM87 gene, herein designated GAM GENE, encodes a GAM87 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM87 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM87 precursor RNA is designated SEQ ID:1111, and is provided hereinbelow with reference to the sequence listing part.

[1091] GAM87 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM87 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1092] An enzyme complex designated DICER COMPLEX, dices the GAM87 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM87 RNA, herein desig-

nated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM87 RNA is designated SEQ ID:1112, and is provided hereinbelow with reference to the sequence listing part.

[1093] GAM87 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM87 target RNA, herein designated GAM TARGET RNA. GAM87 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1094] GAM87 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM87 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM87 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM87 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM87 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1095] The complementary binding of GAM87 RNA, herein designated GAM RNA, to target binding sites on GAM87 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM87 target RNA, herein designated GAM TARGET RNA, into GAM87 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1096] It is appreciated that GAM87 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM87 target genes. The mRNA of each one of this plurality of GAM87 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM87 RNA, herein designated GAM RNA, and which when bound by GAM87 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM87 target proteins.

[1097] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM87 gene, herein designated GAM GENE, on one or more GAM87 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny

RNA world, Science 294,779 (2001)).

[1098] It is yet further appreciated that specific functions, and accordingly utilities, of GAM87 correlate with, and may be deduced from, the identity of the target genes which GAM87 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1099] Nucleotide sequences of the GAM87 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM87 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM87 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM87 are further described hereinbelow with reference to Table Table1.

[1100] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM87 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1101]

[1102] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

88 (GAM88) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1103] GAM88 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM88 was detected is described hereinabove with reference to Figs. 2-8.

[1104] GAM88 gene, herein designated GAM GENE, and GAM88 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1105] GAM88 gene, herein designated GAM GENE, encodes a GAM88 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM88 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM88 precursor RNA is designated SEQ ID:1124, and is provided hereinbelow with reference to the sequence listing part.

[1106] GAM88 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM88 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1107] An enzyme complex designated DICER COMPLEX, dices the GAM88 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM88 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM88 RNA is designated SEQ ID:1125, and is provided hereinbelow with reference to the sequence listing part.

[1108] GAM88 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM88 target RNA, herein designated GAM TARGET RNA. GAM88 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding re-

gion and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1109] GAM88 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM88 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM88 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM88 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM88 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1110] The complementary binding of GAM88 RNA, herein desig-

nated GAM RNA, to target binding sites on GAM88 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM88 target RNA, herein designated GAM TARGET RNA, into GAM88 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1111] It is appreciated that GAM88 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM88 target genes. The mRNA of each one of this plurality of GAM88 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM88 RNA, herein designated GAM RNA, and which when bound by GAM88 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM88 target proteins.

[1112] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM88 gene, herein designated GAM GENE, on one or more GAM88 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As

mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1113] It is yet further appreciated that specific functions, and accordingly utilities, of GAM88 correlate with, and may be deduced from, the identity of the target genes which GAM88 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1114] Nucleotide sequences of the GAM88 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM88 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM88 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM88 are further described hereinbelow with reference to Table Table1.

[1115] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM88 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1116]

[1117] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 89 (GAM89) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1118] GAM89 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM89 was detected is described hereinabove with reference to Figs. 2-8.

[1119] GAM89 gene, herein designated GAM GENE, and GAM89 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1120] GAM89 gene, herein designated GAM GENE, encodes a GAM89 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM89 precursor RNA, herein designated

GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM89 precursor RNA is designated SEQ ID:1140, and is provided hereinbelow with reference to the sequence listing part.

[1121] GAM89 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM89 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1122] An enzyme complex designated DICER COMPLEX, dices the GAM89 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM89 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A

probable (over 56%) nucleotide sequence of GAM89 RNA is designated SEQ ID:1141, and is provided hereinbelow with reference to the sequence listing part.

[1123] GAM89 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM89 target RNA, herein designated GAM TARGET RNA. GAM89 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1124] GAM89 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM89 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM89 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM89 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM89 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1125] The complementary binding of GAM89 RNA, herein designated GAM RNA, to target binding sites on GAM89 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM89 target RNA, herein designated GAM TARGET RNA, into GAM89 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1126] It is appreciated that GAM89 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM89 target genes. The mRNA of each one of this plurality of GAM89 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM89 RNA, herein designated GAM RNA, and which when bound by

GAM89 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM89 target proteins.

[1127] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM89 gene, herein designated GAM GENE, on one or more GAM89 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1128] It is yet further appreciated that specific functions, and accordingly utilities, of GAM89 correlate with, and may be deduced from, the identity of the target genes which GAM89 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[1129] Nucleotide sequences of the GAM89 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM89 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM89 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM89 are further described hereinbelow with reference to Table Table1.

[1130] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM89 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1131]

[1132] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 90 (GAM90) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1133] GAM90 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM90 was detected is described hereinabove with reference to Figs. 2-8.

[1134] GAM90 gene, herein designated GAM GENE, and GAM90 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1135] GAM90 gene, herein designated GAM GENE, encodes a GAM90 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM90 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM90 precursor RNA is designated SEQ ID:1159, and is provided hereinbelow with reference to the sequence listing part.

[1136] GAM90 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM90 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[1137] An enzyme complex designated DICER COMPLEX, dices the GAM90 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM90 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 82%) nucleotide sequence of GAM90 RNA is designated SEQ ID:1160, and is provided hereinbelow with reference to the sequence listing part.

[1138] GAM90 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM90 target RNA, herein designated GAM TARGET RNA. GAM90 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1139] GAM90 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM90 target RNA, herein desig-

nated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM90 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM90 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM90 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1140] The complementary binding of GAM90 RNA, herein designated GAM RNA, to target binding sites on GAM90 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM90 target RNA, herein designated GAM TARGET RNA, into GAM90 target protein, herein desig-

nated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1141] It is appreciated that GAM90 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM90 target genes. The mRNA of each one of this plurality of GAM90 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM90 RNA, herein designated GAM RNA, and which when bound by GAM90 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM90 target proteins.

[1142] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM90 gene, herein designated GAM GENE, on one or more GAM90 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those

skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1143] It is yet further appreciated that specific functions, and accordingly utilities, of GAM90 correlate with, and may be deduced from, the identity of the target genes which GAM90 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1144] Nucleotide sequences of the GAM90 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM90 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM90 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM90 are further described hereinbelow with reference to Table Table1.

[1145] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM90 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1146]

[1147] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 91 (GAM91) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1148] GAM91 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM91 was detected is described hereinabove with reference to Figs. 2-8.

[1149] GAM91 gene, herein designated GAM GENE, and GAM91 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1150] GAM91 gene, herein designated GAM GENE, encodes a GAM91 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM91 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM91 precursor RNA is designated SEQ ID:1168, and is provided hereinbelow with reference to the sequence listing part.

[1151] GAM91 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM91 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1152] An enzyme complex designated DICER COMPLEX, dices the GAM91 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM91 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM91 RNA is designated SEQ ID:1169, and is provided hereinbelow with reference to the sequence listing part.

[1153] GAM91 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM91

target RNA, herein designated GAM TARGET RNA. GAM91 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1154] GAM91 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM91 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM91 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM91 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM91 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites

in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1155] The complementary binding of GAM91 RNA, herein designated GAM RNA, to target binding sites on GAM91 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM91 target RNA, herein designated GAM TARGET RNA, into GAM91 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1156] It is appreciated that GAM91 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM91 target genes. The mRNA of each one of this plurality of GAM91 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM91 RNA, herein designated GAM RNA, and which when bound by GAM91 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM91 target proteins.

[1157] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM91 gene, herein designated GAM GENE, on one or more GAM91 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1158] It is yet further appreciated that specific functions, and accordingly utilities, of GAM91 correlate with, and may be deduced from, the identity of the target genes which GAM91 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1159] Nucleotide sequences of the GAM91 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM91 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM91 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM91 are further described hereinbelow with reference to Table Table1.

[1160] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM91 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1161]

[1162] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 92 (GAM92) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1163] GAM92 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM92 was detected is described hereinabove with reference to Figs. 2-8.

[1164] GAM92 gene, herein designated GAM GENE, and GAM92 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1165] GAM92 gene, herein designated GAM GENE, encodes a GAM92 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM92 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM92 precursor RNA is designated SEQ ID:1171, and is provided hereinbelow with reference to the sequence listing part.

[1166] GAM92 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM92 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1167] An enzyme complex designated DICER COMPLEX, dices the GAM92 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM92 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA seg-

ment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM92 RNA is designated SEQ ID:1172, and is provided hereinbelow with reference to the sequence listing part.

[1168] GAM92 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM92 target RNA, herein designated GAM TARGET RNA. GAM92 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1169] GAM92 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM92 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM92 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM92 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM92 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1170] The complementary binding of GAM92 RNA, herein designated GAM RNA, to target binding sites on GAM92 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM92 target RNA, herein designated GAM TARGET RNA, into GAM92 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1171] It is appreciated that GAM92 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM92 target genes. The mRNA of each one of this plu-

rality of GAM92 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM92 RNA, herein designated GAM RNA, and which when bound by GAM92 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM92 target proteins.

[1172] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM92 gene, herein designated GAM GENE, on one or more GAM92 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1173] It is yet further appreciated that specific functions, and accordingly utilities, of GAM92 correlate with, and may be deduced from, the identity of the target genes which GAM92 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1174] Nucleotide sequences of the GAM92 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM92 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM92 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM92 are further described hereinbelow with reference to Table Table1.

[1175] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM92 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1176]

[1177] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 93 (GAM93) gene, which modulates expression of respec-

tive target genes thereof, the function and utility of which target genes is known in the art.

[1178] GAM93 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM93 was detected is described hereinabove with reference to Figs. 2-8.

[1179] GAM93 gene, herein designated GAM GENE, and GAM93 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1180] GAM93 gene, herein designated GAM GENE, encodes a GAM93 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM93 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM93 precursor RNA is designated SEQ ID:1196, and is provided hereinbelow with reference to the sequence listing part.

[1181] GAM93 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM93 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1182] An enzyme complex designated DICER COMPLEX, dices the GAM93 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM93 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM93 RNA is designated SEQ ID:1197, and is provided hereinbelow with reference to the sequence listing part.

[1183] GAM93 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM93 target RNA, herein designated GAM TARGET RNA. GAM93 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PRO-

TEIN CODING and 3UTR respectively.

[1184] GAM93 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM93 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM93 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM93 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM93 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1185] The complementary binding of GAM93 RNA, herein designated GAM RNA, to target binding sites on GAM93 target

RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM93 target RNA, herein designated GAM TARGET RNA, into GAM93 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1186] It is appreciated that GAM93 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM93 target genes. The mRNA of each one of this plurality of GAM93 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM93 RNA, herein designated GAM RNA, and which when bound by GAM93 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM93 target proteins.

[1187] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM93 gene, herein designated GAM GENE, on one or more GAM93 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background

section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1188] It is yet further appreciated that specific functions, and accordingly utilities, of GAM93 correlate with, and may be deduced from, the identity of the target genes which GAM93 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1189] Nucleotide sequences of the GAM93 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM93 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM93 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM93 are further described hereinbelow with reference to Table Table1.

[1190] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM93 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1191]

[1192] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 94 (GAM94) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1193] GAM94 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM94 was detected is described hereinabove with reference to Figs. 2-8.

[1194] GAM94 gene, herein designated GAM GENE, and GAM94 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1195] GAM94 gene, herein designated GAM GENE, encodes a GAM94 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM94 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nu-

cleotide sequence identical or highly similar to the nucleotide sequence of GAM94 precursor RNA is designated SEQ ID:1206, and is provided hereinbelow with reference to the sequence listing part.

[1196] GAM94 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM94 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1197] An enzyme complex designated DICER COMPLEX, dices the GAM94 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM94 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM94 RNA is

designated SEQ ID:1207, and is provided hereinbelow with reference to the sequence listing part.

[1198] GAM94 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM94 target RNA, herein designated GAM TARGET RNA. GAM94 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1199] GAM94 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM94 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM94 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM94 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM94 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1200] The complementary binding of GAM94 RNA, herein designated GAM RNA, to target binding sites on GAM94 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM94 target RNA, herein designated GAM TARGET RNA, into GAM94 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1201] It is appreciated that GAM94 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM94 target genes. The mRNA of each one of this plurality of GAM94 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM94 RNA, herein designated GAM RNA, and which when bound by GAM94 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM94 target proteins.

[1202] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM94 gene, herein designated GAM GENE, on one or more GAM94 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1203] It is yet further appreciated that specific functions, and accordingly utilities, of GAM94 correlate with, and may be deduced from, the identity of the target genes which GAM94 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1204] Nucleotide sequences of the GAM94 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM94 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM94 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM94 are further described hereinbelow with reference to Table Table1.

[1205] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM94 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1206]

[1207] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 95 (GAM95) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1208] GAM95 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM95 was detected is described hereinabove

with reference to Figs. 2-8.

[1209] GAM95 gene, herein designated GAM GENE, and GAM95 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1210] GAM95 gene, herein designated GAM GENE, encodes a GAM95 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM95 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM95 precursor RNA is designated SEQ ID:1220, and is provided hereinbelow with reference to the sequence listing part.

[1211] GAM95 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM95 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1212] An enzyme complex designated DICER COMPLEX, dices the GAM95 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM95 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 37%) nucleotide sequence of GAM95 RNA is designated SEQ ID:1221, and is provided hereinbelow with reference to the sequence listing part.

[1213] GAM95 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM95 target RNA, herein designated GAM TARGET RNA. GAM95 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1214] GAM95 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM95 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM95 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM95 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM95 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1215] The complementary binding of GAM95 RNA, herein designated GAM RNA, to target binding sites on GAM95 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM95 target RNA, herein designated GAM TARGET RNA, into GAM95 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is there-

fore outlined by a broken line.

[1216] It is appreciated that GAM95 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM95 target genes. The mRNA of each one of this plurality of GAM95 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM95 RNA, herein designated GAM RNA, and which when bound by GAM95 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM95 target proteins.

[1217] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM95 gene, herein designated GAM GENE, on one or more GAM95 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes

by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1218] It is yet further appreciated that specific functions, and accordingly utilities, of GAM95 correlate with, and may be deduced from, the identity of the target genes which GAM95 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1219] Nucleotide sequences of the GAM95 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM95 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM95 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM95 are further described hereinbelow with reference to Table Table1.

[1220] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM95 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1221]

[1222] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 96 (GAM96) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1223] GAM96 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM96 was detected is described hereinabove with reference to Figs. 2-8.

[1224] GAM96 gene, herein designated GAM GENE, and GAM96 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1225] GAM96 gene, herein designated GAM GENE, encodes a GAM96 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM96 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM96 precursor RNA is designated SEQ ID:1227, and is provided hereinbelow with reference to the sequence listing part.

[1226] GAM96 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM96 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1227] An enzyme complex designated DICER COMPLEX, dices the GAM96 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM96 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 34%) nucleotide sequence of GAM96 RNA is designated SEQ ID:1228, and is provided hereinbelow with reference to the sequence listing part.

[1228] GAM96 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM96 target RNA, herein designated GAM TARGET RNA. GAM96

target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1229] GAM96 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM96 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM96 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM96 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM96 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1230] The complementary binding of GAM96 RNA, herein designated GAM RNA, to target binding sites on GAM96 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM96 target RNA, herein designated GAM TARGET RNA, into GAM96 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1231] It is appreciated that GAM96 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM96 target genes. The mRNA of each one of this plurality of GAM96 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM96 RNA, herein designated GAM RNA, and which when bound by GAM96 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM96 target proteins.

[1232] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM96 gene, herein designated GAM GENE, on one or more GAM96 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1233] It is yet further appreciated that specific functions, and accordingly utilities, of GAM96 correlate with, and may be deduced from, the identity of the target genes which GAM96 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1234] Nucleotide sequences of the GAM96 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM96 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM96 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM96 are further described hereinbelow with reference to Table Table1.

[1235] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM96 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1236]

[1237] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 97 (GAM97) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1238] GAM97 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM97 was detected is described hereinabove with reference to Figs. 2-8.

[1239] GAM97 gene, herein designated GAM GENE, and GAM97 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1240] GAM97 gene, herein designated GAM GENE, encodes a

GAM97 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM97 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM97 precursor RNA is designated SEQ ID:1259, and is provided hereinbelow with reference to the sequence listing part.

[1241] GAM97 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM97 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1242] An enzyme complex designated DICER COMPLEX, dices the GAM97 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM97 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured

RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 18%) nucleotide sequence of GAM97 RNA is designated SEQ ID:1260, and is provided hereinbelow with reference to the sequence listing part.

[1243] GAM97 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM97 target RNA, herein designated GAM TARGET RNA. GAM97 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1244] GAM97 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM97 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM97 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM97 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM97 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1245] The complementary binding of GAM97 RNA, herein designated GAM RNA, to target binding sites on GAM97 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM97 target RNA, herein designated GAM TARGET RNA, into GAM97 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1246] It is appreciated that GAM97 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM97 target genes. The mRNA of each one of this plurality of GAM97 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM97 RNA, herein designated GAM RNA, and which when bound by GAM97 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM97 target proteins.

[1247] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM97 gene, herein designated GAM GENE, on one or more GAM97 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1248] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM97 correlate with, and may be deduced from, the identity of the target genes which GAM97 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1249] Nucleotide sequences of the GAM97 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM97 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM97 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM97 are further described hereinbelow with reference to Table Table1.

[1250] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM97 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1251]

[1252] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 98 (GAM98) gene, which modulates expression of respective target genes thereof, the function and utility of which

target genes is known in the art.

[1253] GAM98 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM98 was detected is described hereinabove with reference to Figs. 2-8.

[1254] GAM98 gene, herein designated GAM GENE, and GAM98 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1255] GAM98 gene, herein designated GAM GENE, encodes a GAM98 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM98 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM98 precursor RNA is designated SEQ ID:1276, and is provided hereinbelow with reference to the sequence listing part.

[1256] GAM98 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM98 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1257] An enzyme complex designated DICER COMPLEX, dices the GAM98 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM98 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM98 RNA is designated SEQ ID:1277, and is provided hereinbelow with reference to the sequence listing part.

[1258] GAM98 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM98 target RNA, herein designated GAM TARGET RNA. GAM98 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1259] GAM98 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM98 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM98 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM98 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM98 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1260] The complementary binding of GAM98 RNA, herein designated GAM RNA, to target binding sites on GAM98 target RNA, herein designated GAM TARGET RNA, such as BIND-

ING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM98 target RNA, herein designated GAM TARGET RNA, into GAM98 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1261] It is appreciated that GAM98 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM98 target genes. The mRNA of each one of this plurality of GAM98 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM98 RNA, herein designated GAM RNA, and which when bound by GAM98 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM98 target proteins.

[1262] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM98 gene, herein designated GAM GENE, on one or more GAM98 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site

has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1263] It is yet further appreciated that specific functions, and accordingly utilities, of GAM98 correlate with, and may be deduced from, the identity of the target genes which GAM98 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1264] Nucleotide sequences of the GAM98 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM98 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM98 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM98 are further described hereinbelow with reference to Table Table1.

[1265] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM98 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1266]

[1267] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 99 (GAM99) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1268] GAM99 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM99 was detected is described hereinabove with reference to Figs. 2-8.

[1269] GAM99 gene, herein designated GAM GENE, and GAM99 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1270] GAM99 gene, herein designated GAM GENE, encodes a GAM99 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM99 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM99 precursor RNA is designated SEQ ID:1293, and is provided hereinbelow with reference to the sequence listing part.

[1271] GAM99 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM99 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1272] An enzyme complex designated DICER COMPLEX, dices the GAM99 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM99 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 35%) nucleotide sequence of GAM99 RNA is designated SEQ ID:1294, and is provided hereinbelow with

reference to the sequence listing part.

[1273] GAM99 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM99 target RNA, herein designated GAM TARGET RNA. GAM99 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1274] GAM99 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM99 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM99 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM99 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM99 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1275] The complementary binding of GAM99 RNA, herein designated GAM RNA, to target binding sites on GAM99 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM99 target RNA, herein designated GAM TARGET RNA, into GAM99 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1276] It is appreciated that GAM99 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM99 target genes. The mRNA of each one of this plurality of GAM99 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM99 RNA, herein designated GAM RNA, and which when bound by GAM99 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM99 tar-

get proteins.

[1277] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM99 gene, herein designated GAM GENE, on one or more GAM99 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1278] It is yet further appreciated that specific functions, and accordingly utilities, of GAM99 correlate with, and may be deduced from, the identity of the target genes which GAM99 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1279] Nucleotide sequences of the GAM99 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM99 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM99 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM99 are further described hereinbelow with reference to Table Table1.

[1280] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM99 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1281]

[1282] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 100 (GAM100) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1283] GAM100 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM100 was detected is described hereinabove with reference to Figs. 2-8.

[1284] GAM100 gene, herein designated GAM GENE, and GAM100 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1285] GAM100 gene, herein designated GAM GENE, encodes a GAM100 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM100 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM100 precursor RNA is designated SEQ ID:1303, and is provided hereinbelow with reference to the sequence listing part.

[1286] GAM100 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM100 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1287] An enzyme complex designated DICER COMPLEX, dices

the GAM100 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM100 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM100 RNA is designated SEQ ID:1304, and is provided hereinbelow with reference to the sequence listing part.

[1288] GAM100 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM100 target RNA, herein designated GAM TARGET RNA.

GAM100 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1289] GAM100 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM100 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM100

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM100 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM100 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1290] The complementary binding of GAM100 RNA, herein designated GAM RNA, to target binding sites on GAM100 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM100 target RNA, herein designated GAM TARGET RNA, into GAM100 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1291] It is appreciated that GAM100 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM100 target genes. The mRNA of each one of this plurality of GAM100 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM100 RNA, herein designated GAM RNA, and which when bound by GAM100 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM100 target proteins.

[1292] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM100 gene, herein designated GAM GENE, on one or more GAM100 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1293] It is yet further appreciated that specific functions, and accordingly utilities, of GAM100 correlate with, and may be deduced from, the identity of the target genes which GAM100 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1294] Nucleotide sequences of the GAM100 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM100 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM100 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM100 are further described hereinbelow with reference to Table Table1.

[1295] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM100 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1296]

[1297] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 101 (GAM101) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1298] GAM101 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM101 was detected is described hereinabove with reference to Figs. 2-8.

[1299] GAM101 gene, herein designated GAM GENE, and GAM101 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1300] GAM101 gene, herein designated GAM GENE, encodes a GAM101 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM101 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM101 precursor RNA is designated SEQ ID:1315, and is provided hereinbelow with reference to the sequence listing part.

[1301] GAM101 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM101 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1302] An enzyme complex designated DICER COMPLEX, dices the GAM101 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM101 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM101 RNA is designated SEQ ID:1316, and is provided hereinbelow with reference to the sequence listing part.

[1303] GAM101 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM101 target RNA, herein designated GAM TARGET RNA. GAM101 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1304] GAM101 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM101 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM101 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM101 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM101 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[1305] The complementary binding of GAM101 RNA, herein designated GAM RNA, to target binding sites on GAM101 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM101 target RNA, herein designated GAM TARGET RNA, into GAM101 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1306] It is appreciated that GAM101 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM101 target genes. The mRNA of each one of this plurality of GAM101 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM101 RNA, herein designated GAM RNA, and which when bound by GAM101 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM101 target proteins.

[1307] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM101 gene, herein designated GAM GENE, on one or

more GAM101 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1308] It is yet further appreciated that specific functions, and accordingly utilities, of GAM101 correlate with, and may be deduced from, the identity of the target genes which GAM101 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1309] Nucleotide sequences of the GAM101 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM101 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM101 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM101 are further de-

scribed hereinbelow with reference to Table Table1.

[1310] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM101 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1311]

[1312] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 102 (GAM102) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1313] GAM102 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM102 was detected is described hereinabove with reference to Figs. 2-8.

[1314] GAM102 gene, herein designated GAM GENE, and GAM102 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1315] GAM102 gene, herein designated GAM GENE, encodes a GAM102 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM102 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM102 precursor RNA is designated SEQ ID:1330, and is provided hereinbelow with reference to the sequence listing part.

[1316] GAM102 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM102 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1317] An enzyme complex designated DICER COMPLEX, dices the GAM102 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM102 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM102 RNA is designated SEQ ID:1316, and is provided hereinbelow with reference to the sequence listing part.

[1318] GAM102 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM102 target RNA, herein designated GAM TARGET RNA. GAM102 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1319] GAM102 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM102 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM102 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM102 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM102 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1320] The complementary binding of GAM102 RNA, herein designated GAM RNA, to target binding sites on GAM102 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM102 target RNA, herein designated GAM TARGET RNA, into GAM102 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1321] It is appreciated that GAM102 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM102 target genes. The mRNA of each one of this plurality of GAM102 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM102 RNA, herein designated GAM RNA, and which when bound by GAM102 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM102 target proteins.

[1322] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM102 gene, herein designated GAM GENE, on one or more GAM102 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1323] It is yet further appreciated that specific functions, and accordingly utilities, of GAM102 correlate with, and may

be deduced from, the identity of the target genes which GAM102 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1324] Nucleotide sequences of the GAM102 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM102 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM102 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM102 are further described hereinbelow with reference to Table Table1.

[1325] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM102 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1326]

[1327] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 103 (GAM103) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1328] GAM103 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM103 was detected is described hereinabove with reference to Figs. 2-8.

[1329] GAM103 gene, herein designated GAM GENE, and GAM103 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1330] GAM103 gene, herein designated GAM GENE, encodes a GAM103 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM103 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM103 precursor RNA is designated SEQ ID:1336, and is provided hereinbelow with reference to the sequence listing part.

[1331] GAM103 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM103 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1332] An enzyme complex designated DICER COMPLEX, dices the GAM103 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM103 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM103 RNA is designated SEQ ID:1337, and is provided hereinbelow with reference to the sequence listing part.

[1333] GAM103 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM103 target RNA, herein designated GAM TARGET RNA. GAM103 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1334] GAM103 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM103 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM103 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM103 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM103 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1335] The complementary binding of GAM103 RNA, herein designated GAM RNA, to target binding sites on GAM103 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM103 target RNA, herein designated GAM TARGET RNA, into GAM103 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1336] It is appreciated that GAM103 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM103 target genes. The mRNA of each one of this plurality of GAM103 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM103 RNA, herein designated GAM RNA, and which when bound by GAM103 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM103 target proteins.

[1337] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM103 gene, herein designated GAM GENE, on one or more GAM103 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1338] It is yet further appreciated that specific functions, and accordingly utilities, of GAM103 correlate with, and may be deduced from, the identity of the target genes which GAM103 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1339] Nucleotide sequences of the GAM103 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM103 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM103 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM103 are further described hereinbelow with reference to Table Table1.

[1340] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM103 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1341]

[1342] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 104 (GAM104) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1343] GAM104 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM104 was detected is described hereinabove with reference to Figs. 2-8.

[1344] GAM104 gene, herein designated GAM GENE, and GAM104 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1345] GAM104 gene, herein designated GAM GENE, encodes a GAM104 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM104 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM104 precursor RNA is desig-

nated SEQ ID:1347, and is provided hereinbelow with reference to the sequence listing part.

[1346] GAM104 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM104 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1347] An enzyme complex designated DICER COMPLEX, dices the GAM104 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM104 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 31%) nucleotide sequence of GAM104 RNA is designated SEQ ID:1348, and is provided hereinbelow with reference to the sequence listing part.

[1348] GAM104 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM104 target RNA, herein designated GAM TARGET RNA. GAM104 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1349] GAM104 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM104 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM104 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM104 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM104

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1350] The complementary binding of GAM104 RNA, herein designated GAM RNA, to target binding sites on GAM104 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM104 target RNA, herein designated GAM TARGET RNA, into GAM104 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1351] It is appreciated that GAM104 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM104 target genes. The mRNA of each one of this plurality of GAM104 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM104 RNA, herein designated GAM RNA, and which when bound by GAM104 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM104 target proteins.

[1352] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM104 gene, herein designated GAM GENE, on one or more GAM104 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1353] It is yet further appreciated that specific functions, and accordingly utilities, of GAM104 correlate with, and may be deduced from, the identity of the target genes which GAM104 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1354] Nucleotide sequences of the GAM104 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM104 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM104 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM104 are further described hereinbelow with reference to Table Table1.

[1355] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM104 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1356]

[1357] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 105 (GAM105) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1358] GAM105 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM105 was detected is described hereinabove with reference to Figs. 2-8.

[1359] GAM105 gene, herein designated GAM GENE, and

GAM105 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1360] GAM105 gene, herein designated GAM GENE, encodes a GAM105 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM105 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM105 precursor RNA is designated SEQ ID:1391, and is provided hereinbelow with reference to the sequence listing part.

[1361] GAM105 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM105 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1362] An enzyme complex designated DICER COMPLEX, dices the GAM105 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM105 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM105 RNA is designated SEQ ID:1392, and is provided hereinbelow with reference to the sequence listing part.

[1363] GAM105 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM105 target RNA, herein designated GAM TARGET RNA. GAM105 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1364] GAM105 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM105 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM105 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM105 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM105 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1365] The complementary binding of GAM105 RNA, herein designated GAM RNA, to target binding sites on GAM105 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM105 target RNA, herein designated GAM TARGET RNA, into GAM105 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1366] It is appreciated that GAM105 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM105 target genes. The mRNA of each one of this plurality of GAM105 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM105 RNA, herein designated GAM RNA, and which when bound by GAM105 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM105 target proteins.

[1367] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM105 gene, herein designated GAM GENE, on one or more GAM105 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1368] It is yet further appreciated that specific functions, and accordingly utilities, of GAM105 correlate with, and may be deduced from, the identity of the target genes which GAM105 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1369] Nucleotide sequences of the GAM105 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM105 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM105 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM105 are further described hereinbelow with reference to Table Table1.

[1370] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM105 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1371]

[1372] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 106 (GAM106) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1373] GAM106 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM106 was detected is described hereinabove with reference to Figs. 2-8.

[1374] GAM106 gene, herein designated GAM GENE, and GAM106 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1375] GAM106 gene, herein designated GAM GENE, encodes a GAM106 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM106 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM106 precursor RNA is designated SEQ ID:1403, and is provided hereinbelow with reference to the sequence listing part.

[1376] GAM106 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM106 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1377] An enzyme complex designated DICER COMPLEX, dices the GAM106 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM106 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM106 RNA is designated SEQ ID:1404, and is provided hereinbelow with reference to the sequence listing part.

[1378] GAM106 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM106 target RNA, herein designated GAM TARGET RNA. GAM106 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1379] GAM106 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM106 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM106 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM106 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM106 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1380] The complementary binding of GAM106 RNA, herein designated GAM RNA, to target binding sites on GAM106 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM106 target RNA, herein designated GAM TARGET RNA, into GAM106 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1381] It is appreciated that GAM106 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM106 target genes. The mRNA of each one of this plurality of GAM106 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM106 RNA, herein designated GAM RNA, and which when bound by GAM106 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM106 target proteins.

[1382] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM106 gene, herein designated GAM GENE, on one or more GAM106 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1383] It is yet further appreciated that specific functions, and accordingly utilities, of GAM106 correlate with, and may be deduced from, the identity of the target genes which GAM106 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1384] Nucleotide sequences of the GAM106 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM106 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM106 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM106 are further described hereinbelow with reference to Table Table1.

[1385] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM106 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1386]

[1387] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 107 (GAM107) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1388] GAM107 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM107 was detected is described hereinabove with reference to Figs. 2-8.

[1389] GAM107 gene, herein designated GAM GENE, and GAM107 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1390] GAM107 gene, herein designated GAM GENE, encodes a GAM107 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM107 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM107 precursor RNA is designated SEQ ID:1411, and is provided hereinbelow with reference to the sequence listing part.

[1391] GAM107 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM107 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1392] An enzyme complex designated DICER COMPLEX, dices the GAM107 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM107 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 34%) nucleotide sequence of GAM107 RNA is designated SEQ ID:1412, and is provided hereinbelow with reference to the sequence listing part.

[1393] GAM107 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM107 target RNA, herein designated GAM TARGET RNA. GAM107 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1394] GAM107 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM107 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM107 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM107 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM107 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1395] The complementary binding of GAM107 RNA, herein designated GAM RNA, to target binding sites on GAM107 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM107 target RNA, herein designated GAM TARGET RNA, into GAM107 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1396] It is appreciated that GAM107 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM107 target genes. The mRNA of each one of this plurality of GAM107 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM107 RNA,

herein designated GAM RNA, and which when bound by GAM107 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM107 target proteins.

[1397] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM107 gene, herein designated GAM GENE, on one or more GAM107 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1398] It is yet further appreciated that specific functions, and accordingly utilities, of GAM107 correlate with, and may be deduced from, the identity of the target genes which

GAM107 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1399] Nucleotide sequences of the GAM107 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM107 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM107 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM107 are further described hereinbelow with reference to Table Table1.

[1400] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM107 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1401]

[1402] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 108 (GAM108) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1403] GAM108 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM108 was detected is described hereinabove with reference to Figs. 2-8.

[1404] GAM108 gene, herein designated GAM GENE, and GAM108 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1405] GAM108 gene, herein designated GAM GENE, encodes a GAM108 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM108 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM108 precursor RNA is designated SEQ ID:1428, and is provided hereinbelow with reference to the sequence listing part.

[1406] GAM108 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM108 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[1407] An enzyme complex designated DICER COMPLEX, dices the GAM108 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM108 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM108 RNA is designated SEQ ID:1429, and is provided hereinbelow with reference to the sequence listing part.

[1408] GAM108 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM108 target RNA, herein designated GAM TARGET RNA.
GAM108 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1409] GAM108 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM108 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM108 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM108 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM108 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1410] The complementary binding of GAM108 RNA, herein designated GAM RNA, to target binding sites on GAM108 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM108 target RNA, herein desig-

nated GAM TARGET RNA, into GAM108 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1411] It is appreciated that GAM108 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM108 target genes. The mRNA of each one of this plurality of GAM108 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM108 RNA, herein designated GAM RNA, and which when bound by GAM108 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM108 target proteins.

[1412] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM108 gene, herein designated GAM GENE, on one or more GAM108 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1413] It is yet further appreciated that specific functions, and accordingly utilities, of GAM108 correlate with, and may be deduced from, the identity of the target genes which GAM108 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1414] Nucleotide sequences of the GAM108 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM108 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM108 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM108 are further described hereinbelow with reference to Table Table1.

[1415] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM108 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[1416]

[1417] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 109 (GAM109) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1418] GAM109 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM109 was detected is described hereinabove with reference to Figs. 2-8.

[1419] GAM109 gene, herein designated GAM GENE, and GAM109 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1420] GAM109 gene, herein designated GAM GENE, encodes a GAM109 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM109 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM109 precursor RNA is designated SEQ ID:1443, and is provided hereinbelow with ref-

erence to the sequence listing part.

[1421] GAM109 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM109 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1422] An enzyme complex designated DICER COMPLEX, dices the GAM109 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM109 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM109 RNA is designated SEQ ID:1444, and is provided hereinbelow with reference to the sequence listing part.

[1423] GAM109 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM109 target RNA, herein designated GAM TARGET RNA.

GAM109 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1424] GAM109 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM109 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM109 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM109 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM109 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1425] The complementary binding of GAM109 RNA, herein designated GAM RNA, to target binding sites on GAM109 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM109 target RNA, herein designated GAM TARGET RNA, into GAM109 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1426] It is appreciated that GAM109 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM109 target genes. The mRNA of each one of this plurality of GAM109 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM109 RNA, herein designated GAM RNA, and which when bound by GAM109 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM109 target proteins.

[1427] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM109 gene, herein designated GAM GENE, on one or more GAM109 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1428] It is yet further appreciated that specific functions, and accordingly utilities, of GAM109 correlate with, and may be deduced from, the identity of the target genes which GAM109 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1429] Nucleotide sequences of the GAM109 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM109 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM109 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM109 are further described hereinbelow with reference to Table Table1.

[1430] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM109 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1431]

[1432] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 110 (GAM110) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1433] GAM110 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM110 was detected is described hereinabove with reference to Figs. 2-8.

[1434] GAM110 gene, herein designated GAM GENE, and GAM110 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[1435] GAM110 gene, herein designated GAM GENE, encodes a GAM110 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM110 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM110 precursor RNA is designated SEQ ID:1462, and is provided hereinbelow with reference to the sequence listing part.

[1436] GAM110 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM110 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1437] An enzyme complex designated DICER COMPLEX, dices the GAM110 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM110 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM110 RNA is designated SEQ ID:1463, and is provided hereinbelow with reference to the sequence listing part.

[1438] GAM110 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM110 target RNA, herein designated GAM TARGET RNA. GAM110 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1439] GAM110 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM110 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM110 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM110 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM110 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1440] The complementary binding of GAM110 RNA, herein designated GAM RNA, to target binding sites on GAM110 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM110 target RNA, herein designated GAM TARGET RNA, into GAM110 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1441] It is appreciated that GAM110 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM110 target genes. The mRNA of each one of this plurality of GAM110 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM110 RNA, herein designated GAM RNA, and which when bound by GAM110 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM110 target proteins.

[1442] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM110 gene, herein designated GAM GENE, on one or more GAM110 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[1443] It is yet further appreciated that specific functions, and accordingly utilities, of GAM110 correlate with, and may be deduced from, the identity of the target genes which GAM110 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1444] Nucleotide sequences of the GAM110 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM110 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM110 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM110 are further described hereinbelow with reference to Table Table1.

[1445] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM110 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1446]

[1447] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

111 (GAM111) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1448] GAM111 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM111 was detected is described hereinabove with reference to Figs. 2-8.

[1449] GAM111 gene, herein designated GAM GENE, and GAM111 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1450] GAM111 gene, herein designated GAM GENE, encodes a GAM111 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM111 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM111 precursor RNA is designated SEQ ID:1487, and is provided hereinbelow with reference to the sequence listing part.

[1451] GAM111 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM111 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1452] An enzyme complex designated DICER COMPLEX, dices the GAM111 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM111 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM111 RNA is designated SEQ ID:1488, and is provided hereinbelow with reference to the sequence listing part.

[1453] GAM111 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM111 target RNA, herein designated GAM TARGET RNA. GAM111 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1454] GAM111 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM111 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM111 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM111 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM111 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1455] The complementary binding of GAM111 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM111 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM111 target RNA, herein designated GAM TARGET RNA, into GAM111 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1456] It is appreciated that GAM111 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM111 target genes. The mRNA of each one of this plurality of GAM111 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM111 RNA, herein designated GAM RNA, and which when bound by GAM111 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM111 target proteins.

[1457] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM111 gene, herein designated GAM GENE, on one or more GAM111 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1458] It is yet further appreciated that specific functions, and accordingly utilities, of GAM111 correlate with, and may be deduced from, the identity of the target genes which GAM111 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1459] Nucleotide sequences of the GAM111 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM111 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM111 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM111 are further described hereinbelow with reference to Table Table1.

[1460] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM111 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1461]

[1462] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 112 (GAM112) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1463] GAM112 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM112 was detected is described hereinabove with reference to Figs. 2-8.

[1464] GAM112 gene, herein designated GAM GENE, and GAM112 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1465] GAM112 gene, herein designated GAM GENE, encodes a GAM112 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM112 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM112 precursor RNA is designated SEQ ID:1494, and is provided hereinbelow with reference to the sequence listing part.

[1466] GAM112 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM112 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1467] An enzyme complex designated DICER COMPLEX, dices the GAM112 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM112 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 59%) nucleotide sequence of GAM112 RNA is designated SEQ ID:1495, and is provided hereinbelow with reference to the sequence listing part.

[1468] GAM112 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM112 target RNA, herein designated GAM TARGET RNA. GAM112 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1469] GAM112 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM112 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM112 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM112 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM112 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1470] The complementary binding of GAM112 RNA, herein designated GAM RNA, to target binding sites on GAM112 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM112 target RNA, herein designated GAM TARGET RNA, into GAM112 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1471] It is appreciated that GAM112 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM112 target genes. The mRNA of each one of this plurality of GAM112 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM112 RNA, herein designated GAM RNA, and which when bound by

GAM112 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM112 target proteins.

[1472] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM112 gene, herein designated GAM GENE, on one or more GAM112 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1473] It is yet further appreciated that specific functions, and accordingly utilities, of GAM112 correlate with, and may be deduced from, the identity of the target genes which GAM112 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[1474] Nucleotide sequences of the GAM112 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM112 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM112 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM112 are further described hereinbelow with reference to Table Table1.

[1475] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM112 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1476]

[1477] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 113 (GAM113) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1478] GAM113 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM113 was detected is described hereinabove with reference to Figs. 2-8.

[1479] GAM113 gene, herein designated GAM GENE, and GAM113 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1480] GAM113 gene, herein designated GAM GENE, encodes a GAM113 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM113 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM113 precursor RNA is designated SEQ ID:1515, and is provided hereinbelow with reference to the sequence listing part.

[1481] GAM113 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM113 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[1482] An enzyme complex designated DICER COMPLEX, dices the GAM113 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM113 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM113 RNA is designated SEQ ID:1516, and is provided hereinbelow with reference to the sequence listing part.

[1483] GAM113 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM113 target RNA, herein designated GAM TARGET RNA. GAM113 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1484] GAM113 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM113 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM113 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM113 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM113 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1485] The complementary binding of GAM113 RNA, herein designated GAM RNA, to target binding sites on GAM113 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM113 target RNA, herein designated GAM TARGET RNA, into GAM113 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1486] It is appreciated that GAM113 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM113 target genes. The mRNA of each one of this plurality of GAM113 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM113 RNA, herein designated GAM RNA, and which when bound by GAM113 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM113 target proteins.

[1487] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM113 gene, herein designated GAM GENE, on one or more GAM113 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1488] It is yet further appreciated that specific functions, and accordingly utilities, of GAM113 correlate with, and may be deduced from, the identity of the target genes which GAM113 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1489] Nucleotide sequences of the GAM113 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM113 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM113 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM113 are further described hereinbelow with reference to Table Table1.

[1490] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM113 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1491]

[1492] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 114 (GAM114) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1493] GAM114 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM114 was detected is described hereinabove with reference to Figs. 2-8.

[1494] GAM114 gene, herein designated GAM GENE, and GAM114 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1495] GAM114 gene, herein designated GAM GENE, encodes a GAM114 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM114 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM114 precursor RNA is designated SEQ ID:1525, and is provided hereinbelow with reference to the sequence listing part.

[1496] GAM114 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM114 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1497] An enzyme complex designated DICER COMPLEX, dices the GAM114 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM114 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM114 RNA is designated SEQ ID:1526, and is provided hereinbelow with reference to the sequence listing part.

[1498] GAM114 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM114

target RNA, herein designated GAM TARGET RNA.

GAM114 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1499] GAM114 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM114 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM114 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM114 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM114 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1500] The complementary binding of GAM114 RNA, herein designated GAM RNA, to target binding sites on GAM114 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM114 target RNA, herein designated GAM TARGET RNA, into GAM114 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1501] It is appreciated that GAM114 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM114 target genes. The mRNA of each one of this plurality of GAM114 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM114 RNA, herein designated GAM RNA, and which when bound by GAM114 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM114 target proteins.

[1502] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM114 gene, herein designated GAM GENE, on one or more GAM114 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1503] It is yet further appreciated that specific functions, and accordingly utilities, of GAM114 correlate with, and may be deduced from, the identity of the target genes which GAM114 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1504] Nucleotide sequences of the GAM114 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM114 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM114 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM114 are further described hereinbelow with reference to Table Table1.

[1505] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM114 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1506]

[1507] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 115 (GAM115) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1508] GAM115 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM115 was detected is described hereinabove with reference to Figs. 2-8.

[1509] GAM115 gene, herein designated GAM GENE, and GAM115 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1510] GAM115 gene, herein designated GAM GENE, encodes a GAM115 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM115 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM115 precursor RNA is designated SEQ ID:1575, and is provided hereinbelow with reference to the sequence listing part.

[1511] GAM115 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM115 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1512] An enzyme complex designated DICER COMPLEX, dices the GAM115 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM115 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM115 RNA is designated SEQ ID:1576, and is provided hereinbelow with reference to the sequence listing part.

[1513] GAM115 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM115 target RNA, herein designated GAM TARGET RNA. GAM115 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1514] GAM115 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM115 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM115 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM115 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM115 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1515] The complementary binding of GAM115 RNA, herein designated GAM RNA, to target binding sites on GAM115 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM115 target RNA, herein designated GAM TARGET RNA, into GAM115 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1516] It is appreciated that GAM115 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM115 target genes. The mRNA of each one of this plu-

rality of GAM115 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM115 RNA, herein designated GAM RNA, and which when bound by GAM115 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM115 target proteins.

[1517] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM115 gene, herein designated GAM GENE, on one or more GAM115 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1518] It is yet further appreciated that specific functions, and accordingly utilities, of GAM115 correlate with, and may be deduced from, the identity of the target genes which GAM115 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1519] Nucleotide sequences of the GAM115 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM115 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM115 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM115 are further described hereinbelow with reference to Table Table1.

[1520] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM115 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1521]

[1522] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 116 (GAM116) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[1523] GAM116 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM116 was detected is described hereinabove with reference to Figs. 2-8.

[1524] GAM116 gene, herein designated GAM GENE, and GAM116 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1525] GAM116 gene, herein designated GAM GENE, encodes a GAM116 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM116 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM116 precursor RNA is designated SEQ ID:1589, and is provided hereinbelow with reference to the sequence listing part.

[1526] GAM116 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM116 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1527] An enzyme complex designated DICER COMPLEX, dices the GAM116 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM116 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM116 RNA is designated SEQ ID:1590, and is provided hereinbelow with reference to the sequence listing part.

[1528] GAM116 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM116 target RNA, herein designated GAM TARGET RNA.

GAM116 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[1529] GAM116 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM116 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM116 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM116 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM116 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1530] The complementary binding of GAM116 RNA, herein designated GAM RNA, to target binding sites on GAM116 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM116 target RNA, herein designated GAM TARGET RNA, into GAM116 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1531] It is appreciated that GAM116 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM116 target genes. The mRNA of each one of this plurality of GAM116 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM116 RNA, herein designated GAM RNA, and which when bound by GAM116 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM116 target proteins.

[1532] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM116 gene, herein designated GAM GENE, on one or more GAM116 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1533] It is yet further appreciated that specific functions, and accordingly utilities, of GAM116 correlate with, and may be deduced from, the identity of the target genes which GAM116 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1534] Nucleotide sequences of the GAM116 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM116 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM116 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM116 are further described hereinbelow with reference to Table Table1.

[1535] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM116 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1536]

[1537] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 117 (GAM117) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1538] GAM117 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM117 was detected is described hereinabove with reference to Figs. 2-8.

[1539] GAM117 gene, herein designated GAM GENE, and GAM117 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1540] GAM117 gene, herein designated GAM GENE, encodes a GAM117 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM117 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM117 precursor RNA is designated SEQ ID:1616, and is provided hereinbelow with reference to the sequence listing part.

[1541] GAM117 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM117 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1542] An enzyme complex designated DICER COMPLEX, dices the GAM117 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM117 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of

GAM117 RNA is designated SEQ ID:1617, and is provided hereinbelow with reference to the sequence listing part.

[1543] GAM117 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM117 target RNA, herein designated GAM TARGET RNA. GAM117 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1544] GAM117 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM117 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM117 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM117 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM117 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1545] The complementary binding of GAM117 RNA, herein designated GAM RNA, to target binding sites on GAM117 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM117 target RNA, herein designated GAM TARGET RNA, into GAM117 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1546] It is appreciated that GAM117 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM117 target genes. The mRNA of each one of this plurality of GAM117 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM117 RNA, herein designated GAM RNA, and which when bound by GAM117 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM117 target proteins.

[1547] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM117 gene, herein designated GAM GENE, on one or more GAM117 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1548] It is yet further appreciated that specific functions, and accordingly utilities, of GAM117 correlate with, and may be deduced from, the identity of the target genes which GAM117 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1549] Nucleotide sequences of the GAM117 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM117 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM117 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM117 are further described hereinbelow with reference to Table Table1.

[1550] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM117 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1551]

[1552] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 118 (GAM118) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1553] GAM118 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM118 was detected is described hereinabove

with reference to Figs. 2-8.

[1554] GAM118 gene, herein designated GAM GENE, and GAM118 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1555] GAM118 gene, herein designated GAM GENE, encodes a GAM118 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM118 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM118 precursor RNA is designated SEQ ID:1655, and is provided hereinbelow with reference to the sequence listing part.

[1556] GAM118 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM118 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1557] An enzyme complex designated DICER COMPLEX, dices the GAM118 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM118 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM118 RNA is designated SEQ ID:1617, and is provided hereinbelow with reference to the sequence listing part.

[1558] GAM118 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM118 target RNA, herein designated GAM TARGET RNA. GAM118 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1559] GAM118 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM118 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM118 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM118 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM118 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1560] The complementary binding of GAM118 RNA, herein designated GAM RNA, to target binding sites on GAM118 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM118 target RNA, herein designated GAM TARGET RNA, into GAM118 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[1561] It is appreciated that GAM118 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM118 target genes. The mRNA of each one of this plurality of GAM118 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM118 RNA, herein designated GAM RNA, and which when bound by GAM118 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM118 target proteins.

[1562] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM118 gene, herein designated GAM GENE, on one or more GAM118 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1563] It is yet further appreciated that specific functions, and accordingly utilities, of GAM118 correlate with, and may be deduced from, the identity of the target genes which GAM118 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1564] Nucleotide sequences of the GAM118 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM118 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM118 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM118 are further described hereinbelow with reference to Table Table1.

[1565] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM118 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1566]

[1567] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 119 (GAM119) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1568] GAM119 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM119 was detected is described hereinabove with reference to Figs. 2-8.

[1569] GAM119 gene, herein designated GAM GENE, and GAM119 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1570] GAM119 gene, herein designated GAM GENE, encodes a GAM119 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM119 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM119 precursor RNA is designated SEQ ID:1667, and is provided hereinbelow with reference to the sequence listing part.

[1571] GAM119 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM119 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1572] An enzyme complex designated DICER COMPLEX, dices the GAM119 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM119 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 17%) nucleotide sequence of GAM119 RNA is designated SEQ ID:1668, and is provided hereinbelow with reference to the sequence listing part.

[1573] GAM119 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM119 target RNA, herein designated GAM TARGET RNA.

GAM119 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1574] GAM119 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM119 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM119 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM119 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM119 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1575] The complementary binding of GAM119 RNA, herein designated GAM RNA, to target binding sites on GAM119 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM119 target RNA, herein designated GAM TARGET RNA, into GAM119 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1576] It is appreciated that GAM119 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM119 target genes. The mRNA of each one of this plurality of GAM119 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM119 RNA, herein designated GAM RNA, and which when bound by GAM119 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM119 target proteins.

[1577] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM119 gene, herein designated GAM GENE, on one or more GAM119 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1578] It is yet further appreciated that specific functions, and accordingly utilities, of GAM119 correlate with, and may be deduced from, the identity of the target genes which GAM119 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1579] Nucleotide sequences of the GAM119 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM119 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM119 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM119 are further described hereinbelow with reference to Table Table1.

[1580] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM119 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1581]

[1582] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 120 (GAM120) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1583] GAM120 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM120 was detected is described hereinabove with reference to Figs. 2-8.

[1584] GAM120 gene, herein designated GAM GENE, and GAM120 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1585] GAM120 gene, herein designated GAM GENE, encodes a

GAM120 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM120 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM120 precursor RNA is designated SEQ ID:1684, and is provided hereinbelow with reference to the sequence listing part.

[1586] GAM120 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM120 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1587] An enzyme complex designated DICER COMPLEX, dices the GAM120 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM120 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM120 RNA is designated SEQ ID:1685, and is provided hereinbelow with reference to the sequence listing part.

[1588] GAM120 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM120 target RNA, herein designated GAM TARGET RNA. GAM120 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1589] GAM120 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM120 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM120 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM120 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM120 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1590] The complementary binding of GAM120 RNA, herein designated GAM RNA, to target binding sites on GAM120 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM120 target RNA, herein designated GAM TARGET RNA, into GAM120 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1591] It is appreciated that GAM120 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM120 target genes. The mRNA of each one of this plurality of GAM120 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM120 RNA, herein designated GAM RNA, and which when bound by GAM120 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM120 target proteins.

[1592] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM120 gene, herein designated GAM GENE, on one or more GAM120 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1593] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM120 correlate with, and may be deduced from, the identity of the target genes which GAM120 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1594] Nucleotide sequences of the GAM120 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM120 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM120 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM120 are further described hereinbelow with reference to Table Table1.

[1595] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM120 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1596]

[1597] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 121 (GAM121) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[1598] GAM121 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM121 was detected is described hereinabove with reference to Figs. 2-8.

[1599] GAM121 gene, herein designated GAM GENE, and GAM121 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1600] GAM121 gene, herein designated GAM GENE, encodes a GAM121 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM121 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM121 precursor RNA is designated SEQ ID:1699, and is provided hereinbelow with reference to the sequence listing part.

[1601] GAM121 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM121 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1602] An enzyme complex designated DICER COMPLEX, dices the GAM121 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM121 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM121 RNA is designated SEQ ID:1700, and is provided hereinbelow with reference to the sequence listing part.

[1603] GAM121 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM121 target RNA, herein designated GAM TARGET RNA. GAM121 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1604] GAM121 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM121 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM121 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM121 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM121 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1605] The complementary binding of GAM121 RNA, herein designated GAM RNA, to target binding sites on GAM121 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM121 target RNA, herein designated GAM TARGET RNA, into GAM121 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1606] It is appreciated that GAM121 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM121 target genes. The mRNA of each one of this plurality of GAM121 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM121 RNA, herein designated GAM RNA, and which when bound by GAM121 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM121 target proteins.

[1607] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM121 gene, herein designated GAM GENE, on one or more GAM121 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1608] It is yet further appreciated that specific functions, and accordingly utilities, of GAM121 correlate with, and may be deduced from, the identity of the target genes which GAM121 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1609] Nucleotide sequences of the GAM121 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM121 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM121 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM121 are further described hereinbelow with reference to Table Table1.

[1610] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM121 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1611]

[1612] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 122 (GAM122) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1613] GAM122 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM122 was detected is described hereinabove with reference to Figs. 2-8.

[1614] GAM122 gene, herein designated GAM GENE, and GAM122 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1615] GAM122 gene, herein designated GAM GENE, encodes a GAM122 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM122 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM122 precursor RNA is designated SEQ ID:1702, and is provided hereinbelow with reference to the sequence listing part.

[1616] GAM122 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM122 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1617] An enzyme complex designated DICER COMPLEX, dices the GAM122 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM122 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM122 RNA is designated SEQ ID:1703, and is provided

hereinbelow with reference to the sequence listing part.

[1618] GAM122 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM122 target RNA, herein designated GAM TARGET RNA.
GAM122 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1619] GAM122 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM122 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM122 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM122 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM122 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1620] The complementary binding of GAM122 RNA, herein designated GAM RNA, to target binding sites on GAM122 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM122 target RNA, herein designated GAM TARGET RNA, into GAM122 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1621] It is appreciated that GAM122 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM122 target genes. The mRNA of each one of this plurality of GAM122 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM122 RNA, herein designated GAM RNA, and which when bound by GAM122 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM122 tar-

get proteins.

[1622] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM122 gene, herein designated GAM GENE, on one or more GAM122 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1623] It is yet further appreciated that specific functions, and accordingly utilities, of GAM122 correlate with, and may be deduced from, the identity of the target genes which GAM122 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1624] Nucleotide sequences of the GAM122 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM122 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM122 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM122 are further described hereinbelow with reference to Table Table1.

[1625] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM122 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1626]

[1627] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 123 (GAM123) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1628] GAM123 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM123 was detected is described hereinabove with reference to Figs. 2-8.

[1629] GAM123 gene, herein designated GAM GENE, and GAM123 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1630] GAM123 gene, herein designated GAM GENE, encodes a GAM123 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM123 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM123 precursor RNA is designated SEQ ID:1722, and is provided hereinbelow with reference to the sequence listing part.

[1631] GAM123 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM123 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1632] An enzyme complex designated DICER COMPLEX, dices

the GAM123 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM123 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM123 RNA is designated SEQ ID:1723, and is provided hereinbelow with reference to the sequence listing part.

[1633] GAM123 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM123 target RNA, herein designated GAM TARGET RNA.
GAM123 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1634] GAM123 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM123 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM123

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM123 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM123 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1635] The complementary binding of GAM123 RNA, herein designated GAM RNA, to target binding sites on GAM123 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM123 target RNA, herein designated GAM TARGET RNA, into GAM123 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1636] It is appreciated that GAM123 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM123 target genes. The mRNA of each one of this plurality of GAM123 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM123 RNA, herein designated GAM RNA, and which when bound by GAM123 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM123 target proteins.

[1637] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM123 gene, herein designated GAM GENE, on one or more GAM123 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1638] It is yet further appreciated that specific functions, and accordingly utilities, of GAM123 correlate with, and may be deduced from, the identity of the target genes which GAM123 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1639] Nucleotide sequences of the GAM123 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM123 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM123 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM123 are further described hereinbelow with reference to Table Table1.

[1640] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM123 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1641]

[1642] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 124 (GAM124) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1643] GAM124 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM124 was detected is described hereinabove with reference to Figs. 2-8.

[1644] GAM124 gene, herein designated GAM GENE, and GAM124 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1645] GAM124 gene, herein designated GAM GENE, encodes a GAM124 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM124 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM124 precursor RNA is designated SEQ ID:1734, and is provided hereinbelow with reference to the sequence listing part.

[1646] GAM124 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM124 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1647] An enzyme complex designated DICER COMPLEX, dices the GAM124 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM124 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM124 RNA is designated SEQ ID:1735, and is provided hereinbelow with reference to the sequence listing part.

[1648] GAM124 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM124 target RNA, herein designated GAM TARGET RNA. GAM124 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1649] GAM124 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM124 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM124 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM124 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM124 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[1650] The complementary binding of GAM124 RNA, herein designated GAM RNA, to target binding sites on GAM124 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM124 target RNA, herein designated GAM TARGET RNA, into GAM124 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1651] It is appreciated that GAM124 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM124 target genes. The mRNA of each one of this plurality of GAM124 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM124 RNA, herein designated GAM RNA, and which when bound by GAM124 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM124 target proteins.

[1652] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM124 gene, herein designated GAM GENE, on one or

more GAM124 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1653] It is yet further appreciated that specific functions, and accordingly utilities, of GAM124 correlate with, and may be deduced from, the identity of the target genes which GAM124 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1654] Nucleotide sequences of the GAM124 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM124 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM124 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM124 are further de-

scribed hereinbelow with reference to Table Table1.

[1655] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM124 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1656]

[1657] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 125 (GAM125) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1658] GAM125 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM125 was detected is described hereinabove with reference to Figs. 2-8.

[1659] GAM125 gene, herein designated GAM GENE, and GAM125 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1660] GAM125 gene, herein designated GAM GENE, encodes a GAM125 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM125 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM125 precursor RNA is designated SEQ ID:1741, and is provided hereinbelow with reference to the sequence listing part.

[1661] GAM125 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM125 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1662] An enzyme complex designated DICER COMPLEX, dices the GAM125 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM125 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 21%) nucleotide sequence of GAM125 RNA is designated SEQ ID:1742, and is provided hereinbelow with reference to the sequence listing part.

[1663] GAM125 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM125 target RNA, herein designated GAM TARGET RNA. GAM125 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1664] GAM125 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM125 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM125 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM125 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM125 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1665] The complementary binding of GAM125 RNA, herein designated GAM RNA, to target binding sites on GAM125 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM125 target RNA, herein designated GAM TARGET RNA, into GAM125 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1666] It is appreciated that GAM125 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM125 target genes. The mRNA of each one of this plurality of GAM125 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM125 RNA, herein designated GAM RNA, and which when bound by GAM125 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM125 target proteins.

[1667] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM125 gene, herein designated GAM GENE, on one or more GAM125 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1668] It is yet further appreciated that specific functions, and accordingly utilities, of GAM125 correlate with, and may

be deduced from, the identity of the target genes which GAM125 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1669] Nucleotide sequences of the GAM125 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM125 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM125 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM125 are further described hereinbelow with reference to Table Table1.

[1670] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM125 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1671]

[1672] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 126 (GAM126) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1673] GAM126 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM126 was detected is described hereinabove with reference to Figs. 2-8.

[1674] GAM126 gene, herein designated GAM GENE, and GAM126 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1675] GAM126 gene, herein designated GAM GENE, encodes a GAM126 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM126 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM126 precursor RNA is designated SEQ ID:1770, and is provided hereinbelow with reference to the sequence listing part.

[1676] GAM126 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM126 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1677] An enzyme complex designated DICER COMPLEX, dices the GAM126 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM126 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM126 RNA is designated SEQ ID:1771, and is provided hereinbelow with reference to the sequence listing part.

[1678] GAM126 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM126 target RNA, herein designated GAM TARGET RNA. GAM126 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1679] GAM126 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM126 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM126 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM126 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM126 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1680] The complementary binding of GAM126 RNA, herein designated GAM RNA, to target binding sites on GAM126 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM126 target RNA, herein designated GAM TARGET RNA, into GAM126 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1681] It is appreciated that GAM126 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM126 target genes. The mRNA of each one of this plurality of GAM126 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM126 RNA, herein designated GAM RNA, and which when bound by GAM126 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM126 target proteins.

[1682] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM126 gene, herein designated GAM GENE, on one or more GAM126 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1683] It is yet further appreciated that specific functions, and accordingly utilities, of GAM126 correlate with, and may be deduced from, the identity of the target genes which GAM126 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1684] Nucleotide sequences of the GAM126 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM126 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM126 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM126 are further described hereinbelow with reference to Table Table1.

[1685] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM126 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1686]

[1687] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 127 (GAM127) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1688] GAM127 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM127 was detected is described hereinabove with reference to Figs. 2-8.

[1689] GAM127 gene, herein designated GAM GENE, and GAM127 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1690] GAM127 gene, herein designated GAM GENE, encodes a GAM127 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM127 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM127 precursor RNA is desig-

nated SEQ ID:1780, and is provided hereinbelow with reference to the sequence listing part.

[1691] GAM127 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM127 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1692] An enzyme complex designated DICER COMPLEX, dices the GAM127 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM127 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of GAM127 RNA is designated SEQ ID:1781, and is provided hereinbelow with reference to the sequence listing part.

[1693] GAM127 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM127 target RNA, herein designated GAM TARGET RNA. GAM127 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1694] GAM127 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM127 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM127 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM127 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM127

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1695] The complementary binding of GAM127 RNA, herein designated GAM RNA, to target binding sites on GAM127 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM127 target RNA, herein designated GAM TARGET RNA, into GAM127 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1696] It is appreciated that GAM127 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM127 target genes. The mRNA of each one of this plurality of GAM127 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM127 RNA, herein designated GAM RNA, and which when bound by GAM127 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM127 target proteins.

[1697] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM127 gene, herein designated GAM GENE, on one or more GAM127 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1698] It is yet further appreciated that specific functions, and accordingly utilities, of GAM127 correlate with, and may be deduced from, the identity of the target genes which GAM127 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1699] Nucleotide sequences of the GAM127 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM127 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM127 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM127 are further described hereinbelow with reference to Table Table1.

[1700] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM127 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1701]

[1702] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 128 (GAM128) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1703] GAM128 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM128 was detected is described hereinabove with reference to Figs. 2-8.

[1704] GAM128 gene, herein designated GAM GENE, and

GAM128 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1705] GAM128 gene, herein designated GAM GENE, encodes a GAM128 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM128 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM128 precursor RNA is designated SEQ ID:1786, and is provided hereinbelow with reference to the sequence listing part.

[1706] GAM128 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM128 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1707] An enzyme complex designated DICER COMPLEX, dices the GAM128 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM128 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 19%) nucleotide sequence of GAM128 RNA is designated SEQ ID:1787, and is provided hereinbelow with reference to the sequence listing part.

[1708] GAM128 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM128 target RNA, herein designated GAM TARGET RNA. GAM128 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1709] GAM128 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM128 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM128 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM128 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM128 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1710] The complementary binding of GAM128 RNA, herein designated GAM RNA, to target binding sites on GAM128 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM128 target RNA, herein designated GAM TARGET RNA, into GAM128 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1711] It is appreciated that GAM128 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM128 target genes. The mRNA of each one of this plurality of GAM128 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM128 RNA, herein designated GAM RNA, and which when bound by GAM128 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM128 target proteins.

[1712] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM128 gene, herein designated GAM GENE, on one or more GAM128 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1713] It is yet further appreciated that specific functions, and accordingly utilities, of GAM128 correlate with, and may be deduced from, the identity of the target genes which GAM128 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1714] Nucleotide sequences of the GAM128 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM128 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM128 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM128 are further described hereinbelow with reference to Table Table1.

[1715] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM128 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1716]

[1717] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 129 (GAM129) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1718] GAM129 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM129 was detected is described hereinabove with reference to Figs. 2-8.

[1719] GAM129 gene, herein designated GAM GENE, and GAM129 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1720] GAM129 gene, herein designated GAM GENE, encodes a GAM129 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM129 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM129 precursor RNA is designated SEQ ID:1803, and is provided hereinbelow with reference to the sequence listing part.

[1721] GAM129 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM129 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1722] An enzyme complex designated DICER COMPLEX, dices the GAM129 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM129 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 28%) nucleotide sequence of GAM129 RNA is designated SEQ ID:1804, and is provided hereinbelow with reference to the sequence listing part.

[1723] GAM129 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM129 target RNA, herein designated GAM TARGET RNA. GAM129 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1724] GAM129 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM129 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM129 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM129 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM129 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1725] The complementary binding of GAM129 RNA, herein designated GAM RNA, to target binding sites on GAM129 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM129 target RNA, herein designated GAM TARGET RNA, into GAM129 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1726] It is appreciated that GAM129 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM129 target genes. The mRNA of each one of this plurality of GAM129 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM129 RNA, herein designated GAM RNA, and which when bound by GAM129 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM129 target proteins.

[1727] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM129 gene, herein designated GAM GENE, on one or more GAM129 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1728] It is yet further appreciated that specific functions, and accordingly utilities, of GAM129 correlate with, and may be deduced from, the identity of the target genes which GAM129 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1729] Nucleotide sequences of the GAM129 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM129 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM129 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM129 are further described hereinbelow with reference to Table Table1.

[1730] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM129 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1731]

[1732] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 130 (GAM130) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1733] GAM130 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM130 was detected is described hereinabove with reference to Figs. 2-8.

[1734] GAM130 gene, herein designated GAM GENE, and GAM130 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1735] GAM130 gene, herein designated GAM GENE, encodes a GAM130 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM130 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM130 precursor RNA is designated SEQ ID:1812, and is provided hereinbelow with reference to the sequence listing part.

[1736] GAM130 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM130 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1737] An enzyme complex designated DICER COMPLEX, dices the GAM130 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM130 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM130 RNA is designated SEQ ID:1813, and is provided hereinbelow with reference to the sequence listing part.

[1738] GAM130 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM130 target RNA, herein designated GAM TARGET RNA. GAM130 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1739] GAM130 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM130 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM130 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM130 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM130 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1740] The complementary binding of GAM130 RNA, herein designated GAM RNA, to target binding sites on GAM130 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM130 target RNA, herein designated GAM TARGET RNA, into GAM130 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1741] It is appreciated that GAM130 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM130 target genes. The mRNA of each one of this plurality of GAM130 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM130 RNA,

herein designated GAM RNA, and which when bound by GAM130 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM130 target proteins.

[1742] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM130 gene, herein designated GAM GENE, on one or more GAM130 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1743] It is yet further appreciated that specific functions, and accordingly utilities, of GAM130 correlate with, and may be deduced from, the identity of the target genes which

GAM130 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1744] Nucleotide sequences of the GAM130 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM130 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM130 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM130 are further described hereinbelow with reference to Table Table1.

[1745] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM130 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1746]

[1747] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 131 (GAM131) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1748] GAM131 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM131 was detected is described hereinabove with reference to Figs. 2-8.

[1749] GAM131 gene, herein designated GAM GENE, and GAM131 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1750] GAM131 gene, herein designated GAM GENE, encodes a GAM131 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM131 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM131 precursor RNA is designated SEQ ID:1825, and is provided hereinbelow with reference to the sequence listing part.

[1751] GAM131 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM131 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[1752] An enzyme complex designated DICER COMPLEX, dices the GAM131 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM131 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM131 RNA is designated SEQ ID:1826, and is provided hereinbelow with reference to the sequence listing part.

[1753] GAM131 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM131 target RNA, herein designated GAM TARGET RNA.
GAM131 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1754] GAM131 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM131 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM131 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM131 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM131 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1755] The complementary binding of GAM131 RNA, herein designated GAM RNA, to target binding sites on GAM131 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM131 target RNA, herein desig-

nated GAM TARGET RNA, into GAM131 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1756] It is appreciated that GAM131 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM131 target genes. The mRNA of each one of this plurality of GAM131 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM131 RNA, herein designated GAM RNA, and which when bound by GAM131 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM131 target proteins.

[1757] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM131 gene, herein designated GAM GENE, on one or more GAM131 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1758] It is yet further appreciated that specific functions, and accordingly utilities, of GAM131 correlate with, and may be deduced from, the identity of the target genes which GAM131 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1759] Nucleotide sequences of the GAM131 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM131 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM131 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM131 are further described hereinbelow with reference to Table Table1.

[1760] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM131 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[1761]

[1762] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 132 (GAM132) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1763] GAM132 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM132 was detected is described hereinabove with reference to Figs. 2-8.

[1764] GAM132 gene, herein designated GAM GENE, and GAM132 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1765] GAM132 gene, herein designated GAM GENE, encodes a GAM132 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM132 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM132 precursor RNA is designated SEQ ID:1842, and is provided hereinbelow with ref-

erence to the sequence listing part.

[1766] GAM132 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM132 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1767] An enzyme complex designated DICER COMPLEX, dices the GAM132 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM132 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 20%) nucleotide sequence of GAM132 RNA is designated SEQ ID:1843, and is provided hereinbelow with reference to the sequence listing part.

[1768] GAM132 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM132 target RNA, herein designated GAM TARGET RNA.

GAM132 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1769] GAM132 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM132 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM132 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM132 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM132 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1770] The complementary binding of GAM132 RNA, herein designated GAM RNA, to target binding sites on GAM132 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM132 target RNA, herein designated GAM TARGET RNA, into GAM132 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1771] It is appreciated that GAM132 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM132 target genes. The mRNA of each one of this plurality of GAM132 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM132 RNA, herein designated GAM RNA, and which when bound by GAM132 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM132 target proteins.

[1772] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM132 gene, herein designated GAM GENE, on one or more GAM132 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1773] It is yet further appreciated that specific functions, and accordingly utilities, of GAM132 correlate with, and may be deduced from, the identity of the target genes which GAM132 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1774] Nucleotide sequences of the GAM132 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM132 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM132 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM132 are further described hereinbelow with reference to Table Table1.

[1775] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM132 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1776]

[1777] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 133 (GAM133) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1778] GAM133 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM133 was detected is described hereinabove with reference to Figs. 2-8.

[1779] GAM133 gene, herein designated GAM GENE, and GAM133 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[1780] GAM133 gene, herein designated GAM GENE, encodes a GAM133 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM133 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM133 precursor RNA is designated SEQ ID:1857, and is provided hereinbelow with reference to the sequence listing part.

[1781] GAM133 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM133 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1782] An enzyme complex designated DICER COMPLEX, dices the GAM133 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM133 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM133 RNA is designated SEQ ID:1858, and is provided hereinbelow with reference to the sequence listing part.

[1783] GAM133 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM133 target RNA, herein designated GAM TARGET RNA. GAM133 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1784] GAM133 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM133 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM133 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM133 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM133 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1785] The complementary binding of GAM133 RNA, herein designated GAM RNA, to target binding sites on GAM133 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM133 target RNA, herein designated GAM TARGET RNA, into GAM133 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1786] It is appreciated that GAM133 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM133 target genes. The mRNA of each one of this plurality of GAM133 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM133 RNA, herein designated GAM RNA, and which when bound by GAM133 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM133 target proteins.

[1787] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM133 gene, herein designated GAM GENE, on one or more GAM133 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[1788] It is yet further appreciated that specific functions, and accordingly utilities, of GAM133 correlate with, and may be deduced from, the identity of the target genes which GAM133 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1789] Nucleotide sequences of the GAM133 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM133 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM133 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM133 are further described hereinbelow with reference to Table Table1.

[1790] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM133 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1791]

[1792] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

134 (GAM134) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1793] GAM134 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM134 was detected is described hereinabove with reference to Figs. 2-8.

[1794] GAM134 gene, herein designated GAM GENE, and GAM134 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1795] GAM134 gene, herein designated GAM GENE, encodes a GAM134 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM134 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM134 precursor RNA is designated SEQ ID:1880, and is provided hereinbelow with reference to the sequence listing part.

[1796] GAM134 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM134 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1797] An enzyme complex designated DICER COMPLEX, dices the GAM134 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM134 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM134 RNA is designated SEQ ID:1881, and is provided hereinbelow with reference to the sequence listing part.

[1798] GAM134 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM134 target RNA, herein designated GAM TARGET RNA. GAM134 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1799] GAM134 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM134 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM134 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM134 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM134 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1800] The complementary binding of GAM134 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM134 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM134 target RNA, herein designated GAM TARGET RNA, into GAM134 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1801] It is appreciated that GAM134 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM134 target genes. The mRNA of each one of this plurality of GAM134 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM134 RNA, herein designated GAM RNA, and which when bound by GAM134 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM134 target proteins.

[1802] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM134 gene, herein designated GAM GENE, on one or more GAM134 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1803] It is yet further appreciated that specific functions, and accordingly utilities, of GAM134 correlate with, and may be deduced from, the identity of the target genes which GAM134 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1804] Nucleotide sequences of the GAM134 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM134 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM134 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM134 are further described hereinbelow with reference to Table Table1.

[1805] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM134 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1806]

[1807] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 135 (GAM135) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1808] GAM135 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM135 was detected is described hereinabove with reference to Figs. 2-8.

[1809] GAM135 gene, herein designated GAM GENE, and GAM135 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1810] GAM135 gene, herein designated GAM GENE, encodes a GAM135 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM135 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM135 precursor RNA is designated SEQ ID:1887, and is provided hereinbelow with reference to the sequence listing part.

[1811] GAM135 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM135 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1812] An enzyme complex designated DICER COMPLEX, dices the GAM135 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM135 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 57%) nucleotide sequence of GAM135 RNA is designated SEQ ID:1888, and is provided hereinbelow with reference to the sequence listing part.

[1813] GAM135 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM135 target RNA, herein designated GAM TARGET RNA. GAM135 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1814] GAM135 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM135 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM135 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM135 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM135 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1815] The complementary binding of GAM135 RNA, herein designated GAM RNA, to target binding sites on GAM135 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM135 target RNA, herein designated GAM TARGET RNA, into GAM135 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1816] It is appreciated that GAM135 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM135 target genes. The mRNA of each one of this plurality of GAM135 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM135 RNA, herein designated GAM RNA, and which when bound by

GAM135 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM135 target proteins.

[1817] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM135 gene, herein designated GAM GENE, on one or more GAM135 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1818] It is yet further appreciated that specific functions, and accordingly utilities, of GAM135 correlate with, and may be deduced from, the identity of the target genes which GAM135 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[1819] Nucleotide sequences of the GAM135 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM135 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM135 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM135 are further described hereinbelow with reference to Table Table1.

[1820] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM135 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1821]

[1822] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 136 (GAM136) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1823] GAM136 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM136 was detected is described hereinabove with reference to Figs. 2-8.

[1824] GAM136 gene, herein designated GAM GENE, and GAM136 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1825] GAM136 gene, herein designated GAM GENE, encodes a GAM136 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM136 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM136 precursor RNA is designated SEQ ID:1898, and is provided hereinbelow with reference to the sequence listing part.

[1826] GAM136 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM136 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[1827] An enzyme complex designated DICER COMPLEX, dices the GAM136 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM136 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM136 RNA is designated SEQ ID:1899, and is provided hereinbelow with reference to the sequence listing part.

[1828] GAM136 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM136 target RNA, herein designated GAM TARGET RNA. GAM136 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1829] GAM136 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM136 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM136 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM136 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM136 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1830] The complementary binding of GAM136 RNA, herein designated GAM RNA, to target binding sites on GAM136 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM136 target RNA, herein designated GAM TARGET RNA, into GAM136 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1831] It is appreciated that GAM136 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM136 target genes. The mRNA of each one of this plurality of GAM136 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM136 RNA, herein designated GAM RNA, and which when bound by GAM136 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM136 target proteins.

[1832] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM136 gene, herein designated GAM GENE, on one or more GAM136 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1833] It is yet further appreciated that specific functions, and accordingly utilities, of GAM136 correlate with, and may be deduced from, the identity of the target genes which GAM136 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1834] Nucleotide sequences of the GAM136 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM136 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM136 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM136 are further described hereinbelow with reference to Table Table1.

[1835] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM136 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1836]

[1837] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 137 (GAM137) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1838] GAM137 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM137 was detected is described hereinabove with reference to Figs. 2-8.

[1839] GAM137 gene, herein designated GAM GENE, and GAM137 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1840] GAM137 gene, herein designated GAM GENE, encodes a GAM137 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM137 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM137 precursor RNA is designated SEQ ID:1907, and is provided hereinbelow with reference to the sequence listing part.

[1841] GAM137 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM137 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1842] An enzyme complex designated DICER COMPLEX, dices the GAM137 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM137 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 79%) nucleotide sequence of GAM137 RNA is designated SEQ ID:1908, and is provided hereinbelow with reference to the sequence listing part.

[1843] GAM137 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM137

target RNA, herein designated GAM TARGET RNA.

GAM137 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1844] GAM137 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM137 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM137 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM137 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM137 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1845] The complementary binding of GAM137 RNA, herein designated GAM RNA, to target binding sites on GAM137 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM137 target RNA, herein designated GAM TARGET RNA, into GAM137 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1846] It is appreciated that GAM137 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM137 target genes. The mRNA of each one of this plurality of GAM137 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM137 RNA, herein designated GAM RNA, and which when bound by GAM137 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM137 target proteins.

[1847] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM137 gene, herein designated GAM GENE, on one or more GAM137 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1848] It is yet further appreciated that specific functions, and accordingly utilities, of GAM137 correlate with, and may be deduced from, the identity of the target genes which GAM137 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1849] Nucleotide sequences of the GAM137 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM137 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM137 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM137 are further described hereinbelow with reference to Table Table1.

[1850] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM137 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1851]

[1852] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 138 (GAM138) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1853] GAM138 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM138 was detected is described hereinabove with reference to Figs. 2-8.

[1854] GAM138 gene, herein designated GAM GENE, and GAM138 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1855] GAM138 gene, herein designated GAM GENE, encodes a GAM138 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM138 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM138 precursor RNA is designated SEQ ID:1922, and is provided hereinbelow with reference to the sequence listing part.

[1856] GAM138 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM138 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1857] An enzyme complex designated DICER COMPLEX, dices the GAM138 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM138 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM138 RNA is designated SEQ ID:1923, and is provided hereinbelow with reference to the sequence listing part.

[1858] GAM138 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM138 target RNA, herein designated GAM TARGET RNA. GAM138 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1859] GAM138 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM138 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM138 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM138 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM138 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1860] The complementary binding of GAM138 RNA, herein designated GAM RNA, to target binding sites on GAM138 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM138 target RNA, herein designated GAM TARGET RNA, into GAM138 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1861] It is appreciated that GAM138 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM138 target genes. The mRNA of each one of this plu-

rality of GAM138 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM138 RNA, herein designated GAM RNA, and which when bound by GAM138 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM138 target proteins.

[1862] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM138 gene, herein designated GAM GENE, on one or more GAM138 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1863] It is yet further appreciated that specific functions, and accordingly utilities, of GAM138 correlate with, and may be deduced from, the identity of the target genes which GAM138 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1864] Nucleotide sequences of the GAM138 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM138 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM138 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM138 are further described hereinbelow with reference to Table Table1.

[1865] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM138 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1866]

[1867] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 139 (GAM139) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[1868] GAM139 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM139 was detected is described hereinabove with reference to Figs. 2-8.

[1869] GAM139 gene, herein designated GAM GENE, and GAM139 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1870] GAM139 gene, herein designated GAM GENE, encodes a GAM139 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM139 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM139 precursor RNA is designated SEQ ID:1947, and is provided hereinbelow with reference to the sequence listing part.

[1871] GAM139 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM139 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1872] An enzyme complex designated DICER COMPLEX, dices the GAM139 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM139 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 38%) nucleotide sequence of GAM139 RNA is designated SEQ ID:1948, and is provided hereinbelow with reference to the sequence listing part.

[1873] GAM139 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM139 target RNA, herein designated GAM TARGET RNA.
GAM139 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[1874] GAM139 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM139 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM139 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM139 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM139 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1875] The complementary binding of GAM139 RNA, herein designated GAM RNA, to target binding sites on GAM139 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM139 target RNA, herein designated GAM TARGET RNA, into GAM139 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1876] It is appreciated that GAM139 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM139 target genes. The mRNA of each one of this plurality of GAM139 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM139 RNA, herein designated GAM RNA, and which when bound by GAM139 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM139 target proteins.

[1877] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM139 gene, herein designated GAM GENE, on one or more GAM139 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1878] It is yet further appreciated that specific functions, and accordingly utilities, of GAM139 correlate with, and may be deduced from, the identity of the target genes which GAM139 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1879] Nucleotide sequences of the GAM139 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM139 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM139 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM139 are further described hereinbelow with reference to Table Table1.

[1880] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM139 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1881]

[1882] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 140 (GAM140) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1883] GAM140 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM140 was detected is described hereinabove with reference to Figs. 2-8.

[1884] GAM140 gene, herein designated GAM GENE, and GAM140 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1885] GAM140 gene, herein designated GAM GENE, encodes a GAM140 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM140 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM140 precursor RNA is designated SEQ ID:1954, and is provided hereinbelow with reference to the sequence listing part.

[1886] GAM140 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM140 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1887] An enzyme complex designated DICER COMPLEX, dices the GAM140 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM140 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of

GAM140 RNA is designated SEQ ID:1955, and is provided hereinbelow with reference to the sequence listing part.

[1888] GAM140 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM140 target RNA, herein designated GAM TARGET RNA. GAM140 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1889] GAM140 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM140 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM140 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM140 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM140 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1890] The complementary binding of GAM140 RNA, herein designated GAM RNA, to target binding sites on GAM140 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM140 target RNA, herein designated GAM TARGET RNA, into GAM140 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1891] It is appreciated that GAM140 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM140 target genes. The mRNA of each one of this plurality of GAM140 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM140 RNA, herein designated GAM RNA, and which when bound by GAM140 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM140 target proteins.

[1892] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM140 gene, herein designated GAM GENE, on one or more GAM140 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1893] It is yet further appreciated that specific functions, and accordingly utilities, of GAM140 correlate with, and may be deduced from, the identity of the target genes which GAM140 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1894] Nucleotide sequences of the GAM140 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM140 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM140 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM140 are further described hereinbelow with reference to Table Table1.

[1895] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM140 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1896]

[1897] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 141 (GAM141) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1898] GAM141 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM141 was detected is described hereinabove

with reference to Figs. 2-8.

[1899] GAM141 gene, herein designated GAM GENE, and GAM141 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1900] GAM141 gene, herein designated GAM GENE, encodes a GAM141 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM141 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM141 precursor RNA is designated SEQ ID:1967, and is provided hereinbelow with reference to the sequence listing part.

[1901] GAM141 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM141 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1902] An enzyme complex designated DICER COMPLEX, dices the GAM141 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM141 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM141 RNA is designated SEQ ID:1968, and is provided hereinbelow with reference to the sequence listing part.

[1903] GAM141 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM141 target RNA, herein designated GAM TARGET RNA. GAM141 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1904] GAM141 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM141 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM141 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM141 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM141 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1905] The complementary binding of GAM141 RNA, herein designated GAM RNA, to target binding sites on GAM141 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM141 target RNA, herein designated GAM TARGET RNA, into GAM141 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[1906] It is appreciated that GAM141 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM141 target genes. The mRNA of each one of this plurality of GAM141 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM141 RNA, herein designated GAM RNA, and which when bound by GAM141 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM141 target proteins.

[1907] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM141 gene, herein designated GAM GENE, on one or more GAM141 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1908] It is yet further appreciated that specific functions, and accordingly utilities, of GAM141 correlate with, and may be deduced from, the identity of the target genes which GAM141 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1909] Nucleotide sequences of the GAM141 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM141 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM141 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM141 are further described hereinbelow with reference to Table Table1.

[1910] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM141 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1911]

[1912] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 142 (GAM142) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1913] GAM142 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM142 was detected is described hereinabove with reference to Figs. 2-8.

[1914] GAM142 gene, herein designated GAM GENE, and GAM142 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1915] GAM142 gene, herein designated GAM GENE, encodes a GAM142 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM142 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM142 precursor RNA is designated SEQ ID:1983, and is provided hereinbelow with reference to the sequence listing part.

[1916] GAM142 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM142 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1917] An enzyme complex designated DICER COMPLEX, dices the GAM142 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM142 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM142 RNA is designated SEQ ID:1984, and is provided hereinbelow with reference to the sequence listing part.

[1918] GAM142 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM142 target RNA, herein designated GAM TARGET RNA.

GAM142 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1919] GAM142 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM142 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM142 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM142 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM142 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1920] The complementary binding of GAM142 RNA, herein designated GAM RNA, to target binding sites on GAM142 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM142 target RNA, herein designated GAM TARGET RNA, into GAM142 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1921] It is appreciated that GAM142 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM142 target genes. The mRNA of each one of this plurality of GAM142 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM142 RNA, herein designated GAM RNA, and which when bound by GAM142 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM142 target proteins.

[1922] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM142 gene, herein designated GAM GENE, on one or more GAM142 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1923] It is yet further appreciated that specific functions, and accordingly utilities, of GAM142 correlate with, and may be deduced from, the identity of the target genes which GAM142 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1924] Nucleotide sequences of the GAM142 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM142 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM142 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM142 are further described hereinbelow with reference to Table Table1.

[1925] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM142 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1926]

[1927] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 143 (GAM143) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1928] GAM143 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM143 was detected is described hereinabove with reference to Figs. 2-8.

[1929] GAM143 gene, herein designated GAM GENE, and GAM143 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1930] GAM143 gene, herein designated GAM GENE, encodes a

GAM143 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM143 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM143 precursor RNA is designated SEQ ID:1992, and is provided hereinbelow with reference to the sequence listing part.

[1931] GAM143 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM143 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1932] An enzyme complex designated DICER COMPLEX, dices the GAM143 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM143 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM143 RNA is designated SEQ ID:1993, and is provided hereinbelow with reference to the sequence listing part.

[1933] GAM143 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM143 target RNA, herein designated GAM TARGET RNA. GAM143 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1934] GAM143 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM143 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM143 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM143 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM143 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1935] The complementary binding of GAM143 RNA, herein designated GAM RNA, to target binding sites on GAM143 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM143 target RNA, herein designated GAM TARGET RNA, into GAM143 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1936] It is appreciated that GAM143 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM143 target genes. The mRNA of each one of this plurality of GAM143 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM143 RNA, herein designated GAM RNA, and which when bound by GAM143 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM143 target proteins.

[1937] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM143 gene, herein designated GAM GENE, on one or more GAM143 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1938] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM143 correlate with, and may be deduced from, the identity of the target genes which GAM143 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1939] Nucleotide sequences of the GAM143 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM143 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM143 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM143 are further described hereinbelow with reference to Table Table1.

[1940] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM143 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1941]

[1942] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 144 (GAM144) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[1943] GAM144 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM144 was detected is described hereinabove with reference to Figs. 2-8.

[1944] GAM144 gene, herein designated GAM GENE, and GAM144 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1945] GAM144 gene, herein designated GAM GENE, encodes a GAM144 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM144 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM144 precursor RNA is designated SEQ ID:2008, and is provided hereinbelow with reference to the sequence listing part.

[1946] GAM144 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM144 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1947] An enzyme complex designated DICER COMPLEX, dices the GAM144 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM144 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM144 RNA is designated SEQ ID:2009, and is provided hereinbelow with reference to the sequence listing part.

[1948] GAM144 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM144 target RNA, herein designated GAM TARGET RNA. GAM144 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1949] GAM144 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM144 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM144 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM144 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM144 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1950] The complementary binding of GAM144 RNA, herein designated GAM RNA, to target binding sites on GAM144 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM144 target RNA, herein designated GAM TARGET RNA, into GAM144 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1951] It is appreciated that GAM144 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM144 target genes. The mRNA of each one of this plurality of GAM144 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM144 RNA, herein designated GAM RNA, and which when bound by GAM144 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM144 target proteins.

[1952] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM144 gene, herein designated GAM GENE, on one or more GAM144 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1953] It is yet further appreciated that specific functions, and accordingly utilities, of GAM144 correlate with, and may be deduced from, the identity of the target genes which GAM144 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1954] Nucleotide sequences of the GAM144 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM144 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM144 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM144 are further described hereinbelow with reference to Table Table1.

[1955] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM144 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1956]

[1957] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 145 (GAM145) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1958] GAM145 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM145 was detected is described hereinabove with reference to Figs. 2-8.

[1959] GAM145 gene, herein designated GAM GENE, and GAM145 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1960] GAM145 gene, herein designated GAM GENE, encodes a GAM145 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM145 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM145 precursor RNA is designated SEQ ID:2020, and is provided hereinbelow with reference to the sequence listing part.

[1961] GAM145 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM145 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1962] An enzyme complex designated DICER COMPLEX, dices the GAM145 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM145 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM145 RNA is designated SEQ ID:2021, and is provided

hereinbelow with reference to the sequence listing part.

[1963] GAM145 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM145 target RNA, herein designated GAM TARGET RNA.

GAM145 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1964] GAM145 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM145 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM145 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM145 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM145 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1965] The complementary binding of GAM145 RNA, herein designated GAM RNA, to target binding sites on GAM145 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM145 target RNA, herein designated GAM TARGET RNA, into GAM145 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1966] It is appreciated that GAM145 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM145 target genes. The mRNA of each one of this plurality of GAM145 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM145 RNA, herein designated GAM RNA, and which when bound by GAM145 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM145 tar-

get proteins.

[1967] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM145 gene, herein designated GAM GENE, on one or more GAM145 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1968] It is yet further appreciated that specific functions, and accordingly utilities, of GAM145 correlate with, and may be deduced from, the identity of the target genes which GAM145 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1969] Nucleotide sequences of the GAM145 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM145 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM145 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM145 are further described hereinbelow with reference to Table Table1.

[1970] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM145 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1971]

[1972] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 146 (GAM146) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1973] GAM146 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM146 was detected is described hereinabove with reference to Figs. 2-8.

[1974] GAM146 gene, herein designated GAM GENE, and GAM146 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1975] GAM146 gene, herein designated GAM GENE, encodes a GAM146 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM146 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM146 precursor RNA is designated SEQ ID:2040, and is provided hereinbelow with reference to the sequence listing part.

[1976] GAM146 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM146 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1977] An enzyme complex designated DICER COMPLEX, dices

the GAM146 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM146 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 48%) nucleotide sequence of GAM146 RNA is designated SEQ ID:2041, and is provided hereinbelow with reference to the sequence listing part.

[1978] GAM146 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM146 target RNA, herein designated GAM TARGET RNA. GAM146 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1979] GAM146 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM146 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM146

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM146 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM146 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[1980] The complementary binding of GAM146 RNA, herein designated GAM RNA, to target binding sites on GAM146 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM146 target RNA, herein designated GAM TARGET RNA, into GAM146 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1981] It is appreciated that GAM146 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM146 target genes. The mRNA of each one of this plurality of GAM146 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM146 RNA, herein designated GAM RNA, and which when bound by GAM146 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM146 target proteins.

[1982] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM146 gene, herein designated GAM GENE, on one or more GAM146 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1983] It is yet further appreciated that specific functions, and accordingly utilities, of GAM146 correlate with, and may be deduced from, the identity of the target genes which GAM146 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1984] Nucleotide sequences of the GAM146 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM146 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM146 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM146 are further described hereinbelow with reference to Table Table1.

[1985] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM146 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[1986]

[1987] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 147 (GAM147) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[1988] GAM147 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM147 was detected is described hereinabove with reference to Figs. 2-8.

[1989] GAM147 gene, herein designated GAM GENE, and GAM147 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[1990] GAM147 gene, herein designated GAM GENE, encodes a GAM147 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM147 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM147 precursor RNA is designated SEQ ID:2112, and is provided hereinbelow with reference to the sequence listing part.

[1991] GAM147 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM147 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[1992] An enzyme complex designated DICER COMPLEX, dices the GAM147 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM147 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 22%) nucleotide sequence of GAM147 RNA is designated SEQ ID:2113, and is provided hereinbelow with reference to the sequence listing part.

[1993] GAM147 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM147 target RNA, herein designated GAM TARGET RNA. GAM147 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[1994] GAM147 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM147 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM147 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM147 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM147 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[1995] The complementary binding of GAM147 RNA, herein designated GAM RNA, to target binding sites on GAM147 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM147 target RNA, herein designated GAM TARGET RNA, into GAM147 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[1996] It is appreciated that GAM147 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM147 target genes. The mRNA of each one of this plurality of GAM147 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM147 RNA, herein designated GAM RNA, and which when bound by GAM147 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM147 target proteins.

[1997] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM147 gene, herein designated GAM GENE, on one or

more GAM147 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[1998] It is yet further appreciated that specific functions, and accordingly utilities, of GAM147 correlate with, and may be deduced from, the identity of the target genes which GAM147 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[1999] Nucleotide sequences of the GAM147 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM147 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM147 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM147 are further de-

scribed hereinbelow with reference to Table Table1.

[2000] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM147 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2001]

[2002] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 148 (GAM148) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2003] GAM148 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM148 was detected is described hereinabove with reference to Figs. 2-8.

[2004] GAM148 gene, herein designated GAM GENE, and GAM148 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2005] GAM148 gene, herein designated GAM GENE, encodes a GAM148 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM148 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM148 precursor RNA is designated SEQ ID:2125, and is provided hereinbelow with reference to the sequence listing part.

[2006] GAM148 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM148 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2007] An enzyme complex designated DICER COMPLEX, dices the GAM148 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM148 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM148 RNA is designated SEQ ID:2126, and is provided hereinbelow with reference to the sequence listing part.

[2008] GAM148 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM148 target RNA, herein designated GAM TARGET RNA. GAM148 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2009] GAM148 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM148 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM148 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM148 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM148 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2010] The complementary binding of GAM148 RNA, herein designated GAM RNA, to target binding sites on GAM148 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM148 target RNA, herein designated GAM TARGET RNA, into GAM148 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2011] It is appreciated that GAM148 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM148 target genes. The mRNA of each one of this plurality of GAM148 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM148 RNA, herein designated GAM RNA, and which when bound by GAM148 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM148 target proteins.

[2012] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM148 gene, herein designated GAM GENE, on one or more GAM148 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2013] It is yet further appreciated that specific functions, and accordingly utilities, of GAM148 correlate with, and may

be deduced from, the identity of the target genes which GAM148 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2014] Nucleotide sequences of the GAM148 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM148 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM148 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM148 are further described hereinbelow with reference to Table Table1.

[2015] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM148 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2016]

[2017] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 149 (GAM149) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2018] GAM149 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM149 was detected is described hereinabove with reference to Figs. 2-8.

[2019] GAM149 gene, herein designated GAM GENE, and GAM149 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2020] GAM149 gene, herein designated GAM GENE, encodes a GAM149 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM149 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM149 precursor RNA is designated SEQ ID:2132, and is provided hereinbelow with reference to the sequence listing part.

[2021] GAM149 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM149 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2022] An enzyme complex designated DICER COMPLEX, dices the GAM149 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM149 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM149 RNA is designated SEQ ID:2133, and is provided hereinbelow with reference to the sequence listing part.

[2023] GAM149 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM149 target RNA, herein designated GAM TARGET RNA. GAM149 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2024] GAM149 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM149 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM149 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM149 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM149 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2025] The complementary binding of GAM149 RNA, herein designated GAM RNA, to target binding sites on GAM149 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM149 target RNA, herein designated GAM TARGET RNA, into GAM149 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2026] It is appreciated that GAM149 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM149 target genes. The mRNA of each one of this plurality of GAM149 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM149 RNA, herein designated GAM RNA, and which when bound by GAM149 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM149 target proteins.

[2027] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM149 gene, herein designated GAM GENE, on one or more GAM149 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2028] It is yet further appreciated that specific functions, and accordingly utilities, of GAM149 correlate with, and may be deduced from, the identity of the target genes which GAM149 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2029] Nucleotide sequences of the GAM149 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM149 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM149 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM149 are further described hereinbelow with reference to Table Table1.

[2030] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM149 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2031]

[2032] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 150 (GAM150) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2033] GAM150 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM150 was detected is described hereinabove with reference to Figs. 2-8.

[2034] GAM150 gene, herein designated GAM GENE, and GAM150 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2035] GAM150 gene, herein designated GAM GENE, encodes a GAM150 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM150 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM150 precursor RNA is desig-

nated SEQ ID:2141, and is provided hereinbelow with reference to the sequence listing part.

[2036] GAM150 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM150 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2037] An enzyme complex designated DICER COMPLEX, dices the GAM150 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM150 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM150 RNA is designated SEQ ID:2142, and is provided hereinbelow with reference to the sequence listing part.

[2038] GAM150 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM150 target RNA, herein designated GAM TARGET RNA. GAM150 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2039] GAM150 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM150 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM150 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM150 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM150

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2040] The complementary binding of GAM150 RNA, herein designated GAM RNA, to target binding sites on GAM150 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM150 target RNA, herein designated GAM TARGET RNA, into GAM150 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2041] It is appreciated that GAM150 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM150 target genes. The mRNA of each one of this plurality of GAM150 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM150 RNA, herein designated GAM RNA, and which when bound by GAM150 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM150 target proteins.

[2042] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM150 gene, herein designated GAM GENE, on one or more GAM150 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2043] It is yet further appreciated that specific functions, and accordingly utilities, of GAM150 correlate with, and may be deduced from, the identity of the target genes which GAM150 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2044] Nucleotide sequences of the GAM150 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM150 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM150 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM150 are further described hereinbelow with reference to Table Table1.

[2045] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM150 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2046]

[2047] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 151 (GAM151) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2048] GAM151 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM151 was detected is described hereinabove with reference to Figs. 2-8.

[2049] GAM151 gene, herein designated GAM GENE, and

GAM151 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2050] GAM151 gene, herein designated GAM GENE, encodes a GAM151 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM151 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM151 precursor RNA is designated SEQ ID:2157, and is provided hereinbelow with reference to the sequence listing part.

[2051] GAM151 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM151 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2052] An enzyme complex designated DICER COMPLEX, dices the GAM151 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM151 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM151 RNA is designated SEQ ID:2158, and is provided hereinbelow with reference to the sequence listing part.

[2053] GAM151 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM151 target RNA, herein designated GAM TARGET RNA. GAM151 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2054] GAM151 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM151 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM151 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM151 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM151 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2055] The complementary binding of GAM151 RNA, herein designated GAM RNA, to target binding sites on GAM151 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM151 target RNA, herein designated GAM TARGET RNA, into GAM151 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2056] It is appreciated that GAM151 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM151 target genes. The mRNA of each one of this plurality of GAM151 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM151 RNA, herein designated GAM RNA, and which when bound by GAM151 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM151 target proteins.

[2057] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM151 gene, herein designated GAM GENE, on one or more GAM151 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2058] It is yet further appreciated that specific functions, and accordingly utilities, of GAM151 correlate with, and may be deduced from, the identity of the target genes which GAM151 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2059] Nucleotide sequences of the GAM151 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM151 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM151 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM151 are further described hereinbelow with reference to Table Table1.

[2060] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM151 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2061]

[2062] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 152 (GAM152) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2063] GAM152 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM152 was detected is described hereinabove with reference to Figs. 2-8.

[2064] GAM152 gene, herein designated GAM GENE, and GAM152 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2065] GAM152 gene, herein designated GAM GENE, encodes a GAM152 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM152 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM152 precursor RNA is designated SEQ ID:2170, and is provided hereinbelow with reference to the sequence listing part.

[2066] GAM152 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM152 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2067] An enzyme complex designated DICER COMPLEX, dices the GAM152 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM152 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM152 RNA is designated SEQ ID:2171, and is provided

hereinbelow with reference to the sequence listing part.

[2068] GAM152 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM152 target RNA, herein designated GAM TARGET RNA.
GAM152 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2069] GAM152 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM152 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM152 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM152 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM152 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2070] The complementary binding of GAM152 RNA, herein designated GAM RNA, to target binding sites on GAM152 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM152 target RNA, herein designated GAM TARGET RNA, into GAM152 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2071] It is appreciated that GAM152 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM152 target genes. The mRNA of each one of this plurality of GAM152 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM152 RNA, herein designated GAM RNA, and which when bound by GAM152 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM152 tar-

get proteins.

[2072] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM152 gene, herein designated GAM GENE, on one or more GAM152 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2073] It is yet further appreciated that specific functions, and accordingly utilities, of GAM152 correlate with, and may be deduced from, the identity of the target genes which GAM152 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2074] Nucleotide sequences of the GAM152 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM152 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM152 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM152 are further described hereinbelow with reference to Table Table1.

[2075] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM152 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2076]

[2077] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 153 (GAM153) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2078] GAM153 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM153 was detected is described hereinabove with reference to Figs. 2-8.

[2079] GAM153 gene, herein designated GAM GENE, and GAM153 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2080] GAM153 gene, herein designated GAM GENE, encodes a GAM153 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM153 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM153 precursor RNA is designated SEQ ID:2182, and is provided hereinbelow with reference to the sequence listing part.

[2081] GAM153 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM153 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2082] An enzyme complex designated DICER COMPLEX, dices

the GAM153 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM153 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM153 RNA is designated SEQ ID:2183, and is provided hereinbelow with reference to the sequence listing part.

[2083] GAM153 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM153 target RNA, herein designated GAM TARGET RNA. GAM153 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2084] GAM153 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM153 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM153

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM153 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM153 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2085] The complementary binding of GAM153 RNA, herein designated GAM RNA, to target binding sites on GAM153 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM153 target RNA, herein designated GAM TARGET RNA, into GAM153 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2086] It is appreciated that GAM153 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM153 target genes. The mRNA of each one of this plurality of GAM153 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM153 RNA, herein designated GAM RNA, and which when bound by GAM153 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM153 target proteins.

[2087] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM153 gene, herein designated GAM GENE, on one or more GAM153 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2088] It is yet further appreciated that specific functions, and accordingly utilities, of GAM153 correlate with, and may be deduced from, the identity of the target genes which GAM153 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2089] Nucleotide sequences of the GAM153 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM153 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM153 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM153 are further described hereinbelow with reference to Table Table1.

[2090] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM153 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2091]

[2092] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 154 (GAM154) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2093] GAM154 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM154 was detected is described hereinabove with reference to Figs. 2-8.

[2094] GAM154 gene, herein designated GAM GENE, and GAM154 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2095] GAM154 gene, herein designated GAM GENE, encodes a GAM154 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM154 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM154 precursor RNA is designated SEQ ID:2194, and is provided hereinbelow with reference to the sequence listing part.

[2096] GAM154 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM154 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2097] An enzyme complex designated DICER COMPLEX, dices the GAM154 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM154 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM154 RNA is designated SEQ ID:2195, and is provided hereinbelow with reference to the sequence listing part.

[2098] GAM154 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM154 target RNA, herein designated GAM TARGET RNA. GAM154 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2099] GAM154 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM154 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM154 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM154 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM154 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[2100] The complementary binding of GAM154 RNA, herein designated GAM RNA, to target binding sites on GAM154 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM154 target RNA, herein designated GAM TARGET RNA, into GAM154 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2101] It is appreciated that GAM154 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM154 target genes. The mRNA of each one of this plurality of GAM154 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM154 RNA, herein designated GAM RNA, and which when bound by GAM154 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM154 target proteins.

[2102] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM154 gene, herein designated GAM GENE, on one or

more GAM154 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2103] It is yet further appreciated that specific functions, and accordingly utilities, of GAM154 correlate with, and may be deduced from, the identity of the target genes which GAM154 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2104] Nucleotide sequences of the GAM154 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM154 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM154 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM154 are further de-

scribed hereinbelow with reference to Table Table1.

[2105] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM154 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2106]

[2107] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 155 (GAM155) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2108] GAM155 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM155 was detected is described hereinabove with reference to Figs. 2-8.

[2109] GAM155 gene, herein designated GAM GENE, and GAM155 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2110] GAM155 gene, herein designated GAM GENE, encodes a GAM155 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM155 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM155 precursor RNA is designated SEQ ID:2204, and is provided hereinbelow with reference to the sequence listing part.

[2111] GAM155 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM155 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2112] An enzyme complex designated DICER COMPLEX, dices the GAM155 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM155 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM155 RNA is designated SEQ ID:2205, and is provided hereinbelow with reference to the sequence listing part.

[2113] GAM155 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM155 target RNA, herein designated GAM TARGET RNA. GAM155 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2114] GAM155 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM155 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM155 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM155 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM155 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2115] The complementary binding of GAM155 RNA, herein designated GAM RNA, to target binding sites on GAM155 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM155 target RNA, herein designated GAM TARGET RNA, into GAM155 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2116] It is appreciated that GAM155 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM155 target genes. The mRNA of each one of this plurality of GAM155 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM155 RNA, herein designated GAM RNA, and which when bound by GAM155 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM155 target proteins.

[2117] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM155 gene, herein designated GAM GENE, on one or more GAM155 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2118] It is yet further appreciated that specific functions, and accordingly utilities, of GAM155 correlate with, and may

be deduced from, the identity of the target genes which GAM155 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2119] Nucleotide sequences of the GAM155 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM155 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM155 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM155 are further described hereinbelow with reference to Table Table1.

[2120] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM155 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2121]

[2122] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 156 (GAM156) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2123] GAM156 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM156 was detected is described hereinabove with reference to Figs. 2-8.

[2124] GAM156 gene, herein designated GAM GENE, and GAM156 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2125] GAM156 gene, herein designated GAM GENE, encodes a GAM156 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM156 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM156 precursor RNA is designated SEQ ID:2227, and is provided hereinbelow with reference to the sequence listing part.

[2126] GAM156 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM156 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2127] An enzyme complex designated DICER COMPLEX, dices the GAM156 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM156 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 49%) nucleotide sequence of GAM156 RNA is designated SEQ ID:2228, and is provided hereinbelow with reference to the sequence listing part.

[2128] GAM156 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM156 target RNA, herein designated GAM TARGET RNA. GAM156 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2129] GAM156 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM156 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM156 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM156 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM156 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2130] The complementary binding of GAM156 RNA, herein designated GAM RNA, to target binding sites on GAM156 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM156 target RNA, herein designated GAM TARGET RNA, into GAM156 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2131] It is appreciated that GAM156 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM156 target genes. The mRNA of each one of this plurality of GAM156 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM156 RNA, herein designated GAM RNA, and which when bound by GAM156 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM156 target proteins.

[2132] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM156 gene, herein designated GAM GENE, on one or more GAM156 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2133] It is yet further appreciated that specific functions, and accordingly utilities, of GAM156 correlate with, and may be deduced from, the identity of the target genes which GAM156 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2134] Nucleotide sequences of the GAM156 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM156 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM156 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM156 are further described hereinbelow with reference to Table Table1.

[2135] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM156 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2136]

[2137] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 157 (GAM157) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2138] GAM157 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM157 was detected is described hereinabove with reference to Figs. 2-8.

[2139] GAM157 gene, herein designated GAM GENE, and GAM157 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2140] GAM157 gene, herein designated GAM GENE, encodes a GAM157 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM157 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM157 precursor RNA is desig-

nated SEQ ID:2240, and is provided hereinbelow with reference to the sequence listing part.

[2141] GAM157 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM157 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2142] An enzyme complex designated DICER COMPLEX, dices the GAM157 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM157 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM157 RNA is designated SEQ ID:2241, and is provided hereinbelow with reference to the sequence listing part.

[2143] GAM157 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM157 target RNA, herein designated GAM TARGET RNA. GAM157 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2144] GAM157 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM157 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM157 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM157 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM157

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2145] The complementary binding of GAM157 RNA, herein designated GAM RNA, to target binding sites on GAM157 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM157 target RNA, herein designated GAM TARGET RNA, into GAM157 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2146] It is appreciated that GAM157 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM157 target genes. The mRNA of each one of this plurality of GAM157 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM157 RNA, herein designated GAM RNA, and which when bound by GAM157 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM157 target proteins.

[2147] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM157 gene, herein designated GAM GENE, on one or more GAM157 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2148] It is yet further appreciated that specific functions, and accordingly utilities, of GAM157 correlate with, and may be deduced from, the identity of the target genes which GAM157 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2149] Nucleotide sequences of the GAM157 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM157 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM157 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM157 are further described hereinbelow with reference to Table Table1.

[2150] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM157 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2151]

[2152] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 158 (GAM158) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2153] GAM158 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM158 was detected is described hereinabove with reference to Figs. 2-8.

[2154] GAM158 gene, herein designated GAM GENE, and

GAM158 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2155] GAM158 gene, herein designated GAM GENE, encodes a GAM158 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM158 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM158 precursor RNA is designated SEQ ID:2266, and is provided hereinbelow with reference to the sequence listing part.

[2156] GAM158 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM158 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2157] An enzyme complex designated DICER COMPLEX, dices the GAM158 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM158 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM158 RNA is designated SEQ ID:2267, and is provided hereinbelow with reference to the sequence listing part.

[2158] GAM158 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM158 target RNA, herein designated GAM TARGET RNA. GAM158 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2159] GAM158 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM158 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM158 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM158 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM158 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2160] The complementary binding of GAM158 RNA, herein designated GAM RNA, to target binding sites on GAM158 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM158 target RNA, herein designated GAM TARGET RNA, into GAM158 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2161] It is appreciated that GAM158 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM158 target genes. The mRNA of each one of this plurality of GAM158 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM158 RNA, herein designated GAM RNA, and which when bound by GAM158 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM158 target proteins.

[2162] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM158 gene, herein designated GAM GENE, on one or more GAM158 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2163] It is yet further appreciated that specific functions, and accordingly utilities, of GAM158 correlate with, and may be deduced from, the identity of the target genes which GAM158 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2164] Nucleotide sequences of the GAM158 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM158 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM158 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM158 are further described hereinbelow with reference to Table Table1.

[2165] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM158 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2166]

[2167] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 159 (GAM159) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2168] GAM159 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM159 was detected is described hereinabove with reference to Figs. 2-8.

[2169] GAM159 gene, herein designated GAM GENE, and GAM159 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2170] GAM159 gene, herein designated GAM GENE, encodes a GAM159 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM159 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM159 precursor RNA is designated SEQ ID:2277, and is provided hereinbelow with reference to the sequence listing part.

[2171] GAM159 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM159 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2172] An enzyme complex designated DICER COMPLEX, dices the GAM159 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM159 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM159 RNA is designated SEQ ID:2278, and is provided hereinbelow with reference to the sequence listing part.

[2173] GAM159 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM159 target RNA, herein designated GAM TARGET RNA. GAM159 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2174] GAM159 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM159 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM159 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM159 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM159 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2175] The complementary binding of GAM159 RNA, herein designated GAM RNA, to target binding sites on GAM159 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM159 target RNA, herein designated GAM TARGET RNA, into GAM159 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2176] It is appreciated that GAM159 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM159 target genes. The mRNA of each one of this plurality of GAM159 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM159 RNA, herein designated GAM RNA, and which when bound by GAM159 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM159 target proteins.

[2177] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM159 gene, herein designated GAM GENE, on one or more GAM159 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2178] It is yet further appreciated that specific functions, and accordingly utilities, of GAM159 correlate with, and may be deduced from, the identity of the target genes which GAM159 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2179] Nucleotide sequences of the GAM159 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM159 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM159 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM159 are further described hereinbelow with reference to Table Table1.

[2180] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM159 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2181]

[2182] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 160 (GAM160) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2183] GAM160 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM160 was detected is described hereinabove with reference to Figs. 2-8.

[2184] GAM160 gene, herein designated GAM GENE, and GAM160 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2185] GAM160 gene, herein designated GAM GENE, encodes a GAM160 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM160 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM160 precursor RNA is designated SEQ ID:2290, and is provided hereinbelow with reference to the sequence listing part.

[2186] GAM160 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM160 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2187] An enzyme complex designated DICER COMPLEX, dices the GAM160 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM160 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM160 RNA is designated SEQ ID:2291, and is provided hereinbelow with reference to the sequence listing part.

[2188] GAM160 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM160 target RNA, herein designated GAM TARGET RNA. GAM160 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2189] GAM160 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM160 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM160 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM160 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM160 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2190] The complementary binding of GAM160 RNA, herein designated GAM RNA, to target binding sites on GAM160 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM160 target RNA, herein designated GAM TARGET RNA, into GAM160 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2191] It is appreciated that GAM160 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM160 target genes. The mRNA of each one of this plurality of GAM160 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM160 RNA,

herein designated GAM RNA, and which when bound by GAM160 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM160 target proteins.

[2192] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM160 gene, herein designated GAM GENE, on one or more GAM160 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2193] It is yet further appreciated that specific functions, and accordingly utilities, of GAM160 correlate with, and may be deduced from, the identity of the target genes which

GAM160 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2194] Nucleotide sequences of the GAM160 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM160 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM160 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM160 are further described hereinbelow with reference to Table Table1.

[2195] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM160 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2196]

[2197] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 161 (GAM161) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2198] GAM161 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM161 was detected is described hereinabove with reference to Figs. 2-8.

[2199] GAM161 gene, herein designated GAM GENE, and GAM161 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2200] GAM161 gene, herein designated GAM GENE, encodes a GAM161 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM161 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM161 precursor RNA is designated SEQ ID:2295, and is provided hereinbelow with reference to the sequence listing part.

[2201] GAM161 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM161 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[2202] An enzyme complex designated DICER COMPLEX, dices the GAM161 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM161 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM161 RNA is designated SEQ ID:2296, and is provided hereinbelow with reference to the sequence listing part.

[2203] GAM161 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM161 target RNA, herein designated GAM TARGET RNA.
GAM161 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2204] GAM161 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM161 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM161 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM161 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM161 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2205] The complementary binding of GAM161 RNA, herein designated GAM RNA, to target binding sites on GAM161 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM161 target RNA, herein desig-

nated GAM TARGET RNA, into GAM161 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2206] It is appreciated that GAM161 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM161 target genes. The mRNA of each one of this plurality of GAM161 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM161 RNA, herein designated GAM RNA, and which when bound by GAM161 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM161 target proteins.

[2207] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM161 gene, herein designated GAM GENE, on one or more GAM161 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2208] It is yet further appreciated that specific functions, and accordingly utilities, of GAM161 correlate with, and may be deduced from, the identity of the target genes which GAM161 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2209] Nucleotide sequences of the GAM161 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM161 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM161 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM161 are further described hereinbelow with reference to Table Table1.

[2210] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM161 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[2211]

[2212] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 162 (GAM162) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2213] GAM162 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM162 was detected is described hereinabove with reference to Figs. 2-8.

[2214] GAM162 gene, herein designated GAM GENE, and GAM162 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2215] GAM162 gene, herein designated GAM GENE, encodes a GAM162 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM162 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM162 precursor RNA is designated SEQ ID:2312, and is provided hereinbelow with ref-

erence to the sequence listing part.

[2216] GAM162 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM162 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2217] An enzyme complex designated DICER COMPLEX, dices the GAM162 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM162 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 23%) nucleotide sequence of GAM162 RNA is designated SEQ ID:2313, and is provided hereinbelow with reference to the sequence listing part.

[2218] GAM162 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM162 target RNA, herein designated GAM TARGET RNA.

GAM162 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2219] GAM162 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM162 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM162 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM162 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM162 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2220] The complementary binding of GAM162 RNA, herein designated GAM RNA, to target binding sites on GAM162 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM162 target RNA, herein designated GAM TARGET RNA, into GAM162 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2221] It is appreciated that GAM162 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM162 target genes. The mRNA of each one of this plurality of GAM162 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM162 RNA, herein designated GAM RNA, and which when bound by GAM162 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM162 target proteins.

[2222] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM162 gene, herein designated GAM GENE, on one or more GAM162 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2223] It is yet further appreciated that specific functions, and accordingly utilities, of GAM162 correlate with, and may be deduced from, the identity of the target genes which GAM162 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2224] Nucleotide sequences of the GAM162 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM162 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM162 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM162 are further described hereinbelow with reference to Table Table1.

[2225] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM162 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2226]

[2227] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 163 (GAM163) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2228] GAM163 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM163 was detected is described hereinabove with reference to Figs. 2-8.

[2229] GAM163 gene, herein designated GAM GENE, and GAM163 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[2230] GAM163 gene, herein designated GAM GENE, encodes a GAM163 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM163 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM163 precursor RNA is designated SEQ ID:2328, and is provided hereinbelow with reference to the sequence listing part.

[2231] GAM163 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM163 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2232] An enzyme complex designated DICER COMPLEX, dices the GAM163 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM163 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 42%) nucleotide sequence of GAM163 RNA is designated SEQ ID:2329, and is provided hereinbelow with reference to the sequence listing part.

[2233] GAM163 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM163 target RNA, herein designated GAM TARGET RNA. GAM163 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2234] GAM163 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM163 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM163 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM163 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM163 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2235] The complementary binding of GAM163 RNA, herein designated GAM RNA, to target binding sites on GAM163 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM163 target RNA, herein designated GAM TARGET RNA, into GAM163 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2236] It is appreciated that GAM163 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM163 target genes. The mRNA of each one of this plurality of GAM163 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM163 RNA, herein designated GAM RNA, and which when bound by GAM163 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM163 target proteins.

[2237] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM163 gene, herein designated GAM GENE, on one or more GAM163 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[2238] It is yet further appreciated that specific functions, and accordingly utilities, of GAM163 correlate with, and may be deduced from, the identity of the target genes which GAM163 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2239] Nucleotide sequences of the GAM163 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM163 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM163 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM163 are further described hereinbelow with reference to Table Table1.

[2240] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM163 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2241]

[2242] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

164 (GAM164) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2243] GAM164 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM164 was detected is described hereinabove with reference to Figs. 2-8.

[2244] GAM164 gene, herein designated GAM GENE, and GAM164 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2245] GAM164 gene, herein designated GAM GENE, encodes a GAM164 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM164 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM164 precursor RNA is designated SEQ ID:2345, and is provided hereinbelow with reference to the sequence listing part.

[2246] GAM164 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM164 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2247] An enzyme complex designated DICER COMPLEX, dices the GAM164 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM164 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM164 RNA is designated SEQ ID:2346, and is provided hereinbelow with reference to the sequence listing part.

[2248] GAM164 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM164 target RNA, herein designated GAM TARGET RNA. GAM164 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2249] GAM164 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM164 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM164 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM164 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM164 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2250] The complementary binding of GAM164 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM164 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM164 target RNA, herein designated GAM TARGET RNA, into GAM164 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2251] It is appreciated that GAM164 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM164 target genes. The mRNA of each one of this plurality of GAM164 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM164 RNA, herein designated GAM RNA, and which when bound by GAM164 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM164 target proteins.

[2252] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM164 gene, herein designated GAM GENE, on one or more GAM164 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2253] It is yet further appreciated that specific functions, and accordingly utilities, of GAM164 correlate with, and may be deduced from, the identity of the target genes which GAM164 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2254] Nucleotide sequences of the GAM164 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM164 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM164 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM164 are further described hereinbelow with reference to Table Table1.

[2255] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM164 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2256]

[2257] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 165 (GAM165) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2258] GAM165 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM165 was detected is described hereinabove with reference to Figs. 2-8.

[2259] GAM165 gene, herein designated GAM GENE, and GAM165 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2260] GAM165 gene, herein designated GAM GENE, encodes a GAM165 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM165 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM165 precursor RNA is designated SEQ ID:2363, and is provided hereinbelow with reference to the sequence listing part.

[2261] GAM165 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM165 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2262] An enzyme complex designated DICER COMPLEX, dices the GAM165 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM165 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 64%) nucleotide sequence of GAM165 RNA is designated SEQ ID:2364, and is provided hereinbelow with reference to the sequence listing part.

[2263] GAM165 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM165 target RNA, herein designated GAM TARGET RNA. GAM165 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2264] GAM165 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM165 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM165 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM165 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM165 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2265] The complementary binding of GAM165 RNA, herein designated GAM RNA, to target binding sites on GAM165 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM165 target RNA, herein designated GAM TARGET RNA, into GAM165 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2266] It is appreciated that GAM165 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM165 target genes. The mRNA of each one of this plurality of GAM165 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM165 RNA, herein designated GAM RNA, and which when bound by

GAM165 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM165 target proteins.

[2267] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM165 gene, herein designated GAM GENE, on one or more GAM165 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2268] It is yet further appreciated that specific functions, and accordingly utilities, of GAM165 correlate with, and may be deduced from, the identity of the target genes which GAM165 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[2269] Nucleotide sequences of the GAM165 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM165 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM165 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM165 are further described hereinbelow with reference to Table Table1.

[2270] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM165 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2271]

[2272] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 166 (GAM166) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2273] GAM166 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM166 was detected is described hereinabove with reference to Figs. 2-8.

[2274] GAM166 gene, herein designated GAM GENE, and GAM166 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2275] GAM166 gene, herein designated GAM GENE, encodes a GAM166 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM166 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM166 precursor RNA is designated SEQ ID:2389, and is provided hereinbelow with reference to the sequence listing part.

[2276] GAM166 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM166 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[2277] An enzyme complex designated DICER COMPLEX, dices the GAM166 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM166 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 19%) nucleotide sequence of GAM166 RNA is designated SEQ ID:2390, and is provided hereinbelow with reference to the sequence listing part.

[2278] GAM166 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM166 target RNA, herein designated GAM TARGET RNA. GAM166 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2279] GAM166 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM166 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM166 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM166 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM166 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2280] The complementary binding of GAM166 RNA, herein designated GAM RNA, to target binding sites on GAM166 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM166 target RNA, herein designated GAM TARGET RNA, into GAM166 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2281] It is appreciated that GAM166 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM166 target genes. The mRNA of each one of this plurality of GAM166 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM166 RNA, herein designated GAM RNA, and which when bound by GAM166 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM166 target proteins.

[2282] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM166 gene, herein designated GAM GENE, on one or more GAM166 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2283] It is yet further appreciated that specific functions, and accordingly utilities, of GAM166 correlate with, and may be deduced from, the identity of the target genes which GAM166 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2284] Nucleotide sequences of the GAM166 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM166 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM166 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM166 are further described hereinbelow with reference to Table Table1.

[2285] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM166 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2286]

[2287] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 167 (GAM167) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2288] GAM167 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM167 was detected is described hereinabove with reference to Figs. 2-8.

[2289] GAM167 gene, herein designated GAM GENE, and GAM167 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2290] GAM167 gene, herein designated GAM GENE, encodes a GAM167 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM167 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM167 precursor RNA is designated SEQ ID:2400, and is provided hereinbelow with reference to the sequence listing part.

[2291] GAM167 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM167 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2292] An enzyme complex designated DICER COMPLEX, dices the GAM167 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM167 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 41%) nucleotide sequence of GAM167 RNA is designated SEQ ID:2401, and is provided hereinbelow with reference to the sequence listing part.

[2293] GAM167 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM167

target RNA, herein designated GAM TARGET RNA.

GAM167 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2294] GAM167 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM167 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM167 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM167 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM167 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2295] The complementary binding of GAM167 RNA, herein designated GAM RNA, to target binding sites on GAM167 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM167 target RNA, herein designated GAM TARGET RNA, into GAM167 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2296] It is appreciated that GAM167 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM167 target genes. The mRNA of each one of this plurality of GAM167 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM167 RNA, herein designated GAM RNA, and which when bound by GAM167 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM167 target proteins.

[2297] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM167 gene, herein designated GAM GENE, on one or more GAM167 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2298] It is yet further appreciated that specific functions, and accordingly utilities, of GAM167 correlate with, and may be deduced from, the identity of the target genes which GAM167 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2299] Nucleotide sequences of the GAM167 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM167 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM167 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM167 are further described hereinbelow with reference to Table Table1.

[2300] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM167 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2301]

[2302] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 168 (GAM168) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2303] GAM168 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM168 was detected is described hereinabove with reference to Figs. 2-8.

[2304] GAM168 gene, herein designated GAM GENE, and GAM168 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2305] GAM168 gene, herein designated GAM GENE, encodes a GAM168 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM168 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM168 precursor RNA is designated SEQ ID:2422, and is provided hereinbelow with reference to the sequence listing part.

[2306] GAM168 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM168 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2307] An enzyme complex designated DICER COMPLEX, dices the GAM168 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM168 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM168 RNA is designated SEQ ID:2423, and is provided hereinbelow with reference to the sequence listing part.

[2308] GAM168 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM168 target RNA, herein designated GAM TARGET RNA. GAM168 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2309] GAM168 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM168 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM168 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM168 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM168 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2310] The complementary binding of GAM168 RNA, herein designated GAM RNA, to target binding sites on GAM168 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM168 target RNA, herein designated GAM TARGET RNA, into GAM168 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2311] It is appreciated that GAM168 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM168 target genes. The mRNA of each one of this plu-

rality of GAM168 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM168 RNA, herein designated GAM RNA, and which when bound by GAM168 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM168 target proteins.

[2312] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM168 gene, herein designated GAM GENE, on one or more GAM168 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2313] It is yet further appreciated that specific functions, and accordingly utilities, of GAM168 correlate with, and may be deduced from, the identity of the target genes which GAM168 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2314] Nucleotide sequences of the GAM168 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM168 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM168 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM168 are further described hereinbelow with reference to Table Table1.

[2315] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM168 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2316]

[2317] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 169 (GAM169) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[2318] GAM169 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM169 was detected is described hereinabove with reference to Figs. 2-8.

[2319] GAM169 gene, herein designated GAM GENE, and GAM169 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2320] GAM169 gene, herein designated GAM GENE, encodes a GAM169 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM169 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM169 precursor RNA is designated SEQ ID:2434, and is provided hereinbelow with reference to the sequence listing part.

[2321] GAM169 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM169 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2322] An enzyme complex designated DICER COMPLEX, dices the GAM169 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM169 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 82%) nucleotide sequence of GAM169 RNA is designated SEQ ID:2435, and is provided hereinbelow with reference to the sequence listing part.

[2323] GAM169 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM169 target RNA, herein designated GAM TARGET RNA.

GAM169 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[2324] GAM169 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM169 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM169 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM169 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM169 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2325] The complementary binding of GAM169 RNA, herein designated GAM RNA, to target binding sites on GAM169 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM169 target RNA, herein designated GAM TARGET RNA, into GAM169 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2326] It is appreciated that GAM169 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM169 target genes. The mRNA of each one of this plurality of GAM169 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM169 RNA, herein designated GAM RNA, and which when bound by GAM169 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM169 target proteins.

[2327] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM169 gene, herein designated GAM GENE, on one or more GAM169 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2328] It is yet further appreciated that specific functions, and accordingly utilities, of GAM169 correlate with, and may be deduced from, the identity of the target genes which GAM169 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2329] Nucleotide sequences of the GAM169 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM169 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM169 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM169 are further described hereinbelow with reference to Table Table1.

[2330] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM169 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2331]

[2332] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 170 (GAM170) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2333] GAM170 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM170 was detected is described hereinabove with reference to Figs. 2-8.

[2334] GAM170 gene, herein designated GAM GENE, and GAM170 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2335] GAM170 gene, herein designated GAM GENE, encodes a GAM170 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM170 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM170 precursor RNA is designated SEQ ID:2448, and is provided hereinbelow with reference to the sequence listing part.

[2336] GAM170 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM170 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2337] An enzyme complex designated DICER COMPLEX, dices the GAM170 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM170 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of

GAM170 RNA is designated SEQ ID:2449, and is provided hereinbelow with reference to the sequence listing part.

[2338] GAM170 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM170 target RNA, herein designated GAM TARGET RNA. GAM170 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2339] GAM170 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM170 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM170 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM170 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM170 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2340] The complementary binding of GAM170 RNA, herein designated GAM RNA, to target binding sites on GAM170 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM170 target RNA, herein designated GAM TARGET RNA, into GAM170 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2341] It is appreciated that GAM170 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM170 target genes. The mRNA of each one of this plurality of GAM170 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM170 RNA, herein designated GAM RNA, and which when bound by GAM170 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM170 target proteins.

[2342] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM170 gene, herein designated GAM GENE, on one or more GAM170 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2343] It is yet further appreciated that specific functions, and accordingly utilities, of GAM170 correlate with, and may be deduced from, the identity of the target genes which GAM170 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2344] Nucleotide sequences of the GAM170 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM170 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM170 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM170 are further described hereinbelow with reference to Table Table1.

[2345] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM170 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2346]

[2347] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 171 (GAM171) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2348] GAM171 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM171 was detected is described hereinabove

with reference to Figs. 2-8.

[2349] GAM171 gene, herein designated GAM GENE, and GAM171 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2350] GAM171 gene, herein designated GAM GENE, encodes a GAM171 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM171 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM171 precursor RNA is designated SEQ ID:2458, and is provided hereinbelow with reference to the sequence listing part.

[2351] GAM171 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM171 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2352] An enzyme complex designated DICER COMPLEX, dices the GAM171 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM171 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM171 RNA is designated SEQ ID:2459, and is provided hereinbelow with reference to the sequence listing part.

[2353] GAM171 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM171 target RNA, herein designated GAM TARGET RNA. GAM171 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2354] GAM171 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM171 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM171 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM171 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM171 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2355] The complementary binding of GAM171 RNA, herein designated GAM RNA, to target binding sites on GAM171 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM171 target RNA, herein designated GAM TARGET RNA, into GAM171 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[2356] It is appreciated that GAM171 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM171 target genes. The mRNA of each one of this plurality of GAM171 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM171 RNA, herein designated GAM RNA, and which when bound by GAM171 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM171 target proteins.

[2357] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM171 gene, herein designated GAM GENE, on one or more GAM171 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2358] It is yet further appreciated that specific functions, and accordingly utilities, of GAM171 correlate with, and may be deduced from, the identity of the target genes which GAM171 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2359] Nucleotide sequences of the GAM171 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM171 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM171 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM171 are further described hereinbelow with reference to Table Table1.

[2360] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM171 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2361]

[2362] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 172 (GAM172) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2363] GAM172 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM172 was detected is described hereinabove with reference to Figs. 2-8.

[2364] GAM172 gene, herein designated GAM GENE, and GAM172 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2365] GAM172 gene, herein designated GAM GENE, encodes a GAM172 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM172 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM172 precursor RNA is designated SEQ ID:2477, and is provided hereinbelow with reference to the sequence listing part.

[2366] GAM172 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM172 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2367] An enzyme complex designated DICER COMPLEX, dices the GAM172 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM172 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM172 RNA is designated SEQ ID:2478, and is provided hereinbelow with reference to the sequence listing part.

[2368] GAM172 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM172 target RNA, herein designated GAM TARGET RNA.

GAM172 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2369] GAM172 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM172 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM172 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM172 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM172 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2370] The complementary binding of GAM172 RNA, herein designated GAM RNA, to target binding sites on GAM172 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM172 target RNA, herein designated GAM TARGET RNA, into GAM172 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2371] It is appreciated that GAM172 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM172 target genes. The mRNA of each one of this plurality of GAM172 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM172 RNA, herein designated GAM RNA, and which when bound by GAM172 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM172 target proteins.

[2372] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM172 gene, herein designated GAM GENE, on one or more GAM172 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2373] It is yet further appreciated that specific functions, and accordingly utilities, of GAM172 correlate with, and may be deduced from, the identity of the target genes which GAM172 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2374] Nucleotide sequences of the GAM172 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM172 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM172 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM172 are further described hereinbelow with reference to Table Table1.

[2375] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM172 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2376]

[2377] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 173 (GAM173) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2378] GAM173 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM173 was detected is described hereinabove with reference to Figs. 2-8.

[2379] GAM173 gene, herein designated GAM GENE, and GAM173 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2380] GAM173 gene, herein designated GAM GENE, encodes a

GAM173 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM173 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM173 precursor RNA is designated SEQ ID:2516, and is provided hereinbelow with reference to the sequence listing part.

[2381] GAM173 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM173 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2382] An enzyme complex designated DICER COMPLEX, dices the GAM173 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM173 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM173 RNA is designated SEQ ID:2517, and is provided hereinbelow with reference to the sequence listing part.

[2383] GAM173 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM173 target RNA, herein designated GAM TARGET RNA. GAM173 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2384] GAM173 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM173 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM173 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM173 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM173 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2385] The complementary binding of GAM173 RNA, herein designated GAM RNA, to target binding sites on GAM173 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM173 target RNA, herein designated GAM TARGET RNA, into GAM173 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2386] It is appreciated that GAM173 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM173 target genes. The mRNA of each one of this plurality of GAM173 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM173 RNA, herein designated GAM RNA, and which when bound by GAM173 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM173 target proteins.

[2387] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM173 gene, herein designated GAM GENE, on one or more GAM173 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2388] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM173 correlate with, and may be deduced from, the identity of the target genes which GAM173 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2389] Nucleotide sequences of the GAM173 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM173 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM173 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM173 are further described hereinbelow with reference to Table Table1.

[2390] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM173 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2391]

[2392] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 174 (GAM174) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[2393] GAM174 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM174 was detected is described hereinabove with reference to Figs. 2-8.

[2394] GAM174 gene, herein designated GAM GENE, and GAM174 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2395] GAM174 gene, herein designated GAM GENE, encodes a GAM174 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM174 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM174 precursor RNA is designated SEQ ID:2524, and is provided hereinbelow with reference to the sequence listing part.

[2396] GAM174 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM174 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2397] An enzyme complex designated DICER COMPLEX, dices the GAM174 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM174 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM174 RNA is designated SEQ ID:2525, and is provided hereinbelow with reference to the sequence listing part.

[2398] GAM174 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM174 target RNA, herein designated GAM TARGET RNA. GAM174 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2399] GAM174 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM174 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM174 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM174 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM174 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2400] The complementary binding of GAM174 RNA, herein designated GAM RNA, to target binding sites on GAM174 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM174 target RNA, herein designated GAM TARGET RNA, into GAM174 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2401] It is appreciated that GAM174 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM174 target genes. The mRNA of each one of this plurality of GAM174 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM174 RNA, herein designated GAM RNA, and which when bound by GAM174 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM174 target proteins.

[2402] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM174 gene, herein designated GAM GENE, on one or more GAM174 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2403] It is yet further appreciated that specific functions, and accordingly utilities, of GAM174 correlate with, and may be deduced from, the identity of the target genes which GAM174 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2404] Nucleotide sequences of the GAM174 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM174 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM174 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM174 are further described hereinbelow with reference to Table Table1.

[2405] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM174 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2406]

[2407] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 175 (GAM175) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2408] GAM175 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM175 was detected is described hereinabove with reference to Figs. 2-8.

[2409] GAM175 gene, herein designated GAM GENE, and GAM175 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2410] GAM175 gene, herein designated GAM GENE, encodes a GAM175 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM175 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM175 precursor RNA is designated SEQ ID:2532, and is provided hereinbelow with reference to the sequence listing part.

[2411] GAM175 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM175 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2412] An enzyme complex designated DICER COMPLEX, dices the GAM175 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM175 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM175 RNA is designated SEQ ID:2533, and is provided

hereinbelow with reference to the sequence listing part.

[2413] GAM175 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM175 target RNA, herein designated GAM TARGET RNA.
GAM175 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2414] GAM175 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM175 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM175 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM175 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM175 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2415] The complementary binding of GAM175 RNA, herein designated GAM RNA, to target binding sites on GAM175 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM175 target RNA, herein designated GAM TARGET RNA, into GAM175 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2416] It is appreciated that GAM175 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM175 target genes. The mRNA of each one of this plurality of GAM175 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM175 RNA, herein designated GAM RNA, and which when bound by GAM175 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM175 tar-

get proteins.

[2417] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM175 gene, herein designated GAM GENE, on one or more GAM175 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2418] It is yet further appreciated that specific functions, and accordingly utilities, of GAM175 correlate with, and may be deduced from, the identity of the target genes which GAM175 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2419] Nucleotide sequences of the GAM175 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM175 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM175 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM175 are further described hereinbelow with reference to Table Table1.

[2420] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM175 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2421]

[2422] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 176 (GAM176) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2423] GAM176 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM176 was detected is described hereinabove with reference to Figs. 2-8.

[2424] GAM176 gene, herein designated GAM GENE, and GAM176 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2425] GAM176 gene, herein designated GAM GENE, encodes a GAM176 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM176 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM176 precursor RNA is designated SEQ ID:2540, and is provided hereinbelow with reference to the sequence listing part.

[2426] GAM176 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM176 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2427] An enzyme complex designated DICER COMPLEX, dices

the GAM176 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM176 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM176 RNA is designated SEQ ID:2541, and is provided hereinbelow with reference to the sequence listing part.

[2428] GAM176 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM176 target RNA, herein designated GAM TARGET RNA. GAM176 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2429] GAM176 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM176 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM176

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM176 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM176 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2430] The complementary binding of GAM176 RNA, herein designated GAM RNA, to target binding sites on GAM176 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM176 target RNA, herein designated GAM TARGET RNA, into GAM176 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2431] It is appreciated that GAM176 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM176 target genes. The mRNA of each one of this plurality of GAM176 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM176 RNA, herein designated GAM RNA, and which when bound by GAM176 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM176 target proteins.

[2432] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM176 gene, herein designated GAM GENE, on one or more GAM176 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2433] It is yet further appreciated that specific functions, and accordingly utilities, of GAM176 correlate with, and may be deduced from, the identity of the target genes which GAM176 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2434] Nucleotide sequences of the GAM176 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM176 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM176 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM176 are further described hereinbelow with reference to Table Table1.

[2435] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM176 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2436]

[2437] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 177 (GAM177) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2438] GAM177 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM177 was detected is described hereinabove with reference to Figs. 2-8.

[2439] GAM177 gene, herein designated GAM GENE, and GAM177 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2440] GAM177 gene, herein designated GAM GENE, encodes a GAM177 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM177 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM177 precursor RNA is designated SEQ ID:2564, and is provided hereinbelow with reference to the sequence listing part.

[2441] GAM177 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM177 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2442] An enzyme complex designated DICER COMPLEX, dices the GAM177 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM177 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 35%) nucleotide sequence of GAM177 RNA is designated SEQ ID:2565, and is provided hereinbelow with reference to the sequence listing part.

[2443] GAM177 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM177 target RNA, herein designated GAM TARGET RNA. GAM177 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2444] GAM177 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM177 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM177 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM177 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM177 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[2445] The complementary binding of GAM177 RNA, herein designated GAM RNA, to target binding sites on GAM177 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM177 target RNA, herein designated GAM TARGET RNA, into GAM177 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2446] It is appreciated that GAM177 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM177 target genes. The mRNA of each one of this plurality of GAM177 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM177 RNA, herein designated GAM RNA, and which when bound by GAM177 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM177 target proteins.

[2447] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM177 gene, herein designated GAM GENE, on one or

more GAM177 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2448] It is yet further appreciated that specific functions, and accordingly utilities, of GAM177 correlate with, and may be deduced from, the identity of the target genes which GAM177 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2449] Nucleotide sequences of the GAM177 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM177 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM177 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM177 are further de-

scribed hereinbelow with reference to Table Table1.

[2450] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM177 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2451]

[2452] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 178 (GAM178) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2453] GAM178 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM178 was detected is described hereinabove with reference to Figs. 2-8.

[2454] GAM178 gene, herein designated GAM GENE, and GAM178 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2455] GAM178 gene, herein designated GAM GENE, encodes a GAM178 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM178 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM178 precursor RNA is designated SEQ ID:2583, and is provided hereinbelow with reference to the sequence listing part.

[2456] GAM178 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM178 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2457] An enzyme complex designated DICER COMPLEX, dices the GAM178 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM178 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM178 RNA is designated SEQ ID:2584, and is provided hereinbelow with reference to the sequence listing part.

[2458] GAM178 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM178 target RNA, herein designated GAM TARGET RNA. GAM178 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2459] GAM178 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM178 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM178 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM178 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM178 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2460] The complementary binding of GAM178 RNA, herein designated GAM RNA, to target binding sites on GAM178 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM178 target RNA, herein designated GAM TARGET RNA, into GAM178 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2461] It is appreciated that GAM178 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM178 target genes. The mRNA of each one of this plurality of GAM178 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM178 RNA, herein designated GAM RNA, and which when bound by GAM178 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM178 target proteins.

[2462] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM178 gene, herein designated GAM GENE, on one or more GAM178 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2463] It is yet further appreciated that specific functions, and accordingly utilities, of GAM178 correlate with, and may

be deduced from, the identity of the target genes which GAM178 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2464] Nucleotide sequences of the GAM178 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM178 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM178 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM178 are further described hereinbelow with reference to Table Table1.

[2465] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM178 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2466]

[2467] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 179 (GAM179) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2468] GAM179 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM179 was detected is described hereinabove with reference to Figs. 2-8.

[2469] GAM179 gene, herein designated GAM GENE, and GAM179 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2470] GAM179 gene, herein designated GAM GENE, encodes a GAM179 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM179 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM179 precursor RNA is designated SEQ ID:2595, and is provided hereinbelow with reference to the sequence listing part.

[2471] GAM179 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM179 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2472] An enzyme complex designated DICER COMPLEX, dices the GAM179 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM179 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM179 RNA is designated SEQ ID:2596, and is provided hereinbelow with reference to the sequence listing part.

[2473] GAM179 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM179 target RNA, herein designated GAM TARGET RNA. GAM179 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2474] GAM179 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM179 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM179 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM179 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM179 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2475] The complementary binding of GAM179 RNA, herein designated GAM RNA, to target binding sites on GAM179 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM179 target RNA, herein designated GAM TARGET RNA, into GAM179 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2476] It is appreciated that GAM179 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM179 target genes. The mRNA of each one of this plurality of GAM179 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM179 RNA, herein designated GAM RNA, and which when bound by GAM179 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM179 target proteins.

[2477] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM179 gene, herein designated GAM GENE, on one or more GAM179 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2478] It is yet further appreciated that specific functions, and accordingly utilities, of GAM179 correlate with, and may be deduced from, the identity of the target genes which GAM179 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2479] Nucleotide sequences of the GAM179 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM179 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM179 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM179 are further described hereinbelow with reference to Table Table1.

[2480] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM179 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2481]

[2482] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 180 (GAM180) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2483] GAM180 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM180 was detected is described hereinabove with reference to Figs. 2-8.

[2484] GAM180 gene, herein designated GAM GENE, and GAM180 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2485] GAM180 gene, herein designated GAM GENE, encodes a GAM180 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM180 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM180 precursor RNA is desig-

nated SEQ ID:2600, and is provided hereinbelow with reference to the sequence listing part.

[2486] GAM180 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM180 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2487] An enzyme complex designated DICER COMPLEX, dices the GAM180 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM180 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 46%) nucleotide sequence of GAM180 RNA is designated SEQ ID:2601, and is provided hereinbelow with reference to the sequence listing part.

[2488] GAM180 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM180 target RNA, herein designated GAM TARGET RNA. GAM180 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2489] GAM180 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM180 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM180 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM180 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM180

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2490] The complementary binding of GAM180 RNA, herein designated GAM RNA, to target binding sites on GAM180 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM180 target RNA, herein designated GAM TARGET RNA, into GAM180 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2491] It is appreciated that GAM180 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM180 target genes. The mRNA of each one of this plurality of GAM180 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM180 RNA, herein designated GAM RNA, and which when bound by GAM180 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM180 target proteins.

[2492] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM180 gene, herein designated GAM GENE, on one or more GAM180 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2493] It is yet further appreciated that specific functions, and accordingly utilities, of GAM180 correlate with, and may be deduced from, the identity of the target genes which GAM180 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2494] Nucleotide sequences of the GAM180 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM180 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM180 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM180 are further described hereinbelow with reference to Table Table1.

[2495] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM180 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2496]

[2497] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 181 (GAM181) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2498] GAM181 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM181 was detected is described hereinabove with reference to Figs. 2-8.

[2499] GAM181 gene, herein designated GAM GENE, and

GAM181 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2500] GAM181 gene, herein designated GAM GENE, encodes a GAM181 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM181 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM181 precursor RNA is designated SEQ ID:2619, and is provided hereinbelow with reference to the sequence listing part.

[2501] GAM181 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM181 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2502] An enzyme complex designated DICER COMPLEX, dices the GAM181 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM181 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM181 RNA is designated SEQ ID:2620, and is provided hereinbelow with reference to the sequence listing part.

[2503] GAM181 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM181 target RNA, herein designated GAM TARGET RNA. GAM181 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2504] GAM181 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM181 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM181 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM181 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM181 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2505] The complementary binding of GAM181 RNA, herein designated GAM RNA, to target binding sites on GAM181 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM181 target RNA, herein designated GAM TARGET RNA, into GAM181 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2506] It is appreciated that GAM181 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM181 target genes. The mRNA of each one of this plurality of GAM181 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM181 RNA, herein designated GAM RNA, and which when bound by GAM181 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM181 target proteins.

[2507] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM181 gene, herein designated GAM GENE, on one or more GAM181 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2508] It is yet further appreciated that specific functions, and accordingly utilities, of GAM181 correlate with, and may be deduced from, the identity of the target genes which GAM181 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2509] Nucleotide sequences of the GAM181 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM181 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM181 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM181 are further described hereinbelow with reference to Table Table1.

[2510] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM181 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2511]

[2512] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 182 (GAM182) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2513] GAM182 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM182 was detected is described hereinabove with reference to Figs. 2-8.

[2514] GAM182 gene, herein designated GAM GENE, and GAM182 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2515] GAM182 gene, herein designated GAM GENE, encodes a GAM182 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM182 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM182 precursor RNA is designated SEQ ID:2643, and is provided hereinbelow with reference to the sequence listing part.

[2516] GAM182 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM182 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2517] An enzyme complex designated DICER COMPLEX, dices the GAM182 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM182 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM182 RNA is designated SEQ ID:2644, and is provided hereinbelow with reference to the sequence listing part.

[2518] GAM182 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM182 target RNA, herein designated GAM TARGET RNA. GAM182 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2519] GAM182 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM182 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM182 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM182 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM182 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2520] The complementary binding of GAM182 RNA, herein designated GAM RNA, to target binding sites on GAM182 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM182 target RNA, herein designated GAM TARGET RNA, into GAM182 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2521] It is appreciated that GAM182 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM182 target genes. The mRNA of each one of this plurality of GAM182 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM182 RNA, herein designated GAM RNA, and which when bound by GAM182 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM182 target proteins.

[2522] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM182 gene, herein designated GAM GENE, on one or more GAM182 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2523] It is yet further appreciated that specific functions, and accordingly utilities, of GAM182 correlate with, and may be deduced from, the identity of the target genes which GAM182 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2524] Nucleotide sequences of the GAM182 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM182 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM182 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM182 are further described hereinbelow with reference to Table Table1.

[2525] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM182 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2526]

[2527] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 183 (GAM183) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2528] GAM183 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM183 was detected is described hereinabove with reference to Figs. 2-8.

[2529] GAM183 gene, herein designated GAM GENE, and GAM183 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2530] GAM183 gene, herein designated GAM GENE, encodes a GAM183 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM183 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM183 precursor RNA is designated SEQ ID:2666, and is provided hereinbelow with reference to the sequence listing part.

[2531] GAM183 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM183 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2532] An enzyme complex designated DICER COMPLEX, dices the GAM183 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM183 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 32%) nucleotide sequence of GAM183 RNA is designated SEQ ID:2667, and is provided hereinbelow with reference to the sequence listing part.

[2533] GAM183 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM183 target RNA, herein designated GAM TARGET RNA. GAM183 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2534] GAM183 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM183 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM183 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM183 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM183 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2535] The complementary binding of GAM183 RNA, herein designated GAM RNA, to target binding sites on GAM183 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM183 target RNA, herein designated GAM TARGET RNA, into GAM183 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2536] It is appreciated that GAM183 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM183 target genes. The mRNA of each one of this plurality of GAM183 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM183 RNA,

herein designated GAM RNA, and which when bound by GAM183 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM183 target proteins.

[2537] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM183 gene, herein designated GAM GENE, on one or more GAM183 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2538] It is yet further appreciated that specific functions, and accordingly utilities, of GAM183 correlate with, and may be deduced from, the identity of the target genes which

GAM183 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2539] Nucleotide sequences of the GAM183 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM183 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM183 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM183 are further described hereinbelow with reference to Table Table1.

[2540] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM183 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2541]

[2542] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 184 (GAM184) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2543] GAM184 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM184 was detected is described hereinabove with reference to Figs. 2-8.

[2544] GAM184 gene, herein designated GAM GENE, and GAM184 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2545] GAM184 gene, herein designated GAM GENE, encodes a GAM184 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM184 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM184 precursor RNA is designated SEQ ID:2682, and is provided hereinbelow with reference to the sequence listing part.

[2546] GAM184 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM184 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[2547] An enzyme complex designated DICER COMPLEX, dices the GAM184 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM184 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM184 RNA is designated SEQ ID:2683, and is provided hereinbelow with reference to the sequence listing part.

[2548] GAM184 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM184 target RNA, herein designated GAM TARGET RNA.
GAM184 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2549] GAM184 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM184 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM184 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM184 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM184 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2550] The complementary binding of GAM184 RNA, herein designated GAM RNA, to target binding sites on GAM184 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM184 target RNA, herein desig-

nated GAM TARGET RNA, into GAM184 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2551] It is appreciated that GAM184 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM184 target genes. The mRNA of each one of this plurality of GAM184 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM184 RNA, herein designated GAM RNA, and which when bound by GAM184 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM184 target proteins.

[2552] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM184 gene, herein designated GAM GENE, on one or more GAM184 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2553] It is yet further appreciated that specific functions, and accordingly utilities, of GAM184 correlate with, and may be deduced from, the identity of the target genes which GAM184 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2554] Nucleotide sequences of the GAM184 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM184 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM184 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM184 are further described hereinbelow with reference to Table Table1.

[2555] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM184 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[2556]

[2557] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 185 (GAM185) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2558] GAM185 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM185 was detected is described hereinabove with reference to Figs. 2-8.

[2559] GAM185 gene, herein designated GAM GENE, and GAM185 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2560] GAM185 gene, herein designated GAM GENE, encodes a GAM185 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM185 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM185 precursor RNA is designated SEQ ID:2692, and is provided hereinbelow with ref-

erence to the sequence listing part.

[2561] GAM185 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM185 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2562] An enzyme complex designated DICER COMPLEX, dices the GAM185 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM185 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM185 RNA is designated SEQ ID:2693, and is provided hereinbelow with reference to the sequence listing part.

[2563] GAM185 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM185 target RNA, herein designated GAM TARGET RNA.

GAM185 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2564] GAM185 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM185 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM185 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM185 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM185 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2565] The complementary binding of GAM185 RNA, herein designated GAM RNA, to target binding sites on GAM185 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM185 target RNA, herein designated GAM TARGET RNA, into GAM185 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2566] It is appreciated that GAM185 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM185 target genes. The mRNA of each one of this plurality of GAM185 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM185 RNA, herein designated GAM RNA, and which when bound by GAM185 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM185 target proteins.

[2567] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM185 gene, herein designated GAM GENE, on one or more GAM185 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2568] It is yet further appreciated that specific functions, and accordingly utilities, of GAM185 correlate with, and may be deduced from, the identity of the target genes which GAM185 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2569] Nucleotide sequences of the GAM185 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM185 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM185 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM185 are further described hereinbelow with reference to Table Table1.

[2570] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM185 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2571]

[2572] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 186 (GAM186) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2573] GAM186 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM186 was detected is described hereinabove with reference to Figs. 2-8.

[2574] GAM186 gene, herein designated GAM GENE, and GAM186 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[2575] GAM186 gene, herein designated GAM GENE, encodes a GAM186 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM186 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM186 precursor RNA is designated SEQ ID:2706, and is provided hereinbelow with reference to the sequence listing part.

[2576] GAM186 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM186 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2577] An enzyme complex designated DICER COMPLEX, dices the GAM186 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM186 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM186 RNA is designated SEQ ID:2707, and is provided hereinbelow with reference to the sequence listing part.

[2578] GAM186 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM186 target RNA, herein designated GAM TARGET RNA. GAM186 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2579] GAM186 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM186 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM186 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM186 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM186 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2580] The complementary binding of GAM186 RNA, herein designated GAM RNA, to target binding sites on GAM186 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM186 target RNA, herein designated GAM TARGET RNA, into GAM186 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2581] It is appreciated that GAM186 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM186 target genes. The mRNA of each one of this plurality of GAM186 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM186 RNA, herein designated GAM RNA, and which when bound by GAM186 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM186 target proteins.

[2582] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM186 gene, herein designated GAM GENE, on one or more GAM186 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[2583] It is yet further appreciated that specific functions, and accordingly utilities, of GAM186 correlate with, and may be deduced from, the identity of the target genes which GAM186 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2584] Nucleotide sequences of the GAM186 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM186 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM186 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM186 are further described hereinbelow with reference to Table Table1.

[2585] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM186 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2586]

[2587] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

187 (GAM187) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2588] GAM187 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM187 was detected is described hereinabove with reference to Figs. 2-8.

[2589] GAM187 gene, herein designated GAM GENE, and GAM187 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2590] GAM187 gene, herein designated GAM GENE, encodes a GAM187 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM187 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM187 precursor RNA is designated SEQ ID:2715, and is provided hereinbelow with reference to the sequence listing part.

[2591] GAM187 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM187 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2592] An enzyme complex designated DICER COMPLEX, dices the GAM187 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM187 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM187 RNA is designated SEQ ID:2716, and is provided hereinbelow with reference to the sequence listing part.

[2593] GAM187 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM187 target RNA, herein designated GAM TARGET RNA. GAM187 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2594] GAM187 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM187 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM187 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM187 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM187 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2595] The complementary binding of GAM187 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM187 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM187 target RNA, herein designated GAM TARGET RNA, into GAM187 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2596] It is appreciated that GAM187 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM187 target genes. The mRNA of each one of this plurality of GAM187 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM187 RNA, herein designated GAM RNA, and which when bound by GAM187 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM187 target proteins.

[2597] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM187 gene, herein designated GAM GENE, on one or more GAM187 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2598] It is yet further appreciated that specific functions, and accordingly utilities, of GAM187 correlate with, and may be deduced from, the identity of the target genes which GAM187 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2599] Nucleotide sequences of the GAM187 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM187 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM187 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM187 are further described hereinbelow with reference to Table Table1.

[2600] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM187 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2601]

[2602] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 188 (GAM188) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2603] GAM188 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM188 was detected is described hereinabove with reference to Figs. 2-8.

[2604] GAM188 gene, herein designated GAM GENE, and GAM188 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2605] GAM188 gene, herein designated GAM GENE, encodes a GAM188 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM188 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM188 precursor RNA is designated SEQ ID:2734, and is provided hereinbelow with reference to the sequence listing part.

[2606] GAM188 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM188 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2607] An enzyme complex designated DICER COMPLEX, dices the GAM188 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM188 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 61%) nucleotide sequence of GAM188 RNA is designated SEQ ID:2735, and is provided hereinbelow with reference to the sequence listing part.

[2608] GAM188 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM188 target RNA, herein designated GAM TARGET RNA. GAM188 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2609] GAM188 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM188 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM188 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM188 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM188 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2610] The complementary binding of GAM188 RNA, herein designated GAM RNA, to target binding sites on GAM188 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM188 target RNA, herein designated GAM TARGET RNA, into GAM188 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2611] It is appreciated that GAM188 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM188 target genes. The mRNA of each one of this plurality of GAM188 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM188 RNA, herein designated GAM RNA, and which when bound by

GAM188 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM188 target proteins.

[2612] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM188 gene, herein designated GAM GENE, on one or more GAM188 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2613] It is yet further appreciated that specific functions, and accordingly utilities, of GAM188 correlate with, and may be deduced from, the identity of the target genes which GAM188 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[2614] Nucleotide sequences of the GAM188 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM188 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM188 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM188 are further described hereinbelow with reference to Table Table1.

[2615] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM188 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2616]

[2617] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 189 (GAM189) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2618] GAM189 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM189 was detected is described hereinabove with reference to Figs. 2-8.

[2619] GAM189 gene, herein designated GAM GENE, and GAM189 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2620] GAM189 gene, herein designated GAM GENE, encodes a GAM189 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM189 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM189 precursor RNA is designated SEQ ID:2737, and is provided hereinbelow with reference to the sequence listing part.

[2621] GAM189 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM189 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[2622] An enzyme complex designated DICER COMPLEX, dices the GAM189 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM189 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM189 RNA is designated SEQ ID:2738, and is provided hereinbelow with reference to the sequence listing part.

[2623] GAM189 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM189 target RNA, herein designated GAM TARGET RNA. GAM189 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2624] GAM189 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM189 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM189 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM189 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM189 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2625] The complementary binding of GAM189 RNA, herein designated GAM RNA, to target binding sites on GAM189 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM189 target RNA, herein designated GAM TARGET RNA, into GAM189 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2626] It is appreciated that GAM189 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM189 target genes. The mRNA of each one of this plurality of GAM189 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM189 RNA, herein designated GAM RNA, and which when bound by GAM189 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM189 target proteins.

[2627] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM189 gene, herein designated GAM GENE, on one or more GAM189 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2628] It is yet further appreciated that specific functions, and accordingly utilities, of GAM189 correlate with, and may be deduced from, the identity of the target genes which GAM189 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2629] Nucleotide sequences of the GAM189 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM189 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM189 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM189 are further described hereinbelow with reference to Table Table1.

[2630] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM189 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2631]

[2632] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 190 (GAM190) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2633] GAM190 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM190 was detected is described hereinabove with reference to Figs. 2-8.

[2634] GAM190 gene, herein designated GAM GENE, and GAM190 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2635] GAM190 gene, herein designated GAM GENE, encodes a GAM190 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM190 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM190 precursor RNA is designated SEQ ID:2744, and is provided hereinbelow with reference to the sequence listing part.

[2636] GAM190 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM190 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2637] An enzyme complex designated DICER COMPLEX, dices the GAM190 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM190 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM190 RNA is designated SEQ ID:2745, and is provided hereinbelow with reference to the sequence listing part.

[2638] GAM190 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM190

target RNA, herein designated GAM TARGET RNA.

GAM190 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2639] GAM190 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM190 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM190 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM190 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM190 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2640] The complementary binding of GAM190 RNA, herein designated GAM RNA, to target binding sites on GAM190 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM190 target RNA, herein designated GAM TARGET RNA, into GAM190 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2641] It is appreciated that GAM190 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM190 target genes. The mRNA of each one of this plurality of GAM190 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM190 RNA, herein designated GAM RNA, and which when bound by GAM190 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM190 target proteins.

[2642] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM190 gene, herein designated GAM GENE, on one or more GAM190 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2643] It is yet further appreciated that specific functions, and accordingly utilities, of GAM190 correlate with, and may be deduced from, the identity of the target genes which GAM190 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2644] Nucleotide sequences of the GAM190 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM190 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM190 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM190 are further described hereinbelow with reference to Table Table1.

[2645] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM190 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2646]

[2647] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 191 (GAM191) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2648] GAM191 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM191 was detected is described hereinabove with reference to Figs. 2-8.

[2649] GAM191 gene, herein designated GAM GENE, and GAM191 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2650] GAM191 gene, herein designated GAM GENE, encodes a GAM191 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM191 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM191 precursor RNA is designated SEQ ID:2753, and is provided hereinbelow with reference to the sequence listing part.

[2651] GAM191 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM191 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2652] An enzyme complex designated DICER COMPLEX, dices the GAM191 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM191 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM191 RNA is designated SEQ ID:2754, and is provided hereinbelow with reference to the sequence listing part.

[2653] GAM191 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM191 target RNA, herein designated GAM TARGET RNA. GAM191 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2654] GAM191 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM191 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM191 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM191 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM191 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2655] The complementary binding of GAM191 RNA, herein designated GAM RNA, to target binding sites on GAM191 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM191 target RNA, herein designated GAM TARGET RNA, into GAM191 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2656] It is appreciated that GAM191 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM191 target genes. The mRNA of each one of this plu-

rality of GAM191 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM191 RNA, herein designated GAM RNA, and which when bound by GAM191 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM191 target proteins.

[2657] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM191 gene, herein designated GAM GENE, on one or more GAM191 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2658] It is yet further appreciated that specific functions, and accordingly utilities, of GAM191 correlate with, and may be deduced from, the identity of the target genes which GAM191 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2659] Nucleotide sequences of the GAM191 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM191 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM191 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM191 are further described hereinbelow with reference to Table Table1.

[2660] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM191 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2661]

[2662] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 192 (GAM192) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[2663] GAM192 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM192 was detected is described hereinabove with reference to Figs. 2-8.

[2664] GAM192 gene, herein designated GAM GENE, and GAM192 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2665] GAM192 gene, herein designated GAM GENE, encodes a GAM192 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM192 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM192 precursor RNA is designated SEQ ID:2763, and is provided hereinbelow with reference to the sequence listing part.

[2666] GAM192 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM192 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2667] An enzyme complex designated DICER COMPLEX, dices the GAM192 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM192 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM192 RNA is designated SEQ ID:2764, and is provided hereinbelow with reference to the sequence listing part.

[2668] GAM192 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM192 target RNA, herein designated GAM TARGET RNA.

GAM192 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[2669] GAM192 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM192 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM192 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM192 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM192 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2670] The complementary binding of GAM192 RNA, herein designated GAM RNA, to target binding sites on GAM192 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM192 target RNA, herein designated GAM TARGET RNA, into GAM192 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2671] It is appreciated that GAM192 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM192 target genes. The mRNA of each one of this plurality of GAM192 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM192 RNA, herein designated GAM RNA, and which when bound by GAM192 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM192 target proteins.

[2672] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM192 gene, herein designated GAM GENE, on one or more GAM192 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2673] It is yet further appreciated that specific functions, and accordingly utilities, of GAM192 correlate with, and may be deduced from, the identity of the target genes which GAM192 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2674] Nucleotide sequences of the GAM192 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM192 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM192 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM192 are further described hereinbelow with reference to Table Table1.

[2675] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM192 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2676]

[2677] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 193 (GAM193) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2678] GAM193 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM193 was detected is described hereinabove with reference to Figs. 2-8.

[2679] GAM193 gene, herein designated GAM GENE, and GAM193 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2680] GAM193 gene, herein designated GAM GENE, encodes a GAM193 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM193 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM193 precursor RNA is designated SEQ ID:2783, and is provided hereinbelow with reference to the sequence listing part.

[2681] GAM193 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM193 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2682] An enzyme complex designated DICER COMPLEX, dices the GAM193 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM193 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of

GAM193 RNA is designated SEQ ID:2784, and is provided hereinbelow with reference to the sequence listing part.

[2683] GAM193 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM193 target RNA, herein designated GAM TARGET RNA.

GAM193 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2684] GAM193 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM193 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM193 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM193 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM193 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2685] The complementary binding of GAM193 RNA, herein designated GAM RNA, to target binding sites on GAM193 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM193 target RNA, herein designated GAM TARGET RNA, into GAM193 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2686] It is appreciated that GAM193 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM193 target genes. The mRNA of each one of this plurality of GAM193 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM193 RNA, herein designated GAM RNA, and which when bound by GAM193 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM193 target proteins.

[2687] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM193 gene, herein designated GAM GENE, on one or more GAM193 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2688] It is yet further appreciated that specific functions, and accordingly utilities, of GAM193 correlate with, and may be deduced from, the identity of the target genes which GAM193 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2689] Nucleotide sequences of the GAM193 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM193 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM193 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM193 are further described hereinbelow with reference to Table Table1.

[2690] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM193 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2691]

[2692] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 194 (GAM194) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2693] GAM194 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM194 was detected is described hereinabove

with reference to Figs. 2-8.

[2694] GAM194 gene, herein designated GAM GENE, and GAM194 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2695] GAM194 gene, herein designated GAM GENE, encodes a GAM194 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM194 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM194 precursor RNA is designated SEQ ID:2794, and is provided hereinbelow with reference to the sequence listing part.

[2696] GAM194 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM194 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2697] An enzyme complex designated DICER COMPLEX, dices the GAM194 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM194 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM194 RNA is designated SEQ ID:2795, and is provided hereinbelow with reference to the sequence listing part.

[2698] GAM194 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM194 target RNA, herein designated GAM TARGET RNA. GAM194 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2699] GAM194 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM194 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM194 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM194 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM194 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2700] The complementary binding of GAM194 RNA, herein designated GAM RNA, to target binding sites on GAM194 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM194 target RNA, herein designated GAM TARGET RNA, into GAM194 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[2701] It is appreciated that GAM194 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM194 target genes. The mRNA of each one of this plurality of GAM194 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM194 RNA, herein designated GAM RNA, and which when bound by GAM194 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM194 target proteins.

[2702] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM194 gene, herein designated GAM GENE, on one or more GAM194 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2703] It is yet further appreciated that specific functions, and accordingly utilities, of GAM194 correlate with, and may be deduced from, the identity of the target genes which GAM194 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2704] Nucleotide sequences of the GAM194 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM194 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM194 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM194 are further described hereinbelow with reference to Table Table1.

[2705] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM194 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2706]

[2707] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 195 (GAM195) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2708] GAM195 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM195 was detected is described hereinabove with reference to Figs. 2-8.

[2709] GAM195 gene, herein designated GAM GENE, and GAM195 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2710] GAM195 gene, herein designated GAM GENE, encodes a GAM195 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM195 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM195 precursor RNA is designated SEQ ID:2807, and is provided hereinbelow with reference to the sequence listing part.

[2711] GAM195 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM195 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2712] An enzyme complex designated DICER COMPLEX, dices the GAM195 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM195 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM195 RNA is designated SEQ ID:2808, and is provided hereinbelow with reference to the sequence listing part.

[2713] GAM195 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM195 target RNA, herein designated GAM TARGET RNA.

GAM195 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2714] GAM195 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM195 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM195 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM195 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM195 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2715] The complementary binding of GAM195 RNA, herein designated GAM RNA, to target binding sites on GAM195 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM195 target RNA, herein designated GAM TARGET RNA, into GAM195 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2716] It is appreciated that GAM195 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM195 target genes. The mRNA of each one of this plurality of GAM195 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM195 RNA, herein designated GAM RNA, and which when bound by GAM195 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM195 target proteins.

[2717] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM195 gene, herein designated GAM GENE, on one or more GAM195 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2718] It is yet further appreciated that specific functions, and accordingly utilities, of GAM195 correlate with, and may be deduced from, the identity of the target genes which GAM195 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2719] Nucleotide sequences of the GAM195 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM195 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM195 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM195 are further described hereinbelow with reference to Table Table1.

[2720] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM195 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2721]

[2722] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 196 (GAM196) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2723] GAM196 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM196 was detected is described hereinabove with reference to Figs. 2-8.

[2724] GAM196 gene, herein designated GAM GENE, and GAM196 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2725] GAM196 gene, herein designated GAM GENE, encodes a

GAM196 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM196 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM196 precursor RNA is designated SEQ ID:2826, and is provided hereinbelow with reference to the sequence listing part.

[2726] GAM196 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM196 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2727] An enzyme complex designated DICER COMPLEX, dices the GAM196 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM196 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM196 RNA is designated SEQ ID:2827, and is provided hereinbelow with reference to the sequence listing part.

[2728] GAM196 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM196 target RNA, herein designated GAM TARGET RNA. GAM196 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2729] GAM196 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM196 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM196 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM196 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM196 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2730] The complementary binding of GAM196 RNA, herein designated GAM RNA, to target binding sites on GAM196 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM196 target RNA, herein designated GAM TARGET RNA, into GAM196 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2731] It is appreciated that GAM196 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM196 target genes. The mRNA of each one of this plurality of GAM196 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM196 RNA, herein designated GAM RNA, and which when bound by GAM196 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM196 target proteins.

[2732] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM196 gene, herein designated GAM GENE, on one or more GAM196 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2733] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM196 correlate with, and may be deduced from, the identity of the target genes which GAM196 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2734] Nucleotide sequences of the GAM196 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM196 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM196 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM196 are further described hereinbelow with reference to Table Table1.

[2735] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM196 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2736]

[2737] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 197 (GAM197) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[2738] GAM197 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM197 was detected is described hereinabove with reference to Figs. 2-8.

[2739] GAM197 gene, herein designated GAM GENE, and GAM197 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2740] GAM197 gene, herein designated GAM GENE, encodes a GAM197 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM197 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM197 precursor RNA is designated SEQ ID:2846, and is provided hereinbelow with reference to the sequence listing part.

[2741] GAM197 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM197 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2742] An enzyme complex designated DICER COMPLEX, dices the GAM197 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM197 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM197 RNA is designated SEQ ID:2847, and is provided hereinbelow with reference to the sequence listing part.

[2743] GAM197 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM197 target RNA, herein designated GAM TARGET RNA. GAM197 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2744] GAM197 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM197 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM197 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM197 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM197 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2745] The complementary binding of GAM197 RNA, herein designated GAM RNA, to target binding sites on GAM197 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM197 target RNA, herein designated GAM TARGET RNA, into GAM197 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2746] It is appreciated that GAM197 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM197 target genes. The mRNA of each one of this plurality of GAM197 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM197 RNA, herein designated GAM RNA, and which when bound by GAM197 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM197 target proteins.

[2747] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM197 gene, herein designated GAM GENE, on one or more GAM197 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2748] It is yet further appreciated that specific functions, and accordingly utilities, of GAM197 correlate with, and may be deduced from, the identity of the target genes which GAM197 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2749] Nucleotide sequences of the GAM197 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM197 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM197 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM197 are further described hereinbelow with reference to Table Table1.

[2750] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM197 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2751]

[2752] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 198 (GAM198) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2753] GAM198 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM198 was detected is described hereinabove with reference to Figs. 2-8.

[2754] GAM198 gene, herein designated GAM GENE, and GAM198 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2755] GAM198 gene, herein designated GAM GENE, encodes a GAM198 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM198 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM198 precursor RNA is designated SEQ ID:2858, and is provided hereinbelow with reference to the sequence listing part.

[2756] GAM198 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM198 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2757] An enzyme complex designated DICER COMPLEX, dices the GAM198 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM198 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM198 RNA is designated SEQ ID:2859, and is provided

hereinbelow with reference to the sequence listing part.

[2758] GAM198 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM198 target RNA, herein designated GAM TARGET RNA.

GAM198 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2759] GAM198 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM198 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM198 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM198 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM198 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2760] The complementary binding of GAM198 RNA, herein designated GAM RNA, to target binding sites on GAM198 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM198 target RNA, herein designated GAM TARGET RNA, into GAM198 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2761] It is appreciated that GAM198 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM198 target genes. The mRNA of each one of this plurality of GAM198 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM198 RNA, herein designated GAM RNA, and which when bound by GAM198 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM198 tar-

get proteins.

[2762] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM198 gene, herein designated GAM GENE, on one or more GAM198 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2763] It is yet further appreciated that specific functions, and accordingly utilities, of GAM198 correlate with, and may be deduced from, the identity of the target genes which GAM198 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2764] Nucleotide sequences of the GAM198 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM198 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM198 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM198 are further described hereinbelow with reference to Table Table1.

[2765] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM198 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2766]

[2767] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 199 (GAM199) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2768] GAM199 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM199 was detected is described hereinabove with reference to Figs. 2-8.

[2769] GAM199 gene, herein designated GAM GENE, and GAM199 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2770] GAM199 gene, herein designated GAM GENE, encodes a GAM199 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM199 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM199 precursor RNA is designated SEQ ID:2874, and is provided hereinbelow with reference to the sequence listing part.

[2771] GAM199 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM199 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2772] An enzyme complex designated DICER COMPLEX, dices

the GAM199 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM199 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 42%) nucleotide sequence of GAM199 RNA is designated SEQ ID:2875, and is provided hereinbelow with reference to the sequence listing part.

[2773] GAM199 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM199 target RNA, herein designated GAM TARGET RNA. GAM199 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2774] GAM199 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM199 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM199

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM199 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM199 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2775] The complementary binding of GAM199 RNA, herein designated GAM RNA, to target binding sites on GAM199 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM199 target RNA, herein designated GAM TARGET RNA, into GAM199 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2776] It is appreciated that GAM199 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM199 target genes. The mRNA of each one of this plurality of GAM199 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM199 RNA, herein designated GAM RNA, and which when bound by GAM199 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM199 target proteins.

[2777] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM199 gene, herein designated GAM GENE, on one or more GAM199 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2778] It is yet further appreciated that specific functions, and accordingly utilities, of GAM199 correlate with, and may be deduced from, the identity of the target genes which GAM199 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2779] Nucleotide sequences of the GAM199 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM199 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM199 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM199 are further described hereinbelow with reference to Table Table1.

[2780] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM199 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2781]

[2782] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 200 (GAM200) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2783] GAM200 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM200 was detected is described hereinabove with reference to Figs. 2-8.

[2784] GAM200 gene, herein designated GAM GENE, and GAM200 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2785] GAM200 gene, herein designated GAM GENE, encodes a GAM200 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM200 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM200 precursor RNA is designated SEQ ID:2880, and is provided hereinbelow with reference to the sequence listing part.

[2786] GAM200 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM200 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2787] An enzyme complex designated DICER COMPLEX, dices the GAM200 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM200 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 20%) nucleotide sequence of GAM200 RNA is designated SEQ ID:2881, and is provided hereinbelow with reference to the sequence listing part.

[2788] GAM200 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM200 target RNA, herein designated GAM TARGET RNA. GAM200 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2789] GAM200 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM200 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM200 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM200 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM200 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[2790] The complementary binding of GAM200 RNA, herein designated GAM RNA, to target binding sites on GAM200 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM200 target RNA, herein designated GAM TARGET RNA, into GAM200 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2791] It is appreciated that GAM200 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM200 target genes. The mRNA of each one of this plurality of GAM200 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM200 RNA, herein designated GAM RNA, and which when bound by GAM200 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM200 target proteins.

[2792] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM200 gene, herein designated GAM GENE, on one or

more GAM200 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2793] It is yet further appreciated that specific functions, and accordingly utilities, of GAM200 correlate with, and may be deduced from, the identity of the target genes which GAM200 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2794] Nucleotide sequences of the GAM200 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM200 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM200 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM200 are further de-

scribed hereinbelow with reference to Table Table1.

[2795] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM200 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2796]

[2797] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 201 (GAM201) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2798] GAM201 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM201 was detected is described hereinabove with reference to Figs. 2-8.

[2799] GAM201 gene, herein designated GAM GENE, and GAM201 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2800] GAM201 gene, herein designated GAM GENE, encodes a GAM201 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM201 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM201 precursor RNA is designated SEQ ID:2898, and is provided hereinbelow with reference to the sequence listing part.

[2801] GAM201 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM201 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2802] An enzyme complex designated DICER COMPLEX, dices the GAM201 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM201 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM201 RNA is designated SEQ ID:2899, and is provided hereinbelow with reference to the sequence listing part.

[2803] GAM201 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM201 target RNA, herein designated GAM TARGET RNA. GAM201 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2804] GAM201 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM201 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM201 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM201 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM201 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2805] The complementary binding of GAM201 RNA, herein designated GAM RNA, to target binding sites on GAM201 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM201 target RNA, herein designated GAM TARGET RNA, into GAM201 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2806] It is appreciated that GAM201 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM201 target genes. The mRNA of each one of this plurality of GAM201 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM201 RNA, herein designated GAM RNA, and which when bound by GAM201 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM201 target proteins.

[2807] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM201 gene, herein designated GAM GENE, on one or more GAM201 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2808] It is yet further appreciated that specific functions, and accordingly utilities, of GAM201 correlate with, and may

be deduced from, the identity of the target genes which GAM201 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2809] Nucleotide sequences of the GAM201 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM201 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM201 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM201 are further described hereinbelow with reference to Table Table1.

[2810] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM201 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2811]

[2812] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 202 (GAM202) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2813] GAM202 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM202 was detected is described hereinabove with reference to Figs. 2-8.

[2814] GAM202 gene, herein designated GAM GENE, and GAM202 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2815] GAM202 gene, herein designated GAM GENE, encodes a GAM202 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM202 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM202 precursor RNA is designated SEQ ID:2921, and is provided hereinbelow with reference to the sequence listing part.

[2816] GAM202 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM202 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2817] An enzyme complex designated DICER COMPLEX, dices the GAM202 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM202 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM202 RNA is designated SEQ ID:2922, and is provided hereinbelow with reference to the sequence listing part.

[2818] GAM202 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM202 target RNA, herein designated GAM TARGET RNA. GAM202 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2819] GAM202 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM202 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM202 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM202 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM202 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2820] The complementary binding of GAM202 RNA, herein designated GAM RNA, to target binding sites on GAM202 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM202 target RNA, herein designated GAM TARGET RNA, into GAM202 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2821] It is appreciated that GAM202 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM202 target genes. The mRNA of each one of this plurality of GAM202 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM202 RNA, herein designated GAM RNA, and which when bound by GAM202 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM202 target proteins.

[2822] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM202 gene, herein designated GAM GENE, on one or more GAM202 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2823] It is yet further appreciated that specific functions, and accordingly utilities, of GAM202 correlate with, and may be deduced from, the identity of the target genes which GAM202 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2824] Nucleotide sequences of the GAM202 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM202 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM202 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM202 are further described hereinbelow with reference to Table Table1.

[2825] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM202 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2826]

[2827] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 203 (GAM203) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2828] GAM203 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM203 was detected is described hereinabove with reference to Figs. 2-8.

[2829] GAM203 gene, herein designated GAM GENE, and GAM203 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2830] GAM203 gene, herein designated GAM GENE, encodes a GAM203 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM203 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM203 precursor RNA is desig-

nated SEQ ID:2946, and is provided hereinbelow with reference to the sequence listing part.

[2831] GAM203 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM203 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2832] An enzyme complex designated DICER COMPLEX, dices the GAM203 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM203 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM203 RNA is designated SEQ ID:2947, and is provided hereinbelow with reference to the sequence listing part.

[2833] GAM203 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM203 target RNA, herein designated GAM TARGET RNA. GAM203 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2834] GAM203 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM203 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM203 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM203 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM203

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2835] The complementary binding of GAM203 RNA, herein designated GAM RNA, to target binding sites on GAM203 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM203 target RNA, herein designated GAM TARGET RNA, into GAM203 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2836] It is appreciated that GAM203 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM203 target genes. The mRNA of each one of this plurality of GAM203 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM203 RNA, herein designated GAM RNA, and which when bound by GAM203 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM203 target proteins.

[2837] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM203 gene, herein designated GAM GENE, on one or more GAM203 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2838] It is yet further appreciated that specific functions, and accordingly utilities, of GAM203 correlate with, and may be deduced from, the identity of the target genes which GAM203 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2839] Nucleotide sequences of the GAM203 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM203 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM203 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM203 are further described hereinbelow with reference to Table Table1.

[2840] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM203 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2841]

[2842] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 204 (GAM204) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2843] GAM204 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM204 was detected is described hereinabove with reference to Figs. 2-8.

[2844] GAM204 gene, herein designated GAM GENE, and

GAM204 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2845] GAM204 gene, herein designated GAM GENE, encodes a GAM204 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM204 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM204 precursor RNA is designated SEQ ID:2953, and is provided hereinbelow with reference to the sequence listing part.

[2846] GAM204 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM204 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2847] An enzyme complex designated DICER COMPLEX, dices the GAM204 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM204 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM204 RNA is designated SEQ ID:2954, and is provided hereinbelow with reference to the sequence listing part.

[2848] GAM204 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM204 target RNA, herein designated GAM TARGET RNA. GAM204 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2849] GAM204 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM204 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM204 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM204 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM204 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2850] The complementary binding of GAM204 RNA, herein designated GAM RNA, to target binding sites on GAM204 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM204 target RNA, herein designated GAM TARGET RNA, into GAM204 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2851] It is appreciated that GAM204 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM204 target genes. The mRNA of each one of this plurality of GAM204 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM204 RNA, herein designated GAM RNA, and which when bound by GAM204 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM204 target proteins.

[2852] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM204 gene, herein designated GAM GENE, on one or more GAM204 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2853] It is yet further appreciated that specific functions, and accordingly utilities, of GAM204 correlate with, and may be deduced from, the identity of the target genes which GAM204 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2854] Nucleotide sequences of the GAM204 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM204 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM204 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM204 are further described hereinbelow with reference to Table Table1.

[2855] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM204 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2856]

[2857] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 205 (GAM205) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2858] GAM205 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM205 was detected is described hereinabove with reference to Figs. 2-8.

[2859] GAM205 gene, herein designated GAM GENE, and GAM205 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2860] GAM205 gene, herein designated GAM GENE, encodes a GAM205 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM205 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM205 precursor RNA is designated SEQ ID:2967, and is provided hereinbelow with reference to the sequence listing part.

[2861] GAM205 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM205 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2862] An enzyme complex designated DICER COMPLEX, dices the GAM205 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM205 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM205 RNA is designated SEQ ID:2968, and is provided hereinbelow with reference to the sequence listing part.

[2863] GAM205 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM205 target RNA, herein designated GAM TARGET RNA. GAM205 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2864] GAM205 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM205 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM205 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM205 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM205 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2865] The complementary binding of GAM205 RNA, herein designated GAM RNA, to target binding sites on GAM205 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM205 target RNA, herein designated GAM TARGET RNA, into GAM205 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2866] It is appreciated that GAM205 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM205 target genes. The mRNA of each one of this plurality of GAM205 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM205 RNA, herein designated GAM RNA, and which when bound by GAM205 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM205 target proteins.

[2867] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM205 gene, herein designated GAM GENE, on one or more GAM205 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2868] It is yet further appreciated that specific functions, and accordingly utilities, of GAM205 correlate with, and may be deduced from, the identity of the target genes which GAM205 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2869] Nucleotide sequences of the GAM205 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM205 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM205 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM205 are further described hereinbelow with reference to Table Table1.

[2870] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM205 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2871]

[2872] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 206 (GAM206) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2873] GAM206 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM206 was detected is described hereinabove with reference to Figs. 2-8.

[2874] GAM206 gene, herein designated GAM GENE, and GAM206 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2875] GAM206 gene, herein designated GAM GENE, encodes a GAM206 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM206 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM206 precursor RNA is designated SEQ ID:2973, and is provided hereinbelow with reference to the sequence listing part.

[2876] GAM206 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM206 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2877] An enzyme complex designated DICER COMPLEX, dices the GAM206 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM206 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 82%) nucleotide sequence of GAM206 RNA is designated SEQ ID:2974, and is provided hereinbelow with reference to the sequence listing part.

[2878] GAM206 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM206 target RNA, herein designated GAM TARGET RNA. GAM206 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2879] GAM206 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM206 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM206 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM206 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM206 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2880] The complementary binding of GAM206 RNA, herein designated GAM RNA, to target binding sites on GAM206 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM206 target RNA, herein designated GAM TARGET RNA, into GAM206 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2881] It is appreciated that GAM206 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM206 target genes. The mRNA of each one of this plurality of GAM206 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM206 RNA,

herein designated GAM RNA, and which when bound by GAM206 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM206 target proteins.

[2882] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM206 gene, herein designated GAM GENE, on one or more GAM206 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2883] It is yet further appreciated that specific functions, and accordingly utilities, of GAM206 correlate with, and may be deduced from, the identity of the target genes which

GAM206 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2884] Nucleotide sequences of the GAM206 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM206 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM206 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM206 are further described hereinbelow with reference to Table Table1.

[2885] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM206 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2886]

[2887] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 207 (GAM207) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2888] GAM207 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM207 was detected is described hereinabove with reference to Figs. 2-8.

[2889] GAM207 gene, herein designated GAM GENE, and GAM207 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2890] GAM207 gene, herein designated GAM GENE, encodes a GAM207 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM207 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM207 precursor RNA is designated SEQ ID:3004, and is provided hereinbelow with reference to the sequence listing part.

[2891] GAM207 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM207 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[2892] An enzyme complex designated DICER COMPLEX, dices the GAM207 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM207 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 26%) nucleotide sequence of GAM207 RNA is designated SEQ ID:3005, and is provided hereinbelow with reference to the sequence listing part.

[2893] GAM207 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM207 target RNA, herein designated GAM TARGET RNA.
GAM207 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2894] GAM207 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM207 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM207 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM207 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM207 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2895] The complementary binding of GAM207 RNA, herein designated GAM RNA, to target binding sites on GAM207 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM207 target RNA, herein desig-

nated GAM TARGET RNA, into GAM207 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2896] It is appreciated that GAM207 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM207 target genes. The mRNA of each one of this plurality of GAM207 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM207 RNA, herein designated GAM RNA, and which when bound by GAM207 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM207 target proteins.

[2897] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM207 gene, herein designated GAM GENE, on one or more GAM207 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2898] It is yet further appreciated that specific functions, and accordingly utilities, of GAM207 correlate with, and may be deduced from, the identity of the target genes which GAM207 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2899] Nucleotide sequences of the GAM207 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM207 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM207 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM207 are further described hereinbelow with reference to Table Table1.

[2900] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM207 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[2901]

[2902] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 208 (GAM208) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2903] GAM208 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM208 was detected is described hereinabove with reference to Figs. 2-8.

[2904] GAM208 gene, herein designated GAM GENE, and GAM208 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2905] GAM208 gene, herein designated GAM GENE, encodes a GAM208 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM208 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM208 precursor RNA is designated SEQ ID:3021, and is provided hereinbelow with ref-

erence to the sequence listing part.

[2906] GAM208 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM208 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2907] An enzyme complex designated DICER COMPLEX, dices the GAM208 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM208 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 79%) nucleotide sequence of GAM208 RNA is designated SEQ ID:3022, and is provided hereinbelow with reference to the sequence listing part.

[2908] GAM208 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM208 target RNA, herein designated GAM TARGET RNA.

GAM208 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2909] GAM208 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM208 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM208 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM208 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM208 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2910] The complementary binding of GAM208 RNA, herein designated GAM RNA, to target binding sites on GAM208 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM208 target RNA, herein designated GAM TARGET RNA, into GAM208 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2911] It is appreciated that GAM208 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM208 target genes. The mRNA of each one of this plurality of GAM208 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM208 RNA, herein designated GAM RNA, and which when bound by GAM208 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM208 target proteins.

[2912] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM208 gene, herein designated GAM GENE, on one or more GAM208 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2913] It is yet further appreciated that specific functions, and accordingly utilities, of GAM208 correlate with, and may be deduced from, the identity of the target genes which GAM208 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2914] Nucleotide sequences of the GAM208 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM208 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM208 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM208 are further described hereinbelow with reference to Table Table1.

[2915] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM208 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2916]

[2917] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 209 (GAM209) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2918] GAM209 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM209 was detected is described hereinabove with reference to Figs. 2-8.

[2919] GAM209 gene, herein designated GAM GENE, and GAM209 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[2920] GAM209 gene, herein designated GAM GENE, encodes a GAM209 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM209 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM209 precursor RNA is designated SEQ ID:3030, and is provided hereinbelow with reference to the sequence listing part.

[2921] GAM209 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM209 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2922] An enzyme complex designated DICER COMPLEX, dices the GAM209 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM209 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM209 RNA is designated SEQ ID:3031, and is provided hereinbelow with reference to the sequence listing part.

[2923] GAM209 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM209 target RNA, herein designated GAM TARGET RNA. GAM209 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2924] GAM209 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM209 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM209 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM209 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM209 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2925] The complementary binding of GAM209 RNA, herein designated GAM RNA, to target binding sites on GAM209 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM209 target RNA, herein designated GAM TARGET RNA, into GAM209 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2926] It is appreciated that GAM209 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM209 target genes. The mRNA of each one of this plurality of GAM209 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM209 RNA, herein designated GAM RNA, and which when bound by GAM209 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM209 target proteins.

[2927] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM209 gene, herein designated GAM GENE, on one or more GAM209 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[2928] It is yet further appreciated that specific functions, and accordingly utilities, of GAM209 correlate with, and may be deduced from, the identity of the target genes which GAM209 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2929] Nucleotide sequences of the GAM209 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM209 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM209 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM209 are further described hereinbelow with reference to Table Table1.

[2930] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM209 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2931]

[2932] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

210 (GAM210) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2933] GAM210 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM210 was detected is described hereinabove with reference to Figs. 2-8.

[2934] GAM210 gene, herein designated GAM GENE, and GAM210 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2935] GAM210 gene, herein designated GAM GENE, encodes a GAM210 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM210 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM210 precursor RNA is designated SEQ ID:3043, and is provided hereinbelow with reference to the sequence listing part.

[2936] GAM210 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM210 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2937] An enzyme complex designated DICER COMPLEX, dices the GAM210 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM210 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM210 RNA is designated SEQ ID:3044, and is provided hereinbelow with reference to the sequence listing part.

[2938] GAM210 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM210 target RNA, herein designated GAM TARGET RNA. GAM210 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2939] GAM210 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM210 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM210 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM210 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM210 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2940] The complementary binding of GAM210 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM210 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM210 target RNA, herein designated GAM TARGET RNA, into GAM210 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2941] It is appreciated that GAM210 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM210 target genes. The mRNA of each one of this plurality of GAM210 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM210 RNA, herein designated GAM RNA, and which when bound by GAM210 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM210 target proteins.

[2942] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM210 gene, herein designated GAM GENE, on one or more GAM210 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2943] It is yet further appreciated that specific functions, and accordingly utilities, of GAM210 correlate with, and may be deduced from, the identity of the target genes which GAM210 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2944] Nucleotide sequences of the GAM210 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM210 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM210 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM210 are further described hereinbelow with reference to Table Table1.

[2945] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM210 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2946]

[2947] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 211 (GAM211) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2948] GAM211 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM211 was detected is described hereinabove with reference to Figs. 2-8.

[2949] GAM211 gene, herein designated GAM GENE, and GAM211 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2950] GAM211 gene, herein designated GAM GENE, encodes a GAM211 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM211 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM211 precursor RNA is designated SEQ ID:3049, and is provided hereinbelow with reference to the sequence listing part.

[2951] GAM211 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM211 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2952] An enzyme complex designated DICER COMPLEX, dices the GAM211 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM211 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 66%) nucleotide sequence of GAM211 RNA is designated SEQ ID:3050, and is provided hereinbelow with reference to the sequence listing part.

[2953] GAM211 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM211 target RNA, herein designated GAM TARGET RNA. GAM211 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2954] GAM211 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM211 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM211 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM211 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM211 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2955] The complementary binding of GAM211 RNA, herein designated GAM RNA, to target binding sites on GAM211 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM211 target RNA, herein designated GAM TARGET RNA, into GAM211 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2956] It is appreciated that GAM211 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM211 target genes. The mRNA of each one of this plurality of GAM211 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM211 RNA, herein designated GAM RNA, and which when bound by

GAM211 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM211 target proteins.

[2957] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM211 gene, herein designated GAM GENE, on one or more GAM211 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2958] It is yet further appreciated that specific functions, and accordingly utilities, of GAM211 correlate with, and may be deduced from, the identity of the target genes which GAM211 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[2959] Nucleotide sequences of the GAM211 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM211 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM211 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM211 are further described hereinbelow with reference to Table Table1.

[2960] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM211 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2961]

[2962] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 212 (GAM212) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2963] GAM212 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM212 was detected is described hereinabove with reference to Figs. 2-8.

[2964] GAM212 gene, herein designated GAM GENE, and GAM212 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2965] GAM212 gene, herein designated GAM GENE, encodes a GAM212 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM212 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM212 precursor RNA is designated SEQ ID:3069, and is provided hereinbelow with reference to the sequence listing part.

[2966] GAM212 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM212 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[2967] An enzyme complex designated DICER COMPLEX, dices the GAM212 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM212 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM212 RNA is designated SEQ ID:3070, and is provided hereinbelow with reference to the sequence listing part.

[2968] GAM212 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM212 target RNA, herein designated GAM TARGET RNA. GAM212 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2969] GAM212 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM212 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM212 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM212 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM212 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2970] The complementary binding of GAM212 RNA, herein designated GAM RNA, to target binding sites on GAM212 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM212 target RNA, herein designated GAM TARGET RNA, into GAM212 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2971] It is appreciated that GAM212 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM212 target genes. The mRNA of each one of this plurality of GAM212 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM212 RNA, herein designated GAM RNA, and which when bound by GAM212 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM212 target proteins.

[2972] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM212 gene, herein designated GAM GENE, on one or more GAM212 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2973] It is yet further appreciated that specific functions, and accordingly utilities, of GAM212 correlate with, and may be deduced from, the identity of the target genes which GAM212 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2974] Nucleotide sequences of the GAM212 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM212 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM212 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM212 are further described hereinbelow with reference to Table Table1.

[2975] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM212 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2976]

[2977] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 213 (GAM213) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2978] GAM213 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM213 was detected is described hereinabove with reference to Figs. 2-8.

[2979] GAM213 gene, herein designated GAM GENE, and GAM213 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2980] GAM213 gene, herein designated GAM GENE, encodes a GAM213 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM213 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM213 precursor RNA is designated SEQ ID:3075, and is provided hereinbelow with reference to the sequence listing part.

[2981] GAM213 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM213 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2982] An enzyme complex designated DICER COMPLEX, dices the GAM213 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM213 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM213 RNA is designated SEQ ID:3076, and is provided hereinbelow with reference to the sequence listing part.

[2983] GAM213 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM213

target RNA, herein designated GAM TARGET RNA.

GAM213 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2984] GAM213 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM213 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM213 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM213 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM213 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[2985] The complementary binding of GAM213 RNA, herein designated GAM RNA, to target binding sites on GAM213 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM213 target RNA, herein designated GAM TARGET RNA, into GAM213 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[2986] It is appreciated that GAM213 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM213 target genes. The mRNA of each one of this plurality of GAM213 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM213 RNA, herein designated GAM RNA, and which when bound by GAM213 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM213 target proteins.

[2987] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM213 gene, herein designated GAM GENE, on one or more GAM213 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[2988] It is yet further appreciated that specific functions, and accordingly utilities, of GAM213 correlate with, and may be deduced from, the identity of the target genes which GAM213 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[2989] Nucleotide sequences of the GAM213 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM213 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM213 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM213 are further described hereinbelow with reference to Table Table1.

[2990] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM213 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[2991]

[2992] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 214 (GAM214) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[2993] GAM214 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM214 was detected is described hereinabove with reference to Figs. 2-8.

[2994] GAM214 gene, herein designated GAM GENE, and GAM214 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[2995] GAM214 gene, herein designated GAM GENE, encodes a GAM214 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM214 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM214 precursor RNA is designated SEQ ID:3081, and is provided hereinbelow with reference to the sequence listing part.

[2996] GAM214 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM214 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[2997] An enzyme complex designated DICER COMPLEX, dices the GAM214 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM214 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM214 RNA is designated SEQ ID:3082, and is provided hereinbelow with reference to the sequence listing part.

[2998] GAM214 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM214 target RNA, herein designated GAM TARGET RNA. GAM214 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[2999] GAM214 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM214 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM214 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM214 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM214 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3000] The complementary binding of GAM214 RNA, herein designated GAM RNA, to target binding sites on GAM214 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM214 target RNA, herein designated GAM TARGET RNA, into GAM214 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3001] It is appreciated that GAM214 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM214 target genes. The mRNA of each one of this plu-

rality of GAM214 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM214 RNA, herein designated GAM RNA, and which when bound by GAM214 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM214 target proteins.

[3002] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM214 gene, herein designated GAM GENE, on one or more GAM214 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3003] It is yet further appreciated that specific functions, and accordingly utilities, of GAM214 correlate with, and may be deduced from, the identity of the target genes which GAM214 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3004] Nucleotide sequences of the GAM214 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM214 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM214 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM214 are further described hereinbelow with reference to Table Table1.

[3005] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM214 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3006]

[3007] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 215 (GAM215) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[3008] GAM215 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM215 was detected is described hereinabove with reference to Figs. 2-8.

[3009] GAM215 gene, herein designated GAM GENE, and GAM215 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3010] GAM215 gene, herein designated GAM GENE, encodes a GAM215 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM215 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM215 precursor RNA is designated SEQ ID:3101, and is provided hereinbelow with reference to the sequence listing part.

[3011] GAM215 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM215 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3012] An enzyme complex designated DICER COMPLEX, dices the GAM215 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM215 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM215 RNA is designated SEQ ID:3102, and is provided hereinbelow with reference to the sequence listing part.

[3013] GAM215 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM215 target RNA, herein designated GAM TARGET RNA.
GAM215 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[3014] GAM215 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM215 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM215 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM215 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM215 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3015] The complementary binding of GAM215 RNA, herein designated GAM RNA, to target binding sites on GAM215 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM215 target RNA, herein designated GAM TARGET RNA, into GAM215 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3016] It is appreciated that GAM215 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM215 target genes. The mRNA of each one of this plurality of GAM215 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM215 RNA, herein designated GAM RNA, and which when bound by GAM215 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM215 target proteins.

[3017] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM215 gene, herein designated GAM GENE, on one or more GAM215 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3018] It is yet further appreciated that specific functions, and accordingly utilities, of GAM215 correlate with, and may be deduced from, the identity of the target genes which GAM215 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3019] Nucleotide sequences of the GAM215 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM215 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM215 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM215 are further described hereinbelow with reference to Table Table1.

[3020] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM215 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3021]

[3022] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 216 (GAM216) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3023] GAM216 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM216 was detected is described hereinabove with reference to Figs. 2-8.

[3024] GAM216 gene, herein designated GAM GENE, and GAM216 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3025] GAM216 gene, herein designated GAM GENE, encodes a GAM216 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM216 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM216 precursor RNA is designated SEQ ID:3118, and is provided hereinbelow with reference to the sequence listing part.

[3026] GAM216 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM216 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3027] An enzyme complex designated DICER COMPLEX, dices the GAM216 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM216 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 32%) nucleotide sequence of

GAM216 RNA is designated SEQ ID:3119, and is provided hereinbelow with reference to the sequence listing part.

[3028] GAM216 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM216 target RNA, herein designated GAM TARGET RNA. GAM216 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3029] GAM216 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM216 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM216 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM216 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM216 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3030] The complementary binding of GAM216 RNA, herein designated GAM RNA, to target binding sites on GAM216 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM216 target RNA, herein designated GAM TARGET RNA, into GAM216 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3031] It is appreciated that GAM216 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM216 target genes. The mRNA of each one of this plurality of GAM216 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM216 RNA, herein designated GAM RNA, and which when bound by GAM216 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM216 target proteins.

[3032] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM216 gene, herein designated GAM GENE, on one or more GAM216 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3033] It is yet further appreciated that specific functions, and accordingly utilities, of GAM216 correlate with, and may be deduced from, the identity of the target genes which GAM216 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3034] Nucleotide sequences of the GAM216 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM216 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM216 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM216 are further described hereinbelow with reference to Table Table1.

[3035] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM216 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3036]

[3037] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 217 (GAM217) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3038] GAM217 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM217 was detected is described hereinabove

with reference to Figs. 2-8.

[3039] GAM217 gene, herein designated GAM GENE, and GAM217 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3040] GAM217 gene, herein designated GAM GENE, encodes a GAM217 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM217 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM217 precursor RNA is designated SEQ ID:3132, and is provided hereinbelow with reference to the sequence listing part.

[3041] GAM217 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM217 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3042] An enzyme complex designated DICER COMPLEX, dices the GAM217 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM217 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM217 RNA is designated SEQ ID:3133, and is provided hereinbelow with reference to the sequence listing part.

[3043] GAM217 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM217 target RNA, herein designated GAM TARGET RNA. GAM217 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3044] GAM217 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM217 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM217 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM217 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM217 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3045] The complementary binding of GAM217 RNA, herein designated GAM RNA, to target binding sites on GAM217 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM217 target RNA, herein designated GAM TARGET RNA, into GAM217 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[3046] It is appreciated that GAM217 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM217 target genes. The mRNA of each one of this plurality of GAM217 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM217 RNA, herein designated GAM RNA, and which when bound by GAM217 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM217 target proteins.

[3047] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM217 gene, herein designated GAM GENE, on one or more GAM217 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3048] It is yet further appreciated that specific functions, and accordingly utilities, of GAM217 correlate with, and may be deduced from, the identity of the target genes which GAM217 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3049] Nucleotide sequences of the GAM217 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM217 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM217 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM217 are further described hereinbelow with reference to Table Table1.

[3050] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM217 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3051]

[3052] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 218 (GAM218) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3053] GAM218 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM218 was detected is described hereinabove with reference to Figs. 2-8.

[3054] GAM218 gene, herein designated GAM GENE, and GAM218 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3055] GAM218 gene, herein designated GAM GENE, encodes a GAM218 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM218 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM218 precursor RNA is designated SEQ ID:3172, and is provided hereinbelow with reference to the sequence listing part.

[3056] GAM218 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM218 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3057] An enzyme complex designated DICER COMPLEX, dices the GAM218 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM218 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM218 RNA is designated SEQ ID:3173, and is provided hereinbelow with reference to the sequence listing part.

[3058] GAM218 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM218 target RNA, herein designated GAM TARGET RNA.

GAM218 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3059] GAM218 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM218 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM218 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM218 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM218 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3060] The complementary binding of GAM218 RNA, herein designated GAM RNA, to target binding sites on GAM218 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM218 target RNA, herein designated GAM TARGET RNA, into GAM218 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3061] It is appreciated that GAM218 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM218 target genes. The mRNA of each one of this plurality of GAM218 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM218 RNA, herein designated GAM RNA, and which when bound by GAM218 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM218 target proteins.

[3062] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM218 gene, herein designated GAM GENE, on one or more GAM218 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3063] It is yet further appreciated that specific functions, and accordingly utilities, of GAM218 correlate with, and may be deduced from, the identity of the target genes which GAM218 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3064] Nucleotide sequences of the GAM218 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM218 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM218 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM218 are further described hereinbelow with reference to Table Table1.

[3065] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM218 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3066]

[3067] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 219 (GAM219) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3068] GAM219 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM219 was detected is described hereinabove with reference to Figs. 2-8.

[3069] GAM219 gene, herein designated GAM GENE, and GAM219 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3070] GAM219 gene, herein designated GAM GENE, encodes a

GAM219 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM219 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM219 precursor RNA is designated SEQ ID:3204, and is provided hereinbelow with reference to the sequence listing part.

[3071] GAM219 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM219 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3072] An enzyme complex designated DICER COMPLEX, dices the GAM219 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM219 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM219 RNA is designated SEQ ID:3205, and is provided hereinbelow with reference to the sequence listing part.

[3073] GAM219 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM219 target RNA, herein designated GAM TARGET RNA. GAM219 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3074] GAM219 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM219 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM219 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM219 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM219

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3075] The complementary binding of GAM219 RNA, herein designated GAM RNA, to target binding sites on GAM219 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM219 target RNA, herein designated GAM TARGET RNA, into GAM219 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3076] It is appreciated that GAM219 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM219 target genes. The mRNA of each one of this plurality of GAM219 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM219 RNA, herein designated GAM RNA, and which when bound by GAM219 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM219 target proteins.

[3077] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM219 gene, herein designated GAM GENE, on one or more GAM219 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3078] It is yet further appreciated that specific functions, and accordingly utilities, of GAM219 correlate with, and may be deduced from, the identity of the target genes which GAM219 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3079] Nucleotide sequences of the GAM219 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM219 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM219 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM219 are further described hereinbelow with reference to Table Table1.

[3080] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM219 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3081]

[3082] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 220 (GAM220) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3083] GAM220 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM220 was detected is described hereinabove with reference to Figs. 2-8.

[3084] GAM220 gene, herein designated GAM GENE, and

GAM220 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3085] GAM220 gene, herein designated GAM GENE, encodes a GAM220 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM220 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM220 precursor RNA is designated SEQ ID:3210, and is provided hereinbelow with reference to the sequence listing part.

[3086] GAM220 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM220 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3087] An enzyme complex designated DICER COMPLEX, dices the GAM220 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM220 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM220 RNA is designated SEQ ID:3211, and is provided hereinbelow with reference to the sequence listing part.

[3088] GAM220 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM220 target RNA, herein designated GAM TARGET RNA. GAM220 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3089] GAM220 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM220 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM220 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM220 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM220 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3090] The complementary binding of GAM220 RNA, herein designated GAM RNA, to target binding sites on GAM220 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM220 target RNA, herein designated GAM TARGET RNA, into GAM220 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3091] It is appreciated that GAM220 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM220 target genes. The mRNA of each one of this plurality of GAM220 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM220 RNA, herein designated GAM RNA, and which when bound by GAM220 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM220 target proteins.

[3092] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM220 gene, herein designated GAM GENE, on one or more GAM220 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3093] It is yet further appreciated that specific functions, and accordingly utilities, of GAM220 correlate with, and may be deduced from, the identity of the target genes which GAM220 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3094] Nucleotide sequences of the GAM220 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM220 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM220 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM220 are further described hereinbelow with reference to Table Table1.

[3095] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM220 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3096]

[3097] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 221 (GAM221) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3098] GAM221 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM221 was detected is described hereinabove with reference to Figs. 2-8.

[3099] GAM221 gene, herein designated GAM GENE, and GAM221 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3100] GAM221 gene, herein designated GAM GENE, encodes a GAM221 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM221 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM221 precursor RNA is designated SEQ ID:3247, and is provided hereinbelow with reference to the sequence listing part.

[3101] GAM221 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM221 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3102] An enzyme complex designated DICER COMPLEX, dices the GAM221 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM221 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 33%) nucleotide sequence of GAM221 RNA is designated SEQ ID:3248, and is provided hereinbelow with reference to the sequence listing part.

[3103] GAM221 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM221 target RNA, herein designated GAM TARGET RNA. GAM221 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3104] GAM221 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM221 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM221 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM221 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM221 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3105] The complementary binding of GAM221 RNA, herein designated GAM RNA, to target binding sites on GAM221 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM221 target RNA, herein designated GAM TARGET RNA, into GAM221 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3106] It is appreciated that GAM221 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM221 target genes. The mRNA of each one of this plurality of GAM221 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM221 RNA, herein designated GAM RNA, and which when bound by GAM221 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM221 target proteins.

[3107] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM221 gene, herein designated GAM GENE, on one or more GAM221 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3108] It is yet further appreciated that specific functions, and accordingly utilities, of GAM221 correlate with, and may be deduced from, the identity of the target genes which GAM221 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3109] Nucleotide sequences of the GAM221 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM221 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM221 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM221 are further described hereinbelow with reference to Table Table1.

[3110] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM221 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3111]

[3112] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 222 (GAM222) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3113] GAM222 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM222 was detected is described hereinabove with reference to Figs. 2-8.

[3114] GAM222 gene, herein designated GAM GENE, and GAM222 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3115] GAM222 gene, herein designated GAM GENE, encodes a GAM222 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM222 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM222 precursor RNA is designated SEQ ID:3266, and is provided hereinbelow with reference to the sequence listing part.

[3116] GAM222 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM222 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3117] An enzyme complex designated DICER COMPLEX, dices the GAM222 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM222 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 36%) nucleotide sequence of GAM222 RNA is designated SEQ ID:3267, and is provided hereinbelow with reference to the sequence listing part.

[3118] GAM222 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM222 target RNA, herein designated GAM TARGET RNA. GAM222 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3119] GAM222 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM222 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM222 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM222 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM222 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3120] The complementary binding of GAM222 RNA, herein designated GAM RNA, to target binding sites on GAM222 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM222 target RNA, herein designated GAM TARGET RNA, into GAM222 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3121] It is appreciated that GAM222 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM222 target genes. The mRNA of each one of this plurality of GAM222 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM222 RNA,

herein designated GAM RNA, and which when bound by GAM222 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM222 target proteins.

[3122] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM222 gene, herein designated GAM GENE, on one or more GAM222 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3123] It is yet further appreciated that specific functions, and accordingly utilities, of GAM222 correlate with, and may be deduced from, the identity of the target genes which

GAM222 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3124] Nucleotide sequences of the GAM222 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM222 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM222 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM222 are further described hereinbelow with reference to Table Table1.

[3125] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM222 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3126]

[3127] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 223 (GAM223) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3128] GAM223 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM223 was detected is described hereinabove with reference to Figs. 2-8.

[3129] GAM223 gene, herein designated GAM GENE, and GAM223 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3130] GAM223 gene, herein designated GAM GENE, encodes a GAM223 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM223 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM223 precursor RNA is designated SEQ ID:3279, and is provided hereinbelow with reference to the sequence listing part.

[3131] GAM223 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM223 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[3132] An enzyme complex designated DICER COMPLEX, dices the GAM223 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM223 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 25%) nucleotide sequence of GAM223 RNA is designated SEQ ID:3280, and is provided hereinbelow with reference to the sequence listing part.

[3133] GAM223 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM223 target RNA, herein designated GAM TARGET RNA.
GAM223 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3134] GAM223 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM223 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM223 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM223 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM223 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3135] The complementary binding of GAM223 RNA, herein designated GAM RNA, to target binding sites on GAM223 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM223 target RNA, herein desig-

nated GAM TARGET RNA, into GAM223 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3136] It is appreciated that GAM223 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM223 target genes. The mRNA of each one of this plurality of GAM223 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM223 RNA, herein designated GAM RNA, and which when bound by GAM223 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM223 target proteins.

[3137] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM223 gene, herein designated GAM GENE, on one or more GAM223 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3138] It is yet further appreciated that specific functions, and accordingly utilities, of GAM223 correlate with, and may be deduced from, the identity of the target genes which GAM223 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3139] Nucleotide sequences of the GAM223 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM223 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM223 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM223 are further described hereinbelow with reference to Table Table1.

[3140] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM223 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[3141]

[3142] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 224 (GAM224) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3143] GAM224 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM224 was detected is described hereinabove with reference to Figs. 2-8.

[3144] GAM224 gene, herein designated GAM GENE, and GAM224 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3145] GAM224 gene, herein designated GAM GENE, encodes a GAM224 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM224 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM224 precursor RNA is designated SEQ ID:3287, and is provided hereinbelow with ref-

erence to the sequence listing part.

[3146] GAM224 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM224 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3147] An enzyme complex designated DICER COMPLEX, dices the GAM224 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM224 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM224 RNA is designated SEQ ID:3288, and is provided hereinbelow with reference to the sequence listing part.

[3148] GAM224 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM224 target RNA, herein designated GAM TARGET RNA.

GAM224 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3149] GAM224 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM224 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM224 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM224 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM224 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3150] The complementary binding of GAM224 RNA, herein designated GAM RNA, to target binding sites on GAM224 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM224 target RNA, herein designated GAM TARGET RNA, into GAM224 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3151] It is appreciated that GAM224 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM224 target genes. The mRNA of each one of this plurality of GAM224 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM224 RNA, herein designated GAM RNA, and which when bound by GAM224 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM224 target proteins.

[3152] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM224 gene, herein designated GAM GENE, on one or more GAM224 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3153] It is yet further appreciated that specific functions, and accordingly utilities, of GAM224 correlate with, and may be deduced from, the identity of the target genes which GAM224 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3154] Nucleotide sequences of the GAM224 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM224 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM224 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM224 are further described hereinbelow with reference to Table Table1.

[3155] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM224 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3156]

[3157] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 225 (GAM225) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3158] GAM225 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM225 was detected is described hereinabove with reference to Figs. 2-8.

[3159] GAM225 gene, herein designated GAM GENE, and GAM225 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[3160] GAM225 gene, herein designated GAM GENE, encodes a GAM225 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM225 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM225 precursor RNA is designated SEQ ID:3306, and is provided hereinbelow with reference to the sequence listing part.

[3161] GAM225 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM225 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3162] An enzyme complex designated DICER COMPLEX, dices the GAM225 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM225 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM225 RNA is designated SEQ ID:3307, and is provided hereinbelow with reference to the sequence listing part.

[3163] GAM225 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM225 target RNA, herein designated GAM TARGET RNA. GAM225 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3164] GAM225 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM225 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM225 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM225 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM225 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3165] The complementary binding of GAM225 RNA, herein designated GAM RNA, to target binding sites on GAM225 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM225 target RNA, herein designated GAM TARGET RNA, into GAM225 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3166] It is appreciated that GAM225 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM225 target genes. The mRNA of each one of this plurality of GAM225 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM225 RNA, herein designated GAM RNA, and which when bound by GAM225 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM225 target proteins.

[3167] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM225 gene, herein designated GAM GENE, on one or more GAM225 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[3168] It is yet further appreciated that specific functions, and accordingly utilities, of GAM225 correlate with, and may be deduced from, the identity of the target genes which GAM225 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3169] Nucleotide sequences of the GAM225 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM225 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM225 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM225 are further described hereinbelow with reference to Table Table1.

[3170] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM225 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3171]

[3172] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

226 (GAM226) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3173] GAM226 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM226 was detected is described hereinabove with reference to Figs. 2-8.

[3174] GAM226 gene, herein designated GAM GENE, and GAM226 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3175] GAM226 gene, herein designated GAM GENE, encodes a GAM226 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM226 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM226 precursor RNA is designated SEQ ID:3315, and is provided hereinbelow with reference to the sequence listing part.

[3176] GAM226 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM226 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3177] An enzyme complex designated DICER COMPLEX, dices the GAM226 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM226 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM226 RNA is designated SEQ ID:3316, and is provided hereinbelow with reference to the sequence listing part.

[3178] GAM226 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM226 target RNA, herein designated GAM TARGET RNA. GAM226 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3179] GAM226 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM226 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM226 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM226 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM226 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3180] The complementary binding of GAM226 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM226 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM226 target RNA, herein designated GAM TARGET RNA, into GAM226 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3181] It is appreciated that GAM226 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM226 target genes. The mRNA of each one of this plurality of GAM226 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM226 RNA, herein designated GAM RNA, and which when bound by GAM226 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM226 target proteins.

[3182] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM226 gene, herein designated GAM GENE, on one or more GAM226 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3183] It is yet further appreciated that specific functions, and accordingly utilities, of GAM226 correlate with, and may be deduced from, the identity of the target genes which GAM226 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3184] Nucleotide sequences of the GAM226 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM226 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM226 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM226 are further described hereinbelow with reference to Table Table1.

[3185] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM226 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3186]

[3187] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 227 (GAM227) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3188] GAM227 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM227 was detected is described hereinabove with reference to Figs. 2-8.

[3189] GAM227 gene, herein designated GAM GENE, and GAM227 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3190] GAM227 gene, herein designated GAM GENE, encodes a GAM227 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM227 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM227 precursor RNA is designated SEQ ID:3343, and is provided hereinbelow with reference to the sequence listing part.

[3191] GAM227 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM227 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3192] An enzyme complex designated DICER COMPLEX, dices the GAM227 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM227 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 56%) nucleotide sequence of GAM227 RNA is designated SEQ ID:3344, and is provided hereinbelow with reference to the sequence listing part.

[3193] GAM227 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM227 target RNA, herein designated GAM TARGET RNA. GAM227 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3194] GAM227 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM227 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM227 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM227 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM227 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3195] The complementary binding of GAM227 RNA, herein designated GAM RNA, to target binding sites on GAM227 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM227 target RNA, herein designated GAM TARGET RNA, into GAM227 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3196] It is appreciated that GAM227 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM227 target genes. The mRNA of each one of this plurality of GAM227 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM227 RNA, herein designated GAM RNA, and which when bound by

GAM227 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM227 target proteins.

[3197] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM227 gene, herein designated GAM GENE, on one or more GAM227 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3198] It is yet further appreciated that specific functions, and accordingly utilities, of GAM227 correlate with, and may be deduced from, the identity of the target genes which GAM227 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[3199] Nucleotide sequences of the GAM227 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM227 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM227 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM227 are further described hereinbelow with reference to Table Table1.

[3200] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM227 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3201]

[3202] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 228 (GAM228) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3203] GAM228 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM228 was detected is described hereinabove with reference to Figs. 2-8.

[3204] GAM228 gene, herein designated GAM GENE, and GAM228 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3205] GAM228 gene, herein designated GAM GENE, encodes a GAM228 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM228 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM228 precursor RNA is designated SEQ ID:3350, and is provided hereinbelow with reference to the sequence listing part.

[3206] GAM228 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM228 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[3207] An enzyme complex designated DICER COMPLEX, dices the GAM228 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM228 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM228 RNA is designated SEQ ID:3351, and is provided hereinbelow with reference to the sequence listing part.

[3208] GAM228 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM228 target RNA, herein designated GAM TARGET RNA. GAM228 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3209] GAM228 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM228 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM228 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM228 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM228 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3210] The complementary binding of GAM228 RNA, herein designated GAM RNA, to target binding sites on GAM228 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM228 target RNA, herein designated GAM TARGET RNA, into GAM228 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3211] It is appreciated that GAM228 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM228 target genes. The mRNA of each one of this plurality of GAM228 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM228 RNA, herein designated GAM RNA, and which when bound by GAM228 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM228 target proteins.

[3212] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM228 gene, herein designated GAM GENE, on one or more GAM228 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3213] It is yet further appreciated that specific functions, and accordingly utilities, of GAM228 correlate with, and may be deduced from, the identity of the target genes which GAM228 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3214] Nucleotide sequences of the GAM228 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM228 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM228 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM228 are further described hereinbelow with reference to Table Table1.

[3215] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM228 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3216]

[3217] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 229 (GAM229) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3218] GAM229 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM229 was detected is described hereinabove with reference to Figs. 2-8.

[3219] GAM229 gene, herein designated GAM GENE, and GAM229 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3220] GAM229 gene, herein designated GAM GENE, encodes a GAM229 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM229 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM229 precursor RNA is designated SEQ ID:3375, and is provided hereinbelow with reference to the sequence listing part.

[3221] GAM229 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM229 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3222] An enzyme complex designated DICER COMPLEX, dices the GAM229 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM229 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM229 RNA is designated SEQ ID:3376, and is provided hereinbelow with reference to the sequence listing part.

[3223] GAM229 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM229

target RNA, herein designated GAM TARGET RNA.

GAM229 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3224] GAM229 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM229 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM229 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM229 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM229 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3225] The complementary binding of GAM229 RNA, herein designated GAM RNA, to target binding sites on GAM229 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM229 target RNA, herein designated GAM TARGET RNA, into GAM229 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3226] It is appreciated that GAM229 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM229 target genes. The mRNA of each one of this plurality of GAM229 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM229 RNA, herein designated GAM RNA, and which when bound by GAM229 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM229 target proteins.

[3227] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM229 gene, herein designated GAM GENE, on one or more GAM229 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3228] It is yet further appreciated that specific functions, and accordingly utilities, of GAM229 correlate with, and may be deduced from, the identity of the target genes which GAM229 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3229] Nucleotide sequences of the GAM229 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM229 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM229 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM229 are further described hereinbelow with reference to Table Table1.

[3230] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM229 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3231]

[3232] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 230 (GAM230) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3233] GAM230 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM230 was detected is described hereinabove with reference to Figs. 2-8.

[3234] GAM230 gene, herein designated GAM GENE, and GAM230 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3235] GAM230 gene, herein designated GAM GENE, encodes a GAM230 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM230 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM230 precursor RNA is designated SEQ ID:3409, and is provided hereinbelow with reference to the sequence listing part.

[3236] GAM230 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM230 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3237] An enzyme complex designated DICER COMPLEX, dices the GAM230 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM230 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 93%) nucleotide sequence of GAM230 RNA is designated SEQ ID:3410, and is provided hereinbelow with reference to the sequence listing part.

[3238] GAM230 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM230 target RNA, herein designated GAM TARGET RNA. GAM230 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3239] GAM230 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM230 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM230 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM230 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM230 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3240] The complementary binding of GAM230 RNA, herein designated GAM RNA, to target binding sites on GAM230 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM230 target RNA, herein designated GAM TARGET RNA, into GAM230 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3241] It is appreciated that GAM230 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM230 target genes. The mRNA of each one of this plu-

rality of GAM230 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM230 RNA, herein designated GAM RNA, and which when bound by GAM230 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM230 target proteins.

[3242] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM230 gene, herein designated GAM GENE, on one or more GAM230 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3243] It is yet further appreciated that specific functions, and accordingly utilities, of GAM230 correlate with, and may be deduced from, the identity of the target genes which GAM230 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3244] Nucleotide sequences of the GAM230 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM230 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM230 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM230 are further described hereinbelow with reference to Table Table1.

[3245] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM230 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3246]

[3247] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 231 (GAM231) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[3248] GAM231 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM231 was detected is described hereinabove with reference to Figs. 2-8.

[3249] GAM231 gene, herein designated GAM GENE, and GAM231 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3250] GAM231 gene, herein designated GAM GENE, encodes a GAM231 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM231 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM231 precursor RNA is designated SEQ ID:3419, and is provided hereinbelow with reference to the sequence listing part.

[3251] GAM231 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM231 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3252] An enzyme complex designated DICER COMPLEX, dices the GAM231 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM231 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM231 RNA is designated SEQ ID:3420, and is provided hereinbelow with reference to the sequence listing part.

[3253] GAM231 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM231 target RNA, herein designated GAM TARGET RNA.
GAM231 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[3254] GAM231 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM231 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM231 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM231 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM231 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3255] The complementary binding of GAM231 RNA, herein designated GAM RNA, to target binding sites on GAM231 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM231 target RNA, herein designated GAM TARGET RNA, into GAM231 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3256] It is appreciated that GAM231 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM231 target genes. The mRNA of each one of this plurality of GAM231 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM231 RNA, herein designated GAM RNA, and which when bound by GAM231 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM231 target proteins.

[3257] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM231 gene, herein designated GAM GENE, on one or more GAM231 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3258] It is yet further appreciated that specific functions, and accordingly utilities, of GAM231 correlate with, and may be deduced from, the identity of the target genes which GAM231 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3259] Nucleotide sequences of the GAM231 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM231 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM231 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM231 are further described hereinbelow with reference to Table Table1.

[3260] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM231 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3261]

[3262] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 232 (GAM232) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3263] GAM232 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM232 was detected is described hereinabove with reference to Figs. 2-8.

[3264] GAM232 gene, herein designated GAM GENE, and GAM232 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3265] GAM232 gene, herein designated GAM GENE, encodes a GAM232 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM232 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM232 precursor RNA is designated SEQ ID:3431, and is provided hereinbelow with reference to the sequence listing part.

[3266] GAM232 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM232 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3267] An enzyme complex designated DICER COMPLEX, dices the GAM232 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM232 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of

GAM232 RNA is designated SEQ ID:3432, and is provided hereinbelow with reference to the sequence listing part.

[3268] GAM232 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM232 target RNA, herein designated GAM TARGET RNA. GAM232 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3269] GAM232 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM232 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM232 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM232 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM232 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3270] The complementary binding of GAM232 RNA, herein designated GAM RNA, to target binding sites on GAM232 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM232 target RNA, herein designated GAM TARGET RNA, into GAM232 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3271] It is appreciated that GAM232 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM232 target genes. The mRNA of each one of this plurality of GAM232 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM232 RNA, herein designated GAM RNA, and which when bound by GAM232 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM232 target proteins.

[3272] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM232 gene, herein designated GAM GENE, on one or more GAM232 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3273] It is yet further appreciated that specific functions, and accordingly utilities, of GAM232 correlate with, and may be deduced from, the identity of the target genes which GAM232 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3274] Nucleotide sequences of the GAM232 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM232 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM232 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM232 are further described hereinbelow with reference to Table Table1.

[3275] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM232 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3276]

[3277] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 233 (GAM233) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3278] GAM233 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM233 was detected is described hereinabove

with reference to Figs. 2-8.

[3279] GAM233 gene, herein designated GAM GENE, and GAM233 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3280] GAM233 gene, herein designated GAM GENE, encodes a GAM233 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM233 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM233 precursor RNA is designated SEQ ID:3449, and is provided hereinbelow with reference to the sequence listing part.

[3281] GAM233 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM233 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3282] An enzyme complex designated DICER COMPLEX, dices the GAM233 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM233 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 81%) nucleotide sequence of GAM233 RNA is designated SEQ ID:3450, and is provided hereinbelow with reference to the sequence listing part.

[3283] GAM233 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM233 target RNA, herein designated GAM TARGET RNA. GAM233 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3284] GAM233 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM233 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM233 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM233 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM233 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3285] The complementary binding of GAM233 RNA, herein designated GAM RNA, to target binding sites on GAM233 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM233 target RNA, herein designated GAM TARGET RNA, into GAM233 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[3286] It is appreciated that GAM233 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM233 target genes. The mRNA of each one of this plurality of GAM233 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM233 RNA, herein designated GAM RNA, and which when bound by GAM233 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM233 target proteins.

[3287] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM233 gene, herein designated GAM GENE, on one or more GAM233 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3288] It is yet further appreciated that specific functions, and accordingly utilities, of GAM233 correlate with, and may be deduced from, the identity of the target genes which GAM233 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3289] Nucleotide sequences of the GAM233 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM233 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM233 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM233 are further described hereinbelow with reference to Table Table1.

[3290] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM233 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3291]

[3292] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 234 (GAM234) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3293] GAM234 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM234 was detected is described hereinabove with reference to Figs. 2-8.

[3294] GAM234 gene, herein designated GAM GENE, and GAM234 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3295] GAM234 gene, herein designated GAM GENE, encodes a GAM234 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM234 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM234 precursor RNA is designated SEQ ID:3465, and is provided hereinbelow with reference to the sequence listing part.

[3296] GAM234 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM234 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3297] An enzyme complex designated DICER COMPLEX, dices the GAM234 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM234 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 44%) nucleotide sequence of GAM234 RNA is designated SEQ ID:3466, and is provided hereinbelow with reference to the sequence listing part.

[3298] GAM234 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM234 target RNA, herein designated GAM TARGET RNA.

GAM234 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3299] GAM234 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM234 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM234 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM234 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM234 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3300] The complementary binding of GAM234 RNA, herein designated GAM RNA, to target binding sites on GAM234 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM234 target RNA, herein designated GAM TARGET RNA, into GAM234 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3301] It is appreciated that GAM234 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM234 target genes. The mRNA of each one of this plurality of GAM234 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM234 RNA, herein designated GAM RNA, and which when bound by GAM234 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM234 target proteins.

[3302] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM234 gene, herein designated GAM GENE, on one or more GAM234 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3303] It is yet further appreciated that specific functions, and accordingly utilities, of GAM234 correlate with, and may be deduced from, the identity of the target genes which GAM234 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3304] Nucleotide sequences of the GAM234 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM234 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM234 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM234 are further described hereinbelow with reference to Table Table1.

[3305] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM234 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3306]

[3307] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 235 (GAM235) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3308] GAM235 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM235 was detected is described hereinabove with reference to Figs. 2-8.

[3309] GAM235 gene, herein designated GAM GENE, and GAM235 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3310] GAM235 gene, herein designated GAM GENE, encodes a

GAM235 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM235 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM235 precursor RNA is designated SEQ ID:3469, and is provided hereinbelow with reference to the sequence listing part.

[3311] GAM235 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM235 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3312] An enzyme complex designated DICER COMPLEX, dices the GAM235 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM235 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM235 RNA is designated SEQ ID:3470, and is provided hereinbelow with reference to the sequence listing part.

[3313] GAM235 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM235 target RNA, herein designated GAM TARGET RNA. GAM235 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3314] GAM235 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM235 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM235 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM235 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM235 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3315] The complementary binding of GAM235 RNA, herein designated GAM RNA, to target binding sites on GAM235 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM235 target RNA, herein designated GAM TARGET RNA, into GAM235 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3316] It is appreciated that GAM235 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM235 target genes. The mRNA of each one of this plurality of GAM235 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM235 RNA, herein designated GAM RNA, and which when bound by GAM235 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM235 target proteins.

[3317] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM235 gene, herein designated GAM GENE, on one or more GAM235 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3318] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM235 correlate with, and may be deduced from, the identity of the target genes which GAM235 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3319] Nucleotide sequences of the GAM235 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM235 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM235 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM235 are further described hereinbelow with reference to Table Table1.

[3320] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM235 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3321]

[3322] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 236 (GAM236) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[3323] GAM236 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM236 was detected is described hereinabove with reference to Figs. 2-8.

[3324] GAM236 gene, herein designated GAM GENE, and GAM236 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3325] GAM236 gene, herein designated GAM GENE, encodes a GAM236 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM236 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM236 precursor RNA is designated SEQ ID:3494, and is provided hereinbelow with reference to the sequence listing part.

[3326] GAM236 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM236 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3327] An enzyme complex designated DICER COMPLEX, dices the GAM236 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM236 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM236 RNA is designated SEQ ID:3495, and is provided hereinbelow with reference to the sequence listing part.

[3328] GAM236 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM236 target RNA, herein designated GAM TARGET RNA. GAM236 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3329] GAM236 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM236 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM236 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM236 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM236 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3330] The complementary binding of GAM236 RNA, herein designated GAM RNA, to target binding sites on GAM236 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM236 target RNA, herein designated GAM TARGET RNA, into GAM236 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3331] It is appreciated that GAM236 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM236 target genes. The mRNA of each one of this plurality of GAM236 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM236 RNA, herein designated GAM RNA, and which when bound by GAM236 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM236 target proteins.

[3332] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM236 gene, herein designated GAM GENE, on one or more GAM236 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3333] It is yet further appreciated that specific functions, and accordingly utilities, of GAM236 correlate with, and may be deduced from, the identity of the target genes which GAM236 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3334] Nucleotide sequences of the GAM236 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM236 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM236 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM236 are further described hereinbelow with reference to Table Table1.

[3335] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM236 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3336]

[3337] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 237 (GAM237) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3338] GAM237 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM237 was detected is described hereinabove with reference to Figs. 2-8.

[3339] GAM237 gene, herein designated GAM GENE, and GAM237 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3340] GAM237 gene, herein designated GAM GENE, encodes a GAM237 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM237 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM237 precursor RNA is designated SEQ ID:3506, and is provided hereinbelow with reference to the sequence listing part.

[3341] GAM237 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM237 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3342] An enzyme complex designated DICER COMPLEX, dices the GAM237 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM237 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM237 RNA is designated SEQ ID:3507, and is provided

hereinbelow with reference to the sequence listing part.

[3343] GAM237 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM237 target RNA, herein designated GAM TARGET RNA. GAM237 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3344] GAM237 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM237 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM237 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM237 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM237 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3345] The complementary binding of GAM237 RNA, herein designated GAM RNA, to target binding sites on GAM237 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM237 target RNA, herein designated GAM TARGET RNA, into GAM237 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3346] It is appreciated that GAM237 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM237 target genes. The mRNA of each one of this plurality of GAM237 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM237 RNA, herein designated GAM RNA, and which when bound by GAM237 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM237 tar-

get proteins.

[3347] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM237 gene, herein designated GAM GENE, on one or more GAM237 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3348] It is yet further appreciated that specific functions, and accordingly utilities, of GAM237 correlate with, and may be deduced from, the identity of the target genes which GAM237 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3349] Nucleotide sequences of the GAM237 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM237 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM237 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM237 are further described hereinbelow with reference to Table Table1.

[3350] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM237 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3351]

[3352] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 238 (GAM238) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3353] GAM238 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM238 was detected is described hereinabove with reference to Figs. 2-8.

[3354] GAM238 gene, herein designated GAM GENE, and GAM238 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3355] GAM238 gene, herein designated GAM GENE, encodes a GAM238 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM238 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM238 precursor RNA is designated SEQ ID:3522, and is provided hereinbelow with reference to the sequence listing part.

[3356] GAM238 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM238 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3357] An enzyme complex designated DICER COMPLEX, dices

the GAM238 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM238 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM238 RNA is designated SEQ ID:3523, and is provided hereinbelow with reference to the sequence listing part.

[3358] GAM238 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM238 target RNA, herein designated GAM TARGET RNA. GAM238 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3359] GAM238 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM238 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM238

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM238 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM238 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3360] The complementary binding of GAM238 RNA, herein designated GAM RNA, to target binding sites on GAM238 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM238 target RNA, herein designated GAM TARGET RNA, into GAM238 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3361] It is appreciated that GAM238 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM238 target genes. The mRNA of each one of this plurality of GAM238 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM238 RNA, herein designated GAM RNA, and which when bound by GAM238 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM238 target proteins.

[3362] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM238 gene, herein designated GAM GENE, on one or more GAM238 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3363] It is yet further appreciated that specific functions, and accordingly utilities, of GAM238 correlate with, and may be deduced from, the identity of the target genes which GAM238 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3364] Nucleotide sequences of the GAM238 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM238 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM238 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM238 are further described hereinbelow with reference to Table Table1.

[3365] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM238 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3366]

[3367] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 239 (GAM239) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3368] GAM239 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM239 was detected is described hereinabove with reference to Figs. 2-8.

[3369] GAM239 gene, herein designated GAM GENE, and GAM239 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3370] GAM239 gene, herein designated GAM GENE, encodes a GAM239 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM239 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM239 precursor RNA is designated SEQ ID:3543, and is provided hereinbelow with reference to the sequence listing part.

[3371] GAM239 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM239 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3372] An enzyme complex designated DICER COMPLEX, dices the GAM239 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM239 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM239 RNA is designated SEQ ID:3544, and is provided hereinbelow with reference to the sequence listing part.

[3373] GAM239 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM239 target RNA, herein designated GAM TARGET RNA. GAM239 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3374] GAM239 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM239 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM239 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM239 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM239 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[3375] The complementary binding of GAM239 RNA, herein designated GAM RNA, to target binding sites on GAM239 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM239 target RNA, herein designated GAM TARGET RNA, into GAM239 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3376] It is appreciated that GAM239 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM239 target genes. The mRNA of each one of this plurality of GAM239 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM239 RNA, herein designated GAM RNA, and which when bound by GAM239 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM239 target proteins.

[3377] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM239 gene, herein designated GAM GENE, on one or

more GAM239 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3378] It is yet further appreciated that specific functions, and accordingly utilities, of GAM239 correlate with, and may be deduced from, the identity of the target genes which GAM239 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3379] Nucleotide sequences of the GAM239 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM239 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM239 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM239 are further de-

scribed hereinbelow with reference to Table Table1.

[3380] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM239 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3381]

[3382] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 240 (GAM240) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3383] GAM240 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM240 was detected is described hereinabove with reference to Figs. 2-8.

[3384] GAM240 gene, herein designated GAM GENE, and GAM240 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3385] GAM240 gene, herein designated GAM GENE, encodes a GAM240 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM240 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM240 precursor RNA is designated SEQ ID:3570, and is provided hereinbelow with reference to the sequence listing part.

[3386] GAM240 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM240 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3387] An enzyme complex designated DICER COMPLEX, dices the GAM240 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM240 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM240 RNA is designated SEQ ID:3571, and is provided hereinbelow with reference to the sequence listing part.

[3388] GAM240 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM240 target RNA, herein designated GAM TARGET RNA. GAM240 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3389] GAM240 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM240 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM240 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM240 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM240 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3390] The complementary binding of GAM240 RNA, herein designated GAM RNA, to target binding sites on GAM240 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM240 target RNA, herein designated GAM TARGET RNA, into GAM240 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3391] It is appreciated that GAM240 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM240 target genes. The mRNA of each one of this plurality of GAM240 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM240 RNA, herein designated GAM RNA, and which when bound by GAM240 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM240 target proteins.

[3392] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM240 gene, herein designated GAM GENE, on one or more GAM240 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3393] It is yet further appreciated that specific functions, and accordingly utilities, of GAM240 correlate with, and may

be deduced from, the identity of the target genes which GAM240 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3394] Nucleotide sequences of the GAM240 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM240 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM240 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM240 are further described hereinbelow with reference to Table Table1.

[3395] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM240 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3396]

[3397] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 241 (GAM241) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3398] GAM241 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM241 was detected is described hereinabove with reference to Figs. 2-8.

[3399] GAM241 gene, herein designated GAM GENE, and GAM241 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3400] GAM241 gene, herein designated GAM GENE, encodes a GAM241 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM241 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM241 precursor RNA is designated SEQ ID:3578, and is provided hereinbelow with reference to the sequence listing part.

[3401] GAM241 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM241 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3402] An enzyme complex designated DICER COMPLEX, dices the GAM241 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM241 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM241 RNA is designated SEQ ID:3579, and is provided hereinbelow with reference to the sequence listing part.

[3403] GAM241 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM241 target RNA, herein designated GAM TARGET RNA. GAM241 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3404] GAM241 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM241 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM241 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM241 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM241 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3405] The complementary binding of GAM241 RNA, herein designated GAM RNA, to target binding sites on GAM241 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM241 target RNA, herein designated GAM TARGET RNA, into GAM241 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3406] It is appreciated that GAM241 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM241 target genes. The mRNA of each one of this plurality of GAM241 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM241 RNA, herein designated GAM RNA, and which when bound by GAM241 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM241 target proteins.

[3407] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM241 gene, herein designated GAM GENE, on one or more GAM241 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3408] It is yet further appreciated that specific functions, and accordingly utilities, of GAM241 correlate with, and may be deduced from, the identity of the target genes which GAM241 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3409] Nucleotide sequences of the GAM241 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM241 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM241 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM241 are further described hereinbelow with reference to Table Table1.

[3410] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM241 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3411]

[3412] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 242 (GAM242) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3413] GAM242 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM242 was detected is described hereinabove with reference to Figs. 2-8.

[3414] GAM242 gene, herein designated GAM GENE, and GAM242 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3415] GAM242 gene, herein designated GAM GENE, encodes a GAM242 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM242 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM242 precursor RNA is desig-

nated SEQ ID:3584, and is provided hereinbelow with reference to the sequence listing part.

[3416] GAM242 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM242 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3417] An enzyme complex designated DICER COMPLEX, dices the GAM242 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM242 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 20%) nucleotide sequence of GAM242 RNA is designated SEQ ID:3585, and is provided hereinbelow with reference to the sequence listing part.

[3418] GAM242 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM242 target RNA, herein designated GAM TARGET RNA. GAM242 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3419] GAM242 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM242 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM242 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM242 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM242

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3420] The complementary binding of GAM242 RNA, herein designated GAM RNA, to target binding sites on GAM242 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM242 target RNA, herein designated GAM TARGET RNA, into GAM242 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3421] It is appreciated that GAM242 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM242 target genes. The mRNA of each one of this plurality of GAM242 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM242 RNA, herein designated GAM RNA, and which when bound by GAM242 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM242 target proteins.

[3422] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM242 gene, herein designated GAM GENE, on one or more GAM242 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3423] It is yet further appreciated that specific functions, and accordingly utilities, of GAM242 correlate with, and may be deduced from, the identity of the target genes which GAM242 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3424] Nucleotide sequences of the GAM242 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM242 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM242 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM242 are further described hereinbelow with reference to Table Table1.

[3425] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM242 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3426]

[3427] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 243 (GAM243) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3428] GAM243 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM243 was detected is described hereinabove with reference to Figs. 2-8.

[3429] GAM243 gene, herein designated GAM GENE, and

GAM243 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3430] GAM243 gene, herein designated GAM GENE, encodes a GAM243 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM243 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM243 precursor RNA is designated SEQ ID:3593, and is provided hereinbelow with reference to the sequence listing part.

[3431] GAM243 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM243 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3432] An enzyme complex designated DICER COMPLEX, dices the GAM243 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM243 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 45%) nucleotide sequence of GAM243 RNA is designated SEQ ID:3594, and is provided hereinbelow with reference to the sequence listing part.

[3433] GAM243 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM243 target RNA, herein designated GAM TARGET RNA. GAM243 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3434] GAM243 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM243 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM243 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM243 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM243 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3435] The complementary binding of GAM243 RNA, herein designated GAM RNA, to target binding sites on GAM243 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM243 target RNA, herein designated GAM TARGET RNA, into GAM243 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3436] It is appreciated that GAM243 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM243 target genes. The mRNA of each one of this plurality of GAM243 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM243 RNA, herein designated GAM RNA, and which when bound by GAM243 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM243 target proteins.

[3437] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM243 gene, herein designated GAM GENE, on one or more GAM243 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3438] It is yet further appreciated that specific functions, and accordingly utilities, of GAM243 correlate with, and may be deduced from, the identity of the target genes which GAM243 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3439] Nucleotide sequences of the GAM243 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM243 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM243 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM243 are further described hereinbelow with reference to Table Table1.

[3440] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM243 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3441]

[3442] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 244 (GAM244) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3443] GAM244 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM244 was detected is described hereinabove with reference to Figs. 2-8.

[3444] GAM244 gene, herein designated GAM GENE, and GAM244 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3445] GAM244 gene, herein designated GAM GENE, encodes a GAM244 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM244 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM244 precursor RNA is designated SEQ ID:3622, and is provided hereinbelow with reference to the sequence listing part.

[3446] GAM244 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM244 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3447] An enzyme complex designated DICER COMPLEX, dices the GAM244 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM244 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM244 RNA is designated SEQ ID:3623, and is provided hereinbelow with reference to the sequence listing part.

[3448] GAM244 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM244 target RNA, herein designated GAM TARGET RNA. GAM244 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3449] GAM244 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM244 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM244 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM244 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM244 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3450] The complementary binding of GAM244 RNA, herein designated GAM RNA, to target binding sites on GAM244 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM244 target RNA, herein designated GAM TARGET RNA, into GAM244 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3451] It is appreciated that GAM244 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM244 target genes. The mRNA of each one of this plurality of GAM244 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM244 RNA, herein designated GAM RNA, and which when bound by GAM244 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM244 target proteins.

[3452] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM244 gene, herein designated GAM GENE, on one or more GAM244 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3453] It is yet further appreciated that specific functions, and accordingly utilities, of GAM244 correlate with, and may be deduced from, the identity of the target genes which GAM244 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3454] Nucleotide sequences of the GAM244 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM244 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM244 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM244 are further described hereinbelow with reference to Table Table1.

[3455] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM244 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3456]

[3457] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 245 (GAM245) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3458] GAM245 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM245 was detected is described hereinabove with reference to Figs. 2-8.

[3459] GAM245 gene, herein designated GAM GENE, and GAM245 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3460] GAM245 gene, herein designated GAM GENE, encodes a GAM245 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM245 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM245 precursor RNA is designated SEQ ID:3658, and is provided hereinbelow with reference to the sequence listing part.

[3461] GAM245 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM245 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3462] An enzyme complex designated DICER COMPLEX, dices the GAM245 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM245 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM245 RNA is designated SEQ ID:3623, and is provided hereinbelow with reference to the sequence listing part.

[3463] GAM245 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM245 target RNA, herein designated GAM TARGET RNA. GAM245 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3464] GAM245 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM245 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM245 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM245 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM245 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3465] The complementary binding of GAM245 RNA, herein designated GAM RNA, to target binding sites on GAM245 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM245 target RNA, herein designated GAM TARGET RNA, into GAM245 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3466] It is appreciated that GAM245 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM245 target genes. The mRNA of each one of this plurality of GAM245 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM245 RNA,

herein designated GAM RNA, and which when bound by GAM245 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM245 target proteins.

[3467] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM245 gene, herein designated GAM GENE, on one or more GAM245 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3468] It is yet further appreciated that specific functions, and accordingly utilities, of GAM245 correlate with, and may be deduced from, the identity of the target genes which

GAM245 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3469] Nucleotide sequences of the GAM245 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM245 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM245 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM245 are further described hereinbelow with reference to Table Table1.

[3470] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM245 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3471]

[3472] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 246 (GAM246) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3473] GAM246 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM246 was detected is described hereinabove with reference to Figs. 2-8.

[3474] GAM246 gene, herein designated GAM GENE, and GAM246 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3475] GAM246 gene, herein designated GAM GENE, encodes a GAM246 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM246 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM246 precursor RNA is designated SEQ ID:3660, and is provided hereinbelow with reference to the sequence listing part.

[3476] GAM246 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM246 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[3477] An enzyme complex designated DICER COMPLEX, dices the GAM246 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM246 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM246 RNA is designated SEQ ID:3661, and is provided hereinbelow with reference to the sequence listing part.

[3478] GAM246 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM246 target RNA, herein designated GAM TARGET RNA.
GAM246 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3479] GAM246 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM246 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM246 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM246 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM246 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3480] The complementary binding of GAM246 RNA, herein designated GAM RNA, to target binding sites on GAM246 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM246 target RNA, herein desig-

nated GAM TARGET RNA, into GAM246 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3481] It is appreciated that GAM246 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM246 target genes. The mRNA of each one of this plurality of GAM246 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM246 RNA, herein designated GAM RNA, and which when bound by GAM246 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM246 target proteins.

[3482] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM246 gene, herein designated GAM GENE, on one or more GAM246 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3483] It is yet further appreciated that specific functions, and accordingly utilities, of GAM246 correlate with, and may be deduced from, the identity of the target genes which GAM246 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3484] Nucleotide sequences of the GAM246 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM246 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM246 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM246 are further described hereinbelow with reference to Table Table1.

[3485] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM246 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[3486]

[3487] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 247 (GAM247) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3488] GAM247 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM247 was detected is described hereinabove with reference to Figs. 2-8.

[3489] GAM247 gene, herein designated GAM GENE, and GAM247 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3490] GAM247 gene, herein designated GAM GENE, encodes a GAM247 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM247 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM247 precursor RNA is designated SEQ ID:3675, and is provided hereinbelow with ref-

erence to the sequence listing part.

[3491] GAM247 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM247 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3492] An enzyme complex designated DICER COMPLEX, dices the GAM247 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM247 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM247 RNA is designated SEQ ID:3676, and is provided hereinbelow with reference to the sequence listing part.

[3493] GAM247 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM247 target RNA, herein designated GAM TARGET RNA.

GAM247 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3494] GAM247 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM247 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM247 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM247 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM247 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3495] The complementary binding of GAM247 RNA, herein designated GAM RNA, to target binding sites on GAM247 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM247 target RNA, herein designated GAM TARGET RNA, into GAM247 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3496] It is appreciated that GAM247 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM247 target genes. The mRNA of each one of this plurality of GAM247 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM247 RNA, herein designated GAM RNA, and which when bound by GAM247 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM247 target proteins.

[3497] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM247 gene, herein designated GAM GENE, on one or more GAM247 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3498] It is yet further appreciated that specific functions, and accordingly utilities, of GAM247 correlate with, and may be deduced from, the identity of the target genes which GAM247 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3499] Nucleotide sequences of the GAM247 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM247 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM247 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM247 are further described hereinbelow with reference to Table Table1.

[3500] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM247 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3501]

[3502] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 248 (GAM248) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3503] GAM248 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM248 was detected is described hereinabove with reference to Figs. 2-8.

[3504] GAM248 gene, herein designated GAM GENE, and GAM248 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[3505] GAM248 gene, herein designated GAM GENE, encodes a GAM248 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM248 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM248 precursor RNA is designated SEQ ID:3696, and is provided hereinbelow with reference to the sequence listing part.

[3506] GAM248 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM248 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3507] An enzyme complex designated DICER COMPLEX, dices the GAM248 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM248 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM248 RNA is designated SEQ ID:3697, and is provided hereinbelow with reference to the sequence listing part.

[3508] GAM248 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM248 target RNA, herein designated GAM TARGET RNA. GAM248 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3509] GAM248 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM248 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM248 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM248 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM248 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3510] The complementary binding of GAM248 RNA, herein designated GAM RNA, to target binding sites on GAM248 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM248 target RNA, herein designated GAM TARGET RNA, into GAM248 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3511] It is appreciated that GAM248 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM248 target genes. The mRNA of each one of this plurality of GAM248 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM248 RNA, herein designated GAM RNA, and which when bound by GAM248 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM248 target proteins.

[3512] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM248 gene, herein designated GAM GENE, on one or more GAM248 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[3513] It is yet further appreciated that specific functions, and accordingly utilities, of GAM248 correlate with, and may be deduced from, the identity of the target genes which GAM248 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3514] Nucleotide sequences of the GAM248 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM248 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM248 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM248 are further described hereinbelow with reference to Table Table1.

[3515] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM248 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3516]

[3517] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

249 (GAM249) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3518] GAM249 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM249 was detected is described hereinabove with reference to Figs. 2-8.

[3519] GAM249 gene, herein designated GAM GENE, and GAM249 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3520] GAM249 gene, herein designated GAM GENE, encodes a GAM249 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM249 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM249 precursor RNA is designated SEQ ID:3709, and is provided hereinbelow with reference to the sequence listing part.

[3521] GAM249 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM249 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3522] An enzyme complex designated DICER COMPLEX, dices the GAM249 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM249 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM249 RNA is designated SEQ ID:3710, and is provided hereinbelow with reference to the sequence listing part.

[3523] GAM249 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM249 target RNA, herein designated GAM TARGET RNA. GAM249 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3524] GAM249 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM249 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM249 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM249 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM249 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3525] The complementary binding of GAM249 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM249 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM249 target RNA, herein designated GAM TARGET RNA, into GAM249 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3526] It is appreciated that GAM249 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM249 target genes. The mRNA of each one of this plurality of GAM249 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM249 RNA, herein designated GAM RNA, and which when bound by GAM249 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM249 target proteins.

[3527] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM249 gene, herein designated GAM GENE, on one or more GAM249 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3528] It is yet further appreciated that specific functions, and accordingly utilities, of GAM249 correlate with, and may be deduced from, the identity of the target genes which GAM249 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3529] Nucleotide sequences of the GAM249 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM249 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM249 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM249 are further described hereinbelow with reference to Table Table1.

[3530] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM249 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3531]

[3532] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 250 (GAM250) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3533] GAM250 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM250 was detected is described hereinabove with reference to Figs. 2-8.

[3534] GAM250 gene, herein designated GAM GENE, and GAM250 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3535] GAM250 gene, herein designated GAM GENE, encodes a GAM250 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM250 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM250 precursor RNA is designated SEQ ID:3717, and is provided hereinbelow with reference to the sequence listing part.

[3536] GAM250 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM250 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3537] An enzyme complex designated DICER COMPLEX, dices the GAM250 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM250 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 63%) nucleotide sequence of GAM250 RNA is designated SEQ ID:3718, and is provided hereinbelow with reference to the sequence listing part.

[3538] GAM250 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM250 target RNA, herein designated GAM TARGET RNA. GAM250 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3539] GAM250 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM250 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM250 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM250 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM250 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3540] The complementary binding of GAM250 RNA, herein designated GAM RNA, to target binding sites on GAM250 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM250 target RNA, herein designated GAM TARGET RNA, into GAM250 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3541] It is appreciated that GAM250 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM250 target genes. The mRNA of each one of this plurality of GAM250 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM250 RNA, herein designated GAM RNA, and which when bound by

GAM250 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM250 target proteins.

[3542] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM250 gene, herein designated GAM GENE, on one or more GAM250 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3543] It is yet further appreciated that specific functions, and accordingly utilities, of GAM250 correlate with, and may be deduced from, the identity of the target genes which GAM250 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[3544] Nucleotide sequences of the GAM250 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM250 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM250 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM250 are further described hereinbelow with reference to Table Table1.

[3545] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM250 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3546]

[3547] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 251 (GAM251) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3548] GAM251 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM251 was detected is described hereinabove with reference to Figs. 2-8.

[3549] GAM251 gene, herein designated GAM GENE, and GAM251 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3550] GAM251 gene, herein designated GAM GENE, encodes a GAM251 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM251 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM251 precursor RNA is designated SEQ ID:3735, and is provided hereinbelow with reference to the sequence listing part.

[3551] GAM251 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM251 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[3552] An enzyme complex designated DICER COMPLEX, dices the GAM251 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM251 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 28%) nucleotide sequence of GAM251 RNA is designated SEQ ID:3736, and is provided hereinbelow with reference to the sequence listing part.

[3553] GAM251 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM251 target RNA, herein designated GAM TARGET RNA. GAM251 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3554] GAM251 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM251 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM251 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM251 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM251 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3555] The complementary binding of GAM251 RNA, herein designated GAM RNA, to target binding sites on GAM251 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM251 target RNA, herein designated GAM TARGET RNA, into GAM251 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3556] It is appreciated that GAM251 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM251 target genes. The mRNA of each one of this plurality of GAM251 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM251 RNA, herein designated GAM RNA, and which when bound by GAM251 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM251 target proteins.

[3557] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM251 gene, herein designated GAM GENE, on one or more GAM251 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3558] It is yet further appreciated that specific functions, and accordingly utilities, of GAM251 correlate with, and may be deduced from, the identity of the target genes which GAM251 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3559] Nucleotide sequences of the GAM251 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM251 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM251 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM251 are further described hereinbelow with reference to Table Table1.

[3560] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM251 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3561]

[3562] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 252 (GAM252) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3563] GAM252 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM252 was detected is described hereinabove with reference to Figs. 2-8.

[3564] GAM252 gene, herein designated GAM GENE, and GAM252 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3565] GAM252 gene, herein designated GAM GENE, encodes a GAM252 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM252 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM252 precursor RNA is designated SEQ ID:3745, and is provided hereinbelow with reference to the sequence listing part.

[3566] GAM252 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM252 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3567] An enzyme complex designated DICER COMPLEX, dices the GAM252 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM252 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM252 RNA is designated SEQ ID:3746, and is provided hereinbelow with reference to the sequence listing part.

[3568] GAM252 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM252

target RNA, herein designated GAM TARGET RNA.

GAM252 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3569] GAM252 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM252 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM252 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM252 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM252 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3570] The complementary binding of GAM252 RNA, herein designated GAM RNA, to target binding sites on GAM252 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM252 target RNA, herein designated GAM TARGET RNA, into GAM252 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3571] It is appreciated that GAM252 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM252 target genes. The mRNA of each one of this plurality of GAM252 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM252 RNA, herein designated GAM RNA, and which when bound by GAM252 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM252 target proteins.

[3572] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM252 gene, herein designated GAM GENE, on one or more GAM252 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3573] It is yet further appreciated that specific functions, and accordingly utilities, of GAM252 correlate with, and may be deduced from, the identity of the target genes which GAM252 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3574] Nucleotide sequences of the GAM252 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM252 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM252 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM252 are further described hereinbelow with reference to Table Table1.

[3575] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM252 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3576]

[3577] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 253 (GAM253) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3578] GAM253 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM253 was detected is described hereinabove with reference to Figs. 2-8.

[3579] GAM253 gene, herein designated GAM GENE, and GAM253 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3580] GAM253 gene, herein designated GAM GENE, encodes a GAM253 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM253 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM253 precursor RNA is designated SEQ ID:3761, and is provided hereinbelow with reference to the sequence listing part.

[3581] GAM253 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM253 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3582] An enzyme complex designated DICER COMPLEX, dices the GAM253 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM253 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM253 RNA is designated SEQ ID:3762, and is provided hereinbelow with reference to the sequence listing part.

[3583] GAM253 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM253 target RNA, herein designated GAM TARGET RNA. GAM253 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3584] GAM253 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM253 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM253 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM253 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM253 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3585] The complementary binding of GAM253 RNA, herein designated GAM RNA, to target binding sites on GAM253 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM253 target RNA, herein designated GAM TARGET RNA, into GAM253 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3586] It is appreciated that GAM253 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM253 target genes. The mRNA of each one of this plu-

rality of GAM253 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM253 RNA, herein designated GAM RNA, and which when bound by GAM253 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM253 target proteins.

[3587] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM253 gene, herein designated GAM GENE, on one or more GAM253 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3588] It is yet further appreciated that specific functions, and accordingly utilities, of GAM253 correlate with, and may be deduced from, the identity of the target genes which GAM253 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3589] Nucleotide sequences of the GAM253 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM253 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM253 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM253 are further described hereinbelow with reference to Table Table1.

[3590] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM253 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3591]

[3592] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 254 (GAM254) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[3593] GAM254 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM254 was detected is described hereinabove with reference to Figs. 2-8.

[3594] GAM254 gene, herein designated GAM GENE, and GAM254 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3595] GAM254 gene, herein designated GAM GENE, encodes a GAM254 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM254 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM254 precursor RNA is designated SEQ ID:3779, and is provided hereinbelow with reference to the sequence listing part.

[3596] GAM254 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM254 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3597] An enzyme complex designated DICER COMPLEX, dices the GAM254 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM254 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM254 RNA is designated SEQ ID:3780, and is provided hereinbelow with reference to the sequence listing part.

[3598] GAM254 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM254 target RNA, herein designated GAM TARGET RNA.
GAM254 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[3599] GAM254 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM254 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM254 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM254 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM254 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3600] The complementary binding of GAM254 RNA, herein designated GAM RNA, to target binding sites on GAM254 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM254 target RNA, herein designated GAM TARGET RNA, into GAM254 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3601] It is appreciated that GAM254 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM254 target genes. The mRNA of each one of this plurality of GAM254 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM254 RNA, herein designated GAM RNA, and which when bound by GAM254 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM254 target proteins.

[3602] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM254 gene, herein designated GAM GENE, on one or more GAM254 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3603] It is yet further appreciated that specific functions, and accordingly utilities, of GAM254 correlate with, and may be deduced from, the identity of the target genes which GAM254 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3604] Nucleotide sequences of the GAM254 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM254 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM254 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM254 are further described hereinbelow with reference to Table Table1.

[3605] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM254 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3606]

[3607] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 255 (GAM255) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3608] GAM255 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM255 was detected is described hereinabove with reference to Figs. 2-8.

[3609] GAM255 gene, herein designated GAM GENE, and GAM255 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3610] GAM255 gene, herein designated GAM GENE, encodes a GAM255 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM255 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM255 precursor RNA is designated SEQ ID:3815, and is provided hereinbelow with reference to the sequence listing part.

[3611] GAM255 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM255 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3612] An enzyme complex designated DICER COMPLEX, dices the GAM255 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM255 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of

GAM255 RNA is designated SEQ ID:3816, and is provided hereinbelow with reference to the sequence listing part.

[3613] GAM255 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM255 target RNA, herein designated GAM TARGET RNA. GAM255 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3614] GAM255 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM255 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM255 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM255 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM255 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3615] The complementary binding of GAM255 RNA, herein designated GAM RNA, to target binding sites on GAM255 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM255 target RNA, herein designated GAM TARGET RNA, into GAM255 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3616] It is appreciated that GAM255 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM255 target genes. The mRNA of each one of this plurality of GAM255 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM255 RNA, herein designated GAM RNA, and which when bound by GAM255 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM255 target proteins.

[3617] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM255 gene, herein designated GAM GENE, on one or more GAM255 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3618] It is yet further appreciated that specific functions, and accordingly utilities, of GAM255 correlate with, and may be deduced from, the identity of the target genes which GAM255 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3619] Nucleotide sequences of the GAM255 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM255 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM255 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM255 are further described hereinbelow with reference to Table Table1.

[3620] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM255 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3621]

[3622] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 256 (GAM256) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3623] GAM256 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM256 was detected is described hereinabove

with reference to Figs. 2-8.

[3624] GAM256 gene, herein designated GAM GENE, and GAM256 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3625] GAM256 gene, herein designated GAM GENE, encodes a GAM256 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM256 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM256 precursor RNA is designated SEQ ID:3848, and is provided hereinbelow with reference to the sequence listing part.

[3626] GAM256 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM256 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3627] An enzyme complex designated DICER COMPLEX, dices the GAM256 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM256 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM256 RNA is designated SEQ ID:3849, and is provided hereinbelow with reference to the sequence listing part.

[3628] GAM256 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM256 target RNA, herein designated GAM TARGET RNA. GAM256 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3629] GAM256 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM256 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM256 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM256 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM256 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3630] The complementary binding of GAM256 RNA, herein designated GAM RNA, to target binding sites on GAM256 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM256 target RNA, herein designated GAM TARGET RNA, into GAM256 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[3631] It is appreciated that GAM256 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM256 target genes. The mRNA of each one of this plurality of GAM256 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM256 RNA, herein designated GAM RNA, and which when bound by GAM256 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM256 target proteins.

[3632] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM256 gene, herein designated GAM GENE, on one or more GAM256 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3633] It is yet further appreciated that specific functions, and accordingly utilities, of GAM256 correlate with, and may be deduced from, the identity of the target genes which GAM256 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3634] Nucleotide sequences of the GAM256 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM256 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM256 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM256 are further described hereinbelow with reference to Table Table1.

[3635] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM256 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3636]

[3637] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 257 (GAM257) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3638] GAM257 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM257 was detected is described hereinabove with reference to Figs. 2-8.

[3639] GAM257 gene, herein designated GAM GENE, and GAM257 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3640] GAM257 gene, herein designated GAM GENE, encodes a GAM257 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM257 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM257 precursor RNA is designated SEQ ID:3858, and is provided hereinbelow with reference to the sequence listing part.

[3641] GAM257 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM257 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3642] An enzyme complex designated DICER COMPLEX, dices the GAM257 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM257 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 32%) nucleotide sequence of GAM257 RNA is designated SEQ ID:3859, and is provided hereinbelow with reference to the sequence listing part.

[3643] GAM257 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM257 target RNA, herein designated GAM TARGET RNA.

GAM257 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3644] GAM257 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM257 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM257 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM257 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM257 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3645] The complementary binding of GAM257 RNA, herein designated GAM RNA, to target binding sites on GAM257 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM257 target RNA, herein designated GAM TARGET RNA, into GAM257 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3646] It is appreciated that GAM257 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM257 target genes. The mRNA of each one of this plurality of GAM257 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM257 RNA, herein designated GAM RNA, and which when bound by GAM257 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM257 target proteins.

[3647] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM257 gene, herein designated GAM GENE, on one or more GAM257 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3648] It is yet further appreciated that specific functions, and accordingly utilities, of GAM257 correlate with, and may be deduced from, the identity of the target genes which GAM257 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3649] Nucleotide sequences of the GAM257 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM257 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM257 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM257 are further described hereinbelow with reference to Table Table1.

[3650] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM257 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3651]

[3652] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 258 (GAM258) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3653] GAM258 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM258 was detected is described hereinabove with reference to Figs. 2-8.

[3654] GAM258 gene, herein designated GAM GENE, and GAM258 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3655] GAM258 gene, herein designated GAM GENE, encodes a

GAM258 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM258 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM258 precursor RNA is designated SEQ ID:3884, and is provided hereinbelow with reference to the sequence listing part.

[3656] GAM258 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM258 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3657] An enzyme complex designated DICER COMPLEX, dices the GAM258 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM258 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 25%) nucleotide sequence of GAM258 RNA is designated SEQ ID:3885, and is provided hereinbelow with reference to the sequence listing part.

[3658] GAM258 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM258 target RNA, herein designated GAM TARGET RNA. GAM258 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3659] GAM258 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM258 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM258 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM258 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM258 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3660] The complementary binding of GAM258 RNA, herein designated GAM RNA, to target binding sites on GAM258 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM258 target RNA, herein designated GAM TARGET RNA, into GAM258 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3661] It is appreciated that GAM258 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM258 target genes. The mRNA of each one of this plurality of GAM258 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM258 RNA, herein designated GAM RNA, and which when bound by GAM258 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM258 target proteins.

[3662] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM258 gene, herein designated GAM GENE, on one or more GAM258 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3663] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM258 correlate with, and may be deduced from, the identity of the target genes which GAM258 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3664] Nucleotide sequences of the GAM258 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM258 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM258 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM258 are further described hereinbelow with reference to Table Table1.

[3665] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM258 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3666]

[3667] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 259 (GAM259) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[3668] GAM259 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM259 was detected is described hereinabove with reference to Figs. 2-8.

[3669] GAM259 gene, herein designated GAM GENE, and GAM259 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3670] GAM259 gene, herein designated GAM GENE, encodes a GAM259 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM259 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM259 precursor RNA is designated SEQ ID:3899, and is provided hereinbelow with reference to the sequence listing part.

[3671] GAM259 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM259 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3672] An enzyme complex designated DICER COMPLEX, dices the GAM259 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM259 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM259 RNA is designated SEQ ID:3900, and is provided hereinbelow with reference to the sequence listing part.

[3673] GAM259 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM259 target RNA, herein designated GAM TARGET RNA. GAM259 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3674] GAM259 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM259 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM259 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM259 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM259 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3675] The complementary binding of GAM259 RNA, herein designated GAM RNA, to target binding sites on GAM259 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM259 target RNA, herein designated GAM TARGET RNA, into GAM259 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3676] It is appreciated that GAM259 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM259 target genes. The mRNA of each one of this plurality of GAM259 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM259 RNA, herein designated GAM RNA, and which when bound by GAM259 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM259 target proteins.

[3677] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM259 gene, herein designated GAM GENE, on one or more GAM259 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3678] It is yet further appreciated that specific functions, and accordingly utilities, of GAM259 correlate with, and may be deduced from, the identity of the target genes which GAM259 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3679] Nucleotide sequences of the GAM259 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM259 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM259 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM259 are further described hereinbelow with reference to Table Table1.

[3680] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM259 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3681]

[3682] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 260 (GAM260) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3683] GAM260 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM260 was detected is described hereinabove with reference to Figs. 2-8.

[3684] GAM260 gene, herein designated GAM GENE, and GAM260 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3685] GAM260 gene, herein designated GAM GENE, encodes a GAM260 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM260 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM260 precursor RNA is designated SEQ ID:3909, and is provided hereinbelow with reference to the sequence listing part.

[3686] GAM260 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM260 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3687] An enzyme complex designated DICER COMPLEX, dices the GAM260 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM260 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 81%) nucleotide sequence of GAM260 RNA is designated SEQ ID:3910, and is provided

hereinbelow with reference to the sequence listing part.

[3688] GAM260 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM260 target RNA, herein designated GAM TARGET RNA.
GAM260 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3689] GAM260 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM260 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM260 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM260 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM260 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3690] The complementary binding of GAM260 RNA, herein designated GAM RNA, to target binding sites on GAM260 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM260 target RNA, herein designated GAM TARGET RNA, into GAM260 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3691] It is appreciated that GAM260 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM260 target genes. The mRNA of each one of this plurality of GAM260 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM260 RNA, herein designated GAM RNA, and which when bound by GAM260 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM260 tar-

get proteins.

[3692] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM260 gene, herein designated GAM GENE, on one or more GAM260 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3693] It is yet further appreciated that specific functions, and accordingly utilities, of GAM260 correlate with, and may be deduced from, the identity of the target genes which GAM260 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3694] Nucleotide sequences of the GAM260 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM260 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM260 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM260 are further described hereinbelow with reference to Table Table1.

[3695] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM260 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3696]

[3697] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 261 (GAM261) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3698] GAM261 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM261 was detected is described hereinabove with reference to Figs. 2-8.

[3699] GAM261 gene, herein designated GAM GENE, and GAM261 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3700] GAM261 gene, herein designated GAM GENE, encodes a GAM261 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM261 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM261 precursor RNA is designated SEQ ID:3917, and is provided hereinbelow with reference to the sequence listing part.

[3701] GAM261 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM261 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3702] An enzyme complex designated DICER COMPLEX, dices

the GAM261 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM261 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM261 RNA is designated SEQ ID:3918, and is provided hereinbelow with reference to the sequence listing part.

[3703] GAM261 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM261 target RNA, herein designated GAM TARGET RNA. GAM261 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3704] GAM261 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM261 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM261

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM261 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM261 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3705] The complementary binding of GAM261 RNA, herein designated GAM RNA, to target binding sites on GAM261 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM261 target RNA, herein designated GAM TARGET RNA, into GAM261 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3706] It is appreciated that GAM261 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM261 target genes. The mRNA of each one of this plurality of GAM261 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM261 RNA, herein designated GAM RNA, and which when bound by GAM261 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM261 target proteins.

[3707] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM261 gene, herein designated GAM GENE, on one or more GAM261 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3708] It is yet further appreciated that specific functions, and accordingly utilities, of GAM261 correlate with, and may be deduced from, the identity of the target genes which GAM261 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3709] Nucleotide sequences of the GAM261 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM261 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM261 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM261 are further described hereinbelow with reference to Table Table1.

[3710] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM261 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3711]

[3712] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 262 (GAM262) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3713] GAM262 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM262 was detected is described hereinabove with reference to Figs. 2-8.

[3714] GAM262 gene, herein designated GAM GENE, and GAM262 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3715] GAM262 gene, herein designated GAM GENE, encodes a GAM262 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM262 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM262 precursor RNA is designated SEQ ID:3922, and is provided hereinbelow with reference to the sequence listing part.

[3716] GAM262 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM262 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3717] An enzyme complex designated DICER COMPLEX, dices the GAM262 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM262 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 28%) nucleotide sequence of GAM262 RNA is designated SEQ ID:3923, and is provided hereinbelow with reference to the sequence listing part.

[3718] GAM262 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM262 target RNA, herein designated GAM TARGET RNA. GAM262 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3719] GAM262 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM262 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM262 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM262 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM262 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[3720] The complementary binding of GAM262 RNA, herein designated GAM RNA, to target binding sites on GAM262 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM262 target RNA, herein designated GAM TARGET RNA, into GAM262 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3721] It is appreciated that GAM262 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM262 target genes. The mRNA of each one of this plurality of GAM262 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM262 RNA, herein designated GAM RNA, and which when bound by GAM262 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM262 target proteins.

[3722] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM262 gene, herein designated GAM GENE, on one or

more GAM262 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3723] It is yet further appreciated that specific functions, and accordingly utilities, of GAM262 correlate with, and may be deduced from, the identity of the target genes which GAM262 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3724] Nucleotide sequences of the GAM262 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM262 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM262 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM262 are further de-

scribed hereinbelow with reference to Table Table1.

[3725] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM262 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3726]

[3727] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 263 (GAM263) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3728] GAM263 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM263 was detected is described hereinabove with reference to Figs. 2-8.

[3729] GAM263 gene, herein designated GAM GENE, and GAM263 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3730] GAM263 gene, herein designated GAM GENE, encodes a GAM263 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM263 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM263 precursor RNA is designated SEQ ID:3947, and is provided hereinbelow with reference to the sequence listing part.

[3731] GAM263 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM263 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3732] An enzyme complex designated DICER COMPLEX, dices the GAM263 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM263 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM263 RNA is designated SEQ ID:3948, and is provided hereinbelow with reference to the sequence listing part.

[3733] GAM263 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM263 target RNA, herein designated GAM TARGET RNA. GAM263 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3734] GAM263 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM263 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM263 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM263 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM263 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3735] The complementary binding of GAM263 RNA, herein designated GAM RNA, to target binding sites on GAM263 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM263 target RNA, herein designated GAM TARGET RNA, into GAM263 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3736] It is appreciated that GAM263 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM263 target genes. The mRNA of each one of this plurality of GAM263 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM263 RNA, herein designated GAM RNA, and which when bound by GAM263 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM263 target proteins.

[3737] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM263 gene, herein designated GAM GENE, on one or more GAM263 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3738] It is yet further appreciated that specific functions, and accordingly utilities, of GAM263 correlate with, and may

be deduced from, the identity of the target genes which GAM263 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3739] Nucleotide sequences of the GAM263 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM263 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM263 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM263 are further described hereinbelow with reference to Table Table1.

[3740] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM263 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3741]

[3742] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 264 (GAM264) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3743] GAM264 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM264 was detected is described hereinabove with reference to Figs. 2-8.

[3744] GAM264 gene, herein designated GAM GENE, and GAM264 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3745] GAM264 gene, herein designated GAM GENE, encodes a GAM264 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM264 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM264 precursor RNA is designated SEQ ID:3961, and is provided hereinbelow with reference to the sequence listing part.

[3746] GAM264 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM264 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3747] An enzyme complex designated DICER COMPLEX, dices the GAM264 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM264 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM264 RNA is designated SEQ ID:3962, and is provided hereinbelow with reference to the sequence listing part.

[3748] GAM264 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM264 target RNA, herein designated GAM TARGET RNA. GAM264 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3749] GAM264 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM264 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM264 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM264 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM264 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3750] The complementary binding of GAM264 RNA, herein designated GAM RNA, to target binding sites on GAM264 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM264 target RNA, herein designated GAM TARGET RNA, into GAM264 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3751] It is appreciated that GAM264 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM264 target genes. The mRNA of each one of this plurality of GAM264 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM264 RNA, herein designated GAM RNA, and which when bound by GAM264 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM264 target proteins.

[3752] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM264 gene, herein designated GAM GENE, on one or more GAM264 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3753] It is yet further appreciated that specific functions, and accordingly utilities, of GAM264 correlate with, and may be deduced from, the identity of the target genes which GAM264 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3754] Nucleotide sequences of the GAM264 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM264 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM264 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM264 are further described hereinbelow with reference to Table Table1.

[3755] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM264 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3756]

[3757] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 265 (GAM265) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3758] GAM265 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM265 was detected is described hereinabove with reference to Figs. 2-8.

[3759] GAM265 gene, herein designated GAM GENE, and GAM265 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3760] GAM265 gene, herein designated GAM GENE, encodes a GAM265 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM265 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM265 precursor RNA is desig-

nated SEQ ID:3977, and is provided hereinbelow with reference to the sequence listing part.

[3761] GAM265 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM265 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3762] An enzyme complex designated DICER COMPLEX, dices the GAM265 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM265 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM265 RNA is designated SEQ ID:3978, and is provided hereinbelow with reference to the sequence listing part.

[3763] GAM265 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM265 target RNA, herein designated GAM TARGET RNA. GAM265 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3764] GAM265 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM265 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM265 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM265 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM265

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3765] The complementary binding of GAM265 RNA, herein designated GAM RNA, to target binding sites on GAM265 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM265 target RNA, herein designated GAM TARGET RNA, into GAM265 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3766] It is appreciated that GAM265 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM265 target genes. The mRNA of each one of this plurality of GAM265 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM265 RNA, herein designated GAM RNA, and which when bound by GAM265 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM265 target proteins.

[3767] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM265 gene, herein designated GAM GENE, on one or more GAM265 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3768] It is yet further appreciated that specific functions, and accordingly utilities, of GAM265 correlate with, and may be deduced from, the identity of the target genes which GAM265 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3769] Nucleotide sequences of the GAM265 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM265 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM265 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM265 are further described hereinbelow with reference to Table Table1.

[3770] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM265 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3771]

[3772] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 266 (GAM266) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3773] GAM266 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM266 was detected is described hereinabove with reference to Figs. 2-8.

[3774] GAM266 gene, herein designated GAM GENE, and

GAM266 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3775] GAM266 gene, herein designated GAM GENE, encodes a GAM266 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM266 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM266 precursor RNA is designated SEQ ID:3988, and is provided hereinbelow with reference to the sequence listing part.

[3776] GAM266 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM266 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3777] An enzyme complex designated DICER COMPLEX, dices the GAM266 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM266 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM266 RNA is designated SEQ ID:3989, and is provided hereinbelow with reference to the sequence listing part.

[3778] GAM266 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM266 target RNA, herein designated GAM TARGET RNA. GAM266 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3779] GAM266 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM266 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM266 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM266 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM266 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3780] The complementary binding of GAM266 RNA, herein designated GAM RNA, to target binding sites on GAM266 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM266 target RNA, herein designated GAM TARGET RNA, into GAM266 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3781] It is appreciated that GAM266 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM266 target genes. The mRNA of each one of this plurality of GAM266 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM266 RNA, herein designated GAM RNA, and which when bound by GAM266 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM266 target proteins.

[3782] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM266 gene, herein designated GAM GENE, on one or more GAM266 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3783] It is yet further appreciated that specific functions, and accordingly utilities, of GAM266 correlate with, and may be deduced from, the identity of the target genes which GAM266 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3784] Nucleotide sequences of the GAM266 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM266 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM266 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM266 are further described hereinbelow with reference to Table Table1.

[3785] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM266 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3786]

[3787] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 267 (GAM267) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3788] GAM267 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM267 was detected is described hereinabove with reference to Figs. 2-8.

[3789] GAM267 gene, herein designated GAM GENE, and GAM267 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3790] GAM267 gene, herein designated GAM GENE, encodes a GAM267 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM267 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM267 precursor RNA is designated SEQ ID:4007, and is provided hereinbelow with reference to the sequence listing part.

[3791] GAM267 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM267 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3792] An enzyme complex designated DICER COMPLEX, dices the GAM267 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM267 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM267 RNA is designated SEQ ID:4008, and is provided hereinbelow with reference to the sequence listing part.

[3793] GAM267 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM267 target RNA, herein designated GAM TARGET RNA. GAM267 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3794] GAM267 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM267 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM267 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM267 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM267 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3795] The complementary binding of GAM267 RNA, herein designated GAM RNA, to target binding sites on GAM267 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM267 target RNA, herein designated GAM TARGET RNA, into GAM267 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3796] It is appreciated that GAM267 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM267 target genes. The mRNA of each one of this plurality of GAM267 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM267 RNA, herein designated GAM RNA, and which when bound by GAM267 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM267 target proteins.

[3797] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM267 gene, herein designated GAM GENE, on one or more GAM267 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3798] It is yet further appreciated that specific functions, and accordingly utilities, of GAM267 correlate with, and may be deduced from, the identity of the target genes which GAM267 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3799] Nucleotide sequences of the GAM267 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM267 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM267 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM267 are further described hereinbelow with reference to Table Table1.

[3800] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM267 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3801]

[3802] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 268 (GAM268) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3803] GAM268 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM268 was detected is described hereinabove with reference to Figs. 2-8.

[3804] GAM268 gene, herein designated GAM GENE, and GAM268 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3805] GAM268 gene, herein designated GAM GENE, encodes a GAM268 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM268 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM268 precursor RNA is designated SEQ ID:4018, and is provided hereinbelow with reference to the sequence listing part.

[3806] GAM268 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM268 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3807] An enzyme complex designated DICER COMPLEX, dices the GAM268 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM268 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of GAM268 RNA is designated SEQ ID:4019, and is provided hereinbelow with reference to the sequence listing part.

[3808] GAM268 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM268 target RNA, herein designated GAM TARGET RNA. GAM268 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3809] GAM268 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM268 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM268 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM268 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM268 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3810] The complementary binding of GAM268 RNA, herein designated GAM RNA, to target binding sites on GAM268 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM268 target RNA, herein designated GAM TARGET RNA, into GAM268 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3811] It is appreciated that GAM268 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM268 target genes. The mRNA of each one of this plurality of GAM268 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM268 RNA,

herein designated GAM RNA, and which when bound by GAM268 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM268 target proteins.

[3812] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM268 gene, herein designated GAM GENE, on one or more GAM268 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3813] It is yet further appreciated that specific functions, and accordingly utilities, of GAM268 correlate with, and may be deduced from, the identity of the target genes which

GAM268 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3814] Nucleotide sequences of the GAM268 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM268 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM268 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM268 are further described hereinbelow with reference to Table Table1.

[3815] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM268 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3816]

[3817] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 269 (GAM269) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3818] GAM269 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM269 was detected is described hereinabove with reference to Figs. 2-8.

[3819] GAM269 gene, herein designated GAM GENE, and GAM269 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3820] GAM269 gene, herein designated GAM GENE, encodes a GAM269 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM269 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM269 precursor RNA is designated SEQ ID:4038, and is provided hereinbelow with reference to the sequence listing part.

[3821] GAM269 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM269 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[3822] An enzyme complex designated DICER COMPLEX, dices the GAM269 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM269 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM269 RNA is designated SEQ ID:4039, and is provided hereinbelow with reference to the sequence listing part.

[3823] GAM269 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM269 target RNA, herein designated GAM TARGET RNA.
GAM269 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3824] GAM269 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM269 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM269 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM269 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM269 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3825] The complementary binding of GAM269 RNA, herein designated GAM RNA, to target binding sites on GAM269 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM269 target RNA, herein desig-

nated GAM TARGET RNA, into GAM269 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3826] It is appreciated that GAM269 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM269 target genes. The mRNA of each one of this plurality of GAM269 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM269 RNA, herein designated GAM RNA, and which when bound by GAM269 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM269 target proteins.

[3827] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM269 gene, herein designated GAM GENE, on one or more GAM269 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3828] It is yet further appreciated that specific functions, and accordingly utilities, of GAM269 correlate with, and may be deduced from, the identity of the target genes which GAM269 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3829] Nucleotide sequences of the GAM269 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM269 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM269 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM269 are further described hereinbelow with reference to Table Table1.

[3830] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM269 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[3831]

[3832] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 270 (GAM270) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3833] GAM270 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM270 was detected is described hereinabove with reference to Figs. 2-8.

[3834] GAM270 gene, herein designated GAM GENE, and GAM270 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3835] GAM270 gene, herein designated GAM GENE, encodes a GAM270 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM270 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM270 precursor RNA is designated SEQ ID:4052, and is provided hereinbelow with ref-

erence to the sequence listing part.

[3836] GAM270 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM270 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3837] An enzyme complex designated DICER COMPLEX, dices the GAM270 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM270 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM270 RNA is designated SEQ ID:4053, and is provided hereinbelow with reference to the sequence listing part.

[3838] GAM270 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM270 target RNA, herein designated GAM TARGET RNA.

GAM270 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3839] GAM270 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM270 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM270 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM270 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM270 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3840] The complementary binding of GAM270 RNA, herein designated GAM RNA, to target binding sites on GAM270 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM270 target RNA, herein designated GAM TARGET RNA, into GAM270 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3841] It is appreciated that GAM270 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM270 target genes. The mRNA of each one of this plurality of GAM270 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM270 RNA, herein designated GAM RNA, and which when bound by GAM270 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM270 target proteins.

[3842] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM270 gene, herein designated GAM GENE, on one or more GAM270 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3843] It is yet further appreciated that specific functions, and accordingly utilities, of GAM270 correlate with, and may be deduced from, the identity of the target genes which GAM270 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3844] Nucleotide sequences of the GAM270 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM270 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM270 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM270 are further described hereinbelow with reference to Table Table1.

[3845] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM270 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3846]

[3847] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 271 (GAM271) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3848] GAM271 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM271 was detected is described hereinabove with reference to Figs. 2-8.

[3849] GAM271 gene, herein designated GAM GENE, and GAM271 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[3850] GAM271 gene, herein designated GAM GENE, encodes a GAM271 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM271 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM271 precursor RNA is designated SEQ ID:4068, and is provided hereinbelow with reference to the sequence listing part.

[3851] GAM271 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM271 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3852] An enzyme complex designated DICER COMPLEX, dices the GAM271 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM271 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM271 RNA is designated SEQ ID:4069, and is provided hereinbelow with reference to the sequence listing part.

[3853] GAM271 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM271 target RNA, herein designated GAM TARGET RNA. GAM271 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3854] GAM271 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM271 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM271 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM271 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM271 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3855] The complementary binding of GAM271 RNA, herein designated GAM RNA, to target binding sites on GAM271 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM271 target RNA, herein designated GAM TARGET RNA, into GAM271 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3856] It is appreciated that GAM271 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM271 target genes. The mRNA of each one of this plurality of GAM271 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM271 RNA, herein designated GAM RNA, and which when bound by GAM271 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM271 target proteins.

[3857] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM271 gene, herein designated GAM GENE, on one or more GAM271 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[3858] It is yet further appreciated that specific functions, and accordingly utilities, of GAM271 correlate with, and may be deduced from, the identity of the target genes which GAM271 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3859] Nucleotide sequences of the GAM271 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM271 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM271 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM271 are further described hereinbelow with reference to Table Table1.

[3860] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM271 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3861]

[3862] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

272 (GAM272) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3863] GAM272 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM272 was detected is described hereinabove with reference to Figs. 2-8.

[3864] GAM272 gene, herein designated GAM GENE, and GAM272 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3865] GAM272 gene, herein designated GAM GENE, encodes a GAM272 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM272 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM272 precursor RNA is designated SEQ ID:4104, and is provided hereinbelow with reference to the sequence listing part.

[3866] GAM272 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM272 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3867] An enzyme complex designated DICER COMPLEX, dices the GAM272 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM272 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM272 RNA is designated SEQ ID:4105, and is provided hereinbelow with reference to the sequence listing part.

[3868] GAM272 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM272 target RNA, herein designated GAM TARGET RNA. GAM272 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3869] GAM272 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM272 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM272 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM272 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM272 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3870] The complementary binding of GAM272 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM272 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM272 target RNA, herein designated GAM TARGET RNA, into GAM272 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3871] It is appreciated that GAM272 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM272 target genes. The mRNA of each one of this plurality of GAM272 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM272 RNA, herein designated GAM RNA, and which when bound by GAM272 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM272 target proteins.

[3872] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM272 gene, herein designated GAM GENE, on one or more GAM272 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3873] It is yet further appreciated that specific functions, and accordingly utilities, of GAM272 correlate with, and may be deduced from, the identity of the target genes which GAM272 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3874] Nucleotide sequences of the GAM272 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM272 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM272 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM272 are further described hereinbelow with reference to Table Table1.

[3875] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM272 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3876]

[3877] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 273 (GAM273) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3878] GAM273 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM273 was detected is described hereinabove with reference to Figs. 2-8.

[3879] GAM273 gene, herein designated GAM GENE, and GAM273 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3880] GAM273 gene, herein designated GAM GENE, encodes a GAM273 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM273 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM273 precursor RNA is designated SEQ ID:4125, and is provided hereinbelow with reference to the sequence listing part.

[3881] GAM273 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM273 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3882] An enzyme complex designated DICER COMPLEX, dices the GAM273 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM273 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 54%) nucleotide sequence of GAM273 RNA is designated SEQ ID:4126, and is provided hereinbelow with reference to the sequence listing part.

[3883] GAM273 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM273 target RNA, herein designated GAM TARGET RNA. GAM273 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3884] GAM273 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM273 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM273 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM273 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM273 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3885] The complementary binding of GAM273 RNA, herein designated GAM RNA, to target binding sites on GAM273 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM273 target RNA, herein designated GAM TARGET RNA, into GAM273 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3886] It is appreciated that GAM273 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM273 target genes. The mRNA of each one of this plurality of GAM273 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM273 RNA, herein designated GAM RNA, and which when bound by

GAM273 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM273 target proteins.

[3887] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM273 gene, herein designated GAM GENE, on one or more GAM273 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3888] It is yet further appreciated that specific functions, and accordingly utilities, of GAM273 correlate with, and may be deduced from, the identity of the target genes which GAM273 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[3889] Nucleotide sequences of the GAM273 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM273 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM273 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM273 are further described hereinbelow with reference to Table Table1.

[3890] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM273 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3891]

[3892] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 274 (GAM274) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3893] GAM274 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM274 was detected is described hereinabove with reference to Figs. 2-8.

[3894] GAM274 gene, herein designated GAM GENE, and GAM274 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3895] GAM274 gene, herein designated GAM GENE, encodes a GAM274 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM274 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM274 precursor RNA is designated SEQ ID:4137, and is provided hereinbelow with reference to the sequence listing part.

[3896] GAM274 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM274 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[3897] An enzyme complex designated DICER COMPLEX, dices the GAM274 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM274 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM274 RNA is designated SEQ ID:4138, and is provided hereinbelow with reference to the sequence listing part.

[3898] GAM274 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM274 target RNA, herein designated GAM TARGET RNA. GAM274 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3899] GAM274 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM274 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM274 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM274 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM274 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3900] The complementary binding of GAM274 RNA, herein designated GAM RNA, to target binding sites on GAM274 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM274 target RNA, herein designated GAM TARGET RNA, into GAM274 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3901] It is appreciated that GAM274 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM274 target genes. The mRNA of each one of this plurality of GAM274 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM274 RNA, herein designated GAM RNA, and which when bound by GAM274 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM274 target proteins.

[3902] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM274 gene, herein designated GAM GENE, on one or more GAM274 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3903] It is yet further appreciated that specific functions, and accordingly utilities, of GAM274 correlate with, and may be deduced from, the identity of the target genes which GAM274 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3904] Nucleotide sequences of the GAM274 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM274 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM274 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM274 are further described hereinbelow with reference to Table Table1.

[3905] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM274 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3906]

[3907] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 275 (GAM275) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3908] GAM275 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM275 was detected is described hereinabove with reference to Figs. 2-8.

[3909] GAM275 gene, herein designated GAM GENE, and GAM275 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3910] GAM275 gene, herein designated GAM GENE, encodes a GAM275 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM275 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM275 precursor RNA is designated SEQ ID:4142, and is provided hereinbelow with reference to the sequence listing part.

[3911] GAM275 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM275 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3912] An enzyme complex designated DICER COMPLEX, dices the GAM275 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM275 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM275 RNA is designated SEQ ID:4143, and is provided hereinbelow with reference to the sequence listing part.

[3913] GAM275 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM275

target RNA, herein designated GAM TARGET RNA.

GAM275 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3914] GAM275 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM275 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM275 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM275 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM275 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3915] The complementary binding of GAM275 RNA, herein designated GAM RNA, to target binding sites on GAM275 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM275 target RNA, herein designated GAM TARGET RNA, into GAM275 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3916] It is appreciated that GAM275 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM275 target genes. The mRNA of each one of this plurality of GAM275 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM275 RNA, herein designated GAM RNA, and which when bound by GAM275 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM275 target proteins.

[3917] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM275 gene, herein designated GAM GENE, on one or more GAM275 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3918] It is yet further appreciated that specific functions, and accordingly utilities, of GAM275 correlate with, and may be deduced from, the identity of the target genes which GAM275 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3919] Nucleotide sequences of the GAM275 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM275 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM275 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM275 are further described hereinbelow with reference to Table Table1.

[3920] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM275 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3921]

[3922] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 276 (GAM276) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3923] GAM276 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM276 was detected is described hereinabove with reference to Figs. 2-8.

[3924] GAM276 gene, herein designated GAM GENE, and GAM276 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3925] GAM276 gene, herein designated GAM GENE, encodes a GAM276 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM276 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM276 precursor RNA is designated SEQ ID:4157, and is provided hereinbelow with reference to the sequence listing part.

[3926] GAM276 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM276 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3927] An enzyme complex designated DICER COMPLEX, dices the GAM276 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM276 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 44%) nucleotide sequence of GAM276 RNA is designated SEQ ID:4158, and is provided hereinbelow with reference to the sequence listing part.

[3928] GAM276 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM276 target RNA, herein designated GAM TARGET RNA. GAM276 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3929] GAM276 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM276 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM276 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM276 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM276 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3930] The complementary binding of GAM276 RNA, herein designated GAM RNA, to target binding sites on GAM276 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM276 target RNA, herein designated GAM TARGET RNA, into GAM276 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3931] It is appreciated that GAM276 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM276 target genes. The mRNA of each one of this plu-

rality of GAM276 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM276 RNA, herein designated GAM RNA, and which when bound by GAM276 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM276 target proteins.

[3932] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM276 gene, herein designated GAM GENE, on one or more GAM276 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3933] It is yet further appreciated that specific functions, and accordingly utilities, of GAM276 correlate with, and may be deduced from, the identity of the target genes which GAM276 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3934] Nucleotide sequences of the GAM276 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM276 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM276 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM276 are further described hereinbelow with reference to Table Table1.

[3935] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM276 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3936]

[3937] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 277 (GAM277) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[3938] GAM277 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM277 was detected is described hereinabove with reference to Figs. 2-8.

[3939] GAM277 gene, herein designated GAM GENE, and GAM277 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3940] GAM277 gene, herein designated GAM GENE, encodes a GAM277 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM277 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM277 precursor RNA is designated SEQ ID:4177, and is provided hereinbelow with reference to the sequence listing part.

[3941] GAM277 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM277 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3942] An enzyme complex designated DICER COMPLEX, dices the GAM277 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM277 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM277 RNA is designated SEQ ID:4178, and is provided hereinbelow with reference to the sequence listing part.

[3943] GAM277 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM277 target RNA, herein designated GAM TARGET RNA. GAM277 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[3944] GAM277 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM277 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM277 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM277 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM277 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3945] The complementary binding of GAM277 RNA, herein designated GAM RNA, to target binding sites on GAM277 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM277 target RNA, herein designated GAM TARGET RNA, into GAM277 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3946] It is appreciated that GAM277 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM277 target genes. The mRNA of each one of this plurality of GAM277 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM277 RNA, herein designated GAM RNA, and which when bound by GAM277 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM277 target proteins.

[3947] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM277 gene, herein designated GAM GENE, on one or more GAM277 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3948] It is yet further appreciated that specific functions, and accordingly utilities, of GAM277 correlate with, and may be deduced from, the identity of the target genes which GAM277 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3949] Nucleotide sequences of the GAM277 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM277 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM277 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM277 are further described hereinbelow with reference to Table Table1.

[3950] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM277 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3951]

[3952] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 278 (GAM278) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3953] GAM278 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM278 was detected is described hereinabove with reference to Figs. 2-8.

[3954] GAM278 gene, herein designated GAM GENE, and GAM278 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3955] GAM278 gene, herein designated GAM GENE, encodes a GAM278 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM278 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM278 precursor RNA is designated SEQ ID:4202, and is provided hereinbelow with reference to the sequence listing part.

[3956] GAM278 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM278 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3957] An enzyme complex designated DICER COMPLEX, dices the GAM278 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM278 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of

GAM278 RNA is designated SEQ ID:4203, and is provided hereinbelow with reference to the sequence listing part.

[3958] GAM278 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM278 target RNA, herein designated GAM TARGET RNA. GAM278 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3959] GAM278 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM278 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM278 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM278 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM278 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3960] The complementary binding of GAM278 RNA, herein designated GAM RNA, to target binding sites on GAM278 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM278 target RNA, herein designated GAM TARGET RNA, into GAM278 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3961] It is appreciated that GAM278 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM278 target genes. The mRNA of each one of this plurality of GAM278 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM278 RNA, herein designated GAM RNA, and which when bound by GAM278 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM278 target proteins.

[3962] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM278 gene, herein designated GAM GENE, on one or more GAM278 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3963] It is yet further appreciated that specific functions, and accordingly utilities, of GAM278 correlate with, and may be deduced from, the identity of the target genes which GAM278 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3964] Nucleotide sequences of the GAM278 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM278 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM278 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM278 are further described hereinbelow with reference to Table Table1.

[3965] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM278 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3966]

[3967] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 279 (GAM279) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3968] GAM279 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM279 was detected is described hereinabove

with reference to Figs. 2-8.

[3969] GAM279 gene, herein designated GAM GENE, and GAM279 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3970] GAM279 gene, herein designated GAM GENE, encodes a GAM279 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM279 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM279 precursor RNA is designated SEQ ID:4234, and is provided hereinbelow with reference to the sequence listing part.

[3971] GAM279 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM279 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3972] An enzyme complex designated DICER COMPLEX, dices the GAM279 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM279 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 84%) nucleotide sequence of GAM279 RNA is designated SEQ ID:4235, and is provided hereinbelow with reference to the sequence listing part.

[3973] GAM279 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM279 target RNA, herein designated GAM TARGET RNA. GAM279 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3974] GAM279 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM279 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM279 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM279 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM279 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3975] The complementary binding of GAM279 RNA, herein designated GAM RNA, to target binding sites on GAM279 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM279 target RNA, herein designated GAM TARGET RNA, into GAM279 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[3976] It is appreciated that GAM279 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM279 target genes. The mRNA of each one of this plurality of GAM279 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM279 RNA, herein designated GAM RNA, and which when bound by GAM279 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM279 target proteins.

[3977] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM279 gene, herein designated GAM GENE, on one or more GAM279 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3978] It is yet further appreciated that specific functions, and accordingly utilities, of GAM279 correlate with, and may be deduced from, the identity of the target genes which GAM279 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3979] Nucleotide sequences of the GAM279 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM279 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM279 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM279 are further described hereinbelow with reference to Table Table1.

[3980] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM279 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3981]

[3982] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 280 (GAM280) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3983] GAM280 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM280 was detected is described hereinabove with reference to Figs. 2-8.

[3984] GAM280 gene, herein designated GAM GENE, and GAM280 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[3985] GAM280 gene, herein designated GAM GENE, encodes a GAM280 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM280 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM280 precursor RNA is designated SEQ ID:4245, and is provided hereinbelow with reference to the sequence listing part.

[3986] GAM280 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM280 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[3987] An enzyme complex designated DICER COMPLEX, dices the GAM280 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM280 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM280 RNA is designated SEQ ID:4246, and is provided hereinbelow with reference to the sequence listing part.

[3988] GAM280 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM280 target RNA, herein designated GAM TARGET RNA.

GAM280 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[3989] GAM280 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM280 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM280 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM280 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM280 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[3990] The complementary binding of GAM280 RNA, herein designated GAM RNA, to target binding sites on GAM280 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM280 target RNA, herein designated GAM TARGET RNA, into GAM280 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[3991] It is appreciated that GAM280 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM280 target genes. The mRNA of each one of this plurality of GAM280 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM280 RNA, herein designated GAM RNA, and which when bound by GAM280 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM280 target proteins.

[3992] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM280 gene, herein designated GAM GENE, on one or more GAM280 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[3993] It is yet further appreciated that specific functions, and accordingly utilities, of GAM280 correlate with, and may be deduced from, the identity of the target genes which GAM280 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[3994] Nucleotide sequences of the GAM280 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM280 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM280 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM280 are further described hereinbelow with reference to Table Table1.

[3995] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM280 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[3996]

[3997] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 281 (GAM281) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[3998] GAM281 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM281 was detected is described hereinabove with reference to Figs. 2-8.

[3999] GAM281 gene, herein designated GAM GENE, and GAM281 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4000] GAM281 gene, herein designated GAM GENE, encodes a

GAM281 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM281 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM281 precursor RNA is designated SEQ ID:4261, and is provided hereinbelow with reference to the sequence listing part.

[4001] GAM281 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM281 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4002] An enzyme complex designated DICER COMPLEX, dices the GAM281 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM281 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 22%) nucleotide sequence of GAM281 RNA is designated SEQ ID:4262, and is provided hereinbelow with reference to the sequence listing part.

[4003] GAM281 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM281 target RNA, herein designated GAM TARGET RNA. GAM281 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4004] GAM281 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM281 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM281 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM281 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM281 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4005] The complementary binding of GAM281 RNA, herein designated GAM RNA, to target binding sites on GAM281 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM281 target RNA, herein designated GAM TARGET RNA, into GAM281 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4006] It is appreciated that GAM281 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM281 target genes. The mRNA of each one of this plurality of GAM281 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM281 RNA, herein designated GAM RNA, and which when bound by GAM281 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM281 target proteins.

[4007] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM281 gene, herein designated GAM GENE, on one or more GAM281 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4008] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM281 correlate with, and may be deduced from, the identity of the target genes which GAM281 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4009] Nucleotide sequences of the GAM281 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM281 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM281 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM281 are further described hereinbelow with reference to Table Table1.

[4010] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM281 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4011]

[4012] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 282 (GAM282) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[4013] GAM282 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM282 was detected is described hereinabove with reference to Figs. 2-8.

[4014] GAM282 gene, herein designated GAM GENE, and GAM282 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4015] GAM282 gene, herein designated GAM GENE, encodes a GAM282 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM282 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM282 precursor RNA is designated SEQ ID:4276, and is provided hereinbelow with reference to the sequence listing part.

[4016] GAM282 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM282 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4017] An enzyme complex designated DICER COMPLEX, dices the GAM282 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM282 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM282 RNA is designated SEQ ID:4277, and is provided hereinbelow with reference to the sequence listing part.

[4018] GAM282 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM282 target RNA, herein designated GAM TARGET RNA. GAM282 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4019] GAM282 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM282 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM282 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM282 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM282 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4020] The complementary binding of GAM282 RNA, herein designated GAM RNA, to target binding sites on GAM282 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM282 target RNA, herein designated GAM TARGET RNA, into GAM282 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4021] It is appreciated that GAM282 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM282 target genes. The mRNA of each one of this plurality of GAM282 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM282 RNA, herein designated GAM RNA, and which when bound by GAM282 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM282 target proteins.

[4022] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM282 gene, herein designated GAM GENE, on one or more GAM282 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4023] It is yet further appreciated that specific functions, and accordingly utilities, of GAM282 correlate with, and may be deduced from, the identity of the target genes which GAM282 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4024] Nucleotide sequences of the GAM282 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM282 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM282 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM282 are further described hereinbelow with reference to Table Table1.

[4025] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM282 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4026]

[4027] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 283 (GAM283) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4028] GAM283 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM283 was detected is described hereinabove with reference to Figs. 2-8.

[4029] GAM283 gene, herein designated GAM GENE, and GAM283 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4030] GAM283 gene, herein designated GAM GENE, encodes a GAM283 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM283 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM283 precursor RNA is designated SEQ ID:4290, and is provided hereinbelow with reference to the sequence listing part.

[4031] GAM283 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM283 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4032] An enzyme complex designated DICER COMPLEX, dices the GAM283 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM283 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM283 RNA is designated SEQ ID:4291, and is provided

hereinbelow with reference to the sequence listing part.

[4033] GAM283 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM283 target RNA, herein designated GAM TARGET RNA.

GAM283 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4034] GAM283 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM283 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM283 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM283 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM283 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4035] The complementary binding of GAM283 RNA, herein designated GAM RNA, to target binding sites on GAM283 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM283 target RNA, herein designated GAM TARGET RNA, into GAM283 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4036] It is appreciated that GAM283 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM283 target genes. The mRNA of each one of this plurality of GAM283 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM283 RNA, herein designated GAM RNA, and which when bound by GAM283 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM283 tar-

get proteins.

[4037] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM283 gene, herein designated GAM GENE, on one or more GAM283 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4038] It is yet further appreciated that specific functions, and accordingly utilities, of GAM283 correlate with, and may be deduced from, the identity of the target genes which GAM283 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4039] Nucleotide sequences of the GAM283 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM283 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM283 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM283 are further described hereinbelow with reference to Table Table1.

[4040] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM283 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4041]

[4042] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 284 (GAM284) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4043] GAM284 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM284 was detected is described hereinabove with reference to Figs. 2-8.

[4044] GAM284 gene, herein designated GAM GENE, and GAM284 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4045] GAM284 gene, herein designated GAM GENE, encodes a GAM284 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM284 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM284 precursor RNA is designated SEQ ID:4301, and is provided hereinbelow with reference to the sequence listing part.

[4046] GAM284 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM284 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4047] An enzyme complex designated DICER COMPLEX, dices

the GAM284 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM284 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM284 RNA is designated SEQ ID:4302, and is provided hereinbelow with reference to the sequence listing part.

[4048] GAM284 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM284 target RNA, herein designated GAM TARGET RNA. GAM284 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4049] GAM284 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM284 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM284

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM284 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM284 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4050] The complementary binding of GAM284 RNA, herein designated GAM RNA, to target binding sites on GAM284 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM284 target RNA, herein designated GAM TARGET RNA, into GAM284 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4051] It is appreciated that GAM284 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM284 target genes. The mRNA of each one of this plurality of GAM284 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM284 RNA, herein designated GAM RNA, and which when bound by GAM284 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM284 target proteins.

[4052] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM284 gene, herein designated GAM GENE, on one or more GAM284 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4053] It is yet further appreciated that specific functions, and accordingly utilities, of GAM284 correlate with, and may be deduced from, the identity of the target genes which GAM284 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4054] Nucleotide sequences of the GAM284 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM284 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM284 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM284 are further described hereinbelow with reference to Table Table1.

[4055] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM284 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4056]

[4057] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 285 (GAM285) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4058] GAM285 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM285 was detected is described hereinabove with reference to Figs. 2-8.

[4059] GAM285 gene, herein designated GAM GENE, and GAM285 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4060] GAM285 gene, herein designated GAM GENE, encodes a GAM285 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM285 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM285 precursor RNA is designated SEQ ID:4318, and is provided hereinbelow with reference to the sequence listing part.

[4061] GAM285 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM285 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4062] An enzyme complex designated DICER COMPLEX, dices the GAM285 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM285 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 43%) nucleotide sequence of GAM285 RNA is designated SEQ ID:4319, and is provided hereinbelow with reference to the sequence listing part.

[4063] GAM285 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM285 target RNA, herein designated GAM TARGET RNA. GAM285 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4064] GAM285 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM285 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM285 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM285 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM285 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[4065] The complementary binding of GAM285 RNA, herein designated GAM RNA, to target binding sites on GAM285 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM285 target RNA, herein designated GAM TARGET RNA, into GAM285 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4066] It is appreciated that GAM285 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM285 target genes. The mRNA of each one of this plurality of GAM285 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM285 RNA, herein designated GAM RNA, and which when bound by GAM285 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM285 target proteins.

[4067] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM285 gene, herein designated GAM GENE, on one or

more GAM285 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4068] It is yet further appreciated that specific functions, and accordingly utilities, of GAM285 correlate with, and may be deduced from, the identity of the target genes which GAM285 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4069] Nucleotide sequences of the GAM285 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM285 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM285 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM285 are further de-

scribed hereinbelow with reference to Table Table1.

[4070] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM285 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4071]

[4072] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 286 (GAM286) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4073] GAM286 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM286 was detected is described hereinabove with reference to Figs. 2-8.

[4074] GAM286 gene, herein designated GAM GENE, and GAM286 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4075] GAM286 gene, herein designated GAM GENE, encodes a GAM286 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM286 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM286 precursor RNA is designated SEQ ID:4328, and is provided hereinbelow with reference to the sequence listing part.

[4076] GAM286 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM286 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4077] An enzyme complex designated DICER COMPLEX, dices the GAM286 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM286 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM286 RNA is designated SEQ ID:4329, and is provided hereinbelow with reference to the sequence listing part.

[4078] GAM286 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM286 target RNA, herein designated GAM TARGET RNA. GAM286 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4079] GAM286 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM286 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM286 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM286 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM286 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4080] The complementary binding of GAM286 RNA, herein designated GAM RNA, to target binding sites on GAM286 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM286 target RNA, herein designated GAM TARGET RNA, into GAM286 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4081] It is appreciated that GAM286 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM286 target genes. The mRNA of each one of this plurality of GAM286 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM286 RNA, herein designated GAM RNA, and which when bound by GAM286 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM286 target proteins.

[4082] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM286 gene, herein designated GAM GENE, on one or more GAM286 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4083] It is yet further appreciated that specific functions, and accordingly utilities, of GAM286 correlate with, and may

be deduced from, the identity of the target genes which GAM286 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4084] Nucleotide sequences of the GAM286 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM286 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM286 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM286 are further described hereinbelow with reference to Table Table1.

[4085] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM286 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4086]

[4087] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 287 (GAM287) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4088] GAM287 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM287 was detected is described hereinabove with reference to Figs. 2-8.

[4089] GAM287 gene, herein designated GAM GENE, and GAM287 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4090] GAM287 gene, herein designated GAM GENE, encodes a GAM287 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM287 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM287 precursor RNA is designated SEQ ID:4344, and is provided hereinbelow with reference to the sequence listing part.

[4091] GAM287 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM287 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4092] An enzyme complex designated DICER COMPLEX, dices the GAM287 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM287 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM287 RNA is designated SEQ ID:4345, and is provided hereinbelow with reference to the sequence listing part.

[4093] GAM287 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM287 target RNA, herein designated GAM TARGET RNA. GAM287 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4094] GAM287 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM287 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM287 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM287 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM287 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4095] The complementary binding of GAM287 RNA, herein designated GAM RNA, to target binding sites on GAM287 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM287 target RNA, herein designated GAM TARGET RNA, into GAM287 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[4096] It is appreciated that GAM287 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM287 target genes. The mRNA of each one of this plurality of GAM287 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM287 RNA, herein designated GAM RNA, and which when bound by GAM287 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM287 target proteins.

[4097] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM287 gene, herein designated GAM GENE, on one or more GAM287 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4098] It is yet further appreciated that specific functions, and accordingly utilities, of GAM287 correlate with, and may be deduced from, the identity of the target genes which GAM287 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4099] Nucleotide sequences of the GAM287 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM287 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM287 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM287 are further described hereinbelow with reference to Table Table1.

[4100] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM287 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4101]

[4102] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 288 (GAM288) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4103] GAM288 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM288 was detected is described hereinabove with reference to Figs. 2-8.

[4104] GAM288 gene, herein designated GAM GENE, and GAM288 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4105] GAM288 gene, herein designated GAM GENE, encodes a GAM288 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM288 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM288 precursor RNA is designated SEQ ID:4353, and is provided hereinbelow with reference to the sequence listing part.

[4106] GAM288 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM288 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4107] An enzyme complex designated DICER COMPLEX, dices the GAM288 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM288 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 73%) nucleotide sequence of GAM288 RNA is designated SEQ ID:4354, and is provided hereinbelow with reference to the sequence listing part.

[4108] GAM288 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM288 target RNA, herein designated GAM TARGET RNA.

GAM288 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4109] GAM288 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM288 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM288 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM288 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM288 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4110] The complementary binding of GAM288 RNA, herein designated GAM RNA, to target binding sites on GAM288 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM288 target RNA, herein designated GAM TARGET RNA, into GAM288 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4111] It is appreciated that GAM288 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM288 target genes. The mRNA of each one of this plurality of GAM288 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM288 RNA, herein designated GAM RNA, and which when bound by GAM288 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM288 target proteins.

[4112] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM288 gene, herein designated GAM GENE, on one or more GAM288 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4113] It is yet further appreciated that specific functions, and accordingly utilities, of GAM288 correlate with, and may be deduced from, the identity of the target genes which GAM288 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4114] Nucleotide sequences of the GAM288 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM288 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM288 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM288 are further described hereinbelow with reference to Table Table1.

[4115] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM288 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4116]

[4117] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 289 (GAM289) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4118] GAM289 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM289 was detected is described hereinabove with reference to Figs. 2-8.

[4119] GAM289 gene, herein designated GAM GENE, and GAM289 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4120] GAM289 gene, herein designated GAM GENE, encodes a

GAM289 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM289 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM289 precursor RNA is designated SEQ ID:4369, and is provided hereinbelow with reference to the sequence listing part.

[4121] GAM289 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM289 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4122] An enzyme complex designated DICER COMPLEX, dices the GAM289 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM289 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM289 RNA is designated SEQ ID:4370, and is provided hereinbelow with reference to the sequence listing part.

[4123] GAM289 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM289 target RNA, herein designated GAM TARGET RNA. GAM289 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4124] GAM289 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM289 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM289 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM289 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM289 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4125] The complementary binding of GAM289 RNA, herein designated GAM RNA, to target binding sites on GAM289 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM289 target RNA, herein designated GAM TARGET RNA, into GAM289 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4126] It is appreciated that GAM289 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM289 target genes. The mRNA of each one of this plurality of GAM289 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM289 RNA, herein designated GAM RNA, and which when bound by GAM289 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM289 target proteins.

[4127] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM289 gene, herein designated GAM GENE, on one or more GAM289 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4128] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM289 correlate with, and may be deduced from, the identity of the target genes which GAM289 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4129] Nucleotide sequences of the GAM289 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM289 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM289 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM289 are further described hereinbelow with reference to Table Table1.

[4130] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM289 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4131]

[4132] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 290 (GAM290) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[4133] GAM290 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM290 was detected is described hereinabove with reference to Figs. 2-8.

[4134] GAM290 gene, herein designated GAM GENE, and GAM290 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4135] GAM290 gene, herein designated GAM GENE, encodes a GAM290 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM290 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM290 precursor RNA is designated SEQ ID:4378, and is provided hereinbelow with reference to the sequence listing part.

[4136] GAM290 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM290 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4137] An enzyme complex designated DICER COMPLEX, dices the GAM290 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM290 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 45%) nucleotide sequence of GAM290 RNA is designated SEQ ID:4379, and is provided hereinbelow with reference to the sequence listing part.

[4138] GAM290 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM290 target RNA, herein designated GAM TARGET RNA.
GAM290 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4139] GAM290 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM290 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM290 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM290 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM290 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4140] The complementary binding of GAM290 RNA, herein designated GAM RNA, to target binding sites on GAM290 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM290 target RNA, herein designated GAM TARGET RNA, into GAM290 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4141] It is appreciated that GAM290 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM290 target genes. The mRNA of each one of this plurality of GAM290 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM290 RNA, herein designated GAM RNA, and which when bound by GAM290 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM290 target proteins.

[4142] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM290 gene, herein designated GAM GENE, on one or more GAM290 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4143] It is yet further appreciated that specific functions, and accordingly utilities, of GAM290 correlate with, and may be deduced from, the identity of the target genes which GAM290 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4144] Nucleotide sequences of the GAM290 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM290 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM290 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM290 are further described hereinbelow with reference to Table Table1.

[4145] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM290 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4146]

[4147] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 291 (GAM291) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4148] GAM291 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM291 was detected is described hereinabove with reference to Figs. 2-8.

[4149] GAM291 gene, herein designated GAM GENE, and GAM291 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4150] GAM291 gene, herein designated GAM GENE, encodes a GAM291 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM291 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM291 precursor RNA is designated SEQ ID:4400, and is provided hereinbelow with reference to the sequence listing part.

[4151] GAM291 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM291 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4152] An enzyme complex designated DICER COMPLEX, dices the GAM291 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM291 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 45%) nucleotide sequence of GAM291 RNA is designated SEQ ID:4401, and is provided

hereinbelow with reference to the sequence listing part.

[4153] GAM291 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM291 target RNA, herein designated GAM TARGET RNA. GAM291 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4154] GAM291 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM291 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM291 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM291 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM291 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4155] The complementary binding of GAM291 RNA, herein designated GAM RNA, to target binding sites on GAM291 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM291 target RNA, herein designated GAM TARGET RNA, into GAM291 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4156] It is appreciated that GAM291 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM291 target genes. The mRNA of each one of this plurality of GAM291 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM291 RNA, herein designated GAM RNA, and which when bound by GAM291 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM291 tar-

get proteins.

[4157] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM291 gene, herein designated GAM GENE, on one or more GAM291 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4158] It is yet further appreciated that specific functions, and accordingly utilities, of GAM291 correlate with, and may be deduced from, the identity of the target genes which GAM291 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4159] Nucleotide sequences of the GAM291 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM291 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM291 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM291 are further described hereinbelow with reference to Table Table1.

[4160] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM291 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4161]

[4162] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 292 (GAM292) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4163] GAM292 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM292 was detected is described hereinabove with reference to Figs. 2-8.

[4164] GAM292 gene, herein designated GAM GENE, and GAM292 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4165] GAM292 gene, herein designated GAM GENE, encodes a GAM292 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM292 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM292 precursor RNA is designated SEQ ID:4429, and is provided hereinbelow with reference to the sequence listing part.

[4166] GAM292 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM292 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4167] An enzyme complex designated DICER COMPLEX, dices

the GAM292 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM292 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM292 RNA is designated SEQ ID:4430, and is provided hereinbelow with reference to the sequence listing part.

[4168] GAM292 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM292 target RNA, herein designated GAM TARGET RNA. GAM292 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4169] GAM292 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM292 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM292

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM292 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM292 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4170] The complementary binding of GAM292 RNA, herein designated GAM RNA, to target binding sites on GAM292 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM292 target RNA, herein designated GAM TARGET RNA, into GAM292 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4171] It is appreciated that GAM292 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM292 target genes. The mRNA of each one of this plurality of GAM292 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM292 RNA, herein designated GAM RNA, and which when bound by GAM292 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM292 target proteins.

[4172] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM292 gene, herein designated GAM GENE, on one or more GAM292 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4173] It is yet further appreciated that specific functions, and accordingly utilities, of GAM292 correlate with, and may be deduced from, the identity of the target genes which GAM292 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4174] Nucleotide sequences of the GAM292 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM292 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM292 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM292 are further described hereinbelow with reference to Table Table1.

[4175] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM292 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4176]

[4177] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 293 (GAM293) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4178] GAM293 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM293 was detected is described hereinabove with reference to Figs. 2-8.

[4179] GAM293 gene, herein designated GAM GENE, and GAM293 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4180] GAM293 gene, herein designated GAM GENE, encodes a GAM293 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM293 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM293 precursor RNA is designated SEQ ID:4449, and is provided hereinbelow with reference to the sequence listing part.

[4181] GAM293 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM293 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4182] An enzyme complex designated DICER COMPLEX, dices the GAM293 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM293 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM293 RNA is designated SEQ ID:4450, and is provided hereinbelow with reference to the sequence listing part.

[4183] GAM293 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM293 target RNA, herein designated GAM TARGET RNA. GAM293 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4184] GAM293 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM293 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM293 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM293 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM293 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[4185] The complementary binding of GAM293 RNA, herein designated GAM RNA, to target binding sites on GAM293 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM293 target RNA, herein designated GAM TARGET RNA, into GAM293 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4186] It is appreciated that GAM293 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM293 target genes. The mRNA of each one of this plurality of GAM293 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM293 RNA, herein designated GAM RNA, and which when bound by GAM293 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM293 target proteins.

[4187] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM293 gene, herein designated GAM GENE, on one or

more GAM293 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4188] It is yet further appreciated that specific functions, and accordingly utilities, of GAM293 correlate with, and may be deduced from, the identity of the target genes which GAM293 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4189] Nucleotide sequences of the GAM293 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM293 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM293 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM293 are further de-

scribed hereinbelow with reference to Table Table1.

[4190] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM293 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4191]

[4192] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 294 (GAM294) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4193] GAM294 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM294 was detected is described hereinabove with reference to Figs. 2-8.

[4194] GAM294 gene, herein designated GAM GENE, and GAM294 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4195] GAM294 gene, herein designated GAM GENE, encodes a GAM294 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM294 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM294 precursor RNA is designated SEQ ID:4458, and is provided hereinbelow with reference to the sequence listing part.

[4196] GAM294 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM294 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4197] An enzyme complex designated DICER COMPLEX, dices the GAM294 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM294 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM294 RNA is designated SEQ ID:4459, and is provided hereinbelow with reference to the sequence listing part.

[4198] GAM294 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM294 target RNA, herein designated GAM TARGET RNA. GAM294 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4199] GAM294 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM294 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM294 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM294 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM294 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4200] The complementary binding of GAM294 RNA, herein designated GAM RNA, to target binding sites on GAM294 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM294 target RNA, herein designated GAM TARGET RNA, into GAM294 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4201] It is appreciated that GAM294 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM294 target genes. The mRNA of each one of this plurality of GAM294 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM294 RNA, herein designated GAM RNA, and which when bound by GAM294 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM294 target proteins.

[4202] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM294 gene, herein designated GAM GENE, on one or more GAM294 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4203] It is yet further appreciated that specific functions, and accordingly utilities, of GAM294 correlate with, and may

be deduced from, the identity of the target genes which GAM294 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4204] Nucleotide sequences of the GAM294 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM294 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM294 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM294 are further described hereinbelow with reference to Table Table1.

[4205] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM294 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4206]

[4207] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 295 (GAM295) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4208] GAM295 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM295 was detected is described hereinabove with reference to Figs. 2-8.

[4209] GAM295 gene, herein designated GAM GENE, and GAM295 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4210] GAM295 gene, herein designated GAM GENE, encodes a GAM295 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM295 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM295 precursor RNA is designated SEQ ID:4462, and is provided hereinbelow with reference to the sequence listing part.

[4211] GAM295 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM295 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4212] An enzyme complex designated DICER COMPLEX, dices the GAM295 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM295 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM295 RNA is designated SEQ ID:4463, and is provided hereinbelow with reference to the sequence listing part.

[4213] GAM295 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM295 target RNA, herein designated GAM TARGET RNA. GAM295 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4214] GAM295 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM295 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM295 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM295 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM295 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4215] The complementary binding of GAM295 RNA, herein designated GAM RNA, to target binding sites on GAM295 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM295 target RNA, herein designated GAM TARGET RNA, into GAM295 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4216] It is appreciated that GAM295 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM295 target genes. The mRNA of each one of this plurality of GAM295 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM295 RNA, herein designated GAM RNA, and which when bound by GAM295 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM295 target proteins.

[4217] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM295 gene, herein designated GAM GENE, on one or more GAM295 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4218] It is yet further appreciated that specific functions, and accordingly utilities, of GAM295 correlate with, and may be deduced from, the identity of the target genes which GAM295 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4219] Nucleotide sequences of the GAM295 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM295 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM295 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM295 are further described hereinbelow with reference to Table Table1.

[4220] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM295 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4221]

[4222] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 296 (GAM296) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4223] GAM296 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM296 was detected is described hereinabove with reference to Figs. 2-8.

[4224] GAM296 gene, herein designated GAM GENE, and GAM296 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4225] GAM296 gene, herein designated GAM GENE, encodes a GAM296 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM296 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM296 precursor RNA is desig-

nated SEQ ID:4473, and is provided hereinbelow with reference to the sequence listing part.

[4226] GAM296 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM296 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4227] An enzyme complex designated DICER COMPLEX, dices the GAM296 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM296 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 36%) nucleotide sequence of GAM296 RNA is designated SEQ ID:4474, and is provided hereinbelow with reference to the sequence listing part.

[4228] GAM296 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM296 target RNA, herein designated GAM TARGET RNA. GAM296 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4229] GAM296 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM296 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM296 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM296 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM296

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4230] The complementary binding of GAM296 RNA, herein designated GAM RNA, to target binding sites on GAM296 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM296 target RNA, herein designated GAM TARGET RNA, into GAM296 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4231] It is appreciated that GAM296 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM296 target genes. The mRNA of each one of this plurality of GAM296 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM296 RNA, herein designated GAM RNA, and which when bound by GAM296 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM296 target proteins.

[4232] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM296 gene, herein designated GAM GENE, on one or more GAM296 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4233] It is yet further appreciated that specific functions, and accordingly utilities, of GAM296 correlate with, and may be deduced from, the identity of the target genes which GAM296 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4234] Nucleotide sequences of the GAM296 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM296 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM296 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM296 are further described hereinbelow with reference to Table Table1.

[4235] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM296 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4236]

[4237] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 297 (GAM297) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4238] GAM297 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM297 was detected is described hereinabove with reference to Figs. 2-8.

[4239] GAM297 gene, herein designated GAM GENE, and

GAM297 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4240] GAM297 gene, herein designated GAM GENE, encodes a GAM297 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM297 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM297 precursor RNA is designated SEQ ID:4488, and is provided hereinbelow with reference to the sequence listing part.

[4241] GAM297 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM297 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4242] An enzyme complex designated DICER COMPLEX, dices the GAM297 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM297 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM297 RNA is designated SEQ ID:4489, and is provided hereinbelow with reference to the sequence listing part.

[4243] GAM297 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM297 target RNA, herein designated GAM TARGET RNA. GAM297 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4244] GAM297 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM297 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM297 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM297 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM297 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4245] The complementary binding of GAM297 RNA, herein designated GAM RNA, to target binding sites on GAM297 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM297 target RNA, herein designated GAM TARGET RNA, into GAM297 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4246] It is appreciated that GAM297 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM297 target genes. The mRNA of each one of this plurality of GAM297 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM297 RNA, herein designated GAM RNA, and which when bound by GAM297 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM297 target proteins.

[4247] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM297 gene, herein designated GAM GENE, on one or more GAM297 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4248] It is yet further appreciated that specific functions, and accordingly utilities, of GAM297 correlate with, and may be deduced from, the identity of the target genes which GAM297 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4249] Nucleotide sequences of the GAM297 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM297 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM297 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM297 are further described hereinbelow with reference to Table Table1.

[4250] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM297 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4251]

[4252] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 298 (GAM298) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4253] GAM298 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM298 was detected is described hereinabove with reference to Figs. 2-8.

[4254] GAM298 gene, herein designated GAM GENE, and GAM298 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4255] GAM298 gene, herein designated GAM GENE, encodes a GAM298 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM298 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM298 precursor RNA is designated SEQ ID:4508, and is provided hereinbelow with reference to the sequence listing part.

[4256] GAM298 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM298 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4257] An enzyme complex designated DICER COMPLEX, dices the GAM298 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM298 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM298 RNA is designated SEQ ID:4509, and is provided hereinbelow with reference to the sequence listing part.

[4258] GAM298 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM298 target RNA, herein designated GAM TARGET RNA. GAM298 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4259] GAM298 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM298 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM298 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM298 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM298 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4260] The complementary binding of GAM298 RNA, herein designated GAM RNA, to target binding sites on GAM298 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM298 target RNA, herein designated GAM TARGET RNA, into GAM298 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4261] It is appreciated that GAM298 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM298 target genes. The mRNA of each one of this plurality of GAM298 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM298 RNA, herein designated GAM RNA, and which when bound by GAM298 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM298 target proteins.

[4262] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM298 gene, herein designated GAM GENE, on one or more GAM298 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4263] It is yet further appreciated that specific functions, and accordingly utilities, of GAM298 correlate with, and may be deduced from, the identity of the target genes which GAM298 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4264] Nucleotide sequences of the GAM298 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM298 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM298 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM298 are further described hereinbelow with reference to Table Table1.

[4265] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM298 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4266]

[4267] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 299 (GAM299) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4268] GAM299 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM299 was detected is described hereinabove with reference to Figs. 2-8.

[4269] GAM299 gene, herein designated GAM GENE, and GAM299 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4270] GAM299 gene, herein designated GAM GENE, encodes a GAM299 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM299 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM299 precursor RNA is designated SEQ ID:4521, and is provided hereinbelow with reference to the sequence listing part.

[4271] GAM299 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM299 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4272] An enzyme complex designated DICER COMPLEX, dices the GAM299 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM299 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM299 RNA is designated SEQ ID:4522, and is provided hereinbelow with reference to the sequence listing part.

[4273] GAM299 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM299 target RNA, herein designated GAM TARGET RNA. GAM299 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4274] GAM299 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM299 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM299 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM299 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM299 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4275] The complementary binding of GAM299 RNA, herein designated GAM RNA, to target binding sites on GAM299 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM299 target RNA, herein designated GAM TARGET RNA, into GAM299 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4276] It is appreciated that GAM299 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM299 target genes. The mRNA of each one of this plurality of GAM299 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM299 RNA,

herein designated GAM RNA, and which when bound by GAM299 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM299 target proteins.

[4277] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM299 gene, herein designated GAM GENE, on one or more GAM299 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4278] It is yet further appreciated that specific functions, and accordingly utilities, of GAM299 correlate with, and may be deduced from, the identity of the target genes which

GAM299 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4279] Nucleotide sequences of the GAM299 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM299 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM299 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM299 are further described hereinbelow with reference to Table Table1.

[4280] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM299 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4281]

[4282] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 300 (GAM300) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4283] GAM300 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM300 was detected is described hereinabove with reference to Figs. 2-8.

[4284] GAM300 gene, herein designated GAM GENE, and GAM300 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4285] GAM300 gene, herein designated GAM GENE, encodes a GAM300 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM300 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM300 precursor RNA is designated SEQ ID:4544, and is provided hereinbelow with reference to the sequence listing part.

[4286] GAM300 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM300 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[4287] An enzyme complex designated DICER COMPLEX, dices the GAM300 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM300 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 20%) nucleotide sequence of GAM300 RNA is designated SEQ ID:4545, and is provided hereinbelow with reference to the sequence listing part.

[4288] GAM300 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM300 target RNA, herein designated GAM TARGET RNA.
GAM300 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4289] GAM300 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM300 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM300 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM300 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM300 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4290] The complementary binding of GAM300 RNA, herein designated GAM RNA, to target binding sites on GAM300 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM300 target RNA, herein desig-

nated GAM TARGET RNA, into GAM300 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4291] It is appreciated that GAM300 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM300 target genes. The mRNA of each one of this plurality of GAM300 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM300 RNA, herein designated GAM RNA, and which when bound by GAM300 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM300 target proteins.

[4292] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM300 gene, herein designated GAM GENE, on one or more GAM300 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4293] It is yet further appreciated that specific functions, and accordingly utilities, of GAM300 correlate with, and may be deduced from, the identity of the target genes which GAM300 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4294] Nucleotide sequences of the GAM300 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM300 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM300 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM300 are further described hereinbelow with reference to Table Table1.

[4295] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM300 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[4296]

[4297] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 301 (GAM301) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4298] GAM301 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM301 was detected is described hereinabove with reference to Figs. 2-8.

[4299] GAM301 gene, herein designated GAM GENE, and GAM301 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4300] GAM301 gene, herein designated GAM GENE, encodes a GAM301 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM301 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM301 precursor RNA is designated SEQ ID:4556, and is provided hereinbelow with ref-

erence to the sequence listing part.

[4301] GAM301 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM301 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4302] An enzyme complex designated DICER COMPLEX, dices the GAM301 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM301 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM301 RNA is designated SEQ ID:4557, and is provided hereinbelow with reference to the sequence listing part.

[4303] GAM301 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM301 target RNA, herein designated GAM TARGET RNA.

GAM301 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4304] GAM301 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM301 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM301 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM301 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM301 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4305] The complementary binding of GAM301 RNA, herein designated GAM RNA, to target binding sites on GAM301 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM301 target RNA, herein designated GAM TARGET RNA, into GAM301 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4306] It is appreciated that GAM301 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM301 target genes. The mRNA of each one of this plurality of GAM301 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM301 RNA, herein designated GAM RNA, and which when bound by GAM301 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM301 target proteins.

[4307] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM301 gene, herein designated GAM GENE, on one or more GAM301 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4308] It is yet further appreciated that specific functions, and accordingly utilities, of GAM301 correlate with, and may be deduced from, the identity of the target genes which GAM301 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4309] Nucleotide sequences of the GAM301 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM301 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM301 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM301 are further described hereinbelow with reference to Table Table1.

[4310] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM301 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4311]

[4312] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 302 (GAM302) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4313] GAM302 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM302 was detected is described hereinabove with reference to Figs. 2-8.

[4314] GAM302 gene, herein designated GAM GENE, and GAM302 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[4315] GAM302 gene, herein designated GAM GENE, encodes a GAM302 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM302 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM302 precursor RNA is designated SEQ ID:4578, and is provided hereinbelow with reference to the sequence listing part.

[4316] GAM302 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM302 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4317] An enzyme complex designated DICER COMPLEX, dices the GAM302 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM302 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM302 RNA is designated SEQ ID:4579, and is provided hereinbelow with reference to the sequence listing part.

[4318] GAM302 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM302 target RNA, herein designated GAM TARGET RNA. GAM302 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4319] GAM302 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM302 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM302 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM302 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM302 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4320] The complementary binding of GAM302 RNA, herein designated GAM RNA, to target binding sites on GAM302 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM302 target RNA, herein designated GAM TARGET RNA, into GAM302 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4321] It is appreciated that GAM302 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM302 target genes. The mRNA of each one of this plurality of GAM302 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM302 RNA, herein designated GAM RNA, and which when bound by GAM302 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM302 target proteins.

[4322] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM302 gene, herein designated GAM GENE, on one or more GAM302 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[4323] It is yet further appreciated that specific functions, and accordingly utilities, of GAM302 correlate with, and may be deduced from, the identity of the target genes which GAM302 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4324] Nucleotide sequences of the GAM302 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM302 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM302 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM302 are further described hereinbelow with reference to Table Table1.

[4325] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM302 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4326]

[4327] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

303 (GAM303) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4328] GAM303 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM303 was detected is described hereinabove with reference to Figs. 2-8.

[4329] GAM303 gene, herein designated GAM GENE, and GAM303 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4330] GAM303 gene, herein designated GAM GENE, encodes a GAM303 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM303 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM303 precursor RNA is designated SEQ ID:4591, and is provided hereinbelow with reference to the sequence listing part.

[4331] GAM303 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM303 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4332] An enzyme complex designated DICER COMPLEX, dices the GAM303 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM303 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM303 RNA is designated SEQ ID:4592, and is provided hereinbelow with reference to the sequence listing part.

[4333] GAM303 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM303 target RNA, herein designated GAM TARGET RNA. GAM303 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4334] GAM303 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM303 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM303 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM303 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM303 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4335] The complementary binding of GAM303 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM303 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM303 target RNA, herein designated GAM TARGET RNA, into GAM303 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4336] It is appreciated that GAM303 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM303 target genes. The mRNA of each one of this plurality of GAM303 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM303 RNA, herein designated GAM RNA, and which when bound by GAM303 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM303 target proteins.

[4337] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM303 gene, herein designated GAM GENE, on one or more GAM303 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4338] It is yet further appreciated that specific functions, and accordingly utilities, of GAM303 correlate with, and may be deduced from, the identity of the target genes which GAM303 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4339] Nucleotide sequences of the GAM303 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM303 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM303 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM303 are further described hereinbelow with reference to Table Table1.

[4340] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM303 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4341]

[4342] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 304 (GAM304) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4343] GAM304 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM304 was detected is described hereinabove with reference to Figs. 2-8.

[4344] GAM304 gene, herein designated GAM GENE, and GAM304 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4345] GAM304 gene, herein designated GAM GENE, encodes a GAM304 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM304 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM304 precursor RNA is designated SEQ ID:4618, and is provided hereinbelow with reference to the sequence listing part.

[4346] GAM304 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM304 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4347] An enzyme complex designated DICER COMPLEX, dices the GAM304 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM304 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 52%) nucleotide sequence of GAM304 RNA is designated SEQ ID:4619, and is provided hereinbelow with reference to the sequence listing part.

[4348] GAM304 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM304 target RNA, herein designated GAM TARGET RNA. GAM304 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4349] GAM304 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM304 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM304 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM304 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM304 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4350] The complementary binding of GAM304 RNA, herein designated GAM RNA, to target binding sites on GAM304 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM304 target RNA, herein designated GAM TARGET RNA, into GAM304 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4351] It is appreciated that GAM304 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM304 target genes. The mRNA of each one of this plurality of GAM304 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM304 RNA, herein designated GAM RNA, and which when bound by

GAM304 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM304 target proteins.

[4352] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM304 gene, herein designated GAM GENE, on one or more GAM304 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4353] It is yet further appreciated that specific functions, and accordingly utilities, of GAM304 correlate with, and may be deduced from, the identity of the target genes which GAM304 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[4354] Nucleotide sequences of the GAM304 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM304 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM304 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM304 are further described hereinbelow with reference to Table Table1.

[4355] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM304 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4356]

[4357] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 305 (GAM305) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4358] GAM305 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM305 was detected is described hereinabove with reference to Figs. 2-8.

[4359] GAM305 gene, herein designated GAM GENE, and GAM305 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4360] GAM305 gene, herein designated GAM GENE, encodes a GAM305 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM305 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM305 precursor RNA is designated SEQ ID:4627, and is provided hereinbelow with reference to the sequence listing part.

[4361] GAM305 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM305 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[4362] An enzyme complex designated DICER COMPLEX, dices the GAM305 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM305 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM305 RNA is designated SEQ ID:4628, and is provided hereinbelow with reference to the sequence listing part.

[4363] GAM305 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM305 target RNA, herein designated GAM TARGET RNA. GAM305 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4364] GAM305 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM305 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM305 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM305 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM305 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4365] The complementary binding of GAM305 RNA, herein designated GAM RNA, to target binding sites on GAM305 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM305 target RNA, herein designated GAM TARGET RNA, into GAM305 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4366] It is appreciated that GAM305 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM305 target genes. The mRNA of each one of this plurality of GAM305 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM305 RNA, herein designated GAM RNA, and which when bound by GAM305 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM305 target proteins.

[4367] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM305 gene, herein designated GAM GENE, on one or more GAM305 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4368] It is yet further appreciated that specific functions, and accordingly utilities, of GAM305 correlate with, and may be deduced from, the identity of the target genes which GAM305 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4369] Nucleotide sequences of the GAM305 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM305 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM305 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM305 are further described hereinbelow with reference to Table Table1.

[4370] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM305 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4371]

[4372] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 306 (GAM306) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4373] GAM306 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM306 was detected is described hereinabove with reference to Figs. 2-8.

[4374] GAM306 gene, herein designated GAM GENE, and GAM306 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4375] GAM306 gene, herein designated GAM GENE, encodes a GAM306 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM306 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM306 precursor RNA is designated SEQ ID:4647, and is provided hereinbelow with reference to the sequence listing part.

[4376] GAM306 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM306 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4377] An enzyme complex designated DICER COMPLEX, dices the GAM306 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM306 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM306 RNA is designated SEQ ID:2738, and is provided hereinbelow with reference to the sequence listing part.

[4378] GAM306 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM306

target RNA, herein designated GAM TARGET RNA.

GAM306 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4379] GAM306 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM306 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM306 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM306 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM306 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4380] The complementary binding of GAM306 RNA, herein designated GAM RNA, to target binding sites on GAM306 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM306 target RNA, herein designated GAM TARGET RNA, into GAM306 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4381] It is appreciated that GAM306 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM306 target genes. The mRNA of each one of this plurality of GAM306 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM306 RNA, herein designated GAM RNA, and which when bound by GAM306 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM306 target proteins.

[4382] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM306 gene, herein designated GAM GENE, on one or more GAM306 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4383] It is yet further appreciated that specific functions, and accordingly utilities, of GAM306 correlate with, and may be deduced from, the identity of the target genes which GAM306 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4384] Nucleotide sequences of the GAM306 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM306 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM306 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM306 are further described hereinbelow with reference to Table Table1.

[4385] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM306 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4386]

[4387] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 307 (GAM307) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4388] GAM307 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM307 was detected is described hereinabove with reference to Figs. 2-8.

[4389] GAM307 gene, herein designated GAM GENE, and GAM307 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4390] GAM307 gene, herein designated GAM GENE, encodes a GAM307 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM307 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM307 precursor RNA is designated SEQ ID:4652, and is provided hereinbelow with reference to the sequence listing part.

[4391] GAM307 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM307 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4392] An enzyme complex designated DICER COMPLEX, dices the GAM307 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM307 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 38%) nucleotide sequence of GAM307 RNA is designated SEQ ID:4653, and is provided hereinbelow with reference to the sequence listing part.

[4393] GAM307 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM307 target RNA, herein designated GAM TARGET RNA. GAM307 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4394] GAM307 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM307 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM307 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM307 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM307 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4395] The complementary binding of GAM307 RNA, herein designated GAM RNA, to target binding sites on GAM307 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM307 target RNA, herein designated GAM TARGET RNA, into GAM307 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4396] It is appreciated that GAM307 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM307 target genes. The mRNA of each one of this plu-

rality of GAM307 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM307 RNA, herein designated GAM RNA, and which when bound by GAM307 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM307 target proteins.

[4397] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM307 gene, herein designated GAM GENE, on one or more GAM307 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4398] It is yet further appreciated that specific functions, and accordingly utilities, of GAM307 correlate with, and may be deduced from, the identity of the target genes which GAM307 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4399] Nucleotide sequences of the GAM307 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM307 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM307 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM307 are further described hereinbelow with reference to Table Table1.

[4400] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM307 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4401]

[4402] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 308 (GAM308) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[4403] GAM308 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM308 was detected is described hereinabove with reference to Figs. 2-8.

[4404] GAM308 gene, herein designated GAM GENE, and GAM308 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4405] GAM308 gene, herein designated GAM GENE, encodes a GAM308 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM308 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM308 precursor RNA is designated SEQ ID:4667, and is provided hereinbelow with reference to the sequence listing part.

[4406] GAM308 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM308 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4407] An enzyme complex designated DICER COMPLEX, dices the GAM308 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM308 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM308 RNA is designated SEQ ID:4668, and is provided hereinbelow with reference to the sequence listing part.

[4408] GAM308 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM308 target RNA, herein designated GAM TARGET RNA.
GAM308 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[4409] GAM308 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM308 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM308 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM308 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM308 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4410] The complementary binding of GAM308 RNA, herein designated GAM RNA, to target binding sites on GAM308 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM308 target RNA, herein designated GAM TARGET RNA, into GAM308 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4411] It is appreciated that GAM308 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM308 target genes. The mRNA of each one of this plurality of GAM308 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM308 RNA, herein designated GAM RNA, and which when bound by GAM308 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM308 target proteins.

[4412] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM308 gene, herein designated GAM GENE, on one or more GAM308 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4413] It is yet further appreciated that specific functions, and accordingly utilities, of GAM308 correlate with, and may be deduced from, the identity of the target genes which GAM308 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4414] Nucleotide sequences of the GAM308 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM308 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM308 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM308 are further described hereinbelow with reference to Table Table1.

[4415] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM308 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4416]

[4417] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 309 (GAM309) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4418] GAM309 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM309 was detected is described hereinabove with reference to Figs. 2-8.

[4419] GAM309 gene, herein designated GAM GENE, and GAM309 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4420] GAM309 gene, herein designated GAM GENE, encodes a GAM309 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM309 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM309 precursor RNA is designated SEQ ID:4680, and is provided hereinbelow with reference to the sequence listing part.

[4421] GAM309 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM309 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4422] An enzyme complex designated DICER COMPLEX, dices the GAM309 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM309 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 38%) nucleotide sequence of

GAM309 RNA is designated SEQ ID:4681, and is provided hereinbelow with reference to the sequence listing part.

[4423] GAM309 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM309 target RNA, herein designated GAM TARGET RNA. GAM309 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4424] GAM309 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM309 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM309 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM309 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM309 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4425] The complementary binding of GAM309 RNA, herein designated GAM RNA, to target binding sites on GAM309 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM309 target RNA, herein designated GAM TARGET RNA, into GAM309 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4426] It is appreciated that GAM309 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM309 target genes. The mRNA of each one of this plurality of GAM309 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM309 RNA, herein designated GAM RNA, and which when bound by GAM309 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM309 target proteins.

[4427] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM309 gene, herein designated GAM GENE, on one or more GAM309 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4428] It is yet further appreciated that specific functions, and accordingly utilities, of GAM309 correlate with, and may be deduced from, the identity of the target genes which GAM309 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4429] Nucleotide sequences of the GAM309 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM309 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM309 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM309 are further described hereinbelow with reference to Table Table1.

[4430] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM309 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4431]

[4432] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 310 (GAM310) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4433] GAM310 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM310 was detected is described hereinabove

with reference to Figs. 2-8.

[4434] GAM310 gene, herein designated GAM GENE, and GAM310 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4435] GAM310 gene, herein designated GAM GENE, encodes a GAM310 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM310 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM310 precursor RNA is designated SEQ ID:4701, and is provided hereinbelow with reference to the sequence listing part.

[4436] GAM310 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM310 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4437] An enzyme complex designated DICER COMPLEX, dices the GAM310 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM310 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM310 RNA is designated SEQ ID:4702, and is provided hereinbelow with reference to the sequence listing part.

[4438] GAM310 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM310 target RNA, herein designated GAM TARGET RNA. GAM310 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4439] GAM310 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM310 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM310 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM310 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM310 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4440] The complementary binding of GAM310 RNA, herein designated GAM RNA, to target binding sites on GAM310 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM310 target RNA, herein designated GAM TARGET RNA, into GAM310 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[4441] It is appreciated that GAM310 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM310 target genes. The mRNA of each one of this plurality of GAM310 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM310 RNA, herein designated GAM RNA, and which when bound by GAM310 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM310 target proteins.

[4442] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM310 gene, herein designated GAM GENE, on one or more GAM310 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4443] It is yet further appreciated that specific functions, and accordingly utilities, of GAM310 correlate with, and may be deduced from, the identity of the target genes which GAM310 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4444] Nucleotide sequences of the GAM310 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM310 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM310 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM310 are further described hereinbelow with reference to Table Table1.

[4445] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM310 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4446]

[4447] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 311 (GAM311) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4448] GAM311 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM311 was detected is described hereinabove with reference to Figs. 2-8.

[4449] GAM311 gene, herein designated GAM GENE, and GAM311 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4450] GAM311 gene, herein designated GAM GENE, encodes a GAM311 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM311 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM311 precursor RNA is designated SEQ ID:4719, and is provided hereinbelow with reference to the sequence listing part.

[4451] GAM311 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM311 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4452] An enzyme complex designated DICER COMPLEX, dices the GAM311 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM311 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of GAM311 RNA is designated SEQ ID:4720, and is provided hereinbelow with reference to the sequence listing part.

[4453] GAM311 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM311 target RNA, herein designated GAM TARGET RNA.

GAM311 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4454] GAM311 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM311 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM311 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM311 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM311 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4455] The complementary binding of GAM311 RNA, herein designated GAM RNA, to target binding sites on GAM311 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM311 target RNA, herein designated GAM TARGET RNA, into GAM311 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4456] It is appreciated that GAM311 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM311 target genes. The mRNA of each one of this plurality of GAM311 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM311 RNA, herein designated GAM RNA, and which when bound by GAM311 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM311 target proteins.

[4457] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM311 gene, herein designated GAM GENE, on one or more GAM311 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4458] It is yet further appreciated that specific functions, and accordingly utilities, of GAM311 correlate with, and may be deduced from, the identity of the target genes which GAM311 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4459] Nucleotide sequences of the GAM311 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM311 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM311 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM311 are further described hereinbelow with reference to Table Table1.

[4460] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM311 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4461]

[4462] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 312 (GAM312) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4463] GAM312 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM312 was detected is described hereinabove with reference to Figs. 2-8.

[4464] GAM312 gene, herein designated GAM GENE, and GAM312 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4465] GAM312 gene, herein designated GAM GENE, encodes a

GAM312 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM312 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM312 precursor RNA is designated SEQ ID:4738, and is provided hereinbelow with reference to the sequence listing part.

[4466] GAM312 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM312 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4467] An enzyme complex designated DICER COMPLEX, dices the GAM312 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM312 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM312 RNA is designated SEQ ID:4739, and is provided hereinbelow with reference to the sequence listing part.

[4468] GAM312 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM312 target RNA, herein designated GAM TARGET RNA. GAM312 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4469] GAM312 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM312 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM312 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM312 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM312 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4470] The complementary binding of GAM312 RNA, herein designated GAM RNA, to target binding sites on GAM312 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM312 target RNA, herein designated GAM TARGET RNA, into GAM312 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4471] It is appreciated that GAM312 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM312 target genes. The mRNA of each one of this plurality of GAM312 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM312 RNA, herein designated GAM RNA, and which when bound by GAM312 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM312 target proteins.

[4472] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM312 gene, herein designated GAM GENE, on one or more GAM312 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4473] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM312 correlate with, and may be deduced from, the identity of the target genes which GAM312 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4474] Nucleotide sequences of the GAM312 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM312 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM312 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM312 are further described hereinbelow with reference to Table Table1.

[4475] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM312 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4476]

[4477] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 313 (GAM313) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[4478] GAM313 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM313 was detected is described hereinabove with reference to Figs. 2-8.

[4479] GAM313 gene, herein designated GAM GENE, and GAM313 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4480] GAM313 gene, herein designated GAM GENE, encodes a GAM313 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM313 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM313 precursor RNA is designated SEQ ID:4771, and is provided hereinbelow with reference to the sequence listing part.

[4481] GAM313 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM313 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4482] An enzyme complex designated DICER COMPLEX, dices the GAM313 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM313 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM313 RNA is designated SEQ ID:4772, and is provided hereinbelow with reference to the sequence listing part.

[4483] GAM313 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM313 target RNA, herein designated GAM TARGET RNA. GAM313 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4484] GAM313 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM313 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM313 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM313 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM313 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4485] The complementary binding of GAM313 RNA, herein designated GAM RNA, to target binding sites on GAM313 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM313 target RNA, herein designated GAM TARGET RNA, into GAM313 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4486] It is appreciated that GAM313 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM313 target genes. The mRNA of each one of this plurality of GAM313 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM313 RNA, herein designated GAM RNA, and which when bound by GAM313 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM313 target proteins.

[4487] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM313 gene, herein designated GAM GENE, on one or more GAM313 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4488] It is yet further appreciated that specific functions, and accordingly utilities, of GAM313 correlate with, and may be deduced from, the identity of the target genes which GAM313 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4489] Nucleotide sequences of the GAM313 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM313 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM313 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM313 are further described hereinbelow with reference to Table Table1.

[4490] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM313 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4491]

[4492] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 314 (GAM314) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4493] GAM314 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM314 was detected is described hereinabove with reference to Figs. 2-8.

[4494] GAM314 gene, herein designated GAM GENE, and GAM314 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4495] GAM314 gene, herein designated GAM GENE, encodes a GAM314 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM314 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM314 precursor RNA is designated SEQ ID:4781, and is provided hereinbelow with reference to the sequence listing part.

[4496] GAM314 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM314 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4497] An enzyme complex designated DICER COMPLEX, dices the GAM314 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM314 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 33%) nucleotide sequence of GAM314 RNA is designated SEQ ID:4782, and is provided

hereinbelow with reference to the sequence listing part.

[4498] GAM314 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM314 target RNA, herein designated GAM TARGET RNA. GAM314 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4499] GAM314 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM314 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM314 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM314 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM314 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4500] The complementary binding of GAM314 RNA, herein designated GAM RNA, to target binding sites on GAM314 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM314 target RNA, herein designated GAM TARGET RNA, into GAM314 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4501] It is appreciated that GAM314 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM314 target genes. The mRNA of each one of this plurality of GAM314 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM314 RNA, herein designated GAM RNA, and which when bound by GAM314 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM314 tar-

get proteins.

[4502] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM314 gene, herein designated GAM GENE, on one or more GAM314 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4503] It is yet further appreciated that specific functions, and accordingly utilities, of GAM314 correlate with, and may be deduced from, the identity of the target genes which GAM314 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4504] Nucleotide sequences of the GAM314 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM314 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM314 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM314 are further described hereinbelow with reference to Table Table1.

[4505] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM314 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4506]

[4507] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 315 (GAM315) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4508] GAM315 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM315 was detected is described hereinabove with reference to Figs. 2-8.

[4509] GAM315 gene, herein designated GAM GENE, and GAM315 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4510] GAM315 gene, herein designated GAM GENE, encodes a GAM315 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM315 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM315 precursor RNA is designated SEQ ID:4798, and is provided hereinbelow with reference to the sequence listing part.

[4511] GAM315 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM315 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4512] An enzyme complex designated DICER COMPLEX, dices

the GAM315 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM315 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM315 RNA is designated SEQ ID:4799, and is provided hereinbelow with reference to the sequence listing part.

[4513] GAM315 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM315 target RNA, herein designated GAM TARGET RNA. GAM315 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4514] GAM315 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM315 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM315

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM315 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM315 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4515] The complementary binding of GAM315 RNA, herein designated GAM RNA, to target binding sites on GAM315 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM315 target RNA, herein designated GAM TARGET RNA, into GAM315 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4516] It is appreciated that GAM315 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM315 target genes. The mRNA of each one of this plurality of GAM315 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM315 RNA, herein designated GAM RNA, and which when bound by GAM315 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM315 target proteins.

[4517] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM315 gene, herein designated GAM GENE, on one or more GAM315 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4518] It is yet further appreciated that specific functions, and accordingly utilities, of GAM315 correlate with, and may be deduced from, the identity of the target genes which GAM315 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4519] Nucleotide sequences of the GAM315 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM315 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM315 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM315 are further described hereinbelow with reference to Table Table1.

[4520] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM315 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4521]

[4522] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 316 (GAM316) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4523] GAM316 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM316 was detected is described hereinabove with reference to Figs. 2-8.

[4524] GAM316 gene, herein designated GAM GENE, and GAM316 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4525] GAM316 gene, herein designated GAM GENE, encodes a GAM316 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM316 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM316 precursor RNA is designated SEQ ID:4812, and is provided hereinbelow with reference to the sequence listing part.

[4526] GAM316 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM316 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4527] An enzyme complex designated DICER COMPLEX, dices the GAM316 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM316 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM316 RNA is designated SEQ ID:4813, and is provided hereinbelow with reference to the sequence listing part.

[4528] GAM316 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM316 target RNA, herein designated GAM TARGET RNA. GAM316 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4529] GAM316 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM316 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM316 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM316 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM316 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[4530] The complementary binding of GAM316 RNA, herein designated GAM RNA, to target binding sites on GAM316 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM316 target RNA, herein designated GAM TARGET RNA, into GAM316 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4531] It is appreciated that GAM316 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM316 target genes. The mRNA of each one of this plurality of GAM316 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM316 RNA, herein designated GAM RNA, and which when bound by GAM316 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM316 target proteins.

[4532] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM316 gene, herein designated GAM GENE, on one or

more GAM316 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4533] It is yet further appreciated that specific functions, and accordingly utilities, of GAM316 correlate with, and may be deduced from, the identity of the target genes which GAM316 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4534] Nucleotide sequences of the GAM316 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM316 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM316 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM316 are further de-

scribed hereinbelow with reference to Table Table1.

[4535] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM316 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4536]

[4537] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 317 (GAM317) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4538] GAM317 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM317 was detected is described hereinabove with reference to Figs. 2-8.

[4539] GAM317 gene, herein designated GAM GENE, and GAM317 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4540] GAM317 gene, herein designated GAM GENE, encodes a GAM317 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM317 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM317 precursor RNA is designated SEQ ID:4819, and is provided hereinbelow with reference to the sequence listing part.

[4541] GAM317 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM317 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4542] An enzyme complex designated DICER COMPLEX, dices the GAM317 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM317 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of GAM317 RNA is designated SEQ ID:4820, and is provided hereinbelow with reference to the sequence listing part.

[4543] GAM317 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM317 target RNA, herein designated GAM TARGET RNA. GAM317 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4544] GAM317 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM317 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM317 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM317 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM317 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4545] The complementary binding of GAM317 RNA, herein designated GAM RNA, to target binding sites on GAM317 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM317 target RNA, herein designated GAM TARGET RNA, into GAM317 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4546] It is appreciated that GAM317 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM317 target genes. The mRNA of each one of this plurality of GAM317 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM317 RNA, herein designated GAM RNA, and which when bound by GAM317 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM317 target proteins.

[4547] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM317 gene, herein designated GAM GENE, on one or more GAM317 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4548] It is yet further appreciated that specific functions, and accordingly utilities, of GAM317 correlate with, and may

be deduced from, the identity of the target genes which GAM317 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4549] Nucleotide sequences of the GAM317 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM317 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM317 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM317 are further described hereinbelow with reference to Table Table1.

[4550] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM317 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4551]

[4552] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 318 (GAM318) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4553] GAM318 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM318 was detected is described hereinabove with reference to Figs. 2-8.

[4554] GAM318 gene, herein designated GAM GENE, and GAM318 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4555] GAM318 gene, herein designated GAM GENE, encodes a GAM318 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM318 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM318 precursor RNA is designated SEQ ID:4843, and is provided hereinbelow with reference to the sequence listing part.

[4556] GAM318 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM318 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4557] An enzyme complex designated DICER COMPLEX, dices the GAM318 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM318 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 39%) nucleotide sequence of GAM318 RNA is designated SEQ ID:4844, and is provided hereinbelow with reference to the sequence listing part.

[4558] GAM318 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM318 target RNA, herein designated GAM TARGET RNA. GAM318 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4559] GAM318 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM318 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM318 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM318 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM318 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4560] The complementary binding of GAM318 RNA, herein designated GAM RNA, to target binding sites on GAM318 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM318 target RNA, herein designated GAM TARGET RNA, into GAM318 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4561] It is appreciated that GAM318 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM318 target genes. The mRNA of each one of this plurality of GAM318 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM318 RNA, herein designated GAM RNA, and which when bound by GAM318 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM318 target proteins.

[4562] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM318 gene, herein designated GAM GENE, on one or more GAM318 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4563] It is yet further appreciated that specific functions, and accordingly utilities, of GAM318 correlate with, and may be deduced from, the identity of the target genes which GAM318 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4564] Nucleotide sequences of the GAM318 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM318 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM318 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM318 are further described hereinbelow with reference to Table Table1.

[4565] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM318 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4566]

[4567] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 319 (GAM319) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4568] GAM319 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM319 was detected is described hereinabove with reference to Figs. 2-8.

[4569] GAM319 gene, herein designated GAM GENE, and GAM319 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4570] GAM319 gene, herein designated GAM GENE, encodes a GAM319 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM319 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM319 precursor RNA is desig-

nated SEQ ID:4858, and is provided hereinbelow with reference to the sequence listing part.

[4571] GAM319 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM319 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4572] An enzyme complex designated DICER COMPLEX, dices the GAM319 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM319 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM319 RNA is designated SEQ ID:4859, and is provided hereinbelow with reference to the sequence listing part.

[4573] GAM319 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM319 target RNA, herein designated GAM TARGET RNA. GAM319 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4574] GAM319 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM319 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM319 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM319 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM319

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4575] The complementary binding of GAM319 RNA, herein designated GAM RNA, to target binding sites on GAM319 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM319 target RNA, herein designated GAM TARGET RNA, into GAM319 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4576] It is appreciated that GAM319 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM319 target genes. The mRNA of each one of this plurality of GAM319 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM319 RNA, herein designated GAM RNA, and which when bound by GAM319 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM319 target proteins.

[4577] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM319 gene, herein designated GAM GENE, on one or more GAM319 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4578] It is yet further appreciated that specific functions, and accordingly utilities, of GAM319 correlate with, and may be deduced from, the identity of the target genes which GAM319 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4579] Nucleotide sequences of the GAM319 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM319 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM319 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM319 are further described hereinbelow with reference to Table Table1.

[4580] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM319 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4581]

[4582] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 320 (GAM320) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4583] GAM320 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM320 was detected is described hereinabove with reference to Figs. 2-8.

[4584] GAM320 gene, herein designated GAM GENE, and

GAM320 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4585] GAM320 gene, herein designated GAM GENE, encodes a GAM320 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM320 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM320 precursor RNA is designated SEQ ID:4873, and is provided hereinbelow with reference to the sequence listing part.

[4586] GAM320 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM320 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4587] An enzyme complex designated DICER COMPLEX, dices the GAM320 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM320 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM320 RNA is designated SEQ ID:4874, and is provided hereinbelow with reference to the sequence listing part.

[4588] GAM320 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM320 target RNA, herein designated GAM TARGET RNA. GAM320 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4589] GAM320 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM320 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM320 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM320 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM320 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4590] The complementary binding of GAM320 RNA, herein designated GAM RNA, to target binding sites on GAM320 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM320 target RNA, herein designated GAM TARGET RNA, into GAM320 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4591] It is appreciated that GAM320 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM320 target genes. The mRNA of each one of this plurality of GAM320 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM320 RNA, herein designated GAM RNA, and which when bound by GAM320 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM320 target proteins.

[4592] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM320 gene, herein designated GAM GENE, on one or more GAM320 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4593] It is yet further appreciated that specific functions, and accordingly utilities, of GAM320 correlate with, and may be deduced from, the identity of the target genes which GAM320 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4594] Nucleotide sequences of the GAM320 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM320 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM320 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM320 are further described hereinbelow with reference to Table Table1.

[4595] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM320 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4596]

[4597] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 321 (GAM321) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4598] GAM321 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM321 was detected is described hereinabove with reference to Figs. 2-8.

[4599] GAM321 gene, herein designated GAM GENE, and GAM321 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4600] GAM321 gene, herein designated GAM GENE, encodes a GAM321 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM321 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM321 precursor RNA is designated SEQ ID:4918, and is provided hereinbelow with reference to the sequence listing part.

[4601] GAM321 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM321 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4602] An enzyme complex designated DICER COMPLEX, dices the GAM321 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM321 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM321 RNA is designated SEQ ID:4919, and is provided hereinbelow with reference to the sequence listing part.

[4603] GAM321 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM321 target RNA, herein designated GAM TARGET RNA. GAM321 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4604] GAM321 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM321 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM321 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM321 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM321 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4605] The complementary binding of GAM321 RNA, herein designated GAM RNA, to target binding sites on GAM321 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM321 target RNA, herein designated GAM TARGET RNA, into GAM321 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4606] It is appreciated that GAM321 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM321 target genes. The mRNA of each one of this plurality of GAM321 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM321 RNA, herein designated GAM RNA, and which when bound by GAM321 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM321 target proteins.

[4607] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM321 gene, herein designated GAM GENE, on one or more GAM321 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4608] It is yet further appreciated that specific functions, and accordingly utilities, of GAM321 correlate with, and may be deduced from, the identity of the target genes which GAM321 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4609] Nucleotide sequences of the GAM321 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM321 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM321 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM321 are further described hereinbelow with reference to Table Table1.

[4610] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM321 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4611]

[4612] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 322 (GAM322) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4613] GAM322 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM322 was detected is described hereinabove with reference to Figs. 2-8.

[4614] GAM322 gene, herein designated GAM GENE, and GAM322 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4615] GAM322 gene, herein designated GAM GENE, encodes a GAM322 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM322 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM322 precursor RNA is designated SEQ ID:4923, and is provided hereinbelow with reference to the sequence listing part.

[4616] GAM322 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM322 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4617] An enzyme complex designated DICER COMPLEX, dices the GAM322 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM322 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM322 RNA is designated SEQ ID:4924, and is provided hereinbelow with reference to the sequence listing part.

[4618] GAM322 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM322 target RNA, herein designated GAM TARGET RNA. GAM322 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4619] GAM322 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM322 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM322 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM322 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM322 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4620] The complementary binding of GAM322 RNA, herein designated GAM RNA, to target binding sites on GAM322 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM322 target RNA, herein designated GAM TARGET RNA, into GAM322 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4621] It is appreciated that GAM322 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM322 target genes. The mRNA of each one of this plurality of GAM322 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM322 RNA,

herein designated GAM RNA, and which when bound by GAM322 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM322 target proteins.

[4622] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM322 gene, herein designated GAM GENE, on one or more GAM322 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4623] It is yet further appreciated that specific functions, and accordingly utilities, of GAM322 correlate with, and may be deduced from, the identity of the target genes which

GAM322 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4624] Nucleotide sequences of the GAM322 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM322 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM322 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM322 are further described hereinbelow with reference to Table Table1.

[4625] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM322 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4626]

[4627] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 323 (GAM323) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4628] GAM323 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM323 was detected is described hereinabove with reference to Figs. 2-8.

[4629] GAM323 gene, herein designated GAM GENE, and GAM323 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4630] GAM323 gene, herein designated GAM GENE, encodes a GAM323 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM323 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM323 precursor RNA is designated SEQ ID:4942, and is provided hereinbelow with reference to the sequence listing part.

[4631] GAM323 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM323 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[4632] An enzyme complex designated DICER COMPLEX, dices the GAM323 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM323 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 25%) nucleotide sequence of GAM323 RNA is designated SEQ ID:4943, and is provided hereinbelow with reference to the sequence listing part.

[4633] GAM323 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM323 target RNA, herein designated GAM TARGET RNA.
GAM323 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4634] GAM323 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM323 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM323 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM323 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM323 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4635] The complementary binding of GAM323 RNA, herein designated GAM RNA, to target binding sites on GAM323 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM323 target RNA, herein desig-

nated GAM TARGET RNA, into GAM323 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4636] It is appreciated that GAM323 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM323 target genes. The mRNA of each one of this plurality of GAM323 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM323 RNA, herein designated GAM RNA, and which when bound by GAM323 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM323 target proteins.

[4637] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM323 gene, herein designated GAM GENE, on one or more GAM323 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4638] It is yet further appreciated that specific functions, and accordingly utilities, of GAM323 correlate with, and may be deduced from, the identity of the target genes which GAM323 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4639] Nucleotide sequences of the GAM323 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM323 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM323 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM323 are further described hereinbelow with reference to Table Table1.

[4640] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM323 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[4641]

[4642] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 324 (GAM324) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4643] GAM324 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM324 was detected is described hereinabove with reference to Figs. 2-8.

[4644] GAM324 gene, herein designated GAM GENE, and GAM324 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4645] GAM324 gene, herein designated GAM GENE, encodes a GAM324 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM324 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM324 precursor RNA is designated SEQ ID:4958, and is provided hereinbelow with ref-

erence to the sequence listing part.

[4646] GAM324 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM324 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4647] An enzyme complex designated DICER COMPLEX, dices the GAM324 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM324 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM324 RNA is designated SEQ ID:4959, and is provided hereinbelow with reference to the sequence listing part.

[4648] GAM324 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM324 target RNA, herein designated GAM TARGET RNA.

GAM324 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4649] GAM324 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM324 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM324 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM324 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM324 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4650] The complementary binding of GAM324 RNA, herein designated GAM RNA, to target binding sites on GAM324 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM324 target RNA, herein designated GAM TARGET RNA, into GAM324 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4651] It is appreciated that GAM324 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM324 target genes. The mRNA of each one of this plurality of GAM324 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM324 RNA, herein designated GAM RNA, and which when bound by GAM324 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM324 target proteins.

[4652] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM324 gene, herein designated GAM GENE, on one or more GAM324 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4653] It is yet further appreciated that specific functions, and accordingly utilities, of GAM324 correlate with, and may be deduced from, the identity of the target genes which GAM324 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4654] Nucleotide sequences of the GAM324 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM324 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM324 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM324 are further described hereinbelow with reference to Table Table1.

[4655] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM324 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4656]

[4657] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 325 (GAM325) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4658] GAM325 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM325 was detected is described hereinabove with reference to Figs. 2-8.

[4659] GAM325 gene, herein designated GAM GENE, and GAM325 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[4660] GAM325 gene, herein designated GAM GENE, encodes a GAM325 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM325 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM325 precursor RNA is designated SEQ ID:4970, and is provided hereinbelow with reference to the sequence listing part.

[4661] GAM325 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM325 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4662] An enzyme complex designated DICER COMPLEX, dices the GAM325 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM325 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM325 RNA is designated SEQ ID:4971, and is provided hereinbelow with reference to the sequence listing part.

[4663] GAM325 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM325 target RNA, herein designated GAM TARGET RNA. GAM325 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4664] GAM325 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM325 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM325 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM325 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM325 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4665] The complementary binding of GAM325 RNA, herein designated GAM RNA, to target binding sites on GAM325 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM325 target RNA, herein designated GAM TARGET RNA, into GAM325 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4666] It is appreciated that GAM325 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM325 target genes. The mRNA of each one of this plurality of GAM325 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM325 RNA, herein designated GAM RNA, and which when bound by GAM325 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM325 target proteins.

[4667] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM325 gene, herein designated GAM GENE, on one or more GAM325 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[4668] It is yet further appreciated that specific functions, and accordingly utilities, of GAM325 correlate with, and may be deduced from, the identity of the target genes which GAM325 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4669] Nucleotide sequences of the GAM325 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM325 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM325 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM325 are further described hereinbelow with reference to Table Table1.

[4670] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM325 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4671]

[4672] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

326 (GAM326) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4673] GAM326 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM326 was detected is described hereinabove with reference to Figs. 2-8.

[4674] GAM326 gene, herein designated GAM GENE, and GAM326 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4675] GAM326 gene, herein designated GAM GENE, encodes a GAM326 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM326 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM326 precursor RNA is designated SEQ ID:4974, and is provided hereinbelow with reference to the sequence listing part.

[4676] GAM326 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM326 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4677] An enzyme complex designated DICER COMPLEX, dices the GAM326 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM326 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 48%) nucleotide sequence of GAM326 RNA is designated SEQ ID:4975, and is provided hereinbelow with reference to the sequence listing part.

[4678] GAM326 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM326 target RNA, herein designated GAM TARGET RNA. GAM326 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4679] GAM326 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM326 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM326 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM326 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM326 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4680] The complementary binding of GAM326 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM326 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM326 target RNA, herein designated GAM TARGET RNA, into GAM326 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4681] It is appreciated that GAM326 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM326 target genes. The mRNA of each one of this plurality of GAM326 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM326 RNA, herein designated GAM RNA, and which when bound by GAM326 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM326 target proteins.

[4682] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM326 gene, herein designated GAM GENE, on one or more GAM326 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4683] It is yet further appreciated that specific functions, and accordingly utilities, of GAM326 correlate with, and may be deduced from, the identity of the target genes which GAM326 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4684] Nucleotide sequences of the GAM326 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM326 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM326 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM326 are further described hereinbelow with reference to Table Table1.

[4685] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM326 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4686]

[4687] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 327 (GAM327) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4688] GAM327 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM327 was detected is described hereinabove with reference to Figs. 2-8.

[4689] GAM327 gene, herein designated GAM GENE, and GAM327 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4690] GAM327 gene, herein designated GAM GENE, encodes a GAM327 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM327 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM327 precursor RNA is designated SEQ ID:4979, and is provided hereinbelow with reference to the sequence listing part.

[4691] GAM327 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM327 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4692] An enzyme complex designated DICER COMPLEX, dices the GAM327 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM327 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 51%) nucleotide sequence of GAM327 RNA is designated SEQ ID:4980, and is provided hereinbelow with reference to the sequence listing part.

[4693] GAM327 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM327 target RNA, herein designated GAM TARGET RNA. GAM327 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4694] GAM327 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM327 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM327 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM327 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM327 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4695] The complementary binding of GAM327 RNA, herein designated GAM RNA, to target binding sites on GAM327 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM327 target RNA, herein designated GAM TARGET RNA, into GAM327 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4696] It is appreciated that GAM327 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM327 target genes. The mRNA of each one of this plurality of GAM327 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM327 RNA, herein designated GAM RNA, and which when bound by

GAM327 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM327 target proteins.

[4697] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM327 gene, herein designated GAM GENE, on one or more GAM327 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4698] It is yet further appreciated that specific functions, and accordingly utilities, of GAM327 correlate with, and may be deduced from, the identity of the target genes which GAM327 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[4699] Nucleotide sequences of the GAM327 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM327 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM327 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM327 are further described hereinbelow with reference to Table Table1.

[4700] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM327 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4701]

[4702] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 328 (GAM328) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4703] GAM328 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM328 was detected is described hereinabove with reference to Figs. 2-8.

[4704] GAM328 gene, herein designated GAM GENE, and GAM328 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4705] GAM328 gene, herein designated GAM GENE, encodes a GAM328 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM328 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM328 precursor RNA is designated SEQ ID:4987, and is provided hereinbelow with reference to the sequence listing part.

[4706] GAM328 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM328 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[4707] An enzyme complex designated DICER COMPLEX, dices the GAM328 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM328 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM328 RNA is designated SEQ ID:4988, and is provided hereinbelow with reference to the sequence listing part.

[4708] GAM328 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM328 target RNA, herein designated GAM TARGET RNA. GAM328 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4709] GAM328 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM328 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM328 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM328 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM328 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4710] The complementary binding of GAM328 RNA, herein designated GAM RNA, to target binding sites on GAM328 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM328 target RNA, herein designated GAM TARGET RNA, into GAM328 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4711] It is appreciated that GAM328 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM328 target genes. The mRNA of each one of this plurality of GAM328 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM328 RNA, herein designated GAM RNA, and which when bound by GAM328 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM328 target proteins.

[4712] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM328 gene, herein designated GAM GENE, on one or more GAM328 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4713] It is yet further appreciated that specific functions, and accordingly utilities, of GAM328 correlate with, and may be deduced from, the identity of the target genes which GAM328 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4714] Nucleotide sequences of the GAM328 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM328 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM328 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM328 are further described hereinbelow with reference to Table Table1.

[4715] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM328 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4716]

[4717] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 329 (GAM329) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4718] GAM329 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM329 was detected is described hereinabove with reference to Figs. 2-8.

[4719] GAM329 gene, herein designated GAM GENE, and GAM329 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4720] GAM329 gene, herein designated GAM GENE, encodes a GAM329 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM329 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM329 precursor RNA is designated SEQ ID:5001, and is provided hereinbelow with reference to the sequence listing part.

[4721] GAM329 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM329 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4722] An enzyme complex designated DICER COMPLEX, dices the GAM329 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM329 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM329 RNA is designated SEQ ID:5002, and is provided hereinbelow with reference to the sequence listing part.

[4723] GAM329 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM329

target RNA, herein designated GAM TARGET RNA.

GAM329 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4724] GAM329 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM329 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM329 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM329 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM329 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4725] The complementary binding of GAM329 RNA, herein designated GAM RNA, to target binding sites on GAM329 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM329 target RNA, herein designated GAM TARGET RNA, into GAM329 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4726] It is appreciated that GAM329 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM329 target genes. The mRNA of each one of this plurality of GAM329 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM329 RNA, herein designated GAM RNA, and which when bound by GAM329 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM329 target proteins.

[4727] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM329 gene, herein designated GAM GENE, on one or more GAM329 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4728] It is yet further appreciated that specific functions, and accordingly utilities, of GAM329 correlate with, and may be deduced from, the identity of the target genes which GAM329 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4729] Nucleotide sequences of the GAM329 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM329 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM329 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM329 are further described hereinbelow with reference to Table Table1.

[4730] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM329 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4731]

[4732] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 330 (GAM330) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4733] GAM330 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM330 was detected is described hereinabove with reference to Figs. 2-8.

[4734] GAM330 gene, herein designated GAM GENE, and GAM330 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4735] GAM330 gene, herein designated GAM GENE, encodes a GAM330 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM330 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM330 precursor RNA is designated SEQ ID:5006, and is provided hereinbelow with reference to the sequence listing part.

[4736] GAM330 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM330 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4737] An enzyme complex designated DICER COMPLEX, dices the GAM330 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM330 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM330 RNA is designated SEQ ID:5007, and is provided hereinbelow with reference to the sequence listing part.

[4738] GAM330 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM330 target RNA, herein designated GAM TARGET RNA. GAM330 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4739] GAM330 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM330 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM330 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM330 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM330 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4740] The complementary binding of GAM330 RNA, herein designated GAM RNA, to target binding sites on GAM330 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM330 target RNA, herein designated GAM TARGET RNA, into GAM330 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4741] It is appreciated that GAM330 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM330 target genes. The mRNA of each one of this plu-

rality of GAM330 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM330 RNA, herein designated GAM RNA, and which when bound by GAM330 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM330 target proteins.

[4742] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM330 gene, herein designated GAM GENE, on one or more GAM330 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4743] It is yet further appreciated that specific functions, and accordingly utilities, of GAM330 correlate with, and may be deduced from, the identity of the target genes which GAM330 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4744] Nucleotide sequences of the GAM330 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM330 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM330 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM330 are further described hereinbelow with reference to Table Table1.

[4745] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM330 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4746]

[4747] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 331 (GAM331) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[4748] GAM331 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM331 was detected is described hereinabove with reference to Figs. 2-8.

[4749] GAM331 gene, herein designated GAM GENE, and GAM331 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4750] GAM331 gene, herein designated GAM GENE, encodes a GAM331 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM331 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM331 precursor RNA is designated SEQ ID:5078, and is provided hereinbelow with reference to the sequence listing part.

[4751] GAM331 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM331 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4752] An enzyme complex designated DICER COMPLEX, dices the GAM331 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM331 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 50%) nucleotide sequence of GAM331 RNA is designated SEQ ID:5079, and is provided hereinbelow with reference to the sequence listing part.

[4753] GAM331 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM331 target RNA, herein designated GAM TARGET RNA.
GAM331 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[4754] GAM331 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM331 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM331 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM331 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM331 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4755] The complementary binding of GAM331 RNA, herein designated GAM RNA, to target binding sites on GAM331 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM331 target RNA, herein designated GAM TARGET RNA, into GAM331 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4756] It is appreciated that GAM331 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM331 target genes. The mRNA of each one of this plurality of GAM331 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM331 RNA, herein designated GAM RNA, and which when bound by GAM331 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM331 target proteins.

[4757] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM331 gene, herein designated GAM GENE, on one or more GAM331 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4758] It is yet further appreciated that specific functions, and accordingly utilities, of GAM331 correlate with, and may be deduced from, the identity of the target genes which GAM331 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4759] Nucleotide sequences of the GAM331 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM331 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM331 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM331 are further described hereinbelow with reference to Table Table1.

[4760] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM331 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4761]

[4762] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 332 (GAM332) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4763] GAM332 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM332 was detected is described hereinabove with reference to Figs. 2-8.

[4764] GAM332 gene, herein designated GAM GENE, and GAM332 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4765] GAM332 gene, herein designated GAM GENE, encodes a GAM332 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM332 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM332 precursor RNA is designated SEQ ID:5105, and is provided hereinbelow with reference to the sequence listing part.

[4766] GAM332 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM332 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4767] An enzyme complex designated DICER COMPLEX, dices the GAM332 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM332 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of

GAM332 RNA is designated SEQ ID:5106, and is provided hereinbelow with reference to the sequence listing part.

[4768] GAM332 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM332 target RNA, herein designated GAM TARGET RNA. GAM332 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4769] GAM332 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM332 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM332 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM332 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM332 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4770] The complementary binding of GAM332 RNA, herein designated GAM RNA, to target binding sites on GAM332 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM332 target RNA, herein designated GAM TARGET RNA, into GAM332 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4771] It is appreciated that GAM332 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM332 target genes. The mRNA of each one of this plurality of GAM332 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM332 RNA, herein designated GAM RNA, and which when bound by GAM332 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM332 target proteins.

[4772] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM332 gene, herein designated GAM GENE, on one or more GAM332 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4773] It is yet further appreciated that specific functions, and accordingly utilities, of GAM332 correlate with, and may be deduced from, the identity of the target genes which GAM332 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4774] Nucleotide sequences of the GAM332 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM332 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM332 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM332 are further described hereinbelow with reference to Table Table1.

[4775] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM332 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4776]

[4777] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 333 (GAM333) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4778] GAM333 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM333 was detected is described hereinabove

with reference to Figs. 2-8.

[4779] GAM333 gene, herein designated GAM GENE, and GAM333 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4780] GAM333 gene, herein designated GAM GENE, encodes a GAM333 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM333 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM333 precursor RNA is designated SEQ ID:5123, and is provided hereinbelow with reference to the sequence listing part.

[4781] GAM333 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM333 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4782] An enzyme complex designated DICER COMPLEX, dices the GAM333 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM333 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 26%) nucleotide sequence of GAM333 RNA is designated SEQ ID:5124, and is provided hereinbelow with reference to the sequence listing part.

[4783] GAM333 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM333 target RNA, herein designated GAM TARGET RNA. GAM333 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4784] GAM333 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM333 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM333 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM333 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM333 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4785] The complementary binding of GAM333 RNA, herein designated GAM RNA, to target binding sites on GAM333 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM333 target RNA, herein designated GAM TARGET RNA, into GAM333 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[4786] It is appreciated that GAM333 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM333 target genes. The mRNA of each one of this plurality of GAM333 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM333 RNA, herein designated GAM RNA, and which when bound by GAM333 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM333 target proteins.

[4787] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM333 gene, herein designated GAM GENE, on one or more GAM333 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4788] It is yet further appreciated that specific functions, and accordingly utilities, of GAM333 correlate with, and may be deduced from, the identity of the target genes which GAM333 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4789] Nucleotide sequences of the GAM333 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM333 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM333 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM333 are further described hereinbelow with reference to Table Table1.

[4790] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM333 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4791]

[4792] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 334 (GAM334) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4793] GAM334 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM334 was detected is described hereinabove with reference to Figs. 2-8.

[4794] GAM334 gene, herein designated GAM GENE, and GAM334 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4795] GAM334 gene, herein designated GAM GENE, encodes a GAM334 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM334 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM334 precursor RNA is designated SEQ ID:5135, and is provided hereinbelow with reference to the sequence listing part.

[4796] GAM334 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM334 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4797] An enzyme complex designated DICER COMPLEX, dices the GAM334 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM334 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 38%) nucleotide sequence of GAM334 RNA is designated SEQ ID:5136, and is provided hereinbelow with reference to the sequence listing part.

[4798] GAM334 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM334 target RNA, herein designated GAM TARGET RNA.

GAM334 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4799] GAM334 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM334 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM334 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM334 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM334 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4800] The complementary binding of GAM334 RNA, herein designated GAM RNA, to target binding sites on GAM334 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM334 target RNA, herein designated GAM TARGET RNA, into GAM334 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4801] It is appreciated that GAM334 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM334 target genes. The mRNA of each one of this plurality of GAM334 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM334 RNA, herein designated GAM RNA, and which when bound by GAM334 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM334 target proteins.

[4802] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM334 gene, herein designated GAM GENE, on one or more GAM334 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4803] It is yet further appreciated that specific functions, and accordingly utilities, of GAM334 correlate with, and may be deduced from, the identity of the target genes which GAM334 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4804] Nucleotide sequences of the GAM334 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM334 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM334 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM334 are further described hereinbelow with reference to Table Table1.

[4805] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM334 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4806]

[4807] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 335 (GAM335) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4808] GAM335 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM335 was detected is described hereinabove with reference to Figs. 2-8.

[4809] GAM335 gene, herein designated GAM GENE, and GAM335 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4810] GAM335 gene, herein designated GAM GENE, encodes a

GAM335 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM335 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM335 precursor RNA is designated SEQ ID:5145, and is provided hereinbelow with reference to the sequence listing part.

[4811] GAM335 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM335 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4812] An enzyme complex designated DICER COMPLEX, dices the GAM335 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM335 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM335 RNA is designated SEQ ID:5146, and is provided hereinbelow with reference to the sequence listing part.

[4813] GAM335 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM335 target RNA, herein designated GAM TARGET RNA. GAM335 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4814] GAM335 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM335 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM335 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM335 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM335 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4815] The complementary binding of GAM335 RNA, herein designated GAM RNA, to target binding sites on GAM335 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM335 target RNA, herein designated GAM TARGET RNA, into GAM335 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4816] It is appreciated that GAM335 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM335 target genes. The mRNA of each one of this plurality of GAM335 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM335 RNA, herein designated GAM RNA, and which when bound by GAM335 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM335 target proteins.

[4817] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM335 gene, herein designated GAM GENE, on one or more GAM335 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4818] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM335 correlate with, and may be deduced from, the identity of the target genes which GAM335 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4819] Nucleotide sequences of the GAM335 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM335 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM335 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM335 are further described hereinbelow with reference to Table Table1.

[4820] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM335 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4821]

[4822] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 336 (GAM336) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[4823] GAM336 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM336 was detected is described hereinabove with reference to Figs. 2-8.

[4824] GAM336 gene, herein designated GAM GENE, and GAM336 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4825] GAM336 gene, herein designated GAM GENE, encodes a GAM336 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM336 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM336 precursor RNA is designated SEQ ID:5161, and is provided hereinbelow with reference to the sequence listing part.

[4826] GAM336 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM336 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4827] An enzyme complex designated DICER COMPLEX, dices the GAM336 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM336 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 73%) nucleotide sequence of GAM336 RNA is designated SEQ ID:5162, and is provided hereinbelow with reference to the sequence listing part.

[4828] GAM336 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM336 target RNA, herein designated GAM TARGET RNA.
GAM336 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4829] GAM336 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM336 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM336 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM336 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM336 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4830] The complementary binding of GAM336 RNA, herein designated GAM RNA, to target binding sites on GAM336 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM336 target RNA, herein designated GAM TARGET RNA, into GAM336 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4831] It is appreciated that GAM336 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM336 target genes. The mRNA of each one of this plurality of GAM336 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM336 RNA, herein designated GAM RNA, and which when bound by GAM336 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM336 target proteins.

[4832] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM336 gene, herein designated GAM GENE, on one or more GAM336 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4833] It is yet further appreciated that specific functions, and accordingly utilities, of GAM336 correlate with, and may be deduced from, the identity of the target genes which GAM336 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4834] Nucleotide sequences of the GAM336 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM336 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM336 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM336 are further described hereinbelow with reference to Table Table1.

[4835] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM336 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4836]

[4837] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 337 (GAM337) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4838] GAM337 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM337 was detected is described hereinabove with reference to Figs. 2-8.

[4839] GAM337 gene, herein designated GAM GENE, and GAM337 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4840] GAM337 gene, herein designated GAM GENE, encodes a GAM337 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM337 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM337 precursor RNA is designated SEQ ID:5176, and is provided hereinbelow with reference to the sequence listing part.

[4841] GAM337 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM337 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4842] An enzyme complex designated DICER COMPLEX, dices the GAM337 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM337 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM337 RNA is designated SEQ ID:5177, and is provided

hereinbelow with reference to the sequence listing part.

[4843] GAM337 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM337 target RNA, herein designated GAM TARGET RNA.
GAM337 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4844] GAM337 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM337 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM337 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM337 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM337 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4845] The complementary binding of GAM337 RNA, herein designated GAM RNA, to target binding sites on GAM337 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM337 target RNA, herein designated GAM TARGET RNA, into GAM337 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4846] It is appreciated that GAM337 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM337 target genes. The mRNA of each one of this plurality of GAM337 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM337 RNA, herein designated GAM RNA, and which when bound by GAM337 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM337 tar-

get proteins.

[4847] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM337 gene, herein designated GAM GENE, on one or more GAM337 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4848] It is yet further appreciated that specific functions, and accordingly utilities, of GAM337 correlate with, and may be deduced from, the identity of the target genes which GAM337 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4849] Nucleotide sequences of the GAM337 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM337 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM337 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM337 are further described hereinbelow with reference to Table Table1.

[4850] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM337 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4851]

[4852] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 338 (GAM338) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4853] GAM338 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM338 was detected is described hereinabove with reference to Figs. 2-8.

[4854] GAM338 gene, herein designated GAM GENE, and GAM338 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4855] GAM338 gene, herein designated GAM GENE, encodes a GAM338 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM338 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM338 precursor RNA is designated SEQ ID:5197, and is provided hereinbelow with reference to the sequence listing part.

[4856] GAM338 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM338 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4857] An enzyme complex designated DICER COMPLEX, dices

the GAM338 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM338 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM338 RNA is designated SEQ ID:5198, and is provided hereinbelow with reference to the sequence listing part.

[4858] GAM338 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM338 target RNA, herein designated GAM TARGET RNA. GAM338 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4859] GAM338 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM338 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM338

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM338 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM338 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4860] The complementary binding of GAM338 RNA, herein designated GAM RNA, to target binding sites on GAM338 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM338 target RNA, herein designated GAM TARGET RNA, into GAM338 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4861] It is appreciated that GAM338 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM338 target genes. The mRNA of each one of this plurality of GAM338 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM338 RNA, herein designated GAM RNA, and which when bound by GAM338 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM338 target proteins.

[4862] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM338 gene, herein designated GAM GENE, on one or more GAM338 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4863] It is yet further appreciated that specific functions, and accordingly utilities, of GAM338 correlate with, and may be deduced from, the identity of the target genes which GAM338 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4864] Nucleotide sequences of the GAM338 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM338 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM338 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM338 are further described hereinbelow with reference to Table Table1.

[4865] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM338 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4866]

[4867] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 339 (GAM339) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4868] GAM339 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM339 was detected is described hereinabove with reference to Figs. 2-8.

[4869] GAM339 gene, herein designated GAM GENE, and GAM339 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4870] GAM339 gene, herein designated GAM GENE, encodes a GAM339 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM339 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM339 precursor RNA is designated SEQ ID:5213, and is provided hereinbelow with reference to the sequence listing part.

[4871] GAM339 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM339 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4872] An enzyme complex designated DICER COMPLEX, dices the GAM339 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM339 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM339 RNA is designated SEQ ID:5214, and is provided hereinbelow with reference to the sequence listing part.

[4873] GAM339 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM339 target RNA, herein designated GAM TARGET RNA. GAM339 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4874] GAM339 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM339 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM339 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM339 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM339 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[4875] The complementary binding of GAM339 RNA, herein designated GAM RNA, to target binding sites on GAM339 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM339 target RNA, herein designated GAM TARGET RNA, into GAM339 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4876] It is appreciated that GAM339 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM339 target genes. The mRNA of each one of this plurality of GAM339 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM339 RNA, herein designated GAM RNA, and which when bound by GAM339 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM339 target proteins.

[4877] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM339 gene, herein designated GAM GENE, on one or

more GAM339 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4878] It is yet further appreciated that specific functions, and accordingly utilities, of GAM339 correlate with, and may be deduced from, the identity of the target genes which GAM339 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4879] Nucleotide sequences of the GAM339 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM339 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM339 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM339 are further de-

scribed hereinbelow with reference to Table Table1.

[4880] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM339 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4881]

[4882] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 340 (GAM340) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4883] GAM340 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM340 was detected is described hereinabove with reference to Figs. 2-8.

[4884] GAM340 gene, herein designated GAM GENE, and GAM340 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4885] GAM340 gene, herein designated GAM GENE, encodes a GAM340 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM340 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM340 precursor RNA is designated SEQ ID:5220, and is provided hereinbelow with reference to the sequence listing part.

[4886] GAM340 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM340 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4887] An enzyme complex designated DICER COMPLEX, dices the GAM340 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM340 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM340 RNA is designated SEQ ID:5221, and is provided hereinbelow with reference to the sequence listing part.

[4888] GAM340 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM340 target RNA, herein designated GAM TARGET RNA. GAM340 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4889] GAM340 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM340 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM340 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM340 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM340 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4890] The complementary binding of GAM340 RNA, herein designated GAM RNA, to target binding sites on GAM340 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM340 target RNA, herein designated GAM TARGET RNA, into GAM340 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4891] It is appreciated that GAM340 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM340 target genes. The mRNA of each one of this plurality of GAM340 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM340 RNA, herein designated GAM RNA, and which when bound by GAM340 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM340 target proteins.

[4892] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM340 gene, herein designated GAM GENE, on one or more GAM340 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4893] It is yet further appreciated that specific functions, and accordingly utilities, of GAM340 correlate with, and may

be deduced from, the identity of the target genes which GAM340 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4894] Nucleotide sequences of the GAM340 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM340 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM340 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM340 are further described hereinbelow with reference to Table Table1.

[4895] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM340 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4896]

[4897] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 341 (GAM341) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4898] GAM341 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM341 was detected is described hereinabove with reference to Figs. 2-8.

[4899] GAM341 gene, herein designated GAM GENE, and GAM341 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4900] GAM341 gene, herein designated GAM GENE, encodes a GAM341 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM341 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM341 precursor RNA is designated SEQ ID:5228, and is provided hereinbelow with reference to the sequence listing part.

[4901] GAM341 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM341 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4902] An enzyme complex designated DICER COMPLEX, dices the GAM341 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM341 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM341 RNA is designated SEQ ID:5229, and is provided hereinbelow with reference to the sequence listing part.

[4903] GAM341 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM341 target RNA, herein designated GAM TARGET RNA. GAM341 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4904] GAM341 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM341 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM341 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM341 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM341 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4905] The complementary binding of GAM341 RNA, herein designated GAM RNA, to target binding sites on GAM341 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM341 target RNA, herein designated GAM TARGET RNA, into GAM341 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4906] It is appreciated that GAM341 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM341 target genes. The mRNA of each one of this plurality of GAM341 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM341 RNA, herein designated GAM RNA, and which when bound by GAM341 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM341 target proteins.

[4907] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM341 gene, herein designated GAM GENE, on one or more GAM341 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4908] It is yet further appreciated that specific functions, and accordingly utilities, of GAM341 correlate with, and may be deduced from, the identity of the target genes which GAM341 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4909] Nucleotide sequences of the GAM341 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM341 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM341 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM341 are further described hereinbelow with reference to Table Table1.

[4910] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM341 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4911]

[4912] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 342 (GAM342) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4913] GAM342 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM342 was detected is described hereinabove with reference to Figs. 2-8.

[4914] GAM342 gene, herein designated GAM GENE, and GAM342 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4915] GAM342 gene, herein designated GAM GENE, encodes a GAM342 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM342 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM342 precursor RNA is desig-

nated SEQ ID:5248, and is provided hereinbelow with reference to the sequence listing part.

[4916] GAM342 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM342 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4917] An enzyme complex designated DICER COMPLEX, dices the GAM342 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM342 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM342 RNA is designated SEQ ID:5249, and is provided hereinbelow with reference to the sequence listing part.

[4918] GAM342 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM342 target RNA, herein designated GAM TARGET RNA. GAM342 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4919] GAM342 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM342 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM342 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM342 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM342

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4920] The complementary binding of GAM342 RNA, herein designated GAM RNA, to target binding sites on GAM342 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM342 target RNA, herein designated GAM TARGET RNA, into GAM342 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4921] It is appreciated that GAM342 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM342 target genes. The mRNA of each one of this plurality of GAM342 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM342 RNA, herein designated GAM RNA, and which when bound by GAM342 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM342 target proteins.

[4922] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM342 gene, herein designated GAM GENE, on one or more GAM342 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4923] It is yet further appreciated that specific functions, and accordingly utilities, of GAM342 correlate with, and may be deduced from, the identity of the target genes which GAM342 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4924] Nucleotide sequences of the GAM342 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM342 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM342 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM342 are further described hereinbelow with reference to Table Table1.

[4925] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM342 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4926]

[4927] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 343 (GAM343) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4928] GAM343 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM343 was detected is described hereinabove with reference to Figs. 2-8.

[4929] GAM343 gene, herein designated GAM GENE, and

GAM343 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4930] GAM343 gene, herein designated GAM GENE, encodes a GAM343 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM343 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM343 precursor RNA is designated SEQ ID:5284, and is provided hereinbelow with reference to the sequence listing part.

[4931] GAM343 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM343 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4932] An enzyme complex designated DICER COMPLEX, dices the GAM343 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM343 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM343 RNA is designated SEQ ID:5285, and is provided hereinbelow with reference to the sequence listing part.

[4933] GAM343 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM343 target RNA, herein designated GAM TARGET RNA. GAM343 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4934] GAM343 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM343 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM343 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM343 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM343 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4935] The complementary binding of GAM343 RNA, herein designated GAM RNA, to target binding sites on GAM343 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM343 target RNA, herein designated GAM TARGET RNA, into GAM343 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4936] It is appreciated that GAM343 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM343 target genes. The mRNA of each one of this plurality of GAM343 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM343 RNA, herein designated GAM RNA, and which when bound by GAM343 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM343 target proteins.

[4937] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM343 gene, herein designated GAM GENE, on one or more GAM343 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4938] It is yet further appreciated that specific functions, and accordingly utilities, of GAM343 correlate with, and may be deduced from, the identity of the target genes which GAM343 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4939] Nucleotide sequences of the GAM343 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM343 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM343 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM343 are further described hereinbelow with reference to Table Table1.

[4940] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM343 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4941]

[4942] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 344 (GAM344) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4943] GAM344 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM344 was detected is described hereinabove with reference to Figs. 2-8.

[4944] GAM344 gene, herein designated GAM GENE, and GAM344 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4945] GAM344 gene, herein designated GAM GENE, encodes a GAM344 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM344 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM344 precursor RNA is designated SEQ ID:5296, and is provided hereinbelow with reference to the sequence listing part.

[4946] GAM344 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM344 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4947] An enzyme complex designated DICER COMPLEX, dices the GAM344 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM344 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM344 RNA is designated SEQ ID:5297, and is provided hereinbelow with reference to the sequence listing part.

[4948] GAM344 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM344 target RNA, herein designated GAM TARGET RNA. GAM344 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4949] GAM344 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM344 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM344 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM344 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM344 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4950] The complementary binding of GAM344 RNA, herein designated GAM RNA, to target binding sites on GAM344 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM344 target RNA, herein designated GAM TARGET RNA, into GAM344 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4951] It is appreciated that GAM344 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM344 target genes. The mRNA of each one of this plurality of GAM344 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM344 RNA, herein designated GAM RNA, and which when bound by GAM344 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM344 target proteins.

[4952] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM344 gene, herein designated GAM GENE, on one or more GAM344 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4953] It is yet further appreciated that specific functions, and accordingly utilities, of GAM344 correlate with, and may be deduced from, the identity of the target genes which GAM344 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4954] Nucleotide sequences of the GAM344 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM344 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM344 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM344 are further described hereinbelow with reference to Table Table1.

[4955] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM344 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4956]

[4957] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 345 (GAM345) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4958] GAM345 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM345 was detected is described hereinabove with reference to Figs. 2-8.

[4959] GAM345 gene, herein designated GAM GENE, and GAM345 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4960] GAM345 gene, herein designated GAM GENE, encodes a GAM345 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM345 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM345 precursor RNA is designated SEQ ID:5306, and is provided hereinbelow with reference to the sequence listing part.

[4961] GAM345 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM345 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4962] An enzyme complex designated DICER COMPLEX, dices the GAM345 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM345 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM345 RNA is designated SEQ ID:5307, and is provided hereinbelow with reference to the sequence listing part.

[4963] GAM345 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM345 target RNA, herein designated GAM TARGET RNA. GAM345 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4964] GAM345 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM345 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM345 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM345 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM345 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4965] The complementary binding of GAM345 RNA, herein designated GAM RNA, to target binding sites on GAM345 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM345 target RNA, herein designated GAM TARGET RNA, into GAM345 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4966] It is appreciated that GAM345 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM345 target genes. The mRNA of each one of this plurality of GAM345 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM345 RNA,

herein designated GAM RNA, and which when bound by GAM345 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM345 target proteins.

[4967] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM345 gene, herein designated GAM GENE, on one or more GAM345 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4968] It is yet further appreciated that specific functions, and accordingly utilities, of GAM345 correlate with, and may be deduced from, the identity of the target genes which

GAM345 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4969] Nucleotide sequences of the GAM345 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM345 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM345 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM345 are further described hereinbelow with reference to Table Table1.

[4970] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM345 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[4971]

[4972] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 346 (GAM346) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4973] GAM346 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM346 was detected is described hereinabove with reference to Figs. 2-8.

[4974] GAM346 gene, herein designated GAM GENE, and GAM346 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4975] GAM346 gene, herein designated GAM GENE, encodes a GAM346 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM346 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM346 precursor RNA is designated SEQ ID:5337, and is provided hereinbelow with reference to the sequence listing part.

[4976] GAM346 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM346 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[4977] An enzyme complex designated DICER COMPLEX, dices the GAM346 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM346 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM346 RNA is designated SEQ ID:5338, and is provided hereinbelow with reference to the sequence listing part.

[4978] GAM346 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM346 target RNA, herein designated GAM TARGET RNA.
GAM346 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4979] GAM346 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM346 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM346 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM346 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM346 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4980] The complementary binding of GAM346 RNA, herein designated GAM RNA, to target binding sites on GAM346 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM346 target RNA, herein desig-

nated GAM TARGET RNA, into GAM346 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4981] It is appreciated that GAM346 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM346 target genes. The mRNA of each one of this plurality of GAM346 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM346 RNA, herein designated GAM RNA, and which when bound by GAM346 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM346 target proteins.

[4982] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM346 gene, herein designated GAM GENE, on one or more GAM346 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4983] It is yet further appreciated that specific functions, and accordingly utilities, of GAM346 correlate with, and may be deduced from, the identity of the target genes which GAM346 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4984] Nucleotide sequences of the GAM346 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM346 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM346 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM346 are further described hereinbelow with reference to Table Table1.

[4985] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM346 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[4986]

[4987] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 347 (GAM347) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[4988] GAM347 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM347 was detected is described hereinabove with reference to Figs. 2-8.

[4989] GAM347 gene, herein designated GAM GENE, and GAM347 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[4990] GAM347 gene, herein designated GAM GENE, encodes a GAM347 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM347 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM347 precursor RNA is designated SEQ ID:5348, and is provided hereinbelow with ref-

erence to the sequence listing part.

[4991] GAM347 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM347 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[4992] An enzyme complex designated DICER COMPLEX, dices the GAM347 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM347 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM347 RNA is designated SEQ ID:5349, and is provided hereinbelow with reference to the sequence listing part.

[4993] GAM347 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM347 target RNA, herein designated GAM TARGET RNA.

GAM347 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[4994] GAM347 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM347 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM347 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM347 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM347 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[4995] The complementary binding of GAM347 RNA, herein designated GAM RNA, to target binding sites on GAM347 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM347 target RNA, herein designated GAM TARGET RNA, into GAM347 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[4996] It is appreciated that GAM347 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM347 target genes. The mRNA of each one of this plurality of GAM347 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM347 RNA, herein designated GAM RNA, and which when bound by GAM347 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM347 target proteins.

[4997] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM347 gene, herein designated GAM GENE, on one or more GAM347 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[4998] It is yet further appreciated that specific functions, and accordingly utilities, of GAM347 correlate with, and may be deduced from, the identity of the target genes which GAM347 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[4999] Nucleotide sequences of the GAM347 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM347 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM347 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM347 are further described hereinbelow with reference to Table Table1.

[5000] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM347 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5001]

[5002] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 348 (GAM348) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5003] GAM348 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM348 was detected is described hereinabove with reference to Figs. 2-8.

[5004] GAM348 gene, herein designated GAM GENE, and GAM348 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[5005] GAM348 gene, herein designated GAM GENE, encodes a GAM348 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM348 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM348 precursor RNA is designated SEQ ID:5360, and is provided hereinbelow with reference to the sequence listing part.

[5006] GAM348 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM348 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5007] An enzyme complex designated DICER COMPLEX, dices the GAM348 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM348 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM348 RNA is designated SEQ ID:5361, and is provided hereinbelow with reference to the sequence listing part.

[5008] GAM348 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM348 target RNA, herein designated GAM TARGET RNA. GAM348 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5009] GAM348 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM348 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM348 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM348 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM348 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5010] The complementary binding of GAM348 RNA, herein designated GAM RNA, to target binding sites on GAM348 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM348 target RNA, herein designated GAM TARGET RNA, into GAM348 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5011] It is appreciated that GAM348 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM348 target genes. The mRNA of each one of this plurality of GAM348 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM348 RNA, herein designated GAM RNA, and which when bound by GAM348 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM348 target proteins.

[5012] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM348 gene, herein designated GAM GENE, on one or more GAM348 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[5013] It is yet further appreciated that specific functions, and accordingly utilities, of GAM348 correlate with, and may be deduced from, the identity of the target genes which GAM348 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5014] Nucleotide sequences of the GAM348 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM348 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM348 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM348 are further described hereinbelow with reference to Table Table1.

[5015] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM348 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5016]

[5017] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

349 (GAM349) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5018] GAM349 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM349 was detected is described hereinabove with reference to Figs. 2-8.

[5019] GAM349 gene, herein designated GAM GENE, and GAM349 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5020] GAM349 gene, herein designated GAM GENE, encodes a GAM349 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM349 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM349 precursor RNA is designated SEQ ID:5370, and is provided hereinbelow with reference to the sequence listing part.

[5021] GAM349 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM349 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5022] An enzyme complex designated DICER COMPLEX, dices the GAM349 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM349 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 86%) nucleotide sequence of GAM349 RNA is designated SEQ ID:5371, and is provided hereinbelow with reference to the sequence listing part.

[5023] GAM349 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM349 target RNA, herein designated GAM TARGET RNA. GAM349 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5024] GAM349 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM349 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM349 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM349 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM349 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5025] The complementary binding of GAM349 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM349 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM349 target RNA, herein designated GAM TARGET RNA, into GAM349 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5026] It is appreciated that GAM349 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM349 target genes. The mRNA of each one of this plurality of GAM349 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM349 RNA, herein designated GAM RNA, and which when bound by GAM349 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM349 target proteins.

[5027] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM349 gene, herein designated GAM GENE, on one or more GAM349 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5028] It is yet further appreciated that specific functions, and accordingly utilities, of GAM349 correlate with, and may be deduced from, the identity of the target genes which GAM349 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5029] Nucleotide sequences of the GAM349 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM349 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM349 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM349 are further described hereinbelow with reference to Table Table1.

[5030] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM349 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5031]

[5032] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 350 (GAM350) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5033] GAM350 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM350 was detected is described hereinabove with reference to Figs. 2-8.

[5034] GAM350 gene, herein designated GAM GENE, and GAM350 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5035] GAM350 gene, herein designated GAM GENE, encodes a GAM350 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM350 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM350 precursor RNA is designated SEQ ID:5379, and is provided hereinbelow with reference to the sequence listing part.

[5036] GAM350 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM350 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5037] An enzyme complex designated DICER COMPLEX, dices the GAM350 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM350 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 66%) nucleotide sequence of GAM350 RNA is designated SEQ ID:5380, and is provided hereinbelow with reference to the sequence listing part.

[5038] GAM350 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM350 target RNA, herein designated GAM TARGET RNA. GAM350 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5039] GAM350 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM350 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM350 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM350 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM350 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5040] The complementary binding of GAM350 RNA, herein designated GAM RNA, to target binding sites on GAM350 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM350 target RNA, herein designated GAM TARGET RNA, into GAM350 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5041] It is appreciated that GAM350 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM350 target genes. The mRNA of each one of this plurality of GAM350 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM350 RNA, herein designated GAM RNA, and which when bound by

GAM350 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM350 target proteins.

[5042] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM350 gene, herein designated GAM GENE, on one or more GAM350 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5043] It is yet further appreciated that specific functions, and accordingly utilities, of GAM350 correlate with, and may be deduced from, the identity of the target genes which GAM350 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[5044] Nucleotide sequences of the GAM350 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM350 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM350 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM350 are further described hereinbelow with reference to Table Table1.

[5045] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM350 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5046]

[5047] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 351 (GAM351) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5048] GAM351 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM351 was detected is described hereinabove with reference to Figs. 2-8.

[5049] GAM351 gene, herein designated GAM GENE, and GAM351 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5050] GAM351 gene, herein designated GAM GENE, encodes a GAM351 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM351 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM351 precursor RNA is designated SEQ ID:5384, and is provided hereinbelow with reference to the sequence listing part.

[5051] GAM351 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM351 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[5052] An enzyme complex designated DICER COMPLEX, dices the GAM351 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM351 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM351 RNA is designated SEQ ID:5385, and is provided hereinbelow with reference to the sequence listing part.

[5053] GAM351 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM351 target RNA, herein designated GAM TARGET RNA. GAM351 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5054] GAM351 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM351 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM351 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM351 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM351 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5055] The complementary binding of GAM351 RNA, herein designated GAM RNA, to target binding sites on GAM351 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM351 target RNA, herein designated GAM TARGET RNA, into GAM351 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5056] It is appreciated that GAM351 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM351 target genes. The mRNA of each one of this plurality of GAM351 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM351 RNA, herein designated GAM RNA, and which when bound by GAM351 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM351 target proteins.

[5057] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM351 gene, herein designated GAM GENE, on one or more GAM351 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5058] It is yet further appreciated that specific functions, and accordingly utilities, of GAM351 correlate with, and may be deduced from, the identity of the target genes which GAM351 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5059] Nucleotide sequences of the GAM351 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM351 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM351 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM351 are further described hereinbelow with reference to Table Table1.

[5060] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM351 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5061]

[5062] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 352 (GAM352) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5063] GAM352 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM352 was detected is described hereinabove with reference to Figs. 2-8.

[5064] GAM352 gene, herein designated GAM GENE, and GAM352 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5065] GAM352 gene, herein designated GAM GENE, encodes a GAM352 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM352 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM352 precursor RNA is designated SEQ ID:5401, and is provided hereinbelow with reference to the sequence listing part.

[5066] GAM352 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM352 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5067] An enzyme complex designated DICER COMPLEX, dices the GAM352 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM352 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM352 RNA is designated SEQ ID:5402, and is provided hereinbelow with reference to the sequence listing part.

[5068] GAM352 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM352

target RNA, herein designated GAM TARGET RNA.

GAM352 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5069] GAM352 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM352 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM352 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM352 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM352 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5070] The complementary binding of GAM352 RNA, herein designated GAM RNA, to target binding sites on GAM352 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM352 target RNA, herein designated GAM TARGET RNA, into GAM352 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5071] It is appreciated that GAM352 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM352 target genes. The mRNA of each one of this plurality of GAM352 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM352 RNA, herein designated GAM RNA, and which when bound by GAM352 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM352 target proteins.

[5072] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM352 gene, herein designated GAM GENE, on one or more GAM352 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5073] It is yet further appreciated that specific functions, and accordingly utilities, of GAM352 correlate with, and may be deduced from, the identity of the target genes which GAM352 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5074] Nucleotide sequences of the GAM352 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM352 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM352 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM352 are further described hereinbelow with reference to Table Table1.

[5075] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM352 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5076]

[5077] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 353 (GAM353) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5078] GAM353 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM353 was detected is described hereinabove with reference to Figs. 2-8.

[5079] GAM353 gene, herein designated GAM GENE, and GAM353 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5080] GAM353 gene, herein designated GAM GENE, encodes a GAM353 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM353 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM353 precursor RNA is designated SEQ ID:5418, and is provided hereinbelow with reference to the sequence listing part.

[5081] GAM353 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM353 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5082] An enzyme complex designated DICER COMPLEX, dices the GAM353 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM353 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM353 RNA is designated SEQ ID:5419, and is provided hereinbelow with reference to the sequence listing part.

[5083] GAM353 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM353 target RNA, herein designated GAM TARGET RNA. GAM353 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5084] GAM353 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM353 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM353 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM353 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM353 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5085] The complementary binding of GAM353 RNA, herein designated GAM RNA, to target binding sites on GAM353 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM353 target RNA, herein designated GAM TARGET RNA, into GAM353 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5086] It is appreciated that GAM353 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM353 target genes. The mRNA of each one of this plu-

rality of GAM353 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM353 RNA, herein designated GAM RNA, and which when bound by GAM353 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM353 target proteins.

[5087] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM353 gene, herein designated GAM GENE, on one or more GAM353 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5088] It is yet further appreciated that specific functions, and accordingly utilities, of GAM353 correlate with, and may be deduced from, the identity of the target genes which GAM353 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5089] Nucleotide sequences of the GAM353 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM353 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM353 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM353 are further described hereinbelow with reference to Table Table1.

[5090] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM353 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5091]

[5092] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 354 (GAM354) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[5093] GAM354 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM354 was detected is described hereinabove with reference to Figs. 2-8.

[5094] GAM354 gene, herein designated GAM GENE, and GAM354 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5095] GAM354 gene, herein designated GAM GENE, encodes a GAM354 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM354 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM354 precursor RNA is designated SEQ ID:5459, and is provided hereinbelow with reference to the sequence listing part.

[5096] GAM354 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM354 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5097] An enzyme complex designated DICER COMPLEX, dices the GAM354 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM354 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 85%) nucleotide sequence of GAM354 RNA is designated SEQ ID:5460, and is provided hereinbelow with reference to the sequence listing part.

[5098] GAM354 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM354 target RNA, herein designated GAM TARGET RNA.
GAM354 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[5099] GAM354 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM354 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM354 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM354 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM354 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5100] The complementary binding of GAM354 RNA, herein designated GAM RNA, to target binding sites on GAM354 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM354 target RNA, herein designated GAM TARGET RNA, into GAM354 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5101] It is appreciated that GAM354 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM354 target genes. The mRNA of each one of this plurality of GAM354 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM354 RNA, herein designated GAM RNA, and which when bound by GAM354 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM354 target proteins.

[5102] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM354 gene, herein designated GAM GENE, on one or more GAM354 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5103] It is yet further appreciated that specific functions, and accordingly utilities, of GAM354 correlate with, and may be deduced from, the identity of the target genes which GAM354 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5104] Nucleotide sequences of the GAM354 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM354 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM354 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM354 are further described hereinbelow with reference to Table Table1.

[5105] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM354 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5106]

[5107] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 355 (GAM355) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5108] GAM355 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM355 was detected is described hereinabove with reference to Figs. 2-8.

[5109] GAM355 gene, herein designated GAM GENE, and GAM355 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5110] GAM355 gene, herein designated GAM GENE, encodes a GAM355 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM355 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM355 precursor RNA is designated SEQ ID:5469, and is provided hereinbelow with reference to the sequence listing part.

[5111] GAM355 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM355 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5112] An enzyme complex designated DICER COMPLEX, dices the GAM355 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM355 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 80%) nucleotide sequence of GAM355 RNA is designated SEQ ID:5470, and is provided hereinbelow with reference to the sequence listing part.

[5113] GAM355 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM355

target RNA, herein designated GAM TARGET RNA.

GAM355 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5114] GAM355 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM355 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM355 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM355 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM355 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5115] The complementary binding of GAM355 RNA, herein designated GAM RNA, to target binding sites on GAM355 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM355 target RNA, herein designated GAM TARGET RNA, into GAM355 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5116] It is appreciated that GAM355 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM355 target genes. The mRNA of each one of this plurality of GAM355 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM355 RNA, herein designated GAM RNA, and which when bound by GAM355 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM355 target proteins.

[5117] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM355 gene, herein designated GAM GENE, on one or more GAM355 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5118] It is yet further appreciated that specific functions, and accordingly utilities, of GAM355 correlate with, and may be deduced from, the identity of the target genes which GAM355 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5119] Nucleotide sequences of the GAM355 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM355 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM355 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM355 are further described hereinbelow with reference to Table Table1.

[5120] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM355 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5121]

[5122] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 356 (GAM356) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5123] GAM356 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM356 was detected is described hereinabove with reference to Figs. 2-8.

[5124] GAM356 gene, herein designated GAM GENE, and GAM356 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5125] GAM356 gene, herein designated GAM GENE, encodes a GAM356 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM356 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM356 precursor RNA is designated SEQ ID:5507, and is provided hereinbelow with reference to the sequence listing part.

[5126] GAM356 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM356 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5127] An enzyme complex designated DICER COMPLEX, dices the GAM356 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM356 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM356 RNA is designated SEQ ID:5508, and is provided hereinbelow with reference to the sequence listing part.

[5128] GAM356 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM356 target RNA, herein designated GAM TARGET RNA. GAM356 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5129] GAM356 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM356 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM356 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM356 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM356 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5130] The complementary binding of GAM356 RNA, herein designated GAM RNA, to target binding sites on GAM356 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM356 target RNA, herein designated GAM TARGET RNA, into GAM356 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5131] It is appreciated that GAM356 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM356 target genes. The mRNA of each one of this plu-

rality of GAM356 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM356 RNA, herein designated GAM RNA, and which when bound by GAM356 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM356 target proteins.

[5132] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM356 gene, herein designated GAM GENE, on one or more GAM356 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5133] It is yet further appreciated that specific functions, and accordingly utilities, of GAM356 correlate with, and may be deduced from, the identity of the target genes which GAM356 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5134] Nucleotide sequences of the GAM356 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM356 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM356 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM356 are further described hereinbelow with reference to Table Table1.

[5135] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM356 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5136]

[5137] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 357 (GAM357) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[5138] GAM357 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM357 was detected is described hereinabove with reference to Figs. 2-8.

[5139] GAM357 gene, herein designated GAM GENE, and GAM357 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5140] GAM357 gene, herein designated GAM GENE, encodes a GAM357 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM357 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM357 precursor RNA is designated SEQ ID:5514, and is provided hereinbelow with reference to the sequence listing part.

[5141] GAM357 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM357 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5142] An enzyme complex designated DICER COMPLEX, dices the GAM357 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM357 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 25%) nucleotide sequence of GAM357 RNA is designated SEQ ID:5515, and is provided hereinbelow with reference to the sequence listing part.

[5143] GAM357 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM357 target RNA, herein designated GAM TARGET RNA. GAM357 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[5144] GAM357 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM357 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM357 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM357 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM357 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5145] The complementary binding of GAM357 RNA, herein designated GAM RNA, to target binding sites on GAM357 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM357 target RNA, herein designated GAM TARGET RNA, into GAM357 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5146] It is appreciated that GAM357 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM357 target genes. The mRNA of each one of this plurality of GAM357 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM357 RNA, herein designated GAM RNA, and which when bound by GAM357 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM357 target proteins.

[5147] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM357 gene, herein designated GAM GENE, on one or more GAM357 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5148] It is yet further appreciated that specific functions, and accordingly utilities, of GAM357 correlate with, and may be deduced from, the identity of the target genes which GAM357 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5149] Nucleotide sequences of the GAM357 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM357 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM357 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM357 are further described hereinbelow with reference to Table Table1.

[5150] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM357 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5151]

[5152] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 358 (GAM358) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5153] GAM358 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM358 was detected is described hereinabove with reference to Figs. 2-8.

[5154] GAM358 gene, herein designated GAM GENE, and GAM358 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5155] GAM358 gene, herein designated GAM GENE, encodes a GAM358 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM358 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM358 precursor RNA is designated SEQ ID:5522, and is provided hereinbelow with reference to the sequence listing part.

[5156] GAM358 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM358 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5157] An enzyme complex designated DICER COMPLEX, dices the GAM358 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM358 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of

GAM358 RNA is designated SEQ ID:5523, and is provided hereinbelow with reference to the sequence listing part.

[5158] GAM358 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM358 target RNA, herein designated GAM TARGET RNA. GAM358 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5159] GAM358 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM358 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM358 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM358 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM358 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5160] The complementary binding of GAM358 RNA, herein designated GAM RNA, to target binding sites on GAM358 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM358 target RNA, herein designated GAM TARGET RNA, into GAM358 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5161] It is appreciated that GAM358 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM358 target genes. The mRNA of each one of this plurality of GAM358 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM358 RNA, herein designated GAM RNA, and which when bound by GAM358 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM358 target proteins.

[5162] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM358 gene, herein designated GAM GENE, on one or more GAM358 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5163] It is yet further appreciated that specific functions, and accordingly utilities, of GAM358 correlate with, and may be deduced from, the identity of the target genes which GAM358 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5164] Nucleotide sequences of the GAM358 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM358 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM358 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM358 are further described hereinbelow with reference to Table Table1.

[5165] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM358 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5166]

[5167] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 359 (GAM359) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5168] GAM359 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM359 was detected is described hereinabove

with reference to Figs. 2-8.

[5169] GAM359 gene, herein designated GAM GENE, and GAM359 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5170] GAM359 gene, herein designated GAM GENE, encodes a GAM359 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM359 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM359 precursor RNA is designated SEQ ID:5541, and is provided hereinbelow with reference to the sequence listing part.

[5171] GAM359 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM359 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5172] An enzyme complex designated DICER COMPLEX, dices the GAM359 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM359 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 22%) nucleotide sequence of GAM359 RNA is designated SEQ ID:5542, and is provided hereinbelow with reference to the sequence listing part.

[5173] GAM359 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM359 target RNA, herein designated GAM TARGET RNA. GAM359 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5174] GAM359 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM359 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM359 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM359 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM359 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5175] The complementary binding of GAM359 RNA, herein designated GAM RNA, to target binding sites on GAM359 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM359 target RNA, herein designated GAM TARGET RNA, into GAM359 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[5176] It is appreciated that GAM359 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM359 target genes. The mRNA of each one of this plurality of GAM359 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM359 RNA, herein designated GAM RNA, and which when bound by GAM359 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM359 target proteins.

[5177] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM359 gene, herein designated GAM GENE, on one or more GAM359 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5178] It is yet further appreciated that specific functions, and accordingly utilities, of GAM359 correlate with, and may be deduced from, the identity of the target genes which GAM359 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5179] Nucleotide sequences of the GAM359 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM359 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM359 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM359 are further described hereinbelow with reference to Table Table1.

[5180] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM359 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5181]

[5182] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 360 (GAM360) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5183] GAM360 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM360 was detected is described hereinabove with reference to Figs. 2-8.

[5184] GAM360 gene, herein designated GAM GENE, and GAM360 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5185] GAM360 gene, herein designated GAM GENE, encodes a GAM360 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM360 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM360 precursor RNA is designated SEQ ID:5556, and is provided hereinbelow with reference to the sequence listing part.

[5186] GAM360 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM360 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5187] An enzyme complex designated DICER COMPLEX, dices the GAM360 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM360 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM360 RNA is designated SEQ ID:5557, and is provided hereinbelow with reference to the sequence listing part.

[5188] GAM360 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM360 target RNA, herein designated GAM TARGET RNA.

GAM360 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5189] GAM360 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM360 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM360 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM360 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM360 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5190] The complementary binding of GAM360 RNA, herein designated GAM RNA, to target binding sites on GAM360 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM360 target RNA, herein designated GAM TARGET RNA, into GAM360 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5191] It is appreciated that GAM360 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM360 target genes. The mRNA of each one of this plurality of GAM360 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM360 RNA, herein designated GAM RNA, and which when bound by GAM360 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM360 target proteins.

[5192] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM360 gene, herein designated GAM GENE, on one or more GAM360 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5193] It is yet further appreciated that specific functions, and accordingly utilities, of GAM360 correlate with, and may be deduced from, the identity of the target genes which GAM360 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5194] Nucleotide sequences of the GAM360 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM360 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM360 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM360 are further described hereinbelow with reference to Table Table1.

[5195] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM360 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5196]

[5197] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 361 (GAM361) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5198] GAM361 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM361 was detected is described hereinabove with reference to Figs. 2-8.

[5199] GAM361 gene, herein designated GAM GENE, and GAM361 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5200] GAM361 gene, herein designated GAM GENE, encodes a

GAM361 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM361 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM361 precursor RNA is designated SEQ ID:5566, and is provided hereinbelow with reference to the sequence listing part.

[5201] GAM361 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM361 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5202] An enzyme complex designated DICER COMPLEX, dices the GAM361 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM361 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM361 RNA is designated SEQ ID:5567, and is provided hereinbelow with reference to the sequence listing part.

[5203] GAM361 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM361 target RNA, herein designated GAM TARGET RNA. GAM361 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5204] GAM361 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM361 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM361 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM361 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM361 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5205] The complementary binding of GAM361 RNA, herein designated GAM RNA, to target binding sites on GAM361 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM361 target RNA, herein designated GAM TARGET RNA, into GAM361 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5206] It is appreciated that GAM361 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM361 target genes. The mRNA of each one of this plurality of GAM361 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM361 RNA, herein designated GAM RNA, and which when bound by GAM361 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM361 target proteins.

[5207] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM361 gene, herein designated GAM GENE, on one or more GAM361 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5208] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM361 correlate with, and may be deduced from, the identity of the target genes which GAM361 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5209] Nucleotide sequences of the GAM361 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM361 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM361 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM361 are further described hereinbelow with reference to Table Table1.

[5210] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM361 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5211]

[5212] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 362 (GAM362) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[5213] GAM362 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM362 was detected is described hereinabove with reference to Figs. 2-8.

[5214] GAM362 gene, herein designated GAM GENE, and GAM362 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5215] GAM362 gene, herein designated GAM GENE, encodes a GAM362 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM362 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM362 precursor RNA is designated SEQ ID:5577, and is provided hereinbelow with reference to the sequence listing part.

[5216] GAM362 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM362 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5217] An enzyme complex designated DICER COMPLEX, dices the GAM362 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM362 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM362 RNA is designated SEQ ID:5578, and is provided hereinbelow with reference to the sequence listing part.

[5218] GAM362 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM362 target RNA, herein designated GAM TARGET RNA.
GAM362 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5219] GAM362 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM362 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM362 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM362 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM362 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5220] The complementary binding of GAM362 RNA, herein designated GAM RNA, to target binding sites on GAM362 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM362 target RNA, herein designated GAM TARGET RNA, into GAM362 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5221] It is appreciated that GAM362 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM362 target genes. The mRNA of each one of this plurality of GAM362 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM362 RNA, herein designated GAM RNA, and which when bound by GAM362 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM362 target proteins.

[5222] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM362 gene, herein designated GAM GENE, on one or more GAM362 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5223] It is yet further appreciated that specific functions, and accordingly utilities, of GAM362 correlate with, and may be deduced from, the identity of the target genes which GAM362 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5224] Nucleotide sequences of the GAM362 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM362 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM362 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM362 are further described hereinbelow with reference to Table Table1.

[5225] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM362 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5226]

[5227] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 363 (GAM363) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5228] GAM363 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM363 was detected is described hereinabove with reference to Figs. 2-8.

[5229] GAM363 gene, herein designated GAM GENE, and GAM363 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5230] GAM363 gene, herein designated GAM GENE, encodes a GAM363 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM363 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM363 precursor RNA is designated SEQ ID:5609, and is provided hereinbelow with reference to the sequence listing part.

[5231] GAM363 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM363 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5232] An enzyme complex designated DICER COMPLEX, dices the GAM363 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM363 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM363 RNA is designated SEQ ID:5610, and is provided

hereinbelow with reference to the sequence listing part.

[5233] GAM363 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM363 target RNA, herein designated GAM TARGET RNA.

GAM363 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5234] GAM363 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM363 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM363 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM363 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM363 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5235] The complementary binding of GAM363 RNA, herein designated GAM RNA, to target binding sites on GAM363 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM363 target RNA, herein designated GAM TARGET RNA, into GAM363 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5236] It is appreciated that GAM363 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM363 target genes. The mRNA of each one of this plurality of GAM363 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM363 RNA, herein designated GAM RNA, and which when bound by GAM363 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM363 tar-

get proteins.

[5237] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM363 gene, herein designated GAM GENE, on one or more GAM363 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5238] It is yet further appreciated that specific functions, and accordingly utilities, of GAM363 correlate with, and may be deduced from, the identity of the target genes which GAM363 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5239] Nucleotide sequences of the GAM363 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM363 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM363 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM363 are further described hereinbelow with reference to Table Table1.

[5240] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM363 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5241]

[5242] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 364 (GAM364) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5243] GAM364 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM364 was detected is described hereinabove with reference to Figs. 2-8.

[5244] GAM364 gene, herein designated GAM GENE, and GAM364 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5245] GAM364 gene, herein designated GAM GENE, encodes a GAM364 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM364 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM364 precursor RNA is designated SEQ ID:5617, and is provided hereinbelow with reference to the sequence listing part.

[5246] GAM364 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM364 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5247] An enzyme complex designated DICER COMPLEX, dices

the GAM364 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM364 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM364 RNA is designated SEQ ID:5618, and is provided hereinbelow with reference to the sequence listing part.

[5248] GAM364 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM364 target RNA, herein designated GAM TARGET RNA. GAM364 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5249] GAM364 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM364 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM364

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM364 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM364 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5250] The complementary binding of GAM364 RNA, herein designated GAM RNA, to target binding sites on GAM364 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM364 target RNA, herein designated GAM TARGET RNA, into GAM364 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5251] It is appreciated that GAM364 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM364 target genes. The mRNA of each one of this plurality of GAM364 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM364 RNA, herein designated GAM RNA, and which when bound by GAM364 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM364 target proteins.

[5252] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM364 gene, herein designated GAM GENE, on one or more GAM364 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5253] It is yet further appreciated that specific functions, and accordingly utilities, of GAM364 correlate with, and may be deduced from, the identity of the target genes which GAM364 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5254] Nucleotide sequences of the GAM364 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM364 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM364 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM364 are further described hereinbelow with reference to Table Table1.

[5255] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM364 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5256]

[5257] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 365 (GAM365) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5258] GAM365 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM365 was detected is described hereinabove with reference to Figs. 2-8.

[5259] GAM365 gene, herein designated GAM GENE, and GAM365 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5260] GAM365 gene, herein designated GAM GENE, encodes a GAM365 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM365 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM365 precursor RNA is designated SEQ ID:5631, and is provided hereinbelow with reference to the sequence listing part.

[5261] GAM365 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM365 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5262] An enzyme complex designated DICER COMPLEX, dices the GAM365 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM365 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM365 RNA is designated SEQ ID:5632, and is provided hereinbelow with reference to the sequence listing part.

[5263] GAM365 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM365 target RNA, herein designated GAM TARGET RNA. GAM365 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5264] GAM365 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM365 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM365 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM365 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM365 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[5265] The complementary binding of GAM365 RNA, herein designated GAM RNA, to target binding sites on GAM365 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM365 target RNA, herein designated GAM TARGET RNA, into GAM365 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5266] It is appreciated that GAM365 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM365 target genes. The mRNA of each one of this plurality of GAM365 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM365 RNA, herein designated GAM RNA, and which when bound by GAM365 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM365 target proteins.

[5267] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM365 gene, herein designated GAM GENE, on one or

more GAM365 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5268] It is yet further appreciated that specific functions, and accordingly utilities, of GAM365 correlate with, and may be deduced from, the identity of the target genes which GAM365 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5269] Nucleotide sequences of the GAM365 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM365 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM365 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM365 are further de-

scribed hereinbelow with reference to Table Table1.

[5270] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM365 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5271]

[5272] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 366 (GAM366) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5273] GAM366 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM366 was detected is described hereinabove with reference to Figs. 2-8.

[5274] GAM366 gene, herein designated GAM GENE, and GAM366 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5275] GAM366 gene, herein designated GAM GENE, encodes a GAM366 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM366 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM366 precursor RNA is designated SEQ ID:5639, and is provided hereinbelow with reference to the sequence listing part.

[5276] GAM366 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM366 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5277] An enzyme complex designated DICER COMPLEX, dices the GAM366 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM366 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 76%) nucleotide sequence of GAM366 RNA is designated SEQ ID:5640, and is provided hereinbelow with reference to the sequence listing part.

[5278] GAM366 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM366 target RNA, herein designated GAM TARGET RNA. GAM366 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5279] GAM366 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM366 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM366 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM366 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM366 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5280] The complementary binding of GAM366 RNA, herein designated GAM RNA, to target binding sites on GAM366 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM366 target RNA, herein designated GAM TARGET RNA, into GAM366 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5281] It is appreciated that GAM366 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM366 target genes. The mRNA of each one of this plurality of GAM366 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM366 RNA, herein designated GAM RNA, and which when bound by GAM366 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM366 target proteins.

[5282] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM366 gene, herein designated GAM GENE, on one or more GAM366 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5283] It is yet further appreciated that specific functions, and accordingly utilities, of GAM366 correlate with, and may

be deduced from, the identity of the target genes which GAM366 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5284] Nucleotide sequences of the GAM366 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM366 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM366 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM366 are further described hereinbelow with reference to Table Table1.

[5285] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM366 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5286]

[5287] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 367 (GAM367) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5288] GAM367 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM367 was detected is described hereinabove with reference to Figs. 2-8.

[5289] GAM367 gene, herein designated GAM GENE, and GAM367 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5290] GAM367 gene, herein designated GAM GENE, encodes a GAM367 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM367 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM367 precursor RNA is designated SEQ ID:5660, and is provided hereinbelow with reference to the sequence listing part.

[5291] GAM367 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM367 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5292] An enzyme complex designated DICER COMPLEX, dices the GAM367 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM367 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM367 RNA is designated SEQ ID:5661, and is provided hereinbelow with reference to the sequence listing part.

[5293] GAM367 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM367 target RNA, herein designated GAM TARGET RNA. GAM367 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5294] GAM367 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM367 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM367 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM367 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM367 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5295] The complementary binding of GAM367 RNA, herein designated GAM RNA, to target binding sites on GAM367 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM367 target RNA, herein designated GAM TARGET RNA, into GAM367 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5296] It is appreciated that GAM367 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM367 target genes. The mRNA of each one of this plurality of GAM367 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM367 RNA, herein designated GAM RNA, and which when bound by GAM367 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM367 target proteins.

[5297] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM367 gene, herein designated GAM GENE, on one or more GAM367 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5298] It is yet further appreciated that specific functions, and accordingly utilities, of GAM367 correlate with, and may be deduced from, the identity of the target genes which GAM367 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5299] Nucleotide sequences of the GAM367 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM367 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM367 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM367 are further described hereinbelow with reference to Table Table1.

[5300] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM367 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5301]

[5302] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 368 (GAM368) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5303] GAM368 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM368 was detected is described hereinabove with reference to Figs. 2-8.

[5304] GAM368 gene, herein designated GAM GENE, and GAM368 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5305] GAM368 gene, herein designated GAM GENE, encodes a GAM368 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM368 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM368 precursor RNA is desig-

nated SEQ ID:5676, and is provided hereinbelow with reference to the sequence listing part.

[5306] GAM368 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM368 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5307] An enzyme complex designated DICER COMPLEX, dices the GAM368 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM368 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM368 RNA is designated SEQ ID:5677, and is provided hereinbelow with reference to the sequence listing part.

[5308] GAM368 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM368 target RNA, herein designated GAM TARGET RNA. GAM368 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5309] GAM368 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM368 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM368 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM368 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM368

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5310] The complementary binding of GAM368 RNA, herein designated GAM RNA, to target binding sites on GAM368 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM368 target RNA, herein designated GAM TARGET RNA, into GAM368 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5311] It is appreciated that GAM368 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM368 target genes. The mRNA of each one of this plurality of GAM368 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM368 RNA, herein designated GAM RNA, and which when bound by GAM368 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM368 target proteins.

[5312] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM368 gene, herein designated GAM GENE, on one or more GAM368 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5313] It is yet further appreciated that specific functions, and accordingly utilities, of GAM368 correlate with, and may be deduced from, the identity of the target genes which GAM368 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5314] Nucleotide sequences of the GAM368 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM368 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM368 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM368 are further described hereinbelow with reference to Table Table1.

[5315] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM368 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5316]

[5317] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 369 (GAM369) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5318] GAM369 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM369 was detected is described hereinabove with reference to Figs. 2-8.

[5319] GAM369 gene, herein designated GAM GENE, and

GAM369 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5320] GAM369 gene, herein designated GAM GENE, encodes a GAM369 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM369 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM369 precursor RNA is designated SEQ ID:5703, and is provided hereinbelow with reference to the sequence listing part.

[5321] GAM369 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM369 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5322] An enzyme complex designated DICER COMPLEX, dices the GAM369 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM369 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM369 RNA is designated SEQ ID:5704, and is provided hereinbelow with reference to the sequence listing part.

[5323] GAM369 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM369 target RNA, herein designated GAM TARGET RNA. GAM369 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5324] GAM369 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM369 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM369 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM369 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM369 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5325] The complementary binding of GAM369 RNA, herein designated GAM RNA, to target binding sites on GAM369 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM369 target RNA, herein designated GAM TARGET RNA, into GAM369 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5326] It is appreciated that GAM369 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM369 target genes. The mRNA of each one of this plurality of GAM369 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM369 RNA, herein designated GAM RNA, and which when bound by GAM369 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM369 target proteins.

[5327] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM369 gene, herein designated GAM GENE, on one or more GAM369 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5328] It is yet further appreciated that specific functions, and accordingly utilities, of GAM369 correlate with, and may be deduced from, the identity of the target genes which GAM369 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5329] Nucleotide sequences of the GAM369 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM369 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM369 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM369 are further described hereinbelow with reference to Table Table1.

[5330] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM369 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5331]

[5332] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 370 (GAM370) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5333] GAM370 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM370 was detected is described hereinabove with reference to Figs. 2-8.

[5334] GAM370 gene, herein designated GAM GENE, and GAM370 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5335] GAM370 gene, herein designated GAM GENE, encodes a GAM370 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM370 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM370 precursor RNA is designated SEQ ID:5719, and is provided hereinbelow with reference to the sequence listing part.

[5336] GAM370 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM370 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5337] An enzyme complex designated DICER COMPLEX, dices the GAM370 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM370 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 27%) nucleotide sequence of GAM370 RNA is designated SEQ ID:5720, and is provided hereinbelow with reference to the sequence listing part.

[5338] GAM370 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM370 target RNA, herein designated GAM TARGET RNA. GAM370 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5339] GAM370 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM370 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM370 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM370 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM370 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5340] The complementary binding of GAM370 RNA, herein designated GAM RNA, to target binding sites on GAM370 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM370 target RNA, herein designated GAM TARGET RNA, into GAM370 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5341] It is appreciated that GAM370 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM370 target genes. The mRNA of each one of this plurality of GAM370 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM370 RNA, herein designated GAM RNA, and which when bound by GAM370 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM370 target proteins.

[5342] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM370 gene, herein designated GAM GENE, on one or more GAM370 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5343] It is yet further appreciated that specific functions, and accordingly utilities, of GAM370 correlate with, and may be deduced from, the identity of the target genes which GAM370 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5344] Nucleotide sequences of the GAM370 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM370 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM370 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM370 are further described hereinbelow with reference to Table Table1.

[5345] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM370 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5346]

[5347] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 371 (GAM371) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5348] GAM371 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM371 was detected is described hereinabove with reference to Figs. 2-8.

[5349] GAM371 gene, herein designated GAM GENE, and GAM371 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5350] GAM371 gene, herein designated GAM GENE, encodes a GAM371 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM371 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM371 precursor RNA is designated SEQ ID:5729, and is provided hereinbelow with reference to the sequence listing part.

[5351] GAM371 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM371 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5352] An enzyme complex designated DICER COMPLEX, dices the GAM371 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM371 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM371 RNA is designated SEQ ID:5730, and is provided hereinbelow with reference to the sequence listing part.

[5353] GAM371 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM371 target RNA, herein designated GAM TARGET RNA. GAM371 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5354] GAM371 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM371 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM371 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM371 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM371 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5355] The complementary binding of GAM371 RNA, herein designated GAM RNA, to target binding sites on GAM371 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM371 target RNA, herein designated GAM TARGET RNA, into GAM371 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5356] It is appreciated that GAM371 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM371 target genes. The mRNA of each one of this plurality of GAM371 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM371 RNA,

herein designated GAM RNA, and which when bound by GAM371 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM371 target proteins.

[5357] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM371 gene, herein designated GAM GENE, on one or more GAM371 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5358] It is yet further appreciated that specific functions, and accordingly utilities, of GAM371 correlate with, and may be deduced from, the identity of the target genes which

GAM371 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5359] Nucleotide sequences of the GAM371 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM371 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM371 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM371 are further described hereinbelow with reference to Table Table1.

[5360] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM371 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5361]

[5362] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 372 (GAM372) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5363] GAM372 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM372 was detected is described hereinabove with reference to Figs. 2-8.

[5364] GAM372 gene, herein designated GAM GENE, and GAM372 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5365] GAM372 gene, herein designated GAM GENE, encodes a GAM372 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM372 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM372 precursor RNA is designated SEQ ID:5747, and is provided hereinbelow with reference to the sequence listing part.

[5366] GAM372 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM372 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[5367] An enzyme complex designated DICER COMPLEX, dices the GAM372 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM372 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM372 RNA is designated SEQ ID:5748, and is provided hereinbelow with reference to the sequence listing part.

[5368] GAM372 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM372 target RNA, herein designated GAM TARGET RNA.
GAM372 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5369] GAM372 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM372 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM372 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM372 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM372 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5370] The complementary binding of GAM372 RNA, herein designated GAM RNA, to target binding sites on GAM372 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM372 target RNA, herein desig-

nated GAM TARGET RNA, into GAM372 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5371] It is appreciated that GAM372 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM372 target genes. The mRNA of each one of this plurality of GAM372 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM372 RNA, herein designated GAM RNA, and which when bound by GAM372 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM372 target proteins.

[5372] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM372 gene, herein designated GAM GENE, on one or more GAM372 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5373] It is yet further appreciated that specific functions, and accordingly utilities, of GAM372 correlate with, and may be deduced from, the identity of the target genes which GAM372 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5374] Nucleotide sequences of the GAM372 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM372 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM372 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM372 are further described hereinbelow with reference to Table Table1.

[5375] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM372 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[5376]

[5377] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 373 (GAM373) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5378] GAM373 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM373 was detected is described hereinabove with reference to Figs. 2-8.

[5379] GAM373 gene, herein designated GAM GENE, and GAM373 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5380] GAM373 gene, herein designated GAM GENE, encodes a GAM373 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM373 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM373 precursor RNA is designated SEQ ID:5765, and is provided hereinbelow with ref-

erence to the sequence listing part.

[5381] GAM373 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM373 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5382] An enzyme complex designated DICER COMPLEX, dices the GAM373 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM373 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM373 RNA is designated SEQ ID:5766, and is provided hereinbelow with reference to the sequence listing part.

[5383] GAM373 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM373 target RNA, herein designated GAM TARGET RNA.

GAM373 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5384] GAM373 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM373 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM373 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM373 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM373 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5385] The complementary binding of GAM373 RNA, herein designated GAM RNA, to target binding sites on GAM373 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM373 target RNA, herein designated GAM TARGET RNA, into GAM373 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5386] It is appreciated that GAM373 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM373 target genes. The mRNA of each one of this plurality of GAM373 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM373 RNA, herein designated GAM RNA, and which when bound by GAM373 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM373 target proteins.

[5387] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM373 gene, herein designated GAM GENE, on one or more GAM373 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5388] It is yet further appreciated that specific functions, and accordingly utilities, of GAM373 correlate with, and may be deduced from, the identity of the target genes which GAM373 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5389] Nucleotide sequences of the GAM373 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM373 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM373 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM373 are further described hereinbelow with reference to Table Table1.

[5390] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM373 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5391]

[5392] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 374 (GAM374) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5393] GAM374 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM374 was detected is described hereinabove with reference to Figs. 2-8.

[5394] GAM374 gene, herein designated GAM GENE, and GAM374 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[5395] GAM374 gene, herein designated GAM GENE, encodes a GAM374 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM374 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM374 precursor RNA is designated SEQ ID:5800, and is provided hereinbelow with reference to the sequence listing part.

[5396] GAM374 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM374 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5397] An enzyme complex designated DICER COMPLEX, dices the GAM374 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM374 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM374 RNA is designated SEQ ID:5801, and is provided hereinbelow with reference to the sequence listing part.

[5398] GAM374 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM374 target RNA, herein designated GAM TARGET RNA. GAM374 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5399] GAM374 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM374 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM374 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM374 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM374 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5400] The complementary binding of GAM374 RNA, herein designated GAM RNA, to target binding sites on GAM374 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM374 target RNA, herein designated GAM TARGET RNA, into GAM374 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5401] It is appreciated that GAM374 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM374 target genes. The mRNA of each one of this plurality of GAM374 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM374 RNA, herein designated GAM RNA, and which when bound by GAM374 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM374 target proteins.

[5402] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM374 gene, herein designated GAM GENE, on one or more GAM374 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[5403] It is yet further appreciated that specific functions, and accordingly utilities, of GAM374 correlate with, and may be deduced from, the identity of the target genes which GAM374 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5404] Nucleotide sequences of the GAM374 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM374 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM374 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM374 are further described hereinbelow with reference to Table Table1.

[5405] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM374 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5406]

[5407] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

375 (GAM375) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5408] GAM375 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM375 was detected is described hereinabove with reference to Figs. 2-8.

[5409] GAM375 gene, herein designated GAM GENE, and GAM375 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5410] GAM375 gene, herein designated GAM GENE, encodes a GAM375 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM375 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM375 precursor RNA is designated SEQ ID:5813, and is provided hereinbelow with reference to the sequence listing part.

[5411] GAM375 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM375 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5412] An enzyme complex designated DICER COMPLEX, dices the GAM375 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM375 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM375 RNA is designated SEQ ID:5814, and is provided hereinbelow with reference to the sequence listing part.

[5413] GAM375 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM375 target RNA, herein designated GAM TARGET RNA. GAM375 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5414] GAM375 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM375 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM375 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM375 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM375 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5415] The complementary binding of GAM375 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM375 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM375 target RNA, herein designated GAM TARGET RNA, into GAM375 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5416] It is appreciated that GAM375 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM375 target genes. The mRNA of each one of this plurality of GAM375 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM375 RNA, herein designated GAM RNA, and which when bound by GAM375 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM375 target proteins.

[5417] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM375 gene, herein designated GAM GENE, on one or more GAM375 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5418] It is yet further appreciated that specific functions, and accordingly utilities, of GAM375 correlate with, and may be deduced from, the identity of the target genes which GAM375 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5419] Nucleotide sequences of the GAM375 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM375 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM375 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM375 are further described hereinbelow with reference to Table Table1.

[5420] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM375 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5421]

[5422] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 376 (GAM376) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5423] GAM376 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM376 was detected is described hereinabove with reference to Figs. 2-8.

[5424] GAM376 gene, herein designated GAM GENE, and GAM376 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5425] GAM376 gene, herein designated GAM GENE, encodes a GAM376 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM376 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM376 precursor RNA is designated SEQ ID:5837, and is provided hereinbelow with reference to the sequence listing part.

[5426] GAM376 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM376 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5427] An enzyme complex designated DICER COMPLEX, dices the GAM376 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM376 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 68%) nucleotide sequence of GAM376 RNA is designated SEQ ID:5838, and is provided hereinbelow with reference to the sequence listing part.

[5428] GAM376 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM376 target RNA, herein designated GAM TARGET RNA. GAM376 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5429] GAM376 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM376 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM376 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM376 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM376 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5430] The complementary binding of GAM376 RNA, herein designated GAM RNA, to target binding sites on GAM376 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM376 target RNA, herein designated GAM TARGET RNA, into GAM376 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5431] It is appreciated that GAM376 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM376 target genes. The mRNA of each one of this plurality of GAM376 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM376 RNA, herein designated GAM RNA, and which when bound by

GAM376 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM376 target proteins.

[5432] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM376 gene, herein designated GAM GENE, on one or more GAM376 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5433] It is yet further appreciated that specific functions, and accordingly utilities, of GAM376 correlate with, and may be deduced from, the identity of the target genes which GAM376 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[5434] Nucleotide sequences of the GAM376 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM376 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM376 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM376 are further described hereinbelow with reference to Table Table1.

[5435] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM376 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5436]

[5437] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 377 (GAM377) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5438] GAM377 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM377 was detected is described hereinabove with reference to Figs. 2-8.

[5439] GAM377 gene, herein designated GAM GENE, and GAM377 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5440] GAM377 gene, herein designated GAM GENE, encodes a GAM377 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM377 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM377 precursor RNA is designated SEQ ID:5849, and is provided hereinbelow with reference to the sequence listing part.

[5441] GAM377 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM377 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[5442] An enzyme complex designated DICER COMPLEX, dices the GAM377 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM377 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM377 RNA is designated SEQ ID:5850, and is provided hereinbelow with reference to the sequence listing part.

[5443] GAM377 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM377 target RNA, herein designated GAM TARGET RNA. GAM377 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5444] GAM377 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM377 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM377 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM377 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM377 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5445] The complementary binding of GAM377 RNA, herein designated GAM RNA, to target binding sites on GAM377 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM377 target RNA, herein designated GAM TARGET RNA, into GAM377 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5446] It is appreciated that GAM377 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM377 target genes. The mRNA of each one of this plurality of GAM377 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM377 RNA, herein designated GAM RNA, and which when bound by GAM377 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM377 target proteins.

[5447] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM377 gene, herein designated GAM GENE, on one or more GAM377 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5448] It is yet further appreciated that specific functions, and accordingly utilities, of GAM377 correlate with, and may be deduced from, the identity of the target genes which GAM377 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5449] Nucleotide sequences of the GAM377 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM377 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM377 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM377 are further described hereinbelow with reference to Table Table1.

[5450] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM377 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5451]

[5452] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 378 (GAM378) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5453] GAM378 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM378 was detected is described hereinabove with reference to Figs. 2-8.

[5454] GAM378 gene, herein designated GAM GENE, and GAM378 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5455] GAM378 gene, herein designated GAM GENE, encodes a GAM378 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM378 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM378 precursor RNA is designated SEQ ID:5853, and is provided hereinbelow with reference to the sequence listing part.

[5456] GAM378 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM378 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5457] An enzyme complex designated DICER COMPLEX, dices the GAM378 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM378 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM378 RNA is designated SEQ ID:5854, and is provided hereinbelow with reference to the sequence listing part.

[5458] GAM378 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM378

target RNA, herein designated GAM TARGET RNA.

GAM378 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5459] GAM378 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM378 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM378 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM378 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM378 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5460] The complementary binding of GAM378 RNA, herein designated GAM RNA, to target binding sites on GAM378 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM378 target RNA, herein designated GAM TARGET RNA, into GAM378 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5461] It is appreciated that GAM378 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM378 target genes. The mRNA of each one of this plurality of GAM378 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM378 RNA, herein designated GAM RNA, and which when bound by GAM378 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM378 target proteins.

[5462] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM378 gene, herein designated GAM GENE, on one or more GAM378 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5463] It is yet further appreciated that specific functions, and accordingly utilities, of GAM378 correlate with, and may be deduced from, the identity of the target genes which GAM378 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5464] Nucleotide sequences of the GAM378 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM378 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM378 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM378 are further described hereinbelow with reference to Table Table1.

[5465] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM378 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5466]

[5467] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 379 (GAM379) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5468] GAM379 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM379 was detected is described hereinabove with reference to Figs. 2-8.

[5469] GAM379 gene, herein designated GAM GENE, and GAM379 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5470] GAM379 gene, herein designated GAM GENE, encodes a GAM379 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM379 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM379 precursor RNA is designated SEQ ID:5876, and is provided hereinbelow with reference to the sequence listing part.

[5471] GAM379 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM379 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5472] An enzyme complex designated DICER COMPLEX, dices the GAM379 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM379 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM379 RNA is designated SEQ ID:2525, and is provided hereinbelow with reference to the sequence listing part.

[5473] GAM379 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM379 target RNA, herein designated GAM TARGET RNA. GAM379 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5474] GAM379 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM379 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM379 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM379 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM379 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5475] The complementary binding of GAM379 RNA, herein designated GAM RNA, to target binding sites on GAM379 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM379 target RNA, herein designated GAM TARGET RNA, into GAM379 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5476] It is appreciated that GAM379 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM379 target genes. The mRNA of each one of this plu-

rality of GAM379 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM379 RNA, herein designated GAM RNA, and which when bound by GAM379 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM379 target proteins.

[5477] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM379 gene, herein designated GAM GENE, on one or more GAM379 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5478] It is yet further appreciated that specific functions, and accordingly utilities, of GAM379 correlate with, and may be deduced from, the identity of the target genes which GAM379 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5479] Nucleotide sequences of the GAM379 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM379 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM379 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM379 are further described hereinbelow with reference to Table Table1.

[5480] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM379 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5481]

[5482] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 380 (GAM380) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[5483] GAM380 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM380 was detected is described hereinabove with reference to Figs. 2-8.

[5484] GAM380 gene, herein designated GAM GENE, and GAM380 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5485] GAM380 gene, herein designated GAM GENE, encodes a GAM380 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM380 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM380 precursor RNA is designated SEQ ID:5879, and is provided hereinbelow with reference to the sequence listing part.

[5486] GAM380 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM380 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5487] An enzyme complex designated DICER COMPLEX, dices the GAM380 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM380 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 26%) nucleotide sequence of GAM380 RNA is designated SEQ ID:5880, and is provided hereinbelow with reference to the sequence listing part.

[5488] GAM380 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM380 target RNA, herein designated GAM TARGET RNA.
GAM380 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[5489] GAM380 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM380 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM380 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM380 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM380 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5490] The complementary binding of GAM380 RNA, herein designated GAM RNA, to target binding sites on GAM380 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM380 target RNA, herein designated GAM TARGET RNA, into GAM380 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5491] It is appreciated that GAM380 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM380 target genes. The mRNA of each one of this plurality of GAM380 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM380 RNA, herein designated GAM RNA, and which when bound by GAM380 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM380 target proteins.

[5492] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM380 gene, herein designated GAM GENE, on one or more GAM380 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5493] It is yet further appreciated that specific functions, and accordingly utilities, of GAM380 correlate with, and may be deduced from, the identity of the target genes which GAM380 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5494] Nucleotide sequences of the GAM380 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM380 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM380 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM380 are further described hereinbelow with reference to Table Table1.

[5495] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM380 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5496]

[5497] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 381 (GAM381) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5498] GAM381 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM381 was detected is described hereinabove with reference to Figs. 2-8.

[5499] GAM381 gene, herein designated GAM GENE, and GAM381 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5500] GAM381 gene, herein designated GAM GENE, encodes a GAM381 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM381 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM381 precursor RNA is designated SEQ ID:5892, and is provided hereinbelow with reference to the sequence listing part.

[5501] GAM381 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM381 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5502] An enzyme complex designated DICER COMPLEX, dices the GAM381 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM381 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of

GAM381 RNA is designated SEQ ID:5893, and is provided hereinbelow with reference to the sequence listing part.

[5503] GAM381 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM381 target RNA, herein designated GAM TARGET RNA. GAM381 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5504] GAM381 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM381 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM381 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM381 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM381 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5505] The complementary binding of GAM381 RNA, herein designated GAM RNA, to target binding sites on GAM381 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM381 target RNA, herein designated GAM TARGET RNA, into GAM381 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5506] It is appreciated that GAM381 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM381 target genes. The mRNA of each one of this plurality of GAM381 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM381 RNA, herein designated GAM RNA, and which when bound by GAM381 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM381 target proteins.

[5507] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM381 gene, herein designated GAM GENE, on one or more GAM381 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5508] It is yet further appreciated that specific functions, and accordingly utilities, of GAM381 correlate with, and may be deduced from, the identity of the target genes which GAM381 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5509] Nucleotide sequences of the GAM381 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM381 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM381 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM381 are further described hereinbelow with reference to Table Table1.

[5510] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM381 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5511]

[5512] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 382 (GAM382) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5513] GAM382 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM382 was detected is described hereinabove

with reference to Figs. 2-8.

[5514] GAM382 gene, herein designated GAM GENE, and GAM382 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5515] GAM382 gene, herein designated GAM GENE, encodes a GAM382 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM382 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM382 precursor RNA is designated SEQ ID:5913, and is provided hereinbelow with reference to the sequence listing part.

[5516] GAM382 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM382 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5517] An enzyme complex designated DICER COMPLEX, dices the GAM382 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM382 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM382 RNA is designated SEQ ID:5914, and is provided hereinbelow with reference to the sequence listing part.

[5518] GAM382 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM382 target RNA, herein designated GAM TARGET RNA. GAM382 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5519] GAM382 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM382 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM382 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM382 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM382 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5520] The complementary binding of GAM382 RNA, herein designated GAM RNA, to target binding sites on GAM382 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM382 target RNA, herein designated GAM TARGET RNA, into GAM382 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[5521] It is appreciated that GAM382 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM382 target genes. The mRNA of each one of this plurality of GAM382 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM382 RNA, herein designated GAM RNA, and which when bound by GAM382 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM382 target proteins.

[5522] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM382 gene, herein designated GAM GENE, on one or more GAM382 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5523] It is yet further appreciated that specific functions, and accordingly utilities, of GAM382 correlate with, and may be deduced from, the identity of the target genes which GAM382 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5524] Nucleotide sequences of the GAM382 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM382 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM382 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM382 are further described hereinbelow with reference to Table Table1.

[5525] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM382 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5526]

[5527] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 383 (GAM383) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5528] GAM383 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM383 was detected is described hereinabove with reference to Figs. 2-8.

[5529] GAM383 gene, herein designated GAM GENE, and GAM383 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5530] GAM383 gene, herein designated GAM GENE, encodes a GAM383 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM383 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM383 precursor RNA is designated SEQ ID:5924, and is provided hereinbelow with reference to the sequence listing part.

[5531] GAM383 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM383 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5532] An enzyme complex designated DICER COMPLEX, dices the GAM383 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM383 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM383 RNA is designated SEQ ID:5925, and is provided hereinbelow with reference to the sequence listing part.

[5533] GAM383 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM383 target RNA, herein designated GAM TARGET RNA.

GAM383 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5534] GAM383 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM383 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM383 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM383 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM383 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5535] The complementary binding of GAM383 RNA, herein designated GAM RNA, to target binding sites on GAM383 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM383 target RNA, herein designated GAM TARGET RNA, into GAM383 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5536] It is appreciated that GAM383 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM383 target genes. The mRNA of each one of this plurality of GAM383 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM383 RNA, herein designated GAM RNA, and which when bound by GAM383 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM383 target proteins.

[5537] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM383 gene, herein designated GAM GENE, on one or more GAM383 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5538] It is yet further appreciated that specific functions, and accordingly utilities, of GAM383 correlate with, and may be deduced from, the identity of the target genes which GAM383 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5539] Nucleotide sequences of the GAM383 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM383 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM383 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM383 are further described hereinbelow with reference to Table Table1.

[5540] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM383 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5541]

[5542] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 384 (GAM384) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5543] GAM384 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM384 was detected is described hereinabove with reference to Figs. 2-8.

[5544] GAM384 gene, herein designated GAM GENE, and GAM384 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5545] GAM384 gene, herein designated GAM GENE, encodes a

GAM384 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM384 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM384 precursor RNA is designated SEQ ID:5945, and is provided hereinbelow with reference to the sequence listing part.

[5546] GAM384 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM384 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5547] An enzyme complex designated DICER COMPLEX, dices the GAM384 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM384 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 40%) nucleotide sequence of GAM384 RNA is designated SEQ ID:5946, and is provided hereinbelow with reference to the sequence listing part.

[5548] GAM384 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM384 target RNA, herein designated GAM TARGET RNA. GAM384 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5549] GAM384 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM384 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM384 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM384 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM384 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5550] The complementary binding of GAM384 RNA, herein designated GAM RNA, to target binding sites on GAM384 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM384 target RNA, herein designated GAM TARGET RNA, into GAM384 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5551] It is appreciated that GAM384 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM384 target genes. The mRNA of each one of this plurality of GAM384 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM384 RNA, herein designated GAM RNA, and which when bound by GAM384 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM384 target proteins.

[5552] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM384 gene, herein designated GAM GENE, on one or more GAM384 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5553] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM384 correlate with, and may be deduced from, the identity of the target genes which GAM384 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5554] Nucleotide sequences of the GAM384 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM384 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM384 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM384 are further described hereinbelow with reference to Table Table1.

[5555] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM384 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5556]

[5557] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 385 (GAM385) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[5558] GAM385 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM385 was detected is described hereinabove with reference to Figs. 2-8.

[5559] GAM385 gene, herein designated GAM GENE, and GAM385 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5560] GAM385 gene, herein designated GAM GENE, encodes a GAM385 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM385 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM385 precursor RNA is designated SEQ ID:5969, and is provided hereinbelow with reference to the sequence listing part.

[5561] GAM385 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM385 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5562] An enzyme complex designated DICER COMPLEX, dices the GAM385 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM385 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM385 RNA is designated SEQ ID:5970, and is provided hereinbelow with reference to the sequence listing part.

[5563] GAM385 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM385 target RNA, herein designated GAM TARGET RNA. GAM385 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5564] GAM385 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM385 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM385 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM385 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM385 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5565] The complementary binding of GAM385 RNA, herein designated GAM RNA, to target binding sites on GAM385 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM385 target RNA, herein designated GAM TARGET RNA, into GAM385 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5566] It is appreciated that GAM385 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM385 target genes. The mRNA of each one of this plurality of GAM385 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM385 RNA, herein designated GAM RNA, and which when bound by GAM385 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM385 target proteins.

[5567] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM385 gene, herein designated GAM GENE, on one or more GAM385 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5568] It is yet further appreciated that specific functions, and accordingly utilities, of GAM385 correlate with, and may be deduced from, the identity of the target genes which GAM385 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5569] Nucleotide sequences of the GAM385 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM385 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM385 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM385 are further described hereinbelow with reference to Table Table1.

[5570] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM385 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5571]

[5572] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 386 (GAM386) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5573] GAM386 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM386 was detected is described hereinabove with reference to Figs. 2-8.

[5574] GAM386 gene, herein designated GAM GENE, and GAM386 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5575] GAM386 gene, herein designated GAM GENE, encodes a GAM386 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM386 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM386 precursor RNA is designated SEQ ID:5975, and is provided hereinbelow with reference to the sequence listing part.

[5576] GAM386 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM386 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5577] An enzyme complex designated DICER COMPLEX, dices the GAM386 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM386 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM386 RNA is designated SEQ ID:5976, and is provided

hereinbelow with reference to the sequence listing part.

[5578] GAM386 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM386 target RNA, herein designated GAM TARGET RNA.

GAM386 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5579] GAM386 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM386 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM386 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM386 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM386 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5580] The complementary binding of GAM386 RNA, herein designated GAM RNA, to target binding sites on GAM386 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM386 target RNA, herein designated GAM TARGET RNA, into GAM386 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5581] It is appreciated that GAM386 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM386 target genes. The mRNA of each one of this plurality of GAM386 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM386 RNA, herein designated GAM RNA, and which when bound by GAM386 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM386 tar-

get proteins.

[5582] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM386 gene, herein designated GAM GENE, on one or more GAM386 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5583] It is yet further appreciated that specific functions, and accordingly utilities, of GAM386 correlate with, and may be deduced from, the identity of the target genes which GAM386 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5584] Nucleotide sequences of the GAM386 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM386 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM386 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM386 are further described hereinbelow with reference to Table Table1.

[5585] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM386 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5586]

[5587] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 387 (GAM387) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5588] GAM387 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM387 was detected is described hereinabove with reference to Figs. 2-8.

[5589] GAM387 gene, herein designated GAM GENE, and GAM387 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5590] GAM387 gene, herein designated GAM GENE, encodes a GAM387 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM387 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM387 precursor RNA is designated SEQ ID:5985, and is provided hereinbelow with reference to the sequence listing part.

[5591] GAM387 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM387 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5592] An enzyme complex designated DICER COMPLEX, dices

the GAM387 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM387 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM387 RNA is designated SEQ ID:5986, and is provided hereinbelow with reference to the sequence listing part.

[5593] GAM387 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM387 target RNA, herein designated GAM TARGET RNA. GAM387 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5594] GAM387 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM387 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM387

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM387 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM387 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5595] The complementary binding of GAM387 RNA, herein designated GAM RNA, to target binding sites on GAM387 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM387 target RNA, herein designated GAM TARGET RNA, into GAM387 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5596] It is appreciated that GAM387 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM387 target genes. The mRNA of each one of this plurality of GAM387 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM387 RNA, herein designated GAM RNA, and which when bound by GAM387 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM387 target proteins.

[5597] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM387 gene, herein designated GAM GENE, on one or more GAM387 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5598] It is yet further appreciated that specific functions, and accordingly utilities, of GAM387 correlate with, and may be deduced from, the identity of the target genes which GAM387 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5599] Nucleotide sequences of the GAM387 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM387 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM387 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM387 are further described hereinbelow with reference to Table Table1.

[5600] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM387 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5601]

[5602] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 388 (GAM388) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5603] GAM388 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM388 was detected is described hereinabove with reference to Figs. 2-8.

[5604] GAM388 gene, herein designated GAM GENE, and GAM388 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5605] GAM388 gene, herein designated GAM GENE, encodes a GAM388 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM388 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM388 precursor RNA is designated SEQ ID:6007, and is provided hereinbelow with reference to the sequence listing part.

[5606] GAM388 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM388 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5607] An enzyme complex designated DICER COMPLEX, dices the GAM388 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM388 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM388 RNA is designated SEQ ID:6008, and is provided hereinbelow with reference to the sequence listing part.

[5608] GAM388 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM388 target RNA, herein designated GAM TARGET RNA. GAM388 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5609] GAM388 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM388 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM388 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM388 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM388 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[5610] The complementary binding of GAM388 RNA, herein designated GAM RNA, to target binding sites on GAM388 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM388 target RNA, herein designated GAM TARGET RNA, into GAM388 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5611] It is appreciated that GAM388 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM388 target genes. The mRNA of each one of this plurality of GAM388 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM388 RNA, herein designated GAM RNA, and which when bound by GAM388 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM388 target proteins.

[5612] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM388 gene, herein designated GAM GENE, on one or

more GAM388 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5613] It is yet further appreciated that specific functions, and accordingly utilities, of GAM388 correlate with, and may be deduced from, the identity of the target genes which GAM388 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5614] Nucleotide sequences of the GAM388 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM388 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM388 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM388 are further de-

scribed hereinbelow with reference to Table Table1.

[5615] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM388 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5616]

[5617] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 389 (GAM389) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5618] GAM389 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM389 was detected is described hereinabove with reference to Figs. 2-8.

[5619] GAM389 gene, herein designated GAM GENE, and GAM389 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5620] GAM389 gene, herein designated GAM GENE, encodes a GAM389 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM389 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM389 precursor RNA is designated SEQ ID:6019, and is provided hereinbelow with reference to the sequence listing part.

[5621] GAM389 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM389 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5622] An enzyme complex designated DICER COMPLEX, dices the GAM389 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM389 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM389 RNA is designated SEQ ID:6020, and is provided hereinbelow with reference to the sequence listing part.

[5623] GAM389 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM389 target RNA, herein designated GAM TARGET RNA. GAM389 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5624] GAM389 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM389 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM389 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM389 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM389 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5625] The complementary binding of GAM389 RNA, herein designated GAM RNA, to target binding sites on GAM389 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM389 target RNA, herein designated GAM TARGET RNA, into GAM389 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5626] It is appreciated that GAM389 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM389 target genes. The mRNA of each one of this plurality of GAM389 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM389 RNA, herein designated GAM RNA, and which when bound by GAM389 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM389 target proteins.

[5627] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM389 gene, herein designated GAM GENE, on one or more GAM389 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5628] It is yet further appreciated that specific functions, and accordingly utilities, of GAM389 correlate with, and may

be deduced from, the identity of the target genes which GAM389 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5629] Nucleotide sequences of the GAM389 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM389 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM389 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM389 are further described hereinbelow with reference to Table Table1.

[5630] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM389 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5631]

[5632] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 390 (GAM390) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5633] GAM390 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM390 was detected is described hereinabove with reference to Figs. 2-8.

[5634] GAM390 gene, herein designated GAM GENE, and GAM390 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5635] GAM390 gene, herein designated GAM GENE, encodes a GAM390 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM390 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM390 precursor RNA is designated SEQ ID:6033, and is provided hereinbelow with reference to the sequence listing part.

[5636] GAM390 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM390 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5637] An enzyme complex designated DICER COMPLEX, dices the GAM390 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM390 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM390 RNA is designated SEQ ID:6034, and is provided hereinbelow with reference to the sequence listing part.

[5638] GAM390 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM390 target RNA, herein designated GAM TARGET RNA. GAM390 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5639] GAM390 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM390 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM390 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM390 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM390 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5640] The complementary binding of GAM390 RNA, herein designated GAM RNA, to target binding sites on GAM390 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM390 target RNA, herein designated GAM TARGET RNA, into GAM390 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5641] It is appreciated that GAM390 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM390 target genes. The mRNA of each one of this plurality of GAM390 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM390 RNA, herein designated GAM RNA, and which when bound by GAM390 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM390 target proteins.

[5642] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM390 gene, herein designated GAM GENE, on one or more GAM390 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5643] It is yet further appreciated that specific functions, and accordingly utilities, of GAM390 correlate with, and may be deduced from, the identity of the target genes which GAM390 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5644] Nucleotide sequences of the GAM390 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM390 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM390 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM390 are further described hereinbelow with reference to Table Table1.

[5645] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM390 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5646]

[5647] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 391 (GAM391) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5648] GAM391 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM391 was detected is described hereinabove with reference to Figs. 2-8.

[5649] GAM391 gene, herein designated GAM GENE, and GAM391 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5650] GAM391 gene, herein designated GAM GENE, encodes a GAM391 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM391 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM391 precursor RNA is desig-

nated SEQ ID:6044, and is provided hereinbelow with reference to the sequence listing part.

[5651] GAM391 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM391 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5652] An enzyme complex designated DICER COMPLEX, dices the GAM391 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM391 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 42%) nucleotide sequence of GAM391 RNA is designated SEQ ID:6045, and is provided hereinbelow with reference to the sequence listing part.

[5653] GAM391 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM391 target RNA, herein designated GAM TARGET RNA. GAM391 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5654] GAM391 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM391 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM391 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM391 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM391

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5655] The complementary binding of GAM391 RNA, herein designated GAM RNA, to target binding sites on GAM391 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM391 target RNA, herein designated GAM TARGET RNA, into GAM391 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5656] It is appreciated that GAM391 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM391 target genes. The mRNA of each one of this plurality of GAM391 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM391 RNA, herein designated GAM RNA, and which when bound by GAM391 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM391 target proteins.

[5657] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM391 gene, herein designated GAM GENE, on one or more GAM391 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5658] It is yet further appreciated that specific functions, and accordingly utilities, of GAM391 correlate with, and may be deduced from, the identity of the target genes which GAM391 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5659] Nucleotide sequences of the GAM391 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM391 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM391 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM391 are further described hereinbelow with reference to Table Table1.

[5660] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM391 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5661]

[5662] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 392 (GAM392) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5663] GAM392 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM392 was detected is described hereinabove with reference to Figs. 2-8.

[5664] GAM392 gene, herein designated GAM GENE, and

GAM392 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5665] GAM392 gene, herein designated GAM GENE, encodes a GAM392 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM392 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM392 precursor RNA is designated SEQ ID:6066, and is provided hereinbelow with reference to the sequence listing part.

[5666] GAM392 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM392 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5667] An enzyme complex designated DICER COMPLEX, dices the GAM392 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM392 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 44%) nucleotide sequence of GAM392 RNA is designated SEQ ID:6067, and is provided hereinbelow with reference to the sequence listing part.

[5668] GAM392 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM392 target RNA, herein designated GAM TARGET RNA. GAM392 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5669] GAM392 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM392 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM392 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM392 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM392 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5670] The complementary binding of GAM392 RNA, herein designated GAM RNA, to target binding sites on GAM392 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM392 target RNA, herein designated GAM TARGET RNA, into GAM392 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5671] It is appreciated that GAM392 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM392 target genes. The mRNA of each one of this plurality of GAM392 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM392 RNA, herein designated GAM RNA, and which when bound by GAM392 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM392 target proteins.

[5672] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM392 gene, herein designated GAM GENE, on one or more GAM392 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5673] It is yet further appreciated that specific functions, and accordingly utilities, of GAM392 correlate with, and may be deduced from, the identity of the target genes which GAM392 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5674] Nucleotide sequences of the GAM392 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM392 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM392 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM392 are further described hereinbelow with reference to Table Table1.

[5675] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM392 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5676]

[5677] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 393 (GAM393) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5678] GAM393 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM393 was detected is described hereinabove with reference to Figs. 2-8.

[5679] GAM393 gene, herein designated GAM GENE, and GAM393 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5680] GAM393 gene, herein designated GAM GENE, encodes a GAM393 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM393 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM393 precursor RNA is designated SEQ ID:6072, and is provided hereinbelow with reference to the sequence listing part.

[5681] GAM393 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM393 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5682] An enzyme complex designated DICER COMPLEX, dices the GAM393 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM393 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM393 RNA is designated SEQ ID:6073, and is provided hereinbelow with reference to the sequence listing part.

[5683] GAM393 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM393 target RNA, herein designated GAM TARGET RNA. GAM393 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5684] GAM393 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM393 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM393 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM393 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM393 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5685] The complementary binding of GAM393 RNA, herein designated GAM RNA, to target binding sites on GAM393 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM393 target RNA, herein designated GAM TARGET RNA, into GAM393 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5686] It is appreciated that GAM393 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM393 target genes. The mRNA of each one of this plurality of GAM393 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM393 RNA, herein designated GAM RNA, and which when bound by GAM393 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM393 target proteins.

[5687] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM393 gene, herein designated GAM GENE, on one or more GAM393 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5688] It is yet further appreciated that specific functions, and accordingly utilities, of GAM393 correlate with, and may be deduced from, the identity of the target genes which GAM393 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5689] Nucleotide sequences of the GAM393 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM393 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM393 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM393 are further described hereinbelow with reference to Table Table1.

[5690] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM393 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5691]

[5692] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 394 (GAM394) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5693] GAM394 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM394 was detected is described hereinabove with reference to Figs. 2-8.

[5694] GAM394 gene, herein designated GAM GENE, and GAM394 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5695] GAM394 gene, herein designated GAM GENE, encodes a GAM394 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM394 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM394 precursor RNA is designated SEQ ID:6089, and is provided hereinbelow with reference to the sequence listing part.

[5696] GAM394 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM394 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5697] An enzyme complex designated DICER COMPLEX, dices the GAM394 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM394 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM394 RNA is designated SEQ ID:6090, and is provided hereinbelow with reference to the sequence listing part.

[5698] GAM394 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM394 target RNA, herein designated GAM TARGET RNA. GAM394 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5699] GAM394 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM394 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM394 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM394 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM394 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5700] The complementary binding of GAM394 RNA, herein designated GAM RNA, to target binding sites on GAM394 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM394 target RNA, herein designated GAM TARGET RNA, into GAM394 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5701] It is appreciated that GAM394 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM394 target genes. The mRNA of each one of this plurality of GAM394 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM394 RNA,

herein designated GAM RNA, and which when bound by GAM394 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM394 target proteins.

[5702] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM394 gene, herein designated GAM GENE, on one or more GAM394 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5703] It is yet further appreciated that specific functions, and accordingly utilities, of GAM394 correlate with, and may be deduced from, the identity of the target genes which

GAM394 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5704] Nucleotide sequences of the GAM394 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM394 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM394 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM394 are further described hereinbelow with reference to Table Table1.

[5705] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM394 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5706]

[5707] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 395 (GAM395) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5708] GAM395 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM395 was detected is described hereinabove with reference to Figs. 2-8.

[5709] GAM395 gene, herein designated GAM GENE, and GAM395 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5710] GAM395 gene, herein designated GAM GENE, encodes a GAM395 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM395 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM395 precursor RNA is designated SEQ ID:6103, and is provided hereinbelow with reference to the sequence listing part.

[5711] GAM395 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM395 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[5712] An enzyme complex designated DICER COMPLEX, dices the GAM395 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM395 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM395 RNA is designated SEQ ID:6104, and is provided hereinbelow with reference to the sequence listing part.

[5713] GAM395 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM395 target RNA, herein designated GAM TARGET RNA.
GAM395 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5714] GAM395 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM395 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM395 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM395 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM395 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5715] The complementary binding of GAM395 RNA, herein designated GAM RNA, to target binding sites on GAM395 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM395 target RNA, herein desig-

nated GAM TARGET RNA, into GAM395 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5716] It is appreciated that GAM395 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM395 target genes. The mRNA of each one of this plurality of GAM395 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM395 RNA, herein designated GAM RNA, and which when bound by GAM395 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM395 target proteins.

[5717] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM395 gene, herein designated GAM GENE, on one or more GAM395 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5718] It is yet further appreciated that specific functions, and accordingly utilities, of GAM395 correlate with, and may be deduced from, the identity of the target genes which GAM395 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5719] Nucleotide sequences of the GAM395 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM395 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM395 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM395 are further described hereinbelow with reference to Table Table1.

[5720] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM395 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[5721]

[5722] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 396 (GAM396) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5723] GAM396 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM396 was detected is described hereinabove with reference to Figs. 2-8.

[5724] GAM396 gene, herein designated GAM GENE, and GAM396 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5725] GAM396 gene, herein designated GAM GENE, encodes a GAM396 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM396 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM396 precursor RNA is designated SEQ ID:6117, and is provided hereinbelow with ref-

erence to the sequence listing part.

[5726] GAM396 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM396 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5727] An enzyme complex designated DICER COMPLEX, dices the GAM396 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM396 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM396 RNA is designated SEQ ID:6118, and is provided hereinbelow with reference to the sequence listing part.

[5728] GAM396 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM396 target RNA, herein designated GAM TARGET RNA.

GAM396 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5729] GAM396 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM396 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM396 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM396 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM396 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5730] The complementary binding of GAM396 RNA, herein designated GAM RNA, to target binding sites on GAM396 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM396 target RNA, herein designated GAM TARGET RNA, into GAM396 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5731] It is appreciated that GAM396 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM396 target genes. The mRNA of each one of this plurality of GAM396 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM396 RNA, herein designated GAM RNA, and which when bound by GAM396 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM396 target proteins.

[5732] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM396 gene, herein designated GAM GENE, on one or more GAM396 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5733] It is yet further appreciated that specific functions, and accordingly utilities, of GAM396 correlate with, and may be deduced from, the identity of the target genes which GAM396 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5734] Nucleotide sequences of the GAM396 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM396 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM396 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM396 are further described hereinbelow with reference to Table Table1.

[5735] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM396 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5736]

[5737] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 397 (GAM397) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5738] GAM397 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM397 was detected is described hereinabove with reference to Figs. 2-8.

[5739] GAM397 gene, herein designated GAM GENE, and GAM397 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[5740] GAM397 gene, herein designated GAM GENE, encodes a GAM397 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM397 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM397 precursor RNA is designated SEQ ID:6135, and is provided hereinbelow with reference to the sequence listing part.

[5741] GAM397 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM397 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5742] An enzyme complex designated DICER COMPLEX, dices the GAM397 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM397 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 44%) nucleotide sequence of GAM397 RNA is designated SEQ ID:6136, and is provided hereinbelow with reference to the sequence listing part.

[5743] GAM397 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM397 target RNA, herein designated GAM TARGET RNA. GAM397 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5744] GAM397 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM397 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM397 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM397 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM397 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5745] The complementary binding of GAM397 RNA, herein designated GAM RNA, to target binding sites on GAM397 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM397 target RNA, herein designated GAM TARGET RNA, into GAM397 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5746] It is appreciated that GAM397 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM397 target genes. The mRNA of each one of this plurality of GAM397 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM397 RNA, herein designated GAM RNA, and which when bound by GAM397 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM397 target proteins.

[5747] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM397 gene, herein designated GAM GENE, on one or more GAM397 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[5748] It is yet further appreciated that specific functions, and accordingly utilities, of GAM397 correlate with, and may be deduced from, the identity of the target genes which GAM397 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5749] Nucleotide sequences of the GAM397 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM397 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM397 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM397 are further described hereinbelow with reference to Table Table1.

[5750] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM397 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5751]

[5752] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

398 (GAM398) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5753] GAM398 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM398 was detected is described hereinabove with reference to Figs. 2-8.

[5754] GAM398 gene, herein designated GAM GENE, and GAM398 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5755] GAM398 gene, herein designated GAM GENE, encodes a GAM398 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM398 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM398 precursor RNA is designated SEQ ID:6161, and is provided hereinbelow with reference to the sequence listing part.

[5756] GAM398 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM398 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5757] An enzyme complex designated DICER COMPLEX, dices the GAM398 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM398 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM398 RNA is designated SEQ ID:6162, and is provided hereinbelow with reference to the sequence listing part.

[5758] GAM398 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM398 target RNA, herein designated GAM TARGET RNA. GAM398 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5759] GAM398 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM398 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM398 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM398 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM398 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5760] The complementary binding of GAM398 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM398 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM398 target RNA, herein designated GAM TARGET RNA, into GAM398 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5761] It is appreciated that GAM398 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM398 target genes. The mRNA of each one of this plurality of GAM398 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM398 RNA, herein designated GAM RNA, and which when bound by GAM398 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM398 target proteins.

[5762] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM398 gene, herein designated GAM GENE, on one or more GAM398 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5763] It is yet further appreciated that specific functions, and accordingly utilities, of GAM398 correlate with, and may be deduced from, the identity of the target genes which GAM398 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5764] Nucleotide sequences of the GAM398 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM398 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM398 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM398 are further described hereinbelow with reference to Table Table1.

[5765] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM398 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5766]

[5767] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 399 (GAM399) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5768] GAM399 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM399 was detected is described hereinabove with reference to Figs. 2-8.

[5769] GAM399 gene, herein designated GAM GENE, and GAM399 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5770] GAM399 gene, herein designated GAM GENE, encodes a GAM399 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM399 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM399 precursor RNA is designated SEQ ID:6189, and is provided hereinbelow with reference to the sequence listing part.

[5771] GAM399 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM399 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5772] An enzyme complex designated DICER COMPLEX, dices the GAM399 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM399 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 58%) nucleotide sequence of GAM399 RNA is designated SEQ ID:6190, and is provided hereinbelow with reference to the sequence listing part.

[5773] GAM399 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM399 target RNA, herein designated GAM TARGET RNA. GAM399 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5774] GAM399 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM399 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM399 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM399 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM399 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5775] The complementary binding of GAM399 RNA, herein designated GAM RNA, to target binding sites on GAM399 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM399 target RNA, herein designated GAM TARGET RNA, into GAM399 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5776] It is appreciated that GAM399 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM399 target genes. The mRNA of each one of this plurality of GAM399 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM399 RNA, herein designated GAM RNA, and which when bound by

GAM399 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM399 target proteins.

[5777] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM399 gene, herein designated GAM GENE, on one or more GAM399 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5778] It is yet further appreciated that specific functions, and accordingly utilities, of GAM399 correlate with, and may be deduced from, the identity of the target genes which GAM399 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[5779] Nucleotide sequences of the GAM399 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM399 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM399 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM399 are further described hereinbelow with reference to Table Table1.

[5780] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM399 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5781]

[5782] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 400 (GAM400) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5783] GAM400 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM400 was detected is described hereinabove with reference to Figs. 2-8.

[5784] GAM400 gene, herein designated GAM GENE, and GAM400 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5785] GAM400 gene, herein designated GAM GENE, encodes a GAM400 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM400 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM400 precursor RNA is designated SEQ ID:6197, and is provided hereinbelow with reference to the sequence listing part.

[5786] GAM400 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM400 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[5787] An enzyme complex designated DICER COMPLEX, dices the GAM400 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM400 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM400 RNA is designated SEQ ID:6198, and is provided hereinbelow with reference to the sequence listing part.

[5788] GAM400 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM400 target RNA, herein designated GAM TARGET RNA. GAM400 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5789] GAM400 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM400 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM400 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM400 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM400 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5790] The complementary binding of GAM400 RNA, herein designated GAM RNA, to target binding sites on GAM400 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM400 target RNA, herein designated GAM TARGET RNA, into GAM400 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5791] It is appreciated that GAM400 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM400 target genes. The mRNA of each one of this plurality of GAM400 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM400 RNA, herein designated GAM RNA, and which when bound by GAM400 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM400 target proteins.

[5792] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM400 gene, herein designated GAM GENE, on one or more GAM400 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5793] It is yet further appreciated that specific functions, and accordingly utilities, of GAM400 correlate with, and may be deduced from, the identity of the target genes which GAM400 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5794] Nucleotide sequences of the GAM400 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM400 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM400 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM400 are further described hereinbelow with reference to Table Table1.

[5795] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM400 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5796]

[5797] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 401 (GAM401) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5798] GAM401 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM401 was detected is described hereinabove with reference to Figs. 2-8.

[5799] GAM401 gene, herein designated GAM GENE, and GAM401 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5800] GAM401 gene, herein designated GAM GENE, encodes a GAM401 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM401 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM401 precursor RNA is designated SEQ ID:6201, and is provided hereinbelow with reference to the sequence listing part.

[5801] GAM401 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM401 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5802] An enzyme complex designated DICER COMPLEX, dices the GAM401 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM401 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM401 RNA is designated SEQ ID:6202, and is provided hereinbelow with reference to the sequence listing part.

[5803] GAM401 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM401

target RNA, herein designated GAM TARGET RNA.

GAM401 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5804] GAM401 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM401 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM401 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM401 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM401 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5805] The complementary binding of GAM401 RNA, herein designated GAM RNA, to target binding sites on GAM401 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM401 target RNA, herein designated GAM TARGET RNA, into GAM401 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5806] It is appreciated that GAM401 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM401 target genes. The mRNA of each one of this plurality of GAM401 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM401 RNA, herein designated GAM RNA, and which when bound by GAM401 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM401 target proteins.

[5807] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM401 gene, herein designated GAM GENE, on one or more GAM401 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5808] It is yet further appreciated that specific functions, and accordingly utilities, of GAM401 correlate with, and may be deduced from, the identity of the target genes which GAM401 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5809] Nucleotide sequences of the GAM401 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM401 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM401 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM401 are further described hereinbelow with reference to Table Table1.

[5810] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM401 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5811]

[5812] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 402 (GAM402) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5813] GAM402 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM402 was detected is described hereinabove with reference to Figs. 2-8.

[5814] GAM402 gene, herein designated GAM GENE, and GAM402 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5815] GAM402 gene, herein designated GAM GENE, encodes a GAM402 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM402 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM402 precursor RNA is designated SEQ ID:6210, and is provided hereinbelow with reference to the sequence listing part.

[5816] GAM402 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM402 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5817] An enzyme complex designated DICER COMPLEX, dices the GAM402 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM402 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM402 RNA is designated SEQ ID:6211, and is provided hereinbelow with reference to the sequence listing part.

[5818] GAM402 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM402 target RNA, herein designated GAM TARGET RNA. GAM402 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5819] GAM402 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM402 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM402 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM402 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM402 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5820] The complementary binding of GAM402 RNA, herein designated GAM RNA, to target binding sites on GAM402 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM402 target RNA, herein designated GAM TARGET RNA, into GAM402 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5821] It is appreciated that GAM402 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM402 target genes. The mRNA of each one of this plu-

rality of GAM402 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM402 RNA, herein designated GAM RNA, and which when bound by GAM402 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM402 target proteins.

[5822] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM402 gene, herein designated GAM GENE, on one or more GAM402 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5823] It is yet further appreciated that specific functions, and accordingly utilities, of GAM402 correlate with, and may be deduced from, the identity of the target genes which GAM402 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5824] Nucleotide sequences of the GAM402 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM402 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM402 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM402 are further described hereinbelow with reference to Table Table1.

[5825] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM402 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5826]

[5827] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 403 (GAM403) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[5828] GAM403 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM403 was detected is described hereinabove with reference to Figs. 2-8.

[5829] GAM403 gene, herein designated GAM GENE, and GAM403 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5830] GAM403 gene, herein designated GAM GENE, encodes a GAM403 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM403 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM403 precursor RNA is designated SEQ ID:6262, and is provided hereinbelow with reference to the sequence listing part.

[5831] GAM403 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM403 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5832] An enzyme complex designated DICER COMPLEX, dices the GAM403 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM403 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 39%) nucleotide sequence of GAM403 RNA is designated SEQ ID:6263, and is provided hereinbelow with reference to the sequence listing part.

[5833] GAM403 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM403 target RNA, herein designated GAM TARGET RNA.
GAM403 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[5834] GAM403 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM403 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM403 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM403 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM403 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5835] The complementary binding of GAM403 RNA, herein designated GAM RNA, to target binding sites on GAM403 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM403 target RNA, herein designated GAM TARGET RNA, into GAM403 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5836] It is appreciated that GAM403 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM403 target genes. The mRNA of each one of this plurality of GAM403 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM403 RNA, herein designated GAM RNA, and which when bound by GAM403 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM403 target proteins.

[5837] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM403 gene, herein designated GAM GENE, on one or more GAM403 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5838] It is yet further appreciated that specific functions, and accordingly utilities, of GAM403 correlate with, and may be deduced from, the identity of the target genes which GAM403 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5839] Nucleotide sequences of the GAM403 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM403 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM403 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM403 are further described hereinbelow with reference to Table Table1.

[5840] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM403 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5841]

[5842] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 404 (GAM404) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5843] GAM404 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM404 was detected is described hereinabove with reference to Figs. 2-8.

[5844] GAM404 gene, herein designated GAM GENE, and GAM404 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5845] GAM404 gene, herein designated GAM GENE, encodes a GAM404 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM404 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM404 precursor RNA is designated SEQ ID:6289, and is provided hereinbelow with reference to the sequence listing part.

[5846] GAM404 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM404 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5847] An enzyme complex designated DICER COMPLEX, dices the GAM404 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM404 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of

GAM404 RNA is designated SEQ ID:6290, and is provided hereinbelow with reference to the sequence listing part.

[5848] GAM404 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM404 target RNA, herein designated GAM TARGET RNA. GAM404 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5849] GAM404 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM404 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM404 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM404 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM404 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5850] The complementary binding of GAM404 RNA, herein designated GAM RNA, to target binding sites on GAM404 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM404 target RNA, herein designated GAM TARGET RNA, into GAM404 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5851] It is appreciated that GAM404 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM404 target genes. The mRNA of each one of this plurality of GAM404 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM404 RNA, herein designated GAM RNA, and which when bound by GAM404 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM404 target proteins.

[5852] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM404 gene, herein designated GAM GENE, on one or more GAM404 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5853] It is yet further appreciated that specific functions, and accordingly utilities, of GAM404 correlate with, and may be deduced from, the identity of the target genes which GAM404 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5854] Nucleotide sequences of the GAM404 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM404 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM404 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM404 are further described hereinbelow with reference to Table Table1.

[5855] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM404 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5856]

[5857] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 405 (GAM405) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5858] GAM405 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM405 was detected is described hereinabove

with reference to Figs. 2-8.

[5859] GAM405 gene, herein designated GAM GENE, and GAM405 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5860] GAM405 gene, herein designated GAM GENE, encodes a GAM405 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM405 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM405 precursor RNA is designated SEQ ID:6304, and is provided hereinbelow with reference to the sequence listing part.

[5861] GAM405 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM405 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5862] An enzyme complex designated DICER COMPLEX, dices the GAM405 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM405 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 22%) nucleotide sequence of GAM405 RNA is designated SEQ ID:6305, and is provided hereinbelow with reference to the sequence listing part.

[5863] GAM405 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM405 target RNA, herein designated GAM TARGET RNA. GAM405 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5864] GAM405 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM405 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM405 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM405 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM405 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5865] The complementary binding of GAM405 RNA, herein designated GAM RNA, to target binding sites on GAM405 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM405 target RNA, herein designated GAM TARGET RNA, into GAM405 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[5866] It is appreciated that GAM405 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM405 target genes. The mRNA of each one of this plurality of GAM405 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM405 RNA, herein designated GAM RNA, and which when bound by GAM405 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM405 target proteins.

[5867] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM405 gene, herein designated GAM GENE, on one or more GAM405 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5868] It is yet further appreciated that specific functions, and accordingly utilities, of GAM405 correlate with, and may be deduced from, the identity of the target genes which GAM405 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5869] Nucleotide sequences of the GAM405 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM405 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM405 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM405 are further described hereinbelow with reference to Table Table1.

[5870] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM405 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5871]

[5872] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 406 (GAM406) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5873] GAM406 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM406 was detected is described hereinabove with reference to Figs. 2-8.

[5874] GAM406 gene, herein designated GAM GENE, and GAM406 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5875] GAM406 gene, herein designated GAM GENE, encodes a GAM406 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM406 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM406 precursor RNA is designated SEQ ID:6311, and is provided hereinbelow with reference to the sequence listing part.

[5876] GAM406 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM406 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5877] An enzyme complex designated DICER COMPLEX, dices the GAM406 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM406 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM406 RNA is designated SEQ ID:6312, and is provided hereinbelow with reference to the sequence listing part.

[5878] GAM406 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM406 target RNA, herein designated GAM TARGET RNA.

GAM406 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5879] GAM406 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM406 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM406 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM406 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM406 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5880] The complementary binding of GAM406 RNA, herein designated GAM RNA, to target binding sites on GAM406 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM406 target RNA, herein designated GAM TARGET RNA, into GAM406 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5881] It is appreciated that GAM406 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM406 target genes. The mRNA of each one of this plurality of GAM406 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM406 RNA, herein designated GAM RNA, and which when bound by GAM406 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM406 target proteins.

[5882] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM406 gene, herein designated GAM GENE, on one or more GAM406 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5883] It is yet further appreciated that specific functions, and accordingly utilities, of GAM406 correlate with, and may be deduced from, the identity of the target genes which GAM406 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5884] Nucleotide sequences of the GAM406 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM406 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM406 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM406 are further described hereinbelow with reference to Table Table1.

[5885] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM406 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5886]

[5887] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 407 (GAM407) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5888] GAM407 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM407 was detected is described hereinabove with reference to Figs. 2-8.

[5889] GAM407 gene, herein designated GAM GENE, and GAM407 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5890] GAM407 gene, herein designated GAM GENE, encodes a

GAM407 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM407 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM407 precursor RNA is designated SEQ ID:6329, and is provided hereinbelow with reference to the sequence listing part.

[5891] GAM407 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM407 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5892] An enzyme complex designated DICER COMPLEX, dices the GAM407 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM407 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM407 RNA is designated SEQ ID:6330, and is provided hereinbelow with reference to the sequence listing part.

[5893] GAM407 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM407 target RNA, herein designated GAM TARGET RNA. GAM407 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5894] GAM407 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM407 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM407 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM407 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM407 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5895] The complementary binding of GAM407 RNA, herein designated GAM RNA, to target binding sites on GAM407 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM407 target RNA, herein designated GAM TARGET RNA, into GAM407 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5896] It is appreciated that GAM407 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM407 target genes. The mRNA of each one of this plurality of GAM407 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM407 RNA, herein designated GAM RNA, and which when bound by GAM407 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM407 target proteins.

[5897] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM407 gene, herein designated GAM GENE, on one or more GAM407 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5898] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM407 correlate with, and may be deduced from, the identity of the target genes which GAM407 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5899] Nucleotide sequences of the GAM407 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM407 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM407 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM407 are further described hereinbelow with reference to Table Table1.

[5900] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM407 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5901]

[5902] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 408 (GAM408) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[5903] GAM408 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM408 was detected is described hereinabove with reference to Figs. 2-8.

[5904] GAM408 gene, herein designated GAM GENE, and GAM408 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5905] GAM408 gene, herein designated GAM GENE, encodes a GAM408 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM408 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM408 precursor RNA is designated SEQ ID:6357, and is provided hereinbelow with reference to the sequence listing part.

[5906] GAM408 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM408 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5907] An enzyme complex designated DICER COMPLEX, dices the GAM408 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM408 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM408 RNA is designated SEQ ID:6358, and is provided hereinbelow with reference to the sequence listing part.

[5908] GAM408 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM408 target RNA, herein designated GAM TARGET RNA. GAM408 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5909] GAM408 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM408 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM408 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM408 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM408 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5910] The complementary binding of GAM408 RNA, herein designated GAM RNA, to target binding sites on GAM408 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM408 target RNA, herein designated GAM TARGET RNA, into GAM408 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5911] It is appreciated that GAM408 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM408 target genes. The mRNA of each one of this plurality of GAM408 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM408 RNA, herein designated GAM RNA, and which when bound by GAM408 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM408 target proteins.

[5912] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM408 gene, herein designated GAM GENE, on one or more GAM408 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5913] It is yet further appreciated that specific functions, and accordingly utilities, of GAM408 correlate with, and may be deduced from, the identity of the target genes which GAM408 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5914] Nucleotide sequences of the GAM408 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM408 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM408 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM408 are further described hereinbelow with reference to Table Table1.

[5915] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM408 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5916]

[5917] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 409 (GAM409) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5918] GAM409 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM409 was detected is described hereinabove with reference to Figs. 2-8.

[5919] GAM409 gene, herein designated GAM GENE, and GAM409 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5920] GAM409 gene, herein designated GAM GENE, encodes a GAM409 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM409 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM409 precursor RNA is designated SEQ ID:6372, and is provided hereinbelow with reference to the sequence listing part.

[5921] GAM409 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM409 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5922] An enzyme complex designated DICER COMPLEX, dices the GAM409 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM409 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM409 RNA is designated SEQ ID:6373, and is provided

hereinbelow with reference to the sequence listing part.

[5923] GAM409 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM409 target RNA, herein designated GAM TARGET RNA.
GAM409 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5924] GAM409 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM409 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM409 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM409 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM409 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5925] The complementary binding of GAM409 RNA, herein designated GAM RNA, to target binding sites on GAM409 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM409 target RNA, herein designated GAM TARGET RNA, into GAM409 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5926] It is appreciated that GAM409 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM409 target genes. The mRNA of each one of this plurality of GAM409 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM409 RNA, herein designated GAM RNA, and which when bound by GAM409 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM409 tar-

get proteins.

[5927] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM409 gene, herein designated GAM GENE, on one or more GAM409 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5928] It is yet further appreciated that specific functions, and accordingly utilities, of GAM409 correlate with, and may be deduced from, the identity of the target genes which GAM409 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5929] Nucleotide sequences of the GAM409 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM409 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM409 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM409 are further described hereinbelow with reference to Table Table1.

[5930] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM409 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5931]

[5932] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 410 (GAM410) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5933] GAM410 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM410 was detected is described hereinabove with reference to Figs. 2-8.

[5934] GAM410 gene, herein designated GAM GENE, and GAM410 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5935] GAM410 gene, herein designated GAM GENE, encodes a GAM410 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM410 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM410 precursor RNA is designated SEQ ID:6380, and is provided hereinbelow with reference to the sequence listing part.

[5936] GAM410 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM410 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5937] An enzyme complex designated DICER COMPLEX, dices

the GAM410 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM410 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM410 RNA is designated SEQ ID:6381, and is provided hereinbelow with reference to the sequence listing part.

[5938] GAM410 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM410 target RNA, herein designated GAM TARGET RNA. GAM410 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5939] GAM410 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM410 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM410

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM410 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM410 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5940] The complementary binding of GAM410 RNA, herein designated GAM RNA, to target binding sites on GAM410 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM410 target RNA, herein designated GAM TARGET RNA, into GAM410 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5941] It is appreciated that GAM410 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM410 target genes. The mRNA of each one of this plurality of GAM410 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM410 RNA, herein designated GAM RNA, and which when bound by GAM410 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM410 target proteins.

[5942] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM410 gene, herein designated GAM GENE, on one or more GAM410 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5943] It is yet further appreciated that specific functions, and accordingly utilities, of GAM410 correlate with, and may be deduced from, the identity of the target genes which GAM410 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5944] Nucleotide sequences of the GAM410 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM410 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM410 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM410 are further described hereinbelow with reference to Table Table1.

[5945] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM410 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5946]

[5947] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 411 (GAM411) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5948] GAM411 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM411 was detected is described hereinabove with reference to Figs. 2-8.

[5949] GAM411 gene, herein designated GAM GENE, and GAM411 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5950] GAM411 gene, herein designated GAM GENE, encodes a GAM411 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM411 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM411 precursor RNA is designated SEQ ID:6419, and is provided hereinbelow with reference to the sequence listing part.

[5951] GAM411 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM411 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5952] An enzyme complex designated DICER COMPLEX, dices the GAM411 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM411 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM411 RNA is designated SEQ ID:6420, and is provided hereinbelow with reference to the sequence listing part.

[5953] GAM411 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM411 target RNA, herein designated GAM TARGET RNA. GAM411 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5954] GAM411 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM411 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM411 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM411 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM411 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[5955] The complementary binding of GAM411 RNA, herein designated GAM RNA, to target binding sites on GAM411 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM411 target RNA, herein designated GAM TARGET RNA, into GAM411 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5956] It is appreciated that GAM411 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM411 target genes. The mRNA of each one of this plurality of GAM411 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM411 RNA, herein designated GAM RNA, and which when bound by GAM411 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM411 target proteins.

[5957] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM411 gene, herein designated GAM GENE, on one or

more GAM411 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5958] It is yet further appreciated that specific functions, and accordingly utilities, of GAM411 correlate with, and may be deduced from, the identity of the target genes which GAM411 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5959] Nucleotide sequences of the GAM411 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM411 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM411 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM411 are further de-

scribed hereinbelow with reference to Table Table1.

[5960] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM411 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5961]

[5962] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 412 (GAM412) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5963] GAM412 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM412 was detected is described hereinabove with reference to Figs. 2-8.

[5964] GAM412 gene, herein designated GAM GENE, and GAM412 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5965] GAM412 gene, herein designated GAM GENE, encodes a GAM412 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM412 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM412 precursor RNA is designated SEQ ID:6436, and is provided hereinbelow with reference to the sequence listing part.

[5966] GAM412 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM412 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5967] An enzyme complex designated DICER COMPLEX, dices the GAM412 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM412 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM412 RNA is designated SEQ ID:6437, and is provided hereinbelow with reference to the sequence listing part.

[5968] GAM412 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM412 target RNA, herein designated GAM TARGET RNA. GAM412 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5969] GAM412 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM412 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM412 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM412 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM412 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5970] The complementary binding of GAM412 RNA, herein designated GAM RNA, to target binding sites on GAM412 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM412 target RNA, herein designated GAM TARGET RNA, into GAM412 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5971] It is appreciated that GAM412 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM412 target genes. The mRNA of each one of this plurality of GAM412 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM412 RNA, herein designated GAM RNA, and which when bound by GAM412 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM412 target proteins.

[5972] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM412 gene, herein designated GAM GENE, on one or more GAM412 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5973] It is yet further appreciated that specific functions, and accordingly utilities, of GAM412 correlate with, and may

be deduced from, the identity of the target genes which GAM412 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5974] Nucleotide sequences of the GAM412 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM412 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM412 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM412 are further described hereinbelow with reference to Table Table1.

[5975] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM412 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5976]

[5977] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 413 (GAM413) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5978] GAM413 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM413 was detected is described hereinabove with reference to Figs. 2-8.

[5979] GAM413 gene, herein designated GAM GENE, and GAM413 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5980] GAM413 gene, herein designated GAM GENE, encodes a GAM413 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM413 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM413 precursor RNA is designated SEQ ID:6452, and is provided hereinbelow with reference to the sequence listing part.

[5981] GAM413 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM413 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5982] An enzyme complex designated DICER COMPLEX, dices the GAM413 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM413 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 41%) nucleotide sequence of GAM413 RNA is designated SEQ ID:6453, and is provided hereinbelow with reference to the sequence listing part.

[5983] GAM413 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM413 target RNA, herein designated GAM TARGET RNA. GAM413 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5984] GAM413 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM413 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM413 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM413 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM413 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[5985] The complementary binding of GAM413 RNA, herein designated GAM RNA, to target binding sites on GAM413 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM413 target RNA, herein designated GAM TARGET RNA, into GAM413 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[5986] It is appreciated that GAM413 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM413 target genes. The mRNA of each one of this plurality of GAM413 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM413 RNA, herein designated GAM RNA, and which when bound by GAM413 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM413 target proteins.

[5987] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM413 gene, herein designated GAM GENE, on one or more GAM413 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[5988] It is yet further appreciated that specific functions, and accordingly utilities, of GAM413 correlate with, and may be deduced from, the identity of the target genes which GAM413 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[5989] Nucleotide sequences of the GAM413 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM413 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM413 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM413 are further described hereinbelow with reference to Table Table1.

[5990] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM413 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[5991]

[5992] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 414 (GAM414) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[5993] GAM414 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM414 was detected is described hereinabove with reference to Figs. 2-8.

[5994] GAM414 gene, herein designated GAM GENE, and GAM414 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[5995] GAM414 gene, herein designated GAM GENE, encodes a GAM414 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM414 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM414 precursor RNA is desig-

nated SEQ ID:6474, and is provided hereinbelow with reference to the sequence listing part.

[5996] GAM414 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM414 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[5997] An enzyme complex designated DICER COMPLEX, dices the GAM414 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM414 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM414 RNA is designated SEQ ID:6475, and is provided hereinbelow with reference to the sequence listing part.

[5998] GAM414 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM414 target RNA, herein designated GAM TARGET RNA. GAM414 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[5999] GAM414 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM414 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM414 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM414 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM414

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6000] The complementary binding of GAM414 RNA, herein designated GAM RNA, to target binding sites on GAM414 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM414 target RNA, herein designated GAM TARGET RNA, into GAM414 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6001] It is appreciated that GAM414 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM414 target genes. The mRNA of each one of this plurality of GAM414 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM414 RNA, herein designated GAM RNA, and which when bound by GAM414 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM414 target proteins.

[6002] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM414 gene, herein designated GAM GENE, on one or more GAM414 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6003] It is yet further appreciated that specific functions, and accordingly utilities, of GAM414 correlate with, and may be deduced from, the identity of the target genes which GAM414 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6004] Nucleotide sequences of the GAM414 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM414 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM414 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM414 are further described hereinbelow with reference to Table Table1.

[6005] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM414 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6006]

[6007] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 415 (GAM415) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6008] GAM415 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM415 was detected is described hereinabove with reference to Figs. 2-8.

[6009] GAM415 gene, herein designated GAM GENE, and

GAM415 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6010] GAM415 gene, herein designated GAM GENE, encodes a GAM415 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM415 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM415 precursor RNA is designated SEQ ID:6489, and is provided hereinbelow with reference to the sequence listing part.

[6011] GAM415 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM415 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6012] An enzyme complex designated DICER COMPLEX, dices the GAM415 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM415 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM415 RNA is designated SEQ ID:6490, and is provided hereinbelow with reference to the sequence listing part.

[6013] GAM415 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM415 target RNA, herein designated GAM TARGET RNA. GAM415 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6014] GAM415 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM415 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM415 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM415 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM415 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6015] The complementary binding of GAM415 RNA, herein designated GAM RNA, to target binding sites on GAM415 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM415 target RNA, herein designated GAM TARGET RNA, into GAM415 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6016] It is appreciated that GAM415 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM415 target genes. The mRNA of each one of this plurality of GAM415 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM415 RNA, herein designated GAM RNA, and which when bound by GAM415 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM415 target proteins.

[6017] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM415 gene, herein designated GAM GENE, on one or more GAM415 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6018] It is yet further appreciated that specific functions, and accordingly utilities, of GAM415 correlate with, and may be deduced from, the identity of the target genes which GAM415 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6019] Nucleotide sequences of the GAM415 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM415 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM415 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM415 are further described hereinbelow with reference to Table Table1.

[6020] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM415 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6021]

[6022] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 416 (GAM416) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6023] GAM416 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM416 was detected is described hereinabove with reference to Figs. 2-8.

[6024] GAM416 gene, herein designated GAM GENE, and GAM416 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6025] GAM416 gene, herein designated GAM GENE, encodes a GAM416 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM416 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM416 precursor RNA is designated SEQ ID:6500, and is provided hereinbelow with reference to the sequence listing part.

[6026] GAM416 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM416 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6027] An enzyme complex designated DICER COMPLEX, dices the GAM416 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM416 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM416 RNA is designated SEQ ID:6501, and is provided hereinbelow with reference to the sequence listing part.

[6028] GAM416 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM416 target RNA, herein designated GAM TARGET RNA. GAM416 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6029] GAM416 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM416 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM416 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM416 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM416 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6030] The complementary binding of GAM416 RNA, herein designated GAM RNA, to target binding sites on GAM416 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM416 target RNA, herein designated GAM TARGET RNA, into GAM416 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6031] It is appreciated that GAM416 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM416 target genes. The mRNA of each one of this plurality of GAM416 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM416 RNA, herein designated GAM RNA, and which when bound by GAM416 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM416 target proteins.

[6032] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM416 gene, herein designated GAM GENE, on one or more GAM416 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6033] It is yet further appreciated that specific functions, and accordingly utilities, of GAM416 correlate with, and may be deduced from, the identity of the target genes which GAM416 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6034] Nucleotide sequences of the GAM416 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM416 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM416 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM416 are further described hereinbelow with reference to Table Table1.

[6035] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM416 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6036]

[6037] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 417 (GAM417) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6038] GAM417 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM417 was detected is described hereinabove with reference to Figs. 2-8.

[6039] GAM417 gene, herein designated GAM GENE, and GAM417 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6040] GAM417 gene, herein designated GAM GENE, encodes a GAM417 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM417 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM417 precursor RNA is designated SEQ ID:6514, and is provided hereinbelow with reference to the sequence listing part.

[6041] GAM417 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM417 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6042] An enzyme complex designated DICER COMPLEX, dices the GAM417 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM417 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM417 RNA is designated SEQ ID:6515, and is provided hereinbelow with reference to the sequence listing part.

[6043] GAM417 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM417 target RNA, herein designated GAM TARGET RNA. GAM417 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6044] GAM417 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM417 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM417 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM417 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM417 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6045] The complementary binding of GAM417 RNA, herein designated GAM RNA, to target binding sites on GAM417 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM417 target RNA, herein designated GAM TARGET RNA, into GAM417 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6046] It is appreciated that GAM417 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM417 target genes. The mRNA of each one of this plurality of GAM417 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM417 RNA,

herein designated GAM RNA, and which when bound by GAM417 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM417 target proteins.

[6047] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM417 gene, herein designated GAM GENE, on one or more GAM417 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6048] It is yet further appreciated that specific functions, and accordingly utilities, of GAM417 correlate with, and may be deduced from, the identity of the target genes which

GAM417 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6049] Nucleotide sequences of the GAM417 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM417 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM417 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM417 are further described hereinbelow with reference to Table Table1.

[6050] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM417 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6051]

[6052] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 418 (GAM418) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6053] GAM418 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM418 was detected is described hereinabove with reference to Figs. 2-8.

[6054] GAM418 gene, herein designated GAM GENE, and GAM418 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6055] GAM418 gene, herein designated GAM GENE, encodes a GAM418 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM418 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM418 precursor RNA is designated SEQ ID:6529, and is provided hereinbelow with reference to the sequence listing part.

[6056] GAM418 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM418 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[6057] An enzyme complex designated DICER COMPLEX, dices the GAM418 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM418 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM418 RNA is designated SEQ ID:6530, and is provided hereinbelow with reference to the sequence listing part.

[6058] GAM418 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM418 target RNA, herein designated GAM TARGET RNA. GAM418 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6059] GAM418 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM418 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM418 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM418 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM418 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6060] The complementary binding of GAM418 RNA, herein designated GAM RNA, to target binding sites on GAM418 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM418 target RNA, herein desig-

nated GAM TARGET RNA, into GAM418 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6061] It is appreciated that GAM418 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM418 target genes. The mRNA of each one of this plurality of GAM418 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM418 RNA, herein designated GAM RNA, and which when bound by GAM418 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM418 target proteins.

[6062] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM418 gene, herein designated GAM GENE, on one or more GAM418 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6063] It is yet further appreciated that specific functions, and accordingly utilities, of GAM418 correlate with, and may be deduced from, the identity of the target genes which GAM418 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6064] Nucleotide sequences of the GAM418 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM418 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM418 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM418 are further described hereinbelow with reference to Table Table1.

[6065] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM418 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[6066]

[6067] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 419 (GAM419) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6068] GAM419 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM419 was detected is described hereinabove with reference to Figs. 2-8.

[6069] GAM419 gene, herein designated GAM GENE, and GAM419 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6070] GAM419 gene, herein designated GAM GENE, encodes a GAM419 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM419 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM419 precursor RNA is designated SEQ ID:6547, and is provided hereinbelow with ref-

erence to the sequence listing part.

[6071] GAM419 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM419 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6072] An enzyme complex designated DICER COMPLEX, dices the GAM419 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM419 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM419 RNA is designated SEQ ID:6548, and is provided hereinbelow with reference to the sequence listing part.

[6073] GAM419 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM419 target RNA, herein designated GAM TARGET RNA.

GAM419 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6074] GAM419 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM419 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM419 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM419 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM419 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6075] The complementary binding of GAM419 RNA, herein designated GAM RNA, to target binding sites on GAM419 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM419 target RNA, herein designated GAM TARGET RNA, into GAM419 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6076] It is appreciated that GAM419 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM419 target genes. The mRNA of each one of this plurality of GAM419 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM419 RNA, herein designated GAM RNA, and which when bound by GAM419 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM419 target proteins.

[6077] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM419 gene, herein designated GAM GENE, on one or more GAM419 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6078] It is yet further appreciated that specific functions, and accordingly utilities, of GAM419 correlate with, and may be deduced from, the identity of the target genes which GAM419 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6079] Nucleotide sequences of the GAM419 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM419 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM419 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM419 are further described hereinbelow with reference to Table Table1.

[6080] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM419 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6081]

[6082] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 420 (GAM420) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6083] GAM420 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM420 was detected is described hereinabove with reference to Figs. 2-8.

[6084] GAM420 gene, herein designated GAM GENE, and GAM420 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[6085] GAM420 gene, herein designated GAM GENE, encodes a GAM420 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM420 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM420 precursor RNA is designated SEQ ID:6550, and is provided hereinbelow with reference to the sequence listing part.

[6086] GAM420 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM420 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6087] An enzyme complex designated DICER COMPLEX, dices the GAM420 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM420 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM420 RNA is designated SEQ ID:6551, and is provided hereinbelow with reference to the sequence listing part.

[6088] GAM420 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM420 target RNA, herein designated GAM TARGET RNA. GAM420 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6089] GAM420 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM420 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM420 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM420 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM420 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6090] The complementary binding of GAM420 RNA, herein designated GAM RNA, to target binding sites on GAM420 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM420 target RNA, herein designated GAM TARGET RNA, into GAM420 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6091] It is appreciated that GAM420 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM420 target genes. The mRNA of each one of this plurality of GAM420 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM420 RNA, herein designated GAM RNA, and which when bound by GAM420 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM420 target proteins.

[6092] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM420 gene, herein designated GAM GENE, on one or more GAM420 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[6093] It is yet further appreciated that specific functions, and accordingly utilities, of GAM420 correlate with, and may be deduced from, the identity of the target genes which GAM420 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6094] Nucleotide sequences of the GAM420 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM420 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM420 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM420 are further described hereinbelow with reference to Table Table1.

[6095] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM420 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6096]

[6097] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

421 (GAM421) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6098] GAM421 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM421 was detected is described hereinabove with reference to Figs. 2-8.

[6099] GAM421 gene, herein designated GAM GENE, and GAM421 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6100] GAM421 gene, herein designated GAM GENE, encodes a GAM421 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM421 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM421 precursor RNA is designated SEQ ID:6561, and is provided hereinbelow with reference to the sequence listing part.

[6101] GAM421 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM421 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6102] An enzyme complex designated DICER COMPLEX, dices the GAM421 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM421 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 70%) nucleotide sequence of GAM421 RNA is designated SEQ ID:6562, and is provided hereinbelow with reference to the sequence listing part.

[6103] GAM421 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM421 target RNA, herein designated GAM TARGET RNA. GAM421 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6104] GAM421 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM421 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM421 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM421 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM421 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6105] The complementary binding of GAM421 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM421 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM421 target RNA, herein designated GAM TARGET RNA, into GAM421 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6106] It is appreciated that GAM421 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM421 target genes. The mRNA of each one of this plurality of GAM421 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM421 RNA, herein designated GAM RNA, and which when bound by GAM421 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM421 target proteins.

[6107] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM421 gene, herein designated GAM GENE, on one or more GAM421 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6108] It is yet further appreciated that specific functions, and accordingly utilities, of GAM421 correlate with, and may be deduced from, the identity of the target genes which GAM421 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6109] Nucleotide sequences of the GAM421 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM421 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM421 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM421 are further described hereinbelow with reference to Table Table1.

[6110] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM421 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6111]

[6112] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 422 (GAM422) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6113] GAM422 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM422 was detected is described hereinabove with reference to Figs. 2-8.

[6114] GAM422 gene, herein designated GAM GENE, and GAM422 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6115] GAM422 gene, herein designated GAM GENE, encodes a GAM422 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM422 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM422 precursor RNA is designated SEQ ID:6593, and is provided hereinbelow with reference to the sequence listing part.

[6116] GAM422 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM422 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6117] An enzyme complex designated DICER COMPLEX, dices the GAM422 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM422 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 60%) nucleotide sequence of GAM422 RNA is designated SEQ ID:6594, and is provided hereinbelow with reference to the sequence listing part.

[6118] GAM422 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM422 target RNA, herein designated GAM TARGET RNA. GAM422 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6119] GAM422 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM422 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM422 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM422 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM422 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6120] The complementary binding of GAM422 RNA, herein designated GAM RNA, to target binding sites on GAM422 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM422 target RNA, herein designated GAM TARGET RNA, into GAM422 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6121] It is appreciated that GAM422 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM422 target genes. The mRNA of each one of this plurality of GAM422 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM422 RNA, herein designated GAM RNA, and which when bound by

GAM422 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM422 target proteins.

[6122] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM422 gene, herein designated GAM GENE, on one or more GAM422 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6123] It is yet further appreciated that specific functions, and accordingly utilities, of GAM422 correlate with, and may be deduced from, the identity of the target genes which GAM422 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[6124] Nucleotide sequences of the GAM422 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM422 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM422 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM422 are further described hereinbelow with reference to Table Table1.

[6125] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM422 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6126]

[6127] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 423 (GAM423) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6128] GAM423 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM423 was detected is described hereinabove with reference to Figs. 2-8.

[6129] GAM423 gene, herein designated GAM GENE, and GAM423 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6130] GAM423 gene, herein designated GAM GENE, encodes a GAM423 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM423 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM423 precursor RNA is designated SEQ ID:6611, and is provided hereinbelow with reference to the sequence listing part.

[6131] GAM423 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM423 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6132] An enzyme complex designated DICER COMPLEX, dices the GAM423 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM423 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM423 RNA is designated SEQ ID:6612, and is provided hereinbelow with reference to the sequence listing part.

[6133] GAM423 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM423 target RNA, herein designated GAM TARGET RNA. GAM423 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6134] GAM423 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM423 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM423 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM423 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM423 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6135] The complementary binding of GAM423 RNA, herein designated GAM RNA, to target binding sites on GAM423 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM423 target RNA, herein designated GAM TARGET RNA, into GAM423 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6136] It is appreciated that GAM423 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM423 target genes. The mRNA of each one of this plu-

rality of GAM423 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM423 RNA, herein designated GAM RNA, and which when bound by GAM423 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM423 target proteins.

[6137] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM423 gene, herein designated GAM GENE, on one or more GAM423 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6138] It is yet further appreciated that specific functions, and accordingly utilities, of GAM423 correlate with, and may be deduced from, the identity of the target genes which GAM423 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6139] Nucleotide sequences of the GAM423 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM423 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM423 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM423 are further described hereinbelow with reference to Table Table1.

[6140] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM423 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6141]

[6142] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 424 (GAM424) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[6143] GAM424 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM424 was detected is described hereinabove with reference to Figs. 2-8.

[6144] GAM424 gene, herein designated GAM GENE, and GAM424 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6145] GAM424 gene, herein designated GAM GENE, encodes a GAM424 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM424 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM424 precursor RNA is designated SEQ ID:6623, and is provided hereinbelow with reference to the sequence listing part.

[6146] GAM424 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM424 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6147] An enzyme complex designated DICER COMPLEX, dices the GAM424 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM424 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 20%) nucleotide sequence of GAM424 RNA is designated SEQ ID:6624, and is provided hereinbelow with reference to the sequence listing part.

[6148] GAM424 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM424 target RNA, herein designated GAM TARGET RNA.
GAM424 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[6149] GAM424 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM424 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM424 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM424 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM424 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6150] The complementary binding of GAM424 RNA, herein designated GAM RNA, to target binding sites on GAM424 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM424 target RNA, herein designated GAM TARGET RNA, into GAM424 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6151] It is appreciated that GAM424 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM424 target genes. The mRNA of each one of this plurality of GAM424 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM424 RNA, herein designated GAM RNA, and which when bound by GAM424 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM424 target proteins.

[6152] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM424 gene, herein designated GAM GENE, on one or more GAM424 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6153] It is yet further appreciated that specific functions, and accordingly utilities, of GAM424 correlate with, and may be deduced from, the identity of the target genes which GAM424 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6154] Nucleotide sequences of the GAM424 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM424 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM424 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM424 are further described hereinbelow with reference to Table Table1.

[6155] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM424 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6156]

[6157] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 425 (GAM425) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6158] GAM425 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM425 was detected is described hereinabove with reference to Figs. 2-8.

[6159] GAM425 gene, herein designated GAM GENE, and GAM425 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6160] GAM425 gene, herein designated GAM GENE, encodes a GAM425 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM425 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM425 precursor RNA is designated SEQ ID:6673, and is provided hereinbelow with reference to the sequence listing part.

[6161] GAM425 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM425 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6162] An enzyme complex designated DICER COMPLEX, dices the GAM425 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM425 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of

GAM425 RNA is designated SEQ ID:6674, and is provided hereinbelow with reference to the sequence listing part.

[6163] GAM425 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM425 target RNA, herein designated GAM TARGET RNA. GAM425 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6164] GAM425 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM425 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM425 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM425 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM425 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6165] The complementary binding of GAM425 RNA, herein designated GAM RNA, to target binding sites on GAM425 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM425 target RNA, herein designated GAM TARGET RNA, into GAM425 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6166] It is appreciated that GAM425 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM425 target genes. The mRNA of each one of this plurality of GAM425 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM425 RNA, herein designated GAM RNA, and which when bound by GAM425 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM425 target proteins.

[6167] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM425 gene, herein designated GAM GENE, on one or more GAM425 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6168] It is yet further appreciated that specific functions, and accordingly utilities, of GAM425 correlate with, and may be deduced from, the identity of the target genes which GAM425 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6169] Nucleotide sequences of the GAM425 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM425 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM425 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM425 are further described hereinbelow with reference to Table Table1.

[6170] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM425 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6171]

[6172] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 426 (GAM426) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6173] GAM426 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM426 was detected is described hereinabove

with reference to Figs. 2-8.

[6174] GAM426 gene, herein designated GAM GENE, and GAM426 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6175] GAM426 gene, herein designated GAM GENE, encodes a GAM426 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM426 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM426 precursor RNA is designated SEQ ID:6679, and is provided hereinbelow with reference to the sequence listing part.

[6176] GAM426 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM426 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6177] An enzyme complex designated DICER COMPLEX, dices the GAM426 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM426 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 50%) nucleotide sequence of GAM426 RNA is designated SEQ ID:6680, and is provided hereinbelow with reference to the sequence listing part.

[6178] GAM426 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM426 target RNA, herein designated GAM TARGET RNA. GAM426 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6179] GAM426 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM426 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM426 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM426 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM426 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6180] The complementary binding of GAM426 RNA, herein designated GAM RNA, to target binding sites on GAM426 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM426 target RNA, herein designated GAM TARGET RNA, into GAM426 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[6181] It is appreciated that GAM426 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM426 target genes. The mRNA of each one of this plurality of GAM426 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM426 RNA, herein designated GAM RNA, and which when bound by GAM426 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM426 target proteins.

[6182] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM426 gene, herein designated GAM GENE, on one or more GAM426 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6183] It is yet further appreciated that specific functions, and accordingly utilities, of GAM426 correlate with, and may be deduced from, the identity of the target genes which GAM426 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6184] Nucleotide sequences of the GAM426 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM426 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM426 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM426 are further described hereinbelow with reference to Table Table1.

[6185] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM426 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6186]

[6187] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 427 (GAM427) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6188] GAM427 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM427 was detected is described hereinabove with reference to Figs. 2-8.

[6189] GAM427 gene, herein designated GAM GENE, and GAM427 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6190] GAM427 gene, herein designated GAM GENE, encodes a GAM427 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM427 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM427 precursor RNA is designated SEQ ID:6693, and is provided hereinbelow with reference to the sequence listing part.

[6191] GAM427 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM427 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6192] An enzyme complex designated DICER COMPLEX, dices the GAM427 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM427 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM427 RNA is designated SEQ ID:6694, and is provided hereinbelow with reference to the sequence listing part.

[6193] GAM427 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM427 target RNA, herein designated GAM TARGET RNA.

GAM427 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6194] GAM427 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM427 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM427 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM427 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM427 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6195] The complementary binding of GAM427 RNA, herein designated GAM RNA, to target binding sites on GAM427 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM427 target RNA, herein designated GAM TARGET RNA, into GAM427 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6196] It is appreciated that GAM427 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM427 target genes. The mRNA of each one of this plurality of GAM427 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM427 RNA, herein designated GAM RNA, and which when bound by GAM427 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM427 target proteins.

[6197] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM427 gene, herein designated GAM GENE, on one or more GAM427 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6198] It is yet further appreciated that specific functions, and accordingly utilities, of GAM427 correlate with, and may be deduced from, the identity of the target genes which GAM427 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6199] Nucleotide sequences of the GAM427 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM427 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM427 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM427 are further described hereinbelow with reference to Table Table1.

[6200] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM427 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6201]

[6202] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 428 (GAM428) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6203] GAM428 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM428 was detected is described hereinabove with reference to Figs. 2-8.

[6204] GAM428 gene, herein designated GAM GENE, and GAM428 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6205] GAM428 gene, herein designated GAM GENE, encodes a

GAM428 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM428 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM428 precursor RNA is designated SEQ ID:6716, and is provided hereinbelow with reference to the sequence listing part.

[6206] GAM428 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM428 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6207] An enzyme complex designated DICER COMPLEX, dices the GAM428 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM428 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 73%) nucleotide sequence of GAM428 RNA is designated SEQ ID:6717, and is provided hereinbelow with reference to the sequence listing part.

[6208] GAM428 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM428 target RNA, herein designated GAM TARGET RNA. GAM428 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6209] GAM428 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM428 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM428 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM428 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM428 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6210] The complementary binding of GAM428 RNA, herein designated GAM RNA, to target binding sites on GAM428 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM428 target RNA, herein designated GAM TARGET RNA, into GAM428 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6211] It is appreciated that GAM428 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM428 target genes. The mRNA of each one of this plurality of GAM428 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM428 RNA, herein designated GAM RNA, and which when bound by GAM428 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM428 target proteins.

[6212] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM428 gene, herein designated GAM GENE, on one or more GAM428 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6213] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM428 correlate with, and may be deduced from, the identity of the target genes which GAM428 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6214] Nucleotide sequences of the GAM428 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM428 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM428 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM428 are further described hereinbelow with reference to Table Table1.

[6215] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM428 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6216]

[6217] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 429 (GAM429) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[6218] GAM429 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM429 was detected is described hereinabove with reference to Figs. 2-8.

[6219] GAM429 gene, herein designated GAM GENE, and GAM429 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6220] GAM429 gene, herein designated GAM GENE, encodes a GAM429 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM429 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM429 precursor RNA is designated SEQ ID:6729, and is provided hereinbelow with reference to the sequence listing part.

[6221] GAM429 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM429 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6222] An enzyme complex designated DICER COMPLEX, dices the GAM429 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM429 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM429 RNA is designated SEQ ID:6730, and is provided hereinbelow with reference to the sequence listing part.

[6223] GAM429 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM429 target RNA, herein designated GAM TARGET RNA.
GAM429 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6224] GAM429 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM429 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM429 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM429 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM429 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6225] The complementary binding of GAM429 RNA, herein designated GAM RNA, to target binding sites on GAM429 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM429 target RNA, herein designated GAM TARGET RNA, into GAM429 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6226] It is appreciated that GAM429 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM429 target genes. The mRNA of each one of this plurality of GAM429 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM429 RNA, herein designated GAM RNA, and which when bound by GAM429 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM429 target proteins.

[6227] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM429 gene, herein designated GAM GENE, on one or more GAM429 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6228] It is yet further appreciated that specific functions, and accordingly utilities, of GAM429 correlate with, and may be deduced from, the identity of the target genes which GAM429 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6229] Nucleotide sequences of the GAM429 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM429 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM429 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM429 are further described hereinbelow with reference to Table Table1.

[6230] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM429 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6231]

[6232] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 430 (GAM430) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6233] GAM430 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM430 was detected is described hereinabove with reference to Figs. 2-8.

[6234] GAM430 gene, herein designated GAM GENE, and GAM430 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6235] GAM430 gene, herein designated GAM GENE, encodes a GAM430 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM430 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM430 precursor RNA is designated SEQ ID:6752, and is provided hereinbelow with reference to the sequence listing part.

[6236] GAM430 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM430 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6237] An enzyme complex designated DICER COMPLEX, dices the GAM430 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM430 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM430 RNA is designated SEQ ID:6753, and is provided

hereinbelow with reference to the sequence listing part.

[6238] GAM430 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM430 target RNA, herein designated GAM TARGET RNA.
GAM430 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6239] GAM430 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM430 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM430 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM430 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM430 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6240] The complementary binding of GAM430 RNA, herein designated GAM RNA, to target binding sites on GAM430 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM430 target RNA, herein designated GAM TARGET RNA, into GAM430 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6241] It is appreciated that GAM430 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM430 target genes. The mRNA of each one of this plurality of GAM430 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM430 RNA, herein designated GAM RNA, and which when bound by GAM430 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM430 tar-

get proteins.

[6242] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM430 gene, herein designated GAM GENE, on one or more GAM430 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6243] It is yet further appreciated that specific functions, and accordingly utilities, of GAM430 correlate with, and may be deduced from, the identity of the target genes which GAM430 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6244] Nucleotide sequences of the GAM430 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM430 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM430 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM430 are further described hereinbelow with reference to Table Table1.

[6245] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM430 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6246]

[6247] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 431 (GAM431) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6248] GAM431 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM431 was detected is described hereinabove with reference to Figs. 2-8.

[6249] GAM431 gene, herein designated GAM GENE, and GAM431 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6250] GAM431 gene, herein designated GAM GENE, encodes a GAM431 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM431 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM431 precursor RNA is designated SEQ ID:6759, and is provided hereinbelow with reference to the sequence listing part.

[6251] GAM431 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM431 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6252] An enzyme complex designated DICER COMPLEX, dices

the GAM431 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM431 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 73%) nucleotide sequence of GAM431 RNA is designated SEQ ID:6760, and is provided hereinbelow with reference to the sequence listing part.

[6253] GAM431 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM431 target RNA, herein designated GAM TARGET RNA. GAM431 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6254] GAM431 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM431 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM431

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM431 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM431 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6255] The complementary binding of GAM431 RNA, herein designated GAM RNA, to target binding sites on GAM431 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM431 target RNA, herein designated GAM TARGET RNA, into GAM431 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6256] It is appreciated that GAM431 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM431 target genes. The mRNA of each one of this plurality of GAM431 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM431 RNA, herein designated GAM RNA, and which when bound by GAM431 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM431 target proteins.

[6257] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM431 gene, herein designated GAM GENE, on one or more GAM431 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6258] It is yet further appreciated that specific functions, and accordingly utilities, of GAM431 correlate with, and may be deduced from, the identity of the target genes which GAM431 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6259] Nucleotide sequences of the GAM431 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM431 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM431 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM431 are further described hereinbelow with reference to Table Table1.

[6260] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM431 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6261]

[6262] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 432 (GAM432) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6263] GAM432 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM432 was detected is described hereinabove with reference to Figs. 2-8.

[6264] GAM432 gene, herein designated GAM GENE, and GAM432 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6265] GAM432 gene, herein designated GAM GENE, encodes a GAM432 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM432 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM432 precursor RNA is designated SEQ ID:6766, and is provided hereinbelow with reference to the sequence listing part.

[6266] GAM432 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM432 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6267] An enzyme complex designated DICER COMPLEX, dices the GAM432 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM432 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 36%) nucleotide sequence of GAM432 RNA is designated SEQ ID:6767, and is provided hereinbelow with reference to the sequence listing part.

[6268] GAM432 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM432 target RNA, herein designated GAM TARGET RNA. GAM432 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6269] GAM432 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM432 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM432 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM432 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM432 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[6270] The complementary binding of GAM432 RNA, herein designated GAM RNA, to target binding sites on GAM432 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM432 target RNA, herein designated GAM TARGET RNA, into GAM432 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6271] It is appreciated that GAM432 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM432 target genes. The mRNA of each one of this plurality of GAM432 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM432 RNA, herein designated GAM RNA, and which when bound by GAM432 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM432 target proteins.

[6272] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM432 gene, herein designated GAM GENE, on one or

more GAM432 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6273] It is yet further appreciated that specific functions, and accordingly utilities, of GAM432 correlate with, and may be deduced from, the identity of the target genes which GAM432 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6274] Nucleotide sequences of the GAM432 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM432 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM432 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM432 are further de-

scribed hereinbelow with reference to Table Table1.

[6275] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM432 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6276]

[6277] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 433 (GAM433) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6278] GAM433 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM433 was detected is described hereinabove with reference to Figs. 2-8.

[6279] GAM433 gene, herein designated GAM GENE, and GAM433 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6280] GAM433 gene, herein designated GAM GENE, encodes a GAM433 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM433 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM433 precursor RNA is designated SEQ ID:6788, and is provided hereinbelow with reference to the sequence listing part.

[6281] GAM433 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM433 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6282] An enzyme complex designated DICER COMPLEX, dices the GAM433 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM433 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM433 RNA is designated SEQ ID:6789, and is provided hereinbelow with reference to the sequence listing part.

[6283] GAM433 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM433 target RNA, herein designated GAM TARGET RNA. GAM433 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6284] GAM433 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM433 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM433 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM433 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM433 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6285] The complementary binding of GAM433 RNA, herein designated GAM RNA, to target binding sites on GAM433 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM433 target RNA, herein designated GAM TARGET RNA, into GAM433 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6286] It is appreciated that GAM433 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM433 target genes. The mRNA of each one of this plurality of GAM433 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM433 RNA, herein designated GAM RNA, and which when bound by GAM433 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM433 target proteins.

[6287] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM433 gene, herein designated GAM GENE, on one or more GAM433 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6288] It is yet further appreciated that specific functions, and accordingly utilities, of GAM433 correlate with, and may

be deduced from, the identity of the target genes which GAM433 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6289] Nucleotide sequences of the GAM433 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM433 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM433 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM433 are further described hereinbelow with reference to Table Table1.

[6290] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM433 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6291]

[6292] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 434 (GAM434) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6293] GAM434 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM434 was detected is described hereinabove with reference to Figs. 2-8.

[6294] GAM434 gene, herein designated GAM GENE, and GAM434 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6295] GAM434 gene, herein designated GAM GENE, encodes a GAM434 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM434 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM434 precursor RNA is designated SEQ ID:6806, and is provided hereinbelow with reference to the sequence listing part.

[6296] GAM434 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM434 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6297] An enzyme complex designated DICER COMPLEX, dices the GAM434 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM434 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM434 RNA is designated SEQ ID:6807, and is provided hereinbelow with reference to the sequence listing part.

[6298] GAM434 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM434 target RNA, herein designated GAM TARGET RNA. GAM434 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6299] GAM434 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM434 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM434 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM434 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM434 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6300] The complementary binding of GAM434 RNA, herein designated GAM RNA, to target binding sites on GAM434 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM434 target RNA, herein designated GAM TARGET RNA, into GAM434 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6301] It is appreciated that GAM434 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM434 target genes. The mRNA of each one of this plurality of GAM434 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM434 RNA, herein designated GAM RNA, and which when bound by GAM434 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM434 target proteins.

[6302] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM434 gene, herein designated GAM GENE, on one or more GAM434 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6303] It is yet further appreciated that specific functions, and accordingly utilities, of GAM434 correlate with, and may be deduced from, the identity of the target genes which GAM434 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6304] Nucleotide sequences of the GAM434 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM434 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM434 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM434 are further described hereinbelow with reference to Table Table1.

[6305] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM434 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6306]

[6307] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 435 (GAM435) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6308] GAM435 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM435 was detected is described hereinabove with reference to Figs. 2-8.

[6309] GAM435 gene, herein designated GAM GENE, and GAM435 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6310] GAM435 gene, herein designated GAM GENE, encodes a GAM435 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM435 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM435 precursor RNA is desig-

nated SEQ ID:6817, and is provided hereinbelow with reference to the sequence listing part.

[6311] GAM435 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM435 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6312] An enzyme complex designated DICER COMPLEX, dices the GAM435 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM435 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM435 RNA is designated SEQ ID:6818, and is provided hereinbelow with reference to the sequence listing part.

[6313] GAM435 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM435 target RNA, herein designated GAM TARGET RNA. GAM435 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6314] GAM435 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM435 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM435 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM435 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM435

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6315] The complementary binding of GAM435 RNA, herein designated GAM RNA, to target binding sites on GAM435 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM435 target RNA, herein designated GAM TARGET RNA, into GAM435 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6316] It is appreciated that GAM435 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM435 target genes. The mRNA of each one of this plurality of GAM435 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM435 RNA, herein designated GAM RNA, and which when bound by GAM435 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM435 target proteins.

[6317] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM435 gene, herein designated GAM GENE, on one or more GAM435 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6318] It is yet further appreciated that specific functions, and accordingly utilities, of GAM435 correlate with, and may be deduced from, the identity of the target genes which GAM435 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6319] Nucleotide sequences of the GAM435 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM435 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM435 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM435 are further described hereinbelow with reference to Table Table1.

[6320] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM435 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6321]

[6322] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 436 (GAM436) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6323] GAM436 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM436 was detected is described hereinabove with reference to Figs. 2-8.

[6324] GAM436 gene, herein designated GAM GENE, and

GAM436 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6325] GAM436 gene, herein designated GAM GENE, encodes a GAM436 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM436 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM436 precursor RNA is designated SEQ ID:6826, and is provided hereinbelow with reference to the sequence listing part.

[6326] GAM436 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM436 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6327] An enzyme complex designated DICER COMPLEX, dices the GAM436 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM436 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM436 RNA is designated SEQ ID:6827, and is provided hereinbelow with reference to the sequence listing part.

[6328] GAM436 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM436 target RNA, herein designated GAM TARGET RNA. GAM436 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6329] GAM436 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM436 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM436 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM436 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM436 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6330] The complementary binding of GAM436 RNA, herein designated GAM RNA, to target binding sites on GAM436 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM436 target RNA, herein designated GAM TARGET RNA, into GAM436 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6331] It is appreciated that GAM436 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM436 target genes. The mRNA of each one of this plurality of GAM436 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM436 RNA, herein designated GAM RNA, and which when bound by GAM436 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM436 target proteins.

[6332] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM436 gene, herein designated GAM GENE, on one or more GAM436 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6333] It is yet further appreciated that specific functions, and accordingly utilities, of GAM436 correlate with, and may be deduced from, the identity of the target genes which GAM436 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6334] Nucleotide sequences of the GAM436 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM436 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM436 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM436 are further described hereinbelow with reference to Table Table1.

[6335] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM436 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6336]

[6337] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 437 (GAM437) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6338] GAM437 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM437 was detected is described hereinabove with reference to Figs. 2-8.

[6339] GAM437 gene, herein designated GAM GENE, and GAM437 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6340] GAM437 gene, herein designated GAM GENE, encodes a GAM437 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM437 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM437 precursor RNA is designated SEQ ID:6856, and is provided hereinbelow with reference to the sequence listing part.

[6341] GAM437 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM437 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6342] An enzyme complex designated DICER COMPLEX, dices the GAM437 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM437 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM437 RNA is designated SEQ ID:6857, and is provided hereinbelow with reference to the sequence listing part.

[6343] GAM437 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM437 target RNA, herein designated GAM TARGET RNA. GAM437 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6344] GAM437 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM437 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM437 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM437 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM437 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6345] The complementary binding of GAM437 RNA, herein designated GAM RNA, to target binding sites on GAM437 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM437 target RNA, herein designated GAM TARGET RNA, into GAM437 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6346] It is appreciated that GAM437 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM437 target genes. The mRNA of each one of this plurality of GAM437 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM437 RNA, herein designated GAM RNA, and which when bound by GAM437 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM437 target proteins.

[6347] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM437 gene, herein designated GAM GENE, on one or more GAM437 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6348] It is yet further appreciated that specific functions, and accordingly utilities, of GAM437 correlate with, and may be deduced from, the identity of the target genes which GAM437 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6349] Nucleotide sequences of the GAM437 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM437 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM437 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM437 are further described hereinbelow with reference to Table Table1.

[6350] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM437 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6351]

[6352] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 438 (GAM438) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6353] GAM438 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM438 was detected is described hereinabove with reference to Figs. 2-8.

[6354] GAM438 gene, herein designated GAM GENE, and GAM438 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6355] GAM438 gene, herein designated GAM GENE, encodes a GAM438 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM438 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM438 precursor RNA is designated SEQ ID:6869, and is provided hereinbelow with reference to the sequence listing part.

[6356] GAM438 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM438 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6357] An enzyme complex designated DICER COMPLEX, dices the GAM438 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM438 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM438 RNA is designated SEQ ID:6870, and is provided hereinbelow with reference to the sequence listing part.

[6358] GAM438 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM438 target RNA, herein designated GAM TARGET RNA. GAM438 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6359] GAM438 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM438 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM438 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM438 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM438 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6360] The complementary binding of GAM438 RNA, herein designated GAM RNA, to target binding sites on GAM438 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM438 target RNA, herein designated GAM TARGET RNA, into GAM438 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6361] It is appreciated that GAM438 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM438 target genes. The mRNA of each one of this plurality of GAM438 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM438 RNA,

herein designated GAM RNA, and which when bound by GAM438 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM438 target proteins.

[6362] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM438 gene, herein designated GAM GENE, on one or more GAM438 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6363] It is yet further appreciated that specific functions, and accordingly utilities, of GAM438 correlate with, and may be deduced from, the identity of the target genes which

GAM438 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6364] Nucleotide sequences of the GAM438 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM438 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM438 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM438 are further described hereinbelow with reference to Table Table1.

[6365] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM438 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6366]

[6367] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 439 (GAM439) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6368] GAM439 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM439 was detected is described hereinabove with reference to Figs. 2-8.

[6369] GAM439 gene, herein designated GAM GENE, and GAM439 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6370] GAM439 gene, herein designated GAM GENE, encodes a GAM439 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM439 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM439 precursor RNA is designated SEQ ID:6887, and is provided hereinbelow with reference to the sequence listing part.

[6371] GAM439 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM439 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[6372] An enzyme complex designated DICER COMPLEX, dices the GAM439 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM439 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM439 RNA is designated SEQ ID:6888, and is provided hereinbelow with reference to the sequence listing part.

[6373] GAM439 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM439 target RNA, herein designated GAM TARGET RNA.
GAM439 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6374] GAM439 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM439 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM439 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM439 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM439 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6375] The complementary binding of GAM439 RNA, herein designated GAM RNA, to target binding sites on GAM439 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM439 target RNA, herein desig-

nated GAM TARGET RNA, into GAM439 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6376] It is appreciated that GAM439 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM439 target genes. The mRNA of each one of this plurality of GAM439 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM439 RNA, herein designated GAM RNA, and which when bound by GAM439 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM439 target proteins.

[6377] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM439 gene, herein designated GAM GENE, on one or more GAM439 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6378] It is yet further appreciated that specific functions, and accordingly utilities, of GAM439 correlate with, and may be deduced from, the identity of the target genes which GAM439 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6379] Nucleotide sequences of the GAM439 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM439 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM439 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM439 are further described hereinbelow with reference to Table Table1.

[6380] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM439 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[6381]

[6382] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 440 (GAM440) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6383] GAM440 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM440 was detected is described hereinabove with reference to Figs. 2-8.

[6384] GAM440 gene, herein designated GAM GENE, and GAM440 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6385] GAM440 gene, herein designated GAM GENE, encodes a GAM440 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM440 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM440 precursor RNA is designated SEQ ID:6903, and is provided hereinbelow with ref-

erence to the sequence listing part.

[6386] GAM440 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM440 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6387] An enzyme complex designated DICER COMPLEX, dices the GAM440 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM440 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM440 RNA is designated SEQ ID:6904, and is provided hereinbelow with reference to the sequence listing part.

[6388] GAM440 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM440 target RNA, herein designated GAM TARGET RNA.

GAM440 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6389] GAM440 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM440 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM440 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM440 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM440 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6390] The complementary binding of GAM440 RNA, herein designated GAM RNA, to target binding sites on GAM440 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM440 target RNA, herein designated GAM TARGET RNA, into GAM440 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6391] It is appreciated that GAM440 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM440 target genes. The mRNA of each one of this plurality of GAM440 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM440 RNA, herein designated GAM RNA, and which when bound by GAM440 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM440 target proteins.

[6392] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM440 gene, herein designated GAM GENE, on one or more GAM440 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6393] It is yet further appreciated that specific functions, and accordingly utilities, of GAM440 correlate with, and may be deduced from, the identity of the target genes which GAM440 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6394] Nucleotide sequences of the GAM440 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM440 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM440 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM440 are further described hereinbelow with reference to Table Table1.

[6395] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM440 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6396]

[6397] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 441 (GAM441) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6398] GAM441 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM441 was detected is described hereinabove with reference to Figs. 2-8.

[6399] GAM441 gene, herein designated GAM GENE, and GAM441 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[6400] GAM441 gene, herein designated GAM GENE, encodes a GAM441 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM441 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM441 precursor RNA is designated SEQ ID:6916, and is provided hereinbelow with reference to the sequence listing part.

[6401] GAM441 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM441 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6402] An enzyme complex designated DICER COMPLEX, dices the GAM441 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM441 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 76%) nucleotide sequence of GAM441 RNA is designated SEQ ID:6917, and is provided hereinbelow with reference to the sequence listing part.

[6403] GAM441 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM441 target RNA, herein designated GAM TARGET RNA. GAM441 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6404] GAM441 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM441 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM441 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM441 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM441 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6405] The complementary binding of GAM441 RNA, herein designated GAM RNA, to target binding sites on GAM441 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM441 target RNA, herein designated GAM TARGET RNA, into GAM441 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6406] It is appreciated that GAM441 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM441 target genes. The mRNA of each one of this plurality of GAM441 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM441 RNA, herein designated GAM RNA, and which when bound by GAM441 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM441 target proteins.

[6407] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM441 gene, herein designated GAM GENE, on one or more GAM441 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[6408] It is yet further appreciated that specific functions, and accordingly utilities, of GAM441 correlate with, and may be deduced from, the identity of the target genes which GAM441 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6409] Nucleotide sequences of the GAM441 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM441 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM441 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM441 are further described hereinbelow with reference to Table Table1.

[6410] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM441 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6411]

[6412] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

442 (GAM442) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6413] GAM442 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM442 was detected is described hereinabove with reference to Figs. 2-8.

[6414] GAM442 gene, herein designated GAM GENE, and GAM442 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6415] GAM442 gene, herein designated GAM GENE, encodes a GAM442 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM442 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM442 precursor RNA is designated SEQ ID:6920, and is provided hereinbelow with reference to the sequence listing part.

[6416] GAM442 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM442 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6417] An enzyme complex designated DICER COMPLEX, dices the GAM442 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM442 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM442 RNA is designated SEQ ID:6921, and is provided hereinbelow with reference to the sequence listing part.

[6418] GAM442 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM442 target RNA, herein designated GAM TARGET RNA. GAM442 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6419] GAM442 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM442 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM442 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM442 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM442 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6420] The complementary binding of GAM442 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM442 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM442 target RNA, herein designated GAM TARGET RNA, into GAM442 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6421] It is appreciated that GAM442 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM442 target genes. The mRNA of each one of this plurality of GAM442 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM442 RNA, herein designated GAM RNA, and which when bound by GAM442 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM442 target proteins.

[6422] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM442 gene, herein designated GAM GENE, on one or more GAM442 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6423] It is yet further appreciated that specific functions, and accordingly utilities, of GAM442 correlate with, and may be deduced from, the identity of the target genes which GAM442 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6424] Nucleotide sequences of the GAM442 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM442 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM442 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM442 are further described hereinbelow with reference to Table Table1.

[6425] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM442 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6426]

[6427] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 443 (GAM443) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6428] GAM443 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM443 was detected is described hereinabove with reference to Figs. 2-8.

[6429] GAM443 gene, herein designated GAM GENE, and GAM443 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6430] GAM443 gene, herein designated GAM GENE, encodes a GAM443 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM443 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM443 precursor RNA is designated SEQ ID:6933, and is provided hereinbelow with reference to the sequence listing part.

[6431] GAM443 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM443 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6432] An enzyme complex designated DICER COMPLEX, dices the GAM443 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM443 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 62%) nucleotide sequence of GAM443 RNA is designated SEQ ID:6934, and is provided hereinbelow with reference to the sequence listing part.

[6433] GAM443 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM443 target RNA, herein designated GAM TARGET RNA. GAM443 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6434] GAM443 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM443 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM443 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM443 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM443 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6435] The complementary binding of GAM443 RNA, herein designated GAM RNA, to target binding sites on GAM443 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM443 target RNA, herein designated GAM TARGET RNA, into GAM443 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6436] It is appreciated that GAM443 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM443 target genes. The mRNA of each one of this plurality of GAM443 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM443 RNA, herein designated GAM RNA, and which when bound by

GAM443 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM443 target proteins.

[6437] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM443 gene, herein designated GAM GENE, on one or more GAM443 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6438] It is yet further appreciated that specific functions, and accordingly utilities, of GAM443 correlate with, and may be deduced from, the identity of the target genes which GAM443 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[6439] Nucleotide sequences of the GAM443 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM443 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM443 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM443 are further described hereinbelow with reference to Table Table1.

[6440] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM443 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6441]

[6442] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 444 (GAM444) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6443] GAM444 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM444 was detected is described hereinabove with reference to Figs. 2-8.

[6444] GAM444 gene, herein designated GAM GENE, and GAM444 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6445] GAM444 gene, herein designated GAM GENE, encodes a GAM444 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM444 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM444 precursor RNA is designated SEQ ID:6946, and is provided hereinbelow with reference to the sequence listing part.

[6446] GAM444 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM444 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[6447] An enzyme complex designated DICER COMPLEX, dices the GAM444 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM444 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM444 RNA is designated SEQ ID:6947, and is provided hereinbelow with reference to the sequence listing part.

[6448] GAM444 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM444 target RNA, herein designated GAM TARGET RNA. GAM444 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6449] GAM444 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM444 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM444 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM444 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM444 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6450] The complementary binding of GAM444 RNA, herein designated GAM RNA, to target binding sites on GAM444 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM444 target RNA, herein designated GAM TARGET RNA, into GAM444 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6451] It is appreciated that GAM444 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM444 target genes. The mRNA of each one of this plurality of GAM444 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM444 RNA, herein designated GAM RNA, and which when bound by GAM444 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM444 target proteins.

[6452] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM444 gene, herein designated GAM GENE, on one or more GAM444 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6453] It is yet further appreciated that specific functions, and accordingly utilities, of GAM444 correlate with, and may be deduced from, the identity of the target genes which GAM444 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6454] Nucleotide sequences of the GAM444 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM444 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM444 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM444 are further described hereinbelow with reference to Table Table1.

[6455] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM444 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6456]

[6457] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 445 (GAM445) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6458] GAM445 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM445 was detected is described hereinabove with reference to Figs. 2-8.

[6459] GAM445 gene, herein designated GAM GENE, and GAM445 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6460] GAM445 gene, herein designated GAM GENE, encodes a GAM445 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM445 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM445 precursor RNA is designated SEQ ID:6957, and is provided hereinbelow with reference to the sequence listing part.

[6461] GAM445 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM445 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6462] An enzyme complex designated DICER COMPLEX, dices the GAM445 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM445 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM445 RNA is designated SEQ ID:6958, and is provided hereinbelow with reference to the sequence listing part.

[6463] GAM445 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM445

target RNA, herein designated GAM TARGET RNA.

GAM445 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6464] GAM445 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM445 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM445 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM445 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM445 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6465] The complementary binding of GAM445 RNA, herein designated GAM RNA, to target binding sites on GAM445 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM445 target RNA, herein designated GAM TARGET RNA, into GAM445 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6466] It is appreciated that GAM445 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM445 target genes. The mRNA of each one of this plurality of GAM445 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM445 RNA, herein designated GAM RNA, and which when bound by GAM445 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM445 target proteins.

[6467] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM445 gene, herein designated GAM GENE, on one or more GAM445 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6468] It is yet further appreciated that specific functions, and accordingly utilities, of GAM445 correlate with, and may be deduced from, the identity of the target genes which GAM445 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6469] Nucleotide sequences of the GAM445 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM445 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM445 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM445 are further described hereinbelow with reference to Table Table1.

[6470] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM445 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6471]

[6472] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 446 (GAM446) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6473] GAM446 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM446 was detected is described hereinabove with reference to Figs. 2-8.

[6474] GAM446 gene, herein designated GAM GENE, and GAM446 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6475] GAM446 gene, herein designated GAM GENE, encodes a GAM446 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM446 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM446 precursor RNA is designated SEQ ID:6982, and is provided hereinbelow with reference to the sequence listing part.

[6476] GAM446 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM446 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6477] An enzyme complex designated DICER COMPLEX, dices the GAM446 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM446 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM446 RNA is designated SEQ ID:6983, and is provided hereinbelow with reference to the sequence listing part.

[6478] GAM446 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM446 target RNA, herein designated GAM TARGET RNA. GAM446 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6479] GAM446 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM446 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM446 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM446 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM446 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6480] The complementary binding of GAM446 RNA, herein designated GAM RNA, to target binding sites on GAM446 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM446 target RNA, herein designated GAM TARGET RNA, into GAM446 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6481] It is appreciated that GAM446 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM446 target genes. The mRNA of each one of this plu-

rality of GAM446 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM446 RNA, herein designated GAM RNA, and which when bound by GAM446 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM446 target proteins.

[6482] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM446 gene, herein designated GAM GENE, on one or more GAM446 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6483] It is yet further appreciated that specific functions, and accordingly utilities, of GAM446 correlate with, and may be deduced from, the identity of the target genes which GAM446 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6484] Nucleotide sequences of the GAM446 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM446 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM446 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM446 are further described hereinbelow with reference to Table Table1.

[6485] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM446 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6486]

[6487] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 447 (GAM447) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[6488] GAM447 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM447 was detected is described hereinabove with reference to Figs. 2-8.

[6489] GAM447 gene, herein designated GAM GENE, and GAM447 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6490] GAM447 gene, herein designated GAM GENE, encodes a GAM447 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM447 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM447 precursor RNA is designated SEQ ID:7008, and is provided hereinbelow with reference to the sequence listing part.

[6491] GAM447 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM447 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6492] An enzyme complex designated DICER COMPLEX, dices the GAM447 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM447 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM447 RNA is designated SEQ ID:7009, and is provided hereinbelow with reference to the sequence listing part.

[6493] GAM447 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM447 target RNA, herein designated GAM TARGET RNA.
GAM447 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[6494] GAM447 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM447 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM447 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM447 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM447 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6495] The complementary binding of GAM447 RNA, herein designated GAM RNA, to target binding sites on GAM447 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM447 target RNA, herein designated GAM TARGET RNA, into GAM447 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6496] It is appreciated that GAM447 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM447 target genes. The mRNA of each one of this plurality of GAM447 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM447 RNA, herein designated GAM RNA, and which when bound by GAM447 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM447 target proteins.

[6497] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM447 gene, herein designated GAM GENE, on one or more GAM447 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6498] It is yet further appreciated that specific functions, and accordingly utilities, of GAM447 correlate with, and may be deduced from, the identity of the target genes which GAM447 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6499] Nucleotide sequences of the GAM447 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM447 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM447 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM447 are further described hereinbelow with reference to Table Table1.

[6500] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM447 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6501]

[6502] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 448 (GAM448) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6503] GAM448 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM448 was detected is described hereinabove with reference to Figs. 2-8.

[6504] GAM448 gene, herein designated GAM GENE, and GAM448 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6505] GAM448 gene, herein designated GAM GENE, encodes a GAM448 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM448 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM448 precursor RNA is designated SEQ ID:7057, and is provided hereinbelow with reference to the sequence listing part.

[6506] GAM448 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM448 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6507] An enzyme complex designated DICER COMPLEX, dices the GAM448 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM448 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 50%) nucleotide sequence of

GAM448 RNA is designated SEQ ID:7009, and is provided hereinbelow with reference to the sequence listing part.

[6508] GAM448 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM448 target RNA, herein designated GAM TARGET RNA. GAM448 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6509] GAM448 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM448 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM448 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM448 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM448 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6510] The complementary binding of GAM448 RNA, herein designated GAM RNA, to target binding sites on GAM448 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM448 target RNA, herein designated GAM TARGET RNA, into GAM448 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6511] It is appreciated that GAM448 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM448 target genes. The mRNA of each one of this plurality of GAM448 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM448 RNA, herein designated GAM RNA, and which when bound by GAM448 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM448 target proteins.

[6512] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM448 gene, herein designated GAM GENE, on one or more GAM448 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6513] It is yet further appreciated that specific functions, and accordingly utilities, of GAM448 correlate with, and may be deduced from, the identity of the target genes which GAM448 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6514] Nucleotide sequences of the GAM448 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM448 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM448 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM448 are further described hereinbelow with reference to Table Table1.

[6515] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM448 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6516]

[6517] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 449 (GAM449) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6518] GAM449 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM449 was detected is described hereinabove

with reference to Figs. 2-8.

[6519] GAM449 gene, herein designated GAM GENE, and GAM449 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6520] GAM449 gene, herein designated GAM GENE, encodes a GAM449 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM449 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM449 precursor RNA is designated SEQ ID:7061, and is provided hereinbelow with reference to the sequence listing part.

[6521] GAM449 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM449 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6522] An enzyme complex designated DICER COMPLEX, dices the GAM449 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM449 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM449 RNA is designated SEQ ID:7062, and is provided hereinbelow with reference to the sequence listing part.

[6523] GAM449 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM449 target RNA, herein designated GAM TARGET RNA. GAM449 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6524] GAM449 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM449 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM449 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM449 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM449 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6525] The complementary binding of GAM449 RNA, herein designated GAM RNA, to target binding sites on GAM449 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM449 target RNA, herein designated GAM TARGET RNA, into GAM449 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[6526] It is appreciated that GAM449 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM449 target genes. The mRNA of each one of this plurality of GAM449 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM449 RNA, herein designated GAM RNA, and which when bound by GAM449 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM449 target proteins.

[6527] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM449 gene, herein designated GAM GENE, on one or more GAM449 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6528] It is yet further appreciated that specific functions, and accordingly utilities, of GAM449 correlate with, and may be deduced from, the identity of the target genes which GAM449 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6529] Nucleotide sequences of the GAM449 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM449 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM449 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM449 are further described hereinbelow with reference to Table Table1.

[6530] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM449 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6531]

[6532] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 450 (GAM450) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6533] GAM450 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM450 was detected is described hereinabove with reference to Figs. 2-8.

[6534] GAM450 gene, herein designated GAM GENE, and GAM450 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6535] GAM450 gene, herein designated GAM GENE, encodes a GAM450 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM450 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM450 precursor RNA is designated SEQ ID:7075, and is provided hereinbelow with reference to the sequence listing part.

[6536] GAM450 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM450 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6537] An enzyme complex designated DICER COMPLEX, dices the GAM450 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM450 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 25%) nucleotide sequence of GAM450 RNA is designated SEQ ID:7076, and is provided hereinbelow with reference to the sequence listing part.

[6538] GAM450 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM450 target RNA, herein designated GAM TARGET RNA.

GAM450 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6539] GAM450 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM450 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM450 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM450 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM450 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6540] The complementary binding of GAM450 RNA, herein designated GAM RNA, to target binding sites on GAM450 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM450 target RNA, herein designated GAM TARGET RNA, into GAM450 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6541] It is appreciated that GAM450 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM450 target genes. The mRNA of each one of this plurality of GAM450 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM450 RNA, herein designated GAM RNA, and which when bound by GAM450 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM450 target proteins.

[6542] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM450 gene, herein designated GAM GENE, on one or more GAM450 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6543] It is yet further appreciated that specific functions, and accordingly utilities, of GAM450 correlate with, and may be deduced from, the identity of the target genes which GAM450 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6544] Nucleotide sequences of the GAM450 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM450 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM450 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM450 are further described hereinbelow with reference to Table Table1.

[6545] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM450 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6546]

[6547] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 451 (GAM451) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6548] GAM451 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM451 was detected is described hereinabove with reference to Figs. 2-8.

[6549] GAM451 gene, herein designated GAM GENE, and GAM451 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6550] GAM451 gene, herein designated GAM GENE, encodes a

GAM451 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM451 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM451 precursor RNA is designated SEQ ID:7091, and is provided hereinbelow with reference to the sequence listing part.

[6551] GAM451 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM451 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6552] An enzyme complex designated DICER COMPLEX, dices the GAM451 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM451 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM451 RNA is designated SEQ ID:7092, and is provided hereinbelow with reference to the sequence listing part.

[6553] GAM451 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM451 target RNA, herein designated GAM TARGET RNA. GAM451 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6554] GAM451 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM451 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM451 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM451 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM451 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6555] The complementary binding of GAM451 RNA, herein designated GAM RNA, to target binding sites on GAM451 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM451 target RNA, herein designated GAM TARGET RNA, into GAM451 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6556] It is appreciated that GAM451 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM451 target genes. The mRNA of each one of this plurality of GAM451 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM451 RNA, herein designated GAM RNA, and which when bound by GAM451 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM451 target proteins.

[6557] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM451 gene, herein designated GAM GENE, on one or more GAM451 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6558] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM451 correlate with, and may be deduced from, the identity of the target genes which GAM451 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6559] Nucleotide sequences of the GAM451 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM451 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM451 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM451 are further described hereinbelow with reference to Table Table1.

[6560] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM451 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6561]

[6562] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 452 (GAM452) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[6563] GAM452 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM452 was detected is described hereinabove with reference to Figs. 2-8.

[6564] GAM452 gene, herein designated GAM GENE, and GAM452 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6565] GAM452 gene, herein designated GAM GENE, encodes a GAM452 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM452 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM452 precursor RNA is designated SEQ ID:7099, and is provided hereinbelow with reference to the sequence listing part.

[6566] GAM452 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM452 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6567] An enzyme complex designated DICER COMPLEX, dices the GAM452 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM452 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 24%) nucleotide sequence of GAM452 RNA is designated SEQ ID:7100, and is provided hereinbelow with reference to the sequence listing part.

[6568] GAM452 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM452 target RNA, herein designated GAM TARGET RNA. GAM452 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6569] GAM452 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM452 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM452 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM452 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM452 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6570] The complementary binding of GAM452 RNA, herein designated GAM RNA, to target binding sites on GAM452 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM452 target RNA, herein designated GAM TARGET RNA, into GAM452 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6571] It is appreciated that GAM452 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM452 target genes. The mRNA of each one of this plurality of GAM452 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM452 RNA, herein designated GAM RNA, and which when bound by GAM452 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM452 target proteins.

[6572] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM452 gene, herein designated GAM GENE, on one or more GAM452 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6573] It is yet further appreciated that specific functions, and accordingly utilities, of GAM452 correlate with, and may be deduced from, the identity of the target genes which GAM452 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6574] Nucleotide sequences of the GAM452 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM452 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM452 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM452 are further described hereinbelow with reference to Table Table1.

[6575] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM452 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6576]

[6577] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 453 (GAM453) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6578] GAM453 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM453 was detected is described hereinabove with reference to Figs. 2-8.

[6579] GAM453 gene, herein designated GAM GENE, and GAM453 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6580] GAM453 gene, herein designated GAM GENE, encodes a GAM453 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM453 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM453 precursor RNA is designated SEQ ID:7116, and is provided hereinbelow with reference to the sequence listing part.

[6581] GAM453 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM453 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6582] An enzyme complex designated DICER COMPLEX, dices the GAM453 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM453 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM453 RNA is designated SEQ ID:3102, and is provided

hereinbelow with reference to the sequence listing part.

[6583] GAM453 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM453 target RNA, herein designated GAM TARGET RNA.
GAM453 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6584] GAM453 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM453 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM453 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM453 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM453 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6585] The complementary binding of GAM453 RNA, herein designated GAM RNA, to target binding sites on GAM453 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM453 target RNA, herein designated GAM TARGET RNA, into GAM453 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6586] It is appreciated that GAM453 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM453 target genes. The mRNA of each one of this plurality of GAM453 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM453 RNA, herein designated GAM RNA, and which when bound by GAM453 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM453 tar-

get proteins.

[6587] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM453 gene, herein designated GAM GENE, on one or more GAM453 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6588] It is yet further appreciated that specific functions, and accordingly utilities, of GAM453 correlate with, and may be deduced from, the identity of the target genes which GAM453 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6589] Nucleotide sequences of the GAM453 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM453 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM453 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM453 are further described hereinbelow with reference to Table Table1.

[6590] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM453 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6591]

[6592] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 454 (GAM454) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6593] GAM454 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM454 was detected is described hereinabove with reference to Figs. 2-8.

[6594] GAM454 gene, herein designated GAM GENE, and GAM454 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6595] GAM454 gene, herein designated GAM GENE, encodes a GAM454 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM454 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM454 precursor RNA is designated SEQ ID:7127, and is provided hereinbelow with reference to the sequence listing part.

[6596] GAM454 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM454 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6597] An enzyme complex designated DICER COMPLEX, dices

the GAM454 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM454 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM454 RNA is designated SEQ ID:7128, and is provided hereinbelow with reference to the sequence listing part.

[6598] GAM454 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM454 target RNA, herein designated GAM TARGET RNA. GAM454 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6599] GAM454 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM454 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM454

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM454 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM454 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6600] The complementary binding of GAM454 RNA, herein designated GAM RNA, to target binding sites on GAM454 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM454 target RNA, herein designated GAM TARGET RNA, into GAM454 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6601] It is appreciated that GAM454 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM454 target genes. The mRNA of each one of this plurality of GAM454 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM454 RNA, herein designated GAM RNA, and which when bound by GAM454 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM454 target proteins.

[6602] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM454 gene, herein designated GAM GENE, on one or more GAM454 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6603] It is yet further appreciated that specific functions, and accordingly utilities, of GAM454 correlate with, and may be deduced from, the identity of the target genes which GAM454 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6604] Nucleotide sequences of the GAM454 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM454 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM454 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM454 are further described hereinbelow with reference to Table Table1.

[6605] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM454 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6606]

[6607] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 455 (GAM455) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6608] GAM455 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM455 was detected is described hereinabove with reference to Figs. 2-8.

[6609] GAM455 gene, herein designated GAM GENE, and GAM455 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6610] GAM455 gene, herein designated GAM GENE, encodes a GAM455 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM455 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM455 precursor RNA is designated SEQ ID:7145, and is provided hereinbelow with reference to the sequence listing part.

[6611] GAM455 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM455 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6612] An enzyme complex designated DICER COMPLEX, dices the GAM455 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM455 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM455 RNA is designated SEQ ID:7146, and is provided hereinbelow with reference to the sequence listing part.

[6613] GAM455 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM455 target RNA, herein designated GAM TARGET RNA. GAM455 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6614] GAM455 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM455 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM455 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM455 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM455 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[6615] The complementary binding of GAM455 RNA, herein designated GAM RNA, to target binding sites on GAM455 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM455 target RNA, herein designated GAM TARGET RNA, into GAM455 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6616] It is appreciated that GAM455 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM455 target genes. The mRNA of each one of this plurality of GAM455 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM455 RNA, herein designated GAM RNA, and which when bound by GAM455 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM455 target proteins.

[6617] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM455 gene, herein designated GAM GENE, on one or

more GAM455 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6618] It is yet further appreciated that specific functions, and accordingly utilities, of GAM455 correlate with, and may be deduced from, the identity of the target genes which GAM455 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6619] Nucleotide sequences of the GAM455 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM455 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM455 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM455 are further de-

scribed hereinbelow with reference to Table Table1.

[6620] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM455 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6621]

[6622] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 456 (GAM456) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6623] GAM456 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM456 was detected is described hereinabove with reference to Figs. 2-8.

[6624] GAM456 gene, herein designated GAM GENE, and GAM456 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6625] GAM456 gene, herein designated GAM GENE, encodes a GAM456 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM456 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM456 precursor RNA is designated SEQ ID:7191, and is provided hereinbelow with reference to the sequence listing part.

[6626] GAM456 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM456 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6627] An enzyme complex designated DICER COMPLEX, dices the GAM456 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM456 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM456 RNA is designated SEQ ID:7192, and is provided hereinbelow with reference to the sequence listing part.

[6628] GAM456 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM456 target RNA, herein designated GAM TARGET RNA. GAM456 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6629] GAM456 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM456 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM456 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM456 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM456 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6630] The complementary binding of GAM456 RNA, herein designated GAM RNA, to target binding sites on GAM456 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM456 target RNA, herein designated GAM TARGET RNA, into GAM456 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6631] It is appreciated that GAM456 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM456 target genes. The mRNA of each one of this plurality of GAM456 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM456 RNA, herein designated GAM RNA, and which when bound by GAM456 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM456 target proteins.

[6632] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM456 gene, herein designated GAM GENE, on one or more GAM456 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6633] It is yet further appreciated that specific functions, and accordingly utilities, of GAM456 correlate with, and may

be deduced from, the identity of the target genes which GAM456 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6634] Nucleotide sequences of the GAM456 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM456 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM456 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM456 are further described hereinbelow with reference to Table Table1.

[6635] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM456 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6636]

[6637] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 457 (GAM457) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6638] GAM457 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM457 was detected is described hereinabove with reference to Figs. 2-8.

[6639] GAM457 gene, herein designated GAM GENE, and GAM457 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6640] GAM457 gene, herein designated GAM GENE, encodes a GAM457 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM457 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM457 precursor RNA is designated SEQ ID:7205, and is provided hereinbelow with reference to the sequence listing part.

[6641] GAM457 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM457 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6642] An enzyme complex designated DICER COMPLEX, dices the GAM457 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM457 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 26%) nucleotide sequence of GAM457 RNA is designated SEQ ID:7206, and is provided hereinbelow with reference to the sequence listing part.

[6643] GAM457 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM457 target RNA, herein designated GAM TARGET RNA. GAM457 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6644] GAM457 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM457 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM457 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM457 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM457 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6645] The complementary binding of GAM457 RNA, herein designated GAM RNA, to target binding sites on GAM457 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM457 target RNA, herein designated GAM TARGET RNA, into GAM457 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6646] It is appreciated that GAM457 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM457 target genes. The mRNA of each one of this plurality of GAM457 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM457 RNA, herein designated GAM RNA, and which when bound by GAM457 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM457 target proteins.

[6647] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM457 gene, herein designated GAM GENE, on one or more GAM457 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6648] It is yet further appreciated that specific functions, and accordingly utilities, of GAM457 correlate with, and may be deduced from, the identity of the target genes which GAM457 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6649] Nucleotide sequences of the GAM457 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM457 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM457 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM457 are further described hereinbelow with reference to Table Table1.

[6650] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM457 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6651]

[6652] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 458 (GAM458) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6653] GAM458 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM458 was detected is described hereinabove with reference to Figs. 2-8.

[6654] GAM458 gene, herein designated GAM GENE, and GAM458 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6655] GAM458 gene, herein designated GAM GENE, encodes a GAM458 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM458 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM458 precursor RNA is desig-

nated SEQ ID:7226, and is provided hereinbelow with reference to the sequence listing part.

[6656] GAM458 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM458 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6657] An enzyme complex designated DICER COMPLEX, dices the GAM458 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM458 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM458 RNA is designated SEQ ID:7227, and is provided hereinbelow with reference to the sequence listing part.

[6658] GAM458 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM458 target RNA, herein designated GAM TARGET RNA. GAM458 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6659] GAM458 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM458 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM458 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM458 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM458

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6660] The complementary binding of GAM458 RNA, herein designated GAM RNA, to target binding sites on GAM458 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM458 target RNA, herein designated GAM TARGET RNA, into GAM458 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6661] It is appreciated that GAM458 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM458 target genes. The mRNA of each one of this plurality of GAM458 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM458 RNA, herein designated GAM RNA, and which when bound by GAM458 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM458 target proteins.

[6662] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM458 gene, herein designated GAM GENE, on one or more GAM458 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6663] It is yet further appreciated that specific functions, and accordingly utilities, of GAM458 correlate with, and may be deduced from, the identity of the target genes which GAM458 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6664] Nucleotide sequences of the GAM458 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM458 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM458 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM458 are further described hereinbelow with reference to Table Table1.

[6665] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM458 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6666]

[6667] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 459 (GAM459) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6668] GAM459 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM459 was detected is described hereinabove with reference to Figs. 2-8.

[6669] GAM459 gene, herein designated GAM GENE, and

GAM459 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6670] GAM459 gene, herein designated GAM GENE, encodes a GAM459 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM459 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM459 precursor RNA is designated SEQ ID:7262, and is provided hereinbelow with reference to the sequence listing part.

[6671] GAM459 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM459 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6672] An enzyme complex designated DICER COMPLEX, dices the GAM459 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM459 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM459 RNA is designated SEQ ID:7263, and is provided hereinbelow with reference to the sequence listing part.

[6673] GAM459 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM459 target RNA, herein designated GAM TARGET RNA. GAM459 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6674] GAM459 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM459 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM459 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM459 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM459 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6675] The complementary binding of GAM459 RNA, herein designated GAM RNA, to target binding sites on GAM459 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM459 target RNA, herein designated GAM TARGET RNA, into GAM459 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6676] It is appreciated that GAM459 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM459 target genes. The mRNA of each one of this plurality of GAM459 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM459 RNA, herein designated GAM RNA, and which when bound by GAM459 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM459 target proteins.

[6677] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM459 gene, herein designated GAM GENE, on one or more GAM459 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6678] It is yet further appreciated that specific functions, and accordingly utilities, of GAM459 correlate with, and may be deduced from, the identity of the target genes which GAM459 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6679] Nucleotide sequences of the GAM459 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM459 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM459 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM459 are further described hereinbelow with reference to Table Table1.

[6680] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM459 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6681]

[6682] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 460 (GAM460) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6683] GAM460 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM460 was detected is described hereinabove with reference to Figs. 2-8.

[6684] GAM460 gene, herein designated GAM GENE, and GAM460 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6685] GAM460 gene, herein designated GAM GENE, encodes a GAM460 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM460 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM460 precursor RNA is designated SEQ ID:7273, and is provided hereinbelow with reference to the sequence listing part.

[6686] GAM460 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM460 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6687] An enzyme complex designated DICER COMPLEX, dices the GAM460 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM460 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM460 RNA is designated SEQ ID:7274, and is provided hereinbelow with reference to the sequence listing part.

[6688] GAM460 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM460 target RNA, herein designated GAM TARGET RNA. GAM460 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6689] GAM460 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM460 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM460 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM460 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM460 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6690] The complementary binding of GAM460 RNA, herein designated GAM RNA, to target binding sites on GAM460 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM460 target RNA, herein designated GAM TARGET RNA, into GAM460 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6691] It is appreciated that GAM460 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM460 target genes. The mRNA of each one of this plurality of GAM460 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM460 RNA, herein designated GAM RNA, and which when bound by GAM460 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM460 target proteins.

[6692] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM460 gene, herein designated GAM GENE, on one or more GAM460 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6693] It is yet further appreciated that specific functions, and accordingly utilities, of GAM460 correlate with, and may be deduced from, the identity of the target genes which GAM460 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6694] Nucleotide sequences of the GAM460 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM460 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM460 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM460 are further described hereinbelow with reference to Table Table1.

[6695] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM460 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6696]

[6697] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 461 (GAM461) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6698] GAM461 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM461 was detected is described hereinabove with reference to Figs. 2-8.

[6699] GAM461 gene, herein designated GAM GENE, and GAM461 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6700] GAM461 gene, herein designated GAM GENE, encodes a GAM461 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM461 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM461 precursor RNA is designated SEQ ID:7296, and is provided hereinbelow with reference to the sequence listing part.

[6701] GAM461 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM461 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6702] An enzyme complex designated DICER COMPLEX, dices the GAM461 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM461 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM461 RNA is designated SEQ ID:5801, and is provided hereinbelow with reference to the sequence listing part.

[6703] GAM461 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM461 target RNA, herein designated GAM TARGET RNA. GAM461 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6704] GAM461 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM461 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM461 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM461 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM461 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6705] The complementary binding of GAM461 RNA, herein designated GAM RNA, to target binding sites on GAM461 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM461 target RNA, herein designated GAM TARGET RNA, into GAM461 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6706] It is appreciated that GAM461 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM461 target genes. The mRNA of each one of this plurality of GAM461 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM461 RNA,

herein designated GAM RNA, and which when bound by GAM461 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM461 target proteins.

[6707] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM461 gene, herein designated GAM GENE, on one or more GAM461 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6708] It is yet further appreciated that specific functions, and accordingly utilities, of GAM461 correlate with, and may be deduced from, the identity of the target genes which

GAM461 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6709] Nucleotide sequences of the GAM461 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM461 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM461 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM461 are further described hereinbelow with reference to Table Table1.

[6710] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM461 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6711]

[6712] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 462 (GAM462) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6713] GAM462 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM462 was detected is described hereinabove with reference to Figs. 2-8.

[6714] GAM462 gene, herein designated GAM GENE, and GAM462 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6715] GAM462 gene, herein designated GAM GENE, encodes a GAM462 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM462 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM462 precursor RNA is designated SEQ ID:7301, and is provided hereinbelow with reference to the sequence listing part.

[6716] GAM462 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM462 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[6717] An enzyme complex designated DICER COMPLEX, dices the GAM462 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM462 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM462 RNA is designated SEQ ID:7302, and is provided hereinbelow with reference to the sequence listing part.

[6718] GAM462 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM462 target RNA, herein designated GAM TARGET RNA.
GAM462 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6719] GAM462 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM462 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM462 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM462 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM462 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6720] The complementary binding of GAM462 RNA, herein designated GAM RNA, to target binding sites on GAM462 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM462 target RNA, herein desig-

nated GAM TARGET RNA, into GAM462 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6721] It is appreciated that GAM462 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM462 target genes. The mRNA of each one of this plurality of GAM462 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM462 RNA, herein designated GAM RNA, and which when bound by GAM462 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM462 target proteins.

[6722] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM462 gene, herein designated GAM GENE, on one or more GAM462 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6723] It is yet further appreciated that specific functions, and accordingly utilities, of GAM462 correlate with, and may be deduced from, the identity of the target genes which GAM462 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6724] Nucleotide sequences of the GAM462 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM462 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM462 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM462 are further described hereinbelow with reference to Table Table1.

[6725] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM462 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[6726]

[6727] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 463 (GAM463) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6728] GAM463 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM463 was detected is described hereinabove with reference to Figs. 2-8.

[6729] GAM463 gene, herein designated GAM GENE, and GAM463 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6730] GAM463 gene, herein designated GAM GENE, encodes a GAM463 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM463 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM463 precursor RNA is designated SEQ ID:7309, and is provided hereinbelow with ref-

erence to the sequence listing part.

[6731] GAM463 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM463 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6732] An enzyme complex designated DICER COMPLEX, dices the GAM463 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM463 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM463 RNA is designated SEQ ID:7310, and is provided hereinbelow with reference to the sequence listing part.

[6733] GAM463 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM463 target RNA, herein designated GAM TARGET RNA.

GAM463 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6734] GAM463 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM463 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM463 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM463 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM463 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6735] The complementary binding of GAM463 RNA, herein designated GAM RNA, to target binding sites on GAM463 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM463 target RNA, herein designated GAM TARGET RNA, into GAM463 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6736] It is appreciated that GAM463 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM463 target genes. The mRNA of each one of this plurality of GAM463 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM463 RNA, herein designated GAM RNA, and which when bound by GAM463 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM463 target proteins.

[6737] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM463 gene, herein designated GAM GENE, on one or more GAM463 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6738] It is yet further appreciated that specific functions, and accordingly utilities, of GAM463 correlate with, and may be deduced from, the identity of the target genes which GAM463 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6739] Nucleotide sequences of the GAM463 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM463 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM463 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM463 are further described hereinbelow with reference to Table Table1.

[6740] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM463 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6741]

[6742] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 464 (GAM464) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6743] GAM464 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM464 was detected is described hereinabove with reference to Figs. 2-8.

[6744] GAM464 gene, herein designated GAM GENE, and GAM464 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[6745] GAM464 gene, herein designated GAM GENE, encodes a GAM464 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM464 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM464 precursor RNA is designated SEQ ID:7323, and is provided hereinbelow with reference to the sequence listing part.

[6746] GAM464 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM464 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6747] An enzyme complex designated DICER COMPLEX, dices the GAM464 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM464 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM464 RNA is designated SEQ ID:7324, and is provided hereinbelow with reference to the sequence listing part.

[6748] GAM464 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM464 target RNA, herein designated GAM TARGET RNA. GAM464 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6749] GAM464 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM464 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM464 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM464 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM464 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6750] The complementary binding of GAM464 RNA, herein designated GAM RNA, to target binding sites on GAM464 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM464 target RNA, herein designated GAM TARGET RNA, into GAM464 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6751] It is appreciated that GAM464 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM464 target genes. The mRNA of each one of this plurality of GAM464 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM464 RNA, herein designated GAM RNA, and which when bound by GAM464 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM464 target proteins.

[6752] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM464 gene, herein designated GAM GENE, on one or more GAM464 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[6753] It is yet further appreciated that specific functions, and accordingly utilities, of GAM464 correlate with, and may be deduced from, the identity of the target genes which GAM464 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6754] Nucleotide sequences of the GAM464 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM464 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM464 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM464 are further described hereinbelow with reference to Table Table1.

[6755] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM464 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6756]

[6757] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

465 (GAM465) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6758] GAM465 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM465 was detected is described hereinabove with reference to Figs. 2-8.

[6759] GAM465 gene, herein designated GAM GENE, and GAM465 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6760] GAM465 gene, herein designated GAM GENE, encodes a GAM465 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM465 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM465 precursor RNA is designated SEQ ID:7338, and is provided hereinbelow with reference to the sequence listing part.

[6761] GAM465 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM465 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6762] An enzyme complex designated DICER COMPLEX, dices the GAM465 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM465 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 35%) nucleotide sequence of GAM465 RNA is designated SEQ ID:7339, and is provided hereinbelow with reference to the sequence listing part.

[6763] GAM465 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM465 target RNA, herein designated GAM TARGET RNA. GAM465 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6764] GAM465 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM465 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM465 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM465 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM465 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6765] The complementary binding of GAM465 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM465 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM465 target RNA, herein designated GAM TARGET RNA, into GAM465 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6766] It is appreciated that GAM465 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM465 target genes. The mRNA of each one of this plurality of GAM465 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM465 RNA, herein designated GAM RNA, and which when bound by GAM465 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM465 target proteins.

[6767] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM465 gene, herein designated GAM GENE, on one or more GAM465 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6768] It is yet further appreciated that specific functions, and accordingly utilities, of GAM465 correlate with, and may be deduced from, the identity of the target genes which GAM465 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6769] Nucleotide sequences of the GAM465 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM465 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM465 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM465 are further described hereinbelow with reference to Table Table1.

[6770] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM465 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6771]

[6772] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 466 (GAM466) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6773] GAM466 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM466 was detected is described hereinabove with reference to Figs. 2-8.

[6774] GAM466 gene, herein designated GAM GENE, and GAM466 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6775] GAM466 gene, herein designated GAM GENE, encodes a GAM466 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM466 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM466 precursor RNA is designated SEQ ID:7348, and is provided hereinbelow with reference to the sequence listing part.

[6776] GAM466 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM466 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6777] An enzyme complex designated DICER COMPLEX, dices the GAM466 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM466 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 40%) nucleotide sequence of GAM466 RNA is designated SEQ ID:7349, and is provided hereinbelow with reference to the sequence listing part.

[6778] GAM466 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM466 target RNA, herein designated GAM TARGET RNA. GAM466 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6779] GAM466 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM466 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM466 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM466 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM466 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6780] The complementary binding of GAM466 RNA, herein designated GAM RNA, to target binding sites on GAM466 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM466 target RNA, herein designated GAM TARGET RNA, into GAM466 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6781] It is appreciated that GAM466 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM466 target genes. The mRNA of each one of this plurality of GAM466 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM466 RNA, herein designated GAM RNA, and which when bound by

GAM466 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM466 target proteins.

[6782] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM466 gene, herein designated GAM GENE, on one or more GAM466 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6783] It is yet further appreciated that specific functions, and accordingly utilities, of GAM466 correlate with, and may be deduced from, the identity of the target genes which GAM466 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[6784] Nucleotide sequences of the GAM466 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM466 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM466 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM466 are further described hereinbelow with reference to Table Table1.

[6785] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM466 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6786]

[6787] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 467 (GAM467) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6788] GAM467 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM467 was detected is described hereinabove with reference to Figs. 2-8.

[6789] GAM467 gene, herein designated GAM GENE, and GAM467 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6790] GAM467 gene, herein designated GAM GENE, encodes a GAM467 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM467 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM467 precursor RNA is designated SEQ ID:7361, and is provided hereinbelow with reference to the sequence listing part.

[6791] GAM467 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM467 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[6792] An enzyme complex designated DICER COMPLEX, dices the GAM467 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM467 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM467 RNA is designated SEQ ID:7362, and is provided hereinbelow with reference to the sequence listing part.

[6793] GAM467 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM467 target RNA, herein designated GAM TARGET RNA. GAM467 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6794] GAM467 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM467 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM467 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM467 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM467 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6795] The complementary binding of GAM467 RNA, herein designated GAM RNA, to target binding sites on GAM467 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM467 target RNA, herein designated GAM TARGET RNA, into GAM467 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6796] It is appreciated that GAM467 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM467 target genes. The mRNA of each one of this plurality of GAM467 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM467 RNA, herein designated GAM RNA, and which when bound by GAM467 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM467 target proteins.

[6797] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM467 gene, herein designated GAM GENE, on one or more GAM467 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6798] It is yet further appreciated that specific functions, and accordingly utilities, of GAM467 correlate with, and may be deduced from, the identity of the target genes which GAM467 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6799] Nucleotide sequences of the GAM467 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM467 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM467 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM467 are further described hereinbelow with reference to Table Table1.

[6800] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM467 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6801]

[6802] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 468 (GAM468) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6803] GAM468 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM468 was detected is described hereinabove with reference to Figs. 2-8.

[6804] GAM468 gene, herein designated GAM GENE, and GAM468 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6805] GAM468 gene, herein designated GAM GENE, encodes a GAM468 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM468 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM468 precursor RNA is designated SEQ ID:7380, and is provided hereinbelow with reference to the sequence listing part.

[6806] GAM468 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM468 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6807] An enzyme complex designated DICER COMPLEX, dices the GAM468 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM468 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM468 RNA is designated SEQ ID:7381, and is provided hereinbelow with reference to the sequence listing part.

[6808] GAM468 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM468

target RNA, herein designated GAM TARGET RNA.

GAM468 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6809] GAM468 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM468 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM468 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM468 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM468 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6810] The complementary binding of GAM468 RNA, herein designated GAM RNA, to target binding sites on GAM468 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM468 target RNA, herein designated GAM TARGET RNA, into GAM468 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6811] It is appreciated that GAM468 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM468 target genes. The mRNA of each one of this plurality of GAM468 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM468 RNA, herein designated GAM RNA, and which when bound by GAM468 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM468 target proteins.

[6812] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM468 gene, herein designated GAM GENE, on one or more GAM468 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6813] It is yet further appreciated that specific functions, and accordingly utilities, of GAM468 correlate with, and may be deduced from, the identity of the target genes which GAM468 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6814] Nucleotide sequences of the GAM468 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM468 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM468 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM468 are further described hereinbelow with reference to Table Table1.

[6815] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM468 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6816]

[6817] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 469 (GAM469) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6818] GAM469 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM469 was detected is described hereinabove with reference to Figs. 2-8.

[6819] GAM469 gene, herein designated GAM GENE, and GAM469 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6820] GAM469 gene, herein designated GAM GENE, encodes a GAM469 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM469 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM469 precursor RNA is designated SEQ ID:7395, and is provided hereinbelow with reference to the sequence listing part.

[6821] GAM469 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM469 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6822] An enzyme complex designated DICER COMPLEX, dices the GAM469 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM469 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 69%) nucleotide sequence of GAM469 RNA is designated SEQ ID:7396, and is provided hereinbelow with reference to the sequence listing part.

[6823] GAM469 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM469 target RNA, herein designated GAM TARGET RNA. GAM469 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6824] GAM469 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM469 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM469 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM469 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM469 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6825] The complementary binding of GAM469 RNA, herein designated GAM RNA, to target binding sites on GAM469 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM469 target RNA, herein designated GAM TARGET RNA, into GAM469 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6826] It is appreciated that GAM469 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM469 target genes. The mRNA of each one of this plu-

rality of GAM469 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM469 RNA, herein designated GAM RNA, and which when bound by GAM469 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM469 target proteins.

[6827] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM469 gene, herein designated GAM GENE, on one or more GAM469 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6828] It is yet further appreciated that specific functions, and accordingly utilities, of GAM469 correlate with, and may be deduced from, the identity of the target genes which GAM469 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6829] Nucleotide sequences of the GAM469 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM469 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM469 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM469 are further described hereinbelow with reference to Table Table1.

[6830] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM469 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6831]

[6832] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 470 (GAM470) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[6833] GAM470 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM470 was detected is described hereinabove with reference to Figs. 2-8.

[6834] GAM470 gene, herein designated GAM GENE, and GAM470 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6835] GAM470 gene, herein designated GAM GENE, encodes a GAM470 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM470 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM470 precursor RNA is designated SEQ ID:7429, and is provided hereinbelow with reference to the sequence listing part.

[6836] GAM470 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM470 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6837] An enzyme complex designated DICER COMPLEX, dices the GAM470 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM470 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 39%) nucleotide sequence of GAM470 RNA is designated SEQ ID:7430, and is provided hereinbelow with reference to the sequence listing part.

[6838] GAM470 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM470 target RNA, herein designated GAM TARGET RNA.
GAM470 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[6839] GAM470 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM470 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM470 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM470 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM470 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6840] The complementary binding of GAM470 RNA, herein designated GAM RNA, to target binding sites on GAM470 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM470 target RNA, herein designated GAM TARGET RNA, into GAM470 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6841] It is appreciated that GAM470 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM470 target genes. The mRNA of each one of this plurality of GAM470 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM470 RNA, herein designated GAM RNA, and which when bound by GAM470 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM470 target proteins.

[6842] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM470 gene, herein designated GAM GENE, on one or more GAM470 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6843] It is yet further appreciated that specific functions, and accordingly utilities, of GAM470 correlate with, and may be deduced from, the identity of the target genes which GAM470 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6844] Nucleotide sequences of the GAM470 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM470 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM470 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM470 are further described hereinbelow with reference to Table Table1.

[6845] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM470 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6846]

[6847] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 471 (GAM471) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6848] GAM471 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM471 was detected is described hereinabove with reference to Figs. 2-8.

[6849] GAM471 gene, herein designated GAM GENE, and GAM471 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6850] GAM471 gene, herein designated GAM GENE, encodes a GAM471 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM471 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM471 precursor RNA is designated SEQ ID:7457, and is provided hereinbelow with reference to the sequence listing part.

[6851] GAM471 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM471 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6852] An enzyme complex designated DICER COMPLEX, dices the GAM471 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM471 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 33%) nucleotide sequence of

GAM471 RNA is designated SEQ ID:7458, and is provided hereinbelow with reference to the sequence listing part.

[6853] GAM471 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM471 target RNA, herein designated GAM TARGET RNA. GAM471 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6854] GAM471 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM471 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM471 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM471 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM471 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6855] The complementary binding of GAM471 RNA, herein designated GAM RNA, to target binding sites on GAM471 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM471 target RNA, herein designated GAM TARGET RNA, into GAM471 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6856] It is appreciated that GAM471 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM471 target genes. The mRNA of each one of this plurality of GAM471 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM471 RNA, herein designated GAM RNA, and which when bound by GAM471 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM471 target proteins.

[6857] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM471 gene, herein designated GAM GENE, on one or more GAM471 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6858] It is yet further appreciated that specific functions, and accordingly utilities, of GAM471 correlate with, and may be deduced from, the identity of the target genes which GAM471 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6859] Nucleotide sequences of the GAM471 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM471 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM471 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM471 are further described hereinbelow with reference to Table Table1.

[6860] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM471 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6861]

[6862] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 472 (GAM472) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6863] GAM472 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM472 was detected is described hereinabove

with reference to Figs. 2-8.

[6864] GAM472 gene, herein designated GAM GENE, and GAM472 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6865] GAM472 gene, herein designated GAM GENE, encodes a GAM472 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM472 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM472 precursor RNA is designated SEQ ID:7463, and is provided hereinbelow with reference to the sequence listing part.

[6866] GAM472 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM472 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6867] An enzyme complex designated DICER COMPLEX, dices the GAM472 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM472 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM472 RNA is designated SEQ ID:7464, and is provided hereinbelow with reference to the sequence listing part.

[6868] GAM472 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM472 target RNA, herein designated GAM TARGET RNA. GAM472 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6869] GAM472 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM472 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM472 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM472 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM472 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6870] The complementary binding of GAM472 RNA, herein designated GAM RNA, to target binding sites on GAM472 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM472 target RNA, herein designated GAM TARGET RNA, into GAM472 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[6871] It is appreciated that GAM472 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM472 target genes. The mRNA of each one of this plurality of GAM472 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM472 RNA, herein designated GAM RNA, and which when bound by GAM472 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM472 target proteins.

[6872] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM472 gene, herein designated GAM GENE, on one or more GAM472 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6873] It is yet further appreciated that specific functions, and accordingly utilities, of GAM472 correlate with, and may be deduced from, the identity of the target genes which GAM472 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6874] Nucleotide sequences of the GAM472 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM472 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM472 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM472 are further described hereinbelow with reference to Table Table1.

[6875] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM472 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6876]

[6877] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 473 (GAM473) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6878] GAM473 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM473 was detected is described hereinabove with reference to Figs. 2-8.

[6879] GAM473 gene, herein designated GAM GENE, and GAM473 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6880] GAM473 gene, herein designated GAM GENE, encodes a GAM473 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM473 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM473 precursor RNA is designated SEQ ID:7481, and is provided hereinbelow with reference to the sequence listing part.

[6881] GAM473 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM473 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6882] An enzyme complex designated DICER COMPLEX, dices the GAM473 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM473 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 18%) nucleotide sequence of GAM473 RNA is designated SEQ ID:7482, and is provided hereinbelow with reference to the sequence listing part.

[6883] GAM473 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM473 target RNA, herein designated GAM TARGET RNA.

GAM473 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6884] GAM473 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM473 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM473 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM473 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM473 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6885] The complementary binding of GAM473 RNA, herein designated GAM RNA, to target binding sites on GAM473 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM473 target RNA, herein designated GAM TARGET RNA, into GAM473 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6886] It is appreciated that GAM473 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM473 target genes. The mRNA of each one of this plurality of GAM473 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM473 RNA, herein designated GAM RNA, and which when bound by GAM473 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM473 target proteins.

[6887] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM473 gene, herein designated GAM GENE, on one or more GAM473 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6888] It is yet further appreciated that specific functions, and accordingly utilities, of GAM473 correlate with, and may be deduced from, the identity of the target genes which GAM473 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6889] Nucleotide sequences of the GAM473 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM473 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM473 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM473 are further described hereinbelow with reference to Table Table1.

[6890] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM473 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6891]

[6892] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 474 (GAM474) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6893] GAM474 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM474 was detected is described hereinabove with reference to Figs. 2-8.

[6894] GAM474 gene, herein designated GAM GENE, and GAM474 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6895] GAM474 gene, herein designated GAM GENE, encodes a

GAM474 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM474 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM474 precursor RNA is designated SEQ ID:7493, and is provided hereinbelow with reference to the sequence listing part.

[6896] GAM474 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM474 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6897] An enzyme complex designated DICER COMPLEX, dices the GAM474 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM474 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 43%) nucleotide sequence of GAM474 RNA is designated SEQ ID:7494, and is provided hereinbelow with reference to the sequence listing part.

[6898] GAM474 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM474 target RNA, herein designated GAM TARGET RNA. GAM474 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6899] GAM474 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM474 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM474 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM474 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM474 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6900] The complementary binding of GAM474 RNA, herein designated GAM RNA, to target binding sites on GAM474 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM474 target RNA, herein designated GAM TARGET RNA, into GAM474 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6901] It is appreciated that GAM474 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM474 target genes. The mRNA of each one of this plurality of GAM474 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM474 RNA, herein designated GAM RNA, and which when bound by GAM474 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM474 target proteins.

[6902] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM474 gene, herein designated GAM GENE, on one or more GAM474 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6903] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM474 correlate with, and may be deduced from, the identity of the target genes which GAM474 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6904] Nucleotide sequences of the GAM474 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM474 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM474 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM474 are further described hereinbelow with reference to Table Table1.

[6905] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM474 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6906]

[6907] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 475 (GAM475) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[6908] GAM475 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM475 was detected is described hereinabove with reference to Figs. 2-8.

[6909] GAM475 gene, herein designated GAM GENE, and GAM475 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6910] GAM475 gene, herein designated GAM GENE, encodes a GAM475 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM475 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM475 precursor RNA is designated SEQ ID:7511, and is provided hereinbelow with reference to the sequence listing part.

[6911] GAM475 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM475 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6912] An enzyme complex designated DICER COMPLEX, dices the GAM475 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM475 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 27%) nucleotide sequence of GAM475 RNA is designated SEQ ID:7512, and is provided hereinbelow with reference to the sequence listing part.

[6913] GAM475 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM475 target RNA, herein designated GAM TARGET RNA. GAM475 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6914] GAM475 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM475 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM475 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM475 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM475 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6915] The complementary binding of GAM475 RNA, herein designated GAM RNA, to target binding sites on GAM475 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM475 target RNA, herein designated GAM TARGET RNA, into GAM475 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6916] It is appreciated that GAM475 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM475 target genes. The mRNA of each one of this plurality of GAM475 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM475 RNA, herein designated GAM RNA, and which when bound by GAM475 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM475 target proteins.

[6917] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM475 gene, herein designated GAM GENE, on one or more GAM475 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6918] It is yet further appreciated that specific functions, and accordingly utilities, of GAM475 correlate with, and may be deduced from, the identity of the target genes which GAM475 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6919] Nucleotide sequences of the GAM475 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM475 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM475 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM475 are further described hereinbelow with reference to Table Table1.

[6920] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM475 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6921]

[6922] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 476 (GAM476) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6923] GAM476 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM476 was detected is described hereinabove with reference to Figs. 2-8.

[6924] GAM476 gene, herein designated GAM GENE, and GAM476 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6925] GAM476 gene, herein designated GAM GENE, encodes a GAM476 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM476 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM476 precursor RNA is designated SEQ ID:7541, and is provided hereinbelow with reference to the sequence listing part.

[6926] GAM476 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM476 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6927] An enzyme complex designated DICER COMPLEX, dices the GAM476 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM476 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 55%) nucleotide sequence of GAM476 RNA is designated SEQ ID:7542, and is provided

hereinbelow with reference to the sequence listing part.

[6928] GAM476 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM476 target RNA, herein designated GAM TARGET RNA.

GAM476 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6929] GAM476 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM476 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM476 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM476 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM476 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6930] The complementary binding of GAM476 RNA, herein designated GAM RNA, to target binding sites on GAM476 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM476 target RNA, herein designated GAM TARGET RNA, into GAM476 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6931] It is appreciated that GAM476 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM476 target genes. The mRNA of each one of this plurality of GAM476 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM476 RNA, herein designated GAM RNA, and which when bound by GAM476 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM476 tar-

get proteins.

[6932] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM476 gene, herein designated GAM GENE, on one or more GAM476 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6933] It is yet further appreciated that specific functions, and accordingly utilities, of GAM476 correlate with, and may be deduced from, the identity of the target genes which GAM476 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6934] Nucleotide sequences of the GAM476 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM476 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM476 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM476 are further described hereinbelow with reference to Table Table1.

[6935] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM476 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6936]

[6937] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 477 (GAM477) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6938] GAM477 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM477 was detected is described hereinabove with reference to Figs. 2-8.

[6939] GAM477 gene, herein designated GAM GENE, and GAM477 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6940] GAM477 gene, herein designated GAM GENE, encodes a GAM477 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM477 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM477 precursor RNA is designated SEQ ID:7561, and is provided hereinbelow with reference to the sequence listing part.

[6941] GAM477 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM477 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6942] An enzyme complex designated DICER COMPLEX, dices

the GAM477 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM477 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM477 RNA is designated SEQ ID:7562, and is provided hereinbelow with reference to the sequence listing part.

[6943] GAM477 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM477 target RNA, herein designated GAM TARGET RNA. GAM477 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6944] GAM477 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM477 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM477

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM477 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM477 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6945] The complementary binding of GAM477 RNA, herein designated GAM RNA, to target binding sites on GAM477 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM477 target RNA, herein designated GAM TARGET RNA, into GAM477 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6946] It is appreciated that GAM477 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM477 target genes. The mRNA of each one of this plurality of GAM477 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM477 RNA, herein designated GAM RNA, and which when bound by GAM477 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM477 target proteins.

[6947] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM477 gene, herein designated GAM GENE, on one or more GAM477 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6948] It is yet further appreciated that specific functions, and accordingly utilities, of GAM477 correlate with, and may be deduced from, the identity of the target genes which GAM477 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6949] Nucleotide sequences of the GAM477 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM477 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM477 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM477 are further described hereinbelow with reference to Table Table1.

[6950] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM477 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6951]

[6952] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 478 (GAM478) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6953] GAM478 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM478 was detected is described hereinabove with reference to Figs. 2-8.

[6954] GAM478 gene, herein designated GAM GENE, and GAM478 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6955] GAM478 gene, herein designated GAM GENE, encodes a GAM478 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM478 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM478 precursor RNA is designated SEQ ID:7578, and is provided hereinbelow with reference to the sequence listing part.

[6956] GAM478 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM478 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6957] An enzyme complex designated DICER COMPLEX, dices the GAM478 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM478 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM478 RNA is designated SEQ ID:7579, and is provided hereinbelow with reference to the sequence listing part.

[6958] GAM478 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM478 target RNA, herein designated GAM TARGET RNA. GAM478 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6959] GAM478 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM478 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM478 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM478 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM478 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[6960] The complementary binding of GAM478 RNA, herein designated GAM RNA, to target binding sites on GAM478 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM478 target RNA, herein designated GAM TARGET RNA, into GAM478 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6961] It is appreciated that GAM478 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM478 target genes. The mRNA of each one of this plurality of GAM478 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM478 RNA, herein designated GAM RNA, and which when bound by GAM478 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM478 target proteins.

[6962] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM478 gene, herein designated GAM GENE, on one or

more GAM478 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6963] It is yet further appreciated that specific functions, and accordingly utilities, of GAM478 correlate with, and may be deduced from, the identity of the target genes which GAM478 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6964] Nucleotide sequences of the GAM478 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM478 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM478 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM478 are further de-

scribed hereinbelow with reference to Table Table1.

[6965] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM478 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6966]

[6967] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 479 (GAM479) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6968] GAM479 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM479 was detected is described hereinabove with reference to Figs. 2-8.

[6969] GAM479 gene, herein designated GAM GENE, and GAM479 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6970] GAM479 gene, herein designated GAM GENE, encodes a GAM479 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM479 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM479 precursor RNA is designated SEQ ID:7602, and is provided hereinbelow with reference to the sequence listing part.

[6971] GAM479 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM479 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6972] An enzyme complex designated DICER COMPLEX, dices the GAM479 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM479 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM479 RNA is designated SEQ ID:7603, and is provided hereinbelow with reference to the sequence listing part.

[6973] GAM479 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM479 target RNA, herein designated GAM TARGET RNA. GAM479 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6974] GAM479 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM479 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM479 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM479 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM479 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6975] The complementary binding of GAM479 RNA, herein designated GAM RNA, to target binding sites on GAM479 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM479 target RNA, herein designated GAM TARGET RNA, into GAM479 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6976] It is appreciated that GAM479 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM479 target genes. The mRNA of each one of this plurality of GAM479 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM479 RNA, herein designated GAM RNA, and which when bound by GAM479 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM479 target proteins.

[6977] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM479 gene, herein designated GAM GENE, on one or more GAM479 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6978] It is yet further appreciated that specific functions, and accordingly utilities, of GAM479 correlate with, and may

be deduced from, the identity of the target genes which GAM479 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6979] Nucleotide sequences of the GAM479 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM479 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM479 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM479 are further described hereinbelow with reference to Table Table1.

[6980] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM479 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6981]

[6982] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 480 (GAM480) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6983] GAM480 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM480 was detected is described hereinabove with reference to Figs. 2-8.

[6984] GAM480 gene, herein designated GAM GENE, and GAM480 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[6985] GAM480 gene, herein designated GAM GENE, encodes a GAM480 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM480 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM480 precursor RNA is designated SEQ ID:7622, and is provided hereinbelow with reference to the sequence listing part.

[6986] GAM480 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM480 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[6987] An enzyme complex designated DICER COMPLEX, dices the GAM480 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM480 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 51%) nucleotide sequence of GAM480 RNA is designated SEQ ID:7623, and is provided hereinbelow with reference to the sequence listing part.

[6988] GAM480 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM480 target RNA, herein designated GAM TARGET RNA. GAM480 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[6989] GAM480 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM480 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM480 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM480 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM480 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[6990] The complementary binding of GAM480 RNA, herein designated GAM RNA, to target binding sites on GAM480 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM480 target RNA, herein designated GAM TARGET RNA, into GAM480 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[6991] It is appreciated that GAM480 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM480 target genes. The mRNA of each one of this plurality of GAM480 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM480 RNA, herein designated GAM RNA, and which when bound by GAM480 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM480 target proteins.

[6992] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM480 gene, herein designated GAM GENE, on one or more GAM480 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[6993] It is yet further appreciated that specific functions, and accordingly utilities, of GAM480 correlate with, and may be deduced from, the identity of the target genes which GAM480 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[6994] Nucleotide sequences of the GAM480 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM480 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM480 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM480 are further described hereinbelow with reference to Table Table1.

[6995] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM480 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[6996]

[6997] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 481 (GAM481) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[6998] GAM481 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM481 was detected is described hereinabove with reference to Figs. 2-8.

[6999] GAM481 gene, herein designated GAM GENE, and GAM481 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7000] GAM481 gene, herein designated GAM GENE, encodes a GAM481 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM481 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM481 precursor RNA is desig-

nated SEQ ID:7628, and is provided hereinbelow with reference to the sequence listing part.

[7001] GAM481 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM481 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7002] An enzyme complex designated DICER COMPLEX, dices the GAM481 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM481 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM481 RNA is designated SEQ ID:7629, and is provided hereinbelow with reference to the sequence listing part.

[7003] GAM481 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM481 target RNA, herein designated GAM TARGET RNA. GAM481 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7004] GAM481 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM481 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM481 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM481 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM481

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7005] The complementary binding of GAM481 RNA, herein designated GAM RNA, to target binding sites on GAM481 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM481 target RNA, herein designated GAM TARGET RNA, into GAM481 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7006] It is appreciated that GAM481 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM481 target genes. The mRNA of each one of this plurality of GAM481 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM481 RNA, herein designated GAM RNA, and which when bound by GAM481 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM481 target proteins.

[7007] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM481 gene, herein designated GAM GENE, on one or more GAM481 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7008] It is yet further appreciated that specific functions, and accordingly utilities, of GAM481 correlate with, and may be deduced from, the identity of the target genes which GAM481 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7009] Nucleotide sequences of the GAM481 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM481 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM481 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM481 are further described hereinbelow with reference to Table Table1.

[7010] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM481 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7011]

[7012] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 482 (GAM482) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7013] GAM482 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM482 was detected is described hereinabove with reference to Figs. 2-8.

[7014] GAM482 gene, herein designated GAM GENE, and

GAM482 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7015] GAM482 gene, herein designated GAM GENE, encodes a GAM482 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM482 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM482 precursor RNA is designated SEQ ID:7634, and is provided hereinbelow with reference to the sequence listing part.

[7016] GAM482 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM482 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7017] An enzyme complex designated DICER COMPLEX, dices the GAM482 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM482 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM482 RNA is designated SEQ ID:7635, and is provided hereinbelow with reference to the sequence listing part.

[7018] GAM482 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM482 target RNA, herein designated GAM TARGET RNA. GAM482 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7019] GAM482 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM482 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM482 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM482 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM482 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7020] The complementary binding of GAM482 RNA, herein designated GAM RNA, to target binding sites on GAM482 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM482 target RNA, herein designated GAM TARGET RNA, into GAM482 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7021] It is appreciated that GAM482 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM482 target genes. The mRNA of each one of this plurality of GAM482 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM482 RNA, herein designated GAM RNA, and which when bound by GAM482 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM482 target proteins.

[7022] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM482 gene, herein designated GAM GENE, on one or more GAM482 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7023] It is yet further appreciated that specific functions, and accordingly utilities, of GAM482 correlate with, and may be deduced from, the identity of the target genes which GAM482 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7024] Nucleotide sequences of the GAM482 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM482 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM482 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM482 are further described hereinbelow with reference to Table Table1.

[7025] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM482 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7026]

[7027] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 483 (GAM483) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7028] GAM483 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM483 was detected is described hereinabove with reference to Figs. 2-8.

[7029] GAM483 gene, herein designated GAM GENE, and GAM483 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7030] GAM483 gene, herein designated GAM GENE, encodes a GAM483 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM483 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM483 precursor RNA is designated SEQ ID:7637, and is provided hereinbelow with reference to the sequence listing part.

[7031] GAM483 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM483 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7032] An enzyme complex designated DICER COMPLEX, dices the GAM483 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM483 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM483 RNA is designated SEQ ID:7638, and is provided hereinbelow with reference to the sequence listing part.

[7033] GAM483 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM483 target RNA, herein designated GAM TARGET RNA. GAM483 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7034] GAM483 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM483 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM483 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM483 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM483 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7035] The complementary binding of GAM483 RNA, herein designated GAM RNA, to target binding sites on GAM483 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM483 target RNA, herein designated GAM TARGET RNA, into GAM483 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7036] It is appreciated that GAM483 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM483 target genes. The mRNA of each one of this plurality of GAM483 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM483 RNA, herein designated GAM RNA, and which when bound by GAM483 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM483 target proteins.

[7037] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM483 gene, herein designated GAM GENE, on one or more GAM483 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7038] It is yet further appreciated that specific functions, and accordingly utilities, of GAM483 correlate with, and may be deduced from, the identity of the target genes which GAM483 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7039] Nucleotide sequences of the GAM483 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM483 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM483 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM483 are further described hereinbelow with reference to Table Table1.

[7040] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM483 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7041]

[7042] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 484 (GAM484) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7043] GAM484 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM484 was detected is described hereinabove with reference to Figs. 2-8.

[7044] GAM484 gene, herein designated GAM GENE, and GAM484 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7045] GAM484 gene, herein designated GAM GENE, encodes a GAM484 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM484 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM484 precursor RNA is designated SEQ ID:7649, and is provided hereinbelow with reference to the sequence listing part.

[7046] GAM484 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM484 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7047] An enzyme complex designated DICER COMPLEX, dices the GAM484 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM484 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM484 RNA is designated SEQ ID:7650, and is provided hereinbelow with reference to the sequence listing part.

[7048] GAM484 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM484 target RNA, herein designated GAM TARGET RNA. GAM484 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7049] GAM484 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM484 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM484 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM484 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM484 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7050] The complementary binding of GAM484 RNA, herein designated GAM RNA, to target binding sites on GAM484 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM484 target RNA, herein designated GAM TARGET RNA, into GAM484 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7051] It is appreciated that GAM484 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM484 target genes. The mRNA of each one of this plurality of GAM484 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM484 RNA,

herein designated GAM RNA, and which when bound by GAM484 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM484 target proteins.

[7052] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM484 gene, herein designated GAM GENE, on one or more GAM484 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7053] It is yet further appreciated that specific functions, and accordingly utilities, of GAM484 correlate with, and may be deduced from, the identity of the target genes which

GAM484 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7054] Nucleotide sequences of the GAM484 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM484 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM484 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM484 are further described hereinbelow with reference to Table Table1.

[7055] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM484 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7056]

[7057] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 485 (GAM485) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7058] GAM485 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM485 was detected is described hereinabove with reference to Figs. 2-8.

[7059] GAM485 gene, herein designated GAM GENE, and GAM485 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7060] GAM485 gene, herein designated GAM GENE, encodes a GAM485 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM485 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM485 precursor RNA is designated SEQ ID:7667, and is provided hereinbelow with reference to the sequence listing part.

[7061] GAM485 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM485 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[7062] An enzyme complex designated DICER COMPLEX, dices the GAM485 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM485 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM485 RNA is designated SEQ ID:7668, and is provided hereinbelow with reference to the sequence listing part.

[7063] GAM485 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM485 target RNA, herein designated GAM TARGET RNA.
GAM485 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7064] GAM485 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM485 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM485 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM485 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM485 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7065] The complementary binding of GAM485 RNA, herein designated GAM RNA, to target binding sites on GAM485 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM485 target RNA, herein desig-

nated GAM TARGET RNA, into GAM485 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7066] It is appreciated that GAM485 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM485 target genes. The mRNA of each one of this plurality of GAM485 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM485 RNA, herein designated GAM RNA, and which when bound by GAM485 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM485 target proteins.

[7067] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM485 gene, herein designated GAM GENE, on one or more GAM485 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7068] It is yet further appreciated that specific functions, and accordingly utilities, of GAM485 correlate with, and may be deduced from, the identity of the target genes which GAM485 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7069] Nucleotide sequences of the GAM485 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM485 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM485 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM485 are further described hereinbelow with reference to Table Table1.

[7070] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM485 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[7071]

[7072] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 486 (GAM486) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7073] GAM486 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM486 was detected is described hereinabove with reference to Figs. 2-8.

[7074] GAM486 gene, herein designated GAM GENE, and GAM486 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7075] GAM486 gene, herein designated GAM GENE, encodes a GAM486 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM486 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM486 precursor RNA is designated SEQ ID:7677, and is provided hereinbelow with ref-

erence to the sequence listing part.

[7076] GAM486 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM486 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7077] An enzyme complex designated DICER COMPLEX, dices the GAM486 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM486 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM486 RNA is designated SEQ ID:7678, and is provided hereinbelow with reference to the sequence listing part.

[7078] GAM486 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM486 target RNA, herein designated GAM TARGET RNA.

GAM486 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7079] GAM486 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM486 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM486 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM486 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM486 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7080] The complementary binding of GAM486 RNA, herein designated GAM RNA, to target binding sites on GAM486 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM486 target RNA, herein designated GAM TARGET RNA, into GAM486 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7081] It is appreciated that GAM486 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM486 target genes. The mRNA of each one of this plurality of GAM486 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM486 RNA, herein designated GAM RNA, and which when bound by GAM486 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM486 target proteins.

[7082] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM486 gene, herein designated GAM GENE, on one or more GAM486 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7083] It is yet further appreciated that specific functions, and accordingly utilities, of GAM486 correlate with, and may be deduced from, the identity of the target genes which GAM486 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7084] Nucleotide sequences of the GAM486 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM486 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM486 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM486 are further described hereinbelow with reference to Table Table1.

[7085] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM486 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7086]

[7087] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 487 (GAM487) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7088] GAM487 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM487 was detected is described hereinabove with reference to Figs. 2-8.

[7089] GAM487 gene, herein designated GAM GENE, and GAM487 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[7090] GAM487 gene, herein designated GAM GENE, encodes a GAM487 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM487 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM487 precursor RNA is designated SEQ ID:7689, and is provided hereinbelow with reference to the sequence listing part.

[7091] GAM487 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM487 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7092] An enzyme complex designated DICER COMPLEX, dices the GAM487 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM487 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM487 RNA is designated SEQ ID:7690, and is provided hereinbelow with reference to the sequence listing part.

[7093] GAM487 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM487 target RNA, herein designated GAM TARGET RNA. GAM487 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7094] GAM487 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM487 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM487 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM487 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM487 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7095] The complementary binding of GAM487 RNA, herein designated GAM RNA, to target binding sites on GAM487 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM487 target RNA, herein designated GAM TARGET RNA, into GAM487 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7096] It is appreciated that GAM487 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM487 target genes. The mRNA of each one of this plurality of GAM487 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM487 RNA, herein designated GAM RNA, and which when bound by GAM487 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM487 target proteins.

[7097] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM487 gene, herein designated GAM GENE, on one or more GAM487 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[7098] It is yet further appreciated that specific functions, and accordingly utilities, of GAM487 correlate with, and may be deduced from, the identity of the target genes which GAM487 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7099] Nucleotide sequences of the GAM487 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM487 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM487 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM487 are further described hereinbelow with reference to Table Table1.

[7100] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM487 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7101]

[7102] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

488 (GAM488) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7103] GAM488 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM488 was detected is described hereinabove with reference to Figs. 2-8.

[7104] GAM488 gene, herein designated GAM GENE, and GAM488 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7105] GAM488 gene, herein designated GAM GENE, encodes a GAM488 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM488 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM488 precursor RNA is designated SEQ ID:7704, and is provided hereinbelow with reference to the sequence listing part.

[7106] GAM488 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM488 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7107] An enzyme complex designated DICER COMPLEX, dices the GAM488 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM488 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 23%) nucleotide sequence of GAM488 RNA is designated SEQ ID:7705, and is provided hereinbelow with reference to the sequence listing part.

[7108] GAM488 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM488 target RNA, herein designated GAM TARGET RNA. GAM488 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7109] GAM488 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM488 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM488 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM488 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM488 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7110] The complementary binding of GAM488 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM488 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM488 target RNA, herein designated GAM TARGET RNA, into GAM488 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7111] It is appreciated that GAM488 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM488 target genes. The mRNA of each one of this plurality of GAM488 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM488 RNA, herein designated GAM RNA, and which when bound by GAM488 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM488 target proteins.

[7112] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM488 gene, herein designated GAM GENE, on one or more GAM488 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7113] It is yet further appreciated that specific functions, and accordingly utilities, of GAM488 correlate with, and may be deduced from, the identity of the target genes which GAM488 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7114] Nucleotide sequences of the GAM488 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM488 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM488 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM488 are further described hereinbelow with reference to Table Table1.

[7115] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM488 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7116]

[7117] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 489 (GAM489) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7118] GAM489 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM489 was detected is described hereinabove with reference to Figs. 2-8.

[7119] GAM489 gene, herein designated GAM GENE, and GAM489 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7120] GAM489 gene, herein designated GAM GENE, encodes a GAM489 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM489 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM489 precursor RNA is designated SEQ ID:7719, and is provided hereinbelow with reference to the sequence listing part.

[7121] GAM489 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM489 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7122] An enzyme complex designated DICER COMPLEX, dices the GAM489 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM489 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 79%) nucleotide sequence of GAM489 RNA is designated SEQ ID:7720, and is provided hereinbelow with reference to the sequence listing part.

[7123] GAM489 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM489 target RNA, herein designated GAM TARGET RNA. GAM489 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7124] GAM489 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM489 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM489 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM489 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM489 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7125] The complementary binding of GAM489 RNA, herein designated GAM RNA, to target binding sites on GAM489 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM489 target RNA, herein designated GAM TARGET RNA, into GAM489 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7126] It is appreciated that GAM489 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM489 target genes. The mRNA of each one of this plurality of GAM489 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM489 RNA, herein designated GAM RNA, and which when bound by

GAM489 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM489 target proteins.

[7127] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM489 gene, herein designated GAM GENE, on one or more GAM489 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7128] It is yet further appreciated that specific functions, and accordingly utilities, of GAM489 correlate with, and may be deduced from, the identity of the target genes which GAM489 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[7129] Nucleotide sequences of the GAM489 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM489 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM489 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM489 are further described hereinbelow with reference to Table Table1.

[7130] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM489 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7131]

[7132] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 490 (GAM490) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7133] GAM490 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM490 was detected is described hereinabove with reference to Figs. 2-8.

[7134] GAM490 gene, herein designated GAM GENE, and GAM490 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7135] GAM490 gene, herein designated GAM GENE, encodes a GAM490 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM490 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM490 precursor RNA is designated SEQ ID:7736, and is provided hereinbelow with reference to the sequence listing part.

[7136] GAM490 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM490 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[7137] An enzyme complex designated DICER COMPLEX, dices the GAM490 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM490 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM490 RNA is designated SEQ ID:7737, and is provided hereinbelow with reference to the sequence listing part.

[7138] GAM490 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM490 target RNA, herein designated GAM TARGET RNA. GAM490 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7139] GAM490 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM490 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM490 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM490 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM490 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7140] The complementary binding of GAM490 RNA, herein designated GAM RNA, to target binding sites on GAM490 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM490 target RNA, herein designated GAM TARGET RNA, into GAM490 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7141] It is appreciated that GAM490 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM490 target genes. The mRNA of each one of this plurality of GAM490 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM490 RNA, herein designated GAM RNA, and which when bound by GAM490 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM490 target proteins.

[7142] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM490 gene, herein designated GAM GENE, on one or more GAM490 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7143] It is yet further appreciated that specific functions, and accordingly utilities, of GAM490 correlate with, and may be deduced from, the identity of the target genes which GAM490 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7144] Nucleotide sequences of the GAM490 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM490 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM490 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM490 are further described hereinbelow with reference to Table Table1.

[7145] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM490 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7146]

[7147] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 491 (GAM491) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7148] GAM491 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM491 was detected is described hereinabove with reference to Figs. 2-8.

[7149] GAM491 gene, herein designated GAM GENE, and GAM491 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7150] GAM491 gene, herein designated GAM GENE, encodes a GAM491 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM491 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM491 precursor RNA is designated SEQ ID:7753, and is provided hereinbelow with reference to the sequence listing part.

[7151] GAM491 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM491 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7152] An enzyme complex designated DICER COMPLEX, dices

the GAM491 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM491 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM491 RNA is designated SEQ ID:7754, and is provided hereinbelow with reference to the sequence listing part.

[7153] GAM491 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM491 target RNA, herein designated GAM TARGET RNA. GAM491 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7154] GAM491 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM491 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM491

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM491 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM491 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7155] The complementary binding of GAM491 RNA, herein designated GAM RNA, to target binding sites on GAM491 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM491 target RNA, herein designated GAM TARGET RNA, into GAM491 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7156] It is appreciated that GAM491 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM491 target genes. The mRNA of each one of this plurality of GAM491 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM491 RNA, herein designated GAM RNA, and which when bound by GAM491 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM491 target proteins.

[7157] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM491 gene, herein designated GAM GENE, on one or more GAM491 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7158] It is yet further appreciated that specific functions, and accordingly utilities, of GAM491 correlate with, and may be deduced from, the identity of the target genes which GAM491 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7159] Nucleotide sequences of the GAM491 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM491 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM491 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM491 are further described hereinbelow with reference to Table Table1.

[7160] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM491 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7161]

[7162] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 492 (GAM492) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7163] GAM492 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM492 was detected is described hereinabove with reference to Figs. 2-8.

[7164] GAM492 gene, herein designated GAM GENE, and GAM492 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7165] GAM492 gene, herein designated GAM GENE, encodes a GAM492 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM492 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM492 precursor RNA is designated SEQ ID:7778, and is provided hereinbelow with reference to the sequence listing part.

[7166] GAM492 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM492 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7167] An enzyme complex designated DICER COMPLEX, dices the GAM492 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM492 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM492 RNA is designated SEQ ID:7779, and is provided hereinbelow with reference to the sequence listing part.

[7168] GAM492 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM492 target RNA, herein designated GAM TARGET RNA. GAM492 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7169] GAM492 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM492 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM492 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM492 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM492 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[7170] The complementary binding of GAM492 RNA, herein designated GAM RNA, to target binding sites on GAM492 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM492 target RNA, herein designated GAM TARGET RNA, into GAM492 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7171] It is appreciated that GAM492 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM492 target genes. The mRNA of each one of this plurality of GAM492 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM492 RNA, herein designated GAM RNA, and which when bound by GAM492 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM492 target proteins.

[7172] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM492 gene, herein designated GAM GENE, on one or

more GAM492 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7173] It is yet further appreciated that specific functions, and accordingly utilities, of GAM492 correlate with, and may be deduced from, the identity of the target genes which GAM492 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7174] Nucleotide sequences of the GAM492 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM492 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM492 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM492 are further de-

scribed hereinbelow with reference to Table Table1.

[7175] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM492 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7176]

[7177] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 493 (GAM493) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7178] GAM493 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM493 was detected is described hereinabove with reference to Figs. 2-8.

[7179] GAM493 gene, herein designated GAM GENE, and GAM493 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7180] GAM493 gene, herein designated GAM GENE, encodes a GAM493 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM493 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM493 precursor RNA is designated SEQ ID:7803, and is provided hereinbelow with reference to the sequence listing part.

[7181] GAM493 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM493 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7182] An enzyme complex designated DICER COMPLEX, dices the GAM493 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM493 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 22%) nucleotide sequence of GAM493 RNA is designated SEQ ID:7804, and is provided hereinbelow with reference to the sequence listing part.

[7183] GAM493 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM493 target RNA, herein designated GAM TARGET RNA. GAM493 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7184] GAM493 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM493 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM493 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM493 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM493 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7185] The complementary binding of GAM493 RNA, herein designated GAM RNA, to target binding sites on GAM493 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM493 target RNA, herein designated GAM TARGET RNA, into GAM493 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7186] It is appreciated that GAM493 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM493 target genes. The mRNA of each one of this plurality of GAM493 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM493 RNA, herein designated GAM RNA, and which when bound by GAM493 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM493 target proteins.

[7187] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM493 gene, herein designated GAM GENE, on one or more GAM493 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7188] It is yet further appreciated that specific functions, and accordingly utilities, of GAM493 correlate with, and may

be deduced from, the identity of the target genes which GAM493 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7189] Nucleotide sequences of the GAM493 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM493 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM493 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM493 are further described hereinbelow with reference to Table Table1.

[7190] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM493 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7191]

[7192] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 494 (GAM494) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7193] GAM494 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM494 was detected is described hereinabove with reference to Figs. 2-8.

[7194] GAM494 gene, herein designated GAM GENE, and GAM494 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7195] GAM494 gene, herein designated GAM GENE, encodes a GAM494 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM494 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM494 precursor RNA is designated SEQ ID:7821, and is provided hereinbelow with reference to the sequence listing part.

[7196] GAM494 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM494 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7197] An enzyme complex designated DICER COMPLEX, dices the GAM494 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM494 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 76%) nucleotide sequence of GAM494 RNA is designated SEQ ID:7822, and is provided hereinbelow with reference to the sequence listing part.

[7198] GAM494 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM494 target RNA, herein designated GAM TARGET RNA. GAM494 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7199] GAM494 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM494 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM494 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM494 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM494 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7200] The complementary binding of GAM494 RNA, herein designated GAM RNA, to target binding sites on GAM494 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM494 target RNA, herein designated GAM TARGET RNA, into GAM494 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7201] It is appreciated that GAM494 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM494 target genes. The mRNA of each one of this plurality of GAM494 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM494 RNA, herein designated GAM RNA, and which when bound by GAM494 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM494 target proteins.

[7202] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM494 gene, herein designated GAM GENE, on one or more GAM494 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7203] It is yet further appreciated that specific functions, and accordingly utilities, of GAM494 correlate with, and may be deduced from, the identity of the target genes which GAM494 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7204] Nucleotide sequences of the GAM494 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM494 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM494 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM494 are further described hereinbelow with reference to Table Table1.

[7205] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM494 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7206]

[7207] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 495 (GAM495) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7208] GAM495 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM495 was detected is described hereinabove with reference to Figs. 2-8.

[7209] GAM495 gene, herein designated GAM GENE, and GAM495 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7210] GAM495 gene, herein designated GAM GENE, encodes a GAM495 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM495 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM495 precursor RNA is desig-

nated SEQ ID:7835, and is provided hereinbelow with reference to the sequence listing part.

[7211] GAM495 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM495 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7212] An enzyme complex designated DICER COMPLEX, dices the GAM495 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM495 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM495 RNA is designated SEQ ID:7836, and is provided hereinbelow with reference to the sequence listing part.

[7213] GAM495 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM495 target RNA, herein designated GAM TARGET RNA. GAM495 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7214] GAM495 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM495 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM495 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM495 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM495

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7215] The complementary binding of GAM495 RNA, herein designated GAM RNA, to target binding sites on GAM495 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM495 target RNA, herein designated GAM TARGET RNA, into GAM495 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7216] It is appreciated that GAM495 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM495 target genes. The mRNA of each one of this plurality of GAM495 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM495 RNA, herein designated GAM RNA, and which when bound by GAM495 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM495 target proteins.

[7217] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM495 gene, herein designated GAM GENE, on one or more GAM495 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7218] It is yet further appreciated that specific functions, and accordingly utilities, of GAM495 correlate with, and may be deduced from, the identity of the target genes which GAM495 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7219] Nucleotide sequences of the GAM495 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM495 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM495 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM495 are further described hereinbelow with reference to Table Table1.

[7220] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM495 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7221]

[7222] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 496 (GAM496) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7223] GAM496 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM496 was detected is described hereinabove with reference to Figs. 2-8.

[7224] GAM496 gene, herein designated GAM GENE, and

GAM496 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7225] GAM496 gene, herein designated GAM GENE, encodes a GAM496 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM496 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM496 precursor RNA is designated SEQ ID:7839, and is provided hereinbelow with reference to the sequence listing part.

[7226] GAM496 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM496 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7227] An enzyme complex designated DICER COMPLEX, dices the GAM496 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM496 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM496 RNA is designated SEQ ID:7840, and is provided hereinbelow with reference to the sequence listing part.

[7228] GAM496 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM496 target RNA, herein designated GAM TARGET RNA. GAM496 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7229] GAM496 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM496 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM496 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM496 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM496 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7230] The complementary binding of GAM496 RNA, herein designated GAM RNA, to target binding sites on GAM496 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM496 target RNA, herein designated GAM TARGET RNA, into GAM496 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7231] It is appreciated that GAM496 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM496 target genes. The mRNA of each one of this plurality of GAM496 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM496 RNA, herein designated GAM RNA, and which when bound by GAM496 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM496 target proteins.

[7232] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM496 gene, herein designated GAM GENE, on one or more GAM496 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7233] It is yet further appreciated that specific functions, and accordingly utilities, of GAM496 correlate with, and may be deduced from, the identity of the target genes which GAM496 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7234] Nucleotide sequences of the GAM496 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM496 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM496 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM496 are further described hereinbelow with reference to Table Table1.

[7235] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM496 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7236]

[7237] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 497 (GAM497) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7238] GAM497 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM497 was detected is described hereinabove with reference to Figs. 2-8.

[7239] GAM497 gene, herein designated GAM GENE, and GAM497 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7240] GAM497 gene, herein designated GAM GENE, encodes a GAM497 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM497 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM497 precursor RNA is designated SEQ ID:7858, and is provided hereinbelow with reference to the sequence listing part.

[7241] GAM497 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM497 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7242] An enzyme complex designated DICER COMPLEX, dices the GAM497 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM497 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 83%) nucleotide sequence of GAM497 RNA is designated SEQ ID:7859, and is provided hereinbelow with reference to the sequence listing part.

[7243] GAM497 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM497 target RNA, herein designated GAM TARGET RNA. GAM497 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7244] GAM497 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM497 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM497 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM497 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM497 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7245] The complementary binding of GAM497 RNA, herein designated GAM RNA, to target binding sites on GAM497 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM497 target RNA, herein designated GAM TARGET RNA, into GAM497 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7246] It is appreciated that GAM497 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM497 target genes. The mRNA of each one of this plurality of GAM497 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM497 RNA, herein designated GAM RNA, and which when bound by GAM497 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM497 target proteins.

[7247] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM497 gene, herein designated GAM GENE, on one or more GAM497 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7248] It is yet further appreciated that specific functions, and accordingly utilities, of GAM497 correlate with, and may be deduced from, the identity of the target genes which GAM497 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7249] Nucleotide sequences of the GAM497 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM497 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM497 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM497 are further described hereinbelow with reference to Table Table1.

[7250] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM497 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7251]

[7252] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 498 (GAM498) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7253] GAM498 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM498 was detected is described hereinabove with reference to Figs. 2-8.

[7254] GAM498 gene, herein designated GAM GENE, and GAM498 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7255] GAM498 gene, herein designated GAM GENE, encodes a GAM498 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM498 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM498 precursor RNA is designated SEQ ID:7876, and is provided hereinbelow with reference to the sequence listing part.

[7256] GAM498 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM498 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7257] An enzyme complex designated DICER COMPLEX, dices the GAM498 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM498 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM498 RNA is designated SEQ ID:7877, and is provided hereinbelow with reference to the sequence listing part.

[7258] GAM498 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM498 target RNA, herein designated GAM TARGET RNA. GAM498 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7259] GAM498 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM498 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM498 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM498 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM498 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7260] The complementary binding of GAM498 RNA, herein designated GAM RNA, to target binding sites on GAM498 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM498 target RNA, herein designated GAM TARGET RNA, into GAM498 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7261] It is appreciated that GAM498 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM498 target genes. The mRNA of each one of this plurality of GAM498 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM498 RNA,

herein designated GAM RNA, and which when bound by GAM498 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM498 target proteins.

[7262] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM498 gene, herein designated GAM GENE, on one or more GAM498 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7263] It is yet further appreciated that specific functions, and accordingly utilities, of GAM498 correlate with, and may be deduced from, the identity of the target genes which

GAM498 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7264] Nucleotide sequences of the GAM498 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM498 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM498 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM498 are further described hereinbelow with reference to Table Table1.

[7265] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM498 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7266]

[7267] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 499 (GAM499) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7268] GAM499 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM499 was detected is described hereinabove with reference to Figs. 2-8.

[7269] GAM499 gene, herein designated GAM GENE, and GAM499 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7270] GAM499 gene, herein designated GAM GENE, encodes a GAM499 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM499 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM499 precursor RNA is designated SEQ ID:7890, and is provided hereinbelow with reference to the sequence listing part.

[7271] GAM499 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM499 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[7272] An enzyme complex designated DICER COMPLEX, dices the GAM499 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM499 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 61%) nucleotide sequence of GAM499 RNA is designated SEQ ID:7891, and is provided hereinbelow with reference to the sequence listing part.

[7273] GAM499 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM499 target RNA, herein designated GAM TARGET RNA.
GAM499 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7274] GAM499 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM499 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM499 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM499 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM499 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7275] The complementary binding of GAM499 RNA, herein designated GAM RNA, to target binding sites on GAM499 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM499 target RNA, herein desig-

nated GAM TARGET RNA, into GAM499 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7276] It is appreciated that GAM499 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM499 target genes. The mRNA of each one of this plurality of GAM499 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM499 RNA, herein designated GAM RNA, and which when bound by GAM499 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM499 target proteins.

[7277] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM499 gene, herein designated GAM GENE, on one or more GAM499 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7278] It is yet further appreciated that specific functions, and accordingly utilities, of GAM499 correlate with, and may be deduced from, the identity of the target genes which GAM499 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7279] Nucleotide sequences of the GAM499 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM499 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM499 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM499 are further described hereinbelow with reference to Table Table1.

[7280] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM499 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[7281]

[7282] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 500 (GAM500) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7283] GAM500 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM500 was detected is described hereinabove with reference to Figs. 2-8.

[7284] GAM500 gene, herein designated GAM GENE, and GAM500 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7285] GAM500 gene, herein designated GAM GENE, encodes a GAM500 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM500 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM500 precursor RNA is designated SEQ ID:7896, and is provided hereinbelow with ref-

erence to the sequence listing part.

[7286] GAM500 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM500 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7287] An enzyme complex designated DICER COMPLEX, dices the GAM500 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM500 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM500 RNA is designated SEQ ID:7897, and is provided hereinbelow with reference to the sequence listing part.

[7288] GAM500 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM500 target RNA, herein designated GAM TARGET RNA.

GAM500 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7289] GAM500 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM500 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM500 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM500 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM500 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7290] The complementary binding of GAM500 RNA, herein designated GAM RNA, to target binding sites on GAM500 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM500 target RNA, herein designated GAM TARGET RNA, into GAM500 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7291] It is appreciated that GAM500 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM500 target genes. The mRNA of each one of this plurality of GAM500 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM500 RNA, herein designated GAM RNA, and which when bound by GAM500 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM500 target proteins.

[7292] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM500 gene, herein designated GAM GENE, on one or more GAM500 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7293] It is yet further appreciated that specific functions, and accordingly utilities, of GAM500 correlate with, and may be deduced from, the identity of the target genes which GAM500 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7294] Nucleotide sequences of the GAM500 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM500 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM500 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM500 are further described hereinbelow with reference to Table Table1.

[7295] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM500 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7296]

[7297] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 501 (GAM501) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7298] GAM501 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM501 was detected is described hereinabove with reference to Figs. 2-8.

[7299] GAM501 gene, herein designated GAM GENE, and GAM501 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[7300] GAM501 gene, herein designated GAM GENE, encodes a GAM501 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM501 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM501 precursor RNA is designated SEQ ID:7909, and is provided hereinbelow with reference to the sequence listing part.

[7301] GAM501 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM501 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7302] An enzyme complex designated DICER COMPLEX, dices the GAM501 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM501 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 23%) nucleotide sequence of GAM501 RNA is designated SEQ ID:7910, and is provided hereinbelow with reference to the sequence listing part.

[7303] GAM501 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM501 target RNA, herein designated GAM TARGET RNA. GAM501 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7304] GAM501 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM501 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM501 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM501 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM501 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7305] The complementary binding of GAM501 RNA, herein designated GAM RNA, to target binding sites on GAM501 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM501 target RNA, herein designated GAM TARGET RNA, into GAM501 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7306] It is appreciated that GAM501 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM501 target genes. The mRNA of each one of this plurality of GAM501 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM501 RNA, herein designated GAM RNA, and which when bound by GAM501 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM501 target proteins.

[7307] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM501 gene, herein designated GAM GENE, on one or more GAM501 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[7308] It is yet further appreciated that specific functions, and accordingly utilities, of GAM501 correlate with, and may be deduced from, the identity of the target genes which GAM501 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7309] Nucleotide sequences of the GAM501 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM501 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM501 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM501 are further described hereinbelow with reference to Table Table1.

[7310] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM501 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7311]

[7312] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

502 (GAM502) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7313] GAM502 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM502 was detected is described hereinabove with reference to Figs. 2-8.

[7314] GAM502 gene, herein designated GAM GENE, and GAM502 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7315] GAM502 gene, herein designated GAM GENE, encodes a GAM502 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM502 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM502 precursor RNA is designated SEQ ID:7918, and is provided hereinbelow with reference to the sequence listing part.

[7316] GAM502 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM502 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7317] An enzyme complex designated DICER COMPLEX, dices the GAM502 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM502 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM502 RNA is designated SEQ ID:7919, and is provided hereinbelow with reference to the sequence listing part.

[7318] GAM502 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM502 target RNA, herein designated GAM TARGET RNA. GAM502 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7319] GAM502 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM502 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM502 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM502 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM502 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7320] The complementary binding of GAM502 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM502 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM502 target RNA, herein designated GAM TARGET RNA, into GAM502 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7321] It is appreciated that GAM502 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM502 target genes. The mRNA of each one of this plurality of GAM502 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM502 RNA, herein designated GAM RNA, and which when bound by GAM502 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM502 target proteins.

[7322] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM502 gene, herein designated GAM GENE, on one or more GAM502 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7323] It is yet further appreciated that specific functions, and accordingly utilities, of GAM502 correlate with, and may be deduced from, the identity of the target genes which GAM502 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7324] Nucleotide sequences of the GAM502 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM502 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM502 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM502 are further described hereinbelow with reference to Table Table1.

[7325] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM502 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7326]

[7327] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 503 (GAM503) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7328] GAM503 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM503 was detected is described hereinabove with reference to Figs. 2-8.

[7329] GAM503 gene, herein designated GAM GENE, and GAM503 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7330] GAM503 gene, herein designated GAM GENE, encodes a GAM503 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM503 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM503 precursor RNA is designated SEQ ID:7924, and is provided hereinbelow with reference to the sequence listing part.

[7331] GAM503 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM503 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7332] An enzyme complex designated DICER COMPLEX, dices the GAM503 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM503 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 82%) nucleotide sequence of GAM503 RNA is designated SEQ ID:7925, and is provided hereinbelow with reference to the sequence listing part.

[7333] GAM503 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM503 target RNA, herein designated GAM TARGET RNA. GAM503 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7334] GAM503 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM503 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM503 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM503 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM503 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7335] The complementary binding of GAM503 RNA, herein designated GAM RNA, to target binding sites on GAM503 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM503 target RNA, herein designated GAM TARGET RNA, into GAM503 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7336] It is appreciated that GAM503 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM503 target genes. The mRNA of each one of this plurality of GAM503 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM503 RNA, herein designated GAM RNA, and which when bound by

GAM503 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM503 target proteins.

[7337] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM503 gene, herein designated GAM GENE, on one or more GAM503 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7338] It is yet further appreciated that specific functions, and accordingly utilities, of GAM503 correlate with, and may be deduced from, the identity of the target genes which GAM503 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[7339] Nucleotide sequences of the GAM503 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM503 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM503 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM503 are further described hereinbelow with reference to Table Table1.

[7340] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM503 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7341]

[7342] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 504 (GAM504) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7343] GAM504 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM504 was detected is described hereinabove with reference to Figs. 2-8.

[7344] GAM504 gene, herein designated GAM GENE, and GAM504 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7345] GAM504 gene, herein designated GAM GENE, encodes a GAM504 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM504 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM504 precursor RNA is designated SEQ ID:7952, and is provided hereinbelow with reference to the sequence listing part.

[7346] GAM504 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM504 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[7347] An enzyme complex designated DICER COMPLEX, dices the GAM504 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM504 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM504 RNA is designated SEQ ID:7953, and is provided hereinbelow with reference to the sequence listing part.

[7348] GAM504 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM504 target RNA, herein designated GAM TARGET RNA. GAM504 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7349] GAM504 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM504 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM504 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM504 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM504 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7350] The complementary binding of GAM504 RNA, herein designated GAM RNA, to target binding sites on GAM504 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM504 target RNA, herein designated GAM TARGET RNA, into GAM504 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7351] It is appreciated that GAM504 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM504 target genes. The mRNA of each one of this plurality of GAM504 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM504 RNA, herein designated GAM RNA, and which when bound by GAM504 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM504 target proteins.

[7352] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM504 gene, herein designated GAM GENE, on one or more GAM504 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7353] It is yet further appreciated that specific functions, and accordingly utilities, of GAM504 correlate with, and may be deduced from, the identity of the target genes which GAM504 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7354] Nucleotide sequences of the GAM504 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM504 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM504 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM504 are further described hereinbelow with reference to Table Table1.

[7355] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM504 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7356]

[7357] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 505 (GAM505) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7358] GAM505 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM505 was detected is described hereinabove with reference to Figs. 2-8.

[7359] GAM505 gene, herein designated GAM GENE, and GAM505 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7360] GAM505 gene, herein designated GAM GENE, encodes a GAM505 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM505 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM505 precursor RNA is designated SEQ ID:7992, and is provided hereinbelow with reference to the sequence listing part.

[7361] GAM505 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM505 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7362] An enzyme complex designated DICER COMPLEX, dices the GAM505 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM505 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM505 RNA is designated SEQ ID:7993, and is provided hereinbelow with reference to the sequence listing part.

[7363] GAM505 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM505

target RNA, herein designated GAM TARGET RNA.

GAM505 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7364] GAM505 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM505 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM505 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM505 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM505 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7365] The complementary binding of GAM505 RNA, herein designated GAM RNA, to target binding sites on GAM505 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM505 target RNA, herein designated GAM TARGET RNA, into GAM505 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7366] It is appreciated that GAM505 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM505 target genes. The mRNA of each one of this plurality of GAM505 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM505 RNA, herein designated GAM RNA, and which when bound by GAM505 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM505 target proteins.

[7367] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM505 gene, herein designated GAM GENE, on one or more GAM505 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7368] It is yet further appreciated that specific functions, and accordingly utilities, of GAM505 correlate with, and may be deduced from, the identity of the target genes which GAM505 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7369] Nucleotide sequences of the GAM505 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM505 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM505 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM505 are further described hereinbelow with reference to Table Table1.

[7370] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM505 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7371]

[7372] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 506 (GAM506) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7373] GAM506 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM506 was detected is described hereinabove with reference to Figs. 2-8.

[7374] GAM506 gene, herein designated GAM GENE, and GAM506 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7375] GAM506 gene, herein designated GAM GENE, encodes a GAM506 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM506 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM506 precursor RNA is designated SEQ ID:8026, and is provided hereinbelow with reference to the sequence listing part.

[7376] GAM506 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM506 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7377] An enzyme complex designated DICER COMPLEX, dices the GAM506 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM506 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 23%) nucleotide sequence of GAM506 RNA is designated SEQ ID:8027, and is provided hereinbelow with reference to the sequence listing part.

[7378] GAM506 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM506 target RNA, herein designated GAM TARGET RNA. GAM506 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7379] GAM506 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM506 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM506 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM506 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM506 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7380] The complementary binding of GAM506 RNA, herein designated GAM RNA, to target binding sites on GAM506 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM506 target RNA, herein designated GAM TARGET RNA, into GAM506 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7381] It is appreciated that GAM506 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM506 target genes. The mRNA of each one of this plu-

rality of GAM506 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM506 RNA, herein designated GAM RNA, and which when bound by GAM506 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM506 target proteins.

[7382] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM506 gene, herein designated GAM GENE, on one or more GAM506 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7383] It is yet further appreciated that specific functions, and accordingly utilities, of GAM506 correlate with, and may be deduced from, the identity of the target genes which GAM506 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7384] Nucleotide sequences of the GAM506 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM506 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM506 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM506 are further described hereinbelow with reference to Table Table1.

[7385] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM506 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7386]

[7387] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 507 (GAM507) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[7388] GAM507 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM507 was detected is described hereinabove with reference to Figs. 2-8.

[7389] GAM507 gene, herein designated GAM GENE, and GAM507 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7390] GAM507 gene, herein designated GAM GENE, encodes a GAM507 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM507 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM507 precursor RNA is designated SEQ ID:8051, and is provided hereinbelow with reference to the sequence listing part.

[7391] GAM507 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM507 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7392] An enzyme complex designated DICER COMPLEX, dices the GAM507 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM507 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM507 RNA is designated SEQ ID:8052, and is provided hereinbelow with reference to the sequence listing part.

[7393] GAM507 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM507 target RNA, herein designated GAM TARGET RNA.
GAM507 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[7394] GAM507 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM507 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM507 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM507 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM507 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7395] The complementary binding of GAM507 RNA, herein designated GAM RNA, to target binding sites on GAM507 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM507 target RNA, herein designated GAM TARGET RNA, into GAM507 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7396] It is appreciated that GAM507 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM507 target genes. The mRNA of each one of this plurality of GAM507 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM507 RNA, herein designated GAM RNA, and which when bound by GAM507 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM507 target proteins.

[7397] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM507 gene, herein designated GAM GENE, on one or more GAM507 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7398] It is yet further appreciated that specific functions, and accordingly utilities, of GAM507 correlate with, and may be deduced from, the identity of the target genes which GAM507 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7399] Nucleotide sequences of the GAM507 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM507 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM507 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM507 are further described hereinbelow with reference to Table Table1.

[7400] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM507 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7401]

[7402] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 508 (GAM508) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7403] GAM508 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM508 was detected is described hereinabove with reference to Figs. 2-8.

[7404] GAM508 gene, herein designated GAM GENE, and GAM508 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7405] GAM508 gene, herein designated GAM GENE, encodes a GAM508 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM508 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM508 precursor RNA is designated SEQ ID:8071, and is provided hereinbelow with reference to the sequence listing part.

[7406] GAM508 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM508 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7407] An enzyme complex designated DICER COMPLEX, dices the GAM508 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM508 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of

GAM508 RNA is designated SEQ ID:8072, and is provided hereinbelow with reference to the sequence listing part.

[7408] GAM508 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM508 target RNA, herein designated GAM TARGET RNA. GAM508 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7409] GAM508 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM508 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM508 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM508 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM508 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7410] The complementary binding of GAM508 RNA, herein designated GAM RNA, to target binding sites on GAM508 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM508 target RNA, herein designated GAM TARGET RNA, into GAM508 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7411] It is appreciated that GAM508 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM508 target genes. The mRNA of each one of this plurality of GAM508 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM508 RNA, herein designated GAM RNA, and which when bound by GAM508 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM508 target proteins.

[7412] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM508 gene, herein designated GAM GENE, on one or more GAM508 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7413] It is yet further appreciated that specific functions, and accordingly utilities, of GAM508 correlate with, and may be deduced from, the identity of the target genes which GAM508 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7414] Nucleotide sequences of the GAM508 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM508 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM508 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM508 are further described hereinbelow with reference to Table Table1.

[7415] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM508 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7416]

[7417] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 509 (GAM509) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7418] GAM509 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM509 was detected is described hereinabove

with reference to Figs. 2-8.

[7419] GAM509 gene, herein designated GAM GENE, and GAM509 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7420] GAM509 gene, herein designated GAM GENE, encodes a GAM509 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM509 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM509 precursor RNA is designated SEQ ID:8091, and is provided hereinbelow with reference to the sequence listing part.

[7421] GAM509 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM509 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7422] An enzyme complex designated DICER COMPLEX, dices the GAM509 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM509 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM509 RNA is designated SEQ ID:8092, and is provided hereinbelow with reference to the sequence listing part.

[7423] GAM509 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM509 target RNA, herein designated GAM TARGET RNA. GAM509 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7424] GAM509 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM509 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM509 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM509 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM509 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7425] The complementary binding of GAM509 RNA, herein designated GAM RNA, to target binding sites on GAM509 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM509 target RNA, herein designated GAM TARGET RNA, into GAM509 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[7426] It is appreciated that GAM509 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM509 target genes. The mRNA of each one of this plurality of GAM509 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM509 RNA, herein designated GAM RNA, and which when bound by GAM509 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM509 target proteins.

[7427] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM509 gene, herein designated GAM GENE, on one or more GAM509 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7428] It is yet further appreciated that specific functions, and accordingly utilities, of GAM509 correlate with, and may be deduced from, the identity of the target genes which GAM509 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7429] Nucleotide sequences of the GAM509 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM509 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM509 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM509 are further described hereinbelow with reference to Table Table1.

[7430] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM509 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7431]

[7432] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 510 (GAM510) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7433] GAM510 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM510 was detected is described hereinabove with reference to Figs. 2-8.

[7434] GAM510 gene, herein designated GAM GENE, and GAM510 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7435] GAM510 gene, herein designated GAM GENE, encodes a GAM510 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM510 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM510 precursor RNA is designated SEQ ID:8109, and is provided hereinbelow with reference to the sequence listing part.

[7436] GAM510 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM510 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7437] An enzyme complex designated DICER COMPLEX, dices the GAM510 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM510 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 33%) nucleotide sequence of GAM510 RNA is designated SEQ ID:8110, and is provided hereinbelow with reference to the sequence listing part.

[7438] GAM510 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM510 target RNA, herein designated GAM TARGET RNA.

GAM510 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7439] GAM510 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM510 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM510 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM510 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM510 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7440] The complementary binding of GAM510 RNA, herein designated GAM RNA, to target binding sites on GAM510 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM510 target RNA, herein designated GAM TARGET RNA, into GAM510 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7441] It is appreciated that GAM510 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM510 target genes. The mRNA of each one of this plurality of GAM510 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM510 RNA, herein designated GAM RNA, and which when bound by GAM510 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM510 target proteins.

[7442] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM510 gene, herein designated GAM GENE, on one or more GAM510 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7443] It is yet further appreciated that specific functions, and accordingly utilities, of GAM510 correlate with, and may be deduced from, the identity of the target genes which GAM510 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7444] Nucleotide sequences of the GAM510 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM510 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM510 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM510 are further described hereinbelow with reference to Table Table1.

[7445] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM510 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7446]

[7447] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 511 (GAM511) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7448] GAM511 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM511 was detected is described hereinabove with reference to Figs. 2-8.

[7449] GAM511 gene, herein designated GAM GENE, and GAM511 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7450] GAM511 gene, herein designated GAM GENE, encodes a

GAM511 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM511 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM511 precursor RNA is designated SEQ ID:8136, and is provided hereinbelow with reference to the sequence listing part.

[7451] GAM511 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM511 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7452] An enzyme complex designated DICER COMPLEX, dices the GAM511 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM511 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM511 RNA is designated SEQ ID:8137, and is provided hereinbelow with reference to the sequence listing part.

[7453] GAM511 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM511 target RNA, herein designated GAM TARGET RNA. GAM511 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7454] GAM511 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM511 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM511 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM511 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM511 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7455] The complementary binding of GAM511 RNA, herein designated GAM RNA, to target binding sites on GAM511 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM511 target RNA, herein designated GAM TARGET RNA, into GAM511 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7456] It is appreciated that GAM511 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM511 target genes. The mRNA of each one of this plurality of GAM511 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM511 RNA, herein designated GAM RNA, and which when bound by GAM511 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM511 target proteins.

[7457] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM511 gene, herein designated GAM GENE, on one or more GAM511 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7458] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM511 correlate with, and may be deduced from, the identity of the target genes which GAM511 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7459] Nucleotide sequences of the GAM511 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM511 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM511 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM511 are further described hereinbelow with reference to Table Table1.

[7460] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM511 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7461]

[7462] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 512 (GAM512) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[7463] GAM512 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM512 was detected is described hereinabove with reference to Figs. 2-8.

[7464] GAM512 gene, herein designated GAM GENE, and GAM512 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7465] GAM512 gene, herein designated GAM GENE, encodes a GAM512 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM512 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM512 precursor RNA is designated SEQ ID:8149, and is provided hereinbelow with reference to the sequence listing part.

[7466] GAM512 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM512 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7467] An enzyme complex designated DICER COMPLEX, dices the GAM512 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM512 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 68%) nucleotide sequence of GAM512 RNA is designated SEQ ID:8150, and is provided hereinbelow with reference to the sequence listing part.

[7468] GAM512 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM512 target RNA, herein designated GAM TARGET RNA. GAM512 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7469] GAM512 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM512 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM512 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM512 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM512 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7470] The complementary binding of GAM512 RNA, herein designated GAM RNA, to target binding sites on GAM512 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM512 target RNA, herein designated GAM TARGET RNA, into GAM512 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7471] It is appreciated that GAM512 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM512 target genes. The mRNA of each one of this plurality of GAM512 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM512 RNA, herein designated GAM RNA, and which when bound by GAM512 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM512 target proteins.

[7472] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM512 gene, herein designated GAM GENE, on one or more GAM512 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7473] It is yet further appreciated that specific functions, and accordingly utilities, of GAM512 correlate with, and may be deduced from, the identity of the target genes which GAM512 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7474] Nucleotide sequences of the GAM512 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM512 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM512 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM512 are further described hereinbelow with reference to Table Table1.

[7475] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM512 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7476]

[7477] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 513 (GAM513) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7478] GAM513 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM513 was detected is described hereinabove with reference to Figs. 2-8.

[7479] GAM513 gene, herein designated GAM GENE, and GAM513 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7480] GAM513 gene, herein designated GAM GENE, encodes a GAM513 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM513 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM513 precursor RNA is designated SEQ ID:8162, and is provided hereinbelow with reference to the sequence listing part.

[7481] GAM513 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM513 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7482] An enzyme complex designated DICER COMPLEX, dices the GAM513 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM513 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 80%) nucleotide sequence of GAM513 RNA is designated SEQ ID:8163, and is provided

hereinbelow with reference to the sequence listing part.

[7483] GAM513 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM513 target RNA, herein designated GAM TARGET RNA. GAM513 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7484] GAM513 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM513 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM513 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM513 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM513 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7485] The complementary binding of GAM513 RNA, herein designated GAM RNA, to target binding sites on GAM513 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM513 target RNA, herein designated GAM TARGET RNA, into GAM513 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7486] It is appreciated that GAM513 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM513 target genes. The mRNA of each one of this plurality of GAM513 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM513 RNA, herein designated GAM RNA, and which when bound by GAM513 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM513 tar-

get proteins.

[7487] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM513 gene, herein designated GAM GENE, on one or more GAM513 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7488] It is yet further appreciated that specific functions, and accordingly utilities, of GAM513 correlate with, and may be deduced from, the identity of the target genes which GAM513 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7489] Nucleotide sequences of the GAM513 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM513 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM513 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM513 are further described hereinbelow with reference to Table Table1.

[7490] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM513 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7491]

[7492] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 514 (GAM514) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7493] GAM514 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM514 was detected is described hereinabove with reference to Figs. 2-8.

[7494] GAM514 gene, herein designated GAM GENE, and GAM514 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7495] GAM514 gene, herein designated GAM GENE, encodes a GAM514 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM514 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM514 precursor RNA is designated SEQ ID:8187, and is provided hereinbelow with reference to the sequence listing part.

[7496] GAM514 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM514 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7497] An enzyme complex designated DICER COMPLEX, dices

the GAM514 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM514 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 78%) nucleotide sequence of GAM514 RNA is designated SEQ ID:8188, and is provided hereinbelow with reference to the sequence listing part.

[7498] GAM514 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM514 target RNA, herein designated GAM TARGET RNA. GAM514 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7499] GAM514 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM514 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM514

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM514 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM514 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7500] The complementary binding of GAM514 RNA, herein designated GAM RNA, to target binding sites on GAM514 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM514 target RNA, herein designated GAM TARGET RNA, into GAM514 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7501] It is appreciated that GAM514 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM514 target genes. The mRNA of each one of this plurality of GAM514 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM514 RNA, herein designated GAM RNA, and which when bound by GAM514 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM514 target proteins.

[7502] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM514 gene, herein designated GAM GENE, on one or more GAM514 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7503] It is yet further appreciated that specific functions, and accordingly utilities, of GAM514 correlate with, and may be deduced from, the identity of the target genes which GAM514 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7504] Nucleotide sequences of the GAM514 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM514 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM514 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM514 are further described hereinbelow with reference to Table Table1.

[7505] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM514 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7506]

[7507] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 515 (GAM515) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7508] GAM515 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM515 was detected is described hereinabove with reference to Figs. 2-8.

[7509] GAM515 gene, herein designated GAM GENE, and GAM515 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7510] GAM515 gene, herein designated GAM GENE, encodes a GAM515 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM515 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM515 precursor RNA is designated SEQ ID:8210, and is provided hereinbelow with reference to the sequence listing part.

[7511] GAM515 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM515 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7512] An enzyme complex designated DICER COMPLEX, dices the GAM515 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM515 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 50%) nucleotide sequence of GAM515 RNA is designated SEQ ID:8211, and is provided hereinbelow with reference to the sequence listing part.

[7513] GAM515 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM515 target RNA, herein designated GAM TARGET RNA. GAM515 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7514] GAM515 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM515 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM515 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM515 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM515 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[7515] The complementary binding of GAM515 RNA, herein designated GAM RNA, to target binding sites on GAM515 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM515 target RNA, herein designated GAM TARGET RNA, into GAM515 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7516] It is appreciated that GAM515 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM515 target genes. The mRNA of each one of this plurality of GAM515 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM515 RNA, herein designated GAM RNA, and which when bound by GAM515 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM515 target proteins.

[7517] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM515 gene, herein designated GAM GENE, on one or

more GAM515 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7518] It is yet further appreciated that specific functions, and accordingly utilities, of GAM515 correlate with, and may be deduced from, the identity of the target genes which GAM515 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7519] Nucleotide sequences of the GAM515 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM515 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM515 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM515 are further de-

scribed hereinbelow with reference to Table Table1.

[7520] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM515 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7521]

[7522] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 516 (GAM516) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7523] GAM516 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM516 was detected is described hereinabove with reference to Figs. 2-8.

[7524] GAM516 gene, herein designated GAM GENE, and GAM516 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7525] GAM516 gene, herein designated GAM GENE, encodes a GAM516 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM516 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM516 precursor RNA is designated SEQ ID:8218, and is provided hereinbelow with reference to the sequence listing part.

[7526] GAM516 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM516 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7527] An enzyme complex designated DICER COMPLEX, dices the GAM516 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM516 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM516 RNA is designated SEQ ID:8219, and is provided hereinbelow with reference to the sequence listing part.

[7528] GAM516 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM516 target RNA, herein designated GAM TARGET RNA. GAM516 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7529] GAM516 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM516 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM516 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM516 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM516 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7530] The complementary binding of GAM516 RNA, herein designated GAM RNA, to target binding sites on GAM516 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM516 target RNA, herein designated GAM TARGET RNA, into GAM516 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7531] It is appreciated that GAM516 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM516 target genes. The mRNA of each one of this plurality of GAM516 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM516 RNA, herein designated GAM RNA, and which when bound by GAM516 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM516 target proteins.

[7532] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM516 gene, herein designated GAM GENE, on one or more GAM516 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7533] It is yet further appreciated that specific functions, and accordingly utilities, of GAM516 correlate with, and may

be deduced from, the identity of the target genes which GAM516 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7534] Nucleotide sequences of the GAM516 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM516 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM516 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM516 are further described hereinbelow with reference to Table Table1.

[7535] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM516 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7536]

[7537] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 517 (GAM517) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7538] GAM517 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM517 was detected is described hereinabove with reference to Figs. 2-8.

[7539] GAM517 gene, herein designated GAM GENE, and GAM517 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7540] GAM517 gene, herein designated GAM GENE, encodes a GAM517 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM517 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM517 precursor RNA is designated SEQ ID:8236, and is provided hereinbelow with reference to the sequence listing part.

[7541] GAM517 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM517 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7542] An enzyme complex designated DICER COMPLEX, dices the GAM517 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM517 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 63%) nucleotide sequence of GAM517 RNA is designated SEQ ID:8237, and is provided hereinbelow with reference to the sequence listing part.

[7543] GAM517 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM517 target RNA, herein designated GAM TARGET RNA. GAM517 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7544] GAM517 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM517 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM517 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM517 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM517 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7545] The complementary binding of GAM517 RNA, herein designated GAM RNA, to target binding sites on GAM517 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM517 target RNA, herein designated GAM TARGET RNA, into GAM517 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7546] It is appreciated that GAM517 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM517 target genes. The mRNA of each one of this plurality of GAM517 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM517 RNA, herein designated GAM RNA, and which when bound by GAM517 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM517 target proteins.

[7547] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM517 gene, herein designated GAM GENE, on one or more GAM517 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7548] It is yet further appreciated that specific functions, and accordingly utilities, of GAM517 correlate with, and may be deduced from, the identity of the target genes which GAM517 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7549] Nucleotide sequences of the GAM517 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM517 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM517 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM517 are further described hereinbelow with reference to Table Table1.

[7550] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM517 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7551]

[7552] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 518 (GAM518) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7553] GAM518 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM518 was detected is described hereinabove with reference to Figs. 2-8.

[7554] GAM518 gene, herein designated GAM GENE, and GAM518 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7555] GAM518 gene, herein designated GAM GENE, encodes a GAM518 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM518 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM518 precursor RNA is desig-

nated SEQ ID:8254, and is provided hereinbelow with reference to the sequence listing part.

[7556] GAM518 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM518 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7557] An enzyme complex designated DICER COMPLEX, dices the GAM518 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM518 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 75%) nucleotide sequence of GAM518 RNA is designated SEQ ID:8255, and is provided hereinbelow with reference to the sequence listing part.

[7558] GAM518 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM518 target RNA, herein designated GAM TARGET RNA. GAM518 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7559] GAM518 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM518 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM518 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM518 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM518

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7560] The complementary binding of GAM518 RNA, herein designated GAM RNA, to target binding sites on GAM518 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM518 target RNA, herein designated GAM TARGET RNA, into GAM518 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7561] It is appreciated that GAM518 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM518 target genes. The mRNA of each one of this plurality of GAM518 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM518 RNA, herein designated GAM RNA, and which when bound by GAM518 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM518 target proteins.

[7562] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM518 gene, herein designated GAM GENE, on one or more GAM518 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7563] It is yet further appreciated that specific functions, and accordingly utilities, of GAM518 correlate with, and may be deduced from, the identity of the target genes which GAM518 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7564] Nucleotide sequences of the GAM518 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM518 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM518 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM518 are further described hereinbelow with reference to Table Table1.

[7565] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM518 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7566]

[7567] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 519 (GAM519) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7568] GAM519 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM519 was detected is described hereinabove with reference to Figs. 2-8.

[7569] GAM519 gene, herein designated GAM GENE, and

GAM519 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7570] GAM519 gene, herein designated GAM GENE, encodes a GAM519 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM519 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM519 precursor RNA is designated SEQ ID:8272, and is provided hereinbelow with reference to the sequence listing part.

[7571] GAM519 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM519 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7572] An enzyme complex designated DICER COMPLEX, dices the GAM519 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM519 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 57%) nucleotide sequence of GAM519 RNA is designated SEQ ID:8273, and is provided hereinbelow with reference to the sequence listing part.

[7573] GAM519 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM519 target RNA, herein designated GAM TARGET RNA. GAM519 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7574] GAM519 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM519 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM519 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM519 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM519 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7575] The complementary binding of GAM519 RNA, herein designated GAM RNA, to target binding sites on GAM519 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM519 target RNA, herein designated GAM TARGET RNA, into GAM519 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7576] It is appreciated that GAM519 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM519 target genes. The mRNA of each one of this plurality of GAM519 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM519 RNA, herein designated GAM RNA, and which when bound by GAM519 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM519 target proteins.

[7577] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM519 gene, herein designated GAM GENE, on one or more GAM519 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7578] It is yet further appreciated that specific functions, and accordingly utilities, of GAM519 correlate with, and may be deduced from, the identity of the target genes which GAM519 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7579] Nucleotide sequences of the GAM519 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM519 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM519 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM519 are further described hereinbelow with reference to Table Table1.

[7580] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM519 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7581]

[7582] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 520 (GAM520) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7583] GAM520 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM520 was detected is described hereinabove with reference to Figs. 2-8.

[7584] GAM520 gene, herein designated GAM GENE, and GAM520 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7585] GAM520 gene, herein designated GAM GENE, encodes a GAM520 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM520 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM520 precursor RNA is designated SEQ ID:8283, and is provided hereinbelow with reference to the sequence listing part.

[7586] GAM520 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM520 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7587] An enzyme complex designated DICER COMPLEX, dices the GAM520 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM520 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 29%) nucleotide sequence of GAM520 RNA is designated SEQ ID:8284, and is provided hereinbelow with reference to the sequence listing part.

[7588] GAM520 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM520 target RNA, herein designated GAM TARGET RNA. GAM520 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7589] GAM520 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM520 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM520 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM520 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM520 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7590] The complementary binding of GAM520 RNA, herein designated GAM RNA, to target binding sites on GAM520 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM520 target RNA, herein designated GAM TARGET RNA, into GAM520 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7591] It is appreciated that GAM520 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM520 target genes. The mRNA of each one of this plurality of GAM520 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM520 RNA, herein designated GAM RNA, and which when bound by GAM520 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM520 target proteins.

[7592] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM520 gene, herein designated GAM GENE, on one or more GAM520 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7593] It is yet further appreciated that specific functions, and accordingly utilities, of GAM520 correlate with, and may be deduced from, the identity of the target genes which GAM520 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7594] Nucleotide sequences of the GAM520 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM520 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM520 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM520 are further described hereinbelow with reference to Table Table1.

[7595] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM520 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7596]

[7597] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 521 (GAM521) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7598] GAM521 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM521 was detected is described hereinabove with reference to Figs. 2-8.

[7599] GAM521 gene, herein designated GAM GENE, and GAM521 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7600] GAM521 gene, herein designated GAM GENE, encodes a GAM521 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM521 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM521 precursor RNA is designated SEQ ID:8300, and is provided hereinbelow with reference to the sequence listing part.

[7601] GAM521 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM521 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7602] An enzyme complex designated DICER COMPLEX, dices the GAM521 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM521 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM521 RNA is designated SEQ ID:8301, and is provided hereinbelow with reference to the sequence listing part.

[7603] GAM521 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM521 target RNA, herein designated GAM TARGET RNA. GAM521 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7604] GAM521 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM521 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM521 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM521 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM521 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7605] The complementary binding of GAM521 RNA, herein designated GAM RNA, to target binding sites on GAM521 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM521 target RNA, herein designated GAM TARGET RNA, into GAM521 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7606] It is appreciated that GAM521 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM521 target genes. The mRNA of each one of this plurality of GAM521 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM521 RNA,

herein designated GAM RNA, and which when bound by GAM521 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM521 target proteins.

[7607] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM521 gene, herein designated GAM GENE, on one or more GAM521 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7608] It is yet further appreciated that specific functions, and accordingly utilities, of GAM521 correlate with, and may be deduced from, the identity of the target genes which

GAM521 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7609] Nucleotide sequences of the GAM521 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM521 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM521 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM521 are further described hereinbelow with reference to Table Table1.

[7610] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM521 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7611]

[7612] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 522 (GAM522) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7613] GAM522 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM522 was detected is described hereinabove with reference to Figs. 2-8.

[7614] GAM522 gene, herein designated GAM GENE, and GAM522 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7615] GAM522 gene, herein designated GAM GENE, encodes a GAM522 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM522 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM522 precursor RNA is designated SEQ ID:8319, and is provided hereinbelow with reference to the sequence listing part.

[7616] GAM522 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM522 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[7617] An enzyme complex designated DICER COMPLEX, dices the GAM522 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM522 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 50%) nucleotide sequence of GAM522 RNA is designated SEQ ID:8320, and is provided hereinbelow with reference to the sequence listing part.

[7618] GAM522 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM522 target RNA, herein designated GAM TARGET RNA.
GAM522 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7619] GAM522 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM522 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM522 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM522 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM522 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7620] The complementary binding of GAM522 RNA, herein designated GAM RNA, to target binding sites on GAM522 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM522 target RNA, herein desig-

nated GAM TARGET RNA, into GAM522 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7621] It is appreciated that GAM522 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM522 target genes. The mRNA of each one of this plurality of GAM522 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM522 RNA, herein designated GAM RNA, and which when bound by GAM522 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM522 target proteins.

[7622] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM522 gene, herein designated GAM GENE, on one or more GAM522 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7623] It is yet further appreciated that specific functions, and accordingly utilities, of GAM522 correlate with, and may be deduced from, the identity of the target genes which GAM522 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7624] Nucleotide sequences of the GAM522 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM522 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM522 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM522 are further described hereinbelow with reference to Table Table1.

[7625] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM522 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[7626]

[7627] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 523 (GAM523) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7628] GAM523 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM523 was detected is described hereinabove with reference to Figs. 2-8.

[7629] GAM523 gene, herein designated GAM GENE, and GAM523 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7630] GAM523 gene, herein designated GAM GENE, encodes a GAM523 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM523 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM523 precursor RNA is designated SEQ ID:8335, and is provided hereinbelow with ref-

erence to the sequence listing part.

[7631] GAM523 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM523 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7632] An enzyme complex designated DICER COMPLEX, dices the GAM523 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM523 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 66%) nucleotide sequence of GAM523 RNA is designated SEQ ID:8336, and is provided hereinbelow with reference to the sequence listing part.

[7633] GAM523 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM523 target RNA, herein designated GAM TARGET RNA.

GAM523 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7634] GAM523 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM523 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM523 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM523 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM523 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7635] The complementary binding of GAM523 RNA, herein designated GAM RNA, to target binding sites on GAM523 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM523 target RNA, herein designated GAM TARGET RNA, into GAM523 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7636] It is appreciated that GAM523 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM523 target genes. The mRNA of each one of this plurality of GAM523 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM523 RNA, herein designated GAM RNA, and which when bound by GAM523 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM523 target proteins.

[7637] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM523 gene, herein designated GAM GENE, on one or more GAM523 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7638] It is yet further appreciated that specific functions, and accordingly utilities, of GAM523 correlate with, and may be deduced from, the identity of the target genes which GAM523 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7639] Nucleotide sequences of the GAM523 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM523 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM523 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM523 are further described hereinbelow with reference to Table Table1.

[7640] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM523 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7641]

[7642] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 524 (GAM524) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7643] GAM524 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM524 was detected is described hereinabove with reference to Figs. 2-8.

[7644] GAM524 gene, herein designated GAM GENE, and GAM524 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[7645] GAM524 gene, herein designated GAM GENE, encodes a GAM524 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM524 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM524 precursor RNA is designated SEQ ID:8344, and is provided hereinbelow with reference to the sequence listing part.

[7646] GAM524 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM524 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7647] An enzyme complex designated DICER COMPLEX, dices the GAM524 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM524 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 65%) nucleotide sequence of GAM524 RNA is designated SEQ ID:8345, and is provided hereinbelow with reference to the sequence listing part.

[7648] GAM524 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM524 target RNA, herein designated GAM TARGET RNA. GAM524 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7649] GAM524 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM524 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM524 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM524 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM524 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7650] The complementary binding of GAM524 RNA, herein designated GAM RNA, to target binding sites on GAM524 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM524 target RNA, herein designated GAM TARGET RNA, into GAM524 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7651] It is appreciated that GAM524 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM524 target genes. The mRNA of each one of this plurality of GAM524 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM524 RNA, herein designated GAM RNA, and which when bound by GAM524 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM524 target proteins.

[7652] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM524 gene, herein designated GAM GENE, on one or more GAM524 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[7653] It is yet further appreciated that specific functions, and accordingly utilities, of GAM524 correlate with, and may be deduced from, the identity of the target genes which GAM524 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7654] Nucleotide sequences of the GAM524 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM524 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM524 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM524 are further described hereinbelow with reference to Table Table1.

[7655] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM524 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7656]

[7657] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

525 (GAM525) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7658] GAM525 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM525 was detected is described hereinabove with reference to Figs. 2-8.

[7659] GAM525 gene, herein designated GAM GENE, and GAM525 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7660] GAM525 gene, herein designated GAM GENE, encodes a GAM525 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM525 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM525 precursor RNA is designated SEQ ID:8355, and is provided hereinbelow with reference to the sequence listing part.

[7661] GAM525 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM525 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7662] An enzyme complex designated DICER COMPLEX, dices the GAM525 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM525 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM525 RNA is designated SEQ ID:8356, and is provided hereinbelow with reference to the sequence listing part.

[7663] GAM525 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM525 target RNA, herein designated GAM TARGET RNA. GAM525 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7664] GAM525 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM525 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM525 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM525 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM525 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7665] The complementary binding of GAM525 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM525 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM525 target RNA, herein designated GAM TARGET RNA, into GAM525 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7666] It is appreciated that GAM525 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM525 target genes. The mRNA of each one of this plurality of GAM525 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM525 RNA, herein designated GAM RNA, and which when bound by GAM525 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM525 target proteins.

[7667] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM525 gene, herein designated GAM GENE, on one or more GAM525 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7668] It is yet further appreciated that specific functions, and accordingly utilities, of GAM525 correlate with, and may be deduced from, the identity of the target genes which GAM525 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7669] Nucleotide sequences of the GAM525 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM525 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM525 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM525 are further described hereinbelow with reference to Table Table1.

[7670] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM525 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7671]

[7672] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 526 (GAM526) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7673] GAM526 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM526 was detected is described hereinabove with reference to Figs. 2-8.

[7674] GAM526 gene, herein designated GAM GENE, and GAM526 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7675] GAM526 gene, herein designated GAM GENE, encodes a GAM526 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM526 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM526 precursor RNA is designated SEQ ID:8374, and is provided hereinbelow with reference to the sequence listing part.

[7676] GAM526 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM526 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7677] An enzyme complex designated DICER COMPLEX, dices the GAM526 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM526 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 61%) nucleotide sequence of GAM526 RNA is designated SEQ ID:6904, and is provided hereinbelow with reference to the sequence listing part.

[7678] GAM526 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM526 target RNA, herein designated GAM TARGET RNA. GAM526 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7679] GAM526 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM526 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM526 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM526 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM526 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7680] The complementary binding of GAM526 RNA, herein designated GAM RNA, to target binding sites on GAM526 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM526 target RNA, herein designated GAM TARGET RNA, into GAM526 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7681] It is appreciated that GAM526 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM526 target genes. The mRNA of each one of this plurality of GAM526 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM526 RNA, herein designated GAM RNA, and which when bound by

GAM526 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM526 target proteins.

[7682] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM526 gene, herein designated GAM GENE, on one or more GAM526 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7683] It is yet further appreciated that specific functions, and accordingly utilities, of GAM526 correlate with, and may be deduced from, the identity of the target genes which GAM526 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[7684] Nucleotide sequences of the GAM526 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM526 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM526 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM526 are further described hereinbelow with reference to Table Table1.

[7685] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM526 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7686]

[7687] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 527 (GAM527) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7688] GAM527 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM527 was detected is described hereinabove with reference to Figs. 2-8.

[7689] GAM527 gene, herein designated GAM GENE, and GAM527 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7690] GAM527 gene, herein designated GAM GENE, encodes a GAM527 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM527 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM527 precursor RNA is designated SEQ ID:8390, and is provided hereinbelow with reference to the sequence listing part.

[7691] GAM527 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM527 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[7692] An enzyme complex designated DICER COMPLEX, dices the GAM527 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM527 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 28%) nucleotide sequence of GAM527 RNA is designated SEQ ID:8391, and is provided hereinbelow with reference to the sequence listing part.

[7693] GAM527 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM527 target RNA, herein designated GAM TARGET RNA. GAM527 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7694] GAM527 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM527 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM527 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM527 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM527 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7695] The complementary binding of GAM527 RNA, herein designated GAM RNA, to target binding sites on GAM527 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM527 target RNA, herein designated GAM TARGET RNA, into GAM527 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7696] It is appreciated that GAM527 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM527 target genes. The mRNA of each one of this plurality of GAM527 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM527 RNA, herein designated GAM RNA, and which when bound by GAM527 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM527 target proteins.

[7697] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM527 gene, herein designated GAM GENE, on one or more GAM527 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7698] It is yet further appreciated that specific functions, and accordingly utilities, of GAM527 correlate with, and may be deduced from, the identity of the target genes which GAM527 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7699] Nucleotide sequences of the GAM527 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM527 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM527 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM527 are further described hereinbelow with reference to Table Table1.

[7700] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM527 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7701]

[7702] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 528 (GAM528) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7703] GAM528 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM528 was detected is described hereinabove with reference to Figs. 2-8.

[7704] GAM528 gene, herein designated GAM GENE, and GAM528 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7705] GAM528 gene, herein designated GAM GENE, encodes a GAM528 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM528 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM528 precursor RNA is designated SEQ ID:8394, and is provided hereinbelow with reference to the sequence listing part.

[7706] GAM528 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM528 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7707] An enzyme complex designated DICER COMPLEX, dices the GAM528 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM528 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 59%) nucleotide sequence of GAM528 RNA is designated SEQ ID:8395, and is provided hereinbelow with reference to the sequence listing part.

[7708] GAM528 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM528

target RNA, herein designated GAM TARGET RNA.

GAM528 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7709] GAM528 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM528 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM528 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM528 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM528 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding

sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7710] The complementary binding of GAM528 RNA, herein designated GAM RNA, to target binding sites on GAM528 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM528 target RNA, herein designated GAM TARGET RNA, into GAM528 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7711] It is appreciated that GAM528 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM528 target genes. The mRNA of each one of this plurality of GAM528 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM528 RNA, herein designated GAM RNA, and which when bound by GAM528 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM528 target proteins.

[7712] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with

specific reference to translational inhibition exerted by GAM528 gene, herein designated GAM GENE, on one or more GAM528 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7713] It is yet further appreciated that specific functions, and accordingly utilities, of GAM528 correlate with, and may be deduced from, the identity of the target genes which GAM528 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7714] Nucleotide sequences of the GAM528 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM528 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of

GAM528 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM528 are further described hereinbelow with reference to Table Table1.

[7715] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM528 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7716]

[7717] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 529 (GAM529) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7718] GAM529 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM529 was detected is described hereinabove with reference to Figs. 2-8.

[7719] GAM529 gene, herein designated GAM GENE, and GAM529 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7720] GAM529 gene, herein designated GAM GENE, encodes a GAM529 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM529 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM529 precursor RNA is designated SEQ ID:8403, and is provided hereinbelow with reference to the sequence listing part.

[7721] GAM529 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM529 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7722] An enzyme complex designated DICER COMPLEX, dices the GAM529 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM529 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA

segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 72%) nucleotide sequence of GAM529 RNA is designated SEQ ID:8404, and is provided hereinbelow with reference to the sequence listing part.

[7723] GAM529 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM529 target RNA, herein designated GAM TARGET RNA. GAM529 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7724] GAM529 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM529 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM529 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustra-

tion, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM529 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM529 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7725] The complementary binding of GAM529 RNA, herein designated GAM RNA, to target binding sites on GAM529 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM529 target RNA, herein designated GAM TARGET RNA, into GAM529 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7726] It is appreciated that GAM529 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM529 target genes. The mRNA of each one of this plu-

rality of GAM529 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM529 RNA, herein designated GAM RNA, and which when bound by GAM529 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM529 target proteins.

[7727] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM529 gene, herein designated GAM GENE, on one or more GAM529 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7728] It is yet further appreciated that specific functions, and accordingly utilities, of GAM529 correlate with, and may be deduced from, the identity of the target genes which GAM529 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7729] Nucleotide sequences of the GAM529 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM529 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM529 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM529 are further described hereinbelow with reference to Table Table1.

[7730] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM529 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7731]

[7732] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 530 (GAM530) gene, which modulates expression of re-

spective target genes thereof, the function and utility of which target genes is known in the art.

[7733] GAM530 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM530 was detected is described hereinabove with reference to Figs. 2-8.

[7734] GAM530 gene, herein designated GAM GENE, and GAM530 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7735] GAM530 gene, herein designated GAM GENE, encodes a GAM530 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM530 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM530 precursor RNA is designated SEQ ID:8432, and is provided hereinbelow with reference to the sequence listing part.

[7736] GAM530 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM530 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of

RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7737] An enzyme complex designated DICER COMPLEX, dices the GAM530 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM530 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 67%) nucleotide sequence of GAM530 RNA is designated SEQ ID:8433, and is provided hereinbelow with reference to the sequence listing part.

[7738] GAM530 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM530 target RNA, herein designated GAM TARGET RNA.
GAM530 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated

5UTR, PROTEIN CODING and 3UTR respectively.

[7739] GAM530 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM530 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM530 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM530 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM530 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7740] The complementary binding of GAM530 RNA, herein designated GAM RNA, to target binding sites on GAM530 tar-

get RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM530 target RNA, herein designated GAM TARGET RNA, into GAM530 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7741] It is appreciated that GAM530 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM530 target genes. The mRNA of each one of this plurality of GAM530 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM530 RNA, herein designated GAM RNA, and which when bound by GAM530 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM530 target proteins.

[7742] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM530 gene, herein designated GAM GENE, on one or more GAM530 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the

background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7743] It is yet further appreciated that specific functions, and accordingly utilities, of GAM530 correlate with, and may be deduced from, the identity of the target genes which GAM530 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7744] Nucleotide sequences of the GAM530 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM530 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM530 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM530 are further described hereinbelow with reference to Table Table1.

[7745] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of

Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM530 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7746]

[7747] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 531 (GAM531) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7748] GAM531 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM531 was detected is described hereinabove with reference to Figs. 2-8.

[7749] GAM531 gene, herein designated GAM GENE, and GAM531 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7750] GAM531 gene, herein designated GAM GENE, encodes a GAM531 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM531 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A

nucleotide sequence identical or highly similar to the nucleotide sequence of GAM531 precursor RNA is designated SEQ ID:8440, and is provided hereinbelow with reference to the sequence listing part.

[7751] GAM531 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM531 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7752] An enzyme complex designated DICER COMPLEX, dices the GAM531 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM531 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 41%) nucleotide sequence of

GAM531 RNA is designated SEQ ID:8441, and is provided hereinbelow with reference to the sequence listing part.

[7753] GAM531 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM531 target RNA, herein designated GAM TARGET RNA. GAM531 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7754] GAM531 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM531 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM531 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM531 RNA, herein

designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM531 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7755] The complementary binding of GAM531 RNA, herein designated GAM RNA, to target binding sites on GAM531 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM531 target RNA, herein designated GAM TARGET RNA, into GAM531 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7756] It is appreciated that GAM531 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM531 target genes. The mRNA of each one of this plurality of GAM531 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM531 RNA, herein designated GAM RNA, and which when bound by GAM531 RNA, herein designated GAM RNA, causes inhibi-

tion of translation of respective one or more GAM531 target proteins.

[7757] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM531 gene, herein designated GAM GENE, on one or more GAM531 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7758] It is yet further appreciated that specific functions, and accordingly utilities, of GAM531 correlate with, and may be deduced from, the identity of the target genes which GAM531 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7759] Nucleotide sequences of the GAM531 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM531 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM531 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM531 are further described hereinbelow with reference to Table Table1.

[7760] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM531 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7761]

[7762] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 532 (GAM532) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7763] GAM532 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM532 was detected is described hereinabove

with reference to Figs. 2-8.

[7764] GAM532 gene, herein designated GAM GENE, and GAM532 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7765] GAM532 gene, herein designated GAM GENE, encodes a GAM532 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM532 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM532 precursor RNA is designated SEQ ID:8449, and is provided hereinbelow with reference to the sequence listing part.

[7766] GAM532 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM532 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7767] An enzyme complex designated DICER COMPLEX, dices the GAM532 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM532 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 42%) nucleotide sequence of GAM532 RNA is designated SEQ ID:8441, and is provided hereinbelow with reference to the sequence listing part.

[7768] GAM532 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM532 target RNA, herein designated GAM TARGET RNA. GAM532 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7769] GAM532 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM532 target RNA, herein designated GAM TARGET RNA. This complementary binding is

due to the fact that the nucleotide sequence of GAM532 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM532 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM532 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7770] The complementary binding of GAM532 RNA, herein designated GAM RNA, to target binding sites on GAM532 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM532 target RNA, herein designated GAM TARGET RNA, into GAM532 target protein, herein designated GAM TARGET PROTEIN. GAM target

protein is therefore outlined by a broken line.

[7771] It is appreciated that GAM532 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM532 target genes. The mRNA of each one of this plurality of GAM532 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM532 RNA, herein designated GAM RNA, and which when bound by GAM532 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM532 target proteins.

[7772] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM532 gene, herein designated GAM GENE, on one or more GAM532 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other

genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7773] It is yet further appreciated that specific functions, and accordingly utilities, of GAM532 correlate with, and may be deduced from, the identity of the target genes which GAM532 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7774] Nucleotide sequences of the GAM532 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM532 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM532 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM532 are further described hereinbelow with reference to Table Table1.

[7775] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM532 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7776]

[7777] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 533 (GAM533) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7778] GAM533 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM533 was detected is described hereinabove with reference to Figs. 2-8.

[7779] GAM533 gene, herein designated GAM GENE, and GAM533 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7780] GAM533 gene, herein designated GAM GENE, encodes a GAM533 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM533 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM533 precursor RNA is designated SEQ ID:8469, and is provided hereinbelow with reference to the sequence listing part.

[7781] GAM533 precursor RNA, herein designated GAM PRECUR-

SOR RNA, folds onto itself, forming GAM533 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7782] An enzyme complex designated DICER COMPLEX, dices the GAM533 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM533 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 58%) nucleotide sequence of GAM533 RNA is designated SEQ ID:8470, and is provided hereinbelow with reference to the sequence listing part.

[7783] GAM533 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM533 target RNA, herein designated GAM TARGET RNA.

GAM533 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7784] GAM533 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM533 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM533 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM533 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM533 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only

these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7785] The complementary binding of GAM533 RNA, herein designated GAM RNA, to target binding sites on GAM533 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM533 target RNA, herein designated GAM TARGET RNA, into GAM533 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7786] It is appreciated that GAM533 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM533 target genes. The mRNA of each one of this plurality of GAM533 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM533 RNA, herein designated GAM RNA, and which when bound by GAM533 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM533 target proteins.

[7787] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by

GAM533 gene, herein designated GAM GENE, on one or more GAM533 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7788] It is yet further appreciated that specific functions, and accordingly utilities, of GAM533 correlate with, and may be deduced from, the identity of the target genes which GAM533 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7789] Nucleotide sequences of the GAM533 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM533 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM533 folded precursor RNA, herein designated GAM

FOLDED PRECURSOR RNA, of GAM533 are further described hereinbelow with reference to Table Table1.

[7790] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM533 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7791]

[7792] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 534 (GAM534) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7793] GAM534 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM534 was detected is described hereinabove with reference to Figs. 2-8.

[7794] GAM534 gene, herein designated GAM GENE, and GAM534 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7795] GAM534 gene, herein designated GAM GENE, encodes a

GAM534 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM534 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM534 precursor RNA is designated SEQ ID:8493, and is provided hereinbelow with reference to the sequence listing part.

[7796] GAM534 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM534 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7797] An enzyme complex designated DICER COMPLEX, dices the GAM534 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM534 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin struc-

tured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM534 RNA is designated SEQ ID:8494, and is provided hereinbelow with reference to the sequence listing part.

[7798] GAM534 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM534 target RNA, herein designated GAM TARGET RNA. GAM534 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7799] GAM534 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM534 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM534 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated

BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM534 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM534 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7800] The complementary binding of GAM534 RNA, herein designated GAM RNA, to target binding sites on GAM534 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM534 target RNA, herein designated GAM TARGET RNA, into GAM534 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7801] It is appreciated that GAM534 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM534 target genes. The mRNA of each one of this plurality of GAM534 target genes comprises one or more tar-

get binding sites, each having a nucleotide sequence which is at least partly complementary to GAM534 RNA, herein designated GAM RNA, and which when bound by GAM534 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM534 target proteins.

[7802] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM534 gene, herein designated GAM GENE, on one or more GAM534 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7803] It is yet further appreciated that specific functions, and

accordingly utilities, of GAM534 correlate with, and may be deduced from, the identity of the target genes which GAM534 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7804] Nucleotide sequences of the GAM534 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM534 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM534 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM534 are further described hereinbelow with reference to Table Table1.

[7805] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM534 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7806]

[7807] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 535 (GAM535) gene, which modulates expression of respective target genes thereof, the function and utility of

which target genes is known in the art.

[7808] GAM535 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM535 was detected is described hereinabove with reference to Figs. 2-8.

[7809] GAM535 gene, herein designated GAM GENE, and GAM535 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7810] GAM535 gene, herein designated GAM GENE, encodes a GAM535 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM535 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM535 precursor RNA is designated SEQ ID:8508, and is provided hereinbelow with reference to the sequence listing part.

[7811] GAM535 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM535 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that

the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7812] An enzyme complex designated DICER COMPLEX, dices the GAM535 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM535 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 64%) nucleotide sequence of GAM535 RNA is designated SEQ ID:8509, and is provided hereinbelow with reference to the sequence listing part.

[7813] GAM535 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM535 target RNA, herein designated GAM TARGET RNA. GAM535 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7814] GAM535 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM535 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM535 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM535 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM535 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7815] The complementary binding of GAM535 RNA, herein designated GAM RNA, to target binding sites on GAM535 target RNA, herein designated GAM TARGET RNA, such as

BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM535 target RNA, herein designated GAM TARGET RNA, into GAM535 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7816] It is appreciated that GAM535 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM535 target genes. The mRNA of each one of this plurality of GAM535 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM535 RNA, herein designated GAM RNA, and which when bound by GAM535 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM535 target proteins.

[7817] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM535 gene, herein designated GAM GENE, on one or more GAM535 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary

binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7818] It is yet further appreciated that specific functions, and accordingly utilities, of GAM535 correlate with, and may be deduced from, the identity of the target genes which GAM535 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7819] Nucleotide sequences of the GAM535 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM535 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM535 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM535 are further described hereinbelow with reference to Table Table1.

[7820] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the

complementarity of each of these target binding sites to GAM535 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7821]

[7822] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 536 (GAM536) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7823] GAM536 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM536 was detected is described hereinabove with reference to Figs. 2-8.

[7824] GAM536 gene, herein designated GAM GENE, and GAM536 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7825] GAM536 gene, herein designated GAM GENE, encodes a GAM536 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM536 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nu-

cleotide sequence of GAM536 precursor RNA is designated SEQ ID:8519, and is provided hereinbelow with reference to the sequence listing part.

[7826] GAM536 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM536 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7827] An enzyme complex designated DICER COMPLEX, dices the GAM536 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM536 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 52%) nucleotide sequence of GAM536 RNA is designated SEQ ID:8520, and is provided

hereinbelow with reference to the sequence listing part.

[7828] GAM536 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM536 target RNA, herein designated GAM TARGET RNA.

GAM536 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7829] GAM536 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM536 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM536 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM536 RNA, herein designated GAM RNA, may have a different number of tar-

get binding sites in untranslated regions of a GAM536 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7830] The complementary binding of GAM536 RNA, herein designated GAM RNA, to target binding sites on GAM536 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM536 target RNA, herein designated GAM TARGET RNA, into GAM536 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7831] It is appreciated that GAM536 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM536 target genes. The mRNA of each one of this plurality of GAM536 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM536 RNA, herein designated GAM RNA, and which when bound by GAM536 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM536 tar-

get proteins.

[7832] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM536 gene, herein designated GAM GENE, on one or more GAM536 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7833] It is yet further appreciated that specific functions, and accordingly utilities, of GAM536 correlate with, and may be deduced from, the identity of the target genes which GAM536 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7834] Nucleotide sequences of the GAM536 precursor RNA,

herein designated GAM PRECURSOR RNA, and of the diced GAM536 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM536 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM536 are further described hereinbelow with reference to Table Table1.

[7835] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM536 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7836]

[7837] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 537 (GAM537) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7838] GAM537 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM537 was detected is described hereinabove with reference to Figs. 2-8.

[7839] GAM537 gene, herein designated GAM GENE, and GAM537 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7840] GAM537 gene, herein designated GAM GENE, encodes a GAM537 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM537 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM537 precursor RNA is designated SEQ ID:8528, and is provided hereinbelow with reference to the sequence listing part.

[7841] GAM537 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM537 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7842] An enzyme complex designated DICER COMPLEX, dices

the GAM537 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM537 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 39%) nucleotide sequence of GAM537 RNA is designated SEQ ID:8529, and is provided hereinbelow with reference to the sequence listing part.

[7843] GAM537 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM537 target RNA, herein designated GAM TARGET RNA. GAM537 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7844] GAM537 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM537 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM537

RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM537 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM537 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7845] The complementary binding of GAM537 RNA, herein designated GAM RNA, to target binding sites on GAM537 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM537 target RNA, herein designated GAM TARGET RNA, into GAM537 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7846] It is appreciated that GAM537 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM537 target genes. The mRNA of each one of this plurality of GAM537 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM537 RNA, herein designated GAM RNA, and which when bound by GAM537 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM537 target proteins.

[7847] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM537 gene, herein designated GAM GENE, on one or more GAM537 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific com-

plementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7848] It is yet further appreciated that specific functions, and accordingly utilities, of GAM537 correlate with, and may be deduced from, the identity of the target genes which GAM537 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7849] Nucleotide sequences of the GAM537 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM537 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM537 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM537 are further described hereinbelow with reference to Table Table1.

[7850] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM537 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7851]

[7852] Fig. 8 further provides a conceptual description of another

novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 538 (GAM538) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7853] GAM538 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM538 was detected is described hereinabove with reference to Figs. 2-8.

[7854] GAM538 gene, herein designated GAM GENE, and GAM538 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7855] GAM538 gene, herein designated GAM GENE, encodes a GAM538 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM538 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM538 precursor RNA is designated SEQ ID:8549, and is provided hereinbelow with reference to the sequence listing part.

[7856] GAM538 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM538 folded pre-

cursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7857] An enzyme complex designated DICER COMPLEX, dices the GAM538 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM538 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 28%) nucleotide sequence of GAM538 RNA is designated SEQ ID:8550, and is provided hereinbelow with reference to the sequence listing part.

[7858] GAM538 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM538 target RNA, herein designated GAM TARGET RNA. GAM538 target RNA, herein designated GAM TARGET

RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7859] GAM538 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM538 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM538 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM538 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM538 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR re-

gion, the 5UTR region, or in both 3UTR and 5UTR regions.

[7860] The complementary binding of GAM538 RNA, herein designated GAM RNA, to target binding sites on GAM538 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM538 target RNA, herein designated GAM TARGET RNA, into GAM538 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7861] It is appreciated that GAM538 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM538 target genes. The mRNA of each one of this plurality of GAM538 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM538 RNA, herein designated GAM RNA, and which when bound by GAM538 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM538 target proteins.

[7862] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM538 gene, herein designated GAM GENE, on one or

more GAM538 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7863] It is yet further appreciated that specific functions, and accordingly utilities, of GAM538 correlate with, and may be deduced from, the identity of the target genes which GAM538 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7864] Nucleotide sequences of the GAM538 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM538 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM538 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM538 are further de-

scribed hereinbelow with reference to Table Table1.

[7865] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM538 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7866]

[7867] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 539 (GAM539) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7868] GAM539 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM539 was detected is described hereinabove with reference to Figs. 2-8.

[7869] GAM539 gene, herein designated GAM GENE, and GAM539 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7870] GAM539 gene, herein designated GAM GENE, encodes a GAM539 precursor RNA, herein designated GAM PRECUR-

SOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM539 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM539 precursor RNA is designated SEQ ID:8564, and is provided hereinbelow with reference to the sequence listing part.

[7871] GAM539 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM539 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7872] An enzyme complex designated DICER COMPLEX, dices the GAM539 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM539 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA seg-

ment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 60%) nucleotide sequence of GAM539 RNA is designated SEQ ID:8565, and is provided hereinbelow with reference to the sequence listing part.

[7873] GAM539 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM539 target RNA, herein designated GAM TARGET RNA. GAM539 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7874] GAM539 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM539 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM539 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III re-

spectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM539 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM539 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7875] The complementary binding of GAM539 RNA, herein designated GAM RNA, to target binding sites on GAM539 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM539 target RNA, herein designated GAM TARGET RNA, into GAM539 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7876] It is appreciated that GAM539 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM539 target genes. The mRNA of each one of this plurality of GAM539 target genes comprises one or more target binding sites, each having a nucleotide sequence

which is at least partly complementary to GAM539 RNA, herein designated GAM RNA, and which when bound by GAM539 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM539 target proteins.

[7877] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM539 gene, herein designated GAM GENE, on one or more GAM539 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7878] It is yet further appreciated that specific functions, and accordingly utilities, of GAM539 correlate with, and may

be deduced from, the identity of the target genes which GAM539 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7879] Nucleotide sequences of the GAM539 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM539 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM539 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM539 are further described hereinbelow with reference to Table Table1.

[7880] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM539 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7881]

[7882] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 540 (GAM540) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7883] GAM540 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM540 was detected is described hereinabove with reference to Figs. 2-8.

[7884] GAM540 gene, herein designated GAM GENE, and GAM540 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7885] GAM540 gene, herein designated GAM GENE, encodes a GAM540 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM540 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM540 precursor RNA is designated SEQ ID:8567, and is provided hereinbelow with reference to the sequence listing part.

[7886] GAM540 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM540 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA en-

coded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7887] An enzyme complex designated DICER COMPLEX, dices the GAM540 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM540 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 54%) nucleotide sequence of GAM540 RNA is designated SEQ ID:8568, and is provided hereinbelow with reference to the sequence listing part.

[7888] GAM540 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM540 target RNA, herein designated GAM TARGET RNA. GAM540 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7889] GAM540 RNA, herein designated GAM RNA, binds com-

plementarily to one or more target binding sites located in untranslated regions of GAM540 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM540 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM540 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM540 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7890] The complementary binding of GAM540 RNA, herein designated GAM RNA, to target binding sites on GAM540 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, in-

hibits translation of GAM540 target RNA, herein designated GAM TARGET RNA, into GAM540 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7891] It is appreciated that GAM540 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM540 target genes. The mRNA of each one of this plurality of GAM540 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM540 RNA, herein designated GAM RNA, and which when bound by GAM540 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM540 target proteins.

[7892] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM540 gene, herein designated GAM GENE, on one or more GAM540 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the

known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7893] It is yet further appreciated that specific functions, and accordingly utilities, of GAM540 correlate with, and may be deduced from, the identity of the target genes which GAM540 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7894] Nucleotide sequences of the GAM540 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM540 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM540 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM540 are further described hereinbelow with reference to Table Table1.

[7895] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to

GAM540 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7896]

[7897] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 541 (GAM541) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7898] GAM541 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM541 was detected is described hereinabove with reference to Figs. 2-8.

[7899] GAM541 gene, herein designated GAM GENE, and GAM541 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7900] GAM541 gene, herein designated GAM GENE, encodes a GAM541 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM541 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM541 precursor RNA is desig-

nated SEQ ID:8573, and is provided hereinbelow with reference to the sequence listing part.

[7901] GAM541 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM541 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7902] An enzyme complex designated DICER COMPLEX, dices the GAM541 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM541 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 30%) nucleotide sequence of GAM541 RNA is designated SEQ ID:8574, and is provided hereinbelow with reference to the sequence listing part.

[7903] GAM541 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM541 target RNA, herein designated GAM TARGET RNA. GAM541 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7904] GAM541 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM541 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM541 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM541 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM541

target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7905] The complementary binding of GAM541 RNA, herein designated GAM RNA, to target binding sites on GAM541 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM541 target RNA, herein designated GAM TARGET RNA, into GAM541 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7906] It is appreciated that GAM541 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM541 target genes. The mRNA of each one of this plurality of GAM541 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM541 RNA, herein designated GAM RNA, and which when bound by GAM541 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM541 target proteins.

[7907] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM541 gene, herein designated GAM GENE, on one or more GAM541 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7908] It is yet further appreciated that specific functions, and accordingly utilities, of GAM541 correlate with, and may be deduced from, the identity of the target genes which GAM541 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7909] Nucleotide sequences of the GAM541 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced

GAM541 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM541 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM541 are further described hereinbelow with reference to Table Table1.

[7910] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM541 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7911]

[7912] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 542 (GAM542) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7913] GAM542 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM542 was detected is described hereinabove with reference to Figs. 2-8.

[7914] GAM542 gene, herein designated GAM GENE, and

GAM542 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7915] GAM542 gene, herein designated GAM GENE, encodes a GAM542 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM542 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM542 precursor RNA is designated SEQ ID:8579, and is provided hereinbelow with reference to the sequence listing part.

[7916] GAM542 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM542 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7917] An enzyme complex designated DICER COMPLEX, dices the GAM542 folded precursor RNA, herein designated

GAM FOLDED PRECURSOR RNA, into GAM542 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 62%) nucleotide sequence of GAM542 RNA is designated SEQ ID:8580, and is provided hereinbelow with reference to the sequence listing part.

[7918] GAM542 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM542 target RNA, herein designated GAM TARGET RNA. GAM542 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7919] GAM542 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM542 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM542 RNA, herein designated GAM RNA, is an accurate or a par-

tial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM542 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM542 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7920] The complementary binding of GAM542 RNA, herein designated GAM RNA, to target binding sites on GAM542 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM542 target RNA, herein designated GAM TARGET RNA, into GAM542 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7921] It is appreciated that GAM542 target gene, herein desig-

nated GAM TARGET GENE, in fact represents a plurality of GAM542 target genes. The mRNA of each one of this plurality of GAM542 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM542 RNA, herein designated GAM RNA, and which when bound by GAM542 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM542 target proteins.

[7922] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM542 gene, herein designated GAM GENE, on one or more GAM542 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have

not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7923] It is yet further appreciated that specific functions, and accordingly utilities, of GAM542 correlate with, and may be deduced from, the identity of the target genes which GAM542 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7924] Nucleotide sequences of the GAM542 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM542 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM542 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM542 are further described hereinbelow with reference to Table Table1.

[7925] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM542 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7926]

[7927] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present in-

vention, referred to here as Genomic Address Messenger 543 (GAM543) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7928] GAM543 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM543 was detected is described hereinabove with reference to Figs. 2-8.

[7929] GAM543 gene, herein designated GAM GENE, and GAM543 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7930] GAM543 gene, herein designated GAM GENE, encodes a GAM543 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM543 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM543 precursor RNA is designated SEQ ID:8608, and is provided hereinbelow with reference to the sequence listing part.

[7931] GAM543 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM543 folded precursor RNA, herein designated GAM FOLDED PRECURSOR

RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7932] An enzyme complex designated DICER COMPLEX, dices the GAM543 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM543 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM543 RNA is designated SEQ ID:8609, and is provided hereinbelow with reference to the sequence listing part.

[7933] GAM543 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM543 target RNA, herein designated GAM TARGET RNA. GAM543 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a

protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7934] GAM543 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM543 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM543 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM543 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM543 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7935] The complementary binding of GAM543 RNA, herein designated GAM RNA, to target binding sites on GAM543 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM543 target RNA, herein designated GAM TARGET RNA, into GAM543 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7936] It is appreciated that GAM543 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM543 target genes. The mRNA of each one of this plurality of GAM543 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM543 RNA, herein designated GAM RNA, and which when bound by GAM543 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM543 target proteins.

[7937] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM543 gene, herein designated GAM GENE, on one or more GAM543 target gene, herein designated GAM TAR-

GET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7938] It is yet further appreciated that specific functions, and accordingly utilities, of GAM543 correlate with, and may be deduced from, the identity of the target genes which GAM543 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7939] Nucleotide sequences of the GAM543 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM543 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM543 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM543 are further described hereinbelow with reference to Table Table1.

[7940] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM543 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7941]

[7942] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 544 (GAM544) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7943] GAM544 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM544 was detected is described hereinabove with reference to Figs. 2-8.

[7944] GAM544 gene, herein designated GAM GENE, and GAM544 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7945] GAM544 gene, herein designated GAM GENE, encodes a GAM544 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most

ordinary genes, GAM544 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM544 precursor RNA is designated SEQ ID:8621, and is provided hereinbelow with reference to the sequence listing part.

[7946] GAM544 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM544 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7947] An enzyme complex designated DICER COMPLEX, dices the GAM544 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM544 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an

enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM544 RNA is designated SEQ ID:8622, and is provided hereinbelow with reference to the sequence listing part.

[7948] GAM544 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM544 target RNA, herein designated GAM TARGET RNA. GAM544 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7949] GAM544 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM544 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM544 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target

binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM544 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM544 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7950] The complementary binding of GAM544 RNA, herein designated GAM RNA, to target binding sites on GAM544 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM544 target RNA, herein designated GAM TARGET RNA, into GAM544 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7951] It is appreciated that GAM544 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM544 target genes. The mRNA of each one of this plurality of GAM544 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM544 RNA,

herein designated GAM RNA, and which when bound by GAM544 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM544 target proteins.

[7952] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM544 gene, herein designated GAM GENE, on one or more GAM544 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7953] It is yet further appreciated that specific functions, and accordingly utilities, of GAM544 correlate with, and may be deduced from, the identity of the target genes which

GAM544 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7954] Nucleotide sequences of the GAM544 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM544 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM544 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM544 are further described hereinbelow with reference to Table Table1.

[7955] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM544 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7956]

[7957] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 545 (GAM545) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7958] GAM545 is a novel bioinformatically detected regulatory,

non protein coding, micro RNA (miRNA) gene. The method by which GAM545 was detected is described hereinabove with reference to Figs. 2-8.

[7959] GAM545 gene, herein designated GAM GENE, and GAM545 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7960] GAM545 gene, herein designated GAM GENE, encodes a GAM545 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM545 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM545 precursor RNA is designated SEQ ID:8652, and is provided hereinbelow with reference to the sequence listing part.

[7961] GAM545 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM545 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-

reversed sequence of the nucleotide sequence of the second half thereof.

[7962] An enzyme complex designated DICER COMPLEX, dices the GAM545 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM545 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM545 RNA is designated SEQ ID:8622, and is provided hereinbelow with reference to the sequence listing part.

[7963] GAM545 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM545 target RNA, herein designated GAM TARGET RNA.
GAM545 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7964] GAM545 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in

untranslated regions of GAM545 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM545 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM545 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM545 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7965] The complementary binding of GAM545 RNA, herein designated GAM RNA, to target binding sites on GAM545 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM545 target RNA, herein desig-

nated GAM TARGET RNA, into GAM545 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7966] It is appreciated that GAM545 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM545 target genes. The mRNA of each one of this plurality of GAM545 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM545 RNA, herein designated GAM RNA, and which when bound by GAM545 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM545 target proteins.

[7967] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM545 gene, herein designated GAM GENE, on one or more GAM545 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other

recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7968] It is yet further appreciated that specific functions, and accordingly utilities, of GAM545 correlate with, and may be deduced from, the identity of the target genes which GAM545 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7969] Nucleotide sequences of the GAM545 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM545 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM545 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM545 are further described hereinbelow with reference to Table Table1.

[7970] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM545 RNA, herein designated GAM RNA, are described

hereinbelow with reference to Table Table2.

[7971]

[7972] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 546 (GAM546) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7973] GAM546 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM546 was detected is described hereinabove with reference to Figs. 2-8.

[7974] GAM546 gene, herein designated GAM GENE, and GAM546 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[7975] GAM546 gene, herein designated GAM GENE, encodes a GAM546 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM546 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM546 precursor RNA is designated SEQ ID:8687, and is provided hereinbelow with ref-

erence to the sequence listing part.

[7976] GAM546 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM546 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7977] An enzyme complex designated DICER COMPLEX, dices the GAM546 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM546 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 71%) nucleotide sequence of GAM546 RNA is designated SEQ ID:8609, and is provided hereinbelow with reference to the sequence listing part.

[7978] GAM546 target gene, herein designated GAM TARGET

GENE, encodes a corresponding messenger RNA, GAM546 target RNA, herein designated GAM TARGET RNA.

GAM546 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7979] GAM546 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM546 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM546 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM546 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM546 target RNA, herein designated GAM TARGET RNA. It is fur-

ther appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7980] The complementary binding of GAM546 RNA, herein designated GAM RNA, to target binding sites on GAM546 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM546 target RNA, herein designated GAM TARGET RNA, into GAM546 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7981] It is appreciated that GAM546 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM546 target genes. The mRNA of each one of this plurality of GAM546 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM546 RNA, herein designated GAM RNA, and which when bound by GAM546 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM546 target proteins.

[7982] It is further appreciated by one skilled in the art that the

mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM546 gene, herein designated GAM GENE, on one or more GAM546 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[7983] It is yet further appreciated that specific functions, and accordingly utilities, of GAM546 correlate with, and may be deduced from, the identity of the target genes which GAM546 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7984] Nucleotide sequences of the GAM546 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM546 RNA, herein designated GAM RNA, and a

schematic representation of the secondary folding of GAM546 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM546 are further described hereinbelow with reference to Table Table1.

[7985] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM546 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[7986]

[7987] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 547 (GAM547) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[7988] GAM547 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM547 was detected is described hereinabove with reference to Figs. 2-8.

[7989] GAM547 gene, herein designated GAM GENE, and GAM547 target gene, herein designated GAM TARGET

GENE, are human genes contained in the human genome.

[7990] GAM547 gene, herein designated GAM GENE, encodes a GAM547 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM547 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM547 precursor RNA is designated SEQ ID:8715, and is provided hereinbelow with reference to the sequence listing part.

[7991] GAM547 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM547 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[7992] An enzyme complex designated DICER COMPLEX, dices the GAM547 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM547 RNA, herein

designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 56%) nucleotide sequence of GAM547 RNA is designated SEQ ID:8716, and is provided hereinbelow with reference to the sequence listing part.

[7993] GAM547 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM547 target RNA, herein designated GAM TARGET RNA. GAM547 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[7994] GAM547 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM547 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM547 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide se-

quence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM547 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM547 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[7995] The complementary binding of GAM547 RNA, herein designated GAM RNA, to target binding sites on GAM547 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM547 target RNA, herein designated GAM TARGET RNA, into GAM547 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[7996] It is appreciated that GAM547 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of

GAM547 target genes. The mRNA of each one of this plurality of GAM547 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM547 RNA, herein designated GAM RNA, and which when bound by GAM547 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM547 target proteins.

[7997] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM547 gene, herein designated GAM GENE, on one or more GAM547 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a

tiny RNA world, Science 294,779 (2001)).

[7998] It is yet further appreciated that specific functions, and accordingly utilities, of GAM547 correlate with, and may be deduced from, the identity of the target genes which GAM547 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[7999] Nucleotide sequences of the GAM547 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM547 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM547 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM547 are further described hereinbelow with reference to Table Table1.

[8000] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM547 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[8001]

[8002] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger

548 (GAM548) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[8003] GAM548 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM548 was detected is described hereinabove with reference to Figs. 2-8.

[8004] GAM548 gene, herein designated GAM GENE, and GAM548 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[8005] GAM548 gene, herein designated GAM GENE, encodes a GAM548 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM548 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM548 precursor RNA is designated SEQ ID:8724, and is provided hereinbelow with reference to the sequence listing part.

[8006] GAM548 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM548 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is

well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[8007] An enzyme complex designated DICER COMPLEX, dices the GAM548 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM548 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 53%) nucleotide sequence of GAM548 RNA is designated SEQ ID:8725, and is provided hereinbelow with reference to the sequence listing part.

[8008] GAM548 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM548 target RNA, herein designated GAM TARGET RNA. GAM548 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein

coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[8009] GAM548 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM548 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM548 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM548 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM548 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[8010] The complementary binding of GAM548 RNA, herein des-

ignated GAM RNA, to target binding sites on GAM548 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM548 target RNA, herein designated GAM TARGET RNA, into GAM548 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[8011] It is appreciated that GAM548 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM548 target genes. The mRNA of each one of this plurality of GAM548 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM548 RNA, herein designated GAM RNA, and which when bound by GAM548 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM548 target proteins.

[8012] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM548 gene, herein designated GAM GENE, on one or more GAM548 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA

genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[8013] It is yet further appreciated that specific functions, and accordingly utilities, of GAM548 correlate with, and may be deduced from, the identity of the target genes which GAM548 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[8014] Nucleotide sequences of the GAM548 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM548 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM548 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM548 are further described hereinbelow with reference to Table Table1.

[8015] Nucleotide sequences of target binding sites, such as

BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM548 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[8016]

[8017] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 549 (GAM549) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[8018] GAM549 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method by which GAM549 was detected is described hereinabove with reference to Figs. 2-8.

[8019] GAM549 gene, herein designated GAM GENE, and GAM549 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[8020] GAM549 gene, herein designated GAM GENE, encodes a GAM549 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM549 precursor RNA, herein desig-

nated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM549 precursor RNA is designated SEQ ID:8728, and is provided hereinbelow with reference to the sequence listing part.

[8021] GAM549 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM549 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the second half thereof.

[8022] An enzyme complex designated DICER COMPLEX, dices the GAM549 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM549 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary pro-

teins. A probable (over 49%) nucleotide sequence of GAM549 RNA is designated SEQ ID:8729, and is provided hereinbelow with reference to the sequence listing part.

[8023] GAM549 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM549 target RNA, herein designated GAM TARGET RNA. GAM549 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[8024] GAM549 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM549 target RNA, herein designated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM549 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration

only, and is not meant to be limiting GAM549 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM549 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[8025] The complementary binding of GAM549 RNA, herein designated GAM RNA, to target binding sites on GAM549 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM549 target RNA, herein designated GAM TARGET RNA, into GAM549 target protein, herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[8026] It is appreciated that GAM549 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM549 target genes. The mRNA of each one of this plurality of GAM549 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM549 RNA, herein designated GAM RNA, and which when bound by

GAM549 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM549 target proteins.

[8027] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM549 gene, herein designated GAM GENE, on one or more GAM549 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[8028] It is yet further appreciated that specific functions, and accordingly utilities, of GAM549 correlate with, and may be deduced from, the identity of the target genes which GAM549 binds and inhibits, and the function of these tar-

get genes, as elaborated hereinbelow.

[8029] Nucleotide sequences of the GAM549 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM549 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM549 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM549 are further described hereinbelow with reference to Table Table1.

[8030] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM549 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[8031]

[8032] Fig. 8 further provides a conceptual description of another novel bioinformatically detected gene of the present invention, referred to here as Genomic Address Messenger 550 (GAM550) gene, which modulates expression of respective target genes thereof, the function and utility of which target genes is known in the art.

[8033] GAM550 is a novel bioinformatically detected regulatory, non protein coding, micro RNA (miRNA) gene. The method

by which GAM550 was detected is described hereinabove with reference to Figs. 2-8.

[8034] GAM550 gene, herein designated GAM GENE, and GAM550 target gene, herein designated GAM TARGET GENE, are human genes contained in the human genome.

[8035] GAM550 gene, herein designated GAM GENE, encodes a GAM550 precursor RNA, herein designated GAM PRECURSOR RNA. Similar to other miRNA genes, and unlike most ordinary genes, GAM550 precursor RNA, herein designated GAM PRECURSOR RNA, does not encode a protein. A nucleotide sequence identical or highly similar to the nucleotide sequence of GAM550 precursor RNA is designated SEQ ID:8744, and is provided hereinbelow with reference to the sequence listing part.

[8036] GAM550 precursor RNA, herein designated GAM PRECURSOR RNA, folds onto itself, forming GAM550 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, which has a two-dimensional hairpin structure. As is well known in the art, this hairpin structure, is typical of RNA encoded by miRNA genes, and is due to the fact that the nucleotide sequence of the first half of the RNA encoded by a miRNA gene is an accurate or partial inversed-reversed sequence of the nucleotide sequence of the sec-

ond half thereof.

[8037] An enzyme complex designated DICER COMPLEX, dices the GAM550 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, into GAM550 RNA, herein designated GAM RNA, a single stranded ~22 nt long RNA segment. As is known in the art, dicing of a hairpin structured RNA precursor product into a short ~22nt RNA segment is catalyzed by an enzyme complex comprising an enzyme called Dicer together with other necessary proteins. A probable (over 74%) nucleotide sequence of GAM550 RNA is designated SEQ ID:8745, and is provided hereinbelow with reference to the sequence listing part.

[8038] GAM550 target gene, herein designated GAM TARGET GENE, encodes a corresponding messenger RNA, GAM550 target RNA, herein designated GAM TARGET RNA. GAM550 target RNA, herein designated GAM TARGET RNA, comprises three regions, as is typical of mRNA of a protein coding gene: a 5 untranslated region, a protein coding region and a 3 untranslated region, designated 5UTR, PROTEIN CODING and 3UTR respectively.

[8039] GAM550 RNA, herein designated GAM RNA, binds complementarily to one or more target binding sites located in untranslated regions of GAM550 target RNA, herein des-

ignated GAM TARGET RNA. This complementary binding is due to the fact that the nucleotide sequence of GAM550 RNA, herein designated GAM RNA, is an accurate or a partial inversed-reversed sequence of the nucleotide sequence of each of the target binding sites. As an illustration, Fig. 8 shows 3 such target binding sites, designated BINDING SITE I, BINDING SITE II and BINDING SITE III respectively. It is appreciated that the number of target binding sites shown in Fig. 8 is meant as an illustration only, and is not meant to be limiting GAM550 RNA, herein designated GAM RNA, may have a different number of target binding sites in untranslated regions of a GAM550 target RNA, herein designated GAM TARGET RNA. It is further appreciated that while Fig. 8 depicts target binding sites in the 3UTR region, this is meant as an example only these target binding sites may be located in the 3UTR region, the 5UTR region, or in both 3UTR and 5UTR regions.

[8040] The complementary binding of GAM550 RNA, herein designated GAM RNA, to target binding sites on GAM550 target RNA, herein designated GAM TARGET RNA, such as BINDING SITE I, BINDING SITE II and BINDING SITE III, inhibits translation of GAM550 target RNA, herein designated GAM TARGET RNA, into GAM550 target protein,

herein designated GAM TARGET PROTEIN. GAM target protein is therefore outlined by a broken line.

[8041] It is appreciated that GAM550 target gene, herein designated GAM TARGET GENE, in fact represents a plurality of GAM550 target genes. The mRNA of each one of this plurality of GAM550 target genes comprises one or more target binding sites, each having a nucleotide sequence which is at least partly complementary to GAM550 RNA, herein designated GAM RNA, and which when bound by GAM550 RNA, herein designated GAM RNA, causes inhibition of translation of respective one or more GAM550 target proteins.

[8042] It is further appreciated by one skilled in the art that the mode of translational inhibition illustrated by Fig. 8 with specific reference to translational inhibition exerted by GAM550 gene, herein designated GAM GENE, on one or more GAM550 target gene, herein designated GAM TARGET GENE, is in fact common to other known miRNA genes. As mentioned hereinabove with reference to the background section, although a specific complementary binding site has been demonstrated only for some of the known miRNA genes (primarily Lin-4 and Let-7), all other recently discovered miRNA genes are also believed by

those skilled in the art to modulate expression of other genes by complementary binding, although specific complementary binding sites of these other miRNA genes have not yet been found (Ruvkun G., Perspective: Glimpses of a tiny RNA world, Science 294,779 (2001)).

[8043] It is yet further appreciated that specific functions, and accordingly utilities, of GAM550 correlate with, and may be deduced from, the identity of the target genes which GAM550 binds and inhibits, and the function of these target genes, as elaborated hereinbelow.

[8044] Nucleotide sequences of the GAM550 precursor RNA, herein designated GAM PRECURSOR RNA, and of the diced GAM550 RNA, herein designated GAM RNA, and a schematic representation of the secondary folding of GAM550 folded precursor RNA, herein designated GAM FOLDED PRECURSOR RNA, of GAM550 are further described hereinbelow with reference to Table Table1.

[8045] Nucleotide sequences of target binding sites, such as BINDING SITE-I, BINDING SITE-II and BINDING SITE-III of Fig. 8, found on, and schematic representation of the complementarity of each of these target binding sites to GAM550 RNA, herein designated GAM RNA, are described hereinbelow with reference to Table Table2.

[8046]

[8047] Reference is now made to Fig. 11, which is a simplified diagram describing each of a plurality of novel bioinformatically detected regulatory genes, referred to here as Genomic Record(GR) genes, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8048] GR GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR GENE was detected is described hereinabove with reference to Figs. 6-15.

[8049] GR GENE encodes GR PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8050] GR PRECURSOR RNA folds spatially, forming GR FOLDED PRECURSOR RNA. It is appreciated that GR FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the sec-

ond half thereof, as is well known in the art.

[8051] GR FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into a plurality of separate GAM precursor RNAs, schematically represented by GAM1 PRECURSOR, GAM2 PRECURSOR, GAM3 PRECURSOR, GAM4 PRECURSOR, GAM5 PRECURSOR, GAM6 PRECURSOR, GAM7 PRECURSOR and GAM8 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8052] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, schematically represented as GAM1 RNA, GAM2 RNA, GAM3 RNA, GAM4 RNA, GAM5 RNA, GAM6 RNA, GAM7 RNA and GAM8 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8053] GAM1 RNA binds complimentarily to a target binding site located in an untranslated region of GAM1 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM1 TARGET RNA into GAM1 TARGET PROTEIN, both of Fig. 10.

[8054] GAM2 RNA binds complimentarily to a target binding site

located in an untranslated region of GAM2 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM2 TARGET RNA into GAM2 TARGET PROTEIN, both of Fig. 10.

[8055] GAM3 RNA binds complimentarily to a target binding site located in an untranslated region of GAM3 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM3 TARGET RNA into GAM3 TARGET PROTEIN, both of Fig. 10.

[8056] GAM4 RNA binds complimentarily to a target binding site located in an untranslated region of GAM4 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM4 TARGET RNA into GAM4 TARGET PROTEIN, both of Fig. 10.

[8057] GAM5 RNA binds complimentarily to a target binding site located in an untranslated region of GAM5 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM5 TARGET RNA into GAM5 TARGET PROTEIN, both of Fig. 10.

[8058] GAM6 RNA binds complimentarily to a target binding site located in an untranslated region of GAM6 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM6 TARGET RNA into GAM6 TARGET PROTEIN, both of Fig. 10.

[8059] GAM7 RNA binds complimentarily to a target binding site located in an untranslated region of GAM7 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM7 TARGET RNA into GAM7 TARGET PROTEIN, both of Fig. 10.

[8060] GAM8 RNA binds complimentarily to a target binding site located in an untranslated region of GAM8 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 10, thereby inhibiting translation of GAM8 TARGET RNA into GAM8 TARGET PROTEIN, both of Fig. 10.

[8061] It is appreciated that specific functions, and accordingly utilities, of GR GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR GENE, schematically represented by GAM1 TARGET PRO-

TEIN through GAM8 TARGET PROTEIN.

[8062] Reference is now made to Fig. 551, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 551(GR551) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8063] GR551 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR551 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8064] GR551 GENE encodes GR551 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8065] GR551 PRECURSOR RNA folds spatially, forming GR551 FOLDED PRECURSOR RNA . It is appreciated that GR551 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR551 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[8066] GR551 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8067] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8068] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8069] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8070] It is appreciated that specific functions, and accordingly utilities, of GR551 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR551 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8071] Reference is now made to Fig. 552, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 552(GR552) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8072] GR552 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR552 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[8073] GR552 GENE encodes GR552 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8074] GR552 PRECURSOR RNA folds spatially, forming GR552 FOLDED PRECURSOR RNA . It is appreciated that GR552 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR552 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8075] GR552 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM103 PRECURSOR and GAM312 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8076] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM103 RNA and GAM312 RNA respectively, each of which GAM RNAs

corresponding to GAM RNA of Fig. 8.

[8077] GAM103 RNA binds complimentarily to a target binding site located in an untranslated region of GAM103 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 103A, thereby inhibiting translation of GAM103 TARGET RNA into GAM103 TARGET PROTEIN, both of Fig. 103A.

[8078] GAM312 RNA binds complimentarily to a target binding site located in an untranslated region of GAM312 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 312A, thereby inhibiting translation of GAM312 TARGET RNA into GAM312 TARGET PROTEIN, both of Fig. 312A.

[8079] It is appreciated that specific functions, and accordingly utilities, of GR552 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR552 GENE: GAM103 TARGET PROTEIN and GAM312 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 103D and 312D

[8080] Reference is now made to Fig. 553, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 553(GR553) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8081] GR553 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR553 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8082] GR553 GENE encodes GR553 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8083] GR553 PRECURSOR RNA folds spatially, forming GR553 FOLDED PRECURSOR RNA . It is appreciated that GR553 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR553 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[8084] GR553 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM269 PRECURSOR and GAM372 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8085] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM269 RNA and GAM372 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8086] GAM269 RNA binds complimentarily to a target binding site located in an untranslated region of GAM269 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 269A, thereby inhibiting translation of GAM269 TARGET RNA into GAM269 TARGET PROTEIN, both of Fig. 269A.

[8087] GAM372 RNA binds complimentarily to a target binding site located in an untranslated region of GAM372 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 372A, thereby inhibiting translation of GAM372 TARGET RNA into GAM372 TARGET PROTEIN, both of Fig. 372A.

[8088] It is appreciated that specific functions, and accordingly utilities, of GR553 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR553 GENE: GAM269 TARGET PROTEIN and GAM372 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 269D and 372D

[8089] Reference is now made to Fig. 554, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 554(GR554) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8090] GR554 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR554 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8091] GR554 GENE encodes GR554 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8092] GR554 PRECURSOR RNA folds spatially, forming GR554 FOLDED PRECURSOR RNA . It is appreciated that GR554 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR554 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8093] GR554 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM150 PRECURSOR and GAM542 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8094] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM150 RNA and GAM542 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8095] GAM150 RNA binds complimentarily to a target binding site located in an untranslated region of GAM150 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 150A, thereby inhibiting translation of GAM150 TARGET RNA into GAM150 TARGET PROTEIN, both of Fig. 150A.

[8096] GAM542 RNA binds complimentarily to a target binding site located in an untranslated region of GAM542 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 542A, thereby inhibiting translation of GAM542 TARGET RNA into GAM542 TARGET PROTEIN, both of Fig. 542A.

[8097] It is appreciated that specific functions, and accordingly utilities, of GR554 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR554 GENE: GAM150 TARGET PROTEIN and GAM542 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 150D and 542D

[8098] Reference is now made to Fig. 555, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 555(GR555) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8099] GR555 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR555 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8100] GR555 GENE encodes GR555 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8101] GR555 PRECURSOR RNA folds spatially, forming GR555 FOLDED PRECURSOR RNA . It is appreciated that GR555 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR555 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8102] GR555 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM293 PRECURSOR and GAM368 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8103] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM293 RNA and GAM368 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8104] GAM293 RNA binds complimentarily to a target binding site located in an untranslated region of GAM293 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 293A, thereby inhibiting translation of GAM293 TARGET RNA into GAM293 TARGET PROTEIN, both of Fig. 293A.

[8105] GAM368 RNA binds complimentarily to a target binding site located in an untranslated region of GAM368 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 368A, thereby inhibiting transla-

tion of GAM368 TARGET RNA into GAM368 TARGET PROTEIN, both of Fig. 368A.

[8106] It is appreciated that specific functions, and accordingly utilities, of GR555 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR555 GENE: GAM293 TARGET PROTEIN and GAM368 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 293D and 368D

[8107] Reference is now made to Fig. 556, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 556(GR556) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8108] GR556 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR556 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8109] GR556 GENE encodes GR556 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[8110] GR556 PRECURSOR RNA folds spatially, forming GR556 FOLDED PRECURSOR RNA . It is appreciated that GR556 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR556 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8111] GR556 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM515 PRECURSOR and GAM516 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8112] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM515 RNA and GAM516 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8113] GAM515 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM515 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 515A, thereby inhibiting translation of GAM515 TARGET RNA into GAM515 TARGET PROTEIN, both of Fig. 515A.

[8114] GAM516 RNA binds complimentarily to a target binding site located in an untranslated region of GAM516 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 516A, thereby inhibiting translation of GAM516 TARGET RNA into GAM516 TARGET PROTEIN, both of Fig. 516A.

[8115] It is appreciated that specific functions, and accordingly utilities, of GR556 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR556 GENE: GAM515 TARGET PROTEIN and GAM516 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 515D and 516D

[8116] Reference is now made to Fig. 557, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 557(GR557) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8117] GR557 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR557 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8118] GR557 GENE encodes GR557 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8119] GR557 PRECURSOR RNA folds spatially, forming GR557 FOLDED PRECURSOR RNA . It is appreciated that GR557 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR557 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8120] GR557 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM85 PRECURSOR and GAM121 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8121] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM85 RNA and GAM121 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8122] GAM85 RNA binds complimentarily to a target binding site located in an untranslated region of GAM85 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 85A, thereby inhibiting translation of GAM85 TARGET RNA into GAM85 TARGET PROTEIN, both of Fig. 85A.

[8123] GAM121 RNA binds complimentarily to a target binding site located in an untranslated region of GAM121 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 121A, thereby inhibiting translation of GAM121 TARGET RNA into GAM121 TARGET PRO-

TEIN, both of Fig. 121A.

[8124] It is appreciated that specific functions, and accordingly utilities, of GR557 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR557 GENE: GAM85 TARGET PROTEIN and GAM121 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 85D and 121D

[8125] Reference is now made to Fig. 558, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 558(GR558) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8126] GR558 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR558 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8127] GR558 GENE encodes GR558 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8128] GR558 PRECURSOR RNA folds spatially, forming GR558 FOLDED PRECURSOR RNA . It is appreciated that GR558 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR558 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8129] GR558 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM266 PRECURSOR, GAM357 PRECURSOR and GAM504 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8130] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM266 RNA, GAM357 RNA and GAM504 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8131] GAM266 RNA binds complimentarily to a target binding site located in an untranslated region of GAM266 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 266A, thereby inhibiting translation of GAM266 TARGET RNA into GAM266 TARGET PROTEIN, both of Fig. 266A.

[8132] GAM357 RNA binds complimentarily to a target binding site located in an untranslated region of GAM357 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 357A, thereby inhibiting translation of GAM357 TARGET RNA into GAM357 TARGET PROTEIN, both of Fig. 357A.

[8133] GAM504 RNA binds complimentarily to a target binding site located in an untranslated region of GAM504 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 504A, thereby inhibiting translation of GAM504 TARGET RNA into GAM504 TARGET PROTEIN, both of Fig. 504A.

[8134] It is appreciated that specific functions, and accordingly utilities, of GR558 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR558 GENE: GAM266 TARGET PROTEIN, GAM357 TARGET PROTEIN and GAM504 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 266D, 357D and 504D

[8135] Reference is now made to Fig. 559, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 559(GR559) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8136] GR559 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR559 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8137] GR559 GENE encodes GR559 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8138] GR559 PRECURSOR RNA folds spatially, forming GR559 FOLDED PRECURSOR RNA . It is appreciated that GR559 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR559 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8139] GR559 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM41 PRECURSOR, GAM45 PRECURSOR and GAM179 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8140] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM41 RNA, GAM45 RNA and GAM179 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8141] GAM41 RNA binds complimentarily to a target binding site located in an untranslated region of GAM41 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 41A, thereby inhibiting translation of GAM41 TARGET RNA into GAM41 TARGET PROTEIN, both of Fig. 41A.

[8142] GAM45 RNA binds complimentarily to a target binding site located in an untranslated region of GAM45 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 45A, thereby inhibiting translation of GAM45 TARGET RNA into GAM45 TARGET PROTEIN, both of Fig. 45A.

[8143] GAM179 RNA binds complimentarily to a target binding site located in an untranslated region of GAM179 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 179A, thereby inhibiting translation of GAM179 TARGET RNA into GAM179 TARGET PROTEIN, both of Fig. 179A.

[8144] It is appreciated that specific functions, and accordingly utilities, of GR559 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR559 GENE: GAM41 TARGET PROTEIN, GAM45 TARGET PROTEIN and GAM179 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 41D, 45D and 179D

[8145] Reference is now made to Fig. 560, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 560(GR560) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8146] GR560 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR560 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8147] GR560 GENE encodes GR560 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8148] GR560 PRECURSOR RNA folds spatially, forming GR560 FOLDED PRECURSOR RNA . It is appreciated that GR560 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR560 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8149] GR560 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8150] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8151] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8152] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting transla-

tion of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8153] It is appreciated that specific functions, and accordingly utilities, of GR560 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR560 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8154] Reference is now made to Fig. 561, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 561(GR561) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8155] GR561 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR561 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8156] GR561 GENE encodes GR561 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[8157] GR561 PRECURSOR RNA folds spatially, forming GR561 FOLDED PRECURSOR RNA . It is appreciated that GR561 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR561 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8158] GR561 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM212 PRECURSOR and GAM493 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8159] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM212 RNA and GAM493 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8160] GAM212 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM212 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 212A, thereby inhibiting translation of GAM212 TARGET RNA into GAM212 TARGET PROTEIN, both of Fig. 212A.

[8161] GAM493 RNA binds complimentarily to a target binding site located in an untranslated region of GAM493 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 493A, thereby inhibiting translation of GAM493 TARGET RNA into GAM493 TARGET PROTEIN, both of Fig. 493A.

[8162] It is appreciated that specific functions, and accordingly utilities, of GR561 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR561 GENE: GAM212 TARGET PROTEIN and GAM493 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 212D and 493D

[8163] Reference is now made to Fig. 562, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 562(GR562) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8164] GR562 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR562 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8165] GR562 GENE encodes GR562 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8166] GR562 PRECURSOR RNA folds spatially, forming GR562 FOLDED PRECURSOR RNA . It is appreciated that GR562 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR562 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8167] GR562 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM35 PRECURSOR and GAM513 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8168] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM35 RNA and GAM513 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8169] GAM35 RNA binds complimentarily to a target binding site located in an untranslated region of GAM35 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 35A, thereby inhibiting translation of GAM35 TARGET RNA into GAM35 TARGET PROTEIN, both of Fig. 35A.

[8170] GAM513 RNA binds complimentarily to a target binding site located in an untranslated region of GAM513 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 513A, thereby inhibiting translation of GAM513 TARGET RNA into GAM513 TARGET PROTEIN, both of Fig. 513A.

[8171] It is appreciated that specific functions, and accordingly utilities, of GR562 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR562 GENE: GAM35 TARGET PROTEIN and GAM513 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 35D and 513D

[8172] Reference is now made to Fig. 563, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 563(GR563) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8173] GR563 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR563 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8174] GR563 GENE encodes GR563 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8175] GR563 PRECURSOR RNA folds spatially, forming GR563 FOLDED PRECURSOR RNA . It is appreciated that GR563 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR563 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8176] GR563 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM224 PRECURSOR and GAM262 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8177] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM224 RNA and GAM262 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8178] GAM224 RNA binds complimentarily to a target binding site located in an untranslated region of GAM224 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 224A, thereby inhibiting translation of GAM224 TARGET RNA into GAM224 TARGET PROTEIN, both of Fig. 224A.

[8179] GAM262 RNA binds complimentarily to a target binding site located in an untranslated region of GAM262 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 262A, thereby inhibiting translation of GAM262 TARGET RNA into GAM262 TARGET PROTEIN, both of Fig. 262A.

[8180] It is appreciated that specific functions, and accordingly utilities, of GR563 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR563 GENE: GAM224 TARGET PROTEIN and GAM262 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 224D and 262D

[8181] Reference is now made to Fig. 564, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 564(GR564) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8182] GR564 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR564 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8183] GR564 GENE encodes GR564 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8184] GR564 PRECURSOR RNA folds spatially, forming GR564 FOLDED PRECURSOR RNA . It is appreciated that GR564 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR564 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8185] GR564 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM263 PRECURSOR and GAM385 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8186] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM263 RNA and GAM385 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8187] GAM263 RNA binds complimentarily to a target binding site located in an untranslated region of GAM263 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 263A, thereby inhibiting translation of GAM263 TARGET RNA into GAM263 TARGET PROTEIN, both of Fig. 263A.

[8188] GAM385 RNA binds complimentarily to a target binding site located in an untranslated region of GAM385 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 385A, thereby inhibiting translation of GAM385 TARGET RNA into GAM385 TARGET PROTEIN, both of Fig. 385A.

[8189] It is appreciated that specific functions, and accordingly utilities, of GR564 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR564 GENE: GAM263 TARGET PROTEIN and GAM385

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 263D and 385D

[8190] Reference is now made to Fig. 565, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 565(GR565) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8191] GR565 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR565 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8192] GR565 GENE encodes GR565 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8193] GR565 PRECURSOR RNA folds spatially, forming GR565 FOLDED PRECURSOR RNA . It is appreciated that GR565 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR565 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8194] GR565 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM431 PRECURSOR and GAM531 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8195] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM431 RNA and GAM531 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8196] GAM431 RNA binds complimentarily to a target binding site located in an untranslated region of GAM431 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 431A, thereby inhibiting translation of GAM431 TARGET RNA into GAM431 TARGET PROTEIN, both of Fig. 431A.

[8197] GAM531 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM531 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 531A, thereby inhibiting translation of GAM531 TARGET RNA into GAM531 TARGET PROTEIN, both of Fig. 531A.

[8198] It is appreciated that specific functions, and accordingly utilities, of GR565 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR565 GENE: GAM431 TARGET PROTEIN and GAM531 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 431D and 531D

[8199] Reference is now made to Fig. 566, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 566(GR566) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8200] GR566 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR566 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8201] GR566 GENE encodes GR566 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8202] GR566 PRECURSOR RNA folds spatially, forming GR566 FOLDED PRECURSOR RNA . It is appreciated that GR566 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR566 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8203] GR566 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM473 PRECURSOR and GAM477 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8204] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM473 RNA and GAM477 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8205] GAM473 RNA binds complimentarily to a target binding site located in an untranslated region of GAM473 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 473A, thereby inhibiting translation of GAM473 TARGET RNA into GAM473 TARGET PROTEIN, both of Fig. 473A.

[8206] GAM477 RNA binds complimentarily to a target binding site located in an untranslated region of GAM477 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 477A, thereby inhibiting translation of GAM477 TARGET RNA into GAM477 TARGET PROTEIN, both of Fig. 477A.

[8207] It is appreciated that specific functions, and accordingly utilities, of GR566 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR566 GENE: GAM473 TARGET PROTEIN and GAM477 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 473D and 477D

[8208] Reference is now made to Fig. 567, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 567(GR567) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8209] GR567 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR567 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8210] GR567 GENE encodes GR567 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8211] GR567 PRECURSOR RNA folds spatially, forming GR567 FOLDED PRECURSOR RNA . It is appreciated that GR567 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR567 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8212] GR567 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM480 PRECURSOR and GAM508 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8213] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM480 RNA and GAM508 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8214] GAM480 RNA binds complimentarily to a target binding site located in an untranslated region of GAM480 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 480A, thereby inhibiting translation of GAM480 TARGET RNA into GAM480 TARGET PROTEIN, both of Fig. 480A.

[8215] GAM508 RNA binds complimentarily to a target binding site located in an untranslated region of GAM508 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 508A, thereby inhibiting translation of GAM508 TARGET RNA into GAM508 TARGET PROTEIN, both of Fig. 508A.

[8216] It is appreciated that specific functions, and accordingly utilities, of GR567 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR567 GENE: GAM480 TARGET PROTEIN and GAM508 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 480D and 508D

[8217] Reference is now made to Fig. 568, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 568(GR568) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8218] GR568 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR568 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8219] GR568 GENE encodes GR568 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8220] GR568 PRECURSOR RNA folds spatially, forming GR568 FOLDED PRECURSOR RNA . It is appreciated that GR568 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR568 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8221] GR568 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM163 PRECURSOR and GAM369 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8222] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM163 RNA

and GAM369 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8223] GAM163 RNA binds complimentarily to a target binding site located in an untranslated region of GAM163 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 163A, thereby inhibiting translation of GAM163 TARGET RNA into GAM163 TARGET PROTEIN, both of Fig. 163A.

[8224] GAM369 RNA binds complimentarily to a target binding site located in an untranslated region of GAM369 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 369A, thereby inhibiting translation of GAM369 TARGET RNA into GAM369 TARGET PROTEIN, both of Fig. 369A.

[8225] It is appreciated that specific functions, and accordingly utilities, of GR568 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR568 GENE: GAM163 TARGET PROTEIN and GAM369 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 163D and

[8226] Reference is now made to Fig. 569, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 569(GR569) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8227] GR569 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR569 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8228] GR569 GENE encodes GR569 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8229] GR569 PRECURSOR RNA folds spatially, forming GR569 FOLDED PRECURSOR RNA . It is appreciated that GR569 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR569 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[8230] GR569 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM167 PRECURSOR and GAM499 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8231] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM167 RNA and GAM499 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8232] GAM167 RNA binds complimentarily to a target binding site located in an untranslated region of GAM167 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 167A, thereby inhibiting translation of GAM167 TARGET RNA into GAM167 TARGET PROTEIN, both of Fig. 167A.

[8233] GAM499 RNA binds complimentarily to a target binding site located in an untranslated region of GAM499 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 499A, thereby inhibiting translation of GAM499 TARGET RNA into GAM499 TARGET PROTEIN, both of Fig. 499A.

[8234] It is appreciated that specific functions, and accordingly utilities, of GR569 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR569 GENE: GAM167 TARGET PROTEIN and GAM499 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 167D and 499D

[8235] Reference is now made to Fig. 570, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 570(GR570) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8236] GR570 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR570 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[8237] GR570 GENE encodes GR570 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8238] GR570 PRECURSOR RNA folds spatially, forming GR570 FOLDED PRECURSOR RNA . It is appreciated that GR570 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR570 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8239] GR570 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8240] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs

corresponding to GAM RNA of Fig. 8.

[8241] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8242] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8243] It is appreciated that specific functions, and accordingly utilities, of GR570 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR570 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8244] Reference is now made to Fig. 571, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 571(GR571) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8245] GR571 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR571 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8246] GR571 GENE encodes GR571 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8247] GR571 PRECURSOR RNA folds spatially, forming GR571 FOLDED PRECURSOR RNA . It is appreciated that GR571 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR571 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[8248] GR571 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM318 PRECURSOR and GAM320 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8249] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM318 RNA and GAM320 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8250] GAM318 RNA binds complimentarily to a target binding site located in an untranslated region of GAM318 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 318A, thereby inhibiting translation of GAM318 TARGET RNA into GAM318 TARGET PROTEIN, both of Fig. 318A.

[8251] GAM320 RNA binds complimentarily to a target binding site located in an untranslated region of GAM320 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 320A, thereby inhibiting translation of GAM320 TARGET RNA into GAM320 TARGET PROTEIN, both of Fig. 320A.

[8252] It is appreciated that specific functions, and accordingly utilities, of GR571 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR571 GENE: GAM318 TARGET PROTEIN and GAM320 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 318D and 320D

[8253] Reference is now made to Fig. 572, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 572(GR572) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8254] GR572 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR572 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8255] GR572 GENE encodes GR572 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8256] GR572 PRECURSOR RNA folds spatially, forming GR572 FOLDED PRECURSOR RNA . It is appreciated that GR572 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR572 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8257] GR572 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM166 PRECURSOR, GAM255 PRECURSOR and GAM438 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8258] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM166 RNA, GAM255 RNA and GAM438 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8259] GAM166 RNA binds complimentarily to a target binding site located in an untranslated region of GAM166 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 166A, thereby inhibiting translation of GAM166 TARGET RNA into GAM166 TARGET PROTEIN, both of Fig. 166A.

[8260] GAM255 RNA binds complimentarily to a target binding site located in an untranslated region of GAM255 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 255A, thereby inhibiting translation of GAM255 TARGET RNA into GAM255 TARGET PROTEIN, both of Fig. 255A.

[8261] GAM438 RNA binds complimentarily to a target binding site located in an untranslated region of GAM438 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 438A, thereby inhibiting translation of GAM438 TARGET RNA into GAM438 TARGET PROTEIN, both of Fig. 438A.

[8262] It is appreciated that specific functions, and accordingly utilities, of GR572 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR572 GENE: GAM166 TARGET PROTEIN, GAM255 TARGET PROTEIN and GAM438 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 166D, 255D and 438D

[8263] Reference is now made to Fig. 573, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 573(GR573) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8264] GR573 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR573 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8265] GR573 GENE encodes GR573 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8266] GR573 PRECURSOR RNA folds spatially, forming GR573 FOLDED PRECURSOR RNA . It is appreciated that GR573 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR573 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8267] GR573 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM182 PRECURSOR and GAM287 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8268] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM182 RNA and GAM287 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8269] GAM182 RNA binds complimentarily to a target binding site located in an untranslated region of GAM182 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 182A, thereby inhibiting transla-

tion of GAM182 TARGET RNA into GAM182 TARGET PROTEIN, both of Fig. 182A.

[8270] GAM287 RNA binds complimentarily to a target binding site located in an untranslated region of GAM287 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 287A, thereby inhibiting translation of GAM287 TARGET RNA into GAM287 TARGET PROTEIN, both of Fig. 287A.

[8271] It is appreciated that specific functions, and accordingly utilities, of GR573 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR573 GENE: GAM182 TARGET PROTEIN and GAM287 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 182D and 287D

[8272] Reference is now made to Fig. 574, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 574(GR574) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the

function and utility of which at least one target gene is known in the art.

[8273] GR574 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR574 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8274] GR574 GENE encodes GR574 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8275] GR574 PRECURSOR RNA folds spatially, forming GR574 FOLDED PRECURSOR RNA . It is appreciated that GR574 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR574 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8276] GR574 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM305 PRECURSOR and GAM329 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECUR-

SOR RNA of Fig. 8.

[8277] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM305 RNA and GAM329 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8278] GAM305 RNA binds complimentarily to a target binding site located in an untranslated region of GAM305 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 305A, thereby inhibiting translation of GAM305 TARGET RNA into GAM305 TARGET PROTEIN, both of Fig. 305A.

[8279] GAM329 RNA binds complimentarily to a target binding site located in an untranslated region of GAM329 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 329A, thereby inhibiting translation of GAM329 TARGET RNA into GAM329 TARGET PROTEIN, both of Fig. 329A.

[8280] It is appreciated that specific functions, and accordingly utilities, of GR574 GENE correlate with, and may be deduced from, the identity of the target genes, which are in-

hibited by GAM RNAs comprised in the operon-like cluster of GR574 GENE: GAM305 TARGET PROTEIN and GAM329 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 305D and 329D

[8281] Reference is now made to Fig. 575, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 575(GR575) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8282] GR575 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR575 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8283] GR575 GENE encodes GR575 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8284] GR575 PRECURSOR RNA folds spatially, forming GR575 FOLDED PRECURSOR RNA . It is appreciated that GR575 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin

structures are due to the fact that the nucleotide sequence of GR575 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8285] GR575 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM165 PRECURSOR and GAM506 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8286] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM165 RNA and GAM506 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8287] GAM165 RNA binds complimentarily to a target binding site located in an untranslated region of GAM165 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 165A, thereby inhibiting translation of GAM165 TARGET RNA into GAM165 TARGET PRO-

TEIN, both of Fig. 165A.

[8288] GAM506 RNA binds complimentarily to a target binding site located in an untranslated region of GAM506 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 506A, thereby inhibiting translation of GAM506 TARGET RNA into GAM506 TARGET PROTEIN, both of Fig. 506A.

[8289] It is appreciated that specific functions, and accordingly utilities, of GR575 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR575 GENE: GAM165 TARGET PROTEIN and GAM506 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 165D and 506D

[8290] Reference is now made to Fig. 576, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 576(GR576) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is

known in the art.

[8291] GR576 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR576 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8292] GR576 GENE encodes GR576 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8293] GR576 PRECURSOR RNA folds spatially, forming GR576 FOLDED PRECURSOR RNA . It is appreciated that GR576 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR576 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8294] GR576 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM181 PRECURSOR and GAM392 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8295] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM181 RNA and GAM392 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8296] GAM181 RNA binds complimentarily to a target binding site located in an untranslated region of GAM181 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 181A, thereby inhibiting translation of GAM181 TARGET RNA into GAM181 TARGET PROTEIN, both of Fig. 181A.

[8297] GAM392 RNA binds complimentarily to a target binding site located in an untranslated region of GAM392 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 392A, thereby inhibiting translation of GAM392 TARGET RNA into GAM392 TARGET PROTEIN, both of Fig. 392A.

[8298] It is appreciated that specific functions, and accordingly utilities, of GR576 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR576 GENE: GAM181 TARGET PROTEIN and GAM392 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 181D and 392D

[8299] Reference is now made to Fig. 577, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 577(GR577) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8300] GR577 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR577 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8301] GR577 GENE encodes GR577 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8302] GR577 PRECURSOR RNA folds spatially, forming GR577 FOLDED PRECURSOR RNA . It is appreciated that GR577 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR577 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8303] GR577 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM249 PRECURSOR and GAM384 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8304] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM249 RNA and GAM384 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8305] GAM249 RNA binds complimentarily to a target binding site located in an untranslated region of GAM249 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 249A, thereby inhibiting translation of GAM249 TARGET RNA into GAM249 TARGET PROTEIN, both of Fig. 249A.

[8306] GAM384 RNA binds complimentarily to a target binding site located in an untranslated region of GAM384 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 384A, thereby inhibiting translation of GAM384 TARGET RNA into GAM384 TARGET PROTEIN, both of Fig. 384A.

[8307] It is appreciated that specific functions, and accordingly utilities, of GR577 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR577 GENE: GAM249 TARGET PROTEIN and GAM384 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 249D and 384D

[8308] Reference is now made to Fig. 578, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 578(GR578) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8309] GR578 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR578 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8310] GR578 GENE encodes GR578 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8311] GR578 PRECURSOR RNA folds spatially, forming GR578 FOLDED PRECURSOR RNA . It is appreciated that GR578 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR578 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8312] GR578 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM526 PRECURSOR and GAM541 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8313] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM526 RNA and GAM541 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8314] GAM526 RNA binds complimentarily to a target binding site located in an untranslated region of GAM526 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 526A, thereby inhibiting translation of GAM526 TARGET RNA into GAM526 TARGET PROTEIN, both of Fig. 526A.

[8315] GAM541 RNA binds complimentarily to a target binding site located in an untranslated region of GAM541 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 541A, thereby inhibiting translation of GAM541 TARGET RNA into GAM541 TARGET PROTEIN, both of Fig. 541A.

[8316] It is appreciated that specific functions, and accordingly utilities, of GR578 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR578 GENE: GAM526 TARGET PROTEIN and GAM541

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 526D and 541D

[8317] Reference is now made to Fig. 579, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 579(GR579) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8318] GR579 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR579 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8319] GR579 GENE encodes GR579 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8320] GR579 PRECURSOR RNA folds spatially, forming GR579 FOLDED PRECURSOR RNA . It is appreciated that GR579 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR579 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8321] GR579 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM323 PRECURSOR and GAM529 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8322] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM323 RNA and GAM529 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8323] GAM323 RNA binds complimentarily to a target binding site located in an untranslated region of GAM323 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 323A, thereby inhibiting translation of GAM323 TARGET RNA into GAM323 TARGET PROTEIN, both of Fig. 323A.

[8324] GAM529 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM529 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 529A, thereby inhibiting translation of GAM529 TARGET RNA into GAM529 TARGET PROTEIN, both of Fig. 529A.

[8325] It is appreciated that specific functions, and accordingly utilities, of GR579 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR579 GENE: GAM323 TARGET PROTEIN and GAM529 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 323D and 529D

[8326] Reference is now made to Fig. 580, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 580(GR580) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8327] GR580 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR580 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8328] GR580 GENE encodes GR580 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8329] GR580 PRECURSOR RNA folds spatially, forming GR580 FOLDED PRECURSOR RNA . It is appreciated that GR580 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR580 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8330] GR580 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM91 PRECURSOR and GAM465 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8331] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM91 RNA and GAM465 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8332] GAM91 RNA binds complimentarily to a target binding site located in an untranslated region of GAM91 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 91A, thereby inhibiting translation of GAM91 TARGET RNA into GAM91 TARGET PROTEIN, both of Fig. 91A.

[8333] GAM465 RNA binds complimentarily to a target binding site located in an untranslated region of GAM465 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 465A, thereby inhibiting translation of GAM465 TARGET RNA into GAM465 TARGET PROTEIN, both of Fig. 465A.

[8334] It is appreciated that specific functions, and accordingly utilities, of GR580 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR580 GENE: GAM91 TARGET PROTEIN and GAM465 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 91D and 465D

[8335] Reference is now made to Fig. 581, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 581(GR581) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8336] GR581 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR581 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8337] GR581 GENE encodes GR581 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8338] GR581 PRECURSOR RNA folds spatially, forming GR581 FOLDED PRECURSOR RNA . It is appreciated that GR581 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR581 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8339] GR581 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM44 PRECURSOR and GAM403 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8340] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM44 RNA and GAM403 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8341] GAM44 RNA binds complimentarily to a target binding site located in an untranslated region of GAM44 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 44A, thereby inhibiting translation of GAM44 TARGET RNA into GAM44 TARGET PROTEIN, both of Fig. 44A.

[8342] GAM403 RNA binds complimentarily to a target binding site located in an untranslated region of GAM403 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 403A, thereby inhibiting translation of GAM403 TARGET RNA into GAM403 TARGET PROTEIN, both of Fig. 403A.

[8343] It is appreciated that specific functions, and accordingly utilities, of GR581 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR581 GENE: GAM44 TARGET PROTEIN and GAM403 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 44D and 403D

[8344] Reference is now made to Fig. 582, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 582(GR582) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8345] GR582 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR582 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8346] GR582 GENE encodes GR582 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8347] GR582 PRECURSOR RNA folds spatially, forming GR582 FOLDED PRECURSOR RNA . It is appreciated that GR582 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR582 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8348] GR582 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM67 PRECURSOR, GAM295 PRECURSOR and GAM331 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8349] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM67 RNA,

GAM295 RNA and GAM331 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8350] GAM67 RNA binds complimentarily to a target binding site located in an untranslated region of GAM67 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 67A, thereby inhibiting translation of GAM67 TARGET RNA into GAM67 TARGET PROTEIN, both of Fig. 67A.

[8351] GAM295 RNA binds complimentarily to a target binding site located in an untranslated region of GAM295 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 295A, thereby inhibiting translation of GAM295 TARGET RNA into GAM295 TARGET PROTEIN, both of Fig. 295A.

[8352] GAM331 RNA binds complimentarily to a target binding site located in an untranslated region of GAM331 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 331A, thereby inhibiting translation of GAM331 TARGET RNA into GAM331 TARGET PROTEIN, both of Fig. 331A.

[8353] It is appreciated that specific functions, and accordingly utilities, of GR582 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR582 GENE: GAM67 TARGET PROTEIN, GAM295 TARGET PROTEIN and GAM331 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 67D, 295D and 331D

[8354] Reference is now made to Fig. 583, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 583(GR583) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8355] GR583 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR583 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8356] GR583 GENE encodes GR583 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8357] GR583 PRECURSOR RNA folds spatially, forming GR583

FOLDED PRECURSOR RNA . It is appreciated that GR583 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR583 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8358] GR583 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM189 PRECURSOR and GAM251 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8359] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM189 RNA and GAM251 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8360] GAM189 RNA binds complimentarily to a target binding site located in an untranslated region of GAM189 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 189A, thereby inhibiting translation of GAM189 TARGET RNA into GAM189 TARGET PROTEIN, both of Fig. 189A.

[8361] GAM251 RNA binds complimentarily to a target binding site located in an untranslated region of GAM251 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 251A, thereby inhibiting translation of GAM251 TARGET RNA into GAM251 TARGET PROTEIN, both of Fig. 251A.

[8362] It is appreciated that specific functions, and accordingly utilities, of GR583 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR583 GENE: GAM189 TARGET PROTEIN and GAM251 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 189D and 251D

[8363] Reference is now made to Fig. 584, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 584(GR584) gene, which encodes an operon-like cluster

of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8364] GR584 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR584 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8365] GR584 GENE encodes GR584 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8366] GR584 PRECURSOR RNA folds spatially, forming GR584 FOLDED PRECURSOR RNA . It is appreciated that GR584 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR584 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8367] GR584 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM53 PRECURSOR and GAM442 PRE-

CURSOR, each of which GAM precursor RNAs being a hair-pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8368] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM53 RNA and GAM442 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8369] GAM53 RNA binds complimentarily to a target binding site located in an untranslated region of GAM53 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 53A, thereby inhibiting translation of GAM53 TARGET RNA into GAM53 TARGET PROTEIN, both of Fig. 53A.

[8370] GAM442 RNA binds complimentarily to a target binding site located in an untranslated region of GAM442 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 442A, thereby inhibiting translation of GAM442 TARGET RNA into GAM442 TARGET PROTEIN, both of Fig. 442A.

[8371] It is appreciated that specific functions, and accordingly

utilities, of GR584 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR584 GENE: GAM53 TARGET PROTEIN and GAM442 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 53D and 442D

[8372] Reference is now made to Fig. 585, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 585(GR585) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8373] GR585 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR585 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8374] GR585 GENE encodes GR585 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8375] GR585 PRECURSOR RNA folds spatially, forming GR585 FOLDED PRECURSOR RNA . It is appreciated that GR585

FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR585 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8376] GR585 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM44 PRECURSOR and GAM403 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8377] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM44 RNA and GAM403 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8378] GAM44 RNA binds complimentarily to a target binding site located in an untranslated region of GAM44 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING

SITE III of Fig. 44A, thereby inhibiting translation of GAM44 TARGET RNA into GAM44 TARGET PROTEIN, both of Fig. 44A.

[8379] GAM403 RNA binds complimentarily to a target binding site located in an untranslated region of GAM403 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 403A, thereby inhibiting translation of GAM403 TARGET RNA into GAM403 TARGET PROTEIN, both of Fig. 403A.

[8380] It is appreciated that specific functions, and accordingly utilities, of GR585 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR585 GENE: GAM44 TARGET PROTEIN and GAM403 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 44D and 403D

[8381] Reference is now made to Fig. 586, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 586(GR586) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn

modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8382] GR586 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR586 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8383] GR586 GENE encodes GR586 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8384] GR586 PRECURSOR RNA folds spatially, forming GR586 FOLDED PRECURSOR RNA . It is appreciated that GR586 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR586 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8385] GR586 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM110 PRECURSOR and GAM207 PRECURSOR, each of which GAM precursor RNAs being a hair-

pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8386] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM110 RNA and GAM207 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8387] GAM110 RNA binds complimentarily to a target binding site located in an untranslated region of GAM110 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 110A, thereby inhibiting translation of GAM110 TARGET RNA into GAM110 TARGET PROTEIN, both of Fig. 110A.

[8388] GAM207 RNA binds complimentarily to a target binding site located in an untranslated region of GAM207 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 207A, thereby inhibiting translation of GAM207 TARGET RNA into GAM207 TARGET PROTEIN, both of Fig. 207A.

[8389] It is appreciated that specific functions, and accordingly utilities, of GR586 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR586 GENE: GAM110 TARGET PROTEIN and GAM207 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 110D and 207D

[8390] Reference is now made to Fig. 587, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 587(GR587) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8391] GR587 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR587 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8392] GR587 GENE encodes GR587 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8393] GR587 PRECURSOR RNA folds spatially, forming GR587 FOLDED PRECURSOR RNA . It is appreciated that GR587 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR587 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8394] GR587 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM109 PRECURSOR, GAM215 PRECURSOR and GAM418 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8395] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM109 RNA, GAM215 RNA and GAM418 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8396] GAM109 RNA binds complimentarily to a target binding site located in an untranslated region of GAM109 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 109A, thereby inhibiting transla-

tion of GAM109 TARGET RNA into GAM109 TARGET PROTEIN, both of Fig. 109A.

[8397] GAM215 RNA binds complimentarily to a target binding site located in an untranslated region of GAM215 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 215A, thereby inhibiting translation of GAM215 TARGET RNA into GAM215 TARGET PROTEIN, both of Fig. 215A.

[8398] GAM418 RNA binds complimentarily to a target binding site located in an untranslated region of GAM418 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 418A, thereby inhibiting translation of GAM418 TARGET RNA into GAM418 TARGET PROTEIN, both of Fig. 418A.

[8399] It is appreciated that specific functions, and accordingly utilities, of GR587 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR587 GENE: GAM109 TARGET PROTEIN, GAM215 TARGET PROTEIN and GAM418 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with ref-

erence to Figs. 109D, 215D and 418D

[8400] Reference is now made to Fig. 588, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 588(GR588) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8401] GR588 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR588 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8402] GR588 GENE encodes GR588 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8403] GR588 PRECURSOR RNA folds spatially, forming GR588 FOLDED PRECURSOR RNA . It is appreciated that GR588 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR588 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[8404] GR588 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM120 PRECURSOR and GAM205 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8405] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM120 RNA and GAM205 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8406] GAM120 RNA binds complimentarily to a target binding site located in an untranslated region of GAM120 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 120A, thereby inhibiting translation of GAM120 TARGET RNA into GAM120 TARGET PROTEIN, both of Fig. 120A.

[8407] GAM205 RNA binds complimentarily to a target binding site located in an untranslated region of GAM205 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 205A, thereby inhibiting translation of GAM205 TARGET RNA into GAM205 TARGET PROTEIN, both of Fig. 205A.

[8408] It is appreciated that specific functions, and accordingly utilities, of GR588 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR588 GENE: GAM120 TARGET PROTEIN and GAM205 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 120D and 205D

[8409] Reference is now made to Fig. 589, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 589(GR589) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8410] GR589 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR589 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[8411] GR589 GENE encodes GR589 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8412] GR589 PRECURSOR RNA folds spatially, forming GR589 FOLDED PRECURSOR RNA . It is appreciated that GR589 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR589 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8413] GR589 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM31 PRECURSOR, GAM78 PRECURSOR and GAM177 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8414] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM31 RNA, GAM78 RNA and GAM177 RNA respectively, each of which

GAM RNAs corresponding to GAM RNA of Fig. 8.

[8415] GAM31 RNA binds complimentarily to a target binding site located in an untranslated region of GAM31 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 31A, thereby inhibiting translation of GAM31 TARGET RNA into GAM31 TARGET PROTEIN, both of Fig. 31A.

[8416] GAM78 RNA binds complimentarily to a target binding site located in an untranslated region of GAM78 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 78A, thereby inhibiting translation of GAM78 TARGET RNA into GAM78 TARGET PROTEIN, both of Fig. 78A.

[8417] GAM177 RNA binds complimentarily to a target binding site located in an untranslated region of GAM177 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 177A, thereby inhibiting translation of GAM177 TARGET RNA into GAM177 TARGET PROTEIN, both of Fig. 177A.

[8418] It is appreciated that specific functions, and accordingly

utilities, of GR589 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR589 GENE: GAM31 TARGET PROTEIN, GAM78 TARGET PROTEIN and GAM177 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 31D, 78D and 177D

[8419] Reference is now made to Fig. 590, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 590(GR590) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8420] GR590 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR590 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8421] GR590 GENE encodes GR590 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8422] GR590 PRECURSOR RNA folds spatially, forming GR590 FOLDED PRECURSOR RNA . It is appreciated that GR590

FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR590 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8423] GR590 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM258 PRECURSOR, GAM439 PRECURSOR and GAM537 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8424] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM258 RNA, GAM439 RNA and GAM537 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8425] GAM258 RNA binds complimentarily to a target binding site located in an untranslated region of GAM258 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 258A, thereby inhibiting translation of GAM258 TARGET RNA into GAM258 TARGET PROTEIN, both of Fig. 258A.

[8426] GAM439 RNA binds complimentarily to a target binding site located in an untranslated region of GAM439 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 439A, thereby inhibiting translation of GAM439 TARGET RNA into GAM439 TARGET PROTEIN, both of Fig. 439A.

[8427] GAM537 RNA binds complimentarily to a target binding site located in an untranslated region of GAM537 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 537A, thereby inhibiting translation of GAM537 TARGET RNA into GAM537 TARGET PROTEIN, both of Fig. 537A.

[8428] It is appreciated that specific functions, and accordingly utilities, of GR590 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR590 GENE: GAM258 TARGET PROTEIN, GAM439 TARGET PROTEIN and GAM537 TARGET PROTEIN. The function

of these target genes is elaborated hereinabove with reference to Figs. 258D, 439D and 537D

[8429] Reference is now made to Fig. 591, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 591(GR591) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8430] GR591 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR591 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8431] GR591 GENE encodes GR591 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8432] GR591 PRECURSOR RNA folds spatially, forming GR591 FOLDED PRECURSOR RNA . It is appreciated that GR591 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR591 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8433] GR591 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM209 PRECURSOR and GAM525 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8434] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM209 RNA and GAM525 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8435] GAM209 RNA binds complimentarily to a target binding site located in an untranslated region of GAM209 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 209A, thereby inhibiting translation of GAM209 TARGET RNA into GAM209 TARGET PROTEIN, both of Fig. 209A.

[8436] GAM525 RNA binds complimentarily to a target binding site located in an untranslated region of GAM525 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 525A, thereby inhibiting translation of GAM525 TARGET RNA into GAM525 TARGET PROTEIN, both of Fig. 525A.

[8437] It is appreciated that specific functions, and accordingly utilities, of GR591 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR591 GENE: GAM209 TARGET PROTEIN and GAM525 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 209D and 525D

[8438] Reference is now made to Fig. 592, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 592(GR592) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8439] GR592 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR592 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8440] GR592 GENE encodes GR592 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8441] GR592 PRECURSOR RNA folds spatially, forming GR592 FOLDED PRECURSOR RNA . It is appreciated that GR592 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR592 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8442] GR592 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 4 separate GAM precursor RNAs, GAM27 PRECURSOR, GAM423 PRECURSOR, GAM449 PRECURSOR and GAM459 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8443] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM27 RNA, GAM423 RNA, GAM449 RNA and GAM459 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8444] GAM27 RNA binds complimentarily to a target binding site located in an untranslated region of GAM27 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 27A, thereby inhibiting translation of GAM27 TARGET RNA into GAM27 TARGET PROTEIN, both of Fig. 27A.

[8445] GAM423 RNA binds complimentarily to a target binding site located in an untranslated region of GAM423 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 423A, thereby inhibiting translation of GAM423 TARGET RNA into GAM423 TARGET PROTEIN, both of Fig. 423A.

[8446] GAM449 RNA binds complimentarily to a target binding site located in an untranslated region of GAM449 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 449A, thereby inhibiting transla-

tion of GAM449 TARGET RNA into GAM449 TARGET PROTEIN, both of Fig. 449A.

[8447] GAM459 RNA binds complimentarily to a target binding site located in an untranslated region of GAM459 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 459A, thereby inhibiting translation of GAM459 TARGET RNA into GAM459 TARGET PROTEIN, both of Fig. 459A.

[8448] It is appreciated that specific functions, and accordingly utilities, of GR592 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR592 GENE: GAM27 TARGET PROTEIN, GAM423 TARGET PROTEIN, GAM449 TARGET PROTEIN and GAM459 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 27D, 423D, 449D and 459D

[8449] Reference is now made to Fig. 593, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 593(GR593) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn

modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8450] GR593 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR593 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8451] GR593 GENE encodes GR593 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8452] GR593 PRECURSOR RNA folds spatially, forming GR593 FOLDED PRECURSOR RNA . It is appreciated that GR593 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR593 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8453] GR593 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM32 PRECURSOR and GAM359 PRECURSOR, each of which GAM precursor RNAs being a hair-

pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8454] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM32 RNA and GAM359 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8455] GAM32 RNA binds complimentarily to a target binding site located in an untranslated region of GAM32 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 32A, thereby inhibiting translation of GAM32 TARGET RNA into GAM32 TARGET PROTEIN, both of Fig. 32A.

[8456] GAM359 RNA binds complimentarily to a target binding site located in an untranslated region of GAM359 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 359A, thereby inhibiting translation of GAM359 TARGET RNA into GAM359 TARGET PROTEIN, both of Fig. 359A.

[8457] It is appreciated that specific functions, and accordingly utilities, of GR593 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR593 GENE: GAM32 TARGET PROTEIN and GAM359 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 32D and 359D

[8458] Reference is now made to Fig. 594, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 594(GR594) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8459] GR594 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR594 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8460] GR594 GENE encodes GR594 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8461] GR594 PRECURSOR RNA folds spatially, forming GR594 FOLDED PRECURSOR RNA . It is appreciated that GR594 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR594 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8462] GR594 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 4 separate GAM precursor RNAs, GAM285 PRECURSOR, GAM326 PRECURSOR, GAM436 PRECURSOR and GAM498 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8463] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM285 RNA, GAM326 RNA, GAM436 RNA and GAM498 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8464] GAM285 RNA binds complimentarily to a target binding site located in an untranslated region of GAM285 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 285A, thereby inhibiting translation of GAM285 TARGET RNA into GAM285 TARGET PROTEIN, both of Fig. 285A.

[8465] GAM326 RNA binds complimentarily to a target binding site located in an untranslated region of GAM326 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 326A, thereby inhibiting translation of GAM326 TARGET RNA into GAM326 TARGET PROTEIN, both of Fig. 326A.

[8466] GAM436 RNA binds complimentarily to a target binding site located in an untranslated region of GAM436 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 436A, thereby inhibiting translation of GAM436 TARGET RNA into GAM436 TARGET PROTEIN, both of Fig. 436A.

[8467] GAM498 RNA binds complimentarily to a target binding site located in an untranslated region of GAM498 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 498A, thereby inhibiting transla-

tion of GAM498 TARGET RNA into GAM498 TARGET PROTEIN, both of Fig. 498A.

[8468] It is appreciated that specific functions, and accordingly utilities, of GR594 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR594 GENE: GAM285 TARGET PROTEIN, GAM326 TARGET PROTEIN, GAM436 TARGET PROTEIN and GAM498 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 285D, 326D, 436D and 498D

[8469] Reference is now made to Fig. 595, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 595(GR595) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8470] GR595 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR595 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8471] GR595 GENE encodes GR595 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8472] GR595 PRECURSOR RNA folds spatially, forming GR595 FOLDED PRECURSOR RNA . It is appreciated that GR595 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR595 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8473] GR595 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM101 PRECURSOR, GAM125 PRECURSOR and GAM183 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8474] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM101 RNA, GAM125 RNA and GAM183 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8475] GAM101 RNA binds complimentarily to a target binding site located in an untranslated region of GAM101 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 101A, thereby inhibiting translation of GAM101 TARGET RNA into GAM101 TARGET PROTEIN, both of Fig. 101A.

[8476] GAM125 RNA binds complimentarily to a target binding site located in an untranslated region of GAM125 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 125A, thereby inhibiting translation of GAM125 TARGET RNA into GAM125 TARGET PROTEIN, both of Fig. 125A.

[8477] GAM183 RNA binds complimentarily to a target binding site located in an untranslated region of GAM183 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 183A, thereby inhibiting translation of GAM183 TARGET RNA into GAM183 TARGET PROTEIN, both of Fig. 183A.

[8478] It is appreciated that specific functions, and accordingly utilities, of GR595 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR595 GENE: GAM101 TARGET PROTEIN, GAM125 TARGET PROTEIN and GAM183 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 101D, 125D and 183D

[8479] Reference is now made to Fig. 596, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 596(GR596) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8480] GR596 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR596 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8481] GR596 GENE encodes GR596 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8482] GR596 PRECURSOR RNA folds spatially, forming GR596 FOLDED PRECURSOR RNA . It is appreciated that GR596 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR596 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8483] GR596 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8484] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8485] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting transla-

tion of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8486] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8487] It is appreciated that specific functions, and accordingly utilities, of GR596 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR596 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8488] Reference is now made to Fig. 597, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 597(GR597) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the

function and utility of which at least one target gene is known in the art.

[8489] GR597 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR597 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8490] GR597 GENE encodes GR597 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8491] GR597 PRECURSOR RNA folds spatially, forming GR597 FOLDED PRECURSOR RNA . It is appreciated that GR597 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR597 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8492] GR597 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM155 PRECURSOR and GAM413 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECUR-

SOR RNA of Fig. 8.

[8493] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM155 RNA and GAM413 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8494] GAM155 RNA binds complimentarily to a target binding site located in an untranslated region of GAM155 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 155A, thereby inhibiting translation of GAM155 TARGET RNA into GAM155 TARGET PROTEIN, both of Fig. 155A.

[8495] GAM413 RNA binds complimentarily to a target binding site located in an untranslated region of GAM413 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 413A, thereby inhibiting translation of GAM413 TARGET RNA into GAM413 TARGET PROTEIN, both of Fig. 413A.

[8496] It is appreciated that specific functions, and accordingly utilities, of GR597 GENE correlate with, and may be deduced from, the identity of the target genes, which are in-

hibited by GAM RNAs comprised in the operon-like cluster of GR597 GENE: GAM155 TARGET PROTEIN and GAM413 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 155D and 413D

[8497] Reference is now made to Fig. 598, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 598(GR598) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8498] GR598 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR598 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8499] GR598 GENE encodes GR598 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8500] GR598 PRECURSOR RNA folds spatially, forming GR598 FOLDED PRECURSOR RNA . It is appreciated that GR598 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin

structures are due to the fact that the nucleotide sequence of GR598 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8501] GR598 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM39 PRECURSOR and GAM416 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8502] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM39 RNA and GAM416 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8503] GAM39 RNA binds complimentarily to a target binding site located in an untranslated region of GAM39 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 39A, thereby inhibiting translation of GAM39 TARGET RNA into GAM39 TARGET PROTEIN, both

of Fig. 39A.

[8504] GAM416 RNA binds complimentarily to a target binding site located in an untranslated region of GAM416 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 416A, thereby inhibiting translation of GAM416 TARGET RNA into GAM416 TARGET PROTEIN, both of Fig. 416A.

[8505] It is appreciated that specific functions, and accordingly utilities, of GR598 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR598 GENE: GAM39 TARGET PROTEIN and GAM416 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 39D and 416D

[8506] Reference is now made to Fig. 599, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 599(GR599) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is

known in the art.

[8507] GR599 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR599 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8508] GR599 GENE encodes GR599 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8509] GR599 PRECURSOR RNA folds spatially, forming GR599 FOLDED PRECURSOR RNA . It is appreciated that GR599 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR599 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8510] GR599 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM441 PRECURSOR and GAM527 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8511] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM441 RNA and GAM527 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8512] GAM441 RNA binds complimentarily to a target binding site located in an untranslated region of GAM441 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 441A, thereby inhibiting translation of GAM441 TARGET RNA into GAM441 TARGET PROTEIN, both of Fig. 441A.

[8513] GAM527 RNA binds complimentarily to a target binding site located in an untranslated region of GAM527 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 527A, thereby inhibiting translation of GAM527 TARGET RNA into GAM527 TARGET PROTEIN, both of Fig. 527A.

[8514] It is appreciated that specific functions, and accordingly utilities, of GR599 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR599 GENE: GAM441 TARGET PROTEIN and GAM527 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 441D and 527D

[8515] Reference is now made to Fig. 600, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 600(GR600) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8516] GR600 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR600 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8517] GR600 GENE encodes GR600 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8518] GR600 PRECURSOR RNA folds spatially, forming GR600 FOLDED PRECURSOR RNA . It is appreciated that GR600 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR600 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8519] GR600 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM254 PRECURSOR and GAM495 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8520] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM254 RNA and GAM495 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8521] GAM254 RNA binds complimentarily to a target binding site located in an untranslated region of GAM254 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 254A, thereby inhibiting translation of GAM254 TARGET RNA into GAM254 TARGET PROTEIN, both of Fig. 254A.

[8522] GAM495 RNA binds complimentarily to a target binding site located in an untranslated region of GAM495 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 495A, thereby inhibiting translation of GAM495 TARGET RNA into GAM495 TARGET PROTEIN, both of Fig. 495A.

[8523] It is appreciated that specific functions, and accordingly utilities, of GR600 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR600 GENE: GAM254 TARGET PROTEIN and GAM495 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 254D and 495D

[8524] Reference is now made to Fig. 601, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 601(GR601) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8525] GR601 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR601 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8526] GR601 GENE encodes GR601 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8527] GR601 PRECURSOR RNA folds spatially, forming GR601 FOLDED PRECURSOR RNA . It is appreciated that GR601 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR601 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8528] GR601 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM51 PRECURSOR and GAM298 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8529] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM51 RNA and GAM298 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8530] GAM51 RNA binds complimentarily to a target binding site located in an untranslated region of GAM51 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 51A, thereby inhibiting translation of GAM51 TARGET RNA into GAM51 TARGET PROTEIN, both of Fig. 51A.

[8531] GAM298 RNA binds complimentarily to a target binding site located in an untranslated region of GAM298 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 298A, thereby inhibiting translation of GAM298 TARGET RNA into GAM298 TARGET PROTEIN, both of Fig. 298A.

[8532] It is appreciated that specific functions, and accordingly utilities, of GR601 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR601 GENE: GAM51 TARGET PROTEIN and GAM298

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 51D and 298D

[8533] Reference is now made to Fig. 602, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 602(GR602) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8534] GR602 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR602 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8535] GR602 GENE encodes GR602 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8536] GR602 PRECURSOR RNA folds spatially, forming GR602 FOLDED PRECURSOR RNA . It is appreciated that GR602 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR602 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8537] GR602 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM222 PRECURSOR, GAM341 PRECURSOR and GAM484 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8538] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM222 RNA, GAM341 RNA and GAM484 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8539] GAM222 RNA binds complimentarily to a target binding site located in an untranslated region of GAM222 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 222A, thereby inhibiting translation of GAM222 TARGET RNA into GAM222 TARGET PROTEIN, both of Fig. 222A.

[8540] GAM341 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM341 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 341A, thereby inhibiting translation of GAM341 TARGET RNA into GAM341 TARGET PROTEIN, both of Fig. 341A.

[8541] GAM484 RNA binds complimentarily to a target binding site located in an untranslated region of GAM484 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 484A, thereby inhibiting translation of GAM484 TARGET RNA into GAM484 TARGET PROTEIN, both of Fig. 484A.

[8542] It is appreciated that specific functions, and accordingly utilities, of GR602 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR602 GENE: GAM222 TARGET PROTEIN, GAM341 TARGET PROTEIN and GAM484 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 222D, 341D and 484D

[8543] Reference is now made to Fig. 603, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 603(GR603) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8544] GR603 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR603 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8545] GR603 GENE encodes GR603 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8546] GR603 PRECURSOR RNA folds spatially, forming GR603 FOLDED PRECURSOR RNA . It is appreciated that GR603 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR603 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8547] GR603 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM314 PRECURSOR and GAM433 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8548] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM314 RNA and GAM433 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8549] GAM314 RNA binds complimentarily to a target binding site located in an untranslated region of GAM314 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 314A, thereby inhibiting translation of GAM314 TARGET RNA into GAM314 TARGET PROTEIN, both of Fig. 314A.

[8550] GAM433 RNA binds complimentarily to a target binding site located in an untranslated region of GAM433 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 433A, thereby inhibiting translation of GAM433 TARGET RNA into GAM433 TARGET PRO-

TEIN, both of Fig. 433A.

[8551] It is appreciated that specific functions, and accordingly utilities, of GR603 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR603 GENE: GAM314 TARGET PROTEIN and GAM433 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 314D and 433D

[8552] Reference is now made to Fig. 604, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 604(GR604) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8553] GR604 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR604 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8554] GR604 GENE encodes GR604 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8555] GR604 PRECURSOR RNA folds spatially, forming GR604 FOLDED PRECURSOR RNA . It is appreciated that GR604 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR604 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8556] GR604 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM109 PRECURSOR, GAM215 PRECURSOR and GAM418 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8557] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM109 RNA, GAM215 RNA and GAM418 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8558] GAM109 RNA binds complimentarily to a target binding site located in an untranslated region of GAM109 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 109A, thereby inhibiting translation of GAM109 TARGET RNA into GAM109 TARGET PROTEIN, both of Fig. 109A.

[8559] GAM215 RNA binds complimentarily to a target binding site located in an untranslated region of GAM215 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 215A, thereby inhibiting translation of GAM215 TARGET RNA into GAM215 TARGET PROTEIN, both of Fig. 215A.

[8560] GAM418 RNA binds complimentarily to a target binding site located in an untranslated region of GAM418 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 418A, thereby inhibiting translation of GAM418 TARGET RNA into GAM418 TARGET PROTEIN, both of Fig. 418A.

[8561] It is appreciated that specific functions, and accordingly utilities, of GR604 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR604 GENE: GAM109 TARGET PROTEIN, GAM215 TARGET PROTEIN and GAM418 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 109D, 215D and 418D

[8562] Reference is now made to Fig. 605, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 605(GR605) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8563] GR605 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR605 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8564] GR605 GENE encodes GR605 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8565] GR605 PRECURSOR RNA folds spatially, forming GR605 FOLDED PRECURSOR RNA . It is appreciated that GR605 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR605 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8566] GR605 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM46 PRECURSOR, GAM77 PRECURSOR and GAM382 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8567] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM46 RNA, GAM77 RNA and GAM382 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8568] GAM46 RNA binds complimentarily to a target binding site located in an untranslated region of GAM46 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 46A, thereby inhibiting translation of GAM46 TARGET RNA into GAM46 TARGET PROTEIN, both of Fig. 46A.

[8569] GAM77 RNA binds complimentarily to a target binding site located in an untranslated region of GAM77 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 77A, thereby inhibiting translation of GAM77 TARGET RNA into GAM77 TARGET PROTEIN, both of Fig. 77A.

[8570] GAM382 RNA binds complimentarily to a target binding site located in an untranslated region of GAM382 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 382A, thereby inhibiting translation of GAM382 TARGET RNA into GAM382 TARGET PROTEIN, both of Fig. 382A.

[8571] It is appreciated that specific functions, and accordingly utilities, of GR605 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR605 GENE: GAM46 TARGET PROTEIN, GAM77 TARGET PROTEIN and GAM382 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 46D, 77D and 382D

[8572] Reference is now made to Fig. 606, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 606(GR606) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8573] GR606 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR606 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8574] GR606 GENE encodes GR606 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8575] GR606 PRECURSOR RNA folds spatially, forming GR606 FOLDED PRECURSOR RNA . It is appreciated that GR606 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR606 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8576] GR606 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM447 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8577] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM447 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8578] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8579] GAM447 RNA binds complimentarily to a target binding site located in an untranslated region of GAM447 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 447A, thereby inhibiting transla-

tion of GAM447 TARGET RNA into GAM447 TARGET PROTEIN, both of Fig. 447A.

[8580] It is appreciated that specific functions, and accordingly utilities, of GR606 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR606 GENE: GAM218 TARGET PROTEIN and GAM447 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 447D

[8581] Reference is now made to Fig. 607, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 607(GR607) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8582] GR607 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR607 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8583] GR607 GENE encodes GR607 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[8584] GR607 PRECURSOR RNA folds spatially, forming GR607 FOLDED PRECURSOR RNA . It is appreciated that GR607 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR607 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8585] GR607 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM283 PRECURSOR and GAM297 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8586] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM283 RNA and GAM297 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8587] GAM283 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM283 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 283A, thereby inhibiting translation of GAM283 TARGET RNA into GAM283 TARGET PROTEIN, both of Fig. 283A.

[8588] GAM297 RNA binds complimentarily to a target binding site located in an untranslated region of GAM297 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 297A, thereby inhibiting translation of GAM297 TARGET RNA into GAM297 TARGET PROTEIN, both of Fig. 297A.

[8589] It is appreciated that specific functions, and accordingly utilities, of GR607 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR607 GENE: GAM283 TARGET PROTEIN and GAM297 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 283D and 297D

[8590] Reference is now made to Fig. 608, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 608(GR608) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8591] GR608 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR608 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8592] GR608 GENE encodes GR608 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8593] GR608 PRECURSOR RNA folds spatially, forming GR608 FOLDED PRECURSOR RNA . It is appreciated that GR608 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR608 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8594] GR608 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM38 PRECURSOR and GAM333 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8595] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM38 RNA and GAM333 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8596] GAM38 RNA binds complimentarily to a target binding site located in an untranslated region of GAM38 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 38A, thereby inhibiting translation of GAM38 TARGET RNA into GAM38 TARGET PROTEIN, both of Fig. 38A.

[8597] GAM333 RNA binds complimentarily to a target binding site located in an untranslated region of GAM333 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 333A, thereby inhibiting translation of GAM333 TARGET RNA into GAM333 TARGET PRO-

TEIN, both of Fig. 333A.

[8598] It is appreciated that specific functions, and accordingly utilities, of GR608 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR608 GENE: GAM38 TARGET PROTEIN and GAM333 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 38D and 333D

[8599] Reference is now made to Fig. 609, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 609(GR609) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8600] GR609 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR609 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8601] GR609 GENE encodes GR609 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8602] GR609 PRECURSOR RNA folds spatially, forming GR609 FOLDED PRECURSOR RNA . It is appreciated that GR609 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR609 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8603] GR609 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM265 PRECURSOR and GAM276 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8604] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM265 RNA and GAM276 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8605] GAM265 RNA binds complimentarily to a target binding site located in an untranslated region of GAM265 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 265A, thereby inhibiting translation of GAM265 TARGET RNA into GAM265 TARGET PROTEIN, both of Fig. 265A.

[8606] GAM276 RNA binds complimentarily to a target binding site located in an untranslated region of GAM276 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 276A, thereby inhibiting translation of GAM276 TARGET RNA into GAM276 TARGET PROTEIN, both of Fig. 276A.

[8607] It is appreciated that specific functions, and accordingly utilities, of GR609 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR609 GENE: GAM265 TARGET PROTEIN and GAM276 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 265D and 276D

[8608] Reference is now made to Fig. 610, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

610(GR610) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8609] GR610 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR610 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8610] GR610 GENE encodes GR610 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8611] GR610 PRECURSOR RNA folds spatially, forming GR610 FOLDED PRECURSOR RNA . It is appreciated that GR610 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR610 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8612] GR610 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM

precursor RNAs, GAM251 PRECURSOR, GAM306 PRECURSOR and GAM510 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8613] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM251 RNA, GAM306 RNA and GAM510 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8614] GAM251 RNA binds complimentarily to a target binding site located in an untranslated region of GAM251 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 251A, thereby inhibiting translation of GAM251 TARGET RNA into GAM251 TARGET PROTEIN, both of Fig. 251A.

[8615] GAM306 RNA binds complimentarily to a target binding site located in an untranslated region of GAM306 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 306A, thereby inhibiting translation of GAM306 TARGET RNA into GAM306 TARGET PROTEIN, both of Fig. 306A.

[8616] GAM510 RNA binds complimentarily to a target binding site located in an untranslated region of GAM510 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 510A, thereby inhibiting translation of GAM510 TARGET RNA into GAM510 TARGET PROTEIN, both of Fig. 510A.

[8617] It is appreciated that specific functions, and accordingly utilities, of GR610 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR610 GENE: GAM251 TARGET PROTEIN, GAM306 TARGET PROTEIN and GAM510 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 251D, 306D and 510D

[8618] Reference is now made to Fig. 611, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 611(GR611) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8619] GR611 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR611 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8620] GR611 GENE encodes GR611 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8621] GR611 PRECURSOR RNA folds spatially, forming GR611 FOLDED PRECURSOR RNA . It is appreciated that GR611 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR611 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8622] GR611 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM230 PRECURSOR and GAM307 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8623] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM230 RNA and GAM307 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8624] GAM230 RNA binds complimentarily to a target binding site located in an untranslated region of GAM230 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 230A, thereby inhibiting translation of GAM230 TARGET RNA into GAM230 TARGET PROTEIN, both of Fig. 230A.

[8625] GAM307 RNA binds complimentarily to a target binding site located in an untranslated region of GAM307 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 307A, thereby inhibiting translation of GAM307 TARGET RNA into GAM307 TARGET PROTEIN, both of Fig. 307A.

[8626] It is appreciated that specific functions, and accordingly utilities, of GR611 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR611 GENE: GAM230 TARGET PROTEIN and GAM307

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 230D and 307D

[8627] Reference is now made to Fig. 612, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 612(GR612) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8628] GR612 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR612 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8629] GR612 GENE encodes GR612 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8630] GR612 PRECURSOR RNA folds spatially, forming GR612 FOLDED PRECURSOR RNA . It is appreciated that GR612 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR612 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8631] GR612 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM296 PRECURSOR and GAM303 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8632] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM296 RNA and GAM303 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8633] GAM296 RNA binds complimentarily to a target binding site located in an untranslated region of GAM296 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 296A, thereby inhibiting translation of GAM296 TARGET RNA into GAM296 TARGET PROTEIN, both of Fig. 296A.

[8634] GAM303 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM303 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 303A, thereby inhibiting translation of GAM303 TARGET RNA into GAM303 TARGET PROTEIN, both of Fig. 303A.

[8635] It is appreciated that specific functions, and accordingly utilities, of GR612 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR612 GENE: GAM296 TARGET PROTEIN and GAM303 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 296D and 303D

[8636] Reference is now made to Fig. 613, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 613(GR613) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8637] GR613 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR613 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8638] GR613 GENE encodes GR613 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8639] GR613 PRECURSOR RNA folds spatially, forming GR613 FOLDED PRECURSOR RNA . It is appreciated that GR613 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR613 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8640] GR613 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM199 PRECURSOR, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8641] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM199 RNA, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8642] GAM199 RNA binds complimentarily to a target binding site located in an untranslated region of GAM199 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 199A, thereby inhibiting translation of GAM199 TARGET RNA into GAM199 TARGET PROTEIN, both of Fig. 199A.

[8643] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8644] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PRO-

TEIN, both of Fig. 448A.

[8645] It is appreciated that specific functions, and accordingly utilities, of GR613 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR613 GENE: GAM199 TARGET PROTEIN, GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 199D, 218D and 448D

[8646] Reference is now made to Fig. 614, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 614(GR614) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8647] GR614 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR614 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8648] GR614 GENE encodes GR614 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8649] GR614 PRECURSOR RNA folds spatially, forming GR614 FOLDED PRECURSOR RNA . It is appreciated that GR614 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR614 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8650] GR614 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM526 PRECURSOR and GAM541 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8651] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM526 RNA and GAM541 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8652] GAM526 RNA binds complimentarily to a target binding site located in an untranslated region of GAM526 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 526A, thereby inhibiting translation of GAM526 TARGET RNA into GAM526 TARGET PROTEIN, both of Fig. 526A.

[8653] GAM541 RNA binds complimentarily to a target binding site located in an untranslated region of GAM541 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 541A, thereby inhibiting translation of GAM541 TARGET RNA into GAM541 TARGET PROTEIN, both of Fig. 541A.

[8654] It is appreciated that specific functions, and accordingly utilities, of GR614 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR614 GENE: GAM526 TARGET PROTEIN and GAM541 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 526D and 541D

[8655] Reference is now made to Fig. 615, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

615(GR615) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8656] GR615 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR615 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8657] GR615 GENE encodes GR615 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8658] GR615 PRECURSOR RNA folds spatially, forming GR615 FOLDED PRECURSOR RNA . It is appreciated that GR615 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR615 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8659] GR615 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM

precursor RNAs, GAM113 PRECURSOR and GAM221 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8660] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM113 RNA and GAM221 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8661] GAM113 RNA binds complimentarily to a target binding site located in an untranslated region of GAM113 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 113A, thereby inhibiting translation of GAM113 TARGET RNA into GAM113 TARGET PROTEIN, both of Fig. 113A.

[8662] GAM221 RNA binds complimentarily to a target binding site located in an untranslated region of GAM221 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 221A, thereby inhibiting translation of GAM221 TARGET RNA into GAM221 TARGET PROTEIN, both of Fig. 221A.

[8663] It is appreciated that specific functions, and accordingly utilities, of GR615 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR615 GENE: GAM113 TARGET PROTEIN and GAM221 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 113D and 221D

[8664] Reference is now made to Fig. 616, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 616(GR616) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8665] GR616 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR616 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8666] GR616 GENE encodes GR616 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8667] GR616 PRECURSOR RNA folds spatially, forming GR616

FOLDED PRECURSOR RNA . It is appreciated that GR616 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR616 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8668] GR616 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM94 PRECURSOR and GAM175 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8669] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM94 RNA and GAM175 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8670] GAM94 RNA binds complimentarily to a target binding site located in an untranslated region of GAM94 TARGET RNA, which target binding site corresponds to a target binding

site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 94A, thereby inhibiting translation of GAM94 TARGET RNA into GAM94 TARGET PROTEIN, both of Fig. 94A.

[8671] GAM175 RNA binds complimentarily to a target binding site located in an untranslated region of GAM175 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 175A, thereby inhibiting translation of GAM175 TARGET RNA into GAM175 TARGET PROTEIN, both of Fig. 175A.

[8672] It is appreciated that specific functions, and accordingly utilities, of GR616 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR616 GENE: GAM94 TARGET PROTEIN and GAM175 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 94D and 175D

[8673] Reference is now made to Fig. 617, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 617(GR617) gene, which encodes an operon-like cluster

of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8674] GR617 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR617 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8675] GR617 GENE encodes GR617 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8676] GR617 PRECURSOR RNA folds spatially, forming GR617 FOLDED PRECURSOR RNA . It is appreciated that GR617 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR617 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8677] GR617 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM28 PRECURSOR and GAM274 PRE-

CURSOR, each of which GAM precursor RNAs being a hair-pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8678] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM28 RNA and GAM274 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8679] GAM28 RNA binds complimentarily to a target binding site located in an untranslated region of GAM28 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 28A, thereby inhibiting translation of GAM28 TARGET RNA into GAM28 TARGET PROTEIN, both of Fig. 28A.

[8680] GAM274 RNA binds complimentarily to a target binding site located in an untranslated region of GAM274 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 274A, thereby inhibiting translation of GAM274 TARGET RNA into GAM274 TARGET PROTEIN, both of Fig. 274A.

[8681] It is appreciated that specific functions, and accordingly

utilities, of GR617 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR617 GENE: GAM28 TARGET PROTEIN and GAM274 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 28D and 274D

[8682] Reference is now made to Fig. 618, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 618(GR618) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8683] GR618 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR618 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8684] GR618 GENE encodes GR618 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8685] GR618 PRECURSOR RNA folds spatially, forming GR618 FOLDED PRECURSOR RNA . It is appreciated that GR618

FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR618 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8686] GR618 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM174 PRECURSOR and GAM259 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8687] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM174 RNA and GAM259 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8688] GAM174 RNA binds complimentarily to a target binding site located in an untranslated region of GAM174 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 174A, thereby inhibiting translation of GAM174 TARGET RNA into GAM174 TARGET PROTEIN, both of Fig. 174A.

[8689] GAM259 RNA binds complimentarily to a target binding site located in an untranslated region of GAM259 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 259A, thereby inhibiting translation of GAM259 TARGET RNA into GAM259 TARGET PROTEIN, both of Fig. 259A.

[8690] It is appreciated that specific functions, and accordingly utilities, of GR618 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR618 GENE: GAM174 TARGET PROTEIN and GAM259 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 174D and 259D

[8691] Reference is now made to Fig. 619, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 619(GR619) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn

modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8692] GR619 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR619 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8693] GR619 GENE encodes GR619 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8694] GR619 PRECURSOR RNA folds spatially, forming GR619 FOLDED PRECURSOR RNA . It is appreciated that GR619 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR619 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8695] GR619 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM526 PRECURSOR and GAM541 PRECURSOR, each of which GAM precursor RNAs being a hair-

pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8696] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM526 RNA and GAM541 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8697] GAM526 RNA binds complimentarily to a target binding site located in an untranslated region of GAM526 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 526A, thereby inhibiting translation of GAM526 TARGET RNA into GAM526 TARGET PROTEIN, both of Fig. 526A.

[8698] GAM541 RNA binds complimentarily to a target binding site located in an untranslated region of GAM541 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 541A, thereby inhibiting translation of GAM541 TARGET RNA into GAM541 TARGET PROTEIN, both of Fig. 541A.

[8699] It is appreciated that specific functions, and accordingly utilities, of GR619 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR619 GENE: GAM526 TARGET PROTEIN and GAM541 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 526D and 541D

[8700] Reference is now made to Fig. 620, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 620(GR620) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8701] GR620 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR620 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8702] GR620 GENE encodes GR620 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8703] GR620 PRECURSOR RNA folds spatially, forming GR620 FOLDED PRECURSOR RNA . It is appreciated that GR620 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR620 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8704] GR620 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM172 PRECURSOR and GAM462 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8705] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM172 RNA and GAM462 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8706] GAM172 RNA binds complimentarily to a target binding site located in an untranslated region of GAM172 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 172A, thereby inhibiting transla-

tion of GAM172 TARGET RNA into GAM172 TARGET PROTEIN, both of Fig. 172A.

[8707] GAM462 RNA binds complimentarily to a target binding site located in an untranslated region of GAM462 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 462A, thereby inhibiting translation of GAM462 TARGET RNA into GAM462 TARGET PROTEIN, both of Fig. 462A.

[8708] It is appreciated that specific functions, and accordingly utilities, of GR620 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR620 GENE: GAM172 TARGET PROTEIN and GAM462 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 172D and 462D

[8709] Reference is now made to Fig. 621, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 621(GR621) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the

function and utility of which at least one target gene is known in the art.

[8710] GR621 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR621 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8711] GR621 GENE encodes GR621 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8712] GR621 PRECURSOR RNA folds spatially, forming GR621 FOLDED PRECURSOR RNA . It is appreciated that GR621 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR621 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8713] GR621 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM391 PRECURSOR and GAM500 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECUR-

SOR RNA of Fig. 8.

[8714] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM391 RNA and GAM500 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8715] GAM391 RNA binds complimentarily to a target binding site located in an untranslated region of GAM391 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 391A, thereby inhibiting translation of GAM391 TARGET RNA into GAM391 TARGET PROTEIN, both of Fig. 391A.

[8716] GAM500 RNA binds complimentarily to a target binding site located in an untranslated region of GAM500 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 500A, thereby inhibiting translation of GAM500 TARGET RNA into GAM500 TARGET PROTEIN, both of Fig. 500A.

[8717] It is appreciated that specific functions, and accordingly utilities, of GR621 GENE correlate with, and may be deduced from, the identity of the target genes, which are in-

hibited by GAM RNAs comprised in the operon-like cluster of GR621 GENE: GAM391 TARGET PROTEIN and GAM500 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 391D and 500D

[8718] Reference is now made to Fig. 622, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 622(GR622) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8719] GR622 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR622 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8720] GR622 GENE encodes GR622 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8721] GR622 PRECURSOR RNA folds spatially, forming GR622 FOLDED PRECURSOR RNA . It is appreciated that GR622 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin

structures are due to the fact that the nucleotide sequence of GR622 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8722] GR622 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM139 PRECURSOR, GAM159 PRECURSOR and GAM377 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8723] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM139 RNA, GAM159 RNA and GAM377 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8724] GAM139 RNA binds complimentarily to a target binding site located in an untranslated region of GAM139 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 139A, thereby inhibiting translation of GAM139 TARGET RNA into GAM139 TARGET PRO-

TEIN, both of Fig. 139A.

[8725] GAM159 RNA binds complimentarily to a target binding site located in an untranslated region of GAM159 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 159A, thereby inhibiting translation of GAM159 TARGET RNA into GAM159 TARGET PROTEIN, both of Fig. 159A.

[8726] GAM377 RNA binds complimentarily to a target binding site located in an untranslated region of GAM377 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 377A, thereby inhibiting translation of GAM377 TARGET RNA into GAM377 TARGET PROTEIN, both of Fig. 377A.

[8727] It is appreciated that specific functions, and accordingly utilities, of GR622 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR622 GENE: GAM139 TARGET PROTEIN, GAM159 TARGET PROTEIN and GAM377 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 139D, 159D and 377D

[8728] Reference is now made to Fig. 623, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 623(GR623) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8729] GR623 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR623 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8730] GR623 GENE encodes GR623 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8731] GR623 PRECURSOR RNA folds spatially, forming GR623 FOLDED PRECURSOR RNA . It is appreciated that GR623 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR623 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[8732] GR623 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM236 PRECURSOR and GAM360 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8733] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM236 RNA and GAM360 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8734] GAM236 RNA binds complimentarily to a target binding site located in an untranslated region of GAM236 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 236A, thereby inhibiting translation of GAM236 TARGET RNA into GAM236 TARGET PROTEIN, both of Fig. 236A.

[8735] GAM360 RNA binds complimentarily to a target binding site located in an untranslated region of GAM360 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 360A, thereby inhibiting translation of GAM360 TARGET RNA into GAM360 TARGET PROTEIN, both of Fig. 360A.

[8736] It is appreciated that specific functions, and accordingly utilities, of GR623 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR623 GENE: GAM236 TARGET PROTEIN and GAM360 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 236D and 360D

[8737] Reference is now made to Fig. 624, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 624(GR624) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8738] GR624 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR624 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8739] GR624 GENE encodes GR624 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8740] GR624 PRECURSOR RNA folds spatially, forming GR624 FOLDED PRECURSOR RNA . It is appreciated that GR624 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR624 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8741] GR624 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM66 PRECURSOR and GAM474 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8742] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM66 RNA and GAM474 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8743] GAM66 RNA binds complimentarily to a target binding site located in an untranslated region of GAM66 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 66A, thereby inhibiting translation of GAM66 TARGET RNA into GAM66 TARGET PROTEIN, both of Fig. 66A.

[8744] GAM474 RNA binds complimentarily to a target binding site located in an untranslated region of GAM474 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 474A, thereby inhibiting translation of GAM474 TARGET RNA into GAM474 TARGET PROTEIN, both of Fig. 474A.

[8745] It is appreciated that specific functions, and accordingly utilities, of GR624 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR624 GENE: GAM66 TARGET PROTEIN and GAM474 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 66D and 474D

[8746] Reference is now made to Fig. 625, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 625(GR625) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8747] GR625 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR625 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8748] GR625 GENE encodes GR625 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8749] GR625 PRECURSOR RNA folds spatially, forming GR625 FOLDED PRECURSOR RNA . It is appreciated that GR625 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR625 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8750] GR625 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM291 PRECURSOR and GAM419 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8751] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM291 RNA and GAM419 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8752] GAM291 RNA binds complimentarily to a target binding site located in an untranslated region of GAM291 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 291A, thereby inhibiting translation of GAM291 TARGET RNA into GAM291 TARGET PROTEIN, both of Fig. 291A.

[8753] GAM419 RNA binds complimentarily to a target binding site located in an untranslated region of GAM419 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 419A, thereby inhibiting transla-

tion of GAM419 TARGET RNA into GAM419 TARGET PROTEIN, both of Fig. 419A.

[8754] It is appreciated that specific functions, and accordingly utilities, of GR625 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR625 GENE: GAM291 TARGET PROTEIN and GAM419 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 291D and 419D

[8755] Reference is now made to Fig. 626, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 626(GR626) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8756] GR626 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR626 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8757] GR626 GENE encodes GR626 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[8758] GR626 PRECURSOR RNA folds spatially, forming GR626 FOLDED PRECURSOR RNA . It is appreciated that GR626 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR626 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8759] GR626 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM30 PRECURSOR and GAM227 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8760] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM30 RNA and GAM227 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8761] GAM30 RNA binds complimentarily to a target binding site

located in an untranslated region of GAM30 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 30A, thereby inhibiting translation of GAM30 TARGET RNA into GAM30 TARGET PROTEIN, both of Fig. 30A.

[8762] GAM227 RNA binds complimentarily to a target binding site located in an untranslated region of GAM227 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 227A, thereby inhibiting translation of GAM227 TARGET RNA into GAM227 TARGET PROTEIN, both of Fig. 227A.

[8763] It is appreciated that specific functions, and accordingly utilities, of GR626 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR626 GENE: GAM30 TARGET PROTEIN and GAM227 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 30D and 227D

[8764] Reference is now made to Fig. 627, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 627(GR627) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8765] GR627 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR627 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8766] GR627 GENE encodes GR627 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8767] GR627 PRECURSOR RNA folds spatially, forming GR627 FOLDED PRECURSOR RNA . It is appreciated that GR627 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR627 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8768] GR627 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM196 PRECURSOR, GAM411 PRECURSOR and GAM468 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8769] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM196 RNA, GAM411 RNA and GAM468 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8770] GAM196 RNA binds complimentarily to a target binding site located in an untranslated region of GAM196 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 196A, thereby inhibiting translation of GAM196 TARGET RNA into GAM196 TARGET PROTEIN, both of Fig. 196A.

[8771] GAM411 RNA binds complimentarily to a target binding site located in an untranslated region of GAM411 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 411A, thereby inhibiting translation of GAM411 TARGET RNA into GAM411 TARGET PRO-

TEIN, both of Fig. 411A.

[8772] GAM468 RNA binds complimentarily to a target binding site located in an untranslated region of GAM468 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 468A, thereby inhibiting translation of GAM468 TARGET RNA into GAM468 TARGET PROTEIN, both of Fig. 468A.

[8773] It is appreciated that specific functions, and accordingly utilities, of GR627 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR627 GENE: GAM196 TARGET PROTEIN, GAM411 TARGET PROTEIN and GAM468 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 196D, 411D and 468D

[8774] Reference is now made to Fig. 628, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 628(GR628) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is

known in the art.

[8775] GR628 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR628 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8776] GR628 GENE encodes GR628 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8777] GR628 PRECURSOR RNA folds spatially, forming GR628 FOLDED PRECURSOR RNA . It is appreciated that GR628 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR628 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8778] GR628 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM146 PRECURSOR and GAM410 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8779] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM146 RNA and GAM410 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8780] GAM146 RNA binds complimentarily to a target binding site located in an untranslated region of GAM146 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 146A, thereby inhibiting translation of GAM146 TARGET RNA into GAM146 TARGET PROTEIN, both of Fig. 146A.

[8781] GAM410 RNA binds complimentarily to a target binding site located in an untranslated region of GAM410 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 410A, thereby inhibiting translation of GAM410 TARGET RNA into GAM410 TARGET PROTEIN, both of Fig. 410A.

[8782] It is appreciated that specific functions, and accordingly utilities, of GR628 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR628 GENE: GAM146 TARGET PROTEIN and GAM410 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 146D and 410D

[8783] Reference is now made to Fig. 629, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 629(GR629) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8784] GR629 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR629 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8785] GR629 GENE encodes GR629 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8786] GR629 PRECURSOR RNA folds spatially, forming GR629 FOLDED PRECURSOR RNA . It is appreciated that GR629 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR629 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8787] GR629 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM223 PRECURSOR and GAM356 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8788] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM223 RNA and GAM356 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8789] GAM223 RNA binds complimentarily to a target binding site located in an untranslated region of GAM223 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 223A, thereby inhibiting translation of GAM223 TARGET RNA into GAM223 TARGET PROTEIN, both of Fig. 223A.

[8790] GAM356 RNA binds complimentarily to a target binding site located in an untranslated region of GAM356 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 356A, thereby inhibiting translation of GAM356 TARGET RNA into GAM356 TARGET PROTEIN, both of Fig. 356A.

[8791] It is appreciated that specific functions, and accordingly utilities, of GR629 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR629 GENE: GAM223 TARGET PROTEIN and GAM356 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 223D and 356D

[8792] Reference is now made to Fig. 630, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 630(GR630) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8793] GR630 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR630 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8794] GR630 GENE encodes GR630 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8795] GR630 PRECURSOR RNA folds spatially, forming GR630 FOLDED PRECURSOR RNA . It is appreciated that GR630 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR630 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8796] GR630 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM173 PRECURSOR and GAM351 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8797] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM173 RNA and GAM351 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8798] GAM173 RNA binds complimentarily to a target binding site located in an untranslated region of GAM173 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 173A, thereby inhibiting translation of GAM173 TARGET RNA into GAM173 TARGET PROTEIN, both of Fig. 173A.

[8799] GAM351 RNA binds complimentarily to a target binding site located in an untranslated region of GAM351 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 351A, thereby inhibiting translation of GAM351 TARGET RNA into GAM351 TARGET PROTEIN, both of Fig. 351A.

[8800] It is appreciated that specific functions, and accordingly utilities, of GR630 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR630 GENE: GAM173 TARGET PROTEIN and GAM351

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 173D and 351D

[8801] Reference is now made to Fig. 631, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 631(GR631) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8802] GR631 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR631 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8803] GR631 GENE encodes GR631 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8804] GR631 PRECURSOR RNA folds spatially, forming GR631 FOLDED PRECURSOR RNA . It is appreciated that GR631 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR631 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8805] GR631 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8806] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8807] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8808] GAM448 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8809] It is appreciated that specific functions, and accordingly utilities, of GR631 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR631 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8810] Reference is now made to Fig. 632, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 632(GR632) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8811] GR632 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR632 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8812] GR632 GENE encodes GR632 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8813] GR632 PRECURSOR RNA folds spatially, forming GR632 FOLDED PRECURSOR RNA . It is appreciated that GR632 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR632 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8814] GR632 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM104 PRECURSOR, GAM162 PRECURSOR and GAM497 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8815] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM104 RNA, GAM162 RNA and GAM497 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8816] GAM104 RNA binds complimentarily to a target binding site located in an untranslated region of GAM104 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 104A, thereby inhibiting translation of GAM104 TARGET RNA into GAM104 TARGET PROTEIN, both of Fig. 104A.

[8817] GAM162 RNA binds complimentarily to a target binding site located in an untranslated region of GAM162 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 162A, thereby inhibiting translation of GAM162 TARGET RNA into GAM162 TARGET PROTEIN, both of Fig. 162A.

[8818] GAM497 RNA binds complimentarily to a target binding site located in an untranslated region of GAM497 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 497A, thereby inhibiting translation of GAM497 TARGET RNA into GAM497 TARGET PRO-

TEIN, both of Fig. 497A.

[8819] It is appreciated that specific functions, and accordingly utilities, of GR632 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR632 GENE: GAM104 TARGET PROTEIN, GAM162 TARGET PROTEIN and GAM497 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 104D, 162D and 497D

[8820] Reference is now made to Fig. 633, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 633(GR633) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8821] GR633 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR633 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8822] GR633 GENE encodes GR633 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8823] GR633 PRECURSOR RNA folds spatially, forming GR633 FOLDED PRECURSOR RNA . It is appreciated that GR633 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR633 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8824] GR633 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8825] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8826] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8827] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8828] It is appreciated that specific functions, and accordingly utilities, of GR633 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR633 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8829] Reference is now made to Fig. 634, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

634(GR634) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8830] GR634 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR634 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8831] GR634 GENE encodes GR634 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8832] GR634 PRECURSOR RNA folds spatially, forming GR634 FOLDED PRECURSOR RNA . It is appreciated that GR634 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR634 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8833] GR634 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM

precursor RNAs, GAM52 PRECURSOR, GAM367 PRECURSOR and GAM395 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8834] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM52 RNA, GAM367 RNA and GAM395 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8835] GAM52 RNA binds complimentarily to a target binding site located in an untranslated region of GAM52 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 52A, thereby inhibiting translation of GAM52 TARGET RNA into GAM52 TARGET PROTEIN, both of Fig. 52A.

[8836] GAM367 RNA binds complimentarily to a target binding site located in an untranslated region of GAM367 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 367A, thereby inhibiting translation of GAM367 TARGET RNA into GAM367 TARGET PROTEIN, both of Fig. 367A.

[8837] GAM395 RNA binds complimentarily to a target binding site located in an untranslated region of GAM395 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 395A, thereby inhibiting translation of GAM395 TARGET RNA into GAM395 TARGET PROTEIN, both of Fig. 395A.

[8838] It is appreciated that specific functions, and accordingly utilities, of GR634 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR634 GENE: GAM52 TARGET PROTEIN, GAM367 TARGET PROTEIN and GAM395 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 52D, 367D and 395D

[8839] Reference is now made to Fig. 635, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 635(GR635) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8840] GR635 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR635 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8841] GR635 GENE encodes GR635 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8842] GR635 PRECURSOR RNA folds spatially, forming GR635 FOLDED PRECURSOR RNA . It is appreciated that GR635 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR635 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8843] GR635 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM216 PRECURSOR and GAM286 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8844] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM216 RNA and GAM286 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8845] GAM216 RNA binds complimentarily to a target binding site located in an untranslated region of GAM216 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 216A, thereby inhibiting translation of GAM216 TARGET RNA into GAM216 TARGET PROTEIN, both of Fig. 216A.

[8846] GAM286 RNA binds complimentarily to a target binding site located in an untranslated region of GAM286 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 286A, thereby inhibiting translation of GAM286 TARGET RNA into GAM286 TARGET PROTEIN, both of Fig. 286A.

[8847] It is appreciated that specific functions, and accordingly utilities, of GR635 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR635 GENE: GAM216 TARGET PROTEIN and GAM286

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 216D and 286D

[8848] Reference is now made to Fig. 636, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 636(GR636) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8849] GR636 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR636 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8850] GR636 GENE encodes GR636 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8851] GR636 PRECURSOR RNA folds spatially, forming GR636 FOLDED PRECURSOR RNA . It is appreciated that GR636 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR636 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8852] GR636 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8853] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8854] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8855] GAM448 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8856] It is appreciated that specific functions, and accordingly utilities, of GR636 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR636 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8857] Reference is now made to Fig. 637, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 637(GR637) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8858] GR637 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR637 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8859] GR637 GENE encodes GR637 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8860] GR637 PRECURSOR RNA folds spatially, forming GR637 FOLDED PRECURSOR RNA . It is appreciated that GR637 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR637 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8861] GR637 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM188 PRECURSOR and GAM234 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8862] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM188 RNA and GAM234 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8863] GAM188 RNA binds complimentarily to a target binding site located in an untranslated region of GAM188 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 188A, thereby inhibiting translation of GAM188 TARGET RNA into GAM188 TARGET PROTEIN, both of Fig. 188A.

[8864] GAM234 RNA binds complimentarily to a target binding site located in an untranslated region of GAM234 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 234A, thereby inhibiting translation of GAM234 TARGET RNA into GAM234 TARGET PROTEIN, both of Fig. 234A.

[8865] It is appreciated that specific functions, and accordingly utilities, of GR637 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR637 GENE: GAM188 TARGET PROTEIN and GAM234 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 188D and 234D

[8866] Reference is now made to Fig. 638, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 638(GR638) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8867] GR638 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR638 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8868] GR638 GENE encodes GR638 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8869] GR638 PRECURSOR RNA folds spatially, forming GR638 FOLDED PRECURSOR RNA . It is appreciated that GR638 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR638 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8870] GR638 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8871] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8872] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8873] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[8874] It is appreciated that specific functions, and accordingly utilities, of GR638 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR638 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[8875] Reference is now made to Fig. 639, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 639(GR639) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8876] GR639 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR639 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8877] GR639 GENE encodes GR639 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8878] GR639 PRECURSOR RNA folds spatially, forming GR639 FOLDED PRECURSOR RNA . It is appreciated that GR639 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR639 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8879] GR639 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM102 PRECURSOR and GAM128 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8880] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM102 RNA

and GAM128 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8881] GAM102 RNA binds complimentarily to a target binding site located in an untranslated region of GAM102 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 102A, thereby inhibiting translation of GAM102 TARGET RNA into GAM102 TARGET PROTEIN, both of Fig. 102A.

[8882] GAM128 RNA binds complimentarily to a target binding site located in an untranslated region of GAM128 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 128A, thereby inhibiting translation of GAM128 TARGET RNA into GAM128 TARGET PROTEIN, both of Fig. 128A.

[8883] It is appreciated that specific functions, and accordingly utilities, of GR639 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR639 GENE: GAM102 TARGET PROTEIN and GAM128 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 102D and

[8884] Reference is now made to Fig. 640, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 640(GR640) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8885] GR640 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR640 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8886] GR640 GENE encodes GR640 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8887] GR640 PRECURSOR RNA folds spatially, forming GR640 FOLDED PRECURSOR RNA . It is appreciated that GR640 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR640 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[8888] GR640 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM197 PRECURSOR and GAM475 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8889] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM197 RNA and GAM475 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8890] GAM197 RNA binds complimentarily to a target binding site located in an untranslated region of GAM197 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 197A, thereby inhibiting translation of GAM197 TARGET RNA into GAM197 TARGET PROTEIN, both of Fig. 197A.

[8891] GAM475 RNA binds complimentarily to a target binding site located in an untranslated region of GAM475 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 475A, thereby inhibiting translation of GAM475 TARGET RNA into GAM475 TARGET PROTEIN, both of Fig. 475A.

[8892] It is appreciated that specific functions, and accordingly utilities, of GR640 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR640 GENE: GAM197 TARGET PROTEIN and GAM475 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 197D and 475D

[8893] Reference is now made to Fig. 641, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 641(GR641) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8894] GR641 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR641 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[8895] GR641 GENE encodes GR641 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8896] GR641 PRECURSOR RNA folds spatially, forming GR641 FOLDED PRECURSOR RNA . It is appreciated that GR641 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR641 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8897] GR641 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM515 PRECURSOR and GAM516 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8898] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM515 RNA and GAM516 RNA respectively, each of which GAM RNAs

corresponding to GAM RNA of Fig. 8.

[8899] GAM515 RNA binds complimentarily to a target binding site located in an untranslated region of GAM515 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 515A, thereby inhibiting translation of GAM515 TARGET RNA into GAM515 TARGET PROTEIN, both of Fig. 515A.

[8900] GAM516 RNA binds complimentarily to a target binding site located in an untranslated region of GAM516 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 516A, thereby inhibiting translation of GAM516 TARGET RNA into GAM516 TARGET PROTEIN, both of Fig. 516A.

[8901] It is appreciated that specific functions, and accordingly utilities, of GR641 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR641 GENE: GAM515 TARGET PROTEIN and GAM516 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 515D and 516D

[8902] Reference is now made to Fig. 642, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 642(GR642) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8903] GR642 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR642 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8904] GR642 GENE encodes GR642 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8905] GR642 PRECURSOR RNA folds spatially, forming GR642 FOLDED PRECURSOR RNA . It is appreciated that GR642 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR642 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[8906] GR642 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM81 PRECURSOR and GAM490 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8907] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM81 RNA and GAM490 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8908] GAM81 RNA binds complimentarily to a target binding site located in an untranslated region of GAM81 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 81A, thereby inhibiting translation of GAM81 TARGET RNA into GAM81 TARGET PROTEIN, both of Fig. 81A.

[8909] GAM490 RNA binds complimentarily to a target binding site located in an untranslated region of GAM490 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 490A, thereby inhibiting translation of GAM490 TARGET RNA into GAM490 TARGET PROTEIN, both of Fig. 490A.

[8910] It is appreciated that specific functions, and accordingly utilities, of GR642 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR642 GENE: GAM81 TARGET PROTEIN and GAM490 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 81D and 490D

[8911] Reference is now made to Fig. 643, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 643(GR643) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8912] GR643 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR643 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8913] GR643 GENE encodes GR643 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8914] GR643 PRECURSOR RNA folds spatially, forming GR643 FOLDED PRECURSOR RNA . It is appreciated that GR643 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR643 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8915] GR643 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM364 PRECURSOR and GAM538 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8916] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM364 RNA and GAM538 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8917] GAM364 RNA binds complimentarily to a target binding site located in an untranslated region of GAM364 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 364A, thereby inhibiting translation of GAM364 TARGET RNA into GAM364 TARGET PROTEIN, both of Fig. 364A.

[8918] GAM538 RNA binds complimentarily to a target binding site located in an untranslated region of GAM538 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 538A, thereby inhibiting translation of GAM538 TARGET RNA into GAM538 TARGET PROTEIN, both of Fig. 538A.

[8919] It is appreciated that specific functions, and accordingly utilities, of GR643 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR643 GENE: GAM364 TARGET PROTEIN and GAM538 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 364D and 538D

[8920] Reference is now made to Fig. 644, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 644(GR644) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8921] GR644 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR644 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8922] GR644 GENE encodes GR644 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8923] GR644 PRECURSOR RNA folds spatially, forming GR644 FOLDED PRECURSOR RNA . It is appreciated that GR644 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR644 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8924] GR644 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM321 PRECURSOR and GAM422 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8925] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM321 RNA and GAM422 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8926] GAM321 RNA binds complimentarily to a target binding site located in an untranslated region of GAM321 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 321A, thereby inhibiting translation of GAM321 TARGET RNA into GAM321 TARGET PROTEIN, both of Fig. 321A.

[8927] GAM422 RNA binds complimentarily to a target binding site located in an untranslated region of GAM422 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 422A, thereby inhibiting transla-

tion of GAM422 TARGET RNA into GAM422 TARGET PROTEIN, both of Fig. 422A.

[8928] It is appreciated that specific functions, and accordingly utilities, of GR644 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR644 GENE: GAM321 TARGET PROTEIN and GAM422 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 321D and 422D

[8929] Reference is now made to Fig. 645, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 645(GR645) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8930] GR645 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR645 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8931] GR645 GENE encodes GR645 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[8932] GR645 PRECURSOR RNA folds spatially, forming GR645 FOLDED PRECURSOR RNA . It is appreciated that GR645 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR645 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8933] GR645 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM147 PRECURSOR and GAM361 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8934] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM147 RNA and GAM361 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8935] GAM147 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM147 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 147A, thereby inhibiting translation of GAM147 TARGET RNA into GAM147 TARGET PROTEIN, both of Fig. 147A.

[8936] GAM361 RNA binds complimentarily to a target binding site located in an untranslated region of GAM361 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 361A, thereby inhibiting translation of GAM361 TARGET RNA into GAM361 TARGET PROTEIN, both of Fig. 361A.

[8937] It is appreciated that specific functions, and accordingly utilities, of GR645 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR645 GENE: GAM147 TARGET PROTEIN and GAM361 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 147D and 361D

[8938] Reference is now made to Fig. 646, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 646(GR646) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8939] GR646 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR646 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8940] GR646 GENE encodes GR646 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8941] GR646 PRECURSOR RNA folds spatially, forming GR646 FOLDED PRECURSOR RNA . It is appreciated that GR646 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR646 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8942] GR646 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM230 PRECURSOR and GAM307 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8943] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM230 RNA and GAM307 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8944] GAM230 RNA binds complimentarily to a target binding site located in an untranslated region of GAM230 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 230A, thereby inhibiting translation of GAM230 TARGET RNA into GAM230 TARGET PROTEIN, both of Fig. 230A.

[8945] GAM307 RNA binds complimentarily to a target binding site located in an untranslated region of GAM307 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 307A, thereby inhibiting translation of GAM307 TARGET RNA into GAM307 TARGET PRO-

TEIN, both of Fig. 307A.

[8946] It is appreciated that specific functions, and accordingly utilities, of GR646 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR646 GENE: GAM230 TARGET PROTEIN and GAM307 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 230D and 307D

[8947] Reference is now made to Fig. 647, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 647(GR647) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8948] GR647 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR647 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8949] GR647 GENE encodes GR647 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8950] GR647 PRECURSOR RNA folds spatially, forming GR647 FOLDED PRECURSOR RNA . It is appreciated that GR647 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR647 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8951] GR647 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM60 PRECURSOR and GAM414 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8952] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM60 RNA and GAM414 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8953] GAM60 RNA binds complimentarily to a target binding site located in an untranslated region of GAM60 TARGET RNA,

which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 60A, thereby inhibiting translation of GAM60 TARGET RNA into GAM60 TARGET PROTEIN, both of Fig. 60A.

[8954] GAM414 RNA binds complimentarily to a target binding site located in an untranslated region of GAM414 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 414A, thereby inhibiting translation of GAM414 TARGET RNA into GAM414 TARGET PROTEIN, both of Fig. 414A.

[8955] It is appreciated that specific functions, and accordingly utilities, of GR647 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR647 GENE: GAM60 TARGET PROTEIN and GAM414 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 60D and 414D

[8956] Reference is now made to Fig. 648, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

648(GR648) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8957] GR648 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR648 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8958] GR648 GENE encodes GR648 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8959] GR648 PRECURSOR RNA folds spatially, forming GR648 FOLDED PRECURSOR RNA . It is appreciated that GR648 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR648 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8960] GR648 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM

precursor RNAs, GAM218 PRECURSOR and GAM447 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8961] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM447 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8962] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[8963] GAM447 RNA binds complimentarily to a target binding site located in an untranslated region of GAM447 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 447A, thereby inhibiting translation of GAM447 TARGET RNA into GAM447 TARGET PROTEIN, both of Fig. 447A.

[8964] It is appreciated that specific functions, and accordingly utilities, of GR648 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR648 GENE: GAM218 TARGET PROTEIN and GAM447 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 447D

[8965] Reference is now made to Fig. 649, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 649(GR649) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8966] GR649 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR649 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8967] GR649 GENE encodes GR649 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8968] GR649 PRECURSOR RNA folds spatially, forming GR649

FOLDED PRECURSOR RNA . It is appreciated that GR649 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR649 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8969] GR649 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM88 PRECURSOR and GAM363 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8970] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM88 RNA and GAM363 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8971] GAM88 RNA binds complimentarily to a target binding site located in an untranslated region of GAM88 TARGET RNA, which target binding site corresponds to a target binding

site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 88A, thereby inhibiting translation of GAM88 TARGET RNA into GAM88 TARGET PROTEIN, both of Fig. 88A.

[8972] GAM363 RNA binds complimentarily to a target binding site located in an untranslated region of GAM363 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 363A, thereby inhibiting translation of GAM363 TARGET RNA into GAM363 TARGET PROTEIN, both of Fig. 363A.

[8973] It is appreciated that specific functions, and accordingly utilities, of GR649 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR649 GENE: GAM88 TARGET PROTEIN and GAM363 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 88D and 363D

[8974] Reference is now made to Fig. 650, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 650(GR650) gene, which encodes an operon-like cluster

of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8975] GR650 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR650 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8976] GR650 GENE encodes GR650 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8977] GR650 PRECURSOR RNA folds spatially, forming GR650 FOLDED PRECURSOR RNA . It is appreciated that GR650 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR650 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8978] GR650 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM99 PRECURSOR, GAM434 PRECUR-

SOR and GAM492 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8979] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM99 RNA, GAM434 RNA and GAM492 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8980] GAM99 RNA binds complimentarily to a target binding site located in an untranslated region of GAM99 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 99A, thereby inhibiting translation of GAM99 TARGET RNA into GAM99 TARGET PROTEIN, both of Fig. 99A.

[8981] GAM434 RNA binds complimentarily to a target binding site located in an untranslated region of GAM434 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 434A, thereby inhibiting translation of GAM434 TARGET RNA into GAM434 TARGET PROTEIN, both of Fig. 434A.

[8982] GAM492 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM492 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 492A, thereby inhibiting translation of GAM492 TARGET RNA into GAM492 TARGET PROTEIN, both of Fig. 492A.

[8983] It is appreciated that specific functions, and accordingly utilities, of GR650 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR650 GENE: GAM99 TARGET PROTEIN, GAM434 TARGET PROTEIN and GAM492 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 99D, 434D and 492D

[8984] Reference is now made to Fig. 651, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 651(GR651) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8985] GR651 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR651 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8986] GR651 GENE encodes GR651 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8987] GR651 PRECURSOR RNA folds spatially, forming GR651 FOLDED PRECURSOR RNA . It is appreciated that GR651 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR651 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8988] GR651 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM83 PRECURSOR and GAM132 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8989] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM83 RNA and GAM132 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8990] GAM83 RNA binds complimentarily to a target binding site located in an untranslated region of GAM83 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 83A, thereby inhibiting translation of GAM83 TARGET RNA into GAM83 TARGET PROTEIN, both of Fig. 83A.

[8991] GAM132 RNA binds complimentarily to a target binding site located in an untranslated region of GAM132 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 132A, thereby inhibiting translation of GAM132 TARGET RNA into GAM132 TARGET PROTEIN, both of Fig. 132A.

[8992] It is appreciated that specific functions, and accordingly utilities, of GR651 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR651 GENE: GAM83 TARGET PROTEIN and GAM132 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 83D and 132D

[8993] Reference is now made to Fig. 652, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 652(GR652) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[8994] GR652 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR652 GENE was detected is described hereinabove with reference to Figs. 6-15.

[8995] GR652 GENE encodes GR652 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[8996] GR652 PRECURSOR RNA folds spatially, forming GR652 FOLDED PRECURSOR RNA . It is appreciated that GR652 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR652 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[8997] GR652 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM50 PRECURSOR and GAM300 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[8998] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM50 RNA and GAM300 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[8999] GAM50 RNA binds complimentarily to a target binding site located in an untranslated region of GAM50 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 50A, thereby inhibiting translation of GAM50 TARGET RNA into GAM50 TARGET PROTEIN, both of Fig. 50A.

[9000] GAM300 RNA binds complimentarily to a target binding site located in an untranslated region of GAM300 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 300A, thereby inhibiting translation of GAM300 TARGET RNA into GAM300 TARGET PROTEIN, both of Fig. 300A.

[9001] It is appreciated that specific functions, and accordingly utilities, of GR652 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR652 GENE: GAM50 TARGET PROTEIN and GAM300 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 50D and 300D

[9002] Reference is now made to Fig. 653, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 653(GR653) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9003] GR653 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR653 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9004] GR653 GENE encodes GR653 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9005] GR653 PRECURSOR RNA folds spatially, forming GR653 FOLDED PRECURSOR RNA . It is appreciated that GR653 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR653 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9006] GR653 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM64 PRECURSOR and GAM169 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9007] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM64 RNA

and GAM169 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9008] GAM64 RNA binds complimentarily to a target binding site located in an untranslated region of GAM64 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 64A, thereby inhibiting translation of GAM64 TARGET RNA into GAM64 TARGET PROTEIN, both of Fig. 64A.

[9009] GAM169 RNA binds complimentarily to a target binding site located in an untranslated region of GAM169 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 169A, thereby inhibiting translation of GAM169 TARGET RNA into GAM169 TARGET PROTEIN, both of Fig. 169A.

[9010] It is appreciated that specific functions, and accordingly utilities, of GR653 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR653 GENE: GAM64 TARGET PROTEIN and GAM169 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 64D and

[9011] Reference is now made to Fig. 654, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 654(GR654) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9012] GR654 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR654 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9013] GR654 GENE encodes GR654 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9014] GR654 PRECURSOR RNA folds spatially, forming GR654 FOLDED PRECURSOR RNA . It is appreciated that GR654 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR654 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[9015] GR654 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM133 PRECURSOR and GAM284 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9016] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM133 RNA and GAM284 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9017] GAM133 RNA binds complimentarily to a target binding site located in an untranslated region of GAM133 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 133A, thereby inhibiting translation of GAM133 TARGET RNA into GAM133 TARGET PROTEIN, both of Fig. 133A.

[9018] GAM284 RNA binds complimentarily to a target binding site located in an untranslated region of GAM284 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 284A, thereby inhibiting translation of GAM284 TARGET RNA into GAM284 TARGET PROTEIN, both of Fig. 284A.

[9019] It is appreciated that specific functions, and accordingly utilities, of GR654 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR654 GENE: GAM133 TARGET PROTEIN and GAM284 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 133D and 284D

[9020] Reference is now made to Fig. 655, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 655(GR655) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9021] GR655 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR655 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[9022] GR655 GENE encodes GR655 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9023] GR655 PRECURSOR RNA folds spatially, forming GR655 FOLDED PRECURSOR RNA . It is appreciated that GR655 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR655 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9024] GR655 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM447 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9025] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM447 RNA respectively, each of which GAM RNAs

corresponding to GAM RNA of Fig. 8.

[9026] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[9027] GAM447 RNA binds complimentarily to a target binding site located in an untranslated region of GAM447 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 447A, thereby inhibiting translation of GAM447 TARGET RNA into GAM447 TARGET PROTEIN, both of Fig. 447A.

[9028] It is appreciated that specific functions, and accordingly utilities, of GR655 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR655 GENE: GAM218 TARGET PROTEIN and GAM447 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 447D

[9029] Reference is now made to Fig. 656, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 656(GR656) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9030] GR656 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR656 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9031] GR656 GENE encodes GR656 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9032] GR656 PRECURSOR RNA folds spatially, forming GR656 FOLDED PRECURSOR RNA . It is appreciated that GR656 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR656 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[9033] GR656 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM47 PRECURSOR and GAM430 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9034] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM47 RNA and GAM430 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9035] GAM47 RNA binds complimentarily to a target binding site located in an untranslated region of GAM47 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 47A, thereby inhibiting translation of GAM47 TARGET RNA into GAM47 TARGET PROTEIN, both of Fig. 47A.

[9036] GAM430 RNA binds complimentarily to a target binding site located in an untranslated region of GAM430 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 430A, thereby inhibiting translation of GAM430 TARGET RNA into GAM430 TARGET PROTEIN, both of Fig. 430A.

[9037] It is appreciated that specific functions, and accordingly utilities, of GR656 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR656 GENE: GAM47 TARGET PROTEIN and GAM430 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 47D and 430D

[9038] Reference is now made to Fig. 657, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 657(GR657) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9039] GR657 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR657 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9040] GR657 GENE encodes GR657 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9041] GR657 PRECURSOR RNA folds spatially, forming GR657 FOLDED PRECURSOR RNA . It is appreciated that GR657 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR657 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9042] GR657 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM282 PRECURSOR and GAM290 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9043] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM282 RNA and GAM290 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9044] GAM282 RNA binds complimentarily to a target binding site located in an untranslated region of GAM282 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 282A, thereby inhibiting translation of GAM282 TARGET RNA into GAM282 TARGET PROTEIN, both of Fig. 282A.

[9045] GAM290 RNA binds complimentarily to a target binding site located in an untranslated region of GAM290 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 290A, thereby inhibiting translation of GAM290 TARGET RNA into GAM290 TARGET PROTEIN, both of Fig. 290A.

[9046] It is appreciated that specific functions, and accordingly utilities, of GR657 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR657 GENE: GAM282 TARGET PROTEIN and GAM290 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 282D and 290D

[9047] Reference is now made to Fig. 658, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 658(GR658) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9048] GR658 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR658 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9049] GR658 GENE encodes GR658 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9050] GR658 PRECURSOR RNA folds spatially, forming GR658 FOLDED PRECURSOR RNA . It is appreciated that GR658 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR658 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9051] GR658 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM80 PRECURSOR, GAM424 PRECURSOR and GAM505 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9052] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM80 RNA, GAM424 RNA and GAM505 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9053] GAM80 RNA binds complimentarily to a target binding site located in an untranslated region of GAM80 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 80A, thereby inhibiting translation of GAM80 TARGET RNA into GAM80 TARGET PROTEIN, both of Fig. 80A.

[9054] GAM424 RNA binds complimentarily to a target binding site located in an untranslated region of GAM424 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 424A, thereby inhibiting transla-

tion of GAM424 TARGET RNA into GAM424 TARGET PROTEIN, both of Fig. 424A.

[9055] GAM505 RNA binds complimentarily to a target binding site located in an untranslated region of GAM505 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 505A, thereby inhibiting translation of GAM505 TARGET RNA into GAM505 TARGET PROTEIN, both of Fig. 505A.

[9056] It is appreciated that specific functions, and accordingly utilities, of GR658 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR658 GENE: GAM80 TARGET PROTEIN, GAM424 TARGET PROTEIN and GAM505 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 80D, 424D and 505D

[9057] Reference is now made to Fig. 659, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 659(GR659) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the

function and utility of which at least one target gene is known in the art.

[9058] GR659 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR659 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9059] GR659 GENE encodes GR659 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9060] GR659 PRECURSOR RNA folds spatially, forming GR659 FOLDED PRECURSOR RNA . It is appreciated that GR659 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR659 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9061] GR659 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM218 PRECURSOR and GAM448 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECUR-

SOR RNA of Fig. 8.

[9062] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM218 RNA and GAM448 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9063] GAM218 RNA binds complimentarily to a target binding site located in an untranslated region of GAM218 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 218A, thereby inhibiting translation of GAM218 TARGET RNA into GAM218 TARGET PROTEIN, both of Fig. 218A.

[9064] GAM448 RNA binds complimentarily to a target binding site located in an untranslated region of GAM448 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 448A, thereby inhibiting translation of GAM448 TARGET RNA into GAM448 TARGET PROTEIN, both of Fig. 448A.

[9065] It is appreciated that specific functions, and accordingly utilities, of GR659 GENE correlate with, and may be deduced from, the identity of the target genes, which are in-

hibited by GAM RNAs comprised in the operon-like cluster of GR659 GENE: GAM218 TARGET PROTEIN and GAM448 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 218D and 448D

[9066] Reference is now made to Fig. 660, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 660(GR660) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9067] GR660 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR660 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9068] GR660 GENE encodes GR660 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9069] GR660 PRECURSOR RNA folds spatially, forming GR660 FOLDED PRECURSOR RNA . It is appreciated that GR660 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin

structures are due to the fact that the nucleotide sequence of GR660 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9070] GR660 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM420 PRECURSOR and GAM550 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9071] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM420 RNA and GAM550 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9072] GAM420 RNA binds complimentarily to a target binding site located in an untranslated region of GAM420 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 420A, thereby inhibiting translation of GAM420 TARGET RNA into GAM420 TARGET PRO-

TEIN, both of Fig. 420A.

[9073] GAM550 RNA binds complimentarily to a target binding site located in an untranslated region of GAM550 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 550A, thereby inhibiting translation of GAM550 TARGET RNA into GAM550 TARGET PROTEIN, both of Fig. 550A.

[9074] It is appreciated that specific functions, and accordingly utilities, of GR660 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR660 GENE: GAM420 TARGET PROTEIN and GAM550 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 420D and 550D

[9075] Reference is now made to Fig. 661, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 661(GR661) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is

known in the art.

[9076] GR661 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR661 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9077] GR661 GENE encodes GR661 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9078] GR661 PRECURSOR RNA folds spatially, forming GR661 FOLDED PRECURSOR RNA . It is appreciated that GR661 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR661 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9079] GR661 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM317 PRECURSOR and GAM501 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9080] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM317 RNA and GAM501 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9081] GAM317 RNA binds complimentarily to a target binding site located in an untranslated region of GAM317 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 317A, thereby inhibiting translation of GAM317 TARGET RNA into GAM317 TARGET PROTEIN, both of Fig. 317A.

[9082] GAM501 RNA binds complimentarily to a target binding site located in an untranslated region of GAM501 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 501A, thereby inhibiting translation of GAM501 TARGET RNA into GAM501 TARGET PROTEIN, both of Fig. 501A.

[9083] It is appreciated that specific functions, and accordingly utilities, of GR661 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR661 GENE: GAM317 TARGET PROTEIN and GAM501 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 317D and 501D

[9084] Reference is now made to Fig. 662, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 662(GR662) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9085] GR662 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR662 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9086] GR662 GENE encodes GR662 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9087] GR662 PRECURSOR RNA folds spatially, forming GR662 FOLDED PRECURSOR RNA . It is appreciated that GR662 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR662 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9088] GR662 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM111 PRECURSOR and GAM242 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9089] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM111 RNA and GAM242 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9090] GAM111 RNA binds complimentarily to a target binding site located in an untranslated region of GAM111 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 111A, thereby inhibiting translation of GAM111 TARGET RNA into GAM111 TARGET PROTEIN, both of Fig. 111A.

[9091] GAM242 RNA binds complimentarily to a target binding site located in an untranslated region of GAM242 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 242A, thereby inhibiting translation of GAM242 TARGET RNA into GAM242 TARGET PROTEIN, both of Fig. 242A.

[9092] It is appreciated that specific functions, and accordingly utilities, of GR662 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR662 GENE: GAM111 TARGET PROTEIN and GAM242 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 111D and 242D

[9093] Reference is now made to Fig. 663, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 663(GR663) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9094] GR663 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR663 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9095] GR663 GENE encodes GR663 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9096] GR663 PRECURSOR RNA folds spatially, forming GR663 FOLDED PRECURSOR RNA . It is appreciated that GR663 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR663 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9097] GR663 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM135 PRECURSOR and GAM309 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9098] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM135 RNA and GAM309 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9099] GAM135 RNA binds complimentarily to a target binding site located in an untranslated region of GAM135 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 135A, thereby inhibiting translation of GAM135 TARGET RNA into GAM135 TARGET PROTEIN, both of Fig. 135A.

[9100] GAM309 RNA binds complimentarily to a target binding site located in an untranslated region of GAM309 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 309A, thereby inhibiting translation of GAM309 TARGET RNA into GAM309 TARGET PROTEIN, both of Fig. 309A.

[9101] It is appreciated that specific functions, and accordingly utilities, of GR663 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR663 GENE: GAM135 TARGET PROTEIN and GAM309

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 135D and 309D

[9102] Reference is now made to Fig. 664, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 664(GR664) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9103] GR664 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR664 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9104] GR664 GENE encodes GR664 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9105] GR664 PRECURSOR RNA folds spatially, forming GR664 FOLDED PRECURSOR RNA . It is appreciated that GR664 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR664 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9106] GR664 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM119 PRECURSOR, GAM248 PRECURSOR and GAM257 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9107] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM119 RNA, GAM248 RNA and GAM257 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9108] GAM119 RNA binds complimentarily to a target binding site located in an untranslated region of GAM119 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 119A, thereby inhibiting translation of GAM119 TARGET RNA into GAM119 TARGET PROTEIN, both of Fig. 119A.

[9109] GAM248 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM248 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 248A, thereby inhibiting translation of GAM248 TARGET RNA into GAM248 TARGET PROTEIN, both of Fig. 248A.

[9110] GAM257 RNA binds complimentarily to a target binding site located in an untranslated region of GAM257 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 257A, thereby inhibiting translation of GAM257 TARGET RNA into GAM257 TARGET PROTEIN, both of Fig. 257A.

[9111] It is appreciated that specific functions, and accordingly utilities, of GR664 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR664 GENE: GAM119 TARGET PROTEIN, GAM248 TARGET PROTEIN and GAM257 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 119D, 248D and 257D

[9112] Reference is now made to Fig. 665, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 665(GR665) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9113] GR665 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR665 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9114] GR665 GENE encodes GR665 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9115] GR665 PRECURSOR RNA folds spatially, forming GR665 FOLDED PRECURSOR RNA . It is appreciated that GR665 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR665 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9116] GR665 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM115 PRECURSOR and GAM452 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9117] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM115 RNA and GAM452 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9118] GAM115 RNA binds complimentarily to a target binding site located in an untranslated region of GAM115 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 115A, thereby inhibiting translation of GAM115 TARGET RNA into GAM115 TARGET PROTEIN, both of Fig. 115A.

[9119] GAM452 RNA binds complimentarily to a target binding site located in an untranslated region of GAM452 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 452A, thereby inhibiting translation of GAM452 TARGET RNA into GAM452 TARGET PRO-

TEIN, both of Fig. 452A.

[9120] It is appreciated that specific functions, and accordingly utilities, of GR665 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR665 GENE: GAM115 TARGET PROTEIN and GAM452 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 115D and 452D

[9121] Reference is now made to Fig. 666, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 666(GR666) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9122] GR666 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR666 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9123] GR666 GENE encodes GR666 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9124] GR666 PRECURSOR RNA folds spatially, forming GR666 FOLDED PRECURSOR RNA . It is appreciated that GR666 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR666 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9125] GR666 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM156 PRECURSOR and GAM346 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9126] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM156 RNA and GAM346 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9127] GAM156 RNA binds complimentarily to a target binding site located in an untranslated region of GAM156 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 156A, thereby inhibiting translation of GAM156 TARGET RNA into GAM156 TARGET PROTEIN, both of Fig. 156A.

[9128] GAM346 RNA binds complimentarily to a target binding site located in an untranslated region of GAM346 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 346A, thereby inhibiting translation of GAM346 TARGET RNA into GAM346 TARGET PROTEIN, both of Fig. 346A.

[9129] It is appreciated that specific functions, and accordingly utilities, of GR666 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR666 GENE: GAM156 TARGET PROTEIN and GAM346 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 156D and 346D

[9130] Reference is now made to Fig. 667, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

667(GR667) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9131] GR667 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR667 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9132] GR667 GENE encodes GR667 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9133] GR667 PRECURSOR RNA folds spatially, forming GR667 FOLDED PRECURSOR RNA . It is appreciated that GR667 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR667 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9134] GR667 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM

precursor RNAs, GAM288 PRECURSOR and GAM405 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9135] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM288 RNA and GAM405 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9136] GAM288 RNA binds complimentarily to a target binding site located in an untranslated region of GAM288 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 288A, thereby inhibiting translation of GAM288 TARGET RNA into GAM288 TARGET PROTEIN, both of Fig. 288A.

[9137] GAM405 RNA binds complimentarily to a target binding site located in an untranslated region of GAM405 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 405A, thereby inhibiting translation of GAM405 TARGET RNA into GAM405 TARGET PROTEIN, both of Fig. 405A.

[9138] It is appreciated that specific functions, and accordingly utilities, of GR667 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR667 GENE: GAM288 TARGET PROTEIN and GAM405 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 288D and 405D

[9139] Reference is now made to Fig. 668, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 668(GR668) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9140] GR668 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR668 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9141] GR668 GENE encodes GR668 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9142] GR668 PRECURSOR RNA folds spatially, forming GR668

FOLDED PRECURSOR RNA . It is appreciated that GR668 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR668 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9143] GR668 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM259 PRECURSOR and GAM379 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9144] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM259 RNA and GAM379 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9145] GAM259 RNA binds complimentarily to a target binding site located in an untranslated region of GAM259 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 259A, thereby inhibiting translation of GAM259 TARGET RNA into GAM259 TARGET PROTEIN, both of Fig. 259A.

[9146] GAM379 RNA binds complimentarily to a target binding site located in an untranslated region of GAM379 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 379A, thereby inhibiting translation of GAM379 TARGET RNA into GAM379 TARGET PROTEIN, both of Fig. 379A.

[9147] It is appreciated that specific functions, and accordingly utilities, of GR668 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR668 GENE: GAM259 TARGET PROTEIN and GAM379 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 259D and 379D

[9148] Reference is now made to Fig. 669, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 669(GR669) gene, which encodes an operon-like cluster

of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9149] GR669 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR669 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9150] GR669 GENE encodes GR669 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9151] GR669 PRECURSOR RNA folds spatially, forming GR669 FOLDED PRECURSOR RNA . It is appreciated that GR669 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR669 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9152] GR669 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM68 PRECURSOR and GAM365 PRE-

CURSOR, each of which GAM precursor RNAs being a hair-pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9153] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM68 RNA and GAM365 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9154] GAM68 RNA binds complimentarily to a target binding site located in an untranslated region of GAM68 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 68A, thereby inhibiting translation of GAM68 TARGET RNA into GAM68 TARGET PROTEIN, both of Fig. 68A.

[9155] GAM365 RNA binds complimentarily to a target binding site located in an untranslated region of GAM365 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 365A, thereby inhibiting translation of GAM365 TARGET RNA into GAM365 TARGET PROTEIN, both of Fig. 365A.

[9156] It is appreciated that specific functions, and accordingly

utilities, of GR669 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR669 GENE: GAM68 TARGET PROTEIN and GAM365 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 68D and 365D

[9157] Reference is now made to Fig. 670, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 670(GR670) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9158] GR670 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR670 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9159] GR670 GENE encodes GR670 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9160] GR670 PRECURSOR RNA folds spatially, forming GR670 FOLDED PRECURSOR RNA . It is appreciated that GR670

FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR670 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9161] GR670 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM97 PRECURSOR and GAM316 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9162] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM97 RNA and GAM316 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9163] GAM97 RNA binds complimentarily to a target binding site located in an untranslated region of GAM97 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING

SITE III of Fig. 97A, thereby inhibiting translation of GAM97 TARGET RNA into GAM97 TARGET PROTEIN, both of Fig. 97A.

[9164] GAM316 RNA binds complimentarily to a target binding site located in an untranslated region of GAM316 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 316A, thereby inhibiting translation of GAM316 TARGET RNA into GAM316 TARGET PROTEIN, both of Fig. 316A.

[9165] It is appreciated that specific functions, and accordingly utilities, of GR670 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR670 GENE: GAM97 TARGET PROTEIN and GAM316 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 97D and 316D

[9166] Reference is now made to Fig. 671, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 671(GR671) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn

modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9167] GR671 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR671 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9168] GR671 GENE encodes GR671 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9169] GR671 PRECURSOR RNA folds spatially, forming GR671 FOLDED PRECURSOR RNA . It is appreciated that GR671 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR671 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9170] GR671 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM214 PRECURSOR and GAM450 PRECURSOR, each of which GAM precursor RNAs being a hair-

pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9171] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM214 RNA and GAM450 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9172] GAM214 RNA binds complimentarily to a target binding site located in an untranslated region of GAM214 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 214A, thereby inhibiting translation of GAM214 TARGET RNA into GAM214 TARGET PROTEIN, both of Fig. 214A.

[9173] GAM450 RNA binds complimentarily to a target binding site located in an untranslated region of GAM450 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 450A, thereby inhibiting translation of GAM450 TARGET RNA into GAM450 TARGET PROTEIN, both of Fig. 450A.

[9174] It is appreciated that specific functions, and accordingly utilities, of GR671 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR671 GENE: GAM214 TARGET PROTEIN and GAM450 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 214D and 450D

[9175] Reference is now made to Fig. 672, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 672(GR672) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9176] GR672 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR672 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9177] GR672 GENE encodes GR672 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9178] GR672 PRECURSOR RNA folds spatially, forming GR672 FOLDED PRECURSOR RNA . It is appreciated that GR672 FOLDED PRECURSOR RNA comprises a plurality of what is

known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR672 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9179] GR672 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM334 PRECURSOR and GAM390 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9180] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM334 RNA and GAM390 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9181] GAM334 RNA binds complimentarily to a target binding site located in an untranslated region of GAM334 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 334A, thereby inhibiting transla-

tion of GAM334 TARGET RNA into GAM334 TARGET PROTEIN, both of Fig. 334A.

[9182] GAM390 RNA binds complimentarily to a target binding site located in an untranslated region of GAM390 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 390A, thereby inhibiting translation of GAM390 TARGET RNA into GAM390 TARGET PROTEIN, both of Fig. 390A.

[9183] It is appreciated that specific functions, and accordingly utilities, of GR672 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR672 GENE: GAM334 TARGET PROTEIN and GAM390 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 334D and 390D

[9184] Reference is now made to Fig. 673, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 673(GR673) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the

function and utility of which at least one target gene is known in the art.

[9185] GR673 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR673 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9186] GR673 GENE encodes GR673 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9187] GR673 PRECURSOR RNA folds spatially, forming GR673 FOLDED PRECURSOR RNA . It is appreciated that GR673 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR673 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9188] GR673 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM275 PRECURSOR and GAM415 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9189] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM275 RNA and GAM415 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9190] GAM275 RNA binds complimentarily to a target binding site located in an untranslated region of GAM275 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 275A, thereby inhibiting translation of GAM275 TARGET RNA into GAM275 TARGET PROTEIN, both of Fig. 275A.

[9191] GAM415 RNA binds complimentarily to a target binding site located in an untranslated region of GAM415 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 415A, thereby inhibiting translation of GAM415 TARGET RNA into GAM415 TARGET PROTEIN, both of Fig. 415A.

[9192] It is appreciated that specific functions, and accordingly utilities, of GR673 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR673 GENE: GAM275 TARGET PROTEIN and GAM415 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 275D and 415D

[9193] Reference is now made to Fig. 674, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record

674(GR674) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9194] GR674 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR674 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9195] GR674 GENE encodes GR674 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9196] GR674 PRECURSOR RNA folds spatially, forming GR674 FOLDED PRECURSOR RNA . It is appreciated that GR674 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR674 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9197] GR674 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM

precursor RNAs, GAM44 PRECURSOR, GAM403 PRECURSOR and GAM520 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9198] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM44 RNA, GAM403 RNA and GAM520 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9199] GAM44 RNA binds complimentarily to a target binding site located in an untranslated region of GAM44 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 44A, thereby inhibiting translation of GAM44 TARGET RNA into GAM44 TARGET PROTEIN, both of Fig. 44A.

[9200] GAM403 RNA binds complimentarily to a target binding site located in an untranslated region of GAM403 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 403A, thereby inhibiting translation of GAM403 TARGET RNA into GAM403 TARGET PROTEIN, both of Fig. 403A.

[9201] GAM520 RNA binds complimentarily to a target binding site located in an untranslated region of GAM520 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 520A, thereby inhibiting translation of GAM520 TARGET RNA into GAM520 TARGET PROTEIN, both of Fig. 520A.

[9202] It is appreciated that specific functions, and accordingly utilities, of GR674 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR674 GENE: GAM44 TARGET PROTEIN, GAM403 TARGET PROTEIN and GAM520 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 44D, 403D and 520D

[9203] Reference is now made to Fig. 675, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 675(GR675) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9204] GR675 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR675 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9205] GR675 GENE encodes GR675 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9206] GR675 PRECURSOR RNA folds spatially, forming GR675 FOLDED PRECURSOR RNA . It is appreciated that GR675 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR675 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9207] GR675 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM95 PRECURSOR and GAM261 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9208] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM95 RNA and GAM261 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9209] GAM95 RNA binds complimentarily to a target binding site located in an untranslated region of GAM95 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 95A, thereby inhibiting translation of GAM95 TARGET RNA into GAM95 TARGET PROTEIN, both of Fig. 95A.

[9210] GAM261 RNA binds complimentarily to a target binding site located in an untranslated region of GAM261 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 261A, thereby inhibiting translation of GAM261 TARGET RNA into GAM261 TARGET PROTEIN, both of Fig. 261A.

[9211] It is appreciated that specific functions, and accordingly utilities, of GR675 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR675 GENE: GAM95 TARGET PROTEIN and GAM261

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 95D and 261D

[9212] Reference is now made to Fig. 676, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 676(GR676) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9213] GR676 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR676 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9214] GR676 GENE encodes GR676 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9215] GR676 PRECURSOR RNA folds spatially, forming GR676 FOLDED PRECURSOR RNA . It is appreciated that GR676 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR676 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9216] GR676 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM229 PRECURSOR and GAM370 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9217] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM229 RNA and GAM370 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9218] GAM229 RNA binds complimentarily to a target binding site located in an untranslated region of GAM229 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 229A, thereby inhibiting translation of GAM229 TARGET RNA into GAM229 TARGET PROTEIN, both of Fig. 229A.

[9219] GAM370 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM370 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 370A, thereby inhibiting translation of GAM370 TARGET RNA into GAM370 TARGET PROTEIN, both of Fig. 370A.

[9220] It is appreciated that specific functions, and accordingly utilities, of GR676 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR676 GENE: GAM229 TARGET PROTEIN and GAM370 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 229D and 370D

[9221] Reference is now made to Fig. 677, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 677(GR677) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9222] GR677 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR677 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9223] GR677 GENE encodes GR677 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9224] GR677 PRECURSOR RNA folds spatially, forming GR677 FOLDED PRECURSOR RNA . It is appreciated that GR677 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR677 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9225] GR677 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM158 PRECURSOR and GAM466 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9226] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM158 RNA and GAM466 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9227] GAM158 RNA binds complimentarily to a target binding site located in an untranslated region of GAM158 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 158A, thereby inhibiting translation of GAM158 TARGET RNA into GAM158 TARGET PROTEIN, both of Fig. 158A.

[9228] GAM466 RNA binds complimentarily to a target binding site located in an untranslated region of GAM466 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 466A, thereby inhibiting translation of GAM466 TARGET RNA into GAM466 TARGET PROTEIN, both of Fig. 466A.

[9229] It is appreciated that specific functions, and accordingly utilities, of GR677 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR677 GENE: GAM158 TARGET PROTEIN and GAM466 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 158D and 466D

[9230] Reference is now made to Fig. 678, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 678(GR678) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9231] GR678 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR678 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9232] GR678 GENE encodes GR678 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9233] GR678 PRECURSOR RNA folds spatially, forming GR678 FOLDED PRECURSOR RNA . It is appreciated that GR678 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR678 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9234] GR678 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM350 PRECURSOR and GAM470 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9235] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM350 RNA and GAM470 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9236] GAM350 RNA binds complimentarily to a target binding site located in an untranslated region of GAM350 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 350A, thereby inhibiting translation of GAM350 TARGET RNA into GAM350 TARGET PROTEIN, both of Fig. 350A.

[9237] GAM470 RNA binds complimentarily to a target binding site located in an untranslated region of GAM470 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 470A, thereby inhibiting translation of GAM470 TARGET RNA into GAM470 TARGET PROTEIN, both of Fig. 470A.

[9238] It is appreciated that specific functions, and accordingly utilities, of GR678 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR678 GENE: GAM350 TARGET PROTEIN and GAM470 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 350D and 470D

[9239] Reference is now made to Fig. 679, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 679(GR679) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9240] GR679 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR679 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9241] GR679 GENE encodes GR679 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9242] GR679 PRECURSOR RNA folds spatially, forming GR679 FOLDED PRECURSOR RNA . It is appreciated that GR679 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR679 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9243] GR679 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM143 PRECURSOR and GAM398 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9244] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM143 RNA

and GAM398 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9245] GAM143 RNA binds complimentarily to a target binding site located in an untranslated region of GAM143 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 143A, thereby inhibiting translation of GAM143 TARGET RNA into GAM143 TARGET PROTEIN, both of Fig. 143A.

[9246] GAM398 RNA binds complimentarily to a target binding site located in an untranslated region of GAM398 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 398A, thereby inhibiting translation of GAM398 TARGET RNA into GAM398 TARGET PROTEIN, both of Fig. 398A.

[9247] It is appreciated that specific functions, and accordingly utilities, of GR679 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR679 GENE: GAM143 TARGET PROTEIN and GAM398 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 143D and

[9248] Reference is now made to Fig. 680, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 680(GR680) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9249] GR680 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR680 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9250] GR680 GENE encodes GR680 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9251] GR680 PRECURSOR RNA folds spatially, forming GR680 FOLDED PRECURSOR RNA . It is appreciated that GR680 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR680 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-re-

versed sequence of the second half thereof, as is well known in the art.

[9252] GR680 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM526 PRECURSOR and GAM541 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9253] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM526 RNA and GAM541 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9254] GAM526 RNA binds complimentarily to a target binding site located in an untranslated region of GAM526 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 526A, thereby inhibiting translation of GAM526 TARGET RNA into GAM526 TARGET PROTEIN, both of Fig. 526A.

[9255] GAM541 RNA binds complimentarily to a target binding site located in an untranslated region of GAM541 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 541A, thereby inhibiting translation of GAM541 TARGET RNA into GAM541 TARGET PROTEIN, both of Fig. 541A.

[9256] It is appreciated that specific functions, and accordingly utilities, of GR680 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR680 GENE: GAM526 TARGET PROTEIN and GAM541 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 526D and 541D

[9257] Reference is now made to Fig. 681, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 681(GR681) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9258] GR681 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR681 GENE was detected is described hereinabove with

reference to Figs. 6-15.

[9259] GR681 GENE encodes GR681 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9260] GR681 PRECURSOR RNA folds spatially, forming GR681 FOLDED PRECURSOR RNA . It is appreciated that GR681 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR681 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9261] GR681 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM107 PRECURSOR and GAM264 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9262] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM107 RNA and GAM264 RNA respectively, each of which GAM RNAs

corresponding to GAM RNA of Fig. 8.

[9263] GAM107 RNA binds complimentarily to a target binding site located in an untranslated region of GAM107 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 107A, thereby inhibiting translation of GAM107 TARGET RNA into GAM107 TARGET PROTEIN, both of Fig. 107A.

[9264] GAM264 RNA binds complimentarily to a target binding site located in an untranslated region of GAM264 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 264A, thereby inhibiting translation of GAM264 TARGET RNA into GAM264 TARGET PROTEIN, both of Fig. 264A.

[9265] It is appreciated that specific functions, and accordingly utilities, of GR681 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR681 GENE: GAM107 TARGET PROTEIN and GAM264 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 107D and 264D

[9266] Reference is now made to Fig. 682, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 682(GR682) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9267] GR682 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR682 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9268] GR682 GENE encodes GR682 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9269] GR682 PRECURSOR RNA folds spatially, forming GR682 FOLDED PRECURSOR RNA . It is appreciated that GR682 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR682 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well

known in the art.

[9270] GR682 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM431 PRECURSOR and GAM532 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9271] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM431 RNA and GAM532 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9272] GAM431 RNA binds complimentarily to a target binding site located in an untranslated region of GAM431 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 431A, thereby inhibiting translation of GAM431 TARGET RNA into GAM431 TARGET PROTEIN, both of Fig. 431A.

[9273] GAM532 RNA binds complimentarily to a target binding site located in an untranslated region of GAM532 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 532A, thereby inhibiting translation of GAM532 TARGET RNA into GAM532 TARGET PROTEIN, both of Fig. 532A.

[9274] It is appreciated that specific functions, and accordingly utilities, of GR682 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR682 GENE: GAM431 TARGET PROTEIN and GAM532 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 431D and 532D

[9275] Reference is now made to Fig. 683, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 683(GR683) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9276] GR683 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR683 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9277] GR683 GENE encodes GR683 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9278] GR683 PRECURSOR RNA folds spatially, forming GR683 FOLDED PRECURSOR RNA . It is appreciated that GR683 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR683 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9279] GR683 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM332 PRECURSOR and GAM432 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9280] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM332 RNA and GAM432 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9281] GAM332 RNA binds complimentarily to a target binding site located in an untranslated region of GAM332 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 332A, thereby inhibiting translation of GAM332 TARGET RNA into GAM332 TARGET PROTEIN, both of Fig. 332A.

[9282] GAM432 RNA binds complimentarily to a target binding site located in an untranslated region of GAM432 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 432A, thereby inhibiting translation of GAM432 TARGET RNA into GAM432 TARGET PROTEIN, both of Fig. 432A.

[9283] It is appreciated that specific functions, and accordingly utilities, of GR683 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR683 GENE: GAM332 TARGET PROTEIN and GAM432 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 332D and 432D

[9284] Reference is now made to Fig. 684, which is a simplified

diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 684(GR684) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9285] GR684 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR684 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9286] GR684 GENE encodes GR684 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9287] GR684 PRECURSOR RNA folds spatially, forming GR684 FOLDED PRECURSOR RNA . It is appreciated that GR684 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR684 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9288] GR684 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM352 PRECURSOR and GAM488 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9289] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM352 RNA and GAM488 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9290] GAM352 RNA binds complimentarily to a target binding site located in an untranslated region of GAM352 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 352A, thereby inhibiting translation of GAM352 TARGET RNA into GAM352 TARGET PROTEIN, both of Fig. 352A.

[9291] GAM488 RNA binds complimentarily to a target binding site located in an untranslated region of GAM488 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 488A, thereby inhibiting transla-

tion of GAM488 TARGET RNA into GAM488 TARGET PROTEIN, both of Fig. 488A.

[9292] It is appreciated that specific functions, and accordingly utilities, of GR684 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR684 GENE: GAM352 TARGET PROTEIN and GAM488 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 352D and 488D

[9293] Reference is now made to Fig. 685, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 685(GR685) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9294] GR685 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR685 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9295] GR685 GENE encodes GR685 PRECURSOR RNA, an RNA

molecule, typically several hundred nucleotides long.

[9296] GR685 PRECURSOR RNA folds spatially, forming GR685 FOLDED PRECURSOR RNA . It is appreciated that GR685 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR685 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9297] GR685 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM310 PRECURSOR and GAM507 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9298] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM310 RNA and GAM507 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9299] GAM310 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM310 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 310A, thereby inhibiting translation of GAM310 TARGET RNA into GAM310 TARGET PROTEIN, both of Fig. 310A.

[9300] GAM507 RNA binds complimentarily to a target binding site located in an untranslated region of GAM507 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 507A, thereby inhibiting translation of GAM507 TARGET RNA into GAM507 TARGET PROTEIN, both of Fig. 507A.

[9301] It is appreciated that specific functions, and accordingly utilities, of GR685 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR685 GENE: GAM310 TARGET PROTEIN and GAM507 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 310D and 507D

[9302] Reference is now made to Fig. 686, which is a simplified diagram describing a novel bioinformatically detected

regulatory gene, referred to here as Genomic Record 686(GR686) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9303] GR686 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR686 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9304] GR686 GENE encodes GR686 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9305] GR686 PRECURSOR RNA folds spatially, forming GR686 FOLDED PRECURSOR RNA . It is appreciated that GR686 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR686 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9306] GR686 FOLDED PRECURSOR RNA is naturally processed by

cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM171 PRECURSOR, GAM347 PRECURSOR and GAM457 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9307] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM171 RNA, GAM347 RNA and GAM457 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9308] GAM171 RNA binds complimentarily to a target binding site located in an untranslated region of GAM171 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 171A, thereby inhibiting translation of GAM171 TARGET RNA into GAM171 TARGET PROTEIN, both of Fig. 171A.

[9309] GAM347 RNA binds complimentarily to a target binding site located in an untranslated region of GAM347 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 347A, thereby inhibiting translation of GAM347 TARGET RNA into GAM347 TARGET PRO-

TEIN, both of Fig. 347A.

[9310] GAM457 RNA binds complimentarily to a target binding site located in an untranslated region of GAM457 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 457A, thereby inhibiting translation of GAM457 TARGET RNA into GAM457 TARGET PROTEIN, both of Fig. 457A.

[9311] It is appreciated that specific functions, and accordingly utilities, of GR686 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR686 GENE: GAM171 TARGET PROTEIN, GAM347 TARGET PROTEIN and GAM457 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 171D, 347D and 457D

[9312] Reference is now made to Fig. 687, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 687(GR687) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is

known in the art.

[9313] GR687 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR687 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9314] GR687 GENE encodes GR687 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9315] GR687 PRECURSOR RNA folds spatially, forming GR687 FOLDED PRECURSOR RNA . It is appreciated that GR687 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR687 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9316] GR687 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM96 PRECURSOR and GAM277 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9317] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM96 RNA and GAM277 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9318] GAM96 RNA binds complimentarily to a target binding site located in an untranslated region of GAM96 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 96A, thereby inhibiting translation of GAM96 TARGET RNA into GAM96 TARGET PROTEIN, both of Fig. 96A.

[9319] GAM277 RNA binds complimentarily to a target binding site located in an untranslated region of GAM277 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 277A, thereby inhibiting translation of GAM277 TARGET RNA into GAM277 TARGET PROTEIN, both of Fig. 277A.

[9320] It is appreciated that specific functions, and accordingly utilities, of GR687 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster

of GR687 GENE: GAM96 TARGET PROTEIN and GAM277 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 96D and 277D

[9321] Reference is now made to Fig. 688, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 688(GR688) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9322] GR688 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR688 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9323] GR688 GENE encodes GR688 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9324] GR688 PRECURSOR RNA folds spatially, forming GR688 FOLDED PRECURSOR RNA . It is appreciated that GR688 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence

of GR688 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9325] GR688 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM86 PRECURSOR and GAM202 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9326] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM86 RNA and GAM202 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9327] GAM86 RNA binds complimentarily to a target binding site located in an untranslated region of GAM86 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 86A, thereby inhibiting translation of GAM86 TARGET RNA into GAM86 TARGET PROTEIN, both of Fig. 86A.

[9328] GAM202 RNA binds complimentarily to a target binding site located in an untranslated region of GAM202 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 202A, thereby inhibiting translation of GAM202 TARGET RNA into GAM202 TARGET PROTEIN, both of Fig. 202A.

[9329] It is appreciated that specific functions, and accordingly utilities, of GR688 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR688 GENE: GAM86 TARGET PROTEIN and GAM202 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 86D and 202D

[9330] Reference is now made to Fig. 689, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 689(GR689) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9331] GR689 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR689 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9332] GR689 GENE encodes GR689 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9333] GR689 PRECURSOR RNA folds spatially, forming GR689 FOLDED PRECURSOR RNA . It is appreciated that GR689 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR689 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9334] GR689 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM29 PRECURSOR and GAM180 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9335] The above mentioned GAM precursor RNAs are diced by

DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM29 RNA and GAM180 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9336] GAM29 RNA binds complimentarily to a target binding site located in an untranslated region of GAM29 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 29A, thereby inhibiting translation of GAM29 TARGET RNA into GAM29 TARGET PROTEIN, both of Fig. 29A.

[9337] GAM180 RNA binds complimentarily to a target binding site located in an untranslated region of GAM180 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 180A, thereby inhibiting translation of GAM180 TARGET RNA into GAM180 TARGET PROTEIN, both of Fig. 180A.

[9338] It is appreciated that specific functions, and accordingly utilities, of GR689 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR689 GENE: GAM29 TARGET PROTEIN and GAM180

TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 29D and 180D

[9339] Reference is now made to Fig. 690, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 690(GR690) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9340] GR690 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR690 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9341] GR690 GENE encodes GR690 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9342] GR690 PRECURSOR RNA folds spatially, forming GR690 FOLDED PRECURSOR RNA . It is appreciated that GR690 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR690 PRECURSOR RNA comprises a plurality of seg-

ments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9343] GR690 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM47 PRECURSOR and GAM430 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9344] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM47 RNA and GAM430 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9345] GAM47 RNA binds complimentarily to a target binding site located in an untranslated region of GAM47 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 47A, thereby inhibiting translation of GAM47 TARGET RNA into GAM47 TARGET PROTEIN, both of Fig. 47A.

[9346] GAM430 RNA binds complimentarily to a target binding

site located in an untranslated region of GAM430 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 430A, thereby inhibiting translation of GAM430 TARGET RNA into GAM430 TARGET PROTEIN, both of Fig. 430A.

[9347] It is appreciated that specific functions, and accordingly utilities, of GR690 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR690 GENE: GAM47 TARGET PROTEIN and GAM430 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 47D and 430D

[9348] Reference is now made to Fig. 691, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 691(GR691) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9349] GR691 GENE is a novel bioinformatically detected regula-

tory, non protein coding, RNA gene. The method by which GR691 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9350] GR691 GENE encodes GR691 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9351] GR691 PRECURSOR RNA folds spatially, forming GR691 FOLDED PRECURSOR RNA . It is appreciated that GR691 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR691 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9352] GR691 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM281 PRECURSOR and GAM517 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9353] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA

segments of about 22 nucleotides in length, GAM281 RNA and GAM517 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9354] GAM281 RNA binds complimentarily to a target binding site located in an untranslated region of GAM281 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 281A, thereby inhibiting translation of GAM281 TARGET RNA into GAM281 TARGET PROTEIN, both of Fig. 281A.

[9355] GAM517 RNA binds complimentarily to a target binding site located in an untranslated region of GAM517 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 517A, thereby inhibiting translation of GAM517 TARGET RNA into GAM517 TARGET PROTEIN, both of Fig. 517A.

[9356] It is appreciated that specific functions, and accordingly utilities, of GR691 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR691 GENE: GAM281 TARGET PROTEIN and GAM517 TARGET PROTEIN. The function of these target genes is

elaborated hereinabove with reference to Figs. 281D and 517D

[9357] Reference is now made to Fig. 692, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 692(GR692) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9358] GR692 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR692 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9359] GR692 GENE encodes GR692 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9360] GR692 PRECURSOR RNA folds spatially, forming GR692 FOLDED PRECURSOR RNA . It is appreciated that GR692 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR692 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nu-

cleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9361] GR692 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM98 PRECURSOR and GAM455 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9362] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM98 RNA and GAM455 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9363] GAM98 RNA binds complimentarily to a target binding site located in an untranslated region of GAM98 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 98A, thereby inhibiting translation of GAM98 TARGET RNA into GAM98 TARGET PROTEIN, both of Fig. 98A.

[9364] GAM455 RNA binds complimentarily to a target binding site located in an untranslated region of GAM455 TARGET

RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 455A, thereby inhibiting translation of GAM455 TARGET RNA into GAM455 TARGET PROTEIN, both of Fig. 455A.

[9365] It is appreciated that specific functions, and accordingly utilities, of GR692 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR692 GENE: GAM98 TARGET PROTEIN and GAM455 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 98D and 455D

[9366] Reference is now made to Fig. 693, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 693(GR693) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9367] GR693 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which

GR693 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9368] GR693 GENE encodes GR693 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9369] GR693 PRECURSOR RNA folds spatially, forming GR693 FOLDED PRECURSOR RNA . It is appreciated that GR693 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR693 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9370] GR693 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 3 separate GAM precursor RNAs, GAM74 PRECURSOR, GAM200 PRECURSOR and GAM243 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9371] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM74 RNA,

GAM200 RNA and GAM243 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9372] GAM74 RNA binds complimentarily to a target binding site located in an untranslated region of GAM74 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 74A, thereby inhibiting translation of GAM74 TARGET RNA into GAM74 TARGET PROTEIN, both of Fig. 74A.

[9373] GAM200 RNA binds complimentarily to a target binding site located in an untranslated region of GAM200 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 200A, thereby inhibiting translation of GAM200 TARGET RNA into GAM200 TARGET PROTEIN, both of Fig. 200A.

[9374] GAM243 RNA binds complimentarily to a target binding site located in an untranslated region of GAM243 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 243A, thereby inhibiting translation of GAM243 TARGET RNA into GAM243 TARGET PROTEIN, both of Fig. 243A.

[9375] It is appreciated that specific functions, and accordingly utilities, of GR693 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR693 GENE: GAM74 TARGET PROTEIN, GAM200 TARGET PROTEIN and GAM243 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 74D, 200D and 243D

[9376] Reference is now made to Fig. 694, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 694(GR694) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9377] GR694 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR694 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9378] GR694 GENE encodes GR694 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9379] GR694 PRECURSOR RNA folds spatially, forming GR694

FOLDED PRECURSOR RNA . It is appreciated that GR694 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR694 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9380] GR694 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM129 PRECURSOR and GAM154 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9381] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM129 RNA and GAM154 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9382] GAM129 RNA binds complimentarily to a target binding site located in an untranslated region of GAM129 TARGET RNA, which target binding site corresponds to a target

binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 129A, thereby inhibiting translation of GAM129 TARGET RNA into GAM129 TARGET PROTEIN, both of Fig. 129A.

[9383] GAM154 RNA binds complimentarily to a target binding site located in an untranslated region of GAM154 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 154A, thereby inhibiting translation of GAM154 TARGET RNA into GAM154 TARGET PROTEIN, both of Fig. 154A.

[9384] It is appreciated that specific functions, and accordingly utilities, of GR694 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR694 GENE: GAM129 TARGET PROTEIN and GAM154 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 129D and 154D

[9385] Reference is now made to Fig. 695, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 695(GR695) gene, which encodes an operon-like cluster

of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9386] GR695 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR695 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9387] GR695 GENE encodes GR695 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9388] GR695 PRECURSOR RNA folds spatially, forming GR695 FOLDED PRECURSOR RNA . It is appreciated that GR695 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR695 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9389] GR695 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM336 PRECURSOR and GAM549 PRE-

CURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9390] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM336 RNA and GAM549 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9391] GAM336 RNA binds complimentarily to a target binding site located in an untranslated region of GAM336 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 336A, thereby inhibiting translation of GAM336 TARGET RNA into GAM336 TARGET PROTEIN, both of Fig. 336A.

[9392] GAM549 RNA binds complimentarily to a target binding site located in an untranslated region of GAM549 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 549A, thereby inhibiting translation of GAM549 TARGET RNA into GAM549 TARGET PROTEIN, both of Fig. 549A.

[9393] It is appreciated that specific functions, and accordingly

utilities, of GR695 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR695 GENE: GAM336 TARGET PROTEIN and GAM549 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 336D and 549D

[9394] Reference is now made to Fig. 696, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 696(GR696) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9395] GR696 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR696 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9396] GR696 GENE encodes GR696 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9397] GR696 PRECURSOR RNA folds spatially, forming GR696 FOLDED PRECURSOR RNA . It is appreciated that GR696

FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR696 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9398] GR696 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM426 PRECURSOR and GAM496 PRECURSOR, each of which GAM precursor RNAs being a hairpin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9399] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM426 RNA and GAM496 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9400] GAM426 RNA binds complimentarily to a target binding site located in an untranslated region of GAM426 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or

BINDING SITE III of Fig. 426A, thereby inhibiting translation of GAM426 TARGET RNA into GAM426 TARGET PROTEIN, both of Fig. 426A.

[9401] GAM496 RNA binds complimentarily to a target binding site located in an untranslated region of GAM496 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 496A, thereby inhibiting translation of GAM496 TARGET RNA into GAM496 TARGET PROTEIN, both of Fig. 496A.

[9402] It is appreciated that specific functions, and accordingly utilities, of GR696 GENE correlate with, and may be deduced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR696 GENE: GAM426 TARGET PROTEIN and GAM496 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 426D and 496D

[9403] Reference is now made to Fig. 697, which is a simplified diagram describing a novel bioinformatically detected regulatory gene, referred to here as Genomic Record 697(GR697) gene, which encodes an operon-like cluster of novel micro RNA-like genes, each of which in turn

modulates expression of at least one target gene, the function and utility of which at least one target gene is known in the art.

[9404] GR697 GENE is a novel bioinformatically detected regulatory, non protein coding, RNA gene. The method by which GR697 GENE was detected is described hereinabove with reference to Figs. 6-15.

[9405] GR697 GENE encodes GR697 PRECURSOR RNA, an RNA molecule, typically several hundred nucleotides long.

[9406] GR697 PRECURSOR RNA folds spatially, forming GR697 FOLDED PRECURSOR RNA . It is appreciated that GR697 FOLDED PRECURSOR RNA comprises a plurality of what is known in the art as hairpin structures. These hairpin structures are due to the fact that the nucleotide sequence of GR697 PRECURSOR RNA comprises a plurality of segments, the first half of each such segment having a nucleotide sequence which is at least a partial inversed-reversed sequence of the second half thereof, as is well known in the art.

[9407] GR697 FOLDED PRECURSOR RNA is naturally processed by cellular enzymatic activity into at least 2 separate GAM precursor RNAs, GAM201 PRECURSOR and GAM325 PRECURSOR, each of which GAM precursor RNAs being a hair-

pin shaped RNA segment, corresponding to GAM PRECURSOR RNA of Fig. 8.

[9408] The above mentioned GAM precursor RNAs are diced by DICER COMPLEX of Fig. 8, yielding respective short RNA segments of about 22 nucleotides in length, GAM201 RNA and GAM325 RNA respectively, each of which GAM RNAs corresponding to GAM RNA of Fig. 8.

[9409] GAM201 RNA binds complimentarily to a target binding site located in an untranslated region of GAM201 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 201A, thereby inhibiting translation of GAM201 TARGET RNA into GAM201 TARGET PROTEIN, both of Fig. 201A.

[9410] GAM325 RNA binds complimentarily to a target binding site located in an untranslated region of GAM325 TARGET RNA, which target binding site corresponds to a target binding site such as BINDING SITE I, BINDING SITE II or BINDING SITE III of Fig. 325A, thereby inhibiting translation of GAM325 TARGET RNA into GAM325 TARGET PROTEIN, both of Fig. 325A.

[9411] It is appreciated that specific functions, and accordingly utilities, of GR697 GENE correlate with, and may be de-

duced from, the identity of the target genes, which are inhibited by GAM RNAs comprised in the operon-like cluster of GR697 GENE: GAM201 TARGET PROTEIN and GAM325 TARGET PROTEIN. The function of these target genes is elaborated hereinabove with reference to Figs. 201D and 325D

[9412]

[9413] BIBLIOGRAPHY

[9414] 1.

[9415] Mercken, M.; Takahashi, H.; Honda, T.; Sato, K.; Murayama, M.;

[9416] Nakazato, Y.; Noguchi, K.; Imahori, K.; Takashima, A.: Characterization

[9417] of human presenilin 1 using N-terminal specific monoclonal antibodies:

[9418] evidence that Alzheimer mutations affect proteolytic processing. FEBS

[9419] Lett. 389: 297-303, 1996.

[9420] 2.

[9421] Davies, A. F.; Mirza, G.; Flinter, F.; Ragoussis, J.: An interstitial

[9422] deletion of 6p24-p25 proximal to the FKHL7 locus and including AP-2-alpha
[9423] that affects anterior eye chamber development. J. Med. Genet. 36:
[9424] 708-710, 1999.

[9425] 3.

[9426] Sen Gupta, B.; Friedberg, F.; Detera-Wadleigh, S. D.: Molecular
[9427] analysis of human and rat calmodulin complementary DNA clones: evidence
[9428] for additional active genes in these species. J. Biol. Chem. 262:
[9429] 16663-16670, 1987.

[9430] 4.

[9431] Fishman, G. I.; Moreno, A. P.; Spray, D. C.; Leinwand, L. A.:

[9432] Functional analysis of human cardiac gap junction channel mutants. Proc.
[9433] Nat. Acad. Sci. 88: 3525-3529, 1991.

[9434] 5.

[9435] Hassett, C.; Aicher, L.; Sidhu, J. S.; Omiecinski, C. J.: Human

[9436] microsomal epoxide hydrolase: genetic polymorphism and functional

[9437] expression in vitro of amino acid variants. Hum. Molec. Genet. 3:

[9438] 421-428, 1994.

[9439] 6.

[9440] Henry, I.; Uzan, G.; Weil, D.; Nicolas, H.; Kaplan, J. C.; Marguerie,

[9441] C.; Kahn, A.; Junien, C.: The genes coding for A-alpha-, B-beta-,

[9442] and gamma-chains of fibrinogen map to 4q2. Am. J. Hum. Genet. 36:

[9443] 760-768, 1984.

[9444] 7.

[9445] Sheng, Y.; Herzog, H.; Krilis, S. A.: Cloning and characterization

[9446] of the gene encoding the mouse beta-2-glycoprotein I. Genomics 41:

[9447] 128-130, 1997.

[9448] 8.

[9449] Fields, S.; Jang, S. K.: Presence of a potent transcription

activating

[9450] sequence in the p53 protein. Science 249: 1046-1049, 1990.

[9451] 9.

[9452] Firestein, G. S.; Echeverri, F.; Yeao, M.; Zvaifler, N. J.; Green,

[9453] D. R.: Somatic mutations in the p53 tumor suppressor gene in rheumatoid

[9454] arthritis synovium. Proc. Nat. Acad. Sci. 94: 10895-10900, 1997.

[9455] 10.

[9456] Foster, B. A.; Coffey, H. A.; Morin, M. J.; Rastinejad, F.: Pharmacological

[9457] rescue of mutant p53 conformation and function. Science 286: 2507-2510,

[9458] 1999.

[9459] 11.

[9460] Collod-Beroud, G.; Beroud, C.; Ades, L.; Black, C.; Boxer, M.;

[9461] Brock, D. J. H.; Holman, K. J.; de Paepe, A.; Francke, U.; Grau, U.;

[9462] Hayward, C.; Klein, H.-G.; Liu, W.; Nuytinck, L.; Peltonen,

L.; Alvarez

[9463] Perez, A. B.; Rantamaki, T.; Junien, C.; Boileau, C.: Marfan database

[9464] (third edition): new mutations and new routines for the software. Nucleic

[9465] Acids Res. 26: 229-233, 1998.

[9466] 12.

[9467] Buckway, C. K.; Wilson, E. M.; Ahlsen, M.; Bang, P.; Oh, Y.; Rosenfeld,

[9468] R. G.: Mutation of three critical amino acids of the N-terminal domain

[9469] of IGF-binding protein-3 essential for high affinity IGF binding. J.

[9470] Clin. Endocr. Metab. 86: 4943-4950, 2001.

[9471] 13.

[9472] Vestergaard, P.; Hermann, A.P.; Orskov, H.; Mosekilde, L.; The

[9473] Danish Osteoporosis Prevention Study: Effect of sex hormone replacement

[9474] on the insulin-like growth factor system and bone mineral: a cross-sectional

[9475] and longitudinal study in 595 perimenopausal women

participating in
[9476] the Danish osteoporosis prevention study. J. Clin. Endocr. Metab. 84:
[9477] 2286-2290, 1999.

[9478] 14.

[9479] Holm, C.; Kirchgessner, T. G.; Svenson, K. L.; Fredrikson, G.;
[9480] Nilsson, S.; Miller, C. G.; Shively, J. E.; Heinzmann, C.; Sparkes,
[9481] R. S.; Mohandas, T.; Lusis, A. J.; Belfrage, P.; Schotz, M. C.: Hormone-sensitive
[9482] lipase: sequence, expression, and chromosomal localization to 19cent-q13.3. Science 241:
[9483] 1503-1506, 1988.

[9484] 15.

[9485] Gilgenkrantz, S.; Vigneron, C.; Gregoire, M. J.; Pernot, C.; Raspiller,
[9486] A.: Association of del(11)(p15.1p12), aniridia, catalase deficiency,
[9487] and cardiomyopathy. Am. J. Med. Genet. 13: 39-49, 1982.

[9488] 16.

[9489] Sander, M.; Chavoshan, B.; Harris, S. A.; Iannaccone, S. T.; Stull,
[9490] J. T.; Thomas, G. D.; Victor, R. G.: Functional muscle ischemia in
[9491] neuronal nitric oxide synthase-deficient skeletal muscle of children
[9492] with Duchenne muscular dystrophy. Proc. Nat. Acad. Sci. 97: 13818-13823,
[9493] 2000.

[9494] 17.

[9495] Li, P.; Wood, K.; Mamon, H.; Haser, W.; Roberts, T.: Raf-1: a
[9496] kinase currently without a cause but not lacking in effects. Cell 64:
[9497] 479-482, 1991.

[9498] 18.

[9499] Eiberg, H.; Mohr, J.; Schmiegelow, K.; Nielsen, L. S.; Williamson,
[9500] R.: Linkage relationships of paraoxonase (PON) with other markers:
[9501] indication of PON-cystic fibrosis synteny. Clin. Genet. 28: 265-271,

[9502] 1985.

[9503] 19.

[9504] Berger, J.; Garattini, E.; Hua, J.-C.; Udenfriend, S.: Cloning

[9505] and sequencing of human intestinal alkaline phosphatase cDNA. Proc.

[9506] Nat. Acad. Sci. 84: 695-698, 1987.

[9507] 20.

[9508] Nilsson, I. M.; Tengborn, L.: Impaired fibrinolysis: new evi-dence

[9509] in relation to thrombosis.In: Jespersen, J.; Kluft, C.; Kors-gaard,

[9510] O.: Clinical Aspects of Fibrinolysis and Thrombolysis. Esb-jerg:

[9511] South Jutland Univ. Press (pub.) 1983. Pp. 273-291.

[9512] 21.

[9513] Dihlmann, S.; Gebert, J.; Siermann, A.; Herfarth, C.; von Knebel

[9514] Doeberitz, M.: Dominant negative effect of the APC(1309) mutation:

[9515] a possible explanation for genotype-phenotype correla-tions in familial

[9516] adenomatous polyposis. Cancer Res. 59: 1857-1860, 1999.

[9517] 22.

[9518] Foster, P. L.; Eisenstadt, E.; Miller, J. H.: Base substitution

[9519] mutations induced by metabolically activated aflatoxin B1. Proc.

[9520] Nat. Acad. Sci. 80: 2695-2698, 1983.

[9521] 23.

[9522] Franklin, W. A.; Gazdar, A. F.; Haney, J.; Wistuba, I. I.; La

[9523] Rosa, F. G.; Kennedy, T.; Ritchey, D. M.; Miller, Y. E.: Widely dispersed

[9524] p53 mutation in respiratory epithelium: a novel mecha- nism for field

[9525] carcinogenesis. J. Clin. Invest. 100: 2133-2137, 1997.

[9526] 24.

[9527] Frebourg, T.; Friend, S. H.: Cancer risks from germline P53 mutations. J.

[9528] Clin. Invest. 90: 1637-1641, 1992.

[9529] 25.

[9530] Frebourg, T.; Kassel, J.; Lam, K. T.; Gryka, M. A.; Barbier,

N.;

[9531] Andersen, T. I.; Borresen, A.-L.; Friend, S. H.: Germ-line mutations

[9532] of the p53 tumor suppressor gene in patients with high risk for cancer

[9533] inactivate the p53 protein. Proc. Nat. Acad. Sci. 89: 6413-6417,

[9534] 1992.

[9535] 26.

[9536] Fagin, J. A.; Matsuo, K.; Karmakar, A.; Chen, D. L.; Tang, S.-H.;

[9537] Koeffler, H. P.: High prevalence of mutations of the p53 gene in

[9538] poorly differentiated human thyroid carcinomas. J. Clin. Invest. 91:

[9539] 179-184, 1993.

[9540] 27.

[9541] Farmer, G.; Bargonetti, J.; Zhu, H.; Friedman, P.; Prywes, R.;

[9542] Prives, C.: Wild-type p53 activates transcription in vitro. Nature 358:

[9543] 83-86, 1992.

[9544] 28.

[9545] Felix, C. A.; D'Amico, D.; Mitsudomi, T.; Nau, M. M.; Li, F. P.;

[9546] Fraumeni, J. F., Jr.; Cole, D. E.; McCalla, J.; Reaman, G. H.; Whang-Peng,

[9547] J.; Knutsen, T.; Minna, J. D.; Poplack, D. G.: Absence of hereditary

[9548] p53 mutations in 10 familial leukemia pedigrees. J. Clin. Invest. 90:

[9549] 653-658, 1992.

[9550] 29.

[9551] Felix, C. A.; Nau, M. M.; Takahashi, T.; Mitsudomi, T.; Chiba,

[9552] I.; Poplack, D. G.; Reaman, G. H.; Cole, D. E.; Letterio, J. J.; Whang-Peng,

[9553] J.; Knutsen, T.; Minna, J. D.: Hereditary and acquired p53 gene mutations

[9554] in childhood acute lymphoblastic leukemia. J. Clin. Invest. 89:

[9555] 640-647, 1992.

[9556] 30.

[9557] Fukasawa, K.; Choi, T.; Kuriyama, R.; Rulong, S.; Vande Woude,
[9558] G. F.: Abnormal centrosome amplification in the absence of p53. Science 271:
[9559] 1744-1747, 1996.

[9560] 31.

[9561] Gao, Y.; Ferguson, D. O.; Xie, W.; Manis, J. P.; Sekiguchi, J.;

[9562] Frank, K. M.; Chaudhuri, J.; Horner, J.; DePinho, R. A.; Alt, F. W.
[9563] : Interplay of p53 and DNA-repair protein XRCC4 in tumorigenesis,
[9564] genomic stability and development. Nature 404: 897-900, 2000.
[9565] 32.

[9566] Gu, J.; Kawai, H.; Wiederschain, D.; Yuan, Z.-M.: Mechanism of
[9567] functional inactivation of a Li-Fraumeni syndrome p53 that has a mutation
[9568] outside of the DNA-binding domain. Cancer Res. 61: 1741-1746, 2001.
[9569] 33.

[9570] Guran, S.; Tunca, Y.; Imirzalioglu, N.: Hereditary TP53 codon
[9571] 292 and somatic P16(INK4A) codon 94 mutations in a Li-Fraumeni syndrome
[9572] family. Cancer Genet. Cytogenet. 113: 145-151, 1999.

[9573] 34.

[9574] Hainaut, P.; Soussi, T.; Shomer, B.; Hollstein, M.; Green-blatt,
[9575] M.; Hovig, E.; Harris, C. C.; Montesano, R.: Database of p53 gene
[9576] somatic mutations in human tumors and cell lines: up-dated compilation
[9577] and future prospects. Nucleic Acids Res. 25: 151-157, 1997.

[9578] 35.

[9579] Halevy, O.; Michalovitz, D.; Oren, M.: Different tumor-derived
[9580] p53 mutants exhibit distinct biological activities. Science 250:
[9581] 113-116, 1990.

[9582] 36.

[9583] Harlow, E.; Williamson, N. M.; Ralston, R.; Helfman, D. M.; Adams,
[9584] T. E.: Molecular cloning and in vitro expression of a cDNA clone
[9585] for human cellular tumor antigen p53. Molec. Cell. Biol. 5: 1601-1610,
[9586] 1985.

[9587] 37.

[9588] Harper, J. W.; Adami, G. R.; Wei, N.; Keyomarsi, K.; Elledge,
[9589] S. J.: The p21 Cdk-interacting protein Cip1 is a potent inhibitor
[9590] of G1 cyclin-dependent kinases. Cell 75: 805-816, 1993.

[9591] 38.

[9592] Harris, C. C.: p53: at the crossroads of molecular carcinogenesis
[9593] and risk assessment. Science 262: 1980-1981, 1993.

[9594] 39.

[9595] Harris, C. C.; Hollstein, M.: Clinical implications of the p53
[9596] tumor-suppressor gene. New Eng. J. Med. 329: 1318-1327, 1993.

[9597] 40.

[9598] Hernandez-Boussard, T.; Rodriguez-Tome, P.; Montesano, R.; Hainaut,

[9599] P.: IARC p53 mutation database: a relational database to compile

[9600] and analyze p53 mutations in human tumors and cell lines. Hum. Mutat. 14:

[9601] 1-8, 1999.

[9602] 41.

[9603] Herrmann, M. A.; Hay, I. D.; Bartelt, D. H.; Ritland, S. R., Jr.;

[9604] Dahl, R. J.; Grant, C. S.; Jenkins, R. B.: Cytogenetic and molecular

[9605] genetic studies of follicular and papillary thyroid cancers. J. Clin.

[9606] Invest. 88: 1596-1604, 1991.

[9607] 42.

[9608] Hirao, A.; Kong, Y.-Y.; Matsuoka, S.; Wakeham, A.; Ruland, J.;

[9609] Yoshida, H.; Liu, D.; Elledge, S. J.; Mak, T. W.: DNA dam-age-induced

[9610] activation of p53 by the checkpoint kinase Chk2. Science 287: 1824-1827,
[9611] 2000.

[9612] 43.

[9613] Hollstein, M.; Shomer, B.; Greenblatt, M.; Soussi, T.; Hovig,
[9614] E.; Montesano, R.; Harris, C. C.: Somatic point mutations in the
[9615] p53 gene of human tumors and cell lines: updated compilation. Nucleic
[9616] Acids Res. 24: 141-146, 1996.

[9617] 44.

[9618] Hollstein, M.; Sidransky, D.; Vogelstein, B.; Harris, C. C.:
[9619] p53 mutations in human cancers. Science 253: 49-53, 1991.
[9620] 45.

[9621] Hollstein, M. C.; Metcalf, R. A.; Welsh, J. A.; Montesano, R.;
[9622] Harris, C. C.: Frequent mutation of the p53 gene in human esophageal
[9623] cancer. Proc. Nat. Acad. Sci. 87: 9958-9961, 1990.

[9624] 46.

[9625] Hsu, I. C.; Metcalf, R. A.; Sun, T.; Welsh, J. A.; Wang, N. J.;

[9626] Harris, C. C.: Mutational hotspot in the p53 gene in human hepatocellular

[9627] carcinomas. Nature 350: 427-428, 1991.

[9628] 47.

[9629] Hung, J.; Mims, B.; Lozano, G.; Strong, L.; Harvey, C.; Chen,

[9630] T. T.-Y.; Stastny, V.; Tomlinson, G.: TP53 mutation and haplotype

[9631] analysis of two large African American families. Hum. Mutat. 14:

[9632] 216-221, 1999.

[9633] 48.

[9634] Kaname, T.; Miyauchi, T.; Kuwano, A.; Matsuda, Y.; Muramatsu, T.;

[9635] Kajii, T.: Mapping basigin (BSG), a member of the immunoglobulin

[9636] superfamily, to 19p13.3. Cytogenet. Cell Genet. 64: 195-197, 1993.

[9637] 49.

[9638] Kanekura, T.; Miyauchi, T.; Tashiro, M.; Muramatsu, T.:

Basigin,

[9639] a new member of the immunoglobulin superfamily: genes in different

[9640] mammalian species, glycosylation changes in the molecule from adult

[9641] organs and possible variation in the N-terminal se-quences. Cell Struct.

[9642] Funct. 16: 23-30, 1991.

[9643] 50.

[9644] Kuno, N.; Kadomatsu, K.; Fan, Q.-W.; Hagihara, M.; Senda, T.; Mizutani,

[9645] S.; Muramatsu, T.: Female sterility in mice lacking the basigin gene,

[9646] which encodes a transmembrane glycoprotein belonging to the immunoglobulin

[9647] superfamily. FEBS Lett. 425: 191-194, 1998.

[9648] 51.

[9649] Miyauchi, T.; Kanekura, T.; Yamaoka, A.; Ozawa, M.; Miyazawa, S.;

[9650] Muramatsu, T.: Basigin, a new, broadly distributed mem-ber of the

[9651] immunoglobulin superfamily, has strong homology with

both the immunoglobulin
[9652] V domain and the beta-chain of major histocompatibility complex class
[9653] II antigen. J. Biochem. 107: 316-323, 1990.
[9654] 52.
[9655] Miyauchi, T.; Masuzawa, Y.; Muramatsu, T.: The basigin group of
[9656] the immunoglobulin superfamily: complete conservation of a segment
[9657] in and around transmembrane domains of human and mouse basigin and
[9658] chicken HT7 antigen. J. Biochem. 110: 770-774, 1991.
[9659] 53.
[9660] Naruhashi, K.; Kadomatsu, K.; Igakura, T.; Fan, Q.-W.; Kuno, N.;
[9661] Muramatsu, H.; Miyauchi, T.; Hasegawa, T.; Itoh, A.; Muramatsu, T.;
[9662] Nabeshima, T.: Abnormalities of sensory and memory functions in mice
[9663] lacking Bsg gene. Biochem. Biophsy. Res. Commun. 236: 733-737, 1997.
[9664] 54.

[9665] Snyder, F. F.; Lin, C. C.; Rudd, N. L.; Shearer, J. E.; Heikkila,

[9666] E. M.; Hoo, J. J.: A de novo case of trisomy 10p: gene dosage studies

[9667] of hexokinase, inorganic pyrophosphatase and adenosine kinase. Hum.

[9668] Genet. 67: 187-189, 1984.

[9669] 55.

[9670] Abe, M.; Higuchi, I.; Morisaki, H.; Morisaki, T.; Osame, M.: Myoadenylate

[9671] deaminase deficiency with progressive muscle weakness and atrophy

[9672] caused by new missense mutations in AMPD1 gene: case report in a Japanese

[9673] patient. Neuromusc. Disord. 10: 472-477, 2000.

[9674] 56.

[9675] Fishbein, W. N.: Myoadenylate deaminase deficiency: inherited

[9676] and acquired forms. Biochem. Med. 33: 158-169, 1985.

[9677] 57.

[9678] Patten, J. L.; Smallwood, P. M.; Eil, C.; Johns, D. R.; Valle,

[9679] D.; Steel, G.; Levine, M. A.: An initiator codon mutation in the
[9680] gene encoding the alpha subunit of Gs in pseudohy-poparathyroidism
[9681] type IA (PHP IA). (Abstract) Am. J. Hum. Genet. 45 (suppl.): A212
[9682] only, 1989.

[9683] 58.

[9684] Yu, S.; Yu, D.; Lee, E.; Eckhaus, M.; Lee, R.; Corria, Z.; Ac-cili,
[9685] D.; Westphal, H.; Weinstein, L. S.: Variable and tissue-specific
[9686] hormone resistance in heterotrimeric Gs protein alpha-subunit (Gs-alpha)
[9687] knockout mice is due to tissue-specific imprinting of the Gs-alpha
[9688] gene. Proc. Nat. Acad. Sci. 95: 8715-8720, 1998.

[9689] 59.

[9690] Abdo, Y.; Rousseaux, J.; Dautrevaux, M.: Proalbumin Lille, a new
[9691] variant of human serum albumin. FEBS Lett. 131: 286-288, 1981.

[9692] 60.

[9693] Adams, M. S.: Genetic diversity in serum albumin. J. Med. Genet. 3:

[9694] 198–202, 1966.

[9695] 61.

[9696] Arai, K.; Huss, K.; Madison, J.; Putnam, F. W.; Salzano, F. M.;

[9697] Franco, M. H. L. P.; Santos, S. E. B.; Freitas, M. J. M.: Amino acid

[9698] substitutions in albumin variants found in Brazil. Proc. Nat. Acad.

[9699] Sci. 86: 1821–1825, 1989.

[9700] 62.

[9701] Arai, K.; Ishioka, N.; Huss, K.; Madison, J.; Putnam, F. W.: Identical

[9702] structural changes in inherited albumin variants from dif-ferent populations. Proc.

[9703] Nat. Acad. Sci. 86: 434–438, 1989.

[9704] 63.

[9705] Arai, K.; Madison, J.; Huss, K.; Ishioka, N.; Satoh, C.; Fujita,

[9706] M.; Neel, J. V.; Sakurabayashi, I.; Putnam, F. W.: Point substitutions
[9707] in Japanese alloalbumins. Proc. Nat. Acad. Sci. 86: 6092-6096, 1989.
[9708] 64.
[9709] Arai, K.; Madison, J.; Shimizu, A.; Putnam, F. W.: Point substitutions
[9710] in albumin genetic variants from Asia. Proc. Nat. Acad. Sci. 87:
[9711] 497-501, 1990.
[9712] 65.
[9713] Arends, T.; Gallango, M. L.; Layrisse, M.; Wilbert, J.; Heinen,
[9714] H. D.: Albumin Warao: new type of human alloalbuminemia. Blood 33:
[9715] 414-420, 1969.
[9716] 66.
[9717] Arvan, D.; Blumberg, B.; Melartin, L.: Transient bisalbuminemia
[9718] induced by drugs. Clin. Chim. Acta 22: 211-218, 1968.
[9719] 67.

[9720] Au, H. Y. N.; Brand, S.; Hutchinson, D. W.; Matejtschuk, P.: Albumins

[9721] Warwick 1 and Warwick 2, two human albumin variants. IRCS Med. Sci. 12:

[9722] 56-57, 1984.

[9723] 68.

[9724] Barlow, J. W.; Csicsmann, J. M.; Meinhold, H.; Lim, C.-F.; Stockigt,

[9725] J. R.: Familial dysalbuminaemic hyperthyroxinaemia: studies of albumin

[9726] binding and implications for hormone action. Clin. Endocr. 24: 39-47,

[9727] 1986.

[9728] 69.

[9729] Barlow, J. W.; Csicsmann, J. M.; White, E. L.; Funder, J. W.;

[9730] Stockigt, J. R.: Familial euthyroid thyroxine excess: characterization

[9731] of abnormal intermediate affinity thyroxine binding to albumin. J.

[9732] Clin. Endocr. Metab. 55: 244-250, 1982.

[9733] 70.

[9734] Fishbein, W. N.; Armbrustmacher, V. W.; Griffin, J. L.: Myo-
adenylate
[9735] deaminase deficiency: a new disease of muscle. Science
200: 545-548,
[9736] 1978.

[9737] 71.

[9738] Bennhold, H.; Kallee, E.: Comparative studies on the half-
life
[9739] of I(131) labelled albumins and nonradioactive human
serum albumin
[9740] in a case of analbuminemia. J. Clin. Invest. 38: 863-872,
1959.
[9741] 72.

[9742] St. George-Hyslop, P. H.; Tanzi, R. E.; Polinsky, R. J.;
Haines,
[9743] J. L.; Nee, L.; Watkins, P. C.; Myers, R. H.; Feldman, R. G.;
Pollen,
[9744] D.; Drachman, D.; Growdon, J.; Bruni, A.; Foncin, J.-F.;
Salmon, D.;
[9745] Frommelt, P.; Amaducci, L.; Sorbi, S.; Piacentini, S.; Stew-
art, G.
[9746] D.; Hobbs, W. J.; Conneally, P. M.; Gusella, J. F.: The ge-

netic defect
[9747] causing familial Alzheimer's disease maps on chromo-
some 21. Science 235:
[9748] 885-890, 1987.

[9749] 73.

[9750] Shozu, M.; Akasofu, K.; Harada, T.; Kubota, Y.: A new
cause of
[9751] female pseudohermaphroditism: placental aromatase de-
ficiency. J.
[9752] Clin. Endocr. Metab. 72: 560-566, 1991.

[9753] 74.

[9754] Hadley, T.; Saul, A.; Lamont, G.; Hudson, D. E.; Miller, L.
H.;
[9755] Kidson, C.: Resistance of Melanesian elliptocytes
(ovalocytes) to
[9756] invasion by Plasmodium knowlesi and Plasmodium falci-
parum malaria
[9757] parasites in vitro. J. Clin. Invest. 71: 780-782, 1983.

[9758] 75.

[9759] Drouet, B.; Garcia, L.; Simon-Chazottes, D.; Mattei, M. G.;
Guenet,

[9760] J.-L.; Schwartz, A.; Varadi, G.; Pincon-Raymond, M.: The gene coding

[9761] for the alpha-1 subunit of the skeletal dihydropyridine receptor (Cchl1a3

[9762] = mdg) maps to mouse chromosome 1 and human 1q32. Mammalian Genome 4:

[9763] 499-503, 1993.

[9764] 76.

[9765] Cole, W. G.; Hall, R. K.; Rogers, J. G.: The clinical features

[9766] of spondyloepiphyseal dysplasia congenita resulting from the substitution

[9767] of glycine 997 by serine in the alpha-1(II) chain of type II collagen. J.

[9768] Med. Genet. 30: 27-35, 1993.

[9769] 77.

[9770] Dalla Venezia, N.; Gilsanz, F.; Alloisio, N.; Ducluzeau, M.-T.;

[9771] Benz, E. J., Jr.; Delaunay, J.: Homozygous 4.1(-) hereditary elliptocytosis

[9772] associated with a point mutation in the downstream initiation codon

[9773] of protein 4.1 gene. J. Clin. Invest. 90: 1713-1717, 1992.

[9774] 78.

[9775] Elliott, K. J.; Ellis, S. B.; Berckhan, K. J.; Urrutia, A.; Chavez-
Noriega,
[9776] L. E.; Johnson, E. C.; Velicelebi, G.; Harpold, M. M.: Com-
parative
[9777] structure of human neuronal alpha(2)-alpha(7) and
beta(2)-beta(4)
[9778] nicotinic acetylcholine receptor subunits and functional
expression
[9779] of the alpha(2), alpha(3), alpha(4), alpha(7), beta(2), and
beta(4)
[9780] subunits. J. Molec. Neurosci. 7: 217-228, 1996.

[9781] 79.

[9782] Budarf, M. L.; Korenberg, J. R.; Simon, M.; Emanuel, B. S.:
Regional
[9783] assignment of the guanine nucleotide binding protein,
GNAZ, to chromosome
[9784] 22 (22q11.1-q11.2). (Abstract) Cytogenet. Cell Genet. 58:
2046-2047,
[9785] 1991.

[9786] 80.

[9787] Jayawardena-Wolf, J.; Benlagha, K.; Chiu, Y.-H.; Mehr, R.; Bendelac,

[9788] A.: CD1d endosomal trafficking is independently regulated by an intrinsic

[9789] CD1d-encoded tyrosine motif and by the invariant chain. Immunity 15:

[9790] 897-908, 2001.

[9791] 81.

[9792] Yang, W.-S.; Nevin, D. N.; Peng, R.; Brunzell, J. D.; Deeb, S.

[9793] S.: A mutation in the promoter of the lipoprotein lipase (LPL) gene

[9794] in a patient with familial combined hyperlipidemia and low LPL activity. Proc.

[9795] Nat. Acad. Sci. 92: 4462-4466, 1995.

[9796] 82.

[9797] Lloyd, A.; Modi, W.; Sprenger, H.; Cevario, S.; Oppenheim, J.;

[9798] Kelvin, D.: Assignment of genes for interleukin-8 receptors (IL8R)

[9799] A and B to human chromosome band 2q35. Cytogenet.

Cell Genet. 63:
[9800] 238-240, 1993.

[9801] 83.

[9802] Murgia, C.; Blaikie, P.; Kim, N.; Dans, M.; Petrie, H. T.; Gi-
ancotti,
[9803] F. G.: Cell cycle and adhesion defects in mice carrying a
targeted
[9804] deletion of the integrin beta-4 cytoplasmic domain. EMBO
J. 17:
[9805] 3940-3951, 1998.

[9806] 84.

[9807] Roberts, A. N.; Leighton, B.; Todd, J. A.; Cockburn, D.;
Schofield,
[9808] P. N.; Sutton, R.; Holt, S.; Boyd, Y.; Day, A. J.; Foot, E. A.;
Willis,
[9809] A. C.; Reid, K. B. M.; Cooper, G. J. S.: Molecular and func-
tional
[9810] characterization of amylin, a peptide associated with type
2 diabetes
[9811] mellitus. Proc. Nat. Acad. Sci. 86: 9662-9666, 1989.

[9812] 85.

[9813] Harada, T.; Harada, C.; Nakayama, N.; Okuyama, S.; Yoshida, K.;
[9814] Kohsaka, S.; Matsuda, H.; Wada, K.: Modification of glial-neuronal
[9815] cell interactions prevents photoreceptor apoptosis during light-induced
[9816] retinal degeneration. Neuron 26: 533-541, 2000.

[9817] 86.

[9818] Fakharzadeh, S.; Trusko, S. P.; George, D. L.: Tumorigenic potential
[9819] associated with enhanced expression of a gene that is amplified in
[9820] a mouse tumor cell line. EMBO J. 10: 1565-1569, 1991.

[9821] 87.

[9822] Hoshino, Y.; Horikawa, I.; Oshimura, M.; Yuasa, Y.: Normal human
[9823] chromosome 5, on which a familial adenomatous polypo-sis gene is located,
[9824] has tumor suppressive activity. Biochem. Biophys. Res. Commun. 174:
[9825] 298-304, 1991.

[9826] 88.

[9827] Bonaiti-Pellie, C.; Briard-Guillemot, M. L.; Feingold, J.; Frezal,

[9828] J.: Associated congenital malformations in retinoblastoma. Clin.

[9829] Genet. 7: 37-39, 1975.

[9830] 89.

[9831] Radford, D. M.; Nakai, H.; Pegg, A. E.; Shows, T. B.: Mapping

[9832] genes for rate-limiting enzymes in polyamine biosynthesis. (Abstract) Am.

[9833] J. Hum. Genet. 41: A35 only, 1987.

[9834] 90.

[9835] Zarkowsky, H. S.; Mohandas, N.; Speaker, C. B.; Shohet, S. B.

[9836] : A congenital haemolytic anaemia with thermal sensitivity of the

[9837] erythrocyte membrane. Brit. J. Haemat. 29: 537-543, 1975.

[9838] 91.

[9839] Kogerman, P.; Grimm, T.; Kogerman, L.; Krause, D.; Un-

den, A. B.;

[9840] Sandstedt, B.; Toftgard, R.; Zaphiropoulos, P. G.: Mammalian suppressor-of-fused

[9841] modulates nuclear-cytoplasmic shuttling of GLI-1. Nature Cell Biol. 1:

[9842] 312-319, 1999.

[9843] 92.

[9844] Ruppert, J. M.; Kinzler, K. W.; Wong, A. J.; Bigner, S. H.; Kao,

[9845] F.-T.; Law, M. L.; Seuanez, H. N.; O'Brien, S. J.; Vogelstein, B.

[9846] : The GLI-Kruppel family of human genes. Molec. Cell. Biol. 8: 3104-3113,

[9847] 1988.

[9848] 93.

[9849] Grachtchouk, M.; Mo, R.; Yu, S.; Zhang, X.; Sasaki, H.; Hui, C.;

[9850] Dlugosz, A. A.: Basal cell carcinomas in mice overexpressing Gli2

[9851] in skin. (Letter) Nature Genet. 24: 216-217, 2000.

[9852] 94.

[9853] Matsumoto, N.; Fujimoto, M.; Kato, R.; Niikawa, N.: As-
signment
[9854] of the human GLI2 gene to 2q14 by fluorescence in situ
hybridization. Genomics 36:
[9855] 220-221, 1996.

[9856] 95.

[9857] Ishikawa, T.; Kibe, T.; Wada, Y.: Deletion of small nuclear
ribonucleoprotein
[9858] polypeptide N (SNRPN) in Prader-Willi syndrome detected
by fluorescence
[9859] in situ hybridization: two sibs with the typical phenotype
without
[9860] a cytogenetic deletion in chromosome 15q. Am. J. Med.
Genet. 62:
[9861] 350-352, 1996.

[9862] 96.

[9863] Gubina, E.; Ruiz-Hidalgo, M. J.; Baladron, V.; Laborda, J.:
Assignment
[9864] of DLK1 to human chromosome band 14q32 by in situ hy-
bridization. Cytogenet.
[9865] Cell Genet. 84: 206-207, 1999.

[9866] 97.

[9867] Helman, L. J.; Thiele, C. J.; Linehan, W. M.; Nelkin, B. D.; Baylin,

[9868] S. B.; Israel, M. A.: Molecular markers of neuroendocrine development

[9869] and evidence of environmental regulation. Proc. Nat. Acad. Sci. 84:

[9870] 2336-2339, 1987.

[9871] 98.

[9872] Jensen, C. H.; Krogh, T. N.; Hojrup, P.; Clausen, P. P.; Skjodt,

[9873] K.; Larsson, L.-I.; Enghild, J. J.; Teisner, B.: Protein structure

[9874] of fetal antigen 1 (FA1): a novel circulating human epidermal-growth-factor-like

[9875] protein expressed in neuroendocrine tumors and its relation to the

[9876] gene products of dlk and pG2. Europ. J. Biochem. 225: 83-92, 1994.

[9877] 99.

[9878] Aguzzi, A.: Personal Communication. Zurich, Switzerland 3/20/1997.

[9879] 100.

[9880] Aguzzi, A.; Brandner, S.: The genetics of prions--a con-
tradiction
[9881] in terms? Lancet 354: 22-25, 1999.

[9882] 101.

[9883] Basler, K.; Oesch, B.; Scott, M.; Westaway, D.; Walchli, M.;
Groth,
[9884] D. F.; McKinley, M. P.; Prusiner, S. B.; Weissmann, C.:
Scrapie and
[9885] cellular PrP isoforms are encoded by the same chromoso-
mal gene. Cell 46:
[9886] 417-428, 1986.

[9887] 102.

[9888] Bertoni, J. M.; Brown, P.; Goldfarb, L. G.; Rubenstein, R.;
Gajdusek,
[9889] D. C.: Familial Creutzfeldt-Jakob disease (codon 200 mu-
tation) with
[9890] supranuclear palsy. J.A.M.A. 268: 2413-2415, 1992.

[9891] 103.

[9892] Beyreuther, K.; Masters, C. L.: Catching the culprit prion.
Nature 370:

[9893] 419-420, 1994.

[9894] 104.

[9895] Bosque, P. J.; Vnencak-Jones, C. L.; Johnson, M. D.; Whit-
lock,
[9896] J. A.; McLean, M. J.: A PrP gene codon 178 base substitu-
tion and
[9897] a 24-bp interstitial deletion in familial Creutzfeldt-Jakob
disease. Neurology 42:
[9898] 1864-1870, 1992.

[9899] 105.

[9900] Bounhar, Y.; Zhang, Y.; Goodyer, C. G.; LeBlanc, A.: Prion
protein
[9901] protects human neurons against Bax-mediated apoptosis.
J. Biol. Chem. 276:
[9902] 39145-39149, 2001.

[9903] 106.

[9904] Brown, P.; Galvez, S.; Goldfarb, L. G.; Nieto, A.; Cartier, L.;

[9905] Gibbs, C. J., Jr.; Gajdusek, D. C.: Familial Creutzfeldt-
Jakob disease
[9906] in Chile is associated with the codon 200 mutation of the
PRNP amyloid

[9907] precursor gene on chromosome 20. J. Neurol. Sci. 112: 65-67, 1992.

[9908] 107.

[9909] Brown, P.; Goldfarb, L. G.; Gajdusek, D. C.: The new biol-
ogy

[9910] of spongiform encephalopathy: infectious amyloidoses with a genetic

[9911] twist. Lancet 337: 1019-1022, 1991.

[9912] 108.

[9913] Brown, P.; Goldfarb, L. G.; Kovanen, J.; Haltia, M.; Cathala,

[9914] F.; Sulima, M.; Gibbs, C. J., Jr.; Gajdusek, D. C.: Phenotypic characteristics

[9915] of familial Creutzfeldt-Jakob disease associated with the codon 178-asn

[9916] PRNP mutation. Ann. Neurol. 31: 282-285, 1992.

[9917] 109.

[9918] Brown, P.; Goldfarb, L. G.; McCombie, W. R.; Nieto, A.; Squillacote,

[9919] D.; Sheremata, W.; Little, B. W.; Godec, M. S.; Gibbs, C. J., Jr.;

[9920] Gajdusek, D. C.: Atypical Creutzfeldt-Jakob disease in an

American
[9921] family with an insert mutation in the PRNP amyloid precursor gene. Neurology 42:
[9922] 422-427, 1992.

[9923] 110.

[9924] Bueler, H.; Aguzzi, A.; Sailer, A.; Greiner, R.-A.; Autenried,

[9925] P.; Aguet, M.; Weissman, C.: Mice devoid of PrP are resistant to
[9926] scrapie. Cell 73: 1339-1347, 1993.

[9927] 111.

[9928] Bueler, H.; Raeber, A.; Sailer, A.; Fischer, M.; Aguzzi, A.; Weissmann,
[9929] C.: High prion and PrP(Sc) levels but delayed onset of disease in
[9930] scrapie-inoculated mice heterozygous for a disrupted PrP gene. Molec.
[9931] Med. 1: 19-30, 1994.

[9932] 112.

[9933] Campbell, T. A.; Palmer, M. S.; Will, R. G.; Gibb, W. R. G.; Luthert,
[9934] P. J.; Collinge, J.: A prion disease with a novel 96-base

pair insertional
[9935] mutation in the prion protein gene. Neurology 46:
761-766, 1996.
[9936] 113.

[9937] Carlson, G. A.; Kingsbury, D. T.; Goodman, P. A.; Coleman,
S.;
[9938] Marshall, S. T.; DeArmond, S.; Westaway, D.; Prusiner, S.
B.: Linkage
[9939] of prion protein and scrapie incubation time genes. Cell
46: 503-511,
[9940] 1986.

[9941] 114.

[9942] Chapman, J.; Arlazoroff, A.; Goldfarb, L. G.; Cervenakova,
L.;
[9943] Neufeld, M. Y.; Werber, E.; Herbert, M.; Brown, P.; Gaj-
dusek, D. C.;
[9944] Korczyn, A. D.: Fatal insomnia in a case of familial
Creutzfeldt-Jakob
[9945] disease with the codon 200(lys) mutation. Neurology 46:
758-761,
[9946] 1996.

[9947] 115.

[9948] Chapman, J.; Ben-Israel, J.; Goldhammer, Y.; Korczyn, A. D.:
[9949] The risk of developing Creutzfeldt-Jakob disease in subjects with
[9950] the PRNP gene codon 200 point mutation. Neurology 44: 1683-1686,
[9951] 1994.

[9952] 116.

[9953] Chapman, J.; Brown, P.; Rabey, J. M.; Goldfarb, L. G.; Inzelberg,
[9954] R.; Gibbs, C. J., Jr.; Gajdusek, D. C.; Korczyn, A. D.: Transmission
[9955] of spongiform encephalopathy from a familial Creutzfeldt-Jakob disease
[9956] patient of Jewish Libyan origin carrying the PRNP codon 200 mutation. Neurology 42:
[9957] 1249-1250, 1992.

[9958] 117.

[9959] Chapman, J.; Korczyn, A. D.: Genetic and environmental factors
[9960] determining the development of Creutzfeldt-Jakob disease in Libyan

[9961] Jews. Neuroepidemiology 10: 228–231, 1991.

[9962] 118.

[9963] Chiesa, R.; Drisaldi, B.; Quaglio, E.; Migheli, A.; Piccardo,

[9964] P.; Ghetti, B.; Harris, D. A.: Accumulation of protease-re-
sistant

[9965] prion protein (PrP) and apoptosis of cerebellar granule
cells in transgenic

[9966] mice expressing a PrP insertional mutation. Proc. Nat.
Acad. Sci. 97:

[9967] 5574–5579, 2000.

[9968] 119.

[9969] Chiesa, R.; Piccardo, P.; Ghetti, B.; Harris, D. A.: Neurolog-
ical

[9970] illness in transgenic mice expressing a prion protein with
an insertional

[9971] mutation. Neuron 21: 1339–1351, 1998.

[9972] 120.

[9973] Collinge, J.: Human prion diseases and bovine spongiform
encephalopathy

[9974] (BSE). Hum. Mol. Genet. 6: 1699–1705, 1997.

[9975] 121.

[9976] Collinge, J.; Harding, A. E.; Owen, F.; Poulter, M.; Lofthouse,

[9977] R.; Boughey, A. M.; Shah, T.; Crow, T. J.: Diagnosis of Gerstmann-Straussler

[9978] syndrome in familial dementia with prion protein gene analysis. Lancet II:

[9979] 15-17, 1989.

[9980] 122.

[9981] Collinge, J.; Brown, J.; Hardy, J.; Mullan, M.; Rossor, M. N.;

[9982] Baker, H.; Crow, T. J.; Lofthouse, R.; Poulter, M.; Ridley, R.; Owen,

[9983] F.; Bennett, C.; Dunn, G.; Harding, A. E.; Quinn, N.; Doshi, B.; Roberts,

[9984] G. W.; Honavar, M.; Janota, I.; Lantos, P. L.: Inherited prion disease

[9985] with 144 base pair gene insertion. 2. Clinical and pathological features. Brain 115:

[9986] 687-710, 1992.

[9987] 123.

[9988] Collinge, J.; Owen, F.; Poulter, M.; Leach, M.; Crow, T. J.;

Rossor,
[9989] M. N.; Hardy, J.; Mullan, M. J.; Janota, I.; Lantos, P. L.: Prion
[9990] dementia without characteristic pathology. Lancet 336: 7–9, 1990.
[9991] 124.
[9992] Collinge, J.; Palmer, M. S.; Dryden, A. J.: Genetic predisposition
[9993] to iatrogenic Creutzfeldt–Jakob disease. Lancet 337: 1441–1442,
[9994] 1991.
[9995] 125.
[9996] Collinge, J.; Poulter, M.; Davis, M. B.; Baraitser, M.; Owen,
[9997] F.; Crow, T. J.; Harding, A. E.: Presymptomatic detection or exclusion
[9998] of prion protein gene defects in families with inherited prion diseases. Am.
[9999] J. Hum. Genet. 49: 1351–1354, 1991.
[10000] 126.
[10001] Collinge, J.; Whittington, M.; Sidle, K. C. L.; Smith, C. J.;
[10002] Palmer, M.; Clarke, A. R.; Jefferys, J. G. R.: Prion protein is

necessary
[10003] for normal synaptic function. (Letter) Nature 370: 295-297, 1994.
[10004] 127.
[10005] Schulz, P.; Stucka, R.; Feldmann, H.; Combriato, G.; Klobeck,
[10006] H.-G.; Fittler, F.: Sequence of a cDNA clone encompassing the complete
[10007] mature human prostate specific antigen (PSA) and an un-spliced leader
[10008] sequence. Nucleic Acids Res. 16: 6226 only, 1988.
[10009] 128.
[10010] Sutherland, G. R.; Baker, E.; Hyland, V. J.; Callen, D. F.; Close,
[10011] J. A.; Tregear, G. W.; Evans, B. A.; Richards, R. I.: Human prostate-specific
[10012] antigen (APS) is a member of the glandular kallikrein gene family
[10013] at 19q13. Cytogenet. Cell Genet. 48: 205-207, 1988.
[10014] 129.
[10015] Bookstein, R.; Lee, E. Y.-H. P.; To, H.; Young, L.-J.; Sery, T.

[10016] W.; Hayes, R. C.; Friedmann, T.; Lee, W.-H.: Human retinoblastoma
[10017] susceptibility gene: genomic organization and analysis of heterozygous
[10018] intragenic deletion mutants. Proc. Nat. Acad. Sci. 85: 2210-2214,
[10019] 1988.

[10020] 130.

[10021] Brantley, M. A.; Worley, L.; Harbour, J. W.: Altered expres-sion
[10022] of Rb and p53 in uveal melanomas following plaque ra-diotherapy. Am.
[10023] J. Ophthal. 133: 242-248, 2002.

[10024] 131.

[10025] Bremner, R.; Du, D. C.; Connolly-Wilson, M. J.; Bridge, P.; Ahmad,
[10026] K. F.; Mostachfi, H.; Rushlow, D.; Dunn, J. M.; Gallie, B. L.: Deletion
[10027] of RB exons 24 and 25 causes low-penetrance retinoblas-toma. Am. J.
[10028] Hum. Genet. 61: 556-570, 1997.

[10029] 132.

[10030] Briard-Guillemot, M. L.; Bonaiti-Pellie, C.; Feingold, J.; Frezal,

[10031] J.: Etude genetique du retinoblastome. Humangenetik 24: 271-284,

[10032] 1974.

[10033] 133.

[10034] Brownstein, S.; de Chadarevian, J.-P.; Little, J. M.: Trilateral

[10035] retinoblastoma: report of two cases. Arch. Ophthal. 102: 257-262,

[10036] 1984.

[10037] 134.

[10038] Buchkovich, K.; Duffy, L. A.; Harlow, E.: The retinoblastoma

[10039] protein is phosphorylated during specific phases of the cell cycle. Cell 58:

[10040] 1097-1105, 1989.

[10041] 135.

[10042] Campos-Xavier, A. B.; Saraiva, J. M.; Savarirayan, R.; Verloes,

[10043] A.; Feingold, J.; Faivre, L.; Munnich, A.; Le Merrer, M.; Cormier-Daire,

[10044] V.: Phenotypic variability at the TGF-beta-1 locus in Camurati-Engelmann

[10045] disease. Hum. Genet. 109: 653-658, 2001.

[10046] 136.

[10047] Aguercif, M.; Nigg, O. M.; Lopez, J. M.; Bouvier, C. A.: Congenital

[10048] factor-XIII-activity deficiency with immunologically characterized

[10049] FSF-like-protein. Nouv. Rev. Franc. Hemat. 11: 841-848, 1971.

[10050] 137.

[10051] Amris, C. J.; Ranek, L.: A case of fibrin-stabilizing factor (FSF)

[10052] deficiency. Thromb. Diath. Haemorrh. 14: 332-340, 1965.

[10053] 138.

[10054] Barbui, T.; Rodeghiero, F.; Dini, E.; Mariani, G.; Papa, M. L.;

[10055] De Biasi, R.; Murillo, R. C.; Umana, C. M.: Subunits A and S inheritance

[10056] in four families with congenital factor XIII deficiency. Brit.

J.

[10057] Haemat. 38: 267-271, 1978.

[10058] 139.

[10059] Berliner, S.; Lusky, A.; Zivelin, A.; Modan, M.; Seligsohn, U.

[10060] : Hereditary factor XIII deficiency: report of four families and definition

[10061] of the carrier state. Brit. J. Haemat. 56: 495-505, 1984.

[10062] 140.

[10063] Board, P.; Coggan, M.; Miloszewski, K.: Identification of a point

[10064] mutation in factor XIII A subunit deficiency. Blood 80: 937-941,

[10065] 1992.

[10066] 141.

[10067] Board, P. G.: Genetic polymorphism of the A subunit of human coagulation

[10068] factor XIII. Am. J. Hum. Genet. 31: 116-124, 1979.

[10069] 142.

[10070] Board, P. G.; Chapple, R.; Coggan, M.: Haplotypes of the coagulation

[10071] factor XIII A subunit locus in normal and deficient subjects. Am.

[10072] J. Hum. Genet. 42: 712-717, 1988.

[10073] 143.

[10074] Beutler, E.: Effect of flavin compounds on glutathione reductase

[10075] activity: in vivo and in vitro studies. J. Clin. Invest. 48: 1957-1966,

[10076] 1969.

[10077] 144.

[10078] Blume, K. G.; Gottwik, M.; Lohr, G. W.; Rudiger, H. W.: Familienuntersuchungen

[10079] zum Glutathionreduktasemangel menschlicher Erythrocyten. Humangenetik 6:

[10080] 163-170, 1968.

[10081] 145.

[10082] Carson, P. E.; Brewer, G. J.; Ickes, C.: Decreased glutathione

[10083] reductase with susceptibility to hemolysis. (Abstract) J. Lab. Clin.

[10084] Med. 58: 804 only, 1961.

[10085] 146.

[10086] de la Chapelle, A.; Icen, A.; Aula, P.; Leisti, J.; Turleau, C.;

[10087] de Grouchy, J.: Mapping of the gene for glutathione re-
ductase on

[10088] chromosome 8. Ann. Genet. 19: 253-256, 1976.

[10089] 147.

[10090] de la Chapelle, A.; Vuopio, P.; Icen, A.: Trisomy 8 in the
bone

[10091] marrow associated with high red cell glutathione reduc-
tase activity. Blood 47:

[10092] 815-826, 1976.

[10093] 148.

[10094] Fajnholc, N. E.; Kaminsky, E.; Machtey, I.; De Vries, A.:
Hereditary

[10095] erythrocyte glutathione reductase deficiency. Rev. Europ.
Clin. Biol. 16:

[10096] 987-991, 1971.

[10097] 149.

[10098] Flatz, G.: Population study of erythrocyte glutathione re-
ductase

[10099] activity. I. Stimulation of the enzyme by flavin adenine dinucleotide
[10100] and by riboflavin supplementation. Humangenetik 11: 269-277, 1971.
[10101] 150.

[10102] Flatz, G.: Population study of erythrocyte glutathione reductase
[10103] activity. II. Hematological data of subjects with low enzyme activity
[10104] and stimulation characteristics in their families. Humangenetik 11:
[10105] 278-285, 1971.

[10106] 151.

[10107] George, D. L.; Francke, U.: Gene dose effect: regional mapping
[10108] of human glutathione reductase on chromosome 8. Cytogenet. Cell Genet. 17:
[10109] 282-286, 1976.

[10110] 152.

[10111] Gutensohn, W.; Rodewald, A.; Haas, B.; Schulz, P.; Cleve, H.:

[10112] Refined mapping of the gene for glutathione reductase on human chromosome

[10113] 8. Hum. Genet. 43: 221-224, 1978.

[10114] 153.

[10115] Hampel, K. E.; Lohr, G. W.; Blume, K. G.; Rudiger, H. W.: Spontane

[10116] und chloramphenicolinduzierte Chromosomenmutationen und biochemische

[10117] Befunde bei zwei Faellen mit Glutathionreduktasemangel (NAD(P)H: glutathione

[10118] oxidoreductase, E.C. 1.6.4.2). Humangenetik 7: 305-313, 1969.

[10119] 154.

[10120] Jensen, P. K. A.; Junien, C.; de la Chapelle, A.: Gene for glutathione

[10121] reductase localized to subband 8p21.1. (Abstract) Cyto-genet. Cell

[10122] Genet. 37: 497 only, 1984.

[10123] 155.

[10124] Jensen, P. K. A.; Junien, C.; Despoisse, S.; Bernsen, A.; Thelle,

[10125] T.; Friedrich, U.; de la Chapelle, A.: Inverted tandem duplication
[10126] of the short arm of chromosome 8: a non-random de novo structural
[10127] aberration in man. Localization of the gene for glutathione reductase
[10128] in subband 8p21.1. Ann. Genet. 25: 207-211, 1982.

[10129] 156.

[10130] Kurz, R.; Hohenwallner, W.: Familiaerer Glutathionreduktasemangel
[10131] und Stoerung der Glutathionsynthese im Erythrozyten. Helv. Paediat.
[10132] Acta 25: 542-552, 1970.

[10133] 157.

[10134] Lohr, G. W.: Personal Communication. Marburg, Germany 1963.
[10135] 158.

[10136] Lohr, G. W.; Waller, H. D.: Eine neue enzymopenische haemolytische
[10137] Anaemie mit Glutathionreduktase-Mangel. Med. Klin. 57: 1521-1525,

[10138] 1962.

[10139] 159.

[10140] Lohr, G. W.; Waller, H. D.: Zur Biochemie einiger angeborener

[10141] haemolytischer Anaemien. Folia Haemat. 8: 377-397, 1963.

[10142] 160.

[10143] Long, W. K.: Glutathione reductase in red blood cells: variant

[10144] associated with gout. Science 155: 712-713, 1967.

[10145] 161.

[10146] Long, W. K.: Personal Communication. Austin, Texas 1972.

[10147] 162.

[10148] Loos, J. A.; Roos, D.; Weening, R. S.; Hauwerzijl, J.: Familial

[10149] deficiency of glutathione reductase in human blood cells. Blood 48:

[10150] 53-62, 1976.

[10151] 163.

[10152] Magenis, R. E.; Reiss, J.; Bigley, R.; Champerlin, J.; Lovrien,

[10153] E.: Exclusion of glutathione reductase from 8pter-8p22 and localization
[10154] to 8p21. Cytogenet. Cell Genet. 22: 446-448, 1978.

[10155] 164.

[10156] Nevin, N. C.; Morrison, P. J.; Jones, J.; Reid, M. M.: Inverted
[10157] tandem duplication of 8p12-p23.1 in a child with increased activity
[10158] of glutathione reductase. J. Med. Genet. 27: 135-136, 1990.

[10159] 165.

[10160] Nichols, E. A.; Ruddle, F. H.: Polymorphism and linkage of glutathione
[10161] reductase in Mus musculus. Biochem. Genet. 13: 323-330, 1975.

[10162] 166.

[10163] Oort, M.; Loos, J. A.; Prins, H. K.: Hereditary absence of reduced
[10164] glutathione in the erythrocytes--a new clinical and biochemical entity. Vox
[10165] Sang. 6: 370-373, 1961.

[10166] 167.

[10167] Roos, D.; Weening, R. S.; Voetman, A. A.; van Schaik, M. L. J.;
[10168] Bot, A. A. M.; Meerhof, L. J.; Loos, J. A.: Protection of phagocytic
[10169] leukocytes by endogenous glutathione: studies in a family with glutathione
[10170] reductase deficiency. Blood 53: 851-866, 1979.

[10171] 168.

[10172] Shows, T. B.; Sakaguchi, A. Y.: Gene transfer and gene mapping
[10173] in mammalian cells in culture. In Vitro 16: 55-76, 1980.

[10174] 169.

[10175] Sinet, P. M.; Bresson, J. L.; Couturier, J.; Prieur, M.; Rethore,
[10176] M.-O.; Taillemite, J. L.; Toudec, D.; Jerome, H.; Lejeune, J.: Localisation
[10177] probable du gene de la glutathion reductase (EC 1.6.4.2.) sur la bande
[10178] 8p21. Ann. Genet. 20: 13-17, 1977.

[10179] 170.

[10180] Staal, G. E. J.; Helleman, P. W.; De Wael, J.; Veeger, C.: Pu-

rification
[10181] and properties of an abnormal glutathione reductase from human erythrocytes. Biochim.
[10182] Biophys. Acta 185: 63-69, 1969.

[10183] 171.

[10184] Chu, F.-F.: The human glutathione peroxidase genes GPX2, GPX3,
[10185] and GPX4 map to chromosomes 14, 5, and 19, respectively. Cytogenet.
[10186] Cell Genet. 66: 96-98, 1994.

[10187] 172.

[10188] Beutler, E.; Matsumoto, F.: Ethnic variation in red cell glutathione
[10189] peroxidase activity. Blood 46: 103-110, 1975.

[10190] 173.

[10191] Beutler, E.; West, C.: Red cell glutathione peroxidase polymorphism
[10192] in Afro-Americans. Am. J. Hum. Genet. 26: 255-258, 1974.
[10193] 174.

[10194] Frasier, S. D.; Bashore, R. A.; Mosier, H. D.: Gonadoblas-

toma
[10195] associated with pure gonadal dysgenesis in monozygotic twins. J.
[10196] Pediat. 64: 740-745, 1964.

[10197] 175.

[10198] Justice, M. J.; Silan, C. M.; Ceci, J. D.; Buchberg, A. M.; Copeland,
[10199] N. G.; Jenkins, N. A.: A molecular genetic linkage map of mouse chromosome
[10200] 13 anchored by the beige (bg) and satin (sa) loci. Genomics 6: 341-351,
[10201] 1990.

[10202] 176.

[10203] Terwindt, G. M.; Ophoff, R. A.; Haan, J.; Frants, R. R.; Ferrari,
[10204] M. D.: Familial hemiplegic migraine: a clinical comparison of families
[10205] linked and unlinked to chromosome 19. Cephalalgia 16: 153-155, 1996.

[10206] 177.

[10207] Seldin, M. F.: Personal Communication. Durham, N. C. 3/13/1989.

[10208] 178.

[10209] Weis, J. H.; Morton, C. C.; Bruns, G. A. P.; Weis, J. J.; Klick-

stein,

[10210] L. B.; Wong, W. W.; Fearon, D. T.: A complement receptor locus: genes

[10211] encoding C3b/C4b receptor and C3d/Epstein-Barr virus receptor map

[10212] to 1q32. J. Immun. 138: 312-315, 1987.

[10213] 179.

[10214] Feifel, E.; Prodinger, W. M.; Molgg, M.; Schwaeble, W.; Schonitzer,

[10215] D.; Koistinen, V.; Misasi, R.; Dierich, M. P.: Polymorphism and deficiency

[10216] of human factor H-related proteins p39 and p37. Im-

munogenetics 36:

[10217] 104-109, 1992.

[10218] 180.

[10219] Meyer, C. G.; Skerka, C.; Zipfel, P. F.: Polymorphism of the

human

[10220] factor H-related gene (FHR-1) and of factor H in a West African individual. Immunogenetics 41:

[10221] 335 only, 1995.

[10222] 181.

[10223] Skerka, C.; Horstmann, R. D.; Zipfel, P. F.: Molecular cloning

[10224] of a human serum protein structurally related to complement factor

[10225] H. J. Biol. Chem. 266: 12015-12020, 1991.

[10226] 182.

[10227] Timmann, C.; Leippe, M.; Horstmann, R. D.: Two major serum components

[10228] antigenically related to complement factor H are different glycosylation

[10229] forms of a single protein with no factor H-like complement regulatory

[10230] functions. J. Immun. 146: 1265-1270, 1991.

[10231] 183.

[10232] Zipfel, P. F.; Skerka, C.: Complement factor H and related proteins:

[10233] an expanding family of complement-regulatory proteins? Immun. Today 15:

[10234] 121-126, 1994.

[10235] 184.

[10236] Wroe, S. F.; Kelsey, G.; Skinner, J. A.; Bodle, D.; Ball, S. T.;

[10237] Beechey, C. V.; Peters, J.; Williamson, C. M.: An imprinted transcript,

[10238] antisense to Nesp, adds complexity to the cluster of imprinted genes

[10239] at the mouse Gnas locus. Proc. Nat. Acad. Sci. 97: 3342-3346, 2000.

[10240] 185.

[10241] Wrong, O.: Tegernsee giant. (Letter) Lancet 339: 194 only,

[10242] 1992.

[10243] 186.

[10244] Yang, I.; Park, S.; Ryu, M.; Woo, J.; Kim, S.; Kim, J.; Kim, Y.;

[10245] Choi, Y.: Characteristics of gsp-positive growth hormone-secreting

[10246] pituitary tumors in Korean acromegalic patients. Europ. J. Endocr. 134:

[10247] 720-726, 1996.

[10248] 187.

[10249] Yu, D.; Yu, S.; Schuster, V.; Kruse, K.; Clericuzio, C. L.; Weinstein,

[10250] L. S.: Identification of two novel deletion mutations within the

[10251] Gs-alpha gene (GNAS1) in Albright hereditary osteodys-

trophy. J. Clin.

[10252] Endocr. Metab. 84: 3254-3259, 1999.

[10253] 188.

[10254] Lerman, M. I.; Minna, J. D.: The 630-kb lung cancer ho-

mozygous

[10255] deletion region on human chromosome 3p21.3: identifi-

cation and evaluation

[10256] of the resident candidate tumor suppressor genes. Cancer Res. 60:

[10257] 6116-6133, 2000.

[10258] 189.

[10259] Carver, M. F. H.; Kutlar, A.: International Hemoglobin In-

formation

[10260] Center: variant list. Hemoglobin 19: 37-149, 1995.

[10261] 190.

[10262] Hobbs, H. H.; Leitersdorf, E.; Leffert, C. C.; Cryer, D. R.; Brown,

[10263] M. S.; Goldstein, J. L.: Evidence for a dominant gene that suppresses

[10264] hypercholesterolemia in a family with defective low density lipoprotein

[10265] receptors. J. Clin. Invest. 84: 656-664, 1989.

[10266] 191.

[10267] Alsaleh, Q. A.; Teebi, A. S.: Autosomal recessive epidermolytic

[10268] palmoplantar keratoderma. J. Med. Genet. 27: 519-522, 1990.

[10269] 192.

[10270] Anderson, I. F.; Klintworth, G. K.: Hypovitaminosis-A in a family

[10271] with tylosis and clinodactyly. Brit. Med. J. 1: 1293-1297, 1961.

[10272] 193.

[10273] Bergstrom, C.: Keratodermia palmaris et plantaris. Nord. Med. 78:

[10274] 155-156, 1967.

[10275] 194.

[10276] Berth-Jones, J.; Hutchinson, P. E.: A family with palmo-plantar

[10277] epidermolytic hyperkeratosis. Clin. Exp. Derm. 14: 313-316, 1989.

[10278] 195.

[10279] Blasik, L. G.; Dimond, R. L.; Baughman, R. D.: Hereditary epidermolytic

[10280] palmoplantar keratoderma. Arch. Derm. 117: 229-231, 1981.

[10281] 196.

[10282] Bonifas, J. M.; Matsumura, K.; Chen, M. A.; Berth-Jones, J.; Hutchinson,

[10283] P. E.; Zloczower, M.; Fritsch, P. O.; Epstein, E. H., Jr.: Muta-tions

[10284] of keratin 9 in two families with palmoplantar epider-molytic hyperkeratosis. J.

[10285] Invest. Derm. 103: 474-477, 1994.

[10286] 197.

[10287] Camisa, C.; Williams, H.: Epidermolytic variant of heredi-

tary
[10288] palmoplantar keratoderma. Brit. J. Derm. 112: 221-225, 1985.
[10289] 198.

[10290] Chung, H.-L.: Keratoma palmare et plantare hereditarium, with
[10291] special reference to its mode of inheritance as traced in six and
[10292] seven generations, respectively, in two Chinese families. Arch. Derm.
[10293] Syph. 36: 303-313, 1937.

[10294] 199.

[10295] Covello, S. P.; Irvine, A. D.; McKenna, K. E.; Munro, C. S.; Nevin,
[10296] N. C.; Smith, F. J. D.; Uitto, J.; McLean, W. H. I.: Mutations in
[10297] keratin K9 in kindreds with epidermolytic palmoplantar keratoderma
[10298] and epidemiology in Northern Ireland. J. Invest. Derm. 111: 1207-1209,
[10299] 1998.

[10300] 200.

[10301] Endo, H.; Hatamochi, A.; Shinkai, H.: A novel mutation of a leucine
[10302] residue in coil 1A of keratin 9 in epidermolytic palmo-plantar keratoderma. J.
[10303] Invest. Derm. 109: 113-115, 1997.

[10304] 201.

[10305] Stead, J. D. H.; Jeffreys, A. J.: Allele diversity and germline
[10306] mutation at the insulin minisatellite. Hum. Molec. Genet. 9: 713-723,
[10307] 2000.

[10308] 202.

[10309] Erneux, C.; Roeckel, N.; Takazawa, K.; Mailleux, P.; Vas-sart, G.;
[10310] Mattei, M. G.: Localization of the genes for human inositol 1,4,5-trisphosphate
[10311] 3-kinase A (ITPKA) and B (ITPKB) to chromosome regions 15q14-q21 and
[10312] 1q41-q43, respectively, by in situ hybridization. Ge-nomics 14:
[10313] 546-547, 1992.

[10314] 203.

[10315] Takazawa, K.; Perret, J.; Dumont, J. E.; Erneux, C.: Human brain

[10316] inositol 1,4,5-trisphosphate 3-kinase cDNA sequence. Nucleic Acids

[10317] Res. 18: 7141 only, 1990.

[10318] 204.

[10319] Takazawa, K.; Perret, J.; Dumont, J. E.; Erneux, C.: Molecular

[10320] cloning and expression of a human brain inositol 1,4,5-trisphosphate

[10321] 3-kinase. Biochem. Biophys. Res. Commun. 174: 529-535, 1991.

[10322] 205.

[10323] Takazawa, K.; Perret, J.; Dumont, J. E.; Erneux, C.: Molecular

[10324] cloning and expression of a new putative inositol 1,4,5-trisphosphate

[10325] 3-kinase isoenzyme. Biochem. J. 278: 883-886, 1991.

[10326] 206.

[10327] 't Hart, L. M.; Stolk, R. P.; Dekker, J. M.; Nijpels, G.; Grobbee,

[10328] D. E.; Heine, R. J.; Maassen, J. A.: Prevalence of variants in candidate
[10329] genes for type 2 diabetes mellitus in the Netherlands: the Rotterdam
[10330] study and the Hoorn study. J. Clin. Endocr. Metab. 84: 1002-1006,
[10331] 1999.

[10332] 207.

[10333] Abe, H.; Yamada, N.; Kamata, K.; Kuwaki, T.; Shimada, M.; Osuga,
[10334] J.; Shionoiri, F.; Yahagi, N.; Kadowaki, T.; Tamemoto, H.; Ishibashi,
[10335] S.; Yazaki, Y.; Makuuchi, M.: Hypertension, hypertriglyc- eridemia,
[10336] and impaired endothelium-dependent vascular relaxation in mice lacking
[10337] insulin receptor substrate-1. J. Clin. Invest. 101: 1784-1788, 1998.
[10338] 208.

[10339] Almind, K.; Bjorbaek, C.; Vestergaard, H.; Hansen, T.; Ech- wald,
[10340] S.; Pedersen, O.: Amino acid polymorphisms of insulin re-

ceptor substrate-1
[10341] in non-insulin-dependent diabetes mellitus. Lancet 342: 828-832,
[10342] 1993.

[10343] 209.

[10344] Almind, K.; Inoue, G.; Pedersen, O.; Kahn, C. R.: A com-mon amino
[10345] acid polymorphism in insulin receptor substrate-1 causes impaired
[10346] insulin signaling: evidence from transfection studies. J. Clin. Invest. 97:
[10347] 2569-2575, 1996.

[10348] 210.

[10349] Araki, E.; Sun, X.-J.; Haag, B. L., III; Chuang, L.-M.; Zhang,
[10350] Y.; Yang-Feng, T. L.; White, M. F.; Kahn, C. R.: Human skeletal muscle
[10351] insulin receptor substrate-1: characterization of the cDNA, gene,
[10352] and chromosomal localization. Diabetes 42: 1041-1054, 1993.
[10353] 211.

[10354] Baroni, M. G.; D'Andrea, M. P.; Montali, A.; Pannitteri, G.; Barilla,
[10355] F.; Campagna, F.; Mazzei, E.; Lovari, S.; Seccareccia, F.; Campa,
[10356] P. P.; Ricci, G.; Pozzilli, P.; Urbinati, G.; Arca, M.: A com-
mon
[10357] mutation of the insulin receptor substrate-1 gene is a risk factor
[10358] for coronary artery disease. Arterioscler. Thromb. Vasc. Biol. 19:
[10359] 2975-2980, 1999.

[10360] 212.

[10361] Bohni, R.; Riesgo-Escovar, J.; Oldham, S.; Brogiolo, W.; Stocker,
[10362] H.; Andruss, B. F.; Beckingham, K.; Hafen, E.: Autonomous control
[10363] of cell and organ size by CHICO, a Drosophila homolog of vertebrate
[10364] IRS1-4. Cell 97: 865-875, 1999.

[10365] 213.

[10366] Katzke, S.; Booms, P.; Tiecke, F.; Palz, M.; Pletschacher, A.;

[10367] Turkmen, S.; Neumann, L. M.; Pregla, R.; Leitner, C.; Schramm, C.;

[10368] Lorenz, P.; Hagemeier, C.; Fuchs, J.; Skovby, F.; Rosenberg, T.; Robinson,

[10369] P. N.: TGGE screening of the entire FBN1 coding sequence in 126 individuals

[10370] with Marfan syndrome and related fibrillinopathies. Hum. Mutat. 20:

[10371] 197-208, 2002.

[10372] 214.

[10373] Kielty, C. M.; Raghunath, M.; Siracusa, L. D.; Sherratt, M. J.;

[10374] Peters, R.; Shuttleworth, C. A.; Jimenez, S. A.: The tight skin mouse:

[10375] demonstration of mutant fibrillin-1 production and assembly into abnormal

[10376] microfibrils. J. Cell Biol. 140: 1159-1166, 1998.

[10377] 215.

[10378] Kilpatrick, M. W.; Phylactou, L. A.; Godfrey, M.; Wu, C. H.; Wu,

[10379] G. Y.; Tsipouras, P.: Delivery of a hammerhead ribozyme specifically

[10380] down-regulates the production of fibrillin-1 by cultured dermal fibroblasts. Hum.

[10381] Molec. Genet. 5: 1939-1944, 1996.

[10382] 216.

[10383] Lee, B.; Godfrey, M.; Vitale, E.; Hori, H.; Mattei, M. G.; Sar-

farazi,

[10384] M.; Tsipouras, P.; Ramirez, F.; Hollister, D. W.: Linkage of Marfan

[10385] syndrome and a phenotypically related disorder to two different fibrillin

[10386] genes. Nature 352: 330-334, 1991.

[10387] 217.

[10388] Liu, W.; Qian, C.; Comeau, K.; Brenn, T.; Furthmayr, H.; Francke,

[10389] U.: Mutant fibrillin-1 monomers lacking EGF-like domains disrupt

[10390] microfibril assembly and cause severe Marfan syndrome. Hum. Molec.

[10391] Genet. 5: 1581-1587, 1996.

[10392] 218.

[10393] Liu, W.; Qian, C.; Francke, U.: Silent mutation induces

exon

[10394] skipping of fibrillin-1 gene in Marfan syndrome. (Letter) Nature

[10395] Genet. 16: 328-329, 1997.

[10396] 219.

[10397] Lonnqvist, L.; Child, A.; Kainulainen, K.; Davidson, R.; Puhakka,

[10398] L.; Peltonen, L.: A novel mutation of the fibrillin gene causing

[10399] ectopia lentis. Genomics 19: 573-576, 1994.

[10400] 220.

[10401] Lonnqvist, L.; Reinhardt, D.; Sakai, L.; Peltonen, L.: Evi-dence

[10402] for furin-type activity-mediated C-terminal processing of profibrillin-1

[10403] and interference in the processing by certain mutations. Hum. Molec.

[10404] Genet. 7: 2039-2044, 1998.

[10405] 221.

[10406] Magenis, R. E.; Maslen, C. L.; Smith, L.; Allen, L.; Sakai, L.

[10407] Y.: Localization of the fibrillin (FBN) gene to chromosome

15, band

[10408] q21.1. Genomics 11: 346-351, 1991.

[10409] 222.

[10410] Maquat, L. E.: NASty effects on fibrillin pre-mRNA splicing:

[10411] another case of ESE does it, but proposals for translation-dependent

[10412] splice site choice live on. Genes Dev. 16: 1743-1753, 2002.

[10413] 223.

[10414] Maslen, C. L.; Corson, G. M.; Maddox, B. K.; Glanville, R. W.;

[10415] Sakai, L. Y.: Partial sequence of a candidate gene for the Marfan

[10416] syndrome. Nature 352: 334-337, 1991.

[10417] 224.

[10418] Matyas, G.; De Paepe, A.; Halliday, D.; Boileau, C.; Pals, G.;

[10419] Steinmann, B.: Evaluation and application of denaturing HPLC for

[10420] mutation detection in Marfan syndrome: identification of 20 novel

[10421] mutations and two novel polymorphisms in the FBN1

gene. Hum. Mutat. 19:
[10422] 443-456, 2002.

[10423] 225.

[10424] McGettrick, A. J.; Knott, V.; Willis, A.; Handford, P. A.: Molecular
[10425] effects of calcium binding mutations in Marfan syndrome depend on
[10426] domain context. Hum. Molec. Genet. 9: 1987-1994, 2000.

[10427] 226.

[10428] Milewicz, D. M.; Grossfield, J.; Cao, S.-N.; Kielty, C.; Covitz,
[10429] W.; Jewett, T.: A mutation in FBN1 disrupts profibrillin processing
[10430] and results in isolated skeletal features of the Marfan syn-
drome. J.
[10431] Clin. Invest. 95: 2373-2378, 1995.

[10432] 227.

[10433] Milewicz, D. M.; Pyeritz, R. E.; Crawford, E. S.; Byers, P. H.

[10434] : Marfan syndrome: defective synthesis, secretion, and ex-
tracellular
[10435] matrix formation of fibrillin by cultured dermal fibrob-

lasts. J.

[10436] Clin. Invest. 89: 79-86, 1992.

[10437] 228.

[10438] Montgomery, R. A.; Dietz, H. C.: Inhibition of fibrillin 1 expression

[10439] using U1 snRNA as a vehicle for the presentation of anti-sense targeting

[10440] sequence. Hum. Molec. Genet. 6: 519-525, 1997.

[10441] 229.

[10442] Montgomery, R. A.; Geraghty, M. T.; Bull, E.; Gelb, B. D.; Johnson,

[10443] M.; McIntosh, I.; Francomano, C. A.; Dietz, H. C.: Multiple molecular

[10444] mechanisms underlying subdiagnostic variants of Marfan syndrome. Am.

[10445] J. Hum. Genet. 63: 1703-1711, 1998.

[10446] 230.

[10447] Nijbroek, G.; Sood, S.; McIntosh, I.; Francomano, C. A.; Bull,

[10448] E.; Pereira, L.; Ramirez, F.; Pyeritz, R. E.; Dietz, H. C.: Fif-teen

[10449] novel FBN1 mutations causing Marfan syndrome detected by heteroduplex
[10450] analysis of genomic amplicons. Am. J. Hum. Genet. 57: 8-21, 1995.
[10451] 231.

[10452] Pereira, L.; Andrikopoulos, K.; Tian, J.; Lee, S. Y.; Keene, D.

[10453] R.; Ono, R.; Reinhardt, D. P.; Sakai, L. Y.; Biery, N. J.; Bunton,
[10454] T.; Dietz, H. C.; Ramirez, F.: Targetting of the gene encoding fibrillin-1
[10455] recapitulates the vascular aspect of Marfan syndrome. Nature Genet. 17:
[10456] 218-222, 1997.

[10457] 232.

[10458] Pereira, L.; D'Alessio, M.; Ramirez, F.; Lynch, J. R.; Sykes,

[10459] B.; Pangilinan, T.; Bonadio, J.: Genomic organization of the sequence
[10460] coding for fibrillin, the defective gene product in Marfan syndrome. Hum.
[10461] Molec. Genet. 2: 961-968, 1993.

[10462] 233.

[10463] Raghunath, M.; Superti-Furga, A.; Godfrey, M.; Steinmann, B.:

[10464] Decreased extracellular deposition of fibrillin and decorin in neonatal

[10465] Marfan syndrome fibroblasts. Hum. Genet. 90: 511-515, 1993.

[10466] 234.

[10467] Palz, M.; Tiecke, F.; Booms, P.; Goldner, B.; Rosenberg, T.; Fuchs,

[10468] J.; Skovby, F.; Schumacher, H.; Kaufmann, U. C.; von Kodolitsch, Y.;

[10469] Nienaber, C. A.; Leitner, C.; Katzke, S.; Vetter B.; Hagemeier, C.;

[10470] Robinson, P. N.: Clustering of mutations associated with mild Marfan-like

[10471] phenotypes in the 3-prime region of FBN1 suggests a potential genotype-phenotype

[10472] correlation. Am. J. Med. Genet. 91: 212-221, 2000.

[10473] 235.

[10474] Rantamaki, T.; Raghunath, M.; Karttunen, L.; Lonnqvist, L.; Child,

[10475] A.; Peltonen, L.: Prenatal diagnosis of Marfan syndrome:

identification
[10476] of a fibrillin-1 mutation in chorionic villus sample. Prenatal Diag. 15:
[10477] 1176-1181, 1995.

[10478] 236.

[10479] Reinhardt, D. P.; Ono, R. N.; Notbohm, H.; Muller, P. K.; Bachinger,
[10480] H. P.; Sakai, L. Y.: Mutations in calcium-binding epidermal growth
[10481] factor modules render fibrillin-1 susceptible to proteolysis: a potential
[10482] disease-causing mechanism in Marfan syndrome. J. Biol. Chem. 275:
[10483] 12339-12345, 2000.

[10484] 237.

[10485] Robinson, P. N.; Booms, P.; Katzke, S.; Ladewig, M.; Neumann,
[10486] L.; Palz, M.; Pregla, R.; Tiecke, F.; Rosenberg, T.: Mutations of
[10487] FBN1 and genotype-phenotype correlations in Marfan syndrome and related
[10488] fibrillinopathies. Hum. Mutat. 20: 153-161, 2002.

[10489] 238.

[10490] Robinson, P. N.; Godfrey, M.: The molecular genetics of Marfan

[10491] syndrome and related microfibrillopathies. J. Med. Genet. 37: 9-25,

[10492] 2000.

[10493] 239.

[10494] Sakai, L. Y.; Keene, D. R.; Engvall, E.: Fibrillin, a new 350-kD

[10495] glycoprotein, is a component of extracellular microfibrils. J. Cell

[10496] Biol. 103: 2499-2509, 1986.

[10497] 240.

[10498] Fletcher, W. H.; Britz-Cunningham, S. H.; Zuppan, C. W.: Connexin43

[10499] mutations in sporadic and familial defects of laterality. (Letter) New

[10500] Eng. J. Med. 333: 941-942, 1995.

[10501] 241.

[10502] Gebbia, M.; Towbin, J. A.; Casey, B.: Failure to detect con-nexin43

[10503] mutations in 38 cases of sporadic and familial heterotaxy. Circulation 94:
[10504] 1909-1912, 1996.

[10505] 242.

[10506] Gebbia, M.; Towbin, J. A.; Casey, B.: Connexin43 gene mutations
[10507] and heterotaxy. Response. (Letter) Circulation 97: 118 only, 1998.
[10508] 243.

[10509] Guerrero, P. A.; Schuessler, R. B.; Davis, L. M.; Beyer, E. C.;

[10510] Johnson, C. M.; Yamada, K. A.; Saffitz, J. E.: Slow ventricular conduction
[10511] in mice heterozygous for a connexin43 null mutation. J. Clin. Invest. 99:
[10512] 1991-1998, 1997.

[10513] 244.

[10514] Li, J.-Y.; Hou, X.-E.; Dahlstrom, A.: GAP-43 and its relation
[10515] to autonomic and sensory neurons in sciatic nerve and gastrocnemius
[10516] muscle in the rat. J. Auton. Nerv. Syst. 50: 299-309, 1995.

[10517] 245.

[10518] Liao, Y.; Day, K. H.; Damon, D. N.; Duling, B. R.: Endothe-
lial
[10519] cell-specific knockout of connexin 43 causes hypotension
and bradycardia
[10520] in mice. Proc. Nat. Acad. Sci. 98: 9989-9994, 2001.

[10521] 246.

[10522] Liu, X. Z.; Xia, X. J.; Adams, J.; Chen, Z. Y.; Welch, K. O.;

[10523] Tekin, M.; Ouyang, X. M.; Kristiansen, A.; Pandya, A.;
Balkany, T.;
[10524] Arnos, K. S.; Nance, W. E.: Mutations in GJA1 (connexin
43) are associated
[10525] with non-syndromic autosomal recessive deafness. Hum.
Molec. Genet. 10:
[10526] 2945-2951, 2001.

[10527] 247.

[10528] Reaume, A. G.; de Sousa, P. A.; Kulkarni, S.; Langille, B. L.;

[10529] Zhu, D.; Davies, T. C.; Juneja, S. C.; Kidder, G. M.; Rossant,
J.
[10530] : Cardiac malformation in neonatal mice lacking con-
nexin43. Science 267:

[10531] 1831-1834, 1995.

[10532] 248.

[10533] Splitt, M. P.; Burn, J.; Goodship, J.: Connexin43 mutations in

[10534] sporadic and familial defects of laterality. (Letter) New Eng. J.

[10535] Med. 333: 941, 1995.

[10536] 249.

[10537] Splitt, M. P.; Tsai, M. Y.; Burn, J.; Goodship, J. A.: Absence

[10538] of mutations in the regulatory domain of the gap junction protein

[10539] connexin 43 in patients with visceroatrial heterotaxy. Heart 77:

[10540] 369-370, 1997.

[10541] 250.

[10542] Toth, T.; Hajdu, J.; Marton, T.; Nagy, B.; Papp, Z.: Con-nexin43

[10543] gene mutations and heterotaxy. (Letter) Circulation 97: 117-118,

[10544] 1998.

[10545] 251.

[10546] Ya, J.; Erdstieck-Ernste, E. B. H. W.; de Boer, P. A. J.; van

[10547] Kempen, M. J. A.; Jongsma, H.; Gros, D.; Moorman, A. F. M.; Lamers,

[10548] W. H.: Heart defects in connexin43-deficient mice. Circ. Res. 82:

[10549] 360-366, 1998.

[10550] 252.

[10551] Parker, D.; Ferreri, K.; Nakajima, T.; LaMorte, V. J.; Evans,

[10552] R.; Koerber, S. C.; Hoeger, C.; Montminy, M. R.: Phospho-rylation

[10553] of CREB at ser-133 induces complex formation with CREB-binding protein

[10554] via a direct mechanism. Molec. Cell. Biol. 16: 694-703, 1996.

[10555] 253.

[10556] Radhakrishnan, I.; Perez-Alvarado, G. C.; Parker, D.; Dyson, H.

[10557] J.; Montminy, M. R.; Wright, P. E.: Solution structure of the KIX

[10558] domain of CBP bound to the transactivation domain of

CREB: a model
[10559] for activator:coactivator interactions. Cell 91: 741-752, 1997.
[10560] 254.
[10561] Dhallan, R. S.; Macke, J.; Eddy, R. L.; Shows, T. B.; Reed, R.
[10562] R.; Yau, K.-W.; Nathans, J.: The human rod photoreceptor cyclic GMP
[10563] gated channel gene maps to chromosome 4. (Abstract) Cytogenet. Cell
[10564] Genet. 58: 1886, 1991.
[10565] 255.
[10566] Dhallan, R. S.; Macke, J. P.; Eddy, R. L.; Shows, T. B.; Reed,
[10567] R. R.; Yau, K.-W.; Nathans, J.: Human rod photoreceptor cGMP-gated
[10568] channel: amino acid sequence, gene structure, and functional expression. J.
[10569] Neurosci. 12: 3248-3256, 1992.
[10570] 256.
[10571] Dryja, T. P.; Finn, J. T.; Peng, Y.-W.; McGee, T. L.; Berson, E.
[10572] L.; Yau, K.-W.: Mutations in the gene encoding the alpha

subunit
[10573] of the rod cGMP-gated channel in autosomal recessive retinitis pigmentosa. Proc.
[10574] Nat. Acad. Sci. 92: 10177-10181, 1995.

[10575] 257.

[10576] Griffin, C. A.; Ding, C. L.; Jabs, E. W.; Hawkins, A. L.; Li, X.;

[10577] Levine, M. A.: Human rod cGMP-gated cation channel gene maps to 4p12-to-centromere
[10578] by chromosomal in situ hybridization. Genomics 16: 302-303, 1993.
[10579] 258.

[10580] Kaupp, U. B.; Niidome, T.; Tanabe, T.; Terada, S.; Bonigk, W.;
[10581] Stuhmer, W.; Cook, N. J.; Kangawa, K.; Matsuo, H.; Hirose, T.; Miyata,
[10582] T.; Numa, S.: Primary structure and functional expression from complementary
[10583] DNA of the rod photoreceptor cyclic GMP-gated channel. Nature 342:
[10584] 762-766, 1989.

[10585] 259.

[10586] Pittler, S. J.; Lee, A. K.; Altherr, M. R.; Howard, T. A.; Seldin,
[10587] M. F.; Hurwitz, R. L.; Wasmuth, J. J.; Baehr, W.: Primary structure
[10588] and chromosomal localization of human and mouse rod photoreceptor
[10589] cGMP-gated cation channel. J. Biol. Chem. 267: 6257-6262, 1992.
[10590] 260.

[10591] Meyerson, M.; Enders, G. H.; Wu, C.-L.; Su, L.-K.; Gorka, C.; Nelson,
[10592] C.; Harlow, E.; Tsai, L.-H.: A family of human cdc2-related protein
[10593] kinases. EMBO J. 11: 2909-2917, 1992.

[10594] 261.

[10595] Demetrick, D. J.; Zhang, H.; Beach, D. H.: Chromosomal mapping
[10596] of human CDK2, CDK4, and CDK5 cell cycle kinase genes. Cytogenet.
[10597] Cell Genet. 66: 72-74, 1994.

[10598] 262.

[10599] Harbour, J. W.; Luo, R. X.; Dei Santi, A.; Postigo, A. A.; Dean,
[10600] D. C.: Cdk phosphorylation triggers sequential intramolecular interactions
[10601] that progressively block Rb functions as cells move through G1. Cell 98:
[10602] 859-869, 1999.

[10603] 263.

[10604] Mitchell, E. L. D.; White, G. R. M.; Santibanez-Koref, M. F.; Varley,
[10605] J. M.; Heighway, J.: Mapping of gene loci in the q13-q15 region of
[10606] chromosome 12. Chromosome Res. 3: 261-262, 1995.

[10607] 264.

[10608] Modiano, J. F.; Mayor, J.; Ball, C.; Fuentes, M. K.; Linthicum,
[10609] D. S.: CDK4 expression and activity are required for cytokine responsiveness
[10610] in T cells. J. Immun. 165: 6693-6702, 2000.

[10611] 265.

[10612] Bullrich, F.; MacLachlan, T. K.; Sang, N.; Druck, T.;

Veronese,

[10613] M. L.; Allen, S. L.; Chiorazzi, N.; Koff, A.; Heubner, K.; Croce,

[10614] C. M.; Giordano, A.: Chromosomal mapping of members of the cdc2 family

[10615] of protein kinases, cdk3, cdk6, PISSLRE, and PITALRE, and a cdk inhibitor,

[10616] p27-Kip1, to regions involved in human cancer. Cancer Res. 55: 1199-1205,

[10617] 1995.

[10618] 266.

[10619] Yoshida, N.; Imaoka, S.; Hirata, H.; Matsuda, M.; Asakura, S.

[10620] : Heterozygous abnormal fibrinogen Osaka III with the re-placement

[10621] of gamma-arginine-275 by histidine has an apparently higher molecular

[10622] weight gamma-chain variant. Thromb. Haemost. 68: 534-538, 1992.

[10623] 267.

[10624] Birnbaum, R. A.; O'Marcaigh, A.; Wardak, Z.; Zhang, Y.-Y.; Dranoff,

[10625] G.; Jacks, T.; Clapp, D. W.; Shannon, K. M.: Nf1 and Gmcsf interact

[10626] in myeloid leukemogenesis. Molec. Cell 5: 189-195, 2000.

[10627] 268.

[10628] Cantrell, M. A.; Anderson, D.; Cerretti, D. P.; Price, V.; McKereghan,

[10629] K.; Tushinski, R. J.; Mochizuki, D. Y.; Larsen, A.; Grabstein, K.;

[10630] Gillis, S.; Cosman, D.: Cloning, sequence, and expression of a human

[10631] granulocyte/macrophage colony-stimulating factor. Proc. Nat. Acad.

[10632] Sci. 82: 6250-6254, 1985.

[10633] 269.

[10634] Frolova, E. I.; Dolganov, G. M.; Mazo, I. A.; Smirnov, D. V.; Copeland,

[10635] P.; Stewart, C.; O'Brien, S. J.; Dean, M.: Linkage mapping of the

[10636] human CSF2 and IL3 genes. Proc. Nat. Acad. Sci. 88: 4821-4824, 1991.

[10637] 270.

[10638] Grabstein, K. H.; Urdal, D. L.; Tushinski, R. J.; Mochizuki, D.
[10639] Y.; Price, V. L.; Cantrell, M. A.; Gillis, S.; Conlon, P. J.: Induction
[10640] of macrophage tumoricidal activity by granulocyte-macrophage colony-stimulating
[10641] factor. Science 232: 506-508, 1986.

[10642] 271.

[10643] Huebner, K.; Isobe, M.; Croce, C. M.; Golde, D. W.; Kaufman, S.
[10644] E.; Gasson, J. C.: The human gene encoding GM-CSF is at 5q21-q32,
[10645] the chromosome region deleted in the 5q- anomaly. Science 230: 1282-1285,
[10646] 1985.

[10647] 272.

[10648] Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Larson, R. A.; Rowley,
[10649] J. D.; Gasson, J. C.; Golde, D. W.; Sherr, C. J.: Evidence for the
[10650] involvement of GM-CSF and FMS in the deletion (5q) in myeloid disorders. Science 231:

[10651] 984-987, 1986.

[10652] 273.

[10653] LeVine, A. M.; Reed, J. A.; Kurak, K. E.; Cianciolo, E.; Whit-
sett,

[10654] J. A.: GM-CSF-deficient mice are susceptible to pulmonary
group B

[10655] streptococcal infection. J. Clin. Invest. 103: 563-569,
1999.

[10656] 274.

[10657] Metcalf, D.: The molecular biology and functions of the
granulocyte-macrophage

[10658] colony-stimulating factors. Blood 67: 257-267, 1986.

[10659] 275.

[10660] Pettenati, M. J.; Le Beau, M. M.; Lemons, R. S.; Shima, E. A.;

[10661] Kawasaki, E. S.; Larson, R. A.; Sherr, C. J.; Diaz, M. O.;
Rowley,

[10662] J. D.: Assignment of CSF-1 to 5q33.1: evidence for clus-
tering of

[10663] genes regulating hematopoiesis and for their involvement
in the deletion

[10664] of the long arm of chromosome 5 in myeloid disorders.

Proc. Nat.

[10665] Acad. Sci. 84: 2970-2974, 1987.

[10666] 276.

[10667] Sieff, C. A.; Emerson, S. G.; Donahue, R. E.; Nathan, D. G.; Wang,

[10668] E. A.; Wong, G. G.; Clark, S. C.: Human recombinant granulocyte-macrophage

[10669] colony-stimulating factor: a multilineage hematopoietin. Science 230:

[10670] 1171-1173, 1985.

[10671] 277.

[10672] Thangavelu, M.; Neuman, W. L.; Espinosa, R., III; Nakamura, Y.;

[10673] Westbrook, C. A.; Le Beau, M. M.: A physical and genetic linkage

[10674] map of the distal long arm of human chromosome 5. Cytogenet. Cell

[10675] Genet. 59: 27-30, 1992.

[10676] 278.

[10677] Yang, Y.-C.; Kovacic, S.; Kriz, R.; Wolf, S.; Clark, S. C.; Wellems,

[10678] T. E.; Nienhuis, A.; Epstein, N.: The human genes for GM-CSF and

[10679] IL3 are closely linked in tandem on chromosome 5. Blood 71: 958-961,

[10680] 1988.

[10681] 279.

[10682] Modi, W. S.; Chen, Z.-Q.: Localization of the human CXC chemokine

[10683] subfamily on the long arm of chromosome 4 using radiation hybrids. Genomics 47:

[10684] 136-139, 1998.

[10685] 280.

[10686] Proost, P.; Wuyts, A.; Conings, R.; Lenaerts, J.-P.; Billiau, A.;

[10687] Opdenakker, G.; Van Damme, J.: Human and bovine granulocyte chemotactic

[10688] protein-2: complete amino acid sequence and functional characterization

[10689] as chemokines. Biochemistry 32: 10170-10177, 1993.

[10690] 281.

[10691] Wong, G. G.; Witek, J. S.; Temple, P. A.; Wilkens, K. M.;

Leary,

[10692] A. C.; Luxenberg, D. P.; Jones, S. S.; Brown, E. L.; Kay, R. M.; Orr,

[10693] E. C.; Shoemaker, C.; Golde, D. W.; Kaufman, R. J.; Hewick, R. M.;

[10694] Wang, E. A.; Clark, S. C.: Human GM-CSF: molecular cloning of the

[10695] complementary DNA and purification of the natural and recombinant

[10696] proteins. Science 228: 810-815, 1985.

[10697] 282.

[10698] Rovai, L. E.; Herschman, H. R.; Smith, J. B.: Cloning and characterization

[10699] of the human granulocyte chemotactic protein-2 gene. J. Immun. 158:

[10700] 5257-5266, 1997.

[10701] 283.

[10702] Wuyts, A.; van Osselaer, N.; Haelens, A.; Samson, I.; Herdewijn,

[10703] P.; Ben-Baruch, A.; Oppenheim, J. J.; Proost, P.; van Damme, J.:

[10704] Characterization of synthetic human granulocyte chemo-

tactic protein
[10705] 2: usage of chemokine receptors CXCR1 and CXCR2 and in vivo inflammatory
[10706] properties. Biochemistry 36: 2716-2723, 1997.

[10707] 284.

[10708] Kanda, N.; Fukushige, S.; Murotsu, T.; Yoshida, M. C.; Tsuchiya,
[10709] M.; Asano, S.; Kaziro, Y.; Nagata, S.: Human gene coding for granulocyte-colony
[10710] stimulating factor assigned to the q21-q22 region of chromosome 17. Somat.
[10711] Cell Molec. Genet. 13: 679-684, 1987.

[10712] 285.

[10713] Yoshida, N.; Okuma, M.; Moroi, M.; Matsuda, M.: A lower molecular
[10714] weight gamma-chain variant in a congenital abnormal fibrinogen (Kyoto). Blood 68:
[10715] 703-707, 1986.

[10716] 286.

[10717] Yoshida, N.; Ota, K.; Moroi, M.; Matsuda, M.: An apparently higher

[10718] molecular weight gamma-chain variant in a new congenital abnormal

[10719] fibrinogen Tochigi characterized by the replacement of gamma arginine-275

[10720] by cysteine. Blood 71: 480-487, 1988.

[10721] 287.

[10722] Yoshida, N.; Terukina, S.; Okuma, M.; Moroi, M.; Aoki, N.; Matsuda,

[10723] M.: Characterization of an apparently lower molecular weight gamma-chain

[10724] variant in fibrinogen Kyoto I: the replacement of gamma-asparagine

[10725] 308 by lysine which causes accelerated cleavage of fragment D(1) by

[10726] plasmin and the generation of a new plasmin cleavage site. J. Biol.

[10727] Chem. 263: 13848-13856, 1988.

[10728] 288.

[10729] Abraham, J. A.; Mergia, A.; Whang, J. L.; Tumolo, A.; Friedman,

[10730] J.; Hjerild, K. A.; Gospodarowicz, D.; Fiddes, J. C.: Nucleotide

[10731] sequence of a bovine clone encoding the angiogenic protein, basic
[10732] fibroblast growth factor. Science 233: 545-548, 1986.

[10733] 289.

[10734] Abraham, J. A.; Whang, J. L.; Tumolo, A.; Mergia, A.; Friedman,
[10735] J.; Gospodarowicz, D.; Fiddes, J. C.: Human basic fibroblast growth
[10736] factor: nucleotide sequence and genomic organization. EMBO J. 5:
[10737] 2523-2528, 1986.

[10738] 290.

[10739] Doniach, T.: Basic FGF as an inducer of anteroposterior neural
[10740] pattern. Cell 83: 1067-1070, 1995.

[10741] 291.

[10742] Dono, R.; Texido, G.; Dussel, R.; Ehmke, H.; Zeller, R.: Impaired
[10743] cerebral cortex development and blood pressure regulation in FGF2-deficient
[10744] mice. EMBO J. 17: 4213-4225, 1998.

[10745] 292.

[10746] Fukushima, Y.; Byers, M. G.; Fiddes, J. C.; Shows, T. B.: The

[10747] human basic fibroblast growth factor gene (FGFB) is assigned to chromosome

[10748] 4q25. Cytogenet. Cell Genet. 54: 159-160, 1990.

[10749] 293.

[10750] Gritti, A.; Parati, E. A.; Cova, L.; Frolichsthal, P.; Galli, R.;

[10751] Wanke, E.; Faravelli, L.; Morassutti, D. J.; Roisen, F.; Nickel, D.

[10752] D.; Vescovi, A. L.: Multipotential stem cells from the adult mouse

[10753] brain proliferate and self-renew in response to basic fibroblast growth

[10754] factor. J. Neurosci. 16: 1091-1100, 1996.

[10755] 294.

[10756] Kawaguchi, H.; Nakamura, K.; Tabata, Y.; Ikada, Y.; Aoyama, I.;

[10757] Anzai, J.; Nakamura, T.; Hiyama, Y.; Tamura, M.: Acceleration of

[10758] fracture healing in nonhuman primates by fibroblast growth factor-2. J.

[10759] Clin. Endocr. Metab. 86: 875-880, 2001.

[10760] 295.

[10761] Kurokawa, T.; Sasada, R.; Iwane, M.; Igarashi, K.: Cloning and

[10762] expression of cDNA encoding human basic fibroblast growth factor. FEBS

[10763] Lett. 213: 189-194, 1987.

[10764] 296.

[10765] Lafage-Pochitaloff, M.; Galland, F.; Simonetti, J.; Prats, H.;

[10766] Mattei, M.-G.; Birnbaum, D.: The human basic fibroblast growth factor

[10767] gene is located on the long arm of chromosome 4 at bands q26-q27. Oncogene

[10768] Res. 5: 241-244, 1990.

[10769] 297.

[10770] Mattei, M.-G.; Pebusque, M.-J.; Birnbaum, D.: Chromoso-mal localizations

[10771] of mouse Fgf2 and Fgf5 genes. Mammalian Genome 2: 135-137, 1992.

[10772] 298.

[10773] Montero, A.; Okada, Y.; Tomita, M.; Ito, M.; Tsurukami, H.; Nakamura,
[10774] T.; Doetschman, T.; Coffin, J. D.; Hurley, M. M.: Disruption of the
[10775] fibroblast growth factor-2 gene results in decreased bone mass and
[10776] bone formation. J. Clin. Invest. 105: 1085-1093, 2000.

[10777] 299.

[10778] Ortega, S.; Ittmann, M.; Tsang, S. H.; Ehrlich, M.; Basilico,

[10779] C.: Neuronal defects and delayed wound healing in mice lacking fibroblast
[10780] growth factor 2. Proc. Nat. Acad. Sci. 95: 5672-5677, 1998.
[10781] 300.

[10782] Plotnikov, A. N.; Schlessinger, J.; Hubbard, S. R.; Mohammadi,
[10783] M.: Structural basis for FGF receptor dimerization and activation. Cell 98:
[10784] 641-650, 1999.

[10785] 301.

[10786] Aerssens, J.; Chaffanet, M.; Baens, M.; Matthijs, G.; Van

Den Berghe,
[10787] H.; Cassiman, J.-J.; Marynen, P.: Regional assignment of seven loci
[10788] to 12p13.2-pter by PCR analysis of somatic cell hybrids containing
[10789] the der(12) or the der(X) chromosome from a mesothe-lioma showing t(X;12)(q22;p13). Genomics 20:
[10790] 119-121, 1994.

[10791] 302.

[10792] Angle, B.; Hersh, J. H.; Christensen, K. M.: Molecularly proven
[10793] hypochondroplasia with cloverleaf skull deformity: a novel association. Clin.
[10794] Genet. 54: 417-420, 1998.

[10795] 303.

[10796] Avraham, K. B.; Givol, D.; Avivi, A.; Yayon, A.; Copeland, N. G.;
[10797] Jenkins, N. A.: Mapping of murine fibroblast growth factor receptors
[10798] refines regions of homology between mouse and human chromosomes. Genomics 21:
[10799] 656-658, 1994.

[10800] 304.

[10801] Bellus, G. A.; Gaudenz, K.; Zackai, E. H.; Clarke, L. A.; Sz-
abo,
[10802] J.; Francomano, C. A.; Muenke, M.: Identical mutations in
three different
[10803] fibroblast growth factor receptor genes in autosomal
dominant craniosynostosis
[10804] syndromes. Nature Genet. 14: 174-176, 1996.

[10805] 305.

[10806] Bellus, G. A.; Hefferon, T. W.; Ortiz de Luna, R. I.; Hecht, J.

[10807] T.; Horton, W. A.; Machado, M.; Kaitila, I.; McIntosh, I.;
Francomano,
[10808] C. A.: Achondroplasia is defined by recurrent G380R mu-
tations of
[10809] FGFR3. Am. J. Hum. Genet. 56: 368-373, 1995.

[10810] 306.

[10811] Bellus, G. A.; McIntosh, I.; Smith, E. A.; Aylsworth, A. S.;
Kaitila,
[10812] I.; Horton, W. A.; Greenhaw, G. A.; Hecht, J. T.; Franco-
mano, C. A.
[10813] : A recurrent mutation in the tyrosine kinase domain of fi-

broblast

[10814] growth factor receptor 3 causes hypochondroplasia. Nature Genet. 10:

[10815] 357-359, 1995.

[10816] 307.

[10817] Arai, E.; Ikeuchi, T.; Karasawa, S.; Tamura, A.; Yamamoto, K.;

[10818] Kida, M.; Ichimura, K.; Yuasa, Y.; Tonomura, A.: Constitutional translocation

[10819] t(4;22)(q12;q12.2) associated with neurofibromatosis type 2. Am.

[10820] J. Med. Genet. 44: 163-167, 1992.

[10821] 308.

[10822] Baser, M. E.; Friedman, J. M.; Aeschliman, D.; Joe, H.; Wallace,

[10823] A. J.; Ramsden, R. T.; Evans, D. G. R.: Predictors of the risk of

[10824] mortality in neurofibromatosis 2. Am. J. Hum. Genet. 71: 715-723,

[10825] 2002.

[10826] 309.

[10827] Baser, M. E.; Friedman, J. M.; Evans, D. G. R.: Maternal gene
[10828] effect in neurofibromatosis 2: fact or artefact? (Letter) J. Med.
[10829] Genet. 38: 783-784, 2001.

[10830] 310.

[10831] Bianchi, A. B.; Hara, T.; Ramesh, V.; Gao, J.; Klein-Szanto, A.
[10832] J. P.; Morin, F.; Menon, A. G.; Trofatter, J. A.; Gusella, J. F.;

[10833] Seizinger, B. R.; Kley, N.: Mutations in transcript isoforms of the
[10834] neurofibromatosis 2 gene in multiple human tumour types. Nature Genet. 6:
[10835] 185-192, 1994.

[10836] 311.

[10837] Bianchi, A. B.; Mitsunaga, S.-I.; Cheng, J. Q.; Klein, W. M.; Jhanwar,
[10838] S. C.; Seizinger, B.; Kley, N.; Klein-Szanto, A. J. P.; Testa, J.

[10839] R.: High frequency of inactivating mutations in the neu-rofibromatosis
[10840] type 2 gene (NF2) in primary malignant mesotheliomas.

Proc. Nat.
[10841] Acad. Sci. 92: 10854-10858, 1995.

[10842] 312.

[10843] Drury, A. N.; Szent-Gyorgyi, A.: The physiological activity of
[10844] adenine compounds with especial reference to their action upon the
[10845] mammalian heart. J. Physiol. 68: 213-237, 1929.

[10846] 313.

[10847] Libert, F.; Passage, E.; Parmentier, M.; Simons, M.-J.; Vas-
sart,
[10848] G.; Mattei, M.-G.: Chromosomal mapping of A1 and A2 adenosine receptors,
[10849] VIP receptor, and a new subtype of serotonin receptor. Genomics 11:
[10850] 225-227, 1991.

[10851] 314.

[10852] Stiles, G. L.: Adenosine receptors. J. Biol. Chem. 267:
6451-6454,
[10853] 1992.

[10854] 315.

[10855] Sun, D.; Samuelson, L. C.; Yang, T.; Huang, Y.; Paliege, A.; Saunders,
[10856] T.; Briggs, J.; Schnermann, J.: Mediation of tubu-
loglomerular feedback
[10857] by adenosine: evidence from mice lacking adenosine 1 re-
ceptors. Proc.
[10858] Nat. Acad. Sci. 98: 9983-9988, 2001.

[10859] 316.

[10860] Townsend-Nicholson, A.; Baker, E.; Schofield, P. R.; Sutherland,
[10861] G. R.: Localization of the adenosine A1 receptor subtype gene (ADORA1)
[10862] to chromosome 1q32.1. Genomics 26: 423-425, 1995.

[10863] 317.

[10864] Chen, J.-F.; Huang, Z.; Ma, J.; Zhu, J.; Moratalla, R.; Stan-
daert,
[10865] D.; Moskowitz, M. A.; Fink, J. S.; Schwarzschild, M. A.: A2A adenosine
[10866] receptor deficiency attenuates brain injury induced by transient focal
[10867] ischemia in mice. J. Neurosci. 19: 9192-9200, 1999.

[10868] 318.

[10869] Gaudray, P.: Personal Communication. Nice, France 6/1/1994.

[10870] 319.

[10871] Gusella, J. F.: Personal Communication. Boston, Mass. 4/17/1994.

[10872] 320.

[10873] Le, F.; Townsend-Nicholson, A.; Baker, E.; Sutherland, G. R.; Schofield,

[10874] P. R.: Characterization and chromosomal localization of the human

[10875] A2a adenosine receptor gene: ADORA2A. Biochem. Biophys. Res. Commun. 223:

[10876] 461-467, 1996.

[10877] 321.

[10878] Ledent, C.; Vaugeois, J.-M.; Schiffmann, S. N.; Pedrazzini, T.;

[10879] El Yacoubi, M. E.; Vanderhaeghen, J.-J.; Costentin, J.; Heath, J.

[10880] K.; Vassart, G.; Parmentier, M.: Aggressiveness, hypoalgesia and

[10881] high blood pressure in mice lacking the adenosine A2a receptor. Nature 388:

[10882] 674-678, 1997.

[10883] 322.

[10884] Libert, F.; Passage, E.; Parmentier, M.; Simons, M.-J.; Vassart,

[10885] G.; Mattei, M.-G.: Chromosomal mapping of A1 and A2 adenosine receptors,

[10886] VIP receptor, and a new subtype of serotonin receptor. Genomics 11:

[10887] 225-227, 1991. Note: Erratum: Genomics 23: 305 only, 1994.

[10888] 323.

[10889] MacCollin, M.; Peterfreund, R.; MacDonald, M.; Fink, J. S.; Gusella,

[10890] J.: Mapping of a human A2a adenosine receptor (ADORA2) to chromosome

[10891] 22. Genomics 20: 332-333, 1994.

[10892] 324.

[10893] Ohta, A.; Sitkovsky, M.: Role of G-protein-coupled adenosine receptors

[10894] in downregulation of inflammation and protection from tissue damage. Nature 916-920,

[10895] 2001.

[10896] 325.

[10897] Szepetowski, P.; Perucca-Lostanlen, D.; Gaudray, P.: Mapping genes

[10898] according to their amplification status in tumor cells: contribution

[10899] to the map of 11q13. Genomics 16: 745-750, 1993.

[10900] 326.

[10901] Kudoh, J.; Minoshima, S.; Fukuyama, R.; Maekawa, M.; Neckelmann,

[10902] N.; Wallace, D. C.; Shimizu, Y.; Shimizu, N.: Assignment of ATP synthase

[10903] beta subunit (ATPMB) gene to the p13-qter region of human chromosome

[10904] 12. (Abstract) Cytogenet. Cell Genet. 51: 1026 only, 1989.

[10905] 327.

[10906] Neckelmann, N.; Warner, C. K.; Chung, A.; Kudoh, J.; Minoshima,

[10907] S.; Fukuyama, R.; Maekawa, M.; Shimizu, Y.; Shimizu, N.;

Liu, J. D.;

[10908] Wallace, D. C.: The human ATP synthase beta subunit gene: sequence

[10909] analysis, chromosome assignment, and differential expression. Genomics 5:

[10910] 829-843, 1989.

[10911] 328.

[10912] Neckelmann, N. S.; Chung, A. B.; Warner, C. K.; Hodge, J. A.; Wallace,

[10913] D. C.: The human ATP synthase beta subunit gene has been sequenced

[10914] and shown to be preferentially expressed in heart and skeletal muscle.

[10915] (Abstract) Cytogenet. Cell Genet. 51: 1051 only, 1989.

[10916] 329.

[10917] Ohta, S.; Tomura, H.; Matsuda, K.; Kagawa, Y.: Gene structure

[10918] of the human mitochondrial adenosine triphosphate synthase beta subunit. J.

[10919] Biol. Chem. 263: 11257-11262, 1988.

[10920] 330.

[10921] Hawkins, J. W.; Dugaiczyk, A.: The human serum albumin gene:
[10922] structure of a unique locus. Gene 19: 55-58, 1982.

[10923] 331.

[10924] Huss, K.; Madison, J.; Ishioka, N.; Takahashi, N.; Arai, K.; Putnam,
[10925] F. W.: The same substitution, glutamic acid-to-lysine at position
[10926] 501, occurs in three alloalbumins of Asiatic origin: albumins Vancouver,
[10927] Birmingham, and Adana. Proc. Nat. Acad. Sci. 85: 6692-6696, 1988.
[10928] 332.

[10929] Huss, K.; Putnam, F. W.; Takahashi, N.; Takahashi, Y.; Weaver,
[10930] G. A.; Peters, T., Jr.: Albumin Cooperstown: a serum albumin variant
[10931] with the same (313 lys-to-asn) mutation found in albumins in Italy
[10932] and New Zealand. Clin. Chem. 34: 183-187, 1988.

[10933] 333.

[10934] Hutchinson, D. W.; Matejtschuk, P.; Lord, C.: Albumin Carlisle:
[10935] occurrence and properties of a new human albumin variant. IRCS Med.
[10936] Sci. 14: 1095-1096, 1986.

[10937] 334.

[10938] Jamieson, G. A.; Ganguly, P.: Studies on a genetically determined
[10939] albumin dimer. Biochem. Genet. 3: 403-416, 1969.

[10940] 335.

[10941] Jensen, I. W.; Faber, J.: Familial dysalbuminemic hyperthyroxinemia. Acta
[10942] Med. Scand. 221: 469-473, 1987.

[10943] 336.

[10944] Kaarsalo, E.; Melartin, L.; Blumberg, B. S.: Autosomal linkage
[10945] between the albumin and GC loci in humans. Science 158: 123-125,
[10946] 1967.

[10947] 337.

[10948] Kallee, E.: Bennhold's analbuminemia: a follow-up study of the
[10949] first two cases (1953-1992). J. Lab. Clin. Med. 127: 470-480, 1996.
[10950] 338.

[10951] Kao, F.-T.; Hawkins, J. W.; Law, M. L.; Dugaiczyk, A.: Assignment
[10952] of the structural gene coding for albumin to human chromosome 4. Hum.
[10953] Genet. 62: 337-341, 1982.

[10954] 339.

[10955] Kaur, H.; Franklin, S. G.; Shrivastava, P. K.; Blumberg, B. S.

[10956] : Alloalbuminemia in North India. Am. J. Hum. Genet. 34: 972-979,
[10957] 1982.

[10958] 340.

[10959] Kueppers, F.; Holland, P. V.; Weitkamp, L. R.: Albumin Santa
[10960] Ana: a new inherited variant. Hum. Hered. 19: 378-384, 1969.
[10961] 341.

[10962] Kurnit, D. M.; Philipp, B. W.; Bruns, G. A. P.: Confirmation

[10963] of the mapping assignment of human serum albumin to chromosome 4 using

[10964] a cloned human albumin gene. Cytogenet. Cell Genet. 34: 282-288,

[10965] 1982.

[10966] 342.

[10967] Lalloz, M. R. A.; Byfield, P. G. H.; Himsworth, R. L.: Hyper-thyroxinaemia:

[10968] abnormal binding of T4 by an inherited albumin variant. Clin. Endocr. 18:

[10969] 11-24, 1983.

[10970] 343.

[10971] Lalloz, M. R. A.; Byfield, P. G. H.; Himsworth, R. L.: A new

[10972] and distinctive albumin variant with increased affinities for both

[10973] triiodothyronines and causing hyperthyroxinaemia. Clin. Endocr. 22:

[10974] 521-529, 1985.

[10975] 344.

[10976] Lau, T. J.; Sunderman, F. W., Jr.; Weitkamp, L. R.; Agarwal, S.

[10977] S.; Sutnick, A. I.; Blumberg, B. S.; De Jimenez, R. B. C.: Albumin

[10978] Cartago: a 'new' slow-moving alloalbumin. Am. J. Clin. Path. 57:

[10979] 247-251, 1972.

[10980] 345.

[10981] Laurell, C. B.; Nilehn, J. E.: A new type of inherited serum

[10982] albumin anomaly. J. Clin. Invest. 45: 1935-1945, 1966.

[10983] 346.

[10984] Lavareda de Souza, S.; Frain, M.; Mornet, E.; Sala-Trepat, J.

[10985] M.; Lucotte, G.: Polymorphisms of human albumin gene after DNA restriction

[10986] by HaeIII endonuclease. Hum. Genet. 67: 48-51, 1984.

[10987] 347.

[10988] Lyon, A. W.; Meinert, P.; Bruce, G. A.; Laxdal, V. A.; Salkie,

[10989] M. L.: Influence of methodology on the detection and diagnosis of

[10990] congenital analbuminemia. Clin. Chem. 44: 2365-2367,

1998.

[10991] 348.

[10992] Madison, J.; Arai, K.; Sakamoto, Y.; Feld, R. D.; Kyle, R. A.;

[10993] Watkins, S.; Davis, E.; Matsuda, Y.; Amaki, I.; Putnam, F.
W.: Genetic

[10994] variants of serum albumin in Americans and Japanese.
Proc. Nat. Acad.

[10995] Sci. 88: 9853-9857, 1991.

[10996] 349.

[10997] Madison, J.; Galliano, M.; Watkins, S.; Minchiotti, L.; Porta,

[10998] F.; Rossi, A.; Putnam, F. W.: Genetic variants of human
serum albumin

[10999] in Italy: point mutants and a carboxyl-terminal variant.
Proc. Nat.

[11000] Acad. Sci. 91: 6476-6480, 1994.

[11001] 350.

[11002] Magenis, R. E.; Luo, X. Y.; Dugaiczyk, A.; Ryan, S. C.;
Oosterhuis,

[11003] J. E.: Chromosomal localization of the albumin and alpha-
fetoprotein

[11004] genes in the orangutan (Pongo pygmaeus) and gorilla

(Gorilla gorilla).

[11005] (Abstract) Cytogenet. Cell Genet. 51: 1037 only, 1989.

[11006] 351.

[11007] Melartin, L.: Albumin polymorphism in man: studies on al-
bumin

[11008] variants in North American native populations. Acta Path.
Microbiol.

[11009] Scand. 191 (suppl.): 1-50, 1967.

[11010] 352.

[11011] Melartin, L.; Blumberg, B. S.: Albumin Naskapi: a new vari-
ant

[11012] of serum albumin. Science 153: 1664-1666, 1966.

[11013] 353.

[11014] Melartin, L.; Blumberg, B. S.; Lisker, R.: Albumin Mexico, a

[11015] new variant of serum albumin. Nature 215: 1288-1289,
1967.

[11016] 354.

[11017] Mendel, C. M.; Cavalieri, R. R.; Gavin, L. A.; Pettersson, T.;

[11018] Inoue, M.: Thyroxine transport and distribution in Nagase
analbuminemic

[11019] rats. J. Clin. Invest. 83: 143-148, 1989.

[11020] 355.

[11021] Minchiotti, L.; Galliano, M.; Iadarola, P.; Meloni, M. L.; Ferri,

[11022] G.; Porta, F.; Castellani, A. A.: The molecular defect in a COOH-terminal-modified

[11023] and shortened mutant of human serum albumin. J. Biol. Chem. 264:

[11024] 3385-3389, 1989.

[11025] 356.

[11026] Minchiotti, L.; Galliano, M.; Kragh-Hansen, U.; Watkins, S.; Madison,

[11027] J.; Putnam, F. W.: A genetic variant of albumin (albumin Asola; tyr140-to-cys)

[11028] with no free -SH group but with an additional disulfide bridge. Europ.

[11029] J. Biochem. 228: 155-159, 1995.

[11030] 357.

[11031] Minchiotti, L.; Galliano, M.; Stoppini, M.; Ferri, G.; Cres-peau,

[11032] H.; Rochu, D.; Porta, F.: Two alloalbumins with identical

electrophoretic
[11033] mobility are produced by differently charged amino acid substitutions. Biochim.
[11034] Biophys. Acta 1119: 232-238, 1992.

[11035] 358.

[11036] Minchiotti, L.; Galliano, M.; Zapponi, M. C.; Tenni, R.: The

[11037] structural characterization and bilirubin-binding properties of albumin
[11038] Herborn, a lys240-to-glu albumin mutant. Europ. J. Biochem. 214:
[11039] 437-444, 1993.

[11040] 359.

[11041] Minghetti, P. P.; Law, S. W.; Dugaiczyk, A.: The rate of molecular
[11042] evolution of alpha-fetoprotein approaches that of pseudogenes. Molec.
[11043] Biol. Evol. 2: 347-358, 1985.

[11044] 360.

[11045] Minghetti, P. P.; Ruffner, D. E.; Kuang, W.-J.; Dennison, O. E.;
[11046] Hawkins, J. W.; Beattie, W. G.; Dugaiczyk, A.: Molecular

structure

[11047] of the human albumin gene is revealed by nucleotide sequence within

[11048] q11-22 of chromosome 4. J. Biol. Chem. 261: 6747-6757, 1986.

[11049] 361.

[11050] Murray, J. C.; Demopulos, C. M.; Lawn, R. M.; Motulsky, A. G.

[11051] : Restriction endonuclease study of analbuminemia and polymorphisms

[11052] at the albumin locus. (Abstract) Clin. Res. 31: 456A only, 1983.

[11053] 362.

[11054] Murray, J. C.; Demopulos, C. M.; Lawn, R. M.; Motulsky, A. G.

[11055] : Molecular genetics of human serum albumin: restriction enzyme fragment

[11056] length polymorphisms and analbuminemia. Proc. Nat. Acad. Sci. 80:

[11057] 5951-5955, 1983.

[11058] 363.

[11059] Murray, J. C.; Mills, K. A.; Demopulos, C. M.; Hornung, S.;

Motulsky,
[11060] A. G.: Linkage disequilibrium and evolutionary relationships of DNA
[11061] variants (restriction enzyme fragment length polymorphisms) at the
[11062] serum albumin locus. Proc. Nat. Acad. Sci. 81: 3486-3490, 1984.
[11063] 364.

[11064] Perkoff, G. T.; Nugent, C. A., Jr.; Dolowitz, D. A.; Stephens,

[11065] F. E.; Carnes, W. H.; Tyler, F. H.: A follow-up study of hereditary
[11066] chronic nephritis. Arch. Intern. Med. 102: 733-746, 1958.

[11067] 365.

[11068] Barbry, P.; Champe, M.; Chassande, O.; Munemitsu, S.; Champigny,
[11069] G.; Lingueglia, E.; Maes, P.; Frelin, C.; Tartar, A.; Ullrich, A.;
[11070] Lazdunski, M.: Human kidney amiloride-binding protein: cDNA structure
[11071] and functional expression. Proc. Nat. Acad. Sci. 87: 7347-7351,
[11072] 1990.

[11073] 366.

[11074] Barbry, P.; Simon-Bouy, B.; Mattei, M.-G.; Le Guern, E.; Jaume-Roig,

[11075] B.; Chassande, O.; Ullrich, A.; Lazdunski, M.: Localization of the

[11076] gene for amiloride binding protein on chromosome 7 and RFLP analysis

[11077] in cystic fibrosis families. Hum. Genet. 85: 587-589, 1990.

[11078] 367.

[11079] Chassande, O.; Renard, S.; Barbry, P.; Lazdunski, M.: The human

[11080] gene for diamine oxidase, an amiloride binding protein: molecular

[11081] cloning, sequencing, and characterization of the pro- moter. J. Biol.

[11082] Chem. 269: 14484-14489, 1994.

[11083] 368.

[11084] Novotny, W. F.; Chassande, O.; Baker, M.; Lazdunski, M.; Barbry,

[11085] P.: Diamine oxidase is the amiloride-binding protein and is inhibited

[11086] by amiloride analogues. J. Biol. Chem. 269: 9921-9925, 1994.

[11087] 369.

[11088] Ala-Kapee, M.; Forsberg, U. H.; Jalkanen, S.; Schroder, J.: Mapping

[11089] of gene for human lymphocyte homing receptor to the short arm of chromosome

[11090] 11. (Abstract) Cytogenet. Cell Genet. 51: 948-949, 1989.

[11091] 370.

[11092] Aruffo, A.; Stamenkovic, I.; Melnick, M.; Underhill, C. B.; Seed,

[11093] B.: CD44 is the principal cell surface receptor for hyaluronate. Cell 61:

[11094] 1303-1313, 1990.

[11095] 371.

[11096] Cianfriglia, M.; Viora, M.; Tombesi, M.; Merendino, N.; Es-posito,

[11097] G.; Samoggia, P.; Forsberg, U. H.; Schroder, J.: The gene encoding

[11098] for MC56 determinant (drug-sensitivity marker) is located on the short

[11099] arm of human chromosome 11. Int. J. Cancer 52: 585-587, 1992.

[11100] 372.

[11101] Cywes, C.; Wessels, M. R.: Group A Streptococcus tissue invasion

[11102] by CD44-mediated cell signalling. Nature 414: 648-652, 2001.

[11103] 373.

[11104] Forsberg, U. H.; Ala-Kapee, M. M.; Jalkanen, S.; Anders-

son, L.

[11105] C.; Schroder, J.: The gene for human lymphocyte homing receptor is

[11106] located on chromosome 11. Europ. J. Immun. 19: 409-412, 1989.

[11107] 374.

[11108] Forsberg, U. H.; Jalkanen, S.; Schroder, J.: Assignment of the

[11109] human lymphocyte homing receptor gene to the short arm of chromosome

[11110] 11. Immunogenetics 29: 405-407, 1989.

[11111] 375.

[11112] Haynes, B. F.: Personal Communication. Durham, N. C. 2/28/1986.

[11113] 376.

[11114] Krainer, A. R.; Mayeda, A.; Kozak, D.; Binns, G.: Functional expression

[11115] of cloned human splicing factor SF2: homology to RNA-binding proteins,

[11116] U1 70K, and Drosophila splicing regulators. Cell 66: 383-394, 1991.

[11117] 377.

[11118] Matsumura, Y.; Tarin, D.: Significance of CD44 gene products

[11119] for cancer diagnosis and disease evaluation. Lancet 340: 1053-1058,

[11120] 1992.

[11121] 378.

[11122] Mayer, B.; Jauch, K. W.; Gunthert, U.; Figdor, C. G.; Schildberg,

[11123] F. W.; Funke, I.; Johnson, J. P.: De-novo expression of CD44 and

[11124] survival in gastric cancer. Lancet 342: 1019-1022, 1993.

[11125] 379.

[11126] Omary, M. B.; Trowbridge, I. S.; Letarte, M.; Kagnoff, M. F.;

[11127] Isacke, C. M.: Structural heterogeneity of human Pgp-1 and its relationship

[11128] with p85. Immunogenetics 27: 460-464, 1988.

[11129] 380.

[11130] Schlossman, S. F.; Boumsell, L.; Gilks, W.; Harlan, J. M.; Kishimoto,

[11131] T.; Morimoto, C.; Ritz, J.; Shaw, S.; Silverstein, R. L.; Springer,

[11132] T. A.; Tedder, T. F.; Todd, R. F.: CD antigens 1993. Im-

mun. Today 15:

[11133] 98-99, 1994.

[11134] 381.

[11135] Schmits, R.; Filmus, J.; Gerwin, N.; Senaldi, G.; Kiefer, F.;

[11136] Kundig, T.; Wakeham, A.; Shahinian, A.; Catzavelos, C.; Rak, J.; Furlonger,

[11137] C.; Zakarian, A.; Simard, J. J.; Ohashi, P. S.; Paige, C. J.; Gutierrez-Romas,

[11138] J. C.; Mak, T. W.: CD44 regulates hematopoietic progenitor distribution,

[11139] granuloma formation, and tumorigenicity. Blood 90: 2217-2233, 1997.

[11140] 382.

[11141] Screaton, G. R.; Bell, M. V.; Jackson, D. G.; Cornelis, F. B.;

[11142] Gerth, U.; Bell, J. I.: Genomic structure of DNA encoding the lymphocyte

[11143] homing receptor CD44 reveals at least 12 alternatively spliced exons. Proc.

[11144] Nat. Acad. Sci. 89: 12160-12164, 1992.

[11145] 383.

[11146] Spring, F. A.; Holmes, C. H.; Simpson, K. L.; Mawby, W. J.; Mattes,

[11147] M. J.; Okubo, Y.; Parsons, S. F.: The Ok(a) blood group antigen is

[11148] a marker for the M6 leukocyte activation antigen, the human homolog

[11149] of OX-47 antigen, basigin and neurothelin, an immunoglobulin superfamily

[11150] molecule that is widely expressed in human cells and tissues. Europ.

[11151] J. Immun. 27: 891-897, 1997.

[11152] 384.

[11153] Olives, B.; Neau, P.; Bailly, P.; Hediger, M. A.; Rousselet, G.;

[11154] Cartron, J.-P.; Ripoche, P.: Cloning and functional expres-sion of

[11155] a urea transporter from human bone marrow cells. J. Biol. Chem. 269:

[11156] 31649-31652, 1994.

[11157] 385.

[11158] Pausch, V.; Mayr, W. R.: Analysis of the linkage JK-IGK, MNS-GC

[11159] and of two other possible linkage groups. Hum. Hered. 37: 260-262,

[11160] 1987.

[11161] 386.

[11162] Promeneur, D.; Rousselet, G.; Bankir, L.; et al: Evidence for

[11163] distinct vascular and tubular urea transporters in the rat kidney. J.

[11164] Am. Soc. Nephrol. 7: 852-860, 1996.

[11165] 387.

[11166] Sands, J. M.; Gargus, J. J.; Frohlich, O.; Gunn, R. B.; Kokko,

[11167] J. P.: Urinary concentrating ability in patients with Jk(a/b) blood

[11168] type who lack carrier-mediated urea transport. J. Am. Soc. Nephrol. 2:

[11169] 1689-1696, 1992.

[11170] 388.

[11171] Sherman, S. L.; Simpson, S. P.: Evidence for the location of

[11172] JK and CO on chromosome 2 based on family studies. (Abstract) Cytogenet.

[11173] Cell Genet. 40: 743, 1985.

[11174] 389.

[11175] Shokeir, M. H. K.; Ying, K. L.; Pabello, P.: Deletion of the

[11176] long arm of chromosome no. 7: tentative assignment of the Kidd (Jk)

[11177] locus. Clin. Genet. 4: 360-368, 1973.

[11178] 390.

[11179] Sidoux-Walter, F.; Lucien, N.; Nissinen, R.; Sistonen, P.; Henry,

[11180] S.; Moulds, J.; Cartron, J.-P.; Bailly, P.: Molecular hetero-

geneity

[11181] of the Jk-null phenotype: expression analysis of the Jk(S291P) mutation

[11182] found in Finns. Blood 96: 1566-1573, 2000.

[11183] 391.

[11184] Tsukaguchi, H.; Shayakul, C.; Berger, U. V.; Tokui, T.; Brown,

[11185] D.; Hediger, M. A.: Cloning and characterization of the urea transportation

[11186] UT3: localization in rat kidney and testis. J. Clin. Invest. 99:

[11187] 1506-1515, 1997.

[11188] 392.

[11189] Xu, Y.; Olives, B.; Bailly, P.; et al: Endothelial cells of the

[11190] kidney vasa recta express the urea transporter HUT11. Kidney Int. 51:

[11191] 138-146, 1997.

[11192] 393.

[11193] Grollman, E. F.; Kobata, A.; Ginsburg, V.: An enzymatic basis

[11194] for Lewis blood types in man. J. Clin. Invest. 48:

1489-1494, 1969.

[11195] 394.

[11196] Koda, Y.; Kimura, H.; Mekada, E.: Analysis of Lewis fuco-
syltransferase

[11197] genes from the human gastric mucosa of Lewis-positive
and -negative

[11198] individuals. Blood 82: 2915-2919, 1993.

[11199] 395.

[11200] Koprowski, H.; Blaszczyk, M.; Steplewski, Z.; Brockhaus,
M.; Magnani,

[11201] J.; Ginsburg, V.: Lewis blood-type may affect the incidence
of gastrointestinal

[11202] cancer. Lancet I: 1332-1333, 1982.

[11203] 396.

[11204] Nishihara, S.; Narimatsu, H.; Iwasaki, H.; Yazawa, S.; Aka-
matsu,

[11205] S.; Ando, T.; Seno, T.; Narimatsu, I.: Molecular genetic
analysis

[11206] of the human Lewis histo-blood group system. J. Biol.
Chem. 269:

[11207] 29271-29278, 1994.

[11208] 397.

[11209] Orntoft, T. F.; Vestergaard, E. M.; Holmes, E.; Jakobsen, J. S.;

[11210] Grunnet, N.; Mortensen, M.; Johnson, P.; Bross, P.; Gregersen, N.;

[11211] Skorstengaard, K.; Jensen, U. B.; Bolund, L.; Wolf, H.: Influ-

[11212] ence of Lewis alpha-1-3/4-L-fucosyltransferase (FUT3) gene mutations on

[11213] enzyme activity, erythrocyte phenotyping, and circulating tumor marker

[11214] sialyl-Lewis a levels. J. Biol. Chem. 271: 32260-32268, 1996.

[11215] 398.

[11216] Pang, H.; Liu, Y.; Koda, Y.; Soejima, M.; Jia, J.; Schlaphoff,

[11217] T.; du Toit, E. D.; Kimura, H.: Five novel missense muta-

[11218] tions of the Lewis gene (FUT3) in African (Xhosa) and Caucasian populations

[11219] in South Africa. Hum. Genet. 102: 675-680, 1998.

[11220] 399.

[11221] Reguigne-Arnould, I.; Couillin, P.; Mollicone, R.; Faure, S.; Fletcher,

[11222] A.; Kelly, R. J.; Lowe, J. B.; Oriol, R.: Relative positions of two

[11223] clusters of human alpha-L-fucosyltransferases in 19q (FUT1-FUT2) and

[11224] 19p (FUT6-FUT3-FUT5) within the microsatellite genetic map of chromosome

[11225] 19. Cytogenet. Cell Genet. 71: 158-162, 1995.

[11226] 400.

[11227] Sheinfeld, J.; Schaeffer, A. J.; Cordon-Cardo, C.; Rogatko, A.;

[11228] Fair, W. R.: Association of the Lewis blood-group pheno-type with

[11229] recurrent urinary tract infections in women. New Eng. J. Med. 320:

[11230] 773-777, 1989.

[11231] 401.

[11232] Weitkamp, L. R.; Johnston, E.; Guttormsen, S. A.: Probable genetic

[11233] linkage between the loci for the Lewis blood group and complement

[11234] C3. Cytogenet. Cell Genet. 13: 183-184, 1974.

[11235] 402.

[11236] Yazawa, S.; Oh-Kawara, H.; Nakajima, T.; Hosomi, O.; Akamatsu,

[11237] S.; Kishi, K.: Histo-blood group Lewis genotyping from human hairs

[11238] and blood. Jpn. J. Hum. Genet. 41: 177-188, 1996.

[11239] 403.

[11240] Bandera, C. A.; Muto, M. G.; Schorge, J. O.; Berkowitz, R. S.;

[11241] Rubin, S. C.; Mok, S. C.: BRCA1 gene mutations in women with papillary

[11242] serous carcinoma of the peritoneum. Obstet. Gynec. 92: 596-600,

[11243] 1998.

[11244] 404.

[11245] Bar-Sade, R. B.; Kruglikova, A.; Modan, B.; Gak, E.; Hirsh-Yechezkel,

[11246] G.; Theodor, L.; Novikov, I.; Gershoni-Baruch, R.; Risel, S.; Papa,

[11247] M. Z.; Ben-Baruch, G.; Friedman, E.: The 185delAG BRCA1 mutation

[11248] originated before the dispersion of Jews in the Diaspora and is not

[11249] limited to Ashkenazim. Hum. Molec. Genet. 7: 801-805, 1998.

[11250] 405.

[11251] Bar-Sade, R. B.; Theodor, L.; Gak, E.; Kruglikova, A.; Hirsch-Yechezkel,

[11252] G.; Modan, B.; Kuperstein, G.; Seligsohn, U.; Rechavi, G.; Friedman,

[11253] E.: Could the 185delAG BRCA1 mutation be an ancient Jewish mutation? Europ.

[11254] J. Hum. Genet. 5: 413-416, 1997.

[11255] 406.

[11256] Barker, D. F.; Almeida, E. R. A.; Casey, G.; Fain, P. R.; Liao,

[11257] S. Y.; Masunaka, I.; Noble, B.; Kurosaki, T.; Anton-Culver, H.: BRCA1

[11258] R841W: a strong candidate for a common mutation with moderate phenotype. Genet.

[11259] Epidemiol. 13: 595-604, 1996.

[11260] 407.

[11261] Bennett, L. M.; Haugen-Strano, A.; Cochran, C.; Brownlee, H. A.;

[11262] Fiedorek, F. T., Jr.; Wiseman, R. W.: Isolation of the mouse homologue

[11263] of BRCA1 and genetic mapping to mouse chromosome 11. Genomics 29:

[11264] 576-581, 1995.

[11265] 408.

[11266] Bergman, A.; Einbeigi, Z.; Olofsson, U.; Taib, Z.; Wallgren, A.;

[11267] Karlsson, P.; Wahlstrom, J.; Martinsson, T.; Nordling, M.: The western

[11268] Swedish BRCA1 founder mutation 3171ins5; a 3.7 cM conserved haplotype

[11269] of today is a reminiscence of a 1500-year-old mutation. Europ. J.

[11270] Hum. Genet. 9: 787-793, 2001.

[11271] 409.

[11272] Bochar, D. A.; Wang, L.; Beniya, H.; Kinev, A.; Xue, Y.; Lane,

[11273] W. S.; Wang, W.; Kashanchi, F.; Shiekhattar, R.: BRCA1 is associated

[11274] with a human SWI/SNF-related complex: linking chromatin remodeling

[11275] to breast cancer. Cell 102: 257-265, 2000.

[11276] 410.

[11277] Boyd, J.; Sonoda, Y.; Federici, M. G.; Bogomolniy, F.; Rhei, E.;

[11278] Maresco, D. L.; Saigo, P. E.; Almadrones, L. A.; Barakat, R. R.; Brown,

[11279] C. L.; Chi, D. S.; Curtin, J. P.; Poynor, E. A.; Hoskins, W. J.:

[11280] Clinicopathologic features of BRCA-linked and sporadic ovarian cancer. J.A.M.A. 283:

[11281] 2260-2265, 2000.

[11282] 411.

[11283] BRCA1 Exon 13 Duplication Screening Group: The exon 13 duplication

[11284] in the BRCA1 gene is a founder mutation present in geo-graphically

[11285] diverse populations. Am. J. Hum. Genet. 67: 207-212, 2000.

[11286] 412.

[11287] Breast Cancer Linkage Consortium: Pathology of familial breast

[11288] cancer: differences between breast cancers in carriers of BRCA1 and

[11289] BRCA2 mutations and sporadic cases. Lancet 349: 1505-1510, 1997.

[11290] 413.

[11291] Briki, F.; Busson, B.; Salicru, B.; Esteve, F.; Doucet, J.: Breast-cancer

[11292] diagnosis using hair. (Letter) Nature 400: 220 only, 1999.

[11293] 414.

[11294] Brown, M. A.; Lo, L.-J.; Catteau, A.; Xu, C.-F.; Lindeman, G.

[11295] J.; Hodgson, S.; Solomon, E.: Germline BRCA1 promoter deletions in

[11296] UK and Australian familial breast cancer patients: identification

[11297] of a novel deletion consistent with BRCA1:psi-BRCA1 recombination. Hum.

[11298] Mutat. 19: 435-442, 2002.

[11299] 415.

[11300] Castilla, L. H.; Couch, F. J.; Erdos, M. R.; Hoskins, K. F.; Calzone,

[11301] K.; Garber, J. E.; Boyd, J.; Lubin, M. B.; Deshano, M. L.; Brody,

[11302] L. C.; Collins, F. S.; Weber, B. L.: Mutations in the BRCA1 gene

[11303] in families with early-onset breast and ovarian cancer. Nature Genet. 8:

[11304] 387-391, 1994.

[11305] 416.

[11306] Chen, J.; Silver, D. P.; Walpita, D.; Cantor, S. B.; Gazdar, A.

[11307] F.; Tomlinson, G.; Couch, F. J.; Weber, B. L.; Ashley, T.; Livingston,

[11308] D. M.; Scully, R.: Stable interaction between the products of the

[11309] BRCA1 and BRCA2 tumor suppressor genes in mitotic and meiotic cells. Molec.

[11310] Cell 2: 317-328, 1998.

[11311] 417.

[11312] Chen, Y.; Chen, C.-F.; Riley, D. J.; Allred, D. C.; Chen, P.-L.;

[11313] Von Hoff, D.; Osborne, C. K.; Lee, W.-H.: Aberrant subcellular localization

[11314] of BRCA1 in breast cancer. Science 270: 789-791, 1995.

[11315] 418.

[11316] Chen, Y.; Farmer, A. A.; Chen, C.-F.; Jones, D. C.; Chen, P.-L.;

[11317] Lee, W.-H.: BRCA1 is a 220-kDa nuclear phosphoprotein that is expressed

[11318] and phosphorylated in a cell cycle-dependent manner.

Cancer Res. 56:

[11319] 3168-3172, 1996.

[11320] 419.

[11321] Claus, E. B.; Risch, N.; Thompson, W. D.: Genetic analysis of

[11322] breast cancer in the cancer and steroid hormone study. Am. J. Hum.

[11323] Genet. 48: 232-242, 1991.

[11324] 420.

[11325] Coene, E.; Van Oostveldt, P.; Willems, K.; van Emmelo, J.; De

[11326] Potter, C. R.: BRCA1 is localized in cytoplasmic tube-like invaginations

[11327] in the nucleus. (Letter) Nature Genet. 116: 122-124, 1997.

[11328] 421.

[11329] Cornelis, R. S.; Vasen, H. F. A.; Meijers-Heijboer, H.; Ford,

[11330] D.; van Vliet, M.; van Tilborg, A. A. G.; Cleton, F. J.; Klijn, J.

[11331] G. M.; Menko, F. H.; Khan, P. M.; Cornelisse, C. J.; Devilee, P.:

[11332] Age at diagnosis as an indicator of eligibility for BRCA1

DNA testing
[11333] in familial breast cancer. Hum. Genet. 95: 539-544, 1995.

[11334] 422.

[11335] Couch, F. J.; DeShano, M. L.; Blackwood, M. A.; Calzone, K.; Stopfer,
[11336] J.; Campeau, L.; Ganguly, A.; Rebbeck, T.; Weber, B. L.: BRCA1 mutations
[11337] in women attending clinics that evaluate the risk of breast cancer. New
[11338] Eng. J. Med. 336: 1409-1415, 1997.

[11339] 423.

[11340] Couch, F. J.; Weber, B. L.; Breast Cancer Information Core:

[11341] Mutations and polymorphisms in the familial early-onset breast cancer
[11342] (BRCA1) gene. Hum. Mutat. 8: 8-18, 1996.

[11343] 424.

[11344] De Gregorio, L.; Harshman, K.; Rosenthal, J.; Dragani, T. A.;
[11345] Pierotti, M. A.: Genetic mapping of the Brca1 gene on mouse chromosome
[11346] 11. Mammalian Genome 7: 242, 1996.

[11347] 425.

[11348] De Los Rios, P.; Jack, E.; Kuperstein, G.; Lynch, H.; Lubin-
ski,
[11349] J.; Narod, S. A.: Founder mutations in BRCA1 and BRCA2 in
North American
[11350] families of Polish origin that are affected with breast can-
cer. (Letter) Am.
[11351] J. Hum. Genet. 68: 546 only, 2001.

[11352] 426.

[11353] Dorum, A.; Heimdal, K.; Hovig, E.; Inganas, M.; Moller, P.:
Penetrances
[11354] of BRCA1 1675delA and 1135insA with respect to breast
cancer and ovarian
[11355] cancer. Am. J. Hum. Genet. 65: 671-679, 1999.

[11356] 427.

[11357] Dunning, A. M.; Chiano, M.; Smith, N. R.; Dearden, J.;
Gore, M.;
[11358] Oakes, S.; Wilson, C.; Stratton, M.; Peto, J.; Easton, D.;
Clayton,
[11359] D.; Ponder, B. A. J.: Common BRCA1 variants and suscep-
tibility to

[11360] breast and ovarian cancer in the general population. Hum. Molec.

[11361] Genet. 6: 285-289, 1997.

[11362] 428.

[11363] Dorum, A.; Moller, P.; Kamsteeg, E. J.; Scheffer, H.; Burton,

[11364] M.; Heimdal, K. R.; Maehle, L. O.; Hovig, E.; Trope, C. G.; van der

[11365] Hout, A. H.; van der Meulen, M. A.; Buys, C. H. C. M.; te Meerman,

[11366] G. J.: A BRCA1 founder mutation, identified with haplotype analysis,

[11367] allowing genotype/phenotype determination and predictive testing. Europ.

[11368] J. Cancer 33: 2390-2392, 1997.

[11369] 429.

[11370] Brahe, C.; Bannetta, P.; Meera Khan, P.; Arwert, F.; Serra, A.

[11371] : Assignment of the catechol-O-methyltransferase gene to human chromosome

[11372] 22 in somatic cell hybrids. Hum. Genet. 74: 230-234, 1986.

[11373] 430.

[11374] Brahe, C.; Bannetta, P.; Serra, A.; Arwert, F.: The increased

[11375] COMT activity in Down syndrome patients is not a conse-
quence of dosage

[11376] effect owing to location of the gene on chromosome 21:
further evidence.

[11377] (Letter) Am. J. Med. Genet. 24: 203-204, 1986.

[11378] 431.

[11379] Masuno, M.; Fukao, T.; Song, X.-Q.; Yamaguchi, S.; Orii,
T.; Kondo,

[11380] N.; Imaizumi, K.; Kuroki, Y.: Assignment of the human cy-
tosolic acetoacetyl-coenzyme

[11381] A thiolase (ACAT2) gene to chromosome 6q25.3-q26. Ge-
nomics 36: 217-218,

[11382] 1996.

[11383] 432.

[11384] Morel, Y.; Bristow, J.; Gitelman, S. E.; Miller, W. L.: Tran-
script

[11385] encoded on the opposite strand of the human steroid
21-hydroxylase/complement

[11386] component C4 gene locus. Proc. Nat. Acad. Sci. 86:

6582-6586, 1989.

[11387] 433.

[11388] Song, X.-Q.; Fukao, T.; Yamaguchi, S.; Miyazawa, S.; Hashimoto,

[11389] T.; Orii, T.: Molecular cloning and nucleotide sequence of complementary

[11390] DNA for human hepatic cytosolic acetoacetyl-coenzyme A thiolase. Biochem.

[11391] Biophys. Res. Commun. 201: 478-485, 1994.

[11392] 434.

[11393] Willison, K.; Kelly, A.; Dudley, K.; Goodfellow, P.; Spurr, N.;

[11394] Groves, V.; Gorman, P.; Sheer, D.; Trowsdale, J.: The hu-man homologue

[11395] of the mouse t-complex gene, TCP1, is located on chro-mosome 6 but

[11396] is not near the HLA region. EMBO J. 6: 1967-1974, 1987.

[11397] 435.

[11398] Engel, A. G.; Ohno, K.; Milone, M.; Wang, H.-L.; Nakano, S.; Bouzat,

[11399] C.; Pruitt, J. N., II; Hutchinson, D. O.; Brengman, J. M.; Bren, N.;

[11400] Sieb, J. P.; Sine, S. M.: New mutations in acetylcholine re-
ceptor
[11401] subunit genes reveal heterogeneity in the slow-channel
congenital
[11402] myasthenic syndrome. Hum. Molec. Genet. 5: 1217-1227,
1996.
[11403] 436.

[11404] Lobos, E. A.: Five subunit genes of the human muscle
nicotinic
[11405] acetylcholine receptor are mapped to two linkage groups
on chromosomes
[11406] 2 and 17. Genomics 17: 642-650, 1993.

[11407] 437.

[11408] Deftos, L. J.; Murray, S. S.; Burton, D. W.; Parmer, R. J.;
O'Connor,
[11409] D. T.; Delegeane, A. M.; Mellon, P. L.: A cloned chromo-
granin A (CgA)
[11410] cDNA detects a 2.3kb mRNA in diverse neuroendocrine
tissues. Biochem.
[11411] Biophys. Res. Commun. 137: 418-423, 1986.

[11412] 438.

[11413] Granberg, D.; Stridsberg, M.; Seensalu, R.; Eriksson, B.; Lundqvist,
[11414] G.; Oberg, K.; Skogseid, B.: Plasma chromogranin A in patients with
[11415] multiple endocrine neoplasia type 1. J. Clin. Endocr. Metab. 84:
[11416] 2712-2717, 1999.

[11417] 439.

[11418] Hagn, C.; Schmid, K. W.; Fischer-Colbrie, R.; Winkler, H.: Chromogranin
[11419] A, B, and C in human adrenal medulla and endocrine tissues. Lab.
[11420] Invest. 55: 405-411, 1986.

[11421] 440.

[11422] Kim, T.; Tao-Cheng, J.-H.; Eiden, L. E.; Loh, Y. P.: Chromogranin
[11423] A, an 'on/off' switch controlling dense-core secretory granule biogenesis. Cell 106:
[11424] 499-509, 2001.

[11425] 441.

[11426] Konecki, D. S.; Benedum, U. M.; Gerdes, H.-H.; Huttner, W.

B.:

[11427] The primary structure of human chromogranin A and pancreastatin. J.

[11428] Biol. Chem. 262: 17026-17030, 1987.

[11429] 442.

[11430] Kruggel, W.; O'Connor, D. T.; Lewis, R. V.: The amino terminal

[11431] sequences of bovine and human chromogranin A and secretory protein

[11432] I are identical. Biochem. Biophys. Res. Commun. 127: 380-383, 1985.

[11433] 443.

[11434] Modi, W. S.; Levine, M. A.; Dean, M.; Seuanez, H.; O'Brien, S.

[11435] J.: The chromogranin A gene: chromosome assignment and RFLP analysis.

[11436] (Abstract) Cytogenet. Cell Genet. 51: 1046 only, 1989.

[11437] 444.

[11438] Modi, W. S.; Levine, M. A.; Seuanez, H. N.; Dean, M.; O'Brien,

[11439] S. J.: The human chromogranin A gene: chromosome as-

signment and RFLP
[11440] analysis. Am. J. Hum. Genet. 45: 814-818, 1989.

[11441] 445.

[11442] Murray, S. S.; Deaven, L. L.; Burton, D. W.; O'Connor, D. T.;

[11443] Mellon, P. L.; Deftos, L. J.: The gene for human chromo-
granin A (CgA)
[11444] is located on chromosome 14. Biochem. Biophys. Res.
Commun. 142:
[11445] 141-146, 1987.

[11446] 446.

[11447] Nobels, F. R. E.; Kwekkeboom, D. J.; Coopmans, W.;
Schoenmakers,
[11448] C. H. H.; Lindemans, J.; De Herder, W. W.; Krenning, E. P.;
Bouillon,
[11449] R.; Lamberts, S. W. J.: Chromogranin A as serum marker
for neuroendocrine
[11450] neoplasia: comparison with neuron-specific enolase and
the alpha-subunit
[11451] of glycoprotein hormones. J. Clin. Endocr. Metab. 82:
2622-2628,
[11452] 1997.

[11453] 447.

[11454] O'Connor, D. T.; Deftos, L. J.: Secretion of chromogranin A by

[11455] peptide-producing endocrine neoplasms. New Eng. J. Med. 314: 1145-1151,

[11456] 1986.

[11457] 448.

[11458] Simon-Chazottes, D.; Wu, H.; Parmer, R. J.; Rozansky, D. J.; Szpirer,

[11459] J.; Levan, G.; Kurtz, T. W.; Szpirer, C.; Guenet, J. L.; O'Connor,

[11460] D. T.: Assignment of the chromogranin A (Chga) locus to homologous

[11461] regions on mouse chromosome 12 and rat chromosome 6. Genomics 17:

[11462] 252-255, 1993.

[11463] 449.

[11464] Wu, H.-J.; Rozansky, D. J.; Parmer, R. J.; Gill, B. M.; O'Connor,

[11465] D. T.: Structure and function of the chromogranin A gene: clues to

[11466] evolution and tissue-specific expression. J. Biol. Chem. 266: 13130-13134,

[11467] 1991.

[11468] 450.

[11469] Benedum, U. M.; Lamouroux, A.; Konecki, D. S.; Rosa, P.; Hille,

[11470] A.; Baeuerle, P. A.; Frank, R.; Lottspeich, F.; Mallet, J.; Huttner,

[11471] W. B.: The primary structure of human secretogranin I (chromogranin

[11472] B): comparison with chromogranin A reveals homologous terminal domains

[11473] and a large intervening variable region. EMBO J. 6: 1203-1211,

[11474] 1987.

[11475] 451.

[11476] Higashiyama, S.; Lau, K.; Besner, G. E.; Abraham, J. A.; Klagsbrun,

[11477] M.: Structure of heparin-binding EGF-like growth factor: multiple

[11478] forms, primary structure, and glycosylation of the mature protein. J.

[11479] Biol. Chem. 267: 6205-6212, 1992.

[11480] 452.

[11481] Naglich, J. G.; Metherall, J. E.; Russell, D. W.; Eidels, L.:

[11482] Expression cloning of a diphtheria toxin receptor: identity with a

[11483] heparin-binding EGF-like growth factor precursor. Cell 69: 1051-1061,

[11484] 1992.

[11485] 453.

[11486] Pappenheimer, A. M., Jr.: Diphtheria toxin. Ann. Rev. Biochem. 46:

[11487] 69-94, 1977.

[11488] 454.

[11489] Pappenheimer, A. M., Jr.; Gill, D. M.: Diphtheria. Science 182:

[11490] 353-358, 1973.

[11491] 455.

[11492] Pathak, B. G.; Gilbert, D. J.; Harrison, C. A.; Luetteke, N. C.;

[11493] Chen, X.; Klagsbrun, M.; Plowman, G. D.; Copeland, N. G.;

Jenkins,

[11494] N. A.; Lee, D. C.: Mouse chromosomal location of three EGF receptor

[11495] ligands: amphiregulin (Areg), betacellulin (Btc), and hep-arin-binding

[11496] EGF (Hegfl). Genomics 28: 116-118, 1995.

[11497] 456.

[11498] Roberts, M.; Ruddle, F. H.: The Chinese hamster gene map: assignment

[11499] of four genes (DTS, PGM2, 6PGD, Eno1) to chromosome 2. Exp. Cell

[11500] Res. 127: 47-54, 1980.

[11501] 457.

[11502] Chabas, D.; Baranzini, S. E.; Mitchell, D.; Bernard, C. C. A.;

[11503] Rittling, S. R.; Denhardt, D. T.; Sobel, R. A.; Lock, C.; Karpuj,

[11504] M.; Pedotti, R.; Heller, R.; Oksenberg, J. R.; Steinman, L.: The

[11505] influence of the proinflammatory cytokine, osteopontin, on autoimmune

[11506] demyelinating disease. Science 294: 1731-1735, 2001.

[11507] 458.

[11508] Bowcock, A. M.; Hebert, J. M.; Christiano, A. M.; Wijsman, E.;

[11509] Cavalli-Sforza, L. L.; Boyd, C. D.: The pro alpha 1 (IV) collagen

[11510] gene is linked to the D13S3 locus at the distal end of human chromosome

[11511] 13q. Cytogenet. Cell Genet. 45: 234-236, 1987.

[11512] 459.

[11513] Bowcock, A. M.; Hebert, J. M.; Wijsman, E.; Gadi, I.; Cavalli-Sforza,

[11514] L. L.; Boyd, C. D.: High recombination between two physically close

[11515] human basement membrane collagen genes at the distal end of chromosome

[11516] 13q. Proc. Nat. Acad. Sci. 85: 2701-2705, 1988.

[11517] 460.

[11518] Boyd, C. D.; Weliky, K.; Toth-Fejel, S.; Deak, S. B.; Christiano,

[11519] A. M.; Mackenzie, J. W.; Sandell, L. J.; Tryggvason, K.; Magenis,

[11520] E.: The single copy gene coding for human alpha-1(IV) procollagen

[11521] is located at the terminal end of the long arm of chromosome 13. Hum.

[11522] Genet. 74: 121-125, 1986.

[11523] 461.

[11524] Brinker, J. M.; Gudas, L. J.; Loidl, H. R.; Wang, S.-Y.; Rosenbloom,

[11525] J.; Kefalides, N. A.; Myers, J. C.: Restricted homology between human

[11526] alpha-1 type IV and other procollagen chains. Proc. Nat. Acad. Sci. 82:

[11527] 3649-3653, 1985.

[11528] 462.

[11529] Burbelo, P. D.; Martin, G. R.; Yamada, Y.: Alpha-1(IV) and alpha-2(IV)

[11530] collagen genes are regulated by a bidirectional promoter and a shared

[11531] enhancer. Proc. Nat. Acad. Sci. 85: 9679-9682, 1988.

[11532] 463.

[11533] Crouch, E.; Sage, H.; Bornstein, P.: Structural basis for ap-

parent
[11534] heterogeneity of collagens in human basement membranes: type IV procollagen
[11535] contains two distinct chains. Proc. Nat. Acad. Sci. 77: 745-749,
[11536] 1980.

[11537] 464.

[11538] Cutting, G. R.; Kazazian, H. H., Jr.; Antonarakis, S. E.; Killen,
[11539] P. D.; Yamada, Y.; Francomano, C. A.: Macrorestriction analysis maps
[11540] COL4A1 and COL4A2 collagen genes within a 400 kb region on chromosome
[11541] 13q34. (Abstract) Am. J. Hum. Genet. 41: A163, 1987.

[11542] 465.

[11543] Cutting, G. R.; Kazazian, H. H., Jr.; Antonarakis, S. E.; Killen,
[11544] P. D.; Yamada, Y.; Francomano, C. A.: Macrorestriction mapping of
[11545] COL4A1 and COL4A2 collagen genes on human chromosome 13q34. Genomics 3:
[11546] 256-263, 1988.

[11547] 466.

[11548] Emanuel, B. S.; Sellinger, B. T.; Gudas, L. J.; Myers, J. C.:

[11549] Localization of the human procollagen alpha-1(IV) gene to chromosome

[11550] 13q34 by in situ hybridization. Am. J. Hum. Genet. 38: 38-44, 1986.

[11551] 467.

[11552] Hebert, J. M.; Bowcock, A. M.; Wijsman, E.; Gadi, I.; Boyd, C.;

[11553] Cavalli-Sforza, L. L.: The genes for pro-alpha-1 (IV) collagen, pro-alpha-2

[11554] (IV) collagen and the D13S3 locus are linked at 13q34. (Abstract) Am.

[11555] J. Hum. Genet. 41: A169, 1987.

[11556] 468.

[11557] Kuhn, K.: Personal Communication. Munich, Germany 1/7/1982.

[11558] 469.

[11559] Mayne, R.; Wiedemann, H.; Irwin, M. H.; Sanderson, R. D.; Fitch,

[11560] J. M.; Linsenmayer, T. F.; Kuhn, K.: Monoclonal antibodies

against

[11561] chicken type IV and V collagens: electron microscopic mapping of the

[11562] epitopes after rotary shadowing. J. Cell Biol. 98: 1637-1644, 1984.

[11563] 470.

[11564] Pihlajaniemi, T.; Tryggvason, K.; Myers, J. C.; Kurkinen, M.;

[11565] Lebo, R.; Cheung, M.-C.; Prockop, D. J.; Boyd, C. D.: cDNA clones

[11566] coding for the pro-alpha-1(IV) chain of human type IV procollagen

[11567] reveal an unusual homology of amino acid sequences in two halves of

[11568] the carboxyl terminal domain. J. Biol. Chem. 260: 7681-7687, 1985.

[11569] 471.

[11570] Poschl, E.; Pollner, R.; Kuhn, K.: The genes for the alpha-1(IV)

[11571] and alpha-2(IV) chains of human basement membrane collagen type IV

[11572] are arranged head-to-head and separated by a bidirectional promoter

[11573] of unique structure. EMBO J. 7: 2687-2695, 1988.

[11574] 472.

[11575] Soininen, R.; Chow, L.; Kurkinen, M.; Tryggvason, K.; Prockop,

[11576] D. J.: The gene for the alpha-1(IV) chain of human type IV procollagen:

[11577] the exon structures do not coincide with the two structural subdomains

[11578] in the globular carboxy-terminus of the protein. EMBO J. 5: 2821-2823,

[11579] 1986.

[11580] 473.

[11581] Soininen, R.; Huotari, M.; Hostikka, S. L.; Prockop, D. J.; Tryggvason,

[11582] K.: The structural genes for alpha-1 and alpha-2 chains of human

[11583] type IV collagen are divergently encoded on opposite DNA strands and

[11584] have an overlapping promoter region. J. Biol. Chem. 263: 17217-17220,

[11585] 1988.

[11586] 474.

[11587] Soininen, R.; Tikka, L.; Chow, L.; Pihlajaniemi, T.; Kurki-
nen,
[11588] M.; Prockop, D. J.; Boyd, C. D.; Tryggvason, K.: Large in-
trons in
[11589] the 3-prime end of the gene for the pro-alpha1(IV) chain
of human
[11590] basement membrane collagen. Proc. Nat. Acad. Sci. 83:
1568-1572,
[11591] 1986.

[11592] 475.

[11593] Karoui, M.; Hofmann-Radvanyi, H.; Zimmermann, U.; Cou-
velard, A.;
[11594] Degott, C.; Faridoni-Laurens, L.; Ahomadegbe, J.-C.;
Gazzeri, S.;
[11595] Brambilla, E.; Clerici, T.; Charbonnier, P.; Tresallet, C.; and
8
[11596] others: No evidence of somatic FGFR3 mutation in various
types of
[11597] carcinoma. Oncogene 20: 5059-5061, 2001.

[11598] 476.

[11599] Keegan, K.; Johnson, D. E.; Williams, L. T.; Hayman, M. J.: Isolation
[11600] of an additional member of the fibroblast growth factor receptor family,
[11601] FGFR-3. Proc. Nat. Acad. Sci. 88: 1095-1099, 1991.

[11602] 477.

[11603] Kimura, T.; Suzuki, H.; Ohashi, T.; Kiyota, H.; Eto, Y.: The
[11604] incidence of thanatophoric dysplasia mutations in FGFR3 bladder carcinomas. Cancer 92:
[11605] 2555-2561, 2001.

[11606] 478.

[11607] Kitoh, H.; Brodie, S. G.; Kupke, K. G.; Lachman, R. S.; Wilcox,
[11608] W. R.: Lys650-to-met substitution in the tyrosine kinase domain of
[11609] the fibroblast growth factor receptor gene causes thanatophoric dysplasia
[11610] type I. (Abstract) Hum. Mutat. 12: 362-363, 1998.

[11611] 479.

[11612] Lajeunie, E.; El Ghouzzi, V.; Le Merrer, M.; Munnich, A.; Bonaventure,

[11613] J.; Renier, D.: Sex related expressivity of the phenotype in coronal
[11614] craniosynostosis caused by the recurrent P250R FGFR3 mutation. J.
[11615] Med. Genet. 36: 9-13, 1999.

[11616] 480.

[11617] Lanning, R. W.; Brown, C. A.: An improved methodology for the
[11618] detection of the common mutation in the FGFR3 gene responsible for
[11619] achondroplasia. Hum. Mutat. 10: 496-499, 1997.

[11620] 481.

[11621] Li, C.; Chen, L.; Iwata, T.; Kitagawa, M.; Fu, X.-Y.; Deng, C.-X.
[11622] : A Lys644Glu substitution in fibroblast growth factor receptor 3
[11623] (FGFR3) causes dwarfism in mice by activation of STATs and ink4 cell
[11624] cycle inhibitors. Hum. Molec. Genet. 8: 35-44, 1999.

[11625] 482.

[11626] Lowry, R. B.; Jabs, E. W.; Graham, G. E.; Gerritsen, J.; Flem-

ing,

[11627] J.: Syndrome of coronal craniosynostosis, Klippel-Feil anomaly, and

[11628] Sprengel shoulder with and without pro250arg mutation in the FGFR3

[11629] gene. Am. J. Med. Genet. 104: 112-119, 2001.

[11630] 483.

[11631] McKusick, V. A.; Kelly, T. E.; Dorst, J. P.: Observations suggesting

[11632] allelism of the achondroplasia and hypochondroplasia genes. J. Med.

[11633] Genet. 10: 11-16, 1973.

[11634] 484.

[11635] Moloney, D. M.; Slaney, S. F.; Oldridge, M.; Wall, S. A.; Sahlin,

[11636] P.; Stenman, G.; Wilkie, A. O. M.: Exclusive paternal origin of new

[11637] mutations in Apert syndrome. Nature Genet. 13: 48-53, 1996.

[11638] 485.

[11639] Moloney, D. M.; Wall, S. A.; Ashworth, G. J.; Oldridge, M.;

Glass,
[11640] I. A.; Francomano, C. A.; Muenke, M.; Wilkie, A. O. M.: Prevalence
[11641] of pro250arg mutation of fibroblast growth factor receptor 3 in coronal
[11642] craniosynostosis. Lancet 349: 1059-1062, 1997.

[11643] 486.

[11644] Monsonego-Ornan, E.; Adar, R.; Feferman, T.; Segev, O.; Yayon,
[11645] A.: The transmembrane mutation G380R in fibroblast growth factor
[11646] receptor 3 uncouples ligand-mediated receptor activation from down-regulation. Molec.
[11647] Cell. Biol. 20: 516-522, 2000.

[11648] 487.

[11649] Mortier, G.; Nuytinck, L.; Craen, M.; Renard, J.-P.; Leroy, J.
[11650] G.; De Paepe, A.: Clinical and radiographic features of a family
[11651] with hypochondroplasia owing to a novel asn540ser mutation in the
[11652] fibroblast growth factor receptor 3 gene. J. Med. Genet.

37: 220-224,

[11653] 2000.

[11654] 488.

[11655] Muenke, M.; Gripp, K. W.; McDonald-McGinn, D. M.; Gaudenz, K.;

[11656] Whitaker, L. A.; Bartlett, S. P.; Markowitz, R. I.; Robin, N. H.;

[11657] Nwokoro, N.; Mulvihill, J. J.; Losken, W.; Mulliken, J. B.; Guttmacher,

[11658] A. E.; Wilroy, R. S.; Clarke, L. A.; Hollway, G.; Ades, L. C.; Haan,

[11659] E. A.; Mulley, J. C.; Cohen, M. M., Jr.; Bellus, G. A.; Francomano,

[11660] C. A.; Moloney, D. M.; Wall, S. A.; Wilkie, A. O. M.; Zackai, E. H.

[11661] : A unique point mutation in the fibroblast growth factor receptor

[11662] 3 gene (FGFR3) defines a new craniosynostosis syndrome. Am. J. Hum.

[11663] Genet. 60: 555-564, 1997.

[11664] 489.

[11665] Naski, M. C.; Wang, Q.; Xu, J.; Ornitz, D. M.: Graded acti-

vation
[11666] of fibroblast growth factor receptor 3 by mutations caus-
ing achondroplasia
[11667] and thanatophoric dysplasia. Nature Genet. 13: 233-237,
1996.
[11668] 490.

[11669] Nishimura, G.; Fukushima, Y.; Ohashi, H.; Ikegawa, S.:
Atypical
[11670] radiological findings in achondroplasia with uncommon
mutation of
[11671] the fibroblast growth factor receptor-3 (FGFR-3) gene (gly
to cys
[11672] transition at codon 375). (Letter) Am. J. Med. Genet. 59:
393-395,
[11673] 1995.

[11674] 491.

[11675] Nishimura, G.; Takada, F.: Another observation of atypical
radiologic
[11676] findings in achondroplasia not due to a common mutation
of the FGFR-3
[11677] gene: reply to Dr. Gorlin. (Letter) Am. J. Med. Genet. 70:
203-205,

[11678] 1997.

[11679] 492.

[11680] Orlow, S. J.: Cutaneous findings in craniofacial malforma-
tion
[11681] syndromes. Arch. Derm. 128: 1379-1386, 1992.

[11682] 493.

[11683] Passos-Bueno, M. R.; Wilcox, W. R.; Jabs, E. W.; Sertie, A.
L.;
[11684] Alonso, L. G.; Kitoh, H.: Clinical spectrum of fibroblast
growth
[11685] factor receptor mutations. Hum. Mutat. 14: 115-125,
1999.
[11686] 494.

[11687] Paznekas, W. A.; Cunningham, M. L.; Howard, T. D.; Korf,
B. R.;
[11688] Lipson, M. H.; Grix, A. W.; Feingold, M.; Goldberg, R.;
Borochowitz,
[11689] Z.; Aleck, K.; Mulliken, J.; Yin, M.; Jabs, E. W.: Genetic het-
erogeneity
[11690] of Saethre-Chotzen syndrome, due to TWIST and FGFR
mutations. Am.

[11691] J. Hum. Genet. 62: 1370-1380, 1998.

[11692] 495.

[11693] Perez-Castro, A. V.; Wilson, J.; Altherr, M. R.: Genomic or-
ganization
[11694] of the human fibroblast growth factor receptor 3 (FGFR3)
gene and
[11695] comparative sequence analysis with the mouse Fgfr3
gene. Genomics 41:
[11696] 10-16, 1997.

[11697] 496.

[11698] Pokharel, R. K.; Alimsardjono, H.; Takeshima, Y.; Naka-
mura, H.;
[11699] Naritomi, K.; Hirose, S.; Onishi, S.; Matsuo, M.: Japanese
cases
[11700] of type 1 thanatophoric dysplasia exclusively carry a C to
T transition
[11701] at nucleotide 742 of the fibroblast growth factor receptor
3 gene. Biochem.
[11702] Biophys. Res. Commun. 227: 236-239, 1996.

[11703] 497.

[11704] Prinos, P.; Costa, T.; Sommer, A.; Kilpatrick, M. W.;

Tsipouras,

[11705] P.: A common FGFR3 gene mutation in hypochondroplasia. Hum. Molec.

[11706] Genet. 4: 2097-2101, 1995.

[11707] 498.

[11708] Prinster, C.; Carrera, P.; Del Maschio, M.; Weber, G.; Maghnie,

[11709] M.; Vigone, M. C.; Mora, S.; Tonini, G.; Rigon, F.; Beluffi, G.; Severi,

[11710] F.; Chiumello, G.; Ferrari, M.: Comparison of clinical-radiological

[11711] and molecular findings in hypochondroplasia. Am. J. Med. Gen. 75:

[11712] 109-112, 1998.

[11713] 499.

[11714] Ramaswami, U.; Rumsby, G.; Hindmarsh, P. C.; Brook, C. G. D.:

[11715] Genotype and phenotype in hypochondroplasia. J. Pediat. 133: 99-102,

[11716] 1998.

[11717] 500.

[11718] Rasmussen, T.; Hudlebusch, H. R.; Knudsen, L. M.; Johnsen, H.
[11719] E.: FGFR3 dysregulation in multiple myeloma: frequency and prognostic
[11720] relevance. Br. J. Haematol. 117: 626-628, 2002.

[11721] 501.

[11722] Fritsch, P.; Honigsmann, H.; Jaschke, E.: Epidermolytic hereditary
[11723] palmoplantar keratoderma. Brit. J. Derm. 99: 561-568, 1978.
[11724] 502.

[11725] Jutel, M.; Watanabe, T.; Klunker, S.; Akdis, M.; Thomet, O. A.
[11726] R.; Malolepszy, J.; Zak-Nejmark, T.; Koga, R.; Kobayashi, T.; Blaser,
[11727] K.; Akdis, C. A.: Histamine regulates T-cell and antibody responses
[11728] by differential expression of H1 and H2 receptors. Nature 413: 420-425,
[11729] 2001.

[11730] 503.

[11731] Doenecke, D.; Tonjes, R.: Differential distribution of lysine

[11732] and arginine residues in the closely related histones H1 and H5. Analysis

[11733] of a human H1 gene. J. Molec. Biol. 187: 461-464, 1986.

[11734] 504.

[11735] Gamborg Nielsen, P.: Two different clinical and genetic forms

[11736] of hereditary palmoplantar keratoderma in the northern-most county

[11737] of Sweden. Clin. Genet. 28: 361-366, 1985.

[11738] 505.

[11739] Goette, D. K.: Familial congenital epidermolytic hyperkeratosis

[11740] confined to the palms and soles. Sth. Med. J. 67: 1126-1128, 1974.

[11741] 506.

[11742] Greither, A.: Keratosis extremitatum hereditaria progrediens

[11743] mit dominantem Erbgang. Hautarzt 3: 198-203, 1952.

[11744] 507.

[11745] Hennies, H.-C.; Zehender, D.; Kunze, J.; Kuster, W.; Reis, A.

[11746] : Keratin 9 gene mutational heterogeneity in patients with epidermolytic

[11747] palmoplantar keratoderma. Hum. Genet. 93: 649-654, 1994.

[11748] 508.

[11749] Hennies, H. C.; Kuster, W.; Langbein, L.; Digweed, M.; Mischke,

[11750] D.; Franke, W. W.; Sperling, K.; Reis, A.: Keratin 9 gene mutations

[11751] in epidermolytic palmoplantar keratoderma (EPPK). (Abstract) Am.

[11752] J. Hum. Genet. 53 (suppl.): A211, 1993.

[11753] 509.

[11754] Klaus, S.; Weinstein, G. D.; Frost, P.: Localized epidermolytic

[11755] hyperkeratosis: a form of keratoderma of the palms and soles. Arch.

[11756] Derm. 101: 272-275, 1970.

[11757] 510.

[11758] Klintworth, G. K.; Anderson, I. F.: Tylosis palmaris et plantaris
[11759] familiaris associated with clinodactyly. S. Afr. Med. J. 35: 170-175,
[11760] 1961.

[11761] 511.

[11762] Kobayashi, S.; Tanaka, T.; Matsuyoshi, N.; Imamura, S.: Keratin
[11763] 9 point mutation in the pedigree of epidermolytic hereditary palmoplantar
[11764] keratoderma perturbs keratin intermediate filament network formation. FEBS
[11765] Lett. 386: 149-155, 1996.

[11766] 512.

[11767] Kuster, W.; Becker, A.: Indication for the identity of palmoplantar
[11768] keratoderma type Unna-Thost with type Vorner: Thost's family revisited
[11769] 110 years later. Acta Derm. Venerol. 72: 120-122, 1992.

[11770] 513.

[11771] Moriwaki, S.; Tanaka, T.; Horiguchi, Y.; Danno, K.; Ima-

mura, S.

[11772] : Epidermolytic hereditary palmoplantar keratoderma: histologic, ultrastructural,

[11773] protein-chemical, and DNA analysis in two patients. Arch. Derm. 124:

[11774] 555-559, 1988.

[11775] 514.

[11776] Nogita, T.; Nakagawa, H.; Ishibashi, Y.: Hereditary epidermolytic

[11777] palmoplantar keratoderma with knuckle pad-like lesions over the finger

[11778] joints. (Letter) Brit. J. Derm. 125: 496, 1991.

[11779] 515.

[11780] Reis, A.; Hennies, H.-C.; Langbein, L.; Digweed, M.; Mischke,

[11781] D.; Drechsler, M.; Schrock, E.; Royer-Pokora, B.; Franke, W. W.; Sperling,

[11782] K.; Kuster, W.: Keratin 9 gene mutations in epidermolytic palmoplantar

[11783] keratoderma (EPPK). Nature Genet. 6: 174-179, 1994.

[11784] 516.

[11785] Reis, A.; Kuster, W.; Eckardt, R.; Sperling, K.: Mapping of a

[11786] gene for epidermolytic palmoplantar keratoderma to the region of the

[11787] acidic keratin gene cluster at 17q12-q21. Hum. Genet. 90: 113-116,

[11788] 1992.

[11789] 517.

[11790] Rogaev, E. I.; Rogaeva, E. A.; Ginter, E. K.; Korovaitseva, G.

[11791] I.; Farrer, L. A.; Shlensky, A. B.; Pritkov, A. N.; Mordovtsev, V.

[11792] N.; St. George-Hyslop, P. H.: Identification of the genetic locus

[11793] for keratosis palmaris et plantaris on chromosome 17 near the RARA

[11794] and keratin type I genes. Nature Genet. 5: 158-162, 1993.

[11795] 518.

[11796] Rothnagel, J. A.; Wojcik, S.; Liefer, K. M.; Dominey, A. M.; Huber,

[11797] M.; Hohl, D.; Roop, D. R.: Mutations in the 1A domain of keratin

[11798] 9 in patients with epidermolytic palmoplantar kerato-

derma. J. Invest.

[11799] Derm. 104: 430-433, 1995.

[11800] 519.

[11801] Stevens, H. P.; Rustin, M. H. A.; MacDermot, K.: Keratosis punctata

[11802] palmaris et plantaris: an autosomal dominantly inherited keratoderma

[11803] associated with malignancy. (Abstract) J. Med. Genet. 31: 168 only,

[11804] 1994.

[11805] 520.

[11806] Thost, A.: Ueber erbliche Ichthyosis palmaris et plantaris cornea.

[11807] Dissertation: Heidelberg (pub.) 1880.

[11808] 521.

[11809] Torchard, D.; Blanchet-Bardon, C.; Serova, O.; Langbein, L.; Narod,

[11810] S.; Janin, N.; Goguel, A. F.; Bernheim, A.; Franke, W. W.; Lenoir,

[11811] G. M.; Feunteun, J.: Epidermolytic palmoplantar keratoderma cosegregates

[11812] with a keratin 9 mutation in a pedigree with breast and ovarian cancer. Nature
[11813] Genet. 6: 106-110, 1994.

[11814] 522.

[11815] Unna, P. G.: Ueber das Keratoma palmare et plantare hereditarium. Arch.
[11816] Derm. Syph. 15: 231-270, 1883.

[11817] 523.

[11818] Vorner, H.: Zur Kenntniss des Keratoma hereditarium palmare et
[11819] plantare. Arch. Derm. Syph. 56: 3-31, 1901.

[11820] 524.

[11821] Geurts, J. M. W.; Janssen, R. G. J. H.; van Greevenbroek, M. M.
[11822] J.; van der Kallen, C. J. H.; Cantor, R. M.; Bu, X.; Aouizerat, B.
[11823] E.; Allayee, H.; Rotter, J. I.; de Bruin, T. W. A.: Identification
[11824] of TNFRSF1B as a novel modifier gene in familial combined hyperlipidemia. Hum.
[11825] Molec. Genet. 9: 2067-2074, 2000.

[11826] 525.

[11827] Hatzivassiliou, G.; Miller, I.; Takizawa, J.; Palanisamy, N.; Rao,

[11828] P. H.; Iida, S.; Tagawa, S.; Taniwaki, M.; Russo, J.; Neri, A.; Cattoretti,

[11829] G.; Clynes, R.; Mendelsohn, C.; Chaganti, R. S. K.; Dalla-Favera,

[11830] R.: IRTA1 and IRTA2, novel immunoglobulin superfamily receptors expressed

[11831] in B cells and involved in chromosome 1q21 abnormalities in B cell

[11832] malignancy. Immunity 14: 277-289, 2001.

[11833] 526.

[11834] Mollereau, C.; Muscatelli, F.; Mattei, M.-G.; Vassart, G.; Parmentier,

[11835] M.: The high-affinity interleukin 8 receptor gene (IL8RA) maps to

[11836] the 2q33-q36 region of the human genome: cloning of a pseudogene (IL8RBP)

[11837] for the low-affinity receptor. Genomics 16: 248-251, 1993.

[11838] 527.

[11839] Morris, S. W.; Nelson, N.; Valentine, M. B.; Shapiro, D. N.; Look,
[11840] A. T.; Kozlosky, C. J.; Beckmann, M. P.; Cerretti, D. P.: Assignment
[11841] of the genes encoding human interleukin-8 receptor types 1 and 2 and
[11842] an interleukin-8 receptor pseudogene to chromosome 2q35. Genomics 14:
[11843] 685-691, 1992.

[11844] 528.

[11845] Palter, S. F.; Mulayim, N.; Senturk, L.; Arici, A.: Interleukin-8
[11846] in the human fallopian tube. J. Clin. Endocr. Metab. 86: 2660-2667,
[11847] 2001.

[11848] 529.

[11849] Dixon, M. J.; Gazzard, J.; Chaudhry, S. S.; Sampson, N.; Schulte,
[11850] B. A.; Steel, K. P.: Mutation of the Na-K-Cl co-transporter gene
[11851] Slc12a2 results in deafness in mice. Hum. Molec. Genet. 8: 1579-1584,

[11852] 1999.

[11853] 530.

[11854] Kainulainen, K.; Karttunen, L.; Puhakka, L.; Sakai, L.; Peltonen,

[11855] L.: Mutations in the fibrillin gene responsible for dominant ectopia

[11856] lentis and neonatal Marfan syndrome. Nature Genet. 6: 64-69, 1994.

[11857] 531.

[11858] Li, X.; Pereira, L.; Zhang, H.; Sanguineti, C.; Ramirez, F.; Bonadio,

[11859] J.; Francke, U.: Fibrillin genes map to regions of conserved mouse/human

[11860] synteny on mouse chromosomes 2 and 18. Genomics 18: 667-672, 1993.

[11861] 532.

[11862] Soufir, N.; Avril, M. F.; Chompret, A.; Demenais, F.; Bombled,

[11863] J.; Spatz, A.; Stoppa-Lyonnet, D.; Benard, J.; Bressac-de Paillerets,

[11864] B.: Prevalence of p16 and CDK4 germline mutations in 48 melanoma-prone

[11865] families in France: the French Familial Melanoma Study Group. Hum.

[11866] Molec. Genet. 7: 209-216, 1998.

[11867] 533.

[11868] Dedhar, S.; Rennie, P. S.; Shago, M.; Hagesteijn, C.-Y. L.; Yang,

[11869] H.; Filmus, J.; Hawley, R. G.; Bruchovsky, N.; Cheng, H.; Matusik,

[11870] R. J.; Giguere, V.: Inhibition of nuclear hormone receptor activity

[11871] by calreticulin. Nature 367: 480-483, 1994.

[11872] 534.

[11873] Bastiaens, M. T.; ter Huurne, J. A. C.; Kielich, C.; Gruis, N.

[11874] A.; Westendorp, R. G. J.; Vermeer, B. J.; Bavinck, J. N. B.; Leiden

[11875] Skin Cancer Study Team: Melanocortin-1 receptor gene variants determine

[11876] the risk of nonmelanoma skin cancer independently of fair skin and

[11877] red hair. Am. J. Hum. Genet. 68: 884-894, 2001.

[11878] 535.

[11879] Rose, P. M.; Fernandes, P.; Lynch, J. S.; Frazier, S. T.; Fisher,
[11880] S. M.; Kodukula, K.; Kienzle, B.; Seethala, R.: Cloning and functional
[11881] expression of a cDNA encoding a human type 2 neu-ropeptide Y receptor. J.
[11882] Biol. Chem. 270: 22661-22664, 1995.

[11883] 536.

[11884] Hovnanian, A.; Rebouillat, D.; Mattei, M.-G.; Levy, E. R.; Marie,
[11885] I.; Monaco, A. P.; Hovanessian, A. G.: The human 2-prime,5-prime-oligoadenylate
[11886] synthetase locus is composed of three distinct genes clustered on
[11887] chromosome 12q24.2 encoding the 100-, 69-, and 40-kDa forms. Genomics 52:
[11888] 267-277, 1998.

[11889] 537.

[11890] Frank, M. B.: Personal Communication. Oklahoma City, Okla.
[11891] 6/3/1994.

[11892] 538.

[11893] Itoh, K.; Itoh, Y.; Frank, M. B.: Protein heterogeneity in the

[11894] human Ro/SSA ribonucleoproteins: the 52- and 60-kD Ro/SSA autoantigens

[11895] are encoded by separate genes. J. Clin. Invest. 87: 177-186, 1991.

[11896] 539.

[11897] McCauliffe, D. P.; Lux, F. A.; Lieu, T.-S.; Sanz, I.; Hanke, J.;

[11898] Newkirk, M. M.; Bachinski, L. L.; Itoh, Y.; Siciliano, M. J.; Reichlin,

[11899] M.; Sontheimer, R. D.; Capra, J. D.: Molecular cloning, expression,

[11900] and chromosome 19 localization of a human Ro/SS-A autoantigen. J.

[11901] Clin. Invest. 85: 1379-1391, 1990.

[11902] 540.

[11903] McCauliffe, D. P.; Zappi, E.; Lieu, T.-S.; Michalak, M.; Sontheimer,

[11904] R. D.; Capra, J. D.: A human Ro/SS-A autoantigen is the homologue

[11905] of calreticulin and is highly homologous with onchocercal

RAL-1 antigen
[11906] and an aplysia 'memory molecule.'. J. Clin. Invest. 86: 332-335,
[11907] 1990.

[11908] 541.

[11909] Orth, T.; Dorner, T.; Meyer Zum Buschenfelde, K.-H.; Mayet, W.-J.
[11910] : Complete congenital heart block is associated with increased autoantibody
[11911] titers against calreticulin. Europ. J. Clin. Invest. 26: 205-215,
[11912] 1996.

[11913] 542.

[11914] Rooke, K.; Briquet-Laugier, V.; Xia, Y.-R.; Lusis, A. J.; Doolittle,
[11915] M. H.: Mapping of the gene for calreticulin (Calr) to mouse chromosome
[11916] 8. Mammalian Genome 8: 870-871, 1997.

[11917] 543.

[11918] Frank, M. B.; Itoh, K.; Fujisaku, A.; Pontarotti, P.; Mattei, M.-G.;

[11919] Neas, B. R.: The mapping of the human 52-kD Ro/SSA autoantigen gene
[11920] to human chromosome 11, and its polymorphisms. Am. J. Hum. Genet. 52:
[11921] 183-191, 1993.

[11922] 544.

[11923] Frank, M. B.; Mattei, M.-G.: Mapping of the human 60000 M(r) Ro/SSA
[11924] locus: the genes for three Ro/SSA autoantigens are located on separate
[11925] chromosomes. Immunogenetics 39: 428-431, 1994.

[11926] 545.

[11927] Schoenlebe, J.; Buyon, J. P.; Zitelli, B. J.; Friedman, D.; Greco,
[11928] M. A.; Knisely, A. S.: Neonatal hemochromatosis associated with maternal
[11929] autoantibodies against Ro/SS-A and La/SS-B ribonucleoproteins. Am.
[11930] J. Dis. Child. 147: 1072-1075, 1993.

[11931] 546.

[11932] Maclaren, N. K.; Riley, W. J.: Inherited susceptibility to au-

toimmune

[11933] Addison's disease is linked to human leukocyte antigens-DR3 and/or

[11934] DR4, except when associated with type I autoimmune polyglandular syndrome. J.

[11935] Clin. Endocr. Metab. 62: 455-459, 1986.

[11936] 547.

[11937] Wang, Y. A.; Elson, A.; Leder, P.: Loss of p21 increases sensitivity

[11938] to ionizing radiation and delays the onset of lymphoma in atm-deficient

[11939] mice. Proc. Nat. Acad. Sci. 94: 14590-14595, 1997.

[11940] 548.

[11941] Zakut, R.; Givol, D.: The tumor suppression function of p21(Waf)

[11942] is contained in its N-terminal half ('half-WAF'). Oncogene 11: 393-395,

[11943] 1995.

[11944] 549.

[11945] Boden, P.; Hall, M. D.; Hughes, J.: Cholecystokinin recep-tors. Cell.

[11946] Molec. Neurobiol. 15: 545-559, 1995.

[11947] 550.

[11948] de Weerth, A.; Pisegna, J. R.; Huppi, K.; Wank, S. A.: Molecular

[11949] cloning, functional expression and chromosomal localization of the

[11950] human cholecystokinin type A receptor. Biochem. Biophys. Res. Commun. 194:

[11951] 811-818, 1993.

[11952] 551.

[11953] Funakoshi, A.; Miyasaka, K.; Shinozaki, H.; Masuda, M.; Kawanami,

[11954] T.; Takata, Y.; Kono, A.: An animal model of congenital defect of

[11955] gene expression of cholecystokinin (CCK)-A receptor. Biochem. Biophys.

[11956] Res. Commun. 210: 787-796, 1995.

[11957] 552.

[11958] Hamann, A.; Busing, B.; Munzberg, H.; de Weerth, A.; Hinney, A.;

[11959] Mayer, H.; Siegfried, W.; Hebebrand, J.; Greten, H.: Mis-

sense variants
[11960] in the human cholecystokinin type A receptor gene: no evidence for
[11961] association with early-onset obesity. Horm. Metab. Res. 31: 287-288,
[11962] 1999.

[11963] 553.

[11964] Huppi, K.; Siwarski, D.; Pisegna, J. R.; Wank, S.: Chromo-somal
[11965] localization of the gastric and brain receptors for chole-cystokinin
[11966] (CCKAR and CCKBR) in human and mouse. Genomics 25: 727-729, 1995.
[11967] 554.

[11968] Inoue, H.; Iannotti, C. A.; Welling, C. M.; Veile, R.; Donis-Keller,
[11969] H.; Permutt, M. A.: Human cholecystokinin type A receptor gene: cytogenetic
[11970] localization, physical mapping, and identification of two missense
[11971] variants in patients with obesity and non-in-sulin-dependent diabetes

[11972] mellitus (NIDDM). Genomics 42: 331-335, 1997.

[11973] 555.

[11974] Marchal-Victorion, S.; Vionnet, N.; Escrieut, C.; Dematos,
F.;

[11975] Dina, C.; Dufresne, M.; Vaysse, N.; Pradayrol, L.; Froguel,
P.; Fourmy,

[11976] D.: Genetic, pharmacological and functional analysis of
cholecystokinin-1

[11977] and cholecystokinin-2 receptor polymorphism in type 2
diabetes and

[11978] obese patients. Pharmacogenetics 12: 23-30, 2002.

[11979] 556.

[11980] Miller, L. J.; Holicky, E. L.; Ulrich, C. D.; Wieben, E. D.: Ab-
normal

[11981] processing of the human cholecystokinin receptor gene in
association

[11982] with gallstones and obesity. Gastroenterology 109:
1375-1380, 1995.

[11983] 557.

[11984] Samuelson, L. C.; Isakoff, M. S.; Lacourse, K. A.: Localiza-
tion

[11985] of the murine cholecystokinin A and B receptor genes. Mammalian Genome 6:
[11986] 242-246, 1995.

[11987] 558.

[11988] Ulrich, C. D.; Ferber, I.; Holicky, E.; Hadac, E.; Buell, G.;

[11989] Miller, L. J.: Molecular cloning and functional expression of the
[11990] human gallbladder cholecystokinin A receptor. Biochem. Biophys. Res.
[11991] Commun. 193: 204-211, 1993.

[11992] 559.

[11993] Beinborn, M.; Lee, Y.-M.; McBride, E. W.; Quinn, S. M.; Kopin,
[11994] A. S.: A single amino acid of the cholecystokinin-B/gastrin receptor
[11995] determines specificity for non-peptide antagonists. Na-ture 362:
[11996] 348-350, 1993.

[11997] 560.

[11998] Lee, Y.-M.; Beinborn, M.; McBride, E. W.; Lu, M.; Ko-lakowski, L.

[11999] F., Jr.; Kopin, A. S.: The human brain cholecystokinin-B/gastrin
[12000] receptor: cloning and characterization. J. Biol. Chem. 268: 8164-8169,
[12001] 1993.

[12002] 561.

[12003] Pisegna, J. R.; de Weerth, A.; Huppi, K.; Wank, S. A.: Molecular
[12004] cloning of the human brain and gastric cholecystokinin receptor: structure,
[12005] functional expression and chromosomal localization. Biochem. Biophys.
[12006] Res. Commun. 189: 296-303, 1992.

[12007] 562.

[12008] Samuelson, L. C.; Isakoff, M. S.; Lacourse, K. A.: Localiza-tion
[12009] of the murine cholecystokinin A and B receptor genes. Mammalian
[12010] Genome 6: 242-246, 1995.

[12011] 563.

[12012] Zimonjic, D. B.; Popescu, N. C.; Matsui, T.; Ito, M.; Chihara,

[12013] K.: Localization of the human cholecystokinin-B/gastrin receptor
[12014] gene (CCKBR) to chromosome 11p15.5-p15.4 by fluorescence in situ hybridization.
[12015] Cytogenet. Cell Genet. 65: 184-185, 1994.

[12016] 564.

[12017] Butkowski, R. J.; Langeveld, J. P. M.; Wieslander, J.; Hamilton,
[12018] J.; Hudson, B. G.: Localization of the Goodpasture epitope to a novel
[12019] chain of basement membrane collagen. J. Biol. Chem. 262: 7874-7877,
[12020] 1987.

[12021] 565.

[12022] Butkowski, R. J.; Wieslander, J.; Kleppel, M.; Michael, A. F.;

[12023] Fish, A. J.: Basement membrane collagen in the kidney: regional localization
[12024] of novel chains related to collagen IV. Kidney Int. 35: 1195-1202,
[12025] 1989.

[12026] 566.

[12027] Cosgrove, D.; Meehan, D. T.; Grunkemeyer, J. A.; Kornak, J. M.;

[12028] Sayers, R.; Hunter, W. J.; Samuelson, G. C.: Collagen COL4A3 knockout:

[12029] a mouse model for autosomal Alport syndrome. Genes Dev. 10: 2981-2992,

[12030] 1996.

[12031] 567.

[12032] Finielz, P.; Cartault, F.; Chuet, C.; Genin, R.: Alport syn-drome

[12033] in Reunion Island: phenotypic heterogeneity of the reces-sive-autosomal

[12034] form. (Letter) Nephron 79: 237 only, 1998.

[12035] 568.

[12036] Han, J.; Ohno, N.; Pasco, S.; Monboisse, J.-C.; Borel, J. P.; Kefalides,

[12037] N. A.: A cell binding domain from the alpha-3 chain of type IV collagen

[12038] inhibits proliferation of melanoma cells. J. Biol. Chem. 272: 20395-20401,

[12039] 1997.

[12040] 569.

[12041] Hellmark, T.; Burkhardt, H.; Wieslander, J.: Goodpasture disease:

[12042] characterization of a single conformational epitope as the target

[12043] of pathogenic autoantibodies. J. Biol. Chem. 274: 25862-25868, 1999.

[12044] 570.

[12045] Hudson, B. G.; Kalluri, R.; Gunwar, S.; Weber, M.; Ballester, F.;

[12046] Hudson, J. K.; Noelken, M. E.; Sarras, M.; Richardson, W. R.; Saus,

[12047] J.; Abrahamson, D. R.; Glick, A. D.; Haralson, M. A.; Helderman, J.

[12048] H.; Stone, W. J.; Jacobson, H. R.: The pathogenesis of Alport syndrome

[12049] involves type IV collagen molecules containing the alpha-3(IV) chain:

[12050] evidence from anti-GBM nephritis after renal transplantation. Kidney

[12051] Int. 42: 179-187, 1992.

[12052] 571.

[12053] Kalluri, R.; Gattone, V. H., II; Noelken, M. E.; Hudson, B. G.

[12054] : The alpha-3 chain of type IV collagen induces autoimmune Goodpasture

[12055] syndrome. Proc. Nat. Acad. Sci. 91: 6201-6205, 1994.

[12056] 572.

[12057] Lemmink, H. H.; Schroder, C. H.; Monners, L. A. H.; Smeets, H.

[12058] J. M.: The clinical spectrum of type IV collagen mutations. Hum.

[12059] Mutat. 9: 477-499, 1997.

[12060] 573.

[12061] Lu, W.; Phillips, C. L.; Killen, P. D.; Hlaing, T.; Harrison,

[12062] W. R.; Elder, F. F. B.; Miner, J. H.; Overbeek, P. A.; Meisler, M.

[12063] H.: Insertional mutation of the collagen genes Col4a3 and Col4a4

[12064] in a mouse model of Alport syndrome. Genomics 61: 113-124, 1999.

[12065] 574.

[12066] Maeshima, Y.; Colorado, P. C.; Kalluri, R.: Two RGD-independent

[12067] alpha-V-beta-3 integrin binding sites on tumstatin regulate distinct

[12068] anti-tumor properties. J. Biol. Chem. 275: 23745-23750, 2000.

[12069] 575.

[12070] Maeshima, Y.; Colorado, P. C.; Torre, A.; Holthaus, K. A.; Grunkemeyer,

[12071] J. A.; Ericksen, M. B.; Hopfer, H.; Xiao, Y.; Stillman, I. E.; Kalluri,

[12072] R.: Distinct antitumor properties of a type IV collagen domain derived

[12073] from basement membrane. J. Biol. Chem. 275: 21340-21348, 2000.

[12074] 576.

[12075] Maeshima, Y.; Sudhakar, A.; Lively, J. C.; Ueki, K.; Kharbanda,

[12076] S.; Kahn, C. R.; Sonenberg, N.; Hynes, R. O.; Kalluri, R.: Tumstatin,

[12077] an endothelial cell-specific inhibitor of protein synthesis. Science 295:

[12078] 140-143, 2002.

[12079] 577.

[12080] Maeshima, Y.; Yerramalla, U. L.; Dhanabal, M.; Holthaus, K. A.;

[12081] Barbashov, S.; Kharbanda, S.; Reimer, C.; Manfredi, M.; Dickerson,

[12082] W. M.; Kalluri, R.: Extracellular matrix-derived peptide binds to

[12083] alpha-V-beta-3 integrin and inhibits angiogenesis. J. Biol. Chem. 276:

[12084] 31959-31968, 2001.

[12085] 578.

[12086] Mochizuki, T.; Lemmink, H. H.; Mariyama, M.; Antignac, C.; Gubler,

[12087] M.-C.; Pirson, Y.; Verellen-Dumoulin, C.; Chan, B.; Schroder, C. H.;

[12088] Smeets, H. J.; Reeders, S. T.: Identification of mutations in the

[12089] alpha-3(IV) and alpha-4(IV) collagen genes in autosomal recessive

[12090] Alport syndrome. Nature Genet. 8: 77-81, 1994.

[12091] 579.

[12092] Morrison, K. E.; Mariyama, M.; Yang-Feng, T. L.; Reeders, S. T.

[12093] : Sequence and localization of a partial cDNA encoding the human alpha3

[12094] chain of type IV collagen. Am. J. Hum. Genet. 49: 545-554, 1991.

[12095] 580.

[12096] Quinones, S.; Bernal, D.; Garcia-Sogo, M.; Elena, S. F.; Saus,

[12097] J.: Exon/intron structure of the human alpha-3(IV) gene encompassing

[12098] the Goodpasture antigen (alpha-3(IV)NC1): identification of a potentially

[12099] antigenic region at the triple helix/NC1 domain junction. J. Biol.

[12100] Chem. 267: 19780-19784, 1992.

[12101] 581.

[12102] Saus, J.; Wieslander, J.; Langeveld, J. P. M.; Quinones, S.; Hudson,

[12103] B. G.: Identification of the Goodpasture antigen as the al-pha-3(IV)

[12104] chain of collagen IV. J. Biol. Chem. 263: 13374-13380, 1988.

[12105] 582.

[12106] Shahan, T. A.; Ziaie, Z.; Pasco, S.; Fawzi, A.; Bellon, G.; Monboisse,
[12107] J.-C.; Kefalides, N. A.: Identification of CD47/integrin-associated
[12108] protein and alpha-v-beta-3 as two receptors for the al-pha-3(IV) chain
[12109] of type IV collagen on tumor cells. Cancer Res. 59: 4584-4590, 1999.
[12110] 583.

[12111] Thorner, P. S.; Zheng, K.; Kalluri, R.; Jacobs, R.; Hudson, B.

[12112] G.: Coordinate gene expression of the alpha-3, alpha-4, and alpha-5
[12113] chains of collagen type IV. J. Biol. Chem. 271: 13821-13828, 1996.
[12114] 584.

[12115] Bora, N. S.; Lublin, D. M.; Kumar, B. V.; Hockett, R. D.; Holers,
[12116] V. M.; Atkinson, J. P.: Structural gene for human mem-brane cofactor
[12117] protein (MCP) of complement maps to within 100 kb of the 3-prime end
[12118] of the C3b/C4b receptor gene. J. Exp. Med. 169:

597-602, 1989.

[12119] 585.

[12120] Cui, W.; Hourcade, D.; Post, T.; Greenlund, A. C.; Atkinson, J.

[12121] P.; Kumar, V.: Characterization of the promoter region of the membrane

[12122] cofactor protein (CD46) gene of the human complement system and comparison

[12123] to a membrane cofactor protein-like genetic element. J. Immun. 151:

[12124] 4137-4146, 1993.

[12125] 586.

[12126] Dorig, R. E.; Marcil, A.; Chopra, A.; Richardson, C. D.: The human

[12127] CD46 molecule is a receptor for measles virus (Edmonston strain). Cell 75:

[12128] 295-305, 1993.

[12129] 587.

[12130] Kallstrom, H.; Gill, D. B.; Albiger, B.; Liszewski, M. K.; Atkinson,

[12131] J. P.; Jonsson, A.-B.: Attachment of Neisseria gonorrhoeae

to the
[12132] cellular pilus receptor CD46: identification of domains important
[12133] for bacterial adherence. Cell. Microbiol. 3: 133-143, 2001.
[12134] 588.

[12135] Lublin, D. M.; Liszewski, M. K.; Post, T. W.; Arce, M. A.; LeBeau,
[12136] M. M.; Rebentisch, M. B.; Lemons, R. S.; Seya, T.; Atkinson, J. P.
[12137] : Molecular cloning and chromosomal localization of human complement
[12138] membrane cofactor protein (MCP): evidence for inclusion in the multigene
[12139] family of complement-regulatory proteins. J. Exp. Med. 168: 181-194,
[12140] 1988.

[12141] 589.

[12142] Marie, J. C.; Astier, A. L.; Rivailler, P.; Rabourdin-Combe, C.;
[12143] Wild, T. F.; Horvat, B.: Linking innate and acquired immunity: divergent

[12144] role of CD46 cytoplasmic domains in T cell-induced inflammation. Nature

[12145] Immun. 3: 659-666, 2002.

[12146] 590.

[12147] Post, T. W.; Liszewski, M. K.; Adams, E. M.; Tedja, I.; Miller,

[12148] E. A.; Atkinson, J. P.: Membrane cofactor protein of the complement

[12149] system: alternative splicing of serine/threonine/proline-rich exons

[12150] and cytoplasmic tails produces multiple isoforms that correlate with

[12151] protein phenotype. J. Exp. Med. 174: 93-102, 1991.

[12152] 591.

[12153] Purcell, D. F. J.; Johnstone, R. W.; McKenzie, I. F. C.: Identification

[12154] of four different CD46 (MCP) molecules with anti-peptide antibodies. Biochem.

[12155] Biophys. Res. Commun. 180: 1091-1097, 1991.

[12156] 592.

[12157] Santoro, F.; Kennedy, P. E.; Locatelli, G.; Malnati, M. S.;

Berger,

[12158] E. A.; Lusso, P.: CD46 is a cellular receptor for human herpesvirus

[12159] 6. Cell 99: 817-827, 1999.

[12160] 593.

[12161] Tatsuo, H.; Ono, N.; Tanaka, K.; Yanagi, Y.: SLAM (CDw150) is

[12162] a cellular receptor for measles virus. Nature 406: 893-897, 2000.

[12163] 594.

[12164] Chan, P.; Simon-Chazottes, D.; Mattei, M. G.; Guenet, J. L.; Salier,

[12165] J. P.: Comparative mapping of lipocalin genes in human and mouse:

[12166] the four genes for complement C8 gamma chain, prostaglandin-D-synthase,

[12167] oncogene-24P3, and progestagen-associated endometrial protein map

[12168] to HSA9 and MMU2. Genomics 23: 145-150, 1994.

[12169] 595.

[12170] Dewald, G.; Cichon, S.; Bryant, S. P.; Hemmer, S.; Nothen,

M. M.;

[12171] Spurr, N. K.: The human complement C8G gene, a member of the lipocalin

[12172] gene family: polymorphisms and mapping to chromosome 9q34.3. Ann.

[12173] Hum. Genet. 60: 281-291, 1996.

[12174] 596.

[12175] Haefliger, J.-A.; Jenne, D.; Stanley, K. K.; Tschopp, J.: Structural

[12176] homology of human complement component C8-gamma and plasma protein

[12177] HC: identity of the cysteine bond pattern. Biochem. Biophys. Res.

[12178] Commun. 149: 750-754, 1987.

[12179] 597.

[12180] Hunt, L. T.; Elzanowski, A.; Barker, W. C.: The homology of complement

[12181] factor C8 gamma chain and alpha-1-microglobulin. Biochem. Biophys.

[12182] Res. Commun. 149: 282-288, 1987.

[12183] 598.

[12184] Kaufman, K. M.; Snider, J. V.; Spurr, N. K.; Schwartz, C. E.; Sodetz,

[12185] J. M.: Chromosomal assignment of genes encoding the alpha, beta,

[12186] and gamma subunits of human complement protein C8: identification

[12187] of a close physical linkage between the alpha and the beta loci. Genomics 5:

[12188] 475-480, 1989.

[12189] 599.

[12190] Ng, S. C.; Rao, A. G.; Howard, O. M. Z.; Sodetz, J. M.: The eighth

[12191] component of human complement: evidence that it is an oligomeric serum

[12192] protein assembled from products of three different genes. Biochemistry 26:

[12193] 5229-5233, 1987.

[12194] 600.

[12195] Alvarez, V.; Coto, E.; Setien, F.; Spath, P. J.; Lopez-Larrea,

[12196] C.: Genetic detection of the silent allele (*Q0) in hereditary deficiencies

[12197] of the human complement C6, C7, and C9 components. Am. J. Med. Genet. 55:
[12198] 408-413, 1995.

[12199] 601.

[12200] Coto, E.; Martinez-Naves, E.; Dominguez, O.; DiScipio, R. G.; Urra,
[12201] J. M.; Lopez-Larrea, C.: DNA polymorphism and linkage relationship
[12202] of the human complement component C6, C7, and C9 genes. Immunogenetics 33:
[12203] 184-187, 1991.

[12204] 602.

[12205] Goldfarb, L. G.; Brown, P.; Goldgaber, D.; Asher, D. M.; Rubenstein,
[12206] R.; Brown, W. T.; Piccardo, P.; Kascsak, R. J.; Boellaard, J. W.;
[12207] Gajdusek, D. C.: Creutzfeldt-Jakob disease and kuru patients lack
[12208] a mutation consistently found in the Gerstmann-Straussler-Scheinker
[12209] syndrome. Exp. Neurol. 108: 247-250, 1990.

[12210] 603.

[12211] Manuelidis, L.; Sklaviadis, T.; Manuelidis, E. E.: Evidence suggesting

[12212] that PrP is not the infectious agent in Creutzfeldt-Jakob disease. EMBO

[12213] J. 6: 341-347, 1987.

[12214] 604.

[12215] King, I. A.; Arnemann, J.; Spurr, N. K.; Buxton, R. S.: Cloning

[12216] of the cDNA (DSC1) coding for human type 1 desmocollin and its assignment

[12217] to chromosome 18. Genomics 18: 185-194, 1993.

[12218] 605.

[12219] Troyanovsky, S. M.; Eshkind, L. G.; Troyanovsky, R. B.; Leube,

[12220] R. E.; Franke, W. W.: Contributions of cytoplasmic domains of desmosomal

[12221] cadherins to desmosome assembly and intermediate filament anchorage. Cell 72:

[12222] 561-574, 1993.

[12223] 606.

[12224] Arnemann, J.; Spurr, N. K.; Wheeler, G. N.; Parker, A. E.; Buxton,
[12225] R. S.: Chromosomal assignment of the human genes coding for the major
[12226] proteins of the desmosome junction, desmoglein DGI (DSG), desmocollins
[12227] DGII/III (DSC), desmoplakins DPI/II (DSP), and plakoglobin DPIII (JUP). Genomics 10:
[12228] 640-645, 1991.

[12229] 607.

[12230] Buxton, R. S.; Wheeler, G. N.; Pidsley, S. C.; Marsden, M. D.;
[12231] Adams, M. J.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.: Mouse
[12232] desmocollin (Dsc3) and desmoglein (Dsg1) genes are closely linked
[12233] in the proximal region of chromosome 18. Genomics 21: 510-516, 1994.
[12234] 608.

[12235] Rezaie, T.; Child, A.; Hitchings, R.; Brice, G.; Miller, L.; Coca-Prados,
[12236] M.; Heon, E.; Krupin, T.; Ritch, R.; Kreutzer, D.; Crick, R.

P.; Sarfarazi,

[12237] M.: Adult-onset primary open-angle glaucoma caused by mutations in

[12238] optineurin. Science 295: 1077-1079, 2002.

[12239] 609.

[12240] Dale, D. C.; Person, R. E.; Bolyard, A. A.; Aprikyan, A. G.; Bos,

[12241] C.; Bonilla, M. A.; Boxer, L. A.; Kannourakis, G.; Zeidler, C.; Welte,

[12242] K.; Benson, K. F.; Horwitz, M.: Mutations in the gene en-coding neutrophil

[12243] elastase in congenital and cyclic neutropenia. Blood 96: 2317-2322,

[12244] 2000.

[12245] 610.

[12246] Evans, E.; Cooley, J.; Remold-O'Donnell, E.: Characterization

[12247] and chromosomal localization of ELANH2, the gene encoding human monocyte/neutrophil

[12248] elastase inhibitor. Genomics 28: 235-240, 1995.

[12249] 611.

[12250] Schneider, S. S.; Schick, C.; Fish, K. E.; Miller, E.; Pena, J.

[12251] C.; Treter, S. D.; Hui, S. M.; Silverman, G. A.: A serine proteinase

[12252] inhibitor locus at 18q21.3 contains a tandem duplication of the human

[12253] squamous cell carcinoma antigen gene. Proc. Nat. Acad. Sci. 92:

[12254] 3147-3151, 1995.

[12255] 612.

[12256] Kramer, A.; Yang, F.-C.; Snodgrass, P.; Li, X.; Scammell, T. E.;

[12257] Davis, F. C.; Weitz, C. J.: Regulation of daily locomotor ac-

tivity
[12258] and sleep by hypothalamic EGF receptor signaling. Science 294: 2511-2515,
[12259] 2001.

[12260] 613.

[12261] Lanzetti, L.; Rybin, V.; Malabarba, M. G.; Christoforidis, S.;

[12262] Scita, G.; Zerial, M.; Di Fiore, P. P.: The Eps8 protein coordinates
[12263] EGF receptor signalling through Rac and trafficking through Rab5. Nature 408:
[12264] 374-377, 2000.

[12265] 614.

[12266] Mannick, J. B.; Hausladen, A.; Liu, L.; Hess, D. T.; Zeng, M.;

[12267] Miao, Q. X.; Kane, L. S.; Gow, A. J.; Stamler, J. S.: Fas-induced
[12268] caspase denitrosylation. Science 284: 651-654, 1999.

[12269] 615.

[12270] Volpert, O. V.; Zaichuk, T.; Zhou, W.; Reiher, F.; Ferguson, T.
[12271] A.; Stuart, P. M.; Amin, M.; Bouck, N. P.: Inducer-stimu-

lated Fas

[12272] targets activated endothelium for destruction by anti-angiogenic thrombospondin-1

[12273] and pigment epithelium-derived factor. Nature Med. 8: 349-357, 2002.

[12274] 616.

[12275] Hill, L. L.; Ouhtit, A.; Loughlin, S. M.; Kripke, M. L.; Ananthaswamy,

[12276] H. N.; Owen-Schaub, L. B.: Fas ligand: a sensor for DNA damage critical

[12277] in skin cancer etiology. Science 285: 898-900, 1999.

[12278] 617.

[12279] Xu, S.; Wang, Y.; Roe, B.; Pearson, W. R.: Characterization of

[12280] the human class mu glutathione S-transferase gene cluster and the

[12281] GSTM1 deletion. J. Biol. Chem. 273: 3517-3527, 1998.

[12282] 618.

[12283] Takahashi, Y.; Campbell, E. A.; Hirata, Y.; Takayama, T.; Listowsky,

[12284] I.: A basis for differentiating among the multiple human

mu-glutathione
[12285] S-transferases and molecular cloning of brain GSTM5. J. Biol. Chem. 268:
[12286] 8893-8898, 1993.

[12287] 619.

[12288] Campbell, E.; Takahashi, Y.; Abramovitz, M.; Peretz, M.; Listowsky,
[12289] I.: A distinct human testis and brain mu-class glutathione S-transferase:
[12290] molecular cloning and characterization of a form present even in individuals
[12291] lacking hepatic type mu isoenzymes. J. Biol. Chem. 265: 9188-9193,
[12292] 1990.

[12293] 620.

[12294] Elliott, K. J.; Ellis, S. B.; Berckhan, K. J.; Urrutia, A.; Chavez-Noriega,
[12295] L. E.; Johnson, E. C.; Velicelebi, G.; Harpold, M. M.: Comparative
[12296] structure of human neuronal alpha(2)-alpha(7) and beta(2)-beta(4)
[12297] nicotinic acetylcholine receptor subunits and functional

expression

[12298] of the alpha(2), alpha(3), alpha(4), alpha(7), beta(2), and beta(4)

[12299] subunits. J. Molec. Neurosci. 7: 217-228, 1996.

[12300] 621.

[12301] Solomon, E.; Hiorns, L. R.; Spurr, N.; Kurkinen, M.; Barlow, D.;

[12302] Hogan, B. L. M.; Dalgleish, R.: Chromosomal assignments of the genes

[12303] coding for human types II, III and IV collagen: a dispersed gene family. Proc.

[12304] Nat. Acad. Sci. 82: 3330-3334, 1985.

[12305] 622.

[12306] Haddad, P.; Clement, M.-V.; Bernard, O.; Larsen, C.-J.; De-gos,

[12307] L.; Sasportes, M.; Mathieu-Mahul, D.: Structural organiza-tion of

[12308] the CTLA-1 gene encoding human granzyme B. Gene 87: 265-271, 1990.

[12309] 623.

[12310] Johnson, R.; Jackson, I. J.: Light is a dominant mouse mu-

tation
[12311] resulting in premature cell death. Nature Genet. 1: 226-229, 1992.
[12312] 624.
[12313] Yang, J.; Patil, R. V.; Yu, H.; Gordon, M.; Wax, M. B.: T cell
[12314] subsets and sIL-2R/IL-2 levels in patients with glaucoma. Am. J.
[12315] Ophthal. 131: 421-426, 2001.
[12316] 625.
[12317] Bongcam-Rudloff, E.; Nister, M.; Betsholtz, C.; Wang, J.-L.; Stenman,
[12318] G.; Huebner, K.; Croce, C. M.; Westermark, B.: Human glial fibrillary
[12319] acidic protein: complementary DNA cloning, chromosome localization,
[12320] and messenger RNA expression in human glioma cell lines of various
[12321] phenotypes. Cancer Res. 51: 1553-1560, 1991.
[12322] 626.
[12323] Brenner, M.; Johnson, A. B.; Boespflug-Tanguy, O.; Rodriguez, D.;

[12324] Goldman, J. E.; Messing, A.: Mutations in GFAP, encoding glial fibrillary
[12325] acidic protein, are associated with Alexander disease. Nature Genet. 27:
[12326] 117-120, 2001.

[12327] 627.

[12328] Brownell, E.; Lee, A. S.; Pekar, S. K.; Pravtcheva, D.; Ruddle,
[12329] F. H.; Bayney, R. M.: Glial fibrillary acid protein, an astrocytic-specific
[12330] marker, maps to human chromosome 17. Genomics 10: 1087-1089, 1991.
[12331] 628.

[12332] Gomi, H.; Yokoyama, T.; Fujimoto, K.; Ikeda, T.; Katoh, A.; Itoh,
[12333] T.; Itohara, S.: Mice devoid of the glial fibrillary acidic protein
[12334] develop normally and are susceptible to scrapie prions. Neuron 14:
[12335] 29-41, 1995.

[12336] 629.

[12337] Liedtke, W.; Edelmann, W.; Bieri, P. L.; Chiu, F.-C.; Cowan, N.

[12338] J.; Kucherlapati, R.; Raine, C. S.: GFAP is necessary for the integrity

[12339] of CNS white matter architecture and long-term maintenance of myelination. Neuron 17:

[12340] 607-615, 1996.

[12341] 630.

[12342] Messing, A.; Head, M. W.; Galles, K.; Galbreath, E. J.; Goldman,

[12343] J. E.; Brenner, M.: Fatal encephalopathy with astrocyte inclusions

[12344] in GFAP transgenic mice. Am. J. Path. 152: 391-398, 1998.

[12345] 631.

[12346] Pekny, M.; Leveen, P.; Pekna, M.; Eliasson, C.; Berthold, C.-H.;

[12347] Westermark, B.; Betsholtz, C.: Mice lacking glial fibrillary acidic

[12348] protein display astrocytes devoid of intermediate filaments but develop

[12349] and reproduce normally. EMBO J. 14: 1590-1598, 1995.

[12350] 632.

[12351] Reeves, S. A.; Helman, L. J.; Allison, A.; Israel, M. A.: Molecular
[12352] cloning and primary structure of human glial fibrillary acidic protein. Proc.
[12353] Nat. Acad. Sci. 86: 5178-5182, 1989.

[12354] 633.

[12355] Schuelke, M.; Smeitink, J.; Mariman, E.; Loeffen, J.; Plecko,
[12356] B.; Trijbels, F.; Stockler-Ipsiroglu, S.; van den Heuvel, L.: Mutant
[12357] NDUFV1 subunit of mitochondrial complex I causes leukodystrophy and
[12358] myoclonic epilepsy. (Letter) Nature Genet. 21: 260-261, 1999.

[12359] 634.

[12360] Shibuki, K.; Gomi, H.; Chen, L.; Bao, S.; Kim, J. J.; Wakat-
suki,
[12361] H.; Fujisaki, T.; Fujimoto, K.; Katoh, A.; Ikeda, T.; Chen, C.; Thompson,
[12362] R. F.; Itohara, S.: Deficient cerebellar long-term depres-
sion, impaired

[12363] eyeblink conditioning, and normal motor coordination in GFAP mutant

[12364] mice. Neuron 16: 587-599, 1996.

[12365] 635.

[12366] Rodriguez, D.; Gauthier, F.; Bertini, E.; Bugiani, M.; Brenner,

[12367] M.; N'guyen, S.; Goizet, C.; Gelot, A.; Surtees, R.; Pedespan, J.-M.;

[12368] Hernandorena, X.; Troncoso, M.; Uziel, G.; Messing, A.; Ponsot, G.;

[12369] Pham-Dinh, D.; Dautigny, A.; Boespflug-Tanguy, O.: Infantile Alexander

[12370] disease: spectrum of GFAP mutations and genotype-phenotype correlation. Am.

[12371] J. Hum. Genet. 69: 1134-1140, 2001. Note: Erratum: Am. J. Hum. Genet.

[12372] 69: 1413 only, 2001.

[12373] 636.

[12374] Shiroma, N.; Kanazawa, N.; Izumi, M.; Sugai, K.; Fukumizu, M.;

[12375] Sasaki, M.; Hanaoka, S.; Kaga, M.; Tsujino, S.: Diagnosis of Alexander

[12376] disease in a Japanese patient by molecular genetic analysis. J. Hum.

[12377] Genet. 46: 579-582, 2001.

[12378] 637.

[12379] Groot Kormelink, P. J.; Luyten, W. H. M. L.: Cloning and sequence

[12380] of full-length cDNAs encoding the human neuronal nicotinic acetylcholine

[12381] receptor (nAChR) subunits beta-3 and beta-4 and expression of seven

[12382] nAChR subunits in the human neuroblastoma cell line SH-SY5Y and/or

[12383] IMR-32. FEBS Lett. 400: 309-314, 1997.

[12384] 638.

[12385] Rempel, N.; Heyers, S.; Engels, H.; Sleegers, E.; Steinlein, O.

[12386] K.: The structures of the human neuronal nicotinic acetylcholine

[12387] receptor beta-2- and alpha-3-subunit genes (CHRNB2 and CHRNA3). Hum.

[12388] Genet. 103: 645-653, 1998.

[12389] 639.

[12390] Forman, S. A.; Miller, K. W.; Yellen, G.: A discrete site for

[12391] general anesthetics on a postsynaptic receptor. Molec. Pharm. 48:

[12392] 574-581, 1995.

[12393] 640.

[12394] Forman, S. A.; Yellen, G.; Thiele, E. A.: Alternative mecha-nism

[12395] for pathogenesis of an inherited epilepsy by a nicotinic AChR mutation.

[12396] (Letter) Nature Genet. 13: 396-397, 1996.

[12397] 641.

[12398] Hirose, S.; Iwata, H.; Akiyoshi, H.; Kobayashi, K.; Ito, M.; Wada,

[12399] K.; Kaneko, S.; Mitsudome, A.: A novel mutation of CHRNA4 responsible

[12400] for autosomal dominant nocturnal frontal lobe epilepsy. Neurology 53:

[12401] 1749-1753, 1999.

[12402] 642.

[12403] Marubio, L. M.; del Mar Arroyo-Jimenez, M.; Cordero-Er-
ausquin,
[12404] M.; Lena, C.; Le Novere, N.; de Kerchove d'Exaerde, A.;
Huchet, M.;
[12405] Damaj, M. I.; Changeux, J.-P.: Reduced antinociception in
mice lacking
[12406] neuronal nicotinic receptor subunits. Nature 398:
805-810, 1999.
[12407] 643.

[12408] Monteggia, L. M.; Gopalakrishnan, M.; Touma, E.; Idler, K.
B.;
[12409] Nash, N.; Arneric, S. P.; Sullivan, J. P.; Giordano, T.:
Cloning
[12410] and transient expression of genes encoding the human
alpha-4 and beta-2
[12411] neuronal nicotinic acetylcholine receptor (nAChR) sub-
units. Gene 155:
[12412] 189-193, 1995.

[12413] 644.

[12414] Pilz, A. J.; Willer, E.; Povey, S.; Abbott, C. M.: The genes

[12415] coding for phosphoenolpyruvate carboxykinase-1 (PCK1)
and neuronal

[12416] nicotinic acetylcholine receptor alpha-4 subunit (CHRNA4) map to human

[12417] chromosome 20, extending the known region of homology with mouse chromosome

[12418] 2. Ann. Hum. Genet. 56: 289-293, 1992.

[12419] 645.

[12420] Steinlein, O.; Smigrodzki, R.; Lindstrom, J.; Anand, R.; Kohler,

[12421] M.; Tocharoentanaphol, C.; Vogel, F.: Refinement of the localization

[12422] of the gene for neuronal nicotinic acetylcholine receptor alpha-4

[12423] subunit (CHRNA4) to human chromosome 20q13.2-q13.3. Genomics 22:

[12424] 493-495, 1994.

[12425] 646.

[12426] Steinlein, O.; Weiland, S.; Stoodt, J.; Propping, P.: Exon-intron

[12427] structure of the human neuronal nicotinic acetylcholine receptor alpha-4

[12428] subunit (CHRNA4). Genomics 32: 289-294, 1996.

[12429] 647.

[12430] Steinlein, O. K.; Magnusson, A.; Stoodt, J.; Bertrand, S.; Weiland,

[12431] S.; Berkovic, S. F.; Nakken, K. O.; Propping, P.; Bertrand, D.: An

[12432] insertion mutation of the CHRNA4 gene in a family with autosomal dominant

[12433] nocturnal frontal lobe epilepsy. Hum. Molec. Genet. 6: 943-947,

[12434] 1997.

[12435] 648.

[12436] Steinlein, O. K.; Mulley, J. C.; Propping, P.; Wallace, R. H.;

[12437] Phillips, H. A.; Sutherland, G. R.; Scheffer, I. E.; Berkovic, S.

[12438] F.: A missense mutation in the neuronal nicotinic acetyl-choline receptor

[12439] alpha-4 subunit is associated with autosomal dominant nocturnal frontal

[12440] lobe epilepsy. Nature Genet. 11: 201-203, 1995.

[12441] 649.

[12442] Tsonis, P.; Goetinck, P. F.: The Drosophila homoeotic gene spalt

[12443] is structurally related to collagen alpha-1(IV) chain. (Letter) Collagen

[12444] Rel. Res. 8: 451-452, 1988.

[12445] 650.

[12446] Wieslander, J.; Barr, J. F.; Butkowski, R. J.; Edwards, S. J.;

[12447] Bygren, P.; Heinegard, D.; Hudson, B. G.: Goodpasture antigen of

[12448] the glomerular basement membrane: localization to non-collagenous regions

[12449] of type IV collagen. Proc. Nat. Acad. Sci. 81: 3838-3842, 1984.

[12450] 651.

[12451] Wieslander, J.; Langeveld, J.; Butkowski, R.; Jodlowski, M.; Noelken,

[12452] M.; Hudson, B. G.: Physical and immunochemical studies of the globular

[12453] domain of type IV collagen: cryptic properties of the Goodpasture

[12454] antigen. J. Biol. Chem. 260: 8564-8570, 1985.

[12455] 652.

[12456] Boye, E.; Mollet, G.; Forestier, L.; Cohen-Solal, L.; Heidet,

L.;

[12457] Cochat, P.; Grunfeld, J.-P.; Palcoux, J.-B.; Gubler, M.-C.; Antignac,

[12458] C.: Determination of the genomic structure of the COL4A4 gene and

[12459] of novel mutations causing autosomal recessive Alport syndrome. Am.

[12460] J. Hum. Genet. 63: 1329-1340, 1998.

[12461] 653.

[12462] Gunwar, S.; Saus, J.; Noelken, M. E.; Hudson, B. G.: Glomerular

[12463] basement membrane: identification of a fourth chain, al-pha-4, of type

[12464] IV collagen. J. Biol. Chem. 265: 5466-5469, 1990.

[12465] 654.

[12466] Kamagata, Y.; Mattei, M.-G.; Ninomiya, Y.: Isolation and sequencing

[12467] of cDNAs and genomic DNAs encoding the alpha 4 chain of basement membrane

[12468] collagen type IV and assignment of the gene to the distal long arm

[12469] of human chromosome 2. J. Biol. Chem. 267:

23753-23758, 1992.

[12470] 655.

[12471] Leinonen, A.; Mariyama, M.; Mochizuki, T.; Tryggvason, K.; Reeders,

[12472] S. T.: Complete primary structure of the human type IV collagen alpha-4(IV)

[12473] chain: comparison with structure and expression of the other alpha(IV)

[12474] chains. J. Biol. Chem. 269: 26172-26177, 1994.

[12475] 656.

[12476] Lemmink, H. H.; Nillesen, W. N.; Mochizuki, T.; Schroder, C. H.;

[12477] Brunner, H. G.; van Oost, B. A.; Monnens, L. A. H.; Smeets, H. J.

[12478] M.: Benign familial hematuria due to mutation of the type IV collagen

[12479] alpha-4 gene. J. Clin. Invest. 98: 1114-1118, 1996.

[12480] 657.

[12481] Mariyama, M.; Zheng, K.; Yang-Feng, T. L.; Reeders, S. T.: Colocalization

[12482] of the genes for the alpha-3(IV) and alpha-4(IV) chains of

type IV
[12483] collagen to chromosome 2 bands q35-q37. Genomics 13: 809-813, 1992.
[12484] 658.

[12485] Ahmad, N. N.; Ala-Kokko, L.; Knowlton, R. G.; Jimenez, S. A.; Weaver,
[12486] E. J.; Maguire, J. I.; Tasman, W.; Prockop, D. J.: Stop codon in
[12487] the procollagen II gene (COL2A1) in a family with the Stickler syndrome
[12488] (arthro-ophthalmopathy). Proc. Nat. Acad. Sci. 88: 6624-6627, 1991.
[12489] 659.

[12490] Ahmad, N. N.; Ala-Kokko, L.; Knowlton, R. G.; Weaver, E. J.; Maguire,
[12491] J. I.; Tasman, W.; Prockop, D. J.: A stop codon in the gene for type
[12492] II procollagen (COL2A1) causes one variant of arthro-ophthalmopathy
[12493] (the Stickler syndrome). (Abstract) Am. J. Hum. Genet. 47 (suppl.):
[12494] A206 only, 1990.

[12495] 660.

[12496] Ahmad, N. N.; Dimascio, J.; Knowlton, R. G.; Tasman, W. S.: Stickler

[12497] syndrome: a mutation in the nonhelical 3-prime end of type II procollagen

[12498] gene. Arch. Ophthal. 113: 1454-1457, 1995.

[12499] 661.

[12500] Ahmad, N. N.; McDonald-McGinn, D. M.; Zackai, E. H.; Knowlton,

[12501] R. G.; LaRossa, D.; DiMascio, J.; Prockop, D. J.: A second mutation

[12502] in the type II procollagen gene (COL2A1) causing Stickler syndrome

[12503] (arthro-ophthalmopathy) is also a premature termination codon. Am.

[12504] J. Hum. Genet. 52: 39-45, 1993.

[12505] 662.

[12506] Ala-Kokko, L.; Baldwin, C. T.; Moskowitz, R. W.; Prockop, D. J.

[12507] : Single base mutation in the type II procollagen gene (COL2A1) as

[12508] a cause of primary osteoarthritis associated with a mild chondrodysplasia. Proc.

[12509] Nat. Acad. Sci. 87: 6565-6568, 1990.

[12510] 663.

[12511] Baldwin, C. T.; Reginato, A. M.; Smith, C.; Jimenez, S. A.; Prockop,

[12512] D. J.: Structure of cDNA clones coding for human type II procollagen:

[12513] the alpha-1(II) chain is more similar to the alpha-1(I) chain than

[12514] two other alpha chains of fibrillar collagens. Biochem. J. 262:

[12515] 521-528, 1989.

[12516] 664.

[12517] Beighton, P.; Christy, G.; Learmonth, I. D.: Namaqualand hip dysplasia:

[12518] an autosomal dominant entity. Am. J. Med. Genet. 19: 161-169, 1984.

[12519] 665.

[12520] Bleasel, J. F.; Holderbaum, D.; Brancolini, V.; Moskowitz, R. W.;

[12521] Considine, E. L.; Prockop, D. J.; Devoto, M.; Williams, C. J.: Five
[12522] families with arginine-519-to-cysteine mutation in COL2A1: evidence
[12523] for three distinct founders. Hum. Mutat. 12: 172-176, 1998.
[12524] 666.

[12525] Bogaert, R.; Tiller, G. E.; Weis, M. A.; Gruber, H. E.; Rimoin,
[12526] D. L.; Cohn, D. H.; Eyre, D. R.: An amino acid substitution (gly853-to-glu)
[12527] in the collagen alpha-1(II) chain produces hypochondro-genesis. J.
[12528] Biol. Chem. 267: 22522-22526, 1992.

[12529] 667.

[12530] Brown, D. M.; Nichols, B. E.; Weingeist, T. A.; Sheffield, V.
[12531] C.; Kimura, A. E.; Stone, E. M.: Procollagen II gene muta-tion in
[12532] Stickler syndrome. Arch. Ophthal. 110: 1589-1593, 1992.

[12533] 668.

[12534] Chan, D.; Cole, W. G.; Chow, C. W.; Mundlos, S.; Bateman, J. F.

[12535] : A COL2A1 mutation in achondrogenesis type II results in the replacement

[12536] of type II collagen by type I and III collagens in cartilage. J.

[12537] Biol. Chem. 270: 1747-1753, 1995.

[12538] 669.

[12539] Chan, D.; Taylor, T. K. F.; Cole, W. G.: Characterization of

[12540] an arginine 789 to cysteine substitution in alpha-1(II) collagen chains

[12541] of a patient with spondyloepiphyseal dysplasia. J. Biol. Chem. 268:

[12542] 15238-15245, 1993.

[12543] 670.

[12544] Cheah, K. S. E.; Stoker, N. G.; Griffin, J. R.; Grosveld, F. G.;

[12545] Solomon, E.: Identification and characterization of the human type

[12546] II collagen gene (COL2A1). Proc. Nat. Acad. Sci. 82: 2555-2559,

[12547] 1985.

[12548] 671.

[12549] Eccles, D. M.; van der Luijt, R.; Breukel, C.; Bullman, H.;

Bunyan,

[12550] D.; Fisher, A.; Barber, J.; du Boulay, C.; Primrose, J.; Burn, J.;

[12551] Fodde, R.: Hereditary desmoid disease due to a frameshift mutation

[12552] at codon 1924 of the APC gene. Am. J. Hum. Genet. 59: 1193-1201,

[12553] 1996.

[12554] 672.

[12555] Ishida-Yamamoto, A.; McGrath, J. A.; Lam, H.; Iizuka, H.; Friedman,

[12556] R. A.; Christiano, A. M.: The molecular pathology of pro-

[12557] gressive symmetric erythrokeratoderma: a frameshift mutation in the loricrin

[12558] gene and perturbations in the cornified cell envelope. Am. J. Hum.

[12559] Genet. 61: 581-589, 1997.

[12560] 673.

[12561] van der Schroeff, J. G.; Nijenhuis, L. E.; Meera Khan, P.; Bernini,

[12562] L. F.; Schreuder, G. M. T.; van Loghem, E.; Volkers, W. S.;

Went,

[12563] L. N.: Genetic linkage between erythrokeratodermia variabilis and

[12564] Rh locus. Hum. Genet. 68: 165-168, 1984.

[12565] 674.

[12566] Allende, M. L.; Amsterdam, A.; Becker, T.; Kawakami, K.; Gaiano,

[12567] N.; Hopkins, N.: Insertional mutagenesis in zebrafish identifies

[12568] two novel genes, pescadillo and dead eye, essential for embryonic

[12569] development. Genes Dev. 10: 3141-3155, 1996.

[12570] 675.

[12571] Haque, J.; Boger, S.; Li, J.; Duncan, S. A.: The murine Pes1 gene

[12572] encodes a nuclear protein containing a BRCT domain. Genomics 70:

[12573] 201-210, 2000.

[12574] 676.

[12575] Kinoshita, Y.; Jarell, A. D.; Flaman, J. M.; Foltz, G.; Schuster,

[12576] J.; Sopher, B. L.; Irvin, D. K.; Kanning, K.; Kornblum, H. I.; Nelson,

[12577] P. S.; Hieter, P.; Morrison, R. S.: Pescadillo, a novel cell cycle

[12578] regulatory protein abnormally expressed in malignant cells. J. Biol.

[12579] Chem. 276: 6656-6665, 2001.

[12580] 677.

[12581] Prekeris, R.; Klumperman, J.; Scheller, R. H.: A Rab11/Rip11 protein

[12582] complex regulates apical membrane trafficking via recycling endosomes. Molec.

[12583] Cell 6: 1437-1448, 2000.

[12584] 678.

[12585] Van Hauwe, P.; Everett, L. A.; Coucke, P.; Scott, D. A.; Kraft,

[12586] M. L.; Ris-Stalpers, C.; Bolder, C.; Otten, B.; de Vijlder, J. J.

[12587] M.; Dietrich, N. L.; Ramesh, A.; Srisailapathy, S. C. R.; Parving,

[12588] A.; Cremers, C. W. R. J.; Willems, P. J.; Smith, R. J. H.; Green,

[12589] E. D.; Van Camp, G.: Two frequent missense mutations in Pendred syndrome. Hum.

[12590] Molec. Genet. 7: 1099-1104, 1998.

[12591] 679.

[12592] Cenciarelli, C.; Chiaur, D. S.; Guardavaccaro, D.; Parks, W.; Vidal,

[12593] M.; Pagano, M.: Identification of a family of human F-box proteins. Curr.

[12594] Biol. 9: 1177-1179, 1999.

[12595] 680.

[12596] Winston, J. T.; Koepp, D. M.; Zhu, C.; Elledge, S. J.; Harper,

[12597] J. W.: A family of mammalian F-box proteins. Curr. Biol. 9: 1180-1182,

[12598] 1999.

[12599] 681.

[12600] Gerlach, V. L.; Aravind, L.; Gotway, G.; Schultz, R. A.; Koonin,

[12601] E. V.; Friedberg, E. C.: Human and mouse homologs of Escherichia

[12602] coli DinB (DNA polymerase IV), members of the UmuC/DinB superfamily. Proc.

[12603] Nat. Acad. Sci. 96: 11922-11927, 1999.

[12604] 682.

[12605] Johnson, R. E.; Prakash, S.; Prakash, L.: The human DINB1 gene

[12606] encodes the DNA polymerase Pol-theta. Proc. Nat. Acad. Sci. 97:

[12607] 3838-3843, 2000.

[12608] 683.

[12609] Ogi, T.; Kato, T., Jr.; Kato, T.; Ohmori, H.: Mutation en-hancement

[12610] by DINB1, a mammalian homologue of the Escherichia coli mutagenesis

[12611] protein dinB. Genes Cells 4: 607-618, 1999.

[12612] 684.

[12613] Ohashi, E.; Ogi, T.; Kusumoto, R.; Iwai, S.; Masutani, C.; Hanaoka,

[12614] F.; Ohmori, H.: Error-prone bypass of certain DNA lesions by the

[12615] human DNA polymerase kappa. Genes Dev. 14: 1589-1594, 2000.

[12616] 685.

[12617] Koike, J.; Sagara, N.; Kirikoshi, H.; Takagi, A.; Miwa, T.; Hi-
rai,
[12618] M.; Katoh, M.: Molecular cloning and genomic structure of
the beta-TRCP2
[12619] gene on chromosome 5q35.1. Biochem. Biophys. Res.
Commun. 269: 103-109,
[12620] 2000.

[12621] 686.

[12622] Chiaur, D. S.; Murthy, S.; Cenciarelli, C.; Parks, W.; Loda,
M.;
[12623] Inghirami, G.; Demetrick, D.; Pagano, M.: Five human
genes encoding
[12624] F-box proteins: chromosome mapping and analysis in hu-
man tumors. Cytogenet.
[12625] Cell Genet. 88: 255-258, 2000.

[12626] 687.

[12627] Bauer, H.; Mayer, H.; Marchler-Bauer, A.; Salzer, U.; Pro-
haska,
[12628] R.: Characterization of p40/GPR69A as a peripheral mem-
brane protein
[12629] related to the lantibiotic synthetase component C.
Biochem. Biophys.

[12630] Res. Commun. 275: 69-74, 2000.

[12631] 688.

[12632] Mayer, H.; Bauer, H.; Prohaska, R.: Organization and chro-
mosomal

[12633] localization of the human and mouse genes coding for
LanC-like protein

[12634] 1 (LANCL1). Cytogenet. Cell Genet. 93: 100-104, 2001.

[12635] 689.

[12636] Mayer, H.; Salzer, U.; Breuss, J.; Ziegler, S.; Marchler-
Bauer,

[12637] A.; Prohaska, R.: Isolation, molecular characterization, and
tissue-specific

[12638] expression of a novel putative G protein-coupled recep-
tor. Biochim.

[12639] Biophys. Acta 1395: 301-308, 1998.

[12640] 690.

[12641] Baylin, S. B.; Herman, J. G.: DNA hypermethylation in tu-
morigenesis:

[12642] epigenetics joins genetics. Trends Genet. 16: 168-174,
2000.

[12643] 691.

[12644] Cameron, E. E.; Bachman, K. E.; Myohanen, S.; Herman, J. G.; Baylin,

[12645] S. B.: Synergy of demethylation and histone deacetylase inhibition

[12646] in the re-expression of genes silenced in cancer. Nature Genet. 21:

[12647] 103-107, 1999.

[12648] 692.

[12649] Finch, P. W.; He, X.; Kelley, M. J.; Uren, A.; Schaudies, R. P.;

[12650] Popescu, N. C.; Rudikoff, S.; Aaronson, S. A.; Varmus, H. E.; Rubin,

[12651] J. S.: Purification and molecular cloning of a secreted, friz-zled-related

[12652] antagonist of Wnt action. Proc. Nat. Acad. Sci. 94: 6770-6775, 1997.

[12653] 693.

[12654] Fukuhara, K.; Kariya, M.; Kita, M.; Shime, H.; Kanamori, T.; Kosaka,

[12655] C.; Orii, A.; Fujita, J.; Fujii, S.: Secreted frizzled related protein

[12656] 1 is overexpressed in uterine leiomyomas, associated with a high estrogenic

[12657] environment and unrelated to proliferative activity. J. Clin. Endocr.

[12658] Metab. 87: 1729-1736, 2002.

[12659] 694.

[12660] Melkonyan, H. S.; Chang, W. C.; Shapiro, J. P.; Mahadevappa, M.;

[12661] Fitzpatrick, P. A.; Kiefer, M. C.; Tomei, L. D.; Umansky, S. R.:

[12662] SARPs: a family of secreted apoptosis-related proteins. Proc. Nat.

[12663] Acad. Sci. 94: 13636-13641, 1997.

[12664] 695.

[12665] Suzuki, H.; Gabrielson, E.; Chen, W.; Anbazhagan, R.; van Engeland,

[12666] M.; Weijenberg, M. P.; Herman, J. G.; Baylin, S. B.: A genomic screen

[12667] for genes upregulated by demethylation and histone deacetylase inhibition

[12668] in human colorectal cancer. Nature Genet. 31: 141-149, 2002.

[12669] 696.

[12670] Chang, J. T.; Esumi, N.; Moore, K.; Li, Y.; Zhang, S.; Chew, C.;

[12671] Goodman, B.; Rattner, A.; Moody, S.; Stetten, G.; Campochiaro, P.

[12672] A.; Zack, D. J.: Cloning and characterization of a secreted frizzled-related

[12673] protein that is expressed by the retinal pigment epithelium. Hum.

[12674] Molec. Genet. 8: 575-583, 1999.

[12675] 697.

[12676] Rattner, A.; Hsieh, J.-C.; Smallwood, P. M.; Gilbert, D. J.; Copeland,

[12677] N. G.; Jenkins, N. A.; Nathans, J.: A family of secreted proteins

[12678] contains homology to the cysteine-rich ligand-binding domain of frizzled

[12679] receptors. Proc. Nat. Acad. Sci. 94: 2859-2863, 1997.

[12680] 698.

[12681] Gallagher, M. J.; Burgess, L. H.; Brunden, K. R.: Characterization

[12682] of multiple forms of the human glycine transporter type-2. Molec.

[12683] Brain Res. 70: 101-115, 1999.

[12684] 699.

[12685] Liu, Q. R.; Lopez-Corcuera, B.; Mandiyan, S.; Nelson, H.; Nelson,

[12686] N.: Cloning and expression of a spinal cord- and brain-specific glycine

[12687] transporter with novel structural features. J. Biol. Chem. 268:

[12688] 22802-22808, 1993.

[12689] 700.

[12690] Olaisen, B.; Gedde-Dahl, T., Jr.: GPT-epidermolysis bul-losa simplex

[12691] (EBS Ogna) linkage in man. Hum. Hered. 23: 189-196, 1973.

[12692] 701.

[12693] Olaisen, B.; Gedde-Dahl, T., Jr.: GPT-EBS(1) linkage group: general

[12694] linkage relations. Hum. Hered. 24: 178-185, 1974.

[12695] 702.

[12696] Bach, I.; Galcheva-Gargova, Z.; Mattei, M.-G.; Simon-Chazottes,

[12697] D.; Guenet, J.-L.; Cereghini, S.; Yaniv, M.: Cloning of human hepatic

[12698] nuclear factor 1 (HNF1) and chromosomal localization of its gene in

[12699] man and mouse. Genomics 8: 155-164, 1990.

[12700] 703.

[12701] Byrne, M. M.; Sturis, J.; Menzel, S.; Yamagata, K.; Fajans, S.

[12702] S.; Dronsfield, M. J.; Bain, S. C.; Hattersley, A. T.; Velho, G.;

[12703] Froguel, P.; Bell, G. I.; Polonsky, K. S.: Altered insulin secretory

[12704] responses to glucose in diabetic and nondiabetic subjects with mutations

[12705] in the diabetes susceptibility gene MODY3 on chromosome 12. Diabetes 45:

[12706] 1503-1510, 1996.

[12707] 704.

[12708] Chiu, K. C.; Chuang, L.-M.; Ryu, J. M.; Tsai, G. P.; Saad, M. F.

[12709] : The I27L amino acid polymorphism of hepatic nuclear factor-1-alpha

[12710] is associated with insulin resistance. J. Clin. Endocr.

Metab. 85:
[12711] 2178-2183, 2000.

[12712] 705.

[12713] Collet, C.; et al.; et al.: Prevalence of the missense muta-
tion
[12714] Gly574Ser in the hepatocyte nuclear factor-1-alpha in
Africans with
[12715] diabetes. Diabetes Metab. 28: 39-44, 2002.

[12716] 706.

[12717] Courtois, G.; Morgan, J. G.; Campbell, L. A.; Fourel, G.;
Crabtree,
[12718] G. R.: Interaction of a liver-specific nuclear factor with the
fibrinogen
[12719] and alpha-1-antitrypsin promoters. Science 238:
688-692, 1987.
[12720] 707.

[12721] De Simone, V.; De Magistris, L.; Lazzaro, D.; Gerstner, J.;
Monaci,
[12722] P.; Nicosia, A.; Cortese, R.: LFB3, a heterodimer-forming
homeoprotein
[12723] of the LFB1 family, is expressed in specialized epithelia.

EMBO J. 10:
[12724] 1435-1443, 1991.

[12725] 708.

[12726] Ellard, S.: Hepatocyte nuclear factor 1 alpha
(HNF-1-alpha) mutations
[12727] in maturity-onset diabetes of the young. Hum. Mutat. 16:
377-385,
[12728] 2000.

[12729] 709.

[12730] Frayling, T. M.; Bulman, M. P.; Appleton, M.; Hattersley, A.
T.;
[12731] Ellard, S.: A rapid screening method for hepatocyte nu-
clear factor
[12732] 1 alpha frameshift mutations; prevalence in maturity-on-
set diabetes
[12733] of the young and late-onset non-insulin dependent dia-
betes. Hum.
[12734] Genet. 101: 351-354, 1997.

[12735] 710.

[12736] Frayling, T. M.; Bulman, M. P.; Ellard, S.; Appleton, M.;
Dronsfield,

[12737] M. J.; Mackie, A. D. R.; Baird, J. D.; Kaisaki, P. J.; Yamagata, K.;

[12738] Bell, G. I.; Bain, S. C.; Hattersley, A. T.: Mutations in the hepatocyte

[12739] nuclear factor-1-alpha gene are a common cause of maturity-onset diabetes

[12740] of the young in the U.K. Diabetes 46: 720-725, 1997.

[12741] 711.

[12742] Godart, F.; Bellanne-Chantelot, C.; Clauin, S.; Gragnoli, C.;

[12743] Abderrahmani, A.; Blanche, H.; Boutin, P.; Chevre, J. C.; Froguel,

[12744] P.; Bailleul, B.: Identification of seven novel nucleotide variants

[12745] in the hepatocyte nuclear factor-1-alpha (TCF1) promoter region in

[12746] MODY patients. Hum. Mutat. 15: 173-180, 2000.

[12747] 712.

[12748] Gonzalez, F. J.; Liu, S.-Y.; Kozak, C. A.; Nebert, D. W.: Decreased

[12749] Hnf-1 gene expression in mice homozygous for a 1.2-centimorgan deletion

[12750] on chromosome 7. DNA Cell Biol. 9: 771-776, 1990.

[12751] 713.

[12752] Gragnoli, C.; Lindner, T.; Cockburn, B. N.; Kaisaki, P. J.; Gragnoli,

[12753] F.; Marozzi, G.; Bell, G. I.: Maturity-onset diabetes of the young

[12754] due to a mutation in the hepatocyte nuclear factor-4-alpha binding

[12755] site in the promoter of the hepatocyte nuclear factor-1-alpha gene. Diabetes 46:

[12756] 1648-1651, 1997.

[12757] 714.

[12758] Hansen, T.; Eiberg, H.; Rouard, M.; Vaxillaire, M.; Moller, A.

[12759] M.; Rasmussen, S. K.; Fridberg, M.; Urhammer, S. A.; Holst, J. J.;

[12760] Almind, K.; Echwald, S. M.; Hansen, L.; Bell, G. I.; Pedersen, O.

[12761] : Novel MODY3 mutations in the hepatocyte nuclear factor-1-alpha gene:

[12762] evidence for a hyperexcitability of pancreatic beta-cells to intravenous

[12763] secretagogues in a glucose-tolerant carrier of a P447L mutation. Diabetes 46:

[12764] 726-730, 1997.

[12765] 715.

[12766] Hegele, R. A.; Cao, H.; Harris, S. B.; Hanley, A. J. G.; Zinman,

[12767] B.: The hepatic nuclear factor-1-alpha G319S variant is associated

[12768] with early-onset type 2 diabetes in Canadian Oji-Cree. J. Clin. Endocr.

[12769] Metab. 84: 1077-1082, 1999.

[12770] 716.

[12771] Hegele, R. A.; Cao, H.; Harris, S. B.; Zinman, B.; Hanley, A.

[12772] J. G.; Anderson, C. M.: Peroxisome proliferator-activated receptor-gamma-2

[12773] P12A and type 2 diabetes in Canadian Oji-Cree. J. Clin. Endocr. Metab. 85:

[12774] 2014-2019, 2000.

[12775] 717.

[12776] Hiraiwa, H.; Pan, C.-J.; Lin, B.; Akiyama, T. E.; Gonzalez, F.

[12777] J.; Chou, J. Y.: A molecular link between the common phenotypes of
[12778] type 1 glycogen storage disease and HNF1-alpha-null mice. J. Biol.
[12779] Chem. 276: 7963-7967, 2001.

[12780] 718.

[12781] Hua, Q.-X.; Zhao, M.; Narayana, N.; Nakagawa, S. H.; Jia, W.;
[12782] Weiss, M. A.: Diabetes-associated mutations in a beta-cell transcription
[12783] factor destabilize an antiparallel 'mini-zipper' in a dimerization
[12784] interface. Proc. Nat. Acad. Sci. 97: 1999-2004, 2000.

[12785] 719.

[12786] Kaisaki, P. J.; Menzel, S.; Lindner, T.; Oda, N.; Rjasanowski,
[12787] I.; Sahm, J.; Meincke, G.; Schulze, J.; Schmechel, H.; Petzold, C.;
[12788] Ledermann, H. M.; Sachse, G.; Boriraj, V. V.; Menzel, R.; Kerner,
[12789] W.; Turner, R. C.; Yamagata, K.; Bell, G. I.: Mutations in the hepatocyte

[12790] nuclear factor-1-alpha gene in MODY and early-onset NIDDM: evidence

[12791] for a mutational hotspot in exon 4. Diabetes 46: 528-535, 1997.

[12792] 720.

[12793] Kuo, C. J.; Conley, P. B.; Hsieh, C.-L.; Francke, U.; Crab-tree,

[12794] G. R.: Molecular cloning, functional expression, and chro-mosomal

[12795] localization of mouse hepatocyte nuclear factor 1. Proc. Nat. Acad.

[12796] Sci. 87: 9838-9842, 1990.

[12797] 721.

[12798] Mendel, D. B.; Hansen, L. P.; Graves, M. K.; Conley, P. B.; Crabtree,

[12799] G. R.: HNF-1-alpha and HNF-1-beta (vHNF-1) share dimerization and

[12800] homeo domains, but not activation domains, and form heterodimers in

[12801] vitro. Genes Dev. 5: 1042-1056, 1991.

[12802] 722.

[12803] Miedzybrodzka, Z.; Hattersley, A. T.; Ellard, S.; Pearson, D.;

[12804] de Silva, D.; Harvey, R.; Haites, N.: Non-penetrance in a MODY 3

[12805] family with a mutation in the hepatic nuclear factor 1a gene: implications

[12806] for predictive testing. Europ. J. Hum. Genet. 7: 729-732, 1999.

[12807] 723.

[12808] Pontoglio, M.; Barra, J.; Hadchouel, M.; Doyen, A.; Kress, C.;

[12809] Bach, J. P.; Babinet, C.; Yaniv, M.: Hepatocyte nuclear factor 1

[12810] inactivation results in hepatic dysfunction, phenylketonuria, and

[12811] renal Fanconi syndrome. Cell 84: 575-585, 1996.

[12812] 724.

[12813] Braud, V. M.; Allan, D. S. J.; O'Callaghan, C. A.; Soderstrom,

[12814] K.; D'Andrea, A.; Ogg, G. S.; Lazetic, S.; Young, N. T.; Bell, J.

[12815] I.; Phillips, J. H.; Lanier, L. L.; McMichael, A. J.: HLA-E

binds

[12816] to natural killer cell receptors CD94/NKG2A, B and C. Nature 391:

[12817] 795-799, 1998.

[12818] 725.

[12819] Carroll, M. C.; Katzman, P.; Alicot, E. M.; Koller, B. H.; Geraghty,

[12820] D. E.; Orr, H. T.; Strominger, J. L.; Spies, T.: Linkage map of the

[12821] human major histocompatibility complex including the tumor necrosis

[12822] factor genes. Proc. Nat. Acad. Sci. 84: 8535-8539, 1987.

[12823] 726.

[12824] Botstein, D.; White, R. L.; Skolnick, M.; Davis, R. W.: Construction

[12825] of a genetic linkage map in man using restriction fragment length

[12826] polymorphisms. Am. J. Hum. Genet. 32: 314-331, 1980.

[12827] 727.

[12828] Vernet, C.; Ribouchon, M.-T.; Chimini, G.; Pontarotti, P.: Structure

[12829] and evolution of a member of a new subfamily of GTP-binding proteins

[12830] mapping to the human MHC class I region. Mammalian Genome 5: 100-105,

[12831] 1994.

[12832] 728.

[12833] Mullis, P. E.; Patel, M. S.; Brickell, P. M.; Hindmarsh, P. C.;

[12834] Brook, C. G. D.: Growth characteristics and response to growth hormone

[12835] therapy in patients with hypochondroplasia: genetic link-age of the

[12836] insulin-like growth factor I gene at chromosome 12q23 to the disease

[12837] in a subgroup of these patients. Clin. Endocr. 34: 265-274, 1991.

[12838] 729.

[12839] Waldstreicher, J.; Seminara, S. B.; Jameson, J. L.; Geyer, A.;

[12840] Nachtigall, L. B.; Boepple, P. A.; Holmes, L. B.; Crowley, W.F., Jr.

[12841] : The genetic and clinical heterogeneity of gonadotropin-releasing

[12842] hormone deficiency in the human. J. Clin. Endocr. Metab. 81: 4388-4395,
[12843] 1996.

[12844] 730.

[12845] Baron, J.; Winer, K. K.; Yanovski, J. A.; Cunningham, A. W.; Laue,
[12846] L.; Zimmerman, D.; Cutler, G. B., Jr.: Mutations in the Ca(2+)-sensing
[12847] receptor gene cause autosomal dominant and sporadic hypoparathyroidism. Hum.
[12848] Molec. Genet. 5: 601-606, 1996.

[12849] 731.

[12850] Miki, T.; Kawamata, N.; Arai, A.; Ohashi, K.; Nakamura, Y.; Kato,
[12851] A.; Hirosawa, S.; Aoki, N.: Molecular cloning of the break-point for
[12852] 3q27 translocation in B-cell lymphomas and leukemias. Blood 83:
[12853] 217-222, 1994.

[12854] 732.

[12855] Ferrari, A. C.; Seuanez, H. N.; Hanash, S. M.; Atweh, G. F.:

A
[12856] gene that encodes for a leukemia-associated phospho-protein (p18) maps
[12857] to chromosome bands 1p35-36.1. Genes Chromosomes Cancer 2: 125-129,
[12858] 1990.

[12859] 733.

[12860] Hanash, S. M.; Strahler, J. R.; Kuick, R.; Chu, E. H. Y.; Nichols,
[12861] D.: Identification of a polypeptide associated with the ma-lignant
[12862] phenotype in acute leukemia. J. Biol. Chem. 263: 12813-12815, 1988.
[12863] 734.

[12864] Kumar, R.; Haugen, J. D.: Human and rat osteoblast-like cells
[12865] express stathmin, a growth-regulatory protein. Biochem. Biophys.
[12866] Res. Commun. 201: 861-865, 1994.

[12867] 735.

[12868] Maucuer, A.; Camonis, J. H.; Sobel, A.: Stathmin interac-

tion with

[12869] a putative kinase and coiled-coil-forming protein do-
mains. Proc.

[12870] Nat. Acad. Sci. 92: 3100-3104, 1995.

[12871] 736.

[12872] Mock, B. A.; Krall, M. M.; Padlan, C.; Dosik, J. K.; Schubart,

[12873] U. K.: The gene for Lap18, leukemia-associated phospho-
protein p18

[12874] (metablastin), maps to distal mouse chromosome 4.
Mammalian Genome 4:

[12875] 461-462, 1993.

[12876] 737.

[12877] Okazaki, T.; Yoshida, B. N.; Avraham, K. B.; Wang, H.;
Wuenschell,

[12878] C. W.; Jenkins, N. A.; Copeland, N. G.; Anderson, D. J.;
Mori, N.

[12879] : Molecular diversity of the SCG10/stathmin gene family in
the mouse. Genomics 18:

[12880] 360-373, 1993. Note: Erratum: Genomics 21: 298 only,
1994.

[12881] 738.

[12882] Sobel, A.: Stathmin: a relay phosphoprotein for multiple signal
[12883] transduction? Trends Biochem. Sci. 16: 301-305, 1991.

[12884] 739.

[12885] Sobel, A.; Boutterin, M.-C.; Beretta, L.; Chneiweiss, H.; Doye,
[12886] V.; Peyro-Saint-Paul, H.: Intracellular substrates for extra-cellular
[12887] signaling: characterization of a ubiquitous, neuron-enriched phosphoprotein
[12888] (stathmin). J. Biol. Chem. 264: 3765-3772, 1989.

[12889] 740.

[12890] Zhu, X. X.; Kozarsky, K.; Strahler, J. R.; Eckerskorn, C.; Lottspeich,
[12891] F.; Melhem, R.; Lowe, J.; Fox, D. A.; Hanash, S. M.; Atweh, G. F.
[12892] : Molecular cloning of a novel human leukemia-associated gene: evidence
[12893] of conservation in animal species. J. Biol. Chem. 264: 14556-14560,
[12894] 1989.

[12895] 741.

[12896] Gearing, D. P.; Druck, T.; Huebner, K.; Overhauser, J.; Gilbert,

[12897] D. J.; Copeland, N. G.; Jenkins, N. A.: The leukemia in-

hibitory factor

[12898] receptor (LIFR) gene is located within a cluster of cytokine receptor

[12899] loci on mouse chromosome 15 and human chromosome 5p12-p13. Genomics 18:

[12900] 148-150, 1993.

[12901] 742.

[12902] Delespesse, G.; Sarfati, M.; Peleman, R.: Influence of re-

combinant

[12903] IL-4, IFN-alpha, and IFN-gamma on the production of hu-

man IgE-binding

[12904] factor (soluble CD23). J. Immun. 142: 134-138, 1989.

[12905] 743.

[12906] Ludin, C.; Hofstetter, H.; Sarfati, M.; Levy, C. A.; Suter, U.;

[12907] Alaimo, D.; Kilchherr, E.; Frost, H.; Delespesse, G.: Cloning and

[12908] expression of the cDNA coding for a human lymphocyte

IgE receptor.

[12909] EMBO J. 6: 109-114, 1987.

[12910] 744.

[12911] Wendel-Hansen, V.; Riviere, M.; Uno, M.; Jansson, I.; Szpirer,

[12912] J.; Islam, M. Q.; Levan, G.; Klein, G.; Yodoi, J.; Rosen, A.; Szpirer,

[12913] C.: The gene encoding CD23 leukocyte antigen (FCE2) is located on

[12914] human chromosome 19. Somat. Cell Molec. Genet. 16: 283-286, 1990.

[12915] 745.

[12916] Li, S.-H.; Lam, S.; Cheng, A. L.; Li, X.-J.: Intranuclear huntingtin

[12917] increases the expression of caspase-1 and induces apoptosis. Hum.

[12918] Molec. Genet. 9: 2859-2867, 2000.

[12919] 746.

[12920] Mangiarini, L.; Sathasivam, K.; Seller, M.; Cozens, B.; Harper,

[12921] A.; Hetherington, C.; Lawton, M.; Trottier, Y.; Lehrach, H.;

Davies,
[12922] S. W.; Bates, G. P.: Exon 1 of the HD gene with an ex-
panded CAG repeat
[12923] is sufficient to cause a progressive neurological pheno-
type in transgenic
[12924] mice. Cell 87: 493-506, 1996.

[12925] 747.

[12926] Ona, V. O.; Li, M.; Vonsattel, J. P. G.; Andrews, L. J.; Khan,

[12927] S. Q.; Chung, W. M.; Frey, A. S.; Menon, A. S.; Li, X.-J.;
Stieg,
[12928] P. E.; Yuan, J.; Penney, J. B.; Young, A. B.; Cha, J.-H. J.;
Friedlander,
[12929] R. M.: Inhibition of caspase-1 slows disease progression
in a mouse
[12930] model of Huntington's disease. Nature 399: 263-267,
1999.
[12931] 748.

[12932] Arend, W. P.: Interleukin 1 receptor antagonist: a new
member
[12933] of the interleukin 1 family. J. Clin. Invest. 88: 1445-1451,
1991.
[12934] 749.

[12935] Blakemore, A. I. F.; Cox, A.; Gonzalez, A.-M.; Maskill, J. K.;

[12936] Hughes, M. E.; Wilson, R. M.; Ward, J. D.; Duff, G. W.: In-
terleukin-1

[12937] receptor antagonist allele (IL1RN*2) associated with
nephropathy in

[12938] diabetes mellitus. Hum. Genet. 97: 369-374, 1996.

[12939] 750.

[12940] Blakemore, A. I. F.; Tarlow, J. K.; Cork, M. J.; Gordon, C.;
Emery,

[12941] P.; Duff, G. W.: Interleukin-1 receptor antagonist gene
polymorphism

[12942] as a disease severity factor in systemic lupus erythemato-
sus. Arthritis

[12943] Rheum. 37: 1380-1385, 1994.

[12944] 751.

[12945] Carter, D. B.; Deibel, M. R., Jr.; Dunn, C. J.; Tomich, C.-S.
C.;

[12946] Laborde, A. L.; Slightom, J. L.; Berger, A. E.; Bienkowski, M.
J.;

[12947] Sun, F. F.; McEwan, R. N.; Harris, P. K. W.; Yem, A. W.;
Waszak, G.

[12948] A.; Chosay, J. G.; Sieu, L. C.; Hardee, M. M.; Zurcher-Neely, H. A.;

[12949] Reardon, I. M.; Heinrikson, R. L.; Truesdell, S. E.; Shelly, J. A.;

[12950] Eessalu, T. E.; Taylor, B. M.; Tracey, D. E.: Purification, cloning,

[12951] expression and biological characterization of an inter- leukin-1 receptor

[12952] antagonist protein. Nature 344: 633-638, 1990.

[12953] 752.

[12954] Dinarello, C. A.; Wolff, S. M.: The role of interleukin-1 in disease. New

[12955] Eng. J. Med. 328: 106-113, 1993.

[12956] 753.

[12957] El-Omar, E. M.; Carrington, M.; Chow, W.-H.; McColl, K. E. L.;

[12958] Bream, J. H.; Young, H. A.; Herrera, J.; Lissowska, J.; Yuan, C.-C.;

[12959] Rothman, N.; Lanyon, G.; Martin, M.; Fraumeni, J. F., Jr.; Rabkin,

[12960] C. S.: Interleukin-1 polymorphisms associated with in- creased risk

[12961] of gastric cancer. Nature 404: 398-402, 2000.

[12962] 754.

[12963] Gabay, C.; Smith, M. F., Jr.; Eidlen, D.; Arend, W. P.: Inter-
leukin

[12964] 1 receptor antagonist (IL-1Ra) is an acute-phase protein.
J. Clin.

[12965] Invest. 99: 2930-2940, 1997.

[12966] 755.

[12967] Langdahl, B. L.; Lokke, E.; Carstens, M.; Stenkjaer, L. L.;
Eriksen,

[12968] E. F.: Osteoporotic fractures are associated with an
86-base pair

[12969] repeat polymorphism in the interleukin-1-receptor antag-
onist gene

[12970] but not with polymorphisms in the interleukin-1 beta
gene. J. Bone

[12971] Miner. Res. 15: 402-414, 2000.

[12972] 756.

[12973] Craig, S. P.; Lamouroux, A.; Mallet, J.; Huttner, W.; Craig, I.

[12974] W.: Localisation of the gene for chromogranin B to chro-
mosome 20..

[12975] (Abstract) 7th Int. Cong. Hum. Genet., Berlin 1986.

[12976] 757.

[12977] Craig, S. P.; Lamouroux, A.; Mallet, J.; Huttner, W.; Craig, I.

[12978] W.: Localisation of the human gene for secretogranin 1 (chromogranin

[12979] B) to chromosome 20. (Abstract) Cytogenet. Cell Genet. 46: 600

[12980] only, 1987.

[12981] 758.

[12982] Jenkins, N. A.; Mattei, M.-G.; Gilbert, D. J.; Linard, C. G.; Mbikay,

[12983] M.; Chretien, M.; Copeland, N. G.: Assignment of secretogranin I

[12984] locus to mouse chromosome 2 by in situ hybridization and interspecific

[12985] backcross analysis. Genomics 11: 479-480, 1991.

[12986] 759.

[12987] Kluwe, L.; Mautner, V.; Parry, D. M.; Jacoby, L. B.; Baser, M.;

[12988] Gusella, J.; Davis, K.; Stavrou, D.; MacCollin, M.: The parental

[12989] origin of new mutations in neurofibromatosis 2. Neurogenetics 3:

[12990] 17-24, 2000.

[12991] 760.

[12992] Kluwe, L.; Mautner, V.-F.: Mosaicism in sporadic neurofibromatosis

[12993] 2 patients. Hum. Molec. Genet. 7: 2051-2055, 1998.

[12994] 761.

[12995] Kluwe, L.; Mautner, V.-F.: A missense mutation in the NF2 gene

[12996] results in moderate and mild clinical phenotypes of neurofibromatosis

[12997] type 2. Hum. Genet. 97: 224-227, 1996.

[12998] 762.

[12999] Knudson, A.: Asbestos and mesothelioma: genetic lessons from

[13000] a tragedy. Proc. Nat. Acad. Sci. 92: 10819-10820, 1995.

[13001] 763.

[13002] Kondziolka, D.; Lunsford, L. D.; McLaughlin, M. R.; Flickinger,

[13003] J. C.: Long-term outcomes after radiosurgery for acoustic neuromas. New

[13004] Eng. J. Med. 339: 1426-1433, 1998.

[13005] 764.

[13006] Landau, K.; Dossetor, F. M.; Hoyt, W. F.; Muci-Mendoza, R.: Retinal

[13007] hamartoma in neurofibromatosis 2. Arch. Ophthal. 108: 328-329, 1990.

[13008] 765.

[13009] Lee, D. K.; Abbott, M. L.: Familial central nervous system neoplasia:

[13010] case report of a family with von Recklinghausen's neurofi- bromatosis. Arch.

[13011] Neurol. 20: 154-160, 1969.

[13012] 766.

[13013] Legoix, P.; Legrand, M.-F.; Ollagnon, E.; Lenoir, G.; Thomas,

[13014] G.; Zucman-Rossi, J.: Characterisation of 16 polymorphic markers

[13015] in the NF2 gene: application to hemizygosity detection. Hum. Mutat. 13:

[13016] 290-293, 1999.

[13017] 767.

[13018] MacCollin, M.; Ramesh, V.; Jacoby, L. B.; Louis, D. N.; Ru-
bio,
[13019] M.-P.; Pulaski, K.; Trofatter, J. A.; Short, M. P.; Bove, C.; El-
dridge,
[13020] R.; Parry, D. M.; Gusella, J. F.: Mutational analysis of pa-
tients
[13021] with neurofibromatosis 2. Am. J. Hum. Genet. 55:
314-320, 1994.

[13022] 768.

[13023] Martuza, R. L.; Eldridge, R.: Neurofibromatosis 2 (bilateral
[13024] acoustic neurofibromatosis). New Eng. J. Med. 318:
684-688, 1988.

[13025] 769.

[13026] Martuza, R. L.; Ojemann, R. G.: Bilateral acoustic neuro-
mas:
[13027] clinical aspects, pathogenesis and treatment. Neuro-
surgery 10: 1-12,
[13028] 1982.

[13029] 770.

[13030] Mayfrank, L.; Wullich, B.; Wolff, G.; Finke, J.; Gouzoulis, E.;

[13031] Gilsbach, J. M.: Neurofibromatosis 2: a clinically and ge-
netically

[13032] heterogeneous disease? Report on 10 sporadic cases. Clin.
Genet. 38:

[13033] 362-370, 1990.

[13034] 771.

[13035] Moyes, P. D.: Familial bilateral acoustic neuroma affecting
14

[13036] members from four generations. J. Neurosurg. 29: 78-82,
1968.

[13037] 772.

[13038] Mrazek, J.; Fiser, Z.; Mrazkova, D.: Diagnosis, size, and
operation

[13039] results in 41 acoustic neurinomas. Neoplasma 35:
467-474, 1988.

[13040] 773.

[13041] Merel, P.; Hoang-Xuan, K.; Sanson, M.; Moreau-Aubry, A.;
Bijlsma,

[13042] E. K.; Lazaro, C.; Moisan, J. P.; Resche, F.; Nishisho, I.; Es-
tivill,

[13043] X.; Delattre, J. Y.; Poisson, M.; Theillet, C.; Hulsebos, T.; Delattre,
[13044] O.; Thomas, G.: Predominant occurrence of somatic mutations of the
[13045] NF2 gene in meningiomas and schwannomas. Genes Chromosomes Cancer 13:
[13046] 211-216, 1995.

[13047] 774.

[13048] Myers, R. M.; Fischer, S. G.; Lerman, L. S.; Maniatis, T.: Nearly
[13049] all single base substitutions in DNA fragments joined to a GC-clamp
[13050] can be detected by denaturing gradient gel electrophoresis. Nucleic
[13051] Acids Res. 13: 3131-3145, 1985.

[13052] 775.

[13053] Nager, G. T.: Association of bilateral VIIIth nerve tumors with
[13054] meningioma in von Recklinghausen's disease. Laryngoscope 74: 1220-1261,
[13055] 1964.

[13056] 776.

[13057] Nager, G. T.: Acoustic neuromas: pathology and differen-
tial diagnosis. Arch.
[13058] Otolaryng. 89: 252-279, 1969.

[13059] 777.

[13060] Narod, S. A.; Parry, D. M.; Parboosingh, J.; Lenoir, G. M.;
Ruttledge,
[13061] M.; Fischer, G.; Eldridge, R.; Martuza, R. L.; Frontali, M.;
Haines,
[13062] J.; Gusella, J. F.; Rouleau, G. A.: Neurofibromatosis type 2
appears
[13063] to be a genetically homogeneous disease. Am. J. Hum.
Genet. 51:
[13064] 486-496, 1992.

[13065] 778.

[13066] NIH Consensus Development Conference: Neurofibro-
matosis: conference
[13067] statement. Arch. Neurol. 45: 475-578, 1988.

[13068] 779.

[13069] Niimura, M.: Neurofibromatosis (3). Rinsho Derma (Tokyo)
15:

[13070] 653-663, 1973.

[13071] 780.

[13072] Papi, L.; De Vitis, L. R.; Vitelli, F.; Ammannati, F.; Men-
nonna,
[13073] P.; Montali, E.; Bigozzi, U.: Somatic mutations in the neu-
rofibromatosis
[13074] type 2 gene in sporadic meningiomas. Hum. Genet. 95:
347-351, 1995.
[13075] 781.

[13076] Parry, D. M.; Kaiser-Kupfer, M. I.; Eldridge, R.: Neurofibro-
matosis
[13077] 2 (NF2): lens findings in 40 patients in 5 high risk groups.
(Abstract) Am.
[13078] J. Hum. Genet. 49 (suppl.): 155, 1991.

[13079] 782.

[13080] Parry, D. M.; MacCollin, M. M.; Kaiser-Kupfer, M. I.; Pu-
laski,
[13081] K.; Nicholson, H. S.; Bolesta, M.; Eldridge, R.; Gusella, J. F.:

[13082] Germ-line mutations in the neurofibromatosis 2 gene:
correlations
[13083] with disease severity and retinal abnormalities. Am. J.

Hum. Genet. 59:
[13084] 529-539, 1996.

[13085] 783.

[13086] Pastores, G. M.; Michels, V. V.; Jack, C. R., Jr.: Early child-
hood
[13087] diagnosis of acoustic neuromas in presymptomatic indi-
viduals at risk
[13088] for neurofibromatosis 2. Am. J. Med. Genet. 41: 325-329,
1991.
[13089] 784.

[13090] Pearson-Webb, M. A.; Kaiser-Kupfer, M. I.; Eldridge, R.:
Eye
[13091] findings in bilateral acoustic (central) neurofibromatosis:
association
[13092] with presenile lens opacities and cataracts but absence of
Lisch nodules.
[13093] (Letter) New Eng. J. Med. 315: 1553-1554, 1986.

[13094] 785.

[13095] Perez Demoura, L. F.; Hayden, R. C., Jr.; Conner, G. H.: Bi-
lateral
[13096] acoustic neurinoma and neurofibromatosis. Arch. Oto-

laryng. 90: 28-34,

[13097] 1969.

[13098] 786.

[13099] Pitts, L. H.; Jackler, R. K.: Treatment of acoustic neuromas.

[13100] (Editorial) New Eng. J. Med. 339: 1471-1473, 1998.

[13101] 787.

[13102] Ragge, N. K.; Baser, M. E.; Klein, J.; Nechiporuk, A.; Sainz,

[13103] J.; Pulst, S.-M.; Riccardi, V. M.: Ocular abnormalities in neurofibromatosis

[13104] 2. Am. J. Ophthal. 120: 634-641, 1995.

[13105] 788.

[13106] Rouleau, G. A.; Seizinger, B. R.; Wertelecki, W.; Haines, J. L.;

[13107] Superneau, D. W.; Martuza, R. L.; Gusella, J. F.: Flanking markers

[13108] bracket the neurofibromatosis type 2 (NF2) gene on chro-mosome 22. Am.

[13109] J. Hum. Genet. 46: 323-328, 1990.

[13110] 789.

[13111] Rouleau, G. A.; Wertelecki, W.; Haines, J. L.; Hobbs, W. J.; Trofatter,

[13112] J. A.; Seizinger, B.; Martuza, R. L.; Superneau, D. W.; Conneally,

[13113] P. M.; Gusella, J. F.: Genetic linkage of bilateral acoustic neurofibromatosis

[13114] to DNA markers on chromosome 22. (Abstract) Cytogenet. Cell Genet. 46:

[13115] 684-685, 1987.

[13116] 790.

[13117] Rouleau, G. A.; Wertelecki, W.; Haines, J. L.; Hobbs, W. J.; Trofatter,

[13118] J. A.; Seizinger, B. R.; Martuza, R. L.; Superneau, D. W.; Conneally,

[13119] P. M.; Gusella, J. F.: Genetic linkage of bilateral acoustic neurofibromatosis

[13120] to a DNA marker on chromosome 22. Nature 329: 246-248, 1987.

[13121] 791.

[13122] Goate, A.; Chartier-Harlin, M.-C.; Mullan, M.; Brown, J.; Crawford,

[13123] F.; Fidani, L.; Giuffra, L.; Haynes, A.; Irving, N.; James, L.;

Mant,

[13124] R.; Newton, P.; Rooke, K.; Roques, P.; Talbot, C.; Pericak-Vance,

[13125] M.; Roses, A.; Williamson, R.; Rossor, M.; Owen, M.; Hardy, J.: Segregation

[13126] of a missense mutation in the amyloid precursor protein gene with

[13127] familial Alzheimer's disease. Nature 349: 704-706, 1991.

[13128] 792.

[13129] Anzick, S. L.; Kononen, J.; Walker, R. L.; Azorsa, D. O.; Tanner,

[13130] M. M.; Guan, X.-Y.; Sauter, G.; Kallioniemi, O.-P.; Trent, J. M.;

[13131] Meltzer, P. S.: AIB1, a steroid receptor coactivator amplified in

[13132] breast and ovarian cancer. Science 277: 965-968, 1997.

[13133] 793.

[13134] Bulavin, D. V.; Demidov, O. N.; Saito, S.; Kauraniemi, P.; Phillips,

[13135] C.; Amundson, S. A.; Ambrosino, C.; Sauter, G.; Nebreda, A. R.; Anderson,

[13136] C. W.; Kallioniemi, A.; Fornace, A. J., Jr.; Appella, E.: Am-

plification

[13137] of PPM1D in human tumors abrogates p53 tumor-suppressor activity. Nature

[13138] Genet. 31: 210-215, 2002.

[13139] 794.

[13140] Reguigne, I.; James, M. R.; Richard, C. W., III; Mollicone, R.;

[13141] Seawright, A.; Lowe, J. B.; Oriol, R.; Couillin, P.: The gene encoding

[13142] myeloid alpha-3-fucosyltransferase (FUT4) is located between D11S388

[13143] and D11S919 on 11q21. Cytogenet. Cell Genet. 66: 104-106, 1994.

[13144] 795.

[13145] Tetteroo, P. A. T.; de Heij, H. T.; Van den Eijnden, D. H.; Visser,

[13146] F. J.; Schoenmaker, E.; Geurts van Kessel, A. H. M.: A GDP-fucose:(Gal-beta-1-to-4)GlcNAc

[13147] alpha-1-to-3-fucosyltransferase activity is correlated with the presence

[13148] of human chromosome 11 and the expression of the Le(x), Le(y), and

[13149] sialyl-Le(x) antigens in human-mouse cell hybrids. J. Biol.

Chem. 262:
[13150] 15984-15989, 1987.

[13151] 796.

[13152] Chang, M.-L.; Eddy, R. L.; Shows, T. B.; Lau, J. T. Y.: Three

[13153] genes that encode human beta-galactoside alpha-2,3-sialyltransferases.
[13154] Structural analysis and chromosomal mapping studies. Glycobiology 5:
[13155] 319-325, 1995.

[13156] 797.

[13157] de Heij, H. T.; Tetteroo, P. A. T.; Geurts van Kessel, A. H. M.;
[13158] Schoenmaker, E.; Visser, F. J.; van den Eijnden, D. H.: Spe-cific
[13159] expression of a myeloid-associated CMP-NeuAc:Gal-beta-1-3GalNAc-alpha-R-alpha-2-3-sialyltra nsferase
[13160] and the sialyl-X determinant in myeloid human-mouse cell hybrids containing
[13161] human chromosome 11. Cancer Res. 48: 1489-1493, 1988.

[13162] 798.

[13163] Kitagawa, H.; Mattei, M.-G.; Paulson, J. C.: Genomic orga-
nization

[13164] and chromosomal mapping of the Gal-
beta-1,3GalNAc/Gal-beta-1,4GlcNAc

[13165] alpha-2,3-sialyltransferase. J. Biol. Chem. 271: 931-938,
1996.

[13166] 799.

[13167] Regan, J. W.; Kobilka, T. S.; Yang-Feng, T. L.; Caron, M. G.;
Lefkowitz,

[13168] R. J.; Kobilka, B. K.: Cloning and expression of a human
kidney cDNA

[13169] for an alpha-2-adrenergic receptor subtype. Proc. Nat.
Acad. Sci. 85:

[13170] 6301-6305, 1988.

[13171] 800.

[13172] Heinonen, P.; Koulu, M.; Pesonen, U.; Karvonen, M. K.;
Rissanen,

[13173] A.; Laakso, M.; Valve, R.; Uusitupa, M.; Scheinin, M.: Iden-
tification

[13174] of a three-amino acid deletion in the alpha-2B-adrenergic
receptor

[13175] that is associated with reduced basal metabolic rate in obese subjects. J.

[13176] Clin. Endocr. Metab. 84: 2429-2433, 1999.

[13177] 801.

[13178] Lomasney, J. W.; Lorenz, W.; Allen, L. F.; King, K.; Regan, J.

[13179] W.; Yang-Feng, T. L.; Caron, M. G.; Lefkowitz, R. J.: Expansion of

[13180] the alpha-2-adrenergic receptor family: cloning and characterization

[13181] of a human alpha-2-adrenergic receptor subtype, the gene for which

[13182] is located on chromosome 2. Proc. Nat. Acad. Sci. 87: 5094-5098,

[13183] 1990.

[13184] 802.

[13185] Sellar, G. C.; Jordon, S. A.; Bickmore, W. A.; Fantes, J. A.; van

[13186] Heyningen, V.; Whitehead, A. S.: The human serum amyloid A protein

[13187] (SAA) superfamily gene cluster: mapping to chromosome 11p15.1 by physical

[13188] and genetic linkage analysis. Genomics 19: 221-227, 1994.

[13189] 803.

[13190] Steel, D. M.; Sellar, G. C.; Uhlar, C. M.; Simon, S.; DeBeer, F.

[13191] C.; Whitehead, A. S.: A constitutively expressed serum amyloid A

[13192] protein gene (SAA4) is closely linked to, and shares structural similarities

[13193] with, an acute-phase serum amyloid A protein gene (SAA1). Genomics 16:

[13194] 447-454, 1993.

[13195] 804.

[13196] de Beer, M. C.; de Beer, F. C.; Gerardot, C. J.; Cecil, D. R.;

[13197] Webb, N. R.; Goodson M. L.; Kindy, M. S.: Structure of the mouse

[13198] Saa4 gene and its linkage to the serum amyloid A gene family. Genomics 34:

[13199] 139-142, 1996.

[13200] 805.

[13201] Watson, G.; Coade, S.; Woo, P.: Analysis of the genomic

and derived

[13202] protein structure of a novel human serum amyloid A gene, SAA4. Scand.

[13203] J. Immun. 36: 703-712, 1992.

[13204] 806.

[13205] Whitehead, A. S.; DeBeer, M. C.; Steel, D. M.; Rits, M.; Lelias,

[13206] J. M.; Lane, W. S.; DeBeer, F. C.: Identification of novel members

[13207] of the serum amyloid A protein superfamily as constitutive apolipoproteins

[13208] of high density lipoprotein. J. Biol. Chem. 267: 3862-3867, 1992.

[13209] 807.

[13210] Adler, M. J.; Coronel, C.; Shelton, E.; Seegmiller, J. E.; Dewji,

[13211] N. N.: Increased gene expression of Alzheimer disease beta-amyloid

[13212] precursor protein in senescent cultured fibroblasts. Proc. Nat. Acad.

[13213] Sci. 88: 16-20, 1991.

[13214] 808.

[13215] Ancolio, K.; Dumanchin, C.; Barelli, H.; Warter, J. M.; Brice,

[13216] A.; Campion, D.; Frebourg, T.; Checler, F.: Unusual pheno-
typic alteration

[13217] of beta amyloid precursor protein (beta-APP) maturation
by a new val715-to-met

[13218] beta-APP-770 mutation responsible for probable early-
onset Alzheimer's

[13219] disease. Proc. Nat. Acad. Sci. 96: 4119-4124, 1999.

[13220] 809.

[13221] Baek, S. H.; Ohgi, K. A.; Rose, D. W.; Koo, E. H.; Glass, C.
K.;

[13222] Rosenfeld, M. G.: Exchange of N-CoR corepressor and
Tip60 coactivator

[13223] complexes links gene expression by NF-kappa-B and
beta-amyloid precursor

[13224] protein. Cell 110: 55-67, 2002.

[13225] 810.

[13226] Bakker, E.; van Broeckhoven, C.; Haan, J.; Voorhoeve, E.;
van Hul,

[13227] W.; Levy, E.; Lieberburg, I.; Carman, M. D.; van Ommen, G.
J. B.;

[13228] Frangione, B.; Roos, R. A. C.: DNA diagnosis for hereditary cerebral
[13229] hemorrhage with amyloidosis (Dutch type). Am. J. Hum. Genet. 49:
[13230] 518-521, 1991.

[13231] 811.

[13232] Balbin, M.; Abrahamson, M.; Gustafson, L.; Nilsson, K.; Brun, A.;
[13233] Grubb, A.: A novel mutation in the beta-protein coding region of
[13234] the amyloid beta-protein precursor (APP) gene. Hum. Genet. 89: 580-582,
[13235] 1992.

[13236] 812.

[13237] Bales, K. R.; Verina, T.; Cummins, D. J.; Du, Y.; Dodel, R. C.;
[13238] Saura, J.; Fishman, C. E.; DeLong, C. A.; Piccardo, P.; Pe-
tegnief,
[13239] V.; Ghetti, B.; Paul, S. M.: Apolipoprotein E is essential for amyloid
[13240] deposition in the APP(V717F) transgenic mouse model of Alzheimer's

[13241] disease. Proc. Nat. Acad. Sci. 96: 15233-15238, 1999.

[13242] 813.

[13243] Blanquet, V.; Goldgaber, D.; Turleau, C.; Creau-Goldberg, N.; Delabar,

[13244] J.; Sinet, P. M.; Roudier, M.; de Grouchy, J.: In situ hybridization

[13245] of the beta amyloid protein (APP) cDNA to the vicinity of the interface

[13246] of 21q21 and q22 in normal and Alzheimer's disease individuals.(Abstract) Cytogenet.

[13247] Cell Genet. 46: 582 only, 1987.

[13248] 814.

[13249] Calhoun, M. E.; Wiederhold, K.-H.; Abramowski, D.; Phinney, A.

[13250] L.; Probst, A.; Sturchler-Pierrat, C.; Staufenbiel, M.; Sommer, B.;

[13251] Jucker, M.: Neuron loss in APP transgenic mice. Nature 395: 765-766,

[13252] 1998.

[13253] 815.

[13254] Cao, X.; Sudhof, T. C.: A transcriptively active complex of

APP
[13255] with Fe65 and histone acetyltransferase Tip60. Science 293: 115-120,
[13256] 2001.

[13257] 816.

[13258] Carter, D.; Campion, D.; d'Amato, T.; Jay, M.; Brice, A.; Bellis,
[13259] M.; Mallet, J.; Agid, Y.: No mutation in codon 713 of the amyloid
[13260] precursor gene in schizophrenic patients. Hum. Molec. Genet. 2:
[13261] 321, 1993.

[13262] 817.

[13263] Carter, D. A.; Desmarais, E.; Bellis, M.; Campion, D.; Clerget-Darpoux,
[13264] F.; Brice, A.; Agid, Y.; Jaillard-Serradt, A.; Mallet, J.: More missense
[13265] in amyloid gene.(Letter) Nature Genet. 2: 255-256, 1992.

[13266] 818.

[13267] Chartier-Harlin, M.-C.; Crawford, F.; Houlden, H.; Warren, A.;
[13268] Hughes, D.; Fidani, L.; Goate, A.; Rossor, M.; Roques, P.; Hardy,
[13269] J.; Mullan, M.: Early-onset Alzheimer's disease caused by mutations
[13270] at codon 717 of the beta-amyloid precursor protein gene. Nature 353:
[13271] 844-846, 1991.

[13272] 819.

[13273] Cheng, S. V.; Nadeau, J. H.; Tanzi, R. E.; Watkins, P. C.; Sacchi,
[13274] N.; Neve, R. L.; Gusella, J. F.: Synteny in man and mouse of DNA
[13275] markers from the chromosomal region linked to familial Alzheimer's

[13276] disease and Down syndrome.(Abstract) Am. J. Hum. Genet. 41: A161

[13277] only, 1987.

[13278] 820.

[13279] Citron, M.; Oltersdorf, T.; Haass, C.; McConlogue, L.; Hung, A.

[13280] Y.; Seubert, P.; Vigo-Pelfrey, C.; Lieberburg, I.; Selkoe, D. J.:

[13281] Mutation of the beta-amyloid precursor protein in familial Alzheimer's

[13282] disease increases beta-protein production. Nature 360: 672-674,

[13283] 1992.

[13284] 821.

[13285] Golden, W. L.: Personal Communication. Charlottesville, Va.

[13286] 1/12/1994.

[13287] 822.

[13288] Golden, W. L.; von Kap-herr, C.; Kurth, B.; Wright, R. M.; Flickinger,

[13289] C. J.; Eddy, R.; Shows, T.; Herr, J. C.: Refinement of the lo-

calization

[13290] of the gene for human intraacrosomal protein SP-10 (ACRV1) to the

[13291] junction of bands q23-q24 of chromosome 11 by noniso-topic in situ

[13292] hybridization. Genomics 18: 446-449, 1993.

[13293] 823.

[13294] Herr, J. C.; Wright, R. M.; Flickinger, C. J.; Eddy, R. L.; Shows,

[13295] T. B.: Assignment of the gene for human intra-acrosomal protein SP-10

[13296] (ACRV1) to the p12-q13 region of chromosome 11. (Abstract) Cytogenet.

[13297] Cell Genet. 58: 1963 only, 1991.

[13298] 824.

[13299] Nasir, J.; Lin, B.; Bucan, M.; Koizumi, T.; Nadeau, J. H.; Hayden,

[13300] M. R.: The murine homologues of the Huntington disease gene (Hdh)

[13301] and the alpha-adducin gene (Add1) map to mouse chro-mosome 5 within

[13302] a region of conserved synteny with human chromosome

4p16.3. Genomics 22:
[13303] 198-201, 1994.

[13304] 825.

[13305] Peters, L. L.; Birkenmeier, C. S.; Bronson, R. T.; White, R. A.;
[13306] Lux, S. E.; Otto, E.; Bennett, V.; Higgins, A.; Barker, J. E.: Purkinje
[13307] cell degeneration associated with erythroid ankyrin defi-ciency in
[13308] nb/nb mice. J. Cell Biol. 114: 1233-1241, 1991.

[13309] 826.

[13310] Taylor, S. A. M.; Snell, R. G.; Buckler, A.; Ambrose, C.; Duyao,
[13311] M.; Church, D.; Lin, C. S.; Altherr, M.; Bates, G. P.; Groot, N.;
[13312] Barnes, G.; Shaw, D. J.; Lehrach, H.; Wasmuth, J. J.; Harper, P. S.;
[13313] Housman, D. E.; MacDonald, M. E.; Gusella, J. F.: Cloning of the
[13314] alpha-adducin gene from the Huntington's disease candi-date region
[13315] of chromosome 4 by exon amplification. Nature Genet. 2:

223-227,
[13316] 1992.

[13317] 827.

[13318] Gilligan, D. M.; Lieman, J.; Bennett, V.: Assignment of the human
[13319] beta-adducin gene (ADD2) to 2p13-p14 by in situ hybridization. Genomics 28:
[13320] 610-612, 1995.

[13321] 828.

[13322] Muro, A. F.; Marro, M. L.; Gajovic, S.; Porro, F.; Luzzatto, L.;
[13323] Baralle, F. E.: Mild spherocytic hereditary elliptocytosis and altered
[13324] levels of alpha- and gamma-adducins in beta-adducin-deficient mice. Blood 95:
[13325] 3978-3985, 2000.

[13326] 829.

[13327] Tisminetzky, S.; Devescovi, G.; Tripodi, G.; Muro, A.; Bianchi,
[13328] G.; Colombi, M.; Moro, L.; Barlati, S.; Tuteja, R.; Baralle, F. E.

[13329] : Genomic organisation and chromosomal localisation of the gene encoding
[13330] human beta adducin. Gene 167: 313-316, 1995.

[13331] 830.

[13332] White, R. A.; Angeloni, S. V.; Pasztor, L. M.: Chromosomal localization
[13333] of the beta-adducin gene to mouse chromosome 6 and human chromosome
[13334] 2. Mammalian Genome 6: 741-743, 1995.

[13335] 831.

[13336] Abbott, C. M.; Skidmore, C. J.; Searle, A. G.; Peters, J.: De-ficiency
[13337] of adenosine deaminase in the wasted mouse. Proc. Nat. Acad. Sci. 83:
[13338] 693-695, 1986.

[13339] 832.

[13340] Adrian, G. S.; Wiginton, D. A.; Hutton, J. J.: Characteriza-tion
[13341] of normal and mutant adenosine deaminase messenger RNAs by translation
[13342] and hybridization to a cDNA probe. Hum. Genet. 68:

169-172, 1984.

[13343] 833.

[13344] Adrian, G. S.; Wiginton, D. A.; Hutton, J. J.: Structure of adenosine

[13345] deaminase mRNAs from normal and adenosine deaminase-deficient human

[13346] cell lines. Molec. Cell. Biol. 4: 1712-1717, 1984.

[13347] 834.

[13348] Aitken, D. A.; Ferguson-Smith, M. A.: Investigation of the intrachromosomal

[13349] position of the ADA locus on chromosome 20 by gene dosage studies. Cytogenet.

[13350] Cell Genet. 22: 514-517, 1978.

[13351] 835.

[13352] Aitken, D. A.; Kleijer, W. J.; Niermeijer, M. F.; Herbschleb-Voogt,

[13353] E.; Galjaard, H.: Prenatal detection of a probable heterozygote for

[13354] ADA deficiency and severe combined immunodeficiency disease using

[13355] a microradioassay. Clin. Genet. 17: 293-298, 1980.

[13356] 836.

[13357] Aiuti, A.; Slavin, S.; Aker, M.; Ficara, F.; Deola, S.; Mortel-
laro,
[13358] A.; Morecki, S.; Andolfi, G.; Tabucchi, A.; Carlucci, F.;
Marinello,
[13359] E.; Cattaneo, F.; Vai, S.; Servida, P.; Miniero, R.; Roncarolo,
M.
[13360] G.; Bordignon, C.: Correction of ADA-SCID by stem cell
gene therapy
[13361] combined with nonmyeloablative conditioning. Science
296: 2410-2413,
[13362] 2002.

[13363] 837.

[13364] Akeson, A. L.; Wiginton, D. A.; Dusing, M. R.; States, J. C.;
Hutton,
[13365] J. J.: Mutant human adenosine deaminase alleles and their
expression
[13366] by transfection into fibroblasts. J. Biol. Chem. 263:
16291-16296,
[13367] 1988.

[13368] 838.

[13369] Akeson, A. L.; Wiginton, D. A.; Hutton, J. J.: Normal and mutant
[13370] human adenosine deaminase genes. J. Cell. Biochem. 39: 217-228,
[13371] 1989.

[13372] 839.

[13373] Akeson, A. L.; Wiginton, D. A.; States, J. C.; Perme, C. M.; Dusing,
[13374] M. R.; Hutton, J. J.: Mutations in the human adenosine deaminase
[13375] gene that affect protein structure and RNA splicing. Proc. Nat. Acad.
[13376] Sci. 84: 5947-5951, 1987.

[13377] 840.

[13378] Arredondo-Vega, F. X.; Kurtzberg, J.; Chaffee, S.; Santiste-ban,
[13379] I.; Reisner, E.; Povey, M. S.; Hershfield, M. S.: Paradoxical expression
[13380] of adenosine deaminase in T cells cultured from a patient with adenosine
[13381] deaminase deficiency and combined immunodeficiency. J. Clin. Invest. 86:

[13382] 444-452, 1990.

[13383] 841.

[13384] Arredondo-Vega, F. X.; Santisteban, I.; Kelly, S.; Schloss-
man,
[13385] C. M.; Umetsu, D. T.; Hershfield, M. S.: Correct splicing
despite
[13386] mutation of the invariant first nucleotide of a 5-prime
splice site:
[13387] a possible basis for disparate clinical phenotypes in sib-
lings with
[13388] adenosine deaminase deficiency. Am. J. Hum. Genet. 54:
820-830,
[13389] 1994.

[13390] 842.

[13391] Arredondo-Vega, F. X.; Santisteban, I.; Richard, E.; Bali, P.;

[13392] Koleilat, M.; Loubser, M.; Al-Ghonaium, A.; Al-Helali, M.;
Hershfield,
[13393] M. S.: Adenosine deaminase deficiency with mosaicism for
a 'second-site
[13394] suppressor' of a splicing mutation: decline in revertant T
lymphocytes

[13395] during enzyme replacement therapy. Blood 99: 1005-1013, 2002.

[13396] 843.

[13397] Berkvens, T. M.; Gerritsen, E. J. A.; Oldenburg, M.; Breukel,

[13398] C.; Wijnen, J. T.; van Ormondt, H.; Vossen, J. M.; van der Eb, A.

[13399] J.; Meera Khan, P.: Severe combined immune deficiency due to a homozygous

[13400] 3.2-kb deletion spanning the promoter and first exon of the adenosine

[13401] deaminase gene. Nucleic Acids Res. 15: 9365-9378, 1987.

[13402] 844.

[13403] Salerno, C.; Iotti, S.; Lodi, R.; Crifo, C.; Barbiroli, B.: Failure

[13404] of muscle energy metabolism in a patient with adenylo-succinate lyase

[13405] deficiency: an in vivo study by phosphorus NMR spec-troscopy. Biochim.

[13406] Biophys. Acta 1360: 271-276, 1997.

[13407] 845.

[13408] Sebesta, I.; Krijt, J.; Kmoch, S.; Hartmannova, H.; Wojda,

M.;

[13409] Zeman, J.: Adenylosuccinase deficiency: clinical and bio-
chemical

[13410] findings in 5 Czech patients. J. Inherit. Metab. Dis. 20:
343-344,

[13411] 1997.

[13412] 846.

[13413] Stone, R. L.; Aimi, J.; Barshop, B. A.; Jaeken, J.; Van den
Berghe,

[13414] G.; Zalkin, H.; Dixon, J. E.: A mutation in adenylosuccinate
lyase

[13415] associated with mental retardation and autistic features.
Nature

[13416] Genet. 1: 59-63, 1992.

[13417] 847.

[13418] Van Keuren, M. L.; Hart, I.; Kao, F.-T.; Neve, R. L.; Bruns,
G.

[13419] A. P.; Kurnit, D. M.; Patterson, D.: Human chromosome 22
corrects

[13420] the defect in the CHO mutant (Ade-I) lacking adenylosuc-
cinase activity.

[13421] (Abstract) Am. J. Hum. Genet. 39: A172 only, 1986.

[13422] 848.

[13423] Van Keuren, M. L.; Hart, I. M.; Kao, F.-T.; Neve, R. L.; Bruns,

[13424] G. A. P.; Kurnit, D. M.; Patterson, D.: A somatic cell hybrid with

[13425] a single human chromosome 22 corrects the defect in the CHO mutant

[13426] (Ade-I) lacking adenylosuccinase activity. Cytogenet. Cell Genet. 44:

[13427] 142-147, 1987.

[13428] 849.

[13429] Verginelli, D.; Luckow, B.; Crifo, C.; Salerno, C.; Gross, M.

[13430] : Identification of new mutations in the adenylosuccinate lyase gene

[13431] associated with impaired enzyme activity in lymphocytes and red blood

[13432] cells. Biochim. Biophys. Acta 1406: 81-84, 1998.

[13433] 850.

[13434] Wong, L.-J. C.; O'Brien, W. E.: Characterization of the cDNA

[13435] and the gene encoding murine adenylosuccinate lyase.

Genomics 28:
[13436] 341-343, 1995.

[13437] 851.

[13438] Lai, L.; Hart, I.; Patterson, D.: Human chromosome 1 corrects
[13439] the defect in the CHO mutant (Ade-H) deficient in a branch point enzyme
[13440] in purine de novo biosynthesis. (Abstract) Cytogenet. Cell Genet. 51:
[13441] 1028 only, 1989.

[13442] 852.

[13443] Lai, L.-W.; Hart, I. M.; Patterson, D.: A gene correcting the
[13444] defect in the CHO mutant Ade(-)H, deficient in a branch point enzyme
[13445] (adenylosuccinate synthetase) of de novo purine biosynthesis, is located
[13446] on the long arm of chromosome 1. Genomics 9: 322-328, 1991.

[13447] 853.

[13448] Powell, S. M.; Zalkin, H.; Dixon, J. E.: Cloning and characterization

[13449] of the cDNA encoding human adenylosuccinate synthetase. FEBS Lett. 303:
[13450] 4-10, 1992.

[13451] 854.

[13452] Parma, J.; Stengel, D.; Gannage, M.-H.; Poyard, M.; Barouki, R.;
[13453] Hanoune, J.: Sequence of a human brain adenylyl cyclase partial cDNA:
[13454] evidence for a consensus cyclase domain. Biochem. Biophys. Res. Commun. 179:
[13455] 455-462, 1991.

[13456] 855.

[13457] Stengel, D.; Parma, J.; Gannage, M.-H.; Roeckel, N.; Mattei, M.-G.;
[13458] Barouki, R.; Hanoune, J.: Different chromosomal localization of two
[13459] adenylyl cyclase genes expressed in human brain. Hum. Genet. 90:
[13460] 126-130, 1992.

[13461] 856.

[13462] Edelhoff, S.; Villacres, E. C.; Storm, D. R.; Disteche, C. M.:

[13463] Mapping of adenylyl cyclase genes type I, II, III, IV, V, and VI in

[13464] mouse. Mammalian Genome 6: 111-113, 1995.

[13465] 857.

[13466] Gaudin, C.; Homcy, C. J.; Ishikawa, Y.: Mammalian adenylyl cyclase

[13467] family members are randomly located on different chromosomes. Hum.

[13468] Genet. 94: 527-529, 1994.

[13469] 858.

[13470] Wong, S. T.; Trinh, K.; Hacker, B.; Chan, G. C. K.; Lowe, G.; Gaggar,

[13471] A.; Xia, Z.; Gold, G. H.; Storm, D. R.: Disruption of the type III

[13472] adenylyl cyclase gene leads to peripheral and behavioral anosmia in

[13473] transgenic mice. Neuron 27: 487-497, 2000.

[13474] 859.

[13475] Abdel-Majid, R. M.; Leong, W. L.; Schalkwyk, L. C.; Smallman, D.

[13476] S.; Wong, S. T.; Storm, D. R.; Fine, A.; Dobson, M. J.;

Guernsey,

[13477] D. L.; Neumann, P. E.: Loss of adenylyl cyclase I activity disrupts

[13478] patterning of mouse somatosensory cortex. Nature Genet. 19: 289-291,

[13479] 1998.

[13480] 860.

[13481] Villacres, E. C.; Xia, Z.; Bookbinder, L. H.; Edelhoff, S.; Disteche,

[13482] C. M.; Storm, D. R.: Cloning, chromosomal mapping, and expression

[13483] of human fetal brain type I adenylyl cyclase. Genomics 16: 473-478,

[13484] 1993.

[13485] 861.

[13486] Welker, E.; Armstrong-James, M.; Bronchti, G.; Ourednik, W.; Gheorghita-Baechler,

[13487] F.; Dubois, R.; Guernsey, D. L.; Van der Loos, H.; Neumann, P. E.

[13488] : Altered sensory processing in the somatosensory cortex of the mutant

[13489] mouse barrelless. Science 271: 1864-1867, 1996.

[13490] 862.

[13491] Amor, J. C.; Harrison, D. H.; Kahn, R. A.; Ringe, D.: Structure

[13492] of the human ADP-ribosylation factor 1 complexed with GDP. Nature 372:

[13493] 704-708, 1994.

[13494] 863.

[13495] Bobak, D. A.; Nightingale, M. S.; Murtagh, J. J.; Price, S. R.;

[13496] Moss, J.; Vaughan, M.: Molecular cloning, characterization, and expression

[13497] of human ADP-ribosylation factors: two guanine nucleotide-dependent

[13498] activators of cholera toxin. Proc. Nat. Acad. Sci. 86: 6101-6105,

[13499] 1989.

[13500] 864.

[13501] Hirai, M.; Kusuda, J.; Hashimoto, K.: Assignment of human ADP

[13502] ribosylation factor (ARF) genes ARF1 and ARF3 to chromosomes 1q42

[13503] and 12q13, respectively. Genomics 34: 263-265, 1996.

[13504] 865.

[13505] Lee, C.-M.; Haun, R. S.; Tsai, S.-C.; Moss, J.; Vaughan, M.: Characterization

[13506] of the human gene encoding ADP-ribosylation factor 1, a guanine nucleotide-binding

[13507] activator of cholera toxin. J. Biol. Chem. 267: 9028-9034, 1992.

[13508] 866.

[13509] Mossessova, E.; Gulbis, J. M.; Goldberg, J.: Structure of the

[13510] guanine nucleotide exchange factor Sec7 domain of human Arno and analysis

[13511] of the interaction with ARF GTPase. Cell 92: 415-423, 1998.

[13512] 867.

[13513] Edman, K.; Maret, W.: Alcohol dehydrogenase genes: restriction

[13514] fragment length polymorphisms for ADH4 (pi-ADH) and ADH5 (chi-ADH)

[13515] for construction of haplotypes among different ADH classes. Hum.

[13516] Genet. 90: 395-401, 1992.

[13517] 868.

[13518] Li, T.-K.; Bosron, W. F.; Dafeldecker, W. P.; Lange, L. G.; Vallee,

[13519] B. L.: Isolation of PI-alcohol dehydrogenase of human liver: is it

[13520] a determinant of alcoholism? Proc. Nat. Acad. Sci. 74: 4378-4381,

[13521] 1977.

[13522] 869.

[13523] Mardh, G.; Dingley, A. L.; Auld, D. S.; Vallee, B. L.: Human class

[13524] II (pi) alcohol dehydrogenase has a redox-specific function in norepinephrine

[13525] metabolism. Proc. Nat. Acad. Sci. 83: 8908-8912, 1986.

[13526] 870.

[13527] McPherson, J. D.; Smith, M.; Wagner, C.; Wasmuth, J. J.; Hoog,

[13528] J.-O.: Mapping of the class II alcohol dehydrogenase gene locus to

[13529] 4q22. (Abstract) Cytogenet. Cell Genet. 51: 1043 only, 1989.

[13530] 871.

[13531] Vallee, B. L.; Bazzone, T. J.: Isozymes of human liver alcohol

[13532] dehydrogenase.In: Rattazzi, M. C.; Scandalios, J. G.; Whitt, G. S.

[13533] : Isozymes. Current Topics in Biological and Medical Research. New

[13534] York: Alan R. Liss (pub.) 8: 1983. Pp. 219-244.

[13535] 872.

[13536] Badawi, A. F.; Hirvonen, A.; Bell, D. A.; Lang, N. P.; Kadlubar,

[13537] F. F.: Role of aromatic amine acetyltransferases, NAT1 and NAT2,

[13538] in carcinogen-DNA adduct formation in the human urinary bladder. Cancer

[13539] Res. 55: 5230-5237, 1995.

[13540] 873.

[13541] Bell, D. A.; Badawi, A. F.; Lang, N. P.; Ilett, K. P.; Kadlubar,

[13542] F. F.; Hirvonen, A.: Polymorphism in the N-acetyltransferase 1 (NAT1)

[13543] polyadenylation signal: association of NAT1*10 allele with

higher
[13544] N-acetylation activity in bladder and colon tissue. Cancer Res. 55:
[13545] 5226-5229, 1995.

[13546] 874.

[13547] Bell, D. A.; Stephens, E. A.; Castranio, T.; Umbach, D. M.; Watson,
[13548] M.; Deakin, M.; Elder, J.; Hendrickse, C.; Duncan, H.; Strange, R.
[13549] C.: Polyadenylation polymorphism in the acetyltransferase 1 gene
[13550] (NAT1) increases risk of colorectal cancer. Cancer Res. 55: 3537-3542,
[13551] 1995.

[13552] 875.

[13553] Blum, M.; Grant, D. M.; McBride, W.; Heim, M.; Meyer, U. A.: Human
[13554] arylamine N-acetyltransferase genes: isolation, chromo-somal localization,
[13555] and functional expression. DNA Cell Biol. 9: 193-203, 1990.
[13556] 876.

[13557] Bouchardy, C.; Mitrunen, K.; Wikman, H.; Husgafvel-Pursiainen,
[13558] K.; Dayer, P.; Benhamou, S.; Hirvonen, A.: N-acetyltransferase NAT1
[13559] and NAT2 genotypes and lung cancer risk. Pharmacogenetics 8: 291-298,
[13560] 1998.

[13561] 877.

[13562] Butcher, N. J.; Ilett, K. F.; Minchin, R. F.: Functional polymorphism
[13563] of the human arylamine N-acetyltransferase type 1 gene caused by C190T
[13564] and G560A mutations. Pharmacogenetics 8: 67-72, 1998.

[13565] 878.

[13566] Doll, M. A.; Jiang, W.; Deitz, A. C.; Rustan, T. D.; Hein, D. W.
[13567] : Identification of a novel allele at the human NAT1 acetyltransferase
[13568] locus. Biochem. Biophys. Res. Commun. 233: 584-591, 1997.
[13569] 879.

[13570] Evans, D. A. P.: Personal Communication. Saudi Arabia 6/1/1998.

[13571] 880.

[13572] Evans, D. A. P.: Genetic Factors in Drug Therapy: Clinical and

[13573] Molecular Pharmacogenetics. Cambridge: Cambridge Univ. Press , 1993.

[13574] Pp. 211-305.

[13575] 881.

[13576] Hickman, D.; Risch, A.; Buckle, V.; Spurr, N. K.; Jeremiah, S.

[13577] J.; McCarthy, A.; Sim, E.: Chromosomal localization of human genes

[13578] for arylamine N-acetyltransferase. Biochem. J. 297: 441-445, 1994.

[13579] 882.

[13580] Matas, N.; Thygesen, P.; Stacey, M.; Risch, A.; Sim, E.: Mapping

[13581] AAC1, AAC2 and AACP, the genes for arylamine N-acetyltransferases,

[13582] carcinogen metabolising enzymes on human chromosome 8p22, a region

[13583] frequently deleted in tumours. Cytogenet. Cell Genet. 77: 290-295,
[13584] 1997.

[13585] 883.

[13586] Mattano, S. S.; Erickson, R. P.; Nesbitt, M. N.; Weber, W. W.

[13587] : Linkage of Nat and Es-1 in the mouse and development of strains
[13588] congenic for N-acetyltransferase. J. Hered. 79: 430-433, 1988.
[13589] 884.

[13590] Moisio, A.-L.; Sistonen, P.; Mecklin, J.-P.; Jarvinen, H.; Pel-tomaki,
[13591] P.: Genetic polymorphisms in carcinogen metabolism and their association
[13592] to hereditary nonpolyposis colon cancer. Gastroenterology 115: 1387-1394,
[13593] 1998.

[13594] 885.

[13595] Sim, E.; Payton, M.; Noble, M.; Minchin, R.: An update on genetic,
[13596] structural and functional studies of arylamine N-

acetyltransferases
[13597] in eucaryotes and procaryotes. Hum. Molec. Genet. 9:
2435-2441,
[13598] 2000.

[13599] 886.

[13600] Smelt, V. A.; Upton, A.; Adjaye, J.; Payton, M. A.; Boukou-
vala,
[13601] S.; Johnson, N.; Mardon, H. J.; Sim, E.: Expression of ary-
lamine
[13602] N-acetyltransferases in pre-term placentas and in human
pre-implantation
[13603] embryos. Hum. Molec. Genet. 9: 1101-1107, 2000.

[13604] 887.

[13605] Vatsis, K. P.; Martell, K. J.; Weber, W. W.: Diverse point
mutations
[13606] in the human gene for polymorphic N-acetyltransferase.
Proc. Nat.
[13607] Acad. Sci. 88: 6333-6337, 1991.

[13608] 888.

[13609] Vatsis, K. P.; Weber, W. W.: Structural heterogeneity of
Caucasian

[13610] N-acetyltransferase at the NAT1 gene locus. Arch. Biochem. Biophys. 301:

[13611] 71-76, 1993.

[13612] 889.

[13613] Vatsis, K. P.; Weber, W. W.; Bell, D. A.; Dupret, J.-M.; Price

[13614] Evans, D. A.; Grant, D. M.; Hein, D. W.; Lin, H. J.; Meyer, U. A.;

[13615] Relling, M. V.; Sim, E.; Suzuki, T.; Yamazoe, Y.: Nomencla-ture for

[13616] N-acetyltransferases. Pharmacogenetics 5: 1-17, 1995.

[13617] 890.

[13618] Huebner, K.; Kastury, K.; Druck, T.; Salcini, A. E.; Lanfran-cone,

[13619] L.; Pelicci, G.; Lowenstein, E.; Li, W.; Park, S.-H.; Canniz-zaro,

[13620] L.; Pelicci, P. G.; Schlessinger, J.: Chromosome locations of genes

[13621] encoding human signal transduction adapter proteins, Nck (NCK), Shc

[13622] (SHC1), and Grb2 (GRB2). Genomics 22: 281-287, 1994.

[13623] 891.

[13624] Asimakopoulos, F. A.; White, N. J.; Nacheva, E. P.; Green, A. R.

[13625] : The human CD40 gene lies within chromosome 20q deletions associated

[13626] with myeloid malignancies. Brit. J. Haemat. 92: 127-130, 1996.

[13627] 892.

[13628] Castigli, E.; Alt, F. W.; Davidson, L.; Bottaro, A.; Mizoguchi,

[13629] E.; Bhan, A. K.; Geha, R. S.: CD40-deficient mice generated by recombination-activating

[13630] gene-2-deficient blastocyst complementation. Proc. Nat. Acad. Sci. 91:

[13631] 12135-12139, 1994.

[13632] 893.

[13633] Clark, E. A.: CD40: a cytokine receptor in search of a lig-and. Tissue

[13634] Antigens 35: 33-36, 1990.

[13635] 894.

[13636] Erickson, L. D.; Durell, B. G.; Vogel, L. A.; O'Connor, B. P.;

[13637] Cascalho, M.; Yasui, T.; Kikutani, H.; Noelle, R. J.: Short-circuiting

[13638] long-lived humoral immunity by the heightened engage-
ment of CD40. J.

[13639] Clin. Invest. 109: 613-620, 2002.

[13640] 895.

[13641] Ferrari, S.; Giliani, S.; Insalaco, A.; Al-Ghonaium, A.;
Soresina,

[13642] A. R.; Loubser, M.; Avanzini, M. A.; Marconi, M.; Badolato,
R.; Ugazio,

[13643] A. G.; Levy, Y.; Catalan, N.; Durandy, A.; Tbakhi, A.; No-
tarangelo,

[13644] L. D.; Plebani, A.: Mutations of CD40 gene cause an auto-
somal recessive

[13645] form of immunodeficiency with hyper IgM. Proc. Nat.
Acad. Sci. 98:

[13646] 12614-12619, 2001.

[13647] 896.

[13648] Kato, K.; Cantwell, M. J.; Sharma, S.; Kipps, T. J.: Gene
transfer

[13649] of CD40-ligand induces autologous immune recognition
of chronic lymphocytic

[13650] leukemia B cells. J. Clin. Invest. 101: 1133-1141, 1998.

[13651] 897.

[13652] Kawabe, T.; Naka, T.; Yoshida, K.; Tanaka, T.; Fujiwara, H.; Suematsu,

[13653] S.; Yoshida, N.; Kishimoto, T.; Kikutani, H.: The immune responses

[13654] in CD40-deficient mice: impaired immunoglobulin class switching and

[13655] germinal center formation. Immunity 1: 167-178, 1994.

[13656] 898.

[13657] Lafage-Pochitaloff, M.; Herman, P.; Birg, F.; Galizzi, J.-P.; Simonetti,

[13658] J.; Mannoni, P.; Banchereau, J.: Localization of the human CD40 gene

[13659] to chromosome 20, bands q12-q13.2. Leukemia 8: 1172-1175, 1994.

[13660] 899.

[13661] Mach, F.; Schonbeck, U.; Sukhova, G. K.; Atkinson, E.; Libby, P.

[13662] : Reduction of atherosclerosis in mice by inhibition of CD40 signalling. Nature 394:

[13663] 200-203, 1998.

[13664] 900.

[13665] Schonbeck, U.; Libby, P.: The CD40/CD154 receptor/lig-
and dyad. Cell
[13666] Molec. Life Sci. 58: 4-43, 2001.

[13667] 901.

[13668] Stamenkovic, I.; Clark, E. A.; Seed, B.: A B-lymphocyte ac-
tivation
[13669] molecule related to the nerve growth factor receptor and
induced by
[13670] cytokines in carcinomas. EMBO J. 8: 1403-1410, 1989.

[13671] 902.

[13672] Tan, J.; Town, T.; Paris, D.; Mori, T.; Suo, Z.; Crawford, F.;

[13673] Mattson, M. P.; Flavell, R. A.; Mullan, M.: Microglial activa-
tion
[13674] resulting from CD40-CD40L interaction after beta-
amyloid stimulation. Science 286:
[13675] 2352-2355, 1999.

[13676] 903.

[13677] van Kooten, C.; Banchereau, J.: CD40-CD40 ligand. J.
Leuko.

[13678] Biol. 67: 2-17, 2000.

[13679] 904.

[13680] Gras, M. P.; Laabi, Y.; Linares-Cruz, G.; Blondel, M. O.; Rigaut,

[13681] J. P.; Brouet, J. C.; Leca, G.; Haguenauer-Tsapis, R.; Tsapis, A.

[13682] : BCMAp: an integral membrane protein in the Golgi apparatus of human

[13683] mature B lymphocytes. Int. Immun. 7: 1093 only, 1995.

[13684] 905.

[13685] Hatzoglou, A.; Roussel, J.; Bourgeade, M.-F.; Rogier, E.; Madry,

[13686] C.; Inoue, J.; Devergne, O.; Tsapis, A.: TNF receptor family member

[13687] BCMA (B cell maturation) associates with TNF receptor-associated factor

[13688] (TRAF) 1, TRAF2, and TRAF3 and activates NF-kappa-B, Elk-1, c-Jun

[13689] N-terminal kinase, and p38 mitogen-activated protein kinase. J. Immun. 165:

[13690] 1322-1330, 2000.

[13691] 906.

[13692] Laabi, Y.; Gras, M. P.; Carbonnel, F.; Brouet, J. C.; Berger, R.;

[13693] Larsen, C. J.; Tsapis, A.: A new gene, BCM, on chromo-some 16 is fused

[13694] to the interleukin 2 gene by a t(4;16)(q26;p13) transloca-tion in a

[13695] malignant T cell lymphoma. EMBO J. 11: 3897-3904, 1992.

[13696] 907.

[13697] Bhatia, K.; Huppi, K.; McKeithan, T.; Siwarski, D.; Mushin-ski,

[13698] J. F.; Magrath, I.: Mouse bcl-3: cDNA structure, mapping and stage-dependent

[13699] expression in B lymphocytes. Oncogene 6: 1569-1573, 1991.

[13700] 908.

[13701] Franzoso, G.; Bours, V.; Park, S.; Tomita-Yamaguchi, M.; Kelly,

[13702] K.; Siebenlist, U.: The candidate oncoprotein Bcl-3 is an antagonist

[13703] of p50/NF-kappa-B-mediated inhibition. Nature 359:

338-342, 1992.

[13704] 909.

[13705] Mitchell, T. C.; Hildeman, D.; Kedl, R. M.; Teague, T. K.; Schaefer,

[13706] B. C.; White, J.; Zhu, Y.; Kappler, J.; Marrack, P.: Immunological

[13707] adjuvants promote activated T cell survival via induction of Bcl-3. Nature

[13708] Immun. 2: 397-402, 2001.

[13709] 910.

[13710] Anstee, D. J.: The blood group MNSs-active sialoglycoproteins. Seminars

[13711] Hemat. 18: 13-31, 1981.

[13712] 911.

[13713] Bias, W. B.; Meyers, D. A.: Segregation and linkage analysis of

[13714] the Stoltzfus blood group (SF). (Abstract) Cytogenet. Cell Genet. 25:

[13715] 137, 1979.

[13716] 912.

[13717] Blumenfeld, O. O.; Adamany, A. M.: Structural

(glycophorins) of
[13718] the human erythrocyte membrane. Proc. Nat. Acad. Sci. 75: 2727-2731,
[13719] 1978.

[13720] 913.

[13721] Blumenfeld, O. O.; Huang, C.-H.: Molecular genetics of the glycophorin
[13722] gene family, the antigens for MNSs blood groups: multiple gene rearrangements
[13723] and modulation of splice site usage result in extensive diversification. Hum.
[13724] Mutat. 6: 199-209, 1995.

[13725] 914.

[13726] Cook, P. J. L.; Lindenbaum, R. H.; Salonen, R.; de la Chapelle,
[13727] A.; Daker, M. G.; Buckton, K. E.; Noades, J. E.; Tippett, P.: The
[13728] MNSs blood groups of families with chromosome 4 rearrangements. Ann.
[13729] Hum. Genet. 45: 39-47, 1981.

[13730] 915.

[13731] Cook, P. J. L.; Noades, J. E.; Lomas, C. G.; Buckton, K. E.; Robson,

[13732] E. B.: Exclusion mapping illustrated by the MNSs blood group. Ann.

[13733] Hum. Genet. 44: 61-73, 1980.

[13734] 916.

[13735] Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Slaughter, C. A.;

[13736] Gray, J. E.; Blank, C. E.; James, F. E.; Ridler, M. A. C.; Insley,

[13737] J.; Hulten, M.: Segregation of ABO, AK-1 and ACON-S in families with

[13738] abnormalities of chromosome 9. Ann. Hum. Genet. 41: 365-377, 1978.

[13739] 917.

[13740] Divelbiss, J.; Shiang, R.; German, J.; Moore, J.; Murray, J. C.;

[13741] Patil, S. R.: Refinement of the physical location of glycophorin

[13742] A and beta fibrinogen using in situ hybridization and RFLP analysis.

[13743] (Abstract) Cytogenet. Cell Genet. 51: 991, 1989.

[13744] 918.

[13745] Evans, R. M.: The steroid and thyroid hormone receptor superfamily. Science 240:

[13746] 889-895, 1988.

[13747] 919.

[13748] Bader, B. L.; Rayburn, H.; Crowley, D.; Hynes, R. O.: Extensive

[13749] vasculogenesis, angiogenesis, and organogenesis precede lethality

[13750] in mice lacking all alpha-V integrins. Cell 95: 507-519, 1998.

[13751] 920.

[13752] Fernandez-Ruiz, E.; de Villena, F. P.-M.; Rodriguez de Cordoba,

[13753] S.; Sanchez-Madrid, F.: Regional localization of the human vitronectin

[13754] receptor alpha-subunit gene (VNRA) to chromosome 2q31-q32. Cytogenet.

[13755] Cell Genet. 62: 26-28, 1993.

[13756] 921.

[13757] Sims, M. A.; Field, S. D.; Barnes, M. R.; Shaikh, N.; Elling-

ton,

[13758] K.; Murphy, K. E.; Spurr, N.; Campbell, D. A.: Cloning and characterisation

[13759] of ITGAV, the genomic sequence for human cell adhesion protein (vitronectin)

[13760] receptor alpha subunit, CD51. Cytogenet. Cell Genet. 89: 268-271,

[13761] 2000.

[13762] 922.

[13763] Suzuki, S.; Argraves, W. S.; Pytela, R.; Arai, H.; Krusius, T.;

[13764] Pierschbacher, M. D.; Ruoslahti, E.: cDNA and amino acid sequences

[13765] of the cell adhesion protein receptor recognizing vitronectin reveal

[13766] a transmembrane domain and homologies with other adhesion protein

[13767] receptors. Proc. Nat. Acad. Sci. 83: 8614-8618, 1986.

[13768] 923.

[13769] Blachly-Dyson, E.; Baldini, A.; Litt, M.; McCabe, E. R. B.; Forte,

[13770] M.: Human genes encoding the voltage-dependent anion

channel (VDAC)
[13771] of the outer mitochondrial membrane: mapping and identification of
[13772] two new isoforms. Genomics 20: 62-67, 1994.

[13773] 924.

[13774] Blachly-Dyson, E.; Zambronicz, E. B.; Yu, W. H.; Adams, V.; McCabe,
[13775] E. R. B.; Adelman, J.; Colombini, M.; Forte, M.: Cloning and functional
[13776] expression in yeast of two human isoforms of the outer mitochondrial
[13777] membrane channel, the voltage-dependent anion channel. J. Biol. Chem. 268:
[13778] 1835-1841, 1993.

[13779] 925.

[13780] Huizing, M.; Ruitenbeek, W.; van den Heuvel, L. P.; Dolce, V.;
[13781] Iacobazzi, V.; Smeitink, J. A. M.; Palmieri, F.; Trijbels, J. M. F.
[13782] : Human mitochondrial transmembrane metabolite carriers: tissue distribution
[13783] and its implication for mitochondrial disorders. J. Bioen-

erg. Biomembr. 30:
[13784] 277-284, 1998.

[13785] 926.

[13786] Aubry, M.; Demczuk, S.; Desmaze, C.; Aikem, M.; Aurias, A.; Julien,
[13787] J.-P.; Rouleau, G. A.: Isolation of a zinc finger gene consistently
[13788] deleted in DiGeorge syndrome. Hum. Molec. Genet. 2: 1583-1587, 1993.
[13789] 927.

[13790] Myslinski, E.; Krol, A.; Carbon, P.: ZNF76 and ZNF143 are two
[13791] human homologs of the transcriptional activator Staf. J. Biol. Chem. 273:
[13792] 21998-22006, 1998.

[13793] 928.

[13794] Saleh, M.; Selleri, L.; Evans, G. A.: A novel zinc finger gene
[13795] on human chromosome 1qter that is alternatively spliced in human tissues
[13796] and cell lines. Am. J. Hum. Genet. 52: 192-203, 1993.

[13797] 929.

[13798] Tommerup, N.; Aagaard, L.; Lund, C. L.; Boel, E.; Baxendale, S.;

[13799] Bates, G. P.; Lehrach, H.; Vissing, H.: A zinc-finger gene ZNF141

[13800] mapping at 4p16.3/D4S90 is a candidate gene for the Wolf-Hirschhorn

[13801] (4p-) syndrome. Hum. Molec. Genet. 2: 1571-1575, 1993.

[13802] 930.

[13803] McAlpine, P. J.; Coopland, G.; Guy, C.; James, S.; Komarnicki,

[13804] L.; MacDonald, M.; Stranc, L.; Lewis, M.; Philipps, S.; Coghlan, G.;

[13805] Kaita, H.; Cox, D. W.; Guinto, E. R.; MacGillivray, R.: Mapping the

[13806] genes for erythrocytic alpha-spectrin 1 (SPTA1) and coagulation factor

[13807] V (F5). (Abstract) Cytogenet. Cell Genet. 51: 1042, 1989.

[13808] 931.

[13809] de Silva, H. V.; Harmony, J. A. K.; Stuart, W. D.; Gil, C. M.;

[13810] Robbins, J.: Apolipoprotein J: structure and tissue distribution. Biochemistry 29:

[13811] 5380-5389, 1990.

[13812] 932.

[13813] de Silva, H. V.; Stuart, W. D.; Park, Y. B.; Mao, S. J. T.; Gil,

[13814] C. M.; Wetterau, J. R.; Busch, S. J.; Harmony, J. A. K.: Purifi-
cation

[13815] and characterization of apolipoprotein J. J. Biol. Chem.
265: 14292-14297,

[13816] 1990.

[13817] 933.

[13818] Dietzsch, E.; Murphy, B. F.; Kirszbaum, L.; Walker, I. D.;
Garson,

[13819] O. M.: Regional localization of the gene for clusterin
(SP-40,40;

[13820] gene symbol CLI) to human chromosome 8p12-p21. Cy-
togenet. Cell Genet. 61:

[13821] 178-179, 1992.

[13822] 934.

[13823] Dragunow, M.; Preston, K.; Dodd, J.; Young, D.; Lawlor, P.;
Christie,

[13824] D.: Clusterin accumulates in dying neurons following sta-
tus epilepticus. Molec.

[13825] Brain Res. 32: 279-290, 1995.

[13826] 935.

[13827] Duguid, J. R.; Bohmont, C. W.; Liu, N.; Tourtellotte, W. W.: Changes

[13828] in brain gene expression shared by scrapie and Alzheimer disease. Proc.

[13829] Nat. Acad. Sci. 86: 7260-7264, 1989.

[13830] 936.

[13831] Fink, T. M.; Zimmer, M.; Tschopp, J.; Etienne, J.; Jenne, D. E.;

[13832] Lichter, P.: Human clusterin (CLI) maps to 8p21 in proximity to the

[13833] lipoprotein lipase (LPL) gene. Genomics 16: 526-528, 1993.

[13834] 937.

[13835] Han, B. H.; DeMattos, R. B.; Dugan, L. L.; Kim-Han, J. S.; Brendza,

[13836] R. P.; Fryer, J. D.; Kierson, M.; Cirrito, J.; Quick, K.; Harmony,

[13837] J. A. K.; Aronow, B. J.; Holtzman, D. M.: Clusterin contributes to

[13838] caspase-3-independent brain injury following neonatal hypoxia-ischemia. Nature

[13839] Med. 7: 338-343, 2001.

[13840] 938.

[13841] Jenne, D. E.; Tschopp, J.: Clusterin: the intriguing guises of

[13842] a widely expressed glycoprotein. Trends Biochem. Sci. 17: 154-159,

[13843] 1992.

[13844] 939.

[13845] Kamboh, M. I.; Harmony, J. A. K.; Sepehrnia, B.; Nwankwo, M.;

[13846] Ferrell, R. E.: Genetic studies of human apolipoproteins. XX. Genetic

[13847] polymorphism of apolipoprotein J and its impact on quantitative lipid

[13848] traits in normolipidemic subjects. Am. J. Hum. Genet. 49: 1167-1173,

[13849] 1991.

[13850] 940.

[13851] Kirszbaum, L.; Sharpe, J. A.; Murphy, B.; d'Apice, A. J. F.;

Classon,

[13852] B.; Hudson, P.; Walker, I. D.: Molecular cloning and characterization

[13853] of the novel, human complement-associated protein, SP-40,40: a link

[13854] between the complement and reproductive systems. EMBO J. 8: 711-718,

[13855] 1989.

[13856] 941.

[13857] Murphy, B. F.; Kirszbaum, L.; Walker, I. D.; d'Apice, A. J. F.

[13858] : SP-40,40, a newly identified normal human serum protein found in

[13859] the SC5b-9 complex of complement and in the immune deposits in glomerulonephritis. J.

[13860] Clin. Invest. 81: 1858-1864, 1988.

[13861] 942.

[13862] O'Bryan, M. K.; Baker, H. W. G.; Saunders, J. R.; Kirszbaum, L.;

[13863] Walker, I. D.; Hudson, P.; Liu, D. Y.; Glew, M. D.; d'Apice, A. J.

[13864] F.; Murphy, B. F.: Human seminal clusterin (SP-40,40):

isolation
[13865] and characterization. J. Clin. Invest. 85: 1477-1486, 1990.
[13866] 943.
[13867] Purrello, M.; Bettuzzi, S.; Di Pietro, C.; Mirabile, E.; Di Blasi,
[13868] M.; Rimini, R.; Grzeschik, K.-H.; Ingletti, C.; Corti, A.; Sichel,
[13869] G.: The gene for SP-40,40, human homolog of rat sulfated glycoprotein
[13870] 2, rat clusterin, and rat testosterone-repressed prostate message
[13871] 2, maps to chromosome 8. Genomics 10: 151-156, 1991.
[13872] 944.
[13873] Slawin, K.; Sawczuk, I. S.; Olsson, C. A.; Buttyan, R.: Chromosomal
[13874] assignment of the human homologue encoding SGP-2. Biochem. Biophys.
[13875] Res. Commun. 172: 160-164, 1990.
[13876] 945.
[13877] Alves, S.; Prata, M.-J.; Ferreira, F.; Amorim, A.: Thiopurine

[13878] methyltransferase pharmacogenetics: alternative molecu-
lar diagnosis
[13879] and preliminary data from Northern Portugal. Pharmaco-
genetics 9:
[13880] 257-261, 1999.

[13881] 946.

[13882] Ameyaw, M.-M.; Collie-Duguid, E. S. R.; Powrie, R. H.;
Ofori-Adjei,
[13883] D.; McLeod, H. L.: Thiopurine methyltransferase alleles in
British
[13884] and Ghanaian populations. Hum. Molec. Genet. 8:
367-370, 1999.
[13885] 947.

[13886] Collie-Duguid, E. S. R.; Pritchard, S. C.; Powrie, R. H.;
Sludden,
[13887] J.; Collier, D. A.; Li, T.; McLeod, H. L.: The frequency and
distribution
[13888] of thiopurine methyltransferase alleles in Caucasian and
Asian populations. Pharmacogenetics 9:
[13889] 37-42, 1999.

[13890] 948.

[13891] Evans, W. E.; Horner, M.; Chu, Y. Q.; Kalwinsky, D.; Roberts, W.

[13892] M.: Altered mercaptopurine metabolism, toxic effects, and dosage

[13893] requirement in a thiopurine methyltransferase-deficient child with

[13894] acute lymphocytic leukemia. J. Pediat. 119: 985-989, 1991.

[13895] 949.

[13896] Hon, Y. Y.; Fessing, M. Y.; Pui, C.-H.; Relling, M. V.; Krynetski,

[13897] E. Y.; Evans, W. E.: Polymorphism of the thiopurine S-methyltransferase

[13898] gene in African-Americans. Hum. Molec. Genet. 8: 371-376, 1999.

[13899] 950.

[13900] Honchel, R.; Aksoy, I. A.; Szumlanski, C.; Wood, T. C.; Ot-terness,

[13901] D. M.; Wieben, E. D.; Weinshilboum, R. M.: Human thiop-urine methyltransferase:

[13902] molecular cloning and expression of T84 colon carcinoma cell cDNA. Molec.

[13903] Pharm. 43: 878-887, 1993.

[13904] 951.

[13905] Klemetsdal, B.; Wist, E.; Aarbakke, J.: Gender difference in red

[13906] blood cell thiopurine methyltransferase activity. Scand. J. Clin.

[13907] Lab. Invest. 53: 747-749, 1993.

[13908] 952.

[13909] Krynetski, E. Y.; Schuetz, J. D.; Galpin, A. J.; Pui, C.-H.; Relling,

[13910] M. V.; Evans, W. E.: A single point mutation leading to loss of catalytic

[13911] activity in human thiopurine S-methyltransferase. Proc. Nat. Acad.

[13912] Sci. 92: 949-953, 1995.

[13913] 953.

[13914] Lennard, L.; Lilleyman, J. S.; Van Loon, J.; Weinshilboum, R. M.

[13915] : Genetic variation in response to 6-mercaptopurine for childhood

[13916] acute lymphoblastic leukaemia. Lancet 336: 225-229,

1990.

[13917] 954.

[13918] Lennard, L.; Van Loon, J. A.; Weinshilboum, R. M.: Pharma-
cogenetics

[13919] of acute azathioprine toxicity: relationship to thiopurine
methyltransferase

[13920] genetic polymorphism. Clin. Pharm. Therap. 46: 149-154,
1989.

[13921] 955.

[13922] McLeod, H. L.; Pritchard, S. C.; Githang'a, J.; Indalo, A.;
Ameyaw,

[13923] M.-M.; Powrie, R. H.; Booth, L.; Collie-Duguid, E. S. R.:
Ethnic

[13924] differences in thiopurine methyltransferase pharmacoge-
netics: evidence

[13925] for allele specificity in Caucasian and Kenyan individuals.
Pharmacogenetics 9:

[13926] 773-776, 1999.

[13927] 956.

[13928] Otterness, D.; Szumlanski, C.; Lennard, L. Klemetsdal, B.;
Aarbakke,

[13929] J.; Park-Hah, J. O.; Iven, H.; Schmiegelow, K.; Branum, E.;

O'Brien,

[13930] J.; Weinshilboum, R.: Human thiopurine methyltransferase pharmacogenetics:

[13931] gene sequence polymorphisms. Clin. Pharm. Therap. 62: 60-73, 1997.

[13932] 957.

[13933] Otterness, D. M.; Szumlanski, C. L.; Wood, T. C.; Weinshilboum,

[13934] R. M.: Human thiopurine methyltransferase pharmacogenetics: kindred

[13935] with a terminal exon splice junction mutation that results in loss

[13936] of activity. J. Clin. Invest. 101: 1036-1044, 1998.

[13937] 958.

[13938] Szumlanski, C.; Otterness, D.; Her, C.; Lee, D.; Brandriff, B.;

[13939] Kelsell, D.; Spurr, N.; Lennard, L.; Wieben, E.; Weinshilboum, R.

[13940] : Thiopurine methyltransferase pharmacogenetics: human gene cloning

[13941] and characterization of a common polymorphism. DNA Cell Biol. 15:

[13942] 17-30, 1996.

[13943] 959.

[13944] Tai, H.-L.; Krynetski, E. Y.; Schuetz, E. G.; Yanishevski, Y.;

[13945] Evans, W. E.: Enhanced proteolysis of thiopurine S-
methyltransferase

[13946] (TPMT) encoded by mutant alleles in humans (TPMT*3A,
TPMT*2): mechanisms

[13947] for the genetic polymorphism of TPMT activity. Proc. Nat.
Acad. Sci. 94:

[13948] 6444-6449, 1997.

[13949] 960.

[13950] Tai, H.-L.; Krynetski, E. Y.; Yates, C. R.; Loennechen, T.;
Fessing,

[13951] M. Y.; Krynetskaia, N. F.; Evans, W. E.: Thiopurine S-
methyltransferase

[13952] deficiency: two nucleotide transitions define the most
prevalent mutant

[13953] allele associated with loss of catalytic activity in Cau-
casians. Am.

[13954] J. Hum. Genet. 58: 694-702, 1996.

[13955] 961.

[13956] Weinshilboum, R. M.; Sladek, S. L.: Mercaptopurine pharmacogenetics:

[13957] monogenic inheritance of erythrocyte thiopurine methyltransferase

[13958] activity. Am. J. Hum. Genet. 32: 651-662, 1980.

[13959] 962.

[13960] Heppell-Parton, A.; Cahn, A.; Bench, A.; Lowe, N.; Lehrach, H.;

[13961] Zehetner, G.; Rabbitts, P.: Thioredoxin, a mediator of growth inhibition,

[13962] maps to 9q31. Genomics 26: 379-381, 1995.

[13963] 963.

[13964] Kaghad, M.; Dessarps, F.; Jacquemin-Sablon, H.; Caput, D.; Fradelizi,

[13965] D.; Wollman, E. E.: Genomic cloning of human thioredoxin-encoding

[13966] gene: mapping of the transcription start point and analysis of the

[13967] promoter. Gene 140: 273-278, 1994.

[13968] 964.

[13969] Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Jacobs, P. A.;

Polani,
[13970] P. E.: Segregation of genetic markers in families with chromosome
[13971] polymorphisms and structural rearrangements involving chromosome 1. Ann.
[13972] Hum. Genet. 37: 261-274, 1974.

[13973] 965.

[13974] Herbich, J.; Szilvassy, J.; Schnedl, W.: Gene localisation of

[13975] the PGM-1 enzyme system and the Duffy blood groups on chromosome no.
[13976] 1 by means of a new fragile site at 1p31. Hum. Genet. 70: 178-180,
[13977] 1985.

[13978] 966.

[13979] Bost, K. L.; Smith, E. M.; Blalock, J. E.: Regions of comple-mentarity
[13980] between the messenger RNAs for epidermal growth factor, transferrin,
[13981] interleukin-2 and their respective receptors. Biochem. Biophys. Res.
[13982] Commun. 128: 1373-1380, 1985.

[13983] 967.

[13984] Bowman, B. H.; Yang, F.; Adrian, G. S.: Transferrin: evolu-
tion
[13985] and genetic regulation of expression. Adv. Genet. 25:
1-38, 1988.

[13986] 968.

[13987] Dayhoff, M. O.: Atlas of Protein Sequence and Structure.
Transferrin.
[13988] Washington: National Biomedical Research Foundation
(pub.) 5: 1972.

[13989] Pp. D310 only.

[13990] 969.

[13991] Evans, R. W.: Personal Communication. London, England
5/29/1987.

[13992] 970.

[13993] Evans, R. W.; Williams, J.; Moreton, K.: A variant of human
transferrin
[13994] with abnormal properties. Biochem. J. 201: 19-26, 1982.

[13995] 971.

[13996] Giari, A.; Weidinger, S.; Domenici, R.; Bargagna, M.: Trans-
ferrin

[13997] variants in Tuscany (Italy): evidence for two 'new' Tf alleles. Hum.
[13998] Genet. 69: 284-286, 1985.

[13999] 972.

[14000] Goya, N.; Miyazaki, S.; Kodate, S.; Ushio, B.: A family of congenital
[14001] atransferrinemia. Blood 40: 239-245, 1972.

[14002] 973.

[14003] Hayashi, A.; Wada, Y.; Suzuki, T.; Shimizu, A.: Studies on familial
[14004] hypotransferrinemia: unique clinical course and molecular pathology. Am.
[14005] J. Hum. Genet. 53: 201-213, 1993.

[14006] 974.

[14007] Huerre, C.; Uzan, G.; Grzeschik, K. H.; Weil, D.; Levin, M.; Hors-Cayla,
[14008] M.-C.; Boue, J.; Kahn, A.; Junien, C.: The structural gene for transferrin
[14009] (TF) maps to 3q21-3qter. Ann. Genet. 27: 5-10, 1984.

[14010] 975.

[14011] Jenkins, T.; Bothwell, T. H.; Maier, G.; Laidler, A.: Is transferrin

[14012] normal in idiopathic haemochromatosis? Brit. J. Haemat. 52: 493-495,

[14013] 1982.

[14014] 976.

[14015] Kamboh, M. I.; Ferrell, R. E.: Human transferrin polymorphism. Hum.

[14016] Hered. 37: 65-81, 1987.

[14017] 977.

[14018] King, J.; Robson, E. B.; Edwards, V. H.; Cook, P. J. L.; Buckton,

[14019] K. E.: Miscellaneous contributions to the mapping of chromosome 1.

[14020] (Abstract) Cytogenet. Cell Genet. 25: 172 only, 1979.

[14021] 978.

[14022] Kueppers, F.; Harpel, B. M.: Transferrin C subtypes in US blacks

[14023] and whites. Hum. Hered. 30: 376-382, 1980.

[14024] 979.

[14025] Lee, P. L.; Halloran, C.; Trevino, R.; Felitti, V.; Beutler, E.

[14026] : Human transferrin G277S mutation: a risk factor for iron deficiency

[14027] anaemia. Brit. J. Haemat. 115: 329-333, 2001.

[14028] 980.

[14029] MacGillivray, R. T. A.; Mendez, E.; Sinha, S. K.; Sutton, M. R.;

[14030] Lineback-Zins, J.; Brew, K.: The complete amino acid se-quence of

[14031] human serum transferrin. Proc. Nat. Acad. Sci. 79: 2504-2508, 1982.

[14032] 981.

[14033] McAlpine, P. J.; Allderdice, P. W.; Cox, D. W.; Simpson, N. E.;

[14034] McEachran, M.; Komarnicki, L.: The ordering of TF:CHE1:AHSG and their

[14035] orientation distal to 3q21. (Abstract) Cytogenet. Cell Genet. 46:

[14036] 659 only, 1987.

[14037] 982.

[14038] Namekata, K.; Oyama, F.; Imagawa, M.; Ihara, Y.: Human

transferrin

[14039] (Tf): a single mutation at codon 570 determines Tf C1 or Tf C2 variant. Hum.

[14040] Genet. 100: 457-458, 1997.

[14041] 983.

[14042] Naylor, S. L.; Lalley, P. A.; Elliott, R. W.; Brown, J. A.; Shows,

[14043] T. B.: Evidence for homologous regions of human chromosome 3 and

[14044] mouse chromosome 9 predicts location of human genes. (Abstract) Am.

[14045] J. Hum. Genet. 32: 158A only, 1980.

[14046] 984.

[14047] Pang, H.; Koda, Y.; Soejima, M.; Kimura, H.: Identification of

[14048] a mutation (A1879G) of transferrin from cDNA prepared from peripheral

[14049] blood cells. Ann. Hum. Genet. 62: 271-274, 1998.

[14050] 985.

[14051] Park, I.; Schaeffer, E.; Sidoli, A.; Baralle, F. E.; Cohen, G.

[14052] N.; Zakin, M. M.: Organization of the human transferrin

gene: direct
[14053] evidence that it originated by gene duplication. Proc. Nat. Acad.
[14054] Sci. 82: 3149-3153, 1985.

[14055] 986.

[14056] Parker, W. C.; Bearn, A. G.: Additional genetic variation of

[14057] human serum transferrin. Science 137: 854-856, 1962.

[14058] 987.

[14059] Robson, E. B.; Sutherland, I.; Harris, H.: Evidence for link-age
[14060] between the transferrin locus (TF) and the serum cholinesterase locus
[14061] E(1) in man. Ann. Hum. Genet. 29: 325-336, 1966.

[14062] 988.

[14063] Sass-Kuhn, S. P.; Moqbel, R.; Mackay, J. A.; Cromwell, O.; Kay,
[14064] A. B.: Human granulocyte/pollen-binding protein: recog-nition and
[14065] identification as transferrin. J. Clin. Invest. 73: 202-210, 1984.
[14066] 989.

[14067] Smithies, O.: Variants in human serum B-globulins. Nature 180:

[14068] 1482 only, 1957.

[14069] 990.

[14070] Smithies, O.: Third allele at the serum beta-globulin locus in

[14071] humans. Nature 181: 1203-1204, 1958.

[14072] 991.

[14073] Sutherland, R.; Delia, D.; Schneider, C.; Newman, R.; Kemshead,

[14074] J.; Greaves, M.: Ubiquitous cell-surface glycoprotein on tumor cells

[14075] is proliferation-associated receptor for transferrin. Proc. Nat.

[14076] Acad. Sci. 78: 4515-4519, 1981.

[14077] 992.

[14078] Uzan, G.; Frain, M.; Park, I.; Besmond, C.; Maessen, G.; Trepat,

[14079] J. S.; Zakin, M. M.; Kahn, A.: Molecular cloning and se-quence analysis

[14080] of cDNA for human transferrin. Biochem. Biophys. Res.

Commun. 119:
[14081] 273-281, 1984.

[14082] 993.

[14083] Wang, A.-C.; Sutton, H. E.: Human transferrins C and D(1): chemical
[14084] difference in a peptide. Science 149: 435-437, 1965.

[14085] 994.

[14086] Wang, A.-C.; Sutton, H. E.; Howard, P. N.: Human transfer-rins
[14087] C and D(Chi): an amino-acid difference. Biochem. Genet. 1: 55-60,
[14088] 1967.

[14089] 995.

[14090] Wang, A.-C.; Sutton, H. E.; Riggs, A.: A chemical differ-ence
[14091] between human transferrins B2 and C. Am. J. Hum. Genet. 18: 454-458,
[14092] 1966.

[14093] 996.

[14094] Welch, S.; Langmead, L.: A comparison of the structure

and properties

[14095] of normal human transferrin and a genetic variant of human transferrin. Int.

[14096] J. Biochem. 22: 275-282, 1990.

[14097] 997.

[14098] Yang, F.; Lum, J. B.; McGill, J. R.; Moore, C. M.; Naylor, S.

[14099] L.; van Bragt, P. H.; Baldwin, W. D.; Bowman, B. H.: Human transferrin:

[14100] cDNA characterization and chromosomal localization. Proc. Nat. Acad.

[14101] Sci. 81: 2752-2756, 1984.

[14102] 998.

[14103] Huebner, K.; Druck, T.; LaForgia, S.; Lasota, J.; Croce, C. M.;

[14104] Lanfrancone, L.; Donti, E.; Pengue, G.; La Mantia, G.; Pelicci, P.-G.;

[14105] Lania, L.: Chromosomal localization of four human zinc finger cDNAs. Hum.

[14106] Genet. 91: 217-222, 1993.

[14107] 999.

[14108] Batlle, E.; Sancho, E.; Franci, C.; Dominguez, D.; Monfar,

M.;

[14109] Baulida, J.; Garcia de Herreros, A.: The transition factor Snail

[14110] is a repressor of E-cadherin gene expression in epithelial tumour

[14111] cells. Nature Cell Biol. 2: 84-89, 2000.

[14112] 1000.

[14113] Becker, K.-F.; Atkinson, M. J.; Reich, U.; Becker, I.; Nekarda,

[14114] H.; Siewert, J. R.; Hoefler, H.: E-Cadherin gene mutations provide

[14115] clues to diffuse type gastric carcinomas. Cancer Res. 54: 3845-3852,

[14116] 1994.

[14117] 1001.

[14118] Erbe, R. W.: Cabot case. New Eng. J. Med. 298: 95-101, 1978.

[14119] 1002.

[14120] Feldman, D. E.; Thulasiraman, V.; Ferreyra, R. G.; Frydman, J.

[14121] : Formation of the VHL-elongin BC tumor suppressor

complex is mediated
[14122] by the chaperonin TRiC. Molec. Cell 1051-1061, 1999.

[14123] 1003.

[14124] Fill, W. L.; Lamiell, J. M.; Polk, N. O.: The radiographic manifestations
[14125] of von Hippel-Lindau disease. Radiology 133: 289-295, 1979.

[14126] 1004.

[14127] Fishman, R. S.; Bartholomew, L. G.: Severe pancreatic in-volvement
[14128] in three generations in von Hippel-Lindau disease. Mayo Clin. Proc. 54:
[14129] 329-331, 1979.

[14130] 1005.

[14131] Friedrich, C. A.: Genotype/phenotype correlation in von Hippel-Lindau
[14132] syndrome. Hum. Molec. Genet. 10: 763-767, 2001.

[14133] 1006.

[14134] Fukino, K.; Teramoto, A.; Adachi, K.; Takahashi, H.; Emi, M.:
[14135] A family with hydrocephalus as a complication of cerebel-

lar hemangioblastoma:
[14136] identification of pro157leu mutation in the VHL gene. J. Hum. Genet. 45:
[14137] 47-51, 2000.

[14138] 1007.

[14139] Fulton, J. F.: Harvey Cushing: A Biography. Springfield, Illinois:
[14140] Charles C Thomas (pub.) 1946.

[14141] 1008.

[14142] Funk, K. C.; Heiken, J. P.: Papillary cystadenoma of the broad
[14143] ligament in a patient with von Hippel-Lindau disease. Am. J. Radiol. 153:
[14144] 527-528, 1989.

[14145] 1009.

[14146] Gaffey, M. J.; Mills, S. E.; Boyd, J. C.: Aggressive papillary
[14147] tumor of middle ear/temporal bone and adnexal papillary cystadenoma. Am.
[14148] J. Surg. Path. 18: 1254-1260, 1994.

[14149] 1010.

[14150] Gallou, C.; Joly, D.; Mejean, A.; Staroz, F.; Martin, N.; Tarlet,
[14151] G.; Orfanelli, M. T.; Bouvier, R.; Droz, D.; Chretien, Y.; Marechal,
[14152] J. M.; Richard, S.; Junien, C.; Beroud, C.: Mutations of the VHL
[14153] gene in sporadic renal cell carcinoma: definition of a risk factor
[14154] for VHL patients to develop an RCC. Hum. Mutat. 13: 464-475, 1999.
[14155] 1011.

[14156] Garcia, A.; Matias-Guiu, X.; Cabezas, R.; Chico, A.; Prat, J.;
[14157] Baiget, M.; De Leiva, A.: Molecular diagnosis of von Hippel-Lindau
[14158] disease in a kindred with a predominance of familial phaeochromocytoma. Clin.
[14159] Endocr. 46: 359-363, 1997.

[14160] 1012.

[14161] Gemmill, R. M.; Bemis, L. T.; Lee, J. P.; Sozen, M. A.; Baron,
[14162] A.; Zeng, C.; Erickson, P. F.; Hooper, J. E.; Drabkin, H. A.: The

[14163] TRC8 hereditary kidney cancer gene suppresses growth and functions

[14164] with VHL in a common pathway. Oncogene 21: 3507-3516, 2002.

[14165] 1013.

[14166] Gersell, D. J.; King, T. C.: Papillary cystadenoma of the mesosalpinx

[14167] in von Hippel-Lindau disease. Am. J. Surg. Path. 12: 145-149, 1988.

[14168] 1014.

[14169] Gilcrease, M. Z.; Schmidt, L.; Zbar, B.; Truong, L.; Rut-ledge,

[14170] M.; Wheeler, T. M.: Somatic von Hippel-Lindau mutation in clear cell

[14171] papillary cystadenoma of the epididymis. Hum. Path. 26: 1341-1346,

[14172] 1995.

[14173] 1015.

[14174] Glenn, G. M.; Daniel, L. N.; Choyke, P.; Linehan, W. M.; Oldfield,

[14175] E.; Gorin, M. B.; Hosoe, S.; Latif, F.; Weiss, G.; Walther, M.; Lerman,

[14176] M. I.; Zbar, B.: Von Hippel-Lindau (VHL) disease: distinct phenotypes

[14177] suggest more than one mutant allele at the VHL locus. Hum. Genet. 87:

[14178] 207-210, 1991.

[14179] 1016.

[14180] Glenn, G. M.; Linehan, W. M.; Hosoe, S.; Latif, F.; Yao, M.; Choyke,

[14181] P.; Gorin, M. B.; Chew, E.; Oldfield, E.; Manolatos, C.; Or-

[14182] cutt, M. L.; Walther, M. M.; Weiss, G. H.; Tory, K.; Jensson, O.; Ler-

[14183] man, M. I.; Zbar, B.: Screening for von Hippel-Lindau disease by DNA polymorphism

[14184] analysis. J.A.M.A. 267: 1226-1231, 1992.

[14185] 1017.

[14186] Gnarra, J. R.; Ward, J. M.; Porter, F. D.; Wagner, J. R.; De-

[14187] vor, D. E.; Grinberg, A.; Emmert-Buck, M. R.; Westphal, H.; Klausner, R.

[14188] D.; Marston Linehan, W.: Defective placental vasculogene-

[14189] sis causes

[14189] embryonic lethality in VHL-deficient mice. Proc. Nat. Acad. Sci. 94:
[14190] 9102-9107, 1997.

[14191] 1018.

[14192] Go, R. C. P.; Lamiell, J. M.; Hsia, Y. E.; Yuen, J. W.-M.; Paik,
[14193] Y.: Segregation and linkage analyses of von Hippel Lindau disease
[14194] among 220 descendants from one kindred. Am. J. Hum. Genet. 36: 131-142,
[14195] 1984.

[14196] 1019.

[14197] Goldberg, M. F.; Duke, J. R.: Von Hippel-Lindau disease: histopathologic
[14198] findings in a treated and untreated eye. Am. J. Ophthal. 66: 693-705,
[14199] 1968.

[14200] 1020.

[14201] Graff, J. W.: Personal Communication. Brookline, Mass. 10/4/1998.
[14202] 1021.

[14203] Green, J. S.; Bowmer, M. I.; Johnson, G. J.: Von Hippel-Lindau

[14204] disease in a Newfoundland kindred. Canad. Med. Assoc. J. 134: 133-138

[14205] and 146, 1986.

[14206] 1022.

[14207] Gross, D. J.; Avishai, N.; Meiner, V.; Filon, D.; Zbar, B.; Abeliovich,

[14208] D.: Familial pheochromocytoma associated with a novel mutation in

[14209] the von Hippel-Lindau gene. J. Clin. Endocr. Metab. 81: 147-149,

[14210] 1996.

[14211] 1023.

[14212] Grossman, M.; Melmon, K. L.: Von Hippel-Lindau disease.In: Vinken,

[14213] P. J.; Bruyn, G. W.: Handbook of Clinical Neurology. The Phakomatoses.

[14214] Amsterdam: North Holland (pub.) 14: 1972. Pp. 241-259.

[14215] 1024.

[14216] Haase, V. H.; Glickman, J. N.; Socolovsky, M.; Jaenisch, R.:

[14217] Vascular tumors in livers with targeted inactivation of the von Hippel-Lindau
[14218] tumor suppressor. Proc. Nat. Acad. Sci. 98: 1583-1588, 2001.
[14219] 1025.

[14220] Hagler, W. S.; Hyman, B. N.; Waters, W. C., III: Von Hippel's

[14221] angiomatosis retinae and pheochromocytoma. Trans. Am. Acad. Ophthal.
[14222] Otolaryng. 75: 1022-1034, 1971.

[14223] 1026.

[14224] Hennessy, T. G.; Stern, W. E.; Herrick, S. E.: Cerebellar he-mangioblastoma:
[14225] erythropoietic activity by radioion assay. J. Nucl. Med. 8: 601-606,
[14226] 1967.

[14227] 1027.

[14228] Herman, J. G.; Latif, F.; Weng, Y.; Lerman, M. I.; Zbar, B.; Liu,
[14229] S.; Samid, D.; Duan, D.-S. R.; Guarra, J. R.; Linehan, W. M.; Baylin,
[14230] S. B.: Silencing of the VHL tumor-suppressor gene by DNA

methylation
[14231] in renal carcinomas. Proc. Nat. Acad. Sci. 91: 9700-9704, 1994.
[14232] 1028.
[14233] Hes, F.; Zewald, R.; Peeters, T.; Sijmons, R.; Links, T.; Ver-heij,
[14234] J.; Matthijs, G.; Legius, E.; Mortier, G.; van der Torren, K.; Rosman,
[14235] M.; Lips, C.; Pearson, P.; van der Luijt, R.: Genotype-phe-notype
[14236] correlations in families with deletions in the von Hippel-Lindau (VHL)
[14237] gene. Hum. Genet. 106: 425-431, 2000.
[14238] 1029.
[14239] Hes, F. J.; McKee, S.; Taphoorn, M. J. B.; Rehal, P.; van der
[14240] Luijt, R. B.; McMahon, R.; van der Smagt, J. J.; Dow, D.; Ze-wald,
[14241] R. A.; Whittaker, J.; Lips, C. J. M.; MacDonald, F.; Pearson, P. L.;
[14242] Maher, E. R.: Cryptic von Hippel-Lindau disease: germline mutations
[14243] in patients with haemangioblastoma only. J. Med. Genet.

37: 939-943,

[14244] 2000.

[14245] 1030.

[14246] Hoffman, M. A.; Ohh, M.; Yang, H. Kico, J. M.; Ivan, M.; Kaelin,

[14247] W. G., Jr.: von Hippel-Lindau, protein mutants linked to type 2C

[14248] VHL disease preserve the ability to downregulated HIF. Hum. Molec.

[14249] Genet. 10: 1019-1027, 2001.

[14250] 1031.

[14251] Whitmore, S. A.; Apostolou, S.; Lane, S.; Nancarrow, J. K.; Phillips,

[14252] H. A.; Richards, R. I.; Sutherland, G. R.; Callen, D. F.: Isola-tion

[14253] and characterization of transcribed sequences from a chromosome 16

[14254] hn-cDNA library and the physical mapping of genes and transcribed

[14255] sequences using a high-resolution somatic cell panel of human chromosome

[14256] 16. Genomics 20: 169-175, 1994.

[14257] 1032.

[14258] Bashan, N.; Iancu, T. C.; Lerner, A.; Fraser, D.; Potashnik, R.;

[14259] Moses, S. W.: Glycogenosis due to liver and muscle phosphorylase

[14260] kinase deficiency. Pediat. Res. 15: 299-303, 1981.

[14261] 1033.

[14262] Burwinkel, B.; Maichele, A. J.; Aagenaes, O.; Bakker, H. D.; Lerner,

[14263] A.; Shin, Y. S.; Strachan, J. A.; Kilimann, M. W.: Autosomal glycogenosis

[14264] of liver and muscle due to phosphorylase kinase deficiency is caused

[14265] by mutations in the phosphorylase kinase beta subunit (PHKB). Hum.

[14266] Molec. Genet. 6: 1109-1115, 1997.

[14267] 1034.

[14268] Burwinkel, B.; Moses, S. W.; Kilimann, M. W.: Phosphorylase-kinase-deficient

[14269] liver glycogenosis with an unusual biochemical phenotype in blood

[14270] cells associated with a missense mutation in the beta subunit gene

[14271] (PHKB). Hum. Genet. 101: 170-174, 1997.

[14272] 1035.

[14273] Francke, U.; Darras, B. T.; Zander, N. F.; Kilimann, M. W.: Assignment

[14274] of human genes for phosphorylase kinase subunits alpha (PHKA) to Xq12-q13

[14275] and beta (PHKB) to 16q12-q13. Am. J. Hum. Genet. 45: 276-282, 1989.

[14276] 1036.

[14277] Kilimann, M. W.; Zander, N. F.; Kuhn, C. C.; Crabb, J. W.; Meyer,

[14278] H. E.; Heilmeyer, L. M. G., Jr.: The alpha and beta subunits of phosphorylase

[14279] kinase are homologous: cDNA cloning and primary structure of the beta

[14280] subunit. Proc. Nat. Acad. Sci. 85: 9381-9385, 1988.

[14281] 1037.

[14282] Lederer, B.; Van Hoof, F.; Van den Berghe, G.; Hers, H. G.: Glycogen
[14283] phosphorylase and its converter enzymes in haemolysates of normal
[14284] human subjects and of patients with type VI glycogen storage disease:
[14285] a study of phosphorylase kinase deficiency. Biochem. J. 147: 23-35,
[14286] 1975.

[14287] 1038.

[14288] Wullrich-Schmoll, A.; Kilimann, M. W.: Structure of the hu-man
[14289] gene encoding the phosphorylase kinase beta subunit (PHKB). Europ.
[14290] J. Biochem. 238: 374-380, 1996.

[14291] 1039.

[14292] Zander, N. F.; Meyer, H. E.; Hoffmann-Posorske, E.; Crabb, J. W.;
[14293] Heilmeyer, L. M. G., Jr.; Kilimann, M. W.: cDNA cloning and complete
[14294] primary structure of skeletal muscle phosphorylase kinase (alpha subunit). Proc.

[14295] Nat. Acad. Sci. 85: 2929-2933, 1988.

[14296] 1040.

[14297] Rocchi, M.; Archidiacono, N.; Romeo, G.; Saginati, M.; Zardi, L.

[14298] : Assignment of the gene for human tenascin to the region q32-q34

[14299] of chromosome 9. Hum. Genet. 86: 621-623, 1991.

[14300] 1041.

[14301] Steindler, D. A.; Settles, D.; Erickson, H. P.; Laywell, E. D.;

[14302] Yoshiki, A.; Faissner, A.; Kusakabe, M.: Tenascin knockout mice:

[14303] barrels, boundary molecules, and glial scars. J. Neurosci. 15: 1971-1983,

[14304] 1995.

[14305] 1042.

[14306] Bamford, R. N.; Roessler, E.; Burdine, R. D.; Saplakoglu, U.; dela

[14307] Cruz, J.; Splitt, M.; Towbin, J.; Bowers, P.; Marino, B.; Schier,

[14308] A. F.; Shen, M. M.; Muenke, M.; Casey, B.: Loss-of-function mutations

[14309] in the EGF-CFC gene CFC1 are associated with human left-right laterality

[14310] defects. Nature Genet. 26: 365-369, 2000.

[14311] 1043.

[14312] Ciccodicola, A.; Dono, R.; Obici, S.; Simeone, A.; Zollo, M.; Persico,

[14313] M. G.: Molecular characterization of a gene of the 'EGF family' expressed

[14314] in undifferentiated human NTERA2 teratocarcinoma cells. EMBO J. 8:

[14315] 1987-1991, 1989.

[14316] 1044.

[14317] de la Cruz, J. M.; Bamford, R. N.; Burdine, R. D.; Roessler, E.;

[14318] Barkovich, A. J.; Donnai, D.; Schier, A. F.; Muenke, M.: A loss-of-function

[14319] mutation in the CFC domain of TDGF1 is associated with human forebrain

[14320] defects. Hum. Genet. 110: 422-428, 2002.

[14321] 1045.

[14322] Dono, R.; Montuori, N.; Rocchi, M.; De Ponti-Zilli, L.; Cic-

codicola,

[14323] A.; Persico, M. G.: Isolation and characterization of the CRIPTO

[14324] autosomal gene and its X-linked related sequence. Am. J. Hum. Genet. 49:

[14325] 555-565, 1991.

[14326] 1046.

[14327] Liguori, G.; De Gregorio, L.; Tucci, M.; Lago, C. T.; Barra, A.;

[14328] Dragani, T. A.; Persico, M.: Mapping of the mouse Tdgf1 gene and

[14329] Tdgf pseudogenes. Mammalian Genome 8: 502-505, 1997.

[14330] 1047.

[14331] Liguori, G.; Tucci, M.; Montuori, N.; Dono, R.; Lago, C. T.; Pacifico,

[14332] F.; Armenante, F.; Persico, M. G.: Characterization of the mouse

[14333] Tdgf1 gene and Tdgf pseudogenes. Mammalian Genome 7: 344-348, 1996.

[14334] 1048.

[14335] Saccone, S.; Rapisarda, A.; Motta, S.; Dono, R.; Persico, G.

M.;

[14336] Della Valle, G.: Regional localization of the human EGF-like growth

[14337] factor CRIPTO gene (TDGF-1) to chromosome 3p21. Hum. Genet. 95:

[14338] 229-230, 1995.

[14339] 1049.

[14340] Shen, M. M.; Schier, A. F.: The EGF-CFC gene family in vertebrate

[14341] development. Trends Genet. 16: 303-309, 2000.

[14342] 1050.

[14343] Chang, L. M. S.; Bollum, F. J.: Molecular biology of terminal

[14344] transferase. CRC Crit. Rev. Biochem. 21: 27-52, 1986.

[14345] 1051.

[14346] Epplen, J. T.; Chluba, J.; Hardt, C.; Hinkkanen, A.; Steimle, V.;

[14347] Stockinger, H.: Mammalian T-lymphocyte antigen receptor genes: genetic

[14348] and nongenetic potential to generate variability. Hum. Genet. 75:

[14349] 300-310, 1987.

[14350] 1052.

[14351] Isobe, M.; Huebner, K.; Erikson, J.; Peterson, R. C.; Bollum, F.

[14352] J.; Chang, L. M. S.; Croce, C. M.: Chromosome localization of the

[14353] gene for human terminal deoxynucleotidyltransferase to region 10q23-q25. Proc.

[14354] Nat. Acad. Sci. 82: 5836-5840, 1985.

[14355] 1053.

[14356] Landau, N. R.; St. John, T. P.; Weissman, I. L.; Wolf, S. C.; Silverstone,

[14357] A. E.; Baltimore, D.: Cloning of terminal transferase cDNA by antibody

[14358] screening. Proc. Nat. Acad. Sci. 81: 5836-5840, 1984.

[14359] 1054.

[14360] Riley, L. K.; Morrow, J. K.; Danton, M. J.; Coleman, M. S.: Human

[14361] terminal deoxyribonucleotidyltransferase: molecular cloning and structural

[14362] analysis of the gene and 5-prime flanking region. Proc. Nat. Acad.

[14363] Sci. 85: 2489-2493, 1988.

[14364] 1055.

[14365] Thai, T.-H.; Purugganan, M. M.; Roth, D. B.; Kearney, J. F.: Distinct

[14366] and opposite diversifying activities of terminal transferase splice

[14367] variants. Nature Immun. 3: 457-462, 2002.

[14368] 1056.

[14369] Yang-Feng, T. L.; Landau, N. R.; Baltimore, D.; Francke, U.: The

[14370] terminal deoxynucleotidyltransferase gene is located on human chromosome

[14371] 10 (10q23-q24) and on mouse chromosome 19. Cyto-genet. Cell Genet. 43:

[14372] 121-126, 1986.

[14373] 1057.

[14374] Kasahara, M.; Figueroa, F.; Klein, J.: Random cloning of genes

[14375] from mouse chromosome 17. Proc. Nat. Acad. Sci. 84: 3325-3328, 1987.

[14376] 1058.

[14377] Kasahara, M.; Gutknecht, J.; Brew, K.; Spurr, N.; Goodfellow, P.

[14378] N.: Cloning and mapping of a testis-specific gene with sequence similarity

[14379] to a sperm-coating glycoprotein gene. Genomics 5: 527-534, 1989.

[14380] 1059.

[14381] Kasahara, M.; Passmore, H. C.; Klein, J.: A testis-specific gene

[14382] Tpx-1 maps between Pgk-2 and Mep-1 on mouse chromosome 17. Immunogenetics 29:

[14383] 61-63, 1989.

[14384] 1060.

[14385] Gonzatti-Haces, M.; Seth, A.; Park, M.; Copeland, T.; Oroszlan,

[14386] S.; Vande Woude, G. F.: Characterization of the TPR-MET oncogene

[14387] p65 and the MET protooncogene p140 protein-tyrosine kinases. Proc.

[14388] Nat. Acad. Sci. 85: 21-25, 1988.

[14389] 1061.

[14390] Miranda, C.; Minoletti, F.; Greco, A.; Sozzi, G.; Pierotti, M.

[14391] A.: Refined localization of the human TPR gene to chro-
mosome 1q25

[14392] by in situ hybridization. Genomics 23: 714-715, 1994.

[14393] 1062.

[14394] Cao, Z.; Umek, R. M.; McKnight, S. L.: Regulated expres-
sion of

[14395] three C/EBP isoforms during adipose conversion of
3T3-L1 cells. Genes

[14396] Dev. 5: 1538-1552, 1991.

[14397] 1063.

[14398] Descombes, P.; Chojkier, M.; Lichtsteiner, S.; Falvey, E.;
Schibler,

[14399] U.: LAP, a novel member of the C/EBP gene family, en-
codes a liver-enriched

[14400] transcriptional activator protein. Genes Dev. 4:
1541-1551, 1990.

[14401] 1064.

[14402] Hendricks-Taylor, L. R.; Bachinski, L. L.; Siciliano, M. J.;
Fertitta,

[14403] A.; Trask, B.; de Jong, P. J.; Ledbetter, D. H.; Darlington, G.

J.

[14404] : The CCAAT/enhancer binding protein (C/EBP-alpha) gene (CEBPA) maps

[14405] to human chromosome 19q13.1 and the related nuclear factor NF-IL6

[14406] (C/EBP-beta) gene (CEBPB) maps to human chromosome 20q13.1. Genomics 14:

[14407] 12-17, 1992.

[14408] 1065.

[14409] Jenkins, N. A.; Gilbert, D. J.; Cho, B. C.; Strobel, M. C.; Williams,

[14410] S. C.; Copeland, N. G.; Johnson, P. F.: Mouse chromosomal location

[14411] of the CCAAT/enhancer binding proteins C/EBP-beta (Cebpb), C/EBP-delta

[14412] (Cebpd), and CRP1 (Cebpe). Genomics 28: 333-336, 1995.

[14413] 1066.

[14414] Chang, M.-L.; Artymiuk, P. J.; Wu, X.; Hollan, S.; Lammi, A.;

[14415] Maquat, L. E.: Human triosephosphate isomerase deficiency resulting

[14416] from mutation of phe-240. Am. J. Hum. Genet. 52:

1260-1269, 1993.

[14417] 1067.

[14418] Burchell, B.: Personal Communication. Dundee, Scotland 12/1991.

[14419] 1068.

[14420] Young, S. P.; Bomford, A.; Madden, A. D.; Garratt, R. C.; Williams,

[14421] R.; Evans, R. W.: Abnormal in vitro function of a variant human transferrin. Brit.

[14422] J. Haemat. 56: 581-587, 1984.

[14423] 1069.

[14424] Burchell, B.; Nebert, D. W.; Nelson, D. R.; Bock, K. W.; Iyanagi,

[14425] T.; Jansen, P. L. M.; Lancet, D.; Mulder, G. J.; Roy Chowd-hury, J.;

[14426] Siest, G.; Tephly, T. R.; Mackenzie, P. I.: The UDP glu-curonosyltransferase

[14427] gene superfamily: suggested nomenclature based on evo-lutionary divergence. DNA

[14428] Cell Biol. 10: 487-494, 1991.

[14429] 1070.

[14430] Erps, L. T.; Ritter, J. K.; Hersh, J. H.; Blossom, D.; Martin,

[14431] N. C.; Owens, I. S.: Identification of two single base substitutions

[14432] in the UGT1 gene locus which abolish bilirubin uridine diphosphate

[14433] glucuronosyltransferase activity in vitro. J. Clin. Invest. 93:

[14434] 564-570, 1994.

[14435] 1071.

[14436] Findlay, K. A. B.; Kaptein, E.; Visser, T. J.; Burchell, B.:

[14437] Characterization of the uridine diphosphate-glucuronosyltransferase-catalyzing

[14438] thyroid hormone glucuronidation in man. J. Clin. Endocr. Metab. 85:

[14439] 2879-2883, 2000.

[14440] 1072.

[14441] Francoual, J.; Rivierre, A.; Mokrani, C.; Khrouf, N.; Gottrand,

[14442] F.; Myara, A.; Le Bihan, B.; Capel, L.; Lindenbaum, A.; Labrune, P.

[14443] : Crigler-Najjar syndrome type I in Tunisia may be associated with

[14444] a founder effect related to the Q357R mutation within the UGT1 gene. Hum.

[14445] Mutat. 19: 570-571, 2002.

[14446] 1073.

[14447] Galanello, R.; Perseu, L.; Melis, M. A.; Cipollina, L.; Barella,

[14448] S.; Giagu, N.; Turco, M. P.; Maccioni, O.; Cao, A.: Hyper-bilirubinaemia

[14449] in heterozygous beta-thalassaemia is related to co-inherited Gilbert's

[14450] syndrome. Brit. J. Haemat. 99: 433-436, 1997.

[14451] 1074.

[14452] Gantla, S.; Bakker, C. T. M.; Deocharan, B.; Thummala, N. R.;

[14453] Zweiner, J.; Sinaasappel, M.; Roy Chowdhury, J.; Bosma, P. J.; Roy

[14454] Chowdhury, N.: Splice-site mutations: a novel genetic mechanism of

[14455] Crigler-Najjar syndrome type 1. Am. J. Hum. Genet. 62: 585-592,

[14456] 1998.

[14457] 1075.

[14458] Gollan, J. L.; Huang, S. N.; Billing, B.; Sherlock, S.: Prolonged
[14459] survival in three brothers with severe type 2 Crigler-Najjar syndrome:
[14460] ultrastructural and metabolic studies. Gastroenterology 68: 1543-1555,
[14461] 1975.

[14462] 1076.

[14463] Gong, Q.-H.; Cho, J. W.; Huang, T.; Potter, C.; Gholami, N.; Basu,
[14464] N. K.; Kubota, S.; Carvalho, S.; Pennington, M. W.; Owens, I. S.;
[14465] Popescu, N. C.: Thirteen UDP-glucuronosyltransferase genes are encoded
[14466] at the human UGT1 gene complex locus. Pharmacogenetics 11: 357-368,
[14467] 2001.

[14468] 1077.

[14469] Hagiwara, H.; Takeda, K.; Ikeda, H.; Nakai, H.; Burchell, B.:

[14470] Gene mapping of human bilirubin UDP-glucuronosyl transferase on 1q21-q23

[14471] by a cell sorter and in situ hybridization. Jpn. J. Hum. Genet. 36:
[14472] 189-194, 1991.

[14473] 1078.

[14474] Harding, D.; Fournel-Gigleux, S.; Jackson, M. R.; Burchell, B.
[14475] : Cloning and substrate specificity of a human phenol UDP-glucuronosyltransferase
[14476] expressed in COS-7 cells. Proc. Nat. Acad. Sci. 85: 8381-8385, 1988.
[14477] 1079.

[14478] Harding, D.; Jeremiah, S. J.; Povey, S.; Burchell, B.: Chromosomal
[14479] mapping of a human phenol UDP-glucuronosyltransferase, GNT1. Ann.
[14480] Hum. Genet. 54: 17-21, 1990.

[14481] 1080.

[14482] Harding, D.; Jeremiah, S. J.; Povey, S.; Burchell, B.: Phenol

[14483] UDP-glucuronosyltransferase is coded by a gene on human chromosome
[14484] 2. (Abstract) Cytogenet. Cell Genet. 51: 1011, 1989.

[14485] 1081.

[14486] Iolascon, A.; Faienza, M. F.; Moretti, A.; Perrotta, S.; Miraglia

[14487] del Giudice, E.: UGT1 promoter polymorphism accounts for increased

[14488] neonatal appearance of hereditary spherocytosis.(Letter) Blood 91:

[14489] 1093 only, 1998.

[14490] 1082.

[14491] Iyer, L.; King, C. D.; Whitington, P. F.; Green, M. D.; Roy, S.

[14492] K.; Tephly, T. R.; Coffman, B. L.; Ratain, M. J.: Genetic predisposition

[14493] to the metabolism of irinotecan (CPT-11): role of uridine diphosphate

[14494] glucuronosyltransferase isoform 1A1 in the glucuronidation of its

[14495] active metabolite (SN-38) in human liver microsomes. J. Clin. Invest. 101:

[14496] 847-854, 1998.

[14497] 1083.

[14498] Jackson, M. R.; McCarthy, L. R.; Harding, D.; Wilson, S.;

Coughtrie,
[14499] M. W. H.; Burchell, B.: Cloning of a human liver microsomal UDP-glucuronosyltransferase
[14500] cDNA. Biochem. J. 242: 581-588, 1987.

[14501] 1084.

[14502] Jansen, P. L. M.; Mulder, G. J.; Burchell, B.; Bock, K. W.: New
[14503] developments in glucuronidation research: report of a workshop on
[14504] 'glucuronidation, its role in health and disease.'. Hepatology 15:
[14505] 532-544, 1992.

[14506] 1085.

[14507] Kadakol, A.; Ghosh, S. S.; Sappal, B. S.; Sharma, G.; Roy Chowdhury,
[14508] J.; Roy Chowdhury, N.: Genetic lesions of bilirubin uridine-diphosphoglucuronate
[14509] glucuronosyltransferase (UGT1A1) causing Crigler-Najjar and Gilbert
[14510] syndromes: correlation of genotype to phenotype. Hum. Mutat. 16:
[14511] 297-306, 2000.

[14512] 1086.

[14513] Kadakol, A.; Sappal, B. S.; Ghosh, S. S.; Lowenheim, M.; Chowdhury,

[14514] A.; Chowdhury, S.; Santra, A.; Arias, I. M.; Chowdhury, J. R.; Chowdhury,

[14515] N. R.: Interaction of coding region mutations and the Gilbert-type

[14516] promoter abnormality of the UGT1A1 gene causes moderate degrees of

[14517] unconjugated hyperbilirubinaemia and may lead to neonatal kernicterus.

[14518] (Letter) J. Med. Genet. 38: 244-249, 2001.

[14519] 1087.

[14520] Kaplan, M.; Hammerman, C.; Renbaum, P.; Klein, G.; Levy-Lahad,

[14521] E.: Gilbert's syndrome and hyperbilirubinaemia in ABO-incompatible

[14522] neonates. Lancet 356: 652-653, 2000.

[14523] 1088.

[14524] Kaplan, M.; Renbaum, P.; Levy-Lahad, E.; Hammerman, C.; Lahad,

[14525] A. Beutler, E.: Gilbert syndrome and glucose-6-phosphate dehydrogenase

[14526] deficiency: a dose-dependent genetic interaction crucial to neonatal

[14527] hyperbilirubinemia. Proc. Nat. Acad. Sci. 94: 12128-12132, 1997.

[14528] 1089.

[14529] Kimura, T.; Akaba, K.; Ikegami, T.; Akiba, K.; Kanazawa, C.; Katsuura,

[14530] M.; Shimizu, Y.; Imaizumi, M.; Lin, C.; Hayasaka, K.: Intermittent

[14531] jaundice in patients with acute leukaemia: a common mutation of the

[14532] bilirubin uridine-diphosphate glucuronosyltransferase gene among Asians. J.

[14533] Inherit. Metab. Dis. 22: 747-753, 1999.

[14534] 1090.

[14535] King, C. D.; Rios, G. R.; Tephly, T. R.: UDP-glucuronosyltransferases. Curr.

[14536] Drug Metab. 1: 143-161, 2000.

[14537] 1091.

[14538] Kren, B. T.; Parashar, B.; Bandyopadhyay, P.; Chowdhury, N. R.;

[14539] Chowdhury, J. R.; Steer, C. J.: Correction of the UDP-glucuronosyltransferase

[14540] gene defect in the Gunn rat model of Crigler-Najjar syndrome type

[14541] I with a chimeric oligonucleotide. Proc. Nat. Acad. Sci. 96: 10349-10354,

[14542] 1999.

[14543] 1092.

[14544] Mackenzie, P. I.: The cDNA sequence and expression of a variant

[14545] 17-beta-hydroxysteroid UDP-glucuronosyltransferase. J. Biol. Chem. 265:

[14546] 8699-8703, 1990.

[14547] 1093.

[14548] Clay, S. A.; Shore, N. A.; Landing, B. H.: Triosephosphate isomerase

[14549] deficiency: a case report with neuropathological findings. Am. J.

[14550] Dis. Child. 136: 800-802, 1982.

[14551] 1094.

[14552] Daar, I. O.; Artymiuk, P. J.; Phillips, D. C.; Maquat, L. E.:

[14553] Human triose-phosphate isomerase deficiency: a single amino acid substitution

[14554] results in a thermolabile enzyme. Proc. Nat. Acad. Sci. 83: 7903-7907,

[14555] 1986.

[14556] 1095.

[14557] Decker, R. S.; Mohrenweiser, H. W.: Origin of the triosephosphate

[14558] isomerase isozymes in humans: genetic evidence for the expression

[14559] of a single structural locus. Hum. Genet. 33: 683-691, 1981.

[14560] 1096.

[14561] Eber, S. W.; Dunnwald, M.; Belohradsky, B. H.; Bidling-maier, F.;

[14562] Schievelbein, H.; Weinmann, H. M.; Krietsch, W. K. G.: Hereditary

[14563] deficiency of triosephosphate isomerase in four unrelated families. Europ.

[14564] J. Clin. Invest. 9: 195-202, 1979.

[14565] 1097.

[14566] Eber, S. W.; Dunnwald, M.; Heinemann, G.; Hofstatter, T.; Weinmann,

[14567] H. M.; Belohradsky, B. H.: Prevalence of partial deficiency of red

[14568] cell triosephosphate isomerase in Germany--a study of 3000 people. Hum.

[14569] Genet. 67: 336-339, 1984.

[14570] 1098.

[14571] Eber, S. W.; Pekrun, A.; Bardosi, A.; Gahr, M.; Krietsch, W. K.

[14572] G.; Kruger, J.; Matthei, R.; Schroter, W.: Triosephosphate isomerase

[14573] deficiency: haemolytic anaemia, myopathy with altered mitochondria

[14574] and mental retardation due to a new variant with accelerated enzyme

[14575] catabolism and diminished specific activity. Europ. J. Pediat. 150:

[14576] 761-766, 1991.

[14577] 1099.

[14578] Hendrickson, R. J.; Snapka, R. M.; Sawyer, T. H.; Gracy, R. W.

[14579] : Studies on human triosephosphate isomerase. II. Char-

acterization

[14580] of the enzyme from patients with the cri du chat syn-

drome. Am. J.

[14581] Hum. Genet. 25: 433-438, 1973.

[14582] 1100.

[14583] Hollan, S.; Fujii, H.; Hirono, A.; Hirono, K.; Karro, H.; Miwa,

[14584] S.; Harsanyi, V.; Gyodi, E.; Inselt-Kovacs, M.: Hereditary triosephosphate

[14585] isomerase (TPI) deficiency: two severely affected brothers one with

[14586] and one without neurological symptoms. Hum. Genet. 92: 486-490,

[14587] 1993.

[14588] 1101.

[14589] Hopkinson, D. A.; Harris, H.: Recent work on isozymes in man. Ann.

[14590] Rev. Genet. 5: 5-32, 1971.

[14591] 1102.

[14592] Kwiatowski, J.; Krawczyk, M.; Kornacki, M.; Bailey, K.; Ay-
ala,
[14593] F. J.: Evidence against the exon theory of genes derived
from the
[14594] triose-phosphate isomerase gene. Proc. Nat. Acad. Sci.
92: 8503-8506,
[14595] 1995.

[14596] 1103.

[14597] Logsdon, J. M., Jr.; Tyshenko, M. G.; Dixon, C.; D.-Jafari, J.;

[14598] Walker, V. K.; Palmer, J. D.: Seven newly discovered intron
positions
[14599] in the triose-phosphate isomerase gene: evidence for the
introns-late
[14600] theory. Proc. Nat. Acad. Sci. 92: 8507-8511, 1995.

[14601] 1104.

[14602] Maquat, L. E.; Chilcote, R.; Ryan, P. M.: Human triosephos-
phate
[14603] isomerase cDNA and protein structure: studies of
triosephosphate isomerase
[14604] deficiency in man. J. Biol. Chem. 260: 3748-3753, 1985.

[14605] 1105.

[14606] Merkle, S.; Pretsch, W.: Characterization of triosephos-
phate
[14607] isomerase mutants with reduced enzyme activity in Mus
musculus. Genetics 123:
[14608] 837-844, 1989.

[14609] 1106.

[14610] Mohrenweiser, H. W.: Frequency of enzyme deficiency
variants
[14611] in erythrocytes of newborn infants. Proc. Nat. Acad. Sci.
78: 5046-5050,
[14612] 1981.

[14613] 1107.

[14614] Orosz, F.; Olah, J.; Alvarez, M.; Keseru, G. M.; Szabo, B.;
Wagner,
[14615] G.; Kovari, Z.; Horanyi, M.; Baroti, K.; Martial, J. A.; Hollan,
S.;
[14616] Ovadi, J.: Distinct behavior of mutant triosephosphate iso-
merase
[14617] in hemolysate and in isolated form: molecular basis of en-
zyme deficiency. Blood 98:

[14618] 3106-3112, 2001.

[14619] 1108.

[14620] Pekrun, A.; Neubauer, B. A.; Eber, S. W.; Lakomek, M.; Sei-
del,
[14621] H.; Schroter, W.: Triosephosphate isomerase deficiency:
biochemical
[14622] and molecular genetic analysis for prenatal diagnosis.
Clin. Genet. 47:
[14623] 175-179, 1995.

[14624] 1109.

[14625] Perry, B. A.; Mohrenweiser, H. W.: Human triosephosphate
isomerase:
[14626] substitution of arg for gly at position 122 in a thermola-
bile electromorph
[14627] variant, TPI-Manchester. Hum. Genet. 88: 634-638, 1992.

[14628] 1110.

[14629] Peters, J.; Hopkinson, D. A.; Harris, H.: Genetic and non-
genetic
[14630] variations of triose phosphate isomerase isozymes in hu-
man tissues. Ann.
[14631] Hum. Genet. 36: 297-312, 1973.

[14632] 1111.

[14633] Rethore, M.-O.; Junien, C.; Malpuech, G.; Baccichetti, C.; Tenconi,

[14634] R.; Kaplan, J.-C.; de Romeuf, J.; Lejeune, J.: Localisation du gene

[14635] de la glyceraldehyde 3-phosphate deshydrogenase (G3PD) sur le segment

[14636] distal du bras court du chromosome 12. Ann. Genet. 19: 140-142,

[14637] 1976.

[14638] 1112.

[14639] Rethore, M.-O.; Kaplan, J.-C.; Junien, C.; Lejeune, J.: 12pter-12p12.2:

[14640] Possible assignment of human triosephosphate isomerase. Hum. Genet. 36:

[14641] 235-237, 1977.

[14642] 1113.

[14643] Rosa, R.; Prehu, M.-O.; Calvin, M.-C.; Badoual, J.; Alix, D.;

[14644] Girod, R.: Hereditary triose phosphate isomerase deficiency: seven

[14645] new homozygous cases. Hum. Genet. 71: 235-240, 1985.

[14646] 1114.

[14647] Rudiger, H. W.; Passarge, E.; Hirth, L.; Goedde, H. W.; Blume,

[14648] K. G.; Lohr, G. W.; Benohr, H. C.; Waller, H. D.: Triosephosphate

[14649] isomerase gene not localized on the short arm of chromosome 5 in man.

[14650] (Letter) Nature 228: 1320-1321, 1970.

[14651] 1115.

[14652] Schneider, A.; Forman, L.; Westwood, B.; Yim, C.; Lin, J.; Singh,

[14653] S.; Beutler, E.: The relationship of the -5, -8, and -24 variant

[14654] alleles in African Americans to triosephosphate isomerase (TPI) enzyme

[14655] activity and to TPI deficiency. Blood 92: 2959-2962, 1998.

[14656] 1116.

[14657] Schneider, A. S.; Valentine, W. N.; Hattori, M.; Heins, H. L.,

[14658] Jr.: Hereditary hemolytic anemia with triosephosphate isomerase deficiency. New

[14659] Eng. J. Med. 272: 229-235, 1965.

[14660] 1117.

[14661] Sparkes, R. S.; Carrel, R. E.; Paglia, D. E.: Probable local-
ization
[14662] of a triosephosphate isomerase gene to the short arm of
number 5 human
[14663] chromosome. Nature 224: 367-398, 1969.

[14664] 1118.

[14665] Stoltzfus, A.; Spencer, D. F.; Zuker, M.; Logsdon, J. M., Jr.;

[14666] Doolittle, W. F.: Testing the exon theory of genes: the evi-
dence
[14667] from protein structure. Science 265: 202-207, 1994.

[14668] 1119.

[14669] Vives-Corrons, J.-L.; Rubinson-Skala, H.; Mateo, M.; Es-
tella,
[14670] J.; Feliu, E.; Dreyfus, J.-C.: Triosephosphate isomerase de-
ficiency
[14671] with hemolytic anemia and severe neuromuscular disease:
familial and
[14672] biochemical studies of a case found in Spain. Hum. Genet.
42: 171-180,
[14673] 1978.

[14674] 1120.

[14675] Shprintzen, R. J.; Goldberg, R.; Golding-Kushner, K. J.; Marion,

[14676] R.: Late-onset psychosis in the velo-cardio-facial syndrome. Am.

[14677] J. Med. Genet. 42: 141-142, 1992.

[14678] 1121.

[14679] Sirotkin, H.; Morrow, B.; DasGupta, R.; Goldberg, R.; Patanjali,

[14680] S. R.; Shi, G.; Cannizzaro, L.; Shprintzen, R.; Weissman, S. M.; Kucherlapati,

[14681] R.: Isolation of a new clathrin heavy chain gene with muscle-specific

[14682] expression from the region commonly deleted in velo-cardio-facial

[14683] syndrome. Hum. Molec. Genet. 5: 617-624, 1996.

[14684] 1122.

[14685] Sirotkin, H.; Morrow, B.; Saint-Jore, B.; Puech, A.; Das Gupta,

[14686] R.; Patanjali, S. R.; Skoultchi, A.; Weissman, S. M.; Kucherlapati,

[14687] R.: Identification, characterization, and precise mapping of a human
[14688] gene encoding a novel membrane-spanning protein from the 22q11 region
[14689] deleted in velo-cardio-facial syndrome. Genomics 42: 245-251, 1997.
[14690] 1123.

[14691] Aoyama, T.; Souri, M.; Ueno, I.; Kamijo, T.; Yamaguchi, S.; Rhead,
[14692] W. J.; Tanaka, K.; Hashimoto, T.: Cloning of human very-long-chain
[14693] acyl-coenzyme A dehydrogenase and molecular characterization of its
[14694] deficiency in two patients. Am. J. Hum. Genet. 57: 273-283, 1995.
[14695] 1124.

[14696] Andresen, B. S.; Bross, P.; Vianey-Saban, C.; Divry, P.; Zabot,
[14697] M.-T.; Roe, C. R.; Nada, M. A.; Byskov, A.; Kruse, T. A.; Neve, S.;
[14698] Kristiansen, K.; Knudsen, I.; Corydon, M. J.; Gregersen, N.: Cloning

[14699] and characterization of human very-long-chain acyl-CoA dehydrogenase

[14700] cDNA, chromosomal assignment of the gene and identifi- cation in four

[14701] patients of nine different mutations within the VLCAD gene. Hum.

[14702] Molec. Genet. 5: 461-472, 1996.

[14703] 1125.

[14704] Aoyama, T.; Souri, M.; Ushikubo, S.; Kamijo, T.; Yam-

aguchi, S.;

[14705] Kelley, R. I.; Rhead, W. J.; Uetake, K.; Tanaka, K.; Hashimoto, T.

[14706] : Purification of human very-long-chain acyl-coenzyme A dehydrogenase

[14707] and characterization of its deficiency in seven patients. J. Clin.

[14708] Invest. 95: 2465-2473, 1995.

[14709] 1126.

[14710] Aoyama, T.; Uchida, Y.; Kelley, R. I.; Marble, M.; Hofman, K.;

[14711] Tonsgard, J. H.; Rhead, W. J.; Hashimoto, T.: A novel dis- ease with

[14712] deficiency of mitochondrial very-long-chain acyl-CoA dehydrogenase. Biochem.

[14713] Biophys. Res. Commun. 191: 1369-1372, 1993.

[14714] 1127.

[14715] Bertrand, C.; Largilliere, C.; Zabot, M. T.; Mathieu, M.; Vianey-Saban,

[14716] C.: Very long chain acyl-CoA dehydrogenase deficiency: identification

[14717] of a new inborn error of mitochondrial fatty acid oxidation in fibroblasts. Biochim.

[14718] Biophys. Acta 1180: 327-329, 1993.

[14719] 1128.

[14720] Cox, G. F.; Souri, M.; Aoyama, T.; Rockenmacher, S.; Varvogli,

[14721] L.; Rohr, F.; Hashimoto, T.; Korson, M. S.: Reversal of severe hypertrophic

[14722] cardiomyopathy and excellent neuropsychologic outcome in very-long-chain

[14723] acyl-coenzyme A dehydrogenase deficiency. J. Pediat. 133: 247-253,

[14724] 1998.

[14725] 1129.

[14726] Cox, K. B.; Hamm, D. A.; Millington, D. S.; Matern, D.; Vockley,

[14727] J.; Rinaldo, P.; Pinkert, C. A.; Rhead, W. J.; Lindsey, J. R.; Wood,

[14728] P. A.: Gestational, pathologic and biochemical differences between

[14729] very long-chain acyl-CoA dehydrogenase deficiency and long-chain acyl-CoA

[14730] dehydrogenase deficiency in the mouse. Hum. Molec. Genet. 10: 2069-2077,

[14731] 2001.

[14732] 1130.

[14733] Fukao, T.; Watanabe, H.; Orii, K. E.; Takahashi, Y.; Hirano, A.;

[14734] Kondo, T.; Yamaguchi, S.; Aoyama, T.; Kondo, N.: Myo-pathic form of

[14735] very-long chain acyl-CoA dehydrogenase deficiency: evi-dence for temperature-sensitive

[14736] mild mutations in both mutant alleles in a Japanese girl. Pediat.

[14737] Res. 49: 227-231, 2001.

[14738] 1131.

[14739] Izai, K.; Uchida, Y.; Orii, T.; Yamamoto, S.; Hashimoto, T.:

[14740] Novel fatty acid beta-oxidation enzymes in rat liver mitochondria.

[14741] I. Purification and properties of very-long-chain acyl-coenzyme A

[14742] dehydrogenase. J. Biol. Chem. 267: 1027-1033, 1992.

[14743] 1132.

[14744] Mathur, A.; Sims, H. F.; Gopalakrishnan, D.; Gibson, B.; Rinaldo,

[14745] P.; Vockley, J.; Hug, G.; Strauss, A. W.: Molecular heterogeneity

[14746] in very-long-chain acyl-CoA dehydrogenase deficiency causing pediatric

[14747] cardiomyopathy and sudden death. Circulation 99: 1337-1343, 1999.

[14748] 1133.

[14749] Ogilvie, I.; Pourfarzam, M.; Jackson, S.; Stockdale, C.; Bartlett,

[14750] K.; Turnbull, D. M.: Very long-chain acyl coenzyme A dehydrogenase

[14751] deficiency presenting with exercise-induced myoglobin-
uria. Neurology 44:
[14752] 467-473, 1994.

[14753] 1134.

[14754] Orii, K. O.; Saito-Ohara, F.; Ikeuchi, T.; Orii, T.; Kondo, N.;

[14755] Aoyama, T.; Hashimoto, T.: Assignment of the gene for
very-long-chain
[14756] acyl-CoA dehydrogenase (Acadvl) to mouse chromosome
band 11B2-B5 by
[14757] in situ hybridization. Cytogenet. Cell Genet. 78: 25-26,
1997.
[14758] 1135.

[14759] Parini, R.; Menni, F.; Garavaglia, B.; Fesslova, V.; Melotti,

[14760] D.; Massone, M. L.; Lamantea, E.; Rimoldi, M.; Vizziello, P.;
Gatti,
[14761] R.: Acute, severe cardiomyopathy as main symptom of
late-onset very
[14762] long-chain acyl-coenzyme A dehydrogenase deficiency.
Europ. J. Pediat. 157:
[14763] 992-995, 1998.

[14764] 1136.

[14765] Smelt, A. H. M.; Poorthuis, B. J. H. M.; Onkenhout, W.; Scholte,

[14766] H. R.; Andresen, B. S.; van Duinen, S. G.; Gregersen, N.; Wintzen,

[14767] A. R.: Very long chain acyl-coenzyme A dehydrogenase deficiency with

[14768] adult onset. Ann. Neurol. 43: 540-544, 1998.

[14769] 1137.

[14770] Souri, M.; Aoyama, T.; Orii, K.; Yamaguchi, S.; Hashimoto, T.

[14771] : Mutation analysis of very-long-chain acyl-coenzyme A dehydrogenase

[14772] (VLCAD) deficiency: identification and characterization of mutant

[14773] VLCAD cDNAs from four patients. Am. J. Hum. Genet. 58: 97-106, 1996.

[14774] 1138.

[14775] Strauss, A. W.; Powell, C. K.; Hale, D. E.; Anderson, M. M.; Ahuja,

[14776] A.; Brackett, J. C.; Sims, H. F.: Molecular basis of human mitochondrial

[14777] very-long-chain acyl-CoA dehydrogenase deficiency

causing cardiomyopathy
[14778] and sudden death in childhood. Proc. Nat. Acad. Sci. 92: 10496-10500,
[14779] 1995.

[14780] 1139.

[14781] Tonsgard, J. H.; Stephens, J. K.; Rhead, W. J.; Penn, D.; Horwitz,
[14782] A. L.; Kirschner, B. S.; Whitington, P. F.; Berger, S.; Tripp, M.
[14783] E.: Defect in fatty acid oxidation: laboratory and patho-logic findings
[14784] in a patient. Pediat. Neurol. 7: 125-130, 1991.

[14785] 1140.

[14786] Yamaguchi, S.; Indo, Y.; Coates, P. M.; Hashimoto, T.; Tanaka,
[14787] K.: Identification of very-long-chain acyl-CoA dehydroge-nase deficiency
[14788] in three patients previously diagnosed with long-chain acyl-CoA dehydrogenase
[14789] deficiency. Pediat. Res. 34: 111-113, 1993.

[14790] 1141.

[14791] Berx, G.; Becker, K.-F.; Hofler, H.; van Roy, F.: Mutations of

[14792] the human E-cadherin (CDH1) gene. Hum. Mutat. 12: 226-237, 1998.

[14793] 1142.

[14794] Broeks, A.; Urbanus, J. H. M.; Floore, A. N.; Dahler, E. C.; Klijn,

[14795] J. G. M.; Rutgers, E. J. Th.; Devilee, P.; Russell, N. S.; van Leeuwen,

[14796] F. E.; van't Veer, L. J.: ATM-heterozygous germline muta- tions contribute

[14797] to breast cancer-susceptibility. Am. J. Hum. Genet. 66: 494-500,

[14798] 2000.

[14799] 1143.

[14800] Freund, C. L.; Valle, D.: Cloning and initial characterization

[14801] of the cDNA and gene for human recoverin. (Abstract) Am. J. Hum.

[14802] Genet. 51 (suppl.): A6 only, 1992.

[14803] 1144.

[14804] Heckenlively, J. R.; Fawzi, A. A.; Oversier, J.; Jordan, B. L.;

[14805] Aptsiauri, N.: Autoimmune retinopathy: patients with antirecoverin
[14806] immunoreactivity and panretinal degeneration. Arch. Ophthal. 118:
[14807] 1525-1533, 2000.

[14808] 1145.

[14809] Hurley, J. B.; Dizhoor, A. M.; Ray, S.; Stryer, L.: Recoverin's
[14810] role: conclusion withdrawn. (Letter) Lancet 260: 740 only, 1993.
[14811] 1146.

[14812] McGinnis, J. F.; Lerious, V.; Pazik, J.; Elliott, R. W.: Chromosomal
[14813] assignment of the recoverin gene and cancer-associated retinopathy. Mammalian
[14814] Genome 4: 43-45, 1993.

[14815] 1147.

[14816] McGinnis, J. F.; Leveille, P. J.: Soluble retinal proteins associated
[14817] with photoreceptor cell death in the rd mouse. Curr. Eye Res. 4:
[14818] 1127-1135, 1985.

[14819] 1148.

[14820] McGinnis, J. F.; Stepanik, P. L.; Baehr, W.; Subbaraya, I.; Lerious,

[14821] V.: Cloning and sequencing of the 23 kDa mouse photoreceptor cell-specific

[14822] protein. FEBS Lett. 302: 172-176, 1992.

[14823] 1149.

[14824] Murakami, A.; Yajima, T.; Inana, G.: Isolation of human retinal

[14825] genes: recoverin cDNA and gene. Biochem. Biophys. Res. Commun. 187:

[14826] 234-244, 1992.

[14827] 1150.

[14828] Polans, A. S.; Buczylko, J.; Crabb, J.; Palczewski, K.: A photoreceptor

[14829] calcium binding protein is recognized by autoantibodies obtained from

[14830] patients with cancer-associated retinopathy. J. Cell Biol. 112:

[14831] 981-989, 1991.

[14832] 1151.

[14833] Polans, A. S.; Witkowska, D.; Haley, T. L.; Amundson, D.; Baizer,

[14834] L.; Adamus, G.: Recoverin, a photoreceptor-specific calcium-binding

[14835] protein, is expressed by the tumor of a patient with cancer-associated

[14836] retinopathy. Proc. Nat. Acad. Sci. 92: 9176-9180, 1995.

[14837] 1152.

[14838] Thirkill, C. E.; Roth, A. M.; Keltner, J. L.: Cancer-associated

[14839] retinopathy. Arch. Ophthal. 105: 372-375, 1987.

[14840] 1153.

[14841] Jaffe, E. R.: Personal Communication. Bronx, N. Y. 8/5/1987.

[14842] 1154.

[14843] Bascom, R.; Manara, S.; Gallie, B.; Willard, H.; Kalnins, V.; McInnes,

[14844] R. R.: Identification of a new mammalian photoreceptor-specific gene

[14845] family. (Abstract) Am. J. Hum. Genet. 45: A172 only, 1989.

[14846] 1155.

[14847] Bascom, R. A.; Garcia-Heras, J.; Hsieh, C.-L.; Gerhard, D. S.;

[14848] Jones, C.; Francke, U.; Willard, H. F.; Ledbetter, D. H.; McInnes,

[14849] R. R.: Localization of the photoreceptor gene ROM1 to human chromosome

[14850] 11 and mouse chromosome 19: sublocalization to human 11q13 between

[14851] PGA and PYGM. Am. J. Hum. Genet. 51: 1028-1035, 1992.

[14852] 1156.

[14853] Bascom, R. A.; Liu, L.; Chen, J.; Duncan, A.; Kimberling, W. J.;

[14854] Moller, C. G.; Humphries, P.; Nathans, J.; McInnes, R. R.: ROM1:

[14855] a candidate gene for autosomal dominant retinitis pigmentosa (ADRP),

[14856] Usher syndrome type 1, and Best vitelliform macular dystrophy. (Abstract) Am.

[14857] J. Hum. Genet. 51 (suppl.): A6 only, 1992.

[14858] 1157.

[14859] Finegold, D. N.; Armitage, M. M.; Galiani, M.; Matise, T. C.;

[14860] Pandian, M. R.; Perry, Y. M.; Deka, R.; Ferrell, R. E.: Preliminary
[14861] localization of a gene for autosomal dominant hypoparathyroidism to
[14862] chromosome 3q13. Pediat. Res. 36: 414-417, 1994.

[14863] 1158.

[14864] Goldbloom, R. B.; Gillis, D. A.; Prasad, M.: Hereditary parathyroid
[14865] hyperplasia: a surgical emergency of early infancy. Pediatrics 49:
[14866] 514-523, 1972.

[14867] 1159.

[14868] Ho, C.; Conner, D. A.; Pollak, M. R.; Ladd, D. J.; Kifor, O.;

[14869] Warren, H. B.; Brown, E. M.; Seidman, J. G.; Seidman, C. E.: A mouse
[14870] model of human familial hypocalciuric hypercalcemia and neonatal severe
[14871] hyperparathyroidism. Nature Genet. 11: 389-394, 1995.

[14872] 1160.

[14873] Janicic, N.; Soliman, E.; Pausova, Z.; Seldin, M. F.; Riviere,

[14874] M.; Szpirer, J.; Szpirer, C.; Hendy, G. N.: Mapping of the calcium-sensing

[14875] gene (CASR) to human chromosome 3q13.3-21 by fluorescence in situ

[14876] hybridization, and localization to rat chromosome 11 and mouse chromosome

[14877] 16. Mammalian Genome 6: 798-801, 1995.

[14878] 1161.

[14879] Marx, S. J.; Attie, M. F.; Spiegel, A. M.; Levine, M. A.; Lasker,

[14880] R. D.; Fox, M.: An association between neonatal severe primary hyperparathyroidism

[14881] and familial hypocalciuric hypercalcemia in three kindreds. New Eng.

[14882] J. Med. 306: 257-264, 1982.

[14883] 1162.

[14884] Meeran, K.; Husain, M.; Puccini, M.; Scott, H.; Dionisi-Vici,

[14885] D.; Harvey, D. R.; Lynn, J.; Thakker, R. V.: Neonatal primary hyperparathyroidism

[14886] masked by vitamin D deficiency. Clin. Endocr. 41: 531-534, 1994.

[14887] 1163.

[14888] Pearce, S. H. S.; Trump, D.; Wooding, C.; Besser, G. M.; Chew,

[14889] S. L.; Grant, D. B.; Heath, D. A.; Hughes, I. A.; Paterson, C. R.;

[14890] Whyte, M. P.; Thakker, R. V.: Calcium-sensing receptor mutations

[14891] in familial benign hypercalcemia and neonatal hyper- parathyroidism. J.

[14892] Clin. Invest. 96: 2683-2692, 1995.

[14893] 1164.

[14894] Pollak, M. R.; Brown, E. M.; Chou, Y.-H. W.; Hebert, S. C.; Marx,

[14895] S. J.; Steinmann, B.; Levi, T.; Seidman, C. E.; Seidman, J. G.: Mutations

[14896] in the human Ca(2+)-sensing receptor gene cause familial hypocalciuric

[14897] hypercalcemia and neonatal severe hyperparathyroidism. Cell 75:

[14898] 1297-1303, 1993.

[14899] 1165.

[14900] Pollak, M. R.; Brown, E. M.; Estep, H. L.; McLaine, P. N.; Ki-
for,
[14901] O.; Park, J.; Hebert, S. C.; Seidman, C. E.; Seidman, J. G.:
Autosomal
[14902] dominant hypocalcaemia caused by a Ca(2+)-sensing re-
ceptor gene mutation. Nature
[14903] Genet. 8: 303-307, 1994.

[14904] 1166.

[14905] Pollak, M. R.; Chou, Y.-H. W.; Marx, S. J.; Steinmann, B.;
Cole,
[14906] D. E. C.; Brandi, M. L.; Papapoulos, S. E.; Menko, F. H.;
Hendy, G.
[14907] N.; Brown, E. M.; Seidman, C. E.; Seidman, J. G.: Familial
hypocalciuric
[14908] hypercalcemia and neonatal severe hyperparathyroidism:
effects of
[14909] mutant gene dosage on phenotype. J. Clin. Invest. 93:
1108-1112,
[14910] 1994.

[14911] 1167.

[14912] Pratt, E. L.; Geren, B. B.; Neuhauser, E. B. D.: Hypercal-
cemia

[14913] and idiopathic hyperplasia of the parathyroid glands in an infant. J.

[14914] Pediat. 30: 388-399, 1947.

[14915] 1168.

[14916] Steinmann, B.; Gnehm, H. E.; Rao, V. H.; Kind, H. P.; Prader,

[14917] A.: Neonatal severe primary hyperparathyroidism and alkaptonuria

[14918] in a boy born to related parents with familial hypocalciuric hypercalcemia. Helv.

[14919] Paediat. Acta 39: 171-186, 1984.

[14920] 1169.

[14921] Stadler, H. S.; Padanilam, B. J.; Buetow, K.; Murray, J. C.; Solursh,

[14922] M.: Identification and genetic mapping of a homeobox gene to the

[14923] 4p16.1 region of human chromosome 4. Proc. Nat. Acad. Sci. 89: 11579-11583,

[14924] 1992.

[14925] 1170.

[14926] Wang, W.; Yoshiura, K.; Murray, J.; Lufkin, T.: Assignment

of

[14927] the murine Hmx1 homeobox gene to the proximal region of mouse chromosome

[14928] 5. Mammalian Genome 8: 869-876, 1997.

[14929] 1171.

[14930] Faiella, A.; D'Esposito, M.; Rambaldi, M.; Acampora, D.; Balsofiore,

[14931] S.; Stornaiuolo, A.; Mallamaci, A.; Migliaccio, E.; Gulisano, M.;

[14932] Simeone, A.; Boncinelli, E.: Isolation and mapping of EVX1, a human

[14933] homeobox gene homologous to even-skipped, localized at the 5-prime

[14934] end of HOX1 locus on chromosome 7. Nucleic Acids Res. 19: 6541-6545,

[14935] 1991.

[14936] 1172.

[14937] Raftery, M. J.; Schwab, M.; Eibert, S. M.; Samstag, Y.; Wal-

[14938] czak, H.; Schonrich, G.: Targeting the function of mature dendritic cells

[14939] by human cytomegalovirus: a multilayered viral defense

strategy. Immunity 15:
[14940] 997-1009, 2001.

[14941] 1173.

[14942] Gough, A. C.; Zhong, S.; Wolf, C. R.; Spurr, N. K.: Chromo-
some
[14943] assignment of the human glutathione S-transferase mu-3
gene (GSTM3)
[14944] to chromosome 1 by gene specific polymerase chain reac-
tion. Cytogenet.
[14945] Cell Genet. 65: 111-114, 1994.

[14946] 1174.

[14947] Smith, R. E., Jr.; Jaiyesimi, I. A.; Meza, L. A.; Tchekme-
dyian,
[14948] N. S.; Chan, D.; Griffith, H.; Brosman, S.; Bukowski, R.;
Murdock,
[14949] M.; Rarick, M.; Saven, A.; Colowick, A. B.; Fleishman, A.;
Gayko,
[14950] U.; Glaspy, J.: Novel erythropoiesis stimulating protein
(NESP) for
[14951] the treatment of anaemia of chronic disease associated
with cancer. Brit.
[14952] J. Cancer 84: 24-30, 2001.

[14953] 1175.

[14954] Mangion, J.; Rahman, N.; Mansour, S.; Brice, G.; Ros-
botham, J.;
[14955] Child, A. H.; Murday, V. A.; Mortimer, P. S.; Barfoot, R.;
Sigurdsson,
[14956] A.; Edkins, S.; Sarfarazi, M.; Burnand, K.; Evans, A. L.;
Nunan, T.
[14957] O.; Stratton, M. R.; Jeffery, S.: A gene for lymphedema-dis-
tichiasis
[14958] maps to 16q24.3. Am. J. Hum. Genet. 65: 427-432, 1999.

[14959] 1176.

[14960] Galiegue-Zouitina, S.; Quief, S.; Hildebrand, M.-P.; Denis,
C.;
[14961] Detourmignies, L.; Lai, J.-L.; Kerckaert, J.-P.: Nonrandom
fusion
[14962] of L-plastin (LCP1) and LAZ3 (BCL6) genes by
t(3;13)(q27;q14) chromosome
[14963] translocation in two cases of B-cell non-Hodgkin lym-
phoma. Genes
[14964] Chromosomes Cancer 26: 97-105, 1999.

[14965] 1177.

[14966] Giometti, C. S.; Anderson, N. L.: A variant of human non-muscle
[14967] tropomyosin found in fibroblasts by using two-dimensional electrophoresis. J.
[14968] Biol. Chem. 256: 11840-11846, 1981.

[14969] 1178.

[14970] Hamaguchi, H.; Ohta, A.; Mukai, R.; Yabe, T.; Yamada, M.: Genetic
[14971] analysis of human lymphocyte proteins by two-dimensional gel electrophoresis:
[14972] 1. Detection of genetic variant polypeptides in PHA-stimulated peripheral
[14973] blood lymphocyte. Hum. Genet. 59: 215-220, 1981.

[14974] 1179.

[14975] Hamaguchi, H.; Yamada, M.; Noguchi, A.; Fujii, K.; Shibasaki, M.;
[14976] Mukai, R.; Yabe, T.; Kondo, I.: Genetic analysis of human lymphocyte
[14977] proteins by two-dimensional gel electrophoresis: 2. Genetic polymorphism
[14978] of lymphocyte cytosol 64K polypeptide. Hum. Genet. 60: 176-180,

[14979] 1982.

[14980] 1180.

[14981] Hamaguchi, H.; Yamada, M.; Shibasaki, M.; Kondo, I.: Genetic analysis

[14982] of human lymphocyte proteins by two-dimensional gel electrophoresis:

[14983] 4. Genetic polymorphism of cytosol 100k polypeptide. Hum. Genet. 62:

[14984] 148-151, 1982.

[14985] 1181.

[14986] Hamaguchi, H.; Yamada, M.; Shibasaki, M.; Mukai, R.; Yabe, T.;

[14987] Kondo, I.: Genetic analysis of human lymphocyte proteins by two-dimensional

[14988] gel electrophoresis: 3. Frequent occurrence of genetic variants in

[14989] some abundant polypeptides of PHA-stimulated peripheral blood lymphocytes. Hum.

[14990] Genet. 62: 142-147, 1982.

[14991] 1182.

[14992] Klose, J.: Protein mapping by combined isoelectric focus-

ing and
[14993] electrophoresis of mouse tissue: a novel approach to testing for induced
[14994] point mutations in mammals. Humangenetik 26: 231-243, 1975.
[14995] 1183.

[14996] Klose, J.; Feller, M.: Genetic variability of proteins from plasma
[14997] membranes and cytosols of mouse organs. Biochem. Genet. 19: 859-870,
[14998] 1981.

[14999] 1184.

[15000] Kondo, I.; Hamaguchi, H.: Evidence for the close linkage between
[15001] lymphocyte cytosol polypeptide with molecular weight of 64,000 (LCP1)
[15002] and esterase D. Am. J. Hum. Genet. 37: 1106-1111, 1985.

[15003] 1185.

[15004] Kondo, I.; Hamaguchi, H.: Study of the linkage relationship between
[15005] LCP1 and ESD. (Abstract) Cytogenet. Cell Genet. 40: 672

only, 1985.

[15006] 1186.

[15007] Michelson, A. M.; Russell, E. S.; Harman, P. J.: Dystrophia muscularis:

[15008] a hereditary primary myopathy in the house mouse. Proc. Nat. Acad.

[15009] Sci. 41: 1079-1084, 1955.

[15010] 1187.

[15011] Roberts, R. G.; Freeman, T. C.; Kendall, E.; Vetrie, D. L. P.;

[15012] Dixon, A. K.; Shaw-Smith, C.; Bone, Q.; Bobrow, M.: Characterization

[15013] of DRP2, a novel human dystrophin homologue. Nature Genet. 13: 223-226,

[15014] 1996.

[15015] 1188.

[15016] Xu, H.; Christmas, P.; Wu, X.-R.; Wewer, U. M.; Engvall, E.:

[15017] Defective muscle basement membrane and lack of M-laminin in the dystrophic

[15018] dy/dy mouse. Proc. Nat. Acad. Sci. 91: 5572-5576, 1994.

[15019] 1189.

[15020] Watkins, P. C.; Eddy, R.; Hoffman, N.; Stanislovitis, P.; Beck,
[15021] A. K.; Galli, J.; Vellucci, V.; Gusella, J. F.; Shows, T. B.: Regional
[15022] assignment of the erythropoietin gene to human chromosome region 7pter-q22. Cytogenet.
[15023] Cell Genet. 42: 214-218, 1986.

[15024] 1190.

[15025] Akashi, K.; Traver, D.; Miyamoto, T.; Weissman, I. L.: A clonogenic
[15026] common myeloid progenitor that gives rise to all myeloid lineages. Nature 404:
[15027] 193-197, 2000.

[15028] 1191.

[15029] Arcasoy, M. O.; Degar, B. A.; Harris, K. W.; Forget, B. G.: Familial
[15030] erythrocytosis associated with a short deletion in the erythropoietin
[15031] receptor gene. Blood 89: 4628-4635, 1997.

[15032] 1192.

[15033] Budarf, M.; Huebner, K.; Emanuel, B.; Croce, C. M.;

Copeland, N.

[15034] G.; Jenkins, N. A.; D'Andrea, A. D.: Assignment of the erythropoietin

[15035] receptor (EPOR) gene to mouse chromosome 9 and human chromosome 19. Genomics 8:

[15036] 575-578, 1990.

[15037] 1193.

[15038] D'Andrea, A. D.; Yoshimura, A.; Youssoufian, H.; Zon, L. I.; Koo,

[15039] J.-W.; Lodish, H. F.: The cytoplasmic region of the erythropoietin

[15040] receptor contains nonoverlapping positive and negative growth-regulatory

[15041] domains. Molec. Cell. Biol. 11: 1980-1987, 1991.

[15042] 1194.

[15043] D'Andrea, A. D.; Zon, L. I.: Erythropoietin receptor: subunit

[15044] structure and activation. J. Clin. Invest. 86: 681-687, 1990.

[15045] 1195.

[15046] de la Chapelle, A.; Traskelin, A.-L.; Juvonen, E.: Truncated erythropoietin

[15047] receptor causes dominantly inherited benign human erythrocytosis. Proc.

[15048] Nat. Acad. Sci. 90: 4495-4499, 1993.

[15049] 1196.

[15050] Divoky, V.; Liu, Z.; Ryan, T. M.; Prchal, J. F.; Townes, T. M.;

[15051] Prchal, J. T.: Mouse model of congenital polycythemia: homologous

[15052] replacement of murine gene by mutant human erythropoietin receptor

[15053] gene. Proc. Nat. Acad. Sci. 98: 986-991, 2001.

[15054] 1197.

[15055] Hess, G.; Rose, P.; Gamm, H.; Papadileris, S.; Huber, C.; Seliger,

[15056] B.: Molecular analysis of the erythropoietin receptor system in patients

[15057] with polycythaemia vera. Brit. J. Haemat. 88: 794-802, 1994.

[15058] 1198.

[15059] Jones, S. S.; D'Andrea, A. D.; Haines, L. L.; Wong, G. G.: Human

[15060] erythropoietin receptor: cloning, expression, and biologic

characterization. Blood 76:
[15061] 31-35, 1990.

[15062] 1199.

[15063] Kralovics, R.; Indrak, K.; Stopka, T.; Berman, B. W.; Prchal,

[15064] J. F.; Prchal, J. T.: Two new EPO receptor mutations: truncated EPO
[15065] receptors are most frequently associated with primary familial and
[15066] congenital polycythemias. Blood 90: 2057-2061, 1997.

[15067] 1200.

[15068] Kralovics, R.; Sokol, L.; Prchal, J. T.: Absence of polycythemia
[15069] in a child with a unique erythropoietin receptor mutation in a family
[15070] with autosomal dominant primary polycythemia. J. Clin. Invest. 102:
[15071] 124-129, 1998.

[15072] 1201.

[15073] Le Couedic, J.-P.; Mitjavila, M.-T.; Villeval, J.-L.; Feger, F.;

[15074] Gobert, S.; Mayeux, P.; Casadevall, N.; Vainchenker, W.:

Missense
[15075] mutation of the erythropoietin receptor is a rare event in human erythroid
[15076] malignancies. Blood 87: 1502-1511, 1996.

[15077] 1202.

[15078] Longmore, G. D.; Lodish, H. F.: An activating mutation in the
[15079] murine erythropoietin receptor induces erythroleukemia in mice: a
[15080] cytokine receptor superfamily oncogene. Cell 67: 1089-1102, 1991.
[15081] 1203.

[15082] Maouche, L.; Tournamille, C.; Hattab, C.; Boffa, G.; Cartron,
[15083] J.-P.; Chretien, S.: Cloning of the gene encoding the human erythropoietin
[15084] receptor. Blood 78: 2557-2563, 1991.

[15085] 1204.

[15086] Noguchi, C. T.; Bae, K. S.; Chin, K.; Wada, Y.; Schechter, A.

[15087] N.; Hankins, W. D.: Cloning of the human erythropoietin receptor

[15088] gene. Blood 78: 2548-2556, 1991.

[15089] 1205.

[15090] Penny, L. A.; Forget, B. G.: Genomic organization of the human

[15091] erythropoietin receptor gene. Genomics 11: 974-980, 1991.

[15092] 1206.

[15093] Percy, M. J.; McMullin, M. F.; Roques, A. W. W.; Westwood, N.

[15094] B.; Acharya, J.; Hughes, A. E.; Lappin, T. R. J.; Pearson, T. C.:

[15095] Erythrocytosis due to a mutation in the erythropoietin receptor gene. Brit.

[15096] J. Haemat. 100: 407-410, 1998.

[15097] 1207.

[15098] Sistonen, P.; Traskelin, A.-L.; Lehvaslaiho, H.; de la Chapelle,

[15099] A.: Genetic mapping of the erythropoietin receptor gene. Hum. Genet. 92:

[15100] 299-301, 1993.

[15101] 1208.

[15102] Sokol, L.; Luhovy, M.; Guan, Y.; Prchal, J. F.; Semenza, G. L.;

[15103] Prchal, J. T.: Primary familial polycythemia: a frameshift mutation

[15104] in the erythropoietin receptor gene and increased sensitivity of erythroid

[15105] progenitors to erythropoietin. Blood 86: 15-22, 1995.

[15106] 1209.

[15107] Ward, J. C.; Harris, K. W.; Penny, L. A.; Forget, B. G.; Kitamura,

[15108] T.; Winkelmann, J. C.: A structurally abnormal erythropoietin receptor

[15109] gene in a human erythroleukemia cell line. Exp. Hemat. 20: 371-373,

[15110] 1992.

[15111] 1210.

[15112] Winkelmann, J. C.; Penny, L. A.; Deaven, L. L.; Forget, B. G.;

[15113] Jenkins, R. B.: The gene for the human erythropoietin receptor: analysis

[15114] of the coding sequence and assignment to chromosome 19p. Blood 76:

[15115] 24-30, 1990.

[15116] 1211.

[15117] Yu, X.; Lin, C.-S.; Costantini, F.; Noguchi, C. T.: The hu-
man

[15118] erythropoietin receptor gene rescues erythropoiesis and
developmental

[15119] defects in the erythropoietin receptor null mouse. Blood
98: 475-477,

[15120] 2001.

[15121] 1212.

[15122] Baier, L. J.; Sacchettini, J. C.; Knowler, W. C.; Eads, J.;
Paolisso,

[15123] G.; Tataranni, P. A.; Mochizuki, H.; Bennett, P. H.; Bogar-
dus, C.;

[15124] Prochazka, M.: An amino acid substitution in the human
intestinal

[15125] fatty acid binding protein is associated with increased
fatty acid

[15126] binding, increased fat oxidation, and insulin resistance. J.
Clin.

[15127] Invest. 95: 1281-1287, 1995.

[15128] 1213.

[15129] Carlsson, M.; Orho-Melander, M.; Hedenbro, J.; Almgren, P.; Groop,

[15130] L. C.: The T54 allele of the intestinal fatty acid-binding protein

[15131] 2 is associated with a parental history of stroke. J. Clin. Endocr.

[15132] Metab. 85: 2801-2804, 2000.

[15133] 1214.

[15134] Georgopoulos, A.; Aras, O.; Tsai, M. Y.: Codon-54 polymorphism

[15135] of the fatty acid-binding protein 2 gene is associated with elevation

[15136] of fasting and postprandial triglyceride in type 2 diabetes. J. Clin.

[15137] Endocr. Metab. 85: 3155-3160, 2000.

[15138] 1215.

[15139] Hegele, R. A.; Harris, S. B.; Hanley, A. J. G.; Sadikian, S.; Connelly,

[15140] P. W.; Zinman, B.: Genetic variation of intestinal fatty acid-binding

[15141] protein associated with variation in body mass in aborigi-
nal Canadians. J.

[15142] Clin. Endocr. Metab. 81: 4334-4337, 1996.

[15143] 1216.

[15144] Polymeropoulos, M. H.; Rath, D. S.; Xiao, H.; Merril, C. R.:
Trinucleotide

[15145] repeat polymorphism at the human intestinal fatty acid
binding protein

[15146] gene (FABP2). Nucleic Acids Res. 18: 7198 only, 1991.

[15147] 1217.

[15148] Prochazka, M.; Lillioja, S.; Tait, J. F.; Knowler, W. C.; Mott,

[15149] D. M.; Spraul, M.; Bennett, P. H.; Bogardus, C.: Linkage of
chromosomal

[15150] markers on 4q with a putative gene determining maximal
insulin action

[15151] in Pima Indians. Diabetes 42: 514-519, 1993.

[15152] 1218.

[15153] Sipilainen, R.; Uusitupa, M.; Heikkinen, S.; Rissanen, A.;
Laakso,

[15154] M.: Variants in the human intestinal fatty acid binding
protein 2

[15155] gene in obese subjects. J. Clin. Endocr. Metab. 82:
2629-2632, 1997.
[15156] 1219.
[15157] Sparkes, R. S.; Mohandas, T.; Heinzmann, C.; Gordon, J. I.;
Klisak,
[15158] I.; Zollman, S.; Sweetser, D. A.; Ragunathan, L.; Winokur,
S.; Lusis,
[15159] A. J.: Human fatty acid binding protein assignments in-
testinal to
[15160] 4q28-4q31 and liver to 2p11. (Abstract) Cytogenet. Cell
Genet. 46:
[15161] 697 only, 1987.
[15162] 1220.
[15163] Sweetser, D. A.; Birkenmeier, E. H.; Klisak, I. J.; Zollman,
S.;
[15164] Sparkes, R. S.; Mohandas, T.; Lusis, A. J.; Gordon, J. I.: The
human
[15165] and rodent intestinal fatty acid binding protein genes: a
comparative
[15166] analysis of their structure, expression, and linkage rela-
tionships. J.
[15167] Biol. Chem. 262: 16060-16071, 1987.

[15168] 1221.

[15169] Weber, J. L.; Kwitek, A. E.; May, P. E.; Polymeropoulos, M.:

[15170] Dinucleotide repeat polymorphism at the D12S43 locus. Nucleic Acids

[15171] Res. 18: 4637 only, 1990.

[15172] 1222.

[15173] Weber, J. L.; May, P. E.: Abundant class of human DNA polymorphisms

[15174] which can be typed using the polymerase chain reaction. Am. J. Hum.

[15175] Genet. 44: 388-396, 1989.

[15176] 1223.

[15177] Buckle, V. J.; Fujita, N.; Bateson, A. N.; Darlison, M. G.; Barnard,

[15178] E. A.: Localization of human GABA-A receptor subunit genes to chromosomes

[15179] 4 and 5. (Abstract) Cytogenet. Cell Genet. 51: 972 only, 1989.

[15180] 1224.

[15181] Danciger, M.; Farber, D. B.; Kozak, C. A.: Genetic mapping of

[15182] three GABA-A receptor-subunit genes in the mouse. Genomics 16: 361-365,

[15183] 1993.

[15184] 1225.

[15185] Hadingham, K. L.; Wingrove, P.; Le Bourdelles, B.; Palmer, K. J.;

[15186] Ragan, C. I.; Whiting, P. J.: Cloning of cDNA sequences en-coding

[15187] human alpha-2 and alpha-3 gamma-aminobutyric acid-A receptor subunits

[15188] and characterization of the benzodiazepine pharmacology of recombinant

[15189] alpha-1-, alpha-2-, alpha-3-, and alpha-5-containing human gamma-aminobutyric

[15190] acid-a receptors. Molec. Pharm. 43: 970-975, 1993.

[15191] 1226.

[15192] Low, K.; Crestani, F.; Keist, R.; Benke, D.; Brunig, I.; Benson,

[15193] J. A.; Fritschy, J.-M.; Rulicke, T.; Bluethmann, H.; Mohler, H.; Rudolph,

[15194] U.: Molecular and neuronal substrate for the selective attenuation

[15195] of anxiety. Science 290: 131-134, 2000.

[15196] 1227.

[15197] Glatt, K.; Glatt, H.; Lalande, M.: Structure and organization

[15198] of GABRB3 and GABRA5. Genomics 41: 63-69, 1997.

[15199] 1228.

[15200] Glatt, K. A.; Sinnett, D.; Lalande, M.: Dinucleotide repeat polymorphism

[15201] at the GABA-A receptor alpha-5 (GABRA5) locus at chromosome 15q11-q13. Hum.

[15202] Molec. Genet. 1: 348 only, 1992.

[15203] 1229.

[15204] Knoll, J. H. M.; Sinnett, D.; Wagstaff, J.; Glatt, K.; Wilcox,

[15205] A. S.; Whiting, P.; Wingrove, P.; Sikela, J.; Lalande, M.: FISH ordering

[15206] of DNA markers within the Angelman/Prader-Willi chromosomal regions:

[15207] mapping of a second GABA-A receptor subunit gene, GABRA5. (Abstract) Am.

[15208] J. Hum. Genet. 51 (suppl.): A9 only, 1992.

[15209] 1230.

[15210] Papadimitriou, G. N.; Dikeos, D. G.; Karadima, G.; Avramopoulos,
[15211] D.; Daskalopoulou, E. G.; Vassilopoulos, D.; Stefanis, C. N.: Association
[15212] between the GABA-A receptor alpha-5 subunit gene locus (GABRA5) and
[15213] bipolar affective disorder. Am. J. Med. Genet. 81: 73-80, 1998.
[15214] 1231.

[15215] Ritchie, R. J.; Mattei, M.-G.; Lalande, M.: A large polymor-phic
[15216] repeat in the pericentromeric region of human chromo-some 15q contains
[15217] three partial gene duplications. Hum. Molec. Genet. 7: 1253-1260,
[15218] 1998.

[15219] 1232.

[15220] Russek, S. J.; Farb, D. H.: Mapping of the beta-2 subunit gene
[15221] (GABRB2) to microdissected human chromosome 5q34-q35 defines a gene
[15222] cluster for the most abundant GABA-A receptor isoform.

Genomics 23:
[15223] 528-533, 1994.

[15224] 1233.

[15225] Wingrove, P.; Hadingham, K.; Wafford, K.; Kemp, J. A.; Ragan, C.
[15226] I.; Whiting, P.: Cloning and expression of a cDNA encoding the human
[15227] GABA-A receptor alpha-5 subunit. Biochem. Soc. Trans. 20: 18S only,
[15228] 1991.

[15229] 1234.

[15230] Asuru, A. I.; Mellor, H.; Thomas, N. S. B.; Yu, L.; Chen, J.-J.;
[15231] Crosby, J. S.; Hartson, S. D.; Kimball, S. R.; Jefferson, L. S.; Matts,
[15232] R. L.: Cloning and characterization of cDNAs encoding the epsilon-subunit
[15233] of eukaryotic initiation factor-2B from rabbit and human. Biochim.
[15234] Biophys. Acta 1307: 309-317, 1996.

[15235] 1235.

[15236] Shimizu, F.; Katagiri, T.; Suzuki, M.; Watanabe, T. K.; Okuno,
[15237] S.; Kuga, Y.; Nagata, M.; Fujiwara, T.; Nakamura, Y.; Takahashi, E.
[15238] : Cloning and chromosome assignment to 1q32 of a human cDNA (RAB7L1)
[15239] encoding a small GTP-binding protein, a member of the RAS superfamily. Cytogenet.
[15240] Cell Genet. 77: 261-263, 1997.

[15241] 1236.

[15242] Brenner, R.; Perez, G. J.; Bonev, A. D.; Eckman, D. M.; Kosek,
[15243] J. C.; Wiler, S. W.; Patterson, A. J.; Nelson, M. T.; Aldrich, R.
[15244] W.: Vasoregulation by the beta-1 subunit of the calcium-activated
[15245] potassium channel. Nature 407: 870-876, 2000.

[15246] 1237.

[15247] Jiang, Z.; Wallner, M.; Meera, P.; Toro, L.: Human and rodent
[15248] MaxiK channel beta-subunit genes: cloning and characterization. Genomics 55:

[15249] 57-67, 1999.

[15250] 1238.

[15251] Knaus, H.-G.; Folander, K.; Garcia-Calvo, M.; Garcia, M. L.; Kaczorowski,

[15252] G. J.; Smith, M.; Swanson, R.: Primary sequence and immunological

[15253] characterization of beta-subunit of high conductance Ca(2+)-activated

[15254] K+ channel from smooth muscle. J. Biol. Chem. 269: 17274-17278,

[15255] 1994.

[15256] 1239.

[15257] Meera, P.; Wallner, M.; Jiang, Z.; Toro, L.: A calcium switch

[15258] for the functional coupling between alpha (hslo) and beta subunits

[15259] (K(V), Ca-beta) of maxi K channels. FEBS Lett. 382: 84-88, 1996.

[15260] 1240.

[15261] Tseng-Crank, J.; Godinot, N.; Johansen, T. E.; Ahring, P. K.; Strobaek,

[15262] D.; Mertz, R.; Foster, C. D.; Olesen, S.-P.; Reinhart, P. H.:

Cloning,

[15263] expression, and distribution of a Ca(2+)-activated K+ channel beta-subunit

[15264] from human brain. Proc. Nat. Acad. Sci. 93: 9200-9205, 1996.

[15265] 1241.

[15266] Xie, J.; Black, D. L.: A CaMK IV responsive RNA element mediates

[15267] depolarization-induced alternative splicing of ion channels. Nature 410:

[15268] 936-439, 2001.

[15269] 1242.

[15270] Tsujimoto, A.; Nyunoya, H.; Morita, T.; Sato, T.; Shimotohno, K.

[15271] : Isolation of cDNAs for DNA-binding proteins which specifically bind

[15272] to a tax-responsive enhancer element in the long terminal repeat of

[15273] human T-cell leukemia virus type I. J. Virol. 65: 1420-1426, 1991.

[15274] 1243.

[15275] Griffitts, J. S.; Whitacre, J. L.; Stevens, D. E.; Aroian, R. V.

[15276] : Bt toxin resistance from loss of a putative carbohydrate-modifying

[15277] enzyme. Science 293: 860-864, 2001.

[15278] 1244.

[15279] Isshiki, S.; Togayachi, A.; Kudo, T.; Nishihara, S.; Watan-abe,

[15280] M.; Kubota, T.; Kitajima, M.; Shiraishi, N.; Sasaki, K.; An-doh, T.;

[15281] Narimatsu, H.: Cloning, expression, and characterization of a novel

[15282] UDP-galactose:beta-N-acetylglucosamine beta-1,3-galactosyltransferase

[15283] (beta-3-Gal-T5) responsible for synthesis of type 1 chain in colorectal

[15284] and pancreatic epithelia and tumor cells derived there-from. J. Biol.

[15285] Chem. 274: 12499-12507, 1999.

[15286] 1245.

[15287] Burchett, S. A.; Volk, M. L.; Bannon, M. J.; Granneman, J. G.:

[15288] Regulators of G protein signaling: rapid changes in mRNA abundance

[15289] in response to amphetamine. J. Neurochem. 70: 2216-2219, 1998.

[15290] 1246.

[15291] Chen, C.-K.; Burns, M. E.; He, W.; Wensel, T. G.; Baylor, D. A.;

[15292] Simon, M. I.: Slowed recovery of rod photoresponse in mice lacking

[15293] the GTPase accelerating protein RGS9-1. Nature 403: 557-560, 2000.

[15294] 1247.

[15295] Granneman, J. G.; Zhai, Y.; Zhu, Z.; Bannon, M. J.; Burchett, S.

[15296] A.; Schmidt, C. J.; Andrade, R.; Cooper, J.: Molecular char-

[15297] acterization of human and rat RGS 9L, a novel splice variant enriched in dopamine

[15298] target regions, and chromosomal localization of the RGS 9 gene. Molec.

[15299] Pharm. 54: 687-694, 1998.

[15300] 1248.

[15301] He, W.; Cowan, C. W.; Wensel, T. G.: RGS9, a GTPase ac-

celerator

[15302] for phototransduction. Neuron 20: 95-102, 1998.

[15303] 1249.

[15304] Makino, E. R.; Handy, J. W.; Li, T.; Arshavsky, V. Y.: The

GTPase

[15305] activating factor for transducin in rod photoreceptors is

the complex

[15306] between RGS9 and type 5 G protein beta subunit. Proc.

Nat. Acad.

[15307] Sci. 96: 1947-1952, 1999.

[15308] 1250.

[15309] Goto, K.; Kondo, H.: Molecular cloning and expression of a

90-kDa

[15310] diacylglycerol kinase that predominantly localizes in neu-

rons. Proc.

[15311] Nat. Acad. Sci. 90: 7598-7602, 1993.

[15312] 1251.

[15313] Ding, L.; Traer, E.; McIntyre, T. M.; Zimmerman, G. A.;

Prescott,

[15314] S. M.: The cloning and characterization of a novel human

diacylglycerol
[15315] kinase, DGK-iota. J. Biol. Chem. 273: 32746-32752,
1998.
[15316] 1252.

[15317] Yoshida, K.; Yoshitomo-Nakagawa, K.; Seki, N.; Sasaki, M.;
Sugano,
[15318] S.: Cloning, expression analysis, and chromosomal local-
ization of
[15319] BH-protocadherin (PCDH7), a novel member of the cad-
herin superfamily. Genomics 49:
[15320] 458-461, 1998.

[15321] 1253.

[15322] Nothwang, H. G.; Kim, H. G.; Aoki, J.; Geisterfer, M.;
Kubart,
[15323] S.; Wegner, R. D.; van Moers, A.; Ashworth, L. K.; Haaf, T.;
Bell,
[15324] J.; Arai, H.; Tommerup, N.; Ropers, H. H.; Wirth, J.: Func-
tional
[15325] hemizygosity of PAFAH1B3 due to a PAFAH1B3-CLK2 fu-
sion gene in a female
[15326] with mental retardation, ataxia and atrophy of the brain.
Hum. Molec.

[15327] Genet. 10: 797-806, 2001.

[15328] 1254.

[15329] Katsanis, N.; Fisher, E. M. C.: Identification, expression, and

[15330] chromosomal localization of ubiquitin conjugating enzyme 7 (UBE2G2),

[15331] a human homologue of the Saccharomyces cerevisiae Ubc7 gene. Genomics 51:

[15332] 128-131, 1998.

[15333] 1255.

[15334] Rose, S. A.; Leek, J. P.; Moynihan, T. P.; Ardley, H. C.; Markham,

[15335] A. F.; Robinson, P. A.: Assignment of the ubiquitin conjugating enzyme

[15336] gene, UBE2G2, to human chromosome band 21q22.3 by in situ hybridization. Cytogenet.

[15337] Cell Genet. 83: 98-99, 1998.

[15338] 1256.

[15339] Beisswanger, R.; Corbeil, D.; Vannier, C.; Thiele, C.; Dohrmann,

[15340] U.; Kellner, R.; Ashman, K.; Niehrs, C.; Huttner, W. B.: Ex-

istence

[15341] of distinct tyrosylprotein sulfotransferase genes: molecu-
lar characterization

[15342] of tyrosylprotein sulfotransferase-2. Proc. Nat. Acad. Sci.
95:

[15343] 11134-11139, 1998.

[15344] 1257.

[15345] Ouyang, Y.-B.; Lane, W. S.; Moore, K. L.: Tyrosylprotein
sulfotransferase:

[15346] purification and molecular cloning of an enzyme that cat-
alyzes tyrosine

[15347] O-sulfation, a common posttranslational modification of
eukaryotic

[15348] proteins Proc. Nat. Acad. Sci. 95: 2896-2901, 1998.

[15349] 1258.

[15350] Ju, Y.-T.; Chang, A. C. Y.; She, B.-R.; Tsaur, M.-L.; Hwang,
H.-M.;

[15351] Chao, C. C.-K.; Cohen, S. N.; Lin-Chao, S.: Gas7: a gene
expressed

[15352] preferentially in growth-arrested fibroblasts and termi-
nally differentiated

[15353] Purkinje neurons affects neurite formation. Proc. Nat.

Acad. Sci. 95:
[15354] 11423-11428, 1998.

[15355] 1259.

[15356] Kurtz, A.; Zimmer, A.: Interspecies fluorescence in situ hybridization
[15357] further defines synteny homology between mouse chromosome 11 and human
[15358] chromosome 17. Mammalian Genome 6: 379-380, 1995.

[15359] 1260.

[15360] Megonigal, M. D.; Cheung, N.-K. V.; Rappaport, E. F.; Nowell, P.
[15361] C.; Wilson, R. B.; Jones, D. H.; Addya, K.; Leonard, D. G. B.; Kushner,
[15362] B. H.; Williams, T. M.; Lange, B. J.; Felix, C. A.: Detection of
[15363] leukemia-associated MLL-GAS7 translocation early during chemotherapy
[15364] with DNA topoisomerase II inhibitors. Proc. Nat. Acad. Sci. 97:
[15365] 2814-2819, 2000.

[15366] 1261.

[15367] Chrast, R.; Scott, H. S.; Chen, H.; Kudoh, J.; Rossier, C.;

Minoshima,
[15368] S.; Wang, Y.; Shimizu, N.; Antonarakis, S. E.: Cloning of two human
[15369] homologs of the Drosophila single-minded gene SIM1 on chromosome 6q
[15370] and SIM2 on 21q within the Down syndrome chromosomal region. Genome
[15371] Res. 7: 615-624, 1997.

[15372] 1262.

[15373] Holder, J. L., Jr.; Butte, N. F.; Zinn, A. R.: Profound obesity

[15374] associated with a balanced translocation that disrupts the SIM1 gene. Hum.
[15375] Molec. Genet. 9: 101-108, 2000.

[15376] 1263.

[15377] Michaud, J. L.; Boucher, F.; Melnyk, A.; Gauthier, F.; Goshu, E.;
[15378] Levy, E.; Mitchell, G. A.; Himms-Hagen, J.; Fan, C.-M.: Sim1 haploinsufficiency
[15379] causes hyperphagia, obesity and reduction of the par-aventricular nucleus
[15380] of the hypothalamus. Hum. Molec. Genet. 10: 1465-1473,

2001.

[15381] 1264.

[15382] Fan, C. M.; Kuwana, E.; Bulfone, A.; Fletcher, C. F.; Copeland,

[15383] N. G.; Jenkins, N. A.; Crews, S.; Martinez, S.; Puelles, L.; Rubenstein,

[15384] J. L.; Tessier-Lavigne, M.: Expression patterns of two murine homologs

[15385] of Drosophila single-minded suggest possible roles in embryonic patterning

[15386] and in the pathogenesis of Down syndrome. Molec. Cell. Neurosci. 7:

[15387] 1-16, 1996.

[15388] 1265.

[15389] Duke-Cohan, J. S.; Gu, J.; McLaughlin, D. F.; Xu, Y.; Free-man,

[15390] G. J.; Schlossman, S. F.: Attractin (DPPT-L), a member of the CUB

[15391] family of cell adhesion and guidance proteins, is secreted by activated

[15392] human T lymphocytes and modulates immune cell inter-actions. Proc.

[15393] Nat. Acad. Sci. 95: 11336-11341, 1998.

[15394] 1266.

[15395] Gunn, T. M.; Miller, K. A.; He, L.; Hyman, R. W.; Davis, R. W.;

[15396] Azarani, A.; Schlessman, S. F.; Duke-Cohan, J. S.; Barsh, G. S.:

[15397] The mouse mahogany locus encodes a transmembrane form of human attractin. Nature 398:

[15398] 152-156, 1999.

[15399] 1267.

[15400] He, L.; Gunn, T. M.; Bouley, D. M.; Lu, X.-Y.; Watson, S. J.; Schlossman,

[15401] S. F.; Duke-Cohan, J. S.; Barsh, G. S.: A biochemical function for

[15402] attractin in agouti-induced pigmentation and obesity. Nature Genet. 27:

[15403] 40-47, 2001.

[15404] 1268.

[15405] Tang, W.; Gunn, T. M.; McLaughlin, D. F.; Barsh, G. S.; Schlossman,

[15406] S. F.; Duke-Cohan, J. S.: Secreted and membrane attractin

result
[15407] from alternative splicing of the human ATRN gene. Proc. Nat. Acad.
[15408] Sci. 97: 6025-6030, 2000.

[15409] 1269.

[15410] Fields, S.; Song, O.: A novel genetic system to detect protein-protein
[15411] interactions. (Letter) Nature 340: 245-246, 1989.

[15412] 1270.

[15413] Gordon, D. M.; Shi, Q.; Dancis, A.; Pain, D.: Maturation of frataxin
[15414] within mammalian and yeast mitochondria: one-step processing by matrix
[15415] processing peptidase. Hum. Molec. Genet. 8: 2255-2262, 1999.
[15416] 1271.

[15417] Koutnikova, H.; Campuzano, V.; Koenig, M.: Maturation of wild-type
[15418] and mutated frataxin by the mitochondrial processing peptidase. Hum.
[15419] Molec. Genet. 7: 1485-1489, 1998.

[15420] 1272.

[15421] Lewis, T. M.; Roberts, M. L.; Bretag, A. H.: Immunolabelling for

[15422] VDAC, the mitochondrial voltage-dependent anion channel, on sarcoplasmic

[15423] reticulum from amphibian skeletal muscle. Neurosci. Lett. 181: 83-86,

[15424] 1994.

[15425] 1273.

[15426] Kondo, I.; Ikeuchi, T.; Nishigaki, I.; Takita, H.; Fujiki, K.;

[15427] Takahashi, Y.; Hamaguchi, H.: Assignment of the gene for LCP1 on

[15428] chromosome 13. (Abstract) Cytogenet. Cell Genet. 40: 673 only, 1985.

[15429] 1274.

[15430] Spritz, R. A.: A study in scarlet. Nature Genet. 11: 225-226,

[15431] 1995.

[15432] 1275.

[15433] Xiao, H.; Neuveut, C.; Tiffany, H. L.; Benkirane, M.; Rich, E.

[15434] A.; Murphy, P. M.; Jeang, K.-T.: Selective CXCR4 antagonism by Tat:

[15435] implications for in vivo expansion of coreceptor use by HIV-1. Proc.

[15436] Nat. Acad. Sci. 97: 11466-11471, 2000.

[15437] 1276.

[15438] Akiyama, S.; Endo, H.; Inohara, N.; Ohta, S.; Kagawa, Y.: Gene

[15439] structure and cell type-specific expression of the human ATP synthase

[15440] alpha subunit. Biochim. Biophys. Acta 1219: 129-140, 1994.

[15441] 1277.

[15442] Godbout, R.; Pandita, A.; Beatty, B.; Bie, W.; Squire, J. A.:

[15443] Comparative genomic hybridization analysis of Y79 and FISH mapping

[15444] indicate the amplified human mitochondrial ATP synthase alpha-subunit

[15445] gene (ATP5A) maps to chromosome 18q12-q21. Cytogenet. Cell Genet. 77:

[15446] 253-256, 1997.

[15447] 1278.

[15448] Jabs, E. W.; Thomas, P. J.; Bernstein, M.; Coss, C.; Ferreira,

[15449] G. C.; Pedersen, P. L.: Chromosomal localization of genes required

[15450] for the terminal steps of oxidative metabolism: alpha and gamma subunits

[15451] of ATP synthase and the phosphate carrier. Hum. Genet. 93: 600-602,

[15452] 1994.

[15453] 1279.

[15454] Kataoka, H; Biswas, C.: Nucleotide sequence of a cDNA for the

[15455] alpha subunit of human mitochondrial ATP synthase. Biochim. Biophys.

[15456] Acta 1089: 393-395, 1991.

[15457] 1280.

[15458] Banfi, S.; Servadio, A.; Chung, M.; Kwiatkowski, T. J., Jr.; McCall,

[15459] A. E.; Duvick, L. A.; Shen, Y.; Roth, E. J.; Orr, H. T.; Zoghbi, H.

[15460] Y.: Identification and characterization of the gene causing

type
[15461] 1 spinocerebellar ataxia. Nature Genet. 7: 513-520, 1994.

[15462] 1281.

[15463] Banfi, S.; Servadio, A.; Chung, M.-y.; Capozzoli, F.; Duvick,
L.
[15464] A.; Elde, R.; Zoghbi, H. Y.; Orr, H. T.: Cloning and devel-
opmental
[15465] expression analysis of the murine homolog of the
spinocerebellar ataxia
[15466] type 1 gene (Sca1). Hum. Molec. Genet. 5: 33-40, 1996.

[15467] 1282.

[15468] Burright, E. N.; Clark, H. B.; Servadio, A.; Matilla, T.; Fed-
dersen,
[15469] R. M.; Yunis, W. S.; Duvick, L. A.; Zoghbi, H. Y.; Orr, H. T.:
SCA1
[15470] transgenic mice: a model for neurodegeneration caused
by an expanded
[15471] CAG trinucleotide repeat. Cell 82: 937-948, 1995.

[15472] 1283.

[15473] Chung, M.; Ranum, L. P. W.; Duvick, L. A.; Servadio, A.;
Zoghbi,

[15474] H. Y.; Orr, H. T.: Evidence for a mechanism predisposing to intergenerational

[15475] CAG repeat instability in spinocerebellar ataxia type 1. Nature Genet. 5:

[15476] 254-258, 1993.

[15477] 1284.

[15478] Giunti, P.; Sweeney, M. G.; Spadaro, M.; Jodice, C.; Novelletto,

[15479] A.; Malaspina, P.; Frontali, M.; Harding, A. E.: The trinucleotide

[15480] repeat expansion on chromosome 6p (SCA1) in autosomal dominant cerebellar

[15481] ataxias. Brain 117: 645-649, 1994.

[15482] 1285.

[15483] Raum, D.; Balner, H.; Petersen, B. H.; Alper, C. A.: Genetic

[15484] polymorphism of serum complement components in the chimpanzee. Immunogenetics 10:

[15485] 455-468, 1980.

[15486] 1286.

[15487] Raum, D.; Glass, D.; Carpenter, C. B.; Alper, C. A.; Schur, P.

[15488] H.: The chromosomal order of genes controlling the major histocompatibility

[15489] complex, properdin factor B, and deficiency of the second component

[15490] of complement. J. Clin. Invest. 58: 1240-1248, 1976.

[15491] 1287.

[15492] Raum, D.; Glass, D.; Carpenter, C. B.; Schur, P. H.; Alper, C.

[15493] A.: Mapping for the structural gene for the second component of complement

[15494] with respect to the human major histocompatibility complex. Am. J.

[15495] Hum. Genet. 31: 35-41, 1979.

[15496] 1288.

[15497] Caturegli, P.; Hejazi, M.; Suzuki, K.; Dohan, O.; Carrasco, N.;

[15498] Kohn, L. D.; Rose, N. R.: Hypothyroidism in transgenic mice expressing

[15499] IFN-gamma in the thyroid. Proc. Nat. Acad. Sci. 97: 1719-1724, 2000.

[15500] 1289.

[15501] Kondo, I.; Shin, K.; Honmura, S.; Nakajima, H.; Yamamura, E.;

[15502] Satoh, H.; Terauchi, M.; Usuki, Y.; Takita, H.; Hamaguchi, H.: A

[15503] case report of a patient with retinoblastoma and chromosome 13q deletion:

[15504] assignment of a new gene (gene for LCP1) on human chromosome 13. Hum.

[15505] Genet. 71: 263-266, 1985.

[15506] 1290.

[15507] Lin, C.-S.; Aebersold, R. H.; Kent, S. B.; Varma, M.; Leavitt,

[15508] J.: Molecular cloning and characterization of plastin, a human leukocyte

[15509] protein expressed in transformed human fibroblasts. Molec. Cell.

[15510] Biol. 8: 4659-4668, 1988.

[15511] 1291.

[15512] Lin, C.-S.; Chang, C.-H.; Huynh, T.: The murine L-plastin gene

[15513] promoter: identification and comparison with the human L-plastin gene

[15514] promoter. DNA Cell Biol. 16: 9-16, 1997.

[15515] 1292.

[15516] Lin, C.-S.; Park, T.; Chen, Z. P.; Leavitt, J.: Human plastin

[15517] genes: comparative gene structure, chromosome location,
and differential

[15518] expression in normal and neoplastic cells. J. Biol. Chem.
268: 2781-2792,

[15519] 1993.

[15520] 1293.

[15521] McConkey, E. H.; Taylor, B. J.; Phan, D.: Human heterozy-
gosity:

[15522] a new estimate. Proc. Nat. Acad. Sci. 76: 6500-6504,
1979.

[15523] 1294.

[15524] Murayama, N.; Tanaka, Y.; Hanyu, M.; Kobayashi, K.; Ham-
aguchi,

[15525] H.; Kondo, I.: Assignment of l-plastin to 13q14.3.
(Abstract) Human

[15526] Genome Mapping Workshop 93 27 only, 1993.

[15527] 1295.

[15528] O'Farrell, P. Z.; Goodman, H. M.; O'Farrell, P. H.: High resolution

[15529] two-dimensional electrophoresis of basic as well as acidic proteins. Cell 12:

[15530] 1133-1142, 1977.

[15531] 1296.

[15532] Walton, K. E.; Steyer, D.; Gruenstein, E. I.: Genetic polymorphism

[15533] in normal human fibroblasts as analyzed by two-dimensional polyacrylamide

[15534] gel electrophoresis. J. Biol. Chem. 254: 7951-7960, 1979.

[15535] 1297.

[15536] Zu, Y.; Kohno, M.; Kubota, I.; Nishida, E.; Hanaoka, M.; Namba,

[15537] Y.: Characterization of interleukin 2 stimulated 65-kilodalton phosphoprotein

[15538] in human T cells. Biochemistry 29: 1055-1062, 1990.

[15539] 1298.

[15540] Zu, Y.; Shigesada, K.; Nishida, E.; Kubota, I.; Kohno, M.; Hanaoka,

[15541] M.; Namba, Y.: 65-kilodalton protein phosphorylated by

interleukin

[15542] 2 stimulation bears two putative actin-binding sites and two calcium-binding

[15543] sites. Biochemistry 29: 8319-8324, 1990.

[15544] 1299.

[15545] Aggarwal, B. B.; Eessalu, T. E.; Hass, P. E.: Characterization

[15546] of receptors for human tumour necrosis factor and their regulation

[15547] by gamma-interferon. Nature 318: 665-667, 1985.

[15548] 1300.

[15549] Evans, A. M.; Petersen, J. W.; Sekhon, G. S.; DeMars, R.: Mapping

[15550] of prolactin and tumor necrosis factor-beta genes on human chromosome

[15551] 6p using lymphoblastoid cell deletion mutants. Somat. Cell Molec.

[15552] Genet. 15: 203-213, 1989.

[15553] 1301.

[15554] Camara, V. M.; Harding, J. W.; Prieur, D. J.: Inherited lysozyme

[15555] deficiency in rabbits: the absence of a primary isozyme of

lysozyme
[15556] as the cause of the condition. Lab. Invest. 63: 544-550, 1990.
[15557] 1302.

[15558] Canet, D.; Sunde, M.; Last, A. M.; Miranker, A.; Spencer, A.; Robinson,
[15559] C. V.; Dobson, C. M.: Mechanistic studies of the folding of human
[15560] lysozyme and the origin of amyloidogenic behavior in its disease-related
[15561] variants. Biochemistry 38: 6419-6427, 1999.

[15562] 1303.

[15563] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Lactalbumin
[15564] and Lysozyme. Washington: National Biomedical Research Foundation
[15565] (pub.) 5: 1972. Pp. D133-D140.

[15566] 1304.

[15567] Fleming, A.: On a remarkable bacteriolytic element found in tissues
[15568] and secretions. Proc. Roy. Soc. Ser. B. 93: 306-317, 1922.

[15569] 1305.

[15570] Fleming, A.; Allison, V. D.: Observations on a bacteriolytic substance

[15571] ('lysozyme') found in secretions and tissues. Brit. J. Exp. Path. 3:

[15572] 252-260, 1922.

[15573] 1306.

[15574] Greenwald, R. A.; Cantor, J. O.; Prieur, D. J.; Young, D. M.:

[15575] Composition of cartilage from lysozyme-deficient rabbits. Biochim.

[15576] Biophys. Acta 385: 435-437, 1975.

[15577] 1307.

[15578] Neufeld, E. L.: Personal Communication. Bethesda, Maryland

[15579] 1972.

[15580] 1308.

[15581] Pepys, M. B.; Hawkins, P. N.; Booth, D. R.; Vigushin, D. M.; Tennent,

[15582] G. A.; Soutar, A. K.; Totty, N.; Nguyen, O.; Blake, C. C. F.; Terry,

[15583] C. J.; Feest, T. G.; Zalin, A. M.; Hsuan, J. J.: Human lysozyme gene
[15584] mutations cause hereditary systemic amyloidosis. Nature 362: 553-557,
[15585] 1993.

[15586] 1309.

[15587] Peters, C. W. B.; Kruse, U.; Pollwein, R.; Grzeschik, K.-H.; Sippel,
[15588] A. E.: The human lysozyme gene: sequence organization and chromosomal
[15589] localization. (Abstract) Cytogenet. Cell Genet. 51: 1059 only, 1989.
[15590] 1310.

[15591] Prieur, D. J.: Personal Communication. Pullman, Washing-ton
[15592] 5/13/1975.

[15593] 1311.

[15594] Prieur, D. J.; Olson, H. M.; Young, D. M.: Lysozyme defi-ciency--an
[15595] inherited disorder of rabbits. Am. J. Path. 77: 283-296, 1974.

[15596] 1312.

[15597] Spitznagel, J. K.; Cooper, M. R.; McCall, A. E.; DeChatelet, L.

[15598] R.; Welsh, I. R.: Selective deficiency of granules associated with

[15599] lysozyme and lactoferrin in human polymorphs (PMN) with reduced microbicidal

[15600] capacity. (Abstract) J. Clin. Invest. 51: 93A only, 1972.

[15601] 1313.

[15602] Ku, C. C.; Murakami, M.; Sakamoto, A.; Kappler, J.; Marrack, P.

[15603] : Control of homeostasis of CD8+ memory T cells by opposing cytokines. Science 288:

[15604] 675-678, 2000.

[15605] 1314.

[15606] Leonard, W. J.; Donlon, T. A.; Lebo, R. V.; Greene, W. C.: Localization

[15607] of the gene encoding the human interleukin-2 receptor on chromosome

[15608] 10. Science 228: 1547-1549, 1985.

[15609] 1315.

[15610] Webb, G. C.; Campbell, H. D.; Lee, J. S.; Young, I. G.: Mapping
[15611] the gene for murine T-cell growth factor, Il-2, to bands B-C on chromosome
[15612] 3 and for the alpha chain of the IL2-receptor, Il-2ra, to bands A2-A3
[15613] on chromosome 2. Cytogenet. Cell Genet. 54: 164-168, 1990.
[15614] 1316.

[15615] Du, X.; Williams, D. A.: Interleukin-11: review of molecular,
[15616] cell biology, and clinical use. Blood 89: 3897-3908, 1997.

[15617] 1317.

[15618] Du, X. X.; Williams, D. A.: Interleukin-11: a multifunctional
[15619] growth factor derived from the hematopoietic microenvironment. Blood 83:
[15620] 2023-2030, 1994.

[15621] 1318.

[15622] McKinley, D.; Wu, Q.; Yang-Feng, T.; Yang, Y.-C.: Genomic sequence
[15623] and chromosomal location of human interleukin-11 gene

(IL11). Genomics 13:

[15624] 814-819, 1992.

[15625] 1319.

[15626] Paul, S. R.; Bennett, F.; Calvetti, J. A.; Kelleher, K.; Wood,

[15627] C. R.; O'Hara, R. M., Jr.; Leary, A. C.; Sibley, B.; Clark, S. C.;

[15628] Williams, D. A.; Yang, Y.-C.: Molecular cloning of a cDNA encoding

[15629] interleukin 11, a stromal cell-derived lymphopoietic and hematopoietic

[15630] cytokine. Proc. Nat. Acad. Sci. 87: 7512-7516, 1990.

[15631] 1320.

[15632] Yang-Feng, T. L.; Gibson, L.; Yang, Y. C.: Assignment of the gene

[15633] encoding human interleukin-11 to chromosome 19q13.3-q13.4. (Abstract) Cytogenet.

[15634] Cell Genet. 58: 2027 only, 1991.

[15635] 1321.

[15636] Grunig, G.; Warnock, M.; Wakil, A. E.; Venkayya, R.; Brom-

bacher,

[15637] F.; Rennick, D. M.; Sheppard, D.; Mohrs, M.; Donaldson, D.

D.; Locksley,

[15638] R. M.; Corry, D. B. Requirement for IL-13 independently of IL-4

[15639] in experimental asthma. Science 282: 2261-2263, 1998.

[15640] 1322.

[15641] Heinzmann, H.; Mao, X.-Q.; Akaiwa, M.; Kreomer, R. T.; Gao, P.-S.;

[15642] Ohshima, K.; Umeshita, R.; Abe, Y.; Braun, S.; Yamashita, T.; Roberts,

[15643] M. H.; Sugimoto, R.; and 20 others: Genetic variants of IL-13 signalling

[15644] and human asthma and atopy. Hum. Molec. Genet. 9: 549-559, 2000.

[15645] 1323.

[15646] Howard, T. D.; Koppelman, G. H.; Xu, J.; Zheng, S. L.; Postma,

[15647] D. S.; Meyers, D. A.; Bleecker, E. R.: Gene-gene interaction in asthma:

[15648] IL4RA and IL13 in a Dutch population with asthma. Am. J. Hum. Genet. 70:

[15649] 230-236, 2002.

[15650] 1324.

[15651] Howard, T. D.; Whittaker, P. A.; Zaiman, A. L.; Koppelman, G. H.;

[15652] Xu, J.; Hanley, M. T.; Meyers, D. A.; Postma, D. S.; Bleecker, E.

[15653] R.: Identification and association of polymorphisms in the interleukin-13

[15654] gene with asthma and atopy in a Dutch population. Am. J. Resp. Cell

[15655] Molec. Biol. 25: 377-384, 2001.

[15656] 1325.

[15657] Kuperman, D. A.; Huang, X.; Koth, L. L.; Chang, G. H.; Dolganov,

[15658] G. M.; Zhu, Z.; Elias, J. A.; Sheppard, D.; Erle, D. J.: Direct effects

[15659] of interleukin-13 on epithelial cells cause airway hyperreactivity

[15660] and mucus overproduction in asthma. Nature Med. 8: 885-889, 2002.

[15661] 1326.

[15662] Burshtyn, D. N.; Scharenberg, A. M.; Wagtmann, N.; Rajagopalan,

[15663] S.; Berrada, K.; Yi, T.; Kinet, J.-P.; Long, E. O.: Recruitment

of

[15664] tyrosine phosphatase HCP by the killer cell inhibitor receptor. Immunity 4:

[15665] 77-85, 1996.

[15666] 1327.

[15667] Dean, F. B.; Lian, L.; O'Donnell, M.: cDNA cloning and gene mapping

[15668] of human homologs for Schizosaccharomyces pombe rad17, rad1, and hus1

[15669] and cloning of homologs from mouse, Caenorhabditis elegans, and Drosophila

[15670] melanogaster. Genomics 54: 424-436, 1998.

[15671] 1328.

[15672] Macdonald, D. H. C.; Lahiri, D.; Sampath, A.; Chase, A.; Sohal,

[15673] J.; Cross, N. C. P.: Cloning and characterization of RNF6, a novel

[15674] RING finger gene mapping to 13q12. Genomics 58: 94-97, 1999.

[15675] 1329.

[15676] Strobel, M. C.; Seperack, P. K.; Copeland, N. G.; Jenkins, N.

[15677] A.: Molecular analysis of two mouse dilute locus deletion mutations:
[15678] spontaneous dilute lethal-20J and radiation-induced dilute prenatal
[15679] lethal Aa2 alleles. Molec. Cell. Biol. 10: 501-509, 1990.

[15680] 1330.

[15681] Tanaka, H.; Homma, K.; Iwane, A. H.; Katayama, E.; Ikebe, R.;
[15682] Saito, J.; Yanagida, T.; Ikebe, M.: The motor domain determines the
[15683] large step of myosin-V. Nature 415: 192-195, 2002.

[15684] 1331.

[15685] Walker, M. L.; Burgess, S. A.; Sellers, J. R.; Wang, F.; Hammer,
[15686] J. A., III; Trinick, J.; Knight, P. J.: Two-headed binding of a processive
[15687] myosin to F-actin. Nature 405: 804-807, 2000.

[15688] 1332.

[15689] Yanagida, T.; Iwane, A. H.: A large step for myosin. Proc. Nat.
[15690] Acad. Sci. 97: 9357-9359, 2000.

[15691] 1333.

[15692] Poetter, K.; Jiang, H.; Hassanzadeh, S.; Master, S. R.; Chang,

[15693] A.; Dalakas, M. C.; Rayment, I.; Sellers, J. R.; Fananapazir, L.;

[15694] Epstein, N. D.: Mutations in either the essential or regulatory light

[15695] chains of myosin are associated with a rare myopathy in human heart

[15696] and skeletal muscle. Nature Genet. 13: 63-69, 1996.

[15697] 1334.

[15698] Darras, B. T.; Fodor, B.; Vanin, E.; Francke, U.: A human myosin

[15699] alkali light chain gene mapped to chromosome 3. (Abstract) Cytogenet.

[15700] Cell Genet. 46: 603, 1987.

[15701] 1335.

[15702] Fodor, W. L.; Darras, B.; Seharaseyon, J.; Falkenthal, S.; Francke,

[15703] U.; Vanin, E. F.: Human ventricular/slow twitch myosin alkali light

[15704] chain gene characterization, sequence, and chromosomal location. J.

[15705] Biol. Chem. 264: 2143-2149, 1989.

[15706] 1336.

[15707] Ho, G.; Chisholm, R. L.: Substitution mutations in the myosin

[15708] essential light chain lead to reduced actin-activated AT-Pase activity

[15709] despite stoichiometric binding to the heavy chain. J. Biol. Chem. 272:

[15710] 4522-4527, 1997.

[15711] 1337.

[15712] Hoffmann, E.; Shi, Q. W.; Floroff, M.; Mickle, D. A. G.; Wu, T.-W.;

[15713] Olley, P. M.; Jackowski, G.: Molecular cloning and com-plete nucleotide

[15714] sequence of a human ventricular myosin light chain 1. Nucleic Acids

[15715] Res. 16: 2353, 1988.

[15716] 1338.

[15717] Laugwitz, K.-L.; Moretti, A.; Weig, H.-J.; Gillitzer, A.;

Pinkernell,

[15718] K.; Ott, T.; Pragst, I.; Stadele, C.; Seyfarth, M.; Schomig, A.; Ungerer,

[15719] M.: Blocking caspase-activated apoptosis improves contractility in

[15720] failing myocardium. Human Gene Ther. 12: 2051-2063, 2001.

[15721] 1339.

[15722] Moretti, A.; Weig, H.-J.; Ott, T.; Seyfarth, M.; Holthoff, H.-P.;

[15723] Grewe, D.; Gillitzer, A.; Bott-Flugel, L.; Schomig, A.; Ungerer, M.;

[15724] Laugwitz, K.-L.: Essential myosin light chain as a target for caspase-3

[15725] in failing myocardium. Proc. Nat. Acad. Sci. 99: 11860-11865, 2002.

[15726] 1340.

[15727] Olson, T. M.; Karst, M. L.; Whitby, F. G.; Driscoll, D. J.: Myosin

[15728] light chain mutation causes autosomal recessive cardiomyopathy with

[15729] mid-cavitary hypertrophy and restrictive physiology. Cir-

culation 105:

[15730] 2337-2340, 2002.

[15731] 1341.

[15732] Weber, F. E.; Vaughan, K. T.; Reinach, F. C.; Fischman, D. A.:

[15733] Complete sequence of human fast-type and slow-type muscle myosin-binding-protein

[15734] C (MyBP-C): differential expression, conserved domain structure and

[15735] chromosome assignment. Europ. J. Biochem. 216: 661-669, 1993.

[15736] 1342.

[15737] Pujana, M. A.; Corral, J.; Gratacos, M.; Combarros, O.; Berciano,

[15738] J.; Genis, D.; Banchs, I.; Estivill, X.; Volpini, V.; Ataxia Study

[15739] Group: Spinocerebellar ataxias in Spanish patients: genetic analysis

[15740] of familial and sporadic cases. Hum. Genet. 104: 516-522, 1999.

[15741] 1343.

[15742] Ramesar, R. S.; Bardien, S.; Beighton, P.; Bryer, A.: Ex-

panded
[15743] CAG repeats in spinocerebellar ataxia (SCA1) segregate with distinct
[15744] haplotypes in South African families. Hum. Genet. 100: 131-137,
[15745] 1997.

[15746] 1344.

[15747] Ranum, L. P. W.; Chung, M.; Banfi, S.; Bryer, A.; Schut, L. J.;

[15748] Ramesar, R.; Duvick, L. A.; McCall, A.; Subramony, S. H.; Goldfarb,
[15749] L.; Gomez, C.; Sandkuijl, L. A.; Orr, H. T.; Zoghbi, H. Y.: Molecular
[15750] and clinical correlations in spinocerebellar ataxia type I: evidence
[15751] for familial effects on the age at onset. Am. J. Hum. Genet. 55:
[15752] 244-252, 1994.

[15753] 1345.

[15754] Ranum, L. P. W.; Lundgren, J. K.; Schut, L. J.; Ahrens, M. J.;

[15755] Perlman, S.; Aita, J.; Bird, T. D.; Gomez, C.; Orr, H. T.: Spinocerebellar

[15756] ataxia type 1 and Machado-Joseph disease: incidence of CAG expansions

[15757] among adult-onset ataxia patients from 311 families with dominant,

[15758] recessive, or sporadic ataxia. Am. J. Hum. Genet. 57: 603-608, 1995.

[15759] 1346.

[15760] Servadio, A.; Koshy, B.; Armstrong, D.; Antalffy, B.; Orr, H.

[15761] T.; Zoghbi, H. Y.: Expression analysis of the ataxin-1 protein in

[15762] tissues from normal and spinocerebellar ataxia type 1 individuals. Nature

[15763] Genet. 10: 94-98, 1995.

[15764] 1347.

[15765] Storey, E.; du Sart, D.; Shaw, J. H.; Lorentzos, P.; Kelly, L.;

[15766] Gardner, R. J. M.; Forrest, S. M.; Biros, I.; Nicholson, G. A.: Frequency

[15767] of spinocerebellar ataxia types 1, 2, 3, 6, and 7 in Australian patients

[15768] with spinocerebellar ataxia. Am. J. Med. Genet. 95: 351-357, 2000.

[15769] 1348.

[15770] Suzuki, Y.; Sasaki, H.; Wakisaka, A.; Takada, A.; Yoshiki, T.;

[15771] Iwabuchi, K.; Tashiro, K.; Fukazawa, T.; Hamada, T.: Spinocerebellar

[15772] ataxia 1 (SCA1) in the Japanese: analysis of CAG trinucle-itide (sic)

[15773] repeat expansion and instability of the repeat for paternal transmission. Jpn.

[15774] J. Hum. Genet. 40: 131-143, 1995.

[15775] 1349.

[15776] Volz, A.; Boyle, J. M.; Cann, H. M.; Cottingham, R. W.; Orr, H.

[15777] T.; Ziegler, A.: Report of the second international work-shop on human

[15778] chromosome 6. Genomics 21: 464, 1994.

[15779] 1350.

[15780] Wakisaka, A.; Sasaki, H.; Takada, A.; Fukazawa, T.; Suzuki, Y.;

[15781] Hamada, T.; Iwabuchi, K.; Tashiro, K.; Yoshiki, T.: Spinocerebellar

[15782] ataxia 1 (SCA1) in the Japanese in Hokkaido may derive from a single

[15783] common ancestry. J. Med. Genet. 32: 590-592, 1995.

[15784] 1351.

[15785] Wang, S.; Detera-Wadleigh, S. D.; Coon, H.; Sun, C.; Goldin, L.

[15786] R.; Duffy, D. L.; Byerley, W. F.; Gershon, E. S.; Diehl, S. R.: Evidence

[15787] of linkage disequilibrium between schizophrenia and the SCA1 CAG repeat

[15788] on chromosome 6p23. (Letter) Am. J. Hum. Genet. 59: 731-736, 1996.

[15789] 1352.

[15790] Dionne, C. A.; Kaplan, R.; Seuanez, H.; O'Brien, S. J.; Jaye, M.

[15791] : Chromosome assignment by polymerase chain reaction techniques: assignment

[15792] of the oncogene FGF-5 to human chromosome 4. Biotech-niques 8: 190-194,

[15793] 1990.

[15794] 1353.

[15795] Hebert, J. M.; Rosenquist, T.; Gotz, J.; Martin, G. R.: FGF5 as
[15796] a regulator of the hair growth cycle: evidence from targeted and spontaneous
[15797] mutations. Cell 78: 1017-1025, 1994.

[15798] 1354.

[15799] Nguyen, C.; Roux, D.; Mattei, M.-G.; de Lapeyriere, O.; Goldfarb,
[15800] M.; Birnbaum, D.; Jordan, B. R.: The FGF-related oncogenes hst and
[15801] int.2, and the bcl.1 locus are contained within one megabase in band
[15802] q13 of chromosome 11, while the fgf.5 oncogene maps to 4q21. Oncogene 3:
[15803] 703-708, 1988.

[15804] 1355.

[15805] Zhan, X.; Bates, B.; Hu, X.; Goldfarb, M.: The human FGF-5 oncogene
[15806] encodes a novel protein related to fibroblast growth factors. Molec.
[15807] Cell. Biol. 8: 3487-3495, 1988.

[15808] 1356.

[15809] Kozak, C. A.; Filie, J.; Adamson, M. C.; Chen, Y.; Yu, L.: Murine

[15810] chromosomal location of the mu and kappa opioid receptor genes. Genomics 21:

[15811] 659-661, 1994.

[15812] 1357.

[15813] Bartoshuk, L. M.; Duffy, V. B.; Miller, I. J.: PTC/PROP tasting:

[15814] anatomy, psychophysics, and sex effects. Physiol. Behav. 56: 1165-1171,

[15815] 1994.

[15816] 1358.

[15817] Chautard-Freire-Maia, E. A.: Linkage relationships between 22

[15818] autosomal markers. Ann. Hum. Genet. 38: 191-198, 1974.

[15819] 1359.

[15820] Conneally, P. M.; Dumont-Driscoll, M.; Huntzinger, R. S.; Nance,

[15821] W. E.; Jackson, C. E.: Linkage relations of the loci for Kell

and

[15822] phenylthiocarbamide (PTC) taste sensitivity. Hum. Hered. 26: 267-271,

[15823] 1976.

[15824] 1360.

[15825] Spence, M. A.; Falk, C. T.; Neiswanger, K.; Field, L. L.; Marazita,

[15826] M. L.; Allen, F. H., Jr.; Siervogel, R. M.; Roche, A. F.; Cran-

dall,

[15827] B. F.; Sparkes, R. S.: Estimating the recombination fre-

quency for

[15828] the PTC-Kell linkage. Hum. Genet. 67: 183-186, 1984.

[15829] 1361.

[15830] Sadeghi-Nejad, A.; Senior, B.: Autosomal dominant trans-

mission

[15831] of isolated growth hormone deficiency in iris-dental dys-

plasia (Rieger's

[15832] syndrome). J. Pediat. 85: 644-648, 1974.

[15833] 1362.

[15834] Hing, A. V.; Helms, C.; Slaugh, R.; Burgess, A.; Wang, J. C.; Herman,

[15835] T.; Dowton, S. B.; Donis-Keller, H.: Linkage of preaxial polydactyly

[15836] type 2 to 7q36. Am. J. Med. Genet. 58: 128-135, 1995.

[15837] 1363.

[15838] Kinzler, K. W.; Nilbert, M. C.; Vogelstein, B.; Bryan, T. M.; Levy,

[15839] D. B.; Smith, K. J.; Preisinger, A. C.; Hamilton, S. R.; Hedge, P.;

[15840] Markham, A.; Carlson, M.; Joslyn, G.; Groden, J.; White, R.; Miki,

[15841] Y.; Miyoshi, Y.; Nishisho, I.; Nakamura, Y.: Identification of a

[15842] gene located at chromosome 5q21 that is mutated in colorectal cancers.

[15843] Science 251: 1366-1370, 1991.

[15844] 1364.

[15845] Constantinou, C. D.; Nielsen, K. B.; Prockop, D. J.: A lethal

[15846] variant of osteogenesis imperfecta has a single base mutation that

[15847] substitutes cysteine for glycine 904 of the alpha-1(I) chain of type

[15848] I procollagen: the asymptomatic mother has an unidenti-
fied mutation
[15849] producing an overmodified and unstable type I procolla-
gen. J. Clin.
[15850] Invest. 83: 574-584, 1989.

[15851] 1365.

[15852] Laborda, J.; Sausville, E. A.; Hoffman, T.; Notario, V.: dlk,

[15853] a putative mammalian homeotic gene differentially ex-
pressed in small
[15854] cell lung carcinoma and neuroendocrine tumor cell line. J.
Biol.
[15855] Chem. 268: 3817-3820, 1993.

[15856] 1366.

[15857] Dryja, T. P.; Rapaport, J.; McGee, T. L.; Nork, T. M.;
Schwartz,
[15858] T. L.: Molecular etiology of low-penetrance retinoblas-
toma in two
[15859] pedigrees. Am. J. Hum. Genet. 52: 1122-1128, 1993.

[15860] 1367.

[15861] Dryja, T. P.; Rapaport, J. M.; Joyce, J. M.; Petersen, R. A.:

[15862] Molecular detection of deletions involving band q14 of chromosome
[15863] 13 in retinoblastomas. Proc. Nat. Acad. Sci. 83: 7391-7394, 1986.
[15864] 1368.
[15865] Dryja, T. P.; Rapaport, J. M.; Weichselbaum, R.; Bruns, G. A.
[15866] P.: Chromosome 13 restriction fragment length polymorphisms. Hum.
[15867] Genet. 65: 320-324, 1984.
[15868] 1369.
[15869] Duane, T. B.: Clinical Ophthalmology. Hagerstown: Harper and
[15870] Row (pub.) 3: 1980. Pp. 13 only.
[15871] 1370.
[15872] Duncan, A. M. V.; Morgan, C.; Gallie, B. L.; Phillips, R. A.;
[15873] Squire, J.: Re-evaluation of the sublocalization of esterase D and
[15874] its relation to the retinoblastoma locus by in situ hybridization. Cytogenet.
[15875] Cell Genet. 44: 153-157, 1987.

[15876] 1371.

[15877] Dunn, J. M.; Phillips, R. A.; Becker, A. J.; Gallie, B. L.: Iden-
tification
[15878] of germline and somatic mutations affecting the
retinoblastoma gene. Science 241:
[15879] 1797-1800, 1988.

[15880] 1372.

[15881] Dunn, J. M.; Phillips, R. A.; Zhu, X.; Becker, A.; Gallie, B.

[15882] L.: Mutations in the RB1 gene and their effects on tran-
scription. Molec.
[15883] Cell. Biol. 9: 4596-4604, 1989.

[15884] 1373.

[15885] Ejima, Y.; Sasaki, M. S.; Kaneko, A.; Tanooka, H.: Types,
rates,
[15886] origin and expressivity of chromosome mutations involv-
ing 13q14 in
[15887] retinoblastoma patients. Hum. Genet. 79: 118-123, 1988.

[15888] 1374.

[15889] Ejima, Y.; Sasaki, M. S.; Kaneko, A.; Tanooka, H.; Hara, Y.;
Hida,

[15890] T.; Kinoshita, Y.: Possible inactivation of part of chromo-
some 13
[15891] due to 13qXp translocation associated with retinoblas-
toma. Clin.
[15892] Genet. 21: 357-361, 1982.

[15893] 1375.

[15894] Eldridge, R.; O'Meara, K.; Kitchin, D.: Superior intelligence

[15895] in sighted retinoblastoma patients and their families. J.
Med. Genet. 9:
[15896] 331-335, 1972.

[15897] 1376.

[15898] Falls, H. F.; Neel, J. V.: Genetics of retinoblastoma. Arch.

[15899] Ophthal. 46: 367-389, 1951.

[15900] 1377.

[15901] Fitzgerald, P. H.; Stewart, J.; Suckling, R. D.: Retinoblas-
toma
[15902] mutation rate in New Zealand and support for the two-hit
model. Hum.
[15903] Genet. 64: 128-130, 1983.

[15904] 1378.

[15905] Francke, U.: Retinoblastoma and chromosome 13. Cytogenet. Cell
[15906] Genet. 14: 131-134, 1976.

[15907] 1379.

[15908] Francois, J.: Retinoblastoma and osteogenic sarcoma. Ophthalmologica 175:
[15909] 185-191, 1977.

[15910] 1380.

[15911] Francois, J.: Hereditary malignant tumor of the eye. Congenital
[15912] Anomalies of The Eye. St. Louis: C. V. Mosby Co. (pub.) 1968.
[15913] Pp. 205-246.

[15914] 1381.

[15915] Francois, J.; Matton, M. T.; De Bie, S.; Tanaka, Y.; Vandenbulcke,
[15916] D.: Genesis and genetics of retinoblastoma. Ophthalmologica 170:
[15917] 405-425, 1975.

[15918] 1382.

[15919] Friend, S. H.; Bernards, R.; Rogelj, S.; Weinberg, R. A.; Ra-
paport,
[15920] J. M.; Albert, D. M.; Dryja, T. P.: A human DNA segment
with properties
[15921] of the gene that predisposes to retinoblastoma and os-
teosarcoma. Nature 323:
[15922] 643-646, 1986.

[15923] 1383.

[15924] Friend, S. H.; Dryja, T. P.; Weinberg, R. A.: Oncogenes and
tumor-suppressing
[15925] genes. New Eng. J. Med. 318: 618-622, 1988.

[15926] 1384.

[15927] Friend, S. H.; Horowitz, J. M.; Gerber, M. R.; Wang, X.-F.;
Bogenmann,
[15928] E.; Li, F. P.; Weinberg, R. A.: Deletions of a DNA sequence
in retinoblastomas
[15929] and mesenchymal tumors: organization of the sequence
and its encoded
[15930] protein. Proc. Nat. Acad. Sci. 84: 9059-9063, 1987. Note:
Correction:
[15931] Proc. Nat. Acad. Sci. 85: 2234 only, 1988..

[15932] 1385.

[15933] Fukushima, Y.; Kuroki, Y.; Ito, T.; Kondo, I.; Nishigaki, I.:

[15934] Familial retinoblastoma (mother and son) with 13q14 deletion. Hum.

[15935] Genet. 77: 104-107, 1987.

[15936] 1386.

[15937] Fung, Y.-K. T.; Murphree, A. L.; T'Ang, A.; Qian, J.; Hin-richs,

[15938] S. H.; Benedict, W. F.: Structural evidence for the authen-ticity

[15939] of the human retinoblastoma gene. Science 236: 1657-1661, 1987.

[15940] 1387.

[15941] Gallie, B. L.: Predictive testing for retinoblastoma comes of

[15942] age. (Editorial) Am. J. Hum. Genet. 61: 279-281, 1997.

[15943] 1388.

[15944] Gallie, B. L.; Ellsworth, R. M.; Abramson, D. M.; Phillips, R.

[15945] A.: Retinoma: spontaneous regression of retinoblastoma or benign

[15946] manifestation of the mutation? Brit. J. Cancer 45:

513-521, 1982.

[15947] 1389.

[15948] Gallie, B. L.; Phillips, R. A.: Multiple manifestations of the

[15949] retinoblastoma gene. Birth Defects Orig. Art. Ser. 18(6):
689-701,

[15950] 1982.

[15951] 1390.

[15952] Garcia-Cao, M.; Gonzalo, S.; Dean, D.; Blasco, M. A.: A role

[15953] for the Rb family of proteins in controlling telomere
length. Nature

[15954] Genet. 15Oct.: , 2002. Note: Advance Electronic Publica-
tion.

[15955] 1391.

[15956] Gey, W.: Dq-, multiple Missbildungen und Retinoblastom.
Humangenetik 10:

[15957] 362-365, 1970.

[15958] 1392.

[15959] Cotton, R. G. H.; Rodrigues, N. R.; Campbell, R. D.: Reac-
tivity

[15960] of cytosine and thymine in single-base-pair mismatches

with hydroxylamine

[15961] and osmium tetroxide and its application to the study of mutations. Proc.

[15962] Nat. Acad. Sci. 85: 4397-4401, 1988.

[15963] 1393.

[15964] Huang, Y. Z.; Wang, Q.; Xiong, W. C.; Mei, L.: Erbin is a protein

[15965] concentrated at postsynaptic membranes that interacts with PSD-95. J.

[15966] Biol. Chem. 276: 19318-19326, 2001.

[15967] 1394.

[15968] Aalto-Setala, K.; Helve, E.; Kovanen, P. T.; Kontula, K.: Finnish

[15969] type of low density lipoprotein receptor gene mutation (FH-Helsinki)

[15970] deletes exons encoding the carboxy-terminal part of the receptor and

[15971] creates an internalization-defective phenotype. J. Clin. In-vest. 84:

[15972] 499-505, 1989.

[15973] 1395.

[15974] Aalto-Setala, K.; Koivisto, U.-M.; Miettinen, T. A.; Gylling, H.;

[15975] Kesaniemi, Y. A.; Savolainen, M.; Pyorala, K.; Ebeling, T.; Mononen,

[15976] I.; Turtola, H.; Viikari, J.; Kontula, K.: Prevalence and geo-graphical

[15977] distribution of major LDL receptor gene rearrangements in Finland. J.

[15978] Intern. Med. 231: 227-234, 1992.

[15979] 1396.

[15980] Karibe, A.; Tobacman, L. S.; Strand, J.; Butters, C.; Back, N.;

[15981] Bachinski, L. L.; Arai, A. E.; Ortiz, A.; Roberts, R.; Homsher, E.;

[15982] Fananapazir, L.: Hypertrophic cardiomyopathy caused by a novel alpha-tropomyosin

[15983] mutation (V95A) is associated with mild cardiac pheno-type, abnormal

[15984] calcium binding to troponin, abnormal myosin cycling, and poor prognosis. Circulation 103:

[15985] 65-71, 2001.

[15986] 1397.

[15987] Nordenskjold, A.; Ivarsson, S.-A.: Molecular characterization
[15988] of 5-alpha-reductase type 2 deficiency and fertility in a Swedish
[15989] family. J. Clin. Endocr. Metab. 83: 3236-3238, 1998.

[15990] 1398.

[15991] Nordenskjold, A.; Magnus, O.; Aagenaes, O.; Knudtzon, J.: Homozygous
[15992] mutation (A228T) in the 5-alpha-reductase type 2 gene in a boy with
[15993] 5-alpha-reductase deficiency: genotype-phenotype correlations. Am.
[15994] J. Med. Genet. 80: 269-272, 1998.

[15995] 1399.

[15996] Opitz, J. M.; Simpson, J. L.; Sarto, G. E.; Summitt, R. L.; New,
[15997] M.; German, J.: Pseudovaginal perineoscrotal hypospadias. Clin.
[15998] Genet. 3: 1-26, 1972.

[15999] 1400.

[16000] Peterson, R. E.; Imperato-McGinley, J.; Gautier, T.; Sturla,

E.

[16001] : Male pseudohermaphroditism due to steroid 5-alpha-reductase deficiency. Am.

[16002] J. Med. 62: 170-191, 1977.

[16003] 1401.

[16004] Pinsky, L.; Kaufman, M.; Straisfeld, C.; Zilahi, B.; Hall, C.

[16005] S.-G.: 5-Alpha-reductase activity of genital and nongenital skin

[16006] fibroblasts from patients with 5-alpha-reductase deficiency, androgen

[16007] insensitivity, or unknown forms of pseudohermaphroditism. Am. J.

[16008] Med. Genet. 1: 407-416, 1978.

[16009] 1402.

[16010] Price, P.; Wass, J. A. H.; Griffin, J. E.; Leshin, M.; Savage,

[16011] M. O.; Large, D. M.; Bu'Lock, D. E.; Anderson, D. C.; Wilson, J. D.;

[16012] Besser, G. M.: High dose androgen therapy in male pseudohermaphroditism

[16013] due to 5-alpha-reductase deficiency and disorders of the androgen

[16014] receptor. J. Clin. Invest. 74: 1496-1508, 1984.

[16015] 1403.

[16016] Savage, M. O.; Preece, M. A.; Jeffcoate, S. L.; Ransley, P. G.;

[16017] Rumsby, G.; Mansfield, M. D.; Williams, D. I.: Familial male pseudohermaphroditism

[16018] due to deficiency of 5-alpha-reductase. Clin. Endocr. 12: 397-406,

[16019] 1980.

[16020] 1404.

[16021] Simpson, J. L.; New, M.; Peterson, R. E.; German, J.: Pseudovaginal

[16022] perineoscrotal hypospadias (PPSH) in sibs. Birth Defects Orig. Art.

[16023] Ser. VII(6): 140-144, 1971.

[16024] 1405.

[16025] Soderstrom, T. G.; Bjelfman, C.; Brekkan, E.; Ask, B.; Egevad,

[16026] L.; Norlen, B. J.; Rane, A.: Messenger ribonucleic acid levels of

[16027] steroid 5-alpha-reductase 2 in human prostate predict the enzyme activity. J.

[16028] Clin. Endocr. Metab. 86: 855-858, 2001.

[16029] 1406.

[16030] Thigpen, A. E.; Davis, D. L.; Gautier, T.; Imperato-McGin-
ley,
[16031] J.; Russell, D. W.: The molecular basis of steroid
5-alpha-reductase
[16032] deficiency in a large Dominican kindred. New Eng. J. Med.
327: 1216-1219,
[16033] 1992.

[16034] 1407.

[16035] Thigpen, A. E.; Davis, D. L.; Milatovich, A.; Mendonca, B.
B.;
[16036] Imperato-McGinley, J.; Griffin, J. E.; Francke, U.; Wilson, J.
D.;
[16037] Russell, D. W.: The molecular genetics of steroid
5-alpha-reductase
[16038] 2 deficiency. J. Clin. Invest. 90: 799-809, 1992.

[16039] 1408.

[16040] Vilchis, F.; Canto, P.; Chavez, B.; Ulloa-Aguirre, A.;
Mendez,
[16041] J. P.: Molecular analysis of the 5-alpha-steroid reductase

type 2

[16042] gene in a family with deficiency of the enzyme. Am. J. Med. Genet. 69:

[16043] 69-72, 1997.

[16044] 1409.

[16045] Walsh, P. C.; Madden, J. D.; Harrod, M. J.; Goldstein, J. L.;

[16046] MacDonald, P. C.; Wilson, J. D.: Familial incomplete male pseudohermaphroditism,

[16047] type 2: decreased dihydrotestosterone formation in pseudovaginal perineoscrotal

[16048] hypospadias. New Eng. J. Med. 291: 944-949, 1974.

[16049] 1410.

[16050] Wilson, J. D.: Personal Communication. Dallas, Texas 10/30/1981.

[16051] 1411.

[16052] Wilson, J. D.; Harrod, M. J.; Goldstein, J. L.; Hemsell, D. L.;

[16053] MacDonald, P. C.: Familial incomplete male pseudohermaphroditism

[16054] type I: evidence for androgen resistance in a family with the Reifenstein

[16055] syndrome. New Eng. J. Med. 290: 1097-1103, 1974.

[16056] 1412.

[16057] Albright, F.; Butler, A. M.; Bloomberg, E.: Rickets resistant

[16058] to vitamin D therapy. Am. J. Dis. Child. 54: 529-547, 1937.

[16059] 1413.

[16060] Bouchard, G.; Laberge, C.; Scriver, C. R.: La tyrosinemie hereditaire

[16061] et le rachitisme vitamino-dependant au Saguenay. Un. Med. Canada 114:

[16062] 633-636, 1985.

[16063] 1414.

[16064] Brenza, H. L.; Kimmel-Jehan, C.; Jehan, F.; Shinki, T.; Wakino,

[16065] S.; Anazawa, H.; Suda, T.; DeLuca, H. F.: Parathyroid hormone activation

[16066] of the 25-hydroxyvitamin D3-1-alpha-hydroxylase gene promoter. Proc.

[16067] Nat. Acad. Sci. 95: 1387-1391, 1998.

[16068] 1415.

[16069] Delvin, E. E.; Glorieux, F. H.; Marie, P. J.; Pettifor, J. M.:

[16070] Vitamin D dependency: replacement therapy with cal-
citriol. J. Pediat. 99:
[16071] 26-34, 1981.

[16072] 1416.

[16073] Dent, C. E.; Friedman, M.; Watson, L.: Hereditary pseudo-
vitamin
[16074] D deficiency rickets ('pseudo-mangelrachitis'). J. Bone
Joint Surg. 50B:
[16075] 708-719, 1968.

[16076] 1417.

[16077] Diaz, L.; Sanchez, I.; Avila, E.; Halhali, A.; Vilchis, F.; Lar-
rea,
[16078] F.: Identification of a 25-hydroxyvitamin D3
1-alpha-hydroxylase
[16079] gene transcription product in cultures of human syncy-
tiotrophoblast
[16080] cells. J. Clin. Endocr. Metab. 85: 2543-2549, 2000.

[16081] 1418.

[16082] Fraser, D.; Kooh, S. W.; Kind, H. P.; Holick, M. F.; Tanaka,
Y.;
[16083] DeLuca, H. F.: Pathogenesis of hereditary vitamin-

D-dependent rickets.

[16084] An inborn error of vitamin D metabolism involving defective conversion

[16085] of 25-hydroxyvitamin D to 1-alpha, 25-dihydroxyvitamin D. New Eng.

[16086] J. Med. 289: 817-822, 1973.

[16087] 1419.

[16088] Fraser, D.; Salter, R. B.: The diagnosis and management of the

[16089] various types of rickets. Pediat. Clin. N. Am. 5: 417-441, 1958.

[16090] 1420.

[16091] Lachmann, P. J.; Hobart, M. J.; Woo, P.: Combined genetic deficiency

[16092] of C6 and C7 in man. Clin. Exp. Immun. 33: 193-203, 1978.

[16093] 1421.

[16094] Whitehouse, D. B.: Genetic polymorphism and linkage of the sixth

[16095] and seventh complement components (C6 and C7) in the common marmoset. Biochem.

[16096] Genet. 22: 51-63, 1984.

[16097] 1422.

[16098] Adams, E. M.; Hustead, S.; Rubin, P.; Wagner, R.; Gewurz, A.; Graziano,

[16099] F. M.: Absence of the seventh component of complement in a patient

[16100] with chronic meningococcemia presenting as vasculitis. Ann. Intern.

[16101] Med. 99: 35-38, 1983.

[16102] 1423.

[16103] Boyer, J. T.; Gall, E. P.; Norman, M. E.; Nilsson, U. R.; Zim-merman,

[16104] T. S.: Hereditary deficiency of the seventh component of complement. J.

[16105] Clin. Invest. 56: 905-913, 1975.

[16106] 1424.

[16107] Clough, J. D.; Clough, M. L.; Weinstein, A.; Calabrese, L. H.;

[16108] Mansfield, L. R.; Gulick, P.; Gavin, T.; Braun, W. E.: Familial late

[16109] complement component (C6, C7) deficiency with chronic meningococcemia. Arch.

[16110] Intern. Med. 140: 929-933, 1980.

[16111] 1425.

[16112] Coto, E.; Martinez-Naves, E.; Dominguez, O.; Lopez-Lar-
rea, C.:

[16113] DNA polymorphism of the human complement component
C7 gene in familial

[16114] deficiencies. Hum. Genet. 85: 251-252, 1990.

[16115] 1426.

[16116] Delage, J. M.; Bergeron, P.; Simard, J.; Lehner-Netsch, G.;
Prochazka,

[16117] E.: Hereditary C7 deficiency: diagnosis and HLA studies in
a French-Canadian

[16118] family. J. Clin. Invest. 60: 1061-1069, 1977.

[16119] 1427.

[16120] Egan, L. J.; Orren, A.; Doherty, J.; Wurzner, R.; McCarthy,
C.

[16121] F.: Hereditary deficiency of the seventh component of
complement

[16122] and recurrent meningococcal infection: investigations of
an Irish

[16123] family using a novel haemolytic screening assay for com-

plement activity
[16124] and C7 M/N allotyping. Epidemiol. Infect. 113: 275-281, 1994.
[16125] 1428.

[16126] Eldridge, P. R.; Hobart, M. J.; Lachmann, P. J.: The genetics
[16127] of the sixth and seventh components of complement in the dog: polymorphism,
[16128] linkage, locus duplication, and silent alleles. Biochem. Genet. 21:
[16129] 81-91, 1983.

[16130] 1429.

[16131] Fernie, B. A.; Hobart, M. J.: Complement C7 deficiency: seven
[16132] further molecular defects and their associated marker haplotypes. Hum.
[16133] Genet. 103: 513-519, 1998.

[16134] 1430.

[16135] Fernie, B. A.; Orren, A.; Sheehan, G.; Schlesinger, M.; Ho-
[16136] bart, M. J.: Molecular bases of C7 deficiency: three different defects. J.

[16137] Immun. 159: 1019-1026, 1997.

[16138] 1431.

[16139] Wong, K.; Ren, X.-R.; Huang, Y.-Z.; Xie, Y.; Liu, G.; Saito, H.;

[16140] Tang, H.; Wen, L.; Brady-Kalnay, S. M.; Mei, L.; Wu, J. Y.; Xiong,

[16141] W.-C.; Rao, Y.: Signal transduction in neuronal migration: roles

[16142] of GTPase activating proteins and the small GTPase Cdc42 in the Slit-Robo

[16143] pathway. Cell 107: 209-221, 2001.

[16144] 1432.

[16145] Matesic, L. E.; Yip, R.; Reuss, A. E.; Swing, D. A.; O'Sullivan,

[16146] T. N.; Fletcher, C. F.; Copeland, N. G.; Jenkins, N. A.: Mu-tations

[16147] in Mlph, encoding a member of the Rab effector family, cause the melanosome

[16148] transport defects observed in leaden mice. Proc. Nat. Acad. Sci. 98:

[16149] 10238-10243, 2001.

[16150] 1433.

[16151] Silvers, W. K.: The Coat Colors of Mice. New York: Springer
,
[16152] 1979.

[16153] 1434.

[16154] Gardner, H. P.; Wertheim, G. B. W.; Ha, S. I.; Copeland, N.
G.;
[16155] Gilbert, D. J.; Jenkins, N. A.; Marquis, S. T.; Chodosh, L. A.:
Cloning
[16156] and characterization of Hunk, a novel mammalian
SNF1-related protein
[16157] kinase. Genomics 63: 46-59, 2000.

[16158] 1435.

[16159] Juwana, J.-P.; Henderikx, P.; Mischo, A.; Wadle, A.; Fadle,
N.;
[16160] Gerlach, K.; Arends, J. W.; Hoogenboom, H.; Pfreundschuh,
M.; Renner,
[16161] C.: EB/RP gene family encodes tubulin binding proteins.
Int. J.
[16162] Cancer 81: 275-284, 1999.

[16163] 1436.

[16164] Renner, C.; Pfitzenmeier, J.-P.; Gerlach, K.; Held, G.;

Ohnesorge,
[16165] S.; Sahin, U.; Bauer, S.; Pfreundschuh, M.: RP1, a new member of
[16166] the adenomatous polyposis coli-binding EB1-like gene family, is differently
[16167] expressed in activated T cells. J. Immun. 159: 1276-1283, 1997.
[16168] 1437.

[16169] Wadle, A.; Thiel, G.; Mischo, A.; Jung, V.; Pfreundschuh, M.; Renner,
[16170] C.: Chromosomal localization and promoter analysis of the adenomatous
[16171] polyposis coli binding protein RP1. Oncogene 20: 5920-5929, 2001.
[16172] 1438.

[16173] Iwasaki, K.; Toyonaga, R.: The Rab3 GDP/GTP exchange factor homolog
[16174] AEX-3 has a dual function in synaptic transmission. EMBO J. 19:
[16175] 4806-4816, 2000.

[16176] 1439.

[16177] Awasthi, S.; Cheng, J.; Singhal, S. S.; Saini, M. K.; Pandya,

U.;

[16178] Pikula, S.; Bandorowicz-Pikula, J.; Singh, S. V.; Zimniak, P.; Awasthi,

[16179] Y. C.: Novel function of human RLIP76: ATP-dependent transport of

[16180] glutathione conjugates and doxorubicin. Biochemistry 39: 9327-9334,

[16181] 2000.

[16182] 1440.

[16183] Jullien-Flores, V.; Dorseuil, O.; Romero, R.; Letourneur, F.; Saragosti,

[16184] S.; Berger, R.; Tavitian, A.; Gacon, G.; Camonis, J. H.: Bridging

[16185] Ral GTPase to Rho pathways: RLIP76, a Ral effector with CDC42/Rac

[16186] GTPase-activating protein activity. J. Biol. Chem. 270: 22473-22477,

[16187] 1995.

[16188] 1441.

[16189] Cacheux, V.; Dastot-Le Moal, F.; Kaariainen, H.; Bon-durand, N.;

[16190] Rintala, R.; Boissier, B.; Wilson, M.; Mowat, D.; Goossens,

M.: Loss-of-function
[16191] mutations in SIP1 Smad interacting protein 1 result in a syndromic
[16192] Hirschsprung disease. Hum. Molec. Genet. 10: 1503-1510, 2001.
[16193] 1442.

[16194] Comijn, J.; Berx, G.; Vermassen, P.; Verschueren, K.; van Grunsven,
[16195] L.; Bruyneel, E.; Mareel, M.; Huylebroeck, D.; van Roy, F.: The two-handed
[16196] E box binding zinc finger protein SIP1 downregulates E-cadherin and
[16197] induces invasion. Molec. Cell 7: 1267-1278, 2001.

[16198] 1443.

[16199] Remacle, J. E.; Kraft, H.; Lerchner, W.; Wuytens, G.; Collart,

[16200] C.; Verschueren, K.; Smith, J. C.; Huylebroeck, D.: New mode of DNA
[16201] binding of multi-zinc finger transcription factors: deltaEF1 family
[16202] members bind with two hands to two target sites. EMBO J. 18: 5073-5084,
[16203] 1999.

[16204] 1444.

[16205] Verschueren, K.; Remacle, J. E.; Collart, C.; Kraft, H.; Baker,

[16206] B. S.; Tylzanowski, P.; Nelles, L.; Wuytens, G.; Su, M.-T.; Bodmer,

[16207] R.; Smith, J. C.; Huylebroeck, D.: SIP1, a novel zing finger/ homeodomain

[16208] repressor, interacts with Smad proteins and binds to 5-prime-CACCT

[16209] sequences in candidate target genes. J. Biol. Chem. 274: 20489-20498,

[16210] 1999.

[16211] 1445.

[16212] Mas, C.; Bourgeois, F.; Bulfone, A.; Levacher, B.; Mugnier, C.;

[16213] Simonneau, M.: Cloning and expression analysis of a novel gene, RP42,

[16214] mapping to an autism susceptibility locus on 6q16. Ge- nomics 65:

[16215] 70-74, 2000.

[16216] 1446.

[16217] Faulkner, G.; Pallavicini, A.; Formentin, E.; Comelli, A.; Iev-

olella,

[16218] C.; Trevisan, S.; Bortoletto, G.; Scannapieco, P.; Salamon, M.; Mouly,

[16219] V.; Valle, G.; Lanfranchi, G.: ZASP: a new Z-band alternatively spliced

[16220] PDZ-motif protein. J. Cell Biol. 146: 465-475, 1999.

[16221] 1447.

[16222] Zhou, Q.; Ruiz-Lozano, P.; Martone, M. E.; Chen, J.: Cypher, a

[16223] striated muscle-restricted PDZ and LIM domain-containing protein,

[16224] binds to alpha-actinin-2 and protein kinase C. J. Biol. Chem. 274:

[16225] 19807-19813, 1999.

[16226] 1448.

[16227] Ben-Zeev, B.; Levy-Nissenbaum, E.; Lahat, H.; Anikster, Y.; Shinar,

[16228] Y.; Brand, N.; Gross-Tzur, V.; MacGregor, D.; Sidi, R.; Kleta, R.;

[16229] Frydman, M.; Pras, E.: Megalencephalic leukoencephalopathy with subcortical

[16230] cysts; a founder effect in Israeli patients and a higher than

expected
[16231] carrier rate among Libyan Jews. Hum. Genet. 111:
214-218, 2002.
[16232] 1449.

[16233] Leegwater, P. A. J.; Boor, P. K. I.; Yuan, B. Q.; van der
Steen,
[16234] J.; Visser, A.; Konst, A. A. M.; Oudejans, C. B. M.; Schut-
gens, R.
[16235] B. H.; Pronk, J. C.; van der Knaap, M. S.: Identification of
novel
[16236] mutations in MLC1 responsible for megalencephalic
leukoencephalopathy
[16237] with subcortical cysts. Hum. Genet. 110: 279-283, 2002.

[16238] 1450.

[16239] Rowe, P. S. N.; de Zoysa, P. A.; Dong, R.; Wang, H. R.;
White,
[16240] K. E.; Econs, M. J.; Oudet, C. L.: MEPE, a new gene ex-
pressed in
[16241] bone marrow and causing osteomalacia. Genomics 67:
54-68, 2000.
[16242] 1451.

[16243] Feng, S.-L. Y.; Guo, Y.; Factor, V. M.; Thorgeirsson, S. S.;

Bell,

[16244] D. W.; Testa, J. R.; Peifley, K. A.; Winkles, J. A.: The Fn14 immediate-early

[16245] response gene is induced during liver regeneration and highly expressed

[16246] in both human and murine hepatocellular carcinomas. Am. J. Path. 156:

[16247] 1253-1261, 2000.

[16248] 1452.

[16249] Meighan-Mantha, R. L.; Hsu, D. K. W.; Guo, Y.; Brown, S. A. N.;

[16250] Feng, S.-L. Y.; Peifley, K. A.; Alberts, G. F.; Copeland, N. G.; Gilbert,

[16251] D. J.; Jenkins, N. A.; Richards, C. M.; Winkles, J. A.: The mitogen-inducible

[16252] Fn14 gene encodes a type I transmembrane protein that modulates fibroblast

[16253] adhesion and migration. J. Biol. Chem. 274:

33166-33176, 1999.

[16254] 1453.

[16255] Ikeda, H.; Lethe, B.; Lehmann, F.; van Baren, N.; Baurain, J.-F.;

[16256] de Smet, C.; Chambost, H.; Vitale, M.; Moretta, A.; Boon, T.; Coulie,

[16257] P. G.: Characterization of an antigen that is recognized on a melanoma

[16258] showing partial HLA loss by CTL expressing an NK inhibitory receptor. Immunity 6:

[16259] 199-208, 1997.

[16260] 1454.

[16261] van Baren, N.; Chambost, H.; Ferrant, A.; Michaux, L.; Ikeda, H.;

[16262] Millard, I.; Olive, D.; Boon, T.; Coulie, P. G.: PRAME, a gene encoding

[16263] an antigen recognized on a human melanoma by cytolytic T cells, is

[16264] expressed in acute leukaemia cells. Brit. J. Haemat. 102: 1376-1379,

[16265] 1998.

[16266] 1455.

[16267] Burgener, R.; Wolf, M.; Ganz, T.; Baggiolini, M.: Purification

[16268] and characterization of a major phosphatidylserine-binding phosphoprotein

[16269] from human platelets. Biochem. J. 269: 729-734, 1990.

[16270] 1456.

[16271] Gustincich, S.; Schneider, C.: Serum deprivation response gene

[16272] is induced by serum starvation but not by contact inhibition. Cell

[16273] Growth Diff. 4: 753-760, 1993.

[16274] 1457.

[16275] Gustincich, S.; Vatta, P.; Goruppi, S.; Wolf, M.; Saccone, S.;

[16276] Della Valle, G.; Baggiolini, M.; Schneider, C.: The human serum deprivation

[16277] response gene (SDPR) maps to 2q32-q33 and codes for a phosphatidylserine-binding

[16278] protein. Genomics 57: 120-129, 1999.

[16279] 1458.

[16280] van Eenennaam, H.; Pruijn, G. J. M.; van Venrooij, W. J.: hPop4:

[16281] a new protein subunit of the human RNase MRP and RNase P ribonucleoprotein

[16282] complexes. Nucleic Acids Res. 27: 2465-2472, 1999.

[16283] 1459.

[16284] Eder, P. S.; Kekuda, R.; Stolc, V.; Altman, S.: Characterization

[16285] of two scleroderma autoimmune antigens that copurify with human ribonuclease

[16286] P. Proc. Nat. Acad. Sci. 94: 1101-1106, 1997.

[16287] 1460.

[16288] Huizing, M.; Anikster, Y.; Fitzpatrick, D. L.; Jeong, A. B.; D'Souza,

[16289] M.; Rausche, M.; Toro, J. R.; Kaiser-Kupfer, M. I.; White, J. G.;

[16290] Gahl, W. A.: Hermansky-Pudlak syndrome type 3 in Ashkenazi Jews and

[16291] other non-Puerto Rican patients with hypopigmentation and platelet

[16292] storage-pool deficiency. Am. J. Hum. Genet. 69: 1022-1032, 2001.

[16293] 1461.

[16294] Suzuki, T.; Li, W.; Zhang, Q.; Novak, E. K.; Sviderskaya, E. V.;

[16295] Wilson, A.; Bennett, D. C.; Roe, B. A.; Swank, R. T.; Spritz, R. A.

[16296] : The gene mutated in cocoa mice, carrying a defect of organelle biogenesis,
[16297] is a homologue of the human Hermansky-Pudlak syndrome-3 gene. Genomics 78:
[16298] 30-37, 2001.

[16299] 1462.

[16300] Bascom, R. A.; Liu, L.; Humphries, P.; Fishman, G. A.; Murray,
[16301] J. C.; McInnes, R. R.: Polymorphisms and rare sequence variants at
[16302] the ROM1 locus. Hum. Molec. Genet. 2: 1975-1977, 1993.

[16303] 1463.

[16304] Clarke, G.; Goldberg, A. F. X.; Vidgen, D.; Collins, L.; Ploder,
[16305] L.; Schwarz, L.; Molday, L. L.; Rossant, J.; Szel, A.; Molday, R.
[16306] S.; Birch, D. G.; McInnes, R. R.: Rom-1 is required for rod photoreceptor
[16307] viability and the regulation of disk morphogenesis. Nature Genet. 25:
[16308] 67-73, 2000.

[16309] 1464.

[16310] Kajiwara, K.; Berson, E. L.; Dryja, T. P.: Digenic retinitis pigmentosa

[16311] due to mutations at the unlinked peripherin/RDS and ROM1 loci. Science 264:

[16312] 1604-1608, 1994.

[16313] 1465.

[16314] Nadeau, J. H.: Personal Communication. Bar Harbor, Me. 10/19/1994.

[16315] 1466.

[16316] Nathans, J.: Personal Communication. Baltimore, Md. 12/6/1993.

[16317] 1467.

[16318] Stone, E. M.; Nichols, B. E.; McInnes, R. R.; Bascom, R. A.; Kimura,

[16319] A. E.; Streb, L. M.; Sheffield, V. C.: The gene for Best's disease

[16320] (vitelliform macular dystrophy) maps to chromosome 11q13. (Abstract) Am.

[16321] J. Hum. Genet. 51 (suppl.): A35 only, 1992.

[16322] 1468.

[16323] Taylor, B. A.; Phillips, S. J.; Bascom, R. A.; McInnes, R. R.

[16324] : The retinal outer segment membrane protein-1 gene (Rom1) maps to

[16325] the proximal end of mouse chromosome 19. Genomics 23: 510-511, 1994.

[16326] 1469.

[16327] Barton, D. E.; Spritz, R. A.; Francke, U.: RPU1 encoding the 68kDa

[16328] U1 snRNP-associated protein is located on chromosome 19. (Abstract) Cytogenet.

[16329] Cell Genet. 46: 577 only, 1987.

[16330] 1470.

[16331] Du, H.; Rosbash, M.: The U1 snRNP protein U1C recognizes the 5-prime

[16332] splice site in the absence of base pairing. Nature 419: 86-90, 2002.

[16333] 1471.

[16334] Montzka, K. A.; Steitz, J. A.: Additional low-abundance human

[16335] small nuclear ribonucleoproteins: U11, U12, etc. Proc. Nat. Acad.

[16336] Sci. 85: 8885-8889, 1988.

[16337] 1472.

[16338] Nelissen, R. L.; Sillekens, P. T.; Beijer, R. P.; Geurts van Kessel,

[16339] A. H.; van Venrooij, W. J.: Structure, chromosomal local-ization and

[16340] evolutionary conservation of the gene encoding human U1 snRNP-specific

[16341] A protein. Gene 102: 189-196, 1991.

[16342] 1473.

[16343] Spritz, R. A.; Strunk, K.; Surowy, C. S.; Hoch, S. O.; Barton,

[16344] D. E.; Francke, U.: The human 70-kD SnRNP protein: cDNA cloning,

[16345] chromosomal localization, and expression. (Abstract) Am. J. Hum.

[16346] Genet. 41: A239 only, 1987.

[16347] 1474.

[16348] Spritz, R. A.; Strunk, K.; Surowy, C. S.; Hoch, S. O.; Barton,

[16349] D. E.; Francke, U.: Human U1-70K snRNP protein: cDNA cloning, chromosomal

[16350] localization, expression, alternative splicing and RNA-binding. Nucleic

[16351] Acids Res. 15: 10373-10391, 1987.

[16352] 1475.

[16353] Bain, P. A.; Meisler, M. H.; Taylor, B. A.; Payne, A. H.: The

[16354] genes encoding gonadal and nongonadal forms of 3-beta-hydroxysteroid

[16355] dehydrogenase/delta-5-delta-4 isomerase are closely linked on mouse

[16356] chromosome 3. Genomics 16: 219-223, 1993.

[16357] 1476.

[16358] Berube, D.; Luu The, V.; Lachance, Y.; Gagne, R.; Labrie, F.:

[16359] The gene encoding human 3-beta hydroxysteroid dehydrogenase/isomerase

[16360] mapped to the p13 band of chromosome 1. (Abstract) Cytogenet. Cell

[16361] Genet. 51: 962 only, 1989.

[16362] 1477.

[16363] Ruiz-Perez, V. L.; Ide, S. E.; Strom, T. M.; Lorenz, B.; Wil-
son,
[16364] D.; Woods, K.; King, L.; Francomano, C.; Freisinger, P.;
Spranger,
[16365] S.; Marino, B.; Dallapiccola, B.; Wright, M.; Meitinger, T.;
Polymeropoulos,
[16366] M. H.; Goodship, J.: Mutations in a new gene in Ellis-van
Creveld
[16367] syndrome and Weyers acrodental dysostosis. Nature
Genet. 24: 283-286,
[16368] 2000. Note: Erratum: Nature Genet. 25: 125 only, 2000.

[16369] 1478.

[16370] Slavotinek, A. M.; Stone, E. M.; Mykytyn, K.; Heckenlively,
J.
[16371] R.; Green, J. S.; Heon, E.; Musarella, M. A.; Parfrey, P. S.;
Sheffield,
[16372] V. C.; Biesecker, L. G.: Mutations in MKKS cause Bardet-
Biedl syndrome. Nature
[16373] Genet. 26: 15-16, 2000.

[16374] 1479.

[16375] Alos, N.; Moisan, A.-M.; Ward, L.; Desrochers, M.; Legault, L.;
[16376] Leboeuf, G.; Van Vliet, G.; Simard, J.: A novel A10E homozygous mutation
[16377] in the HSD3B2 gene causing severe salt-wasting 3-beta-hydroxysteroid
[16378] dehydrogenase deficiency in 46,XX and 46,XY French-Canadians: evaluation
[16379] of gonadal function after puberty. J. Clin. Endocr. Metab. 85: 1968-1974,
[16380] 2000.

[16381] 1480.

[16382] Douhan, J., III; Hauber, I.; Eibl, M. M.; Glimcher, L. H.: Genetic
[16383] evidence for a new type of major histocompatibility complex class
[16384] II combined immunodeficiency characterized by a dysco-ordinate regulation
[16385] of HLA-D alpha and beta chains. J. Exp. Med. 183: 1063-1069, 1996.
[16386] 1481.

[16387] Moulard, B.; Guipponi, M.; Chaigne, D.; Mouthon, D.; Bu-

resi, C.;

[16388] Malafosse, A.: Identification of a new locus for generalized epilepsy

[16389] with febrile seizures plus (GEFS+) on chromosome 2q24-q33. Am. J.

[16390] Hum. Genet. 65: 1396-1400, 1999.

[16391] 1482.

[16392] Ohmori, I.; Ouchida, M.; Ohtsuka, Y.; Oka, E.; Shimizu, K.: Significant

[16393] correlation of the SCN1A mutations and severe myoclonic epilepsy in

[16394] infancy. Biochem. Biophys. Res. Commun. 295: 17-23, 2002.

[16395] 1483.

[16396] Sugawara, T.; Mazaki-Miyazaki, E.; Fukushima, K.; Shimo-mura, J.;

[16397] Fujiwara, T.; Hamano, S.; Inoue, Y.; Yamakawa, K.: Fre-quent mutations

[16398] of SCN1A in severe myoclonic epilepsy in infancy. Neurol-ogy 58:

[16399] 1122-1124, 2002.

[16400] 1484.

[16401] Luo, X.; Ikeda, Y.; Parker, K. L.: A cell-specific nuclear receptor
[16402] is essential for adrenal and gonadal development and sexual differentiation. Cell 77:
[16403] 481-490, 1994.

[16404] 1485.

[16405] Morohashi, K.: Gonadal and extragonadal functions of Ad4BP/SF-1:
[16406] developmental aspects. Trends Endocr. Metab. 10: 169-173, 1999.
[16407] 1486.

[16408] Nachtigal, M. W.; Hirokawa, Y.; Enyeart-VanHouten, D. L.; Flanagan,
[16409] J. N.; Hammer, G. D.; Ingraham, H. A.: Wilms' tumor 1 and Dax-1 modulate
[16410] the orphan nuclear receptor SF-1 in sex-specific gene expression. Cell 93:
[16411] 445-454, 1998.

[16412] 1487.

[16413] Wallace, R. H.; Scheffer, I. E.; Barnett, S.; Richards, M.; Dibbens,

[16414] L.; Desai, R. R.; Lerman-Sagie, T.; Lev, D.; Mazarib, A.; Brand, N.;

[16415] Ben-Zeev, B.; Goikhman, I.; Singh, R.; Kremmidiotis, G.; Gardner,

[16416] A.; Sutherland, G. R.; George, A. L., Jr.; Mulley, J. C.; Berkovic,

[16417] S. F.: Neuronal sodium-channel alpha-1-subunit mutations in generalized

[16418] epilepsy with febrile seizures plus. Am. J. Hum. Genet. 68: 859-865,

[16419] 2001.

[16420] 1488.

[16421] Ahmed, C. M. I.; Ware, D. H.; Lee, S. C.; Patten, C. D.; Ferrer-Montiel,

[16422] A. V.; Schinder, A. F.; McPherson, J. D.; Wagner-McPherson, C. B.;

[16423] Wasmuth, J. J.; Evans, G. A.; Montal, M.: Primary structure, chromosomal

[16424] localization, and functional expression of a voltage-gated sodium

[16425] channel from human brain. Proc. Nat. Acad. Sci. 89: 8220-8224, 1992.

[16426] 1489.

[16427] Green, R. M.; Ananthanarayanan, M.; Suchy, F. J.; Beier, D. R.

[16428] : Genetic mapping of the Na(+)-taurocholate cotransporting polypeptide

[16429] to mouse chromosome 12. Mammalian Genome 9: 598-600, 1998.

[16430] 1490.

[16431] Hagenbuch, B.; Meier, P. J.: Molecular cloning, chromosomal localization,

[16432] and functional characterization of a human liver Na(+)/bile acid cotransporter. J.

[16433] Clin. Invest. 93: 1326-1331, 1994.

[16434] 1491.

[16435] Hallen, S.; Mareninova, O.; Branden, M.; Sachs, G.: Organization

[16436] of the membrane domain of the human liver sodium/bile acid cotransporter. Biochemistry 41:

[16437] 7253-7266, 2002.

[16438] 1492.

[16439] Shiao, T.; Iwahashi, M.; Fortune, J.; Quattrochi, L.; Bow-

man, S.;

[16440] Wick, M.; Qadri, I.; Simon, F. R.: Structural and functional characterization

[16441] of liver cell-specific activity of the human sodium/ taurocholate cotransporter. Genomics 69:

[16442] 203-213, 2000.

[16443] 1493.

[16444] Hammer, G. D.; Krylova, I.; Zhang, Y.; Darimont, B. D.; Simpson,

[16445] K.; Weigel, N. L.; Ingraham, H. A.: Phosphorylation of the nuclear

[16446] receptor SF-1 modulates cofactor recruitment: integration of hormone

[16447] signaling in reproduction and stress. Molec. Cell 3: 521-526, 1999.

[16448] 1494.

[16449] Harris, A. N.; Mellon, P. L.: The basic helix-loop-helix, leucine

[16450] zipper transcription factor, USF (upstream stimulatory factor), is

[16451] a key regulator of SF-1 (steroidogenic factor-1) gene expression in

[16452] pituitary gonadotrope and steroidogenic cells. Molec. En-
docr. 12:
[16453] 714-726, 1998.

[16454] 1495.

[16455] Lala, D. S.; Rice, D. A.; Parker, K. L.: Steroidogenic factor

[16456] 1, a key regulator of steroidogenic enzyme expression, is
the mouse
[16457] homolog of fushi tarazu-factor 1. Molec. Endocr. 6:
1249-1258, 1992.
[16458] 1496.

[16459] Calvo, R. M.; Asuncion, M.; Telleria, D.; Sancho, J.; San
Millan,
[16460] J. L.; Escobar-Morreale, H. F.: Screening for mutations in
the steroidogenic
[16461] acute regulatory protein and steroidogenic factor-1
genes, and in
[16462] CYP11A and dosage-sensitive sex reversal-adrenal hy-
poplasia gene on
[16463] the X chromosome, gene-1 (DAX-1), in hyperandrogenic
hirsute women. J.
[16464] Clin. Endocr. Metab. 86: 1746-1749, 2001.

[16465] 1497.

[16466] Oba, K.; Yanase, T.; Nomura. M.; Morohashi, K.;
Takayanagi, R.;
[16467] Nawata, H.: Structural characterization of human Ad4bp
(SF-1) gene. Biochem.
[16468] Biophys. Res. Commun. 226: 261-267, 1996.

[16469] 1498.

[16470] Shen, W.-H.; Moore, C. C. D.; Ikeda, Y.; Parker, K. L.; In-
graham,
[16471] H. A.: Nuclear receptor steroidogenic factor 1 regulates
the Mullerian
[16472] inhibiting substance gene: a link to the sex determination
cascade. Cell 77:
[16473] 651-661, 1994.

[16474] 1499.

[16475] Swift, S.; Ashworth, A.: The mouse Ftzf1 gene required for
gonadal
[16476] and adrenal development maps to mouse chromosome 2.
Genomics 28:
[16477] 609-610, 1995.

[16478] 1500.

[16479] Taketo, M.; Parker, K. L.; Howard, T. A.; Tsukiyama, T.; Wong,
[16480] M.; Niwa, O.; Morton, C. C.; Miron, P. M.; Seldin, M. F.: Homologs
[16481] of Drosophila fushi-tarazu factor 1 map to mouse chromosome 2 and
[16482] human chromosome 9q33. Genomics 25: 565-567, 1995.

[16483] 1501.

[16484] Tremblay, A.; Tremblay, G. B.; Labrie, F.; Giguere, V.: Ligand-independent
[16485] recruitment of SRC-1 to estrogen receptor beta through phosphorylation
[16486] of activation function AF-1. Molec. Cell 3: 513-519, 1999.

[16487] 1502.

[16488] Wilhelm, D.; Englert, C.: The Wilms tumor suppressor WT1 regulates
[16489] early gonadal development by activation of Sf1. Genes Dev. 16: 1839-1851,
[16490] 2002.

[16491] 1503.

[16492] Nagase, H.; Barrett, A. J.; Woessner, J. F., Jr.: Nomenclature

[16493] and glossary of the matrix metalloproteinases. Matrix Suppl. 1:
[16494] 421-424, 1992.

[16495] 1504.

[16496] Bertrand, P.; Poirier, J.; Oda, T.; Finch, C. E.; Pasinetti, G.

[16497] M.: Association of apolipoprotein E genotype with brain levels of
[16498] apolipoprotein E and apolipoprotein J (clusterin) in Alzheimer disease. Molec.
[16499] Brain Res. 33: 174-178, 1995.

[16500] 1505.

[16501] Birkenmeier, E. H.; Letts, V. A.; Frankel, W. N.; Magen-heimer,
[16502] B. S.; Calvet, J. P.: Sulfated glycoprotein-2 (Sgp-2) maps to mouse
[16503] chromosome 14. Mammalian Genome 4: 131-132, 1993.

[16504] 1506.

[16505] Danik, M.; Chabot, J.-G.; Hassan-Gonzalez, D.; Suh, M.; Quirion,
[16506] R.: Localization of sulfated glycoprotein-2/clusterin mRNA in the

[16507] rat brain by in situ hybridization. J. Comp. Neurol. 334: 209-227,
[16508] 1993.

[16509] 1507.

[16510] Eldadah, Z. A.; Hamosh, A.; Biery, N. J.; Montgomery, R. A.; Duke,
[16511] M.; Elkins, R.; Dietz, H. C.: Familial tetralogy of Fallot caused
[16512] by mutation in the jagged1 gene. Hum. Molec. Genet. 10: 163-169,
[16513] 2001.

[16514] 1508.

[16515] Pan, H.; Yin, C.; Dyson, N. J.; Harlow, E.; Yamasaki, L.; Van

[16516] Dyke, T.: Key roles for E2F1 in signaling p53-dependent apoptosis
[16517] and in cell division within developing tumors. Molec. Cell 2: 283-292,
[16518] 1998.

[16519] 1509.

[16520] Phillips, A. C.; Ernst, M. K.; Bates, S.; Rice, N. R.; Vousden,

[16521] K. H.: E2F-1 potentiates cell death by blocking antiapop-
totic signaling
[16522] pathways. Molec. Cell 4: 771-781, 1999.

[16523] 1510.

[16524] Saenz Robles, M. T.; Symonds, H.; Chen, J.; Van Dyke, T.:
Induction
[16525] versus progression of brain tumor development: differen-
tial functions
[16526] for the pRB- and p53-targeting domains of simian virus
40 T antigen. Molec.
[16527] Cell. Biol. 14: 2686-2698, 1994.

[16528] 1511.

[16529] Sherr, C. J.: Tumor surveillance via the ARF-p53 pathway.
Genes
[16530] Dev. 12: 2984-2991, 1998.

[16531] 1512.

[16532] Tsai, K. Y.; Hu, Y.; Macleod, K. F.; Crowley, D.; Yamasaki,
L.;
[16533] Jacks, T.: Mutation of E2f-1 suppresses apoptosis and in-
appropriate
[16534] S phase entry and extends survival of Rb-deficient mouse

embryos. Molec.
[16535] Cell 2: 293-304, 1998.

[16536] 1513.

[16537] Weinberg, R. A.: E2F and cell proliferation: a world turned upside
[16538] down. Cell 85: 457-459, 1996.

[16539] 1514.

[16540] Wu, L.; Timmers, C.; Maiti, B.; Saavedra, H. I.; Sang, L.; Chong,
[16541] G. T.; Nuckolls, F.; Giangrande, P.; Wright, F. A.; Field, S. J.;
[16542] Greenberg, M. E.; Orkin, S.; Nevins, J. R.; Robinson, M. L.; Leone,
[16543] G.: The E2F1-3 transcription factors are essential for cellular proliferation. Nature 414:
[16544] 457-462, 2001.

[16545] 1515.

[16546] Yamasaki, L.; Jacks, T.; Bronson, R.; Goillot, E.; Harlow, E.;

[16547] Dyson, N. J.: Tumor induction and tissue atrophy in mice lacking
[16548] E2F-1. Cell 85: 537-548, 1996.

[16549] 1516.

[16550] Zhang, H. S.; Postigo, A. A.; Dean, D. C.: Active transcrip-
tional

[16551] repression by the Rb-E2F complex mediates G1 arrest
triggered by p16(INK4a),

[16552] TGF-beta, and contact inhibition. Cell 97: 53-61, 1999.

[16553] 1517.

[16554] Zhang, Y.; Chellappan, S. P.: Cloning and characterization
of

[16555] human DP2, a novel dimerization partner of E2F. Onco-
gene 10: 2085-2093,

[16556] 1995.

[16557] 1518.

[16558] Agami, R.; Blandino, G.; Oren, M.; Shaul, Y.: Interaction of
c-Abl

[16559] and p73-alpha and their collaboration to induce apopto-
sis. Nature 399:

[16560] 809-813, 1999.

[16561] 1519.

[16562] Arico, M.; Valsecchi, M. G.; Camitta, B.; Schrappe, M.;
Chessells,

[16563] J.; Baruchel, A.; Gaynon, P.; Silverman, L.; Janka-Schaub, G.; Kamps,
[16564] W.; Pui, C.-H.; Masera, G.: Outcome of treatment in children with
[16565] Philadelphia chromosome-positive acute lymphoblastic leukemia. N.
[16566] Eng. J. Med. 342: 998-1006, 2000.

[16567] 1520.

[16568] Barila, D.; Superti-Furga, G.: An intramolecular SH3-domain interaction
[16569] regulates c-Abl activity. Nature Genet. 18: 280-282, 1998.
[16570] 1521.

[16571] Watanabe, M.; Zingg, B. C.; Mohrenweiser, H. W.: Molecular analysis
[16572] of a series of alleles in humans with reduced activity at the triosephosphate
[16573] isomerase locus. Am. J. Hum. Genet. 58: 308-316, 1996.

[16574] 1522.

[16575] FitzGerald, M. G.; Bean, J. M.; Hedge, S. R.; Unsal, H.; MacDonald,

[16576] D. J.; Harkin, D. P.; Finkelstein, D. M.: Isselbacher, K. J.; Haber,
[16577] D. A.: Heterozygous ATM mutations do not contribute to early onset
[16578] of breast cancer. Nature Genet. 15: 307-310, 1997.

[16579] 1523.

[16580] Yuan, P. M.; Dewan, R. N.; Zaun, M.; Thompson, R. E.; Gracy, R.
[16581] W.: Isolation and characterization of triosephosphate isomerase isozymes
[16582] from human placenta. Arch. Biochem. Biophys. 198: 42-52, 1979.

[16583] 1524.

[16584] Zanella, A.; Mariani, M.; Colombo, M. B.; Borgna-Pignatti, C.;
[16585] De Stefano, P.; Morgese, G.; Sirchia, G.: Triosephosphate isomerase
[16586] deficiency: 2 new cases. Scand. J. Haemat. 34: 417-424, 1985.

[16587] 1525.

[16588] Balci, S.; Demirtas, M.; Civelek, B.; Piskin, M.; Sensoz, O.; Akarsu,

[16589] A. N.: Phenotypic variability of triphalangeal thumb-polysyndactyly

[16590] syndrome linked to chromosome 7q36. Am. J. Med. Genet. 87: 399-406,

[16591] 1999.

[16592] 1526.

[16593] Heus, H. C.; Hing, A.; van Baren, M. J.; Joosse, M.; Breed-veld,

[16594] G. J.; Wang, J. C.; Burgess, A.; Donnis-Keller, H.; Berglund, C.;

[16595] Zguricas, J.; Scherer, S. W.; Rommens, J. M.; Oostra, B. A.; Heutink,

[16596] P.: A physical and transcriptional map of the preaxial polydactyly

[16597] locus on chromosome 7q36. Genomics 57: 342-351, 1999.

[16598] 1527.

[16599] Heutink, P.; Zguricas, J.; van Oosterhout, L.; Breedveld, G. J.;

[16600] Testers, L.; Sandkuijl, L. A.; Snijders, P. J. L. M.; Weis-senbach,

[16601] J.; Lindhout, D.; Hovius, S. E. R.; Oostra, B. A.: The gene

for triphalangeal
[16602] thumb maps to the subtelomeric region of chromosome 7q. Nature Genet. 6:
[16603] 287-292, 1994.

[16604] 1528.

[16605] Kantaputra, P. N.; Chalidapong, P.: Are triphalangeal thumb-polysyndactyly
[16606] syndrome (TPTPS) and tibial hemimelia-polysyndactyly-triphalangeal
[16607] thumb syndrome (THPTTS) identical?: A father with TPTPS and his daughter
[16608] with THPTTS in a Thai family. Am. J. Med. Genet. 93: 126-131, 2000.
[16609] 1529.

[16610] Nicolai, J.-P. A.; Hamel, B. C. J.: A family with complex bilateral
[16611] polysyndactyly. J. Hand Surg. 13A: 405-407, 1988.

[16612] 1530.

[16613] Seidman, C.: Personal Communication. Boston, Mass. 3/17/1994.
[16614] 1531.

[16615] Temtamy, S. A.: Personal Communication. Cairo, Egypt 4/25/1994.

[16616] 1532.

[16617] Tsukurov, O.; Boehmer, A.; Flynn, J.; Nicolai, J.-P.; Hamel, B.

[16618] C. J.; Traill, S.; Zaleske, D.; Mankin, H. J.; Yeon, H.; Ho, C.; Tabin,

[16619] C.; Seidman, J. G.; Seidman, C.: A complex bilateral polysyndactyly

[16620] disease locus maps to chromosome 7q36. Nature Genet. 6: 282-286,

[16621] 1994.

[16622] 1533.

[16623] Zguricas, J.; Heus, H.; Morales-Peralta, E.; Breedveld, G.; Kuyt,

[16624] B.; Mumcu, E. F.; Bakker, W.; Akarsu, N.; Kay, S. P. J.; Hov-

ius, S.

[16625] E. R.; Heredero-Baute, L.; Oostra, B. A.; Heutink, P.: Clini-

cal and

[16626] genetic studies on 12 preaxial polydactyly families and re-

finement

[16627] of the localisation of the gene responsible to a 1.9 cM re-

gion on
[16628] chromosome 7q36. J. Med. Genet. 36: 32-40, 1999.

[16629] 1534.

[16630] Zguricas, J.; Snijders, P. J. L. M.; Hovius, S. E. R.; Heutink,

[16631] P.; Oostra, B. A.; Lindhout, D.: Phenotypic analysis of triphalangeal
[16632] thumb and associated hand malformations. J. Med. Genet. 31: 462-467,
[16633] 1994.

[16634] 1535.

[16635] Altafaj, X.; Dierssen, M.; Baamonde, C.; Marti, E.; Visa, J.; Guimera,
[16636] J.; Oset, M.; Gonzalez, J. R.; Florez, J.; Fillat, C.; Estivill, X.

[16637] : Neurodevelopmental delay, motor abnormalities and cognitive deficits
[16638] in transgenic mice overexpressing Dyrk1A (minibrain), a murine model
[16639] of Down's syndrome. Hum. Molec. Genet. 10: 1915-1923, 2001.

[16640] 1536.

[16641] Tanaka, T.; Nakahara, K.; Kato, N.; Imai, T.; Yamazaki, T.;

Tomita,

[16642] H.; Shimokawa, H.; Matsuhashi, H.; Sato, N.; Matsui, M.; Kihira, S.;

[16643] Shimizu, A.; Sano, T.; Haneda, N.; Kino, M.; Miyakita, Y.; Matsuoka,

[16644] R.; Nagai, R.; Yazaki, Y.; Nakamura, Y.: Genetic linkage analyses

[16645] of Romano-Ward syndrome (RWS) in 13 Japanese families. Hum. Genet. 94:

[16646] 380-384, 1994.

[16647] 1537.

[16648] Towbin, J. A.; Li, H.; Taggart, T. R.; Lehmann, M. H.; Schwartz,

[16649] P. J.; Satler, C. A.; Ayyagari, R.; Robinson, J. L.; Moss, A.; Hejtmancik,

[16650] J. F.: Evidence of genetic heterogeneity in Romano-Ward long QT syndrome:

[16651] analysis of 23 families. Circulation 90: 2635-2644, 1994.

[16652] 1538.

[16653] Tye, K.-H.; Desser, K. B.; Benchimol, A.: Survival following

[16654] spontaneous ventricular flutter-fibrillation associated with

QT syndrome:
[16655] documentation during ambulatory monitoring. Arch. In-
tern. Med. 140:
[16656] 255-256, 1980.

[16657] 1539.

[16658] Tyson, J.; Tranebjaerg, L.; McEntagart, M.; Larsen, L. A.;
Christiansen,
[16659] M.; Whiteford, M. L.; Bathen, J.; Aslaksen, B.; Sorland, S. J.;
Lund,
[16660] O.; Pembrey, M. E.; Malcolm, S.; Bitner-Glindzicz, M.: Mu-
tational
[16661] spectrum in the cardioauditory syndrome of Jervell and
Lange-Nielsen. Hum.
[16662] Genet. 107: 499-503, 2000.

[16663] 1540.

[16664] Van der Straaten, P. J. C.; Bruins, C. L. D.: A family with
heritable
[16665] electrocardiographic Q-T prolongation. J. Med. Genet. 10:
158-160,
[16666] 1973.

[16667] 1541.

[16668] Vincent, G. M.: The heart rate of Romano-Ward syndrome patients. Am.
[16669] Heart J. 112: 61-64, 1986.

[16670] 1542.

[16671] Vincent, G. M.; Timothy, K. W.; Leppert, M.; Keating, M.: The
[16672] spectrum of symptoms and QT intervals in carriers of the gene for
[16673] the long-QT syndrome. New Eng. J. Med. 327: 846-852, 1992.

[16674] 1543.

[16675] Ward, O. C.: A new familial cardiac syndrome in children. J.
[16676] Irish Med. Assoc. 54: 103-106, 1964.

[16677] 1544.

[16678] Weitkamp, L. R.; Moss, A. J.: The long QT (Romano-Ward) syndrome
[16679] locus, LQT, is probably linked to the HLA loci. (Abstract) Cytogenet.
[16680] Cell Genet. 40: 775 only, 1985.

[16681] 1545.

[16682] Weitkamp, L. R.; Moss, A. J.; Lewis, R. A.; Hall, W. J.; Mac-Cluer,

[16683] J. W.; Schwartz, P. J.; Locati, E. H.; Tzivoni, D.; Vincent, G. M.;

[16684] Robinson, J. L.; Guttormsen, S. A.: Analysis of HLA and disease susceptibility:

[16685] chromosome 6 genes and sex influence long-QT pheno-type. Am. J. Hum.

[16686] Genet. 55: 1230-1241, 1994.

[16687] 1546.

[16688] Weitkamp, L. R.; Moss, A. J.; Schwartz, P. J.; Locati, E.; Tzivoni,

[16689] D.; Vincent, G. M.; Robinson, J.; Guttormsen, S.: Analysis of HLA

[16690] haplotypes in long QT syndrome: withdrawal of the pre-liminary assignment

[16691] of LQT to the HLA linkage group. (Abstract) Cytogenet. Cell Genet. 51:

[16692] 1106 only, 1989.

[16693] 1547.

[16694] Yang, W.-P.; Levesque, P. C.; Little, W. A.; Conder, M. L.; Shalaby,

[16695] F. Y.; Blanar, M. A.: KvLQT1, a voltage-gated potassium channel responsible

[16696] for human cardiac arrhythmias. Proc. Nat. Acad. Sci. 94: 4017-4021,

[16697] 1997.

[16698] 1548.

[16699] Laudet, V.; Begue, A.; Henry-Duthoit, C.; Joubel, A.; Martin, P.;

[16700] Stehelin, D.; Saule, S.: Genomic organization of the human thyroid

[16701] hormone alpha (c-erbA-1) gene. Nucleic Acids Res. 19: 1105-1112,

[16702] 1991.

[16703] 1549.

[16704] Kluwe, L.; Friedrich, R. E.; Hagel, C.; Lindenau, M.; Mautner,

[16705] V.-F.: Mutations and allelic loss of the NF2 gene in neurofibromatosis

[16706] 2-associated skin tumors. J. Invest. Derm. 114: 1017-1021, 2000.

[16707] 1550.

[16708] Schleutermann, D. A.; Bias, W. B.; Murdoch, J. L.; McKu-
sick, V.
[16709] A.: Linkage of the loci for the nail-patella syndrome and
adenylate
[16710] kinase. Am. J. Hum. Genet. 21: 606-630, 1969.

[16711] 1551.

[16712] Scott, J.; Knott, T. J.; Shaw, D. J.; Brook, J. D.: Localization

[16713] of genes encoding apolipoproteins CI, CII, and E to the
p13-cen region
[16714] of human chromosome 19. Hum. Genet. 71: 144-146,
1985.
[16715] 1552.

[16716] Hildebrand, C. E.; Gonzalez, F. J.; McBride, O. W.; Nebert,
D.
[16717] W.: Assignment of the human
2,3,7,8-tetrachlorodibenzo-p-dioxin-inducible
[16718] cytochrome P1-450 gene to chromosome 15. Nucleic
Acids Res. 13:
[16719] 2009-2016, 1985.

[16720] 1553.

[16721] Olives, B.; Merriman, M.; Bailly, P.; Bain, S.; Barnett, A.;

Todd,

[16722] J.; Cartron, J.-P.; Merriman, T.: The molecular basis of the Kidd

[16723] blood group polymorphism and its lack of association with type 1 diabetes

[16724] susceptibility. Hum. Molec. Genet. 6: 1017-1020, 1997.

[16725] 1554.

[16726] Bahar, A. Y.; Taylor, P. J.; Andrews, L.; Proos, A.; Burnett, L.;

[16727] Tucker, K.; Friedlander, M.; Buckley, M. F.: The frequency of founder

[16728] mutations in the BRCA1, BRCA2, and APC genes in Aus-

tralian Ashkenazi

[16729] Jews: implications for the generality of U.S. population data. Cancer 92:

[16730] 440-445, 2001.

[16731] 1555.

[16732] Cohn, D. V.; Zangerle, R.; Fischer-Colbrie, R. R.; Chu, L. L. H.;

[16733] Elting, J. J.; Hamilton, J. W.; Winkler, H.: Similarity of secre-

tory

[16734] protein I from parathyroid gland to chromogranin A from

the adrenal

[16735] medulla. Proc. Nat. Acad. Sci. 79: 6056-6059, 1982.

[16736] 1556.

[16737] Stover, M. L.; Primorac, D.; Liu, S. C.; McKinstry, M. B.; Rowe,

[16738] D. W.: Defective splicing of mRNA from one COL1A1 allele of type

[16739] I collagen in nondeforming (type I) osteogenesis imperfecta. J. Clin.

[16740] Invest. 92: 1994-2002, 1993.

[16741] 1557.

[16742] Sundar Raj, C. V.; Church, R. L.; Klobutcher, L. A.; Ruddle, F.

[16743] H.: Genetics of the connective tissue proteins: assignment of the

[16744] gene for human type I procollagen to chromosome 17 by analysis of

[16745] cell hybrids and microcell hybrids. Proc. Nat. Acad. Sci. 74: 4444-4448,

[16746] 1977.

[16747] 1558.

[16748] Tromp, G.; Kuivaniemi, H.; Stacey, A.; Shikata, H.; Baldwin, C.

[16749] T.; Jaenisch, R.; Prockop, D. J.: Structure of a full-length cDNA

[16750] clone for the prepro-alpha-1(I) chain of human type I procollagen. Biochem.

[16751] J. 253: 919-922, 1988.

[16752] 1559.

[16753] Tsipouras, P.; Schwartz, R. C.; Phillips, J. A., III; Willard,

[16754] H. F.: A centromere-based linkage group on the long arm of human

[16755] chromosome 17. Cytogenet. Cell Genet. 47: 109-110, 1988.

[16756]

[16757] 100. Tsuneyoshi, T.; Westerhausen, A.; Constantinou, C. D.; Prockop,

[16758] D. J.: Substitutions for glycine alpha-1-637 and glycine alpha-2-694

[16759] of type I procollagen in lethal osteogenesis imperfecta: the conformational

[16760] strain on the triple helix introduced by a glycine substitution can

[16761] be transmitted along the helix. J. Biol. Chem. 266: 15608-15613,

[16762] 1991.

[16763]

[16764] 101. Uitterlinden, A. G.; Burger, H.; Huang, Q.; Yue, F.; McGuigan,

[16765] F. E. A.; Grant, S. F. A.; Hofman, A.; van Leeuwen, J. P. T. M.; Pols,

[16766] H. A. P.; Ralston, S. H.: Relation of alleles of the collagen type

[16767] I-alpha-1 gene to bone density and the risk of osteo-porotic fractures

[16768] in postmenopausal women. New Eng. J. Med. 338: 1016-1021, 1998.

[16769]

[16770] 102. Uitterlinden, A. G.; Weel, A. E. A. M.; Burger, H.; Fang, Y.;

[16771] Van Duijn, C. M.; Hofman, A.; Van Leeuwen, J. P. T. M.; Pols, H. A.

[16772] P.: Interaction between the vitamin D receptor gene and collagen

[16773] type I-alpha-1 gene in susceptibility for fracture. J. Bone

Miner.

[16774] Res. 16: 379-385, 2001.

[16775]

[16776] 103. Valli, M.; Mottes, M.; Tenni, R.; Sangalli, A.; Gomez Lira, M.;

[16777] Rossi, A.; Antoniazzi, F.; Cetta, G.; Pignatti, P. F.: A de novo

[16778] G to T transversion in a pro-alpha-1(I) collagen gene for a moderate

[16779] case of osteogenesis imperfecta: substitution of cysteine for glycine

[16780] 178 in the triple helical domain. J. Biol. Chem. 266: 1872-1878,

[16781] 1991.

[16782]

[16783] 104. Vogel, B. E.; Doelz, R.; Kadler, K. E.; Hojima, Y.; Engel, J.;

[16784] Prockop, D. J.: A substitution of cysteine for glycine 748 of the

[16785] alpha-1 chain produces a kink at this site in the procollagen I molecule

[16786] and an altered N-proteinase cleavage site over 225 nm

away. J. Biol.
[16787] Chem. 263: 19249-19255, 1988.

[16788]

[16789] 105. Vogel, B. E.; Minor, R. R.; Freund, M.; Prockop, D. J.: A point
[16790] mutation in a type I procollagen gene converts glycine 748 of the
[16791] alpha-1 chain to cysteine and destabilizes the triple helix in a lethal
[16792] variant of osteogenesis imperfecta. J. Biol. Chem. 262: 14737-14744,
[16793] 1987.

[16794]

[16795] 106. Wallis, G. A.; Starman, B. J.; Byers, P. H.: Clinical het-erogeneity
[16796] of OI explained by molecular heterogeneity and somatic mosaicism.(Abstract) Am.
[16797] J. Hum. Genet. 45: A228, 1989.

[16798]

[16799] 107. Wallis, G. A.; Starman, B. J.; Zinn, A. B.; Byers, P. H.: Variable

[16800] expression of osteogenesis imperfecta in a nuclear family is explained

[16801] by somatic mosaicism for a lethal point mutation in the alpha-1(I)

[16802] gene (COL1A1) of type I collagen in a parent. Am. J. Hum. Genet. 46:

[16803] 1034-1040, 1990.

[16804]

[16805] 108. Wallis, G. A.; Sykes, B.; Byers, P. H.; Mathew, C. G.; Viljoen,

[16806] D.; Beighton, P.: Osteogenesis imperfecta type III: mutations in

[16807] the type I collagen structural genes, COL1A1 and COL1A2, are not necessarily

[16808] responsible. J. Med. Genet. 30: 492-496, 1993.

[16809]

[16810] 109. Wang, Q.; Forlino, A.; Marini, J. C.: Alternative splicing in

[16811] COL1A1 mRNA leads to a partial null allele and two in-frame forms

[16812] with structural defects in non-lethal osteogenesis imperfecta. J.

[16813] Biol. Chem. 271: 28617-28623, 1996.

[16814]

[16815] 110. Weil, D.; D'Alessio, M.; Ramirez, F.; de Wet, W.; Cole, W. G.;

[16816] Chan, D.; Bateman, J. F.: A base substitution in the exon of a collagen

[16817] gene causes alternative splicing and generates a structurally abnormal

[16818] polypeptide in a patient with Ehlers-Danlos syndrome type VII. EMBO

[16819] J. 8: 1705-1710, 1989.

[16820]

[16821] 111. Weisinger, B.; Glassman, E.; Spencer, F. C.; Berger, A.: Successful

[16822] aortic valve replacement for aortic regurgitation associated with

[16823] osteogenesis imperfecta. Brit. Heart J. 37: 475-477, 1975.

[16824]

[16825] 112. Willing, M. C.; Cohn, D. H.; Byers, P. H.: Frameshift mutation

[16826] near the 3-prime end of the COL1A1 gene of type I colla-

gen predicts

[16827] an elongated pro-alpha-1(I) chain and results in osteogenesis imperfecta

[16828] type I. J. Clin. Invest. 85: 282-290, 1990.

[16829]

[16830] 113. Willing, M. C.; Deschenes, S. P.; Scott, D. A.; Byers, P. H.;

[16831] Slayton, R. L.; Pitts, S. H.; Arikat, H.; Roberts, E. J.: Osteogenesis

[16832] imperfecta type I: molecular heterogeneity for COL1A1 null alleles

[16833] of type I collagen. Am. J. Hum. Genet. 55: 638-647, 1994.

[16834]

[16835] 114. Willing, M. C.; Deschenes, S. P.; Slayton, R. L.; Roberts, E.

[16836] J.: Premature chain termination is a unifying mechanism for COL1A1

[16837] null alleles in osteogenesis imperfecta type I cell strains. Am.

[16838] J. Hum. Genet. 59: 799-809, 1996.

[16839]

[16840] 115. Willing, M. C.; Pruchno, C. J.; Atkinson, M.; Byers, P. H.:
[16841] Osteogensis (sic) imperfecta type I is commonly due to a COL1A1 null
[16842] allele of type I collagen. Am. J. Hum. Genet. 51: 508-515, 1992.
[16843]

[16844] 116. Willing, M. C.; Pruchno, C. J.; Byers, P. H.: Molecular heterogeneity
[16845] in osteogenesis imperfecta type I. Am. J. Med. Genet. 45: 223-227,
[16846] 1993.

[16847]

[16848] 117. Willing, M. C.; Slayton, R. L.; Pitts, S. H.; Deschenes, S. P.
[16849] : Absence of mutations in the promoter of the COL1A1 gene of type
[16850] I collagen in patients with osteogenesis imperfecta type I. J. Med.
[16851] Genet. 32: 697-700, 1995.

[16852]

[16853] 118. Zhuang, J.; Constantinou, C. D.; Ganguly, A.; Prockop, D. J.

[16854] : A single base mutation in type I procollagen (COL1A1) that converts

[16855] glycine alpha(1)-541 to aspartate in a lethal variant of osteogenesis

[16856] imperfecta: detection of the mutation with a carbodiimide reaction

[16857] of DNA heteroduplexes and direct sequencing of products of the PCR. Am.

[16858] J. Hum. Genet. 48: 1186-1191, 1991.

[16859]

[16860] 119. Zhuang, J.; Tromp, G.; Kuivaniemi, H.; Castells, S.; Prockop,

[16861] D. J.: Substitution of arginine for glycine at position 154 of the

[16862] alpha-1 chain of type I collagen in a variant of osteogenesis imperfecta:

[16863] comparison to previous cases with the same mutation. Am. J. Med.

[16864] Genet. 61: 111-116, 1996.

[16865] 1560.

[16866] Bianchi, G.; Tripodi, G.; Casari, G.; Salardi, S.; Barber, B. R.;

[16867] Garcia, R.; Leoni, P.; Torielli, L.; Cusi, D.; Ferrandi, M.; Pinna,

[16868] L. A.; Baralle, F. E.; Ferrari, P.: Two point mutations within the

[16869] adducin genes are involved in blood pressure variation. Proc. Nat.

[16870] Acad. Sci. 91: 3999-4003, 1994.

[16871] 1561.

[16872] Cusi, D.; Barlassina, C.; Azzani, T.; Casari, G.; Citterio, L.;

[16873] Devoto, M.; Glorioso, N.; Lanzani, C.; Manunta, P.; Righetti, M.;

[16874] Rivera, R.; Stella, P.; Troffa, C.; Zagato, L.; Bianchi, G.: Polymorphisms

[16875] of alpha-adducin and salt sensitivity in patients with essential hypertension. Lancet 349:

[16876] 1353-1357, 1997.

[16877] 1562.

[16878] Gardner, K.; Bennett, V.: A new erythrocyte membrane-associated

[16879] protein with calmodulin binding activity: identification and

purification. J.

[16880] Biol. Chem. 261: 1339-1348, 1986.

[16881] 1563.

[16882] Goldberg, Y. P.; Lin, B.-Y.; Andrew, S. E.; Nasir, J.; Graham,

[16883] R.; Glaves, M. L.; Hutchinson, G.; Theilmann, J.; Ginzinger, D. G.;

[16884] Schappert, K.; Clarke, L.; Rommens, J. M.; Hayden, M. R.: Cloning

[16885] and mapping of the alpha-adducin gene close to D4S95 and assessment

[16886] of its relationship to Huntington disease. Hum. Molec. Genet. 1:

[16887] 669-675, 1992.

[16888] 1564.

[16889] Hayden, M. R.: Huntington's Chorea. New York: Springer-Verlag

[16890] (pub.) 1981.

[16891] 1565.

[16892] Joshi, R.; Bennett, V.: Mapping the domain structure of human

[16893] erythrocyte adducin. J. Biol. Chem. 265: 13130-13136,

1990.

[16894] 1566.

[16895] Joshi, R.; Gilligan, D. M.; Otto, E.; McLaughlin, T.; Bennett,

[16896] V.: Primary structure and domain organization of human alpha and

[16897] beta adducin. J. Cell Biol. 115: 665-675, 1991.

[16898] 1567.

[16899] Seger, J.; Tchen, P.; Feingold, N.; Grenand, F.; Bois, E.: Ho-mozygosity

[16900] of adenylate kinase allele 3: two cases. Hum. Genet. 43: 337-339,

[16901] 1978.

[16902] 1568.

[16903] Singer, J. D.; Brock, D. J.: Half-normal adenylate kinase activity

[16904] in three generations. Ann. Hum. Genet. 35: 109-114, 1971.

[16905] 1569.

[16906] Szeinberg, A.; Gavendo, S.; Cahane, D.: Erythrocyte adenylate-kinase

[16907] deficiency. (Letter) Lancet I: 315-316, 1969.

[16908] 1570.

[16909] Szeinberg, A.; Kahana, D.; Gavendo, S.; Zaidman, J.; Ben-Ezzer,

[16910] J.: Hereditary deficiency of adenylate kinase in red blood cells. Acta

[16911] Haemat. 42: 111-126, 1969.

[16912] 1571.

[16913] Toren, A.; Brok-Simoni, F.; Ben-Bassat, I.; Holtzman, F.; Mandel,

[16914] M.; Neumann, Y.; Ramot, B.; Rechavi, G.; Kende, G.: Con-genital haemolytic

[16915] anaemia associated with adenylate kinase deficiency. Brit. J. Haemat. 87:

[16916] 376-380, 1994.

[16917] 1572.

[16918] Weitkamp, L. R.; Sing, C. F.; Shreffler, D. C.; Guttormsen, S.

[16919] A.: The genetic linkage relations of adenylate kinase: fur-ther data

[16920] on the ABO-AK linkage group. Am. J. Hum. Genet. 21: 600-605, 1969.

[16921] 1573.

[16922] Westerveld, A.; Jongsma, A. P. M.; Meera Khan, P.; Van Someren,
[16923] H.; Bootsma, D.: Assignment of the AK(1): Np: AKO linkage group to
[16924] human chromosome 9. Proc. Nat. Acad. Sci. 73: 895-899, 1976.

[16925] 1574.

[16926] Zuffardi, O.; Caiulo, A.; Maraschio, P.; Tupler, R.; Bianchi,
[16927] E.; Amisano, P.; Beluffi, G.; Moratti, R.; Liguri, G.: Regional assignment
[16928] of the loci for adenylate kinase to 9q32 and for alpha(1)-acid glycoprotein
[16929] to 9q31-q32: a locus for Goltz syndrome in region 9q32-qter? Hum.
[16930] Genet. 82: 17-19, 1989.

[16931] 1575.

[16932] Cook, P. J. L.; Buckton, K. E.; Spowart, G.: Family studies on
[16933] chromosome 9. Cytogenet. Cell Genet. 16: 284-288, 1976.

[16934] 1576.

[16935] Mohandas, T.; Sparkes, R. S.; Sparkes, M. C.; Shulkin, J. D.; Toomey,
[16936] K. E.; Funderburk, S. J.: Regional localization of human gene loci
[16937] on chromosome 9: studies of somatic cell hybrids con-taining human
[16938] translocation. Am. J. Hum. Genet. 31: 586-600, 1979.

[16939] 1577.

[16940] Pilz, A.; Woodward, K.; Povey, S.; Abbott, C.: Comparative mapping
[16941] of 50 human chromosome 9 loci in the laboratory mouse. Genomics 25:
[16942] 139-149, 1995.

[16943] 1578.

[16944] Povey, S.; Slaughter, C. A.; Wilson, D. E.; Gormley, I. P.; Buckton,
[16945] K. E.; Perry, P.; Bobrow, M.: Evidence for the assignment of the
[16946] loci AK 1, AK 3 and ACON to chromosome 9 in man. Ann. Hum. Genet. 39:

[16947] 413-422, 1976.

[16948] 1579.

[16949] Robson, E. B.; Meera Khan, P.: Report of the committee on the

[16950] genetic constitution of chromosomes 7, 8, and 9. Cyto-genet. Cell

[16951] Genet. 32: 144-152, 1982.

[16952] 1580.

[16953] Steinbach, P.; Benz, R.: Demonstration of gene dosage ef-fects

[16954] for AK3 and GALT in fibroblasts from a fetus with 9p tri-somy. Hum.

[16955] Genet. 63: 290-291, 1983.

[16956] 1581.

[16957] Viskochil, D.; Buchberg, A. M.; Xu, G.; Cawthon, R. M.; Stevens,

[16958] J.; Wolff, R. K.; Culver, M.; Carey, J. C.; Copeland, N. G.; Jenkins,

[16959] N. A.; White, R.; O'Connell, P.: Deletions and a transloca-tion interrupt

[16960] a cloned gene at the neurofibromatosis type 1 locus. Cell

62: 187-192,

[16961] 1990.

[16962] 1582.

[16963] Wilson, D. E., Jr.; Povey, S.; Harris, H.: Adenylate kinases in

[16964] man: evidence for a third locus. Ann. Hum. Genet. 39: 305-313, 1976.

[16965] 1583.

[16966] Xu, G.; O'Connell, P.; Stevens, J.; White, R.: Characterization

[16967] of human adenylate kinase 3 (AK3) cDNA and mapping of the AK3 pseudogene

[16968] to an intron of the NF1 gene. Genomics 13: 537-542, 1992.

[16969] 1584.

[16970] Budarf, M. L.; Emanuel, B. S.; Collins, J.; Fibison, W.; Barshop,

[16971] B. A.: Isolation and regional localization of the human adenylosuccinate

[16972] lyase gene. (Abstract) Cytogenet. Cell Genet. 58: 2046 only, 1991.

[16973] 1585.

[16974] Fon, E. A.; Demczuk, S.; Delattre, O.; Thomas, G.; Rouleau, G.

[16975] A.: Mapping of the human adenylosuccinate lyase (ADSL) gene to chromosome

[16976] 22q13.1-q13.2. Cytogenet. Cell Genet. 64: 201-203, 1993.

[16977] 1586.

[16978] Jaeken, J.; Van den Bergh, F.; Vincent, M. F.; Casaer, P.; Van

[16979] den Berghe, G.: Adenylosuccinase deficiency: a newly recognized variant. J.

[16980] Inherit. Metab. Dis. 15: 416-418, 1992.

[16981] 1587.

[16982] Jaeken, J.; Van den Berghe, G.: An infantile autistic syndrome

[16983] characterised by the presence of succinylpurines in body fluids. Lancet II:

[16984] 1058-1061, 1984.

[16985] 1588.

[16986] Jaeken, J.; Wadman, S. K.; Duran, M.; van Sprang, F. J.; Beemer,

[16987] F. A.; Holl, R. A.; Theunissen, P. M.; de Cock, P.; van den Bergh,
[16988] F.; Vincent, M. F.; van den Berghe, G.: Adenylosuccinase deficiency:
[16989] an inborn error of purine nucleotide synthesis. Europ. J. Pediat. 148:
[16990] 126-131, 1988.

[16991] 1589.

[16992] Kmoch, S.; Hartmannova, H.; Krijt, J.; Valik, D.; Jones, J. D.;

[16993] Sebesta, L.: Genetic heterogeneity in adenylosuccinate lyase deficiency. Clin.
[16994] Biochem. 30: 264 only, 1997.

[16995] 1590.

[16996] Kmoch, S.; Hartmannova, H.; Stiburkova, B.; Krijt, J.; Zikanova,
[16997] M.; Sebesta, I.: Human adenylosuccinate lyase (ADSL), cloning and
[16998] characterization of full-length cDNA and its isoform, gene structure
[16999] and molecular basis for ADSL deficiency in six patients. Hum. Molec.

[17000] Genet. 9: 1501-1513, 2000.

[17001] 1591.

[17002] Maaswinkel-Mooij, P. D.; Laan, L. A. E. M.; Onkenhout, W.; Brouwer,

[17003] O. F.; Jaeken, J.; Poorthuis, B. J. H. M.: Adenylosuccinase deficiency

[17004] presenting with epilepsy in early infancy. J. Inherit. Metab. 20:

[17005] 606-607, 1997.

[17006] 1592.

[17007] Maddocks, J.; Reed, T.: Urine test for adenylosuccinase deficiency

[17008] in autistic children. (Letter) Lancet I: 158-159, 1989.

[17009] 1593.

[17010] Marie, S.; Cuppens, H.; Heuterspreute, M.; Jaspers, M.; Tola,

[17011] E. Z.; Gu, X. X.; Legius, E.; Vincent, M.-F.; Jaeken, J.; Cassi-man,

[17012] J.-J.; Van den Berghe, G.: Mutation analysis in adenylosuc-cinate

[17013] lyase deficiency: eight novel mutations in the re-evaluated

full ADSL
[17014] coding sequence. Hum. Mutat. 13: 197-202, 1999.

[17015] 1594.

[17016] Marie, S.; Race, V.; Nassogne, M.-C.; Vincent, M.-F.; Van den
[17017] Berghe, G.: Mutation of a nuclear respiratory factor 2 binding site
[17018] in the 5-prime untranslated region of the ADSL gene in three patients
[17019] with adenylosuccinate lyase deficiency. Am. J. Hum. Genet. 71: 14-21,
[17020] 2002.

[17021] 1595.

[17022] Salerno, C.; Crifo, C.; Giardini, O.: Adenylosuccinase defi-ciency:
[17023] a patient with impaired erythrocyte activity and anomalous response
[17024] to intravenous fructose. J. Inherit. Metab. Dis. 18: 602-608, 1995.
[17025] 1596.

[17026] Adinolfi, A.; Adinolfi, M.; Hopkinson, D. A.: Immunological

and

[17027] biochemical characterization of the human alcohol dehydrogenase chi-ADH

[17028] isozyme. Ann. Hum. Genet. 48: 1-10, 1984.

[17029] 1597.

[17030] Beisswenger, T. B.; Holmquist, B.; Vallee, B. L.: Chi-ADH is the

[17031] sole alcohol dehydrogenase isozyme of mammalian brains: implications

[17032] and inferences. Proc. Nat. Acad. Sci. 82: 8369-8373, 1985.

[17033] 1598.

[17034] Carlock, L.; Hiroshige, S.; Wasmuth, J.; Smith, M.: Assignment

[17035] of the gene coding for class III ADH to human chromosome 4: 4q21-4q25.

[17036] (Abstract) Cytogenet. Cell Genet. 40: 598 only, 1985.

[17037] 1599.

[17038] Engeland, K.; Hoog, J.-O.; Holmquist, B.; Estonius, M.; Jornvall,

[17039] H.; Vallee, B. L.: Mutation of arg-115 of human class III al-

cohol

[17040] dehydrogenase: a binding site required for formaldehyde dehydrogenase

[17041] activity and fatty acid activation. Proc. Nat. Acad. Sci. 90: 2491-2494,

[17042] 1993.

[17043] 1600.

[17044] Giri, P.; Krug, J. F.; Kozak, C.; Moretti, T.; O'Brien, S. J.;

[17045] Seuanez, H. N.; Goldman, D.: Cloning and comparative mapping of a

[17046] human class III (chi) alcohol dehydrogenase cDNA. Biochem. Biophys.

[17047] Res. Commun. 164: 453-460, 1989.

[17048] 1601.

[17049] Goldman, D.; RathnaGiri, P.; Moretti, T. R.; Krug, J. F.; Kozak,

[17050] C.; Dean, M.; Seuanez, H. N.; O'Brien, S. J.: Class III alcohol dehydrogenase

[17051] (ADH5): widespread expression and synteny with other ADHs in both

[17052] mouse and man. (Abstract) Am. J. Hum. Genet. 45

(suppl.): A141

[17053] only, 1989.

[17054] 1602.

[17055] Matsuo, Y.; Yokoyama, S.: Cloning and sequencing of a processed

[17056] pseudogene derived from a human class III alcohol dehydrogenase gene.

[17057] Am. J. Hum. Genet. 46: 85-91, 1990.

[17058] 1603.

[17059] Smith, M.: Genetics of human alcohol and aldehyde dehydrogenases.

[17060] Adv. Hum. Genet. 15: 249-290, 1986.

[17061] 1604.

[17062] Herz, J.; Hamann, U.; Rogne, S.; Myklebost, O.; Gausepohl, H.;

[17063] Stanley, K. K.: Surface location and high affinity for calcium of

[17064] a 500 kd liver membrane protein closely related to the LDL-receptor

[17065] suggest a physiological role as lipoprotein receptor. EMBO J. 7:

[17066] 4119-4127, 1988.

[17067] 1605.

[17068] Cailleau-Thomas, A.; Coullin, P.; Candelier, J.-J.; Bal-
anzino,
[17069] L.; Mennesson, B.; Oriol, R.; Mollicone, R.: FUT4 and FUT9
genes
[17070] are expressed early in human embryogenesis. Glycobiol-
ogy 10: 789-802,
[17071] 2000.

[17072] 1606.

[17073] Couillin, P.; Mollicone, R.; Grisard, M. C.; Gibaud, A.;
Ravise,
[17074] N.; Feingold, J.; Oriol, R.: Chromosome 11q localization of
one of
[17075] the three expected genes for the human alpha-
3-fucosyltransferases,
[17076] by somatic hybridization. Cytogenet. Cell Genet. 56:
108-111, 1991.

[17077] 1607.

[17078] Geurts van Kessel, A.; Tetteroo, P.; van Agthoven, T.;
Paulussen,

[17079] R.; van Dongen, J.; Hagemeijer, A.; von dem Borne, A.: Localization

[17080] of human myeloid-associated surface antigen detected by a panel of

[17081] 20 monoclonal antibodies to the q12-qter region of chromosome 11. J.

[17082] Immun. 133: 1265-1269, 1984.

[17083] 1608.

[17084] Cazeneuve, C.; Ajrapetyan, H.; Papin, S.; Roudot-Thoraval, F.;

[17085] Genevieve, D.; Mndjoyan, E.; Papazian, M.; Sarkisian, A.; Babloyan,

[17086] A.; Boissier, B.; Duquesnoy, P.; Kouyoumdjian, J.-C.; Girodon-Boulandet,

[17087] E.; Grateau, G.; Sarkisian, T.; Amselem, S.: Identification of MEFV-independent

[17088] modifying genetic factors for familial Mediterranean fever. Am. J.

[17089] Hum. Genet. 67: 1136-1143, 2000.

[17090] 1609.

[17091] Sack, G. H., Jr.: Serum amyloid A (SAA) gene variations in familial

[17092] Mediterranean fever. Molec. Biol. Med. 5: 61-67, 1988.

[17093] 1610.

[17094] Azem, A.; Kessel, M.; Goloubinoff, P.: Characterization of a functional

[17095] GroEL-14(GroES-7)-2 chaperonin hetero-oligomer. Science 265: 653-656,

[17096] 1994.

[17097] 1611.

[17098] Cheng, M. Y.; Hartl, F.-U.; Martin, J.; Pollock, R. A.; Kalousek,

[17099] F.; Neupert, W.; Hallberg, E. M.; Hallberg, R. L.; Horwich, A. L.

[17100] : Mitochondrial heat-shock protein hsp60 is essential for assembly

[17101] of proteins imported into yeast mitochondria. Nature 337: 620-625,

[17102] 1989.

[17103] 1612.

[17104] Ellis, R. J.: The molecular chaperone concept. Semin. Cell Biol. 1:

[17105] 1-9, 1990.

[17106] 1613.

[17107] Fontaine, B.; Davoine, C.-S.; Durr, A.; Paternotte, C.; Feki, I.;

[17108] Weissenbach, J.; Hazan, J.; Brice, A.: A new locus for auto-somal

[17109] dominant pure spastic paraplegia, on chromosome 2q24-q34. Am. J.

[17110] Hum. Genet. 66: 702-707, 2000.

[17111] 1614.

[17112] Rothman, J. E.: Polypeptide chain binding proteins: cata-lysts

[17113] of protein folding and related processes in cells. Cell 59: 591-601,

[17114] 1989.

[17115] 1615.

[17116] Saibil, H.; Dong, Z.; Wood, S.; auf der Mauer, A.: Binding of

[17117] chaperonins. Nature 353: 25-26, 1991.

[17118] 1616.

[17119] Schmidt, M.; Rutkat, K.; Rachel, R.; Pfeifer, G.; Jaenicke, R.;

[17120] Viitanen, P.; Lorimer, G.; Buchner, J.: Symmetric com-

plexes of GroE

[17121] chaperonins as part of the functional cycle. Science 265: 656-659,

[17122] 1994.

[17123] 1617.

[17124] Venner, T. J.; Singh, B.; Gupta, R. S.: Nucleotide sequences

[17125] and novel structural features of human and Chinese hamster hsp60 (chaperonin)

[17126] gene families. DNA Cell Biol. 9: 545-552, 1990.

[17127] 1618.

[17128] Smit, M.; van der Kooij-Meijs, E.; Frants, R. R.; Havekes, L.;

[17129] Klasen, E. C.: Apolipoprotein gene cluster on chromosome 19: definite

[17130] localization of the APOC2 gene and the polymorphic HpaI site associated

[17131] with type III hyperlipoproteinemia. Hum. Genet. 78: 90-93, 1988.

[17132] 1619.

[17133] Smit, M.; van der Kooij-Meijs, E.; Woudt, L. P.; Havekes, L. M.;

[17134] Frants, R. R.: Exact localization of the familial dysbetal-

ipoproteinemia
[17135] associated HpaI restriction site in the promoter region of the APOC1
[17136] gene. Biochem. Biophys. Res. Commun. 152: 1282-1288, 1988.
[17137] 1620.

[17138] Tata, F.; Henry, I.; Markham, A. F.; Wallis, S. C.; Weil, D.;

[17139] Grzeschik, K. H.; Junien, C.; Williamson, R.; Humphries, S. E.: Isolation
[17140] and characterisation of a cDNA clone for human apolipoprotein CI and
[17141] assignment of the gene to chromosome 19. Hum. Genet. 69: 345-349,
[17142] 1985.

[17143] 1621.

[17144] Coste, H.; Rodriguez, J. C.: Orphan nuclear hormone re-ceptor Rev-erb-alpha
[17145] regulates the human apolipoprotein CIII promoter. J. Biol. Chem. 277:
[17146] 27120-27129, 2002.

[17147] 1622.

[17148] Dammerman, M.; Sandkuijl, L. A.; Halaas, J. L.; Chung, W.; Breslow,

[17149] J. L.: An apolipoprotein CIII haplotype protective against hypertriglyceridemia

[17150] is specified by promoter and 3-prime untranslated region polymorphisms. Proc.

[17151] Nat. Acad. Sci. 90: 4562-4566, 1993.

[17152] 1623.

[17153] Henderson, H. E.; Landon, S. V.; Michie, J.; Berger, G. M. B.:

[17154] Association of a DNA polymorphism in the apolipoprotein C-III gene

[17155] with diverse hyperlipidaemic phenotypes. Hum. Genet. 75: 62-65,

[17156] 1987.

[17157] 1624.

[17158] Karathanasis, S. K.; McPherson, J.; Zannis, V. I.; Breslow, J.

[17159] L.: Linkage of human apolipoproteins A-I and C-III genes. Nature 304:

[17160] 371-373, 1984.

[17161] 1625.

[17162] Li, W. W.; Dammerman, M. M.; Smith, J. D.; Metzger, S.; Breslow,
[17163] J. L.; Leff, T.: Common genetic variation in the promoter of the
[17164] human apo CIII gene abolishes regulation by insulin and may contribute
[17165] to hypertriglyceridemia. J. Clin. Invest. 96: 2601-2605, 1995.
[17166] 1626.

[17167] Maeda, H.; Hashimoto, R. K.; Oguro, T.; Hiraga, S.; Uzawa, H.:
[17168] Molecular cloning of a human apoC-III variant: thr 74-to-ala 74 mutation
[17169] prevents O-glycosylation. J. Lipid Res. 28: 1405-1409, 1987.
[17170] 1627.

[17171] Maeda, H.; Uzawa, H.; Kamei, R.: Unusual familial lipopro-tein
[17172] C-III associated with apolipoprotein C-III-0 preponder-ance. Biochim.
[17173] Biophys. Acta 665: 578-585, 1981.

[17174] 1628.

[17175] Oettgen, P.; Antonarakis, S. E.; Karathanasis, S. K.: PvuII polymorphic
[17176] site upstream to the human apoCIII gene. Nucleic Acids Res. 14:
[17177] 5571, 1986.

[17178] 1629.

[17179] von Eckardstein, A.; Holz, H.; Sandkamp, M.; Weng, W.; Funke,
[17180] H.; Assmann, G.: Apolipoprotein C-III(lys58-to-glu): identification
[17181] of an apolipoprotein C-III variant in a family with hyperalphalipoproteinemia. J.
[17182] Clin. Invest. 87: 1724-1731, 1991.

[17183] 1630.

[17184] Vu-Dac, N.; Gervois, P.; Torra, I. P.; Fruchart, J.-C.; Kosykh,
[17185] V.; Kooistra, T.; Princen, H. M. G.; Dallongeville, J.; Staels, B.
[17186] : Retinoids increase human Apo C-III expression at the transcriptional
[17187] level via the retinoid X receptor: contribution to the hypertriglyceridemic

[17188] action of retinoids. J. Clin. Invest. 102: 625-632, 1998.

[17189] 1631.

[17190] Zannis, V. I.; Kan, H. Y.; Kritis, A.; Zanni, E. E.; Kardassis,

[17191] D.: Transcriptional regulatory mechanisms of the human apolipoprotein

[17192] genes in vitro and in vivo. Curr. Opin. Lipid. 12: 181-207, 2001.

[17193] 1632.

[17194] Jaiswal, A. K.; Gonzalez, F. J.; Nebert, D. W.: Comparison of

[17195] human mouse P(1)450 upstream regulatory sequences in liver- and nonliver-derived

[17196] cell lines. Molec. Endocr. 1: 312-320, 1987.

[17197] 1633.

[17198] Jaiswal, A. K.; Gonzalez, F. J.; Nebert, D. W.: Human P(1)-450

[17199] gene sequence and correlation of mRNA with genetic dif- ferences in

[17200] benzo(a)pyrene metabolism. Nucleic Acids Res. 13: 4503-4520, 1985.

[17201] 1634.

[17202] Jaiswal, A. K.; Gonzalez, F. J.; Nebert, D. W.: Human dioxin-inducible

[17203] cytochrome P1-450: complementary DNA and amino acid sequence. Science 228:

[17204] 80-83, 1985.

[17205] 1635.

[17206] Jaiswal, A. K.; Nebert, D. W.: Two RFLPs associated with the

[17207] human P(1)450 gene linked to the MPI locus on chromosome 15 (HGM8

[17208] D15S8). Nucleic Acids Res. 14: 4376, 1986.

[17209] 1636.

[17210] Jaiswal, A. K.; Nebert, D. W.; Gonzalez, F. J.: Human P(3)450:

[17211] cDNA and complete amino acid sequence. Nucleic Acids Res. 14: 6773-6774,

[17212] 1986.

[17213] 1637.

[17214] Jones, S. N.; Jones, P. G.; Ibarguen, H.; Caskey, C. T.; Craigen,

[17215] W. J.: Induction of the Cyp1a-1 dioxin-responsive en-

hancer in transgenic
[17216] mice. Nucleic Acids Res. 19: 6547-6551, 1991.

[17217] 1638.

[17218] Kawajiri, K.; Eguchi, H.; Nakachi, K.; Sekiya, T.; Yamamoto, M.
[17219] : Association of CYP1A1 germ line polymorphisms with mutations of
[17220] the p53 gene in lung cancer. Cancer Res. 56: 72-76, 1996.
[17221] 1639.

[17222] Kawajiri, K.; Nakachi, K.; Imai, K.; Yoshii, A.; Shinoda, N.;

[17223] Watanabe, J.: Identification of genetically high risk individuals
[17224] to lung cancer by DNA polymorphisms of the cytochrome P450IA1 gene. FEBS
[17225] Lett. 263: 131-133, 1990.

[17226] 1640.

[17227] Kawajiri, K.; Watanabe, J.; Gotoh, O.; Tagashira, Y.; Sogawa,
[17228] K.; Fujii-Kuriyama, Y.: Structure and drug inducibility of the human

[17229] cytochrome P-450c gene. Europ. J. Biochem. 159:
219-225, 1986.

[17230] 1641.

[17231] Kouri, R. E.; McKinney, C. E.; Slomiany, D. J.; Snodgrass, D.

[17232] R.; Wray, N. P.; McLemore, T. L.: Positive correlation be-
tween high

[17233] aryl hydrocarbon hydroxylase activity and primary lung
cancer as analyzed

[17234] in cryopreserved lymphocytes. Cancer Res. 42:
5030-5037, 1982.

[17235] 1642.

[17236] McBride, O. W.: Personal Communication. Bethesda, Md.
9/16/1985.

[17237] 1643.

[17238] Mooney, L. A.; Bell, D. A.; Santella, R. M.; Van Bennekum,
A.

[17239] M.; Ottman, R.; Paik, M.; Blaner, W. S.; Lucier, G. W.;
Covey, L.;

[17240] Young, T. L.; Cooper, T. B.; Glassman, A. H.; Perera, F. P.:
Contribution

[17241] of genetic and nutritional factors to DNA damage in heavy
smokers. Carcinogenesis 18:

[17242] 503-509, 1997.

[17243] 1644.

[17244] Nakachi, K.; Imai, K.; Hayashi, S.; Watanabe, J.; Kawajiri, K.

[17245] : Genetic susceptibility to squamous cell carcinoma of the lung in

[17246] relation to cigarette smoking dose. Cancer Res. 51: 5177-5180, 1991.

[17247] 1645.

[17248] Nebert, D. W.: Personal Communication. Bethesda, Md. 2/1/1988.

[17249] 1646.

[17250] Ocraft, K. P.; Muskett, J. M.; Brown, S.: Localization of the

[17251] human arylhydrocarbon hydroxylase gene to the 2q31-2pter region of

[17252] chromosome 2. Ann. Hum. Genet. 49: 237-239, 1985.

[17253] 1647.

[17254] Perera, F. P.: Environment and cancer: who are suscepti-

ble? Science 278:

[17255] 1068-1073, 1997.

[17256] 1648.

[17257] Petersen, D. D.; McKinney, C. E.; Ikeya, K.; Smith, H. H.; Bale,
[17258] A. E.; McBride, O. W.; Nebert, D. W.: Human CYP1A1 gene: cosegregation
[17259] of the enzyme inducibility phenotype and an RFLP. Am. J. Hum. Genet. 48:
[17260] 720-725, 1991.

[17261] 1649.

[17262] Quattrochi, L. C.; Okino, S. T.; Pendurthi, U. R.; Tukey, R. H.
[17263] : Cloning and isolation of human cytochrome P-450 cD-NAs homologous
[17264] to dioxin-inducible rabbit mRNAs encoding P-450 4 and P-450 6. DNA 4:
[17265] 395-400, 1985.

[17266] 1650.

[17267] Tukey, R. H.; Lalley, P. A.; Nebert, D. W.: Localization of cytochrome
[17268] P1-450 and P3-450 genes to mouse chromosome 9. Proc. Nat. Acad. Sci. 81:
[17269] 3163-3166, 1984.

[17270] 1651.

[17271] Wang, X.; Zuckerman, B.; Pearson, C.; Kaufman, G.; Chen, C.; Wang,

[17272] G.; Niu, T.; Wise, P. H.; Bauchner, H.; Xu, X.: Maternal cigarette

[17273] smoking, metabolic gene polymorphism, and infant birth weight. J.A.M.A. 287:

[17274] 195-202, 2002.

[17275] 1652.

[17276] Wiebel, F. J.; Hlavica, P.; Grzeschik, K. H.: Expression of aromatic

[17277] polycyclic hydrocarbon-induced monooxygenase (aryl hydrocarbon hydroxylase)

[17278] in man-mouse hybrids is associated with human chromosome 2. Hum.

[17279] Genet. 59: 277-280, 1981.

[17280] 1653.

[17281] Xu, X.; Kelsey, K. T.; Wiencke, J. K.; Wain, J. C.; Christiani,

[17282] D. C.: Cytochrome P450 CYP1A1 MspI polymorphism and lung cancer susceptibility. Cancer

[17283] Epidemiol. Biomarkers Prev. 5: 687-692, 1996.

[17284] 1654.

[17285] Berne, R. M.: Cardiac nucleotides in hypoxia: possible role in

[17286] regulation of coronary blood flow. Am. J. Physiol. 204: 317-322,

[17287] 1963.

[17288] 1655.

[17289] Harper, M. E.; Dugaiczyk, A.: Linkage of the evolutionar-ily-related

[17290] serum albumin and alpha-fetoprotein genes within q11-22 of human chromosome

[17291] 4. Am. J. Hum. Genet. 35: 565-572, 1983.

[17292] 1656.

[17293] Mohandas, N.; Winardi, R.; Knowles, D.; Leung, A.; Parra, M.;

[17294] George, E.; Conboy, J.; Chasis, J.: Molecular basis for membrane

[17295] rigidity of hereditary ovalocytosis: a novel mechanism in-volving the

[17296] cytoplasmic domain of band 3. J. Clin. Invest. 89: 686-692, 1992.

[17297] 1657.

[17298] Livingstone, F. B.: The Duffy blood groups, vivax malaria, and

[17299] malaria selection in human populations: a review. Hum. Biol. 56:

[17300] 413-425, 1984.

[17301] 1658.

[17302] Lefort, A.; Passage, E.; Libert, F.; Szpirer, J.; Vassart, G.;

[17303] Mattei, M.-G.: Localization of human calcyphosine gene (CAPS) to

[17304] the p13.3 region of chromosome 19 by in situ hybridization. Cytogenet.

[17305] Cell Genet. 54: 154-155, 1990.

[17306] 1659.

[17307] Yang-Feng, T. L.; Naiman, T.; Kopatz, I.; Eli, D.; Dafni, N.; Canaani,

[17308] D.: Assignment of the human casein kinase II alpha-prime subunit

[17309] gene (CSNK2A1) to chromosome 16p13.2-p13.3. Genomics 19: 173 only,

[17310] 1994.

[17311] 1660.

[17312] Bonner, T. I.; Young, A. C.; Brann, M. R.; Buckley, N. J.: Cloning

[17313] and expression of the human and rat m5 muscarinic acetylcholine genes. Neuron 1:

[17314] 403-410, 1988.

[17315] 1661.

[17316] Rosipal, R.; Lamoril, J.; Puy, H.; Da Silva, V.; Gouya, L.; De

[17317] Rooij, F. W. M.; Te Velde, K.; Nordmann, Y.; Martasek, P.; Deybach,

[17318] J. C.: Systematic analysis of coproporphyrinogen oxidase gene defects

[17319] in hereditary coproporphyria and mutation update. Hum. Mutat. 13:

[17320] 44-53, 1999.

[17321] 1662.

[17322] Jones, M. E. E.; Thorburn, A. W.; Britt, K. L.; Hewitt, K. N.;

[17323] Wreford, N. G.; Proietto, J.; Oz, O. K.; Leury, B. J.; Robert-son,

[17324] K. M.; Yao, S.; Simpson, E. R.: Aromatase-deficient (ArKO) mice have

[17325] a phenotype of increased adiposity. Proc. Nat. Acad. Sci. 97: 12735-12740,

[17326] 2000.

[17327] 1663.

[17328] Sakai, T.; Johnson, K. J.; Murozono, M.; Sakai, K.; Magnuson,

[17329] M. A.; Wieloch, T.; Cronberg, T.; Isshiki, A.; Erickson, H. P.; Fassler,

[17330] R.: Plasma fibronectin supports neuronal survival and reduces brain

[17331] injury following transient focal cerebral ischemia but is not essential

[17332] for skin-wound healing and hemostasis. Nature Med. 7: 324-330, 2001.

[17333] 1664.

[17334] Dausset, J.; Ivanyi, P.; Colombani, J.; Feingold, N.; Legrand,

[17335] L.: Le systeme Hu-1: etudes genetiques de population et de familles. Nouv.

[17336] Rev. Franc. Hemat. 7: 897-899, 1967.

[17337] 1665.

[17338] Geurts, J. M.; Schoenmakers, E. F.; Roijer, E.; Stenman, G.;

Van

[17339] de Ven, W. J. M.: Expression of reciprocal hybrid transcripts of
[17340] HMGIC and FHIT in a pleomorphic adenoma of the parotid gland. Cancer
[17341] Res. 57: 13-17, 1997.

[17342] 1666.

[17343] Tan, J. C.; Indelicato, S. R.; Narula, S. K.; Zavodny, P. J.; Chou,
[17344] C.-C.: Characterization of interleukin-10 receptors on human and
[17345] mouse cells. J. Biol. Chem. 268: 21053-21059, 1993.

[17346] 1667.

[17347] Michaud, J.; Wu, F.; Osato, M.; Cottles, G. M.; Yanagida, M.;
[17348] Asou, N.; Shigesada, K.; Ito, Y.; Benson, K. F.; Raskind, W. H.; Rossier,
[17349] C.; Antonarakis, S. E.; Israels, S.; McNicol, A.; Weiss, H.; Horwitz,
[17350] M.; Scott, H. S.: In vitro analyses of known and novel RUNX1/AML1
[17351] mutations in dominant familial platelet disorder with predisposition

[17352] to acute myelogenous leukemia: implications for mechanisms of pathogenesis. Blood 99:
[17353] 1364-1372, 2002.

[17354] 1668.

[17355] Baumann, P.; West, S. C.: DNA end-joining catalyzed by human cell-free
[17356] extracts. Proc. Nat. Acad. Sci. 95: 14066-14070, 1998.

[17357] 1669.

[17358] Lage, H.; Dietel, M.: Cloning of a human cDNA encoding a protein
[17359] with high homology to yeast methionyl-tRNA synthetase. Gene 178:
[17360] 187-189, 1996.

[17361] 1670.

[17362] O'Hara, B.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Shows,
[17363] T. B.; Eddy, R. L.; Bohlen, P.; Kovesdi, I.: Chromosomal assignment
[17364] of the heparin-binding cytokine genes MDK and PTN in mouse and man. Cytogenet.
[17365] Cell Genet. 69: 40-43, 1995.

[17366] 1671.

[17367] Tsukamoto, A. S.; Grosschedl, R.; Guzman, R. C.; Parslow, T.; Varmus,

[17368] H. E.: Expression of the int-1 gene in transgenic mice is associated

[17369] with mammary gland hyperplasia and adenocarcinomas in male and female

[17370] mice. Cell 55: 619-625, 1988.

[17371] 1672.

[17372] Kawagishi, J.; Kumabe, T.; Yoshimoto, T.; Yamamoto, T.: Structure,

[17373] organization, and transcription units of the human alpha-platelet-derived

[17374] growth factor receptor gene, PDGFRA. Genomics 30: 224-232, 1995.

[17375] 1673.

[17376] Watts, J. D.; Brabb, T.; Bures, E. J.; Wange, R. L.; Samelson,

[17377] L. E.; Aebersold, R.: Identification and characterization of a substrate

[17378] specific for the T cell protein tyrosine kinase ZAP-70. FEBS Lett. 398:

[17379] 217-222, 1996.

[17380] 1674.

[17381] Pearson, C. A.; Pearson, D.; Shibahara, S.; Hofsteenge, J.; Chiquet-Ehrismann,

[17382] R.: Tenascin: cDNA cloning and induction by TGF-beta. EMBO J. 7:

[17383] 2977-2981, 1988.

[17384] 1675.

[17385] Abou-Khalil, B.; Ge, Q.; Desai, R.; Ryther, R.; Bazyk, A.; Bailey,

[17386] R.; Haines, J. L.; Sutcliffe, J. S.; George, A. L., Jr.: Partial

[17387] and generalized epilepsy with febrile seizures plus and a novel SCN1A

[17388] mutation. Neurology 57: 2265-2272, 2001.

[17389] 1676.

[17390] Baulac, S.; Gourfinkel-An, I.; Picard, F.; Rosenberg-Bour-gin, M.;

[17391] Prud'homme, J.-F.; Baulac, M.; Brice, A.; LeGuern, E.: A second locus

[17392] for familial generalized epilepsy with febrile seizures plus maps

[17393] to chromosome 2q21-q33. Am. J. Hum. Genet. 65: 1078-1085, 1999.

[17394] 1677.

[17395] Blanchard, B. J.; Ingram, V. M.: The gene for the sodium channel

[17396] I-alpha subunit is located on chromosome 21. (Abstract) Cytogenet.

[17397] Cell Genet. 58: 2034 only, 1991.

[17398] 1678.

[17399] Claes, L.; Del-Favero, J.; Ceulemans, B.; Lagae, L.; Van Broeckhoven,

[17400] C.; De Jonghe, P.: De novo mutations in the sodium-channel gene SCN1A

[17401] cause severe myoclonic epilepsy of infancy. Am. J. Hum. Genet. 68:

[17402] 1327-1332, 2001.

[17403] 1679.

[17404] Escayg, A.; Heils, A.; MacDonald, B. T.; Haug, K.; Sander, T.;

[17405] Meisler, M. H.: A novel SCN1A mutation associated with generalized

[17406] epilepsy with febrile seizures plus--and prevalence of variants in
[17407] patients with epilepsy. Am. J. Hum. Genet. 68: 866-873, 2001.
[17408] 1680.

[17409] Escayg, A.; MacDonald, B. T.; Meisler, M. H.; Baulac, S.; Huberfeld,
[17410] G.; An-Gourfinkel, I.; Brice, A.; LeGuern, E.; Moulard, B.; Chaigne,
[17411] D.; Buresi, C.; Malafosse, A.: Mutations of SCN1A, encoding a neuronal
[17412] sodium channel, in two families with GEFS+2. Nature Genet. 24: 343-345,
[17413] 2000.

[17414] 1681.

[17415] Goldin, A. L.; Snutch, T.; Lubbert, H.; Dowsett, A.; Marshall,
[17416] J.; Auld, V.; Downey, W.; Fritz, L. C.; Lester, H. A.; Dunn, R.; Catterall,
[17417] W. A.; Davidson, N.: Messenger RNA coding for only the alpha subunit
[17418] of the rat brain Na channel is sufficient for expression of

functional
[17419] channels in Xenopus oocytes. Proc. Nat. Acad. Sci. 83:
7503-7507,
[17420] 1986.

[17421] 1682.

[17422] Isom, L. L.: The role of sodium channels in cell adhesion.
Front.
[17423] Biosci. 7: 12-23, 2002.

[17424] 1683.

[17425] Lossin, C.; Wang, D. W.; Rhodes, T. H.; Vanoye, C. G.;
George,
[17426] A. L., Jr.: Molecular basis of an inherited epilepsy. Neuron
34:
[17427] 877-884, 2002.

[17428] 1684.

[17429] Malo, D.; Schurr, E.; Dorfman, J.; Canfield, V.; Levenson,
R.;
[17430] Gros, P.: Three brain sodium channel alpha-subunit genes
are clustered
[17431] on the proximal segment of mouse chromosome 2. Ge-
nomics 10: 666-672,

[17432] 1991.

[17433] 1685.

[17434] Malo, M.: Personal Communication. Pearl River, N. Y. 12/6/1993.

[17435] 1686.

[17436] Malo, M. S.; Blanchard, B. J.; Andresen, J. M.; Srivastava, K.;

[17437] Chen, X.-N.; Li, X.; Jabs, E. W.; Korenberg, J. R.; Ingram, V. M.

[17438] : Localization of a putative human brain sodium channel gene (SCN1A)

[17439] to chromosome band 2q24. Cytogenet. Cell Genet. 67: 178-186, 1994.

[17440] 1687.

[17441] Malo, M. S.; Srivastava, K.; Andresen, J. M.; Chen, X.-N.; Korenberg,

[17442] J. R.; Ingram, V. M.: Targeted gene walking by low strin-gency polymerase

[17443] chain reaction: assignment of a putative human brain sodium channel

[17444] gene (SCN3A) to chromosome 2q24-31. Proc. Nat. Acad. Sci. 91: 2975-2979,

[17445] 1994.

[17446] 1688.

[17447] Dickinson, D. P.; Ridall, A. L.; Levine, M. J.: Human sub-
mandibular

[17448] gland statherin and basic histidine-rich peptide are en-
coded by highly

[17449] abundant mRNA's derived from a common ancestral se-
quence. Biochem.

[17450] Biophys. Res. Commun. 149: 784-790, 1987.

[17451] 1689.

[17452] Sabatini, L. M.; Carlock, L. R.; Johnson, G. W.; Azen, E. A.:

[17453] cDNA cloning and chromosomal localization of a gene for
statherin,

[17454] a regulator of calcium in saliva. (Abstract) Am. J. Hum.
Genet. 39:

[17455] A217 only, 1986.

[17456] 1690.

[17457] Sabatini, L. M.; Carlock, L. R.; Johnson, G. W.; Azen, E. A.:

[17458] cDNA cloning and chromosomal localization (4q11-13) of
a gene for

[17459] statherin, a regulator of calcium in saliva. Am. J. Hum. Genet. 41:

[17460] 1048-1060, 1987.

[17461] 1691.

[17462] Carey, A. H.; Chan, K. L.; Short, F.; White, D.; Williamson, R.;

[17463] Franks, S.: Evidence for a single gene effect causing poly-cystic

[17464] ovaries and male pattern baldness. Clin. Endocr. 38: 653-658, 1993.

[17465] 1692.

[17466] Carey, A. H.; Waterworth, D.; Patel, K.; White, D.; Little, J.;

[17467] Novelli, P.; Franks, S.; Williamson, R.: Polycystic ovaries and premature

[17468] male pattern baldness are associated with one allele of the steroid

[17469] metabolism gene CYP17. Hum. Molec. Genet. 3: 1873-1876, 1994.

[17470] 1693.

[17471] Ikeda, H.; Mitani, T.; Ohnuma, M.; Haga, H.; Ohtzuka, S.; Kato,

[17472] T.; Nakase, T.; Sekiguchi, S.: A new platelet-specific antigen, Nak(a),

[17473] involved in the refractoriness of HLA-matched platelet transfusion. Vox

[17474] Sang. 57: 213-217, 1989.

[17475] 1694.

[17476] Kashiwagi, H.; Honda, S.; Tomiyama, Y.; Mizutani, H.; Take, H.;

[17477] Honda, Y.; Kosugi, S.; Kanayama, Y.; Kurata, Y.; Matsuzawa, Y.: A

[17478] novel polymorphism in glycoprotein IV (replacement of proline-90 by

[17479] serine) predominates in subjects with platelet GPIV deficiency. Thromb.

[17480] Haemost. 69: 481-484, 1993.

[17481] 1695.

[17482] Kashiwagi, H.; Tomiyama, Y.; Honda, S.; Kosugi, S.; Shiraga, M.;

[17483] Nagao, N.; Sekiguchi, S.; Kanayama, Y.; Kurata, Y.; Matsuzawa, Y.

[17484] : Molecular basis of CD36 deficiency: evidence that a 478-C-to-T substitution

[17485] (proline90-to-serine) in CD36 cDNA accounts for CD36 deficiency. J.

[17486] Clin. Invest. 95: 1040-1046, 1995.

[17487] 1696.

[17488] Kashiwagi, H.; Tomiyama, Y.; Nozaki, S.; Kiyoi, T.; Ta-

dokoro,

[17489] S.; Matsumoto, K.; Honda, S.; Kosugi, S.; Kurata, Y.; Mat-

suzawa, Y.

[17490] : Analyses of genetic abnormalities in type I CD36 defi-

ciency in Japan:

[17491] identification and cell biological characterization of two novel mutations

[17492] that cause CD36 deficiency in man. Hum. Genet. 108: 459-466, 2001.

[17493] 1697.

[17494] Lee, K.; Godeau, B.; Fromont, P.; Plonquet, A.; Debili, N.; Bachir,

[17495] D.; Reviron, D.; Gourin, J.; Fernandez, E.; Galacteros, F.; Bierling,

[17496] P.: CD36 deficiency is frequent and can cause platelet im-

munization

[17497] in Africans. Transfusion 39: 873-879, 1999.

[17498] 1698.

[17499] Pravenec, M.; Landa, V.; Zidek, V.; Musilova, A.; Kren, V.; Kazdova,

[17500] L.; Aitman, T. J.; Glazier, A. M.; Ibrahimi, A.; Abumrad, N. A.; Qi,

[17501] N.; Wang, J.-M.; St. Lezin, E. M.; Kurtz, T. W.: Transgenic rescue

[17502] of defective Cd36 ameliorates insulin resistance in spontaneously

[17503] hypertensive rats. Nature Genet. 27: 156-158, 2001.

[17504] 1699.

[17505] Pravenec, M.; Zidek, V.; Simakova, M.; Kren, V.; Krenova, D.;

[17506] Horky, K.; Jachymova, M.; Mikova, B.; Kazdova, L.; Aitman, T. J.;

[17507] Churchill, P. C.; Webb, R. C.; Hingarh, N. H.; Yang, Y.; Wang, J.-M.;

[17508] St. Lezin, E. M.; Kurtz, T. W.: Genetics of Cd36 and the clustering

[17509] of multiple cardiovascular risk factors in spontaneous hypertension. J.

[17510] Clin. Invest. 103: 1651-1657, 1999.

[17511] 1700.

[17512] Reaven, G. M.; Lithell, H.; Landsberg, L.: Hypertension and associated

[17513] metabolic abnormalities: the role of insulin resistance and the sympathoadrenal

[17514] system. New Eng. J. Med. 334: 374-381, 1996.

[17515] 1701.

[17516] Savill, J.; Hogg, N.; Ren, Y.; Haslett, C.: Thrombospondin cooperates

[17517] with CD36 and the vitronectin receptor in macrophage recognition of

[17518] neutrophils undergoing apoptosis. J. Clin. Invest. 90: 1513-1522,

[17519] 1992.

[17520] 1702.

[17521] Tanaka, T.; Sohmiya, K.; Kawamura, K.: Is CD36 deficiency an

[17522] etiology of hereditary hypertrophic cardiomyopathy? J. Molec. Cell.

[17523] Cardiol. 29: 121-127, 1997.

[17524] 1703.

[17525] Tandon, N. N.; Kralisz, U.; Jamieson, G. A.: Identification of
[17526] glycoprotein IV (CD36) as a primary receptor for platelet-collagen
[17527] adhesion. J. Biol. Chem. 264: 7576-7583, 1989.

[17528] 1704.

[17529] Tandon, N. N.; Lipsky, R. H.; Burgess, W. H.; Jamieson, G. A.
[17530] : Isolation and characterization of platelet glycoprotein IV (CD36). J.
[17531] Biol. Chem. 264: 7570-7575, 1989.

[17532] 1705.

[17533] Tomiyama, Y.; Take, H.; Ikeda, H.; Mitani, T.; Furubayashi, T.;
[17534] Mizutani, H.; Yamamoto, N.; Tandon, N. N.; Sekiguchi, S.; Jamieson,
[17535] G. A.; Kurata, Y.; Yonezawa, T.; Tarui, S.: Identification of the
[17536] platelet-specific alloantigen, Nak(a), on platelet mem-brane glycoprotein
[17537] IV. Blood 75: 684-687, 1990.

[17538] 1706.

[17539] Urwijitaroon, Y.; Barusrux, S.; Romphruk, A.; Puapairoj, C.:

[17540] Frequency of human platelet antigens among blood donors in northeastern

[17541] Thailand. Transfusion 35: 868-870, 1995.

[17542] 1707.

[17543] van Schravendijk, M. R.; Handunnetti, S. M.; Barnwell, J. W.;

[17544] Howard, R. J.: Normal human erythrocytes express CD36, an adhesion

[17545] molecule of monocytes, platelets, and endothelial cells. Blood 80:

[17546] 2105-2114, 1992.

[17547] 1708.

[17548] Yamamoto, N.; Akamatsu, N.; Sakuraba, H.; Yamazaki, H.; Tanoue,

[17549] K.: Platelet glycoprotein IV (CD36) deficiency is associated with

[17550] the absence (type I) or the presence (type II) of glycopro-tein IV

[17551] on monocytes. Blood 83: 392-397, 1994.

[17552] 1709.

[17553] Yamamoto, N.; Ikeda, H.; Tandon, N. N.; Herman, J.; Tomiyama,

[17554] Y.; Mitani, T.; Sekiguchi, S.; Lipsky, R.; Kralisz, U.; Jamieson,

[17555] G. A.: A platelet membrane glycoprotein (GP) deficiency in healthy

[17556] blood donors: Nak(a-) platelets lack detectable GPIV (CD36). Blood 76:

[17557] 1698-1703, 1990.

[17558] 1710.

[17559] Yufu, Y.; Ideguchi, H.; Narishige, T.; Suematsu, E.; Toyoda, K.;

[17560] Nishimura, J.; Nawata, H.; Oda, S.: Familial macrothrom-bocytopenia

[17561] associated with decreased glycosylation of platelet mem-brane glycoprotein

[17562] IV. Am. J. Hemat. 33: 271-273, 1990.

[17563] 1711.

[17564] Clemetson, J. M.; Kyrle, P. A.; Brenner, B.; Clemetson, K. J.:

[17565] Variant Bernard-Soulier syndrome associated with a ho-

mozygous mutation
[17566] in the leucine-rich domain of glycoprotein IX. Blood 84: 1124-1131,
[17567] 1994.

[17568] 1712.

[17569] Hickey, M. J.; Deaven, L. L.; Roth, G. J.: Human platelet glycoprotein
[17570] IX: characterization of cDNA and localization of the gene to chromosome
[17571] 3. FEBS Lett. 274: 189-192, 1990.

[17572] 1713.

[17573] Hickey, M. J.; Roth, G. J.: Characterization of the gene encoding
[17574] human platelet glycoprotein IX. (Abstract) Clin. Res. 39: 337A only,
[17575] 1991.

[17576] 1714.

[17577] Tunnacliffe, A.; McGuire, R. S.: A physical linkage group in
[17578] human chromosome band 11q23 covering a region implicated in leukocyte
[17579] neoplasia. Genomics 8: 447-453, 1990.

[17580] 1715.

[17581] Lomedico, P.; Rosenthal, N.; Efstratiadis, A.; Gilbert, W.; Koladner,

[17582] R.; Tizard, R.: The structure and evolution of the two non-allelic

[17583] rat preproinsulin genes. Cell 18: 545-558, 1979.

[17584] 1716.

[17585] Allore, R.; O'Hanlon, D.; Price, R.; Neilson, K.; Willard, H. F.;

[17586] Cox, D. R.; Marks, A.; Dunn, R. J.: Gene encoding the beta-subunit

[17587] of S100 protein is on chromosome 21: implications for Down syndrome. Science 239:

[17588] 1311-1313, 1988.

[17589] 1717.

[17590] Lafage-Pochitaloff-Huvale, M.; Shaw, A.; Dessarps, F.; Mannoni,

[17591] P.; Fradelizi, D.; Wollman, E. E.: The gene for human thioredoxin

[17592] maps on the short arm of chromosome 3 at bands 3p11-p12. FEBS Lett. 255:

[17593] 89-91, 1989.

[17594] 1718.

[17595] Mandrup-Poulsen, T.; Owerbach, D.; Mortensen, S. A.; Jo-
hansen,

[17596] K.; Meinertz, H.; Sorensen, H.; Nerup, J.: DNA sequences
flanking

[17597] the insulin gene on chromosome 11 confer risk of
atherosclerosis. Lancet I:

[17598] 250-254, 1984.

[17599] 1719.

[17600] Meyers, D. A.; Beaty, T. H.; Maestri, N. E.; Antonarakis, S.
E.;

[17601] Kazazian, H. H., Jr.: Multipoint mapping studies of the
beta-globin,

[17602] insulin and C-HA-RAS-1 loci on 11p. (Letter) Am. J. Hum.
Genet. 39:

[17603] 539-541, 1986.

[17604] 1720.

[17605] Michalova, K.; Bucchini, D.; Ripoche, M.-A.; Pictet, R.;
Jami,

[17606] J.: Chromosome localization of the human insulin gene in

transgenic

[17607] mouse lines. Hum. Genet. 80: 247-252, 1988.

[17608] 1721.

[17609] Nanjo, K.; Sanke, T.; Miyano, M.; Okai, K.; Sowa, R.;
Kondo, M.;

[17610] Nishimura, S.; Iwo, K.; Miyamura, K.; Given, B. D.; Chan, S.
J.; Tager,

[17611] H. S.; Steiner, D. F.; Rubenstein, A. H.: Diabetes due to se-
cretion

[17612] of a structurally abnormal insulin (insulin Wakayama):
clinical and

[17613] functional characteristics of (leu-A3) insulin. J. Clin. In-
vest. 77:

[17614] 514-519, 1986.

[17615] 1722.

[17616] Olansky, L.; Welling, C.; Giddings, S.; Adler, S.; Bourey, R.;

[17617] Dowse, G.; Serjeantson, S.; Zimmet, P.; Permutt, M. A.: A
variant

[17618] insulin promoter in non-insulin-dependent diabetes mel-
litus. J. Clin.

[17619] Invest. 89: 1596-1602, 1992.

[17620] 1723.

[17621] Osawa, H.; Onuma, H.; Murakami, A.; Ochi, M.; Nishimiya, T.; Kato,

[17622] K.; Shimizu, I.; Fujii, Y.; Ohashi, J.; Makino, H.: Systematic search

[17623] for single nucleotide polymorphisms in the insulin gene: evidence

[17624] for a high frequency of -23T-A in Japanese subjects. Biochem. Biophys.

[17625] Res. Commun. 286: 451-455, 2001.

[17626] 1724.

[17627] Owerbach, D.; Bell, G. I.; Rutter, W. J.; Shows, T. B.: The in-sulin

[17628] gene is located on chromosome 11 in human. Nature 286: 82-84, 1980.

[17629] 1725.

[17630] Pugliese, A.; Zeller, M.; Fernandez, A., Jr.; Zalcberg, L. J.;

[17631] Bartlett, R. J.; Ricordi, C.; Pietropaolo, M.; Eisenbarth, G. S.;

[17632] Bennett, S. T.; Patel, D. D.: The insulin gene is transcribed in

[17633] the human thymus and transcription levels correlate with

allelic variation
[17634] at the INS VNTR-IDDM2 susceptibility locus for type 1 diabetes. Nature 15:
[17635] 293-297, 1997.

[17636] 1726.

[17637] Robbins, D. C.; Blix, P. M.; Rubenstein, A. H.; Kanazawa, Y.;
[17638] Kosaka, K.; Tager, H. S.: A human proinsulin variant at arginine
[17639] 65. Nature 291: 679-681, 1981.

[17640] 1727.

[17641] Robbins, D. C.; Shoelson, S. E.; Rubenstein, A. H.; Tager, H.
[17642] S.: Familial hyperproinsulinemia: two cohorts secreting indistinguishable
[17643] type II intermediates of proinsulin conversion. J. Clin. Invest. 73:
[17644] 714-719, 1984.

[17645] 1728.

[17646] Robbins, D. C.; Tager, H. S.; Rubenstein, A. H.: Biologic and

[17647] clinical importance of proinsulin. New Eng. J. Med. 310:
1165-1175,
[17648] 1984.

[17649] 1729.

[17650] Roder, M. E.; Vissing, H.; Nauck, M. A.: Hyperproinsuline-
mia
[17651] in a three-generation Caucasian family due to mutant
proinsulin (Arg(65)->His)
[17652] not associated with impaired glucose tolerance: The con-
tribution of
[17653] mutant proinsulin to insulin bioactivity. J. Clin. Endocr.
Metab. 81:
[17654] 1634-1640, 1996.

[17655] 1730.

[17656] Rotwein, P.; Chyn, R.; Chirgwin, J.; Cordell, B.; Goodman,
H.
[17657] M.; Permutt, M. A.: Polymorphism in the 5-prime-flanking
region of
[17658] the human insulin gene and its possible relation to type 2
diabetes. Science 213:
[17659] 1117-1120, 1981.

[17660] 1731.

[17661] Rotwein, P.; Yokoyama, S.; Didier, D. K.; Chirgwin, J. M.: Genetic

[17662] analysis of the hypervariable region flanking the human insulin gene. Am.

[17663] J. Hum. Genet. 39: 291-299, 1986.

[17664] 1732.

[17665] Rubenstein, A. H.: Personal Communication. Chicago, Ill. 6/17/1983.

[17666] 1733.

[17667] Sakura, H.; Iwamoto, Y.; Sakamoto, Y.; Kuzuya, T.; Hirata, H.

[17668] : Structurally abnormal insulin in a diabetic patient: characterization

[17669] of the mutant insulin A3 (val-to-leu) isolated from the pancreas. J.

[17670] Clin. Invest. 78: 1666-1672, 1986.

[17671] 1734.

[17672] Schwartz, G. P.; Burke, G. T.; Katsoyannis, P. G.: A super-active

[17673] insulin: [B10-aspartic acid]insulin(human). Proc. Nat.

Acad. Sci. 84:
[17674] 6408-6411, 1987.

[17675] 1735.

[17676] Seino, S.; Funakoshi, A.; Fu, Z. Z.; Vinik, A.: Identification
[17677] of insulin variants in patients with hyperinsulinemia by re-
versed-phase
[17678] high-performance liquid chromatography. Diabetes 34:
1-7, 1985.
[17679] 1736.

[17680] Selden, R. F.; Skoskiewicz, M. J.; Russell, P. S.; Goodman,
H.
[17681] M.: Regulation of insulin-gene expression: implications
for gene
[17682] therapy. New Eng. J. Med. 317: 1067-1076, 1987.

[17683] 1737.

[17684] Shibasaki, Y.; Kawakami, T.; Kanazawa, Y.; Akanuma, Y.;
Takaku,
[17685] F.: Posttranslational cleavage of proinsulin is blocked by a
point
[17686] mutation in familial hyperproinsulinemia. J. Clin. Invest.
76: 378-380,

[17687] 1985.

[17688] 1738.

[17689] Shoelson, S.; Fickova, M.; Haneda, M.; Nahum, A.; Musso, G.; Kaiser,

[17690] E. T.; Rubenstein, A. H.; Tager, H.: Identification of a mutant human

[17691] insulin predicted to contain a serine-for-phenylalanine substitution. Proc.

[17692] Nat. Acad. Sci. 80: 7390-7394, 1983.

[17693] 1739.

[17694] Shoelson, S.; Haneda, M.; Blix, P.; Nanjo, A.; Sanke, T.; Inouye,

[17695] K.; Steiner, D.; Rubenstein, A.; Tager, H.: Three mutant insulins

[17696] in man. Nature 302: 540-543, 1983.

[17697] 1740.

[17698] Soares, M. B.; Schon, E.; Henderson, A.; Karathanasis, S. K.;

[17699] Cate, R.; Zeitlin, S.; Chirgwin, J.; Efstratiadis, A.: RNA-mediated

[17700] gene duplication: the rat preproinsulin I gene is a func-

tional retroposon. Molec.
[17701] Cell. Biol. 5: 2090-2103, 1985.

[17702] 1741.

[17703] Steiner, D. F.: Errors in insulin biosynthesis. (Editorial) New

[17704] Eng. J. Med. 294: 952-953, 1976.

[17705] 1742.

[17706] Steiner, D. F.; Chan, S. J.; Welsh, J. M.; Kwok, S. C. M.: Structure
[17707] and evolution of the insulin gene. Ann. Rev. Genet. 19: 463-484,
[17708] 1985.

[17709] 1743.

[17710] Steiner, D. F.; Oyer, P. E.: The biosynthesis of insulin and

[17711] a probable precursor of insulin by a human islet cell ade-
noma. Proc.
[17712] Nat. Acad. Sci. 57: 473-480, 1967.

[17713] 1744.

[17714] Sures, I.; Goeddel, D. V.; Gray, A.; Ullrich, A.: Nucleotide

[17715] sequences of human preproinsulin complementation DNA.

Science 208:
[17716] 57-59, 1980.

[17717] 1745.

[17718] Tager, H.; Given, B.; Baldwin, D.; Mako, M.; Markese, J.; Rubenstein,
[17719] A.; Olefsky, J.; Kobayashi, M.; Kolterman, O.; Poucher, R.: A structurally
[17720] abnormal insulin causing human diabetes. Nature 281: 122-125, 1979.
[17721] 1746.

[17722] Todd, S.; Yoshida, M. C.; Fang, X. E.; McDonald, L.; Jacobs, J.;
[17723] Heinrich, G.; Bell, G. I.; Naylor, S. L.; Sakaguchi, A. Y.: Genes
[17724] for insulin I and II, parathyroid hormone, and calcitonin are on rat
[17725] chromosome 1. Biochem. Biophys. Res. Commun. 131: 1175-1180, 1985.
[17726] 1747.

[17727] Ullrich, A.; Dull, T. J.; Gray, A.: Genetic variation in the

[17728] human insulin gene. Science 209: 612-615, 1980.

[17729] 1748.

[17730] Vafiadis, P.; Bennett, S. T.; Todd, J. A.; Nadeau, J.; Grabs,

[17731] R.; Goodyer, C. G.; Wickramasinghe, S.; Colle, E.; Poly-
chronakos,

[17732] C.: Insulin expression in the human thymus is modulated
by INS VNTR

[17733] alleles at the IDDM2 locus. Nature 15: 289-292, 1997.

[17734] 1749.

[17735] Karet, F. E.; Gainza, F. J.; Gyory, A. Z.; Unwin, R. J.; Wrong,

[17736] O.; Tanner, M. J. A.; Nayir, A.; Alpay, H.; Santos, F.; Hulton,
S.

[17737] A.; Bakkaloglu, A.; Ozen, S.; Cunningham, M. J.; di Pietro,
A.; Walker,

[17738] W. G.; Lifton, R. P.: Mutations in the chloride-bicarbonate
exchanger

[17739] gene AE1 cause autosomal dominant but not autosomal
recessive distal

[17740] renal tubular acidosis. Proc. Nat. Acad. Sci. 95:
6337-6342, 1998.

[17741] 1750.

[17742] Richards, P.; Wrong, O. M.: Dominant inheritance in a

family
[17743] with familial renal tubular acidosis. Lancet II: 998-999, 1972.
[17744] 1751.
[17745] Allore, R. J.; Friend, W. C.; O'Hanlon, D.; Neilson, K. M.; Baumal,
[17746] R.; Dunn, R. J.; Marks, A.: Cloning and expression of the human S100-beta
[17747] gene. J. Biol. Chem. 265: 15537-15543, 1990.
[17748] 1752.
[17749] Duncan, A. M. V.; Higgins, J.; Dunn, R. J.; Allore, R.; Marks,
[17750] A.: Refined sublocalization of the human gene encoding the beta subunit
[17751] of the S100 protein (S100B) and confirmation of a subtle t(9;21) translocation
[17752] using in situ hybridization. Cytogenet. Cell Genet. 50: 234-235,
[17753] 1989.
[17754] 1753.
[17755] Reeves, R. H.; Yao, J.; Crowley, M. R.; Buck, S.; Zhang, X.; Yarowsky,

[17756] P.; Gearhart, J. D.; Hilt, D. C.: Astrocytosis and axonal proliferation

[17757] in the hippocampus of S100b transgenic mice. Proc. Nat. Acad. Sci. 91:

[17758] 5359-5363, 1994.

[17759] 1754.

[17760] Petrij, F.; Dauwerse, H. G.; Blough, R. I.; Giles, R. H.; van

[17761] der Smagt, J. J.; Wallerstein, R.; Maaswinkel-Mooy, P. D.; van Karnebeek,

[17762] C. D.; van Ommen, G.-J. B.; van Haeringen, A.; Rubinstein, J. H.;

[17763] Saal, H. M.; Hennekam, R. C. M.; Peters, D. J. M.; Breuning, M. H.

[17764] : Diagnostic analysis of the Rubinstein-Taybi syndrome: five cosmids

[17765] should be used for microdeletion detection and low number of protein

[17766] truncating mutations. J. Med. Genet. 37: 168-176, 2000.

[17767] 1755.

[17768] Petrij, F.; Giles, R. H.; Dauwerse, H. G.; Saris, J. J.; Hennekam,

[17769] R. C. M.; Masuno, M.; Tommerup, N.; van Ommen, G.-J. B.; Goodman,

[17770] R. H.; Peters, D. J. M.; Breuning, M. H.: Rubinstein-Taybi syndrome

[17771] caused by mutations in the transcriptional co-activator CBP. Nature 376:

[17772] 348-351, 1995.

[17773] 1756.

[17774] Wessels, J. W.; Mollevanger, P.; Dauwerse, J. G.; Cluitmans, F.

[17775] H. M.; Breuning, M. H.; Beverstock, G. C.: Two distinct loci on the

[17776] short arm of chromosome 16 are involved in myeloid leukemia. Blood 77:

[17777] 1555-1559, 1991.

[17778] 1757.

[17779] Ruttledge, M. H.; Andermann, A. A.; Phelan, C. M.; Claudio, J.

[17780] O.; Han, F.; Chretien, N.; Rangaratnam, S.; MacCollin, M.; Short,

[17781] P.; Parry, D.; Michels, V.; Riccardi, V. M.; Weksberg, R.; Kitamura,

[17782] K.; Bradburn, J. M.; Hall, B. D.; Propping, P.; Rouleau, G. A.: Type
[17783] of mutation in the neurofibromatosis type 2 gene (NF2) frequently
[17784] determines severity of disease. Am. J. Hum. Genet. 59: 331-342,
[17785] 1996.

[17786] 1758.

[17787] Citron, M.; Vigo-Pelfrey, C.; Teplow, D. B.; Miller, C.; Schenk,
[17788] D.; Johnston, J.; Winblad, B.; Venizelos, N.; Lannfelt, L.; Selkoe,
[17789] D. J.: Excessive production of amyloid beta-protein by pe-ripheral
[17790] cells of symptomatic and presymptomatic patients carry-ing the Swedish
[17791] familial Alzheimer disease mutation. Proc. Nat. Acad. Sci. 91: 11993-11997,
[17792] 1994.

[17793] 1759.

[17794] Easton, D. F.; Bishop, D. T.; Ford, D.; Crockford, G. P.; Breast

[17795] Cancer Linkage Consortium: Genetic linkage analysis in familial
[17796] breast and ovarian cancer: results from 214 families. Am. J. Hum.
[17797] Genet. 52: 678-701, 1993.

[17798] 1760.

[17799] Cloninger, C.: The psychobiological regulation of social cooperation. Nature
[17800] Med. 1: 623-625, 1995.

[17801] 1761.

[17802] Kurlan, R.; Behr, J.; Medved, L.; Shoulson, I.; Pauls, D.; Kidd,
[17803] J. R.; Kidd, K. K.: Familial Tourette's syndrome: report of a large
[17804] pedigree and potential for linkage analysis. Neurology 36: 772-776,
[17805] 1986.

[17806] 1762.

[17807] Nakae, J.; Biggs, W. H., III; Kitamura, T.; Cavenee, W. K.; Wright,
[17808] C. V. E.; Arden, K. C.; Accili, D.: Regulation of insulin ac-

tion

[17809] and pancreatic beta-cell function by mutated alleles of the gene encoding

[17810] forkhead transcription factor Foxo1. Nature Genet. 32: 245-253,

[17811] 2002.

[17812] 1763.

[17813] Kamoun, P.; Aral, B.; Saudubray, J.-M.: A new inherited metabolic

[17814] disease: delta-1-pyrroline 5-carboxylate synthetase deficiency. Bull.

[17815] Nat. Acad. Med. 182: 131-139, 1998.

[17816] 1764.

[17817] Fischer, J. A.; Egert, F.; Werder, E.; Born, W.: An inherited

[17818] mutation associated with functional deficiency of the alpha-subunit

[17819] of the guanine nucleotide-binding protein Gs in pseudo- and pseudopseudohypoparathyroidism. J.

[17820] Clin. Endocr. Metab. 83: 935-938, 1998.

[17821] 1765.

[17822] Ophoff, R. A.; Terwindt, G. M.; Vergouwe, M. N.; van Eijk,

R.;

[17823] Oefner, P. J.; Hoffman, S. M. G.; Lamerdin, J. E.; Mohren-
weiser, H.

[17824] W.; Bulman, D. E.; Ferrari, M.; Haan, J.; Lindhout, D.; van
Ommen,

[17825] G.-J. B.; Hofker, M. H.; Ferrari, M. D.; Frants, R. R.: Familial

[17826] hemiplegic migraine and episodic ataxia type-2 are
caused by mutations

[17827] in the Ca(2+) channel gene CACNL1A4. Cell 87: 543-552,
1996.

[17828] 1766.

[17829] Zhuang, Y.; Soriano, P.; Weintraub, H.: The helix-
loop-helix

[17830] gene E2A is required for B cell formation. Cell 79:
875-884, 1994.

[17831] 1767.

[17832] Gladwin, A. J.; Dixon, J.; Loftus, S. K.; Edwards, S.; Was-
muth,

[17833] J. J.; Hennekam, R. C. M.; Dixon, M. J.: Treacher Collins
syndrome

[17834] may result from insertions, deletions, or splicing muta-
tions, which

[17835] introduce a termination codon into the gene. Hum. Molec. Genet. 5:

[17836] 1533-1538, 1996.

[17837] 1768.

[17838] Haworth, K. E.; Islam, I.; Breen, M.; Putt, W.; Makrinou, E.;

[17839] Binns, M.; Hopkinson, D.; Edwards, Y.: Canine TCOF1; cloning, chromosome

[17840] assignment and genetic analysis in dogs with different head types. Mammalian

[17841] Genome 12: 622-629, 2001.

[17842] 1769.

[17843] Jabs, E. W.; Li, X.; Coss, C. A.; Taylor, E. W.; Meyers, D. A.;

[17844] Weber, J. L.: Mapping the Treacher Collins syndrome locus to 5q31.3-q33.3. Genomics 11:

[17845] 193-198, 1991.

[17846] 1770.

[17847] Jones, N. C.; Farlie, P. G.; Minichiello, J.; Newgreen, D. F.

[17848] : Detection of an appropriate kinase activity in branchial arches

[17849] I and II that coincides with peak expression of the

Treacher Collins
[17850] syndrome gene product, treacle. Hum. Mol. Genet. 8: 2239-2245, 1999.
[17851] 1771.

[17852] Neville, M. J.; Campbell, R. D.: A new member of the Ig superfamily
[17853] and a V-ATPase G subunit are among the predicted products of novel
[17854] genes close to the TNF locus in the human MHC. J. Immun. 162: 4745-4754,
[17855] 1999.

[17856] 1772.

[17857] Yang, X.-F.; Wu, C. J.; McLaughlin, S.; Chillemi, A.; Wang, K.
[17858] S.; Canning, C.; Alyea, E. P.; Kantoff, P.; Soiffer, R. J.; Dranoff,
[17859] G.; Ritz, J.: CML66, a broadly immunogenic tumor antigen, elicits
[17860] a humoral immune response associated with remission of chronic myelogenous
[17861] leukemia. Proc. Nat. Acad. Sci. 98: 7492-7497, 2001.

[17862] 1773.

[17863] Jarrous, N.; Eder, P. S.; Wesolowski, D.; Altman, S.: Rpp14 and

[17864] Rpp29, two protein subunits of human ribonuclease P. RNA 5: 153-157,

[17865] 1999.

[17866] 1774.

[17867] Jarrous, N.; Eder, P. S.; Guerrier-Takada, C.; Hoog, C.; Altman,

[17868] S.: Autoantigenic properties of some protein subunits of catalytically

[17869] active complexes of human ribonuclease P. RNA 4: 407-417, 1998.

[17870] 1775.

[17871] MacArthur, C. A.; Shackleford, G. M.: Npm3: A novel, widely expressed

[17872] gene encoding a protein related to the molecular chaperones nucleoplasmin

[17873] and nucleophosmin. Genomics 42: 137-140, 1997.

[17874] 1776.

[17875] Shackleford, G. M.; Ganguly, A.; MacArthur, C. A.: Cloning, expression

[17876] and nuclear localization of human NPM3, a member of the nucleophosmin/nucleoplasmin

[17877] family of nuclear chaperones. BMC Genomics 2: 8, 2001. Note: Electronic

[17878] Article.

[17879] 1777.

[17880] Ishiguro, H.; Tsunoda, T.; Tanaka, T.; Fujii, Y.; Nakamura, Y.;

[17881] Furukawa, Y.: Identification of AXUD1, a novel human gene induced

[17882] by AXIN1 and its reduced expression in human carcinomas of the lung,

[17883] liver, colon and kidney. Oncogene 20: 5062-5066, 2001.

[17884] 1778.

[17885] Zhu, Y.; Qi, C.; Cao, W.-Q.; Yeldandi, A. V.; Rao, M. S.; Reddy,

[17886] J. K.: Cloning and characterization of PIMT, a protein with a methyltransferase

[17887] domain, which interacts with and enhances nuclear receptor coactivator

[17888] PRIP function. Proc. Nat. Acad. Sci. 98: 10380-10385, 2001.

[17889] 1779.

[17890] Hauf, S.; Waizenegger, I. C.; Peters, J.-M.: Cohesin cleav-
age
[17891] by separase required for anaphase and cytokinesis in hu-
man cells. Science 293:
[17892] 1320-1323, 2001.

[17893] 1780.

[17894] Hoque, M. T.; Ishikawa, F.: Human chromatid cohesin
component
[17895] hRad21 is phosphorylated in M phase and associated with
metaphase
[17896] centromeres. J. Biol. Chem. 276: 5059-5067, 2001.

[17897] 1781.

[17898] McKay, M. J.; Troelstra, C.; van der Spek, P.; Kanaar, R.;
Smit,
[17899] B.; Hagemeijer, A.; Bootsma, D.; Hoeijmakers, J. H. J.: Se-
quence
[17900] conservation of the rad21 Schizosaccharomyces pombe
DNA double-strand
[17901] break repair gene in human and mouse. Genomics 36:
305-315, 1996.

[17902] 1782.

[17903] Sadano, H.; Sugimoto, H.; Sakai, F.; Nomura, N.; Osumi, T.: NXP-1,

[17904] a human protein related to Rad21/Scc1/Mcd1, is a component of the

[17905] nuclear matrix. Biochem. Biophys. Res. Commun. 267: 418-422, 2000.

[17906] 1783.

[17907] Fukamachi, S.; Shimada, A.; Shima, A.: Mutations in the gene encoding

[17908] B, a novel transporter protein, reduce melanin content in medaka. Nature

[17909] Genet. 28: 381-385, 2001.

[17910] 1784.

[17911] Harada, M.; Li, Y. F.; El-Gamil, M.; Rosenberg, S. A.; Robbins,

[17912] P. F.: Use of an in vitro immunoselected tumor line to identify shared

[17913] melanoma antigens recognized by HLA-A*0201-restricted T cells. Cancer

[17914] Res. 61: 1089-1094, 2001.

[17915] 1785.

[17916] Newton, J. M.; Cohen-Barak, O.; Hagiwara, N.; Gardner, J. M.; Davisson,

[17917] M. T.; King, R. A.; Brilliant, M. H.: Mutations in the human orthologue

[17918] of the mouse underwhite gene (uw) underlie a new form of oculocutaneous

[17919] albinism, OCA4. Am. J. Hum. Genet. 69: 981-988, 2001.

[17920] 1786.

[17921] Gu, H.; Saito, K.; Klaman, L. D.; Shen, J.; Fleming, T.; Wang,

[17922] Y.-P.; Pratt, J. C.; Lin, G.; Lim, B.; Kinet, J.-P.; Neel, B. G.:

[17923] Essential role for Gab2 in the allergic response. Nature 412: 186-190,

[17924] 2001.

[17925] 1787.

[17926] Appel, S.; Reichwald, K.; Zimmermann, W.; Reis, A.; Rosenthal,

[17927] A.; Hennies, H. C.: Identification and localization of a new human

[17928] myotubularin-related protein gene, MTMR8, on

8p22-p23. Genomics 75:
[17929] 6-8, 2001.

[17930] 1788.

[17931] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Mandel, J.-L.: The myotubularin
[17932] family: from genetic disease to phosphoinositide metabolism. Trends
[17933] Genet. 17: 221-228, 2001.

[17934] 1789.

[17935] Ariizumi, K.; Shen, G.-L.; Shikano, S.; Xu, S.; Ritter, R., III;

[17936] Kumamoto, T.; Edelbaum, D.; Morita, A.; Bergstresser, P. R.; Takashima,
[17937] A.: Identification of a novel, dendritic cell-associated molecule,
[17938] dectin-1, by subtractive cDNA cloning. J. Biol. Chem. 275: 20157-20167,
[17939] 2000.

[17940] 1790.

[17941] Brown, G. D.; Gordon, S.: A new receptor for beta-glucans. Nature 413:
[17942] 36-37, 2001.

[17943] 1791.

[17944] Hernanz-Falcon, P.; Arce, I.; Roda-Navarro, P.; Fernandez-Ruiz,

[17945] E.: Cloning of human dectin-1, a novel C-type lectin-like receptor

[17946] gene expressed on dendritic cells. Immunogenetics 53: 288-295, 2001.

[17947] 1792.

[17948] Yokota, K.; Takashima, A.; Bergstresser, P. R.; Ariizumi, K.:

[17949] Identification of a human homologue of the dendritic cell-associated

[17950] C-type lectin-1, dectin-1. Gene 272: 51-60, 2001.

[17951] 1793.

[17952] Hagen, G.; Muller, S.; Beato, M.; Suske, G.: Sp1-mediated transcriptional

[17953] activation is repressed by Sp3. EMBO J. 13: 3843-3851, 1994.

[17954] 1794.

[17955] Kalff-Suske, M.; Kunz, J.; Grzeschik, K.-H.; Suske, G.: Human

[17956] Sp3 transcriptional regulator gene (SP3) maps to chromo-

some 2q31. Genomics 37:

[17957] 410-412, 1996.

[17958] 1795.

[17959] Kingsley, C.; Winoto, A.: Cloning of GT box-binding proteins:

[17960] a novel Sp1 multigene family regulating T-cell receptor gene expression. Molec.

[17961] Cell. Biol. 12: 4251-44261, 1992.

[17962] 1796.

[17963] Carmeci, C.; Thompson, D. A.; Ring, H. Z.; Francke, U.; Weigel,

[17964] R. J.: Identification of a gene (GPR30) with homology to the G-protein-coupled

[17965] receptor superfamily associated with estrogen receptor expression

[17966] in breast cancer. Genomics 45: 607-617, 1997.

[17967] 1797.

[17968] Feng, Y.; Gregor, P.: Cloning of a novel member of the G protein-coupled

[17969] receptor family related to peptide receptors. Biochem. Biophys. Res.

[17970] Commun. 231: 651-654, 1997.

[17971] 1798.

[17972] Kvingedal, A. M.; Smeland, E. B.: A novel putative G-protein-coupled

[17973] receptor expressed in lung, heart and lymphoid tissue. FEBS Lett. 407:

[17974] 59-62, 1997.

[17975] 1799.

[17976] Owman, C.; Blay, P.; Nilsson, C.; Lolait, S. J.: Cloning of human

[17977] cDNA encoding a novel heptahelix receptor expressed in Burkitt's lymphoma

[17978] and widely distributed in brain and peripheral tissues. Biochem.

[17979] Biophys. Res. Commun. 228: 285-292, 1996.

[17980] 1800.

[17981] Takada, Y.; Kato, C.; Kondo, S.; Korenaga, R.; Ando, J.: Cloning

[17982] of cDNAs encoding G protein-coupled receptor expressed in human endothelial

[17983] cells exposed to fluid shear stress. Biochem. Biophys. Res.

Commun. 240;-737-741,

[17984] 1997.

[17985] 1801.

[17986] Cossins, J.; Dudgeon, T. J.; Catlin, G.; Gearing, A J. H.; Clements,

[17987] J. M.: Identification of MMP-18, a putative novel human matrix metalloproteinase. Biochem.

[17988] Biophys. Res. Commun. 228: 494-498, 1996.

[17989] 1802.

[17990] Pendas, A. M.; Knauper, V.; Puente, X. S.; Llano, E.; Mattei, M.-G.;

[17991] Apte, S.; Murphy, G.; Lopez-Otin, C.: Identification and characterization

[17992] of a novel human matrix metalloproteinase with unique structural characteristics,

[17993] chromosomal location, and tissue distribution. J. Biol. Chem. 272:

[17994] 4281-4286, 1997.

[17995] 1803.

[17996] Liu, X.; Zou, H.; Slaughter, C.; Wang, X.: DFF, a het-erodimeric

[17997] protein that functions downstream of caspase-3 to trigger DNA fragmentation

[17998] during apoptosis. Cell 89: 175-184, 1997.

[17999] 1804.

[18000] Judson, H.; van Roy, N.; Strain, L.; Vandesompele, J.; Van Gele,

[18001] M.; Speleman, F.; Bonthron, D. T.: Structure and mutation analysis

[18002] of the gene encoding DNA fragmentation factor 40 (caspase-activated

[18003] nuclease), a candidate neuroblastoma tumour suppressor gene. Hum.

[18004] Genet. 106: 406-413, 2000.

[18005] 1805.

[18006] Liu, X.; Li, P.; Widlak, P.; Zou, H.; Luo, X.; Garrard, W. T.;

[18007] Wang, X.: The 40-kDa subunit of DNA fragmentation fac- tor induces

[18008] DNA fragmentation and chromatin condensation during apoptosis. Proc.

[18009] Nat. Acad. Sci. 95: 8461-8466, 1998.

[18010] 1806.

[18011] Mukae, N.; Enari, M.; Sakahira, H.; Fukuda, Y.; Inazawa, J.; Toh,

[18012] H.; Nagata, S.: Molecular cloning and characterization of human caspase-activated

[18013] DNase. Proc. Nat. Acad. Sci. 95: 9123-9128, 1998.

[18014] 1807.

[18015] Little, R. D.; Carulli, J. P.; Del Mastro, R. G.; Dupuis, J.; Osborne,

[18016] M.; Folz, C.; Manning, S. P.; Swain, P. M.; Zhao, S.-C.; Eustace,

[18017] B.; Lappe, M. M.; Spitzer, L.; and 23 others: A mutation in the

[18018] LDL receptor-related protein 5 gene results in the autosomal dominant

[18019] high-bone-mass trait. Am. J. Hum. Genet. 70: 11-19, 2002.

[18020] 1808.

[18021] Sherman, P. M.; Sun, H.; Macke, J. P.; Williams, J.; Smallwood,

[18022] P. M.; Nathans, J.: Identification and characterization of a conserved

[18023] family of protein serine/threonine phosphatases homolo-

gous to Drosophila

[18024] retinal degeneration C (rdgC). Proc. Nat. Acad. Sci. 94: 11639-11644,

[18025] 1997.

[18026] 1809.

[18027] Ohira, M.; Ootsuyama A.; Suzuki, E.; Ichikawa, H.; Seki, N.; Nagase,

[18028] T.; Monura, N.; Ohki, M.: Identification of a novel human gene containing

[18029] the tetratricopeptide repeat domain from the Down syndrome region

[18030] of chromosome 21. DNA Res. 3: 9-16, 1996.

[18031] 1810.

[18032] Tsukahara, F.; Hattori, M.; Muraki, T.; Sakaki, Y.: Identification

[18033] and cloning of a novel cDNA belonging to tetratricopeptide repeat

[18034] gene family from Down syndrome-critical region 21q22.2. J. Biochem. 120:

[18035] 820-827, 1996.

[18036] 1811.

[18037] Albrecht, U.; Sun, Z. S.; Eichele, G.; Lee, C. C.: A differen-
tial
[18038] response to two putative mammalian circadian regulators,
mper1 and
[18039] mper2, to light. Cell 91: 1055-1064, 1997.

[18040] 1812.

[18041] Bai, J.; Uehara, Y.; Montell, D. J.: Regulation of invasive cell

[18042] behavior by taiman, a Drosophila protein related to AIB1,
a steroid
[18043] receptor coactivator amplified in breast cancer. Cell 103:
1047-1058,
[18044] 2000.

[18045] 1813.

[18046] Chen, H.; Lin, R. J.; Schiltz, R. L.; Chakravarti, D.; Nash, A.;

[18047] Nagy, L.; Privalsky, M. L.; Nakatani, Y.; Evans, R. M.: Nu-
clear receptor
[18048] coactivator ACTR is a novel histone acetyltransferase and
forms a
[18049] multimeric activation complex with P/CAF and CBP/p300.
Cell 90:
[18050] 569-580, 1997.

[18051] 1814.

[18052] Guan, X.-Y.; Xu, J.; Anzick, S. L.; Zhang, H.; Trent, J. M.; Meltzer,

[18053] P. S.: Hybrid selection of transcribed sequences from microdissected

[18054] DNA: isolation of genes within amplified region at 20q11-q13.2 in

[18055] breast cancer. Cancer Res. 56: 3446-3450, 1996.

[18056] 1815.

[18057] Shirazi, S. K.; Bober, M. A.; Coetzee, G. A.: Polymorphic exonic

[18058] CAG microsatellites in the gene amplified in breast cancer (AIB1 gene). Clin.

[18059] Genet. 54: 102-103, 1998.

[18060] 1816.

[18061] Takeshita, A.; Cardona, G. R.; Koibuchi, N.; Suen, C.-S.; Chin,

[18062] W. W.: TRAM-1, a novel 160-kDa thyroid hormone receptor activator

[18063] molecule, exhibits distinct properties from steroid receptor coactivator-1. J.

[18064] Biol. Chem. 272: 27629-27634, 1997.

[18065] 1817.

[18066] Jay, P.; Sahly, I.; Goze, C.; Taviaux, S.; Poulat, F.; Couly, G.;

[18067] Abitbol, M.; Berta, P.: SOX22 is a new member of the SOX gene family,

[18068] mainly expressed in human nervous tissue. Hum. Molec. Genet. 6:

[18069] 1069-1077, 1997.

[18070] 1818.

[18071] Arrigo, G.; Gherzi, R.; Bonaglia, M. C.; Leprini, A.; Zuffardi,

[18072] O.; Zardi, L.: Assignment of the tenascin-R gene (Tnr) to mouse chromosome

[18073] 4 band E2 by fluorescence in situ hybridization; refine-ment of the

[18074] human TNR location to chromosome 1q24. Cytogenet. Cell Genet. 78:

[18075] 145-146, 1997.

[18076] 1819.

[18077] Carnemolla, B.; Leprini, A.; Borsi, L.; Querze, G.; Urbini, S.;

[18078] Zardi, L.: Human tenascin-R: complete primary structure,

pre-mRNA
[18079] alternative splicing and gene localization on chromosome 1q23-q24. J.
[18080] Biol. Chem. 271: 8157-8160, 1996.

[18081] 1820.

[18082] Erickson, H. P.: Tenascin-C, tenascin-R and tenascin-X: a family
[18083] of talented proteins in search of functions. Curr. Opin. Cell Biol. 5:
[18084] 869-876, 1993.

[18085] 1821.

[18086] Leprini, A.; Gherzi, R.; Siri, A.; Querze, G.; Viti, F.; Zardi,

[18087] L.: The human tenascin-R gene. J. Biol. Chem. 271: 31251-31254,
[18088] 1996.

[18089] 1822.

[18090] Williams, H.; Schachner, M.; Wang, B.; Kenwrick, S.: Radia-tion
[18091] hybrid mapping of the genes for tenascin-R (TNR), phos-ducin (PDC),
[18092] Laminin C1 (LAMC1), and TAX in 1q25-q32. Genomics 46:

165-166, 1997.

[18093] 1823.

[18094] Giguere, V.; Yang, N.; Segui, P.; Evans, R. M.: Identification

[18095] of a new class of steroid hormone receptors. Nature 331: 91-94,

[18096] 1988.

[18097] 1824.

[18098] Sladek, R.; Bader, J.-A.; Giguere, V.: The orphan nuclear receptor

[18099] estrogen-related receptor alpha is a transcriptional regulator of

[18100] the human medium-chain acyl coenzyme A dehydrogenase gene. Molec.

[18101] Cell. Biol. 17: 5400-5409, 1997.

[18102] 1825.

[18103] Sladek, R.; Beatty, B.; Squire, J.; Copeland, N. G.; Gilbert, D.

[18104] J.; Jenkins, N. A.; Giguere, V.: Chromosomal mapping of the human

[18105] and murine orphan receptors ERR-alpha (ESRRA) and ERR-beta (ESRRB)

[18106] and identification of a novel human ERR-alpha-related pseudogene. Genomics 45:
[18107] 320-326, 1997.

[18108] 1826.

[18109] Bengtson, P.; Larson, C.; Lundblad, A.; Larson, G.; Pahlsson, P.
[18110] : Identification of a missense mutation (G329A; arg110-to-gln) in
[18111] the human FUT7 gene. J. Biol. Chem. 276: 31575-31582, 2001.
[18112] 1827.

[18113] Mahdavi, J.; Sonden, B.; Hurtig, M.; Olfat, F. O.; Forsberg, L.;
[18114] Roche, N.; Angstrom, J.; Larsson, T.; Teneberg, S.; Karlsson, K.-A.;
[18115] Altraja, S.; Wadstrom, T.; and 11 others: Helicobacter pylori SabA
[18116] adhesin in persistent infection and chronic inflammation. Science 297:
[18117] 573-578, 2002.

[18118] 1828.

[18119] Natsuka, S.; Gersten, K. M.; Zenita, K.; Kannagi, R.; Lowe, J.

[18120] B.: Molecular cloning of a cDNA encoding a novel human leukocyte

[18121] alpha-1,3-fucosyltransferase capable of synthesizing the sialyl Lewis

[18122] x determinant. J. Biol. Chem. 269: 16789-16794, 1994.

[18123] 1829.

[18124] Sasaki, K.; Kurata, K.; Funayama, K.; Nagata, M.; Watan-

abe, E.;

[18125] Ohta, S.; Hannai, N.; Nishi, T.: Expression cloning of a novel alpha-1,3-fucosyltransferase

[18126] that is involved in biosynthesis of the sialyl Lewis x carbo-

hydrate

[18127] determinants in leukocytes. J. Biol. Chem. 269: 14730-14737, 1994.

[18128] 1830.

[18129] Eudy, J. D.; Ma-Edmonds, M.; Yao, S.-F.; Talmadge, C. B.; Kelley,

[18130] P. M.; Weston, M. D.; Kimberling, W. J.; Sumegi, J.: Isola-

tion of

[18131] a novel human homologue of the gene coding for echino-

derm microtubule-associated
[18132] protein (EMAP) from the Usher syndrome type 1a locus at 14q32. Genomics 43:
[18133] 104-106, 1997.

[18134] 1831.

[18135] Bale, T. L.; Contarino, A.; Smith, G. W.; Chan, R.; Gold, L. H.;
[18136] Sawchenko, P. E.; Koob, G. F.; Vale, W. W.; Lee, K.-F.: Mice deficient
[18137] for corticotropin-releasing hormone receptor-2 display anxiety-like
[18138] behaviour and are hypersensitive to stress. Nature Genet. 24: 410-414,
[18139] 2000.

[18140] 1832.

[18141] Coste, S. C.; Kesterson, R. A.; Heldwein, K. A.; Stevens, S. L.;
[18142] Heard, A. D.; Hollis, J. H.; Murray, S. E.; Hill, J. K.; Pantely,
[18143] G. A.; Hohimer, A. R.; Hatton, D. C.; Phillips, T. J.; Finn, D. A.;
[18144] Low, M. J.; Rittenberg, M. B.; Stenzel, P.; Stenzel-Poore, M.

P.:

[18145] Abnormal adaptations to stress and impaired cardiovascular function

[18146] in mice lacking corticotropin-releasing hormone receptor-2. Nature

[18147] Genet. 24: 403-409, 2000.

[18148] 1833.

[18149] Hsu, S. Y.; Hsueh, A. J. W.: Human stresscopin and stresscopin-related

[18150] peptide are selective ligands for the type 2 corticotropin-releasing

[18151] hormone receptor. Nature Med. 7: 605-611, 2001.

[18152] 1834.

[18153] Kishimoto, T.; Radulovic, J.; Radulovic, M.; Lin, C. R.; Schrick,

[18154] C.; Hooshmand, F.; Hermanson, O.; Rosenfeld, M. G.; Spiess, J.: Deletion

[18155] of Crhr2 reveals an anxiolytic role for corticotropin-releasing hormone

[18156] receptor-2. Nature Genet. 24: 415-419, 2000.

[18157] 1835.

[18158] Kostich, W. A.; Chen, A.; Sperle, K.; Largent, B. L.: Molecu-
lar
[18159] identification and analysis of a novel human corti-
cotropin-releasing
[18160] factor (CRF) receptor: the CRF2-gamma receptor. Molec.
Endocr. 12:
[18161] 1077-1085, 1998.

[18162] 1836.

[18163] Lesh, J. S.; Burrows, H. L.; Seasholtz, A. F.; Camper, S. A.:

[18164] Mapping of the mouse corticotropin-releasing hormone
receptor 2 gene
[18165] (Crhr2) to chromosome 6. Mammalian Genome 8:
944-945, 1997.
[18166] 1837.

[18167] Liaw, C. W.; Lovenberg, T. W.; Barry, G.; Oltersdorf, T.;
Grigoriadis,
[18168] D. E.; De Souza, E. B.: Cloning and characterization of the
human
[18169] corticotropin-releasing factor-2 receptor complementary
deoxyribonucleic
[18170] acid. Endocrinology 137: 72-77, 1996.

[18171] 1838.

[18172] Meyer, A. H.; Ullmer, C.; Schmuck, K.; Morel, C.; Wishart, W.;

[18173] Lubbert, H.; Engels, P.: Localization of the human CRF2 receptor

[18174] to 7p21-p15 by radiation hybrid mapping and FISH analysis. Genomics 40:

[18175] 189-190, 1997.

[18176] 1839.

[18177] Dallery, E.; Galiegue-Zouitina, S.; Collyn-d'Hooghe, M.; Quief,

[18178] S.; Denis, C.; Hildebrand, M.-P.; Lantoine, D.; Deweindt, C.; Tilly,

[18179] H.; Bastard, C.; Kerckaert, J.-P.: TTF, a gene encoding a novel small

[18180] G protein, fuses to the lymphoma-associated LAZ3 gene by t(3;4) chromosomal

[18181] translocation. Oncogene 10: 2171-2178, 1995.

[18182] 1840.

[18183] Dallery-Prudhomme, E.; Roumier, C.; Denis, C.; Preud-homme, C.;

[18184] Kerckaert, J.-P.; Galiegue-Zouitina, S.: Genomic structure and assignment
[18185] of the RhoH/TTF small GTPase gene (ARHH) to 4p13 by in situ hybridization. Genomics 43:
[18186] 89-94, 1997.

[18187] 1841.

[18188] Aaltonen, J.; Bjorses, P.; Sandkuijl, L.; Perheentupa, J.; Pel-tonen,
[18189] L.: An autosomal locus causing autoimmune disease: au-toimmune polyglandular
[18190] disease type I assigned to chromosome 21. Nature Genet. 8: 83-87,
[18191] 1994.

[18192] 1842.

[18193] Aaltonen, J.; Horelli-Kuitunen, N.; Fan, J.-B.; Bjorses, P.; Perheentupa,
[18194] J.; Myers, R.; Palotie, A.; Peltonen, L.: High-resolution physical
[18195] and transcriptional mapping of the autoimmune polyen-docrinopathy-candidiasis-ectodermal
[18196] dystrophy locus on chromosome 21q22.3 by FISH. Genome Res. 7: 820-829,

[18197] 1997.

[18198] 1843.

[18199] Ahonen, P.: Autoimmune polyendocrinopathy-
--candidosis--ectodermal
[18200] dystrophy (APECED): autosomal recessive inheritance.
Clin. Genet. 27:
[18201] 535-542, 1985.

[18202] 1844.

[18203] Ahonen, P.; Myllarniemi, S.; Sipila, I.; Perheentupa, J.: Clin-
ical
[18204] variation of autoimmune polyendocrinopathy-candidia-
sis-ectodermal
[18205] dystrophy (APECED) in a series of 68 patients. New Eng. J.
Med. 322:
[18206] 1829-1836, 1990.

[18207] 1845.

[18208] Arulanantham, K.; Kwyer, J. M.; Genel, M.: Evidence for
defective
[18209] immunoregulation in the syndrome of familial candidiasis
endocrinopathy. New
[18210] Eng. J. Med. 300: 164-168, 1979.

[18211] 1846.

[18212] Betterle, C.; Greggio, N. A.; Volpato, M.: Autoimmune polyglandular

[18213] syndrome type 1. J. Clin. Endocr. Metab. 83: 1049-1055, 1998.

[18214] 1847.

[18215] Bjorses, P.; Aaltonen, J.; Vikman, A.; Perheentupa, J.; Ben-Zion,

[18216] G.; Chiumello, G.; Dahl, N.; Heideman, P.; Hoorweg-Nijman, J. J. G.;

[18217] Mathivon, L.; Mullis, P. E.; Pohl, M.; Ritzen, M.; Romeo, G.; Shapiro,

[18218] M. S.; Smith, C. S.; Solyom, J.; Zlotogora, J.; Peltonen, L.: Genetic

[18219] homogeneity of autoimmune polyglandular disease type I. Am. J. Hum.

[18220] Genet. 59: 879-886, 1996.

[18221] 1848.

[18222] Bjorses, P.; Halonen, M.; Palvimo, J. J.; Kolmer, M.; Aaltonen,

[18223] J.; Ellonen, P.; Perheentupa, J.; Ulmanen, I.; Peltonen, L.: Mutations

[18224] in the AIRE gene: effects on subcellular location and transactivation

[18225] function of the autoimmune polyendocrinopathy-candidiasis-ectodermal

[18226] dystrophy protein. Am. J. Hum. Genet. 66: 378-392, 2000.

[18227] 1849.

[18228] Bjorses, P.; Pelto-Huikko, M.; Kaukonen, J.; Aaltonen, J.; Peltonen,

[18229] L.; Ulmanen, I.: Localization of the APECED protein in distinct nuclear

[18230] structures. Hum. Molec. Genet. 8: 259-266, 1999.

[18231] 1850.

[18232] Blizzard, R. M.; Kyle, M. A.: Studies of the adrenal antigens

[18233] and antibodies in Addison's disease. J. Clin. Invest. 42: 1653-1660,

[18234] 1963.

[18235] 1851.

[18236] Castells, S.; Fikrig, S.; Inamdar, S.; Orti, E.: Familial moniliasis,

[18237] defective delayed hypersensitivity, and adrenocorti-

cotropic hormone
[18238] deficiency. J. Pediat. 79: 72-79, 1971.

[18239] 1852.

[18240] Kim, S. H.; Moores, J. C.; David, D.; Respess, J. G.; Jolly, D.

[18241] J.; Friedmann, T.: The organization of the human HPRT gene. Nucleic
[18242] Acids Res. 14: 3103-3118, 1986.

[18243] 1853.

[18244] Alkhayat, A. H.; Kraemer, S. A.; Leipprandt, J. R.; Macek, M.;
[18245] Kleijer, W. J.; Friderici, K. H.: Human beta-mannosidase cDNA characterization
[18246] and first identification of a mutation associated with human beta-mannosidosis. Hum.
[18247] Molec. Genet. 7: 75-83, 1998.

[18248] 1854.

[18249] Cooper, A.; Hatton, C.; Thornley, M.; Sardharwalla, I. B.: Human
[18250] beta-mannosidase deficiency: biochemical findings in plasma, fibroblasts,
[18251] white cells and urine. J. Inherit. Metab. Dis. 11: 17-29,

1988.

[18252] 1855.

[18253] Cooper, A.; Sardharwalla, I. B.; Roberts, M. M.: Human beta-mannosidase

[18254] deficiency. (Letter) New Eng. J. Med. 315: 1231, 1986.

[18255] 1856.

[18256] Dorland, L.; Duran, M.; Hoefnagels, F. E. T.; Breg, J. N.; Fabery

[18257] de Jonge, H.; Cransberg, K.; van Sprang, F. J.; van Digge-

[18258] len, O. P.

[18259] : Beta-mannosidosis in two brothers with hearing loss. J. Inherit. Metab. Dis. 11 (suppl. 2): 255-258, 1988.

[18260] 1857.

[18261] Hu, P.; Wenger, D. A.; van Diggelen, O. P.; Klejer, W. J.: Complementation

[18262] studies in human and caprine beta-mannosidosis. J. In-

[18263] herit. Metab. Dis. 14: 13, 1990.

[18264] 1858.

[18265] Jones, M. Z.; Abbitt, B.: Animal model of human disease:

bovine

[18266] beta-mannosidosis. Am. J. Path. 142: 957-960, 1993.

[18267] 1859.

[18268] Jones, M. Z.; Cunningham, J. G.; Dade, A. W.; Alessi, D. M.; Mostosky,

[18269] U. V.; Vorro, J. R.; Benitez, J. T.; Lovell, K. L.: Caprine beta-mannosidosis:

[18270] clinical and pathological features. J. Neuropath. Exp. Neurol. 42:

[18271] 268-285, 1983.

[18272] 1860.

[18273] Jones, M. Z.; Dawson, G.: Caprine beta-mannosidosis: inherited

[18274] deficiency of beta-D-mannosidase. J. Biol. Chem. 256: 5185-5188,

[18275] 1981.

[18276] 1861.

[18277] Kleijer, W. J.; Hu, P.; Thoomes, R.; Boer, M.; Huijmans, J. G.

[18278] M.; Blom, W.; Van Diggelen, O. P.; Seemanova, E.; Macek, M.: Beta-mannosidase

[18279] deficiency: heterogeneous manifestation in the first fe-

male patient

[18280] and her brother. J. Inherit. Metab. Dis. 13: 867-872, 1990.

[18281] 1862.

[18282] Leipprandt, J. R.; Kraemer, S. A.; Haithcock, B. E.; Chen, H.;

[18283] Dyme, J. L.; Cavanagh, K. T.; Friderici, K. H.; Jones, M. Z.: Caprine

[18284] beta-mannosidase: sequencing and characterization of the cDNA and

[18285] identification of the molecular defect of caprine beta-mannosidosis. Genomics 37:

[18286] 51-56, 1996.

[18287] 1863.

[18288] Levade, T.; Graber, D.; Flurin, V.; Delisle, M.-B.; Pieraggi,

[18289] M.-T.; Testut, M.-F.; Carriere, J.-P.; Salvayre, R.: Human beta-mannosidase

[18290] deficiency associated with peripheral neuropathy. Ann. Neurol. 35:

[18291] 116-119, 1994.

[18292] 1864.

[18293] Lovell, K. L.; Jones, M. Z.: Distribution of central nervous

[18294] system lesions in beta-mannosidosis. Acta Neuropath. 62: 121-126,
[18295] 1983.

[18296] 1865.

[18297] Lundin, L.-G.: A gene (Bmn) controlling beta-mannosidase activity
[18298] in the mouse is located in the distal part of chromosome 3. Biochem.
[18299] Genet. 25: 603-610, 1987.

[18300] 1866.

[18301] Malachowski, J. A.; Jones, M. Z.: Beta-mannosidosis: lesions
[18302] of the distal peripheral nervous system. Acta Neuropath. 61: 95-100,
[18303] 1983.

[18304] 1867.

[18305] Poenaru, L.; Akli, S.; Rocchiccioli, F.; Eydoux, P.; Zamet, P.

[18306] : Human beta-mannosidosis: a 3-year-old boy with speech impairment
[18307] and emotional instability. Clin. Genet. 41: 331-334, 1992.

[18308] 1868.

[18309] Rodriguez-Serna, M.; Botella-Estrada, R.; Chabas, A.; Coll, M.-J.;

[18310] Oliver, V.; Febrer, M.-I.; Aliaga, A.: Angiokeratoma cor-poris diffusum

[18311] associated with beta-mannosidase deficiency. Arch. Derm. 132: 1219-1222,

[18312] 1996.

[18313] 1869.

[18314] Wenger, D. A.; Sujansky, E.; Fennessey, P. V.; Thompson, J. N.

[18315] : Human beta-mannosidase deficiency. New Eng. J. Med. 315: 1201-1205,

[18316] 1986.

[18317] 1870.

[18318] Bloom, G. E.; Zarkowsky, H. S.: Heterogeneity of the en-zyme defect

[18319] in congenital methemoglobinemia. New Eng. J. Med. 281: 919-922,

[18320] 1970.

[18321] 1871.

[18322] Aalfs, C. M.; Salieb-Beugelaar, G. B.; Wanders, R. J. A.; Mannens,
[18323] M. M. A. M.; Wijburg, F. A.: A case of methemoglobinemia type II
[18324] due to NADH-cytochrome b5 reductase deficiency: determination of the
[18325] molecular basis. Hum. Mutat. 16: 18-22, 2000.

[18326] 1872.

[18327] Balsamo, P.; Hardy, W. R.; Scott, E. M.: Hereditary methemoglobinemia
[18328] due to diaphorase deficiency in Navajo Indians. J. Pediat. 65: 928-930,
[18329] 1964.

[18330] 1873.

[18331] Board, P. G.; Pidcock, M. E.: Methaemoglobinaemia resulting from
[18332] heterozygosity for two NADH-methaemoglobin reductase variants: characterization
[18333] as NADH-ferricyanide reductase. Brit. J. Haemat. 47: 361-370, 1981.
[18334] 1874.

[18335] Bull, P. C.; Shephard, E. A.; Povey, S.; Santisteban, I.; Phillips,
[18336] I. R.: Cloning and chromosomal mapping of human cytochrome b(5) reductase
[18337] (DIA1). Ann. Hum. Genet. 52: 263-268, 1988.

[18338] 1875.

[18339] Cawein, M. J.; Behlen, C. H.; Lappat, E. J.; Cohn, J. E.: Hereditary
[18340] diaphorase deficiency and methemoglobinemia. Arch. Intern. Med. 113:
[18341] 578-585, 1964.

[18342] 1876.

[18343] Choury, D.; Leroux, A.; Kaplan, J.-C.: Membrane-bound cytochrome
[18344] b5 reductase (methemoglobin reductase) in human erythrocytes: study
[18345] in normal and methemoglobinemic subjects. J. Clin. Invest. 67: 149-155,
[18346] 1981.

[18347] 1877.

[18348] Cohen, R. J.; Sachs, J. R.; Wicker, D. J.; Conrad, M. E.:

Methemoglobinemia
[18349] provoked by malarial chemoprophylaxis in Vietnam. New Eng. J. Med. 279:
[18350] 1127-1131, 1968.

[18351] 1878.

[18352] Comly, H. H.: Cyanosis in infants caused by nitrates in well water. J.A.M.A. 129:
[18353] 112-116, 1945.

[18354] 1879.

[18355] Dekker, J.; Eppink, M. H. M.; van Zwieten, R.; de Rijk, R.; Remacha,
[18356] A. F.; Law, L. K.; Li, A. M.; Cheung, K. L.; van Berkel, W. J. H.;
[18357] Roos, D.: Seven new mutations in the nicotinamide adenine dinucleotide

[18358] reduced-cytochrome b5 reductase gene leading to methemoglobinemia

[18359] type I. Blood 97: 1106-1114, 2001.

[18360] 1880.

[18361] Du, M.; Shirabe, K.; Takeshita, M.: Identification of alternative

[18362] first exons of NADH-cytochrome b5 reductase gene expressed ubiquitously

[18363] in human cells. Biochem. Biophys. Res. Commun. 235: 779-783, 1997.

[18364] 1881.

[18365] Fialkow, P. J.; Browder, J. A.; Sparkes, R. S.; Motulsky, A. G.

[18366] : Mental retardation in methemoglobinemia due to diaphorase deficiency. New

[18367] Eng. J. Med. 273: 840-845, 1965.

[18368] 1882.

[18369] Fisher, R. A.; Povey, S.; Bobrow, M.; Solomon, E.; Boyd, Y.; Carritt,

[18370] B.: Assignment of the DIA-1 locus to chromosome 22. Ann. Hum. Genet. 41:

[18371] 151-155, 1977.

[18372] 1883.

[18373] Gibson, Q. H.: Methemoglobinemia--long ago and far away. Am.

[18374] J. Hemat. 42: 3-6, 1993.

[18375] 1884.

[18376] Gonzalez, R.; Estrada, M.; Wade, M.; de la Torre, E.; Svarch,

[18377] E.; Fernandez, O.; Oritz, R.; Guzman, E.; Colombo, B.: Het-

[18378] erogeneity of hereditary methaemoglobinaemia: a study of 4 Cuban families with

[18379] NADH-methaemoglobin reductase deficiency including a new variant (Santiage

[18380] de Cuba variant). Scand. J. Haemat. 20: 385-393, 1978.

[18381] 1885.

[18382] Hirano, M.; Matsuki, T.; Tanishima, K.; Takeshita, M.; Shimizu,

[18383] S.; Nagamura, Y.; Yoneyama, Y.: Congenital methaemoglobinaemia due

[18384] to NADH methaemoglobin reductase deficiency: success-

[18385] ful treatment

[18385] with oral riboflavin. Brit. J. Haemat. 47: 353-359, 1981.

[18386] 1886.

[18387] Moerman, P.; Fryns, J.-P.; Devlieger, H.; Van Assche, A.; Lauweryns,

[18388] J.: Congenital eventration of the diaphragm: an unusual cause of

[18389] intractable neonatal respiratory distress with variable etiology. Am.

[18390] J. Med. Genet. 27: 213-218, 1987.

[18391] 1887.

[18392] Tsai, H.-M.: Physiologic cleavage of von Willebrand factor by

[18393] a plasma protease is dependent on its conformation and requires calcium

[18394] ion. Blood 87: 4235-4244, 1996.

[18395] 1888.

[18396] Tsai, H.-M.; Lian, E. C.-Y.: Antibodies to von Willebrand factor-cleaving

[18397] protease in acute thrombotic thrombocytopenic purpura. New Eng. J.

[18398] Med. 339: 1585-1594, 1998.

[18399] 1889.

[18400] Baek, S. J.; Lee, K.-D.; Shen, R.-F.: Genomic structure and polymorphism

[18401] of the human thromboxane synthase-encoding gene. Gene 173: 251-256,

[18402] 1996.

[18403] 1890.

[18404] Chase, M. B.; Baek, S. J.; Purtell, D. C.; Schwartz, S.; Shen,

[18405] R.-F.: Mapping of the human thromboxane synthase gene (TBXAS1) to

[18406] chromosome 7q34-q35 by two-color fluorescence in situ hybridization. Genomics 16:

[18407] 771-773, 1993.

[18408] 1891.

[18409] Metsel, F.; Oetliker, O.; Beck, E.; Felix, R.; Imbach, P.; Wagner,

[18410] H.-P.: Severe bleeding associated with defective throm-boxane synthetase.

[18411] (Letter) Lancet I: 157, 1980.

[18412] 1892.

[18413] Miyata, A.; Yokoyama, C.; Ihara, H.; Bandoh, S.; Takeda, O.; Takahashi,
[18414] E.; Tanabe, T.: Characterization of the human gene (TBXAS1) encoding
[18415] thromboxane synthase. Europ. J. Biochem. 224: 273-279, 1994.
[18416] 1893.

[18417] Nebert, D. W.: Personal Communication. Cincinnati, Ohio 5/9/1994.
[18418] 1894.

[18419] Pareti, F. I.; Mannucci, P. M.; D'Angelo, A.; Smith, J. B.; Sautebin,
[18420] L.; Galli, G.: Congenital deficiency of thromboxane and prostacyclin. Lancet I:
[18421] 898-900, 1980.

[18422] 1895.

[18423] Weiss, H. J.; Lages, B. A.: Possible congenital defect in platelet
[18424] thromboxane synthetase. (Letter) Lancet II: 760-761, 1977.
[18425] 1896.

[18426] Yokoyama, C.; Miyata, A.; Ihara, H.; Ullrich, V.; Tanabe, T.:

[18427] Molecular cloning of human platelet thromboxane A syn-
thase. Biochem.

[18428] Biophys. Res. Commun. 178: 1479-1484, 1991.

[18429] 1897.

[18430] Zhang, L.; Xiao, H.; Schultz, R. A.; Shen, R.-F.: Genomic
organization,

[18431] chromosomal localization, and expression of the murine
thromboxane

[18432] synthase gene (Tbxas1). Genomics 45: 519-528, 1997.

[18433] 1898.

[18434] Nagase, T.; Ishikawa, K.; Nakajima, D.; Ohira, M.; Seki, N.;
Miyajima,

[18435] N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Predic-
tion of

[18436] the coding sequences of unidentified human genes. VII.
The complete

[18437] sequences of 100 new cDNA clones from brain which can
code for large

[18438] proteins in vitro. DNA Res. 4: 141-150, 1997.

[18439] 1899.

[18440] Di Como, C. J.; Arndt, K. T.: Nutrients, via the Tor proteins,
[18441] stimulate the association of Tap42 with type 2A phosphatases. Genes
[18442] Dev. 10: 1904-1916, 1996.

[18443] 1900.

[18444] Inui, S.; Kuwahara, K.; Mizutani, J.; Maeda, K.; Kawai, T.; Nakayasu,
[18445] H.; Sakaguchi, N.: Molecular cloning of a cDNA clone encoding a phosphoprotein
[18446] component related to the Ig receptor-mediated signal transduction. J.
[18447] Immun. 154: 2714-2723, 1995.

[18448] 1901.

[18449] Onda, M.; Inui, S.; Maeda, K.; Suzuki, M.; Takahashi, E.; Sakaguchi,
[18450] N.: Expression and chromosomal localization of the human alpha-4/IGBP1
[18451] gene, the structure of which is closely related to the yeast TAP42
[18452] protein of the rapamycin-sensitive signal transduction pathway. Genomics 46:

[18453] 373-378, 1997.

[18454] 1902.

[18455] Prueitt, R. L.; Ross, J. L.; Zinn, A. R.: Physical mapping of

[18456] nine Xq translocation breakpoints and identification of XPNPEP2 as

[18457] a premature ovarian failure candidate gene. Cytogenet. Cell Genet. 89:

[18458] 44-50, 2000.

[18459] 1903.

[18460] Sprinkle, T. J.; Stone, A. A.; Venema, R. C.; Denslow, N. D.; Caldwell,

[18461] C.; Ryan, J. W.: Assignment of the membrane-bound human aminopeptidase

[18462] P gene (XPNPEP2) to chromosome Xq25. Genomics 50: 114-116, 1998.

[18463] 1904.

[18464] Venema, R. C.; Ju, H.; Zou, R.; Venema, V. J.; Ryan, J. W.: Cloning

[18465] and tissue distribution of human membrane-bound aminopeptidase P. Biochim.

[18466] Biophys. Acta 1354: 45-48, 1997.

[18467] 1905.

[18468] Dear, N.; Matena, K.; Vingron, M.; Boehm, T.: A new sub-
family

[18469] of vertebrate calpains lacking a calmodulin-like domain:
implications

[18470] for calpain regulation and evolution. Genomics 45:
175-184, 1997.

[18471] 1906.

[18472] Whiting, P. J.; Bonnert, T. P.; McKernan, R. M.; Farrar, S.; le

[18473] Bourdelles, B.; Heavens, R. P.; Smith, D. W.; Hewson, L.;
Rigby, M.

[18474] R.; Sirinathsinghji, D. J. S.; Thompson, S. A.; Wafford, K.
A.: Molecular

[18475] and functional diversity of the expanding GABA-A recep-
tor gene family. Ann.

[18476] N.Y. Acad. Sci. 645-653, 1999.

[18477] 1907.

[18478] Langnaese, K.; Colleaux, L.; Kloos, D. U.; Fontes, M.;
Wieacker,

[18479] P.: Cloning of Z39Ig, a novel gene with immunoglobulin-
like domains

[18480] located on human chromosome X. Biochim. Biophys. Acta 1492: 522-525,

[18481] 2000.

[18482] 1908.

[18483] Abo, A.; Qu, J.; Cammarano, M. S.; Dan, C.; Fritsch, A.; Baud,

[18484] V.; Belisle, B.; Minden, A.: PAK4, a novel effector for Cdc42Hs,

[18485] is implicated in the reorganization of the actin cytoskeleton and

[18486] in the formation of filopodia. EMBO J. 17: 6527-6540, 1998.

[18487] 1909.

[18488] Zhang, F.; Zhang, W.; Liu, L.; Fisher, C. L.; Hui, D.; Childs,

[18489] S.; Dorovini-Zis, K.; Ling, V.: Characterization of ABCB9, an ATP

[18490] binding cassette protein associated with lysosomes. J. Biol. Chem. 275:

[18491] 23287-23294, 2000.

[18492] 1910.

[18493] Seki, N.; Yoshikawa, T.; Hattori, A.; Miyajima, N.; Mura-

matsu,
[18494] M.; Saito, T.: cDNA cloning of a human RAB26-related gene encoding
[18495] a Ras-like GTP-binding protein on chromosome 16p13.3 region. J. Hum.
[18496] Genet. 45: 309-314, 2000.

[18497] 1911.

[18498] Lee, J.; Ho, W.-H.; Maruoka, M.; Corpuz, R. T.; Baldwin, D. T.;
[18499] Foster, J. S.; Goddard, A. D.; Yansura, D. G.; Vandlen, R. L.; Wood,
[18500] W. L.; Gurney, A. L.: IL-17E, a novel proinflammatory lig- and for
[18501] the IL-17 receptor homolog IL-17Rh1. J. Biol. Chem. 276: 1660-1664,
[18502] 2001.

[18503] 1912.

[18504] Tian, E.; Sawyer, J. R.; Largaespada, D. A.; Jenkins, N. A.; Copeland,
[18505] N. G.; Shaughnessy, J. D., Jr.: Evi27 encodes a novel mem- brane protein
[18506] with homology to the IL17 receptor. Oncogene 19:

2098-2109, 2000.

[18507] 1913.

[18508] Buanne, P.; Corrente, G.; Micheli, L.; Palena, A.; Lavia, P.; Spadafora,

[18509] C.; Lakshmana, M. K.; Rinaldi, A.; Banfi, S.; Quarto, M.; Bulfone,

[18510] A.; Tirone, F.: Cloning of PC3B, a novel member of the PC3/BTG/TOB

[18511] family of growth inhibitory genes, highly expressed in the olfactory

[18512] epithelium. Genomics 68: 253-263, 2000.

[18513] 1914.

[18514] Yoshida, Y.; Matsuda, S.; Ikematsu, N.; Kawamura-Tsuzuku, J.; Inazawa,

[18515] J.; Umemori, H.; Yamamoto, T.: ANA, a novel member of Tob/BTG1 family,

[18516] is expressed in the ventricular zone of the developing central nervous

[18517] system. Oncogene 16: 2687-2693, 1998.

[18518] 1915.

[18519] Ueki, N.; Oda, T.; Kondo, M.; Yano, K.; Noguchi, T.; Mura-

matsu,

[18520] M.: Selection system for genes encoding nuclear-targeted proteins. Nat.

[18521] Biotech. 16: 1338-1342, 1998.

[18522] 1916.

[18523] Ueki, N.; Seki, N.; Yano, K.; Masuho, Y.; Saito, T.; Muramatsu,

[18524] M.: Isolation and characterization of a novel human gene (HFB30)

[18525] which encodes a protein with a RING finger motif. Biochim. Biophys.

[18526] Acta 232-236, 1999.

[18527] 1917.

[18528] Oikawa, E.; Iijima, H.; Suzuki, T.; Sasano, H.; Sato, H.; Kamataki,

[18529] A.; Nagura, H.; Kang, M. J.; Fujino, T.; Suzuki, H.; Yamamoto, T.

[18530] T.: A novel acyl-CoA synthetase, ACS5, expressed in intestinal epithelial

[18531] cells and proliferating preadipocytes. J. Biochem. 124: 679-685,

[18532] 1998.

[18533] 1918.

[18534] Yamashita, Y.; Kumabe, T.; Cho, Y.-Y.; Watanabe, M.;
Kawagishi,
[18535] J.; Yoshimoto, T.; Fujino, T.; Kang, M.-J.; Yamamoto, T. T.
Fatty
[18536] acid induced glioma cell growth is mediated by the acyl-
CoA synthetase
[18537] 5 gene located on chromosome 10q25.1-q25.2, a region
frequently deleted
[18538] in malignant gliomas. Oncogene 19: 5919-5925, 2000.

[18539] 1919.

[18540] Meng, X.; Lu, X.; Li, Z.; Green, E. D.; Massa, H.; Trask, B. J.;

[18541] Morris, C. A.; Keating, M. T.: Complete physical map of
the common
[18542] deletion region in Williams syndrome and identification
and characterization
[18543] of three novel genes. Hum. Genet. 103: 590-599, 1998.

[18544] 1920.

[18545] Jones, M. H.; Hamana, N.; Nezu, J.; Shimane, M.: A novel
family
[18546] of bromodomain genes. Genomics 63: 40-45, 2000.

[18547] 1921.

[18548] Lu, X.; Meng, X.; Morris, C. A.; Keating, M. T.: A novel hu-
man
[18549] gene, WSTF, is deleted in Williams syndrome. Genomics
54: 241-249,
[18550] 1998.

[18551] 1922.

[18552] Peoples, R. J.; Cisco, M. J.; Kaplan, P.; Francke, U.: Identifi-
cation
[18553] of the WBSCR9 gene, encoding a novel transcriptional reg-
ulator, in
[18554] the Williams-Beuren syndrome deletion at 7q11.23. Cyto-
genet. Cell
[18555] Genet. 82: 238-246, 1998.

[18556] 1923.

[18557] Yano, H.; Wang, C.-Z.; Yamashita, S.; Yokoyama, Y.; Yokoi,
N.;
[18558] Seino, S.: Assignment of the human solute carrier family
4, sodium
[18559] bicarbonate cotransporter-like, member 10 gene
(SLC4A10) to 2q23-q24

[18560] by in situ hybridization and radiation hybrid mapping. Cytogenet.

[18561] Cell Genet. 89: 276-277, 2000.

[18562] 1924.

[18563] Mitchelmore, C.; Troelsen, J. T.; Sjostrom, H.; Noren, O.: The

[18564] HOXC11 homeodomain protein interacts with the lactose-phlorizin hydrolase

[18565] promoter and stimulates HNF1-alpha-dependent transcription. J. Biol.

[18566] Chem. 273: 13297-13306, 1998.

[18567] 1925.

[18568] Watson, M. A.; Darrow, C.; Zimonjic, D. B.; Popescu, N.; Fleming,

[18569] T. P.: Structure and transcriptional regulation of the human mammaglobin

[18570] gene, a breast cancer associated member of the uteroglobin gene family

[18571] localized to chromosome 11q13. Oncogene 16: 817-824, 1998.

[18572] 1926.

[18573] Watson, M. A.; Fleming, T. P.: Mammaglobin, a mammary-specific
[18574] member of the uteroglobin gene family, is overexpressed in human breast
[18575] cancer. Cancer Res. 56: 860-865, 1996.

[18576] 1927.

[18577] Watson, M. A.; Fleming, T. P.: Isolation of differentially expressed
[18578] sequence tags from human breast cancer. Cancer Res. 54: 4598-4602,
[18579] 1994.

[18580] 1928.

[18581] Seki, N., Hattori, A.; Hayashi, A.; Kozuma, S.; Ohira, M.; Hori,
[18582] T.; Saito, T.: Structure, expression profile and chromosomal location
[18583] of an isolog of DNA-PKcs interacting protein (KIP) gene. Biochim.
[18584] Biophys. Acta 1444: 143-147, 1999.

[18585] 1929.

[18586] Scott, D. A.; Drury, S.; Sundstrom, R. A.; Bishop, J.;

Swiderski,
[18587] R. E.; Carmi, R.; Ramesh, A.; Elbedour, K.; Srisailapathy, C. R. S.;
[18588] Keats, B. J.; Sheffield, V. C.; Smith, R. J. H.: Refining the DFNB7-DFNB11
[18589] deafness locus using intragenic polymorphisms in a novel gene, TMEM2. Gene 246:
[18590] 265-274, 2000.

[18591] 1930.

[18592] Lancaster, J. M.; Wooster, R.; Mangion, J.; Phelan, C. M.; Cochran,
[18593] C.; Gumbs, C.; Seal, S.; Barfoot, R.; Collins, N.; Bignell, G.; Patel,
[18594] S.; Hamoudi, R.; Larsson, C.; Wiseman, R. W.; Berchuck, A.; Iglehart,
[18595] J. D.; Marks, J. R.; Ashworth, A.; Stratton, M. R.; Futreal, P. A.
[18596] : BRCA2 mutations in primary breast and ovarian cancers. Nature Genet. 13:
[18597] 238-240, 1996.

[18598] 1931.

[18599] Lee, H.; Trainer, A. H.; Friedman, L. S.; Thistlethwaite, F.

C.;

[18600] Evans, M. J.; Ponder, B. A. J.; Venkitaraman, A. R.: Mitotic checkpoint

[18601] inactivation fosters transformation in cells lacking the breast cancer

[18602] susceptibility gene, Brca2. Molec. Cell 4: 1-10, 1999.

[18603] 1932.

[18604] Lerer, I.; Wang, T.; Peretz, T.; Sagi, M.; Kaduri, L.; Orr-Urtreger,

[18605] A.; Stadler, J.; Gutman, H.; Abeliovich, D.: The 8765delAG mutation

[18606] in BRCA2 is common among Jews of Yemenite extraction. (Letter) Am.

[18607] J. Hum. Genet. 63: 272-274, 1998.

[18608] 1933.

[18609] Miki, Y.; Katagiri, T.; Kasumi, F.; Yoshimoto, T.; Nakamura, Y.

[18610] : Mutation analysis in the BRCA2 gene in primary breast cancers. Nature

[18611] Genet. 13: 245-247, 1996.

[18612] 1934.

[18613] Milner, J.; Ponder, B.; Hughes-Davies, L.; Seltmann, M.; Kouzarides,
[18614] T.: Transcriptional activation functions in BRCA2. (Letter) Nature 386:
[18615] 772-773, 1997.

[18616] 1935.

[18617] Moynahan, M. E.; Pierce, A. J.; Jasin, M.: BRCA2 is required
[18618] for homology-directed repair of chromosomal breaks. Molec. Cell 7:
[18619] 263-272, 2001.

[18620] 1936.

[18621] Nastiuk, K. L.; Mansukhani, M.; Terry, M. B.; Kularatne, P.; Rubin,
[18622] M. A.; Melamed, J.; Gammon, M. D.; Ittmann, M.; Krolewski, J. J.:
[18623] Common mutations in BRCA1 and BRCA2 do not con-
tribute to early prostate
[18624] cancer in Jewish men. Prostate 40: 172-177, 1999.

[18625] 1937.

[18626] Neuhausen, S.; Gilewski, T.; Norton, L.; Tran, T.; McGuire, P.;

[18627] Swensen, J.; Hampel, H.; Borgen, P.; Brown, K.; Skolnick, M.; Shattuck-Eidens,

[18628] D.; Jhanwar, S.; Goldgar, D.; Offit, K.: Recurrent BRCA2 6174delT

[18629] mutations in Ashkenazi Jewish women affected by breast cancer. Nature

[18630] Genet. 13: 126-128, 1996.

[18631] 1938.

[18632] Neuhausen, S. L.; Godwin, A. K.; Gershoni-Baruch, R.; Schubert,

[18633] E.; Garbert, J.; Stoppa-Lyonnet, D.; Olah, E.; Csokay, B.; Serova,

[18634] O.; Lalloo, F.; Osorio, A.; Stratton, M.; and 18 others: Haplotype

[18635] and phenotype analysis of nine recurrent BRCA2 mutations in 111 families:

[18636] Results of an international study. Am. J. Hum. Genet. 62: 1381-1388,

[18637] 1998.

[18638] 1939.

[18639] Neuhausen, S. L.; Godwin, A. K.; Gershoni-Baruch, R.; Shubert,

[18640] E.; Garber, J.; Stoppa-Lyonnet, D.; Olah, E.; Csokay, B.; Serova,
[18641] O.; Lalloo, F.; Osorio, A.; Stratton, M.; and 18 others: Haplotype
[18642] and phenotype analysis of nine recurrent BRCA2 mutations in 111 families:
[18643] results of an international study. Am. J. Hum. Genet. 62: 1381-1388,
[18644] 1998.

[18645] 1940.

[18646] Oddoux, C.; Struewing, J. P.; Clayton, C. M.; Neuhausen, S.; Brody,
[18647] L. C.; Kaback, M.; Haas, B.; Norton, L.; Borgen, P.; Jhanwar, S.;
[18648] Goldgar, D.; Ostrer, H.; Offit, K.: The carrier frequency of the
[18649] BRCA2 6174delT mutation among Ashkenazi Jewish individuals is approximately
[18650] 1%. Nature Genet. 14: 188-190, 1996.

[18651] 1941.

[18652] Ozcelik, H.; Schmocker, B.; Di Nicola, N.; Shi, X.-H.; Langer,

[18653] B.; Moore, M.; Taylor, B. R.; Narod, S. A.; Darlington, G.; Andrulis,

[18654] I. L.; Gallinger, S.; Redston, M.: Germline BRCA2 6174delT mutations

[18655] in Ashkenazi Jewish pancreatic cancer patients. (Letter) Nature Genet. 16:

[18656] 17-18, 1997.

[18657] 1942.

[18658] Patel, K. J.; Yu, V. P. C. C.; Lee, H.; Corcoran, A.; Thistlethwaite,

[18659] F. C.; Evans, M. J.; Colledge, W. H.; Friedman, L. S.; Ponder, B.

[18660] A. J.; Venkitaraman, A. R.: Involvement of Brca2 in DNA repair. Molec.

[18661] Cell 1: 347-357, 1998.

[18662] 1943.

[18663] Hu, X.; Ray, P. N.; Murphy, E. G.; Thompson, M. W.; Worton, R.

[18664] G.: Duplicational mutation at the Duchenne muscular dystrophy locus:

[18665] its frequency, distribution, origin, and phenotype-genotype correlation. Am.

[18666] J. Hum. Genet. 46: 682-695, 1990.

[18667] 1944.

[18668] Hu, X.; Ray, P. N.; Worton, R. G.: Mechanisms of tandem duplication

[18669] in the Duchenne muscular dystrophy gene include both homologous and

[18670] nonhomologous intrachromosomal recombination. EMBO J. 10: 2471-2477,

[18671] 1991.

[18672] 1945.

[18673] Hu, X.; Worton, R. G.: Partial gene duplication as a cause of

[18674] human disease. Hum. Mutat. 1: 3-12, 1992.

[18675] 1946.

[18676] Ingram, V. M.: Gene evolution and the haemoglobins. Na-ture 189:

[18677] 704-708, 1961.

[18678] 1947.

[18679] Itagaki, Y.; Saida, K.; Iwamura, K.: Regenerative capacity of

[18680] mdx mouse muscles after repeated applications of myo-

necrotic bupivacaine. Acta
[18681] Neuropath. 89: 380-384, 1995.

[18682] 1948.

[18683] Kaplan, J.-C.; Kahn, A.; Chelly, J.: Illegitimate transcrip-
tion:
[18684] its use in the study of inherited disease. Hum. Mutat. 1:
357-360,
[18685] 1992.

[18686] 1949.

[18687] Kavaslar, G. N.; Telatar, M.; Serdaroglu, P.; Deymeer, F.;
Ozdemir,
[18688] C.; Tolun, A.: Identification of a one-basepair deletion in
exon
[18689] 6 of the dystrophin gene. Hum. Mutat. 6: 85-86, 1995.

[18690] 1950.

[18691] Kilimann, M. W.; Pizzuti, A.; Grompe, M.; Caskey, C. T.:
Point
[18692] mutations and polymorphisms in the human dystrophin
gene identified
[18693] in genomic DNA sequences amplified by multiplex PCR.
Hum. Genet. 89:

[18694] 253-258, 1992.

[18695] 1951.

[18696] Kim, T.-W.; Wu, K.; Black, I. B.: Deficiency of brain synaptic

[18697] dystrophin in human Duchenne muscular dystrophy. Ann. Neurol. 38:

[18698] 446-449, 1995.

[18699] 1952.

[18700] Kneppers, A. L. J.; Deutz-Terlouw, P. P.; van Ommen, G. J. B.;

[18701] Bakker, E.: Point mutation screening for Duchenne muscular dystrophy

[18702] (DMD) by SSCP-analysis of multiplex PCR products by use of the PhastSystem

[18703] (TM). Am. J. Hum. Genet. Suppl. 53: Abstract-1493, 1993.

[18704] 1953.

[18705] Ostendorff, H. P.; Bossenz, M.; Mincheva, A.; Copeland, N. G.;

[18706] Gilbert, D. J.; Jenkins, N. A.; Lichter, P.; Bach, I.: Functional

[18707] characterization of the gene encoding RLIM, the corepressor of LIM

[18708] homeodomain factors. Genomics 69: 120-130, 2000.

[18709] 1954.

[18710] Ostendorff, H. P.; Peirano, R. I.; Peters, M. A.; Schluter, A.;

[18711] Bossenz, M.; Scheffner, M.; Bach, I.: Ubiquitination-depen-
dent cofactor

[18712] exchange on LIM homeodomain transcription factors. Na-
ture 416: 99-103,

[18713] 2002.

[18714] 1955.

[18715] Koenig, M.: Personal Communication. Boston, Mass.
10/8/1987.

[18716]

[18717] 100. Koenig, M.; Beggs, A. H.; Moyer, M.; Scherpf, S.;
Heindrich,

[18718] K.; Bettecken, T.; Meng, G.; Muller, C. R.; Lindlof, M.;
Kaariainen,

[18719] H.; de la Chapelle, A.; Kiuru, A.; and 24 others: The
molecular

[18720] basis for Duchenne versus Becker muscular dystrophy:
correlation of

[18721] severity with type of deletion. Am. J. Hum. Genet. 45:

498-506,
[18722] 1989.

[18723]

[18724] 101. Koenig, M.; Bertelson, C. J.; Monaco, A. P.; Hoffman, E.; Feener,
[18725] C. C.; Kunkel, L. M.: Complete cloning of the Duchenne muscular dystrophy
[18726] cDNA and an analysis of the entire DMD locus. (Abstract) Am. J. Hum.
[18727] Genet. 41: A222, 1987.

[18728]

[18729] 102. Koenig, M.; Hoffman, E. P.; Bertelson, C. J.; Monaco, A. P.;
[18730] Feener, C.; Kunkel, L. M.: Complete cloning of the Duchenne muscular
[18731] dystrophy (DMD) cDNA and preliminary genomic organi-zation of the DMD
[18732] gene in normal and affected individuals. Cell 50: 509-517, 1987.

[18733]

[18734] 103. Koenig, M.; Monaco, A. P.; Kunkel, L. M.: The com-

plete sequence
[18735] of dystrophin predicts a rod-shaped cytoskeletal protein. Cell 53:
[18736] 219-228, 1988.

[18737]

[18738] 104. Koh, J.; Bartlett, R. J.; Pericak-Vance, M. A.; Speer, M. C.;
[18739] Yamaoka, L. H.; Phillips, K.; Hung, W.-Y.; Ray, P. N.; Wor-
ton, R.
[18740] G.; Gilbert, J. R.; Lee, J. E.; Siddique, T.; Kandt, R. S.; Roses,
[18741] A. D.: Inherited deletion at Duchenne dystrophy locus in normal male.
[18742] (Letter) Lancet II: 1154-1155, 1987.

[18743]

[18744] 105. Kunkel, L. M.: Analysis of deletions in DNA from pa-
tients with
[18745] Becker and Duchenne muscular dystrophy. Nature 322:
73-77, 1986.

[18746]

[18747] 106. Kunkel, L. M.; Monaco, A. P.; Middlesworth, W.; Ochs,

H. D.;

[18748] Latt, S. A.: Specific cloning of DNA fragments absent from the DNA

[18749] of a male patient with an X chromosome deletion. Proc. Nat. Acad.

[18750] Sci. 82: 4778-4782, 1985.

[18751]

[18752] 107. Laing, N. G.; Layton, M. G.; Johnsen, R. D.; Chandler, D. C.;

[18753] Mears, M. E.; Goldblatt, J.; Kakulas, B. A.: Two distinct mutations

[18754] in a single dystrophin gene: chance occurrence or premutation? Am.

[18755] J. Med. Genet. 42: 688-692, 1992.

[18756]

[18757] 108. Lederfein, D.; Levy, Z.; Augier, N.; Mornet, D.; Morris, G.;

[18758] Fuchs, O.; Yaffe, D.; Nudel, U.: A 71-kilodalton protein is a major

[18759] product of the Duchenne muscular dystrophy gene in brain and other

[18760] nonmuscle tissues. Proc. Nat. Acad. Sci. 89: 5346-5350,

1992.

[18761]

[18762] 109. Lederfein, D.; Yaffe, D.; Nudel, U.: A housekeeping type promoter,

[18763] located in the 3-prime region of the Duchenne muscular dystrophy gene,

[18764] controls the expression of Dp71, a major product of the gene. Hum.

[18765] Molec. Genet. 2: 1883-1888, 1993.

[18766]

[18767] 110. Lee, C. C.; Pearlman, J. A.; Chamberlain, J. S.; Caskey, C. T.

[18768] : Expression of recombinant dystrophin and its localization to the

[18769] cell membrane. Nature 349: 334-336, 1991.

[18770]

[18771] 111. Lee, G.-H.; Badorff, C.; Knowlton, K. U.: Dissociation of sarcoglycans

[18772] and the dystrophin carboxyl terminus from the sarcolemma in enteroviral

[18773] cardiomyopathy. Circ. Res. 87: 489-495, 2000.

[18774]

[18775] 112. Lenk, U.; Hanke, R.; Kraft, U.; Grade, K.; Grunewald, I.; Speer,

[18776] A.: Non-isotopic analysis of single strand conformation polymorphism

[18777] (SSCP) in the exon 13 region of the human dystrophin gene. J. Med.

[18778] Genet. 30: 951-954, 1993.

[18779]

[18780] 113. Lenk, U.; Hanke, R.; Speer, A.: Carrier detection in DMD families

[18781] with point mutations, using PCR-SSCP and direct sequencing. Neuromusc.

[18782] Disord. 4: 411-418, 1994.

[18783]

[18784] 114. Lenk, U.; Hanke, R.; Thiele, H.; Speer, A.: Point mutations

[18785] at the carboxy terminus of the human dystrophin gene: implications

[18786] for an association with mental retardation in DMD patients. Hum.

[18787] Molec. Genet. 2: 1877-1881, 1993.

[18788]

[18789] 115. Lenk, U.; Oexle, K.; Voit, T.; Ancker, U.; Hellner, K.-A.; Speer,

[18790] A.; Hubner, C.: A cysteine 3340 substitution in the dystro-glycan-binding

[18791] domain of dystrophin associated with Duchenne muscular dystrophy,

[18792] mental retardation and absence of the ERG b-wave. Hum. Molec. Genet. 973-975,

[18793] 1996.

[18794]

[18795] 116. Liechti-Gallati, S.; Braga, S.; Hirsiger, H.; Moser, H.: Familial

[18796] deletion in Becker type muscular dystrophy within the pXJ region. Hum.

[18797] Genet. 77: 267-268, 1987.

[18798]

[18799] 117. Lindlof, M.; Kaariainen, H.; van Ommen, G. J. B.; de la Chapelle,

[18800] A.: Microdeletions in patients with X-linked muscular dys-

trophy:
molecular-clinical correlations. Clin. Genet. 33: 131-139, 1988.

118. Lindlof, M.; Kiuru, A.; Kaariainen, H.; Kalimo, H.; Lang, H.;
Pihko, H.; Rapola, J.; Somer, H.; Somer, M.; Savontaus, M.-L.; de
la Chapelle, A.: Gene deletions in X-linked muscular dystrophy. Am.
J. Hum. Genet. 44: 496-503, 1989.

119. Mankin, A. S.; Liebman, S. W.: Baby, don't stop! Nature Genet. 23:
8-10, 1999.

120. Mao, Y.; Cremer, M.: Detection of Duchenne muscular dystrophy
carriers by dosage analysis using the DMD cDNA clone 8. Hum. Genet. 81:
193-195, 1989.

[18814]

[18815] 121. Matsuo, M.; Masumura, T.; Nakajima, T.; Kitoh, Y.; Takumi, T.;

[18816] Nishio, H.; Koga, J.; Nakamura, H.: A very small frame-shifting deletion

[18817] within exon 19 of the Duchenne muscular dystrophy gene. Biochem.

[18818] Biophys. Res. Commun. 170: 963-967, 1990.

[18819]

[18820] 122. Matsuo, M.; Masumura, T.; Nishio, H.; Nakajima, T.; Kitoh, Y.;

[18821] Takumi, T.; Koga, J.; Nakamura, H.: Exon skipping during splicing

[18822] of dystrophin mRNA precursor due to an intraexon dele-tion in the dystrophin

[18823] gene of Duchenne muscular dystrophy. J. Clin. Invest. 87: 2127-2131,

[18824] 1991.

[18825]

[18826] 123. McArdle, A.; Edwards, R. H. T.; Jackson, M. J.: Time course

[18827] of changes in plasma membrane permeability in the dys-
trophin-deficient

[18828] mdx mouse. Muscle Nerve 17: 1378-1384, 1994.

[18829]

[18830] 124. McCabe, E. R. B.; Towbin, J.; Chamberlain, J.; Baum-
bach, L.;

[18831] Witkowski, J.; van Ommen, G. J. B.; Koenig, M.; Kunkel, L.
M.; Seltzer,

[18832] W. K.: Complementary DNA probes for the Duchenne
muscular dystrophy

[18833] locus demonstrate a previously undetectable deletion in a
patient

[18834] with dystrophic myopathy, glycerol kinase deficiency, and
congenital

[18835] adrenal hypoplasia. J. Clin. Invest. 83: 95-99, 1989.

[18836]

[18837] 125. Milasin, J.; Muntoni, F.; Severini, G. M.; Bartoloni, L.;
Vatta,

[18838] M.; Krajinovic, M.; Mateddu, A.; Angelini, C.; Camerini, F.;
Falaschi,

[18839] A.; Mestroni, L.; Giacca, M.; Heart Muscle Disease Study
Group:

[18840] A point mutation in the 5-prime splice site of the dystrophin gene

[18841] first intron responsible for X-linked dilated cardiomyopathy. Hum.

[18842] Molec. Genet. 5: 73-79, 1996.

[18843]

[18844] 126. Minetti, C.; Bonilla, E.: Mosaic expression of dystrophin in

[18845] carriers of Becker's muscular dystrophy and the X-linked syndrome

[18846] of myalgia and cramps. (Letter) New Eng. J. Med. 327: 1100, 1992.

[18847]

[18848] 127. Moizard, M.-P.; Toutain, A.; Fournier, D.; Berret, F.; Raynaud,

[18849] M.; Billard, C.; Andres, C.; Moraine, C.: Severe cognitive impairment

[18850] in DMD: obvious clinical indication for Dp71 isoform point mutation

[18851] screening. Europ. J. Hum. Genet. 8: 552-556, 2000.

[18852]

[18853] 128. Monaco, A. P.; Bertelson, C. J.; Liechti-Gallati, S.; Moser,
[18854] H.; Kunkel, L. M.: An explanation for phenotypic differences between
[18855] patients bearing partial deletions of DMD locus. Genomics 2: 90-95,
[18856] 1988.

[18857]

[18858] 129. Monaco, A. P.; Kunkel, L. M.: A giant locus for the Duchenne
[18859] and Becker muscular dystrophy gene. Trends Genet. 3: 33-37, 1987.

[18860]

[18861] 130. Monaco, A. P.; Neve, R. L.; Colletti-Feener, C.; Bertelson, C.
[18862] J.; Kurnit, D. M.; Kunkel, L. M.: Isolation of candidate cDNAs for
[18863] portions of the Duchenne muscular dystrophy gene. Nature 323: 646-650,
[18864] 1986.

[18865]

[18866] 131. Muntoni, F.; Cau, M.; Ganau, A.; Congiu, R.; Arvedi, G.; Mateddu,
[18867] A.; Marrosu, M. G.; Cianchetti, C.; Realdi, G.; Cao, A.; Melis, M.
[18868] A.: Deletion of the dystrophin muscle-promoter region associated
[18869] with X-linked dilated cardiomyopathy. New Eng. J. Med. 329: 921-925,
[18870] 1993.

[18871]

[18872] 132. Muntoni, F.; Melis, M. A.; Ganau, A.; Dubowitz, V.: Transcription
[18873] of the dystrophin gene in normal tissues and in skeletal muscle of
[18874] a family with X-linked dilated cardiomyopathy. Am. J. Hum. Genet. 56:
[18875] 151-157, 1995.

[18876]

[18877] 133. Muntoni, F.; Wilson, L.; Marrosu, G.; Marrosu, M. G.; Cianchetti,
[18878] C.; Mestroni, L.; Ganau, A.; Dubowitz, V.; Sewry, C.: A mutation

[18879] in the dystrophin gene selectively affecting dystrophin expression

[18880] in the heart. J. Clin. Invest. 96: 693-699, 1995.

[18881]

[18882] 134. Nevin, N. C.; Hughes, A. E.; Calwell, M.; Lim, J. H. K.: Duchenne

[18883] muscular dystrophy in a female with a translocation involving Xp21. J.

[18884] Med. Genet. 23: 171-187, 1986.

[18885]

[18886] 135. Nigro, V.; Politano, L.; Nigro, G.; Romano, S. C.; Molinari,

[18887] A. M.; Puca, G. A.: Detection of a nonsense mutation in the dystrophin

[18888] gene by multiple SSCP. Hum. Molec. Genet. 1: 517-520, 1992.

[18889]

[18890] 136. Nobile, C.; Marchi, J.; Nigro, V.; Roberts, R. G.; Danieli, G.

[18891] A.: Exon-intron organization of the human dystrophin gene. Genomics 45:

[18892] 421-424, 1997.

[18893]

[18894] 137. Nobile, C.; Toffolatti, L.; Rizzi, F.; Simionati, B.; Ni-
gro,
[18895] V.; Cardazzo, B.; Patarnello, T.; Valle, G.; Danieli, G. A.:
Analysis
[18896] of 22 deletion breakpoints in dystrophin intron 49. Hum.
Genet. 110:
[18897] 418-421, 2002.

[18898]

[18899] 138. Norman, A.; Harper, P.: A survey of manifesting car-
riers of
[18900] Duchenne and Becker muscular dystrophy in Wales. Clin.
Genet. 36:
[18901] 31-37, 1989.

[18902]

[18903] 139. Ohno, S.: Evolution by Gene Duplication. Berlin:
Springer-Verlag
[18904] (pub.) 1970.

[18905]

[18906] 140. Ortiz-Lopez, R.; Li, H.; Su, J.; Goytia, V.; Towbin, J. A.:
[18907] Evidence for a dystrophin missense mutation as a cause of X-linked
[18908] dilated cardiomyopathy. Circulation 95: 2434-2440, 1997.
[18909]

[18910] 141. Palmucci, L.; Doriguzzi, C.; Mongini, T.; Restagno, G.; Chiado-Piat,
[18911] L.; Maniscalco, M.: Unusual expression and very mild course of Xp21
[18912] muscular dystrophy (Becker type) in a 60-year-old man with 26 percent
[18913] deletion of the dystrophin gene. Neurology 44: 541-543, 1994.
[18914]

[18915] 142. Passos-Bueno, M. R.; Bakker, E.; Kneppers, A. L. J.; Takata,
[18916] R. I.; Rapaport, D.; den Dunnen, J. T.; Zatz, M.; van Om-
men, G. J.
[18917] B.: Different mosaicism frequencies for proximal and dis-
tal Duchenne

[18918] muscular dystrophy (DMD) mutations indicate difference in etiology
[18919] and recurrence risk. Am. J. Hum. Genet. 51: 1150-1155, 1992.
[18920]

[18921] 143. Paulson, K. E.; Deka, N.; Schmid, C. W.; Misra, R.; Schindler,
[18922] C. W.; Rush, M. G.; Kadyk, L.; Leinwand, L.: A transposon-like element
[18923] in human DNA. Nature 316: 359-361, 1985.

[18924]

[18925] 144. Pernelle, J.-J.; Chafey, P.; Chelly, J.; Wahrmann, J. P.; Kaplan,
[18926] J.-C.; Tome, F.; Fardeau, M.: Nebulin seen in DMD males including
[18927] one patient with a large DNA deletion encompassing the DMD gene. Hum.
[18928] Genet. 78: 285, 1988.

[18929]

[18930] 145. Pillers, D.-A. M.; Fitzgerald, K. M.; Duncan, N. M.; Rash, S.

[18931] M.; White, R. A.; Dwinnell, S. J.; Powell, B. R.; Schnur, R. E.; Ray,

[18932] P. N.; Cibis, G. W.; Weleber, R. G.: Duchenne/Becker muscular dystrophy:

[18933] correlation of phenotype by electroretinography with sites of dystrophin

[18934] mutations. Hum. Genet. 105: 2-9, 1999.

[18935]

[18936] 146. Pizzuti, A.; Pieretti, M.; Fenwick, R. G.; Gibbs, R. A.; Caskey,

[18937] C. T.: A transposon-like element in the deletion-prone region of

[18938] the dystrophin gene. Genomics 13: 594-600, 1992.

[18939]

[18940] 147. Porter, J. D.; Khanna, S.; Kaminski, H. J.; Rao, J. S.; Merriam,

[18941] A. P.; Richmonds, C. R.; Leahy, P.; Li, J.; Guo, W.; Andrade, F. H.

[18942] : A chronic inflammatory response dominates the skeletal muscle molecular

[18943] signature in dystrophin-deficient mdx mice. Hum. Molec. Genet. 11:

[18944] 263-272, 2002.

[18945]

[18946] 148. Prior, T. W.; Papp, A. C.; Snyder, P. J.; Burghes, A. H. M.;

[18947] Bartolo, C.; Sedra, M. S.; Western, L. M.; Mendell, J. R.: A missense

[18948] mutation in the dystrophin gene in a Duchenne muscular dystrophy patient. Nature

[18949] Genet. 4: 357-360, 1993.

[18950]

[18951] 149. Prior, T. W.; Papp, A. C.; Snyder, P. J.; Burghes, A. H. M.;

[18952] Sedra, M. S.; Western, L. M.; Bartello, C.; Mendell, J. R.: Identification

[18953] of two point mutations and a one base deletion in exon 19 of the dystrophin

[18954] gene by heteroduplex formation. Hum. Molec. Genet. 2: 311-313, 1993.

[18955]

[18956] 150. Prior, T. W.; Papp, A. C.; Snyder, P. J.; Burghes, A. H. M.;

[18957] Sedra, M. S.; Western, L. M.; Bartolo, C.; Mendell, J. R.: Exon 44

[18958] nonsense mutation in two-Duchenne muscular dystrophy brothers detected

[18959] by heteroduplex analysis. Hum. Mutat. 2: 192-195, 1993.

[18960]

[18961] 151. Prior, T. W.; Papp, A. C.; Snyder, P. J.; Sedra, M. S.; Western,

[18962] L. M.; Bartolo, C.; Moxley, R. T.; Mendell, J. R.: Heterodu-plex analysis

[18963] of the dystrophin gene: application to point mutation and carrier

[18964] detection. Am. J. Med. Genet. 50: 68-73, 1994.

[18965]

[18966] 152. Rafael, J. A.; Sunada, Y.; Cole, N. M.; Campbell, K. P.; Faulkner,

[18967] J. A.; Chamberlain, J. S.: Prevention of dystrophic pathol-ogy in

[18968] mdx mice by a truncated dystrophin isoform. Hum. Molec. Genet. 3:

[18969] 1725-1733, 1994.

[18970]

[18971] 153. Rafael, J. A.; Townsend, E. R.; Squire, S. E.; Potter, A. C.;

[18972] Chamberlain, J. S.; Davies, K. E.: Dystrophin and utrophin influence

[18973] fiber type composition and post-synaptic membrane structure. Hum.

[18974] Molec. Genet. 9: 1357-1367, 2000.

[18975]

[18976] 154. Ray, P. N.; Belfall, B.; Duff, C.; Logan, C.; Kean, V.; Thompson,

[18977] M. W.; Sylvester, J. E.; Gorski, J. L.; Schmickel, R. D.; Wor-

[18978] ton, R. G.: Cloning of the breakpoint of an X;21 translocation associated

[18979] with Duchenne muscular dystrophy. Nature 318: 672-675, 1985.

[18980]

[18981] 155. Read, A. P.; Mountford, R. C.; Forrest, S. M.; Ken-

[18982] wrick, S. J.; Davies, K. E.; Harris, R.: Patterns of exon deletions in Duchenne

[18983] and Becker muscular dystrophy. Hum. Genet. 80:
152-156, 1988.

[18984]

[18985] 156. Rininsland, F.; Hahn, A.; Niemann-Seyde, S.; Slomski,
R.; Hanefeld,
[18986] F.; Reiss, J.: Identification of a new DMD gene deletion by
ectopic
[18987] transcript analysis. J. Med. Genet. 29: 647-651, 1992.

[18988]

[18989] 157. Roberts, R. G.; Bentley, D. R.; Bobrow, M.: Infidelity in
the
[18990] structure of ectopic transcripts: a novel exon in lympho-
cyte dystrophin
[18991] transcripts. Hum. Mutat. 2: 293-299, 1993.

[18992]

[18993] 158. Roberts, R. G.; Bobrow, M.; Bentley, D. R.: The spec-
trum of
[18994] mild X-linked recessive muscular dystrophy. Arch. Neurol.
34: 408-416,
[18995] 1992.

[18996]

[18997] 159. Roberts, R. G.; Bobrow, M.; Bentley, D. R.: Point mutations
[18998] in the dystrophin gene. Proc. Nat. Acad. Sci. 89: 2331-2335, 1992.
[18999]

[19000] 160. Roberts, R. G.; Gardner, R. J.; Bobrow, M.: Searching for the
[19001] 1 in 2,400,000: a review of dystrophin gene point mutations. Hum.
[19002] Mutat. 4: 1-11, 1994.
[19003]

[19004] 161. Roberts, R. G.; Passos-Bueno, M. R.; Bobrow, M.; Vainzof, M.;
[19005] Zatz, M.: Point mutation in a Becker muscular dystrophy patient. Hum.
[19006] Molec. Genet. 2: 75-77, 1992.
[19007]

[19008] 162. Rowland, L. P.: Biochemistry of muscle membranes in Duchenne
[19009] muscular dystrophy. Muscle Nerve 3: 3-20, 1980.
[19010]

[19011] 163. Ryder-Cook, A. S.; Sicinski, P.; Thomas, K.; Davies, K. E.; Worton,
[19012] R. G.; Barnard, E. A.; Darlison, M. G.; Barnard, P. J.: Local-
ization
[19013] of the mdx mutation within the mouse dystrophin gene. EMBO J. 7:
[19014] 3017-3021, 1988.

[19015]

[19016] 164. Saad, F. A.; Vita, G.; Mora, M.; Morandi. L.; Vitiello, L.; Oliviero,
[19017] S.; Danieli, G. A.: A novel nonsense mutation in the human dystrophin
[19018] gene. Hum. Mutat. 2: 314-316, 1993.

[19019]

[19020] 165. Saad, F. A.; Vita, G.; Toffolatti, L.; Danieli, G. A.: A possible
[19021] missense mutation detected in the dystrophin gene by double strand
[19022] conformation analysis (DSCA). Neuromusc. Disord. 4:
335-341, 1994.

[19023]

[19024] 166. Sakamoto, M.; Yuasa, K.; Yoshimura, M.; Yokota, T.; Ikemoto,
[19025] T.; Suzuki, M.; Dickson, G.; Miyagoe-Suzuki, Y.; Takeda, S.: Micro-dystrophin
[19026] cDNA ameliorates dystrophic phenotypes when introduced into mdx mice
[19027] as a transgene. Biochem. Biophys. Res. Commun. 293: 1265-1272, 2002.
[19028]

[19029] 167. Sarig, R.; Mezger-Lallemand, V.; Gitelman, I.; Davis, C.; Fuchs,
[19030] O.; Yaffe, D.; Nudel, U.: Targeted inactivation of Dp71, the major
[19031] non-muscle product of the DMD gene: differential activity of the Dp71
[19032] promoter during development. Hum. Molec. Genet. 8: 1-10, 1999.
[19033]

[19034] 168. Sarkar, G.; Sommer, S. S.: Access to a messenger RNA sequence
[19035] or its protein product is not limited by tissue or species specificity. Science 244:

[19036] 331-334, 1989.

[19037]

[19038] 169. Schwartz, L. S.; Tarleton, J.; Popovich, B.; Seltzer, W. K.;

[19039] Hoffman, E. P.: Fluorescent multiplex linkage analysis and carrier

[19040] detection for Duchenne/Becker muscular dystrophy. Am. J. Hum. Genet. 51:

[19041] 721-729, 1992.

[19042]

[19043] 170. Scott, M. O.; Sylvester, J. E.; Heiman-Patterson, T.; Shi, Y.-J.;

[19044] Fieles, W.; Stedman, H.; Burghes, A.; Ray, P.; Worton, R.; Fischbeck,

[19045] K. H.: Duchenne muscular dystrophy gene expression in normal and

[19046] diseased human muscle. Science 239: 1418-1420, 1988.

[19047]

[19048] 171. Sharp, N. J. H.; Kornegay, J. N.; Van Camp, S. D.; Herbstreith,

[19049] M. H.; Secore, S. L.; Kettle, S.; Hung, W.-Y.; Constantinou,

C. D.;

[19050] Dykstra, M. J.; Roses, A. D.; Bartlett, R. J.: An error in dystrophin

[19051] mRNA processing in golden retriever muscular dystrophy, an animal

[19052] homologue of Duchenne muscular dystrophy. Genomics 13: 115-121,

[19053] 1992.

[19054]

[19055] 172. Shiga, N.; Takeshima, Y.; Sakamoto, H.; Inoue, K.; Yokota, Y.;

[19056] Yokoyama, M.; Matsuo, M.: Disruption of the splicing enhancer sequence

[19057] within exon 27 of the dystrophin gene by a nonsense mutation induces

[19058] partial skipping of the exon and is responsible for Becker muscular

[19059] dystrophy. J. Clin. Invest. 100: 2204-2210, 1997.

[19060]

[19061] 173. Sicinski, P.; Geng, Y.; Ryder-Cook, A. S.; Barnard, E. A.; Darlison,

[19062] M. G.; Barnard, P. J.: The molecular basis of muscular dys-

trophy
[19063] in the mdx mouse: a point mutation. Science 244:
1578-1580, 1989.
[19064]

[19065] 174. Smithies, O.; Connell, G. E.; Dixon, G. H.: Chromoso-
mal rearrangements
[19066] and the evolution of haptoglobin genes. Nature 196:
232-236, 1962.
[19067]

[19068] 175. Southern, E. M.: Detection of specific sequences
among DNA fragments
[19069] separated by gel electrophoresis. J. Molec. Biol. 98:
503-517, 1975.
[19070]

[19071] 176. Stratford-Perricaudet, L. D.; Makeh, I.; Perricaudet,
M.; Briand,
[19072] P.: Widespread long-term gene transfer to mouse skeletal
muscles
[19073] and heart. J. Clin. Invest. 90: 626-630, 1992.

[19074]

[19075] 177. Takeshima, Y.; Nishio, H.; Narita, N.; Wada, H.;

Ishikawa, Y.;

[19076] Ishikawa, Y.; Minami, R.; Nakamura, H.; Matsuo, M.: Amino-terminal

[19077] deletion of 53% of dystrophin results in an intermediate Duchenne-Becker

[19078] muscular dystrophy phenotype. Neurology 44: 1648-1651, 1994.

[19079]

[19080] 178. Tennyson, C. N.; Klamut, H. J.; Worton, R. G.: The human dystrophin

[19081] gene requires 16 hours to be transcribed and is cotranscriptionally

[19082] spliced. Nature Genet. 9: 184-190, 1995.

[19083]

[19084] 179. Tinsley, J. M.; Blake, D. J.; Davies, K. E.: Apodystrophin-3:

[19085] a 2.2kb transcript from the DMD locus encoding the dystrophin glycoprotein

[19086] binding site. Hum. Molec. Genet. 2: 521-524, 1993.

[19087]

[19088] 180. Tinsley, J. M.; Potter, A. C.; Phelps, S. R.; Fisher, R.;

Trickett,
[19089] J. I.; Davies, K. E.: Amelioration of the dystrophic pheno-
type of
[19090] mdx mice using a truncated utrophin transgene. Nature
384: 349-353,
[19091] 1996.

[19092]

[19093] 181. Todorova, A.; Danieli, G. A.: Large majority of single-
nucleotide
[19094] mutations along the dystrophin gene can be explained by
more than
[19095] one mechanism of mutagenesis. Hum. Mutat. 9: 537-547,
1997.

[19096]

[19097] 182. Torelli, S.; Muntoni, F.: Alternative splicing of dys-
trophin
[19098] exon 4 in normal human muscle. Hum. Genet. 97:
521-523, 1996.

[19099]

[19100] 183. Towbin, J. A.; Hejtmancik, J. F.; Brink, P.; Gelb, B.;
Zhu, X.
[19101] M.; Chamberlain, J. S.; McCabe, E. R. B.; Swift, M.: X-linked

dilated

[19102] cardiomyopathy: molecular genetic evidence of linkage to the Duchenne

[19103] muscular dystrophy (dystrophin) gene at the Xp21 locus. Circulation 87:

[19104] 1854-1865, 1993.

[19105]

[19106] 184. Towbin, J. A.; Ortiz-Lopez, R.: X-linked dilated car-diomyopathy.

[19107] (Letter) New Eng. J. Med. 330: 369-370, 1994.

[19108]

[19109] 185. Towbin, J. A.; Zhu, X. M.; Gelb, B.; Bies, R.; Chamber-lain, J.;

[19110] Maichele, A.; Ohlendieck, K.; Campbell, K.; McCabe, E. R. B.; Swift,

[19111] M.: X-linked dilated cardiomyopathy (XLCM): molecular characterization.

[19112] (Abstract) Am. J. Hum. Genet. 49 (suppl.): 421, 1991.

[19113]

[19114] 186. Tuffery, S.; Lenk, U.; Roberts, R. G.; Coubes, C.; De-maille,

[19115] J.; Claustres, M.: Protein truncation test: analysis of two novel

[19116] point mutations at the carboxy-terminus of the human dystrophin gene

[19117] associated with mental retardation. Hum. Mutat. 6: 126-135, 1995.

[19118]

[19119] 187. Valentine, B. A.; Winand, N. J.; Pradhan, D.; Moise, N. S.; de

[19120] Lahunta, A.; Kornegay, J. N.; Cooper, B. J.: Canine X-linked muscular

[19121] dystrophy as an animal model of Duchenne muscular dystrophy: a review. Am.

[19122] J. Med. Genet. 42: 352-356, 1992.

[19123]

[19124] 188. Verellen-Dumoulin, C.; Freund, M.; De Meyer, R.; Laterre, C.;

[19125] Frederic, J.; Thompson, M. W.; Markovic, V. D.; Worton, R. G.: Expression

[19126] of an X-linked muscular dystrophy in a female due to translocation

[19127] involving Xp21 and non-random inactivation of the nor-

mal X chromosome. Hum.
[19128] Genet. 67: 115-119, 1984.

[19129]

[19130] 189. Wehling, M.; Spencer, M. J.; Tidball, J. G.: A nitric ox-
ide
[19131] synthase transgene ameliorates muscular dystrophy in
mdx mice. J.
[19132] Cell Biol. 155: 123-131, 2001.

[19133]

[19134] 190. Werner, W.; Spiegler, A. W. J.: Inherited deletion of
subband
[19135] Xp21.13 in a male with Duchenne muscular dystrophy. J.
Med. Genet. 25:
[19136] 377-382, 1988.

[19137]

[19138] 191. Wilton, S. D.; Chandler, D. C.; Kakulas, B. A.; Laing,
N. G.
[19139] : Identification of a point mutation and germinal mo-
saicism in a Duchenne
[19140] muscular dystrophy family. Hum. Mutat. 3: 133-140,
1994.

[19141]

[19142] 192. Wilton, S. D.; Johnsen, R. D.; Pedretti, J. R.; Laing, N. G.

[19143] : Two distinct mutations in a single dystrophin gene: identification

[19144] of an altered splice-site as the primary Becker muscular dystrophy

[19145] mutation. Am. J. Med. Genet. 46: 563-569, 1993.

[19146]

[19147] 193. Winnard, A. V.; Jia-Hsu, Y.; Gibbs, R. A.; Mendell, J. R.; Burghes,

[19148] A. H. M.: Identification of a 2 base pair nonsense mutation causing

[19149] a cryptic splice site in a DMD patient. Hum. Molec. Genet. 1: 645-646,

[19150] 1992.

[19151]

[19152] 194. Wood, D. S.; Zeviani, M.; Prelle, A.; Bonilla, E.; Salviati,

[19153] G.; Miranda, A. F.; DiMauro, S.; Rowland, L. P.: Is nebulin the defective

[19154] gene product in Duchenne muscular dystrophy? (Letter) New Eng. J.
[19155] Med. 316: 107-108, 1987.

[19156]

[19157] 195. Worton, R. G.: Dystrophin: the long and short of it. (Editorial) J.
[19158] Clin. Invest. 93: 4, 1994.

[19159]

[19160] 196. Worton, R. G.: Personal Communication. Toronto, Ontario, Canada
[19161] 9/12/1987.

[19162]

[19163] 197. Xiong, D.; Lee, G.-H.; Badorff, C.; Dorner, A.; Lee, S.; Wolf,
[19164] P.; Knowlton, K. U.: Dystrophin deficiency markedly in-
creases enterovirus-induced
[19165] cardiomyopathy: a genetic predisposition to viral heart disease. Nature
[19166] Med. 8: 872-877, 2002.

[19167]

[19168] 198. Yang, T. P.; Patel, P. I.; Chinault, A. C.; Stout, J. T.; Jackson,

[19169] L. G.; Hildebrand, B. M.; Caskey, C. T.: Molecular evidence for new

[19170] mutation at the HPRT locus in Lesch-Nyhan patients. Nature 310:

[19171] 412-414, 1984.

[19172]

[19173] 199. Yoshida, K.; Ikeda, S.; Nakamura, A.; Kagoshima, M.; Takeda,

[19174] S.; Shoji, S.; Yanagisawa, N.: Molecular analysis of the Duchenne

[19175] muscular dystrophy gene in patients with Becker muscular dystrophy

[19176] presenting with dilated cardiomyopathy. Muscle Nerve 16: 1161-1166,

[19177] 1993.

[19178]

[19179] 200. Yoshida, K.; Nakamura, A.; Yazaki, M.; Ikeda, S.; Takeda, S.

[19180] : Insertional mutation by transposable element, L1, in the DMD gene

[19181] results in X-linked dilated cardiomyopathy. Hum. Molec. Genet. 7:

[19182] 1129-1132, 1998.

[19183]

[19184] 201. Zubrzycka-Gaarn, E. E.; Bulman, D. E.; Karpati, G.; Burghes,

[19185] A. H. M.; Belfall, B.; Klamut, H. J.; Talbot, J.; Hodges, R. S.; Ray,

[19186] P. N.; Worton, R. G.: The Duchenne muscular dystrophy gene product

[19187] is localized in sarcolemma of human skeletal muscle. Na- ture 333:

[19188] 466-469, 1988.

[19189] 1956.

[19190] Brinkhous, K. M.; Davis, P. D.; Graham, J. B.; Dodds, W. J.:

[19191] Expression and linkage of genes for X-linked hemophilia A and B in

[19192] the dog. Blood 41: 577-585, 1973.

[19193] 1957.

[19194] Abinun, M.; Spickett, G.; Appleton, A. L.; Flood, T.; Cant, A.

[19195] J.: Anhidrotic ectodermal dysplasia associated with specific antibody
[19196] deficiency. Europ. J. Pediat. 155: 146-147, 1995.

[19197] 1958.

[19198] Bayes, M.; Hartung, A. J.; Ezer, S.; Pispa, J.; Thesleff, I.; Srivastava,
[19199] A. K.; Kere, J.: The anhidrotic ectodermal dysplasia gene (EDA) undergoes
[19200] alternative splicing and encodes ectodysplasin-A with deletion mutations
[19201] in collagenous repeats. Hum. Molec. Genet. 7: 1661-1669, 1998.

[19202] 1959.

[19203] Blaschko, A.: Die Nervenverteilung in der Haut in ihrer Beziehung
[19204] zu den Erkrankungen der Haut. Wein and Leipzig: Braunmuller (pub.)

[19205] 1901.

[19206] 1960.

[19207] Blecher, S. R.; Kapalanga, J.; Lalonde, D.: Induction of sweat

[19208] glands by epidermal growth factor in murine X-linked anhidrotic ectodermal
[19209] dysplasia. Nature 345: 542-544, 1990.

[19210] 1961.

[19211] Bowen, R.: Hereditary ectodermal dysplasia of the anhidrotic type. Sth.
[19212] Med. J. 50: 1018-1021, 1957.

[19213] 1962.

[19214] Burck, U.; Held, K. R.: Athelia in a female infant heterozygous
[19215] for anhidrotic ectodermal dysplasia. Clin. Genet. 19: 117-121, 1981.
[19216] 1963.

[19217] Casal, M. L.; Jezyk, P. F.; Greek, J. M.; Goldschmidt, M. H.; Patterson,
[19218] D. F.: X-linked ectodermal dysplasia in the dog. J. Hered. 88:
[19219] 513-516, 1997.

[19220] 1964.

[19221] Chautard-Freire-Maia, E. A.; Primo-Parmo, S. L.; Pinheiro, M.;

[19222] Freire-Maia, N.: Further evidence against linkage between Christ-Siemens-Touraine
[19223] (CST) and Xg loci. Hum. Genet. 57: 205-206, 1981.

[19224] 1965.

[19225] Chen, Y.; Molloy, S. S.; Thomas, L.; Gambee, J.; Bachinger, H.
[19226] P.; Ferguson, B.; Zonana, J.; Thomas, G.; Morris, N. P.: Mutations
[19227] within a furin consensus sequence block proteolytic release of ectodysplasin-A
[19228] and cause X-linked hypohidrotic ectodermal dysplasia. Proc. Nat.
[19229] Acad. Sci. 98: 7218-7223, 2001.

[19230] 1966.

[19231] Clark, R. P.; Goff, M. R.; MacDermot, K. D.: Identification of
[19232] functioning sweat pores and visualization of skin temperature patterns
[19233] in X-linked hypohidrotic ectodermal dysplasia by whole body thermography. Hum.
[19234] Genet. 86: 7-13, 1990.

[19235] 1967.

[19236] Brinkhous, K. M.; Graham, J. B.: Hemophilia in the female dog. Science 111:

[19237] 723-724, 1950.

[19238] 1968.

[19239] Brocker-Vriends, A. H. J. T.; Briet, E.; Dreesen, J. C. F. M.;

[19240] Bakker, B.; Reitsma, P.; Pannekoek, H.; van de Kamp, J. J. P.; Pearson,

[19241] P. L.: Somatic origin of inherited haemophilia A. Hum. Genet. 85:

[19242] 288-292, 1990.

[19243] 1969.

[19244] Brocker-Vriends, A. H. J. T.; Rosendaal, F. R.; van Houwelingen,

[19245] J. C.; Bakker, E.; van Ommen, G. J. B.; van de Kamp, J. J. P.; Briet,

[19246] E.: Sex ratio of the mutation frequencies in haemophilia A: coagulation

[19247] assays and RFLP analysis. J. Med. Genet. 28: 672-680, 1991.

[19248] 1970.

[19249] Casarino, L.; Pecorara, M.; Mori, P. G.; Morfini, M.; Mancuso,
[19250] G.; Scrivano, L.; Molinari, A. C.; Lanza, T.; Giavarella, G.; Loi,
[19251] A.; Perseu, L.; Cao, A.; Pirastu, M.: Molecular basis of hemophilia
[19252] A in Italians. Ric. Clin. Lab. 16: 227, 1986.

[19253] 1971.

[19254] Casula, L.; Murru, S.; Pecorara, M.; Ristaldi, M. S.; Restagno,
[19255] G.; Mancuso, G.; Morfini, M.; De Biasi, R.; Baudo, F.; Carbonara,
[19256] A.; Mori, P. G.; Cao, A.; Pirastu, M.: Recurrent mutations and three
[19257] novel rearrangements in the factor VIII gene of hemophilia A patients
[19258] of Italian descent. Blood 75: 662-670, 1990.

[19259] 1972.

[19260] Chan, V.; Chan, T. K.; Tong, T. M.; Todd, D.: A novel missense
[19261] mutation in exon 4 of the factor VIII:C gene resulting in moderately

[19262] severe hemophilia A. Blood 74: 2688-2691, 1989.

[19263] 1973.

[19264] Chediak, J.; Telfer, M. C.; Jaojaroenkul, T.; Green, D.: Lower

[19265] factor VIII coagulant activity in daughters of subjects with hemophilia

[19266] A compared to other obligate carriers. Blood 55: 552-558, 1980.

[19267] 1974.

[19268] Coleman, R.; Genet, S. A.; Harper, J. I.; Wilkie, A. O. M.: In-teraction

[19269] of incontinentia pigmenti and factor VIII mutations in a fe-male with

[19270] biased X inactivation, resulting in haemophilia. J. Med. Genet. 30:

[19271] 497-500, 1993.

[19272] 1975.

[19273] Cone, T. E., Jr.: A case of hereditary hemorrhagic tendency (hemophilia)

[19274] reported in 1832 by a physician practicing in Simpsonville, Kentucky. Pediatrics 64:

[19275] 291, 1979.

[19276] 1976.

[19277] Cooper, H. A.; Wagner, R. H.: The defect in hemophilic and Von

[19278] Willebrand's disease. Plasmas studied by a recombination technique. J.

[19279] Clin. Invest. 54: 1093-1099, 1974.

[19280] 1977.

[19281] Cutler, J. A.; Mitchell, M. J.; Smith, M. P.; Savidge, G. F.:

[19282] The identification and classification of 41 novel mutations in the

[19283] factor VIII gene (F8C). Hum. Mutat. 19: 274-278, 2002.

[19284] 1978.

[19285] David, D.; Moreira, I.; Lalloz, M. R.; Rosa, H. A.; Schwaab, R.;

[19286] Morais, S.; Diniz, M. J.; de Deus, G.; Campos, M.; Lavinha, J.: Analysis

[19287] of the essential sequences of the factor VIII gene in twelve haemophilia

[19288] A patients by single-stranded conformation polymor-phism. Blood Coagul.

[19289] Fibrinolysis 5: 257-264, 1994.

[19290] 1979.

[19291] Denson, K. W. E.: Two forms of haemophilia? (Letter) Lancet II:

[19292] 222-223, 1968.

[19293] 1980.

[19294] Denson, K. W. E.; Biggs, R.; Haddon, M. E.; Borrett, R.; Cobb,

[19295] K.: Two types of haemophilia (A+ and A-): a study of 48 cases. Brit.

[19296] J. Haemat. 17: 163-171, 1969.

[19297] 1981.

[19298] Diamond, C.; Kogan, S.; Levinson, B.; Gitschier, J.: Amino acid

[19299] substitutions in conserved domains of factor VIII and related proteins:

[19300] study of patients with mild and moderately severe hemophilia A. Hum.

[19301] Mutat. 1: 248-257, 1992.

[19302] 1982.

[19303] Din, N.; Schwartz, M.; Kruse, T.; Vestertgaard, S. R.; Ahrens,
[19304] P.; Scheibel, E.; Nordfang, O.; Ezban, M.: Factor VIII gene-specific
[19305] probes used to study heritage and molecular defects in hemophilia
[19306] A. (Abstract) Ric. Clin. Lab. 16: 182 only, 1986.

[19307] 1983.

[19308] Dombroski, B. A.; Mathias, S. L.; Nanthakumar, E. J.; Scott, A.
[19309] F.; Kazazian, H. H., Jr.: Isolation of the L1 gene responsi-ble for
[19310] a retrotransposition event in man. (Abstract) Am. J. Hum. Genet. 49
[19311] (suppl.): 403 only, 1991.

[19312] 1984.

[19313] Dwarki, V. J.; Belloni, P.; Nijjar, T.; Smith, J.; Couto, L.;

[19314] Rabier, M.; Clift, S.; Berns, A.; Cohen, L. K.: Gene therapy for
[19315] hemophilia A: production of therapeutic levels of human factor VIII

[19316] in vivo in mice. Proc. Nat. Acad. Sci. 92: 1023-1027,
1995.
[19317] 1985.
[19318] Economou, E. P.; Kazazian, H. H., Jr.; Antonarakis, S. E.:
Detection
[19319] of mutations in the factor VIII gene using single-stranded
conformational
[19320] polymorphism (SSCP). Genomics 13: 909-911, 1992.
[19321] 1986.
[19322] Edgell, C.-J. S.; Kirkman, H. N.; Clemons, E.; Buchanan, P.
D.;
[19323] Miller, C. H.: Prenatal diagnosis by linkage: hemophilia A
and polymorphic
[19324] glucose-6-phosphate dehydrogenase. Am. J. Hum. Genet.
30: 80-84,
[19325] 1978.
[19326] 1987.
[19327] Erlich, H. A.; Gelfand, D. H.; Saiki, R. K.: Specific DNA am-
plification. Nature 331:
[19328] 461-462, 1988.
[19329] 1988.

[19330] Favier, R.; Lavergne, J.-M.; Costa, J.-M.; Caron, C.; Mazurier,
[19331] C.; Viemont, M.; Delpech, M.; Valleix, S.: Unbalanced X-chromosome
[19332] inactivation with a novel FVIII gene mutation resulting in severe
[19333] hemophilia A in a female. Blood 96: 4373-4375, 2000.

[19334] 1989.

[19335] Fay, P. J.; Chavin, S. I.; Schroeder, D.; Young, F. E.; Marder,

[19336] V. J.: Purification and characterization of a highly purified human
[19337] factor VIII consisting of a single type of polypeptide chain. Proc.
[19338] Nat. Acad. Sci. 79: 7200-7204, 1982.

[19339] 1990.

[19340] Feinstein, D.; Chong, M. N. Y.; Kasper, C. K.; Rapaport, S. I.
[19341] : Hemophilia A: polymorphism detectable by a factor VIII antibody. Science 163:
[19342] 1071-1072, 1969.

[19343] 1991.

[19344] Figueiredo, M. S.; Bernardi, F.; Zago, M. A.: A novel deletion
[19345] of FVIII gene associated with variable levels of FVIII inhibitor. Europ.
[19346] J. Haemat. 48: 152-154, 1992.

[19347] 1992.

[19348] Filippi, G.; Mannucci, P. M.; Coppola, R.; Farris, A.; Rinaldi,
[19349] A.; Siniscalco, M.: Studies on hemophilia A in Sardinia bearing on
[19350] the problems of multiple allelism, carrier detection, and differential
[19351] mutation rate in the two sexes. Am. J. Hum. Genet. 36: 44-71, 1984.
[19352] 1993.

[19353] Firshein, S. I.; Hoyer, L. W.; Lazarchick, J.; Forget, B. G.;
[19354] Hobbins, J. C.; Clyne, L. P.; Pitlick, F. A.; Muir, W. A.; Merkatz,
[19355] I. R.; Mahoney, M. J.: Prenatal diagnosis of classic hemophilia. New
[19356] Eng. J. Med. 300: 937-941, 1979.

[19357] 1994.

[19358] Freije, D.; Schlessinger, D.: A 1.6-Mb contig of yeast artificial
[19359] chromosomes around the human factor VIII gene reveals three regions
[19360] homologous to probes for the DXS115 locus and two for the DXYS64 locus. Am.
[19361] J. Hum. Genet. 51: 66-80, 1992.

[19362] 1995.

[19363] Frommel, D.; Muller, J. Y.; Prou-Wartelle, O.; Allain, J. P.:

[19364] Possible linkage between the major histocompatibility complex and
[19365] the immune response to factor VIII in classic haemophilia. Vox Sang. 33:
[19366] 270-272, 1977.

[19367] 1996.

[19368] Zankl, A.; Addor, M.-C.; Cousin, P.; Gaide, A.-C.; Gudinchet,
[19369] F.; Schorderet, D. F.: Fatal outcome in a female monozygotic twin
[19370] with X-linked hypohydrotic (sic) ectodermal dysplasia (XLHED) due

[19371] to a de novo t(X;9) translocation with probable disruption of the

[19372] EDA gene. Europ. J. Pediat. 160: 296-299, 2001.

[19373] 1997.

[19374] Tanner, S.; Stagljar, I.; Georgiev, O.; Schaffner, W.; Bourquin,

[19375] J.-P.: A novel SR-related protein specifically interacts with the

[19376] carboxy-terminal domain (CTD) of RNA polymerase II through a conserved

[19377] interaction domain. Biol. Chem. 378: 565-571, 1997.

[19378] 1998.

[19379] Zhang, W.-J.; Wu, J. Y.: Sip1, a novel RS domain-con-
taining protein
[19380] essential for pre-mRNA splicing. Molec. Cell. Biol. 18:
676-684,
[19381] 1998.

[19382] 1999.

[19383] Puder, M.; Barnard, G. F.; Staniunas, R. J.; Steele, G. D., Jr.;

[19384] Chen, L. B.: Nucleotide and deduced amino acid sequence
of human
[19385] ribosomal protein L18. Biochim. Biophys. Acta 1216:
134-136, 1993.
[19386] 2000.

[19387] Johnson, K. R.: Characterization of cDNA clones encoding
the human
[19388] homologue of Saccharomyces cerevisiae ribosomal protein
L30. Gene 123:
[19389] 283-285, 1993.

[19390] 2001.

[19391] Tsukaguchi, H.; Tokui, T.; Mackenzie, B.; Berger, U. V.; Chen,
[19392] X.-Z.; Wang, Y.; Brubaker, R. F.; Hediger, M. A.: A family of mammalian
[19393] Na(+)-dependent L-ascorbic acid transporters. Nature 399: 70-75,
[19394] 1999.

[19395] 2002.

[19396] Li, X.; Tedder, T. F.: CHST1 and CHST2 sulfotransferases expressed
[19397] by human vascular endothelial cells: cDNA cloning, ex- pression, and
[19398] chromosomal localization. Genomics 55: 345-347, 1999.

[19399] 2003.

[19400] Uchimura, K.; Muramatsu, H.; Kadomatsu, K.; Fan, Q.-W.; Kurosawa,
[19401] N.; Mitsuoka, C.; Kannagi, R.; Habuchi, O.; Muramatsu, T.: Molecular
[19402] cloning and characterization of an N- acetylglucosamine-6-O-sulfotransferase. J.
[19403] Biol. Chem. 273: 22577-22583, 1998.

[19404] 2004.

[19405] Uchimura, K.; Muramatsu, H.; Kaname, T.; Ogawa, H.; Ya-
makawa, T.;

[19406] Fan, Q.-W.; Mitsuoka, C.; Kannagi, R.; Habuchi, O.;
Yokoyama, I.;

[19407] Yamamura, K.; Ozaki, T.; Nakagawara, A.; Kadomatsu, K.;
Muramatsu,

[19408] T.: Human N-acetylglucosamine-6-O-sulfotransferase in-
volved in the

[19409] biosynthesis of 6-sulfo sialyl Lewis X: molecular cloning,
chromosomal

[19410] mapping, and expression in various organs and tumor
cells. J. Biochem. 124:

[19411] 670-678, 1998.

[19412] 2005.

[19413] Ayres, J. A.; Shum, L.; Akarsu, A. N.; Dashner, R.; Taka-
hashi,

[19414] K.; Ikura, T.; Slavkin, H. C.; Nuckolls, G. H.: DACH: ge-
nomic characterization,

[19415] evaluation as a candidate for postaxial polydactyly type
A2, and developmental

[19416] expression pattern of the mouse homologue. Genomics

77: 18-26, 2001.

[19417] 2006.

[19418] Davis, R. J.; Shen, W.; Sandler, Y. I.; Amoui, M.; Purcell, P.;

[19419] Maas, R.; Ou, C.-N.; Vogel, H.; Beaudet, A. L.; Mardon, G.: Dach1

[19420] mutant mice bear no gross abnormalities in eye, limb, and brain development

[19421] and exhibit postnatal lethality. Molec. Cell. Biol. 21: 1484-1490,

[19422] 2001.

[19423] 2007.

[19424] Hammond, K. L.; Hanson, I. M.; Brown, A. G.; Lettice, L. A.; Hill,

[19425] R. E.: Mammalian and Drosophila dachshund genes are related to the

[19426] Ski proto-oncogene and are expressed in eye and limb. Mech. Dev. 74:

[19427] 121-131, 1998.

[19428] 2008.

[19429] Hammond, K. L.; Lettice, L. A.; Hill, R. E.; Lee, M.; Boyle, S.;

[19430] Hanson, I. M.: Human (DACH) and mouse (Dach) homo-

logues of Drosophila
[19431] dachshund map to chromosomes 13q22 and 14E3, respectively. Genomics 55:
[19432] 252-253, 1999.

[19433] 2009.

[19434] Li, X.; Perissi, V.; Liu, F.; Rose, D. W.; Rosenfeld, M. G.: Tissue-specific
[19435] regulation of retinal and pituitary precursor cell proliferation. Science 297:
[19436] 1180-1183, 2002.

[19437] 2010.

[19438] Blanco, G.; Irving, N. G.; Brown, S. D. M.; Miller, C. C. J.; McLoughlin,
[19439] D. M.: Mapping of the human and murine X11-like genes (APBA2 and
[19440] Apba2), the murine Fe65 gene (Apbb1), and the human Fe65-like gene
[19441] (APBB2): genes encoding phosphotyrosine-binding domain proteins that
[19442] interact with the Alzheimer's disease amyloid precursor protein. Mammalian
[19443] Genome 9: 473-475, 1998.

[19444] 2011.

[19445] Butz, S.; Okamoto, M.; Sudhof, T. C.: A tripartite protein complex

[19446] with the potential to couple synaptic vesicle exocytosis to cell adhesion

[19447] in brain. Cell 94: 773-782, 1998.

[19448] 2012.

[19449] Chen, W.-J.; Goldstein, J. L.; Brown, M. S.: NPXY, a sequence

[19450] often found in cytoplasmic tails, is required for coated pit-mediated

[19451] internalization of the low density lipoprotein receptor. J. Biol.

[19452] Chem. 265: 3116-3123, 1990.

[19453] 2013.

[19454] Duclos, F.; Boschert, U.; Sirugo, G.; Mandel, J.-L.; Hen, R.; Koenig,

[19455] M.: Gene in the region of the Friedreich ataxia locus encodes a putative

[19456] transmembrane protein expressed in the nervous system. Proc. Nat.

[19457] Acad. Sci. 90: 109-113, 1993.

[19458] 2014.

[19459] Duclos, F.; Koenig, M.: Comparison of primary structure of a neuron-specific

[19460] protein, X11, between human and mouse. Mammalian Genome 6: 57-58,

[19461] 1995.

[19462] 2015.

[19463] Okamoto, M.; Sudhof, T. C.: Mints, Munc18-interacting proteins

[19464] in synaptic vesicle exocytosis. J. Biol. Chem. 272: 31459-31464,

[19465] 1997.

[19466] 2016.

[19467] van der Geer, P.; Pawson, T.: The PTB domain: a new protein module

[19468] implicated in signal transduction. Trends Biochem. Sci. 20: 277-280,

[19469] 1995.

[19470] 2017.

[19471] Loh, N. Y.; Ambrose, H. J.; Guay-Woodford, L. M.; Das-Gupta, S.;
[19472] Nawrotzki, R. A.; Blake, D. J.; Davies, K. E.: Genomic organization
[19473] and refined mapping of the mouse beta-dystrobrevin gene. Mammalian
[19474] Genome 9: 857-862, 1998.

[19475] 2018.

[19476] Rodius, F.; Duclos, F.; Wrogemann, K.; Le Paslier, D.; Ougen, P.;
[19477] Billault, A.; Belal, S.; Musenger, C.; Brice, A.; Durr, A.; Mignard,
[19478] C.; Sirugo, G.; Weissenbach, J.; Cohen, D.; Hentati, F.; Ben Hamida,
[19479] M.; Mandel, J.-L.; Koenig, M.: Recombinations in individuals homozygous
[19480] by descent localize the Friedreich ataxia locus in a cloned 450-kb
[19481] interval. Am. J. Hum. Genet. 54: 1050-1059, 1994.

[19482] 2019.

[19483] Peters, M. F.; O'Brien, K. F.; Sadoulet-Puccio, H. M.; Kunkel,

[19484] L. M.; Adams, M. E.; Froehner, S. C.: Beta-dystrobrevin, a new member
[19485] of the dystrophin family: identification, cloning, and protein associations. J.
[19486] Biol. Chem. 272: 50:-31561-31569, 1997.

[19487] 2020.

[19488] Morris, M. E.; Viswanathan, N.; Kuhlman, S.; Davis, F. C.; Weitz,
[19489] C. J.: A screen for genes induced in the suprachiasmatic nucleus
[19490] by light. Science 279: 1544-1547, 1998.

[19491] 2021.

[19492] Patwardhan, S.; Gashler, A.; Siegel, M. G.; Chang, L. C.; Joseph,
[19493] L. J.; Shows, T. B.; Le Beau, M. M.; Sukhatme, V. P.: EGR3, a novel
[19494] member of the Egr family of genes encoding immediate-early transcription
[19495] factors. Oncogene 6: 917-928, 1991.

[19496] 2022.

[19497] Tourtellotte, W. G.; Milbrandt, J.: Sensory ataxia and mus-

cle
[19498] spindle agenesis in mice lacking the transcription factor Egr3. Nature
[19499] Genet. 20: 87-91, 1998.

[19500] 2023.

[19501] Anand, R.; Ogilvie, D. J.; Butler, R.; Riley, J. H.; Finniear,

[19502] R. S.; Powell, S. J.; Smith, J. C.; Markham, A. F.: A yeast artificial
[19503] chromosome contig encompassing the cystic fibrosis locus. Genomics 9:
[19504] 124-130, 1991.

[19505] 2024.

[19506] Antinolo, G.; Borrego, S.; Gili, M.; Dapena, J.; Alfageme, I.;

[19507] Reina, F.: Genotype-phenotype relationship in 12 patients carrying
[19508] cystic fibrosis mutation R334W. J. Med. Genet. 34: 89-91, 1997.
[19509] 2025.

[19510] Audrezet, M. P.; Mercier, B.; Guillermit, H.; Quere, I.; Verlingue,
[19511] C.; Rault, G.; Ferec, C.: Identification of 12 novel muta-

tions in
[19512] the CFTR gene. Hum. Molec. Genet. 2: 51-54, 1993.

[19513] 2026.

[19514] Dorval, I.; Odent, S.; Jezequel, P.; Journel, H.; Chauvel, B.;

[19515] Dabadie, A.; Roussey, M.; Le Gall, J. Y.; Le Marec, B.; David, V.;
[19516] Blayau, M.: Analysis of 160 CF chromosomes: detection of a novel
[19517] mutation in exon 20. Hum. Genet. 91: 254-256, 1993.

[19518] 2027.

[19519] Dork, T.; Macek, M., Jr.; Mekus, F.; Tummler, B.; Tzountzouris,
[19520] J.; Casals, T.; Krebsova, A.; Koudova, M.; Sakmaryova, I.; Macek,
[19521] M., Sr.; Vavrova, V.; Zemkova, D.; and 64 others: Characterization
[19522] of a novel 21-kb deletion, CFTRdele2,3(21 kb), in the CFTR gene: a
[19523] cystic fibrosis mutation of Slavic origin common in Central and East
[19524] Europe. Hum. Genet. 106: 259-268, 2000.

[19525] 2028.

[19526] Ellsworth, R. E.; Jamison, D. C.; Touchman, J. W.; Chissoe, S.

[19527] L.; Maduro, V. V. B.; Bouffard, G. G.; Dietrich, N. L.; Beck-strom-Sternberg,

[19528] S. M.; Iyer, L. M.; Weintraub, L. A.; Cotton, M.; Courtney, L.; and

[19529] 18 others: Comparative genomic sequence analysis of the human and

[19530] mouse cystic fibrosis transmembrane conductance regu-lator genes. Proc.

[19531] Nat. Acad. Sci. 97: 1172-1177, 2000.

[19532] 2029.

[19533] Ferrie, R. M.; Schwarz, M. J.; Robertson, N. H.; Vaudin, S.; Super,

[19534] M.; Malone, G.; Little, S.: Development, multiplexing, and application

[19535] of ARMS tests for common mutations in the CFTR gene. Am. J. Hum.

[19536] Genet. 51: 251-262, 1992.

[19537] 2030.

[19538] Fonknechten, N.; Chelly, J.; Lepercq, J.; Kahn, A.; Kaplan, J.-C.;

[19539] Kitzis, A.; Chomel, J.-C.: CFTR illegitimate transcription in lymphoid

[19540] cells: quantification and applications to the investigation of pathological

[19541] transcripts. Hum. Genet. 88: 508-512, 1992.

[19542] 2031.

[19543] French, P. J.; van Doorninck, J. H.; Peters, R. H. P. C.; Ver-beek,

[19544] E.; Ameen, N. A.; Marino, C. R.; de Jonge, H. R.; Bijman, J.; Scholte,

[19545] B. J.: A delta-F508 mutation in mouse cystic fibrosis transmembrane

[19546] conductance regulator results in a temperature-sensitive processing

[19547] defect in vivo. J. Clin. Invest. 98: 1304-1312, 1996.

[19548] 2032.

[19549] Gasparini, P.; Borgo, G.; Mastella, G.; Bonizzato, A.; Dognini,

[19550] M.; Pignatti, P. F.: Nine cystic fibrosis patients homozy-gous for

[19551] the CFTR nonsense mutation R1162X have mild or moderate lung disease. J.

[19552] Med. Genet. 29: 558-562, 1992.

[19553] 2033.

[19554] Gasparini, P.; Marigo, C.; Bisceglia, G.; Nicolis, E.; Zelante,

[19555] L.; Bombieri, C.; Borgo, G.; Pignatti, P. F.; Cabrini, G.: Screening

[19556] of 62 mutations in the cohort of cystic fibrosis patients from north

[19557] eastern Italy: their incidence and clinical features of defined genotypes. Hum.

[19558] Mutat. 2: 389-394, 1993.

[19559] 2034.

[19560] Gasparini, P.; Nunes, V.; Savoia, A.; Dognini, M.; Morral, N.;

[19561] Gaona, A.; Bonizzato, A.; Chillon, M.; Sangiuolo, F.; Novelli, G.;

[19562] Dallapiccola, B.; Pignatti, P. F.; Estivill, X.: The search for south

[19563] European cystic fibrosis mutations: identification of two new mutations,

[19564] four variants, and intronic sequences. Genomics 10: 193-200, 1991.

[19565] 2035.

[19566] Gervais, R.; Dumur, V.; Rigot, J.-M.; Lafitte, J.-J.; Roussel,

[19567] P.; Claustres, M.; Demaille, J.: High frequency of the R117H cystic

[19568] fibrosis mutation in patients with congenital absence of the vas deferens.

[19569] (Letter) New Eng. J. Med. 328: 328-329, 1993.

[19570] 2036.

[19571] Ghanem, N.; Costes, B.; Girodon, E.; Martin, J.; Fanen, P.; Goossens,

[19572] M.: Identification of eight mutations and three sequence variations

[19573] in the cystic fibrosis transmembrane conductance regulator (CFTR)

[19574] gene. Genomics 21: 434-436, 1994.

[19575] 2037.

[19576] Gille, C.; Grade, K.; Coutelle, C.: A pooling strategy for heterozygote

[19577] screening of the delta-F508 cystic fibrosis mutation. Hum.

Genet. 86:
[19578] 289-291, 1991.

[19579] 2038.

[19580] Girodon, E.; Cazeneuve, C.; Lebargy, F.; Chinet, T.; Costes, B.;
[19581] Ghanem, N.; Martin, J.; Lemay, S.; Scheid, P.; Housset, B.; Bignon,
[19582] J.; Goossens, M.: CFTR gene mutations in adults with dis-seminated
[19583] bronchiectasis. Europ. J. Hum. Genet. 5: 149-155, 1997.

[19584] 2039.

[19585] Gomez Lira, M.; Benetazzo, M. G.; Marzari, M. G.; Bombieri, C.;
[19586] Belpinati, F.; Castellani, C.; Cavallini, G. C.; Mastella, G.; Pignatti,
[19587] P. F.: High frequency of cystic fibrosis transmembrane regulator
[19588] mutation L997F in patients with recurrent idiopathic pan-creatitis
[19589] and in newborns with hypertrypsinemia. (Letter) Am. J. Hum. Genet. 66:
[19590] 2013-2014, 2000.

[19591] 2040.

[19592] Graham, C. A.; Goon, P. K.-C.; Hill, A. J. M.; Nevin, N. C.:

[19593] Identification of a frameshift mutation (557 del T) in exon 4 of the

[19594] CFTR gene. Genomics 12: 854-855, 1992.

[19595] 2041.

[19596] Granell, R.; Solera, J.; Carrasco, S.; Molano, J.: Identification

[19597] of a nonframeshift 84-bp deletion in exon 13 of the cystic fibrosis

[19598] gene. Am. J. Hum. Genet. 50: 1022-1026, 1992.

[19599] 2042.

[19600] Grebe, T. A.; Doane, W. W.; Richter, S. F.; Clericuzio, C.; Norman,

[19601] R. A.; Seltzer, W. K.; Rhodes, S. N.; Goldberg, B. E.; Hernried, L.

[19602] S.; McClure, M.; Kaplan, G.: Mutation analysis of the cystic fibrosis

[19603] transmembrane regulator gene in Native American populations of the

[19604] Southwest. Am. J. Hum. Genet. 51: 736-740, 1992.

[19605] 2043.

[19606] Grebe, T. A.; Seltzer, W. K.; DeMarchi, J.; Silva, D. K.; Doane,

[19607] W. W.; Gozal, D.; Richter, S. F.; Bowman, C. M.; Norman, R. A.; Rhodes,

[19608] S. N.; Hernried, L. S.; Murphy, S.; Harwood, I. R.; Accurso, F. J.;

[19609] Jain, K. D.: Genetic analysis of Hispanic individuals with cystic

[19610] fibrosis. Am. J. Hum. Genet. 54: 443-446, 1994.

[19611] 2044.

[19612] Cavailles, V.; Dauvois, S.; Horset, L. F.; Lopez, G.; Hoare, S.;

[19613] Kushner, P. J.; Parker, M. G.: Nuclear factor RIP140 modulates transcriptional

[19614] activation by the estrogen receptor. EMBO J. 14: 3741-3751, 1995.

[19615] 2045.

[19616] Katsanis, N.; Ives, J. H.; Groet, J.; Nizetic, D.; Fisher, E. M.

[19617] C.: Localisation of receptor interacting protein 140 (RIP140) within

[19618] 100 kb of D21S13 on 21q11, a gene-poor region of the human genome. Hum.

[19619] Genet. 102: 221-223, 1998.

[19620] 2046.

[19621] Brown, J. L.; Stowers, L.; Baer, M.; Trejo, J.; Coughlin, S.; Chant,

[19622] J.: Human Ste20 homologue hPAK1 links GTPases to the JNK MAP kinase

[19623] pathway. Curr. Biol. 6: 598-605, 1996.

[19624] 2047.

[19625] Knaus, U. G.; Morris, S.; Dong, H.-J.; Chernoff, J.; Bokoch, G.

[19626] M.: Regulation of human leukocyte p21-activated kinases through G

[19627] protein-coupled receptors. Science 269: 221-223, 1995.

[19628] 2048.

[19629] Lei, M.; Lu, W.; Meng, W.; Parrini, M.-C.; Eck, M. J.; Mayer, B.

[19630] J.; Harrison, S. C.: Structure of PAK1 in an autoinhibited conformation

[19631] reveals a multistage activation switch. Cell 102: 387-397,

2000.

[19632] 2049.

[19633] Parrini, M. C.; Lei, M.; Harrison, S. C.; Mayer, B. J.: Pak1 ki-
nase

[19634] homodimers are autoinhibited in trans and dissociated
upon activation

[19635] by Cdc42 and Rac1. Molec. Cell 9: 73-83, 2002.

[19636] 2050.

[19637] Sanders, L. C.; Matsumura, F.; Bokoch, G. M.; de Lanerolle,
P.

[19638] : Inhibition of myosin light chain kinase by p21-activated
kinase. Science 283:

[19639] 2083-2085, 1999.

[19640] 2051.

[19641] Fischer, U.; Liu, Q.; Dreyfuss, G.: The SMN-SIP1 complex
has an

[19642] essential role in spliceosomal snRNP biogenesis. Cell 90:
1023-1029,

[19643] 1997.

[19644] 2052.

[19645] Hannus, S.; Buhler, D.; Romano, M.; Seraphin, B.; Fischer,

U.:

[19646] The Schizosaccharomyces pombe protein Yab8p and a novel factor, Yip1p,

[19647] share structural and functional similarity with the spinal muscular

[19648] atrophy-associated proteins SMN and SIP1. Hum. Molec. Genet. 9:

[19649] 663-674, 2000.

[19650] 2053.

[19651] Helmken, C.; Wetter, A.; Rudnik-Schoneborn, S.; Liehr, T.; Zerres,

[19652] K.; Wirth, B.: An essential SMN interacting protein (SIP1) is not

[19653] involved in the phenotypic variability of spinal muscular atrophy

[19654] (SMA). Europ. J. Hum. Genet. 8: 493-499, 2000.

[19655] 2054.

[19656] Jablonka, S.; Bandilla, M.; Wiese, S.; Buhler, D.; Wirth, B.; Sendtner,

[19657] M.; Fischer, U.: Co-regulation of survival of motor neuron (SMN)

[19658] protein and its interactor SIP1 during development and in

spinal muscular
[19659] atrophy. Hum. Molec. Genet. 10: 497-505, 2001.

[19660] 2055.

[19661] Young, P. J.; Man, N.; Lorson, C. L.; Le, T. T.; Androphy, E. J.;
[19662] Burghes, A. H. M.; Morris, G. E.: The exon 2b region of the spinal
[19663] muscular atrophy protein, SMN, is involved in self-association and
[19664] SIP1 binding. Hum. Molec. Genet. 9: 2869-2877, 2000.

[19665] 2056.

[19666] Arany, Z.; Sellers, W. R.; Livingston, D. M.; Eckner, R.: E1A-associated
[19667] p300 and CREB-associated CBP belong to a conserved family of coactivators.
[19668] (Letter) Cell 77: 799-800, 1994.

[19669] 2057.

[19670] Gayther, S. A.; Batley, S. J.; Linger, L.; Bannister, A.; Thorpe,
[19671] K.; Chin, S.-F.; Daigo, Y.; Russell, P.; Wilson, A.; Sowter, H. M.;

[19672] Delhanty, J. D. A.; Ponder, B. A. J.; Kouzarides, T.; Caldas, C.:

[19673] Mutations truncating the EP300 acetylase in human cancers. Nature

[19674] Genet. 24: 300-303, 2000.

[19675] 2058.

[19676] Ida, K.; Kitabayashi, I.; Taki, T.; Taniwaki, M.; Noro, K.; Yamamoto,

[19677] M.; Ohki, M.; Hayashi, Y.: Adenoviral E1A-associated protein p300

[19678] is involved in acute myeloid leukemia with t(11;22)(q23;q13). Blood 90:

[19679] 4699-4704, 1997.

[19680] 2059.

[19681] Muraoka, M.; Konishi, M.; Kikuchi-Yanoshita, R.; Tanaka, K.; Shitara,

[19682] N.; Chong, J.-M.; Iwama, T.; Miyaki, M.: p300 gene alterations in

[19683] colorectal and gastric carcinomas. Oncogene 12: 1565-1569, 1996.

[19684] 2060.

[19685] Yao, T. P.; Oh, S. P.; Fuchs, M.; Zhou, N.-D.; Ch'ng, L.-E.; Newsome,
[19686] D.; Bronson, R. T.; Li, E.; Livingston, D. M.; Eckner, R.: Gene dosage-dependent
[19687] embryonic development and proliferation defects in mice lacking the
[19688] transcriptional integrator p300. Cell 93: 361-372, 1998.

[19689] 2061.

[19690] Parant, J.; Chavez-Reyes, A.; Little, N. A.; Yan, W.; Reinke, V.;
[19691] Jochemsen, A. G.; Lozano, G.: Rescue of embryonic lethality in Mdm4-null
[19692] mice by loss of Trp53 suggests a nonoverlapping pathway with MDM2
[19693] to regulate p53. Nature Genet. 29: 92-95, 2001.

[19694] 2062.

[19695] Shvarts, A.; Bazuine, M.; Dekker, P.; Ramos, Y. F. M.; Steegenga,
[19696] W. T.; Merckx, G.; van Ham, R. C. A.; van der Houven van Oordt, W.;
[19697] van der Eb, A. J.; Jochemsen, A. G.: Isolation and identification

[19698] of the human homolog of a new p53-binding protein, Mdmx. Genomics 43:

[19699] 34-42, 1997.

[19700] 2063.

[19701] Hosaka, M.; Sudhof, T. C.: Synapsin III, a novel synapsin with

[19702] an unusual regulation by Ca(2)+. J. Biol. Chem. 273: 13371-13374,

[19703] 1998.

[19704] 2064.

[19705] Kao, H.-T.; Porton, B.; Czernik, A. J.; Feng, J.; Yiu, G.; Har-

ing,

[19706] M.; Benfenati, F.; Greengard, P.: A third member of the synapsin

[19707] gene family. Proc. Nat. Acad. Sci. 95: 4667-4672, 1998.

[19708] 2065.

[19709] Schizophrenia Collaborative Linkage Group (Chromosome 22): A

[19710] combined analysis of D22S278 marker alleles in affected sib-pairs:

[19711] support for a susceptibility locus for schizophrenia at

chromosome
[19712] 22q12. Am. J. Med. Genet. 67: 40-45, 1996.

[19713] 2066.

[19714] Tsai, M.-T.; Hung, C.-C.; Tsai, C.-Y.; Liu, M.-Y.; Su, Y.-C.; Chen,
[19715] Y.-H.; Hsaio, K.-J.; Chen, C.-H.: Mutation analysis of synapsin III
[19716] gene in schizophrenia. Am. J. Med. Genet. (Neuropsychiat. Genet.) 114:
[19717] 79-83, 2002.

[19718] 2067.

[19719] Kull, F. J.; Sablin, E. P.; Lau, R.; Fletterick, R. J.; Vale, R.
[19720] D.: Crystal structure of the kinesin motor domain reveals a structural
[19721] similarity to myosin. Nature 380: 550-555, 1996.

[19722] 2068.

[19723] Navone, F.; Niclas, J.; Hom-Booher, N.; Sparks, L.; Bern-stein,
[19724] H. D.; McCaffrey, G.; Vale, R. D.: Cloning and expression of a human
[19725] kinesin heavy chain gene: interaction of the COOH-ter-

minal domain
[19726] with cytoplasmic microtubules in transfected CV-1 cells. J. Cell
[19727] Biol. 117: 1263-1275, 1992.

[19728] 2069.

[19729] Niclas, J.; Navone, F.; Hom-Booher, N.; Vale, R. D.: Cloning and
[19730] localization of a conventional kinesin motor expressed exclusively
[19731] in neurons. Neuron 12: 1059-1072, 1994.

[19732] 2070.

[19733] Tanaka, Y.; Kanai, Y.; Okada, Y.; Nonaka, S.; Takeda, S.; Harada,
[19734] A.; Hirokawa, N.: Targeted disruption of mouse conven-
tional kinesin
[19735] heavy chain, kif5B, results in abnormal perinuclear clus-
tering of
[19736] mitochondria. Cell 93: 1147-1158, 1998.

[19737] 2071.

[19738] Wang, H.; Huang, Z.-Q.; Xia, L.; Feng, Q.; Erdjument-Bro-
mage, H.;

[19739] Strahl, B. D.; Briggs, S. D.; Allis, C. D.; Wong, J.; Tempst, P.;

[19740] Zhang, Y.: Methylation of histone H4 at arginine 3 facili-tating transcriptional

[19741] activation by nuclear hormone receptor. Science 293: 853-857, 2001.

[19742] 2072.

[19743] Albala, J. S.; Thelen, M. P.; Prange, C.; Fan, W.; Chris-tensen,

[19744] M.; Thompson, L. H.; Lennon, G. G.: Identification of a novel human

[19745] RAD51 homolog, RAD51B. Genomics 46: 476-479, 1997.

[19746] 2073.

[19747] Cartwright, R.; Dunn, A. M.; Simpson, P. J.; Tambini, C. E.; Thacker,

[19748] J.: Isolation of novel human and mouse genes of the recA/RAD51 recombination-repair

[19749] gene family. Nucleic Acids Res. 26: 1653-1659, 1998.

[19750] 2074.

[19751] Rice, M. C.; Smith, S. T.; Bullrich, F.; Havre, P.; Kmiec, E. B.

[19752] : Isolation of human and mouse genes based on homology

to REC2, a
[19753] recombinational repair gene from the fungus Ustilago maydis. Proc.
[19754] Nat. Acad. Sci. 94: 7417-7422, 1997.

[19755] 2075.

[19756] Abramovich, C.; Yakobson, B.; Chebath, J.; Revel, M.: A protein-arginine
[19757] methyltransferase binds to the intracytoplasmic domain of the IFNAR1
[19758] chain in the type I interferon receptor. EMBO J. 16: 260-266, 1997.

[19759] 2076.

[19760] Lin, W.-J.; Gary, J. D.; Yang, M. C.; Clarke, S.; Herschman, H.
[19761] R.: The mammalian immediate-early TIS21 protein and the leukemia-associated
[19762] BTG1 protein interact with a protein-arginine N-methyltransferase. J.
[19763] Biol. Chem. 271: 15034-15044, 1996.

[19764] 2077.

[19765] Mowen, K. A.; Tang, J.; Zhu, W.; Schurter, B. T.; Shuai, K.;

Herschman,
[19766] H. R.; David, M.: Arginine methylation of STAT1 modulates IFN-alpha/beta-induce
[19767] d transcription. Cell 104: 731-741, 2001.

[19768] 2078.

[19769] Nikawa, J.; Nakano, H.; Ohi, N.: Structural and functional conservation
[19770] of human and yeast HCP1 genes which can suppress the growth defect
[19771] of the Saccharomyces cerevisiae ire15 mutant. Gene 171: 107-111,
[19772] 1996.

[19773] 2079.

[19774] Pawlak, M. R.; Scherer, C. A.; Chen, J.; Roshon, M. J.; Ruley,
[19775] H. E.: Arginine N-methyltransferase 1 is required for early postimplantation
[19776] mouse development, but cells deficient in the enzyme are viable. Molec.
[19777] Cell. Biol. 20: 4859-4869, 2000.

[19778] 2080.

[19779] Scorilas, A.; Black, M. H.; Talieri, M.; Diamandis, E. P.: Ge-

nomic

[19780] organization, physical mapping, and expression analysis of the human

[19781] protein arginine methyltransferase 1 gene. Biochem. Biophys. Res.

[19782] Commun. 278: 349-359, 2000.

[19783] 2081.

[19784] Baxendale, S.; MacDonald, M. E.; Mott, R.; Francis, F.; Lin, C.;

[19785] Kirby, S. F.; James, M.; Zehetner, G.; Hummerich, H.; Valdes, J.;

[19786] Collins, F. S.; Deaven, L. J.; Gusella, J. F.; Lehrach, H.; Bates,

[19787] G. P.: A cosmid contig and high resolution restriction map of the

[19788] 2 megabase region containing the Huntington's disease gene. Nature

[19789] Genet. 4: 181-186, 1993.

[19790] 2082.

[19791] Richelda, R.; Ronchetti, D.; Baldini, L.; Cro, L.; Viggiano, L.;

[19792] Marzella, R.; Rocchi, M.; Otsuki, T.; Lombardi, L.; Maiolo,

A. T.;

[19793] Neri, A.: A novel chromosomal translocation t(4;14)(p16.3;q32) in

[19794] multiple myeloma involves the fibroblast growth-factor receptor 3

[19795] gene. Blood 90: 4062-4070, 1997.

[19796] 2083.

[19797] Wright, T. J.; Ricke, D. O.; Denison, K.; Abmayr, S.; Cotter, P.

[19798] D.; Hirschhorn, K.; Keinanen, M.; McDonald-McGinn, D.; Somer, M.;

[19799] Spinner, N.; Yang-Feng, T.; Zackai, E.; Altherr, M. R.: A transcript

[19800] map of the newly defined 165 kb Wolf-Hirschhorn syndrome critical

[19801] region. Hum. Molec. Genet. 6: 317-324, 1997.

[19802] 2084.

[19803] Chen, H.; Rossier, C.; Lalioti, M. D.; Lynn, A.; Chakravarti, A.;

[19804] Perrin, G.; Antonarakis, S. E.: Cloning of the cDNA for a human homologue

[19805] of the Drosophila white gene and mapping to chromo-

some 21q22.3. Am.

[19806] J. Hum. Genet. 59: 66-75, 1996.

[19807] 2085.

[19808] Croop, J. M.; Tiller, G. E.; Fletcher, J. A.; Lux, M. L.; Raab,

[19809] E.; Goldenson, D.; Son, D.; Arciniegas, S.; Wu, R. L.: Isola-
tion

[19810] and characterization of a mammalian homolog of the
Drosophila white

[19811] gene. Gene 185: 77-85, 1997.

[19812] 2086.

[19813] Klucken, J.; Buchler, C.; Orso, E.; Kaminski, W. E.; Porsch-
Ozcurumez,

[19814] M.; Liebisch, G.; Kapinsky, M.; Diederich, W.; Drobnik, W.;
Dean,

[19815] M.; Allikmets, R.; Schmitz, G.: ABCG1 (ABC8), the human
homolog of

[19816] the Drosophila white gene, is a regulator of macrophage
cholesterol

[19817] and phospholipid transport. Proc. Nat. Acad. Sci. 97:
817-822, 2000.

[19818] 2087.

[19819] Langmann, T.; Porsch-Ozcurumez, M.; Unkelbach, U.; Klucken, J.;

[19820] Schmitz, G.: Genomic organization and characterization of the promoter

[19821] of the human ATP-binding cassette transporter-G1 (ABCG1) gene. Biochim.

[19822] Biophys. Acta 1494: 175-180, 2000.

[19823] 2088.

[19824] Lorkowski, S.; Rust, S.; Engel, T.; Jung, E.; Tegelkamp, K.; Galinski,

[19825] E. A.; Assmann, G.; Cullen, P.: Genomic sequence and structure of

[19826] the human ABCG1 (ABC8) gene. Biochem. Biophys. Res. Commun. 280:

[19827] 121-131, 2001.

[19828] 2089.

[19829] Savary, S.; Denizot, F.; Luciani, M.-F.; Mattei, M.-G.; Chi-mini,

[19830] G.: Molecular cloning of a mammalian ABC transporter homologous to

[19831] Drosophila white gene. Mammalian Genome 7: 673-676, 1996.

[19832] 2090.

[19833] Borden, L. A.; Smith, K. E.; Gustafson, E. L.; Branchek, T. A.;

[19834] Weinshank, R. L.: Cloning and expression of a betaine/ GABA transporter

[19835] from human brain. J. Neurochem. 64: 977-984, 1995.

[19836] 2091.

[19837] Rasola, A.; Galietta, L. J. V.; Barone, V.; Romeo, G.; Bag-nasco,

[19838] S.: Molecular cloning and functional characterization of a GABA/betaine

[19839] transporter from human kidney. FEBS Lett. 373: 229-233, 1995.

[19840] 2092.

[19841] Yamauchi, A.; Uchida, S.; Kwon, H. M.; Preston, A. S.; Robey, R.

[19842] B.; Garcia-Perez, A.; Burg, M. B.; Handler, J. S.: Cloning of a Na(+)

[19843] and Cl(-)-dependent betaine transporter that is regulated by hypertonicity. J.

[19844] Biol. Chem. 267: 649-652, 1992.

[19845] 2093.

[19846] Schwienbacher, C.; Sabbioni, S.; Campi, M.; Veronese, A.; Bernardi,

[19847] G.; Menegatti, A.; Hatada, I.; Mukai, T.; Ohashi, H.; Barbanti-Brodano,

[19848] G; Croce, C. M.; Negrini, M.: Transcriptional map of 170-kb region

[19849] at chromosome 11p15.5: identification and mutational analysis of the

[19850] BWR1A gene reveals the presence of mutations in tumor samples. Proc.

[19851] Nat. Acad. Sci. 95: 3873-3878, 1998.

[19852] 2094.

[19853] Hirono, Y.; Fushida, S.; Yonemura, Y.; Yamamoto, H.; Watanabe,

[19854] H.; Raz, A.: Expression of autocrine motility factor receptor correlates

[19855] with disease progression in human gastric cancer. Brit. J. Cancer 74:

[19856] 2003-2007, 1996.

[19857] 2095.

[19858] Huang, B.; Xie, Y.; Raz, A.: Identification of an upstream region
[19859] that controls the transcription of the human autocrine motility factor
[19860] receptor. Biochem. Biophys. Res. Commun. 212: 727-742, 1995.
[19861] 2096.

[19862] Silletti, S.; Yao, J.; Sanford, J.; Mohammed, A. N.; Otto, T.;

[19863] Wolman, S. R.; Raz, A.: Autocrine motility factor receptor in human
[19864] bladder carcinoma: gene expression, loss of cell-contact regulation
[19865] and chromosomal mapping. Int. J. Oncol. 3: 801-807, 1993.
[19866] 2097.

[19867] Watanabe, H.; Carmi, P.; Hogan, V.; Raz, T.; Silletti, S.; Nabi,
[19868] I. R.; Raz, A.: Purification of human tumor cell autocrine motility
[19869] factor and molecular cloning of its receptor. J. Biol. Chem. 266:
[19870] 13442-13448, 1991.

[19871] 2098.

[19872] Amlal, H.; Burnham, C. E.; Soleimani, M.: Characterization of

[19873] Na(+)/HCO(3-) cotransporter isoform NBC-3. Am. J. Physiol. 276:

[19874] F903-F913, 1999.

[19875] 2099.

[19876] Burnham, C. E.; Wang, Z.; Soleimani, M.: Personal Communication. Cincinnati,

[19877] Oh. 6/1/2000.

[19878] 2100.

[19879] Choi, I.; Aalkjaer, C.; Boulpaep, E. L.; Boron, W. F.: An electroneutral

[19880] sodium/bicarbonate cotransporter NBCn1 and associated sodium channel. Nature 405:

[19881] 571-575, 2000.

[19882] 2101.

[19883] Ishibashi, K.; Sasaki, S.; Marumo, F.: Molecular cloning of a

[19884] new sodium bicarbonate cotransporter cDNA from human retina. Biochem.

[19885] Biophys. Res. Commun. 246: 535-538, 1998.

[19886] 2102.

[19887] Pushkin, A.; Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Kurtz,

[19888] I.: Mapping of the human NBC3 (SLC4A7) gene to chromosome 3p22. Genomics 57:

[19889] 321-322, 1999. Note: Correction: Genomics 58: 216 and 321-322, 1999.

[19890] 2103.

[19891] Pushkin, A.; Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Kurtz,

[19892] I.: Cloning, tissue distribution, genomic organization, and functional

[19893] characterization of NBC3, a new member of the sodium bicarbonate cotransporter

[19894] family. J. Biol. Chem. 274: 16569-16575, 1999.

[19895] 2104.

[19896] Brown, M. S.; Goldstein, J. L.: The SREBP pathway: regulation

[19897] of cholesterol metabolism by proteolysis of a membrane-bound transcription

[19898] factor. Cell 89: 331-340, 1997.

[19899] 2105.

[19900] Duncan, E. A.; Brown, M. S.; Goldstein, J. L.; Sakai, J.: Cleavage

[19901] site for sterol-regulated protease localized to a leu-ser bond in

[19902] the lumenal loop of sterol regulatory element-binding protein-2. J.

[19903] Biol. Chem. 272: 12778-12785, 1997.

[19904] 2106.

[19905] Nagase, T.; Miyajima, N.; Tanaka, A.; Sazuka, T.; Seki, N.; Sato,

[19906] S.; Tabata, S.; Ishikawa, K.; Kawarabayasi, Y.; Kotani, H.; Nomura,

[19907] N.: Prediction of the coding sequences of unidentified hu-man genes.

[19908] III. The coding sequences of 40 new genes (KIAA0081-KIAA0120) deduced

[19909] by analysis of cDNA clones from human cell line KG-1. DNA Res. 2:

[19910] 37-43, 1995.

[19911] 2107.

[19912] Nakajima, T.; Iwaki, K.; Kodama, T.; Inazawa, J.; Emi, M.: Genomic

[19913] structure and chromosomal mapping of the human site-1 protease (S1P)

[19914] gene. J. Hum. Genet. 45: 212-217, 2000.

[19915] 2108.

[19916] Sakai, J.; Duncan, E. A.; Rawson, R. B.; Hua, X.; Brown, M. S.;

[19917] Goldstein, J. L.: Sterol-regulated release of SREBP-2 from cell membranes

[19918] requires two sequential cleavages, one within a trans-membrane segment. Cell 85:

[19919] 1037-1046, 1996.

[19920] 2109.

[19921] Sakai, J.; Rawson, R. B.; Espenshade, P. J.; Cheng, D.; Seegmiller,

[19922] A. C.; Goldstein, J. L.; Brown, M. S.: Molecular identification of

[19923] the sterol-regulated luminal protease that cleaves SREBPs and controls

[19924] lipid composition of animal cells. Molec. Cell 2: 505-514, 1998.

[19925] 2110.

[19926] Watt, S. M.; Buhring, H.-J.; Rappold, I.; Chan, J. Y.-H.; Lee-Prudhoe,

[19927] J.; Jones, T.; Zannettino, A. C. W.; Simmons, P. J.; Doyon-nas, R.;

[19928] Sheer, D.; Butler, L. H.: CD164, a novel sialomucin on CD34+ and

[19929] erythroid subsets, is located on human chromosome 6q21. Blood 92:

[19930] 849-866, 1998.

[19931] 2111.

[19932] Zannettino, A. C. W.; Buhring, H.-J.; Niutta, S.; Watt, S. M.;

[19933] Benton, M. A.; Simmons, P. J.: The sialomucin CD164 (MGC-24v) is

[19934] an adhesive glycoprotein expressed by human hematopoietic progenitors

[19935] and bone marrow stromal cells that serves as a potent negative regulator

[19936] of hematopoiesis. Blood 92: 2613-2628, 1998.

[19937] 2112.

[19938] Yagasaki, F.; Jinnai, I.; Yoshida, S.; Yokoyama, Y.; Matsuda,16.

[19939] Yagasaki, F.; Jinnai, I.; Yoshida, S.; Yokoyama, Y.; Matsuda, A.;

[19940] Kusumoto, S.; Kobayashi, H.; Terasaki, H.; Ohyashiki, K.; Asou, N.;

[19941] Murohashi, I.; Bessho, M.; Hirashima, K.: Fusion of TEL/ETV6 to a

[19942] novel ACS2 in myelodysplastic syndrome and acute myelogenous leukemia

[19943] with t(5;12)(q31;p13). Genes Chromosomes Cancer 26: 192-202, 1999.

[19944] 2113.

[19945] Abts, H. F.; Breuhahn, K.; Michel, G.; Kohrer, K.; Esser, P.; Ruzicka,

[19946] T.: Analysis of UVB modulated gene expression in human keratinocytes

[19947] by mRNA differential display polymerase chain reaction. Photochem.

[19948] Photobiol. 66: 363-367, 1997.

[19949] 2114.

[19950] Abts, H. F.; Welss, T.; Mirmohammadsadegh, A.; Kohrer, K.; Michel,

[19951] G.; Ruzicka, T.: Cloning and characterization of hurpin (protease

[19952] inhibitor 13): a new skin-specific, UV-repressible serine proteinase

[19953] inhibitor of the ovalbumin serpin family. J. Molec. Biol. 293: 29-39,

[19954] 1999.

[19955] 2115.

[19956] Nakashima, T.; Pak, S. C.; Silverman, G. A.; Spring, P. M.; Frederick,

[19957] M. J.; Clayman, G. L.: Genomic cloning, mapping, structure and promoter

[19958] analysis of HEADPIN, a serpin which is down-regulated in head and

[19959] neck cancer cells. Biochim. Biophys. Acta 1492: 441-446, 2000.

[19960] 2116.

[19961] Spring, P.; Nakashima, T.; Frederick, M.; Henderson, Y.; Clayman,

[19962] G.: Identification and cDNA cloning of headpin, a novel

differentially
[19963] expressed serpin that maps to chromosome 18q.
Biochem. Biophys. Res.
[19964] Commun. 264: 299-304, 1999.

[19965] 2117.

[19966] Alimova-Kost, M. V.; Imreh, S.; Buchman, V. L.; Ninkina, N.
N.
[19967] : Assignment of phosphotriesterase-related gene (PTER)
to human chromosome
[19968] band 10p12 by in situ hybridization. Cytogenet. Cell
Genet. 83:
[19969] 16-17, 1998.

[19970] 2118.

[19971] Davies, J. A.; Buchman, V. L.; Krylova, O.; Ninkina, N. N.:
Molecular
[19972] cloning and expression pattern of rpr-1, a resinifera-
toxin-binding,
[19973] phosphotriesterase-related protein, expressed in rat kid-
ney tubules. FEBS
[19974] Lett. 410: 378-382, 1997.

[19975] 2119.

[19976] Jones, P. G.; Lombardi, S. J.; Cockett, M. I.: Cloning and tissue
[19977] distribution of the human G protein beta-5 cDNA. Biochim. Biophys.
[19978] Acta 1402: 288-291, 1998.

[19979] 2120.

[19980] Watson, A. J.; Aragay, A. M.; Slepak, V. Z.; Simon, M. I.: A novel
[19981] form of the G protein beta subunit G-beta-5 is specifically expressed
[19982] in the vertebrate retina. J. Biol. Chem. 271: 28154-28160, 1996.
[19983] 2121.

[19984] Watson, A. J.; Katz, A.; Simon, M. I.: A fifth member of the mammalian
[19985] G-protein beta subunit family: expression in brain and ac-tivation
[19986] of the beta-2 isotype of phospholipase C. J. Biol. Chem. 269: 22150-22156,
[19987] 1994.

[19988] 2122.

[19989] Dixon, B.; Sahely, B.; Liu, L.; Pohajdak, B.: Cloning a cDNA from
[19990] human NK/T cells which codes for an unusual leucine zip-per containing
[19991] protein. Biochim. Biophys. Acta 1216: 321-324, 1993.

[19992] 2123.

[19993] Kim, H.-S.: Assignment of the human B3-1 gene (PSCDBP) to chromosome
[19994] 2 band q11.2 by radiation hybrid mapping. Cytogenet. Cell Genet. 84:
[19995] 95 only, 1999.

[19996] 2124.

[19997] Tang, P.; Cheng, T. P.; Agnello, D.; Wu, C.-Y.; Hissong, B. D.;
[19998] Watford, W. T.; Ahn, H.-J.; Galon, J.; Moss, J.; Vaughan, M.; O'Shea,
[19999] J. J.; Gadina, M.: Cybr, a cytokine-inducible protein that binds
[20000] cytohesin-1 and regulates its activity. Proc. Nat. Acad. Sci. 99:
[20001] 2625-2629, 2002.

[20002] 2125.

[20003] Nakayama, Y.; Goebl, M.; O'Brine Greco, B.; Lemmon, S.; Pingchang

[20004] Chow, E.; Kirchhausen, T.: The medium chains of the mammalian clathrin-associated

[20005] proteins have a homolog in yeast. Europ. J. Biochem. 202: 569-574,

[20006] 1991.

[20007] 2126.

[20008] Hiramoto, T.; Nakanishi, T.; Sumiyoshi, T.; Fukuda, T.; Matsuura,

[20009] S.; Tauchi, H.; Komatsu, K.; Shibasaki, Y.; Inui, H.; Watatani, M.;

[20010] Yasutomi, M.; Sumii, K.; Kajiyama, G.; Kamada, N.; Miya-gawa, K.; Kamiya,

[20011] K.: Mutations of a novel human RAD54 homologue, RAD54B, in primary

[20012] cancer. Oncogene 18: 3422-3426, 1999.

[20013] 2127.

[20014] Miyagawa, K.; Tsuruga, T.; Kinomura, A.; Usui, K.; Katsura, M.;

[20015] Tashiro, S.; Mishima, H.; Tanaka, K.: A role for RAD54B in homologous

[20016] recombination in human cells. EMBO J. 21: 175-180, 2002.

[20017] 2128.

[20018] Tanaka, K.; Hiramoto, T.; Fukuda, T.; Miyagawa, K.: A novel human

[20019] Rad54 homologue, Rad54B, associates with Rad51. J. Biol. Chem. 275:

[20020] 26316-26321, 2000.

[20021] 2129.

[20022] Pillutla, R. C.; Shimamoto, A.; Furuichi, Y.; Shatkin, A. J.:

[20023] Human mRNA capping enzyme (RNGTT) and cap methyl-transferase (RNMT)

[20024] map to 6q16 and 18p11.22-p11.23, respectively. Ge-nomics 54: 351-353,

[20025] 1998.

[20026] 2130.

[20027] Tsukamoto, T.; Shibagaki, Y.; Murakoshi, T.; Suzuki, M.; Nakamura,

[20028] A.; Gotoh, H.; Mizumoto, K.: Cloning and characterization

of two
[20029] human cDNAs encoding the mRNA capping enzyme. Biochem. Biophys. Res.
[20030] Commun. 243: 101-108, 1998.

[20031] 2131.

[20032] Yamada-Okabe, T.; Doi, R.; Shimmi, O.; Arisawa, M.; Ya-mada-Okabe,
[20033] H.: Isolation and characterization of a human cDNA for mRNA 5-prime-capping
[20034] enzyme. Nucleic Acids Res. 26: 1700-1706, 1998.

[20035] 2132.

[20036] Yue, Z.; Maldonado, E.; Pillutla, R.; Cho, H.; Reinberg, D.; Shatkin,
[20037] A. J.: Mammalian capping enzyme complements mutant Saccharomyces
[20038] cerevisiae lacking mRNA guanylyltransferase and selec-tively binds
[20039] the elongating form of RNA polymerase II. Proc. Nat. Acad. Sci. 94:
[20040] 12898-12903, 1997.

[20041] 2133.

[20042] Ishikawa, K.; Nagase, T.; Nakajima, D.; Seki, N.; Ohira, M.; Miyajima,
[20043] N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of
[20044] the coding sequences of unidentified human genes. VIII. 78 new cDNA
[20045] clones from brain which code for large proteins in vitro. DNA Res. 4:
[20046] 307-313, 1997.

[20047] 2134.

[20048] Pillutla, R. C.; Yue, Z.; Maldonado, E.; Shatkin, A. J.: Recombinant
[20049] human mRNA cap methyltransferase binds capping enzyme/RNA polymerase
[20050] IIo complexes. J. Biol. Chem. 273: 21443-21446, 1998.

[20051] 2135.

[20052] Tsukamoto, T.; Shibagaki, Y.; Niikura, Y.; Mizumoto, K.: Cloning
[20053] and characterization of three human cDNAs encoding mRNA (guanine-7)-methyltransferase,
[20054] an mRNA cap methylase. Biochem. Biophys. Res. Commun. 251: 27-34,

[20055] 1998.

[20056] 2136.

[20057] Misra-Press, A.; Cooke, N. E.; Liebhaber, S. A.: Complex alternative

[20058] splicing partially inactivates the human chorionic somatomammotropin-like

[20059] (hCS-L) gene. J. Biol. Chem. 269: 23220-23229, 1994.

[20060] 2137.

[20061] Sawamura, T.; Kume, N.; Aoyama, T.; Moriwaki, H.; Hoshikawa, H.;

[20062] Aiba, Y.; Tanaka, T.; Miwa, S.; Katsura, Y.; Kita, T.; Masaki, T.

[20063] : An endothelial receptor for oxidized low-density lipoprotein. Nature 386:

[20064] 73-77, 1997.

[20065] 2138.

[20066] Yamanaka, S.; Zhang, X.-Y.; Miura, K.; Kim, S.; Iwao, H.: The

[20067] human gene encoding the lectin-type oxidized LDL receptor (OLR1) is

[20068] a novel member of the natural killer gene complex with a

unique expression
[20069] profile. Genomics 54: 191-199, 1998.

[20070] 2139.

[20071] Zhao, X.-F.; Colaizzo-Anas, T.; Nowak, N. J.; Shows, T. B.; Elliott,
[20072] R. W.; Aplan, P. D.: The mammalian homologue of mago nashi encodes
[20073] a serum-inducible protein. Genomics 47: 319-322, 1998.

[20074] 2140.

[20075] Gorboulev, V.; Ulzheimer, J. C.; Akhoundova, A.; Ulzheimer-Teuber,
[20076] I.; Karbach, U.; Quester, S.; Baumann, C.; Lang, F.; Busch, A. E.;
[20077] Koepsell, H.: Cloning and characterization of two human polyspecific
[20078] organic cation transporters. DNA Cell Biol. 16: 871-881, 1997.
[20079] 2141.

[20080] Grundemann, D.; Gorboulev, V.; Gambaryan, S.; Veyhl, M.; Koepsell,
[20081] H.: Drug excretion mediated by a new prototype of

polyspecific transporter. Nature 372:
[20082] 549-552, 1994.

[20083] 2142.

[20084] Hayer, M.; Bonisch, H.; Bruss, M.: Molecular cloning, functional
[20085] characterization and genomic organization of four alternatively spliced
[20086] isoforms of the human organic cation transporter 1 (hOCT1/SLC22A1). Ann.
[20087] Hum. Genet. 63: 473-482, 1999.

[20088] 2143.

[20089] Koehler, M. R.; Wissinger, B.; Gorboulev, V.; Koepsell, H.; Schmid,
[20090] M.: The two human organic cation transporter genes SLC22A1 and SLC22A2
[20091] are located on chromosome 6q26. Cytogenet. Cell Genet. 79: 198-200,
[20092] 1997.

[20093] 2144.

[20094] Zhang, L.; Dresser, M. J.; Gray, A. T.; Yost, S. C.; Terashita,

[20095] S.; Giacomini, K. M.: Cloning and functional expression of

a human
[20096] liver organic cation transporter. Molec. Pharm. 51: 913-921, 1997.
[20097] 2145.
[20098] Grundemann, D.; Schomig, E.: Gene structures of the human non-neuronal
[20099] monoamine transporters EMT and OCT2. Hum. Genet. 106: 627-635, 2000.
[20100] 2146.
[20101] McLoughlin, D. M.; Miller, C. C. J.: The intracellular cytoplasmic
[20102] domain of the Alzheimer's disease amyloid precursor protein interacts
[20103] with phosphotyrosine-binding domain proteins in the yeast two-hybrid
[20104] system. FEBS Lett. 397: 197-200, 1996.
[20105] 2147.
[20106] Guenette, S. Y.; Chen, J.; Jondro, P. D.; Tanzi, R. E.: Association
[20107] of a novel human FE65-like protein with the cytoplasmic domain of
[20108] the beta-amyloid precursor protein. Proc. Nat. Acad. Sci.

93: 10832-10837,

[20109] 1996.

[20110] 2148.

[20111] Duilio, A.; Faraonio, R.; Minopoli, G.; Zambrano, N.; Russo, T.

[20112] : Fe65L2: a new member of the Fe65 protein family interacting with

[20113] the intracellular domain of the Alzheimer's beta-amyloid precursor

[20114] protein. Biochem. J. 330: 513-519, 1998.

[20115] 2149.

[20116] Tanahashi, H.; Tabira, T.: Genome structure and chromosomal mapping

[20117] of the gene for Fe65L2 interacting with Alzheimer's beta-amyloid precursor

[20118] protein. Biochem. Biophys. Res. Commun. 258: 385-389, 1999.

[20119] 2150.

[20120] Tanahashi, H.; Tabira, T.: Molecular cloning of human Fe65L2 and

[20121] its interaction with the Alzheimer's beta-amyloid precur-

sor protein. Neurosci.

[20122] Lett. 261: 143-146, 1999.

[20123] 2151.

[20124] Howard, L.; Nelson, K. K.; Maciewicz, R. A.; Blobel, C. P.: Interaction

[20125] of the metalloprotease disintegrins MDC9 and MDC15 with two SH3 domain-containing

[20126] proteins, endophilin I and SH3PX1. J. Biol. Chem. 274: 31693-31699,

[20127] 1999.

[20128] 2152.

[20129] Galliano, M.-F.; Huet, C.; Frygelius, J.; Polgren, A.; Wewer, U.

[20130] M.; Engvall, E.: Binding of ADAM12, a marker of skeletal muscle regeneration,

[20131] to the muscle-specific actin-binding protein, alpha-actinin-2, is

[20132] required for myoblast fusion. J. Biol. Chem. 275: 13933-13939, 2000.

[20133] 2153.

[20134] Gilpin, B. J.; Loechel, F.; Mattei, M.-G.; Engvall, E.; Al-

brechtsen,
[20135] R.; Wewer, U. M.: A novel, secreted form of human ADAM 12 (meltrin
[20136] alpha) provokes myogenesis in vivo. J. Biol. Chem. 273: 157-166,
[20137] 1998.

[20138] 2154.

[20139] Yagami-Hiromasa, T.; Sato, T.; Kurisaki, T.; Kamijo, K.; Nabeshima,
[20140] Y.; Fujisawa-Sehara, A.: A metalloprotease-disintegrin participating
[20141] in myoblast fusion. Nature 377: 652-656, 1995.

[20142] 2155.

[20143] Brown, R. A.; Ho, L. K. F.; Weber-Hall, S. J.; Shipley, J. M.;

[20144] Fry, M. J.: Identification and cDNA cloning of a novel mammalian
[20145] C2 domain-containing phosphoinositide 3-kinase, HsC2-PI3K. Biochem.
[20146] Biophys. Res. Commun. 233: 537-544, 1997.

[20147] 2156.

[20148] Chantry, D.; Vojtek, A.; Kashishian, A.; Holtzman, D. A.;

Wood,
[20149] C.; Gray, P. W.; Cooper, J. A.; Hoekstra, M. F.: p110-delta, a novel
[20150] phosphatidylinositol 3-kinase catalytic subunit that associates with
[20151] p85 and is expressed predominantly in leukocytes. J. Biol. Chem. 272:
[20152] 19236-19241, 1997.

[20153] 2157.

[20154] Okkenhaug, K.; Bilancio, A.; Farjot, G.; Priddle, H.; Sancho, S.;
[20155] Peskett, E.; Pearce, W.; Meek, S. E.; Salpekar, A.; Waterfield, M.
[20156] D.; Smith, A. J. H.; Vanhaesebroeck, B.: Impaired B and T cell antigen
[20157] receptor signaling in p110-delta PI 3-kinase mutant mice. Science 297:
[20158] 1031-1034, 2002.

[20159] 2158.

[20160] Seki, N.; Nimura, Y.; Ohira, M.; Saito, T.; Ichimiya, S.; Nomura,
[20161] N.; Nakagawara, A.: Identification and chromosome as-

signment of a

[20162] human gene encoding a novel phosphatidylinositol-3 ki-
nase. DNA Res. 4:

[20163] 355-358, 1997.

[20164] 2159.

[20165] Vanhaesebroeck, B.; Welham, M. J.; Kotani, K.; Stein, R.;
Warne,

[20166] P. H.; Zvelebil, M. J.; Higashi, K.; Volinia, S.; Downward, J.;
Waterfield,

[20167] M. D.: p110-delta, a novel phosphoinositide 3-kinase in
leukocytes. Proc.

[20168] Nat. Acad. Sci. 94: 4330-4335, 1997.

[20169] 2160.

[20170] Bowman, M. R.; Crimmins, M. A. V.; Yetz-Aldape, J.; Kriz,
R.; Kelleher,

[20171] K.; Herrmann, S.: The cloning of CD70 and its identifica-
tion as the

[20172] ligand for CD27. J. Immun. 152: 1756-1761, 1994.

[20173] 2161.

[20174] Slaugenhaupt, S. A.; Blumenfeld, A.; Gill, S. P.; Leyne, M.;
Mull,

[20175] J.; Cuajungco, M. P.; Liebert, C. B.; Chadwick, B.; Idelson, M.; Reznik,

[20176] L.; Robbins, C. M.; Makalowska, I.; Brownstein, M. J.; Krappmann,

[20177] D.; Scheidereit, C.; Maayan, C.; Axelrod, F. B.; Gusella, J. F.:

[20178] Tissue-specific expression of a splicing mutation in the IKBKAP gene

[20179] causes familial dysautonomia. Am. J. Hum. Genet. 68: 598-605, 2001.

[20180] 2162.

[20181] Okubo, M.; Kanda, F.; Horinishi, A.; Takahashi, K.; Okuda, S.;

[20182] Chihara, K.; Murase, T.: Glycogen storage disease type IIIa: first

[20183] report of a causative missense mutation (G1448R) of the glycogen debranching

[20184] enzyme gene found in a homozygous patient. (Abstract) Hum. Mutat. 14:

[20185] 542-543, 1999.

[20186] 2163.

[20187] Parvari, R.; Moses, S.; Shen, J.; Hershkovitz, E.; Lerner, A.;

[20188] Chen, Y.-T.: A single-base deletion in the 3-prime coding region
[20189] of glycogen-debranching enzyme is prevalent in glycogen storage disease
[20190] type IIIA in a population of North African Jewish patients. Europ.
[20191] J. Hum. Genet. 5: 266-270, 1997.

[20192] 2164.

[20193] Cheng, G.; Ye, Z.-S.; Baltimore, D.: Binding of Bruton's tyrosine
[20194] kinase to Fyn, Lyn, or Hck through a Src homology 3 domain-mediated
[20195] interaction. Proc. Nat. Acad. Sci. 91: 8152-8155, 1994.

[20196] 2165.

[20197] Gibson, S.; Leung, B.; Squire, J. A.; Hill, M.; Arima, N.; Goss,
[20198] P.; Hogg, D.; Mills, G. B.: Identification, cloning, and characterization
[20199] of a novel human T-cell-specific tyrosine kinase located at the hematopoietin
[20200] complex on chromosome 5q. Blood 82: 1561-1572, 1993.

[20201] 2166.

[20202] Janis, E. M.; Siliciano, J. D.; Isaac, D. D.; Griffin, C. A.; Hawkins,

[20203] A. L.; Kozak, C. A.; Desiderio, S.: Mapping of the gene for the tyrosine

[20204] kinase Itk to a region of conserved synteny between mouse chromosome

[20205] 11 and human chromosome 5q. Genomics 23: 269-271, 1994.

[20206] 2167.

[20207] Schaeffer, E. M.; Debnath, J.; Yap, G.; McVicar, D.; Liao, X. C.;

[20208] Littman, D. R.; Sher, A.; Varmus, H. E.; Lenardo, M. J.; Schwartzberg,

[20209] P. L.: Requirement for Tec kinases Rlk and Itk in T cell re-

ceptor

[20210] signaling and immunity. Science 284: 638-641, 1999.

[20211] 2168.

[20212] Woods, M. L.; Kivens, W. J.; Adelsman, M. A.; Qiu, Y.; Au-

gust,

[20213] A.; Shimizu, Y.: A novel function for the Tec family tyro-

sine kinase

[20214] Itk in activation of beta-1 integrins by the T-cell receptor. EMBO
[20215] J. 20: 1232-1244, 2001.

[20216] 2169.

[20217] Blanche, H.; Wright, L. G.; Vergnaud, G.; de Gouyon, B.; Lauthier,
[20218] V.; Silver, L. M.; Dausset, J.; Cann, H. M.; Spielman, R. S.: Genetic
[20219] mapping of three human homologues of murine t-complex genes localizes
[20220] TCP10 to 6q27, 15 cM distal to TCP1 and PLG. Genomics 12: 826-828,
[20221] 1992.

[20222] 2170.

[20223] Spencer, N.; Hopkinson, D. A.: Biochemical genetics of the pentose
[20224] phosphate cycle: human ribose 5-phosphate isomerase (RPI) and ribulose
[20225] 5-phosphate 3-epimerase (RPE). Ann. Hum. Genet. 43: 335-342, 1980.
[20226] 2171.

[20227] Elliott, R. W.; Samuelson, L. C.; Lambert, M. S.; Meisler, M. H.

[20228] : Assignment of pancreatic ribonuclease gene to mouse chromosome 14. Cytogenet.

[20229] Cell Genet. 42: 110-112, 1986.

[20230] 2172.

[20231] Piccoli, R.; Di Gaetano, S.; De Lorenzo, C.; Grauso, M.; Monaco,

[20232] C.; Spalletti-Cernia, D.; Laccetti, P.; Cinatl, J.; Matousek, J.;

[20233] D'Alessio, G.: A dimeric mutant of human pancreatic ri-bonuclease

[20234] with selective cytotoxicity toward malignant cells. Proc. Nat. Acad.

[20235] Sci. 96: 7768-7773, 1999.

[20236] 2173.

[20237] Zhang, J.; Zhang, Y.; Rosenberg, H. F.: Adaptive evolution of

[20238] a duplicated pancreatic ribonuclease gene in a leaf-eating monkey. Nature

[20239] Genet. 30: 416-420, 2002.

[20240] 2174.

[20241] Lilja, H.; Abrahamsson, P.-A.; Lundwall, A.: Semenogelin, the
[20242] predominant protein in human semen: primary structure and identification
[20243] of closely related proteins in the male accessory sex glands and on
[20244] the spermatozoa. J. Biol. Chem. 264: 1894-1900, 1989.

[20245] 2175.

[20246] Loeffler, C.; Rao, V. V. N. G.; Schnittger, S.; Pfau, H. P.; Lundwall,
[20247] A.; Schaefer, R.; Stolz, F. M.; Hansmann, I.: Personal Communication.
[20248] 8/18/1991.

[20249] 2176.

[20250] Ulvsback, M.; Lazure, C.; Lilja, H.; Spurr, N. K.; Rao, V. V.;

[20251] Loffler, C.; Hansmann, I.; Lundwall, A.: Gene structure of semenogelin
[20252] I and II: the predominant proteins in human semen are encoded by two
[20253] homologous genes on chromosome 20. J. Biol. Chem. 267: 18080-18084,

[20254] 1992.

[20255] 2177.

[20256] Byrne, P. C.; Shipley, J. M.; Chave, K. J.; Sanders, P. G.; Snell,

[20257] K.: Characterisation of a human serine hydroxymethyl-transferase pseudogene

[20258] and its localisation of 1p32.3-33. Hum. Genet. 97: 340-344, 1996.

[20259] 2178.

[20260] Elsea, S. H.; Juyal, R. C.; Jiralerspong, S.; Finucane, B. M.;

[20261] Pandolfo, M.; Greenberg, F.; Baldini, A.; Stover, P.; Patel, P. I.

[20262] : Haploinsufficiency of cytosolic serine hydroxymethyl-transferase

[20263] in the Smith-Magenis syndrome. Am. J. Hum. Genet. 57: 1342-1350,

[20264] 1995.

[20265] 2179.

[20266] Baecher, C. M.; Dorfman, K. S.; Mattei, M.-G.; Frelinger, J. G.

[20267] : cDNA cloning and localization of the mouse leukosialin

gene (Ly48)
[20268] to chromosome 7. Immunogenetics 31: 307-314, 1990.

[20269] 2180.

[20270] Bazil, V.; Strominger, J. L.: CD43, the major sialoglycopro-
tein
[20271] of human leukocytes, is proteolytically cleaved from the
surface of
[20272] stimulated lymphocytes and granulocytes. Proc. Nat. Acad.
Sci. 90:
[20273] 3792-3796, 1993.

[20274] 2181.

[20275] Delon, J.; Kaibuchi, K.; Germain, R. N.: Exclusion of CD43
from
[20276] the immunological synapse is mediated by phosphoryla-
tion-regulated
[20277] relocation of the cytoskeletal adaptor moesin. Immunity
15: 691-701,
[20278] 2001.

[20279] 2182.

[20280] Pallant, A.; Eskenazi, A.; Mattei, M.-G.; Fournier, R. E. K.;
Carlsson,

[20281] S. R.; Fukuda, M.; Frelinger, J. G.: Characterization of cD-
NAs encoding
[20282] human leukosialin and localization of the leukosialin gene
to chromosome
[20283] 16. Proc. Nat. Acad. Sci. 86: 1328-1332, 1989.

[20284] 2183.

[20285] Park, J. K.; Rosenstein, Y. J.; Remold-O'Donnell, E.; Bierer,
B.
[20286] E.; Rosen, F. S.; Burakoff, S. J.: Enhancement of T-cell acti-
vation
[20287] by the CD43 molecule whose expression is defective in
Wiskott-Aldrich
[20288] syndrome. Nature 350: 706-709, 1991.

[20289] 2184.

[20290] Rosenstein, Y.; Park, J. K.; Hahn, W. C.; Rosen, F. S.; Bierer,

[20291] B. E.; Burakoff, S. J.: CD43, a molecule defective in
Wiskott-Aldrich
[20292] syndrome, binds ICAM-1. Nature 354: 233-235, 1991.

[20293] 2185.

[20294] Schmid, K.; Hediger, M. A.; Brossmer, R.; Collins, J. H.;
Haupt,

[20295] H.; Marti, T.; Offner, G. D.; Schaller, J.; Takagaki, K.; Walsh, M.

[20296] T.; Schwick, H. G.; Rosen, F. S.; Remold-O'Donnell, E.: Amino acid

[20297] sequence of human plasma galactoglycoprotein: identity with the extracellular

[20298] region of CD43 (sialophorin). Proc. Nat. Acad. Sci. 89: 663-667,

[20299] 1992.

[20300] 2186.

[20301] Shelley, C. S.; Carroll, M. C.; Davis, A. E., III; Bruns, G. A.

[20302] P.; Whitehead, A. S.; Finberg, R.; Remold-O'Donnell, E.; Rosen, F.

[20303] S.: Human sialophorin: cDNA cloning and gene location. (Abstract) FASEB

[20304] J. 2: A1659 only, 1988.

[20305] 2187.

[20306] Shelley, C. S.; Remold-O'Donnell, E.; Davis, A. E., III; Bruns,

[20307] G. A. P.; Rosen, F. S.; Carroll, M. C.; Whitehead, A. S.: Molecular

[20308] characterization of sialophorin (CD43), the lymphocyte

surface sialoglycoprotein
[20309] defective in Wiskott-Aldrich syndrome. Proc. Nat. Acad. Sci. 86:
[20310] 2819-2823, 1989.

[20311] 2188.

[20312] Shelley, C. S.; Remold-O'Donnell, E.; Rosen, F. S.; White-head,
[20313] A. S.: Structure of the human sialophorin (CD43) gene: identification
[20314] of features atypical of genes encoding integral membrane proteins. Biochem.
[20315] J. 270: 569-576, 1990.

[20316] 2189.

[20317] van den Berg, T. K.; Nath, D.; Ziltener, H. J.; Vestweber, D.;

[20318] Fukuda, M.; van Die, I.; Crocker, P. R.: Cutting edge: CD43 functions
[20319] as a T cell counterreceptor for the macrophage adhesion receptor sialoadhesin
[20320] (Siglec-1). J. Immun. 166: 3637-3640, 2001.

[20321] 2190.

[20322] Ianakiev, P.; Kilpatrick, M. W.; Toudjarska, I.; Basel, D.;

Beighton,
[20323] P.; Tsipouras, P.: Split-hand/split-foot malformation is caused by
[20324] mutations in the p63 gene on 3q27. Am. J. Hum. Genet. 67: 59-66,
[20325] 2000.

[20326] 2191.

[20327] Allen, R. C.; Webster, A. R.; Sui, R.; Brown, J.; Taylor, C. M.;
[20328] Stone, E. M.: Molecular characterization and ophthalmic investigation
[20329] of a large family with type 2A von Hippel-Lindau disease. Arch. Ophthal. 119:
[20330] 1659-1665, 2001.

[20331] 2192.

[20332] Boissy, R. E.; Zhao, H.; Oetting, W. S.; Austin, L. M.; Wildenberg,
[20333] S. C.; Boissy, Y. L.; Zhao, Y.; Sturm, R. A.; Hearing, V. J.; King,
[20334] R. A.; Nordlund, J. J.: Mutation in and lack of expression of tyrosinase-related
[20335] protein-1 (TRP-1) in melanocytes from an individual with

brown oculocutaneous
[20336] albinism: a new subtype of albinism classified as 'OCA3.' Am. J.
[20337] Hum. Genet. 58: 1145-1156, 1996.

[20338] 2193.

[20339] Anikster, Y.; Huizing, M.; White, J.; Shevchenko, Y. O.; Fitzpatrick,
[20340] D. L.; Touchman, J. W.; Compton, J. G.; Bale, S. J.; Swank, R. T.;
[20341] Gahl, W. A.; Toro, J. R.: Mutation of a new gene causes a unique
[20342] form of Hermansky-Pudlak syndrome in a genetic isolate of central
[20343] Puerto Rico. Nature Genet. 28: 376-380, 2001.

[20344] 2194.

[20345] Feng, G. H.; Bailin, T.; Oh, J.; Spritz, R. A.: Mouse pale ear
[20346] (ep) is homologous to human Hermansky-Pudlak syndrome and contains
[20347] a rare 'AT-AC' intron. Hum. Molec. Genet. 6: 793-797, 1997.
[20348] 2195.

[20349] Gahl, W. A.; Brantly, M.; Kaiser-Kupfer, M. I.; Iwata, F.; Hazelwood,
[20350] S.; Shotelersuk, V.; Duffy, L. F.; Kuehl, E. M.; Troendle, J.; Bernardini,
[20351] I.: Genetic defects and clinical characteristics of patients with
[20352] a form of oculocutaneous albinism (Hermansky-Pudlak syndrome). New
[20353] Eng. J. Med. 338: 1258-1264, 1998.

[20354] 2196.

[20355] Oh, J.; Bailin, T.; Fukai, K.; Feng, G. H.; Ho, L.; Mao, J.; Frenk,
[20356] E.; Tamura, N.; Spritz, R. A.: Positional cloning of a gene for Hermansky-Pudlak
[20357] syndrome, a disorder of cytoplasmic organelles. Nature Genet. 14:
[20358] 300-306, 1996.

[20359] 2197.

[20360] Escamilla, M. A.; DeMille, M. C.; Benavides, E.; Roche, E.; Almasy,
[20361] L.; Pittman, S.; Hauser, J.; Lew, D. F.; Freimer, N. B.; Whit-
tle,

[20362] M. R.: A minimalist approach to gene mapping: locating the gene for
[20363] acheiropodia, by homozygosity analysis. Am. J. Hum. Genet. 66: 1995-2000,
[20364] 2000.

[20365] 2198.

[20366] Ianakiev, P.; van Baren, M. J.; Daly, M. J.; Toledo, S. P. A.;

[20367] Cavalcanti, M. G.; Neto, J. C.; Silveira, E. L.; Freire-Maia, A.;

[20368] Heutink, P.; Kilpatrick, M. W.; Tsipouras, P.: Acheiropodia is caused
[20369] by a genomic deletion in C7orf2, the human orthologue of the Lmbr1
[20370] gene. Am. J. Hum. Genet. 68: 38-45, 2001.

[20371] 2199.

[20372] Beebe, S. J.; Oyen, O.; Sandberg, M.; Froysa, A.; Hansson, V.;
[20373] Jahnsen, T.: Molecular cloning of a tissue-specific protein kinase
[20374] (C gamma) from human testis--representing a third iso-form for the
[20375] catalytic subunit of cAMP-dependent protein kinase.

Molec. Endocr. 4:
[20376] 465-475, 1990.

[20377] 2200.

[20378] Foss, K. B.; Berube, D.; Simard, J.; Beebe, S. J.; Sandberg, M.;
[20379] Grzeschik, K.-H.; Gagne, R.; Hansson, V.; Jahnsen, T.: Localization
[20380] of the catalytic subunit C-gamma of cAMP-dependent protein kinase
[20381] on human chromosome 9q13. (Abstract) Cytogenet. Cell Genet. 58:
[20382] 1937-1938, 1991.

[20383] 2201.

[20384] Foss, K. B.; Simard, J.; Berube, D.; Beebe, S. J.; Sandberg, M.;
[20385] Grzeschik, K.-H.; Gagne, R.; Hansson, V.; Jahnsen, T.: Localization
[20386] of the catalytic subunit C-gamma of the cAMP-dependent protein kinase
[20387] gene (PRKACG) to human chromosome region 9q13. Cytogenet. Cell Genet. 60:
[20388] 22-25, 1992.

[20389] 2202.

[20390] Reinton, N.; Haugen, T. B.; Orstavik, S.; Skalhegg, B. S.; Hansson,

[20391] V.; Jahnsen, T.; Tasken, K.: The gene encoding the C gamma catalytic

[20392] subunit of cAMP-dependent protein kinase is a transcribed retroposon. Genomics

[20393] 49: 290-297, 1998.

[20394] 2203.

[20395] Orstavik, S.; Natarajan, V.; Tasken, K.; Jahnsen, T.; Sandberg,

[20396] M.: Characterization of the human gene encoding the type I-alpha
[20397] and type I-beta cGMP-dependent protein kinase (PRKG1). Genomics 42:
[20398] 311-318, 1997.

[20399] 2204.

[20400] Orstavik, S.; Sandberg, M.; Berube, D.; Natarajan, V.; Simard,
[20401] J.; Walter, U.; Gagne, R.; Hansson, V.; Jahnsen, T.: Local-ization
[20402] of the human gene for type I cyclic GMP-dependent pro-tein kinase to
[20403] chromosome 10. Cytogenet. Cell Genet. 59: 270-273, 1992.
[20404] 2205.

[20405] Osborne, K. A; Robichon, A.; Burgess, E.; Butland, S.; Shaw, R.
[20406] A.; Coulthard, A.; Pereira, H. S.; Greenspan, R. J.; Sokolowski, M.
[20407] B.: Natural behavior polymorphism due to a cGMP-dependent protein
[20408] kinase of Drosophila. Science 277: 834-836, 1997.

[20409] 2206.

[20410] Pfeifer, A.; Klatt, P.; Massberg, S.; Ny, L.; Sausbier, M.;
Hirneill,
[20411] C.; Wang, G.-X.; Korth, M.; Aszodi, A.; Andersson, K.-E.;
Krombach,
[20412] F.; Mayerhofer, A.; Ruth, P.; Fassler, R.; Hofmann, F.: De-
fective
[20413] smooth muscle regulation in cGMP kinase I-deficient
mice. EMBO J. 17:
[20414] 3045-3051, 1998.

[20415] 2207.

[20416] Sandberg, M.; Natarajan, V.; Ronander, I.; Kalderon, D.;
Walter,
[20417] U.; Lohmann, S. M.; Jahnsen, T.: Molecular cloning and
predicted
[20418] full-length amino acid sequence of the type I beta
isozyme of cGMP-dependent
[20419] protein kinase from human placenta: tissue distribution
and developmental
[20420] changes in rat. FEBS Lett. 255: 321-329, 1989.

[20421] 2208.

[20422] Tamura, N.; Itoh, H.; Ogawa, Y.; Nakagawa, O.; Harada, M.; Chun,
[20423] T.-H.; Suga, S.; Yoshimasa, T.; Nakao, K.: cDNA cloning and gene
[20424] expression of human type I-alpha cGMP-dependent protein kinase. Hypertension 27:
[20425] 552-557, 1996.

[20426] 2209.

[20427] Oyen, O.; Myklebust, F.; Scott, J. D.; Hansson, V.; Jahnsen, T.
[20428] : Human testis cDNA for the regulatory subunit RII alpha of cAMP-dependent
[20429] protein kinase encodes an alternate amino-terminal region. FEBS Lett. 246:
[20430] 57-64, 1989.

[20431] 2210.

[20432] Tasken, K.; Naylor, S. L.; Solberg, R.; Jahnsen, T.: Mapping of
[20433] the gene encoding the regulatory subunit RII-alpha of cAMP-dependent
[20434] protein kinase (locus PRKAR2A) to human chromosome region 3p21.3-p21.2. Genomics 50:

[20435] 378-381, 1998.

[20436] 2211.

[20437] Solberg, R.; Sistonen, P.; Traskelin, A.-L.; Berube, D.; Simard,

[20438] J.; Krajci, P.; Jahnsen, T.; de la Chapelle, A.: Mapping of the regulatory

[20439] subunits RI-beta and RII-beta of cAMP-dependent protein kinase genes

[20440] on human chromosome 7. Genomics 14: 63-69, 1992.

[20441] 2212.

[20442] Cummings, D. E.; Brandon, E. P.; Planas, J. V.; Motamed, K.; Idzerda,

[20443] R. L.; McKnight, G. S.: Genetically lean mice result from targeted

[20444] disruption of the RII-beta subunit of protein kinase A. Na- ture 382:

[20445] :622-626, 1996.

[20446] 2213.

[20447] Scambler, P.; Oyen, O.; Wainwright, B.; Farrall, M.; Law, H.-Y.;

[20448] Estivill, X.; Sandberg, M.; Williamson, R.; Jahnsen, T.: Ex-

clusion

[20449] of catalytic and regulatory subunits of cAMP-dependent protein kinase

[20450] as candidate genes for the defect causing cystic fibrosis. Am. J.

[20451] Hum. Genet. 41: 925-932, 1987.

[20452] 2214.

[20453] Wainwright, B.; Lench, N.; Davies, K.; Scambler, P.; Kruyer, H.;

[20454] Williamson, R.; Jahnsen, T.; Farrall, M.: A human regulatory subunit

[20455] of type II cAMP-dependent protein kinase localized by its linkage

[20456] relationship to several cloned chromosome 7q markers. Cytogenet.

[20457] Cell Genet. 45: 237-239, 1987.

[20458] 2215.

[20459] Norman, S. A.; Mott, D. M.: Molecular cloning and chromosomal

[20460] localization of a human skeletal muscle PP-1-gamma-1 cDNA. Mammalian

[20461] Genome 5: 41-45, 1994.

[20462] 2216.

[20463] Sugawara, S.; Uehara, A.; Nochi, T.; Yamaguchi, T.; Ueda, H.;

[20464] Sugiyama, A.; Hanzawa, K.; Kumagai, K.; Okamura, H.; Takada, H.:

[20465] Neutrophil proteinase 3-mediated induction of bioactive IL-18 secretion

[20466] by human oral epithelial cells. J. Immun. 167: 6568-6575, 2001.

[20467] 2217.

[20468] Akiyama, K.; Yokota, K.; Kagawa, S.; Shimbara, N.; Tamura, T.;

[20469] Akioka, H.; Nothwang, H. G.; Noda, C.; Tanaka, K.; Ichi-hara, A.:

[20470] cDNA cloning and interferon gamma down-regulation of proteasomal subunits

[20471] X and Y. Science 265: 1231-1234, 1994.

[20472] 2218.

[20473] Driscoll, J.; Brown, M. G.; Finley, D.; Monaco, J. J.: MHC-linked

[20474] LMP gene products specifically alter peptidase activities of the proteasome. Nature 365:

[20475] 262-264, 1993.

[20476] 2219.

[20477] Gaczynska, M.; Rock, K. L.; Goldberg, A. L.: Gamma-in-
terferon
[20478] and expression of MHC genes regulate peptide hydrolysis
by proteasomes. Nature 365:
[20479] 264-267, 1993.

[20480] 2220.

[20481] Glynne, R.; Powis, S. H.; Beck, S.; Kelly, A.; Kerr, L. A.;
Trowsdale,
[20482] J.: A proteasome-related gene between the two ABC
transporter loci
[20483] in the class II region of the human MHC. Nature 353:
357-360, 1991.
[20484] 2221.

[20485] Gray, P. W.; Barrett, K.; Chantry, D.; Turner, M.; Feldmann,
M.
[20486] : Cloning of human tumor necrosis factor (TNF) receptor
cDNA and expression
[20487] of recombinant soluble TNF-binding protein. Proc. Nat.
Acad. Sci. 87:

[20488] 7380-7384, 1990.

[20489] 2222.

[20490] Hohmann, H.-P.; Remy, R.; Brockhaus, M.; van Loon, A. P. G. M.

[20491] : Two different cell types have different major receptors for human

[20492] tumor necrosis factor (TNF-alpha). J. Biol. Chem. 264: 14927-14934,

[20493] 1989.

[20494] 2223.

[20495] Karenko, L.; Pettersson, T.; Roberts, P.: Autosomal domi-nant

[20496] 'Mediterranean fever' in a Finnish family. J. Int. Med. 232: 365-369,

[20497] 1992.

[20498] 2224.

[20499] Loetscher, H.; Pan, Y.-C. E.; Lahm, H.-W.; Gentz, R.; Brockhaus,

[20500] M.; Tabuchi, H.; Lesslauer, W.: Molecular cloning and ex-pression

[20501] of the human 55 kd tumor necrosis factor receptor. Cell

61: 351-359,

[20502] 1990.

[20503] 2225.

[20504] McDermott, M. F.; Aksentijevich, I.; Galon, J.; McDermott,
E.

[20505] M.; Ogunkolade, B. W.; Centola, M.; Mansfield, E.; Gadina,
M.; Karenko,

[20506] L.; Pettersson, T.; McCarthy, J.; Frucht, D. M.; and 21 oth-
ers:

[20507] Germline mutations in the extracellular domains of the 55
kDa TNF

[20508] receptor, TNFR1, define a family of dominantly inherited
autoinflammatory

[20509] syndromes. Cell 97: 133-144, 1999.

[20510] 2226.

[20511] McDermott, M. F.; Ogunkolade, B. W.; McDermott, E. M.;
Jones,

[20512] L. C.; Wan, Y.; Quane, K. A.; McCarthy, J.; Phelan, M.; Mol-
loy, M.

[20513] G.; Powell, R. J.; Amos, C. I.; Hitman, G. A.: Linkage of fa-
milial

[20514] Hibernian fever to chromosome 12p13. Am. J. Hum.

Genet. 62: 1446-1451,

[20515] 1998.

[20516] 2227.

[20517] Milatovich, A.; Song, K.; Heller, R. A.; Francke, U.: The genes

[20518] for two tumor necrosis factor receptors, TNFR1 and TNFR2, map to human

[20519] chromosomes 12 (12pter-cen) and 1 (1pter-p32). (Abstract) Cytogenet.

[20520] Cell Genet. 58: 1980 only, 1991.

[20521] 2228.

[20522] Milatovich, A.; Song, K.; Heller, R. A.; Francke, U.: Tumor necrosis

[20523] factor receptor genes, TNFR1 and TNFR2, on human chromosomes 12 and

[20524] 1. Somat. Cell Molec. Genet. 17: 519-523, 1991.

[20525] 2229.

[20526] Nophar, Y.; Kemper, O.; Brakebusch, C.; Engelmann, H.; Zwang,

[20527] R.; Aderka, D.; Holtmann, H.; Wallach, D.: Soluble forms of tumor

[20528] necrosis factor receptors (TNF-Rs): the cDNA for the type I TNF-R,

[20529] cloned using amino acid sequence data of its soluble form, encodes

[20530] both the cell surface and a soluble form of the receptor. EMBO J. 9:

[20531] 3269-3278, 1990.

[20532] 2230.

[20533] Qian, Y.; Dekaris, I.; Yamagami, S.; Dana, R.: Topical soluble

[20534] tumor necrosis factor receptor type I suppresses ocular chemokine

[20535] gene expression and rejection of allogeneic corneal transplants. Arch.

[20536] Ophthal. 118: 1666-1671, 2000.

[20537] 2231.

[20538] Rothe, J.; Lesslauer, W.; Lotscher, H.; Lang, Y.; Koebel, P.;

[20539] Kontgen, F.; Althage, A.; Zinkernagel, R.; Steinmetz, M.; Bluethmann,

[20540] H.: Mice lacking the tumour necrosis factor receptor 1 are resistant

[20541] to TNF-mediated toxicity but highly susceptible to infection by Listeria

[20542] monocytogenes. Nature 364: 798-802, 1993.

[20543] 2232.

[20544] Schall, T. J.; Lewis, M.; Koller, K. J.; Lee, A.; Rice, G. C.;

[20545] Wong, G. H. W.; Gatanaga, T.; Granger, G. A.; Lentz, R.; Raab, H.;

[20546] Kohr, W. J.; Goeddel, D. V.: Molecular cloning and expression of

[20547] a receptor for human tumor necrosis factor. Cell 61: 361-370, 1990.

[20548] 2233.

[20549] Schievella, A. R.; Chen, J. H.; Graham, J. R.; Lin, L.-L.: MADD,

[20550] a novel death domain protein that interacts with the type 1 tumor

[20551] necrosis factor receptor and activates mitogen-activated protein kinase. J.

[20552] Biol. Chem. 272: 12069-12075, 1997.

[20553] 2234.

[20554] Smith, C. A.; Davis, T.; Anderson, D.; Solam, L.; Beckmann,

M.

[20555] P.; Jerzy, R.; Dower, S. K.; Cosman, D.; Goodwin, R. G.: A receptor

[20556] for tumor necrosis factor defines an unusual family of cellular and

[20557] viral proteins. Science 248: 1019-1023, 1990.

[20558] 2235.

[20559] Stauber, G. B.; Aiyer, R. A.; Aggarwal, B. B.: Human tumor necrosis

[20560] factor-alpha receptor: purification by immunoaffinity chromatography

[20561] and initial characterization. J. Biol. Chem. 263: 19098-19104, 1988.

[20562] 2236.

[20563] Beltinger, C. P.; White, P. S.; Maris, J. M.; Sulman, E. P.; Jensen,

[20564] S. J.; LePaslier, D.; Stallard, B. J.; Goeddel, D. V.; de Sauvage,

[20565] F. J.; Brodeur, G. M.: Physical mapping and genomic structure of

[20566] the human TNFR2 gene. Genomics 35: 94-100, 1996.

[20567] 2237.

[20568] Kaufman, B. A.; White, P. S.; Steinbrueck, T.; Donis-Keller, H.;

[20569] Brodeur, G. M.: Linkage mapping of the tumor necrosis factor receptor

[20570] 2 (TNFR2) gene to 1p36.2 using the single-strand conformation polymorphism

[20571] technique. Hum. Genet. 94: 418-422, 1994.

[20572] 2238.

[20573] Kemper, O.; Derre, J.; Cherif, D.; Engelmann, H.; Wallach, D.;

[20574] Berger, R.: The gene for the type II (p75) tumor necrosis factor

[20575] receptor (TNF-RII) is localized on band 1p36.2-p36.3. Hum. Genet. 87:

[20576] 623-624, 1991.

[20577] 2239.

[20578] Li, X.; Yang, Y.; Ashwell, J. D.: TNF-RII and c-IAP1 mediate ubiquitination

[20579] and degradation of TRAF2. Nature 416: 345-349, 2002.

[20580] 2240.

[20581] Santee, S. M.; Owen-Schaub, L. B.: Human tumor necrosis

factor
[20582] receptor p75/80 (CD120b) gene structure and promoter characterization. J.
[20583] Biol. Chem. 271: 21151-21159, 1996.

[20584] 2241.

[20585] Hon, W.-C.; Wilson, M. I.; Harlos, K.; Claridge, T. D. W.; Schofield,
[20586] C. J.; Pugh, C. W.; Maxwell, P. H.; Ratcliffe, P. J.; Stuart, D. I.;
[20587] Jones, E. Y.: Structural basis for the recognition of hydroxyproline
[20588] in HIF-1-alpha by pVHL. Nature 417: 975-978, 2002.

[20589] 2242.

[20590] Godfrey, M.; Hollister, D. W.: Type II achondrogenesis-hypochondrogenesis:
[20591] identification of abnormal type II collagen. Am. J. Hum. Genet. 43:
[20592] 904-913, 1988.

[20593] 2243.

[20594] Takenaka, M.; Noguchi, T.; Sadahiro, S.; Hirai, H.; Yamada, K.;

[20595] Matsuda, T.; Imai, E.; Tanaka, T.: Isolation and characterization
[20596] of the human pyruvate kinase M gene. Europ. J. Biochem. 198: 101-106,
[20597] 1991.

[20598] 2244.

[20599] Tani, K.; Yoshida, M. C.; Satoh, H.; Mitamura, K.; Noguchi, T.;
[20600] Tanaka, T.; Fujii, H.; Miwa, S.: Human M(2)-type pyruvate kinase:
[20601] cDNA cloning, chromosomal assignment and expression in hepatoma. Gene 73:
[20602] 509-516, 1988.

[20603] 2245.

[20604] Tsutsumi, H.; Tani, K.; Fujii, H.; Miwa, S.: Expression of L-
[20605] and M-type pyruvate kinase in human tissues. Genomics 2: 86-89,
[20606] 1988.

[20607] 2246.

[20608] Williams, J. M.; Chen, G.-C.; Zhu, L.; Rest, R. F.: Using the

[20609] yeast two-hybrid system to identify human epithelial cell proteins

[20610] that bind gonococcal Opa proteins: intracellular gonococci bind pyruvate

[20611] kinase via their Opa proteins and require host pyruvate for growth. Molec.

[20612] Microbiol. 27: 171-186, 1998.

[20613] 2247.

[20614] Westerveld, A.; Van Someren, H.; Van Henegouwen, H. M. A.; Oosterbaan,

[20615] R. A.: Synteny relationship between the human loci for hexosaminidase-A,

[20616] mannose phosphate isomerase, and pyruvate kinase-3: studies in man-Chinese

[20617] hamster somatic cell hybrids. Birth Defects Orig. Art. Ser. XI(3):

[20618] 283-284, 1975. Note: Alternate: Cytogenet. Cell Genet. 14: 453-454,

[20619] 1975.

[20620] 2248.

[20621] Seabra, M. C.; Brown, M. S.; Goldstein, J. L.: Retinal degeneration

[20622] in choroideremia: deficiency of Rab geranylgeranyl transferase. Science 259:

[20623] 377-381, 1993.

[20624] 2249.

[20625] Seabra, M. C.; Brown, M. S.; Slaughter, C. A.; Sudhof, T. C.; Goldstein,

[20626] J. L.: Purification of component A of Rab geranylgeranyl transferase:

[20627] possible identity with the choroideremia gene product. Cell 70:

[20628] 1049-1057, 1992.

[20629] 2250.

[20630] Seabra, M. C.; Reiss, Y.; Casey, P. J.; Brown, M. S.; Goldstein,

[20631] J. L.: Protein farnesyltransferase and geranylgeranyltransferase

[20632] share a common alpha subunit. Cell 65: 429-434, 1991.

[20633] 2251.

[20634] van Bokhoven, H.; Rawson, R. B.; Merkx, G. F. M.; Cremers, F. P.

[20635] M.; Seabra, M. C.: cDNA cloning and chromosomal local-

ization of the
[20636] genes encoding the alpha- and beta-subunits of human Rab geranylgeranyl
[20637] transferase: the 3-prime end of the alpha-subunit gene overlaps with
[20638] the transglutaminase 1 gene promoter. Genomics 38: 133-140, 1996.
[20639] 2252.

[20640] Koken, M. H. M.; Smit, E. M. E.; Jaspers-Dekker, I.; Oostra, B.
[20641] A.; Hagemeijer, A.; Bootsma, D.; Hoeijmakers, J. H. J.: Localization
[20642] of two human homologs, HHR6A and HHR6B, of the yeast DNA repair gene
[20643] RAD6 to chromosomes Xq24-q25 and 5q23-q31. Genomics 12: 447-453,
[20644] 1992.

[20645] 2253.

[20646] Lench, N. J.; Thompson, J.; Markham, A. F.; Robinson, P. A.: (CGG)
[20647] trinucleotide repeat polymorphism in the 5-prime region of the HHR6B

[20648] gene: the human homolog of the yeast DNA repair gene RAD6. Hum. Genet. 96:

[20649] 369-370, 1995.

[20650] 2254.

[20651] Roest, H. P.; van Klaveren, J.; de Wit, J.; van Gurp, C. G.; Koken,

[20652] M. H. M.; Vermey, M.; van Roijen, J. H.; Hoogerbrugge, J. W.; Vreeburg,

[20653] J. T. M.; Baarends, W. M.; Bootsma, D.; Grootegoed, J. A.; Hoeijmakers,

[20654] J. H. J.: Inactivation of the HR6B ubiquitin-conjugating DNA repair

[20655] enzyme in mice causes male sterility associated with chromatin modification. Cell 86:

[20656] 799-810, 1996.

[20657] 2255.

[20658] Roller, M. L.; Lossie, A. C.; Koken, M. H. M.; Smit, E. M. E.;

[20659] Hagemeijer, A.; Camper, S. A.: Localization of sequences related

[20660] to the human RAD6 DNA repair gene on mouse chromosomes 11 and 13. Mammalian

[20661] Genome 6: 305-306, 1995.

[20662] 2256.

[20663] Baur, M. P.; Rittner, C.: Application of a computer program for

[20664] the mapping of a gene locus to the disputed PGM-3 localization on

[20665] human chromosome 6. Tissue Antigens 12: 341-349, 1978.

[20666] 2257.

[20667] Burgess, R. M.; Sutton, J. G.: An improved method of typing hair

[20668] sheath cells using the PGM-3 locus following starch gel electrophoresis. Hum.

[20669] Genet. 56: 391-393, 1981.

[20670] 2258.

[20671] Hopkinson, D. A.; Harris, H.: A third phosphoglucomutase locus

[20672] in man. Ann. Hum. Genet. 31: 359-368, 1968.

[20673] 2259.

[20674] Jahannsmann, R.; Schwinger, E.; Grzeschik, K. H.: Assignment of

[20675] the gene locus for human phosphoglucomutase 3 to chromosome 6q12-qter. Ann.
[20676] Genet. 23: 12-14, 1980.

[20677] 2260.

[20678] Jongsma, A.; Van Someren, H.; Westerveld, A.; Hagemeijer, A.; Pearson,
[20679] P. L.: Localization of genes on human chromosomes by studies of human-Chinese
[20680] hamster somatic cell hybrids. Assignment of PGM to chromosome C6 and
[20681] regional mapping of the PGD, PGM, and pep-C genes on chromosome A1. Humangenetik 20:
[20682] 195-202, 1973.

[20683] 2261.

[20684] Kompf, J.; Bissbort, S.; Gohler, F.; Schunter, F.; Wernet, P.:

[20685] Mapping of the linkage group GLO-Bf-HLA-B, C, A-PGM. I. Recombination
[20686] frequencies. Hum. Genet. 44: 313-319, 1978.

[20687] 2262.

[20688] Lamm, L. U.; Jorgensen, F.; Kissmeyer-Nielsen, F.: On the mapping

[20689] of PGM-3 in relation to HLA. Tissue Antigens 17: 245-246, 1981.

[20690] 2263.

[20691] Lamm, L. U.; Kissmeyer-Nielsen, F.; Henningsen, K.: Linkage and

[20692] association studies of two phosphoglucomutase loci (PGM-1 and PGM-3)

[20693] to eighteen other markers. Hum. Hered. 20: 305-318, 1970.

[20694] 2264.

[20695] Li, C.; Rodriguez, M.; Banerjee, D.: Cloning and characterization

[20696] of complementary DNA encoding human N-acetylglucosamine-phosphate

[20697] mutase protein. Gene 242: 97-103, 2000.

[20698] 2265.

[20699] McCaw, B. K.; Hecht, F.; Linder, D.; Lovrien, E. W.; Wyandt, H.;

[20700] Bacon, D.; Clark, B.; Lea, N.: Ovarian teratomas: cytologic data. Cytogenet.

[20701] Cell Genet. 16: 391-395, 1976.

[20702] 2266.

[20703] Mohandas, T.; Sparkes, R. S.; Shulkin, J. D.; Sparkes, M. C.;

[20704] Moedjono, S.: Assignment of PGM3 to the long arm of hu-
man chromosome

[20705] 6: studies using Chinese hamster-human cell hybrids
containing a human

[20706] 6-15 translocation. Cytogenet. Cell Genet. 28: 116-120,
1980.

[20707] 2267.

[20708] Nadeau, J. H.; Kompf, J.; Siebert, G.; Taylor, B. A.: Linkage

[20709] of Pgm-3 in the house mouse and homologies of three
phosphoglucomutase

[20710] loci in mouse and man. Biochem. Genet. 19: 465-474,
1981.

[20711] 2268.

[20712] Pang, H.; Koda, Y.; Soejima, M.; Kimura, H.: Identification
of

[20713] human phosphoglucomutase 3 (PGM3) as N-
acetylglucosamine-phosphate

[20714] mutase (AGM1). Ann. Hum. Genet. 66: 139-144, 2002.

[20715] 2269.

[20716] Parrington, J.; West, L.; Povey, S.: Gene mapping from ovarian
[20717] teratomas. (Abstract) Cytogenet. Cell Genet. 25: 196 only, 1979.
[20718] 2270.

[20719] Van Someren, H.; Van Henegouwen, H. B.; Los, W.; Wurzer-Figurelli,
[20720] E.; Doppert, B.; Vervloet, M.; Meera Khan, P.: Enzyme electrophoresis
[20721] on cellulose acetate gel: zymogram patterns in man-Chinese hamster
[20722] somatic cell hybrids. Humangenetik 25: 189-201, 1974.

[20723] 2271.

[20724] Hickey, M. J.; Williams, S. A.; Roth, G. J.: Human platelet glycoprotein
[20725] IX: an adhesive prototype of leucine-rich glycoproteins with flank-center-flank
[20726] structures. Proc. Nat. Acad. Sci. 86: 6773-6777, 1989.

[20727] 2272.

[20728] Noris, P.; Arbustini, E.; Spedini, P.; Belletti, S.; Balduini,

[20729] C. L.: A new variant of Bernard-Soulier syndrome charac-

terized by

[20730] dysfunctional glycoprotein (GP) Ib and severely reduced amounts of

[20731] GPIX and GPV. Brit. J. Haemat. 103: 1004-1013, 1998.

[20732] 2273.

[20733] Noris, P.; Simsek, S.; Stibbe, J.; von dem Borne, A. E. G. K.:

[20734] A phenylalanine-55 to serine amino-acid substitution in the human

[20735] glycoprotein IX leucine-rich repeat is associated with Bernard-Soulier

[20736] syndrome. Brit. J. Haemat. 97: 312-320, 1997.

[20737] 2274.

[20738] Rivera, C. E.; Villagra, J.; Riordan, M.; Williams, S.; Lind-

strom,

[20739] K. J.; Rick, M. E.: Identification of a new mutation in platelet

[20740] glycoprotein IX (GPIX) in a patient with Bernard-Soulier syndrome. Brit.

[20741] J. Haemat. 112: 105-108, 2001.

[20742] 2275.

[20743] Wright, S. D.; Michaelides, K.; Johnson, D. J. D.; West, N.

C.;

[20744] Tuddenham, E. G. D.: Double heterozygosity for mutations in the platelet

[20745] glycoprotein IX gene in three siblings with Bernard-Soulier syndrome. Blood 81:

[20746] 2339-2347, 1993.

[20747] 2276.

[20748] Tobe, T.; Minoshima, S.; Yamase, S.; Choi, N.-H.; Tomita, M.;

[20749] Shimizu, N.: Assignment of a human serum glycoprotein SP-40,40 gene

[20750] (CLI) to chromosome 8. Cytogenet. Cell Genet. 57: 193-195, 1991.

[20751] 2277.

[20752] Wong, P.; Pineault, J.; Lakins, J.; Taillefer, D.; Leger, J.;

[20753] Wang, C.; Tenniswood, M.: Genomic organization and expression of

[20754] the rat TRPM-2 (clusterin) gene, a gene implicated in apoptosis. J.

[20755] Biol. Chem. 268: 5021-5031, 1993.

[20756] 2278.

[20757] Wong, P.; Taillefer, D.; Lakins, J.; Pineault, J.; Chader, G.;

[20758] Tenniswood, M.: Molecular characterization of human TRPM-2/clusterin,

[20759] a gene associated with sperm maturation, apoptosis and neurodegeneration. Europ.

[20760] J. Biochem. 221: 917-925, 1994.

[20761] 2279.

[20762] de Ronde, A.; Mannens, M.; Slater, R. M.; Hoovers, J.; Heyting,

[20763] C.; Bleeker-Wagemakers, E. M.; Leschot, N. J.; Van Strien, A.; Sol,

[20764] C. J. A.; Ter Schegget, J.; van der Noordaa, J.: Morphologi-cal transformation

[20765] by early region human polyomavirus BK DNA of human fi-broblasts with

[20766] deletions in the short arm of one chromosome 11. J. Gen. Virol. 69:

[20767] 467-471, 1988.

[20768] 2280.

[20769] Seilhamer, J. J.; Randall, T. L.; Johnson, L. K.; Heinzmann, C.;

[20770] Klisak, I.; Sparkes, R. S.; Lusis, A. J.: Novel gene exon homologous
[20771] to pancreatic phospholipase A(2): sequence and chromosomal mapping
[20772] of both human genes. J. Cell. Biochem. 39: 327-337, 1989.
[20773] 2281.

[20774] Cormier, R. T.; Hong, K. H.; Halberg, R. B.; Hawkins, T. L.; Richardson,
[20775] P.; Mulherkar, R.; Dove, W. F.; Lander, E. S.: Secretory phospholipase
[20776] Pla2g2a confers resistance to intestinal tumorigenesis. Nature Genet. 17:
[20777] 88-91, 1997.

[20778] 2282.

[20779] Dennis, E. A.: Diversity of group types, regulation, and function
[20780] of phospholipase A2. J. Biol. Chem. 269: 13057-13060, 1994.
[20781] 2283.

[20782] Dietrich, W. F.; Lander, E. S.; Smith, J. S.; Moser, A. R.; Gould,

[20783] K. A.; Luongo, C.; Borenstein, N.; Dove, W.: Genetic identi-
fication
[20784] of Mom-1, a major modifier locus affecting Min-induced
intestinal
[20785] neoplasia in the mouse. Cell 75: 631-639, 1993.

[20786] 2284.

[20787] Dobbie, Z.; Heinimann, K.; Bishop, D. T.; Muller, H.; Scott,
R.
[20788] J.: Identification of a modifier gene locus on chromosome
1p35-36
[20789] in familial adenomatous polyposis. Hum. Genet. 99:
653-657, 1997.
[20790] 2285.

[20791] Dragani, T. A.; Manenti, G.: Mom1 leads the pack. Nature
Genet. 17:
[20792] 7-8, 1997.

[20793] 2286.

[20794] Haapamaki, M. M.; Gronroos, J. M.; Nurmi, H.; Alanen, K.;
Kallajoki,
[20795] M.; Nevalainen, T. J.: Gene expression of group II phos-
pholipase

[20796] A2 in intestine in ulcerative colitis. Gut 40: 95-101, 1997.

[20797] 2287.

[20798] Johnson, L. K.; Frank, S.; Vades, P.; Pruzanski, W.; Lusis, A.

[20799] J.; Seilhamer, J. J.: Localization and evolution of two hu-
man phospholipase

[20800] A2 genes and two related genetic elements.In: Wong, P.
Y.-K.; Dennis,

[20801] E. A.: Phospholipase A2. New York: Plenum Press (pub.)
1990.

[20802] Pp. 17-34.

[20803] 2288.

[20804] Johnson, L. K.; Seilhamer, J. J.; Frank, S.; Lusis, A.; Vadas,

[20805] P.; Pruzanski, W.: Synovial fluid phospholipase A(2): chro-
mosomal

[20806] co-localization with homologous genes may provide dis-
ease-related

[20807] PLA(2) diversity. (Abstract) Arthritis Rheum. 33 (suppl.):
S79,

[20808] 1990.

[20809] 2289.

[20810] MacPhee, M.; Chepenik, K. P.; Liddell, R. A.; Nelson, K. K.; Siracusa,
[20811] L. D.; Buchberg, A. M.: The secretory phospholipase A2 gene is a
[20812] candidate for the Mom1 locus, a major modifier of Apc(Min)-induced
[20813] intestinal neoplasia. Cell 81: 957-966, 1995.

[20814] 2290.

[20815] Masharani, U.; Coleman, R. T.; Johnson, L. K.; Seilhamer, J. J.
[20816] : EcoRI and NsiI RFLPs at a human PLA2 gene on chromosome 1. Nucleic
[20817] Acids Res. 16: 9073, 1988.

[20818] 2291.

[20819] Moser, A. R.; Dove, W. F.; Roth, K. A.; Gordon, J. I.: The Min
[20820] (multiple intestinal neoplasia) mutation: its effect on gut epithelial
[20821] cell differentiation and interaction with a modifier system. J. Cell
[20822] Biol. 116: 1517-1526, 1992.

[20823] 2292.

[20824] Moser, A. R.; Mattes, E. M.; Dove, W. F.; Lindstrom, M. J.; Haag,

[20825] J. D.; Gould, M. N.: Apc(Min), a mutation in the murine Apc gene,

[20826] predisposes to mammary carcinomas and focal alveolar hyperplasias. Proc.

[20827] Nat. Acad. Sci. 90: 8977-8981, 1993.

[20828] 2293.

[20829] Moser, A. R.; Pitot, H. C.; Dove, W. F.: A dominant muta-tion

[20830] that predisposes to multiple intestinal neoplasia in the mouse. Science 247:

[20831] 322-324, 1990.

[20832] 2294.

[20833] Nadeau, J. H.: Modifier genes in mice and humans. Nature Rev. 2:

[20834] 165-174, 2001.

[20835] 2295.

[20836] Nimmrich, I.; Friedl, W.; Kruse, R.; Pietsch, S.; Hentsch, S.;

[20837] Deuter, R.; Winde, G.; Muller, O.: Loss of the PLA2G2A gene in a
[20838] sporadic colorectal tumor of a patient with a PLA2G2A germline mutation
[20839] and absence of PLA2G2A germline alterations in patients with FAP. Hum.
[20840] Genet. 100: 345-349, 1997.

[20841] 2296.

[20842] Seilhamer, J. J.; Plant, S.; Pruzanski, W.; Schilling, J.; Ste-
fanski,
[20843] E.; Vadas, P.; Johnson, L. K.: Multiple forms of phospholi-
pase A(2)
[20844] in arthritic synovial fluid. J. Biochem. 106: 38-42, 1989.

[20845] 2297.

[20846] Seilhamer, J. J.; Pruzanski, W.; Vadas, P.; Plant, S.; Miller,
[20847] J. A.; Kloss, J.; Johnson, L. K.: Cloning and recombinant expression
[20848] of phospholipase A(2) present in rheumatoid arthritic syn-
ovial fluid. J.
[20849] Biol. Chem. 264: 5335-5338, 1989.

[20850] 2298.

[20851] Seilhamer, J. J.; Randall, T. L.; Johnson, L. K.; Lusis, A.; Sparkes,

[20852] R. S.; Heinzman, C.: Chromosomal mapping of human pancreatic PLA2

[20853] and a homologous PLA2 exon. (Abstract) J. Cell. Biochem. (Suppl.

[20854] 12E): 55 only, 1988.

[20855] 2299.

[20856] Su, L.-K.; Kinzler, K. W.; Vogelstein, B.; Preisinger, A. C.;

[20857] Moser, A. R.; Luongo, C.; Gould, K. A.; Dove, W. F.: Multi-ple intestinal

[20858] neoplasia caused by a mutation in the murine homolog of the APC gene. Science 256:

[20859] 668-670, 1992.

[20860] 2300.

[20861] Bristol, A.; Hall, S. M.; Kriz, R. W.; Stahl, M. L.; Fan, Y. S.;

[20862] Byers, M. G.; Eddy, R. L.; Shows, T. B.; Knopf, J. L.: Phos-pholipase

[20863] C-148: chromosomal location and deletion mapping of functional domains. Cold

[20864] Spring Harbor Symp. Quant. Biol. 53: 915-920, 1988.

[20865] 2301.

[20866] Nelson, K. K.; Knopf, J. L.; Siracusa, L. D.: Localization of

[20867] phospholipase C-gamma 1 to mouse chromosome 2.
Mammalian Genome 3:

[20868] 597-600, 1992.

[20869] 2302.

[20870] Rothschild, C. B.; Akots, G.; Fajans, S. S.; Bowden, D. W.: A

[20871] microsatellite polymorphism associated with the PLC1
(phospholipase

[20872] C) locus: identification, mapping, and linkage to the
MODY locus on

[20873] chromosome 20. Genomics 13: 560-564, 1992.

[20874] 2303.

[20875] Stahl, M. L.; Ferenz, C. R.; Kelleher, K. L.; Kriz, R. W.;
Knopf,

[20876] J. L.: Sequence similarity of phospholipase C with the non-
catalytic

[20877] region of src. Nature 332: 269-272, 1988.

[20878] 2304.

[20879] Phillips, S. A.; Barr, V. A.; Haft, D. H.; Taylor, S. I.; Haft,

[20880] C. R.: Identification and characterization of SNX15, a novel sorting
[20881] nexin involved in protein trafficking. J. Biol. Chem. 276: 5074-5084,
[20882] 2001.

[20883] 2305.

[20884] Roberts, W. M.; Look, A. T.; Ruossel, M. F.; Sherr, C. J.: Tandem
[20885] linkage of human CSF-1 receptor (c-fms) and PDGF re-ceptor genes. Cell 55:
[20886] 655-661, 1988.

[20887] 2306.

[20888] Smith, E. A.; Seldin, M. F.; Martinez, L.; Watson, M. L.; Ghosh
[20889] Choudhury, G.; Lalley, P. A.; Pierce, J.; Aaronson, S.; Barker, J.;
[20890] Naylor, S. L.; Sakaguchi, A. Y.: Mouse platelet-derived growth factor
[20891] receptor alpha gene is deleted in W-19H and patch muta-tions on chromosome
[20892] 5. Proc. Nat. Acad. Sci. 88: 4811-4815, 1991.

[20893] 2307.

[20894] Stenman, G.; Eriksson, A.; Claesson-Welsh, L.: Human PDGFA receptor

[20895] gene maps to the same region on chromosome 4 as the KIT oncogene. Genes

[20896] Chromosomes Cancer 1: 155-158, 1989.

[20897] 2308.

[20898] Stephenson, D. A.; Mercola, M.; Anderson, E.; Wang, C.; Stiles,

[20899] C. D.; Bowen-Pope, D. F.; Chapman, V. M.: Platelet-de-rived growth

[20900] factor receptor alpha-subunit gene (Pdgfra) is deleted in the mouse

[20901] patch (Ph) mutation. Proc. Nat. Acad. Sci. 88: 6-10, 1991.

[20902] 2309.

[20903] Xie, J.; Aszterbaum, M.; Zhang, X.; Bonifas, J. M.; Zachary, C.;

[20904] Epstein, E.; McCormick, F.: A role of PDGFR-alpha in basal cell carcinoma

[20905] proliferation. Proc. Nat. Acad. Sci. 98: 9255-9259, 2001.

[20906] 2310.

[20907] Aitman, T. J.; Cooper, L. D.; Norsworthy, P. J.; Wahid, F. N.;

[20908] Gray, J. K.; Curtis, B. R.; McKeigue, P. M.; Kwiatkowski, D.; Greenwood,

[20909] B. M.; Snow, R. W.; Hill, A. V.; Scott, J.: Malaria susceptibil-ity

[20910] and CD36 mutation. Nature 405: 1015-1016, 2000.

[20911] 2311.

[20912] Aitman, T. J.; Glazier, A. M.; Wallace, C. A.; Cooper, L. D.; Norsworthy,

[20913] P. J.; Wahid, F. N.; Al-Majali, K. M.; Trembling, P. M.; Mann, C.

[20914] J.; Shoulders, C. C.; Graf, D.; St. Lezin, E.; Kurtz, T. W.; Kren,

[20915] V.; Pravenec, M.; Ibrahimi, A.; Abumrad, N. A.; Stanton, L. W.; Scott,

[20916] J.: Identification of Cd36 (Fat) as an insulin-resistance gene causing

[20917] defective fatty acid and glucose metabolism in hyperten-sive rats. Nature

[20918] Genet. 21: 76-83, 1999.

[20919] 2312.

[20920] Aitman, T. J.; Godsland, I. F.; Farren, B.; Crook, D.; Wong, H.

[20921] J.; Scott, J.: Defects of insulin action on fatty acid and carbohydrate

[20922] metabolism in familial combined hyperlipidemia. Arterioscler. Thromb.

[20923] Vasc. Biol. 17: 748-754, 1997.

[20924] 2313.

[20925] Armesilla, A. L.; Vega, M. A.: Structural organization of the

[20926] gene for human CD36 glycoprotein. J. Biol. Chem. 269: 18985-18991,

[20927] 1994.

[20928] 2314.

[20929] Coburn, C. T.; Knapp, F. F., Jr.; Febbraio, M.; Beets, A. L.; Silverstein,

[20930] R. L.; Abumrad, N. A.: Defective uptake and utilization of long chain

[20931] fatty acids in muscle and adipose tissues of CD36 knock-out mice. J.

[20932] Biol. Chem. 275: 32523-32529, 2000.

[20933] 2315.

[20934] Curtis, B. R.; Aster, R. H.: Incidence of the Nak(a)-negative

[20935] platelet phenotype in African Americans is similar to that of Asians. Transfusion 36:

[20936] 331-334, 1996.

[20937] 2316.

[20938] Fernandez-Ruiz, E.; Armesilla, A. L.; Sanchez-Madrid, F.; Vega,

[20939] M. A.: Gene encoding the collagen type I and thrombospondin receptor

[20940] CD36 is located on chromosome 7q11.2. Genomics 17: 759-761, 1993.

[20941] 2317.

[20942] Gotoda, T.; Iizuka, Y.; Kato, N.; Osuga, J.; Bihoreau, M.-T.; Murakami,

[20943] T.; Yamori, Y.; Shimano, H.; Ishibashi, S.; Yamada, N.: Absence of

[20944] Cd36 mutation in the original spontaneously hypertensive rats with

[20945] insulin resistance. (Letter) Nature Genet. 22: 226-228, 1999.

[20946] 2318.

[20947] Greenwalt, D. E.; Lipsky, R. H.; Ockenhouse, C. F.; Ikeda, H.;
[20948] Tandon, N. N.; Jamieson, G. A.: Membrane glycoprotein CD36: a review
[20949] of its roles in adherence, signal transduction, and transfusion medicine. Blood 80:
[20950] 1105-1115, 1992.

[20951] 2319.

[20952] Griffin, E.; Re, A.; Hamel, N.; Fu, C.; Bush, H.; McCaffrey, T.;
[20953] Asch, A. S.: A link between diabetes and atherosclerosis: glucose
[20954] regulates expression of CD36 at the level of translation. Nature
[20955] Med. 7: 840-846, 2001.

[20956] 2320.

[20957] Groop, L. C.; Bonadonna, R. C.; DelPrato, S.; Ratheiser, K.; Zyck,
[20958] K.; Ferrannini, E.; DeFronzo, R. A.: Glucose and free fatty acid
[20959] metabolism in non-insulin-dependent diabetes mellitus: evidence for

[20960] multiple sites of insulin resistance. J. Clin. Invest. 84: 205-213,

[20961] 1989.

[20962] 2321.

[20963] Berube, D.; Luu The, V.; Lachance, Y.; Gagne, R.; Labrie, F.:

[20964] Assignment of the human 3 beta-hydroxysteroid dehydrogenase gene (HSDB3)

[20965] to the p13 band of chromosome 1. Cytogenet. Cell Genet. 52: 199-200,

[20966] 1989.

[20967] 2322.

[20968] Nekrep, N.; Jabrane-Ferrat, N.; Wolf, H. M.; Eibl, M. M.; Geyer,

[20969] M.; Peterlin, B. M.: Mutation in a winged-helix DNA-binding motif

[20970] causes atypical bare lymphocyte syndrome. Nature Immun. 30Sept,

[20971] 2002. Note: Advance Electronic Publication.

[20972] 2323.

[20973] Spencer, S.; Dowbenko, D.; Cheng, J.; Li, W.; Brush, J.; Utzig,

[20974] S.; Simanis, V.; Lasky, L. A.: PSTPIP: a tyrosine phosphorylated
[20975] cleavage furrow-associated protein that is a substrate for PEST tyrosine
[20976] phosphatase. J. Cell Biol. 138: 845-860, 1997.

[20977] 2324.

[20978] Trakhtenbrot, L.; Kelman, Z.; Rotter, V.; Haran-Ghera, N.: Chromosomal
[20979] mapping of the murine c-abl proto-oncogene by in situ hybridization. Leukemia 4:
[20980] 136-137, 1990.

[20981] 2325.

[20982] Tybulewicz, V. L. J.; Crawford, C. E.; Jackson, P. K.; Bronson,
[20983] R. T.; Mulligan, R. C.: Neonatal lethality and lymphopenia in mice
[20984] with a homozygous disruption of the c-abl proto-oncogene. Cell 65:
[20985] 1153-1163, 1991.

[20986] 2326.

[20987] Welch, P. J.; Wang, J. Y. J.: A C-terminal protein-binding

domain
[20988] in the retinoblastoma protein regulates nuclear c-abl tyrosine kinase
[20989] in the cell cycle. Cell 75: 779-790, 1993.

[20990] 2327.

[20991] Westbrook, C. A.; Le Beau, M. M.; Diaz, M. O.; Groffen, J.; Rowley,
[20992] J. D.: Chromosomal localization and characterization of c-abl in
[20993] the t(6;9) of acute nonlymphocytic leukemia. Proc. Nat. Acad. Sci. 82:
[20994] 8742-8746, 1985.

[20995] 2328.

[20996] Yuan, Z.-M.; Shioya, H.; Ishiko, T.; Sun, X.; Gu, J.; Huang, Y.
[20997] Y.; Lu, H.; Kharbanda, S.; Weichselbaum, R.; Kufe, D.: p73 is regulated
[20998] by tyrosine kinase c-Abl in the apoptotic response to DNA damage. Nature 399:
[20999] 814-817, 1999.

[21000] 2329.

[21001] Barletta, C.; Pelicci, P.-G.; Kenyon, L. C.; Smith, S. D.; Dalla-Favera,

[21002] R.: Relationship between the c-myb locus and the 6q- chromosomal

[21003] aberration in leukemias and lymphomas. Science 235: 1064-1067, 1987.

[21004] 2330.

[21005] Dalla-Favera, R.; Franchini, G.; Martinotti, S.; Wong-Staal, F.;

[21006] Gallo, R. C.; Croce, C. M.: Chromosomal assignment of the human homologues

[21007] of feline sarcoma virus and avian myeloblastosis virus onc genes. Proc.

[21008] Nat. Acad. Sci. 79: 4714-4717, 1982.

[21009] 2331.

[21010] Franchini, G.; Wong-Staal, F.; Baluda, M. A.; Lengel, C.; Tronick,

[21011] S. R.: Structural organization and expression of human DNA sequences

[21012] related to the transforming gene of avian myeloblastosis virus. Proc.

[21013] Nat. Acad. Sci. 80: 7385-7389, 1983.

[21014] 2332.

[21015] Harper, M. E.; Franchini, G.; Love, J.; Simon, M. I.; Gallo, R.

[21016] C.; Wong-Staal, F.: Chromosomal sublocalization of human c-myb and

[21017] c-fes cellular onc genes. Nature 304: 169-171, 1983.

[21018] 2333.

[21019] Hawkins, B. R.; Dawkins, R. L.; Hockey, A.; Houliston, J. B.; Kirk,

[21020] R. L.: Evidence for linkage between HLA and malignant melanoma. Tissue

[21021] Antigens 17: 540-541, 1981.

[21022] 2334.

[21023] Janssen, J. W. G.; Vernole, P.; de Boer, P. A. J.; Oosterhuis,

[21024] J. W.; Collard, J. G.: Sublocalization of c-myb to 6q21-q23 by in

[21025] situ hybridization and c-myb expression in a human teratocarcinoma

[21026] with 6q rearrangements. Cytogenet. Cell Genet. 41: 129-135, 1986.

[21027] 2335.

[21028] Majello, B.; Kenyon, L. C.; Dalla-Favera, R.: Human c-myb protooncogene:

[21029] nucleotide sequence of cDNA and organization of the genomic locus. Proc.

[21030] Nat. Acad. Sci. 83: 9636-9640, 1986.

[21031] 2336.

[21032] Manak, J. R.; Mitiku, N.; Lipsick, J. S.: Mutation of the Drosophila

[21033] homologue of the Myb protooncogene causes genomic instability. Proc.

[21034] Nat. Acad. Sci. 99: 7438-7443, 2002.

[21035] 2337.

[21036] McBride, O. W.; Swan, D. C.; Tronick, S. R.; Gol, R.; Klimanis,

[21037] D.; Moore, D. E.; Aaronson, S. A.: Regional chromosomal localization

[21038] of N-ras, K-ras-1, K-ras-2 and myb oncogenes in human cells. Nucleic

[21039] Acids Res. 11: 8221-8236, 1983.

[21040] 2338.

[21041] Meese, E.; Meltzer, P. S.; Witkowski, C. M.; Trent, J. M.:

Molecular
[21042] mapping of the oncogene MYB and rearrangements in malignant melanoma. Genes
[21043] Chromosomes Cancer 1: 88-94, 1989.

[21044] 2339.

[21045] Pelicci, P.-G.; Lanfrancone, L.; Brathwaite, M. D.; Wolman, S.
[21046] R.; Dalla-Favera, R.: Amplification of the c-myb oncogene in a case
[21047] of human acute myelogenous leukemia. Science 224: 1117-1121, 1984.
[21048] 2340.

[21049] Ratajczak, M. Z.; Kant, J. A.; Luger, S. M.; Hijiya, N.; Zhang,

[21050] J.; Zon, G.; Gewirtz, A. M.: In vivo treatment of human leukemia
[21051] in a scid mouse model with c-myb antisense oligodeoxynucleotides. Proc.
[21052] Nat. Acad. Sci. 89: 11823-11827, 1992.

[21053] 2341.

[21054] Rushlow, K. E.; Lautenberger, J. A.; Papas, T. S.; Baluda, M.

[21055] A.; Perbal, B.; Chirikjian, J. G.; Reddy, E. P.: Nucleotide se-

quence

[21056] of the transforming gene of avian myeloblastosis virus. Science 216:

[21057] 1421-1423, 1982.

[21058] 2342.

[21059] Westin, E. H.; Gallo, R. C.; Arya, S. K.; Eva, A.; Souza, L. M.;

[21060] Baluda, M. A.; Aaronson, S. A.; Wong-Staal, F.: Differential expression

[21061] of the amv gene in human hematopoietic cells. Proc. Nat. Acad. Sci. 79:

[21062] 2194-2198, 1982.

[21063] 2343.

[21064] Yokota, J.; Tsunetsugu-Yokota, Y.; Battifora, H.; Le Fevre, C.;

[21065] Cline, M. J.: Alterations of myc, myb, and ras(Ha) proto-oncogenes

[21066] in cancers are frequent and show clinical correlation. Science 231:

[21067] 261-265, 1986.

[21068] 2344.

[21069] Asada-Senju, M.; Maeda, T.; Sakata, T.; Hayashi, A.;

Suzuki, T.

[21070] : Molecular analysis of the transferrin gene in a patient with hereditary

[21071] hypotransferrinemia. J. Hum. Genet. 47: 355-359, 2002.

[21072] 2345.

[21073] Baranov, V. S.; Schwartzman, A. L.; Gorbunova, V. N.; Gaitskhoki,

[21074] V. S.; Rubtsov, N. B.; Timchenko, N. A.; Neifakh, S. A.: Chromosomal

[21075] localization of ceruloplasmin and transferrin genes in lab-oratory

[21076] rats, mice and in man by hybridization with specific DNA probes. Chromosoma 96:

[21077] 60-66, 1987.

[21078] 2346.

[21079] Beutler, E.; Gelbart, T.; Lee, P.; Trevino, R.; Fernandez, M. A.;

[21080] Fairbanks, V. F.: Molecular characterization of a case of atransferrinemia. Blood 96:

[21081] 4071-4074, 2000.

[21082] 2347.

[21083] Ahmed, Z. M.; Riazuddin, S.; Bernstein, S. L.; Ahmed, Z.; Khan,
[21084] S.; Griffith, A. J.; Morell, R. J.; Friedman, T. B.; Riazuddin, S.;
[21085] Wilcox, E. R.: Mutations of the protocadherin gene PCDH15 cause Usher
[21086] syndrome type 1F. Am. J. Hum. Genet. 69: 25-34, 2001.

[21087] 2348.

[21088] Falkenberg, M.; Tom, C.; DeYoung, M. B.; Wen, S.; Linne-
mann, R.;
[21089] Dichek, D. A.: Increased expression of urokinase during atherosclerotic
[21090] lesion development causes arterial constriction and lumen loss, and
[21091] accelerates lesion growth. Proc. Nat. Acad. Sci. 99: 10665-10670,
[21092] 2002.

[21093] 2349.

[21094] Lijnen, H. R.; Van Hoef, B.; Nelles, L.; Holmes, W. E.; Collen,
[21095] D.: Enzymatic properties of single-chain and two-chain forms of a

[21096] lys(158)-to-glu(158) mutant of urokinase-type plasminogen activator. Europ.

[21097] J. Biochem. 172: 185-188, 1988.

[21098] 2350.

[21099] Nagai, M.; Hiramatsu, R.; Kaneda, T.; Hayasuke, N.; Arimura, H.;

[21100] Nishida, M.; Suyama, T.: Molecular cloning of cDNA coding for human

[21101] preprourokinase. Gene 36: 183-188, 1985.

[21102] 2351.

[21103] Nelles, L.; Lijnen, H. R.; Collen, D.; Holmes, W. E.: Characterization

[21104] of recombinant human single chain urokinase-type plasminogen activator

[21105] mutants produced by site-specific mutagenesis of lysine 158. J. Biol.

[21106] Chem. 262: 5682-5689, 1987.

[21107] 2352.

[21108] Riccio, A.; Grimaldi, G.; Verde, P.; Sebastio, G.; Boast, S.; Blasi,

[21109] F.: The human urokinase-plasminogen activator gene and

its promoter. Nucleic
[21110] Acids Res. 13: 1759-2771, 1985.

[21111] 2353.

[21112] Salerno, G.; Verde, P.; Nolli, M. L.; Corti, A.; Szots, H.; Meo,
[21113] T.; Johnson, J.; Bullock, S.; Cassani, G.; Blasi, F.: Mono-clonal
[21114] antibodies to human urokinase identify the single-chain pro-urokinase
[21115] precursor. Proc. Nat. Acad. Sci. 81: 110-114, 1984.

[21116] 2354.

[21117] Tripputi, P.; Blasi, F.; Verde, P.; Cannizzaro, L. A.; Emanuel,
[21118] B. S.; Croce, C. M.: Human urokinase gene is located on the long
[21119] arm of chromosome 10. Proc. Nat. Acad. Sci. 82: 4448-4452, 1985.
[21120] 2355.

[21121] Deng, Z.; Johnson, K.; Engleward, B. P.; Lane, S.; Callen, D. F.;
[21122] Samson, L. D.; Davisson, M. T.; Siciliano, M. J.: New re-

gions of
[21123] conserved synteny and linkage between human chromosome 16p12-p13 and
[21124] mouse chromosomes 16 and 11. (Abstract) Cytogenet. Cell Genet. 68:
[21125] 180 only, 1995.

[21126] 2356.

[21127] Fukuoka, S.-I.; Matsuda, Y.: Assignment of the Tamm-Horsfall protein/uromodulin
[21128] gene (Umod) to mouse chromosome bands 7F1-F2 and rat chromosome bands
[21129] 1q36-q37 by in situ hybridization. Cytogenet. Cell Genet. 79: 241-242,
[21130] 1997.

[21131] 2357.

[21132] Jeanpierre, C.; Whitmore, S. A.; Austruy, E.; Cohen-Salmon, M.;
[21133] Callen, D. F.; Junien, C.: Chromosomal assignment of the uromodulin
[21134] gene (UMOD) to 16p13.11. Cytogenet. Cell Genet. 62: 185-187, 1993.
[21135] 2358.

[21136] Muchmore, A. V.; Decker, J. M.: Uromodulin: a unique 85-kilodalton
[21137] immunosuppressive glycoprotein isolated from urine of pregnant women. Science 229:
[21138] 479-481, 1985.

[21139] 2359.

[21140] Pennica, D.; Kohr, W. J.; Kuang, W.-J.; Glaister, D.; Aggar-wal,
[21141] B. B.; Chen, E. Y.; Goeddel, D. V.: Identification of human uromodulin
[21142] as the Tamm-Horsfall urinary glycoprotein. Science 236: 83-88, 1987.
[21143] 2360.

[21144] Pook, M. A.; Jeremiah, S.; Scheinman, S. J.; Povey, S.; Thakker,
[21145] R. V.: Localization of the Tamm-Horsfall glycoprotein (uromodulin)
[21146] gene to chromosome 16p12.3-16p13.11. Ann. Hum. Genet. 57: 285-290,
[21147] 1993.

[21148] 2361.

[21149] Tamm, I.; Horsfall, F. L., Jr.: A mucoprotein derived from human
[21150] urine which reacts with influenza, mumps, and Newcastle disease viruses. J.
[21151] Exp. Med. 95: 71-97, 1952.

[21152] 2362.

[21153] Hoffman, H. M.; Mueller, J. L.; Broide, D. H.; Wanderer, A. A.;
[21154] Kolodner, R. D.: Mutation of a new gene encoding a putative pyrin-like
[21155] protein causes familial cold autoinflammatory syndrome and Muckle-Wells
[21156] syndrome. Nature Genet. 29: 301-305, 2001.

[21157] 2363.

[21158] Davies, P. O.; Poirier, C.; Deltour, L.; Montagutelli, X.: Genetic
[21159] reassignment of the insulin-1 (Ins1) gene to distal mouse chromosome
[21160] 19. Genomics 21: 665-667, 1994.

[21161] 2364.

[21162] Dayhoff, M. O.: Atlas of Protein Sequence and Structure.

Proinsulin.

[21163] Washington: National Biomedical Research Foundation (pub.) 5: 1972.

[21164] Pp. D208 only.

[21165] 2365.

[21166] Deltour, L.; Leduque, P.; Blume, N.; Madsen, O.; Dubois, P.; Jami,

[21167] J.; Bucchini, D.: Differential expression of the two nonal-lelic proinsulin

[21168] genes in the developing mouse embryo. Proc. Nat. Acad. Sci. 90:

[21169] 527-531, 1993.

[21170] 2366.

[21171] Elbein, S. C.; Gruppuso, P.; Schwartz, R.; Skolnick, M.; Per-mutt,

[21172] M. A.: Hyperproinsulinemia in a family with a proposed defect in

[21173] conversion is linked to the insulin gene. Diabetes 34: 821-824,

[21174] 1985.

[21175] 2367.

[21176] Gabbay, K. H.: The insulinopathies. (Editorial) New Eng. J.

[21177] Med. 302: 165-167, 1980.

[21178] 2368.

[21179] Gabbay, K. H.; Bergenstal, R. M.; Wolff, J.; Mako, M. E.; Rubenstein,

[21180] A. H.: Familial hyperproinsulinemia: partial characterization of

[21181] circulating proinsulin-like material. Proc. Nat. Acad. Sci. 76:

[21182] 2881-2885, 1979.

[21183] 2369.

[21184] Gabbay, K. H.; DeLuca, K.; Fisher, J. N., Jr.; Mako, M. E.; Rubenstein,

[21185] A. H.: Familial hyperproinsulinemia: an autosomal dominant defect. New

[21186] Eng. J. Med. 294: 911-915, 1976.

[21187] 2370.

[21188] Given, B. D.; Mako, M. E.; Tager, H. S.; Baldwin, D.; Markese,

[21189] J.; Rubenstein, A. H.; Olefsky, J.; Kobayashi, M.; Kolterman, O.;

[21190] Poucher, R.: Diabetes due to secretion of an abnormal in-
sulin. New
[21191] Eng. J. Med. 302: 129-135, 1980.

[21192] 2371.

[21193] Gruppuso, P. A.; Gorden, P.; Kahn, C. R.; Cornblath, M.;
Zeller,
[21194] W. P.; Schwartz, R.: Familial hyperproinsulinemia due to a
proposed
[21195] defect in conversion of proinsulin to insulin. New Eng. J.
Med. 311:
[21196] 629-634, 1984.

[21197] 2372.

[21198] Haneda, M.; Kobayashi, M.; Maegawa, H.; Shigeta, Y.: Low
frequency
[21199] of the large insertion in the human insulin gene in
Japanese. Diabetes 35:
[21200] 115-118, 1986.

[21201] 2373.

[21202] Haneda, M.; Polonsky, K. S.; Bergenstal, R. M.; Jaspan, J. B.;

[21203] Shoelson, S. E.; Blix, P. M.; Chan, S. J.; Kwok, S. C. M.;
Wishner,

[21204] W. B.; Zeidler, A.; Olefsky, J. M.; Freidenberg, G.; Tager, H. S.;

[21205] Steiner, D. F.; Rubenstein, A. H.: Familial hyperinsulinemia due

[21206] to a structurally abnormal insulin: definition of an emerging new

[21207] clinical syndrome. New Eng. J. Med. 310: 1288-1294, 1984.

[21208] 2374.

[21209] Harper, M. E.; Saunders, G. F.: Chromosomal localization of human

[21210] insulin gene, placental lactogen-growth hormone genes, and other single

[21211] copy genes by in situ hybridization. (Abstract) Am. J. Hum. Genet. 33:

[21212] 105A, 1981.

[21213] 2375.

[21214] Harper, M. E.; Ullrich, A.; Saunders, G. F.: Localization of

[21215] the human insulin gene to the distal end of the short arm of chromosome

[21216] 11. Proc. Nat. Acad. Sci. 78: 4458-4460, 1981.

[21217] 2376.

[21218] Hua, Q. X.; Shoelson, S. E.; Inouye, K.; Weiss, M. A.: Paradoxical

[21219] structure and function in a mutant human insulin associated with diabetes

[21220] mellitus. Proc. Nat. Acad. Sci. 90: 582-586, 1993.

[21221] 2377.

[21222] Huerre, C.; Gilgenkrantz, S.; Leonard, C.; Pictet, R.; Kaplan,

[21223] J. C.; Junien, C.: Regional assignment of the structural gene for

[21224] insulin to 11p15.1-11p15.5 by deletion mapping. (Abstract) Cytogenet.

[21225] Cell Genet. 37: 495, 1984.

[21226] 2378.

[21227] Ibanez, L.; Ong, K.; Potau, N.; Marcos, M. V.; de Zegher, F.;

[21228] Dunger, D.: Insulin gene variable number of tandem repeat genotype

[21229] and the low birth weight, precocious pubarche, and hyperinsulinism

[21230] sequence. J. Clin. Endocr. Metab. 86: 5788-5793, 2001.

[21231] 2379.

[21232] Jaquet, D.; Gaboriau, A.; Czernichow, P.; Levy-Marchal, C.: Insulin

[21233] resistance early in adulthood in subjects born with intrauterine growth

[21234] retardation. J. Clin. Endocr. Metab. 85: 1401-1406, 2000.

[21235] 2380.

[21236] Kanazawa, Y.; Hayashi, M.; Ikeuchi, M.; Hiramatsu, K.; Kosaka,

[21237] K.: Familial proinsulinemia: a possible cause of abnormal glucose

[21238] tolerance. (Abstract) Europ. J. Clin. Invest. 8: 327, 1978.

[21239] 2381.

[21240] Leibiger, B.; Leibiger, I. B.; Moede, T.; Kemper, S.; Kulkarni,

[21241] R. N.; Kahn, C. R.; de Vargas, L. M.; Berggren, P.-O.: Selective

[21242] insulin signaling through A and B insulin receptors regulates transcription

[21243] of insulin and glucokinase genes in pancreatic beta cells. Molec.

[21244] Cell 7: 559-570, 2001.

[21245] 2382.

[21246] Bucciantini, M.; Giannoni, E.; Chiti, F.; Baroni, F.; Formigli,

[21247] L.; Zurdo, J.; Taddei, N.; Ramponi, G.; Dobson, C. M.; Ste-
fani, M.

[21248] : Inherent toxicity of aggregates implies a common mech-
anism for protein

[21249] misfolding diseases. Nature 416: 507-511, 2002.

[21250] 2383.

[21251] Campion, D.; Dumanchin, C.; Hannequin, D.; Dubois, B.;
Belliard,

[21252] S.; Puel, M.; Thomas-Anterion, C.; Michon, A.; Martin, C.;
Charbonnier,

[21253] F.; Raux, G.; Camuzat, A.; Penet, C.; Mesnage, V.; Mar-
tinez, M.; Clerget-Darpoux,

[21254] F.; Brice, A.; Frebourg, T.: Early-onset autosomal domi-
nant Alzheimer

[21255] disease: prevalence, genetic heterogeneity, and mutation
spectrum. Am.

[21256] J. Hum. Genet. 65: 664-670, 1999.

[21257] 2384.

[21258] DeMattos, R. B.; Bales, K. R.; Cummins, D. J.; Paul, S. M.; Holtzman,

[21259] D. M.: Brain to plasma amyloid-beta efflux: a measure of brain amyloid

[21260] burden in a mouse model of Alzheimer's disease. Science 295: 2264-2267,

[21261] 2002.

[21262] 2385.

[21263] Games, D.; Adams, D.; Alessandrini, R.; Barbour, R.; Berthelette,

[21264] P.; Blackwell, C.; Carr, T.; Clemens, J.; Donaldson, T.; Gillespie,

[21265] F.; Guido, T.; Hagoplan, S.; and 22 others: Alzheimer-type neuropathology

[21266] in transgenic mice overexpressing V717F beta-amyloid precursor protein. Nature 373:

[21267] 523-527, 1995.

[21268] 2386.

[21269] Coon, H.; Hoff, M.; Holik, J.; Delisi, L. E.; Crowe, T.; Freed-man,

[21270] R.; Shields, G.; Boccio, A. M.; Lerman, M.; Gershon, E. S.; Gejman,

[21271] P. V.; Leppert, M.; Byerley, W.: C to T nucleotide substitution in
[21272] codon 713 of amyloid precursor protein gene not found in 86 unrelated
[21273] schizophrenics from multiplex families. Am. J. Med. Genet. 48: 36-39,
[21274] 1993.

[21275] 2387.

[21276] Cras, P.; van Harskamp, F.; Hendriks, L.; Ceuterick, C.; van Duijn,
[21277] C. M.; Stefanko, S. Z.; Hofman, A.; Kros, J. M.; Van Broeckhoven,
[21278] C.; Martin, J. J.: Presenile Alzheimer dementia characterized by
[21279] amyloid angiopathy and large amyloid core type senile plaques in the
[21280] APP 692ala-gly mutation. Acta Neuropath. 96: 253-260, 1998.
[21281] 2388.

[21282] De Jonghe, C.; Esselens, C.; Kumar-Singh, S.; Craessaerts, K.;
[21283] Serneels, S.; Checler, F.; Annaert, W.; Van Broeckhoven, C.;

De Strooper,

[21284] B.: Pathogenic APP mutations near the gamma-secretase cleavage site

[21285] differentially affect A-beta secretion and APP C-terminal fragment

[21286] stability. Hum. Molec. Genet. 10: 1665-1671, 2001.

[21287] 2389.

[21288] De Jonghe, C.; Zehr, C.; Yager, D.; Prada, C.-M.; Younkin, S.;

[21289] Hendriks, L.; Van Broeckhoven, C.; Eckman, C. B.: Flemish and Dutch

[21290] mutations in amyloid beta precursor protein have different effects

[21291] on amyloid beta secretion. Neurobiol. Dis. 5: 281-286, 1998.

[21292] 2390.

[21293] Delabar, J.-M.; Goldgaber, D.; Lamour, Y.; Nicole, A.; Huret,

[21294] J.-L.; de Grouchy, J.; Brown, P.; Gajdusek, D. C.; Sinet, P.-M.:

[21295] Beta-amyloid gene duplication in Alzheimer's disease and karyotypically

[21296] normal Down syndrome. Science 235: 1390-1392, 1987.

[21297] 2391.

[21298] Di Luca, M.; Pastorino, L.; Bianchetti, A.; Perez, J.; Vignolo,

[21299] L. A.; Lenzi, G. L.; Trabucchi, M.; Cattabeni, F.; Padovani, A.:

[21300] Differential level of platelet amyloid beta precursor protein isoforms:

[21301] an early marker for Alzheimer disease. Arch. Neurol. 55: 1195-1200,

[21302] 1998.

[21303] 2392.

[21304] Dodart, J.-C.; Bales, K. R.; Gannon, K. S.; Greene, S. J.; DeMattos,

[21305] R. B.; Mathis, C.; DeLong, C. A.; Wu, S.; Wu, X.; Holtzman, D. M.;

[21306] Paul, S. M.: Immunization reverses memory deficits without reducing

[21307] brain A-beta burden in Alzheimer's disease model. Nature Neurosci. 5:

[21308] 452-457, 2002.

[21309] 2393.

[21310] Eckman, C. B.; Mehta, N. D.; Crook, R.; Perez-tur, J.; Prihar,
[21311] G.; Pfeiffer, E.; Graff-Radford, N.; Hinder, P.; Yager, D.; Zenk,
[21312] B.; Refolo, L. M.; Prada, C. M.; Younkin, S. G.; Hutton, M.; Hardy,
[21313] J.: A new pathogenic mutation in the APP gene (I716V) increases the
[21314] relative proportion of A-beta-42(43). Hum. Molec. Genet. 6: 2087-2089,
[21315] 1997.

[21316] 2394.

[21317] Farlow, M.; Murrell, J.; Ghetti, B.; Unverzagt, F.; Zeldenrust,
[21318] S.; Benson, M.: Clinical characteristics in a kindred with early-onset
[21319] Alzheimer's disease and their linkage to a G-to-T change at position
[21320] 2149 of the amyloid precursor protein gene. Neurology 44: 105-111,
[21321] 1994.

[21322] 2395.

[21323] Fernandez-Madrid, I.; Levy, E.; Marder, K.; Frangione, B.: Codon

[21324] 618 variant of Alzheimer amyloid gene associated with inherited cerebral

[21325] hemorrhage. Ann. Neurol. 30: 730-733, 1991.

[21326] 2396.

[21327] Gervais, F. G.; Xu, D.; Robertson, G. S.; Vaillancourt, J. P.;

[21328] Zhu, Y.; Huang, J.; LeBlanc, A.; Smith, D.; Rigby, M.; Shearman, M.

[21329] S.; Clarke, E. E.; Zheng, H.; Van Der Ploeg, L. H. T.; Ruffolo, S.

[21330] C.; Thornberry, N. A.; Xanthoudakis, S.; Zamboni, R. J.; Roy, S.;

[21331] Nicholson, D. W.: Involvement of caspases in proteolytic cleavage

[21332] of Alzheimer's amyloid-beta precursor protein and amyloidogenic A-beta

[21333] peptide formation. Cell 97: 395-406, 1999.

[21334] 2397.

[21335] Gotz, J.; Chen, F.; van Dorpe, J.; Nitsch, R. M.: Formation of

[21336] neurofibrillary tangles in P301L tau transgenic mice induced by A-beta42
[21337] fibrils. Science 293: 1491-1495, 2001.

[21338] 2398.

[21339] Grabowski, T. J.; Cho, H. S.; Vonsattel, J. P. G.; Rebeck, G.

[21340] W.; Greenberg, S. M.: Novel amyloid precursor protein mutation in
[21341] an Iowa family with dementia and severe cerebral amyloid angiopathy. Ann.
[21342] Neurol. 49: 697-705, 2001.

[21343] 2399.

[21344] Graffagnino, C.; Herbstreith, M. H.; Roses, A. D.; Alberts, M.
[21345] J.: A molecular genetic study of intracerebral hemorrhage. Arch.
[21346] Neurol. 51: 981-984, 1994.

[21347] 2400.

[21348] Haan, J.; Lanser, J. B. K.; Zijderveld, I.; van der Does, I. G.

[21349] F.; Roos, R. A. C.: Dementia in hereditary cerebral hemorrhage with

[21350] amyloidosis--Dutch type. Arch. Neurol. 47: 965-967, 1990.

[21351] 2401.

[21352] Haass, C.; Lemere, C. A.; Capell, A.; Citron, M.; Seubert, P.;

[21353] Schenk, D.; Lannfelt, L.; Selkoe, D. J.: The Swedish muta-

[21354] tion causes early-onset Alzheimer's disease by beta-secretase cleav-

[21355] age within the secretory pathway. Nature Med. 1: 1291-1296, 1995.

[21356] 2402.

[21357] Hendriks, L.; van Duijn, C. M.; Cras, P.; Cruts, M.; Van Hul,

[21358] W.; van Harskamp, F.; Warren, A.; McInnis, M. G.; An-

[21359] tonarakis, S. E.; Martin, J.-J.; Hofman, A.; Van Broeckhoven, C.: Prese-

[21360] nile dementia and cerebral haemorrhage linked to a mutation at codon

[21361] 692 of the beta-amyloid precursor protein gene. Nature Genet. 1: 218-221, 1992.

[21362] 2403.

[21363] Borrow, J.; Shearman, A. M.; Stanton, V. P., Jr.; Becher, R.;

Collins,

[21364] T.; Williams, A. J.; Dube, I.; Katz, F.; Kwong, Y. L.; Morris, C.;

[21365] Ohyashiki, K.; Toyama, K.; Rowley, J.; Housman, D. E.: The t(7;11)(p15;p15)

[21366] translocation in acute myeloid leukaemia fuses the genes for nucleoporin

[21367] NUP98 and class I homeoprotein HOXA9. Nature Genet. 12: 159-167,

[21368] 1996.

[21369] 2404.

[21370] Eisen, A.; Weber, B. L.: Prophylactic mastectomy for women with

[21371] BRCA1 and BRCA2 mutations--facts and controversy. (Editorial) New

[21372] Eng. J. Med. 345: 207-208, 2001.

[21373] 2405.

[21374] Fodor, F. H.; Weston, A.; Bleiweiss, I. J.; McCurdy, L. D.; Walsh,

[21375] M. M.; Tartter, P. I.; Brower, S. T.; Eng, C. M.: Frequency and carrier

[21376] risk associated with common BRCA1 and BRCA2 mutations

in Ashkenazi
[21377] Jewish breast cancer patients. Am. J. Hum. Genet. 63:
45-51, 1998.
[21378] 2406.
[21379] Ford, D.; Easton, D. F.; Stratton, M.; Narod, S.; Goldgar, D.;
[21380] Devilee, P.; Bishop, D. T.; Weber, B.; Lenoir, G.; Chang-
Claude, J.;
[21381] Sobol, H.; Teare, M. D.; and 27 others: Genetic hetero-
geneity and
[21382] penetrance analysis of the BRCA1 and BRCA2 genes in
breast cancer
[21383] families. Am. J. Hum. Genet. 62: 676-689, 1998.
[21384] 2407.
[21385] Ford, J. M.; Hanawalt, P. C.: Li-Fraumeni syndrome fibrob-
lasts
[21386] homozygous for p53 mutations are deficient in global
DNA repair but
[21387] exhibit normal transcription-coupled repair and enhanced
UV resistance. Proc.
[21388] Nat. Acad. Sci. 92: 8876-8880, 1995.
[21389] 2408.

[21390] Ford, J. M.; Hanawalt, P. C.: Expression of wild-type p53 is

[21391] required for efficient global genomic nucleotide excision repair in

[21392] UV-irradiated human fibroblasts. J. Biol. Chem. 272: 28073-28080,

[21393] 1997.

[21394] 2409.

[21395] Berry, R.; Stevens, T. J.; Walter, N. A. R.; Wilcox, A. S.; Rubano,

[21396] T.; Hopkins, J. A.; Weber, J.; Goold, R.; Soares, M. B.; Sikela, J.

[21397] M.: Gene-based sequence-tagged-sites (STSs) as the basis for a human

[21398] gene map. Nature Genet. 10: 415-423, 1995.

[21399] 2410.

[21400] Renouf, S.; Beullens, M.; Wera, S.; Van Eynde, A.; Sikela, J.;

[21401] Stalmans, W.; Bollen, M.: Molecular cloning of a human polypeptide

[21402] related to yeast sds22, a regulator of protein phosphatase-1. FEBS

[21403] Lett. 375: 75-78, 1995.

[21404] 2411.

[21405] Palmiter, R. D.; Cole, T. B.; Quaife, C. J.; Findley, S. D.: ZnT-3,

[21406] a putative transporter of zinc into synaptic vesicles. Proc. Nat.

[21407] Acad. Sci. 93: 14934-14939, 1996.

[21408] 2412.

[21409] Hatzfeld, M.; Kristjansson, G. I.; Plessmann, U.; Weber, K.: Band

[21410] 6 protein, a major constituent of desmosomes from stratified epithelia,

[21411] is a novel member of the armadillo multigene family. J. Cell Sci. 107:

[21412] 2259-2270, 1994.

[21413] 2413.

[21414] Heid, H. W.; Schmidt, A.; Zimbelmann, R.; Schafer, S.; Winter-Simanowski,

[21415] S.; Stumpp, S.; Keith, M.; Figge, U.; Schnolzer, M.; Franke, W. W.

[21416] : Cell type-specific desmosomal plaque proteins of the plakoglobin

[21417] family: plakophilin 1 (band 6 protein). Differentiation 58: 113-131,

[21418] 1994.

[21419] 2414.

[21420] McGrath, J. A.; McMillan, J. R.; Shemanko, C. S.; Runswick, S.

[21421] K.; Leigh, I. M.; Lane, E. B.; Garrod, D. R.; Eady, R. A. J.: Mutations

[21422] in the plakophilin 1 gene result in ectodermal dysplasia/ skin fragility

[21423] syndrome. Nature Genet. 17: 240-244, 1997.

[21424] 2415.

[21425] Schmidt, A.; Langbein, L.; Pratzel, S.; Rode, M.; Rackwitz, H.-R.;

[21426] Franke, W. W.: Plakophilin 3--a novel cell-type-specific desmosomal

[21427] plaque protein. Differentiation 64: 291-306, 1999.

[21428] 2416.

[21429] Schmidt, A.; Langbein, L.; Rode, M.; Pratzel, S.; Zimbel-
mann, R.;

[21430] Franke, W. W.: Plakophilins 1a and 1b: widespread nuclear
proteins

[21431] recruited in specific epithelial cells as desmosomal plaque
components. Cell

[21432] Tissue Res. 290: 481-499, 1997.

[21433] 2417.

[21434] Whittock, N. V.; Haftek, M.; Angoulvant, N.; Wolf, F.; Per-
rot,

[21435] H.; Eady, R. A. J.; McGrath, J. A.: Genomic amplification of
the

[21436] human plakophilin 1 gene and detection of a new muta-
tion in ectodermal

[21437] dysplasia/skin fragility syndrome. J. Invest. Derm. 115:
368-374,

[21438] 2000.

[21439] 2418.

[21440] Rosenberg, H. F.; Dyer, K. D.: Molecular cloning and char-
acterization

[21441] of a novel human ribonuclease (RNase k6): increasing diversity in
[21442] the enlarging ribonuclease gene family. Nucleic Acids Res. 24: 3507-3513,
[21443] 1996.

[21444] 2419.

[21445] Aburatani, H.; Hippo, Y.; Ishida, T.; Takashima, R.; Matsuba, C.;
[21446] Kodama, T.; Takao, M.; Yasui, A.; Yamamoto, K.; Asano, M.; Fukasawa,
[21447] K.; Yoshinari, T.; Inoue, H.; Ohtsuka, E.; Nishimura, S.: Cloning
[21448] and characterization of mammalian 8-hydroxyguanine-specific DNA glycosylase/apurinic,
[21449] apyrimidinic lyase, a functional mutM homologue. Cancer Res. 57:
[21450] 2151-2156, 1997.

[21451] 2420.

[21452] Arai, K.; Morishita, K.; Shinmura, K.; Kohno, T.; Kim, S.-R.; Nohmi,
[21453] T.; Taniwaki, M.; Ohwada, S.; Yokota, J.: Cloning of a human homolog

[21454] of the yeast OGG1 gene that is involved in the repair of oxidative
[21455] DNA damage. Oncogene 14: 2857-2861, 1997.

[21456] 2421.

[21457] Audebert, M.; Chevillard, S.; Levalois, C.; Gyapay, G.; Vieillefond,
[21458] A.; Klijanienko, J.; Vielh, P.; El Naggar, A. K.; Oudard, S.; Boiteux,
[21459] S.; Radicella, J. P.: Alterations of the DNA repair gene OGG1 in
[21460] human clear cell carcinomas of the kidney. Cancer Res. 60: 4740-4744,
[21461] 2000.

[21462] 2422.

[21463] Bjoras, M.; Luna, L.; Johnsen, B.; Hoff, E.; Haug, T.; Rognes,
[21464] T.; Seeberg, E.: Opposite base-dependent reactions of a human base
[21465] excision repair enzyme on DNA containing 7,8-dihydro-8-oxoguanine
[21466] and abasic sites. EMBO J. 16: 6314-6322, 1997.

[21467] 2423.

[21468] Ishida, T.; Hippo, Y.; Nakahori, Y.; Matsushita, I.; Kodama, T.;

[21469] Nishimura, S.; Aburatani, H.: Structure and chromosome location of

[21470] human OGG1. Cytogenet. Cell Genet. 85: 232-236, 1999.

[21471] 2424.

[21472] Kohno, T.; Shinmura, K.; Tosaka, M.; Tani, M.; Kim, S.-R.; Sugimura,

[21473] H.; Nohmi, T.; Kasai, H.; Yokota, J.: Genetic polymor-

phisms and alternative

[21474] splicing of the hOGG1 gene, that is involved in the repair of 8-hydroxyguanine

[21475] in damaged DNA. Oncogene 16: 3219-3225, 1998.

[21476] 2425.

[21477] Kuo, F. C.; Sklar, J.: Augmented expression of a human gene for

[21478] 8-oxoguanine DNA glycosylase (MutM) in B lymphocytes of the dark zone

[21479] in lymph node germinal centers. J. Exp. Med. 186: 1547-1556, 1997.

[21480] 2426.

[21481] Lu, R.; Nash, H. M.; Verdine, G. L.: A mammalian DNA re-
pair enzyme
[21482] that excises oxidatively damaged guanines maps to a lo-
cus frequently
[21483] lost in lung cancer. Curr. Biol. 7: 397-407, 1997.

[21484] 2427.

[21485] Radicella, J. P.; Dherin, C.; Desmaze, C.; Fox, M. S.; Boi-
teux,
[21486] S.: Cloning and characterization of hOGG1, a human ho-
molog of the
[21487] OGG1 gene of Saccharomyces cerevisiae. Proc. Nat. Acad.
Sci. 94:
[21488] 8010-8015, 1997.

[21489] 2428.

[21490] Guo, S.; Yamaguchi, Y.; Schilbach, S.; Wada, T.; Lee, J.;
Goddard,
[21491] A.; French, D.; Handa, H.; Rosenthal, A.: A regulator of
transcriptional
[21492] elongation controls vertebrate neuronal development. Na-
ture 408:

[21493] 366-369, 2000.

[21494] 2429.

[21495] DiLeone, R. J.; King, J. A.; Storm, E. E.; Copeland, N. G.; Jenkins,

[21496] N. A.; Kingsley, D. M.: The Bmp8 gene is expressed in de-veloping

[21497] skeletal tissue and maps near the achondroplasia locus on mouse chromosome

[21498] 4. Genomics 40: 196-198, 1997.

[21499] 2430.

[21500] Burtis, K. C.; Baker, B. S.: Drosophila doublesex gene con-trols

[21501] somatic sexual differentiation by producing alternatively spliced

[21502] mRNAs encoding related sex-specific polypeptides. Cell 56: 997-1010,

[21503] 1989.

[21504] 2431.

[21505] Moog-Lutz, C.; Bouillet, P.; Regnier, C. H.; Tomasetto, C.; Mattei,

[21506] M. G.; Chenard, M. P.; Anglard, P.; Rio, M. C.; Basset, P.:

Comparative
[21507] expression of the psoriasin (S100A7) and S100C genes in breast carcinoma
[21508] and co-localization to human chromosome 1q21-q22. Int. J. Cancer 63:
[21509] 297-303, 1995.

[21510] 2432.

[21511] Tanaka, M.; Adzuma, K.; Iwami, M.; Yoshimoto, K.; Monden, Y.; Itakura,
[21512] M.: Human calgizzarin: one colorectal cancer-related gene selected
[21513] by a large scale random cDNA sequencing and northern blot analysis. Cancer
[21514] Lett. 89: 195-200, 1995.

[21515] 2433.

[21516] Todoroki, H.; Kobayashi, R.; Watanabe, M.; Minami, H.; Hidaka,
[21517] H.: Purification, characterization, and partial sequence analysis
[21518] of a newly identified EF-hand type 13-kDa Ca(2+)-binding protein from
[21519] smooth muscle and non-muscle tissues. J. Biol. Chem.

266: 18668-18673,
[21520] 1991.

[21521] 2434.

[21522] Watanabe, M.; Ando, Y.; Todoroki, H.; Minami, H.; Hidaka,
H.:
[21523] Molecular cloning and sequencing of a cDNA clone encod-
ing a new calcium
[21524] binding protein, named calgizzarin, from rabbit lung.
Biochem. Biophys.
[21525] Res. Commun. 181: 644-649, 1991.

[21526] 2435.

[21527] Kimura, S. H.; Tsuruga, H.; Yabuta, N.; Endo, Y.; Nojima,
H.:
[21528] Structure, expression, and chromosomal localization of
human GAK. Genomics 44:
[21529] 179-187, 1997.

[21530] 2436.

[21531] Chieffo, C.; Garvey, N.; Gong, W.; Roe, B.; Zhang, G.; Sil-
ver,
[21532] L.; Emanuel, B. S.; Budarf, M. L.: Isolation and characteri-
zation

[21533] of a gene from the DiGeorge chromosomal region homol-
ogous to the mouse
[21534] Tbx1 gene. Genomics 43: 267-277, 1997.

[21535] 2437.

[21536] Bensch, K. W.; Raida, M.; Magert, H. J.; Schulz-Knappe, P.;
Forssmann,
[21537] W. G.: hBD-1: a novel beta-defensin from human plasma.
FEBS Lett. 368:
[21538] 331-335, 1995.

[21539] 2438.

[21540] Stachora, A. A.; Schafer, R. E.; Pohlmeier, M.; Maier, G.;
Ponstingl,
[21541] H.: Human Supt5h protein, a putative modulator of chro-
matin structure,
[21542] is reversibly phosphorylated in mitosis. FEBS Lett. 409:
74-78,
[21543] 1997.

[21544] 2439.

[21545] Lafreniere, R. G.; Kibar, Z.; Rochefort, D. L.; Han, F.-Y.;
Fon,
[21546] E. A.; Dube, M.-P.; Kang, X.; Baird, S.; Korneluk, R. G.;

Rommens,
[21547] J. M.; Rouleau, G. A.: Genomic structure of the human GT334 (EHOC-1)
[21548] gene mapping to 21q22.3. Gene 198: 313-321, 1997.

[21549] 2440.

[21550] Nagamine, K.; Kudoh, J.; Kawasaki, K.; Minoshima, S.; Asakawa,
[21551] S.; Ito, F.; Shimizu, N.: Genomic organization and complete nucleotide
[21552] sequence of the TMEM1 gene on human chromosome 21q22.3. Biochem.
[21553] Biophys. Res. Commun. 235: 185-190, 1997.

[21554] 2441.

[21555] Yamakawa, K.; Mitchell, S.; Hubert, R.; Chen, X.-N.; Colbern, S.;
[21556] Huo, Y.-K.; Gadomski, C.; Kim, U.-J.; Korenberg, J. R.: Isolation
[21557] and characterization of a candidate gene for progressive myoclonus
[21558] epilepsy on 21q22.3. Hum. Molec. Genet. 4: 709-716, 1995.
[21559] 2442.

[21560] Bell, S. M.; Shaw, M.; Jou, Y.-S.; Myers, R. M.; Knowles, M. A.

[21561] : Identification and characterization of the human homologue of SH3BP2,

[21562] an SH3 binding domain protein within a common region of deletion at

[21563] 4p16.3 involved in bladder cancer. Genomics 44: 163-170, 1997.

[21564] 2443.

[21565] Gosset, P.; Ghezala, G. A.; Korn, B.; Yaspo, M.-L.; Poutska, A.;

[21566] Lehrach, H.; Sinet, P.-M.; Creau, N.: A new inward rectifier potassium

[21567] channel gene (KCNJ15) localized on chromosome 21 in the Down syndrome

[21568] chromosome region 1 (DCR1). Genomics 44: 237-241, 1997.

[21569] 2444.

[21570] Williams, J. B.; Lanahan, A. A.: A mammalian delayed-early response

[21571] gene encodes HNP36, a novel, conserved nucleolar protein. Biochem.

[21572] Biophys. Res. Commun. 213: 325-333, 1995.

[21573] 2445.

[21574] Williams, J. B.; Rexer, B.; Sirripurapu, S.; John, S.; Gold-
stein,
[21575] R.; Phillips, J. A., III; Haley, L. L.; Sait, S. N. J.; Shows, T.
[21576] B.; Smith, C. M.; Gerhard, D. S.: The human HNP36 gene is
localized
[21577] to chromosome 11q13 and produces alternative tran-
scripts that are
[21578] not mutated in multiple endocrine neoplasia, type 1 (MEN
I) syndrome. Genomics 42:
[21579] 325-330, 1997.

[21580] 2446.

[21581] Loeffen, J.; van den Heuvel, L.; Smeets, R.; Triepels, R.;
Sengers,
[21582] R.; Trijbels, F.; Smeitink, J.: cDNA sequence and chromo-
somal localization
[21583] of the remaining three human nuclear encoded iron sul-
phur protein
[21584] (IP) subunits of complex I: the human IP fraction is com-
pleted. Biochem.

[21585] Biophys. Res. Commun. 247: 751-758, 1998.

[21586] 2447.

[21587] Procaccio, V.; de Sury, R.; Martinez, P.; Depetris, D.; Rabil-
loud,
[21588] T.; Soularue, P.; Lunardi, J.; Issartel, J.-P.: Mapping to
1q23 of
[21589] the human gene (NDUFS2) encoding the 49-kDa subunit
of the mitochondrial
[21590] respiratory complex I and immunodetection of the mature
protein in
[21591] mitochondria. Mammalian Genome 9: 482-484, 1998.

[21592] 2448.

[21593] Schenker, T.; Lach, C.; Kessler, B.; Calderara, S.; Trueb, B.:

[21594] A novel GTP-binding protein which is selectively repressed
in SV40
[21595] transformed fibroblasts. J. Biol. Chem. 269:
25447-25453, 1994.
[21596] 2449.

[21597] Daigo, Y.; Suzuki, K.; Maruyama, O.; Miyoshi, Y.; Yasuda,
T.; Kabuto,
[21598] T.; Imaoka, S.; Fujiwara, T.; Takahashi, E.; Fujino, M. A.;

Nakamura,

[21599] Y.: Isolation, mapping, and mutation analysis of a human cDNA homologous

[21600] to the doc-1 gene of the Chinese hamster, a candidate tumor suppressor

[21601] for oral cancer. Genes Chromosomes Cancer 20: 204-207, 1997.

[21602] 2450.

[21603] Todd, R.; McBride, J.; Tsuji, T.; Donoff, R. B.; Nagai, M.; Chou,

[21604] M. Y.; Chiang, T.; Wong, D. T. W.: Deleted in oral cancer-1 (doc-1),

[21605] a novel oral tumor suppressor gene. FASEB J. 1362-1370, 1995.

[21606] 2451.

[21607] Yamagata, T.; Tsuru, T.; Momoi, M. Y.; Suwa, K.; Nozaki, Y.; Mukasa,

[21608] T.; Ohashi, H.; Fukushima, Y.; Momoi, T.: Genome organization of

[21609] human 48-kDa oligosaccharyltransferase (DDOST). Genomics 45: 535-540,

[21610] 1997.

[21611] 2452.

[21612] Aral, B.; de Saint Basile, G.; Al-Garawi, S.; Kamoun, P.; Ceballos-Picot,

[21613] I.: Novel nonsense mutation in the hypoxanthine guanine phosphoribosyltransferase

[21614] gene and nonrandom X-inactivation causing Lesch-Nyhan syndrome in

[21615] a female patient. Hum. Mutat. 7: 52-58, 1996.

[21616] 2453.

[21617] Benke, P. J.; Hebert, A.; Herrick, N.: In vitro effects of magnesium

[21618] ions on mutant cells from patients with the Lesch-Nyhan syndrome. New

[21619] Eng. J. Med. 289: 446-450, 1973.

[21620] 2454.

[21621] Benke, P. J.; Herrick, N.; Hebert, A.: Hypoxanthine-guanine phosphoribosyltransferase

[21622] variant associated with accelerated purine synthesis. J. Clin. Invest. 52:

[21623] 2234-2240, 1973.

[21624] 2455.

[21625] Kelley, W. N.; Greene, M. L.; Rosenbloom, F. M.; Henderson, J.
[21626] F.; Seegmiller, J. E.: Hypoxanthine-guanine phosphoribosyltransferase
[21627] deficiency in gout. Ann. Intern. Med. 70: 155-206, 1969.

[21628] 2456.

[21629] Ballabio, A.; Parenti, G.; Tippett, P.; Mondello, C.; Di Maio,
[21630] S.; Tenore, A.; Andria, G.: X-linked ichthyosis, due to steroid sulphatase
[21631] deficiency, associated with Kallmann syndrome (hypogonadotropic hypogonadism
[21632] and anosmia): linkage relationships with Xg and cloned DNA sequences
[21633] from the distal short arm of the X chromosome. Hum. Genet. 72: 237-240,
[21634] 1986.

[21635] 2457.

[21636] Caplen, N. J.; Taylor, J. P.; Statham, V. S.; Tanaka, F.; Fire,
[21637] A.; Morgan, R. A.: Rescue of polyglutamine-mediated cytotoxicity
[21638] by double-stranded RNA-mediated RNA interference.

Hum. Molec. Genet. 11:
[21639] 175-184, 2002.

[21640] 2458.

[21641] La Spada, A. R.; Wilson, E. M.; Lubahn, D. B.; Harding, A. E.;
[21642] Fischbeck, K. H.: Androgen receptor gene mutations in X-linked spinal
[21643] and bulbar muscular atrophy. Nature 352: 77-79, 1991.

[21644] 2459.

[21645] Rosenberg, A. L.; Bergstrom, L.; Troost, B. T.; Bartholomew, B.
[21646] A.: Hyperuricemia and neurologic deficits: a family study. New Eng.
[21647] J. Med. 282: 992-997, 1970.

[21648] 2460.

[21649] Simmonds, H. A.; Webster, D. R.; Wilson, J.; Lingham, S.: An
[21650] X-linked syndrome characterised by hyperuricaemia, deafness, and neurodevelopmental
[21651] abnormalities. Lancet II: 68-70, 1982.

[21652] 2461.

[21653] Sperling, O.; Eliam, G.; Persky-Brosh, S.; De Vries, A.: Accelerated
[21654] erythrocyte 5-phosphoribosyl-1-pyrophosphate synthesis: a familial
[21655] abnormality associated with excessive uric acid production and gout. Biochem.
[21656] Med. 6: 310-316, 1972.

[21657] 2462.

[21658] Sperling, O.; Persky-Brosh, S.; Boer, P.; De Vries, A.: Human
[21659] erythrocyte phosphoribosylpyrophosphate synthetase mutationally altered
[21660] in regulatory properties. Biochem. Med. 7: 389-395, 1973.
[21661] 2463.

[21662] Takeuchi, F.; Hanaoka, F.; Yano, E.; Yamada, M.; Horiuchi, Y.;
[21663] Akaoka, I.: The mode of genetic transmission of a gouty family with
[21664] increased phosphoribosylpyrophosphate synthetase activity. Hum. Genet. 58:
[21665] 322-330, 1981.

[21666] 2464.

[21667] Wada, Y.; Nishimura, Y.; Tanabu, M.; Yoshimura, Y.;
Iinuma, K.;
[21668] Yoshida, T.; Arakawa, T.: Hypouricemic, mentally retarded
infant
[21669] with a defect of 5-phosphoribosyl-1-pyrophosphate syn-
thetase of erythrocytes. Tohoku
[21670] J. Exp. Med. 113: 149-157, 1974.

[21671] 2465.

[21672] Yen, R. C. K.; Adams, W. B.; Lazar, C.; Becker, M. A.: Evi-
dence
[21673] for X-linkage of human phosphoribosylpyrophosphate
synthetase. Proc.
[21674] Nat. Acad. Sci. 75: 482-485, 1978.

[21675] 2466.

[21676] Zoref, E.; De Vries, A.; Sperling, O.: Mutant feedback-re-
sistant
[21677] phosphoribosylpyrophosphate synthetase associated with
purine overproduction
[21678] and gout: phosphoribosylpyrophosphate and purine
metabolism in cultured

[21679] fibroblasts. J. Clin. Invest. 56: 1093-1099, 1975.

[21680] 2467.

[21681] Zoref, E.; De Vries, A.; Sperling, O.: Metabolic cooperation

[21682] between human fibroblasts with normal and with mutant superactive

[21683] phosphoribosylpyrophosphate synthetase. Nature 260: 786-788, 1976.

[21684] 2468.

[21685] Zoref, E.; De Vries, A.; Sperling, O.: Evidence for X-linkage

[21686] of phosphoribosylpyrophosphate synthetase in man: studies with cultured

[21687] fibroblasts from a gouty family with mutant feedback-resistant enzyme. Hum.

[21688] Hered. 27: 73-80, 1977.

[21689] 2469.

[21690] Iizasa, T.; Taira, M.; Shimada, H.; Ishijima, S.; Tatibana, M.

[21691] : Molecular cloning and sequencing of human cDNA for phosphoribosyl

[21692] pyrophosphate synthetase subunit II. FEBS Lett. 244: 47-50, 1989.

[21693] 2470.

[21694] Taira, M.; Ishijima, S.; Kita, K.; Yamada, K.; Iizasa, T.; Tati-
bana,

[21695] M.: Nucleotide and deduced amino acid sequences of two
distinct cDNAs

[21696] for rat phosphoribosylpyrophosphate synthetase. J. Biol.
Chem. 262:

[21697] 14867-14870, 1987.

[21698] 2471.

[21699] Taira, M.; Kudoh, J.; Minoshima, S.; Iizasa, T.; Shimada, H.;
Shimizu,

[21700] Y.; Tatibana, M.; Shimizu, N.: Localization of human phos-
phoribosylpyrophosphate

[21701] synthetase subunit I and II genes (PRPS1 and PRPS2) to
different regions

[21702] of the X chromosome and assignment of two
PRPS1-related genes to autosomes. Somat.

[21703] Cell Molec. Genet. 15: 29-37, 1989.

[21704] 2472.

[21705] Wang, J. C.; Passage, M. B.; Ellison, J.; Becker, M. A.; Yen,
P.

[21706] H.; Shapiro, L. J.; Mohandas, T. K.: Physical mapping of loci in
[21707] the distal half of the short arm of the human X chromosome: implications
[21708] for the spreading of X-chromosome inactivation. Somat. Cell Molec.
[21709] Genet. 18: 195-200, 1992.

[21710] 2473.

[21711] Barnard, P. J.; Derry, J. M. J.; Ryder-Cook, A. S.; Zander, N.

[21712] F.; Kilimann, M. W.: Mapping of the phosphorylase kinase alpha subunit
[21713] gene on the mouse X chromosome. Cytogenet. Cell Genet. 53: 91-94,
[21714] 1990.

[21715] 2474.

[21716] Bruno, C.; Manfredi, G.; Andreu, A. L.; Shanske, S.; Krishna, S.;
[21717] Ilse, W. K.; DiMauro, S.: A splice junction mutation in the alpha-M
[21718] gene of phosphorylase kinase in a patient with myopathy. Biochem.

[21719] Biophys. Res. Commun. 249: 648-651, 1998.

[21720] 2475.

[21721] Clemens, P. R.; Yamamoto, M.; Engel, A. G.: Adult phos-
phorylase
[21722] b kinase deficiency. Ann. Neurol. 28: 529-538, 1990.

[21723] 2476.

[21724] Wehner, M.; Clemens, P. R.; Engel, A. G.; Kilimann, M. W.:
Human
[21725] muscle glycogenosis due to phosphorylase kinase defi-
ciency associated
[21726] with a nonsense mutation in the muscle isoform of the al-
pha subunit. Hum.
[21727] Molec. Genet. 3: 1983-1987, 1994.

[21728] 2477.

[21729] Wilkinson, D. A.; Tonin, P.; Shanske, S.; Lombes, A.; Carl-
son,
[21730] G. M.; DiMauro, S.: Clinical and biochemical features of 10
adult
[21731] patients with muscle phosphorylase kinase deficiency.
Neurology 44:
[21732] 461-466, 1994.

[21733] 2478.

[21734] Sullivan, J. L.; Byron, K. S.; Brewster, F. E.; Baker, S. M.;

[21735] Ochs, H. D.: X-linked lymphoproliferative syndrome: nat-
ural history

[21736] of the immunodeficiency. J. Clin. Invest. 71: 1765-1778,
1983.

[21737] 2479.

[21738] Sullivan, J. L.; Byron, K. S.; Brewster, F. E.; Purtilo, D. T.

[21739] : Deficient natural killer cell activity in X-linked lympho-
proliferative

[21740] syndrome. Science 210: 543-545, 1980.

[21741] 2480.

[21742] Sumazaki, R.; Kanegane, H.; Osaki, M.; Fukushima, T.;
Tsuchida,

[21743] M.; Matsukura, H.; Shinozaki, K.; Kimura, H.; Matsui, A.;
Miyawaki,

[21744] T.: SH2D1A mutations in Japanese males with severe Ep-
stein-Barr virus-associated

[21745] illnesses. Blood 98: 1268-1270, 2001.

[21746] 2481.

[21747] Sumegi, J.; Gross, T. G.; Seemayer, T. A.: The molecular genetics
[21748] of X-linked lymphoproliferative (Duncan's) disease. Cancer J. Sci.
[21749] Am. 5: 57-62, 1999.

[21750] 2482.

[21751] Sumegi, J.; Huang, D.; Lanyi, A.; Davis, J. D.; Seemayer, T. A.;
[21752] Maeda, A.; Klein, G.; Seri, M.; Wakiguchi, H.; Purtilo, D. T.; Gross,
[21753] T. G.: Correlation of mutations of the SH2D1A gene and Epstein-Barr
[21754] virus infection with clinical phenotype and outcome in X-linked lymphoproliferative
[21755] disease. Blood 96: 3118-3125, 2000.

[21756] 2483.

[21757] Xiang, X.; Benson, K. F.; Chada, K.: Mini-mouse: disrup-tion of
[21758] the pygmy locus in a transgenic insertional mutant. Sci-ence 247:
[21759] 967-969, 1990.

[21760] 2484.

[21761] Scriver, C. R.: Vitamin B6 deficiency and dependency in man. Am.

[21762] J. Dis. Child. 113: 109-114, 1967.

[21763] 2485.

[21764] Scriver, C. R.; Hutchison, J. H.: The vitamin B6 deficiency syndrome

[21765] in human infancy: biochemical and clinical observations. Pediatrics 31:

[21766] 240-250, 1963.

[21767] 2486.

[21768] Wolpert, S. M.; Barnes, P. D.: MRI in Pediatric Neuroradiology. Mosby

[21769] Year Book, St. Louis , 1992.

[21770] 2487.

[21771] Matsumoto, M.; Saito, T.; Takasaki, J.; Kamohara, M.; Sugimoto,

[21772] T.; Kobayashi, M.; Tadokoro, M.; Matsumoto, S.; Ohishi, T.; Furuichi,

[21773] K.: An evolutionarily conserved G-protein coupled receptor family,

[21774] SREB, expressed in the central nervous system. Biochem. Biophys.

[21775] Res. Commun. 272: 576-582, 2000.

[21776] 2488.

[21777] Aagaard, L.; Laible, G.; Selenko, P.; Schmid, M.; Dorn, R.; Schotta,

[21778] G.; Kuhfittig, S.; Wolf, A.; Lebersorger, A.; Singh, P. B.; Reuter,

[21779] G.; Jenuwein, T.: Functional mammalian homologues of the Drosophila

[21780] PEV-modifier Su(var)3-9 encode centromere-associated proteins which

[21781] complex with the heterochromatin component M31. EMBO J. 18: 1923-1938,

[21782] 1999.

[21783] 2489.

[21784] Bannister, A. J.; Zegerman, P.; Partridge, J. F.; Miska, E. A.;

[21785] Thomas, J. O.; Allshire, R. C.; Kouzarides, T.: Selective recognition

[21786] of methylated lysine 9 on histone H3 by the HP1 chromo domain. Nature 410:

[21787] 120-124, 2001.

[21788] 2490.

[21789] Fabrizi, G. M.; Rizzuto, R.; Nakase, H.; Mita, S.; Kaden-
bach, B.;

[21790] Schon, E. A.: Sequence of a cDNA specifying subunit VIa of
human

[21791] cytochrome c oxidase. Nucleic Acids Res. 17: 6409 only,
1989.

[21792] 2491.

[21793] Hey, Y.; Hoggard, N.; Burt, E.; James, L. A.; Varley, J. M.:
Assignment

[21794] of COX6A1 to 6p21 and a pseudogene (COX6A1P) to
1p31.1 by in situ

[21795] hybridization and somatic cell hybrids. Cytogenet. Cell
Genet. 77:

[21796] 167-168, 1997.

[21797] 2492.

[21798] Connors, T. D.; Van Raay, T. J.; Petry, L. R.; Klinger, K. W.;

[21799] Landes, G. M.; Burn, T. C.: The cloning of a human ABC
gene (ABC3)

[21800] mapping to chromosome 16p13.3. Genomics 39:

231-234, 1997.

[21801] 2493.

[21802] Klugbauer, N.; Hofmann, F.: Primary structure of a novel ABC transporter

[21803] with a chromosomal localization on the band encoding the multidrug

[21804] resistance-associated protein. FEBS Lett. 391: 61-65, 1996.

[21805] 2494.

[21806] Wu, Y.-C.; Horvitz, H. R.: The C. elegans cell corpse en-gulfment

[21807] gene ced-7 encodes a protein similar to ABC transporters. Cell 93:

[21808] 951-960, 1998.

[21809] 2495.

[21810] Ahn, D.; Kourakis, M. J.; Rohde, L. A.; Silver, L. M.; Ho, R. K.

[21811] : T-box gene tbx5 is essential for formation of the pec-toral limb

[21812] bud. Nature 417: 754-758, 2002.

[21813] 2496.

[21814] Bruneau, B. G.; Nemer, G.; Schmitt, J. P.; Charron, F.; Robitaille,
[21815] L.; Caron, S.; Conner, D. A.; Gessler, M.; Nemer, M.; Seidman, C.
[21816] E.; Seidman, J. G.: A murine model of Holt-Oram syndrome defines
[21817] roles of the T-box transcription factor Tbx5 in cardiogenesis and
[21818] disease. Cell 106: 709-721, 2001.

[21819] 2497.

[21820] Ghosh, T. K.; Packham, E. A.; Bonser, A. J.; Robinson, T. E.; Cross,
[21821] S. J.; Brook, J. D.: Characterization of the TBX5 binding site and
[21822] analysis of mutations that cause Holt-Oram syndrome. Hum. Molec.
[21823] Genet. 10: 1983-1994, 2001.

[21824] 2498.

[21825] Koshiba-Takeuchi, K.; Takeuchi, J. K.; Matsumoto, K.; Momose, T.;
[21826] Uno, K.; Hoepker, V.; Ogura, K.; Takahashi, N.; Nakamura, H.; Yasuda,

[21827] K.; Ogura, T.: Tbx5 and the retinotectum projection. Science 287:

[21828] 134-137, 2000.

[21829] 2499.

[21830] Logan, M.; Tabin, C. J.: Role of Pitx1 upstream of Tbx4 in specification

[21831] of hindlimb identity. Science 283: 1736-1739, 1999.

[21832] 2500.

[21833] McCormick, M. B.; Tamimi, R. M.; Snider, L.; Asakura, A.; Bergstrom,

[21834] D.; Tapscott, S. J.: neuroD2 and neuroD3: distinct expression patterns

[21835] and transcriptional activation potentials within the neuroD gene family. Molec.

[21836] Cell. Biol. 16: 5792-5800, 1996.

[21837] 2501.

[21838] Tamimi, R. M.; Steingrimsson, E.; Montgomery-Dyer, K.; Copeland,

[21839] N. G.; Jenkins, N. A.; Tapscott, S. J.: NEUROD2 and NEUROD3 genes

[21840] map to human chromosomes 17q12 and 5q23-q31 and

mouse chromosomes

[21841] 11 and 13, respectively. Genomics 40: 355-357, 1997.

[21842] 2502.

[21843] Birck, A.; Ahrenkiel, V.; Zeuthen, J.; Hou-Jensen, K.; Guld-berg,

[21844] P.: Mutation and allelic loss of the PTEN/MMAC1 gene in primary and

[21845] metastatic melanoma biopsies. J. Invest. Derm. 114: 277-280, 2000.

[21846] Di Cristofano and Pandolfi (2000) reviewed the multiple roles

[21847] of PTEN in tumor suppression.

[21848] 2503.

[21849] Bonneau, D.; Longy, M.: Mutations of the human PTEN gene. Hum.

[21850] Mutat. 16: 109-122, 2000.

[21851] 2504.

[21852] Cairns, P.; Okami, K.; Halachmi, S.; Halachmi, N.; Esteller, M.;

[21853] Herman, J. G.; Jen, J.; Isaacs, W. B.; Bova, G. S.; Sidransky, D.

[21854] : Frequent inactivation of PTEN/MMAC1 in primary prostate cancer. Cancer

[21855] Res. 57: 4997-5000, 1997.

[21856] 2505.

[21857] Cantley, L. C.; Neel, B. G.: New insights into tumor suppression:

[21858] PTEN suppresses tumor formation by restraining the phosphoinositide

[21859] 3-kinase/AKT pathway. Proc. Nat. Acad. Sci. 96: 4240-4245, 1999.

[21860] 2506.

[21861] Carethers, J. M.; Furnari, F. B.; Zigman, A. F.; Lavine, J. E.;

[21862] Jones, M. C.; Graham, G. E.; Teebi, A. S.; Huang, H.-J. S.;

Ha, H.

[21863] T.; Chauhan, D. P.; Chang, C. L.; Cavenee, W. K.; Boland, C. R.:

[21864] Absence of PTEN/MMAC1 germ-line mutations in sporadic Bannayan-Riley-Ruvalcaba

[21865] syndrome. Cancer. Res. 58: 2724-2726, 1998.

[21866] 2507.

[21867] Celebi, J. T.; Shendrik, I.; Silvers, D. N.; Peacocke, M.:

Identification
[21868] of PTEN mutations in metastatic melanoma specimens. J. Med. Genet. 37:
[21869] 653-657, 2000.

[21870] 2508.

[21871] Dahia, P. L. M.; FitzGerald, M. G.; Zhang, X.; Marsh, D. J.; Zheng,
[21872] Z.; Pietsch, T.; von Deimling, A.; Haluska, F. G.; Haber, D. A.; Eng,
[21873] C.: A highly conserved processed PTEN pseudogene is lo-cated on chromosome
[21874] band 9p21. Oncogene 16: 2403-2406, 1998.

[21875] 2509.

[21876] De Vivo, I.; Gertig, D. M.; Nagase, S.; Hankinson, S. E.; O'Brien,
[21877] R.; Speizer, F. E.; Parsons, R.; Hunter, D. J.: Novel germline mutations
[21878] in the PTEN tumour suppressor gene found in women with multiple cancers. J.
[21879] Med. Genet. 37: 336-341, 2000.

[21880] 2510.

[21881] Di Cristofano, A.; De Acetis, M.; Koff, A.; Cordon-Cardo, C.;

[21882] Pandolfi, P. P.: Pten and p27(KIP1) cooperate in prostate cancer

[21883] tumor suppression in the mouse. Nature Genet. 27: 222-224, 2001.

[21884] 2511.

[21885] Di Cristofano, A.; Kotsi, P.; Peng, Y. F.; Cordon-Cardo, C.; Elkon,

[21886] K. B.; Pandolfi, P. P.: Impaired Fas response and autoimmunity in

[21887] Pten +/- mice. Science 285: 2122-2125, 1999.

[21888] 2512.

[21889] Di Cristofano, A.; Pandolfi, P. P.: The multiple roles of PTEN

[21890] in tumor suppression. Cell 100: 387-390, 2000.

[21891] 2513.

[21892] Di Cristofano, A.; Pesce, B.; Cordon-Cardo, C.; Pandolfi, P. P.

[21893] : Pten is essential for embryonic development and tumour suppression. Nature

[21894] Genet. 19: 348-355, 1998.

[21895] 2514.

[21896] Hirota, T.; Morisaki, T.; Nishiyama, Y.; Marumoto, T.;
Tada, K.;

[21897] Hara, T.; Masuko, N.; Inagaki, M.; Hatakeyama, K.; Saya,
H.: Zyxin,

[21898] a regulator of actin filament assembly, targets the mitotic
apparatus

[21899] by interacting with h-warts/LATS1 tumor suppressor. J.
Cell Biol. 149:

[21900] 1073-1086, 2000.

[21901] 2515.

[21902] Sobek-Klocke, I.; Disque-Kochem, C.; Ronsiek, M.; Klocke,
R.; Jockusch,

[21903] H.; Breuning, A.; Ponstingl, H.; Rojas, K.; Overhauser, J.;
Eichenlaub-Ritter,

[21904] U.: The human gene ZFP161 on 18p11.21-pter encodes a
putative c-myc

[21905] repressor and is homologous to murine Zfp161 (chr 17)
and Zfp161-rs1

[21906] (X chr). Genomics 43: 156-164, 1997. Note: Erratum: Ge-
nomics 45:

[21907] 633 only, 1997.

[21908] 2516.

[21909] Sugiura, K.; Muro, Y.; Nagai, Y.; Kamimoto, T.; Wak-
abayashi, T.;

[21910] Ohashi, M.; Hagiwara, M.: Expression cloning and intracel-
lular localization

[21911] of a human ZF5 homologue. Biochim. Biophys. Acta 1352:
23-26, 1997.

[21912] 2517.

[21913] Engelen, J. J. M.; Esterling, L. E.; Albrechts, J. C. M.; De-
tera-Wadleigh,

[21914] S. D.; van Eys, G. J. J. M.: Assignment of the human gene
for smoothelin

[21915] (SMTN) to chromosome 22q12 by fluorescence in situ hy-
bridization and

[21916] radiation hybrid mapping. Genomics 43: 245-247, 1997.

[21917] 2518.

[21918] Rensen, S.; Merkx, G.; Doevendans, P.; Geurts van Kessel,
A.; van

[21919] Eys, G.: Structure and chromosome location of Smtn, the
mouse smoothelin

[21920] gene. Cytogenet. Cell Genet. 89: 225-229, 2000.

[21921] 2519.

[21922] van der Loop, F. T. L.; Schaart, G.; Timmer, E. D. J.; Ra-
maekers,
[21923] F. C. S.; van Eys, G. J. J. M.: Smoothelin, a novel cytoskele-
tal
[21924] protein specific for smooth muscle cells. J. Cell Biol. 134:
401-411,
[21925] 1996.

[21926] 2520.

[21927] Zucman-Rossi, J.; Legoix, P.; Thomas, G.: Identification of
new
[21928] members of the Gas2 and Ras families in the 22q12 chro-
mosome region. Genomics 38:
[21929] 247-254, 1996.

[21930] 2521.

[21931] Frank, D.; Fortino, W.; Clark, L.; Musalo, R.; Wang, W.;
Saxena,
[21932] A.; Li, C.-M.; Reik, W.; Ludwig, T.; Tycko, B.: Placental
overgrowth
[21933] in mice lacking the imprinted gene Ipl. Proc. Nat. Acad.

Sci. 99:
[21934] 7490-7495, 2002.

[21935] 2522.

[21936] Frank, D.; Mendelsohn, C. L.; Ciccone, E.; Svensson, K.; Ohlsson,
[21937] R.; Tycko, B. :Mammalian Genome 10: 1150-1159, 1999.

[21938] 2523.

[21939] Muller, S.; van den Boom, D.; Zirkel, D.; Koster, H.; Berthold,
[21940] F.; Schwab, M.; Westphal, M.; Zumkeller, W.: Retention of imprinting
[21941] of the human apoptosis-related gene TSSC3 in human brain tumors. Hum.
[21942] Molec. Genet. 9: 757-763, 2000.

[21943] 2524.

[21944] Qian, N.; Frank, D.; O'Keefe, D.; Dao, D.; Zhao, L.; Yuan, L.;
[21945] Wang, Q.; Keating, M.; Walsh, C.; Tycko, B.: The IPL gene on chromosome
[21946] 11p15.5 is imprinted in humans and mice and is similar to TDAG51,

[21947] implicated in Fas expression and apoptosis. Hum. Molec. Genet. 6:

[21948] 2021-2029, 1997.

[21949] 2525.

[21950] Toshima, J.; Ohashi, K.; Okano, I.; Nunoue, K.; Kishioka, M.; Kuma,

[21951] K.; Miyata, T.; Hirai, M.; Baba, T.; Mizuno, K.: Identification and

[21952] characterization of a novel protein kinase, TESK1, specifically expressed

[21953] in testicular germ cells. J. Biol. Chem. 270: 31331-31337, 1995.

[21954] 2526.

[21955] Toshima, J.; Toshima, J. Y.; Suzuki, M.; Noda, T.; Mizuno, K.:

[21956] Cell-type-specific expression of a TESK1 promoter-linked lacZ gene

[21957] in transgenic mice. Biochem. Biophys. Res. Commun. 286: 566-573,

[21958] 2001.

[21959] 2527.

[21960] Garcia-Anoveros, J.; Derfler, B.; Neville-Golden, J.; Hyman, B.

[21961] T.; Corey, D. P.: BNaC1 and BNaC2 constitute a new family of human

[21962] neuronal sodium channels related to degenerins and epithelial sodium

[21963] channels. Proc. Nat. Acad. Sci. 94: 1459-1464, 1997.

[21964] 2528.

[21965] Price, M. P.; Lewin, G. R.; McIlwrath, S. L.; Cheng, C.; Xie, J.;

[21966] Heppenstall, P. A.; Stucky, C. L.; Mannsfeldt, A. G.; Brennan, T.

[21967] J.; Drummond, H. A.; Qiao, J.; Benson, C. J.; Tarr, D. E.; Hrstka,

[21968] R. F.; Yang, B.; Williamson, R. A.; Welsh, M. J.: The mammalian sodium

[21969] channel BNC1 is required for normal touch sensation. Nature 407:

[21970] 1007-1011, 2000.

[21971] 2529.

[21972] Price, M. P.; Snyder, P. M.; Welsh, M. J.: Cloning and expression

[21973] of a novel human brain Na+ channel. J. Biol. Chem. 271: 7879-7882,
[21974] 1996.

[21975] 2530.

[21976] Waldmann, R.; Champigny, G.; Voilley, N.; Lauritzen, I.; Lazdunski,
[21977] M.: The mammalian degenerin MDEG, an amiloride-sensitive cation channel
[21978] activated by mutations causing neurodegeneration in Caenorhabditis
[21979] elegans. J. Biol. Chem. 271: 10433-10436, 1996.

[21980] 2531.

[21981] Waldmann, R.; Voilley, N.; Mattei, M.-G.; Lazdunski, M.: The human
[21982] degenerin MDEG, an amiloride-sensitive neuronal cation channel, is
[21983] localized on chromosome 17q11.2-17q12 close to the microsatellite
[21984] D17S798. Genomics 37: 269-270, 1996.

[21985] 2532.

[21986] Bjursell, C.; Erlandson, A.; Nordling, M.; Nilsson, S.;

Wahlstrom,
[21987] J.; Stibler, H.; Kristiansson, B.; Martinsson, T.: PMM2 mutation
[21988] spectrum, including 10 novel mutations, in a large CDG type 1A family
[21989] material with a focus on Scandinavian families. Hum. Mutat. 16:
[21990] 395-400, 2000.

[21991] 2533.

[21992] Darlington, T. K.; Wager-Smith, K.; Ceriani, M. F.; Staknis, D.;
[21993] Gekakis, N.; Steeves, T. D. L.; Weitz, C. J.; Takahashi, J. S.; Kay,
[21994] S. A.: Closing the circadian loop: CLOCK-induced transcription of
[21995] its own inhibitors per and tim. Science 280: 1599-1603, 1998.
[21996] 2534.

[21997] Gekakis, N.; Staknis, D.; Nguyen, H. B.; Davis, F. C.; Wilsbacher,
[21998] L. D.; King, D. P.; Takahashi, J. S.; Weitz, C. J.: Role of the CLOCK

[21999] protein in the mammalian circadian mechanism. Science 280: 1564-1569,

[22000] 1998.

[22001] 2535.

[22002] Rutter, J.; Reick, M.; Wu, L. C.; McKnight, S. L.: Regulation

[22003] of Clock and NPAS2 DNA binding by the redox state of NAD cofactors. Science 293:

[22004] 510-514, 2001.

[22005] 2536.

[22006] Shearman, L. P.; Sriram, S.; Weaver, D. R.; Maywood, E. S.; Chaves,

[22007] I.; Zheng, B.; Kume, K.; Lee, C. C.; van der Horst, G. T. J.; Hastings,

[22008] M. H.; Reppert, S. M.: Interacting molecular loops in the mammalian

[22009] circadian clock. Science 288: 1013-1019, 2000.

[22010] 2537.

[22011] Odermatt, A.; Taschner, P. E. M.; Scherer, S. W.; Beatty, B.; Khanna,

[22012] V. K.; Cornblath, D. R.; Chaudhry, V.; Yee, W.-C.; Schrank, B.; Karpati,

[22013] G.; Breuning, M. H.; Knoers, N.; MacLennan, D. H.: Characterization

[22014] of the gene encoding human sarcolipin (SLN), a proteolipid associated

[22015] with SERCA1: absence of structural mutations in five patients with

[22016] Brody disease. Genomics 45: 541-553, 1997.

[22017] 2538.

[22018] Bowden, P. E.; Hainey, S. D.; Parker, G.; Jones, D. O.; Zimonjic,

[22019] D.; Popescu, N.; Hodgins, M. B.: Characterization and chromosomal

[22020] localization of human hair-specific keratin genes and comparative

[22021] expression during the hair growth cycle. J. Invest. Derm. 110: 158-164,

[22022] 1998.

[22023] 2539.

[22024] Heid, H. W.; Werner, E.; Franke, W. W.: The complement of native

[22025] alpha-keratin polypeptides of hair-forming cells: a subset of eight

[22026] polypeptides that differ from epithelial cytokeratins. Differentiation 32:

[22027] 101-119, 1986.

[22028] 2540.

[22029] Langbein, L.; Rogers, M. A.; Winter, H.; Silke, P.; Beckhaus, U.;

[22030] Rackwitz, H.-R.; Schweizer, J.: The catalog of human hair keratins.

[22031] I. Expression of the nine type I members in the hair follicle. J.

[22032] Biol. Chem. 274: 19874-19884, 1999.

[22033] 2541.

[22034] Rogers, M. A.; Langbein, L.; Praetzel, S.; Moll, I.; Krieg, T.;

[22035] Winter, H.; Schweizer, J.: Sequences and differential expression

[22036] of three novel human type-II hair keratins. Differentiation 61:

[22037] 187-194, 1997.

[22038] 2542.

[22039] Rogers, M. A.; Nischt, R.; Korge, B.; Krieg, T.; Fink, T. M.; Lichter,

[22040] P.; Winter, H.; Schweizer, J.: Sequence data and chromoso-
mal localization
[22041] of human type I and type II hair keratin genes. Exp. Cell
Res. 220:
[22042] 357-362, 1995.

[22043] 2543.

[22044] Rogers, M. A.; Winter, H.; Wolf, C.; Heck, M.; Schweizer, J.:

[22045] Characterization of a 190-kilobase pair domain of human
type I hair
[22046] keratin genes. J. Biol. Chem. 273: 26683-26691, 1998.

[22047] 2544.

[22048] Kaiser, P.; Seufert, W.; Hofferer, L.; Kofler, B.; Sachsen-
maier,
[22049] C.; Herzog, H.; Jentsch, S.; Schweiger, M.; Schneider, R.:
Human
[22050] ubiquitin-conjugating enzyme homologous to yeast UBC8.
J. Biol. Chem. 269:
[22051] 8797-8802, 1994.

[22052] 2545.

[22053] Giannakudis, J.; Ropke, A.; Kujat, A.; Krajewska-Walasek,
M.; Hughes,

[22054] H.; Fryns, J.-P.; Bankier, A.; Amor, D.; Schlicker, M.; Hansmann,

[22055] I.: Parental mosaicism of JAG1 mutations in families with Alagille

[22056] syndrome. Europ. J. Hum. Genet. 9: 209-216, 2001.

[22057] 2546.

[22058] Gray, G. E.; Mann, R. S.; Mitsiadis, E.; Henrique, D.; Carcangiu,

[22059] M.-L.; Banks, A.; Leiman, J.; Ward, D.; Ish-Horowitz, D.; Artavanis-Tsakonas,

[22060] S.: Human ligands of the Notch receptor. Am. J. Path. 154: 785-794,

[22061] 1999.

[22062] 2547.

[22063] Heritage, M. L.; MacMillan, J. C.; Colliton, R. P.; Genin, A.;

[22064] Spinner, N. B.; Anderson, G. J.: Jagged1 (JAG1) mutation detection

[22065] in an Australian Alagille syndrome population. Hum. Mutat. 16: 408-416,

[22066] 2000.

[22067] 2548.

[22068] Jones, E. A.; Clement-Jones, M.; Wilson, D. I.: JAGGED1 expression

[22069] in human embryos: correlation with the Alagille syndrome phenotype. J.

[22070] Med. Genet. 37: 658-662, 2000.

[22071] 2549.

[22072] Kiernan, A. E.; Ahituv, N.; Fuchs, H.; Balling, R.; Avraham, K.

[22073] B.; Steel, K. P.; Hrabe de Angelis, M.: The Notch ligand Jagged1

[22074] is required for inner ear sensory development. Proc. Nat. Acad. Sci. 98:

[22075] 3873-3878, 2001.

[22076] 2550.

[22077] Krantz, I. D.; Colliton, R. P.; Genin, A.; Rand, E. B.; Li, L.;

[22078] Piccoli, D. A.; Spinner, N. B.: Spectrum and frequency of Jagged1

[22079] (JAG1) mutations in Alagille syndrome patients and their families. Am.

[22080] J. Hum. Genet. 62: 1361-1369, 1998.

[22081] 2551.

[22082] Krantz, I. D.; Smith, R.; Colliton, R. P.; Tinkel, H.; Zackai,

[22083] E. H.; Piccoli, D. A.; Goldmuntz, E.; Spinner, N. B.: Jagged1 mutations

[22084] in patients ascertained with isolated congenital heart de-fects. Am.

[22085] J. Med. Genet. 84: 56-60, 1999.

[22086] 2552.

[22087] Le Caignec, C.; Lefevre, M.; Schott, J. J.; Chaventre, A.; Gayet,

[22088] M.; Calais, C.; Moisan, J. P.: Familial deafness, congenital heart

[22089] defects, and posterior embryotoxon caused by cysteine substitution

[22090] in the first epidermal-growth-factor-like domain of Jagged 1. Am.

[22091] J. Hum. Genet. 71: 180-186, 2002.

[22092] 2553.

[22093] Li, L.; Krantz, I. D.; Deng, Y.; Genin, A.; Banta, A. B.; Collins,

[22094] C. C.; Qi, M.; Trask, B. J.; Kuo, W. L.; Cochran, J.; Costa, T.; Pierpont,

[22095] M. E. M.; Rand, E. B.; Piccoli, D. A.; Hood, L.; Spinner, N. B.:

[22096] Alagille syndrome is caused by mutations in human Jagged1, which encodes

[22097] a ligand for Notch1. Nature Genet. 16: 243-251, 1997.

[22098] 2554.

[22099] Lindsell, C. E.; Shawber, C. J.; Boulter, J.; Weinmaster, G.:

[22100] Jagged: a mammalian ligand that activates Notch1. Cell 80: 909-917,

[22101] 1995.

[22102] 2555.

[22103] Loomes, K. M.; Underkoffler, L. A.; Morabito, J.; Gottlieb, S.;

[22104] Piccoli, D. A.; Spinner, N. B.; Baldwin, H. S.; Oakey, R. J.: The

[22105] expression of Jagged1 in the developing mammalian heart correlates

[22106] with cardiovascular disease in Alagille syndrome. Hum. Molec. Genet. 8:

[22107] 2443-2449, 1999.

[22108] 2556.

[22109] Morrissette, J. J. D.; Colliton, R. P.; Spinner, N. B.: Defective

[22110] intracellular transport and processing of JAG1 missense mutations

[22111] in Alagille syndrome. Hum. Molec. Genet. 10: 405-413, 2001.

[22112] 2557.

[22113] Oda, T.; Elkahloun, A. G.; Meltzer, P. S.; Chan-drasekharappa,

[22114] S. C.: Identification and cloning of the human homolog (JAGL1) of

[22115] the rat Jagged gene from the Alagille syndrome critical re-gion at

[22116] 20p12. Genomics 43: 376-379, 1997.

[22117] 2558.

[22118] Oda, T.; Elkahloun, A. G.; Pike, B. L.; Okajima, K.; Krantz, I.

[22119] D.; Genin, A.; Piccoli, D. A.; Meltzer, P. S.; Spinner, N. B.; Collins,

[22120] F. S.; Chandrasekharappa, S. C.: Mutations in the human Jagged1 gene

[22121] are responsible for Alagille syndrome. Nature Genet. 16: 235-242,

[22122] 1997.

[22123] 2559.

[22124] Raas-Rothschild, A.; Shteyer, E.; Lerer, I.; Nir, A.; Granot,

[22125] E.; Rein, A. J. J. T.: Jagged1 gene mutation for abdominal coarctation

[22126] of the aorta in Alagille syndrome. Am. J. Med. Genet. 112: 75-78,

[22127] 2002.

[22128] 2560.

[22129] Spinner, N. B.; Colliton, R. P.; Crosnier, C.; Krantz, I. D.;

[22130] Hadchouel, M.; Meunier-Rotival, M.: Jagged1 mutations in Alagille

[22131] syndrome. Hum. Mutat. 17: 18-33, 2001.

[22132] 2561.

[22133] Tsai, H.; Hardisty, R. E.; Rhodes, C.; Kiernan, A. E.; Roby, P.;

[22134] Tymowska-Lalanne, Z.; Mburu, P.; Rastan, S.; Hunter, A. J.; Brown,

[22135] S. D. M.; Steel, K. P.: The mouse slalom mutant demonstrates a role

[22136] for Jagged1 in neuroepithelial patterning in the organ of Corti. Hum.

[22137] Molec. Genet. 10: 507-512, 2001.

[22138] 2562.

[22139] Yuan, Z. R.; Okaniwa, M.; Nagata, I.; Tazawa, Y.; Ito, M.; Kawarazaki,

[22140] K.; Inomata, Y.; Okano, S.; Yoshida, T.; Kobayashi, N.; Kohsaka, T.

[22141] : The DSL domain in mutant JAG1 ligand is essential for the severity

[22142] of the liver defect in Alagille syndrome. Clin. Genet. 59: 330-337,

[22143] 2001.

[22144] 2563.

[22145] Penninger, J. M.; Woodgett, J.: PTEN--coupling tumor suppression

[22146] to stem cells? Science 294: 2116-2118, 2001.

[22147] 2564.

[22148] Raizis, A. M.; Ferguson, M. M.; Nicholls, D. T.; Goodisson, D.

[22149] W.; George, P. M.: A novel 5-prime (40^41insA) mutation

in a patient
[22150] with numerous manifestations of Cowden disease. J. In-
vest. Derm. 114:
[22151] 597-598, 2000.

[22152] 2565.

[22153] Ramaswamy, S.; Nakamura, N.; Vazquez, F.; Batt, D. B.;
Perera,
[22154] S.; Roberts, T. M.; Sellers, W. R.: Regulation of G1 pro-
gression
[22155] by the PTEN tumor suppressor protein is linked to inhibi-
tion of the
[22156] phosphatidylinositol 3-kinase/Akt pathway. Proc. Nat.
Acad. Sci. 96:
[22157] 2110-2115, 1999.

[22158] 2566.

[22159] Reardon, W.; Zhou, X.-P.; Eng, C.: A novel germline muta-
tion
[22160] of the PTEN gene in a patient with macrocephaly, ventric-
ular dilatation,
[22161] and features of VATER association. J. Med. Genet. 38:
820-823, 2001.
[22162] 2567.

[22163] Shugart, Y. Y.; Cour, C.; Renard, H.; Lenoir, G.; Goldgar, D.;
[22164] Teare, D.; Easton, D.; Rahman, N.; Gusterton, R.; Seal, S.; Barfoot,
[22165] R.; Stratton, M.; and 10 others: Linkage analysis of 56 multiplex
[22166] families excludes the Cowden disease gene PTEN as a major contributor
[22167] to familial breast cancer. J. Med. Genet. 36: 720-721, 1999.
[22168] 2568.

[22169] Stambolic, V.; MacPherson, D.; Sas, D.; Lin, Y.; Snow, B.; Jang,
[22170] Y.; Benchimol, S.; Mak, T. W.: Regulation of PTEN transcription by
[22171] p53. Molec. Cell 8: 317-325, 2001.

[22172] 2569.

[22173] Stambolic, V.; Suzuki, A.; de la Pompa, J. L.; Brothers, G. M.;
[22174] Mirtsos, C.; Sasaki, T.; Ruland, J.; Penninger, J. M.; Siderovski,
[22175] D. P.; Mak, T. W.: Negative regulation of PKB/

Akt-dependent cell
[22176] survival by the tumor suppressor PTEN. Cell 95: 29-39, 1998.
[22177] 2570.
[22178] Sutphen, R.; Diamond, T. M.; Minton, S. E.; Peacocke, M.; Tsou,
[22179] H. C.; Root, A. W.: Severe Lhermitte-Duclos disease with unique germline
[22180] mutation of PTEN. Am. J. Med. Genet. 82: 290-293, 1999.
[22181] 2571.
[22182] Tamura, M.; Gu, J.; Matsumoto, K.; Aota, S.; Parsons, R.; Yamada,
[22183] K. M.: Inhibition of cell migration, spreading, and focal adhesions
[22184] by tumor suppressor PTEN. Science 280: 1614-1617, 1998.
[22185] 2572.
[22186] Tsou, H. C.; Ping, X. L.; Xie, X. X.; Gruener, A. C.; Zhang, H.;
[22187] Nini, R.; Swisshelm, K.; Sybert, V.; Diamond, T. M.; Sut-phen, R.;
[22188] Peacocke, M.: The genetic basis of Cowden's syndrome:

three novel
[22189] mutations in PTEN/MMAC1/TEP1. Hum. Genet. 102: 467-473, 1998.
[22190] 2573.
[22191] Wen, S.; Stolarov, J.; Myers, M. P.; Su, J. D.; Wigler, M. H.;
[22192] Tonks, N. K.; Durden, D. L.: PTEN controls tumor-induced angiogenesis. Proc.
[22193] Nat. Acad. Sci. 98: 4622-4627, 2001.
[22194] 2574.
[22195] Steck, P. A.; Pershouse, M. A.; Jasser, S. A.; Yung, W. K. A.;
[22196] Lin, H.; Ligon, A. H.; Langford, L. A.; Baumgard, M. L.; Hattier,
[22197] T.; Davis, T.; Frye, C.; Hu, R.; Swedlund, B.; Teng, D. H. F.; Tavtigian,
[22198] S. V.: Identification of a candidate tumour suppressor gene, MMAC1,
[22199] at chromosome 10q23.3 that is mutated in multiple advanced cancers. Nature
[22200] Genet. 15: 356-362, 1997.
[22201] 2575.
[22202] Tsou, H. C.; Teng, D. H.-F.; Ping, X. L.; Brancolini, V.;

Davis,
[22203] T.; Hu, R.; Xie, X. X.; Gruener, A. C.; Schrager, C. A. Christiano,
[22204] A. M.; Eng, C.; Steck, P.; Ott, J.; Tavtigian, S. V.; Peacocke, M.
[22205] : The role of MMAC1 mutations in early-onset breast cancer: causative
[22206] in association with Cowden syndrome and excluded in BRCA1-negative
[22207] cases. Am. J. Hum. Genet. 61: 1036-1043, 1997.

[22208] 2576.

[22209] Weng, L.-P.; Brown, J. L.; Eng, C.: PTEN coordinates G1 arrest
[22210] by down-regulating cyclin D1 via its protein phosphatase activity
[22211] and up-regulating p27 via its lipid phosphatase activity in a breast
[22212] cancer model. Hum. Molec. Genet. 10: 599-604, 2001.

[22213] 2577.

[22214] Weng, L.-P.; Gimm, O.; Kum, J. B.; Smith, W. M.; Zhou, X.-P.;
[22215] Wynford-Thomas, D.; Leone, G.; Eng, C.: Transient ectopic

expression
[22216] of PTEN in thyroid cancer cell lines induces cell cycle arrest and
[22217] cell type-dependent cell death. Hum. Molec. Genet. 10: 251-258,
[22218] 2001.

[22219] 2578.

[22220] Weng, L.-P.; Smith, W. M.; Brown, J. L.; Eng, C.: PTEN inhibits
[22221] insulin-stimulated MEK/MAPK activation and cell growth by blocking
[22222] IRS-1 phosphorylation and IRS-1/Grb-2/Sos complex formation in a breast
[22223] cancer model. Hum. Molec. Genet. 10: 605-616, 2001.

[22224] 2579.

[22225] Weng, L.-P.; Smith, W. M.; Dahia, P. L. M.; Ziebold, U.; Gil,

[22226] E.; Lees, J. A.; Eng, C.: PTEN suppresses breast cancer cell growth
[22227] by phosphatase activity-dependent G1 arrest followed by cell death. Cancer
[22228] Res. 59: 5808-5814, 1999.

[22229] 2580.

[22230] Wishart, M. J.; Taylor, G. S.; Slama, J. T.; Dixon, J. E.: PTEN

[22231] and myotubularin phosphoinositide phosphatases: bring-
ing bioinformatics

[22232] to the lab bench. Cell Biol. 13: 172-181, 2001.

[22233] 2581.

[22234] Zori, R. T.; Marsh, D. J.; Graham, G. E.; Marliss, E. B.; Eng,

[22235] C.: Germline PTEN mutation in a family with Cowden syn-
drome and Bannayan-Riley-Ruvalcaba

[22236] syndrome. Am. J. Med. Genet. 80: 399-402, 1998.

[22237] 2582.

[22238] Zundel, W.; Schindler, C.; Haas-Kogan, D.; Koong, A.;
Kaper, F.;

[22239] Chen, E.; Gottschalk, A. R.; Ryan, H. E.; Johnson, R. S.; Jef-
ferson,

[22240] A. B.; Stokoe, D.; Giaccia, A. J.: Loss of PTEN facilitates
HIF-1-mediated

[22241] gene expression. Genes Dev. 14: 391-396, 2000.

[22242] 2583.

[22243] Jacquemin, P.; Depetris, D.; Mattei, M.-G.; Martial, J. A.;

Davidson,
[22244] I.: Localization of human transcription factor TEF-4 and TEF-5 (TEAD2,
[22245] TEAD3) genes to chromosomes 19q13.3 and 6p21.2 using fluorescence
[22246] in situ hybridization and radiation hybrid analysis. Genomics 55:
[22247] 127-129, 1999.

[22248] 2584.

[22249] Kai, M.; Sakane, F.; Imai, S.; Wada, I.; Kanoh, H.: Molecular

[22250] cloning of a diacylglycerol kinase isozyme predominantly expressed
[22251] in human retina with a truncated and inactive enzyme expression in
[22252] most other human cells. J. Biol. Chem. 269:
18492-18498, 1994.
[22253] 2585.

[22254] Masai, I.; Okazaki, A.; Hosoya, T.; Hotta, Y.: Drosophila retinal
[22255] degeneration A gene encodes an eye-specific diacylglycerol kinase
[22256] with cysteine-rich zinc-finger motifs and ankyrin repeats.

Proc.
[22257] Nat. Acad. Sci. 90: 11157-11161, 1993.

[22258] 2586.

[22259] Stohr, H.; Klein, J.; Gehrig, A.; Koehler, M. R.; Jurklies, B.;

[22260] Kellner, U.; Leo-Kottler, B.; Schmid, M.; Weber, B. H. F.: Mapping
[22261] and genomic characterization of the gene encoding diacylglycerol kinase
[22262] gamma (DAGK3): assessment of its role in dominant optic atrophy (OPA1). Hum.
[22263] Genet. 104: 99-105, 1999.

[22264] 2587.

[22265] Hart, M. J.; Sharma, S.; elMasry, N.; Qiu, R.-G.; McCabe, P.; Polakis,
[22266] P.; Bollag, G.: Identification of a novel guanine nucleotide exchange
[22267] factor for the rho GTPase. J. Biol. Chem. 271: 25452-25458, 1996.

[22268] 2588.

[22269] Gripp, K. W.; Zackai, E. H.; Stolle, C. A.: Mutations in the human

[22270] TWIST gene. Hum. Mutat. 15: 150-155, 2000.

[22271] 2589.

[22272] Valve, R.; Sivenius, K.; Miettinen, R.; Pihlajamaki, J.; Rissa-
nen,
[22273] A.; Deeb, S. S.; Auwerx, J.; Uusitupa, M.; Laakso, M.: Two
polymorphisms
[22274] in the peroxisome proliferator-activated receptor-gamma
gene are associated
[22275] with severe overweight among obese women. J. Clin. En-
docr. Metab. 84:
[22276] 3708-3712, 1999.

[22277] 2590.

[22278] Wang, X. L.; Oosterhof, J.; Duarte, N.: Peroxisome prolifer-
ator-activated
[22279] receptor gamma C161-T polymorphism and coronary
artery disease. Cardiovasc.
[22280] Res. 44: 588-594, 1999.

[22281] 2591.

[22282] Yen, C.-J.; Beamer, B. A.; Negri, C.; Silver, K.; Brown, K. A.;

[22283] Yarnall, D. P.; Burns, D. K.; Roth, J.; Shuldiner, A. R.:
Molecular

[22284] scanning of the human peroxisome proliferator activated receptor gamma

[22285] (hPPAR-gamma) gene in diabetic Caucasians: identification of a pro12ala

[22286] PPAR-gamma-2 missense mutation. Biochem. Biophys. Res. Commun. 241:

[22287] 270-274, 1997.

[22288] 2592.

[22289] Bravo, J.; Karathanassis, D.; Pacold, C. M.; Pacold, M. E.; Ellson,

[22290] C. D.; Anderson, K. E.; Butler, P. J. G.; Lavenir, I.; Perisic, O.;

[22291] Hawkins, P. T.; Stephens, L.; Williams, R. L.: The crystal structure

[22292] of the PX domain from p40-phox bound to phosphatidylinositol 3-phosphate. Molec.

[22293] Cell 829-839, 2001.

[22294] 2593.

[22295] Zhan, S.; Vazquez, N.; Zhan, S.; Wientjes, F. B.; Budarf, M. L.;

[22296] Schrock, E.; Ried, T.; Green, E. D.; Chanock, S. J.: Genomic structure,

[22297] chromosomal localization, start of transcription, and tissue expression
[22298] of the human p40-phox, a new component of the nicotinamide adenine
[22299] dinucleotide phosphate-oxidase complex. Blood 88: 2714-2721, 1996.
[22300] 2594.

[22301] Chan, J. Y.; Han, X.-L.; Kan, Y. W.: Isolation of cDNA encoding
[22302] the human NF-E2 protein. Proc. Nat. Acad. Sci. 90: 11366-11370,
[22303] 1993.

[22304] 2595.

[22305] Peters, L. L.; Andrews, N. C.; Eicher, E. M.; Davidson, M. B.;

[22306] Orkin, S. H.; Lux, S. E.: Mouse microcytic anaemia caused by a defect
[22307] in the gene encoding the globin enhancer-binding protein NF-E2. Nature 362:
[22308] 768-770, 1993.

[22309] 2596.

[22310] Peters, L. L.; Bishop, T. R.; Andrews, N. C.: Globin-en-

hancer
[22311] binding factor NF-E2 is implicated in the regulation of heme biosynthesis
[22312] and iron uptake in mk/mk mice. (Abstract) Blood 82 (suppl. 1): 179a,
[22313] 1993.

[22314] 2597.

[22315] Shivdasani, R. A.; Orkin, S. H.: Erythropoiesis and globin gene
[22316] expression in mice lacking the transcription factor NF-E2. Proc.
[22317] Nat. Acad. Sci. 92: 8690-8694, 1995.

[22318] 2598.

[22319] Shivdasani, R. A.; Rosenblatt, M. F.; Zucker-Franklin, D.; Jackson,
[22320] C. W.; Hunt, P.; Saris, C. J. M.; Orkin, S. H.: Transcription factor
[22321] NF-E2 is required for platelet formation independent of the actions
[22322] of thrombopoietin/MGDF in megakaryocyte development. Cell 81: 695-704,
[22323] 1995.

[22324] 2599.

[22325] Weremowicz, S.; Andrews, N. C.; Orkin, S. H.; Morton, C. C.: Mapping

[22326] the p45 subunit of human NFE2 to 12q13. (Abstract) Human Genome Mapping

[22327] Workshop 93 25, 1993.

[22328] 2600.

[22329] Natowicz, M. R.; Short, M. P.; Wang, Y.; Dickersin, G. R.; Gebhardt,

[22330] M. C.; Rosenthal, D. I.; Sims, K. B.; Rosenberg, A. E.: Clinical

[22331] and biochemical manifestations of hyaluronidase deficiency. New Eng.

[22332] J. Med. 335: 1029-1033, 1996.

[22333] 2601.

[22334] Triggs-Raine, B.; Salo, T. J.; Zhang, H.; Wicklow, B. A.; Natowicz,

[22335] M. R.: Mutations in HYAL1, a member of a tandemly distributed multigene

[22336] family encoding disparate hyaluronidase activities, cause a newly

[22337] described lysosomal disorder, mucopolysaccharidosis IX. Proc. Nat.
[22338] Acad. Sci. 96: 6296-6300, 1999.

[22339] 2602.

[22340] Carlson, H.; Ota, S.; Campbell, C. E.; Hurlin, P. J.: A dominant
[22341] repression domain in Tbx3 mediates transcriptional repression and
[22342] cell immortalization: relevance to mutations in Tbx3 that cause ulnar-mammary
[22343] syndrome. Hum. Molec. Genet. 10: 2403-2413, 2001.

[22344] 2603.

[22345] Bourgeois, P.; Bolcato-Bellemin, A.-L.; Danse, J.-M.; Bloch-Zupan,
[22346] A.; Yoshiba, K.; Stoetzel, C.; Perrin-Schmitt, F.: The variable expressivity
[22347] and incomplete penetrance of the twist-null heterozygous mouse phenotype
[22348] resemble those of human Saethre-Chotzen syndrome. Hum. Molec. Genet. 7:
[22349] 945-957, 1998.

[22350] 2604.

[22351] Bamshad, M.; Le, T.; Watkins, W. S.; Dixon, M. E.; Kramer, B. E.;

[22352] Roeder, A. D.; Carey, J. C.; Root, S.; Schinzel, A.; Van Maldergem,

[22353] L.; Gardner, R. J. M.; Lin, R. C.; Seidman, C. E.; Seidman, J. G.;

[22354] Wallerstein, R.; Moran, E.; Sutphen, R.; Campbell, C. E.; Jorde, L.

[22355] B.: The spectrum of mutations in TBX3: genotype/phe-

notype relationship

[22356] in ulnar-mammary syndrome. Am. J. Hum. Genet. 64: 1550-1562, 1999.

[22357] 2605.

[22358] Hamamori, Y.; Sartorelli, V.; Ogryzko, V.; Puri, P. L.; Wu, H.-Y.;

[22359] Wang, J. Y. J.; Nakatani, Y.; Kedes, L.: Regulation of his-

tone acetyltransferases

[22360] p300 and PCAF by the bHLH protein Twist and adenoviral oncoprotein

[22361] E1A. Cell 96: 405-413, 1999.

[22362] 2606.

[22363] Krebs, I.; Weis, I.; Hudler, M.; Rommens, J. M.; Roth, H.; Scherer,

[22364] S. W.; Tsui, L.-C.; Fuchtbauer, E.-M.; Grzeschik, K.-H.; Tsuji, K.;

[22365] Kunz, J.: Translocation breakpoint maps 5 kb 3-prime from TWIST in

[22366] a patient affected with Saethre-Chotzen syndrome. Hum. Molec. Genet. 6:

[22367] 1079-1086, 1997.

[22368] 2607.

[22369] Shiratori, H.; Sakuma, R.; Watanabe, M.; Hashiguchi, H.; Mochida,

[22370] K.; Sakai, Y.; Nishino, J.; Saijoh, Y.; Whitman, M.; Hamada, H.:

[22371] Two-step regulation of left-right asymmetric expression of Pitx2:

[22372] initiation by Nodal signaling and maintenance by Nkx2. Molec. Cell 7:

[22373] 137-149, 2001.

[22374] 2608.

[22375] Yoshioka, H.; Meno, C.; Koshiba, K.; Sugihara, M.; Itoh, H.; Ishimaru,

[22376] Y.; Inoue, T.; Ohuchi, H.; Semina, E. V.; Murray, J. C.; Hamada, H.;

[22377] Noji, S.: Pitx2, a bicoid-type homeobox gene, is involved in a lefty-signaling

[22378] pathway in determination of left-right asymmetry. Cell 94: 299-305,

[22379] 1998.

[22380] 2609.

[22381] Cahana, A.; Escamez, T.; Nowakowski, R. S.; Hayes, N. L.; Giacobini,

[22382] M.; von Holst, A.; Shmueli, O.; Sapir, T.; McConnell, S. K.; Wurst,

[22383] W.; Martinez, S.; Reiner, O.: Targeted mutagenesis of Lis1 disrupts

[22384] cortical development and LIS1 homodimerization. Proc. Nat. Acad.

[22385] Sci. 98: 6429-6434, 2001.

[22386] 2610.

[22387] Cardoso, C.; Leventer, R. J.; Dowling, J. J.; Ward, H. L.; Chung,

[22388] J.; Petras, K. S.; Roseberry, J. A.; Weiss, A. M.; Das, S.; Martin,

[22389] C. L.; Pilz, D. T.; Dobyns, W. B.; Ledbetter, D. H.: Clinical and
[22390] molecular basis of classical lissencephaly: mutations in the LIS1
[22391] gene (PAFAH1B1). Hum. Mutat. 19: 4-15, 2002.

[22392] 2611.

[22393] Cardoso, C.; Leventer, R. J.; Matsumoto, N.; Kuc, J. A.; Ramocki,
[22394] M. B.; Mewborn, S. K.; Dudlicek, L. L.; May, L. F.; Mills, P. L.;
[22395] Das, S.; Pilz, D. T.; Dobyns, W. B.; Ledbetter, D. H.: The lo-
cation
[22396] and type of mutation predict malformation severity in iso-
lated lissencephaly
[22397] caused by abnormalities within the LIS1 gene. Hum. Molec. Genet. 9:
[22398] 3019-3028, 2000.

[22399] 2612.

[22400] Chong, S. S.; Pack, S. D.; Roschke, A. V.; Tanigami, A.; Carrozzo,
[22401] R.; Smith, A. C. M.; Dobyns, W. B.; Ledbetter, D. H.: A revi-
sion

[22402] of the lissencephaly and Miller-Dieker syndrome critical regions in
[22403] chromosome 17p13.3. Hum. Molec. Genet. 6: 147-155, 1997.
[22404] 2613.

[22405] Faulkner, N. E.; Dujardin, D. L.; Tai, C.-Y.; Vaughan, K. T.; O'Connell,
[22406] C. B.; Wang, Y.; Vallee, R. B.: A role for the lissencephaly gene
[22407] LIS1 in mitosis and cytoplasmic dynein function. Nature Cell Biol. 2:
[22408] 784-791, 2000.

[22409] 2614.

[22410] Fogli, A.; Giglio, S.; Arrigo, G.; Lo Nigro, C.; Zollo, M.; Vig-giano,
[22411] L.; Rocchi, M.; Archidiacono, N.; Zuffardi, O.; Carrozzo, R.: Identification
[22412] of two paralogous regions mapping to the short and long arms of human
[22413] chromosome 2 comprising LIS1 pseudogenes. Cytogenet. Cell Genet. 86:
[22414] 225-232, 1999.

[22415] 2615.

[22416] Grunewald, S.; Schollen, E.; Van Schaftingen, E.; Jaeken, J.; Matthijs,
[22417] G.: High residual activity of PMM2 in patients' fibroblasts: possible
[22418] pitfall in the diagnosis of CDG-Ia (phosphomannomutase deficiency). Am.
[22419] J. Hum. Genet. 68: 347-354, 2001.

[22420] 2616.

[22421] Kjaergaard, S.; Skovby, F.; Schwartz, M.: Carbohydrate-defi-cient
[22422] glycoprotein syndrome type 1A: expression and charac-terisation of
[22423] wild type and mutant PMM2 in E. coli. Europ. J. Hum. Genet. 7: 884-888,
[22424] 1999.

[22425] 2617.

[22426] Kjaergaard, S.; Skovby, F.; Schwartz, M.: Absence of ho-mozygosity
[22427] for predominant mutations in PMM2 in Danish patients with carbohydrate-deficient

[22428] glycoprotein syndrome type 1. Europ. J. Hum. Genet. 6: 331-336,

[22429] 1998.

[22430] 2618.

[22431] Kondo, I.; Mizugishi, K.; Yoneda, Y.; Hashimoto, T.; Kuwa-jima,

[22432] K.; Yuasa, L.; Shigemoto, K.; Kuroda, Y.: Missense muta-tions in phosphomannomutase

[22433] 2 gene in two Japanese families with carbohydrate-defi-cient glycoprotein

[22434] syndrome type 1. Clin. Genet. 55: 50-54, 1999.

[22435] 2619.

[22436] Matthijs, G.; Schollen, E.; Bjursell, C.; Erlandson, A.; Freeze,

[22437] H.; Imtiaz, F.; Kjaergaard, S.; Martinsson, T.; Schwartz, M.; Seta,

[22438] N.; Vuillaumier-Barrot, S.; Westphal, V.; Winchester, B.: Mutations

[22439] in PMM2 that cause congenital disorders of glycosylation, type Ia

[22440] (CDG-Ia). Hum. Mutat. 16: 386-394, 2000.

[22441] 2620.

[22442] Matthijs, G.; Schollen, E.; Van Schaftingen, E.; Cassiman, J.-J.;

[22443] Jaeken, J.: Lack of homozygotes for the most frequent disease allele

[22444] in carbohydrate-deficient glycoprotein syndrome type 1A. Am. J. Hum.

[22445] Genet. 62: 542-550, 1998.

[22446] 2621.

[22447] Schollen, E.; Kjaergaard, S.; Legius, E.; Schwartz, M.;

Matthijs,
[22448] G.: Lack of Hardy-Weinberg equilibrium for the most prevalent PMM2
[22449] mutation in CDG-Ia (congenital disorders of glycosylation type Ia). Europ.
[22450] J. Hum. Genet. 8: 367-371, 2000.

[22451] 2622.

[22452] Schollen, E.; Pardon, E.; Heykants, L.; Renard, J.; Doggett, N.
[22453] A.; Callen, D. F.; Cassiman, J. J.; Matthijs, G.: Comparative analysis
[22454] of the phosphomannomutase genes PMM1, PMM2 and PMM2-psi: the sequence
[22455] variation in the processed pseudogene is a reflection of the mutations
[22456] found in the functional gene. Hum. Molec. Genet. 7: 157-164, 1998.
[22457] 2623.

[22458] Vuillaumier-Barrot, S.; Hetet, G.; Barnier, A.; Dupre, T.; Cuer,
[22459] M.; de Lonlay, P.; Cormier-Daire, V.; Durand, G.; Grand-champ, B.;

[22460] Seta, N.: Identification of four novel PMM2 mutations in congenital
[22461] disorders of glycosylation (CDG) Ia French patients. J. Med. Genet. 37:
[22462] 579-580, 2000.

[22463] 2624.

[22464] Westphal, V.; Kjaergaard, S.; Schollen, E.; Martens, K.; Grunewald,
[22465] S.; Schwartz, M.; Matthijs, G.; Freeze, H. H.: A frequent mild mutation
[22466] in ALG6 may exacerbate the clinical severity of patients with congenital
[22467] disorder of glycosylation Ia (CDG-Ia) caused by phospho-mannomutase
[22468] deficiency. Hum. Molec. Genet. 11: 599-604, 2002.

[22469] 2625.

[22470] Dubrovskaya, V.; Lavigne, A.-C.; Davidson, I.; Acker, J.; Staub,
[22471] A.; Tora, L.: Distinct domains of hTAFII100 are required for functional
[22472] interaction with transcription factor TFIIF-beta (RAP30) and incorporation

[22473] into the TFIID complex. EMBO J. 15: 3702-3712, 1996.

[22474] 2626.

[22475] Dubrovskaya, V.; Mattei, M.-G.; Tora, L.: Localization of the

[22476] gene (TAF2D) encoding the 100-kDa subunit (hTAFII100) of the human

[22477] TFIID complex to chromosome 10 band q24-q25.2. Genomics 36: 556-557,

[22478] 1996.

[22479] 2627.

[22480] Tanese, N.; Saluja, D.; Vassallo, M. F.; Chen, J.-L.; Admon, A.

[22481] : Molecular cloning and analysis of two subunits of the human TFIID

[22482] complex: hTAFII130 and hTAFII100. Proc. Nat. Acad. Sci. 93: 13611-13616,

[22483] 1996.

[22484] 2628.

[22485] Tao, Y.; Guermah, M.; Martinez, E.; Oelgeschlager, T.; Hasegawa,

[22486] S.; Takada, R.; Yamamoto, T.; Horikoshi, M.; Roeder, R. G.:

Specific
[22487] interactions and potential functions of human TAF(II)100. J. Biol.
[22488] Chem. 272: 6714-6721, 1997.

[22489] 2629.

[22490] Charlier, C.; Coppieters, W.; Farnir, F.; Grobet, L.; Leroy, P.

[22491] L.; Michaux, C.; Mni, M.; Schwers, A.; Vanmanshoven, P.; Hanset, R.;
[22492] Georges, M.: The mh gene causing double-muscling in cattle maps to
[22493] bovine chromosome 2. Mammalian Genome 6: 788-792, 1995.
[22494] 2630.

[22495] Ferrell, R. E.; Conte, V.; Lawrence, E. C.; Roth, S. M.; Hag-berg,
[22496] J. M.; Hurley, B. F.: Frequent sequence variation in the hu-man myostatin
[22497] (GDF8) gene as a marker for analysis of muscle-related phenotypes. Genomics 62:
[22498] 203-207, 1999.

[22499] 2631.

[22500] Gonzalez-Cadavid, N. F.; Taylor, W. E.; Yarasheski, K.; Sinha-Hikim,

[22501] I.; Ma, K.; Ezzat, S.; Shen, R.; Lalani, R.; Asa, S.; Mamita, M.;

[22502] Nair, G.; Arver, S.; Bhasin, S.: Organization of the human myostatin

[22503] gene and expression in healthy men and HIV-infected men with muscle

[22504] wasting. Proc. Nat. Acad. Sci. 95: 14938-14943, 1998.

[22505] 2632.

[22506] Grobet, L.; Martin, L. J. R.; Poncelet, D.; Pirottin, D.; Brouwers,

[22507] B.; Riquet, J.; Schoeberlein, A.; Dunner, S.; Menissier, F.; Massabanda,

[22508] J.; Fries, R.; Hanset, R.; Georges, M.: A deletion in the bovine

[22509] myostatin gene causes the double-muscled phenotype in cattle. Nature

[22510] Genet. 17: 71-74, 1997.

[22511] 2633.

[22512] Orstavik, S.; Solberg, R.; Tasken, K.; Nordahl, M.; Altherr, M.

[22513] R.; Hansson, V.; Jahnsen, T.; Sandberg, M.: Molecular cloning, cDNA
[22514] structure, and chromosomal localization of the human type II cGMP-dependent
[22515] protein kinase. Biochem. Biophys. Res. Commun. 220: 759-765, 1996.
[22516] 2634.

[22517] Pfeifer, A.; Aszodi, A.; Seidler, U.; Ruth, P.; Hofmann, F.; Fassler,
[22518] R.: Intestinal secretory defects and dwarfism in mice lacking cGMP-dependent
[22519] protein kinase II. Science 274: 2082-2084, 1996.

[22520] 2635.

[22521] Buckel, A.; Beeson, D.; James, M.; Vincent, A.: Cloning of cDNA
[22522] encoding human rapsyn and mapping of the RAPSN gene locus to chromosome
[22523] 11p11.2-p11.1. Genomics 35: 613-616, 1996.

[22524] 2636.

[22525] Frail, D.; McLaughlin, L.; Mudd, J.; Merlie, J.: Identification

[22526] of the mouse muscle 43,000-Dalton acetylcholine recep-

tor-associated
[22527] protein (RAPsyn) by cDNA cloning. J. Biol. Chem. 263:
15602-15607,
[22528] 1988.

[22529] 2637.

[22530] Ohno, K.; Engel, A. G.; Shen, X.-M.; Selcen, D.; Brengman,
J.;
[22531] Harper, C. M.; Tsujino, A.; Milone, M.: Rapsyn mutations in
humans
[22532] cause endplate acetylcholine-receptor deficiency and
myasthenic syndrome. Am.
[22533] J. Hum. Genet. 70: 875-885, 2002.

[22534] 2638.

[22535] Bradbury, A.; Possenti, R.; Shooter, E. M.; Tirone, F.:
Molecular
[22536] cloning of PC3, a putatively secreted protein whose mRNA
is induced
[22537] by nerve growth factor and depolarization. Proc. Nat.
Acad. Sci. 88:
[22538] 3353-3357, 1991.

[22539] 2639.

[22540] Duriez, C.; Falette, N.; Audoynaud, C.; Moyret-Lalle, C.; Bensaad,
[22541] K.; Courtois, S.; Wang, Q.; Soussi, T.; Puisieux, A.: The human BTG2/TIS21/PC3
[22542] gene: genomic structure, transcriptional regulation and evaluation
[22543] as a candidate tumor suppressor gene. Gene 282: 207-214, 2002.
[22544] 2640.

[22545] Montagnoli, A.; Guardavaccaro, D.; Starace, G.; Tirone, F.: Overexpression
[22546] of the nerve growth factor-inducible PC3 immediate early gene is associated
[22547] with growth inhibition. Cell Growth Differ. 7: 1327-1336, 1996.
[22548] 2641.

[22549] Clarke, E. P.; Sanwal, B. D.: Cloning of a human collagen-binding
[22550] protein, and its homology with rat gp46, chick hsp47 and mouse J6
[22551] proteins. Biochim. Biophys. Acta 1129: 246-248, 1992.

[22552] 2642.

[22553] Ikegawa, S.; Nakamura, Y.: Structure of the gene encoding human
[22554] colligin-2 (CBP2). Gene 194: 301-303, 1997.

[22555] 2643.

[22556] Ikegawa, S.; Sudo, K.; Okui, K.; Nakamura, Y.: Isolation, characterization
[22557] and chromosomal assignment of human colligin-2 gene (CBP2). Cytogenet.
[22558] Cell Genet. 71: 182-186, 1995.

[22559] 2644.

[22560] Delplanque, J.; Vasseur, F.; Durand, E.; Abderrahmani, A.; Dina,
[22561] C.; Waeber, G.; Guy-Grand, B.; Clement, K.; Weill, J.; Boutin, P.;
[22562] Froguel, P.: Mutation screening of the urocortin gene: identification
[22563] of new single nucleotide polymorphisms and association studies with
[22564] obesity in French Caucasians. J. Clin. Endocr. Metab. 87: 867-869,
[22565] 2002.

[22566] 2645.

[22567] Donaldson, C. J.; Sutton, S. W.; Perrin, M. H.; Corrigan, A. Z.;

[22568] Lewis, K. A.; Rivier, J. E.; Vaughan, J. M.; Vale, W. W.: Cloning

[22569] and characterization of human urocortin. Endocrinology 137: 2167-2170,

[22570] 1996.

[22571] 2646.

[22572] Spina, M.; Merlo-Pich, E.; Chan, R. K. W.; Basso, A. M.; Riv-

[22573] ier, J.; Vale, W.; Koob, G. F.: Appetite-suppressing effects of urocortin,

[22574] a CRF-related neuropeptide. Science 273: 1561-1564, 1996.

[22575] 2647.

[22576] Vaughan, J.; Donaldson, C.; Rittencourt, J.; Porrin, M. H.; Lewis,

[22577] K.; Sutton, S.; Chan, R.; Turnbull, A. V.; Lovejoy, D.; Rivier, C.;

[22578] Rivier, J.; Sawchenko, P. E.; Vale, W.: The mammalian neu-

ropeptide

[22579] urocortin, related to fish urotensin I and to corticotropin-releasing

[22580] factor. Nature 378: 287-292, 1995.

[22581] 2648.

[22582] Vetter, D. E.; Li, C.; Zhao, L.; Contarino, A.; Liberman, M. C.;

[22583] Smith, G. W.; Marchuk, Y.; Koob, G. F.; Heinemann, S. F.; Vale, W.;

[22584] Lee, K.-F.: Urocortin-deficient mice show hearing impairment and

[22585] increased anxiety-like behavior. Nature Genet. 31: 363-369, 2002.

[22586] 2649.

[22587] Amit, T.; Bergman, T.; Dastot, F.; Youdim, M. B. H.; Amselem, S.;

[22588] Hochberg, Z.: A membrane-fixed, truncated isoform of the human growth

[22589] hormone receptor. J. Clin. Endocr. Metab. 82: 3813-3817, 1997.

[22590] 2650.

[22591] Amselem, S.; Duquesnoy, P.; Attree, O.; Novelli, G.; Bousnina,

[22592] S.; Postel-Vinay, M.-C.; Goossens, M.: Laron dwarfism and mutations
[22593] of the growth hormone-receptor gene. New Eng. J. Med. 321: 989-995,
[22594] 1989.

[22595] 2651.

[22596] Amselem, S.; Sobrier, M.-L.; Duquesnoy, P.; Rappaport, R.; Postel-Vinay,
[22597] M.-C.; Gourmelen, M.; Dallapiccola, B.; Goossens, M.: Re-current nonsense
[22598] mutations in the growth hormone receptor from patients with Laron
[22599] dwarfism. J. Clin. Invest. 87: 1098-1102, 1991.

[22600] 2652.

[22601] Amselem, S.; Sobrier, M. L.; Duquesnoy, P.; Dallapiccola, B.; Gourmelen,
[22602] M.; Rappaport, R.; Goossens, M.: Recurrent nonsense mu-tations in
[22603] the human growth hormone-receptor gene. (Abstract) Am. J. Hum. Genet. 47
[22604] (suppl.): A107, 1990.

[22605] 2653.

[22606] Ayling, R. M.; Ross, R.; Towner, P.; Von Laue, S.; Finidori, J.;

[22607] Moutoussamy, S.; Buchanan, C. R.; Clayton, P. E.; Norman, M. R.:

[22608] A dominant-negative mutation of the growth hormone receptor causes

[22609] familial short stature. (Letter) Nature Genet. 16: 13-14, 1997.

[22610] 2654.

[22611] Berg, M. A.; Guevara-Aguirre, J.; Rosenbloom, A. L.; Rosenfeld,

[22612] R. G.; Francke, U.: Mutation creating a new splice site in the growth

[22613] hormone receptor genes of 37 Ecuadorean patients with Laron syndrome. Hum.

[22614] Mutat. 1: 24-34, 1992.

[22615] 2655.

[22616] Duquesnoy, P.; Sobrier, M. L.; Amselem, S.; Goossens, M.: Defective

[22617] membrane expression of human growth hormone (GH) receptor causes Laron-type

[22618] GH insensitivity syndrome. Proc. Nat. Acad. Sci. 88: 10272-10276,
[22619] 1991.

[22620] 2656.

[22621] Edery, M.; Rozakis-Adcock, M.; Goujon, L.; Finidori, J.; Levi-Meyrueis,
[22622] C.; Paly, J.; Djiane, J.; Postel-Vinay, M.-C.; Kelly, P. A.: Lack
[22623] of hormone binding in COS-7 cells expressing a mutated growth hormone
[22624] receptor found in Laron dwarfism. J. Clin. Invest. 91: 838-844,
[22625] 1993.

[22626] 2657.

[22627] Furukawa, K.; Soejima, H.; Niikawa, N.; Shiku, H.; Furukawa, K.
[22628] : Genomic organization and chromosomal assignment of the human beta-1,4-N-acetylgalactosaminyltransferase
[22629] gene. J. Biol. Chem. 271: 20836-20844, 1996.

[22630] 2658.

[22631] Hamlin, P. J.; Jones, P. F.; Leek, J. P.; Bransfield, K.; Lench,

[22632] N. J.; Aldersley, M. A.; Howdle, P. D.; Markham, A. F.; Robinson,

[22633] P. A.: Assignment of GALGT encoding beta-1,4N-acetylgalactosaminyl-transferase

[22634] (GalNAc-T) and KIF5A encoding neuronal kinesin (D12S1889) to human

[22635] chromosome band 12q13 by assignment to ICI YAC 26EG10 and in situ

[22636] hybridization. Cytogenet. Cell Genet. 82: 267-268, 1998.

[22637] 2659.

[22638] Nagata, Y.; Yamashiro, S.; Yodoi, J.; Lloyd, K. O.; Shiku, H.;

[22639] Furukawa, K.: Expression cloning of beta-1,4-N-acetylgalactosaminyltransferase

[22640] cDNAs that determine the expression of G(M2) and G(D2) gangliosides. J.

[22641] Biol. Chem. 267: 12082-12089, 1992.

[22642] 2660.

[22643] Kosaki, K.; Bassi, M. T.; Kosaki, R.; Lewin, M.; Belmont, J.; Schauer,

[22644] G.; Casey, B.: Characterization and mutation analysis of human LEFTY

[22645] A and LEFTY B, homologues of murine genes implicated in left-right

[22646] axis development. Am. J. Hum. Genet. 64: 712-721, 1999.

[22647] 2661.

[22648] Kothapalli, R.; Buyuksal, I.; Wu, S.-Q.; Chegini, N.; Tabibzadeh,

[22649] S.: Detection of ebaf, a novel human gene of the trans-forming growth

[22650] factor beta superfamily: association of gene expression with endometrial

[22651] bleeding. J. Clin. Invest. 99: 2342-2350, 1997.

[22652] 2662.

[22653] Meno, C.; Itoh, Y.; Saijoh, Y.; Matsuda, Y.; Tashiro, K.; Kuhara,

[22654] S.; Hamada, H.: Two closely-related left-right asymmetri-cally expressed

[22655] genes, lefty-1 and lefty-2: their distinct expression do-mains, chromosomal

[22656] linkage and direct neuralizing activity in Xenopus em-bryos. Genes

[22657] Cells 2: 513-524, 1997.

[22658] 2663.

[22659] Meno, C.; Saijoh, Y.; Fujii, H.; Ikeda, M.; Yokoyama, T.; Yokoyama,

[22660] M.; Toyoda, Y.; Hamada, H.: Left-right asymmetric expression of the

[22661] TGF-beta-family member lefty in mouse embryos. Nature 381: 151-155,

[22662] 1996.

[22663] 2664.

[22664] Meno, C.; Shimono, A.; Saijoh, Y.; Yashiro, K.; Mochida, K.; Ohishi,

[22665] S.; Noji, S.; Kondoh, H.; Hamada, H.: Lefty-1 is required for left-right

[22666] determination as a regulator of lefty-2 and nodal. Cell 94: 287-297,

[22667] 1998.

[22668] 2665.

[22669] Tabibzadeh, S.; Mason, J. M.; Shea, W.; Cai, Y.; Murray, M. J.;

[22670] Lessey, B.: Dysregulated expression of ebaf, a novel molecular defect

[22671] in the endometria of patients with infertility. J. Clin. Endocr.

[22672] Metab. 85: 2526-2536, 2000.

[22673] 2666.

[22674] Caterina, M. J.; Leffler, A.; Malmberg, A. B.; Martin, W. J.; Trafton,

[22675] J.; Petersen-Zeltz, K. R.; Koltzenburg, M.; Basbaum, A. I.; Julius,

[22676] D.: Impaired nociception and pain sensation in mice lacking the capsaicin

[22677] receptor. Science 288: 306-313, 2000.

[22678] 2667.

[22679] Caterina, M. J.; Schumacher, M. A.; Tominaga, M.; Rosen, T. A.;

[22680] Levine, J. D.; Julius, D.: The capsaicin receptor: a heat-activated

[22681] ion channel in the pain pathway. Nature 389: 816-824, 1997.

[22682] 2668.

[22683] Jordt, S.-E.; Julius, D.: Molecular basis for species-specific

[22684] sensitivity to 'hot' chili peppers. Cell 108: 421-430, 2002.

[22685] 2669.

[22686] Liedtke, W.; Choe, Y.; Marti-Renom, M. A.; Bell, A. M.; De-
nis,
[22687] C. S.; Sali, A.; Hudspeth, A. J.; Friedman, J. M.; Heller, S.:
Vanilloid
[22688] receptor-related osmotically activated channel (VR-OAC),
a candidate
[22689] vertebrate osmoreceptor. Cell 103: 525-535, 2000.

[22690] 2670.

[22691] Liu, L.; Simon, S. A.: Similarities and differences in the
currents
[22692] activated by capsaicin, piperine, and zingerone in rat
trigeminal
[22693] ganglion cells. J. Neurophysiol. 76: 1858-1869, 1996.

[22694] 2671.

[22695] Prescott, J.; Stevenson, R. J.: Psychophysical responses to
single
[22696] and multiple presentations of the oral irritant zingerone:
relationship
[22697] to frequency of chili consumption. Physiol. Behav. 60:
617-624,

[22698] 1996.

[22699] 2672.

[22700] Stevenson, R. J.; Prescott, J.: The effects of prior experi-
ence

[22701] with capsaicin on ratings of its burn. Chem. Senses 19:
651-656,

[22702] 1994.

[22703] 2673.

[22704] Stevenson, R. J.; Yeomans, M. R.: Differences in ratings of
intensity

[22705] and pleasantness for the capsaicin burn between chili lik-
ers and non-likers:

[22706] implications for liking development. Chem. Senses 18:
471-482, 1993.

[22707] 2674.

[22708] Trevisani, M.; Smart, D.; Gunthorpe, M. J.; Tognetto, M.;
Barbieri,

[22709] M.; Campi, B.; Amadesi, S.; Gray, J.; Jerman, J. C.; Brough,
S. J.;

[22710] Owen, D.; Smith, G. D.; Randall, A. D.; Harrison, S.;
Bianchi, A.;

[22711] Davis, J. B.; Geppetti, P.: Ethanol elicits and potentiates nociceptor
[22712] responses via the vanilloid receptor-1. Nature Neurosci. 5: 546-551,
[22713] 2002.

[22714] 2675.

[22715] de Boer, M.; de Klein, A.; Hossle, J.-P.; Seger, R.; Corbeel, L.;
[22716] Weening, R. S.; Roos, D.: Cytochrome b(558)-negative, autosomal recessive
[22717] chronic granulomatous disease: two new mutations in the cytochrome
[22718] b(558) light chain of the NADPH oxidase (p22-phox). Am. J. Hum. Genet. 51:
[22719] 1127-1135, 1992.

[22720] 2676.

[22721] Dinauer, M. C.; Pierce, E. A.; Bruns, G. A. P.; Curnutte, J. T.;
[22722] Orkin, S. H.: Human neutrophil cytochrome b light chain (p22-phox):
[22723] gene structure, chromosomal location, and mutations in cytochrome-negative

[22724] autosomal recessive chronic granulomatous disease. J. Clin. Invest. 86:
[22725] 1729-1737, 1990.

[22726] 2677.

[22727] Dinauer, M. C.; Pierce, E. A.; Erickson, R. W.; Muhlebach, T. J.;
[22728] Messner, H.; Orkin, S. H.; Seger, R. A.; Curnutte, J. T.: Point mutation
[22729] in the cytoplasmic domain of the neutrophil p22-phox cytochrome b
[22730] subunit is associated with a nonfunctional NADPH oxidase and chronic
[22731] granulomatous disease. Proc. Nat. Acad. Sci. 88: 11231-11235, 1991.
[22732] 2678.

[22733] Fukui, T.; Lassegue, B.; Kai, H.; Alexander, R. W.; Griendling,
[22734] K. K.: Cytochrome b-558 alpha-subunit cloning and expression in rat
[22735] aortic smooth muscle cells. Biochim. Biophys. Acta 1231: 215-219,
[22736] 1995.

[22737] 2679.

[22738] Hamers, M. N.; de Boer, M.; Meerhof, L. J.; Weening, R. S.; Roos,

[22739] D.: Complementation in monocyte hybrids revealing genetic heterogeneity

[22740] in chronic granulomatous disease. Nature 307: 553-555, 1984.

[22741] 2680.

[22742] Inoue, N.; Kawashima, S.; Kanazawa, K.; Yamada, S.; Akita, H.;

[22743] Yokoyama, M.: Polymorphism of the NADH/NADPH oxidase p22 phox gene

[22744] in patients with coronary artery disease. Circulation 97: 135-137,

[22745] 1998.

[22746] 2681.

[22747] Li, A.; Prasad, A.; Mincemoyer, R.; Satorius, C.; Epstein, N.;

[22748] Finkel, T.; Quyyumi, A. A.: Relationship of the C242T p22phox gene

[22749] polymorphism to angiographic coronary artery disease and endothelial

[22750] function. Am. J. Med. Genet. 86: 57-61, 1999.

[22751] 2682.

[22752] Bothwell, T. H.; Cohen, I.; Abrahams, O. L.; Perold, S. M.: A

[22753] familial study in idiopathic hemochromatosis. Am. J. Med. 27: 730-738,

[22754] 1959.

[22755] 2683.

[22756] Brittenham, G. M.; Franks, A. L.; Rickles, F. R.: Research priorities

[22757] in hereditary hemochromatosis. Ann. Intern. Med. 129: 993-996, 1998.

[22758] 2684.

[22759] Brown, A. S.; Gwinn, M.; Cogswell, M. E.; Khoury, M. J.: Hemochromatosis-associated

[22760] morbidity in the United States: an analysis of the National Hospital

[22761] Discharge Survey, 1979-1997. Genet. Med. 3: 109-111, 2001.

[22762] 2685.

[22763] Bulaj, Z. J.; Ajioka, R. S.; Phillips, J. D.; LaSalle, B. A.;

[22764] Jorde, L. B.; Griffen, L. M.; Edwards, C. Q.; Kushner, J. P.: Disease-related

[22765] conditions in relatives of patients with hemochromatosis. New Eng.

[22766] J. Med. 343: 1529-1535, 2000.

[22767] 2686.

[22768] Bulaj, Z. J.; Griffen, L. M.; Jorde, L. B.; Edwards, C. Q.; Kushner,

[22769] J. P.: Clinical and biochemical abnormalities in people heterozygous

[22770] for hemochromatosis. New Eng. J. Med. 335: 1799-1805, 1996.

[22771] 2687.

[22772] Bullen, J. J.; Spalding,P. B.; Ward, C. G.; Gutteridge, J. M.

[22773] : Hemochromatosis, iron and septicemia caused by Vibrio vulnificus. Arch.

[22774] Intern. Med. 151: 1606-1609, 1991.

[22775] 2688.

[22776] Burt, M. J.; George, P. M.; Upton, J. D.; Collett, J. A.; Frampton,

[22777] C. M. A.; Chapman, T. M.; Walmsley, T. A.; Chapman, B.

A.: The significance
[22778] of haemochromatosis gene mutations in the general population: implications
[22779] for screening. Gut 43: 830-836, 1998.

[22780] 2689.

[22781] Cairo, G.; Conte, D.; Bianchi, L.; Fraquelli, M.; Recalcati, S.

[22782] : Reduced serum ceruloplasmin levels in hereditary haemochromatosis. Brit.
[22783] J. Haemat. 114: 226-229, 2001.

[22784] 2690.

[22785] Camaschella, C.; Roetto, A.; Cicilano, M.; Pasquero, P.; Bosio,
[22786] S.; Gubetta, L.; Di Vito, F.; Girelli, D.; Totaro, A.; Carella, M.;
[22787] Grifa, A.; Gasparini, P.: Juvenile and adult hemochromatosis are
[22788] distinct genetic disorders. Europ. J. Hum. Genet. 5: 371-375, 1997.

[22789] 2691.

[22790] Carella, M.; D'Ambrosio, L.; Totaro, A.; Grifa, A.; Valentino,

[22791] M. A.; Piperno, A.; Girelli, D.; Roetto, A.; Franco, B.; Gas-

parini,

[22792] P.; Camaschella, C.: Mutation analysis of the HLA-H gene in Italian

[22793] hemochromatosis patients. Am. J. Hum. Genet. 60: 828-832, 1997.

[22794] 2692.

[22795] Cartwright, G. E.; Edwards, C. Q.; Kravitz, K.; Skolnick, M.;

[22796] Amos, D. B.; Johnson, A.; Bushjaer, L.: Hereditary hemochromatosis:

[22797] phenotypic expression of the disease. New Eng. J. Med. 301: 175-179,

[22798] 1979.

[22799] 2693.

[22800] Cartwright, G. E.; Skolnick, M.; Amos, D. B.; Edwards, C. Q.;

[22801] Kravitz, K.; Johnson, A.: Inheritance of hemochromatosis: linkage

[22802] to HLA. Trans. Assoc. Am. Phys. 91: 273-281, 1978.

[22803] 2694.

[22804] Cazzola, M.; Ascari, E.; Barosi, G.; Claudiani, G.; Dacco, M.;

[22805] Kaltwasser, J. P.; Panaiotopoulos, N.; Schalk, K. P.; Werner,

E. E.

[22806] : Juvenile idiopathic haemochromatosis: a life-threatening disorder

[22807] presenting as hypogonadotropic hypogonadism. Hum. Genet. 65: 149-154,

[22808] 1983.

[22809] 2695.

[22810] Charlton, R. W.; Abrahams, C.; Bothwell, T. H.: Idiopathic hemochromatosis

[22811] in young subjects. Arch. Path. 83: 132-140, 1967.

[22812] 2696.

[22813] Cox, T.: Haemochromatosis: strike while the iron is hot. Nature

[22814] Genet. 13: 386-388, 1996.

[22815] 2697.

[22816] Cox, T. M.; Peters, T. J.: Uptake of iron by duodenal biopsy

[22817] specimens from patients with iron-deficiency anaemia and primary haemochromatosis. Lancet I:

[22818] 123-124, 1978.

[22819] 2698.

[22820] Cragg, S. J.; Darke, C.; Worwood, M.: HLA class I and H ferritin
[22821] gene polymorphisms in normal subjects and patients with haemochromatosis. Hum.
[22822] Genet. 80: 63-68, 1988.

[22823] 2699.

[22824] Cullen, L. M.; Gao, X.; Easteal, S.; Jazwinska, E. C.: The hemochromatosis
[22825] 845 G-to-A and 187 C-to-G mutations: prevalence in non-Caucasian populations. Am.
[22826] J. Hum. Genet. 62: 1403-1407, 1998.

[22827] 2700.

[22828] Cutler, D. J.; Isner, J. M.; Bracey, A. W.; Hufnagel, C. A.; Conrad,
[22829] P. W.; Roberts, W. C.; Kerwin, D. M.; Weintraub, A. M.: Hemochromatosis
[22830] heart disease: an unemphasized cause of potentially re-versible restrictive
[22831] cardiomyopathy. Am. J. Med. 69: 923-928, 1980.

[22832] 2701.

[22833] Dadone, M.; Skolnick, M.; Edwards, C.: Linkage between

hereditary

[22834] hemochromatosis and HLA. (Abstract) Cytogenet. Cell Genet. 32: 261-262,

[22835] 1982.

[22836] 2702.

[22837] Dadone, M. M.; Kushner, J. P.; Edwards, C. Q.; Bishop, D. T.;

[22838] Skolnick, M. H.: Hereditary hemochromatosis: analysis of laboratory

[22839] expression of the disease by genotype in 18 pedigrees. Am. J. Clin.

[22840] Path. 78: 196-207, 1982.

[22841] 2703.

[22842] David, V.; Papadopoulos, P.; Yaouanq, J.; Blayau, M.; Abel, L.;

[22843] Zappone, E.; Perichon, M.; Drysdale, J.; Le Gall, J.-Y.; Simon, M.

[22844] : Ferritin H gene polymorphism in idiopathic hemochromatosis. Hum.

[22845] Genet. 81: 123-126, 1989.

[22846] 2704.

[22847] David, V.; Paul, P.; Simon, M.; Le Gall, J.-Y.; Fauchet, R.; Gicquel,

[22848] I.; Dugast, I.; Le Mignon, L.; Yaouanq, J.; Cohen, D.; Bourel, M.

[22849] : DNA polymorphism related to the idiopathic hemochromatosis gene:

[22850] evidence in a recombinant family. Hum. Genet. 74: 113-120, 1986.

[22851] 2705.

[22852] David, V.; Paul, P.; Yaouanq, J.; Blayau, M.; Fauchet, R.; Cohen,

[22853] D.; Le Gall, J. Y.; Simon, M.: Molecular genetic approach to the

[22854] hemochromatosis gene. (Abstract) Cytogenet. Cell Genet. 46: 604,

[22855] 1987.

[22856] 2706.

[22857] De Braekeleer, M.: A prevalence and fertility study of haemochromatosis

[22858] in Saguenay-Lac-Saint-Jean. Ann. Hum. Biol. 20: 501-505, 1993.

[22859] 2707.

[22860] Debre, R.; Dreyfus, J.-C.; Frezal, J.; Labie, D.; Lamy, M.; Maroteaux,
[22861] P.; Schapira, F.; Schapira, G.: Genetics of haemochromatosis. Ann.
[22862] Hum. Genet. 23: 16-30, 1958.

[22863] 2708.

[22864] de Juan, M.; Reta, A.; Castiella, A.; Pozueta, J.; Prada, A.;

[22865] Cudrado, E.: HFE gene mutations analysis in Basque hereditary haemochromatosis
[22866] patients and controls. Europ. J. Hum. Genet. 9: 961-964, 2001.
[22867] 2709.

[22868] Deugnier, Y. M.; Guyader, D.; Crantock, L.; Lopez, J.-M.; Turlin,
[22869] B.; Yaouanq, J.; Jouanolle, H.; Campion, J.-P.; Launois, B.; Halliday,
[22870] J. W.; Powell, L. W.; Brissot, P.: Primary liver cancer in genetic
[22871] hemochromatosis: a clinical, pathological, and pathogenetic study
[22872] of 54 cases. Gastroenterology 104: 228-234, 1993.

[22873] 2710.

[22874] Dormer, A. E.; Hale, J. F.: Familial Mediterranean fever, a cause
[22875] of periodic fever. Brit. Med. J. 1: 87-89, 1962.

[22876] 2711.

[22877] Dupont, M.; Dross, C.; Smaoui, N.; Nedelec, B.; Grateau, G.; Clepet,
[22878] C.; Gourdier, I.; Kone-Paut, I.; Delpech, M.; Demaille, J.; Touitou,
[22879] I.: Genotypic diagnosis of familial Mediterranean fever (FMF) using
[22880] new microsatellite markers: example of two extensive non-Ashkenazi
[22881] Jewish pedigrees. J. Med. Genet. 34: 375-381, 1997.

[22882] 2712.

[22883] Ehrenfeld, E. N.; Eliakim, M.; Rachmilewitz, M.: Recurrent polyserositis
[22884] (familial Mediterranean fever: periodic disease): a report of fifty-five
[22885] cases. Am. J. Med. 31: 107-123, 1961.

[22886] 2713.

[22887] Eshel, G.; Zemer, D.; Bar-Yochai, A.: Acute orchitis in familial

[22888] Mediterranean fever. Ann. Intern. Med. 109: 164-165, 1988.

[22889] 2714.

[22890] Fischel-Ghodsian, N.; Bu, X.; Oztas, S.; Prezant, T. R.; Danon,

[22891] Y.; Shohat, T.; Rotter, J. I.; Shohat, M.: The gene for familial

[22892] Mediterranean fever maps to the short arm of chromosome 16 in both

[22893] Armenians and non-Ashkenazi Jews. (Abstract) Am. J. Hum. Genet. 51

[22894] (suppl.): A187, 1992.

[22895] 2715.

[22896] Fischel-Ghodsian, N.; Bu, X.; Prezant, T. R.; Oeztas, S.; Huang,

[22897] Z.-S.; Bohlman, M. C.; Rotter, J. I.; Shohat, M.: Regional mapping

[22898] of the gene for familial Mediterranean fever on human chromosome 16p13. Am.

[22899] J. Med. Genet. 46: 689-693, 1993.

[22900] 2716.

[22901] Flatau, E.; Kohn, D.; Schiller, D.; Lurie, M.; Levy, E.: Schon-
lein-Henoch

[22902] syndrome in patients with familial Mediterranean fever.
Arthritis

[22903] Rheum. 25: 42-47, 1982.

[22904] 2717.

[22905] French FMF Consortium: A candidate gene for familial
Mediterranean

[22906] fever. Nature Genet. 17: 25-31, 1997.

[22907] 2718.

[22908] French FMF Consortium: Localization of the familial
Mediterranean

[22909] fever gene (FMF) to a 250-kb interval in non-Ashkenazi
Jewish founder

[22910] haplotypes. Am. J. Hum. Genet. 59: 603-612, 1996.

[22911] 2719.

[22912] Gershoni-Baruch, R.; Brik, R.; Shinawi, M.; Livneh, A.: The
differential

[22913] contribution of MEFV mutant alleles to the clinical profile
of familial

[22914] Mediterranean fever. Europ. J. Hum. Genet. 10: 145-149, 2002.

[22915] 2720.

[22916] Gershoni-Baruch, R.; Shinawi, M.; Leah, K.; Badarnah, K.; Brik,

[22917] R.: Familial Mediterranean fever: prevalence, penetrance and genetic

[22918] drift. Europ. J. Hum. Genet. 9: 634-637, 2001.

[22919] 2721.

[22920] Gershoni-Baruch, R.; Shinawi, M.; Shamaly, H.; Katsinetz, L.;

[22921] Brik, R.: Familial Mediterranean fever: The segregation of four different

[22922] mutations in 13 individuals from one inbred family: geno-type-phenotype

[22923] correlation and intrafamilial variability. Am. J. Med. Genet. 109:

[22924] 198-201, 2002.

[22925] 2722.

[22926] Goldfinger, S. E.: Colchicine for familial Mediterranean fever.

[22927] (Letter) New Eng. J. Med. 287: 1302, 1972.

[22928] 2723.

[22929] Gruberg, L.; Aksentijevich, I.; Balow, J.; Dean, M.; Kovo, M.;

[22930] Pras, M.; Kastner, D. L.: Exclusion of candidate genes in familial

[22931] Mediterranean fever. (Abstract) Cytogenet. Cell Genet. 58: 2113,

[22932] 1991.

[22933] 2724.

[22934] Heller, H.; Sohar, E.; Gafni, J.; Heller, J.: Amyloidosis in

[22935] familial Mediterranean fever. Arch. Intern. Med. 107: 539-550, 1961.

[22936] 2725.

[22937] Hurwich, B. J.; Schwartz, J.; Goldfarb, S.: Record survival of

[22938] siblings with familial Mediterranean fever, phenotypes 1 and 2: case

[22939] of renal vein thrombosis complicating the amyloidosis (phenotype 2). Arch.

[22940] Intern. Med. 125: 308-311, 1970.

[22941] 2726.

[22942] Ilfeld, D.; Kuperman, O.: Correction of a suppressor cell deficiency

[22943] in four patients with familial Mediterranean fever by in vitro or

[22944] in vivo colchicine. Clin. Exp. Immun. 50: 99-106, 1982.

[22945] 2727.

[22946] International FMF Consortium: Ancient missense muta-tions in

[22947] a new member of the RoRet gene family are likely to cause familial

[22948] Mediterranean fever. Cell 90: 797-807, 1997.

[22949] 2728.

[22950] Janeway, T. C.; Mosenthal, H. O.: An unusual paroxysmal syndrome,

[22951] probably allied to recurrent vomiting, with a study of the nitrogen

[22952] metabolism. Trans. Assoc. Am. Phys. 23: 504-518, 1908.

[22953] 2729.

[22954] Jones, M. B.; Adams, J. M.; Passer, J. A.: Amyloidosis in a renal

[22955] allograft in familial Mediterranean fever. Ann. Intern. Med.

87:

[22956] 579-581, 1977.

[22957] 2730.

[22958] Kastner, D. L.; Aksentijevich, I.; Gruberg, L.; Balow, J.; Dean,

[22959] M.; Hampsch, K.; Gazit, E.; Kovo, M.; Pras, M.: Familial Mediterranean

[22960] fever: a 90 marker exclusion map and evidence for linkage to chromosome

[22961] 17. (Abstract) Cytogenet. Cell Genet. 58: 2115, 1991.

[22962] 2731.

[22963] Kees, S.; Langevitz, P.; Zemer, D.; Padeh, S.; Pras, M.; Livneh,

[22964] A.: Attacks of pericarditis as a manifestation of familial Mediterranean

[22965] fever (FMF). Quart. J. Med. 90: 643-647, 1997.

[22966] 2732.

[22967] Khachadurian, A. K.; Armenian, H. K.: Familial paroxysmal polyserositis

[22968] (familial Mediterranean fever): incidence of amyloidosis and mode

[22969] of inheritance. Birth Defects Orig. Art. Ser. X(4): 62-66, 1974.

[22970] 2733.

[22971] Knecht, A.; de Beer, F. C.; Pras, M.: Serum amyloid A protein

[22972] in familial Mediterranean fever. Ann. Intern. Med. 102: 71-72, 1985.

[22973] 2734.

[22974] Kogan, A.; Shinar, Y.; Lidar, M.; Revivo, A.; Langevitz, P.; Padeh,

[22975] S.; Pras, M.; Livneh, A.: Common MEFV mutations among Jewish ethnic

[22976] groups in Israel: high frequency of carrier and phenotype III states

[22977] and absence of a perceptible biological advantage for the carrier

[22978] state. Am. J. Med. Genet. 102: 272-276, 2001.

[22979] 2735.

[22980] Lawrence, J. S.; Mellinkoff, S. M.: Familial Mediterranean fever. Trans.

[22981] Assoc. Am. Phys. 72: 111-121, 1959.

[22982] 2736.

[22983] Livneh, A.; Aksentijevich, I.; Langevitz, P.; Torosyan, Y.; G-
Shoham,
[22984] N.; Shinar, Y.; Pras, E.; Zaks, N.; Padeh, S.; Kastner, D. L.;
Pras,
[22985] M.: A single mutated MEFV allele in Israeli parents suffer-
ing from
[22986] familial Mediterranean fever and Behcet's disease
(FMF-BD). Europ.
[22987] J. Hum. Genet. 9: 191-196, 2001.

[22988] 2737.

[22989] Ludomirsky, A.; Passwell, J.; Boichis, H.: Amyloidosis in
children
[22990] with familial Mediterranean fever. Arch. Dis. Child. 56:
464-467,
[22991] 1981.

[22992] 2738.

[22993] Wang, Y.; Macke, J. P.; Abella, B. S.; Andreasson, K.; Wor-
ley,
[22994] P.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Nathans,
J.:

[22995] A large family of putative transmembrane receptors homologous to the

[22996] product of the Drosophila tissue polarity gene frizzled. J. Biol.

[22997] Chem. 271: 4468-4476, 1996.

[22998] 2739.

[22999] Chioza, B.; Wilkie, H.; Nashef, L.; Blower, J.; McCormick, D.;

[23000] Sham, P.; Asherson, P.; Makoff, A. J.: Association between the alpha-1A

[23001] calcium channel gene CACNA1A and idiopathic generalized epilepsy. Neurology 56:

[23002] 1245-1246, 2001.

[23003] 2740.

[23004] Bell, S. P.; Learned, R. M.; Jantzen, H.-M.; Tjian, R.: Functional

[23005] cooperativity between transcription factors UBF1 and SL1 mediates

[23006] human ribosomal RNA synthesis. Science 241: 1192-1197, 1988.

[23007] 2741.

[23008] Chan, E. K. L.; Imai, H.; Hamel, J. C.; Tan, E. M.: Human autoantibody
[23009] to RNA polymerase I transcription factor hUBF: molecular identity
[23010] of nucleolus organizer region autoantigen NOR-90 and ribosomal RNA
[23011] transcription upstream binding factor. J. Exp. Med. 174: 1239-1244,
[23012] 1991.

[23013] 2742.

[23014] Hisatake, K.; Nishimura, T.; Maeda, Y.; Hanada, K.; Song, C.-Z.;
[23015] Muramatsu, M.: Cloning and structural analysis of cDNA and the gene
[23016] for mouse transcription factor UBF. Nuc. Acids Res. 19: 4631-4637,
[23017] 1991.

[23018] 2743.

[23019] Jantzen, H.-M.; Admon, A.; Bell, S. P.; Tjian, R.: Nucleolar transcription
[23020] factor hUBF contains a DNA-binding motif with homology to HMG proteins. Nature 344:

[23021] 830-836, 1990.

[23022] 2744.

[23023] Jones, K. A.; Black, D. M.; Griffiths, B. L.; Solomon, E.: Localization

[23024] of the human RNA polymerase I transcription factor gene (UBTF) to

[23025] the S17S183 locus on chromosome 17q21 and construction of a long-range

[23026] restriction map of the region. Genomics 30: 602-604, 1995.

[23027] 2745.

[23028] Matera, A. G.; Wu, W.; Imai, H.; O'Keefe, C. L.; Chan, E. K. L.

[23029] : Molecular cloning of the RNA polymerase I transcription factor hUBF/NOR-90

[23030] (UBTF) gene and localization to 17q21.3 by fluorescence in situ hybridization

[23031] and radiation hybrid mapping. Genomics 41: 135-138, 1997.

[23032] 2746.

[23033] O'Mahony, D. J.; Rothblum, L. I.: Identification of two forms

[23034] of the RNA polymerase I transcription factor UBF. Proc. Nat. Acad.

[23035] Sci. 88: 3180-3184, 1991.

[23036] 2747.

[23037] Konishi, H.; Tsutsui, H.; Murakami, T.; Yumikura-Futat-sugi, S.;

[23038] Yamanaka, K.; Tanaka, M.; Iwakura, Y.; Suzuki, N.; Takeda, K.; Akira,

[23039] S.; Nakanishi, K.; Mizutani, H.: IL-18 contributes to the spontaneous

[23040] development of atopic dermatitis-like inflammatory skin lesion independently

[23041] of IgE/stat6 under specific pathogen-free conditions. Proc. Nat.

[23042] Acad. Sci. 99: 11340-11345, 2002.

[23043] 2748.

[23044] Lazner, F.; Gowen, M.; Pavasovic, D.; Kola, I.: Osteopetro-sis

[23045] and osteoporosis: two sides of the same coin. Hum. Molec. Genet. 8:

[23046] 1839-1846, 1999.

[23047] 2749.

[23048] Norman, C. H., Jr.; Dubowy, J.: Pycnodysostosis with splenomegaly

[23049] and anemia. NY State J. Med. 71: 2419-2421, 1971.

[23050] 2750.

[23051] Rantakokko, J.; Aro, H. T.; Savontaus, M.; Vuorio, E.: Mouse

[23052] cathepsin K: cDNA cloning and predominant expression of the gene in

[23053] osteoclasts, and in some hypertrophying chondrocytes during mouse

[23054] development. FEBS Lett. 393: 307-313, 1996.

[23055] 2751.

[23056] Rood, J. A.; Van Horn, S.; Drake, F. H.; Gowen, M.; De-

bouck, C.

[23057] : Genomic organization and chromosome localization of the human cathepsin

[23058] K gene (CTSK). Genomics 41: 169-176, 1997.

[23059] 2752.

[23060] Saftig, P.; Hunziker, E.; Wehmeyer, O.; Jones, S.; Boyde, A.;

[23061] Rommerskirch, W.; Moritz, J. D.; Schu, P.; von Figura, K.: Impaired
[23062] osteoclastic bone resorption leads to osteopetrosis in cathepsin-K-deficient
[23063] mice. Proc. Nat. Acad. Sci. 95: 13453-13458, 1998.

[23064] 2753.

[23065] Shi, G.-P.; Chapman, H. A.; Bhairi, S. M.; DeLeeuw, C.; Reddy,
[23066] V. Y.; Weiss, S. J.: Molecular cloning of human cathepsin O, a novel
[23067] endoproteinase and homologue of rabbit OC2. FEBS Lett. 357: 129-134,
[23068] 1995.

[23069] 2754.

[23070] Evers, R.; Kool, M.; van Deemter, L.; Janssen, H.; Calafat, J.;
[23071] Oomen, L. C. J. M.; Paulusma, C. C.; Oude Elferink, R. P. J.; Baas,
[23072] F.; Schinkel, A. H.; Borst, P.: Drug export activity of the human
[23073] canalicular multispecific organic anion transporter in polarized kidney

[23074] MDCK cells expressing cMOAT (MRP2) cDNA. J. Clin. In-
vest. 101: 1310-1319,
[23075] 1998.

[23076] 2755.

[23077] Gopalan, G.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.;
Donovan,
[23078] P. J.: Chromosome localization of two new mammalian ki-
nases related
[23079] to yeast and fly chromosome segregation-regulators.
Mammalian Genome 9:
[23080] 86-96, 1998.

[23081] 2756.

[23082] Kajihara, S.; Hisatomi, A.; Mizuta, T.; Hara, T.; Ozaki, I.;
Wada,
[23083] I.; Yamamoto, K.: A splice mutation in the human canalic-
ular multispecific
[23084] organic anion transporter gene causes Dubin-Johnson
syndrome. Biochem.
[23085] Biophys. Res. Commun. 253: 454-457, 1998.

[23086] 2757.

[23087] Paulusma, C. C.; Bosma, P. J.; Zaman, G. J. R.; Bakker, C. T.

M.;

[23088] Otter, M.; Scheffer, G. L.; Scheper, R. J.; Borst, P.; Oude Elferink,

[23089] R. P. J.: Congenital jaundice in rats with a mutation in a multidrug

[23090] resistance-associated protein gene. Science 271: 1126-1128, 1996.

[23091] 2758.

[23092] Taniguchi, K.; Wada, M.; Kohno, K.; Nakamura, T.; Kawabe, T.; Kawakami,

[23093] M.; Kagotani, K.; Okumura, K.; Akiyama, S.; Kuwano, M.: A human canalicular

[23094] multispecific organic anion transporter (cMOAT) gene is overexpressed

[23095] in cisplatin-resistant human cancer cell lines with de-creased drug

[23096] accumulation. Cancer Res. 56: 4124-4129, 1996.

[23097] 2759.

[23098] Toh, S.; Wada, M.; Uchiumi, T.; Inokuchi, A.; Makino, Y.; Horie,

[23099] Y.; Adachi, Y.; Sakisaka, S.; Kuwano, M.: Genomic struc-ture of the

[23100] canalicular multispecific organic anion-transporter gene (MRP2/cMOAT)

[23101] and mutations in the ATP-binding-cassette region in Du-bin-Johnson

[23102] syndrome. Am. J. Hum. Genet. 64: 739-746, 1999.

[23103] 2760.

[23104] Zimniak, P.: Dubin-Johnson and Rotor syndromes: molec-ular basis

[23105] and pathogenesis. Semin. Liver Dis.. 13: 248-260, 1993.

[23106] 2761.

[23107] Nolan, K. F.; Greaves, D. R.; Waldmann, H.: The human in-terleukin

[23108] 18 gene IL18 maps to 11q22.2-q22.3, closely linked to the DRD2 gene

[23109] locus and distinct from mapped IDDM loci. Genomics 51: 161-163,

[23110] 1998.

[23111] 2762.

[23112] Okamoto, I.; Kohno, K.; Tanimoto, T.; Iwaki, K.; Ishihara, T.;

[23113] Akamatsu, S.; Ikegami, H.; Kurimoto, M.: IL-18 prevents

the development
[23114] of chronic graft-versus-host disease in mice. J. Immun. 164: 6067-6074,
[23115] 2000.

[23116] 2763.

[23117] Okamura, H.; Tsutsui, H.; Komatsu, T.; Yutsudo, M.; Hakura, A.;
[23118] Tanimoto, T.; Torigoe, K.; Okura, T.; Nukada, Y.; Hattori, K.; Akita,
[23119] K.; Namba, M.; Tanabe, F.; Konishi, K.; Fukuda, S.; Kurimoto, M.:
[23120] Cloning of a new cytokine that induces IFN-gamma production by T cells. Nature 378:
[23121] 88-91, 1995.

[23122] 2764.

[23123] Reddy, P.; Teshima, T.; Kukuruga, M.; Ordemann, R.; Liu, C.; Lowler,
[23124] K.; Ferrara, J. L. M.: Interleukin-18 regulates acute graft-versus-host
[23125] disease by enhancing Fas-mediated donor T cell apoptosis. J. Exp.
[23126] Med. 194: 1433-1440, 2001.

[23127] 2765.

[23128] Rothe, H.; Jenkins, N. A.; Copeland, N. G.; Kolb, H.: Active stage

[23129] of autoimmune diabetes is associated with the expression of a novel

[23130] cytokine, IGIF, which is located near Idd2. J. Clin. Invest. 99:

[23131] 469-474, 1997.

[23132] 2766.

[23133] Sarvetnick, N.: IFN-gamma, IGIF, and IDDM. (Editorial) J. Clin.

[23134] Invest. 99: 371-372, 1997.

[23135] 2767.

[23136] Shida, K.; Shiratori, I.; Matsumoto, M.; Fukumori, Y.; Mat-suhisa,

[23137] A.; Kikkawa, S.; Tsuji, S.; Okamura, H.; Toyoshima, K.; Seya, T.:

[23138] An alternative form of IL-18 in human blood plasma: complex formation

[23139] with IgM defined by monoclonal antibodies. J. Immun. 166: 6671-6679,

[23140] 2001.

[23141] 2768.

[23142] Zhao, J.; Dynlacht, B.; Imai, T.; Hori, T.; Harlow, E.: Ex-
pression
[23143] of NPAT, a novel substrate of cyclin E-CDK2, promotes S-
phase entry. Genes
[23144] Dev. 12: 456-461, 1998.

[23145] 2769.

[23146] Margolis, R. L.; Stine, O. C.; McInnis, M. G.; Ranen, N. G.;
Rubinsztein,
[23147] D. C.; Leggo, J.; Brando, L. V. J.; Kidwai, A. S.; Loev, S. J.;
Breschel,
[23148] T. S.; Callahan, C.; Simpson, S. G.; and 12 others: cDNA
cloning
[23149] of a human homologue of the Caenorhabditis elegans cell
fate-determining
[23150] gene mab-21: expression, chromosomal localization and
analysis of
[23151] a highly polymorphic (CAG)n trinucleotide repeat. Hum.
Molec. Genet. 5:
[23152] 607-616, 1996.

[23153] 2770.

[23154] Margolis, R. L.; Stine, O. C.; Ward, C. M.; Franz, M. L.; Rosenblatt,
[23155] A.; Callahan, C.; Sherr, M.; Ross, C. A.; Potter, N. T.: Unstable
[23156] expansion of the CAG trinucleotide repeat in MAB21L1: report of a
[23157] second pedigree and effect on protein expression. J. Med. Genet. 36:
[23158] 62-64, 1999.

[23159] 2771.

[23160] Mariani, M.; Baldessari, D.; Francisconi, S.; Viggiano, L.; Rocchi,
[23161] M.; Zappavigna, V.; Malgaretti, N.; Consalez, G. G.: Two murine and
[23162] human homologs of mab-21, a cell fate determination gene involved
[23163] in Caenorhabditis elegans neural development. Hum. Molec. Genet. 8:
[23164] 2397-2406, 1999.

[23165] 2772.

[23166] Potter, N. T.: Meiotic instability associated with the CAGR1 trinucleotide
[23167] repeat at 13q13. J. Med. Genet. 34: 411-413, 1997.

[23168] 2773.

[23169] Altshuler, D.; Daly, M.; Kruglyak, L.: Guilt by association. Nature
[23170] Genet. 26: 135-137, 2000.

[23171] 2774.

[23172] Cox, N. J.: Challenges in identifying genetic variation af-fecting
[23173] susceptibility to type 2 diabetes: examples from studies of the calpain-10
[23174] gene. Hum. Molec. Genet. 10: 2301-2305, 2001.

[23175] 2775.

[23176] Horikawa, Y.; Oda, N.; Cox, N. J.; Li, X.; Orho-Melander, M.; Hara,
[23177] M.; Hinokio, Y.; Lindner, T. H.; Mashima, H.; Schwarz, P. E. H.; del
[23178] Bosque-Plata, L.; Horikawa, Y.; and 14 others: Genetic variation
[23179] in the gene encoding calpain-10 is associated with type 2

diabetes
[23180] mellitus. Nature Genet. 26: 163-175, 2000.

[23181] 2776.

[23182] Futreal, P. A.; Cochran, C.; Rosenthal, J.; Miki, Y.; Swenson,

[23183] J.; Hobbs, M.; Bennett, L. M.; Haugen-Strano, A.; Marks, J.; Barrett,

[23184] J. C.; Tavtigian, S. V.; Shattuck-Eidens, D.; Kamb, A.; Skol-

nick,

[23185] M.; Wiseman, R. W.: Isolation of a diverged homeobox gene, MOX1,

[23186] from the BRCA1 region on 17q21 by solution hybrid capture. Hum. Molec.

[23187] Genet. 3: 1359-1364, 1994.

[23188] 2777.

[23189] Jones, K. A.; Black, D. M.; Brown, M. A.; Griffiths, B. L.; Nicolai,

[23190] H. M.; Chambers, J. A.; Bonjardim, M.; Xu, C.-F.; Boyd, M.; McFarlane,

[23191] R.; Korn, B.; Poustka, A.; North, M. A.; Schalkwyk, L.; Lehrach, H.;

[23192] Solomon, E.: The detailed characterisation of a 400 kb

cosmid walk

[23193] in the BRCA1 region: identification and localisation of 10 genes including

[23194] a dual-specificity phosphatase. Hum. Molec. Genet. 3: 1927-1934,

[23195] 1994.

[23196] 2778.

[23197] Horton, Y. M.; Sullivan, M.; Houslay, M. D.: Molecular cloning

[23198] of a novel splice variant of human type IVA (PDE-IVA) cyclic AMP phosphodiesterase

[23199] and localization of the gene to the p13.2-q12 region of human chromosome

[23200] 19. Biochem. J. 308: 683-691, 1995.

[23201] 2779.

[23202] Huston, E.; Pooley, L.; Julien, P.; Scotland, G.; McPhee, I.; Sullivan,

[23203] M.; Bolger, G.; Houslay, M. D.: The human cyclic AMP-specific phosphodiesterase

[23204] PDE-46 (HSPDE4A4B) expressed in transfected COS7 cells occurs as both

[23205] particulate and cytosolic species that exhibit distinct ki-

netics of
[23206] inhibition by the antidepressant rolipram. J. Biol. Chem. 271: 31334-31344,
[23207] 1996.

[23208] 2780.

[23209] Livi, G. P.; Kmetz, P.; McHale, M. M.; Cieslinski, L. B.; Sathe,
[23210] G. M.; Taylor, D. P.; Davis, R. L.; Torphy, T. J.; Balcarek, J. M.
[23211] : Cloning and expression of cDNA for a human low-K(m), rolipram-sensitive
[23212] cyclic AMP phosphodiesterase. Molec. Cell. Biol. 10: 2678-2686,
[23213] 1990.

[23214] 2781.

[23215] Obernolte, R.; Bhakta, S.; Alvarez, R.; Bach, C.; Zuppan, P.; Mulkins,
[23216] M.; Jarnagin, K.; Shelton, E. R.: The cDNA of a human lymphocyte
[23217] cyclic-AMP phosphodiesterase (PDE IV) reveals a multigene family. Gene 129:
[23218] 239-247, 1993.

[23219] 2782.

[23220] Sullivan, M.; Egerton, M.; Shakur, Y.; Marquardsen, A.; Houslay,

[23221] M. D.: Molecular cloning and expression, in both COS-1 cells and

[23222] S. cerevisiae, of a human cytosolic type-IVA, cyclic AMP specific

[23223] phosphodiesterase (hPDE-IVA-h6.1). Cell. Signal. 6: 793-812, 1994.

[23224] 2783.

[23225] Sullivan, M.; Rena, G.; Begg, F.; Gordon, L.; Olsen, A. S.; Houslay,

[23226] M. D.: Identification and characterization of the human homologue

[23227] of the short PDE4A cAMP-specific phosphodiesterase RD1 (PDE4A1) by

[23228] analysis of the human HSPDE4A gene locus located at chromosome 19p13.2. Biochem.

[23229] J. 333: 693-703, 1998.

[23230] 2784.

[23231] Wilson, M.; Sullivan, M.; Brown, N.; Houslay, M. D.: Purifi-cation,

[23232] characterization and analysis of rolipram inhibition of a human-type

[23233] IVA cyclic AMP-specific phosphodiesterase expressed in yeast. Biochem.

[23234] J. 304: 407-415, 1994.

[23235] 2785.

[23236] Huston, E.; Lumb, S.; Russell, A.; Catterall, C.; Ross, A. H.;

[23237] Steele, M. R.; Bolger, G. B.; Perry, M. J.; Owens, R. J.; Hous-lay,

[23238] M. D.: Molecular cloning and transient expression in COS7 cells of

[23239] a novel human PDE4B cAMP-specific phosphodiesterase, HSPDE4B3. Biochem.

[23240] J. 328: 549-558, 1997.

[23241] 2786.

[23242] Szpirer, C.; Szpirer, J.; Riviere, M.; Swinnen, J.; Vicini, E.;

[23243] Conti, M.: Chromosomal localization of the human and rat genes (PDE4D

[23244] and PDE4B) encoding the cAMP-specific phosphodi-esterases 3 and 4. Cytogenet.

[23245] Cell Genet. 69: 11-14, 1995.

[23246] 2787.

[23247] Xu, R. X.; Hassell, A. M.; Vanderwall, D.; Lambert, M. H.; Holmes,

[23248] W. D.; Luther, M. A.; Rocque, W. J.; Milburn, M. V.; Zhao, Y.; Ke,

[23249] H.; Nolte, R. T.: Atomic structure of PDE4: insights into phosphodiesterase

[23250] mechanism and specificity. Science 288: 1822-1825, 2000.

[23251] 2788.

[23252] Bolger, G. B.; Erdogan, S.; Jones, R. E.; Loughney, K.; Scot-land,

[23253] G.; Hoffmann, R.; Wilkinson, I.; Farrell, C.; Houslay, M. D.: Characterization

[23254] of five different proteins produced by alternatively spliced mRNAs

[23255] from the human cAMP-specific phosphodiesterase PDE4D gene. Biochem.

[23256] J. 328: 539-548, 1997.

[23257] 2789.

[23258] Dudai, Y.; Jan, Y.-N.; Byers, D.; Quinn, W. G.; Benzer, S.: dunce,

[23259] a mutant of Drosophila deficient in learning. Proc. Nat. Acad. Sci. 73:
[23260] 1684-1688, 1976.

[23261] 2790.

[23262] Hansen, G.; Jin, S.-L. C.; Umetsu, D. T.; Conti, M.: Absence of
[23263] muscarinic cholinergic airway responses in mice deficient in the cyclic
[23264] nucleotide phosphodiesterase PDE4D. Proc. Nat. Acad. Sci. 97: 6751-6756,
[23265] 2000.

[23266] 2791.

[23267] Jin, S.-L. C.; Richard, F. J.; Kuo, W.-P.; D'Ercole, A. J.; Conti,
[23268] M.: Impaired growth and fertility of cAMP-specific phos-phodiesterase
[23269] PDE4D-deficient mice. Proc. Nat. Acad. Sci. 96: 11998-12003, 1999.
[23270] 2792.

[23271] Partanen, J.; Armstrong, E.; Makela, T. P.; Korhonen, J.; Sandberg,

[23272] M.; Renkonen, R.; Knuutila, S.; Huebner, K.; Alitalo, K.: A novel
[23273] endothelial cell surface receptor tyrosine kinase with extracellular
[23274] epidermal growth factor homology domains. Molec. Cell. Biol. 12:
[23275] 1698-1707, 1992.

[23276] 2793.

[23277] Ramsden, J. D.; Cocks, H. C.; Shams, M.; Nijjar, S.; Watkinson,
[23278] J. C.; Sheppard, M. C.; Ahmed, A.; Eggo, M. C.: Tie-2 is expressed
[23279] on thyroid follicular cells, is increased in goiter, and is regulated
[23280] by thyrotropin through cyclic adenosine 3-prime,5-prime-monophosphate. J.
[23281] Clin. Endocr. Metab. 86: 2709-2716, 2001.

[23282] 2794.

[23283] Zhao, J.; Kennedy, B. K.; Lawrence, B. D.; Barbie, D. A.; Matera,
[23284] A. G.; Fletcher, J. A.; Harlow, E.: NPAT links cyclin E-Cdk2 to the

[23285] regulation of replication-dependent histone gene transcription. Genes

[23286] Dev. 14: 2283-2297, 2000.

[23287] 2795.

[23288] Kalaydjieva, L.; Gresham, D.; Gooding, R.; Heather, L.; Baas, F.;

[23289] de Jonge, R.; Blechschmidt, K.; Angelicheva, D.; Chandler, D.; Worsley,

[23290] P.; Rosenthal, A.; King, R. H. M.; Thomas, P. K.: N-myc downstream-regulated

[23291] gene 1 is mutated in hereditary motor and sensory neuropathy-Lom. Am.

[23292] J. Hum. Genet. 67: 47-58, 2000.

[23293] 2796.

[23294] Kalaydjieva, L.; Hallmayer, J.; Chandler, D.; Savov, A.; Nikolova,

[23295] A.; Angelicheva, D.; King, R. H. H.; Ishpekova, B.; Honeyman, K.;

[23296] Calafell, F.; Shmarov, A.; Petrova, J.; Turnev, I.; Hristova, A.;

[23297] Moskov, M.; Stancheva, S.; Petkova, I.; Bittles, A. H.; Georgieva,

[23298] V.; Middleton, L.; Thomas, P. K.: Gene mapping in Gypsies identifies
[23299] a novel demyelinating neuropathy on chromosome 8q24. Nature Genet. 14:
[23300] 214-217, 1996.

[23301] 2797.

[23302] Tsujikawa, M.; Kurahashi, H.; Tanaka, T.; Okada, M.; Ya-
mamoto,
[23303] S.; Maeda, N.; Watanabe, H.; Inoue, Y.; Kiridoshi, A.; Mat-
sumoto,
[23304] K.; Ohashi, Y.; Kinoshita, S.; Shimomura, Y.; Nakamura, Y.; Tano,
[23305] Y.: Homozygosity mapping of a gene responsible for gelatinous drop-like
[23306] corneal dystrophy to chromosome 1p. Am. J. Hum. Genet. 63: 1073-1077,
[23307] 1998.

[23308] 2798.

[23309] Yamamoto, S.; Okada, M.; Tsujikawa, M.; Shimomura, Y.; Nishida,
[23310] K.; Inoue, Y.; Watanabe, H.; Maeda, N.; Kurahashi, H.; Ki-
noshita,

[23311] S.; Nakamura, Y.; Tano, Y.: A kerato-epithelin (beta-ig-h3) mutation
[23312] in lattice corneal dystrophy type IIIA. (Letter) Am. J. Hum. Genet. 62:
[23313] 719-722, 1998.

[23314] 2799.

[23315] FitzGerald, M. G.; MacDonald, D. J.; Krainer, M.; Hoover, I.;

[23316] O'Neil, E.; Unsal, H.; Silva-Arrieto, S.; Finkelstein, D. M.; Beer-Romero,
[23317] P.; Englert, C.; Sgroi, D. C.; Smith, B. L.; Younger, J. W.; Garber,
[23318] J. E.; Duda, R. B.; Mayzel, K. A.; Isselbacher, K. J.; Friend, S.

[23319] H.; Haber, D. A.: Germ-line BRCA1 mutations in Jewish and non-Jewish
[23320] women with early-onset breast cancer. New Eng. J. Med. 334: 143-149,
[23321] 1996.

[23322] 2800.

[23323] Franceschi, S.; Parazzini, F.; Negri, E.; Booth, M.; La Vec-chia,
[23324] C.; Beral, V.; Tzonov, A.; Trichopoulos, D.: Pooled analysis

of 3

[23325] European case-control studies of epithelial ovarian cancer. III. Oral

[23326] contraceptive use. Int. J. Cancer 49: 61-65, 1991.

[23327] 2801.

[23328] Friedman, E.; Bar-Sade, R. B.; Kruglikova, A.; Risel, S.; Levy-Lahad,

[23329] E.; Halle, D.; Bar-On, E.; Gershoni-Baruch, R.; Dagan, E.; Kepten,

[23330] I.; Peretz, T.; Lerer, I.; Wienberg, N.; Shushan, A.; Abe-liovich,

[23331] D.: Double heterozygotes for the Ashkenazi founder mu-tations in BRCA1

[23332] and BRCA2 genes. (Letter) Am. J. Hum. Genet. 63: 1224-1227, 1998.

[23333] 2802.

[23334] Friedman, L. S.; Gayther, S. A.; Kurosaki, T.; Gordon, D.; Noble,

[23335] B.; Casey, G.; Ponder, B. A. J.; Anton-Culver, H.: Mutation analysis

[23336] of BRCA1 and BRCA2 in a male breast cancer population. Am. J. Hum.

[23337] Genet. 60: 313-319, 1997.

[23338] 2803.

[23339] Friedman, L. S.; Ostermeyer, E. A.; Szabo, C. I.; Dowd, P.; Lynch,

[23340] E. D.; Rowell, S. E.; King, M.-C.: Confirmation of BRCA1 by analysis

[23341] of germline mutations linked to breast and ovarian cancer in ten families. Nature

[23342] Genet. 8: 399-404, 1994.

[23343] 2804.

[23344] Friedman, L. S.; Szabo, C. I.; Ostermeyer, E. A.; Dowd, P.; Butler,

[23345] L.; Park, T.; Lee, M. K.; Goode, E. L.; Rowell, S. E.; King, M.-C.

[23346] : Novel inherited mutations and variable expressivity of BRCA1 alleles,

[23347] including the founder mutation 185delAG in Ashkenazi Jewish families. Am.

[23348] J. Hum. Genet. 57: 1284-1297, 1995.

[23349] 2805.

[23350] Futreal, P. A.; Liu, Q.; Shattuck-Eidens, D.; Cochran, C.;

Harshman,
[23351] K.; Tavtigian, S.; Bennett, L. M.; Haugen-Strano, A.; Swensen, J.;
[23352] Miki, Y.; Eddington, K.; McClure, M.; and 15 others: BRCA1 mutation
[23353] in primary breast and ovarian carcinomas. Science 266: 120-122,
[23354] 1994.

[23355] 2806.

[23356] Gayther, S. A.; Harrington, P.; Russell, P.; Kharkevich, G.; Garkavtseva,
[23357] R. F.; Ponder, B. A. J.: Frequently occurring germ-line mu-tations
[23358] of the BRCA1 gene in ovarian cancer families from Russia. (Letter) Am.
[23359] J. Hum. Genet. 60: 1239-1242, 1997.

[23360] 2807.

[23361] Gayther, S. A.; Harrington, P.; Russell, P.; Kharkevich, G.; Garkavtseva,
[23362] R. F.; Ponder, B. A. J.; UKCCCR Familial Ovarian Cancer Study Group
[23363] : Rapid detection of regionally clustered germ-line BRCA1

mutations
[23364] by multiplex heteroduplex analysis. Am. J. Hum. Genet. 58: 451-456,
[23365] 1996.

[23366] 2808.

[23367] Gayther, S. A.; Warren, W.; Mazoyer, S.; Russell, P. A.; Harrington,
[23368] P. A.; Chiano, M.; Seal, S.; Hamoudi, R.; van Rensburg, E. J.; Dunning,
[23369] A. M.; Love, R.; Evans, G.; Easton, D.; Clayton, D.; Stratton, M.
[23370] R.; Ponder, B. A. J.: Germline mutations of the BRCA1 gene in breast
[23371] and ovarian cancer families provide evidence for a genotype-phenotype
[23372] correlation. Nature Genet. 11: 428-433, 1995.

[23373] 2809.

[23374] Goldgar, D. E.; Fields, P.; Lewis, C. M.; Cannon-Albright, L.

[23375] A.; Linker, G.; Tran, T.; Skolnick, M.: A large kindred with 17q-linked
[23376] susceptibility to breast and ovarian cancer: relationship

between
[23377] genotype and phenotype. (Abstract) Am. J. Hum. Genet. 51 (suppl.):
[23378] A27, 1992.

[23379] 2810.

[23380] Goldgar, D. E.; Reilly, P. R.: A common BRCA1 mutation in the
[23381] Ashkenazim. Nature Genet. 11: 113-114, 1995.

[23382] 2811.

[23383] Gorski, B.; Byrski, T.; Huzarski, T.; Jakubowska, A.; Menkiszak,
[23384] J.; Gronwald, J.; Pluzanska, A.; Bebenek, M.; Fischer-Maliszewska,
[23385] L.; Grzybowska, E.; Narod, S. A.; Lubinski, J.: Founder mu-tations
[23386] in the BRCA1 gene in Polish families with breast-ovarian cancer. Am.
[23387] J. Hum. Genet. 66: 1963-1968, 2000.

[23388] 2812.

[23389] Gowen, L. C.; Avrutskaya, A. V.; Latour, A. M.; Koller, B. H.;

[23390] Leadon, S. A.: BRCA1 required for transcription-coupled

repair of
[23391] oxidative DNA damage. Science 281: 1009-1012, 1998.

[23392] 2813.

[23393] Gowen, L. C.; Johnson, B. L.; Latour, A. M.; Sulik, K. K.; Koller,
[23394] B. H.: Brca1 deficiency results in early embryonic lethality characterized
[23395] by neuroepithelial abnormalities. Nature Genet. 12: 191-194, 1996.
[23396] 2814.

[23397] Hacia, J. G.; Brody, L. C.; Chee, M. S.; Fodor, S. P. A.; Collins,
[23398] F. S.: Detection of heterozygous mutations in BRCA1 using high density
[23399] oligonucleotide arrays and two-colour fluorescence analysis. Nature
[23400] Genet. 14: 441-447, 1996.

[23401] 2815.

[23402] Hakem, R.; de la Pompa, J. L.; Elia, A.; Potter, J.; Mak, T. W.

[23403] : Partial rescue of Brca1(5-6) early embryonic lethality by p53 or

[23404] p21 null mutation. Nature Genet. 16: 298-302, 1997.

[23405] 2816.

[23406] Hakem, R.; de la Pompa, J. L.; Sirard, C.; Mo, R.; Woo, M.; Hakem,

[23407] A.; Wakeham, A.; Potter, J.; Reitmair, A.; Billia, F.; Firpo, E.;

[23408] Hui, C. C.; Roberts, J.; Rossant, J.; Mak, T. W.: The tumor suppressor

[23409] gene Brca1 is required for embryonic cellular proliferation in the

[23410] mouse. Cell 85: 1009-1023, 1996.

[23411] 2817.

[23412] Hall, J. M.; Friedman, L.; Guenther, C.; Lee, M. K.; Weber, J.

[23413] L.; Black, D. M.; King, M.-C.: Closing in on a breast cancer gene

[23414] on chromosome 17q. Am. J. Hum. Genet. 50: 1235-1242, 1992.

[23415] 2818.

[23416] Berkvens, T. M.; van Ormondt, H.; Gerritsen, E. J. A.; Meera Khan,

[23417] P.; van der Eb, A. J.: Identical 3250-bp deletion between two AluI

[23418] repeats in the ADA genes of unrelated ADA-SCID patients. Genomics 7:
[23419] 486-490, 1990.

[23420] 2819.

[23421] Peng, Z. G.; Calvert, I.; Clark, J.; Helman, L.; Kahn, R.; Kung,
[23422] H. F.: Molecular cloning, sequence analysis and mRNA expression of
[23423] human ADP-ribosylation factor. Biofactors 2: 45-49, 1989.
[23424] 2820.

[23425] Sanford, J. P.; Eddy, R. L.; Doyle, D.; Shows, T. B.: Assignment
[23426] of human asialoglycoprotein receptor gene (ASGR1) to chromosome 17p11-13. Genomics 11:
[23427] 779-781, 1991.

[23428] 2821.

[23429] Wulczyn, F. G.; Naumann, M.; Scheidereit, C.: Candidate proto-oncogene
[23430] bcl-3 encodes a subunit-specific inhibitor of transcription factor

[23431] NF-kappa-B. Nature 358: 597-599, 1992.

[23432] 2822.

[23433] Falk, C. T.: New family data supporting the MN/GC linkage. (Abstract) Cytogenet.

[23434] Cell Genet. 37: 466, 1984.

[23435] 2823.

[23436] Hogan, K.: To fire the train: a second malignant-hyperthermia

[23437] gene. (Editorial) Am. J. Hum. Genet. 1303-1308, 1997.

[23438] 2824.

[23439] Monnier, N.; Procaccio, V.; Stieglitz, P.; Lunardi, J.: Malignant-hyperthermia

[23440] susceptibility is associated with a mutation of the alpha-1-subunit

[23441] of the human dihydropyridine-sensitive L-type voltage-dependent calcium-channel

[23442] receptor in skeletal muscle. Am. J. Hum. Genet. 60: 1316-1325, 1997.

[23443] 2825.

[23444] Sanford, J. P.; Elliott, R. W.; Doyle, D.: Asialoglycoprotein

[23445] receptor genes are linked on chromosome 11 in the mouse. DNA 7:

[23446] 721-728, 1988.

[23447] 2826.

[23448] Spiess, M.; Schwartz, A. L.; Lodish, H. F.: Sequence of human

[23449] asialoglycoprotein receptor cDNA: an internal signal sequence for

[23450] membrane insertion. J. Biol. Chem. 260: 1979-1982, 1985.

[23451] 2827.

[23452] Spiess, M.; Lodish, H. F.: Sequence of a second human asialoglycoprotein

[23453] receptor: conservation of two receptor genes during evo-
lution. Proc.
[23454] Nat. Acad. Sci. 82: 6465-6469, 1985.

[23455] 2828.

[23456] Arfin, S. M.; Cirullo, R. E.; Arredondo-Vega, F. X.; Smith,
M.
[23457] : Assignment of the structural gene for asparagine syn-
thetase to human
[23458] chromosome 7. Somat. Cell Genet. 9: 517-531, 1983.

[23459] 2829.

[23460] Greco, A.; Ittmann, M.; Basilico, C.: Molecular cloning of a
gene
[23461] that is necessary for G(1) progression in mammalian cells.
Proc.
[23462] Nat. Acad. Sci. 84: 1565-1569, 1987.

[23463] 2830.

[23464] Heng, H. H. Q.; Shi, X.-M.; Scherer, S. W.; Andrulis, I. L.;
Tsui,
[23465] L.-C.: Refined localization of the asparagine synthetase
gene (ASNS)
[23466] to chromosome 7, region q21.3, and characterization of

the somatic
[23467] cell hybrid line 4AF/106/KO15. Cytogenet. Cell Genet. 66: 135-138,
[23468] 1994.

[23469] 2831.

[23470] Lambert, M. A.; Cairney, A. E. L.; Ray, P. N.; Weksberg, R.; Andrulis,
[23471] I. L.: Genomic characterization of the human asparagine synthetase
[23472] gene. (Abstract) Am. J. Hum. Genet. 39: A207 only, 1986.

[23473] 2832.

[23474] Zhang, Y. P.; Lambert, M. A.; Cairney, A. E. L.; Wills, D.; Ray,
[23475] P. N.; Andrulis, I. L.: Molecular structure of the human asparagine
[23476] synthetase gene. Genomics 4: 259-265, 1989.

[23477] 2833.

[23478] Cirullo, R. E.; Arredondo-Vega, F. X.; Smith, M.; Wasmuth, J. J.
[23479] : Isolation and characterization of interspecific heat-resistant hybrids

[23480] between a temperature-sensitive Chinese hamster cell asparaginyl-tRNA

[23481] synthetase mutant and normal human leukocytes: assignment of human

[23482] asnS gene to chromosome 18. Somat. Cell Genet. 9: 215-233, 1983.

[23483] 2834.

[23484] Shows, T. B.: Personal Communication. Buffalo, N. Y. 1/11/1983.

[23485] 2835.

[23486] Akasaka, T.; Miura, I.; Takahashi, N.; Akasaka, H.; Yonetani, N.;

[23487] Ohno, H.; Fukuhara, S.; Okuma, M.: A recurring translocation, t(3;6)(q27;p21),

[23488] in non-Hodgkin's lymphoma results in replacement of the 5-prime regulatory

[23489] region of BCL6 with a novel H4 histone gene. Cancer Res. 57: 7-12,

[23490] 1997.

[23491] 2836.

[23492] Baron, B. W.; Nucifora, G.; McCabe, N.; Espinosa, R., III; Le Beau,

[23493] M. M.; McKeithan, T. W.: Identification of the gene associated with

[23494] the recurring chromosomal translocations t(3;14)(q27;q32) and t(3;22)(q27;q11)

[23495] in B-cell lymphomas. Proc. Nat. Acad. Sci. 90: 5262-5266, 1993.

[23496] 2837.

[23497] Capello, D.; Carbone, A.; Pastore, C.; Gloghini, A.; Saglio, G.;

[23498] Gaidano, G.: Point mutations of the BCL-6 gene in Burkitt's syndrome. Brit.

[23499] J. Haemat. 99: 168-170, 1997.

[23500] 2838.

[23501] Cattoretti, G.; Chang, C.-C.; Cechova, K.; Zhang, J.; Ye, B. H.;

[23502] Falini, B.; Louie, D. C.; Offit, K.; Chaganti, R. S. K.; Dalla-Favera,

[23503] R.: BCL-6 protein is expressed in germinal-center B cells. Blood 86:

[23504] 45-53, 1995.

[23505] 2839.

[23506] Cesarman, E.; Chadburn, A.; Liu, Y.-F.; Migliazza, A.; Dalla-Favera,
[23507] R.; Knowles, D. M.: BCL-6 gene mutations in posttrans-plantation lymphoproliferative
[23508] disorders predict response to therapy and clinical out-come. Blood 92:
[23509] 2294-2302, 1998.

[23510] 2840.

[23511] Chaganti, S. R.; Chen, W.; Parsa, N.; Offit, K.; Louie, D. C.;

[23512] Dalla-Favera, R.; Chaganti, R. S. K.: Involvement of BCL6 in chromosomal
[23513] aberrations affecting band 3q27 in B-cell non-Hodgkin lymphoma. Genes
[23514] Chromosomes Cancer 23: 323-327, 1998.

[23515] 2841.

[23516] Chaganti, S. R.; Rao, P. H.; Chen, W.; Dyomin, V.; Jhanwar, S.
[23517] C.; Parsa, N. Z.; Dalla-Favera, R.; Chaganti, R. S. K.: Deregulation
[23518] of BCL6 in non-Hodgkin lymphoma by insertion of IGH sequences in complex

[23519] translocations involving band 3q27. Genes Chromosomes Cancer 23:

[23520] 328-336, 1998.

[23521] 2842.

[23522] Chang, C.-C.; Ye, B. H.; Chaganti, R. S. K.; Dalla-Favera, R.:

[23523] BCL-6, a POZ/zinc-finger protein, is a sequence-specific transcriptional

[23524] repressor. Proc. Nat. Acad. Sci. 93: 6947-6952, 1996.

[23525] 2843.

[23526] Dhordain, P.; Albagli, O.; Honore, N.; Guidez, F.; Lantoine, D.;

[23527] Schmid, M.; De The, H.; Zelent, A.; Koken, M. H. M.: Colo-calization

[23528] and heteromerization between the two human oncogene POZ/zinc finger

[23529] proteins, LAZ3 (BCL6) and PLZF. Oncogene 19: 6240-6250, 2000.

[23530] 2844.

[23531] Gaidano, G.; Capello, D.; Gloghini, A.; Fassone, L.; Vivenza,

[23532] D.; Ariatti, C.; Migliazza, A.; Saglio, G.; Carbone, A.: Frequent

[23533] mutation of bcl-6 proto-oncogene in high grade, but not low grade,

[23534] MALT lymphomas of the gastrointestinal tract. Haematologica 84:

[23535] 582-588, 1999.

[23536] 2845.

[23537] Hamblin, T. J.; Davis, Z.; Gardiner, A.; Oscier, D. G.; Stevenson,

[23538] F. K.: Unmutated Ig V(H) genes are associated with a more aggressive

[23539] form of chronic lymphocytic leukemia. Blood 94: 1848-1854, 1999.

[23540] 2846.

[23541] Hosokawa, Y.; Maeda, Y.; Ichinohasama, R.; Miura, I.; Taniwaki,

[23542] M.; Seto, M.: The Ikaros gene, a central regulator of lymphoid differentiation,

[23543] fuses to the BCL6 gene as a result of t(3;7)(q27;p12) translocation

[23544] in a patient with diffuse large B-cell lymphoma. Blood 95:

2719-2721,

[23545] 2000.

[23546] 2847.

[23547] Ichii, H.; Sakamoto, A.; Hatano, M.; Okada, S.; Toyama, H.; Taki,

[23548] S.; Arima, M.; Kuroda, Y.; Tokuhisa, T.: Role of Bcl-6 in the generation

[23549] and maintenance of memory CD8+ T cells. Nature Immun. 3: 558-563,

[23550] 2002.

[23551] 2848.

[23552] Ichinohasama, R.; Miura, I.; Funato, T.; Sato, I.; Suzuki, C.;

[23553] Saito, Y.; Decoteau, J. F.; Myers, J. B.; Kadin, M. E.; Sawai, T.;

[23554] Ooya, K.: A recurrent nonrandom translocation (3;7)(q27;p12) associated

[23555] with BCL-6 gene rearrangement in B-cell diffuse large cell lymphoma. Cancer

[23556] Genet. Cytogenet. 104: 19-27, 1998.

[23557] 2849.

[23558] Kerckaert, J.-P.; Deweindt, C.; Tilly, H.; Quief, S.; Lecocq,

[23559] G.; Bastard, C.: LAZ3, a novel zinc-finger encoding gene, is disrupted

[23560] by recurring chromosome 3q27 translocations in human lymphomas. Nature

[23561] Genet. 5: 66-70, 1993.

[23562] 2850.

[23563] Liao, X.; Gilbert, D. J.; Dent, A.; Staudt, L. M.; Jenkins, N.

[23564] A.; Copeland, N. G.: Mapping of the mouse Bcl6 gene to chromosome

[23565] 16. Mammalian Genome 7: 621-622, 1996.

[23566] 2851.

[23567] Migliazza, A.; Martinotti, S.; Chen, W.; Fusco, C.; Ye, B. H.;

[23568] Knowles, D. M.; Offit, K.; Changanti, R. S. K.; Dalla-Favera, R.:

[23569] Frequent somatic hypermutation of the 5-prime noncoding region of

[23570] the BCL6 gene in B-cell lymphoma. Proc. Nat. Acad. Sci. 92: 12520-12524,

[23571] 1995.

[23572] 2852.

[23573] Miki, T.; Kawamata, N.; Hirosawa, S.; Aoki, N.: Gene involved

[23574] in the 3q27 translocation associated with B-cell lymphoma, BCL5, encodes

[23575] a Kruppel-like zinc-finger protein. Blood 83: 26-32, 1994.

[23576] 2853.

[23577] Sahota, S. S.; Davis, Z.; Hamblin, T. J.; Stevenson, F. K.: Somatic

[23578] mutation of bcl-6 genes can occur in the absence of V-H mutations

[23579] in chronic lymphocytic leukemia. Blood 95: 3534-3540, 2000.

[23580] 2854.

[23581] Shaffer, A. L.; Yu, X.; He, Y.; Boldrick, J.; Chan, E. P.; Staudt,

[23582] L. M.: BCL-6 represses genes that function in lymphocyte differentiation,

[23583] inflammation, and cell cycle control. Immunity 13: 199-212, 2000.

[23584] 2855.

[23585] Shen, H. M.; Peters, A.; Baron, B.; Zhu, X.; Storb, U.: Muta-

tion

[23586] of BCL-6 gene in normal B cells by the process of somatic hypermutation

[23587] of Ig genes. Science 280: 1750-1752, 1998.

[23588] 2856.

[23589] Ueda, C.; Akasaka, T.; Kurata, M.; Maesako, Y.; Nishikori, M.;

[23590] Ichinohasama, R.; Imada, K.; Uchiyama, T.; Ohno, H.: The gene for

[23591] interleukin-21 receptor is the partner of BCL6 in t(13;16)(q27;p11),

[23592] which is recurrently observed in diffuse large B-cell lymphoma. Oncogene 21:

[23593] 368-376, 2002.

[23594] 2857.

[23595] Ye, B. H.; Cattoretti, G.; Shen, Q.; Zhang, J.; Hawe, N.; de Waard,

[23596] R.; Leung, C.; Nouri-Shirazi, M.; Orazi, A.; Chaganti, R. S. K.; Rothman,

[23597] P.; Stall, A. M.; Pandolfi, P.-P.; Dalla-Favera, R.: The BCL-6 proto-oncogene

[23598] controls germinal-centre formation and Th2-type inflam-

mation. Nature
[23599] Genet. 16: 161-170, 1997.

[23600] 2858.

[23601] Ye, B. H.; Lista, F.; Lo Coco, F.; Knowles, D. M.; Offit, K.;

[23602] Chaganti, R. S. K.; Dalla-Favera, R.: Alterations of a zinc finger-encoding
[23603] gene, BCL-6, in diffuse large-cell lymphoma. Science 262: 747-750,
[23604] 1993.

[23605] 2859.

[23606] Ye, B. H.; Rao, P. H.; Chaganti, R. S. K.; Dalla-Favera, R.:

[23607] Cloning of bcl-6, the locus involved in chromosome translocations
[23608] affecting band 3q27 in B-cell lymphoma. Cancer Res. 53: 2732-2735,
[23609] 1993.

[23610] 2860.

[23611] Rimokh, R.; Rouault, J. P.; Wahbi, K.; Gadoux, M.; Lafage, M.;
[23612] Archimbaud, E.; Charrin, C.; Gentilhomme, O.; Germain,

D.; Samarut,
[23613] J.; Magaud, J. P.: A chromosome 12 coding region is juxtaposed to
[23614] the MYC protooncogene locus in a t(8;12)(q24;q22) translocation in
[23615] a case of B-cell chronic lymphocytic leukemia. Genes Chromosomes
[23616] Cancer 3: 24-36, 1991.

[23617] 2861.

[23618] Gejman, P. V.; Weinstein, L. S.; Martinez, M.; Spiegel, A. M.;
[23619] Cao, Q.; Hsieh, W.-T.; Hoehe, M. R.; Gershon, E. S.: Genetic mapping
[23620] of the Gs-alpha subunit gene (GNAS1) to the distal long arm of chromosome
[23621] 20 using a polymorphism detected by denaturing gradient gel electrophoresis. Genomics 9:
[23622] 782-783, 1991.

[23623] 2862.

[23624] Fragoso, M. C. B. V.; Latronico, A. C.; Carvalho, F. M.; Zerbini,
[23625] M. C. N.; Marcondes, J. A. M.; Araujo, L. M. B.; Lando, V. S.;

Frazzatto,
[23626] E. T.; Mendonca, B. B.; Villares, S. M. F.: Activating mutation of
[23627] the stimulatory G protein (gsp) as a putative cause of ovarian and
[23628] testicular human stromal Leydig cell tumors. J. Clin. Endocr. Metab. 83:
[23629] 2074-2078, 1998.

[23630] 2863.

[23631] Gopal Rao, V. V. N.; Schnittger, S.; Hansmann, I.: G protein

[23632] Gs-alpha (GNAS1), the probable candidate gene for Albright hereditary
[23633] osteodystrophy, is assigned to human chromosome 20q12-q13.2. Genomics 10:
[23634] 257-261, 1991.

[23635] 2864.

[23636] Gorelov, V. N.; Dumon, K.; Barteneva, N. S.; Palm, D.; Roher,
[23637] H.-D.; Goretzki, P. E.: Overexpression of Gs-alpha subunit in thyroid
[23638] tumors bearing a mutated Gs-alpha gene. J. Cancer Res.

Clin. Oncol. 121:
[23639] 219-224, 1995.

[23640] 2865.

[23641] Happle, R.: The McCune-Albright syndrome: a lethal gene surviving
[23642] by mosaicism. Clin. Genet. 29: 321-324, 1986.

[23643] 2866.

[23644] Harris, B. A.; Robishaw, J. D.; Mumby, S. M.; Gilman, A. G.:

[23645] Molecular cloning of complementary DNA for the alpha subunit of the
[23646] G protein that stimulates adenylate cyclase. Science 229: 1274-1277,
[23647] 1985.

[23648] 2867.

[23649] Hayward, B.; Kamiya, M.; Takada, S.; Moran, V.; Strain, L.; Hayashizaki,
[23650] Y.; Bonthron, D. T.: XL alpha s is a paternally derived pro-tein product
[23651] of the human GNAS1 gene. (Abstract) Europ. J. Hum. Genet. 6 (suppl.
[23652] 1): 36 only, 1998.

[23653] 2868.

[23654] Hayward, B. E.; Bonthron, D. T.: An imprinted antisense transcript

[23655] at the human GNAS1 locus. Hum. Molec. Genet. 9: 835-841, 2000.

[23656] 2869.

[23657] Hayward, B. E.; Kamiya, M.; Strain, L.; Moran, V.; Campbell, R.;

[23658] Hayashizaki, Y.; Bonthron, D. T.: The human GNAS1 gene is imprinted

[23659] and encodes distinct paternally and biallelically expressed G proteins. Proc.

[23660] Nat. Acad. Sci. 95: 10038-10043, 1998.

[23661] 2870.

[23662] Hayward, B. E.; Moran, V.; Strain, L.; Bonthron, D. T.: Bidirectional

[23663] imprinting of a single gene: GNAS1 encodes maternally, paternally,

[23664] and biallelically derived proteins. Proc. Nat. Acad. Sci. 95: 15475-15480,

[23665] 1998.

[23666] 2871.

[23667] Hurowitz, E. H.; Melnyk, J. M.; Chen, Y.-J.; Kouros-Mehr, H.;

[23668] Simon, M. I.; Shizuya, H.: Genomic characterization of the human

[23669] heterotrimeric G protein alpha, beta, and gamma subunit genes. DNA

[23670] Res. 7: 111-120, 2000.

[23671] 2872.

[23672] Iiri, T.; Farfel, Z.; Bourne, H. R.: Conditional activation de-fect

[23673] of a human Gs-alpha mutant. Proc. Nat. Acad. Sci. 94: 5656-5661,

[23674] 1997.

[23675] 2873.

[23676] Iiri, T.; Herzmark, P.; Nakamoto, J. M.; Van Dop, C.; Bourne,

[23677] H. R.: Rapid GDP release from Gs-alpha in patients with gain and

[23678] loss of endocrine function. Nature 371: 164-168, 1994.

[23679] 2874.

[23680] Ishikawa, Y.; Tajima, T.; Nakae, J.; Nagashima, T.; Satoh, K.;

[23681] Okuhara, K.; Fujieda, K.: Two mutations of the Gs-alpha gene in two

[23682] Japanese patients with sporadic pseudohypoparathy-roidism type Ia. J.

[23683] Hum. Genet. 46: 426-430, 2001.

[23684] 2875.

[23685] Jia, H.; Hingorani, A. D.; Sharma, P.; Hopper, R.; Dicker-son,

[23686] C.; Trutwein, D.; Lloyd, D. D.; Brown, M. J.: Association of the

[23687] G(s)-alpha gene with essential hypertension and response to beta-blockade. Hypertension 34:

[23688] 8-14, 1999.

[23689] 2876.

[23690] Kehlenbach, R. H.; Matthey, J.; Huttner, W. B.: XL alpha S is

[23691] a new type of G protein. Nature 372: 804-809, 1994.

[23692] 2877.

[23693] Kikyo, N.; Williamson, C. M.; John, R. M.; Barton, S. C.; Beechey,

[23694] C. V.; Ball, S. T.; Cattanach, B. M.; Surani, M. A.; Peters, J.:

[23695] Genetic and functional analysis of neuronatin in mice with maternal

[23696] or paternal duplication of distal chr. 2. Dev. Biol. 190: 66-77,

[23697] 1997.

[23698] 2878.

[23699] Kinard, R. E.; Walton, J. E.; Buckwalter, J. A.: Pseudohypoparathyroidism. Arch.

[23700] Intern. Med. 139: 204-207, 1979.

[23701] 2879.

[23702] Kozasa, T.; Itoh, H.; Tsukamoto, T.; Kaziro, Y.: Isolation and

[23703] characterization of the human Gs-alpha gene. Proc. Nat. Acad. Sci. 85:

[23704] 2081-2085, 1988.

[23705] 2880.

[23706] Juppner, H.; Schipani, E.; Bastepe, M.; Cole, D. E. C.; Lawson,

[23707] M. L.; Mannstadt, M.; Hendy, G. N.; Plotkin, H.; Koshiyama, H.; Koh,

[23708] T.; Crawford, J. D.; Olsen, B. R.; Vikkula, M.: The gene responsible
[23709] for pseudohypoparathyroidism type Ib is paternally imprinted and maps
[23710] in four unrelated kindreds to chromosome 20q13.3. Proc. Nat. Acad.
[23711] Sci. 95: 11798-11803, 1998.

[23712] 2881.

[23713] Landis, C. A.; Masters, S. B.; Spada, A.; Pace, A. M.; Bourne,
[23714] H. R.; Vallar, L.: GTPase inhibiting mutations activate the alpha
[23715] chain of Gs and stimulate adenylyl cyclase in human pituitary tumours. Nature 340:
[23716] 692-696, 1989.

[23717] 2882.

[23718] Levine, M. A.; Deily, J. R.: Identification of multiple mutations
[23719] in the gene encoding the alpha subunit of Gs in patients with pseudohypoparathyroidism
[23720] type IA. (Abstract) J. Bone Miner. Res. 5: S142 only, 1990.

[23721] 2883.

[23722] Levine, M. A.; Modi, W. S.; O'Brien, S. J.: Mapping of the gene

[23723] encoding the alpha subunit of the stimulatory G protein of adenylyl

[23724] cyclase (GNAS1) to 20q13.2-q13.3 in human by in situ hybridization. Genomics 11:

[23725] 478-479, 1991.

[23726] 2884.

[23727] Levine, M. A.; Vechio, J. D.: Personal Communication. Baltimore,

[23728] Md. 8/1/1990.

[23729] 2885.

[23730] Lin, C. K.; Hakakha, M. J.; Nakamoto, J. M.; Englund, A. T.; Brickman,

[23731] A. S.; Scott, M. L.; Van Dop, C.: Prevalence of three mutations in

[23732] the Gs-alpha gene among 24 families with pseudohypoparathyroidism

[23733] type Ia. Biochem. Biophys. Res. Commun. 189: 343-349, 1992.

[23734] 2886.

[23735] Linglart, A.; Carel, J. C.; Garabedian, M.; Le, T.; Mallet, E.;

[23736] Kottler, M. L.: GNAS1 lesions in pseudohypoparathy-
roidism Ia and

[23737] Ic: genotype phenotype relationship and evidence of the
maternal transmission

[23738] of the hormonal resistance. J. Clin. Endocr. Metab. 87:
189-197,

[23739] 2002.

[23740] 2887.

[23741] Malchoff, C. D.; Reardon, G.; MacGillivray, D. C.; Yamase,
H.;

[23742] Rogol, A. D.; Malchoff, D. M.: An unusual presentation of
McCune-Albright

[23743] syndrome confirmed by an activating mutation of the Gs
alpha-subunit

[23744] from a bone lesion. J. Clin. Endocr. Metab. 78: 803-806,
1994.

[23745] 2888.

[23746] Mantovani, G.; Romoli, R.; Weber, G.; Brunelli, V.; De Me-
nis,

[23747] E.; Beccio, S.; Beck-Peccoz, P.; Spada, A.: Mutational analysis of
[23748] GNAS1 in patients with pseudohypoparathyroidism: identification of
[23749] two novel mutations. J. Clin. Endocr. Metab. 85: 4243-4248, 2000.
[23750] 2889.

[23751] Falk, C. T.; Martin, M. D.; Walker, M. E.; Chen, T.; Rubinstein,
[23752] P.; Allen, F. H., Jr.: Family data suggesting a linkage between MN
[23753] and Gc. (Abstract) Cytogenet. Cell Genet. 25: 152, 1979.

[23754] 2890.

[23755] Furthmayr, H.; Metaxas, M. N.; Metaxas-Buhler, M.: M(g) and M(c):
[23756] mutations within the amino-terminal region of glycophorin A. Proc.
[23757] Nat. Acad. Sci. 78: 631-635, 1981.

[23758] 2891.

[23759] Gedde-Dahl, T., Jr.; Olaisen, B.: MN:Ss--GC more likely than

[23760] Ss:MN--GC? (Abstract) Cytogenet. Cell Genet. 32:
277-278, 1981.

[23761] 2892.

[23762] German, J.; Chaganti, R. S. K.: Mapping human autosomes:
assignment

[23763] of the MN locus to a specific segment in the long arm of
chromosome

[23764] no. 2. Science 182: 1261-1262, 1973.

[23765] 2893.

[23766] German, J.; Metaxas, M. N.; Metaxas-Buhler, M.; Louie, E.;
Chaganti,

[23767] R. S. K.: Further evaluation of a child with the M(k) pheno-
type and

[23768] a translocation affecting the long arms of chromosomes 2
and 4. (Abstract) Cytogenet.

[23769] Cell Genet. 25: 160, 1979.

[23770] 2894.

[23771] German, J.; Walker, M. E.; Steifel, F. H.; Allen, F. H., Jr.:

[23772] Autoradiographic studies of human chromosomes. II. Data
concerning

[23773] the position of the MN locus. Vox Sang. 16: 130-145,

1969.

[23774] 2895.

[23775] German, J.; Walker, M. E.; Stiefel, F. H.; Allen, F. H., Jr.:

[23776] MN blood-group locus: data concerning the possible chromosomal location. Science 162:

[23777] 1014-1015, 1968.

[23778] 2896.

[23779] Grant, S. G.; Bigbee, W. L.: Bone marrow somatic mutation after

[23780] genotoxic cancer therapy. (Letter) Lancet 343:
1507-1508, 1994.

[23781] 2897.

[23782] Heiberg, A.; Berg, K.: Linkage data on the MNSs blood group-red

[23783] cell acid phosphatase relationship. Hum. Hered. 25:
93-94, 1975.

[23784] 2898.

[23785] Huang, C.-H.; Chen, Y.; Blumenfeld, O. O.: A novel St(a) glycophorin

[23786] produced via gene conversion of pseudoexon III from gly-
cophorin E

[23787] to glycophorin A gene. Hum. Mutat. 15: 533-540, 2000.

[23788] 2899.

[23789] Huang, C.-H.; Guizzo, M. L.; Kikuchi, M.; Blumenfeld, O. O.:

[23790] Molecular genetic analysis of a hybrid gene encoding St(a) glycophorin

[23791] of the human erythrocyte membrane. Blood 74: 836-843, 1989.

[23792] 2900.

[23793] Huang, C.-H.; Puglia, K. V.; Bigbee, W. L.; Guizzo, M. L.; Hoffman,

[23794] M.; Blumenfeld, O. O.: A family study of multiple mutations of alpha

[23795] and delta glycophorins (glycophorins A and B). Hum. Genet. 81: 26-30,

[23796] 1988.

[23797] 2901.

[23798] Huang, C.-H.; Reid, M.; Daniels, G.; Blumenfeld, O. O.: Alteration

[23799] of splice site selection by an exon mutation in the human glycophorin

[23800] A gene. J. Biol. Chem. 268: 25902-25908, 1993.

[23801] 2902.

[23802] Huang, C.-H.; Spruell, P.; Moulds, J. J.; Blumenfeld, O. O.:

[23803] Molecular basis for the human erythrocyte glycophorin specifying the

[23804] Miltenberger class I (Mil) phenotype. Blood 80: 257-263, 1992.

[23805] 2903.

[23806] Kudo, S.; Chagnovich, D.; Rearden, A.; Mattei, M. G.; Fukuda,

[23807] M.: Molecular analysis of a hybrid gene encoding human glycophorin

[23808] variant Miltenberger V-like molecule. J. Biol. Chem. 265: 13825-13829,

[23809] 1990.

[23810] 2904.

[23811] Langlois, R. G.; Bigbee, W. L.; Jensen, R. H.: Measurements of

[23812] the frequency of human erythrocytes with gene expres- sion loss phenotypes

[23813] at the glycophorin A locus. Hum. Genet. 74: 353-362,

1986.

[23814] 2905.

[23815] Langlois, R. G.; Bigbee, W. L.; Kyoizumi, S.; Nakamura, N.; Bean,

[23816] M. A.; Akiyama, M.; Jensen, R. H.: Evidence for increased somatic

[23817] cell mutations at the glycophorin A locus in atomic bomb survivors. Science 236:

[23818] 445-448, 1987.

[23819] 2906.

[23820] Mattei, M. G.; London, J.; Rahuel, C.; d'Auriol, L.; Colin, Y.;

[23821] Le Van Kim, C.; Mattei, J. F.; Galibert, F.; Cartron, J. P.: Chromosome

[23822] localization by in situ hybridization of the gene for human erythrocyte

[23823] glycophorin to region 4q28-q31. (Abstract) Cytogenet. Cell Genet. 46:

[23824] 658, 1987.

[23825] 2907.

[23826] Mawby, W. J.; Anstee, D. J.; Tanner, M. J. A.: Immuno-chemical

[23827] evidence for hybrid sialoglycoproteins of human erythrocytes. Nature 291:
[23828] 161-162, 1981.

[23829] 2908.

[23830] Okubo, Y.; Daniels, G. L.; Parsons, S. F.; Anstee, D. J.; Yamaguchi,
[23831] H.; Tomita, T.; Seno, T.: A Japanese family with two sisters apparently
[23832] homozygous for M(k). Vox Sang. 54: 107-111, 1988.

[23833] 2909.

[23834] Onda, M.; Fukuda, M.: Detailed physical mapping of the genes
[23835] encoding glycophorins A, B, and E, as revealed by P1 plasmids containing
[23836] human genomic DNA. Gene 159: 225-230, 1995.

[23837] 2910.

[23838] Pasvol, G.; Wainscoat, J. S.; Weatherall, D. J.: Erythrocytes

[23839] deficient in glycophorin resist invasion by the malarial parasite
[23840] Plasmodium falciparum. Nature 297: 64-66, 1982.

[23841] 2911.

[23842] Prohaska, R.; Koerner, T. A. W., Jr.; Armitage, I. M.; Furth-
mayr,
[23843] H.: Chemical and carbon-13 nuclear magnetic resonance
studies of
[23844] the blood group M and N active sialoglycopeptides from
human glycophorin
[23845] A. J. Biol. Chem. 256: 5781-5791, 1986.

[23846] 2912.

[23847] Rahuel, C.; London, J.; d'Auriol, L.; Mattei, M.-G.; Tour-
namille,
[23848] C.; Skrzynia, C.; Lebouc, Y.; Galibert, F.; Cartron, J.-P.:
Characterization
[23849] of cDNA clones for human glycophorin A: use for gene lo-
calization
[23850] and for analysis of normal of glycophorin-A-deficient
(Finnish type)
[23851] genomic DNA. Europ. J. Biochem. 172: 147-153, 1988.

[23852] 2913.

[23853] Rahuel, C.; London, J.; Vignal, A.; Cherif-Zahar, B.; Colin,
Y.;

[23854] Siebert, P.; Fukuda, M.; Cartron, J.-P.: Alteration of the genes
[23855] for glycophorin A and B in glycophorin-A-deficient individuals. Europ.
[23856] J. Biochem. 177: 605-614, 1988.

[23857] 2914.

[23858] Rothman, N.; Haas, R.; Hayes, R. B.; Li, G.-L.; Wiemels, J.; Campleman,
[23859] S.; Quintana, P. J. E.; Xi, L.-J.; Dosemeci, M.; Titenko-Holland,
[23860] N.; Meyer, K. B.; Lu, W.; Zhang, L. P.; Bechtold, W.; Wang, Y.-Z.;
[23861] Kolachana, P.; Yin, S.-N.; Blot, W.; Smith, M. T.: Benzene induces
[23862] gene-duplicating but not gene-inactivating mutations at the glycophorin
[23863] A locus in exposed humans. Proc. Nat. Acad. Sci. 92: 4069-4073,
[23864] 1995.

[23865] 2915.

[23866] Siebert, P. D.; Fukuda, M.: Isolation and characterization of

[23867] human glycophorin A cDNA clones by a synthetic oligonu-
cleotide approach:
[23868] nucleotide sequence and mRNA structure. Proc. Nat. Acad.
Sci. 83:
[23869] 1665-1669, 1986.

[23870] 2916.

[23871] Spence, M. A.; Field, L. L.; Marazita, M. L.; Joseph, J.;
Sparkes,
[23872] M.; Crist, M.; Crandall, B. F.; Anderson, C. E.; Bateman, J.
B.; Rotter,
[23873] J. I.; Kidd, K. K.; Hodge, S. E.; Sparkes, R. S.: Estimating the
[23874] recombination frequency for the MN and the Ss loci. Hum.
Hered. 34:
[23875] 343-347, 1984.

[23876] 2917.

[23877] Springer, G. F.; Tegtmeyer, H.: Further evidence that car-
bohydrates
[23878] are the immunodeterminant structures of blood group M
and N specificities. Immun.
[23879] Commun. 10: 157-171, 1981.

[23880] 2918.

[23881] Egan, M. F.; Goldberg, T. E.; Kolachana, B. S.; Callicott, J. H.;
[23882] Mazzanti, C. M.; Straub, R. E.; Goldman, D.; Weinberger, D. R.: Effect
[23883] of COMT val(108/158)met genotype on frontal lobe function and risk
[23884] for schizophrenia. Proc. Nat. Acad. Sci. 98: 6917-6922, 2001.
[23885] 2919.

[23886] Floderus, Y.; Iselius, L.; Lindsten, J.; Wetterberg, L.: Evidence
[23887] for a major locus as well as a multifactorial component in the regulation
[23888] of human red blood cell catechol-O-methyl-transferase activity. Hum.
[23889] Hered. 32: 76-79, 1982.

[23890] 2920.

[23891] Floderus, Y.; Wetterberg, L.: The inheritance of human erythrocyte
[23892] catechol-O-methyltransferase activity. Clin. Genet. 19: 392-395,
[23893] 1981.

[23894] 2921.

[23895] Gogos, J. A.; Morgan, M.; Luine, V.; Santha, M.; Ogawa, S.; Pfaff,

[23896] D.; Karayiorgou, M.: Catechol-

0-methyltransferase-deficient mice

[23897] exhibit sexually dimorphic changes in catecholamine lev-

els and behavior. Proc.

[23898] Nat. Acad. Sci. 95: 9991-9996, 1998.

[23899] 2922.

[23900] Graf, W. D.; Unis, A. S.; Yates, C. M.; Sulzbacher, S.; Dinu-

los,

[23901] M. B.; Jack, R. M.; Dugaw, K. A.; Paddock, M. N.; Parson, W.

W.:

[23902] Catecholamines in patients with 22q11.2 deletion syn-

drome and the

[23903] low-activity COMT polymorphism. Neurology 57:

410-416, 2001.

[23904] 2923.

[23905] Grossman, M. H.; Emanuel, B. S.; Budarf, M. L.: Chromoso-

mal mapping

[23906] of the human catechol-O-methyltransferase gene to

22q11.1-q11.2. Genomics 12:

[23907] 822-825, 1992.

[23908] 2924.

[23909] Grossman, M. H.; Littrell, J.; Weinstein, R.; Punnett, H. H.;

[23910] Emanuel, B. S.; Budarf, M.: The gene for human catechol-
O-methyltransferase

[23911] (COMT) maps to 22pter-22q11.1. (Abstract) Cytogenet.
Cell Genet. 58:

[23912] 2048 only, 1991.

[23913] 2925.

[23914] Gustavson, K. H.; Floderus, Y.; Jagell, S.; Wetterberg, L.;
Ross,

[23915] S. B.: Catechol-O-methyltransferase activity in erythro-
cytes in Down's

[23916] syndrome: family studies. Clin. Genet. 22: 22-24, 1982.

[23917] 2926.

[23918] Gustavson, K. H.; Wetterberg, L.; Backstrom, M.; Ross, S.
B.:

[23919] Catechol-O-methyltransferase activity in erythrocytes in
Down's syndrome. Clin.

[23920] Genet. 4: 279-280, 1973.

[23921] 2927.

[23922] Hoda, F.; Nicholl, D.; Bennett, P.; Arranz, M.; Aitchison, K.

[23923] J.; Al-Chalabi, A.; Kunugi, H.; Vallada, H.; Leigh, P. N.; Chaudhuri,

[23924] K. R.; Collier, D. A.: No association between Parkinson's disease

[23925] and low-activity alleles of catechol O-methyltransferase. Biochem.

[23926] Biophys. Res. Commun. 228: 780-784, 1996.

[23927] 2928.

[23928] Karayiorgou, M.; Altemus, M.; Galke, B. L.; Goldman, D. Murphy,

[23929] D. L.; Ott, J.; Gogos, J. A.: Genotype determining low catechol-O-methyltransferase

[23930] activity as a risk factor for obsessive-compulsive disorder. Proc.

[23931] Nat. Acad. Sci. 94: 4572-4575, 1997.

[23932] 2929.

[23933] Lachman, H. M.; Morrow, B.; Shprintzen, R.; Veit, S.; Parsia,

[23934] S. S.; Faedda, G.; Goldberg, R.; Kucherlapati, R.; Papolos,

D. F.

[23935] : Association of codon 108/158 catechol-
O-methyltransferase gene polymorphism

[23936] with the psychiatric manifestations of velo-cardio-facial
syndrome. Am.

[23937] J. Med. Genet. 67: 468-472, 1996.

[23938] 2930.

[23939] Levitt, M.; Baron, M.: Human erythrocyte catechol-
O-transferase:

[23940] variation in thermal lability. (Abstract) Sixth Int. Cong.
Hum. Genet.,

[23941] Jerusalem 21 only, 1981.

[23942] 2931.

[23943] Lundstrom, K.; Salminen, M.; Jalanko, A.; Savolainen, R.;
Ulmanen,

[23944] I.: Cloning and characterization of human placental cate-
chol-O-methyltransferase

[23945] cDNA. DNA Cell Biol. 10: 181-189, 1991.

[23946] 2932.

[23947] Scanlon, P. D.; Raymond, F. A.; Weinshilboum, R. M.: Cate-
chol-O-methyltransferase:

[23948] thermolabile enzyme in erythrocytes of subjects homozygous for allele

[23949] for low activity. Science 203: 63-65, 1979.

[23950] 2933.

[23951] Siervogel, R. M.; Weinshilboum, R.; Wilson, A. F.; Elston, R.

[23952] C.: Major gene model for the inheritance of catechol-O-methyltransferase

[23953] activity in five large families. Am. J. Med. Genet. 19: 315-323,

[23954] 1984.

[23955] 2934.

[23956] Spielman, R. S.; Weinshilboum, R. M.: Genetics of red cell COMT

[23957] activity: analysis of thermal stability and family data. Am. J. Med.

[23958] Genet. 10: 279-290, 1981.

[23959] 2935.

[23960] Syvanen, A.-C.; Tilgmann, C.; Rinne, J.; Ulmanen, I.: Genetic

[23961] polymorphism of catechol-O-methyltransferase (COMT): correlation of

[23962] genotype with individual variation of S-COMT activity and comparison

[23963] of the allele frequencies in the normal population and parkinsonian

[23964] patients in Finland. Pharmacogenetics 7: 65-71, 1997.

[23965] 2936.

[23966] Weinshilboum, R.; Dunnette, J.: Thermal stability and the biochemical

[23967] genetics of erythrocyte catechol-O-methyltransferase and plasma dopamine-beta-hydroxylase. Clin.

[23968] Genet. 19: 426-437, 1981.

[23969] 2937.

[23970] Weinshilboum, R. M.; Raymond, F. A.: Inheritance of low erythrocyte

[23971] catechol-O-methyltransferase activity in man. Am. J. Hum. Genet. 29:

[23972] 125-135, 1977.

[23973] 2938.

[23974] Rouault, J.-P.; Rimokh, R.; Tessa, C.; Paranhos, G.; Ffrench, M.;

[23975] Duret, L.; Garoccio, M.; Germain, D.; Samarut, J.; Magaud,

J.-P.:

[23976] BTG1, a member of a new family of antiproliferative genes. EMBO J. 11:

[23977] 1663-1670, 1992.

[23978] 2939.

[23979] Hall, J. M.; Lee, M. K.; Newman, B.; Morrow, J. E.; Ander-son,

[23980] L. A.; Huey, B.; King, M.-C.: Linkage of early-onset familial breast

[23981] cancer to chromosome 17q21. Science 250: 1684-1689, 1990.

[23982] 2940.

[23983] Lucker, G. P. H.; van de Kerkhof, P. C. M.; Steijlen, P. M.: The

[23984] hereditary palmoplantar keratoses: an updated review and classification. Brit.

[23985] J. Derm. 131: 1-14, 1994.

[23986] 2941.

[23987] DiCroce, P. A.; Krontiris, T. G.: The BCL2 major breakpoint region

[23988] is a sequence- and cell-cycle-specific binding site of the

Ku antigen. Proc.
[23989] Nat. Acad. Sci. 92: 10137-10141, 1995.

[23990] 2942.

[23991] Seidman, C. E.: Personal Communication. Boston, Mass. 3/4/1992.

[23992] 2943.

[23993] Bachman, E. S.; Dhillon, H.; Zhang, C.-Y.; Cinti, S.; Bianco, A.
[23994] C.; Kobilka, B. K.; Lowell, B. B.: Beta-AR signaling required for
[23995] diet-induced thermogenesis and obesity resistance. Science 297:
[23996] 843-845, 2002.

[23997] 2944.

[23998] Benovic, J. L.; Stone, W. C.; Huebner, K.; Croce, C.; Caron, M.
[23999] G.; Lefkowitz, R. J.: cDNA cloning and chromosomal localization of
[24000] the human beta-adrenergic receptor kinase. FEBS Lett. 283: 122-126,
[24001] 1991.

[24002] 2945.

[24003] Harding, V. B.; Jones, L. R.; Lefkowitz, R. J.; Koch, W. J.; Rockman,

[24004] H. A.: Cardiac beta-ARK1 inhibition prolongs survival and augments

[24005] beta blocker therapy in a mouse model of severe heart failure. Proc.

[24006] Nat. Acad. Sci. 98: 5809-5814, 2001.

[24007] 2946.

[24008] Penn, R. B.; Benovic, J. L.: Structure of the human gene en-coding

[24009] the beta-adrenergic receptor kinase. J. Biol. Chem. 269: 14924-14930,

[24010] 1994.

[24011] 2947.

[24012] Rockman, H. A.; Chien, K. R.; Choi, D.-J.; Iaccarino, G.; Hunter,

[24013] J. J.; Ross, J., Jr.; Lefkowitz, R. J.; Koch, W. J.: Expression of

[24014] a beta-adrenergic receptor kinase 1 inhibitor prevents the development

[24015] of myocardial failure in gene-targeted mice. Proc. Nat.

Acad. Sci. 95:
[24016] 7000-7005, 1998.

[24017] 2948.

[24018] Spurney, R. F.; Flannery, P. J.; Garner, S. C.; Athirakul, K.;

[24019] Liu, S.; Guilak, F.; Quarles, L. D.: Anabolic effects of a G protein-coupled
[24020] receptor kinase inhibitor expressed in osteoblasts. J. Clin. Invest. 109:
[24021] 1361-1371, 2002.

[24022] 2949.

[24023] Hamann, U.; Haner, M.; Stosiek, U.; Bastert, G.; Scott, R. J.

[24024] : Low frequency of BRCA1 germline mutations in 45 Ger-man breast/ovarian
[24025] cancer families. J. Med. Genet. 34: 884-888, 1997.

[24026] 2950.

[24027] Harkin, D. P.; Bean, J. M.; Miklos, D.; Song, Y.-H.; Truong, V.
[24028] B.; Englert, C.; Christians, F. C.; Ellisen, L. W.; Maheswaran, S.;
[24029] Oliner, J. D.; Haber, D. A.: Induction of GADD45 and JNK/

SAPK-dependent

[24030] apoptosis following inducible expression of BRCA1. Cell 97: 575-586,

[24031] 1999.

[24032] 2951.

[24033] Hartman, A.-R.; Ford, J. M.: BRCA1 induces DNA damage recognition

[24034] factors and enhances nucleotide excision repair. Nature Genet. 32:

[24035] 180-184, 2002.

[24036] 2952.

[24037] Hedenfalk, I.; Duggan, D.; Chen, Y.; Radmacher, M.; Bit-tner, M.;

[24038] Simon, R.; Meltzer, P.; Gusterson, B.; Esteller, M.; Kallion-iemi,

[24039] O.-P.; Wilfond, B.; Borg, A.; Trent, J.: Gene-expression profiles

[24040] in hereditary breast cancer. New Eng. J. Med. 344: 539-548, 2001.

[24041] 2953.

[24042] Hogervorst, F. B. L.; Cornelis, R. S.; Bout, M.; van Vliet, M.;

[24043] Oosterwijk, J. C.; Olmer, R.; Bakker, B.; Klijn, J. G. M.; Vasen,
[24044] H. F. A.; Meijers-Heijboer, H.; Menko, F. H.; Cornelisse, C. J.; den
[24045] Dunnen, J. T.; Devilee, P.; van Ommen, G.-J. B.: Rapid detection
[24046] of BRCA1 mutations by the protein truncation test. Nature Genet. 10:
[24047] 208-212, 1995.

[24048] 2954.

[24049] Holt, J. T.; Thompson, M. E.; Szabo, C.; Robinson-Benion, C.;
[24050] Arteaga, C. L.; King, M.-C.; Jensen, R. A.: Growth retardation and
[24051] tumour inhibition by BRCA1. Nature Genet. 12: 298-302, 1996.
[24052] 2955.

[24053] Huttley, G. A.; Easteal, S.; Southey, M. C.; Tesoriero, A.; Giles,
[24054] G. G.; McCredie, M. R. E.; Hopper, J. L.; Venter, D. J.; Australian
[24055] Breast Cancer Family Study: Adaptive evolution of the tu-

mour suppressor

[24056] BRCA1 in humans and chimpanzees. Nature Genet. 25: 410-413, 2000.

[24057] 2956.

[24058] Isaacs, S. D.; Kiemeney, L. A. L. M.; Baffoe-Bonnie, A.; Beaty,

[24059] T. H.; Walsh, P. C.: Risk of cancer in relatives of prostate cancer

[24060] probands. J. Nat. Cancer Inst. 87: 991-996, 1995.

[24061] 2957.

[24062] Ishioka, C.; Suzuki, T.; Fitzgerald, M.; Krainer, M.; Shi-modaira,

[24063] H.; Shimada, A.; Nomizu, T.; Isselbacher, K. J.; Haber, D.; Kanamaru,

[24064] R.: Detection of heterozygous truncating mutations in the BRCA1 and

[24065] APC genes by using a rapid screening assay in yeast. Proc. Nat. Acad.

[24066] Sci. 94: 2449-2453, 1997.

[24067] 2958.

[24068] James, V.; Kearsley, J.; Irving, T.; Amemiya, Y.; Cookson,

D.

[24069] : Using hair to screen for breast cancer. Nature 398: 33-34, 1999.

[24070] 2959.

[24071] Janezic, S. A.; Ziogas, A.; Krumroy, L. M.; Krasner, M.; Plummer,

[24072] S. J.; Cohen, P.; Gildea, M.; Barker, D.; Haile, R.; Casey, G.; Anton-Culver,

[24073] H.: Germline BRCA1 alterations in a population-based series of ovarian

[24074] cancer cases. Hum. Molec. Genet. 8: 889-897, 1999.

[24075] 2960.

[24076] Jensen, R. A.; Thompson, M. E.; Jetton, T. L.; Szabo, C. I.; van

[24077] der Meer, R.; Helou, B.; Tronick, S. R.; Page, D. L.; King, M.-C.;

[24078] Holt, J. T.: BRCA1 is secreted and exhibits properties of a granin. Nature

[24079] Genet. 12: 303-308, 1996.

[24080] 2961.

[24081] Jin, Y.; Xu, X. L.; Yang, M.-C. W.; Wei, F.; Ayi, T.-C.; Bow-

cock,

[24082] A. M.; Baer, R.: Cell cycle-dependent colocalization of BARD1 and

[24083] BRCA1 proteins in discrete nuclear domains. Proc. Nat. Acad. Sci. 94:

[24084] 12075-12080, 1997.

[24085] 2962.

[24086] Johannsson, O.; Ostermeyer, E. A.; Hakansson, S.; Friedman, L.

[24087] S.; Johansson, U.; Sellberg, G.; Brondum-Nielsen, K.; Sele, V.; Olsson,

[24088] H.; King, M.-C.; Borg, A.: Founding BRCA1 mutations in hereditary

[24089] breast and ovarian cancer in Southern Sweden. Am. J. Hum. Genet. 58:

[24090] 441-450, 1996.

[24091] 2963.

[24092] Kauff, N. D.; Satagopan, J. M.; Robson, M. E.; Scheuer, L.; Hensley,

[24093] M.; Hudis, C. A.; Ellis, N. A.; Boyd, J.; Borgen, P. I.; Barakat,

[24094] R. R.; Norton, L.; Offit, K.: Risk-reducing salpingo-

oophorectomy

[24095] in women with a BRCA1 or BRCA2 mutation. New Eng. J. Med. 346: 1609-1615,

[24096] 2002.

[24097] 2964.

[24098] Kelsell, D. P.; Black, D. M.; Bishop, D. T.; Spurr, N. K.: Genetic

[24099] analysis of the BRCA1 region in a large breast/ovarian family: refinement

[24100] of the minimal region containing BRCA1. Hum. Molec. Genet. 2: 1823-1828,

[24101] 1993.

[24102] 2965.

[24103] Khavari, P. A.; Peterson, C. L.; Tamkun, J. W.; Mendel, D. B.;

[24104] Crabtree, G. R.: BRG1 contains a conserved domain of the SWI2/SNF2

[24105] family necessary for normal mitotic growth and transcription. Nature 366:

[24106] 170-174, 1993.

[24107] 2966.

[24108] Krainer, M.; Silva-Arrieta, S.; Fitzgerald, M. G.; Shimada, A.;
[24109] Ishioka, C.; Kanamaru, R.; MacDonald, D. J.; Unsal, H.; Finkelstein,
[24110] D. M.; Bowcock, A.; Isselbacher, K. J.; Haber, D. A.: Differ-ential
[24111] contributions of BRCA1 and BRCA2 to early-onset breast cancer. New
[24112] Eng. J. Med. 336: 1416-1421, 1997.

[24113] 2967.

[24114] Langston, A. A.; Malone, K. E.; Thompson, J. D.; Daling, J. R.;
[24115] Ostrander, E. A.: BRCA1 mutations in a population-based sample of
[24116] young women with breast cancer. New Eng. J. Med. 334: 137-142, 1996.
[24117] 2968.

[24118] Langston, A. A.; Stanford, J. L.; Wicklund, K. G.; Thomp-son, J.
[24119] D.; Blazej, R. G.; Ostrander, E. A.: Germ-line BRCA1 muta-tions in
[24120] selected men with prostate cancer. (Letter) Am. J. Hum.

Genet. 58:
[24121] 881-885, 1996.

[24122] 2969.

[24123] Lee, C. G.; Hurwitz, J.: Human RNA helicase A is homolo-
gous to
[24124] the maleless protein of Drosophila. J. Biol. Chem. 268:
16822-16830,
[24125] 1993.

[24126] 2970.

[24127] Lee, J.-S.; Collins, K. M.; Brown, A. L.; Lee, C.-H.; Chung, J.

[24128] H.: hCds1-mediated phosphorylation of BRCA1 regulates
the DNA damage
[24129] response. Nature 404: 201-204, 2000.

[24130] 2971.

[24131] Li, S.; Ting, N. S. Y.; Zheng, L.; Chen, P.-L.; Ziv, Y.; Shiloh,

[24132] Y.; Lee, E. Y.-H. P.; Lee, W.-H.: Functional link of BRCA1
and ataxia
[24133] telangiectasia gene product in DNA damage response.
Nature 406:
[24134] 210-215, 2000.

[24135] 2972.

[24136] Liede, A.; Malik, I. A.; Aziz, Z.; de los Rios, P.; Kwan, E.;

[24137] Narod, S. A.: Contribution of BRCA1 and BRCA2 mutations to breast

[24138] and ovarian cancer in Pakistan. Am. J. Hum. Genet. 71: 595-606,

[24139] 2002.

[24140] 2973.

[24141] Liede, A.; Rehal, P.; Vesprini, D.; Jack, E.; Abrahamson, J.;

[24142] Narod, S. A.: A breast cancer patient of Scottish descent with germ-line

[24143] mutations in BRCA1 and BRCA2. (Letter) Am. J. Hum. Genet. 62: 1543-1544,

[24144] 1998.

[24145] 2974.

[24146] Liu, H.-X.; Cartegni, L.; Zhang, M. Q.; Krainer, A. R.: A mechanism

[24147] for exon skipping caused by nonsense or missense muta-tions in BRCA1

[24148] and other genes. Nature Genet. 27: 55-58, 2001.

[24149] 2975.

[24150] Lorick, K. L.; Jensen, J. P.; Fang, S.; Ong, A. M.;
Hatakeyama,
[24151] S.; Weissman, A. M.: RING fingers mediate ubiquitin-
conjugating enzyme
[24152] (E2)-dependent ubiquitination. Proc. Nat. Acad. Sci. 96:
11364-11369,
[24153] 1999.

[24154] 2976.

[24155] Ludwig, T.; Chapman, D. L.; Papaioannou, V. E.; Efstra-
tiadis,
[24156] A.: Targeted mutations of breast cancer susceptibility
gene homologs
[24157] in mice: lethal phenotypes of Brca1, Brca2, Brca1/Brca2,
Brca1/p53,
[24158] and Brca2/p53 nullizygous embryos. Genes Dev. 11:
1226-1241, 1997.
[24159] 2977.

[24160] Lynch, H. T.; Watson, P.: Genetic counselling and heredi-
tary
[24161] breast/ovarian cancer. (Letter) Lancet 339: 1181 only,
1992.

[24162] 2978.

[24163] Liu, G.; Maunoury, C.; Kamoun, P.; Aral, B.: Assignment of the

[24164] human gene encoding the delta-1-pyrroline-5-carboxylate synthetase

[24165] (P5CS) to 10q24.3 by in situ hybridization. Genomics 37: 145-146,

[24166] 1996.

[24167] 2979.

[24168] Rabier, D.; Nuttin, C.; Poggi, F.; Padovani, J. P.; Abdo, K.; Bardet,

[24169] J.; Parvy, P.; Kamoun, P.; Saudubray, J. M.: Familial joint hyperlaxity,

[24170] skin hyperelasticity, cataract and mental retardation with hyperammonemia

[24171] and low citrulline, ornithine and proline. A new disorder of collagen

[24172] metabolism?In: Abstracts of Free Communications, 30th Annual Symposium

[24173] Leuven, Sept. 8-11, 1992. Pp. 61. SSIEM: The Society for the Study

[24174] of Inborn Errors of Metabolism, Leuven. 1992.

[24175] 2980.

[24176] Hess, S. D.; Daggett, L. P.; Crona, J.; Deal, C.; Lu, C.-C.; Urrutia,

[24177] A.; Chavez-Noriega, L.; Ellis, S. B.; Johnson, E. C.; Velicelebi,

[24178] G.: Cloning and functional characterization of human heteromeric

[24179] N-methyl-D-aspartate receptors. J. Pharm. Exp. Ther. 278: 808-816,

[24180] 1996.

[24181] 2981.

[24182] Setou, M.; Nakagawa, T.; Seog, D.-H.; Hirokawa, N.: Kinesin superfamily

[24183] motor protein KIF17 and mLin-10 in NMDA receptor-containing vesicle

[24184] transport. Science 288: 1796-1802, 2000.

[24185] 2982.

[24186] Sprengel, R.; Suchanek, B.; Amico, C.; Brusa, R.; Burnashev, N.;

[24187] Rozov, A.; Hvalby, O.; Jensen, V.; Paulsen, O.; Andersen, P.; Kim,

[24188] J. J.; Thompson, R. F.; Sun, W.; Webster, L. C.; Grant, S. G. N.;

[24189] Eilers, J.; Konnerth, A.; Li, J.; McNamara, J. O.; Seeburg, P. H.

[24190] : Importance of the intracellular domain of NR2 subunits for NMDA

[24191] receptor function in vivo. Cell 92: 279-289, 1998.

[24192] 2983.

[24193] Ishii, T.; Moriyoshi, K.; Sugihara, H.; Sakurada, K.; Kadotani,

[24194] H.; Yokoi, M.; Akazawa, C.; Shigemoto, R.; Mizuno, N.; Masu, M.; Nakanishi,

[24195] S.: Molecular characterization of the family of the N-methyl-D-aspartate

[24196] receptor subunits. J. Biol. Chem. 268: 2836-2843, 1993.

[24197] 2984.

[24198] Kalsi, G.; Whiting, P.; Le Bourdelles, B.; Callen, D.; Barnard,

[24199] E. A.; Gurling, H.: Localization of the human NMDAR2D receptor subunit

[24200] gene (GRIN2D) to 19q13.1-qter, the NMDAR2A subunit gene to 16p13.2

[24201] (GRIN2A), and the NMDAR2C subunit gene (GRIN2C) to 17q24-q25 using
[24202] somatic cell hybrid and radiation hybrid mapping panels. Genomics 47:
[24203] 423-425, 1998.

[24204] 2985.

[24205] Monyer, H.; Sprengel, R.; Schoepfer, R.; Herb, A.; Higuchi, M.;
[24206] Lomeli, H.; Burnashev, N.; Sakmann, B.; Seeburg, P. H.: Heteromeric
[24207] NMDA receptors: molecular and functional distinction of subtypes. Science 256:
[24208] 1217-1221, 1992.

[24209] 2986.

[24210] Sakimura, K.; Kutsuwada, T.; Ito, I.; Manabe, T.; Takayama, C.;
[24211] Kushiya, E.; Yagi, T.; Aizawa, S.; Inoue, Y.; Sugiyama, H.; Mishina,
[24212] M.: Reduced hippocampal LTP and spatial learning in mice lacking
[24213] NMDA receptor epsilon 1 subunit. Nature 373: 151-155, 1995.

[24214] 2987.

[24215] Wakui, K.; Nishida, T.; Masuda, J.; Itoh, T.; Katsumata, D.; Ohno,

[24216] T.; Fukushima, Y.: De novo interstitial deletion of 4q[46,XX,del(4)(q27q28.2)]

[24217] with intact blood group-MN locus, confining its locus to 4q28.2-4q31.1. Jpn.

[24218] J. Hum. Genet. 36: 149-153, 1991.

[24219] 2988.

[24220] Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Weinshilboum,

[24221] R.; Ward, L. J.: Linkage relationships between a major gene for catechol-O-methyltransferase

[24222] activity and 25 polymorphic marker systems. Am. J. Med. Genet. 19:

[24223] 525-532, 1984.

[24224] 2989.

[24225] Mattera, R.; Graziano, M. P.; Yatani, A.; Zhou, Z.; Graf, R.;

[24226] Codina, J.; Birnbaumer, L.; Gilman, A. G.; Brown, A. M.: Splice variants

[24227] of the alpha subunit of the G protein G(8) activate both adenylyl

[24228] cyclase and calcium channels. Science 243: 804-807, 1989.

[24229] 2990.

[24230] Mehlmann, L. M.; Jones, T. L. Z.; Jaffe, L. A.: Meiotic arrest

[24231] in the mouse follicle maintained by a GS protein in the oocyte. Science 297:

[24232] 1343-1345, 2002.

[24233] 2991.

[24234] Nerlich, A.; Peschel, O.; Lohrs, U.; Parsche, F.; Betz, P.: Juvenile

[24235] gigantism plus polyostotic fibrous dysplasia in the Tegernsee giant.

[24236] (Letter) Lancet 338: 886-887, 1991.

[24237] 2992.

[24238] Patten, J. L.; Johns, D. R.; Valle, D.; Eil, C.; Gruppuso, P.

[24239] A.; Steele, G.; Smallwood, P. M.; Levine, M. A.: Mutation in the

[24240] gene encoding the stimulatory G protein of adenylate cyclase in Albright's

[24241] hereditary osteodystrophy. New Eng. J. Med. 322: 1412-1419, 1990.

[24242] 2993.

[24243] Persani, L.; Borgato, S.; Lania, A.; Filopanti, M.; Mantovani,

[24244] G.; Conti, M.; Spada, A.: Relevant cAMP-specific phospho-
diesterase

[24245] isoforms in human pituitary: effect of Gs-alpha mutations.
J. Clin.

[24246] Endocr. Metab. 86: 3795-3800, 2001.

[24247] 2994.

[24248] Peters, J.; Beechey, C. V.; Ball, S. T.; Evans, E. P.: Mapping

[24249] studies of the distal imprinting region of mouse chromo-
some 2. Genet.

[24250] Res. 63: 169-174, 1994.

[24251] 2995.

[24252] Premawardhana, L. D. K. E.; Vora, J. P.; Mills, R.; Scanlon,
M.

[24253] F.: Acromegaly and its treatment in the McCune-Albright
syndrome. Clin.

[24254] Endocr. 36: 605-608, 1992.

[24255] 2996.

[24256] Riminucci, M.; Fisher, L. W.; Majolagbe, A.; Corsi, A.; Lala,

[24257] R.; De Sanctis, C.; Robey, P. G.; Bianco, P.: A novel GNAS1 mutation,

[24258] R201G, in McCune-Albright syndrome. J. Bone Miner. Res. 14: 1987-1989,

[24259] 1999.

[24260] 2997.

[24261] Schwindinger, W. F.; Francomano, C. A.; Levine, M. A.: Identification

[24262] of a mutation in the gene encoding the alpha subunit of the stimulatory

[24263] G-protein of adenylyl cyclase in McCune-Albright syndrome. Proc.

[24264] Nat. Acad. Sci. 89: 5152-5156, 1992.

[24265] 2998.

[24266] Schwindinger, W. F.; Francomano, C. A.; Levine, M. A.; McKusick,

[24267] V. A.: DNA light on the Tegernsee giant. (Letter) Lancet 338: 1454-1455,

[24268] 1991.

[24269] 2999.

[24270] Shapira, H.; Mouallem, M.; Shapiro, M. S.; Weisman, Y.;

Farfel,
[24271] Z.: Pseudohypoparathyroidism type Ia: two new heterozygous frameshift
[24272] mutations in exons 5 and 10 of the Gs alpha gene. Human Genet. 97:
[24273] 73-75, 1995.

[24274] 3000.

[24275] Shenker, A.; Chanson, P.; Weinstein, L. S.; Chi, P.; Spiegel,

[24276] A. M.; Lomri, A.; Marie, P. J.: Osteoblastic cells derived from isolated
[24277] lesions of fibrous dysplasia contain activating somatic mutations
[24278] of the G(S)-alpha gene. Hum. Molec. Genet. 4:
1675-1676, 1995.
[24279] 3001.

[24280] Shenker, A.; Weinstein, L. S.; Moran, A.; Pescovitz, O. H.; Charest,
[24281] N. J.; Boney, C. M.; Van Wyk, J. J.; Merino, M. J.; Feuillan, P. P.;
[24282] Spiegel, A. M.: Severe endocrine and nonendocrine manifestations
[24283] of the McCune-Albright syndrome associated with activat-

ing mutations
[24284] of stimulatory G protein Gs. J. Pediat. 123: 509-518, 1993.
[24285] 3002.
[24286] Shore, E. M.; Ahn, J.; Jan de Beur, S.; Li, M.; Xu, M.; Gar-diner,
[24287] R. J. M.; Zasloff, M. A.; Whyte, M. P.; Levine, M. A.; Kaplan, F.
[24288] S.: Paternally inherited inactivating mutations of the GNAS1 gene
[24289] in progressive osseous heteroplasia. New Eng. J. Med. 346: 99-106,
[24290] 2002.
[24291] 3003.
[24292] Shore, E. M.; Kaplan, F. S.; Levine, M. A.: GNAS1 mutations and
[24293] progressive osseous heteroplasia. (Letter) New Eng. J. Med. 346:
[24294] 1670-1671, 2002.
[24295] 3004.
[24296] Sparkes, R. S.; Cohn, V. H.; Mohandas, T.; Zollman, S.;

Cire-Eversole,
[24297] P.; Amatruda, T. T.; Reed, R. R.; Lochrie, M. A.; Simon, M. I.: Mapping
[24298] of genes encoding the subunits of guanine nucleotide-binding protein
[24299] (G-proteins) in humans. (Abstract) Cytogenet. Cell Genet. 46: 696
[24300] only, 1987.

[24301] 3005.

[24302] Tinschert, S.; Gerl, H.; Gewies, A.; Jung, H.-P.; Nurnberg, P.
[24303] : McCune-Albright syndrome: clinical and molecular evidence of mosaicism
[24304] in an unusual giant patient. Am. J. Med. Genet. 83: 100-108, 1999.
[24305] 3006.

[24306] Vallar, L.; Spada, A.; Giannattasio, G.: Altered Gs and adenylate
[24307] cyclase activity in human GH-secreting pituitary adenomas. Nature 330:
[24308] 566-568, 1987.

[24309] 3007.

[24310] Warner, D. R.; Gejman, P. V.; Collins, R. M.; Weinstein, L. S.

[24311] : A novel mutation adjacent to the switch III domain of Gs-alpha in

[24312] a patient with pseudohypoparathyroidism. Molec. Endocr. 11: 1718-1727,

[24313] 1997.

[24314] 3008.

[24315] Warner, D. R.; Weinstein, L. S.: A mutation in the heterotrimeric

[24316] stimulatory guanine nucleotide binding protein alpha-subunit with

[24317] impaired receptor-mediated activation because of elevated GTPase activity. Proc.

[24318] Nat. Acad. Sci. 96: 4268-4272, 1999.

[24319] 3009.

[24320] Warner, D. R.; Weng, G.; Yu, S.; Matalon, R.; Weinstein, L. S.

[24321] : A novel mutation in the switch 3 region of Gs-alpha in a patient

[24322] with Albright hereditary osteodystrophy impairs GDP binding and receptor

[24323] activation. J. Biol. Chem. 273: 23976-23983, 1998.

[24324] 3010.

[24325] Weinstein, L. S.; Gejman, P. V.; de Mazancourt, P.; Ameri-
can,
[24326] N.; Spiegel, A. M.: A heterozygous 4-bp deletion mutation
in the
[24327] Gs-alpha gene (GNAS1) in a patient with Albright heredi-
tary osteodystrophy. Genomics 13:
[24328] 1319-1321, 1992.

[24329] 3011.

[24330] Weinstein, L. S.; Gejman, P. V.; Friedman, E.; Kadowaki, T.;
Collins,
[24331] R. M.; Gershon, E. S.; Spiegel, A. M.: Mutations of the Gs
alpha-subunit
[24332] gene in Albright hereditary osteodystrophy detected by
denaturing
[24333] gradient gel electrophoresis. Proc. Nat. Acad. Sci. 87:
8287-8290,
[24334] 1990.

[24335] 3012.

[24336] Weinstein, L. S.; Shenker, A.; Gejman, P. V.; Merino, M. J.;

Friedman,
[24337] E.; Spiegel, A. M.: Activating mutations of the stimulatory G protein
[24338] in the McCune-Albright syndrome. New Eng. J. Med. 325: 1688-1695,
[24339] 1991.

[24340] 3013.

[24341] Weinstein, L. S.; Yu, S.: The role of genomic imprinting of GS-alpha
[24342] in the pathogenesis of Albright hereditary osteodystrophy. Trends
[24343] Endocr. Metab. 10: 81-85, 1999.

[24344] 3014.

[24345] Walker, M. E.; Rubinstein, P.; Allen, F. H., Jr.: Biochemical

[24346] genetics of MN. Vox Sang. 32: 111-120, 1977.

[24347] 3015.

[24348] Weitkamp, L. R.; Adams, M. S.; Rowley, P. T.: Linkage be-tween
[24349] the MN and Hb beta loci. Hum. Hered. 22: 566-572, 1972.

[24350] 3016.

[24351] Weitkamp, L. R.; Lovrien, E. W.; Olaisen, B.; Fenger, K.; Gedde-Dahl,

[24352] T., Jr.; Sorensen, S. A.; Conneally, P. M.; Bias, W. B.; Ott, J.:

[24353] Linkage relations of the loci for the MN blood group and red cell

[24354] phosphate. Birth Defects Orig. Art. Ser. XI(3): 276-280, 1975. Note:

[24355] Alternate: Cytogenet. Cell Genet. 14: 446-450, 1975.....

[24356] 3017.

[24357] Hansen, R. L.; Marx, J. J.; Ptacek, L. J.; Roberts, R. C.: Immunological

[24358] studies on an aberrant form of ataxia-telangiectasia. Am. J. Dis.

[24359] Child. 131: 518-521, 1977.

[24360] 3018.

[24361] Brown, K. D.; Barlow, C.; Wynshaw-Boris, A.: Multiple ATM-dependent

[24362] pathways: an explanation for pleiotropy. (Editorial) Am. J. Hum.

[24363] Genet. 64: 46-50, 1999.

[24364] 3019.

[24365] Brown, K. D.; Ziv, Y.; Sadanandan, S. N.; Chessa, L.; Collins,

[24366] F. S.; Shiloh, Y.; Tagle, D. A.: The ataxia-telangiectasia gene product,

[24367] a constitutively expressed nuclear protein that is not up-regulated

[24368] following genome damage. Proc. Nat. Acad. Sci. 94: 1840-1845, 1997.

[24369] 3020.

[24370] Brzoska, P. M.; Chen, H.; Zhu, Y.; Levin, N. A.; Disatnik, M.-H.;

[24371] Mochly-Rosen, D.; Murnane, J. P.; Christman, M. F.: The product of

[24372] the ataxia-telangiectasia group D complementing gene, ATDC, interacts

[24373] with a protein kinase C substrate and inhibitor. Proc. Nat. Acad.

[24374] Sci. 92: 7824-7828, 1995.

[24375] 3021.

[24376] Byrd, P. J.; McConville, C. M.; Cooper, P.; Parkhill, J.; Stankovic,

[24377] T.; McGuire, G. M.; Thick, J. A.; Taylor, A. M. R.: Mutations

revealed

[24378] by sequencing the 5-prime half of the gene for ataxia telangiectasia. Hum.

[24379] Molec. Genet. 5: 145-149, 1996.

[24380] 3022.

[24381] Byrne, E.; Hallpike, J. F.; Manson, J. F.; Sutherland, G. R.;

[24382] Thong, Y. H.: Ataxia-without-telangiectasia: progressive multisystem

[24383] degeneration with IgE deficiency and chromosome instability. J. Neurol.

[24384] Sci. 66: 307-317, 1984.

[24385] 3023.

[24386] Canman, C. E.; Lim, D.-S.; Cimprich, K. A.; Taya, Y.; Tamai, K.;

[24387] Sakaguchi, K.; Appella, E.; Kastan, M. B.; Siliciano, J. D.: Activation

[24388] of the ATM kinase by ionizing radiation and phosphorylation of p53. Science 281:

[24389] 1677-1679, 1998.

[24390] 3024.

[24391] Carbonari, M.; Cherchi, M.; Paganelli, R.; Giannini, G.;

Galli,
[24392] E.; Gaetano, C.; Papetti, C.; Fiorilli, M.: Relative increase of
[24393] T cells expressing the gamma/delta rather than the al-
pha/beta receptor
[24394] in ataxia-telangiectasia. New Eng. J. Med. 322: 73-76,
1990.
[24395] 3025.
[24396] Castellvi-Bel, S.; Sheikhavandi, S.; Telatar, M.; Tai, L.-Q.;
[24397] Hwang, M.; Wang, Z.; Yang, Z.; Cheng, R.; Gatti, R. A.: New
mutations,
[24398] polymorphisms, and rare variants in the ATM gene de-
tected by a novel
[24399] SSCP strategy. Hum. Mutat. 14: 156-162, 1999.
[24400] 3026.
[24401] Chen, P.; Imray, F. P.; Kidson, C.: Gene dosage and com-
plementation
[24402] analysis of ataxia telangiectasia lymphoblastoid cell lines
assayed
[24403] by induced chromosome aberrations. Mutat. Res. 129:
165-172, 1984.
[24404] 3027.

[24405] Chong, M. J.; Murray, M. R.; Gosink, E. C.; Russell, H. R. C.;

[24406] Srinivasan, A.; Kapsetaki, M.; Korsmeyer, S. J.; McKinnon,
P. J.:

[24407] Atm and Bax cooperate in ionizing radiation-induced
apoptosis in the

[24408] central nervous system. Proc. Nat. Acad. Sci. 97: 889-894,
2000.

[24409] 3028.

[24410] Chou, P. Y.; Fasman, G. D.: Prediction of the secondary
structure

[24411] of proteins from their amino acid sequence. Adv. Enzymol.
Relat.

[24412] Areas Molec. Biol. 47: 45-148, 1978.

[24413] 3029.

[24414] Cohen, M. M.; Sagi, M.; Ben-Zur, Z.; Schaap, T.; Voss, R.;
Kohn,

[24415] G.; Ben-Bassat, H.: Ataxia telangiectasia: chromosomal
stability

[24416] in continuous lymphoblastoid cell lines. Cytogenet. Cell
Genet. 23:

[24417] 44-52, 1979.

[24418] 3030.

[24419] Cohen, M. M.; Shaham, M.; Dagan, J.; Shmueli, E.; Kohn, G.: Cytogenetic

[24420] investigations in families with ataxia-telangiectasia. Cyto-genet.

[24421] Cell Genet. 15: 338-356, 1975.

[24422] 3031.

[24423] Concannon, P.; Gatti, R. A.: Diversity of ATM gene muta-tions

[24424] detected in patients with ataxia-telangiectasia. Hum. Mu-tat. 10:

[24425] 100-107, 1997.

[24426] 3032.

[24427] Cooper, D. N.; Youssoufian, H.: The CpG dinucleotide and human

[24428] genetic disease. Hum. Genet. 78: 151-155, 1988. 1. Al Saadi, A.;

[24429] Palutke, M.; Krishna Kumar, G.: Evolution of chromosomal abnormalities

[24430] in sequential cytogenetic studies of ataxia telangiectasia. Hum.

[24431] Genet. 55: 23-29, 1980.

[24432] 3033.

[24433] Cornforth, M. N.; Bedford, J. S.: On the nature of a defect in

[24434] cells from individuals with ataxia-telangiectasia. Science 227:

[24435] 1589-1591, 1985.

[24436] 3034.

[24437] Cortez, D.; Wang, Y.; Qin, J.; Elledge, S. J.: Requirement of

[24438] ATM-dependent phosphorylation of Brca1 in the DNA damage response

[24439] to double-strand breaks. Science 286: 1162-1166, 1999.

[24440] 3035.

[24441] Cox, R.; Hosking, G. P.; Wilson, J.: Ataxia telangiectasia: evaluation

[24442] of radiosensitivity in cultured skin fibroblasts as a diag- nostic test. Arch.

[24443] Dis. Child. 53: 386-390, 1978.

[24444] 3036.

[24445] Curry, C. J. R.; Tsai, J.; Hutchinson, H. T.; Jaspers, N. G. J.;

[24446] Wara, D.; Gatti, R. A.: AT-Fresno: a phenotype linking ataxia-telangiectasia
[24447] with the Nijmegen breakage syndrome. Am. J. Hum. Genet. 45: 270-275,
[24448] 1989.

[24449] 3037.

[24450] DeLeon, G. A.; Grover, W. D.; Huff, D. S.: Neuropathologic changes
[24451] in ataxia-telangiectasia. Neurology 26: 947-951, 1976.

[24452] 3038.

[24453] Easton, D. F.: Cancer risks in A-T heterozygotes. Int. J. Rad.
[24454] Biol. 66: S177-S182, 1994.

[24455] 3039.

[24456] Ejima, Y.; Oshimura, M.; Sasaki, M. S.: Establishment of a novel
[24457] immortalized cell line from ataxia-telangiectasia fibrob-lasts and
[24458] its use for the chromosomal assignment of radiosensitivity gene. Int.
[24459] J. Radiat. Biol. 58: 989-997, 1990.

[24460] 3040.

[24461] Ejima, Y.; Sasaki, M. S.: Mutations of the ATM gene de-
tected
[24462] in Japanese ataxia-telangiectasia patients: possible pre-
ponderance
[24463] of the two founder mutations 4612del165 and 7883del5.
Hum. Genet. 102:
[24464] 403-408, 1998.

[24465] 3041.

[24466] Elson, A.; Wang, Y.; Daugherty, C. J.; Morton, C. C.; Zhou,
F.;
[24467] Campos-Torres, J.; Leder, P.: Pleiotropic defects in ataxia-
telangiectasia
[24468] protein-deficient mice. Proc. Nat. Acad. Sci. 93:
13084-13089, 1996.
[24469] 3042.

[24470] Falck, J.; Mailand, N.; Syljuasen, R. G.; Bartek, J.; Lukas, J.

[24471] : The ATM-Chk2-Cdc25A checkpoint pathway guards
against radioresistant

[24472] DNA synthesis. Nature 410: 842-847, 2001.

[24473] 3043.

[24474] Falck, J.; Petrini, J. H. J.; Williams, B. R.; Lukas, J.; Bartek,

[24475] J.: The DNA damage-dependent intra-S phase checkpoint is regulated

[24476] by parallel pathways. Nature Genet. 30: 290-294, 2002.

[24477] 3044.

[24478] Feigin, R. D.; Vietti, T. J.; Wyatt, R. G.; Kaufman, D. G.; Smith,

[24479] C. H.: Ataxia telangiectasia with granulocytopenia. J. Pediat. 77:

[24480] 431-438, 1970.

[24481] 3045.

[24482] Fiorilli, M.; Antonelli, A.; Russo, G.; Crescenzi, M.; Carbonari,

[24483] M.; Petrinelli, P.: Variant of ataxia-telangiectasia with low-level

[24484] radiosensitivity. Hum. Genet. 70: 274-277, 1985.

[24485] 3046.

[24486] Fiorilli, M.; Businco, L.; Pandolfi, F.; Paganelli, R.; Russo,

[24487] G.; Aiuti, F.: Heterogeneity of immunological abnormalities in ataxia-telangiectasia. J.

[24488] Clin. Immun. 3: 135-141, 1983.

[24489] 3047.

[24490] Kanters, J. K.; Larsen, L. A.; Orholm, M.; Agner, E.; Anderson,

[24491] P. S.; Vuust, J.; Christiansen, M.: Novel donor splice site mutation

[24492] in the KVLQT1 gene is associated with the long QT syndrome. J. Cardiovasc.

[24493] Electrophysiol. 9: 620-624, 1998.

[24494] 3048.

[24495] Itoh, T.; Kikuchi, K.; Odagawa, Y.; Takata, S.; Yano, K.; Okada,

[24496] S.; Haneda, N.; Ogawa, S.; Nakano, O.; Kawahara, Y.; Kasai, H.; Nakayama,

[24497] T.; Fukutomi, T.; Sakurada, H.; Shimizu, A.; Yazaki, Y.; Nagai, R.;

[24498] Nakamura, Y.; Tanaka, T.: Correlation of genetic etiology with response

[24499] to beta-adrenergic blockade among symptomatic patients with familial

[24500] long-QT syndrome. J. Hum. Genet. 46: 38-40, 2001.

[24501] 3049.

[24502] Keating, M.: Response to Benhorin et al. (1993). Science 260:

[24503] 1962 only, 1993.

[24504] 3050.

[24505] Keating, M.; Atkinson, D.; Dunn, C.; Timothy, K.; Vincent, G.

[24506] M.; Leppert, M.: Linkage of a cardiac arrhythmia, the long QT syndrome,

[24507] and the Harvey RAS-1 gene. Science 252: 704-706, 1991.

[24508] 3051.

[24509] Keating, M.; Atkinson, D.; Dunn, C.; Timothy, K.; Vincent, G.

[24510] M.; Leppert, M.: Long QT syndrome is closely linked to the Harvey

[24511] RAS-1 gene on chromosome 11. (Abstract) Clin. Res. 39: 317A only,

[24512] 1991.

[24513] 3052.

[24514] Keating, M.; Dunn, C.; Atkinson, D.; Timothy, K.; Vincent, G.

[24515] M.; Leppert, M.: Consistent linkage of the long-QT syndrome to the

[24516] Harvey Ras-1 locus on chromosome 11. Am. J. Hum. Genet. 49: 1335-1339,

[24517] 1991.

[24518] 3053.

[24519] Kerem, B.; Benhorin, J.; Kalman, Y. M.; Medina, A.; Dyer, T. D.;

[24520] Blangero, J.; MacCluer, J. W.: Evidence for genetic heterogeneity

[24521] in the long QT syndrome. (Abstract) Am. J. Hum. Genet. 51 (suppl.):

[24522] A192 only, 1992.

[24523] 3054.

[24524] Klein, H. O.; Levi, A.; Kaplinsky, E.; Di Segni, E.; David, D.

[24525] : Congenital long-QT syndrome: deleterious effect of long-term high-rate

[24526] ventricular pacing and definitive treatment by cardiac transplantation. Am.

[24527] Heart J. 132: 1079-1081, 1996.

[24528] 3055.

[24529] Ko, Y.-L.; Chen, S.-A.; Tang, T. K.; Lin, J.-L.; Chiang, C.-E.;

[24530] Chen, J.-J.; Teng, M.-S.; Chang, M.-S.; Lien, W.-P.; Wu, C.-W.: No

[24531] evidence for linkage of long QT syndrome and chromosome 11p15.5 markers

[24532] in a Chinese family: evidence for genetic heterogeneity. Hum. Genet. 94:

[24533] 364-366, 1994.

[24534] 3056.

[24535] Kukolich, M. K.; Telsey, A.; Ott, J.; Motulsky, A. G.: Sudden

[24536] infant death syndrome: normal QT intervals on ECGs of relatives. Pediatrics 60:

[24537] 51-54, 1977.

[24538] 3057.

[24539] Larsen, L. A.; Fosdal, I.; Andersen, P. S.; Kanters, J. K.; Vuust,

[24540] J.; Wettrell, G.; Christiansen, M.: Recessive Romano-Ward syndrome

[24541] associated with compound heterozygosity for two mutations in the KVLQT1
[24542] gene. Europ. J. Hum. Genet. 7: 724-728, 1999.

[24543] 3058.

[24544] Lee, M. P.; et al.; et al.: Loss of imprinting of a paternally
[24545] expressed transcript, with antisense orientation to KVLQT1, occurs
[24546] frequently in Beckwith-Wiedemann syndrome and is independent of insulin-like
[24547] growth factor II imprinting. Proc. Nat. Acad. Sci. 96: 5203-5208,
[24548] 1999.

[24549] 3059.

[24550] Lee, M. P.; Hu, R.-J.; Johnson, L. A.; Feinberg, A. P.: Human
[24551] KVLQT1 gene shows tissue-specific imprinting and encompasses Beckwith-Wiedemann
[24552] syndrome chromosomal rearrangements. Nature Genet. 15: 181-185,
[24553] 1997.

[24554] 3060.

[24555] Li, H.; Chen, Q.; Moss, A. J.; Robinson, J.; Goytia, V.; Perry,

[24556] J. C.; Vincent, G. M.; Priori, S. G.; Lehmann, M. H.; Den-
field, S.

[24557] W.; Duff, D.; Kaine, S.; Shimizu, W.; Schwartz, P. J.; Wang,
Q.; Towbin,

[24558] J. A.: New mutations in the KVLQT1 potassium channel
that cause long

[24559] QT syndrome. Circulation 97: 1264-1269, 1998.

[24560] 3061.

[24561] Locati, E. H.; Zareba, W.; Moss, A. J.; Schwartz, P. J.; Vin-
cent,

[24562] G. M.; Lehmann, M. H.; Towbin, J. A.; Priori, S. G.; Napoli-
tano, C.;

[24563] Robinson, J. L.; Andrews, M.; Timothy, K.; Hall, W. J.: Age-
and

[24564] sex-related differences in clinical manifestations in pa-
tients with

[24565] congenital long-QT syndrome. Circulation 97:
2237-2244, 1998.

[24566] 3062.

[24567] Mannens, M.; Wilde, A.: KVLQT1, the rhythm of imprint-
ing. Nature

[24568] Genet. 15: 113-115, 1997.

[24569] 3063.

[24570] Martini, B.: Personal Communication. Thiene, Italy 11/9/1998.

[24571] 3064.

[24572] Melki, J.; Kaplan, J.; Lucet, V.; Halley, L.; Clemenceau, S.;

[24573] Kaplan, C.; Baule, M. S.; Frezal, J.: Long QT (Romano-Ward) syndrome:

[24574] further definition of the linkage to the HLA loci. (Abstract) Cytogenet.

[24575] Cell Genet. 46: 661 only, 1987.

[24576] 3065.

[24577] Milne, J. R.; Ward, D. E.; Spurrell, R. A. J.; Camm, A. J.: The

[24578] long QT syndrome: effects of drugs and left stellate ganglion block. Am.

[24579] Heart J. 104: 194-198, 1982.

[24580] 3066.

[24581] Mitsutake, A.; Takeshita, A.; Kuroiwa, A.; Nakamura, M.: Usefulness

[24582] of the Valsalva maneuver in management of the long QT

syndrome. Circulation 63:
[24583] 1029-1035, 1981.

[24584] 3067.

[24585] Moss, A. J.; McDonald, J.: Unilateral cervico-thoracic sym-
pathetic
[24586] ganglionectomy for the treatment of long Q-T interval
syndrome. New
[24587] Eng. J. Med. 285: 903-904, 1971.

[24588] 3068.

[24589] Moss, A. J.; Schwartz, P. J.: Sudden death and the idio-
pathic
[24590] long Q-T syndrome. (Editorial) Am. J. Med. 66: 6-7, 1979.

[24591] 3069.

[24592] Moss, A. J.; Schwartz, P. J.; Crampton, R. S.; Tzivoni, D.;
Locati,
[24593] E. H.; MacCluer, J.; Hall, W. J.; Weitkamp, L.; Vincent, G. M.;
Garson,
[24594] A., Jr.; Robinson, J. L.; Benhorin, J.; Choi, S.: The long QT
syndrome:
[24595] prospective longitudinal study of 328 families. Circulation
84:

[24596] 1136-1144, 1991.

[24597] 3070.

[24598] Murray, A.; Donger, C.; Fenske, C.; Spillman, I.; Richard, P.;

[24599] Dong, Y. B.; Neyroud, N.; Chevalier, P.; Denjoy, I.; Carter, N.; Syrris,

[24600] P.; Afzal, A. P.; Patton, M. A.; Guicheney, P.; Jeffery, S.: Splicing

[24601] mutations in KCNQ1: a mutation hot spot at codon 344 that produces

[24602] in frame transcripts. Circulation 100: 1077-1084, 1999.

[24603] 3071.

[24604] Neyroud, N.; Denjoy, I.; Donger, C.; Gary, F.; Villain, E.; Leenhardt,

[24605] A.; Benali, K.; Schwartz, K.; Coumel, P.; Guicheney, P.: Heterozygous

[24606] mutation in the pore of potassium channel gene KvLQT1 causes an apparently

[24607] normal phenotype in long QT syndrome. Europ. J. Hum. Genet. 6: 129-133,

[24608] 1998.

[24609] 3072.

[24610] Neyroud, N.; Richard, P.; Vignier, N.; Donger, C.; Denjoy, I.;

[24611] Demay, L.; Shkolnikova, M.; Pesce, R.; Chevalier, P.; Hainque, B.;

[24612] Coumel, P.; Schwartz, K.; Guicheney, P.: Genomic organi-zation of

[24613] the KCNQ1 K+ channel gene and identification of C-terminal mutations

[24614] in the long-QT syndrome. Circ. Res. 84: 290-297, 1999.

[24615] 3073.

[24616] Sato, T.; Saito, H.; Swensen, J.; Olifant, A.; Wood, C.; Dan-ner,

[24617] D.; Sakamoto, T.; Takita, K.; Kasumi, F.; Miki, Y.; Skolnick, M.;

[24618] Nakamura, Y.: The human prohibitin gene located on chromosome 17q21

[24619] is mutated in sporadic breast cancer. Cancer Res. 52: 1643-1646,

[24620] 1992.

[24621] 3074.

[24622] Heald, R.; McLoughlin, M.; McKeon, F.: Human wee1 maintains mitotic
[24623] timing by protecting the nucleus from cytoplasmically activated cdc2
[24624] kinase. Cell 74: 463-474, 1993.

[24625] 3075.

[24626] Igarashi, M.; Nagata, A.; Jinno, S.; Suto, K.; Okayama, H.: Wee1(+)-like
[24627] gene in human cells. Nature 353: 80-83, 1991.

[24628] 3076.

[24629] Spranger, S.; Tariverdian, G.: Symptomatic heterozygosity in the
[24630] Ellis-van-Creveld syndrome? Clin. Genet. 47: 217-220, 1995.
[24631] 3077.

[24632] Jupe, E. R.; Badgett, A. A.; Neas, B. R.; Craft, M. A.; Mitchell,
[24633] D. S.; Resta, R.; Mulvihill, J. J.; Aston, C. E.; Thompson, L. F.
[24634] : Single nucleotide polymorphism in prohibitin 3-prime untranslated

[24635] region and breast-cancer susceptibility. Lancet 357:
1588-1589,
[24636] 2001.

[24637] 3078.

[24638] Jupe, E. R.; Liu, X. T.; Kiehlbauch, J. L.; et al.: Prohibitin

[24639] in breast cancer cell lines: loss of antiproliferative activity
is
[24640] linked to 3-prime untranslated region mutations. Cell
Growth Diff. 7:
[24641] 871-878, 1996.

[24642] 3079.

[24643] Sato, T.; Sakamoto, T.; Takita, K.-I.; Saito, H.; Okui, K.;
Nakamura,
[24644] Y.: The human prohibitin (PHB) gene family and its so-
matic mutations
[24645] in human tumors. Genomics 17: 762-764, 1993.

[24646] 3080.

[24647] White, J. J.; Ledbetter, D. H.; Eddy, R. L., Jr.; Shows, T. B.;

[24648] Stewart, D. A.; Nuell, M. J.; Friedman, V.; Wood, C. M.;
Owens, G.

[24649] A.; McClung, J. K.; Danner, D. B.; Morton, C. C.: Assign-
ment of the
[24650] human prohibition (sic) gene (PHB) to chromosome 17 and
identification
[24651] of a DNA polymorphism. Genomics 11: 228-230, 1991.

[24652] 3081.

[24653] Annunen, P.; Helaakoski, T.; Myllyharju, J.; Veijola, J.; Pih-
lajaniemi,
[24654] T.; Kivirikko, K. I.: Cloning of the human prolyl
4-hydroxylase alpha
[24655] subunit isoform alpha(II) and characterization of the type
II enzyme
[24656] tetramer: the alpha(I) and alpha(II) subunits do not form a
mixed
[24657] alpha(I)alpha(II)beta2 tetramer. J. Biol. Chem. 272:
17342-17348,
[24658] 1997.

[24659] 3082.

[24660] Friedman, L.; Higgin, J. J.; Moulder, G.; Barstead, R.;
Raines,
[24661] R. T.; Kimble, J.: Prolyl 4-hydroxylase is required for via-
bility

[24662] and morphogenesis in Caenorhabditis elegans. Proc. Nat. Acad. Sci. 97:

[24663] 4736-4741, 2000.

[24664] 3083.

[24665] Helaakoski, T.; Annunen, P.; Vuori, K.; MacNeil, I. A.; Pihlajaniemi,

[24666] T.; Kivirikko, K. I.: Cloning, baculovirus expression, and characterization

[24667] of a second mouse prolyl 4-hydroxylase alpha-subunit isoform: formation

[24668] of an alpha(2)beta(2) tetramer with the protein disulfide-isomerase/beta

[24669] subunit. Proc. Nat. Acad. Sci. 92: 4427-4431, 1995.

[24670] 3084.

[24671] Helaakoski, T.; Vuori, K.; Myllyla, R.; Kivirikko, K. I.; Pihlajaniemi,

[24672] T.: Molecular cloning of the alpha-subunit of human prolyl 4-hydroxylase:

[24673] the complete cDNA-derived amino acid sequence and evidence for alternative

[24674] splicing of RNA transcripts. Proc. Nat. Acad. Sci. 86: 4392-4396,

[24675] 1989.

[24676] 3085.

[24677] Pajunen, L.; Jones, T. A.; Helaakoski, T.; Pihlajaniemi, T.; Solomon,

[24678] E.; Sheer, D.; Kivirikko, K. I.: Assignment of the gene coding for

[24679] the alpha-subunit of prolyl 4-hydroxylase to human chromosome region

[24680] 10q21.3-23.1. Am. J. Hum. Genet. 45: 829-834, 1989.

[24681] 3086.

[24682] Helaakoski, T.; Veijola, J.; Vuori, K.; Rehn, M.; Chow, L. T.;

[24683] Taillon-Miller, P.; Kivirikko, K. I.; Pihlajaniemi, T.: Structure

[24684] and expression of the human gene for the alpha subunit of prolyl 4-hydroxylase:

[24685] the two alternatively spliced types of mRNA correspond to two homologous

[24686] exons the sequences of which are expressed in a variety of tissues. J.

[24687] Biol. Chem. 269: 27847-27854, 1994.

[24688] 3087.

[24689] Zelinski, T.; Coghlan, G.; Myal, Y.; Shiu, R. P. C.; Philipps,

[24690] S.; White, L.; Lewis, M.: Genetic linkage between the Kell blood

[24691] group system and prolactin-inducible protein loci: provisional assignment

[24692] of KEL to chromosome 7. Ann. Hum. Genet. 55: 137-140, 1991.

[24693] 3088.

[24694] Abney, M.; Ober, C.; McPeek, M. S.: Quantitative-trait homozygosity

[24695] and association mapping and empirical genomewide significance in large,

[24696] complex pedigrees: fasting serum-insulin level in the Hutterites. Am.

[24697] J. Hum. Genet. 70: 920-934, 2002.

[24698] 3089.

[24699] Barbetti, F.; Raben, N.; Kadowaki, T.; Cama, A.; Accili, D.; Gabbay,

[24700] K. H.; Merenich, J. A.; Taylor, S. I.; Roth, J.: Two unrelated patients

[24701] with familial hyperproinsulinemia due to a mutation substituting histidine

[24702] for arginine at position 65 in the proinsulin molecule: identification

[24703] of the mutation by direct sequencing of genomic deoxyribonucleic acid

[24704] amplified by polymerase chain reaction. J. Clin. Endocr. Metab. 71:

[24705] 164-169, 1990.

[24706] 3090.

[24707] Bell, G. I.; Pictet, R. L.; Rutter, W. J.; Cordell, B.; Tischer,

[24708] E.; Goodman, H. M.: Sequence of the human insulin gene. Nature 284:

[24709] 26-32, 1980.

[24710] 3091.

[24711] Bell, G. I.; Swain, W. F.; Pictet, R.; Cordell, B.; Goodman, H.

[24712] M.; Rutter, W. J.: Nucleotide sequence of cDNA clone encoding human

[24713] preproinsulin. Nature 282: 525-527, 1979.

[24714] 3092.

[24715] Carroll, R. J.; Hammer, R. E.; Chan, S. J.; Swift, H. H.; Rubenstein,

[24716] A. H.; Steiner, D. F.: A mutant human proinsulin is secreted from
[24717] islets of Langerhans in increased amounts via an unregulated pathway. Proc.
[24718] Nat. Acad. Sci. 85: 8943-8947, 1988.

[24719] 3093.

[24720] Chaganti, R. S. K.; Jhanwar, S. C.; Antonarakis, S. E.; Hayward,
[24721] W. S.: Germ-line chromosomal localization of genes in chromosome
[24722] 11p linkage; parathyroid hormone, beta-globin, c-Ha-ras-1, and insulin. Somat.
[24723] Cell Molec. Genet. 11: 197-202, 1985.

[24724] 3094.

[24725] Chan, S. J.; Seino, S.; Gruppuso, P. A.; Schwartz, R.; Steiner,
[24726] D. F.: A mutation in the B chain coding region is associated with
[24727] impaired proinsulin conversion in a family with hyperproinsulinemia. Proc.
[24728] Nat. Acad. Sci. 84: 2194-2197, 1987.

[24729] 3095.

[24730] Collinet, M.; Berthelon, M.; Benit, P.; Laborde, K.; Des-
buquois,
[24731] B.; Munnich, A.; Robert, J. J.: Familial hyperproinsuli-
naemia due
[24732] to a mutation substituting histidine for arginine at posi-
tion 65 in
[24733] proinsulin: identification of the mutation by restriction
enzyme mapping. Europ.
[24734] J. Pediat. 157: 456-460, 1998.

[24735] 3096.

[24736] Dandona, P.; Aljada, A.; Mohanty, P.; Ghanim, H.;
Hamouda, W.;
[24737] Assian, E.; Ahmad, S.: Insulin inhibits intranuclear nuclear
factor
[24738] kappa-B and stimulates I-kappa-B in mononuclear cells in
obese subjects:
[24739] evidence for an anti-inflammatory effect? J. Clin. Endocr.
Metab. 86:
[24740] 3257-3265, 2001.

[24741] 3097.

[24742] Bhalla, U. S.; Ram, P. T.; Iyengar, R.: MAP kinase phos-
phatase
[24743] as a locus of flexibility in a mitogen-activated protein ki-
nase signaling
[24744] network. Science 297: 1018-1023, 2002.

[24745] 3098.

[24746] Boulton, T. G.; Nye, S. H.; Robbins, D. J.; Ip, N. Y.;
Radziejewska,
[24747] E.; Morgenbesser, S. D.; DePinho, R. A.; Panayotatos, N.;
Cobb, M.
[24748] H.; Yancopoulos, G. D.: ERKs: a family of protein-ser-
ine/threonine
[24749] kinases that are activated and tyrosine phosphorylated in
response
[24750] to insulin and NGF. Cell 65: 663-675, 1991.

[24751] 3099.

[24752] Cobb, M. H.; Boulton, T. G.; Robbins, D. J.: Extracellular
signal-regulated
[24753] kinases: ERKs in progress. Cell Regul. 2: 965-978, 1991.

[24754] 3100.

[24755] Di Cristo, G.; Berardi, N.; Cancedda, L.; Pizzorusso, T.;

Putignano,
[24756] E.; Ratto, G. M.; Maffei, L.: Requirement of ERK activation for visual
[24757] cortical plasticity. Science 292: 2337-2340, 2001.

[24758] 3101.

[24759] Forcet, C.; Stein, E.; Pays, L.; Corset, V.; Llambi, F.; Tessier-Lavigne,
[24760] M.; Mehlen, P.: Netrin-1-mediated axon outgrowth requires deleted
[24761] in colorectal cancer-dependent MAPK activation. Nature 417: 443-447,
[24762] 2002.

[24763] 3102.

[24764] Khokhlatchev, A. V.; Canagarajah, B.; Wilsbacher, J.; Robinson,
[24765] M.; Atkinson, M.; Goldsmith, E.; Cobb, M. H.: Phosphorylation of
[24766] the MAP kinase ERK2 promotes its homodimerization and nuclear translocation. Cell 93:
[24767] 605-615, 1998.

[24768] 3103.

[24769] Li, L.; Wysk, M.; Gonzalez, F. A.; Davis, R. J.: Genomic loci

[24770] of human mitogen-activated protein kinases. Oncogene 9: 647-649,

[24771] 1994.

[24772] 3104.

[24773] Owaki, H.; Makar, R.; Boulton, T. G.; Cobb, M. H.; Geppert, T.

[24774] D.: Extracellular signal-regulated kinases in T cells: characterization

[24775] of human ERK1 and ERK2 cDNAs. Biochem. Biophys. Res. Commun. 182:

[24776] 1416-1422, 1992.

[24777] 3105.

[24778] Saba-El-Leil, M. K.; Malo, D.; Meloche, S.: Chromosomal localization

[24779] of the mouse genes encoding the ERK1 and ERK2 isoforms of MAP kinases. Mammalian

[24780] Genome 8: 141-142, 1997.

[24781] 3106.

[24782] Stefanovsky, V. Y.; Pelletier, G.; Hannan, R.; Gagnon-Kugler,

[24783] T.; Rothblum, L. I.; Moss, T.: An immediate response of ri-
bosomal
[24784] transcription to growth factor stimulation in mammals is
mediated
[24785] by ERK phosphorylation of UBF. Molec. Cell 8: 1063-1073,
2001.
[24786] 3107.

[24787] Thomas, G.: MAP kinase by any other name smells just as
sweet. Cell 68:
[24788] 3-6, 1992.

[24789] 3108.

[24790] Garcia, J. I.; Zalba, G.; Detera-Wadleigh, S. D.; de Miguel,
C.
[24791] : Isolation of a cDNA encoding the rat MAP-kinase ho-
molog of human
[24792] p63mapk. Mammalian Genome 7: 810-814, 1996.

[24793] 3109.

[24794] Gonzalez, F. A.; Raden, D. L.; Rigby, M. R.; Davis, R. J.:
Heterogeneous
[24795] expression of four MAP kinase isoforms in human tissues.
FEBS Lett. 304:

[24796] 170-178, 1992.

[24797] 3110.

[24798] Zhu, A. X.; Zhao, Y.; Moller, D. E.; Flier, J. S.: Cloning and

[24799] characterization of p97(MAPK), a novel human homolog of rat ERK-3. Molec.

[24800] Cell. Biol. 14: 8202-8211, 1994.

[24801] 3111.

[24802] Linnenbach, A. J.; Huebner, K.; Reddy, E. P.; Herlyn, M.; Parmiter,

[24803] A. H.; Nowell, P. C.; Koprowski, H.: Structural alteration in the

[24804] MYB protooncogene and deletion within the gene encoding alpha-type

[24805] protein kinase C in human melanoma cell lines. Proc. Nat. Acad. Sci. 85:

[24806] 74-78, 1988.

[24807] 3112.

[24808] Bayes, M.; Valverde, D.; Balcells, S.; Grinberg, D.; Vilageliu,

[24809] L.; Benitez, J.; Ayuso, C.; Beneyto, M.; Baiget, M.; Gonzalez-Duarte,

[24810] R.: Evidence against involvement of recoverin in autosomal recessive
[24811] retinitis pigmentosa in 42 Spanish families. Hum. Genet. 96: 89-94,
[24812] 1995.

[24813] 3113.

[24814] Dizhoor, A. M.; Ray, S.; Kumar, S.; Niemi, G.; Spencer, M.; Brolley,
[24815] D.; Walsh, K. A.; Philipov, P. P.; Hurley, J. B.; Stryer, L.: Recoverin:
[24816] a calcium sensitive activator of retinal rod guanylate cyclase. Science 251:
[24817] 915-918, 1991.

[24818] 3114.

[24819] Kanazawa, Y.; Hayashi, M.; Ikeuchi, M.; Kasuga, M.; Oka, Y.; Sato,
[24820] H.; Hiramatsu, K.; Kosaka, K.: Familial proinsulinemia: a rare disorder
[24821] of insulin biosynthesis.In: Baba, S.; Kaneko, T.; Yanaihara, N.:
[24822] Proc. Symp. on proinsulin, insulin, and C-peptide. Tokushima, Japan,

[24823] 1978. Amsterdam: Excerpta Medica (pub.) 1979. Pp. 262-269.

[24824] 3115.

[24825] Kayo, T.; Koizumi, A.: Mapping of murine diabetogenic gene Mody

[24826] on chromosome 7 at D7Mit258 and its involvement in pancreatic islet

[24827] and beta cell development during the perinatal period. J. Clin. Invest. 101:

[24828] 2112-2118, 1998.

[24829] 3116.

[24830] Kwok, S. C. M.; Chan, S. J.; Rubenstein, A. H.; Poucher, R.; Steiner,

[24831] D. F.: Loss of a restriction endonuclease cleavage site in the gene

[24832] of a structurally abnormal human insulin. Biochem. Biophys. Res.

[24833] Commun. 98: 844-849, 1981.

[24834] 3117.

[24835] Kwok, S. C. M.; Steiner, D. F.; Rubenstein, A. H.; Tager, H. S.

[24836] : Identification of a point mutation in the human insulin gene giving
[24837] rise to a structurally abnormal insulin (insulin Chicago). Diabetes 32:
[24838] 872-875, 1983.

[24839] 3118.

[24840] Lebo, R. V.; Chakravarti, A.; Buetow, K. H.; Cheung, M.-C.; Cann,
[24841] H.; Cordell, B.; Goodman, H.: Recombination within and between the
[24842] human insulin and beta-globin gene loci. Proc. Nat. Acad. Sci. 80:
[24843] 4808-4812, 1983.

[24844] 3119.

[24845] Lebo, R. V.; Kan, Y. W.; Cheung, M. C.; Carrano, A. V.; Yu, L.-C.;
[24846] Chang, J. C.; Cordell, B.; Goodman, H. M.: Assigning the polymorphic
[24847] human insulin gene to the short arm of chromosome 11 by chromosome
[24848] sorting. Hum. Genet. 60: 10-15, 1982.

[24849] 3120.

[24850] Lebo, R. V.; Kan, Y. W.; Cheung, M. C.; Cordell, B.; Goodman,

[24851] H. M.; Law, M. L.; Jones, C.; Kao, F. T.: Assignment of the human

[24852] insulin gene to chromosome 11 band p11 and linkage analysis with the

[24853] beta-globin locus. (Abstract) Am. J. Hum. Genet. 33: 150A, 1981.

[24854] 3121.

[24855] Le Stunff, C.; Fallin, D.; Schork, N. J.; Bougneres, P.: The

[24856] insulin gene VNTR is associated with fasting insulin levels and development

[24857] of juvenile obesity. Nature Genet. 26: 444-446, 2000. Note: Erratum:

[24858] Nature Genet. 28: 97 only, 2001.

[24859] 3122.

[24860] Lichter, P.; Tang, C. C.; Call, K.; Hermanson, G.; Evans, G. A.;

[24861] Housman, D.; Ward, D. C.: High-resolution mapping of human chromosome

[24862] 11 by in situ hybridization with cosmid clones. Science 247: 64-69,

[24863] 1990.

[24864] 3123.

[24865] Premkumar, L. S.; Ahern, G. P.: Induction of vanilloid receptor

[24866] channel activity by protein kinase C. Nature 408: 985-990, 2000.

[24867] 3124.

[24868] Huppi, K.; Siwarski, D.; Goodnight, J.; Mischak, H.: Assignment

[24869] of the protein kinase C delta-polypeptide gene (PRKCD) to human chromosome

[24870] 3 and mouse chromosome 14. Genomics 19: 161-162, 1994.

[24871] 3125.

[24872] Mecklenbrauker, I.; Saijo, K.; Zheng, N. Y.; Leitges, M.; Tarakhovsky,

[24873] A.: Protein kinase C-delta controls self-antigen-induced B-cell tolerance. Nature 416:

[24874] 860-865, 2002.

[24875] 3126.

[24876] Mischak, H.; Bodenteich, A.; Kolch, W.; Goodnight, J.; Hofer, F.;

[24877] Mushinski, J. F.: Mouse protein kinase C-delta, the major isoform

[24878] expressed in mouse hemopoietic cells: sequence of the cDNA, expression

[24879] patterns and characterization of the protein. Biochemistry 30: 7925-7931,

[24880] 1991.

[24881] 3127.

[24882] Miyamoto, A.; Nakayama, K.; Imaki, H.; Hirose, S.; Jiang, Y.; Abe,

[24883] M.; Tsukiyama, T.; Nagahama, H.; Ohno, S.; Hatakeyama, S.; Nakayama,

[24884] K. I.: Increased proliferation of B cells and auto-immunity in mice

[24885] lacking protein kinase C-delta. Nature 416: 865-869, 2002.

[24886] 3128.

[24887] Gong, J.; Xu, J.; Bezanilla, M.; van Huizen, R.; Derin, R.; Li,

[24888] M.: Differential stimulation of PKC phosphorylation of potassium
[24889] channels by ZIP1 and ZIP2. Science 285: 1565-1569, 1999.
[24890] 3129.

[24891] Spritz, R. A.; Strunk, K.; Surowy, C. S.; Mohrenweiser, H. W.:
[24892] Human U1-70K ribonucleoprotein antigen gene: organization, nucleotide
[24893] sequence, and mapping to locus 19q13.3. Genomics 8: 371-379, 1990.
[24894] 3130.

[24895] Ford, M. D.; Lavin, M. F.: Ataxia telangiectasia: an anomaly
[24896] in DNA replication after irradiation. Nucleic Acids Res. 9: 1395-1404,
[24897] 1981.

[24898] 3131.

[24899] Frais, M. A.: Gastric adenocarcinoma due to ataxia-telangiectasia
[24900] (Louis-Bar syndrome). J. Med. Genet. 16: 160-161, 1979.

[24901] 3132.

[24902] Garnier, J.; Osguthorpe, D. J.; Robson, B.: Analysis of the accuracy
[24903] and implications of simple methods for predicting the secondary structure
[24904] of globular proteins. J. Molec. Biol. 120: 97-120, 1978.

[24905] 3133.

[24906] Epstein, A. C. R.; Gleadle, J. M.; McNeill, L. A.; Hewitson, K.
[24907] S.; O'Rourke, J.; Mole, D. R.; Mukherji, M.; Metzen, E.; Wil-son, M.
[24908] I.; Dhanda, A.; Tian, Y.-M.; Masson, N.; Hamilton, D. L.; Jaakkola,
[24909] P.; Barstead, R.; Hodgkin, J.; Maxwell, P. H.; Pugh, C. W.; Schofield,
[24910] C. J.; Ratcliffe, P. J.: C. elegans EGL-9 and mammalian ho-mologs
[24911] define a family of dioxygenases that regulate HIF by prolyl hydroxylation. Cell
[24912] 107: 43-54, 2001.

[24913] 3134.

[24914] Foroud, T.; Wei, S.; Ziv, Y.; Sobel, E.; Lange, E.; Chao, A.;
[24915] Goradia, T.; Huo, Y.; Tolun, A.; Chessa, L.; Charmley, P.;

Sanal,
[24916] O.; Salman, N.; Julier, C.; Concannon, P.; McConville, C.; Taylor,
[24917] A. M. R.; Shiloh, Y.; Lange, K.; Gatti, R. A.: Localization of an
[24918] ataxia-telangiectasia locus to a 3-cM interval on chromosome 11q23:
[24919] linkage analysis of 111 families by an international consortium. Am.
[24920] J. Hum. Genet. 49: 1263-1279, 1991.

[24921] 3135.

[24922] Gatei, M.; Young, D.; Cerosaletti, K. M.; Desai-Mehta, A.; Spring,
[24923] K.; Kozlov, S.; Lavin, M. F.; Gatti, R. A.; Concannon, P.; Khanna,
[24924] K.: ATM-dependent phosphorylation of nibrin in response to radiation
[24925] exposure. Nature Genet. 25: 115-119, 2000.

[24926] 3136.

[24927] Gatti, R. A.: Personal Communication. Los Angeles, Calif.

[24928] 6/1990.

[24929] 3137.

[24930] Gatti, R. A.: Personal Communication. Los Angeles, Calif.

[24931] 6/13/1989.

[24932] 3138.

[24933] Gatti, R. A.; Aurias, A.; Griscelli, C.; Sparkes, R. S.: Translocations

[24934] involving chromosomes 2p and 22q in ataxia-telangiectasia. Dis. Markers 3:

[24935] 169-195, 1985.

[24936] 3139.

[24937] Gatti, R. A.; Bick, M.; Tam, C. F.; Medici, M. A.; Oxelius, V.-A.;

[24938] Holland, M.; Goldstein, A. L.; Boder, E.: Ataxia-telangiectasia:

[24939] a multiparameter analysis of eight families. Clin. Immun. Immunopath. 23:

[24940] 501-516, 1982.

[24941] 3140.

[24942] Gatti, R. A.; Boder, E.; Vinters, H. V.; Sparkes, R. S.; Norman,

[24943] A.; Lange, K.: Ataxia-telangiectasia: an interdisciplinary approach
[24944] to pathogenesis. Medicine 70: 99-117, 1991.

[24945] 3141.

[24946] Gatti, R. A.; Peterson, K. L.; Novak, J.; Chen, X.; Yang-Chen,
[24947] L.; Liang, T.; Lange, E.; Lange, K.: Prenatal genotyping of ataxia-telangiectasia.
[24948] (Letter) Lancet 342: 376, 1993.

[24949] 3142.

[24950] Gatti, R. A.; Swift, M.: Ataxia-telangiectasia: Genetics, Neuropathology,
[24951] and Immunology of a Degenerative Disease of Childhood. New York:
[24952] Alan R. Liss (pub.) 1985.

[24953] 3143.

[24954] Gatti, R. A.; Berkel, I.; Boder, E.; Braedt, G.; Charmley, P.;

[24955] Concannon, P.; Ersoy, R.; Foroud, T.; Jaspers, N. G. J.; Lange, K.;
[24956] Lathrop, G. M.; Leppert, M.; Nakamura, Y.; O'Connell, P.; Paterson,

[24957] M.; Salser, W.; Sanal, O.; Silver, J.; Sparkes, R. S.; Susi, E.; Weeks,
[24958] D. E.; Wei, S.; White, R.; Yoder, F.: Localization of an ataxia-telangiectasia
[24959] gene to chromosome 11q22-23. Nature 336: 577-580, 1988.
[24960] 3144.

[24961] Gilad, S.; Bar-Shira, A.; Harnik, R.; Shkedy, D.; Ziv, Y.; Khosravi,
[24962] R.; Brown, K.; Vanagaite, L.; Xu, G.; Frydman, M.; Lavin, M. F.; Hill,
[24963] D.; Tagle, D. A.; Shiloh, Y.: Ataxia-telangiectasia: founder effect
[24964] among North African Jews. Hum. Molec. Genet. 5: 2033-2037, 1996.
[24965] 3145.

[24966] Gilad, S.; Chessa, L.; Khosravi, R.; Russell, P.; Galanty, Y.;

[24967] Piane, M.; Gatti, R. A.; Jorgensen, T. J.; Shiloh, Y.; Bar-Shira,
[24968] A.: Genotype-phenotype relationships in ataxia-telangiectasia and
[24969] variants. Am. J. Hum. Genet. 62: 551-561, 1998.

[24970] 3146.

[24971] Gilad, S.; Khosravi, R.; Shkedy, D.; Uziel, T.; Ziv, Y.; Savit-
sky,
[24972] K.; Rotman, G.; Smith, S.; Chessa, L.; Jorgensen, T. J.;
Harnik, R.;
[24973] Frydman, M.; Sanal, O.; Portnoi, S.; Goldwicz, Z.; Jaspers,
N. G.
[24974] J.; Gatti, R. A.; Lenoir, G.; Lavin, M. F.; Tatsumi, K.; Weg-
ner, R.
[24975] D.; Shiloh, Y.; Bar-Shira, A.: Predominance of null muta-
tions in
[24976] ataxia-telangiectasia. Hum. Molec. Genet. 5: 433-439,
1996.
[24977] 3147.

[24978] Ginter, D. N.; Tallapragada, R.: Ataxia-telangiectasia. Birth

[24979] Defects Orig. Art. Ser. XI(2): 408-409, 1975.

[24980] 3148.

[24981] Greenwell, P. W.; Kronmal, S. L.; Porter, S. E.; Gassbenhu-
ber,
[24982] J.; Obermaier, B.; Petes, T. D.: TEL1, a gene involved in
controlling

[24983] telomere length in S. cerevisiae, is homologous to the human ataxia

[24984] telangiectasia gene. Cell 82: 823-829, 1995.

[24985] 3149.

[24986] Haerer, A. F.; Jackson, J. F.; Evers, C. G.: Ataxia-telangiectasia

[24987] with gastric adenocarcinoma. J.A.M.A. 210: 1884-1887, 1969.

[24988] 3150.

[24989] Hagberg, A.; Hansson, O.; Liden, S.; Nilsson, K.: Familial ataxic

[24990] diplegia with deficient cellular immunity: a new clinical entity. Acta

[24991] Paediat. Scand. 59: 545-550, 1970.

[24992] 3151.

[24993] Hande, M. P.; Balajee, A. S.; Tchirkov, A.; Wynshaw-Boris, A.;

[24994] Lansdorp, P. M.: Extra-chromosomal telomeric DNA in cells from Atm-/-

[24995] mice and patients with ataxia-telangiectasia. Hum. Molec. Genet. 10:

[24996] 519-528, 2001.

[24997] 3152.

[24998] Crawford, R. J.; Hudson, P.; Shine, J.; Niall, H. D.; Eddy, R.

[24999] L.; Shows, T. B.: Two human relaxin genes are on chromosome 9. EMBO

[25000] J. 3: 2341-2345, 1984.

[25001] 3153.

[25002] Hsu, S. Y.; Nakabayashi, K.; Nishi, S.; Kumagai, J.; Kudo, M.;

[25003] Sherwood, O. D.; Hsueh, A. J. W.: Activation of orphan receptors

[25004] by the hormone relaxin. Science 295: 671-674, 2002.

[25005] 3154.

[25006] Hudson, P.; Haley, J.; Cronk, M.; Shine, J.; Niall, H.: Molecular

[25007] cloning and characterization of cDNA sequences coding for rat relaxin. Nature 291:

[25008] 127-131, 1981.

[25009] 3155.

[25010] Hudson, P.; Haley, J.; John, M.; Cronk, M.; Crawford, R.;

Haralambidis,
[25011] J.; Tregear, G.; Shine, J.; Niall, H.: Structure of a genomic clone
[25012] encoding biologically active human relaxin. Nature 301: 628-831,
[25013] 1983.

[25014] 3156.

[25015] Hudson, P.; John, M.; Crawford, R.; Haralambidis, J.; Scan-
lon,
[25016] D.; Gorman, J.; Tregear, G.; Shine, J.; Niall, H.: Relaxin gene expression
[25017] in human ovaries and the predicted structure of a human preprorelaxin
[25018] by analysis of cDNA clones. EMBO J. 3: 2333-2339, 1984.

[25019] 3157.

[25020] Naggert, J. K.; Mu, J.-L.: The mouse very low density lipoprotein
[25021] receptor (Vldlr) gene maps to chromosome 19. Mam-
malian Genome 5:
[25022] 453-455, 1994.

[25023] 3158.

[25024] Schwabe, C.; Bullesbach, E. E.: Relaxin: structures, functions,
[25025] promises, and nonevolution. FASEB J. 8: 1152-1160, 1994.
[25026] 3159.
[25027] Zhao, L.; Roche, P. J.; Gunnerson, J. M.; Hammond, V. E.; Tregear,
[25028] G. W.; Wintour, E. M.; Beck, F.: Mice without a functional relaxin
[25029] gene are unable to deliver milk to their pups. Endocrinology 140:
[25030] 445-453, 1999.
[25031] 3160.
[25032] Meloni, A. M.; Dobbs, R. M.; Pontes, J. E.; Sandberg, A. A.: Translocation
[25033] (X;1) in papillary renal cell carcinoma: a new cytogenetic subtype. Cancer
[25034] Genet. Cytogenet. 65: 1-6, 1993.
[25035] 3161.
[25036] Zbar, B.; Tory, K.; Merino, M.; Schmidt, L.; Glenn, G.; Choyke,

[25037] P.; Walther, M. M.; Lerman, M.; Linehan, W. M.: Hereditary papillary
[25038] renal cell carcinoma. J. Urol. 151: 561-566, 1994.

[25039] 3162.

[25040] Adachi, H.; Tawaragi, Y.; Inuzuka, C.; Kubota, I.; Tsuji-moto, M.;
[25041] Nishihara, T.; Nakazato, H.: Primary structure of human microsomal
[25042] dipeptidase deduced from molecular cloning. J. Biol. Chem. 265:
[25043] 3992-3995, 1990.

[25044] 3163.

[25045] Austruy, E.; Jeanpierre, C.; Antignac, C.; Whitmore, S. A.; Van
[25046] Cong, N.; Bernheim, A.; Callen, D. F.; Junien, C.: Physical and genetic
[25047] mapping of the dipeptidase gene DPEP1 to 16q24.3. Ge-nomics 15: 684-687,
[25048] 1993.

[25049] 3164.

[25050] Campbell, B. J.; Forrester, L. J.; Zahler, W. L.; Burks, M.:

Beta-lactamase

[25051] activity of purified and partially characterized human renal dipeptidase. J.

[25052] Biol. Chem. 259: 14586-14590, 1984.

[25053] 3165.

[25054] Kozak, E. M.; Tate, S. S.: Glutathione-degrading enzymes of microvillus

[25055] membranes. J. Biol. Chem. 257: 6322-6327, 1982.

[25056] 3166.

[25057] Nakagawa, H.; Inazawa, J.; Inoue, K.; Misawa, S.; Kashima, K.;

[25058] Adachi, H.; Nakazato, H.; Abe, T.: Assignment of the human renal

[25059] dipeptidase gene (DPEP1) to band q24 of chromosome 16. (Abstract) Cytogenet.

[25060] Cell Genet. 58: 2002 only, 1991.

[25061] 3167.

[25062] Bruce, L. J.; Cope, D. L.; Jones, G. K.; Schofield, A. E.; Burley,

[25063] M.; Povey, S.; Unwin, R. J.; Wrong, O.; Tanner, M. J. A.: Familial

[25064] distal renal tubular acidosis is associated with mutations in the
[25065] red cell anion exchanger (band 3, AE1) gene. J. Clin. In-
vest. 100:
[25066] 1693-1707, 1997.

[25067] 3168.

[25068] Kunz, J.; Hudler, M.; Fritz, B.; Gillessen-Kaesbach, G.; Pas-
sarge,
[25069] E.: Identification of a frameshift mutation in the gene
TWIST in
[25070] a family affected with Robinow-Sorauf syndrome. J. Med.
Genet. 36:
[25071] 650-652, 1999.

[25072] 3169.

[25073] Adachi, T.; Ohta, H.; Yamada, H.; Futenma, A.; Kato, K.;
Hirano,
[25074] K.: Quantitative analysis of extracellular-superoxide dis-
mutase in
[25075] serum and urine by ELISA with monoclonal antibody. Clin.
Chim. Acta 212:
[25076] 89-102, 1992.

[25077] 3170.

[25078] Folz, R. J.; Crapo, J. D.: Extracellular superoxide dismutase

[25079] (SOD3): tissue-specific expression, genomic characterization, and

[25080] computer-assisted sequence analysis of the human EC SOD gene. Genomics 22:

[25081] 162-171, 1994.

[25082] 3171.

[25083] Hendrickson, D. J.; Fisher, J. H.; Jones, C.; Ho, Y.-S.: Regional

[25084] localization of human extracellular superoxide dismutase gene to 4pter-q21.

[25085] Genomics 8: 736-738, 1990.

[25086] 3172.

[25087] Hjalmarsson, K.; Marklund, S. L.; Engstrom, A.; Edlund, T.: Isolation

[25088] and sequence of complementary DNA encoding human extracellular superoxide

[25089] dismutase. Proc. Nat. Acad. Sci. 84: 6340-6344, 1987.

[25090] 3173.

[25091] Marklund, S. L.: Extracellular superoxide dismutase in human tissues
[25092] and human cell lines. J. Clin. Invest. 74: 1398-1403, 1984.
[25093] 3174.

[25094] Sandstrom, J.; Nilsson, P.; Karlsson, K.; Marklund, S. L.: 10-fold
[25095] increase in human plasma extracellular superoxide dismutase content
[25096] caused by a mutation in heparin-binding domain. J. Biol. Chem. 269:
[25097] 19163-19166, 1994.

[25098] 3175.

[25099] Yamada, H.; Yamada, Y.; Adachi, T.; Goto, H.; Ogasawara, N.; Futenma,
[25100] A.; Kitano, M.; Hirano, K.; Kato, K.: Molecular analysis of extracellular-superoxide
[25101] dismutase gene associated with high level in serum. Jpn. J. Hum.
[25102] Genet. 40: 177-184, 1995.

[25103] 3176.

[25104] McNew, J. A.; Parlati, F.; Fukuda, R.; Johnston, R. J.; Paz, K.;
[25105] Paumet, F.; Sollner, T. H.; Rothman, J. E.: Compartmental specificity
[25106] of cellular membrane fusion encoded in SNARE proteins. Nature 407:
[25107] 153-159, 2000.

[25108] 3177.

[25109] Weber, T.; Zemelman, B. V.; McNew, J. A.; Westermann, B.; Gmachi,
[25110] M.; Parlati, F.; Sollner, T. H.; Rothman, J. E.: SNAREpins: minimal
[25111] machinery for membrane fusion. Cell 92: 759-772, 1998.

[25112] 3178.

[25113] Bodmer, W. F.: Monoclonal antibodies: their role in human genetics.
[25114] (Abstract) Sixth Int. Cong. Hum. Genet., Jerusalem 112 only, 1981.
[25115] 3179.

[25116] Bongiovanni, A. M.: The adrenogenital syndrome with de-
ficiency

[25117] of 3-beta-hydroxysteroid dehydrogenase. J. Clin. Invest. 41: 2086-2092,

[25118] 1962.

[25119] 3180.

[25120] Chang, Y. T.; Kappy, M. S.; Iwamoto, K.; Wang, J.; Yang, X.; Pang,

[25121] S.: Mutations in the type II 3-beta-hydroxysteroid dehydrogenase

[25122] gene in a patient with classic salt-wasting 3-beta-hydroxysteroid

[25123] dehydrogenase deficiency congenital adrenal hyperplasia. Pediat.

[25124] Res. 34: 698-700, 1993.

[25125] 3181.

[25126] Cravioto, M. D.; Ulloa-Aguirre, A.; Bermudez, J. A.; Herrera, J.;

[25127] Lisker, R.; Mendez, J. P.; Perez-Palacios, G.: A new inherited variant

[25128] of the 3-beta-hydroxysteroid dehydrogenase-isomerase deficiency syndrome:

[25129] evidence for the existence of two isoenzymes. J. Clin. Endocr. Metab. 63:

[25130] 360-367, 1986.

[25131] 3182.

[25132] Hamilton, W.; Brush, M. G.: Four clinical variants of con-
genital
[25133] adrenal hyperplasia. Arch. Dis. Child. 39: 66-72, 1964.

[25134] 3183.

[25135] Mendonca, B. B.; Bloise, W.; Arnhold, I. J. P.; Batista, M. C.;

[25136] Toledo, S. P. A.; Drummond, M. C. F.; Nicolau, W.; Mattar,
E.: Male
[25137] pseudohermaphroditism due to nonsalt-losing
3-beta-hydroxysteroid
[25138] dehydrogenase deficiency: gender role change and ab-
sence of gynecomastia
[25139] at puberty. J. Steroid Biochem. 28: 669-675, 1987.

[25140] 3184.

[25141] Morrison, N.; Nickson, D. A.; McBride, M.; Mueller, U. W.;
Boyd,
[25142] E.; Sutcliffe, R. G.: Regional chromosomal assignment of
the human
[25143] 3 beta-hydroxysteroid dehydrogenase/delta 5-4 iso-
merase (HSDB3) to

[25144] 1p13.1. (Abstract) Cytogenet. Cell Genet. 58: 1860 only, 1991.

[25145] 3185.

[25146] Morrison, N.; Nickson, D. A.; McBride, M. W.; Mueller, U. W.;

[25147] Boyd, E.; Sutcliffe, R. G.: Regional chromosomal assign-

ment of human

[25148] 3-beta-hydroxy-5-ene steroid dehydrogenase to 1p13.1 by non-isotopic

[25149] in situ hybridisation. Hum. Genet. 87: 223-225, 1991.

[25150] 3186.

[25151] Nishi, Y.; Tezuka, T.: Mild adrenal 3-beta-hydroxysteroid dehydrogenase

[25152] deficiency in children with accelerated growth, premature pubarche

[25153] and/or hirsutism. Europ. J. Pediat. 151: 19-23, 1992.

[25154] 3187.

[25155] Rheaume, E.; Leblanc, J. F.; Lachance, Y.; Labrie, F.; Simard,

[25156] J.: Detection of frequent BglII polymorphism by poly-

merase chain

[25157] reaction and TaqI restriction fragment length polymorphism for 3-beta-hydroxysteroid

[25158] dehydrogenase/delta-5-delta-4 isomerase at the human HSDB3 locus (1p11-p13). Hum.

[25159] Genet. 87: 753-754, 1991.

[25160] 3188.

[25161] Rheaume, E.; Simard, J.; Morel, Y.; Mebarki, F.; Zachmann, M.;

[25162] Forest, M. G.; New, M. I.; Labrie, F.: Congenital adrenal hyperplasia

[25163] due to point mutations in the type II 3-beta-hydroxysteroid dehydrogenase

[25164] gene. Nature Genet. 1: 239-245, 1992.

[25165] 3189.

[25166] Simard, J.; Durocher, F.; Mebarki, F.; Turgeon, C.; Sanchez, R.;

[25167] Labrie, Y.; Couet, J.; Trudel, C.; Rheaume, E.; Morel, Y.; Luu-The,

[25168] V.; Labrie, F.: Molecular biology and genetics of the 3-beta-hydroxysteroid

[25169] dehydrogenase/delta-5-delta-4 isomerase gene family. J. Endocrin. 150:

[25170] S189-S207, 1996.

[25171] 3190.

[25172] Stalvey, J. R. D.; Meisler, M. H.; Payne, A. H.: Evidence that

[25173] the same structural gene encodes testicular and adrenal 3-beta-hydroxysteroid

[25174] dehydrogenase-isomerase. Biochem. Genet. 25: 181-190, 1987.

[25175] 3191.

[25176] Zachmann, M.; Forest, M. G.; De Peretti, E.: 3 beta-hydroxysteroid

[25177] dehydrogenase deficiency: follow-up study in a girl with pubertal

[25178] bone age. Hormone Res. 11: 292-302, 1979.

[25179] 3192.

[25180] Aguiar-Oliveira, M. H.; Gill, M. S.; de A. Barretto, E. S.; Al-cantara,

[25181] M. R. S.; Miraki-Moud, F.; Menezes, C. A.; Souza, A. H. O.; Martinelli,

[25182] C. E.; Pereira, F. A.; Salvatori, R.; Levine, M. A.; Shalet, S. M.;

[25183] Camacho-Hubner, C.; Clayton, P. E.: Effect of severe

growth hormone

[25184] (GH) deficiency due to a mutation in the GH-releasing hormone receptor

[25185] on insulin-like growth factors (IGFs), IGF-binding proteins, and ternary

[25186] complex formation throughout life. J. Clin. Endocr. Metab. 84: 4118-4126,

[25187] 1999.

[25188] 3193.

[25189] DeAlmeida, V. I.; Mayo, K. E.: Identification and binding domains

[25190] of the growth hormone-releasing hormone receptor by analysis of mutant

[25191] and chimeric receptor proteins. Molec. Endocr. 12: 750-765, 1998.

[25192] 3194.

[25193] Gaylinn, B. D.; Harrison, J. K.; Zysk, J. R.; Lyons, C. E.; Lynch,

[25194] K. R.; Thorner, M. O.: Molecular cloning and expression of a human

[25195] anterior pituitary receptor for growth hormone-releasing hormone. Molec.

[25196] Endocr. 7: 77-84, 1993.

[25197] 3195.

[25198] Gaylinn, B. D.; von Kap-Herr, C.; Golden, W. L.; Thorner, M. O.

[25199] : Assignment of the human growth hormone-releasing hormone receptor

[25200] gene (GHRHR) to 7p14 by in situ hybridization. Genomics 19: 193-195,

[25201] 1994.

[25202] 3196.

[25203] Godfrey, P.; Rahal, J. O.; Beamer, W. G.; Copeland, N. G.; Jenkins,

[25204] N. A.; Mayo, K. E.: GHRH receptor of little mice contains a missense

[25205] mutation in the extracellular domain that disrupts receptor function. Nature

[25206] Genet. 4: 227-232, 1993.

[25207] 3197.

[25208] Gondo, R. G.; Aguiar-Oliveira, M. H.; Hayashida, C. Y.; Toledo,

[25209] S. P. A.; Abelin, N.; Levine, M. A.; Bowers, C. Y.; Souza, A.

H. O.;

[25210] Pereira, R. M. C.; Santos, N. L.; Salvatori, R.: Growth hormone-releasing

[25211] peptide-2 stimulates GH secretion in GH-deficient patients with mutated

[25212] GH-releasing hormone receptor. J. Clin. Endocr. Metab. 86: 3279-3283,

[25213] 2001.

[25214] 3198.

[25215] Lin, S.-C.; Lin, C. R.; Gukovsky, I.; Lusis, A. J.; Sawchenko,

[25216] P. E.; Rosenfeld, M. G.: Molecular basis of the little mouse phenotype

[25217] and implications for cell type-specific growth. Nature 364: 208-213,

[25218] 1993.

[25219] 3199.

[25220] Maheshwari, H. G.; Silverman, B. L.; Dupuis, J.; Baumann, G.:

[25221] Phenotype and genetic analysis of a syndrome caused by an inactivating

[25222] mutation in the growth hormone-releasing hormone re-

ceptor: dwarfism
[25223] of Sindh. J. Clin. Endocr. Metab. 83: 4065-4074, 1998.

[25224] 3200.

[25225] Netchine, I.; Talon, P.; Dastot, F.; Vitaux, F.; Goossens, M.;

[25226] Amselem, S.: Extensive phenotypic analysis of a family with growth
[25227] hormone (GH) deficiency caused by a mutation in the GH-releasing hormone
[25228] receptor gene. J. Clin. Endocr. Metab. 83: 432-436, 1998.

[25229] 3201.

[25230] Petersenn, S.; Rasch, A. C.; Heyens, M.; Schulte, H. M.: Structure
[25231] and regulation of the human growth hormone-releasing hormone receptor
[25232] gene. Molec. Endocr. 12: 233-247, 1998.

[25233] 3202.

[25234] Roelfsema, F.; Biermasz, N. R.; Veldman, R. G.; Veldhuis, J. D.;
[25235] Frolich, M.; Stovkis-Brantsma, W. H.; Wit, J.-M.: Growth hormone
[25236] (GH) secretion in patients with an inactivating defect of

the GH-releasing
[25237] hormone (GHRH) receptor is pulsatile: evidence for a role for non-GHRH
[25238] inputs into the generation of GH pulses. J. Clin. Endocr. Metab. 86:
[25239] 2459-2464, 2001.

[25240] 3203.

[25241] Salvatori, R.; Fan, X.; Mullis, P. E.; Haile, A.; Levine, M. A.

[25242] : Decreased expression of the GHRH receptor gene due to a mutation
[25243] in a Pit-1 binding site. Molec. Endocr. 16: 450-458, 2002.

[25244] 3204.

[25245] Salvatori, R.; Fan, X.; Phillips, J. A., III; Espigares-Martin,

[25246] R.; de Lara, I. M.; Freeman, K. L.; Plotnick, L.; Al-Ashwal, A.; Levine,
[25247] M. A.: Three new mutations in the gene for the growth hormone (GH)-releasing
[25248] hormone receptor in familial isolated GH deficiency type IB. J. Clin.
[25249] Endocr. Metab. 86: 273-279, 2001.

[25250] 3205.

[25251] Salvatori, R.; Hayashida, C. Y.; Aguiar-Oliveira, M. H.; Phillips,

[25252] J. A., III; Souza, A. H. O.; Gondo, R. G.; Toledo, S. P. A.; Conceicao,

[25253] M. M.; Prince, M.; Maheshwari, H. G.; Baumann, G.; Levine, M. A.

[25254] : Familial dwarfism due to a novel mutation of the growth hormone-releasing

[25255] hormone receptor gene. J. Clin. Endocr. Metab. 84: 917-923, 1999.

[25256] 3206.

[25257] Vamvakopoulos, N. C.; Kunz, J.; Olberding, U.; Scherer, S. W.;

[25258] Sioutopoulou, T. O.; Schneider, V.; Durkin, A. S.; Nierman, W. C.

[25259] : Mapping the human growth hormone-releasing hormone receptor (GHRHR)

[25260] gene to the short arm of chromosome 7 (7p13-p21) near the epidermal

[25261] growth factor receptor (EGFR) gene. Genomics 20: 338-340, 1994.

[25262] 3207.

[25263] Wajnrajch, M. P.; Gertner, J. M.; Harbison, M. D.; Chua, S. C.,

[25264] Jr.; Leibel, R. L.: Nonsense mutation in the human growth hormone-releasing

[25265] hormone receptor causes growth failure analogous to the little (lit)

[25266] mouse. Nature Genet. 12: 88-90, 1996.

[25267] 3208.

[25268] Baier, L.; The Pima Diabetes Genes Group: Suggestive linkage

[25269] of genetic markers on chromosome 4q12 to NIDDM and insulin action

[25270] in Pima Indians: new evidence to extend associations re-ported in other

[25271] populations. (Abstract) Diabetes 45 (suppl. 2): 30A only, 1996.

[25272] 3209.

[25273] Baier, L. J.; Dobberfuhl, A. M.; Pratley, R. E.; Hanson, R. L.;

[25274] Bogardus, C.: Variations in the vitamin D-binding protein (Gc locus)

[25275] are associated with oral glucose tolerance in nondiabetic Pima Indians. J.

[25276] Clin. Endocr. Metab. 83: 2993-2996, 1998.

[25277] 3210.

[25278] Bearn, A. G.; Bowman, B. H.; Kitchin, F. D.: Genetic and biochemical

[25279] consideration of the serum group-specific component. Cold Spring

[25280] Harbor Symp. Quant. Biol. 29: 435-442, 1964.

[25281] 3211.

[25282] Bowman, B. H.; Brune, J. L.; McCombs, J. L.; Moore, C. M.; Lum,

[25283] J. B.; Wieder, K.; Barnett, D. R.; Yang, F.: Human group-specific

[25284] component: a member of the albumin and alpha-fe-toprotein gene family.

[25285] (Abstract) Am. J. Hum. Genet. 37: A145 only, 1985.

[25286] 3212.

[25287] Braun, A.; Bichlmaier, R.; Cleve, H.: Molecular analysis of the

[25288] gene for the human vitamin-D-binding protein (group-specific component):

[25289] allelic differences of the common genetic GC types. Hum.

Genet. 89:

[25290] 401-406, 1992.

[25291] 3213.

[25292] Borsani, G.; Bassi, M. T.; Sperandeo, M. P.; De Grandi, A.; Buoninconti,

[25293] A.; Riboni, M.; Manzoni, M.; Incerti, B.; Pepe, A.; Andria, G.; Ballabio,

[25294] A.; Sebastio, G.: SLC7A7, encoding a putative permease-related protein,

[25295] is mutated in patients with lysinuric protein intolerance. Nature

[25296] Genet. 21: 297-301, 1999.

[25297] 3214.

[25298] Itoh, H.; Tomita, M.; Uchino, H.; Kobayashi, T.; Kataoka, H.; Sekiya,

[25299] R.; Nawa, Y.: cDNA cloning of rat pS2 peptide and expression of trefoil

[25300] peptides in acetic acid-induced colitis. Biochem. J. 318: 939-944,

[25301] 1996.

[25302] 3215.

[25303] Seib, T.; Blin, N.; Hilgert, K.; Seifert, M.; Theisinger, B.; Engel,
[25304] M.; Dooley, S.; Zang, K.-D.; Welter, C.: The three human trefoil
[25305] genes TFF1, TFF2, and TFF3 are located within a region of 55 kb on
[25306] chromosome 21q22.3. Genomics 40: 200-202, 1997.

[25307] 3216.

[25308] Aagenaes, O.: Hereditary spastic paraplegia. Acta Psychiat. Neurol.
[25309] Scand. 34: 489-494, 1959.

[25310] 3217.

[25311] Bayley, W. D.: Hereditary spastic paraplegia. J. Nerv. Ment.

[25312] Dis. 24: 697-701, 1897.

[25313] 3218.

[25314] Behan, W. M. H.; Maia, M.: Strumpell's familial spastic paraplegia:
[25315] genetics and neuropathology. J. Neurol. Neurosurg. Psychiat. 37:
[25316] 8-20, 1974.

[25317] 3219.

[25318] Boustany, R.-M. N.; Fleischnick, E.; Alper, C. A.; Marazita, M.

[25319] L.; Spence, M. A.; Martin, J. B.; Kolodny, E. H.: The autosomal dominant

[25320] form of 'pure' familial spastic paraplegia: clinical findings and

[25321] linkage analysis of a large pedigree. Neurology 37: 910-915, 1987.

[25322] 3220.

[25323] Burdick, A. B.; Owens, L. A.; Peterson, C. R.: Slowly progressive

[25324] autosomal dominant spastic paraplegia with late onset, variable expression

[25325] and reduced penetrance: a basis for diagnosis and counseling. Clin.

[25326] Genet. 19: 1-7, 1981.

[25327] 3221.

[25328] Cooley, W. C.; Melkonian, G.; Moses, C.; Moeschler, J. B.: Autosomal

[25329] dominant familial spastic paraplegia: description of a large New England

[25330] family and a study of management. Dev. Med. Child Neu-
rol. 32: 1087-1104,
[25331] 1990.

[25332] 3222.

[25333] Cooley, W. C.; Rawnsley, E.; Melkonian, G.; Moses, C.; Mc-
Cann,
[25334] D.; Virgin, B.; Coughlan, J.; Moeschler, J. B.: Autosomal
dominant
[25335] familial spastic paraplegia: report of a large New England
family. Clin.
[25336] Genet. 38: 57-68, 1990.

[25337] 3223.

[25338] Dube, M.-P.; Mlodzienski, M. A.; Kibar, Z.; Farlow, M. R.;
Ebers,
[25339] G.; Harper, P.; Kolodny, E. H.; Rouleau, G. A.; Figlewicz, D.
A.:
[25340] Hereditary spastic paraplegia: LOD-score considerations
for confirmation
[25341] of linkage in a heterogeneous trait. Am. J. Hum. Genet.
60: 625-629,
[25342] 1997.

[25343] 3224.

[25344] Durr, A.; Brice, A.; Serdaru, M.; Rancurel, G.; Derouesne, C.;

[25345] Lyon-Caen, O.; Agid, Y.; Fontaine, B.: The phenotype of 'pure' autosomal

[25346] dominant spastic paraplegia. Neurology 44: 1274-1277, 1994.

[25347] 3225.

[25348] Fink, J. K.: Advances in hereditary spastic paraplegia. Curr.

[25349] Opin. Neurol. 10: 313-318, 1997.

[25350] 3226.

[25351] Fink, J. K.; Heiman-Patterson, T.; Bird, T.; Cambi, F.; Dube,

[25352] M.-P.; Figlewicz, D. A.; Haines, J. L.; Hentati, A.; Pericak-Vance,

[25353] M. A.; Raskind, W.; Rouleau, G. A.; Siddique, T.: Hereditary spastic

[25354] paraplegia: advances in genetic research. Neurology 46: 1507-1514,

[25355] 1996.

[25356] 3227.

[25357] Garland, H. G.; Astley, C. E.: Hereditary spastic paraplegia

[25358] with amyotrophy and pes cavus. J. Neurol. Neurosurg. Psychiat. 13:

[25359] 130-133, 1950.

[25360] 3228.

[25361] Gispert, S.; Santos, N.; Damen, R.; Voit, T.; Schulz, J.; Klockgether,

[25362] T.; Orozco, G.; Kreuz, F.; Weissenbach, J.; Auburger, G.: Autosomal

[25363] dominant familial spastic paraplegia: reduction of the FSP1 candidate

[25364] region on chromosome 14q to 7 cM and locus hetero- geneity. Am. J.

[25365] Hum. Genet. 56: 183-187, 1995.

[25366] 3229.

[25367] Harding, A. E.: Hereditary 'pure' spastic paraplegia: a clini- cal

[25368] and genetic study of 22 families. J. Neurol. Neurosurg. Psychiat. 44:

[25369] 871-883, 1981.

[25370] 3230.

[25371] Hariga, J.; Matthys, E.: De la paraplegie spasmodique de Strumpell-Lorrain
[25372] a l'amyotrophie de Charcot-Marie-Tooth: (etude d'une famille). J.
[25373] Genet. Hum. 10: 326-337, 1961.

[25374] 3231.

[25375] Hazan, J.; Lamy, C.; Melki, J.; Munnich, A.; de Recondo, J.; Weissenbach,
[25376] J.: Autosomal dominant familial spastic paraplegia is ge-netically
[25377] heterogeneous and one locus maps to chromosome 14q. Nature Genet. 5:
[25378] 163-167, 1993.

[25379] 3232.

[25380] Hohmann, H.: Die Diplegia spastica infantilis hereditaria und
[25381] ihre Beziehungen zur familiaeren spastischen Spinalparal-yse. Nervenarzt 28:
[25382] 323-325, 1957.

[25383] 3233.

[25384] Huang, S.; Zhuyu; Li, H.; Labu; Baizhu; Lo, W. H. Y.; Fis-

cher,
[25385] C.; Vogel, F.: Another pedigree with pure autosomal dominant spastic
[25386] paraplegia (AD-FSP) from Tibet mapping to 14q11.2-q24.3. Hum. Genet. 100:
[25387] 620-623, 1997.

[25388] 3234.

[25389] McKusick, V. A.: Personal Communication. Baltimore, Md. 1965.
[25390] 3235.

[25391] McLeod, J. G.; Morgan, J. A.; Reye, C.: Electrophysiological
[25392] studies in familial spastic paraplegia. J. Neurol. Neurosurg. Psychiat. 40:
[25393] 611-615, 1977.

[25394] 3236.

[25395] Muglia, M.; Magariello, A.; Nicoletti, G.; Patitucci, A.; Gabriele,
[25396] A. L.; Conforti, F. L.; Mazzei, R.; Caracciolo, M.; Ardito, B.; Lastilla,
[25397] M.; Tedeschi, G.; Quattrone, A.: Further evidence that SPG3A gene

[25398] mutations cause autosomal dominant hereditary spastic paraplegia. Ann.
[25399] Neurol. 51: 794-795, 2002.

[25400] 3237.

[25401] Opjordsmoen, S.; Nyberg-Hansen, R.: Hereditary spastic paraplegia
[25402] with neurogenic bladder disturbances and syndactyly. Acta Neurol.
[25403] Scand. 61: 35-41, 1980.

[25404] 3238.

[25405] Polo, J. M.; Calleja, J.; Combarros, O.; Berciano, J.: Heredi-tary
[25406] 'pure' spastic paraplegia: a study of nine families. J. Neu-rol. Neurosurg.
[25407] Psychiat. 56: 175-181, 1993.

[25408] 3239.

[25409] Rainier, S.; Hedera, P.; Alvarado, D.; Zhao, X.; Kleopa, K. A.;
[25410] Heiman-Patterson, T.; Fink, J. K.: Hereditary spastic para-plegia
[25411] linked to chromosome 14q11-q21: reduction of the SPG3

locus interval

[25412] from 5.3 to 2.7 cM. J. Med. Genet. 39: , 2001. Note: Electronic

[25413] Article.

[25414] 3240.

[25415] Brose, N.; Petrenko, A. G.; Sudhof, T. C.; Jahn, R.: Synaptotagmin:

[25416] a calcium sensor on the synaptic vesicle surface. Science 256: 1021-1025,

[25417] 1992.

[25418] 3241.

[25419] Fernandez-Chacon, R.; Konigstorfer, A.; Gerber, S. H.; Garcia,

[25420] J.; Matos, M. F.; Stevens, C. F.; Brose, N.; Rizo, J.; Rosenmund,

[25421] C.; Sudhof, T. C.: Synaptotagmin I functions as a calcium regulator

[25422] of release probability. Nature 410: 41-49, 2001.

[25423] 3242.

[25424] Geppert, M.; Archer, B. T., III; Sudhof, T. C.: Synaptotagmin

[25425] II: a novel differentially distributed form of synaptotagmin. J.

[25426] Biol. Chem. 266: 13548-13552, 1991.

[25427] 3243.

[25428] Hilbush, B. S.; Morgan, J. I.: A third synaptotagmin gene, Syt3,

[25429] in the mouse. Proc. Nat. Acad. Sci. 91: 8195-8199, 1994.

[25430] 3244.

[25431] Jones, J. M.; Popma, S. J.; Mizuta, M.; Seino, S.; Meisler, M.

[25432] H.: Synaptotagmin genes on mouse chromosomes 1, 7, and 10 and human

[25433] chromosome 19. Mammalian Genome 6: 212-213, 1995.

[25434] 3245.

[25435] Kwon, O.-J.; Adamson, M. C.; Chin, H.; Kozak, C. A.: Genetic mapping

[25436] of five mouse genes encoding synaptotagmins. Mammalian Genome 6:

[25437] 880-881, 1995.

[25438] 3246.

[25439] Mackler, J. M.; Drummond, J. A.; Loewen, C. A.; Robinson,

I. M.;

[25440] Reist, N. E.: The C2B Ca(2+)-binding motif of synaptotagmin is required

[25441] for synaptic transmission in vivo. Nature 418: 340-344, 2002.

[25442] 3247.

[25443] Perin, M. S.; Fried, V. A.; Mignery, G. A.; Jahn, R.; Sudhof, T.

[25444] C.: Phospholipid binding by a synaptic vesicle protein homologous

[25445] to the regulatory region of protein kinase C. Nature 345: 260-263,

[25446] 1990.

[25447] 3248.

[25448] Perin, M. S.; Johnston, P. A.; Ozcelik, T.; Jahn, R.; Francke,

[25449] U.; Sudhof, T. C.: Structural and functional conservation of synaptotagmin

[25450] (p65) in Drosophila and humans. J. Biol. Chem. 266: 615-622, 1991.

[25451] 3249.

[25452] Robinson, I. M.; Ranjan, R.; Schwarz, T. L.: Synaptotagmins

I

[25453] and IV promote transmitter release independently of Ca(2+) binding

[25454] in the C2A domain. Nature 418: 336-340, 2002.

[25455] 3250.

[25456] Shin, O.-H.; Rizo, J.; Sudhof, T. C.: Synaptotagmin function

[25457] in dense core vesicle exocytosis studied in cracked PC12 cells. Nature

[25458] Neurosci. 5: 649-656, 2002.

[25459] 3251.

[25460] Wang, C.-T.; Grishanin, R.; Earles, C. A.; Chang, P. Y.; Martin,

[25461] T. F. J.; Chapman, E. R.; Jackson, M. B.: Synaptotagmin modulation

[25462] of fusion pore kinetics in regulated exocytosis of dense-core vesicles. Science 294:

[25463] 1111-1115, 2001.

[25464] 3252.

[25465] Magoulas, C.; Fried, M.: The Surf-6 gene of the mouse surfeit

[25466] locus encodes a novel nucleolar protein. DNA Cell Biol. 15: 305-316,
[25467] 1996.

[25468] 3253.

[25469] Boehm, T.; Baer, R.; Lavenir, I.; Forster, A.; Waters, J. J.; Nacheva,
[25470] E.; Rabbitts, T. H.: The mechanism of chromosomal translocation t(11;14)
[25471] involving the T-cell receptor C-delta locus on human chromosome 14q11
[25472] and a transcribed region of chromosome 11p15. EMBO J. 7: 385-394,
[25473] 1988.

[25474] 3254.

[25475] Begley, C. G.; Aplan, P. D.; Davey, M. P.; Nakahara, K.; Tchorz,
[25476] K.; Kurtzberg, J.; Hershfield, M. S.; Haynes, B. F.; Cohen, D. I.;
[25477] Waldmann, T. A.; Kirsch, I. R.: Chromosomal translocation in a human
[25478] leukemic stem-cell line disrupts the T-cell antigen receptor delta-chain

[25479] diversity region and results in a previously unreported fusion transcript. Proc.

[25480] Nat. Acad. Sci. 86: 2031-2035, 1989.

[25481] 3255.

[25482] Aruffo, A.; Seed, B.: Molecular cloning of two CD7 (T-cell leukemia

[25483] antigen) cDNAs by a COS cell expression system. EMBO J. 6: 3313-3316,

[25484] 1987.

[25485] 3256.

[25486] Baker, E.; Sandrin, M. S.; Garson, O. M.; Sutherland, G. R.; McKenzie,

[25487] I. F. C.; Webber, L. M.: Localization of the cell surface antigen

[25488] CD7 by chromosomal in situ hybridization. Immunogenetics 31: 412-413,

[25489] 1990.

[25490] 3257.

[25491] Bonilla, F. A.; Kokron, C. M.; Swinton, P.; Geha, R. S.: Targeted
[25492] gene disruption of murine CD7. Int. Immun. 9: 1875-1883, 1997.
[25493] 3258.

[25494] Jung, L. K. L.; Fu, S. M.; Hara, T.; Kapoor, N.; Good, R. A.:
[25495] Defective expression of T cell-associated glycoprotein in severe combined
[25496] immunodeficiency. J. Clin. Invest. 77: 940-946, 1986.

[25497] 3259.

[25498] Osada, S.; Utsumi, K. R.; Ueda, R.; Akao, Y.; Tsuge, I.; Nishida,
[25499] K.; Okada, J.; Matsuoka, H.; Takahashi, T.: Assignment of a gene
[25500] coding for a human T-cell antigen with a molecular weight of 40,000
[25501] daltons to chromosome 17. Cytogenet. Cell Genet. 47: 8-10, 1988.
[25502] 3260.

[25503] Schanberg, L. E.; Fleenor, D. E.; Kurtzberg, J.; Haynes, B. F.;

[25504] Kaufman, R. E.: Isolation and characterization of the genomic human

[25505] CD7 gene: structural similarity with the murine Thy-1 gene. Proc.

[25506] Nat. Acad. Sci. 88: 603-607, 1991.

[25507] 3261.

[25508] Yoshikawa, K.; Seto, M.; Ueda, R.; Obata, Y.; Notake, K.; Yokochi,

[25509] T.; Takahashi, T.: Molecular cloning of the gene coding for the human

[25510] T cell differentiation antigen CD7. Immunogenetics 33: 352-360,

[25511] 1991.

[25512] 3262.

[25513] Simon, S.; Awdeh, Z.; Campbell, R. D.; Ronco, P., II; Brink, S.

[25514] J.; Eisenbarth, G. S.; Yunis, E. J.; Alper, C. A.: A restriction

[25515] fragment of the C2 gene is a unique marker for C2 deficiency and the

[25516] uncommon C2 allele C2*B (a marker for type 1 diabetes). J. Clin.

[25517] Invest. 88: 2142-2145, 1991.

[25518] 3263.

[25519] Thong, Y. H.; Simpson, D. A.; Muller-Eberhard, H. J.: Ho-
mozygous
[25520] deficiency of the second component of complement pre-
senting with recurrent
[25521] bacterial meningitis. Arch. Dis. Child. 55: 471-473, 1980.

[25522] 3264.

[25523] Tokunaga, K.; Araki, C.; Juji, T.; Omoto, K.: Polymorphism
of
[25524] properdin factor B in Japanese: description of a rare vari-
ant and
[25525] data of association with HLA and C2. Hum. Genet. 60:
42-45, 1982.
[25526] 3265.

[25527] Tokunaga, K.; Araki, C.; Juji, T.; Omoto, K.: Genetic poly-
morphism
[25528] of the complement C2 in Japanese. Hum. Genet. 58:
213-216, 1981.
[25529] 3266.

[25530] Wahl, R.; Meo, T.; Shreffler, D.; Miller, W.; Atkinson, J. P.;

[25531] Schultz, J.; Osterland, C. K.: C2 deficiency and a lupus erythematosus-like
[25532] illness: family re-evaluation. (Letter) Ann. Intern. Med. 90: 717-718,
[25533] 1979.

[25534] 3267.

[25535] Wetsel, R. A.; Kulics, J.; Lokki, M.-L.; Kiepiela, P.; Akama,
[25536] H.; Johnson, C. A. C.; Densen, P.; Colten, H. R.: Type II human complement
[25537] C2 deficiency: allele-specific amino acid substitutions (ser189-to-phe;
[25538] gly444-to-arg) cause impaired C2 secretion. J. Biol. Chem. 271:
[25539] 5824-5831, 1996.

[25540] 3268.

[25541] WHO-IUIS Nomenclature Sub-Committee: Nomenclature for human
[25542] complement component C2*2. Europ. J. Immunogenet. 20: 299-300, 1993.
[25543] 3269.

[25544] Wolski, K. P.; Schmid, F. R.; Mittal, K. K.: Genetic linkage

[25545] between the HL-A system and a deficit of the second component (C2)

[25546] of complement in four generations of a family. Tissue Antigens 7:

[25547] 35-38, 1976.

[25548] 3270.

[25549] Kikuchi, M.; Tayama, T.; Hayakawa, H.; Takahashi, I.; Hoshino,

[25550] H.; Ohsaka, A.: Familial thrombocytosis. Brit. J. Haemat. 89: 900-902,

[25551] 1995.

[25552] 3271.

[25553] Wiestner, A.; Schlemper, R. J.; van der Maas, A. P. C.; Skoda,

[25554] R. C.: An activating splice donor mutation in the throm-bopoietin

[25555] gene causes hereditary thrombocythaemia. Nature Genet. 18: 49-52,

[25556] 1998.

[25557] 3272.

[25558] European Working Group on CF Genetics: Gradient of dis-

tribution
[25559] in Europe of the major CF mutation and of its associated haplotype. Hum.
[25560] Genet. 85: 436-445, 1990.

[25561] 3273.

[25562] Fanen, P.; Ghanem, N.; Vidaud, M.; Besmond, C.; Martin, J.; Costes,
[25563] B.; Plassa, F.; Goossens, M.: Molecular characterization of cystic
[25564] fibrosis: 16 novel mutations identified by analysis of the whole cystic
[25565] fibrosis conductance transmembrane regulator (CFTR) coding regions
[25566] and splice site junctions. Genomics 13: 770-776, 1992.

[25567] 3274.

[25568] Ruivenkamp, C. A. L.; van Wezel, T.; Zanon, C.; Stassen, A. P.
[25569] M.; Vlcek, C.; Csikos, T.; Klous, A. M.; Tripodis, N.; Per-rakis, A.;
[25570] Boerrigter, L.; Groot, P. C.; Lindeman, J.; Mooi, W. J.; Meij-jer,
[25571] G. A.; Scholten, G.; Dauwerse, H.; Paces, V.; van Zandwijk,

N.; van

[25572] Ommen, G. J. B.; Demant, P.: Ptprj is a candidate for the mouse colon-cancer

[25573] susceptibility locus Scc1 and is frequently deleted in human cancers. Nature

[25574] Genet. 31: 295-300, 2002.

[25575] 3275.

[25576] Watanabe, T.; Mukouyama, Y.; Rhodes, M.; Thomas, M.; Kume, T.;

[25577] Oishi, M.: Chromosomal location of murine protein tyrosine phosphatase

[25578] (Ptprj and Ptpre) genes. Genomics 29: 793-795, 1995.

[25579] 3276.

[25580] Elson, A.; Kozak, C. A.; Morton, C. C.; Weremowicz, S.; Leder,

[25581] P.: The protein tyrosine phosphatase epsilon gene maps to mouse chromosome

[25582] 7 and human chromosome 10q26. Genomics 31: 373-375, 1996.

[25583] 3277.

[25584] Krueger, N. X.; Streuli, M.; Saito, H.: Structural diversity

and
[25585] evolution of human receptor-like protein tyrosine phos-
phatases. EMBO
[25586] J. 9: 3241-3252, 1990.

[25587] 3278.

[25588] Melhado, I. G.; Anderson, L. L.; Duncan, A. M. V.; Jirik, F.
R.
[25589] : The gene encoding protein-tyrosine phosphatase RPTP-
epsilon (PTPRE)
[25590] is assigned to human chromosome 10q26. Cytogenet. Cell
Genet. 73:
[25591] 168-170, 1996.

[25592] 3279.

[25593] Banerjee, P.; Kleyn, P. W.; Knowles, J. A.; Lewis, C. A.;
Ross,
[25594] B. M.; Parano, E.; Kovats, S. G.; Lee, J. J.; Penchaszadeh, G.
K.;
[25595] Ott, J.; Jacobson, S. G.; Gilliam, T. C.: TULP1 mutation in
two extended
[25596] Dominican kindreds with autosomal recessive retinitis
pigmentosa. Nature
[25597] Genet. 18: 177-179, 1998.

[25598] 3280.

[25599] Hagstrom, S. A.; North, M. A.; Nishina, P. M.; Berson, E. L.; Dryja,

[25600] T. P.: Recessive mutations in the gene encoding the tubby-like protein

[25601] TULP1 in patients with retinitis pigmentosa. Nature Genet. 18: 174-176,

[25602] 1998.

[25603] 3281.

[25604] Knowles, J. A.; Shugart, Y.; Banerjee, P.; Gilliam, T. C.; Lewis,

[25605] C. A.; Jacobson, S. G.; Ott, J.: Identification of a locus, dis-tinct

[25606] from RDS-peripherin, for autosomal recessive retinitis pigmentosa

[25607] on chromosome 6p. Hum. Molec. Genet. 3: 1401-1403, 1994.

[25608] 3282.

[25609] Iivanainen, A.; Sainio, K.; Sariola, H.; Tryggvason, K.: Pri-mary

[25610] structure and expression of a novel human laminin alpha-4 chain. FEBS

[25611] Lett. 365: 183-188, 1995.

[25612] 3283.

[25613] Richards, A. J.; Al-Imara, L.; Carter, N. P.; Lloyd, J. C.; Leversha,

[25614] M. A.; Pope, F. M.: Localization of the gene (LAMA4) to chromosome

[25615] 6q21 and isolation of a partial cDNA encoding a variant laminin A

[25616] chain. Genomics 22: 237-239, 1994.

[25617] 3284.

[25618] Richards, A. J.; Al-Imara, L.; Carter, N. P.; Lloyd, J. C.; Pope,

[25619] F. M.: A laminin A variant gene (LAMA3) is present on chromosome

[25620] 6q21 (Abstract) J. Med. Genet. 31: 164, 1994.

[25621] 3285.

[25622] Dorow, D. S.; Devereux, L.; Dietzsch, E.; De Kretser, T.: Identification

[25623] of a new family of human epithelial protein kinases con-taining two

[25624] leucine/isoleucine-zipper domains. Europ. J. Biochem.

213: 701-710,

[25625] 1993.

[25626] 3286.

[25627] Hanks, S. K.: Eukaryotic protein kinases. Curr. Opin. Struct.

[25628] Biol. 1: 369-383, 1991.

[25629] 3287.

[25630] Cairns, P.; Polascik, T. J.; Eby, Y.; Tokino, K.; Califano, J.;

[25631] Merlo, A.; Mao, L.; Herath, J.; Jenkins, R.; Westra, W.; Rut-ter, J.

[25632] L.; Buckler, A.; Gabrielson, E.; Tockman, M.; Cho, K. R.; Hedrick,

[25633] L.; Bova, G. S.; Isaacs, W.; Koch, W.; Schwab, D.; Sidransky, D.:

[25634] Frequency of homozygous deletion at p16/CDKN2 in pri-mary human tumours. Nature

[25635] Genet. 11: 210-212, 1995.

[25636] 3288.

[25637] Ciotti, P.; Strigini, P.; Bianchi-Scarra, G.: Familial melanoma

[25638] and pancreatic cancer.(Letter) New Eng. J. Med. 334:

469-470, 1996.

[25639] 3289.

[25640] Ciotti, P.; Struewing, J. P.; Mantelli, M.; Chompret, A.; Avril,

[25641] M.-F.; Santi, P. L.; Tucker, M. A.; Bianchi-Scarra, G.; Bres-
sac-de

[25642] Paillerets, B.; Goldstein, A. M.: A single genetic origin for
the

[25643] G101W CDKN2A mutation in 20 melanoma-prone fami-
lies. Am. J. Hum.

[25644] Genet. 67: 311-319, 2000.

[25645] 3290.

[25646] Cowan, J. M.; Halaban, R.; Francke, U.: Cytogenetic analy-
sis

[25647] of melanocytes from premalignant nevi and melanomas. J.
Nat. Cancer

[25648] Inst. 80: 1159-1164, 1988.

[25649] 3291.

[25650] Della Torre, G.; Pasini, B.; Frigerio, S.; Donghi, R.; Rovini,

[25651] D.; Delia, D.; Peters, G.; Huot, T. J. G.; Bianchi-Scarra, G.;
Lantieri,

[25652] F.; Rodolfo, M.; Parmiani, G.; Pierotti, M. A.: CDKN2A and

CDK4 mutation
[25653] analysis in Italian melanoma-prone families: functional characterization
[25654] of a novel CDKN2A germ line mutation. Brit. J. Cancer 85: 836-844,
[25655] 2001.

[25656] 3292.

[25657] Dracopoli, N. C.; Fountain, J. W.: CDKN2 mutations in melanoma. Cancer
[25658] Surv. 26: 115-132, 1996.

[25659] 3293.

[25660] Fargnoli, M. C.; Chimenti, S.; Keller, G.; Soyer, H. P.; Dal Pozzo,
[25661] V.; Hofler, H.; Peris, K.: CDKN2a/p16(INK4a) mutations and lack of
[25662] p19(ARF) involvement in familial melanoma kindreds. J. Invest. Derm. 111:
[25663] 1202-1206, 1998.

[25664] 3294.

[25665] FitzGerald, M. G.; Harkin, D. P.; Silva-Arrieta, S.; MacDon-ald,

[25666] D. J.; Lucchina, L. C.; Unsal, H.; O'Neil, E.; Koh, J.; Finkel-
stein,
[25667] D. M.; Isselbacher, K. J.; Sober, A. J.; Haber, D. A.: Preva-
lence
[25668] of germ-line mutations in p16, p19ARF, and CDK4 in fa-
milial melanoma:
[25669] analysis of a clinic-based population. Proc. Nat. Acad. Sci.
93:
[25670] 8541-8545, 1996.

[25671] 3295.

[25672] Flores, J. F.; Pollock, P. M.; Walker, G. J.; Glendening, J. M.;

[25673] Lin, A. H.; Palmer, J. M.; Walters, M. K.; Hayward, N. K.;
Fountain,
[25674] J. W.: Analysis of the CDKN2A, CDKN2B and CDK4 genes
in 48 Australian
[25675] melanoma kindreds. Oncogene 15: 2999-3005, 1997.

[25676] 3296.

[25677] Goldstein, A. M.; Fraser, M. C.; Struewing J. P.; Hussussian,

[25678] C. J.; Ranade, K.; Zametkin, D. P.; Fontaine, L. S.; Organic,
S. M.;
[25679] Dracopoli, N. C.; Clark, W. H., Jr.; Tucker, M. A.: Increased

risk
[25680] of pancreatic cancer in melanoma-prone kindreds with p16 (INK4) mutations. New
[25681] Eng. J. Med. 333: 970-974, 1995.

[25682] 3297.

[25683] Goldstein, A. M.; Liu, L.; Shennan, M. G.; Hogg, D.; Tucker, M.
[25684] A.; Struewing, J. P.: A common founder for the V126D CDKN2A mutation
[25685] in seven North American melanoma-prone families. Brit. J. Cancer 85:
[25686] 527-530, 2001.

[25687] 3298.

[25688] Gruis, N. A.; van der Velden, P. A.; Sandkuijl, L. A.; Prins,
[25689] D. E.; Weaver-Feldhaus, J.; Kamb, A.; Bergman, W.; Frants, R. R.:
[25690] Homozygotes for CDKN2 (p16) germline mutation in Dutch familial melanoma
[25691] kindreds. Nature Genet. 10: 351-353, 1995.

[25692] 3299.

[25693] Harland, M.; Meloni, R.; Gruis, N.; Pinney, E.; Brookes, S.;

Spurr,
[25694] N. K.; Frischauf, A.-M.; Bataille, V.; Peters, G.; Cuzick, J.; Selby,
[25695] P.; Bishop, D. T.; Bishop, J. N.: Germline mutations of the CDKN2
[25696] gene in UK melanoma families. Hum. Molec. Genet. 6: 2061-2067, 1997.
[25697] 3300.

[25698] Harland, M.; Mistry, S.; Bishop, D. T.; Bishop, J. A. N.: A deep
[25699] intronic mutation in CDKN2A is associated with disease in a subset
[25700] of melanoma pedigrees. Hum. Molec. Genet. 10: 2679-2686, 2001.
[25701] 3301.

[25702] Hashemi, J.; Bendahl, P.-O.; Sandberg, T.; Platz, A.; Linder,

[25703] S.; Stierner, U.; Olsson, H.; Ingvar, C.; Hansson, J.; Borg, A.:
[25704] Haplotype analysis and age estimation of the 113insR CDKN2A founder
[25705] mutation in Swedish melanoma families. Genes Chromo-somes Cancer 31:

[25706] 107-116, 2001.

[25707] 3302.

[25708] Hayward, N. K.: The current situation with regard to hu-
man melanoma
[25709] and genetic inferences. Curr. Opin. Oncol. 8: 136-142,
1996.
[25710] 3303.

[25711] Hebert, J.; Cayuela, J. M.; Berkeley, J.; Sigaux, F.: Candidate

[25712] tumor-suppressor genes MTS1 (p16INK4A) and MTS2
(p15INK4B) display
[25713] frequent homozygous deletions in primary cells from T-
but not from
[25714] B-cell lineage acute lymphoblastic leukemias. Blood 84:
4038-4044,
[25715] 1994.

[25716] 3304.

[25717] Holland, E. A.; Beaton, S. C.; Becker, T. M.; Grulet, O. M.;
Peters,
[25718] B. A.; Rizos, H.; Kefford, R. F.; Mann, G. J.: Analysis of the
p16
[25719] gene, CDKN2, in 17 Australian melanoma kindreds. Onco-

gene 11: 2289-2294,

[25720] 1995.

[25721] 3305.

[25722] Hussussian, C. J.; Struewing, J. P.; Goldstein, A. M.; Hig-

gins,

[25723] P. A. T.; Ally, D. S.; Sheahan, M. D.; Clark, W. H., Jr.;

Tucker,

[25724] M. A.; Dracopoli, N. C.: Germline p16 mutations in familial

melanoma. Nature

[25725] Genet. 8: 15-21, 1994.

[25726] 3306.

[25727] Igaki, H.; Sasaki, H.; Kishi, T.; Sakamoto, H.; Tachimori, Y.;

[25728] Kato, H.; Watanabe, H.; Sugimura, T.; Terada, M.: Highly

frequent

[25729] homozygous deletion of the p16 gene in esophageal can-

cer cell lines. Biochem.

[25730] Biophys. Res. Commun. 203: 1090-1095, 1994.

[25731] 3307.

[25732] Balczon, R.; Bao, L.; Zimmer, W. E.: PCM-1, a 228-kD cen-

trosome

[25733] autoantigen with a distinct cell cycle distribution. J. Cell

Biol. 124:

[25734] 783-793, 1994.

[25735] 3308.

[25736] Tapia, C.; Featherstone, T.; Gomez, C.; Taillon-Miller, P.; Allende,

[25737] C. C.; Allende, J. E.: Cloning and chromosomal localization of the

[25738] gene coding for human protein kinase CK1. FEBS Lett. 349: 307-312,

[25739] 1994.

[25740] 3309.

[25741] Cardon, L. R.; Smith, S. D.; Fulker, D. W.; Kimberling, W. J.;

[25742] Pennington, B. F.; DeFries, J. C.: Quantitative trait locus for reading

[25743] disability on chromosome 6. Science 266: 276-279, 1994. Note: Erratum:

[25744] Science 268: 1553 only, 1995.

[25745] 3310.

[25746] Field, L. L.; Kaplan, B. J.: Absence of linkage of phonologi-cal

[25747] coding dyslexia to chromosome 6p23-p21.3 in a large

family data set. Am.

[25748] J. Hum. Genet. 63: 1448-1456, 1998.

[25749] 3311.

[25750] Fisher, S. E.; Francks, C.; Marlow, A. J.; MacPhie, I. L.; Newbury,

[25751] D. F.; Cardon, L. R.; Ishikawa-Brush, Y.; Richardson, A. J.; Talcott,

[25752] J. B.; Gayan, J.; Olson, R. K.; Pennington, B. F.; Smith, S. D.; DeFries,

[25753] J. C.; Stein, J. F.; Monaco, A. P.: Independent genome-wide scans

[25754] identify a chromosome 18 quantitative-trait locus influencing dyslexia. Nature

[25755] Genet. 30: 86-91, 2002.

[25756] 3312.

[25757] Fisher, S. E.; Marlow, A. J.; Lamb, J.; Maestrini, E.; Williams,

[25758] D. F.; Richardson, A. J.; Weeks, D. E.; Stein, J. F.; Monaco, A. P.

[25759] : A quantitative-trait locus on chromosome 6p influences different

[25760] aspects of developmental dyslexia. Am. J. Hum. Genet. 64:

146-156,

[25761] 1999.

[25762] 3313.

[25763] Gayan, J.; Smith, S. D.; Cherny, S. S.; Cardon, L. R.; Fulker,

[25764] D. W.; Brower, A. M.; Olson, R. K.; Pennington, B. F.; De-

Fries, J.

[25765] C.: Quantitative-trait locus for specific language and

reading deficits

[25766] on chromosome 6p. Am. J. Hum. Genet. 64: 157-164,

1999.

[25767] 3314.

[25768] Kaplan, D. E.; Gayan, J.; Ahn, J.; Won, T.-W.; Pauls, D.; Ol-

son,

[25769] R. K.; DeFries, J. C.; Wood, F.; Pennington, B. F.; Page, G.

P.; Smith,

[25770] S. D.; Gruen, J. R.: Evidence for linkage and association

with reading

[25771] disability, on 6p21.3-22. Am. J. Hum. Genet. 70:

1287-1298, 2002.

[25772] 3315.

[25773] Smith, S. D.; Kimberling, W. J.; Pennington, B. F.: Screening

[25774] for multiple genes influencing dyslexia. Reading Writing 3: 285-298,
[25775] 1991.

[25776] 3316.

[25777] Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.; Miki,
[25778] T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the epidermal
[25779] growth factor receptor kinase, enhances EGF-dependent mitogenic signals. EMBO
[25780] J. 12: 3799-3808, 1993.

[25781] 3317.

[25782] Corvi, R.; Berger, N.; Balczon, R.; Romeo, G.: RET/PCM-1: a novel
[25783] fusion gene in papillary thyroid carcinoma. Oncogene 19: 4236-4242,
[25784] 2000.

[25785] 3318.

[25786] Ohata, H.; Fujiwara, Y.; Koyama, K.; Nakamura, Y.: Map-ping of
[25787] the human autoantigen pericentriolar material 1 (PCM1)

gene to chromosome
[25788] 8p21.3-p22. Genomics 24: 404-406, 1994.

[25789] 3319.

[25790] Aoki, M.; Lin, C. L.; Rothstein, J. D.; Geller, B. A.; Hosler,

[25791] B. A.; Munsat, T. L.; Horvitz, H. R.; Brown, R. H.: Mutations in
[25792] the glutamate transporter EAAT2 gene do not cause abnormal EAAT2 transcripts
[25793] in amyotrophic lateral sclerosis. Ann. Neurol. 43: 645-653, 1998.
[25794] 3320.

[25795] Flowers, J. M.; Powell, J. F.; Leigh, P. N.; Andersen, P.; Shaw,
[25796] C. E.: Intron 7 retention and exon 9 skipping EAAT2 mRNA variants
[25797] are not associated with amyotrophic lateral sclerosis. Ann. Neurol. 49:
[25798] 643-649, 2001.

[25799] 3321.

[25800] Herault, J.; Martineau, J.; Petit, E.; Perrot, A.; Sauvage, D.;

[25801] Barthelemy, C.; Mallet, J.; Muh, J. P.; Lelord, G.: Genetic

markers

[25802] in autism: association study on short arm of chromosome 11. J. Autism

[25803] Dev. Dis. 24: 233-236, 1994.

[25804] 3322.

[25805] Honig, L. S.; Chambliss, D. D.; Bigio, E. H.; Carroll, S. L.; Elliott,

[25806] J. L.: Glutamate transporter EAAT2 splice variants occur not only

[25807] in ALS, but also in AD and controls. Neurology 55: 1082-1088, 2000.

[25808] 3323.

[25809] Kanai, Y.; Smith, C. P.; Hediger, M. A.: The elusive trans-

porters

[25810] with a high affinity for glutamate. Trends Neurosci. 16: 365-730,

[25811] 1993.

[25812] 3324.

[25813] Kirschner, M. A.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[25814] A.; Amara, S. G.: Mouse excitatory amino acid transporter EAAT2:

[25815] isolation, characterization, and proximity to neuroex-
citability loci
[25816] on mouse chromosome 2. Genomics 24: 218-224, 1994.

[25817] 3325.

[25818] Krishnan, S. N.; Desai, T.; Wyman, R. J.; Haddad, G. G.:
Cloning
[25819] of a glutamate transporter from human brain. Soc. Neu-
rosci. Abstr. 19:
[25820] 219 only, 1993.

[25821] 3326.

[25822] Li, X.; Francke, U.: Assignment of the gene SLC1A2 coding
for
[25823] the human glutamate transporter EAAT2 to human chro-
mosome 11 bands
[25824] p13-p12. Cytogenet. Cell Genet. 71: 212-213, 1995.

[25825] 3327.

[25826] Lin, C.-L. G.; Bristol, L. A.; Jin, L.; Dykes-Hoberg, M.;
Crawford,
[25827] T.; Clawson, L.; Rothstein, J. D.: Aberrant RNA processing
in a neurodegenerative
[25828] disease: the cause for absent EAAT2, a glutamate trans-

porter, in amyotrophic
[25829] lateral sclerosis. Neuron 20: 589-602, 1998.

[25830] 3328.

[25831] Meyer, T.; Munch, C.; Volkel, H.; Booms, P.; Ludolph, A. C.:

[25832] The EAAT2 (GLT-1) gene in motor neuron disease: ab-
sence of mutations
[25833] in amyotrophic lateral sclerosis and a point mutation in
patients
[25834] with hereditary spastic paraplegia. J. Neurol. Neurosurg.
Psychiat. 65:
[25835] 594-596, 1998.

[25836] 3329.

[25837] Takai, S.; Kawakami, H.; Nakayama, T.; Yamada, K.; Naka-
mura, S.
[25838] : Localization of the gene encoding the human L-
glutamate transporter
[25839] (GLT-1) to 11p11.2-p13 by fluorescence in situ hy-
bridization. Hum.
[25840] Genet. 97: 387-389, 1996.

[25841] 3330.

[25842] Tanaka, K.; Watase, K.; Manabe, T.; Yamada, K.; Watanabe,

M.;

[25843] Takahashi, K.; Iwama, H.; Nishikawa, T.; Ichihara, N.; Kikuchi, T.;

[25844] Okuyama, S.; Kawashima, N.; Hori, S.; Takimoto, M.; Wada, K.: Epilepsy

[25845] and exacerbation of brain injury in mice lacking the glutamate transporter

[25846] GLT-1. Science 276: 1699-1702, 1997.

[25847] 3331.

[25848] Trotti, D.; Aoki, M.; Pasinelli, P.; Berger, U. V.; Danbolt, N.

[25849] C.; Brown, R. H., Jr.; Hediger, M. A.: Amyotrophic lateral sclerosis-linked

[25850] glutamate transporter mutant has impaired glutamate clearance capacity. J.

[25851] Biol. Chem. 276: 576-582, 2001.

[25852] 3332.

[25853] Andresen, B. S.; Christensen, E.; Corydon, T. J.; Bross, P.; Pilgaard,

[25854] B.; Wanders, R. J. A.; Ruiter, J. P. N.; Simonsen, H.; Winter, V.;

[25855] Knudsen, I.; Schroeder, L. D.; Gregersen, N.; Skovby, F.:

Isolated
[25856] 2-methylbutyrylglycinuria caused by short/
branched-chain acyl-CoA
[25857] dehydrogenase deficiency: identification of a new enzyme
defect, resolution
[25858] of its molecular basis, and evidence for distinct acyl-CoA
dehydrogenases
[25859] in isoleucine and valine metabolism. Am. J. Hum. Genet.
67: 1095-1103,
[25860] 2000.

[25861] 3333.

[25862] Arden, K. C.; Viars, C. S.; Fu, K.; Rozen, R.: Localization of

[25863] short/branched chain acyl-CoA dehydrogenase (ACADSB)
to human chromosome
[25864] 10. Genomics 25: 743-745, 1995.

[25865] 3334.

[25866] Gibson, K. M.; Burlingame, T. G.; Hogema, B.; Jakobs, C.;
Schutgens,
[25867] R. B. H.; Millington, D.; Roe, C. R.; Roe, D. S.; Sweetman,
L.; Steiner,
[25868] R. D.; Linck, L.; Pohowalla, P.; Sacks, M.; Kiss, D.; Rinaldo,

P.;

[25869] Vockley, J.: 2-Methylbutyryl-coenzyme A dehydrogenase deficiency:

[25870] a new inborn error of L-isoleucine metabolism. Pediat. Res. 47:

[25871] 830-833, 2000.

[25872] 3335.

[25873] Rozen, R.; Vockley, J.; Zhou, L.; Milos, R.; Willard, J.; Fu, K.;

[25874] Vicanek, C.; Low-Nang, L.; Torban, E.; Fournier, B.: Isola-tion and

[25875] expression of a cDNA encoding the precursor for a novel member (ACADSB)

[25876] of the acyl-CoA dehydrogenase gene family. Genomics 24: 280-287,

[25877] 1994.

[25878] 3336.

[25879] Takai, S.; Tanaka, M.; Sugimura, H.; Yamada, K.; Naito, Y.; Kino,

[25880] I.; Matsuda, M.: Mapping of the human C3G gene coding a guanine nucleotide

[25881] releasing protein for Ras family to 9q34.3 by fluorescence

in situ

[25882] hybridization. Hum. Genet. 94: 549-550, 1994.

[25883] 3337.

[25884] Tanaka, S.; Morishita, T.; Hashimoto, Y.; Hattori, S.; Naka-

mura,

[25885] S.; Shibuya, M.; Matsuoka, K.; Takenawa, T.; Kurata, T.;

Nagashima,

[25886] K.; Matsuda, M.: C3G, a guanine nucleotide-releasing pro-

tein expressed

[25887] ubiquitously, binds to the Src homology 3 domains of CRK

and GRB2/ASH

[25888] proteins. Proc. Nat. Acad. Sci. 91: 3443-3447, 1994.

[25889] 3338.

[25890] McCusker, D.; Jones, T.; Sheer, D.; Trowsdale, J.: Genetic

relationships

[25891] of the genes encoding the human proteasome beta sub-

units and the proteasome

[25892] PA28 complex. Genomics 45: 362-367, 1997.

[25893] 3339.

[25894] Kumar, S.; Tomooka, Y.; Noda, M.: Identification of a set

of genes

[25895] with developmentally down-regulated expression in the mouse brain. Biochem.

[25896] Biophys. Res. Commun. 185: 1155-1161, 1992.

[25897] 3340.

[25898] Carmi, R.; Rokhlina, T.; Kwitek-Black, A. E.; Elbedour, K.; Nishimura,

[25899] D.; Stone, E. M.; Sheffield, V. C.: Use of a DNA pooling strategy

[25900] to identify a human obesity syndrome locus on chromosome 15. Hum.

[25901] Molec. Genet. 4: 9-13, 1995.

[25902] 3341.

[25903] Dallapiccola, B.: Familial translocation t(2p-; 17p+). Ann. Genet. 14:

[25904] 153-155, 1971.

[25905] 3342.

[25906] Katsanis, N.; Eichers, E. R.; Ansley, S. J.; Lewis, R. A.; Kayserili,

[25907] H.; Hoskins, B. E.; Scambler, P. J.; Beales, P. L.; Lupski, J. R.

[25908] : BBS4 is a minor contributor to Bardet-Biedl syndrome and may also

[25909] participate in triallelic inheritance. Am. J. Hum. Genet. 71: 22-29,

[25910] 2002.

[25911] 3343.

[25912] Losson, R.; Lacroute, F.: Interference of nonsense mutations with

[25913] eukaryotic messenger RNA stability. Proc. Nat. Acad. Sci. 76: 5134-5137,

[25914] 1979.

[25915] 3344.

[25916] Mykytyn, K. Braun, T.; Carmi, R.; Haider, N. B.; Searsby, C. C.;

[25917] Shastri, M.; Beck, G.; Wright, A. F.; Iannaccone, A.; Elbedour, K.;

[25918] Riise, R.; Baldi, A.; Raas-Rothschild, A.; Gorman, S. W.; Duhl, D.

[25919] M.; Jacobson, S. G.; Casavant, T.; Stone, E. M.; Sheffield, V. C.

[25920] : Identification of the gene that, when mutated, causes the human

[25921] obesity syndrome BBS4. Nature Genet. 28: 188-191, 2001.

[25922] 3345.

[25923] Braybrooke, J. P.; Spink, K. G.; Thacker, J.; Hickson, I. D.:

[25924] The RAD51 family member, RAD51L3, is a DNA-stimulated ATPase that

[25925] forms a complex with XRCC2. J. Biol. Chem. 275: 29100-29106, 2000.

[25926] 3346.

[25927] Johnson, R. D.; Liu, N.; Jasin, M.: Mammalian XRCC2 pro-motes the

[25928] repair of DNA double-strand breaks by homologous re-combination. Nature 401:

[25929] 397-399, 1999.

[25930] 3347.

[25931] Jones, N. J.; Cox, R.; Thacker, J.: Isolation and cross-sensitivity

[25932] of x-ray-sensitive mutants of V79-4 hamster cells. Mutat. Res. 183:

[25933] 279-286, 1987.

[25934] 3348.

[25935] Jones, N. J.; Zhao, Y.; Siciliano, M. J.; Thompson, L. H.: As-signment

[25936] of the XRCC2 human DNA repair gene to chromosome 7q36 by complementation

[25937] analysis. Genomics 26: 619-622, 1995.

[25938] 3349.

[25939] Kurumizaka, H.; Ikawa, S.; Nakada, M.; Enomoto, R.; Ka-

gawa, W.;

[25940] Kinebuchi, T.; Yamazoe, M.; Yokoyama, S.; Shibata, T.: Homologous

[25941] pairing and ring and filament structure formation activi-

ties of the

[25942] human Xrcc2-Rad51D complex. J. Biol. Chem. 277: 14315-14320, 2002.

[25943] 3350.

[25944] Liu, N.; Schild, D.; Thelen, M. P.; Thompson, L. H.: Involve-

ment

[25945] of Rad51C in two distinct protein complexes of Rad51 paralogs in human

[25946] cells. Nucleic Acids Res. 30: 1009-1015, 2002.

[25947] 3351.

[25948] Masson, J.-Y.; Tarsounas, M. C.; Stasiak, A. Z.; Stasiak, A.; Shah,

[25949] R.; McIlwraith, M. J.; Benson, F. E.; West, S. C.: Identification

[25950] and purification of two distinct complexes containing the five RAD51

[25951] paralogs. Genes Dev. 15: 3296-3307, 2001.

[25952] 3352.

[25953] Tambini, C. E.; George, A. M.; Rommens, J. M.; Tsui, L.-C.; Scherer,

[25954] S. W.; Thacker, J.: The XRCC2 DNA repair gene: identification of

[25955] a positional candidate. Genomics 41: 84-92, 1997.

[25956] 3353.

[25957] Thacker, J.; Tambini, C. E.; Simpson, P. J.; Tsui, L.-C.; Scherer,

[25958] S. W.: Localization to chromosome 7q36.1 of the human XRCC2 gene,

[25959] determining sensitivity to DNA-damaging agents. Hum. Molec. Genet. 4:

[25960] 113-120, 1995.

[25961] 3354.

[25962] Pollock, P. M.; Spurr, N.; Bishop, T.; Newton-Bishop, J.;

Gruis,
[25963] N.; van der Velden, P. A.; Goldstein, A. M.; Tucker, M. A.; Foulkes,
[25964] W. D.; Barnhill, R.; Haber, D.; Fountain, J.; Hayward, N. K.: Haplotype
[25965] analysis of two recurrent CDKN2A mutations in 10 melanoma families:
[25966] evidence for common founders and independent muta-tions. Hum. Mutat. 11:
[25967] 424-431, 1998.

[25968] 3355.

[25969] Orimo, A.; Inoue, S.; Minowa, O.; Tominaga, N.; Tomioka, Y.; Sato,
[25970] M.; Kuno, J.; Hiroi, H.; Shimizu, Y.; Suzuki, M.; Noda, T.; Muramatsu,
[25971] M.: Underdeveloped uterus and reduced estrogen respon-siveness in
[25972] mice with disruption of the estrogen-responsive finger protein gene,
[25973] which is a direct target of estrogen receptor alpha. Proc. Nat. Acad.
[25974] Sci. 96: 12027-12032, 1999.

[25975] 3356.

[25976] Urano, T.; Saito, T.; Tsukui, T.; Fujita, M.; Hosoi, T.; Mura-
matsu,

[25977] M.; Ouchi, Y.; Inoue, S.: Efp targets 14-3-3-sigma for
proteolysis

[25978] and promotes breast tumour growth. Nature 417:
871-875, 2002.

[25979] 3357.

[25980] Grun, F.; Hirose, Y.; Kawauchi, S.; Ogura, T.; Umesono, K.:
Aldehyde

[25981] dehydrogenase 6, a cytosolic retinaldehyde dehydroge-
nase prominently

[25982] expressed in sensory neuroepithelia during development.
J. Biol.

[25983] Chem. 275: 41210-41218, 2000.

[25984] 3358.

[25985] Hsu, L. C.; Chang, W.-C.; Hiraoka, L.; Hsieh, C.-L.: Molec-
ular

[25986] cloning, genomic organization, and chromosomal local-
ization of an

[25987] additional human aldehyde dehydrogenase gene, ALDH6.
Genomics 24:

[25988] 333-341, 1994.

[25989] 3359.

[25990] Kapfhamer, D.; Miller, D. E.; Lambert, S.; Bennett, V.; Glover,
[25991] T. W.; Burmeister, M.: Chromosomal localization of the ankyrin-G
[25992] gene (ANK3/Ank3) to human 10q21 and mouse 10. Genomics 27: 189-191,
[25993] 1995.

[25994] 3360.

[25995] Bennett, M.; Reed, R.: Correspondence between a mammalian spliceosome
[25996] component and an essential yeast splicing factor. Science 262: 105-108,
[25997] 1993.

[25998] 3361.

[25999] Dresser, D. W.; Hacker, A.; Lovell-Badge, R.; Guerrier, D.: The
[26000] genes for a spliceosome protein (SAP62) and the anti-Mullerian hormone
[26001] (AMH) are contiguous. Hum. Molec. Genet. 4: 1613-1618,

1995.

[26002] 3362.

[26003] Burks, D. J.; Font de Mora, J.; Schubert, M.; Withers, D. J.; Myers,

[26004] M. G.; Towery, H. H.; Altamuro, S. L.; Flint, C. L.; White, M. F.

[26005] : IRS-2 pathways integrate female reproduction and energy homeostasis. Nature 407:

[26006] 377-382, 2000.

[26007] 3363.

[26008] Fritsche, A.; Madaus, A.; Renn, W.; Tschritter, O.; Teigeler, A.;

[26009] Weisser, M.; Maerker, E.; Machicao, F.; Haring, H.; Stumvoll, M.:

[26010] The prevalent Gly1057Asp polymorphism in the insulin receptor substrate-2

[26011] gene is not associated with impaired insulin secretion. J. Clin.

[26012] Endocr. Metab. 86: 4822-4825, 2001.

[26013] 3364.

[26014] Kubota, N.; Tobe, K.; Terauchi, Y.; Eto, K.; Yamauchi, T.;

Suzuki,
[26015] R.; Tsubamoto, Y.; Komeda, K.; Nakano, R.; Miki, H.; Satoh, S.; Sekihara,
[26016] H.; Sciacchitano, S.; Lesniak, M.; Aizawa, S.; Nagai, R.; Kimura,
[26017] S.; Akanuma, Y.; Taylor, S. I.; Kadowaki, T.: Disruption of insulin
[26018] receptor substrate 2 causes type 2 diabetes because of liver insulin
[26019] resistance and lack of compensatory beta-cell hyperplasia. Diabetes 49:
[26020] :-1880-1889, 2000.

[26021] 3365.

[26022] Sun, X. J.; Wang, L.-M.; Zhang, Y.; Yenush, L.; Myers, M. G., Jr.;
[26023] Glasheen, E.; Lane, W. S.; Pierce, J. H.; White, M. F.: Role of IRS-2
[26024] in insulin and cytokine signalling. Nature 377: 173-177, 1995.
[26025] 3366.

[26026] Withers, D. J.; Burks, D. J.; Towery, H. H.; Altamuro, S. L.;

[26027] Flint, C. L.; White, M. F.: Irs-2 coordinates Igf-1 receptor-

mediated
[26028] beta-cell development and peripheral insulin signalling. Nature Genet. 23:
[26029] 32-40, 1999.

[26030] 3367.

[26031] Withers, D. J.; Gutierrez, J. S.; Towery, H.; Burks, D. J.; Ren,

[26032] J.-M.; Previs, S.; Zhang, Y.; Bernal, D.; Pons, S.; Shulman, G. I.;
[26033] Bonner-Weir, S.; White, M. F.: Disruption of IRS-2 causes type 2
[26034] diabetes in mice. Nature 391: 900-902, 1998.

[26035] 3368.

[26036] Eberle, F.; Dubreuil, P.; Mattei, M.-G.; Devilard, E.; Lopez, M.
[26037] : The human PRR2 gene, related to the human poliovirus receptor gene
[26038] (PVR), is the true homolog of the murine Mph gene. Gene 159: 267-272,
[26039] 1995.

[26040] 3369.

[26041] Morrison, M. E.; Racaniello, V. R.: Molecular cloning and

expression
[26042] of a murine homolog of the human poliovirus receptor gene. J. Virol. 66:
[26043] 2807-2813, 1992.

[26044] 3370.

[26045] Hanks, S. K.; Quinn, A. M.; Hunter, T.: The protein kinase family:
[26046] conserved features and deduced phylogeny of the catalytic domains. Science 241:
[26047] 42-52, 1988.

[26048] 3371.

[26049] Machado, R. D.; Pauciulo, M. W.; Thomson, J. R.; Lane, K. B.; Morgan,
[26050] N. V.; Wheeler, L.; Phillips, J. A., III; Newman, J.; Williams, D.;
[26051] Galie, N.; Manes, A.; McNeil, K.; and 11 others: BMPR2 haploinsufficiency
[26052] as the inherited molecular mechanism for primary pul- monary hypertension. Am.
[26053] J. Hum. Genet. 68: 92-102, 2001.

[26054] 3372.

[26055] Wong, W. T.; Carlomagno, F.; Druck, T.; Barletta, C.; Croce, C.

[26056] M.; Huebner, K.; Kraus, M. H.; Di Fiore, P. P.: Evolutionary conservation

[26057] of the EPS8 gene and its mapping to human chromosome 12q23-q24. Oncogene 9:

[26058] 3057-3061, 1994.

[26059] 3373.

[26060] Kobayashi, M.; Takamatsu, K.; Fujishiro, M.; Saitoh, S.; Noguchi,

[26061] T.: Molecular cloning of a novel calcium-binding protein structurally

[26062] related to hippocalcin from human brain and chromoso-mal mapping of

[26063] its gene. Biochim. Biophys. Acta 1222: 515-518, 1994.

[26064] 3374.

[26065] Avraham, K. B.; Levanon, D.; Negreanu, V.; Bernstein, Y.; Groner,

[26066] Y.; Copeland, N. G.; Jenkins, N. A.: Mapping of the mouse homolog

[26067] of the human runt domain gene, AML2, to the distal re-gion of mouse

[26068] chromosome 4. Genomics 25: 603-605, 1995.

[26069] 3375.

[26070] Bae, S.-C.; Takahashi, E.; Zhang, Y. W.; Ogawa, E.; Shige-
sada,
[26071] K.; Namba, Y.; Satake, M.; Ito, Y.: Cloning, mapping and
expression
[26072] of PEBP2-alpha-C, a third gene encoding the mammalian
Runt domain. Gene 159:
[26073] 245-248, 1995.

[26074] 3376.

[26075] Inoue, K,; Ozaki, S.; Shiga, T.; Ito, K.; Masuda, T.; Okado,
N.;
[26076] Iseda, T.; Kawaguchi, S.; Ogawa, M.; Bae, S.-C.; Yamashita,
N.; Itohara,
[26077] S.; Kudo, N.; Ito, Y.: Runx3 controls the axonal projection
of proprioceptive
[26078] dorsal root ganglion neurons. Nature Neurosci. 23Sept,
2002. Note:
[26079] Advance Electronic Publication.

[26080] 3377.

[26081] Li, Q.-L.; Ito, K.; Sakakura, C.; Fukamachi, H.; Inoue, K.;

Chi,
[26082] X.-Z.; Lee, K.-Y.; Nomura, S.; Lee, C.-W.; Han, S.-B.; Kim, H.-M.;
[26083] Kim, W.-J.; and 15 others: Causal relationship between the loss
[26084] of RUNX3 expression and gastric cancer. Cell 109: 113-124, 2002.
[26085] 3378.

[26086] Wijmenga, C.; Speck, N. A.; Dracopoli, N. C.; Hofker, M. H.; Liu,
[26087] P.; Collins, F. S.: Identification of a new murine runt do-main-containing
[26088] gene, Cbfa3, and localization of the human homolog, CBFA3, to chromosome
[26089] 1p35-pter. Genomics 26: 611-614, 1995.

[26090] 3379.

[26091] Cody, J. D.; Hale, D. E.; Brkanac, Z.; Kaye, C. I.; Leach, R. J.

[26092] : Growth hormone insufficiency associated with haploin-sufficiency
[26093] at 18q23. Am. J. Med. Genet. 71: 420-425, 1997.

[26094] 3380.

[26095] Habert-Ortoli, E.; Amiranoff, B.; Loquet, I.; Laburthe, M.; Mayaux,

[26096] J.-F.: Molecular cloning of a functional human galanin receptor. Proc.

[26097] Nat. Acad. Sci. 91: 9780-9783, 1994.

[26098] 3381.

[26099] Hecht, G.; Marrero, J. A.; Danilkovich, A.; Matkowskyj, K. A.;

[26100] Savkovic, S. D.; Koutsouris, A.; Benya, R. V.: Pathogenic Escherichia

[26101] coli increase Cl- secretion from intestinal epithelia by up-regulating

[26102] galanin-1 receptor expression. J. Clin. Invest. 104: 253-262, 1999.

[26103] 3382.

[26104] Scoggin, C. H.; Fisher, J. H.; Shoemaker, S. A.; Morse, H.; Leigh,

[26105] T.; Riccardi, V. M.: The E7-associated cell-surface antigen: a marker

[26106] for the 11p13 chromosomal deletion associated with aniridia-Wilms

[26107] tumor. Am. J. Hum. Genet. 37: 883-889, 1985.

[26108] 3383.

[26109] Avramopoulos, D.; Cox, T.; Blaschak, J. E.; Chakravarti, A.; Antonarakis,

[26110] S. E.: Linkage mapping of the AML1 gene on human chromosome 21 using

[26111] a DNA polymorphism in the 3-prime untranslated region. Genomics 14:

[26112] 506-507, 1992.

[26113] 3384.

[26114] Bernardin, F.; Yang, Y.; Cleaves, R.; Zahurak, M.; Cheng, L.; Civin,

[26115] C. I.; Friedman, A. D.: TEL-AML1, expressed from t(12;21) in human

[26116] acute lymphocytic leukemia, induces acute leukemia in mice. Cancer

[26117] Res. 62: 3904-3908, 2002.

[26118] 3385.

[26119] Cleary, M. L.: A new angle on a pervasive oncogene. Nature Genet. 23:

[26120] 134-135, 1999.

[26121] 3386.

[26122] Cuenco, G. M.; Nucifora, G.; Ren, R.: Human AML1/MDS1/EVI1 fusion
[26123] protein induces an acute myelogenous leukemia (AML) in mice: a model
[26124] for human AML. Proc. Nat. Acad. Sci. 97: 1760-1765, 2000.
[26125] 3387.

[26126] Gamou, T.; Kitamura, E.; Hosoda, F.; Shimizu, K.; Shinohara, K.;
[26127] Hayashi, Y.; Nagase, T.; Yokoyama, Y.; Ohki, M.: The partner gene
[26128] of AML1 in t(16;21) myeloid malignancies is a novel member of the
[26129] MTG8(ETO) family. Blood 91: 4028-4037, 1998.

[26130] 3388.

[26131] Johansson, B.; Mertens, F.; Mitelman, F.: Geographic heterogeneity
[26132] of neoplasia-associated chromosome aberrations. Genes Chromosomes
[26133] Cancer 3: 1-7, 1991.

[26134] 3389.

[26135] Levanon, D.; Negreanu, V.; Bernstein, Y.; Bar-Am, I.; Avivi, L.;

[26136] Groner, Y.: AML1, AML2, and AML3, the human members of the runt domain

[26137] gene-family: cDNA structure, expression, and chromosomal localization. Genomics 23:

[26138] 425-432, 1994.

[26139] 3390.

[26140] Look, A. T.: Oncogenic transcription factors in the human acute

[26141] leukemias. Science 278: 1059-1064, 1997.

[26142] 3391.

[26143] McNeil, S.; Zeng, C.; Harrington, K. S.; Hiebert, S.; Lian, J.

[26144] B.; Stein, J. L.; van Wijnen, A. J.; Stein, G. S.: The t(8;21) chromosomal

[26145] translocation in acute myelogenous leukemia modifies intranuclear

[26146] targeting of the AML1/CBF-alpha2 transcription factor. Proc. Nat.

[26147] Acad. Sci. 96: 14882-14887, 1999.

[26148] 3392.

[26149] Burt, R. W.; Berenson, M. M.; Lee, R. G.; Tolman, K. G.; Freston,
[26150] J. W.; Gardner, E. J.: Upper gastrointestinal polyps in Gardner's
[26151] syndrome. Gastroenterology 86: 295-301, 1984.

[26152] 3393.

[26153] Butson, A. R. C.: Familial multiple polyposis coli with multiple
[26154] associated tumors. Dis. Colon Rectum 26: 578-582, 1983.

[26155] 3394.

[26156] Cachon-Gonzalez, M. B.; Delhanty, J. D. A.; Burn, J.; Tsioupra,
[26157] K.; Davis, M. B.; Attwood, J.; Chapman, P.: Linkage analysis in adenomatous
[26158] polyposis coli: the use of four closely linked DNA probes in 20 UK
[26159] families. J. Med. Genet. 28: 681-685, 1991.

[26160] 3395.

[26161] Calin, G.; Wijnen, J.; van der Klift, H.; Ionita, A.; Mulder,

[26162] A.; Breukel, C.; Smits, R.; Dauwerse, H.; Hansson, K.; Calin, S.;

[26163] Stefanescu, D.; Oproiu, A.; Fodde, R.: Marfan-like habitus and familial

[26164] adenomatous polyposis in two unrelated males: a significant association? Europ.

[26165] J. Hum. Genet. 7: 609-614, 1999.

[26166] 3396.

[26167] Cama, A.; Esposito, D. L.; Palmirotta, R.; Curia, M. C.; Ranieri,

[26168] A.; Ficari, F.; Valanzano, R.; Modesti, A.; Battista, P.; Tonelli,

[26169] F.; Mariani-Costantini, R.: A novel mutation at the splice junction

[26170] of exon 9 of the APC gene in familial adenomatous polyposis. Hum.

[26171] Mutat. 3: 305-308, 1994.

[26172] 3397.

[26173] Camiel, M. R.; Mule, J. E.; Alexander, L. L.; Benninghoff, D.

[26174] L.: Association of thyroid carcinoma with Gardner's syndrome in siblings. New

[26175] Eng. J. Med. 278: 1056-1058, 1968.

[26176] 3398.

[26177] Cao, X.; Eu, K. W.; Seow-Choen, F.; Zao, Y.; Cheah, P. Y.: APC
[26178] mutation and phenotypic spectrum of Singapore familial adenomatous
[26179] polyposis patients. Europ. J. Hum. Genet. 8: 42-48, 2000.

[26180] 3399.

[26181] Cao, X.; Eu, K. W.; Seow-Choen, F.; Zhao, Y.; Cheah, P. Y.: Topoisomerase-I-
[26182] and Alu-mediated genomic deletions of the APC gene in familial adenomatous
[26183] polyposis. Hum. Genet. 108: 436-442, 2001.

[26184] 3400.

[26185] Capps, W. F., Jr.; Lewis, M. I.; Gazzaniga, D. A.: Carcinoma
[26186] of the colon, ampulla of Vater and urinary bladder associated with
[26187] familial multiple polyposis: a case report. Dis. Colon Rectum 11:
[26188] 298-304, 1968.

[26189] 3401.

[26190] Carl, W.; Herrera, L.: Dental and bone abnormalities in patients

[26191] with familial polyposis coli. Sem. Surg. Oncol 3: 77-83, 1987.

[26192] 3402.

[26193] Caspari, R.; Friedl, W.; Mandl, M.; Moslein, G.; Kadmon, M.; Knapp,

[26194] M.; Jacobasch, K.-H.; Ecker, K.-W.; Kreissler-Haag, D.; Timmermanns,

[26195] G.; Propping, P.: Familial adenomatous polyposis: mutation at codon

[26196] 1309 and early onset of colon cancer. Lancet 343: 629-632, 1994.

[26197] 3403.

[26198] Caspari, R.; Olschwang, S.; Friedl, W.; Mandl, M.; Boisson, C.;

[26199] Boker, T.; Augustin, A.; Kadmon, M.; Moslein, G.; Thomas, G.; Propping,

[26200] P.: Familial adenomatous polyposis: desmoid tumours and lack of ophthalmic

[26201] lesions (CHRPE) associated with APC mutations beyond codon 1444. Hum.

[26202] Molec. Genet. 4: 337-340, 1995.

[26203] 3404.

[26204] Chang, C. H.; Platt, E. D.; Thomas, K. E.; Watne, A. L.: Bone

[26205] abnormalities in Gardner's syndrome. Am. J. Roentgen.
103: 645-652,

[26206] 1968.

[26207] 3405.

[26208] Chapman, P. D.; Church, W.; Burn, J.; Gunn, A.: Congenital
hypertrophy

[26209] of retinal pigment epithelium: a sign of familial adenoma-
tous polyposis. Brit.

[26210] Med. J. 298: 353-354, 1989.

[26211] 3406.

[26212] Chau, I.; Cunningham, D.: Cyclooxygenase inhibition in
cancer

[26213] - a blind alley or a new therapeutic reality? (Editorial) New
Eng.

[26214] J. Med. 346: 1085-1087, 2002.

[26215] 3407.

[26216] Chen, S.; Kazim, D.; Kraveka, J.; Pollack, R. E.: Skin fibrob-
lasts

[26217] from individuals hemizygous for the familial adenopoly-
posis susceptibility

[26218] gene show delayed crisis in vitro. Proc. Nat. Acad. Sci. 86: 2008-2012,

[26219] 1989.

[26220] 3408.

[26221] Clark, S. K.; Neale, K. F.; Landgrebe, J. C.; Phillips, R. K.

[26222] S.: Desmoid tumours complicating familial adenomatous polyposis. Brit.

[26223] J. Surg. 86: 1185-1189, 1999.

[26224] 3409.

[26225] Cottrell, S.; Bicknell, D.; Kaklamanis, L.; Bodmer, W. F.: Molecular

[26226] analysis of APC mutations in familial adenomatous poly- posis and sporadic

[26227] colon carcinomas. Lancet 340: 626-630, 1992.

[26228] 3410.

[26229] Couture, J.; Mitri, A.; Lagace, R.; Smits, R.; Berk, T.; Bouchard,

[26230] H.-L.; Fodde, R.; Alman, B.; Bapat, B.: A germline mutation at the

[26231] extreme 3-prime end of the APC gene results in a severe desmoid phenotype

[26232] and is associated with overexpression of beta-catenin in the desmoid
[26233] tumor. Clin. Genet. 57: 205-212, 2000.

[26234] 3411.

[26235] Crabtree, M. D.; Tomlinson, I. P. M.; Talbot, I. C.; Phillips,

[26236] R. K. S.: Variability in the severity of colonic disease in fa-
milial
[26237] adenomatous polyposis results from differences in tumour initiation
[26238] rather than progression and depends relatively little on patient age. Gut 49:
[26239] 540-543, 2001.

[26240] 3412.

[26241] Cross, I.; Delhanty, J.; Chapman, P.; Bowles, L. V.; Griffin,

[26242] D.; Wolstenholme, J.; Bradburn, M.; Brown, J.; Wood, C.; Gunn, A.;
[26243] Burn, J.: An intrachromosomal insertion causing 5q22 deletion and
[26244] familial adenomatous polyposis coli in two generations. J. Med. Genet. 29:
[26245] 175-179, 1992.

[26246] 3413.

[26247] Curia, M. C.; Esposito, D. L.; Aceto, G.; Palmirotta, R.; Crognale,

[26248] S.; Valanzano, R.; Ficari, F.; Tonelli, F.; Battista, P.; Mariani-Costantini,

[26249] R.; Cama, A.: Transcript dosage effect in familial adenomatous polyposis:

[26250] model offered by two kindreds with exon 9 APC gene mutations. Hum.

[26251] Mutat. 11: 197-201, 1998.

[26252] 3414.

[26253] Danes, B. S.: The Gardner syndrome: increased tetraploidy in

[26254] cultured skin fibroblast. J. Med. Genet. 13: 52-56, 1976.

[26255] 3415.

[26256] Danes, B. S.: The Gardner syndrome. Cancer 36: 2327-2333, 1975.

[26257] 3416.

[26258] Danes, B. S.; Gardner, E. J.: The Gardner syndrome: a cell culture

[26259] study of kindred 109. J. Med. Genet. 15: 346-551, 1978.

[26260] 3417.

[26261] Davies, D. R.; Armstrong, J. G.; Thakker, N.; Horner, K.; Guy,

[26262] S. P.; Clancy, T.; Sloan, P.; Blair, V.; Dodd, C.; Warnes, T. W.;

[26263] Harris, R.; Evans, D. G. R.: Severe Gardner syndrome in families

[26264] with mutations restricted to a specific region of the APC gene. Am.

[26265] J. Hum. Genet. 57: 1151-1158, 1995.

[26266] 3418.

[26267] Denzler, T. B.; Harned, R. K.; Pergam, C. J.: Gastric polyps

[26268] in familial polyposis coli. Radiology 130: 63-66, 1979.

[26269] 3419.

[26270] Deuter, R.; Muller, O.: Detection of APC mutations in stool DNA

[26271] of patients with colorectal cancer by HD-PCR. Hum. Mutat. 11: 84-89,

[26272] 1998.

[26273] 3420.

[26274] De Vries, E. M. G.; Ricke, D. O.; De Vries, T. N.; Hartmann, A.;
[26275] Blaszyk, H.; Liao, D.; Soussi, T.; Kovach, J. S.; Sommer, S. S.:
[26276] Database of mutations in the p53 and APC tumor suppressor genes designed
[26277] to facilitate molecular epidemiological analyses. Hum. Mutat. 7:
[26278] 202-213, 1996.

[26279] 3421.

[26280] Dhaliwal, M. K.; Hughes, J. I.; Jackson, G. L.; Pathak, S.: Multiple
[26281] polyposis coli associated with Gardner's syndrome and chromosomal
[26282] mosaicism: a family analysis. Am. J. Gastroenterol. 85: 880-883,
[26283] 1990.

[26284] 3422.

[26285] Diaz-Llopis, M.; Menezo, J. L.: Congenital hypertrophy of the
[26286] retinal pigment epithelium in familial adenomatous polyposis. Arch.

[26287] Ophthal. 106: 412-413, 1988.

[26288] 3423.

[26289] Bielinska, B.; Blaydes, S. M.; Buiting, K.; Yang, T.; Krajew-
[26290] ska-Walasek, M.; Horsthemke, B.; Brannan, C. I.: De novo deletions of SNRPN exon
[26291] 1 in early human and mouse embryos result in a paternal to maternal
[26292] imprint switch. Nature Genet. 25: 74-78, 2000. Note: Er-
[26293] ratum: Nature Genet. 25: 241 only, 2000.

[26294] 3424.

[26295] Lorenzo, F.; del Giudice, E. M.; Alloisio, N.; Morle, L.;
[26296] Forissier, A.; Perrotta, S.; Sciarratta, G.; Iolascon, A.; Delaunay, J.: Severe
[26297] poikilocytosis associated with a de novo alpha-28 arg-
[26298] to-cys mutation in spectrin. Brit. J. Haemat. 83: 152-157, 1993.

[26299] 3425.

[26300] Marchesi, S. L.: Personal Communication. New Haven,

Conn.

[26301] 3/1990.

[26302] 3426.

[26303] Marchesi, S. L.; Agre, P. A.; Speicher, D. W.; Tse, W. T.; Forget,

[26304] B. G.: Mutant spectrin alpha-II domain in recessively inherited spherocytosis.

[26305] (Abstract) Blood 74: 182A, 1989.

[26306] 3427.

[26307] Marchesi, S. L.; Knowles, W. J.; Morrow, J. S.; Bologna, M.; Marchesi,

[26308] V. T.: Abnormal spectrin in hereditary elliptocytosis. Blood 67:

[26309] 141-151, 1986.

[26310] 3428.

[26311] Marchesi, S. L.; Letsinger, J. T.; Speicher, D. W.; Marchesi,

[26312] V. T.; Agre, P.; Hyun, B.; Gulati, G.: Mutant forms of spectrin alpha-subunits

[26313] in hereditary elliptocytosis. J. Clin. Invest. 80: 191-198, 1987.

[26314] 3429.

[26315] Middleton-Price, H.; Gendler, S.; Malcolm, S.: Close linkage
[26316] of PUM and SPTA within chromosome band 1q21. Ann. Hum. Genet. 52:
[26317] 273-278, 1988.

[26318] 3430.

[26319] Moon, R. T.; Lazarides, E.: Beta-spectrin limits alpha-spectrin
[26320] assembly on membranes following synthesis in a chicken erythroid cell
[26321] lysate. Nature 305: 62-65, 1983.

[26322] 3431.

[26323] Morle, L.; Alloisio, N.; Ducluzeau, M. T.; Pothier, B.; Bilbech,
[26324] R.; Kastally, R.; Delaunay, J.: Spectrin Tunis (alpha-I/78): a new
[26325] alpha-I variant that causes asymptomatic hereditary elliptocytosis
[26326] in the heterozygous state. Blood 71: 508-511, 1988.

[26327] 3432.

[26328] Morle, L.; Morle, F.; Roux, A. F.; Godet, J.; Forget, B. G.;

Denoroy,

[26329] L.; Garbarz, M.; Dhermy, D.; Kastally, R.; Delaunay, J.: Spectrin

[26330] Tunis (Sp-alpha-I/78), an elliptocytogenic variant, is due to the

[26331] CGG-to-TGG codon change (arg-to-trp) at position 35 of the alpha-I

[26332] domain. Blood 74: 828-832, 1989.

[26333] 3433.

[26334] Morle, L.; Roux, A.-F.; Alloisio, N.; Pothier, B.; Starck, J.;

[26335] Denoroy, J.; Morle, F.; Rudigoz, R.-C.; Forget, B. G.; De-launay, J.;

[26336] Godet, J.: Two elliptocytogenic alpha-I/74 variants of the spectrin

[26337] alpha-I domain: spectrin Culoz and spectrin Lyon, mu-tated at position

[26338] 40 (GGT-to-GTT; gly-to-val) and at position 43 (CTT-to-TTT; leu-to-phe),

[26339] respectively. (Abstract) Blood 74: 60A, 1989.

[26340] 3434.

[26341] Morle, L.; Roux, A.-F.; Alloisio, N.; Pothier, B.; Starck, J.;

[26342] Denoroy, L.; Morle, F.; Rudigoz, R.-C.; Forget, B. G.; De-
launay, J.;
[26343] Godet, J.: Two elliptocytogenic alpha(I/74) variants of the
spectrin
[26344] alpha-I domain: spectrin Culoz (GGT-to-GTT; alpha-I 40
gly-to-val)
[26345] and spectrin Lyon (CTT-to-TTT; alpha-I 43 leu-to-phe). J.
Clin. Invest. 86:
[26346] 548-554, 1990.

[26347] 3435.

[26348] Motulsky, A. G.; Singer, K.; Crosby, W. H.; Smith, V.: The
life
[26349] span of the elliptocyte: hereditary elliptocytosis and its re-
lationship
[26350] to other familial hemolytic diseases. Blood 9: 57-72,
1954.
[26351] 3436.

[26352] Pakstis, A. J.; Miki, T.; Kidd, J. R.; Kidd, K. K.: D1S75 is

[26353] polymorphic in Caucasians as well as Japanese and maps
between AT3
[26354] and SPTA1 on chromosome 1q. (Abstract) Cytogenet. Cell
Genet. 51:

[26355] 1057, 1989.

[26356] 3437.

[26357] Palek, J.; Coetzer, T.: Clinical expression of alpha-spectrin

[26358] mutants in hereditary elliptocytosis. Blood Cells 13: 237-250, 1987.

[26359] 3438.

[26360] Parquet, N.; Devaux, I.; Boulanger, L.; Galand, C.; Boivin, P.;

[26361] Lecomte, M.-C.; Dhermy, D.; Garbarz, M.: Identification of three

[26362] novel spectrin alphaI/74 mutations in hereditary elliptocytosis: further

[26363] support for a triple-stranded folding unit model of the spectrin heterodimer

[26364] contact site. Blood 84: 303-308, 1994.

[26365] 3439.

[26366] Perrotta, P.; del Giudice, E. M.; Alloisio, N.; Sciarratta, G.;

[26367] Pinto, L.; Delaunay, J.; Cutillo, S.; Iolascon, A.: Mild elliptocytosis

[26368] associated with the alpha-34 arg-to-trp mutation in spectrin Genova

[26369] (alpha-I/74). Blood 83: 3346-3349, 1994.

[26370] 3440.

[26371] Raeymaekers, P.; Van Broeckhoven, C.; Backhovens, H.; Wehnert,

[26372] A.; Muylle, L.; De Jonghe, P.; Gheuens, J.; Vandenberghe, A.: The

[26373] Duffy blood group is linked to the alpha-spectrin locus in a large

[26374] pedigree with autosomal dominant inheritance of Char-cot-Marie-Tooth

[26375] disease type 1. Hum. Genet. 78: 76-78, 1988.

[26376] 3441.

[26377] Randon, J.; Boulanger, L.; Marechal, J.; Garbarz, M.; Vallier,

[26378] A.; Ribeiro, L.; Tamagnini, G.; Dhermy, D.; Delaunay, J.: A variant

[26379] of spectrin low-expression allele alpha-LELY carrying a hereditary

[26380] elliptocytosis mutation in codon 28. Brit. J. Haemat. 88: 534-540,

[26381] 1994.

[26382] 3442.

[26383] Ravindranath, Y.; Johnson, R. M.: Altered spectrin association
[26384] and membrane fragility without abnormal spectrin heat sensitivity
[26385] in a case of congenital hemolytic anemia. Am. J. Hemat. 20: 53-65,
[26386] 1985.

[26387] 3443.

[26388] Roux, A.-F.; Morle, F.; Guetarni, D.; Colonna, P.; Sahr, K.; Forget,
[26389] B. G.; Delaunay, J.; Godet, J.: Molecular basis of Sp alpha(I/65)
[26390] hereditary elliptocytosis in North Africa: insertion of a TTG triplet
[26391] between codons 147 and 149 in the alpha-spectrin gene from five unrelated
[26392] families. Blood 73: 2196-2201, 1989.

[26393] 3444.

[26394] Sahr, K. E.; Laurila, P.; Kotula, L.; Scarpa, A. L.; Coupal, E.;
[26395] Leto, T. L.; Linnenbach, A. J.; Winkelmann, J. C.; Speicher, D. W.;

[26396] Marchesi, V. T.; Curtis, P. J.; Forget, B. G.: The complete cDNA
[26397] and polypeptide sequences of human erythroid alpha-spectrin. J. Biol.
[26398] Chem. 265: 4434-4443, 1990.

[26399] 3445.

[26400] Sahr, K. E.; Tobe, T.; Scarpa, A.; Laughinghouse, K.; Marchesi,
[26401] S. L.; Agre, P.; Linnenbach, A. J.; Marchesi, V. T.; Forget, B. G.
[26402] : Sequence and exon-intron organization of the DNA encoding the alpha-I
[26403] domain of human spectrin: application to the study of mutations causing
[26404] hereditary elliptocytosis. J. Clin. Invest. 84: 1243-1252, 1989.
[26405] 3446.

[26406] Speicher, D. W.; Weglarz, L.; DeSilva, T. M.: Properties of human
[26407] red cell spectrin heterodimer (side-to-side) assembly and identification
[26408] of an essential nucleation site. J. Biol. Chem. 267:

14775-14782,

[26409] 1992.

[26410] 3447.

[26411] Eagle, L. R.; Yin, X.; Brothman, A. R.; Williams, B. J.; Atkin,

[26412] N. B.; Prochownik, E. V.: Mutation of the MXI1 gene in prostate cancer. Nature

[26413] Genet. 9: 249-255, 1995.

[26414] 3448.

[26415] Edwards, A.; Hammond, H. A.; Jin, L.; Caskey, C. T.; Chakraborty,

[26416] R.: Genetic variation at five trimeric and tetrameric tandem repeat

[26417] loci in four human population groups. Genomics 12: 241-253, 1992.

[26418] 3449.

[26419] Ichikawa, T.; Ichikawa, Y.; Dong, J.; Hawkins, A. L.; Griffin,

[26420] C. A.; Isaacs, W. B.; Oshimura, M.; Barrett, J. C.; Isaacs, J. T.

[26421] : Localization of metastasis suppressor gene(s) for pro-static cancer

[26422] to the short arm of human chromosome 11. Cancer Res.

52: 3486-3490,
[26423] 1992.

[26424] 3450.

[26425] Ichikawa, T.; Ichikawa, Y.; Isaacs, J. T.: Genetic factors and

[26426] suppression of metastatic ability of prostatic cancer. Cancer Res. 51:
[26427] 3788-3792, 1991.

[26428] 3451.

[26429] Irvine, R. A.; Yu, M. C.; Ross, R. K.; Coetzee, G. A.: The CAG
[26430] and GGC microsatellites of the androgen receptor gene are in linkage
[26431] disequilibrium in men with prostate cancer. Cancer. Res. 55: 1937-1940,
[26432] 1995.

[26433] 3452.

[26434] Peters, M. A.; Ostrander, E. A.: Prostate cancer: more than two
[26435] to tango. Nature Genet. 27: 134-135, 2001.

[26436] 3453.

[26437] Bansal, A.; Murray, D. K.; Wu, J. T.; Stephenson, R. A.; Middleton,

[26438] R. G.; Meikle, A. W.: Heritability of prostate-specific antigen and

[26439] relationship with zonal prostate volumes in aging twins. J. Clin.

[26440] Endocr. Metab. 85: 1272-1276, 2000.

[26441] 3454.

[26442] Cleutjens, K. B. J. M.; van der Korput, H. A. G. M.; van Eekelen,

[26443] C. C. E. M.; van Rooij, H. C. J.; Faber, P. W.; Trapman, J.: An androgen

[26444] response element in a far upstream enhancer region is essential for

[26445] high, androgen-regulated activity of the prostate-specific antigen

[26446] promoter. Molec. Endocr. 11: 148-161, 1997.

[26447] 3455.

[26448] Nelson, R. J.; Demas, G. E.; Huang, P. L.; Fishman, M. C.; Dawson,

[26449] V. L.; Dawson, T. M.; Snyder, S. H.: Behavioural abnormalities in

[26450] male mice lacking neuronal nitric oxide synthase. Nature 378: 383-386,

[26451] 1995.

[26452] 3456.

[26453] Henry, J. G.; Mitnick, M.; Dann, P. R.; Stewart, A. F.: Parathyroid

[26454] hormone-related protein-(1-36) is biologically active when administered

[26455] subcutaneously to humans. J. Clin. Endocr. Metab. 82: 900-906, 1997.

[26456] 3457.

[26457] Magnaghi, P.; Agazzi, A.; Semino, O.; Ferrari, M.; Barbui, T.;

[26458] D'Angelo, A.; Taramelli, R.: A recombination event in the closely

[26459] linked plasminogen and apolipoprotein(a) gene loci. Clin. Genet. 47:

[26460] 285-289, 1995.

[26461] 3458.

[26462] Schwarz, H. P.; Fischer, M.; Hopmeier, P.; Batard, M. A.; Griffin,

[26463] J. H.: Plasma protein S deficiency in familial thrombotic disease. Blood 64:

[26464] 1297-1300, 1984.

[26465] 3459.

[26466] Kan, S.; Elanko, N.; Johnson, D.; Cornejo-Roldan, L.; Cook, J.;

[26467] Reich, E. W.; Tomkins, S.; Verloes, A.; Twigg, S. R. F.; Rannan-Eliya,

[26468] S.; McDonald-McGinn, D. M.; Zackai, E. H.; Wall, S. A.; Muenke, M.;

[26469] Wilkie, A. O. M.: Genomic screening of fibroblast growth-factor receptor

[26470] 2 reveals a wide spectrum of mutations in patients with syndromic

[26471] craniosynostosis. Am. J. Hum. Genet. 70: 472-486, 2002.

[26472] 3460.

[26473] Zhuchenko, O.; Bailey, J.; Bonnen, P.; Ashizawa, T.; Stockton,

[26474] D. W.; Amos, C.; Dobyns, W. B.; Subramony, S. H.; Zoghbi, H. Y.; Lee,

[26475] C. C.: Autosomal dominant cerebellar ataxia (SCA6) associated with

[26476] small polyglutamine expansions in the al-
pha(1A)-voltage-dependent
[26477] calcium channel. Nature Genet. 15: 62-69, 1997.

[26478] 3461.

[26479] Imbert, G.; Saudou, F.; Yvert, G.; Devys, D.; Trottier, Y.;
Garnier,
[26480] J.-M.; Weber, C.; Mandel, J.-L.; Cancel, G.; Abbas, N.; Durr,
A.;
[26481] Didierjean, O.; Stevanin, G.; Agid, Y.; Brice, A.: Cloning of
the
[26482] gene for spinocerebellar ataxia 2 reveals a locus with high
sensitivity
[26483] to expanded CAG/glutamine repeats. Nature Genet. 14:
285-291, 1996.
[26484] 3462.

[26485] Holick, M. F.; Ray, S.; Chen, T. C.; Tian, X.; Persons, K. S.:

[26486] A parathyroid hormone antagonist stimulates epidermal
proliferation
[26487] and hair growth in mice. Proc. Nat. Acad. Sci. 91:
8014-8016, 1994.
[26488] 3463.

[26489] Lanske, B.; Amling, M.; Neff, L.; Guiducci, J.; Baron, R.; Kronenberg,
[26490] H. M.: Ablation of the PTHrP gene or the PTH/PTHrP receptor gene
[26491] leads to distinct abnormalities in bone development. J. Clin. Invest. 104:
[26492] 399-407, 1999.

[26493] 3464.

[26494] Mangin, M.; Ikeda, K.; Dreyer, B. E.; Broadus, A. E.: Isolation
[26495] and characterization of the human parathyroid hormone-like peptide
[26496] gene. Proc. Nat. Acad. Sci. 86: 2408-2412, 1989.

[26497] 3465.

[26498] Mangin, M.; Webb, A. C.; Dreyer, B. E.; Posillico, J. T.; Ikeda,
[26499] K.; Weir, E. C.; Stewart, A. F.; Bander, N. H.; Milstone, L.; Barton,
[26500] D. E.; Francke, U.; Broadus, A. E.: Identification of a cDNA encoding
[26501] a parathyroid hormone-like peptide from a human tumor associated with

[26502] humoral hypercalcemia of malignancy. Proc. Nat. Acad. Sci. 85: 597-601,

[26503] 1988.

[26504] 3466.

[26505] Moseley, J. M.; Kubota, M.; Diefenbach-Jagger, H.; Wettenhall,

[26506] R. E. H.; Kemp, B. E.; Suva, L. J.; Rodda, C. P.; Ebeling, P. R.;

[26507] Hudson, P. J.; Zajac, J. D.; Martin, T. J.: Parathyroid hormone-related

[26508] protein purified from a human lung cancer cell line. Proc. Nat. Acad.

[26509] Sci. 84: 5048-5052, 1987.

[26510] 3467.

[26511] Philbrick, W. M.; Dreyer, B. E.; Nakchbandi, I. A.; Karaplis,

[26512] A. C.: Parathyroid hormone-related protein is required for tooth

[26513] eruption. Proc. Nat. Acad. Sci. 95: 11846-11851, 1998.

[26514] 3468.

[26515] Strewler, G. J.: The physiology of parathyroid hormone-related

[26516] protein. New Eng. J. Med. 342: 177-185, 2000.

[26517] 3469.

[26518] Suva, L. J.; Winslow, G. A.; Wettenhall, R. E. H.; Hammonds, R.

[26519] G.; Moseley, J. M.; Diefenbach-Jagger, H.; Rodda, C. P.; Kemp, B.

[26520] E.; Rodriguez, H.; Chen, E. Y.; Hudson, P. J.; Martin, T. J.; Wood,

[26521] W. I.: A parathyroid hormone-related protein implicated in malignant

[26522] hypercalcemia: cloning and expression. Science 237: 893-896, 1987.

[26523] 3470.

[26524] Vortkamp, A.; Lee, K.; Lanske, B.; Segre, G. V.; Kronenberg, H.

[26525] M.; Tabin, C. J.: Regulation of rate of cartilage differentiation

[26526] by Indian hedgehog and PTH-related protein. Science 273: 613-622,

[26527] 1996.

[26528] 3471.

[26529] Wysolmerski, J. J.; Cormier, S.; Philbrick, W. M.; Dann, P.; Zhang,

[26530] J.-P.; Roume, J.; Delezoide, A.-L.; Silve, C.: Absence of functional

[26531] type 1 parathyroid hormone (PTH)/PTH-related protein receptors in

[26532] humans is associated with abnormal breast development and tooth impaction. J.

[26533] Clin. Endocr. Metab. 86: 1788-1794, 2001.

[26534] 3472.

[26535] Wysolmerski, J. J.; Philbrick, W. M.; Dunbar, M. E.; Lanske, B.;

[26536] Kronenberg, H.; Broadus, A. E.: Rescue of the parathyroid hormone-related

[26537] protein knockout mouse demonstrates that parathyroid hormone-related
[26538] protein is essential for mammary gland development. De-velopment 125:
[26539] 1285-1294, 1998.

[26540] 3473.

[26541] Yasuda, T.; Banville, D.; Hendy, G. N.; Goltzman, D.: Char-acterization
[26542] of the human parathyroid hormone-like peptide gene: functional and
[26543] evolutionary aspects. J. Biol. Chem. 264: 7720-7725, 1989.
[26544] 3474.

[26545] Mavrogiannis, L. A.; Antonopoulou, I.; Baxova, A.; Kutilek, S.;
[26546] Kim, C. A.; Sugayama, S. M.; Salamanca, A.; Wall, S. A.; Morriss-Kay,
[26547] G. M.; Wilkie, A. O. M.: Haploinsufficiency of the human homeobox
[26548] gene ALX4 causes skull ossification defects. Nature Genet. 27: 17-18,
[26549] 2001.

[26550] 3475.

[26551] Qu, S.; Niswender, K. D.; Ji, Q.; van der Meer, R.; Keeney, D.;

[26552] Magnuson, M. A.; Wisdom, R.: Polydactyly and ectopic ZPA formation

[26553] in Alx-4 mutant mice. Development 124: 3999-4008, 1997.

[26554] 3476.

[26555] Wu, Y.-Q.; Badano, J. L.; McCaskill, C.; Vogel, H.; Potocki, L.;

[26556] Shaffer, L. G.: Haploinsufficiency of ALX4 as a potential cause of

[26557] parietal foramina in the 11p11.2 contiguous gene-deletion syndrome. Am.

[26558] J. Hum. Genet. 67: 1327-1332, 2000.

[26559] 3477.

[26560] Wuyts, W.; Cleiren, E.; Homfray, T.; Rasore-Quartino, A.; Vanhoenacker,

[26561] F.; Van Hul, W.: The ALX4 homeobox gene is mutated in patients with

[26562] ossification defects of the skull (foramina parietalia per-magna, OMIM

[26563] 168500). J. Med. Genet. 37: 916-920, 2000.

[26564] 3478.

[26565] Fischer, M. B.; Roeckl, C.; Parizek, P.; Schwarz, H. P.; Aguzzi,

[26566] A.: Binding of disease-associated prion protein to plasminogen. Nature 408:

[26567] 479-483, 2000.

[26568] 3479.

[26569] Cerone, R.; Holme, E.; Schiaffino, M. C.; Caruso, U.; Maritano,

[26570] L.; Romano, C.: Tyrosinemia type III: diagnosis and ten-year follow-up. Acta

[26571] Paediat. 86: 1013-1015, 1997.

[26572] 3480.

[26573] Lemyre, E.; Russo, P.; Melancon, S. B.; Gagne, R.; Potier, M.;

[26574] Lambert, M.: Clinical spectrum of infantile free sialic acid storage

[26575] disease. Am. J. Med. Genet. 82: 385-391, 1999.

[26576] 3481.

[26577] Mancini, G. M. S.; Beerens, C. E. M. T.; Aula, P. P.; Verheijen,
[26578] F. W.: Sialic acid storage diseases: a multiple lysosomal transport
[26579] defect for acidic monosaccharides. J. Clin. Invest. 87: 1329-1335,
[26580] 1991.

[26581] 3482.

[26582] Schleutker, J.; Laine, A.-P.; Haataja, L.; Renlund, M.; Weissenbach,
[26583] J.; Aula, P.; Peltonen, L.: Linkage disequilibrium utilized to establish
[26584] a refined genetic position of the Salla disease locus on 6q14-q15. Genomics 27:
[26585] 286-292, 1995.

[26586] 3483.

[26587] Schleutker, J.; Leppanen, P.; Mansson, J.-E.; Erikson, A.; Weissenbach,
[26588] J.; Peltonen, L.; Aula, P.: Lysosomal free sialic acid storage disorders
[26589] with different phenotypic presentations--infantile-form sialic acid

[26590] storage disease and Salla disease--represent allelic disorders on

[26591] 6q14-15. Am. J. Hum. Genet. 57: 893-901, 1995.

[26592] 3484.

[26593] Tondeur, M.; Libert, J.; Vamos, E.; Van Hoof, F.; Thomas, G. H.;

[26594] Strecker, G.: Infantile form of sialic acid storage disorder: clinical,

[26595] ultrastructural, and biochemical studies in two siblings. Europ.

[26596] J. Pediat. 139: 142-147, 1982.

[26597] 3485.

[26598] Verheijen, F. W.; Verbeek, E.; Aula, N.; Beerens, C. E. M. T.;

[26599] Havelaar, A. C.; Joosse, M.; Peltonen, L.; Aula, P.; Galjaard, H.;

[26600] van der Spek, P. J.; Mancini, G. M. S.: A new gene, encoding an anion

[26601] transporter, is mutated in sialic acid storage diseases. Nature Genet. 23:

[26602] 462-465, 1999.

[26603] 3486.

[26604] Krasnewich, D. M.; Tietze, F.; Krause, W.; Pretzlaff, R.; Wenger,
[26605] D. A.; Diwadkar, V.; Gahl, W. A.: Clinical and biochemical studies
[26606] in an American child with sialuria. Biochem. Med. Metab. Biol. 49:
[26607] 90-96, 1993.

[26608] 3487.

[26609] Leroy, J. G.; Seppala, R.; Huizing, M.; Dacremont, G.; De Simpel,
[26610] H.; Van Coster, R. N.; Orvisky, E.; Krasnewich, D. M.; Gahl, W. A.
[26611] : Dominant inheritance of sialuria, an inborn error of feedback inhibition. Am.
[26612] J. Hum. Genet. 68: 1419-1427, 2001.

[26613] 3488.

[26614] Seppala, R.; Lehto, V.-P.; Gahl, W. A.: Mutations in the hu-man
[26615] UDP-N-acetylglucosamine 2-epimerase gene define the disease sialuria
[26616] and the allosteric site of the enzyme. Am. J. Hum. Genet. 64: 1563-1569,

[26617] 1999.

[26618] 3489.

[26619] Cho, Y.-Y.; Kang, M.-J.; Sone, H.; Suzuki, T.; Abe, M.; Igarashi,

[26620] M.; Tokunaga, T.; Ogawa, S.; Takei, Y. A.; Miyazawa, T.; Sasano, H.;

[26621] Fujino, T.; Yamamoto, T. T.: Abnormal uterus with poly-cysts, accumulation

[26622] of uterine prostaglandins, and reduced fertility in mice heterozygous

[26623] for acyl-CoA synthetase 4 deficiency. Biochem. Biophys. Res. Commun. 284:

[26624] 993-997, 2001.

[26625] 3490.

[26626] Kay, M. A.; High, K.: Gene therapy for the hemophilias. (Commentary) Proc.

[26627] Nat. Acad. Sci. 96: 9973-9975, 1999.

[26628]

[26629] 100. Kazazian, H. H., Jr.; Wong, C.; Youssoufian, H.; Scott, A. F.;

[26630] Phillips, D. G.; Antonarakis, S. E.: Haemophilia A resulting

from
de novo insertion of L1 sequences represents a novel mechanism for
mutation in man. Nature 332: 164-166, 1988.

101. Kemball-Cook, G.; Tuddenham, E. G. D.: The factor VIII mutation
database on the World Wide Web: the haemophilia A mutation, search,
test and resource site. HAMSTeRS update (version 3.0). Nucleic Acids
Res. 25: 128-132, 1997.

102. Kenwrick, S.; Gitschier, J.: A contiguous, 3-Mb physical map
of Xq28 extending from the colorblindness locus to DXS15. Am. J.
Hum. Genet. 45: 873-882, 1989.

103. Kenwrick, S.; Levinson, B.; Taylor, S.; Shapiro, A.; Gitschier,

[26644] J.: Isolation and sequence of two genes associated with a CpG island

[26645] 5-prime of the factor VIII gene. Hum. Molec. Genet. 1: 179-186,

[26646] 1992.

[26647]

[26648] 104. Kitchens, C. S.: Occult hemophilia. Johns Hopkins Med. J. 146:

[26649] 255-259, 1980.

[26650]

[26651] 105. Klein, H. G.; Aledort, L. M.; Bouma, B. N.; Hoyer, L. W.; Zimmerman,

[26652] T. S.; De Mets, D. L.: A co-operative study for the detection of

[26653] the carrier state of classic hemophilia. New Eng. J. Med. 296: 959-962,

[26654] 1977.

[26655]

[26656] 106. Kogan, S.; Gitschier, J.: Mutations and a polymorphism in the

[26657] factor VIII gene discovered by denaturing gradient gel

electrophoresis. Proc.
[26658] Nat. Acad. Sci. 87: 2092-2096, 1990.

[26659]

[26660] 107. Kogan, S. C.; Doherty, M.; Gitschier, J.: An improved method
[26661] for prenatal diagnosis of genetic diseases by analysis of amplified
[26662] DNA sequences: application to hemophilia A. New Eng. J. Med. 317:
[26663] 985-990, 1987.

[26664]

[26665] 108. Krepelova, A.; Varlova, Z.; Zavadil, J.; Brdicka, R.: Factor
[26666] VIII gene deletions in haemophilia A patients in Czechoslovakia. Brit.
[26667] J. Haemat. 86: 271-276, 1992.

[26668]

[26669] 109. Lacroix-Desmazes, S.; Bayry, J.; Misra, N.; Horn, M. P.; Villard,
[26670] S.; Pashov, A.; Stieltjes, N.; d'Oiron, R.; Saint-Remy, J.-M.; Hoebeke,

[26671] J.; Kazatchkine, M. D.; Kaveri, S. V.: The prevalence of pro-
teolytic
[26672] antibodies against factor VIII in hemophilia A. New Eng. J.
Med. 346:
[26673] 662-667, 2002.

[26674]

[26675] 110. Lakich, D.; Kazazian, H. H., Jr.; Antonarakis, S. E.;
Gitschier,
[26676] J.: Inversions disrupting the factor VIII gene are a common
cause
[26677] of severe haemophilia A. Nature Genet. 5: 236-241, 1993.

[26678]

[26679] 111. Lavergne, J. M.; Bahnak, B. R.; Vidaud, M.; Laurian, Y.;
Meyer,
[26680] D.: A directed search for mutations in hemophilia A using
restriction
[26681] enzyme analysis and denaturing gradient gel elec-
trophoresis: a study
[26682] of seven exons in the factor VIII gene of 170 cases. Nouv.
Rev. Fr.
[26683] Hematol. 34: 85-91, 1992.

[26684]

[26685] 112. Lawn, R. M.: The molecular genetics of hemophilia: blood clotting

[26686] factors VIII and IX. Cell 42: 405-406, 1985.

[26687]

[26688] 113. Leuer, M.; Oldenburg, J.; Lavergne, J.-M.; Ludwig, M.; Fregin,

[26689] A.; Eigel, A.; Ljung, R.; Goodeve, A.; Peake, I.; Olek, K.: So-matic

[26690] mosaicism in hemophilia A: a fairly common event. Am. J. Hum. Genet. 69:

[26691] 75-87, 2001.

[26692]

[26693] 114. Levin, J.: Personal Communication. Baltimore, Md. 12/1979.

[26694]

[26695] 115. Levinson, B.; Janco, R.; Phillips, J., III; Gitschier, J.: A

[26696] novel missense mutation in the factor VIII gene identified by analysis

[26697] of amplified hemophilia DNA sequences. Nucleic Acids Res. 15: 9797-9805,

[26698] 1987.

[26699]

[26700] 116. Levinson, B.; Kenwrick, S.; Lakich, D.; Hammonds, G.; Gitschier,

[26701] J.: A transcribed gene in an intron of the human factor VIII gene. Genomics 7:

[26702] 1-11, 1990.

[26703]

[26704] 117. Levinson, B.; Lehesjoki, A.-E.; de la Chapelle, A.; Gitschier,

[26705] J.: Molecular analysis of hemophilia A mutations in the Finnish population. Am.

[26706] J. Hum. Genet. 46: 53-62, 1990.

[26707]

[26708] 118. Lewis, J. H.; Bontempo, F. A.; Spero, J. A.; Ragni, M. V.; Starzl,

[26709] T. E.: Liver transplantation in a hemophiliac. New Eng. J. Med. 312:

[26710] 1189-1190, 1985.

[26711]

[26712] 119. Lillicrap, D. P.; Taylor, S. A. M.; Grover, H.; Teitel, J.; Giles,
[26713] A. R.; Holden, J. J. A.; White, B. N.: Genetic analysis in hemophilia
[26714] A: identification of a large F.VIII gene deletion in a patient with
[26715] high titre antibodies to human and porcine F.VIII. (Abstract) Blood 68:
[26716] 337a only, 1986.

[26717]

[26718] 120. Lin, S.-W.; Lin, S.-R.; Shen, M.-C.: Characterization of genetic
[26719] defects of hemophilia A in patients of Chinese origin. Genomics 18:
[26720] 496-504, 1993.

[26721]

[26722] 121. Mannucci, P. M.; Tuddenham, E. G. D.: The hemophilias--from
[26723] royal genes to gene therapy. New Eng. J. Med. 344: 1773-1779, 2001.

[26724]

[26725] 122. Marchesi, S. L.; Shulman, N. R.; Gralnick, H. R.: Studies on
[26726] the purification and characterization of human factor VIII. J. Clin.
[26727] Invest. 51: 2151-2161, 1972.

[26728]

[26729] 123. McGinniss, M. J.; Kazazian, H. H., Jr.; Hoyer, L. W.; Bi, L.;
[26730] Inaba, H.; Antonarakis, S. E.: Spectrum of mutations in CRM-positive
[26731] and CRM-reduced hemophilia A. Genomics 15: 392-398, 1993.

[26732]

[26733] 124. McKusick, V. A.: The royal hemophilia. Sci. Am. 213(2): 88-95,
[26734] 1965.

[26735]

[26736] 125. McKusick, V. A.: On the X Chromosome of Man. Washington, D.
[26737] C.: Am. Inst. Biol. Sci. (pub.) 1964.

[26738]

[26739] 126. McKusick, V. A.: Hemophilia in early New England: a follow-up
[26740] of four kindreds in which hemophilia occurred in pre-Revolutionary
[26741] period. J. Hist. Med. 17: 342-365, 1962.

[26742]

[26743] 127. McKusick, V. A.: The earliest record of hemophilia in America? Blood 19:
[26744] 243-244, 1962.

[26745]

[26746] 128. Mibashan, R. S.; Peake, I. R.; Rodeck, C. H.; Thumpston, J. K.;
[26747] Furlong, R. A.; Gorer, R.; Bains, L.; Bloom, A. L.: Dual diagnosis
[26748] of prenatal haemophilia A by measurement of fetal factor VIIIC and
[26749] VIIIC antigen (VIIICAg). Lancet II: 994-997, 1980.

[26750]

[26751] 129. Migeon, B. R.; Axelman, J.; Jan de Beur, S.; Valle, D.; Mitchell,
[26752] G. A.; Rosenbaum, K. N.: Selection against lethal alleles in

females

[26753] heterozygous for incontinentia pigmenti. Am. J. Hum. Genet. 44:

[26754] 100-106, 1989.

[26755]

[26756] 130. Migeon, B. R.; McGinniss, M. J.; Antonarakis, S. E.; Axelman,

[26757] J.; Stasiowski, B. A.; Youssoufian, H.; Kearns, W. G.; Chung, A.;

[26758] Pearson, P. L.; Kazazian, H. H., Jr.; Muneer, R. S.: Severe hemophilia

[26759] A in a female by cryptic translocation: order and orientation of factor

[26760] VIII within Xq28. Genomics 16: 20-25, 1993.

[26761]

[26762] 131. Mikami, S.: Gene analysis in haemophilia A--restriction fragment

[26763] length polymorphism and molecular defects in the factor VIII gene. Acta

[26764] Haemat. Jpn. 51: 1680-1688, 1988.

[26765]

[26766] 132. Millar, D. S.; Steinbrecher, R. A.; Wieland, K.; Grundy, C. B.;
[26767] Martinowitz, U.; Krawczak, M.; Zoll, B.; Whitmore, D.; Stephenson,
[26768] J.; Mibashan, R. S.; Kakkar, V. V.; Cooper, D. N.: The molecular
[26769] genetic analysis of haemophilia A; characterization of six partial
[26770] deletions in the factor VIII gene. Hum. Genet. 86: 219-227, 1990.
[26771]

[26772] 133. Millar, D. S.; Steinbrecher, R. A.; Wieland, K.; Grundy, C. B.;
[26773] Martinowitz, U.; Krawczak, M.; Zoll, B.; Whitmore, D.; Stephenson,
[26774] J.; Mibashan, R. S.; Kakkar, V. V.; Cooper, D. N.: The molecular
[26775] genetic analysis of haemophilia A: characterization of six partial
[26776] deletions in the factor VIII gene. Hum. Genet. 86: 219-227, 1990.
[26777]

[26778] 134. Millar, D. S.; Zoll, B.; Martinowitz, U.; Kakkar, V. V.; Cooper,
[26779] D. N.: The molecular genetics of haemophilia A: screening for point
[26780] mutations in the factor VIII gene using the restriction enzyme TaqI. Hum.
[26781] Genet. 87: 607-612, 1991.

[26782]

[26783] 135. Mori, P. G.; Pasino, M.; Vadala, C. R.; Bisogni, M. C.; Tonini,
[26784] G. P.; Scarabicchi, S.: Haemophilia 'A' in a 46,X,i(Xq) female. Brit.
[26785] J. Haemat. 43: 143-147, 1979.

[26786]

[26787] 136. Muneer, R. S.; Coffman, M. A.; Thompson, L. M.; Sexauer, C. L.;
[26788] Rennert, O. M.: Classic hemophilia in a female with X/17 complex
[26789] translocation and partial deletion of the long arm X chromosome (Xq11-13). Am.
[26790] J. Hum. Genet. 39: A126, 1986.

[26791]

[26792] 137. Murru, S.; Casula, L.; Pecorara, M.; Mori, P.; Cao, A.; Pirastu,

[26793] M.: Illegitimate recombination produced a duplication within the

[26794] FVIII gene in a patient with mild hemophilia A. Genomics 7: 115-118,

[26795] 1990.

[26796]

[26797] 138. Myers, R. M.; Fischer, S. G.; Lerman, L. S.; Maniatis, T.: Nearly

[26798] all single base substitutions in DNA fragments joined to a GC-clamp

[26799] can be detected by denaturing gradient gel electrophore-sis. Nucleic

[26800] Acids Res. 13: 3131-3145, 1985.

[26801]

[26802] 139. Myers, R. M.; Fischer, S. G.; Maniatis, T.; Lerman, L. S.: Modification

[26803] of the melting properties of duplex DNA by attachment of a GC-rich

[26804] DNA sequence as determined by denaturing gradient gel electrophoresis. Nucleic
[26805] Acids Res. 13: 3111-3129, 1985.

[26806]

[26807] 140. Nafa, K.; Bundis, M.; Deburgrave, N.; Bardin, J.-M.; Sultan,
[26808] Y.; Kaplan, J.-C.; Delpech, M.: A novel mutation (Arg-to-Leu in exon
[26809] 18) in factor VIII gene responsible for moderate hemophilia A. Hum.
[26810] Mutat. 1: 77-78, 1992.

[26811]

[26812] 141. Nafa, K.; Meriane, F.; Reghis, A.; Benabadji, M.; De-menais, F.;
[26813] Guilloud-Bataille, M.; Sultan, Y.; Kaplan, J. C.; Delpech, M.: Investigation
[26814] of factor VIIIC gene restriction fragment length polymor-phisms and
[26815] search for deletions in hemophiliac subjects in Algeria. Hum. Genet. 84:
[26816] 401-405, 1990.

[26817]

[26818] 142. Naylor, J.; Brinke, A.; Hassock, S.; Green, P. M.; Gian-
nelli,

[26819] F.: Characteristic mRNA abnormality found in half the pa-
tients with

[26820] severe haemophilia A is due to large DNA inversions.
Hum. Molec.

[26821] Genet. 2: 1773-1778, 1993.

[26822]

[26823] 143. Naylor, J. A.; Green, P. M.; Montandon, A. J.; Rizza, C.
R.;

[26824] Giannelli, F.: Detection of three novel mutations in two
haemophilia

[26825] A patients by rapid screening of whole essential region of
factor

[26826] VIII gene. Lancet 337: 635-639, 1991.

[26827]

[26828] 144. Naylor, J. A.; Green, P. M.; Rizza, C. R.; Giannelli, F.:
Factor

[26829] VIII gene explains all cases of haemophilia A. Lancet 340:
1066-1067,

[26830] 1992.

[26831]

[26832] 145. Naylor, J. A.; Green, P. M.; Rizza, C. R.; Giannelli, F.: Analysis

[26833] of factor VIII mRNA reveals defects in everyone of 28 haemophilia

[26834] A patients. Hum. Molec. Genet. 2: 11-17, 1993.

[26835]

[26836] 146. Nilsson, I. M.; Berntorp, E.; Zettervall, O.: Induction of immune

[26837] tolerance in patients with hemophilia and antibodies to factor VIII

[26838] by combined treatment with intravenous IgG, cyclophos-phamide, and

[26839] factor VIII. New Eng. J. Med. 318: 947-950, 1988.

[26840]

[26841] 147. Nilsson, I. M.; Blomback, M.; Ramgren, O.: Investiga-tions on

[26842] hemophilia A and B carriers. Bibl. Haemat. 26: 26-29, 1966.

[26843]

[26844] 148. Nilsson, I. M.; Blomback, M.; Ramgren, O.; Von Francken, I.:
[26845] Haemophilia in Sweden. II. Carriers of haemophilia A and B. Acta
[26846] Med. Scand. 171: 223-235, 1962.

[26847]

[26848] 149. Nilsson, I. M.; Blomback, M.; Von Francken, I.: On an inherited
[26849] autosomal hemorrhagic diathesis with antihemophilic globulin (AHG)
[26850] deficiency and prolonged bleeding time. Acta Med. Scand. 159: 35-57,
[26851] 1957.

[26852]

[26853] 150. Nilsson, I. M.; Lamme, S.: On acquired hemophilia A: a survey
[26854] of 11 cases. Acta Med. Scand. 208: 5-12, 1980.

[26855]

[26856] 151. Nisen, P. D.; Waber, P. G.: Nonrandom X chromo-some DNA methylation
[26857] patterns in hemophiliac females. J. Clin. Invest. 83:

1400-1403,
[26858] 1989.

[26859]

[26860] 152. Norman, J. C.; Covelli, V. H.; Sise, H. S.: Transplanta-
tion
[26861] of the spleen. (Editorial) Ann. Intern. Med. 78: 700-704,
1968.
[26862]

[26863] 153. O'Brien, D. P.; Pattinson, J. K.; Tuddenham, E. G. D.:
Purification
[26864] and characterization of factor VIII 372-cys: a hypofunc-
tional cofactor
[26865] from a patient with moderately severe hemophilia A. Blood
75: 1664-1672,
[26866] 1990.

[26867]

[26868] 154. Oberle, I.; Camerino, G.; Heilig, R.; Grunebaum, L.;
Cazenave,
[26869] J.-P.; Crapanzano, C.; Mannucci, P. M.; Mandel, J.-L.: Ge-
netic screening
[26870] for hemophilia A (classic hemophilia) with a polymorphic

DNA probe. New
[26871] Eng. J. Med. 312: 682-686, 1985.

[26872]

[26873] 155. Oberle, I.; Drayna, D.; Camerino, G.; White, R.; Man-
del, J.-L.
[26874] : The telomeric region of the human X chromosome long
arm: presence
[26875] of a highly polymorphic DNA marker and analysis of re-
combination frequency. Proc.
[26876] Nat. Acad. Sci. 82: 2824-2828, 1985.

[26877]

[26878] 156. Oldenburg, J.; Rost, S.; El-Maarri, O.; Leuer, M.; Olek,
K.;
[26879] Muller, C. R.; Schwaab, R.: De novo factor VIII gene intron
22 inversion
[26880] in a female carrier presents as a somatic mosaicism. Blood
96: 2905-2906,
[26881] 2000.

[26882]

[26883] 157. Pasi, K. J.: Gene therapy for haemophilia. Brit. J.
Haemat. 115:

[26884] 744-757, 2001.

[26885]

[26886] 158. Patterson, M.; Gitschier, J.; Bloomfield, J.; Bell, M.; Dorkins,

[26887] H.; Froster-Iskenius, U.; Sommer, S.; Sobell, J.; Schaid, D.; Thibodeau,

[26888] S.; Davies, K. E.: An intronic region within the human factor VIII

[26889] gene is duplicated within Xq28 and is homologous to the polymorphic

[26890] locus DXS115 (767). Am. J. Hum. Genet. 44: 679-685, 1989.

[26891]

[26892] 159. Patterson, M.; Schwartz, C.; Bell, M.; Sauer, S.; Hofker, M.;

[26893] Trask, B.; van den Engh, G.; Davies, K. E.: Physical mapping studies

[26894] on the human X chromosome in the region Xq27-Xqter. Genomics 1:

[26895] 297-306, 1987.

[26896]

[26897] 160. Pattinson, J. K.; Millar, D. S.; McVey, J. H.; Grundy, C. B.;
[26898] Wieland, K.; Mibashan, R. S.; Martinowitz, U.; Tan-Un, K.; Vidaud,
[26899] M.; Goossens, M.; Sampietro, M.; Mannucci, P. M.; and 17 others
[26900] : The molecular genetic analysis of hemophilia A: a di-rected search
[26901] strategy for the detection of point mutations in the human factor
[26902] VIII gene. Blood 76: 2242-2248, 1990.

[26903]

[26904] 161. Paynton, C.; Sarkar, G.; Sommer, S. S.: Identification of mutations
[26905] in two families with sporadic hemophilia A. Hum. Genet. 87: 397-400,
[26906] 1991.

[26907]

[26908] 162. Peake, I. R.; Lillicrap, D. P.; Liddell, M. B.; Matthews, R.
[26909] J.; Bloom, A. L.: Linked and intragenic probes for haemophilia A. Lancet II:

[26910] 1003-1004, 1985.

[26911]

[26912] 163. Pecorara, M.; Casarino, L.; Mori, P. G.; Morfini, M.; Mancuso,

[26913] G.; Scrivano, A. M.; Boeri, E.; Molinari, A. C.; De Biasi, R.; Ciavarella,

[26914] N.; Bencivelli, F.; Repa, T.; Barbujani, G.; Loi, A.; Perseu, L.;

[26915] Cao, A.; Pirastu, M.: Hemophilia A: carrier detection and prenatal

[26916] diagnosis by DNA analysis. Blood 70: 531-535, 1987.

[26917]

[26918] 164. Pieneman, W. C.; Reitsma, P. H.; Briet, E.: Double strand conformation

[26919] polymorphism (DSCP) detects two point mutations at codon 280 (AAC-ATC)

[26920] and at codon 431 (TAC-AAC) of the blood coagulation factor VIII gene. Thromb.

[26921] Haemost. 69: 473-475, 1993.

[26922]

[26923] 165. Pignone, A.; Matucci-Cerinic, M.; Morfini, M.; Lombardi, A.;

[26924] Rossi Ferrini, P. L.; Cagnoni, M.: Suppression of autoanti-
bodies
[26925] to factor VIII and correction of factor VIII deficiency with a
combined
[26926] steroid-cyclophosphamide-porcine factor VIII treatment in
a patient
[26927] with rheumatoid arthritis. J. Intern. Med. 231: 617-619,
1992.
[26928]

[26929] 166. Pola, V.; Svojitka, J.: Klassische Haemophilie bei
Frauen. Folia
[26930] Haemat. 75: 43-51, 1957.

[26931]

[26932] 167. Poustka, A.; Dietrich, A.; Langenstein, G.; Toniolo, D.;
Warren,
[26933] S. T.; Lehrach, H.: Physical map of human Xq27-qter: lo-
calizing the
[26934] region of the fragile X mutation. Proc. Nat. Acad. Sci. 88:
8302-8306,
[26935] 1991.

[26936]

[26937] 168. Pratt, K. P.; Shen, B. W.; Takeshima, K.; Davie, E. W.; Fujikawa,
[26938] K.; Stoddard, B. L.: Structure of the C2 domain of human factor VIII
[26939] at 1.5 angstrom resolution. Nature 402: 439-442, 1999.

[26940]

[26941] 169. Rapaport, S. I.; Patch, M. J.; Moore, F. J.: Anti-hemophilic
[26942] globulin levels in carriers of hemophilia A. J. Clin. Invest. 39:
[26943] 1619-1625, 1960.

[26944]

[26945] 170. Ratnoff, O. D.: Antihemophilic factor (factor VIII). Ann. Intern.
[26946] Med. 88: 403-409, 1978.

[26947]

[26948] 171. Ratnoff, O. D.; Bennett, B.: The genetics of hereditary disorders
[26949] of blood coagulation. Science 179: 1291-1298, 1973.

[26950]

[26951] 172. Ratnoff, O. D.; Lewis, J. H.: Heckathorn's disease: variable
[26952] functional deficiency of antihemophilic factor (factor VIII). Blood 46:
[26953] 161-173, 1975.

[26954]

[26955] 173. Reiner, A. P.; Stray, S. M.; Thompson, A. R.: Three missense
[26956] mutations in Arg codons of the factor VIII gene of mild to moderately
[26957] severe hemophilia A patients. Thromb. Res. 66: 93-99, 1992.

[26958]

[26959] 174. Reiner, A. P.; Thompson, A. R.: Screening for non-sense mutations
[26960] in patients with severe hemophilia A can provide rapid, direct carrier
[26961] detection. Hum. Genet. 89: 88-94, 1992.

[26962]

[26963] 175. Richardson, D. W.; Robinson, A. G.: Desmopressin. Ann. Intern.

[26964] Med. 103: 228-239, 1985.

[26965]

[26966] 176. Roberts, D. F.: The genetic basis of variation in factor VIII

[26967] levels among haemophiliacs. J. Med. Genet. 8: 136-139, 1971.

[26968]

[26969] 177. Robertson, J. H.; Trueman, R. G.: Combined hemophilia and Christmas

[26970] disease. Blood 24: 281-288, 1964.

[26971]

[26972] 178. Rosendaal, F. R.; Briet, E.; Stibbe, J.; van Herpen, G.; Gevers

[26973] Leuven, J. A.; Hofman, A.; Vandenbroucke, J. P.: Haemophilia protects

[26974] against ischaemic heart disease: a study of risk factors. Brit. J.

[26975] Haemat. 75: 525-530, 1990.

[26976]

[26977] 179. Rosendaal, F. R.; Brocker-Vriends, A. H. J. T.; van Houwelingen,

[26978] J. C.; Smit, C.; Varekamp, I.; van Dijck, H.; Suurmeijer, T. P. B.

[26979] M.; Vandenbroucke, J. P.; Briet, E.: Sex ratio of the muta-tion frequencies

[26980] in haemophilia A: estimation and meta-analysis. Hum. Genet. 86:

[26981] 139-146, 1990.

[26982]

[26983] 180. Rossiter, J. P.; Young, M.; Kimberland, M. L.; Hutter, P.; Ketterling,

[26984] R. P.; Gitschier, J.; Horst, J.; Morris, M. A.; Schaid, D. J.; de

[26985] Moerloose, P.; Sommer, S. S.; Kazazian, H. H., Jr.; An-tonarakis, S.

[26986] E.: Factor VIII gene inversions causing severe hemophilia A originate

[26987] almost exclusively in male germ cells. Hum. Molec. Genet. 3: 1035-1039,

[26988] 1994.

[26989]

[26990] 181. Roth, D. A.; Tawa, N. E., Jr.; O'Brien, J. M.; Treco, D. A.;

[26991] Selden, R. F.: Nonviral transfer of the gene encoding coag-
ulation
[26992] factor VIII in patients with severe hemophilia A. New Eng.
J. Med. 344:
[26993] 1735-1742, 2001.

[26994]

[26995] 182. Saiki, R. K.; Scharf, S.; Faloona, F.; Mullis, K. B.; Horn,
G.
[26996] T.; Erlich, H. A.; Arnheim, N.: Enzymatic amplification of
beta-globin
[26997] genomic sequences and restriction site analysis for diag-
nosis of sickle
[26998] cell anemia. Science 230: 1350-1354, 1985.

[26999]

[27000] 183. Samama, M.; Perrotez, C.; Houissa, R.; Hafsia, A.;
Seger, J.
[27001] : Hemophilic A feminine avec deletion d'une partie du bras
long d'un
[27002] chromosome X. Path. Biol. (Paris) 25 (suppl.): 10-17,
1977.

[27003]

[27004] 184. Schiffman, S.; Rapaport, S. I.: Increased factor VIII levels

[27005] in suspected carriers of hemophilia A: taking contraceptives by mouth. New

[27006] Eng. J. Med. 275: 599, 1966.

[27007]

[27008] 185. Schwaab, R.; Brackmann, H.-H.; Meyer, C.; Seehafer, J.; Kirchgesser,

[27009] M.; Haack, A.; Olek, K.; Tuddenham, E. G. D.; Oldenburg, J.: Haemophilia

[27010] A: mutation type determines risk of inhibitor formation. Thromb.

[27011] Haemat. 74: 1402-1406, 1995.

[27012]

[27013] 186. Schwaab, R.; Oldenburg, J.; Tuddenham, E. G. D.; Brackmann, H.

[27014] H.; Olek, K.: Mutations in haemophilia A. Brit. J. Haemat. 83:

[27015] 450-458, 1993.

[27016]

[27017] 187. Schwartz, R. S.; Abildgaard, C. F.; Aledort, L. M.;

Arkin, S.;

[27018] Bloom, A. L.; Brackmann, H. H.; Brettler, D. B.; Fukui, H.; Hilgartner,

[27019] M. W.; Inwood, M. J.; Kasper, C. K.; Kernoff, P. B. A.; Levine, P.

[27020] H.; Lusher, J. M.; Mannucci, P. M.; Scharrer, I.; MacKenzie, M. A.;

[27021] Pancham, N.; Kuo, H. S.; Allred, R. U.; the Recombinant Factor VIII

[27022] Study Group: Human recombinant DNA-derived antihe-mophilic factor

[27023] (factor VIII) in the treatment of hemophilia A. New Eng. J. Med. 323:

[27024] 1800-1805, 1990.

[27025]

[27026] 188. Seligsohn, U.; Zivelin, A.; Perez, C.; Modan, M. A.: Detection

[27027] of hemophilia A carriers by replicate factor VIII activity and factor

[27028] VIII antigenicity determinations. Brit. J. Haemat. 42: 433-439,

[27029] 1979.

[27030]

[27031] 189. Shima, M.; Ware, J.; Yoshioka, A.; Fukui, H.; Fulcher, C. A.

[27032] : An arginine to cysteine amino acid substitution at a critical thrombin

[27033] cleavage site in a dysfunctional factor VIII molecule. Blood 74:

[27034] 1612-1617, 1989.

[27035]

[27036] 190. Sie, P.; Caranobe, C.; Benalioua, M.; Boneu, B.: Homozygous

[27037] hemophilia A in a female. (Letter) Thromb. Haemost. 54: 728 only,

[27038] 1985.

[27039]

[27040] 191. Smith, C. A. B.: Personal Communication. London 1968.

[27041]

[27042] 192. Soucie, J. M.; Evatt, B.; Jackson, D.; Hemophilia Surveillance

[27043] System Project Investigators: Occurrence of hemophilia in

the United
[27044] States. Am. J. Hemat. 59: 288-294, 1998.

[27045]

[27046] 193. Stevens, R. F.: The history of haemophilia in the royal families
[27047] of Europe. Brit. J. Haemat. 105: 25-32, 1999.

[27048]

[27049] 194. Stites, D. P.; Hershgold, E. J.; Perlman, J. D.; Fudenberg, H.
[27050] H.: Factor VIII detection by hemagglutination inhibition: hemophilia
[27051] A and von Willebrand's disease. Science 171: 196-197, 1971.

[27052]

[27053] 195. Stoneking, M.; Melton, T.; Nott, J.; Barritt, S.; Roby, R.; Holland,
[27054] M.; Weedn, V.; Gill, P.; Kimpton, C.; Aliston-Greiner, R.; Sullivan,
[27055] K.: Establishing the identity of Anna Anderson Manahan. (Letter) Nature
[27056] Genet. 9: 9-10, 1995.

[27057]

[27058] 196. Tantravahi, U.; Murty, V. V. V. S.; Jhanwar, S. C.; Toole, J.

[27059] J.; Woozney, J. M.; Chaganti, R. S. K.; Latt, S. A.: Physical mapping

[27060] of the factor VIII gene proximal to two polymorphic DNA probes in

[27061] human chromosome band Xq28: implications for factor VIII gene segregation

[27062] analysis. Cytogenet. Cell Genet. 42: 75-79, 1986.

[27063]

[27064] 197. Toole, J. J.; Knopf, J. L.; Wozney, J. M.; Sultzman, L. A.; Buecker,

[27065] J. L.; Pittman, D. D.; Kaufman, R. J.; Brown, E.; Shoemaker, C.; Orr,

[27066] E. C.; Amphlett, G. W.; Foster, W. B.; Coe, M. L.; Knutson, G. J.;

[27067] Fass, D. N.; Hewick, R. M.: Molecular cloning of a cDNA encoding

[27068] human antihaemophilic factor. Nature 312: 342-347, 1984.

[27069]

[27070] 198. Traystman, M. D.; Higuchi, M.; Kasper, C. K.; An-
tonarakis, S.
[27071] E.; Kazazian, H. H., Jr.: Use of denaturing gradient gel
electrophoresis
[27072] to detect point mutations in the factor VIII gene. Genomics
6: 293-301,
[27073] 1990.

[27074]

[27075] 199. Tuddenham, E. G. D.; Lazarchick, J.; Hoyer, L. W.:
Synthesis
[27076] and release of factor VIII by cultured human endothelial
cells. Brit.
[27077] J. Haemat. 47: 617-626, 1981.

[27078]

[27079] 200. VandenDriessche, T.; Vanslembrouck, V.; Goovaerts,
I.; Zwinnen,
[27080] H.; Vanderhaeghen, M.-L.; Collen, D.; Chuah, M. K. L.:
Long-term
[27081] expression of human coagulation factor VIII and correc-
tion of hemophilia
[27082] A after in vivo retroviral gene transfer in factor VIII-
deficient

[27083] mice. Proc. Nat. Acad. Sci. 96: 10379-10384, 1999.

[27084]

[27085] 201. Vidaud, D.; Vidaud, M.; Plassa, F.; Gazengel, C.; Noel, B.; Goossens,

[27086] M.: Father-to-son transmission of hemophilia A due to uniparental

[27087] disomy. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A226 only, 1989.

[27088]

[27089] 202. Vogel, F.: A probable sex difference in some mutation rates. Am.

[27090] J. Hum. Genet. 29: 312-319, 1977.

[27091]

[27092] 203. Wacey, A. I.; Kemball-Cook, G.; Kazazian, H. H.; Antonarakis,

[27093] S. E.; Schwaab, R.; Lindley, P.; Tuddenham, E. G. D.: The haemophilia

[27094] A mutation search test and resource site, home page of the factor

[27095] VIII mutation database: HAMSTeRS. Nucleic Acids Res. 24: 100-102,

[27096] 1996.

[27097]

[27098] 204. Wehnert, M.; Herrmann, F. H.; Wulff, K.: Partial deletions of

[27099] factor VIII gene as molecular diagnostic markers in hemophilia A. Dis.

[27100] Markers 7: 113-117, 1989.

[27101]

[27102] 205. Windsor, S.; Lyng, A.; Taylor, S. A. M.; Ewenstein, B. M.; Neufeld,

[27103] E. J.; Lillicrap, D.: Severe haemophilia A in a female resulting

[27104] from two de novo factor VIII mutations. Brit. J. Haemat. 90: 906-909,

[27105] 1995.

[27106]

[27107] 206. Winter, R. M.; Tuddenham, E. G. D.; Goldman, E.; Matthews, K.

[27108] B.: A maximum likelihood estimate of the sex ratio of mutation rates

[27109] in haemophilia A. Hum. Genet. 64: 156-159, 1983.

[27110]

[27111] 207. Wood, W. I.; Capon, D. J.; Simonsen, C. C.; Eaton, D. L.; Gitschier,

[27112] J.; Keyt, B.; Seeburg, P. H.; Smith, D. H.; Hollingshead, P.; Wion,

[27113] K. L.; Delwart, E.; Tuddenham, E. G. D.; Vehar, G. A.; Lawn, R. M.

[27114] : Expression of active human factor VIII from recombinant DNA clones. Nature 312:

[27115] 330-337, 1984.

[27116]

[27117] 208. Woodliff, H. J.; Jackson, J. M.: Combined haemophilia and Christmas

[27118] disease: a genetic study of a patient and his relatives. Med. J.

[27119] Aust. 53: 658-661, 1966.

[27120]

[27121] 209. Woods-Samuels, P.; Wong, C.; Mathias, S. L.; Scott, A. F.; Kazazian,

[27122] H. H., Jr.; Antonarakis, S. E.: Characterization of a non-deleterious

[27123] L1 insertion in an intron of the human factor VIII gene and further
[27124] evidence of open reading frames in functional L1 elements. Genomics 4:
[27125] 290-296, 1989.

[27126]

[27127] 210. Woolf, L. I.: Gene expression in heterozygotes. Nature 194:
[27128] 609-610, 1962.

[27129]

[27130] 211. Young, M.; Inaba, H.; Hoyer, L. W.; Higuchi, M.; Kazazian, H.
[27131] H., Jr.; Antonarakis, S. E.: Partial correction of a severe molecular
[27132] defect in hemophilia A, because of errors during expression of the
[27133] factor VIII gene. Am. J. Hum. Genet. 60: 565-573, 1997.

[27134]

[27135] 212. Youssoufian, H.; Antonarakis, S. E.; Aronis, S.; Tsiftis, G.;
[27136] Phillips, D. G.; Kazazian, H. H., Jr.: Characterization of five

partial
[27137] deletions of the factor VIII gene. Proc. Nat. Acad. Sci. 84:
3772-3776,
[27138] 1987.

[27139]

[27140] 213. Youssoufian, H.; Antonarakis, S. E.; Bell, W.; Griffin,
A. M.;
[27141] Kazazian, H. H., Jr.: Nonsense and missense mutations in
hemophilia
[27142] A: estimate of the relative mutation rate at CG dinu-
cleotides. Am.
[27143] J. Hum. Genet. 42: 718-725, 1988.

[27144]

[27145] 214. Youssoufian, H.; Antonarakis, S. E.; Kasper, C. K.;
Phillips,
[27146] D. G.; Kazazian, H. H., Jr.: The spectrum and origin of mu-
tations
[27147] in hemophilia A. (Abstract) Am. J. Hum. Genet. 41: A249
only, 1987.

[27148]

[27149] 215. Youssoufian, H.; Kasper, C. K.; Phillips, D. G.;

Kazazian, H.

[27150] H., Jr.; Antonarakis, S. E.: Restriction endonuclease mapping of

[27151] six novel deletions of the factor VIII gene in hemophilia A. Hum.

[27152] Genet. 80: 143-148, 1988.

[27153]

[27154] 216. Youssoufian, H.; Kazazian, H. H., Jr.; Phillips, D. G.; Aronis,

[27155] S.; Tsiftis, G.; Brown, V. A.; Antonarakis, S. E.: Recurrent mutations

[27156] in haemophilia A give evidence for CpG mutation hotspots. Nature 324:

[27157] 380-382, 1986.

[27158]

[27159] 217. Youssoufian, H.; Wong, C.; Aronis, S.; Platokoukis, H.; Kazazian,

[27160] H. H., Jr.; Antonarakis, S. E.: Moderately severe hemophilia A resulting

[27161] from glu-to-gly substitution in exon 7 of the factor VIII gene. Am.

[27162] J. Hum. Genet. 42: 867-871, 1988.

[27163]

[27164] 218. Zacharski, L. R.; Bowie, E. J. W.; Titus, J. L.; Owen, C. A.,

[27165] Jr.: Synthesis of antihemophilic factor (factor VIII) by leukocytes:

[27166] preliminary report. Mayo Clin. Proc. 43: 617-619, 1968.

[27167]

[27168] 219. Zimmerman, T. S.; Ratnoff, O. D.; Littell, A. S.: Detection

[27169] of carriers of classic hemophilia using an immunologic assay for antihemophilic

[27170] factor (factor VIII). J. Clin. Invest. 50: 255-258, 1971.

[27171]

[27172] 220. Zimmerman, T. S.; Ratnoff, O. D.; Powell, A. E.: Immunologic

[27173] differentiation of classic hemophilia (factor VIII deficiency) and

[27174] von Willebrand's disease, with observations on combined deficiencies

[27175] of antihemophilic factor and proaccelerin (factor V) and on an acquired

[27176] circulating anticoagulant against antihemophilic factor. J. Clin.

[27177] Invest. 50: 244-245, 1971.

[27178] 3491.

[27179] Endo, F.; Katoh, H.; Matsuda, I.: Putative genetic deficiency

[27180] of 4-hydroxyphenylpyruvic acid dioxygenase in mice: a murine model

[27181] for hereditary tyrosinaemia type III. J. Inherit. Metab. Dis. 13:

[27182] 780-782, 1990.

[27183] 3492.

[27184] Endo, F.; Katoh, H.; Yamamoto, S.; Matsuda, I.: A murine model

[27185] for type III tyrosinemia: lack of immunologically detectable 4-hydroxyphenylpyruvic

[27186] acid dioxygenase enzyme protein in a novel mouse strain with hypertyrosinemia. Am.

[27187] J. Hum. Genet. 48: 704-709, 1991.

[27188] 3493.

[27189] Endo, F.; Kitano, A.; Uehara, I.; Nagata, N.; Matsuda, I.; Shinka,

[27190] T.; Kuhara, T.; Matsumoto, I.: 4-Hydroxyphenylpyruvic acid oxidase
[27191] deficiency with normal fumarylacetoacetase: a new variant form of
[27192] hereditary hypertyrosinemia. Pediat. Res. 17: 92-96, 1983.
[27193] 3494.

[27194] Giardini, O.; Cantani, A.; Kennaway, N. G.; D'Eufemia, P.: Chronic
[27195] tyrosinemia associated with 4-hydroxyphenylpyruvate dioxygenase deficiency
[27196] with acute intermittent ataxia and without visceral and bone involvement. Pediat.
[27197] Res. 17: 25-29, 1983.

[27198] 3495.

[27199] Ruetschi, U.; Cerone, R.; Perez-Cerda, C.; Schiaffino, M. C.; Standing,
[27200] S.; Ugarte, M.; Holme, E.: Mutations in the 4-hydroxyphenylpyruvate
[27201] dioxygenase gene (HPD) in patients with tyrosinemia type III. Hum.
[27202] Genet. 106: 654-662, 2000.

[27203] 3496.

[27204] Ruetschi, U.; Dellsen, A.; Sahlin, P.; Stenman, G.; Rymo, L.; Lindstedt,

[27205] S.: Human 4-hydroxyphenylpyruvate dioxygenase: pri-mary structure

[27206] and chromosomal localization of the gene. Europ. J. Biochem. 213:

[27207] 1081-1089, 1993.

[27208] 3497.

[27209] Ruetschi, U.; Rymo, L.; Lindstedt, S.: Human 4-hydroxyphenylpyruvate

[27210] dioxygenase gene (HPD). Genomics 44: 292-299, 1997.

[27211] 3498.

[27212] Stenman, G.; Roijer, E.; Ruetschi, U.; Dellsen, A.; Rymo, L.;

[27213] Lindstedt, S.: Regional assignment of the human 4-hydroxyphenylpyruvate

[27214] dioxygenase gene (HPD) to 12q24-qter by fluorescence in situ hybridization. Cytogenet.

[27215] Cell Genet. 71: 374-376, 1995.

[27216] 3499.

[27217] Manga, P.; Kromberg, J. G. R.; Box, N. F.; Sturm, R. A.; Jenkins,
[27218] T.; Ramsay, M.: Rufous oculocutaneous albinism in southern African
[27219] blacks is caused by mutations in the TYRP1 gene. Am. J. Hum. Genet. 61:
[27220] 1095-1101, 1997.

[27221] 3500.

[27222] Cleaver, J. E.: Personal Communication. San Francisco, Calif.
[27223] 1/15/1993.

[27224] 3501.

[27225] Murrell, J. R.; Spillantini, M. G.; Zolo, P.; Guazzelli, M.; Smith,
[27226] M. J.; Hasegawa, M.; Redi, F.; Crowther, R. A.; Pietrini, P.; Ghetti,
[27227] B.; Goedert, M.: Tau gene mutation G389R causes a tauopathy with
[27228] abundant Pick body-like inclusions and axonal deposits. J. Neuropath.
[27229] Exp. Neurol. 58: 1207-1226, 1999.

[27230] 3502.

[27231] Nagase, T.; Ishikawa, K.; Kikuno, R.; Hirosawa, M.; No-
mura, N.;

[27232] Ohara, O.: Prediction of the coding sequences of unidenti-
fied human

[27233] genes. XV. The complete sequences of 100 new cDNA
clones from brain

[27234] which code for large proteins in vitro. DNA Res. 6:
337-345, 1999.

[27235] 3503.

[27236] Neumann, M.; Schulz-Schaeffer, W.; Crowther, R. A.;
Smith, M.

[27237] J.; Spillantini, M. G.; Goedert, M.; Kretzschmar, H. A.:
Pick's disease

[27238] associated with the novel tau gene mutation K369I. Ann.
Neurol. 50:

[27239] 503-513, 2001.

[27240] 3504.

[27241] Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon,

[27242] T. A.: Identification of cDNA clones for the human micro-
tubule-associated

[27243] protein tau and chromosomal localization of the genes for tau and

[27244] microtubule-associated protein 2. Molec. Brain Res. 1: 271-280,

[27245] 1986.

[27246] 3505.

[27247] Pickering-Brown, S.; Baker, M.; Yen, S.-H.; Liu, W.-K.; Hasegawa,

[27248] M.; Cairns, N.; Lantos, P. L.; Rossor, M.; Iwatsubo, T.; Davies, Y.;

[27249] Allsop, D.; Furlong, R.; Owen, F.; Hardy, J.; Mann, D.; Hut-

ton, M.

[27250] : Pick's disease is associated with mutations in the tau gene. Ann.

[27251] Neurol. 48: 859-867, 2000.

[27252] 3506.

[27253] Poorkaj, P.: Personal Communication. Seattle, Wash. 11/10/1998.

[27254] 3507.

[27255] Poorkaj, P.; Bird, T. D.; Wijsman, E.; Nemens, E.; Garruto, R.

[27256] M.; Anderson, L.; Andreadis, A.; Wiederholt, W. C.; Raskind, M.; Schellenberg,

[27257] G. D.: Tau is a candidate gene for chromosome 17 frontotemporal dementia. Ann.

[27258] Neurol. 43: 815-825, 1998.

[27259] 3508.

[27260] Poorkaj, P.; Kas, A.; D'Souza, I.; Zhou, Y.; Pham, Q.; Stone,

[27261] M.; Olson, M. V.; Schellenberg, G. D.: A genomic sequence analysis

[27262] of the mouse and human microtubule-associated protein tau. Mammalian

[27263] Genome 12: 700-712, 2001.

[27264] 3509.

[27265] Rapoport, M.; Dawson, H. N.; Binder, L. I.; Vitek, M. P.; Ferreira,

[27266] A.: Tau is essential to beta-amyloid-induced neurotoxicity. Proc.

[27267] Nat. Acad. Sci. 99: 6364-6369, 2002.

[27268] 3510.

[27269] Rizzu, P.; Van Swieten, J. C.; Joosse, M.; Hasegawa, M.; Stevens,

[27270] M.; Tibben, A.; Niermeijer, M. F.; Hillebrand, M.; Ravid, R.; Oostra,
[27271] B. A.; Goedert, M.; van Duijn, C. M.; Heutink, P.: High prevalence
[27272] of mutations in the microtubule-associated protein tau in a population
[27273] study of frontotemporal dementia in the Netherlands. Am. J. Hum.
[27274] Genet. 64: 414-421, 1999.

[27275] 3511.

[27276] Spillantini, M. G.; Murrell, J. R.; Goedert, M.; Farlow, M. R.;

[27277] Klug, A.; Ghetti, B.: Mutation in the tau gene in familial multiple
[27278] system tauopathy with presenile dementia. Proc. Nat. Acad. Sci. 95:
[27279] 7737-7741, 1998.

[27280] 3512.

[27281] Spillantini, M. G.; Yoshida, H.; Rizzini, C.; Lantos, P. L.; Khan,
[27282] N.; Rossor, M. N.; Goedert, M.; Brown, J.: A novel tau mutation (N296N)

[27283] in familial dementia with swollen achromatic neurons and corticobasal
[27284] inclusion bodies. Ann. Neurol. 48: 939-943, 2000.

[27285] 3513.

[27286] Stambolic, V.; Ruel, L.; Woodgett, J. R.: Lithium inhibits glycogen
[27287] synthase kinase-3 activity and mimics Wingless signalling in intact
[27288] cells. Curr. Biol. 6: 1664-1668, 1996. Note: Erratum: Curr. Biol.
[27289] 7: 196 only, 1997.

[27290] 3514.

[27291] Stamer, K.; Vogel, R.; Thies, E.; Mandelkow, E.; Man-
delkow, E.-M.
[27292] : Tau blocks traffic of organelles, neurofilaments, and APP vesicles
[27293] in neurons and enhances oxidative stress. J. Cell Biol. 156: 1051-1063,
[27294] 2002.

[27295] 3515.

[27296] Varani, L.; Hasegawa, M.; Spillantini, M. G.; Smith, M. J.;

Murrell,
[27297] J. R.; Ghetti, B.; Klug, A.; Goedert, M.; Varani, G.: Structure of
[27298] tau exon 10 splicing regulatory element RNA and destabilization by
[27299] mutations of frontotemporal dementia and parkinsonism linked to chromosome
[27300] 17. Proc. Nat. Acad. Sci. 96: 8229-8234, 1999.

[27301] 3516.

[27302] Verpillat, P.; Camuzat, A.; Hannequin, D.; Thomas-Anterion, C.;
[27303] Puel, M.; Belliard, S.; Dubois, B.; Didic, M.; Michel, B.-F.; Lacomblez,
[27304] L.; Moreaud, O.; Sellal, F.; Golfier, V.; Campion, D.; Clerget-Darpoux,
[27305] F.; Brice, A.: Association between the extended tau haplotype and
[27306] frontotemporal dementia. Arch. Neurol. 59: 935-939, 2002.
[27307] 3517.

[27308] Wilhelmsen, K. C.; Lynch, T.; Pavlou, E.; Higgins, M.; Hygaard,

[27309] T. G.: Localization of disinhibition-demen-
tia-parkinsonism-amyotrophy
[27310] complex to 17q21-22. Am. J. Hum. Genet. 55:
1159-1165, 1994.
[27311] 3518.

[27312] Yasuda, M.; Yokoyama, K.; Nakayasu, T.; Nishimura, Y.;
Matsui,
[27313] M.; Yokoyama, T.; Miyoshi, K.; Tanaka, C.: A Japanese pa-
tient with
[27314] frontotemporal dementia and parkinsonism by a tau
P301S mutation. Neurology 55:
[27315] 1224-1227, 2000.

[27316] 3519.

[27317] Zhukareva, V.; Mann, D.; Pickering-Brown, S.; Uryu, K.;
Shuck,
[27318] T.; Shah, K.; Grossman, M.; Miller, B. L.; Hulette, C. M.; Fe-
instein,
[27319] S. C.; Trojanowski, J. Q.; Lee, V. M.-Y.: Sporadic Pick's dis-
ease:
[27320] a tauopathy characterized by a spectrum of pathological
tau isoforms
[27321] in gray and white matter. Ann. Neurol. 51: 730-739,

2002.

[27322] 3520.

[27323] Amack, J. D.; Mahadevan, M. S.: The myotonic dystrophy expanded

[27324] CUG repeat tract is necessary but not sufficient to disrupt C2C12

[27325] myoblast differentiation. Hum. Molec. Genet. 10: 1879-1887, 2001.

[27326] 3521.

[27327] Chen, S. L.; Dowhan, D. H.; Hosking, B. M.; Muscat, G. E. O.:

[27328] The steroid receptor coactivator, GRIP-1, is necessary for MEF-2C-dependent

[27329] gene expression and skeletal muscle differentiation. Genes Dev. 14:

[27330] 1209-1228, 2000.

[27331] 3522.

[27332] Hasty, P.; Bradley, A.; Morris, J. H.; Edmondson, D. G.; Venuti,

[27333] J. M.; Olson, E. N.; Klein, W. H.: Muscle deficiency and neonatal

[27334] death in mice with a targeted mutation in the myogenin

gene. Nature 364:
[27335] 501-506, 1993.

[27336] 3523.

[27337] Nabeshima, Y.; Hanaoka, K.; Hayasaka, M.; Esumi, E.; Li, S.; Nonaka,
[27338] I.; Nabeshima, Y.-I.: Myogenin gene disruption results in perinatal
[27339] lethality because of severe muscle defect. Nature 364: 532-535,
[27340] 1993.

[27341] 3524.

[27342] Braun, T.; Bober, E.; Winter, B.; Rosenthal, N.; Arnold, H. H.
[27343] : Myf-6, a new member of the human gene family of myo-genic determination
[27344] factors: evidence for a gene cluster on chromosome 12. EMBO J. 9:
[27345] 821-831, 1990.

[27346] 3525.

[27347] Olson, E.; Edmondson, D.; Wright, W. E.; Lin, V. K.; Guenet, J.-L.;

[27348] Simon-Chazottes, D.; Thompson, L. H.; Stallings, R. L.; Schroeder,
[27349] W. T.; Duvic, M.; Brock, D.; Helin, D.; Siciliano, M. J.: Myogenin
[27350] is in an evolutionarily conserved linkage group on human chromosome
[27351] 1q31-q41 and unlinked to other mapped muscle regulatory factor genes. Genomics 8:
[27352] 427-434, 1990.

[27353] 3526.

[27354] Cupelli, L.; Renault, B.; Leblanc-Straceski, J.; Banks, A.; Ward,
[27355] D.; Kucherlapati, R. S.; Krauter, K.: Assignment of the human myogenic
[27356] factors 5 and 6 (MYF5, MYF6) gene cluster to 12q21 by in situ hybridization
[27357] and physical mapping of the locus between D12S350 and D12S106. Cytogenet.
[27358] Cell Genet. 72: 250-251, 1996.

[27359] 3527.

[27360] Kerst, B.; Mennerich, D.; Schuelke, M.; Stoltenburg-Didinger, G.;

[27361] von Moers, A.; Gossrau, R.; van Landeghem, F. K. H.; Speer, A.; Braun,
[27362] T.; Hubner, C.: Heterozygous myogenic factor 6 mutation associated
[27363] with myopathy and severe course of Becker muscular dystrophy. Neuromusc.
[27364] Disord. 10: 572-577, 2000.

[27365] 3528.

[27366] Miner, J. H.; Wold, B.: Herculin: a fourth member of the MyoD
[27367] family of myogenic regulatory genes. Proc. Nat. Acad. Sci. 87: 1089-1093,
[27368] 1990.

[27369] 3529.

[27370] Rhodes, S. J.; Konieczny, S. F.: Identification of MRF4: a new
[27371] member of the muscle regulatory factor gene family. Genes Dev. 3:
[27372] 2050-2061, 1989.

[27373] 3530.

[27374] Yun, K.; Wold, B.: Skeletal muscle determination and dif-

ferentiation:
[27375] story of a core regulatory network and its context. Curr. Opin. Cell
[27376] Biol. 8: 877-889, 1996.

[27377] 3531.

[27378] Akaboshi, E.: Cloning of the human myoglobin gene. Gene 33:
[27379] 241-249, 1985.

[27380] 3532.

[27381] Boulton, F. E.; Huntsman, R. G.; Lehmann, H.; Lorkin, P. A.; Romero-Herrera,
[27382] A. E.: Myoglobin variants. (Abstract) Biochem. J. 118: 39P only,
[27383] 1970.

[27384] 3533.

[27385] Boulton, F. E.; Huntsman, R. G.; Lorkin, P. A.; Lehmann, H.: Abnormal
[27386] human myoglobin: 53(D4) glutamic acid lysine. Nature 223: 832-833,
[27387] 1969.

[27388] 3534.

[27389] Boulton, F. E.; Huntsman, R. G.; Yawson, G. I.; Romero-Herrera,
[27390] A. E.; Lorkin, P. A.: The second variant of human myoglobin: 138(H16)
[27391] arginine to glutamine. Brit. J. Haemat. 20A: 69-74, 1971.

[27392] 3535.

[27393] Boyer, S. H.: Similar incidence and non-randomness among human
[27394] myoglobin and hemoglobin mutants in general populations: implications
[27395] for the study of myoglobin in muscle disease.In: Pathogenesis of Human
[27396] Muscular Dystrophies. Proc. Vth Int. Cong. of Muscular Dystrophy Assoc.,
[27397] Durango, Colo., June 21-25, 1976. Amsterdam: Excerpta Medica (pub.)
[27398] 1977.

[27399] 3536.

[27400] Boyer, S. H.; Fainer, D. C.; Naughton, M. A.: Myoglobin inherited
[27401] structural variation in man. Science 140: 1228-1231, 1963.

[27402] 3537.

[27403] Dozier, C.; Walbaum, S.; Leprince, D.; Stehelin, D.: EcoRI RFLP

[27404] linked to the human myb gene. Nucleic Acids Res. 14: 1928 only,

[27405] 1986.

[27406] 3538.

[27407] Garry, D. J.; Ordway, G. A.; Lorenz, J. N.; Radford, N. B.; Chin,

[27408] E. R.; Grange, R. W.; Bassel-Duby, R.; Williams, R. S.: Mice without

[27409] myoglobin. Nature 395: 905-908, 1998.

[27410] 3539.

[27411] Godecke, A.; Flogel, U.; Zanger, K.; Ding, Z.; Hirchenhain, J.;

[27412] Decking, U. K. M.; Schrader, J.: Disruption of myoglobin in mice

[27413] induces multiple compensatory mechanisms. Proc. Nat. Acad. Sci. 96:

[27414] 10495-10500, 1999.

[27415] 3540.

[27416] Jeffreys, A.; Wilson, V.; Thein, S.: Hypervariable 'minisatellite'

[27417] regions in human DNA. Science 314: 67-73, 1985.

[27418] 3541.

[27419] Jeffreys, A. J.; Wilson, V.; Blanchetot, A.; Weller, P.; Geurts

[27420] van Kessel, A.; Spurr, N.; Solomon, E.; Goodfellow, P.: The human

[27421] myoglobin gene: a third dispersed globin locus in the human genome. Nucleic

[27422] Acids Res. 12: 3235-3243, 1984.

[27423] 3542.

[27424] McKeithan, T. W.; Rowley, J. D.; Shows, T. B.; Diaz, M. O.: Cloning

[27425] of the chromosome translocation breakpoint junction of the t(14;19)

[27426] in chronic lymphocytic leukemia. Proc. Nat. Acad. Sci. 84: 9257-9260,

[27427] 1987.

[27428] 3543.

[27429] de Groot, N. G.; Otting, N.; Doxiadis, G. G. M.; Balla-Jhagjhoorsingh,

[27430] S. S.; Heeney, J. L.; van Rood, J. J.; Gagneux, P.; Bontrop, R. E.

[27431] : Evidence for an ancient selective sweep in the MHC class I gene

[27432] repertoire of chimpanzees. Proc. Nat. Acad. Sci. 99: 11748-11753,

[27433] 2002.

[27434] 3544.

[27435] Del Pozzo, G.; Guardiola, J.: Mummy DNA fragment identified.

[27436] (Letter) Nature 339: 431-432, 1989.

[27437] 3545.

[27438] Dorf, M. E.; Balner, H.; DeGroot, M. L.; Benacerraf, B.: Histocompatibility-linked

[27439] immune-response genes in the Rhesus monkey. Transplant. Proc. 6:

[27440] 119-124, 1974.

[27441] 3546.

[27442] Edwards, J. H.; Allen, F. H.; Glenn, K. P.; Lamm, L. U.; Robson,

[27443] E. B.: The linkage relationships of HL-A.In: Histocompati-

bility Testing.

[27444] Baltimore: Williams and Wilkins (pub.) 1973.

[27445] 3547.

[27446] Engelfriet, C. P.; Britten, A.: The cytotoxic test for leuco-
cyte

[27447] antibodies. A simple and reliable technique. Vox Sang. 10:
660-674,

[27448] 1965.

[27449] 3548.

[27450] Erlich, H.; Lee, J. S.; Petersen, J. W.; Bugawan, T.; DeMars,

[27451] R.: Molecular analysis of HLA class I and class II antigen
loss mutants

[27452] reveals a homozygous deletion of the DR, DQ, and part of
the DP region:

[27453] implications for class II gene order. Hum. Immun. 16:
205-219, 1986.

[27454] 3549.

[27455] Erlich, H. A.; Stetler, D.; Sheng-Dong, R.; Ness, D.;
Grumet,

[27456] C.: Segregation and mapping analysis of polymorphic HLA
class I restriction

[27457] fragments: detection of a novel fragment. Science 222: 72-74, 1983.

[27458] 3550.

[27459] Fellous, M.; Dausset, J.: Probable haploid expression of HL-A

[27460] antigens on human spermatozoan. Nature 225: 191-193, 1970.

[27461] 3551.

[27462] Figueroa, F.; Gunther, E.; Klein, J.: MHC polymorphism pre-dating

[27463] speciation. (Letter) Nature 335: 265-267, 1988.

[27464] 3552.

[27465] Francke, U.; Pellegrino, M. A.: Assignment of the major histocompatibility

[27466] complex to a region of the short arm of human chromosome 6. Proc.

[27467] Nat. Acad. Sci. 74: 5776 only, 1977.

[27468] 3553.

[27469] Francke, U.; Pellegrino, M. A.: Assignment of the major histocompatibility

[27470] complex to a region of the short arm of human chromo-

some 6. Proc.

[27471] Nat. Acad. Sci. 74: 1147-1151, 1977.

[27472] 3554.

[27473] Gill, T. J., III; Cramer, D. V.; Kunz, H. W.: The major histo-compatibility

[27474] complex--comparison in the mouse, man, and the rat: a review. Am.

[27475] J. Path. 90: 735-777, 1978.

[27476] 3555.

[27477] Gladstone, P.; Fueresz, L.; Pious, D.: Gene dosage and gene expression

[27478] in the HLA region: evidence from deletion variants. Proc. Nat. Acad.

[27479] Sci. 79: 1235-1239, 1982.

[27480] 3556.

[27481] Gluecksohn-Waelsch, S.; Erickson, R. P.: The T-locus of the mouse:

[27482] implications for mechanisms of development. Curr. Top. Dev. Biol. 5:

[27483] 281-316, 1970.

[27484] 3557.

[27485] Goodfellow, P. N.; Jones, E.; Van Heyningen, V.; Solomon, E.;

[27486] Kennett, R.; Bobrow, M.; Bodmer, W. F.: Linkage relationships of

[27487] the HL-A system and beta-2-microglobulin. Birth Defects Orig. Art.

[27488] Ser. 11(3): 162-167, 1975. Note: Alternate: Cytogenet. Cell Genet.

[27489] 14: 332-337, 1975.

[27490] 3558.

[27491] Gruen, J. R.; Nalabolu, S. R.; Chu, T. W.; Bowlus, C.; Fan, W.

[27492] F.; Goei, V. L.; Wei, H.; Sivakamasundari, R.; Liu, Y.-C.; Xu, H.

[27493] X.; Parimoo, S.; Nallur, G.; Ajioka, R.; Shukla, H.; Bray-Ward, P.;

[27494] Pan, J.; Weissman, S. M.: A transcription map of the major histocompatibility

[27495] complex (MHC) class I region. Genomics 36: 70-85, 1996.

[27496] 3559.

[27497] Hedrick, S. M.: Dawn of the hunt for nonclassical MHC function. Cell 70:

[27498] 177-180, 1992.

[27499] 3560.

[27500] Helenius, A.; Morein, B.; Fries, E.; Simons, K.; Robinson, P.;

[27501] Schirrmacher, V.; Terhorst, C.; Strominger, J. L.: Human (HLA-A and

[27502] HLA-B) and murine (H-2K and H-2D) histocompatibility antigens are

[27503] cell surface receptors for Semliki Forest virus. Proc. Nat. Acad.

[27504] Sci. 75: 3846-3850, 1978.

[27505] 3561.

[27506] Hirai, M.; Takahashi, E.; Ishida, T.; Hori, T.: Chromosomal localization

[27507] of the major histocompatibility complex (MHC) in the rhe-sus monkey

[27508] and chimpanzee by fluorescence in situ hybridization. Cy-togenet.

[27509] Cell Genet. 57: 204-205, 1991.

[27510] 3562.

[27511] Hood, L.; Steinmetz, M.; Goodenow, R.: Genes of the ma-jor histocompatibility

[27512] complex. Cell 28: 685-687, 1982.

[27513] 3563.

[27514] Jordan, B. R.; Bregegere, F.; Kourilsky, P.: Human HLA gene segment

[27515] isolated by hybridization with mouse H-2 cDNA probes. Nature 290:

[27516] 521-523, 1981.

[27517] 3564.

[27518] Kissmeyer-Nielsen, F.; Jorgensen, F.; Lamm, L. U.: The HL-A system

[27519] in clinical medicine. Johns Hopkins Med. J. 131: 385-400, 1972.

[27520] 3565.

[27521] Kissmeyer-Nielsen, F.; Svejgaard, A.; Ahrons, S.; Nielsen, L.

[27522] S.: Crossing-over within the HL-A system. Nature 224: 75-76, 1969.

[27523] 3566.

[27524] Kissmeyer-Nielsen, F.; Svejgaard, A.; Hauge, M.: Genetics of

[27525] the human HL-A transplantation system. Nature 219:

1116–1119, 1968.

[27526] 3567.

[27527] Kissmeyer-Nielsen, F.; Thorsby, E.: Human transplantation antigens. Transplant.

[27528] Rev. 4: 1–176, 1970.

[27529] 3568.

[27530] Klein, J.: The major histocompatibility complex of the mouse. Science 203:

[27531] 516-521, 1979.

[27532] 3569.

[27533] Klein, J.; Figueroa, F.: Evolution of the major histocompatibility

[27534] complex. CRC Crit. Rev. Immun. 6: 295-386, 1986.

[27535] 3570.

[27536] Kompf, J.; Bissbort, S.; Gohler, F.; Schunter, F.; Wernet, P.

[27537] : Mapping of the linkage group GLO-Bf-HLA-B, C, A-PGM. I. Recombination

[27538] frequencies. Hum. Genet. 44: 313-319, 1978.

[27539] 3571.

[27540] Lamm, L. U.; Friedrich, U.; Petersen, G. B.; Jorgensen, J.; Nielsen,

[27541] J.; Therkelsen, A. J.; Kissmeyer-Nielsen, F.: Assignment of the major

[27542] histocompatibility complex to chromosome no. 6 in a family with a

[27543] pericentric inversion. Hum. Hered. 24: 273-284, 1974.

[27544] 3572.

[27545] Lamm, L. U.; Kissmeyer-Nielsen, F.; Svejgaard, A.; Pe-
tersen, G.
[27546] B.; Thorsby, E.; Mayr, W.; Hogman, C.: On the orientation
of the
[27547] HL-A region and the PGM(3) locus in the chromosome.
Tissue Antigens 2:
[27548] 205-214, 1972.

[27549] 3573.

[27550] Lamm, L. U.; Svejgaard, A.; Kissmeyer-Nielsen, F.: Further
evidence
[27551] for PGM(3): HL-A is another linkage in man. Nature N.B.
231: 109-110,
[27552] 1971.

[27553] 3574.

[27554] Lamm, L. U.; Thorsen, I.-L.; Petersen, G. B.; Jorgensen, J.;
Henningsen,
[27555] K.; Bech, B.; Kissmeyer-Nielsen, F.: Data on the HL-A link-
age group. Ann.
[27556] Hum. Genet. 38: 383-390, 1975.

[27557] 3575.

[27558] Lawlor, D. A.; Dickel, C. D.; Hauswirth, W. W.; Parham, P.: Ancient

[27559] HLA genes from 7,500-year-old archaeological remains. Nature 349:

[27560] 785-788, 1991.

[27561] 3576.

[27562] Lawlor, D. A.; Ward, F. E.; Ennis, P. D.; Jackson, A. P.; Parham,

[27563] P.: HLA-A and B polymorphisms predate the divergence of humans and

[27564] chimpanzees. (Letter) Nature 335: 268-271, 1988.

[27565] 3577.

[27566] Lawrance, S. K.; Smith, C. L.; Srivastava, R.; Cantor, C. R.;

[27567] Weissman, S. M.: Megabase-scale mapping of the HLA gene complex by

[27568] pulsed field gel electrophoresis. Science 235: 1387-1390, 1987.

[27569] 3578.

[27570] Levine, F.; Erlich, H.; Mach, B.; Leach, R.; White, R.; Pious,

[27571] D.: Deletion mapping of HLA and chromosome 6p genes. Proc. Nat.

[27572] Acad. Sci. 82: 3741-3745, 1985.

[27573] 3579.

[27574] Slate, D. L.; Ruddle, F. H.: Genetics of the interferon system. Pharm.

[27575] Therap. 4: 221-230, 1979.

[27576] 3580.

[27577] Stock, A. D.; Hsu, T. C.: Evolutionary conservatism in arrangement

[27578] of genetic material. A comparative analysis of chromosome banding

[27579] between the Rhesus macaque (2n=42, 84 arms) and the African green

[27580] monkey (2n=60, 120 arms). Chromosoma 43: 211-224, 1973.

[27581] 3581.

[27582] Streuli, M.; Nagata, S.; Weissmann, C.: At least three human

[27583] type alpha interferons: structure of alpha-2. Science 209: 1343-1347,

[27584] 1980.

[27585] 3582.

[27586] Szabo, S. J.; Sullivan, B. M.; Stemmann, C.; Satoskar, A. R.;

[27587] Sleckman, B. P.; Glimcher, L. H.: Distinct effects of T-bet in T(H)1

[27588] lineage commitment and IFN-gamma production in CD4 and CD8 T cells. Science 295:

[27589] 338-342, 2002.

[27590] 3583.

[27591] Takayanagi, H.; Ogasawara, K.; Hida, S.; Chiba, T.; Murata, S.;

[27592] Sato, K.; Takaoka, A.; Yokochi, T.; Oda, H.; Tanaka, K.; Nakamura,

[27593] K.; Taniguchi, T.: T-cell-mediated regulation of osteoclastogenesis

[27594] by signalling cross-talk between RANKL and IFN-gamma. Nature 408:

[27595] 600-605, 2000.

[27596] 3584.

[27597] Tan, Y. H.; Creagan, R. P.; Ruddle, F. H.: Assignment of the

[27598] genes of the human interferon system to chromosomes 2 and 5. Cytogenet.

[27599] Cell Genet. 13: 155-157, 1974.

[27600] 3585.

[27601] Tan, Y. H.; Ke, Y. H.; Armstrong, J. A.; Ho, M.: The regulation

[27602] of cellular interferon production: enhancement by antimetabolites. Proc.

[27603] Nat. Acad. Sci. 67: 464-471, 1970.

[27604] 3586.

[27605] Trent, J. M.; Olson, S.; Lawn, R. M.: Chromosomal localization

[27606] of human leukocyte, fibroblast, and immune interferon genes by means

[27607] of in situ hybridization. Proc. Nat. Acad. Sci. 79: 7809-7813, 1982.

[27608] 3587.

[27609] Tzoneva, M.; Ganev, V.; Galabov, A.; Georgieva, K.: Selective

[27610] immunodeficiency with defect in interferon-gamma induction in two

[27611] sibs with recurrent infections. Clin. Genet. 33: 454-456, 1988.

[27612] 3588.

[27613] White, A. C., Jr.; Robinson, P.; Okhuysen, P. C.; Lewis, D. E.;

[27614] Shahab, I.; Lahoti, S.; DuPont, H. L.; Chappell, C. L.: Interferon-gamma

[27615] expression in jejunal biopsies in experimental human cryptosporidiosis

[27616] correlates with prior sensitization and control of oocyst excretion. J.

[27617] Infect. Dis. 181: 701-709, 2000.

[27618] 3589.

[27619] Yip, Y. K.; Barrowclough, B. S.; Urban, C.; Vilcek, J.: Purification

[27620] of two subspecies of human gamma (immune) interferon. Proc. Nat.

[27621] Acad. Sci. 79: 1820-1824, 1982.

[27622] 3590.

[27623] Zimonjic, D. B.; Rezanka, L. J.; Evans, C. H.; Polymeropoulos,

[27624] M. H.; Trent, J. M.; Popescu, N. C.: Mapping of the immune interferon
[27625] gamma gene (IFNG) to chromosome band 12q14 by fluorescence in situ
[27626] hybridization. Cytogenet. Cell Genet. 71: 247-248, 1995.

[27627] 3591.

[27628] Zohlnhofer, D.; Richter, T.; Neumann, F.-J.; Nuhrenberg, T.; Wessely,
[27629] R.; Brandl, R.; Murr, A.; Klein, C. A.; Baeuerle, P. A.: Transcriptome
[27630] analysis reveals a role of interferon-gamma in human neointoma formation. Molec.
[27631] Cell 7: 1059-1069, 2001.

[27632] 3592.

[27633] Zoon, K. C.; Smith, M. E.; Bridgen, P. J.; Anfinsen, C. B.; Hunkapiller,
[27634] M. W.; Hood, L. E.: Amino terminal sequence of the major component
[27635] of human lymphoblastoid interferon. Science 207: 527-528, 1980.

[27636] 3593.

[27637] Schindler, C.; Fu, X.-Y.; Improta, T.; Aebersold, R.; Darnell,

[27638] J. E., Jr.: Proteins of transcription factor ISGF-3: one gene encodes

[27639] the 91- and 84-kDa ISGF-3 proteins that are activated by interferon

[27640] alpha. Proc. Nat. Acad. Sci. 89: 7836-7839, 1992.

[27641] 3594.

[27642] Eason, D. D.; Shepherd, A. T.; Blanck, G.: Interferon regu-latory

[27643] factor 1 tryptophan 11 to arginine point mutation abol-ishes DNA binding. Biochim.

[27644] Biophys. Acta 1446: 140-144, 1999.

[27645] 3595.

[27646] Savtchenko, E. S.; Tomic, M.; Ivker, R.; Blumenberg, M.: Three

[27647] parallel linkage groups of human acidic keratin genes. Genomics 7:

[27648] 394-407, 1990.

[27649] 3596.

[27650] Rosenberg, M.; Fuchs, E.; Le Beau, M. M.; Eddy, R. L.; Shows, T.

[27651] B.: Three epidermal and one simple epithelial type II keratin genes
[27652] map to human chromosome 12. Cytogenet. Cell Genet. 57: 33-38, 1991.
[27653] 3597.

[27654] Wang, N.; Perkins, K. L.: Involvement of band 3p14 in t(3;8)
[27655] hereditary renal carcinoma. Cancer Genet. Cytogenet. 11: 479-481,
[27656] 1984.

[27657] 3598.

[27658] Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J. M.:
[27659] Allelic knockout of novel splice variants of human recombination repair
[27660] gene RAD51B in t(12;14) uterine leiomyomas. Cancer Res. 59: 19-23,
[27661] 1999.

[27662] 3599.

[27663] Funk, C. D.; Funk, L. B.; FitzGerald, G. A.; Samuelsson, B.: Characterization

[27664] of human 12-lipoxygenase genes. Proc. Nat. Acad. Sci. 89: 3962-3966,

[27665] 1992.

[27666] 3600.

[27667] Carlsson, B.; Lindell, K.; Gabrielsson, B.; Karlsson, C.; Bjar-

[27668] nason, R.; Westphal, O.; Karlsson, U.; Sjostrom, L.; Carlsson, L. M. S.:

[27669] Obese (ob) gene defects are rare in human obesity. Obe-

[27670] sity Res. 5: 30-35, 1997.

[27671] 3601.

[27672] Yoshida, M. C.; Sasaki, M.; Mise, K.; Semba, K.; Nishizawa, M.;

[27673] Yamamoto, T.; Toyoshima, K.: Regional mapping of the human proto-oncogene

[27674] c-yes-1 to chromosome 18 at band q21.3. Jpn. J. Cancer Res. 76:

[27675] 559-562, 1985.

[27676] 3602.

[27677] Chehab, F. F.; Lim, M. E.; Lu, R.: Correction of the sterility

[27678] defect in homozygous obese female mice by treatment with the human

[27679] recombinant leptin. Nature Genet. 12: 318–320, 1996.

[27680] 3603.

[27681] Chen, G.; Koyama, K.; Yuan, X.; Lee, Y.; Zhou, Y.-T.; O'Doherty,

[27682] R.; Newgard, C. B.; Unger, R. H.: Disappearance of body fat in normal

[27683] rats induced by adenovirus-mediated leptin gene therapy. Proc. Nat.

[27684] Acad. Sci. 93: 14795–14799, 1996.

[27685] 3604.

[27686] Cooke, N. E.; Szpirer, C.; Levan, G.: The related genes en-coding

[27687] growth hormone and prolactin have been dispersed to chromosomes 10

[27688] and 17 in the rat. Endocrinology 119: 2451–2454, 1986.

[27689] 3605.

[27690] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Hormones,

[27691] active peptides and toxins. Washington: Biomedical Re-

search Foundation

[27692] (pub.) 5: 1972. Pp. D202 only.

[27693] 3606.

[27694] de Boer, H.; van der Veen, E. A.: Why retest young adults with

[27695] childhood-onset growth hormone deficiency? (Editorial) J. Clin. Endocr.

[27696] Metab. 82: 2032-2036, 1997.

[27697] 3607.

[27698] Stoffel, M.; Froguel, P.; Takeda, J.; Zouali, H.; Vionnet, N.;

[27699] Nishi, S.; Weber, I. T.; Harrison, R. W.; Pilkis, S. J.; Lesage, S.;

[27700] Vaxillaire, M.; Velho, G.; Sun, F.; Iris, F.; Passa, P.; Cohen, D.;

[27701] Bell, G. I.: Human glucokinase gene: isolation, characteri-zation,

[27702] and identification of two missense mutations linked to early-onset

[27703] non-insulin-dependent (type 2) diabetes mellitus. Proc. Nat. Acad.

[27704] Sci. 89: 7698-7702, 1992.

[27705] 3608.

[27706] Cuneo, R. C.; Judd, S.; Wallace, J. D.; Perry-Keene, D.; Burger,

[27707] H.; Lim-Tio, S.; Strauss, B.; Stockigt, J.; Topliss, D.; Alford, F.;

[27708] Hew, L.; Bode, H.; Conway, A.; Handelsman, D.; Dunn, S.; Boyages,

[27709] S.; Cheung, N. W.; Hurley, D.: The Australian multicenter trial of

[27710] growth hormone (GH) treatment in GH-deficient adults. J. Clin. Endocr.

[27711] Metab. 83: 107-116, 1998.

[27712] 3609.

[27713] Duquesnoy, P.; Amselem, S.; Gourmelen, M.; LeBouc, Y.; Goossens,

[27714] M.: A frameshift mutation causing isolated growth hormone deficiency

[27715] type 1A. (Abstract) Am. J. Hum. Genet. 47: A110, 1990.

[27716] 3610.

[27717] Fiddes, J. C.; Seeburg, P. H.; DeNoto, F. M.; Hallewell, R. A.;

[27718] Baxter, J. D.; Goodman, H. M.: Structure of genes for human growth
[27719] hormone and chorionic somatomammotropin. Proc. Nat. Acad. Sci. 76:
[27720] 4294-4298, 1979.

[27721] 3611.

[27722] George, D. L.; Phillips, J. A., III; Francke, U.; Seeburg, P.
[27723] H.: The genes for growth hormone and chorionic somatomammotropin
[27724] are on the long arm of human chromosome 17 in region q21-to-qter. Hum.
[27725] Genet. 57: 138-141, 1981.

[27726] 3612.

[27727] Gibney, J.; Wallace, J. D.; Spinks, T.; Schnorr, L.; Ranicar,
[27728] A.; Cuneo, R. C.; Lockhart, S.; Burnand, K. G.; Salomon, F.; Sonksen,
[27729] P. H.; Russell-Jones, D.: The effects of 10 years of recombinant
[27730] human growth hormone (GH) in adult GH-deficient patients. J. Clin.
[27731] Endocr. Metab. 84: 2596-2602, 1999.

[27732] 3613.

[27733] Goossens, M.; Brauner, R.; Czernichow, P.; Duquesnoy, P.; Rappaport,

[27734] R.: Isolated growth hormone (GH) deficiency type 1A associated with

[27735] a double deletion in the human GH gene cluster. J. Clin. Endocr.

[27736] Metab. 62: 712-716, 1986.

[27737] 3614.

[27738] Grumbach, M. M.: Personal Communication. San Francisco, Calif.

[27739] 1981.

[27740] 3615.

[27741] Guyda, H. J.: Four decades of growth hormone therapy for short

[27742] children: what have we achieved? J. Clin. Endocr. Metab. 84: 4307-4316,

[27743] 1999.

[27744] 3616.

[27745] Harper, M. E.; Barrera-Saldana, H. A.; Saunders, G. F.: Chromosomal

[27746] localization of the human placental lactogen-growth hormone gene cluster

[27747] to 17q22-24. Am. J. Hum. Genet. 34: 227-234, 1982.

[27748] 3617.

[27749] Hasegawa, Y.; Fujii, K.; Yamada, M.; Igarashi, Y.; Tachibana,

[27750] K.; Tanaka, T.; Onigata, K.; Nishi, Y.; Kato, S.; Hasegawa, T.: Identification

[27751] of novel human GH-1 gene polymorphisms that are associated with growth

[27752] hormone secretion and height. J. Clin. Endocr. Metab. 85: 1290-1295,

[27753] 2000.

[27754] 3618.

[27755] Hayashi, Y.; Yamamoto, M.; Ohmori, S.; Kamijo, T.; Ogawa, M.;

[27756] Seo, H.: Inhibition of growth hormone (GH) secretion by a mutant

[27757] GH-I gene product in neuroendocrine cells containing secretory granules:

[27758] an implication for isolated GH deficiency inherited in an autosomal

[27759] dominant manner. J. Clin. Endocr. Metab. 84: 2134-2139, 1999.

[27760] 3619.

[27761] He, Y. A.; Chen, S. S.; Wang, Y. X.; Lin, X. Y.; Wang, D. F.:

[27762] A Chinese familial growth hormone deficiency with a deletion of 7.1

[27763] kb of DNA. J. Med. Genet. 27: 151-154, 1990.

[27764] 3620.

[27765] Hindmarsh, P. C.; Dennison, E.; Pincus, S. M.; Cooper, C.; Fall,

[27766] C. H. D.; Matthews, D. R.; Pringle, P. J.; Brook, C. G. D.: A sexually

[27767] dimorphic pattern of growth hormone secretion in the elderly. J.

[27768] Clin. Endocr. Metab. 84: 2679-2685, 1999.

[27769] 3621.

[27770] Munier, F. L.; Korvatska, E.; Djemai, A.; Le Paslier, D.; Zografos,

[27771] L.; Pescia, G.; Schorderet, D. F.: Kerato-epithelin mutations in

[27772] four 5q31-linked corneal dystrophies. Nature Genet. 15:

247-251,
[27773] 1997.

[27774] 3622.

[27775] Irvine, A. D.; Corden, L. D.; Swensson, O.; Swensson, B.; Moore,
[27776] J. E.; Frazer, D. G.; Smith, F. J. D.; Knowlton, R. G.; Christophers,
[27777] E.; Rochels, R.; Uitto, J.; McLean, W. H. I.: Mutations in cornea-specific
[27778] keratin K3 or K12 genes cause Meesmann's corneal dystrophy. Nature
[27779] Genet. 16: 184-187, 1997.

[27780] 3623.

[27781] Bateman, J. F.; Lamande, S. R.; Dahl, H.-H. M.; Chan, D.; Mascara,
[27782] T.; Cole, W. G.: A frameshift mutation results in a truncated nonfunctional
[27783] carboxyl-terminal pro-alpha-1(I) propeptide of type I collagen in
[27784] osteogenesis imperfecta. J. Biol. Chem. 264: 10960-10964, 1989.
[27785] 3624.

[27786] Bateman, J. F.; Mascara, T.; Chan, D.; Cole, W. G.: A structural
[27787] mutation of the collagen alpha-1(I)CB7 peptide in lethal perinatal
[27788] osteogenesis imperfecta. J. Biol. Chem. 262: 4445-4451, 1987.
[27789] 3625.

[27790] Boedtker, H.; Fuller, F.; Tate, V.: The structure of collagen
[27791] genes. Int. Rev. Connect. Tissue Res. 10: 1-63, 1983.

[27792] 3626.

[27793] Bonadio, J.; Patterson, E.; Smiley, E.: DNA sequence analysis
[27794] of perinatal lethal OI mutations.(Abstract) Collagen Rel. Res. 8:
[27795] 506A, 1988.

[27796] 3627.

[27797] Bonaventure, J.; Cohen-Solal, L.; Lasselin, C.; Maroteaux, P.
[27798] : A dominant mutation in the COL1A1 gene that substitutes glycine
[27799] for valine causes recurrent lethal osteogenesis imperfecta.

Hum.

[27800] Genet. 89: 640-646, 1992.

[27801] 3628.

[27802] Brookes, A. J.; Sykes, B.; Solomon, E.: Single base pair al-

terations

[27803] as the predominant category of mutation in type I osteo-

genesis imperfecta.(Letter) J.

[27804] Med. Genet. 26: 410, 1989.

[27805] 3629.

[27806] Byers, P. H.: Personal Communication. Seattle, Wash.

9/23/1993.

[27807] 3630.

[27808] Byers, P. H.: Personal Communication. Seattle, Wash.

1990.

[27809] 3631.

[27810] Byers, P. H.; Starman, B. J.; Cohn, D. H.; Horwitz, A. L.: A

[27811] novel mutation causes a perinatal lethal form of osteoge-

nesis imperfecta:

[27812] an insertion in one alpha-1(I) collagen allele (COL1A1). J.

Biol.

[27813] Chem. 263: 7855-7861, 1988.

[27814] 3632.

[27815] Byers, P. H.; Duvic, M.; Atkinson, M.; Robinow, M.; Smith, L.

[27816] T.; Krane, S. M.; Greally, M. T.; Ludman, M.; Matalon, R.; Pauker,

[27817] S.; Quanbeck, D.; Schwarze, U.: Ehlers-Danlos syndrome type VIIA

[27818] and VIIB result from splice-junction mutations or genomic deletions

[27819] that involve exon 6 in the COL1A1 and COL1A2 genes of type I collagen. Am.

[27820] J. Med. Genet. 72: 94-105, 1997.

[27821] 3633.

[27822] Byrne, D. E. S.; Church, R. L.: Assignment of the genes for mouse

[27823] type I procollagen to chromosome 16 using mouse fi-broblast-Chinese

[27824] hamster somatic cell hybrids. Somat. Cell Genet. 9: 313-331, 1983.

[27825] 3634.

[27826] Chu, M.-L.; de Wet, W.; Bernard, M.; Ramirez, F.: Fine structural

[27827] analysis of the human pro-alpha-1(I) collagen gene: promoter structure,

[27828] AluI repeats, and polymorphic transcripts. J. Biol. Chem. 260: 2315-2320,

[27829] 1985.

[27830] 3635.

[27831] Church, R. L.; SundarRaj, N.; Rohrbach, D. H.: Gene mapping of

[27832] human ocular connective tissue proteins: assignment of the structural

[27833] gene for corneal type I procollagen to human chromosome 7 in human

[27834] corneal stroma-mouse fibroblast somatic cell hybrids. Invest. Ophthal. 21:

[27835] 73-79, 1981.

[27836] 3636.

[27837] Cohn, D. H.; Apone, S.; Eyre, D. R.; Starman, B. J.; Andreassen,

[27838] P.; Charbonneau, H.; Nicholls, A. C.; Pope, F. M.; Byers, P. H.:

[27839] Substitution of cysteine for glycine within the carboxyl-terminal

[27840] telopeptide of the alpha-1 chain of type I collagen produces mild

[27841] osteogenesis imperfecta. J. Biol. Chem. 263: 14605-14607, 1988.

[27842] 3637.

[27843] Cohn, D. H.; Zhang, X.; Byers, P. H.: Homology-mediated recombination

[27844] between type I collagen gene exons results in an internal tandem duplication

[27845] and lethal osteogenesis imperfecta. Hum. Mutat. 2: 21-27, 1993.

[27846] 3638.

[27847] Cole, W. G.; Campbell, P. E.; Rogers, J. G.; Bateman, J. F.:

[27848] The clinical features of osteogenesis imperfecta resulting from a

[27849] non-functional carboxy terminal pro-alpha-1(I) propeptide of type

[27850] I procollagen and a severe deficiency of normal type I collagen in

[27851] tissues. J. Med. Genet. 27: 545-551, 1990.

[27852] 3639.

[27853] Cole, W. G.; Chow, C. W.; Bateman, J. F.; Sillence, D. O.: The

[27854] phenotypic features of osteogenesis imperfecta resulting from a mutation

[27855] of the carboxyl-terminal pro-alpha-1(I) propeptide that impairs the

[27856] assembly of type I procollagen and formation of the ex-tracellular

[27857] matrix. J. Med. Genet. 33: 965-967, 1996.

[27858] 3640.

[27859] Cole, W. G.; Chow, C. W.; Rogers, J. G.; Bateman, J. F.: The

[27860] clinical features of three babies with osteogenesis imper-fecta resulting

[27861] from the substitution of glycine by arginine in the pro al-pha-1(I)

[27862] chain of type I procollagen. J. Med. Genet. 27: 228-235, 1990.

[27863] 3641.

[27864] Cole, W. G.; Evans, R.; Sillence, D. O.: The clinical features

[27865] of Ehlers-Danlos syndrome type VII due to a deletion of 24 amino acids

[27866] from the pro-alpha-1(I) chain of type I procollagen. J. Med. Genet. 24:

[27867] 698-701, 1987.

[27868] 3642.

[27869] Cole, W. G.; Patterson, E.; Bonadio, J.; Campbell, P. E.; For-tune,

[27870] D. W.: The clinicopathological features of three babies with osteogenesis

[27871] imperfecta resulting from the substitution of glycine by valine in

[27872] the pro alpha-1(I) chain of type I procollagen. J. Med. Genet. 29:

[27873] 112-118, 1992.

[27874] 3643.

[27875] Constantinou, C. D.; Pack, M.; Young, S. B.; Prockop, D. J.:

[27876] Phenotypic heterogeneity in osteogenesis imperfecta: the mildly affected

[27877] mother of a proband with a lethal variant had the same mutation substituting

[27878] cysteine for alpha-1-glycine 904 in a type I procollagen gene (COL1A1). Am.

[27879] J. Hum. Genet. 47: 670-679, 1990.

[27880] 3644.

[27881] D'Alessio, M.; Ramirez, F.; Blumberg, B. D.; Wirtz, M. K.; Rao,

[27882] V. H.; Godfrey, M. D.; Hollister, D. W.: Characterization of a COL1A1

[27883] splicing defect in a case of Ehlers-Danlos syndrome type VII: further

[27884] evidence of molecular homogeneity. Am. J. Hum. Genet. 49: 400-406,

[27885] 1991.

[27886] 3645.

[27887] Stoffel, M.; Patel, P.; Lo, Y.-M. D.; Hattersley, A. T.; Lucassen,

[27888] A. M.; Page, R.; Bell, J. I.; Bell, G. I.; Turner, R. C.; Wainscoat,

[27889] J. S.: Missense glucokinase mutation in maturity-onset diabetes of

[27890] the young and mutation screening in late-onset diabetes. Nature Genet. 2:

[27891] 153-156, 1992.

[27892] 3646.

[27893] Stone, L. M.; Kahn, S. E.; Fujimoto, W. Y.; Deeb, S. S.; Porte,

[27894] D. Jr.: A variation at position -30 of the beta-cell glucokinase

[27895] gene promoter is associated with reduced beta-cell function in middle-aged

[27896] Japanese-American men. Diabetes 45: 422-428, 1996.

[27897] 3647.

[27898] Sun, F.; Knebelmann, B.; Pueyo, M. E.; Zouali, H.; Lesage, S.;

[27899] Vaxillaire, M.; Passa, P.; Cohen, D.; Velho, G.; Antignac, C.; Froguel,

[27900] P.: Deletion of the donor splice site of intron 4 in the glucokinase

[27901] gene causes maturity-onset diabetes of the young. J. Clin. Invest. 92:

[27902] 1174-1180, 1993.

[27903] 3648.

[27904] Thornton, P. S.; Satin-Smith, M. S.; Herold, K.; Glaser, B.; Chiu,

[27905] K. C.; Nestorowicz, A.; Permutt, M. A.; Baker, L.; Stanley, C. A.

[27906] : Familial hyperinsulinism with apparent autosomal dominant inheritance:

[27907] clinical and genetic differences from the autosomal recessive variant. J.

[27908] Pediat. 132: 9-14, 1998.

[27909] 3649.

[27910] Velho, G.; Blanche, H.; Vaxillaire, M.; Bellane-Chantelot, C.;

[27911] Pardini, V. C.; Timsit, J.; Passa, P.; Deschamps, I.; Robert, J.-J.;

[27912] Weber, I. T.; Marotta, D.; Pilkis, S. J.; Lipkind, G. M.; Bell, G.

[27913] I.; Froguel, P.: Identification of 14 new glucokinase mutations and

[27914] description of the clinical profile of 42 MODY-2 families. Diabetologia 40:

[27915] 217-224, 1997.

[27916] 3650.

[27917] Velho, G.; Froguel, P.; Clement, K.; Pueyo, M. E.; Rakotoambinina,

[27918] B.; Zouali, H.; Passa, P.; Cohen, D.; Robert, J.-J.: Primary pancreatic
[27919] beta-cell secretory defect caused by mutations in glucokinase gene
[27920] in kindreds of maturity onset diabetes of the young. Lancet 340:
[27921] 444-448, 1992.

[27922] 3651.

[27923] Velho, G.; Petersen, K. F.; Perseghin, G.; Hwang, J.-H.; Rothman,
[27924] D. L.; Pueyo, M. E.; Cline, G. W.; Froguel, P.; Shulman, G. I.: Impaired
[27925] hepatic glycogen synthesis in glucokinase-deficient (MODY-2) subjects. J.
[27926] Clin. Invest. 98: 1755-1761, 1996.

[27927] 3652.

[27928] Hendershot, L. M.; Valentine, V. A.; Lee, A. S.; Morris, S. W.;
[27929] Shapiro, D. N.: Localization of the gene encoding human BiP/GRP78,
[27930] the endoplasmic reticulum cognate of the HSP70 family, to chromosome

[27931] 9q34. Genomics 20: 281-284, 1994.

[27932] 3653.

[27933] Law, M. L.; Seeliger, M. B.; Lee, A. S.; Kao, F. T.: Genetic mapping

[27934] of the structural gene coding for a glucose-regulated protein (GRP78)

[27935] of 78k-dalton to the long arm of human chromosome 9. (Abstract) Cytogenet.

[27936] Cell Genet. 37: 518-519, 1984.

[27937] 3654.

[27938] Lee, A. S.; Delegeane, A.; Scharff, D.: Highly conserved glucose-regulated

[27939] protein in hamster and chicken cells: preliminary characterization

[27940] of its cDNA clone. Proc. Nat. Acad. Sci. 78: 4922-4925, 1981.

[27941] 3655.

[27942] Lee, A. S.; Delegeane, A. M.; Baker, V.; Chow, P. C.: Transcriptional

[27943] regulation of two genes specifically induced by glucose starvation

[27944] in hamster mutant fibroblast cell line. J. Biol. Chem. 258: 597-603,

[27945] 1983.

[27946] 3656.

[27947] Muresan, Z.; Arvan, P.: Enhanced binding to the molecular chaperone

[27948] BiP slows thyroglobulin export from the endoplasmic reticulum. Molec.

[27949] Endocr. 12: 458-467, 1998.

[27950] 3657.

[27951] Anagnou, N. P.; Seuanez, H.; Modi, W.; O'Brien, S. J.; Pap-

matheakis,

[27952] J.; Moschonas, N.: Chromosomal mapping of the human glutamate dehydrogenase

[27953] (GLUD) genes to chromosomes 10q21.1-21.2 and Xq26-28. (Abstract) Am.

[27954] J. Hum. Genet. 45 (suppl.): A170 only, 1989.

[27955] 3658.

[27956] Barbeau, A.; Charbonneau, M.; Cloutier, T.: Leucocyte glu-

tamate

[27957] dehydrogenase in various hereditary ataxias. Canad. J.

Neurol. Sci. 7:

[27958] 421-424, 1980.

[27959] 3659.

[27960] Colon, A. D.; Plaitakis, A.; Perakis, A.; Berl, S.; Clarke, D.

[27961] D.: Purification and characterization of a soluble and a particulate

[27962] glutamate dehydrogenase from rat brain. J. Neurochem. 46: 1811-1819,

[27963] 1986.

[27964] 3660.

[27965] Deloukas, P.; Dauwerse, J. G.; Moschonas, N. K.; van Om-

men, G.

[27966] J. B.; van Loon, A. P. G. M.: Three human glutamate dehy-

drogenase

[27967] genes (GLUD1, GLUDP2, and GLUDP3) are located on chromosome 10q, but

[27968] are not closely physically linked. Genomics 17: 676-681, 1993.

[27969] 3661.

[27970] Duvoisin, R. C.; Chokroverty, S.; Lepore, F.; Nicklas, W. J.:

[27971] Glutamate dehydrogenase deficiency in patients with

olivopontocerebellar
[27972] atrophy. Neurology 33: 1322-1326, 1983.

[27973] 3662.

[27974] Hanauer, A.; Mandel, J. L.; Mattei, M. G.: X-linked and autosomal
[27975] sequences corresponding to glutamate dehydrogenase (GLUD) and to an
[27976] anonymous cDNA. (Abstract) Cytogenet. Cell Genet. 40: 647-648, 1985.
[27977] 3663.

[27978] Dalgleish, R.: The human type I collagen mutation database. Nucleic
[27979] Acids Res. 25: 181-187, 1997.

[27980] 3664.

[27981] Dolphin, C.; Shephard, E. A.; Povey, S.; Palmer, C. N. A.; Ziegler,
[27982] D. M.; Ayesh, R.; Smith, R. L.; Phillips, I. R.: Cloning, primary
[27983] sequence, and chromosomal mapping of a human flavin-containing monooxygenase
[27984] (FMO1). J. Biol. Chem. 266: 12379-12385, 1991.

[27985] 3665.

[27986] Shephard, E. A.; Dolphin, C.; Povey, S.; Smith, R.; Phillips,
I.

[27987] : The genes encoding the flavin-containing monooxyge-
nases FMO1 and

[27988] FMO2 are located on human chromosome 1q. (Abstract)
Cytogenet. Cell

[27989] Genet. 58: 1863 only, 1991.

[27990] 3666.

[27991] Shephard, E. A.; Dolphin, C. T.; Fox, M. F.; Povey, S.;
Smith,

[27992] R.; Phillips, I. R.: Localization of genes encoding three dis-
tinct

[27993] flavin-containing monooxygenases to human chromo-
some 1q. Genomics 16:

[27994] 85-89, 1993.

[27995] 3667.

[27996] Ziegler, D. M.: Flavin-containing monooxygenases: en-
zymes adapted

[27997] for multisubstrate specificity. Trends Pharm. Sci. 11:
321-324,

[27998] 1990.

[27999] 3668.

[28000] McCombie, R. R.; Dolphin, C. T.; Povey, S.; Phillips, I. R.; Shephard,

[28001] E. A.: Localization of human flavin-containing monooxy-genase genes

[28002] FMO2 and FMO5 to chromosome 1q. Genomics 34: 426-429, 1996.

[28003] 3669.

[28004] Barnes, D. E.; Johnston, L. H.; Kodama, K.; Tomkinson, A. E.; Lasko,

[28005] D. D.; Lindahl, T.: Human DNA ligase I cDNA: cloning and functional

[28006] expression in Saccharomyces cerevisiae. Proc. Nat. Acad. Sci. 87:

[28007] 6679-6683, 1990.

[28008] 3670.

[28009] Barnes, D. E.; Kodama, K.-I.; Tynan, K.; Trask, B. J.; Chris-tensen,

[28010] M.; de Jong, P. J.; Spurr, N. K.; Lindahl, T.; Mohrenweiser, H. W.

[28011] : Assignment of the gene-encoding DNA ligase I to human chromosome

[28012] 19q13.2-13.3. Genomics 12: 164-166, 1992.

[28013] 3671.

[28014] Barnes, D. E.; Tomkinson, A. E.; Lehmann, A. R.; Webster, A. D.

[28015] B.; Lindahl, T.: Mutations in the DNA ligase I gene of an individual

[28016] with immunodeficiencies and cellular hypersensitivity to DNA-damaging

[28017] agents. Cell 69: 495-503, 1992.

[28018] 3672.

[28019] Bentley, D. J.; Selfridge, J.; Millar, J. K.; Samuel, K.; Hole,

[28020] N.; Ansell, J. D.; Melton, D. W.: DNA ligase I is required for fetal

[28021] liver erythropoiesis but is not essential for mammalian cell viability. Nature

[28022] Genet. 13: 489-491, 1996.

[28023] 3673.

[28024] Gariboldi, M.; Montecucco, A.; Columbano, A.; Ledda-Columbano,

[28025] G. M.; Savini, E.; Manenti, G.; Pierotti, M. A.; Dragani, T. A.:

[28026] Genetic mapping and expression analysis of the murine DNA ligase gene. Molec.

[28027] Carcinog. 14: 71-74, 1995.

[28028] 3674.

[28029] Lindahl, T.; Barnes, D. E.: Mammalian DNA ligases. Annu. Rev.

[28030] Biochem. 61: 251-281, 1992.

[28031] 3675.

[28032] Noguiez, P.; Barnes, D. E.; Mohrenweiser, H. W.; Lindahl, T.:

[28033] Structure of the human DNA ligase I gene. Nucleic Acids Res. 20:

[28034] 3845-3850, 1992.

[28035] 3676.

[28036] Souttou, B.; Juhl, H.; Hackenbruck, J.; Rockseisen, M.; Klomp,

[28037] H. J.; Raulais, D.; Vigny, M.; Wellstein, A.: Relationship be-tween

[28038] serum concentrations of the growth factor pleiotrophin and pleiotrophin-positive

[28039] tumors. J. Nat. Cancer Inst. 90: 1468-1473, 1998.

[28040] 3677.

[28041] Weber, D.; Klomp, H.-J.; Czubayko, F.; Wellstein, A.; Juhl, H.

[28042] : Pleiotrophin can be rate-limiting for pancreatic cancer cell growth. Cancer

[28043] Res. 60: 5284-5288, 2000.

[28044] 3678.

[28045] Zhang, N.; Yeh, H.-J.; Zhong, R.; Li, Y.-S.; Deuel, T. F.: A dominant-negative

[28046] pleiotrophin mutant introduced by homologous recombination leads to

[28047] germ-cell apoptosis in male mice. Proc. Nat. Acad. Sci. 96: 6734-6738,

[28048] 1999.

[28049] 3679.

[28050] Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.;

[28051] Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of neurite

[28052] outgrowth-promoting proteins: heparin-binding neurite outgrowth promoting

[28053] factor maps to 11p11-11p13. (Abstract) Cytogenet. Cell Genet. 58:

[28054] 1958 only, 1991.

[28055] 3680.

[28056] Kaname, T.; Kuwano, A.; Murano, I.; Uehara, K.; Mura-

matsu, T.;

[28057] Kajii, T.: Midkine gene (MDK), a gene for prenatal differ-

entiation

[28058] and neuroregulation, maps to band 11p11.2 by fluores-

cence in situ

[28059] hybridization. Genomics 17: 514-515, 1993.

[28060] 3681.

[28061] Simon-Chazottes, D.; Matsubara, S.; Miyauchi, T.; Mura-

matsu, T.;

[28062] Guenet, J.-L.: Chromosomal localization of two cell sur-

face-associated

[28063] molecules of potential importance in development: mid-

kine (Mdk) and

[28064] basigin (Bsg). Mammalian Genome 2: 269-271, 1992.

[28065] 3682.

[28066] Airaksinen, E. M.; Iivanainen, M.; Karli, P.; Sainio, K.; Hal-

tia,

[28067] M.: Hereditary recurrent brachial plexus neuropathy with dysmorphic

[28068] features. Acta Neurol. Scand. 71: 309-316, 1985.

[28069] 3683.

[28070] Chance, P. F.; Lensch, M. W.; Lipe, H.; Brown, R. H., Sr.; Brown,

[28071] R. H., Jr.; Bird, T. D.: Hereditary neuralgic amyotrophy and hereditary

[28072] neuropathy with liability to pressure palsies: two distinct genetic

[28073] disorders. Neurology 44: 2253-2257, 1994.

[28074] 3684.

[28075] Dreschfeld, J.: On some rarer forms of muscular atrophies. Brain 79:

[28076] 226-232, 1886.

[28077] 3685.

[28078] Gardner, J. H.; Maloney, W.: Hereditary brachial and cranial neuritis

[28079] genetically linked with ocular hypotelorism and syn-dactyly. (Abstract) Neurology 18:

[28080] 278, 1968.

[28081] 3686.

[28082] Geiger, L. R.; Mancall, E. L.; Penn, A. S.; Tucker, S. H.: Fa-
milial
[28083] neuralgic amyotrophy: report of three families with review
of the
[28084] literature. Brain 97: 87-102, 1974.

[28085] 3687.

[28086] Gouider, R.; LeGuern, E.; Emile, J.; Tardieu, S.; Cabon, F.;
Samid,
[28087] M.; Weissenbach, J.; Agid, Y.; Bouche, P.; Brice, A.: Heredi-
tary
[28088] neuralgic amyotrophy and hereditary neuropathy with lia-
bility to pressure
[28089] palsies: two distinct clinical, electrophysiologic, and ge-
netic entities. Neurology 44:
[28090] 2250-2252, 1994.

[28091] 3688.

[28092] Guillozet, N.; Mercer, R. D.: Hereditary recurrent brachial
neuropathy. Am.
[28093] J. Dis. Child. 125: 884-887, 1973.

[28094] 3689.

[28095] Jacob, J. C.; Andermann, F.; Robb, J. P.: Heredofamilial neuritis

[28096] with brachial predilection. Neurology 11: 1025-1033, 1961.

[28097] 3690.

[28098] Jeannet, P.-Y.; Watts, G. D. J.; Bird, T. D.; Chance, P. F.: Craniofacial

[28099] and cutaneous findings expand the phenotype of hereditary neuralgic

[28100] amyotrophy. Neurology 57: 1963-1968, 2001.

[28101] 3691.

[28102] Matthews, W. B.; Esiri, M.: The migrant sensory neuritis of Wartenberg. J.

[28103] Neurol. Neurosurg. Psychiat. 46: 1-4, 1983.

[28104] 3692.

[28105] Meuleman, J.; Kuhlenbaumer, G.; Audenaert, D.; Hunermund, G.;

[28106] Hor, H.; Young, P.; Stogbauer, F.; Ringelstein, E. B.; Van Broeckhoven,

[28107] C.; De Jonghe, P.; Timmerman, V.: Mutation analysis of 4

candidate
[28108] genes for hereditary neuralgic amyotrophy (HNA). Hum. Genet. 108:
[28109] 390-393, 2001.

[28110] 3693.

[28111] Meuleman, J.; Timmerman, V.; Van Broeckhoven, C.; De Jonghe, P.
[28112] : Hereditary neuralgic amyotrophy. Neurogenetics 3: 115-118, 2001.
[28113] 3694.

[28114] Orstavik, K.; Ro, H.; Orstavik, K. H.: Recurrent brachial plexus
[28115] neuropathy in a family with subtle dysmorphic features--a case of
[28116] hereditary neuralgic amyotrophy. Clin. Genet. 51: 421-425, 1997.
[28117] 3695.

[28118] Pellegrino, J. E.; George, R. A. V.; Biegel, J.; Farlow, M. R.;

[28119] Gardner, K.; Caress, J.; Brown, M. J.; Rebbeck, T. R.; Bird, T. D.;
[28120] Chance, P. F.: Hereditary neuralgic amyotrophy: evidence

for genetic
[28121] homogeneity and mapping to chromosome 17q25. Hum. Genet. 101: 277-283,
[28122] 1997.

[28123] 3696.

[28124] Pellegrino, J. E.; Rebbeck, T. R.; Brown, M. J.; Bird, T. D.;

[28125] Chance, P. F.: Mapping of hereditary neuralgic amyotrophy (familial
[28126] brachial plexus neuropathy) to distal chromosome 17q. Neurology 46:
[28127] 1128-1132, 1996.

[28128] 3697.

[28129] Phillips, L. H., II: Familial long thoracic nerve palsy: a man-ifestation
[28130] of brachial plexus neuropathy. Neurology 36: 1251-1253, 1986.
[28131] 3698.

[28132] Smith, B. H.; Ramakrishna, T.; Schlagenhauff, R. R.: Famil-ial
[28133] brachial neuropathy: two case reports with discussion. Neurology 21:

[28134] 941-945, 1971.

[28135] 3699.

[28136] Stogbauer, F.; Young, P.; Timmerman, V.; Spoelders, P.; Bernd

[28137] Ringelstein, E.; Van Broeckhoven, C.; Kurlemann, G.: Re-finement of

[28138] the hereditary neuralgic amyotrophy (HNA) locus to chro-mosome 17q24-q25. Hum.

[28139] Genet. 99: 685-687, 1997.

[28140] 3700.

[28141] Taylor, R. A.: Heredofamilial mononeuritis multiplex with brachial

[28142] predilection. Brain 83: 113-137, 1960.

[28143] 3701.

[28144] Thomas, P. K.; Ormerod, I. E. C.: Hereditary neuralgic amyotrophy

[28145] associated with a relapsing multifocal sensory neuropathy. J. Neurol.

[28146] Neurosurg. Psychiat. 56: 107-109, 1993.

[28147] 3702.

[28148] Tsairis, P.; Dyck, P. J.; Mulder, D. W.: Natural history of brachial
[28149] plexus neuropathy. Arch. Neurol. 27: 109-117, 1972.

[28150] 3703.

[28151] Watts, G. D. J.; O'Briant, K. C.; Chance, P. F.: Evidence of
[28152] a founder effect and refinement of the hereditary neuralgic amyotrophy
[28153] (HNA) locus on 17q25 in American families. Hum. Genet. 110: 166-172,
[28154] 2002.

[28155] 3704.

[28156] Windebank, A. J.: Inherited recurrent focal neuropathies.In:
[28157] Dyck, P. J.; Thomas, P. K.; Griffin, J. W.; Low, P. A.; Poduslo, J.
[28158] F. (eds.): Peripheral Neuropathy. Volume 2. Philadelphia: W. B.
[28159] Saunders Company (3rd ed.): 1993. Pp. 1137-1148.

[28160] 3705.

[28161] Windebank, A. J.; Schenone, A.; Dewald, G. W.: Hereditary neuropathy

[28162] with liability to pressure palsies and inherited brachial plexus neuropathy:

[28163] two genetically distinct disorders. Mayo Clin. Proc. 70: 743-746,

[28164] 1995.

[28165] 3706.

[28166] Cool, D. R.; Normant, E.; Shen, F.; Chen, H.-C.; Pannell, L.; Zhang,

[28167] Y.; Loh, Y. P.: Carboxypeptidase E is a regulated secretory pathway

[28168] sorting receptor: genetic obliteration leads to endocrine disorders

[28169] in Cpe(fat) mice. Cell 88: 73-83, 1997.

[28170] 3707.

[28171] Beg, A. A.; Sha, W. C.; Bronson, R. T.; Ghosh, S.; Baltimore, D.

[28172] : Embryonic lethality and liver degeneration in mice lacking the RelA

[28173] component of NF-kappa B. Nature 376: 167-170, 1995.

[28174] 3708.

[28175] Chen, L.; Fischle, W.; Verdin, E.; Greene, W. C.: Duration of

[28176] nuclear NF-kappa-B action regulated by reversible acetylation. Science 293:

[28177] 1653-1657, 2001.

[28178] 3709.

[28179] Deloukas, P.; Dauwerse, J. G.; van Ommen, G. J. B.; van Loon, A.

[28180] P. G. M.: The human NFKB3 gene encoding the p65 subunit of transcription

[28181] factor NF-kappa-B is located on chromosome 11q12. Genomics 19: 592-594,

[28182] 1994.

[28183] 3710.

[28184] Deloukas, P.; van Loon, A. P. G. M.: Genomic organization of the

[28185] gene encoding the p65 subunit of NF-kappa-B: multiple variants of

[28186] the p65 protein may be generated by alternative splicing. Hum. Molec.

[28187] Genet. 2: 1895-1900, 1993.

[28188] 3711.

[28189] Neurath, M. F.; Pettersson, S.; Myer zum Buschenfelde,

K.-H.;

[28190] Strober, W.: Local administration of antisense phospho-
rothioate oligonucleotides

[28191] to the p65 subunit of NF-kappa-B abrogates established
experimental

[28192] colitis in mice. Nature Med. 2: 998-1004, 1996.

[28193] 3712.

[28194] Zhong, H.; Voll, R. E.; Ghosh, S.: Phosphorylation of NF-
kappa

[28195] B by PKA stimulates transcriptional activity by promoting a
novel

[28196] bivalent interaction with the coactivator CBP/p300. Molec.
Cell 1:

[28197] 661-671, 1998.

[28198] 3713.

[28199] Biamonti, G.; Buvoli, M.; Bassi, M. T.; Morandi, C.; Co-
bianchi,

[28200] F.; Riva, S.: Isolation of an active gene encoding human
hnRNP protein

[28201] A1: evidence for alternative splicing. J. Molec. Biol. 207:
491-503,

[28202] 1989.

[28203] 3714.

[28204] Buvoli, M.; Biamonti, G.; Tsoulfas, P.; Bassi, M. T.; Ghetti, A.;

[28205] Riva, S.; Morandi, C.: cDNA cloning of human hnRNP protein A1 reveals

[28206] the existence of multiple mRNA isoforms. Nucleic Acids Res. 16:

[28207] 3751-3770, 1988.

[28208] 3715.

[28209] Michael, W. M.; Choi, M.; Dreyfuss, G.: A nuclear export signal

[28210] in hnRNP A1: a signal-mediated, temperature-dependent nuclear protein

[28211] export pathway. Cell 83: 415-422, 1995.

[28212] 3716.

[28213] Saccone, S.; Biamonti, G.; Maugeri, S.; Bassi, M. T.; Bunone, G.;

[28214] Riva, S.; Della Valle, G.: Assignment of the human heterogeneous

[28215] nuclear ribonucleoprotein A1 gene (HNRPA1) to chromosome 12q13.1 by

[28216] cDNA competitive in situ hybridization. Genomics 12: 171-174, 1992.

[28217] 3717.

[28218] Pollard, A. J.; Sparey, C.; Robson, S. C.; Krainer, A. R.; Europe-Finner,

[28219] G. N.: Spatio-temporal expression of the trans-acting splicing factors

[28220] SF2/ASF and heterogeneous ribonuclear proteins A1/A1B in the myometrium

[28221] of the pregnant human uterus: a molecular mechanism for regulating

[28222] regional protein isoform expression in vivo. J. Clin. Endocr. Metab. 85:

[28223] 1928-1936, 2000.

[28224] 3718.

[28225] Turc-Carel, C.; Pietrzak, E.; Kakati, S.; Kinniburgh, A. J.; Sandberg,

[28226] A. A.: INT1 maps to 12q12-12q13. (Abstract) Cytogenet. Cell Genet. 46:

[28227] 706 only, 1987.

[28228] 3719.

[28229] van't Veer, L. J.; Geurts van Kessel, A. H. M.; van Heerikhuizen,
[28230] H.; van Ooyen, A.; Nusse, R.: Molecular cloning and chromosomal assignment
[28231] of the human homolog of int-1, a mouse gene implicated in mammary
[28232] tumorigenesis. Molec. Cell. Biol. 4: 2532-2534, 1984.

[28233] 3720.

[28234] van Ooyen, A.; Kwee, V.; Nusse, R.: The nucleotide sequence of
[28235] the human int-1 mammary oncogene; evolutionary conservation of coding
[28236] and non-coding sequences. EMBO J. 4: 2905-2909, 1985.

[28237] 3721.

[28238] Wolda, S. L.; Moon, R. T.: Cloning and developmental expression
[28239] in Xenopus laevis of seven additional members of the Wnt family. Oncogene 7:
[28240] 1941-1947, 1992.

[28241] 3722.

[28242] Armstrong, B. C.; Krystal, G. W.: Isolation and characteri-

zation

[28243] of complementary DNA for N-cym, a gene encoded by the DNA strand opposite

[28244] to N-myc. Cell Growth Differ. 3: 385-390, 1992.

[28245] 3723.

[28246] Brodeur, G. M.; Seeger, R. C.: Gene amplification in human neuroblastomas:

[28247] basic mechanisms and clinical implications. Cancer Genet. Cytogenet. 19:

[28248] 101-111, 1986.

[28249] 3724.

[28250] Brodeur, G. M.; Seeger, R. C.; Schwab, M.; Varmus, H. E.; Bishop,

[28251] J. M.: Amplification of N-myc in untreated human neurob-lastomas correlates

[28252] with advanced disease stage. Science 224: 1121-1124, 1984.

[28253] 3725.

[28254] Campbell, G. R.; Zimmerman, K.; Blank, R. D.; Alt, F. W.; D'Eustachio,

[28255] P.: Chromosomal location of N-myc and L-myc genes in

the mouse. Oncogene
[28256] Res. 4: 47-54, 1989.

[28257] 3726.

[28258] Corvi, R.; Amler, L. C.; Savelyeva, L.; Gehring, M.; Schwab, M.
[28259] : MYCN is retained in single copy at chromosome 2 band p23-24 during
[28260] amplification in human neuroblastoma cells. Proc. Nat. Acad. Sci. 91:
[28261] 5523-5527, 1994.

[28262] 3727.

[28263] Emanuel, B. S.; Balaban, G.; Boyd, J. P.; Grossman, A.; Negishi,
[28264] M.; Parmiter, A.; Glick, M. C.: N-myc amplification in mul-tiple homogeneously
[28265] staining regions in two human neuroblastomas. Proc. Nat. Acad. Sci. 82:
[28266] 3736-3740, 1985.

[28267] 3728.

[28268] Garson, J. A.; van den Berghe, J. A.; Kemshead, J. T.: Novel non-isotopic

[28269] in situ hybridization technique detects small (1 kb) unique sequences

[28270] in routinely G-banded human chromosomes: fine mapping of N-myc and

[28271] beta-NGF genes. Nucleic Acids Res. 15: 4761-4770, 1987.

[28272] 3729.

[28273] Guo, C.; White, P. S.; Weiss, M. J.; Hogarty, M. D.; Thompson,

[28274] P. M.; Stram, D. O.; Gerbing, R.; Matthay, K. K.; Seeger, R. C.; Brodeur,

[28275] G. M.; Maris, J. M.: Allelic deletion at 11q23 is common in MYCN

[28276] single copy neuroblastomas. Oncogene 18: 4948-4957, 1999.

[28277] 3730.

[28278] Kanda, N.; Schreck, R.; Alt, F.; Bruns, G.; Baltimore, D.; Latt,

[28279] S.: Isolation of amplified DNA sequences from IMR-32 human neuroblastoma

[28280] cells: facilitation by fluorescence-activated flow sorting of metaphase

[28281] chromosomes. Proc. Nat. Acad. Sci. 80: 4069-4073, 1983.

[28282] 3731.

[28283] Kohl, N. E.; Gee, C. E.; Alt, F. W.: Activated expression of

[28284] the N-myc gene in human neuroblastomas and related tu-
mors. Science 226:

[28285] 1335-1337, 1984.

[28286] 3732.

[28287] Kohl, N. E.; Kanda, N.; Schreck, R. R.; Bruns, G.; Latt, S. A.;

[28288] Gilbert, F.; Alt, F. W.: Transposition and amplification of
oncogene-related

[28289] sequences in human neuroblastomas. Cell 35: 359-367,
1983.

[28290] 3733.

[28291] Kohl, N. E.; Legouy, E.; DePinho, R. A.; Nisen, P. D.; Smith,

[28292] R. K.; Gee, C. E.; Alt, F. W.: Human N-myc is closely re-
lated in

[28293] organization and nucleotide sequence to c-myc. Nature
319: 73-77,

[28294] 1986.

[28295] 3734.

[28296] Lee, W.-H.; Murphree, A. L.; Benedict, W. F.: Expression and
[28297] amplification of the N-myc gene in primary retinoblas-
toma. Nature 309:
[28298] 458-460, 1984.

[28299] 3735.

[28300] Michitsch, R. W.; Melera, P. W.: Nucleotide sequence of the 3-prime
[28301] exon of the human N-myc gene. Nucleic Acids Res. 13:
2545-2558,
[28302] 1985.

[28303] 3736.

[28304] Ramsay, G.; Stanton, L.; Schwab, M.; Bishop, J. M.: Human proto-oncogene
[28305] N-myc encodes nuclear proteins that bind DNA. Molec.
Cell. Biol. 6:
[28306] 4450-4457, 1986.

[28307] 3737.

[28308] Reiter, J. L.; Brodeur, G. M.: High-resolution mapping of a 130-kb
[28309] core region of the MYCN amplicon in neuroblastomas. Ge-

nomics 32:
[28310] 97-103, 1996.

[28311] 3738.

[28312] Schwab, M.: Amplification of N-myc in human neuroblastomas. Trends
[28313] Genet. 1: 271-275, 1985.

[28314] 3739.

[28315] Schwab, M.; Alitalo, K.; Klempnauer, K.-H.; Varmus, H. E.; Bishop,
[28316] J. M.; Gilbert, F.; Brodeur, G.; Goldstein, M.; Trent, J.: Amplified
[28317] DNA with limited homology to myc cellular oncogene is shared by human
[28318] neuroblastoma cell lines and a neuroblastoma tumour. Nature 305:
[28319] 245-248, 1983.

[28320] 3740.

[28321] Schwab, M.; Ellison, J.; Busch, M.; Rosenau, W.; Varmus, H. E.;
[28322] Bishop, J. M.: Enhanced expression of the human gene N-myc consequent

[28323] to amplification of DNA may contribute to malignant progression of
[28324] neuroblastoma. Proc. Nat. Acad. Sci. 81: 4940-4944, 1984.
[28325] 3741.

[28326] Schwab, M.; Varmus, H. E.; Bishop, J. M.; Grzeschik, K.-H.; Naylor,
[28327] S. L.; Sakaguchi, A. Y.; Brodeur, G.; Trent, J.: Chromosome localization
[28328] in normal human cells and neuroblastomas of a gene related to c-myc. Nature 308:
[28329] 288-291, 1984.

[28330] 3742.

[28331] Seeger, R. C.; Brodeur, G. M.; Sather, H.; Dalton, A.; Siegel,
[28332] S. E.; Wong, K. Y.; Hammond, D.: Association of multiple copies of
[28333] the N-myc oncogene with rapid progression of neuroblastomas. New
[28334] Eng. J. Med. 313: 1111-1116, 1985.

[28335] 3743.

[28336] Shiloh, Y.; Shipley, J.; Brodeur, G. M.; Bruns, G.; Korf, B.;

[28337] Donlon, T.; Schreck, R. R.; Seeger, R.; Sakai, K.; Latt, S. A.: Differential
[28338] amplification, assembly, and relocation of multiple DNA sequences
[28339] in human neuroblastomas and neuroblastoma cell lines. Proc. Nat.
[28340] Acad. Sci. 82: 3761-3765, 1985.

[28341] 3744.

[28342] Grant, S. F. A.; Reid, D. M.; Blake, G.; Herd, R.; Fogelman, I.;
[28343] Ralston, S. H.: Reduced bone density and osteoporosis as-sociated
[28344] with a polymorphic Sp1 binding site in the collagen type I-alpha 1
[28345] gene. Nature Genet. 14: 203-205, 1996.

[28346] 3745.

[28347] Pocock, N. A.; Eisman, J. A.; Hopper, J. L.; Yeates, M. G.; Sambrook,
[28348] P. N.; Eberl, S.: Genetic determinants of bone mass in adults: a
[28349] twin study. J. Clin. Invest. 80: 706-710, 1987.

[28350] 3746.

[28351] Lusis, A. J.; Heinzmann, C.; Sparkes, R. S.; Scott, J.; Knott,

[28352] T. J.; Geller, R.; Sparkes, M. C.; Mohandas, T.: Regional mapping

[28353] of human chromosome 19: organization of genes for plasma lipid transport

[28354] (APOC1, -C2, and -E and LDLR) and the genes C3, PEPD, and GPI. Proc.

[28355] Nat. Acad. Sci. 83: 3929-3933, 1986.

[28356] 3747.

[28357] Martiniuk, F.; Ellenbogen, A.; Hirschhorn, K.; Hirschhorn, R.

[28358] : Further regional localization of the genes for human acid alpha

[28359] glucosidase (GAA), peptidase D (PEPD), and alpha mannosidase B (MANB)

[28360] by somatic cell hybridization. Hum. Genet. 69: 109-111, 1985.

[28361] 3748.

[28362] Siciliano, M. J.; Stallings, R. L.; Adair, G. M.; Humphrey, R.

[28363] M.; Siciliano, J.: Provisional assignment of TPI, GPI, and

PEPD to
[28364] Chinese hamster autosomes 8 and 9: a cytogenetic basis for functional
[28365] haploidy of an autosomal linkage group in CHO cells. Cy-
togenet. Cell
[28366] Genet. 35: 15-20, 1983.

[28367] 3749.

[28368] Taketo, M.; Matsui, M.; Rochelle, J. M.; Yodoi, J.; Seldin, M.

[28369] F.: Mouse thioredoxin gene maps on chromosome 4, whereas its pseudogene
[28370] maps on chromosome 1. Genomics 21: 251-253, 1994.

[28371] 3750.

[28372] Tonissen, K. F.; Wells, J. R. E.: Isolation and characteriza-
tion
[28373] of human thioredoxin-encoding genes. Gene 102:
221-228, 1991.
[28374] 3751.

[28375] Wollman, E. E.; d'Auriol, L.; Rimsky, L.; Shaw, A.; Jacquot, J.-P.;
[28376] Wingfield, P.; Graber, P.; Dessarps, F.; Robin, P.; Galibert, F.;

[28377] Bertoglio, J.; Fradelizi, D.: Cloning and expression of a cDNA for
[28378] human thioredoxin. J. Biol. Chem. 263: 15506-15512, 1988.
[28379] 3752.

[28380] Caen, J. P.; Castaldi, P. A.; Leclerc, J. C.; Inceman, S.; Lar-
rieu,
[28381] M. J.; Probst, M.; Bernard, J.: Congenital bleeding disor-
ders with
[28382] long bleeding time and normal platelet count. I. Glanz-
mann's thrombasthenia
[28383] (report of fifteen patients). Am. J. Med. 41: 4-26, 1966.

[28384] 3753.

[28385] Ruggeri, Z. M.; Bader, R.; de Marco, L.: Glanzmann throm-
basthenia:
[28386] deficient binding of von Willebrand factor to thrombin-
stimulated
[28387] platelets. Proc. Nat. Acad. Sci. 79: 6038-6041, 1982.

[28388] 3754.

[28389] Bahou, W. F.; Nierman, W. C.; Durkin, A. S.; Potter, C. L.;
Demetrick,

[28390] D. J.: Chromosomal assignment of the human thrombin receptor gene:
[28391] localization to region q13 of chromosome 5. Blood 82: 1532-1537,
[28392] 1993.

[28393] 3755.

[28394] Coughlin, S. R.; Vu, T.-K. H.; Hung, D. T.; Wheaton, V. I.: Characterization
[28395] of a functional thrombin receptor: issues and opportuni- ties. J. Clin.
[28396] Invest. 89: 351-355, 1992.

[28397] 3756.

[28398] Griffin, C. T.; Srinivasan, Y.; Zheng, Y.-W.; Huang, W.; Coughlin,
[28399] S. R.: A role for thrombin receptor signaling in endothelial cells
[28400] during embryonic development. Science 293: 1666-1670, 2001.
[28401] 3757.

[28402] Poirier, C.; O'Brien, E. P.; Bueno Brunialti, A. L.; Chambard,

[28403] J.-C.; Swank, R. T.; Guenet, J.-L.: The gene encoding the

thrombin

[28404] receptor (Cf2r) maps to mouse chromosome 13. Mam-

malian Genome 7:

[28405] 322, 1996.

[28406] 3758.

[28407] Schmidt, V. A.; Nierman, W. C.; Feldblyum, T. V.; Maglott,

D. R.;

[28408] Bahou, W. F.: The human thrombin receptor and pro-

teinase activated

[28409] receptor-2 genes are tightly linked on chromosome 5q13.

Brit. J.

[28410] Haemat. 97: 523-529, 1997.

[28411] 3759.

[28412] Schmidt, V. A.; Vitale, E.; Bahou, W. F.: Genomic cloning

and

[28413] characterization of the human thrombin receptor gene:

structural similarity

[28414] to the proteinase activated receptor-2 gene. J. Biol. Chem.

271:

[28415] 9307-9312, 1996.

[28416] 3760.

[28417] Vu, T.-K. H.; Hung, D. T.; Wheaton, V. I.; Coughlin, S. R.: Molecular

[28418] cloning of a functional thrombin receptor reveals a novel proteolytic

[28419] mechanism of receptor activation. Cell 64: 1057-1068, 1991.

[28420] 3761.

[28421] Dean, G.: The Porphyrias. A Story of Inheritance and Environment.

[28422] Philadelphia: J. B. Lippincott (pub.) (2nd ed.): 1972.

[28423] 3762.

[28424] Hukshorn, C. J.; Saris, W. H. M.; Westerterp-Plantenga, M. S.;

[28425] Farid, A. R.; Smith, F. J.; Campfield, L. A.: Weekly subcutaneous

[28426] pegylated recombinant native human leptin (PEG-OB) administration

[28427] in obese men. J. Clin. Endocr. Metab. 85: 4003-4009, 2000.

[28428] 3763.

[28429] Vafiadis, P.; Ounissi-Benkalha, H.; Palumbo, M.; Grabs, R.; Rousseau,

[28430] M.; Goodyer, C. G.; Polychronakos, C.: Class III alleles of the variable
[28431] number of tandem repeat insulin polymorphism associated with silencing
[28432] of thymic insulin predispose to type 1 diabetes. J. Clin. Endocr.
[28433] Metab. 86: 3705-3710, 2001.

[28434] 3764.

[28435] Bennett, M. K.; Calakos, N.; Scheller, R. H.: Syntaxin: a synaptic
[28436] protein implicated in docking of synaptic vesicles at presynaptic
[28437] active zones. Science 257: 255-259, 1992.

[28438] 3765.

[28439] Bennett, M. K.; Garcia-Arraras, J. E.; Elferink, L. A.; Peterson,
[28440] K.; Fleming, A. M.; Hazuka, C. D.; Scheller, R. H.: The syntaxin
[28441] family of vesicular transport receptors. Cell 74: 863-873, 1993.
[28442] 3766.

[28443] Fernandez, I.; Ubach, J.; Dulubova, I.; Zhang, X.; Sudof, T. C.;

[28444] Rizo, J.: Three-dimensional structure of an evolutionarily conserved

[28445] N-terminal domain of syntaxin 1A. Cell 94: 841-849, 1998.

[28446] 3767.

[28447] Fisher, R. J.; Pevsner, J.; Burgoyne, R. D.: Control of fusion

[28448] pore dynamics during exocytosis by Munc18. Science 291: 875-878,

[28449] 2001.

[28450] 3768.

[28451] Martin-Martin, B.; Nabokina, S. M.; Lazo, P. A.; Mollinedo, F.

[28452] : Co-expression of several human syntaxin genes in neutrophils and

[28453] differentiating HL-60 cells: various isoforms and detection of syntaxin

[28454] 1. J. Leuko. Biol. 65: 397-406, 1999.

[28455] 3769.

[28456] Nakayama, T.; Fujiwara, T.; Miyazawa, A.; Asakawa, S.;

Shimizu,
[28457] N.; Shimizu, Y.; Mikoshiba, K.; Akagawa, K.: Mapping of the human
[28458] HPC-1/syntaxin 1A gene (STX1A) to chromosome 7 band q11.2. Genomics 42:
[28459] 173-176, 1997.

[28460] 3770.

[28461] Naren, A. P.; Di, A.; Cormet-Boyaka, E.; Boyaka, P. N.; McGhee,
[28462] J. R.; Zhou, W.; Akagawa, K.; Fujiwara, T.; Thome, U.; En-gelhardt,
[28463] J. F.; Nelson, D. J.; Kirk, K. L.: Syntaxin 1A is expressed in airway
[28464] epithelial cells, where it modulates CFTR C1- currents. J. Clin.
[28465] Invest. 105: 377-386, 2000.

[28466] 3771.

[28467] Richmond, J. E.; Weimer, R. M.; Jorgensen, E. M.: An open form
[28468] of syntaxin bypasses the requirement for UNC-13 in vesi-cle priming. Nature 412:
[28469] 338-341, 2001.

[28470] 3772.

[28471] Zhang, R.; Maksymowych, A. B.; Simpson, L. L.: Cloning and sequence

[28472] analysis of a cDNA encoding human syntaxin 1A, a polypeptide essential

[28473] for exocytosis. Gene 159: 293-294, 1995.

[28474] 3773.

[28475] Shepley, M. P.; Sherry, B.; Weiner, H. L.: Monoclonal antibody

[28476] identification of a 100-kDa membrane protein in HeLa cells and human

[28477] spinal cord involved in poliovirus attachment. Proc. Nat. Acad. Sci. 85:

[28478] 7743-7747, 1988.

[28479] 3774.

[28480] Siddique, T.; Bartlett, R. J.; McKinney, R.; Hung, W.-Y.; Bruns,

[28481] G.; Wilfert, C.; Roses, A. D.: The poliovirus sensitivity (PVS) is

[28482] on chromosome 19q13-qter. (Abstract) Cytogenet. Cell Genet. 40:

[28483] 745 only, 1985.

[28484] 3775.

[28485] Siddique, T.; McKinney, R.; Hung, W.-Y.; Bartlett, R. J.; Bruns,

[28486] G.; Mohandas, T. K.; Ropers, H.-H.; Wilfert, C.; Roses, A. D.: The

[28487] poliovirus sensitivity (PVS) gene is on chromosome 19q12-q13.2. Genomics 3:

[28488] 156-160, 1988.

[28489] 3776.

[28490] Solecki, D. J.; Gromeier, M.; Mueller, S.; Bernhardt, G.; Wimmer,

[28491] E.: Expression of the human poliovirus receptor/CD155 gene is activated

[28492] by Sonic hedgehog. J. Biol. Chem. 277: 25697-25702, 2002.

[28493] 3777.

[28494] Morris, C. M.; Bodger, M. P.: Localization of the human poly(A)-binding

[28495] protein gene (PAB1) to chromosomal regions 3q22-q25, 12q13-q14, and

[28496] 13q12-q13 by in situ hybridization. Genomics 15: 209-211, 1993.

[28497] 3778.

[28498] Alkhatib, H. M.; Chen, D.; Cherney, B.; Bhatia, K.; Notario, V.;

[28499] Giri, C.; Stein, G.; Slattery, E.; Roeder, R. G.; Smulson, M. E.:

[28500] Cloning and expression of cDNA for human poly(ADP-ribose) polymerase. Proc.

[28501] Nat. Acad. Sci. 84: 1224-1228, 1987.

[28502] 3779.

[28503] Auer, B.; Nagl, U.; Herzog, H.; Schneider, R.; Schweiger, M.:

[28504] Human nuclear NAD+ ADP-ribosyltrans-ferase(polymerizing): organization

[28505] of the gene. DNA 8: 575-580, 1989.

[28506] 3780.

[28507] Baumgartner, M.; Schneider, R.; Auer, B.; Herzog, H.; Schweiger,

[28508] M.; Hirsch-Kauffmann, M.: Fluorescence in situ mapping of the human

[28509] nuclear NAD+ ADP-ribosyltransferase gene (ADPRT) and two secondary

[28510] sites to human chromosomal bands 1q42, 13q34, and 14q24. Cytogenet.

[28511] Cell Genet. 61: 172-174, 1992.

[28512] 3781.

[28513] Bhatia, K. G.; Cherney, B. W.; Huppi, K.; Magrath, I. T.; Cossman,

[28514] J.; Sausville, E.; Barriga, F.; Johnson, B.; Gause, B.; Bonney, G.;

[28515] Neequayi, J.; DeBernardi, M.; Smulson, M.: A deletion linked to a

[28516] poly(ADP-ribose) polymerase gene on chromosome 3q33-qter occurs frequently

[28517] in the normal black population as well as in multiple tumor DNA. Cancer

[28518] Res. 50: 5406-5413, 1990.

[28519] 3782.

[28520] Cherney, B. W.; McBride, O. W.; Chen, D.; Alkhatib, H.; Bhatia,

[28521] K.; Hensley, P.; Smulson, M. E.: cDNA sequence, protein structure,

[28522] and chromosomal location of the human gene for poly(ADP-ribose) polymerase. Proc.
[28523] Nat. Acad. Sci. 84: 8370-8374, 1987.

[28524] 3783.

[28525] d'Adda di Fagagna, F.; Hande, M. P.; Tong, W.-M.; Lansdorp, P.
[28526] M.; Wang, Z.-Q.; Jackson, S. P.: Functions of poly(ADP-ribose) polymerase
[28527] in controlling telomere length and chromosomal stability. Nature
[28528] Genet. 23: 76-80, 1999.

[28529] 3784.

[28530] Doll, J. A.; Suarez, B. K.; Donis-Keller, H.: Association be-tween
[28531] prostate cancer in black Americans and an allele of the PADPRP pseudogene
[28532] locus on chromosome 13. (Letter) Am. J. Hum. Genet. 58: 425-428,
[28533] 1996.

[28534] 3785.

[28535] Flick, K.; Schneider, R.; Auer, B.; Hirsch-Kauffmann, M.;

Schweiger,

[28536] M.: No abnormalities in the NAD(+) ADP-ribosyltransferase (polymerizing)

[28537] gene of transformed cells from a Fanconi's anemia patient. (Letter) Hum.

[28538] Genet. 89: 690-691, 1992.

[28539] 3786.

[28540] Grube, K.; Burkle, A.: Poly(ADP-ribose) polymerase activity in

[28541] mononuclear leukocytes of 13 mammalian species correlates with species-specific

[28542] life span. Proc. Nat. Acad. Sci. 89: 11759-11763, 1992.

[28543] 3787.

[28544] Herzog, H.; Zabel, B.; Schneider, R.; Auer, B.; Hirsch-Kauffmann,

[28545] M.; Schweiger, M.: Human NAD(+):protein ADP ribosyltransferase (ADPRT):

[28546] synthesis of active enzyme in E. coli, sequence of its cDNA from Hela

[28547] cells and chromosomal localization. Biol. Chem. Hoppe-Seyler 369:

[28548] 836-837, 1988.

[28549] 3788.

[28550] Herzog, H.; Zabel, B. U.; Schneider, R.; Auer, B.; Hirsch-
Kauffmann,
[28551] M.; Schweiger, M.: Human nuclear NAD(+) ADP-
ribosyltransferase: localization
[28552] of the gene on chromosome 1q41-q42 and expression of
an active human
[28553] enzyme in Escherichia coli. Proc. Nat. Acad. Sci. 86:
3514-3518,
[28554] 1989.

[28555] 3789.

[28556] Keijzer, W.; Stefanini, M.; Bootsma, D.; Verkerk, A.; Geurts
van
[28557] Kessel, A. H. M.; Jongkind, J. F.; Westerveld, A.: Localiza-
tion of
[28558] a gene involved in complementation of the defect in xero-
derma pigmentosum
[28559] group A cells on human chromosome 1. Exp. Cell Res.
169: 490-501,
[28560] 1987.

[28561] 3790.

[28562] Kurosaki, T.; Ushiro, H.; Mitsuuchi, Y.; Suzuki, S.; Mat-
suda,
[28563] M.; Matsuda, Y.; Katunuma, N.; Kangawa, K.; Matsuo, H.;
Hirose, T.;
[28564] Inayama, S.; Shizuta, Y.: Primary structure of human
poly(ADP-ribose)
[28565] synthetase as deduced from cDNA sequence. J. Biol.
Chem. 262: 15990-15997,
[28566] 1987.

[28567] 3791.

[28568] Loetscher, P.; Alvarez-Gonzalez, R.; Althaus, F. R.:
Poly(ADP-ribose)
[28569] may signal changing metabolic conditions to the chro-
matin of mammalian
[28570] cells. Proc. Nat. Acad. Sci. 84: 1286-1289, 1987.

[28571] 3792.

[28572] Lyn, D.; Cherney, B. W.; Lalande, M.; Berenson, J. R.; Licht-
enstein,
[28573] A.; Lupold, S.; Bhatia, K. G.; Smulson, M.: A duplicated re-
gion is

[28574] responsible for the poly(ADP-ribose) polymerase polymorphism, on chromosome

[28575] 13, associated with a predisposition to cancer. Am. J. Hum. Genet. 52:

[28576] 124-134, 1993.

[28577] 3793.

[28578] McBride, O. W.; Cherney, B.; Takourdin, C.; Smulson, M. E.: Human

[28579] poly(ADP-ribose) polymerase sequences are located on chromosomes 1,

[28580] 13, and 14. (Abstract) Cytogenet. Cell Genet. 46: 659-660, 1987.

[28581] 3794.

[28582] Pieper, A. A.; Brat, D. J.; Krug, D. K.; Watkins, C. C.; Gupta,

[28583] A.; Blackshaw, S.; Verma, A.; Wang, Z.-Q.; Snyder, S. H.: Poly(ADP-ribose)

[28584] polymerase-deficient mice are protected from streptozotocin-induced

[28585] diabetes. Proc. Nat. Acad. Sci. 96: 3059-3064, 1999.

[28586] 3795.

[28587] Schweiger, M.; Auer, B.; Burtscher, H. J.; Hirsch-

Kauffmann, M.;

[28588] Klocker, H.; Schneider, R.: DNA repair in human cells: bio-
chemistry

[28589] of the hereditary diseases Fanconi's anaemia and Cock-
ayne syndrome. Europ.

[28590] J. Biochem. 165: 235-242, 1987.

[28591] 3796.

[28592] Simbulan-Rosenthal, C. M.; Haddad, B. R.; Rosenthal, D.
S.; Weaver,

[28593] Z.; Coleman, A.; Luo, R.; Young, H. M.; Wang, Z.-Q.; Ried,
T.; Smulson,

[28594] M. E.: Chromosomal aberrations in PARP -/- mice:
genome stabilization

[28595] in immortalized cells by reintroduction of
poly(ADP-ribose) polymerase

[28596] cDNA. Proc. Nat. Acad. Sci. 96: 13191-13196, 1999.

[28597] 3797.

[28598] Smithies, O.; Gregg, R. G.; Boggs, S. S.; Koralewski, M. A.;
Kucherlapati,

[28599] R. S. :Nature (London) 317: 230-234, 1985.

[28600] 3798.

[28601] Vinik, A.; Bell, G.: Mutant insulin syndromes. Hormone Metab.

[28602] Res. 20: 1-10, 1988.

[28603] 3799.

[28604] Vinik, A. I.; Seino, S.; Funakoshi, A.; Schwartz, J.; Mat-sumoto,

[28605] M.; Schteingart, D. E.; Fu, Z.-Z.; Tsai, S.-T.: Familial hy-perinsulinemia

[28606] associated with secretion of an abnormal insulin, and co-existence

[28607] of insulin resistance in the propositus. J. Clin. Endocr. Metab. 62:

[28608] 645-652, 1986.

[28609] 3800.

[28610] Wang, J.; Takeuchi, T.; Tanaka, S.; Kubo, S.-K.; Kayo, T.; Lu,

[28611] D.; Takata, K.; Koizumi, A.; Izumi, T.: A mutation in the insulin

[28612] 2 gene induces diabetes with severe pancreatic beta-cell dysfunction

[28613] in the Mody mouse. J. Clin. Invest. 103: 27-37, 1999.

[28614] 3801.

[28615] Warren-Perry, M. G.; Manley, S. E.; Ostrega, D.; Polonsky, K.;

[28616] Mussett, S.; Brown, P.; Turner, R. C.: A novel point mutation in

[28617] the insulin gene giving rise to hyperproinsulinemia. J. Clin. Endocr.

[28618] Metab. 82: 1629-1631, 1997.

[28619] 3802.

[28620] White, R.; Leppert, M.; Bishop, D. T.; Barker, D.; Berkowitz,

[28621] J.; Brown, C.; Callahan, P.; Holm, T.; Jerominski, L.: Construction

[28622] of linkage maps with DNA markers for human chromosomes. Nature 313:

[28623] 101-105, 1985.

[28624] 3803.

[28625] Williams, L. G.; Jowett, N. I.; Vella, M. A.; Humphries, S.; Galton,

[28626] D. J.: Allelic variation adjacent to the human insulin and apolipoprotein

[28627] C-II genes in different ethnic groups. Hum. Genet. 71:

227-230,

[28628] 1985.

[28629] 3804.

[28630] Yano, H.; Kitano, N.; Morimoto, M.; Polonsky, K. S.; Imura, H.;

[28631] Seino, Y.: A novel point mutation in the human insulin gene giving

[28632] rise to hyperproinsulinemia (proinsulin Kyoto). J. Clin. Invest. 89:

[28633] 1902-1907, 1992.

[28634] 3805.

[28635] Yoshioka, M.; Kayo, T.; Ikeda, T.; Koizumi, A.: A novel locus,

[28636] Mody4, distal to D7Mit189 on chromosome 7 determines early-onset NIDDM

[28637] in nonobese C57BL/6 (Akita) mutant mice. Diabetes 46: 887-894, 1997.

[28638] 3806.

[28639] Hasan, S.; Hassa, P. O.; Imhof, R.; Hottiger, M. O.: Transcription

[28640] coactivator p300 binds PCNA and may have a role in DNA

repair synthesis. Nature 410:
[28641] 387-391, 2001.

[28642] 3807.

[28643] Hoege, C.; Pfander, B.; Moldovan, G.-L.; Pyrowolakis, G.; Jentsch,
[28644] S.: RAD6-dependent DNA repair is linked to modification of PCNA by
[28645] ubiquitin and SUMO. Nature 419: 135-141, 2002.

[28646] 3808.

[28647] Fonatsch, C.; Duchrow, M.; Rieder, H.; Schluter, C.; Gerdes, J.
[28648] : Assignment of the human Ki-67 gene (MKI67) to 10q25-qter. Genomics 11:
[28649] 476-477, 1991.

[28650] 3809.

[28651] Schluter, C.; Duchrow, M.; Wohlenberg, C.; Becker, M. H. G; Key,
[28652] G.; Flad, H.-D.; Gerdes, J.: The cell proliferation-associated antigen
[28653] of antibody Ki-67: a very large, ubiquitous nuclear protein with numerous

[28654] repeated elements, representing a new kind of cell cycle-maintaining
[28655] proteins. J. Cell. Biol. 123: 513-522, 1993.

[28656] 3810.

[28657] Schonk, D. M.; Kuijpers, H. J. H.; vanDrunen, E.; vanDalen, C.
[28658] H.; Geurts van Kessel, A. H. M.; Verheijen, R.; Ramaekers, F. C. S.
[28659] : Assignment of the gene(s) involved in the expression of the proliferation-related
[28660] Ki-67 antigen to human chromosome 10. Hum. Genet. 83: 297-299, 1989.

[28661] 3811.

[28662] Traut, W.; Scholzen, T.; Winking, H.; Kubbutat, M. H. G.; Gerdes,
[28663] J.: Assignment of the murine Ki-67 gene (Mki67) to chromosome band
[28664] 7F3-F5 by in situ hybridization. Cytogenet. Cell Genet. 83: 12-13,
[28665] 1998.

[28666] 3812.

[28667] Cooke, N. E.; Baxter, J. D.: Structural analysis of the pro-
lactin
[28668] gene suggests a separate origin for its 5-prime end. Na-
ture 297:
[28669] 603-606, 1982.

[28670] 3813.

[28671] Cooke, N. E.; Coit, D.; Shine, J.; Baxter, J. D.; Martial, J. A.

[28672] : Human prolactin: cDNA structural analysis and evolu-
tionary comparisons. J.
[28673] Biol. Chem. 256: 4007-4016, 1981.

[28674] 3814.

[28675] D'Alessandro, E.; Santiemma, V.; Lo Re, M. L.; Ligas, C.;
Del Porto,
[28676] G.: 6p23 deletion mosaicism in a woman with recurrent
abortions and
[28677] idiopathic hypoprolactinemia. Am. J. Med. Genet. 44:
220-222, 1992.
[28678] 3815.

[28679] DiMattia, G. E.: Personal Communication. London, Ontario,
Canada
[28680] 3/23/1998.

[28681] 3816.

[28682] DiMattia, G. E.; Gellersen, B.; Duckworth, M. L.; Friesen, H. G.

[28683] : Human prolactin gene expression: the use of an alternative noncoding

[28684] exon in decidua and the IM-9-P3 lymphoblast cell line. J. Biol. Chem. 265:

[28685] 16412-16421, 1990.

[28686] 3817.

[28687] Evans, A. M.; Petersen, J. W.; Sekhon, G. S.; DeMars, R. I.: Use

[28688] of human lymphoblastoid deletion mutants to map the prolactin gene

[28689] on human chromosome 6p. (Abstract) Am. J. Hum. Genet. 43: A143 only,

[28690] 1988.

[28691] 3818.

[28692] Farid, N. R.; Noel, E. P.; Sampson, L.; Russell, N. A.: Prolactin-secreting

[28693] adenomata are possibly associated with HLA-B8. Tissue Antigens 15:

[28694] 333-335, 1980.

[28695] 3819.

[28696] Gellersen, B.; Kempf, R.; Telgmann, R.; DiMattia, G. E.: Nonpituitary

[28697] human prolactin gene transcription is independent of Pit-1 and differentially

[28698] controlled in lymphocytes and in endometrial stroma. Molec. Endocr. 8:

[28699] 356-373, 1994.

[28700] 3820.

[28701] Rouleau, G. A.; Bazanowski, A.; Gusella, J. F.; Haines, J. L.

[28702] : A genetic map of chromosome 1: comparison of different data sets

[28703] and linkage programs. Genomics 7: 313-318, 1990.

[28704] 3821.

[28705] Erdile, L. F.; Heyer, W.-D.; Kolodner, R.; Kelly, T. J.: Characterization

[28706] of a cDNA encoding the 70-kDa single-stranded DNA-binding subunit

[28707] of human replication protein A and the role of the protein in DNA

[28708] replication. J. Biol. Chem. 266: 12090-12098, 1991.

[28709] 3822.

[28710] Gomes, X. V.; Wold, M. S.: Functional domains of the 70-kilodalton

[28711] subunit of human replication protein A. Biochemistry 35: 10558-10568,

[28712] 1996.

[28713] 3823.

[28714] Shen, L. X.; Basilion, J. P.; Stanton, V. P., Jr.: Single-nucleotide

[28715] polymorphisms can cause different structural folds of mRNA. Proc.

[28716] Nat. Acad. Sci. 96: 7871-7876, 1999.

[28717] 3824.

[28718] Umbricht, C. B.; Erdile, L. F.; Jabs, E. W.; Kelly, T. J.: Cloning,

[28719] overexpression, and genomic mapping of the 14-kDa subunit of human

[28720] replication protein A. J. Biol. Chem. 268: 6131-6138, 1993.

[28721] 3825.

[28722] Umbricht, C. B.; Griffin, C. A.; Hawkins, A. L.; Grzeschik, K.

[28723] H.; O'Connell, P.; Leach, R.; Green, E. D.; Kelly, T. J.: High-resolution

[28724] genomic mapping of the three human replication protein A genes (RPA1,

[28725] RPA2, and RPA3). Genomics 20: 249-257, 1994.

[28726] 3826.

[28727] Fukata, M.; Watanabe, T.; Noritake, J.; Nakagawa, M.; Ya-maga, M.;

[28728] Kuroda, S.; Matsuura, Y.; Iwamatsu, A.; Perez, F.; Kaibuchi, K.:

[28729] Rac1 and Cdc42 capture microtubules through IQGAP1 and CLIP-170. Cell 109:

[28730] 873-885, 2002.

[28731] 3827.

[28732] Meyaard, L.; van der Vuurst de Vries, A.-R.; de Ruiter, T.; Lanier,

[28733] L. L.; Phillips, J. H.; Clevers, H.: The epithelial cellular ad-hesion

[28734] molecule (Ep-CAM) is a ligand for the leukocyte-associ-ated immunoglobulin-like

[28735] receptor (LAIR). J. Exp. Med. 194: 107-112, 2001.

[28736] 3828.

[28737] Hitchins, M. P.; Monk, D.; Bell, G. M.; Ali, Z.; Preece, M. A.;

[28738] Stanier, P.; Moore, G. E.: Maternal repression of the human GRB10

[28739] gene in the developing central nervous system; evaluation of the role

[28740] for GRB10 in Silver-Russell syndrome. Europ. J. Hum. Genet. 9: 82-90,

[28741] 2001.

[28742] 3829.

[28743] McCann, J. A.; Zheng, H.; Islam, A.; Goodyer, C. G.; Poly-chronakos,

[28744] C.: Evidence against GRB10 as the gene responsible for Silver-Russell

[28745] syndrome. Biochem. Biophys. Res. Commun. 286: 943-948, 2001.

[28746] 3830.

[28747] Miyoshi, N.; Kuroiwa, Y.; Kohda, T.; Shitara, H.; Yonekawa, H.;

[28748] Kawabe, T.; Hasegawa, H.; Barton, S. C.; Surani, M. A.;

Kaneko-Ishino,
[28749] T.; Ishino, F.: Identification of the Meg1/Grb10 imprinted gene on
[28750] mouse proximal chromosome 11, a candidate for the Silver-Russell syndrome
[28751] gene. Proc. Nat. Acad. Sci. 95: 1102-1107, 1998.

[28752] 3831.

[28753] Yoshihashi, H.; Maeyama, K.; Kosaki, R.; Ogata, T.; Tsukahara,
[28754] M.; Goto, Y.; Hata, J.; Matsuo, N.; Smith, R. J.; Kosaki, K.: Imprinting
[28755] of human GRB10 and its mutations in two patients with Russell-Silver
[28756] syndrome. Am. J. Hum. Genet. 67: 476-482, 2000.

[28757] 3832.

[28758] Grob, P. M.; Ross, A. H.; Koprowski, H.; Bothwell, M.: Characterization
[28759] of the human melanoma nerve growth factor receptor. J. Biol. Chem. 260:
[28760] 8044-8049, 1985.

[28761] 3833.

[28762] Stanton, L. W.; Schwab, M.; Bishop, J. M.: Nucleotide se-
quence
[28763] of the human N-myc gene. Proc. Nat. Acad. Sci. 83:
1772-1776, 1986.
[28764] 3834.

[28765] Brown, M. A.; Nicolai, H.; Xu, C.-F.; Griffiths, B. L.; Jones,
[28766] K. A.; Solomon, E.; Hosking, L.; Trowsdale, J.; Black, D. M.;
McFarlane,
[28767] R.: Regulation of BRCA1. (Letter) Nature 372: 733 only,
1994.
[28768] 3835.

[28769] Brown, M. A.; Xu, C.-F.; Nicolai, H.; Griffiths, B.; Cham-
bers,
[28770] J. A.; Black, D.; Solomon, E.: The 5-prime end of the
BRCA1 gene
[28771] lies within a duplicated region of human chromosome
17q21. Oncogene 12:
[28772] 2507-2513, 1996.

[28773] 3836.

[28774] Campbell, I. G.; Nicolai, H. M.; Foulkes, W. D.; Senger, G.;
Stamp,

[28775] G. W.; Allan, G.; Boyer, C.; Jones, K.; Bast, R. C., Jr.; Solomon,
[28776] E.; Trowsdale, J.; Black, D. M.: A novel gene encoding a B-box protein
[28777] within the BRCA1 region at 17q21.1. Hum. Molec. Genet. 3: 589-594,
[28778] 1994.

[28779] 3837.

[28780] Kawashima, K.; Shikama, H.; Imoto, K.; Izawa, M.; Naruke, T.; Okabayashi,
[28781] K.; Nishimura, S.: Close correlation between restriction fragment
[28782] length polymorphism of the L-MYC gene and metastasis of human lung
[28783] cancer to the lymph nodes and other organs. Proc. Nat. Acad. Sci. 85:
[28784] 2353-2356, 1988.

[28785] 3838.

[28786] Kaye, F.; Battey, J.; Nau, M.; Brooks, B.; Seifter, E.; De Greve,
[28787] J.; Birrer, M.; Sausville, E.; Minna, J.: Structure and expres-sion

[28788] of the human L-myc gene reveal a complex pattern of alternative mRNA

[28789] processing. Molec. Cell. Biol. 8: 186-195, 1988.

[28790] 3839.

[28791] McBride, O. W.; Kirsch, I.; Hollis, G.; Nau, M.; Battey, J.; Minna,

[28792] J.: Human L-myc (MYCL) proto-oncogene is on chromosome 1p32.(Abstract) Cytogenet.

[28793] Cell Genet. 40: 694 only, 1985.

[28794] 3840.

[28795] Nau, M. M.; Brooks, B. J.; Battey, J.; Sausville, E.; Gazdar, A.

[28796] F.; Kirsch, I. R.; McBride, O. W.; Bertness, V.; Hollis, G. F.; Minna,

[28797] J. D.: L-myc, a new myc-related gene amplified and expressed in human

[28798] small cell lung cancer. Nature 318: 69-73, 1985.

[28799] 3841.

[28800] Rouleau, G. A.; Bazanowski, A.; Gusella, J. F.; Haines, J. L.:

[28801] A genetic map of chromosome 1: comparison of different data sets and

[28802] linkage programs. Genomics 7: 313-318, 1990.

[28803] 3842.

[28804] Speleman, F.; Van Camp, G.; Van Roy, N.: Reassignment of MYCL1

[28805] to human chromosome 1p34.3 by fluorescence in situ hybridization. Cytogenet.

[28806] Cell Genet. 72: 189-190, 1996.

[28807] 3843.

[28808] Van Roy, N.; Cheng, N. C.; Laureys, G.; Opdenakker, G.; Versteeg,

[28809] R.; Speleman, F.: Molecular cytogenetic analysis of 1;17 translocations

[28810] in neuroblastoma. Europ. J. Cancer 31A: 530-535, 1995.

[28811] 3844.

[28812] Bottaro, D. P.; Rubin, J. S.; Faletto, D. L.; Chan, A. M.-L.; Kmiecik,

[28813] T. E.; Vande Woude, G. F.; Aaronson, S. A.: Identification of the

[28814] hepatocyte growth factor receptor as the c-met proto-oncogene product. Science 251:

[28815] 802-804, 1991.

[28816] 3845.

[28817] Cooper, C. S.: The met oncogene: from detection by transfection

[28818] to transmembrane receptor for hepatocyte growth factor. Oncogene 7:

[28819] 3-7, 1992.

[28820] 3846.

[28821] Cooper, C. S.; Park, M.; Blair, D. G.; Tainsky, M. A.; Hueb-ner,

[28822] K.; Croce, C. M.; Vande Woude, G. F.: Molecular cloning of a new

[28823] transforming gene from a chemically transformed human cell line. Nature 311:

[28824] 29-33, 1984.

[28825] 3847.

[28826] Dean, M.; Kozak, C.; Robbins, J.; Callahan, R.; O'Brien, S.; Vande

[28827] Woude, G. F.: Chromosomal localization of the met proto-oncogene

[28828] in the mouse and cat genome. Genomics 1: 167-173, 1987.

[28829] 3848.

[28830] Dean, M.; Park, M.; Le Beau, M. M.; Robins, T. S.; Diaz, M. O.;

[28831] Rowley, J. D.; Blair, D. G.; Vande Woude, G. F.: The human met oncogene

[28832] is related to the tyrosine kinase oncogenes. Nature 318: 385-388,

[28833] 1985.

[28834] 3849.

[28835] Giordano, S.; Bardelli, A.; Zhen, Z.; Menard, S.; Ponzetto, C.;

[28836] Comoglio, P. M.: A point mutation in the MET oncogene abrogates metastasis

[28837] without affecting transformation. Proc. Nat. Acad. Sci. 94: 13868-13872,

[28838] 1997.

[28839] 3850.

[28840] Giordano, S.; Corso, S.; Conrotto, P.; Artigiani, S.; Gilestro,

[28841] G.; Barberis, D.; Tamagnone, L.; Comoglio, P. M.: The Semaphorin

[28842] 4D receptor controls invasive growth by coupling with

Met. Nature
[28843] Cell Biol. 4: 720-724, 2002.

[28844] 3851.

[28845] Jeffers, M.; Fiscella, M.; Webb, C. P.; Anver, M.; Koochekpour,
[28846] S.; Vande Woude, G. F.: The mutationally activated Met receptor mediates
[28847] motility and metastasis. Proc. Nat. Acad. Sci. 95: 14417-14422,
[28848] 1998.

[28849] 3852.

[28850] Jeffers, M.; Schmidt, L.; Nakaigawa, N.; Webb, C. P.; Weirich,
[28851] G.; Kishida, T.; Zbar, B.; Vande Woude, G. F.: Activating mutations
[28852] for the Met tyrosine receptor in human cancer. Proc. Nat. Acad. Sci. 94:
[28853] 11445-11450, 1997.

[28854] 3853.

[28855] Kovacs, G.: Molecular cytogenetics of renal cell tumors. Adv.

[28856] Cancer Res. 62 89-124, 1993.

[28857] 3854.

[28858] Maina, F.; Casagranda, F.; Audero, E.; Simeone, A.; Co-
moglio,

[28859] P. M.; Klein, R.; Ponzetto, C.: Uncoupling of Grb2 from the
Met receptor

[28860] in vivo reveals complex roles in muscle development. Cell
87: 531-542,

[28861] 1996.

[28862] 3855.

[28863] Maina, F.; Pante, G.; Helmbacher, F.; Andres, R.; Porthin,
A.;

[28864] Davies, A. M.; Ponzetto, C.; Klein, R.: Coupling Met to spe-
cific

[28865] pathways results in distinct developmental outcomes.
Molec. Cell 7:

[28866] 1293-1306, 2001.

[28867] 3856.

[28868] Park, M.; Dean, M.; Kaul, K.; Braun, M. J.; Gonda, M. A.;
Vande

[28869] Woude, G.: Sequence of MET protooncogene cDNA has

features characteristic
[28870] of the tyrosine kinase family of growth-factor receptors. Proc. Nat.
[28871] Acad. Sci. 84: 6379-6383, 1987.

[28872] 3857.

[28873] Schmidt, L.; Duh, F.-M.; Chen, F.; Kishida, T.; Glenn, G.; Choyke,
[28874] P.; Scherer, S. W.; Zhuang, Z.; Lubensky, I.; Dean, M.; Al-likmets,
[28875] R.; Chidambaram, A.; and 23 others: Germline and so-matic mutations
[28876] in the tyrosine kinase domain of the MET proto-oncogene in papillary
[28877] renal carcinoma. Nature Genet. 16: 68-73, 1997.

[28878] 3858.

[28879] Park, W. S.; Dong, S. M.; Kim, S. Y.; Na, E. Y.; Shin, M. S.;

[28880] Pi, J. H.; Kim, B. J.; Bae, J. H.; Hong, Y. K.; Lee, K. S.; Lee, S.

[28881] H.; Yoo, N. J.; Jang, J. J.; Pack, S.; Zhuang, Z.; Schmidt, L.; Zbar,
[28882] B.; Lee, J. Y.: Somatic mutations in the kinase domain of the Met/hepatocyte

[28883] growth factor receptor gene in childhood hepatocellular carcinomas. Cancer
[28884] Res. 59: 307-310, 1999.

[28885] 3859.

[28886] Zbar, B.; Glenn, G.; Lubensky, I.; Choyke, P.; Walther, M. M.;
[28887] Magnusson, G.; Bergerheim, U. S. R.; Pettersson, S.; Amin, M.; Hurley,
[28888] K.; Linehan, W. M.: Hereditary papillary renal cell carci-noma: clinical
[28889] studies in 10 families. J. Urol. 153: 907-912, 1995.

[28890] 3860.

[28891] Zhuang, Z.; Park, W.-S.; Pack, S.; Schmidt, L.; Vortmeyer, A.
[28892] O.; Pak, E.; Pham, T.; Weil, R. J.; Candidus, S.; Lubensky, I. A.;
[28893] Linehan, W. M.; Zbar, B.; Weirich, G.: Trisomy 7-harbouring non-random
[28894] duplication of the mutant MET allele in hereditary papillary renal
[28895] carcinomas. Nature Genet. 20: 66-69, 1998.

[28896] 3861.

[28897] Akiyama, T.; Sudo, C.; Ogawara, H.; Toyoshima, K.; Ya-
mamoto, T.

[28898] : The product of the human c-erbB-2 gene: a
185-kilodalton glycoprotein

[28899] with tyrosine kinase activity. Science 232: 1644-1646,
1986.

[28900] 3862.

[28901] Ameyaw, M.-M.; Tayeb, M.; Thornton, N.; Folayan, G.;
Tariq, M.;

[28902] Mobarek, A.; Evans, D. A. P.; Ofori-Adjei, D.; McLeod, H.
L.: Ethnic

[28903] variation in the HER-2 codon 655 genetic polymorphism
previously associated

[28904] with breast cancer. J. Hum. Genet. 47: 172-175, 2002.

[28905] 3863.

[28906] Chan, J. Y. C.; Lerman, M. I.; Prabhakar, B. S.; Isozaki, O.;
Santisteban,

[28907] P.; Kuppers, R. C.; Oates, E. L.; Notkins, A. L.; Kohn, L. D.:
Cloning

[28908] and characterization of a cDNA that encodes a 70-kDa
novel human thyroid

[28909] autoantigen. J. Biol. Chem. 264: 3651-3654, 1989.

[28910] 3864.

[28911] Goedecke, W.; Eijpe, M.; Offenberg, H. H.; van Aalderen, M.; Heyting,

[28912] C.: Mre11 and Ku70 interact in somatic cells, but are differentially

[28913] expressed in early meiosis. Nature Genet. 23: 194-198, 1999.

[28914] 3865.

[28915] Koike, M.; Matsuda, Y.; Mimori, T.; Harada, Y.-N.; Shiomi, N.;

[28916] Shiomi, T.: Chromosomal localization of the mouse and rat DNA double-strand

[28917] break repair genes Ku p70 and Ku p80/XRCC5 and their mRNA expression

[28918] in various mouse tissues. Genomics 38: 38-44, 1996.

[28919] 3866.

[28920] Takata, M.; Sasaki, M. S.; Sonoda, E.; Morrison, C.; Hashimoto,

[28921] M.; Utsumi, H.; Yamaguchi-Iwai, Y.; Shinohara, A.; Takeda, S.: Homologous

[28922] recombination and non-homologous end-joining path-
ways of DNA double-strand
[28923] break repair have overlapping roles in the maintenance of
chromosomal
[28924] integrity in vertebrate cells. EMBO J. 17: 5497-5508,
1998.
[28925] 3867.

[28926] Tuteja, N.; Tuteja, R.; Ochem, A.; Taneja, P.; Huang, N. W.;
Simoncsits,
[28927] A.; Susic, S.; Rahman, K.; Marusic, L.; Chen, J.; Zhang, J.;
Wang,
[28928] S.; Pongor, S.; Falaschi, A.: Human DNA helicase II: a novel
DNA
[28929] unwinding enzyme identified as the Ku autoantigen. EMBO
J. 13: 4991-5001,
[28930] 1994.

[28931] 3868.

[28932] Walker, J. R.; Corpina, R. A.; Goldberg, J.: Structure of the

[28933] Ku heterodimer bound to DNA and its implications for
double-strand
[28934] break repair. Nature 412: 607-614, 2001.

[28935] 3869.

[28936] Balestrazzi, P.; Baeteman, M. A.; Mattei, M. G.; Mattei, J. F.

[28937] : Franceschetti syndrome in a child with a de novo balanced translocation

[28938] (5;13)(q11;p11) and significant decrease of hexosaminidase B. Hum.

[28939] Genet. 64: 305-308, 1983.

[28940] 3870.

[28941] Dixon, M. J.: Treacher Collins syndrome. Hum. Molec. Genet. 1996:

[28942] 1391-1396, 1996.

[28943] 3871.

[28944] Dixon, M. J.; Dixon, J.; Houseal, T.; Bhatt, M.; Ward, D. C.; Klinger,

[28945] K.; Landes, G. M.: Narrowing the position of the Treacher Collins

[28946] syndrome locus to a small interval between three new microsatellite

[28947] markers at 5q32-33.1. Am. J. Hum. Genet. 52: 907-914, 1993.

[28948] 3872.

[28949] Dixon, M. J.; Dixon, J.; Raskova, D.; Le Beau, M. M.; Williamson,
[28950] R.; Klinger, K.; Landes, G. M.: Genetic and physical mapping of the
[28951] Treacher Collins syndrome locus: refinement of the localization to
[28952] chromosome 5q32-33.2. Hum. Molec. Genet. 1: 249-253, 1992.
[28953] 3873.

[28954] Edery, P.; Manach, Y.; Le Merrer, M.; Till, M.; Vignal, A.; Lyonnet,
[28955] S.; Munnich, A.: Apparent genetic homogeneity of the Treacher Collins-Franceschetti
[28956] syndrome. Am. J. Med. Genet. 52: 174-177, 1994.

[28957] 3874.

[28958] Arakawa, T.; Narisawa, K.; Tanno, K.; Ohara, K.; Higashi, O.; Honda,
[28959] Y.; Tamura, T.; Wada, Y.; Mizuno, T.; Hayashi, T.; Hirooka, Y.; Ohno,
[28960] T.; Ikeda, M.: Megaloblastic anemia and mental retardation associated
[28961] with hyperfolic-acidemia: probably due to N-

5-methyltetrahydrofolate
[28962] transferase deficiency. Tohoku J. Exp. Med. 93: 1-22, 1967.
[28963] 3875.
[28964] Chen, L. H.; Liu, M.-L.; Hwang, H.-Y.; Chen, L.-S.; Koren-berg,
[28965] J.; Shane, B.: Human methionine synthase: cDNA cloning, gene localization,
[28966] and expression. J. Biol. Chem. 272: 3628, and 3634, 1997.
[28967] 3876.
[28968] Christensen, B.; Arbour, L.; Tran, P.; Leclerc, D.; Sab-baghian,
[28969] N.; Platt, R.; Gilfix, B. M.; Rosenblatt, D. S.; Gravel, R. A.; Forbes,
[28970] P.; Rozen, R.: Genetic polymorphisms in methylenete-trahydrofolate
[28971] reductase and methionine synthase, folate levels in red blood cells,
[28972] and risk of neural tube defects. Am. J. Med. Genet. 84: 151-157,
[28973] 1999.

[28974] 3877.

[28975] Gulati, S.; Baker, P.; Li, Y. N.; Fowler, B.; Kruger, W.; Brody,

[28976] L. C.; Banerjee, R.: Defects in human methionine synthase in cblG

[28977] patients. Hum. Molec. Genet. 5: 1859-1865, 1996.

[28978] 3878.

[28979] Kvittingen, E. A.; Spangen, S.; Lindemans, J.; Fowler, B.: Methionine

[28980] synthase deficiency without megaloblastic anaemia. Eu-rop. J. Pediat. 156:

[28981] 925-930, 1997.

[28982] 3879.

[28983] Leclerc, D.; Campeau, E.; Goyette, P.; Adjalla, C. E.; Chris-tensen,

[28984] B.; Ross, M.; Eydoux, P.; Rosenblatt, D. S.; Rozen, R.; Gravel, R.

[28985] A.: Human methionine synthase: cDNA cloning and identi-fication of

[28986] mutations in patients of the cblG complementation group of folate/cobalamin

[28987] disorders. Hum. Molec. Genet. 5: 1867-1874, 1996.

[28988] 3880.

[28989] Li, Y. N.; Gulati, S.; Baker, P. J.; Brody, L. C.; Banerjee, R.;

[28990] Kruger, W. D.: Cloning, mapping and RNA analysis of the human methionine

[28991] synthase gene. Hum. Molec. Genet. 5: 1851-1858, 1996.

[28992] 3881.

[28993] Mellman, I. S.; Lin, P.-F.; Ruddle, F. H.; Rosenberg, L. E.: Genetic

[28994] control of cobalamin binding in normal and mutant cells: assignment

[28995] of the gene for 5-methyltetrahydrofolate: L-homocysteine S-methyltransferase

[28996] to human chromosome 1. Proc. Nat. Acad. Sci. 76: 405-409, 1979.

[28997] 3882.

[28998] Mudd, S. H.: Cobalamin-responsive genetic disorders.In: Shimazono,

[28999] N.; Arakawa, T. (eds.): Nutritional deficiency secondary to inborn

[29000] errors of metabolism its relationship to physical and mental development.

[29001] Ichikawa: Jap. Panel Maln. U.S.-Japan Coop. Med. Sci. Program 1977.

[29002] 3883.

[29003] Paz, M. F.; Avila, S.; Fraga, M. F.; Pollan, M.; Capella, G.;

[29004] Peinado, M. A.; Sanchez-Cespedes, M.; Herman, J. G.; Esteller, M.

[29005] : Germ-line variants in methyl-group metabolism genes and susceptibility

[29006] to DNA methylation in normal tissues and human primary tumors. Cancer

[29007] Res. 62: 4519-4524, 2002.

[29008] 3884.

[29009] Van Someren, H.; Westerveld, A.; Hagemeijer, A.; Mees, J. R.;

[29010] Meera Khan, P.; Zaalberg, O. B.: Human antigen and enzyme markers

[29011] in man-Chinese hamster somatic cell hybrids: evidence for synteny

[29012] between the HL-A, PGM-3, ME-1, and IPO-B loci. Proc. Nat. Acad. Sci. 71:

[29013] 962-965, 1974.

[29014] 3885.

[29015] Jakoi, E. R.; Brown, A. L.; Ho, Y.-S.; Snyderman, R.: Molec-
ular
[29016] cloning of the cDNA for ligatin. J. Cell Sci. 93: 227-232,
1989.
[29017] 3886.

[29018] Malnar-Dragojevic, D.; Trachtulec, Z.; Vincek, V.: Assign-
ment
[29019] of the mouse ligatin gene (Lgtn) to chromosome 1F by
fluorescence
[29020] in situ hybridization. Genomics 40: 192-193, 1997.

[29021] 3887.

[29022] Dombroski, B. A.; Mathias, S. L.; Nanthakumar, E.; Scott,
A. F.;
[29023] Kazazian, H. H., Jr.: Isolation of an active human trans-
posable element. Science 254:
[29024] 1805-1808, 1991.

[29025] 3888.

[29026] Longy, M.; Coulon, V.; Duboue, B.; David, A.; Larregue, M.;
Eng,
[29027] C.; Amati, P.; Kraimps, J.-L.; Bottani, A.; Lacombe, D.;

Bonneau,
[29028] D.: Mutations of PTEN in patients with Bannayan-Riley-Ruvalcaba phenotype. J.
[29029] Med. Genet. 35: 886-889, 1998.

[29030] 3889.

[29031] Marsh, D. J.; Dahia, P. L. M.; Zheng, Z.; Liaw, D.; Parsons, R.;
[29032] Gorlin, R. J.; Eng, C.: Germline mutations in PTEN are present in
[29033] Bannayan-Zonana syndrome. (Letter) Nature Genet. 16: 333-334, 1997.
[29034] 3890.

[29035] Marsh, D. J.; Kum, J. B.; Lunetta, K. L.; Bennett, M. J.; Gorlin,
[29036] R. J.; Ahmed, S. F.; Bodurtha, J.; Crowe, C.; Curtis, M. A.; Dasouki,
[29037] M.; Dunn, T.; Feit, H.; and 20 others: PTEN mutation spectrum and
[29038] genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba syndrome
[29039] suggest a single entity with Cowden syndrome. Hum. Molec. Genet. 8:

[29040] 1461-1472, 1999.

[29041] 3891.

[29042] Kurt-Jones, E. A.; Popova, L.; Kwinn, L.; Haynes, L. M.; Jones,

[29043] L. P.; Tripp, R. A.; Walsh, E. E.; Freeman, M. W.; Golenbock, D. T.;

[29044] Anderson, L. J.; Finberg, R. W.: Pattern recognition recep-tors TLR4

[29045] and CD14 mediate response to respiratory syncytial virus. Nature

[29046] Immun. 1: 398-401, 2000.

[29047] 3892.

[29048] Daikh, D. I.; Douglass, J. O.; Adelman, J. P.: Structure and expression

[29049] of the human motilin gene. DNA 8: 615-621, 1989.

[29050] 3893.

[29051] Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[29052] M.; Yano, H.; Seino, Y.; Fujita, J.; Yamada, Y.; Inagaki, N.; Takeda,

[29053] J.; Imura, H.; Bell, G. I.; Shows, T. B.: Localization of the human

[29054] motilin gene (MLN) to chromosome 6p21.2. (Abstract) Cytogenet. Cell

[29055] Genet. 51: 977 only, 1989.

[29056] 3894.

[29057] Feighner, S. D.; Tan, C. P.; McKee, K. K.; Palyha, O. C.; Hreniuk,

[29058] D. L.; Pong, S.-S.; Austin, C. P.; Figueroa, D.; MacNeil, D.; Cascieri,

[29059] M. A.; Nargund, R.; Bakshi, R.; Abramovitz, M.; Stocco, R.; Kargman,

[29060] S.; O'Neill, G.; Van Der Ploeg, L. H. T.; Evans, J.; Patchett, A.

[29061] A.; Smith, R. G.; Howard, A. D.: Receptor for motilin identified

[29062] in the human gastrointestinal system. Science 284: 2184-2188, 1999.

[29063] 3895.

[29064] Gasparini, P.; Grifa, A.; Savasta, S.; Merlo, I.; Bisceglia, L.;

[29065] Totaro, A.; Zelante, L.: The motilin gene: subregional localisation,

[29066] tissue expression, DNA polymorphisms and exclusion as a candidate

[29067] gene for the HLA-associated immotile cilia syndrome. Hum. Genet. 94:
[29068] 671-674, 1994.

[29069] 3896.

[29070] Isnard, P.; Depetris, D.; Mattei, M.-G.; Ferrier, P.; Djabali,

[29071] M.: cDNA cloning, expression and chromosomal localization of the
[29072] murine AF-4 gene involved in human leukemia. Mammalian Genome 9:
[29073] 1065-1068, 1998.

[29074] 3897.

[29075] Lovett, B. D.; Lo Nigro, L.; Rappaport, E. F.; Blair, I. A.; Osheroff,
[29076] N.; Zheng, N.; Megonigal, M. D.; Williams, W. R.; Nowell, P. C.; Felix,
[29077] C. A.: Near-precise interchromosomal recombination and functional
[29078] DNA topoisomerase II cleavage sites at MLL and AF-4 genomic breakpoints
[29079] in treatment-related acute lymphoblastic leukemia with t(4;11) translocation. Proc.

[29080] Nat. Acad. Sci. 98: 9802-9807, 2001.

[29081] 3898.

[29082] Uckun, F. M.; Herman-Hatten, K.; Crotty, M.-L.; Sensel, M. G.;

[29083] Sather, H. N.; Tuel-Ahlgren, L.; Sarquis, M. B.; Bostrom, B.; Nachman,

[29084] J. B.; Steinherz, P. G.; Gaynon, P. S.; Heerema, N.: Clinical significance

[29085] of MLL-AF4 fusion transcript expression in the absence of a cytogenetically

[29086] detectable t(4;11)(q21;q23) chromosomal translocation. Blood 92:

[29087] 810-821, 1998.

[29088] 3899.

[29089] Prasad, R.; Gu, Y.; Alder, H.; Nakamura, T.; Canaani, O.; Saito,

[29090] H.; Huebner, K.; Gale, R. P.; Nowell, P. C.; Kuriyama, K.; Miyazaki,

[29091] Y.; Croce, C. M.; Canaani, E.: Cloning of the ALL-1 fusion partner,

[29092] the AF-6 gene, involved in acute myeloid leukemias with the t(6;11)

[29093] chromosome translocation. Cancer Res. 53: 5624-5628, 1993.

[29094] 3900.

[29095] Saha, V.; Lillington, D. M.; Shelling, A. N.; Chaplin, T.; Yaspo,

[29096] M.-L.; Ganesan, T. S.; Young, B. D.: AF6 gene on chromosome band

[29097] 6q27 maps distal to the minimal region of deletion in epithelial ovarian

[29098] cancer. Genes Chromosomes Cancer 14: 220-222, 1995.

[29099] 3901.

[29100] Gerling, I. C.; Karlsen, A. E.; Chapman, H. D.; Andersen, H. U.;

[29101] Boel, E.; Cunningham, J. M.; Nerup, J.; Leiter, E. H.: The inducible

[29102] nitric oxide synthase gene, Nos2, maps to mouse chromosome 11. Mammalian

[29103] Genome 5: 318-320, 1994.

[29104] 3902.

[29105] Van Cong, N.; Moullec, J.: Linkage probable entre les groupes

[29106] de phosphatase acide des globules rouges et le systeme Lewis. Ann.

[29107] Genet. 14: 121-125, 1971.

[29108] 3903.

[29109] Carmeliet, P.; Moons, L.; Lijnen, R.; Janssens, S.; Lupu, F.; Collen,

[29110] D.; Gerard, R. D.: Inhibitory role of plasminogen activator inhibitor-1

[29111] in arterial wound healing and neointima formation: a gene targeting

[29112] and gene transfer study in mice. Circulation 96: 3180-3191, 1997.

[29113] 3904.

[29114] Crandall, D. L.; Busler, D. E.; McHendry-Rinde, B.; Groel-ing, T.

[29115] M.; Kral, J. G.: Autocrine regulation of human preadipocyte migration

[29116] by plasminogen activator inhibitor-1. J. Clin. Endrocr. Metab. 85:

[29117] 2609-2614, 2000.

[29118] 3905.

[29119] Dawson, S. J.; Wiman, B.; Hamsten, A.; Green, F.; Humphries, S.;

[29120] Henney, A. M.: The two allele sequences of a common polymorphism

[29121] in the promoter of the plasminogen activator inhibitor-1 (PAI-1) gene

[29122] respond differently to interleukin-1 in HepG2 cells. J. Biol. Chem. 268:

[29123] 10739-10745, 1993.

[29124] 3906.

[29125] Engesser, L.; Brommer, E. J. P.; Kluft, C.; Briet, E.: Elevated

[29126] plasminogen activator inhibitor (PAI), a cause of thrombophilia? A

[29127] study in 203 patients with familial or sporadic venous thrombophilia. Thromb.

[29128] Haemost. 62: 673-680, 1989.

[29129] 3907.

[29130] Erickson, L. A.; Fici, G. J.; Lund, J. E.; Boyle, T. P.; Polites,

[29131] H. G.; Marotti, K. R.: Development of venous occlusions in mice transgenic

[29132] for the plasminogen activator inhibitor-1 gene. Nature

346: 74-76,

[29133] 1990.

[29134] 3908.

[29135] Eriksson, P.; Kallin, B.; van t'Hooft, F. M.; Bavhenolm. P.; Hamsten,

[29136] A.: Allelic-specific increase in basal transcription of the plasminogen-activator

[29137] inhibitor 1 gene is associated with myocardial infarction. Proc.

[29138] Nat. Acad. Sci. 92: 1851-1855, 1995.

[29139] 3909.

[29140] Fay, W. P.; Parker, A. C.; Condrey, L. R.; Shapiro, A. D.: Human

[29141] plasminogen activator inhibitor-1 (PAI-1) deficiency: characterization

[29142] of a large kindred with a null mutation in the PAI-1 gene. Blood 90:

[29143] 204-208, 1997.

[29144] 3910.

[29145] Fay, W. P.; Shapiro, A. D.; Shih, J. L.; Schleef, R. R.; Ginsburg,

[29146] D.: Complete deficiency of plasminogen-activator inhibitor type I

[29147] due to a frameshift mutation. New Eng. J. Med. 327: 1729-1733, 1992.

[29148] 3911.

[29149] Ginsburg, D.; Zeheb, R.; Yang, A. Y.; Rafferty, U. M.; Andreasen,

[29150] P. A.; Nielsen, L.; Dano, K.; Lebo, R. V.; Gelehrter, T. D.: cDNA

[29151] cloning of human plasminogen activator-inhibitor from endothelial

[29152] cells. J. Clin. Invest. 78: 1673-1680, 1986.

[29153] 3912.

[29154] Glueck, C. J.; Glueck, H. I.; Mieczkowski, L.; Tracy, T.; Speirs,

[29155] J.; Stroop, D.: Familial high plasminogen activator inhibitor with

[29156] hypofibrinolysis, a new pathophysiologic cause of osteonecrosis? Thromb.

[29157] Haemost. 69: 460-465, 1993.

[29158] 3913.

[29159] Halleux, C. M.; Declerck, P. J.; Tran, S. L.; Detry, R.; Brichard,

[29160] S. M.: Hormonal control of plasminogen activator inhibitor-1 gene

[29161] expression and production in human adipose tissue: stimulation by

[29162] glucocorticoids and inhibition by catecholamines. J. Clin. Endocr.

[29163] Metab. 84: 4097-4105, 1999.

[29164] 3914.

[29165] Hermans, P. W. M.; Hibberd, M. L.; Booy, R.; Daramola, O.; Hazelzet,

[29166] J. A.; de Groot, R.; Levin, M.; Meningococcal Research Group: 4G/5G

[29167] promoter polymorphism in the plasminogen-activator-inhibitor-1 gene

[29168] and outcome of meningococcal disease. Lancet 354: 556-560, 1999.

[29169] 3915.

[29170] Klinger, K. W.; Winqvist, R.; Andreasen, P. A.; Stuart, N.; Stanislovitis,

[29171] P.; Schwartz, R.; Doherty, R.; Alitalo, K.; Blasi, F.; Dano, K.:

[29172] Regional mapping of the plasminogen activator inhibitor type 1 gene

[29173] to 7q21.3-q22 and assignment to the CF linkage group. (Abstract) Cytogenet.

[29174] Cell Genet. 46: 640 only, 1987.

[29175] 3916.

[29176] Klinger, K. W.; Winqvist, R.; Andreasen, P. A.; Stuart, N.; Stanislovitis,

[29177] P.; Watkins, P.; Schwartz, R.; Doherty, R.; Alitalo, K.; Blasi, F.;

[29178] Dano, K.: Assignment of the human plasminogen activator inhibitor

[29179] type 1 (PAI-1) gene to 7q21.3-q22 and genetic linkage to markers on

[29180] chromosome 7. (Abstract) Am. J. Hum. Genet. 41: A172 only, 1987.

[29181] 3917.

[29182] Klinger, K. W.; Winqvist, R.; Riccio, A.; Andreasen, P. A.; Sartorio,

[29183] R.; Nielsen, L. S.; Stuart, N.; Stanislovitis, P.; Watkins, P.; Douglas,

[29184] R.; Grzeschik, K.-H.; Alitalo, K.; Blasi, F.; Dano, K.: Plas-

minogen
[29185] activator inhibitor type 1 gene is located at region q21.3-q22 of
[29186] chromosome 7 and genetically linked with cystic fibrosis. Proc. Nat.
[29187] Acad. Sci. 84: 8548-8552, 1987.

[29188] 3918.

[29189] Kohler, H. P.; Grant, P. J.: Plasminogen-activator inhibitor

[29190] type 1 and coronary artery disease. New Eng. J. Med. 342: 1792-1801,
[29191] 2000.

[29192] 3919.

[29193] Loskutoff, D. J.; Linders, M.; Keijer, J.; Veerman, H.; van Heerikhuizen,
[29194] H.; Pannekoek, H.: Structure of the human plasminogen activator inhibitor
[29195] 1 gene: nonrandom distribution of introns. Biochemistry 26: 3763-3768,
[29196] 1987.

[29197] 3920.

[29198] Margaglione, M.; Cappucci, G.; Colaizzo, D.; Giuliani, N.;

Vecchione,
[29199] G.; Grandone, E.; Pennelli, O.; Di Minno, G.: The PAI-1 gene locus
[29200] 4G/5G polymorphism is associated with a family history of coronary
[29201] artery disease. Arterioscler. Thromb. Vasc. Biol. 18: 152-156, 1998.
[29202] 3921.

[29203] Nilsson, I. M.; Ljungner, H.; Tengborn, L.: Two different mechanisms
[29204] in patients with venous thrombosis and defective fibrinol-ysis: low
[29205] concentration of plasminogen activator or increased con-centration
[29206] of plasminogen activator inhibitor. Brit. Med. J. 290: 1453-1456,
[29207] 1985.

[29208] 3922.

[29209] Wakita, Y.; Narahara, K.; Takahashi, Y.; Kikkawa, K.; Kimura,
[29210] S.; Oda, M.; Kimoto, H.: Duplication of 2p25: confirmation of the

[29211] assignment of soluble acid phosphatase (ACP1) locus to 2p25. Hum.

[29212] Genet. 71: 259-260, 1985.

[29213] 3923.

[29214] Weitkamp, L. R.; Janzen, M. K.; Guttormsen, S. A.; Ger-

showitz,

[29215] H.: Inherited pericentric inversion of chromosome number two: a linkage

[29216] study. Ann. Hum. Genet. 33: 53-59, 1969.

[29217] 3924.

[29218] Weitkamp, L. R.; Lovrien, E. W.; Olaisen, B.; Fenger, K.; Gedde-Dahl,

[29219] T., Jr.; Sorensen, S. A.; Conneally, P. M.; Bias, W. B.; Ott, J.:

[29220] Linkage relations of the loci for the MN blood group and red cell

[29221] phosphate. Birth Defects Orig. Art. Ser. 11(3): 276-280, 1975. Note:

[29222] Alternate: Cytogenet. Cell Genet. 14: 446-450, 1975...

[29223] 3925.

[29224] Wo, Y.-Y. P.; McCormack, A. L.; Shabanowitz, J.; Hunt, D. F.;

[29225] Davis, J. P.; Mitchell, G. L.; Van Etten, R. L.: Sequencing, cloning,
[29226] and expression of human red cell-type acid phosphatase, a cytoplasmic
[29227] phosphotyrosyl protein phosphatase. J. Biol. Chem. 267: 10856-10865,
[29228] 1992.

[29229] 3926.

[29230] Yoshihara, C. M.; Mohrenweiser, H. W.: Characterization of ACP1(TIC-1),
[29231] an electrophoretic variant of erythrocyte acid phosphatase restricted
[29232] to the Ticuna Indians of Central Amazonas. Am. J. Hum. Genet. 32:
[29233] 898-907, 1980.

[29234] 3927.

[29235] Allen, B. S.; Ketcham, C. M.; Roberts, R. M.; Nick, H. S.; Ostrer,
[29236] H.: Localization of the human type 5, tartrate-resistant acid phosphatase
[29237] gene by in situ hybridization. Genomics 4: 597-600, 1989.

[29238] 3928.

[29239] Bune, A. J.; Hayman, A. R.; Evans, M. J.; Cox, T. M.: Mice lacking

[29240] tartrate-resistant acid phosphatase (Acp 5) have disordered macrophage

[29241] inflammatory responses and reduced clearance of the pathogen, Staphylococcus

[29242] aureus. Immunology 102: 103-113, 2001.

[29243] 3929.

[29244] Grimes, R.; Reddy, S. V.; Leach, R. J.; Scarcez, T.; Roodman, G.

[29245] D.; Sakaguchi, A. Y.; Lalley, P. A.; Windle, J. J.: Assignment of

[29246] the mouse tartrate-resistant acid phosphatase gene (Acp5) to chromosome

[29247] 9. Genomics 15: 421-422, 1993.

[29248] 3930.

[29249] Hayman, A. R.; Jones, S. J.; Boyde, A.; Foster, D.; Colledge, W.

[29250] H.; Carlton, M. B.; Evans, M. J.; Cox, T. M.: Mice lacking tartrate-resistant

[29251] acid phosphatase (Acp 5) have disrupted endochondral ossification
[29252] and mild osteopetrosis. Development 122: 3151-3162, 1996.
[29253] 3931.
[29254] Ketcham, C. M.; Roberts, R. M.; Simmen, R. C.; Nick, H. S.: Molecular
[29255] cloning of the type 5, iron-containing, tartrate-resistant acid phosphatase
[29256] from human placenta. J. Biol. Chem. 264: 557-563, 1988.
[29257] 3932.
[29258] Leach, R. J.; Reus, B. E.; Hundley, J. E.; Johnson-Pais, T. L.;
[29259] Windle, J. J.: Confirmation of the assignment of the human tartrate-resistant
[29260] acid phosphatase gene (ACP5) to chromosome 19. Genomics 19: 180-181,
[29261] 1994.
[29262] 3933.
[29263] Lord, D. K.; Cross, N. C. P.; Bevilacqua, M. A.; Rider, S. H.;
[29264] Gorman, P. A.; Groves, A. V.; Moss, D. W.; Sheer, D.; Cox, T. M.:

[29265] Type 5 acid phosphatase: sequence, expression and chromosomal localization
[29266] of a differentiation-associated protein of the human macrophage. Europ.
[29267] J. Biochem. 189: 287-293, 1990.

[29268] 3934.

[29269] Griffiths, L. R.: Authors response to 'Chromosome 17 and the inducible
[29270] nitric oxide synthase gene in human essential hypertension' by Rutherford
[29271] et al., Human Genetics, published on-line September 2001. (Letter) Hum.
[29272] Genet. 110: 100-103, 2002.

[29273] 3935.

[29274] Jenkins, N. A.; Rothe, H.; Gilbert, D. J.; Copeland, N. G.; Kolb,
[29275] H.: Mapping of the gene for inducible nitric oxide (NO) synthase
[29276] of mouse macrophages to chromosome 11, close to Evi-2, nu, and Idd-4. Genomics 19:
[29277] 402-404, 1994.

[29278] 3936.

[29279] Johannesen, J.; Pie, A.; Pociot, F.; Kristiansen, O. P.; Karlsen,

[29280] A. E.; Nerup, J. The Danish Study Group of Diabetes in Childhood;

[29281] The Danish Insulin-Dependent Diabetes Mellitus Epidemiology and Genetics

[29282] Group: Linkage of the human inducible nitric oxide synthase gene

[29283] to type 1 diabetes. J. Clin. Endocr. Metab. 86: 2792-2796, 2001.

[29284] 3937.

[29285] Lowenstein, C. J.; Glatt, C. S.; Bredt, D. S.; Snyder, S. H.:

[29286] Cloned and expressed macrophage nitric oxide synthase contrasts with

[29287] the brain enzyme. Proc. Nat. Acad. Sci. 89: 6711-6715, 1992.

[29288] 3938.

[29289] Marsden, P. A.; Heng, H. H. Q.; Duff, C. L.; Shi, X.-M.; Tsui,

[29290] L.-C.; Hall, A. V.: Localization of the human gene for inducible

[29291] nitric oxide synthase (NOS2) to chromosome 17q11.2-q12. Genomics 19:
[29292] 183-185, 1994.

[29293] 3939.

[29294] Mehrabian, M.; Xia, Y.-R.; Wen, P.-Z.; Warden, C. H.; Herschman,
[29295] H. R.; Lusis, A. J.: Localization of murine macrophage inducible
[29296] nitric oxide synthase to mouse chromosome 11. Genomics 22: 646-647,
[29297] 1994.

[29298] 3940.

[29299] Morris, B. J.: Critique of 'Chromosome 17 and the inducible nitric
[29300] oxide synthase gene in human essential hypertension' by Rutherford
[29301] et al., Human Genetics, published online September 2001. (Letter) Hum.
[29302] Genet. 110: 98-99, 2002.

[29303] 3941.

[29304] Nathan, C.; Shiloh, M. U.: Reactive oxygen and nitrogen

intermediates
[29305] in the relationship between mammalian hosts and microbial pathogens. Proc.
[29306] Nat. Acad. Sci. 97: 8841-8848, 2000.

[29307] 3942.

[29308] Nicholson, S.; Bonecini-Almeida Mda, G; Lapa e Silva, J. R.; Nathan,
[29309] C.; Xie, Q. W.; Mumford, R.; Weidner, J. R.; Calaycay, J.; Geng, J.;
[29310] Boechat, N.; et al.; et al.: Inducible nitric oxide synthase in pulmonary
[29311] alveolar macrophages from patients with tuberculosis. J. Exp. Med. 183:
[29312] 2293-2302, 1996.

[29313] 3943.

[29314] Nozaki, Y.; Hasegawa, Y.; Ichiyama, S.; Nakashima, I.; Shimokata,
[29315] K.: Mechanism of nitric oxide-dependent killing of Mycobacterium
[29316] bovis BCG in human alveolar macrophages. Infect. Immun. 65: 3644-3647,
[29317] 1997.

[29318] 3944.

[29319] Vouldoukis, I.; Riveros-Moreno, V.; Dugas, B.; Ouaaz, F.; Becherel,

[29320] P.; Debre, P.; Moncada, S.; Mossalayi, M. D.: The killing of Leishmania

[29321] major by human macrophages is mediated by nitric oxide induced after

[29322] ligation of the Fc-epsilon-RII/CD23 surface antigen. Proc. Nat. Acad.

[29323] Sci. 92: 7804-7808, 1995.

[29324] 3945.

[29325] Xu, W.; Charles, I. G.; Liu, L.; Moncada, S.; Emson, P.: Molecular

[29326] cloning and structural organization of the human inducible nitric

[29327] oxide synthase gene (NOS2). Biochem. Biophys. Res. Commun. 219:

[29328] 784-788, 1996.

[29329] 3946.

[29330] Xu, W.; Charles, I. G.; Moncada, S.; Gorman, P.; Sheer, D.; Liu,

[29331] L.; Emson, P.: Mapping of the genes encoding human inducible and
[29332] endothelial nitric oxide synthase (NOS2 and NOS3) to the pericentric
[29333] region of chromosome 17 and to chromosome 7, respectively. Genomics 21:
[29334] 419-422, 1994.

[29335] 3947.

[29336] Bredt, D. S.; Hwang, P. M.; Glatt, C. E.; Lowenstein, C.; Reed,
[29337] R. R.; Snyder, S. H.: Cloned and expressed nitric oxide synthase
[29338] structurally resembles cytochrome P-450 reductase. Nature 351: 714-718,
[29339] 1991.

[29340] 3948.

[29341] Brenman, J. E.; Chao, D. S.; Xia, H.; Aldape, K.; Bredt, D. S.

[29342] : Nitric oxide synthase complexed with dystrophin and absent from
[29343] skeletal muscle sarcolemma in Duchenne muscular dystrophy. Cell 82:

[29344] 743-752, 1995.

[29345] 3949.

[29346] Burnett, A. L.; Lowenstein, C. J.; Bredt, D. S.; Chang, T. S. K.;

[29347] Snyder, S. H.: Nitric oxide: a physiologic mediator of penile erection. Science 257:

[29348] 401-403, 1992.

[29349] 3950.

[29350] Day, B. J.; Patel, M.; Calavetta, L.; Chang, L.-Y.; Stamler, J.

[29351] S.: A mechanism of paraquat toxicity involving nitric oxide synthase. Proc.

[29352] Nat. Acad. Sci. 96: 12760-12765, 1999.

[29353] 3951.

[29354] Deans, Z.; Dawson, S. J.; Xie, J.; Young, A. P.; Wallace, D.; Latchman,

[29355] D. S.: Differential regulation of the two neuronal nitric-oxide synthase

[29356] gene promoters by the Oct-2 transcription factor. J. Biol. Chem. 271:

[29357] 32153-32158, 1996.

[29358] 3952.

[29359] Gu, Z.; Kaul, M.; Yan, B.; Kridel, S. J.; Cui, J.; Strongin, A.;

[29360] Smith, J. W.; Liddington, R. C.; Lipton, S. A.: S-nitrosylation of

[29361] matrix metalloproteinases: signaling pathway to neuronal cell death. Science 2

[29362] 97: 1186-1190, 2002.

[29363] 3953.

[29364] Kharazia, V. N.; Schmidt, H. H. H. W.; Weinberg, R. J.: Type I

[29365] nitric oxide synthase fully accounts for NADPH-di-aphorase in rat striatum,

[29366] but not cortex. Neuroscience 62: 983-987, 1994.

[29367] 3954.

[29368] Kishimoto, J.; Spurr, N.; Liao, M.; Lizhi, L.; Emson, P.; Xu, W.

[29369] : Localization of brain nitric oxide synthase (NOS) to human chromosome

[29370] 12. Genomics 14: 802-804, 1992.

[29371] 3955.

[29372] Kuo, R. C.; Baxter, G. T.; Thompson, S. H.; Stricker, S. A.; Patton,
[29373] C.; Bonaventura, J.; Epel, D.: NO is necessary and sufficient for
[29374] egg activation at fertilization. Nature 406: 633-636, 2000.
[29375] 3956.

[29376] Lee, C. G. L.; Gregg, A. R.; O'Brien, W. E.: Localization of
[29377] the neuronal form of nitric oxide synthase to mouse chro-mosome 5. Mammalian
[29378] Genome 6: 56-57, 1995.

[29379] 3957.

[29380] Magee, T.; Fuentes, A. M.; Garban, H.; Rajavashisth, T.; Marquez,
[29381] D.; Rodriguez, J. A.; Rajfer, J.; Gonzalez-Cadavid, N. F.: Cloning
[29382] of a novel neuronal nitric oxide synthase expressed in pe-nis and lower
[29383] urinary tract. Biochem. Biophys. Res. Commun. 226: 145-151, 1996.
[29384] 3958.

[29385] Servidei, S.; Riepe, R.; Langston, C.; Tani, L. Y.; Bricker, J.

[29386] T.; Crisp Lindgren, N.; Travers, H.; Armstrong, D.; Di-
Mauro, S.:

[29387] Severe cardiopathy in branching enzyme deficiency. J. Pe-
diat. 111:

[29388] 51-56, 1987.

[29389] 3959.

[29390] Shin, Y. S.; Steiguber, H.; Klemm, P.; Endres, W.; Schwab,
O.;

[29391] Wolff, G.: Branching enzyme in erythrocytes: detection of
type IV

[29392] glycogenosis homozygotes and heterozygotes. J. Inherit.
Metab. Dis. 11

[29393] (suppl. 2): 252-254, 1988.

[29394] 3960.

[29395] Sidbury, J. B., Jr.; Mason, J.; Burns, W. B., Jr.; Ruebner, B.

[29396] H.: Type IV glycogenosis: report of a case proven by char-
acterization

[29397] of glycogen and studied at necropsy. Bull. Johns Hopkins
Hosp. 111:

[29398] 157-181, 1962.

[29399] 3961.

[29400] Tang, T. T.; Segura, A. D.; Chen, Y.-T.; Ricci, L. M.; Fran-
ciosi,

[29401] R. A.; Splaingard, M. L.; Lubinsky, M. S.: Neonatal hypoto-
nia and

[29402] cardiomyopathy secondary to type IV glycogenosis. Acta
Neuropath. 87:

[29403] 531-536, 1994.

[29404] 3962.

[29405] Thon, V. J.; Khalil, M.; Cannon, J. F.: Isolation of human
glycogen

[29406] branching enzyme cDNAs by screening complementation
in yeast. J.

[29407] Biol. Chem. 268: 7509-7513, 1993.

[29408] 3963.

[29409] Zellweger, H.; Mueller, S.; Ionasescu, V.; Schochet, S. S.;
McCormick,

[29410] W. F.: Glycogenosis. IV. A new cause of infantile hypoto-
nia. J.

[29411] Pediat. 80: 842-844, 1972.

[29412] 3964.

[29413] Abarbanel, J. M.; Potashnik, R.; Frisher, S.; Moses, S. W.; Osimani,
[29414] A.; Herishanu, Y.: Myophosphorylase deficiency: the course of an
[29415] unusual congenital myopathy. Neurology 37: 316-318, 1987.
[29416] 3965.

[29417] Bartram, C.; Edwards, R. H. T.; Clague, J.; Beynon, R. J.: McArdle's
[29418] disease: a nonsense mutation in exon 1 of the muscle glycogen phosphorylase
[29419] gene explains some but not all cases. Hum. Molec. Genet. 2: 1291-1293,
[29420] 1993.

[29421] 3966.

[29422] Bogusky, R. T.; Taylor, R. G.; Anderson, L. J.; Angelos, K. L.;
[29423] Lieberman, J. S.; Walsh, D. A.: McArdle's disease heterozy-gotes:
[29424] metabolic adaptation assessed using (31)P-nuclear mag-netic resonance. J.
[29425] Clin. Invest. 77: 1881-1887, 1986.

[29426] 3967.

[29427] Braakhekke, J. P.; de Bruin, M. I.; Stegeman, D. F.; Wevers, R.

[29428] A.; Binkhorst, R. A.; Joosten, E. M. G.: The second wind phenomenon

[29429] in McArdle's disease. Brain 109: 1087-1101, 1986.

[29430] 3968.

[29431] Bruno, C.; Lofberg, M.; Tamburino, L.; Jankala, H.; Hadji-georgiou,

[29432] G. M.; Andreu, A. L.; Shanske, S.; Somer, H.; DiMauro, S.: Molecular

[29433] characterization of McArdle's disease in two large Finnish families. J.

[29434] Neurol. Sci. 165: 121-125, 1999.

[29435] 3969.

[29436] Burke, J.; Hwang, P.; Anderson, L.; Lebo, R.; Gorin, F.; Flet-terick,

[29437] R.: Intron/exon structure of the human gene for the mus-cle isozyme

[29438] of glycogen phosphorylase. Protein 2: 177-187, 1987.

[29439] 3970.

[29440] Cerri, C. G.; Willner, J. H.: Phosphorylation of McArdle phosphorylase
[29441] induces activity. Proc. Nat. Acad. Sci. 78: 2688-2692, 1981.
[29442] 3971.

[29443] Chui, L. A.; Munsat, T. L.: Dominant inheritance of McArdle syndrome. Arch.
[29444] Neurol. 33: 636-641, 1976.

[29445] 3972.

[29446] Cochran, P.; Huges, R. R.; Buxton, P. H.; Yorke, R. A.: Myophosphorylase
[29447] deficiency (McArdle's disease) in two interrelated families. J. Neurol.
[29448] Neurosurg. Psychiat. 36: 217-224, 1973.

[29449] 3973.

[29450] Daegelen, D.; Gautron, S.; Mennecier, F.; Dreyfus, J.-C.; Kahn,
[29451] A.: Molecular heterogeneity of McArdle disease. J. Inherit. Metab.
[29452] Dis. 9: 287-290, 1986.

[29453] 3974.

[29454] Daegelen-Proux, D.; Kahn, A.; Marie, J.; Dreyfus, J.-C.: Research
[29455] on molecular mechanisms of McArdle's disease (muscle glycogen phosphorylase
[29456] deficiency): use of new protein mapping and immunological techniques. Ann.
[29457] Hum. Genet. 45: 113-120, 1981.

[29458] 3975.

[29459] Dawson, D. M.; Spong, F. L.; Harrington, J. F.: McArdle's disease:
[29460] lack of muscle phosphorylase. Ann. Intern. Med. 69: 229-236, 1968.
[29461] 3976.

[29462] DiMauro, S.; Hartlage, P. L.: Fatal infantile form of muscle
[29463] phosphorylase deficiency. Neurology 28: 1124-1129, 1978.
[29464] 3977.

[29465] Di Sant'Agnese, P. A.; Anderson, D. H.; Metcalf, K. M.: Glycogen
[29466] storage disease of the muscles. J. Pediat. 61: 438-442, 1962.

[29467] 3978.

[29468] Engel, W. K.; Eyerman, E. L.; Williams, H. E.: Late-onset type

[29469] of skeletal-muscle phosphorylase deficiency: a new familial variety

[29470] with completely and partially affected subjects. New Eng. J. Med. 268:

[29471] 135-137, 1963.

[29472] 3979.

[29473] Fernandez, R.; Navarro, C.; Andreu, A. L.; et al.: A novel missense

[29474] mutation (W797R) in the myophosphorylase gene in Spanish patients

[29475] with McArdle disease. Arch. Neurol. 57: 217-219, 2000.

[29476] 3980.

[29477] Gautron, S.; Daegelen, D.; Mennecier, F.; Dubocq, D.; Kahn, A.;

[29478] Dreyfus, J.-C.: Molecular mechanisms of McArdle's disease (muscle

[29479] glycogen phosphorylase deficiency). J. Clin. Invest. 79: 275-281,

[29480] 1987.

[29481] 3981.

[29482] Grunfeld, J.-P.; Ganeval, D.; Chanard, J.; Fardeau, M.; Dreyfus,

[29483] J.-C.: Acute renal failure in McArdle's disease: report of two cases. New

[29484] Eng. J. Med. 286: 1237-1241, 1972.

[29485] 3982.

[29486] Haller, R. G.; Dempsey, W. B.; Feit, H.; Cook, J. D.; Knochel,

[29487] J. P.: Low muscle levels of pyridoxine in McArdle's syn-drome. Am.

[29488] J. Med. 74: 217-220, 1983.

[29489] 3983.

[29490] Iwasaki, H.; Stewart, P. W.; Dilley, W. G.; Holt, M. S.; Stein-brueck,

[29491] T. D.; Wells, S. A., Jr.; Donis-Keller, H.: A minisatellite and a

[29492] microsatellite polymorphism within 1.5 kb at the human muscle glycogen

[29493] phosphorylase (PYGM) locus can be amplified by PCR and have combined

[29494] informativeness of PIC 0.95. Genomics 13: 7-15, 1992.

[29495] 3984.

[29496] Iyengar, S.; Kalinsky, H.; Weiss, S.; Korostishevsky, M.; Sadeh,

[29497] M.; Zhao, Y.; Kidd, K. K.; Bonne-Tamir, B.: Homozygosity by descent

[29498] for a rare mutation in the myophosphorylase gene is associated with

[29499] variable phenotypes in a Druze family with McArdle disease. J. Med.

[29500] Genet. 34: 391-394, 1997.

[29501] 3985.

[29502] Devlin, R.; Henderson, H.; Monsalve, V.; Brunzell, J.; Deeb, S.;

[29503] Hayden, M. R.: The molecular biology of hypertriglyceridemia: characterization

[29504] of mutations in patients with lipoprotein lipase deficiency.(Abstract) Am.

[29505] J. Hum. Genet. 45 (suppl.): A4, 1989.

[29506] 3986.

[29507] Pomponio, R. J.; Yamaguchi, A.; Arashima, S.; Hymes, J.;

Wolf,
[29508] B.: Mutation in a putative glycosylation site (N489T) of bi-
otinidase
[29509] in the only known Japanese child with biotinidase defi-
ciency. Molec.
[29510] Genet. Metab. 64: 152-154, 1998.

[29511] 3987.

[29512] Chynn, E. W.; Walton, D. S.; Hahn, L. B.; Dryja, T. P.: Norrie

[29513] disease: diagnosis of a simplex case by DNA analysis.
Arch. Ophthal. 114:
[29514] 1136-1138, 1996.

[29515] 3988.

[29516] Thomas, S. A.; Palmiter, R. D.: Impaired maternal behavior
in
[29517] mice lacking norepinephrine and epinephrine. Cell 91:
583-592, 1997.
[29518] 3989.

[29519] Thomas, S. A.; Palmiter, R. D.: Thermoregulatory and
metabolic
[29520] phenotypes of mice lacking noradrenaline and adrenaline.
Nature 387:

[29521] 94-97, 1997.

[29522] 3990.

[29523] Weinshilboum, R. M.: Catecholamine biochemical genetics in human

[29524] populations.In: Breakefield, X. O.: Neurogenetics: Genetic Approaches

[29525] to the Nervous System. New York: Elsevier/North Holland (pub.)

[29526] 1979. Pp. 257-282.

[29527] 3991.

[29528] Weinshilboum, R. M.; Schrott, H. G.; Raymond, F. A.; Weid-man,

[29529] W. H.; Elveback, L. R.: Inheritance of very low serum dopamine-beta-hydroxylase

[29530] activity. Am. J. Hum. Genet. 27: 573-585, 1975.

[29531] 3992.

[29532] Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Tran, L. D.:

[29533] Linkage of a gene regulating dopamine-beta-hydroxylase activity and

[29534] the ABO blood group locus. Am. J. Hum. Genet. 42: 160-166, 1988.

[29535] 3993.

[29536] Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Tran, L. D.:

[29537] Linkage of a gene regulating dopamine-beta-hydroxylase activity and

[29538] the ABO blood group locus. (Abstract) Am. J. Hum. Genet. 41: A191

[29539] only, 1987.

[29540] 3994.

[29541] Zabetian, C. P.; Anderson, G. M.; Buxbaum, S. G.; Elston, R. C.;

[29542] Ichinose, H.; Nagatsu, T.; Kim, K.-S.; Kim, C.-H.; Malison, R. T.;

[29543] Gelernter, J.; Cubells, J. F.: A quantitative-trait analysis of human

[29544] plasma-dopamine beta-hydroxylase activity: evidence for a major functional

[29545] polymorphism at the DBH locus. Am. J. Hum. Genet. 68: 515-522, 2001.

[29546] 3995.

[29547] Schulz, P. E.; Weiner, S. P.; Belmont, J. W.; Fishman, M. A.:

[29548] Basal ganglia calcifications in a case of biotinidase defi-

ciency. Neurology 38:
[29549] 1326-1328, 1988.

[29550] 3996.

[29551] Suormala, T.; Wick, H.; Bonjour, J.-P.; Baumgartner, E. R.: Intestinal
[29552] absorption and renal excretion of biotin in patients with biotinidase
[29553] deficiency. Europ. J. Pediat. 144: 21-26, 1985.

[29554] 3997.

[29555] Suormala, T. M.; Baumgartner, E. R.; Wick, H.; Scheibenre-iter,
[29556] S.; Schweitzer, S.: Comparison of patients with complete and partial
[29557] biotinidase deficiency: biochemical studies. J. Inherit. Metab. Dis. 13:
[29558] 76-92, 1990.

[29559] 3998.

[29560] Swango, K. L.; Demirkol, M.; Huner, G.; Pronicka, E.; Sykut-Cegielska,
[29561] J.; Schulze, A.; Wolf, B.: Partial biotinidase deficiency is usually

[29562] due to the D444H mutation in the biotinidase gene. Hum. Genet. 102:

[29563] 571-575, 1998.

[29564] 3999.

[29565] Taitz, L. S.; Green, A.; Strachan, I.; Bartlett, K.; Bennet, M.

[29566] : Biotinidase deficiency and the eye and ear. (Letter) Lancet II:

[29567] 918 only, 1983.

[29568] 4000.

[29569] Thoene, J.; Wolf, B.: Biotinidase deficiency in juvenile mul-
tiple
[29570] carboxylase deficiency. (Letter) Lancet II: 398 only, 1983.

[29571] 4001.

[29572] Thoene, J. G.; Lemons, R. M.; Borysko, K. Z.; Baker, H.: Ju-
venile
[29573] multiple carboxylase deficiency: evidence of impaired bi-
otin absorption.
[29574] (Abstract) Pediat. Res. 16: 179A only, 1982.

[29575] 4002.

[29576] Thuy, L. P.; Zielinska, B.; Zammarchi, E.; Pavari, E.;
Vierucci,
[29577] A.; Sweetman, F.; Sweetman, L.; Nyhan, W. L.: Multiple
carboxylase
[29578] deficiency due to deficiency of biotinidase. J. Neurogenet.
3: 357-363,
[29579] 1986.

[29580] 4003.

[29581] Tokatli, A.; Coskun, T.; Ozalp, I.; Gunay, M.: The major presenting

[29582] symptom in a biotinidase-deficient patient: laryngeal stridor. J.

[29583] Inherit. Metab. Dis. 15: 281-282, 1992.

[29584] 4004.

[29585] Wallace, S. J.: Biotinidase deficiency: presymptomatic treatment. Arch.

[29586] Dis. Child. 60: 574-575, 1985.

[29587] 4005.

[29588] Wastell, H. J.; Bartlett, K.; Dale, G.; Shein, A.: Biotinidase

[29589] deficiency: a survey of 10 cases. Arch. Dis. Child. 63: 1244-1249,

[29590] 1988.

[29591] 4006.

[29592] Weissbecker, K. A.; Nance, W. E.; Eaves, L. J.; Piussan, C.; Wolf,

[29593] B.: Statistical approaches for the detection of heterozygotes for

[29594] biotinidase deficiency. Am. J. Med. Genet. 39: 385-390, 1991.

[29595] 4007.

[29596] Wolf, B.; Grier, R. E.; Heard, G. S.: Hearing loss in bio-
tinidase
[29597] deficiency. (Letter) Lancet II: 1365-1366, 1983.

[29598] 4008.

[29599] Wolf, B.; Grier, R. E.; Parker, W. D., Jr.; Goodman, S. I.;
Allen,
[29600] R. J.: Deficient biotinidase activity in late-onset multiple
carboxylase
[29601] deficiency. (Letter) New Eng. J. Med. 308: 161, 1983.

[29602] 4009.

[29603] Wolf, B.; Grier, R. E.; Secor McVoy, J. R.; Heard, G. S.: Bio-
tinidase
[29604] deficiency: a novel vitamin recycling defect. J. Inherit.
Metab.
[29605] Dis. 8 (suppl. 1): 53-58, 1985.

[29606] 4010.

[29607] Wolf, B.; Heard, G. S.; Jefferson, L. G.; Proud, V. K.; Nance,

[29608] W. E.; Weissbecker, K. A.: Clinical findings in four children
with

[29609] biotinidase deficiency detected through a statewide neonatal screening

[29610] program. New Eng. J. Med. 313: 16-19, 1985.

[29611] 4011.

[29612] Wolf, B.; Heard, G. S.; Weissbecker, K. A.; Secor McVoy, J. R.;

[29613] Grier, R. E.; Leshner, R. T.: Biotinidase deficiency: initial clinical

[29614] features and rapid diagnosis. Ann. Neurol. 18: 614-617, 1985.

[29615] 4012.

[29616] Wolf, B.; Norrgard, K.; Pomponio, R. J.; Mock, D. M.; McVoy, J.

[29617] R. S.; Fleischhauer, K. Shapiro, S.; Blitzer, M. G.; Hymes, J.: Profound

[29618] biotinidase deficiency in two asymptomatic adults. Am. J. Med. Genet. 73:

[29619] 5-9, 1997.

[29620] 4013.

[29621] Aoki, Y.; Li, X.; Sakamoto, O.; Hiratsuka, M.; Akaishi, H.; Xu,

[29622] L.; Briones, P.; Suormala, T.; Baumgartner, E. R.; Suzuki, Y.; Narisawa,
[29623] K.: Identification and characterization of mutations in patients
[29624] with holocarboxylase synthetase deficiency. Hum. Genet. 104: 143-148,
[29625] 1999.

[29626] 4014.

[29627] Bartlett, K.; Ghneim, H. K.; Stirk, H.-J.; Wastell, H.: Enzyme
[29628] studies in biotin-responsive disorders. J. Inherit. Metab. Dis. 8
[29629] (suppl. 1): 46-52, 1985.

[29630] 4015.

[29631] Burri, B. J.; Sweetman, L.; Nyhan, W. L.: Mutant holocarboxylase
[29632] synthetase: evidence for the enzyme defect in early infantile biotin-responsive
[29633] multiple carboxylase deficiency. J. Clin. Invest. 68: 1491-1495,
[29634] 1981.

[29635] 4016.

[29636] Burri, B. J.; Sweetman, L.; Nyhan, W. L.: Heterogeneity of holocarboxylase
[29637] synthetase in patients with biotin-responsive multiple carboxylase
[29638] deficiency. Am. J. Hum. Genet. 37: 326-337, 1985.

[29639] 4017.

[29640] Dupuis, L.; Leon-Del-Rio, A.; Leclerc, D.; Campeau, E.; Sweetman,
[29641] L.; Saudubray, J. M.; Herman, G.; Gibson, K. M.; Gravel, R. A.: Clustering
[29642] of mutations in the biotin-binding region of holocarboxy-lase synthetase
[29643] in biotin-responsive multiple carboxylase deficiency. Hum. Molec.
[29644] Genet. 5: 1011-1016, 1996.

[29645] 4018.

[29646] Feldman, G. L.; Hsia, Y. E.; Wolf, B.: Biochemical character-ization
[29647] of biotin-responsive multiple carboxylase deficiency: het-erogeneity
[29648] within the bio genetic complementation group. Am. J. Hum. Genet. 33:

[29649] 692-701, 1981.

[29650] 4019.

[29651] Leon-Del-Rio, A.; Leclerc, D.; Akerman, B.; Wakamatsu, N.; Gravel,

[29652] R. A.: Isolation of a cDNA encoding human holocarboxy-lase synthetase

[29653] by functional complementation of a biotin auxotroph of Escherichia

[29654] coli. Proc. Nat. Acad. Sci. 92: 4626-4630, 1995.

[29655] 4020.

[29656] Mock, D. M.; deLorimer, A. A.; Liebman, W. M.; Sweetman, L.; Baker,

[29657] H.: Biotin deficiency: an unusual complication of par-enteral alimentation. New

[29658] Eng. J. Med. 304: 820-822, 1981.

[29659] 4021.

[29660] Morrone, A.; Malvagia, S.; Donati, M. A.; Funghini, S.; Ciani,

[29661] F.; Pela, I.; Boneh, A.; Peters, H.; Pasquini, E.; Zammarchi, E.:

[29662] Clinical findings and biochemical and molecular analysis

of four patients
[29663] with holocarboxylase synthetase deficiency. Am. J. Med. Genet. 111:
[29664] 10-18, 2002.

[29665] 4022.

[29666] Munnich, A.; Saudubray, J. M.; Carre, G.; Coude, F. X.; Ogier,
[29667] H.; Charpentier, C.; Frezal, J.: Defective biotin absorption in multiple
[29668] carboxylase deficiency. (Letter) Lancet II: 263, 1981.

[29669] 4023.

[29670] Narisawa, K.; Arai, N.; Igarashi, Y.; Satoh, T.; Tada, K.: Clinical
[29671] and biochemical findings on a child with multiple biotin-responsive
[29672] carboxylase deficiencies. J. Inherit. Metab. Dis. 5: 67-68, 1982.
[29673] 4024.

[29674] Packman, S.; Caswell, N.; Gonzalez-Rios, M. C.; Kadlecek, T.;
[29675] Cann, H.; Rassin, D.; McKay, C.: Acetyl CoA carboxylase in

cultured
[29676] fibroblasts: differential biotin dependence in the two types of biotin-responsive
[29677] multiple carboxylase deficiency. Am. J. Hum. Genet. 36: 80-92, 1984.
[29678] 4025.

[29679] Monros, E.; Molto, M. D.; Martinez, F.; Canizares, J.; Blanca,
[29680] J.; Vilchez, J. J.; Prieto, F.; de Frutos, R.; Palau, F.: Pheno-type
[29681] correlation and intergenerational dynamics of the Friedre-ich ataxia
[29682] GAA trinucleotide repeat. Am. J. Hum. Genet. 61: 101-110, 1997.
[29683] 4026.

[29684] Montermini, L.; Andermann, E.; Labuda, M.; Richter, A.; Pandolfo,
[29685] M.; Cavalcanti, F.; Pianese, L.; Iodice, L.; Farina, G.; Monti-celli,
[29686] A.; Turano, M.; Filla, A.; De Michele, G.; Cocozza, S.: The Friedreich
[29687] ataxia GAA triplet repeat: premutation and normal alleles.

Hum. Molec.
[29688] Genet. 6: 1261-1266, 1997.

[29689] 4027.

[29690] Ohshima, K.; Montermini, L.; Wells, R. D.; Pandolfo, M.: In-
hibitory
[29691] effects of expanded GAA-TTC triplet repeats from intron I
of the Friedreich
[29692] ataxia gene on transcription and replication in vivo. J. Biol.
Chem. 273:
[29693] 14588-14595, 1998.

[29694] 4028.

[29695] Pianese, L.; Cavalcanti, F.; De Michele, G.; Filla, A.; Cam-
panella,
[29696] G.; Calabrese, O.; Castaldo, I.; Monticelli, A.; Cocozza, S.:
The
[29697] effect of parental gender on the GAA dynamic mutation in
the FRDA
[29698] gene. (Letter) Am. J. Hum. Genet. 60: 460-463, 1997.

[29699] 4029.

[29700] Puccio, H.; Koenig, M.: Recent advances in the molecular
pathogenesis

[29701] of Friedreich ataxia. Hum. Molec. Genet. 9: 887-892, 2000.

[29702] 4030.

[29703] Rodius, F.; Duclos, F.; Wrogemann, K.; Le Paslier, D.; Ougen,

[29704] P.; Billault, A.; Belal, S.; Musenger, C.; Brice, A.; Durr, A.; Mignard,

[29705] C.; Sirugo, G.; Weissenbach, J.; Cohen, D.; Hentati, F.; Ben Hamida,

[29706] M.; Mandel, J.-L.; Koenig, M.: Recombinations in individu-als homozygous

[29707] by descent localize the Friedreich ataxia locus in a cloned 450-kb

[29708] interval. Am. J. Hum. Genet. 54: 1050-1059, 1994.

[29709] 4031.

[29710] Roca, B.; Calabuig, C.; Sastre, J.; Arenas, M.: Hereditary xanthinuria

[29711] and Ehlers-Danlos syndrome. J. Inherit. Metab. Dis. 15: 881-882,

[29712] 1992.

[29713] 4032.

[29714] Rosenfeld, E. L.; Popova, I. A.; Chibisov, I. V.: Some cases

[29715] of type III glycogen storage disease. Clin. Chim. Acta 67: 123-130,

[29716] 1976.

[29717] 4033.

[29718] Santer, R.; Kinner, M.; Steuerwald, U.; Kjaergaard, S.; Skovby,

[29719] F.; Simonsen, H.; Shaiu, W.-L.; Chen, Y.-T.; Schneppenheim, R.; Schaub,

[29720] J.: Molecular genetic basis and prevalence of glycogen storage disease

[29721] type IIIA in the Faroe Islands. Europ. J. Hum. Genet. 9: 388-391,

[29722] 2001.

[29723] 4034.

[29724] Shaiu, W.-L.; Kishnani, P. S.; Shen, J.; Liu, H.-M.; Chen, Y.-T.

[29725] : Genotype-phenotype correlation in two frequent mutations and mutation

[29726] update in type III glycogen storage disease. Molec. Genet. Metab. 69:

[29727] 16-23, 2000.

[29728] 4035.

[29729] Shen, J,; Bao, Y.; Chen, Y.-T.: A nonsense mutation due to a

[29730] single base insertion in the 3-prime-coding region of glycogen debranching

[29731] enzyme gene associated with a severe phenotype in a patient with glycogen

[29732] storage disease type IIIa. Hum. Mutat. 9: 37-40, 1997.

[29733] 4036.

[29734] Shen, J.; Bao, Y.; Liu, H.-M.; Lee, P.; Leonard, J. V.; Chen,

[29735] Y.-T.: Mutations in exon 3 of the glycogen debranching enzyme gene

[29736] are associated with glycogen storage disease type III that is differentially

[29737] expressed in liver and muscle. J. Clin. Invest. 98: 352-357, 1996.

[29738] 4037.

[29739] Shen, J.; Liu, H.-M.; Bao, Y.; Chen, Y.-T.: Polymorphic markers

[29740] of the glycogen debranching enzyme gene allowing link-

age analysis
[29741] in families with glycogen storage disease type III. J. Med. Genet. 34:
[29742] 34-38, 1997.

[29743] 4038.

[29744] Slonim, A. E.; Weisberg, C.; Benke, P.; Evans, O. B.; Burr, I.

[29745] M.: Reversal of debrancher deficiency myopathy by the use of high-protein
[29746] nutrition. Ann. Neurol. 11: 420-422, 1982.

[29747] 4039.

[29748] van Creveld, S.: Over een bijzondere stoornis in de kool-hydraatstof-Wisseling
[29749] in den kinderleeftijd. Nederl. Maandschr. Geneesk 8: 349-359, 1928.

[29750] 4040.

[29751] von Gierke, E.: Hepato-nephromegalia glykogenica (Glykogenspeicherkrankheit
[29752] der Leber und Nieren.). Beitr. Path. Anat. 82: 497-513, 1929.

[29753] 4041.

[29754] Waaler, P. E.; Garatun-Tjeldsto, O.; Moe, P. J.: Genetic

studies
[29755] in glycogen storage disease type III. Acta Paediat. Scand. 59: 529-535,
[29756] 1970.

[29757] 4042.

[29758] Yang, B.-Z.; Ding, J.-H.; Bao, Y.; Eason, J. F. M.; Chen, Y.-T.
[29759] : Molecular basis of the enzymatic variability in type III glycogen
[29760] storage disease (GSD-III). (Abstract) Am. J. Hum. Genet. 51 (suppl.):
[29761] A28, 1992.

[29762] 4043.

[29763] Yang, B. Z.; Ding, J. H.; Enghild, J. J.; Bao, Y.; Chen, Y. T.
[29764] : Molecular cloning and nucleotide sequence of cDNA encoding human
[29765] muscle glycogen debranching enzyme. J. Biol. Chem. 267: 9294-9299,
[29766] 1992.

[29767] 4044.

[29768] Yang-Feng, T. L.; Zheng, K.; Yu, J.; Yang, B.-Z.; Chen,

Y.-T.;

[29769] Kao, F.-T.: Assignment of the human glycogen debrancher gene to chromosome

[29770] 1p21. Genomics 13: 931-934, 1992.

[29771] 4045.

[29772] Alegria, A.; Martins, E.; Dias, M.; Cunha, A.; Cardoso, M. L.;

[29773] Maire, I.: Glycogen storage disease type IV presenting as hydrops

[29774] fetalis. J. Inherit. Metab. Dis. 22: 330-332, 1999.

[29775] 4046.

[29776] Bannayan, G. A.; Dean, W. J.; Howell, R. R.: Type IV glycogen-storage

[29777] disease: light-microscopic and enzymatic study. Am. J. Clin. Path. 66:

[29778] 702-709, 1976.

[29779] 4047.

[29780] Bao, Y.; Kishnani, P.; Wu, J.-Y.; Chen, Y.-T.: Hepatic and neuromuscular

[29781] forms of glycogen storage disease type IV caused by mutations in the

[29782] same glycogen-branching enzyme gene. J. Clin. Invest. 97: 941-948,
[29783] 1996.

[29784] 4048.

[29785] Brown, B. I.; Brown, D. H.: Branching enzyme activity of cultured
[29786] amniocytes and chorionic villi: prenatal testing for type IV glycogen
[29787] storage disease. Am. J. Hum. Genet. 44: 378-381, 1989.

[29788] 4049.

[29789] Bruno, C.; DiRocco, M.; Lamba, L. D.; Bado, M.; Marino, C.; Tsujino,
[29790] S.; Shanske, S.; Stella, G.; Minetti, C.; van Diggelen, O. P.; DiMauro,
[29791] S.: A novel missense mutation in the glycogen branching enzyme gene
[29792] in a child with myopathy and hepatopathy. Neuromusc. Disord. 9:
[29793] 403-407, 1999.

[29794] 4050.

[29795] Cox, P. M.; Brueton, L. A.; Murphy, K. W.; Worthington, V.

C.;

[29796] Bjelogrlic, P.; Lazda, E. J.; Sabire, N. J.; Sewry, C. A.: Early-onset

[29797] fetal hydrops and muscle degeneration in siblings due to a novel variant

[29798] of type IV glycogenosis. Am. J. Med. Genet. 86: 187-193, 1999.

[29799] 4051.

[29800] Ferguson, I. T.; Mahon, M.; Cumming, W. J. K.: An adult case of

[29801] Andersen's disease--type IV glycogenosis: a clinical, his-tochemical,

[29802] ultrastructural and biochemical study. J. Neurol. Sci. 60: 337-351,

[29803] 1983.

[29804] 4052.

[29805] Greene, G. M.; Weldon, D. C.; Ferrans, V. J.; Cheatham, J. P.;

[29806] McComb, R. D.; Brown, B. I.; Gumbiner, C. H.; Vanderhoff, J. A.; Itkin,

[29807] P. G.; McManus, B. M.: Juvenile polysaccharidosis with car-dioskeletal

[29808] myopathy. Arch. Path. Lab. Med. 111: 977-982, 1987.

[29809] 4053.

[29810] Guerra, A. S.; van Diggelen, O. P.; Carneiro, F.; Tsou, R. M.;

[29811] Simoes, S.; Santos, N. T.: A juvenile variant of glycogenosis IV

[29812] (Andersen disease). Europ. J. Pediat. 145: 179-181, 1986.

[29813] 4054.

[29814] Howell, R. R.: Continuing lessons from glycogen storage diseases.

[29815] (Editorial) New Eng. J. Med. 324: 55-56, 1991.

[29816] 4055.

[29817] Howell, R. R.; Kaback, M. M.; Brown, B. I.: Type IV glycogen

[29818] storage disease: branching enzyme deficiency in skin fibroblasts and

[29819] possible heterozygote detection. J. Pediat. 78: 638-642, 1971.

[29820] 4056.

[29821] Levin, B.; Burgess, E. A.; Mortimer, P. E.: Glycogen storage

[29822] disease type IV, amylopectinosis. Arch. Dis. Child. 43:

548-555,

[29823] 1968.

[29824] 4057.

[29825] McMaster, K. R.; Powers, J. M.; Hennigar, G. R., Jr.; Wohltmann,

[29826] H. J.; Farr, G. H., Jr.: Nervous system involvement in type IV glycogenosis. Arch.

[29827] Path. Lab. Med. 103: 105-111, 1979.

[29828] 4058.

[29829] Reusche, E.; Aksu, F.; Goebel, H. H.; Shin, Y. S.; Yokota, T.;

[29830] Reichmann, H.: A mild juvenile variant of type IV glycogenosis. Brain

[29831] Dev. 14: 36-43, 1992.

[29832] 4059.

[29833] Schochet, S. S., Jr.; McCormick, W. F.; Zellweger, H.: Type IV

[29834] glycogenosis (amylopectinosis): light and electron microscopic observations. Arch.

[29835] Path. 90: 354-363, 1970.

[29836] 4060.

[29837] Schroder, J. M.; May, R.; Shin, Y. S.; Sigmund, M.; Nase-Huppmeier,

[29838] S.: Juvenile hereditary polyglucosan body disease with complete branching

[29839] enzyme deficiency (type IV glycogenosis). Acta Neuropath. 85: 419-430,

[29840] 1993.

[29841] 4061.

[29842] Selby, R.; Starzl, T. E.; Yunis, E.; Brown, B. I.; Kendall, R.

[29843] S.; Tzakis, A.: Liver transplantation for type IV glycogen storage

[29844] disease. New Eng. J. Med. 324: 39-42, 1991.

[29845] 4062.

[29846] Lorenzi, M. V.; Horii, Y.; Yamanaka, R.; Sakaguchi, K.; Miki, T.

[29847] : FRAG1, a gene that potently activates fibroblast growth factor receptor

[29848] by C-terminal fusion through chromosomal rearrange-ment. Proc. Nat.

[29849] Acad. Sci. 93: 8956-8961, 1996.

[29850] 4063.

[29851] Macdonald, D.; Aguiar, R. C.; Mason, P. J.; Goldman, J. M.; Cross,
[29852] N. C.: A new myeloproliferative disorder associated with chromosomal
[29853] translocations involving 8p11: a review. Leukemia 9: 1628-1630,
[29854] 1995.

[29855] 4064.

[29856] Muenke, M.; Schell, U.; Hehr, A.; Robin, N. H.; Losken, H. W.;
[29857] Schinzel, A.; Pulleyn, L. J.; Rutland, P.; Reardon, W.; Malcolm, S.;
[29858] Winter, R. M.: A common mutation in the fibroblast growth factor
[29859] receptor 1 gene in Pfeiffer syndrome. Nature Genet. 8: 269-274,
[29860] 1994.

[29861] 4065.

[29862] Popovici, C.; Adelaide, J.; Ollendorff, V.; Chaffanet, M.; Guasch,
[29863] G.; Jacrot, M.; Leroux, D.; Birnbaum, D.; Pebusque, M.-J.: Fibroblast

[29864] growth factor receptor 1 is fused to FIM in stem-cell myeloproliferative

[29865] disorder with t(8;13)(p12;q12). Proc. Nat. Acad. Sci. 95: 5712-5717,

[29866] 1998.

[29867] 4066.

[29868] Popovici, C.; Zhang, B.; Gregoire, M.-J.; Jonveaux, P.; Lafage-Pochitaloff,

[29869] M.; Birnbaum, D.; Pebusque, M.-J.: The t(6;8)(q27;p11) translocation

[29870] in a stem cell myeloproliferative disorder fuses a novel gene, FOP,

[29871] to fibroblast growth factor receptor 1. Blood 93: 1381-1389, 1999.

[29872] 4067.

[29873] Robin, N. H.; Feldman, G. J.; Mitchell, H. F.; Lorenz, P.; Wilroy,

[29874] R. S.; Zackai, E. H.; Allanson, J. E.; Reich, E. W.; Pfeiffer, R.

[29875] A.; Clarke, L. A.; Warman, M. L.; Mulliken, J. B.; Brueton, L. A.;

[29876] Winter, R. M.; Price, R. A.; Gasser, D. L.; Muenke, M.: Linkage of

[29877] Pfeiffer syndrome to chromosome 8 centromere and evi-
dence for genetic
[29878] heterogeneity. Hum. Molec. Genet. 3: 2153-2158, 1994.

[29879] 4068.

[29880] Ruta, M.; Burgess, W.; Givol, D.; Epstein, J.; Neiger, N.;
Kaplow,
[29881] J.; Crumley, G.; Dionne, C.; Jaye, M.; Schlessinger, J.: Re-
ceptor
[29882] for acidic fibroblast growth factor is related to the tyrosine
kinase
[29883] encoded by the FMS-like gene (FLG). Proc. Nat. Acad. Sci.
86: 8722-8726,
[29884] 1989.

[29885] 4069.

[29886] Ruta, M.; Howk, R.; Ricca, G.; Drohan, W.; Zabelshansky,
M.; Laureys,
[29887] G.; Barton, D. E.; Francke, U.; Schlessinger, J.; Givol, D.: A
novel
[29888] protein tyrosine kinase gene whose expression is modu-
lated during
[29889] endothelial cell differentiation. Oncogene 3: 9-15, 1988.

[29890] 4070.

[29891] Sohal, J.; Chase, A.; Mould, S.; Corcoran, M.; Oscier, D.; Iqbal,
[29892] S.; Parker, S.; Welborn, J.; Harris, R. I.; Martinelli, G.; Montefusco,
[29893] V.; Sinclair, P.; Wilkins, B. S.; van den Berg, H.; Vanstraelen, D.;
[29894] Goldman, J. M.; Cross, N. C. P.: Identification of four new translocations
[29895] involving FGFR1 in myeloid disorders. Genes Chromosomes Cancer 32:
[29896] 155-163, 2001.

[29897] 4071.

[29898] Thomas, W.; Rubenstein, M.; Goto, M.; Drayna, D.: A genetic analysis
[29899] of the Werner syndrome region on human chromosome 8p. Genomics 16:
[29900] 685-690, 1993.

[29901] 4072.

[29902] Twigg, S. R. F.; Burns, H. D.; Oldridge, M.; Heath, J. K.; Wilkie,

[29903] A. O. M.: Conserved use of a non-canonical 5-prime splice site (/GA)
[29904] in alternative splicing by fibroblast growth factor receptors 1, 2
[29905] and 3. Hum. Molec. Genet. 7: 685-691, 1998.

[29906] 4073.

[29907] Wang, L.-Y.; Edenson, S. P.; Yu, Y.-L.; Senderowicz, L.; Turck,
[29908] C. W.: A natural kinase-deficient variant of fibroblast growth factor
[29909] receptor 1. Biochemistry 35: 10134-10142, 1996.

[29910] 4074.

[29911] Welm, B. E.; Freeman, K. W.; Chen, M.; Contreras, A.; Spencer,
[29912] D. M.; Rosen, J. M.: Inducible dimerization of FGFR1: development
[29913] of a mouse model to analyze progressive transformation of the mammary
[29914] gland. J. Cell Biol. 157: 703-714, 2002.

[29915] 4075.

[29916] Wilkie, A. O. M.; Morriss-Kay, G. M.: Genetics of craniofa-

cial
[29917] development and malformation. Nature Rev. Genet. 2:
458-468, 2001.
[29918] 4076.

[29919] Wilkie, A. O. M.; Patey, S. J.; Kan, S.; van den Ouweland, A.

[29920] M. W.; Hamel, B. C. J.: FGFs, their receptors, and human
limb malformations:
[29921] clinical and molecular correlations. Am. J. Med. Genet.
112: 266-278,
[29922] 2002.

[29923] 4077.

[29924] Wood, S.; Schertzer, M.; Yaremko, M. L.: Sequence identity
locates
[29925] CEBPD and FGFR1 to mapped human loci within proximal
8p. Cytogenet.
[29926] Cell Genet. 70: 188-191, 1995.

[29927] 4078.

[29928] Xiao, S.; Nalabolu, S. R.; Aster, J. C.; Ma, J.; Abruzzo, L.;

[29929] Jaffe, E. S.; Stone, R.; Weissman, S. M.; Hudson, T. J.;
Fletcher,
[29930] J. A.: FGFR1 is fused with a novel zinc-finger gene,

ZNF198, in the
[29931] t(8;13) leukaemia/lymphoma syndrome. Nature Genet. 18: 84-87, 1998.
[29932] 4079.
[29933] Zhou, Y.-X.; Xu, X.; Chen, L.; Li, C.; Brodie, S. G.; Deng, C.-X.
[29934] : A Pro250Arg substitution in mouse Fgfr1 causes increased expression
[29935] of Cbfa1 and premature fusion of calvarial sutures. Hum. Molec. Genet. 9:
[29936] 2001-2008, 2000.
[29937] 4080.
[29938] Wang, D.; Kranz-Eble, P.; De Vivo, D. C.: Mutational analysis
[29939] of GLUT1 (SLC2A1) in Glut-1 deficiency syndrome. Hum. Mutat. 16:
[29940] 224-231, 2000.
[29941] 4081.
[29942] Xiang, K.; Cox, N. J.; Karam, J. H.; Bell, G. I.: Bgl II RFLP
[29943] at the human erythrocyte/HepG2-type glucose transporter (GLUT) locus

[29944] on chromosome 1. Nucleic Acids Res. 15: 9101 only,
1987.

[29945] 4082.

[29946] Andersson, S. M.; Pispa, J. P.: Purification and properties
of

[29947] human liver tyrosine aminotransferase. Clin. Chim. Acta
125: 117-123,

[29948] 1982.

[29949] 4083.

[29950] Barton, D. E.; Yang-Feng, T. L.; Francke, U.: The human
tyrosine

[29951] aminotransferase gene mapped to the long arm of chro-
mosome 16 (region

[29952] 16q22-q24) by somatic cell hybrid analysis and in situ
hybridization.

[29953] Hum. Genet. 72: 221-224, 1986.

[29954] 4084.

[29955] Chen, S.-H.; Giblett, E. R.: Genetic variation of soluble
glutamic-oxaloacetic

[29956] transaminase in man. Am. J. Hum. Genet. 23: 419-424,
1971.

[29957] 4085.

[29958] Craig, I. W.; Tolley, E.; Bobrow, M.; van Heyningen, V.: As-
signment
[29959] of a gene necessary for the expression of mitochondrial
glutamic-oxaloacetic
[29960] transaminase in human-mouse hybrid cells. Cytogenet.
Cell Genet. 22:
[29961] 190-194, 1978.

[29962] 4086.

[29963] Davidson, R. G.; Cortner, J. A.; Rattazzi, M. C.; Ruddle, F.
H.;
[29964] Lubs, H. A.: Genetic polymorphisms of human mitochon-
drial glutamic
[29965] oxaloacetic transaminase. Science 169: 391-392, 1970.

[29966] 4087.

[29967] DeLorenzo, R. J.; Ruddle, F. H.: Glutamate transaminase
(GOT)
[29968] genetics in mus musculus: linkage, polymorphism, and
phenotypes of
[29969] the GOT-2 and GOT-1 loci. Biochem. Genet. 4: 259-273,
1970.

[29970] 4088.

[29971] Francke, U.; Weitkamp, L. R.: Report of the committee on the genetic

[29972] constitution of chromosome 6. Cytogenet. Cell Genet. 25: 32-38,

[29973] 1979.

[29974] 4089.

[29975] Jeremiah, S. J.; Povey, S.; Burley, M. W.; Kielty, C.; Lee, M.;

[29976] Spowart, G.; Corney, G.; Cook, P. J. L.: Mapping studies on human

[29977] mitochondrial glutamate oxaloacetate transaminase. Ann. Hum. Genet. 46:

[29978] 145-152, 1982.

[29979] 4090.

[29980] Pol, S.; Bousquet-Lemercier, B.; Pave-Preux, M.; Bulle, F.; Passage,

[29981] E.; Hanoune, J.; Mattei, M. G.; Barouki, R.: Chromosomal localization

[29982] of human aspartate aminotransferase genes by in situ hybridization.

[29983] Hum. Genet. 83: 159-164, 1989.

[29984] 4091.

[29985] Pol, S.; Bousquet-Lemercier, B.; Pave-Preux, M.; Pawlak, A.; Nalpas,

[29986] B.; Berthelot, P.; Hanoune, J.; Barouki, R.: Nucleotide se-quence

[29987] and tissue distribution of the human mitochondrial aspar-tate aminotransferase

[29988] mRNA. Biochem. Biophys. Res. Commun. 157: 1309-1315, 1988.

[29989] 4092.

[29990] Tolley, E.; van Heyningen, V.; Brown, R.; Bobrow, M.; Craig, I.

[29991] W.: Assignment to chromosome 16 of a gene necessary for the expression

[29992] of human mitochondrial glutamate oxaloacetate transami-nase (aspartate

[29993] aminotransferase) (E.C. 2.6.1.1). Biochem. Genet. 18: 947-954,

[29994] 1980.

[29995] 4093.

[29996] Toyomasu, T.; Sakakibara, S.; Kagamiyama, H.; Mat-sumoto, H.:

[29997] Genetic polymorphism of mitochondrial glutamate-oxaloacetate transaminase
[29998] in Japanese. Hum. Genet. 66: 90-91, 1984.

[29999] 4094.

[30000] Akagi, M.; Inui, K.; Nakajima, S.; Shima, M.; Nishigaki, T.; Muramatsu,
[30001] T.; Kokubu, C.; Tsukamoto, H.; Sakai, N.; Okada, S.: Mutation analysis
[30002] of two Japanese patients with Fanconi-Bickel syndrome. J. Hum. Genet. 45:
[30003] 60-62, 2000.

[30004] 4095.

[30005] Aperia, A.; Bergqvist, G.; Linne, T.; Zetterstrom, R.: Familial
[30006] Fanconi syndrome with malabsorption and galactose intolerance, normal
[30007] kinase and transferase activity: a report on two siblings. Acta Paediat.
[30008] Scand. 70: 527-533, 1981.

[30009] 4096.

[30010] Baroni, M. G.; Alcolado, J. C.; Pozzilli, P.; Cavallo, M. G.; Li,

[30011] S. R.; Galton, D. J.: Polymorphisms at the GLUT2 (beta-cell/liver)
[30012] glucose transporter gene and non-insulin-dependent diabetes mellitus
[30013] (NIDDM): analysis in affected pedigree members. Clin. Genet. 41:
[30014] 229-234, 1992.

[30015] 4097.

[30016] Berry, G. T.; Baker, L.; Kaplan, F. S.; Witzleben, C. L.: Diabetes-like
[30017] renal glomerular disease in Fanconi-Bickel syndrome. Pediat. Nephrol. 9:
[30018] 287-291, 1995.

[30019] 4098.

[30020] Burwinkel, B.; Sanjad, S. A.; Al-Sabban, E.; Al-Abbad, A.; Kilimann,
[30021] M. W.: A mutation in GLUT2, not in phosphorylase kinase subunits,
[30022] in hepato-renal glycogenosis with Fanconi syndrome and low phosphorylase
[30023] kinase activity. Hum. Genet. 105: 240-243, 1999.

[30024] 4099.

[30025] Efrat, S.: Making sense of glucose sensing. Nature Genet. 17:

[30026] 249-250, 1997.

[30027] 4100.

[30028] Fanconi, G.; Bickel, H.: Die chronische Aminoazidurie (Aminoaeurendiabetes

[30029] oder nephrotisch-glukosurischer Zwergwuchs) bei der Glykogenose und

[30030] der Cystinkrankheit. Helv. Paediat. Acta 4: 359-396, 1949.

[30031] 4101.

[30032] Fukumoto, H.; Seino, S.; Imura, H.; Seino, Y.; Eddy, R. L.; Fukushima,

[30033] Y.; Byers, M. G.; Shows, T. B.; Bell, G. I.: Sequence, tissue distribution,

[30034] and chromosomal localization of mRNA encoding a human glucose transporter-like

[30035] protein. Proc. Nat. Acad. Sci. 85: 5434-5438, 1988.

[30036] 4102.

[30037] Garcia, J. C.; Strube, M.; Leingang, K.; Keller, K.; Mueckler,

[30038] M.: Amino acid substitutions at tryptophan 388 and tryptophan 412

[30039] of the HepG2 (Glut1) glucose transporter inhibit transport activity

[30040] and targeting to the plasma membrane in Xenopus oocytes. J. Biol.

[30041] Chem. 267: 7770-7776, 1992.

[30042] 4103.

[30043] Gitzelmann, R.: Glukagonprobleme bei den Glykogenspeicherkrankheiten. Helv.

[30044] Paediat. Acta 12: 425-479, 1957.

[30045] 4104.

[30046] Guillam, M.-T.; Hummler, E.; Schaerer, E.; Wu, J.-Y.; Birnbaum,

[30047] M. J.; Beermann, F.; Schmidt, A.; Deriaz, N.; Thorens, B.: Early

[30048] diabetes and abnormal postnatal pancreatic islet development in mice

[30049] lacking Glut-2. Nature Genet. 17: 327-330, 1997.

[30050] 4105.

[30051] Holman, G. D.: Side-specific photolabelling of the hexose

transporter. Biochem.

[30052] Soc. Trans. 17: 438-440, 1989.

[30053] 4106.

[30054] Ishihara, H.; Asano, T.; Katagiri, H.; Lin, J.-L.; Tsukuda, K.;

[30055] Shibasaki, Y.; Yazaki, Y.; Oka, Y.: The glucose transport activity

[30056] of GLUT1 is markedly decreased by substitution of a sin-gle amino acid

[30057] with a different charge at residue 415. Biochem. Biophys. Res. Commun. 176:

[30058] 922-930, 1991.

[30059] 4107.

[30060] Katagiri, H.; Asano, T.; Shibasaki, Y.; Lin, J.-L.; Tsukuda, K.;

[30061] Ishihara, H.; Akanuma, Y.; Takaku, F.; Oka, Y.: Substitution of leucine

[30062] for tryptophan 412 does not abolish cytochalasin B la-belling but markedly

[30063] decreases the intrinsic activity of GLUT1 glucose trans-porter. J.

[30064] Biol. Chem. 266: 7769-7773, 1991.

[30065] 4108.

[30066] Lee, P. J.; Van't Hoff, W. G.; Leonard, J. V.: Catch-up growth

[30067] in Fanconi-Bickel syndrome with uncooked cornstarch. J. Inherit.

[30068] Metab. Dis. 18: 153-156, 1995.

[30069] 4109.

[30070] Maiden, M. C. J.; Davis, E. O.; Baldwin, S. A.; Moore, D. C. M.;

[30071] Henderson, P. J. F.: Mammalian and bacterial sugar trans-port proteins

[30072] are homologous. Nature 325: 641-643, 1987.

[30073] 4110.

[30074] Rimoin, D. L.; Phillips, J. A., III: Genetic disorders of the

[30075] pituitary gland.In: Rimoin, D. L.; Connor, J. M.; Pyeritz, R. E. (eds.)

[30076] : Principles and Practice of Medical Genetics. Vol. I. New York:

[30077] Churchill Livingstone (3rd ed.): 1997. Pp. 1331-1364.

[30078] 4111.

[30079] Rosenfeld, R. G: Editorial: is growth hormone deficiency a viable
[30080] diagnosis? J. Clin. Endocr. Metab. 82: 349-351, 1997.

[30081] 4112.

[30082] Ruddle, F. H.: Personal Communication. New Haven, Conn. 2/7/1982.
[30083] 4113.

[30084] Saitoh, H.; Fukushima, T.; Kamoda, T.; Tanae, A.; Kamijo, T.;
[30085] Yamamoto, M.; Ogawa, M.; Hayashi, Y.; Ohmori, S.; Seo, H.: A Japanese
[30086] family with autosomal dominant growth hormone deficiency. Europ.
[30087] J. Pediat. 158: 624-627, 1999.

[30088] 4114.

[30089] Sirand-Pugnet, P.; Durosay, P.; Brody, E.; Marie, J.: An intronic
[30090] (A/U)GGG repeat enhances the splicing of an alternative intron of
[30091] the chicken beta-tropomyosin pre-mRNA. Nucleic Acids Res. 23: 3501-3507,

[30092] 1995.

[30093] 4115.

[30094] Smith, L. E. H.; Kopchick, J. J.; Chen, W.; Knapp, J.; Kinose,

[30095] F.; Daley, D.; Foley, E.; Smith, R. G.; Schaeffer, J. M.: Es-
sential

[30096] role of growth hormone in ischemia-induced retinal neo-
vascularization. Science 276:

[30097] 1706-1709, 1997.

[30098] 4116.

[30099] Sundstrom, M.; Lundqvist, T.; Rodin, J.; Giebel, L. B.; Milli-
gan,

[30100] D.; Norstedt, G.: Crystal structure of an antagonist mutant
of human

[30101] growth hormone, G120R, in complex with its receptor at
2.9 angstrom

[30102] resolution. J. Biol. Chem. 271: 32197-32203, 1996.

[30103] 4117.

[30104] Takahashi, I.; Takahashi, T.; Komatsu, M.; Sato, T.;
Takada, G.

[30105] : An exonic mutation of the GH-1 gene causing familial
isolated growth

[30106] hormone deficiency type II. Clin. Genet. 61: 222-225, 2002.

[30107] 4118.

[30108] Takahashi, Y.; Kaji, H.; Okimura, Y.; Goji, K.; Abe, H.; Chi-

hara,

[30109] K.: Short stature caused by a mutant growth hormone. New Eng. J.

[30110] Med. 334: 432-436, 1996. Note: Erratum: New Eng. J. Med. 334: 1207

[30111] only, 1996.

[30112] 4119.

[30113] Takahashi, Y.; Shirono, H.; Arisaka, O.; Takahashi, K.; Yagi,

[30114] T.; Koga, J.; Kaji, H.; Okimura, Y.; Abe, H.; Tanaka, T.; Chi-

hara,

[30115] K.: Biologically inactive growth hormone caused by an amino acid

[30116] substitution. J. Clin. Invest. 100: 1159-1165, 1997.

[30117] 4120.

[30118] Tillmann, V.; Buckler, J. M. H.; Kibirige, M. S.; Price, D. A.;

[30119] Shalet, S. M.; Wales, J. K. H.; Addison, M. G.; Gill, M. S.;

Whatmore,
[30120] A. J.; Clayton, P. E.: Biochemical tests in the diagnosis of childhood
[30121] growth hormone deficiency. J. Clin. Endocr. Metab. 82: 531-535,
[30122] 1997.

[30123] 4121.

[30124] Tobiume, H.; Kanzaki, S.; Hida, S.; Ono, T.; Moriwake, T.; Yamauchi,
[30125] S.; Tanaka, H.; Seino, Y.: Serum bone alkaline phosphatase isoenzyme
[30126] levels in normal children and children with growth hormone (GH) deficiency:
[30127] a potential marker for bone formation and response to GH therapy. J.
[30128] Clin. Endocr. Metab. 82: 2056-2061, 1997.

[30129] 4122.

[30130] Valenta, L. J.; Sigel, M. B.; Lesniak, M. A.; Elias, A. N.; Lewis,
[30131] U. J.; Friesen, H. G.; Kershnar, A. K.: Pituitary dwarfism in a patient
[30132] with circulating abnormal growth hormone polymers. New

Eng. J. Med. 312:
[30133] 214-217, 1985.

[30134] 4123.

[30135] Vnencak-Jones, C. L.; Phillips, J. A., III; Chen, E. Y.; See-
burg,
[30136] P. H.: Molecular basis of human growth hormone gene
deletions. Proc.
[30137] Nat. Acad. Sci. 85: 5615-5619, 1988.

[30138] 4124.

[30139] Tsuji, S.; Kobayashi, H.; Uchida, Y.; Ihara, Y.; Miyatake, T.:

[30140] Molecular cloning of human growth inhibitory factor cDNA
and its down-regulation
[30141] in Alzheimer's disease. EMBO J. 11: 4843-4850, 1992.

[30142] 4125.

[30143] Uchida, Y.; Takio, K.; Titani, K.; Ihara, Y.; Tomonaga, M.:
The
[30144] growth inhibitory factor that is deficient in the Alzheimer's
disease
[30145] brain is a 68 amino acid metallothionein-like protein.
Neuron 7:
[30146] 337-347, 1991.

[30147] 4126.

[30148] Hoshino, S.; Miyazawa, H.; Enomoto, T.; Hanaoka, F.; Kikuchi, Y.;

[30149] Kikuchi, A.; Ui, M.: A human homologue of the yeast GST1 gene codes

[30150] for a GTP-binding protein and is expressed in a prolifera- tion-dependent

[30151] manner in mammalian cells. EMBO J. 8: 3807-3814, 1989.

[30152] 4127.

[30153] Kikuchi, Y.; Shimatake, H.; Kikucki, A.: A yeast gene re- quired

[30154] for the G1-to-S transition encodes a protein containing an A-kinase

[30155] target site and GTPase domain. EMBO J. 7: 1175-1182, 1988.

[30156] 4128.

[30157] Ozawa, K.; Murakami, Y.; Eki, T.; Yokoyama, K.; Soeda, E.; Hoshino,

[30158] S.; Ui, M.; Hanaoka, F.: Mapping of the human GSPT1 gene, a human

[30159] homolog of the yeast GST1 gene, to chromosomal band 16p13.1. Somat.

[30160] Cell Molec. Genet. 18: 189-194, 1992.

[30161] 4129.

[30162] Harris, H.; Hopkinson, D. A.; Robson, E. B.: The incidence of

[30163] rare alleles determining electrophoretic variants: data on 43 enzyme

[30164] loci in man. Ann. Hum. Genet. 37: 237-253, 1974.

[30165] 4130.

[30166] Brady, W. A.; Kokoris, M. S.; Fitzgibbon, M.; Black, M. E.: Cloning,

[30167] characterization, and modeling of mouse and human guanylate kinases. J.

[30168] Biol. Chem. 271: 16734-16740, 1996.

[30169] 4131.

[30170] Dallapiccola, B.; Lungarotti, M. S.; Falorni, A.; Magnani, M.;

[30171] Dacha, M.: Evidence for the assignment of GUK1 gene locus to 1q32-q43

[30172] segment from gene dosage effect. Ann. Genet. 23: 83-85, 1980.

[30173] 4132.

[30174] Molchanova, T. P.; Postnikov, Y. V.; Gu, L.-H.; Huisman, T. H.

[30175] J.: Hb A(2)-Grovetown or alpha (2) delta (2) 75 (E19) leu-to-val.

[30176] (Letter) Hemoglobin 17: 289-291, 1993.

[30177] 4133.

[30178] Neeb, H.; Beiboer, J. L.; Jonxis, J. H.; Kaars-Sijpesteijn, J.

[30179] A.; Muller, C. J.: Homozygous Lepore haemoglobin disease appearing

[30180] as thalassaemia major in two Papuan siblings. Trop. Ge-ogr. Med. 13:

[30181] 207-215, 1961.

[30182] 4134.

[30183] Kajiwara, K.; Berson, E. L.; Dryja, T. P.: Digenic retinitis

[30184] pigmentosa due to mutations at the unlinked peripherin/RDS and ROM1

[30185] loci. Science 264: 1604-1608, 1994.

[30186] 4135.

[30187] Kajiwara, K.; Hahn, L. B.; Mukai, S.; Travis, G. H.; Berson, E.

[30188] L.; Dryja, T. P.: Mutations in the human retinal degenera-
tion slow
[30189] gene in autosomal dominant retinitis pigmentosa. Nature
354: 480-483,
[30190] 1991.

[30191] 4136.

[30192] Kajiwara, K.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.: A

[30193] null mutation in the human RDS/peripherin gene in a
family with autosomal
[30194] dominant retinitis punctata albescens. (Abstract) Am. J.
Hum. Genet. 51
[30195] (suppl.): A6 only, 1992.

[30196] 4137.

[30197] Kajiwara, K.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.: A

[30198] null mutation in the human peripherin/RDS gene in a
family with autosomal
[30199] dominant retinitis punctata albescens. Nature Genet. 3:
208-212,
[30200] 1993.

[30201] 4138.

[30202] Kedzierski, W.; Nusinowitz, S.; Birch, D.; Clarke, G.; McInnes,
[30203] R. R.; Bok, D.; Travis, G. H.: Deficiency of rds/peripherin causes
[30204] photoreceptor death in mouse models of digenic and dominant retinitis
[30205] pigmentosa. Proc. Nat. Acad. Sci. 98: 7718-7723, 2001.

[30206] 4139.

[30207] Keen, T. J.; Inglehearn, C. F.: Mutations and polymor-phisms in
[30208] the human peripherin-RDS gene and their involvement in inherited retinal
[30209] degeneration. Hum. Mutat. 8: 297-303, 1996.

[30210] 4140.

[30211] Kikawa, E.; Nakazawa, M.; Chida, Y.; Shiono, T.; Tamai, M.: A
[30212] novel mutation (asn244-to-lys) in the peripherin/RDS gene causing
[30213] autosomal dominant retinitis pigmentosa associated with bull's-eye
[30214] maculopathy detected by nonradioisotopic SSCP. Ge-nomics 20: 137-139,

[30215] 1994.

[30216] 4141.

[30217] Kim, R. Y.; Dollfus, H.; Keen, T. J.; Fitzke, F. W.; Arden, G.

[30218] B.; Bhattacharya, S. S.; Bird, A. C.: Autosomal dominant pattern

[30219] dystrophy of the retina associated with a 4-base pair insertion at

[30220] codon 140 in the peripherin/RDS gene. Arch. Ophthal. 113: 451-455,

[30221] 1995.

[30222] 4142.

[30223] Kohl, S.; Christ-Adler, M.; Apfelstedt-Sylla, E.; Kellner, U.;

[30224] Eckstein, A.; Zrenner, E.; Wissinger, B.: RDS/peripherin gene mutations

[30225] are frequent causes of central retinal dystrophies. J. Med. Genet. 34:

[30226] 620-626, 1997.

[30227] 4143.

[30228] Ma, J.; Norton, J. C.; Allen, A. C.; Burns, J. B.; Hasel, K. W.;

[30229] Burns, J. L.; Sutcliffe, J. G.; Travis, G. H.: Retinal degenera-

tion

[30230] slow (rds) in mouse results from simple insertion of a t haplotype-specific

[30231] element into protein-coding exon II. Genomics 28: 212-219, 1995.

[30232] 4144.

[30233] Nichols, B. E.; Drack, A. V.; Vandenburgh, K.; Kimura, A. E.;

[30234] Sheffield, V. C.; Stone, E. M.: A 2 base pair deletion in the RDS

[30235] gene associated with butterfly-shaped pigment dystrophy of the fovea. Hum.

[30236] Molec. Genet. 2: 601-603, 1993.

[30237] 4145.

[30238] Nichols, B. E.; Sheffield, V. C.; Vandenburgh, K.; Drack, A. V.;

[30239] Kimura, A. E.; Stone, E. M.: Butterfly-shaped pigment dys- trophy of

[30240] the fovea caused by a point mutation in codon 167 of the RDS gene. Nature

[30241] Genet. 3: 202-207, 1993.

[30242] 4146.

[30243] Payne, A. M.; Downes, S. M.; Bessant, D. A. R.; Bird, A. C.; Bhattacharya,
[30244] S. S.: Founder effect, seen in the British population, of the 172
[30245] peripherin/RDS mutation--and further refinement of genetic positioning
[30246] of the peripherin/RDS gene. (Letter) Am. J. Hum. Genet. 192-195,
[30247] 1998.

[30248] 4147.

[30249] Pendleton, J. W.; Violette, S. M.; Hunihan, L. W.; Greene, L.
[30250] A.; Ruddle, F. H.: The peripherin gene maps to mouse chromosome 15. Genomics 9:
[30251] 369-372, 1991.

[30252] 4148.

[30253] Reig, C.; Serra, A.; Gean, E.; Vidal, M.; Arumi, J.; De la Calzada,
[30254] M. D.; Antich, J.; Carballo, M.: A point mutation in the RDS-peripherin
[30255] gene in a Spanish family with central areolar choroidal dystrophy. Ophthal.

[30256] Genet. 16: 35-44, 1995.

[30257] 4149.

[30258] Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bainbridge, J. W. B.;

[30259] Smith, A. J.; Thrasher, A. J.; Ali, R. R.: Gene replacement therapy

[30260] in the retinal degeneration slow (rds) mouse: the effect on retinal

[30261] degeneration following partial transduction of the retina. Hum. Molec.

[30262] Genet. 10: 2353-2361, 2001.

[30263] 4150.

[30264] Travis, G. H.; Brennan, M. B.; Danielson, P. E.; Kozak, C. A.;

[30265] Sutcliffe, J. G.: Identification of a photoreceptor-specific mRNA

[30266] encoded by the gene responsible for retinal degeneration slow (rds). Nature 338:

[30267] 70-73, 1989.

[30268] 4151.

[30269] Travis, G. H.; Christerson, L.; Danielson, P. E.; Klisak, I.;

[30270] Sparkes, R. S.; Hahn, L. B.; Dryja, T. P.; Sutcliffe, J. G.: The

[30271] human retinal degeneration slow (RDS) gene: chromosome assignment

[30272] and structure of the mRNA. Genomics 10: 733-739, 1991.

[30273] 4152.

[30274] Travis, G. H.; Hepler, J. E.: A medley of retinal dystrophies. Nature

[30275] Genet. 3: 191-192, 1993.

[30276] 4153.

[30277] Cramer, P.; Bushnell, D. A.; Fu, J.; Gnatt, A. L.; Maier-Davis,

[30278] B.; Thompson, N. E.; Burgess, R. R.; Edwards, A. M.; David, P. R.;

[30279] Kornberg, R. D.: Architecture of RNA polymerase II and implications

[30280] for the transcription mechanism. Science 288: 640-648, 2000.

[30281] 4154.

[30282] Yudkovsky, N.; Ranish, J. A.; Hahn, S.: A transcription reinitiation

[30283] intermediate that is stabilized by activator. Nature 408:

225-229,
[30284] 2000.

[30285] 4155.

[30286] Dammann, R.; Pfeifer, G. P.: Cloning and characterization of the
[30287] human RNA polymerase I subunit hRPA40. Biochim. Biophys. Acta 1396:
[30288] 153-157, 1998.

[30289] 4156.

[30290] Fanciulli, M.; Bruno, T.; Di Padova, M.; De Angelis, R.; Lovari,
[30291] S.; Floridi, A.; Passananti, C.: The interacting RNA polymerase II
[30292] subunits, hRPB11 and hRPB3, are coordinately expressed in adult human
[30293] tissues and down-regulated by doxorubicin. FEBS Lett. 427: 236-240,
[30294] 1998.

[30295] 4157.

[30296] Pati, U. K.; Weissman, S. M.: The amino acid sequence of the human

[30297] RNA polymerase II 33-kDa subunit hRPB 33 is highly conserved among

[30298] eukaryotes. J. Biol. Chem. 265: 8400-8403, 1990.

[30299] 4158.

[30300] Nelissen, R. L. H.; Sillekens, P. T. G.; Beijer, R. P.; Geurts

[30301] van Kessel, A. H. M.; van Venrooij, W. J.: Structure, chromosomal

[30302] localization and evolutionary conservation of the gene encoding human

[30303] U1 snRNP-specific A protein. Gene 102: 189-196, 1991.

[30304] 4159.

[30305] Henrion, A. A.; Martinez, A.; Mattei, M.-G.; Kahn, A.; Raymondjean,

[30306] M.: Structure, sequence, and chromosomal location of the gene for

[30307] USF2 transcription factors in mouse. Genomics 25: 36-43, 1995.

[30308] 4160.

[30309] Jakimiuk, A. J.; Weitsman, S. R.; Magoffin, D. A.: 5-alpha-reductase

[30310] activity in women with polycystic ovary syndrome. J. Clin.

Endocr.
[30311] Metab. 84: 2414-2418, 1999.

[30312] 4161.

[30313] Reddi, A. H.: BMP-1: resurrection as procollagen C-proteinase. Science 271:
[30314] 5-6, 1996.

[30315] 4162.

[30316] Takahara, K.; Lee, S.; Wood, S.; Greenspan, D. S.: Structural
[30317] organization and genetic localization of the human bone morphogenetic
[30318] protein 1/mammalian tolloid gene. Genomics 29: 9-15, 1995.
[30319] 4163.

[30320] Takahara, K.; Lyons, G. E.; Greenspan, D. S.: Bone morphogenetic
[30321] protein-1 and a mammalian tolloid homologue (mTld) are encoded by
[30322] alternatively spliced transcripts which are differentially expressed
[30323] in some tissues. J. Biol. Chem. 269: 32572-32578, 1994.

[30324] 4164.

[30325] Scott, I. C.; Blitz, I. L.; Pappano, W. N.; Imamura, Y.; Clark,

[30326] T. G.; Steiglitz, B. M.; Thomas, C. L.; Maas, S. A.; Takahara, K.;

[30327] Cho, K. W. Y.; Greenspan, D. S.: Mammalian BMP-1/Tolloid-related

[30328] metalloproteinases, including novel family member mammalian Tolloid-like

[30329] 2, have differential enzymatic activities and distributions of expression

[30330] relevant to patterning and skeletogenesis. Dev. Biol. 213: 283-300,

[30331] 1999.

[30332] 4165.

[30333] Yoshiura, K.; Tamura, T.; Hong, H.-S.; Ohta, T.; Soejima, H.;

[30334] Kishino, T.; Jinno, Y.; Niikawa, N.: Mapping of the bone morphogenetic

[30335] protein 1 gene (BMP1) to 8p21: removal of BMP1 from candidacy for

[30336] the bone disorder in Langer-Giedion syndrome. Cytogenet. Cell Genet. 64:

[30337] 208-209, 1993.

[30338] 4166.

[30339] Hahn, G. V.; Cohen, R. B.; Wozney, J. M.; Levitz, C. L.; Shore,

[30340] E. M.; Zasloff, M. A.; Kaplan, F. S.: A bone morphogenetic protein

[30341] subfamily: chromosomal localization of human genes for BMP5, BMP6,

[30342] and BMP7. Genomics 14: 759-762, 1992.

[30343] 4167.

[30344] Lyons, K.; Graycar, J. L.; Lee, A.; Hashmi, S.; Lindquist, P. B.;

[30345] Chen, E. Y.; Hogan, B. L. M.; Derynck, R.: Vgr-1, a mammalian gene

[30346] related to Xenopus Vg-1, is a member of the transforming growth factor

[30347] beta gene superfamily. Proc. Nat. Acad. Sci. 86: 4554-4558, 1989.

[30348] 4168.

[30349] Rickard, D. J.; Hofbauer, L. C.; Bonde, S. K.; Gori, F.; Spelsberg,

[30350] T. C.; Riggs, B. L.: Bone morphogenetic protein-6 production in human
[30351] osteoblastic cell lines: selective regulation by estrogen. J. Clin.
[30352] Invest. 101: 413-422, 1998.

[30353] 4169.

[30354] Modi, W. S.; Dean, M.; Pollock, D. D.; Seuanez, H. N.; Christakos,
[30355] S.: Chromosomal localization of the calbindin gene. (Abstract) Cytogenet.
[30356] Cell Genet. 58: 1930 only, 1991.

[30357] 4170.

[30358] Parmentier, M.; De Vijlder, J. J. M.; Muir, E.; Szpirer, C.; Islam,
[30359] M. Q.; Geurts van Kessel, A.; Lawson, D. E. M.; Vassart, G.: The
[30360] human calbindin 27 kDa gene: structural organization of the 5-prime
[30361] and 3-prime regions, chromosomal assignment and restriction fragment
[30362] length polymorphism. Genomics 4: 309-319, 1989.

[30363] 4171.

[30364] Parmentier, M.; Lawson, D. E. M.; Vassart, G.: Human
27-kDa calbindin
[30365] complementary DNA sequence: evolutionary and func-
tional implications. Europ.
[30366] J. Biochem. 170: 207-215, 1987.

[30367] 4172.

[30368] Parmentier, M.; Passage, E.; Vassart, G.; Mattei, M.-G.: The
human
[30369] calbindin D28k (CALB1) and calretinin (CALB2) genes are
located at
[30370] 8q21.3-q22.1 and 16q22-q23, respectively, suggesting a
common duplication
[30371] with the carbonic anhydrase isozyme loci. Cytogenet. Cell
Genet. 57:
[30372] 41-43, 1991.

[30373] 4173.

[30374] Parmentier, M.; Vassart, G.: HindIII RFLP on chromosome 8
detected
[30375] with a calbindin 27 kDa cDNA probe, HBSC21. Nucleic
Acids Res. 16:

[30376] 9373 only, 1988.

[30377] 4174.

[30378] Seto-Ohshima, A.; Emson, P. C.; Lawson, E.; Mountjoy, C. Q.; Carrasco,

[30379] L. H.: Loss of matrix calcium-binding protein-containing neurons

[30380] in Huntington's disease. Lancet I: 1252-1254, 1988.

[30381] 4175.

[30382] Parmentier, M.; Szpirer, J.; Levan, G.; Vassart, G.: The human

[30383] genes for calbindin 27 and 29 kDa proteins are located on chromosomes

[30384] 8 and 16, respectively. Cytogenet. Cell Genet. 52: 85-87, 1989.

[30385] 4176.

[30386] Schiffmann, S. N.; Cheron, G.; Lohof, A.; d'Alcantara, P.; Meyer,

[30387] M.; Parmentier, M.; Schurmans, S.: Impaired motor coordination and

[30388] Purkinje cell excitability in mice lacking calretinin. Proc. Nat.

[30389] Acad. Sci. 96: 5257-5262, 1999.

[30390] 4177.

[30391] Davies, A. A.; Moss, S. E.; Crompton, M. R.; Jones, T. A.; Spurr,

[30392] N. K.; Sheer, D.; Kozak, C.; Crumpton, M. J.: The gene coding for

[30393] the p68 calcium-binding protein is localized to bands q32-q34 of human

[30394] chromosome 5, and to mouse chromosome 11. Hum. Genet. 82: 234-238,

[30395] 1989.

[30396] 4178.

[30397] Smith, P. D.; Davies, A.; Crumpton, M. J.; Moss, S. E.: Structure

[30398] of the human annexin VI gene. Proc. Nat. Acad. Sci. 91: 2713-2717,

[30399] 1994.

[30400] 4179.

[30401] Sudhof, T. C.; Slaughter, C. A.; Leznicki, I.; Barjon, P.; Reynolds,

[30402] G. A.: Human 67-kDa calelectrin contains a duplication of

four repeats
[30403] found in 35-kDa lipocortins. Proc. Nat. Acad. Sci. 85:
664-668,
[30404] 1988.

[30405] 4180.

[30406] Blow, J. J.; Laskey, R. A.: A role for the nuclear envelope in

[30407] controlling DNA replication within the cell cycle. Nature
332: 546-548,
[30408] 1988.

[30409] 4181.

[30410] Burkhart, R.; Schulte, D.; Hu, D.; Musahl, C.; Gohring, F.;
Knippers,
[30411] R.: Interactions of human nuclear proteins P1Mcm3 and
P1Cdc46. Europ.
[30412] J. Biochem. 228: 431-438, 1995.

[30413] 4182.

[30414] Chong, J. P.; Mahbubani, H. M.; Khoo, C. Y.; Blow, J. J.: Pu-
rification
[30415] of an MCM-containing complex as a component of the
DNA replication
[30416] licensing system. Nature 375: 418-421, 1995.

[30417] 4183.

[30418] Hu, B.; Burkhart, R.; Schulte, D.; Musahl, C.; Knippers, R.: The

[30419] P1 family: a new class of nuclear mammalian proteins re-

lated to the

[30420] yeast Mcm replication proteins. Nucleic Acids Res. 21:

5289-5293,

[30421] 1993.

[30422] 4184.

[30423] Kubota, Y.; Mimura, S.; Nishimoto, S.; Takisawa, H.; No-

jima, H.

[30424] : Identification of the yeast MCM3-related protein as a

component

[30425] of Xenopus DNA replication licensing factor. Cell 81:

601-609, 1995.

[30426] 4185.

[30427] Labib, K.; Tercero, J. A.; Diffley, J. F. X.: DNA replication

[30428] fork progression requires uninterrupted MCM2-7 func-

tion. Science 288:

[30429] 1643-1647, 2000.

[30430] 4186.

[30431] Madine, M. A.; Khoo, C. Y.; Mills, A. D.; Laskey, R. A.: MCM3
[30432] complex required for cell cycle regulation of DNA replication in vertebrate
[30433] cells. Nature 375: 421-424, 1995.

[30434] 4187.

[30435] Mincheva, A.; Todorov, I.; Werner, D.; Fink, T. M.; Lichter, P.
[30436] : The human gene for nuclear protein BM28 (CDCL1), a new member of
[30437] the early S-phase family of proteins, maps to chromosome band 3q21. Cytogenet.
[30438] Cell Genet. 65: 276-277, 1994.

[30439] 4188.

[30440] Tsuruga, H.; Yabuta, N.; Hashizume, K.; Ikeda, M.; Endo, Y.; Nojima,
[30441] H.: Expression, nuclear localization and interactions of human MCM/P1
[30442] proteins. Biochem. Biophys. Res. Commun. 236: 118-125, 1997.
[30443] 4189.

[30444] Adams, M. D.; Dubnick, M.; Kerlavage, A. R.; Moreno, R.; Kelley,

[30445] J. M.; Utterback, T. R.; Nagle, J. W.; Fields, C.; Venter, J. C.:

[30446] Sequence identification of 2,375 human brain genes. Na-

ture 355:

[30447] 632-634, 1992.

[30448] 4190.

[30449] Tugendreich, S.; Boguski, M. S.; Seldin, M. S.; Hieter, P.: Linking

[30450] yeast genetics to mammalian genomes: identification and mapping of

[30451] the human homolog of CDC27 via the expressed se-

quence tag (EST) data

[30452] base. Proc. Nat. Acad. Sci. 90: 10031-10035, 1993.

[30453] 4191.

[30454] Demetrick, D. J.; Beach, D. H.: Chromosome mapping of human CDC25A

[30455] and CDC25B phosphatases. Genomics 18: 144-147, 1993.

[30456] 4192.

[30457] Plon, S. E.; Leppig, K. A.; Do, H.-N.; Groudine, M.: Cloning of

[30458] the human homolog of the CDC34 cell cycle gene by complementation
[30459] in yeast. Proc. Nat. Acad. Sci. 90: 10484-10488, 1993.

[30460] 4193.

[30461] Lane, S. A.; Baker, E.; Sutherland, G. R.; Tonks, I.; Hayward,
[30462] N.; Ellem, K.: The human cell cycle gene CDC25B is located at 20p13. Genomics 15:
[30463] 693-694, 1993.

[30464] 4194.

[30465] Lincoln, A. J.; Wickramasinghe, D.; Stein, P.; Schultz, R. M.;

[30466] Palko, M. E.; De Miguel, M. P.; Tessarollo, L.; Donovan, P. J.: Cdc25b
[30467] phosphatase is required for resumption of meiosis during oocyte maturation. Nature
[30468] Genet. 30: 446-449, 2002.

[30469] 4195.

[30470] Bentley, K. L.; Ferguson-Smith, A. C.; Miki, T.; Kidd, K. K.; Ruddle,
[30471] F. H.: Physical linkage of Hox 2.1 and nerve growth factor receptor.

[30472] (Abstract) Cytogenet. Cell Genet. 51: 961 only, 1989.

[30473] 4196.

[30474] Bibel, M.; Barde, Y.-A.: Neurotrophins: key regulators of cell

[30475] fate and cell shape in the vertebrate nervous system. Genes Dev. 14:

[30476] 2919-2937, 2000.

[30477] 4197.

[30478] Bothwell, M.: p75(NTR): a receptor after all. Science 272: 506-507,

[30479] 1996.

[30480] 4198.

[30481] Carter, B. D.; Kaltschmidt, C.; Kaltschmidt, B.; Offen-hauser, N.;

[30482] Bohm-Matthaei, R.; Baeuerle, P. A.; Barde, Y.-A.: Selective activation

[30483] of NF-kappa-B by nerve growth factor through the neu-rotrophin receptor

[30484] p75. Science 272: 542-545, 1996.

[30485] 4199.

[30486] Carter, B. D.; Lewin, G. R.: Neurotrophins live or let die: does

[30487] p75(NTR) decide? Neuron 18: 187–190, 1997.

[30488] 4200.

[30489] Chao, M. V.; Bothwell, M. A.; Ross, A. H.; Koprowski, H.; Lanahan,

[30490] A. A.; Buck, C. R.; Sehgal, A.: Gene transfer and molecular cloning

[30491] of the human NGF receptor. Science 232: 518–521, 1986.

[30492] 4201.

[30493] Dobrowsky, R. T.; Werner, M. H.; Castellino, A. M.; Chao, M. V.;

[30494] Hannun, Y. A.: Activation of the sphingomyelin cycle through the

[30495] low-affinity neurotrophin receptor. Science 265: 1596–1599, 1994.

[30496] 4202.

[30497] Colonna, M.; Bresnahan, M.; Bahram, S.; Strominger, J. L.; Spies,

[30498] T.: Allelic variants of the human putative peptide trans-porter involved

[30499] in antigen processing. Proc. Nat. Acad. Sci. 89:
3932-3936, 1992.

[30500] 4203.

[30501] Huh, G. S.; Boulanger, L. M.; Du, H.; Riquelme, P. A.; Brotz,
T.

[30502] M.; Shatz, C. J.: Functional requirement for class I MHC in
CNS development

[30503] and plasticity. Science 290: 2155-2159, 2000.

[30504] 4204.

[30505] Karttunen, J. T.; Lehner, P. J.; Gupta, S. S.; Hewitt, E. W.;
Cresswell,

[30506] P.: Distinct functions and cooperative interaction of the
subunits

[30507] of the transporter associated with antigen processing
(TAP). Proc.

[30508] Nat. Acad. Sci. 98: 7431-7436, 2001.

[30509] 4205.

[30510] Cullen, M.; Erlich, H.; Klitz, W.; Carrington, M.: Molecular
mapping

[30511] of a recombination hotspot located in the second intron of
the human

[30512] TAP2 locus. Am. J. Hum. Genet. 56: 1350-1358, 1995.

[30513] 4206.

[30514] de la Salle, H.; Donato, L.; Zimmer, J; Plebani, A.; Hanau, D.;

[30515] Bonneville, M.; Tongio, M.-M.: HLA class I deficiencies.In: Ochs,

[30516] H. D.; Smith, C. I. E.; Puck, J. M. (eds.): Primary Immunod-eficiency

[30517] Diseases: A Molecular and Genetic Approach. New York: Oxford University

[30518] Press 1999. Pp. 181-188.

[30519] 4207.

[30520] de la Salle, H.; Hanau, D.; Fricker, D.; Urlacher, A.; Kelly, A.;

[30521] Salamero, J.; Powis, S. H.; Donato, L.; Bausinger, H.; Laforet, M.;

[30522] Jeras, M.; Spehner, D.; Bieber, T.; Falkenrodt, A.; Cazenave, J.-P.;

[30523] Trowsdale, J.; Tongio, M.-M.: Homozygous human TAP peptide transporter

[30524] mutation in HLA class I deficiency. Science 265: 237-241, 1994.

[30525] 4208.

[30526] Jeffreys, A. J.; Ritchie, A.; Neumann, R.: High resolution analysis

[30527] of haplotype diversity and meiotic crossover in the human TAP2 recombination

[30528] hotspot. Hum. Molec. Genet. 9: 725-733, 2000.

[30529] 4209.

[30530] Powis, S. H.; Mockridge, I.; Kelly, A.; Kerr, L.-A.; Glynne, R.;

[30531] Gileadi, U.; Beck, S.; Trowsdale, J.: Polymorphism in a sec-ond ABC

[30532] transporter gene located within the class II region of the human major

[30533] histocompatibility complex. Proc. Nat. Acad. Sci. 89: 1463-1467,

[30534] 1992.

[30535] 4210.

[30536] Jackson, R. S.; Creemers, J. W. M.; Ohagi, S.; Raffin-San-son, M.-L.;

[30537] Sanders, L.; Montague, C. T.; Hutton, J. C.; O'Rahilly, S.: Obesity

[30538] and impaired prohormone processing associated with mutations in the
[30539] human prohormone convertase 1 gene. Nature Genet. 16: 303-306, 1997.
[30540] 4211.

[30541] Naggert, J. K.; Fricker, L. D.; Varlamov, O.; Nishina, P. M.; Rouille,
[30542] Y.; Steiner, D. F.; Carroll, R. J.; Paigen, B. J.; Leiter, E. H.:
[30543] Hyperproinsulinaemia in obese fat/fat mice associated with a carboxypeptidase
[30544] E mutation which reduces enzyme activity. Nature Genet. 10: 135-142,
[30545] 1995.

[30546] 4212.

[30547] O'Rahilly, S.; Gray, H.; Humphreys, P. J.; Krook, A.; Polonsky,
[30548] K. S.; White, A.; Gibson, S.; Taylor, K.; Carr, C.: Brief report:
[30549] impaired processing of prohormones associated with abnormalities of
[30550] glucose homeostasis and adrenal function. New Eng. J. Med. 333:

[30551] 1386-1390, 1995.

[30552] 4213.

[30553] Ohagi, S.; Sakaguchi, H.; Sanke, T.; Tatsuta, H.; Hanabusa, T.;

[30554] Nanjo, K.: Human prohormone convertase 3 gene: exon-intron organization

[30555] and molecular scanning for mutations in Japanese subjects with NIDDM. Diabetes 45:

[30556] 897-901, 1996.

[30557] 4214.

[30558] Furuta, M.; Carroll, R.; Martin, S.; Swift, H.; Ravazzola, M.;

[30559] Orci, L.; Steiner, D.: Incomplete processing of proinsulin to insulin

[30560] accompanied by elevation of Des-31,32 proinsulin intermediates in

[30561] islets of mice lacking active PC2. J. Biol. Chem. 273: 3431-3437,

[30562] 1998.

[30563] 4215.

[30564] Furuta, M.; Yano, H.; Zhou, A.; Rouille, Y.; Holst, J.; Caroll,

[30565] R.; Ravazzola, M.; Orci, L.; Furuta, H.; Steiner, D.: Defec-
tive prohormone
[30566] processing and altered pancreatic islet morphology in
mice lacking
[30567] active SPC2. Proc. Nat. Acad. Sci. 94: 6646-6651, 1997.

[30568] 4216.

[30569] Furuta, M.; Zhou, A.; Webb, G.; Carroll, R.; Ravazzola, M.;
Orci,
[30570] L.; Steiner, D. F.: Severe defect in proglucagon processing
in islet
[30571] A-cells of prohormone convertase 2 null mice. J. Biol.
Chem. 276:
[30572] 27197-27202, 2001.

[30573] 4217.

[30574] Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Sei-
dah,
[30575] N. G.; Van de Loo, J.-W.; Van de Ven, W. J. M.; Martens, G.
J. M.
[30576] and van Leeuwen, F. W.: Attenuation of the polypeptide
7B2, prohormone
[30577] convertase PC2, and vasopressin in the hypothalamus of
some Prader-Willi

[30578] patients: indications for a processing defect. J. Clin. Endocr. Metab. 83:

[30579] 591-599, 1998.

[30580] 4218.

[30581] Maglott, D. R.; Feldblyum, T. V.; Durkin, A. S.; Nierman, W. C.

[30582] : Radiation hybrid mapping of SNAP, PCSK2, and THBD (human chromosome

[30583] 20p). Mammalian Genome 7: 400-401, 1996.

[30584] 4219.

[30585] Ohagi, S.; LaMendola, J.; LeBeau, M. M.; Espinosa, R., III; Takeda,

[30586] J.; Smeekens, S. P.; Chan, S. J.; Steiner, D. F.: Identification

[30587] and analysis of the gene encoding human PC2, a prohor-
mone convertase

[30588] expressed in neuroendocrine tissues. Proc. Nat. Acad. Sci.
89: 4977-4981,

[30589] 1992.

[30590] 4220.

[30591] Taylor, N. A.; Shennan, K. I. J.; Cutler, D. F.; Docherty, K.:

[30592] Mutations within the propeptide, the primary cleavage site
or the

[30593] catalytic site, or deletion of C-terminal sequences, pre-
vents secretion

[30594] of proPC2 from transfected COS-7 cells. Biochem. J. 321:
367-373,

[30595] 1997.

[30596] 4221.

[30597] Abeliovich, D.; Gelman-Kohan, Z.; Silverstein, S.; Lerer, I.;
Chemke,

[30598] J.; Merlin, S.; Zlotogora, J.: Familial cafe au lait spots: a
variant

[30599] of neurofibromatosis type 1. J. Med. Genet. 32: 985-986,

1995.

[30600] 4222.

[30601] Abernathy, C. R.; Rasmussen, S. A.; Stalker, H. J.; Zori, R.; Driscoll,

[30602] D. J.; Williams, C. A.; Kousseff, B. G.; Wallace, M. R.: NF1 mutation

[30603] analysis using a combined heteroduplex/SSCP approach. Hum. Mutat. 9:

[30604] 548-554, 1997.

[30605] 4223.

[30606] Ablon, J.: Parents' responses to their child's diagnosis of neurofibromatosis

[30607] 1. Am. J. Med. Genet. 93: 136-142, 2000.

[30608] 4224.

[30609] Adornato, B. T.; Berg, B.: Diencephalic syndrome and von Recklinghausen's

[30610] disease. Ann. Neurol. 2: 159-160, 1977.

[30611] 4225.

[30612] Ainsworth, P. J.; Rodenhiser, D. I.; Costa, M. T.: Identifica-tion

[30613] and characterization of sporadic and inherited mutations

in exon 31
[30614] of the neurofibromatosis (NF1) gene. Hum. Genet. 91: 151-156, 1993.
[30615] 4226.
[30616] Allan, T. N. K.; Davies, E. R.: Neurofibromatosis of the renal
[30617] artery. Brit. J. Radiol. 43: 906-908, 1970.
[30618] 4227.
[30619] Ars, E.; Kruyer, H.; Gaona, A.; Casquero, P.; Rosell, J.; Volpini,
[30620] V.; Serra, E.; Lazaro, C.; Estivill, X.: A clinical variant of neurofibromatosis
[30621] type 1: familial spinal neurofibromatosis with a frameshift mutation
[30622] in the NF1 gene. Am. J. Hum. Genet. 62: 834-841, 1998.
[30623] 4228.
[30624] Ars, E.; Serra, E.; Garcia, J.; Kruyer, H.; Gaona, A.; Lazaro,
[30625] C.; Estivill, X.: Mutations affecting mRNA splicing are the most
[30626] common molecular defects in patients with neurofibromatosis type 1. Hum.

[30627] Molec. Genet. 9: 237-247, 2000.

[30628] 4229.

[30629] Bahuau, M.; Pelet, A.; Vidaud, D.; Lamireau, T.; Le Bail, B.; Munnich,

[30630] A.; Vidaud, M.; Lyonnet, S.; Lacombe, D.: GDNF as a candidate modifier

[30631] in a type 1 neurofibromatosis (NF1) enteric phenotype. (Letter) J.

[30632] Med. Genet. 38: 638-643, 2001.

[30633] 4230.

[30634] Balcer, L. J.; Liu, G. T.; Heller, G.; Bilaniuk, L.; Volpe, N.

[30635] J.; Galetta, S. L.; Molloy, P. T.; Phillips, P. C.; Janss, A. J.;

[30636] Vaughn, S.; Maguire, M. G.: Visual loss in children with neurofibromatosis

[30637] type I and optic pathway gliomas: relation to tumor location by magnetic

[30638] resonance imaging. Am. J. Ophthal. 131: 442-445, 2001.

[30639] 4231.

[30640] Dutta, S.; Akey, I. V.; Dingwall, C.; Hartman, K. L.; Laue, T.;

[30641] Nolte, R. T.; Head, J. F.; Akey, C. W.: The crystal structure

of
[30642] nucleoplasmin-core: implications for histone binding and nucleosome
[30643] assembly. Molec. Cell 8: 841-853, 2001.

[30644] 4232.

[30645] Wilson, J. G.: Personal Communication. Boston, Mass. 10/25/1982.

[30646] 4233.

[30647] Wilson, J. G.; Andriopoulos, N. A.; Fearon, D. T.: CR1 and the
[30648] cell membrane proteins that bind C3 and C4: a basic and clinical review. Immun.
[30649] Res. 6: 192-209, 1987.

[30650] 4234.

[30651] Wilson, J. G.; Jack, R. M.; Wong, W. W.; Schur, P. H.; Fearon,

[30652] D. T.: Autoantibody to the C3b/C4b receptor and absence of this receptor
[30653] from erythrocytes of a patient with systemic lupus erythematosus. J.
[30654] Clin. Invest. 76: 182-190, 1985.

[30655] 4235.

[30656] Wilson, J. G.; Murphy, E. E.; Wong, W. W.; Klickstein, L. B.;

[30657] Weis, J. H.; Fearon, D. T.: Identification of a restriction fragment

[30658] length polymorphism by a CR1 cDNA that correlates with the number

[30659] of CR1 on erythrocytes. J. Exp. Med. 164: 50-59, 1986.

[30660] 4236.

[30661] Wilson, J. G.; Wong, W. W.; Schur, P. H.; Fearon, D. T.: Mode

[30662] of inheritance of decreased C3b receptors on erythrocytes of patients

[30663] with systemic lupus erythematosus. New Eng. J. Med. 307: 981-986,

[30664] 1982.

[30665] 4237.

[30666] Wong, W. W.; Cahill, J. M.; Rosen, M. D.; Kennedy, C. A.; Bonaccio,

[30667] E. T.; Morris, M. J.; Wilson, J. G.; Klickstein, L. B.; Fearon, D.

[30668] T.: Structure of the human CR1 gene: molecular basis of the structural

[30669] and quantitative polymorphisms and identification of a new CR1-like

[30670] allele. J. Exp. Med. 169: 847-863, 1989.

[30671] 4238.

[30672] Wong, W. W.; Klickstein, L. B.; Smith, J. A.; Weis, J. H.; Fearon,

[30673] D. T.: Identification of a partial cDNA clone for the human receptor

[30674] for complement fragments C3b/C4b. Proc. Nat. Acad. Sci. 82: 7711-7715,

[30675] 1985.

[30676] 4239.

[30677] Alper, C. A.; Abramson, N.; Johnston, R. B., Jr.; Jandl, J. H.;

[30678] Rosen, F. S.: Increased susceptibility to infection associated with

[30679] abnormalities of complement-mediated functions and of the third component

[30680] of complement (C3). New Eng. J. Med. 282: 349-354, 1970.

[30681] 4240.

[30682] Alper, C. A.; Abramson, N.; Johnston, R. B., Jr.; Jandl, J. H.;

[30683] Rosen, F. S.: Studies in vivo and in vitro on an abnormality in the
[30684] metabolism of C3 in a patient with increased susceptibility to infection. J.
[30685] Clin. Invest. 49: 1975-1985, 1970.

[30686] 4241.

[30687] Alper, C. A.; Colten, H. R.; Gear, J. S. S.; Rabson, A. R.; Rosen,
[30688] F. S.: Homozygous human C3 deficiency: the role of C3 in antibody
[30689] production, C1s-induced vasopermeability, and cobra venom-induced
[30690] passive hemolysis. J. Clin. Invest. 57: 222-229, 1976.

[30691] 4242.

[30692] Alper, C. A.; Colten, H. R.; Rosen, S. F.; Rabson, A. R.; MacNab,
[30693] G. M.; Gear, J. S. S.: Homozygous deficiency of C3 in a pa-tient with
[30694] repeated infections. Lancet II: 1179-1181, 1972.

[30695] 4243.

[30696] Alper, C. A.; Propp, R. P.: Genetic polymorphism of the

third
[30697] component of human complement (C-prime-3). J. Clin. Invest. 47:
[30698] 2181-2192, 1968.

[30699] 4244.

[30700] Alper, C. A.; Propp, R. P.; Klemperer, M. R.; Rosen, F. S.: Inherited
[30701] deficiency of the third component of human complement (C-prime-3). J.
[30702] Clin. Invest. 48: 553-557, 1969.

[30703] 4245.

[30704] Alper, C. A.; Rosen, F. S.: Studies of a hypomorphic variant of
[30705] human C3. J. Clin. Invest. 50: 324-326, 1971.

[30706] 4246.

[30707] Arvilommi, H.; Berg, K.; Eriksson, A. W.: C3 types and their inheritance
[30708] in Finnish Lapps, Maris (Cheremisses) and Greenland Eskimos. Humangenetik 18:
[30709] 253-259, 1973.

[30710] 4247.

[30711] Azen, E. A.; Smithies, O.: Genetic polymorphism of C-prime-3 (beta-1C-globulin)
[30712] in human serum. Science 162: 905-907, 1968.

[30713] 4248.

[30714] Ball, S.; Buckton, K. E.; Corney, G.; Fey, G.; Monteiro, M.; Noades,
[30715] J. E.; Pym, B.; Robson, E. B.; Tippett, P.: Mapping studies with
[30716] peptidase D (PEPD). (Abstract) Cytogenet. Cell Genet. 37: 411 only,
[30717] 1984.

[30718] 4249.

[30719] Ballow, M.; Shira, J. E.; Harden, L.; Yang, S. Y.; Day, N. K.

[30720] : Complete absence of the third component of complement in man. J.
[30721] Clin. Invest. 56: 703-710, 1975.

[30722] 4250.

[30723] Bauer, J.: Constitutional principles in clinical medicine. Harvey
[30724] Lect. 28: 37-55, 1932.

[30725] 4251.

[30726] Berg, K.; Heiberg, A.: Linkage between familial hyperc-
holesterolemia
[30727] with xanthomatosis and the C3 polymorphism confirmed.
Cytogenet.
[30728] Cell Genet. 22: 621-623, 1978.

[30729] 4252.

[30730] Berg, K.; Heiberg, A.: Linkage studies on familial hyperc-
holesterolemia
[30731] with xanthomatosis: normal lipoprotein markers and the
C3 polymorphism. Cytogenet.
[30732] Cell Genet. 16: 266-270, 1976.

[30733] 4253.

[30734] Berger, M.; Balow, J. E.; Wilson, C. B.; Frank, M. M.: Circu-
lating
[30735] immune complexes and glomerulonephritis in a patient
with congenital
[30736] absence of the third component of complement. New Eng.
J. Med. 308:
[30737] 1009-1012, 1983.

[30738] 4254.

[30739] Borzy, M. S.; Gewurz, A.; Wolff, L.; Houghton, D.; Lovrien, E.

[30740] : Inherited C3 deficiency with recurrent infections and glomerulonephritis. Am.

[30741] J. Dis. Child. 142: 79-83, 1988.

[30742] 4255.

[30743] Botto, M.; Fong, K. Y.; So, A. K.; Barlow, R.; Routier, R.; Morley,

[30744] B. J.; Walport, M. J.: Homozygous hereditary C3 deficiency due to

[30745] a partial gene deletion. Proc. Nat. Acad. Sci. 89: 1957-1961, 1992.

[30746] 4256.

[30747] Arbiser, J. L.; Morton, C. C.; Bruns, G. A. P.; Majzoub, J. A.

[30748] : Human corticotropin releasing hormone gene is located on the long

[30749] arm of chromosome 8. Cytogenet. Cell Genet. 47: 113-116, 1988.

[30750] 4257.

[30751] Behan, D. P.; Heinrichs, S. C.; Troncoso, J. C.; Liu, X.-J.; Kawas,

[30752] C. H.; Ling, N.; De Souza, E. B.: Displacement of corti-
cotropin releasing
[30753] factor from its binding protein as a possible treatment for
Alzheimer's
[30754] disease. Nature 378: 284-287, 1995.

[30755] 4258.

[30756] Cheng, Y.-H.; Nicholson, R. C.; King, B.; Chan, E.-C.; Fit-
ter,
[30757] J. T.; Smith, R.: Corticotropin-releasing hormone gene ex-
pression
[30758] in primary placental cells is modulated by cyclic adenosine
3-prime,5-prime-monophosphate. J.
[30759] Clin. Endocr. Metab. 85: 1239-1244, 2000.

[30760] 4259.

[30761] Habib, K. E.; Weld, K. P.; Rice, K. C.; Pushkas, J.; Cham-
poux,
[30762] M.; Listwak, S.; Webster, E. L.; Atkinson, A. J.; Schulkin, J.;
Contoreggi,
[30763] C.; Chrousos, G. P.; McCann, S. M.; Suomi, S. J.; Higley, J.
D.; Gold,
[30764] P. W.: Oral administration of a corticotropin-releasing
hormone receptor

[30765] antagonist significantly attenuates behavioral, neuroendocrine, and
[30766] autonomic responses to stress in primates. Proc. Nat. Acad. Sci. 97:
[30767] 6079-6084, 2000.

[30768] 4260.

[30769] Inder, W. J.; Prickett, T. C. R.; Ellis, M. J.; Hull, L.; Reid,

[30770] R.; Benny, P. S.; Livesey, J. H.; Donald, R. A.: The utility of plasma
[30771] CRH as a predictor of preterm delivery. J. Clin. Endocr. Metab. 86:
[30772] 5706-5710, 2001.

[30773] 4261.

[30774] Kellogg, J.; Luty, J. A.; Thompson, R.; Luo, X. Y.; Magenis, R.
[30775] E.; Litt, M.: Corticotropin releasing hormone (CRH) maps to human
[30776] chromosome 8 and identifies a TaqI RFLP. Cytogenet. Cell Genet. 51:
[30777] 1022, 1989.

[30778] 4262.

[30779] Knapp, L. T.; Keegan, C. E.; Seasholtz, A. F.; Camper, S. A.:

[30780] Corticotropin-releasing hormone (Crh) maps to mouse chromosome 3. Mammalian

[30781] Genome 4: 615-617, 1993.

[30782] 4263.

[30783] Kyllo, J. H.; Collins, M. M.; Vetter, K. L.; Cuttler, L.; Rosen-field,

[30784] R. L.; Donohoue, P. A.: Linkage of congenital isolated adrenocorticotropic

[30785] hormone deficiency to the corticotropin releasing hor-mone locus using

[30786] simple sequence repeat polymorphisms. Am. J. Med. Genet. 62: 262-267,

[30787] 1996.

[30788] 4264.

[30789] Majzoub, J. A.: Personal Communication. Boston, Mass. 3/3/1995.

[30790] 4265.

[30791] Mandel, H.; Berant, M.; Gotfried, E.; Hochberg, Z.: Autoso-mal

[30792] recessive hypothalamic corticotropin deficiency: a new en-

tity and
[30793] its metabolic consequences. (Abstract) Am. J. Hum. Genet. 47 (suppl.):
[30794] A66, 1990.

[30795] 4266.

[30796] McLean, M.; Bisits, A.; Davies, J.; Woods, R.; Lowry, P.; Smith,
[30797] R.: A placental clock controlling the length of human pregnancy. Nature
[30798] Med. 1: 460-463, 1995.

[30799] 4267.

[30800] Muglia, L.; Jacobson, L.; Dikkes, P.; Majzoub, J. A.: Corti-cotropin-releasing
[30801] hormone deficiency reveals major fetal but not adult glu-cocorticoid
[30802] need. Nature 373: 427-432, 1995.

[30803] 4268.

[30804] Robinson, B. G.; Emanuel, R. L.; Frim, D. M.; Majzoub, J. A.:
[30805] Glucocorticoid stimulates expression of corticotropin-releas-ing hormone

[30806] gene in human placenta. Proc. Nat. Acad. Sci. 85:
5244-5248, 1988.
[30807] 4269.
[30808] Scatena, C. D.; Adler, S.: Trans-acting factors dictate the
species-specific
[30809] placental expression of corticotropin-releasing factor
genes in choriocarcinoma
[30810] cell lines. Endocrinology 137: 3000-3008, 1996.
[30811] 4270.
[30812] Scatena, C. D.; Adler, S.: Characterization of a human-
specific
[30813] regulator of placental corticotropin-releasing hormone.
Molec. Endocr. 12:
[30814] 1228-1240, 1998.
[30815] 4271.
[30816] Shibahara, S.; Morimoto, Y.; Furutani, Y.; Notake, M.;
Takahashi,
[30817] H.; Shimizu, S.; Horikawa, S.; Numa, S.: Isolation and se-
quence analysis
[30818] of the human corticotropin-releasing factor precursor
gene. EMBO

[30819] J. 2: 775-779, 1983.

[30820] 4272.

[30821] Stratakis, C. A.; Chrousos, G. P.: Neuroendocrinology and pathophysiology

[30822] of the stress system. Ann. N.Y. Acad. Sci. 771: 1-18, 1995.

[30823] 4273.

[30824] Stratakis, C. A.; Sarlis, N. J.; Berrettini, W. H.; Badner, J.

[30825] A.; Chrousos, G. P.; Gershon, E. S.; Detera-Wadleigh, S. D.: Lack

[30826] of linkage between the corticotropin-releasing hormone (CRH) gene

[30827] and bipolar affective disorder. Molec. Psychiat. 2: 483-485, 1997.

[30828] 4274.

[30829] Xu, B.; Sano, T.; Yamada, S.; Li, C. C.; Hirokawa, M.: Ex-pression

[30830] of corticotropin-releasing hormone messenger ribonucleic acid in human

[30831] pituitary corticotroph adenomas associated with prolifera-tive potential. J.

[30832] Clin. Endocr. Metab. 85: 1220-1225, 2000.

[30833] 4275.

[30834] Zouboulis, C. C.; Seltmann, H.; Hiroi, N.; Chen, W.; Young, M.;

[30835] Oeff, M.; Scherbaum, W. A.; Orfanos, C. E.; McCann, S. M.; Bornstein,

[30836] S. R.: Corticotropin-releasing hormone: an autocrine hormone that

[30837] promotes lipogenesis in human sebocytes. Proc. Nat. Acad. Sci. 99:

[30838] 7148-7153, 2002.

[30839] 4276.

[30840] Chen, R.; Lewis, K. A.; Perrin, M. H.; Vale, W. W.: Expression

[30841] cloning of a human corticotropin-releasing-factor receptor. Proc.

[30842] Nat. Acad. Sci. 90: 8967-8971, 1993.

[30843] 4277.

[30844] Dieterich, K. D.; Gundelfinger, E. D.; Ludecke, D. K.; Lehnert,

[30845] H.: Mutation and expression analysis of corticotropin-releas-

ing factor
[30846] 1 receptor in adrenocorticotropin-secreting pituitary adenomas. J.
[30847] Clin. Endocr. Metab. 83: 3327-3331, 1998.
[30848] 4278.
[30849] Grammatopoulos, D.; Dai, Y.; Chen, J.; Karteris, E.; Papadopoulou,
[30850] N.; Easton, A. J.; Hillhouse, E. W.: Human corticotropin-releasing
[30851] hormone receptor: differences in subtype expression between pregnant
[30852] and nonpregnant myometria. J. Clin. Endocr. Metab. 83: 2539-2544,
[30853] 1998.
[30854] 4279.
[30855] Leproult, R.; Colecchia, E. F.; L'Hermite-Baleriaux, M.; Van Cauter,
[30856] E.: Transition from dim to bright light in the morning induces an
[30857] immediate elevation of cortisol levels. J. Clin. Endocr. Metab. 86:
[30858] 151-157, 2001.

[30859] 4280.

[30860] Liaw, C. W.; Grigoriadis, D. E.; Lovenberg, T. W.; De Souza, E.

[30861] B.; Maki, R. A.: Localization of ligand-binding domains of human

[30862] corticotropin-releasing factor receptor: a chimeric receptor approach. Molec.

[30863] Endocr. 11: 980-985, 1997.

[30864] 4281.

[30865] Ion, A.; Crosby, A. H.; Kremer, H.; Kenmochi, N.; Van Reen, M.;

[30866] Fenske, C.; Van Der Burgt, I.; Brunner, H. G.; Montgomery, K.: Detailed

[30867] mapping, mutation analysis, and intragenic polymorphism identification

[30868] in candidate Noonan syndrome genes MYL2, DCN, EPS8, and RPL6. J.

[30869] Med. Genet. 37: 884-886, 2000.

[30870] 4282.

[30871] Comer, K. A.; Falany, J. L.; Falany, C. N.: Cloning and expression

[30872] of human liver dehydroepiandrosterone sulphotransferase. Biochem.
[30873] J. 289: 233-240, 1993.
[30874] 4283.
[30875] Durocher, F.; Morissette, J.; Dufort, I.; Simard, J.; Luu-The,
[30876] V.: Genetic linkage mapping of the dehydroepiandrosterone sulfotransferase
[30877] (STD) gene on the chromosome 19q13.3 region. Genomics 29: 781-783,
[30878] 1995.
[30879] 4284.
[30880] Kong, A.-N. T.; Yang, L.; Ma, M.; Tao, D.; Bjornsson, T. D.: Molecular
[30881] cloning of the alcohol/hydroxysteroid form (hSTa) of sulfotransferase
[30882] from human liver. Biochem. Biophys. Res. Commun. 187: 448-454, 1992.
[30883] 4285.
[30884] Otterness, D.; Mohrenweiser, H. W.; Brandriff, B. F.; Weinshilboum,
[30885] R. M.: Dehydroepiandrosterone sulfotransferase gene

(STD): localization
[30886] to human chromosome 19q13.3. Cytogenet. Cell Genet. 70: 45-47, 1995.
[30887] 4286.
[30888] Otterness, D. M.; Her, C.; Aksoy, S.; Kimura, S.; Wieben, E. D.;
[30889] Weinshilboum, R. M.: Human dehydroepiandrosterone sulfotransferase
[30890] gene: molecular cloning and structural characterization. DNA Cell
[30891] Biol. 14: 331-341, 1995.
[30892] 4287.
[30893] Otterness, D. M.; Wieben, E. D.; Wood, T. C.; Watson, W. G.; Madden,
[30894] B. J.; McCormick, D. J.; Weinshilboum, R. M.: Human liver dehydroepiandrosterone
[30895] sulfotransferase: molecular cloning and expression of cDNA. Molec.
[30896] Pharm. 41: 865-872, 1992.
[30897] 4288.
[30898] Schulte-Korne, G.; Grimm, T.; Nothen, M. M.; Muller-

Myhsok, B.;

[30899] Cichon, S.; Vogt, I. R.; Propping, P.; Remschmidt, H.: Evi-
dence for

[30900] linkage of spelling disability to chromosome 15. (Letter)
Am. J.

[30901] Hum. Genet. 63: 279-282, 1998.

[30902] 4289.

[30903] Yoon, J.-W.; Yoon, C.-S.; Lim, H.-W.; Huang, Q. Q.; Kang,
Y.;

[30904] Pyun, K. H.; Hirasawa, K.; Sherwin, R. S.; Jun, H.-S.: Con-
trol of

[30905] autoimmune diabetes in NOD mice by GAD expression or
suppression in

[30906] beta cells. Science 284: 1183-1187, 1999.

[30907] 4290.

[30908] Elgadi, K. M.; Meguid, R. A.; Qian, M.; Souba, W. W.; Ab-
couwer,

[30909] S. F.: Cloning and analysis of unique human glutaminase
isoforms

[30910] generated by tissue-specific alternative splicing. Physiol.
Genomics 31:

[30911] 51-62, 1999.

[30912] 4291.

[30913] Mock, B.; Kozak, C.; Seldin, M. F.; Ruff, N.; D'Hoostelaere, L.;

[30914] Szpirer, C.; Levan, G.; Seuanez, H.; O'Brien, S.; Banner, C.: A glutaminase

[30915] (Gls) gene maps to mouse chromosome 1, rat chromosome 9 and human

[30916] chromosome 2. Genomics 5: 291-297, 1989. Note: Erratum: Genomics

[30917] 5: 957 only, 1989.

[30918] 4292.

[30919] Modi, W. S.; Pollock, D. D.; Mock, B. A.; Banner, C.; Renauld,

[30920] J.-C.; Van Snick, J.: Regional localization of the human glutaminase

[30921] (GLS) and interleukin-9 (IL9) genes by in situ hybridization. Cytogenet.

[30922] Cell Genet. 57: 114-116, 1991.

[30923] 4293.

[30924] Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hirosawa, M.;

[30925] Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Oharo, O.: Prediction
[30926] of the coding sequences of unidentified human genes. XII. The complete
[30927] sequences of 100 new cDNA clones from brain which code for large proteins
[30928] in vitro. DNA Res. 5: 355-364, 1998.

[30929] 4294.

[30930] Prusiner, S. B.: Disorders of glutamate metabolism and neurological
[30931] dysfunction. Ann. Rev. Med. 32: 512-542, 1981.

[30932] 4295.

[30933] Sahai, S.: Glutaminase in human platelets. Clin. Chim. Acta 127:
[30934] 197-203, 1983.

[30935] 4296.

[30936] Sahai, S.; Vogel, F.: Genetic control of platelet glutami-
nase:
[30937] a twin study. Hum. Genet. 63: 292-293, 1983.

[30938] 4297.

[30939] Clancy, K. P.; Berger, R.; Cox, M.; Bleskan, J.; Walton, K. A.;

[30940] Hart, I.; Patterson, D.: Localization of the L-glutamine synthetase

[30941] gene to chromosome 1q23. Genomics 38: 418-420, 1996.

[30942] 4298.

[30943] Gibbs, C. S.; Campbell, K. E.; Wilson, R. H.: Sequence of a human

[30944] glutamine synthetase cDNA. Nucleic Acids Res. 15: 6293 only, 1987.

[30945] 4299.

[30946] Gunnersen, D.; Haley, B.: Detection of glutamine synthetase in

[30947] the cerebrospinal fluid of Alzheimer diseased patients: a potential

[30948] diagnostic biochemical marker. Proc. Nat. Acad. Sci. 89: 11949-11953,

[30949] 1992.

[30950] 4300.

[30951] Helou, K.; Das, A. T.; Lamers, W. H.; Hoovers, J. M. N.; Szpirer,

[30952] C.; Szpirer, J.; Klinga-Levan, K.; Levan, G.: FISH mapping of three
[30953] ammonia metabolism genes (Glul, Cps1, Glud1) in rat, and the chromosomal
[30954] localization of GLUL in human and Cps1 in mouse. Mammalian Genome 8:
[30955] 362-364, 1997.

[30956] 4301.

[30957] Pesole, G.; Bozzetti, M. P.; Lanave, C.; Preparata, G.; Saccone,
[30958] C.: Glutamine synthetase gene evolution: a good molecular clock. Proc.
[30959] Nat. Acad. Sci. 88: 522-526, 1991.

[30960] 4302.

[30961] Wang, Y.; Kudoh, J.; Kubota, R.; Asakawa, S.; Minoshima, S.; Shimizu,
[30962] N.: Chromosomal mapping of a family of human glutamine synthetase
[30963] genes: functional gene (GLUL) on 1q25, pseudogene (GLULP) on 9p13,
[30964] and three related genes (GLULL1, GLULL2, GLULL3) on 5q33, 11p15, and

[30965] 11q24. Genomics 37: 195-199, 1996.

[30966] 4303.

[30967] Oki, T.; Yamazaki, K.; Kuromitsu, J.; Okada, M.; Tanaka, I.: cDNA

[30968] cloning and mapping of a novel subtype of glutamine:fructose-6-phosphate

[30969] amidotransferase (GFAT2) in human and mouse. Genomics 57: 227-234,

[30970] 1999.

[30971] 4304.

[30972] Cerini, C.; Kerjan, P.; Astier, M.; Gratecos, D.; Mirande, M.;

[30973] Semeriva, M.: A component of the multisynthetase complex is a multifunctional

[30974] aminoacyl-tRNA synthetase. EMBO J. 10: 4267-4277, 1991.

[30975] 4305.

[30976] Fett, R.; Knippers, R.: The primary structure of human glutaminyl-tRNA

[30977] synthetase: a highly conserved core, amino acid repeat regions, and

[30978] homologies with translation elongation factors. J. Biol.

Chem. 266:
[30979] 1448-1455, 1991.

[30980] 4306.

[30981] Kaiser, E.; Hu, B.; Becher, S.; Eberhard, D.; Schray, B.; Baack,
[30982] M.; Hameister, H.; Knippers, R.: The human EPRS locus (formerly the
[30983] QARS locus): a gene encoding a class I and a class II aminoacyl-tRNA
[30984] synthetase. Genomics 19: 280-290, 1994.

[30985] 4307.

[30986] Kunze, N.; Bittler, E.; Fett, R.; Schray, B.; Hameister, H.; Wiedorn,
[30987] K. H.; Knippers, R.: The human QARS locus: assignment of the human
[30988] gene for glutaminyl-tRNA synthetase to chromosome 1q32-42. Hum. Genet. 85:
[30989] 527-530, 1990.

[30990] 4308.

[30991] Hoglund, P.; Haila, S.; Socha, J.; Tomaszewski, L.; Saari-alho-Kere,

[30992] U.; Karjalainen-Lindsberg, M.-L.; Airola, K.; Holmberg, C.; de la

[30993] Chapelle, A.; Kere, J.: Mutations of the down-regulated in adenoma

[30994] (DRA) gene cause congenital chloride diarrhoea. Nature Genet. 14:

[30995] 316-319, 1996.

[30996] 4309.

[30997] Horisberger, M. A.; Wathelet, M.; Szpirer, J.; Szpirer, C.; Islam,

[30998] Q.; Levan, G.; Huez, G.; Content, J.: cDNA cloning and assignment

[30999] to chromosome 21 of IFI-78K gene, the human equivalent of murine Mx

[31000] gene. Somat. Cell Molec. Genet. 14: 123-131, 1988.

[31001] 4310.

[31002] Huber, P.; Aebi, M.; Grob, R.; Pravtcheva, D.; Ruddle, F.; Haller,

[31003] O.: Chromosomal localization of two human Mx genes. (Abstract) Experientia 44:

[31004] A84 only, 1988.

[31005] 4311.

[31006] Li, Y.; Youssoufian, H.: MxA overexpression reveals a common genetic

[31007] link in four Fanconi anemia complementation groups. J. Clin. Invest. 100:

[31008] 2873-2880, 1997.

[31009] 4312.

[31010] Hoglund, P.; Sormaala, M.; Haila, S.; Socha, J.; Rajaram, U.; Scheurlen,

[31011] W.; Sinaasappel, M.; de Jonge, H.; Holmberg, C.; Yoshikawa, H.; Kere,

[31012] J.: Identification of seven novel mutations including the first two

[31013] genomic rearrangements in SLC26A3 mutated in congeni- tal chloride diarrhea. Hum.

[31014] Mutat. 18: 233-242, 2001.

[31015] 4313.

[31016] Schweinfest, C. W.; Henderson, K. W.; Suster, S.; Kondoh, N.; Papas,

[31017] T. S.: Identification of a colon mucosa gene that is down- regulated

[31018] in colon adenomas and adenocarcinomas. Proc. Nat. Acad. Sci. 90:
[31019] 4166-4170, 1993.

[31020] 4314.

[31021] Taguchi, T.; Testa, J. R.; Papas, T. S.; Schweinfest, C.: Localization
[31022] of a candidate colon tumor-suppressor gene (DRA) to 7q22-q31.1 by
[31023] fluorescence in situ hybridization. Genomics 20: 146-147, 1994.
[31024] 4315.

[31025] Yoshikawa, H.; Watanabe, T.; Abe, T.; Sato, M.; Oda, Y.: Japanese
[31026] siblings with congenital chloride diarrhea. Pediat. Int. 42: 313-315,
[31027] 2000.

[31028] 4316.

[31029] McAlpine, P. J.; Kaita, H.; Lewis, M.: Is the DIA-1 locus linked
[31030] to the P blood group locus? Cytogenet. Cell Genet. 22: 629-632,

[31031] 1978.

[31032] 4317.

[31033] Winqvist, R.; Lundstrom, K.; Salminen, M.; Laatikainen, M.; Ulmanen,

[31034] I.: Mapping of human catechol-O-methyltransferase gene to 22q11.2

[31035] and detection of a frequent RFLP with BglI. (Abstract) Cytogenet.

[31036] Cell Genet. 58: 2051 only, 1991.

[31037] 4318.

[31038] Bu, L.; Jin, Y.; Shi, Y.; Chu, R.; Ban, A.; Eiberg, H.; Andres,

[31039] L.; Jiang, H.; Zheng, G.; Qian, M.; Cui, B.; Xia, Y.; Liu, J.; Hu,

[31040] L.; Zhao, G.; Hayden, M. R.; Kong, X.: Mutant DNA-binding domain

[31041] of HSF4 is associated with autosomal dominant lamellar and Marner

[31042] cataract. Nature Genet. 31: 276-278, 2002.

[31043] 4319.

[31044] Eiberg, H.; Marner, E.; Rosenberg, T.; Mohr, J.: Marner's cataract

[31045] (CAM) assigned to chromosome 16: linkage to haptoglobin. Clin. Genet. 34:
[31046] 272-275, 1988.

[31047] 4320.

[31048] Marner, E.: A family with eight generations of hereditary cataract. Acta
[31049] Ophthal. 27: 537-551, 1949.

[31050] 4321.

[31051] Furukawa, Y.; Nakatsuru, S.; Nagafuchi, A.; Tsukita, S.; Muto,
[31052] T.; Nakamura, Y.; Horii, A.: Structure, expression and chromosome
[31053] assignment of the human catenin (cadherin-associated protein) alpha
[31054] 1 gene (CTNNA1). Cytogenet. Cell Genet. 65: 74-78, 1994.
[31055] 4322.

[31056] Hirano, S.; Kimoto, N.; Shimoyama, Y.; Hirohashi, S.; Takeichi,
[31057] M.: Identification of a neural alpha-catenin as a key regulator of

[31058] cadherin function and multicellular organization. Cell 70: 293-301,

[31059] 1992.

[31060] 4323.

[31061] McPherson, J. D.; Morton, R. A.; Ewing, C. M.; Wasmuth, J. J.;

[31062] Overhauser, J.; Nagafuchi, A.; Tsukita, S.; Isaacs, W. B.: Assignment

[31063] of the human alpha-catenin gene (CTNNA1) to chromosome 5q21-q22. Genomics 19:

[31064] 188-190, 1994.

[31065] 4324.

[31066] Nagafuchi, A.; Takeichi, M.; Tsukita, S.: The 102 kd cadherin-associated

[31067] protein: similarity to vinculin and posttranscriptional regulation

[31068] of expression. Cell 65: 849-857, 1991.

[31069] 4325.

[31070] Nollet, F.; van Hengel, J.; Berx, G.; Molemans, F.; van Roy, F.

[31071] : Isolation and characterization of a human pseudogene

(CTNNAP1) for

[31072] alpha-E-catenin (CTNNA1): assignment of the pseudo-gene to 5q22 and

[31073] the alpha-E-catenin gene to 5q31. Genomics 26: 410-413, 1995.

[31074] 4326.

[31075] Oda, T.; Kanai, Y.; Shimoyama, Y.; Nagafuchi, A.; Tsukita, S.;

[31076] Hirohashi, S.: Cloning of the human alpha-catenin cDNA and its aberrant

[31077] mRNA in a human cancer cell line. Biochem. Biophys. Res. Commun. 193:

[31078] 897-904, 1993.

[31079] 4327.

[31080] Pokutta, S.; Weis, W. I.: Structure of the dimerization and beta-catenin-binding

[31081] region of alpha-catenin. Molec. Cell 5: 533-543, 2000.

[31082] 4328.

[31083] Shimoyama, Y.; Nagafuchi, A.; Fujita, S.; Gotoh, M.; Take-ichi,

[31084] M.; Tsukita, S.; Hirohashi, S.: Cadherin dysfunction in a

human cancer
[31085] cell line: possible involvement of loss of alpha-catenin expression
[31086] in reduced cell-cell adhesiveness. Cancer Res. 52: 5770-5774, 1992.
[31087] 4329.
[31088] Vasioukhin, V.; Bauer, C.; Degenstein, L.; Wise, B.; Fuchs, E.
[31089] : Hyperproliferation and defects in epithelial polarity upon conditional
[31090] ablation of alpha-catenin in skin. Cell 104: 605-617, 2001.
[31091] 4330.
[31092] Bailey, A.; Norris, A. L.; Leek, J. P.; Clissold, P. M.; Carr,
[31093] I. M.; Ogilvie, D. J.; Morrison, J. F. J.; Meredith, D. M.; Markham,
[31094] A. F.: Yeast artificial chromosome cloning of the beta-catenin locus
[31095] on human chromosome 3p21-22. Chromosome Res. 3: 201-203, 1995.
[31096] 4331.
[31097] Blaker, H.; Hofmann, W. J.; Rieker, R. J.; Penzel, R.; Graf,

M.;

[31098] Otto, H. F.: Beta-catenin accumulation and mutation of the CTNNB1

[31099] gene in hepatoblastoma. Genes Chromosomes Cancer 25: 399-402, 1999.

[31100] 4332.

[31101] Chan, T. A.; Wang, Z.; Dang, L. H.; Vogelstein, B.; Kinzler, K.

[31102] W.: Targeted inactivation of CTNNB1 reveals unexpected effects of

[31103] beta-catenin mutation. Proc. Nat. Acad. Sci. 99: 8265-8270, 2002.

[31104] 4333.

[31105] Chenn, A.; Walsh, C. A.: Regulation of cerebral cortical size

[31106] by control of cell cycle exit in neural precursors. Science 297:

[31107] 365-369, 2002.

[31108] 4334.

[31109] Eastman, Q.; Grosschedl, R.: Regulation of LEF-1/TCF transcription

[31110] factors by Wnt and other signals. Curr. Opin. Cell Biol. 11:

233-240,

[31111] 1999.

[31112] 4335.

[31113] Gat, U.; DasGupta, R.; Degenstein, L.; Fuchs, E.: De novo hair

[31114] follicle morphogenesis and hair tumors in mice express-ing a truncated

[31115] beta-catenin in skin. Cell 95: 605-614, 1998.

[31116] 4336.

[31117] Groden, J.; Thliveris, A.; Samowitz, W.; Carlson, M.; Gel-bert,

[31118] L.; Albertsen, H.; Joslyn, G.; Stevens, J.; Spirio, L.; Robert-son,

[31119] M.; Sargeant, L.; Krapcho, K.; Wolff, E.; Burt, R.; Hughes, J. P.;

[31120] Warrington, J.; McPherson, J.; Wasmuth, J.; Le Paslier, D.; Abderrahim,

[31121] H.; Cohen, D.; Leppert, M.; White, R.: Identification and characterization

[31122] of the familial adenomatous polyposis coli gene. Cell 66: 589-600,

[31123] 1991.

[31124] 4337.

[31125] Walport, M. J.: Complement (first of two parts). New Eng. J.

[31126] Med. 344: 1058-1066, 2001.

[31127] 4338.

[31128] White, R. T.; Damm, D.; Hancock, N.; Rosen, B. S.; Lowell,
B.

[31129] B.; Usher, P.; Flier, J. S.; Spiegelman, B. M.: Human adipsin
is

[31130] identical to complement factor D and is expressed at high
levels in

[31131] adipose tissue. J. Biol. Chem. 267: 9210-9213, 1992.

[31132] 4339.

[31133] Diaz-Guillen, M. A.; Rodriguez de Cordoba, S.; Heine-
Suner, D.

[31134] : A radiation hybrid map of complement factor H and fac-
tor H-related

[31135] genes. Immunogenetics 49: 549-552, 1999.

[31136] 4340.

[31137] Atkinson, J. P.: Personal Communication. St. Louis, Mo.
3/7/1983.

[31138] 4341.

[31139] Curry, R. A.; Dierich, M. P.; Pellegrino, M. A.; Hoch, H. A.:

[31140] Evidence for linkage between HLA antigens and receptors for complement

[31141] components C3b and C3d in human-mouse hybrids. Immunogenetics 3:

[31142] 465-471, 1976.

[31143] 4342.

[31144] Dykman, T. R.; Cole, J. L.; Iida, K.; Atkinson, J. P.: Polymorphism

[31145] of human erythrocyte C3b/C4b receptor. Proc. Nat. Acad. Sci. 80:

[31146] 1698-1702, 1983.

[31147] 4343.

[31148] Dykman, T. R.; Cole, J. L.; Iida, K.; Atkinson, J. P.: Structural

[31149] heterogeneity of the C3b/C4b receptor (CR1) on human peripheral blood

[31150] cells. J. Exp. Med. 157: 2160-2165, 1983.

[31151] 4344.

[31152] Dykman, T. R.; Hatch, J. A.; Atkinson, J. P.: Polymorphism of
[31153] the human C3b/C4b receptor: identification of a third allele and analysis
[31154] of receptor phenotypes in families and patients with systemic lupus
[31155] erythematosus. J. Exp. Med. 159: 691-703, 1984.

[31156] 4345.

[31157] Gerdes, J.; Hansmann, M.-L.; Stein, H.; Naiem, M.; Mason, D. Y.
[31158] : Ultrastructural localization of human complement C3b receptors in
[31159] the human kidney as determined by immunoperoxidase staining with the
[31160] monoclonal antibody C3RTo5. Virchows Arch. B 40: 1-7, 1982.
[31161] 4346.

[31162] Hing, S.; Day, A. J.; Linton, S. J.; Ripoche, J.; Sim, R. B.; Reid,
[31163] K. B.; Solomon, E.: Assignment of complement components C4 binding
[31164] protein (C4BP) and factor H (FH) to human chromosome

1q, using cDNA
[31165] probes. Ann. Hum. Genet. 52: 117-122, 1988.

[31166] 4347.

[31167] Holers, V. M.; Chaplin, D. D.; Leykam, J. F.; Gruner, B. A.; Kumar,
[31168] V.; Atkinson, J. P.: Human complement C3b/C4b receptor (CR1) mRNA
[31169] polymorphism that correlates with the CR1 allelic molecular weight
[31170] polymorphism. Proc. Nat. Acad. Sci. 84: 2459-2463, 1987.
[31171] 4348.

[31172] Iida, K.; Momaghi, R.; Nussenzweig, V.: Complement receptor (CR1)
[31173] deficiency in erythrocytes from patients with systemic lupus erythematosus. J.
[31174] Exp. Med. 155: 1427-1438, 1982.

[31175] 4349.

[31176] Jozsi, M.; Prechl, J.; Bajtay, Z.; Erdei, A.: Complement receptor
[31177] type 1 (CD35) mediates inhibitory signals in human B lym-

phocytes. J.

[31178] Immun. 168: 2782-2788, 2002.

[31179] 4350.

[31180] Miyakawa, Y.; Yamada, A.; Kosaka, K.; Tsuda, F.; Kosugi, E.; Mayumi,

[31181] M.: Defective immune-adherence (C3b) receptor on erythrocytes from

[31182] patients with systemic lupus erythematosus. Lancet II: 493-497,

[31183] 1981.

[31184] 4351.

[31185] Moldenhauer, F.; David, J.; Fielder, A. H. L.; Lachmann, P. J.;

[31186] Walport, M. J.: Inherited deficiency of erythrocyte complement receptor

[31187] type 1 does not cause susceptibility to systemic lupus erythematosus. Arthritis

[31188] Rheum. 30: 961-966, 1987.

[31189] 4352.

[31190] Nelson, R. A.: The immune-adherence phenomenon: an immunologically

[31191] specific reaction between microorganisms and erythrocytes leading
[31192] to enhanced phagocytosis. Science 118: 733-737, 1953.

[31193] 4353.

[31194] Nowak, J. S.: Genetic variability of complement receptor on human
[31195] erythrocytes. J. Genet. 66: 133-138, 1987.

[31196] 4354.

[31197] Ohi, H.; Ikezawa, T.; Watanabe, S.; Seki, M.; Mizutani, Y.; Nawa,
[31198] N.; Hatano, M.: Two cases of mesangiocapillary glomerulonephritis
[31199] with CR1 deficiency. (Letter) Nephron 43: 307 only, 1986.

[31200] 4355.

[31201] Smith, B. O.; Mallin, R. L.; Krych-Goldberg, M.; Wang, X.; Hauhart,
[31202] R. E.; Bromek, K.; Uhrin, D.; Atkinson, J. P.; Barlow, P. N.: Structure
[31203] of the C3b binding site of CR1 (CD35), the immune adherence receptor. Cell 108:
[31204] 769-780, 2002.

[31205] 4356.

[31206] Asakura, H.; Zwain, I. H.; Yen, S. S. C.: Expression of genes

[31207] encoding corticotropin-releasing factor (CRF), type 1 CRF receptor,

[31208] and CRF-binding protein and localization of the gene products in the

[31209] human ovary. J. Clin. Endocr. Metab. 82: 2720-2725, 1997.

[31210] 4357.

[31211] Lynch, E. D.; Lee, M. K.; Morrow, J. E.; Welcsh, P. L.; Leon, P.

[31212] E.; King, M.-C.: Nonsyndromic deafness DFNA1 associated with mutation

[31213] of the human homolog of the Drosophila gene di-aphanous. Science 278:

[31214] 1315-1318, 1997.

[31215] 4358.

[31216] Rowley, P. T.: Familial hearing loss associated with branchial

[31217] fistulas. Pediatrics 44: 978-985, 1969.

[31218] 4359.

[31219] Ganz, T.; Lehrer, R. I.: Defensins. Pharm. Ther. 66:
191-205,
[31220] 1995.

[31221] 4360.

[31222] Liu, L.; Zhao, C.; Heng, H. H. Q.; Ganz, T.: The human
beta-defensin-1
[31223] and alpha-defensins are encoded by adjacent genes: two
peptide families
[31224] with differing disulfide topology share a common ances-
try. Genomics 43:
[31225] 316-320, 1997.

[31226] 4361.

[31227] Beischlag, T. V.; Marchese, A.; Meador-Woodruff, J. H.;
Damask,
[31228] S. P.; O'Dowd, B. F.; Tyndale, R. F.; Van Tol, H. H. M.; See-
man, P.;
[31229] Niznik, H. B.: The human dopamine D5 receptor gene:
cloning and characterization
[31230] of the 5-prime flanking and promoter region. Biochem-
istry 34: 5960-5970,
[31231] 1995.

[31232] 4362.

[31233] Eubanks, J. H.; Altherr, M.; Wagner-McPherson, C.;
McPherson, J.
[31234] D.; Wasmuth, J. J.; Evans, G. A.: Localization of the D5
dopamine
[31235] receptor gene to human chromosome 4p15.1-p15.3, cen-
tromeric to the
[31236] Huntington's disease locus. Genomics 12: 510-516, 1992.

[31237] 4363.

[31238] Grandy, D. K.; Allen, L. J.; Zhang, Y.; Magenis, R. E.; Civelli,
[31239] O.: Chromosomal localization of three human D5
dopamine receptor
[31240] genes. Genomics 13: 968-973, 1992.

[31241] 4364.

[31242] Grandy, D. K.; Zhang, Y.; Bouvier, C.; Zhou, Q.-Y.; John-
son, R.
[31243] A.; Allen, L.; Buck, K.; Bunzow, J. R.; Salon, J.; Civelli, O.:
Multiple
[31244] human D5 dopamine receptor genes: a functional receptor
and two pseudogenes. Proc.
[31245] Nat. Acad. Sci. 88: 9175-9179, 1991.

[31246] 4365.

[31247] Grosson, C. L. S.; MacDonald, M. E.; Duyao, M. P.; Am-
brose, C.
[31248] M.; Roffler-Tarlov, S.; Gusella, J. F.: Synteny conservation
of the
[31249] Huntington's disease gene and surrounding loci on mouse
chromosome
[31250] 5. Mammalian Genome 5: 424-428, 1994.

[31251] 4366.

[31252] Liu, F.; Wan, Q.; Pristupa, Z. B.; Yu, X.-M.; Wang, Y. T.;
Niznik,
[31253] H. B.: Direct protein-protein coupling enables cross-talk
between
[31254] dopamine D5 and gamma-aminobutyric acid A receptors.
Nature 403:
[31255] 274-280, 2000.

[31256] 4367.

[31257] Misbahuddin, A.; Placzek, M. R.; Chaudhuri, K. R.; Wood,
N. W.;
[31258] Bhatia, K. P.; Warner, T. T.: A polymorphism in the
dopamine receptor

[31259] DRD5 is associated with blepharospasm. Neurology 58: 124-126, 2002.

[31260] 4368.

[31261] Placzek, M. R.; Misbahuddin, A.; Chaudhuri, K. R.; Wood, N. W.;

[31262] Bhatia, K. P.; Warner, T. T.: Cervical dystonia is associated with

[31263] a polymorphism in the dopamine (D5) receptor gene. J. Neurol. Neurosurg.

[31264] Psychiat. 71: 262-264, 2001.

[31265] 4369.

[31266] Polymeropoulos, M. H.; Xiao, H.; Merril, C. R.: The human D5 dopamine

[31267] receptor (DRD5) maps on chromosome 4. Genomics 11: 777-778, 1991.

[31268] 4370.

[31269] Sherrington, R.; Mankoo, B.; Attwood, J.; Kalsi, G.; Curtis, D.;

[31270] Buetow, K.; Povey, S.; Gurling, H.: Cloning of the human dopamine

[31271] D5 receptor gene and identification of a highly polymorphic microsatellite

[31272] for the DRD5 locus that shows tight linkage to the chromosome 4p reference

[31273] marker RAF1P1. Genomics 18: 423-425, 1993.

[31274] 4371.

[31275] Tiberi, M.; Jarvie, K. R.; Silvia, C.; Falardeau, P.; Gingrich,

[31276] J. A.; Godinot, N.; Bertrand, L.; Yang-Feng, T. L.; Fremeau, R. T.,

[31277] Jr.; Caron, M. G.: Cloning, molecular characterization, and chromosomal

[31278] assignment of a gene encoding a second D-1 dopamine receptor subtype:

[31279] differential expression pattern in rat brain compared with the D-1A

[31280] receptor. Proc. Nat. Acad. Sci. 88: 7491-7495, 1991.

[31281] 4372.

[31282] Weinshank, R. L.; Adham, N.; Macchi, M.; Olsen, M. A.; Branchek,

[31283] T. A.; Hartig, P. R.: Molecular cloning and characterization of a

[31284] high affinity dopamine receptor (D-1-beta) and its pseudogene. J.

[31285] Biol. Chem. 266: 22427-22435, 1991.

[31286] 4373.

[31287] Byerley, W.; Hoff, M.; Holik, J.; Caron, M. G.; Giros, B.: VNTR

[31288] polymorphism for the human dopamine transporter gene (DAT1). Hum.

[31289] Molec. Genet. 2: 335, 1993.

[31290] 4374.

[31291] Cook, E. H., Jr.; Stein, M. A.; Krasowski, M. D.; Cox, N. J.; Olkon,

[31292] D. M.; Kieffer, J. E.; Leventhal, B. L.: Association of atten-tion-deficit

[31293] disorder and the dopamine transporter gene. Am. J. Hum. Genet. 56:

[31294] 993-998, 1995.

[31295] 4375.

[31296] Doucette-Stamm, L.; Blakely, D. J.; Tian, J.; Mockus, S.; Mao,

[31297] J.: Population genetic study of the human dopamine transporter gene

[31298] (DAT1). Genet. Epidemiol. 12: 303-308, 1995.

[31299] 4376.

[31300] Gainetdinov, R. R.; Wetsel, W. C.; Jones, S. R.; Levin, E. D.;

[31301] Jaber, M.; Caron, M. G.: Role of serotonin in the paradoxi-
cal calming

[31302] effect of psychostimulants on hyperactivity. Science 283:
397-401,

[31303] 1999.

[31304] 4377.

[31305] Gelernter, J.; Vandenbergh, D.; Kruger, S. D.; Pauls, D. L.;
Kurlan,

[31306] R.; Pakstis, A. J.; Kidd, K. K.; Uhl, G.: The dopamine trans-
porter

[31307] protein gene (SLC6A3): primary linkage mapping and link-
age studies

[31308] in Tourette syndrome. Genomics 30: 459-463, 1995.

[31309] 4378.

[31310] Gill, M.; Daly, G.; Heron, S.; Hawi, Z.; Fitgerald, M.: Confir-
mation

[31311] of association between attention deficit hyperactivity dis-
order and

[31312] a dopamine transporter polymorphism. Molec. Psychiat. 2:

311-313,

[31313] 1997.

[31314] 4379.

[31315] Giros, B.; El Mestikawy, S.; Godinot, N.; Zheng, K.; Han, H.; Yang-Feng,

[31316] T.; Caron, M. G.: Cloning, pharmacological characterization, and

[31317] chromosome assignment of the human dopamine transporter. Molec. Pharm. 42:

[31318] 383-390, 1992.

[31319] 4380.

[31320] Giros, B.; Jaber, M.; Jones, S. R.; Wightman, R. M.; Caron, M.

[31321] G.: Hyperlocomotion and indifference to cocaine and amphetamine in

[31322] mice lacking the dopamine receptor. Nature 370:
606-612, 1996.

[31323] 4381.

[31324] Goldman, D.: Dopamine transporter, alcoholism and other diseases. Nature

[31325] Med. 1: 624-625, 1995.

[31326] 4382.

[31327] Korn, W. T.; Schatzki, S. C.; DiSciullo, A. J.; Scully, R. E.

[31328] : Papillary cystadenoma of the broad ligament in von Hip-
pel-Lindau
[31329] disease. Am. J. Obstet. Gynec. 163: 596-598, 1990.

[31330] 4383.

[31331] Isnard, R.; Kalotka, H.; Durr, A.; Cossee, M.; Schmitt, M.;
Pousset,
[31332] F.; Thomas, D.; Brice, A.; Koenig, M.; Komajda, M.: Corre-
lation between
[31333] left ventricular hypertrophy and GAA trinucleotide repeat
length in
[31334] Friedreich's ataxia. Circulation 95: 2247-2249, 1997.

[31335] 4384.

[31336] Koutnikova, H.; Campuzano, V.; Foury, F.; Dolle, P.; Caz-
zalini,
[31337] O.; Koenig, M.: Studies of human, mouse and yeast homo-
logues indicate
[31338] a mitochondrial function for frataxin. Nature Genet. 16:
345-351,
[31339] 1997.

[31340] 4385.

[31341] Litt, M.; Luty, J. A.: A hypervariable microsatellite revealed

[31342] by in vitro amplification of a dinucleotide repeat within the cardiac

[31343] muscle actin gene. Am. J. Hum. Genet. 44: 397-401, 1989.

[31344] 4386.

[31345] Baker, E.; Chen, L. Z.; Smith, C. A.; Callen, D. F.; Goodwin, R.;

[31346] Sutherland, G. R.: Chromosomal location of the human tumor necrosis

[31347] factor receptor genes. Cytogenet. Cell Genet. 57: 117-118, 1991.

[31348] 4387.

[31349] Brockhaus, M.; Schoenfeld, H.-J.; Schlaeger, E.-J.; Hunziker, W.;

[31350] Lesslauer, W.; Loetscher, H.: Identification of two types of tumor

[31351] necrosis factor receptors on human cell lines by monoclonal antibodies. Proc.

[31352] Nat. Acad. Sci. 87: 3127-3131, 1990.

[31353] 4388.

[31354] Castellino, A. M.; Parker, G. J.; Boronenkov, I. V.; Ander-
son,
[31355] R. A.; Chao, M. V.: A novel interaction between the jux-
tamembrane
[31356] region of the p55 tumor necrosis factor receptor and
phosphatidylinositol-4-phosphate
[31357] 5-kinase. J. Biol. Chem. 272: 5861-5870, 1997.

[31358] 4389.

[31359] Chan, F. K.-M.; Chun, H. J.; Zheng, L.; Siegel, R. M.; Bui, K.

[31360] L.; Lenardo, M. J.: A domain in TNF receptors that medi-
ates ligand-independent
[31361] receptor assembly and signaling. Science 288:
2351-2354, 2000.

[31362] 4390.

[31363] Aksentijevich, I.; Galon, J.; Soares, M.; Mansfield, E.; Hull,

[31364] K.; Oh, H.-H.; Goldbach-Mansky, R.; Dean, J.; Athreya, B.;
Reginato,
[31365] A. J.; Henrickson, M.; Pons-Estel, B.; O'Shea, J. J.; Kastner,
D.
[31366] L.: The tumor-necrosis-factor receptor-associated peri-

odic syndrome:

[31367] new mutations in TNFRSF1A, ancestral origins, genotype-phenotype studies,

[31368] and evidence for further genetic heterogeneity of periodic fevers. Am.

[31369] J. Hum. Genet. 69: 301-314, 2001. Note: Erratum: Am. J. Hum. Genet.

[31370] 69: 1160 only, 2001.

[31371] 4391.

[31372] Derre, J.; Kemper, O.; Cherif, D.; Nophar, Y.; Berger, R.; Wallach,

[31373] D.: The gene for the type 1 tumor necrosis factor receptor (TNF-R1)

[31374] is localized on band 12p13. Hum. Genet. 87: 231-233, 1991.

[31375] 4392.

[31376] Dode, C.; Papo, T.; Fieschi, C. Pecheux, C.; Dion, E.; Pi-card,

[31377] F.; Godeau, P.; Bienvenu, J.; Piette, J.-C.; Delpech, M.; Grateau,

[31378] G.: A novel missense mutation (C30S) in the gene encod-ing tumour

[31379] necrosis factor receptor 1 linked to autosomal-dominant recurrent
[31380] fever with localized myositis in a French family. Arthritis Rheum. 43:
[31381] 1535-1542, 2000.

[31382] 4393.

[31383] Engelmann, H.; Novick, D.; Wallach, D.: Two tumor necrosis factor-binding
[31384] proteins purified from human urine: evidence for immunological cross-reactivity
[31385] with cell surface tumor necrosis factor receptors. J. Biol. Chem. 265:
[31386] 1531-1536, 1990.

[31387] 4394.

[31388] Fuchs, P.; Strehl, S.; Dworzak, M.; Himmler, A.; Ambros, P. F.
[31389] : Structure of the human TNF receptor 1 (p60) gene (TNFR1) and localization
[31390] to chromosome 12p13. Genomics 13: 219-224, 1992.

[31391] 4395.

[31392] Atuk, N. O.; McDonald, T.; Wood, T.; Carpenter, J. T.;

Walzak,
[31393] M. P.; Donaldson, M.; Gillenwater, J. Y.; Turner, S. M.; Westfall,
[31394] V.: Familial pheochromocytoma, hypercalcemia, and von Hippel-Lindau
[31395] disease: a ten-year study of a large family. Medicine 58: 209-218,
[31396] 1979.

[31397] 4396.

[31398] Bender, B. U.; Altehofer, C.; Januszewicz, A.; Gartner, R.; Schmidt,
[31399] H.; Hoffmann, M. M.; Heidemann, P. H.; Neumann, H. P. H.: Functioning
[31400] thoracic paraganglioma: association with Von Hippel-Lindau syndrome. J.
[31401] Clin. Endocr. Metab. 82: 3356-3360, 1997.

[31402] 4397.

[31403] Bender, B. U.; Eng, C.; Olschewski, M.; Berger, D. P.; Laubenberger,
[31404] J.; Altehofer, C.; Kirste, G.; Orszagh, M.; van Velthoven, V.; Miosczka,
[31405] H.; Schmidt, D.; Neumann, H. P. H.: VHL c.505 T-C muta-

tion confers
[31406] a high age related penetrance but no increased overall mortality. J.
[31407] Med. Genet. 38: 508-514, 2001.

[31408] 4398.

[31409] Bender, B. U.; Gutsche, M.; Glasker, S.; Muller, B.; Kirste, G.;
[31410] Eng, C.; Neumann, H. P. H.: Differential genetic alterations in von
[31411] Hippel-Lindau syndrome-associated and sporadic pheochromocytomas. J.
[31412] Clin. Endocr. Metab. 85: 4568-4574, 2000.

[31413] 4399.

[31414] Bonnet, P.; Dechaume, J.; Blanc, E.: L'anevrisme cirsoide de la
[31415] retine l'anevrisme racemeux, ses relations avec l'anevrisme cirsoide
[31416] de la face et l'anevrisme cirsoid du cerveau. Bull. Soc. Ophtal.
[31417] Franc. 51: 521-524, 1938.

[31418] 4400.

[31419] Bradley, J. F.; Collins, D. L.; Schimke, R. N.; Parrott, H. N.;

[31420] Rothberg, P. G.: Two distinct phenotypes caused by two different

[31421] missense mutations in the same codon of the VHL gene. Am. J. Med.

[31422] Genet. 87: 163-167, 1999.

[31423] 4401.

[31424] Brauch, H.; Kishida, T.; Glavac, D.; Chen, F.; Pausch, F.; Hofler,

[31425] H.; Latif, F.; Lerman, M. I.; Zbar, B.; Neumann, H. P. H.: Von Hippel-Lindau

[31426] (VHL) disease with pheochromocytoma in the Black Forest region of

[31427] Germany: evidence for a founder effect. Hum. Genet. 95: 551-556,

[31428] 1995.

[31429] 4402.

[31430] Brown, D. G.; Hilal, S. K.; Tenner, M. S.: Wyburn-Mason syndrome. Arch.

[31431] Neurol. 28: 67-68, 1973.

[31432] 4403.

[31433] Cahill, G. F.; Melicow, M. M.; Guerry, D.: The renal lesions

[31434] in von Hippel-Lindau's disease. Trans. Am. Assoc. Genitourinary Surg. 35:

[31435] 271-281, 1942.

[31436] 4404.

[31437] Cameron, S. J.; Doig, A.: Cerebellar tumors presented with clinical

[31438] features of phaeochromocytoma. Lancet I: 492-494, 1970.

[31439] 4405.

[31440] Chapman, R. C.; Diaz-Perez, R.: Pheochromocytoma associated with

[31441] cerebellar hemangioblastoma. J.A.M.A. 182: 1014-1017, 1962.

[31442] 4406.

[31443] Chen, F.; Slife, L.; Kishida, T.; Mulvihill, J.; Tisherman, S.

[31444] E.; Zbar, B.: Genotype-phenotype correlation in von Hippel-Lindau

[31445] disease: identification of a mutation associated with VHL type 2A. J.

[31446] Med. Genet. 33: 716-717, 1996.

[31447] 4407.

[31448] Christoferson, L. A.; Gustafson, M. B.; Petersen, A. G.: Von

[31449] Hippel-Lindau's disease. J.A.M.A. 178: 280-282, 1961.

[31450] 4408.

[31451] Chen, F.; Kishida, T.; Yao, M.; Hustad, T.; Glavac, D.; Dean,

[31452] M.; Gnarra, J. R.; Orcutt, M. L.; Duh, F. M.; Glenn, G.; Green, J.;

[31453] Hsia, Y. E.; Lamiell, J.; Li, H.; Wei, M. H.; Schmidt, L.; Tory, K.;

[31454] Kuzman, I.; Stackhouse, T.; Latif, F.; Linehan, W. M.; Ler-

man, M.;

[31455] Zbar, B.: Germline mutations in the von Hippel-Lindau disease tumor

[31456] suppressor gene: correlations with phenotype. Hum. Mu-

tat. 5: 66-75,

[31457] 1995.

[31458] 4409.

[31459] Collins, E. T.: Intra-ocular growths (two cases, brother and

[31460] sister, with peculiar vascular new growth, probably pri-

marily retinal,

[31461] affecting both eyes). Trans. Ophthal. Soc. U.K. 14: 141-149, 1894.

[31462] 4410.

[31463] Crossey, P. A.; Foster, K.; Richards, F. M.; Phipps, M. E.; Latif,

[31464] F.; Tory, K.; Jones, M. H.; Bentley, E.; Kumar, R.; Lerman, M. I.;

[31465] Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.; Maher, E. R.: Molecular

[31466] genetic investigations of the mechanism of tumourigenesis in von Hippel-Lindau

[31467] disease: analysis of allele loss in VHL tumours. Hum. Genet. 93:

[31468] 53-58, 1994.

[31469] 4411.

[31470] Crossey, P. A.; Richards, F. M.; Foster, K.; Green, J. S.; Prowse,

[31471] A.; Latif, F.; Lerman, M. I.; Zbar, B.; Affara, N. A.; Ferguson-Smith,

[31472] M. A.; Maher, E. R.: Identification of intragenic mutations in the

[31473] von Hippel-Lindau disease tumour suppressor gene and

correlation with
[31474] disease phenotype. Hum. Molec. Genet. 3: 1303-1308, 1994.
[31475] 4412.
[31476] Cushing, H.; Bailey, P.: Hemangiomas of cerebellum and retina
[31477] (Lindau's disease), with the report of a case. Arch. Ophthal. 57:
[31478] 447-463, 1928.
[31479] 4413.
[31480] Davies, D. R.; Norman, A. M.; Whitehouse, R. W.; Evans, D. G.
[31481] R.: Non-expression of von Hippel-Lindau phenotype in an obligate
[31482] gene carrier. Clin. Genet. 45: 104-106, 1994.
[31483] 4414.
[31484] Decker, H.-J. H.; Neumann, H. P. H.; Walter, T. A.; Sandberg,
[31485] A. A.: 3p involvement in a renal cell carcinoma in von Hippel-Lindau
[31486] syndrome: region of tumor breakpoint clustering on 3p?

Cancer Genet.

[31487] Cytogenet. 33: 59-65, 1988.

[31488] 4415.

[31489] Duan, D. R.; Humphrey, J. S.; Chen, D. Y. T.; Weng, Y.; Sukegawa,

[31490] J.; Lee, S.; Gnarra, J. R.; Linehan, W. M.; Klausner, R. D.: Characterization

[31491] of the VHL tumor suppressor gene product: localization, complex formation,

[31492] and the effect of natural inactivating mutations. Proc. Nat. Acad.

[31493] Sci. 92: 6459-6463, 1995.

[31494] 4416.

[31495] Duan, D. R.; Pause, A.; Burgess, W. H.; Aso, T.; Chen, D. Y. T.;

[31496] Garrett, K. P.; Conaway, R. C.; Conaway, J. W.; Linehan, W. M.; Klausner,

[31497] R. D.: Inhibition of transcription elongation by the VHL tu-mor suppressor

[31498] protein. Science 269: 1402-1406, 1995.

[31499] 4417.

[31500] Hurvitz, J. R.; Suwairi, W. M.; Van Hul, W.; El-Shanti, H.; Superti-Furga,
[31501] A.; Roudier, J.; Holderbaum, D.; Pauli, R. M.; Herd, J. K.; Van Hul,
[31502] E.; Rezai-Delui, H.; Legius, E.; Le Merrer, M.; Al-Alami, J.; Bahabri,
[31503] S. A.; Warman, M. L.: Mutations in the CCN gene family member WISP3
[31504] cause progressive pseudorheumatoid dysplasia. Nature Genet. 23:
[31505] 94-98, 1999.

[31506] 4418.

[31507] Spranger, J.; Albert, C.; Schilling, F.; Bartsocas, C.; Stoss,

[31508] H.: Progressive pseudorheumatoid arthritis of childhood (PPAC): a
[31509] hereditary disorder simulating rheumatoid arthritis. Europ. J. Pediat. 140:
[31510] 34-40, 1983.

[31511] 4419.

[31512] Bahabri, S. A.; Suwairi, W. M.; Laxer, R. M.; Polinkovsky, A.;

[31513] Dalaan, A. A.; Warman, M. L.: The camptodactyly-arthropa-

thy-coxa
[31514] vara-pericarditis syndrome: clinical features and genetic mapping
[31515] to human chromosome 1. Arthritis Rheum. 41: 730-735, 1998.
[31516] 4420.

[31517] Marcelino, J.; Carpten, J. D.; Suwairi, W. M.; Gutierrez, O. M.;
[31518] Schwartz, S.; Robbins, C.; Sood, R.; Makalowska, I.; Baxevanis, A.;
[31519] Johnstone, B.; Laxer, R. M.; Zemel, L.; and 13 others: CACP, encoding
[31520] a secreted proteoglycan, is mutated in camptodactyly-arthropathy-coxa
[31521] vara-pericarditis syndrome. Nature Genet. 23: 319-322, 1999.
[31522] 4421.

[31523] Drolet, D. W.; Scully, K. M.; Simmons, D. M.; Wegner, M.; Chu,
[31524] K.; Swanson, L. W.; Rosenfeld, M. G.: TEF, a transcription factor
[31525] expressed specifically in the anterior pituitary during em-

bryogenesis,

[31526] defines a new class of leucine zipper proteins. Genes Dev. 5: 1739-1753,

[31527] 1991.

[31528] 4422.

[31529] Andrechek, E. R.; Hardy, W. R.; Girgis-Gabardo, A. A.; Perry, R.

[31530] L. S.; Butler, R.; Graham, F. L.; Kahn, R. C.; Rudnicki, M. A.; Muller,

[31531] W. J.: ErbB2 is required for muscle spindle and myoblast cell survival. Molec.

[31532] Cell. Biol. 22: 4714-4722, 2002.

[31533] 4423.

[31534] Bargmann, C. I.; Hung, M.-C.; Weinberg, R. A.: The NEU oncogene

[31535] encodes an epidermal growth factor receptor-related protein. Nature 319:

[31536] 226-230, 1986.

[31537] 4424.

[31538] Chan, R.; Hardy, W. R.; Laing, M. A.; Hardy, S. E.; Muller, W.

[31539] J.: The catalytic activity of the ErbB-2 receptor tyrosine ki-
nase
[31540] is essential for embryonic development. Molec. Cell. Biol.
22: 1073-1078,
[31541] 2002.

[31542] 4425.

[31543] Coussens, L.; Yang-Feng, T. L.; Liao, Y.-C.; Chen, E.; Gray,
A.;
[31544] McGrath, J.; Seeburg, P. H.; Libermann, T. A.; Schlessinger,
J.; Francke,
[31545] U.; Levinson, A.; Ullrich, A.: Tyrosine kinase receptor with
extensive
[31546] homology to EGF receptor shares chromosomal location
with NEU oncogene. Science 230:
[31547] 1132-1139, 1985.

[31548] 4426.

[31549] Crone, S. A.; Zhao, Y.-Y.; Fan, L.; Gu, Y.; Minamisawa, S.;
Liu,
[31550] Y.; Peterson, K. L.; Chen, J.; Kahn, R.; Condorelli, G.; Ross,
J.,
[31551] Jr.; Chien, K. R.; Lee, K.-F.: ErbB2 is essential in the pre-
vention

[31552] of dilated cardiomyopathy. Nature Med. 8: 459-465, 2002.

[31553] 4427.

[31554] Dankort, D.; Maslikowski, B.; Warner, N.; Kanno, N.; Kim, H.; Wang,

[31555] Z.; Moran, M. F.; Oshima, R. G.; Cardiff, R. D.; Muller, W. J.: Grb2

[31556] and Shc adapter proteins play distinct roles in Neu (ErbB-2) induced

[31557] mammary tumorigenesis: implications for human breast cancer. Molec.

[31558] Cell. Biol. 21: 1540-1551, 2001.

[31559] 4428.

[31560] De Boer, J. G.: A new mutator phenotype in breast cancer? (Commentary) Proc.

[31561] Nat. Acad. Sci. 99: 3368-3369, 2002.

[31562] 4429.

[31563] De Placido, S.; Carlomagno, C.; De Laurentiis, M.; Bianco, A.

[31564] R.: C-erbB2 expression predicts tamoxifen efficacy in breast cancer

[31565] patients. Breast Cancer Res. Treat. 52: 55-64, 1998.

[31566] 4430.

[31567] Di Fiore, P. P.; Pierce, J. H.; Kraus, M. H.; Segatto, O.; King,

[31568] C. R.; Aaronson, S. A.: erbB-2 is a potent oncogene when overexpressed

[31569] in NIH/3T3 cells. Science 237: 178-182, 1987.

[31570] 4431.

[31571] Doherty, J. K.; Bond, C.; Jardim, A.; Adelman, J. P.; Clinton,

[31572] G. M.: The HER-2/neu receptor tyrosine kinase gene encodes a secreted

[31573] autoinhibitor. Proc. Nat. Acad. Sci. 96: 10869-10874, 1999.

[31574] 4432.

[31575] Ehsani, A.; Low, J.; Wallace, R. B.; Wu, A. M.: Characterization

[31576] of a new allele of the human ERBB2 gene by allele-specific competition

[31577] hybridization. Genomics 15: 426-429, 1993.

[31578] 4433.

[31579] Francke, U.: Personal Communication. New Haven, Con-

necticut

[31580] 4/1988.

[31581] 4434.

[31582] Fukushige, S.-I.; Matsubara, K.-I.; Yoshida, M.; Sasaki, M.; Suzuki,

[31583] T.; Semba, K.; Toyoshima, K.; Yamamoto, T.: Localization of a novel

[31584] v-erbB-related gene, c-erbB-2, on human chromosome 17 and its amplification

[31585] in a gastric cancer cell line. Molec. Cell. Biol. 6: 955-958, 1986.

[31586] 4435.

[31587] Kaneko, Y.; Homma, C.; Maseki, N.; Sakurai, M.; Toyoshima, K.;

[31588] Yamamoto, T.: Human c-erbB-2 remains on chromosome 17 in band q21

[31589] in the 15;17 translocation associated with acute promye-locytic leukemia. Jpn.

[31590] J. Cancer Res. 78: 16-19, 1987.

[31591] 4436.

[31592] Lin, W.; Sanchez, H. B.; Deerinck, T.; Morris, J. K.; Ellisman,

[31593] M.; Lee, K. F.: Aberrant development of motor axons and neuromuscular

[31594] synapses in erbB2-deficient mice. Proc. Nat. Acad. Sci. 97: 1299-1304,

[31595] 2000.

[31596] 4437.

[31597] Liu, S.; Liu, W.; Jakubczak, J. L.; Erexson, G. L.; Tindall, K.

[31598] R.; Chan, R.; Muller, W. J.; Adhya, S.; Garges, S.; Merlino, G.:

[31599] Genetic instability favoring transversions associated with ErbB2-induced

[31600] mammary tumorigenesis. Proc. Nat. Acad. Sci. 99: 3770-3775, 2002.

[31601] 4438.

[31602] Mehta, R. R.; McDermott, J. H.; Hieken, T. J.; Marler, K. C.;

[31603] Patel, M. K.; Wild, L. D.; Das Gupta, T. K.: Plasma c-erbB2 levels

[31604] in breast cancer patients: prognostic significance in predicting response

[31605] to chemotherapy. J. Clin. Oncol. 16: 2409-2416, 1998.

[31606] 4439.

[31607] Morris, J. K.; Lin, W.; Hauser, C.; Marchuk, Y.; Getman, D.; Lee,

[31608] K.-F.: Rescue of the cardiac defect in ErbB2 mutant mice reveals

[31609] essential roles of ErbB2 in peripheral nervous system development. Neuron 23:

[31610] 273-283, 1999.

[31611] 4440.

[31612] Muleris, M.; Almeida, A.; Malfoy, B.; Dutrillaux, B.: Assignment

[31613] of v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2

[31614] (ERBB2) to human chromosome band 17q21.1 by in situ hybridization. Cytogenet.

[31615] Cell Genet. 76: 34-35, 1997.

[31616] 4441.

[31617] Ozcelik, C.; Erdmann, B.; Pilz, B.; Wettschureck, N.; Britsch,

[31618] S.; Hubner, N.; Chien, K. R.; Birchmeier, C.; Garratt, A. N.:

Conditional
[31619] mutation of the ErbB2 (HER2) receptor in cardiomyocytes leads to dilated
[31620] cardiomyopathy. Proc. Nat. Acad. Sci. 99: 8880-8885, 2002.
[31621] 4442.

[31622] Papewalis, J.; Nikitin, A. Y.; Rajewsky, M. F.: G to A poly-morphism
[31623] at amino acid codon 655 of the human erbB-2/HER2 gene. Nucleic Acids
[31624] Res. 19: 5452 only, 1991.

[31625] 4443.

[31626] Pegram, M. D.; Finn, R. S.; Arzoo, K.; Beryt, M.; Pietras, R.

[31627] J.; Slamon, D. J.: The effect of HER-2/neu overexpression on chemotherapeutic
[31628] drug sensitivity in human breast and ovarian cancer cells. Oncogene 15:
[31629] 537-547, 1997.

[31630] 4444.

[31631] Pietras, R. J.; Pegram, M. D.; Finn, R. S.; Maneval, D. A.; Slamon,

[31632] D. J.: Remission of human breast cancer xenografts on therapy with
[31633] humanized monoclonal antibody to HER-2 receptor and DNA-reactive drugs. Oncogene 17:
[31634] 2235-2249, 1998.

[31635] 4445.

[31636] Popescu, N. C.; King, C. R.; Kraus, M. H.: Localization of the
[31637] human erbB-2 gene on normal and rearranged chromosomes 17 to bands
[31638] q12-21.32. Genomics 4: 362-366, 1989.

[31639] 4446.

[31640] Qiu, Y.; Ravi, L.; Kung, H.-J.: Requirement of ErbB2 for signalling
[31641] by interleukin-6 in prostate carcinoma cells. Nature 393: 83-85,
[31642] 1998.

[31643] 4447.

[31644] Semba, K.; Kamata, N.; Toyoshima, K.; Yamamoto, T.: A v-erbB-related
[31645] protooncogene, c-erbB-2, is distinct from the c-

erbB-1/epidermal growth
[31646] factor-receptor gene and is amplified in a human salivary gland adenocarcinoma. Proc.
[31647] Nat. Acad. Sci. 82: 6497-6501, 1985.

[31648] 4448.

[31649] Slamon, D. J.; Godolphin, W.; Jones, L. A.; Holt, J. A.; Wong,
[31650] S. G.; Keith, D. E.; Levin, W. J.; Stuart, S. G.; Udove, J.; Ullrich,
[31651] A.; Press, M. F.: Studies of the HER-2/neu proto-oncogene in human
[31652] breast and ovarian cancer. Science 244: 707-712, 1989.

[31653] 4449.

[31654] Slamon, D. J.; Leyland-Jones, B.; Shak, S.; Fuchs, H.; Paton,
[31655] V.; Bajamonde, A.; Fleming, T.; Eiermann, W.; Wolter, J.; Pegram,
[31656] M.; Baselga, J.; Norton, L.: Use of chemotherapy plus a monoclonal
[31657] antibody against HER2 for metastatic breast cancer that overexpresses
[31658] HER2. New Eng. J. Med. 344: 783-792, 2001.

[31659] 4450.

[31660] Beutler, E.; Kuhl, W.; Gelbart, T.:
6-Phosphogluconolactonase
[31661] deficiency, a hereditary erythrocyte enzyme deficiency:
possible interaction
[31662] with glucose-6-phosphate dehydrogenase deficiency.
Proc. Nat. Acad.
[31663] Sci. 82: 3876-3878, 1985.

[31664] 4451.

[31665] Benkmann, H.-G.; Paik, Y. K.; Chen, L. Z.; Goedde, H. W.:
Polymorphism
[31666] of 6-PGD in South Korea: a new genetic variant 6-PGD
Korea. Hum.
[31667] Genet. 74: 204-205, 1986.

[31668] 4452.

[31669] Blake, N. M.; Kirk, R. L.: New genetic variant of
6-phosphogluconate
[31670] dehydrogenase in Australian aborigines. Nature 221: 278
only, 1969.
[31671] 4453.

[31672] Bowman, J. E.; Carson, P. E.; Frischer, H.; De Garay, A. L.:

Genetics

[31673] of starch-gel electrophoretic variants of human 6-phosphogluconic

[31674] dehydrogenase: population and family studies in the United States

[31675] and in Mexico. Nature 210: 811-812, 1966.

[31676] 4454.

[31677] Brewer, G. J.; Dern, R. J.: A new inherited enzymatic defi-ciency

[31678] of human erythrocytes: 6-phosphogluconate dehydroge-nase deficiency. Am.

[31679] J. Hum. Genet. 16: 472-476, 1964.

[31680] 4455.

[31681] Burgerhout, W.; Van Someren, H.; Bootsma, D.: Cytological mapping

[31682] of the genes assigned to the human A1 chromosome by use of radiation-induced

[31683] chromosome breakage in a human-Chinese hamster hy-brid cell line. Humangenetik 20:

[31684] 159-162, 1973.

[31685] 4456.

[31686] Davidson, R. G.: Electrophoretic variants of human 6-phosphogluconate
[31687] dehydrogenase: population and family studies and description of a
[31688] new variant. Ann. Hum. Genet. 30: 355-362, 1967.

[31689] 4457.

[31690] Dern, R. J.; Brewer, G. J.; Tashian, R. E.; Shows, T. B.: Hereditary
[31691] variation of erythrocytic 6-phosphogluconate dehydrogenase. J. Lab.
[31692] Clin. Med. 67: 255-264, 1966.

[31693] 4458.

[31694] Douglas, G. R.; McAlpine, P. J.; Hamerton, J. L.: Regional localization
[31695] of loci for human PGM-1 and 6PGD on human chromosome 1 by use of hybrids
[31696] of Chinese hamster-human somatic cells. Proc. Nat. Acad. Sci. 70:
[31697] 2737-2740, 1973.

[31698] 4459.

[31699] Fildes, R. A.; Parr, C. W.: Human red-cell phosphoglu-

conate dehydrogenases. Nature 200:
[31700] 890-891, 1963.

[31701] 4460.

[31702] Nelson, M. S.: Biochemical and genetic characterization of the
[31703] Lowell variant: a new phenotype of 6-phosphogluconate dehydrogenase. Hum.
[31704] Genet. 62: 333-336, 1982.

[31705] 4461.

[31706] Nevo, S.: A new rare PGD variant, PGD Mediterranean. Hum. Genet. 81:
[31707] 199 only, 1989.

[31708] 4462.

[31709] Parr, C. W.: Erythrocyte phosphogluconate dehydrogenase polymorphism. Nature 210:
[31710] 487-489, 1966.

[31711] 4463.

[31712] Parr, C. W.; Fitch, L. I.: Inherited quantitative variations
[31713] of human phosphogluconate dehydrogenase. Ann. Hum. Genet. 30: 339-353,

[31714] 1967.

[31715] 4464.

[31716] Ritter, H.; Toriverdiau, G.; Wendt, G. G.; Zilch, I.: Genetic

[31717] and linkage analysis on 6-PGD. Humangenetik 14: 73-75, 1971.

[31718] 4465.

[31719] Tariverdian, G.; Ropers, H.-H.; Op't Hof, J.; Ritter, H.: Zur

[31720] Genetik der 6-Phosphogluconatdehydrogenase (EC: 1.1.1.44): Eine neue

[31721] Variante F (Freiburg). Humangenetik 10: 355-357, 1970.

[31722] 4466.

[31723] Weitkamp, L. R.: Genetic linkage relationships of the ADA and

[31724] 6-PGD loci in 'Humangenetik.' (Letter) Humangenetik 15: 359-360,

[31725] 1972.

[31726] 4467.

[31727] Weitkamp, L. R.; Guttormsen, S. A.; Greendyke, R. M.: Genetic

[31728] linkage between a locus for 6-PGD and the Rh locus: eval-

uation of

possible heterogeneity in the recombination fraction between sexes

and among families. Am. J. Hum. Genet. 23: 462-470, 1971.

4468.

Weitkamp, L. R.; Guttormsen, S. A.; Shreffler, D. C.; Sing, C.

F.; Napier, J. A.: Genetic linkage relations of the loci for 6-phosphogluconate

dehydrogenase and adenosine deaminase in man. Am. J. Hum. Genet. 22:

216-220, 1970.

4469.

Westerveld, A.; Meera Khan, P.: Evidence for linkage between

human loci for 6-phosphogluconate dehydrogenase and phosphoglucomutase(1)

in man-Chinese hamster somatic cell hybrids. Nature 236: 30-32,

1972.

4470.

[31742] Charles, D. J.; Pretsch, W.: Linear dose-response relation-
ship
[31743] of erythrocyte enzyme-activity mutations in offspring of
ehylnitrosourea-treated
[31744] mice. Mutat. Res. 176: 81-91, 1987.

[31745] 4471.

[31746] Chen, S.-H.; Anderson, J.; Giblett, E. R.; Lewis, M.: Phos-
phoglyceric
[31747] acid mutase: rare genetic variants and tissue distribution.
Am. J.
[31748] Hum. Genet. 26: 73-77, 1974.

[31749] 4472.

[31750] DiMauro, S.; Miranda, A. F.; Sakoda, S.; Schon, E. A.;
Servidei,
[31751] S.; Shanske, S.; Zeviani, M.: Metabolic myopathies. Am. J.
Med.
[31752] Genet. 25: 635-651, 1986.

[31753] 4473.

[31754] Fundele, R.; Bucher, T.; Gropp, A.; Winking, H.: Enzyme
patterns
[31755] in trisomy 19 of the mouse. Dev. Genet. 2: 291-303,

1981.

[31756] 4474.

[31757] Gerald, P. S.; Grzeschik, K. H.: Report of the committee on the

[31758] genetic constitution of chromosomes 10, 11 and 12. Cytogenet. Cell

[31759] Genet. 37: 103-126, 1984.

[31760] 4475.

[31761] Pretsch, W.; Favor, J.: A precise localization of a mouse gene

[31762] encoding increased phosphoglycerate mutase activity (Pgam1e1) on chromosome

[31763] 19. Mammalian Genome 7: 619 only, 1996.

[31764] 4476.

[31765] Sakoda, S.; Shanske, S.; DiMauro, S.; Schon, E. A.: Isolation

[31766] of a cDNA encoding the B isozyme of human phosphoglycerate mutase

[31767] (PGAM) and characterization of the PGAM gene family. J. Biol. Chem. 263:

[31768] 16899-16905, 1988.

[31769] 4477.

[31770] Blanche, H.; Szabo, P.; Cohen, D.; Dausset, J.; Cann, H. M.: A

[31771] subclone of the autosomal phosphoglycerate kinase pseudogene (HGM8

[31772] gene symbol PGK1P2), localized to 6p23-q12, detects moderately polymorphic

[31773] RFLP. Nucleic Acids Res. 15: 3941 only, 1987.

[31774] 4478.

[31775] Thomas, K. R.; Capecchi, M. R. :Cell 51: 503-512, 1987.

[31776] 4479.

[31777] Berdichevskii, F. B.; Chumakov, I. M.; Kiselev, L. L.: Determination

[31778] of the nucleotide sequence of the son3 fragment of the human genome:

[31779] identification of a new protein with an unusual structure and homology

[31780] with DNA-binding proteins. Molec. Biol. (Mosk.) 22: 794-801, 1988.

[31781] 4480.

[31782] Khan, I. M.; Fisher, R. A.; Johnson, K. J.; Bailey, M. E. S.; Siciliano,

[31783] M. J.; Kessling, A. M.; Farrer, M.; Carritt, B.; Kamalati, T.; Buluwela,
[31784] L.: The SON gene encodes a conserved DNA binding protein mapping
[31785] to human chromosome 21. Ann. Hum. Genet. 58: 25-34, 1994.
[31786] 4481.

[31787] Carr, I. M.: Personal Communication. Leeds, England 3/1/1999.
[31788] 4482.

[31789] Carr, I. M.; Markham, A. F.: Molecular genetic analysis of the
[31790] human sorbitol dehydrogenase gene. Mammalian Genome 6: 645-652,
[31791] 1995.

[31792] 4483.

[31793] Carr, I. M.; Markham, A. F.; Coletta, P. L.: Identification and
[31794] characterisation of a sequence related to human sorbitol dehydrogenase. Europ.
[31795] J. Biochem. 245: 760-767, 1997.

[31796] 4484.

[31797] Carr, I. M.; Whitehouse, A.; Coletta, P. L.; Markham, A. F.: Structural
[31798] and evolutionary characterization of the human sorbitol dehydrogenase
[31799] gene duplication. Mammalian Genome 9: 1042-1048, 1998.

[31800] 4485.

[31801] Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of the
[31802] sorbitol dehydrogenase locus to human chromosome 15pter-q21. Biochem.
[31803] Genet. 18: 425-431, 1980.

[31804] 4486.

[31805] Iwata, T.; Popescu, N. C.; Zimonjic, D. B.; Karlsson, C.; Hoog,
[31806] J.-O.; Vaca, G.; Rodriguez, I. R.; Carper, D.: Structural or-
ganization
[31807] of the human sorbitol dehydrogenase gene (SORD). Ge-
nomics 26: 55-62,
[31808] 1995.

[31809] 4487.

[31810] Lee, F. K.; Cheung, M. C.; Chung, S.: The human sorbitol dehydrogenase

[31811] gene: cDNA cloning, sequence determination, and mapping by fluorescence

[31812] in situ hybridization. Genomics 21: 354-358, 1994.

[31813] 4488.

[31814] Op't Hof, J.: Isoenzymes and population genetics of sorbitol dehydrogenase

[31815] (EC: 1.1.1.14) in swine (Sus scrofa). Humangenetik 7: 258-259, 1969.

[31816] 4489.

[31817] Shin, Y. S.; Rieth, M.; Endres, W.: Sorbitol dehydrogenase deficiency

[31818] in a family with congenital cataracts. J. Inherit. Metab. Dis. 7

[31819] (suppl. 2): 151-152, 1984.

[31820] 4490.

[31821] Vaca, G.; Ibarra, B.; Bracamontes, M.; Garcia-Cruz, D.; Sanchez-Corona,

[31822] J.; Medina, C.; Wunsch, C.; Gonzalez-Quiroga, G.; Cantu, J.

M.: Red
[31823] blood cell sorbitol dehydrogenase deficiency in a family with cataracts. Hum.
[31824] Genet. 61: 338-341, 1982.

[31825] 4491.

[31826] Bowtell, D.; Fu, P.; Simon, M. A.; Senior, P.: Identification

[31827] of murine homologues of the Drosophila Son of sevenless gene: potential
[31828] activators of ras. Proc. Nat. Acad. Sci. 89: 6511-6515, 1992.
[31829] 4492.

[31830] Chardin, P.; Camonis, J. H.; Gale, N. W.; Van Aelst, L.; Schlessinger,
[31831] J.; Wigler, M. H.; Bar-Sagi, D.: Human Sos1: a guanine nucleotide
[31832] exchange factor for Ras that binds to GRB2. Science 260: 1338-1343,
[31833] 1993.

[31834] 4493.

[31835] Scita, G.; Nordstrom, J.; Carbone, R.; Tenca, P.; Giardina, G.;

[31836] Gutkind, S.; Bjarnegard, M.; Betsholtz, C.; Di Fiore, P. P.: EPS8
[31837] and E3B1 transduce signals from Ras to Rac. Nature 401: 290-293,
[31838] 1999.

[31839] 4494.

[31840] Soisson, S. M.; Nimnual, A. S.; Uy, M.; Bar-Sagi, D.; Kuriyan,
[31841] J.: Crystal structure of the Dbl and pleckstrin homology domains
[31842] from the human Son of sevenless protein. Cell 95: 259-268, 1998.
[31843] 4495.

[31844] Webb, G. C.; Jenkins, N. A.; Largaespada, D. A.; Copeland, N. G.;
[31845] Fernandez, C. S.; Bowtell, D. D. L.: Mammalian homo-logues of the
[31846] Drosophila Son of sevenless gene map to murine chromo-somes 17 and
[31847] 12 and to human chromosomes 2 and 14, respectively. Genomics 18:
[31848] 14-19, 1993.

[31849] 4496.

[31850] Sarraf, P.; Mueller, E.; Smith, W. M.; Wright, H. M.; Kum, J.

[31851] B.; Aaltonen, L. A.; de la Chapelle, A.; Spiegelman, B. M.;
Eng, C.

[31852] : Loss-of-function mutations in PPAR-gamma associated
with human colon

[31853] cancer. Molec. Cell 3: 799-804, 1999.

[31854] 4497.

[31855] Scheidler, S.; Fredericks, W. J.; Rauscher, F. J., III; Barr,

[31856] F. G.; Vogt, P. K.: The hybrid PAX3-FKHR fusion protein of
alveolar

[31857] rhabdomyosarcoma transforms fibroblasts in culture.
Proc. Nat. Acad.

[31858] Sci. 93: 9805-9809, 1996.

[31859] 4498.

[31860] Sublett, J. E.; Jeon, I-S.; Shapiro, D. N.: The alveolar rhab-
domyosarcoma

[31861] PAX3/FKHR fusion protein is a transcriptional activator.
Oncogene 11:

[31862] 545-552, 1995.

[31863] 4499.

[31864] Whang-Peng, J.; Knutsen, T.; Theil, K.; Horowitz, M. E.; Triche,

[31865] T.: Cytogenetic studies in subgroups of rhabdomyosar-coma. Genes

[31866] Chromosomes Cancer 5: 299-310, 1992.

[31867] 4500.

[31868] Richard, I.; Broux, O.; Chiannilkulchai, N.; Fougerousse, F.; Allamand,

[31869] V.; Bourg, N.; Brenguier, L.; Devaud, C.; Pasturaud, P.; Roudaut,

[31870] C.; Lorenzo, F.; Sebastiani-Kabatchis, C.; Schultz, R. A.; Polymeropoulos,

[31871] M. H.; Gyapay, G.; Auffray, C.; Beckmann, J. S.: Regional localization

[31872] of human chromosome 15 loci. Genomics 23: 619-627, 1994.

[31873] 4501.

[31874] Zuniga, A.; Haramis, A.-P. G.; McMahon, A. P.; Zeller, R.: Signal

[31875] relay by BMP antagonism controls the SHH/FGF4 feedback loop in vertebrate

[31876] limb buds. Nature 401: 598-602, 1999.

[31877] 4502.

[31878] Bao, L.; Gerard, N. P.; Eddy, R. L., Jr.; Shows, T. B.; Gerard,

[31879] C.: Mapping of genes for the human C5a receptor (C5AR), human FMLP

[31880] receptor (FPR), and two FMLP receptor homologue orphan receptors (FPRH1,

[31881] FPRH2) to chromosome 19. Genomics 13: 437-440, 1992.

[31882] 4503.

[31883] Ozcelik, T.; Murphy, P. M.; Francke, U.: Chromosomal as-signment

[31884] of genes for a formyl peptide receptor (FPR1), a structural homologue

[31885] of the formyl peptide receptor (FPRL1) and a low affinity interleukin

[31886] 8 receptor (IL8RA) in human. (Abstract) Cytogenet. Cell Genet. 58:

[31887] 2023-2024, 1991.

[31888] 4504.

[31889] Walther, A.; Riehemann, K.; Gerke, V.: A novel ligand of the formyl

[31890] peptide receptor: annexin I regulates neutrophil extravasation by
[31891] interacting with the FPR. Molec. Cell 5: 831-840, 2000.

[31892] 4505.

[31893] Klein, C.; Paul, J. I.; Sauve, K.; Schmidt, M. M.; Arcangeli, L.;
[31894] Ransom, J.; Trueheart, J.; Manfredi, J. P., Broach, J. R.; Murphy,
[31895] A. J.: Identification of surrogate agonists for the human FPRL-1
[31896] receptor by autocrine selection in yeast. Nature Biotech. 16: 1334-1337,
[31897] 1998.

[31898] 4506.

[31899] Le, Y.; Yazawa, H.; Gong, W.; Yu, Z.; Ferrans, V. J.; Murphy, P.
[31900] M.; Wang, J. M.: Cutting edge: the neurotoxic prion peptide fragment
[31901] PrP(106-126) is a chemotactic agonist for the G protein-coupled receptor
[31902] formyl peptide receptor-like 1. J. Immun. 166: 1448-1451, 2001.

[31903] 4507.

[31904] Murphy, P. M.; Ozcelik, T.; Kenney, R. T.; Tiffany, H. L.; McDermott,

[31905] D.; Francke, U.: A structural homologue of the N-formyl peptide receptor:

[31906] characterization and chromosome mapping of a peptide chemoattractant

[31907] receptor family. J. Biol. Chem. 267: 7637-7643, 1992.

[31908] 4508.

[31909] Margaritte, P.; Bonaiti-Pellie, C.; King, M.-C.; Clerget-Darpoux,

[31910] F.: Linkage of familial breast cancer to chromosome 17q21 may not

[31911] be restricted to early-onset disease. Am. J. Hum. Genet. 50: 1231-1234,

[31912] 1992.

[31913] 4509.

[31914] Mefford, H. C.; Baumbach, L.; Panguluri, R. C. K.; Whit-field-Broome,

[31915] C.; Szabo, C.; Smith, S.; King, M.-C.; Dunston, G.; Stoppa-Lyonnet,

[31916] D.; Arena, F.: Evidence for a BRCA1 founder mutation in families
[31917] of West African ancestry. (Letter) Am. J. Hum. Genet. 65: 575-578,
[31918] 1999.

[31919] 4510.

[31920] Meijers-Heijboer, E. J.; Verhoog, L. C.; Brekelmans, C. T. M.;
[31921] Seynaeve, C.; Tilanus-Linthorst, M. M. A.; Wagner, A.; Dukel, L.;
[31922] Devilee, P.; van den Ouweland, A. M. W.; van Geel, A. N.; Klijn, J.
[31923] G. M.: Presymptomatic DNA testing and prophylactic surgery in families
[31924] with a BRCA1 or BRCA2 mutation. Lancet 355: 2015-2020, 2000.
[31925] 4511.

[31926] Meijers-Heijboer, H.; van Geel, B.; van Putten, W. L. J.; Henzen-Logmans,
[31927] S. C.; Seynaeve, C.; Menke-Pluymers, M. B. E.; Bartels, C. C. M.;
[31928] Verhoog, L. C.; van den Ouweland, A. M. W.; Niermeijer,

M. F.; Brekelmans,
[31929] C. T. M.; Klijn, J. G. M.: Breast cancer after prophylactic bi-
lateral
[31930] mastectomy in women with a BRCA1 or BRCA2 mutation.
New Eng. J. Med. 345:
[31931] 159-164, 2001.

[31932] 4512.

[31933] Merajver, S. D.; Pham, T. M.; Caduff, R. F.; Chen, M.; Poy,
E.
[31934] L.; Cooney, K. A.; Weber, B. L.; Collins, F. S.; Johnston, C.;
Frank,
[31935] T. S.: Somatic mutations in the BRCA1 gene in sporadic
ovarian tumours. Nature
[31936] Genet. 9: 439-443, 1995.

[31937] 4513.

[31938] Miki, Y.; Swensen, J.; Shattuck-Eidens, D.; Futreal, P. A.;
Harshman,
[31939] K.; Tavtigian, S.; Liu, Q.; Cochran, C.; Bennett, L. M.; Ding,
W.;
[31940] Bell, R.; Rosenthal, J.; and 33 others: A strong candidate
for
[31941] the breast and ovarian cancer susceptibility gene BRCA1.

Science 266:
[31942] 66-71, 1994.

[31943]

[31944] 100. Moynahan, M. E.; Chiu, J. W.; Koller, B. H.; Jasin, M.: Brca1
[31945] controls homology-directed DNA repair. Molec. Cell 4: 511-518, 1999.

[31946]

[31947] 101. Narod, S.; Feunteun, J.; Lynch, H.; Watson, P.; Conway, T.; Lynch,
[31948] J.; Lenoir, G.: A breast-ovarian cancer locus on chromosome 17. (Abstract) Am.
[31949] J. Hum. Genet. 49 (suppl.): 352, 1991.

[31950]

[31951] 102. Narod, S.; Ford, D.; Devilee, P.; Barkardottir, R. B.; Eyfjord,
[31952] J.; Lenoir, G.; Serova, O.; Easton, D.; Goldgar, D.; Breast Cancer
[31953] Linkage Consortium: Genetic heterogeneity of breast-ovarian cancer
[31954] revisited. (Letter) Am. J. Hum. Genet. 57: 957-958, 1995.

[31955]

[31956] 103. Narod, S.; Tonin, P.; Lynch, H.; Watson, P.; Feunteun, J.; Lenoir,

[31957] G.: Histology of BRCA1-associated ovarian tumours. (Letter) Lancet 343:

[31958] 236 only, 1994.

[31959]

[31960] 104. Narod, S. A.; Feunteun, J.; Lynch, H. T.; Watson, P.; Conway,

[31961] T.; Lynch, J.; Lenoir, G. M.: Familial breast-ovarian cancer locus

[31962] on chromosome 17q12-q23. Lancet 338: 82-83, 1991.

[31963]

[31964] 105. Narod, S. A.; Ford, D.; Devilee, P.; Barkardottir, R. B.; Lynch,

[31965] H. T.; Smith, S. A.; Ponder, B. A. J.; Weber, B. L.; Garber, J. E.;

[31966] Birch, J. M.; Cornelis, R. S.; Kelsell, D. P.; and 17 others: An

[31967] evaluation of genetic heterogeneity in 145 breast-ovarian cancer families. Am.

[31968] J. Hum. Genet. 56: 254-264, 1995.

[31969]

[31970] 106. Narod, S. A.; Risch, H.; Moslehi, R.; Dorum, A.;
Neuhausen, S.;
[31971] Olsson, H.; Provencher, D.; Radice, P.; Evans, G.; Bishop,
S.; Brunet,
[31972] J.-S.; Ponder, B. A. J.; Hereditary Ovarian Cancer Clinical
Study
[31973] Group: Oral contraceptives and the risk of hereditary
ovarian cancer. New
[31974] Eng. J. Med. 339: 424-428, 1998.

[31975]

[31976] 107. Nastiuk, K. L.; Mansukhani, M.; Terry, M. B.; Ku-
laratne, P.;
[31977] Rubin, M. A.; Melamed, J.; Gammon, M. D.; Ittmann, M.;
Krolewski,
[31978] J. J.: Common mutations in BRCA1 and BRCA2 do not con-
tribute to early
[31979] prostate cancer in Jewish men. Prostate 40: 172-177,
1999.

[31980]

[31981] 108. Neuhausen, S. L.; Mazoyer, S.; Friedman, L.; Stratton, M.; Offit,
[31982] K.; Caligo, A.; Tomlinson, G.; Cannon-Albright, L.; Bishop, T.; Kelsell,
[31983] D.; Solomon, E.; Weber, B.; Couch, F.; Struewing, J.; Tonin, P.; Durocher,
[31984] F.; Narod, S.; Skolnick, M. H.; Lenoir, G.; Serova, O.; Ponder, B.;
[31985] Stoppa-Lyonnet, D.; Easton, D.; King, M.-C.; Goldgar, D. E.: Haplotype
[31986] and phenotype analysis of six recurrent BRCA1 mutations in 61 families:
[31987] results of an international study. Am. J. Hum. Genet. 58: 271-280,
[31988] 1996.

[31989]

[31990] 109. O'Connell, P.; Albertsen, H.; Matsunami, N.; Taylor, T.; Hundley,
[31991] J. E.; Johnson-Pais, T. L.; Reus, B.; Lawrence, E.; Ballard, L.; White,
[31992] R.; Leach, R. J.: A radiation hybrid map of the BRCA1 region. Am.

[31993] J. Hum. Genet. 54: 526-534, 1994.

[31994]

[31995] 110. Pace, P.; Johnson, M.; Tan, W. M.; Mosedale, G.; Sng, C.; Hoatlin,

[31996] M.; de Winter, J.; Joenje, H.; Gergely, F.; Patel, K. J.: FANCE:

[31997] the link between Fanconi anaemia complex assembly and activity. EMBO

[31998] J. 21: 3414-3423, 2002.

[31999]

[32000] 111. Paull, T. T.; Cortez, D.; Bowers, B.; Elledge, S. J.; Gellert,

[32001] M.: Direct DNA binding by Brca1. Proc. Nat. Acad. Sci. 98: 6086-6091,

[32002] 2001.

[32003]

[32004] 112. Petrij-Bosch, A.; Peelen, T.; van Vliet, M.; van Eijk, R.; Olmer,

[32005] R.; Drusedau, M.; Hogervorst, F. B. L.; Hageman, S.; Arts, P. J. W.;

[32006] Ligtenberg, M. J. L.; Meijers-Heijboer, H.; Klijn, J. G. M.; Vasen,

[32007] H. F. A.; Cornelisse, C. J.; van't Veer, L. J.; Bakker, E.; van Ommen,
[32008] G.-J. B.; Devilee, P.: BRCA1 genomic deletions are major founder
[32009] mutations in Dutch breast cancer patients. Nature Genet. 17: 341-345,
[32010] 1997.

[32011]

[32012] 113. Phelan, C. M.; Rebbeck, T. R.; Weber, B. L.; Devilee, P.; Ruttledge,
[32013] M. H.; Lynch, H. T.; Lenoir, G. M.; Stratton, M. R.; Easton, D. F.;
[32014] Ponder, B. A. J.; Cannon-Albright, L.; Larsson, C.; Goldgar, D. E.;
[32015] Narod, S. A.: Ovarian cancer risk in BRCA1 carriers is modified by
[32016] the HRAS1 variable number of tandem repeat (VNTR) lo-cus. Nature Genet. 12:
[32017] 309-311, 1996.

[32018]

[32019] 114. Piver, M. S.; Baker, T. R.; Jishi, M. F.; Sandecki, A. M.; Tsukada,

[32020] Y.; Natarajan, N.; Mettlin, C. J.; Blake, C. A.: Familial ovarian
[32021] cancer: a report of 658 families from the Gilda Radner Familial Ovarian
[32022] Cancer Registry 1981-1991. Cancer 71: 582-588, 1993.
[32023]
[32024] 115. Puget, N.; Gad, S.; Perrin-Vidoz, L.; Sinilnikova, O. M.; Stoppa-Lyonnet,
[32025] D.; Lenoir, G. M.; Mazoyer, S.: Distinct BRCA1 rearrangements involving
[32026] the BRCA1 pseudogene suggest the existence of a recombination hot
[32027] spot. Am. J. Hum. Genet. 70: 858-865, 2002.
[32028]
[32029] 116. Puget, N.; Sinilnikova, O. M.; Stoppa-Lyonnet, D.; Audoynaud,
[32030] C.; Pages, S.; Lynch, H. T.; Goldgar, D.; Lenoir, G. M.; Mazoyer,
[32031] S.: An Alu-mediated 6-kb duplication in the BRCA1 gene: a new founder
[32032] mutation? Am. J. Hum. Genet. 64: 300-302, 1999.

[32033]

[32034] 117. Puget, N.; Stoppa-Lyonnet, D.; Sinilnikova, O. M.; Pages, S.;

[32035] Lynch, H. T.; Lenoir, G. M.; Mazoyer, S.: Screening for germline

[32036] rearrangements and regulatory mutations in BRCA1 led to the identification

[32037] of four new deletions. Cancer Res. 59: 455-461, 1999.

[32038]

[32039] 118. Ramus, S. J.; Friedman, L. S.; Gayther, S. A.; Ponder, B. A.

[32040] J.; Bobrow, L. G.; van der Looji, M.; Papp, J.; Olah, E.: A breast/ovarian

[32041] cancer patient with germline mutations in both BRCA1 and BRCA2. (Letter) Nature

[32042] Genet. 15: 14-15, 1997.

[32043]

[32044] 119. Rebbeck, T. R.; Couch, F. J.; Kant, J.; Calzone, K.; De-Shano,

[32045] M.; Peng, Y.; Chen, K.; Garber, J. E.; Weber, B. L.: Genetic heterogeneity

[32046] in hereditary breast cancer: role of BRCA1 and BRCA2. Am. J. Hum.

[32047] Genet. 59: 547-553, 1996.

[32048]

[32049] 120. Rebbeck, T. R.; Lynch, H. T.; Neuhausen, S. L.; Narod, S. A.;

[32050] van't Veer, L.; Garber, J. E.; Evans, G.; Isaacs, C.; Daly, M. B.;

[32051] Matloff, E.; Olopade, O. I.; Weber, B. L. Prophylactic oophorectomy

[32052] in carriers of BRCA1 or BRCA2 mutations. New Eng. J. Med. 346: 1616-1622,

[32053] 2002.

[32054]

[32055] 121. Risch, H. A.; McLaughlin, J. R.; Cole, D. E. C.; Rosen, B.; Bradley,

[32056] L.; Kwan, E.; Jack, E.; Vesprini, D. J.; Kuperstein, G.; Abra-hamson,

[32057] J. L. A.; Fan, I.; Wong, B.; Narod, S. A.: Prevalence and pen-etrance

[32058] of germline BRCA1 and BRCA2 mutations in a population series of 649

[32059] women with ovarian cancer. Am. J. Hum. Genet. 68:
700-710, 2001.

[32060]

[32061] 122. Roa, B. B.; Boyd, A. A.; Volcik, K.; Richards, C. S.:
Ashkenazi

[32062] Jewish population frequencies for common mutations in
BRCA1 and BRCA2. Nature

[32063] Genet. 14: 185-187, 1996.

[32064]

[32065] 123. Ruffner, H.; Jiang, W.; Craig, A. G.; Hunter, T.; Verma,
I. M.

[32066] : BRCA1 is phosphorylated at serine 1497 in vivo at a cy-
clin-dependent

[32067] kinase 2 phosphorylation site. Molec. Cell. Biol. 19:
4843-4854,

[32068] 1999.

[32069]

[32070] 124. Sarantaus, L.; Huusko, P.; Eerola, H.; Launonen, V.;
Vehmanen,

[32071] P.; Rapakko, K.; Gillanders, E.; Syrjakoski, K.; Kainu, T.;
Vahteristo,

[32072] P.; Krahe, R.; Paakkonen, K.; and 14 others: Multiple founder effects

[32073] and geographical clustering of BRCA1 and BRCA2 families in Finland. Europ.

[32074] J. Hum. Genet. 8: 757-763, 2000.

[32075]

[32076] 125. Sarantaus, L.; Vahteristo, P.; Bloom, E.; Tamminen, A.; Unkila-Kallio,

[32077] L.; Butzow, R.; Nevanlinna, H. BRCA1 and BRCA2 mutations among

[32078] 233 unselected Finnish ovarian carcinoma patients. Europ. J. Hum.

[32079] Genet. 9: 424-430, 2001.

[32080]

[32081] 126. Schorge, J. O.; Muto, M. G.; Welch, W. R.; Bandera, C. A.; Rubin,

[32082] S. C.; Bell, D. A.; Berkowitz, R. S.; Mok, S. C.: Molecular evidence

[32083] for multifocal papillary serous carcinoma of the peritoneum in patients

[32084] with germline BRCA1 mutations. J. Nat. Cancer Inst. 90: 841-845,

[32085] 1998.

[32086]

[32087] 127. Schrock, E.; Badger, P.; Larson, D.; Erdos, M.; Wyn-
shaw-Boris,
[32088] A.; Ried, T.; Brody, L.: The murine homolog of the human
breast and
[32089] ovarian cancer susceptibility gene Brca1 maps to mouse
chromosome
[32090] 11D. Hum. Genet. 97: 256-259, 1996.

[32091]

[32092] 128. Schuyer, M.; Berns, E. M.: Is TP53 dysfunction re-
quired for
[32093] BRCA1-associated carcinogenesis? Molec. Cell. Endocrin.
155: 143-152,
[32094] 1999.

[32095]

[32096] 129. Scully, R.; Anderson, S. F.; Chao, D. M.; Wei, W.; Ye,
L.; Young,
[32097] R. A.; Livingston, D. M.; Parvin, J. D.: BRCA1 is a compo-
nent of
[32098] the RNA polymerase II holoenzyme. Proc. Nat. Acad. Sci.

94: 5605-5610,
[32099] 1997.

[32100]

[32101] 130. Scully, R.; Ganesan, S.; Brown, M.; De Caprio, J. A.; Cannistra,
[32102] S. A.; Feunteun, J.; Schnitt, S.; Livingston, D. M.: Location of
[32103] BRCA1 in human breast and ovarian cancer cells. Science 272: 123-125,
[32104] 1996.

[32105]

[32106] 131. Scully, R.; Ganesan, S.; Vlasakova, K.; Chen, J.; So-colovsky,
[32107] M.; Livingston, D. M.: Genetic analysis of BRCA1 function in a defined
[32108] tumor cell line. Molec. Cell 4: 1093-1099, 1999.

[32109]

[32110] 132. Serova, O.; Montagna, M.; Torchard, D.; Narod, S. A.; Tonin,
[32111] P.; Sylla, B.; Lynch, H. T.; Feunteun, J.; Lenoir, G. M.: A high

[32112] incidence of BRCA1 mutations in 20 breast-ovarian cancer families. Am.

[32113] J. Hum. Genet. 58: 42-51, 1996.

[32114]

[32115] 133. Simard, J.; Feunteun, J.; Lenoir, G.; Tonin, P.; Nor-

mand, T.;

[32116] The, V. L.; Vivier, A.; Lasko, D.; Morgan, K.; Rouleau, G. A.; Lynch,

[32117] H.; Labrie, F.; Narod, S. A.: Genetic mapping of the breast-ovarian

[32118] cancer syndrome to a small interval on chromosome 17q12-21: exclusion

[32119] of candidate genes EDH17B2 and RARA. Hum. Molec. Genet. 2: 1193-1199,

[32120] 1993.

[32121]

[32122] 134. Simard, J.; Tonin, P.; Durocher, F.; Morgan, K.; Rom-

mens, J.;

[32123] Gingras, S.; Samson, C.; Leblanc, J.-F.; Belanger, C.; Dion, F.; Liu,

[32124] Q.; Skolnick, M.; Goldgar, D.; Shattuck-Eidens, D.; Labrie, F.; Narod,

[32125] S. A.: Common origins of BRCA1 mutations in Canadian breast and ovarian
[32126] cancer families. Nature Genet. 8: 392-398, 1994.

[32127]

[32128] 135. Skolnick, M. H.; Cannon-Albright, L. A.; Goldgar, D. E.; Ward,
[32129] J. H.; Marshall, C. J.; Schumann, G. B.; Hogle, H.; McWhorter, W.
[32130] P.; Wright, E. C.; Tran, T. D.; Bishop, D. T.; Kushner, J. P.; Eyre,
[32131] H. J.: Inheritance of proliferative breast disease in breast cancer
[32132] kindreds. Science 250: 1715-1720, 1990.

[32133]

[32134] 136. Smith, S. A.; Easton, D. F.; Evans, D. G. R.; Ponder, B. A. J.
[32135] : Allele losses in the region 17q12-21 in familial breast and ovarian
[32136] cancer involve the wild-type chromosome. Nature Genet. 2: 128-131,
[32137] 1992.

[32138]

[32139] 137. Smith, T. M.; Lee, M. K.; Szabo, C. I.; Jerome, N.;
McEuen, M.;

[32140] Taylor, M.; Hood, L.; King, M.-C.: Complete genomic se-
quence and

[32141] analysis of 117 kb of human DNA containing the gene
BRCA1. Genome

[32142] Res. 6: 1029-1049, 1996.

[32143]

[32144] 138. Sobol, H.; Mazoyer, S.; Narod, S. A.; Smith, S. A.;
Black, D.

[32145] M.; Kerbrat, P.; Jamot, B.; Solomon, E.; Ponder, B. A. J.;
Guerin,

[32146] D.: Genetic heterogeneity of early-onset familial breast
cancer. Hum.

[32147] Genet. 89: 381-383, 1992.

[32148]

[32149] 139. Steeg, P. S.: Granin expectations in breast cancer?
Nature

[32150] Genet. 12: 223-225, 1996.

[32151]

[32152] 140. Stoppa-Lyonnet, D.; Fricker, J. P.; Essioux, L.; Pages, S.; Limacher,
[32153] J. M.; Sobol, H.; Laurent-Puig, P.; Thomas, G.: Segregation of two
[32154] BRCA1 mutations in a single family. (Letter) Am. J. Hum. Genet. 59:
[32155] 479-481, 1996.

[32156]

[32157] 141. Stratton, M. R.; Ford, D.; Neuhasen, S.; Seal, S.; Wooster, R.;
[32158] Friedman, L. S.; King, M.-C.; Egilsson, V.; Devilee, P.; Mc-Manus,
[32159] R.; Daly, P. A.; Smyth, E.; Ponder, B. A. J.; Peto, J.; Cannon-Albright,
[32160] L.; Easton, D. F.; Goldgar, D. E.: Familial male breast cancer is
[32161] not linked to the BRCA1 locus on chromosome 17q. Nature Genet. 7:
[32162] 103-107, 1994.

[32163]

[32164] 142. Struewing, J. P.; Abeliovich, D.; Peretz, T.; Avishai, N.; Kaback,

[32165] M. M.; Collins, F. S.; Brody, L. C.: The carrier frequency of the

[32166] BRCA1 185delAG mutation is approximately 1 percent in Ashkenazi Jewish

[32167] individuals. Nature Genet. 11: 198-200, 1995.

[32168]

[32169] 143. Struewing, J. P.; Brody, L. C.; Erdos, M. R.; Kase, R. G.; Giambarresi,

[32170] T. R.; Smith, S. A.; Collins, F. S.; Tucker, M.A.: Detection of eight

[32171] BRCA1 mutations in 10 breast/ovarian cancer families, in-cluding 1

[32172] family with male breast cancer. Am. J. Hum. Genet. 57: 1-7, 1995.

[32173]

[32174] 144. Struewing, J. P.; Hartge, P.; Wacholder, S.; Baker, S. M.; Berlin,

[32175] M.; McAdams, M.; Timmerman, M. M.; Brody, L. C.; Tucker, M. A.: The

[32176] risk of cancer associated with specific mutations of BRCA1 and BRCA2

[32177] among Ashkenazi Jews. New Eng. J. Med. 336:

1401-1408, 1997.

[32178]

[32179] 145. Szabo, C. I.; King, M.-C.: Population genetics of BRCA1 and

[32180] BRCA2. (Editorial) Am. J. Hum. Genet. 60: 1013-1020, 1997.

[32181]

[32182] 146. Tesoriero, A.; Andersen, C.; Southey, M.; Somers, G.; McKay,

[32183] M.; Armes, J.; McCredie, M.; Giles, G.; Hopper, J. L.; Ven-

ter, D.

[32184] : De novo BRCA1 mutation in a patient with breast cancer and an inherited

[32185] BRCA2 mutation. (Letter) Am. J. Hum. Genet. 65: 567-569, 1999.

[32186]

[32187] 147. Thompson, M. E.; Jensen, R. A.; Obermiller, P. S.; Page, D. L.;

[32188] Holt, J. T.: Decreased expression of BRCA1 accelerates growth and

[32189] is often present during sporadic breast cancer progres-

sion. Nature

[32190] Genet. 9: 444-450, 1995.

[32191]

[32192] 148. Tonin, P.; Moslehi, R.; Green, R.; Rosen, B.; Cole, D.; Boyd,

[32193] N.; Cutler, C.; Margolese, R.; Carter, R.; McGillivray, B.; Ives,

[32194] E.; Labrie, F.; Gilchrist, D.; Morgan, K.; Simard, J.; Narod, S. A.

[32195] : Linkage analysis of 26 Canadian breast and breast-ovarian cancer

[32196] families. Hum. Genet. 95: 545-550, 1995.

[32197]

[32198] 149. Tonin, P. N.; Mes-Masson, A.-M.; Futreal, P. A.; Morgan, K.;

[32199] Mahon, M.; Foulkes, W. D.; Cole, D. E. C.; Provencher, D.; Ghadirian,

[32200] P.; Narod, S. A.: Founder BRCA1 and BRCA2 mutations in French Canadian

[32201] breast ovarian cancer families. Am. J. Hum. Genet. 63: 1341-1351,

[32202] 1998.

[32203]

[32204] 150. Vallon-Christersson, J.; Cayanan, C.; Haraldsson, K.; Loman,

[32205] N.; Bergthorsson, J. T.; Brondum-Nielsen, K.; Gerdes, A.-M.; Moller,

[32206] P.; Kristoffersson, U.; Olsson, H.; Borg, A.; Monteiro, A. N. A.:

[32207] Functional analysis of BRCA1 C-terminal missense muta-tions identified

[32208] in breast and ovarian cancer families. Hum. Molec. Genet. 10: 353-360,

[32209] 2001.

[32210]

[32211] 151. van Orsouw, N. J.; Dhanda, R. K.; Elhaji, Y.; Narod, S. A.; Li,

[32212] F. P.; Eng, C.; Vijg, J.: A highly accurate, low cost test for BRCA1

[32213] mutations. J. Med. Genet. 36: 747-753, 1999.

[32214]

[32215] 152. Vazina, A.; Baniel, J.; Yaacobi, Y.; Shtriker, A.; Engel-stein,

[32216] D.; Leibovitz, I.; Zehavi, M.; Sidi, A. A.; Ramon, Y.; Tischler, T.;

[32217] Livne, P. M.; Friedman, E.: The rate of the founder Jewish mutations

[32218] in BRCA1 and BRCA2 in prostate cancer patients in Israel. Brit. J.

[32219] Cancer 83: 463-466, 2000.

[32220]

[32221] 153. Vega, A.; Campos, B.; Bressac-de Paillerets, B.; Bond, P.; Janin,

[32222] N.; Douglas, F.; Domenech, M.; Baena, M.; Pericay, C.; Alonso, M.

[32223] C.; Carracedo, A.; Baiget, M.; Diez, O.: The R71G BRCA1 is a founder

[32224] Spanish mutation and leads to aberrant splicing of the transcript. Hum.

[32225] Mutat. 17: 520-521, 2001.

[32226]

[32227] 154. Vega, A.; Torres, M.; Martinez, J. I.; Ruiz-Ponte, C.; Barros,

[32228] F.; Carracedo, A.: Analysis of BRCA1 and BRCA2 in breast and breast/ovarian

[32229] cancer families shows population substructure in the Iberian peninsula. Ann.
[32230] Hum. Genet. 66: 29-36, 2002.

[32231]

[32232] 155. Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin,
[32233] J.: BASC, a super complex of BRCA1-associated proteins involved in
[32234] the recognition and repair of aberrant DNA structures. Genes Dev. 14:
[32235] 927-939, 2000.

[32236]

[32237] 156. Welcsh, P. L.; King, M.-C.: BRCA1 and BRCA2 and the genetics
[32238] of breast and ovarian cancer. Hum. Molec. Genet. 10: 705-713, 2001.

[32239]

[32240] 157. Whittemore, A. S.; Harris, R.; Itnyre, J.; Collaborative Ovarian
[32241] Cancer Group: Characteristics relating to ovarian cancer risk: collaborative

[32242] analysis of 12 U.S. case-control studies. II. Invasive epithelial

[32243] ovarian cancers in white women. Am. J. Epidemiol. 136: 1184-1203,

[32244] 1992.

[32245]

[32246] 158. Wilson, C. A.; Ramos, L.; Villasenor, M. R.; Anders, K. H.; Press,

[32247] M. F.; Clarke, K.; Karlan, B.; Chen, J.-J.; Scully, R.; Livingston,

[32248] D.; Zuch, R. H.; Kanter, M. H.; Cohen, S.; Calzone, F. J.; Slamon,

[32249] D. J.: Localization of human BRCA1 and its loss in high-grade, non-inherited

[32250] breast carcinomas. Nature Genet. 21: 236-240, 1999.

[32251]

[32252] 159. Wu, L. C.; Wang, Z. W.; Tsan, J. T.; Spillman, M. A.; Phung,

[32253] A.; Xu, X. L.; Yang, M.-C. W.; Hwang, L.-Y.; Bowcock, A. M.; Baer,

[32254] R.: Identification of a RING protein that can interact in vivo with

[32255] the BRCA1 gene product. Nature Genet. 14: 430-440, 1996.

[32256]

[32257] 160. Xu, X.; Qiao, W.; Linke, S. P.; Cao, L.; Li, W.-M.; Furth, P.

[32258] A.; Harris, C. C.; Deng, C.-X.: Genetic interactions between tumor

[32259] suppressors Brca1 and p53 in apoptosis, cell cycle and tumorigenesis. Nature

[32260] Genet. 28: 266-271, 2001.

[32261]

[32262] 161. Xu, X.; Weaver, Z.; Linke, S. P.; Li, C.; Gotay, J.; Wang, X.-W.;

[32263] Harris, C. C.; Ried, T.; Deng, C.-X.: Centrosome amplification and

[32264] a defective G(2)-M cell cycle checkpoint induce genetic instability

[32265] in BRCA1 exon 11 isoform-deficient cells. Molec. Cell 3: 389-395,

[32266] 1999.

[32267]

[32268] 162. Yarden, R. I.; Brody, L. C.: BRCA1 interacts with components
[32269] of the histone deacetylase complex. Proc. Nat. Acad. Sci. 96: 4983-4988,
[32270] 1999.

[32271]

[32272] 163. Yarden, R. I.; Pardo-Reoyo, S.; Sgagias, M.; Cowan, K. H.; Brody,
[32273] L. C.: BRCA1 regulates the G2/M checkpoint by activating Chk1 kinase
[32274] upon DNA damage. Nature Genet. 30: 265-269, 2002.

[32275]

[32276] 164. Zhang, S.; Grosse, F.: Domain structure of human nuclear DNA
[32277] helicase II (RNA helicase A). J. Biol. Chem. 272: 11487-11494, 1997.

[32278]

[32279] 165. Zheng, L.; Pan, H.; Li, S.; Flesken-Nikitin, A.; Chen, P.-L.;
[32280] Boyer, T. G.; Lee, W.-H.: Sequence-specific transcriptional corepressor

[32281] function for BRCA1 through a novel zinc finger protein, ZBRK1. Molec.

[32282] Cell 6: 757-768, 2000.

[32283]

[32284] 166. Zhong, Q.; Chen, C.-F.; Li, S.; Chen, Y.; Wang, C.-C.; Xiao,

[32285] J.; Chen, P.-L.; Sharp, Z. D.; Lee, W.-H.: Association of BRCA1 with

[32286] the hRad50-hMre11-p95 complex and the DNA damage response. Science 285:

[32287] 747-750, 1999.

[32288] 4514.

[32289] Ponnambalam, S.; Jackson, A. P.; LeBeau, M. M.; Pravtcheva, D.;

[32290] Ruddle, F. H.; Alibert, C.; Parham, P.: Chromosomal loca-tion and

[32291] some structural features of human clathrin light-chain genes (CLTA

[32292] and CLTB). Genomics 24: 440-444, 1994.

[32293] 4515.

[32294] Andersen, T. I.; Wooster, R.; Laake, K.; Collins, N.; Warren,

W.;

[32295] Skrede, M.; Eeles, R.; Tveit, K. M.; Johnston, S. R. D.; Dowsett,

[32296] M.; Olsen, A. O.; Moller, P.; Stratton, M. R.; Borresen-Dale, A.-L.

[32297] : Screening for ESR mutations in breast and ovarian cancer patients. Hum.

[32298] Mutat. 9: 531-536, 1997.

[32299] 4516.

[32300] Auboeuf, D.; Honig, A.; Berget, S. M.; O'Malley, B. W.: Co-ordinate

[32301] regulation of transcription and splicing by steroid receptor coregulators. Science 298:

[32302] 416-419, 2002.

[32303] 4517.

[32304] Balleine, R. L.; Hunt, S. M. N.; Clarke, C. L.: Coexpression of

[32305] alternatively spliced estrogen and progesterone receptor transcripts

[32306] in human breast cancer. J. Clin. Endocr. Metab. 84: 1370-1377, 1999.

[32307] 4518.

[32308] Becherini, L.; Gennari, L.; Masi, L.; Mansani, R.; Massart, F.;

[32309] Morelli, A.; Falchetti, A.; Gonnelli, S.; Fiorelli, G.; Tanini, A.;

[32310] Brandi, M. L.: Evidence of a linkage disequilibrium be-

tween polymorphisms

[32311] in the human estrogen receptor-alpha gene and their re-

lationship to

[32312] bone mass variation in postmenopausal Italian women. Hum. Molec.

[32313] Genet. 9: 2043-2050, 2000.

[32314] 4519.

[32315] Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen re-

ceptors

[32316] alpha and beta are differentially expressed in developing human bone. J.

[32317] Clin. Endocr. Metab. 86: 2309-2314, 2001.

[32318] 4520.

[32319] Castagnoli, A.; Maestri, I.; Bernardi, F.; Del Senno, L.: PvuII

[32320] RFLP inside the human estrogen receptor gene. Nucleic Acids Res. 15:

[32321] 866 only, 1987.

[32322] 4521.

[32323] Chaidarun, S. S.; Alexander, J. M.: A tumor-specific truncated

[32324] estrogen receptor splice variant enhances estrogen-stimulated gene

[32325] expression. Molec. Endocr. 12: 1355-1366, 1998.

[32326] 4522.

[32327] Chiang, C.-H.; Cheng, K. W.; Igarashi, S.; Nathwani, P. S.; Leung,

[32328] P. C. K.: Hormonal regulation of estrogen receptor alpha and beta

[32329] gene expression in human granulosa-luteal cells in vitro. J. Clin.

[32330] Endocr. Metab. 85: 3828-3839, 2000.

[32331] 4523.

[32332] Clark, G. M.; McGuire, W. L.: Steroid receptors and other prognostic

[32333] factors in primary breast cancer. Semin. Oncol. 15 (suppl. 1): 20-25,

[32334] 1988.

[32335] 4524.

[32336] Couse, J. F.; Hewitt, S. C.; Bunch, D. O.; Sar, M.; Walker, V.

[32337] R.; Davis, B. J.; Korach, K. S.: Postnatal sex reversal of the ovaries

[32338] in mice lacking estrogen receptors alpha and beta. Science 286:

[32339] 2328-2331, 1999.

[32340] 4525.

[32341] Davis, V. L.; Chan, C.-C.; Schoen, T. J.; Couse, J. F.; Chader,

[32342] G. J.; Korach, K. S.: An estrogen receptor repressor in-

duces cataract

[32343] formation in transgenic mice. Proc. Nat. Acad. Sci. 99:

9427-9432,

[32344] 2002.

[32345] 4526.

[32346] Esmaeli, B.; Harvey, J. T.; Hewlett, B.: Immunohistochemi-

cal

[32347] evidence for estrogen receptors in meibomian glands.

Ophthalmology 107:

[32348] 180-184, 2000.

[32349] 4527.

[32350] Fan, S.; Wang, J.-A.; Yuan, R.; Ma, Y.; Meng, Q.; Erdos, M. R.;

[32351] Pestell, R. G.; Yuan, F.; Auborn, K. J.; Goldberg, I. D.; Rosen, E.

[32352] M.: BRCA1 inhibition of estrogen receptor signaling in transfected

[32353] cells. Science 284: 1354-1356, 1999.

[32354] 4528.

[32355] Fuqua, S. A. W.; Chamness, G. C.; McGuire, W. L.: Estrogen receptor

[32356] mutations in breast cancer. J. Cell. Biochem. 51: 135-139, 1993.

[32357] 4529.

[32358] Gosden, J. R.; Middleton, P. G.; Rout, D.: Localization of the

[32359] human oestrogen receptor gene to chromosome 6q24-q27 by in situ hybridization. Cytogenet.

[32360] Cell Genet. 43: 218-220, 1986.

[32361] 4530.

[32362] Green, S.; Walter, P.; Kumar, V.; Krust, A.; Bornert, J.-M.;

Argos,

[32363] P.; Chambon, P.: Human oestrogen receptor cDNA: sequence, expression

[32364] and homology to v-erb-A. Nature 320: 134-139, 1986.

[32365] 4531.

[32366] Greene, G. L.; Gilna, P.; Waterfield, M.; Baker, A.; Hort, Y.;

[32367] Shine, J.: Sequence and expression of human estrogen receptor complementary

[32368] DNA. Science 231: 1150-1154, 1986.

[32369] 4532.

[32370] Heine, P. A.; Taylor, J. A.; Iwamoto, G. A.; Lubahn, D. B.; Cooke,

[32371] P. S.: Increased adipose tissue in male and female estrogen receptor-alpha

[32372] knockout mice. Proc. Nat. Acad. Sci. 97: 12729-12734, 2000.

[32373] 4533.

[32374] Herrington, D. M.; Howard, T. D.; Hawkins, G. A.; Reboussin, D.

[32375] M.; Xu, J.; Zheng, S. L.; Brosnihan, K. B.; Meyers, D. A.; Bleecker,

[32376] E. R.: Estrogen-receptor polymorphisms and effects of es-
trogen replacement
[32377] on high-density lipoprotein cholesterol in women with
coronary disease. New
[32378] Eng. J. Med. 346: 967-974, 2002.

[32379] 4534.

[32380] Issa, J.-P. J.; Ottaviano, Y. L.; Celano, P.; Hamilton, S. R.;

[32381] Davidson, N. E.; Baylin, S. B.: Methylation of the oestrogen
receptor
[32382] CpG island links ageing and neoplasia in human colon.
Nature Genet. 7:
[32383] 536-540, 1994.

[32384] 4535.

[32385] Jeltsch, J. M.; Roberts, M.; Schatz, C.; Garnier, J. M.; Brown,

[32386] A. M. C.; Chambon, P.: Structure of the human oestrogen-
responsive
[32387] gene pS2. Nucleic Acids Res. 15: 1401-1414, 1987.

[32388] 4536.

[32389] Bing, D. H.; Almeda, S.; Isliker, H.; Lahav, J.; Hynes, R. O.:

[32390] Fibronectin binds to the C1q component of complement.

Proc. Nat.
[32391] Acad. Sci. 79: 4198-4201, 1982.

[32392] 4537.

[32393] Bittner, M.; Meltzer, P.; Chen, Y.; Jiang, Y.; Seftor, E.; Hen-
drix,
[32394] M.; Radmacher, M.; Simon, R.; Yakhini, Z.; Ben-Dor, A.;
Sampas, N.;
[32395] Dougherty, E.; and 16 others: Molecular classification of
cutaneous
[32396] malignant melanoma by gene expression profiling. Nature
406: 536-540,
[32397] 2000.

[32398] 4538.

[32399] Clark, E. A.; Golub, T. R.; Lander, E. S.; Hynes, R. O.: Ge-
nomic
[32400] analysis of metastasis reveals an essential role for RhoC.
Nature 406:
[32401] 532-535, 2000.

[32402] 4539.

[32403] Clemmensen, I.: Fibronectin and its role in connective tis-
sue

[32404] diseases. (Editorial) Europ. J. Clin. Invest. 11: 145-146, 1981.

[32405] 4540.

[32406] Croce, C. M.: Personal Communication. Philadelphia, Pa. 1/12/1983.

[32407] 4541.

[32408] Eun, C. K.; Klinger, H. P.: Human chromosome 11 affects the expression

[32409] of fibronectin fibers in human-times-mouse cell hybrids. Cytogenet.

[32410] Cell Genet. 27: 57-65, 1980.

[32411] 4542.

[32412] Gutman, A.; Kornblihtt, A. R.: Identification of a third re-gion

[32413] of cell-specific alternative splicing in human fibronectin mRNA. Proc.

[32414] Nat. Acad. Sci. 84: 7179-7182, 1987.

[32415] 4543.

[32416] Henry, I.; Jeanpierre, M.; Weil, D.; Grzeschik, K. H.; Ramirez,

[32417] F.; Junien, C.: The structural gene for fibronectin (FN)

maps to
[32418] 2q323-qter. (Abstract) Cytogenet. Cell Genet. 40: 650 only, 1985.
[32419] 4544.
[32420] Hirano, H.; Yamada, Y.; Sullivan, M.; de Crombrugghe, B.; Pastan,
[32421] I.; Yamada, K. M.: Isolation of genomic DNA clones spanning the entire
[32422] fibronectin gene. Proc. Nat. Acad. Sci. 80: 46-50, 1983.
[32423] 4545.
[32424] Jhanwar, S. C.; Jensen, J. T.; Kaelbling, M.; Chaganti, R. S.
[32425] K.; Klinger, H. P.: In situ localization of human fibronectin (FN)
[32426] genes to chromosome regions 2p14-p16, 2q34-q36, and 11q12.1-q13.5
[32427] in germ line cells, but to chromosome 2 sites only in somatic cells. Cytogenet.
[32428] Cell Genet. 41: 47-53, 1986.
[32429] 4546.
[32430] Koch, G. A.; Schoen, R. C.; Klebe, R. J.; Shows, T. B.: Assignment

[32431] of a fibronectin gene to human chromosome 2 using monoclonal antibodies. Exp.

[32432] Cell Res. 141: 293-302, 1982.

[32433] 4547.

[32434] Kornblihtt, A. R.; Umezawa, K.; Vibe-Pedersen, K.; Baralle, F.

[32435] E.: Primary structure of human fibronectin: differential splicing

[32436] may generate at least 10 polypeptides from a single gene. EMBO J. 4:

[32437] 1755-1759, 1985.

[32438] 4548.

[32439] Kornblihtt, A. R.; Vibe-Pedersen, K.; Baralle, F. E.: Isolation

[32440] and characterization of cDNA clones for human and bovine fibronectins. Proc.

[32441] Nat. Acad. Sci. 80: 3218-3222, 1983.

[32442] 4549.

[32443] Kornblihtt, A. R.; Vibe-Pedersen, K.; Baralle, F. E.: Human fibronectin:

[32444] molecular cloning evidence for two mRNA species differing by an internal

[32445] segment coding for a structural domain. EMBO J. 3:
221-226, 1984.
[32446] 4550.

[32447] Kurkinen, M.; Vartio, T.; Vaheri, A.: Polypeptides of human plasma
[32448] fibronectin are similar but not identical. Biochim. Biophys. Acta 624:
[32449] 490-498, 1980.

[32450] 4551.

[32451] Matsuura, H.; Takio, K.; Titani, K.; Greene, T.; Levery, S. B.;

[32452] Salyan, M. E. K.; Hakomori, S.: The oncofetal structure of human
[32453] fibronectin defined by monoclonal antibody FDC-6. J. Biol. Chem. 263:
[32454] 3314-3322, 1988.

[32455] 4552.

[32456] McDonagh, J.: Fibronectin: a molecular glue. Arch. Path. Lab.
[32457] Med. 105: 393-396, 1981.

[32458] 4553.

[32459] Mosesson, M. W.; Amrani, D. L.: The structure and biologic activities
[32460] of plasma fibronectin. Blood 56: 145-158, 1980.

[32461] 4554.

[32462] Odermatt, E.; Tamkun, J. W.; Hynes, R. O.: Repeating modular
[32463] structure of the fibronectin gene: relationship to protein structure
[32464] and subunit variation. Proc. Nat. Acad. Sci. 82: 6571-6575, 1985.

[32465] 4555.

[32466] Owerbach, D.; Doyle, D.; Shows, T. B.: Genetics of the large,
[32467] external, transformation-sensitive (LETS) protein: assignment of a
[32468] gene coding for expression of LETS to human chromosome 8. Proc. Nat.
[32469] Acad. Sci. 75: 5640-5644, 1978.

[32470] 4556.

[32471] Prowse, K.; Tricoli, J.; Klebe, R.; Shows, T.: Chromosome 2 assignment

[32472] of the structural gene for fibronectin (FN) using a cloned probe.

[32473] (Abstract) Cytogenet. Cell Genet. 40: 724 only, 1985.

[32474] 4557.

[32475] Prowse, K. R.; Tricoli, J. V.; Klebe, R. J.; Shows, T. B.: Assignment

[32476] of the human fibronectin structural gene to chromosome 2. Cytogenet.

[32477] Cell Genet. 41: 42-46, 1986.

[32478] 4558.

[32479] Rennard, S. I.; Church, R. L.; Rohrbach, D. H.; Shupp, D. E.;

[32480] Abe, S.; Hewitt, A. T.; Murray, J. C.; Martin, G. R.: Localization

[32481] of the human fibronectin (FN) gene on chromosome 8 by a specific enzyme

[32482] immunoassay. Biochem. Genet. 19: 551-566, 1981.

[32483] 4559.

[32484] Ridley, A.: Molecular switches in metastasis. Nature 406: 466-467,

[32485] 2000.

[32486] 4560.

[32487] Li, Q. J.; Ashraf, M. F.; Shen, D.; Green, W. R.; Stark, W. J.;

[32488] Chan, C.-C; O'Brien, T. P.: The role of apoptosis in the pathogenesis

[32489] of Fuchs endothelial dystrophy of the cornea. Arch. Oph-thal. 119:

[32490] 1597-1604, 2001.

[32491] 4561.

[32492] Paigen, K.: Acid hydrolases as models of genetic control. Ann.

[32493] Rev. Genet. 13: 417-466, 1979.

[32494] 4562.

[32495] McCurley, R. S.; Recinos, A., III; Olsen, A. S.; Gingrich, J. C.;

[32496] Szczepaniak, D.; Cameron, H. S.; Krauss, R.; Weston, B. W.: Physical

[32497] maps of human alpha(1,3)fucosyltransferase genes FUT3-FUT6 on chromosomes

[32498] 19p13.3 and 11q21. Genomics 26: 142-146, 1995.

[32499] 4563.

[32500] Weston, B. W.; Nair, R. P.; Larsen, R. D.; Lowe, J. B.: Isola-

tion
[32501] of a novel human alpha(1,3)fucosyltransferase gene and molecular comparison
[32502] to the human Lewis blood group alpha(1,3/1,4)fucosyltransferase gene:
[32503] syntenic, homologous, nonallelic genes encoding enzymes with distinct
[32504] acceptor substrate specificities. J. Biol. Chem. 267: 4152-4160,
[32505] 1992.

[32506] 4564.

[32507] Weston, B. W.; Smith, P. L.; Kelly, R. J.; Lowe, J. B.: Molecular
[32508] cloning of a fourth member of a human alpha(1,3)fucosyltransferase
[32509] gene family: multiple homologous sequences that determine expression
[32510] of the Lewis x, sialyl Lewis x, and difucosyl sialyl Lewis x epitopes. J.
[32511] Biol. Chem. 267: 24575-24584, 1992.

[32512] 4565.

[32513] Brinkman-Van der Linden, E. C. M.; Mollicone, R.; Oriol,

R.; Larson,
[32514] G.; Van den Eijnden, D. H.; Van Dijk, W.: A missense mutation in
[32515] the FUT6 gene results in total absence of alpha-3-fucosylation of
[32516] human alpha-1-acid glycoprotein. J. Biol. Chem. 271: 14492-14495,
[32517] 1996.

[32518] 4566.

[32519] Herbschleb-Voogt, E.; Monteba-van Heuvel, M.; Wijnen, L. M. M.;
[32520] Westerveld, A.; Pearson, P. L.; Meera Khan, P.: Chromosomal assignment
[32521] and regional localization of CS, ENO-2, GAPDH, LDH-B, PEPB, and TPI
[32522] in man-rodent cell hybrids. Cytogenet. Cell Genet. 22: 482-486,
[32523] 1978.

[32524] 4567.

[32525] Brook, J. D.; Shaw, D. J.; Meredith, A. L.; Worwood, M.; Cowell,
[32526] J.; Scott, J.; Knott, T. J.; Litt, M.; Bufton, L.; Harper, P. S.:

[32527] A somatic cell hybrid panel for chromosome 19: localization of known

[32528] genes and RFLPs and orientation of the linkage group. (Abstract) Cytogenet.

[32529] Cell Genet. 40: 590-591, 1985.

[32530] 4568.

[32531] Smith, J. S.; Tachibana, I.; Pohl, U.; Lee, H. K.; Thanarajasingam,

[32532] U.; Portier, B. P.; Ueki, K.; Ramaswamy, S.; Billings, S. J.; Mohrenweiser,

[32533] H. W.; Louis, D. N.; Jenkins, R. B.: A transcript map of the chromosome

[32534] 19q-arm glioma tumor suppressor region. Genomics 64: 44-50, 2000.

[32535] 4569.

[32536] Staal, F. J. T.; van der Luijt, R. B.; Baert, M. R. M.; van Drunen,

[32537] J.; van Bakel, H.; Peters, E.; de Valk, I.; van Amstel, H. K. P.;

[32538] Taphoorn, M. J. B.; Jansen, G. H.; van Veelen, C. W. M.; Burgering,

[32539] B.; Staal, G. E. J.: A novel germline mutation of PTEN associated

[32540] with brain tumours of multiple lineages. Brit. J. Cancer 86: 1586-1591,

[32541] 2002.

[32542] 4570.

[32543] Zhou, X.-P.; Smith, W. M.; Gimm, O.; Mueller, E.; Gao, X.; Sarraf,

[32544] P.; Prior, T. W.; Plass, C.; van Deimling, A.; Black, P. M.; Yates,

[32545] A. J.; Eng, C.: Over-representation of PPAR-gamma se-

quence variants

[32546] in sporadic cases of glioblastoma multiforme: preliminary evidence

[32547] for common low penetrance modifiers for brain tumour risk in the general

[32548] population. J. Med. Genet. 37: 410-414, 2000.

[32549] 4571.

[32550] Bingham, C.; Bulman, M. P.; Ellard, S.; Allen, L. I. S.; Lipkin,

[32551] G. W.; van't Hoff, W. G.; Woolf, A. S.; Rizzoni, G.; Novelli, G.;

[32552] Nicholls, A. J.; Hattersley, A. T.: Mutations in the hepato-

cyte nuclear

[32553] factor-1-beta gene are associated with familial hypoplastic glomerulocystic
[32554] kidney disease. Am. J. Hum. Genet. 68: 219-224, 2001.

[32555] 4572.

[32556] Kaplan, B. S.; Gordon, I.; Pincott, J.; Barratt, T. M.: Familial

[32557] hypoplastic glomerulocystic kidney disease: a definite entity with
[32558] dominant inheritance. Am. J. Med. Genet. 34: 569-573, 1989.

[32559] 4573.

[32560] Rizzoni, G.; Loirat, C.; Levy, M.; Milanesi, C.; Zachello, G.;

[32561] Mathieu, H.: Familial hypoplastic glomerulocystic kidney: a new entity? Clin.
[32562] Nephrol. 18: 263-268, 1982.

[32563] 4574.

[32564] Fong, H. K. W.; Yoshimoto, K. K.; Eversole-Cire, P.; Simon, M.
[32565] I.: Identification of a GTP-binding protein alpha subunit that lacks
[32566] an apparent ADP-ribosylation site for pertussis toxin. Proc. Nat.

[32567] Acad. Sci. 85: 3066-3070, 1988.

[32568] 4575.

[32569] Matsuoka, M.; Itoh, H.; Kozasa, T.; Kaziro, Y.: Sequence analysis

[32570] of cDNA and genomic DNA for a putative pertussis toxin-insensitive

[32571] guanine nucleotide-binding regulatory protein alpha sub-unit. Proc.

[32572] Nat. Acad. Sci. 85: 5384-5388, 1988.

[32573] 4576.

[32574] Wilkie, T. M.; Gilbert, D. J.; Olsen, A. S.; Chen, X.-N.; Ama-truda,

[32575] T. T.; Korenberg, J. R.; Trask, B. J.; de Jong, P.; Reed, R. R.; Simon,

[32576] M. I.; Jenkins, N. A.; Copeland, N. G.: Evolution of the mammalian

[32577] G protein alpha subunit multigene family. Nature Genet. 1: 85-91,

[32578] 1992.

[32579] 4577.

[32580] Magovcevic, I.; Ang, S.-L.; Seidman, J. G.; Tolman, C. J.;

Neer,
[32581] E. J.; Morton, C. C.: Regional localization of the human G protein
[32582] alpha(i2) (GNAI2) gene: assignment to 3p21 and a related sequence
[32583] (GNAI2L) to 12p12-p13. Genomics 12: 125-129, 1992.

[32584] 4578.

[32585] Colombo, M. P.; Martinotti, A.; Howard, T. A.; Schneider, C.; D'Eustachio,
[32586] P.; Seldin, M. F.: Localization of growth arrest-specific genes on
[32587] mouse chromosomes 1, 7, 8, 11, 13, and 16. Mammalian Genome 2: 130-134,
[32588] 1992.

[32589] 4579.

[32590] Del Sal, G.; Collavin, L.; Ruaro, M. E.; Edomi, P.; Saccone, S.;
[32591] Della Valle, G.; Schneider, C.: Structure, function, and chromosome
[32592] mapping of the growth-suppressing human homologue of the murine gas1
[32593] gene. Proc. Nat. Acad. Sci. 91: 1848-1852, 1994.

[32594] 4580.

[32595] Del Sal, G.; Ruaro, M. E.; Philipson, L.; Schneider, C.: The growth

[32596] arrest-specific gene, gas1, is involved in growth suppression. Cell 70:

[32597] 595-607, 1992.

[32598] 4581.

[32599] Evdokiou, A.; Webb, G. C.; Peters, G. B.; Dobrovic, A.; O'Keefe,

[32600] D. S.; Forbes, I. J.; Cowled, P. A.: Localization of the human growth

[32601] arrest-specific gene (GAS1) to chromosome bands 9q21.3-q22, a region

[32602] frequently deleted in myeloid malignancies. Genomics 18: 731-733,

[32603] 1993.

[32604] 4582.

[32605] Schneider, C.; King, R. M.; Philipson, L.: Genes specifically

[32606] expressed at growth arrest of mammalian cells. Cell 54: 787-793,

[32607] 1988.

[32608] 4583.

[32609] Webb, G. C.; Cowled, P. A.; Evdokiou, A.; Ford, J. H.; Forbes,

[32610] I. J.: Assignment, by in situ hybridization, of the growth arrest-specific

[32611] gene, Gas-1, to mouse chromosome 13, bands B3-C2. Genomics 14: 548-549,

[32612] 1992.

[32613] 4584.

[32614] Lebo, R. V.; Cheung, M.-C.; Bruce, B. D.: Rapid gene map-ping by

[32615] dual laser chromosome sorting and spot blot DNA analy-sis. (Abstract) Am.

[32616] J. Hum. Genet. 36: 101S only, 1984.

[32617] 4585.

[32618] Kudo, J.; Chao, L.-Y.; Narni, F.; Saunders, G. F.: Structure of

[32619] the human gene encoding the invariant gamma-chain of class II histocompatibility

[32620] antigens. Nucleic Acids Res. 13: 8827-8841, 1985.

[32621] 4586.

[32622] O'Sullivan, D. M.; Larhammar, D.; Wilson, M. C.; Peterson, P. A.;

[32623] Quaranta, V.: Structure of the human Ia-associated invariant (gamma)-chain

[32624] gene: identification of 5-prime sequences shared with major histocompatibility

[32625] complex class II genes. Proc. Nat. Acad. Sci. 83: 4484-4488, 1986.

[32626] 4587.

[32627] Riese, R. J.; Shi, G.-P.; Villadangos, J.; Stetson, D.; Driessen,

[32628] C.; Lennon-Dumenil, A.-M.; Chu, C.-L.; Naumov, Y.; Be-har, S. M.; Ploegh,

[32629] H.; Locksley, R.; Chapman, H. A.: Regulation of CD1 func-tion and

[32630] NK1.1+ T cell selection and maturation by cathepsin S. Immunity 15:

[32631] 909-919, 2001.

[32632] 4588.

[32633] Topilski, I.; Harmelin, A.; Flavell, R. A.; Levo, Y.; Shachar,

[32634] I.: Preferential Th1 immune response in invariant chain-deficient

[32635] mice. J. Immun. 168: 1610-1617, 2002.

[32636] 4589.

[32637] Abderrahim, H.; Sambucy, J.-L.; Iris, F.; Ougen, P.; Billault,

[32638] A.; Chumakov, I. M.; Dausset, J.; Cohen, D.; Le Paslier, D.: Cloning
[32639] the human major histocompatibility complex in YACs. Genomics 23:
[32640] 520-527, 1994.

[32641] 4590.

[32642] Adman, R.; Pious, D. A.: Isoantigenic variants: isolation from
[32643] human diploid cells in culture. Science 168: 370-372, 1970.
[32644] 4591.

[32645] Arnaiz-Villena, A.; Festenstein, H.: HLA genotyping by using spermatozoa:
[32646] evidence for haploid gene expression. Lancet II: 707-709, 1976.
[32647] 4592.

[32648] Auffray, C.; Strominger, J. L.: Molecular genetics of the human
[32649] major histocompatibility complex. Adv. Hum. Genet. 15: 197-247,
[32650] 1986.

[32651] 4593.

[32652] Awdeh, Z. L.; Raum, D.; Yunis, E. J.; Alper, C. A.: Extended HLA/complement

[32653] allele haplotypes: evidence for T/t-like complex in man. Proc. Nat.

[32654] Acad. Sci. 80: 259-263, 1983.

[32655] 4594.

[32656] Bach, F. H.; Amos, D. B.: Hu-1 major histocompatibility locus

[32657] in man. Science 156: 1506-1508, 1967.

[32658] 4595.

[32659] Bach, M. L.; Bach, F. H.: The genetics of histocompatibility. Hosp.

[32660] Practice 5(8): 33-44, 1970.

[32661] 4596.

[32662] Balner, H.: The major histocompatibility complex of pri-mates:

[32663] evolutionary aspects and comparative histogenetics. Phil. Trans.

[32664] Roy. Soc. London B 292: 109-119, 1981.

[32665] 4597.

[32666] Balner, H.; D'Amaro, J.; Visser, T. P.: Tissue typing of chimpanzees:

[32667] I. Evidence for two allelic series of leukocyte antigens. Transplant.

[32668] Proc. 6: 141-149, 1974.

[32669] 4598.

[32670] Berger, R.; Bernheim, A.; Sasportes, M.; Hauptmann, G.; Hors,

[32671] J.; Legrand, L.; Fellous, M.: Regional mapping of the HLA on the

[32672] short arm of chromosome 6. Clin. Genet. 15: 245-251, 1979.

[32673] 4599.

[32674] Bernard, J.: La decouverte du systeme principal d'histocompatibilite

[32675] de l'homme. (Editorial) Presse Med. 75: 2369 only, 1967.

[32676] 4600.

[32677] Bodmer, W. F.: Personal Communication. London, England 9/26/1986.

[32678] 4601.

[32679] Bodmer, W. F.; Bodmer, J. G.; Adler, S.; Payne, R.; Bialek, J.

[32680] : Genetics of '4' and 'LA' human leukocyte groups. Ann. N.Y. Acad.

[32681] Sci. 129: 473–489, 1966.

[32682] 4602.

[32683] Bodmer, W. F.; Bodmer, J. G.; Tripp, M.: Recombination between

[32684] the LA and 4 loci of the HL-A system.In: Histocompatibil-ity Testing

[32685] 1970. Copenhagen: Munksgaard (pub.) 1970. Pp. 187–191.

[32686] 4603.

[32687] Bodmer, W. F. (ed.): Evolutionary significance of the HL-A system. Nature 237:

[32688] 139–145, 1972.

[32689] 4604.

[32690] Bodmer, W. F. (ed.): The HLA System (17 papers). Brit. Med.

[32691] Bull. 34(3): 213–316, 1978.

[32692] 4605.

[32693] Borgaonkar, D. S.; Bias, W. B.: HL-A loci and chromosome 6. Birth

[32694] Defects Orig. Art. Ser. X(3): 67-68, 1974.

[32695] 4606.

[32696] Borgaonkar, D. S.; Bias, W. B.; Chase, G. A.; Sadasivan, G.; Herr,

[32697] H. M.; Golomb, H. M.; Bahr, G. F.; Kunkel, L. M.: Identification

[32698] of C6-G21 translocation chromosome by the Q-M and Giemsa banding techniques

[32699] in a patient with Down's syndrome, with possible assignment of Gm

[32700] locus. Clin. Genet. 4: 53-57, 1973.

[32701] 4607.

[32702] Breuning, M. H.; van den Berg-Loonen, E. M.; Bernini, L. F.; Bijlsma,

[32703] J. B.; van Loghem, E.; Meera Khan, P.; Nijenhuis, L. E.: Localization

[32704] of HLA on the short arm of chromosome 6. Hum. Genet. 37: 131-139,

[32705] 1977.

[32706] 4608.

[32707] Bull, R. W.; Benson, J. W.; Pearson, G.; Mann, J. D.: HLA substantiation

[32708] of a trisomic human chromosome 6. Transplant. Proc. 10: 747-748,

[32709] 1978.

[32710] 4609.

[32711] Campbell, R. D.; Trowsdale, J.: Map of the human MHC. Immun.

[32712] Today 14: 349-352, 1993.

[32713] 4610.

[32714] Cardoso, C. S.; Alves, H.; Mascarenhas, M.; Goncalves, R.; Oliveira,

[32715] P.; Rodrigues, P.; Cruz, E.; de Sousa, M.; Porto, G.: Co-selection

[32716] of the H63D mutation and the HLA-A29 allele: a new paradigm of linkage

[32717] disequilibrium? Immunogenetics 53: 1002-1008, 2002.

[32718] 4611.

[32719] Carrington, M.; Nelson, G. W.; Martin, M. P.; Kissner, T.; Vlahov,

[32720] D.; Goedert, J. J.; Kaslow, R.; Buchbinder, S.; Hoots, K.; O'Brien,

[32721] S. J.: HLA and HIV-1: heterozygote advantage and B*35-Cw*04 disadvantage. Science 283:

[32722] 1748-1752, 1999.

[32723] 4612.

[32724] Ceppellini, R.; Van Rood, J. J.: The HL-A system. I. Genet-ics

[32725] and molecular biology. Seminars Hemat. 11: 233-252, 1974.

[32726] 4613.

[32727] Cohen, D.; Paul, P.; Font, M.-P.; Cohen, O.; Sayagh, B.; Marcadet,

[32728] A.; Busson, M.; Mahouy, G.; Cann, H. M.; Dausset, J.: Anal-ysis of

[32729] HLA class I genes with restriction endonuclease frag-ments: implications

[32730] for polymorphism of the human major histocompatibility complex. Proc.

[32731] Nat. Acad. Sci. 80: 6289-6292, 1983.

[32732] 4614.

[32733] Colombani, J.: Changing the name of the major histocompatibility
[32734] complex. Res. Immun. 143: 411-417, 1992.

[32735] 4615.

[32736] Dausset, J.: The major histocompatibility complex in man: past,
[32737] present, and future concepts. Science 213: 1469-1474, 1981.
[32738] 4616.

[32739] Dausset, J.: Personal Communication. Paris, France 1/12/1983.
[32740] 4617.

[32741] Dausset, J.: Similarities between the HL-A system and other immunogenetic
[32742] systems. (Editorial) Vox Sang. 23: 153-164, 1972.

[32743] 4618.

[32744] Asakawa, J.; Satoh, C.; Takahashi, N.; Fujita, M.; Kaneko, J.;
[32745] Goriki, K.; Hazama, R.; Kageoka, T.: Electrophoretic variants of
[32746] blood proteins in Japanese: III. Triosephosphate iso-

merase. Hum.
[32747] Genet. 68: 185-188, 1984.

[32748] 4619.

[32749] Ationu, A.; Humphries, A.; Wild, B.; Carr, T.; Will, A.; Arya,

[32750] R.; Layton, D. M.: Towards enzyme-replacement treatment in triosephosphate
[32751] isomerase deficiency. Lancet 353: 1155-1156, 1999.

[32752] 4620.

[32753] Bellingham, A. J.; Lestas, A. N.: Prenatal diagnosis of triose

[32754] phosphate isomerase deficiency. (Letter) Lancet 335: 230, 1990.
[32755] 4621.

[32756] Bellingham, A. J.; Lestas, A. N.; Williams, L. H. P.; Nicolaides,
[32757] K. H.: Prenatal diagnosis of a red-cell enzymopathy: triose phosphate
[32758] isomerase deficiency. Lancet II: 419-421, 1989.

[32759] 4622.

[32760] Boyer, T. G.; Krug, J. R.; Maquat, L. E.: Transcriptional regulatory

[32761] sequences of the housekeeping gene for human triosephosphate isomerase. J.

[32762] Biol. Chem. 264: 5177-5187, 1989.

[32763] 4623.

[32764] Boyer, T. G.; Maquat, L. E.: Minimal sequence and factor requirements

[32765] for the initiation of transcription from an atypical, TATATAA box-containing

[32766] housekeeping promoter. J. Biol. Chem. 265: 20524-20532, 1990.

[32767] 4624.

[32768] Brock, D. J. H.; Singer, J. D.: Red cell triosephosphate isomerase

[32769] gene. (Letter) Lancet II: 1136, 1970.

[32770] 4625.

[32771] Matsushime, H.; Jinno, A.; Takagi, N.; Shibuya, M.: A novel mammalian

[32772] protein kinase gene (mak) is highly expressed in testicular germ cells

[32773] at and after meiosis. Molec. Cell. Biol. 10: 2261-2268, 1990.

[32774] 4626.

[32775] Taketo, M.; Jinno, A.; Yamaguchi, S.; Matushime, H.;
Shibuya, M.;

[32776] Seldin, M. F.: Mouse Mak gene for male germ cell-
associated kinase

[32777] maps to chromosome 13. Genomics 19: 397-398, 1994.

[32778] 4627.

[32779] Jiang, C.; Atkinson, D.; Towbin, J. A.; Splawski, I.;
Lehmann,

[32780] M. H.; Li, H.; Timothy, K.; Taggart, R. T.; Schwartz, P. J.;
Vincent,

[32781] G. M.; Moss, A. J.; Keating, M. T.: Two long QT syndrome
loci map

[32782] to chromosomes 3 and 7 with evidence for further hetero-
geneity. Nature

[32783] Genet. 8: 141-147, 1994.

[32784] 4628.

[32785] Jongbloed, R. J. E.; Wilde, A. A. M.; Geelen, J. L. M. C.; Do-
evendans,

[32786] P.; Schaap, C.; Van Langen, I.; van Tintelen, J. P.; Cobben,
J. M.;

[32787] Beaufort-Krol, G. C. M.; Geraedts, J. P. M.; Smeets, H. J. M.: Novel

[32788] KCNQ1 and HERG missense mutations in Dutch long-QT families. Hum.

[32789] Mutat. 13: 301-310, 1999.

[32790] 4629.

[32791] Priori, S. G.; Barhanin, J.; Hauer, R. N. W.; Haverkamp, W.; Jongsma,

[32792] H. J.; Kleber, A. G.; McKenna, W. J.; Roden, D. M.; Rudy, Y.; Schwartz,

[32793] K.; Schwartz, P. J.; Towbin, J. A.; Wilde, A. M.: Genetic and molecular

[32794] basis of cardiac arrhythmias: impact on clinical manage- ment. Parts

[32795] I and II. Circulation 99: 518-528, 1999.

[32796] 4630.

[32797] Priori, S. G.; Barhanin, J.; Hauer, R. N. W.; Haverkamp, W.; Jongsma,

[32798] H. J.; Kleber, A. G.; McKenna, W. J.; Roden, D. M.; Rudy, Y.; Schwartz,

[32799] K.; Schwartz, P. J.; Towbin, J. A.; Wilde, A. M.: Genetic and molecular

[32800] basis of cardiac arrhythmias: impact on clinical manage-
ment. Part
[32801] III. Circulation 99: 674-681, 1999.

[32802] 4631.

[32803] Priori, S. G.; Napolitano, C.; Schwartz, P. J.: Low pene-
trance
[32804] in the long QT syndrome: clinical impact. Circulation 99:
529-533,
[32805] 1999.

[32806] 4632.

[32807] Splawski, I.; Shen, J.; Timothy, K. W.; Vincent, G. M.;
Lehmann,
[32808] M. H.; Keating, M. T.: Genomic structure of three long QT
syndrome
[32809] genes: KVLQT1, HERG, and KCNE1. Genomics 51: 86-97,
1998.
[32810] 4633.

[32811] Tanaka, T.; Nagai, R.; Tomoike, H.; Takata, S.; Yano, K.;
Yabuta,
[32812] K.; Haneda, N. Nakano, O.; Shibata, A.; Sawayama, T.; Ka-
sai, H.; Yazaki,

[32813] Y.; Nakamura, Y.: Four novel KVLQT1 and four novel HERG mutations
[32814] in familial long-QT syndrome. Circulation 95: 565-567, 1997.
[32815] 4634.
[32816] Zareba, W.; Moss, A. J.; Schwartz, P. J.; Vincent, G. M.; Robinson,
[32817] J. L.; Priori, S. G.; Benhorin, J.; Locati, E. H.; Towbin, J. A.;
[32818] Keating, M. T.; Lehmann, M. H.; Hall, W. J.; International Long-QT
[32819] Syndrome Registry Research Group: Influence of the genotype on the
[32820] clinical course of the long-QT syndrome. New Eng. J. Med. 339: 960-965,
[32821] 1998.
[32822] 4635.
[32823] Caffey, J. P.; Christensen, W. R.: Personal Communication. Pittsburgh,
[32824] Pa. and Salt Lake City, Utah 1963.
[32825] 4636.
[32826] Jacenko, O.; Lu Valle, P. A.; Olsen, B. R.: Spondylometa-

physeal
[32827] dysplasia in mice carrying a dominant negative mutation in a matrix
[32828] protein specific for cartilage-to-bone transition. Nature 365: 56-61,
[32829] 1993.

[32830] 4637.

[32831] Stephens, F. E.: An achondroplastic mutation and the nature of
[32832] its inheritance. J. Hered. 34: 229-235, 1943.

[32833] 4638.

[32834] Wallis, G. A.; Rash, B.; Sykes, B.; Bonaventure, J.; Maroteaux,
[32835] P.; Zabel, B.; Wynne-Davies, R.; Grant, M. E.; Boot-Handford, R. P.
[32836] : Mutations within the gene encoding the alpha-1(X) chain of type
[32837] X collagen (COL1A1) cause metaphyseal chondrodysplasia type Schmid
[32838] but not several other forms of metaphyseal chondrodysplasia. J. Med.
[32839] Genet. 33: 450-457, 1996.

[32840] 4639.

[32841] Warman, M. L.; Abbott, M.; Apte, S. S.; Hefferon, T.; McIn-
tosh,

[32842] I.; Cohn, D. H.; Hecht, J. T.; Olsen, B. R.; Francomano, C.
A.: A

[32843] type X collagen mutation causes Schmid metaphyseal
chondrodysplasia. Nature

[32844] Genet. 5: 79-82, 1993.

[32845] 4640.

[32846] Boyes, J.; Bird, A.: DNA methylation inhibits transcription
indirectly

[32847] via a methyl-CpG binding protein. Cell 64: 1123-1134,
1991.

[32848] 4641.

[32849] Cross, S. H.; Meehan, R. R.; Nan, X.; Bird, A.: A component
of

[32850] the transcriptional repressor MeCP1 shares a motif with
DNA methyltransferase

[32851] and HRX proteins. Nature Genet. 16: 256-259, 1997.

[32852] 4642.

[32853] Hendrich, B.; Abbott, C.; McQueen, H.; Chambers, D.;

Cross, S.;

[32854] Bird, A.: Genomic structure and chromosomal mapping of the murine

[32855] and human Mbd1, Mbd2, Mbd3, and Mbd4 genes. Mammalian Genome 10:

[32856] 906-912, 1999.

[32857] 4643.

[32858] Jobsis, G. J.; Bolhuis, P. A.; Boers, J. M.; Baas, F.; Wolterman,

[32859] R. A.; Hensels, G. W.; de Visser, M.: Genetic localization of Bethlem

[32860] myopathy. Neurology 46: 779-782, 1996.

[32861] 4644.

[32862] Jobsis, G. J.; Keizers, H.; Vreijling, J. P.; de Visser, M.; Speer,

[32863] M. C.; Wolterman, R. A.; Baas, F.; Bohlhuis, P. A.: Type VI collagen

[32864] mutations in Bethlem myopathy, an autosomal dominant myopathy with

[32865] contractures. Nature Genet. 14: 113-115, 1996.

[32866] 4645.

[32867] Hempstead, B. L.; Martin-Zanca, D.; Kaplan, D. R.; Parada, L.
[32868] F.; Chao, M. V.: High-affinity NGF binding requires coexpression
[32869] of the trk proto-oncogene and the low-affinity NGF receptor. Nature 350:
[32870] 678-683, 1991.

[32871] 4646.

[32872] Huebner, K.; Isobe, M.; Chao, M.; Bothwell, M.; Ross, A. H.; Finan,
[32873] J.; Hoxie, J. A.; Sehgal, A.; Buck, C. R.; Lanahan, A.; Nowell, P.
[32874] C.; Koprowski, H.; Croce, C. M.: The nerve growth factor receptor
[32875] gene is at human chromosome region 17q12-17q22, distal to the chromosome
[32876] 17 breakpoint in acute leukemias. Proc. Nat. Acad. Sci. 83: 1403-1407,
[32877] 1986.

[32878] 4647.

[32879] Ip, N. Y.; Stitt, T. N.; Tapley, P.; Klein, R.; Glass, D. J.;

[32880] Fandl, J.; Greene, L. A.; Barbacid, M.; Yancopoulos, G. D.: Similarities

[32881] and differences in the way neurotrophins interact with the Trk receptors

[32882] in neuronal and nonneuronal cells. Neuron 10: 137-149, 1993.

[32883] 4648.

[32884] Johnson, D.; Lanahan, A.; Buck, C. R.; Sehgal, A.; Morgan, C.;

[32885] Mercer, E.; Bothwell, M.; Chao, M.: Expression and structure of the

[32886] human NGF receptor. Cell 47: 545-554, 1986.

[32887] 4649.

[32888] Lee, K. F.; Li, E.; Huber, J.; Landis, S. C.; Sharpe, A. H.; Chao,

[32889] M. V.; Jaenisch, R.: Targeted mutation of the gene encoding the low

[32890] affinity NGF receptor p75 leads to deficits in the peripheral sensory

[32891] nervous system. Cell 69: 737-749, 1992.

[32892] 4650.

[32893] Mischel, P. S.; Smith, S. G.; Vining, E. R.; Valletta, J. S.;

[32894] Mobley, W. C.; Reichardt, L. F.: The extracellular domain of p75(NTR)

[32895] is necessary to inhibit neurotrophin-3 signaling through TrkA. J.

[32896] Biol. Chem. 276: 11294-11301, 2001.

[32897] 4651.

[32898] Rettig, W. J.; Thomson, T. M.; Spengler, B. A.; Biedler, J. L.;

[32899] Old, L. J.: Assignment of human nerve growth factor receptor gene

[32900] to chromosome 17 and regulation of receptor expression in somatic

[32901] cell hybrids. Somat. Cell Molec. Genet. 12: 441-447, 1986.

[32902] 4652.

[32903] Tuffereau, C.; Benejean, J.; Blondel, D.; Kieffer, B.; Flamand,

[32904] A.: Low-affinity nerve-growth factor receptor (p75NTR) can serve

[32905] as a receptor for rabies virus. EMBO J. 17: 7250-7259, 1998.

[32906] 4653.

[32907] Welcher, A. A.; Bitler, C. M.; Radeke, M. J.; Shooter, E. M.:

[32908] Nerve growth factor binding domain of the nerve growth factor receptor. Proc.

[32909] Nat. Acad. Sci. 88: 159-163, 1991.

[32910] 4654.

[32911] Chuang, H.; Prescott, E. D.; Kong, H.; Shields, S.; Jordt, S.-E.;

[32912] Basbaum, A. I.; Chao, M. V.; Julius, D.: Bradykinin and nerve growth

[32913] factor release the capsaicin receptor from Pt-dIns(4,5)P2-mediated

[32914] inhibition. Nature 411: 957-962, 2001.

[32915] 4655.

[32916] Darby, J. K.; Feder, J.; Selby, M.; Riccardi, V.; Ferrell, R.;

[32917] Siao, D.; Goslin, K.; Rutter, W.; Shooter, E. M.; Cavalli-Sforza,

[32918] L. L.: A discordant sibship analysis between beta-NGF and neurofibromatosis. Am.

[32919] J. Hum. Genet. 37: 52-59, 1985.

[32920] 4656.

[32921] Mitchell, E. L. D.; Jones, D.; White, G. R. M.; Varley, J. M.;

[32922] Santibanez Koref, M. F.: Determination of the gene order of the three

[32923] loci CD2, NGFB, and NRAS at human chromosome band 1p13 and refinement

[32924] of their localisation at the subband level by fluorescence in situ

[32925] hybridization. Cytogenet. Cell Genet. 70: 183-185, 1995.

[32926] 4657.

[32927] Beauchemin, N.; Draber, P.; Dveksler, G.; Gold, P.; Gray-Owen,

[32928] S.; Grunert, F.; Hammarstrom, S.; Holmes, K. V.; Karlsson, A.; Kuroki,

[32929] M.; Lin, S.-H.; Lucka, L.; and 13 others: Redefined nomen-clature

[32930] for members of the carcinoembryonic antigen family. Exp. Cell Res. 252:

[32931] 243-249, 1999.

[32932] 4658.

[32933] Willcocks, T. C.; Craig, S. P.; Craig, I. W.: Assignment of

the

[32934] coding sequence for carcinoembryonic antigen (CEA) and normal cross-reacting

[32935] antigen (NCA) to human chromosome 19q13. Ann. Hum. Genet. 53: 141-148,

[32936] 1989.

[32937] 4659.

[32938] Qian, F.; Kruse, U.; Lichter, P.; Sippel, A. E.: Chromosomal localization

[32939] of the four genes (NFIA, B, C, and X) for the human tran-scription

[32940] factor nuclear factor I by FISH. Genomics 28: 66-73, 1995.

[32941] 4660.

[32942] Auphan, N.; DiDonato, J.A.; Rosette, C.; Helmberg, A.; Karin, M.

[32943] : Immunosuppression by glucocorticoids: inhibition of NF-kappa-B activity

[32944] through induction of I-kappa-B synthesis. Science 270: 286-290,

[32945] 1995.

[32946] 4661.

[32947] Baeuerle, P. A.: I-kappa-B--NF-kappa-B structures: at the interface
[32948] of inflammation control. Cell 95: 729-731, 1998.

[32949] 4662.

[32950] Haskill, S.; Beg, A. A.; Tompkins, S. M.; Morris, J. S.; Yurochko,
[32951] A. D.; Sampson-Johannes, A.; Mondal, K.; Ralph, P.; Baldwin, A. S.,
[32952] Jr.: Characterization of an immediate-early gene induced in adherent
[32953] monocytes that encodes I-kappa-B-like activity. Cell 65: 1281-1289,
[32954] 1991.

[32955] 4663.

[32956] Huxford, T.; Huang, D.-B.; Malek, S.; Ghosh, G.: The crystal structure
[32957] of the I-kappa-B-alpha/NF-kappa-B complex reveals mechanisms of NF-kappa-B
[32958] inactivation. Cell 95: 759-770, 1998.

[32959] 4664.

[32960] Ito, C. Y.; Adey, N.; Bautch, V. L.; Baldwin, A. S., Jr.: Struc-

ture

[32961] and evolution of the human IKBA gene. Genomics 29: 490-495, 1995.

[32962] 4665.

[32963] Jacobs, M. D.; Harrison, S. C.: Structure of an I-kappa-B-alpha/NF-kappa-B

[32964] complex. Cell 95: 749-758, 1998.

[32965] 4666.

[32966] Jung, M.; Zhang, Y.; Lee, S.; Dritschilo, A.: Correction of radiation

[32967] sensitivity in ataxia telangiectasia cells by a truncated I-kappa-B-alpha. Science 268:

[32968] 1619-1621, 1995.

[32969] 4667.

[32970] Le Beau, M. M.; Ito, C.; Cogswell, P.; Espinosa, R., III; Fer-nald,

[32971] A. A.; Baldwin, A. S., Jr.: Chromosomal localization of the genes

[32972] encoding the p50/p105 subunits of NF-kappa-B (NFKB2) and the I-kappa-B/MAD-3

[32973] (NFKBI) inhibitor of NF-kappa-B to 4q24 and 14q13, re-

spectively. Genomics 14:
[32974] 529-531, 1992.

[32975] 4668.

[32976] Neish, A. S..; Gewirtz, A. T.; Zeng, H.; Young, A. N.; Hobert,
[32977] M. E.; Karmali, V.; Rao, A. S.; Madara, J. L.: Prokaryotic regulation
[32978] of epithelial responses by inhibition of I-kappa-B-alpha ubiquitination. Science 289:
[32979] 1560-1563, 2000.

[32980] 4669.

[32981] Rupec, R. A.; Poujol, D.; Grosgeorge, J.; Carle, G. F.; Livolsi,
[32982] A.; Peyron, J.-F.; Schmid, R. M.; Baeuerle, P. A.; Messer, G.: Structural
[32983] analysis, expression, and chromosomal localization of the mouse ikba
[32984] gene. Immunogenetics 49: 395-403, 1999.

[32985] 4670.

[32986] Scheinman, R. I.; Cogswell, P. C.; Lofquist, A. K.; Baldwin, A.

[32987] S., Jr.: Role of transcriptional activation of I-
kappa-B-alpha in
[32988] mediation of immunosuppression by glucocorticoids. Sci-
ence 270:
[32989] 283-286, 1995.

[32990] 4671.

[32991] Compton, D. A.; Szilak, I.; Cleveland, D. W.: Primary struc-
ture
[32992] of NuMA, an intranuclear protein that defines a novel
pathway for
[32993] segregation of proteins at mitosis. J. Cell Biol. 116:
1395-1408,
[32994] 1992.

[32995] 4672.

[32996] Lydersen, B. K.; Pettijohn, D. E.: Human-specific nuclear
protein
[32997] that associates with the polar region of the mitotic appa-
ratus: distribution
[32998] in a human/hamster hybrid cell. Cell 22: 489-499, 1980.

[32999] 4673.

[33000] Merdes, A.; Ramyar, K.; Vechio, J. D.; Cleveland, D. W.: A

complex
[33001] of NuMA and cytoplasmic dynein is essential for mitotic spindle assembly. Cell 87:
[33002] 447-458, 1996.

[33003] 4674.

[33004] Sparks, C. A.; Bangs, P. L.; McNeil, G. P.; Lawrence, J. B.; Fey,
[33005] E. G.: Assignment of the nuclear mitotic apparatus protein NuMA gene
[33006] to human chromosome 11q13. Genomics 17: 222-224, 1993.
[33007] 4675.

[33008] Wells, R. A.; Catzavelos, C.; Kamel-Reid, S.: Fusion of retinoic
[33009] acid receptor alpha to NuMA, the nuclear mitotic appara-tus protein,
[33010] by a variant translocation in acute promyelocytic leukaemia. Nature
[33011] Genet. 17: 109-113, 1997.

[33012] 4676.

[33013] Wiese, C.; Wilde, A.; Moore, M. S.; Adam, S. A.; Merdes, A.;

Zheng,
[33014] Y.: Role of importin-beta in coupling Ran to downstream targets in
[33015] microtubule assembly. Science 291: 653-656, 2001.

[33016] 4677.

[33017] Yang, C. H.; Lambie, E. J.; Snyder, M.: NuMA: an unusually long
[33018] coiled-coil related protein in the mammalian nucleus. J. Cell Biol. 116:
[33019] 1303-1317, 1992.

[33020] 4678.

[33021] Halila, R.; Apostolou, S.; Winqvist, R.; Callen, D.; Prockop, D.
[33022] J.; Peltonen, L.: Isolation and genomic assignment of a candidate
[33023] cDNA clone for type III procollagen N-proteinase. (Abstract) Am.
[33024] J. Hum. Genet. 51 (suppl.): A128 only, 1992.

[33025] 4679.

[33026] Halila, R.; Peltonen, L.; Prockop, D. J.: Isolation of a candi-
date

[33027] cDNA clone for type III procollagen N-proteinase from human placental

[33028] cDNA library. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A192 only,

[33029] 1989.

[33030] 4680.

[33031] Nomura, N.; Nagase, T.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Sato,

[33032] S.; Seki, N.; Kawarabayasi, Y.; Ishikawa, K.; Tabata, S.: Prediction

[33033] of the coding sequences of unidentified human genes. II. The coding

[33034] sequences of 40 new genes (KIAA0041-KIAA0080) deduced by analysis

[33035] of cDNA clones from human cell line KG-1. DNA Res. 1: 223-229, 1994.

[33036] 4681.

[33037] Freedman, D. A.; Epstein, C. B.; Roth, J. C.; Levine, A. J.: A

[33038] genetic approach to mapping the p53 binding site in the MDM2 protein. Molec.

[33039] Med. 3: 248-259, 1997.

[33040] 4682.

[33041] Fuchs, S. Y.; Adler, V.; Buschmann, T.; Yin, Z.; Wu, X.; Jones,

[33042] S. N.; Ronai, Z.: JNK targets p53 ubiquitination and degra-dation

[33043] in nonstressed cells. Genes Dev. 12: 2658-2663, 1998.

[33044] 4683.

[33045] Mayo, L. D.; Donner, D. B.: A phosphatidylinositol 3-kinase/Akt

[33046] pathway promotes translocation of Mdm2 from the cyto-plasm to the nucleus. Proc.

[33047] Nat. Acad. Sci. 98: 11598-11603, 2001.

[33048] 4684.

[33049] Mitchell, E. L. D.; White, G. R. M.; Santibanez-Koref, M. F.;

[33050] Varley, J. M.; Heighway, J.: Mapping of gene loci in the q13-q15

[33051] region of chromosome 12. Chromosome Res. 3: 261-262, 1995.

[33052] 4685.

[33053] Momand, J.; Zambetti, G. P.; Olson, D. C.; George, D. L.; Levine,

[33054] A. J.: The mdm-2 oncogene product forms a complex with the p53 protein
[33055] and inhibits p53-mediated transactivation. Cell 69: 1237-1245, 1992.
[33056] 4686.

[33057] Oliner, J. D.; Kinzler, K. W.; Meltzer, P. S.; George, D. L.;

[33058] Vogelstein, B.: Amplification of a gene encoding a p53-associated
[33059] protein in human sarcomas. Nature 358: 80-83, 1992.

[33060] 4687.

[33061] Ries, S.; Biederer, C.; Woods, D.; Shifman, O.; Shirasawa, S.;
[33062] Sasazuki, T.; McMahon, M.; Oren, M.; McCormick, F.: Op-posing effects
[33063] of Ras on p53: transcriptional activation of mdm2 and in-duction of
[33064] p19(ARF). Cell 103: 321-330, 2000.

[33065] 4688.

[33066] Shenoy, S. K.; McDonald, P. H.; Kohout, T. A.; Lefkowitz, R. J.
[33067] : Regulation of receptor fate by ubiquitination of activated

beta-2-adrenergic
[33068] receptor and beta-arrestin. Science 294: 1307-1313,
2001.
[33069] 4689.

[33070] Shieh, S.-Y.; Ikeda, M.; Taya, Y.; Prives, C.: DNA damage-
induced
[33071] phosphorylation of p53 alleviates inhibition by MDM2. Cell
91: 325-334,
[33072] 1997.

[33073] 4690.

[33074] Testa, J. R.; Bellacosa, A.: AKT plays a central role in tu-
morigenesis. Proc.
[33075] Nat. Acad. Sci. 98: 10983-10985, 2001.

[33076] 4691.

[33077] Xiao, Z.-X.; Chen, J.; Levine, A. J.; Modjtahedi, N.; Xing, J.;

[33078] Sellers, W. R.; Livingston, D. M.: Interaction between the
retinoblastoma
[33079] protein and the oncoprotein MDM2. Nature 375: 694-698,
1995.
[33080] 4692.

[33081] Zhang, Y.; Xiong, Y.: A p53 amino-terminal nuclear export

signal

[33082] inhibited by DNA damage-induced phosphorylation. Science 292: 1910-1915,

[33083] 2001.

[33084] 4693.

[33085] Ashkar, S.; Weber, G. F.; Panoutsakopoulou, V.; Sanchirico, M.

[33086] E.; Jansson, M.; Zawaideh, S.; Rittling, S. R.; Denhardt, D. T.; Glimchar,

[33087] M. J.; Cantor, H.: Eta-1 (osteopontin): an early component of type-1

[33088] (cell-mediated) immunity. Science 287: 860-864, 2000.

[33089] 4694.

[33090] Baccarini-Contri, M.; Taparelli, F.; Pasquali-Ronchetti, I.: Osteopontin

[33091] is a constitutive component of normal elastic fibers in human skin

[33092] and aorta. Matrix Biol. 14: 553-560, 1994.

[33093] 4695.

[33094] Beck, G. R., Jr.; Zerler, B.; Moran, E.: Phosphate is a specific

[33095] signal for induction of osteopontin gene expression. Proc. Nat. Acad.

[33096] Sci. 97: 8352-8357, 2000.

[33097] 4696.

[33098] Crosby, A. H.; Lyu, M. S.; Lin, K.; McBride, O. W.; Kerr, J. M.;

[33099] Aplin, H. M.; Fisher, L. W.; Young, M. F.; Kozak, C. A.; Dixon, M.

[33100] J.: Mapping of the human and mouse bone sialoprotein and osteopontin

[33101] loci. Mammalian Genome 7: 149-151, 1996.

[33102] 4697.

[33103] Fisher, L. W.; McBride, O. W.; Termine, J. D.; Young, M. F.: Human

[33104] bone sialoprotein: deduced protein sequence and chromosomal localization. J.

[33105] Biol. Chem. 265: 2347-2351, 1990.

[33106] 4698.

[33107] Graf, K.; Do, Y. S.; Ashizawa, N.; Meehan, W. P.; Giachelli, C.

[33108] M.; Marboe, C. C.; Fleck, E.; Hsueh, W. A.: Myocardial os-

teopontin
[33109] expression is associated with left ventricular hypertrophy. Circulation 96:
[33110] 3063-3071, 1997.

[33111] 4699.

[33112] Kiefer, M. C.; Bauer, D. M.; Barr, P. J.: The cDNA and de-
rived
[33113] amino acid sequence for human osteopontin. Nucleic Acids Res. 17:
[33114] 3306, 1989.

[33115] 4700.

[33116] Kim, J.-H.; Skates, S. J.; Uede, T.; Wong, K.; Schorge, J. O.;

[33117] Feltmate, C. M.; Berkowitz, R. S.; Cramer, D. W.; Mok, S. C.: Osteopontin
[33118] as a potential diagnostic biomarker for ovarian cancer. J.A.M.A. 287:
[33119] 1671-1679, 2002.

[33120] 4701.

[33121] Kohri, K.; Nomura, S.; Kitamura, Y.; Nagata, T.; Yoshioka, K.;
[33122] Iguchi, M.; Yamate, T.; Umekawa, T.; Suzuki, Y.; Sinohara,

H.; Kurita,

[33123] T.: Structure and expression of the mRNA encoding urinary stone protein

[33124] (osteopontin). J. Biol. Chem. 268: 15180-15184, 1993.

[33125] 4702.

[33126] Kohri, K.; Suzuki, Y.; Yoshida, K.; Yamamoto, K.; Amasaki, N.;

[33127] Yamate, T.; Umekawa, T.; Iguchi, M.; Sinohara, H.; Kurita, T.: Molecular

[33128] cloning and sequencing of cDNA encoding urinary stone protein, which

[33129] is identical to osteopontin. Biochem. Biophys. Res. Commun. 184:

[33130] 859-864, 1992.

[33131] 4703.

[33132] Liaw, L.; Birk, D. E.; Ballas, C. B.; Whitsitt, J. S.; Davidson,

[33133] J. M.; Hogan, B. L. M.: Altered wound healing in mice lacking a functional

[33134] osteopontin gene (spp1). J. Clin. Invest. 101: 1468-1478, 1998.

[33135] 4704.

[33136] Morimoto, I.; Sasaki, Y.; Ishida, S.; Imai, K.; Tokino, T.: Identification
[33137] of the osteopontin gene as a direct target of TP53. Genes Chromosomes
[33138] Cancer 33: 270-278, 2002.

[33139] 4705.

[33140] Reinholt, F. P.; Hultenby, K.; Oldberg, A.; Heinegard, D.: Osteopontin--a
[33141] possible anchor of osteoclasts to bone. Proc. Nat. Acad. Sci. 87:
[33142] 4473-4475, 1990.

[33143] 4706.

[33144] Singh, K.; Balligand, J.-L.; Fischer, T. A.; Smith, T. W.; Kelly,
[33145] R. A.: Glucocorticoids increase osteopontin expression in cardiac
[33146] myocytes and microvascular endothelial cells: role in reg-ulation of
[33147] inducible nitric oxide synthase. J. Biol. Chem. 270: 28471-28478,
[33148] 1995.

[33149] 4707.

[33150] Weber, G. F.; Ashkar, S.; Glimcher, M. J.; Cantor, H.: Receptor-ligand

[33151] interaction between CD44 and osteopontin (Eta-1). Science 271: 509-512,

[33152] 1996.

[33153] 4708.

[33154] Staal, A.; van Wijnen, A. J.; Birkenhager, J. C.; Pols, H. A.

[33155] P.; Prahl, J.; DeLuca, H.; Gaub, M.-P.; Lian, J. B.; Stein, G. S.;

[33156] van Leeuwen, J. P. T. M.; Stein, J. L.: Distinct conformations of

[33157] vitamin D receptor/retinoid X receptor-alpha heterodimers are specified

[33158] by dinucleotide differences in the vitamin D-responsive elements of

[33159] the osteocalcin and osteopontin genes. Molec. Endocr. 10: 1444-1456,

[33160] 1996.

[33161] 4709.

[33162] Young, M. F.; Kerr, J. M.; Termine, J. D.; Wewer, U. M.;

Wang,
[33163] M. G.; McBride, O. W.; Fisher, L. W.: cDNA cloning, mRNA distribution
[33164] and heterogeneity, chromosomal location and RFLP analysis of human
[33165] osteopontin. Genomics 7: 491-502, 1990.

[33166] 4710.

[33167] Vogelstein, B.; Fearon, E. R.; Hamilton, S. R.; Kern, S. E.; Preisinger,
[33168] A. C.; Leppert, M.; Nakamura, Y.; White, R.; Smits, A. M. M.; Bos,
[33169] J. L.: Genetic alterations during colorectal-tumor development. New
[33170] Eng. J. Med. 319: 525-532, 1988.

[33171] 4711.

[33172] Zhou, X.-P.; Kuismanen, S.; Nystrom-Lahti, M.; Peltomaki, P.;
[33173] Eng, C.: Distinct PTEN mutational spectra in hereditary non-polyposis
[33174] colon cancer syndrome-related endometrial carcinomas compared to sporadic
[33175] microsatellite unstable tumors. Hum. Molec. Genet. 11:

445-450,

[33176] 2002.

[33177] 4712.

[33178] Zhao, C.; Takita, J.; Tanaka, Y.; Setou, M.; Nakagawa, T.; Takeda,

[33179] S.; Yang, H. W.; Terada, S.; Nakata, T.; Takei, Y.; Saito, M.; Tsuji,

[33180] S.; Hayashi, Y.; Hirokawa, N.: Charcot-Marie-Tooth dis-ease type 2A

[33181] caused by mutation in a microtubule motor KIF1B-beta. Cell 105:

[33182] 587-597, 2001.

[33183] 4713.

[33184] Maggi, M.; Barni, T.; Fantoni, G.; Mancina, R.; Pupilli, C.; Luconi,

[33185] M.; Crescioli, C.; Serio, M.; Vannelli, G. B.: Expression and biological

[33186] effects of endothelin-1 in human gonadotropin-releasing hormone-secreting

[33187] neurons. J. Clin. Endocr. Metab. 85: 1658-1665, 2000.

[33188] 4714.

[33189] Napolitano, M.; Miceli, F.; Calce, A.; Vacca, A.; Gulino, A.;

[33190] Apa, R.; Lanzone, A.: Expression and relationship between endothelin-1

[33191] messenger ribonucleic acid (mRNA) and inducible/endothelial nitric

[33192] oxide synthase mRNA isoforms from normal and preeclamptic placentas. J.

[33193] Clin. Endocr. Metab. 85: 2318-2323, 2000.

[33194] 4715.

[33195] Okafor, M. C.; Delamere, N. A.: The inhibitory influence of endothelin

[33196] on active sodium-potassium transport in porcine lens. Invest. Ophthal.

[33197] Vis. Sci. 42: 1018-1023, 2001.

[33198] 4716.

[33199] Zeidel, M. L.; Brady, H. R.; Kone, B. C.; Gullans, S. R.; Brenner,

[33200] B. M.: Endothelin, a peptide inhibitor of Na(+)-K(+)-ATPase in intact

[33201] renal tubular epithelial cells. Am. J. Physiol. 257: C1101-C1107,

[33202] 1989.

[33203] 4717.

[33204] Baynash, A. G.; Hosoda, K.; Giaid, A.; Richardson, J. A.; Emoto,

[33205] N.; Hammer, R. E.; Yanagisawa, M.: Interaction of endothelin-3 with

[33206] endothelin-B receptor is essential for development of epidermal melanocytes

[33207] and enteric neurons. Cell 79: 1277-1285, 1994.

[33208] 4718.

[33209] Bidaud, C.; Salomon, R.; Van Camp, G.; Pelet, A.; Attie, T.; Eng,

[33210] C.; Bonduelle, M.; Amiel, J.; Nihoul-Fekete, C.; Willems, P. J.; Munnich,

[33211] A.; Lyonnet, S.: Endothelin-3 gene mutations in isolated and syndromic

[33212] Hirschsprung disease. Europ. J. Hum. Genet. 5: 247-251, 1997.

[33213] 4719.

[33214] Bloch, K.; Eddy, R. L.; Shows, T. B.; Quertermous, T.: cDNA cloning

[33215] and chromosomal assignment of the gene encoding endothelin 3. J.

[33216] Biol. Chem. 264: 18156-18161, 1989.

[33217] 4720.

[33218] Bolk, S.; Angrist, M.; Xie, J.; Yanagisawa, M.; Silvestri, J. M.;

[33219] Weese-Mayer, D. E.; Chakravarti, A.: Endothelin-3 frameshift mutation

[33220] in congenital central hypoventilation syndrome. (Letter) Nature Genet. 13:

[33221] 395-396, 1996.

[33222] 4721.

[33223] Dupin, E.; Glavieux, C.; Vaigot, P.; Le Douarin, N. M.: Endothelin

[33224] 3 induces the reversion of melanocytes to glia through a neural crest-derived

[33225] glial-melanocytic progenitor. Proc. Nat. Acad. Sci. 97: 7882-7887,

[33226] 2000.

[33227] 4722.

[33228] Edery, P.; Attie, T.; Amiel, J.; Pelet, A.; Eng, C.; Hofstra, R.

[33229] M. W.; Martelli, H.; Bidaud, C.; Munnich, A.; Lyonnet, S.: Mutation
[33230] of the endothelin-3 gene in the Waardenburg-Hirschsprung disease (Shah-Waardenburg
[33231] syndrome). Nature Genet. 12: 442-444, 1996.

[33232] 4723.

[33233] Gopal Rao, V. V. N.; Loffler, C.; Hansmann, I.: The gene for the
[33234] novel vasoactive peptide endothelin 3 (EDN3) is localized to human
[33235] chromosome 20q13.2-qter. Genomics 10: 840-841, 1991.

[33236] 4724.

[33237] Marger, R. S.; Marger, D.: Carcinoma of the esophagus and tylosis:
[33238] a lethal genetic combination. Cancer 72: 17-19, 1993.

[33239] 4725.

[33240] Risk, J. M.; Field, E. A.; Field, J. K.; Whittaker, J.; Fryer,
[33241] A.; Ellis, A.; Shaw, J. M.; Friedmann, P. S.; Bishop, D. T.; Bodmer,
[33242] J.; Leigh, I. M.: Tylosis oesophageal cancer mapped.

(Letter) Nature
[33243] Genet. 8: 319-321, 1994.

[33244] 4726.

[33245] Shine, I.; Allison, P. R.: Carcinoma of the esophagus with tylosis. Lancet I:
[33246] 951-953, 1966.

[33247] 4727.

[33248] Stevens, H. P.; Kelsell, D. P.; Bryant, S. P.; Bishop, D. T.;

[33249] Spurr, N. K.; Weissenbach, J.; Marger, D.; Marger, R. S.; Leigh, I.
[33250] M.: Linkage of an American pedigree with palmoplantar keratoderma
[33251] and malignancy (palmoplantar ectodermal dysplasia type III) to 17q24:
[33252] literature survey and proposed updated classification of the keratodermas. Arch.
[33253] Derm. 132: 640-651, 1996.

[33254] 4728.

[33255] Tyldesley, W. R.: Oral leukoplakia associated with tylosis and
[33256] esophageal carcinoma. J. Oral Path. 3: 62-70, 1974.

[33257] 4729.

[33258] von Brevern, M.; Hollstein, M. C.; Risk, J. M.; Garde, J.; Bennett,

[33259] W. P.; Harris, C. C.; Muehlbauer, K.-R.; Field, J. K.: Loss of heterozygosity

[33260] in sporadic oesophageal tumors in the tylosis oesophageal cancer (TOC)

[33261] gene region of chromosome 17q. Oncogene 17: 2101-2105, 1998.

[33262] 4730.

[33263] Yesudian, P.; Premalatha, S.; Thambiah, A. S.: Genetic ty-losis

[33264] with malignancy: a study of a South Indian pedigree. Brit. J. Derm. 102:

[33265] 597-600, 1980.

[33266] 4731.

[33267] Lerman, C.; Caporaso, N. E.; Audrain, J.; Main, D.; Bow-man, E.

[33268] D.; Lockshin, B.; Boyd, N. R.; Shields, P. G.: Evidence sug-gesting

[33269] the role of specific genetic factors in cigarette smoking. Health

[33270] Psych. 18: 14-20, 1999.

[33271] 4732.

[33272] Lossie, A. C.; Vandenbergh, D. J.; Uhl, G. R.; Camper, S. A.:

[33273] Localization of the dopamine transporter gene, Dat1, on mouse chromosome 4733.

[33274] Mammalian Genome 5: 117-118, 1994.

[33275]

[33276] 13. Sabol, S. Z.; Nelson, M. L.; Fisher, C.; Gunzerath, L.; Brody,

[33277] C. L.; Hu, S.; Sirota, L. A.; Marcus, S. E.; Greenberg, B. D.; Lucas,

[33278] F. R., IV; Benjamin, J.; Murphy, D. L.; Hamer, D. H.: A genetic association

[33279] for cigarette smoking behavior. Health Psych. 18: 7-13, 1999.

[33280] 4734.

[33281] Tiihonen, J.; Kuikka, J.; Bergstrom, K.; Hakola, P.; Karhu, J.;

[33282] Ryynanen, O.-P.; Fohr, J.: Altered striatal dopamine re-uptake site

[33283] densities in habitually violent and non-violent alcoholics. Nature

[33284] Med. 1: 654-657, 1995.

[33285] 4735.

[33286] Vandenbergh, D. J.; Persico, A. M.; Hawkins, A. L.; Griffin, C.

[33287] A.; Li, X.; Jabs, E. W.; Uhl, G. R.: Human dopamine transporter gene

[33288] (DAT1) maps to chromosome 5p15.3 and displays a VNTR. Genomics 14:

[33289] 1104-1106, 1992.

[33290] 4736.

[33291] Vandenbergh, D. J.; Persico, A. M.; Uhl, G. R.: A human dopamine

[33292] transporter cDNA predicts reduced glycosylation, displays a novel

[33293] repetitive element and provides racially-dimorphic TaqI RFLPs. Molec.

[33294] Brain Res. 15: 161-166, 1992.

[33295] 4737.

[33296] Waldman, I. D.; Rowe, D. C.; Abramowitz, A.; Kozel, S. T.; Mohr,

[33297] J. H.; Sherman, S. L.; Cleveland, H. H.; Sanders, M. L.;

Gard, J.

[33298] M. C.; Stever, C.: Association and linkage of the dopamine transporter

[33299] gene and attention-deficit hyperactivity disorder in children: heterogeneity

[33300] owing to diagnostic subtype and severity. Am. J. Hum. Genet. 63:

[33301] 1767-1776, 1998.

[33302] 4738.

[33303] Haila, S.; Hoglund, P.; Scherer, S. W.; Lee, J. R.; Kristo, P.;

[33304] Coyle, B.; Trembath, R.; Holmberg, C.; de la Chapelle, A.; Kere, J.

[33305] : Genomic structure of the human congenital chloride diarrhea (CLD)

[33306] gene. Gene 214: 87-93, 1998.

[33307] 4739.

[33308] Hoglund, P.; Haila, S.; Gustavson, K.-H.; Taipale, M.; Hannula,

[33309] K.; Popinska, K.; Holmberg, C.; Socha, J.; de la Chapelle, A.; Kere,

[33310] J.: Clustering of private mutations in the congenital chlo-

ride diarrhea/down-regulated
[33311] in adenoma gene. Hum. Mutat. 11: 321-327, 1998.

[33312] 4740.

[33313] Huang, H.; Mahler-Araujo, B. M.; Sankila, A.; Chimelli, L.; Yonekawa,
[33314] Y.; Kleihues, P.; Ohgaki, H.: APC mutations in sporadic medulloblastomas. Am.
[33315] J. Path. 156: 433-437, 2000.

[33316] 4741.

[33317] von Boehmer, H.; Sarukhan, A.: GAD, a single autoantigen for
[33318] diabetes. Science 284: 1135-1136, 1999.

[33319] 4742.

[33320] Williamson, E. A.; Ince, P. G.; Harrison, D.; Kendall-Taylor,
[33321] P.; Harris, P. E.: G-protein mutations in human pituitary adrenocorticotrophic
[33322] hormone-secreting adenomas. Europ. J. Clin. Invest. 25: 128-131,
[33323] 1995.

[33324] 4743.

[33325] Farlie, P. G.; Dringen, R.; Rees, S. M.; Kannourakis, G.; Bernard,
[33326] O.: bcl-2 transgene expression can protect neurons against developmental
[33327] and induced cell death. Proc. Nat. Acad. Sci. 92: 4397-4401, 1995.
[33328] 4744.

[33329] Fesus, L.; Davies, P. J. A.; Piacentini, M.: Apoptosis: molec-ular
[33330] mechanisms in programmed cell death. Europ. J. Cell Biol. 56: 170-177,
[33331] 1991.

[33332] 4745.

[33333] Frankel, S. R.: Correction: location of bcl-2 oncogene. (Letter) New
[33334] Eng. J. Med. 328: 141, 1993.

[33335] 4746.

[33336] Graninger, W. B.; Seto, M.; Boutain, B.; Goldman, P.; Ko-rsmeyer,
[33337] S. J.: Expression of Bcl-2 and Bcl-2-Ig fusion transcripts in normal

[33338] and neoplastic cells. J. Clin. Invest. 80: 1512-1515, 1987.

[33339] 4747.

[33340] Haluska, F. G.; Tsujimoto, Y.; Croce, C. M.: Mechanisms of chromosome

[33341] translocation in B- and T-cell neoplasia. Trends Genet. 3: 11-15,

[33342] 1987.

[33343] 4748.

[33344] Hengartner, M. O.; Horvitz, H. R.: C. elegans cell survival gene

[33345] ced-9 encodes a functional homolog of the mammalian proto-oncogene

[33346] bcl-2. Cell 76: 665-676, 1994.

[33347] 4749.

[33348] Hockenbery, D.; Nunez, G.; Milliman, C.; Schreiber, R. D.; Korsmeyer,

[33349] S. J.: Bcl-2 is an inner mitochondrial membrane protein that blocks

[33350] programmed cell death. Nature 348: 334-336, 1990.

[33351] 4750.

[33352] Jacobson, M. D.; Burne, J. F.; King, M. P.; Miyashita, T.; Reed,
[33353] J. C.; Raff, M. C.: Bcl-2 blocks apoptosis in cells lacking mitochondrial
[33354] DNA. Nature 361: 365-369, 1993.
[33355] 4751.
[33356] Ji, L.; Mochon, E.; Arcinas, M.; Boxer, L. M.: CREB proteins
[33357] function as positive regulators of the translocated bcl-2 allele in
[33358] t(14;18) lymphomas. J. Biol. Chem. 271: 22687-22691, 1996.
[33359] 4752.
[33360] Korsmeyer, S. J.: Bcl-2 initiates a new category of onco-genes:
[33361] regulators of cell death. Blood 80: 879-886, 1992.
[33362] 4753.
[33363] Lee, M.-S.; Chang, K.-S.; Cabanillas, F.; Freireich, E. J.; Trujillo,
[33364] J. M.; Stass, S. A.: Detection of minimal residual cells car-rying
[33365] the t(14;18) by DNA sequence amplification. Science 237:

175-178,

[33366] 1987.

[33367] 4754.

[33368] Limana, F.; Urbanek, K.; Chimenti, S.; Quaini, F.; Leri, A.; Kajstura,

[33369] J.; Nadal-Ginard, B.; Izumo, S.; Anversa, P.: bcl-2 overex-pression

[33370] promotes myocyte proliferation. Proc. Nat. Acad. Sci. 99: 6257-6262,

[33371] 2002.

[33372] 4755.

[33373] Liu, Y.; Hernandez, A. M.; Shibata, D.; Cortopassi, G. A.: BCL2

[33374] translocation frequency rises with age in humans. Proc. Nat. Acad.

[33375] Sci. 91: 8910-8914, 1994.

[33376] 4756.

[33377] Marsden, V. S.; O'Connor, L.; O'Reilly, L. A.; Silke, J.; Met-calf,

[33378] D.; Ekert, P. G.; Huang, D. C. S.; Cecconi, F.; Kuida, K.; Tomaselli,

[33379] K. J.; Roy, S.; Nicholson, D. W.; Vaux, D. L.; Bouillet, P.; Adams,
[33380] J. M.; Strasser, A.: Apoptosis initiated by Bcl-2-regulated caspase
[33381] activation independently of the cytochrome c/ Apaf-1/caspase-9 apoptosome. Nature 419:
[33382] 634-637, 2002.

[33383] 4757.

[33384] Martinou, J.-C.; Dubois-Dauphin, M.; Staple, J. K.; Rodriguez,
[33385] I.; Frankowski, H.; Missotten, M.; Albertini, P.; Talabot, D.; Catsicas,
[33386] S.; Pietra, C.; Huarte, J.: Overexpression of BCL-2 in transgenic
[33387] mice protects neurons from naturally occurring cell death and experimental
[33388] ischemia. Neuron 13: 1017-1030, 1994.

[33389] 4758.

[33390] McDonnell, T. J.; Deane, N.; Platt, F. M.; Nunez, G.; Jaeger,
[33391] U.; McKearn, J. P.; Korsmeyer, S. J.: Bcl-2-immunoglobulin transgenic

[33392] mice demonstrate extended B cell survival and follicular lymphoproliferation. Cell 57:

[33393] 79-88, 1989.

[33394] 4759.

[33395] McGill, G. G.; Horstmann, M.; Widlund, H. R.; Du, J.; Motyckova,

[33396] G.; Nishimura, E. K.; Lin, Y.-L.; Ramaswamy, S.; Avery, W.; Ding,

[33397] H.-F.; Jordan, S. A.; Jackson, I. J.; Korsmeyer, S. J.; Golub, T.

[33398] R.; Fisher, D. E.: Bcl2 regulation by the melanocyte master regulator

[33399] Mitf modulates lineage survival and melanoma cell viability. Cell 109:

[33400] 707-718, 2002.

[33401] 4760.

[33402] Migheli, A.; Cavalla, P.; Piva, R.; Giordana, M. T.; Schiffer,

[33403] D.: Bcl-2 protein expression in aged brain and neurodegenerative

[33404] diseases. Neuroreport 5: 1906-1908, 1994.

[33405] 4761.

[33406] Huelsken, J.; Vogel, R.; Erdmann, B.; Cotsarelis, G.; Birch-
meier,
[33407] W.: Beta-catenin controls hair follicle morphogenesis and
stem cell
[33408] differentiation in the skin. Cell 105: 533-545, 2001.

[33409] 4762.

[33410] Ilyas, M.; Tomlinson, I. P. M.; Rowan, A.; Pignatelli, M.;
Bodmer,
[33411] W. F.: Beta-catenin mutations in cell lines established
from human
[33412] colorectal cancers. Proc. Nat. Acad. Sci. 94: 10330-10334,
1997.
[33413] 4763.

[33414] Iwao, K.; Nakamori, S.; Kameyama, M.; Imaoka, S.; Ki-
noshita, M.;
[33415] Fukui, T.; Ishiguro, S.; Nakamura, Y.; Miyoshi, Y.: Activa-
tion of
[33416] the beta-catenin gene by interstitial deletions involving
exon 3 in
[33417] primary colorectal carcinomas without adenomatous poly-
posis coli mutations. Cancer
[33418] Res. 58: 1021-1026, 1998.

[33419] 4764.

[33420] Kawasaki, Y.; Senda, T.; Ishidata, T.; Koyama, R.; Mor-
ishita,
[33421] T.; Iwayama, Y.; Higuchi, O.; Akiyama, T.: Asef, a link be-
tween the
[33422] tumor suppressor APC and G-protein signaling. Science
289: 1194-1197,
[33423] 2000.

[33424] 4765.

[33425] Koch, A.; Denkhaus, D.; Albrecht, S.; Leuschner, I.; von
Schweinitz,
[33426] D.; Pietsch, T.: Childhood hepatoblastomas frequently
carry a mutated
[33427] degradation targeting box of the beta-catenin gene. Can-
cer Res. 59:
[33428] 269-273, 1999.

[33429] 4766.

[33430] Korinek, V.; Barker, N.; Morin, P. J.; van Wichen, D.; de
Weger,
[33431] R.; Kinzler, K. W.; Vogelstein, B.; Clevers, H.: Constitutive
transcriptional

[33432] activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma. Science 275:
[33433] 1784-1787, 1997.

[33434] 4767.

[33435] Kraus, C.; Liehr, T.; Hulsken, J.; Behrens, J.; Birchmeier, W.;

[33436] Grzeschik, K.-H.; Ballhausen, W. G.: Localization of the human beta-catenin
[33437] gene (CTNNB1) to 3p21: a region implicated in tumor development. Genomics 23:
[33438] 272-274, 1994.

[33439] 4768.

[33440] Legoix, P.; Bluteau, O.; Bayer, J.; Perret, C.; Balabaud, C.;

[33441] Belghiti, J.; Franco, D.; Thomas, G.; Laurent-Puig, P.; Zucman-Rossi,
[33442] J.: Beta-catenin mutations in hepatocellular carcinoma correlate
[33443] with a low rate of loss of heterozygosity. Oncogene 18: 4044-4046,
[33444] 1999.

[33445] 4769.

[33446] Lin, S.-Y.; Xia, W.; Wang, J. C.; Kwong, K. Y.; Spohn, B.; Wen,
[33447] Y.; Pestell, R. G.; Hung, M.-C.: Beta-catenin, a novel prog-nostic
[33448] marker for breast cancer: its roles in cyclin D1 expression and cancer
[33449] progression. Proc. Nat. Acad. Sci. 97: 4262-4266, 2000.

[33450] 4770.

[33451] McCrea, P. D.; Turck, C. W.; Gumbiner, B.: A homolog of the armadillo
[33452] protein in Drosophila (plakoglobin) associated with E-cadherin. Science 254:
[33453] 1359-1361, 1991.

[33454] 4771.

[33455] Morin, P. J.; Sparks, A. B.; Korinek, V.; Barker, N.; Clevers,
[33456] H.; Vogelstein, B.; Kinzler, K. W.: Activation of beta-catenin-Tcf
[33457] signaling in colon cancer by mutations in beta-catenin or APC. Science 275:
[33458] 1787-1790, 1997.

[33459] 4772.

[33460] Murase, S.; Mosser, E.; Schuman, E. M.: Depolarization drives
[33461] beta-catenin into neuronal spines promoting changes in synaptic structure
[33462] and function. Neuron 35: 91-105, 2002.

[33463] 4773.

[33464] Nollet, F.; Berx, G.; Molemans, F.; van Roy, F.: Genomic organization
[33465] of the human beta-catenin gene (CTNNB1). Genomics 32: 413-424, 1996.
[33466] 4774.

[33467] Peifer, M.: Cancer, catenins, and cuticle pattern: a com-plex
[33468] connection. Science 262: 1667-1668, 1993.

[33469] 4775.

[33470] Rodova, M.; Islam, M. R.; Maser, R. L.; Calvet, J. P.: The polycystic
[33471] kidney disease-1 promoter is a target of the beta-catenin/T-cell factor
[33472] pathway. J. Biol. Chem. 277: 29577-29583, 2002.

[33473] 4776.

[33474] Roose, J.; Huls, G.; van Beest, M.; Moerer, P.; van der Horn,

[33475] K.; Goldschmeding, R.; Logtenberg, T.; Clevers, H.: Synergy between

[33476] tumor suppressor APC and the beta-catenin-Tcf4 target Tcf1. Science 285:

[33477] 1923-1926, 1999.

[33478] 4777.

[33479] Rubinfeld, B.; Robbins, P.; El-Gamil, M.; Albert, I.; Porfiri,

[33480] E.; Polakis, P.: Stabilization of beta-catenin by genetic defects

[33481] in melanoma cell lines. Science 275: 1790-1792, 1997.

[33482] 4778.

[33483] Saadi-Kheddouci, S.; Berrebi, D.; Romagnolo, B.; Cluzeaud, F.;

[33484] Peuchmaur, M.; Kahn, A.; Vandewalle, A.; Perret, C.: Early development

[33485] of polycystic kidney disease in transgenic mice expressing an activated

[33486] mutant of the beta-catenin gene. Oncogene 20: 5972-5981, 2001.

[33487] 4779.

[33488] Sagae, S.; Kobayashi, K.; Nishioka, Y.; Sugimura, M.; Ishioka,
[33489] S.; Nagata, M.; Terasawa, K.; Tokino, T.; Kudo, R.: Mutational analysis
[33490] of beta-catenin gene in Japanese ovarian carcinomas: frequent mutations
[33491] in endometrioid carcinomas. Jpn. J. Cancer Res. 90: 510-515, 1999.
[33492] 4780.

[33493] Trent, J. M.; Wiltshire, R.; Su, L.-K.; Nicolaides, N. C.; Vogelstein,
[33494] B.; Kinzler, K. W.: The gene for the APC-binding protein beta-catenin
[33495] (CTNNB1) maps to chromosome 3p22, a region frequently altered in human
[33496] malignancies. Cytogenet. Cell Genet. 71: 343-344, 1995.

[33497] 4781.

[33498] van Hengel, J.; Nollet, F.; Berx, G.; van Roy, N.; Speleman, F.;
[33499] van Roy, F.: Assignment of the human beta-catenin gene (CTNNB1) to
[33500] 3p22-p21.3 by fluorescence in situ hybridization. Cyto-

genet. Cell
[33501] Genet. 70: 68-70, 1995.

[33502] 4782.

[33503] Chan, S. J.; San Segundo, B.; McCormick, M. B.; Steiner, D. F.
[33504] : Nucleotide and predicted amino acid sequences of cloned human and
[33505] mouse preprocathepsin B cDNAs. Proc. Nat. Acad. Sci. 83: 7721-7725,
[33506] 1986.

[33507] 4783.

[33508] Deussing, J.; Roth, W.; Rommerskirch, W.; Wiederanders, B.; von
[33509] Figura, K.; Peters, C.: The genes of the lysosomal cysteine proteinases
[33510] cathepsin B, H, L, and S map to different mouse chromo-somes. Mammalian
[33511] Genome 8: 241-245, 1997.

[33512] 4784.

[33513] Shigemitsu, K.; Sekido, Y.; Usami, N.; Mori, S.; Sato, M.; Horio,

[33514] Y.; Hasegawa, Y.; Bader, S. A.; Gazdar, A. F.; Minna, J. D.; Hida,

[33515] T.; Yoshioka, H.; Imaizumi, M.; Ueda, Y.; Takahashi, M.; Shimokata,

[33516] K.: Genetic alteration of the beta-catenin gene (CTNNB1) in human

[33517] lung cancer and malignant mesothelioma and identification of a new

[33518] 3p21.3 homozygous deletion. Oncogene 20: 4249-4257, 2001.

[33519] 4785.

[33520] Takahashi, M.; Arita, Y.; Yamagata, K.; Matsukawa, Y.; Okutomi,

[33521] K.; Horie, M.; Shimomura, I.; Hotta, K.; Kuriyama, H.; Kihara, S.;

[33522] Nakamura, T.; Yamashita, S.; Funahashi, T,; Matsuzawa, Y.: Genomic

[33523] structure and mutations in adipose-specific gene, adiponectin. Int.

[33524] J. Obes. 24: 861-868, 2000.

[33525] 4786.

[33526] Yang, W.-S.; Lee, W.-J.; Funahashi, T.; Tanaka, S.; Mat-

suzawa,
[33527] Y.; Chao, C.-L.; Chen, C.-L.; Tai, T.-Y.; Chuang, L.-M.: Weight reduction
[33528] increases plasma levels of an adipose-derived anti-inflammatory protein,
[33529] adiponectin. J. Clin. Endocr. Metab. 86: 3815-3819, 2001.

[33530] 4787.

[33531] Camenisch, T. D.; Koller, B. H.; Earp, H. S.; Matsushima, G. K.
[33532] : A novel receptor tyrosine kinase, mer, inhibits TNF-alpha production
[33533] and lipopolysaccharide-induced endotoxic shock. J. Immun. 162: 3498-3503,
[33534] 1999.

[33535] 4788.

[33536] D'Cruz, P. M.; Yasumura, D.; Weir, J.; Matthes, M. T.; Abderrahim,
[33537] H.; LaVail, M. M.; Vollrath, D.: Mutation of the receptor tyrosine
[33538] kinase gene Mertk in the retinal dystrophic RCS rat. Hum. Molec.
[33539] Genet. 9: 645-651, 2000.

[33540] 4789.

[33541] Gal, A.; Li, Y.; Thompson, D. A.; Weir, J.; Orth, U.; Jacob-
son,
[33542] S. G.; Apfelstedt-Sylla, E.; Vollrath, D.: Mutations in
MERTK, the
[33543] human orthologue of the RCS rat retinal dystrophy gene,
cause retinitis
[33544] pigmentosa. Nature Genet. 26: 270-271, 2000.

[33545] 4790.

[33546] Graham, D. K.; Dawson, T. L.; Mullaney, D. L.; Snodgrass,
H. R.;
[33547] Earp, H. S.: Cloning and mRNA expression analysis of a
novel human
[33548] protooncogene, c-mer. Cell Growth Differ. 5: 647-657,
1994.
[33549] 4791.

[33550] Lu, Q.; Gore, M.; Zhang, Q.; Camenisch, T.; Boast, S.;
Casagranda,
[33551] F.; Lai, C.; Skinner, M. K.; Klein, R.; Matsushima, G. K.;
Earp, H.
[33552] S.; Goff, S. P.; Lemke, G.: Tyro-3 family receptors are es-
sential

[33553] regulators of mammalian spermatogenesis. Nature 398: 723-728, 1999.

[33554] 4792.

[33555] Scott, R. S.; McMahon, E. J.; Pop, S. M.; Reap, E. A.; Caric-

[33556] chio, R.; Cohen, P. L.; Earp, H. S.; Matsushima, G. K.: Phagocy-

[33557] tosis and clearance of apoptotic cells is mediated by MER. Nature 411: 207-211,

[33558] 2001.

[33559] 4793.

[33560] Vollrath, D.; Feng, W.; Duncan, J. L.; Yasumura, D.; D'Cruz,

[33561] P. M.; Chappelow, A.; Matthes, M. T.; Kay, M. A.; LaVail, M.

[33562] M.: Correction of the retinal dystrophy phenotype of the RCS rat by viral

[33563] gene transfer of Mertk. Proc. Nat. Acad. Sci. 98: 12584-12589, 2001.

[33564] 4794.

[33565] Weier, H.-U. G.; Fung, J.; Lersch, R. A.: Assignment of pro-

[33566] tooncogene

[33566] MERTK (a.k.a. c-mer) to human chromosome 2q14.1 by in situ hybridization. Cytogenet.

[33567] Cell Genet. 84: 91-92, 1999.

[33568] 4795.

[33569] Adler, H. J.; Winnicki, R. S.; Gong, T.-W. L.; Lomax, M. I.: A

[33570] gene upregulated in the acoustically damaged chick basilar papilla

[33571] encodes a novel WD40 repeat protein. Genomics 56: 59-69, 1999.

[33572] 4796.

[33573] Geck, P.; Sonnenschein, C.; Soto, A. M.: The D13S171 marker, misannotated

[33574] to BRCA2, links the AS3 gene to various cancers. (Letter) Am. J.

[33575] Hum. Genet. 69: 461-463, 2001.

[33576] 4797.

[33577] Geck, P.; Szelei, J.; Jimenez, J.; Sonnenschein, C.; Soto, A. M.

[33578] : Early gene expression during androgen-induced inhibition of proliferation

[33579] of prostate cancer cells: a new suppressor candidate on

chromosome
[33580] 13, in the BRCA2-Rb1 locus. J. Steroid Biochem. Molec. Biol. 68:
[33581] 41-50, 1999.

[33582] 4798.

[33583] Bontekoe, C. J. M.; McIlwain, K. L.; Nieuwenhuizen, I. M.; Yuva-Paylor,
[33584] L. A.; Nellis, A.; Willemsen, R.; Fang, Z.; Kirkpatrick, L.; Bakker,
[33585] C. E.; McAninch, R.; Cheng, N. C.; Merriweather, M.; Hoogeveen, A.
[33586] T.; Nelson, D.; Paylor, R.; Oostra, B. A.: Knockout mouse model for
[33587] Fxr2: a model for mental retardation. Hum. Molec. Genet. 11: 487-498,
[33588] 2002.

[33589] 4799.

[33590] Tamanini, F.; Willemsen, R.; van Unen, L.; Bontekoe, C.; Galjaard,
[33591] H.; Oostra, B. A.; Hoogeveen, A. T.: Differential expression of FMR1,
[33592] FXR1 and FXR2 proteins in human brain and testis. Hum.

Molec. Genet. 6:
[33593] 1315-1322, 1997.

[33594] 4800.

[33595] Zhang, Y.; O'Connor, J. P.; Siomi, M. C.; Srinivasan, S.; Dutra,
[33596] A.; Nussbaum, R. L.; Dreyfuss, G.: The fragile X mental retardation
[33597] syndrome protein interacts with novel homologs FXR1 and FXR2. EMBO
[33598] J. 14: 5358-5366, 1995.

[33599] 4801.

[33600] Samaridis, J.; Colonna, M.: Cloning of novel immunoglobulin superfamily
[33601] receptors expressed on human myeloid and lymphoid cells: structural
[33602] evidence for new stimulatory and inhibitory pathways. Europ. J. Immun. 27:
[33603] 660-665, 1997.

[33604] 4802.

[33605] Wagtmann, N.; Rojo, S.; Eichler, E.; Mohrenweiser, H.; Long, E.

[33606] O.: A new human gene complex encoding the killer cell inhibitory
[33607] receptors and related monocyte/macrophage receptors. Curr. Biol. 7:
[33608] 615-618, 1997.

[33609] 4803.

[33610] Chang, C. C.; Ciubotariu, R.; Manavalan, J. S.; Yuan, J.; Colovai,
[33611] A. I.; Piazza, F.; Lederman, S.; Colonna, M.; Cortesini, R.; Dalla-Favera,
[33612] R.; Suciu-Foca, N.: Tolerization of dendritic cells by Ts cells:
[33613] the crucial role of inhibitory receptors ILT3 and ILT4. Na-ture Immun. 3:
[33614] 237-243, 2002.

[33615] 4804.

[33616] Lee, J. K.; Bhakta, S.; Rosen, S. D.; Hemmerich, S.: Cloning and
[33617] characterization of a mammalian N-acetylglucosamine-6-sulfotransferase
[33618] that is highly restricted to intestinal tissue. Biochem. Bio-phys.

[33619] Res. Commun. 263: 543-549, 1999.

[33620] 4805.

[33621] Arm, J. P.; Nwankwo, C.; Austen, K. F.: Molecular identification

[33622] of a novel family of human Ig superfamily members that possess immunoreceptor

[33623] tyrosine-based inhibition motifs and homology to the mouse gp49B1

[33624] inhibitory receptor. J. Immun. 159: 2342-2349, 1997.

[33625] 4806.

[33626] Cella, M.; Dohring, C.; Samaridis, J.; Dessing, M.; Brockhaus,

[33627] M.; Lanzavecchia, A.; Colonna, M.: A novel inhibitory receptor (ILT3)

[33628] expressed on monocytes, macrophages, and dendritic cells involved

[33629] in antigen processing. J. Exp. Med. 185: 1743-1751, 1997.

[33630] 4807.

[33631] Raymond, C. S.; Parker, E. D.; Kettlewell, J. R.; Brown, L. G.;

[33632] Page, D. C.; Kusz, K.; Jaruzelska, J.; Reinberg, Y.; Fletjer,

W. L.;

[33633] Bardwell, V. J.; Hirsch, B.; Zarkower, D.: A region of human chromosome

[33634] 9p required for testis development contains two genes re-lated to known

[33635] sexual regulators. Hum. Molec. Genet. 8: 989-996, 1999.

[33636] 4808.

[33637] Ancian, P.; Lambeau, G.; Mattei, M.-G.; Lazdunski, M.: The human

[33638] 180-kDa receptor for secretory phospholipases A2: molecular cloning,

[33639] identification of a secreted soluble form, expression, and chromosomal

[33640] localization. J. Biol. Chem. 270: 8963-8970, 1995.

[33641] 4809.

[33642] Okazaki, I.; Kinoshita, K.; Muramatsu, M.; Yoshikawa, K.; Honjo,

[33643] T.: The AID enzyme induces class switch recombination in fibroblasts. Nature 416:
[33644] 340-345, 2002.

[33645] 4810.

[33646] Petersen-Mahrt, S. K.; Harris, R. S.; Neuberger, M. S.: AID mutates
[33647] E. coli suggesting a DNA deamination mechanism for antibody diversification. Nature 418:
[33648] 99-104, 2002.

[33649] 4811.

[33650] Revy, P.; Muto, T.; Levy, Y.; Geissmann, F.; Plebani, A.; Sanal,
[33651] O.; Catalan, N.; Forveille, M.; Dufourcq-Lagelouse, R.; Gennery, A.;
[33652] Tezcan, I.; Ersoy, F.; and 9 others: Activation-induced cytidine
[33653] deaminase (AID) deficiency causes the autosomal recessive form of
[33654] the hyper-IgM syndrome (HIGM2). Cell 102: 565-575, 2000.
[33655] 4812.

[33656] Yoshikawa, K.; Okazaki, I.; Eto, T.; Kinoshita, K.; Muramatsu,

[33657] M.; Nagaoka, H.; Honjo, T.: AID enzyme-induced hypermutation in an

[33658] actively transcribed gene in fibroblasts. Science 296: 2033-2036,

[33659] 2002.

[33660] 4813.

[33661] Geiszt, M.; Kopp, J. B.; Varnai, P.; Leto, T. L.: Identification

[33662] of Renox, an NAD(P)H oxidase in kidney. Proc. Nat. Acad. Sci. 97:

[33663] 8010-8014, 2000.

[33664] 4814.

[33665] Shiose, A.; Kuroda, J.; Tsuruya, K.; Hirai, M.; Hirakata, H.; Naito,

[33666] S.; Hattori, M.; Sakaki Y.; Sumimoto, H.: A novel superoxide-producing

[33667] NAD(P)H oxidase in kidney. J. Biol. Chem. 276: 1417-1423, 2001.

[33668] 4815.

[33669] Kokame, K.; Kato, H.; Miyata, T.: Homocysteine-respon-

dent genes

[33670] in vascular endothelial cells identified by differential dis-
play analysis:

[33671] GRP78/BiP and novel genes. J. Biol. Chem. 271:
29659-29665, 1996.

[33672] 4816.

[33673] Park, H,; Adams, M. A.; Lachat, P.; Bosman, F.; Pang, S. C.;
Graham,

[33674] C. H.: Hypoxia induces the expression of a 43-kDa protein
(PROXY-1)

[33675] in normal and malignant cells. Biochem. Biophys. Res.
Commun. 276:

[33676] 321-328, 2000.

[33677] 4817.

[33678] Shimono, A.; Okuda, T.; Kondoh, H.: N-myc-dependent
repression

[33679] of Ndr1, a gene identified by direct subtraction of whole
mouse embryo

[33680] cDNAs between wild type and N-myc mutant. Mech. Dev.
83: 39-52,

[33681] 1999.

[33682] 4818.

[33683] van Belzen, N.; Dinjens, W. N. M.; Diesveld, M. P. G.; Groen, N.

[33684] A.; van der Made, A. C. J.; Nozawa, Y.; Vliestra, R.; Trapman, J.;

[33685] Bosman, F. T.: A novel gene which is up-regulated during colon epithelial

[33686] cell differentiation and down-regulated in colorectal neoplasms. Lab.

[33687] Invest. 77: 85-92, 1997.

[33688] 4819.

[33689] Zhou, D.; Salnikow, K.; Costa, M.: Cap43, a novel gene specifically

[33690] induced by Ni2+ compounds. Cancer Res. 58: 2182-2189, 1998.

[33691] 4820.

[33692] Ludwig, J.; Kerscher, S.; Brandt, U.; Pfeiffer, K.; Getlawi, F.;

[33693] Apps, D. K.; Schagger, H.: Identification and characterization of

[33694] a novel 9.2-kDa membrane sector-associated protein of vacuolar proton-ATPase

[33695] from chromaffin granules. J. Biol. Chem. 273: 10939-10947, 1998.

[33696] 4821.

[33697] Nanji, M.; Coronado, V. A.; Cox, D. W.: ATP6H, a subunit of vacuolar

[33698] ATPase involved in metal transport: evaluation in canine copper toxicosis. Mammalian

[33699] Genome 12: 617-621, 2001.

[33700] 4822.

[33701] Arama, E.; Yanai, A.; Kilfin, G.; Bernstein, A.; Motro, B.: Murine

[33702] NIMA-related kinases are expressed in patterns suggesting distinct

[33703] functions in gametogenesis and a role in the nervous system. Oncogene 16:

[33704] 1813-1823, 1998.

[33705] 4823.

[33706] Fry, A. M.; Meraldi, P.; Nigg, E. A.: A centrosomal function for

[33707] the human Nek2 protein kinase, a member of the NIMA family of cell

[33708] cycle regulators. EMBO J. 17: 470-481, 1998.

[33709] 4824.

[33710] Schultz, S. J.; Fry, A. M.; Sutterlin, C.; Ried, T.; Nigg, E. A.

[33711] : Cell cycle-dependent expression of Nek2, a novel human protein kinase

[33712] related to the NIMA mitotic regulator of Aspergillus nidulans. Cell

[33713] Growth Diff. 5: 625-635, 1994.

[33714] 4825.

[33715] Schultz, S. J.; Nigg, E. A.: Identification of 21 novel human

[33716] protein kinases, including 3 members of a family related to the cell

[33717] cycle regulator nimA of Aspergillus nidulans. Cell Growth Diff. 4:

[33718] 821-830, 1993.

[33719] 4826.

[33720] Tamari, M.; Daigo, Y.; Nakamura, Y.: Isolation and characterization

[33721] of a novel serine threonine kinase gene on chromosome 3q22-21.3. J.

[33722] Hum. Genet. 44: 116-120, 1999.

[33723] 4827.

[33724] Tamari, M.; Daigo, Y.; Ishikawa, S.; Nakamura, Y.: Genomic structure
[33725] of a novel human gene (XYLB) on chromosome 3p22-p21.3 encoding a xylulokinase-like
[33726] protein. Cytogenet. Cell Genet. 82: 101-104, 1998.

[33727] 4828.

[33728] Daigo, Y.; Nishiwaki, T.; Kawasoe, T.; Tamari, M.; Tsuchiya, E.;
[33729] Nakamura, Y.: Molecular cloning of a candidate tumor suppressor gene,
[33730] DLC1, from chromosome 3p21.3. Cancer Res. 59: 1966-1972, 1999.
[33731] 4829.

[33732] Migone, T.-S.; Zhang, J.; Luo, X.; Zhuang, L.; Chen, C.; Hu, B.;
[33733] Hong, J. S.; Perry, J. W.; Chen, S.-F.; Zhou, J. X. H.; Cho, Y. H.;
[33734] Ullrich, S.; and 14 others: TL1A is a TNF-like ligand for DR3 and
[33735] TR6/DcR3 and functions as a T cell costimulator. Immu- nity 16: 479-492,
[33736] 2002.

[33737] 4830.

[33738] Zhai, Y.; Ni, J.; Jiang, G.-W.; Lu, J.; Xing, L.; Lincoln, C.;

[33739] Carter, K. C.; Janat, F.; Kozak, D.; Xu, S.; Rojas, L.; Aggar-
wal,

[33740] B. B.; Ruben, S.; Li, L.-Y.; Gentz, R.; Yu, G.-L.: VEGI, a
novel

[33741] cytokine of the tumor necrosis factor family, is an angio-
genesis inhibitor

[33742] that suppresses the growth of colon carcinomas in vivo.
FASEB J. 13:

[33743] 181-189, 1999.

[33744] 4831.

[33745] Yue, T.-L.; Ni, J.; Romanic, A. M.; Gu, J.-L.; Keller, P.;
Wang,

[33746] C.; Kumar, S.; Yu, G.; Hart, T. K.; Wang, X.; Xia, Z.; DeWolf,
W.

[33747] E., Jr.; Feuerstein, G. Z.: TL1, a novel tumor necrosis fac-
tor-like

[33748] cytokine, induces apoptosis in endothelial cells: involve-
ment of activation

[33749] of stress protein kinases (stress-activated protein kinase
and p38

[33750] mitogen-activated protein kinase) and caspase-3-like protease. J.

[33751] Biol. Chem. 274: 1479-1486, 1999.

[33752] 4832.

[33753] Gu, Z.; Flemington, C.; Chittenden, T.; Zambetti, G. P.: Ei24,

[33754] a p53 response gene involved in growth suppression and apoptosis. Molec.

[33755] Cell. Biol. 20: 233-241, 2000.

[33756] 4833.

[33757] Gu, Z.; Gilbert, D. J.; Valentine, V. A.; Jenkins, N. A.; Copeland,

[33758] N. G.; Zambetti, G. P.: The p53-inducible gene EI24/PIG8 localizes

[33759] to human chromosome 11q23 and the proximal region of mouse chromosome

[33760] 9. Cytogenet. Cell Genet. 89: 230-233, 2000.

[33761] 4834.

[33762] Polyak, K.; Xia, Y.; Zweier, J. L.; Kinzler, K. W.; Vogelstein,

[33763] B.: A model for p53-induced apoptosis. Nature 389: 300-305, 1997.

[33764] 4835.

[33765] Forsberg, L.; Leeb, L.; Thoren, S.; Morgenstern, R.; Jakobs-
son,
[33766] P.-J.: Human glutathione dependent prostaglandin E syn-
thase: gene
[33767] structure and regulation. FEBS Lett. 471: 78-82, 2000.

[33768] 4836.

[33769] Jakobsson, P.-J.; Morgenstern, R.; Mancini, J.; Ford-
Hutchinson,
[33770] A.; Persson, B.: Common structural features of MAPEG: a
widespread
[33771] superfamily of membrane associated proteins with highly
divergent
[33772] functions in eicosanoid and glutathione metabolism. Pro-
tein Sci. 8:
[33773] 689-692, 1999.

[33774] 4837.

[33775] Jakobsson, P.-J.; Thoren, S.; Morgenstern, R.; Samuelsson,
B.:
[33776] Identification of human prostaglandin E synthase: a mi-
crosomal, glutathione-dependent,

[33777] inducible enzyme, constituting a potential novel drug target. Proc.

[33778] Nat. Acad. Sci. 96: 7220-7225, 1999.

[33779] 4838.

[33780] Hernandez, M.-C.; Andres-Barquin, P. J.; Holt, I.; Israel, M. A.

[33781] : Cloning of human ENC-1 and evaluation of its expression and regulation

[33782] in nervous system tumors. Exp. Cell Res. 242: 470-477, 1998.

[33783] 4839.

[33784] Hernandez, M.-C.; Andres-Barquin, P. J.; Israel, M. A.: Assignment

[33785] of the ectodermal-neural cortex 1 gene (Enc1) to mouse chromosome

[33786] band 13D1 by fluorescence in situ hybridization. Cytogenet. Cell

[33787] Genet. 89: 158-159, 2000.

[33788] 4840.

[33789] Hernandez, M.-C.; Andres-Barquin, P. J.; Kuo, W. L.; Israel, M.

[33790] A.: Assignment of the ectodermal-neural cortex 1 gene (ENC1) to human

[33791] chromosome band 5q13 by in situ hybridization. Cyto-

genet. Cell Genet. 87:

[33792] 89-90, 1999.

[33793] 4841.

[33794] Kim, T.-A.; Lim, J.; Ota, S.; Raja, S.; Rogers, R.; Rivnay, B.;

[33795] Avraham, H.; Avraham, S.: NRP/B, a novel nuclear matrix protein,

[33796] associates with p110(RB) and is involved in neuronal dif-

ferentiation. J.

[33797] Cell Biol. 141: 553-566, 1998.

[33798] 4842.

[33799] Rosenfeld, M. G.; Mermod, J.-J.; Amara, S. G.; Swanson, L. W.;

[33800] Sawchenko, P. E.; Rivier, J.; Vale, W. W.; Evans, R. M.: Pro-

duction

[33801] of a novel neuropeptide encoded by the calcitonin gene via tissue-specific

[33802] RNA processing. Nature 304: 129-135, 1983.

[33803] 4843.

[33804] Saller, B.; Feldmann, G.; Haupt, K.; Broecker, M.; Janssen, O.

[33805] E.; Roggendorf, M.; Mann, K.; Lu, M.: RT-PCR-based detection of circulating

[33806] calcitonin-producing cells in patients with advanced medullary thyroid

[33807] cancer. J. Clin. Endocr. Metab. 87: 292-296, 2002.

[33808] 4844.

[33809] Simpson, N. E.; Goodfellow, P. J.; Riddell, D. C.; Hamerton, J.

[33810] L.; Holden, J. J. A.; White, B. N.: Assignment of the calcitonin

[33811] gene to chromosome 11 and probable exclusion of linkage between the

[33812] gene and the locus for multiple endocrine neoplasia type 2. (Abstract) Am.

[33813] J. Hum. Genet. 36: 153S, 1984.

[33814] 4845.

[33815] Struthers, A. D.; Brown, M. J.; Macdonald, D. W. R.; Beacham,

[33816] J. L.; Stevenson, J. C.; Morris, H. R.; MacIntyre, I.: Human calcitonin

[33817] gene related peptide: a potent endogenous vasodilator in man. Clin.

[33818] Sci. 70: 389–393, 1986.

[33819] 4846.

[33820] Testa, J. R.: Personal Communication. Baltimore, Md. 3/1984.

[33821] 4847.

[33822] Tiller-Borcich, J. K.; Capili, H.; Gordan, G. S.: Human brain

[33823] calcitonin gene-related peptide (CGRP) is concentrated in the locus

[33824] caeruleus. Neuropeptides 11: 55–61, 1988.

[33825] 4848.

[33826] Tippins, J. R.: CGRP: a novel neuropeptide from the calci-tonin

[33827] gene is the most potent vasodilator known. J. Hyperten-sion 4 (suppl.

[33828] 5): S102–S105, 1986.

[33829] 4849.

[33830] Tschopp, F. A.; Henke, H.; Petermann, J. B.; Tobler, P. H.; Janzer,

[33831] R.; Hokfelt, T.; Lundberg, J. M.; Cuello, C.; Fischer, J. A.: Calcitonin
[33832] gene-related peptide and its binding sites in the human central nervous
[33833] system and pituitary. Proc. Nat. Acad. Sci. 82: 248-252, 1985.
[33834] 4850.

[33835] Christopoulos, G.; Perry, K. J.; Morfis, M.; Tilakaratne, N.; Gao,
[33836] Y.; Fraser, N. J.; Main, M. J.; Foord, S. M.; Sexton, P. M.: Multiple
[33837] amylin receptors arise from receptor activity-modifying protein interaction
[33838] with the calcitonin receptor gene product. Molec. Pharm. 56: 235-242,
[33839] 1999.

[33840] 4851.

[33841] Bourgeois, P.; Stoetzel, C.; Bolcato-Bellemin, A. L.; Mattei, M.
[33842] G.; Perrin-Schmitt, F.: The human H-twist gene is located at 7p21
[33843] and encodes a B-HLH protein that is 96% similar to its

murine M-twist
[33844] counterpart. Mammalian Genome 7: 915-917, 1996.

[33845] 4852.

[33846] Dollfus, H.; Biswas, P.; Kumaramanickavel, G.; Stoetzel, C.; Quillet,
[33847] R.; Biswas, J.; Lajeunie, E.; Renier, D.; Perrin-Schmitt, F.: Saethre-Chotzen
[33848] syndrome: notable intrafamilial phenotypic variability in a large
[33849] family with Q28X TWIST mutation. Am. J. Med. Genet. 109: 218-225,
[33850] 2002.

[33851] 4853.

[33852] Dollfus, H.; Kumaramanickavel, G.; Biswas, P.; Stoetzel, C.; Quillet,
[33853] R.; Denton, M.; Maw, M.; Perrin-Schmitt, F.: Identification of a
[33854] new TWIST mutation (7p21) with variable eyelid manifes-tations supports
[33855] locus homogeneity of BPES at 3q22. (Letter) J. Med. Genet. 58: 470-471,
[33856] 2001.

[33857] 4854.

[33858] El Ghouzzi, V.; Le Merrer, M.; Perrin-Schmitt, F.; Lajeunie, E.;

[33859] Benit, P.; Renier, D.; Bourgeois, P.; Bolcato-Bellemin, A.-L.; Munnich,

[33860] A.; Bonaventure, J.: Mutations of the TWIST gene in the Saethre-Chotzen

[33861] syndrome. Nature Genet. 15: 42-46, 1997.

[33862] 4855.

[33863] Howard, T. D.; Paznekas, W. A.; Green, E. D.; Chiang, L. C.; Ma,

[33864] N.; Ortiz De Luna, R. I.; Delgado, C. G.; Gonzalez-Ramos, M.; Kline,

[33865] A. D.; Jabs, E. W.: Mutations in TWIST, a basic helix-loop-helix

[33866] transcription factor, in Saethre-Chotzen syndrome. Nature Genet. 15:

[33867] 36-41, 1997.

[33868] 4856.

[33869] Maw, M.; Kar, B.; Biswas, J.; Biswas, P.; Nancarrow, D.; Denton,

[33870] M.; Bridges, R.; Kumaramanickavel, G.; Badrinath, S. S.: Linkage
[33871] of blepharophimosis syndrome in a large Indian pedigree to chromosome
[33872] 7p. Hum. Molec. Genet. 5: 2049-2054, 1996.

[33873] 4857.

[33874] Hagen, F. S.; Grant, F. J.; Kuijper, J. L.; Slaughter, C. A.; Moomaw,
[33875] C. R.; Orth, K.; O'Hara, P. J.; Munford, R. S.: Expression and characterization
[33876] of recombinant human acyloxyacyl hydrolase, a leukocyte enzyme that
[33877] deacylates bacterial lipopolysaccharides. Biochemistry 30: 8415-8423,
[33878] 1991.

[33879] 4858.

[33880] Whitmore, T. E.; Mathewes, S. L.; O'Hara, P. J.; Durnam, D. M.
[33881] : Chromosomal localization of the acyloxyacyl hydrolase (AOAH) gene
[33882] to 7p14-p12 using fluorescence in situ hybridization. Genomics 21:

[33883] 457-458, 1994.

[33884] 4859.

[33885] Morinaga, T.; Yasuda, H.; Hashimoto, T.; Higashio, K.; Tamaoki,

[33886] T.: A human alpha-fetoprotein enhancer-binding protein, ATBF1, contains

[33887] four homeodomains and seventeen zinc fingers. Molec. Cell. Biol. 11:

[33888] 6041-6049, 1991.

[33889] 4860.

[33890] Yamada, K.; Ma, D.; Miura, Y.; Ido, A.; Tamaoki, T.; Yoshida, M.

[33891] C.: Assignment of the ATBF1 transcription factor gene (Atbf1) to

[33892] mouse chromosome band 8E1 by in situ hybridization. Cytogenet. Cell

[33893] Genet. 75: 30-31, 1996.

[33894] 4861.

[33895] Yamada, K.; Mirua, Y.; Scheidl, T.; Yoshida, M. C.; Tamaoki, T.

[33896] : Assignment of the human ATBF1 transcription factor

gene to chromosome
[33897] 16q22.3-q23.1. Genomics 29: 552-553, 1995.

[33898] 4862.

[33899] Abbott, C. A.; Baker, E.; Sutherland, G. R.; McCaughan, G. W.:
[33900] Genomic organization, exact localization, and tissue expression of
[33901] the human CD26 (dipeptidyl peptidase IV) gene. Immunogenetics 40:
[33902] 331-338, 1994.

[33903] 4863.

[33904] Callebaut, C.; Krust, B.; Jacotot, E.; Hovanessian, A. G.: T cell
[33905] activation antigen, CD26, as a cofactor for entry of HIV in CD4+ cells. Science 262:
[33906] 2045-2050, 1993.

[33907] 4864.

[33908] Darmoul, D.; Lacasa, M.; Chantret, I.; Swallow, D. M.; Trugnan,
[33909] G.: Isolation of a cDNA probe for the human intestinal dipeptidylpeptidase

[33910] IV and assignment of the gene locus DPP4 to chromosome 2. Ann. Hum.

[33911] Genet. 54: 191-197, 1990.

[33912] 4865.

[33913] Herbschleb-Voogt, E.; Grzeschik, K.-H.; Pearson, P. L.; Meera Khan,

[33914] P.: Assignment of adenosine deaminase complexing protein (ADCP) gene(s)

[33915] to human chromosome 2 in rodent-human somatic cell hybrids. Hum.

[33916] Genet. 59: 317-323, 1981.

[33917] 4866.

[33918] Kameoka, J.; Tanaka, T.; Nojima, Y.; Schlossman, S. F.; Morimoto,

[33919] C.: Direct association of adenosine deaminase with a T cell activation

[33920] antigen, CD26. Science 261: 466-469, 1993.

[33921] 4867.

[33922] Koch, G. A.; Shows, T. B.: Genes on human chromosomes 2 and 6

[33923] are required for expression of the adenosine deaminase

complexing
[33924] protein (ADCP) in human-mouse somatic cell hybrids. (Abstract) Cytogenet.
[33925] Cell Genet. 25: 174, 1979.

[33926] 4868.

[33927] Marguet, D.; Baggio, L.; Kobayashi, T.; Bernard, A.-M.; Pierres,
[33928] M.; Nielsen, P. F.; Ribel, U.; Watanabe, T.; Drucker, D. J.; Wagtmann,
[33929] N.: Enhanced insulin secretion and improved glucose tol-erance in
[33930] mice lacking CD26. Proc. Nat. Acad. Sci. 97: 6874-6879, 2000.
[33931] 4869.

[33932] Mathew, S.; Morrison, M. E.; Murty, V. V. V. S.; Houghton, A. N.;
[33933] Chaganti, R. S. K.: Assignment of the DPP4 gene encoding adenosine
[33934] deaminase binding protein (CD26/dipeptidylpeptidase IV) to 2q23. Genomics 22:
[33935] 211-212, 1994.

[33936] 4870.

[33937] Misumi, Y.; Hayashi, Y.; Arakawa, F.; Ikehara, Y.: Molecular

[33938] cloning and sequence analysis of human dipeptidyl pepti-
dase IV, a

[33939] serine proteinase on the cell surface. Biochim. Biophys.
Acta 1131:

[33940] 333-336, 1992.

[33941] 4871.

[33942] Morrison, M. E.; Vijayasaradhi, S.; Engelstein, D.; Albino,
A.

[33943] P.; Houghton, A. N.: A marker for neoplastic progression
of human

[33944] melanocytes is a cell surface ectopeptidase. J. Exp. Med.
117: 1135-1143,

[33945] 1993.

[33946] 4872.

[33947] Van Cong, N.; Weil, D.; Gross, M.-S.; Foubert, C.; Jami, J.;
Frezal,

[33948] J.: Controle genetique et epigenetique de l'expression de
l'adenosine

[33949] deaminase. Analyse des cellules humaines et hybrides
homme-rongeur. Ann.

[33950] Genet. 24: 141-147, 1981.

[33951] 4873.

[33952] Bundey, S.: Recent views on genetic factors in retinoblas-
toma.
[33953] (Abstract) J. Med. Genet. 17: 386-387, 1980.

[33954] 4874.

[33955] Bundey, S.; Morten, J. E. N.: An unusual pedigree with
retinoblastoma.
[33956] Does it shed light on the delayed mutation and host resis-
tance theories? Hum.
[33957] Genet. 59: 434-436, 1981.

[33958] 4875.

[33959] Cance, W. G.; Brennan, M. F.; Dudas, M. E.; Huang, C.-M.;
Cordon-Cardo,
[33960] C.: Altered expression of the retinoblastoma gene product
in human
[33961] sarcomas. New Eng. J. Med. 323: 1457-1462, 1990.

[33962] 4876.

[33963] Canning, S.; Dryja, T. P.: Short, direct repeats at the
breakpoints

[33964] of deletions of the retinoblastoma gene. Proc. Nat. Acad. Sci. 86:
[33965] 5044-5048, 1989.

[33966] 4877.

[33967] Carlson, E. A.; Desnick, R. J.: Mutational mosaicism and genetic
[33968] counseling in retinoblastoma. Am. J. Med. Genet. 4: 365-381, 1979.
[33969] 4878.

[33970] Cavenee, W. K.: The genetic basis of neoplasia: the retinoblastoma
[33971] paradigm. Trends Genet. 2: 299-300, 1986.

[33972] 4879.

[33973] Cavenee, W. K.; Dryja, T. P.; Phillips, R. A.; Benedict, W. F.;

[33974] Godbout, R.; Gallie, B. L.; Murphree, A. L.; Strong, L. C.; White,
[33975] R. L.: Expression of recessive alleles by chromosomal mechanisms
[33976] in retinoblastoma. Nature 305: 779-784, 1983.

[33977] 4880.

[33978] Cavenee, W. K.; Hansen, M. F.; Nordenskjold, M.; Kock, E.; Maumenee,
[33979] I.; Squire, J. A.; Phillips, R. A.; Gallie, B. L.: Genetic origin
[33980] of mutations predisposing to retinoblastoma. Science 228: 501-503,
[33981] 1985.
[33982] 4881.
[33983] Cavenee, W. K.; Murphree, A. L.; Shull, M. M.; Benedict, W. F.;
[33984] Sparkes, R. S.; Kock, E.; Nordenskjold, M.: Prediction of familial
[33985] predisposition to retinoblastoma. New Eng. J. Med. 314: 1201-1207,
[33986] 1986.
[33987] 4882.
[33988] Chano, T.; Ikegawa, S.; Kontani, K.; Okabe, H.; Baldini, N.; Saeki,
[33989] Y.: Identification of RB1CC1, a novel human gene that can induce
[33990] RB1 in various human cells. Oncogene 21: 1295-1298, 2002.

[33991] 4883.

[33992] Chauveinc, L.; Mosseri, V.; Quintana, E.; Desjardins, L.; Schlienger,

[33993] P.; Doz, F.; Dutrillaux, B.: Osteosarcoma following retinoblastoma:

[33994] age at onset and latency period. Ophthalmic Genet. 22: 77-88, 2001.

[33995] 4884.

[33996] Chen, P.-L.; Scully, P.; Shew, J.-Y.; Wang, J. Y. J.; Lee, W.-H.

[33997] : Phosphorylation of the retinoblastoma gene product is modulated

[33998] during the cell cycle and cellular differentiation. Cell 58: 1193-1198,

[33999] 1989.

[34000] 4885.

[34001] Connolly, M. J.; Payne, R. H.; Johnson, G.; Gallie, B. L.; Allderdice,

[34002] P. W.; Marshall, W. H.; Lawton, R. D.: Familial, EsD-linked, retinoblastoma

[34003] with reduced penetrance and variable expressivity. Hum. Genet. 65:

[34004] 122-124, 1983.

[34005] 4886.

[34006] Cowell, J. K.; Bia, B.: A novel missense mutation in patients

[34007] from a retinoblastoma pedigree showing only mild ex-
pression of the

[34008] tumor phenotype. Oncogene 16: 3211-3213, 1998.

[34009] 4887.

[34010] Cowell, J. K.; Rutland, P.; Hungerford, J.; Jay, M.: Deletion

[34011] of chromosome region 13q14 is transmissible and does
not always predispose

[34012] to retinoblastoma. Hum. Genet. 80: 43-45, 1988.

[34013] 4888.

[34014] Cowell, J. K.; Rutland, P.; Jay, M.; Hungerford, J.: Deletions

[34015] of the esterase D locus from a survey of 200 retinoblas-
toma patients. Hum.

[34016] Genet. 72: 164-167, 1986.

[34017] 4889.

[34018] Cowell, J. K.; Smith, T.; Bia, B.: Frequent constitutional C

[34019] to T mutations in CGA-arginine codons in the RB1 gene

produce premature
[34020] stop codons in patients with bilateral (hereditary) retinoblastoma. Europ.
[34021] J. Hum. Genet. 2: 281-290, 1994.

[34022] 4890.

[34023] Dahiya, A.; Wong, S.; Gonzalo, S.; Gavin, M.; Dean, D. C.: Linking
[34024] the Rb and Polycomb pathways. Molec. Cell 8: 557-568, 2001.
[34025] 4891.

[34026] Davison, E. V.; Gibbons, B.; Aherne, G. E. S.; Roberts, D. F.

[34027] : Chromosomes in retinoblastoma patients. Clin. Genet. 15: 505-508,
[34028] 1979.

[34029] 4892.

[34030] DeCaprio, J. A.; Ludlow, J. W.; Lynch, D.; Furukawa, Y.; Griffin,
[34031] J.; Piwnica-Worms, H.; Huang, C.-M.; Livingston, D. M.: The product
[34032] of the retinoblastoma susceptibility gene has properties of a cell

[34033] cycle regulatory element. Cell 58: 1085-1095, 1989.

[34034] 4893.

[34035] de Grouchy, J.; Turleau, C.; Cabanis, M. O.; Richardet, J. M.

[34036] : Retinoblastome et deletion intercalaire du chromosome 13. Arch.

[34037] Franc. Pediat. 37: 531-535, 1980.

[34038] 4894.

[34039] Dryja, T.; Cavenee, W.; Epstein, J.; Rapaport, J.; Goorin, A.;

[34040] Koufos, A.: Chromosome 13 homozygosity in osteogenic sarcoma without

[34041] retinoblastoma. (Abstract) Am. J. Hum. Genet. 36: 28S, 1984.

[34042] 4895.

[34043] Dryja, T. P.; Bruns, G. A. P.; Gallie, B.; Petersen, R.; Green,

[34044] W.; Rapaport, J. M.; Albert, D. M.; Gerald, P. S.: Low inci-dence

[34045] of deletion of the esterase D locus in retinoblastoma pa-tients. Hum.

[34046] Genet. 64: 151-155, 1983.

[34047] 4896.

[34048] Dryja, T. P.; Cavenee, W.; White, R.; Rapaport, J. M.; Petersen,
[34049] R.; Albert, D. M.; Bruns, G. A. P.: Homozygosity of chromosome 13
[34050] in retinoblastoma. New Eng. J. Med. 310: 550-553, 1984.

[34051] 4897.

[34052] Dryja, T. P.; Friend, S.; Weinberg, R. A.: Isolation of a cDNA
[34053] fragment derived from human retina mRNA which detects a locus within
[34054] 13q14 often deleted in retinoblastomas. (Abstract) Am. J. Hum. Genet. 39:
[34055] A29, 1986.

[34056] 4898.

[34057] Dryja, T. P.; Mukai, S.; Petersen, R.; Rapaport, J. M.; Walton,
[34058] D.; Yandell, D. W.: Parental origin of mutations of the retinoblastoma
[34059] gene. Nature 339: 556-558, 1989.

[34060] 4899.

[34061] Dryja, T. P.; Mukai, S.; Rapaport, J. M.; Yandell, D. W.: Parental

[34062] origin of mutations of the retinoblastoma gene. (Abstract) Am. J.

[34063] Hum. Genet. 45 (suppl.): A19, 1989.

[34064] 4900.

[34065] Messer Peters, P. G.; Kamarck, M. E.; Hemler, M. E.; Strominger,

[34066] J. L.; Ruddle, F. H.: Genetic and biochemical characterization of

[34067] a human surface determinant on somatic cell hybrids: the 4F2 antigen. Somat.

[34068] Cell Genet. 8: 825-834, 1982.

[34069] 4901.

[34070] Markert, C. L.; Silvers, W. K.: The effects of genotype and cell

[34071] environment on melanoblast differentiation in the house mouse. Genetics 41:

[34072] 429-450, 1956.

[34073] 4902.

[34074] Koch, M. C.; Baumbach, K.; George, A. L.; Ricker, K.: Paramyotonia

[34075] congenita without paralysis on exposure to cold: a novel

mutation
[34076] in the SCN4A gene (val1293ile). Neuroreport 6:
2001-2004, 1995.
[34077] 4903.

[34078] Bell, L. R.; Maine, E. M.; Schedl, P.; Cline, T. W.: Sex-lethal,

[34079] a Drosophila sex determination switch gene, exhibits sex-
specific
[34080] RNA splicing and sequence similarity to RNA binding pro-
teins. Cell 55:
[34081] 1037-1046, 1988.

[34082] 4904.

[34083] Fletcher, C. F.; Okano, H. J.; Gilbert, D. J.; Yang, Y.; Yang,

[34084] C.; Copeland, N. G.; Jenkins, N. A.; Darnell, R. B.: Mouse
chromosomal
[34085] locations of nine genes encoding homologs of human
paraneoplastic
[34086] neurologic disorder antigens. Genomics 45: 313-319,
1997.
[34087] 4905.

[34088] King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.:

[34089] Mammalian homologs of Drosophila elav localized to a

neuronal subset
[34090] can bind in vitro to the 3-prime UTR of mRNA encoding the Id transcriptional
[34091] repressor. J. Neurosci. 14: 1943-1952, 1994.

[34092] 4906.

[34093] Levine, T. D.; Gao, F.; King, P. H.; Andrews, L. G.; Keene, J.

[34094] D.: He1-N1: an autoimmune RNA-binding protein with specificity for
[34095] 3-prime uridylate-rich untranslated regions of growth factor mRNAs. Molec.
[34096] Cell. Biol. 13: 3494-3504, 1993.

[34097] 4907.

[34098] Muresu, R.; Baldini, A.; Gress, T.; Posner, J. B.; Furneaux, H.
[34099] M.; Siniscalco, M.: Mapping of the gene coding for a para-neoplastic
[34100] encephalomyelitis antigen (HuD) to human chromosome site 1p34. Cytogenet.
[34101] Cell Genet. 65: 177-178, 1994.

[34102] 4908.

[34103] Robinow, S.; Campos, A. R.; Yao, K.-M.; White, K.: The elav

gene
[34104] product of Drosophila, required in neurons, has three RNP consensus
[34105] motifs. Science 242: 1570-1572, 1988.

[34106] 4909.

[34107] Szabo, A.; Dalmau, J.; Manley, G.; Rosenfeld, M.; Wong, E.; Henson,
[34108] J.; Posner, J. B.; Furneaux, H. M.: HuD, a paraneoplastic encephalomyelitis
[34109] antigen, contains RNA-binding domains and is homologous to elav and
[34110] sex-lethal. Cell 67: 325-333, 1991.

[34111] 4910.

[34112] Arnold, A.; Kim, H. G.; Gaz, R. D.; Eddy, R. L.; Fukushima, Y.;
[34113] Byers, M. G.; Shows, T. B.; Kronenberg, H. M.: Molecular cloning
[34114] and chromosomal mapping of DNA rearranged with the parathyroid hormone
[34115] gene in a parathyroid adenoma. J. Clin. Invest. 83: 2034-2040, 1989.
[34116] 4911.

[34117] Mehta, A. D.; Rock, R. S.; Rief, M.; Spudich, J. A.; Mooseker,
[34118] M. S.; Cheney, R. E.: Myosin-V is a processive actin-based motor. Nature 400:
[34119] 590-593, 1999.

[34120] 4912.

[34121] Mercer, J. A.; Seperack, P. K.; Strobel, M. C.; Copeland, N. G.;
[34122] Jenkins, N. A.: Novel myosin heavy chain encoded by murine dilute
[34123] coat colour locus. Nature 349: 709-713, 1991.

[34124] 4913.

[34125] Moore, K. J.; Testa, J. R.; Francke, U.; Milatovich, A.; Copeland,
[34126] N. G.; Jenkins, N. A.: Cloning and regional assignment of the human
[34127] myosin heavy chain 12 (MYH12) gene to chromosome band 15q21. Cytogenet.
[34128] Cell Genet. 69: 53-58, 1995.

[34129] 4914.

[34130] Pastural, E.; Barrat, F. J.; Dufourcq-Lagelouse, R.; Certain,

[34131] S.; Sanal, O.; Jabado, N.; Seger, R.; Griscelli, C.; Fischer, A.;

[34132] de Saint Basile, G.: Griscelli disease maps to chromosome 15q21 and

[34133] is associated with mutations in the myosin-Va gene. Nature Genet. 16:

[34134] 289-292, 1997. Note: Erratum: Nature Genet. 23: 373 only, 1999.

[34135] 4915.

[34136] Rief, M.; Rock, R. S.; Mehta, A. D.; Mooseker, M. S.; Cheney,

[34137] R. E.; Spudich, J. A.: Myosin-V stepping kinetics: a molecular model

[34138] for processivity. Proc. Nat. Acad. Sci. 97: 9482-9486, 2000.

[34139] 4916.

[34140] Russell, E. S.: A quantitative histological study of the pigment

[34141] found in the coat-color mutants of the house mouse. IV. The nature

[34142] of the effects of genic substitution in five major allelic series. Genetics 34:

[34143] 146-166, 1949.

[34144] 4917.

[34145] Kitada, T.; Asakawa, S.; Minoshima, S.; Mizuno, Y.;
Shimizu, N.

[34146] : Molecular cloning, gene expression, and identification of
a splicing

[34147] variant of the mouse parkin gene. Mammalian Genome
11: 417-421,

[34148] 2000.

[34149] 4918.

[34150] Lucking, C. B.; Abbas, N.; Durr, A.; Bonifati, V.; Bonnet,
A.-M.;

[34151] de Broucker, T.; De Michele, G.; Wood, N. W.; Agid, Y.;
Brice, A.

[34152] : Homozygous deletions in parkin gene in European and
North African

[34153] families with autosomal recessive juvenile parkinsonism.
Lancet 352:

[34154] 1355-1356, 1998.

[34155] 4919.

[34156] Musahl, C.; Schulte, D.; Burkhart, R.; Knippers, R.: A hu-
man homologue

[34157] of the yeast replication protein Cdc21: interactions with other Mcm

[34158] proteins. Europ. J. Biochem. 230: 1096-1101, 1995.

[34159] 4920.

[34160] Anderson, D. M.; Maraskovsky, E.; Billingsley, W. L.; Dougall,

[34161] W. C.; Tometsko, M. E.; Roux, E. R.; Teepe, M. C.; DuBose, R. F.;

[34162] Cosman, D.; Galibert, L.: A homologue of the TNF receptor and its

[34163] ligand enhance T-cell growth and dendritic-cell function. Nature 390:

[34164] 175-179, 1997.

[34165] 4921.

[34166] Nykjaer, A.; Fyfe, J. C.; Kozyraki, R.; Leheste, J.-R.; Jacob-sen,

[34167] C.; Nielsen, M. S.; Verroust, P. J.; Aminoff, M.; de la Chapelle,

[34168] A.; Moestrup, S. K.; Ray, R.; Gliemann, J.; Willnow, T. E.; Christensen,

[34169] E. I.: Cubilin dysfunction causes abnormal metabolism of the steroid

[34170] hormone 25(OH) vitamin D3. Proc. Nat. Acad. Sci. 98: 13895-13900,
[34171] 2001.

[34172] 4922.

[34173] Udell, C. M.; Lee, S. K.; Davey, S.: HRAD1 and MRAD1 encode mammalian
[34174] homologues of the fission yeast rad1+ cell cycle checkpoint control
[34175] gene. Nucleic Acids Res. 26: 3971-3976, 1998.

[34176] 4923.

[34177] Carvajal, J. J.; Pook, M. A.; dos Santos, M.; Doudney, K.; Hillermann,
[34178] R.; Minogue, S.; Williamson, R.; Hsuan, J. J.; Chamberlain, S.: The
[34179] Friedreich's ataxia gene encodes a novel phosphatidylinositol-4-phosphate
[34180] 5-kinase. Nature Genet. 14: 157-162, 1996.

[34181] 4924.

[34182] Carvajal, J. J.; Pook, M. A.; Doudney, K.; Hillermann, R.; Wilkes,
[34183] D.; Al-Mahdawi, S.; Williamson, R.; Chamberlain, S.:

Friedreich's
[34184] ataxia: a defect in signal transduction? Hum. Molec. Genet. 4: 1411-1419,
[34185] 1995.

[34186] 4925.

[34187] Pook, M. A.; Carvajal, J. J; Doudney, K.; Hillermann, R.; Chamberlain,
[34188] S.: Exon-intron structure of a 2.7-kb transcript of the STM7 gene
[34189] with phosphatidylinositol-4-phosphate 5-kinase activity. Genomics 42:
[34190] 170-172, 1997.

[34191] 4926.

[34192] Carfi, A.; Willis, S. H.; Whitbeck, J. C.; Krummenacher, C.; Cohen,
[34193] G. H.; Eisenberg, R. J.; Wiley, D. C.: Herpes simplex virus glycoprotein
[34194] D bound to the human receptor HveA. Molec. Cell 8: 169-179, 2001.
[34195] 4927.

[34196] Hsu, H.; Solovyev, I.; Colombero, A.; Elliott, R.; Kelley, M.;

[34197] Boyle, W. J.: ATAR, a novel tumor necrosis factor receptor family

[34198] member, signals through TRAF2 and TRAF5. J. Biol. Chem. 272: 13471-13474,

[34199] 1997.

[34200] 4928.

[34201] Marsters, S. A.; Ayres, T. M.; Skubatch, M.; Gray, C. L.; Rothe,

[34202] M.; Ashkenazi, A.: Herpesvirus entry mediator, a member of the tumor

[34203] necrosis factor receptor (TNFR) family, interacts with members of

[34204] the TNFR-associated factor family and activates the transcription

[34205] factors NF-kappa-B and AP-1. J. Biol. Chem. 272: 14029-14032, 1997.

[34206] 4929.

[34207] Montgomery, R. I.; Warner, M. S.; Lum, B. J.; Spear, P. G.: Herpes

[34208] simplex virus-1 entry into cells mediated by a novel member of the

[34209] TNF/NGF receptor family. Cell 87: 427-436, 1996.

[34210] 4930.

[34211] Kwon, B. S.; Tan, K. B.; Ni, J.; Oh, K.-O.; Lee, Z. H.; Kim, K.

[34212] K.; Kim, Y.-J.; Wang, S.; Gentz, R.; Yu, G.-L.; Harrop, J.;
Lyn, S.

[34213] D.; Silverman, C.; Porter, T. G.; Truneh, A.; Young, P. R.: A
newly

[34214] identified member of the tumor necrosis factor receptor
superfamily

[34215] with a wide tissue distribution and involvement in lym-
phocyte activation. J.

[34216] Biol. Chem. 272: 14272-14276, 1997.

[34217] 4931.

[34218] Morel, Y.; Schiano de Colella, J.-M.; Harrop, J.; Deen, K. C.;

[34219] Holmes, S. D.; Wattam, T. A.; Khandekar, S. S.; Truneh, A.;
Sweet,

[34220] R. W.; Gastaut, J.-A.; Olive, D.; Costello, R. T.: Reciprocal
expression

[34221] of the TNF family receptor herpes virus entry mediator
and its ligand

[34222] LIGHT on activated T cells: LIGHT down-regulates its own
receptor. J.

[34223] Immun. 165: 4397-4404, 2000.

[34224] 4932.

[34225] Guan, K.-L.; Butch, E.: Isolation and characterization of a novel

[34226] dual specific phosphatase, HVH2, which selectively de-phosphorylates

[34227] the mitogen-activated protein kinase. J. Biol. Chem. 270: 7197-7203,

[34228] 1995.

[34229] 4933.

[34230] Smith, A.; Price, C.; Cullen, M.; Muda, M.; King, A.; Ozanne, B.;

[34231] Arkinstall, S.; Ashworth, A.: Chromosomal localization of three human

[34232] dual specificity phosphatase genes (DUSP4, DUSP6, and DUSP7). Genomics 42:

[34233] 524-527, 1997.

[34234] 4934.

[34235] Furukawa, T.; Yatsuoka, T.; Youssef, E. M.; Abe, T.; Yokoyama,

[34236] T.; Fukushige, S.; Soeda, E.; Hoshi, M.; Hayashi, Y.; Suna-

mura, M.;

[34237] Kobari, M.; Horii, A.: Genomic analysis of DUSP6, a dual specificity

[34238] MAP kinase phosphatase, in pancreatic cancer. Cytogenet. Cell Genet. 82:

[34239] 156-159, 1998.

[34240] 4935.

[34241] Groom, L. A.; Sneddon, A. A.; Alessi, D. R.; Dowd, S.; Keyse, S.

[34242] M.: Differential regulation of the MAP, SAP and RK/p38 kinases by

[34243] Pyst1, a novel cytosolic dual-specificity phosphatase. EMBO J. 15:

[34244] 3621-3632, 1996.

[34245] 4936.

[34246] Muda, M.; Boschert, U.; Dickinson, R.; Martinou, J.-C.; Martinou,

[34247] I.; Camps, M.; Schlegel, W.; Arkinstall, S.: MKP-3, a novel cytosolic

[34248] protein-tyrosine phosphatase that exemplifies a new class of mitogen-activated

[34249] protein kinase phosphatase. J. Biol. Chem. 271:

4319-4326, 1996.

[34250] 4937.

[34251] Smith, A.; Price, C.; Cullen, M.; Muda, M.; King, A.;
Ozanne, B.;

[34252] Arkinstall, S.; Ahsworth, A.: Chromosomal localization of
three human

[34253] dual specificity phosphatase genes (DUSP4, DUSP6, and
DUSP7). Genomics 42:

[34254] 524-527, 1997.

[34255] 4938.

[34256] Nishihira, J.; Fujinaga, M.; Kuriyama, T.; Suzuki, M.; Sugi-
moto,

[34257] H.; Nakagawa, A.; Tanaka, I.; Sakai, M.: Molecular cloning
of human

[34258] D-dopachrome tautomerase cDNA: N-terminal proline is
essential for

[34259] enzyme activation. Biochem. Biophys. Res. Commun. 243:
538-544,

[34260] 1998.

[34261] 4939.

[34262] Blouin, J.-L.; Sail, G. D.; Guipponi, M.; Rossier, C.; Pap-

pasavas,

[34263] M.-P.; Antonarakis, S. E.: Isolation of the human BACH1 transcription

[34264] regulator gene, which maps to chromosome 21q22.1. Hum. Genet. 102:

[34265] 282-288, 1998.

[34266] 4940.

[34267] Strehl, S.; Glatt, K.; Liu, Q. M.; Glatt, H.; Lalande, M.: Characterization

[34268] of two novel protocadherins (PCDH8 and PCDH9) localized on human chromosome

[34269] 13 and mouse chromosome 14. Genomics 53: 81-89, 1998.

[34270] 4941.

[34271] Ohbayashi, N.; Hoshikawa, M.; Kimura, S.; Yamasaki, M.; Fukui,

[34272] S.; Itoh, N.: Structure and expression of the mRNA encoding a novel

[34273] fibroblast growth factor, FGF-18. J. Biol. Chem. 273: 18161-18164,

[34274] 1998.

[34275] 4942.

[34276] Jackson, A.; Panayiotidis, P.; Foroni, L.: The human homologue
[34277] of the Drosophila tailless gene (TLX): characterization and mapping
[34278] to a region of common deletion in human lymphoid leukemia on chromosome
[34279] 6q21. Genomics 50: 34-43, 1998.

[34280] 4943.

[34281] Monaghan, A. P.; Bock, D.; Gass, P.; Schwager, A.; Wolfer, D. P.;
[34282] Lipp, H.-P.; Schutz, G.: Defective limbic system in mice lacking
[34283] the tailless gene. Nature 390: 515-517, 1997.

[34284] 4944.

[34285] Yu, R. T.; McKeown, M.; Evans, R. M.; Umesono, K.: Relationship
[34286] between Drosophila gap gene tailless and a vertebrate nuclear receptor
[34287] Tlx. Nature 370: 375-379, 1994.

[34288] 4945.

[34289] Plougastel, B.; Trowsdale, J.: Cloning of NKG2-F, a new

member

[34290] of the NKG2 family of human natural killer cell receptor genes. Europ.

[34291] J. Immun. 27: 2835-2839, 1997.

[34292] 4946.

[34293] Sutherland, C. L.; Chalupny, N. J.; Schooley, K.; Vanden-Bos, T.;

[34294] Kubin, M.; Cosman, D.: UL16-binding proteins, novel MHC class I-related

[34295] proteins, bind to NKG2D and activate multiple signaling pathways in

[34296] primary NK cells. J. Immun. 168: 671-679, 2002.

[34297] 4947.

[34298] Wu, J.; Song, Y.; Bakker, A. B. H.; Bauer, S.; Spies, T.; Lanier,

[34299] L. L.; Phillips, J. H.: An activating immunoreceptor com-plex formed

[34300] by NKG2D and DAP10. Science 285: 730-732, 1999.

[34301] 4948.

[34302] Chang, C.; Rodriguez, A.; Carretero, M.; Lopez-Botet, M.; Phillips,

[34303] J. H.; Lanier, L. L.: Molecular characterization of human CD94: a

[34304] type II membrane glycoprotein related to the C-type lectin superfamily. Europ.

[34305] J. Immun. 25: 2433-2437, 1995.

[34306] 4949.

[34307] Lazetic, S.; Chang, C.; Houchins, J. P.; Lanier, L. L.; Phillips,

[34308] J. H.: Human natural killer cell receptors involved in MHC class

[34309] I recognition are disulfide-linked heterodimers of CD94 and NKG2 subunits. J.

[34310] Immun. 157: 4741-4745, 1996.

[34311] 4950.

[34312] Rodriguez, A.; Carretero, M.; Glienke, J.; Bellon, T.; Ramirez,

[34313] A.; Lehrach, H.; Francis, F.; Lopez-Botet, M.: Structure of the human

[34314] CD94 C-type lectin gene. Immunogenetics 47: 305-309, 1998.

[34315] 4951.

[34316] Volkmer, E.; Karnitz, L. M.: Human homologs of

Schizosaccharomyces

[34317] pombe Rad1, Hus1, and Rad9 form a DNA damage-responsive protein complex. J.

[34318] Biol. Chem. 274: 567-570, 1999.

[34319] 4952.

[34320] Brandner, J. M.; Reidenbach, S.; Franke, W. W.: Evidence that

[34321] 'pinin,' reportedly a differentiation-specific desmosomal protein,

[34322] is actually a widespread nuclear protein. Differentiation 62: 119-127,

[34323] 1997.

[34324] 4953.

[34325] Brandner, J. M.; Reidenbach, S.; Kuhn, C.; Franke, W. W.: Identification

[34326] and characterization of a novel kind of nuclear protein occurring

[34327] free in nucleoplasm and in ribonucleoprotein structures of the 'speckle'

[34328] type. Europ. J. Cell Biol. 75: 295-308, 1998.

[34329] 4954.

[34330] Degen, W. G. J.; Agterbos, M. A.; Muyrers, J. P. P.; Bloemers,
[34331] H. P. J.; Swart, G. W. M.: memA/DRS, a putative mediator of multiprotein
[34332] complexes, is overexpressed in the metastasizing human melanoma cell
[34333] lines BLM and MV3. Biochim. Biophys. Acta 1444: 384-394, 1999.
[34334] 4955.

[34335] Ouyang, P.; Sugrue, S. P.: Characterization of pinin, a novel
[34336] protein associated with the desmosome-intermediate filament complex. J.
[34337] Cell Biol. 135: 1027-1042, 1996.

[34338] 4956.

[34339] Puente, X. S.; Lopez-Otin, C.: Cloning and expression analysis
[34340] of a novel human serine hydrolase with sequence similarity to prokaryotic
[34341] enzymes involved in the degradation of aromatic compounds. J. Biol.
[34342] Chem. 270: 12926-12932, 1995.

[34343] 4957.

[34344] Puente, X. S.; Pendas, A. M.; Lopez-Otin, C.: Structural characterization
[34345] and chromosomal localization of the gene encoding human biphenyl hydrolase-related
[34346] protein (BPHL). Genomics 51: 459-462, 1998.

[34347] 4958.

[34348] Janssen, J. W. G.; Schleithoff, L.; Bartram, C. R.; Schulz, A.

[34349] S.: An oncogenic fusion product of the phosphatidylinositol 3-kinase
[34350] p85-beta subunit and HUMORF8, a putative deubiquitinating enzyme. Oncogene 16:
[34351] 1767-1772, 1998.

[34352] 4959.

[34353] Chadwick, B. P.; Williamson, J.; Sheer, D.; Frischauf, A.-M.:

[34354] cDNA cloning and chromosomal mapping of a mouse gene with homology
[34355] to NTPases. Mammalian Genome 9: 162-164, 1998.

[34356] 4960.

[34357] Yeung, G.; Mulero, J. J.; McGowan, D. W.; Bajwa, S. S.;

Ford, J.

[34358] E.: CD39L2, a gene encoding a human nucleoside diphosphatase, predominantly

[34359] expressed in the heart. Biochemistry 39: 12916-12923, 2000.

[34360] 4961.

[34361] Mulero, J. J.; Yeung, G.; Nelken, S. T.; Bright, J. M.; McGowan,

[34362] D. W.; Ford, J. E.: Biochemical characterization of CD39L4. Biochemistry 39:

[34363] 12924-12928, 2000.

[34364] 4962.

[34365] Mulero, J. J.; Yeung, G.; Nelken, S. T.; Ford, J. E.: CD39-L4

[34366] is a secreted human apyrase, specific for the hydrolysis of nucleoside

[34367] diphosphates. J. Biol. Chem. 274: 20064-20067, 1999.

[34368] 4963.

[34369] Ghaedi, K.; Honsho, M.; Shimozawa, N.; Suzuki, Y.; Kondo, N.; Fujiki,

[34370] Y.: PEX3 is the causal gene responsible for peroxisome membrane assembly-defective

[34371] Zellweger syndrome of complementation group G. Am. J. Hum. Genet. 67:
[34372] 976-981, 2000.

[34373] 4964.

[34374] Kammerer, S.; Holzinger, A.; Welsch, U.; Roscher, A. A.: Cloning
[34375] and characterization of the gene encoding the human peroxisomal assembly
[34376] protein Pex3p. FEBS Lett. 429: 53-60, 1998.

[34377] 4965.

[34378] Muntau, A. C.; Holzinger, A.; Mayerhofer, P. U.; Gartner, J.; Roscher,
[34379] A. A.; Kammerer, S.: The human PEX3 gene encoding a peroxisomal assembly
[34380] protein: genomic organization, positional mapping, and mutation analysis
[34381] in candidate phenotypes. Biochem. Biophys. Res. Com-
mun. 268: 704-710,
[34382] 2000.

[34383] 4966.

[34384] Muntau, A. C.; Mayerhofer, P. U.; Paton, B. C.; Kammerer,

S.; Roscher,
[34385] A. A.: Defective peroxisome membrane synthesis due to mutations in
[34386] human PEX3 causes Zellweger syndrome, complementation group G. Am.
[34387] J. Hum. Genet. 67: 967-975, 2000.

[34388] 4967.

[34389] Seetharam, B.; Christensen, E. I.; Moestrup, S. K.; Hammond, T.
[34390] G.; Verroust, P. J.: Identification of rat yolk sac target protein
[34391] of teratogenic antibodies, gp280, as intrinsic factor-cobalamin receptor. J.
[34392] Clin. Invest. 99: 2317-2322, 1997.

[34393] 4968.

[34394] Xu, D.; Kozyraki, R.; Newman, T. C.; Fyfe, J. C.: Genetic evidence
[34395] of an accessory activity required specifically for cubilin brush-border
[34396] expression and intrinsic factor-cobalamin absorption. Blood 94:
[34397] 3604-3606, 1999.

[34398] 4969.

[34399] Doherty, M. J.; Young, P. R.; Cohen, P. T. W.: Amino acid sequence

[34400] of a novel protein phosphatase 1 binding protein (R5) which is related

[34401] to the liver- and muscle-specific glycogen binding sub-units of protein

[34402] phosphatase 1. FEBS Lett. 399: 339-343, 1996.

[34403] 4970.

[34404] Seroussi, E.; Pan, H.-Q.; Kedra, D.; Roe, B. A.; Dumanski, J. P.

[34405] : Characterization of the human NIPSNAP1 gene from 22q12: a member

[34406] of a novel gene family. Gene 212: 13-20, 1998.

[34407] 4971.

[34408] Holzinger, A.; Kammerer, S.; Roscher, A. A.: Primary structure

[34409] of human PMP69, a putative peroxisomal ABC-transporter. Biochem.

[34410] Biophys. Res. Commun. 237: 152-157, 1997.

[34411] 4972.

[34412] Holzinger, A.; Roscher, A. A.; Landgraf, P.; Lichtner, P.; Kammerer,

[34413] S.: Genomic organization and chromosomal localization of the human

[34414] peroxisomal membrane protein-1-like protein (PXMP1-L) gene encoding

[34415] a peroxisomal ABC transporter. FEBS Lett. 426: 238-242, 1998.

[34416] 4973.

[34417] Shani, N.; Jimenez-Sanchez, G.; Steel, G.; Dean, M.; Valle, D.

[34418] : Identification of a fourth half ABC transporter in the human peroxisomal

[34419] membrane. Hum. Molec. Genet. 6: 1925-1931, 1997.

[34420] 4974.

[34421] Wang, H.; Frelin, L.; Pevsner, J.: Human syntaxin 7: a Pep12p/Vps6p

[34422] homologue implicated in vesicle trafficking to lysosomes. Gene 199:

[34423] 39-48, 1997.

[34424] 4975.

[34425] Fink, M.; Duprat, F.; Lesage, F.; Reyes, R.; Romey, G.; Heurteaux,
[34426] C.; Lazdunski, M.: Cloning, functional expression and brain localization
[34427] of a novel unconventional outward rectifier K+ channel. EMBO J. 15:
[34428] 6854-6862, 1996.

[34429] 4976.

[34430] Lesage, F.; Lazdunski, M.: Mapping of human potassium channel
[34431] genes TREK-1 (KCNK2) and TASK (KCNK3) to chromosomes 1q41 and 2p23. Genomics 51:
[34432] 478-479, 1998.

[34433] 4977.

[34434] Duprat, F.; Lesage, F.; Fink, M.; Reyes, R.; Heurteaux, C.; Lazdunski,
[34435] M.: TASK, a human background K+ channel to sense external pH variations
[34436] near physiological pH. EMBO J. 16: 5464-5471, 1997.

[34437] 4978.

[34438] Manjunath, N. A.; Bray-Ward, P.; Goldstein, S. A. N.; Gal-

lagher,

[34439] P. G.: Assignment of the 2P domain, acid-sensitive potassium channel

[34440] OAT1 gene KCNK3 to human chromosome bands 2p24.1-p23.3 and murine

[34441] 5B by in situ hybridization. Cytogenet. Cell Genet. 86: 242-243,

[34442] 1999.

[34443] 4979.

[34444] Hacker, B. M.; Tomlinson, J. E.; Wayman, G. A.; Sultana, R.; Chan,

[34445] G.; Villacres, E.; Disteche, C.; Storm, D. R.: Cloning, chromosomal

[34446] mapping, and regulatory properties of the human type 9 adenylyl cyclase

[34447] (ADCY9). Genomics 50: 97-104, 1998.

[34448] 4980.

[34449] Paterson, J. M.; Smith, S. M.; Harmar, A. J.; Antoni, F. A.: Control

[34450] of a novel adenylyl cyclase by calcineurin. Biochem. Biophys. Res.

[34451] Commun. 214: 1000-1008, 1995.

[34452] 4981.

[34453] Premont, R. T.; Matsuoka, I.; Mattei, M. G.; Pouille, Y.; De-
fer,
[34454] N.; Hanoune, J.: Identification and characterization of a
widely
[34455] expressed form of adenylyl cyclase. J. Biol. Chem. 271:
13900-13907,
[34456] 1996.

[34457] 4982.

[34458] Toyota, T.; Hattori, E.; Meerabux, J.; Yamada, K.; Saito, K.;
Shibuya,
[34459] H.; Nankai, M.; Yoshikawa, T.: Molecular analysis, muta-
tion screening,
[34460] and association study of adenylate cyclase type 9 gene
(ADCY9) in
[34461] mood disorders. Am. J. Med. Genet. (Neuropsychiat.
Genet.) 114:
[34462] 84-92, 2002.

[34463] 4983.

[34464] Chi, N.-W.; Lodish, H. F.: Tankyrase is a Golgi-associated
mitogen-activated

[34465] protein kinase substrate that interacts with IRAP in GLUT4 vesicles. J.

[34466] Biol. Chem. 275: 38437-38444, 2000.

[34467] 4984.

[34468] Seimiya, H.; Smith, S.: The telomeric poly(ADP-ribose) polymerase,

[34469] tankyrase 1, contains multiple binding sites for telomeric repeat

[34470] binding factor 1 (TRF1) and a novel acceptor, 182-kDa tankyrase-binding

[34471] protein (TAB182). J. Biol. Chem. 277: 14116-14126, 2002.

[34472] 4985.

[34473] Smith, S.; Giriat, I.; Schmitt, A.; de Lange, T.: Tankyrase, a

[34474] poly(ADP-ribose) polymerase at human telomeres. Science 282: 1484-1487,

[34475] 1998.

[34476] 4986.

[34477] Zhu, L.; Smith, S.; de Lange, T.; Seldin, M. F.: Chromoso-mal mapping

[34478] of the tankyrase gene in human and mouse. Genomics 57:

320-321,

[34479] 1999.

[34480] 4987.

[34481] Hidai, H.; Bardales, R.; Goodwin, R.; Quertermous, T.; Quertermous,

[34482] E. E.: Cloning of capsulin, a basic helix-loop-helix factor expressed

[34483] in progenitor cells of the pericardium and the coronary arteries. Mech.

[34484] Dev. 73: 33-43, 1998.

[34485] 4988.

[34486] Lu, J.; Chang, P.; Richardson, J. A.; Gan, L.; Weiler, H.; Ol-son,

[34487] E. N.: The basic helix-loop-helix transcription factor cap-sulin controls

[34488] spleen organogenesis. Proc. Nat. Acad. Sci. 97: 9525-9530, 2000.

[34489] 4989.

[34490] Lu, J.; Richardson, J. A.; Olson, E. N.: Capsulin: a novel bHLH

[34491] transcription factor expressed in epicardial progenitors

and mesenchyme
[34492] of visceral organs. Mech. Dev. 73: 23-32, 1998.

[34493] 4990.

[34494] Quaggin, S. E.; Vanden Heuvel, G. B.; Igarashi, P.: Pod-1, a mesoderm-specific
[34495] basic-helix-loop-helix protein expressed in mesenchymal and glomerular
[34496] epithelial cells in the developing kidney. Mech. Dev. 71: 37-48,
[34497] 1998.

[34498] 4991.

[34499] Robb, L.; Mifsud, L.; Hartley, L.; Biben, C.; Copeland, N. G.;

[34500] Gilbert, D. J.; Jenkins, N. A.; Harvey, R. P.: Epicardin: a novel
[34501] basic helix-loop-helix transcription factor gene expressed in epicardium,
[34502] branchial arch myoblasts, and mesenchyme of developing lung, gut,
[34503] kidney, and gonads. Dev. Dyn. 213: 105-113, 1998.

[34504] 4992.

[34505] Stacey, P.; Rulten, S.; Dapling, A.; Phillips, S. C.: Molecular

[34506] cloning and expression of human cGMP-binding cGMP-specific phosphodiesterase
[34507] (PDE5). Biochem. Biophys. Res. Commun. 247: 249-254, 1998.
[34508] 4993.

[34509] Yanaka, N.; Kotera, J.; Ohtsuka, A.; Akatsuka, H.; Imai, Y.; Michibata,
[34510] H.; Fujishige, K.; Kawai, E.; Takebayashi, S.-I.; Okumura, K.; Omori,
[34511] K.: Expression, structure and chromosomal localization of the human
[34512] cGMP-binding cGMP-specific phosphodiesterase PDE5A gene. Eur. J.
[34513] Biochem. 255: 391-399, 1998.

[34514] 4994.

[34515] Athanassiadou, A.; Voutsinas, G.; Psiouri, L.; Leroy, E.; Polymeropoulos,
[34516] M. H.; Ilias, A.; Maniatis, G. M.; Papapetropoulos, T.: Genetic analysis
[34517] of families with Parkinson disease that carry the ala53-to-thr mutation
[34518] in the gene encoding alpha-synuclein. (Letter) Am. J. Hum.

Genet. 65:
[34519] 555-558, 1999.

[34520] 4995.

[34521] Auluck, P. K.; Chan, H. Y. E.; Trojanowski, J. Q.; Lee, V. M.-Y.;
[34522] Bonini, N. M.: Chaperone suppression of alpha-synuclein toxicity
[34523] in a Drosophila model for Parkinson's disease. Science 295: 865-868,
[34524] 2002.

[34525] 4996.

[34526] Campion, D.; Martin, C.; Heilig, R.; Charbonnier, F.; Moreau, V.;
[34527] Flaman, J. M.; Petit, J. L.; Hannequin, D.; Brice, A.; Fre-
bourg, T.
[34528] : The NACP/synuclein gene: chromosomal assignment and screening for
[34529] alterations in Alzheimer disease. Genomics 26: 254-257, 1995.
[34530] 4997.

[34531] Chen, X.; Rohan de Silva, H. A.; Pettenati, M. J.; Rao, P. N.;

[34532] St. George-Hyslop, P.; Roses, A. D.; Xia, Y.; Horsburgh, K.; Ueda,
[34533] K.; Saitoh, T.: The human NACP/alpha-synuclein gene: chromosome assignment
[34534] to 4q21.3-q22 and TaqI RFLP analysis. Genomics 26: 425-427, 1995.
[34535] 4998.

[34536] Conway, K. A.; Lee, S.-J.; Rochet, J.-C.; Ding, T. T.; Williamson,
[34537] R. E.; Lansbury, P. T., Jr.: Acceleration of oligomerization, not
[34538] fibrillization, is a shared property of both alpha-synuclein mutations
[34539] linked to early-onset Parkinson's disease: implications for pathogenesis
[34540] and therapy. Proc. Nat. Acad. Sci. 97: 571-576, 2000.

[34541] 4999.

[34542] Engelender, S.; Kaminsky, Z.; Guo, X.; Sharp, A. H.; Amar-
avi, R.
[34543] K.; Kleiderlein, J. J.; Margolis, R. L.; Troncoso, J. C.; Lana-
han,
[34544] A. A.; Worley, P. F.; Dawson, V. L.; Dawson, T. M.; Ross, C.

A.:

[34545] Synphilin-1 associates with alpha-synuclein and promotes the formation

[34546] of cytosolic inclusions. Nature Genet. 22: 110-114, 1999.

[34547] 5000.

[34548] Farrer, M.; Wavrant-De Vrieze, F.; Crook, R.; Boles, L.; Perez-Tur,

[34549] J.; Hardy, J.; Johnson, W. G.; Steele, J.; Maraganore, D.; Gwinn,

[34550] K.; Lynch, T.: Low frequency of alpha-synuclein mutations in familial

[34551] Parkinson's disease. Ann. Neurol. 43: 394-397, 1998.

[34552] 5001.

[34553] Feany, M. B.; Bender, W. W.: A Drosophila model of Parkinson's

[34554] disease. Nature 404: 394-398, 2000.

[34555] 5002.

[34556] George, J. M.; Jin, H.; Woods, W. S.; Clayton, D. F.: Characterization

[34557] of a novel protein regulated during the critical period for song learning

[34558] in the zebra finch. Neuron 15: 361-372, 1995.

[34559] 5003.

[34560] Giasson, B. I.; Duda, J. E.; Murray, I. V. J.; Chen, Q.; Souza,

[34561] J. M.; Hurtig, H. I.; Ischiropoulos, H.; Trojanowski, J. Q.;
Lee,

[34562] V. M.-Y.: Oxidative damage linked to neurodegeneration
by selective

[34563] alpha-synuclein nitration in synucleinopathy lesions. Sci-
ence 290:

[34564] 985-989, 2000.

[34565] 5004.

[34566] Goedert, M.: The awakening of alpha-synuclein. Nature
388:

[34567] 232-233, 1997.

[34568] 5005.

[34569] Ho, S.-L.; Kung, M. H. W.: G209A mutation in the alpha-
synuclein

[34570] gene is rare and not associated with sporadic Parkinson's
disease. Mov.

[34571] Disord. 13: 970-971, 1998.

[34572] 5006.

[34573] Jakes, R.; Spillantini, M. G.; Goedert, M.: Identification of

[34574] two distinct synucleins from human brain. FEBS Lett. 345:
27-32,

[34575] 1994.

[34576] 5007.

[34577] Kruger, R.; Kuhn, W.; Leenders, K. L.; Sprengelmeyer, R.;
Muller,

[34578] T.; Woitalla, D.; Portman, A. T.; Maguire, R. P.; Veenma, L.;
Schroder,

[34579] U.; Schols, L.; Epplen, J. T.; Riess, O.; Przuntek, H.: Familial

[34580] parkinsonism with synuclein pathology: clinical and PET
studies of

[34581] A30P mutation carriers. Neurology 56: 1355-1362, 2001.

[34582] 5008.

[34583] Kruger, R.; Kuhn, W.; Muller, T.; Woitalla, D.; Graeber, M.;
Kosel,

[34584] S.; Przuntek, H.; Epplen, J. T.; Schols, L.; Riess, O.:
Ala30-to-pro

[34585] mutation in the gene encoding alpha-synuclein in Parkin-
son's disease.

[34586] (Letter) Nature Genet. 18: 106-108, 1998.

[34587] 5009.

[34588] Mark, J.; Dahlenfors, R.; Ekedahl, C.; Stenman, G.: The mixed

[34589] salivary gland tumor--a normally benign human neoplasm frequently

[34590] showing specific chromosomal abnormalities. Cancer Genet. Cytogenet. 2:

[34591] 231-234, 1980.

[34592] 5010.

[34593] Mark, J.; Dahlenfors, R.; Ekedahl, C.; Stenman, G.: Chro-mosomal

[34594] patterns in a benign human neoplasm, the mixed salivary gland tumour. Hereditas 96:

[34595] 141-148, 1982.

[34596] 5011.

[34597] Rapp, U. R.; Goldsborough, M. D.; Mark, G. E.; Bonner, T. I.;

[34598] Groffen, J.; Reynolds, F. H., Jr.; Stephenson, J. R.: Structure and

[34599] biological activity of v-raf, a unique oncogene transduced

by a retrovirus. Proc.
[34600] Nat. Acad. Sci. 80: 4218-4222, 1983.

[34601] 5012.

[34602] Shimizu, K.; Nakatsu, Y.; Sekiguchi, M.; Hokamura, K.; Tanaka,
[34603] K.; Terada, M.; Sugimura, T.: Molecular cloning of an activated human
[34604] oncogene, homologous to v-raf, from primary stomach cancer. Proc.
[34605] Nat. Acad. Sci. 82: 5641-5645, 1985.

[34606] 5013.

[34607] Tory, K.; Latif, F.; Modi, W.; Schmidt, L.; Wei, M. H.; Li, H.;

[34608] Cobler, P.; Orcutt, M. L.; Delisio, J.; Geil, L.; Zbar, B.; Lerman,
[34609] M. I.: A genetic linkage map of 96 loci on the short arm of human
[34610] chromosome 3. Genomics 13: 275-286, 1992.

[34611] 5014.

[34612] Amiel, J.; Salomon, R.; Attie, T.; Pelet, A.; Trang, H.; Mokhtari,
[34613] M.; Gaultier, C.; Munnich, A.; Lyonnet, S.: Mutations of the

RET-GDNF
[34614] signaling pathway in Ondine's curse. (Letter) Am. J. Hum. Genet. 62:
[34615] 715-717, 1998.

[34616] 5015.

[34617] Angrist, M.; Bolk, S.; Thiel, B.; Puffenberger, E. G.; Hofstra,

[34618] R. M.; Buys, C. H. C. M.; Cass, D. T.; Chakravarti, A.: Muta-
tion
[34619] analysis of the RET receptor tyrosine kinase in Hirschsprung disease. Hum.
[34620] Molec. Genet. 4: 821-830, 1995.

[34621] 5016.

[34622] Antinolo, G.; Marcos, I.; Fernandez, R. M.; Romero, M.; Borrego,
[34623] S.: A novel germline point mutation, c.2304G(T, in codon 768 of the
[34624] RET proto-oncogene in a patient with medullary thyroid carcinoma.
[34625] (Letter) Am. J. Med. Genet. 110: 85-87, 2002.

[34626] 5017.

[34627] Attie, T.; Pelet, A.; Edery, P.; Eng, C.; Mulligan, L. M.;

Amiel,

[34628] J.; Boutrand, L.; Beldjord, C.; Nihoul-Fekete, C.; Munnich, A.; Ponder,

[34629] B. A. J.; Lyonnet, S.: Diversity of RET proto-oncogene mutations

[34630] in familial and sporadic Hirschsprung disease. Hum. Molec. Genet. 4:

[34631] 1381-1386, 1995.

[34632] 5018.

[34633] Attie-Bitach, T.; Abitbol, M.; Gerard, M.; Delezoide, A.-L.; Auge,

[34634] J.; Pelet, A.; Amiel, J.; Pachnis, V.; Munnich, A.; Lyonnet, S.; Vekemans,

[34635] M.: Expression of the RET proto-oncogene in human embryos. Am. J.

[34636] Med. Genet. 80: 481-486, 1998.

[34637] 5019.

[34638] Auricchio, A.; Griseri, P.; Carpentieri, M. L.; Betsos, N.; Staiano,

[34639] A.; Tozzi, A.; Priolo, M.; Thompson, H.; Bocciardi, R.; Romeo, G.;

[34640] Ballabio, A.; Ceccherini, I.: Double heterozygosity for a

RET substitution

[34641] interfering with splicing and an EDNRB missense mutation in Hirschsprung

[34642] disease. (Letter) Am. J. Hum. Genet. 64: 1216-1221, 1999.

[34643] 5020.

[34644] Batourina, E.; Choi, C.; Paragas, N.; Bello, N.; Hensle, T.; Costantini,

[34645] F. D.; Schuchardt, A.; Bacallao, R. L.; Mendelsohn, C. L.: Distal

[34646] ureter morphogenesis depends on epithelial cell remodeling mediated

[34647] by vitamin A and Ret. Nature Genet. 32: 109-115, 2002. Note: Erratum:

[34648] Nature Genet. 32: 331 only, 2002.

[34649] 5021.

[34650] Batourina, E.; et al; et al: Vitamin A controls epithelial/mesenchymal

[34651] interactions through Ret expression. Nature Genet. 27: 74-78, 2001.

[34652] 5022.

[34653] Berndt, I.; Reuter, M.; Saller, B.; Frank-Raue, K.; Groth, P.:

[34654] Grubendorf, M.; Raue, F.; Ritter, M. M.; Hoppner, W.: A new hot spot

[34655] for mutations in the ret protooncogene causing familial medullary

[34656] thyroid carcinoma and multiple endocrine neoplasia type 2A. J. Clin.

[34657] Endocr. Metab. 83: 770-774, 1998.

[34658] 5023.

[34659] Boccia, L. M.; Green, J. S.; Joyce, C.; Eng, C.; Taylor, S. A.

[34660] M.; Mulligan, L. M.: Mutation of RET codon 768 is associated with

[34661] the FMTC phenotype. Clin. Genet. 51: 81-85, 1997.

[34662] 5024.

[34663] Bolino, A.; Schuffenecker, I.; Luo, Y.; Seri, M.; Silengo, M.;

[34664] Tocco, T.; Chabrier, G.; Houdent, C.; Murat, A.; Schlum-
berger, M.;

[34665] Tourniaire, J.; Lenoir, G. M.; Romeo, G.: RET mutations in
exons

[34666] 13 and 14 of FMTC patients. Oncogene 10: 2415-2419,
1995.

[34667] 5025.

[34668] Bolk, S.; Angrist, M.; Schwartz, S.; Silvestri, J. M.; Weese-
Mayer,

[34669] D. E.; Chakravarti, A.: Congenital central hypoventilation
syndrome:

[34670] mutation analysis of the receptor tyrosine kinase RET. Am.
J. Med.

[34671] Genet. 63: 603-609, 1996.

[34672] 5026.

[34673] Bolk Gabriel, S.; Salomon, R.; Pelet, A.; Angrist, M.; Amiel,

[34674] J.; Fornage, M.; Attie-Bitach, T.; Olson, J. M.; Hofstra, R.;
Buys,

[34675] C.; Steffann, J.; Munnich, A.; Lyonnet, S.; Chakravarti, A.:

Segregation
[34676] at three loci explains familial and population risk in Hirschsprung
[34677] disease. Nature Genet. 31: 89-93, 2002.

[34678] 5027.

[34679] Ceccherini, I.; Hofstra, R. M.; Luo, Y.; Stulp, R. P.; Barone,

[34680] V.; Stelwagen, T.; Bocciardi, R.; Nijveen, H.; Bolino, A.; Seri, M.;
[34681] Ronchetto, P.; Pasini, B.; Bozzano, M.; Buys, C. H. C. M.; Romeo,
[34682] G.: DNA polymorphisms and conditions for SSCP analysis of the 20
[34683] exons of the Ret proto-oncogene. Oncogene 9: 3025-3029, 1994.
[34684] 5028.

[34685] Cohn, D. H.; Byers, P. H.; Steinmann, B.; Gelinas, R. E.: Lethal
[34686] osteogenesis imperfecta resulting from a single nucleotide change
[34687] in one human pro-alpha-1(I) collagen allele. Proc. Nat. Acad. Sci. 83:
[34688] 6045-6047, 1986.

[34689] 5029.

[34690] Genovese, C.; Brufsky, A.; Shapiro, J.; Rowe, D.: Detection of

[34691] mutations in human type I collagen mRNA in osteogenesis imperfecta

[34692] by indirect RNase protection. J. Biol. Chem. 264: 9632-9637, 1989.

[34693] 5030.

[34694] Furlong, C. E.; Richter, R. J.; Chapline, C.; Crabb, J. W.: Purification

[34695] of rabbit and human serum paraoxonase. Biochemistry 30: 10133-10140,

[34696] 1991.

[34697] 5031.

[34698] Furlong, C. E.; Richter, R. J.; Seidel, S. L.; Motulsky, A. G.

[34699] : Role of genetic polymorphism of human plasma paraoxonase/arylesterase

[34700] in hydrolysis of the insecticide metabolites chlorpyrifos oxon and

[34701] paraoxon. Am. J. Hum. Genet. 43: 230-238, 1988.

[34702] 5032.

[34703] Garin, M.-C. B.; James, R. W.; Dussoix, P.; Blanche, H.; Passa,
[34704] P.; Froguel, P.; Ruiz, J.: Paraoxonase polymorphism met-leu54 is
[34705] associated with modified serum concentrations of the enzyme: a possible
[34706] link between the paraoxonase gene and increased risk of cardiovascular
[34707] disease in diabetes. J. Clin. Invest. 99: 62-66, 1997.

[34708] 5033.

[34709] Geldmacher-von Mallinckrodt, M.: Polymorphism of human serum
[34710] paraoxonase. Hum. Genet. 45 (suppl. 1): 65-68, 1978.

[34711] 5034.

[34712] Geldmacher-von Mallinckrodt, M.; Lindorft, H. H.; Petenyi, M.;
[34713] Flugel, M.; Fischer, T.; Hiller, T.: Genetisch determinierter Polymorphismus
[34714] de menschlichen Serum-Paroxonase (E.C.3.1.1.2). Humangenetik 17:
[34715] 331-335, 1973.

[34716] 5035.

[34717] Hassett, C.; Richter, R. J.; Humbert, R.; Chapline, C.; Crabb,

[34718] J. W.; Omiecinski, C. J.; Furlong, C. E.: Characterization of cDNA

[34719] clones encoding rabbit and human serum paraoxonase: the mature protein

[34720] retains its signal sequence. Biochemistry 30: 10141-10149, 1991.

[34721] 5036.

[34722] Humbert, R.; Adler, D. A.; Disteche, C. M.; Hassett, C.; Omiecinski,

[34723] C. J.; Furlong, C. E.: The molecular basis of the human serum paraoxonase

[34724] activity polymorphism. Nature Genet. 3: 73-76, 1993.

[34725] 5037.

[34726] Ito, T.; Yasue, H.; Yoshimura, M.; Nakamura, S.; Nakayama, M.;

[34727] Shimasaki, Y.; Harada, E.; Mizuno, Y.; Kawano, H.; Ogawa, H.: Paraoxonase

[34728] gene gln192-to-arg (Q192R) polymorphism is associated with coronary

[34729] artery spasm. Hum. Genet. 110: 89-94, 2002.

[34730] 5038.

[34731] Kao, Y.-L.; Donaghue, K.; Chan, A.; Knight, J.; Silink, M.: A

[34732] variant of paraoxonase (PON1) gene is associated with di-
abetic retinopathy

[34733] in IDDM. J. Clin. Endocr. Metab. 83: 2589-2592, 1998.

[34734] 5039.

[34735] La Du, B. N.: The human serum paraoxonase/arylesterase
polymorphism.

[34736] (Editorial) Am. J. Hum. Genet. 43: 227-229, 1988.

[34737] 5040.

[34738] Li, W.-F.; Furlong, C. E.; Costa, L. G.: Paraoxonase protects

[34739] against chlorpyrifos toxicity in mice. Toxic. Lett. 76:
219-226,

[34740] 1995.

[34741] 5041.

[34742] Li, W.-F.; Matthews, C.; Disteche, C. M.; Costa, L. G.; Fur-
long,

[34743] C. E.: Paraoxonase (Pon1) gene in mice: sequencing, chro-

mosomal localization
[34744] and developmental expression. Pharmacogenetics 7: 137-144, 1997.
[34745] 5042.
[34746] Mackness, B.; Mackness, M. I.; Arrol, S.; Turkie, W.; Durrington,
[34747] P. N.: Effect of the human serum paraoxonase 55 and 192 genetic polymorphisms
[34748] on the protection by high density lipoprotein against low density
[34749] lipoprotein oxidative modification. FEBS Lett. 423: 57-60, 1998.
[34750] 5043.
[34751] Mochizuki, H.; Scherer, S. W.; Xi, T.; Nickle, D. C.; Majer, M.;
[34752] Huizenga, J. J.; Tsui, L.-C.; Prochazka, M.: Human PON2 gene at 7q21.3:
[34753] cloning, multiple mRNA forms, and missense polymorphisms in the coding
[34754] sequence. Gene 213: 149-157, 1998.
[34755] 5044.
[34756] Mueller, R. F.; Hornung, S.; Furlong, C. E.; Anderson, J.;

Giblett,
[34757] E. R.; Motulsky, A. G.: Plasma paraoxonase polymorphism: a new enzyme
[34758] assay, population, family, biochemical, and linkage studies. Am.
[34759] J. Hum. Genet. 35: 393-408, 1983.

[34760] 5045.

[34761] Navab, M.; Hama-Levy, S.; Van Lenten, B. J.; Fonarow, G. C.; Cardinez,
[34762] C. J.; Castellani, L. W.; Brennan, M.-L.; Lusis, A. J.; Fogelman,
[34763] A. M.: Mildly oxidized LDL induces an increased apolipoprotein J/paraoxonase
[34764] ratio. J. Clin. Invest. 99: 2005-2019, 1997.

[34765] 5046.

[34766] Neel, J. V.; Tanis, R. J.; Migliazza, E. C.; Spielman, R. S.;

[34767] Salzano, F. M.; Oliver, W. J.; Morrow, M.; Bachofer, S.: Genetic
[34768] studies of the Macushi and Wapishana Indians. I. Rare genetic variants
[34769] and a 'private polymorphism' of esterase A. Hum. Genet.

36: 81-108,

[34770] 1977.

[34771] 5047.

[34772] Nielsen, A.; Eiberg, H.; Mohr, J.: Number of loci responsi-
ble

[34773] for the inheritance of high and low activity of paraox-
onase. Clin.

[34774] Genet. 29: 216-221, 1986.

[34775] 5048.

[34776] Odawara, M.; Tachi, Y.; Yamashita, K.: Paraoxonase poly-
morphism

[34777] Gln192-Arg is associated with coronary heart disease in
Japanese noninsulin-dependent

[34778] diabetes mellitus. J. Clin. Endocr. Metab. 82: 2257-2260,
1997.

[34779] 5049.

[34780] Ortigoza-Ferado, J.; Richter, R. J.; Hornung, S. K.; Motul-
sky,

[34781] A. G.; Furlong, C. E.: Paraoxon hydrolysis in human serum
mediated

[34782] by a genetically variable arylesterase and albumin. Am. J.

Hum. Genet. 36:
[34783] 295-305, 1984.

[34784] 5050.

[34785] Paolisso, G.; Manzella, D.; Tagliamonte, M. R.; Barbieri, M.;

[34786] Marfella, R.; Zito, G.; Bonafe, M.; Giugliano, D.; Franceschi, C.;
[34787] Varricchio, M.: The BB-paraoxonase genotype is associated with impaired
[34788] brachial reactivity after acute hypertriglyceridemia in healthy subjects. J.
[34789] Clin. Endocr. Metab. 86: 1078-1082, 2001.

[34790] 5051.

[34791] Playfer, J. R.; Eze, L. C.; Bullen, M. F.; Evans, D. A. P.: Genetic
[34792] polymorphism and interethnic variability of plasma paroxonase activity. J.
[34793] Med. Genet. 13: 337-342, 1976.

[34794] 5052.

[34795] Schmiegelow, K.; Eiberg, H.; Tsui, L.-C.; Buchwald, M.; Phelan,
[34796] P. D.; Williamson, R.; Warwick, W.; Niebuhr, E.; Mohr, J.;

Schwartz,

[34797] M.; Koch, C.: Linkage between the loci for cystic fibrosis and paraoxonase. Clin.

[34798] Genet. 29: 374-377, 1986.

[34799] 5053.

[34800] Serrato, M.; Marian, A. J.: A variant of human paraoxonase/arylesterase

[34801] (HUMPONA) gene is a risk factor for coronary artery disease. J. Clin.

[34802] Invest. 96: 3005-3008, 1995.

[34803] 5054.

[34804] Shih, D. M.; Gu, L.; Xia, Y.-R.; Navab, M.; Li, W.-F.; Hama, S.;

[34805] Castellani, L. W.; Furlong, C. E.; Costa, L. G.; Fogelman, A. M.;

[34806] Lusis, A. J.: Mice lacking serum paraoxonase are susceptible to organophosphate

[34807] toxicity and atherosclerosis. Nature 394: 284-287, 1998.

[34808] 5055.

[34809] Simpson, N. E.: Serum arylesterase levels of activity in twins

[34810] and their parents. Am. J. Hum. Genet. 23: 375-382, 1971.

[34811] 5056.

[34812] Sorenson, R. C.; Primo-Parmo, S. L.; Camper, S.; La Du, B. N.

[34813] : The genetic mapping and gene structure of mouse paraoxonase/arylesterase. Genomics 30:

[34814] 431-438, 1995.

[34815] 5057.

[34816] Tashian, R. E.: Genetic variation and evolution of the car-boxylic

[34817] esterases and carbonic anhydrases of primate erythro-cytes. Am. J.

[34818] Hum. Genet. 17: 257-272, 1965.

[34819] 5058.

[34820] Tashian, R. E.; Shaw, M. W.: Inheritance of an erythrocyte acetylesterase

[34821] variant of man. Am. J. Hum. Genet. 14: 295-300, 1962.

[34822] 5059.

[34823] Tsui, L.-C.; Buchwald, M.; Barker, D.; Braman, J. C.; Knowlton,

[34824] R.; Schumm, J. W.; Eiberg, H.; Mohr, J.; Kennedy, D.; Plavsic, N.;

[34825] Zsiga, M.; Markiewicz, D.; Akots, G.; Brown, V.; Helms, C.; Gravius,

[34826] T.; Parker, C.; Rediker, K.; Donis-Keller, H.: Cystic fibrosis locus

[34827] defined by a genetically linked polymorphic DNA marker. Science 230:

[34828] 1054-1057, 1985.

[34829] 5060.

[34830] Gogolin, K. J.; Wray, L. K.; Slaughter, C. A.; Harris, H.: A monoclonal

[34831] antibody that reacts with nonallelic enzyme glycoproteins. Science 216:

[34832] 59-61, 1982.

[34833] 5061.

[34834] Goldstein, D. J.; Rogers, C. E.; Harris, H.: Expression of al-kaline

[34835] phosphatase loci in mammalian tissues. Proc. Nat. Acad. Sci. 77:

[34836] 2857-2860, 1980.

[34837] 5062.

[34838] Gould, B. S.: Studies on the source of serum phosphatase: the

[34839] nature of the increased serum phosphatase in rats after fat feeding. Arch.

[34840] Biochem. 4: 175-181, 1944.

[34841] 5063.

[34842] Griffin, C. A.; Smith, M.; Henthorn, P. S.; Harris, H.; Weiss,

[34843] M. J.; Raducha, M.; Emanuel, B. S.: Human placental and intestinal

[34844] alkaline phosphatase genes map to 2q34-q37. Am. J. Hum. Genet. 41:

[34845] 1025-1034, 1987.

[34846] 5064.

[34847] Henthorn, P. S.; Raducha, M.; Edwards, Y. H.; Weiss, M. J.; Slaughter,

[34848] C.; Lafferty, M. A.; Harris, H.: Nucleotide and amino acid sequences

[34849] of human intestinal alkaline phosphatase: close homology to placental

[34850] alkaline phosphatase. Proc. Nat. Acad. Sci. 84:

1234-1238, 1987.

[34851] 5065.

[34852] Henthorn, P. S.; Raducha, M.; Kadesch, T.; Weiss, M. J.; Harris,

[34853] H.: Sequence and characterization of the human intestinal alkaline

[34854] phosphatase gene. J. Biol. Chem. 263: 12011-12019, 1988.

[34855] 5066.

[34856] Langman, M. J. S.; Leuthold, E.; Robson, E. B.; Harris, J.; Luffman,

[34857] J. E.; Harris, H.: Influence of diet on the 'intestinal' compo-nent

[34858] of serum alkaline phosphatase in people of different ABO blood groups

[34859] and secretor status. Nature 212: 41-43, 1966.

[34860] 5067.

[34861] Lehmann, F.-G.: Human alkaline phosphatases: evidence of three

[34862] isoenzymes (placental, intestinal and liver-bone-kidney-type) by lectin-binding

[34863] affinity and immunological specificity. Biochim. Biophys.

Acta 616:
[34864] 41-59, 1980.

[34865] 5068.

[34866] Fedde, K. N.; Whyte, M. P.: Alkaline phosphatase (tissue-nonspecific
[34867] isoenzyme) is a phosphoethanolamine and pyridoxal-5-prime-phosphate
[34868] ectophosphatase: normal and hypophosphatasia fibroblast study. Am.
[34869] J. Hum. Genet. 47: 767-775, 1990.

[34870] 5069.

[34871] Goseki-Sone, M.; Orimo, H.; Iimura, T.; Miyazaki, H.; Oda, K.;
[34872] Shibata, H.; Yanagishita, M.; Takagi, Y.; Watanabe, H.; Shimada, T.;
[34873] Oida, S.: Expression of the mutant (1735T-DEL) tissue-nonspecific
[34874] alkaline phosphatase gene from hypophosphatasia patients. J. Bone
[34875] Miner. Res. 13: 1827-1834, 1998.

[34876] 5070.

[34877] Greenberg, C. R.; Evans, J. A.; McKendry-Smith, S.; Redekopp, S.;
[34878] Haworth, J. C.; Mulivor, R.; Chodirker, B. N.: Infantile hypophosphatasia:
[34879] localization within chromosome region 1p36.1-34 and prenatal diagnosis
[34880] using linked DNA markers. Am. J. Hum. Genet. 46: 286-292, 1990.
[34881] 5071.

[34882] Greenberg, C. R.; Taylor, C. L. D.; Haworth, J. C.; Seargeant,
[34883] L. E.; Philipps, S.; Triggs-Raine, B.; Chodirker, B. N.: A homoallelic
[34884] gly317-to-asp mutation in ALPL causes the perinatal (lethal) form
[34885] of hypophosphatasia in Canadian Mennonites. Genomics 17: 215-217,
[34886] 1993.

[34887] 5072.

[34888] Henthorn, P. S.; Whyte, M. P.: Missense mutations of the tissue-nonspecific
[34889] alkaline phosphatase gene in hypophosphatasia. Clin.

Chem. 38: 2501-2505,

[34890] 1992.

[34891] 5073.

[34892] Hu, J. C. C.; Plaetke, R.; Mornet, E.; Zhang, C.; Sun, X.; Thomas,

[34893] H. F.; Simmer, J. P.: Case report of a family with dominant hypophosphatasia. Eur.

[34894] J. Oral Sci. 108: 189-194, 2000.

[34895] 5074.

[34896] Hua, J.-C.; Berger, J.; Pan, Y.-C. E.; Hulmes, J. D.; Uden-friend,

[34897] S.: Partial sequencing of human adult, human fetal, and bovine intestinal

[34898] alkaline phosphatases: comparison with the human placental and liver

[34899] isozymes. Proc. Nat. Acad. Sci. 83: 2368-2372, 1986.

[34900] 5075.

[34901] Kishi, F.; Matsuura, S.; Murano, I.; Akita, A.; Kajii, T.: Prenatal

[34902] diagnosis of infantile hypophosphatasia. Prenatal Diag. 11: 305-309,

[34903] 1991.

[34904] 5076.

[34905] Lia-Baldini, A. S.; Muller, F.; Taillandier, A.; Gibrat, J. F.;

[34906] Mouchard, M.; Robin, B.; Simon-Bouy, B.; Serre, J. L.;
Aylsworth,

[34907] A. S.; Bieth, E.; Delanote, S.; Freisinger, P.; Hu, J. C.-C.;
Krohn,

[34908] H.-P.; Nunes, M. E.; Mornet, E.: A molecular approach to
dominance

[34909] in hypophosphatasia. Hum. Genet. 109: 99-108, 2001.

[34910] 5077.

[34911] Matsuura, S.; Kishi, F.; Kajii, T.: Characterization of a
5-prime-flanking

[34912] region of the human liver/bone/kidney alkaline phos-
phatase gene: two

[34913] kinds of mRNA from a single gene. Biochem. Biophys. Res.
Commun. 168:

[34914] 993-1000, 1990.

[34915] 5078.

[34916] Ny, T.; Sawdey, M.; Lawrence, D.; Millan, J. L.; Loskutoff,
D.

[34917] J.: Cloning and sequence of a cDNA coding for the human beta-migrating
[34918] endothelial-cell-type plasminogen activator inhibitor. Proc. Nat.
[34919] Acad. Sci. 83: 6776-6780, 1986.

[34920] 5079.

[34921] Pannekoek, H.; Veerman, H.; Lambers, H.; Diergaarde, P.; Verweij,
[34922] C. L.; van Zonneveld, A.-J.; van Mourik, J. A.: Endothelial plasminogen
[34923] activator inhibitor (PAI): a new member of the serpin gene family. EMBO
[34924] J. 5: 2539-2544, 1986.

[34925] 5080.

[34926] Pastinen, T.; Perola, M.; Niini, P.; Terwilliger, J.; Salomaa,
[34927] V.; Vartiainen, E.; Peltonen, L.; Syvanen, A.-C.: Array-based multiplex
[34928] analysis of candidate genes reveals two independent and additive genetic
[34929] risk factors for myocardial infarction in the Finnish popu-lation. Hum.

[34930] Molec. Genet. 7: 1453-1462, 1998.

[34931] 5081.

[34932] Patrassi, G. M.; Sartori, M. T.; Saggiorato, G.; Boeri, G.; Girolami,

[34933] A.: Familial thrombophila (sic) associated with high levels of plasminogen

[34934] activator inhibitor. Fibrinolysis 6: 99-103, 1992.

[34935] 5082.

[34936] Schleef, R. R.; Higgins, D. L.; Pillemer, E.; Levitt, L. J.:

[34937] Bleeding diathesis due to decreased functional activity of type 1

[34938] plasminogen activator inhibitor. J. Clin. Invest. 83: 1747-1752,

[34939] 1989.

[34940] 5083.

[34941] Schwartz, C. E.; Stanislovitis, P.; Phelan, M. C.; Klinger, K.;

[34942] Taylor, H. A.; Stevenson, R. E.: Deletion mapping of plas-minogen

[34943] activator inhibitor, type I (PLANH1) and beta-glucuronidase (GUSB)

[34944] in 7q21-q22. Cytogenet. Cell Genet. 51: 152-153, 1991.

[34945] 5084.

[34946] van Zonneveld, A.-J.; Curriden, S. A.; Loskutoff, D. J.: Type

[34947] 1 plasminogen activator inhibitor gene: functional analysis and glucocorticoid

[34948] regulation of its promoter. Proc. Nat. Acad. Sci. 85: 5525-5529,

[34949] 1988.

[34950] 5085.

[34951] Westendorp, R. G. J.; Hottenga, J.-J.; Slagboom, P. E.: Vari-ation

[34952] in plasminogen-activator-inhibitor-1 gene and risk of meningococcal

[34953] septic shock. Lancet 354: 561-563, 1999.

[34954] 5086.

[34955] Yamada, N.; Arinami, T.; Yamakawa-Kobayashi, K.; Watanabe, H.;

[34956] Sohda, S.; Hamada, H.; Kubo, T.; Hamaguchi, H.: The 4G/5G polymorphism

[34957] of the plasminogen activator inhibitor-1 gene is associ-ated with severe

[34958] preeclampsia. J. Hum. Genet. 45: 138-141, 2000.

[34959] 5087.

[34960] Yamamoto, K.; Takeshita, K.; Shimokawa, T.; Yi, H.; Isobe, K.;

[34961] Loskutoff, D. J.; Saito, H.: Plasminogen activator inhibitor-1 is

[34962] a major stress-regulated gene: implications for stress-induced thrombosis

[34963] in aged individuals. Proc. Nat. Acad. Sci. 99: 890-895, 2002.

[34964] 5088.

[34965] Aznar, J.; Estelles, A.; Vila, V.; Reganon, E.; Espana, F.; Villa,

[34966] P.: Inherited fibrinolytic disorder due to an enhanced plasminogen

[34967] activator level. Thromb. Haemost. 52: 196-200, 1984.

[34968] 5089.

[34969] Benham, F. J.; Spurr, N.; Povey, S.; Brinton, B. T.; Solomon, E.;

[34970] Goodfellow, P. N.; Harris, T. J. R.: Tissue-type plasminogen activator

[34971] (PLAT) maps to chromosome 8 and there is a common restriction fragment
[34972] length polymorphism within the gene. (Abstract) Cytogenet. Cell Genet. 40:
[34973] 581 only, 1985.

[34974] 5090.

[34975] Booth, N. A.; Bennett, B.; Wijngaards, G.; Grieve, J. H. K.: A

[34976] new life-long hemorrhagic disorder due to excess plasminogen activator. Blood 61:
[34977] 267-275, 1983.

[34978] 5091.

[34979] Bell, L. D.; Smith, J. C.; Derbyshire, R.; Finlay, M.; Johnson,

[34980] I.; Gilbert, R.; Slocombe, P.; Cook, E.; Richards, H.; Clissold, P.;
[34981] Meredith, D.; Powell-Jones, C. H.; Dawson, K. M.; Carter, B. L.; McCullagh,
[34982] K. G.: Chemical synthesis, cloning and expression in mammalian cells
[34983] of a gene coding for human tissue-type plasminogen activator. Gene 63:
[34984] 155-163, 1988.

[34985] 5092.

[34986] Browne, M. J.; Tyrrell, A. W. R.; Chapman, C. G.; Carey, J. E.;

[34987] Glover, D. M.; Grosveld, F. G.; Dodd, I.; Robinson, J. H.: Isolation

[34988] of a human tissue-type plasminogen-activator genomic DNA clone and

[34989] its expression in mouse L cells. Gene 33: 279-284, 1985.

[34990] 5093.

[34991] Edlund, T.; Ny, T.; Ranby, M.; Heden, L.-O.; Palm, G.; Holmgren,

[34992] E.; Josephson, S.: Isolation of cDNA sequences coding for a part

[34993] of human tissue plasminogen activator. Proc. Nat. Acad. Sci. 80:

[34994] 349-352, 1983.

[34995] 5094.

[34996] Johansson, L.; Hedner, U.; Nilsson, I. M.: A family with thromboembolic

[34997] disease associated with deficient fibrinolytic activity in vessel

[34998] wall. Acta Med. Scand. 203: 477-480, 1978.

[34999] 5095.

[35000] Jorgensen, M.; Mortensen, J. Z.; Madsen, A. G.; Thorsen, S.; Jacobsen,

[35001] B.: A family with reduced plasminogen activator activity in blood

[35002] associated with recurrent venous thrombosis. Scand. J. Haemat. 29:

[35003] 217-223, 1982.

[35004] 5096.

[35005] Kucherlapati, R.; Tepper, R.; Granelli-Piperno, A.; Reich, E. W.

[35006] : Modulation and mapping of a human plasminogen activator by cell

[35007] fusion. Cell 15: 1331-1340, 1978.

[35008] 5097.

[35009] Loscalzo, J.; Braunwald, E.: Tissue plasminogen activator. New

[35010] Eng. J. Med. 319: 925-931, 1988.

[35011] 5098.

[35012] Ludwig, M.; Wohn, K.-D.; Schleuning, W.-D.; Olek, K.: Allelic
[35013] dimorphism in the human tissue-type plasminogen activator (TPA) gene
[35014] as a result of an Alu insertion/deletion event. Hum. Genet. 88:
[35015] 388-392, 1992.

[35016] 5099.

[35017] MacDonald, M. E.; van Zonneveld, A.-J.; Pannekoek, H.: Functional
[35018] analysis of the human tissue-type plasminogen activator protein: the
[35019] light chain. Gene 42: 59-67, 1986.

[35020] 5100.

[35021] Ny, T.; Elgh, F.; Lund, B.: The structure of the human tissue-type
[35022] plasminogen activator gene: correlation of intron and exon structures
[35023] to functional and structural domains. Proc. Nat. Acad. Sci. 81:
[35024] 5355-5359, 1984.

[35025] 5101.

[35026] Pennica, D.; Holmes, W. E.; Kohr, W. J.; Harkins, R. N.; Ve-
har,
[35027] G. A.; Ward, C. A.; Bennett, W. F.; Yelverton, E.; Seeburg, P.
H.;
[35028] Heyneker, H. L.; Goeddel, D. V.; Collen, D.: Cloning and
expression
[35029] of human tissue-type plasminogen activator cDNA in E.
coli. Nature 301:
[35030] 214-221, 1983.

[35031] 5102.

[35032] Petaja, J.; Rasi, V.; Vahtera, E.; Myllyla, G.: Familial cluster-
ing
[35033] of defective release of t-PA. Brit. J. Haemat. 79: 291-295,
1991.
[35034] 5103.

[35035] Rajput, B.; Degen, S. F.; Reich, E.; Eddy, R. L.; Shows, T. B.

[35036] : Mapping of tissue plasminogen activator (PLAT) to chro-
mosome 8 and
[35037] urokinase (PLAU) to chromosome 10 in humans. (Abstract)
Cytogenet.

[35038] Cell Genet. 40: 728-729, 1985.

[35039] 5104.

[35040] Rajput, B.; Degen, S. F.; Reich, E.; Waller, E. K.; Axelrod, J.;

[35041] Eddy, R. L.; Shows, T. B.: Chromosomal locations of hu-
man tissue

[35042] plasminogen activator and urokinase genes. Science 230:
672-674,

[35043] 1985.

[35044] 5105.

[35045] Distante, S.; Nasioulas, S.; Somers, G. R.; Cameron, D. J.
S.;

[35046] Young, M. A.; Forrest, S. M.; Gardner, R. J. M.: Familial
adenomatous

[35047] polyposis in a 5 year old child: a clinical, pathological, and
molecular

[35048] genetic study. J. Med. Genet. 33: 157-160, 1996.

[35049] 5106.

[35050] Duhamel, J.; Berthon, G.; Dubarry, J. J.: Etude mathema-
tique

[35051] de l'heredite de la polypose recto-colique. J. Genet. Hum.
9: 65-77,

[35052] 1960.

[35053] 5107.

[35054] Dunlop, M. G.; Wyllie, A. H.; Nakamura, Y.; Steel, C. M.; Evans,

[35055] H. J.; White, R. L.; Bird, C. C.: Genetic linkage map of six polymorphic

[35056] DNA markers around the gene for familial adenomatous polyposis on

[35057] chromosome 5. Am. J. Hum. Genet. 47: 982-987, 1990.

[35058] 5108.

[35059] Dunlop, M. G.; Wyllie, A. H.; Steel, C. M.; Piris, J.; Evans,

[35060] H. J.: Linked DNA markers for presymptomatic diagnosis of familial

[35061] adenomatous polyposis. Lancet 337: 313-316, 1991.

[35062] 5109.

[35063] Eccles, D. M.; Lunt, P. W.; Wallis, Y.; Griffiths, M.; Sandhu,

[35064] B.; McKay, S.; Morton, D.; Shea-Simonds, J.; Macdonald, F.: An unusually

[35065] severe phenotype for familial adenomatous polyposis. Arch. Dis. Child. 77:

[35066] 431-435, 1997.

[35067] 5110.

[35068] Endo, A.; Kasukawa, T.: Gardner syndrome and interstitial chromosome

[35069] deletion. (Letter) Am. J. Med. Genet. 28: 511-512, 1987.

[35070] 5111.

[35071] Erbe, R. W.; Welch, W. R.: Case records of the Mas-sachusetts

[35072] General Hospital. Weekly clinicopathological exercises. Case 47-1978. New

[35073] Eng. J. Med. 299: 1237-1245, 1978.

[35074] 5112.

[35075] Evans, D. G. R.; Guy, S. P.; Thakker, N.; Armstrong, J. G.; Dodd,

[35076] C.; Davies, D. R.; Babbs, C.; Clancy, T.; Warnes, T.; Sloan, P.; Taylor,

[35077] T. V.; Harris, R.: Non-penetrance and late appearance of polyps in

[35078] families with familial adenomatous polyposis. Gut 34: 1389-1393,

[35079] 1993.

[35080] 5113.

[35081] Fader, M.; Kline, S. N.; Spatz, S. S.; Zubrow, H. J.: Gardner's

[35082] syndrome (intestinal polyposis, osteomas, sebaceous cysts) and a new

[35083] dental discovery. Oral Surg. 15: 153-172, 1962.

[35084] 5114.

[35085] Fearnhead, N. S.; Britton, M. P.; Bodmer, W. F.: The ABC of APC. Hum.

[35086] Molec. Genet. 10: 721-733, 2001.

[35087] 5115.

[35088] Fialkow, P. J.: Clonal origin of human tumors. Ann. Rev. Med. 30:

[35089] 135-143, 1979.

[35090] 5116.

[35091] Fineman, R. M.; Morgan, M.; Burt, R. W.; Gardner, E. J.: Failure

[35092] to demonstrate a chromosome 2 deletion in adenomatous colorectal polyposis

[35093] patients. Cancer 53: 317-318, 1984.

[35094] 5117.

[35095] Fodde, R.; Edelmann, W.; Yang, K.; van Leeuwen, C.; Carl-
son, C.;
[35096] Renault, B.; Breukel, C.; Alt, E.; Lipkin, M.; Meera Khan, P.;
Kucherlapati,
[35097] R.: A targeted chain-termination mutation in the mouse
Apc gene results
[35098] in multiple intestinal tumors. Proc. Nat. Acad. Sci. 91:
8969-8973,
[35099] 1994.

[35100] 5118.

[35101] Fodde, R.; Kuipers, J.; Rosenberg, C.; Smits, R.; Kielman,
M.;
[35102] Gaspar, C.; van Es, J. H.; Breukel, C.; Wiegant, J.; Giles, R.
H.;
[35103] Clevers, H.: Mutations in the APC tumour suppressor gene
cause chromosomal
[35104] instability. Nature Cell Biol. 3: 433-438, 2001.

[35105] 5119.

[35106] Fodde, R.; van der Luijt, R.; Wijnen, J.; Tops, C.; van der
Klift,
[35107] H.; van Leeuwen-Cornelisse, I.; Griffioen, G.; Vasen, H.;
Meera Khan,

[35108] P.: Eight novel inactivating germ line mutations at the APC gene
[35109] identified by denaturing gradient gel electrophoresis. Genomics 13:
[35110] 1162-1168, 1992.

[35111] 5120.

[35112] Fraumeni, J. F., Jr.; Vogel, C. L.; Easton, J. M.: Sarcomas and
[35113] multiple polyposis in a kindred. A genetic variety of hereditary polyposis? Arch.
[35114] Intern. Med. 121: 57-61, 1968.

[35115] 5121.

[35116] Frayling, I. M.; Beck, N. E.; Ilyas, M.; Dove-Edwin, I.; Goodman,
[35117] P.; Pack, K.; Bell, J. A.; Williams, C. B.; Hodgson, S. V.; Thomas,
[35118] H. J. W.; Talbot, I. C.; Bodmer, W. F.; Tomlinson, I. P. M.: The
[35119] APC variants I1307K and E1317Q are associated with colorectal tumors,
[35120] but not always with a family history. Proc. Nat. Acad. Sci. 95:

[35121] 10722-10727, 1998.

[35122] 5122.

[35123] Friedl, W.; Meuschel, S.; Caspari, R.; Lamberti, C.; Krieger,

[35124] S.; Sengteller, M.; Propping, P.: Attenuated familial adeno-
matous

[35125] polyposis due to a mutation in the 3-prime part of the
APC gene: a

[35126] clue for understanding the function of the APC protein.
Hum. Genet. 97:

[35127] 579-584, 1996.

[35128] 5123.

[35129] Garber, J. E.; Li, F. P.; Kingston, J. E.; Krush, A. J.; Strong,

[35130] L. C.; Finegold, M. J.; Bertario, L.; Bulow, S.; Filippone, A.,
Jr.;

[35131] Gedde-Dahl, T., Jr.; Jarvinen, H. J.: Hepatoblastoma and
familial

[35132] adenomatous polyposis. J. Nat. Cancer Inst. 80:
1626-1628, 1988.

[35133] 5124.

[35134] Gardner, E. J.: A genetic and clinical study of intestinal
polyposis,

[35135] a predisposing factor for carcinoma of the colon and rectum. Am.

[35136] J. Hum. Genet. 3: 167-176, 1951.

[35137] 5125.

[35138] Gardner, E. J.: Discovery of the Gardner syndrome. Birth Defects

[35139] Orig. Art. Ser. VIII(2): 48-51, 1972.

[35140] 5126.

[35141] Gardner, E. J.: Follow-up study of a family group exhibiting

[35142] dominant inheritance for a syndrome including intestinal polyps, osteomas,

[35143] fibromas and epidermal cysts. Am. J. Hum. Genet. 14: 376-390, 1962.

[35144] 5127.

[35145] Gardner, E. J.; Plenk, H. P.: Hereditary pattern for multiple

[35146] osteomas in a family group. Am. J. Hum. Genet. 4: 31-36, 1952.

[35147] 5128.

[35148] Gardner, E. J.; Rogers, S. W.; Woodward, S.: Numerical and structural

[35149] chromosome aberrations in cultured lymphocytes and cu-
taneous fibroblasts
[35150] of patients with multiple adenomas of the colorectum.
Cancer 49:
[35151] 1413-1419, 1982.

[35152] 5129.

[35153] Gayther, S. A.; Wells, D.; SenGupta, S. B.; Chapman, P.;
Neale,
[35154] K.; Tsioupra, K.; Delhanty, J. D. A.: Regionally clustered
APC mutations
[35155] are associated with a severe phenotype and occur at a
high frequency
[35156] in new mutation cases of adenomatous polyposis coli.
Hum. Mol. Genet. 3:
[35157] 53-56, 1994.

[35158] 5130.

[35159] Giardiello, F. M.; Brensinger, J. D.; Petersen, G. M.; Luce,
M.
[35160] C.; Hylind, L. M.; Bacon, J. A.; Booker, S. V.; Parker, R. D.;
Hamilton,
[35161] S. R.: The use and interpretation of commercial APC gene
testing

[35162] for familial adenomatous polyposis. New Eng. J. Med. 336: 823-827,

[35163] 1997.

[35164] 5131.

[35165] Giardiello, F. M.; Hamilton, S. R.; Krush, A. J.; Piantadosi,

[35166] S.; Hylind, L. M.; Celano, P.; Booker, S. V.; Robinson, C. R.; Offerhaus,

[35167] G. J. A.: Treatment of colonic and rectal adenomas with sulindac

[35168] in familial adenomatous polyposis. New Eng. J. Med. 328: 1313-1316,

[35169] 1993.

[35170] 5132.

[35171] Giardiello, F. M.; Offerhaus, G. J. A.; Krush, A. J.; Booker,

[35172] S. V.; Tersmette, A. C.; Mulder, J.-W. R.; Kelley, C. N.; Hamilton,

[35173] S. R.: Risk of hepatoblastoma in familial adenomatous polyposis. J.

[35174] Pediat. 119: 766-768, 1991.

[35175] 5133.

[35176] Zuppan, P. J.; Hall, J. M.; Ponglikitmongkol, M.; Spielman, R.;
[35177] King, M. C.: Polymorphisms at the estrogen receptor (ESR) locus and
[35178] linkage relationships on chromosome 6q. (Abstract) Cyto-genet. Cell
[35179] Genet. 51: 1116 only, 1989.

[35180] 5134.

[35181] Apte, S. S.; Mattei, M.-G.; Olsen, B. R.: Cloning of the cDNA
[35182] encoding human tissue inhibitor of metalloproteinases-3 (TIMP-3) and
[35183] mapping of the TIMP3 gene to chromosome 22. Genomics 19: 86-90,
[35184] 1994.

[35185] 5135.

[35186] Janssens, K.; Gershoni-Baruch, R.; Guanabens, N.; Migone, N.;
[35187] Ralston, S.; Bonduelle, M.; Lissens, W.; Van Maldergem, L.; Vanhoenacker,
[35188] F.; Verbruggen, L.; Van Hul, W.: Mutations in the gene en-coding the

[35189] latency-associated peptide of TGF-beta-1 cause Camu-
rati-Engelmann
[35190] disease. Nature Genet. 26: 273-275, 2000.

[35191] 5136.

[35192] Lang, F.; Klingel, K.; Wagner, C. A.; Stegen, C.; Warntges,
S.;
[35193] Friedrich, B.; Lanzendorfer, M.; Melzig, J.; Moschen, I.;
Steuer,
[35194] S.; Waldegger, S.; Sauter, M.; and 9 others: Deranged tran-
scriptional
[35195] regulation of cell-volume-sensitive kinase hSGK in dia-
betic nephropathy. Proc.
[35196] Nat. Acad. Sci. 97: 8157-8162, 2000.

[35197] 5137.

[35198] Li, B.; Khanna, A.; Sharma, V.; Singh, T.; Suthanthiran, M.;
August,
[35199] P.: TGF-beta-1 DNA polymorphisms, protein levels, and
blood pressure. Hypertension 33:
[35200] 271-275, 1999.

[35201] 5138.

[35202] Marquardt, H.; Lioubin, M. N.; Ikeda, T.: Complete amino

acid
[35203] sequence of human transforming growth factor type beta-2. J. Biol.
[35204] Chem. 262: 12127-12131, 1987.

[35205] 5139.

[35206] Reeves, W. B.; Andreoli, T. E.: Transforming growth factor beta
[35207] contributes to progressive diabetic nephropathy. Proc. Nat. Acad.
[35208] Sci. 97: 7667-7669, 2000.

[35209] 5140.

[35210] Roberts, A. B.; Sporn, M. B.; Assoian, R. K.; Smith, J. M.; Roche,
[35211] N. S.; Wakefield, L. M.; Heine, U. I.; Liotta, L. A.; Falanga, V.;
[35212] Kehrl, J. H.; Fauci, A. S.: Transforming growth factor type beta:
[35213] rapid induction of fibrosis and angiogenesis in vivo and stimulation
[35214] of collagen formation in vitro. Proc. Nat. Acad. Sci. 83: 4167-4171,
[35215] 1986.

[35216] 5141.

[35217] Sharma, K.; Jin, Y.; Guo, J.; Ziyadeh, F. N. :Diabetes 45:

[35218] 522-530, 1996.

[35219] 5142.

[35220] Shenkar, R.; Coulson, W. F.; Abraham, E.: Hemorrhage and resuscitation

[35221] induce alterations in cytokine expression and the development of acute

[35222] lung injury. Am. J. Resp. Cell. Mol. Biol. 10: 290-297, 1994.

[35223] 5143.

[35224] Sporn, M. B.; Roberts, A. B.; Wakefield, L. M.; Assoian, R. K.

[35225] : Transforming growth factor-beta: biological function and chemical

[35226] structure. Science 233: 532-534, 1986.

[35227] 5144.

[35228] Suthanthiran, M.; Khanna, A.; Cukran, D.; Adhikarla, R.; Sharma,

[35229] V. K.; Singh, T.; August, P.: Transforming growth factor-beta-1 hyperexpression

[35230] in African American end-stage renal disease patients. Kidney Int. 53:

[35231] 639-644, 1998.

[35232] 5145.

[35233] Suthanthiran, M.; Li, B.; Song, J. O.; Ding, R.; Sharma, V. K.;

[35234] Schwartz, J. E.; August, P.: Transforming growth factor-beta-1 hyperexpression

[35235] in African-American hypertensives: a novel mediator of hypertension

[35236] and/or target organ damage. Proc. Nat. Acad. Sci. 97: 3479-3484,

[35237] 2000.

[35238] 5146.

[35239] Waldegger, S.; Klingel, K.; Barth, P.; Sauter, M.; Lanzendorfer,

[35240] M.; Kandolf, R.; Lang, F.: h-sgk serine-threonine protein kinase

[35241] gene as transcriptional target of transforming growth factor beta

[35242] in human intestine. Gastroenterology 116: 1081-1088, 1999.

[35243] 5147.

[35244] Watanabe, Y.; Kinoshita, A.; Yamada, T.; Ohta, T.; Kishino, T.;

[35245] Matsumoto, N.; Ishikawa, M.; Niikawa, N.; Yoshiura, K.-I.: A catalog

[35246] of 106 single-nucleotide polymorphisms (SNPs) and 11 other types of

[35247] variations in genes for transforming growth factor-beta-1 (TGF-beta-1)

[35248] and its signaling pathway. J. Hum. Genet. 47: 478-483, 2002.

[35249] 5148.

[35250] Wyss-Coray, T.; Lin, C.; Yan, F.; Yu, G.; Rohde, M.; Mc-Conlogue,

[35251] L.; Masliah, E.; Mucke, L.: TGF-beta-1 promotes microglial amyloid-beta

[35252] clearance and reduces plaque burden in transgenic mice. Nature Med. 7:

[35253] 612-618, 2001.

[35254] 5149.

[35255] Ziyadeh, F. N.; Hoffman, B. B.; Han, D. C.; Iglesias-de la Cruz,

[35256] M. C.; Hong, S. W.; Isono, M.; Chen, S.; McGowan, T. A.; Sharma, K.

[35257] : Long-term prevention of renal insufficiency, excess matrix gene

[35258] expression, and glomerular mesangial matrix expansion by treatment

[35259] with monoclonal antitransforming growth factor-beta antibody in db/db

[35260] diabetic mice. Proc. Nat. Acad. Sci. 97: 8015-8020, 2000.

[35261] 5150.

[35262] Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocytoplasmic shuttling

[35263] of Smads 2, 3, and 4 permits sensing of TGF-beta receptor activity. Molec.

[35264] Cell 10: 283-294, 2002.

[35265] 5151.

[35266] Johnson, D. W.; Qumsiyeh, M.; Benkhalifa, M.; Marchuk, D. A.:

[35267] Assignment of human transforming growth factor-beta type I and type

[35268] III receptor genes (TGFBR1 and TGFBR3) to 9q33-q34 and 1p32-p33, respectively. Genomics 28:

[35269] 356-357, 1995.

[35270] 5152.

[35271] Dallapiccola, B.; Novelli, G.; Giannotti, A.: Deletion 2q31.3-2q33.3:

[35272] gene dosage effect of ribulose 5-phosphate 3-epimerase. Hum. Genet. 79:

[35273] 92 only, 1988.

[35274] 5153.

[35275] Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of the

[35276] gene for ribulose-5-phosphate 3-epimerase (RPE) to human chromosome

[35277] 2. (Abstract) Cytogenet. Cell Genet. 32: 268 only, 1982.

[35278] 5154.

[35279] Donald, L. J.; Wang, H. S.; Hamerton, J. L.: A ribulose-5-phosphate-3-epimerase

[35280] (RPE) locus is on human chromosome 2. Cytogenet. Cell Genet. 33:

[35281] 261-263, 1982.

[35282] 5155.

[35283] Gross, M.-S.; Weil, D.; Van Cong, N.; Finaz, C.; Foubert, C.; Cochet,

[35284] C.; Parisi, I.; de Grouchy, J.; Frezal, J.: Localisation du gene

[35285] de la ribulose-5-phosphate-3-epimerase (RPE) sur le segment 2q32-2qter

[35286] par hybridation cellulaire interspecifique. Ann. Genet. 25: 87-91,

[35287] 1982.

[35288] 5156.

[35289] Miyazaki, K.; Yamanaka, T.; Ogasawara, N.: Interstitial deletion

[35290] 2q32.1-q34 in a child with half normal activity of ribulose 5-phosphate

[35291] 3-epimerase (RPE). J. Med. Genet. 25: 850-851, 1988.

[35292] 5157.

[35293] Helfgott, S. M.; Mosciscki, R. A.; San Martin, J.; Lorenzo, C.;

[35294] Kieval, R.; McKenna, M.; Nadol, J.; Trentham, D. E.: Correlation

[35295] between antibodies to type II collagen and treatment outcome in bilateral

[35296] progressive sensorineural hearing loss. Lancet 337:
387-389, 1991.

[35297] 5158.

[35298] Lee, B.; Vissing, H.; Ramirez, F.; Rogers, D.; Rimoin, D.:
Identification

[35299] of the molecular defect in a family with spondyloepiphy-
seal dysplasia. Science 244:

[35300] 978-980, 1989.

[35301] 5159.

[35302] Tiller, G. E.; Rimoin, D. L.; Murray, L. W.; Cohn, D. H.: Tan-
dem

[35303] duplication within a type II collagen gene (COL2A1) exon
in an individual

[35304] with spondyloepiphyseal dysplasia. Proc. Nat. Acad. Sci.
87: 3889-3893,

[35305] 1990.

[35306] 5160.

[35307] Yoo, T. J.; Tomoda, K.; Stuart, J. M.; Cremer, M. A.;
Townes,

[35308] A. S.; Kang, A. H.: Type II collagen-induced autoimmune
sensorineural

[35309] hearing loss and vestibular dysfunction in rats. Ann. Otol. Rhinol.

[35310] Laryng. 92: 267-271, 1983.

[35311] 5161.

[35312] Begley, C. G.; Visvader, J.; Green, A. R.; Aplan, P. D.; Met-

[35313] calf, D.; Kirsch, I. R.; Gough, N. M.: Molecular cloning and chromosomal

[35314] localization of the murine homolog of the human helix-loop-helix gene

[35315] SCL. Proc. Nat. Acad. Sci. 88: 869-873, 1991.

[35316] 5162.

[35317] Finger, L. R.; Kagan, J.; Christopher, G.; Kurtzberg, J.; Her-

[35318] shfield, M. S.; Nowell, P. C.; Croce, C. M.: Involvement of the TCL5 gene

[35319] on human chromosome 1 in T-cell leukemia and melanoma. Proc. Nat.

[35320] Acad. Sci. 86: 5039-5043, 1989.

[35321] 5163.

[35322] Gottgens, B.; Barton, L. M.; Chapman, M. A.; Sinclair, A.

M.; Knudsen,
[35323] B.; Grafham, D.; Gilbert, J. G. R.; Rogers, J.; Bentley, D. R.; Green,
[35324] A. R.: Transcriptional regulation of the stem cell leukemia gene
[35325] (SCL)--comparative analysis of five vertebrate SCL loci. Genome Res. 12:
[35326] 749-759, 2002.

[35327] 5164.

[35328] Gottgens, B.; Gilbert, J. G.; Barton, L. M.; Grafham, D.; Rogers,
[35329] J.; Bentley, D. R.; Green, A. R.: Long-range comparison of human
[35330] and mouse SCL loci: localized regions of sensitivity to restriction
[35331] endonucleases correspond precisely with peaks of conserved noncoding
[35332] sequences. Genome Res. 11: 87-97, 2001.

[35333] 5165.

[35334] Kocher, O.; Cheresh, P.; Lee, S. W.: Identification and partial
[35335] characterization of a novel membrane-associated protein

(MAP17) up-regulated
[35336] in human carcinomas and modulating cell replication and tumor growth. Am.
[35337] J. Path. 149: 493-500, 1996.

[35338] 5166.

[35339] Kozak, M.: An analysis of 5-prime-noncoding sequences from 699
[35340] vertebrate messenger RNAs. Nucleic Acids Res. 15: 8125-8148, 1987.
[35341] 5167.

[35342] Kurtzberg, J.; Bigner, S. H.; Hershfield, M. S.: Establish-ment
[35343] of the DU.528 human lymphohemopoietic stem cell line. J. Exp. Med. 162:
[35344] 1561-1578, 1985.

[35345] 5168.

[35346] Robb, L.; Lyons, I.; Li, R.; Hartley, L.; Kontgen, F.; Harvey,

[35347] R. P.; Metcalf, D.; Begley, C. G.: Absence of yolk sac hematopoiesis
[35348] from mice with a targeted disruption of the scl gene. Proc. Nat.

[35349] Acad. Sci. 92: 7075-7079, 1995.

[35350] 5169.

[35351] Shivdasani, R. A.; Mayer, E. L.; Orkin, S. H.: Absence of blood

[35352] formation in mice lacking the T-cell leukaemia oncoprotein tal-1/SCL. Nature 373:

[35353] 432-434, 1995.

[35354] 5170.

[35355] Sinclair, A. M.; Bench, A. J.; Bloor, A. J. C.; Li, J.; Gottgens,

[35356] B.; Stanley, M. L.; Miller, J.; Piltz, S.; Hunter, S.; Nacheva, E.

[35357] P.; Sanchez, M.-J.; Green, A. R.: Rescue of the lethal scl-/- phenotype

[35358] by the human SCL locus. Blood 99: 3931-3938, 2002.

[35359] 5171.

[35360] Williams, M.; Lyu, M. S.; Yang, Y. L.; Lin, E. P.; Dunbrack, R.;

[35361] Birren, B.; Cunningham, J.; Hunter, K.: Ier5, a novel member of the

[35362] slow-kinetics immediate-early genes. Genomics 44: 327-334, 1999.

[35363] 5172.

[35364] Xia, Y.; Brown, L.; Yang, C. Y.-C.; Tsou Tsan, J.; Siciliano,

[35365] M. J.; Espinosa, R., III; Le Beau, M. M.; Baer, R. J.: TAL2, a helix-loop-helix

[35366] gene activated by the (7;9)(q34;q32) translocation in human T-cell

[35367] leukemia. Proc. Nat. Acad. Sci. 88: 11416-11420, 1991.

[35368] 5173.

[35369] Artandi, S. E.; Chang, S.; Lee, S.-L.; Alson, S.; Gottlieb, G.

[35370] J.; Chin, L.; DePinho, R. A.: Telomere dysfunction promotes non-reciprocal

[35371] translocations and epithelial cancers in mice. Nature 406: 641-645,

[35372] 2000.

[35373] 5174.

[35374] Carrero-Valenzuela, R. D.; Klein, M. L.; Weleber, R. G.; Murphey,

[35375] W. H.; Litt, M.: Sorsby fundus dystrophy: a family with the ser181cys

[35376] mutation of the tissue inhibitor of metalloproteinases 3. Arch. Ophthal. 114:

[35377] 737-738, 1996.

[35378] 5175.

[35379] Felbor, U.; Stohr, H.; Amann, T.; Schonherr, U.; Weber, B. H. F.

[35380] : A novel ser156cys mutation in the tissue inhibitor of metalloproteinase-3

[35381] (TIMP3) in Sorsby's fundus dystrophy with unusual clinical features. Hum.

[35382] Molec. Genet. 4: 2415-2416, 1995.

[35383] 5176.

[35384] Langton, K. P.; Barker, M. D.; McKie, N.: Localization of the

[35385] functional domains of human tissue inhibitor of metallo-proteinases-3

[35386] and the effects of a Sorsby's fundus dystrophy mutation. J. Biol.

[35387] Chem. 273: 16778-16781, 1998.

[35388] 5177.

[35389] Langton, K. P.; McKie, N.; Curtis, A.; Goodship, J. A.; Bond, P.

[35390] M.; Barker, M. D.; Clarke, M.: A novel tissue inhibitor of metalloproteinases-3

[35391] mutation reveals a common molecular phenotype in Sorsby's fundus dystrophy. J.

[35392] Biol. Chem. 275: 27027-27031, 2000.

[35393] 5178.

[35394] Osman, M.; Tortorella, M.; Londei, M.; Quaratino, S.: Ex-

pression

[35395] of matrix metalloproteinases and tissue inhibitors of met-

alloproteinases

[35396] define the migratory characteristics of human monocyte-de-

rived dendritic

[35397] cells. Immunology 105: 73-82, 2002.

[35398] 5179.

[35399] Stohr, H.; Roomp, K.; Felbor, U.; Weber, B. H. F.: Genomic organization

[35400] of the human tissue inhibitor of metalloproteinases-3 (TIMP3). Genome

[35401] Res. 5: 483-487, 1995.

[35402] 5180.

[35403] Wijesuriya, S. D.; Evans, K.; Jay, M. R.; Davison, C.; Weber,

[35404] B. H. F.; Bird, A. C.; Bhattacharya, S. S.; Gregory, C. Y.: Sorsby's

[35405] fundus dystrophy in the British Isles: demonstration of a striking
[35406] founder effect by microsatellite-generated haplotypes. Genome Res. 6:
[35407] 92-101, 1996.

[35408] 5181.

[35409] Wilde, C. G.; Hawkins, P. R.; Coleman, R. T.; Levine, W. B.; Delegeane,
[35410] A. M.; Okamoto, P. M.; Ito, L. Y.; Scott, R. W.; Seilhamer, J. J.
[35411] : Cloning and characterization of human tissue inhibitor of metalloproteinases-3. DNA
[35412] Cell Biol. 13: 711-718, 1994.

[35413] 5182.

[35414] Durick, K.; Gill, G. N.; Taylor, S. S.: Shc and Enigma are both
[35415] required for mitogenic signaling by Ret/ptc2. Molec. Cell. Biol. 18:
[35416] 2298-2308, 1998.

[35417] 5183.

[35418] Saadat, M.; Nomoto, K.; Mizuno, Y.; Kikuchi, K.; Yoshida,

M. C.

[35419] : Assignment of the gene encoding type 1-gamma protein phosphatase

[35420] catalytic subunit (PPP1CC) on human, rat, and mouse chromosomes. Jpn.

[35421] J. Hum. Genet. 41: 159-165, 1996.

[35422] 5184.

[35423] Cohen, P.; Cohen, P. T. W.: Protein phosphatases come of age. J.

[35424] Biol. Chem. 264: 21435-21438, 1989.

[35425] 5185.

[35426] Groves, M. R.; Hanlon, N.; Turowski, P.; Hemmings, B. A.; Barford,

[35427] D.: The structure of the protein phosphatase 2A PR65/A subunit reveals

[35428] the conformation of its 15 tandemly repeated HEAT mo- tifs. Cell 96:

[35429] 99-110, 1999.

[35430] 5186.

[35431] Jones, T. A.; Barker, H. M.; da Cruz e Silva, E. F.; Mayer- Jaekel,

[35432] R. E.; Hemmings, B. A.; Spurr, N. K.; Sheer, D.; Cohen, P. T. W.:

[35433] Localization of the genes encoding the catalytic subunits of protein

[35434] phosphatase 2A to human chromosome bands 5q23-q31 and 8p12-p11.2,

[35435] respectively. Cytogenet. Cell Genet. 63: 35-41, 1993.

[35436] 5187.

[35437] Stone, S. R.; Mayer, R.; Wernet, W.; Maurer, F.; Hofsteenge, J.;

[35438] Hemmings, B. A.: The nucleotide sequence of the cDNA encoding the

[35439] human lung protein phosphatase 2A alpha catalytic sub-unit. Nucleic

[35440] Acids Res. 16: 11365 only, 1988.

[35441] 5188.

[35442] Imbert, A.; Chaffanet, M.; Essioux, L.; Noguchi, T.; Ade-laide,

[35443] J.; Kerangueven, F.; Le Paslier, D.; Bonaiti-Pellie, C.; Sobol, H.;

[35444] Birnbaum, D.; Pebusque, M.-J.: Integrated map of the chromosome 8p12-p21

[35445] region, a region involved in human cancers and Werner syndrome. Genomics 32:
[35446] 29-38, 1996.

[35447] 5189.

[35448] Lu, S.; Day, N. E.; Degos, L.; Lepage, V.; Wang, P.-C.; Chan, S.-H.;
[35449] Simons, M.; McKnight, B.; Easton, D.; Zeng, Y.; de-The, G.: Linkage
[35450] of a nasopharyngeal carcinoma susceptibility locus to the HLA region. Nature 346:
[35451] 470-471, 1990.

[35452] 5190.

[35453] Adamkiewicz, T. V.; McSherry, C.; Bach, F. H.; Houchins, J. P.
[35454] : Natural killer lectin-like receptors have divergent car-boxy-termini,
[35455] distinct from C-type lectins. Immunogenetics 39: 218 only, 1994.
[35456] 5191.

[35457] Houchins, J. P.; Yabe, T.; McSherry, C.; Bach, F. H.: DNA sequence

[35458] analysis of NKG2, a family of related cDNA clones encoding type II
[35459] integral membrane proteins on human natural killer cells. J. Exp.
[35460] Med. 173: 1017-1020, 1991.

[35461] 5192.

[35462] Plougastel, B.; Jones, T.; Trowsdale, J.: Genomic structure, chromosome
[35463] location, and alternative splicing of the human NKG2A gene. Immunogenetics 44:
[35464] 286-291, 1996.

[35465] 5193.

[35466] Plougastel, B.; Trowsdale, J.: Sequence analysis of a 62-kb region
[35467] overlapping the human KLRC cluster of genes. Genomics 49: 193-199,
[35468] 1998.

[35469] 5194.

[35470] Renedo, M.; Arce, I.; Rodriguez, A.; Carretero, M.; Lanier, L.
[35471] L.; Lopez-Botet, M.; Fernandez-Ruiz, E.: The human natu-

ral killer
[35472] gene complex is located on chromosome 12p12-p13. Immunogenetics 46:
[35473] 307-311, 1997.

[35474] 5195.

[35475] Yabe, T.; McSherry, C.; Bach, F. H.; Fisch, P.; Schall, R. P.;

[35476] Sondel, P. M.; Houchins, J. P.: A multigene family on human chromosome
[35477] 12 encodes natural killer-cell lectins. Immunogenetics 37: 455-460,
[35478] 1993.

[35479] 5196.

[35480] Allamand, V.; Sunada, Y.; Salih, M. A. M.; Straub, V.; Ozo, C.
[35481] O.; Al-Turaiki, M. H. S.; Akbar, M.; Kolo, T.; Colognato, H.; Zhang,
[35482] X.; Sorokin, L. M.; Yurchenco, P. D.; Tryggvason, K.; Campbell, K.
[35483] P.: Mild congenital muscular dystrophy in two patients with an internally
[35484] deleted laminin alpha-2-chain. Hum. Molec. Genet. 6:

747-752, 1997.

[35485] 5197.

[35486] Bushby, K.; Anderson, L. V. B.; Pollitt, C.; Naom, I.; Muntoni,

[35487] F.; Bindoff, L.: Abnormal merosin in adults: a new form of late onset

[35488] muscular dystrophy not linked to chromosome 6q2. Brain 121: 581-588,

[35489] 1998.

[35490] 5198.

[35491] D'Alessandro, M.; Naom, I.; Ferlini, A.; Sewry, C.; Dubowitz, V.;

[35492] Muntoni, F.: Is there selection in favour of heterozygotes in families

[35493] with merosin-deficient congenital muscular dystrophy? Hum. Genet. 105:

[35494] 308-313, 1999.

[35495] 5199.

[35496] Ehrig, K.; Leivo, I.; Argraves, W. S.; Ruoslahti, E.; Engvall,

[35497] E.: Merosin, a tissue-specific basement membrane protein, is a laminin-like

[35498] protein. Proc. Nat. Acad. Sci. 87: 3264-3268, 1990.

[35499] 5200.

[35500] Hayashi, Y. K.; Koga, R.; Tsukahara, T.; Ishii, H.; Matsuishi,

[35501] T.; Yamashita, Y.; Nonaka, I.; Arahata, K.: Deficiency of laminin

[35502] alpha-2-chain mRNA in muscle in a patient with merosin-negative congenital

[35503] muscular dystrophy. Muscle Nerve 18: 1027-1030, 1995.

[35504] 5201.

[35505] Helbling-Leclerc, A.; Zhang, X.; Topaloglu, H.; Cruaud, C.; Tesson,

[35506] F.; Weissenbach, J.; Tome, F. M. S.; Schwartz, K.; Fardeau, M.; Tryggvason,

[35507] K.; Guicheney, P.: Mutations in the laminin alpha-2-chain gene (LAMA2)

[35508] cause merosin-deficient congenital muscular dystrophy. Nature Genet. 11:

[35509] 216-218, 1995.

[35510] 5202.

[35511] Hillaire, D.; Leclerc, A.; Faure, S.; Topaloglu, H.; Chian-nilkulchai,

[35512] N.; Guicheney, P.; Grinas, L.; Legos, P.; Philpot, J.; Evangelista,

[35513] T.; Routon, M.-C.; Mayer, M.; Pellissier, J.-F.; Estournet, B.; Barois,

[35514] A.; Hentati, F.; Feingold, N.; Beckmann, J. S.; Dubowitz, V.; Tome,

[35515] F. M. S.; Fardeau, M.: Localization of merosin-negative congenital

[35516] muscular dystrophy to chromosome 6q2 by homozygosity mapping. Hum.

[35517] Molec. Genet. 3: 1657-1661, 1994.

[35518] 5203.

[35519] Jones, K. J.; Morgan, G.; Johnston, H.; Tobias, V.; Ouvrier, R.

[35520] A.; Wilkinson, I.; North, K. N.: The expanding phenotype of laminin

[35521] alpha-2 chain (merosin) abnormalities: case series and review. J.

[35522] Med. Genet. 38: 649-657, 2001.

[35523] 5204.

[35524] Kuang, W.; Xu, H.; Vachon, P. H.; Liu, L.; Loechel, F.; Wewer,

[35525] U. M.; Engvall, E.: Merosin-deficient congenital muscular dystrophy:
[35526] partial genetic correction in two mouse models. J. Clin. Invest. 102:
[35527] 844-852, 1998.

[35528] 5205.

[35529] Mercuri, E.; Dubowitz, L.; Berardinelli, A.; Pennock, J.; Jongmans,
[35530] M.; Henderson, S.; Muntoni, F.; Sewry, C.; Philpot, J.; Dubowitz,
[35531] V.: Minor neurological and perceptuo-motor deficits in children with
[35532] congenital muscular dystrophy: correlation with brain MRI changes. Neuropediatrics 26:
[35533] 156-162, 1995.

[35534] 5206.

[35535] Mercuri, E.; Muntoni, F.; Berardinelli, A.; Pennock, J.; Sewry,
[35536] C.; Philpot, J.; Dubowitz, V.: Somatosensory and visual evoked potentials
[35537] in congenital muscular dystrophy: correlation with MRI changes and

[35538] muscle merosin status. Neuropediatrics 26: 3-7, 1995.

[35539] 5207.

[35540] Moll, J.; Barzaghi, P.; Lin, S.; Bezakova, G.; Lochmuller, H.;

[35541] Engvall, E.; Muller, U.; Ruegg, M. A.: An agrin minigene rescues

[35542] dystrophic symptoms in a mouse model for congenital muscular dystrophy. Nature 413:

[35543] 302-307, 2001.

[35544] 5208.

[35545] Naom, I.; D'Alessandro, M.; Sewry, C.; Ferlini, A.; Topaloglu,

[35546] H.; Helbling-Leclerc, A.; Guicheney, P.; Schwartz, K.; Ak-coren, Z.;

[35547] Dubowitz, V.; Muntoni, F.: The role of immunocytochem-istry and linkage

[35548] analysis in the prenatal diagnosis of merosin-deficient congenital

[35549] muscular dystrophy. Hum. Genet. 99: 535-540, 1997.

[35550] 5209.

[35551] Ng, V.; Zanazzi, G.; Timpl, R.; Talts, J. F.; Salzer, J. L.; Brennan,

[35552] P. J.; Rambukkana, A.: Role of the cell wall phenolic gly-
colipid-1

[35553] in the peripheral nerve predilection of Mycobacterium lep-
rae. Cell 103:

[35554] 511-524, 2000.

[35555] 5210.

[35556] Philpot, J.; Cowan, F.; Pennock, J.; Sewry, C.; Dubowitz, V.;

[35557] Bydder, G.; Muntoni, F.: Merosin-deficient congenital
muscular dystrophy:

[35558] the spectrum of brain involvement on magnetic resonance
imaging. Neuromusc.

[35559] Disord. 9: 81-85, 1999.

[35560] 5211.

[35561] Sallinen, R.; Kuang, W.; Engvall, E.; Palotie, A.; Wessman,
M.;

[35562] Horelli-Kuitunen, N.: Assignment of laminin alpha 2-chain
gene (Lama2)

[35563] to mouse chromosome 10A4-B1 by fluorescence in situ
hybridization. Cytogenet.

[35564] Cell Genet. 87: 195-196, 1999.

[35565] 5212.

[35566] Shorer, Z.; Philpot, J.; Muntoni, F.; Sewry, C.; Dubowitz, V.

[35567] : Demyelinating peripheral neuropathy in merosin-deficient congenital

[35568] muscular dystrophy. J. Child Neurol. 10: 472-475, 1995.

[35569] 5213.

[35570] Sunada, Y.; Bernier, S. M.; Kozak, C. A.; Yamada, Y.; Campbell,

[35571] K. P.: Deficiency of merosin in dystrophic dy mice and genetic linkage

[35572] of laminin M chain gene to dy locus. J. Biol. Chem. 269: 13729-13732,

[35573] 1994.

[35574] 5214.

[35575] Sunada, Y.; Edgar, T. S.; Lotz, B. P.; Rust, R. S.; Campbell,

[35576] K. P.: Merosin-negative congenital muscular dystrophy associated

[35577] with extensive brain abnormalities. Neurology 45: 2084-2089, 1995.

[35578] 5215.

[35579] Taratuto, A. L.; Lubieniecki, F.; Diaz, D.; Schultz, M.; Ruggieri,

[35580] V.; Saccoliti, M.; Dubrovsky, A.: Merosin-deficient congenital muscular
[35581] dystrophy associated with abnormal cerebral cortical gyration: an
[35582] autopsy study. Neuromusc. Disord. 9: 86-94, 1999.

[35583] 5216.

[35584] Tome, F. M. S.; Evangelista, T.; Leclerc, A.; Sunada, Y.; Manole,
[35585] E.; Estournet, B.; Barois, A.; Campbell, K. P.; Fardeau, M.: Congenital
[35586] muscular dystrophy with merosin deficiency. Comp. Rend. Acad. Sci.
[35587] (Paris) 317: 351-357, 1994.

[35588] 5217.

[35589] Vuolteenaho, R.; Nissinen, M.; Sainio, K.; Byers, M.; Eddy, R.;
[35590] Hirvonen, H.; Shows, T. B.; Sariola, H.; Engvall, E.; Tryggvason,
[35591] K.: Human laminin M chain (merosin): complete primary structure,
[35592] chromosomal assignment, and expression of the M and A chain in human

[35593] fetal tissues. J. Cell Biol. 124: 381-394, 1994.

[35594] 5218.

[35595] Lindner, T. H.; Njolstad, P. R.; Horikawa, Y.; Bostad, L.; Bell,

[35596] G. I.; Sovik, O.: A novel syndrome of diabetes mellitus, re-

nal dysfunction

[35597] and genital malformation associated with a partial dele-

tion of the

[35598] pseudo-POU domain of hepatocyte nuclear factor-1-beta.

Hum. Molec.

[35599] Genet. 8: 2001-2008, 1999.

[35600] 5219.

[35601] Anderson, L. A.; Hall, J. M.; Lee, M. K.; Lebo, R. V.; King, M.

[35602] C.: Polymorphisms near chromosome 1cen. (Abstract) Cy-

togenet. Cell

[35603] Genet. 51: 951, 1989.

[35604] 5220.

[35605] Gendler, S. J.; Lancaster, C. A.; Taylor-Papadimitriou, J.; Duhig,

[35606] T.; Peat, N.; Burchell, J.; Pemberton, L.; Lalani, E.-N.; Wil-

son,

[35607] D.: Molecular cloning and expression of human tumor-associated polymorphic

[35608] epithelial mucin. J. Biol. Chem. 265: 15286-15293, 1990.

[35609] 5221.

[35610] Karlsson, S.; Swallow, D. M.; Griffiths, B.; Corney, G.; Hopkinson,

[35611] D. A.; Dawnay, A.; Cartron, J. P.: A genetic polymorphism of a human

[35612] urinary mucin. Ann. Hum. Genet. 47: 263-269, 1983.

[35613] 5222.

[35614] Kingsmore, S. F.; Spicer, A. P.; Gendler, S. J.; Seldin, M. F.

[35615] : Genetic mapping of the tumor-associated mucin 1 gene on mouse chromosome

[35616] 3. Mammalian Genome 6: 378, 1995.

[35617] 5223.

[35618] Lan, M. S.; Batra, S. K.; Qi, W.-N.; Metzgar, R. S.; Hollingsworth,

[35619] M. A.: Cloning and sequencing of a human pancreatic tumor mucin cDNA. J.

[35620] Biol. Chem. 265: 15294-15299, 1990.

[35621] 5224.

[35622] Ligtenberg, M. J.; Gennissen, A. M.; Vos, H. L.; Hilkens, J.:

[35623] A single nucleotide polymorphism in an exon dictates allele dependent

[35624] differential splicing of episialin mRNA. Nucleic Acids Res. 19:

[35625] 297-301, 1991.

[35626] 5225.

[35627] Ligtenberg, M. J. L.; Vos, H. L.; Gennissen, A. M. C.; Hilkens,

[35628] J.: Episialin, a carcinoma-associated mucin, is generated by a polymorphic

[35629] gene encoding splice variants with alternative amino termini. J.

[35630] Biol. Chem. 265: 5573-5578, 1990.

[35631] 5226.

[35632] Middleton-Price, H.; Gendler, S.; Malcolm, S.: Close linkage of

[35633] PUM and SPTA within chromosome band 1q21. Ann. Hum. Genet. 52: 273-278,

[35634] 1988.

[35635] 5227.

[35636] Pratt, W. S.; Islam, I.; Swallow, D. M.: Two additional polymorphisms

[35637] within the hypervariable MUC1 gene: association of alleles either

[35638] side of the VNTR region. Ann. Hum. Genet. 60: 21-28, 1996.

[35639] 5228.

[35640] Silva, F.; Carvalho, F.; Peixoto, A.; Seixas, M.; Almeida, R.;

[35641] Carneiro, F.; Mequita, P.; Figueiredo, C.; Nogueira, C.; Swallow,

[35642] D. M.; Amorim, A.; David, L.: MUC1 gene polymorphism in the gastric

[35643] carcinogenesis pathway. Europ. J. Hum. Genet. 9: 548-552, 2001.

[35644] 5229.

[35645] Swallow, D. M.; Gendler, S.; Griffiths, B.; Corney, G.; Taylor-Papadimitriou

[35646] , J.; Bramwell, M. E.: The human tumour-associated epithelial mucins

[35647] are coded by an expressed hypervariable gene locus PUM. Nature 328:

[35648] 82-84, 1987.

[35649] 5230.

[35650] Swallow, D. M.; Gendler, S.; Griffiths, B.; Kearney, A.; Povey,

[35651] S.; Sheer, D.; Palmer, R. W.; Taylor-Papadimitriou, J.: The hypervariable

[35652] gene locus PUM, which codes for the tumour associated epithelial mucins,

[35653] is located on chromosome 1, within the region 1q21-24. (Abstract) Cytogenet.

[35654] Cell Genet. 46: 701, 1987.

[35655] 5231.

[35656] Swallow, D. M.; Gendler, S.; Griffiths, B.; Kearney, A.; Povey,

[35657] S.; Sheer, D.; Palmer, R. W.; Taylor-Papadimitriou, J.: The hypervariable

[35658] gene locus PUM, which codes for the tumour associated epithelial mucins,

[35659] is located on chromosome 1, within the region 1q21-q24. Ann. Hum.

[35660] Genet. 51: 289-294, 1987.

[35661] 5232.

[35662] Swallow, D. M.; Griffiths, B.; Noades, J.; Corney, G.: Link-
age
[35663] between the expressed hypervariable gene locus PUM and
the gene coding
[35664] for the Duffy blood group FY. Ann. Hum. Genet. 52:
269-271, 1988.

[35665] 5233.

[35666] Cole, S. P. C.; Bhardwaj, G.; Gerlach, J. H.; Mackie, J. E.;
Grant,
[35667] C. E.; Almquist, K. C.; Stewart, A. J.; Kurz, E. U.; Duncan, A.
M.
[35668] V.; Deeley, R. G.: Overexpression of a transporter gene in
a multidrug-resistant
[35669] human lung cancer cell line. Science 258: 1650-1654,
1992.

[35670] 5234.

[35671] Conrad, S.; Kauffmann, H.-M.; Ito, K.; Deeley, R. G.; Cole,
S.
[35672] P. C.; Schrenk, D.: Identification of human multidrug re-
sistance
[35673] protein 1 (MRP1) mutations and characterization of a

G671V substitution. J.
[35674] Hum. Genet. 46: 656-663, 2001.

[35675] 5235.

[35676] Grant, C. E.; Kurz, E. U.; Cole, S. P. C.; Deeley, R. G.: Anal-
ysis
[35677] of the intron-exon organization of the human multidrug-
resistance
[35678] protein gene (MRP) and alternative splicing of its mRNA.
Genomics 45:
[35679] 368-378, 1997.

[35680] 5236.

[35681] Lorico, A.; Bertola, A.; Baum, C.; Fodstad, O.; Rappa, G.: Role
[35682] of multidrug resistance protein 1 in protection from heavy metal oxyanions:
[35683] investigations in vitro and in Mrp1-deficient mice. Biochem. Biophys.
[35684] Res. Commun. 291: 617-622, 2002.

[35685] 5237.

[35686] Robbiani, D. F.; Finch, R. A.; Jager, D.; Muller, W. A.; Sar-
torelli,
[35687] A. C.; Randolph, G. J.: The leukotriene C4 transporter MRP1 regulates
[35688] CCL19 (MIP-3-beta, ELC)-dependent mobilization of den-
dritic cells
[35689] to lymph nodes. Cell 103: 757-768, 2000.

[35690] 5238.

[35691] Schultz, M. J.; Wijnholds, J.; Peppelenbosch, M. P.; Vervo-
ordeldonk,
[35692] M. J. B. M.; Speelman, P.; van Deventer, S. J. H.; Borst, P.; van
[35693] der Poll, T.: Mice lacking the multidrug resistance protein 1 are

[35694] resistant to Streptococcus pneumoniae-induced pneumo-
nia. J. Immun. 166:
[35695] 4059-4064, 2001.

[35696] 5239.

[35697] Zaman, G. J. R.; Flens, M. J.; van Leusden, M. R.; de Haas,
M.;
[35698] Mulder, H. S.; Lankelma, J.; Pinedo, H. M.; Scheper, R. J.;
Baas,
[35699] F.; Broxterman, H. J.; Borst, P.: The human multidrug re-
sistance-associated
[35700] protein MRP is a plasma membrane drug-efflux pump.
Proc. Nat. Acad.
[35701] Sci. 91: 8822-8826, 1994.

[35702] 5240.

[35703] le Gallic, L.; Fort, P.: Structure of the human ARHG locus
encoding
[35704] the Rho/Rac-like RhoG GTPase. Genomics 42: 157-160,
1997.
[35705] 5241.

[35706] Taviaux, S. A.; Vincent, S.; Fort, P.; Demaille, J. G.: Local-
ization

[35707] of ARHG, a member of the RAS homolog gene family, to 11p15.5-11p15.4
[35708] by fluorescence in situ hybridization. Genomics 16: 788-790, 1993.
[35709] 5242.
[35710] Vincent, S.; Jeanteur, P.; Fort, P.: Growth-regulated expression
[35711] of rhoG, a new member of the ras homolog gene family. Molec. Cell.
[35712] Biol. 12: 3138-3148, 1992.
[35713] 5243.
[35714] Allan, B. B.; Moyer, B. D.; Balch, W. E.: Rab1 recruitment of
[35715] p115 into a cis-SNARE complex: programming budding COPII vesicles
[35716] for fusion. Science 289: 444-448, 2000.
[35717] 5244.
[35718] Wedemeyer, N.; Lengeling, A.; Ronsiek, M.; Korthaus, D.; Baer,
[35719] K.; Wuttke, M.; Jockusch, H.: YAC contigs of the Rab1 and wobbler
[35720] (wr) spinal muscular atrophy gene region on proximal

mouse chromosome
[35721] 11 and of the homologous region on human chromosome 2p. Genomics 32:
[35722] 447-454, 1996.

[35723] 5245.

[35724] Rousseau-Merck, M. F.; Zahraoui, A.; Touchot, N.; Tavitian, A.;
[35725] Berger, R.: Chromosome assignment of four RAS-related RAB genes. Hum.
[35726] Genet. 86: 350-354, 1991.

[35727] 5246.

[35728] Rousseau-Merck, M.-F.; Zahraoui, A.; Touchot, N.; Tavitian, A.;
[35729] Berger, R.: Chromosome assignment of four RAS-related RAB genes. Hum.
[35730] Genet. 86: 350-354, 1991.

[35731] 5247.

[35732] Bucci, C.; Parton, R. G.; Mather, I. H.; Stunnenberg, H.; Simons,
[35733] K.; Hoflack, B.; Zerial, M.: The small GTPase rab5 functions as a

[35734] regulatory factor in the early endocytic pathway. Cell 70: 715-728,

[35735] 1992.

[35736] 5248.

[35737] Stenmark, H.; Vitale, G.; Ullrich, O.; Zerial, M.: Rabaptin-5

[35738] is a direct effector of the small GTPase Rab5 in endocytic membrane

[35739] fusion. Cell 83: 423-432, 1995.

[35740] 5249.

[35741] Xiao, G.-H.; Shoarinejad, F.; Jin, F.; Golemis, E. A.; Yeung, R.

[35742] S.: The tuberous sclerosis 2 gene product, tuberin, functions as

[35743] a Rab5 GTPase activating protein (GAP) in modulating endocytosis. J.

[35744] Biol. Chem. 272: 6097-6100, 1997.

[35745] 5250.

[35746] Aita, N.; Ishii, K.; Akamatsu, Y.; Ogasawara, Y.; Tanabe, S.:

[35747] Cloning and expression of human liver rhodanese cDNA. Biochem. Biophys.

[35748] Res. Commun. 231: 56-60, 1997.

[35749] 5251.

[35750] Cagianut, B.; Rhyner, K.; Furrer, W.; Schnebli, H. P.: Thio-
sulphate-sulphur
[35751] transferase (rhodanese) deficiency in Leber's hereditary
optic atrophy.
[35752] (Letter) Lancet II: 981-982, 1981.

[35753] 5252.

[35754] Nikoskelainen, E.: New aspects of the genetic, etiologic,
and
[35755] clinical puzzle of Leber's disease. Neurology 34:
1482-1484, 1984.
[35756] 5253.

[35757] Scott, E. M.; Wright, R. C.: Genetic polymorphism of rho-
danese
[35758] from human erythrocytes. Am. J. Hum. Genet. 32:
112-114, 1980.
[35759] 5254.

[35760] Weng, L.; Heinrikson, R. L.; Westley, J.: Active site cysteinyl

[35761] and arginyl residues of rhodanese. J. Biol. Chem. 253:
8109-8119,

[35762] 1978.

[35763] 5255.

[35764] Whitehouse, D. B.; Pilz, A. J.; Porta, G.; Hopkinson, D. A.: Rhodanese

[35765] isozymes in human tissues. Ann. Hum. Genet. 52: 1-10, 1988.

[35766] 5256.

[35767] Whitehouse, D. B.; Poole, C. J. M.; Kind, P. R. N.; Hopkin-son,

[35768] D. A.: Rhodanese isozymes in three subjects with Leber's optic neuropathy. J.

[35769] Med. Genet. 26: 113-115, 1989.

[35770] 5257.

[35771] al-Jandal, N.; Farrar, G. J.; Kiang, A.-S.; Humphries, M. M.; Bannon,

[35772] N.; Findlay, J. B. C.; Humphries, P.; Kenna, P. F.: A novel mutation

[35773] within the rhodopsin gene (thr94 to ile) causing autoso-mal dominant

[35774] congenital stationary night blindness. Hum. Mutat. 13: 75-81, 1999.

[35775] 5258.

[35776] Alloway, P. G.; Howard, L.; Dolph, P. J.: The formation of stable

[35777] rhodopsin-arrestin complexes induces apoptosis and photoreceptor cell

[35778] degeneration. Neuron 28: 129-138, 2000.

[35779] 5259.

[35780] Berson, E. L.; Rosner, B.; Sandberg, M. A.; Dryja, T. P.: Ocular

[35781] findings in patients with autosomal dominant retinitis pigmentosa

[35782] and a rhodopsin gene defect (pro23-to-his). Arch. Ophthal. 109:

[35783] 92-101, 1991.

[35784] 5260.

[35785] Berson, E. L.; Rosner, B.; Sandberg, M. A.; Hayes, K. C.; Nicholson,

[35786] B. W.; Weigel-DiFranco, C.; Willett, W.: A randomized trial of vitamin

[35787] A and vitamin E supplementation for retinitis pigmentosa. Arch. Ophthal. 111:

[35788] 761-772, 1993.

[35789] 5261.

[35790] Borhan, B.; Souto, M. L.; Imai, H.; Shichida, Y.; Nakanishi, K.

[35791] : Movement of retinal along the visual transduction path. Science 288:

[35792] 2209-2212, 2000.

[35793] 5262.

[35794] Bourne, H. R.; Meng, E. C.: Rhodopsin sees the light. Science 289:

[35795] 733-734, 2000.

[35796] 5263.

[35797] Bradley, D. G.; Farrar, G. J.; Sharp, E.; Kenna, P.; Humphries,

[35798] M. M.; McWilliam, P.; Humphries, P.: Exclusion mapping of the autosomal

[35799] dominant retinitis pigmentosa (RP1) locus from the short arm of chromosome

[35800] 1. (Abstract) Cytogenet. Cell Genet. 51: 968, 1989.

[35801] 5264.

[35802] Bouchard, B.; Del Marmol, V.; Jackson, I. J.; Cherif, D.; Dubertret,
[35803] L.: Molecular characterization of a human tyrosinase-related-protein-2
[35804] cDNA: patterns of expression in melanocytic cells. Europ. J. Biochem. 219:
[35805] 127-134, 1994.

[35806] 5265.

[35807] Budd, P. S.; Jackson, I. J.: Structure of the mouse tyrosinase-related
[35808] protein-2/dopachrome tautomerase (Tyrp2/Dct) gene and sequence of
[35809] two novel slaty alleles. Genomics 29: 35-43, 1995.

[35810] 5266.

[35811] Cassady, J. L.; Sturm, R. A.: Sequence of the human dopachrome
[35812] tautomerase-encoding TRP-2 cDNA. Gene 143: 295-298, 1994.
[35813] 5267.

[35814] Jackson, I. J.; Chambers, D. M.; Tsukamoto, K.; Copeland, N. G.;

[35815] Gilbert, D. J.; Jenkins, N. A.; Hearing, V.: A second tyrosinase-related

[35816] protein, TRP-2, maps to and is mutated at the mouse slaty locus. EMBO

[35817] J. 11: 527-535, 1992.

[35818] 5268.

[35819] Khong, H. T.; Rosenberg, S. A.: Pre-existing immunity to tyrosinase-related

[35820] protein (TRP)-2, a new TRP-2 isoform, and the NY-ESO-1 melanoma antigen

[35821] in a patient with a dramatic response to immunotherapy. J. Immun. 168:

[35822] 951-956, 2002.

[35823] 5269.

[35824] Kwon, B. S.: Pigmentation genes: the tyrosinase gene family and

[35825] the pmel 17 gene family. J. Invest. Derm. 100: 134S-140S, 1993.

[35826] 5270.

[35827] Sturm, R. A.; Baker, E.; Sutherland, G. R.: Assignment of the

[35828] tyrosinase-related protein-2 gene (TYRP2) to human chromosome 13q31-q32

[35829] by fluorescence in situ hybridization: extended synteny with mouse

[35830] chromosome 14. Genomics 21: 293-296, 1994.

[35831] 5271.

[35832] Sturm, R. A.; O'Sullivan B. J.; Box, N. F.; Smith, A. G.; Smit,

[35833] S. E.; Puttick, E. R. J.; Parsons, P. G.; Dunn, I. S.: Chromo-somal

[35834] structure of the human TYRP1 and TYRP2 loci and com-parison of the

[35835] tyrosinase-related protein gene family. Genomics 29: 24-34, 1995.

[35836] 5272.

[35837] Somlo, S.; Wirth, B.; Germino, G. G.; Weinstat-Saslow, D.; Gillespie,

[35838] G. A. J.; Himmelbauer, H.; Steevens, L.; Coucke, P.; Willems, P.;

[35839] Bachner, L.; Coto, E.; Lopez-Larrea, C.; Peral, B.; San Mil-lan, J.

[35840] L.; Saris, J. J.; Breuning, M. H.; Frischauf, A.-M.; Reeders, S. T.

[35841] : Fine genetic localization of the gene for autosomal dominant polycystic

[35842] kidney disease (PKD1) with respect to physically mapped markers. Genomics 13:

[35843] 152-158, 1992.

[35844] 5273.

[35845] Turco, A. E.; Rossetti, S.; Bresin, E.; Corra, S.; Gammaro, L.;

[35846] Maschio, G.; Pignatti, P. F.: A novel nonsense mutation in the PKD1

[35847] gene (C3817T) is associated with autosomal dominant polycystic kidney

[35848] disease (ADPKD) in a large three-generation Italian family. Hum.

[35849] Molec. Genet. 4: 1331-1335, 1995.

[35850] 5274.

[35851] Ward, C. J.; Turley, H.; Ong, A. C. M.; Comley, M.; Biddolph,

[35852] S.; Chetty, R.; Ratcliffe, P. J.; Gatter, K.; Harris, P. C.: Polycystin,

[35853] the polycystic kidney disease 1 protein, is expressed by epithelial

[35854] cells in fetal, adult, and polycystic kidney. Proc. Nat. Acad. Sci. 93:
[35855] 1524-1528, 1996.

[35856] 5275.

[35857] Watson, M. L.; Wright, A. F.; Macnicol, A. M.; Allan, P. L.; Clayton,
[35858] J. F.; Dempster, M.; Jeremiah, S. J.; Corney, G.; Hopkinson, D. A.
[35859] : Studies of genetic linkage between adult polycystic kidney disease
[35860] and three markers on chromosome 16. J. Med. Genet. 24: 457-461,
[35861] 1987.

[35862] 5276.

[35863] Bachner, L.; Vinet, M. C.; Lacave, R.; Babron, M. C.; Rondeau,
[35864] E.; Sraer, J. D.; Chevet, D.; Kaplan, J.-C.: Linkage study of a large
[35865] family with autosomal dominant polycystic kidney disease with reduced
[35866] expression: absence of linkage to the PKD1 locus. Hum. Genet. 85:

[35867] 221-227, 1990.

[35868] 5277.

[35869] Bhunia, A. K.; Piontek, K.; Boletta, A.; Liu, L.; Qian, F.; Xu,

[35870] P.-N.; Germino, F. J.; Germino, G. G.: PKD1 induces
p21-waf1 and

[35871] regulation of the cell cycle via direct activation of the JAK-
STAT

[35872] signaling pathway in a process requiring PKD2. Cell 109:
157-168,

[35873] 2002.

[35874] 5278.

[35875] Greenberg, A. S.; Egan, J. J.; Wek, S. A.; Moos, M. C., Jr.;
Londos,

[35876] C.; Kimmel, A. R.: Isolation of cDNAs for perilipins A and
B: sequence

[35877] and expression of lipid droplet-associated proteins of
adipocytes. Proc.

[35878] Nat. Acad. Sci. 90: 12035-12039, 1993.

[35879] 5279.

[35880] Greenberg, A. S.; Egan, J. J.; Wek, S. A.; Takeda, T.; Lon-
dos,

[35881] C.; Kimmel, A. K.: Perilipin, a lipid droplet-associated, adipocyte
[35882] specific protein; cDNA cloning and expression. (Abstract) Clin. Res. 39:
[35883] 287A only, 1991.

[35884] 5280.

[35885] Lu, X.; Gruia-Gray, J.; Copeland, N. G.; Gilbert, D. J.; Jenk-ins,
[35886] N. A.; Londos, C.; Kimmel, A. R.: The murine perilipin gene: the
[35887] lipid droplet-associated perilipins derive from tissue-specific, mRNA
[35888] splice variants and define a gene family of ancient origin. Mammalian
[35889] Genome 12: 741-749, 2001.

[35890] 5281.

[35891] Martinez-Botas, J.; Anderson, J. B.; Tessier, D.; Lapillonne, A.;
[35892] Chang, B. H.-J.; Quast, M. J.; Gorenstein, D.; Chen, K.-H.; Chan,
[35893] L.: Absence of perilipin results in leanness and reverses obesity

[35894] in Lepr(db/db) mice. Nature Genet. 26: 474-479, 2000.

[35895] 5282.

[35896] Nishiu, J.; Tanaka, T.; Nakamura, Y.: Isolation and chro-
mosomal

[35897] mapping of the human homolog of perilipin (PLIN), a rat
adipose tissue-specific

[35898] gene, by differential display method. Genomics 48:
254-257, 1998.

[35899] 5283.

[35900] Tansey, J. T.; Sztalryd, C.; Gruia-Gray, J.; Roush, D. L.; Zee,

[35901] J. V.; Gavrilova, O.; Reitman, M. L.; Deng, C.-X.; Li, C.;
Kimmel,

[35902] A. R.; Londos, C.: Perilipin ablation results in a lean mouse
with

[35903] aberrant adipocyte lipolysis, enhanced leptin production,
and resistance

[35904] to diet-induced obesity. Proc. Nat. Acad. Sci. 98:
6494-6499, 2001.

[35905] 5284.

[35906] King, C. R.; Schimke, R. N.; Arthur, T.; Davoren, B.; Collins,

[35907] D.: Proximal 3p deletion in renal cell carcinoma cells from

a patient
[35908] with von Hippel-Lindau disease. Cancer Genet. Cytogenet. 27: 345-348,
[35909] 1987.

[35910] 5285.

[35911] Li, F. P.; Decker, H.-J. H.; Zbar, B.; Stanton, V. P., Jr.; Ko-
vacs,
[35912] G.; Seizinger, B. R.; Aburatani, H.; Sandberg, A. A.; Berg, S.; Hosoe,
[35913] S.; Brown, R. S.: Clinical and genetic studies of renal cell carcinomas
[35914] in a family with a constitutional chromosome 3;8 translo-
cation: genetics
[35915] of familial renal carcinoma. Ann. Intern. Med. 118: 106-111, 1993.
[35916] 5286.

[35917] Ohta, M.; Inoue, H.; Cotticelli, M. G.; Kastury, K.; Baffa, R.;

[35918] Palazzo, J.; Siprashvili, Z.; Mori, M.; McCue, P.; Druck, T.; Croce,
[35919] C. M.; Huebner, K.: The FHIT gene, spanning the chromo-
some 3p14.2
[35920] fragile site and renal carcinoma-associated t(3;8) break-

point, is
[35921] abnormal in digestive tract cancers. Cell 84: 587-597, 1996.
[35922] 5287.
[35923] Komure, O.; Sano, A.; Nishino, N.; Yamauchi, N.; Ueno, S.; Kondoh,
[35924] K.; Sano, N.; Takahashi, M.; Murayama, N.; Kondo, I.; Nagafuchi, S.;
[35925] Yamada, M.; Kanazawa, I.: DNA analysis in hereditary dentatorubral-pallidoluysian
[35926] atrophy: correlation between CAG repeat length and phenotypic variation
[35927] and the molecular basis of anticipation. Neurology 45: 143-149,
[35928] 1995.
[35929] 5288.
[35930] Nagafuchi, S.; Yanagisawa, H.; Sato, K.; Shirayama, T.; Ohsaki,
[35931] E.; Bundo, M.; Takeda, T.; Tadokoro, K.; Kondo, I.; Murayama, N.;
[35932] Tanaka, Y.; Kikushima, H.; Umino, K.; Kurosawa, H.; Furukawa, T.;

[35933] Nihei, K.; Inoue, T.; Sano, A.; Komure, O.; Takahashi, M.; Yoshizawa,

[35934] T.; Kanazawa, I.; Yamada, M.: Dentatorubral and pallidoluysian atrophy

[35935] expansion of an unstable CAG trinucleotide on chromosome 12p. Nature

[35936] Genet. 6: 14-18, 1994.

[35937] 5289.

[35938] Norremolle, A.; Nielsen, J. E.; Sorensen, S. A.; Hasholt, L.:

[35939] Elongated CAG repeats of the B37 gene in a Danish family with dentato-rubro-pallido-luysian

[35940] atrophy. Hum. Genet. 95: 313-318, 1995.

[35941] 5290.

[35942] Oyake, M.; Onodera, O.; Shiroishi, T.; Takano, H.; Takahashi,

[35943] Y.; Kominami, R.; Moriwaki, K.; Ikeuchi, T.; Igarashi, S.; Tanaka,

[35944] H.; Tsuji, S.: Molecular cloning of murine homologue dentatorubral-pallidoluysian

[35945] atrophy (DRPLA) cDNA: strong conservation of a polymorphic CAG repeat

[35946] in the murine gene. Genomics 40: 205-207, 1997.

[35947] 5291.

[35948] Potter, N. T.: The relationship between (CAG)n repeat number

[35949] and age of onset in a family with dentatorubral-palli-doluysian atrophy

[35950] (DRPLA): diagnostic implications of confirmatory and pre-dictive testing. J.

[35951] Med. Genet. 33: 168-170, 1996.

[35952] 5292.

[35953] Roses, A. D.: From genes to mechanisms to therapies: lessons

[35954] to be learned from neurological disorders. Nature Med. 2: 267-269,

[35955] 1996.

[35956] 5293.

[35957] Saitoh, S.; Momoi, M. Y.; Yamagata, T.; Miyao, M.; Suwa, K.:

[35958] Clinical and electroencephalographic findings in juvenile type DRPLA. Pediat.

[35959] Neurol. 18: 265-268, 1998.

[35960] 5294.

[35961] Sano, A.; Yamauchi, N.; Kakimoto, Y.; Komure, O.; Kawai, J.; Hazama,

[35962] F.; Kuzume, K.; Sano, N.; Kondo, I.: Anticipation in hereditary dentatorubral-pallidoluysian

[35963] atrophy. Hum. Genet. 93: 699-702, 1994.

[35964] 5295.

[35965] Sato, K.; Kashihara, K.; Okada, S.; Ikeuchi, T.; Tsuji, S.; Shomori,

[35966] T.; Morimoto, K.; Hayabara, T.: Does homozygosity advance the onset

[35967] of dentatorubral-pallidoluysian atrophy? Neurology 45: 1934-1936,

[35968] 1995.

[35969] 5296.

[35970] Sato, T.; Oyake, M.; Nakamura, K.; Nakao, K.; Fukusima, Y.; Onodera,

[35971] O.; Igarashi, S.; Takano, H.; Kikugawa, K.; Ishida, Y.; Shimohata,

[35972] T.; Koide, R.; and 15 others: Transgenic mice harboring a full-length

[35973] human mutant DRPLA gene exhibit age-dependent inter-
generational and
[35974] somatic instabilities of CAG repeats comparable with
those in DRPLA
[35975] patients. Hum. Molec. Genet. 8: 99-106, 1999.

[35976] 5297.

[35977] Schmitt, I.; Epplen, J. T.; Riess, O.: Predominant neuronal
expression
[35978] of the gene responsible for dentatorubral-pallidoluysian
atrophy (DRPLA)
[35979] in rat. Hum. Molec. Genet. 4: 1619-1624, 1995.

[35980] 5298.

[35981] Shimojo, Y.; Osawa, Y.; Fukumizu, M.; Hanaoka, S.;
Tanaka, H.;
[35982] Ogata, F.; Sasaki, M; Sugai, K.: Severe infantile denta-
torubral pallidoluysian
[35983] atrophy with extreme expansion of CAG repeats. Neurol-
ogy 56: 277-278,
[35984] 2001.

[35985] 5299.

[35986] Sisodia, S. S.: Nuclear inclusions in glutamine repeat dis-

orders:

[35987] are they pernicious, coincidental, or beneficial? Cell 95: 1-4,

[35988] 1998.

[35989] 5300.

[35990] Smith, J. K.: Dentatorubropallidoluysian atrophy.In: Vinken,

[35991] P. J.; Bruyn, G. W. (eds): Handbook of Clinical Neurology. Part I.

[35992] System Disorders and Atrophies. New York: Elsevier Publishing Co.

[35993] Vol. 21: 1975. Pp. 519-534.

[35994] 5301.

[35995] Smith, J. K.; Conda, V. E.; Malamud, N.: Unusual form of cerebellar

[35996] ataxia: combined dentato-rubral and pallido-Luysian degeneration. Neurology 8:

[35997] 205-209, 1958.

[35998] 5302.

[35999] Takahashi, H.; Ohama, E.; Naito, H.; Takeda, S.; Nakashima, S.;

[36000] Makifuchi, T.; Ikuta, F.: Hereditary dentato-
rubral-pallidoluysian
[36001] atrophy: clinical and pathologic variants in a family. Neu-
rology 38:
[36002] 1065-1070, 1988.

[36003] 5303.

[36004] Takano, T.; Yamanouchi, Y.; Nagafuchi, S.; Yamada, M.:
Assignment
[36005] of the dentatorubral and pallidoluysian atrophy (DRPLA)
gene to 12p13.31
[36006] by fluorescence in situ hybridization. Genomics 32:
171-172, 1996.
[36007] 5304.

[36008] Takiyama, Y.; Sakoe, K.; Amaike, M.; Soutome, M.; Ogawa,
T.; Nakano,
[36009] I.; Nishizawa, M.: Single sperm analysis of the CAG repeats
in the
[36010] gene for dentatorubral-pallidoluysian atrophy (DRPLA):
the instability
[36011] of the CAG repeats in the DRPLA gene is prominent
among the CAG repeat
[36012] diseases. Hum. Molec. Genet. 8: 453-457, 1999.

[36013] 5305.

[36014] Tomoda, A.; Ikezawa, M.; Ohtani, Y.; Miike, T.; Kumamoto, T.:

[36015] Progressive myoclonus epilepsy: dentato-rubro-pallido-luysian atrophy

[36016] (DRPLA) in childhood. Brain Dev. 13: 266-269, 1991.

[36017] 5306.

[36018] Tsuji, S.: Personal Communication. Niigata, Japan 11/15/1993.

[36019] 5307.

[36020] Ueno, S.; Kondoh, K.; Kotani, Y.; Komure, O.; Kuno, S.; Kawai,

[36021] J.; Hazama, F.; Sano, A.: Somatic mosaicism of CAG repeat in dentatorubral-pallidoluysian

[36022] atrophy (DRPLA). Hum. Molec. Genet. 4: 663-666, 1995.

[36023] 5308.

[36024] Warner, T. T.; Williams, L.; Harding, A. E.: DRPLA in Eu-rope.

[36025] (Letter) Nature Genet. 6: 225, 1994.

[36026] 5309.

[36027] Watanabe, H.; Tanaka, F.; Matsumoto, M.; Doyu, M.; Ando, T.; Mitsuma,
[36028] T.; Sobue, G.: Frequency analysis of autosomal dominant cerebellar
[36029] ataxias in Japanese patients and clinical characterization of spinocerebellar
[36030] ataxia type 6. Clin. Genet. 53: 13-19, 1998.

[36031] 5310.

[36032] Wood, J. D.; Yuan, J.; Margolis, R. L.; Colomer, V.; Duan, K.;
[36033] Kushi, J.; Kaminsky, Z.; Kleiderlein, J. J., Jr.; Sharp, A. H.; Ross,
[36034] C. A.: Atrophin-1, the DRPLA gene product, interacts with two families
[36035] of WW domain-containing proteins. Molec. Cell. Neurosci. 11: 149-160,
[36036] 1998.

[36037] 5311.

[36038] Aksoy, M.; Erdem, S.: Combination of hereditary elliptocy-tosis
[36039] and heterozygous beta-thalassemia: a family study. J. Med. Genet. 5:

[36040] 298-301, 1968.

[36041] 5312.

[36042] Alloisio, N.; Dorleac, E.; Delaunay, J.; Girot, R.; Galand, C.;

[36043] Boivin, P.: A shortened variant of red cell membrane protein 4.1. Blood 60:

[36044] 265-267, 1982.

[36045] 5313.

[36046] Alloisio, N.; Dorleac, E.; Girot, R.; Delaunay, J.: Analysis of

[36047] red cell membrane in a family with hereditary elliptocytosis: total

[36048] or partial absence of protein 4.1. Hum. Genet. 59: 68-71, 1981.

[36049] 5314.

[36050] Alloisio, N.; Morle, L.; Dorleac, E.; Gentilhomme, O.; Bachir,

[36051] D.; Guetarni, D.; Colonna, P.; Bost, M.; Zouaoui, Z.; Roda, L.; Roussel,

[36052] D.; Delaunay, J.: The heterozygous form of 4.1(-) hereditary elliptocytosis

[36053] [the 4.1(-) trait]. Blood 65: 46-51, 1985.

[36054] 5315.

[36055] Bahary, N.; Zorich, G.; Pachter, J. E.; Leibel, R. L.; Friedman,

[36056] J. M.: Molecular genetic linkage maps of mouse chromosomes 4 and 5316.

[36057] Genomics 11: 33-47, 1991.

[36058]

[36059] 6. Baklouti, F.; Huang, S.-C.; Vulliamy, T. J.; Delaunay, J.; Benz,

[36060] E. J., Jr.: Organization of the human protein 4.1 genomic locus:

[36061] new insights into the tissue-specific alternative splicing of the

[36062] pre-mRNA. Genomics 39: 289-302, 1997.

[36063] 5317.

[36064] Bannerman, R. M.; Renwick, J. H.: The hereditary elliptocytoses:

[36065] clinical and linkage data. Ann. Hum. Genet. 26: 23-38, 1962.

[36066] 5318.

[36067] Clarke, C. A.; Donohoe, W. T. A.; Finn, R.; McConnell, R. B.;

Sheppard,
[36068] P. M.; Nicol, D. S. H.: Data on linkage in man: ovalocytosis, sickling
[36069] and the Rhesus blood group complex. Ann. Hum. Genet. 24: 283-287,
[36070] 1960.

[36071] 5319.

[36072] Conboy, J.; Kan, Y. W.; Shohet, S. B.; Mohandas, N.: Molec-ular
[36073] cloning of protein 4.1, a major structural element of the human erythrocyte
[36074] membrane skeleton. Proc. Nat. Acad. Sci. 83: 9512-9516, 1986.
[36075] 5320.

[36076] Conboy, J.; Marchesi, S.; Kim, R.; Agre, P.; Kan, Y. W.; Mo-handas,
[36077] N.: Molecular analysis of insertion/deletion mutations in protein
[36078] 4.1 in elliptocytosis. II. Determination of molecular genetic origins
[36079] of rearrangements. J. Clin. Invest. 86: 524-530, 1990.

[36080] 5321.

[36081] Conboy, J.; Mohandas, N.; Tchernia, G.; Kan, Y. W.: Molec-
ular
[36082] basis of hereditary elliptocytosis due to protein 4.1 defi-
ciency. New
[36083] Eng. J. Med. 315: 680-685, 1986.

[36084] 5322.

[36085] Conboy, J. G.: Structure, function, and molecular genetics
of
[36086] erythroid membrane skeletal protein 4.1 in normal and
abnormal red
[36087] blood cells. Seminars Hemat. 30: 58-73, 1993.

[36088] 5323.

[36089] Conboy, J. G.; Chan, J.; Mohandas, N.; Kan, Y. W.: Multiple
protein
[36090] 4.1 isoforms produced by alternative splicing in human
erythroid cells. Proc.
[36091] Nat. Acad. Sci. 85: 9062-9065, 1988.

[36092] 5324.

[36093] Conboy, J. G.; Chasis, J. A.; Winardi, R.; Tchernia, G.; Kan,

[36094] Y. W.; Mohandas, N.: An isoform-specific mutation in the
protein

[36095] 4.1 gene results in hereditary elliptocytosis and complete deficiency
[36096] of protein 4.1 in erythrocytes but not in nonerythroid cells. J.
[36097] Clin. Invest. 91: 77-82, 1993.

[36098] 5325.

[36099] Conboy, J. G.; Mohandas, N.; Wang, C.; Tchernia, G.; Shohet, S.
[36100] B.; Kan, Y. W.: Molecular cloning and characterization of the gene
[36101] coding for red cell membrane skeletal protein 4.1. (Abstract) Blood 66
[36102] (suppl. 1): 31A, 1985.

[36103] 5326.

[36104] Cook, P. J. L.; Noades, J. E.; Newton, M. S.; de Mey, R.: On
[36105] the orientation of the Rh:E1-1 linkage group. Ann. Hum. Genet. 41:
[36106] 157-162, 1977.

[36107] 5327.

[36108] Puffenberger, E. G.; Hosoda, K.; Washington, S. S.; Nakao, K.;

[36109] deWit, D.; Yanagisawa, M.; Chakravarti, A.: A missense mutation of
[36110] the endothelin-B receptor gene in multigenic Hirschsprung's disease. Cell 79:
[36111] 1257-1266, 1994.

[36112] 5328.

[36113] Svensson, P.-J.; Anvret, M.; Molander, M.-L.; Norden-
skjold, A.
[36114] : Phenotypic variation in a family with mutations in two Hirschsprung-related
[36115] genes (RET and endothelin receptor B). Hum. Genet. 103: 145-148,
[36116] 1998.

[36117] 5329.

[36118] Hanks, M.; Wurst, W.; Anson-Cartwright, L.; Auerbach, A. B.; Joyner,
[36119] A. L.: Rescue of the En-1 mutant phenotype by replace-
ment of En-1
[36120] with En-2. Science 269: 679-682, 1995.

[36121] 5330.

[36122] Johnson, R. L.; Tabin, C. J.: Molecular models for verte-

brate
[36123] limb development. Cell 90: 979-990, 1997.

[36124] 5331.

[36125] Kohler, A.; Logan, C.; Joyner, A. L.; Muenke, M.: Regional assignment
[36126] of the human homeobox-containing gene EN1 to chromosome 2q13-q21. Genomics 15:
[36127] 233-235, 1993.

[36128] 5332.

[36129] Logan, C.; Hanks, M. C.; Noble-Topham, S.; Nallainathan, D.; Provart,
[36130] N. J.; Joyner, A. L.: Cloning and sequence comparison of the mouse,
[36131] human, and chicken engrailed genes reveal potential functional domains
[36132] and regulatory regions. Dev. Genet. 13: 345-358, 1992.

[36133] 5333.

[36134] Logan, C.; Willard, H. F.; Rommens, J. M.; Joyner, A. L.: Chromosomal
[36135] localization of the human homeo box-containing genes, EN1 and EN2. Genomics 4:

[36136] 206-209, 1989.

[36137] 5334.

[36138] Loomis, C. A.; Harris, E.; Michaud, J.; Wurst, W.; Hanks, M.; Joyner,

[36139] A. L.: The mouse engrailed-1 gene and ventral limb patterning. Nature 382:

[36140] 360-363, 1996.

[36141] 5335.

[36142] Martin, G. R.; Richman, M.; Reinsch, S.; Nadeau, J. H.; Joyner,

[36143] A.: Mapping of the two mouse engrailed-like genes: close linkage

[36144] of En-1 to dominant hemimelia (Dh) on chromosome 1 and of En-2 to

[36145] hemimelic extra-toes (Hx) on chromosome 5. Genomics 6: 302-308,

[36146] 1990.

[36147] 5336.

[36148] Matsui, T.; Hirai, M.; Hirano, M.; Kurosawa, Y.: The HOX complex

[36149] neighbored by the EVX gene, as well as two other home-

obox-containing

[36150] genes, the GBX-class and the EN class, are located on the same chromosomes

[36151] 2 and 7 in humans. FEBS Lett. 336: 107-110, 1993.

[36152] 5337.

[36153] Wurst, W.; Auerbach, A. B.; Joyner, A. L.: Multiple developmental

[36154] defects in Engrailed-1 mutant mice: an early mid-hindbrain deletion

[36155] and patterning defects in forelimbs and sternum. Development 120:

[36156] 2065-2075, 1994.

[36157] 5338.

[36158] Kohl, S.; Baumann, B.; Rosenberg, T.; Kellner, U.; Lorenz, B.;

[36159] Vadala, M.; Jacobson, S. G.; Wissinger, B.: Mutations in the cone

[36160] photoreceptor G-protein alpha-subunit gene GNAT2 in patients with

[36161] achromatopsia. Am. J. Hum. Genet. 71: 422-425, 2002.

[36162] 5339.

[36163] Morris, T. A.; Fong, S.-L.: Characterization of the gene encoding
[36164] human cone transducin alpha-subunit (GNAT2). Genomics 17: 442-448,
[36165] 1993.

[36166] 5340.

[36167] Sparkes, R. S.; Cohn, V. H.; Cire-Eversole, P.; Blatt, C.; Amatruda,
[36168] T. T.; Weiner, L. P.; Nesbitt, M.; Reed, R. R.; Lochrie, M. A.; Fournier,
[36169] R. E. K.; Simon, M. I.: Mapping of genes encoding the subunits of
[36170] guanine nucleotide-binding proteins (G-proteins) in the mouse. (Abstract) Cytogenet.
[36171] Cell Genet. 46: 696 only, 1987.

[36172] 5341.

[36173] Compton, J. G.: Epidermal disease: faulty keratin filaments take
[36174] their toll. Nature Genet. 6: 6-7, 1994.

[36175] 5342.

[36176] Kimonis, V.; DiGiovanna, J. J.; Yang, J.-M.; Doyle, S. Z.;

Bale,

[36177] S. J.; Compton, J. G.: A mutation in the V1 end domain of keratin

[36178] 1 in non-epidermolytic palmar-plantar keratoderma. J. Invest. Derm. 103:

[36179] 764-769, 1994.

[36180] 5343.

[36181] Le Marchand-Brustel, Y.; Gremeaux, T.; Ballotti, R.; van Obberghen,

[36182] E.: Insulin receptor tyrosine kinase is defective in skeletal muscle

[36183] of insulin-resistant obese mice. Nature 315: 676-679, 1985.

[36184] 5344.

[36185] Leme, C. E.; Wajchenberg, B. L.; Lerario, A. C.; Goldman, J.;

[36186] Borges, J. L. C.: Acanthosis nigricans, hirsutism, insulin resistance

[36187] and insulin receptor defect. Clin. Endocr. 17: 43-49, 1982.

[36188] 5345.

[36189] Longo, N.; Langley, S. D.; Griffin, L. D.; Elsas, L. J.: Activa-

tion

[36190] of glucose transport by a natural mutation in the human insulin receptor. Proc.

[36191] Nat. Acad. Sci. 90: 60-64, 1993.

[36192] 5346.

[36193] Longo, N.; Langley, S. D.; Griffin, L. D.; Elsas, L. J., II:

[36194] Reduced mRNA and a nonsense mutation in the insulin-receptor gene

[36195] produce heritable severe insulin resistance. Am. J. Hum. Genet. 50:

[36196] 998-1007, 1992.

[36197] 5347.

[36198] Maassen, J. A.; Klinkhamer, M. P.; Odink, R. J. H.; Sips, H.;

[36199] van der Zon, G. C. M.; Wieringa, T.; Krans, H. M. J.; Moller, W.:

[36200] Improper expression of insulin receptors on fibroblasts from a leprechaun

[36201] patient. Europ. J. Biochem. 172: 725-729, 1988.

[36202] 5348.

[36203] Mariani, S.; Pedone, A.; Meschi, F.; Di Natale, B.; Caputo,

R.;

[36204] Broggi, U.; Chiumello, G.: Insulin resistance in a child with acanthosis

[36205] nigricans type A. Acta Paediat. Scand. 71: 667-670, 1982.

[36206] 5349.

[36207] Michael, M. D.; Kulkarni, R. N.; Postic, C.; Previs, S. F.; Shulman,

[36208] G. I.; Magnuson, M. A.; Kahn, C. R.: Loss of insulin signal-ing in

[36209] hepatocytes leads to severe insulin resistance and pro-gressive hepatic

[36210] dysfunction. Molec. Cell 6: 87-97, 2000.

[36211] 5350.

[36212] Moller, D. E.; Cohen, O.; Yamaguchi, Y.; Assiz, R.; Grig-orescu,

[36213] F.; Eberle, A.; Morrow, L. A.; Moses, A. C.; Flier, J. S.: Prevalence

[36214] of mutations in the insulin receptor gene in subjects with features

[36215] of the type A syndrome of insulin resistance. Diabetes 43: 247-255,

[36216] 1994.

[36217] 5351.

[36218] Moller, D. E.; Flier, J. S.: Insulin resistance--mechanisms,

[36219] syndromes, and implications. New Eng. J. Med. 325:
938-948, 1991.

[36220] 5352.

[36221] Moller, D. E.; Flier, J. S.: Detection of an alteration in the

[36222] insulin-receptor gene in a patient with insulin resistance,
acanthosis

[36223] nigricans, and the polycystic ovary syndrome (type A in-
sulin resistance). New

[36224] Eng. J. Med. 319: 1526-1529, 1988.

[36225] 5353.

[36226] Moller, D. E.; Yokota, A.; Ginsberg-Fellner, F.; Flier, J. S.

[36227] : Functional properties of a naturally occurring
trp(1200)-to-ser(1200)

[36228] mutation of the insulin receptor. Molec. Endocr. 4:
1183-1191, 1990.

[36229] 5354.

[36230] Moller, D. E.; Yokota, A.; Pazianos, A.; Flier, J. S.: A mis-
sense

[36231] mutation in one allele of the tyrosine kinase domain of the insulin

[36232] receptor gene is associated with dominantly inherited insulin resistance.

[36233] (Abstract) Clin. Res. 38: 435A only, 1990.

[36234] 5355.

[36235] Moller, D. E.; Yokota, A.; White, M. F.; Pazianos, A. G.; Flier,

[36236] J. S.: A naturally occurring mutation of insulin receptor alanine

[36237] 1134 impairs tyrosine kinase function and is associated with dominantly

[36238] inherited insulin resistance. J. Biol. Chem. 265: 14979-14985, 1990.

[36239] 5356.

[36240] Moncada, V. Y.; Hedo, J. A.; Serrano-Rios, M.; Taylor, S. I.:

[36241] Insulin-receptor biosynthesis in cultured lymphocytes from an insulin-resistant

[36242] patient (Rabson-Mendenhall syndrome): evidence for defect before insertion

[36243] of receptor into plasma membrane. Diabetes 35: 802-807, 1986.

[36244] 5357.

[36245] Norton, K. I.; Glicklich, M.; Kupchik, G.; Gray, C. E.; Lud-
man,
[36246] M.: Leprechaunism: a case report with radiographic fea-
tures. Dysmorph.
[36247] Clin. Genet. 4: 57-62, 1990.

[36248] 5358.

[36249] Odawara, M.; Kadowaki, T.; Yamamoto, R.; Shibasaki, Y.;
Tobe,
[36250] K.; Accili, D.; Bevins, C.; Mikami, Y.; Matsuura, N.;
Akanuma, Y.;
[36251] Takaku, F.; Taylor, S. I.; Kasuga, M.: Human diabetes as-
sociated
[36252] with a mutation in the tyrosine kinase domain of the in-
sulin receptor. Science 245:
[36253] 66-68, 1989.

[36254] 5359.

[36255] Ojamaa, K.; Hedo, J. A.; Roberts, C. T., Jr.; Moncada, V. Y.;

[36256] Gorden, P.; Ullrich, A.; Taylor, S. I.: Defects in human in-
sulin
[36257] receptor gene expression. Molec. Endocr. 2: 242-247,

1988.

[36258] 5360.

[36259] Prince, M. J.; Smith, F. E.; Peters, E. J.; Stuart, C. A.: Functional

[36260] characteristics of decreased insulin receptors on fibroblasts obtained

[36261] from a subject with severe insulin resistance and acanthosis nigricans. Diabetes 35:

[36262] 148-154, 1986.

[36263] 5361.

[36264] Quin, J. D.; Fisher, B. M.; Paterson, K. R.; Inoue, A.; Beastall,

[36265] G. H.; MacCuish, A. C.: Acute response to recombinant insulin-like

[36266] growth factor-I in a patient with Mendenhall's syndrome. New Eng.

[36267] J. Med. 323: 1425-1426, 1991.

[36268] 5362.

[36269] Quon, M. J.; Guerre-Millo, M.; Zarnowski, M. J.; Butte, A. J.;

[36270] Em, M.; Cushman, S. W.; Taylor, S. I.: Tyrosine kinase-deficient

[36271] mutant human insulin receptors (met1153-to-ile) overex-
pressed in transfected

[36272] rat adipose cells fail to mediate translocation of epitope-
tagged

[36273] GLUT4. Proc. Nat. Acad. Sci. 91: 5587-5591, 1994.

[36274] 5363.

[36275] Rabson, S. M.; Mendenhall, E. N.: Familial hypertrophy of
pineal

[36276] body, hyperplasia of adrenal cortex and diabetes mellitus.
Am. J.

[36277] Clin. Path. 26: 283-290, 1956.

[36278] 5364.

[36279] Rajala, R. V. S.; Anderson, R. E.: Interaction of the insulin

[36280] receptor beta-subunit with phosphatidylinositol 3-kinase
in bovine

[36281] ROS. Invest. Ophthal. Vis. Sci. 42: 3110-3117, 2001.

[36282] 5365.

[36283] Roth, R. A.; Cassell, D. J.: Insulin receptor: evidence that

[36284] it is a protein kinase. Science 219: 299-301, 1983.

[36285] 5366.

[36286] Rubin, C. S.: Personal Communication. Bronx, N. Y. 12/8/1984.

[36287] 5367.

[36288] Rudiger, H. W.; Ahrens, P.; Dreyer, M.; Frorath, B.; Loffel, C.;

[36289] Schmidt-Preuss, U.: Impaired insulin-induced RNA synthesis secondary

[36290] to a genetically defective insulin receptor. Hum. Genet. 69: 76-78,

[36291] 1985.

[36292] 5368.

[36293] Rudiger, H. W.; Dreyer, M.; Kuhnau, J.; Bartelheimer, H.: Familial

[36294] insulin-resistant diabetes secondary to an affinity defect of the

[36295] insulin receptor. Hum. Genet. 64: 407-411, 1983.

[36296] 5369.

[36297] Salmeen, A.; Andersen, J. N.; Myers, M. P.; Tonks, N. K.; Barford,

[36298] D.: Molecular basis for the dephosphorylation of the activation segment

[36299] of the insulin receptor by protein tyrosine phosphatase 1B. Molec.

[36300] Cell 6: 1401-1412, 2000.

[36301] 5370.

[36302] Scarlett, J. A.; Kolterman, O. G.; Moore, P.; Saekow, M.; In-

[36303] sel, J.; Griffin, J.; Mako, M.; Rubenstein, A. H.; Olefsky, J. M.: Insulin

[36304] resistance and diabetes due to a genetic defect in insulin receptors. J.

[36305] Clin. Endocr. Metab. 55: 123-132, 1982.

[36306] 5371.

[36307] Seino, S.; Seino, M.; Nishi, S.; Bell, G. I.: Structure of the

[36308] human insulin receptor gene and characterization of its promoter. Proc.

[36309] Nat. Acad. Sci. 86: 114-118, 1989.

[36310] 5372.

[36311] Botto, M.; Fong, K. Y.; So, A. K.; Koch, C.; Walport, M. J.:

[36312] Molecular basis of polymorphisms of human complement component C3. J.

[36313] Exp. Med. 172: 1011-1017, 1990.

[36314] 5373.

[36315] Polymeropoulos, M. H.; Torres, R.; Yanovski, J. A.; Chan-
drasekharappa,
[36316] S. C.; Ledbetter, D. H.: The human corticotropin-releasing
factor
[36317] receptor (CRHR) gene maps to chromosome 17q12-q22.
Genomics 28:
[36318] 123-124, 1995.

[36319] 5374.

[36320] Greenspan, D. S.; Northrup, H.; Au, K.-S.; McAllister, K. A.;
Francomano,
[36321] C. A.; Wenstrup, R. J.; Marchuk, D. A.; Kwiatkowski, D. J.:
COL5A1:
[36322] fine genetic mapping and exclusion as candidate gene in
families with
[36323] nail-patella syndrome, tuberous sclerosis 1, hereditary
hemorrhagic
[36324] telangiectasia, and Ehlers-Danlos syndrome type II. Ge-
nomics 25:
[36325] 737-739, 1995.

[36326] 5375.

[36327] Comings, D. E.: Evidence for ancient tetraploidy and con-
servation
[36328] of linkage groups in mammalian chromosomes. Nature
238: 455-467,
[36329] 1972.

[36330] 5376.

[36331] Deol, M. S.: Genetical studies on the skeleton of the
mouse. XXVIII.
[36332] Tail-short. Proc. Roy. Soc. Ser. B. 155: 78-95, 1961.

[36333] 5377.

[36334] Hart, C. P.; Awgulewitsch, A.; Fainsod, A.; McGinnis, W.;
Ruddle,
[36335] F. H.: Homeo box gene complex on mouse chromosome
11: molecular cloning,
[36336] expression in embryogenesis, and homology to a human
homeo box locus. Cell 43:
[36337] 9-18, 1985.

[36338] 5378.

[36339] Hauser, C. A.; Joyner, A. L.; Klein, R. D.; Learned, T. K.;
Martin,

[36340] G. R.; Tjian, R.: Expression of homologous homeo-
box-containing genes
[36341] in differentiated human teratocarcinoma cells and mouse
embryos. Cell 43:
[36342] 19-28, 1985.

[36343] 5379.

[36344] Levine, M.; Rubin, G. M.; Tjian, R.: Human DNA sequences
homologous
[36345] to a protein coding region conserved between homeotic
genes of Drosophila. Cell 38:
[36346] 667-673, 1984.

[36347] 5380.

[36348] Manley, J. L.; Levine, M. S.: The homeo box and mam-
malian development. Cell 43:
[36349] 1-2, 1985.

[36350] 5381.

[36351] Meijlink, F.; de Laaf, R.; Verrijzer, P.; Destree, O.; Kroezen,

[36352] V.; Hilkens, J.; Deschamps, J.: A mouse homeobox con-
taining gene
[36353] on chromosome 11: sequence and tissue-specific expres-
sion. Nucleic

[36354] Acids Res. 15: 6773-6786, 1987.

[36355] 5382.

[36356] Miki, T.; Murphy, P. D.; Pletcher, B. A.; Kidd, J. R.; Ferguson-Smith,

[36357] A. C.; Ruddle, F. H.; Kidd, K. K.: HOX2 maps to 17q near PPY and

[36358] NGFR. (Abstract) Cytogenet. Cell Genet. 46: 662 only, 1987.

[36359] 5383.

[36360] Munke, M.; Cox, D. R.; Jackson, I. J.; Hogan, B. L. M.; Francke,

[36361] U.: The murine Hox-2 cluster of homeo box containing genes maps distal

[36362] on chromosome 11 near the tail-short (Ts) locus. Cytogenet. Cell

[36363] Genet. 42: 236-240, 1986.

[36364] 5384.

[36365] Ruddle, F. H.: Personal Communication. New Haven, Conn. 1987.

[36366] 5385.

[36367] Ruddle, F. H.: Personal Communication. New Haven, Conn.

1985.

[36368] 5386.

[36369] Schughart, K.; Utset, M. F.; Awgulewitsch, A.; Ruddle, F. H.:

[36370] Structure and expression of Hox-2.2, a murine homeobox-containing

[36371] gene. Proc. Nat. Acad. Sci. 85: 5582-5586, 1988.

[36372] 5387.

[36373] Trainor, P. A.; Ariza-McNaughton, L.; Krumlauf, R.: Role of the

[36374] isthmus and FGFs in resolving the paradox of neural crest plasticity

[36375] and prepatterning. Science 295: 1288-1291, 2002.

[36376] 5388.

[36377] Kaur, S.; Singh, G.; Stock, J. L.; Schreiner, C. M.; Kier, A. B.;

[36378] Yager, K. L.; Mucenski, M. L.; Scott, W. J., Jr.; Potter, S. S.:

[36379] Dominant mutation of the murine Hox-2.2 gene results in developmental

[36380] abnormalities. J. Exp. Zool. 264: 323-336, 1992.

[36381] 5389.

[36382] Lill, M. C.; Fuller, J. F.; Herzig, R.; Crooks, G. M.; Gasson,

[36383] J. C.: The role of the homeobox gene, HOX B7, in human myelomonocytic

[36384] differentiation. Blood 85: 692-697, 1995.

[36385] 5390.

[36386] Simeone, A.; Mavilio, F.; Acampora, D.; Giampaolo, A.; Faiella,

[36387] A.; Zappavigna, V.; D'Esposito, M.; Pannese, M.; Russo, G.; Boncinelli,

[36388] E.; Peschle, C.: Two human homeobox genes, c1 and c8: structure analysis

[36389] and expression in embryonic development. Proc. Nat. Acad. Sci. 84:

[36390] 4914-4918, 1987.

[36391] 5391.

[36392] Yaron, Y.; McAdara, J. K.; Lynch, M.; Hughes, E.; Gasson, J. C.

[36393] : Identification of novel functional regions important for the activity

[36394] of HOXB7 in mammalian cells. J. Immun. 166: 5058-5067, 2001.

[36395] 5392.

[36396] Botto, M.; Fong, K. Y.; So, A. K.; Rudge, A.; Walport, M. J.:

[36397] Molecular basis of hereditary C3 deficiency. J. Clin. Invest. 86:

[36398] 1158-1163, 1990.

[36399] 5393.

[36400] Brook, J. D.; Shaw, D. J.; Meredith, L.; Bruns, G. A. P.; Harper,

[36401] P. S.: Localisation of genetic markers and orientation of the linkage

[36402] group on chromosome 19. Hum. Genet. 68: 282-285, 1984.

[36403] 5394.

[36404] Davies, K. E.; Williamson, R.; Ball, S.; Sarfarazi, M.; Mered-

[36405] ith, L.; Fey, G.; Harper, P. S.: C3 DNA sequence and protein polymorphisms

[36406] in linkage analysis of myotonic dystrophy. (Abstract) Cy-

[36407] togenet. Cell Genet. 37: 447 only, 1984.

[36408] 5395.

[36409] de Bruijn, M. H. L.; Fey, G. H.: Human complement component C3:
[36410] cDNA coding sequence and derived primary structure. Proc. Nat. Acad.
[36411] Sci. 82: 708-712, 1985.

[36412] 5396.

[36413] Donald, J. A.; Ball, S. P.: Approximate linkage equilibrium between
[36414] two polymorphic sites within the gene for human complement component
[36415] 3. Ann. Hum. Genet. 48: 269-273, 1984.

[36416] 5397.

[36417] Drouin, S. M.; Corry, D. B.; Kildsgaard, J.; Wetsel, R. A.: Cutting
[36418] edge: the absence of C3 demonstrates a role for complement in Th2
[36419] effector functions in a murine model of pulmonary allergy. J. Immun. 167:
[36420] 4141-4145, 2001.

[36421] 5398.

[36422] Einstein, L. P.; Hansen, P. J.; Ballow, M.; Davis, A. E., III;

[36423] Davis, J. S., IV; Alper, C. A.; Rosen, F. S.; Colten, H. R.: Biosynthesis
[36424] of the third component of complement (C3) in vitro by monocytes from
[36425] both normal and homozygous C3-deficient humans. J. Clin. Invest. 60:
[36426] 963-969, 1977.

[36427] 5399.

[36428] Fong, K. Y.; Botto, M.; Walport, M. J.; So, A. K.: Genomic organization
[36429] of human complement component C3. Genomics 7: 579-586, 1990.
[36430] 5400.

[36431] Gedde-Dahl, T., Jr.; Teisberg, P.; Thorsby, E.: C(3) polymorphism:
[36432] genetic linkage relations. Clin. Genet. 6: 66-72, 1974.

[36433] 5401.

[36434] Goedde, H. W.; Benkmann, H.-G.; Hirth, L.: Genetic polymorphism
[36435] of C-prime-3(beta-1C-globulin) component of complement in a German

[36436] and a Spanish population. Humangenetik 10: 231-234, 1970.

[36437] 5402.

[36438] Grace, H. J.; Brereton-Stiles, G. G.; Vos, G. H.; Schonland, M.

[36439] : A family with partial and total deficiency of complement C3. S.

[36440] Afr. Med. J. 50: 139-140, 1976.

[36441] 5403.

[36442] Hoppe, H. H.; Goedde, H. W.; Agarwal, D. P.; Benkmann, H.-G.;

[36443] Hirth, L.; Janssen, W.: A silent (C-prime-3) producing par-tial deficiency

[36444] of the third component of human complement. Hum. Hered. 28: 141-146,

[36445] 1978.

[36446] 5404.

[36447] Johnson, J. P.; McLean, R. H.; Cork, L. C.; Winkelstein, J. A.

[36448] : Genetic analysis of an inherited deficiency of the third component

[36449] of complement in Brittany spaniel dogs. Am. J. Med.

Genet. 25: 557-562,
[36450] 1986.

[36451] 5405.

[36452] Koch, C.; Behrendt, N.: A novel polymorphism of human complement
[36453] component C3 detected by means of a monoclonal antibody. Immunogenetics 23:
[36454] 322-325, 1986.

[36455] 5406.

[36456] McLean, R. H.; Bryan, R. K.; Winkelstein, J.: Hypomorphic variant
[36457] of the slow allele of C3 associated with hypocomplementemia and hematuria. Am.
[36458] J. Med. 78: 865-868, 1985.

[36459] 5407.

[36460] McLean, R. H.; Siegel, N. J.; Kashgarian, M.: Activation of the
[36461] classic complement pathway in patients with the C3 nephritic factors. Nephron 25:
[36462] 57-64, 1980.

[36463] 5408.

[36464] McLean, R. H.; Wienstein, A.; Chapitis, J.; Lowenstein, M.; Rothfield,

[36465] N. F.: Familial partial deficiency of the third component of complement

[36466] (C3) and the hypocomplementemic cutaneous vasculitis syndrome. Am.

[36467] J. Med. 68: 549-558, 1980.

[36468] 5409.

[36469] Muller-Eberhard, H. J.: Chemistry and reaction mechanisms of

[36470] complement. Adv. Immun. 8: 1-80, 1968.

[36471] 5410.

[36472] Nilsson, U. R.; Nilsson, B.; Storm, K.-E.; Sjolin-Forsberg, G.;

[36473] Hallgren, R.: Hereditary dysfunction of the third component of complement

[36474] associated with a systemic lupus erythematosus-like syndrome and meningococcal

[36475] meningitis. Arthritis Rheum. 35: 580-586, 1992.

[36476] 5411.

[36477] Osofsky, S. G.; Thompson, B. H.; Gewurz, H.; Schmid, F.

R.; Mittal,
[36478] K. K.: Evidence for lack of linkage between HLA and C3 deficiency
[36479] in man. Immunogenetics 4: 195-198, 1977.

[36480] 5412.

[36481] Osofsky, S. G.; Thompson, B. H.; Lint, T. F.; Gewurz, H.: Hereditary
[36482] deficiency of 3rd component of complement in a child with fever, skin
[36483] rash, and arthralgias--response to transfusion of whole blood. J.
[36484] Pediat. 90: 180-186, 1977.

[36485] 5413.

[36486] Power, D. A.; Ng, Y. C.; Simpson, J. G.: Familial incidence of
[36487] C3 nephritic factor, partial lipodystrophy and membra-
noproliferative
[36488] glomerulonephritis. Quart. J. Med. 75: 387-398, 1990.

[36489] 5414.

[36490] Pratt, J. R.; Basheer, S. A.; Sacks, S. H.: Local synthesis of

[36491] complement component C3 regulates acute renal trans-

plant rejection. Nature
[36492] Med. 8: 582-587, 2002.

[36493] 5415.

[36494] Pussell, B. A.; Bourke, E.; Nayef, M.; Morris, S.; Peters, D.

[36495] K.: Complement deficiency and nephritis: a report of a family. Lancet I:
[36496] 675-677, 1980.

[36497] 5416.

[36498] Raum, D.; Donaldson, V. H.; Rosen, F. S.; Alper, C. A.: Genetics
[36499] of complement. Curr. Top. Hemat. 3: 111-174, 1980.

[36500] 5417.

[36501] Sanders, M. F.; Crandall, J.; Huey, B.; Leung, R.; King, M. C.

[36502] : Possible synteny of LE, SE, and C3. (Abstract) Cytogenet. Cell
[36503] Genet. 37: 575 only, 1984.

[36504] 5418.

[36505] Sakai, K.; Yamada, M.; Horiba, N.; Wakui, M.; Demura, H.; Suda,

[36506] T.: The genomic organization of the human corticotropin-releasing
[36507] factor type-1 receptor. Gene 219: 125-130, 1998.

[36508] 5419.

[36509] Sillaber, I.; Rammes, G.; Zimmermann, S.; Mahal, B.; Zieglgansberger,
[36510] W.; Wurst, W.; Holsboer, F.; Spanagel, R.: Enhanced and delayed stress-induced
[36511] alcohol drinking in mice lacking functional CRH1 receptors. Science 296:
[36512] 931-933, 2002.

[36513] 5420.

[36514] Smith, G. W.; Aubry, J.-M.; Dellu, F.; Contarino, A.; Bilezikjian,
[36515] L. M.; Gold, L. H.; Chen, R.; Marchuk, Y.; Hauser, C.; Bentley, C.
[36516] A.; Sawchenko, P. E.; Koob, G. F.; Vale, W.; Lee, K.-F.: Corticotropin
[36517] releasing factor receptor 1-deficient mice display decreased anxiety,
[36518] impaired stress response, and aberrant neuroendocrine development. Neuron 20:

[36519] 1093-1102, 1998.

[36520] 5421.

[36521] Timpl, P.; Spanagel, R.; Sillaber, I.; Kresse, A.; Reul, J. M.

[36522] H. M.; Stalla, G. K.; Blanquet, V.; Steckler, T.; Holsboer, F.; Wurst,

[36523] W.: Impaired stress response and reduced anxiety in mice lacking

[36524] a functional corticotropin-releasing hormone receptor 1. Nature Genet. 19:

[36525] 162-166, 1998.

[36526] 5422.

[36527] Nebert, D. W.: Personal Communication. Bethesda, Md. 1994.

[36528] 5423.

[36529] Yamano, S.; Tatsuno, J.; Gonzalez, F. J.: The CYP2A3 gene product

[36530] catalyzes coumarin 7-hydroxylation in human liver microsomes. Biochemistry 29:

[36531] 1322-1329, 1990.

[36532] 5424.

[36533] Daigo, S.; Takahashi, Y.; Fujieda, M.; Ariyoshi, N.; Ya-
mazaki,
[36534] H.; Koizumi, W.; Tanabe, S.; Saigenji, K.; Nagayama, S.;
Ikeda, K.;
[36535] Nishioka, Y.; Kamataki, T.: A novel mutant allele of the
CYP2A6 gene
[36536] (CYP2A6*11) found in a cancer patient who showed poor
metabolic phenotype
[36537] towards tegafur. Pharmacogenetics 12: 299-306, 2002.

[36538] 5425.

[36539] Ding, S.; Lake, B. G.; Friedberg, T.; Wolf, C. R.: Expression

[36540] and alternative splicing of the cytochrome P-450 CYP2A7.
Biochem.
[36541] J. 306: 161-166, 1995.

[36542] 5426.

[36543] Fernandez-Salguero, P.; Hoffman, S. M. G.; Cholerton, S.;
Mohrenweiser,
[36544] H.; Raunio, H.; Rautio, A.; Pelkonen, O.; Huang, J.; Evans,
W. E.;
[36545] Idle, J. R.; Gonzalez, F. J.: A genetic polymorphism in
coumarin

[36546] 7-hydroxylation: sequence of the human CYP2A genes and identification

[36547] of variant CYP2A6 alleles. Am. J. Hum. Genet. 57: 651-660, 1995.

[36548] 5427.

[36549] Gu, D. F.; Hinks, L. J.; Morton, N. E.; Day, I. N. M.: The use

[36550] of long PCR to confirm three common alleles at the CYP2A6 locus and

[36551] the relationship between genotype and smoking habit. Ann. Hum. Genet. 64:

[36552] 383-390, 2000.

[36553] 5428.

[36554] Azim, A. C.; Knoll, J. H. M.; Beggs, A. H.; Chisti, A. H.: Iso-form

[36555] cloning, actin binding, and chromosomal localization of human erythroid

[36556] dematin, a member of the villin superfamily. J. Biol. Chem. 270:

[36557] 17407-17413, 1995.

[36558] 5429.

[36559] Azim, A. C.; Marfatia, S. M.; Korsgren, C.; Dotimas, E.; Co-

hen,

[36560] C. M.; Chishti, A. H.: Human erythrocyte dematin and protein 4.2

[36561] (pallidin) are ATP binding proteins. Biochemistry 35: 3001-3006,

[36562] 1996.

[36563] 5430.

[36564] Chishti, A. H.; Faquin, W.; Wu, C.-C.; Branton, D.: Purification

[36565] of erythrocyte dematin (protein 4.9) reveals an endogenous protein

[36566] kinase that modulates actin-bundling activity. J. Biol. Chem. 264:

[36567] 8985-8991, 1989.

[36568] 5431.

[36569] Gilligan, D. M.; Bennett, V.: The junctional complex of the membrane

[36570] skeleton. Seminars Hemat. 30: 74-83, 1993.

[36571] 5432.

[36572] Khanna, R.; Chang, S. H.; Andrabi, S.; Azam, M.; Kim, A.; Rivera,

[36573] A.; Brugnara, C.; Low, P. S.; Liu, S.-C.; Chishti, A. H.: Headpiece
[36574] domain of dematin is required for the stability of the erythrocyte
[36575] membrane. Proc. Nat. Acad. Sci. 99: 6637-6642, 2002.

[36576] 5433.

[36577] Peters, L. L.; Eicher, E. M.; Azim, A. C.; Chishti, A. H.: The

[36578] gene encoding the erythrocyte membrane skeleton protein dematin (Epb4.9)
[36579] maps to mouse chromosome 14. Genomics 26: 634-635, 1995.
[36580] 5434.

[36581] Rana, A. P.; Ruff, P.; Maalouf, G. J.; Speicher, D. W.; Chishti,
[36582] A. H.: Cloning of human erythroid dematin reveals another member
[36583] of the villin family. Proc. Nat. Acad. Sci. 90: 6651-6655, 1993.
[36584] 5435.

[36585] Loughlin, J.; Irven, C.; Hardwick, L. J.; Butcher, S.; Walsh, S.;

[36586] Wordsworth, P.; Sykes, B.: Linkage of the gene that encodes the alpha-1
[36587] chain of type V collagen (COL5A1) to type II Ehlers-Danlos syndrome
[36588] (EDS II). Hum. Molec. Genet. 4: 1649-1651, 1995.

[36589] 5436.

[36590] Correas, I.; Speicher, D. W.; Marchesi, V. T.: Structure of the
[36591] spectrin-actin binding site of erythrocyte protein 4.1. J. Biol.
[36592] Chem. 261: 13362-13366, 1986.

[36593] 5437.

[36594] Albig, W.; Drabent, B.; Kunz, J.; Kalff-Suske, M.; Grzeschik, K.-H.;
[36595] Doenecke, D.: All known human H1 histone genes except the H1(0) gene
[36596] are clustered on chromosome 6. Genomics 16: 649-654, 1993.
[36597] 5438.

[36598] Tanguay, R. M.; Berube, D.; Gagne, R.: Localization of histone

[36599] genes to chromosomes 6, 12, and 1 by in situ hybridization. (Abstract) Cytogenet.

[36600] Cell Genet. 46: 702 only, 1987.

[36601] 5439.

[36602] Cheng, J.; Baumhueter, S.; Cacalano, G.; Carver-Moore, K.; Thibodeaux,

[36603] H.; Thomas, R.; Broxmeyer, H. E.; Cooper, S.; Hague, N.; Moore, M.;

[36604] Lasky, L. A.: Hematopoietic defects in mice lacking the sialomucin

[36605] CD34. Blood 87: 479-490, 1996.

[36606] 5440.

[36607] He, X.-Y.; Antao, V. P.; Basila, D.; Marx, J. C.; Davis, B. R.

[36608] : Isolation and molecular characterization of the human CD34 gene. Blood 79:

[36609] 2296-2302, 1992.

[36610] 5441.

[36611] Howell, S. M.; Molgaard, H. V.; Greaves, M. F.; Spurr, N. K.:

[36612] Localisation of the gene coding for the haemopoietic stem cell antigen

[36613] CD34 to chromosome 1q32. Hum. Genet. 87: 625-627, 1991.

[36614] 5442.

[36615] Okuno, Y.; Iwasaki, H.; Huettner, C. S.; Radomska, H. S.; Gonzalez,

[36616] D. A.; Tenen, D. G.; Akashi, K.: Differential regulation of the human

[36617] and murine CD34 genes in hematopoietic stem cells. Proc. Nat. Acad.

[36618] Sci. 99: 6246-6251, 2002.

[36619] 5443.

[36620] Satterthwaite, A. B.; Burn, T. C.; Le Beau, M. M.; Tenen, D. G.

[36621] : Structure of the gene encoding CD34, a human hematopoietic stem

[36622] cell antigen. Genomics 12: 788-794, 1992.

[36623] 5444.

[36624] Simmons, D. L.; Satterthwaite, A. B.; Tenen, D. G.; Seed, B.:

[36625] Molecular cloning of a cDNA encoding CD34, a sialomucin of human hematopoietic

[36626] stem cells. J. Immun. 148: 267-271, 1992.

[36627] 5445.

[36628] Sutherland, D. R.; Stewart, A. K.; Keating, A.: CD34 anti-
gen:

[36629] molecular features and potential clinical applications.
Stem Cells 11

[36630] (suppl. 3): 50-57, 1993.

[36631] 5446.

[36632] Sutherland, D. R.; Watt, S. M.; Dowden, G.; Karhi, K.;
Baker, M.

[36633] A.; Greaves, M. F.; Smart, J. E.: Structural and partial amino
acid

[36634] sequence analysis of the human hemopoietic progenitor
cell antigen

[36635] CD34. Leukemia 2: 793-803, 1988.

[36636] 5447.

[36637] Tenen, D. G.; Satterthwaite, A. B.; Borson, R.; Simmons, D.;
Eddy,

[36638] R. L.; Shows, T. B.: Chromosome 1 localization of the gene
for CD34,

[36639] a surface antigen of human stem cells. Cytogenet. Cell

Genet. 53:
[36640] 55-57, 1990.

[36641] 5448.

[36642] Clegg, J. B.: Can the product of the theta gene be a real globin? Nature 329:
[36643] 465-466, 1987.

[36644] 5449.

[36645] Fei, Y. J.; Fujita, S.; Huisman, T. H. J.: Two different theta

[36646] globin gene deletions observed among black newborn babies. Brit.
[36647] J. Haemat. 68: 249-253, 1988.

[36648] 5450.

[36649] Hsu, S.-L.; Marks, J.; Shaw, J.-P.; Tam, M.; Higgs, D. R.; Shen,
[36650] C. C.; Shen, C.-K. J.: Structure and expression of the human theta-1
[36651] globin gene. Nature 331: 94-96, 1988.

[36652] 5451.

[36653] Leung, S.; Proudfoot, N. J.; Whitelaw, E.: The gene for theta-globin

[36654] is transcribed in human fetal erythroid tissues. Nature 329: 551-554,

[36655] 1987.

[36656] 5452.

[36657] Marks, J.; Shaw, J.-P.; Perez-Stabile, C.; Hu, W.-S.; Ayres, T.

[36658] M.; Shen, C.; Shen, C.-K. J.: The primate alpha-globin gene family:

[36659] a paradigm of the fluid genome. Cold Spring Harbor Symp. Quant. Biol. 51:

[36660] 499-508, 1986.

[36661] 5453.

[36662] Marks, J.; Shaw, J.-P.; Shen, C.-K. J.: Sequence organiza-tion

[36663] and genomic complexity of primary theta-1 globin gene, a novel alpha-globin-like

[36664] gene. Nature 321: 785-788, 1986.

[36665] 5454.

[36666] Peschle, C.; Mavilio, F.; Care, A.; Migliaccio, G.; Migliaccio,

[36667] A. R.; Salvo, G.; Samoggia, P.; Petti, S.; Guerriero, R.; Mar-inucci,

[36668] M.; Lazzaro, D.; Russo, G.; Mastroberardino, G.: Haemoglobin switching
[36669] in human embryos: asynchrony of zeta-to-alpha- and epsilon-to-gamma-globin
[36670] switches in primitive and definitive erythropoietic lineage. Nature 313:
[36671] 235-238, 1985.

[36672] 5455.

[36673] Utsch, B.; Albers, N.; Dame, C.; Bartmann, P.; Lentze, M. J.; Ludwig,
[36674] M.: Homozygous alpha-thalassemia associated with hypospadias: SEA-type
[36675] deletion does not affect expression of the -14 gene and loss of the
[36676] theta-1-globin gene on 16p13.3 is compensated by its duplicate theta-2
[36677] on chromosome 10. (Letter) Am. J. Med. Genet. 101: 286-287, 2001.
[36678] 5456.

[36679] Pogue-Geile, K.; Geiser, J. R.; Shu, M.; Miller, C.; Wool, I. G.;
[36680] Meisler, A. I.; Pipas, J. M.: Ribosomal protein genes are

overexpressed
[36681] in colorectal cancer: isolation of a cDNA clone encoding the human
[36682] S3 ribosomal protein. Molec. Cell. Biol. 11: 3842-3849, 1991.
[36683] 5457.

[36684] Barton, D. E.; Crimaudo, C.; Hortsch, M.; Francke, U.: The genes
[36685] for ribophorins I and II are on human chromosomes 3q and 20 and mouse
[36686] chromosomes 6 and 12, respectively. (Abstract) Cyto- genet. Cell Genet. 46:
[36687] 577 only, 1987.

[36688] 5458.

[36689] Crimaudo, C.; Hortsch, M.; Gausepohl, H.; Meyer, D. I.: Human
[36690] ribophorins I and II: the primary structure and membrane topology
[36691] of two highly conserved round endoplasmic reticulum-spe- cific glycoproteins. EMBO
[36692] J. 6: 75-82, 1987.

[36693] 5459.

[36694] Elsasser, S.; Gali, R. R.; Schwickart, M.; Larsen, C. N.; Leggett,

[36695] D. S.; Muller, B.; Feng, M. T.; Tubing, F.; Dittmar, G. A. G.; Finley,

[36696] D.: Proteasome subunit Rpn1 binds ubiquitin-like protein domains. Nature

[36697] Cell Biol. 4: 725-730, 2002.

[36698] 5460.

[36699] Kelleher, D. J.; Kreibich, G.; Gilmore, R.: Oligosaccharyl-transferase

[36700] activity is associated with a protein complex composed of ribophorins

[36701] I and II and a 48 kd protein. Cell 69: 55-65, 1992.

[36702] 5461.

[36703] Horbach, J. M. L. M.; Brenninkmeyer, S. J.; van de Velde, C. J.

[36704] H.; Nieuwenhuyzen Kruseman, A. C.: A forme fruste of von Hippel-Lindau

[36705] disease--a combination of adrenal pheochromocytoma and ipsilateral

[36706] renal cell carcinoma: a case report. Surgery 105: 436-441, 1989.

[36707] 5462.

[36708] Horton, W. A.; Wong, V.; Eldridge, R.: Von Hippel-Lindau disease--clinical

[36709] and pathological manifestations in 9 families with 50 af- fected members. Arch.

[36710] Intern. Med. 136: 769-777, 1976.

[36711] 5463.

[36712] Hull, M. T.; Roth, L. M.; Glover, J. L.; Walker, P. D.: Metastatic

[36713] carotid body paraganglioma in von Hippel-Lindau disease: an electron
[36714] microscopic study. Arch. Path. Lab. Med. 106: 235-239, 1982.
[36715] 5464.
[36716] Hosoe, S.; Brauch, H.; Latif, F.; Glenn, G.; Daniel, L.; Bale,
[36717] S.; Choyke, P.; Gorin, M.; Oldfield, E.; Berman, A.; Goodman, J.;
[36718] Orcutt, M. L.; Hampsch, K.; Delisio, J.; Modi, W.; McBride, W.; Anglard,
[36719] P.; Weiss, G.; Walther, M. M.; Linehan, W. M.; Lerman, M. I.; Zbar,
[36720] B.: Localization of the von Hippel-Lindau disease gene to a small
[36721] region of chromosome 3. Genomics 8: 634-640, 1990.
[36722] 5465.
[36723] Huson, S. M.; Harper, P. S.; Hourihan, M. D.; Cole, G.; Weeks,
[36724] R. D.; Compston, D. A. S.: Cerebellar haemangioblastoma and von Hippel-Lindau
[36725] disease. Brain 109: 1297-1310, 1986.

[36726] 5466.

[36727] Iliopoulos, O.; Kibel, A.; Gray, S.; Kaelin, W. G., Jr.: Tumour

[36728] suppression by the human von Hippel-Lindau gene prod-
uct. Nature Med. 1:

[36729] 822-826, 1995.

[36730] 5467.

[36731] Iliopoulos, O.; Levy, A. P.; Jiang, C.; Kaelin, W. G., Jr.;
Goldberg,

[36732] M. A.: Negative regulation of hypoxia-inducible genes by
the von

[36733] Hippel-Lindau protein. Proc. Nat. Acad. Sci. 93:
10595-10599, 1996.

[36734] 5468.

[36735] Iliopoulos, O.; Ohh, M.; Kaelin, W. G., Jr.: pVHL(19) is a bi-
ologically

[36736] active product of the von Hippel-Lindau gene arising from
internal

[36737] translation initiation. Proc. Nat. Acad. Sci. 95:
11661-11666, 1998.

[36738] 5469.

[36739] Ivan, M.; Kondo, K.; Yang, H.; Kim, W.; Valiando, J.; Ohh,

M.;

[36740] Salic, A.; Asara, J. M.; Lane, W. S.; Kaelin, W. G., Jr.: HIF-alpha

[36741] targeted for VHL-mediated destruction by proline hydroxylation: implications

[36742] for O(2) sensing. Science 292: 464-468, 2001.

[36743] 5470.

[36744] Ivanov, S. V.; Kuzmin, I.; Wei, M.-H.; Pack, S.; Geil, L.; Johnson,

[36745] B. E.; Stanbridge, E. J.; Lerman, M. I.: Down-regulation of transmembrane

[36746] carbonic anhydrases in renal cell carcinoma cell lines by wild-type

[36747] von Hippel-Lindau transgenes. Proc. Nat. Acad. Sci. 95: 12596-12601,

[36748] 1998.

[36749] 5471.

[36750] Jaakkola, P.; Mole, D. R.; Tian, Y.-M.; Wilson, M. I.; Gielbert,

[36751] J.; Gaskell, S. J.; von Kriegsheim, A.; Hebestreit, H. F.; Mukherji,

[36752] M.; Schofield, C. J.; Maxwell, P. H.; Pugh, C. W.; Ratcliffe, P.

J.

[36753] : Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex

[36754] by O(2)-regulated prolyl hydroxylation. Science 292: 468-472, 2001.

[36755] 5472.

[36756] James, G. P.: Personal Communication. Springfield, Ohio 8/15/1998.

[36757] 5473.

[36758] Jennings, A. M.; Smith, C.; Cole, D. R.; Jennings, C.; Shortland,

[36759] J. R.; Williams, J. L.; Brown, C. B.: Von Hippel-Lindau disease in

[36760] a large British family: clinicopathological features and recommendations

[36761] for screening and follow-up. Quart. J. Med. 66: 233-249, 1988.

[36762] 5474.

[36763] Kanno, H.; Kondo, K.; Ito, S.; Yamamoto, I.; Fujii, S.; Torigoe,

[36764] S.; Sakai, N.; Hosaka, M.; Shuin, T.; Yao, M.: Somatic mutations

[36765] of the von Hippel-Lindau tumor suppressor gene in sporadic central
[36766] nervous system hemangioblastomas. Cancer Res. 54: 4845-4847, 1994.
[36767] 5475.

[36768] Kanno, H.; Saljooque, F.; Yamamoto, I.; Hattori, S.; Yao, M.;
[36769] Shuin, T.; U, H.-S.: Role of the von Hippel-Lindau tumor suppressor
[36770] protein during neuronal differentiation. Cancer Res. 60: 2820-2824,
[36771] 2000.

[36772] 5476.

[36773] Kaplan, C.; Sayre, G. P.; Greene, L. F.: Bilateral nephrogenic
[36774] carcinomas in Lindau-von Hippel disease. J. Urol. 86: 36-42, 1961.
[36775] 5477.

[36776] Karsdorp, N.; Elderson, A.; Wittebol-Post, D.; Hene, R. J.; Vos,
[36777] J.; Feldberg, M. A. M.; van Gils, A. P. G.; Jansen-Schillhorn van

[36778] Veen, J. M.; Vroom, T. M.; Hoppener, J. W. M.; Lips, C. J. M.: Von

[36779] Hippel-Lindau disease: new strategies in early detection and treatment. Am.

[36780] J. Med. 97: 158-168, 1994.

[36781] 5478.

[36782] Keeler, L. L., III; Klauber, G. T.: von Hippel-Lindau disease

[36783] and renal cell carcinoma in a 16-year-old boy. J. Urol. 147: 1588-1591,

[36784] 1992.

[36785] 5479.

[36786] Kenck, C.; Wilhelm, M.; Bugert, P.; Staehler, G.; Kovacs, G.:

[36787] Mutation of the VHL gene is associated exclusively with the development

[36788] of non-papillary renal cell carcinomas. J. Path. 179: 157-161, 1996.

[36789] 5480.

[36790] Kerr, D. J.; Scheithauer, B. W.; Miller, G. M.; Ebersold, M. J.;

[36791] McPhee, T. J.: Hemangioblastoma of the optic nerve: case report. Neurosurgery 36:

[36792] 573-581, 1995.

[36793] 5481.

[36794] Kibel, A.; Iliopoulos, O.; DeCaprio, J. A.; Kaelin, W. G., Jr.

[36795] : Binding of the von Hippel-Lindau tumor suppressor protein to elongin

[36796] B and C. Science 269: 1444-1446, 1995.

[36797] 5482.

[36798] Kiechle-Schwarz, M.; Neumann, H. P. H.; Decker, H.-J. H.; Dietrich,

[36799] C.; Wullich, B.; Schempp, W.: Cytogenetic studies on three pheochromocytomas

[36800] derived from patients with von Hippel-Lindau syndrome. Hum. Genet. 82:

[36801] 127-130, 1989.

[36802] 5483.

[36803] Spranger, M.; Schapera, J.: Anomalous inheritance in a kindred

[36804] with split hand, split foot malformation. Europ. J. Pediat. 147:

[36805] 202-205, 1988.

[36806] 5484.

[36807] Brown, M. H.; Gorman, P. A.; Sewell, W. A.; Spurr, N. K.; Sheer,

[36808] D.; Crumpton, M. J.: The gene coding for the human T-lymphocyte CD2

[36809] antigen is located on chromosome 1p. Hum. Genet. 76: 191-195, 1987.

[36810] 5485.

[36811] Clayton, L. K.; Ramachandran, H.; Pravtcheva, D.; Chen, Y.-F.;

[36812] Diamond, D. J.; Ruddle, F. H.; Reinherz, E. L.: The gene for T11

[36813] (CD2) maps to chromosome 1 in humans and to chromosome 3 in mice. J.

[36814] Immun. 140: 3617-3621, 1988.

[36815] 5486.

[36816] Diamond, D. J.; Clayton, L. K.; Sayre, P. H.; Reinherz, E. L.:

[36817] Exon-intron organization and sequence comparison of human and murine

[36818] T11 (CD2) genes. Proc. Nat. Acad. Sci. 85: 1615-1619, 1988.

[36819] 5487.

[36820] Festenstein, R.; Tolaini, M.; Corbella, P.; Mamalaki, C.; Parrington,

[36821] J.; Fox, M.; Miliou, A.; Jones, M.; Kioussis, D.: Locus control region

[36822] function and heterochromatin-induced position effect variegation. Science 271:

[36823] 1123-1125, 1996.

[36824] 5488.

[36825] Kingsmore, S. F.; Watson, M. L.; Moseley, W. S.; Seldin, M. F.

[36826] : Physical linkage of genes encoding the lymphocyte adhesion molecules

[36827] CD2 and its ligand LFA-3. Immunogenetics 30: 123-125, 1989.

[36828] 5489.

[36829] Lang, G.; Wotton, D.; Owen, M. J.; Sewell, W. A.; Brown, M. H.;

[36830] Mason, D. Y.; Crumpton, M. J.; Kioussis, D.: The structure of the

[36831] human CD2 gene and its expression in transgenic mice. EMBO J. 7:

[36832] 1675-1682, 1988.

[36833] 5490.

[36834] Richardson, N. E.; Chang, H.-C.; Brown, N. R.; Hussey, R. E.; Sayre,

[36835] P. H.; Reinherz, E. L.: Adhesion domain of human T11 (CD2) is encoded

[36836] by a single exon. Proc. Nat. Acad. Sci. 85: 5176-5180, 1988.

[36837] 5491.

[36838] Seed, B.; Aruffo, A.: Molecular cloning of the CD2 antigen, the

[36839] T-cell erythrocyte receptor, by a rapid immunoselection procedure. Proc.

[36840] Nat. Acad. Sci. 84: 3365-3369, 1987.

[36841] 5492.

[36842] Sewell, W. A.; Brown, M. H.; Dunne, J.; Owen, M. J.; Crumpton,

[36843] M. J.: Molecular cloning of the human T-lymphocyte sur- face CD2 (T11)

[36844] antigen. Proc. Nat. Acad. Sci. 83: 8718-8722, 1986.

[36845] 5493.

[36846] Wang, J.; Smolyar, A.; Tan, K.; Liu, J.; Kim, M.; Sun, Z. J.;

[36847] Wagner, G.; Reinherz, E. L.: Structure of a heterophilic adhesion

[36848] complex between the human CD2 and CD58 (LFA-3) counterreceptors. Cell 97:

[36849] 791-803, 1999.

[36850] 5494.

[36851] Bibbins, K. B.; Tsai, J.-Y.; Schimenti, J.; Sarvetnick, N.; Zoghbi,

[36852] H. Y.; Goodfellow, P.; Silver, L. M.: Human homologs of two testes-expressed

[36853] loci on mouse chromosome 17 map to opposite arms of chromosome 6.

[36854] Genomics 5: 139-143, 1989.

[36855] 5495.

[36856] Blanche, H.; de Gouyon, B.; Kolaga, V.; Dausset, J.; Silver, L.;

[36857] Cann, H.; Spielman, R.: Three loci homologous with murine t complex

[36858] genes are localized in a 4 point linkage group on the long arm of

[36859] chromosome 6 (6q). (Abstract) Cytogenet. Cell Genet. 51: 963 only,
[36860] 1989.

[36861] 5496.

[36862] Gogolin, K. J.; Wright, L. G.; Kolaga, V. J.; Silver, L. M.; Spielman,
[36863] R. S.: RFLPs detected with the human TCP10 gene, a homologue of a
[36864] mouse t-complex gene. Nucleic Acids Res. 19: 4313 only, 1991.
[36865] 5497.

[36866] Islam, S. D.; Pilder, S. H.; Decker, C. L.; Cebra-Thomas, J. A.;
[36867] Silver, L. M.: The human homolog of a candidate mouse t complex responder
[36868] gene: conserved motifs and evolution with punctuated equilibria. Hum.
[36869] Molec. Genet. 2: 2075-2079, 1993.

[36870] 5498.

[36871] Silver, L. M.: The peculiar journey of a selfish chromosome: mouse

[36872] t haplogypes and meiotic drive. Trends Genet. 9: 250-254, 1993.

[36873] 5499.

[36874] Brissenden, J. E.; Derynck, R.; Francke, U.: Transforming growth

[36875] factor alpha gene (TGFA) maps to human chromosome 2 close to the breakpoint

[36876] of the t(2;8) variant translocation in Burkitt lymphoma. (Abstract) Cytogenet.

[36877] Cell Genet. 40: 589 only, 1985.

[36878] 5500.

[36879] Collin, G. B.; Marshall, J. D.; Naggert, J. K.; Nishina, P. M.

[36880] : TGFA: exon-intron structure and evaluation as a candidate gene for

[36881] Alstrom syndrome. (Letter) Clin. Genet. 55: 61-62, 1999.

[36882] 5501.

[36883] Ellis, D. L.; Kafka, S. P.; Chow, J. C.; Nanney, L. B.; Inman,

[36884] W. H.; McCadden, M. E.; King, L. E., Jr.: Melanoma, growth factors,

[36885] acanthosis nigricans, the sign of Leser-Trelat, and multiple acrochordons:

[36886] a possible role for alpha-transforming growth factor in cutaneous
[36887] paraneoplastic syndromes. New Eng. J. Med. 317: 1582-1587, 1987.
[36888] 5502.

[36889] Fernandez-Larrea, J.; Merlos-Suarez, A.; Urena, J. M.; Baselga,
[36890] J.; Arribas, J.: A role for a PDZ protein in the early secre-tory
[36891] pathway for the targeting of proTGF-alpha to the cell sur-face. Molec.
[36892] Cell 3: 423-433, 1999.

[36893] 5503.

[36894] Fowler, K. J.; Mann, G. B.; Dunn, A. R.: Linkage of the murine
[36895] transforming growth factor-alpha gene with Igk, Ly-2, and Fabp1 on
[36896] chromosome 6. Genomics 16: 782-784, 1993.

[36897] 5504.

[36898] Tam, J. P.; Scheikh, M. A.; Solomon, D. S.; Ossowski, L.: Efficient

[36899] synthesis of human type alpha transforming growth factor: its physical
[36900] and biological characterization. Proc. Nat. Acad. Sci. 83: 8082-8086,
[36901] 1986.

[36902] 5505.

[36903] Tricoli, J. V.; Nakai, H.; Byers, M. G.; Rall, L. B.; Bell, G.

[36904] I.; Shows, T. B.: Assignment of the gene coding for human TGF-alpha
[36905] to chromosome 2p13. (Abstract) Cytogenet. Cell Genet. 40: 762 only,
[36906] 1985.

[36907] 5506.

[36908] Tricoli, J. V.; Nakai, H.; Byers, M. G.; Rall, L. B.; Bell, G.

[36909] I.; Shows, T. B.: The gene for human transforming growth factor alpha
[36910] is on the short arm of chromosome 2. Cytogenet. Cell Genet. 42:
[36911] 94-98, 1986.

[36912] 5507.

[36913] Awad, M. R.; El-Gamel, A.; Hasleton, P.; Turner, D. M.; Sinnott,
[36914] P. J.; Hutchinson, I. V.: Genotypic variation in the trans-forming
[36915] growth factor-beta-1 gene: association with transforming growth factor-beta-1
[36916] production, fibrotic lung disease, and graft fibrosis after lung transplantation. Transplantation 66:
[36917] 1014-1020, 1998.

[36918] 5508.

[36919] Azar, S. T.; Salti, I.; Zantout, M. S.; Major, S.: Alterations

[36920] in plasma transforming growth factor beta in normoalbu-minuric type
[36921] 1 and type 2 diabetic patients. J. Clin. Endocr. Metab. 85: 4680-4682,
[36922] 2000.

[36923] 5509.

[36924] Bernasconi, P.; Torchiana, E.; Confalonieri, P.; Brugnoni, R.;
[36925] Barresi, R.; Mora, M.; Conelio, F.; Morandi, L.; Mantegazza, R.:

[36926] Expression of transforming growth factor-beta-1 in dystrophic patient

[36927] muscles correlates with fibrosis: pathogenetic role of a fibrogenic

[36928] cytokine. J. Clin. Invest. 96: 1137-1144, 1995.

[36929] 5510.

[36930] Blobe, G. C.; Schiemann, W. P.; Lodish, H. F.: Role of transforming

[36931] growth factor beta in human disease. New Eng. J. Med. 342: 1350-1358,

[36932] 2000.

[36933] 5511.

[36934] Border, W. A.; Noble, N. A.: Transforming growth factor beta in

[36935] tissue fibrosis. New Eng. J. Med. 331: 1286-1292, 1994.

[36936] 5512.

[36937] Border, W. A.; Noble, N. A.: Fibrosis linked to TGF-beta in yet

[36938] another disease. (Editorial) J. Clin. Invest. 96: 655-656, 1995.

[36939] 5513.

[36940] Clouthier, D. E.; Comerford, S. A.; Hammer, R. E.: Hepatic fibrosis,
[36941] glomerulosclerosis, and lipodystrophy-like PEPCK-TGF-beta-1 transgenic
[36942] mice. J. Clin. Invest. 100: 2697-2713, 1997.

[36943] 5514.

[36944] Crawford, S. E.; Stellmach, V.; Murphy-Ullrich, J. E.; Ribeiro,
[36945] S. M. F.; Lawler, J.; Hynes, R. O.; Boivin, G. P.; Bouck, N.: Thrombospondin-1
[36946] is a major activator of TGF-beta-1 in vivo. Cell 93: 1159-1170,
[36947] 1998.

[36948] 5515.

[36949] Derynck, R.; Akhurst, R. J.; Balmain, A.: TGF-beta signaling
[36950] in tumor suppression and cancer progression. Nature Genet. 29: 117-129,
[36951] 2001. Note: Erratum: Nature Genet. 29: 351 only, 2001.

[36952] 5516.

[36953] Derynck, R.; Jarrett, J. A.; Chen, E. Y.; Eaton, D. H.; Bell,

[36954] J. R.; Assoian, R. K.; Roberts, A. B.; Sporn, M. B.; Goeddel, D. V.

[36955] : Human transforming growth factor-beta complementary DNA sequence

[36956] and expression in normal and transformed cells. Nature 316: 701-705,

[36957] 1985.

[36958] 5517.

[36959] Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;

[36960] ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt-Clermont, P.

[36961] J.: Deficient Smad7 expression: a putative molecular defect in scleroderma. Pr

[36962] oc. Nat. Acad. Sci. 99: 3908-3913, 2002.

[36963] 5518.

[36964] Dubois, C. M.; Laprise, M.-H.; Blanchette, F.; Gentry, L. E.;

[36965] Leduc, R.: Processing of transforming growth factor beta-1 precursor

[36966] by human furin convertase. J. Biol. Chem. 270: 10618-10624, 1995.

[36967] 5519.

[36968] Fujii, D.; Brissenden, J. E.; Derynck, R.; Francke, U.: Transforming

[36969] growth factor beta gene maps to human chromosome 19 long arm and to

[36970] mouse chromosome 7. Somat. Cell Molec. Genet. 12: 281-288, 1986.

[36971] 5520.

[36972] Fujii, D. M.; Brissenden, J. E.; Derynck, R.; Francke, U.: Transforming

[36973] growth factor beta gene (TGFB) maps to human chromosome 19. (Abstract) Cytogenet.

[36974] Cell Genet. 40: 632, 1985.

[36975] 5521.

[36976] Grainger, D. J.; Heathcote, K.; Chiano, M.; Snieder, H.; Kemp,

[36977] P. R.; Metcalfe, J. C.; Carter, N. D.; Spector, T. D.: Genetic control

[36978] of the circulating concentration of transforming growth factor type

[36979] beta-1. Hum. Molec. Genet. 8: 93-97, 1999.

[36980] 5522.

[36981] Dickinson, M. E.; Kobrin, M. S.; Silan, C. M.; Kingsley, D. M.;

[36982] Justice, M. J.; Miller, D. A.; Ceci, J. D.; Lock, L. F.; Lee, A.;

[36983] Buchberg, A. M.; Siracusa, L. D.; Lyons, K. M.; Derynck, R.; Hogan,

[36984] B. L. M.; Copeland, N. G.; Jenkins, N. A.: Chromosomal lo-

calization

[36985] of seven members of the murine TGF-beta superfamily suggests close

[36986] linkage to several morphogenetic mutant loci. Genomics 6: 505-520,

[36987] 1990.

[36988] 5523.

[36989] Heldin, C.-H.; Miyazono, K.; ten Dijke, P.: TGF-beta sig-

nalling

[36990] from cell membrane to nucleus through SMAD proteins. Nature 390:

[36991] 465-471, 1997.

[36992] 5524.

[36993] Thomas, S. A.; Matsumoto, A. M.; Palmiter, R. D.: Nora-

drenaline

[36994] is essential for mouse fetal development. Nature 374:
643-646, 1995.

[36995] 5525.

[36996] Drebin, J. A.; Hartzell, S. W.; Griffin, C.; Campbell, M. J.;
Niederhuber,

[36997] J. E.: Molecular cloning and chromosomal localization of
the human

[36998] homologue of a B-lymphocyte specific protein tyrosine ki-
nase (blk). Oncogene 10:

[36999] 477-486, 1995.

[37000] 5526.

[37001] Dymecki, S.; Niederhuber, J.; Desiderio, S.: Specific ex-
pression

[37002] of a novel tyrosine kinase gene, Blk, in B lymphoid cells.
Science 247:

[37003] 332-336, 1990.

[37004] 5527.

[37005] Islam, K. B.; Rabbani, H.; Larsson, C.; Sanders, R.; Smith,
C.

[37006] I. E.: Molecular cloning, characterization, and chromoso-
mal localization

[37007] of a human lymphoid tyrosine kinase related to murine Blk. J. Immun. 154:

[37008] 1265-1272, 1995.

[37009] 5528.

[37010] Kozak, C. A.; Dymecki, S. M.; Niederhuber, J. E.; Desiderio, S.

[37011] V.: Genetic mapping of the gene for a novel tyrosine kinase, Blk,

[37012] to mouse chromosome 14. Genomics 9: 762-764, 1991.

[37013] 5529.

[37014] Aiello, L. P.; Pierce, E. A.; Foley, E. D.; Takagi, H.; Chen, H.;

[37015] Riddle, L.; Ferrara, N.; King, G. L.; Smith, L. E. H.: Suppression

[37016] of retinal neovascularization in vivo by inhibition of vascular endothelial

[37017] growth factor (VEGF) using soluble VEGF-receptor chimeric proteins. Proc.

[37018] Nat. Acad. Sci. 92: 10457-10461, 1995.

[37019] 5530.

[37020] Kikuchi, S.; Hata, M.; Fukumoto, K.; Yamane, Y.; Matsui, T.; Tamura,

[37021] A.; Yonemura, S.; Yamagishi, H.; Keppler, D.; Tsukita, S.; Tsukita,
[37022] S.: Radixin deficiency causes conjugated hyperbilirubine-mia with
[37023] loss of Mrp2 from bile canalicular membranes. Nature Genet. 31:
[37024] 320-325, 2002.

[37025] 5531.

[37026] Wilgenbus, K. K.; Milatovich, A.; Francke, U.; Furthmayr, H.:
[37027] Molecular cloning, cDNA sequence, and chromosomal as-signment of the
[37028] human radixin gene and two dispersed pseudogenes. Ge-nomics 16: 199-206,
[37029] 1993.

[37030] 5532.

[37031] Geppert, M.; Goda, Y.; Stevens, C. F.; Sudhof, T. C.: The small
[37032] GTP-binding protein Rab3A regulates a late step in synaptic vesicle
[37033] fusion. Nature 387: 810-814, 1997.

[37034] 5533.

[37035] Kapfhamer, D.; Valladares, O.; Sun, Y.; Nolan, P. M.; Rux, J. J.;

[37036] Arnold, S. E.; Veasey, S. C.; Bucan, M.: Mutations in Rab3a alter

[37037] circadian period and homeostatic response to sleep loss in the mouse. Nature

[37038] Genet. 32: 290-295, 2002.

[37039] 5534.

[37040] Rousseau-Merck, M. F.; Zahraoui, A.; Bernheim, A.; Touchot, N.;

[37041] Miglierina, R.; Tavitian, A.; Berger, R.: Chromosome mapping of the

[37042] human RAS related RAB3A and RAB3B genes to chromosomes 19p13.2 and

[37043] 1p31-p32, respectively. (Abstract) Cytogenet. Cell Genet. 51: 1070

[37044] only, 1989.

[37045] 5535.

[37046] Rousseau-Merck, M. F.; Zahraoui, A.; Bernheim, A.; Touchot, N.;

[37047] Miglierina, R.; Tavitian, A.; Berger, R.: Chromosome mapping of the
[37048] human RAS-related RAB3A gene to 19p13.2. Genomics 5: 694-698, 1989.
[37049] 5536.
[37050] Zahraoui, A.; Touchot, N.; Chardin, P.; Tavitian, A.: The human
[37051] rab genes encode a family of GTP-binding proteins related to yeast
[37052] YPT1 and SEC4 products involved in secretion. J. Biol. Chem. 264:
[37053] 12394-12401, 1989.
[37054] 5537.
[37055] Hussey, D. J.; Nicola, M.; Moore, S.; Peters, G. B.; Dobrovic,
[37056] A.: The (4;11)(q21;p15) translocation fuses the NUP98 and RAP1GDS1
[37057] genes and is recurrent in T-cell acute lymphocytic leukemia. Blood 94:
[37058] 2072-2079, 1999.
[37059] 5538.
[37060] Cheng, G.-X.; Zhu, X.-H.; Men, X.-Q.; Wang, L.; Huang,

Q.-H.;

[37061] Jin, X. L.; Xiong, S.-M.; Zhu, J.; Guo, W.-M.; Chen, J.-Q.; Xu, S.-F.;

[37062] So, E.; Chan, L.-C.; Waxman, S.; Zelent, A.; Chen, G.-Q.; Dong, S.;

[37063] Liu, J.-X.; Chen, S.-J.: Distinct leukemia phenotypes in transgenic

[37064] mice and different corepressor interactions generated by promyelocytic

[37065] leukemia variant fusion genes PLZF-RAR-alpha and NPM-RAR-alpha. Proc.

[37066] Nat. Acad. Sci. 96: 6318-6323, 1999.

[37067] 5539.

[37068] de The, H.; Chomienne, C.; Lanotte, M.; Degos, L.; Dejean, A.

[37069] : The t(15;17) translocation of acute promyelocytic leukaemia fuses

[37070] the retinoic acid receptor alpha gene to a novel tran-scribed locus. Nature 347:

[37071] 558-561, 1990.

[37072] 5540.

[37073] Germain, P.; Iyer, J.; Zechel, C.; Gronemeyer, H.: Co-

regulator
[37074] recruitment and the mechanism of retinoic acid receptor synergy. Nature 415:
[37075] 187-192, 2002.

[37076] 5541.

[37077] Giguere, V.; Ong, E. S.; Segui, P.; Evans, R. M.: Identification
[37078] of a receptor for the morphogen retinoic acid. Nature 330: 624-629,
[37079] 1987.

[37080] 5542.

[37081] He, L.-Z.; Bhaumik, M.; Tribioli, C.; Rego, E. M.; Ivins, S.;

[37082] Zelent, A.; Pandolfi, P. P.: Two critical hits for promyelocytic
[37083] leukemia. Molec. Cell 6: 1131-1141, 2000.

[37084] 5543.

[37085] Hiorns, L. R.; Min, T.; Swansbury, G. J.; Zelent, A.; Dyer, M.

[37086] J. S.; Catovsky, D.: Interstitial insertion of retinoic acid receptor-alpha
[37087] gene in acute promyelocytic leukemia with normal chro-

mosomes 15 and 5544.

[37088] Blood 83: 2946-2951, 1994.

[37089]

[37090] 17. Hjalt, T. A. H.; Murray, J. C.: Genomic structure of the human

[37091] retinoic acid receptor-alpha-1 gene. Mammalian Genome 10: 528-259,

[37092] 1999.

[37093] 5545.

[37094] Kanno, K.; Suzuki, Y.; Yang, X.; Yamada, A.; Aoki, Y.; Kure, S.;

[37095] Matsubara, Y.: Lack of evidence for a significant associa-tion between

[37096] nonsyndromic cleft lip with or without cleft palate and the retinoic

[37097] acid receptor alpha gene in the Japanese population. J. Hum. Genet. 47:

[37098] 269-274, 2002.

[37099] 5546.

[37100] Lemons, R. S.; Eilender, D.; Waldmann, R. A.; Rebentisch, M.;

[37101] Frej, A.-K.; Ledbetter, D. H.; Willman, C.; McConnell, T.; O'Connell,
[37102] P.: Cloning and characterization of the t(15;17) translocation breakpoint
[37103] region in acute promyelocytic leukemia. Genes Chromosomes Cancer 2:
[37104] 79-87, 1990.

[37105] 5547.

[37106] Lin, R. J.; Evans, R. M.: Acquisition of oncogenic potential

[37107] by RAR chimeras in acute promyelocytic leukemia through formation
[37108] of homodimers. Molec. Cell 5: 821-830, 2000.

[37109] 5548.

[37110] Mattei, M.-G.; Petkovich, M.; Mattei, J.-F.; Brand, N.; Chambon,
[37111] P.: Mapping of the human retinoic acid receptor to the q21 band of
[37112] chromosome 17. Hum. Genet. 80: 186-188, 1988.

[37113] 5549.

[37114] McNamara, P.; Seo, S.; Rudic, R. D.; Sehgal, A.; Chakravarti,

[37115] D.; FitzGerald, G. A.: Regulation of CLOCK and MOP4 by nuclear hormone

[37116] receptors in the vasculature: a humoral mechanism to re-set a peripheral

[37117] clock. Cell 105: 877-889, 2001.

[37118] 5550.

[37119] Pandolfi, P. P.: Oncogenes and tumor suppressors in the molecular

[37120] pathogenesis of acute promyelocytic leukemia. Hum. Molec. Genet. 10:

[37121] 769-775, 2001.

[37122] 5551.

[37123] Petkovich, M.; Brand, N. J.; Krust, A.; Chambon, P.: A human

[37124] retinoic acid receptor which belongs to the family of nuclear receptors. Nature 330:

[37125] 444-450, 1987.

[37126] 5552.

[37127] Rassool, F.; Martiat, P.; Klisak, I.; Taj, A.; Goldman, J.: Interstitial

[37128] insertion of varying amounts of ABL-containing genetic

material into
[37129] chromosome 22 in Ph-negative CML. Leukemia 4:
273-277, 1990.
[37130] 5553.

[37131] Robertson, M.: Retinoic acid receptor: towards a biochem-
istry
[37132] of morphogenesis. Nature 330: 420-421, 1987.

[37133] 5554.

[37134] Zelent, A.; Guidez, F.; Melnick, A.; Waxman, S.; Licht, J. D.

[37135] : Translocations of the RAR-alpha gene in acute promye-
locytic leukemia. Oncogene 20:
[37136] 7186-7203, 2001.

[37137] 5555.

[37138] Almasan, A.; Mangelsdorf, D. J.; Ong, E. S.; Wahl, G. M.;
Evans,
[37139] R. M.: Chromosomal localization of the human retinoid X
receptors. Genomics 20:
[37140] 397-403, 1994.

[37141] 5556.

[37142] Claudel, T.; Leibowitz, M. D.; Fievet, C.; Tailleux, A.; Wag-

ner,
[37143] B.; Repa, J. J.; Torpier, G.; Lobaccaro, J.-M.; Paterniti, J. R.;
[37144] Mangelsdorf, D. J.; Heyman, R. A.; Auwerx, J.: Reduction of atherosclerosis
[37145] in apolipoprotein E knockout mice by activation of the retinoid X
[37146] receptor. Proc. Nat. Acad. Sci. 98: 2610-2615, 2001.
[37147] 5557.
[37148] de Urquiza, A. M.; Liu, S.; Sjoberg, M.; Zetterstrom, R. H.; Griffiths,
[37149] W.; Sjovall, J.; Perlmann, T.: Docosahexaenoic acid, a lig-and for
[37150] the retinoid X receptor in mouse brain. Science 290: 2140-2144,
[37151] 2000.
[37152] 5558.
[37153] Thomas, G. D.; Sander, M.; Lau, K. S.; Huang, P. L.; Stull, J.
[37154] T.; Victor, R. G.: Impaired metabolic modulation of alpha-adrenergic
[37155] vasoconstriction in dystrophin-deficient skeletal muscle. Proc. Nat.

[37156] Acad. Sci. 95: 15090-15095, 1998.

[37157] 5559.

[37158] Xie, J.; Roddy, P.; Rife, T. K.; Murad, F.; Young, A. P.: Two

[37159] closely linked but separable promoters for human neu-
ronal nitric oxide

[37160] synthase gene transcription. Proc. Nat. Acad. Sci. 92:
1242-1246,

[37161] 1995.

[37162] 5560.

[37163] Xu, W.; Gorman, P.; Sheer, D.; Bates, G.; Kishimoto, J.;
Lizhi,

[37164] L.; Emson, P.: Regional localization of the gene coding for
human

[37165] brain nitric oxide synthase (NOS1) to 12q24.2-24.31 by
fluorescent

[37166] in situ hybridization. Cytogenet. Cell Genet. 64: 62-63,
1993.

[37167] 5561.

[37168] Abeliovich, A.; Schmitz, Y.; Farinas, I.; Choi-Lundberg, D.;
Ho,

[37169] W.-H.; Castillo, P. E.; Shinsky, N.; Verdugo, J. M. G.; Ar-

manini,

[37170] M.; Ryan, A.; Hynes, M.; Phillips, H.; Sulzer, D.; Rosenthal, A.:

[37171] Mice lacking alpha-synuclein display functional deficits in the nigrostriatal

[37172] dopamine system. Neuron 25: 239-252, 2000.

[37173] 5562.

[37174] Teyssier, J. R.; Henry, I.; Dozier, C.; Ferre, D.; Adnet, J. J.;

[37175] Pluot, M.: Recurrent deletion of the short arm of chromosome 3 in

[37176] human renal cell carcinoma: shift of the c-raf 1 locus. J. Nat. Cancer

[37177] Inst. 77: 1187-1191, 1986.

[37178] 5563.

[37179] Taniyama, T.; Takai, S.; Miyazaki, E.; Fukumura, R.; Sato, J.;

[37180] Kobayashi, Y.; Hirakawa, T.; Moore, K. W.; Yamada, K.: The human

[37181] interleukin-10 receptor gene maps to chromosome 11q23.3. Hum. Genet. 95:

[37182] 99-101, 1995.

[37183] 5564.

[37184] Bono, M. R.; Alcaide-Loridan, C.; Couillin, P.; Letouze, B.; Grisard,

[37185] M. C.; Jouin, H.; Fellous, M.: Human chromosome 16 encodes a factor

[37186] involved in induction of class II major histocompatibility antigens

[37187] by interferon gamma. Proc. Nat. Acad. Sci. 88: 6077-6081, 1991.

[37188] 5565.

[37189] Bono, R.; Hatat, D.; Couillin, P.; Grisard, M. C.; Van Cong, N.;

[37190] Fisher, D.; Fellous, M.: Receptor for human gamma interferon is specified

[37191] by human chromosomes 6 and 21. (Abstract) Cytogenet. Cell Genet. 46:

[37192] 584, 1987.

[37193] 5566.

[37194] Dorman, S. E.; Holland, S. M.: Mutation in the signal-transducing

[37195] chain of the interferon-gamma receptor and susceptibility to mycobacterial

[37196] infection. J. Clin. Invest. 101: 2364-2369, 1998.

[37197] 5567.

[37198] Jung, V.; Rashidbaigi, A.; Jones, C.; Tischfield, J. A.; Shows,

[37199] T. B.; Pestka, S.: Human chromosomes 6 and 21 are re-
quired for sensitivity

[37200] to human interferon gamma. Proc. Nat. Acad. Sci. 84:
4151-4155,

[37201] 1987.

[37202] 5568.

[37203] Langer, J. A.; Rashidbaigi, A.; Lai, L.-W.; Patterson, D.;
Jones,

[37204] C.: Sublocalization on chromosome 21 of human inter-
feron-alpha receptor

[37205] gene and the gene for an interferon-gamma response
protein. Somat.

[37206] Cell Molec. Genet. 16: 231-240, 1990.

[37207] 5569.

[37208] Mariano, T. M.; Muthukumaran, G.; Donnelly, R. J.; Wang,
N.; Adamson,

[37209] M. C.; Pestka, S.; Kozak, C. A.: Genetic mapping of the
gene for

[37210] the mouse interferon-gamma receptor signaling subunit to the distal

[37211] end of chromosome 16. Mammalian Genome 7: 321-322, 1996.

[37212] 5570.

[37213] Rhee, S.; Ebensperger, C.; Dembic, Z.; Pestka, S.: The structure

[37214] of the gene for the second chain of the human interferon-gamma receptor. J.

[37215] Biol. Chem. 271: 28947-28952, 1996.

[37216] 5571.

[37217] Soh, J.; Donnelly, R. J.; Mariano, T. M.; Cook, J. R.; Schwartz,

[37218] B.; Pestka, S.: Identification of a yeast artificial chromosome clone

[37219] encoding an accessory factor for the human interferon gamma receptor:

[37220] evidence for multiple accessory factors. Proc. Nat. Acad. Sci. 90:

[37221] 8737-8741, 1993.

[37222] 5572.

[37223] Badovinac, V. P.; Tvinnereim, A. R.; Harty, J. T.: Regulation

[37224] of antigen-specific CD8(+) T cell homeostasis by perforin and interferon-gamma. Science 290:

[37225] 1354-1357, 2000.

[37226] 5573.

[37227] Ben-Asouli, Y.; Banai, Y.; Pel-Or, Y.; Shir, A.; Kaempfer, R.:

[37228] Human interferon-gamma mRNA autoregulates its translation through

[37229] a pseudoknot that activates the interferon-inducible protein kinase

[37230] PKR. Cell 108: 221-232, 2002.

[37231] 5574.

[37232] Binder, G. K.; Griffin, D. E.: Interferon-gamma-mediated site-specific

[37233] clearance of alphavirus from CNS neurons. Science 293: 303-306,

[37234] 2001.

[37235] 5575.

[37236] Blalock, J. E.; Smith, E. M.: Human leukocyte interferon: structural

[37237] and biological relatedness to adrenocorticotropic hormone and endorphins. Proc.
[37238] Nat. Acad. Sci. 77: 5972-5974, 1980.

[37239] 5576.

[37240] Bureau, J. F.; Bihl, F.; Brahic, M.; Le Paslier, D.: The gene

[37241] coding for interferon-gamma is linked to the D12S335 and D12S313 microsatellites
[37242] and to the MDM2 gene. Genomics 28: 109-112, 1995.

[37243] 5577.

[37244] Burke, D. C.: The status of interferon. Sci. Am. 236(4): 42-50,
[37245] 1977.

[37246] 5578.

[37247] Cassingena, R.; Chany, C.; Vignal, M.; Suarex, H.; Estrade, S.;
[37248] Lazar, P.: Use of monkey-mouse hybrid cells for the study of the
[37249] cellular regulation of interferon production and action. Proc. Nat.
[37250] Acad. Sci. 68: 580-584, 1971.

[37251] 5579.

[37252] Creagan, R. P.; Tan, Y. H.; Chen, S.-H.; Ruddle, F. H.: So-
matic
[37253] cell genetic analysis of the interferon system. Fed. Proc.
34: 2222-2226,
[37254] 1975.

[37255] 5580.

[37256] Devos, R.; Cheroutre, H.; Taya, Y.; Degrave, W.; Van
Heuverswyn,
[37257] H.; Fiers, W.: Molecular cloning of human immune inter-
feron cDNA
[37258] and its expression in eukaryotic cells. Nucleic Acids Res.
10: 2487-2502,
[37259] 1982.

[37260] 5581.

[37261] Diaz, M. O.; Bohlander, S.; Allen, G.: Nomenclature of the
human
[37262] interferon genes. J. Interferon Res. 13: 443-444, 1993.

[37263] 5582.

[37264] Diefenbach, A.; Schindler, H.; Rollinghoff, M.; Yokoyama,
W. M.;

[37265] Bogdan, C.: Requirement for type 2 NO synthase for IL-12 signaling
[37266] in innate immunity. Science 284: 951-955, 1999.

[37267] 5583.

[37268] Gray, P. W.; Goeddel, D. V.: Structure of the human immune interferon
[37269] gene. Nature 298: 859-863, 1982.

[37270] 5584.

[37271] Isaacs, A.; Lindenmann, J.: Virus interference. I. The interferon. Proc.
[37272] Roy. Soc. London 147B: 258-267, 1957.

[37273] 5585.

[37274] Isaacs, A.; Lindenmann, J.; Valentine, R. C.: Virus interference.
[37275] II. Some properties of interferon. Proc. Roy. Soc. London 147B:
[37276] 268-273, 1957.

[37277] 5586.

[37278] Knight, E., Jr.: Human fibroblast interferon: amino acid analysis

[37279] and amino terminal amino acid sequence. Science 207: 525-526, 1980.

[37280] 5587.

[37281] Lipinski, M.; Virelizier, J. L.; Tursz, T.; Griscelli, C.: Natural

[37282] killer and killer cell activities in patients with primary immunodeficiencies

[37283] or defects in immune interferon production. Europ. J. Immun. 10:

[37284] 246-249, 1980.

[37285] 5588.

[37286] Maeda, S.; McCandliss, R.; Gross, M.; Sloma, A.; Familletti, P.

[37287] C.; Tabor, J. M.; Evinger, M.; Levy, W. P.; Pestka, S.: Construction

[37288] and identification of bacterial plasmids containing nucleotide sequence

[37289] for human leukocyte interferon. Proc. Nat. Acad. Sci. 77: 7010-7013,

[37290] 1980.

[37291] 5589.

[37292] Mantei, N.; Schwarzstein, M.; Streuli, M.; Panem, S.; Na-

gata,

[37293] S.; Weissmann, C.: The nucleotide sequence of a cloned human leukocyte

[37294] interferon cDNA. Gene 10: 1-10, 1980.

[37295] 5590.

[37296] Nagata, S.; Mantei, N.; Weissmann, C.: The structure of one of

[37297] the eight or more distinct chromosomal genes for human interferon-alpha. Nature 287:

[37298] 401-408, 1980.

[37299] 5591.

[37300] Nagata, S.; Taira, H.; Hall, A.; Johnstrud, L.; Streuli, M.; Escodi,

[37301] J.; Boll, W.; Cantell, K.; Weissmann, C.: Synthesis in E. coli of

[37302] a polypeptide with human leukocyte interferon activity. Nature 284:

[37303] 316-320, 1980.

[37304] 5592.

[37305] Nathan, C. F.; Murray, H. W.; Wiebe, M. E.; Rubin, B. Y.: Identification

[37306] of interferon-gamma as the lymphokine that activates human macrophage
[37307] oxidative metabolism and antimicrobial activity. J. Exp. Med. 158:
[37308] 670-689, 1983.

[37309] 5593.

[37310] Naylor, S. L.; Sakaguchi, A. Y.; Shows, T. B.; Law, M. L.; Goeddel,
[37311] D. V.; Gray, P. W.: Human immune interferon gene is located on chromosome
[37312] 12. J. Exp. Med. 57: 1020-1027, 1983.

[37313] 5594.

[37314] Shimizu, A.; Sakai, Y.; Ohno, K.; Masaki, S.; Kuwano, R.; Takahashi,
[37315] Y.; Miyashita, N.; Watanabe, T.: A molecular genetic linkage map
[37316] of mouse chromosome 10, including the Myb, S100b, Pah, Sl, and Ifg
[37317] genes. Biochem. Genet. 30: 529-535, 1992.

[37318] 5595.

[37319] Bader, B. L.; Jahn, L.; Franke, W. W.: Low level expression

of

[37320] cytokeratins 8, 18 and 19 in vascular smooth muscle cells of human

[37321] umbilical cord and in cultured cells derived therefrom, with an analysis

[37322] of the chromosomal locus containing the cytokeratin 19 gene. Europ.

[37323] J. Cell Biol. 47: 300-319, 1988.

[37324] 5596.

[37325] Bader, B. L.; Magin, T. M.; Hatzfield, M.; Franke, W. W.: Amino

[37326] acid sequence and gene organization of cytokeratin no. 19, an exceptional

[37327] tail-less intermediate filament protein. EMBO J. 5: 1865-1875, 1986.

[37328] 5597.

[37329] Blumenberg, M.: Concerted gene duplications in the two keratin

[37330] gene families. J. Molec. Evol. 27: 203-211, 1988.

[37331] 5598.

[37332] Ceratto, N.; Dobkin, C.; Carter, M.; Jenkins, E.; Yao, X.-L.;

Cassiman,

[37333] J.-J.; Aly, M. S.; Bosco, P.; Leube, R.; Langbein, L.; Feo, S.; Romano,

[37334] V.: Human type I cytokeratin genes are a compact cluster. Cytogenet.

[37335] Cell Genet. 77: 169-174, 1997.

[37336] 5599.

[37337] Eckert, R. L.: Sequence of the human 40-kDa keratin reveals an

[37338] unusual structure with very high sequence identity to the corresponding

[37339] bovine keratin. Proc. Nat. Acad. Sci. 85: 1114-1118, 1988.

[37340] 5600.

[37341] Lussier, M.; Filion, M.; Compton, J. G.; Nadeau, J. H.; Lapointe,

[37342] L.; Royal, A.: The mouse keratin 19-encoding gene: sequence, structure

[37343] and chromosomal assignment. Gene 95: 203-213, 1990.

[37344] 5601.

[37345] Savtchenko, E. S.; Schiff, T. A.; Jiang, C.-K.; Freedberg, I.

M.;

[37346] Blumenberg, M.: Embryonic expression of the human 40-kD keratin:

[37347] evidence from a processed pseudogene sequence. Am. J. Hum. Genet. 43:

[37348] 630-637, 1988.

[37349] 5602.

[37350] Stasiak, P. C.; Lane, E. B.: Sequence of cDNA coding for human

[37351] keratin #19. Nucleic Acids Res. 15: 10058 only, 1987.

[37352] 5603.

[37353] Blumenberg, M.; Savtchenko, E. S.: Linkage of human keratin genes. Cytogenet.

[37354] Cell Genet. 42: 65-71, 1986.

[37355] 5604.

[37356] Fuchs, E.; Coppock, S.; Green, H.; Cleveland, D.: Two distinct

[37357] classes of keratin genes and their evolutionary significance. Cell 27:

[37358] 75-84, 1981.

[37359] 5605.

[37360] Fuchs, E.; Green, H.: Changes in keratin gene expression during
[37361] terminal differentiation of the keratinocyte. Cell 19: 1033-1042,
[37362] 1980.

[37363] 5606.

[37364] Hanukoglu, I.; Fuchs, E.: The cDNA sequence of a human epidermal
[37365] keratin: divergence of sequence but conservation of structure among
[37366] intermediate filament proteins. Cell 31: 243-252, 1982.

[37367] 5607.

[37368] Lee, L. D.; Baden, H. P.: Organisation of the polypeptide chains
[37369] in mammalian keratin. Nature 264: 377-379, 1976.

[37370] 5608.

[37371] Marchuk, D.; McCrohon, S.; Fuchs, E.: Complete sequence of a gene
[37372] encoding a human type I keratin: sequences homologous to enhancer
[37373] elements in the regulatory region of the gene. Proc. Nat.

Acad. Sci. 82:
[37374] 1609-1613, 1985.

[37375] 5609.

[37376] Nadeau, J. H.; Berger, F. G.; Cox, D. R.; Crosby, J. L.; Davisson,
[37377] M. T.; Ferrara, D.; Fuchs, E.; Hart, C.; Hunihan, L.; Lalley, P. A.;
[37378] Langley, S. H.; Martin, G. R.; Nichols, L.; Phillips, S. J.; Rod-
erick,
[37379] T. H.; Roop, D. R.; Ruddle, F. H.; Skow, L. C.; Compton, J. G.: A
[37380] family of type I keratin genes and the homeobox-2 gene complex are
[37381] closely linked to the rex locus on mouse chromosome 11. Genomics 5:
[37382] 454-462, 1989.

[37383] 5610.

[37384] RayChaudhury, A.; Marchuk, D.; Lindhurst, M.; Fuchs, E.: Three
[37385] tightly linked genes encoding human type I keratins: con-
servation
[37386] of sequence in the 5-prime-untranslated leader and

5-prime-upstream
[37387] regions of coexpressed keratin genes. Molec. Cell. Biol. 6: 539-548,
[37388] 1986.

[37389] 5611.

[37390] Romano, V.; Bosco, P.; Raimondi, E.; Feo, S.; Leube, R.; Franke,
[37391] W.; Ceratto, N.: Chromosomal mapping and physical linkage analysis
[37392] of human acidic cytokeratin genes. (Abstract) Cytogenet. Cell Genet. 58:
[37393] 2009-2010, 1991.

[37394] 5612.

[37395] Heim, S.; Nilbert, M.; Vanni, R.; Floderus, U.-M.; Mandahl, N.;
[37396] Liedgren, S.; Lecca, U.; Mitelman, F.: A specific translocation,
[37397] t(12;14)(q14-15;q23-24), characterizes a subgroup of uterine leiomyomas. Cancer
[37398] Genet. Cytogenet. 32: 13-17, 1988.

[37399] 5613.

[37400] Minucci, S.; Maccarana, M.; Cioce, M.; De Luca, P.; Gel-
metti,
[37401] V.; Segalla, S.; Di Croce, L.; Giavara, S.; Matteucci, C.;
Gobbi,
[37402] A.; Bianchini, A.; Colombo, E.; Schiavoni, I.; Badaracco, G.;
Hu,
[37403] X.; Lazar, M. A.; Landsberger, N.; Nervi, C.; Pelicci, P. G.:
Oligomerization
[37404] of RAR and AML1 transcription factors as a novel mecha-
nism of oncogenic
[37405] activation. Molec. Cell 5: 811-820, 2000.

[37406] 5614.

[37407] Nucifora, G.; Begy, C. R.; Kobayashi, H.; Roulston, D.;
Claxton,
[37408] D.; Pedersen-Bjergaard, J.; Parganas, E.; Ihle, J. N.; Rowley,
J.
[37409] D.: Consistent intergenic splicing and production of mul-
tiple transcripts
[37410] between AML1 at 21q22 and unrelated genes at 3q26 in
(3;21)(q26;q22)
[37411] translocations. Proc. Nat. Acad. Sci. 91: 4004-4008, 1994.

[37412] 5615.

[37413] Nucifora, G.; Rowley, J. D.: AML1 and the 8;21 and 3;21 translocations
[37414] in acute and chronic myeloid leukemia. Blood 86: 1-14, 1995.
[37415] 5616.

[37416] Okuda, T.; Takeda, K.; Fujita, Y.; Nishimura, M.; Yagyu, S.; Yoshida,
[37417] M.; Akira, S.; Downing, J. R.; Abe, T.: Biological characteristics
[37418] of the leukemia-associated transcriptional factor AML1 disclosed by
[37419] hematopoietic rescue of AML1-deficient embryonic stem cells by using
[37420] a knock-in strategy. Molec. Cell. Biol. 20: 319-328, 2000.

[37421] 5617.

[37422] Okuda, T.; van Deursen, J.; Hiebert, S. W.; Grosveld, G.; Downing,
[37423] J. R.: AML1, the target of multiple chromosomal translocations in
[37424] human leukemia, is essential for normal fetal liver hematopoiesis. Cell 84:
[37425] 321-330, 1996.

[37426] 5618.

[37427] Osato, M.; Asou, N.; Abdalla, E.; Hoshino, K.; Yamasaki, H.; Okubo,

[37428] T.; Suzushima, H.; Takatsuki, K.; Kanno, T.; Shigesada, K.; Ito, Y.

[37429] : Biallelic and heterozygous point mutations in the runt domain of

[37430] the AML1/PEBP2-alpha-B gene associated with myeloblastic leukemias. Blood 93:

[37431] 1817-1824, 1999.

[37432] 5619.

[37433] Pui, C.-H.: Childhood leukemias. New Eng. J. Med. 332: 1618-1630,

[37434] 1995.

[37435] 5620.

[37436] Rowley, J. D.: Recurring chromosome abnormalities in leukemia

[37437] and lymphoma. Seminars Hemat. 27: 122-136, 1990.

[37438] 5621.

[37439] Song, W.-J.; Sullivan, M. G.; Legare, R. D.; Hutchings, S.; Tan,

[37440] X.; Kufrin, D.; Ratajczak, J.; Resende, I. C.; Haworth, C.; Hock,
[37441] R.; Loh, M.; Felix, C.; and 12 others: Haploinsufficiency of CBFA2
[37442] causes familial thrombocytopenia with propensity to develop acute
[37443] myelogenous leukaemia. Nature Genet. 23: 166-175, 1999.
[37444] 5622.

[37445] Walker, L. C.; Stevens, J.; Campbell, H.; Corbett, R.; Spearing,
[37446] R.; Heaton, D.; Macdonald, D. H.; Morris, C. M.; Ganly, P.: A novel
[37447] inherited mutation of the transcription factor RUNX1 causes thrombocytopenia
[37448] and may predispose to acute myeloid leukaemia. Brit. J. Haemat. 117:
[37449] 878-881, 2002.

[37450] 5623.

[37451] Wang, Q.; Stacy, T.; Binder, M.; Marin-Padilla, M.; Sharpe, A.
[37452] H.; Speck, N. A.: Disruption of the Cbfa2 gene causes

necrosis and
[37453] hemorrhaging in the central nervous system and blocks definitive hematopoiesis. Proc.
[37454] Nat. Acad. Sci. 93: 3444-3449, 1996.

[37455] 5624.

[37456] Zipursky, A.; Poon, A.; Doyle, J.: Leukemia in Down syndrome:
[37457] a review. Pediat. Hemat. Oncol. 9: 139-149, 1992.

[37458] 5625.

[37459] Erikson, J.; Williams, D. L.; Finan, J.; Nowell, P. C.; Croce,
[37460] C. M.: Locus of the alpha-chain of the T-cell receptor is split by
[37461] chromosome translocation in T-cell leukemias. Science 229: 784-786,
[37462] 1985.

[37463] 5626.

[37464] Das, R.; Reddy, E. P.; Chatterjee, D.; Andrews, D. W.: Identification
[37465] of a novel Bcl-2 related gene, BRAG-1, in human glioma. Oncogene 12:
[37466] 947-951, 1996. Note: Erratum: Oncogene 16: 429 only,

1998.

[37467] 5627.

[37468] Deng, G.; Podack, E. R.: Suppression of apoptosis in a cy-
totoxic

[37469] T-cell line by interleukin 2-mediated gene transcription
and deregulated

[37470] expression of the protooncogene bcl-2. Proc. Nat. Acad.
Sci. 90:

[37471] 2189-2193, 1993.

[37472] 5628.

[37473] Barker, D.; Wright, E.; Nguyen, K.; Cannon, L.; Fain, P.;
Goldgar,

[37474] D.; Bishop, D. T.; Carey, J.; Baty, B.; Kivlin, J.; Willard, H.;
Waye,

[37475] J. S.; Greig, G.; Leinwand, L.; Nakamura, Y.; O'Connell, P.;
Leppert,

[37476] M.; Lalouel, J.-M.; White, R.; Skolnick, M.: Gene for von
Recklinghausen

[37477] neurofibromatosis is in the pericentromeric region of
chromosome 17. Science 236:

[37478] 1100-1102, 1987.

[37479] 5629.

[37480] Costa, R. M.; Yang, T.; Huynh, D. P.; Pulst, S. M.; Viskochil,

[37481] D. H.; Silva, A. J.; Brannan, C. I.: Learning deficits, but nor-
mal

[37482] development and tumor predisposition, in mice lacking
exon 23a of

[37483] Nf1. Nature Genet. 27: 399-405, 2001.

[37484] 5630.

[37485] Tan, M.; Jing, T.; Lan, K.-H.; Neal, C. L.; Li, P.; Lee, S.;
Fang,

[37486] D.; Nagata, Y.; Liu, J.; Arlinghaus, R.; Hung, M.-C.; Yu, D.:
Phosphorylation

[37487] on tyrosine-15 of p34(Cdc2) by ErbB2 inhibits p34(Cdc2)
activation

[37488] and is involved in resistance to taxol-induced apoptosis.
Molec.

[37489] Cell 9: 993-1004, 2002.

[37490] 5631.

[37491] Gromeier, M.; Solecki, D.; Patel, D. D.; Wimmer, E.: Ex-
pression

[37492] of the human poliovirus receptor/CD155 gene during de-
velopment of

[37493] the central nervous system: implications for the pathogenesis of poliomyelitis. Virology 273:
[37494] 248-257, 2000.

[37495] 5632.

[37496] He, Y.; Bowman, V. D.; Mueller, S.; Bator, C. M.; Bella, J.; Peng,
[37497] X.; Baker, T. S.; Wimmer, E.; Kuhn, R. J.; Rossmann, M. G.: Interaction
[37498] of the poliovirus receptor with poliovirus. Proc. Nat. Acad. Sci. 97:
[37499] 79-84, 2000.

[37500] 5633.

[37501] Koike, S.; Horie, H.; Ise, I.; Okitsu, A.; Yoshida, M.; Iizuka,
[37502] N.; Takeuchi, K.; Takegami, T.; Nomoto, A.: The poliovirus receptor
[37503] protein is produced both as membrane-bound and secreted forms. EMBO
[37504] J. 9: 3217-3224, 1990.

[37505] 5634.

[37506] Koike, S.; Taya, C.; Kurata, T.; Abe, S.; Ise, I.; Yonekawa, H.;

[37507] Nomoto, A.: Transgenic mice susceptible to poliovirus. Proc. Nat.

[37508] Acad. Sci. 88: 951-955, 1991.

[37509] 5635.

[37510] Mendelsohn, C.; Johnson, B.; Lionetti, K. A.; Nobis, P.; Wimmer,

[37511] E.; Racaniello, V. R.: Transformation of a human poliovirus receptor

[37512] gene into mouse cells. Proc. Nat. Acad. Sci. 83: 7845-7849, 1986.

[37513] 5636.

[37514] Mendelsohn, C. L.; Wimmer, E.; Racaniello, V. R.: Cellular receptor

[37515] for poliovirus: molecular cloning, nucleotide sequence, and expression

[37516] of a new member of the immunoglobulin superfamily. Cell 56: 855-865,

[37517] 1989.

[37518] 5637.

[37519] Miller, D. A.; Miller, O. J.; Dev, V. G.; Hashmi, S.; Tantravahi,

[37520] R. R.; Medrano, L.; Green, H.: Human chromosome 19 carries a poliovirus
[37521] receptor gene. Cell 1: 167-174, 1974.

[37522] 5638.

[37523] Larrea, F.; Escorza, A.; Granados, J.; Valencia, X.; Valero, A.;
[37524] Cravioto, M. C.; Perez-Palacios, G.: Familial occurrence of big-big
[37525] prolactin as the predominant immunoreactive human prolactin species
[37526] in blood. Fertil. Steril. 47: 956-963, 1987.

[37527] 5639.

[37528] Michelson, A. M.; Bruns, G. A. P.; Morton, C. C.; Orkin, S. H.
[37529] : The human phosphoglycerate kinase multigene family: HLA-associated
[37530] sequences and an X-linked locus containing a processed pseudogene
[37531] and its functional counterpart. J. Biol. Chem. 260: 6982-6992, 1985.
[37532] 5640.

[37533] Singer-Sam, J.; Simmer, R. L.; Keith, D. H.; Shively, L.; Teplitz,
[37534] M.; Itakura, K.; Gartler, S. M.; Riggs, A. D.: Isolation of a cDNA
[37535] clone for human X-linked 3-phosphoglycerate kinase by use of a mixture
[37536] of synthetic oligodeoxyribonucleotides as a detection probe. Proc.
[37537] Nat. Acad. Sci. 80: 802-806, 1983.

[37538] 5641.

[37539] Willard, H. F.; Goss, S. J.; Holmes, M. T.; Munroe, D. L.: Re-gional
[37540] localization of the phosphoglycerate kinase gene and pseudogene on
[37541] the human X chromosome and assignment of a related DNA sequence to
[37542] chromosome 19. Hum. Genet. 71: 138-143, 1985.

[37543] 5642.

[37544] Killian, J. M.; Kloepfer, H. W.: Homozygous expression of a dominant
[37545] gene for Charcot-Marie-Tooth neuropathy. Ann. Neurol. 5: 515-522,

[37546] 1979.

[37547] 5643.

[37548] Lupski, J. R.; Montes de Oca-Luna, R.; Slaugenhaupt, S.; Pentao,

[37549] L.; Guzzetta, V.; Trask, B. J.; Saucedo-Cardenas, O.; Barker, D. F.;

[37550] Killian, J. M.; Garcia, C. A.; Chakravarti, A.; Patel, P. I.: DNA

[37551] duplication associated with Charcot-Marie-Tooth disease type 1A. Cell 66:

[37552] 219-232, 1991.

[37553] 5644.

[37554] Nelis, E.; Haites, N.; Van Broeckhoven, C.: Mutations in the

[37555] peripheral myelin genes and associated genes in inherited peripheral

[37556] neuropathies. Hum. Mutat. 13: 11-28, 1999.

[37557] 5645.

[37558] Ouvrier, R. A.; McLeod, J. G.; Conchin, T. E.: The hyper-trophic

[37559] forms of hereditary motor and sensory neuropathy: a study of hypertrophic

[37560] Charcot-Marie-Tooth disease (HMSN type I) and Dejerine-Sottas disease
[37561] (HMSN type III) in childhood. Brain 110: 121-148, 1987.
[37562] 5646.
[37563] Spicer, A. P.; Seldin, M. F.; Olsen, A. S.; Brown, N.; Wells, D.
[37564] E.; Doggett, N. A.; Itano, N.; Kimata, K.; Inazawa, J.; Mc-Donald,
[37565] J. A.: Chromosomal localization of the human and mouse hyaluronan
[37566] synthase genes. Genomics 41: 493-497, 1997.
[37567] 5647.
[37568] Watanabe, K.; Yamaguchi, Y.: Molecular identification of a putative
[37569] human hyaluronan synthase. J. Biol. Chem. 271: 22945-22948, 1996.
[37570] 5648.
[37571] Cahill, D. P.; da Costa, L. T.; Carson-Walter, E. B.; Kinzler,
[37572] K. W.; Vogelstein, B.; Lengauer, C.: Characterization of MAD2B and
[37573] other mitotic spindle checkpoint genes. Genomics 58:

181-187, 1999.

[37574] 5649.

[37575] Luo, X.; Tang, Z.; Rizo, J.; Yu, H.: The Mad2 spindle check-
point

[37576] protein undergoes similar major conformational changes
upon binding

[37577] to either Mad1 or Cdc20. Molec. Cell 9: 59-71, 2002.

[37578] 5650.

[37579] Nelson, K. K.; Schlondorff, J.; Blobel, C. P.: Evidence for an

[37580] interaction of the metalloprotease-disintegrin tumour
necrosis factor

[37581] alpha convertase (TACE) with mitotic arrest deficient 2
(MAD2), and

[37582] of the metalloprotease-disintegrin MDC9 with a novel
MAD2-related

[37583] protein, MAD2-beta. Biochem. J. 343: 673-680, 1999.

[37584] 5651.

[37585] Liu, Y.; Chiu, I.-M.: Assignment of FGF12, the human FGF
homologous

[37586] factor 1 gene, to chromosome 3q29-3qter by fluores-
cence in situ hybridization. Cytogenet.

[37587] Cell Genet. 78: 48-49, 1997.

[37588] 5652.

[37589] Caslini, C.; Spinelli, O.; Cazzaniga, G.; Golay, J.; De Gioia,

[37590] L.; Pedretti, A.; Breviario, F.; Amaru, R.; Barbui, T.; Biondi, A.;

[37591] Introna, M.; Rambaldi, A.: Identification of two novel iso-forms of

[37592] the ZNF162 gene: a growing family of signal transduction and activator

[37593] of RNA proteins. Genomics 42: 268-277, 1997.

[37594] 5653.

[37595] Kramer, A.; Quentin, M.; Mulhauser, F.: Diverse modes of alternative

[37596] splicing of human splicing factor SF1 deduced from the exon-intron

[37597] structure of the gene. Gene 211: 29-37, 1998.

[37598] 5654.

[37599] Gong, Y.; Chitayat, D.; Kerr, B.; Chen, T.; Babul-Hirji, R.; Pal,

[37600] A.; Reiss, M.; Warman, M. L.: Brachydactyly type B: clinical description,

[37601] genetic mapping to chromosome 9q, and evidence for a shared ancestral
[37602] mutation. Am. J. Hum. Genet. 64: 578-585, 1999.

[37603] 5655.

[37604] Innis, J. W.; Mortlock, D. P.: Limb development: molecular dysmorphology
[37605] is at hand! Clin. Genet. 53: 337-348, 1998.

[37606] 5656.

[37607] Manouvrier-Hanu, S.; Holder-Espinasse, M.; Lyonnet, S.: Genetics
[37608] of limb anomalies in humans. Trends Genet. 15: 409-417, 1999.
[37609] 5657.

[37610] Oishi, I.; Takeuchi, S.; Hashimoto, R.; Nagabukuro, A.; Ueda, T.;
[37611] Liu, Z.-J.; Hatta, T.; Akira, S.; Matsuda, Y.; Yamamura, H.; Otani,
[37612] H.; Minami, Y.: Spatio-temporally regulated expression of receptor
[37613] tyrosine kinases, mRor1, mRor2, during mouse develop- ment: implications

[37614] in development and function of the nervous system.
Genes Cells 4:
[37615] 41-56, 1999.

[37616] 5658.

[37617] Schwabe, G. C.; Tinschert, S.; Buschow, C.; Meinecke, P.;
Wolff,
[37618] G.; Gillessen-Kaesbach, G.; Oldridge, M.; Wilkie, A. O. M.;
Komec,
[37619] R.; Mundlos, S.: Distinct mutations in the receptor tyrosine
kinase
[37620] gene ROR2 cause brachydactyly type B. Am. J. Hum.
Genet. 67: 822-831,
[37621] 2000.

[37622] 5659.

[37623] Takeuchi, S.; Takeda, K.; Oishi, I.; Nomi, M.; Ikeya, M.;
Itoh,
[37624] K.; Tamura, S.; Ueda, T.; Hatta, T.; Otani, H.; Terashima,
T.; Takada,
[37625] S.; Yamamura, H.; Akira, S.; Minami, Y.: Mouse Ror2 re-
ceptor tyrosine
[37626] kinase is required for the heart development and limb for-
mation. Genes

[37627] Cells 5: 71-78, 2000.

[37628] 5660.

[37629] van Bokhoven, H.; Brunner, H. G.: Splitting p63. Am. J. Hum.

[37630] Genet. 71: 1-13, 2002.

[37631] 5661.

[37632] Jaffrey, S. R.; Snyder, S. H.: PIN: an associated protein inhibitor

[37633] of neuronal nitric oxide synthase. Science 274: 774-777, 1996.

[37634] 5662.

[37635] Garkavtsev, I.; Demetrick, D.; Riabowol, K.: Cellular localization

[37636] and chromosome mapping of a novel candidate tumor suppressor gene

[37637] (ING1). Cytogenet. Cell Genet. 76: 176-178, 1997.

[37638] 5663.

[37639] Garkavtsev, I.; Kazarov, A.; Gudkov, A.; Riabowol, K.: Suppression

[37640] of the novel growth inhibitor p33(ING1) promotes neoplastic transformation. Nature

[37641] Genet. 14: 415-420, 1996. Note: Erratum: Nature Genet. 23: 373 only,
[37642] 1999.

[37643] 5664.

[37644] Saito, A.; Furukawa, T.; Fukushige, S.; Koyama, S.; Hoshi, M.;
[37645] Hayashi, Y.; Horii, A.: p24/ING1-ALT1 and p47/ING1-ALT2, distinct
[37646] alternative transcripts of p33/ING1. J. Hum. Genet. 45: 177-181,
[37647] 2000.

[37648] 5665.

[37649] Zeremski, M.; Horrigan, S. K.; Grigorian, I. A.; Westbrook, C.
[37650] A.; Gudkov, A. V.: Localization of the candidate tumor suppressor
[37651] gene ING1 to human chromosome 13q34. Somat. Cell Molec. Genet. 23:
[37652] 233-236, 1997.

[37653] 5666.

[37654] Gilligan, D. M.; Lozovatsky, L.; Gwynn, B.; Brugnara, C.;

Mohandas,
[37655] N.; Peters, L. L.: Targeted disruption of the beta adducin gene (Add2)
[37656] causes red blood cell spherocytosis in mice. Proc. Nat. Acad. Sci. 96:
[37657] 10717-10722, 1999.

[37658] 5667.

[37659] Katagiri, T.; Ozaki, K.; Fujiwara, T.; Shimizu, F.; Kawai, A.;

[37660] Okuno, S.; Suzuki, M.; Nakamura, Y.; Takahashi, E.; Hirai, Y.: Cloning,
[37661] expression and chromosome mapping of adducin-like 70 (ADDL), a human
[37662] cDNA highly homologous to human erythrocyte adducin. Cytogenet. Cell
[37663] Genet. 74: 90-95, 1996.

[37664] 5668.

[37665] Adamson, M. C.; Dennis, C.; Delaney, S.; Christiansen, J.; Monkley,
[37666] S.; Kozak, C. A.; Wainwright, B.: Isolation and genetic mapping of
[37667] two novel members of the murine Wnt gene family, Wnt11

and Wnt12,

[37668] and the mapping of Wnt5a and Wnt7a. Genomics 24: 9-13, 1994.

[37669] 5669.

[37670] Bui, T. D.; Lako, M.; Lejeune, S.; Curtis, A. R. J.; Strachan,

[37671] T.; Lindsay, S.; Harris, A. L.: Isolation of a full-length hu-man

[37672] WNT7A gene implicated in limb development and cell transformation,

[37673] and mapping to chromosome 3p25. Gene 189: 25-29, 1997.

[37674] 5670.

[37675] Ikegawa, S.; Kumano, Y.; Okui, K.; Fujiwara, T.; Takahashi, E.;

[37676] Nakamura, Y.: Isolation, characterization and chromoso-mal assignment

[37677] of the human WNT7A gene. Cytogenet. Cell Genet. 74: 149-152, 1996.

[37678] 5671.

[37679] Li, S.; Chiang, T.-C.; Davis, G. R.; Williams, R. M.; Wilson, V.

[37680] P.; McLachlan, J. A.: Decreased expression of Wnt7a mRNA

is inversely
[37681] associated with the expression of estrogen receptor-alpha in human
[37682] uterine leiomyoma. J. Clin. Endocr. Metab. 86: 454-457, 2001.
[37683] 5672.

[37684] Walder, K.; Norman, R. A.; Hanson, R. L.; Schrauwen, P.; Neverova,
[37685] M.; Jenkinson, C. P.; Easlick, J.; Warden, C. H.; Pecqueur, C.; Raimbault,
[37686] S.; Ricquier, D.; Harper, M.; Silver, K.; Shuldiner, A. R.; Solanes,
[37687] G.; Lowell, B. B.; Chung, W. K.; Leibel, R. L.; Pratley, R.; Ravussin,
[37688] E.: Association between uncoupling protein polymor-phisms (UCP2-UCP3)
[37689] and energy metabolism/obesity in Pima Indians. Hum. Molec. Genet. 7:
[37690] 1431-1435, 1998.

[37691] 5673.

[37692] Lindsay, E. A.; Rizzu, P.; Antonacci, R.; Jurecic, V.; Delmas-Mata,

[37693] J.; Lee, C.-C.; Kim, U.-J.; Scambler, P. J.; Baldini, A.: A transcription
[37694] map in the CATCH22 critical region: identification, mapping, and ordering
[37695] of four novel transcripts expressed in heart. Genomics 32: 104-112,
[37696] 1996.

[37697] 5674.

[37698] Takai, S.; Hinoda, Y.; Adachi, T.; Imai, K.; Oshima, M.: A human
[37699] UPD (sic)-GalNAc: polypeptide, N-acetylgalactosaminyltransferase type
[37700] 1 gene is located at the chromosomal region 18q12.1. Hum. Genet. 99:
[37701] 293-294, 1997.

[37702] 5675.

[37703] White, T.; Bennett, E. P.; Takio, K.; Sorensen, T.; Bonding, N.;
[37704] Clausen, H.: Purification and cDNA cloning of a human UDP-N-acetyl-alpha-D-galactosamine:polypeptide
[37705] N-acetylgalactosaminyltransferase. J. Biol. Chem. 270: 24156-24165,

[37706] 1995.

[37707] 5676.

[37708] Candau, R.; Moore, P. A.; Wang, L.; Barlev, N.; Ying, C. Y.; Rosen,

[37709] C. A.; Berger, S. L.: Identification of human proteins functionally

[37710] conserved with the yeast putative adaptors ADA2 and GCN5. Molec.

[37711] Cell. Biol. 16: 593-602, 1996.

[37712] 5677.

[37713] Carter, K. C.; Wang, L.; Shell, B. K.; Zamir, I.; Berger, S. L.;

[37714] Moore, P. A.: The human transcriptional adaptor genes TADA2L and

[37715] GCN5L2 colocalize to chromosome 17q12-q21 and display a similar tissue

[37716] expression pattern. Genomics 40: 497-500, 1997.

[37717] 5678.

[37718] Ogryzko, V. V.; Kotani, T.; Zhang, X.; Schiltz, R. L.; Howard,

[37719] T.; Yang, X.-J.; Howard, B. H.; Qin, J.; Nakatani, Y.: His-
tone-like
[37720] TAFs within the PCAF histone acetylase complex. Cell 94:
35-44,
[37721] 1998.

[37722] 5679.

[37723] Struhl, K.; Moqtaderi, Z.: The TAFs in the HAT. Cell 94:
1-4,
[37724] 1998.

[37725] 5680.

[37726] Goldwurm, S.; Menzies, M. L.; Banyer, J. L.; Powell, B. L.
W.;
[37727] Jazwinska, E. C.: Identification of a novel Krueppel-related
zinc
[37728] finger gene (ZNF184) mapping to 6p21.3. Genomics 40:
486-489, 1997.
[37729] 5681.

[37730] Imhof, M. O.; McDonnell, D. P.: Yeast RSP5 and its human
homolog
[37731] hRPF1 potentiate hormone-dependent activation of tran-
scription by

[37732] human progesterone and glucocorticoid receptors. Molec. Cell. Biol. 16:

[37733] 2594-2605, 1996.

[37734] 5682.

[37735] Kumar, S.; Harvey, K. F.; Kinoshita, M.; Copeland, N. G.; Noda,

[37736] M.; Jenkins, N. A.: cDNA cloning, expression analysis, and mapping

[37737] of the mouse Nedd4 gene. Genomics 40: 435-443, 1997.

[37738] 5683.

[37739] Dikstein, R.; Zhou, S.; Tjian, R.: Human TAF(II)105 is a cell

[37740] type-specific TFIID subunit related to hTAF(II)130. Cell 87: 137-146,

[37741] 1996.

[37742] 5684.

[37743] Freiman, R. N.; Albright, S. R.; Zheng, S.; Sha, W. C.; Hammer,

[37744] R. E.; Tjian, R.: Requirement of tissue-selective TBP-associated

[37745] factor TAFII105 in ovarian development. Science 293: 2084-2087,

[37746] 2001.

[37747] 5685.

[37748] Allikmets, R.: Simple and complex ABCR: genetic predis-
position
[37749] to retinal disease. (Editorial) Am. J. Hum. Genet. 67:
793-799,
[37750] 2000.

[37751] 5686.

[37752] Allikmets, R.; Gerrard, B.; Hutchinson, A.; Dean, M.: Char-
acterization
[37753] of the human ABC superfamily: isolation and mapping of
21 new genes
[37754] using the expressed sequence tags database. Hum. Molec.
Genet. 5:
[37755] 1649-1655, 1996.

[37756] 5687.

[37757] Allikmets, R.; Singh, N.; Sun, H.; Shroyer, N. F.; Hutchin-
son,
[37758] A.; Chidambaram, A.; Gerrard, B.; Baird, L.; Stauffer, D.;
Peiffer,
[37759] A.; Rattner, A.; Smallwood, P.; Li, Y.; Anderson, K. L.;

Lewis, R.

[37760] A.; Nathans, J.; Leppert, M.; Dean, M.; Lupski, J. R.: A photoreceptor

[37761] cell-specific ATP-binding transporter gene (ABCR) is mutated in recessive

[37762] Stargardt macular dystrophy. Nature Genet. 15: 236-246, 1997. Note:

[37763] Erratum: Nature Genet. 17:122 only, 1997.

[37764] 5688.

[37765] Allikmets, R.; the International ABCR Screening Consortium:

[37766] Further evidence for an association of ABCR alleles with age-related

[37767] macular degeneration. Am. J. Hum. Genet. 67: 487-491, 2000.

[37768] 5689.

[37769] Allikmets, R.; Wasserman, W. W.; Hutchinson, A.; Smallwood, P.;

[37770] Nathans, J.; Rogan, P. K.; Schneider, T. D.; Dean, M.: Organization

[37771] of the ABCR gene: analysis of promoter and splice junction sequences. Gene 215:

[37772] 111-122, 1998.

[37773] 5690.

[37774] Azarian, S. M.; Travis, G. H.: The photoreceptor rim pro-
tein is

[37775] an ABC transporter encoded by the gene for recessive
Stargardt's disease

[37776] (ABCR). FEBS Lett. 409: 247-252, 1997.

[37777] 5691.

[37778] Birnbach, C. D.; Jarvelainen, M.; Possin, D. E.; Milam, A. H.:

[37779] Histopathology and immunocytochemistry of the neu-
rosensory retina

[37780] in fundus flavimaculatus. Ophthalmology 101:
1211-1219, 1994.

[37781] 5692.

[37782] Fumagalli, A.; Ferrari, M.; Soriani, N.; Gessi, A.; Foglieni,

[37783] B.; Martina, E.; Manitto, M. P.; Brancato, R.; Dean, M.; Al-
likmets,

[37784] R.; Cremonesi, L.: Mutational scanning of the ABCR gene
with double-gradient

[37785] denaturing-gradient gel electrophoresis (DG-DGGE) in
Italian Stargardt

[37786] disease patients. Hum. Genet. 109: 326-338, 2001.

[37787] 5693.

[37788] Gerber, S.; Rozet, J. M.; van de Pol, T. J. R.; Hoyng, C. B.;

[37789] Munnich, A.; Blankenagel, A.; Kaplan, J.; Cremers, F. P. M.: Complete

[37790] exon-intron structure of the retina-specific ATP binding transporter

[37791] gene (ABCR) allows the identification of novel mutations underlying

[37792] Stargardt disease. Genomics 48: 139-142, 1998.

[37793] 5694.

[37794] Rizzu, P.; Lindsay, E. A.; Taylor, C.; O'Donnell, H.; Levy, A.;

[37795] Scambler, P.; Baldini, A.: Cloning and comparative mapping of a gene

[37796] from the commonly deleted region of DiGeorge and velocardiofacial

[37797] syndromes conserved in C. elegans. Mammalian Genome 7: 639-643,

[37798] 1996.

[37799] 5695.

[37800] Cremers, F. P. M.; van de Pol, D. J. R.; van Driel, M.; den Hollander,

[37801] A. I.; van Haren, F. J. J.; Knoers, N. V. A. M.; Tijmes, N.; Bergen,

[37802] A. A. B.; Rohrschneider, K.; Blankenagel, A.; Pinckers, A. J. L. G.;

[37803] Deutman, A. F.; Hoyng, C. B.: Autosomal recessive retinitis pigmentosa

[37804] and cone-rod dystrophy caused by splice site mutations in the Stargardt's

[37805] disease gene ABCR. Hum. Molec. Genet. 7: 355-362, 1998.

[37806] 5696.

[37807] Bennett, E. P.; Hassan, H,.; Clausen, H.: cDNA cloning and expression

[37808] of a novel human UDP-N-acetyl-alpha-D-galactosamine. J. Biol. Chem. 271:

[37809] 17006-17012, 1996.

[37810] 5697.

[37811] Bennett, E. P.; Weghuis, D. O.; Merkx, G.; Geurts van Kessel, A.;

[37812] Eiberg, H.; Clausen, H.: Genomic organization and chro-

mosomal localization
[37813] of three members of the UDP-
N-acetylgalactosamine:polypeptide N-
acetylgalactosaminyltransferase
[37814] family. Glycobiology 8: 547-555, 1998.

[37815] 5698.

[37816] Zara, J.; Hagen, F. K.; Ten Hagen, K. G.; Van Wuyckhuyse,
B. C.;
[37817] Tabak, L. A.: Cloning and expression of mouse UDP-
GalNAc:polypeptide
[37818] N-acetylgalactosaminyltransferase-T3. Biochem. Biophys.
Res. Commun. 228:
[37819] 38-44, 1996.

[37820] 5699.

[37821] Zhao, Y.-Y.; Liu, Y.; Stan, R.-V.; Fan, L.; Gu, Y.; Dalton, N.;

[37822] Chu, P.-H.; Peterson, K.; Ross, J., Jr.; Chien, K. R.: Defects
in
[37823] caveolin-1 cause dilated cardiomyopathy and pulmonary
hypertension
[37824] in knockout mice. Proc. Nat. Acad. Sci. 99: 11375-11380,
2002.

[37825] 5700.

[37826] Mok, S. C.; Wong, K.-K.; Chan, R. K. W.; Lau, C. C.; Tsao, S.-W.;

[37827] Knapp, R. C.; Berkowitz, R. S.: Molecular cloning of differ-

[37828] entially expressed genes in human epithelial ovarian cancer. Gy-

[37829] necol. Oncol. 52: 247-252, 1994.

[37830] 5701.

[37831] Xu, X.-X.; Yang, W.; Jackowski, S.; Rock, C. O.: Cloning of a

[37832] novel phosphoprotein regulated by colony-stimulating factor 1 shares

[37833] a domain with the Drosophila disabled gene product. J.

[37834] Biol. Chem. 270: 14184-14191, 1995.

[37835] 5702.

[37836] Campbell, H. D.; Webb, G. C.; Fountain, S.; Young, I. G.: The

[37837] human PIN1 peptidyl-prolyl cis/trans isomerase gene maps to human

[37838] chromosome 19p13 and the closely related PIN1L gene to 1p31. Genomics 44:

[37839] 157-162, 1997.

[37840] 5703.

[37841] Liou, Y.-C.; Ryo, A.; Huang, H.-K.; Lu, P.-J.; Bronson, R.; Fujimori,

[37842] F.; Uchida, T.; Hunter, T.; Lu, K. P.: Loss of Pin1 function in the

[37843] mouse causes phenotypes resembling cyclin D1-null phenotypes. Proc.

[37844] Nat. Acad. Sci. 99: 1335-1340, 2002.

[37845] 5704.

[37846] Lu, K. P.; Hanes, S. D.; Hunter, T.: A human peptidyl-prolyl isomerase

[37847] essential for regulation of mitosis. Nature 380: 544-547, 1996.

[37848] 5705.

[37849] Lu, P.-J.; Wulf, G.; Zhou, X. Z.; Davies, P.; Lu, K. P.: The prolyl

[37850] isomerase Pin1 restores the function of Alzheimer-associated phosphorylated

[37851] tau protein. Nature 399: 784-788, 1999.

[37852] 5706.

[37853] Maleszka, R.; Hanes, S. D.; Hackett, R. L.; de Couet, H. G.; Gabor

[37854] Miklos, G. L.: The Drosophila melanogaster dodo (dod) gene, conserved

[37855] in humans, is functionally interchangeable with the ESS1 cell division

[37856] gene of Saccharomyces cerevisiae. Proc. Nat. Acad. Sci. 93: 447-451,

[37857] 1996.

[37858] 5707.

[37859] Winkler, K. E.; Swanson, K. I.; Kornbluth, S.; Means, A. R.: Requirement

[37860] of the prolyl isomerase Pin1 for the replication check-point. Science 287:

[37861] 1644-1647, 2000.

[37862] 5708.

[37863] Takahashi, T.; Fournier, A.; Nakamura, F.; Wang, L.-H.; Murakami,

[37864] Y.; Kalb, R. G.; Fujisawa, H.; Strittmatter, S. M.: Plexin-

neuropilin-1

[37865] complexes form functional semaphorin-3A receptors. Cell 99: 59-69,

[37866] 1999.

[37867] 5709.

[37868] Santagata, S.; Boggon, T. J.; Baird, C. L.; Gomez, C. A.; Zhao,

[37869] J.; Shan, W. S.; Myszka, D. G.; Shapiro, L.: G-protein signaling

[37870] through tubby proteins. Science 292: 2041-2050, 2001.

[37871] 5710.

[37872] Sasaki, T.; Irie-Sasaki, J.; Horie, Y.; Bachmaier, K.; Fata, J.

[37873] E.; Li, M.; Suzuki, A.; Bouchard, D.; Ho, A.; Redston, M.; Gallinger,

[37874] S.; Khokha, R.; Mak, T. W.; Hawkins, P. T.; Stephens, L.; Scherer,

[37875] S. W.; Tsao, M.; Penninger, J. M.: Colorectal carcinomas in mice

[37876] lacking the catalytic subunit of PI(3)K-gamma. Nature 406: 897-902,

[37877] 2000.

[37878] 5711.

[37879] Sasaki, T.; Irie-Sasaki, J.; Jones, R. G.; Oliveira-dos-Santos,

[37880] A. J.; Stanford, W. L.; Bolon, B.; Wakeham, A.; Itie, A.; Bouchard,

[37881] D.; Kozieradzki, I.; Joza, N.; Mak, T. W.; Ohashi, P. S.; Suzuki,

[37882] A.; Penninger, J. M.: Function of PI3K-gamma in thymo-cyte development,

[37883] T cell activation, and neutrophil migration. Science 287: 1040-1046,

[37884] 2000.

[37885] 5712.

[37886] Stoyanov, B.; Volinia, S.; Hanck, T.; Rubio, I.; Loubtchenkov,

[37887] M.; Malek, D.; Stoyanova, S.; Vanhaesebroeck, B.; Dhand, R.; Nurnberg,

[37888] B.; Gierschik, P.; Seedorf, K.; Hsuan, J. J.; Waterfield, M. D.; Wetzker,

[37889] R.: Cloning and characterization of a G protein-activated human phosphoinositide-3

[37890] kinase. Science 269: 690-693, 1995.

[37891] 5713.

[37892] Hotten, G.; Neidhardt, H.; Schneider, C.; Pohl, J.: Cloning of

[37893] a new member of the TGF-beta family: a putative new activin betaC

[37894] chain. Biochem. Biophys. Res. Commun. 206: 608-613, 1995.

[37895] 5714.

[37896] Ling, N.; Ying, S. Y.; Ueno, N.; Shimasaki, S.; Esch, F.; Hotta,

[37897] M.; Guillemin, R.: Pituitary FSH is released by a heterodimer of

[37898] the beta-subunits from the two forms of inhibin. Nature 321: 779-782,

[37899] 1986.

[37900] 5715.

[37901] Schmitt, J.; Hotten, G.; Jenkins, N. A.; Gilbert, D. J.; Copeland,

[37902] N. G.; Pohl, J.; Schrewe, H.: Structure, chromosomal localization,

[37903] and expression analysis of the mouse inhibin/activin betaC (Inhbc)

[37904] gene. Genomics 32: 358-366, 1996.

[37905] 5716.

[37906] Savitsky, K.; Ziv, Y.; Bar-Shira, A.; Gilad, S.; Tagle, D. A.;

[37907] Smith, S.; Uziel, T.; Sfez, S.; Nahmias, J.; Sartiel, A.; Eddy,
R.

[37908] L.; Shows, T. B.; Collins, F. S.; Shiloh, Y.; Rotman, G.: A
human

[37909] gene (DDX10) encoding a putative DEAD-box RNA heli-
case at 11q22-q23. Genomics 33:

[37910] 199-206, 1996.

[37911] 5717.

[37912] Albertsen, H. M.; Smith, S. A.; Melis, R.; Williams, B.; Holik,

[37913] P.; Stevens, J.; White, R.: Sequence, genomic structure, and
chromosomal

[37914] assignment of human DOC-2. Genomics 33: 207-213,
1996.

[37915] 5718.

[37916] Mok, S. C.; Chan, W. Y.; Wong, K. K.; Cheung, K. K.; Lau, C.
C.;

[37917] Ng, S. W.; Baldini, A.; Colitti, C. V.; Rock, C. O.; Berkowitz,
R.

[37918] S.: DOC-2, a candidate tumor suppressor gene in human epithelial

[37919] ovarian cancer. Oncogene 16: 2381-2387, 1998.

[37920] 5719.

[37921] Kimura, T.; Arakawa, Y.; Inoue, S.; Fukushima, Y.; Kondo, I.; Koyama,

[37922] K.; Hosoi, T.; Orimo, A.; Muramatsu, M.; Nakamura, Y.; Abe, T.; Inazawa,

[37923] J.: The brain finger protein gene (ZNF179), a member of the RING

[37924] finger family, maps within the Smith-Magenis syndrome region at 17p11.2. Am.

[37925] J. Med. Genet. 69: 320-324, 1997.

[37926] 5720.

[37927] Matsuda, Y.; Inue, S.; Seki, N.; Hosoi, T.; Orimo, A.; Mura-matsu,

[37928] M.; Hori, T.: Chromosome mapping of human (ZNF179), mouse, and rat

[37929] genes for brain finger protein (bfp), a member of the RING finger

[37930] family. Genomics 33: 325-327, 1996.

[37931] 5721.

[37932] Ambrose, H. J.; Blake, D. J.; Nawrotzki, R. A.; Davies, K. E.:

[37933] Genomic organization of the mouse dystrobrevin gene:
comparative analysis

[37934] with the dystrophin gene. Genomics 39: 359-369, 1997.

[37935] 5722.

[37936] Blake, D. J.; Nawrotzki, R.; Loh, N. Y.; Gorecki, D. C.;
Davies,

[37937] K. E.: Beta-dystrobrevin, a member of the dystrophin-re-
lated protein

[37938] family. Proc. Nat. Acad. Sci. 95: 241-246, 1998.

[37939] 5723.

[37940] Ichida, F.; Tsubata, S.; Bowles, K. R.; Haneda, N.; Uese, K.;
Miyawaki,

[37941] T.; Dreyer, W. J.; Messina, J.; Li, H.; Bowles, N. E.; Towbin,
J.

[37942] A.: Novel gene mutations in patients with left ventricular
noncompaction

[37943] or Barth syndrome. Circulation 103: 1256-1263, 2001.

[37944] 5724.

[37945] England, S. K.; Uebele, V. N.; Shear, H.; Kodali, J.; Bennett,

[37946] P. B.; Tamkun, M. M.: Characterization of a voltage-gated K+ channel

[37947] beta subunit expressed in human heart. Proc. Nat. Acad. Sci. 92:

[37948] 6309-6313, 1995.

[37949] 5725.

[37950] Jones, J. M.; Bentley, E.; Meisler, M. H.; Darling, S. M.: Genetic

[37951] mapping of the voltage-gated shaker potassium channel beta subunit

[37952] Kcnab1 to mouse chromosome 3. Mammalian Genome 9: 260 only, 1998.

[37953] 5726.

[37954] Leicher, T.; Bahring, R.; Isbrandt, D.; Pongs, O.: Coexpression

[37955] of the KCNA3B gene product with Kv1.5 leads to a novel A-type potassium

[37956] channel. J. Biol. Chem. 273: 35095-35101, 1998.

[37957] 5727.

[37958] Leicher, T. Roeper, J.; Weber, K.; Wang, X.; Pongs, O.:

Structural

[37959] and functional characterization of human potassium channel subunit

[37960] beta-1 (KCNA1B). Neuropharmacology 35: 787-795, 1996.

[37961] 5728.

[37962] McCormack, K.; McCormack, T.; Tanouye, M.; Rudy, B.; Stuhmer, W.

[37963] : Alternative splicing of the human Shaker K+ channel beta-1 gene

[37964] and functional expression of the beta-2 gene product. FEBS Lett. 370:

[37965] 32-36, 1995.

[37966] 5729.

[37967] Schultz, D.; Litt, M.; Smith, L.; Thayer, M.; McCormack, K.: Localization

[37968] of two potassium channel beta subunit genes, KCNA1B and KCNA2B. Genomics 31:

[37969] 389-391, 1996.

[37970] 5730.

[37971] Collin, G. B.; Nishina, P. M.; Marshall, J. D.; Naggert, J. K.

[37972] : Human DCTN1: genomic structure and evaluation as a candidate for

[37973] Alstrom syndrome. Genomics 53: 359-364, 1998.

[37974] 5731.

[37975] Holzbaur, E. L. F.; Tokito, M. K.: Localization of the DCTN1 gene

[37976] encoding p150(Glued) to human chromosome 2p13 by fluorescence in situ

[37977] hybridization. Genomics 31: 398-399, 1996.

[37978] 5732.

[37979] Holzbaur, E. L. F.; Vallee, R. B.: Dyneins: molecular structure

[37980] and cellular function. Ann. Rev. Cell Biol. 10: 339-372, 1994.

[37981] 5733.

[37982] Jang, W.; Weber, J. S.; Tokito, M. K.; Holzbaur, E. L. F.; Meisler,

[37983] M. H.: Mouse p150(Glued) (dynactin 1) cDNA sequence and evaluation

[37984] as a candidate for the neuromuscular disease mutation mnd2. Biochem.

[37985] Biophys. Res. Commun. 231: 344-347, 1997.

[37986] 5734.

[37987] Korthaus, D.; Wedemeyer, N.; Lengeling, A.; Ronsiek, M.; Jockusch,

[37988] H.; Schmitt-John, T.: Integrated radiation hybrid map of human chromosome

[37989] 2p13: possible involvement of dynactin in neuromuscular diseases. Genomics 43:

[37990] 242-244, 1997.

[37991] 5735.

[37992] Pushkin, A.; Abuladze, N.; Newman, D.; Tatishchev, S.; Kurtz, I.

[37993] : Genomic organization of the DCTN1-SLC4A5 locus encoding both NBC4

[37994] and p150(Glued). Cytogenet. Cell Genet. 95: 163-168, 2001.

[37995] 5736.

[37996] Miyata, A.; Hara, S.; Yokoyama, C.; Inoue, H.; Ullrich, V.; Tanabe,

[37997] T.: Molecular cloning and expression of human prostacyclin synthase. Biochem.

[37998] Biophys. Res. Commun. 200: 1728-1734, 1994.

[37999] 5737.

[38000] Inoue, H.; Ishii, H.; Alder, H.; Snyder, E.; Druck, T.; Hueb-
ner,
[38001] K.; Croce, C. M.: Sequence of the FRA3B common fragile
region: implications
[38002] for the mechanism of FHIT deletion. Proc. Nat. Acad. Sci.
94: 14584-14589,
[38003] 1997.

[38004] 5738.

[38005] Morikawa, H.; Nakagawa, Y.; Hashimoto, K.; Niki, M.;
Egashira,
[38006] Y.; Hirata, I.; Katsu, K.; Akao, Y.: Frequent altered expres-
sion
[38007] of fragile histidine triad protein in human colorectal ade-
nomas. Biochem.
[38008] Biophys. Res. Commun. 278: 205-210, 2000.

[38009] 5739.

[38010] Nelson, D. R.; Koymans, L.; Kamataki, T.; Stegeman, J. J.;
Feyereisen,
[38011] R.; Waxman, D. J.; Waterman, M. R.; Gotoh, O.; Coon, M. J.;

Estabrook,
[38012] R. W.; Gunsalus, I. C.; Nebert, D. W.: P450 superfamily: update on
[38013] new sequences, gene mapping, accession numbers and nomenclature. Pharmacogenetics 6:
[38014] 1-42, 1996.

[38015] 5740.

[38016] Wang, L.-H.; Chen, L.: Organization of the gene encoding human
[38017] prostacyclin synthase. Biochem. Biophys. Res. Commun. 226: 631-637,
[38018] 1996.

[38019] 5741.

[38020] Yokoyama, C.; Yabuki, T.; Inoue, H.; Tone, Y.; Hara, S.; Hatae,
[38021] T.; Nagata, M.; Takahashi, E.-I.; Tanabe, T.: Human gene encoding
[38022] prostacyclin synthase (PTGIS): genomic organization, chromosomal localization,
[38023] and promoter activity. Genomics 36: 296-304, 1996.

[38024] 5742.

[38025] Brindle, N. P. J.; Holt, M. R.; Davies, J. E.; Price, C. J.; Critchley,
[38026] D. R.: The focal-adhesion vasodilator-stimulated phosphoprotein (VASP)
[38027] binds to the proline-rich domain in vinculin. Biochem. J. 318: 753-757,
[38028] 1996.

[38029] 5743.

[38030] Haffner, C.; Jarchau, T.; Reinhard, M.; Hoppe, J.; Lohmann, S.
[38031] M.; Walter, V.: Molecular cloning, structural analysis and functional
[38032] expression of the proline-rich focal adhesion and microfilament-associated
[38033] protein VASP. EMBO J. 14: 19-27, 1995.

[38034] 5744.

[38035] Zimmer, M.; Fink, T.; Fischer, L.; Hauser, W.; Scherer, K.; Lichter,
[38036] P.; Walter, U.: Cloning of the VASP (vasodilator-stimulated phosphoprotein)
[38037] genes in human and mouse: structure, sequence, and chromosomal localization. Genomics 36:

[38038] 227-233, 1996.

[38039] 5745.

[38040] Hackam, A. S.; Yassa, A. S.; Singaraja, R.; Metzler, M.; Gutekunst,

[38041] C.-A.; Gan, L.; Warby, S.; Wellington, C. L.; Vaillancourt, J.; Chen,

[38042] N.; Gervais, F. G.; Raymond, L.; Nicholson, D. W.; Hayden, M. R.:

[38043] Huntingtin interacting protein 1 induces apoptosis via a novel caspase-dependent

[38044] death effector domain. J. Biol. Chem. 275: 41299-41308, 2000.

[38045] 5746.

[38046] Himmelbauer, H.; Wedemeyer, N.; Haaf, T.; Wanker, E. E.; Schalkwyk,

[38047] L. C.; Lehrach, H.: IRS-PCR-based genetic mapping of the huntingtin

[38048] interacting protein gene (HIP1) on mouse chromosome 5. Mammalian

[38049] Genome 9: 26-31, 1998.

[38050] 5747.

[38051] Manabe, T.; Noda, Y.; Mamiya, T.; Katagiri, H.; Houtani, T.; Nishi,

[38052] M.; Noda, T.; Takahashi, T.; Sugimoto, T.; Nabeshima, T.; Takeshima,

[38053] H.: Facilitation of long-term potentiation and memory in mice lacking

[38054] nociceptin receptors. Nature 394: 577-581, 1998.

[38055] 5748.

[38056] Mollereau, C.; Simons, M.-J.; Soularue, P.; Liners, F.; Vas-

[38057] sart, G.; Meunier, J.-C.; Parmentier, M.: Structure, tissue distri-

[38058] bution, and chromosomal localization of the prepronociceptin gene. Proc.

[38059] Nat. Acad. Sci. 93: 8666-8700, 1996.

[38060] 5749.

[38061] Nishi, M.; Houtani, T.; Noda, Y.; Mamiya, T.; Sato, K.; Doi, T.;

[38062] Kuno, J.; Takeshima, H.; Nukada, T.; Nabeshima, T.; Ya-

[38063] mashita, T.; Noda, T.; Sugimoto, T.: Unrestrained nociceptive response and disregulation

[38064] of hearing ability in mice lacking the nociceptin/orphanin FQ receptor. EMBO
[38065] J. 16: 1858-1864, 1997.

[38066] 5750.

[38067] Okuda-Ashitaka, E.; Minami, T.; Tachibana, S.; Yoshihara, Y.; Nishiuchi,
[38068] Y.; Kimura, T.; Ito, S.: Nocistatin, a peptide that blocks nociceptin
[38069] action in pain transmission. Nature 392: 286-289, 1998.

[38070] 5751.

[38071] Pan, Z. Z.; Hirakawa, N.; Fields, H. L.: A cellular mechanism
[38072] for the bidirectional pain-modulating actions of orphanin FQ/nociceptin. Neuron 26:
[38073] 515-522, 2000.

[38074] 5752.

[38075] Serhan, C. N.; Fierro, I. M.; Chiang, N.; Pouliot, M.: Cutting
[38076] edge: nociceptin stimulates neutrophil chemotaxis and recruitment:
[38077] inhibition by aspirin-triggered-15-epi-lipoxin A(4). J. Immun. 166:

[38078] 3650-3654, 2001.

[38079] 5753.

[38080] Kanai, N.; Lu, R.; Satriano, J. A.; Bao, Y.; Wolkoff, A. W.; Schuster,

[38081] V. L.: Identification and characterization of a prostaglandin transporter. Science 268:

[38082] 866-869, 1995.

[38083] 5754.

[38084] Lu, R.; Kanai, N.; Bao, Y.; Schuster, V. L.: Cloning, in vitro

[38085] expression, and tissue distribution of a human prostaglandin transporter

[38086] cDNA (hPGT). J. Clin. Invest. 98: 1142-1149, 1996.

[38087] 5755.

[38088] Lu, R.; Schuster, V. L.: Molecular cloning of the gene for the

[38089] human prostaglandin transporter hPGT: gene organiza-tion, promoter

[38090] activity, and chromosomal localization. Biochem. Biophys. Res. Commun. 246:

[38091] 805-812, 1998.

[38092] 5756.

[38093] Funamoto, S.; Meili, R.; Lee, S.; Parry, L.; Firtel, R. A.: Spa-
tial
[38094] and temporal regulation of 3-phosphoinositides by PI
3-kinase and
[38095] PTEN mediates chemotaxis. Cell 109: 611-623, 2002.

[38096] 5757.

[38097] Gimm, O.; Attie-Bitach, T.; Lees, J. A.; Vekemans, M.; Eng,
C.
[38098] : Expression of the PTEN tumour suppressor protein dur-
ing human development. Hum.
[38099] Molec. Genet. 9: 1633-1639, 2000.

[38100] 5758.

[38101] Groszer, M.; Erickson, R.; Scripture-Adams, D. D.; Lesche,
R.;
[38102] Trumpp, A.; Zack, J. A.; Kornblum, H. I.; Liu, X.; Wu, H.:
Negative
[38103] regulation of neural stem/progenitor cell proliferation by
the Pten
[38104] tumor suppressor gene in vivo. Science 294: 2186-2189,
2001.

[38105] 5759.

[38106] Forrest, M. S.; Edwards, S. M.; Hamoudi, R. A.; Dearnaley, D.

[38107] P.; Arden-Jones, A.; Dowe, A.; Murkin, A.; Kelly, J.; Teare, M. D.;

[38108] Easton, D. F.; Knowles, M. A.; Bishop, D. T.; Eeles, R. A.; CRC/BPG

[38109] UK Familial Prostate Cancer Study Collaborators; EC Biomed Familial

[38110] Prostate Cancer Study Collaborators: No evidence of germline PTEN

[38111] mutations in familial prostate cancer. J. Med. Genet. 37: 210-212,

[38112] 2000.

[38113] 5760.

[38114] Hansen, G. M.; Justice, M. J.: Pten, a candidate tumor suppressor

[38115] gene, maps to mouse chromosome 19. Mammalian Genome 9: 88-90, 1998.

[38116] 5761.

[38117] Iijima, M.; Devreotes, P.: Tumor suppressor PTEN mediates sensing

[38118] of chemoattractant gradients. Cell 109: 599-610, 2002.

[38119] 5762.

[38120] Kim, S. K.; Su, L.-K.; Oh, Y.; Kemp, B. L.; Hong, W. K.; Mao,

[38121] L.: Alterations of PTEN/MMAC1, a candidate tumor sup-
pressor gene,

[38122] and its homologue, PTH2, in small cell lung cancer cell
lines. Oncogene 16:

[38123] 89-93, 1998.

[38124] 5763.

[38125] Kong, D.; Suzuki, A.; Zou, T.-T.; Sakurada, A.; Kemp, L.
W.; Wakatsuki,

[38126] S.; Yokoyama, T.; Yamakawa, H.; Furukawa, T.; Sato, M.;
Ohuchi, N.;

[38127] Sato, S.; Yin, J.; Wang, S.; Abraham, J. M.; Souza, R. F.;
Smolinski,

[38128] K. N.; Meltzer, S. J.; Horii, A.: PTEN1 is frequently mutated
in

[38129] primary endometrial carcinomas. (Letter) Nature Genet.
17: 143-144,

[38130] 1997.

[38131] 5764.

[38132] Kurose, K.; Gilley, K.; Matsumoto, S.; Watson, P. H.; Zhou, X.-P.;

[38133] Eng, C.: Frequent somatic mutations in PTEN and TP53 are mutually

[38134] exclusive in the stroma of breast carcinomas. Nature Genet. 15Oct.:

[38135] , 2002. Note: Advance Electronic Publication.

[38136] 5765.

[38137] Kurose, K.; Zhou, X.-P.; Araki, T.; Eng, C.: Biallelic inacti- vating

[38138] mutations and an occult germline mutation of PTEN in primary cervical

[38139] carcinomas. Genes Chromosomes Cancer 29: 166-172, 2000.

[38140] 5766.

[38141] Kwabi-Addo, B.; Giri, D.; Schmidt, K.; Podsypanina, K.; Parsons,

[38142] R.; Greenberg, N.; Ittmann, M.: Haploinsufficiency of the Pten tumor

[38143] suppressor gene promotes prostate cancer progression. Proc. Nat.

[38144] Acad. Sci. 98: 11563-11568, 2001.

[38145] 5767.

[38146] Lee, J.-O.; Yang, H.; Georgescu, M.-M.; Di Cristofano, A.; Maehama,

[38147] T.; Shi, Y.; Dixon, J. E.; Pandolfi, P.; Pavletich, N. P.: Crystal

[38148] structure of the PTEN tumor suppressor: implications for its phosphoinositide

[38149] phosphatase activity and membrane association. Cell 99: 323-334,

[38150] 1999.

[38151] 5768.

[38152] Li, D.-M.; Sun, H.: PTEN/MMAC1/TEP1 suppresses the tu-morigenicity

[38153] and induces G1 cell cycle arrest in human glioblastoma cells. Proc.

[38154] Nat. Acad. Sci. 95: 15406-15411, 1998.

[38155] 5769.

[38156] Li, J.; Yen, C.; Liaw, D.; Podsypanina, K.; Bose, S.; Wang, S.

[38157] I.; Puc, J.; Miliaresis, C.; Rodgers, L.; McCombie, R.; Bigner, S.

[38158] H.; Giovanella, B. C.; Ittmann, M.; Tycko, B.; Hibshoosh, H.; Wigler,

[38159] M. H.; Parsons, R.: PTEN, a putative protein tyrosine phosphatase
[38160] gene mutated in human brain, breast, and prostate cancer. Science 275:
[38161] 1943-1946, 1997.

[38162] 5770.

[38163] Li, L.; Ernsting, B. R.; Wishart, M. J.; Lohse, D. L.; Dixon,
[38164] J. E.: A family of putative tumor suppressors is structurally and
[38165] functionally conserved in humans and yeast. J. Biol. Chem. 272:
[38166] 29403-29406, 1997.

[38167] 5771.

[38168] Marsh, D. J.; Coulon, V.; Lunetta, K. L.; Rocca-Serra, P.; Dahia,
[38169] P. L. M.; Zheng, Z.; Liaw, D.; Caron, S.; Duboue, B.; Lin, A. Y.;
[38170] Richardson, A.-L.; Bonnetblanc, J.-M.; and 25 others: Mutation
[38171] spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana

[38172] syndrome, two hamartoma syndromes with germline PTEN mutation. Hum.

[38173] Molec. Genet. 7: 507-515, 1998.

[38174] 5772.

[38175] Mutter, G. L.; Lin, M.-C.; Fitzgerald, J. T.; Kum, J. B.; Eng,

[38176] C.: Changes in endometrial PTEN expression throughout the human menstrual

[38177] cycle. J. Clin. Endocr. Metab. 85: 2334-2338, 2000.

[38178] 5773.

[38179] Loughna, S.; Sato, T. N.: A combinatorial role of angiopoi-etin-1

[38180] and orphan receptor TIE1 pathways in establishing vascu-lar polarity

[38181] during angiogenesis. Molec. Cell 7: 233-239, 2001.

[38182] 5774.

[38183] Bandmann, O.; Nygaard, T. G.; Surtees, R.; Marsden, C. D.; Wood,

[38184] N. W.; Harding, A. E.: Dopa-responsive dystonia in British patients:

[38185] new mutations of the GTP-cyclohydrolase I gene and evi-dence for genetic

[38186] heterogeneity. Hum. Molec. Genet. 5: 403-406, 1996.

[38187] 5775.

[38188] Blau, N.; Ichinose, H.; Nagatsu, T.; Heizmann, C. W.; Zac-
chello,
[38189] F.; Burlina, A. B.: A missense mutation in a patient with
guanosine
[38190] triphosphate cyclohydrolase I deficiency missed in the
newborn screening
[38191] program. J. Pediat. 126: 401-405, 1995.

[38192] 5776.

[38193] Bode, V. C.; McDonald, J. D.; Guenet, J.-L.; Simon, D.:
Hph-1:
[38194] a mouse mutant with hereditary hyperphenylalaninemia
induced by ethylnitrosourea
[38195] mutagenesis. Genetics 118: 299-305, 1988.

[38196] 5777.

[38197] Brique, S.; Destee, A.; Lambert, J.-C.; Mouroux, V.; Dela-
courte,
[38198] A.; Amouyel, P.; Chartier-Harlin, M.-C.: A new GTP-
cyclohydrolase
[38199] I mutation in an unusual dopa-responsive dystonia, famil-

ial form. Neuroreport 10:
[38200] 487-491, 1999.

[38201] 5778.

[38202] Furukawa, Y.; Kish, S. J.; Bebin, E. M.; Jacobson, R. D.; Fry-
burg,
[38203] J. S.; Wilson, W. G.; Shimadzu, M.; Hyland, K.; Trugman, J.
M.: Dystonia
[38204] with motor delay in compound heterozygotes for GTP-
cyclohydrolase
[38205] I gene mutations. Ann. Neurol. 44: 10-16, 1998.

[38206] 5779.

[38207] Hahn, H.; Trant, M. R.; Brownstein, M. J.; Harper, R. A.;
Milstien,
[38208] S.; Butler, I. J.: Neurologic and psychiatric manifestations
in a
[38209] family with a mutation in exon 2 of the guanosine
triphosphate-cyclohydrolase
[38210] gene. Arch. Neurol. 58: 749-755, 2001.

[38211] 5780.

[38212] Hirano, M.; Imaiso, Y.; Ueno, S.: Differential splicing of the

[38213] GTP cyclohydrolase I RNA in dopa-responsive dystonia.

Biochem. Biophys.
[38214] Res. Commun. 234: 316-319, 1997.

[38215] 5781.

[38216] Hong, K.-M.; Kim, Y.-S.; Paik, M.-K.: A novel nonsense mutation
[38217] of the GTP cyclohydrolase I gene in a family with dopa-responsive
[38218] dystonia. Hum. Hered. 52: 59-60, 2001.

[38219] 5782.

[38220] Hwu, W.-L.; Wang, P.-J.; Hsiao, K.-J.; Wang, T.-R.; Chiou, Y.-W.;
[38221] Lee, Y.-M.: Dopa-responsive dystonia induced by a recessive GTP cyclohydrolase
[38222] I mutation. Hum. Genet. 105: 226-230, 1999.

[38223] 5783.

[38224] Hyland, K.; Nygaard, T. G.; Trugman, J. M.; Swoboda, K. J.; Arnold,
[38225] L. A.; Sparagana, S. P.: Oral phenylalanine loading profiles in symptomatic
[38226] and asymptomatic gene carriers with dopa-responsive dystonia due to

[38227] dominantly inherited GTP cyclohydrolase deficiency. J. Inherit. Metab.

[38228] Dis. 22: 213-215, 1999.

[38229] 5784.

[38230] Ichinose, H.; Ohye, T.; Matsuda, Y.; Hori, T.; Blau, N.; Burlina,

[38231] A.; Rouse, B.; Matalon, R.; Fujita, K.; Nagatsu, T.: Characterization

[38232] of mouse and human GTP cyclohydrolase I genes: mutations in patients

[38233] with GTP cyclohydrolase I deficiency. J. Biol. Chem. 270: 10062-10071,

[38234] 1995.

[38235] 5785.

[38236] Inagaki, H.; Ohye, T.; Suzuki, T.; Segawa, M.; Nomura, Y.; Nagatsu,

[38237] T.; Ichinose, H.: Decrease in GTP cyclohydrolase I gene expression

[38238] caused by inactivation of one allele in hereditary progressive dystonia

[38239] with marked diurnal fluctuation. Biochem. Biophys. Res. Commun. 260:

[38240] 747-751, 1999.

[38241] 5786.

[38242] McDonald, J. D.; Cotton, R. J. H.; Jennings, I.; Ledley, F. D.;

[38243] Woo, S. L. C.; Bode, V. C.: Biochemical defect of hph-1 mouse mutant

[38244] is a deficiency in GTP-cyclohydrolase activity. J. Neu- rochem. 50:

[38245] 655-657, 1988.

[38246] 5787.

[38247] Montanez, C. S.; McDonald, J. D.: Linkage analysis of the hph-1

[38248] mutation and the GTP cyclohydrolase I structural gene. Molec. Genet.

[38249] Metab. 68: 91-92, 1999.

[38250] 5788.

[38251] Skrygan, M.; Bartholome, B.; Bonafe, L.; Blau, N.; Bartholome,

[38252] K.: A splice mutation in the GTP cyclohydrolase I gene causes dopa-responsive

[38253] dystonia by exon skipping. J. Inherit. Metab. Dis. 24: 345-351,

[38254] 2001.

[38255] 5789.

[38256] Steinberger, D.; Topka, H.; Fischer, D.; Muller, U.: GCH1 mutation

[38257] in a patient with adult-onset oromandibular dystonia. Neurology 52:

[38258] 877-879, 1999.

[38259] 5790.

[38260] Tamaru, Y.; Hirano, M.; Ito, H.; Kawamura, J.; Matsumoto, S.;

[38261] Imai, T.; Ueno, S.: Clinical similarities of hereditary pro-gressive/dopa

[38262] responsive dystonia caused by different types of muta-tions in the

[38263] GTP cyclohydrolase I gene. J. Neurol. Neurosurg. Psychiat. 64: 469-473,

[38264] 1998.

[38265] 5791.

[38266] Furuichi, M.; Yoshida, M. C.; Oda, H.; Tajiri, T.; Nakabeppu, Y.;

[38267] Tsuzuki, T.; Sekiguchi, M.: Genomic structure and chro-

mosome location

[38268] of the human mutT homologue gene MTH1 encoding 8-oxo-dGTPase for prevention

[38269] of A:T to C:G transversion. Genomics 24: 485-490, 1994.

[38270] 5792.

[38271] Igarashi, H.; Tsuzuki, T.; Kakuma, T.; Tominaga, Y.; Sekiguchi,

[38272] M.: Organization and expression of the mouse MTH1 gene for preventing

[38273] transversion mutation. J. Biol. Chem. 272: 3766-3772, 1997.

[38274] 5793.

[38275] Tsuzuki, T.; Egashira, A.; Igarashi, H.; Iwakuma, T.; Nakat-suru,

[38276] Y.; Tominaga, Y.; Kawate, H.; Nakao, K.; Nakamura, K.; Ide, F.; Kura,

[38277] S.; Nakabeppu, Y.; Katsuki, M.; Ishikawa, T.; Sekiguchi, M.: Spontaneous

[38278] tumorigenesis in mice defective in the MTH1 gene encod-ing 8-oxo-dGTPase. Proc.

[38279] Nat. Acad. Sci. 98: 11456-11461, 2001.

[38280] 5794.

[38281] Tafuri, A.; Shahinian, A.; Bladt, F.; Yoshinaga, S. K.; Jordana,
[38282] M.; Wakeham, A.; Boucher, L.-M.; Bouchard, D.; Chan, V. S. F.; Duncan,
[38283] G.; Odermatt, B.; Ho, A.; Itie, A.; Horan, T.; Whoriskey, J. S.; Pawson,
[38284] T.; Penninger, J. M.; Ohashi, P. S.; Mak, T. W.: ICOS is es-
sential
[38285] for effective T-helper-cell responses. Nature 409: 105-109, 2001.
[38286] 5795.
[38287] Yoshinaga, S. K.; Whoriskey, J. S.; Khare, S. D.; Sarmiento, U.;
[38288] Guo, J.; Horan, T.; Shih, G.; Zhang, M.; Coccia, M. A.; Kohno, T.;
[38289] Tafuri-Bladt, A.; Brankow, D.; and 14 others: T-cell co-
stimulation
[38290] through B7RP-1 and ICOS. Nature 402: 827-832, 1999.
[38291] 5796.
[38292] Clark, S. W.; Staub, O.; Clark, I. B.; Holzbaur, E. L. F.; Paschal,
[38293] B. M.; Vallee, R. B.; Meyer, D. I.: Beta-centractin: charac-

terization
[38294] and distribution of a new member of the centractin family of actin-related
[38295] proteins. Molec. Biol. Cell 5: 1301-1310, 1994.

[38296] 5797.

[38297] Elsea, S. H.; Clark, I. B.; Juyal, R. C.; Meyer, D. J.; Meyer,

[38298] D. I.; Patel, P. I.: Assignment of beta-centractin (CTRN2) to human
[38299] chromosome 2 bands q11.1-q11.2 with somatic cell hybrids and in situ
[38300] hybridization. Cytogenet. Cell Genet. 84: 48-49, 1999.

[38301] 5798.

[38302] Ho, A. M.; Johnson, M. D.; Kingsley, D. M.: Role of the mouse
[38303] ank gene in control of tissue calcification and arthritis. Science 289:
[38304] 265-270, 2000.

[38305] 5799.

[38306] Yu, H.; Peters, J.-M.; King, R. W.; Page, A. M.; Hieter, P.; Kirschner,
[38307] M. W.: Identification of a cullin homology region in a sub-

unit of
[38308] the anaphase-promoting complex. Science 279:
1219-1222, 1998.
[38309] 5800.

[38310] Zhao, N.; Lai, F.; Fernald, A. A.; Eisenbart, J. D.; Espinosa,

[38311] R., III.; Wang, P. W.; Le Beau, M. M.: Human CDC23: cDNA
cloning,
[38312] mapping to 5q31, genomic structure, and evaluation as a
candidate
[38313] tumor suppressor gene in myeloid leukemias. Genomics
53: 184-190,
[38314] 1998.

[38315] 5801.

[38316] Brambilla, R.; Draetta, G.: Molecular cloning of PISSLRE, a
novel
[38317] putative member of the cdk family of protein serine/
threonine kinases. Oncogene 9:
[38318] 3037-3041, 1994.

[38319] 5802.

[38320] Grana, X.; Claudio, P. P.; De Luca, A.; Sang, N.; Giordano,
A.

[38321] : PISSLRE, a human novel CDC2-related protein kinase. Oncogene 9:

[38322] 2097-2103, 1994.

[38323] 5803.

[38324] de Winter, J. P.; Rooimans, M. A.; van der Weel, L.; van Berkel,

[38325] C. G. M.; Alon, N.; Bosnoyan-Collins, L.; de Groot, J.; Zhi, Y.; Waisfisz,

[38326] Q.; Pronk, J. C.; Arwert, F.; Mathew, C. G.; Scheper, R. J.; Hoatlin,

[38327] M. E.; Buchwald, M.; Joenje, H.: The Fanconi anaemia gene FANCF encodes

[38328] a novel protein with homology to ROM. (Letter) Nature Genet. 24:

[38329] 15-16, 2000.

[38330] 5804.

[38331] de Winter, J. P.; van der Weel, L.; de Groot, J.; Stone, S.; Waisfisz,

[38332] Q.; Arwert, F.; Scheper, R. J.; Kruyt, F. A. E.; Hoatlin, M. E.; Joenje,

[38333] H.: The Fanconi anemia protein FANCF forms a nuclear complex with

[38334] FANCA, FANCC and FANCG. Hum. Molec. Genet. 9: 2665-2674, 2000.

[38335] 5805.

[38336] Beaudet, A. L.; O'Brien, W. E.; Bock, H.-G. O.; Freytag, S. O.;

[38337] Su, T.-S.: The human argininosuccinate synthetase locus and citrullinemia. Adv.

[38338] Hum. Genet. 15: 161-196, 1986.

[38339] 5806.

[38340] Beaudet, A. L.; Su, T.-S.; Bock, H.-G.; D'Eustachio, P.; Rud-dle,

[38341] F. H.; O'Brien, W. E.: Use of a cloned cDNA to study human argininosuccinate

[38342] synthetase. (Abstract) Am. J. Hum. Genet. 33: 36A only, 1981.

[38343] 5807.

[38344] Beaudet, A. L.; Su, T.-S.; O'Brien, W. E.; D'Eustachio, P.; Barker,

[38345] P. E.; Ruddle, F. H.: Dispersion of argininosuccinate-syn-thetase-like

[38346] human genes to multiple autosomes and the X chromo-some. Cell 30:

[38347] 287-293, 1982.

[38348] 5808.

[38349] Daiger, S. P.; Wildin, R. S.; Su, T.-S.: Sequences on the hu-
man

[38350] Y chromosome homologous to the autosomal gene for
argininosuccinate

[38351] synthetase. Nature 298: 682-684, 1982.

[38352] 5809.

[38353] Underwood, K. W.; Song, C.; Kriz, R. W.; Chang, X. J.;
Knopf, J.

[38354] L.; Lin, L.-L.: A novel calcium-independent phospholipase
A(2), cPLA(2)-gamma,

[38355] that is prenylated and contains homology to cPLA(2). J.
Biol. Chem. 273:

[38356] 21926-21932, 1998.

[38357] 5810.

[38358] Cupillard, L.; Koumanov, K.; Mattei, M.-G.; Lazdunski, M.;
Lambeau,

[38359] G.: Cloning, chromosomal mapping, and expression of a
novel human

[38360] secretory phospholipase A(2). J. Biol. Chem. 272:

15745-15752, 1997.

[38361] 5811.

[38362] Marsters, S. A.; Sheridan, J. P.; Pitti, R. M.; Huang, A.; Sku-
batch,

[38363] M.; Baldwin, D.; Yuan, J.; Gurney, A.; Goddard, A. D.;
Godowski, P.;

[38364] Ashkenazi, A.: A novel receptor for Apo2L/TRAIL contains
a truncated

[38365] death domain. Curr. Biol. 7: 1003-1006, 1997.

[38366] 5812.

[38367] Pan, G.; O'Rourke, K.; Chinnaiyan, A. M.; Gentz, R.; Ebner,
R.;

[38368] Ni, J.; Dixit, V. M.: The receptor for the cytotoxic ligand
TRAIL. Science 276:

[38369] 111-113, 1997.

[38370] 5813.

[38371] MacFarlane, M.; Ahmad, M.; Srinivasula, S. M.; Fernandes-Al-
nemri,

[38372] T.; Cohen, G. M.; Alnemri, E. S.: Identification and molecu-
lar cloning

[38373] of two novel receptors for the cytotoxic ligand TRAIL. J.

Biol. Chem. 272:

[38374] 25417-25420, 1997.

[38375] 5814.

[38376] Pai, S. I.; Wu, G. S.; Ozoren, N.; Wu, L.; Jen, J.; Sidransky,

[38377] D.; El-Deiry, W. S.: Rare loss-of-function mutation of a death receptor

[38378] gene in head and neck cancer. Cancer Res. 58: 3513-3518, 1998.

[38379] 5815.

[38380] Pan, G.; Ni, J.; Wei, Y.-F.; Yu, G.; Gentz, R.; Dixit, V. M.:

[38381] An antagonist decoy receptor and a death domain-containing receptor

[38382] for TRAIL. Science 277: 815-818, 1997.

[38383] 5816.

[38384] Schneider, P.; Bodmer, J.-L.; Thome, M.; Hofmann, K.; Holler, N.;

[38385] Tschopp, J.: Characterization of two receptors for TRAIL. FEBS Lett. 416:

[38386] 329-334, 1997.

[38387] 5817.

[38388] Screaton, G. R.; Mongkolsapaya, J.; Xu, X.-N.; Cowper, A. E.; McMichael,
[38389] A. J.; Bell, J. I.: TRICK2, a new alternatively spliced receptor
[38390] that transduces the cytotoxic signal from TRAIL. Curr. Biol. 7:
[38391] 693-696, 1997.

[38392] 5818.

[38393] Sheridan, J. P.; Marsters, S. A.; Pitti, R. M.; Gurney, A.; Skubatch,
[38394] M.; Baldwin, D.; Ramakrishnan, L.; Gray, C. L.; Baker, K.; Wood, W.
[38395] I.; Goddard, A. D.; Godowski, P.; Ashkenazi, A.: Control of TRAIL-induced
[38396] apoptosis by a family of signaling and decoy receptors. Science 277:
[38397] 818-821, 1997.

[38398] 5819.

[38399] Boeglin, W. E.; Kim, R. B.; Brash, A. R.: A 12R-lipoxygenase in
[38400] human skin: mechanistic evidence, molecular cloning, and expression. Proc.

[38401] Nat. Acad. Sci. 95: 6744-6749, 1998.

[38402] 5820.

[38403] Sun, D.; McDonnell, M.; Chen, X.-S.; Lakkis, M. M.; Li, H.; Isaacs,

[38404] S. N.; Elsea, S. H.; Patel, P. I.; Funk, C. D.: Human 12(R)-lipoxygenase

[38405] and the mouse ortholog: molecular cloning, expression, and gene chromosomal

[38406] assignment. J. Biol. Chem. 273: 33540-33547, 1998.

[38407] 5821.

[38408] Duchateau, P. N.; Pullinger, C. R.; Cho, M. H.; Eng, C.; Kane,

[38409] J. P.: Apolipoprotein L gene family: tissue-specific expres- sion,

[38410] splicing, promoter regions; discovery of a new gene. J. Lipid Res. 42:

[38411] 620-630, 2001.

[38412] 5822.

[38413] Duchateau, P. N.; Pullinger, C. R.; Orellana, R. E.; Kunitake,

[38414] S. T.; Naya-Vigne, J.; O'Connor, P. M.; Malloy, M. J.; Kane, J. P.

[38415] : Apolipoprotein L, a new human high density lipoprotein apolipoprotein

[38416] expressed by the pancreas: identification, cloning, characterization,

[38417] and plasma distribution of apolipoprotein L. J. Biol. Chem. 272:

[38418] 25576-25582, 1997.

[38419] 5823.

[38420] Monajemi, H.; Fontijn, R. D.; Pannekoek, H.; Horrevoets, A. J.

[38421] G.: The apolipoprotein L gene cluster has emerged recently in evolution

[38422] and is expressed in human vascular tissue. Genomics 79: 539-546,

[38423] 2002.

[38424] 5824.

[38425] Page, N. M.; Butlin, D. J.; Lomthaisong, K.; Lowry, P. J.: The

[38426] human apolipoprotein L gene cluster: identification, classification,

[38427] and sites of distribution. Genomics 74: 71-78, 2001.

[38428] 5825.

[38429] Lin, C.-Y.; Huang, P.-H.; Liao, W.-L.; Cheng, H.-J.; Huang, C.-F.;

[38430] Kuo, J.-C.; Patton, W. A.; Massenburg, D.; Moss, J.; Lee, F.-J. S.

[38431] : ARL4, an ARF-like protein that is developmentally regulated and

[38432] localized to nuclei and nucleoli. J. Biol. Chem. 275: 37815-37823,

[38433] 2000.

[38434] 5826.

[38435] Schurmann, A.; Breiner, M.; Becker, W.; Huppertz, C.; Kainulainen,

[38436] H.; Kentrup, H.; Joost, H.-G.: Cloning of two novel ADP-ribosylation

[38437] factor-like proteins and characterization of their differential expression

[38438] in 3T3-L1 cells. J. Biol. Chem. 269: 15683-15688, 1994.

[38439] 5827.

[38440] Schurmann, A.; Koling, S.; Jacobs, S.; Saftig, P.; Kraub, S.; Wennemuth,

[38441] G.; Kluge, R.; Joost, H.-G.: Reduced sperm count and normal fertility

[38442] in male mice with targeted disruption of the ADP-ribosylation factor-like
[38443] 4 (Arl4) gene. Molec. Cell. Biol. 22: 2761-2768, 2002.

[38444] 5828.

[38445] Lehnert, K.; Ni, J.; Leung, E.; Gough, S. M.; Weaver, A.; Yao,
[38446] W.-P.; Liu, D.; Wang, S.-X.; Morris, C. M.; Krissansen, G. W.: Cloning,
[38447] sequence analysis, and chromosomal localization of the novel human
[38448] integrin alpha-11 subunit (ITGA11). Genomics 60: 179-187, 1999.
[38449] 5829.

[38450] Velling, T.; Kusche-Gullberg, M.; Sejersen, T.; Gullberg, D.:
[38451] cDNA cloning and chromosomal localization of human alpha-11 integrin:
[38452] a collagen-binding, I domain-containing, beta-1-associated integrin
[38453] alpha-chain present in muscle tissues. J. Biol. Chem. 274: 25735-25742,
[38454] 1999.

[38455] 5830.

[38456] Liao, W.; Hong, S.-H.; Chan, B. H.-J.; Rudolph, F. B.; Clark,
S.

[38457] C.; Chan, L.: APOBEC-2, a cardiac- and skeletal muscle-
specific member

[38458] of the cytidine deaminase supergene family. Biochem.
Biophys. Res.

[38459] Commun. 260: 398-404, 1999.

[38460] 5831.

[38461] Smith, S. J.; Cases, S.; Jensen, D. R.; Chen, H. C.; Sande, E.;

[38462] Tow, B.; Sanan, D. A.; Raber, J.; Eckel, R. H.; Farese, R. V.,
Jr.

[38463] : Obesity resistance and multiple mechanisms of triglyc-
eride synthesis

[38464] in mice lacking Dgat. Nature Genet. 25: 87-90, 2000.

[38465] 5832.

[38466] Mital, R.; Kobayashi, R.; Hernandez, N.: RNA polymerase III
transcription

[38467] from the human U6 and adenovirus type 2 VAI promoters
has different

[38468] requirements for human BRF, a subunit of human TFIIIB.

Molec. Cell.

[38469] Biol. 16: 7031-7042, 1996.

[38470] 5833.

[38471] Wang, Z.; Roeder, R. G.: Structure and function of a human transcription

[38472] factor TFIIIB subunit that is evolutionarily conserved and contains

[38473] both TFIIB- and high-mobility-group protein 2-related domains. Proc.

[38474] Nat. Acad. Sci. 92: 7026-7030, 1995.

[38475] 5834.

[38476] Comai, L.; Zomerdijk, J. C. B. M.; Beckmann, H.; Zhou, S.; Admon,

[38477] A.; Tjian, R.: Reconstitution of transcription factor SL1: exclusive

[38478] binding of TBP by SL1 or TFIID subunits. Science 266: 1966-1972,

[38479] 1994.

[38480] 5835.

[38481] Di Pietro, C.; Rapisarda, A.; Amico, V.; Bonaiuto, C.; Viola, A.;

[38482] Scalia, M.; Motta, S.; Amato, A.; Engel, H.; Messina, A.; Sichel,
[38483] G.; Grzeschik, K.-H.; Purrello, M.: Genomic localization of the human
[38484] genes TAF1A, TAF1B and TAF1C, encoding TAFI48, TAFI63 and TAFI110
[38485] subunits of class I general transcription initiation factor SL1. Cytogenet.
[38486] Cell Genet. 89: 133-136, 2000.

[38487] 5836.

[38488] Kloeker, S.; Wadzinski, B. E.: Purification and identification
[38489] of a novel subunit of protein serine/threonine phos-phatase 4. J.
[38490] Biol. Chem. 274: 5339-5347, 1999.

[38491] 5837.

[38492] Nakamura, T.; Takeuchi, K.; Muraoka, S.; Takezoe, H.; Takahashi,
[38493] N.; Mori, N.: A neurally enriched coronin-like protein, ClipnC, is
[38494] a novel candidate for an actin cytoskeleton-cortical mem-brane-linking

[38495] protein. J. Biol. Chem. 274: 13322-13327, 1999.

[38496] 5838.

[38497] Bennett, E. P.; Hassan, H.; Hollingsworth, M. A.; Clausen, H.:

[38498] A novel human UDP-N-acetyl-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase,

[38499] GalNAc-T7, with specificity for partial GalNAc-glycosylated acceptor

[38500] substrates. FEBS Lett. 460: 226-230, 1999.

[38501] 5839.

[38502] Saitoh, T.; Moriwaki, J.; Koike, J.; Takagi, A.; Miwa, T.; Shiokawa,

[38503] K.; Katoh, M.: Molecular cloning and characterization of FRAT2, encoding

[38504] a positive regulator of the WNT signaling pathway. Biochem. Biophys.

[38505] Res. Commun. 281: 815-820, 2001.

[38506] 5840.

[38507] Abbaszade, I.; Liu, R.-Q.; Yang, F.; Rosenfeld, S. A.; Ross, O.

[38508] H.; Link, J. R.; Ellis, D. M.; Tortorella, M. D.; Pratta, M. A.; Hollis,

[38509] J. M.; Wynn, R.; Duke, J. L.; and 15 others: Cloning and characterization

[38510] of ADAMTS11, an aggrecanase from the ADAMTS family. J. Biol. Chem. 274:

[38511] 23443-23450, 1999.

[38512] 5841.

[38513] Chen, D.; Guo, J.; Miki, T.; Tachibana, M.; Gahl, W. A.: Molecular

[38514] cloning and characterization of rab27a and rab27b, novel human rab

[38515] proteins shared by melanocytes and platelets. Biochem. Molec. Med. 60:

[38516] 27-37, 1997.

[38517] 5842.

[38518] Nagata, K.; Itoh, H.; Katada, T.; Takenaka, K.; Ui, M.; Kaziro,

[38519] Y.; Nozawa, Y.: Purification, identification, and characteri-zation

[38520] of two GTP-binding proteins with molecular weights of 25,000 and 21,000

[38521] in human platelet cytosol: one is the rap1/smg21/Krev-1 protein and
[38522] the other is a novel GTP-binding protein. J. Biol. Chem. 264: 17000-17005,
[38523] 1989.

[38524] 5843.

[38525] Nagata, K.; Satoh, T.; Itoh, H.; Kozasa, T.; Okano, Y.; Doi, T.;
[38526] Kaziro, Y.; Nozawa, Y.: The ram: a novel low molecular weight GTP-binding
[38527] protein cDNA from a rat megakaryocyte library. FEBS Lett. 275: 29-32,
[38528] 1990.

[38529] 5844.

[38530] Seabra, M. C.; Ho, Y. K.; Anant, J. S.: Deficient geranylger-anylation
[38531] of Ram/Rab27 in choroideremia. J. Biol. Chem. 270: 24420-24427,
[38532] 1995.

[38533] 5845.

[38534] Tolmachova, T.; Ramalho, J. S.; Anant, J. S.; Schultz, R. A.;

Huxley,
[38535] C. M.; Seabra, M. C.: Cloning, mapping and characterization of the
[38536] human RAB27A gene. Gene 239: 109-116, 1999.

[38537] 5846.

[38538] Wilson, S. M.; Yip, R.; Swing, D. A.; O'Sullivan, T. N.; Zhang,
[38539] Y.; Novak, E. K.; Swank, R. T.; Russell, L. B.; Copeland, N. G.; Jenkins,
[38540] N. A.: A mutation in Rab27a causes the vesicle transport defects
[38541] observed in ashen mice. Proc. Nat. Acad. Sci. 97: 7933-7938, 2000.

[38542] 5847.

[38543] Kallmann, F. J.; Schoenfeld, W. A.; Barrera, S. E.: The genetic
[38544] aspects of primary eunuchoidism. Am. J. Ment. Defic. 48: 203-236,
[38545] 1944.

[38546] 5848.

[38547] De Boulle, K.; Verkerk, A. J. M. H.; Reyniers, E.; Vits, L.;

Hendrickx,
[38548] J.; Van Roy, B.; Van Den Bos, F.; de Graaff, E.; Oostra, B. A.; Willems,
[38549] P. J.: A point mutation in the FMR-1 gene associated with fragile
[38550] X mental retardation. Nature Genet. 3: 31-35, 1993.
[38551] 5849.
[38552] de Vries, B. B. A.; Fryns, J.-P.; Butler, M. G.; Canziani, F.;
[38553] Wesby-van Swaay, E.; van Hemel, J. O.; Oostra, B. A.; Halley, D. J.
[38554] J.; Niermeijer, M. F.: Clinical and molecular studies in fragile
[38555] X patients with a Prader-Willi-like phenotype. J. Med. Genet. 30:
[38556] 761-766, 1993.
[38557] 5850.
[38558] de Vries, B. B. A.; Robinson, H.; Stolte-Dijkstra, I.; Tjon Pian
[38559] Gi, C. V.; Dijkstra, P. F.; van Doorn, J.; Halley, D. J. J.; Oostra,
[38560] B. A.; Turner, G.; Niermeijer, M. F.: General overgrowth in

the fragile
[38561] X syndrome: variability in the phenotypic expression of the FMR1 gene
[38562] mutation. J. Med. Genet. 32: 764-769, 1995.

[38563] 5851.

[38564] Devys, D.; Biancalana, V.; Rousseau, F.; Boue, J.; Mandel, J.

[38565] L.; Oberle, I.: Analysis of full fragile X mutations in fetal tissues
[38566] and monozygotic twins indicate that abnormal methyla- tion and somatic
[38567] heterogeneity are established early in development. Am. J. Med. Genet. 43:
[38568] 208-216, 1992.

[38569] 5852.

[38570] Devys, D.; Lutz, Y.; Rouyer, N.; Bellocq, J.-P.; Mandel, J.-L.

[38571] : The FMR-1 protein is cytoplasmic, most abundant in neurons and appears
[38572] normal in carriers of a fragile X premutation. Nature Genet. 4:
[38573] 335-340, 1993.

[38574] 5853.

[38575] Dombrowski, C.; Levesque, S.; Morel, M. L.; Rouillard, P.; Morgan,
[38576] K.; Rousseau, F.: Premutation and intermediate-size FMR1 alleles
[38577] in 10 572 males from the general population: loss of an AGG interruption
[38578] is a late event in the generation of fragile X syndrome al-
leles. Hum.
[38579] Molec. Genet. 11: 371-378, 2002.

[38580] 5854.

[38581] Dreesen, J. C. F. M.; Geraedts, J. P. M.; Dumoulin, J. C. M.;

[38582] Evers, J. L. H.; Pieters, M. H. E. C.: RS46 (DXS548) geno-
typing of
[38583] reproductive cells: approaching preimplantation testing of the fragile-X
[38584] syndrome. Hum. Genet. 96: 323-329, 1995.

[38585] 5855.

[38586] Dutch-Belgian Fragile X Consortium: Fmr1 knockout mice: a model
[38587] to study fragile X mental retardation. Cell 78: 23-33, 1994.

[38588] 5856.

[38589] Eberhart, D. E.; Malter, H. E.; Feng, Y.; Warren, S. T.: The

[38590] fragile X mental retardation protein is a ribonucleoprotein containing

[38591] both nuclear localization and nuclear export signals. Hum. Molec.

[38592] Genet. 5: 1083-1091, 1996.

[38593] 5857.

[38594] Eichler, E. E.; Nelson, D. L.: Genetic variation and evolutionary

[38595] stability of the FMR1 CGG repeat in six closed human populations. Am.

[38596] J. Med. Genet. 64: 220-225, 1996.

[38597] 5858.

[38598] Eichler, E. E.; Richards, S.; Gibbs, R. A.; Nelson, D. L.: Fine

[38599] structure of the human FMR1 gene. Hum. Molec. Genet. 2: 1147-1153,

[38600] 1993.

[38601] 5859.

[38602] Escalante, J. A.; Grunspun, H.; Frota-Pessoa, O.: Severe

sex-linked
[38603] mental retardation. J. Genet. Hum. 19: 137-140, 1971.

[38604] 5860.

[38605] Faust, C. J.; Verkerk, A. J. M. H.; Wilson, P. J.; Morris, C.

[38606] P.; Hopwood, J. J.; Oostra, B. A.; Herman, G. E.: Genetic mapping
[38607] on the mouse X chromosome of human cDNA clones for the fragile X and
[38608] Hunter syndromes. Genomics 12: 814-817, 1992.

[38609] 5861.

[38610] Feng, Y.; Absher, D.; Eberhart, D. E.; Brown, V.; Malter, H. E.;
[38611] Warren, S. T.: FMRP associates with polyribosomes as an mRNP, and
[38612] the I304N mutation of severe fragile X syndrome abolishes this association. Molec.
[38613] Cell 1: 109-118, 1997.

[38614] 5862.

[38615] Filippi, G.; Arslanian, A.; Dagna-Bricarelli, F.; Pierluigi, M.;

[38616] Grasso, M.; Rinaldi, A.; Rocchi, M.; Siniscalco, M.: Premu-

tation

[38617] for the Martin-Bell syndrome analyzed in a large pedigree segregating

[38618] also for G6PD-deficiency. I. A working hypothesis on the nature of

[38619] the FRAX-mutations. Am. J. Med. Genet. 40: 387-394, 1991.

[38620] 5863.

[38621] Filippi, G.; Rinaldi, A.; Archidiacono, N.; Rocchi, M.; Balazs,

[38622] I.; Siniscalco, M.: Linkage between G6PD and fragile-X syndrome. Am.

[38623] J. Med. Genet. 15: 113-119, 1983.

[38624] 5864.

[38625] Follette, P. J.; Laird, C. D.: Estimating the stability of the

[38626] proposed imprinted state of the fragile-X mutation when transmitted

[38627] by females. Hum. Genet. 88: 335-343, 1992.

[38628] 5865.

[38629] Froster-Iskenius, U.; Schulze, A.; Schwinger, E.: Transmis-sion

[38630] of the marker X syndrome trait by unaffected males: con-

clusions from
[38631] studies of large families. Hum. Genet. 67: 419-427, 1984.

[38632] 5866.

[38633] Fryns, J. P.: The fragile X syndrome: a study of 83 families. Clin.
[38634] Genet. 26: 497-528, 1984.

[38635] 5867.

[38636] Fryns, J. P.: Personal Communication. Leuven, Belgium 5/28/1993.
[38637] 5868.

[38638] Fryns, J. P.; Van Den Berghe, H.: Transmission of fragile(X)(q27)
[38639] from normal male(s). Hum. Genet. 61: 262-263, 1982.

[38640] 5869.

[38641] Fu, Y.-H.; Kuhl, D. P. A.; Pizzuti, A.; Pieretti, M.; Sutcliffe,
[38642] J. S.; Richards, S.; Verkerk, A. J. M. H.; Holden, J. J. A.; Fenwick,
[38643] R. G., Jr.; Warren, S. T.; Oostra, B. A.; Nelson, D. L.; Caskey, C.
[38644] T.: Variation of the CGG repeat at the fragile X site results

in
[38645] genetic instability: resolution of the Sherman paradox. Cell 67:
[38646] 1047-1058, 1991.

[38647] 5870.

[38648] Fulchignoni-Lataud, M. C.; Olchwang, S.; Serre, J. L.: The fragile
[38649] X CGG repeat shows a marked level of instability in hereditary non-polyposis
[38650] colorectal cancer patients. Europ. J. Hum. Genet. 5: 89-93, 1997.
[38651] 5871.

[38652] Gedeon, A. K.; Baker, E.; Robinson, H.; Partington, M. W.; Gross,
[38653] B.; Manca, A.; Korn, B.; Poustka, A.; Yu, S.; Sutherland, G. R.; Mulley,
[38654] J. C.: Fragile X syndrome without CCG amplification has an FMR1 deletion. Nature
[38655] Genet. 1: 341-344, 1992.

[38656] 5872.

[38657] Gerald, P. S.: Personal Communication. Boston, Mass.

8/12/1983.

[38658] 5873.

[38659] Gerald, P. S.: X-linked mental retardation and an X-chromosome

[38660] marker. (Editorial) New Eng. J. Med. 303: 696-697, 1980.

[38661] 5874.

[38662] Giannelli, F.; Morris, A. H.; Garrett, C.; Daker, M.; Thurston,

[38663] C.; Smith, C. A. B.: Genetic heterogeneity of X-linked mental retardation

[38664] with fragile X: association of tight linkage to factor IX and incomplete

[38665] penetrance in males. Ann. Hum. Genet. 51: 107-124, 1987.

[38666] 5875.

[38667] Giraud, F.; Ayme, S.; Mattei, J. F.; Mattei, M. G.: Constitu-tional

[38668] chromosomal breakage. Hum. Genet. 34: 125-136, 1976.

[38669] 5876.

[38670] Glover, T. W.: FUdR induction of the X-chromosome frag-ile site:

[38671] evidence for the mechanism of folic acid and thymidine inhibition. Am.

[38672] J. Hum. Genet. 33: 234-242, 1981.

[38673] 5877.

[38674] Goldman, A.; Krause, A.; Jenkins, T.: Fragile X syndrome occurs

[38675] in the South African black population. (Editorial) S. Afr. Med. J. 87:

[38676] 418-420, 1997.

[38677] 5878.

[38678] Goodman, R. M.; Strauss, S.; Friedman, E.; Chaki, R.: Ovar-ian

[38679] size in the fragile X mental retardation syndrome. Am. J. Med. Genet. 26:

[38680] 17-18, 1987.

[38681] 5879.

[38682] Gould, E. L.; Loesch, D. Z.; Martin, M. J.; Hagerman, R. J.; Armstrong,

[38683] S. M.; Huggins, R. M.: Melatonin profiles and sleep charac-teristics

[38684] in boys with fragile X syndrome: a preliminary study. Am.

J. Med.

[38685] Genet. 95: 307-315, 2000.

[38686] 5880.

[38687] Ellison, J.; Passage, M.; Yu, L.-C.; Yen, P.; Mohandas, T. K.;

[38688] Shapiro, L.: Directed isolation of human genes that escape X inactivation. Somat.

[38689] Cell Molec. Genet. 18: 259-268, 1992.

[38690] 5881.

[38691] Ellison, J. W.; Ramos, C.; Yen, P. H.; Shapiro, L. J.: Structure

[38692] and expression of the human pseudoautosomal gene XE7. Hum. Molec.

[38693] Genet. 1: 691-696, 1992.

[38694] 5882.

[38695] Amar, L. C.; Dandolo, L.; Hanauer, A.; Cook, A. R.; Arnaud, D.;

[38696] Mandel, J. L.; Avner, P.: Conservation and reorganization of loci

[38697] on the mammalian X chromosome: a molecular framework for the identification

[38698] of homologous subchromosomal regions in man and

mouse. Genomics 2:
[38699] 220-230, 1988.

[38700] 5883.

[38701] Chin, L.-S.; Li, L.; Ferreira, A.; Kosik, K. S.; Greengard, P.

[38702] : Impairment of axonal development and of synaptogene-
sis in hippocampal
[38703] neurons of synapsin I-deficient mice. Proc. Nat. Acad. Sci.
92:
[38704] 9230-9234, 1995.

[38705] 5884.

[38706] Klagges, B. R. E.; Heimbeck, G.; Godenschwege, T. A.;
Hofbauer,
[38707] A.; Pflugfelder, G. O.; Reifegerste, R.; Reisch, D.; Schaupp,
M.;
[38708] Buchner, S.; Buchner, E.: Invertebrate synapsins: a single
gene codes
[38709] for several isoforms in Drosophila. J. Neurosci. 16:
3154-3165,
[38710] 1996.

[38711] 5885.

[38712] Li, L.; Chin, L.-S.; Shupliakov, O.; Brodin, L.; Sihra, T. S.;

[38713] Hvalby, O.; Jensen, V.; Zheng, D.; McNamara, J. O.; Greengard, P.;
[38714] Andersen, P.: Impairment of synaptic vesicle clustering and of synaptic
[38715] transmission, and increased seizure propensity, in synapsin I-deficient
[38716] mice. Proc. Nat. Acad. Sci. 92: 9235-9239, 1995.

[38717] 5886.

[38718] Sudhof, T. C.: The structure of the human synapsin I gene and
[38719] protein. J. Biol. Chem. 265: 7849-7852, 1990.

[38720] 5887.

[38721] Banting, G. S.; Pym, B.; Goodfellow, P. N.: Biochemical analysis
[38722] of an antigen produced by both human sex chromosomes. EMBO J. 4:
[38723] 1967-1972, 1985.

[38724] 5888.

[38725] Harada, T.; Harada, C.; Watanabe, M.; Inoue, Y.; Sakagawa, T.;
[38726] Nakayama, N.; Sasaki, S.; Okuyama, S.; Watase, K.; Wada,

K.; Tanaka,

[38727] K.: Functions of the two glutamate transporters GLAST and GLT-1 in

[38728] the retina. Proc. Nat. Acad. Sci. 95: 4663-4666, 1998.

[38729] 5889.

[38730] Agulnik, A. I.; Mitchell, M. J.; Lerner, J. L.; Woods, D. R.; Bishop,

[38731] C. E.: A mouse Y chromosome gene encoded by a region essential for

[38732] spermatogenesis and expression of male-specific minor histocompatibility

[38733] antigens. Hum. Molec. Genet. 3: 873-878, 1994.

[38734] 5890.

[38735] Agulnik, A. I.; Mitchell, M. J.; Mattei, M.-G.; Borsani, G.; Avner,

[38736] P. A.; Lerner, J. L.; Bishop, C. E.: A novel X gene with a widely

[38737] transcribed Y-linked homologue escapes X-inactivation in mouse and

[38738] human. Hum. Molec. Genet. 3: 879-884, 1994.

[38739] 5891.

[38740] Lingenfelter, P. A.; Adler, D. A.; Poslinski, D.; Thomas, S.; Elliott,

[38741] R. W.; Chapman, V. M.; Disteche, C. M.: Escape from X in-

activation

[38742] of Smcx is preceded by silencing during mouse develop-

ment. Nature

[38743] Genet. 18: 212-213, 1998.

[38744] 5892.

[38745] Wu, J.; Ellison, J.; Salido, E.; Yen, P.; Mohandas, T.; Shapiro,

[38746] L. J.: Isolation and characterization of XE169, a novel hu-

man gene

[38747] that escapes X-inactivation. Hum. Molec. Genet. 3:

153-160, 1994.

[38748] 5893.

[38749] Wu, J.; Salido, E. C.; Yen, P. H.; Mohandas, T. K.; Heng, H.

H.

[38750] Q.; Tsui, L.-C.; Park, J.; Chapman, V. M.; Shapiro, L. J.: The

murine

[38751] Xe169 gene escapes X-inactivation like its human homo-

logue. Nature
[38752] Genet. 7: 491-496, 1994.

[38753] 5894.

[38754] Evans, H. J.; Buckton, K. E.; Spowart, G.; Carothers, A. D.:

[38755] Heteromorphic X chromosomes in 46,XX males: evidence for the involvement
[38756] of X-Y interchange. Hum. Genet. 49: 11-31, 1979.

[38757] 5895.

[38758] Hu, X.; Burghes, A. H. M.; Ray, P. N.; Thompson, M. W.; Murphy,
[38759] E. G.; Worton, R. G.: Partial gene duplication in Duchenne and Becker
[38760] muscular dystrophies. J. Med. Genet. 25: 369-376, 1988.

[38761] 5896.

[38762] Bode, H. H.; Crawford, J. D.: Nephrogenic diabetes in-sipidus
[38763] in North America--the Hopewell hypothesis. New Eng. J. Med. 280:
[38764] 750-754, 1969.

[38765] 5897.

[38766] Mills, E. L.; Rholl, K. S.; Quie, P. G.: X-linked inheritance

[38767] in females with chronic granulomatous disease. J. Clin. In-

[38768] vest. 66: 332-340, 1980.

[38769] 5898.

[38770] Muller, C. R.; Migl, B.; Ropers, H.-H.: X-linked steroid sul-

[38771] fatase: evidence for different gene-dosage in males and females.

[38772] Hum. Genet. 54: 197-199, 1980.

[38773] 5899.

[38774] Berger, W.; van de Pol, D.; Warburg, M.; Gal, A.; Bleeker-

[38775] Wagemakers, L.; de Silva, H.; Meindl, A.; Meitinger, T.; Cremers, F.; Rop-

[38776] ers, H.-H.: Mutations in the candidate gene for Norrie disease.

[38777] Hum. Molec. Genet. 1: 461-465, 1992.

[38778] 5900.

[38779] Bleeker-Wagemakers, L. M.; Friedrich, U.; Gal, A.; Wienker, T.

[38780] F.; Warburg, M.; Ropers, H.-H.: Close linkage between Norrie disease,
[38781] a cloned DNA sequence from the proximal short arm, and the centromere
[38782] of the X chromosome. Hum. Genet. 71: 211-214, 1985.

[38783] 5901.

[38784] Chen, Z.-Y.; Battinelli, E. M.; Hendriks, R. W.; Powell, J. F.;

[38785] Middleton-Price, H.; Sims, K. B.; Breakefield, X. O.; Craig, I. W.
[38786] : Norrie disease gene: characterization of deletions and possible
[38787] function. Genomics 16: 533-535, 1993.

[38788] 5902.

[38789] Chen, Z.-Y.; Battinelli, E. M.; Woodruff, G.; Young, I.; Breakefield,
[38790] X. O.; Craig, I. W.: Characterization of a mutation within the NDP
[38791] gene in a family with a manifesting female carrier. Hum. Molec. Genet. 2:
[38792] 1727-1729, 1993.

[38793] 5903.

[38794] Chen, Z.-Y.; Hendriks, R. W.; Jobling, M. A.; Powell, J. F.; Breakefield,
[38795] X. O.; Sims, K. B.; Craig, I. W.: Isolation and characterization
[38796] of a candidate gene for Norrie disease. Nature Genet. 1: 204-208,
[38797] 1992.

[38798] 5904.

[38799] Chen, Z.-Y.; Sims, K. B.; Coleman, M.; Donnai, D.; Monaco, A.;
[38800] Breakefield, X. O.; Davies, K. E.; Craig, I. W.: Characterization
[38801] of a YAC containing part or all of the Norrie disease locus. Hum.
[38802] Molec. Genet. 1: 161-164, 1992.

[38803] 5905.

[38804] Clarke, E.: 'Pseudo-glioma' in both eyes. Trans. Ophthal. Soc.
[38805] U.K. 18: 136-138, 1898.

[38806] 5906.

[38807] Dahlberg-Parrow, R.: Congenital sex-linked pseudoglioma

and grave
[38808] mental deficiency. Acta Ophthal. 34: 250-254, 1956.

[38809] 5907.

[38810] Diergaarde, P. J.; Wieringa, B.; Bleeker-Wagemakers, E. M.; Sims,
[38811] K. B.; Breakefield, X. O.; Ropers, H.-H.: Physical fine-mapping of
[38812] a deletion spanning the Norrie gene. Hum. Genet. 84: 22-26, 1989.
[38813] 5908.

[38814] Donnai, D.; Mountford, R. C.; Read, A. P.: Norrie disease resulting
[38815] from a gene deletion: clinical features and DNA studies. J. Med.
[38816] Genet. 25: 73-78, 1988.

[38817] 5909.

[38818] Duke-Elder, J. R.: Pseudoglioma in children: aspects of clinical
[38819] and pathological diagnosis. Sth. Med. J. 51: 754-759, 1958.
[38820] 5910.

[38821] Forssman, H.: Mental deficiency and pseudoglioma, a syn-
drome
[38822] inherited as an X-linked recessive. Am. J. Ment. Defic. 64:
984-987,
[38823] 1960.

[38824] 5911.

[38825] Fuchs, S.; Kellner, U.; Wedemann, H.; Gal, A.: Missense
mutation
[38826] (Arg121Trp) in the Norrie disease gene associated with X-
linked exudative
[38827] vitreoretinopathy. Hum. Mutat. 6: 257-259, 1995.

[38828] 5912.

[38829] Fuchs, S.; Xu, S. Y.; Caballero, M.; Salcedo, M.; La O, A.;
Wedemann,
[38830] H.; Gal, A.: A missense point mutation (Leu13Arg) of the
Norrie disease
[38831] gene in a large Cuban kindred with Norrie disease. Hum.
Molec. Genet. 3:
[38832] 655-656, 1994.

[38833] 5913.

[38834] Collins, F. A.; Murphy, D. L.; Reiss, A. L.; Sims, K. B.; Lewis,

[38835] J. G.; Freund, L.; Karoum, F.; Zhu, D.; Maumenee, I. H.; Antonarakis,
[38836] S. E.: Clinical, biochemical, and neuropsychiatric evaluation of
[38837] a patient with a contiguous gene syndrome due to a microdeletion Xp11.3
[38838] including the Norrie disease locus and monoamine oxidase (MAOA and
[38839] MAOB) genes. Am. J. Med. Genet. 42: 127-134, 1992.

[38840] 5914.

[38841] Fullwood, P.; Jones, J.; Bundey, S.; Dudgeon, J.; Fielder, A.
[38842] R.; Kilpatrick, M. W.: X linked exudative vitreoretinopathy: clinical
[38843] features and genetic linkage analysis. Brit. J. Ophthal. 77: 168-170,
[38844] 1993.

[38845] 5915.

[38846] Gal, A.; Bleeker-Wagemakers, L.; Wienker, T. F.; Warburg, M.;
[38847] Ropers, H.-H.: Localization of the gene for Norrie disease by linkage

[38848] to the DXS7 locus. (Abstract) Cytogenet. Cell Genet. 40: 633, 1985.

[38849] 5916.

[38850] Gal, A.; Stolzenberger, C.; Wienker, T.; Wieacker, P.; Ropers,

[38851] H.-H.; Friedrich, U.; Bleeker-Wagemakers, L.; Pearson, P.; Warburg,

[38852] M.: Norrie's disease: close linkage with genetic markers from the

[38853] proximal short arm of the X chromosome. Clin. Genet. 27: 282-283,

[38854] 1985.

[38855] 5917.

[38856] Gal, A.; Uhlhaas, S.; Glaser, D.; Grimm, T.: Prenatal exclusion

[38857] of Norrie disease with flanking DNA markers. Am. J. Med. Genet. 31:

[38858] 449-453, 1988.

[38859] 5918.

[38860] Harendra de Silva, D. G.; de Silva, D. B. K.: Norrie's disease

[38861] in an Asian family. Brit. J. Ophthal. 72: 62-64, 1988.

[38862] 5919.

[38863] Bode, H. H.; Miettinen, O. S.: Nephrogenic diabetes in-
sipidus:

[38864] absence of close linkage with Xg. Am. J. Hum. Genet. 22:
221-227,

[38865] 1970.

[38866] 5920.

[38867] Calmont, A.; Reichwald, K.; Ronco, P.; Rossert, J.: Identifi-
cation

[38868] of a short cis-acting element in the human vasopressin
type 2 receptor

[38869] gene which confers high-level expression of a reporter
gene specifically

[38870] in collecting duct cells. Molec. Endocr. 14: 1682-1695,
2000. 1.

[38871] Abelson, H.: Nephrogenic diabetes insipidus. Pediat. Res.
2: 271-282,

[38872] 1968.

[38873] 5921.

[38874] Cannon, J. F.: Diabetes insipidus: clinical and experimental

[38875] studies with consideration of genetic relationships. Arch.

Intern.
[38876] Med. 96: 215-272, 1955.

[38877] 5922.

[38878] Carter, C.; Simpkiss, M. J.: The carrier state in nephrogenic

[38879] diabetes insipidus. Lancet II: 1069-1073, 1956.

[38880] 5923.

[38881] Cheong, H. I.; Park, H. W.; Ha, I. S.; Moon, H. N.; Choi, Y.;

[38882] Ko, K. W.; Jun, J. K.: Six novel mutations in the vasopressin V2

[38883] receptor gene causing nephrogenic diabetes insipidus. Nephron 75:

[38884] 431-437, 1997.

[38885] 5924.

[38886] Crawford, J. D.; Kennedy, G. C.: Chlorothiazid in diabetes insipidus. Nature 183:

[38887] 891-892, 1959.

[38888] 5925.

[38889] Friedman, E.; Bale, A. E.; Carson, E.; Boson, W. L.; Norden-skjold,

[38890] M.; Ritzen, M.; Ferriera, P. C.; Jammal, A.; De Marco, L.: Nephrogenic
[38891] diabetes insipidus: an X chromosome-linked dominant inheritance pattern
[38892] with a vasopressin type 2 receptor gene that is structurally normal. Proc.
[38893] Nat. Acad. Sci. 91: 8457-8461, 1994.

[38894] 5926.

[38895] Holtzman, E. J.; Harris, H. W., Jr.; Kolakowski, L. F., Jr.; Guay-Woodford,
[38896] L. M.; Botelho, B.; Ausiello, D. A.: A molecular defect in the vasopressin
[38897] V2-receptor gene causing nephrogenic diabetes insipidus. New Eng.
[38898] J. Med. 328: 1534-1537, 1993.

[38899] 5927.

[38900] Holtzman, E. J.; Kolakowski, L. F., Jr.; O'Brien, D.; Crawford,
[38901] J. D.; Ausiello, D. A.: A null mutation in the vasopressin V2 receptor
[38902] gene (AVPR2) associated with nephrogenic diabetes insipidus in the

[38903] Hopewell kindred. Hum. Molec. Genet. 2: 1201-1204, 1993.

[38904] 5928.

[38905] Inaba, S.; Hatakeyama, H.; Taniguchi, N.; Miyamori, I.: The property

[38906] of a novel V2 receptor mutant in a patient with nephrogenic diabetes

[38907] insipidus. J. Clin. Endocr. Metab. 86: 381-385, 2001.

[38908] 5929.

[38909] Jans, D. A.; van Oost, B. A.; Ropers, H. H.; Fahrenholz, F.:

[38910] Derivatives of somatic cell hybrids which carry the human gene locus

[38911] for nephrogenic diabetes insipidus (NDI) express functional vasopressin

[38912] renal V2-type receptors. J. Biol. Chem. 265: 15379-15382, 1990.

[38913] 5930.

[38914] Kambouris, M.; Dlouhy, S. R.; Trofatter, J. A.; Conneally, P.

[38915] M.; Hodes, M. E.: Localization of the gene for X-linked nephrogenic

[38916] diabetes insipidus to Xq28. Am. J. Med. Genet. 29:

239-246, 1988.

[38917] 5931.

[38918] Kambouris, M.; Hodes, M. E.; Conneally, P. M.; Trofatter, J.;

[38919] Dlouhy, S. R.: Nephrogenic diabetes insipidus linked to Xq28. (Abstract) Cytogenet.

[38920] Cell Genet. 46: 636 only, 1987.

[38921] 5932.

[38922] Knoers, N.; van den Heyden, H.; van Oost, B.; Monnens, L.; Willems,

[38923] J.; Ropers, H.: Tight linkage between nephrogenic diabetes insipidus

[38924] and DXS52. (Abstract) Cytogenet. Cell Genet. 46: 640 only, 1987.

[38925] 5933.

[38926] Knoers, N.; van den Heyden, H.; van Oost, B. A.; Monnens, L.;

[38927] Willems, J.; Ropers, H. H.: Three-point linkage analysis using multiple

[38928] DNA polymorphic markers in families with X-linked nephrogenic diabetes

[38929] insipidus. Genomics 4: 434-437, 1989.

[38930] 5934.

[38931] Knoers, N.; van der Heyden, H.; van Oost, B. A.; Ropers, H. H.;

[38932] Monnens, L.; Willems, J.: Nephrogenic diabetes insipidus: close linkage

[38933] with markers from the distal long arm of the human X chromosome. Hum.

[38934] Genet. 80: 31-38, 1988.

[38935] 5935.

[38936] Kobrinsky, N. L.; Doyle, J. J.; Israels, E. D.; Winter, J. S.

[38937] D.; Cheang, M. S.; Walker, R. D.; Bishop, A. J.: Absent factor VIII

[38938] response to synthetic vasopressin analogue (DDAVP) in nephrogenic

[38939] diabetes insipidus. Lancet I: 1293-1294, 1985.

[38940] 5936.

[38941] Libber, S.; Harrison, H.; Spector, D.: Treatment of nephrogenic

[38942] diabetes insipidus with prostaglandin synthesis inhibitors. J. Pediat. 108:

[38943] 305-311, 1986.

[38944] 5937.

[38945] Lolait, S. J.; O'Carroll, A.-M.; McBride, O. W.; Konig, M.; Morel,

[38946] A.; Brownstein, M. J.: Cloning and characterization of a va-

sopressin

[38947] V2 receptor and possible link to nephrogenic diabetes in-

sipidus. Nature 357:

[38948] 336-339, 1992.

[38949] 5938.

[38950] Mattei, M.-G.; Matterson, J.; Chen, J. W.; Williams, M. A.; Fukuda,

[38951] M.: Two human lysosomal membrane glycoproteins, h-

lamp-1 and h-lamp-2,

[38952] are encoded by genes localized to chromosome 13q34 and chromosome

[38953] Xq24-25, respectively. J. Biol. Chem. 265: 7548-7551, 1990.

[38954] 5939.

[38955] Schleutker, J.; Haataja, L.; Renlund, M.; Puhakka, L.; Vi-

itala,

[38956] J.; Peltonen, L.; Aula, P.: Confirmation of the chromosomal localization

[38957] of human lamp genes and their exclusion as candidate genes for Salla

[38958] disease. Hum. Genet. 88: 95-97, 1991.

[38959] 5940.

[38960] Andreae, S.; Piras, F.; Burdin, N.; Triebel, F.: Maturation and

[38961] activation of dendritic cells induced by lymphocyte activation gene-3

[38962] (CD223). J. Immun. 168: 3874-3880, 2002.

[38963] 5941.

[38964] Triebel, F.; Jitsukawa, S.; Baixeras, E.; Roman-Roman, S.; Genevee,

[38965] C.; Viegas-Pequignot, E.; Hercend, T.: LAG-3, a novel lymphocyte

[38966] activation gene closely related to CD4. J. Exp. Med. 171: 1393-1405,

[38967] 1990.

[38968] 5942.

[38969] Bierer, B. E.; Nishimura, Y.; Burakoff, S. J.; Smith, B. R.: Phenotypic

[38970] and functional characterization of human cytolytic T cells

lacking
[38971] expression of CD5. J. Clin. Invest. 81: 1390-1397, 1988.

[38972] 5943.

[38973] Gershon, R. K.: Personal Communication. New Haven, Conn. 12/4/1978.

[38974] 5944.

[38975] Hecht, B. K.; Kipps, T.; Johnston, N. K.; Cannizzaro, L. A.: Leu
[38976] 1 (CD5) cell surface antigen mapped to 11q13 by in situ hybridization.
[38977] (Abstract) Cytogenet. Cell Genet. 51: 1012 only, 1989.

[38978] 5945.

[38979] Jones, N. H.; Clabby, M. L.; Dialynas, D. P.; Huang, H.-J. S.;

[38980] Herzenberg, L. A.; Strominger, J. L.: Isolation of comple-
mentary
[38981] DNA clones encoding the human lymphocyte glycoprotein T1/Leu-1. Nature 323:
[38982] 346-349, 1986.

[38983] 5946.

[38984] LeClair, K. P.; Rabin, M.; Nesbitt, M. N.; Pravtcheva, D.;

Ruddle,
[38985] F. H.; Palfree, R. G. E.; Bothwell, A.: Murine Ly-6 multigene family
[38986] is located on chromosome 15. Proc. Nat. Acad. Sci. 84: 1638-1642,
[38987] 1987.

[38988] 5947.

[38989] Ravetch, J. V.; Lanier, L. L.: Immune inhibitory receptors. Science 290:
[38990] 84-89, 2000.

[38991] 5948.

[38992] van Dongen, J. J. M.; Wolvers-Tettero, I. L. M.; Versnel, M. A.;
[38993] Westerveld, A.; Geurts van Kessel, A. H. M.: Assignment of the genes
[38994] coding for the T-cell antigens CD7 (Tp41), CD5 (T1) and CD4 (T4) to
[38995] human chromosome 17, 11, 12 respectively. (Abstract) Cytogenet. Cell
[38996] Genet. 40: 767 only, 1985.

[38997] 5949.

[38998] Fischer, A.; Durandy, A.; Sterkers, G.; Griscelli, C.: Role of

[38999] the LFA-1 molecule in cellular interactions required for antibody

[39000] production in humans. J. Immun. 136: 3198-3203, 1986.

[39001] 5950.

[39002] Marlin, S. D.; Morton, C. C.; Anderson, D. C.; Springer, T. A.

[39003] : LFA-1 immunodeficiency disease: definition of the genetic defect

[39004] and chromosomal mapping of alpha and beta subunits of the lymphocyte

[39005] function-associated antigen 1 (LFA-1) by complementation in hybrid

[39006] cells. J. Exp. Med. 164: 855-867, 1986.

[39007] 5951.

[39008] Annunen, S.; Korkko, J.; Czarny, M.; Warman, M. L.; Brunner, H.

[39009] G.; Kaariainen, H.; Mulliken, J. B.; Tranebjaerg, L.; Brooks, D. G.;

[39010] Cox, G. F.; Cruysberg, J. R.; Curtis, M. A.; and 13 others: Splicing

[39011] mutations of 54-bp exons in the COL11A1 gene cause Marshall syndrome,

[39012] but other mutations cause overlapping Marshall/Stickler phenotypes. Am.

[39013] J. Hum. Genet. 65: 974-983, 1999.

[39014] 5952.

[39015] Shanske, A. L.; Bogdanow, A.; Shprintzen, R. J.; Marion, R. W.

[39016] : The Marshall syndrome: report of a new family and re-view of the

[39017] literature. Am. J. Med. Genet. 70: 52-57, 1997.

[39018] 5953.

[39019] Warman, M. L.; Tiller, G. E.; Griffith, A. J.: Reply to Shanske

[39020] et al. (Letter) Am. J. Hum. Genet. 63: 1559-1561, 1998.

[39021] 5954.

[39022] Futscher, B. W.; Oshiro, M. M.; Wozniak, R. J.; Holtan, N.; Hanigan,

[39023] C. L.; Duan, H.; Domann, F. E.: Role for DNA methylation in the control

[39024] of cell type-specific maspin expression. Nature Genet. 31: 175-179,

[39025] 2002.

[39026] 5955.

[39027] Ngamkitidechakul, C.; Burke, J. M.; O'Brien, W. J.; Twining, S.

[39028] S.: Maspin: synthesis by human cornea and regulation of in vitro

[39029] stromal cell adhesion to extracellular matrix. Invest. Ophthal. Vis.

[39030] Sci. 42: 3135-3141, 2001.

[39031] 5956.

[39032] Zou, Z.; Anisowicz, A.; Hendrix, M. J. C.; Thor, A.; Neveu, M.;

[39033] Sheng, S.; Rafidi, K.; Seftor, E.; Sager, R.: Maspin, a serpin with

[39034] tumor-suppressing activity in human mammary epithelial cells. Science 263:

[39035] 526-529, 1994.

[39036] 5957.

[39037] Johnson, T. L.; Sakaguchi, A. Y.; Lalley, P. A.; Leach, R. J.:

[39038] Chromosomal assignment in mouse of matrix GLA protein and bone GLA

[39039] protein genes. Genomics 11: 770-772, 1991.

[39040] 5958.

[39041] Blackwood, E.; Eisenman, R. N.: Max: a helix-loop-helix zipper

[39042] protein that forms a sequence-specific DNA-binding complex with Myc. Science 251:

[39043] 1211-1217, 1991.

[39044] 5959.

[39045] Eisenman, R. N.: Personal Communication. Seattle, Wash. 7/27/1994.

[39046] 5960.

[39047] Gilladoga, A. D.; Edelhoff, S.; Blackwood, E. M.; Eisenman, R.

[39048] N.; Disteche, C. M.: Mapping of MAX to human chromosome 14 and mouse

[39049] chromosome 12 by in situ hybridization. Oncogene 7: 1249-1251, 1992.

[39050] 5961.

[39051] Loeffen, J. L. C. M.; Triepels, R. H.; van den Heuvel, L. P.; Schuelke,

[39052] M.; Buskens, C. A. F.; Smeets, R. J. P.; Trijbels, J. M. F.;

Smeitink,

[39053] J. A. M.: cDNA of eight nuclear encoded subunits of NADH:ubiquinone

[39054] oxidoreductase: human complex I cDNA characterization completed. Biochem.

[39055] Biophys. Res. Commun. 253: 415-422, 1998.

[39056] 5962.

[39057] Bulavin, D. V.; Higashimoto, Y.; Popoff, I. J.; Gaarde, W. A.;

[39058] Basrur, V.; Potapova, O.; Appella, E.; Fornace, A. J., Jr.: Initiation

[39059] of a G2/M checkpoint after ultraviolet radiation requires p38 kinase. Nature 411:

[39060] 102-107, 2001.

[39061] 5963.

[39062] Gould, K. L.; Moreno, S.; Tonks, N. K.; Nurse, P.: Complementation

[39063] of the mitotic activator, p80(cdc25), by a human protein-tyrosine

[39064] phosphatase. Science 250: 1573-1576, 1990.

[39065] 5964.

[39066] Moreno, S.; Nurse, P.: Clues to action of cdc25 protein.

(Letter) Nature 351:

[39067] 194 only, 1991.

[39068] 5965.

[39069] Raj, K.; Ogston, P.; Beard, P.: Virus-mediated killing of cells

[39070] that lack p53 activity. Nature 412: 914-917, 2001. Note: Addendum:

[39071] Nature 416: 202 only, 2002.

[39072] 5966.

[39073] Sadhu, K.; Reed, S. I.; Richardson, H.; Russell, P.: Human homolog

[39074] of fission yeast cdc25 mitotic inducer is predominantly expressed

[39075] in G(2). Proc. Nat. Acad. Sci. 87: 5139-5143, 1990.

[39076] 5967.

[39077] Strausfeld, U.; Labbe, J. C.; Fesquet, D.; Cavadore, J. C.; Picard,

[39078] A.; Sadhu, K.; Russell, P.; Doree, M.: Dephosphorylation and activation

[39079] of a p34(cdc2)/cyclin B complex in vitro by human CDC25 protein. Nature 351:

[39080] 242-245, 1991.

[39081] 5968.

[39082] Taviaux, S. A.; Demaille, J. G.: Localization of human cell cycle

[39083] regulatory genes CDC25C to 5q31 and WEE1 to 11p15.3-11p.15.1 by fluorescence

[39084] in situ hybridization. Genomics 15: 194-196, 1993.

[39085] 5969.

[39086] Willman, C. L.; Sever, C. E.; Pallavicini, M. G.; Harada, H.; Tanaka,

[39087] N.; Slovak, M. L.; Yamamoto, H.; Harada, K.; Meeker, T. C.; List,

[39088] A. F.; Taniguchi, T.: Deletion of IRF-1, mapping to chromosome 5q31.1,

[39089] in human leukemia and preleukemic myelodysplasia. Science 259: 968-971,

[39090] 1993.

[39091] 5970.

[39092] Yazawa, I.; Nukina, N.; Hashida, H.; Goto, J.; Yamada, M.; Kanazawa,

[39093] I.: Abnormal gene product identified in hereditary denta-

torubral-pallidoluysian
[39094] atrophy (DRPLA) brain. Nature Genet. 10: 99-103, 1995.

[39095] 5971.

[39096] Board, P. G.; Coggan, M.: Polymorphism of the A subunit of coagulation
[39097] factor XIII in the Pacific region: description of new phenotypes. Hum.
[39098] Genet. 59: 135-136, 1981.

[39099] 5972.

[39100] Beutler, E.; West, C.; Beutler, B.: Electrophoretic polymorphism
[39101] of glutathione peroxidase. Ann. Hum. Genet. 38: 163-169, 1974.
[39102] 5973.

[39103] Belin, V.; Cusin, V.; Viot, G.; Girlich, D.; Toutain, A.; Moncla,
[39104] A.; Vekemans, M.; Le Merrer, M.; Munnich, A.; Cormier-Daire, V.:
[39105] SHOX mutations in dyschondrosteosis (Leri-Weill syndrome). Nature
[39106] Genet. 19: 67-69, 1998.

[39107] 5974.

[39108] Fujimoto, M.; Kantaputra, P. N.; Ikegawa, S.; Fukushima, Y.; Sonta,

[39109] S.; Matsuo, M.; Ishida, T.; Matsumoto, T.; Kondo, S.; Tomita, H.;

[39110] Deng, H.-X.; D'urso, M.; Rinaldi, M. M.; Ventruto, V.; Takagi, T.;

[39111] Nakamura, Y.; Niikawa, N.: The gene for mesomelic dysplasia Kantaputra

[39112] type is mapped to chromosome 2q24-q32. J. Hum. Genet. 43: 32-36,

[39113] 1998.

[39114] 5975.

[39115] Grigelioniene, G.; Eklof, O.; Ivarsson, S. A.; Westphal, O.; Neumeyer,

[39116] L.; Kedra, D.; Dumanski, J.; Hagenas, L.: Mutations in short stature

[39117] homeobox containing gene (SHOX) in dyschondrosteosis but not in hypochondroplasia. Hum.

[39118] Genet. 107: 145-149, 2000.

[39119] 5976.

[39120] Huber, C.; Cusin, V.; Le Merrer, M.; Mathieu, M.; Sulmont, V.;
[39121] Dagoneau, N.; Munnich, A.; Cormier-Daire, V.: SHOX point mutations
[39122] in dyschondrosteosis. J. Med. Genet. 38: 323 only, 2001.

[39123] 5977.

[39124] Ross, J. L.; Scott, C., Jr.; Marttila, P.; Kowal, K.; Nass, A.;
[39125] Papenhausen, P.; Abboudi, J.; Osterman, L.; Kushner, H.; Carter, P.;
[39126] Ezaki, M.; Elder, F.; Wei, F.; Chen, H.; Zinn, A. R.: Phenotypes
[39127] associated with SHOX deficiency. J. Clin. Endocr. Metab. 86: 5674-5680,
[39128] 2001.

[39129] 5978.

[39130] Shears, D. J.; Vassal, H. J.; Goodman, F. R.; Palmer, R. W.; Reardon,
[39131] W.; Superti-Furga, A.; Scambler, P. J.; Winter, R. M.: Mutation and
[39132] deletion of the pseudoautosomal gene SHOX cause Leri-Weill dyschondrosteosis. Nature

[39133] Genet. 19: 70-73, 1998.

[39134] 5979.

[39135] Ventruto, V.; Pisciotta, R.; Renda, S.; Fosta, B.; Rinaldi, M.

[39136] M.; Stabile, M.; Cavaliere, M. L.; Esposito, M.: Multiple skeletal

[39137] familial abnormalities associated with balanced reciprocal translocation

[39138] 2;8(q32;p13). Am. J. Med. Genet. 16: 589-594, 1983.

[39139] 5980.

[39140] Agarwal, S. K.; Guru, S. C.; Heppner, C.; Erdos, M. R.; Collins,

[39141] R. M.; Park, S. Y.; Saggar, S.; Chandrasekharappa, S. C.; Collins,

[39142] F. S.; Spiegel, A. M.; Marx, S. J.; Burns, A. L.: Menin inter-acts

[39143] with the AP1 transcription factor JunD and represses JunD-activated

[39144] transcription. Cell 96: 143-152, 1999.

[39145] 5981.

[39146] Agarwal, S. K.; Kester, M. B.; Debelenko, L. V.; Heppner, C.; Emmert-Buck,

[39147] M. R.; Skarulis, M. C.; Doppman, J. L.; Kim, Y. S.; Lubensky, I. A.;

[39148] Zhuang, Z.; Green, J. S.; Guru, S. C.; Manickam, P.; Olufemi, S.-E.;

[39149] Liotta, L. A.; Chandrasekharappa, S. C.; Collins, F. S.; Spiegel,

[39150] A. M.; Burns, A. L.; Marx, S. J.: Germline mutations of the MEN1

[39151] gene in familial multiple endocrine neoplasia type 1 and related states. Hum.

[39152] Molec. Genet. 7: 1169-1175, 1997.

[39153] 5982.

[39154] Agarwal, S. K.; Kester, M. B.; Debelenko, L. V.; Heppner, C.; Emmert-Buck,

[39155] M. R.; Skarulis, M. C.; Doppman, J. L.; Kim, Y. S.; Lubensky, I. A.;

[39156] Zhuang, Z.; Green, J. S.; Guru, S. C.; Manickam, P.; Olufemi, S.-E.;

[39157] Liotta, L. A.; Chandrasekharappa, S. C.; Collins, F. S.; Spiegel,

[39158] A. M.; Burns, A. L.; Marx, S. J.: Germline mutations of the MEN1

[39159] gene in familial multiple endocrine neoplasia type 1 and related states. Hum.

[39160] Molec. Genet. 6: 1169-1175, 1997.

[39161] 5983.

[39162] Bahn, R. S.; Scheithauer, B. W.; van Heerden, J. A.; Laws, E. R.,

[39163] Jr.; Horvath, E.; Gharib, H.: Nonidentical expressions of multiple

[39164] endocrine neoplasia, type I, in identical twins. Mayo Clin. Proc. 61:

[39165] 689-696, 1986.

[39166] 5984.

[39167] Bale, A. E.; Friedman, E.; Sakaguchi, K.; Nakamura, Y.; McBride,

[39168] O. W.; Spiegel, A. M.; Aurbach, G. D.; Marx, S. J.: Allelic loss

[39169] in hereditary and sporadic tumors associated with MEN1. (Abstract) Am.

[39170] J. Hum. Genet. 45 (suppl.): A15, 1989.

[39171] 5985.

[39172] Bale, S. J.; Bale, A. E.; Marx, S. J.: Mapping of multiple en-

docrine
[39173] neoplasia type 1 with markers on chromosome 11q. (Abstract) Cytogenet.
[39174] Cell Genet. 51: 956, 1989.

[39175] 5986.

[39176] Bale, S. J.; Bale, A. E.; Stewart, K.; Dachowski, L.; McBride,

[39177] O. W.; Glaser, T.; Green, J. E., III; Mulvihill, J. J.; Brandi, M.

[39178] L.; Sakaguchi, K.; Aurbach, G. D.; Marx, S. J.: Linkage anal-
ysis
[39179] of multiple endocrine neoplasia type 1 with INT2 and other markers
[39180] on chromosome 11. Genomics 4: 320-322, 1989.

[39181] 5987.

[39182] Bale, S. J.; Marx, S. J.; Langfield, D.; Bale, A. E.; Mulvihill,

[39183] J.; Green, J., III: Multiple endocrine neoplasia type 1 (MEN1): genetic
[39184] linkage studies. (Abstract) Cytogenet. Cell Genet. 46: 575, 1987.
[39185] 5988.

[39186] Ballard, H. S.; Frame, B.; Hartsock, R. J.: Familial multiple

[39187] endocrine adenoma-peptic ulcer complex. Medicine 43: 481-516, 1964.

[39188] 5989.

[39189] Bassett, J. H. D.; Forbes, S. A.; Pannett, A. A. J.; Lloyd, S.

[39190] E.; Christie, P. T.; Wooding, C.; Harding, B.; Besser, G. M.; Edwards,

[39191] C. R.; Monson, J. P.; Sampson, J.; Wass, J. A. H.; Wheeler, M. H.;

[39192] Thakker, R. V.: Characterization of mutations in patients with multiple

[39193] endocrine neoplasia type 1. Am. J. Hum. Genet. 62: 232-244, 1998.

[39194] 5990.

[39195] Bear, J. C.; Briones-Urbina, R.; Fahey, J. F.; Farid, N. R.:

[39196] Variant multiple endocrine neoplasia I (MEN I-Burin): fur- ther studies

[39197] and non-linkage to HLA. Hum. Hered. 35: 15-20, 1985.

[39198] 5991.

[39199] Betts, J. B.; O'Malley, B. P.; Rosenthal, F. D.: Hyperparathy- roidism:

[39200] a prerequisite for Zollinger-Ellison syndrome in multiple

endocrine
[39201] adenomatosis type 1--report of a further family and a re-
view of the
[39202] literature. Quart. J. Med. 49: 69-76, 1980.

[39203] 5992.

[39204] Boni, R.; Vortmeyer, A. O.; Pack, S.; Park, W.-S.; Burg, G.;
Hofbauer,
[39205] G.; Darling, T.; Liotta, L.; Zhuang, Z.: Somatic mutations of
the
[39206] MEN1 tumor suppressor gene detected in sporadic an-
giofibromas. (Letter) J.
[39207] Invest. Derm. 111: 539-540, 1998.

[39208] 5993.

[39209] Bordi, C.; Corleto, V. D.; Azzoni, C.; Pizzi, S.; Ferraro, G.;

[39210] Gibril, F.; Delle Fave, G.; Jensen, R. T.: The antral mucosa
as a
[39211] new site for endocrine tumors in multiple endocrine neo-
plasia type
[39212] 1 and Zollinger-Ellison syndromes. J. Clin. Endocr. Metab.
86: 2236-2242,
[39213] 2001.

[39214] 5994.

[39215] Kinoshita, A.; Saito, T.; Tomita, H.; Makita, Y.; Yoshida, K.;

[39216] Ghadami, M.; Yamada, K.; Kondo, S.; Ikegawa, S.;
Nishimura, G.; Fukushima,

[39217] Y.; Nakagomi, T.; Saito, H.; Sugimoto, T.; Kamegaya, M.;
Hisa, K.;

[39218] Murray, J. C.; Taniguchi, N.; Niikawa, N.; Yoshiura, K.: Do-
main-specific

[39219] mutations in TGFB1 result in Camurati-Engelmann dis-
ease. Nature Genet. 26:

[39220] 19-20, 2000.

[39221] 5995.

[39222] Joyner, A. L.; Herrup, K.; Auerbach, B. A.; Davis, C. A.;
Rossant,

[39223] J.: Subtle cerebellar phenotype in mice homozygous for a
targeted

[39224] deletion of the En-2 homeobox. Science 251: 1239-1243,
1991.

[39225] 5996.

[39226] Joyner, A. L.; Skarnes, W. C.; Rossant, J.: Production of a
mutation

[39227] in mouse En-2 gene by homologous recombination in embryonic stem cells. Nature 338:
[39228] 153-156, 1989.

[39229] 5997.

[39230] Poole, S. J.; Law, M. L.; Kao, F.-T.; Lau, Y.-F.: Isolation and

[39231] chromosomal localization of the human En-2 gene. Genomics 4: 225-231,
[39232] 1989.

[39233] 5998.

[39234] Board, P. G.; Reid, M.; Serjeantson, S.: The gene for coagulation
[39235] factor XIII A subunit (F13A) is distal to HLA on chromosome 6. Hum.
[39236] Genet. 67: 406-408, 1984.

[39237] 5999.

[39238] Board, P. G.; Webb, G. C.; McKee, J.; Ichinose, A.: Localization
[39239] of the coagulation factor XIII A subunit gene (F13A) to chromosome
[39240] bands 6p24-p25. Cytogenet. Cell Genet. 48: 25-27, 1988.

[39241] 6000.

[39242] Butten, A. F. H.: Congenital deficiency of factor XIII (fibrin-stabilizing

[39243] factor): report of a case and review of the literature. Am. J. Med. 43:

[39244] 751-761, 1967.

[39245] 6001.

[39246] Castle, S.; Board, P. G.; Anderson, R. A. M.: Genetic het-erogeneity

[39247] of factor XIII deficiency: first description of unstable A subunits. Brit.

[39248] J. Haemat. 48: 337-342, 1981.

[39249] 6002.

[39250] Catto, A. J.; Kohler, H. P.; Bannan, S.; Stickland, M.; Carter,

[39251] A.; Grant, P. J.: Factor XIII val34-to-leu: a novel associa-tion with

[39252] primary intracerebral hemorrhage. Stroke 29: 813-816, 1998.

[39253] 6003.

[39254] Coggan, M.; Baker, R.; Miloszewski, K.; Woodfield, G.; Board,

[39255] P.: Mutations causing coagulation factor XIII subunit A deficiency:
[39256] characterization of the mutant proteins after expression in yeast. Blood 85:
[39257] 2455-2460, 1995.

[39258] 6004.

[39259] Duckert, F.: Factor XIII Deficiency. Proc. 10th Int. Cong. Soc.
[39260] Haematol., Stockholm. (pub.) 1964.

[39261] 6005.

[39262] Duckert, F.; Jung, E.; Shmerling, D. H.: A hitherto undescribed
[39263] congenital haemorrhagic diathesis probably due to fibrin stabilizing
[39264] factor deficiency. Thromb. Diath. Haemorrh. 5: 179-186, 1960.
[39265] 6006.

[39266] Eiberg, H.; Mohr, J.; Nielsen, L. S.: Linkage relationships of
[39267] human coagulation factor XIIIB. (Abstract) Cytogenet. Cell Genet. 37:
[39268] 463 only, 1984.

[39269] 6007.

[39270] Eiberg, H.; Nielsen, L. S.; Mohr, J.: Confirmation of F13A assignment

[39271] and sequence information concerning F13A-HLA-GLO. Clin. Genet. 26:

[39272] 385-388, 1984.

[39273] 6008.

[39274] Fisher, S.; Rikover, M.; Naor, S.: Factor XIII deficiency with

[39275] severe hemorrhagic diathesis. Blood 28: 34-39, 1966.

[39276] 6009.

[39277] Forman, W. B.; Byer, R.; Hadady, M.; Krill, C.; Lubin, A.: Congenital

[39278] fibrin stabilizing factor deficiency (FSF, XIII): evidence for dys-FSF.

[39279] (Abstract) Blood 50 (suppl. 1): 266 only, 1977.

[39280] 6010.

[39281] Francis, J. L.; Todd, P. J.: Factor XIII deficiency: a family

[39282] study by measurement of factor XIII subunits A and S. Acta Haemat. 62:

[39283] 167-172, 1979.

[39284] 6011.

[39285] Franco, R. F.; Reitsma, P. H.; Lourenco, D.; Maffei, F. H.; Morelli,

[39286] V.; Tavella, M. H.; Araujo, A. G.; Piccinato, C. E.; Zago, M. A.:

[39287] Factor XIII val34-to-leu is a genetic factor involved in the aetiology

[39288] of venous thrombosis. Thromb. Haemost. 81: 676-679, 1999.

[39289] 6012.

[39290] Fried, K.; Kaufman, S.; Beer, S.: Factor XIII deficiency. Clin.

[39291] Genet. 20: 455-457, 1981.

[39292] 6013.

[39293] Frydman, M.; Bonne-Tamir, B.; Braude, E.; Zamir, R.; Creter, D.

[39294] : Male fertility in factor XIII deficiency. Fertil. Steril. 45:

[39295] 729-731, 1986.

[39296] 6014.

[39297] Frydman, M.; Farrer, L. A.; Bonne-Tamir, B.; Zamir, R.: The locus

[39298] for factor XIII deficiency linked to the HLA region on chromosome

[39299] 6p. (Abstract) Am. J. Hum. Genet. 37: A53 only, 1985.

[39300] 6015.

[39301] Girolami, A.; Burrul, A.; Sticchi, A.: Congenital deficiency

[39302] of factor XIII with normal subunits S and lack of subunit A: report

[39303] of a new family. Acta Haemat. 58: 17-26, 1977.

[39304] 6016.

[39305] Girolami, A.; Cappellato, M. G.; Vicarioto, M. A.: Congenital

[39306] factor XIII deficiency: type I and type II disease. (Letter) Brit.

[39307] J. Haemat. 60: 375-377, 1985.

[39308] 6017.

[39309] Gomez Garcia, E. B.; Poort, S. R.; Stibbe, J.; Sturk, A.; Schaap,

[39310] M. C. L.; Kappers, M.; Bertina, R. M.: Two novel and one recurrent

[39311] missense mutation in the factor XIII A gene in two Dutch patients

[39312] with factor XIII deficiency. Brit. J. Haemat. 112: 513-518, 2001.

[39313] 6018.

[39314] Graham, J. B.; Edgell, C.-J. S.; Fleming, H.; Namboodiri, K. K.;

[39315] Keats, B. J. B.; Elston, R. C.: Coagulation factor XIII: a use-ful

[39316] polymorphic genetic marker. Hum. Genet. 67: 132-135, 1984.

[39317] 6019.

[39318] Grundmann, U.; Amann, E.; Zettlmeissl, G.; Kupper, H. A.: Characterization

[39319] of cDNA coding for human factor XIIIa. Proc. Nat. Acad. Sci. 83:

[39320] 8024-8028, 1986.

[39321] 6020.

[39322] Ichinose, A.; Davie, E. W.: Characterization of the gene for

[39323] the a subunit of human factor XIII (plasma transglutami-nase), a blood

[39324] coagulation factor. Proc. Nat. Acad. Sci. 85: 5829-5833, 1988.

[39325] 6021.

[39326] Ichinose, A.; Hendrickson, L. E.; Fujikawa, K.; Davie, E. W.:

[39327] Amino acid sequence of the a subunit of human factor XIII. Biochemistry 25:

[39328] 6900-6906, 1986.

[39329] 6022.

[39330] Ichinose, A.; Izumi, T.; Hashiguchi, T.: The normal and ab-normal

[39331] genes of the a and b subunits in coagulation factor XIII. Semin.

[39332] Thromb. Hemost. 22: 385-391, 1996.

[39333] 6023.

[39334] Ichinose, A.; Souri, M.; Izumi, T.; Takahashi, N.: Molecular

[39335] and genetic mechanisms of factor XIII A subunit defi-ciency. Semin.

[39336] Thromb. Hemost. 26: 5-10, 2000.

[39337] 6024.

[39338] Ikkala, E.; Nevanlinna, H. R.: Congenital deficiency of fibrin

[39339] stabilizing factor. Thromb. Diath. Haemorrh. 7: 567-571,

1962.

[39340] 6025.

[39341] Israels, E. D.; Paraskevas, F.; Israels, L. G.: Immunological

[39342] studies of coagulation factor XIII. J. Clin. Invest. 52:
2398-2403,

[39343] 1973.

[39344] 6026.

[39345] Kamboh, M. I.; Ferrell, R. E.: Three FXIIIA gene loci? (Letter)
Hum.

[39346] Genet. 84: 102 only, 1989.

[39347] 6027.

[39348] Kamura, T.; Okamura, T.; Murakawa, M.; Tsuda, H.;
Teshima, T.;

[39349] Shibuya, T.; Harada, M.; Niho, Y.: Deficiency of coagula-
tion factor

[39350] XIII A subunit caused by the dinucleotide deletion at the
5-prime

[39351] end of exon III. J. Clin. Invest. 90: 315-319, 1992.

[39352] 6028.

[39353] Kangsadalampai, S.; Chelvanayagam, G.; Baker, R.; Tiede-

mann, K.;
[39354] Kuperan, P.; Board, P. G.: Identification and characterization of
[39355] two missense mutations causing factor XIIIA deficiency. Brit. J.
[39356] Haemat. 104: 37-43, 1999.

[39357] 6029.

[39358] Kera, Y.; Nishimukai, H.; Yamasawa, K.: Genetic polymorphism
[39359] of the B subunit of human coagulation factor XIII: another classification. Hum.
[39360] Genet. 59: 360-364, 1981.

[39361] 6030.

[39362] Kitchens, C. S.; Newcomb, T. F.: Factor XIII. Medicine 58:

[39363] 413-429, 1979.

[39364] 6031.

[39365] Kohler, H. P.; Stickland, M. H.; Ossei-Gerning, N.; Carter, A.;
[39366] Mikkola, H.; Grant, P. J.: Association of a common polymorphism in
[39367] the factor XIII gene with myocardial infarction. Thromb.

Haemost. 79:
[39368] 8-13, 1998.

[39369] 6032.

[39370] Kompf, J.; Schunter, F.; Wernet, P.; Ritter, H.: Linkage be-
tween
[39371] the loci for mitochondrial malic enzyme (ME2) and coagu-
lation factor
[39372] XIIIA subunit (F13A). Hum. Genet. 70: 43-44, 1985.

[39373] 6033.

[39374] Lorand, L.; Urayama, T.; Atencio, A. C.; Hsia, D. Y.-Y.: In-
heritance
[39375] of deficiency of fibrin-stabilizing factor (factor XIII). Am. J.
[39376] Hum. Genet. 22: 89-95, 1970.

[39377] 6034.

[39378] Board, P. G.: Further electrophoretic studies of erythrocyte
glutathione
[39379] peroxidase. Am. J. Hum. Genet. 35: 914-918, 1983.

[39380] 6035.

[39381] Bock, A.; Forchhammer, K.; Heider, J.; Leinfelder, W.; Saw-
ers,

[39382] G.; Veprek, B.; Zinoni, F.: Selenocysteine: the 21st amino acid. Molec.
[39383] Microbiol. 5: 515-520, 1991.

[39384] 6036.

[39385] Boivin, P.; Galand, C.; Hakim, J.: Anemie hemolytique avec deficit
[39386] en glutathion-peroxydase chez un adulte. Enzym. Biol. Clin. 10:
[39387] 68-80, 1969.

[39388] 6037.

[39389] Chada, S.; Le Beau, M. M.; Casey, L.; Newburger, P. E.: Isolation
[39390] and chromosomal localization of the human glutathione peroxidase gene. Genomics 6:
[39391] 268-271, 1990.

[39392] 6038.

[39393] Cohen, H. J.; Chovaniec, M. E.; Mistretta, O.; Baker, S. S.: Selenium
[39394] repletion and glutathione peroxidase-differential effects on plasma
[39395] and red cell enzyme activity. Am. J. Clin. Nutr. 41:

735-747, 1985.

[39396] 6039.

[39397] Feaster, W. W.; Kwok, L. W.; Epstein, C. J.: Dosage effects for

[39398] superoxide dismutase-1 in nucleated cells aneuploid for chromosome

[39399] 21. Am. J. Hum. Genet. 29: 563-570, 1977.

[39400] 6040.

[39401] Forsberg, L.; de Faire, U.; Morgenstern, R.: Low yield of polymorphisms

[39402] from EST Blast searching: analysis of genes related to oxidative stress

[39403] and verification of the P197L polymorphism in GPX1. Hum. Mutat. 13:

[39404] 294-300, 1999.

[39405] 6041.

[39406] de Haan, J. B.; Bladier, C.; Griffiths, P.; Kelner, M.; O'Shea,

[39407] R. D.; Cheung, N. S.; Bronson, R. T.; Silvestro, M. J.; Wild, S.;

[39408] Zheng, S. S.; Beart, P. M.; Hertzog, P. J.; Kola, I.: Mice with a

[39409] homozygous null mutation for the most abundant glutathione peroxidase,
[39410] Gpx1, show increased susceptibility to the oxidative stress-inducing
[39411] agents paraquat and hydrogen peroxide. J. Biol. Chem. 273: 22528-22536,
[39412] 1998.

[39413] 6042.

[39414] Golan, R.; Ezzer, J. B.; Szeinberg, A.: Red cell glutathione

[39415] peroxidase in various Jewish ethnic groups in Israel. Hum. Hered. 30:
[39416] 136-141, 1980.

[39417] 6043.

[39418] Johannsmann, R.; Hellkuhl, B.; Grzeschik, K.-H.: Regional assignment
[39419] of a gene for glutathione peroxidase on human chromosome 3. (Abstract) Cytogenet.
[39420] Cell Genet. 25: 167 only, 1979.

[39421] 6044.

[39422] Johannsmann, R.; Hellkuhl, B.; Grzeschik, K.-H.: Regional mapping

[39423] of human chromosome 3: assignment of a glutathione peroxidase-1 gene
[39424] to 3p13-3q12. Hum. Genet. 56: 361-363, 1981.

[39425] 6045.

[39426] Kiss, C.; Li, J.; Szeles, A.; Gizatullin, R. Z.; Kashuba, V. I.;

[39427] Lushnikova, T.; Protopopov, A. I.; Kelve, M.; Kiss, H.; Kholodnyuk,
[39428] I. D.; Imreh, S.; Klein, G.; Zabarovsky, E. R.: Assignment of the
[39429] ARHA and GPX1 genes to human chromosome bands 3p21.3 by in situ hybridization
[39430] and with somatic cell hybrids. Cytogenet. Cell Genet. 79: 228-230,
[39431] 1997.

[39432] 6046.

[39433] Le Beau, M. M.: Personal Communication. Chicago, Ill. 1/23/1989.
[39434] 6047.

[39435] Le Beau, M. M.; Chada, S.; Casey, L.; Newburger, P. E.: Chromosomal
[39436] sublocalization of the human glutathione peroxidase

gene. (Abstract) Cytogenet.

[39437] Cell Genet. 51: 1030 only, 1989.

[39438] 6048.

[39439] McBride, O. W.; Mitchell, A.; Lee, B. J.; Mullenbach, G.; Hatfield,

[39440] D.: Gene for selenium-dependent glutathione peroxidase maps to human

[39441] chromosomes 3, 21 and X. BioFactors 1: 285-292, 1988.

[39442] 6049.

[39443] Meera Khan, P.; Verma, C.; Wijnen, L. M. M.; Jairaj, S.: Red

[39444] cell glutathione peroxidase (GPX1) variation in Afro-Jamaican, Asiatic

[39445] Indian, and Dutch populations: is the GPX1*2 allele of 'Thomas' variant

[39446] an African marker? Hum. Genet. 66: 352-355, 1984.

[39447] 6050.

[39448] Meera Khan, P.; Verma, C.; Wijnen, L. M. M.; Wijnen, J. T.; Prins,

[39449] H. K.; Nijenhuis, L. E.: Electrotypes and formal genetics of red

[39450] cell glutathione peroxidase (GPX1) in the Djuka of Suri-

nam. Am. J.

[39451] Hum. Genet. 38: 712-723, 1986.

[39452] 6051.

[39453] Mehdizadeh, S.; Warden, C. H.; Wen, P.-Z.; Xia, Y.-R.; Mehrabian,

[39454] M.; Lusis, A. J.: The glutathione peroxidase gene, Gpx1, maps to

[39455] mouse chromosome 9. Mammalian Genome 7: 465-466, 1996.

[39456] 6052.

[39457] Necheles, T. F.; Boles, T. A.; Allen, D. M.: Erythrocyte glu-tathione-peroxidase

[39458] deficiency and hemolytic disease of the newborn infant. J. Pediat. 72:

[39459] 319-324, 1968.

[39460] 6053.

[39461] Necheles, T. F.; Maldonado, N. I.; Barquet-Chediak, A.; Allen,

[39462] D. M.: Homozygous erythrocyte glutathione-peroxidase deficiency:

[39463] clinical and biochemical studies. Blood 33: 164-169,

1969.

[39464] 6054.

[39465] Necheles, T. F.; Steinberg, M. H.; Cameron, D.: Erythrocyte glutathione-peroxidase

[39466] deficiency. Brit. J. Haemat. 19: 605-612, 1970.

[39467] 6055.

[39468] Nishimura, Y.; Chida, N.; Hayashi, T.; Arakawa, T. S.: Homozygous

[39469] glutathione-peroxidase deficiency of erythrocytes and leukocytes. Tohoku

[39470] J. Exp. Med. 108: 207-218, 1972.

[39471] 6056.

[39472] Paglia, D. E.: Erythroenzymopathies: 1960s in retrospect. Curr.

[39473] Contents 32: 17 only, 1989.

[39474] 6057.

[39475] Paglia, D. E.; Valentine, W. N.: Studies on the quantitative

[39476] and qualitative characterization of erythrocyte glutathione peroxidase. J.

[39477] Lab. Clin. Med. 70: 158-169, 1967.

[39478] 6058.

[39479] Perona, G.; Guidi, G. C.; Piga, A.; Cellerino, R.; Milani, G.;

[39480] Colautti, P.; Moschini, G.; Stievano, B. M.: Neonatal ery-
throcyte

[39481] glutathione peroxidase deficiency as a consequence of se-
lenium imbalance

[39482] during pregnancy. Brit. J. Haemat. 42: 567-574, 1979.

[39483] 6059.

[39484] Reddy, V. N.; Giblin, F. J.; Lin, L.-R.; Dang, L.; Unakar, N.

[39485] J.; Musch, D. C.; Boyle, D. L.; Takemoto, L. J.; Ho, Y.-S.;
Knoernschild,

[39486] T.; Juenemann, A.; Lutjen-Drecoll, E.: Glutathione peroxi-
dase-1 deficiency

[39487] leads to increased nuclear light scattering, membrane
damage, and

[39488] cataract formation in gene-knockout mice. Invest. Oph-
thal. Vis. Sci. 42:

[39489] 3247-3255, 2001.

[39490] 6060.

[39491] Shen, Q.; Townes, P. L.; Padden, C.; Newburger, P. E.: An
in-frame

[39492] trinucleotide repeat in the coding region of the human cellular glutathione

[39493] peroxidase (GPX1) gene: in vivo polymorphism and in vitro instability. Genomics 23:

[39494] 292-294, 1994.

[39495] 6061.

[39496] Sinet, P. M.; Michelson, A. M.; Bazin, A.; Lejeune, J.; Jerome,

[39497] H.: Increase in glutathione peroxidase activity in erythro-cytes from

[39498] trisomy 21 subjects. Biochem. Biophys. Res. Commun. 67: 910-915,

[39499] 1975.

[39500] 6062.

[39501] Steinberg, M. H.; Brauer, M. J.; Necheles, T. F.: Acute hemolytic

[39502] anemia associated with erythrocyte glutathione-peroxi-dase deficiency. Arch.

[39503] Intern. Med. 125: 302-303, 1970.

[39504] 6063.

[39505] Steinberg, M. H.; Necheles, T. F.: Erythrocyte glutathione

peroxidase
[39506] deficiency: biochemical studies on the mechanisms of drug-induced
[39507] hemolysis. Am. J. Med. 50: 542-546, 1971.

[39508] 6064.

[39509] Sukenaga, Y.; Ishida, K.; Takeda, T.; Takagi, K.: cDNA se-quence
[39510] coding for human glutathione peroxidase. Nucleic Acids Res. 15:
[39511] 7178 only, 1987.

[39512] 6065.

[39513] Takahashi, K.; Avissar, N.; Whitin, J.; Cohen, H.: Purifica-tion
[39514] and characterization of human plasma glutathione peroxi-dase: a selenoglycoprotein
[39515] distinct from the known cellular enzyme. Arch. Biochem. Biophys. 256:
[39516] 677-686, 1987.

[39517] 6066.

[39518] Danciger, M.; Farber, D. B.; Peyser, M.; Kozak, C. A.: The gene

[39519] for the beta-subunit of retinal transducin (Gnb-1) maps to distal
[39520] mouse chromosome 4, and related sequences map to mouse chromosomes
[39521] 5 and 8. Genomics 6: 428-435, 1990.

[39522] 6067.

[39523] Hurowitz, E. H.; Melnyk, J. M.; Chen, Y.-J.; Kouros-Mehr, H.; Simon,
[39524] M. I.; Shizuya, H.: Genomic characterization of the human heterotrimeric
[39525] G protein alpha, beta, and gamma subunit genes. DNA Res. 7: 111-120,
[39526] 2000.

[39527] 6068.

[39528] Modi, W. S.; O'Brien, S. J.; Levine, M. A.: Chromosomal as-signment
[39529] of 2 GTP binding protein subunit genes: the alpha subunit of adenylyl
[39530] cyclase (GNAS) and the beta 1 polypeptide (GNB). (Abstract) Cytogenet.
[39531] Cell Genet. 58: 1860 only, 1991.

[39532] 6069.

[39533] Gao, B.; Gilman, A. G.; Robishaw, J. D.: A second form of the

[39534] beta subunit of signal-transducing G proteins. Proc. Nat. Acad. Sci. 84:

[39535] 6122-6125, 1987.

[39536] 6070.

[39537] Kere, J.; Taillon-Miller, P.; Lovett, M.; de la Chapelle, A.; Donis-Keller,

[39538] H.: Construction of human chromosome 7 YAC contigs around the COL1A2

[39539] and EPO genes and mapping of the GNB2 gene. (Abstract) Cytogenet.

[39540] Cell Genet. 58: 1922-1923, 1991.

[39541] 6071.

[39542] Lovett, M.; Kere, J.; Hinton, L. M.: Direct selection: a method

[39543] for the isolation of cDNAs encoded by large genomic re-gions. Proc.

[39544] Nat. Acad. Sci. 88: 9628-9632, 1991.

[39545] 6072.

[39546] Parimoo, S.; Patanjali, S. R.; Shukla, H.; Chaplin, D. D.; Weissman,
[39547] S. M.: cDNA selection: efficient PCR approach for the selection of
[39548] cDNAs encoded in large chromosomal DNA fragments. Proc. Nat. Acad.
[39549] Sci. 88: 9623-9627, 1991.

[39550] 6073.

[39551] Peng, Y.-W.; Robishaw, J. D.; Levine, M. A.; Yau, K.-W.: Retinal
[39552] rods and cones have distinct G protein beta and gamma subunits. Proc.
[39553] Nat. Acad. Sci. 89: 10882-10886, 1992.

[39554] 6074.

[39555] Ma, J. J.; Nishimura, M.; Mine, H.; Kuroki, S.; Nukina, M.; Ohta,
[39556] M.; Saji, H.; Obayashi, H.; Kawakami, H.; Saida, T.; Uchiyama, T.
[39557] : Genetic contribution of the tumor necrosis factor region in Guillain-Barre
[39558] syndrome. Ann. Neurol. 44: 815-818, 1998.

[39559] 6075.

[39560] Giuili, G.; Roechel, N.; Scholl, U.; Mattei, M.-G.; Guellaen, G.

[39561] : Colocalization of the genes coding for the alpha-3 and beta-3 subunits

[39562] of soluble guanylyl cyclase to human chromosome 4 at q31.3-q33. Hum.

[39563] Genet. 91: 257-260, 1993.

[39564] 6076.

[39565] Cameron, H. S.; Szczepaniak, D.; Weston, B. W.: Expression of

[39566] human chromosome 19p alpha-(1,3)-fucosyltransferase genes in normal

[39567] tissues: alternative splicing, polyadenylation, and isoforms. J.

[39568] Biol. Chem. 270: 20112-20122, 1995.

[39569] 6077.

[39570] Koszdin, K. L.; Bowen, B. R.: The cloning and expression of a

[39571] human alpha-1,3 fucosyltransferase capable of forming the E-selectin

[39572] ligand. Biochem. Biophys. Res. Commun. 187: 152-157, 1992.

[39573] 6078.

[39574] Mollicone, R.; Reguigne, I.; Fletcher, A.; Aziz, A.; Rustam, M.;

[39575] Weston, B. W.; Kelly, R. J.; Lowe, J. B.; Oriol, R.: Molecular basis

[39576] for plasma alpha(1,3)-fucosyltransferase gene deficiency (FUT6). J.

[39577] Biol. Chem. 269: 12662-12671, 1994.

[39578] 6079.

[39579] Pang, H.; Koda, Y.; Soejima, M.; Schlaphoff, T.; Du Toit, E. D.;

[39580] Kimura, H.: Allelic diversity of the human plasma al-pha(1,3)fucosyltransferase

[39581] gene (FUT6). Ann. Hum. Genet. 63: 277-284, 1999.

[39582] 6080.

[39583] Schnyder-Candrian, S.; Borsig, L.; Moser, R.; Berger, E. G.: Localization

[39584] of alpha-1,3-fucosyltransferase VI in Weibel-Palade bod-ies of human

[39585] endothelial cells. Proc. Nat. Acad. Sci. 97: 8369-8374, 2000.

[39586] 6081.

[39587] Allen, S. J.; O'Donnell, A.; Alexander, N. D. E.; Alpers, M. P.;

[39588] Peto, T. E. A.; Clegg, J. B.; Weatherall, D. J.: Alpha(+)-thalassemia

[39589] protects children against disease caused by other infections as well

[39590] as malaria. Proc. Nat. Acad. Sci. 94: 14736-14741, 1997.

[39591] 6082.

[39592] Keats, B. J. B.; Morton, N. E.; Rao, D. C.: Possible linkages

[39593] (lod score over 1.5) and a tentative map of the Jk-Km linkage group. Cytogenet.

[39594] Cell Genet. 22: 304-308, 1978.

[39595] 6083.

[39596] Keats, B. J. B.; Morton, N. E.; Rao, D. C.: Likely linkage: InV

[39597] with Jk. Hum. Genet. 39: 157-159, 1977.

[39598] 6084.

[39599] Cannizzaro, L. A.; Croce, C. M.; Griffin, C. A.; Simeone, A.;

Boncinelli,
[39600] E.; Huebner, K.: Human homeo box-containing genes located at chromosome
[39601] regions 2q31-2q37 and 12q12-12q13. Am. J. Hum. Genet. 41: 1-15,
[39602] 1987.

[39603] 6085.

[39604] Boncinelli, E.; Acampora, D.; Pannese, M.; D'Esposito, M.; Somma,
[39605] R.; Gaudino, G.; Stornaiuolo, A.; Cafiero, M.; Faiella, A.; Simeone,
[39606] A.: Organization of human class I homeobox genes. Genome 31: 745-756,
[39607] 1989.

[39608] 6086.

[39609] Chariot, A.; Castronovo, V.; Le, P.; Gillet, C.; Sobel, M. E.;

[39610] Gielen, J.: Cloning and expression of a new HOXC6 transcript encoding
[39611] a repressing protein. Biochem. J. 319: 91-97, 1996.

[39612] 6087.

[39613] Semina, E. V.; Reiter, R.; Leysens, N. J.; Alward, W. L. M.;

Small,
[39614] K. W.; Datson, N. A.; Siegel-Bartelt, J.; Bierke-Nelson, D.; Bitoun,
[39615] P.; Zabel, B. U.; Carey, J. C.; Murray, J. C.: Cloning and characterization
[39616] of a novel bicoid-related homeobox transcription factor gene, RIEG,
[39617] involved in Rieger syndrome. Nature Genet. 14: 392-399, 1996.
[39618] 6088.

[39619] Zhou, X.-P.; Marsh, D. J.; Hampel, H.; Mulliken, J. B.; Gimm,
[39620] O.; Eng, C.: Germline and germline mosaic PTEN mutations associated
[39621] with a Proteus-like syndrome of hemihypertrophy, lower limb asymmetry,
[39622] arteriovenous malformations and lipomatosis. Hum. Molec. Genet. 9:
[39623] 765-768, 2000.

[39624] 6089.

[39625] Gibbs, S.; Fijneman, R.; Wiegant, J.; Geurts van Kessel, A.; van

[39626] de Putte, P.; Backendorf, C.: Molecular characterization and evolution
[39627] of the SPRR family of keratinocyte differentiation markers encoding
[39628] small proline-rich proteins. Genomics 16: 630-637, 1993.

[39629] 6090.

[39630] Bressler, J.; Tsai, T.-F.; Wu, M.-Y.; Tsai, S.-F.; Ramirez, M.

[39631] A.; Armstrong, D.; Beaudet, A. L.: The SNRPN promoter is not required
[39632] for genomic imprinting of the Prader-Willi/Angelman domain in mice. Nature
[39633] Genet. 28: 232-240, 2001.

[39634] 6091.

[39635] Buiting, K.; Saitoh, S.; Gross, S.; Dittrich, B.; Schwartz, S.;

[39636] Nicholls, R. D.; Horsthemke, B.: Inherited microdeletions in the
[39637] Angelman and Prader-Willi syndromes define an imprinting centre on
[39638] human chromosome 15. Nature Genet. 9: 395-400, 1995.

[39639] 6092.

[39640] Dittrich, B.; Buiting, K.; Korn, B.; Rickard, S.; Buxton, J.; Saitoh,
[39641] S.; Nicholls, R. D.; Poustka, A.; Winterpacht, A.; Zabel, B.; Horsthemke,
[39642] B.: Imprint switching on human chromosome 15 may involve alternative
[39643] transcripts of the SNRPN gene. Nature Genet. 14: 163-170, 1996.
[39644] 6093.
[39645] Gallagher, R. C.; Pils, B.; Albalwi, M.; Francke, U.: Evidence
[39646] for the role of PWCR1/HBII-85 C/D box small nucleolar RNAs in Prader-Willi
[39647] syndrome. Am. J. Hum. Genet. 71: 669-678, 2002.
[39648] 6094.
[39649] Glenn, C. C.; Porter, K. A.; Jong, M. T. C.; Nicholls, R. D.; Driscoll,
[39650] D. J.: Functional imprinting and epigenetic modification of the human
[39651] SNRPN gene. Hum. Molec. Genet. 2: 2001-2005, 1993.
[39652] 6095.
[39653] Gray, T. A.; Saitoh, S.; Nicholls, R. D.: An imprinted, mam-

malian

[39654] bicistronic transcript encodes two independent proteins. Proc. Nat.

[39655] Acad. Sci. 96: 5616-5621, 1999.

[39656] 6096.

[39657] Kuslich, C. D.; Kobori, J. A.; Mohapatra, G.; Gregorio-King, C.;

[39658] Donlon, T. A.: Prader-Willi syndrome is caused by disruption of the

[39659] SNRPN gene. Am. J. Hum. Genet. 64: 70-76, 1999.

[39660] 6097.

[39661] Leff, S. E.; Brannan, C. I.; Reed, M. L.; Ozcelik, T.; Francke,

[39662] U.; Copeland, N. G.; Jenkins, N. A.: Maternal imprinting of the mouse

[39663] Snrpn gene and conserved linkage homology with the human Prader-Willi

[39664] syndrome region. Nature Genet. 2: 259-264, 1992.

[39665] 6098.

[39666] Li, S.; Klein, E. S.; Russo, A. F.; Simmons, D. M.; Rosenfeld,

[39667] M. G.: Isolation of cDNA clones encoding small nuclear ri-

bonucleoparticle-associated
[39668] proteins with different tissue specificities. Proc. Nat. Acad. Sci. 86:
[39669] 9778-9782, 1989.

[39670] 6099.

[39671] Lyko, F.; Buiting, K.; Horsthemke, B.; Paro, R.: Identification
[39672] of a silencing element in the human 15q11-q13 imprinting center by
[39673] using transgenic Drosophila. Proc. Nat. Acad. Sci. 95: 1698-1702,
[39674] 1998.

[39675] 6100.

[39676] McAllister, G.; Amara, S. G.; Lerner, M. R.: Tissue-specific
[39677] expression and cDNA cloning of small nuclear ribonucleoprotein-associated
[39678] polypeptide N. Proc. Nat. Acad. Sci. 85: 5296-5300, 1988.

[39679] 6101.

[39680] Mutirangura, A.; Jayakumar, A.; Sutcliffe, J. S.; Nakao, M.; McKinney,
[39681] M. J.; Buiting, K.; Horsthemke, B.; Beaudet, A. L.; Chinault,

A. C.;

[39682] Ledbetter, D. H.: A complete YAC contig of the Prader-Willi/Angelman

[39683] chromosome region (15q11-q13) and refined localization of the SNRPN

[39684] gene. Genomics 18: 546-552, 1993.

[39685] 6102.

[39686] Ozcelik, T.; Leff, S.; Robinson, W.; Donlon, T.; Lalande, M.;

[39687] Sanjines, E.; Schinzel, A.; Francke, U.: Small nuclear ribonucleoprotein

[39688] polypeptide N (SNRPN), an expressed gene in the Prader-Willi syndrome

[39689] critical region. Nature Genet. 2: 265-269, 1992.

[39690] 6103.

[39691] Reed, M. L.; Leff, S. E.: Maternal imprinting of human SNRPN,

[39692] a gene deleted in Prader-Willi syndrome. Nature Genet. 6: 163-167,

[39693] 1994.

[39694] 6104.

[39695] Saitoh, S.; Wada, T.: Parent-of-origin specific histone

acetylation
[39696] and reactivation of a key imprinted gene locus in Prader-Willi syndrome. Am.
[39697] J. Hum. Genet. 66: 1958-1962, 2000.

[39698] 6105.

[39699] Tiller, G. E.; Polumbo, P. A.; Weis, M. A.; Bogaert, R.; Lachman,
[39700] R. S.; Cohn, D. H.; Rimoin, D. L.; Eyre, D. R.: Dominant mutations
[39701] in the type II collagen gene, COL2A1, produce spondyloepimetaphyseal
[39702] dysplasia, Strudwick type. Nature Genet. 11: 87-89, 1995.

[39703] 6106.

[39704] Tiller, G. E.; Weis, M. A.; Lachman, R. S.; Cohn, D. H.; Rimoin,
[39705] D. L.; Eyre, D. R.: A dominant mutation in the type II collagen gene
[39706] (COL2A1) produces spondyloepimetaphyseal dysplasia (SEMD), Strudwick
[39707] type. (Abstract) Am. J. Hum. Genet. 53 (suppl.): A209 only, 1993.

[39708] 6107.

[39709] Xu, H.; Wu, X.-R.; Wewer, U. M.; Engvall, E.: Murine mus-
cular
[39710] dystrophy caused by a mutation in the laminin alpha-2
(Lama2) gene. Nature
[39711] Genet. 8: 297-302, 1994.

[39712] 6108.

[39713] Zhang, X.; Vuolteenaho, R.; Tryggvason, K.: Structure of
the
[39714] human laminin alpha-2-chain gene (LAMA2), which is af-
fected in congenital
[39715] muscular dystrophy. J. Biol. Chem. 271: 27664-27669,
1996.
[39716] 6109.

[39717] Kantaputra, P. N.; Gorlin, R. J.; Langer, L. O., Jr.: Dominant

[39718] mesomelic dysplasia, ankle, carpal, and tarsal synostosis
type: a
[39719] new autosomal dominant bone disorder. Am. J. Med.
Genet. 44: 730-737,
[39720] 1992.

[39721] 6110.

[39722] Baser, M. E.; De Rienzo, A.; Altomare, D.; Balsara, B. R.;

Hedrick,
[39723] N. M.; Gutmann, D. H.; Pitts, L. H.; Jackler, R. K.; Testa, J. R.
[39724] : Neurofibromatosis 2 and malignant mesothelioma. Neurology 59:
[39725] 290-291, 2002.

[39726] 6111.

[39727] Backman, S. A.; Stambolic, V.; Suzuki, A.; Haight, J.; Elia, A.;
[39728] Pretorius, J.; Tsao, M.-S.; Shannon, P.; Bolon, B.; Ivy, G. O.; Mak,
[39729] T. W.: Deletion of Pten in mouse brain causes seizures, ataxia and
[39730] defects in soma size resembling Lhermitte-Duclos disease. Nature
[39731] Genet. 29: 396-403, 2001.

[39732] 6112.

[39733] Fackenthal, J. D.; Marsh, D. J.; Richardson, A.-L.; Cummings,
[39734] S. A.; Eng, C.; Robinson, B. G.; Olopade, O. I.: Male breast cancer
[39735] in Cowden syndrome patients with germline PTEN muta-

tions. J. Med.

[39736] Genet. 38: 159-164, 2001.

[39737] 6113.

[39738] Kwon, C.-H.; Zhu, X.; Zhang, J.; Knoop, L. L.; Tharp, R.; Smeyne,

[39739] R. J.; Eberhart, C. G.; Burger, P. C.; Baker, S. J.: Pten regu-lates

[39740] neuronal soma size: a mouse model of Lhermitte-Duclos disease. Nature

[39741] Genet. 29: 404-411, 2001.

[39742] 6114.

[39743] Liaw, D.; Marsh, D. J.; Li, J.; Dahia, P. L. M.; Wang, S. I.;

[39744] Zheng, Z.; Bose, S.; Call, K. M.; Tsou, H. C.; Peacocke, M.; Eng,

[39745] C.; Parsons, R.: Germline mutations of the PTEN gene in Cowden disease,

[39746] an inherited breast and thyroid cancer syndrome. Nature Genet. 16:

[39747] 64-67, 1997.

[39748] 6115.

[39749] Marsh, D. J.; Dahia, P. L. M.; Caron, S.; Kum, J. B.; Frayling,

[39750] I. M.; Tomlinson, I. P. M.; Hughes, K. S.; Eeles, R. A.; Hodgson,
[39751] S. V.; Murday, V. A.; Houlston, R.; Eng, C.: Germline PTEN mutations
[39752] in Cowden syndrome-like families. J. Med. Genet. 35: 881-885, 1998.
[39753] 6116.

[39754] Labeit, S.; Kolmerer, B.: The complete primary structure of human
[39755] nebulin and its correlation to muscle structure. J. Molec. Biol. 248:
[39756] 308-315, 1995.

[39757] 6117.

[39758] Limongi, M. Z.; Pelliccia, F.; Rocchi, A.: Assignment of the human

[39759] nebulin gene (NEB) to chromosome band 2q24.2 and the alpha-1 (III)

[39760] collagen gene (COL3A1) to chromosome band 2q32.2 by in situ hybridization:

[39761] the FRA2G common fragile site lies between the two genes in the 2q31

[39762] band. Cytogenet. Cell Genet. 77: 259-260, 1997.

[39763] 6118.

[39764] Schurr, E.; Skamene, E.; Gros, P.: Mapping of the gene coding

[39765] for the muscle protein nebulin (Neb) to the proximal region of mouse

[39766] chromosome 2. Cytogenet. Cell Genet. 57: 214-216, 1991.

[39767] 6119.

[39768] Stedman, H.; Browning, K.; Oliver, N.; Oronzi-Scott, M.; Fischbeck,

[39769] K.; Sarkar, S.; Sylvester, J.; Schmickel, R.; Wang, K.: Nebulin cDNAs

[39770] detect a 25-kilobase transcript in skeletal muscle and localize to

[39771] human chromosome 2. Genomics 2: 1-7, 1988.

[39772] 6120.

[39773] Wang, K.; Knipfer, M.; Huang, Q.-Q.; van Heerden, A.; Hsu, L. C.-L.;

[39774] Gutierrez, G.; Quian, X.-L.; Stedman, H.: Human skeletal muscle nebulin

[39775] sequence encodes a blueprint for thin filament architecture: sequence

[39776] motifs and affinity profiles of tandem repeats and terminal SH3. J.

[39777] Biol. Chem. 271: 4304-4314, 1996.

[39778] 6121.

[39779] Pelin, K.; Hilpela, P.; Donner, K.; Sewry, C.; Akkari, P. A.; Wilton,

[39780] S. D.; Wattanasirichaigoon, D.; Bang, M.-L.; Centner, T.; Hanefeld,

[39781] F.; Odent, S.; Fardeau, M.; Urtizberea, J. A.; Muntoni, F.; Dubowitz,

[39782] V.; Beggs, A. H.; Laing, N. G.; Labeit, S.; de la Chapelle, A.; Wallgren-Pettersson,

[39783] C.: Mutations in the nebulin gene associated with autosomal recessive
[39784] nemaline myopathy. Proc. Nat. Acad. Sci. 96: 2305-2310, 1999.
[39785] 6122.

[39786] Pelin, K.; Ridanpaa, M.; Donner, K.; Wilton, S.; Krishnarajah,
[39787] J.; Laing, N.; Kolmerer, B.; Millevoi, S.; Labeit, S.; de la Chapelle,
[39788] A.; Wallgren-Pettersson, C.: Refined localisation of the genes for
[39789] nebulin and titin on chromosome 2q allows the assignment of nebulin
[39790] as a candidate gene for autosomal recessive nemaline myopathy. Europ.
[39791] J. Hum. Genet. 5: 229-234, 1997.

[39792] 6123.

[39793] Zeviani, M.; Darras, B. T.; Rizzuto, R.; Salviati, G.; Betto, R.;

[39794] Bonilla, E.; Miranda, A. F.; Du, J.; Samitt, C.; Dickson, G.; Walsh,
[39795] F. S.; DiMauro, S.; Francke, U.; Schon, E. A.: Cloning and expression

[39796] of human nebulin cDNAs and assignment of the gene to chromosome 2q31-q32. Genomics 2:
[39797] 249-256, 1988.

[39798] 6124.

[39799] Laing, N. G.; Wilton, S. D.; Akkari, P. A.; Dorosz, S.; Boundy,
[39800] K.; Kneebone, C.; Blumbergs, P.; White, S.; Watkins, H.; Love, D.
[39801] R.; Haan, E.: A mutation in the alpha tropomyosin gene TPM3 associated
[39802] with autosomal dominant nemaline myopathy. Nature Genet. 9: 75-79,
[39803] 1995.

[39804] 6125.

[39805] Shirakawa, H.; Yoshida, M.: Structure of a gene coding for human
[39806] HMG2 protein. J. Biol. Chem. 267: 6641-6645, 1992.

[39807] 6126.

[39808] Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.; Bartnitzke,
[39809] S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the hu-

man HMG2
[39810] gene to 4q31. Genomics 31: 264-265, 1996.

[39811] 6127.

[39812] Bustin, M.: Regulation of DNA-dependent activities by the functional
[39813] motifs of the high-mobility-group chromosomal proteins. Molec. Cell.
[39814] Biol. 19: 5237-5246, 1999.

[39815] 6128.

[39816] Landsman, D.; Bustin, M.: Chromosomal proteins HMG-14 and HMG-17:
[39817] distinct multigene families coding for similar types of transcripts. J.
[39818] Biol. Chem. 261: 16087-16091, 1986.

[39819] 6129.

[39820] Landsman, D.; Soares, N.; Gonzalez, F. J.; Bustin, M.: Chromosomal
[39821] protein HMG-17: complete human cDNA sequence and evidence for a multigene
[39822] family. J. Biol. Chem. 261: 7479-7484, 1986.

[39823] 6130.

[39824] Mitchell, A.; McBride, W.; Landsman, D.; Bustin, M.: Chromosomal
[39825] mapping of HMG-17 gene to human chromosome 1p. (Abstract) Am. J.
[39826] Hum. Genet. 43: A152 only, 1988.

[39827] 6131.

[39828] Mitchell, A. L.; Bale, A. E.; Bustin, M.; Landsman, D.; Popescu,
[39829] N.; McBride, O. W.: Localization of HMG17 gene to chromosome 1p35-36.1.
[39830] (Abstract) Cytogenet. Cell Genet. 51: 1045 only, 1989.

[39831] 6132.

[39832] Popescu, N.; Landsman, D.; Bustin, M.: Mapping the human gene
[39833] coding for chromosomal protein HMG-17. Hum. Genet. 85: 376-378,
[39834] 1990.

[39835] 6133.

[39836] Porkka, K.; Laakkonen, P.; Hoffman, J. A.; Bernasconi, M.; Ruoslahti,
[39837] E.: A fragment of the HMGN2 protein homes to the nuclei

of tumor
[39838] cells and tumor endothelial cells in vivo. Proc. Nat. Acad. Sci. 99:
[39839] 7444-7449, 2002.

[39840] 6134.

[39841] Srikantha, J.; Landsman, D.; Bustin, M.: Retropseudogenes for
[39842] human chromosomal protein HMG-17. J. Molec. Biol. 197: 405-413,
[39843] 1987.

[39844] 6135.

[39845] Murakami, K.; Blei, F.; Tilton, W.; Seaman, C.; Piomelli, S.:

[39846] An isozyme of hexokinase specific for the human red blood cell (HK-R). Blood 75:
[39847] 770-775, 1990.

[39848] 6136.

[39849] Murakami, K.; Piomelli, S.: Identification of the cDNA for human
[39850] red blood cell-specific hexokinase isozyme. Blood 89: 762-766, 1997.
[39851] 6137.

[39852] Deal, C.; Ma, J.; Wilkin, F.; Paquette, J.; Rozen, F.; Ge, B.;

[39853] Hudson, T.; Stampfer, M.; Pollak, M.: Novel promoter polymorphism

[39854] in insulin-like growth factor-binding protein-3: correlation with

[39855] serum levels and interaction with known regulators. J. Clin. Endocr.

[39856] Metab. 86: 1274-1280, 2001.

[39857] 6138.

[39858] Ehrenborg, E.; Larsson, C.; Stern, I.; Janson, M.; Powell, D. R.;

[39859] Luthman, H.: Contiguous localization of the genes encoding human

[39860] insulin-like growth factor-binding proteins 1 (IGBP1) and 3 (IGBP3)

[39861] on chromosome 7. Genomics 12: 497-502, 1992.

[39862] 6139.

[39863] Ferry, R. J., Jr.; Cerri, R. W.; Cohen, P.: Insulin-like growth

[39864] factor binding proteins: new proteins, new functions. Horm. Res. 51:

[39865] 53-67, 1999.

[39866] 6140.

[39867] Fraser, H. M.; Lunn, S. F.; Kim, H.; Duncan, W. C.; Rodger, F.

[39868] E.; Illingworth, P. J.; Erickson, G. F.: Changes in insulin-like

[39869] growth factor-binding protein-3 messenger ribonucleic acid in endothelial

[39870] cells of the human corpus luteum: a possible role in luteal development

[39871] and rescue. J. Clin. Endocr. Metab. 85: 1672-1677, 2000.

[39872] 6141.

[39873] Kiefer, M. C.; Ioh, R. S.; Bauer, D. M.; Zapf, J.: Molecular cloning

[39874] of a new human insulin-like growth factor binding pro-tein. Biochem.

[39875] Biophys. Res. Commun. 176: 219-225, 1991.

[39876] 6142.

[39877] Weinzimer, S. A.; Gibson, T. B.; Collett-Solberg, P. F.; Khare,

[39878] A.; Liu, B.; Cohen, P.: Transferrin is an insulin-like growth factor-binding

[39879] protein-3 binding protein. J. Clin. Endocr. Metab. 86: 1806-1813,
[39880] 2001.

[39881] 6143.

[39882] Wood, W. I.; Cachianes, G.; Henzel, W. J.; Winslow, G. A.; Spencer,
[39883] S. A.; Hellmiss, R.; Martin, J. L.; Baxter, R. C.: Cloning and expression
[39884] of the GH dependent IGF binding protein. Molec. Endocr. 2: 1176-1185,
[39885] 1988.

[39886] 6144.

[39887] Bajalica, S.; Allander, S. V.; Ehrenborg, E.; Brondum-Nielsen,
[39888] K.; Luthman, H.; Larsson, C.: Localization of the human insulin-like
[39889] growth-factor-binding protein 4 gene to chromosomal region 17q12-21.1. Hum.
[39890] Genet. 89: 234-236, 1992.

[39891] 6145.

[39892] Kiefer, M. C.; Schmid, C.; Waldvogel, M.; Schlapfer, I.;

Futo,
[39893] E.; Masiarz, F. R.; Green, K.; Barr, P. J.; Zapf, J.: Characterization
[39894] of recombinant human insulin-like growth factor binding proteins 4,
[39895] 5, and 6 produced in yeast. J. Biol. Chem. 267: 12692-12699, 1992.
[39896] 6146.

[39897] Shimasaki, S.; Uchiyama, F.; Shimonaka, M.; Ling, N.: Molecular
[39898] cloning of the cDNAs encoding a novel insulin-like growth factor-binding
[39899] protein from rat and human. Molec. Endocr. 4: 1451-1458, 1990.
[39900] 6147.

[39901] Tonin, P.; Ehrenborg, E.; Lenoir, G.; Feunteun, J.; Lynch, H.;
[39902] Morgan, K.; Zazzi, H.; Vivier, A.; Pollak, M.; Huynh, H.; Luthman,
[39903] H.; Larsson, C.; Narod, S.: The human insulin-like growth factor-binding
[39904] protein 4 gene maps to chromosome region 17q12-q21.1

and is close

[39905] to the gene for hereditary breast-ovarian cancer. Genomics 18: 414-417,

[39906] 1993.

[39907] 6148.

[39908] Zazzi, H.; Nikoshkov, A.; Hall, K.; Luthman, H.: Structure and

[39909] transcription regulation of the human insulin-like growth factor binding

[39910] protein 4 gene (IGFBP4). Genomics 49: 401-410, 1998.

[39911] 6149.

[39912] Kou, K.; James, P. L.; Clemmons, D. R.; Copeland, N. G.; Gilbert,

[39913] D. J.; Jenkins, N. A.; Rotwein, P.: Identification of two clusters

[39914] of mouse insulin-like growth factor binding protein genes on chromosomes

[39915] 1 and 11. Genomics 21: 653-655, 1994.

[39916] 6150.

[39917] Kato, M.; Ishizaki, A.; Hellman, U.; Wernstedt, C.; Kyogoku, M.;

[39918] Miyazono, K.; Heldin, C.-H.; Funa, K.: A human keratinocyte cell

[39919] line produces two autocrine growth inhibitors, transforming growth

[39920] factor-beta and insulin-like growth factor binding protein-6, in a

[39921] calcium- and cell density-dependent manner. J. Biol. Chem. 270:

[39922] 12373-12379, 1995.

[39923] 6151.

[39924] Shimasaki, S.; Gao, L.; Shimonaka, M.; Ling, N.: Isolation and

[39925] molecular cloning of insulin-like growth factor-binding protein-6. Molec.

[39926] Endocr. 5: 938-948, 1991.

[39927] 6152.

[39928] Brink, P. A.; Steyn, L. T.; Coetzee, G. A.; van der Westhuyzen,

[39929] D. R.: Familial hypercholesterolemia in South African Afrikaners:

[39930] PvuII and StuI DNA polymorphisms in the LDL-receptor gene consistent

[39931] with a predominating founder gene effect. Hum. Genet. 77: 32-35,

[39932] 1987.

[39933] 6153.

[39934] Brown, M. S.; Goldstein, J. L.: Analysis of a mutant strain of

[39935] human fibroblasts with a defect in the internalization of receptor-bound

[39936] low density lipoproteins. Cell 9: 663-674, 1976.

[39937] 6154.

[39938] Brown, M. S.; Goldstein, J. L.: Familial hypercholes-

[39939] terolemia: defective binding of lipoproteins to cultured fibroblasts associated

[39940] with impaired regulation of 3-hydroxy-3-methylglutaryl coenzyme at

[39941] reductase activity. Proc. Nat. Acad. Sci. 71: 788-792, 1974.

[39942] 6155.

[39943] Davis, C. G.; Lehrman, M. A.; Russell, D. W.; Anderson, R. G.

[39944] W.; Brown, M. S.; Goldstein, J. L.: The J.D. mutation in familial
[39945] hypercholesterolemia: amino acid substitution in cytoplasmic domain
[39946] impedes internalization of LDL receptors. Cell 45: 15-24, 1986.
[39947] 6156.
[39948] De Braekeleer, M.: Hereditary disorders in Saguenay-Lac-St-Jean
[39949] (Quebec, Canada). Hum. Hered. 41: 141-146, 1991.
[39950] 6157.
[39951] Defesche, J. C.; Kastelein, J. J. P.: Molecular epidemiology
[39952] of familial hypercholesterolaemia. (Letter) Lancet 352: 1643-1644,
[39953] 1998.
[39954] 6158.
[39955] Durst, R.; Colombo, R.; Shpitzen, S.; Ben Avi, L.; Friedlander,
[39956] Y.; Wexler, R.; Raal, F. J.; Marais, D. A.; Defesche, J. C.; Mandelshtam,
[39957] M. Y.; Kotze, M. J.; Leitersdorf, E.; Meiner, V.: Recent ori-

gin and

[39958] spread of a common Lithuanian mutation, G197del LDLR, causing familial

[39959] hypercholesterolemia: positive selection is not always necessary to

[39960] account for disease incidence among Ashkenazi Jews. Am. J. Hum. Genet. 68:

[39961] 1172-1188, 2001.

[39962] 6159.

[39963] Ekstrom, U.; Abrahamson, M.; Floren, C.-H.; Tollig, H.; Wettrell,

[39964] G.; Nilsson, G.; Sun, X.-M.; Soutar, A. K.; Nilsson-Ehle, P.: An

[39965] individual with a healthy phenotype in spite of a pathogenic LDL receptor

[39966] mutation (C240F). Clin. Genet. 55: 332-339, 1999.

[39967] 6160.

[39968] Elston, R. C.; Namboodiri, K. K.; Go, R. C. P.; Siervogel, R.

[39969] M.; Glueck, C. J.: Probable linkage between essential familial hypercholesterolemia

[39970] and third complement component (C3). Cytogenet. Cell

Genet. 16:
[39971] 294-297, 1976.

[39972] 6161.

[39973] Feussner, G.; Dobmeyer, J.; Nissen, H.; Hansen, T. S.: Un-
usual
[39974] Xanthomas in a young patient with heterozygous familial
hypercholesterolemia
[39975] and type III hyperlipoproteinemia. Am. J. Med. Genet. 65:
149-154,
[39976] 1996.

[39977] 6162.

[39978] Francke, U.; Brown, M. S.; Goldstein, J. L.: Assignment of
the
[39979] human gene for the low density lipoprotein receptor to
chromosome
[39980] 19: synteny of a receptor, a ligand, and a genetic disease.
Proc.
[39981] Nat. Acad. Sci. 81: 2826-2830, 1984.

[39982] 6163.

[39983] Frank, S. L.; Taylor, B. A.; Lusis, A. J.: Linkage of the
mouse

[39984] LDL receptor gene on chromosome 9. Genomics 5: 646-648, 1989.

[39985] 6164.

[39986] Copeland, N. G.; Silan, C. M.; Kingsley, D. M.; Jenkins, N. A.;

[39987] Cannizzaro, L. A.; Croce, C. M.; Huebner, K.; Sims, J. E.: Chromosomal

[39988] location of murine and human IL-1 receptor genes. Genomics 9: 44-50,

[39989] 1991.

[39990] 6165.

[39991] Dale, M.; Nicklin, M. J.: Interleukin-1 receptor cluster: gene

[39992] organization of IL1R2, IL1R1, IL1RL2 (IL-1Rrp2), IL1RL1 (T1/ST2),

[39993] and IL18R1 (IL-1Rrp) on human chromosome 2q. Genomics 57: 177-179,

[39994] 1999.

[39995] 6166.

[39996] Dower, S. K.; Kronheim, S. R.; Hopp, T. P.; Cantrell, M.; Deeley,

[39997] M.; Gillis, S.; Henney, C. S.; Urdal, D. L.: The cell surface receptors

[39998] for interleukin-1(alpha) and interleukin-1(beta) are identical. Nature 324:

[39999] 266-268, 1986.

[40000] 6167.

[40001] Sims, J. E.; Acres, R. B.; Grubin, C. E.; McMahan, C. J.; Wignall,

[40002] J. M.; March, C. J.; Dower, S. K.: Cloning the interleukin 1 receptor

[40003] from human T cells. Proc. Nat. Acad. Sci. 86: 8946-8950, 1989.

[40004] 6168.

[40005] Habets, G. G. M.; van der Kammen, R. A.; Willemsen, V.; Balemans,

[40006] M.; Wiegant, J.; Collard, J. G.: Sublocalization of an invasion-inducing

[40007] locus and other genes on human chromosome 7. Cytogenet. Cell Genet. 60:

[40008] 200-205, 1992.

[40009] 6169.

[40010] Hill, A. V. S.: The immunogenetics of resistance to malaria. Proc.

[40011] Assoc. Am. Phys. 111: 272-277, 1999.

[40012] 6170.

[40013] Katz, F. E.; Parkar, M.; Stanley, K.; Murray, L. J.; Clark, E.

[40014] A.; Greaves, M. F.: Chromosome mapping of cell membrane antigens

[40015] expressed on activated B cells. Europ. J. Immun. 15: 103-106, 1985.

[40016] 6171.

[40017] Gedde-Dahl, T., Jr.; Dupuy, B. M.; Jonassen, R.; Winberg, J.-O.;

[40018] Anton-Lamprecht, I.; Olaisen, B.: Junctional epidermolysis bullosa

[40019] inversa (locus EBR2A) assigned to 1q31 by linkage and association

[40020] of LAMC1. Hum. Molec. Genet. 3: 1387-1391, 1994.

[40021] 6172.

[40022] Kallunki, T.; Ikonen, J.; Chow, L. T.; Kallunki, P.; Tryggvason,

[40023] K.: Structure of the human laminin B2 chain gene reveals

extensive
[40024] divergence from the laminin B1 chain gene. J. Biol. Chem. 266: 221-228,
[40025] 1991.

[40026] 6173.

[40027] Kallunki, T.; Pikkarainen, T.; Tryggvason, K.; Savolainen, E.-R.
[40028] : A Pst I polymorphism in the human laminin B2 chain gene on 1q25-q31. Nucleic
[40029] Acids Res. 17: 4423 only, 1989.

[40030] 6174.

[40031] Mattei, M.-G.; Weil, D.; Passage, E.; Van Cong, N.; Pribula-Conway,
[40032] D.; Timpl, R.; Chu, M.-L.: Human gene for laminin B2 chain (LAMB2)
[40033] maps to the long arm of chromosome 1. (Abstract) Cyto-genet. Cell
[40034] Genet. 46: 659 only, 1987.

[40035] 6175.

[40036] Mattei, M.-G.; Weil, D.; Pribula-Conway, D.; Bernard, M. P.; Passage,

[40037] E.; Van Cong, N.; Timpl, R.; Chu, M.-L.: cDNA cloning, ex-
pression
[40038] and mapping of human laminin B2 gene to chromosome
1q31. Hum. Genet. 79:
[40039] 235-241, 1988.

[40040] 6176.

[40041] Miner, J. H.; Patton, B. L.; Lentz, S. I.; Gilbert, D. J.; Jenkins,

[40042] N. A.; Copeland, N. G.; Sanes, J. R.: The laminin alpha
chains: expression,
[40043] developmental transitions, and chromosomal locations of
alpha1-5,
[40044] identification of heterodimeric laminins 8-11, and cloning
of a novel
[40045] alpha3 isoform. J. Cell Biol. 137: 685-701, 1997.

[40046] 6177.

[40047] Braun, A.; Kofler, A.; Morawietz, S.; Cleve, H.: Sequence
and
[40048] organization of the human vitamin D-binding protein
gene. Biochim.
[40049] Biophys. Acta 1216: 385-394, 1993.

[40050] 6178.

[40051] Chautard-Freire-Maia, E. A.: Concerning the linkage relationships
[40052] of the Gc and MNSs loci (Hum. Genet. 43: 215-220, 1978): disentangling
[40053] part of the data overlap. (Letter) Hum. Genet. 49: 115-116, 1979.
[40054] 6179.

[40055] Cleve, H.; Kirk, R. L.; Gajdusek, D. C.; Guiart, J.: On the distribution
[40056] of the Gc variant Gc Aborigine in Melanesian populations: determination
[40057] of Gc-types in sera from Tongariki Island, New Hebrides. Acta Genet.
[40058] Statist. Med. 17: 511-517, 1967.

[40059] 6180.

[40060] Cleve, H.; Kirk, R. L.; Parker, W. C.; Bearn, A. G.; Schacht,
[40061] L. E.; Kleinman, H.; Horsfall, W. R.: Two genetic variants of the
[40062] group-specific component of human serum: Gc Chippewa and Gc Aborigine. Am.
[40063] J. Hum. Genet. 15: 368-379, 1963.

[40064] 6181.

[40065] Cleve, H.; Patutschnick, W.: The vitamin D binding of the common

[40066] rare variants of the group-specific component (Gc): an autoradiographic

[40067] study. Hum. Genet. 38: 289-296, 1977.

[40068] 6182.

[40069] Constans, J.; Cleve, H.; Dykes, D.; Fischer, M.; Kirk, R. L.;

[40070] Papiha, S. S.; Scheffran, W.; Scherz, R.; Thymann, M.; We-

ber, W.:

[40071] The polymorphism of the vitamin D-binding protein (Gc); isoelectric

[40072] focusing in 3 M urea as additional method for identifica-

tion of genetic

[40073] variants. Hum. Genet. 65: 176-180, 1983.

[40074] 6183.

[40075] Constans, J.; Viau, M.: Group-specific component: evi-

dence for

[40076] two subtypes of the Gc(1) gene. Science 198: 1070-1071, 1977.

[40077] 6184.

[40078] Cooke, N. E.; David, E. V.: Serum vitamin D-binding protein is
[40079] a third member of the albumin and alpha fetoprotein gene family. J.
[40080] Clin. Invest. 76: 2420-2424, 1985.

[40081] 6185.

[40082] Cooke, N. E.; Willard, H. F.; David, E. V.; George, D. L.: Direct
[40083] regional assignment of the gene for vitamin D binding protein (Gc-globulin)
[40084] to human chromosome 4q11-q13 and identification of an associated DNA
[40085] polymorphism. Hum. Genet. 73: 225-229, 1986.

[40086] 6186.

[40087] Daiger, S. P.; Brewton, G. W.; Rios, A. A.; Mansell, P. W. A.;

[40088] Reuben, J. M.: Genetic susceptibility to AIDS: absence of an association
[40089] with group-specific component (Gc). (Letter) New Eng. J. Med. 317:
[40090] 631-632, 1987.

[40091] 6187.

[40092] Daiger, S. P.; Cavalli-Sforza, L. L.: Detection of genetic variation
[40093] with radioactive ligands. II. Genetic variants of vitamin D-labeled
[40094] group-specific component (Gc) proteins. Am. J. Hum. Genet. 29: 593-604,
[40095] 1977.

[40096] 6188.

[40097] Daiger, S. P.; Miller, M.; Chakraborty, R.: Heritability of quantitative
[40098] variation at the group-specific component (Gc) locus. Am. J. Hum.
[40099] Genet. 36: 663-676, 1984.

[40100] 6189.

[40101] Daiger, S. P.; Schanfield, M. S.; Cavalli-Sforza, L. L.: Group-specific
[40102] component (Gc) proteins bind vitamin D and 25-hydroxyvitamin D. Proc.
[40103] Nat. Acad. Sci. 72: 2076-2080, 1975.

[40104] 6190.

[40105] Dykes, D.; Copouls, B.; Polesky, H.: Description of six new

Gc
[40106] variants. Hum. Genet. 63: 35-37, 1983.

[40107] 6191.

[40108] Dykes, D. D.; Polesky, H. F.: Gc1C12: a new Gc variant. Hum.
[40109] Hered. 32: 136-138, 1982.

[40110] 6192.

[40111] Eales, L.-J.; Nye, K. E.; Parkin, J. M.; Weber, J. N.; Forster,

[40112] S. M.; Harris, J. R. W.; Pinching, A. J.: Association of different
[40113] allelic forms of group specific component with susceptibility to and
[40114] clinical manifestation of human immunodeficiency virus infection. Lancet I:
[40115] 999-1002, 1987.

[40116] 6193.

[40117] Falk, C. T.; Martin, M. D.; Walker, M. E.; Chen, T.; Rubin-
[40118] stein, P.; Allen, F. H., Jr.: Family data suggesting a linkage between MN
[40119] and Gc. (Abstract) Cytogenet. Cell Genet. 25: 152 only,

1979.

[40120] 6194.

[40121] German, J. L.; Walker, M. E.; Stiefel, F. H.; Allen, F. H., Jr.

[40122] : Autoradiographic studies of human chromosomes. II. Data concerning

[40123] the position of the MN locus. Vox Sang. 16: 130-145, 1969.

[40124] 6195.

[40125] Gilles, K.; Louie, L.; Newman, B.; Crandall, J.; King, M.-C.:

[40126] Genetic susceptibility to AIDS: absence of an association with group-specific

[40127] component (Gc). (Letter) New Eng. J. Med. 317: 630-631, 1987.

[40128] 6196.

[40129] Henningsen, K.; Jacobsen, P.; Mikkelsen, M.: B-F chromosome translocation

[40130] associated with father-child incompatibility within the Gc-system. Hum.

[40131] Hered. 19: 283-287, 1969.

[40132] 6197.

[40133] Hirschfeld, J.: The Gc-system: immuno-electrophoretic

studies
[40134] of normal human sera with special reference to a new genetically determined
[40135] serum system (GC). Prog. Allergy 6: 155-186, 1962.

[40136] 6198.

[40137] Hirschfeld, J.: Immune-electrophoretic demonstration of qualitative
[40138] differences in human sera and their relation to the haptoglobins. Acta
[40139] Path. Microbiol. Scand. 47: 160-168, 1959.

[40140] 6199.

[40141] Johnson, A. M.; Cleve, H.; Alper, C. A.: Variants of the group-specified
[40142] component system as demonstrated by immunofixation electrophoresis:
[40143] report of a new variant, Gc Boston (Gc B). Am. J. Hum. Genet. 27:
[40144] 728-736, 1975.

[40145] 6200.

[40146] Karlsson, S.; Arnason, A.; Thordarson, G.; Olaisen, B.: Frequency

[40147] of Gc alleles and a variant Gc allele in Iceland. Hum. Hered. 30:
[40148] 119-121, 1980.

[40149] 6201.

[40150] Kofler, A.; Braun, A.; Jenkins, T.; Serjeantson, S. W.; Cleve,
[40151] H.: Characterization of mutants of the vitamin-D-binding protein/group
[40152] specific component: GC Aborigine (1A1) from Australian Aborigines
[40153] and South African blacks, and 2A9 from South Germany. Vox Sang. 68:
[40154] 50-54, 1995.

[40155] 6202.

[40156] Magenis, R. E.; Eoff, J. S.; Toth-Fejel, S.; Lovrien, E.: Proba-
[40157] ble linkage of GC to a chromosome 4 inversion and localiza-
[40158] tion to 4q12. (Abstract) Cytogenet. Cell Genet. 40: 684 only, 1985.

[40159] 6203.

[40160] Mars, M.; Farrant, S.; Roberts, G. J.: Dentinogenesis imper-
fecta,

[40161] report of a 5-generation family. Brit. Dent. J. 140:
206-209, 1976.

[40162] 6204.

[40163] McCombs, J. L.; Yang, F.; Bowman, B. H.; McGill, J. R.;
Moore,

[40164] C. M.: Chromosomal localization of group-specific com-
ponent by in

[40165] situ hybridization. Cytogenet. Cell Genet. 42: 62-64,
1986.

[40166] 6205.

[40167] Mikkelsen, M.; Jacobsen, P.; Henningsen, K.: Possible lo-
calization

[40168] of Gc-system on chromosome 4. Loss of long arm 4 ma-
terial associated

[40169] with father-child incompatibility within the Gc-system.
Hum. Hered. 27:

[40170] 105-107, 1977.

[40171] 6206.

[40172] Mourant, A. E.; Tills, D.; Domaniewska-Sobczak, K.: Sun-
shine

[40173] and the geographical distribution of the alleles of the Gc
system

[40174] of plasma proteins. Hum. Genet. 33: 307-314, 1976.

[40175] 6207.

[40176] Parker, W. C.; Cleve, H.; Bearn, A. G.: Determination of phenotypes

[40177] in the human group-specific component (Gc) system by starch gel electrophoresis. Am.

[40178] J. Hum. Genet. 15: 353-367, 1963.

[40179] 6208.

[40180] Petrini, M.; Emerson, D. L.; Galbraith, R. M.: Linkage between

[40181] surface immunoglobulin and cytoskeleton of B lymphocytes may involve

[40182] Gc protein. Nature 306: 73-74, 1983.

[40183] 6209.

[40184] Cleaver, J. E.; Thompson, L. H.; Richardson, A. S.; States, J.

[40185] C.: A summary of mutations in the UV-sensitive disorders: xeroderma

[40186] pigmentosum, Cockayne syndrome, and trichothiodystrophy. Hum. Mutat. 14:

[40187] 9-22, 1999.

[40188] 6210.

[40189] Coetzer, T.; Zail, S. S.: Tryptic digestion of spectrin in variants

[40190] of hereditary elliptocytosis. J. Clin. Invest. 67: 1241-1248, 1981.

[40191] 6211.

[40192] Evans, J. P. M.; Baines, A. J.; Hann, I. M.; Al-Hakim, I.; Knowles,

[40193] S. M.; Hoffbrand, A. V.: Defective spectrin dimer-dimer association

[40194] in a family with transfusion dependent homozygous hereditary elliptocytosis. Brit.

[40195] J. Haemat. 54: 163-172, 1983.

[40196] 6212.

[40197] Keats, B. J. B.: Another elliptocytosis locus on chromo-some 1? Hum.

[40198] Genet. 50: 227-230, 1979.

[40199] 6213.

[40200] Rao, D. C.; Keats, B. J.; Lalouel, J. M.; Morton, N. E.; Yee,

[40201] S.: A maximum likelihood map of chromosome 1. Am. J. Hum. Genet. 31:

[40202] 680-696, 1979.

[40203] 6214.

[40204] Kaneda, Y.; Yoshida, M. C.; Kohno, K.; Uchida, T.; Okada, Y.:

[40205] Chromosomal assignment of the gene for human elongation factor-2. Proc.

[40206] Nat. Acad. Sci. 81: 3158-3162, 1984.

[40207] 6215.

[40208] Rapp, G.; Klaudiny, J.; Hagendorff, G.; Luck, M. R.; Scheit, K.

[40209] H.: Complete sequence of the coding region of human elongation factor

[40210] 2 (EF-2) by enzymatic amplification of cDNA from human ovarian granulosa

[40211] cells. Biol. Chem. Hoppe-Seyler 370: 1071-1075, 1989.

[40212] 6216.

[40213] Kenmochi, N.; Kawaguchi, T.; Rozen, S.; Davis, E.; Goodman, N.;

[40214] Hudson, T. J.; Tanaka, T.; Page, D. C.: A map of 75 human ribosomal

[40215] protein genes. Genome Res. 8: 509-523, 1998.

[40216] 6217.

[40217] Alders, M.; Ryan, A.; Hodges, M.; Bliek, J.; Feinberg, A. P.; Privitera,

[40218] O.; Westerveld, A.; Little, P. F. R.; Mannens, M.: Disruption of

[40219] a novel imprinted zinc-finger gene, ZNF215, in Beckwith-Wiedemann

[40220] syndrome. Am. J. Hum. Genet. 66: 1473-1484, 2000.

[40221] 6218.

[40222] Algar, E. M.; Deeble, G. J.; Smith, P. J.: CDKN1C expression in

[40223] Beckwith-Wiedemann syndrome patients with allele im-balance. J. Med.

[40224] Genet. 36: 524-531, 1999.

[40225] 6219.

[40226] Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp,

[40227] N.; Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins,

[40228] N. A.: A genetic linkage map of mouse chromosome 10: localization

[40229] of eighteen molecular markers using a single interspecific

backcross. Genetics 125:
[40230] 855-866, 1990.

[40231] 6220.

[40232] King, M. C.: Personal Communication. Berkeley, Calif. 5/1996.

[40233] 6221.

[40234] Korach, K. S.: Insights from the study of animals lacking functional
[40235] estrogen receptor. Science 266: 1524-1527, 1994.

[40236] 6222.

[40237] Kos, M.; Reid, G.; Denger, S.; Gannon, F.: Minireview: genomic
[40238] organization of the human ER-alpha gene promoter region. Molec. Endocr. 15:
[40239] 2057-2063, 2001.

[40240] 6223.

[40241] Kumar, R.; Wang, R.-A.; Mazumdar, A.; Talukder, A. H.; Mandal,
[40242] M.; Yang, Z.; Bagheri-Yarmand, R.; Sahin, A.; Hortobagyi, G.; Adam,
[40243] L.; Barnes, C. J.; Vadlamudi, R. K.: A naturally occurring

MTA1 variant
[40244] sequesters oestrogen receptor-alpha in the cytoplasm. Nature 418:
[40245] 654-657, 2002.

[40246] 6224.

[40247] Lawson, J. S.; Field, A. S.; Champion, S.; Tran, D.; Ishikura,

[40248] H.; Trichopoulos, D.: Low oestrogen receptor alpha expression in
[40249] normal breast tissue underlies low breast cancer incidence in Japan.
[40250] (Letter) Lancet 354: 1787-1788, 1999.

[40251] 6225.

[40252] Lonard, D. M.; Nawaz, Z.; Smith, C. L.; O'Malley, B. W.: The

[40253] 26S proteasome is required for estrogen receptor-alpha and coactivator
[40254] turnover and for efficient estrogen receptor-alpha transactivation. Molec.
[40255] Cell 5: 939-948, 2000.

[40256] 6226.

[40257] Lorentzon, M.; Lorentzon, R.; Backstrom, T.; Nordstrom,

P.: Estrogen
[40258] receptor gene polymorphism, but not estradiol levels, is related to
[40259] bone density in healthy adolescent boys: a cross-sectional and longitudinal
[40260] study. J. Clin. Endocr. Metab. 84: 4597-4601, 1999.

[40261] 6227.

[40262] Mader, S.; Kumar, V.; de Verneuil, H.; Chambon, P.: Three amino
[40263] acids of the oestrogen receptor are essential to its ability to distinguish
[40264] an oestrogen from a glucocorticoid-responsive element. Nature 338:
[40265] 271-274, 1989.

[40266] 6228.

[40267] McGuire, W. L.; Chamness, G. C.; Fuqua, S. A. W.: Estrogen receptor
[40268] variants in clinical breast cancer. Molec. Endocr. 5: 1571-1577,
[40269] 1991.

[40270] 6229.

[40271] McGuire, W. L.; Chamness, G. C.; Fuqua, S. A. W.: Abnormal estrogen

[40272] receptor in clinical breast cancer. J. Steroid Biochem. Molec. Biol. 43:

[40273] 243-247, 1992.

[40274] 6230.

[40275] McInerney, E. M.; Ince, B. A.; Shapiro, D. J.; Katzenellenbogen,

[40276] B. S.: A transcriptionally active estrogen receptor mutant is a novel

[40277] type of dominant negative inhibitor of estrogen action. Molec. Endocr. 10:

[40278] 1519-1526, 1996.

[40279] 6231.

[40280] Menasce, L. P.; White, G. R. M.; Harrison, C. J.; Boyle, J. M.

[40281] : Localization of the estrogen receptor locus (ESR) to chromosome

[40282] 6q25.1 by FISH and a simple post-FISH banding technique. Genomics 17:

[40283] 263-265, 1993.

[40284] 6232.

[40285] Metzger, D.; White, J. H.; Chambon, P.: The human oestrogen receptor
[40286] functions in yeast. Nature 334: 31-36, 1988.

[40287] 6233.

[40288] Murphy, L. C.; Wang, M.; Coutt, A.; Dotzlaw, H.: Novel mutations
[40289] in the estrogen receptor messenger RNA in human breast cancers. J.
[40290] Clin. Endocr. Metab. 81: 1420-1427, 1996.

[40291] 6234.

[40292] Pelletier, G.; El-Alfy, M.: Immunocytochemical localization of
[40293] estrogen receptors alpha and beta in the human reproductive organs. J.
[40294] Clin. Endocr. Metab. 85: 4835-4840, 2000.

[40295] 6235.

[40296] Ponglikitmongkol, M.; Green, S.; Chambon, P.: Genomic organization
[40297] of the human oestrogen receptor gene. EMBO J. 7: 3385-3388, 1988.

[40298] 6236.

[40299] Reese, J. C.; Katzenellenbogen, B. S.: Characterization of a

[40300] temperature-sensitive mutation in the hormone binding domain of the

[40301] human estrogen receptor: studies in cell extracts and intact cells

[40302] and their implications for hormone-dependent transcriptional activation. J.

[40303] Biol. Chem. 267: 9868-9873, 1992.

[40304] 6237.

[40305] Reese, J. C.; Katzenellenbogen, B. S.: Mutagenesis of cysteines

[40306] in the hormone binding domain of the human estrogen receptor: alterations

[40307] in binding and transcriptional activation by covalently and reversibly

[40308] attaching ligands. J. Biol. Chem. 266: 10880-10887, 1991.

[40309] 6238.

[40310] Scott, G. K.; Kushner, P.; Vigne, J.-L.; Benz, C. C.: Truncated

[40311] forms of DNA-binding estrogen receptors in human breast cancer. J.

[40312] Clin. Invest. 88: 700-706, 1991.

[40313] 6239.

[40314] Shang, Y.; Brown, M.: Molecular determinants for the tis-
sue specificity
[40315] of SERMs. Science 295: 2465-2468, 2002.

[40316] 6240.

[40317] Shiau, A. K.; Barstad, D.; Loria, P. M.; Cheng, L.; Kushner,
P.
[40318] J.; Agard, D. A.; Greene, G. L.: The structural basis of es-
trogen
[40319] receptor/coactivator recognition and the antagonism of
this interaction
[40320] by tamoxifen. Cell 95: 927-937, 1998.

[40321] 6241.

[40322] Shupnik, M. A.; Pitt, L. K.; Soh, A. Y.; Anderson, A.; Lopes,

[40323] M. B.; Laws, E. R., Jr.: Selective expression of estrogen re-
ceptor
[40324] alpha and beta isoforms in human pituitary tumors. J.
Clin. Endocr.
[40325] Metab. 83: 3965-3972, 1998.

[40326] 6242.

[40327] Simoncini, T.; Hafezi-Moghadam, A.; Brazil, D. P.; Ley, K.; Chin,

[40328] W. W.; Liao, J. K.: Interaction of oestrogen receptor with the regulatory

[40329] subunit of phosphatidylinositol-3-OH kinase. Nature 407: 538-541,

[40330] 2000.

[40331] 6243.

[40332] Sluyser, M.: Mutations in the estrogen receptor gene. Hum. Mutat. 6:

[40333] 97-103, 1995.

[40334] 6244.

[40335] Sluyser, M.; Mester, J.: Oncogenes homologous to steroid receptors?

[40336] (Letter) Nature 315: 546 only, 1985.

[40337] 6245.

[40338] Smith, E. P.; Boyd, J.; Frank, G. R.; Takahashi, H.; Cohen, R.

[40339] M.; Specker, B.; Williams, T. C.; Lubahn, D. B.; Korach, K. S.: Estrogen

[40340] resistance caused by a mutation in the estrogen-receptor gene in a
[40341] man. New Eng. J. Med. 331: 1056-1061, 1994.

[40342] 6246.

[40343] Sudhir, K.; Chou, T. M.; Chatterjee, K.; Smith, E. P.; Williams,
[40344] T. C.; Kane, J. P.; Malloy, M. J.; Korach, K. S.; Rubanyi, G. M.:
[40345] Premature coronary artery disease associated with a disruptive mutation
[40346] in the estrogen receptor gene in a man. Circulation 96: 3774-3777,
[40347] 1997.

[40348] 6247.

[40349] Takeyama, J.; Suzuki, T.; Inoue, S.; Kaneko, C.; Nagura, H.; Harada,
[40350] N.; Sasano, H.: Expression and cellular localization of estrogen
[40351] receptors alpha and beta in the human fetus. J. Clin. Endocr. Metab. 86:
[40352] 2258-2262, 2001.

[40353] 6248.

[40354] Walter, P.; Green, S.; Greene, G.; Krust, A.; Bornert, J.-M.;

[40355] Jeltsch, J.-M.; Staub, A.; Jensen, E.; Scrace, G.; Waterfield, M.;

[40356] Chambon, P.: Cloning of the human estrogen receptor cDNA. Proc.

[40357] Nat. Acad. Sci. 82: 7889-7893, 1985.

[40358] 6249.

[40359] Weel, A. E. A. M.; Uitterlinden, A. G.; Westendorp, I. C. D.;

[40360] Burger, H.; Schuit, S. C. E.; Hofman, A.; Helmerhorst, T. J. M.; van

[40361] Leeuwen, J. P. T. M.; Pols, H. A. P.: Estrogen receptor polymorphism

[40362] predicts the onset of natural and surgical menopause. J. Clin. Endocr.

[40363] Metab. 84: 3146-3150, 1999.

[40364] 6250.

[40365] Weis, K. E.; Ekena, K.; Thomas, J. A.; Lazennec, G.; Katzenellenbogen,

[40366] B. S.: Constitutively active human estrogen receptors containing

[40367] amino acid substitutions for tyrosine 537 in the receptor protein. Molec.

[40368] Endocr. 10: 1388-1398, 1996.

[40369] 6251.

[40370] Barton, D. E.; Arquint, M.; Roder, J.; Dunn, R.; Francke, U.:

[40371] The myelin-associated glycoprotein gene: mapping to human chromosome

[40372] 19 and mouse chromosome 7 and expression in quivering mice. Genomics 1:

[40373] 107-112, 1987.

[40374] 6252.

[40375] Liu, B. P.; Fournier, A.; GrandPre, T.; Strittmatter, S. M.: Myelin-associated

[40376] glycoprotein as a functional ligand for the Nogo-66 receptor. Science 297:

[40377] 1190-1193, 2002.

[40378] 6253.

[40379] Barker, D. F.; Wright, E.; Nguyen, K.; Cannon, L.; Fain, P.; Goldgar,

[40380] D.; Bishop, D. T.; Carey, J.; Baty, B.; Kivlin, J.; Willard, H.; Waye,

[40381] J. S.; Greig, G.; Leinwand, L.; Nakamura, Y.; O'Connell, P.; Leppert,
[40382] M.; Lalouel, J.-M.; White, R.; Skolnick, M.: The gene for NF1 (von
[40383] Recklinghausen neurofibromatosis) is on chromosome 17 near the centromere.
[40384] (Abstract) Cytogenet. Cell Genet. 46: 576, 1987.

[40385] 6254.

[40386] Barrall, J. L.; Summers, C. G.: Ocular ischemic syndrome in a
[40387] child with moyamoya disease and neurofibromatosis. Surv. Ophthal. 40:
[40388] 500-504, 1996.

[40389] 6255.

[40390] Basu, T. N.; Gutmann, D. H.; Fletcher, J. A.; Glover, T. W.; Collins,
[40391] F. S.; Downward, J.: Aberrant regulation of ras proteins in malignant
[40392] tumour cells from type 1 neurofibromatosis patients. Na- ture 356:
[40393] 713-715, 1992.

[40394] 6256.

[40395] Benatar, M. G.: Intracranial fusiform aneurysms in von Recklinghausen's

[40396] disease: case report and literature review. (Letter) J. Neurol. Neurosurg.

[40397] Psychiat. 63: 1279-1280, 1994.

[40398] 6257.

[40399] Benedict, P. H.; Szabo, G.; Fitzpatrick, T. B.; Sinesi, S. J.

[40400] : Melanotic macules in Albright's syndrome and in neurofibromatosis. J.A.M.A. 205:

[40401] 618-626, 1968.

[40402] 6258.

[40403] Bidot-Lopez, P.; Frankel, J. W.: Enhanced viral transformation

[40404] of skin fibroblasts from neurofibromatosis patients. Ann. Clin. Lab.

[40405] Sci. 13: 27-32, 1983.

[40406] 6259.

[40407] Borberg, A.: Clinical and genetic investigations into tuberous

[40408] sclerosis and Recklinghausen's neurofibromatosis: contribution to

[40409] elucidation of interrelationship and eugenics of the syndromes. Acta

[40410] Psychiat. Neurol. 71 (suppl.): 1-239, 1951.

[40411] 6260.

[40412] Boudin, G.; Pepin, B.; Vernant, C.: Les tumeurs multiples du

[40413] systeme nerveux au cours de la maladie de Recklinghausen: a propos

[40414] d'une observation anatomo-clinique avec adenome chromophobe de l'hypophyse. Presse

[40415] Med. 78: 1427-1430, 1970.

[40416] 6261.

[40417] Brunner, H.; Stacher, G.; Bankl, H.; Grabner, G.: Chronic mesenteric

[40418] arterial insufficiency caused by vascular neurofibromatosis: a case

[40419] report. Am. J. Gastroent. 62: 442-447, 1974.

[40420] 6262.

[40421] Buchberg, A. M.; Cleveland, L. S.; Jenkins, N. A.; Copeland,

N.

[40422] G.: Sequence homology shared by neurofibromatosis type-1 gene and

[40423] IRA-1 and IRA-2 negative regulators of the RAS cyclic AMP pathway. Nature 347:

[40424] 291-294, 1990.

[40425] 6263.

[40426] Buntin, P. T.; Fitzgerald, J. F.: Gastrointestinal neurofibromatosis:

[40427] a rare cause of chronic anemia. Am. J. Dis. Child. 119: 521-523,

[40428] 1970.

[40429] 6264.

[40430] Cappione, A. J.; French, B. L.; Skuse, G. R.: A potential role

[40431] for NF1 mRNA editing in the pathogenesis of NF1 tumors. Am. J. Hum.

[40432] Genet. 60: 305-312, 1997.

[40433] 6265.

[40434] Cawthon, R. M.; Viskochil, D.; O'Connell, P.; Buchberg, A.; Andersen,

[40435] L.; Wallace, M.; Marchuk, D.; Stevens, J.; Culver, M.; Xu, G.;

Collins,
[40436] F. S.; Jenkins, N.; Copeland, N.; White, R.: Identification and characterization
[40437] of several genes between or nearby neurofibromatosis type I translocation
[40438] breakpoint. (Abstract) Am. J. Hum. Genet. 47 (suppl.): A109, 1990.
[40439] 6266.

[40440] Cawthon, R. M.; Weiss, R.; Xu, G.; Viskochil, D.; Culver, M.;

[40441] Stevens, J.; Robertson, M.; Dunn, D.; Gesteland, R.; O'Connell, P.;
[40442] White, R.: A major segment of the neurofibromatosis type 1 gene:
[40443] cDNA sequence, genomic structure, and point mutations. Cell 62:
[40444] 193-201, 1990.

[40445] 6267.

[40446] Charron, J. W.; Gariepy, G.: Neurofibromatosis of bladder: case
[40447] report and review of the literature. Canad. J. Surg. 13: 303-306,
[40448] 1970.

[40449] 6268.

[40450] Chu, M.-H.; Lee, H.-C.; Shen, E.-Y.; Wang, N.-L.; Yeung, C.-Y.;

[40451] Chen, B.-F.; Shih, S.-L.: Gastro-intestinal bleeding caused by leiomyoma

[40452] of the small intestine in a child with neurofibromatosis. Europ.

[40453] J. Pediat. 158: 460-462, 1999.

[40454] 6269.

[40455] Cichowski, K.; Shih, T. S.; Schmitt, E.; Santiago, S.; Reilly,

[40456] K.; McLaughlin, M. E.; Bronson, R. T.; Jacks, T.: Mouse models of

[40457] tumor development in neurofibromatosis type 1. Science 286: 2172-2176,

[40458] 1999.

[40459] 6270.

[40460] Clark, R. D.; Hutter, J. J., Jr.: Familial neurofibromatosis

[40461] and juvenile chronic myelogenous leukemia. Hum. Genet. 60: 230-232,

[40462] 1982.

[40463] 6271.

[40464] Clark, S. S.; Marlett, M.; Prudencio, R.; Dasgupta, T. K.: Neurofibromatosis

[40465] of the bladder in children: case report and literature review. J.

[40466] Urol. 118: 654-656, 1977.

[40467] 6272.

[40468] Clementi, M.; Barbujani, G.; Turolla, L.; Tenconi, R.: Neurofibromatosis-1:

[40469] a maximum likelihood estimation of mutation rate. Hum. Genet. 84:

[40470] 116-118, 1990.

[40471] 6273.

[40472] Clementi, M.; Milani, S.; Mammi, I.; Boni, S.; Monciotti, C.;

[40473] Tenconi, R.: Neurofibromatosis type 1 growth charts. Am. J. Med.

[40474] Genet. 87: 317-323, 1999.

[40475] 6274.

[40476] Auwerx, J.; Staels, B.: Leptin. Lancet 351: 737-742, 1998.

[40477] 6275.

[40478] Bado, A.; Levasseur, S.; Attoub, S.; Karmorgant, S.; Laigneau,
[40479] J.-P.; Bortoluzzi, M.-N.; Moizo, L.; Lehy, T.; Guerre-Millo, M.; Le
[40480] Marchand-Brustel, Y.; Lewin, M. J. M.: The stomach is a source of
[40481] leptin. Nature 394: 290-293, 1998.

[40482] 6276.

[40483] Cnossen, M. H.; Moons, K. G. M.; Garssen, M. P. J.; Pas-
[40484] M. T.; de Goede-Bolder, A.; Niermeijer, M. F.; Grobbee, D. E.; Neurofibromatosis
[40485] Team of Sophia Children's Hospital: Minor disease features in neurofibromatosis
[40486] type 1 (NF1) and their possible value in diagnosis of NF1 in children
[40487] less than 6 years and clinically suspected of having NF1. J. Med.
[40488] Genet. 35: 624-627, 1998.

[40489] 6277.

[40490] Cnossen, M. H.; van der Est, M. N.; Breuning, M. H.; van Asperen,

[40491] C. J.; Breslau-Siderius, E. J.; van der Ploeg, A. T.; de Goede-Bolder,
[40492] A.; van den Ouweland, A. M. W.; Halley, D. J. J.; Niermeijer, M. F.
[40493] : Deletions spanning the neurofibromatosis type 1 gene: implications
[40494] for genotype-phenotype correlations in neurofibromatosis type 1? Hum.
[40495] Mutat. 9: 458-464, 1997.

[40496] 6278.

[40497] Cohen, M. M., Jr.: Further diagnostic thoughts about the Elephant
[40498] Man. Am. J. Med. Genet. 29: 777-782, 1988.

[40499] 6279.

[40500] Colman, S. D.; Rasmussen, S. A.; Ho, V. T.; Abernathy, C. R.;
[40501] Wallace, M. R.: Somatic mosaicism in a patient with neu-rofibromatosis
[40502] type 1. Am. J. Hum. Genet. 58: 484-490, 1996.

[40503] 6280.

[40504] Colman, S. D.; Williams, C. A.; Wallace, M. R.: Benign neu-

rofibromas
[40505] in type 1 neurofibromatosis (NF1) show somatic deletions of the NF1
[40506] gene. Nature Genet. 11: 90-92, 1995.

[40507] 6281.

[40508] Combemale, P.; Abitan, R.; Kanitakis, J.: Segmental neurofibromatosis:
[40509] report of two cases and critical review of the literature. Europ.
[40510] J. Derm. 4: 194-201, 1994.

[40511] 6282.

[40512] Correa, C. L.; Brems, H.; Lazaro, C.; Marynen, P.; Legius, E.

[40513] : Unequal meiotic crossover: a frequent cause of NF1 microdeletions. Am.
[40514] J. Hum. Genet. 66: 1969-1974, 2000.

[40515] 6283.

[40516] Costa, R. M.; Federov, N. B.; Kogan, J. H.; Murphy, G. G.; Stern,
[40517] J.; Ohno, M.; Kucherlapati, R.; Jacks, T.; Silva, A. J.: Mechanism
[40518] for the learning deficits in a mouse model of neurofibro-

matosis type

[40519] 1. Nature 415: 526-530, 2002.

[40520] 6284.

[40521] Blain, H.; Vuillemin, A.; Guillemin, F.; Durant, R.; Hanesse, B.;

[40522] de Talance, N.; Doucet, B.; Jeandel, C.: Serum leptin level is a

[40523] predictor of bone mineral density in postmenopausal women. J. Clin.

[40524] Endocr. Metab. 87: 1030-1035, 2002.

[40525] 6285.

[40526] Campfield, L. A.; Smith, F. J.; Guisez, Y.; Devos, R.; Burn, P.

[40527] : Recombinant mouse OB protein: evidence for a peripheral signal linking

[40528] adiposity and central neural networks. Science 269: 546-549, 1995.

[40529] 6286.

[40530] Cao, R.; Brakenhielm, E.; Wahlestedt, C.; Thyberg, J.; Cao, Y.

[40531] : Leptin induces vascular permeability and synergistically

stimulates
[40532] angiogenesis with FGF-2 and VEGF. Proc. Nat. Acad. Sci. 98: 6390-6395,
[40533] 2001.

[40534] 6287.

[40535] van de Vijver, M. J.; Peterse, J. L.; Mooi, W. J.; Wisman, P.;

[40536] Lomans, J.; Dalesio, O.; Nusse, R.: NEU-protein overex-pression in
[40537] breast cancer: association with comedo-type ductal carci-noma in situ
[40538] and limited prognostic value in stage II breast cancer. New Eng.
[40539] J. Med. 319: 1239-1245, 1988.

[40540] 6288.

[40541] Xie, D.; Shu, X. O.; Deng, Z.; Wen, W.-Q.; Creek, K. E.; Dai,

[40542] Q., Gao, Y.-T.; Jin, F.; Zheng, W.: Population-based, case-control
[40543] study of HER2 genetic polymorphism and breast cancer risk. J. Nat.
[40544] Cancer Inst. 92: 412-417, 2000.

[40545] 6289.

[40546] Yamamoto, T.; Ikawa, S.; Akiyama, T.; Semba, K.; Nomura, N.; Miyajima,

[40547] N.; Saito, T.; Toyoshima, K.: Similarity of protein encoded by the

[40548] human c-erb-B-2 gene to epidermal growth factor receptor. Nature 319:

[40549] 230-234, 1986.

[40550] 6290.

[40551] Yang-Feng, T. L.; Schechter, A. L.; Weinberg, R. A.; Francke,

[40552] U.: Oncogene from rat neuro/glioblastomas (human gene symbol NGL)

[40553] is located on the proximal long arm of human chromosome 17 and EGFR

[40554] is confirmed at 7p13-q11.2. (Abstract) Cytogenet. Cell Genet. 40:

[40555] 784 only, 1985.

[40556] 6291.

[40557] Yu, D.; Jing, T.; Liu, B.; Yao, J.; Tan, M.; McDonnell, T. J.;

[40558] Hung, M.-C.: Overexpression of ErbB2 blocks Taxol-induced apoptosis

[40559] by upregulation of p21Cip1, which inhibits p34Cdc2 kinase. Molec.

[40560] Cell 2: 581-591, 1998.

[40561] 6292.

[40562] Klappacher, G. W.; Lunyak, V. V.; Sykes, D. B.; Sawka-Verhelle,

[40563] D.; Sage, J.; Brard, G.; Ngo, S. D.; Gangadharan, D.; Jacks, T.; Kamps,

[40564] M. P.; Rose, D. W.; Rosenfeld, M. G.: An induced Ets repressor complex

[40565] regulates growth arrest during terminal macrophage differentiation. Cell 109:

[40566] 169-180, 2002.

[40567] 6293.

[40568] Klemsz, M.; Hromas, R.; Raskind, W.; Bruno, E.; Hoffman, R.: PE-1,

[40569] a novel ETS oncogene family member, localizes to chromosome 1q21-q23. Genomics 20:

[40570] 291-294, 1994.

[40571] 6294.

[40572] Kastury, K.; Li, J.; Druck, T.; Su, H.; Vogt, P. K.; Croce, C.

[40573] M.; Huebner, K.: The human homologue of the retroviral oncogene qin

[40574] maps to chromosome 14q13. Proc. Nat. Acad. Sci. 91: 3616-3618, 1994.

[40575] 6295.

[40576] Li, J.; Chang, H. W.; Lai, E.; Parker, E. J.; Vogt, P. K.: The

[40577] oncogene qin codes for a transcriptional repressor. Cancer Res. 55:

[40578] 5540-5544, 1995.

[40579] 6296.

[40580] Li, J.; Vogt, P. K.: The retroviral oncogene qin belongs to the

[40581] transcription factor family that includes the homeotic gene fork head. Proc.

[40582] Nat. Acad. Sci. 90: 4490-4494, 1993.

[40583] 6297.

[40584] Murphy, D. B.; Wiese, S.; Burfeind, P.; Schmundt, D.; Mattei, M.-G.;

[40585] Schulz-Schaeffer, W.; Thies, U.: Human brain factor 1, a new member

[40586] of the fork head gene family. Genomics 21: 551-557,

1994.

[40587] 6298.

[40588] Wiese, S.; Murphy, D. B.; Schlung, A.; Burfeind, P.; Schmundt,

[40589] D.; Schnulle, V.; Mattei, M.-G.; Thies, U.: The genes for human brain

[40590] factor 1 and 2, members of the fork head gene family, are clustered

[40591] on chromosome 14q. Biochim. Biophys. Acta 1262: 105-112, 1995.

[40592] 6299.

[40593] Bustelo, X. R.: Regulatory and signaling properties of the Vav

[40594] family. Molec. Cell. Biol. 20: 1461-1477, 2000.

[40595] 6300.

[40596] Bustelo, X. R.; Barbacid, M.: Tyrosine phosphorylation of the

[40597] VAV proto-oncogene product in activated B cells. Science 256: 1196-1199,

[40598] 1992.

[40599] 6301.

[40600] Denkinger, D. J.; Borges, C. R.; Butler, C. L.; Cushman, A. M.;

[40601] Kawahara, R. S.: Genomic organization and regulation of the vav proto-oncogene. Biochim.

[40602] Biophys. Acta 1491: 253-262, 2000.

[40603] 6302.

[40604] Fackler, O. T.; Luo, W.; Geyer, M.; Alberts, A. S.; Peterlin, B.

[40605] M.: Activation of Vav by Nef induces cytoskeletal rearrangements

[40606] and downstream effector functions. Molec. Cell 3: 729-739, 1999.

[40607] 6303.

[40608] Fischer, K.-D.; Zmuidzinas, A.; Gardner, S.; Barbacid, M.; Bernstein,

[40609] A.; Guidos, C.: Defective T-cell receptor signalling and positive

[40610] selection of Vav-deficient CD4(+) CD8(+) thymocytes. Nature 374:

[40611] 474-477, 1995.

[40612] 6304.

[40613] Katzav, S.; Martin-Zanca, D.; Barbacid, M.: VAV, a novel human
[40614] oncogene derived from a locus ubiquitously expressed in hematopoietic
[40615] cells. EMBO J. 8: 2283-2290, 1989.

[40616] 6305.

[40617] Martinerie, C.; Cannizzaro, L. A.; Croce, C. M.; Huebner, K.; Katzav,
[40618] S.; Barbacid, M.: The human VAV proto-oncogene maps to chromosome
[40619] region 19p12-19p13.2. Hum. Genet. 86: 65-68, 1990.

[40620] 6306.

[40621] Moores, S. L.; Selfors, L. M.; Fredericks, J.; Breit, T.; Fu-jikawa,
[40622] K.; Alt, F. W.; Brugge, J. S.; Swat, W.: Vav family proteins couple
[40623] to diverse cell surface receptors. Molec. Cell. Biol. 20: 6364-6373,
[40624] 2000.

[40625] 6307.

[40626] Tarakhovsky, A.; Turner, M.; Schaal, S.; Mee, P. J.; Duddy,

L.

[40627] P.; Rajewsky, K.; Tybulewicz, V. L. J.: Defective antigen receptor-mediated

[40628] proliferation of B and T cells in the absence of Vav. Nature 374:

[40629] 467-470, 1995.

[40630] 6308.

[40631] Zhang, R.; Alt, F. W.; Davidson, L.; Orkin, S. H.; Swat, W.:

[40632] Defective signalling through the T- and B-cell antigen receptors in

[40633] lymphoid cells lacking the vav proto-oncogene. Nature 374: 470-473,

[40634] 1995.

[40635] 6309.

[40636] Fukuhara, S.; Rowley, J. D.; Variakojis, D.; Sweet, D. L.: Chromosome

[40637] abnormalities in poorly differentiated lymphocytic lymphoma. Cancer

[40638] Res. 39: 3119-3128, 1979.

[40639] 6310.

[40640] Ohno, H.; Fukuhara, S.; Takahashi, R.; Mihara, K.;

Sugiyama, T.;

[40641] Doi, S.; Uchino, H.; Toyoshima, K.: c-yes and bcl-2 genes located

[40642] on 18q21.3 in a follicular lymphoma cell line carrying a t(14;18)

[40643] chromosomal translocation. Int. J. Cancer 39: 785-788, 1987.

[40644] 6311.

[40645] Semba, K.; Nishizawa, M.; Satoh, H.; Fukushige, S.; Yoshida, M.

[40646] C.; Sasaki, M.; Matsubara, K.; Yamamoto, T.; Toyoshima, K.: Nucleotide

[40647] sequence and chromosomal mapping of the human c-yes-2 gene. Jpn.

[40648] J. Cancer Res. 79: 710-717, 1988.

[40649] 6312.

[40650] Semba, K.; Yamanashi, Y.; Nishizawa, M.; Sukegawa, J.; Yoshida,

[40651] M.; Sasaki, M.; Yamamoto, T.; Toyoshima, K.: Location of the c-yes

[40652] gene on the human chromosome and its expression in various tissues. Science 227:

[40653] 1038-1040, 1985.

[40654] 6313.

[40655] Silverman, G. A.; Kuo, W.-L.; Taillon-Miller, P.; Gray, J. W.:

[40656] Chromosomal reassignment: YACs containing both YES1 and thymidylate

[40657] synthase map to the short arm of chromosome 18. Genomics 15: 442-445,

[40658] 1993.

[40659] 6314.

[40660] Sukegawa, J.; Semba, K.; Yamanashi, Y.; Nishizawa, M.; Miyajima,

[40661] N.; Yamamoto, T.; Toyoshima, K.: Characterization of cDNA clones

[40662] for the human c-yes gene. Molec. Cell. Biol. 7: 41-47, 1987.

[40663] 6315.

[40664] Alves, S. E.; Lopez, V.; McEwen, B. S.; Weiland, N. G.: Differential

[40665] colocalization of estrogen receptor beta (ER-beta) with oxytocin and

[40666] vasopressin in the paraventricular and supraoptic nuclei of

the female
[40667] rat brain: an immunocytochemical study. Proc. Nat. Acad. Sci. 95:
[40668] 3281-3286, 1998.

[40669] 6316.

[40670] Brownstein, M. J.; Russell, J. T.; Gainer, H.: Synthesis, transport,
[40671] and release of posterior pituitary hormones. Science 207: 373-378,
[40672] 1980.

[40673] 6317.

[40674] Gopal Rao, V. V. N.; Loffler, C.; Battey, J.; Hansmann, I.: The
[40675] human gene for oxytocin-neurophysin I (OXT) is physi-cally mapped to
[40676] chromosome 20p13 by in situ hybridization. Cytogenet. Cell Genet. 61:
[40677] 271-273, 1992.

[40678] 6318.

[40679] Marini, J. C.; Nelson, K. K.; Battey, J.; Siracusa, L. D.: The

[40680] pituitary hormones arginine vasopressin-neurophysin II

and oxytocin-neurophysin

[40681] I show close linkage with interleukin-1 on mouse chro-
mosome 2. Genomics 15:

[40682] 200-202, 1993.

[40683] 6319.

[40684] Inoue, T.; Kimura, T.; Azuma, C.; Inazawa, J.; Takemura,
M.; Kikuchi,

[40685] T.; Kubota, Y.; Ogita, K.; Saji, F.: Structural organization of
the

[40686] human oxytocin receptor gene. J. Biol. Chem. 269:
32451-32456, 1994.

[40687] 6320.

[40688] Kimura, T.; Tanizawa, O.; Mori, K.; Brownstein, M. J.;
Okayama,

[40689] H.: Structure and expression of a human oxytocin recep-
tor. Nature 356:

[40690] 526-529, 1992.

[40691] 6321.

[40692] Simmons, C. F., Jr.; Clancy, T. E.; Quan, R.; Knoll, J. H. M.:

[40693] The oxytocin receptor gene (OXTR) localizes to human
chromosome 3p25

[40694] by fluorescence in situ hybridization and PCR analysis of somatic
[40695] cell hybrids. Genomics 26: 623-625, 1995.

[40696] 6322.

[40697] Yang, M.; Wang, W.; Zhong, M.; Philippi, A.; Lichtarge, O.; Sanborn,
[40698] B. M.: Lysine 270 in the third intracellular domain of the oxytocin
[40699] receptor is an important determinant for G-alpha-q coupling specificity. Molec.
[40700] Endocr. 16: 814-823, 2002.

[40701] 6323.

[40702] Cotlier, E.: Cafe-au-lait spots of the fundus in neurofibromatosis. Arch.
[40703] Ophthal. 95: 1990-1992, 1977.

[40704] 6324.

[40705] Craddock, G. R., Jr.; Challa, V. R.; Dean, R. H.: Neurofibromatosis
[40706] and renal artery stenosis: a case of familial incidence. J. Vasc.
[40707] Surg. 8: 489-494, 1988.

[40708] 6325.

[40709] Crawford, A. H.: Neurofibromatosis in children. Acta Orthop.

[40710] Scand. 57 (suppl. 218): 9–60, 1986.

[40711] 6326.

[40712] Crowe, F. W.: Axillary freckling as a diagnostic aid in neurofibromatosis. Ann.

[40713] Intern. Med. 61: 1142–1143, 1964.

[40714] 6327.

[40715] Crowe, F. W.; Schull, W. J.; Neel, J. V.: A Clinical, Pathological

[40716] and Genetic Study of Multiple Neurofibromatosis. Springfield, Ill.:

[40717] Charles C Thomas (pub.) 1956.

[40718] 6328.

[40719] D'Agostino, A. N.; Soule, E. H.; Miller, R. H.: Sarcomas of the

[40720] peripheral nerves and somatic soft tissue associated with multiple

[40721] neurofibromatosis. Cancer 16: 1015–1027, 1963.

[40722] 6329.

[40723] Daston, M. M.; Scrable, H.; Nordlund, M.; Sturbaum, A. K.; Nissen,

[40724] L. M.; Ratner, N.: The protein product of the neurofibromatosis type

[40725] 1 gene is expressed at highest abundance in neurons, Schwann cells,

[40726] and oligodendrocytes. Neuron 8: 415-428, 1992.

[40727] 6330.

[40728] DeBella, K.; Szudek, J.; Friedman, J. M.: Use of the National

[40729] Institutes of Health Criteria for Diagnosis of Neurofibromatosis 1

[40730] in children. Pediatrics 105: 608-614, 2000.

[40731] 6331.

[40732] DeClue, J. E.; Cohen, B. D.; Lowy, D. R.: Identification and

[40733] characterization of the neurofibromatosis type 1 protein product. Proc.

[40734] Nat. Acad. Sci. 88: 9914-9918, 1991.

[40735] 6332.

[40736] DeClue, J. E.; Papageorge, A. G.; Fletcher, J. A.; Diehl, S. R.;

[40737] Ratner, N.; Vass, W. C.; Lowy, D. R.: Abnormal regulation of mammalian

[40738] p21(ras) contributes to malignant tumor growth in von Recklinghausen

[40739] (type 1) neurofibromatosis. Cell 69: 265-273, 1992.

[40740] 6333.

[40741] Dent, C. E.: Rickets and osteomalacia from renal tubular defects. J.

[40742] Bone Joint Surg. 34B: 266-274, 1952.

[40743] 6334.

[40744] Diehl, S. R.; Boehnke, M.; Erickson, R. P.; Ploughman, L. M.;

[40745] Seiler, K. A.; Lieberman, J. L.; Clarke, H. B.; Bruce, M. A.; Schorry,

[40746] E. K.; Pericak-Vance, M.; O'Connell, P.; Collins, F. S.: A re-fined

[40747] genetic map of the region of chromosome 17 surrounding the von Recklinghausen

[40748] neurofibromatosis (NF1) gene. Am. J. Hum. Genet. 44: 33-37, 1989.

[40749] 6335.

[40750] Diekmann, L.; Huther, W.; Pfeiffer, R. A.: Ungewoehnliche Erscheinungsformen

[40751] der Neurofibromatose (von Recklinghausensche Krankheit) im Kindesalter. Z.

[40752] Kinderheilk. 101: 191-222, 1967.

[40753] 6336.

[40754] Dietz, J. N.; Robbins, T.; Schwartz, C.; Cannon, L.; McLel-

[40755] lan, T.; Williamson, R.; Carey, J.; Johnson, J.; Kivlin, J.; Skolnick,

[40756] M. H.: Linkage analysis of neurofibromatosis: chromo-

[40757] somes 4 and 19. (Abstract) Am. J. Hum. Genet. 37: A193, 1985.

[40758] 6337.

[40759] DiLiberti, J. H.; Buist, N. R. M.; Rosenberg, N.; Andrews, T.

[40760] : Myotonic dystrophy, hyperparathyroidism and multiple endocrine adenomatosis

[40761] type 2A. (Abstract) Am. J. Hum. Genet. 34: 86A, 1982.

[40762] 6338.

[40763] Dorschner, M. O.; Sybert, V. P.; Weaver, M.; Pletcher, B. A.;

[40764] Stephens, K.: NF1 microdeletion breakpoints are clustered

at flanking
[40765] repetitive sequences. Hum. Molec. Genet. 9: 35-46, 2000.

[40766] 6339.

[40767] Dugoff, L.; Sujansky, E.: Neurofibromatosis type 1 and pregnancy. Am.
[40768] J. Med. Genet. 66: 7-10, 1996.

[40769] 6340.

[40770] Duncan, A. M. V.; Partington, M. W.; Soudek, D.: Neurofi-bromatosis
[40771] in a man with a ring 22: in situ hybridization studies. Can-cer Genet.
[40772] Cytogenet. 25: 169-174, 1987.

[40773] 6341.

[40774] Dunn, B. G.; Ferrell, R. E.; Riccardi, V. M.: A genetic linkage
[40775] study in 15 families of individuals with von Reckling-hausen neurofibromatosis. Am.

[40776] J. Med. Genet. 22: 403-407, 1985.

[40777] 6342.

[40778] Dunn, F. G.; DeCarvalho, J. G. R.; Kem, D. C.; Higgins, J. R.;

[40779] Frohlich, E. D.: Pheochromocytoma crisis induced by sar-
alasin: relation

[40780] of angiotensin analogue to catecholamine release. New
Eng. J. Med. 295:

[40781] 605-607, 1976.

[40782] 6343.

[40783] Easton, D. F.; Ponder, M. A.; Huson, S. M.; Ponder, B. A. J.:

[40784] An analysis of variation in expression of neurofibromato-
sis (NF) type

[40785] I (NF1): evidence for modifying genes. Am. J. Hum. Genet.
53: 305-313,

[40786] 1993.

[40787] 6344.

[40788] Edwards, J. H.: Personal Communication. Oxford, England
10/1990.

[40789] 6345.

[40790] Eichhorn, C.; Wendt, G.; Staudte, H.-W.; Gilsbach, J.-M.:

Dural
[40791] ectasia in Von Recklinghausen's disease of the lumbar spine: a case
[40792] report. J. Bone Joint Surg. 77: B834-B835, 1995.

[40793] 6346.

[40794] Eisenbarth, I.; Beyer, K.; Krone, W.; Assum, G.: Toward a survey
[40795] of somatic mutation of the NF1 gene in benign neurofibromas of patients
[40796] with neurofibromatosis type 1. Am. J. Hum. Genet. 66: 393-401, 2000.

[40797] 6347.

[40798] Erickson, R. P.; Woolliscroft, J.; Allen, R. J.: Familial occurrence
[40799] of intracranial arterial occlusive disease (moyamoya) in neurofibromatosis. Clin.
[40800] Genet. 18: 191-196, 1980.

[40801] 6348.

[40802] Estivill, X.; Lazaro, C.; Casals, T.; Ravella, A.: Recurrence
[40803] of a nonsense mutation in the NF1 gene causing classical neurofibromatosis

[40804] type 1. Hum. Genet. 88: 185-188, 1991.

[40805] 6349.

[40806] Fabricant, R. N.; Todaro, G. J.: Increased serum levels of nerve

[40807] growth factor in von Recklinghausen's disease. Arch. Neurol. 38:

[40808] 401-405, 1981.

[40809] 6350.

[40810] Owerbach, D.; Rutter, W. J.; Cooke, N. E.; Martial, J. A.; Shows,

[40811] T. B.: The prolactin gene is located on chromosome 6 in humans. Science 212:

[40812] 815-816, 1981.

[40813] 6351.

[40814] Shome, B.; Parlow, A. F.: Human pituitary prolactin (hPRL): the

[40815] entire linear amino acid sequence. J. Clin. Endocr. Metab. 45: 1112-1115,

[40816] 1977.

[40817] 6352.

[40818] Sun, Z.; Lee, M. S.; Rhee, H. K.; Arrandale, J. M.; Dannies, P.

[40819] S.: Inefficient secretion of human H27A-prolactin, a mutant that

[40820] does not bind Zn(2+). Molec. Endocr. 11: 1544-1551, 1997.

[40821] 6353.

[40822] Taggart, R. T.; Mohandas, T. K.; Bell, G. I.: Assignment of the

[40823] human preprogastricsin (PGC) to chromosome 6 and regional localization

[40824] of PGC (6pter-p21.1), prolactin PRL (6pter-p21.1). (Abstract) Cytogenet.

[40825] Cell Genet. 46: 701-702, 1987.

[40826] 6354.

[40827] Truong, A. T.; Duez, C.; Belayew, A.; Renard, A.; Pictet, R.;

[40828] Bell, G. I.; Martial, J. A.: Isolation and characterization of the

[40829] human prolactin gene. EMBO J. 3: 429-437, 1984.

[40830] 6355.

[40831] Slate, D. L.; Ruddle, F. H.: Fibroblast interferon in man is

[40832] coded by two loci on separate chromosomes. Cell 16: 171-180, 1979.

[40833] 6356.

[40834] Rosenberg, M.; RayChaudhury, A.; Shows, T. B.; Le Beau, M. M.;

[40835] Fuchs, E.: A group of type I keratin genes on human chro-

[40836] mosome 17: characterization and expression. Molec. Cell. Biol. 8: 722-736,

[40837] 1988.

[40838] 6357.

[40839] Gross, J. A.; Johnston, J.; Mudri, S.; Enselman, R.; Dillon, S.

[40840] R.; Madden, K.; Xu, W.; Parrish-Novak, J.; Foster, D.; Lofton-Day,

[40841] C.; Moore, M.; Littau, A.; Grossman, A.; Haugen, H.; Foley, K.; Blumberg,

[40842] H.; Harrison, K.; Kindsvogel, W.; Clegg, C. H.: TACI and BCMA are

[40843] receptors for a TNF homologue implicated in B-cell au-

[40844] toimmune disease. Nature 404: 995-999, 2000.

[40845] 6358.

[40846] Watkins, D.; Rosenblatt, D. S.: Genetic heterogeneity among patients

[40847] with methylcobalamin deficiency. J. Clin. Invest. 81: 1690-1694,

[40848] 1988.

[40849] 6359.

[40850] Sato, S.; Fujita, N.; Kurihara, T.; Kuwano, R.; Sakimura, K.; Takahashi,

[40851] Y.; Miyatake, T.: cDNA cloning and amino acid sequence for human

[40852] myelin-associated glycoprotein. Biochem. Biophys. Res. Commun. 163:

[40853] 1473-1480, 1989.

[40854] 6360.

[40855] Lafage, M.; Clauss, I.; Couez, D.; Simonetti, J.; Wathelet, M.

[40856] G.; Huez, G.: The interferon- and virus-inducible IFI-56K and IFI-54K

[40857] genes are located on human chromosome 10 at bands q23-q24. Genomics 13:

[40858] 458-460, 1992.

[40859] 6361.

[40860] de Wit, T. P. M.; Morton, H. C.; Capel, P. J. A.; van de Winkel,

[40861] J. G. J.: Structure of the gene for the human myeloid IgA Fc receptor

[40862] (CD89). J. Immun. 155: 1203-1209, 1995.

[40863] 6362.

[40864] Kremer, E. J.; Kalatzis, V.; Baker, E.; Callen, D. F.; Sutherland,

[40865] G. R.; Maliszewski, C. R.: The gene for the human IgA Fc receptor

[40866] maps to 19q13.4. Hum. Genet. 89: 107-108, 1992.

[40867] 6363.

[40868] Maliszewski, C. R.; March, C. J.; Schoenborn, M. A.; Gimpel, S.;

[40869] Shen, L.: Expression cloning of a human Fc receptor for IgA. J.

[40870] Exp. Med. 172: 1665-1672, 1990.

[40871] 6364.

[40872] Monteiro, R. C.; Hostoffer, R. W.; Cooper, M. D.; Bonner, J. R.;

[40873] Gartland, G. L.; Kubagawa, H.: Definition of immunoglob-
ulin A receptors
[40874] on eosinophils and their enhanced expression in allergic
individuals. J.
[40875] Clin. Invest. 92: 1681-1685, 1993.

[40876] 6365.

[40877] Narita, I.; Goto, S.; Saito, N.; Sakatsume, M.; Jin, S.; Omori,

[40878] K.; Gejyo, F.: Genetic polymorphisms in the promoter and
5-prime
[40879] UTR region of the Fc alpha receptor (CD89) are not asso-
ciated with
[40880] a risk of IgA nephropathy. J. Hum. Genet. 46: 694-698,
2001.
[40881] 6366.

[40882] Pleass, R. J.; Andrews, P. D.; Kerr, M. A.; Woof, J. M.: Alter-
native
[40883] splicing of the human IgA Fc receptor CD89 in neutrophils
and eosinophils. Biochem.
[40884] J. 318: 771-777, 1996.

[40885] 6367.

[40886] Shimokawa, T.; Tsuge, T.; Okumura, K.; Ra, C.: Identifica-

tion
[40887] and characterization of the promoter for the gene encoding the human
[40888] myeloid IgA Fc receptor (Fc-alpha-R, CD89). Immunogenetics 51: 945-954,
[40889] 2000.

[40890] 6368.

[40891] Tsuge, T.; Shimokawa, T.; Horikoshi, S.; Tomino, Y.; Ra, C.: Polymorphism
[40892] in promoter region of Fc-alpha receptor gene in patients with IgA
[40893] nephropathy. Hum. Genet. 108: 128-133, 2001.

[40894] 6369.

[40895] Lalley, P. A.; Sakaguchi, A. Y.; Eddy, R. L.; Honey, N. H.; Bell,
[40896] G. I.; Shen, L.-P.; Rutter, W. J.; Jacobs, J. W.; Heinrich, G.; Chin,
[40897] W. W.; Naylor, S. L.: Mapping polypeptide hormone genes in the mouse:
[40898] somatostatin, glucagon, calcitonin, and parathyroid hormone. Cytogenet.
[40899] Cell Genet. 44: 92-97, 1987.

[40900] 6370.

[40901] Dillon, J. S.; Tanizawa, Y.; Wheeler, M. B.; Leng, X.-H.; Ligon,

[40902] B. B.; Rabin, D. U.; Yoo-Warren, H.; Permutt, M. A.; Boyd, A. E.,

[40903] III: Cloning and functional expression of the human glucagon-like

[40904] peptide-1 (GLP-1) receptor. Endocrinology 133: 1907-1910, 1993.

[40905] 6371.

[40906] Kershaw, E. E.; Chua, S. C., Jr.; Leibel, R. L.: Localization

[40907] of a (CA)n repeat in glucagon-like peptide-1 receptor gene (Glp1r)

[40908] to proximal mouse chromosome 17 and its linkage to other markers. Mammalian

[40909] Genome 6: 301-303, 1995.

[40910] 6372.

[40911] Stoffel, M.; Espinosa, R., III; Le Beau, M. M.; Bell, G. I.: Human

[40912] glucagon-like peptide-1 receptor gene: localization to chromosome

[40913] band 6p21 by fluorescence in situ hybridization and linkage of a highly

[40914] polymorphic simple tandem repeat DNA polymorphism to other markers

[40915] on chromosome 6. Diabetes 42: 1215-1218, 1993.

[40916] 6373.

[40917] Thorens, B.: Expression cloning of the pancreatic beta cell receptor

[40918] for the gluco-incretin hormone glucagon-like peptide 1. Proc. Nat.

[40919] Acad. Sci. 89: 8641-8645, 1992.

[40920] 6374.

[40921] Cohen, P.; Miyazaki, M.; Socci, N. D.; Hagge-Greenberg, A. Liedtke,

[40922] W.; Soukas, A. A.; Sharma, R.; Hudgins, L. C.; Ntambi, J. M.; Friedman,

[40923] J. M.: Role for stearoyl-CoA desaturase-1 in leptin-mediated weight

[40924] loss. Science 297: 240-243, 2002.

[40925] 6375.

[40926] Coleman, D. L.: Obesity genes: beneficial effects in het-

erozygous

[40927] mice. Science 203: 663-665, 1979.

[40928] 6376.

[40929] Comuzzie, A. G.; Hixson, J. E.; Almasy, L.; Mitchell, B. D.; Mahaney,

[40930] M. C.; Dyer, T. D.; Stern, M. P.; MacCluer, J. W.; Blangero, J.:

[40931] A major quantitative trait locus determining serum leptin levels and

[40932] fat mass is located on human chromosome 2. Nature Genet. 15: 273-276,

[40933] 1997.

[40934] 6377.

[40935] Fahsold, R.; Hoffmeyer, S.; Mischung, C.; Gille, C.; Ehlers, C.;

[40936] Kucukceylan, N.; Abdel-Nour, M.; Gewies, A.; Peters, H.; Kaufmann,

[40937] D.; Buske, A.; Tinschert, S.; Nurnberg, P.: Minor lesion mutational

[40938] spectrum of the entire NF1 gene does not explain its high mutability

[40939] but points to a functional domain upstream of the GAP-

related domain. Am.

[40940] J. Hum. Genet. 66: 790-818, 2000.

[40941] 6378.

[40942] Considine, R. V.; Considine, E. L.; Williams, C. J.; Nyce, M.

[40943] R.; Magosin, S. A.; Bauer, T. L.; Rosato, E. L.; Colberg, J.; Caro,

[40944] J. F.: Evidence against either a premature stop codon or the absence

[40945] of obese gene mRNA in human obesity. J. Clin. Invest. 95: 2986-2988,

[40946] 1995.

[40947] 6379.

[40948] Ducy, P.; Amling, M.; Takeda, S.; Priemel, M.; Schilling, A. F.;

[40949] Beil, F. T.; Shen, J.; Vinson, C.; Rueger, J. M., Karsenty, G.: Leptin

[40950] inhibits bone formation through a hypothalamic relay: a central control

[40951] of bone mass. Cell 100: 197-207, 2000.

[40952] 6380.

[40953] Duggirala, R.; Stern, M. P.; Mitchell, B. D.; Reinhart, L. J.;

[40954] Shipman, P. A.; Uresandi, O. C.; Chung, W. K.; Leibel, R. L.; Hales,
[40955] C. N.; O'Connell, P.; Blangero, J.: Quantitative variation in obesity-related
[40956] traits and insulin precursors linked to the OB gene region on human
[40957] chromosome 7. Am. J. Hum. Genet. 59: 694-703, 1996.

[40958] 6381.

[40959] Faggioni, R.; Jones-Carson, J.; Reed, D. A.; Dinarello, C. A.;

[40960] Feingold, K. R.; Grunfeld, C.; Fantuzzi, G.: Leptin-deficient (ob/ob)
[40961] mice are protected from T cell-mediated hepatotoxicity: role of tumor
[40962] necrosis factor alpha and IL-18. Proc. Nat. Acad. Sci. 97: 2367-2372,
[40963] 2000.

[40964] 6382.

[40965] Farooqi, I. S.; Jebb, S. A.; Langmack, G.; Lawrence, E.; Cheetham,
[40966] C. H.; Prentice, A. M.; Hughes, I. A.; McCamish, M. A.; O'Rahilly,

[40967] S.: Effects of recombinant leptin therapy in a child with congenital
[40968] leptin deficiency. New Eng. J. Med. 341: 879–884, 1999.

[40969] 6383.

[40970] Farooqi, I. S.; Keogh, J. M.; Kamath, S.; Jones, S.; Gibson, W.
[40971] T.; Trussell, R.; Jebb, S. A.; Lip, G. Y. H.; O'Rahilly, S.: Partial
[40972] leptin deficiency and human adiposity. Nature 414: 34–35, 2001.

[40973] 6384.

[40974] Forbes, S.; Bui, S.; Robinson, B. R.; Hochgeschwender, U.; Brennan,
[40975] M. B.: Integrated control of appetite and fat metabolism by the leptin–proopiomelanocortin
[40976] pathway. Proc. Nat. Acad. Sci. 98: 4233–4237, 2001.

[40977] 6385.

[40978] Frederich, R. C.; Lollmann, B.; Hamann, A.; Napolitano–Rosen,
[40979] A.; Kahn, B. B.; Lowell, B. B.; Flier, J. S.: Expression of ob mRNA

[40980] and its encoded protein in rodents: impact of nutrition and obesity. J.

[40981] Clin. Invest. 96: 1658-1663, 1995.

[40982] 6386.

[40983] Friedman, J. M.: Personal Communication. New York, N. Y. 12/2/1994.

[40984] 6387.

[40985] Friedman, J. M.; Halaas, J. L.: Leptin and the regulation of

[40986] body weight in mammals. Nature 395: 763-770, 1998.

[40987] 6388.

[40988] Friedman, J. M.; Leibel, R. L.; Siegel, D. S.; Walsh, J.; Ba-hary,

[40989] N.: Molecular mapping of the mouse ob mutation. Ge-nomics 11: 1054-1062,

[40990] 1991.

[40991] 6389.

[40992] Geffroy, S.; De Vos, P.; Staels, B.; Duban, B.; Auwerx, J.; de

[40993] Martinville, B.: Localization of the human OB gene (OBS) to chromosome

[40994] 7q32 by fluorescence in situ hybridization. Genomics 28:

603-604,

[40995] 1995.

[40996] 6390.

[40997] Gloaguen, I.; Costa, P.; Demartis, A.; Lazzaro, D.; Di Marco,

[40998] A.; Graziani, R.; Paonessa, G.; Chen, F.; Rosenblum, C. I.; Van der

[40999] Ploeg, L. H. T.; Cortese, R.; Ciliberto, G.; Laufer, R.: Ciliary

[41000] neurotrophic factor corrects obesity and diabetes associated with

[41001] leptin deficiency and resistance. Proc. Nat. Acad. Sci. 94: 6456-6461,

[41002] 1997.

[41003] 6391.

[41004] Gong, D.-W.; Bi, S.; Pratley, R. E.; Weintraub, B. D.: Genomic

[41005] structure and promoter analysis of the human obese gene. J. Biol.

[41006] Chem. 271: 3971-3974, 1996.

[41007] 6392.

[41008] Green, E. D.; Maffei, M.; Braden, V. V.; Proenca, R.; DeSilva,

[41009] U.; Zhang, Y.; Chua, S. C., Jr.; Leibel, R. L.; Weissenbach, J.; Friedman,
[41010] J. M.: The human obese (OB) gene: RNA expression pattern and mapping
[41011] on the physical, cytogenetic, and genetic maps of chromosome 7. Genome
[41012] Res. 5: 5-12, 1995.

[41013] 6393.

[41014] Haffner, S. M.; Miettinen, H.; Karhapaa, P.; Mykkanen, L.; Laakso,
[41015] M.: Leptin concentrations, sex hormones, and cortisol in nondiabetic
[41016] men. J. Clin. Endocr. Metab. 82: 1807-1809, 1997.

[41017] 6394.

[41018] Hager, J.; Francke, S.; Clement, K.; Dina, C.; Basdevant, A.;

[41019] Guy-Grand, B.; Froguel, P.: A polymorphism in the 5-prime UTR region
[41020] of the human OB gene is associated with morbid obesity and low leptin
[41021] levels. (Abstract) Medizinische Genetik 9: 10 only, 1997.

[41022] 6395.

[41023] Halaas, J. L.; Gajiwala, K. S.; Maffei, M.; Cohen, S. L.; Chait,

[41024] B. T.; Rabinowitz, D.; Lallone, R. L.; Burley, S. K.; Friedman, J.

[41025] M.: Weight-reducing effects on the plasma protein encoded by the

[41026] obese gene. Science 269: 543-546, 1995.

[41027] 6396.

[41028] Hamilton, B. S.; Paglia, D.; Kwan, A. Y. M.; Deitel, M.: Increased

[41029] obese mRNA expression in omental fat cells from massively obese humans. Nature

[41030] Med. 1: 953-956, 1995.

[41031] 6397.

[41032] Harigaya, A.; Nagashima, K.; Nako, Y.; Morikawa, A.: Relationship

[41033] between concentration of serum leptin and fetal growth. J. Clin.

[41034] Endocr. Metab. 82: 3281-3284, 1997.

[41035] 6398.

[41036] He, Y.; Chen, H.; Quon, M. J.; Reitman, M.: The mouse 'obese'

[41037] gene: genomic organization, promoter activity, and activation by CCAAT/enhancer-binding
[41038] protein-alpha. J. Biol. Chem. 270: 28887-28891, 1995.

[41039] 6399.

[41040] Jabs, E. W.; Coss, C. A.; Hayflick, S. J.; Whitmore, T. E.; Pauli,
[41041] R. M.; Kirkpatrick, S. J.; Meyers, D. A.; Goldberg, R.; Day, D. W.;
[41042] Rosenbaum, K. N.: Chromosomal deletion 4p15.32-p14 in a Treacher
[41043] Collins syndrome patient: exclusion of the disease locus from and
[41044] mapping of anonymous DNA sequences to this region. Genomics 11:
[41045] 188-192, 1991.

[41046] 6400.

[41047] Marsh, K. L.; Dixon, J.; Dixon, M. J.: Mutations in the Treacher
[41048] Collins syndrome gene lead to mislocalization of the nucleolar protein
[41049] treacle. Hum. Molec. Genet. 7: 1795-1800, 1998.

[41050] 6401.

[41051] Splendore, A.; Silva, E. O.; Alonso, L. G.; Richieri-Costa, A.;

[41052] Alonso, N.; Rosa, A.; Carakushanky, G.; Cavalcanti, D. P.;
Brunoni,

[41053] D.; Passos-Bueno, M. R.: High mutation detection rate in
TCOF1 among

[41054] Treacher Collins syndrome patients reveals clustering of
mutations

[41055] and 16 novel pathogenic changes. Hum. Mutat. 16:
315-322, 2000.

[41056] 6402.

[41057] Wise, C. A.; Chiang, L. C.; Paznekas, W. A.; Sharma, M.;
Musy,

[41058] M. M.; Ashley, J. A.; Lovett, M.; Jabs, E. W.: TCOF1 gene
encodes

[41059] a putative nucleolar phosphoprotein that exhibits muta-
tions in Treacher

[41060] Collins syndrome throughout its coding region. Proc. Nat.
Acad. Sci. 94:

[41061] 3110-3115, 1997.

[41062] 6403.

[41063] Dahms, N. M.; Lobel, P.; Breitmeyer, J.; Chirgwin, J. M.; Kornfeld,
[41064] S.: 46 kd mannose 6-phosphate receptor: cloning, expression, and
[41065] homology to the 215 kd mannose 6-phosphate receptor. Cell 50: 181-192,
[41066] 1987.

[41067] 6404.

[41068] Ludwig, T.; Ruther, U.; Metzger, R.; Copeland, N. G.; Jenkins,
[41069] N. A.; Lobel, P.; Hoflack, B.: Gene and pseudogene of the mouse cation-dependent
[41070] mannose 6-phosphate receptor: genomic organization, expression, and
[41071] chromosomal localization. J. Biol. Chem. 267: 12211-12219, 1992.
[41072] 6405.

[41073] Pohlmann, R.; Boeker, M. W. C.; von Figura, K.: The two mannose
[41074] 6-phosphate receptors transport distinct complements of lysosomal
[41075] proteins. J. Biol. Chem. 270: 27311-27318, 1995.

[41076] 6406.

[41077] Pohlmann, R.; Nagel, G.; Schmidt, B.; Stein, M.; Lorkowski, G.;

[41078] Krentler, C.; Cully, J.; Meyer, H. E.; Grzeschik, K.-H.; Mersmann,

[41079] G.; Hasilik, A.; von Figura, K.: Cloning of a cDNA encoding the human

[41080] cation-dependent mannose 6-phosphate-specific receptor. Proc. Nat.

[41081] Acad. Sci. 84: 5575-5579, 1987.

[41082] 6407.

[41083] Roberts, D. L.; Weix, D. J.; Dahms, N. M.; Kim, J.-J. P.: Molecular

[41084] basis of lysosomal enzyme recognition: three-dimensional structure

[41085] of the cation-dependent mannose 6-phosphate receptor. Cell 93: 639-648,

[41086] 1998.

[41087] 6408.

[41088] Watanabe, H.; Grubb, J. H.; Sly, W. S.: The overexpressed human

[41089] 46-kDa mannose 6-phosphate receptor mediates endocytosis and sorting
[41090] of beta-glucuronidase. Proc. Nat. Acad. Sci. 87: 8036-8040, 1990.
[41091] 6409.

[41092] Bax, W. A.; Cluysenaer, O. J. J.; Bartelink, A. K. M.; Aerts, P.

[41093] C.; Ezekowitz, R. A. B.; van Dijk, H.: Association of familial deficiency
[41094] of mannose-binding lectin and meningococcal disease. Lancet 354:
[41095] 1094-1095, 1999.

[41096] 6410.

[41097] Davies, J.; Neth, O.; Alton, E.; Klein, N.; Turner, M.: Differential
[41098] binding of mannose-binding lectin to respiratory pathogens in cystic
[41099] fibrosis. Lancet 355: 1885-1886, 2000.

[41100] 6411.

[41101] Ezekowitz, R. A.; Day, L. E.; Herman, G. A.: A human mannose-binding
[41102] protein is an acute phase reactant that shares sequence

homology with
[41103] other vertebrate lectins. J. Exp. Med. 167: 1034-1046, 1988.
[41104] 6412.
[41105] Gabolde, M.; Hubert, D.; Guilloud-Bataille, M.; Lenaerts, C.; Feingold,
[41106] J.; Besmond, C.: The mannose binding lectin gene influences the severity
[41107] of chronic liver disease in cystic fibrosis. J. Med. Genet. 38:
[41108] 310-311, 2001.
[41109] 6413.
[41110] Garred, P.; Madsen, H. O.; Balslev, U.; Hofmann, B.; Pedersen,
[41111] C.; Gerstoft, J.; Svejgaard, A.: Susceptibility to HIV infection
[41112] and progression of AIDs in relation to variant alleles of mannose-binding
[41113] lectin. Lancet 349: 236-240, 1997.
[41114] 6414.
[41115] Garred, P.; Madsen, H. O.; Hofmann, B.; Svejgaard, A.: In-

creased
[41116] frequency of homozygosity of abnormal mannan-binding-protein alleles
[41117] in patients with suspected immunodeficiency. Lancet 346: 941-943,
[41118] 1995.

[41119] 6415.

[41120] Garred, P.; Pressler, T.; Madsen, H. O.; Frederiksen, B.; Svejgaard,
[41121] A.; Hoiby, N.; Schwartz, M.; Koch, C.: Association of mannose-binding
[41122] lectin gene heterogeneity with severity of lung disease and survival
[41123] in cystic fibrosis. J. Clin. Invest. 104: 431-437, 1999.

[41124] 6416.

[41125] Garred, P.; Thiel, S.; Madsen, H. O.; Ryder, L. P.; Jensenius,
[41126] J. C.; Svejgaard, A.: Gene frequency and partial protein characterization
[41127] of an allelic variant of mannan binding protein associated with low
[41128] serum concentrations. Clin. Exp. Immun. 90: 517-521,

1992.

[41129] 6417.

[41130] Wildin, R. S.; Scott, C. R.: Cbl-G: presentation, treatment,

[41131] and prolonged followup in a patient with absence of methionine synthase.

[41132] (Abstract) Am. J. Hum. Genet. 51: A357 only, 1992.

[41133] 6418.

[41134] Wilson, A.; Leclerc, D.; Saberi, F.; Campeau, E.; Hwang, H. Y.;

[41135] Shane, B.; Phillips, J. A., III; Rosenblatt, D. S.; Gravel, R. A.

[41136] : Functionally null mutations in patients with the cblG-variant form

[41137] of methionine synthase deficiency. Am. J. Hum. Genet. 63: 409-414,

[41138] 1998.

[41139] 6419.

[41140] Zhang, Z.-X.; Leclerc, D.; Gravel, R.; Rozen, R.: The methionine

[41141] synthase (Mtr) gene maps to proximal mouse chromosome 13. Mammalian

[41142] Genome 8: 787-797, 1997.

[41143] 6420.

[41144] Watkins, D.; Ru, M.; Hwang, H.-Y.; Kim, C. D.; Murray, A.; Philip,

[41145] N. S.; Kim, W.; Legakis, H.; Wai, T.; Hilton, J. F.; Ge, B.; Dore,

[41146] C.; Hosack, A.; Wilson, A.; Gravel, R. A.; Shane, B.; Hudson, T. J.;

[41147] Rosenblatt, D. S.: Hyperhomocysteinemia due to methionine synthase

[41148] deficiency, cblG: structure of the MTR gene, genotype diversity, and

[41149] recognition of a common mutation, P1173L. Am. J. Hum. Genet. 71:

[41150] 143-153, 2002.

[41151] 6421.

[41152] Spagnol, G.; Williams, M.; Srinivasan, J.; Golier, J.; Bauer, D.;

[41153] Lebo, R. V.; Latov, N.: Molecular cloning of human myelin-associated

[41154] glycoprotein. J. Neurosci. Res. 24: 137-142, 1989.

[41155] 6422.

[41156] Pham-Dinh, D.: Personal Communication. Paris, France 3/15/1994.

[41157] 6423.

[41158] Pham-Dinh, D.; Gaspera, B. D.; De Rosbo, N. K.; Dautigny, A.:

[41159] Structure of the human myelin/oligodendrocyte glycopro-tein gene and

[41160] multiple alternative spliced isoforms. Genomics 29: 345-352, 1995.

[41161] 6424.

[41162] Pham-Dinh, D.; Jones, E. P; Pitiot, G.; Della Gaspera, B.; Daubas,

[41163] P.; Mallet, J.; Le Paslier, D.; Lindahl, C. F.; Dautigny, A.: Physical

[41164] mapping of the human and mouse MOG gene at the distal end of the MHC

[41165] class Ib region. Immunogenetics 42: 386-391, 1995.

[41166] 6425.

[41167] Pham-Dinh, D.; Mattei, M. G.; Nussbaum, J. L.; Roussel, G.; Pontarotti,

[41168] P.; Roeckel, N.; Mather, I. H.; Artzt, K.; Lindahl, K. F.; Dautigny,

[41169] A.: Myelin/oligodendrocyte glycoprotein is a member of a subset of
[41170] the immunoglobulin superfamily encoded within the major histocompatibility
[41171] complex. Proc. nat. Acad. Sci. 90: 7990-7994, 1993.

[41172] 6426.

[41173] Roth, M.-P.; Malfroy, L.; Offer, C.; Sevin, J.; Enault, G.; Borot,
[41174] N.; Pontarotti, P.; Coppin, H.: The human myelin oligodendrocyte
[41175] glycoprotein (MOG) gene: complete nucleotide sequence and structural
[41176] characterization. Genomics 28: 241-250, 1995.

[41177] 6427.

[41178] Ballmaier, M.; Germeshausen, M.; Schulze, H.; Cherkaoui, K.; Lang,
[41179] S.; Gaudig, A.; Krukemeier, S.; Eilers, M.; Straub, G.; Welte, K.
[41180] : c-mpl mutations are the cause of congenital amegakaryocytic thrombocytopenia. Blood 97:
[41181] 139-146, 2001.

[41182] 6428.

[41183] Cwirla, S. E.; Balasubramanian, P.; Duffin, D. J.; Wagstrom, C.

[41184] R.; Gates, C. M.; Singer, S. C.; Davis, A. M.; Tansik, R. L.; Mattheakis,

[41185] L. C.; Boytos, C. M.; Schatz, P. J.; Baccanari, D. P.; Wrighton, N.

[41186] C.; Barrett, R. W.; Dower, W. J.: Peptide agonist of the thrombopoietin

[41187] receptor as potent as the natural cytokine. Science 276: 1696-1699,

[41188] 1997.

[41189] 6429.

[41190] Gurney, A. L.; Carver-Moore, K.; de Sauvage, F. J.; Moore, M. W.

[41191] : Thrombocytopenia in c-mpl-deficient mice. Science 265: 1445-1447,

[41192] 1994.

[41193] 6430.

[41194] Ihara, K.; Ishii, E.; Eguchi, M.; Takada, H.; Suminoe, A.; Good,

[41195] R. A.; Hara, T.: Identification of mutations in the c-mpl gene in
[41196] congenital amegakaryocytic thrombocytopenia. Proc. Nat. Acad. Sci. 96:
[41197] 3132-3136, 1999.

[41198] 6431.

[41199] Kimura, S.; Roberts, A. W.; Metcalf, D.; Alexander, W. S.: Hematopoietic
[41200] stem cell deficiencies in mice lacking c-Mpl, the receptor for thrombopoietin. Proc.
[41201] Nat. Acad. Sci. 95: 1195-1200, 1998.

[41202] 6432.

[41203] Le Coniat, M.; Souyri, M.; Vigon, I.; Wendling, F.; Tam-
bourin,
[41204] P.; Berger, R.: The human homolog of the myeloprolifera-
tive virus
[41205] maps to chromosome band 1p34. Hum. Genet. 83:
194-196, 1989.
[41206] 6433.

[41207] Mignotte, V.; Vigon, I.; Boucher de Crevecoeur, E.; Romeo, P.-H.;

[41208] Lemarchandel, V.; Chretien, S.: Structure and transcription of the
[41209] human c-mpl gene (MPL). Genomics 20: 5-12, 1994.

[41210] 6434.

[41211] Moliterno, A. R.; Hankins, W. D.; Spivak, J. L.: Impaired expression
[41212] of the thrombopoietin receptor by platelets from patients with polycythemia
[41213] vera. New Eng. J. Med. 338: 572-580, 1998.

[41214] 6435.

[41215] Muraoka, K.; Ishii, E.; Tsuji, K.; Yamamoto, S.; Yamaguchi, H.;
[41216] Hara, T.; Koga, H.; Nakahata, T.; Miyazaki, S.: Defective response
[41217] to thrombopoietin and impaired expression of c-mpl mRNA of bone marrow
[41218] cells in congenital amegakaryocytic thrombocytopenia. Brit. J. Haemat. 96:
[41219] 287-292, 1997.

[41220] 6436.

[41221] Penciolelli, J. F.; Wendling, F.; Robert-Lezenes, J.; Barque,

[41222] J. F.; Tambourin, P.; Gisselbrecht, S.: Genetic analysis of myeloproliferative
[41223] leukemia virus, a novel acute leukemogenic replication-defective retrovirus. J.
[41224] Virol. 61: 579-583, 1987.

[41225] 6437.

[41226] Tonelli, R.; Scardovi, A. L.; Pession, A.; Strippoli, P.; Bonsi,
[41227] L.; Vitale, L.; Prete, A.; Locatelli, F.; Bagnara, G. P.; Paolucci,
[41228] G.: Compound heterozygosity for two different amino-acid substitution
[41229] mutations in the thrombopoietin receptor (c-mpl gene) in congenital
[41230] amegakaryocytic thrombocytopenia (CAMT). Hum. Genet. 107: 225-233,
[41231] 2000.

[41232] 6438.

[41233] van den Oudenrijn, S.; Bruin, M.; Folman, C. C.; Peters, M.; Faulkner,
[41234] L. B.; de Haas, M.; van dem Borne, A. E. G. K.: Mutations in the

[41235] thrombopoietin receptor, Mpl, in children with congenital amegakaryocytic
[41236] thrombocytopenia. Brit. J. Haemat. 110: 441-448, 2000.

[41237] 6439.

[41238] Vigon, I.; Mornon, J.-P.; Cocault, L.; Mitjavila, M.-T.; Tambourin,
[41239] P.; Gisselbrecht, S.; Souyri, M.: Molecular cloning and characterization
[41240] of MPL, the human homolog of the v-mpl oncogene: identification of
[41241] a member of the hematopoietic growth factor receptor superfamily. Proc.
[41242] Nat. Acad. Sci. 89: 5640-5644, 1992.

[41243] 6440.

[41244] Bamberger, A.-M.; Jenatschke, S.; Schulte, H. M.; Loning, T.; Bamberger,
[41245] C. M.: Leukemia inhibitory factor (LIF) stimulates the human HLA-G
[41246] promoter in JEG3 choriocarcinoma cells. J. Clin. Endocr. Metab. 85:
[41247] 3932-3936, 2000.

[41248] 6441.

[41249] Barasch, J.; Yang, J.; Ware, C. B.; Taga, T.; Yoshida, K.; Erdjument-Bromage,

[41250] H.; Tempst, P.; Parravicini, E.; Malach, S.; Aranoff, T.; Oliver,

[41251] J. A.: Mesenchymal to epithelial conversion in rat metanephros is

[41252] induced by LIF. Cell 99: 377-386, 1999.

[41253] 6442.

[41254] Bucan, M.; Gatalica, B.; Nolan, P.; Chung, A.; Leroux, A.; Grossman,

[41255] M. H.; Nadeau, J. H.; Emanuel, B. S.; Budarf, M.: Compara-

tive mapping

[41256] of 9 human chromosome 22q loci in the laboratory mouse. Hum. Molec.

[41257] Genet. 2: 1245-1252, 1993.

[41258] 6443.

[41259] Budarf, M.; Emanuel, B. S.; Mohandas, T.; Goeddel, D. V.; Lowe,

[41260] D. G.: Human differentiation-stimulating factor (leukemia inhibitory

[41261] factor, human interleukin DA) gene maps distal to the Ewing sarcoma
[41262] breakpoint on 22q. Cytogenet. Cell Genet. 52: 19-22, 1989.
[41263] 6444.

[41264] Doolittle, D.: Personal Communication. Bar Harbor, Me. 3/10/1992.
[41265] 6445.

[41266] Giovannini, M.; Djabali, M.; McElligott, D.; Selleri, L.; Evans,
[41267] G. A.: Tandem linkage of genes coding for leukemia inhibitory factor
[41268] (LIF) and oncostatin M (OSM) on human chromosome 22. Cytogenet. Cell
[41269] Genet. 64: 240-244, 1993.

[41270] 6446.

[41271] Gough, N. M.; Gearing, D. P.; King, J. A.; Willson, T. A.; Hilton,
[41272] D. J.; Nicola, N. A.; Metcalf, D.: Molecular cloning and expression
[41273] of the human homologue of the murine gene encoding myeloid leukemia-inhibitory

[41274] factor. Proc. Nat. Acad. Sci. 85: 2623-2627, 1988.

[41275] 6447.

[41276] Hanauer, A.; Mattei, M. G.; Mandel, J. L.: Presence of a TaqI

[41277] polymorphism in the human glutamate dehydrogenase (GLUD) gene on chromosome

[41278] 10. Nucleic Acids Res. 15: 6308 only, 1987.

[41279] 6448.

[41280] Fain, P. R.; Goldgar, D. E.; Wright, E. C.; Andersen, W. L.; Willard,

[41281] H. F.; van Tuinen, P.; Ledbetter, D. H.; Barker, D. F.: Mark-

ers tightly

[41282] linked to the NF1 locus on chromosome 17. (Abstract) Cy-

togenet. Cell

[41283] Genet. 51: 996, 1989.

[41284] 6449.

[41285] Fain, P. R.; Wright, E.; Willard, H. F.; Stephens, K.; Barker,

[41286] D. F.: The order of loci in the pericentric region of chro-

mosome

[41287] 17, based on evidence from physical and genetic break-

points. Am.

[41288] J. Hum. Genet. 44: 68-72, 1989.

[41289] 6450.

[41290] Fanconi, G.: Hochgradige Galaktose-intoleranz
(Galaktose-Diabetes)
[41291] bei einem Kinde mit Neurofibromatosis Recklinghausen.
Jahrb. Kinderheilk. 138:
[41292] 1-8, 1933.

[41293] 6451.

[41294] Faravelli, F.; Upadhyaya, M.; Osborn, M.; Huson, S. M.;
Hayward,
[41295] R.; Winter, R.: Unusual clustering of brain tumours in a
family with
[41296] NF1 and variable expression of cutaneous features. J.
Med. Genet. 36:
[41297] 893-896, 1999.

[41298] 6452.

[41299] Fialkow, P. J.; Sagebiel, R. W.; Gartler, S. M.; Rimoin, D. L.

[41300] : Multiple cell origin of hereditary neurofibromas. New
Eng. J. Med. 284:
[41301] 298-300, 1971.

[41302] 6453.

[41303] Fienman, N. L.; Yakovac, W. C.: Neurofibromatosis in childhood. J.

[41304] Pediat. 76: 339-346, 1970.

[41305] 6454.

[41306] Finley, J. L.; Dabbs, D. J.: Renal vascular smooth muscle proliferation

[41307] in neurofibromatosis. Hum. Path. 19: 107-110, 1988.

[41308] 6455.

[41309] Fitzpatrick, A. P.; Emanuel, R. W.: Familial neurofibro-matosis

[41310] and hypertrophic cardiomyopathy. Brit. Heart J. 60: 247-251, 1988.

[41311] 6456.

[41312] Fountain, J. W.; Wallace, M. R.; Brereton, A. M.; O'Connell, P.;

[41313] White, R. L.; Rich, D. C.; Ledbetter, D. H.; Leach, R. J.; Fournier,

[41314] R. E. K.; Menon, A. G.; Gusella, J. F.; Barker, D.; Stephens, K.;

[41315] Collins, F. S.: Physical mapping of the von Recklinghausen

neurofibromatosis
[41316] region on chromosome 17. Am. J. Hum. Genet. 44: 58-67, 1989.
[41317] 6457.
[41318] Fountain, J. W.; Wallace, M. R.; Bruce, M. A.; Seizinger, B. R.;
[41319] Menon, A. G.; Gusella, J. F.; Michels, V. V.: Schmidt, M. A.; Dewald,
[41320] G. W. and Collins, F. S.: Physical mapping of a translocation breakpoint
[41321] in neurofibromatosis. Science 244: 1085-1087, 1989.
[41322] 6458.
[41323] Friedman, J. M.; Birch, P.; Greene, C.; NNFF International Database
[41324] Participants: National Neurofibromatosis Foundation International
[41325] Database. Am. J. Med. Genet. 45: 88-91, 1993.
[41326] 6459.
[41327] Friedman, J. M.; Birch, P. H.: Type 1 neurofibromatosis: a descriptive
[41328] analysis of the disorder in 1,728 patients. Am. J. Med.

Genet. 70:
[41329] 138-143, 1997.

[41330] 6460.

[41331] Friedman, J. M.; Fialkow, P. J.; Greene, G. L.; Weinberg, M.
N.
[41332] : Probable clonal origin of neurofibrosarcoma in a patient
with hereditary
[41333] neurofibromatosis. J. Nat. Cancer Inst. 69: 1289-1292,
1982.
[41334] 6461.

[41335] Garty, B. Z.; Laor, A.; Danon, Y. L.: Neurofibromatosis type
[41336] 1 in Israel: survey of young adults. J. Med. Genet. 31:
853-857,
[41337] 1994.

[41338] 6462.

[41339] Gervasini, C.; Bentivegna, A.; Venturin, M.; Corrado, L.;
Larizza,
[41340] L.; Riva, P.: Tandem duplication of the NF1 gene detected
by high-resolution
[41341] FISH in the 17q11.2 region. Hum. Genet. 110: 314-321,
2002.

[41342] 6463.

[41343] Gitzelmann, R.: Deficiency of erythrocyte galactokinase in a

[41344] patient with galactose diabetes. Lancet II: 670-671, 1965.

[41345] 6464.

[41346] Goldgar, D. E.; Green, P.; Parry, D. M.; Mulvihill, J. J.: Multi-point

[41347] linkage analysis in neurofibromatosis type 1: an international collaboration. Am.

[41348] J. Hum. Genet. 44: 6-12, 1989.

[41349] 6465.

[41350] Graham, P. W.; Oehlschlaeger, F. H.: Articulating the Elephant

[41351] Man: Joseph Merrick and His Interpreters. Baltimore: Johns Hopkins

[41352] Univ. Press (pub.) 1992.

[41353] 6466.

[41354] Grant, W. M.; Walton, D. S.: Distinctive gonioscopic findings

[41355] in glaucoma due to neurofibromatosis. Arch. Ophthal. 79: 127-134,

[41356] 1968.

[41357] 6467.

[41358] Griffiths, D. F. R.; Williams, G. T.; Williams, E. D.: Duodenal

[41359] carcinoid tumours, phaeochromocytoma and neurofibro-
matosis: islet

[41360] cell tumour, phaeochromocytoma and the von Hippel-
Lindau complex:

[41361] two distinctive neuroendocrine syndromes. Quart. J. Med.
245: 769-782,

[41362] 1987.

[41363] 6468.

[41364] Guo, H.-F.; The, I.; Hannan, F.; Bernards, A.; Zhong, Y.:
Requirement

[41365] of Drosophila NF1 for activation of adenylyl cyclase by
PACAP38-like

[41366] neuropeptides. Science 276: 795-798, 1997.

[41367] 6469.

[41368] Gusella, J. F.: Personal Communication. Boston, Mass.
9/12/1987.

[41369] 6470.

[41370] Gutmann, D.; Aylsworth, A.; Carley, J.; Korf, B.; Marks, J.; Pyeritz,
[41371] R.; Rubenstein, A.; et al.: The diagnostic evaluation and multidisciplinary
[41372] management of neurofibromatosis 1 and neurofibromatosis 2. J.A.M.A. 278:
[41373] 51-57, 1997.

[41374] 6471.

[41375] Gutmann, D. H.: The neurofibromatoses: when less is more. Hum
[41376] Molec. Genet. 10: 747-755, 2001.

[41377] 6472.

[41378] Gutmann, D. H.; Loehr, A.; Zhang, Y.; Kim, J.; Henkemeyer, M.;
[41379] Cashen, A.: Haploinsufficiency for the neurofibromatosis 1 (NF1)
[41380] tumor supressor results in increased astrocyte proliferation. Oncogene 18:
[41381] 4450-4459, 1999.

[41382] 6473.

[41383] Gutmann, D. H.; Wood, D. L.; Collins, F. S.: Identification

of
[41384] the neurofibromatosis type 1 gene product. Proc. Nat. Acad. Sci. 88:
[41385] 9658-9662, 1991.

[41386]

[41387] 100. Gutmann, D. H.; Wu, Y. L.; Hedrick, N. M.; Zhu, Y.; Guha, A.;
[41388] Parada, L. F.: Heterozygosity for the neurofibromatosis 1 (NF1) tumor
[41389] suppressor results in abnormalities in cell attachment, spreading
[41390] and motility in astrocytes. Hum. Molec. Genet. 10: 3009-3016, 2001.

[41391]

[41392] 101. Gutzmer, R.; Herbst, R. A.; Mommert, S.; Kiehl, P.; Matiaske,
[41393] F.; Rutten, A.; Kapp, A.; Weiss, J.: Allelic loss at the neu-
rofibromatosis
[41394] type 1 (NF1) gene locus is frequent in desmoplastic neu-
rotropic melanoma. Hum.
[41395] Genet. 107: 357-361, 2000.

[41396]

[41397] 102. Hamilton, S. J.; Allard, M. F.; Friedman, J. M.: Cardiac findings
[41398] in an individual with neurofibromatosis 1 and sudden death. Am. J.
[41399] Med. Genet. 100: 95-99, 2001.

[41400]

[41401] 103. Hamilton, S. J.; Friedman, J. M.: Insights into the pathogenesis
[41402] of neurofibromatosis 1 vasculopathy. Clin. Genet. 58: 341-344, 2000.

[41403]

[41404] 104. Hashemian, H.: Familial fibromatosis of small intes-tine. Brit.
[41405] J. Surg. 40: 346-350, 1953.

[41406]

[41407] 105. Hayes, D. M.; Spurr, C. L.; Felts, J. H.; Miller, E. C., Jr.
[41408] : Von Recklinghausen's disease with massive intra-abdominal tumor
[41409] and spontaneous hypoglycemia: metabolic studies before and after perfusion
[41410] of abdominal cavity with nitrogen mustard. Metabolism

10: 183-199,
[41411] 1961.

[41412]

[41413] 106. Heim, R. A.; Kam-Morgan, L. N. W.; Binnie, C. G.;
Corns, D. D.;
[41414] Cayouette, M. C.; Farber, R. A.; Aylsworth, A. S.; Silverman,
L. M.;
[41415] Luce, M. C.: Distribution of 13 truncating mutations in the
neurofibromatosis
[41416] 1 gene. Hum. Molec. Genet. 4: 975-981, 1995.

[41417]

[41418] 107. Heim, R. A.; Silverman, L. M.; Farber, R. A.; Kam-
Morgan, L.
[41419] N. W.; Luce, M. C.: Screening for truncated NF1 proteins.
(Letter) Nature
[41420] Genet. 8: 218-219, 1994.

[41421]

[41422] 108. Herrmann, J.: Sarcomatous transformation in multi-
ple neurofibromatosis
[41423] (von Recklinghausen's disease). Ann. Surg. 131: 206-217,
1950.

[41424]

[41425] 109. Hinrichs, S. H.; Nerenberg, M.; Reynolds, R. K.; Khoury, G.;

[41426] Jay, G.: A transgenic mouse model for human neurofibromatosis. Science 237:

[41427] 1340-1343, 1987.

[41428]

[41429] 110. Hochberg, F. H.; DaSilva, A. B.; Galdabini, J.; Richardson, E.

[41430] P., Jr.: Gastrointestinal involvement in von Recklinghausen's neurofibromatosis. Neurology 24:

[41431] 1144-1151, 1974.

[41432]

[41433] 111. Hoffmeyer, S.; Nurnberg, P.; Ritter, H.; Fahsold, R.; Leistner,

[41434] W.; Kaufmann, D.; Krone, W.: Nearby stop codons in exons of the neurofibromatosis

[41435] type 1 gene are disparate splice effectors. Am. J. Hum. Genet. 62:

[41436] 269-277, 1998.

[41437]

[41438] 112. Hofman, K. J.; Harris, E. L.; Bryan, R. N.; Denckla, M. B.:
[41439] Neurofibromatosis type 1: the cognitive phenotype. J. Pediat. 124:
[41440] S1-S8, 1994.

[41441]

[41442] 113. Holt, J. F.: Neurofibromatosis in children. Am. J. Roentgen. 130:
[41443] 615-639, 1978.

[41444]

[41445] 114. Horiuchi, T.; Hatta, N.; Matsumoto, M.; Ohtsuka, H.; Collins,
[41446] F. S.; Kobayashi, Y.; Fujita, S.: Nonsense mutations at arg-1947
[41447] in two cases of familial neurofibromatosis type 1 in Japanese. Hum.
[41448] Genet. 93: 81-83, 1994.

[41449]

[41450] 115. Horwich, A.; Riccardi, V. M.; Francke, U.: Aqueductal stenosis
[41451] leading to hydrocephalus--an unusual manifestation of

neurofibromatosis. Am.
[41452] J. Med. Genet. 14: 577-581, 1983.

[41453]

[41454] 116. Howell, M.; Ford, P.: The True History of the Elephant Man.
[41455] New York: Penguin (pub.) 1980.

[41456]

[41457] 117. Hull, J.; Shackleton, S.; Harris, A.: The stop mutation R553X
[41458] in the CFTR gene results in exon skipping. Genomics 19: 362-364,
[41459] 1994.

[41460]

[41461] 118. Hunerbein, M.; Dohmoto, M.; Haensch, W.; Schlag, P. M.: Biliary
[41462] obstruction caused by a multifocal duodenal neurosar- coma in a patient
[41463] with von Recklinghausen's disease. Encoscopy 28: 405 only, 1996.

[41464]

[41465] 119. Huson, S. M.; Compston, D. A. S.; Clark, P.; Harper, P.

S.:
[41466] A genetic study of von Recklinghausen neurofibromatosis in south east
[41467] Wales. I. Prevalence, fitness, mutation rate, and effect of parental
[41468] transmission on severity. J. Med. Genet. 26: 704-711, 1989.
[41469]

[41470] 120. Huson, S. M.; Hughes, R. A. C.: The Neurofibro-matoses: a Pathogenetic
[41471] and Clinical Overview. London: Chapman and Hall Medical , 1994.
[41472]

[41473] 121. Huson, S. M.; Meredith, A. L.; Sarfarazi, M.; Shaw, D. J.; Compston,
[41474] D. A. S.; Harper, P. S.: Linkage analysis of peripheral neu-rofibromatosis
[41475] (Von Recklinghausen disease) and chromosome 19 mark-ers linked to myotonic
[41476] dystrophy. J. Med. Genet. 23: 55-57, 1986.
[41477]

[41478] 122. Ingordo, V.; D'Andria, G.; Mendecini, S.; Grecucci, M.;

Baglivo,

[41479] A.: Segmental neurofibromatosis: is it uncommon or underdiagnosed? Arch.

[41480] Derm. 131: 959-960, 1995.

[41481]

[41482] 123. Ingram, D. A.; Yang, F.-C.; Travers, J. B.; Wenning, M. J.; Hiatt,

[41483] K.; New, S.; Hood, A.; Shannon, K.; Williams, D. A.; Clapp, D. W.

[41484] : Genetic and biochemical evidence that haploinsufficiency of the

[41485] Nf1 tumor suppressor gene modulates melanocyte and mast cell fates

[41486] in vivo. J. Exp. Med. 191: 181-187, 2000.

[41487]

[41488] 124. Izumi, A. K.; Rosato, F. E.; Wood, M. G.: Von Recklinghausen's

[41489] disease associated with multiple neurolemmomas. Arch. Derm. 104:

[41490] 172-176, 1971.

[41491]

[41492] 125. Jadayel, D.; Fain, P.; Upadhyaya, M.; Ponder, M. A.;
Huson, S.
[41493] M.; Carey, J.; Fryer, A.; Mathew, C. G. P.; Barker, D. F.;
Ponder,
[41494] B. A. J.: Paternal origin of new mutations in Von Reckling-
hausen
[41495] neurofibromatosis. Nature 343: 558-559, 1990.

[41496]

[41497] 126. Jenne, D. E.; Tinschert, S.; Reimann, H.; Lasinger, W.;
Thiel,
[41498] G.; Hameister, H.; Kehrer-Sawatzki, H.: Molecular charac-
terization
[41499] and gene content of breakpoint boundaries in patients
with neurofibromatosis
[41500] type 1 with 17q11.2 microdeletions. Am. J. Hum. Genet.
69: 516-527,
[41501] 2001.

[41502]

[41503] 127. John, A. M.; Ruggieri, M.; Ferner, R.; Upadhyaya, M.:
A search
[41504] for evidence of somatic mutations in the NF1 gene. J. Med.
Genet. 37:

[41505] 44-49, 2000.

[41506]

[41507] 128. Johnson, B. L.; Charneco, D. R.: Cafe-au-lait spot in neurofibromatosis

[41508] and in normal individuals. Arch. Derm. 102: 442-446, 1970.

[41509]

[41510] 129. Johnson, M. R.; Look, A. T.; DeClue, J. E.; Valentine, M. B.;

[41511] Lowy, D. R.: Inactivation of the NF1 gene in human melanoma and neuroblastoma

[41512] cell lines without impaired regulation of GTP-Ras. Proc. Nat. Acad.

[41513] Sci. 90: 5539-5543, 1993.

[41514]

[41515] 130. Kalff, V.; Shapiro, B.; Lloyd, R.; Sisson, J. C.; Holland, K.;

[41516] Nakajo, M.; Beierwaltes, W. H.: The spectrum of pheochromocytoma

[41517] in hypertensive patients with neurofibromatosis. Arch. Intern. Med. 142:

[41518] 2092-2098, 1982.

[41519]

[41520] 131. Kaneko, Y.; Maseki, N.; Sakurai, M.; Shibuya, A.; Shi-
nohara,
[41521] T.; Fujimoto, T.; Kanno, H.; Nishikawa, A.: Chromosome
pattern in
[41522] juvenile chronic myelogenous leukemia, myelodysplastic
syndrome, and
[41523] acute leukemia associated with neurofibromatosis.
Leukemia 3: 36-41,
[41524] 1989.

[41525]

[41526] 132. Kanter, W. R.; Eldridge, R.; Fabricant, R.; Allen, J. C.;
Koerber,
[41527] T.: Central neurofibromatosis with bilateral acoustic neu-
roma: genetic,
[41528] clinical and biochemical distinctions from peripheral neu-
rofibromatosis. Neurology 30:
[41529] 851-859, 1980.

[41530]

[41531] 133. Kaplan, J.; Cushing, B.; Chang, C.-H.; Poland, R.;

Roscamp, J.;

[41532] Perrin, E.; Bhaya, N.: Familial T-cell lymphoblastic lymphoma: association

[41533] with von Recklinghausen neurofibromatosis and Gardner syndrome. Am.

[41534] J. Hemat. 12: 247-250, 1982.

[41535]

[41536] 134. Kaufmann, D.; Muller, R.; Bartelt, B.; Wolf, M.; Kunzi-Rapp,

[41537] K.; Hanemann, C. O.; Fahsold, R.; Hein, C.; Vogel, W.; Assum, G.:

[41538] Spinal neurofibromatosis without cafe-au-lait macules in two families

[41539] with null mutations of the NF1 gene. Am. J. Hum. Genet. 69: 1395-1400,

[41540] 2001.

[41541]

[41542] 135. Kayes, L. M.; Burke, W.; Riccardi, V. M.; Bennett, R.; Ehrlich,

[41543] P.; Rubenstein, A.; Stephens, K.: Deletions spanning the neurofibromatosis

[41544] I gene: identification and phenotype of five patients. Am.

J. Hum.

[41545] Genet. 54: 424-436, 1994.

[41546]

[41547] 136. King, A. A.; DeBaun, M. R.; Riccardi, V. M.; Gutmann, D. H.:

[41548] Malignant peripheral nerve sheath tumors in neurofibromatosis 1. Am.

[41549] J. Med. Genet. 93: 388-392, 2000.

[41550]

[41551] 137. Kittur, S. D.; Bagdon, M. M.; Lubs, M. L.; Phillips, J. A., III;

[41552] Murray, J. C.; Slaugenhaupt, S. A.; Chakravarti, A.; Adler, W. H.

[41553] : Linkage analysis of neurofibromatosis type 1, using chromosome 17

[41554] DNA markers. Am. J. Hum. Genet. 44: 48-50, 1989.

[41555]

[41556] 138. Klose, A.; Ahmadian, M. R.; Schuelke, M.; Scheffzek, K.; Hoffmeyer,

[41557] S.; Gewies, A.; Schmitz, F.; Kaufmann, D.; Peters, H.; Wittinghofer,

[41558] A.; Nurnberg, P.: Selective disactivation of neurofibromin GAP activity
[41559] in neurofibromatosis type 1 (NF1). Hum. Molec. Genet. 7: 1261-1268,
[41560] 1998.

[41561]

[41562] 139. Kluwe, L.; Friedrich, R. E.; Mautner, V. F.: Allelic loss of
[41563] the NF1 gene in NF1-associated plexiform neurofibromas. Cancer Genet.
[41564] Cytogenet. 113: 65-69, 1999.

[41565]

[41566] 140. Knight, W. A., III; Murphy, W. K.; Gottlieb, J. A.: Neu-rofibromatosis
[41567] associated with malignant neurofibromas. Arch. Derm. 107: 747-750,
[41568] 1973.

[41569]

[41570] 141. Kohn, S. R.: The elephant man. (Letter) New Eng. J. Med. 301:
[41571] 947, 1979.

[41572]

[41573] 142. Koivunen, J.; Yla-Outinen, H.; Korkiamaki, T.; Karvo-
nen, S.-L.;
[41574] Poyhonen, M.; Laato, M.; Karvonen, J.; Peltonen, S.; Pelto-
nen, J.
[41575] : New function for NF1 tumor suppressor. J. Invest. Derm.
114: 473-479,
[41576] 2000.

[41577]

[41578] 143. Konishi, K.; Nakamura, M.; Yamakawa, H.; Suzuki, H.;
Saruta,
[41579] T.; Hanaoka, H.; Davatchi, F.: Case report: hypophos-
phatemic osteomalacia
[41580] in von Recklinghausen neurofibromatosis. Am. J. Med. Sci.
301: 322-328,
[41581] 1991.

[41582]

[41583] 144. Korenberg, J. R.; Barker, D.; Fain, P.; Graham, J. M.;
Pribyl,
[41584] T.; Pulst, S.-M.: Achondroplasia is not tightly linked to the
locus

[41585] for neurofibromatosis 1. (Abstract) Cytogenet. Cell Genet. 51: 1025,
[41586] 1989.

[41587]

[41588] 145. Kousseff, B. G.; Gilbert-Barness, E. F.: 'Vascular neurofibromatosis'
[41589] and infantile gangrene. Am. J. Med. Genet. 34: 221-226, 1989.
[41590]

[41591] 146. Krone, W.; Hogemann, I.: Cell culture studies on neurofibromatosis
[41592] (von Recklinghausen): V. Monosomy 22 and other chromosomal anomalies
[41593] in cultures from peripheral neurofibromas. Hum. Genet. 74: 453-455,
[41594] 1986.

[41595]

[41596] 147. Kurotaki, H.; Kamata, Y.; Kimura, M.; Nagai, K.: Multiple papillary
[41597] adenomas of type II pneumocytes found in a 13-year-old boy with von

[41598] Recklinghausen's disease. Virchows Arch. A 423:
319-322, 1993.

[41599]

[41600] 148. Lazaro, C.; Kruyer, H.; Gaona, A.; Estivill, X.: Two fur-
ther

[41601] cases of mutation R1947X in the NF1 gene: screening for
a relatively

[41602] common recurrent mutation. Hum. Genet. 96: 361-363,
1995.

[41603]

[41604] 149. Lazaro, C.; Ravella, A.; Gaona, A.; Volpini, V.; Estivill,
X.

[41605] : Neurofibromatosis type 1 due to germ-line mosaicism in
a clinically

[41606] normal father. New Eng. J. Med. 331: 1403-1407, 1994.

[41607]

[41608] 150. Ledbetter, D. H.; Rich, D. C.; O'Connell, P.; Leppert,
M.; Carey,

[41609] J. C.: Precise localization of NF1 to 17q11.2 by balanced
translocation. Am.

[41610] J. Hum. Genet. 44: 20-24, 1989.

[41611]

[41612] 151. Legius, E.; Descheemaeker, M.-J.; Spaepen, A.; Casaer, P.; Fryns,

[41613] J.-P.: Neuropsychological profile in children with NF1. (Abstract) Genet.

[41614] Counsel. 5: 213-214, 1994.

[41615]

[41616] 152. Legius, E.; Descheemaeker, M. J.; Steyaert, J.; Spaepen, A.;

[41617] Vlietinck, R.; Casaer, P.; Demaerel, P.; Fryns, J. P.: Neurofi-bromatosis

[41618] type 1 in childhood: correlation of MRI findings with intel-ligence. J.

[41619] Neurol. Neurosurg. Psychiat. 59: 638-640, 1995.

[41620]

[41621] 153. Legius, E.; Marchuk, D. A.; Collins, F. S.; Glover, T. W.: Somatic

[41622] deletion of the neurofibromatosis type 1 gene in a neu-rofibrosarcoma

[41623] supports a tumour suppressor gene hypothesis. Nature Genet. 3: 122-125,

[41624] 1993.

[41625]

[41626] 154. Legius, E.; Marchuk, D. A.; Hall, B. K.; Andersen, L. B.; Wallace,

[41627] M. R.; Collins, F. S.; Glover, T. W.: NF1-related locus on chromosome

[41628] 15. Genomics 13: 1316-1318, 1992.

[41629]

[41630] 155. Lehrnbecher, T.; Gassel, A. M.; Rauh, V.; Kirchner, T.; Huppertz,

[41631] H.-I.: Neurofibromatosis presenting as a severe systemic vasculopathy. Europ.

[41632] J. Pediat. 153: 107-109, 1994.

[41633]

[41634] 156. Leppig, K. A.; Viskochil, D.; Neil, S.; Rubenstein, A.; Johnson,

[41635] V. P.; Zhu, X. L.; Brothman, A. R.; Stephens, K.: The detec-tion of

[41636] contiguous gene deletions at the neurofibromatosis 1 lo-cus with fluorescence

[41637] in situ hybridization. Cytogenet. Cell Genet. 72: 95-98,

1996.

[41638]

[41639] 157. Li, Y.; Bollag, G.; Clark, R.; Stevens, J.; Conroy, L.; Fults,

[41640] D.; Ward, K.; Friedman, E.; Samowitz, W.; Robertson, M.; Bradley,

[41641] P.; McCormick, F.; White, R.; Cawthon, R.: Somatic mutations in the

[41642] neurofibromatosis 1 gene in human tumors. Cell 69: 275-281, 1992.

[41643]

[41644] 158. Li, Y.; O'Connell, P.; Huntsman Breidenbach, H.; Cawthon, R.;

[41645] Stevens, J.; Xu, G.; Neil, S.; Robertson, M.; White, R.; Viskochil,

[41646] D.: Genomic organization of the neurofibromatosis 1 gene (NF1). Genomics 25:

[41647] 9-18, 1995.

[41648]

[41649] 159. Lin, A. E.; Birch, P. H.; Korf, B. R.; Tenconi, R.; Nimura,

[41650] M.; Poyhonen, M.; Uhas, K. A.; Sigorini, M.; Virdis, R.; Ro-

mano, C.;

[41651] Bonioli, E.; Wolkenstein, P.; Pivnick. E. K.; Lawrence, M.; Friedman,

[41652] J. M.; NNFF International Database Participants: Cardiovas-

cular

[41653] malformations and other cardiovascular abnormalities in neurofibromatosis

[41654] 1. Am. J. Med. Genet. 95: 108-117, 2000.

[41655]

[41656] 160. Lisch, K.: Ueber Beteiligung der Augen, insbesondere das Vorkommen

[41657] von Irisknotchen bei der Neurofibromatose (Recklinghausen). Augenheilkunde 93:

[41658] 137-143, 1937.

[41659]

[41660] 161. Listernick, R.; Charrow, J.; Greenwald, M.; Mets, M.: Natural

[41661] history of optic pathway tumors in children with neurofi-

bromatosis

[41662] type 1: a longitudinal study. J. Pediat. 125: 63-66, 1994.

[41663]

[41664] 162. Listernick, R.; Darling, C.; Greenwald, M.; Strauss, L.; Charrow,
[41665] J.: Optic pathway tumors in children: the effect of neurofibromatosis
[41666] type 1 on clinical manifestations and natural history. J. Pediat. 127:
[41667] 718-722, 1995.

[41668]

[41669] 163. Littler, M.; Morton, N. E.: Segregation analysis of peripheral
[41670] neurofibromatosis (NF1). J. Med. Genet. 27: 307-310, 1990.

[41671]

[41672] 164. Lopez-Correa, C.; Dorschner, M.; Brems, H.; Lazaro, C.; Clementi,
[41673] M.; Upadhyaya, M.; Dooijes, D.; Moog, U.; Kehrer-Sawatzki, H.; Rutkowski,
[41674] J. L.; Fryns, J.-P.; Marynen, P.; Stephens, K.; Legius, E.: Recombination
[41675] hotspot in NF1 microdeletion patients. Hum. Molec. Genet. 10: 1387-1392,
[41676] 2001.

[41677]

[41678] 165. Luijten, M.; Redeker, S.; Minoshima, S.; Shimizu, N.; Westerveld,

[41679] A.; Hulsebos, T. J. M.: Duplication and transposition of the NF1

[41680] pseudogene regions on chromosomes 2, 14, and 22. Hum. Genet. 109:

[41681] 109-116, 2001.

[41682]

[41683] 166. Lund, A. M.; Skovby, F.: Optic gliomas in children with neurofibromatosis

[41684] type 1. Europ. J. Pediat. 150: 835-838, 1991.

[41685]

[41686] 167. Manley, K. A.; Skyring, A. P.: Some heritable causes of gastrointestinal

[41687] disease: special reference to hemorrhage. Arch. Intern. Med. 107:

[41688] 182-203, 1961.

[41689]

[41690] 168. Marchuk, D. A.; Saulino, A. M.; Tavakkol, R.; Swa-
roop, M.; Wallace,

[41691] M. R.; Andersen, L. B.; Mitchell, A. L.; Gutmann, D. H.; Boguski,

[41692] M.; Collins, F. S.: cDNA cloning of the type 1 neurofibromatosis

[41693] gene: complete sequence of the NF1 gene product. Genomics 11: 931-940,

[41694] 1991.

[41695]

[41696] 169. Massaro, D.; Katz, S.: Fibrosing alveolitis: its occurrence,

[41697] roentgenographic, and pathologic features in von Recklinghausen's

[41698] neurofibromatosis. Am. Rev. Resp. Dis. 93: 934-942, 1966.

[41699]

[41700] 170. Mathew, C. G. P.; Thorpe, K.; Easton, D. F.; Chin, K. S.; Jadayel,

[41701] D.; Ponder, M.; Moore, G.; Wallis, C. E.; Slater, C. P.; De Jong,

[41702] G.; O'Connell, P.; White, R.; Barker, D.; Ponder, B. A. J.: Linkage

[41703] analysis of chromosome 17 markers in British and South

African families
[41704] with neurofibromatosis type 1. Am. J. Hum. Genet. 44:
38-40, 1989.
[41705]

[41706] 171. Maynard, J.; Krawczak, M.; Upadhyaya, M.: Charac-
terization and
[41707] significance of nine novel mutations in exon 16 of the
neurofibromatosis
[41708] type 1 (NF1) gene. Hum. Genet. 99: 674-676, 1997.

[41709]

[41710] 172. McGaughran, J. M.; Harris, D. I.; Donnai, D.; Teare,
D.; MacLeod,
[41711] R.; Westerbeek, R.; Kingston, H.; Super, M.; Harris, R.;
Evans, D.
[41712] G. R.: A clinical study of type I neurofibromatosis in north
west
[41713] England. J. Med. Genet. 36: 197-203, 1999.

[41714]

[41715] 173. Menon, A. G.; Anderson, K. M.; Riccardi, V. M.;
Chung, R. Y.;
[41716] Whaley, J. M.; Yandell, D. W.; Farmer, G. E.; Freiman, R. N.;

Lee,

[41717] J. K.; Li, F. P.; Barker, D. F.; Ledbetter, D. H.; Kleider, A.; Martuza,

[41718] R. L.; Gusella, J. F.; Seizinger, B. R.: Chromosome 17p deletions

[41719] and p53 gene mutations associated with the formation of malignant

[41720] neurofibrosarcomas in von Recklinghausen neurofibro- matosis. Proc.

[41721] Nat. Acad. Sci. 87: 5435-5439, 1990.

[41722]

[41723] 174. Menon, A. G.; Ledbetter, D. H.; Rich, D. C.; Seizinger, B. R.;

[41724] Rouleau, G. A.; Michels, V. F.; Schmidt, M. A.; Dewald, G.; DallaTorre,

[41725] C. M.; Haines, J. L.; Gusella, J. F.: Characterization of a translocation

[41726] within the von Recklinghausen neurofibromatosis region of chromosome

[41727] 17. Genomics 5: 245-249, 1989.

[41728]

[41729] 175. Messiaen, L. M.; Callens, T.; Mortier, G.; Beysen, D.;

Vandenbroucke,
[41730] I.; Van Roy, N.; Speleman, F.; De Paepe, A.: Exhaustive mutation
[41731] analysis of the NF1 gene allows identification of 95% of mutations
[41732] and reveals a high frequency of unusual splicing defects. Hum. Mutat. 15:
[41733] 541-555, 2000.

[41734]

[41735] 176. Messiaen, L. M.; Callens, T.; Roux, K. J.; Mortier, G. R.; De
[41736] Paepe, A.; Abramowicz, M.; Pericak-Vance, M. A.; Vance, J. M.; Wallace,
[41737] M. R.: Exon 10b of the NF1 gene represents a mutational hotspot and
[41738] harbors a recurrent missense mutation Y489C associated with aberrant
[41739] splicing. Genet. Med. 1: 248-253, 1999.

[41740]

[41741] 177. Miles, J.; Pennybacker, J.; Sheldon, P.: Intrathoracic meningocele.
[41742] Its development and association with neurofibromatosis. J.

Neurol.
[41743] Neurosurg. Psychiat. 32: 99-110, 1969.

[41744]

[41745] 178. Miller, M.; Hall, J. G.: Possible maternal effect of severity
[41746] of neurofibromatosis. Lancet II: 1071-1073, 1978.

[41747]

[41748] 179. Miller, M. E.; Hall, J. G.: Maternal effect in neurofibro-matosis.
[41749] (Abstract) Am. J. Hum. Genet. 30: 60A, 1978.

[41750]

[41751] 180. Miller, R. M.; Sparkes, R. S.: Segmental neurofibro-matosis. Arch.
[41752] Derm. 113: 837-838, 1977.

[41753]

[41754] 181. Molloy, P. T.; Bilaniuk, L. T.; Vaughan, S. N.; Needle, M. N.;
[41755] Liu, G. T.; Zackai, E. H.; Phillips, P. C.: Brainstem tumors in patients
[41756] with neurofibromatosis type 1: a distinct clinical entity.

Neurology 45:
[41757] 1897-1902, 1995.

[41758]

[41759] 182. Mukhopadhyay, D.; Anant, S.; Lee, R. M.; Kennedy, S.; Viskochil,
[41760] D.; Davidson, N. O.: C-U editing of neurofibromatosis 1 mRNA occurs
[41761] in tumors that express both the type II transcript and apobec-1, the
[41762] catalytic subunit of the apolipoprotein B mRNA-editing enzyme. Am.
[41763] J. Hum. Genet. 70: 38-50, 2002.

[41764]

[41765] 183. Mukonoweshuro, W.; Griffiths, P. D.; Blaser, S.: Neu-rofibromatosis
[41766] type 1: the role of neuroradiology. Neuropediatrics 30: 111-119,
[41767] 1999.

[41768]

[41769] 184. Muller-Wiefel, D. E.: Renovaskulare Hypertension bei Neurofibromatose

[41770] von Recklinghausen. Mschr. Kinderheilk. 126: 113-118, 1978.

[41771]

[41772] 185. Nager, G. T.: Association of bilateral VIIIth nerve tumors with

[41773] meningiomas in von Recklinghausen's disease. Laryngoscope 74: 1220-1261,

[41774] 1964.

[41775]

[41776] 186. Nakafuku, M.; Nagamine, M.; Ohtoshi, A.; Tanaka, K.; Toh-e, A.;

[41777] Kaziro, Y.: Suppression of oncogenic Ras by mutant neurofibromatosis

[41778] type 1 genes with single amino acid substitutions. Proc. Nat. Acad.

[41779] Sci. 90: 6706-6710, 1993.

[41780]

[41781] 187. National Institutes of Health Consensus Development Conference

[41782] : Neurofibromatosis: conference statement. Arch. Neurol. 45: 575-578,

[41783] 1988.

[41784]

[41785] 188. Newman, A.; So, S. K.: Bilateral neurofibroma of the intrathoracic

[41786] vagus associated with von Recklinghausen's disease. Am. J. Roentgen. 112:

[41787] 389-392, 1971.

[41788]

[41789] 189. Nicolls, E. M.: Somatic variation and multiple neurofibromatosis. Hum.

[41790] Hered. 19: 473-479, 1969.

[41791]

[41792] 190. Nopajaroonsri, C.; Lurie, A. A.: Venous aneurysm, arterial dysplasia,

[41793] and near-fatal hemorrhages in neurofibromatosis type 1. Hum. Path. 27:

[41794] 982-985, 1996.

[41795]

[41796] 191. O'Connell, P.; Leach, R.; Cawthon, R. M.; Culver, M.; Stevens,

[41797] J.; Viskochil, D.; Fournier, R. E. K.; Rich, D. C.; Ledbetter, D.

[41798] H.; White, R.: Two NF1 translocations map within a 600-kilobase segment

[41799] of 17q11.2. Science 244: 1087-1088, 1989.

[41800]

[41801] 192. O'Connell, P.; Leach, R. J.; Ledbetter, D. H.; Cawthon, R. M.;

[41802] Culver, M.; Eldridge, J. R.; Frej, A.-K.; Holm, T. R.; Wolff, E.;

[41803] Thayer, M. J.; Schafer, A. J.; Fountain, J. W.; Wallace, M. R.; Collins,

[41804] F. S.; Skolnick, M. H.; Rich, D. C.; Fournier, R. E. K.; Baty, B.

[41805] J.; Carey, J. C.; Leppert, M. F.; Lathrop, G. M.; Lalouel, J.-M.;

[41806] White, R.: Fine structure DNA mapping studies of the chromosomal

[41807] region harboring the genetic defect in neurofibromatosis type 1. Am.

[41808] J. Hum. Genet. 44: 51-57, 1989.

[41809]

[41810] 193. Obringer, A. C.; Meadows, A. T.; Zackai, E. H.: The diagnosis

[41811] of neurofibromatosis-1 in the child under the age of 6 years. Am.

[41812] J. Dis. Child. 143: 717-719, 1989.

[41813]

[41814] 194. Osborn, M. J.; Upadhyaya, M.: Evaluation of the protein truncation

[41815] test and mutation detection in the NF1 gene: mutational analysis of

[41816] 15 known and 40 unknown mutations. Hum. Genet. 105: 327-332, 1999.

[41817]

[41818] 195. Otsuka, F.; Kawashima, T.; Imakado, S.; Usuki, Y.; Hon-mura,

[41819] S.: Lisch nodules and skin manifestation in neurofibromatosis type

[41820] 1. Arch. Derm. 137: 232-233, 2001.

[41821]

[41822] 196. Parazzini, C.; Triulzi, F.; Bianchini, E.; Agnetti, V.; Conti,
[41823] M.; Zanolini, C.; Maninetti, M. M.; Rossi, L. N.; Scotti, G.: Spontaneous
[41824] involution of optic pathway lesions in neurofibromatosis type 1: serial
[41825] contrast MR evaluation. Am. J. Neuroradiol. 16: 1711-1718, 1995.
[41826]

[41827] 197. Park, K. C.; Choi, H. O.; Park, K. H.; Kim, K. H.; Eun, H. C.
[41828] : A nonsense mutation at arg-1947 in the NF1 gene in a case of neurofibromatosis
[41829] type 1 in a Korean patient. J. Hum. Genet. 45: 84-85, 2000.
[41830]

[41831] 198. Park, V. M.; Pivnick, E. K.: Neurofibromatosis type 1 (NF1):
[41832] a protein truncation assay yielding identification of muta-tions in
[41833] 73% of patients. J. Med. Genet. 35: 813-820, 1998.
[41834]

[41835] 199. Parsa, C. F.; Hoyt, C. S.; Lesser, R. L.; Weinstein, J. M.; Strother,

[41836] C. M.; Muci-Mendoza, R.; Ramella, M.; Manor, R. S.; Fletcher, W. A.;

[41837] Repka, M. X.; Garrity, J. A.; Ebner, R. N.; Monteiro, M. L. R.; McFadzean,

[41838] R. M.; Rubtsova, I. V.; Hoyt, W. F.: Spontaneous regression of optic

[41839] gliomas: thirteen cases documented by serial neuroimaging. Arch.

[41840] Ophthal. 119: 516-529, 2001.

[41841]

[41842] 200. Pellock, J. M.; Kleinman, P. K.; McDonald, B. M.; Wixson, D.

[41843] : Childhood hypertensive stroke with neurofibromatosis. Neurology 30:

[41844] 656-659, 1980.

[41845]

[41846] 201. Perry, H. D.; Font, R. L.: Iris nodules in von Recklinghausen's

[41847] neurofibromatosis: electron microscopic confirmation of their melanocytic

[41848] origin. Arch. Ophthal. 100: 1635-1640, 1982.

[41849]

[41850] 202. Philippart, M.: Neurofibromatose hereditaire a large spectre

[41851] phenotypique (famille SN). J. Genet. Hum. 10: 338-346, 1961.

[41852]

[41853] 203. Porterfield, J. K.; Pyeritz, R. E.; Traill, T. A.: Pulmonary

[41854] hypertension and interstitial fibrosis in von Reckling-hausen neurofibromatosis. Am.

[41855] J. Med. Genet. 25: 531-535, 1986.

[41856]

[41857] 204. Poyhonen, M.; Kytola, S.; Leisti, J.: Epidemiology of neurofibromatosis

[41858] type 1 (NF1) in northern Finland. (Letter) J. Med. Genet. 37: 632-636,

[41859] 2000.

[41860]

[41861] 205. Poyhonen, M.; Leisti, E.-L.; Kytola, S.; Leisti, J.: Hereditary

[41862] spinal neurofibromatosis: a rare form of NF1? J. Med. Genet. 34:

[41863] 184-187, 1997.

[41864]

[41865] 206. Pulst, S.-M.; Graham, J. M., Jr.; Fain, P.; Barker, D.; Pribyl,

[41866] T.; Korenberg, J. R.: The achondroplasia gene is not linked to the

[41867] locus for neurofibromatosis 1 on chromosome 17. Hum. Genet. 85:

[41868] 12-14, 1990.

[41869]

[41870] 207. Pulst, S.-M.; Pribyl, T.; Barker, D. F.; Riccardi, V. M.; Ren,

[41871] M.; Yaari, H.; Korenberg, J. R.: Molecular analysis of a pa-tient

[41872] with neurofibromatosis 1 and achondroplasia. Am. J. Med. Genet. 40:

[41873] 84-87, 1991.

[41874]

[41875] 208. Pulst, S.-M.; Riccardi, V. M.; Fain, P.; Korenberg, J. R.:

Familial
[41876] spinal neurofibromatosis: clinical and DNA linkage analysis. Neurology 41:
[41877] 1923-1927, 1991.

[41878]

[41879] 209. Purandare, S. M.; Lanyon, W. G.; Arngrimsson, R.; Connor, J.
[41880] M.: Characterisation of a novel splice donor mutation affecting position
[41881] +1 in intron 18 of the NF-1 gene. Hum. Molec. Genet. 4: 767-768,
[41882] 1995.

[41883]

[41884] 210. Pyeritz, R. E.: Personal Communication. Baltimore, Md. 7/20/1987.

[41885]

[41886] 211. Ragge, N. K.; Falk, R. E.; Cohen, W. E.; Murphree, A. L.: Images
[41887] of Lisch nodules across the spectrum. Eye 7: 95-101, 1993.

[41888]

[41889] 212. Rasmussen, S. A.; Colman, S. D.; Ho, V. T.; Abernathy, C. R.;
[41890] Arn, P. H.; Weiss, L.; Schwartz, C.; Saul, R. A.; Wallace, M. R.:
[41891] Constitutional and mosaic large NF1 gene deletions in neurofibromatosis
[41892] type 1. J. Med. Genet. 35: 468-471, 1998.

[41893]

[41894] 213. Rasmussen, S. A.; Yang, Q.; Friedman, J. M.: Mortality in neurofibromatosis
[41895] 1: an analysis using U.S. death certificates. Am. J. Hum. Genet. 68:
[41896] 1110-1118, 2001.

[41897]

[41898] 214. Reilly, K. M.; Loisel, D. A.; Bronson, R. T.; McLaughlin, M.
[41899] E.; Jacks, T.: Nf1;Trp53 mutant mice develop glioblastoma with evidence
[41900] of strain-specific effects. Nature Genet. 26: 109-113, 2000.

[41901]

[41902] 215. Reubi, F.: Neurofibromatose et lesions vasculaires. Schweiz.

[41903] Med. Wochenschr. 75: 463-465, 1945.

[41904]

[41905] 216. Riccardi, V. M.: The vasculopathy of NF1 and histo-genesis control

[41906] genes. Clin. Genet. 58: 345-347, 2000.

[41907]

[41908] 217. Riccardi, V. M.: Von Recklinghausen neurofibromato-sis. New

[41909] Eng. J. Med. 305: 1617-1626, 1981.

[41910]

[41911] 218. Riccardi, V. M.: Personal Communication. Houston, Tex. 1982.

[41912]

[41913] 219. Riccardi, V. M.: Neurofibromatosis: clinical hetero-geneity. Curr.

[41914] Probl. Cancer 7: 1-34, 1982.

[41915]

[41916] 220. Riccardi, V. M.; Dobson, C. E., II; Chakraborty, R.;

Bontke,
[41917] C.: The pathophysiology of neurofibromatosis: IX. Paternal age as
[41918] a factor in the origin of new mutations. Am. J. Med. Genet. 18:
[41919] 169-176, 1984.

[41920]

[41921] 221. Riccardi, V. M.; Eichner, J. E.: Neurofibromatosis: Phenotype,
[41922] Natural History and Pathogenesis. Baltimore: Johns Hopkins Univ.
[41923] Press (pub.) 1986.

[41924]

[41925] 222. Riccardi, V. M.; Mulvihill, J. J.: Neurofibromatosis (von Recklinghausen
[41926] Disease): Genetics, Cell Biology, and Biochemistry. Advances in Neurology.
[41927] New York: Raven Press (pub.) 29: 1981. Pp. 304 only.

[41928]

[41929] 223. Ricciardone, M. D.; Ozcelik, T.; Cevher, B.; Ozdag, H.; Tuncer,

[41930] M.; Gurgey, A.; Uzunalimoglu, O.; Cetinkaya, H.; Tanyeli, A.; Erken,
[41931] E.; Ozturk, M.: Human MLH1 deficiency predisposes to hematological
[41932] malignancy and neurofibromatosis type 1. Cancer Res. 59: 290-293,
[41933] 1999.

[41934]

[41935] 224. Ritter, J. L.; Riccardi, V. M.: Von Recklinghausen neu-
rofibromatosis
[41936] (NF-I): an argument for very high penetrance and a com-
parison of sporadic
[41937] and inherited cases. (Abstract) Am. J. Hum. Genet. 37: A135, 1985.

[41938]

[41939] 225. Riva, P.; Castorina, P.; Manoukian, S.; Dalpra, L.; Doneda, L.;
[41940] Marini, G.; den Dunnen, J.; Larizza, L.: Characterization of a cytogenetic
[41941] 17q11.2 deletion in an NF1 patient with a contiguous gene syndrome. Hum.
[41942] Genet. 98: 646-650, 1996.

[41943]

[41944] 226. Robertson, D.: Personal Communication. Nashville, Tenn. 8/1979.

[41945]

[41946] 227. Robinson, P. N.; Buske, A.; Neumann, R.; Tinschert, S.; Nurnberg,

[41947] P.: Recurrent 2-bp deletion in exon 10c of the NF1 gene in two cases

[41948] of von Recklinghausen neurofibromatosis. Hum. Mutat. 7: 85-88, 1996.

[41949]

[41950] 228. Rockower, S.; McKay, D.; Nason, S.: Dislocation of the spine

[41951] in neurofibromatosis: a report of two cases. J. Bone Joint Surg. 64:

[41952] 1240-1242, 1982.

[41953]

[41954] 229. Rosenquist, G. C.; Krovetz, L. J.; Haller, J. A., Jr.; Si-mon,

[41955] A. L.; Bannayan, G. A.: Acquired right ventricular outflow obstruction

[41956] in a child with neurofibromatosis. Am. Heart J. 79: 103-108, 1970.

[41957]

[41958] 230. Rutkowski, J. L.; Wu, K.; Gutmann, D. H.; Boyer, P. J.; Legius,

[41959] E.: Genetic and cellular defects contributing to benign tumor formation

[41960] in neurofibromatosis type 1. Hum. Molec. Genet. 9: 1059-1066, 2000.

[41961]

[41962] 231. Sakaguchi, N.; Sano, K.; Ito, M.; Baba, T.; Fukuzawa, M.; Hotchi,

[41963] M.: A case of von Recklinghausen's disease with bilateral pheochromocytoma-malignant

[41964] peripheral nerve sheath tumors of the adrenal and gastrointestinal

[41965] autonomic nerve tumors. Am. J. Surg. Path. 20: 889-897, 1996.

[41966]

[41967] 232. Sakurai, T.: Multiple neurofibroma patient showing multiple

[41968] flecks on the anterior surface of the iris. Acta Soc. Oph-

thal. Jpn. 39:
[41969] 87-93, 1935.

[41970]

[41971] 233. Salyer, W. R.; Salyer, D. C.: The vascular lesions of neurofibromatosis. Angiology 25:
[41972] 510-519, 1974.

[41973]

[41974] 234. Samuelsson, B.; Akesson, H. O.: Relative fertility and mutation
[41975] rate in neurofibromatosis. Hereditas 108: 169-171, 1988.

[41976]

[41977] 235. Sands, M. J.; McDonough, M. T.; Cohen, A. M.; Rutenberg, H. L.;
[41978] Eisner, J. W.: Fatal malignant degeneration in multiple neurofibromatosis. J.A.M.A. 233:
[41979] 1381-1382, 1975.

[41980]

[41981] 236. Satran, L.; Letson, R. D.; Seljeskog, E. L.: Neurofibro-matosis
[41982] with congenital glaucoma and buphthalmos in a newborn.

Am. J. Dis.
[41983] Child. 134: 182-183, 1980.

[41984]

[41985] 237. Sawada, S.; Florell, S.; Purandare, S. M.; Ota, M.; Stephens,
[41986] K.; Viskochil, D.: Identification of NF1 mutations in both alleles
[41987] of a dermal neurofibroma. Nature Genet. 14: 110-112, 1996.

[41988]

[41989] 238. Sayed, A. K.; Bernhardt, B.; Perez-Atayde, A. R.; Ban-nerman,
[41990] R. M.: Malignant schwannoma in siblings with neurofibro-matosis. Cancer 59:
[41991] 829-835, 1987.

[41992]

[41993] 239. Schenkein, I.; Bueker, E. D.; Helson, L.; Axelrod, F.; Dancis,
[41994] J.: Increased nerve-growth stimulating activity in dissemi-nated neurofibromatosis. New
[41995] Eng. J. Med. 290: 613-614, 1974.

[41996]

[41997] 240. Schmidt, M. A.; Michels, V. V.; Dewald, G. W.: Cases of neurofibromatosis

[41998] with rearrangements of chromosome 17 involving band 17q11.2. Am.

[41999] J. Med. Genet. 28: 771-777, 1987.

[42000]

[42001] 241. Schotland, H. M.; Eldridge, R.; Sommer, S. S.; Malawar, M.:

[42002] Neurofibromatosis 1 and osseous fibrous dysplasia in a family. Am.

[42003] J. Med. Genet. 43: 815-822, 1992.

[42004]

[42005] 242. Seizinger, B. R.; Farmer, G. E.; Haines, J. L.; Ozelius, L. J.;

[42006] Anderson, K.; Korf, B. R.; Parry, D. M.; Pericak-Vance, M. A.; Mulvihill,

[42007] J. J.; Menon, A.; Hobbs, W. J.; Martuza, R. L.; Gusella, J. F.: Flanking

[42008] markers for the gene causing von Recklinghausen neu- rofibromatosis

[42009] (NF1). Am. J. Hum. Genet. 44: 30-32, 1989.

[42010]

[42011] 243. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane, A. H.;

[42012] Faryniarz, A. G.; Chao, M. V.; Huson, S.; Korf, B. R.; Parry, D. M.;

[42013] Pericak-Vance, M. A.; and 21 others: Genetic linkage of von Recklinghausen

[42014] neurofibromatosis to the nerve growth factor receptor gene. Cell 49:

[42015] 589-594, 1987.

[42016]

[42017] 244. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane, A. H.;

[42018] Faryniarz, A. G.; Chao, M. V.; Huson, S.; Korf, B. R.; Parry, D. M.;

[42019] Pericak-Vance, M. A.; Collins, F. S.; Hobbs, W. J.; and 21 others

[42020] : Genetic linkage of von Recklinghausen neurofibromato-sis to the nerve

[42021] growth factor receptor gene. (Abstract) Cytogenet. Cell Genet. 46:

[42022] 690 only, 1987.

[42023]

[42024] 245. Senveli, E.; Altinors, N.; Kars, Z.; Arda, N.; Turker, A.; Cinar,

[42025] N.; Yalniz, Z.: Association of von Recklinghausen's neurofibromatosis

[42026] and aqueduct stenosis. Neurosurgery 24: 99-101, 1989.

[42027]

[42028] 246. Serra, E.; Ars, E.; Ravella, A.; Sanchez, A.; Puig, S.; Rosenbaum,

[42029] T.; Estivill, X.; Lazaro, C.: Somatic NF1 mutational spectrum in

[42030] benign neurofibromas: mRNA splice defects are common among point mutations. Hum.

[42031] Genet. 108: 416-429, 2001.

[42032]

[42033] 247. Serra, E.; Puig, S.; Otero, D.; Gaona, A.; Kruyer, H.; Ars, E.;

[42034] Estivill, X.; Lazaro, C.: Confirmation of a double-hit model for

[42035] the NF1 gene in benign neurofibromas. Am. J. Hum.

Genet. 61: 512-519,
[42036] 1997.

[42037]

[42038] 248. Serra, E.; Rosenbaum, T.; Winner, U.; Aledo, R.; Ars,
E.; Estivill,
[42039] X.; Lenard, H.-G.; Lazaro, C.: Schwann cells harbor the so-
matic NF1
[42040] mutation in neurofibromas: evidence of two different
Schwann cell
[42041] subpopulations. Hum. Molec. Genet. 9: 3055-3064, 2000.

[42042]

[42043] 249. Shannon, K. M.; O'Connell, P.; Martin, G. A.;
Paderanga, D.;
[42044] Olson, K.; Dinndorf, P.; McCormick, F.: Loss of the normal
NF1 allele
[42045] from the bone marrow of children with type 1 neurofibro-
matosis and
[42046] malignant myeloid disorders. New Eng. J. Med. 330:
597-601, 1994.

[42047]

[42048] 250. Shannon, K. M.; Watterson, J.; Johnson, P.; O'Connell,

P.; Lange,
[42049] B.; Shah, N.; Steinherz, P.; Kan, Y. W.; Priest, J. R.: Monosomy
[42050] 7 myeloproliferative disease in children with neurofibromatosis, type
[42051] 1: epidemiology and molecular analysis. Blood 79: 1311-1318, 1992.
[42052]

[42053] 251. Shen, M. H.; Harper, P. S.; Upadhyaya, M.: Molecular genetics
[42054] of neurofibromatosis type 1 (NF1). J. Med. Genet. 33: 2-17, 1996.
[42055]

[42056] 252. Side, L.; Taylor, B.; Cayouette, M.; Conner, E.; Thompson, P.;
[42057] Luce, M.; Shannon, K.: Homozygous inactivation of the NF1 gene in
[42058] bone marrow cells from children with neurofibromatosis type 1 and
[42059] malignant myeloid disorders. New Eng. J. Med. 336: 1713-1720, 1997.
[42060]

[42061] 253. Side, L. E.; Emanuel, P. D.; Taylor, B.; Franklin, J.; Thompson,
[42062] P.; Castleberry, R. P.; Shannon, K. M.: Mutations of the NF1 gene
[42063] in children with juvenile myelomonocytic leukemia without clinical
[42064] evidence of neurofibromatosis, type 1. Blood 92: 267-272, 1998.
[42065]

[42066] 254. Siggers, D. C.; Boyer, S. H.; Eldridge, R.: Nerve-growth factor
[42067] in disseminated neurofibromatosis. (Letter) New Eng. J. Med. 292:
[42068] 1134, 1975.

[42069]

[42070] 255. Silva, A. J.; Frankland, P. W.; Marowitz, Z.; Friedman, E.; Lazlo,
[42071] G.; Cioffi, D.; Jacks, T.; Bourtchuladze, R.: A mouse model for the
[42072] learning and memory deficits associated with neurofibro-matosis type
[42073] I. Nature Genet. 15: 281-284, 1997.

[42074]

[42075] 256. Singhal, S.; Birch, J. M.; Kerr, B.; Lashford, L.; Evans, D.

[42076] G. R.: Neurofibromatosis type 1 and sporadic optic gliomas. Arch.

[42077] Dis. Child. 87: 65-70, 2002.

[42078]

[42079] 257. Skuse, G. R.; Cappione, A. J.: RNA processing and clinical variability

[42080] in neurofibromatosis type 1 (NF1). Hum. Mol. Genet. 6: 1707-1712,

[42081] 1997.

[42082]

[42083] 258. Skuse, G. R.; Cappione, A. J.; Sowden, M.; Metheny, L. J.; Smith,

[42084] H. C.: The neurofibromatosis type I messenger RNA undergoes base-modification

[42085] RNA editing. Nucleic Acids Res. 24: 478-485, 1996.

[42086]

[42087] 259. Skuse, G. R.; Kosciolek, B. A.; Rowley, P. T.: Molecular genetic

[42088] analysis of tumors in von Recklinghausen neurofibro-
matosis: loss of
[42089] heterozygosity for chromosome 17. Genes Chromosomes
Cancer 1: 36-41,
[42090] 1989.

[42091]

[42092] 260. Skuse, G. R.; Kosciolek, B. A.; Rowley, P. T.: The neu-
rofibroma
[42093] in von Recklinghausen neurofibromatosis has a unicellular
origin. Am.
[42094] J. Hum. Genet. 49: 600-607, 1991.

[42095]

[42096] 261. Smith, C. J.; Hatch, F. E.; Johnson, J. G.; Kelly, B. J.:
Renal
[42097] artery dysplasia as a cause of hypertension in neurofibro-
matosis. Arch.
[42098] Intern. Med. 125: 1022-1026, 1970.

[42099]

[42100] 262. Sorensen, S. A.; Mulvihill, J. J.; Nielsen, A.: Long-term
follow-up
[42101] of von Recklinghausen neurofibromatosis: survival and

malignant neoplasms. New
[42102] Eng. J. Med. 314: 1010-1015, 1986.

[42103]

[42104] 263. Stambolian, D.; Zackai, E. H.: Gene location in neu-
rofibromatosis.
[42105] (Letter) Am. J. Med. Genet. 29: 963-965, 1988.

[42106]

[42107] 264. Stanley, J. C.: Arterial fibrodysplasia. Arch. Surg. 110:

[42108] 561-566, 1975.

[42109]

[42110] 265. Stanley, J. C.; Fry, W. J.: Pediatric renal artery occlu-
sive
[42111] disease and renovascular hypertension: etiology, diagno-
sis, and operative
[42112] treatment. Arch. Surg. 116: 669-676, 1981.

[42113]

[42114] 266. Stark, M.; Assum, G.; Kaufmann, D.; Kehrer, H.;
Krone, W.: Analysis
[42115] of segregation and expression of an identified mutation at
the neurofibromatosis

[42116] type 1 locus. Hum. Genet. 90: 356-359, 1992.

[42117]

[42118] 267. Stark, M.; Assum, G.; Krone, W.: Single-cell PCR per-
formed with

[42119] neurofibroma Schwann cells reveals the presence of both
alleles of

[42120] the neurofibromatosis type 1 (NF1) gene. Hum. Genet. 96:
619-623,

[42121] 1995.

[42122]

[42123] 268. Stark, M.; Assum, G.; Krone, W.: A small deletion and
an adjacent

[42124] base exchange in a potential stem-loop region of the
neurofibromatosis

[42125] 1 gene. Hum. Genet. 87: 685-687, 1991.

[42126]

[42127] 269. Stephens, K.; Green, P.; Riccardi, V. M.; Ng, S.; Rising,
M.;

[42128] Barker, D.; Darby, J. K.; Falls, K. M.; Collins, F. S.; Willard,
H.

[42129] F.; Donis-Keller, H.: Genetic analysis of eight loci tightly

linked
[42130] to neurofibromatosis 1. Am. J. Hum. Genet. 44: 13-19,
1989.
[42131]

[42132] 270. Stephens, K.; Kayes, L.; Riccardi, V. M.; Rising, M.;
Sybert,
[42133] V. P.; Pagon, R. A.: Preferential mutation of the neurofi-
bromatosis
[42134] type 1 gene in paternally derived chromosomes. Hum.
Genet. 88: 279-282,
[42135] 1992.

[42136]

[42137] 271. Stevenson, D. A.; Birch, P. H.; Friedman, J. M.;
Viskochil, D.
[42138] H.; Balestrazzi, P.; Boni, S.; Buske, A.; Korf, B. R.; Niimura,
M.;
[42139] Pivnick, E. K.; Schorry, E. K.; Short, M. P.; Tenconi, R.; Ton-
sgard,
[42140] J. H.; Carey, J. C.: Descriptive analysis of tibial pseu-
doarthrosis
[42141] in patients with neurofibromatosis 1. Am. J. Med. Genet.
84: 413-419,

[42142] 1999.

[42143]

[42144] 272. Streubel, B.; Latta, E.; Kehrer-Sawatzki, H.; Hoff-
mann, G. F.;

[42145] Fonatsch, C.; Rehder, H.: Somatic mosaicism of a greater
than 1.7-Mb

[42146] deletion of genomic DNA involving the entire NF1 gene as
verified

[42147] by FISH: further evidence for a contiguous gene syndrome
in 17q11.2. Am.

[42148] J. Med. Genet. 87: 12-16, 1999.

[42149]

[42150] 273. Stumpf, D. A.; Alksne, J. F.; Annegers, J. F.; Brown, S.
S.;

[42151] Conneally, P. M.; Housman, D.; Leppert, M. F.; Miller, J. P.;
Moss,

[42152] M. L.; Pileggi, A. J.; Rapin, I.; Strohman, R. C.; Swanson, L.
W.;

[42153] Zimmerman, A.: Neurofibromatosis: conference state-
ment. Arch. Neurol. 45:

[42154] 575-578, 1988.

[42155]

[42156] 274. Sutphen, R.; Galan-Gomez, E.; Kousseff, B. G.: Cli-
toromegaly

[42157] in neurofibromatosis. Am. J. Med. Genet. 55: 325-330,
1995.

[42158]

[42159] 275. Swinburn, B. A.; Yeong, M. L.; Lane, M. R.; Nicholson,
G. I.;

[42160] Holdaway, I. M.: Neurofibromatosis associated with so-
matostatinoma:

[42161] a report of two patients. Clin. Endocr. 28: 353-359, 1988.

[42162]

[42163] 276. Szudek, J.; Birch, P.; Friedman, J. M.; The National
Neurofibromatosis

[42164] Foundation International Database Participants: Growth in
North

[42165] American white children with neurofibromatosis 1 (NF1). J.
Med. Genet. 37:

[42166] 933-938, 2000.

[42167]

[42168] 277. Tassabehji, M.; Strachan, T.; Sharland, M.; Colley, A.;

Donnai,

[42169] D.; Harris, R.; Thakker, N.: Tandem duplication within a neurofibromatosis

[42170] type I (NF1) gene exon in a family with features of Watson syndrome

[42171] and Noonan syndrome. Am. J. Hum. Genet. 53: 90-95, 1993.

[42172]

[42173] 278. Taylor, P. E.: Encapsulated glioma of the sylvian fissure associated

[42174] with neurofibromatosis. Report of a case with histopathological comparison

[42175] of surgical lesion and autopsy specimen following recurrence. J.

[42176] Neuropath. Exp. Neurol. 21: 566-578, 1962.

[42177]

[42178] 279. The, I.; Hannigan, G. E.; Cowley, G. S.; Reginald, S.; Zhong,

[42179] Y.; Gusella, J. F.; Hariharan, I. K.; Bernards, A.: Rescue of a Drosophila

[42180] NF1 mutant phenotype by protein kinase A. Science 276: 791-794,

[42181] 1997.

[42182]

[42183] 280. Tinschert, S.; Naumann, I.; Stegmann, E.; Buske, A.; Kaufmann,

[42184] D.; Thiel, G.; Jenne, D. E.: Segmental neurofibromatosis is caused

[42185] by somatic mutation of the neurofibromatosis type 1 (NF1) gene. Europ.

[42186] J. Hum. Genet. 8: 455-459, 2000.

[42187]

[42188] 281. Tomsick, T. A.; Lukin, R. R.; Chambers, A. A.; Benton, C.: Neurofibromatosis

[42189] and intracranial arterial occlusive disease. Neuroradiology 11:

[42190] 229-234, 1976.

[42191]

[42192] 282. Tonsgard, J. H.; Yelavarthi, K. K.; Cushner, S.; Short, M. P.;

[42193] Lindgren, V.: Do NF1 gene deletions result in a character-istic phenotype? Am.

[42194] J. Med. Genet. 73: 80-86, 1997.

[42195]

[42196] 283. Upadhyaya, M.; Cheryson, A.; Broadhead, W.; Fryer, A.; Shaw,

[42197] D. J.; Huson, S.; Wallace, M. R.; Andersen, L. B.; Marchuk, D. A.;

[42198] Viskochil, D.; Black, D.; O'Connell, P.; Collins, F. S.; Harper, P.

[42199] S.: A 90 kb DNA deletion associated with neurofibromato-sis type 1. J.

[42200] Med. Genet. 27: 738-741, 1990.

[42201]

[42202] 284. Upadhyaya, M.; Cooper, D. N. (eds.): Neurofibro-matosis Type

[42203] 1: from Genotype to Phenotype. Oxford: BIOS Scientific Publ. , 1998.

[42204]

[42205] 285. Upadhyaya, M.; Maynard, J.; Osborn, M.; Huson, S. M.; Ponder,

[42206] M.; Ponder, B. A. J.; Harper, P. S.: Characterisation of germline

[42207] mutations in the neurofibromatosis type 1 (NF1) gene. J. Med. Genet. 32:

[42208] 706-710, 1995.

[42209]

[42210] 286. Upadhyaya, M.; Osborn, M. J.; Maynard, J.; Kim, M. R.; Tamanoi,

[42211] F.; Cooper, D. N.: Mutational and functional analysis of the neurofibromatosis

[42212] type 1 (NF1) gene. Hum. Genet. 99: 88-92, 1997.

[42213]

[42214] 287. Upadhyaya, M.; Roberts, S. H.; Maynard, J.; Sorour, E.; Thompson,

[42215] P. W.; Vaughan, M.; Wilkie, A. O. M.; Hughes, H. E.: A cyto-genetic

[42216] deletion, del(17)(q11.22q21.1), in a patient with sporadic neurofibromatosis

[42217] type 1 (NF1) associated with dysmorphism and develop-mental delay. J.

[42218] Med. Genet. 33: 148-152, 1996.

[42219]

[42220] 288. Upadhyaya, M.; Ruggieri, M.; Maynard, J.; Osborn, M.; Hartog,

[42221] C.; Mudd, S.; Penttinen, M.; Cordeiro, I.; Ponder, M.; Pon-

der, B.

[42222] A. J.; Krawczak, M.; Cooper, D. N.: Gross deletions of the neurofibromatosis

[42223] type 1 (NF1) gene are predominantly of maternal origin and commonly

[42224] associated with a learning disability, dysmorphic features and developmental

[42225] delay. Hum. Genet. 102: 591-597, 1998.

[42226]

[42227] 289. Upadhyaya, M.; Sarfarazi, M.; Huson, S. M.; Broad-head, W.; Fryer,

[42228] A.; Harper, P. S.: Close flanking markers for neurofibro-matosis type

[42229] 1 (NF1). Am. J. Hum. Genet. 44: 41-47, 1989.

[42230]

[42231] 290. Upadhyaya, M.; Sarfarazi, M.; Huson, S. M.; Broad-head, W.; Fryer,

[42232] A. E.; Harper, P. S.: Genetic linkage mapping of chromo-some 17 markers

[42233] and neurofibromatosis type I (NF1). (Abstract) Cytogenet. Cell Genet. 51:

[42234] 1094, 1989.

[42235]

[42236] 291. Upadhyaya, M.; Shen, M.; Cherryson, A.; Farnham, J.; Maynard,

[42237] J.; Huson, S. M.; Harper, P. S.: Analysis of mutations at the neurofibromatosis

[42238] 1 (NF1) locus. Hum. Molec. Genet. 1: 735-740, 1992.

[42239]

[42240] 292. Uren, N.; Been, M.; Guzman, F.: Congenital left atrial wall

[42241] aneurysm in a patient with neurofibromatosis. Brit. Heart J. 59:

[42242] 391-394, 1988.

[42243]

[42244] 293. Valero, M. C.; Velasco, E.; Moreno, F.; Hernandez-Chico, C.:

[42245] Characterization of four mutations in the neurofibromato-sis type 1

[42246] gene by denaturing gradient gel electrophoresis (DGGE). Hum. Molec.

[42247] Genet. 3: 639-641, 1994.

[42248]

[42249] 294. Vance, J. M.; Pericak-Vance, M. A.; Yamaoka, L. H.; Speer, M.

[42250] C.; Rosenwasser, G. O. D.; Small, K.; Gaskell, P. C., Jr.; Hung, W.-Y.;

[42251] Alberts, M. J.; Haynes, C. S.; Gilbert, J. R.; Aylsworth, A. S.; Roses,

[42252] A. D.: Genetic linkage mapping of chromosome 17 markers and neurofibromatosis

[42253] type 1. Am. J. Med. Genet. 44: 25-29, 1989.

[42254]

[42255] 295. Viskochil, D.; Buchberg, A. M.; Xu, G.; Cawthon, R. M.; Stevens,

[42256] J.; Wolff, R. K.; Culver, M.; Carey, J. C.; Copeland, N. G.; Jenkins,

[42257] N. A.; White, R.; O'Connell, P.: Deletions and a translocation interrupt

[42258] a cloned gene at the neurofibromatosis type 1 locus. Cell 62: 187-192,

[42259] 1990.

[42260]

[42261] 296. Vogel, K. S.; Brannan, C. I.; Jenkins, N. A.; Copeland, N. G.;

[42262] Parada, L. F.: Loss of neurofibromin results in neu-
rotrophin-independent
[42263] survival of embryonic sensory and sympathetic neurons.
Cell 82:
[42264] 733-742, 1995.

[42265]

[42266] 297. Vogel, K. S.; Klesse, L. J.; Velasco-Miguel, S.; Meyers,
K.;
[42267] Rushing, E. J.; Parada, L. F.: Mouse tumor model for neu-
rofibromatosis
[42268] type 1. Science 286: 2176-2179, 1999.

[42269]

[42270] 298. von Recklinghausen, F.: Ueber die multiplen Fibrome
der Haut
[42271] und ihre Beziehung zu den multiplen Neuromen. Berlin:
August Hirschwald
[42272] (pub.) 1882.

[42273]

[42274] 299. Voutsinas, S.; Wynne-Davies, R.: The infrequency of
malignant
[42275] disease in diaphyseal aclasis and neurofibromatosis. J.

Med. Genet. 20:
[42276] 345-349, 1983.

[42277]

[42278] 300. Waggoner, D. J.; Towbin, J.; Gottesman, G.; Gutmann, D. H.:
[42279] Clinic-based study of plexiform neurofibromas in neurofibromatosis
[42280] 1. Am. J. Med. Genet. 92: 132-135, 2000.

[42281]

[42282] 301. Wallace, M.; Fountain, J.; Ledbetter, D. H.; Leach, R.; Fournier,
[42283] R. E. K.; O'Connell, P.; White, R.; Seizinger, B.; Menon, A.; Gusella,
[42284] J. F.; Collins, F. S.: A NotI linking clone detects the 17q11.2 breakpoints
[42285] in two patients with von Recklinghausen neurofibromatosis (NF1). (Abstract) Cytogenet.
[42286] Cell Genet. 51: 1101, 1989.

[42287]

[42288] 302. Wallace, M. R.; Andersen, L. B.; Marchuk, D. A.; Mitchell, A.

[42289] L.; Letcher, R.; Odeh, H.; Saulino, A. M.; Brereton, A.; Collins,
[42290] F. S.: Candidate genes in the von Recklinghausen neurofibromatosis
[42291] (NF1) region. (Abstract) Am. J. Hum. Genet. 47 (suppl.): A240, 1990.
[42292]

[42293] 303. Wallace, M. R.; Andersen, L. B.; Saulino, A. M.; Gregory, P.
[42294] E.; Glover, T. W.; Collins, F. S.: A de novo Alu insertion results
[42295] in neurofibromatosis type 1. Nature 353: 864-866, 1991.
[42296]

[42297] 304. Wallace, M. R.; Marchuk, D. A.; Andersen, L. B.; Letcher, R.;
[42298] Odeh, H. M.; Saulino, A. M.; Fountain, J. W.; Brereton, A.; Nicholson,
[42299] J.; Mitchell, A. L.; Brownstein, B. H.; Collins, F. S.: Type 1 neurofibromatosis
[42300] gene: identification of a large transcript disrupted in three NF1
[42301] patients. Science 249: 181-186, 1990.

[42302]

[42303] 305. Wallis, K.; Deutsch, V.; Azizi, E.: Hypertension in a case of

[42304] Von Recklinghausen's neurofibromatosis. Helv. Paediat. Acta 25:

[42305] 147-153, 1970.

[42306]

[42307] 306. Wang, Q.; Lasset, C.; Desseigne, F.; Frappaz, D.; Bergeron, C.;

[42308] Navarro, C.; Ruano, E.; Puisieux, A.: Neurofibromatosis and early

[42309] onset of cancers in hMLH1-deficient children. Cancer Res. 59: 294-297,

[42310] 1999.

[42311]

[42312] 307. Ward, K.; O'Connell, P.; Carey, J. C.; Leppert, M.; Jol-ley, S.;

[42313] Plaetke, R.; Ogden, B.; White, R.: Diagnosis of neurofibro-matosis

[42314] 1 by using tightly linked, flanking DNA markers. Am. J. Hum. Genet. 46:

[42315] 943-949, 1990.

[42316]

[42317] 308. Weiming, X.; Yu, Q.; Lizhi, L.; Ponder, M.; Wallace, M.; Gangfeng,

[42318] X.; Ponder, B.: Molecular analysis of neurofibromatosis type 1 mutations. Hum.

[42319] Mutat. 1: 474-477, 1992.

[42320]

[42321] 309. Westerhof, W.; Delleman, J. W.; Wolters, E. C. M. J.; Dijkstra,

[42322] P. F.: Neurofibromatosis and hypertelorism. Brit. J. Derm. 109:

[42323] 475-477, 1983.

[42324]

[42325] 310. Winter, R. B.: Spontaneous dislocation of a vertebra in a patient

[42326] who had neurofibromatosis: report of a case with dural ectasia. J.

[42327] Bone Joint Surg. 73A: 1402-1404, 1991.

[42328]

[42329] 311. Wolkenstein, P.; Mahmoudi, A.; Zeller, J.; Revuz, J.: More on
[42330] the frequency of segmental neurofibromatosis. (Letter) Arch. Derm. 131:
[42331] 1465, 1995.

[42332]

[42333] 312. Woody, R. C.; Perrot, L. J.; Beck, S. A.: Neurofibro-matosis
[42334] cerebral vasculopathy in an infant: clinical, neuroradio-graphic, and
[42335] neuropathologic studies. Pediat. Path. 12: 613-619, 1992.

[42336]

[42337] 313. Wu, B.-L.; Austin, M. A.; Schneider, G. H.; Boles, R. G.; Korf,
[42338] B. R.: Deletion of the entire NF1 gene detected by FISH: four deletion
[42339] patients associated with severe manifestations. Am. J. Med. Genet. 59:
[42340] 528-535, 1995.

[42341]

[42342] 314. Wu, B.-L.; Schneider, G. H.; Korf, B. R.: Deletion of

the entire

[42343] NF1 gene causing distinct manifestations in a family. Am. J. Med.

[42344] Genet. 69: 98-101, 1997.

[42345]

[42346] 315. Wu, B. L.; Boles, R. G.; Yaari, H.; Weremowicz, S.; Schneider,

[42347] G. H.; Korf, B. R.: Somatic mosaicism for deletion of the entire

[42348] NF1 gene identified by FISH. Hum. Genet. 99: 209-213, 1997.

[42349]

[42350] 316. Wu, R.; Legius, E.; Robberecht, W.; Dumoulin, M.; Cassiman, J.-J.;

[42351] Fryns, J.-P.: Neurofibromatosis type I gene mutation in a patient

[42352] with features of LEOPARD syndrome. Hum. Mutat. 8: 51-56, 1996.

[42353]

[42354] 317. Xu, G.; O'Connell, P.; Stevens, J.; White, R.: Charac-terization

[42355] of human adenylate kinase 3 (AK3) cDNA and mapping of

the AK3 pseudogene

[42356] to an intron of the NF1 gene. Genomics 13: 537-542, 1992.

[42357]

[42358] 318. Xu, G.; O'Connell, P.; Viskochil, D.; Cawthon, R.; Robertson,

[42359] M.; Culver, M.; Dunn, D.; Stevens, J.; Gesteland, R.; White, R.; Weiss,

[42360] R.: The neurofibromatosis type 1 gene encodes a protein related to

[42361] GAP. Cell 62: 599-608, 1990.

[42362]

[42363] 319. Xu, W.; Lizhi, L.; Yuengde, B.; Yagle, M.; Solomon, E.: Enrichment

[42364] of regional cosmid clones from a human chromosome 17 transfectant

[42365] PLT-6B constructed by chromosome mediated gene transfer. (Abstract) Cytogenet.

[42366] Cell Genet. 51: 1112, 1989.

[42367]

[42368] 320. Yagle, M.; Rider, S. H.; Parruti, G.; Nicolai, H.; Borrow,

J.;

[42369] Mathew, C.; Ponder, B.; Sheer, D.; Ledbetter, D.; Solomon, E.: Isolation

[42370] of cosmids flanking the translocations for acute promyelocytic leukaemia

[42371] (APL) and von Recklinghausen neurofibromatosis (NF1). (Abstract) Cytogenet.

[42372] Cell Genet. 51: 1112, 1989.

[42373]

[42374] 321. Yagle, M. K.; Parruti, G.; Xu, W.; Ponder, B. A. J.; Solomon,

[42375] E.: Genetic and physical map of the von Recklinghausen neurofibromatosis

[42376] (NF1) region on chromosome 17. Proc. Nat. Acad. Sci. 87: 7255-7259,

[42377] 1990.

[42378]

[42379] 322. Yamauchi, T.; Tada, M.; Houkin, K.; Tanaka, T.; Nakamura, Y.;

[42380] Kuroda, S.; Abe, H.; Inoue, T.; Ikezaki, K.; Matsushima, T.; Fukui,

[42381] M.: Linkage of familial moyamoya disease (spontaneous

occlusion of
[42382] the circle of Willis) to chromosome 17q25. Stroke 31:
930-935, 2000.
[42383]

[42384] 323. Yasunari, T.; Shiraki, K.; Hattori, H.; Miki, T.: Fre-
quency
[42385] of choroidal abnormalities in neurofibromatosis type 1.
Lancet 356:
[42386] 988-992, 2000.

[42387]

[42388] 324. Zacharin, M.: Precocious puberty in two children with
neurofibromatosis
[42389] type I in the absence of optic chiasmal glioma. J. Pediat.
130:
[42390] 155-157, 1997.

[42391]

[42392] 325. Zehavi, C.; Romano, A.; Goodman, R. M.: Iris (Lisch)
nodules
[42393] in neurofibromatosis. Clin. Genet. 29: 51-55, 1986.

[42394]

[42395] 326. Zhu, Y.; Ghosh, P.; Charnay, P.; Burns, D. K.; Parada,

L. F.

[42396] : Neurofibromas in NF1: Schwann cell origin and role of tumor environment. Science 296:

[42397] 920-922, 2002.

[42398]

[42399] 327. Zochodne, D.: Von Recklinghausen's vasculopathy. Am. J. Med.

[42400] Sci. 287: 64-65, 1984.

[42401]

[42402] 328. Zonana, J.; Weleber, R. G.: Segmental neurofibromatosis and

[42403] iris hamartomata (Lisch nodules). (Abstract) Proc. Greenwood Genet.

[42404] Center 3: 140-141, 1984.

[42405] 6474.

[42406] Pulst, S.-M.; Riccardi, V. M.; Fain, P.; Korenberg, J. R.: Familial

[42407] spinal neurofibromatosis: clinical and DNA linkage analysis. Neurology 41:

[42408] 1923-1927, 1991.

[42409] 6475.

[42410] Sano, Y.; Nishimukai, H.; Kitamura, H.; Nagaki, K.; Inai, S.;

[42411] Hamasaki, Y.; Maruyama, I.; Igata, A.: Hereditary defi-
ciency of the

[42412] third component of complement in two sisters with sys-
temic lupus erythematosus-like

[42413] symptoms. Arthritis Rheum. 24: 1255-1260, 1981.

[42414] 6476.

[42415] Hadidi, H.; Zahlsen, K.; Idle, J. R.; Cholerton, S.: A single

[42416] amino acid substitution (leu160his) in cytochrome P450
CYP2A6 causes

[42417] switching from 7-hydroxylation to 3-hydroxylation of
coumarin. Food

[42418] Chem. Toxicol. 35: 903-907, 1997.

[42419] 6477.

[42420] Kanazawa, I.: Dentatorubral-pallidoluysian atrophy or
Naito-Oyanagi

[42421] disease. Neurogenetics 2: 1-17, 1998.

[42422] 6478.

[42423] Porjesz, B.; Almasy, L.; Edenberg, H. J.; Wang, K.; Chorlian,
D.

[42424] B.; Foroud, T.; Goate, A.; Rice, J. P.; O'Connor, S. J.; Rohrbaugh,

[42425] J.; Kuperman, S.; Bauer, L. O.; Crowe, R. R.; Schuckit, M. A.; Hesselbrock,

[42426] V.; Conneally, P. M.; Tischfield, J. A.; Li, T.-K.; Reich, T.; Begleiter,

[42427] H.: Linkage disequilibrium between the beta frequency of the human

[42428] EEG and a GABA-A receptor gene locus. Proc. Nat. Acad. Sci. 99:

[42429] 3729-3733, 2002.

[42430] 6479.

[42431] Yoon, S.-J.; LeBlanc-Straceski, J.; Ward, D.; Krauter, K.; Kucherlapati,

[42432] R.: Organization of the human keratin type II gene cluster at 12q13. Genomics 24:

[42433] 502-508, 1994.

[42434] 6480.

[42435] Hermouet, S.; Merendino, J. J., Jr.; Gutkind, J. S.; Spiegel, A.

[42436] M.: Activating and inactivating mutations of the alpha subunit of

[42437] G(i2) protein have opposite effects on proliferation of NIH 3T3 cells. Proc.

[42438] Nat. Acad. Sci. 88: 10455-10459, 1991.

[42439] 6481.

[42440] Lerman, B. B.; Dong, B.; Stein, K. M.; Markowitz, S. M.; Lin-den,

[42441] J.; Catanzaro, D. F.: Right ventricular outflow tract tachy-cardia

[42442] due to a somatic cell mutation in G protein subunit-al-pha-i2. J.

[42443] Clin. Invest. 101: 2862-2868, 1998.

[42444] 6482.

[42445] Lyons, J.; Landis, C. A.; Harsh, G.; Vallar, L.; Grunewald, K.;

[42446] Feichtinger, H.; Duh, Q.-Y.; Clark, O. H.; Kawasaki, E.; Bourne, H.

[42447] R.; McCormick, F.: Two G protein oncogenes in human en-docrine tumors. Science 249:

[42448] 655-659, 1990.

[42449] 6483.

[42450] Rudolph, U.; Finegold, M. J.; Rich, S. S.; Harriman, G. R.; Srinivasan,

[42451] Y.; Brabet, P.; Boulay, G.; Bradley, A.; Birnbaumer, L.: Ulcerative
[42452] colitis and adenocarcinoma of the colon in G alpha(i2)-deficient mice. Nature
[42453] Genet. 10: 143-150, 1995.

[42454] 6484.

[42455] Baron, B.; Fernandez, M. A.; Toledo, F.; Le Roscouet, D.; Mayau,
[42456] V.; Martin, N.; Buttin, G.; Debatisse, M.: The highly conserved Chinese
[42457] hamster GNAI3 gene maps less than 60 kb from the AMPD2 gene and lacks
[42458] the intronic U6 snRNA present in its human counterpart. Genomics 24:
[42459] 288-294, 1994.

[42460] 6485.

[42461] Sissons, J. G. P.; West, R. J.; Fallows, J.; Williams, D. G.;

[42462] Boucher, B. J.; Amos, N.; Peters, D. K.: The complement abnormalities
[42463] of lipodystrophy. New Eng. J. Med. 294: 461-465, 1976.

[42464] 6486.

[42465] Teisberg, P.: New variants in the C3 system. Hum. Hered. 20:
[42466] 631-637, 1970.

[42467] 6487.

[42468] Teisberg, P.: Another variant in the C3 system. Clin. Genet. 2:
[42469] 298-302, 1971.

[42470] 6488.

[42471] Whitehead, A. S.; Sim, R. B.; Bodmer, W. F.: A monoclonal antibody
[42472] against human complement component C3: the production of C3 by human
[42473] cells in vitro. Europ. J. Immun. 11: 140-146, 1981.

[42474] 6489.

[42475] Whitehead, A. S.; Solomon, E.; Chambers, S.; Bodmer, W. F.; Povey,
[42476] S.; Fey, G.: Assignment of the structural gene for the third component
[42477] of human complement to chromosome 19. Proc. Nat. Acad. Sci. 79:
[42478] 5021-5025, 1982.

[42479] 6490.

[42480] Whitehead, A. S.; Solomon, E.; Chambers, S. P.; Povey, S.; Bodmer,

[42481] W. F.: Assignment of the gene for the third component of human complement

[42482] (C3) to chromosome 19 using human-mouse somatic cell hybrids. (Abstract) Cytogenet.

[42483] Cell Genet. 32: 326-327, 1982.

[42484] 6491.

[42485] Wieme, R. J.; Demeulenaere, L.: Genetically determined electrophoretic

[42486] variant of the human complement component C-prime-3. Nature 214:

[42487] 1042-1043, 1967.

[42488] 6492.

[42489] Williamson, R.: Personal Communication. London, England 8/25/1983.

[42490] 6493.

[42491] Winkelstein, J. A.; Cork, L. C.; Griffin, D. E.; Griffin, J. W.;

[42492] Adams, R. J.; Price, D. L.: Genetically determined deficiency of

[42493] the third component of complement in the dog. Science 212: 1169-1170,
[42494] 1981.
[42495] 6494.
[42496] Kratz, F.: Coumarin-7-hydroxylase activity in microsomes from
[42497] needle biopsies of normal and diseased human liver. Eu-rop. J. Clin.
[42498] Pharm. 10: 133-137, 1976.
[42499] 6495.
[42500] London, S. J.; Idle, J. R.; Daly, A. K.; Coetzee, G. A.: Genetic
[42501] variation of CYP2A6, smoking, and risk of cancer. Lancet 353: 898-899,
[42502] 1999.
[42503] 6496.
[42504] Miyamoto, M.; Umetsu, Y.; Dosaka-Akita, H.; Sawamura, Y.; Yokota,
[42505] J.; Kunitoh, H.; Nemoto, N.; Sato, K.; Ariyoshi, N.; Ka-mataki, T.
[42506] : CYP2A6 gene deletion reduces susceptibility to lung cancer. Biochem.

[42507] Biophys. Res. Commun. 261: 658-660, 1999.

[42508] 6497.

[42509] Nakajima, M.; Yamamoto, T.; Nunoya, K.; Yokoi, T.; Na-

gashima, K.;

[42510] Inoue, K.; Funae, Y.; Shimada, N.; Kamataki, T.; Kuroiwa,

Y.: Role

[42511] of human cytochrome P4502A6 in C-oxidation of nico-

tine. Drug Metab.

[42512] Dispos. 24: 1212-1217, 1996.

[42513] 6498.

[42514] Oscarson, M.; Gullsten, H.; Rautio, A.; Bernal, M. L.; Sin-

ues,

[42515] B.; Dahl, M.-L.; Stengard, J. H.; Pelkonen, O.; Raunio, H.;

Ingelman-Sundberg,

[42516] M.: Genotyping of human cytochrome P450 2A6 (CYP2A6),

a nicotine

[42517] C-oxidase. FEBS Lett. 438: 201-205, 1998.

[42518] 6499.

[42519] Oscarson, M.; McLellan, R. A.; Gullsten, H.; Yue, Q.-Y.;

Lang,

[42520] M. A.; Bernal, M. L.; Sinues, B.; Hirvonen, A.; Raunio, H.;

Pelkonen,
[42521] O.; Ingelman-Sundberg, M.: Characterisation and PCR-based detection
[42522] of a CYP2A6 gene deletion found at a high frequency in a Chinese population. FEBS
[42523] Lett. 448: 105-110, 1999.

[42524] 6500.

[42525] Pianezza, M. L.; Sellers, E. M.; Tyndale, R. F.: Nicotine metabolism
[42526] defect reduces smoking. Nature 393: 750 only, 1998.

[42527] 6501.

[42528] Sabol, S. Z.; Hamer, D. H.: An improved assay shows no association
[42529] between the CYP2A6 gene and cigarette smoking behavior. Behav. Genet. 29:
[42530] 257-261, 1999.

[42531] 6502.

[42532] Wood, A. W.; Conney, A. H.: Genetic variation in coumarin hydroxylase
[42533] activity in mouse (Mus musculus). Science 185: 612-614, 1974.

[42534] 6503.

[42535] Yamano, S.; Nagata, K.; Yamazoe, Y.; Kato, R.; Gelboin, H. V.;

[42536] Gonzalez, F. J.: cDNA and deduced amino acid sequences of human P450

[42537] IIA3 (CYP2A3). Nucleic Acids Res. 17: 4888 only, 1989.

[42538] 6504.

[42539] Yokoi, T.; Kamataki, T.: Genetic polymorphism of drug metabolizing

[42540] enzymes: new mutations in CYP2D6 and CYP2A6 gene in Japanese. Pharm.

[42541] Res. 15: 517-524, 1998.

[42542] 6505.

[42543] Koide, R.; Ikeuchi, T.; Onodera, O.; Tanaka, H.; Igarashi, S.;

[42544] Endo, K.; Takahashi, H.; Kondo, R.; Ishikawa, A.; Hayashi, T.; Saito,

[42545] M.; Tomoda, A.; Miike, T.; Naito, H.; Ikuta, F.; Tsuji, S.: Unstable

[42546] expansion of CAG repeat in hereditary dentatorubral-pal-lidoluysian

[42547] atrophy (DRPLA). Nature Genet. 6: 9-13, 1994.

[42548] 6506.

[42549] Kondo, I.; Ohta, H.; Yazaki, M.; Ikeda, J.-E.; Gusella, J. F.;

[42550] Kanazawa, I.: Exclusion mapping of the hereditary denta-
torubropallidoluysian

[42551] atrophy gene from the Huntington's disease locus. J. Med.
Genet. 27:

[42552] 105-108, 1990.

[42553] 6507.

[42554] Kuwano, A.; Morimoto, Y.; Nagai, T.; Fukushima, Y.;
Ohashi, H.;

[42555] Hasegawa, T.; Kondo, I.: Precise chromosomal locations of
the genes

[42556] for dentatorubral-pallidoluysian atrophy (DRPLA), von
Willebrand factor

[42557] (F8vWF) and parathyroid hormone-like hormone (PTHLH)
in human chromosome

[42558] 12p by deletion mapping. Hum. Genet. 97: 95-98, 1996.

[42559] 6508.

[42560] Li, S.-H.; McInnis, M. G.; Margolis, R. L.; Antonarakis, S. E.;

[42561] Ross, C. A.: Novel triplet repeat containing genes in human brain:
[42562] cloning, expression, and length polymorphisms. Genomics 16: 572-579,
[42563] 1993.

[42564] 6509.

[42565] Naito, H.; Oyanagi, S.: Familial myoclonus epilepsy and choreoathetosis:
[42566] hereditary dentatorubral-pallidoluysian atrophy. Neurology 32: 798-807,
[42567] 1982.

[42568] 6510.

[42569] Arcioni, L.; Simeone, A.; Guazzi, S.; Zappavigna, V.; Boncinelli,
[42570] E.; Mavilio, F.: The upstream region of the human homeobox gene HOX3D
[42571] is a target for regulation by retinoic acid and HOX homeoproteins. EMBO
[42572] J. 11: 265-277, 1992.

[42573] 6511.

[42574] Godwin, A. R.; Capecchi, M. R.: Hoxc13 mutant mice lack

external

[42575] hair. Genes Dev. 12: 11-20, 1998.

[42576] 6512.

[42577] Goodman, F. R.: Limb malformations and the human HOX genes. Am.

[42578] J. Med. Genet. 112: 256-265, 2002.

[42579] 6513.

[42580] Magli, M. C.; Barba, P.; Celetti, A.; De Vita, G.; Cillo, C.; Boncinelli,

[42581] E.: Coordinate regulation of HOX genes in human hematopoietic cells. Proc.

[42582] Nat. Acad. Sci. 88: 6348-6352, 1991.

[42583] 6514.

[42584] Pravtcheva, D.; Newman, M.; Hunihan, L.; Lonai, P.; Rud-

dle, F.

[42585] H.: Chromosome assignment of the murine Hox-4.1 gene. Genomics 5:

[42586] 541-545, 1989.

[42587] 6515.

[42588] Taniguchi, Y.; Komatsu, N.; Moriuchi, T.: Overexpression

of the
[42589] HOX4A (HOXD3) homeobox gene in human ery-
throleukemia HEL cells results
[42590] in altered adhesive properties. Blood 85: 2786-2794,
1995.
[42591] 6516.

[42592] Zakany, J.; Duboule, D.: Hox genes and the making of
sphincters. Nature 401:
[42593] 761-762, 1999.

[42594] 6517.

[42595] Mavilio, F.; Simeone, A.; Giampaolo, A.; Faiella, A.; Zap-
pavigna,
[42596] V.; Acampora, D.; Poiana, G.; Russo, G.; Peschle, C.;
Boncinelli,
[42597] E.: Differential and stage-related expression in embryonic
tissues
[42598] of a new human homoeobox gene. Nature 324: 664-668,
1986.
[42599] 6518.

[42600] Oliver, G.; Sidell, N.; Fiske, W.; Heinzmann, C.; Mohandas,
T.;
[42601] Sparkes, R. S.; De Robertis, E. M.: Complementary homeo

protein gradients

[42602] in developing limb buds. Genes Dev. 3: 641-650, 1989.

[42603] 6519.

[42604] Shaw, D. J.; Meredith, A. L.; Brook, J. D.; Sarfarazi, M.; Harley,

[42605] H. G.; Huson, S. M.; Bell, G. I.; Harper, P. S.: Linkage relationships

[42606] of the insulin receptor gene with the complement component 3, LDL

[42607] receptor, apolipoprotein C2 and myotonic dystrophy loci on chromosome

[42608] 19. Hum. Genet. 74: 267-269, 1986.

[42609] 6520.

[42610] Blangy, A.; Lane, H. A.; d'Herin, P.; Harper, M.; Kress, M.; Nigg,

[42611] E. A.: Phosphorylation by p34(cdc2) regulates spindle association

[42612] of human Eg5, a kinesin-related motor essential for bipolar spindle

[42613] formation in vivo. Cell 83: 1159-1169, 1995.

[42614] 6521.

[42615] Mock, B. A.; Givol, D.; D'Hoostelaere, L. A.; Huppi, K.; Seldin,
[42616] M. F.; Gurfinkel, N.; Unger, T.; Potter, M.; Mushinski, J. F.: Mapping
[42617] of the bcl-2 oncogene on mouse chromosome 1. Cytogenet. Cell Genet. 47:
[42618] 11-15, 1988.

[42619] 6522.

[42620] Shimada, F.; Suzuki, Y.; Taira, M.; Hashimoto, N.; Nozaki, O.;
[42621] Makino, H.; Yoshida, S.: Abnormal messenger ribonucleic acid (mRNA)
[42622] transcribed from a mutant insulin receptor gene in a patient with
[42623] type A insulin resistance. Diabetologia 35: 639-644, 1992.
[42624] 6523.

[42625] Shimada, F.; Taira, M.; Suzuki, Y.; Hashimoto, N.; Nozaki, O.;
[42626] Taira, M.; Tatibana, M.; Ebina, Y.; Tawata, M.; Onaya, T.; Makino,
[42627] H.; Yoshida, S.: Insulin-resistant diabetes associated with

partial
[42628] deletion of insulin-receptor gene. Lancet 335:
1179-1181, 1990.
[42629] 6524.

[42630] Taira, M.; Taira, M.; Hashimoto, N.; Shimada, F.; Suzuki,
Y.;
[42631] Kanatsuka, A.; Nakamura, F.; Ebina, Y.; Tatibana, M.;
Makino, H.;
[42632] Yoshida, S.: Human diabetes associated with a deletion of
the tyrosine
[42633] kinase domain of the insulin receptor. Science 245:
63-66, 1989.
[42634] 6525.

[42635] Takahashi, Y.; Kadowaki, H.; Ando, A.; Quin, J. D.; Mac-
Cuish,
[42636] A. C.; Yazaki, Y.; Akanuma, Y.; Kadowaki, T.: Two aberrant
splicings
[42637] caused by mutations in the insulin receptor gene in cul-
tured lymphocytes
[42638] from a patient with Rabson-Mendenhall's syndrome. J.
Clin. Invest. 101:
[42639] 588-594, 1998.

[42640] 6526.

[42641] Taylor, S. I.; Cama, A.; Accili, D.; Barbetti, F.; Imano, E.;

[42642] Kadowaki, H.; Kadowaki, T.: Genetic basis of endocrine disease 1:

[42643] molecular genetics of insulin resistant diabetes mellitus. J. Clin.

[42644] Endocr. Metab. 73: 1158-1163, 1991.

[42645] 6527.

[42646] Taylor, S. I.; Marcus-Samuels, B.; Ryan-Young, J.; Leven-thal,

[42647] S.; Elders, M. J.: Genetics of the insulin receptor defect in a patient

[42648] with extreme insulin resistance. J. Clin. Endocr. Metab. 62: 1130-1135,

[42649] 1986.

[42650] 6528.

[42651] Taylor, S. I.; Underhill, L. H.; Hedo, J. A.; Roth, J.; Serrano

[42652] Rios, M.; Blizzard, R. M.: Decreased insulin binding to cul-tured

[42653] cells from a patient with the Rabson-Mendenhall syn-drome: dichotomy

[42654] between studies with cultured lymphocytes and cultured fibroblasts. J.

[42655] Clin. Endocr. Metab. 56: 856-861, 1983.

[42656] 6529.

[42657] Tevaarwerk, G. J. M.; Strickland, K. P.; Lin, C. H.; Hudson, A.

[42658] J.: Studies on insulin resistance and insulin receptor binding in

[42659] myotonia dystrophica. J. Clin. Endocr. Metab. 49: 216-222, 1979.

[42660] 6530.

[42661] Ullrich, A.; Bell, J. R.; Chen, E. Y.; Herrera, R.; Petruzzelli,

[42662] L. M.; Dull, T. J.; Gray, A.; Coussens, L.; Liao, Y.-C.; Tsubokawa,

[42663] M.; Mason, A.; Seeburg, P. H.; Grunfeld, C.; Rosen, O. M.; Ramachandran,

[42664] J.: Human insulin receptor and its relationship to the tyrosine kinase

[42665] family of oncogenes. Nature 313: 756-761, 1985.

[42666] 6531.

[42667] van der Vorm, E. R.; Kuipers, A.; Kielkopf-Renner, S.;

Krans,

[42668] H. M. J.; Moller, W.; Maassen, J. A.: A mutation in the insulin receptor

[42669] that impairs proreceptor processing but not insulin binding. J. Biol.

[42670] Chem. 269: 14297-14302, 1994.

[42671] 6532.

[42672] van der Vorm, E. R.; van der Zon, G. C. M.; Moller, W.; Krans,

[42673] H. M. J.; Lindhout, D.; Maassen, J. A.: An arg for gly substitution

[42674] at position 31 in the insulin receptor, linked to insulin resistance,

[42675] inhibits receptor processing and transport. J. Biol. Chem. 267:

[42676] 66-71, 1992.

[42677] 6533.

[42678] Ward, G. M.; Harrison, L. C.: Structure of the human erythrocyte

[42679] insulin receptor. Diabetes 35: 101-105, 1986.

[42680] 6534.

[42681] Wertheimer, E.; Lu, S.-P.; Backeljauw, P. F.; Davenport, M. L.;

[42682] Taylor, S. I.: Homozygous deletion of the human insulin receptor

[42683] gene results in leprechaunism. Nature Genet. 5: 71-73, 1993.

[42684] 6535.

[42685] Williams, D. L.; Look, A. T.; Melvin, S. L.; Roberson, P. K.;

[42686] Dahl, G.; Flake, T.; Stass, S.: New chromosomal translocations correlate

[42687] with specific immunophenotypes of childhood acute lymphoblastic leukemia. Cell 36:

[42688] 101-109, 1984.

[42689] 6536.

[42690] Williams, J. F.; McClain, D. A.; Dull, T. J.; Ullrich, A.; Olefsky,

[42691] J. M.: Characterization of an insulin receptor mutant lacking the

[42692] subunit processing site. J. Biol. Chem. 265: 8463-8469, 1990.

[42693] 6537.

[42694] Yamamoto-Honda, R.; Koshio, O.; Tobe, K.; Shibasaki, Y.; Momomura,
[42695] K.; Odawara, M.; Kadowaki, T.; Takaku, F.; Akanuma, Y.; Kasuga, M.
[42696] : Phosphorylation state and biological function of a mutant human
[42697] insulin receptor val(996). J. Biol. Chem. 265: 14777-14783, 1990.
[42698] 6538.

[42699] Yang-Feng, T. L.; Francke, U.; Ullrich, A.: Gene for human insulin
[42700] receptor: localization to site on chromosome 19 involved in pre-B-cell
[42701] leukemia. Science 228: 728-731, 1985.

[42702] 6539.

[42703] Yoshimasa, Y.; Seino, S.; Whittaker, J.; Kakehi, T.; Kosaki, A.;
[42704] Kuzuya, H.; Imura, H.; Bell, G. I.; Steiner, D. F.: Insulin-resistant
[42705] diabetes due to a point mutation that prevents insulin proreceptor
[42706] processing. Science 240: 784-787, 1988.

[42707] 6540.

[42708] Kurachi, H.; Jobo, K.; Ohta, M.; Kawasaki, T.; Itoh, N.: A new

[42709] member of the insulin receptor family, insulin receptor-related receptor,

[42710] is expressed preferentially in the kidney. Biochem. Biophys. Res.

[42711] Commun. 187: 934-939, 1992.

[42712] 6541.

[42713] Shier, P.; Watt, V. M.: Primary structure of a putative receptor

[42714] for a ligand of the insulin family. J. Biol. Chem. 264: 14605-14608,

[42715] 1989.

[42716] 6542.

[42717] Shier, P.; Willard, H. F.; Watt, V. M.: Localization of the insulin

[42718] receptor-related receptor gene to human chromosome 1. Cytogenet.

[42719] Cell Genet. 54: 80-81, 1990.

[42720] 6543.

[42721] Whitmore, T. E.; Maurer, M. F.; Day, H. L.; Jelmberg, A. C.; Dasovich,

[42722] M. M.; Sundborg, L. M.; Burkhead, S. K.; Heipel, M. D.; Madden, K.

[42723] L.; Kramer, J. M.; Kuijper, J. L.; Xu, W. F.; Jaspers, S. R.; Holly,

[42724] R. D.; Lok, S.: The assignment of the human insulin receptor-related

[42725] receptor gene (INSRR) to chromosome 1q21-q23 by the use of radiation

[42726] hybrid mapping. Cytogenet. Cell Genet. 87: 93-94, 1999.

[42727] 6544.

[42728] Erickson, J. C.; Hollopeter, G.; Palmiter, R. D.: Attenuation

[42729] of the obesity syndrome of ob/ob mice by the loss of neuropeptide

[42730] Y. Science 274: 1704-1706, 1996.

[42731] 6545.

[42732] Lutz, C. M.; Frankel, W. N.; Richards, J. E.; Thompson, D. A.:

[42733] Neuropeptide Y receptor genes on human chromosome 4q31-q32 map to

[42734] conserved linkage groups on mouse chromosomes 3 and 8. Genomics 41:

[42735] 498-500, 1997.

[42736] 6546.

[42737] Ammar, D. A.; Eadie, D. M.; Wong, D. J.; Ma, Y.-Y.; Ko-lakowski,

[42738] L. F., Jr.; Yang-Feng, T. L.; Thompson, D. A.: Characteriza-tion of

[42739] the human type 2 neuropeptide Y receptor gene (NPY2R) and localization

[42740] to the chromosome 4q region containing the type 1 neu-ropeptide Y receptor

[42741] gene. Genomics 38: 392-398, 1996.

[42742] 6547.

[42743] Gerald, C.; Walker, M. W.; Vaysse, P. J.-J.; He, C.; Branchek,

[42744] T. A.; Weinshank, R. L.: Expression cloning and pharmaco-logical characterization

[42745] of a human hippocampal neuropeptide Y/peptide YY Y2 receptor subtype. J.

[42746] Biol. Chem. 270: 26758-26761, 1995.

[42747] 6548.

[42748] Solomon, S. D.; Geisterfer-Lowrance, A. A. T.; Vosberg, H.-P.;

[42749] Hiller, G.; Jarcho, J. A.; Morton, C. C.; McBride, W. O.; Mitchell,

[42750] A. L.; Bale, A. E.; McKenna, W. J.; Seidman, J. G.; Seidman, C. E.

[42751] : A locus for familial hypertrophic cardiomyopathy is closely linked

[42752] to the cardiac myosin heavy chain genes, CRI-L436, and CRI-L329 on

[42753] chromosome 14 at q11-q12. Am. J. Hum. Genet. 47: 389-394, 1990.

[42754] 6549.

[42755] Tanigawa, G.; Jarcho, J. A.; Kass, S.; Solomon, S. D.; Vosberg,

[42756] H.-P.; Seidman, J. G.; Seidman, C. E.: A molecular basis for familial

[42757] hypertrophic cardiomyopathy: an alpha/beta cardiac myosin heavy chain

[42758] hybrid gene. Cell 62: 991-998, 1990.

[42759] 6550.

[42760] Teare, D.: Asymmetrical hypertrophy of the heart in young

adults. Brit.

[42761] Heart J. 20: 1-8, 1958.

[42762] 6551.

[42763] Weydert, A.; Daubas, P.; Lazaridis, I.; Barton, P.; Garner, I.;

[42764] Leader, D. P.; Bonhomme, F.; Catalan, J.; Simon, D.;
Guenet, J. L.;

[42765] Gros, F.; Buckingham, M. E.: Genes for skeletal muscle
myosin heavy

[42766] chains are clustered and are not located on the same
mouse chromosome

[42767] as a cardiac myosin heavy chain gene. Proc. Nat. Acad. Sci.
82:

[42768] 7183-7187, 1985.

[42769] 6552.

[42770] Yamauchi-Takihara, K.; Sole, M. J.; Liew, J.; Ing, D.; Liew,
C.

[42771] C.: Characterization of human cardiac myosin heavy chain
genes. Proc.

[42772] Nat. Acad. Sci. 86: 3504-3508, 1989.

[42773] 6553.

[42774] Stewart, R. J.; Pesavento, P. A.; Woerpel, D. N.; Goldstein,

L.

[42775] S. B.: Identification and partial characterization of six members

[42776] of the kinesin superfamily in Drosophila. Proc. Nat. Acad. Sci. 88:

[42777] 8470-8474, 1991.

[42778] 6554.

[42779] Tihy, F.; Kress, M.; Harper, M.; Dutrillaux, B.; Lemieux, N.:

[42780] Localization of the human kinesin-related gene to band 10q24 by fluorescence

[42781] in situ hybridization. Genomics 13: 1371-1372, 1992.

[42782] 6555.

[42783] Pasteris, N. G.; Trask, B.; Sheldon, S.; Gorski, J. L.: A chromosome

[42784] deletion 2q35-36 spanning loci HuP2 and COL4A3 results in Waardenburg

[42785] syndrome type III (Klein-Waardenburg syndrome). (Abstract) Am. J.

[42786] Hum. Genet. 51 (suppl.): A224, 1992.

[42787] 6556.

[42788] Lee, M. K.; Stirling, W.; Xu, Y.; Xu, X.; Qui, D.; Mandir, A.

[42789] S.; Dawson, T. M.; Copeland, N. G.; Jenkins, N. A.; Price, D. L.:
[42790] Human alpha-synuclein-harboring familial Parkinson's disease-linked
[42791] ala53-to-thr mutation causes neurodegenerative disease with alpha-synuclein
[42792] aggregation in transgenic mice. Proc. Nat. Acad. Sci. 99: 8968-8973,
[42793] 2002.

[42794] 6557.

[42795] Markopoulou, K.; Wszolek, Z. K.; Pfeiffer, R. F.: A Greek-American
[42796] kindred with autosomal dominant, levodopa-responsive Parkinsonism
[42797] and anticipation. Ann. Neurol. 38: 373-378, 1995.

[42798] 6558.

[42799] Maroteaux, L.; Scheller, R. H.: The rat brain synucleins: family
[42800] of proteins transiently associated with neuronal mem-brane. Molec.
[42801] Brain Res. 11: 335-343, 1991.

[42802] 6559.

[42803] Masliah, E.; Rockenstein, E.; Veinbergs, I.; Mallory, M.; Hashimoto,

[42804] M.; Takeda, A.; Sagara, Y.; Sisk, A.; Mucke, L.: Dopaminergic loss

[42805] and inclusion body formation in alpha-synuclein mice: implications

[42806] for neurodegenerative disorders. Science 287: 1265-1269, 2000.

[42807] 6560.

[42808] Masliah, E.; Rockenstein, E.; Veinbergs, I.; Sagara, Y.; Mallory,

[42809] M.; Hashimoto, M.; Mucke, L.: Beta-amyloid peptides enhance alpha-synuclein

[42810] accumulation and neuronal deficits in a transgenic mouse model linking

[42811] Alzheimer's disease and Parkinson's disease. Proc. Nat. Acad. Sci. 98:

[42812] 12245-12250, 2001.

[42813] 6561.

[42814] Narhi, L.; Wood, S. J.; Steavenson, S.; Jiang, Y.; Wu, G. M.;

[42815] Anafi, D.; Kaufman, S. A.; Martin, F.; Sitney, K.; Denis, P.;
Louis,

[42816] J.-C.; Wypych, J.; Biere, A. L.; Citron, M.: Both familial
Parkinson's

[42817] disease mutations accelerate alpha-synuclein aggrega-
tion. J. Biol.

[42818] Chem. 274: 9843-9846, 1999.

[42819] 6562.

[42820] Polymeropoulos, M. H.; Higgins, J. J.; Golbe, L. I.; Johnson,

[42821] W. G.; Ide, S. E.; Di Iorio, G.; Sanges, G.; Stenroos, E. S.;
Pho,

[42822] L. T.; Schaffer, A. A.; Lazzarini, A. M.; Nussbaum, R. L.;
Duvoisin,

[42823] R. C.: Mapping of a gene for Parkinson's disease to chro-
mosome 4q21-q23. Science 274:

[42824] 1197-1198, 1996.

[42825] 6563.

[42826] Polymeropoulos, M. H.; Lavedan, C.; Leroy, E.; Ide, S. E.;
Dehejia,

[42827] A.; Dutra, A.; Pike, B.; Root, H.; Rubenstein, J.; Boyer, R.;

Stenroos,
[42828] E. S.; Chandrasekharappa, S.; Athanassiadou, A.; Papepetropoulos,
[42829] T.; Johnson, W. G.; Lazzarini, A. M.; Duvoisin, R. C.; Di Iorio, G.;
[42830] Golbe, L. I.; Nussbaum, R. L.: Mutation in the alpha-synuclein gene
[42831] identified in families with Parkinson's disease. Science 276: 2045-2047,
[42832] 1997.

[42833] 6564.

[42834] Scott, W. K.; Stajich, J. M.; Yamaoka, L. H.; Speer, M. C.; Vance,
[42835] J. M.; Roses, A. D.; Pericak-Vance, M. A.; Deane Laboratory Parkinson
[42836] Disease Research Group: Genetic complexity and Parkinson's disease.
[42837] (Letter) Science 277: 387-388, 1997.

[42838] 6565.

[42839] Scott, W. K.; Yamaoka, L. H.; Stajich, J. M.; Scott, B. L.; Vance,
[42840] J. M.; Roses, A. D.; Pericak-Vance, M. A.; Watts, R. L.;

Nance, M.;

[42841] Hubble, J.; Koller, W.; Stern, M. B.; and 15 others: The alpha-synuclein

[42842] gene is not a major risk factor in familial Parkinson disease. (Letter) Neurogenetics 2:

[42843] 191-192, 1999.

[42844] 6566.

[42845] Shibasaki, Y.; Baillie, D. A. M.; St. Clair, D.; Brookes, A. J.

[42846] : High-resolution mapping of SNCA encoding a-synuclein, the non-A-beta

[42847] component of Alzheimer's disease amyloid precursor, to human chromosome

[42848] 4q21.3-q22 by fluorescence in situ hybridization. Cytogenet. Cell

[42849] Genet. 71: 54-55, 1995.

[42850] 6567.

[42851] Shimura, H.; Schlossmacher, M. G.; Hattori, N.; Frosch, M. P.;

[42852] Trockenbacher, A.; Schneider, R.; Mizuno, Y.; Kosik, K. S.; Selkoe,

[42853] D. J.: Ubiquitination of a new form of alpha-synuclein by

parkin
[42854] from human brain: implications for Parkinson's disease. Science 293:
[42855] 263-269, 2001.

[42856] 6568.

[42857] Spillantini, M. G.; Divane, A.; Goedert, M.: Assignment of human
[42858] alpha-synuclein (SNCA) and beta-synuclein (SNCB) genes to chromosomes
[42859] 4q21 and 5q35. Genomics 27: 379-381, 1995.

[42860] 6569.

[42861] Spillantini, M. G.; Schmidt, M. L.; Lee, V. M.-Y.; Tro-
janowski,
[42862] J. Q.; Jakes, R.; Goedert, M.: Alpha-synuclein in Lewy bod-
ies. Nature 388:
[42863] 839-840, 1997.

[42864] 6570.

[42865] Tabrizi, S. J.; Orth, M.; Wilkinson, J. M.; Taanman, J.-W.;
Warner,
[42866] T. T.; Cooper, J. M.; Schapira, A. H. V.: Expression of mu-
tant alpha-synuclein

[42867] causes increased susceptibility to dopamine toxicity. Hum. Molec.

[42868] Genet. 9: 2683-2689, 2000.

[42869] 6571.

[42870] Tanaka, Y.; Engelender, S.; Igarashi, S.; Rao, R. K.; Wanner,

[42871] T.; Tanzi, R. E.; Sawa, A.; Dawson, V. L.; Dawson, T. M.; Ross, C.

[42872] A.: Inducible expression of mutant alpha-synuclein de-

creases proteasome

[42873] activity and increases sensitivity to mitochondria-depen-

dent apoptosis. Hum.

[42874] Molec. Genet. 10: 919-926, 2001.

[42875] 6572.

[42876] Ueda, K.; Fukushima, H.; Masliah, E.; Xia, Y.; Iwai, A.; Yoshimoto,

[42877] M.; Otero, D. A. C.; Kondo, J.; Ihara, Y.; Saitoh, T.: Molecu-

lar

[42878] cloning of cDNA encoding an unrecognized component of amyloid in Alzheimer

[42879] disease. Proc. Nat. Acad. Sci. 90: 11282-11286, 1993.

[42880] 6573.

[42881] Vaughan, J. R.; Farrer, M. J.; Wszolek, Z. K.; Gasser, T.; Durr,
[42882] A.; Agid, Y.; Bonifati, V.; DeMichele, G.; Volpe, G.; Lincoln, S.;
[42883] Breteler, M.; Meco, G.; Brice, A.; Marsden, C. D.; Hardy, J.; Wood,
[42884] N. W.; European Consortium on Genetic Susceptibility in Parkinson's
[42885] Disease (GSPD): Sequencing of the alpha-synuclein gene in a large
[42886] series of cases of familial Parkinson's disease fails to reveal any
[42887] further mutations. Hum. Molec. Genet. 7: 751-753, 1998.

[42888] 6574.

[42889] Xu, J.; Kao, S.-Y.; Lee, F. J. S.; Song, W.; Jin, L.-W.; Yankner,
[42890] B. A.: Dopamine-dependent neurotoxicity of alpha-synuclein: a mechanism
[42891] for selective neurodegeneration in Parkinson disease. Nature Med. 8:
[42892] 600-606, 2002.

[42893] 6575.

[42894] Cetta, F.; Chiappetta, G.; Melillo, R. M.; Petracci, M.; Montalto,
[42895] G.; Santoro, M.; Fusco, A.: The ret/ptc1 oncogene is activated in
[42896] familial adenomatous polyposis-associated thyroid papillary carcinomas. J.
[42897] Clin. Endocr. Metab. 83: 1003-1006, 1998.

[42898] 6576.

[42899] Decker, R. A.; Peacock, M. L.; Watson, P.: Hirschsprung disease
[42900] in MEN 2A: increased spectrum of RET exon 10 genotypes and strong
[42901] genotype-phenotype correlation. Hum. Molec. Genet. 7: 129-134, 1998.
[42902] 6577.

[42903] Donis-Keller, H.; Dou, S.; Chi, D.; Carlson, K. M.; Toshima, K.;
[42904] Lairmore, T. C.; Howe, J. R.; Moley, J. F.; Goodfellow, P.; Wells,
[42905] S. A., Jr.: Mutations in the RET proto-oncogene are associated with
[42906] MEN 2A and FMTC. Hum. Molec. Genet. 2: 851-856, 1993.

[42907] 6578.

[42908] Doray, B.; Salomon, R.; Amiel, J.; Pelet, A.; Touraine, R.; Billaud,

[42909] M.; Attie, T.; Bachy, B.; Munnich, A.; Lyonnet, S.: Mutation of the

[42910] RET ligand, neurturin, supports multigenic inheritance in Hirschsprung

[42911] disease. Hum. Molec. Genet. 7: 1449-1452, 1998.

[42912] 6579.

[42913] Edery, P.; Lyonnet, S.; Mulligan, L. M.; Pelet, A.; Dow, E.; Abel,

[42914] L.; Holder, S.; Nihoul-Fekete, C.; Ponder, B. A. J.; Munnich, A.:

[42915] Mutations of the RET proto-oncogene in Hirschsprung's disease. Nature 367:

[42916] 378-380, 1994.

[42917] 6580.

[42918] Eng, C.: The RET proto-oncogene in multiple endocrine neoplasia

[42919] type 2 and Hirschsprung's disease. New Eng. J. Med. 335: 943-951,

[42920] 1996.

[42921] 6581.

[42922] Eng, C.; Crossey, P. A.; Mulligan, L. M.; Healey, C. S.;
Houghton,
[42923] C.; Prowse, A.; Chew, S. L.; Dahia, P. L. M.; O'Riordan, J. L.
H.;
[42924] Toledo, S. P. A.; Smith, D. P.; Maher, E. R.; Ponder, B. A. J.:
Mutations
[42925] in the RET proto-oncogene and the von Hippel-Lindau
disease tumour
[42926] suppressor gene in sporadic and syndromic phaeochro-
mocytomas. J.
[42927] Clin. Genet. 32: 934-937, 1995.

[42928] 6582.

[42929] Eng, C.; Mulligan, L. M.: Mutations of the RET proto-
oncogene
[42930] in the multiple endocrine neoplasia type 2 syndromes, re-
lated sporadic
[42931] tumours, and Hirschsprung disease. Hum. Mutat. 9:
97-109, 1997.

[42932] 6583.

[42933] Eng, C.; Mulligan, L. M.; Smith, D. P.; Healey, C. S.; Frilling,

[42934] A.; Raue, F.; Neumann, H. P. H.; Pfragner, R.; Behmel, A.; Lorenzo,

[42935] M. J.; Stonehouse, T. J.; Ponder, M. A.; Ponder, B. A. J.: Mutation

[42936] of the RET protooncogene in sporadic medullary thyroid carcinoma. Genes

[42937] Chromosomes Cancer 12: 209-212, 1995.

[42938] 6584.

[42939] Eng, C.; Smith, D. P.; Mulligan, L. M.; Healey, C. S.; Zvelebil,

[42940] M. J.; Stonehouse, T. J.; Ponder, M. A.; Jackson, C. E.; Waterfield,

[42941] M. D.; Ponder, B. A. J.: A novel point mutation in the tyrosine kinase

[42942] domain of the RET proto-oncogene in sporadic medullary thyroid carcinoma

[42943] and in a family with FMTC. Oncogene 10: 509-513, 1995.

[42944] 6585.

[42945] Eng, C.; Smith, D. P.; Mulligan, L. M.; Nagai, M. A.; Healey,

[42946] C. S.; Ponder, M. A.; Gardner, E.; Scheumann, G. F. W.;

Jackson, C.

[42947] E.; Tunnacliffe, A.; Ponder, B. A. J.: Point mutation within the

[42948] tyrosine kinase domain of the RET proto-oncogene in multiple endocrine

[42949] neoplasia type 2B and related sporadic tumors. Hum. Molec. Genet. 3:

[42950] 237-241, 1994.

[42951] 6586.

[42952] Fearon, E. R.: Human cancer syndromes: clues to the origin and

[42953] nature of cancer. Science 278: 1043-1050, 1997.

[42954] 6587.

[42955] Fitze, G.; Schreiber, M.; Kuhlisch, E.; Schackert, H. K.; Roesner,

[42956] D.: Association of RET protooncogene codon 45 polymorphism with Hirschsprung

[42957] disease. (Letter) Am. J. Hum. Genet. 65: 1469-1473, 1999.

[42958] 6588.

[42959] Gardner, E.; Mulligan, L. M.; Eng, C.; Healey, C. S.; Kwok, J.

[42960] B. J.; Ponder, M. A.; Ponder, B. A. J.: Haplotype analysis of MEN
[42961] 2 mutations. Hum. Molec. Genet. 3: 1771-1774, 1994.
[42962] 6589.
[42963] Grieco, M.; Santoro, M.; Berlingieri, M. T.; Melillo, R. M.; Donghi,
[42964] R.; Bongarzone, I.; Pierotti, M. A.; Della Porta, G.; Fusco, A.; Vecchio,
[42965] G.: PTC is a novel rearranged form of the ret proto-oncogene and
[42966] is frequently detected in vivo in human thyroid papillary carcinomas. Cell 60:
[42967] 557-563, 1990.
[42968] 6590.
[42969] Hofstra, R. M. W.; Landsvater, R. M.; Ceccherini, I.; Stulp, R.
[42970] P.; Stelwagen, T.; Luo, Y.; Pasini, B.; Hoppener, J. W. M.; Ploos
[42971] van Amstel, H. K.; Romeo, G.; Lips, C. J. M.; Buys, C. H. C. M.:
[42972] A mutation in the RET proto-oncogene associated with multiple endocrine

[42973] neoplasia type 2B and sporadic medullary thyroid carcinoma. Nature 367:
[42974] 375-376, 1994.

[42975] 6591.

[42976] Hoppener, J. W. M.; Lips, C. J. M.: RET receptor tyrosine kinase
[42977] gene mutations: molecular biological, physiological and clinical aspects. Europ.
[42978] J. Clin. Invest. 26: 613-624, 1996.

[42979] 6592.

[42980] Hoppner, W.; Ritter, M. M.: A duplication of 12 bp in the critical
[42981] cysteine rich domain of the RET proto-oncogene results in a distinct
[42982] phenotype of multiple endocrine neoplasia type 2A. Hum. Molec. Genet. 6:
[42983] 587-590, 1997.

[42984] 6593.

[42985] Ikeda, I.; Ishizaka, Y.; Tahira, T.; Suzuki, T.; Onda, M.; Sugimura,
[42986] T.; Nagao, M.: Specific expression of the ret proto-

oncogene in human
[42987] neuroblastoma cell lines. Oncogene 5: 1291-1296, 1990.

[42988] 6594.

[42989] Ishizaka, Y.; Itoh, F.; Tahira, T.; Ikeda, I.; Sugimura, T.; Tucker,
[42990] J.; Fertitta, A.; Carrano, A. V.; Nagao, M.: Human ret proto-oncogene
[42991] mapped to chromosome 10q11.2. Oncogene 4: 1519-1521, 1989.
[42992] 6595.

[42993] Japon, M. A.; Urbano, A. G.; Saez, C.; Segura, D. I.; Cerro, A.
[42994] L.; Dieguez, C.; Alvarez, C. V.: Glial-derived neurotropic factor
[42995] and RET gene expression in normal human anterior pituitary cell types
[42996] and in pituitary tumors. J. Clin. Endocr. Metab. 87: 1879-1884,
[42997] 2002.

[42998] 6596.

[42999] Julies, M. G.; Moore, S. W.; Kotze, M. J.; du Plessis, L.:

Novel
[43000] RET mutations in Hirschsprung's disease patients from the diverse
[43001] South African population. Europ. J. Hum. Genet. 9: 419-423, 2001.
[43002] 6597.
[43003] Klugbauer, S.; Demidchik, E. P.; Lengfelder, E.; Rabes, H. M.
[43004] : Detection of a novel type of RET rearrangement (PTC5) in thyroid
[43005] carcinomas after Chernobyl and analysis of the involved RET-fused
[43006] gene RFG5. Cancer Res. 58: 198-203, 1998.
[43007] 6598.
[43008] Shapiro, J. R.; Stover, M. L.; Burn, V. E.; McKinstry, M. B.;
[43009] Burshell, A. L.; Chipman, S. D.; Rowe, D. W.: An osteopenic nonfracture
[43010] syndrome with features of mild osteogenesis imperfecta associated
[43011] with the substitution of a cysteine for glycine at triple he-lix position
[43012] 43 in the pro-alpha-1(I) chain of type I collagen. J. Clin.

Invest. 89:

[43013] 567-573, 1992.

[43014] 6599.

[43015] Steinmann, B.; Nicholls, A.; Pope, F. M.: Clinical variability

[43016] of osteogenesis imperfecta reflecting molecular hetero-
geneity: cysteine

[43017] substitutions in the alpha-1(I) collagen chain producing
lethal and

[43018] mild forms. J. Biol. Chem. 261: 8958-8964, 1986.

[43019] 6600.

[43020] de Oca Luna, R. M.; Tabor, A. D.; Eberspaecher, H.; Hul-
boy, D.

[43021] L.; Worth, L. L.; Colman, M. S.; Finlay, C. A.; Lozano, G.:
The organization

[43022] and expression of the mdm2 gene. Genomics 33:
352-357, 1996.

[43023] 6601.

[43024] Watson, C. E.; Draganov, D. I.; Billecke, S. S.; Bisgaier, C.

[43025] L.; La Du, B. N.: Rabbits possess a serum paraoxonase
polymorphism

[43026] similar to the human Q192R. Pharmacogenetics 11:

123-134, 2001.

[43027] 6602.

[43028] Mornet, E.: Hypophosphatasia: the mutations in the tissue-nonspecific

[43029] alkaline phosphatase gene. Hum. Mutat. 15: 309-315, 2000.

[43030] 6603.

[43031] Rajput, B.; Marshall, A.; Killary, A. M.; Lalley, P. A.; Naylor,

[43032] S. L.; Belin, D.; Rickles, R. J.; Strickland, S.: Chromosomal assignments

[43033] of genes for tissue plasminogen activator and urokinase in mouse. Somat.

[43034] Cell Molec. Genet. 13: 581-586, 1987.

[43035] 6604.

[43036] Giardiello, F. M.; Petersen, G. M.; Piantadosi, S.; Gruber, S.

[43037] B.; Traboulsi, E. I.; Offerhaus, G. J. A.; Muro, K.; Krush, A. J.;

[43038] Booker, S. V.; Luce, M. C.; Laken, S. J.; Kinzler, K. W.; Vo-gelstein,

[43039] B.; Hamilton, S. R.: APC gene mutations and extraintestinal phenotype

[43040] of familial adenomatous polyposis. Gut 40: 521-525, 1997.

[43041] 6605.

[43042] Yamasaki, Y.; Sakamoto, K.; Watada, H.; Kajimoto, Y.; Hori, M.

[43043] : The arg-192 isoform of paraoxonase with low sarin-hydrolyzing activity

[43044] is dominant in the Japanese. Hum. Genet. 101: 67-68, 1997.

[43045] 6606.

[43046] Mornet, E.; Taillandier, A.; Peyramaure, S.; Kaper, F.; Muller,

[43047] F.; Brenner, R.; Bussiere, P.; Freisinger, P.; Godard, J.; Le Merrer,

[43048] M.; Oury, J. F.; Plauchu, H.; Puddu, R.; Rival, J. M.; Superti-Furga,

[43049] A.; Touraine, R. L.; Serre, J. L.; Simon-Bouy, B.: Identification

[43050] of fifteen novel mutations in the tissue-nonspecific alkaline phosphatase

[43051] (TNSALP) gene in European patients with severe hypophosphatasia. Europ.

[43052] J. Hum. Genet. 6: 308-314, 1998.

[43053] 6607.

[43054] Orimo, H.; Hayashi, Z.; Watanabe, A.; Hirayama, T.; Hi-
rayama,
[43055] T.; Shimada, T.: Novel missense and frameshift mutations
in the tissue-nonspecific
[43056] alkaline phosphatase gene in a Japanese patient with hy-
pophosphatasia. Hum.
[43057] Molec. Genet. 3: 1683-1684, 1994.

[43058] 6608.

[43059] Ozono, K.; Yamagata, M.; Michigami, T.; Nakajima, S.;
Sakai, N.;
[43060] Cai, G.; Satomura, K.; Yasui, N.; Okada, S.; Nakayama, M.:
Identification
[43061] of novel missense mutations (phe310leu and gly439arg)
in a neonatal
[43062] case of hypophosphatasia. J. Clin. Endocr. Metab. 81:
4458-4461,
[43063] 1996.

[43064] 6609.

[43065] Sergi, C.; Mornet, E.; Troeger, J.; Voigtlaender, T.: Perinatal

[43066] hypophosphatasia: radiology, pathology and molecular bi-
ology studies
[43067] in a family harboring a splicing mutation (648+1A) and a
novel missense
[43068] mutation (N400S) in the tissue-nonspecific alkaline phos-
phatase (TNSALP)
[43069] gene. Am. J. Med. Genet. 103: 235-240, 2001.

[43070] 6610.

[43071] Smith, M.; Weiss, M. J.; Griffin, C. A.; Murray, J. C.; Buetow,

[43072] K. H.; Emanuel, B. S.; Henthorn, P. S.; Harris, H.: Regional
assignment
[43073] of the gene for human liver/bone/kidney alkaline phos-
phatase to chromosome
[43074] 1p36.1-p34. Genomics 2: 139-143, 1988.

[43075] 6611.

[43076] Swallow, D. M.; Povey, S.; Parkar, M.; Andrews, P. W.; Har-
ris,
[43077] H.; Pym, B.; Goodfellow, P.: Mapping of the gene coding
for the human
[43078] liver/bone/kidney isozyme of alkaline phosphatase to
chromosome 1. Ann.

[43079] Hum. Genet. 50: 229-235, 1986.

[43080] 6612.

[43081] Waymire, K. G.; Mahuren, J. D.; Jaje, J. M.; Guilarte, T. R.;

[43082] Coburn, S. P.; MacGregor, G. R.: Mice lacking tissue non-
specific

[43083] alkaline phosphatase die from seizures due to defective
metabolism

[43084] of vitamin B-6. Nature Genet. 11: 45-51, 1995.

[43085] 6613.

[43086] Weiss, M.; Smith, M.; Griffin, C.; Nussbaum, R.; Murray, J.;
Emanuel,

[43087] B.; Harris, H.: Assignment of the gene encoding the liver/
bone/kidney

[43088] form of alkaline phosphatase ALPL to the region
1p34-p36.1. (Abstract) Cytogenet.

[43089] Cell Genet. 46: 714 only, 1987.

[43090] 6614.

[43091] Weiss, M. J.; Cole, D. E. C.; Ray, K.; Whyte, M. P.; Lafferty,

[43092] M. A.; Mulivor, R. A.; Harris, H.: A missense mutation in
the human

[43093] liver/bone/kidney alkaline phosphatase gene causing a lethal form

[43094] of hypophosphatasia. Proc. Nat. Acad. Sci. 85: 7666-7669, 1988.

[43095] 6615.

[43096] Weiss, M. J.; Henthorn, P. S.; Lafferty, M. A.; Slaughter, C.;

[43097] Raducha, M.; Harris, H.: Isolation and characterization of a cDNA

[43098] encoding a human liver/bone/kidney-type alkaline phosphatase. Proc.

[43099] Nat. Acad. Sci. 83: 7182-7186, 1986.

[43100] 6616.

[43101] Weiss, M. J.; Ray, K.; Henthorn, P. S.; Lamb, B.; Kadesch, T.;

[43102] Harris, H.: Structure of the human liver/bone/kidney alkaline phosphatase

[43103] gene. J. Biol. Chem. 263: 12002-12010, 1988.

[43104] 6617.

[43105] Weiss, M. J.; Spielman, R. S.; Harris, H.: A high-frequency RFLP

[43106] at the human liver/bone/kidney-type alkaline phos-

phatase locus. Nucleic
[43107] Acids Res. 15: 860 only, 1987.

[43108] 6618.

[43109] Zurutuza, L.; Muller, F.; Gibrat, J. F.; Taillandier, A.; Si-
mon-Bouy,
[43110] B.; Serre, J. L.; Mornet, E.: Correlations of genotype and
phenotype
[43111] in hypophosphatasia. Hum. Molec. Genet. 8: 1039-1046,
1999.
[43112] 6619.

[43113] Whyte, M. P.; Mahuren, J. D.; Fedde, K. N.; Cole, F. S.; Mc-
Cabe,
[43114] E. R.; Coburn, S. P.: Perinatal hypophosphatasia: tissue
levels of
[43115] vitamin B-6 are unremarkable despite markedly increased
circulating
[43116] concentrations of pyridoxal-5-prime-phosphate: evidence
for an ectoenzyme
[43117] role for tissue-nonspecific alkaline phosphatase. J. Clin.
Invest. 81:
[43118] 1234-1239, 1988.

[43119] 6620.

[43120] Badger, K. S.; Sussman, H. H.: Structural evidence that human
[43121] liver and placental alkaline phosphatase isoenzymes are coded by different
[43122] genes. Proc. Nat. Acad. Sci. 73: 2201-2205, 1976.

[43123] 6621.

[43124] Beckman, L.; Beckman, G.; Christodoulou, C.; Ifekwunigwe, A.:
[43125] Variations in human placental alkaline phosphatase. Acta Genet. Statist.
[43126] Med. 17: 406-412, 1967.

[43127] 6622.

[43128] Beckman, L.; Bjorling, G.; Christodoulou, C.: Pregnancy enzymes
[43129] and placental polymorphism: alkaline phosphatase. Acta Genet. Statist.
[43130] Med. 16: 59-73, 1966.

[43131] 6623.

[43132] Boyer, S. H.: Alkaline phosphatase in human sera and placenta. Science 134:
[43133] 1002-1004, 1961.

[43134] 6624.

[43135] Giardiello, F. M.; Yang, V. W.; Hylind, L. M.; Krush, A. J.; Petersen,

[43136] G. M.; Trimbath, J. D.; Piantadosi, S.; Garrett, E.; Geiman, D. E.;

[43137] Hubbard, W.; Offerhaus, G. J. A.; Hamilton, S. R.: Primary chemoprevention

[43138] of familial adenomatous polyposis with sulindac. New Eng. J. Med. 346:

[43139] 1054-1059, 2002.

[43140] 6625.

[43141] Giovannucci, E.; Rimm, E. B.; Stampfer, M. J.; Colditz, G. A.;

[43142] Ascherio, A.; Willett, W. C.: Aspirin use and the risk for colorectal

[43143] cancer and adenoma in male health professionals. Ann. Intern. Med. 121:

[43144] 241-246, 1994.

[43145] 6626.

[43146] Gorlin, R. J.; Chaudhry, A. P.: Multiple osteomatosis, fibro-mas,

[43147] lipomas and fibrosarcomas of the skin and mesentery, epidermoid inclusion
[43148] cysts of the skin, leiomyomas and multiple intestinal polyposis: an
[43149] heritable disorder of connective tissue. New Eng. J. Med. 263: 1151-1158,
[43150] 1960.

[43151] 6627.

[43152] Goss, K. H.; Trzepacz, C.; Tuohy, T. M. F.; Groden, J.: Attenuated
[43153] APC alleles produce functional protein from internal translation initiation. Proc.
[43154] Nat. Acad. Sci. 99: 8161-8166, 2002.

[43155] 6628.

[43156] Greer, J. A., Jr.; Devine, K. D.; Dahlin, D. C.: Gardner's syndrome
[43157] and chondrosarcoma of the hyoid bone. Arch. Otolaryng. 103: 425-427,
[43158] 1977.

[43159] 6629.

[43160] Gruber, S. B.; Petersen, G. M.; Kinzler, K. W.; Vogelstein, B.

[43161] : Cancer, crash sites, and the new genetics of neoplasia. (Editorial) Gastroenterology 116:
[43162] 210-212, 1999.

[43163] 6630.

[43164] Gryfe, R.; Di Nicola, N.; Gallinger, S.; Redston, M.: Somatic

[43165] instability of the APC I1307K allele in colorectal neoplasia. Cancer
[43166] Res. 58: 4040-4043, 1998.

[43167] 6631.

[43168] Gryfe, R.; Di Nicola, N.; Lal, G.; Gallinger, S.; Redston, M.

[43169] : Inherited colorectal polyposis and cancer risk of the APC I1307K
[43170] polymorphism. Am. J. Hum. Genet. 64: 378-384, 1999.

[43171] 6632.

[43172] Haggitt, R. C.; Booth, J. L.: Bilateral fibromatosis of the breast
[43173] in Gardner's syndrome. Cancer 25: 161-166, 1970.

[43174] 6633.

[43175] Halberg, R. B.; Katzung, D. S.; Hoff, P. D.; Moser, A. R.;

Cole,
[43176] C. E.; Lubet, R. A.; Donehower, L. A.; Jacoby, R. F.; Dove, W. F.
[43177] : Tumorigenesis in the multiple intestinal neoplasia mouse: redundancy
[43178] of negative regulators and specificity of modifiers. Proc. Nat. Acad.
[43179] Sci. 97: 3461-3466, 2000.

[43180] 6634.

[43181] Halling, K. C.; Lazzaro, C. R.; Honchel, R.; Bufill, J. A.; Powell,
[43182] S. M.; Arndt, C. A. S.; Lindor, N. M.: Hereditary desmoid disease
[43183] in a family with a germline AluI repeat mutation of the APC gene. Hum.
[43184] Hered. 49: 97-102, 1999.

[43185] 6635.

[43186] Hampton, G. M.; Ward, J. R. T. J.; Cottrell, S.; Howe, K.; Thomas,
[43187] H. J. W.; Ballhausen, W. G.; Jones, T.; Sheer, D.; Solomon, E.; Frischauf,
[43188] A.-M.; Bodmer, W. F.: Yeast artificial chromosomes for the

molecular
[43189] analysis of the familial polyposis APC gene region. Proc. Nat. Acad.
[43190] Sci. 89: 8249-8253, 1992.

[43191]

[43192] 100. Hardwick, J. C. H.; van den Brink, G. R.; Offerhaus, G. J.; van
[43193] Deventer, S. J. H.; Peppelenbosch, M. P.: NF-kappaB, p38 MAPK and
[43194] JNK are highly expressed and active in the stroma of human colonic
[43195] adenomatous polyps. Oncogene 20: 819-827, 2001.

[43196]

[43197] 101. Harned, R. K.; Williams, S. M.: Familial polyposis coli and
[43198] periampullary malignancy. Dis. Colon Rectum 25: 227-229, 1982.

[43199]

[43200] 102. He, T.-C.; Sparks, A. B.; Rago, C.; Hermeking, H.; Zawel, L.;
[43201] da Costa, L. T.; Morin, P. J.; Vogelstein, B.; Kinzler, K. W.:

Identification

[43202] of c-MYC as a target of the APC pathway. Science 281: 1509-1512,

[43203] 1998.

[43204]

[43205] 103. Heimann, A.; White, P. F.; Riely, C. A.; Ritchey, A. K.; Flye,

[43206] M. W.; Barwick, K. W.: Hepatoblastoma presenting as iso-sexual precocity:

[43207] the clinical importance of histologic and serologic parameters. J.

[43208] Clin. Gastroent. 9: 105-110, 1987.

[43209]

[43210] 104. Herrera, L.; Kakati, S.; Gibas, L.; Pietrzak, E.; Sandberg, A.

[43211] A.: Gardner syndrome in a man with an interstitial deletion of 5q. Am.

[43212] J. Med. Genet. 25: 473-476, 1986.

[43213]

[43214] 105. Herrmann, C.; Block, C.; Geisen, C.; Haas, K.; Weber, C.; Winde,

[43215] G.; Moroy, T.; Muller, O.: Sulindac sulfide inhibits Ras signaling. Oncogene 17:
[43216] 1769-1776, 1998.

[43217]

[43218] 106. Herve, R.; Farret, O.; Mayaudon, H.; Helie, C.; Denee, J. M.;
[43219] Bauduceau, B.; Molinie, C.: Association syndrome de Gardner et carcinome
[43220] thyroidien. (Letter) Presse Med. 24: 415 only, 1995.

[43221]

[43222] 107. Hockey, K. A.; Mulcahy, M. T.; Montgomery, P.; Levitt, S.: Deletion
[43223] of chromosome 5q and familial adenomatous polyposis. J. Med. Genet. 26:
[43224] 61-62, 1989.

[43225]

[43226] 108. Hodgson, S. V.; Coonar, A. S.; Hanson, P. J. V.; Cottrell, S.;
[43227] Scriven, P. N.; Jones, T.; Hawley, P. R.; Wilkinson, M. L.: Two cases
[43228] of 5q deletions in patients with familial adenomatous

polyposis: possible
[43229] link with Caroli's disease. J. Med. Genet. 30: 369-375, 1993.
[43230]

[43231] 109. Hodgson, S. V.; Fagg, N. L. K.; Talbot, I. C.; Wilkinson, M.
[43232] : Deletions of the entire APC gene are associated with sessile colonic
[43233] adenomas. (Letter) J. Med. Genet. 31: 426 only, 1994.
[43234]

[43235] 110. Hoffmann, D. C.; Brooke, B. N.: Familial sarcoma of bone in
[43236] a polyposis coli family. Dis. Colon Rectum 13: 119-120, 1970.
[43237]

[43238] 111. Homfray, T. F. R.; Cottrell, S. E.; Ilyas, M.; Rowan, A.; Talbot,
[43239] I. C.; Bodmer, W. F.; Tomlinson, I. P. M.: Defects in mismatch repair
[43240] occur after APC mutations in the pathogenesis of sporadic colorectal
[43241] tumours. Hum. Mutat. 11: 114-120, 1998.

[43242]

[43243] 112. Horii, A.; Nakatsuru, S.; Ichii, S.; Nagase, H.; Nakamura, Y.

[43244] : Multiple forms of the APC gene transcripts and their tissue-specific

[43245] expression. Hum. Molec. Genet. 2: 283-287, 1993.

[43246]

[43247] 113. Horii, A.; Nakatsuru, S.; Miyoshi, Y.; Ichii, S.; Nagase, H.;

[43248] Kato, Y.; Yanagisawa, A.; Nakamura, Y.: The APC gene, responsible

[43249] for familial adenomatous polyposis, is mutated in human gastric cancer. Cancer

[43250] Res. 52: 3231-3233, 1992.

[43251]

[43252] 114. Houlston, R.; Crabtree, M.; Phillips, R.; Crabtree, M.; Tomlinson,

[43253] I.: Explaining differences in the severity of familial adenomatous

[43254] polyposis and the search for modifier genes. Gut 48: 1-5, 2001.

[43255]

[43256] 115. Houlston, R. S.; Fallon, T.; Harocopos, C.; Williams, C. B.;

[43257] Davey, C.; Slack, J.: Congenital hypertrophy of retinal pigment epithelium

[43258] in patients with colonic polyps associated with cancer family syndrome. Clin.

[43259] Genet. 42: 16-18, 1992.

[43260]

[43261] 116. Hsu, S. H.; Luk, G. D.; Krush, A. J.; Hamilton, S. R.; Hoover,

[43262] H. H., Jr.: Multiclonal origin of polyps in Gardner syndrome. Science 221:

[43263] 951-953, 1983.

[43264]

[43265] 117. Huang, H.; Mahler-Araujo, B. M.; Sankila, A.; Chimelli, L.; Yonekawa,

[43266] Y.; Kleihues, P.; Ohgaki, H.: APC mutations in sporadic medulloblastomas. Am.

[43267] J. Path. 156: 433-437, 2000.

[43268]

[43269] 118. Hughes, L. J.; Michels, V. V.: Risk of hepatoblastoma in familial
[43270] adenomatous polyposis. Am. J. Med. Genet. 43: 1023-1025, 1992.
[43271]

[43272] 119. Hyson, E. A.; Burrell, M.: Familial gastric polyposis (cont.)
[43273] (Letter) New Eng. J. Med. 295: 905 only, 1976.
[43274]

[43275] 120. Iida, M.; Yao, T.; Itoh, H.; Watanabe, H.; Matsui, T.; Iwashita,
[43276] A.; Fujishima, M.: Natural history of gastric adenomas in patients
[43277] with familial adenomatosis coli/Gardner's syndrome. Cancer 61: 605-611,
[43278] 1988.
[43279]

[43280] 121. Ingram, J. T.; Oldfield, M. C.: Hereditary sebaceous cysts. Brit.
[43281] Med. J. 1: 960-963, 1937.
[43282]

[43283] 122. Jagelman, D. G.; DeCosse, J. J.; Bussey, H. J. R.; the Leeds
[43284] Castle Polyposis Group: Upper gastrointestinal cancer in familial
[43285] adenomatous polyposis. Lancet I: 1149–1151, 1988.

[43286]

[43287] 123. Jarvinen, H. J.; Peltokallio, P.; Landtman, M.; Wolf, J.: Gardner's
[43288] stigmas in patients with familial adenomatosis coli. Brit. J. Surg. 69:
[43289] 718–721, 1982.

[43290]

[43291] 124. Jones, T. R.; Nance, F. C.: Periampullary malignancy in Gardner's
[43292] syndrome. Ann. Surg. 185: 565–573, 1977.

[43293]

[43294] 125. Joslyn, G.; Carlson, M.; Thliveris, A.; Albertsen, H.; Gelbert,
[43295] L.; Samowitz, W.; Groden, J.; Stevens, J.; Spirio, L.; Robert-
son,
[43296] M.; Sargeant, L.; Krapcho, K.; Wolff, E.; Burt, R.; Hughes, J.

P.;

[43297] Warrington, J.; McPherson, J.; Wasmuth, J.; Le Paslier, D.; Abderrahim,

[43298] H.; Cohen, D.; Leppert, M.; White, R.: Identification of deletion

[43299] mutations and three new genes at the familial polyposis locus. Cell 66:

[43300] 601-613, 1991.

[43301]

[43302] 126. Kaplan, J.; Cushing, B.; Chang, C.-H.; Poland, R.; Roscamp, J.;

[43303] Perrin, E.; Bhaya, N.: Familial T-cell lymphoblastic lym-

phoma: association

[43304] with von Recklinghausen neurofibromatosis and Gardner syndrome. Am.

[43305] J. Hemat. 12: 247-250, 1982.

[43306]

[43307] 127. Kaplan, K. B.; Burds, A. A.; Swedlow, J. R.; Bekir, S. S.; Sorger,

[43308] P. K.; Nathke, I. S.: A role for the adenomatous polyposis coli protein

[43309] in chromosome segregation. Nature Cell Biol. 3: 492-432,

2001.

[43310]

[43311] 128. Kartheuser, A.; Walon, C.; West, S.; Breukel, C.; Detry, R.;

[43312] Gribomont, A.-C.; Hamzehloei, T.; Hoang, P.; Maiter, D.; Pringot,

[43313] J.; Rahier, J.; Khan, P. M.; Curtis, A.; Burn, J.; Fodde, R.; Verellen-Dumoulin,

[43314] C.: Familial adenomatous polyposis associated with multi-

ple adrenal

[43315] adenomas in a patient with a rare 3-prime APC mutation. J. Med. Genet. 36:

[43316] 65-67, 1999.

[43317]

[43318] 129. Kassirer, J. P.: What role for nurse practitioner in pri-

mary

[43319] care? (Editorial) New Eng. J. Med. 330: 204-205, 1994.

[43320]

[43321] 130. Kasukawa, T.; Watanabe, T.; Endo, A.: Absence of heteromorphism

[43322] of chromosome number 2 homologues in patients with

hereditary adenomatosis
[43323] of the colon and rectum. Cancer Genet. Cytogenet. 9:
283-295, 1983.
[43324]

[43325] 131. Kawasaki, Y.; Senda, T.; Ishidata, T.; Koyama, R.;
Morishita,
[43326] T.; Iwayama, Y.; Higuchi, O.; Akiyama, T.: Asef, a link be-
tween the
[43327] tumor suppressor APC and G-protein signaling. Science
289: 1194-1197,
[43328] 2000.

[43329]

[43330] 132. Kelly, P. B.; McKinnon, D. A.: Familial multiple poly-
posis of
[43331] the colon: review and description of a large kindred.
McGill Med.
[43332] J. 30: 67-85, 1961.

[43333]

[43334] 133. Kingston, J. E.; Draper, G. J.; Mann, J. R.: Hepatoblas-
toma
[43335] and polyposis coli. (Letter) Lancet I: 457 only, 1982.

[43336]

[43337] 134. Kinzler, K. W.: Personal Communication. Baltimore, Md. 9/30/1991.

[43338]

[43339] 135. Kinzler, K. W.; Nilbert, M. C.; Su, L.-K.; Vogelstein, B.; Bryan,

[43340] T. M.; Levy, D. B.; Smith, K. J.; Preisinger, A. C.; Hedge, P.; McKechnie,

[43341] D.; Finniear, R.; Markham, A.; Groffen, J.; Boguski, M. S.; Altschul,

[43342] S. F.; Horii, A.; Ando, H.; Miyoshi, Y.; Miki, Y.; Nishisho, I.; Nakamura,

[43343] Y.: Identification of FAP locus genes from chromosome 5q21. Science 253:

[43344] 661-665, 1991.

[43345]

[43346] 136. Kinzler, K. W.; Vogelstein, B.: Lessons from hereditary colorectal

[43347] cancer. Cell 87: 159-170, 1996.

[43348]

[43349] 137. Klemmer, S.; Pascoe, L.; DeCosse, J.: Occurrence of

desmoids
[43350] in patients with familial adenomatous polyposis of the colon. Am.
[43351] J. Med. Genet. 28: 385-392, 1987.

[43352]

[43353] 138. Kobayashi, T.; Narahara, K.; Yokoyama, Y.; Ueyama, S.; Mohri,
[43354] O.; Fujii, T.; Fujimoto, M.; Ohtsuki, S.; Tsuji, K.; Seino, Y.: Gardner
[43355] syndrome in a boy with interstitial deletion of the long arm of chromosome
[43356] 5. Am. J. Med. Genet. 41: 460-463, 1991.

[43357]

[43358] 139. Krush, A. J.: Personal Communication. Baltimore, Md. 4/1/1988.

[43359]

[43360] 140. Krush, A. J.; Traboulsi, E. I.; Offerhaus, G. J. A.; Maumenee,
[43361] I. H.; Yardley, J. H.; Levin, L. S.: Hepatoblastoma, pigmented ocular
[43362] fundus lesions and jaw lesions in Gardner syndrome. Am.

J. Med. Genet. 29:
[43363] 323-332, 1988.

[43364]

[43365] 141. Laken, S. J.; Papadopoulos, N.; Petersen, G. M.; Gru-
ber, S. B.;
[43366] Hamilton, S. R.; Giardiello, F. M.; Brensinger, J. D.; Vogel-
stein,
[43367] B.; Kinzler, K. W.: Analysis of masked mutations in familial
adenomatous
[43368] polyposis. Proc. Nat. Acad. Sci. 96: 2322-2326, 1999.

[43369]

[43370] 142. Laken, S. J.; Petersen, G. M.; Gruber, S. B.; Oddoux,
C.; Ostrer,
[43371] H.; Giardiello, F. M.; Hamilton, S. R.; Hampel, H.;
Markowitz, A.;
[43372] Klimstra, D.; Jhanwar, S.; Winawer, S.; Offit, K.; Luce, M.
C.; Kinzler,
[43373] K. W.; Vogelstein, B.: Familial colorectal cancer in Ashke-
nazim due
[43374] to a hypermutable tract in APC. Nature Genet. 17: 79-83,
1997.

[43375]

[43376] 143. Lal, G.; Ash, C.; Hay, K.; Redston, M.; Kwong, E.; Hancock, B.;
[43377] Mak, T.; Kargman, S.; Evans, J. F.; Gallinger, S.: Suppression of
[43378] intestinal polyps in Msh2-deficient and non-Msh2-deficient multiple
[43379] intestinal neoplasia mice by a specific cyclooxygenase-2 inhibitor
[43380] and by a dual cyclooxygenase-1/2 inhibitor. Cancer Res. 61: 6131-6136,
[43381] 2001.

[43382]

[43383] 144. Lambertz, S.; Ballhausen, W. G.: Identification of an alternative
[43384] 5-prime untranslated region of the adenomatous polyposis coli gene. Hum.
[43385] Genet. 90: 650-652, 1993.

[43386]

[43387] 145. Lamlum, H.; Al Tassan, N.; Jaeger, E.; Frayling, I.; Sieber,
[43388] O.; Bin Reza, F.; Eckert, M.; Rowan, A.; Barclay, E.; Atkin, W.; Williams,

[43389] C.; Gilbert, J.; Cheadle, J.; Bell, J.; Houlston, R.; Bodmer, W.;

[43390] Sampson, J.; Tomlinson, I.: Germline APC variants in patients with

[43391] multiple colorectal adenomas, with evidence for the particular importance

[43392] of E1317Q. Hum. Molec. Genet. 9: 2215-2221, 2000.

[43393]

[43394] 146. Lamlum, H.; Ilyas, M.; Rowan, A.; Clark, S.; Johnson, V.; Bell,

[43395] J.; Frayling, I.; Efstathiou, J.; Pack, K.; Payne, S.; Roylance, R.;

[43396] Gorman, P.; Sheer, D.; Neale, K.; Phillips, R.; Talbot, I.; Bodmer,

[43397] W.; Tomlinson, I.: The type of somatic mutation at APC in familial

[43398] adenomatous polyposis is determined by the site of the germline mutation:

[43399] a new facet of Knudson's 'two-hit' hypothesis. Nature Med. 5: 1071-1075,

[43400] 1999.

[43401]

[43402] 147. Lander, E.: Personal Communication. Cambridge, Mass. 10/1991.

[43403]

[43404] 148. Leffall, L. D.; Chung, E. B.; Dewitty, R. L.; Cornwell, E. E.;

[43405] Blakey, T. M.: Familial polyposis coli in Black patients. Ann. Surg. 186:

[43406] 324-333, 1977.

[43407]

[43408] 149. Leppard, B.; Bussey, H. J. R.: Epidermoid cysts, poly-posis coli

[43409] and Gardner's syndrome. Brit. J. Surg. 62: 387-393, 1975.

[43410]

[43411] 150. Leppert, M.; Burt, R.; Hughes, J. P.; Samowitz, W.; Nakamura,

[43412] Y.; Woodward, S.; Gardner, E.; Lalouel, J.-M.; White, R.: Genetic

[43413] analysis of an inherited predisposition to colon cancer in a family

[43414] with a variable number of adenomatous polyps. New Eng. J. Med. 322:

[43415] 904-908, 1990.

[43416]

[43417] 151. Leppert, M.; Dobbs, M.; Scambler, P.; O'Connell, P.; Nakamura,

[43418] Y.; Stauffer, D.; Woodward, S.; Burt, R.; Hughes, J.; Gardner, E.;

[43419] Lathrop, M.; Wasmuth, J.; Lalouel, J.-M.; White, R.: The gene for

[43420] familial polyposis coli maps to the long arm of chromosome 5. Science 238:

[43421] 1411-1413, 1987.

[43422]

[43423] 152. Leppert, M.; Dobbs, M.; Scambler, P.; O'Connell, P.; Nakamura,

[43424] Y.; Stauffer, D.; Woodward, S.; Burt, R.; Hughes, J. P.; Gardner,

[43425] E.; Lathrop, M.; Wasmuth, J.; Lalouel, J. M.; White, R.: The gene

[43426] for familial polyposis maps to the long arm of chromosome 5. (Abstract) Cytogenet.

[43427] Cell Genet. 46: 647 only, 1987.

[43428]

[43429] 153. Lewis, R. A.; Crowder, W. E.; Eierman, L. A.; Nuss-
baum, R. L.;
[43430] Ferrell, R. E.: The Gardner syndrome: significance of ocu-
lar features. Ophthalmology 91:
[43431] 916-925, 1984.

[43432]

[43433] 154. Lewis, R. J.; Mitchell, J. C.: Basal cell carcinoma in
Gardner's
[43434] syndrome. Acta Derm. Venerol. 51: 67-68, 1971.

[43435]

[43436] 155. Li, F. P.; Thurber, W. A.; Seddon, J.; Holmes, G. E.:
Hepatoblastoma
[43437] in families with polyposis coli. J.A.M.A. 257: 2475-2477,
1987.

[43438]

[43439] 156. Lindgren, V.; Bryke, C. R.; Ozcelik, T.; Yang-Feng, T.
L.; Francke,
[43440] U.: Phenotypic, cytogenetic, and molecular studies of
three patients
[43441] with constitutional deletions of chromosome 5 in the re-

gion of the
[43442] gene for familial adenomatous polyposis. Am. J. Hum. Genet. 50:
[43443] 988-997, 1992.

[43444]

[43445] 157. Luk, G. D.; Baylin, S. B.: Ornithine decarboxylase as a biologic
[43446] marker in familial colonic polyposis. New Eng. J. Med. 311: 80-83,
[43447] 1984.

[43448]

[43449] 158. Luongo, C.; Gould, K. A.; Su, L.-K.; Kinzler, K. W.; Vogelstein,
[43450] B.; Dietrich, W.; Lander, E. S.; Moser, A. R.: Mapping of multiple
[43451] intestinal neoplasia (Min) to proximal chromosome 18 of the mouse. Genomics 15:
[43452] 3-8, 1993.

[43453]

[43454] 159. Lynch, H. T.: Desmoid tumors: genotype-phenotype differences

[43455] in familial adenomatous polyposis: a nosological dilemma. (Editorial) Am.
[43456] J. Hum. Genet. 59: 1184-1185, 1996.

[43457]

[43458] 160. Lynch, H. T.; Smyrk, T. C.: Classification of familial adenomatous
[43459] polyposis: a diagnostic nightmare. (Editorial) Am. J. Hum. Genet. 62:
[43460] 1288-1289, 1998.

[43461]

[43462] 161. Lynch, H. T.; Smyrk, T. C.; Watson, P.; Lanspa, S. J.; Lynch,
[43463] P. M.; Jenkins, J. X.; Rouse, J.; Cavalieri, J.; Howard, L.; Lynch,
[43464] J.: Hereditary flat adenoma syndrome: a variant of familial adenomatous
[43465] polyposis? Dis. Colon Rectum 35: 411-421, 1992.

[43466]

[43467] 162. Lyons, L. A.; Lewis, R. A.; Strong, L. C.; Zuckerbrod, S.; Ferrell,
[43468] R. E.: A genetic study of Gardner syndrome and congenital

hypertrophy
[43469] of the retinal pigment epithelium. Am. J. Hum. Genet. 42: 290-296,
[43470] 1988.

[43471]

[43472] 163. MacDonald, J. M.; Davis, W. C.; Crago, H. R.; Berk, A. D.: Gardner's
[43473] syndrome and periampullary malignancy. Am. J. Surg. 113: 425-430,
[43474] 1967.

[43475]

[43476] 164. Maher, E. R.; Barton, D. E.; Slatter, R.; Koch, D. J.; Jones,
[43477] M. H.; Nagase, H.; Payne, S. J.; Charles, S. J.; Moore, A. T.; Nakamura,
[43478] Y.; Ferguson-Smith, M. A.: Evaluation of molecular genetic diagnosis
[43479] in the management of familial adenomatous polyposis coli: a population
[43480] based study. J. Med. Genet. 30: 675-678, 1993.

[43481]

[43482] 165. Maher, E. R.; Morson, B.; Beach, R.; Hodgson, S. V.: Phenotypic
[43483] variation in hereditary nonpolyposis colon cancer syndrome: association
[43484] with infiltrative fibromatosis (desmoid tumor). Cancer 69: 2049-2051,
[43485] 1992.

[43486]

[43487] 166. Mahmoud, N. N.; Bilinski, R. T.; Churchill, M. R.; Edelmann,
[43488] W.; Kucherlapati, R.; Bertagnolli, M. M.: Genotype-phenotype correlation
[43489] in murine Apc mutation: differences in enterocyte migration and response
[43490] to sulindac. Cancer Res. 59: 353-359, 1999.

[43491]

[43492] 167. Mahmoud, N. N.; Boolbol, S. K.; Bilinski, R. T.; Martucci, C.
[43493] M.; Chadburn, A.; Bertagnolli, M. M.: Apc gene mutation is associated
[43494] with a dominant-negative effect upon intestinal cell migration. Cancer

[43495] Res. 57: 5045-5050, 1997.

[43496]

[43497] 168. Marcus, A. J.: Aspirin as prophylaxis against colorec-
tal cancer. New.
[43498] Eng. J. Med. 333: 656-658, 1995.

[43499]

[43500] 169. Marshall, B.; Isidro, G.; Carvalhas, R.; Boavida, M. G.:
Germline
[43501] versus somatic mutations of the APC gene: evidence for
mechanistic
[43502] differences. (Letter) Hum. Mutat. 9: 286-288, 1997.

[43503]

[43504] 170. Marshall, W. H.; Martin, F. I. R.; MacKay, I. R.: Gard-
ner's
[43505] syndrome with adrenal carcinoma. Aust. Ann. Med. 16:
242-244, 1967.

[43506]

[43507] 171. Martin-Denavit, T.; Duthel, S.; Giraud, S.; Olschwang,
S.; Saurin,
[43508] J. C.; Plauchu, H.: Phenotype variability of two FAP families
with

[43509] an identical APC germline mutation at codon 1465: a po-
tential modifier
[43510] effect? Clin. Genet. 60: 125-131, 2001.

[43511]

[43512] 172. Matsumine, A.; Ogai, A.; Senda, T.; Okumura, N.;
Satoh, K.; Baeg,
[43513] G.-H.; Kawahara, T.; Kobayashi, S.; Okada, M.; Toyoshima,
K.; Akiyama,
[43514] T.: Binding of APC to the human homolog of the
Drosophila discs large
[43515] tumor suppressor protein. Science 272: 1020-1023,
1996.

[43516]

[43517] 173. Matsumoto, T.; Iida, M.; Mizuno, M.; Nakamura, S.;
Hizawa, K.;
[43518] Yao, T.: Serrated adenoma in familial adenomatous poly-
posis: relation
[43519] to germline APC gene mutation. Gut 50: 402-404, 2002.

[43520]

[43521] 174. Maule, W. F.: Screening for colorectal cancer by nurse
endoscopists. New

[43522] Eng. J. Med. 330: 183-187, 1994.

[43523]

[43524] 175. McKusick, V. A.: Genetic factors in intestinal polypo-
sis. J.A.M.A. 182:
[43525] 271-277, 1962.

[43526]

[43527] 176. Meera Khan, P.; Tops, C. M. J.; van den Broek, M.;
Breukel, C.;
[43528] Wijnen, J. T.; Oldenburg, M.; v.d. Bos, J.; van Leeuwen-
Cornelisse,
[43529] I. S. J.; Vasen, H. F. A.; Griffioen, G.; Verspaget, H. M.; den
Hartog
[43530] Jager, F. C. A.; Lamers, C. B. H. W.: Close linkage of a
highly polymorphic
[43531] marker (D5S37) to familial adenomatous polyposis (FAP)
and confirmation
[43532] of FAP localization on chromosome 5q21-q22. Hum.
Genet. 79: 183-185,
[43533] 1988.

[43534]

[43535] 177. Middleton, S. B.; Frayling, I. M.; Phillips, R. K. S.:

Desmoids
[43536] in familial adenomatous polyposis are monoclonal prolif-
erations. Brit.
[43537] J. Cancer 82: 827-832, 2000.

[43538]

[43539] 178. Midgley, C. A.; White, S.; Howitt, R.; Save, V.; Dunlop,
M. G.;
[43540] Hall, P. A.; Lane, D. P.; Wyllie, A. H.; Bubb, V. J.: APC ex-
pression
[43541] in normal human tissues. J. Path. 181: 426-433, 1997.

[43542]

[43543] 179. Miyashiro, I.; Senda, T.; Matsumine, A.; Baeg, G.-H.;
Kuroda,
[43544] T.; Shimano, T.; Miura, S.; Noda, T.; Kobayashi, S.; Mon-
den, M.; Toyoshima,
[43545] K.; Akiyama, T.: Subcellular localization of the APC pro-
tein: immunoelectron
[43546] microscopic study of the association of the APC protein
with catenin. Oncogene 11:
[43547] 89-96, 1995.

[43548]

[43549] 180. Miyoshi, Y.; Ando, H.; Nagase, H.; Nishisho, I.; Horii, A.; Miki,
[43550] Y.; Mori, T.; Utsunomiya, J.; Baba, S.; Petersen, G.; Hamilton, S.
[43551] R.; Kinzler, K. W.; Vogelstein, B.; Nakamura, Y.: Germ-line mutations
[43552] of the APC gene in 53 familial adenomatous polyposis patients. Proc.
[43553] Nat. Acad. Sci. 89: 4452-4456, 1992.

[43554]

[43555] 181. Miyoshi, Y.; Nagase, H.; Ando, H.; Horii, A.; Ichii, S.; Nakatsuru,
[43556] S.; Aoki, T.; Miki, Y.; Mori, T.; Nakamura, Y.: Somatic mutations
[43557] of the APC gene in colorectal tumors: mutation cluster region in the
[43558] APC gene. Hum. Molec. Genet. 1: 229-233, 1992.

[43559]

[43560] 182. Morton, D. G.; Macdonald, F.; Cachon-Gonzales, M. B.; Rindl,
[43561] P. M.; Neoptolemos, J. P.; Keighley, M. R. B.; Delhanty, J. D. A.;

[43562] McKeown, C. M.; Kilpatrick, M.; Hulten, M. A.: The use of DNA from
[43563] paraffin wax preserved tissue for predictive diagnosis in familial
[43564] adenomatous polyposis. J. Med. Genet. 29: 571-573, 1992.
[43565]

[43566] 183. Moser, A. R.; Mattes, E. M.; Dove, W. F.; Lindstrom, M. J.; Haag,
[43567] J. D.; Gould, M. N.: Apc(Min), a mutation in the murine Apc gene,
[43568] predisposes to mammary carcinomas and focal alveolar hyperplasias. Proc.
[43569] Nat. Acad. Sci. 90: 8977-8981, 1993.
[43570]

[43571] 184. Murphy, E. A.; Krush, A. J.; Dietz, M.; Rohde, C. A.: Hereditary
[43572] polyposis coli. III. Genetic and evolutionary fitness. Am. J. Hum.
[43573] Genet. 32: 700-713, 1980.
[43574]

[43575] 185. Nagase, H.; Miyoshi, Y.; Horii, A.; Aoki, T.; Ogawa, M.; Utsunomiya,
[43576] J.; Baba, S.; Sasazuki, T.; Nakamura, Y.: Correlation between the
[43577] location of germ-line mutations in the APC gene and the number of
[43578] colorectal polyps in familial adenomatous polyposis patients. Cancer
[43579] Res. 52: 4055-4057, 1992.

[43580]

[43581] 186. Nagase, H.; Miyoshi, Y.; Horii, A.; Aoki, T.; Petersen, G. M.;
[43582] Vogelstein, B.; Maher, E.; Ogawa, M.; Maruyama, M.; Utsunomiya, J.;
[43583] Baba, S.; Nakamura, Y.: Screening for germ-line mutations in familial
[43584] adenomatous polyposis patients: 61 new patients and a summary of 150
[43585] unrelated patients. Hum. Mutat. 1: 467-473, 1992.

[43586]

[43587] 187. Nagase, H.; Nakamura, Y.: Mutations of the APC (adenomatous

[43588] polyposis coli) gene. Hum. Mutat. 2: 425-434, 1993.

[43589]

[43590] 188. Nakagawa, H.; Murata, Y.; Koyama, K.; Fujiyama, A.; Miyoshi,

[43591] Y.; Monden, M.; Akiyama, T.; Nakamura, Y.: Identification of a brain-specific

[43592] APC homologue, APCL, and its interaction with beta-catenin. Cancer

[43593] Res. 58: 5176-5181, 1998.

[43594]

[43595] 189. Nakamura, Y.; Lathrop, M.; Leppert, M.; Dobbs, M.; Wasmuth, J.;

[43596] Wolff, E.; Carlson, M.; Fujimoto, E.; Krapcho, K.; Sears, T.; Woodward,

[43597] S.; Hughes, J.; Burt, R.; Gardner, E.; Lalouel, J.-M.; White, R.:

[43598] Localization of the genetic defect in familial adenomatous polyposis

[43599] within a small region of chromosome 5. Am. J. Hum. Genet. 43: 638-644,

[43600] 1988.

[43601]

[43602] 190. Naylor, E. W.; Gardner, E. J.: Penetrance and expres-
sivity of
[43603] the gene responsible for the Gardner syndrome. Clin.
Genet. 11:
[43604] 381-393, 1977.

[43605]

[43606] 191. Naylor, E. W.; Gardner, E. J.: Adrenal adenomas in a
patient
[43607] with Gardner's syndrome. Clin. Genet. 20: 67-73, 1981.

[43608]

[43609] 192. Naylor, E. W.; Lebenthal, E.: Gardner's syndrome: re-
cent developments
[43610] in research and management. Digest. Dis. Sci. 25:
945-959, 1980.

[43611]

[43612] 193. Neufeld, K. L.; Nix, D. A.; Bogerd, H.; Kang, Y.; Beck-
erle, M.
[43613] C.; Cullen, B. R.; White, R. L.: Adenomatous polyposis coli
protein
[43614] contains two nuclear export signals and shuttles between

the nucleus
[43615] and cytoplasm. Proc. Nat. Acad. Sci. 97: 12085-12090, 2000.
[43616]

[43617] 194. Neufeld, K. L.; White, R. L.: Nuclear and cytoplasmic localizations
[43618] of the adenomatous polyposis coli protein. Proc. Nat. Acad. Sci. 94:
[43619] 3034-3039, 1997.

[43620]

[43621] 195. Nishisho, I.; Nakamura, Y.; Miyoshi, Y.; Miki, Y.; Ando, H.;
[43622] Horii, A.; Koyama, K.; Utsunomiya, J.; Baba, S.; Hedge, P.; Markham,
[43623] A.; Krush, A. J.; Petersen, G.; Hamilton, S. R.; Nilbert, M. C.; Levy,
[43624] D. B.; Bryan, T. M.; Preisinger, A. C.; Smith, K. J.; Su, L.-K.; Kinzler,
[43625] K. W.; Vogelstein, B.: Mutations of chromosome 5q21 genes in FAP
[43626] and colorectal cancer patients. Science 253: 665-669, 1991.

[43627]

[43628] 196. Novelli, M. R.; Williamson, J. A.; Tomlinson, I. P. M.; Elia,

[43629] G.; Hodgson, S. V.; Talbot, I. C.; Bodmer, W. F.; Wright, N. A.:

[43630] Polyclonal origin of colonic adenomas in an XO/XY patient with FAP. Science 272:

[43631] 1187-1190, 1996.

[43632]

[43633] 197. Oda, H.; Imai, Y.; Nakatsuru, Y.; Hata, J.; Ishikawa, T.: Somatic

[43634] mutations of the APC gene in sporadic hepatoblastomas. Cancer Res. 56:

[43635] 3320-3323, 1996.

[43636]

[43637] 198. Offerhaus, G. J. A.; Giardiello, F. M.; Krush, A. J.; Booker,

[43638] S. V.; Tersmette, A. C.; Kelley, N. C.; Hamilton, S. R.: The risk

[43639] of upper gastrointestinal cancer in familial adenomatous polyposis. Gastroenterology 102:

[43640] 1980-1982, 1992.

[43641]

[43642] 199. Okamoto, M.; Sato, C.; Kohno, Y.; Mori, T.; Iwama, T.; Tonomura,

[43643] A.; Miki, Y.; Utsunomiya, J.; Nakamura, Y.; White, R.; Miyaki, M.

[43644] : Molecular nature of chromosome 5q loss in colorectal tumors and

[43645] desmoids from patients with familial adenomatous poly-posis. Hum.

[43646] Genet. 85: 595-599, 1990.

[43647]

[43648] 200. Olschwang, S.; Tiret, A.; Laurent-Puig, P.; Muleris, M.; Parc,

[43649] R.; Thomas, G.: Restriction of ocular fundus lesions to a specific

[43650] subgroup of APC mutations in adenomatous polyposis coli patients. Cell 75:

[43651] 959-968, 1993.

[43652]

[43653] 201. Oshima, M.; Dinchuk, J. E.; Kargman, S. L.; Oshima,

H.; Hancock,

[43654] B.; Kwong, E.; Trzaskos, J. M.; Evans, J. F.; Taketo, M. M.: Suppression

[43655] of intestinal polyposis in Apc(delta-716) knockout mice by inhibition

[43656] of cyclooxygenase 2 (COX-2). Cell 87: 803-809, 1996.

[43657]

[43658] 202. Oshima, M.; Oshima, H.; Kitagawa, K.; Kobayashi, M.; Itakura,

[43659] C.; Taketo, M.: Loss of Apc heterozygosity and abnormal tissue building

[43660] in nascent intestinal polyps in mice carrying a truncated Apc gene. Proc.

[43661] Nat. Acad. Sci. 92: 4482-4486, 1995.

[43662]

[43663] 203. Pandey, J. P.; Ebbesen, P.; Bulow, S.; Svendsen, L. B.; Fudenberg,

[43664] H. H.: IgG heavy-chain (Gm) allotypes in familial polyposis coli. Am.

[43665] J. Hum. Genet. 39: 133-136, 1986.

[43666]

[43667] 204. Papadopoulos, N.; Leach, F. S.; Kinzler, K. W.; Vogel-
stein, B.
[43668] : Monoallelic mutation analysis (MAMA) for identifying
germline mutations. Nature
[43669] Genet. 11: 99-102, 1995.

[43670]

[43671] 205. Patael, Y.; Figer, A.; Gershoni-Baruch, R.; Papa, M. Z.;
Risel,
[43672] S.; Shtoyerman-Chen, R.; Karasik, A.; Theodor, L.; Fried-
man, E.:
[43673] Common origin of the I1307K APC polymorphism in
Ashkenazi and non-Ashkenazi
[43674] Jews. Europ. J. Hum. Genet. 7: 555-559, 1999.

[43675]

[43676] 206. Pauli, R. M.; Pauli, M. E.; Hall, J. G.: Gardner syn-
drome and
[43677] periampullary malignancy. Am. J. Med. Genet. 6: 205-219,
1980.

[43678]

[43679] 207. Pavlides, G. P.; Milligan, F. D.; Clark, D. N.; Cohen, S.
B.;

[43680] Wennstrom, C. J.; Burbige, E. J.; Krush, A. J.; Murphy, E. A.: Hereditary
[43681] polyposis coli. I. The diagnostic value of colonoscopy, barium enema,
[43682] and fecal occult blood. Cancer 40: 2632-2639, 1977.

[43683]

[43684] 208. Peifer, M.: Cancer, catenins, and cuticle pattern: a complex
[43685] connection. Science 262: 1667-1668, 1993.

[43686]

[43687] 209. Petersen, G. M.; Francomano, C.; Kinzler, K.; Naka-
mura, Y.:
[43688] Presymptomatic direct detection of adenomatous polypo-
sis coli (APC)
[43689] gene mutations in familial adenomatous polyposis. Hum.
Genet. 91:
[43690] 307-311, 1993.

[43691]

[43692] 210. Petersen, G. M.; Parmigiani, G.; Thomas, D.: Missense mutations
[43693] in disease genes: a Bayesian approach to evaluate causal-

ity. Am.
[43694] J. Hum. Genet. 62: 1516-1524, 1998.

[43695]

[43696] 211. Petersen, G. M.; Shohat, T.; Brown, J.; Nakamura, Y.: Genetic
[43697] counseling for familial adenomatous polyposis (FAP) with chromosome
[43698] 5q linkage information. (Abstract) Am. J. Hum. Genet. 45 (suppl.):
[43699] A125 only, 1989.

[43700]

[43701] 212. Phillips, L. G., Jr.: Polyposis and carcinoma of the small bowel
[43702] and familial colonic polyposis. Dis. Colon Rectum 24: 478-481, 1981.

[43703]

[43704] 213. Pierce, E. R.: Pleiotropism and heterogeneity in hereditary
[43705] intestinal polyposis. Birth Defects Orig. Art. Ser. VII(2): 52-62,
[43706] 1972.

[43707]

[43708] 214. Pierce, E. R.; Weisbord, T.; McKusick, V. A.: Gardner's syndrome:

[43709] formal genetics and statistical analysis of a large Canadian kindred. Clin.

[43710] Genet. 1: 65-80, 1970.

[43711]

[43712] 215. Pilarski, R. T.; Brothman, A. R.; Benn, P.; Rosengren, S. S.

[43713] : Attenuated familial adenomatous polyposis in a man with an interstitial

[43714] deletion of chromosome arm 5q. Am. J. Med. Genet. 86: 321-324, 1999.

[43715]

[43716] 216. Polakis, P.: Mutations in the APC gene and their im-plications

[43717] for protein structure and function. Curr. Opin. Genet. Dev. 5: 66-71,

[43718] 1995.

[43719]

[43720] 217. Powell, S. M.; Petersen, G. M.; Krush, A. J.; Booker, S.;

Jen,

[43721] J.; Giardiello, F. M.; Hamilton, S. R.; Vogelstein, B.; Kinzler, K.

[43722] W.: Molecular diagnosis of familial adenomatous polyposis. New Eng.

[43723] J. Med. 329: 1982-1987, 1993.

[43724]

[43725] 218. Powell, S. M.; Zilz, N.; Beazer-Barclay, Y.; Bryan, T. M.; Hamilton,

[43726] S. R.; Thibodeau, S. N.; Vogelstein, B.; Kinzler, K. W.: APC mutations

[43727] occur early during colorectal tumorigenesis. Nature 359: 235-237,

[43728] 1992.

[43729]

[43730] 219. Prescott, S. M.; White, R. L.: Self-promotion? Intimate connections

[43731] between APC and prostaglandin H synthase-2. Cell 87: 783-786, 1996.

[43732]

[43733] 220. Prior, T. W.; Chadwick, R. B.; Papp, A. C.; Arcot, A. N.;

Isa,

[43734] A. M.; Pearl, D. K.; Stemmermann, G.; Percesepe, A.; Loukola, A.;

[43735] Aaltonen, L. A.; De La Chapelle, A.: The I1307K polymorphism of the

[43736] APC gene in colorectal cancer. Gastroenterology 116: 58-63, 1999.

[43737]

[43738] 221. Rasheed, S.; Rhim, J. S.; Gardner, E. J.: Inherited susceptibility

[43739] to retrovirus-induced transformation of Gardner syndrome cells. Am.

[43740] J. Hum. Genet. 35: 919-931, 1983.

[43741]

[43742] 222. Redston, M.; Nathanson, K. L.; Yuan, Z. Q.; Neuhausen, S. L.;

[43743] Satagopan, J.; Wong, N.; Yang, D.; Nafa, D.; Abrahamson, J.; Ozcelik,

[43744] H.; Antin-Ozerkis, D.; Andrulis, I.; and 13 others: The APC I1307K

[43745] allele and breast cancer risk. Nature Genet. 20: 13-14, 1998.

[43746]

[43747] 223. Rider, S. H.; Mazzullo, H. A.; Davis, M. B.; Delhanty, J. D.

[43748] A.: Familial polyposis coli: growth characteristics of karyotypically

[43749] variable cultured fibroblasts, response to epidermal growth factor

[43750] and the tumour promoter 12-0-tetradecanoyl phorbol-13-acetate. J.

[43751] Med. Genet. 23: 131-144, 1986.

[43752]

[43753] 224. Roose, J.; Huls, G.; van Beest, M.; Moerer, P.; van der Horn,

[43754] K.; Goldschmeding, R.; Logtenberg, T.; Clevers, H.: Synergy between

[43755] tumor suppressor APC and the beta-catenin-Tcf4 target Tcf1. Science 285:

[43756] 1923-1926, 1999.

[43757]

[43758] 225. Rosin-Arbesfeld, R.; Townsley, F.; Bienz, M.: The APC tumour

[43759] suppressor has a nuclear export function. Nature 406:
1009-1012,
[43760] 2000.

[43761]

[43762] 226. Rowan, A. J.; Lamlum, H.; Ilyas, M.; Wheeler, J.;
Straub, J.;
[43763] Papadopoulou, A.; Bicknell, D.; Bodmer, W. F.; Tomlinson,
I. P. M.
[43764] : APC mutations in sporadic colorectal tumors: a muta-
tional 'hotspot'
[43765] and interdependence of the 'two hits.' Proc. Nat. Acad. Sci.
97:
[43766] 3352-3357, 2000.

[43767]

[43768] 227. Rozen, P.; Naiman, T.; Strul, H.; Taussky, P.; Karmin-
sky, N.;
[43769] Shomrat, R.; Samuel, Z.; Yaron, Y.; Orr-Urtreger, A.: Clini-
cal and
[43770] screening implications of the I1307K adenomatous poly-
posis coli gene
[43771] variant in Israeli Ashkenazi Jews with familial colorectal
neoplasia:

[43772] evidence for a founder effect. Cancer 94: 2561-2568, 2002.

[43773]

[43774] 228. Rozen, P.; Samuel, Z.; Shomrat, R.; Legum, C.: Notable intrafamilial

[43775] phenotypic variability in a kindred with familial adenomatous polyposis

[43776] and an APC mutation in exon 9. Gut 45: 829-833, 1999.

[43777]

[43778] 229. Rozen, P.; Shomrat, R.; Strul, H.; Naiman, T.; Karminsky, N.;

[43779] Legum, C.; Orr-Urtreger, A.: Prevalence of the I1307K APC gene variant

[43780] in Israeli Jews of differing ethnic origin and risk for colorectal

[43781] cancer. Gastroenterology 116: 54-57, 1999.

[43782]

[43783] 230. Rubinfeld, B.; Albert, I.; Porfiri, E.; Fiol, C.; Munemitsu,

[43784] S.; Polakis, P.: Binding of GSK3-beta to the APC-beta-catenin complex

[43785] and regulation of complex assembly. Science 272: 1023-1026, 1996.

[43786]

[43787] 231. Rubinfeld, B.; Souza, B.; Albert, I.; Muller, O.; Cham-berlain,

[43788] S. H.; Masiarz, F. R.; Munemitsu, S.; Polakis, P.: Associa-tion of

[43789] the APC gene product with beta-catenin. Science 262: 1731-1734,

[43790] 1993.

[43791]

[43792] 232. Samowitz, W. S.; Burt, R. W.; Leppert, M.: Gastroin-testinal

[43793] polyposis and nonpolyposis syndromes. (Letter) New Eng. J. Med. 332:

[43794] 1518 only, 1995.

[43795]

[43796] 233. Sanchez, M. A.; Zali, M. R.; Khalil, A. A.; Ponce, R.; Font,

[43797] R. G.: Be aware of Gardner's syndrome: a review of the lit-erature. Am.

[43798] J. Gastroent. 71: 68-73, 1979.

[43799]

[43800] 234. Savage, P. T.: Polyposis coli associated with multiple tumours
[43801] in other parts of the body (Gardner's syndrome). Proc. Roy. Soc.
[43802] Med. 57: 402-403, 1964.

[43803]

[43804] 235. Schneider, N. R.; Cubilla, A. L.; Chaganti, R. S. K.: Association
[43805] of endocrine neoplasia with multiple polyposis of the colon. Cancer 51:
[43806] 1171-1175, 1983.

[43807]

[43808] 236. Schnitzler, M.; Dwight, T.; Robinson, B. G.: Sulindac increases
[43809] the expression of APC mRNA in malignant colonic epithelial cells:
[43810] an in vitro study. Gut 38: 707-713, 1996.

[43811]

[43812] 237. Schnur, P. L.; David, E.; Brown, P. W., Jr.; Beahrs, O. H.; Remine,
[43813] W. H.; Harrison, E. G., Jr.: Duodenal cancer and Gardner syndrome. J.A.M.A. 223:
[43814] 1229-1232, 1973.

[43815]

[43816] 238. Scott, R. J.; Froggart, N. J.; Trembath, R. C.; Evans, D. G.
[43817] R.; Hodgson, S. V.; Maher, E. R.: Familial infiltrative fibro-matosis
[43818] (desmoid tumours) (MIM135290) caused by a recurrent 3-prime APC gene
[43819] mutation. Hum. Molec. Genet. 5: 1921-1924, 1996.

[43820]

[43821] 239. Seki, M.; Tanaka, K.; Kikuchi-Yanoshita, R.; Konishi, M.; Fukunari,
[43822] H.; Iwama, T.; Miyaki, M.: Loss of normal allele of the APC gene
[43823] in an adrenocortical carcinoma from a patient with familial adenomatous
[43824] polyposis. Hum. Genet. 89: 298-300, 1992.

[43825]

[43826] 240. Sen-Gupta, S.; Van der Luijt, R. B.; Bowles, L. V.; Meera Khan,

[43827] P.; Delhanty, J. D. A.: Somatic mutation of APC gene in desmoid tumour

[43828] in familial adenomatous polyposis. (Letter) Lancet 342: 552-553,

[43829] 1993.

[43830]

[43831] 241. Shaoul, R.; Papsin, B.; Cutz, E.; Durie, P.: Congenital cholesteatoma

[43832] in a child carrying a gene mutation for adenomatous polyposis coli. J.

[43833] Pediat. Gastroent. Nutr. 28: 100-103, 1999.

[43834]

[43835] 242. Shemesh, E.: Adenomatous polyp of the common bile duct in familial

[43836] polyposis coli. Israel J. Med. Sci. 21: 701-702, 1985.

[43837]

[43838] 243. Shields, J. A.; Shields, C. L.; Eagle, R. C., Jr.; Singh, A.

[43839] D.: Adenocarcinoma arising from congenital hypertrophy of retinal

[43840] pigment epithelium. Arch. Ophthal. 119: 597-602, 2001.

[43841]

[43842] 244. Shields, J. A.; Shields, C. L.; Singh, A. D.: Acquired tumors

[43843] arising from congenital hypertrophy of the retinal pigment epithelium. Arch.

[43844] Ophthal. 118: 637-641, 2000.

[43845]

[43846] 245. Shih, I.-M.; Wang, T.-L.; Traverso, G.; Romans, K.; Hamilton,

[43847] S. R.; Ben-Sasson, S.; Kinzler, K. W.; Vogelstein, B.: Top-

down morphogenesis
[43848] of colorectal tumors. Proc. Nat. Acad. Sci. 98: 2640-2645, 2001.
[43849]

[43850] 246. Shneider, B. L.; Haque, S.; van Hoff, J.; Touloukian, R. J.;
[43851] West, A. B.: Familial adenomatous polyposis following liver transplantation
[43852] for a virilizing hepatoblastoma. J. Pediat. Gastroent. Nutr. 15:
[43853] 198-201, 1992.

[43854]

[43855] 247. Shull, L. N., Jr.; Fitts, C. T.: Lymphoid polyposis asso-ciated
[43856] with familial polyposis and Gardner's syndrome. Ann. Surg. 180:
[43857] 319-322, 1974.

[43858]

[43859] 248. Sieber, O. M.; Lamlum, H.; Crabtree, M. D.; Rowan, A. J.; Barclay,
[43860] E.; Lipton, L.; Hodgson, S.; Thomas, H. J. W.; Neale, K.;

Phillips,
[43861] R. K. S.; Farrington, S. M.; Dunlop, M. G.; and 9 others: Whole-gene
[43862] APC deletions cause classical familial adenomatous poly-posis, but
[43863] not attenuated polyposis or 'multiple' colorectal adeno-mas. Proc.
[43864] Nat. Acad. Sci. 99: 2954-2958, 2002.

[43865]

[43866] 249. Silverberg, M. S.; Clelland, C.; Murphy, J. E.; Steinhart, A.
[43867] H.; McLeod, R. S.; Greenberg, G. R.; Cohen, Z.; Simi-novitch, K. A.
[43868] : Carrier rate of APC I1307K is not increased in inflamma-tory bowel
[43869] disease patients of Ashkenazi Jewish origin. Hum. Genet. 108: 205-210,
[43870] 2001.

[43871]

[43872] 250. Simpson, R. D.; Harrison, E. G., Jr.; Mayo, C. W.: Mesenteric
[43873] fibromatosis in familial polyposis: a variant of Gardner's

syndrome. Cancer 17:
[43874] 526-534, 1964.

[43875]

[43876] 251. Sivak, M. V.; Jagelman, D. G.: Upper gastrointestinal endoscopy
[43877] in polyposis syndromes: familial polyposis coli and Gardner's syndrome. Gastrointest.
[43878] Endosc. 30: 102-104, 1984.

[43879]

[43880] 252. Smith, G.; Carey, F. A.; Beattie, J.; Wilkie, M. J. V.; Lightfoot,
[43881] T. J.; Coxhead, J.; Garner, R. C.; Steele, R. J. C.; Wolf, C. R.:
[43882] Mutations in APC, Kirsten-ras, and p53--alternative genetic pathways
[43883] to colorectal cancer. Proc. Nat. Acad. Sci. 99: 9433-9438, 2002.

[43884]

[43885] 253. Smith, K. J.; Johnson, K. A.; Bryan, T. M.; Hill, D. E.; Markowitz,
[43886] S.; Willson, J. K. V.; Paraskeva, C.; Petersen, G. M.; Hamilton, S.

[43887] R.; Vogelstein, B.; Kinzler, K. W.: The APC gene product in normal
[43888] and tumor cells. Proc. Nat. Acad. Sci. 90: 2846-2850, 1993.
[43889]
[43890] 254. Smith, W. G.: Multiple polyposis, Gardner's syndrome and desmoid
[43891] tumors. Dis. Colon Rectum 1: 323-332, 1958.
[43892]
[43893] 255. Smith, W. G.: Familial multiple polyposis: research tool for
[43894] investigating the etiology of carcinoma of the colon? Dis. Colon
[43895] Rectum 11: 17-31, 1968.
[43896]
[43897] 256. Solomon, E.; Voss, R.; Hall, V.; Bodmer, W. F.; Jass, J. R.;
[43898] Jeffreys, A. J.; Lucibello, F. C.; Patel, I.; Rider, S. H.: Chro-mosome
[43899] 5 allele loss in human colorectal carcinomas. Nature 328: 616-619,

[43900] 1987.

[43901]

[43902] 257. Soravia, C.; Berk, T.; Madlensky, L.; Mitri, A.; Cheng, H.; Gallinger,

[43903] S.; Cohen, Z.; Bapat, B.: Genotype-phenotype correlations in attenuated

[43904] adenomatous polyposis coli. Am. J. Hum. Genet. 62: 1290-1301, 1998.

[43905]

[43906] 258. Soravia, C.; Sugg, S. L.; Berk, T.; Mitri, A.; Cheng, H.; Gallinger,

[43907] S.; Cohen, Z.; Asa, S. L.; Bapat, B. V.: Familial adenomatous polyposis-associated

[43908] thyroid cancer: a clinical, pathological, and molecular genetics study. Am.

[43909] J. Path. 154: 127-135, 1999.

[43910]

[43911] 259. Spirio, L.; Green, J.; Robertson, J.; Robertson, M.; Otterud,

[43912] B.; Sheldon, J.; Howse, E.; Groden, J.; White, R.; Leppert, M.: The

[43913] identical 5-prime splice-site acceptor mutation in five attenuated

[43914] APC families from Newfoundland demonstrates a founder effect. Hum.

[43915] Genet. 105: 388-398, 1999.

[43916]

[43917] 260. Spirio, L.; Olschwang, S.; Groden, J.; Robertson, M.; Samowitz,

[43918] W.; Joslyn, G.; Gelbert, L.; Thliveris, A.; Carlson, M.; Otterud,

[43919] B.; Lynch, H.; Watson, P.; Lynch, P.; Laurent-Puig, P.; Burt, R.;

[43920] Hughes, J. P.; Thomas, G.; Leppert, M.; White, R.: Alleles of the

[43921] APC gene: an attenuated form of familial polyposis. Cell 75: 951-957,

[43922] 1993.

[43923]

[43924] 261. Spirio, L. N.; Samowitz, W.; Robertson, J.; Robertson, M.; Burt,

[43925] R. W.; Leppert, M.; White, R.: Alleles of APC modulate the frequency

[43926] and classes of mutations that lead to colon polyps. Nature Genet. 20:

[43927] 385-388, 1998.

[43928]

[43929] 262. Steinbach, G.; Lynch, P. M.; Phillips, R. K. S.; Wallace, M.

[43930] H.; Hawk, E.; Gordon, G. B.; Wakabayashi, N.; Saunders, B.; Shen,

[43931] Y.; Fujimura, T.; Su, L.-K.; Levin, B.: The effect of cele-coxib,

[43932] a cyclooxygenase-2 inhibitor, in familial adenomatous polyposis. New

[43933] Eng. J. Med. 342: 1946-1952, 2000.

[43934]

[43935] 263. Stella, A.; Resta, N.; Gentile, M.; Susca, F.; Mareni, C.; Montera,

[43936] M. P.; Guanti, G.: Exclusion of the APC gene as the cause of a variant

[43937] form of familial adenomatous polyposis (FAP). Am. J. Hum. Genet. 53:

[43938] 1031-1037, 1993.

[43939]

[43940] 264. Su, L.-K.; Abdalla, E. K.; Law, C. H. L.; Kohlmann, W.; Rashid,

[43941] A.; Vauthey, J.-N.: Biallelic inactivation of the APC gene is associated

[43942] with hepatocellular carcinoma in familial adenomatous polyposis coli. Cancer 92:

[43943] 332-339, 2001.

[43944]

[43945] 265. Su, L.-K.; Barnes, C. J.; Yao, W.; Qi, Y.; Lynch, P. M.; Steinbach,

[43946] G.: Inactivation of germline mutant APC alleles by attenuated somatic

[43947] mutations: a molecular genetic mosaicism for attenuated familial adenomatous

[43948] polyposis. Am. J. Hum. Genet. 67: 582-590, 2000. Note: Erratum:

[43949] Am. J. Hum. Genet. 67: 1045 only, 2000.

[43950]

[43951] 266. Su, L.-K.; Kinzler, K. W.; Vogelstein, B.; Preisinger, A. C.;

[43952] Moser, A. R.; Luongo, C.; Gould, K. A.; Dove, W. F.: Multi-
ple intestinal
[43953] neoplasia caused by a mutation in the murine homolog of
the APC gene. Science 256:
[43954] 668-670, 1992.

[43955]

[43956] 267. Su, L.-K.; Kohlmann, W.; Ward, P. A.; Lynch, P. M.:
Different
[43957] familial adenomatous polyposis phenotypes resulting
from deletions
[43958] of the entire APC exon 15. Hum. Genet. 111 88-95, 2002.

[43959]

[43960] 268. Su, L.-K.; Steinbach, G.; Sawyer, J. C.; Hindi, M.;
Ward, P.
[43961] A.; Lynch, P. M.: Genomic rearrangements of the APC tu-
mor-suppressor
[43962] gene in familial adenomatous polyposis. Hum. Genet. 106:
101-107,
[43963] 2000.

[43964]

[43965] 269. Su, L.-K.; Vogelstein, B.; Kinzler, K. W.: Association

of the
[43966] APC tumor suppressor protein with catenins. Science 262:
1734-1737,
[43967] 1993.

[43968]

[43969] 270. Sugihara, K.; Muto, T.; Kamiya, J.; Konishi, F.;
Sawada, T.;
[43970] Morioka, Y.: Gardner's syndrome associated with peri-
ampullary carcinoma,
[43971] duodenal and gastric adenomatosis: report of a case. Dis.
Colon Rectum 25:
[43972] 766-771, 1982.

[43973]

[43974] 271. Sulekova, Z.; Ballhausen, W. G.: A novel coding exon
of the
[43975] human adenomatous polyposis coli gene. Hum. Genet. 96:
469-471,
[43976] 1995.

[43977]

[43978] 272. Takaku, K.; Oshima, M.; Miyoshi, H.; Matsui, M.;
Seldin, M. F.;

[43979] Taketo, M. M.: Intestinal tumorigenesis in compound mutant mice of

[43980] both Dpc4 (Smad4) and Apc genes. Cell 92: 645-656, 1998.

[43981]

[43982] 273. Thakker, N.; Davies, R.; Horner, K.; Armstrong, J.; Clancy, T.;

[43983] Guy, S.; Harris, R.; Sloan, P.; Evans, G.: The dental phenotype in

[43984] familial adenomatous polyposis: diagnostic application of a weighted

[43985] scoring system for changes on dental panoramic radiographs. J. Med.

[43986] Genet. 32: 458-464, 1995.

[43987]

[43988] 274. Thompson, J. S.; Harned, R. K.; Anderson, J. C.; Hodgson, P.

[43989] E.: Papillary carcinoma of the thyroid and familial polyposis coli. Dis.

[43990] Colon Rectum 26: 583-585, 1983.

[43991]

[43992] 275. Tops, C. M. J.; van der Klift, H. M.; van der Luijt, R. B.; Griffioen,

[43993] G.; Taal, B. G.; Vasen, H. F. A.; Khan, P. M.: Non-allelic heterogeneity

[43994] of familial adenomatous polyposis. Am. J. Med. Genet. 47: 563-567,

[43995] 1993.

[43996]

[43997] 276. Tops, C. M. J.; Vasen, H. F. A.; van Berge Henegouwen, G.; Simoons,

[43998] P. P.; van de Klift, H. M.; van Leeuwen, I. S. J.; Breukel, C.; Fodde,

[43999] R.; den Hartog Jager, F. C. A.; Nagengast, F. M.; Griffioen, G.; Meera

[44000] Khan, P.: Genetic evidence that Turcot syndrome is not allelic to

[44001] familial adenomatous polyposis. Am. J. Med. Genet. 43: 888-893,

[44002] 1992.

[44003]

[44004] 277. Tops, C. M. J.; Wijnen, J. T.; Griffioen, G.; van Leeuwen, I.

[44005] S. J.; Vasen, H. F. A.; den Hartog Jager, F. C. A.; Breukel, C.; Nagengast,

[44006] F. M.; van der Klift, H. M.; Lamers, C. B. H. W.; Meera Khan, P.:

[44007] Presymptomatic diagnosis of familial adenomatous polyposis by bridging

[44008] DNA markers. Lancet II: 1361-1363, 1989.

[44009]

[44010] 278. Traboulsi, E. I.; Krush, A. J.; Gardner, E. J.; Booker, S. V.;

[44011] Offerhaus, G. J. A.; Yardley, J. H.; Hamilton, S. R.; Luk, G. D.;

[44012] Giardiello, F. M.; Welsh, S. B.; Hughes, J. P.; Maumenee, I. H.:

[44013] Prevalence and importance of pigmented ocular fundus lesions in Gardner's

[44014] syndrome. New Eng. J. Med. 316: 661-667, 1987.

[44015]

[44016] 279. Traverso, G.; Shuber, A.; Levin, B.; Johnson, C.; Olsson, L.;

[44017] Schoetz, D. J., Jr.; Hamilton, S. R.; Boynton, K.; Kinzler, K. W.;

[44018] Vogelstein, B.: Detection of APC mutations in fecal DNA from patients
[44019] with colorectal tumors. New Eng. J. Med. 346: 311-320, 2002.
[44020]

[44021] 280. Tsujii, M.; DuBois, R. N.: Alterations in cellular adhe-sion
[44022] and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide
[44023] synthase 2. Cell 83: 493-501, 1995.
[44024]

[44025] 281. Turcot, J.; Despres, J. P.; St. Pierre, F.: Malignant tu-mors
[44026] of the central nervous system associated with familial polyposis of
[44027] the colon: report of two cases. Dis. Colon Rectum 2: 465-468, 1959.
[44028]

[44029] 282. Utsunomiya, J.; Nakamura, T.: The occult osteoma-tous changes
[44030] in the mandible in patients with familial polyposis coli.. Brit.

[44031] J. Surg. 62: 45-51, 1975.

[44032]

[44033] 283. van der Luijt, R.; Meera Khan, P.; Vasen, H.; van Leeuwen, C.;

[44034] Tops, C.; Roest, P.; den Dunnen, J.; Fodde, R.: Rapid de-

[44035] tection of translation-terminating mutations at the adenomatous polyposis coli

[44036] (APC) gene by direct protein truncation test. Genomics 20: 1-4,

[44037] 1994.

[44038]

[44039] 284. van der Luijt, R. B.; Khan, P. M.; Vasen, H. F. A.; Breukel,

[44040] C.; Tops, C. M. J.; Scott, R. J.; Fodde, R.: Germline muta-

[44041] tions in the 3-prime part of APC exon 15 do not result in trun-

[44042] cated proteins and are associated with attenuated adenomatous polypo-

[44043] sis coli. Hum. Genet. 98: 727-734, 1996.

[44044]

[44045] 285. van der Luijt, R. B.; Khan, P. M.; Vasen, H. F. A.; Tops, C.

[44046] M. J.; van Leeuwen-Cornelisse, I. S. J.; Wijnen, J. T.; van der Klift,

[44047] H. M.; Plug, R. J.; Griffioen, G.; Fodde, R.: Molecular analysis

[44048] of the APC gene in 105 Dutch kindreds with familial adenomatous polyposis:

[44049] 67 germline mutations identified by DGGE, PTT, and Southern analysis. Hum.

[44050] Mutat. 9: 7-16, 1997.

[44051]

[44052] 286. van der Luijt, R. B.; Tops, C. M. J.; Khan, P. M.; van der Klift,

[44053] H. M.; Breukel, C.; van Leeuwen-Cornelisse, I. S. J.; Dauwerse, H.

[44054] G.; Beverstock, G. C.; van Noort, E.; Snel, P.; Slors, F. J. M.; Vasen,

[44055] H. F. A.; Fodde, R.: Molecular, cytogenetic, and phenotypic studies

[44056] of a constitutional reciprocal translocation

t(5;10)(q22;q25) responsible

[44057] for familial adenomatous polyposis in a Dutch pedigree. Genes Chromosomes

[44058] Cancer 13: 192-202, 1995.

[44059]

[44060] 287. van Es, J.; Kirkpatrick, C.; van de Wetering, M.; Molenaar, M.;

[44061] Miles, A.; Kuipers, J.; Destree, O.; Peifer, M.; Clevers, H.: Identification

[44062] of APC2, a homologue of the adenomatous polyposis coli tumour suppressor. Curr.

[44063] Biol. 9: 105-108, 1999.

[44064]

[44065] 288. Vanhoutte, J. J.: Polypoid lymphoid hyperplasia of the terminal

[44066] ileum in patients with familial polyposis coli and with Gardner's

[44067] syndrome. Am. J. Roentgen. 110: 340-342, 1970.

[44068]

[44069] 289. Varesco, L.; Thomas, H. J. W.; Cottrell, S.; Murday, V.; Fennell,

[44070] S. J.; Williams, S.; Searle, S.; Sheer, D.; Bodmer, W. F.; Frischauf,

[44071] A.-M.; Solomon, E.: CpG island clones from a deletion en- compassing

[44072] the gene for adenomatous polyposis coli. Proc. Nat. Acad. Sci. 86:

[44073] 10118-10122, 1989.

[44074]

[44075] 290. Veale, A. M.: Clinical and genetic problems in familial intestinal

[44076] polyposis. Gut 1: 285-290, 1960.

[44077]

[44078] 291. Venkitachalam, P. S.; Hirsch, E.; Elguezabal, A.; Littman, L.

[44079] : Multiple lymphoid polyposis and familial polyposis of the colon:

[44080] a genetic relationship. Dis. Colon Rectum 21: 336-341, 1978.

[44081]

[44082] 292. Waddell, W. R.; Loughry, R. W.: Sulindac for polyposis of the

[44083] colon. J. Surg. Oncol. 24: 83-87, 1983.

[44084]

[44085] 293. Wallis, Y. L.; Macdonald, F.; Hulten, M.; Morton, J. E. V.; McKeown,

[44086] C. M.; Neoptolemos, J. P.; Keighley, M.; Morton, D. G.: Genotype-phenotype

[44087] correlation between position of constitutional APC gene mutation and

[44088] CHRPE expression in familial adenomatous polyposis. Hum. Genet. 94:

[44089] 543-548, 1994.

[44090]

[44091] 294. Wallis, Y. L.; Morton, D. G.; McKeown, C. M.; Macdonald, F.:

[44092] Molecular analysis of the APC gene in 205 families: extended genotype-phenotype

[44093] correlations in FAP and evidence for the role of APC amino acid changes

[44094] in colorectal cancer predisposition. J. Med. Genet. 36: 14-20, 1999.

[44095]

[44096] 295. Walsh, N.; Qizilbash, A.; Banerjee, R.; Waugh, G. A.: Biliary
[44097] neoplasia in Gardner's syndrome. Arch. Path. Lab. Med. 111: 76-77,
[44098] 1987.

[44099]

[44100] 296. Wasan, H. S.; Novelli, M.; Bee, J.; Bodmer, W. F.: Di-etary fat
[44101] influences on polyp phenotype in multiple intestinal neo-plasia mice. Proc.
[44102] Nat. Acad. Sci. 94: 3308-3313, 1997.

[44103]

[44104] 297. Watanabe, H.; Enjoji, M.; Yao, T.; Ohsato, K.: Gastric lesions
[44105] in familial adenomatosis coli: their incidence and histo-logic analysis. Hum.
[44106] Path. 9: 269-283, 1978.

[44107]

[44108] 298. Westbrook, C. A.; Chmura, S. J.; Arenas, R. B.; Kim, S. Y.; Otto,
[44109] G.: Human APC gene expression in rodent colonic epithe-

lium in vivo
[44110] using liposomal gene delivery. Hum. Molec. Genet. 3:
2005-2010,
[44111] 1994.

[44112]

[44113] 299. Wheeler, J. M. D.; Warren, B. F.; Mortensen, N. J. M.;
Kim, H.
[44114] C.; Biddolph, S. C.; Elia, G.; Beck, N. E.; Williams, G. T.;
Shepherd,
[44115] N. A.; Bateman, A. C.; Bodmer, W. F.: An insight into the
genetic
[44116] pathway of adenocarcinoma of the small intestine. Gut 50:
218-223,
[44117] 2002.

[44118]

[44119] 300. Won, Y.-J.; Park, K. J.; Kwon, H.-J.; Lee, J.-H.; Kim,
J.-H.;
[44120] Kim, Y. J.; Chun, S. H.; Han, H.-J.; Park, J.-G.: Germline
mutations
[44121] of the APC gene in Korean familial adenomatous polyposis
patients. J.
[44122] Hum. Genet. 44: 103-108, 1999.

[44123]

[44124] 301. Woodage, T.; King, S. M.; Wacholder, S.; Hartge, P.; Struewing,

[44125] J. P.; McAdams, M.; Laken, S. J.; Tucker, M. A.; Brody, L. C.: The

[44126] APC I1307K allele and cancer risk in a community-based study of Ashkenazi

[44127] Jews. Nature Genet. 20: 62-65, 1998.

[44128]

[44129] 302. Xia, L.; St. Denis, K. A.; Bapat, B.: Evidence for a novel exon

[44130] in the coding region of the adenomatous polyposis coli (APC) gene. Genomics 28:

[44131] 589-591, 1995.

[44132]

[44133] 303. Yan, H.; Dobbie, Z.; Gruber, S. B.; Markowitz, S.; Romans, K.;

[44134] Giardiello, F. M.; Kinzler, K. W.; Vogelstein, B.: Small changes

[44135] in expression affect predisposition to tumorigenesis. Nature Genet. 30:

[44136] 25-26, 2002.

[44137]

[44138] 304. Yan, H.; Papadopoulos, N.; Marra, G.; Perrera, C.; Jir-

icny, J.;

[44139] Boland, C. R.; Lynch, H. T.; Chadwick, R. B.; de la Chapelle,

A.;

[44140] Berg, K.; Eshleman, J. R.; Yuan, W.; Markowitz, S.; Laken, S.

J.;

[44141] Lengauer, C.; Kinzler, K. W.; Vogelstein, B.: Conversion of

diploidy

[44142] to haploidy. Nature 403: 723-724, 2000.

[44143]

[44144] 305. Yonemoto, R. H.; Slayback, J. B.; Byron, R. L., Jr.;

Rosen, R.

[44145] B.: Familial polyposis of the entire gastrointestinal tract.

Arch.

[44146] Surg. 99: 427-434, 1969.

[44147]

[44148] 306. Young, J.; Simms, L. A.; Tarish, J.; Buttenshaw, R.;

Knight,

[44149] N.; Anderson, G. J.; Bell, A.; Leggett, B.: A family with at-

tenuated

[44150] familial adenomatous polyposis due to a mutation in the alternatively

[44151] spliced region of APC exon 9. Hum. Mutat. 11: 450-455, 1998.

[44152]

[44153] 307. Yuan, Z. Q.; Kasprzak, L.; Gordon, P. H.; Pinsky, L.; Foulkes,

[44154] W. D.: I1307K APC and hMLH1 mutations in a non-Jewish family with

[44155] hereditary non-polyposis colorectal cancer. Clin. Genet. 54: 368-370,

[44156] 1998.

[44157] 6636.

[44158] Riccio, A.; Grimaldi, G.; Verde, P.; Sebastio, G.; Boast, S.;

[44159] Blasi, F.: The human urokinase-plasminogen activator gene and its

[44160] promoter. Nucleic Acids Res. 13: 2759-2771, 1985.

[44161] 6637.

[44162] Seeds, N. W.; Basham, M. E.; Haffke, S. P.: Neuronal migration

[44163] is retarded in mice lacking the tissue plasminogen activator gene. Proc.

[44164] Nat. Acad. Sci. 96: 14118-14123, 1999.

[44165] 6638.

[44166] Seeds, N. W.; Williams, B. L.; Bickford, P. C.: Tissue plasminogen

[44167] activator induction in Purkinje neurons after cerebellar motor learning. Science 270:

[44168] 1992-1994, 1996.

[44169] 6639.

[44170] Stead, N. W.; Bauer, K. A.; Kinney, T. R.; Lewis, J. G.; Campbell,

[44171] E. E.; Shifman, M. A.; Rosenberg, R. D.; Pizzo, S. V.: Venous thrombosis

[44172] in a family with defective release of vascular plasminogen activator

[44173] and elevated plasma factor VIII/von Willebrand's factor. Am. J. Med. 74:

[44174] 33-39, 1983.

[44175] 6640.

[44176] Tishkoff, S. A.; Pakstis, A. J.; Stoneking, M.; Kidd, J. R.; De-

stro-Bisol,

[44177] G.; Sanjantila, A.; Lu, R.; Deinard, A. S.; Sirugo, G.; Jenkins, T.;

[44178] Kidd, K. K.; Clark, A. G.: Short tandem-repeat polymorphism/Alu haplotype

[44179] variation at the PLAT locus: implications for modern human origins. Am.

[44180] J. Hum. Genet. 67: 901-925, 2000.

[44181] 6641.

[44182] Tripputi, P.; Blasi, F.; Emanuel, B. S.; Letofsky, J.; Croce,

[44183] C. M.: Tissue-type plasminogen activator gene is on chromosome 8. Cytogenet.

[44184] Cell Genet. 42: 24-28, 1986.

[44185] 6642.

[44186] Verheijen, J. H.; Visse, R.; Wijnen, J. T.; Chang, G. T. G.; Kluft,

[44187] C.; Meera Khan, P.: Assignment of the human tissue-type plasminogen

[44188] activator gene (PLAT) to chromosome 8. Hum. Genet. 72: 153-156,

[44189] 1986.

[44190] 6643.

[44191] Visse, R.; Chang, G. T. G.; Wijnen, J. T.; Verheijen, J. H.; Kluft,

[44192] C.; Meera Khan, P.: Provisional assignment of human tissue-type plasminogen

[44193] activator (PLAT) to chromosome 8. (Abstract) Cytogenet. Cell Genet. 40:

[44194] 771 only, 1985.

[44195] 6644.

[44196] Yang-Feng, T. L.; Opdenakker, G.; Volckaert, G.; Francke, U.:

[44197] Mapping of the human tissue-type plasminogen activator (PLAT) gene

[44198] to chromosome 8 (8p12-q11.2). (Abstract) Cytogenet. Cell Genet. 40:

[44199] 784 only, 1985.

[44200] 6645.

[44201] Yang-Feng, T. L.; Opdenakker, G.; Volckaert, G.; Francke, U.:

[44202] Human tissue-type plasminogen activator gene located near chromosomal

[44203] breakpoint in myeloproliferative disorder. Am. J. Hum. Genet. 39:

[44204] 79-87, 1986.

[44205] 6646.

[44206] Rijneveld, A. W.; Levi, M.; Florquin, S.; Speelman, P.; Carmeliet,

[44207] P.; van der Poll, T.: Urokinase receptor is necessary for adequate

[44208] host defense against pneumococcal pneumonia. J. Immun. 168: 3507-3511,

[44209] 2002.

[44210] 6647.

[44211] Aitchison, K.; Ogilvie, D.; Honeyman, M.; Thompson, E.; Sykes,

[44212] B.: Homozygous osteogenesis imperfecta unlinked to collagen I genes. Hum.

[44213] Genet. 78: 233-236, 1988.

[44214] 6648.

[44215] Barsh, G. S.; Byers, P. H.: Reduced secretion of structurally

[44216] abnormal type I procollagen in a form of osteogenesis imperfecta. Proc.

[44217] Nat. Acad. Sci. 78: 5142-5146, 1981.

[44218] 6649.

[44219] Bateman, J. F.; Chan, D.; Walker, I. D.; Rogers, J. G.; Cole, W.

[44220] G.: Lethal perinatal osteogenesis imperfecta due to the substitution

[44221] of arginine for glycine at residue 391 of the alpha-1(I) chain of

[44222] type I collagen. J. Biol. Chem. 262: 7021-7027, 1987.

[44223] 6650.

[44224] Bonadio, J.: Personal Communication. Ann Arbor, Mich. 3/1990.

[44225] 6651.

[44226] Bonadio, J.: Personal Communication. Ann Arbor, Mich. 2/7/1992.

[44227] 6652.

[44228] Bonadio, J.; Ramirez, F.; Barr, M.: An intron mutation in the

[44229] human alpha-1(I) collagen gene alters the efficiency of pre-mRNA splicing

[44230] and is associated with osteogenesis imperfecta type II. J.

Biol.

[44231] Chem. 265: 2262-2268, 1990.

[44232] 6653.

[44233] Cohn, D. H.; Starman, B. J.; Blumberg, B.; Byers, P. H.: Re-
currence

[44234] of lethal osteogenesis imperfecta due to parental mo-
saicism for a

[44235] dominant mutation in a human type I collagen gene
(COL1A1). Am. J.

[44236] Hum. Genet. 46: 591-601, 1990.

[44237] 6654.

[44238] Brecher, R.; Bird, A. C.: Adult vitelliform macular dystro-
phy. Eye 4:

[44239] 210-215, 1990.

[44240] 6655.

[44241] Cullen, T. S.: Embryology, Anatomy, and Diseases of the
Umbilicus

[44242] Together with Diseases of the Urachus. Philadelphia: W. B.
Saunders

[44243] (pub.) 1916.

[44244] 6656.

[44245] Feingold, M.; Shiere, F.; Fogels, H. R.; Donaldson, D. D.: Rieger's
[44246] syndrome. Pediatrics 44: 564-569, 1969.

[44247] 6657.

[44248] Friedman, J. M.: Umbilical dysmorphology: the importance of contemplating
[44249] the belly button. Clin. Genet. 28: 343-347, 1985.

[44250] 6658.

[44251] Chu, X.; Thompson, D.; Yee, L. J.; Sung, L. A.: Genomic or-
ganization
[44252] of mouse and human erythrocyte tropomodulin genes en-
coding the pointed
[44253] end capping protein for the actin filaments. Gene 256:
271-281,
[44254] 2000.

[44255] 6659.

[44256] Fowler, V. M.; Sussmann, M. A.; Miller, P. G.; Flucher, B. E.;

[44257] Daniels, M. P.: Tropomodulin is associated with the free
(pointed)
[44258] ends of the thin filaments in rat skeletal muscle. J. Cell
Biol. 120:

[44259] 411-420, 1993.

[44260] 6660.

[44261] Gilligan, D. M.; Bennett, V.: The junctional complex of the membrane

[44262] skeleton. Semin. Hemat. 30: 74-83, 1993.

[44263] 6661.

[44264] Lench, N. J.; Telford, E. A.; Andersen, S. E.; Moynihan, T. P.;

[44265] Robinson, P. A.; Markham, A. F.: An EST and STS-based YAC contig

[44266] map of human chromosome 9q22.3 Genomics 38: 199-205, 1996.

[44267] 6662.

[44268] Sung, L. A.; Fan, Y.-S.; Lin, C. C.: Gene assignment, expression,

[44269] and homology of human tropomodulin. Genomics 34: 92-96, 1996.

[44270] 6663.

[44271] Sung, L. A.; Fan, Y. S.; Lambert, K.; Fowler, V.; Chien, S.; Lin,

[44272] C.: Assignment of human erythrocyte tropomodulin gene

to q22 of chromosome
[44273] 9. (Abstract) Cytogenet. Cell Genet. 58: 1944, 1991.

[44274] 6664.

[44275] Sung, L. A.; Fowler, V. M.; Lambert, K.; Chien, S.: Molecu-
lar
[44276] cloning of human erythroid tropomodulin. (Abstract)
FASEB J. 5:
[44277] A1625, 1991.

[44278] 6665.

[44279] Sung, L. A.; Fowler, V. M.; Lambert, K.; Sussman, M. A.;
Karr,
[44280] D.; Chien, S.: Molecular cloning and characterization of
human fetal
[44281] liver tropomodulin: a tropomyosin-binding protein. J. Biol.
Chem. 267:
[44282] 2616-2621, 1992.

[44283] 6666.

[44284] White, R. A.; Dowler, L. L.; Woo, M.; Adkison, L. R.; Pal, S.;

[44285] Gershon, D.; Fowler, V. M.: The tropomodulin (Tmod) gene
maps to
[44286] chromosome 4, closely linked to Mup1. Mammalian

Genome 6: 332-333,
[44287] 1995.

[44288] 6667.

[44289] Hunt, C. C. J.; Eyre, H. J.; Akkari, P. A.; Meredith, C.;
Dorosz,
[44290] S. M.; Wilton, S. D.; Callen, D. F.; Laing, N. G.; Baker, E.:
Assignment
[44291] of the human beta tropomyosin gene (TPM2) to band
9p13 by fluorescence
[44292] in situ hybridization. Cytogenet. Cell Genet. 71: 94-95,
1995.
[44293] 6668.

[44294] Sri Widada, J.; Ferraz, C.; Capony, J.-P.; Liautard, J.-P.:
Complete
[44295] nucleotide sequence of the adult skeletal isoform of hu-
man skeletal
[44296] muscle beta-tropomyosin. Nucleic Acids Res. 16: 3109
only, 1988.
[44297] 6669.

[44298] Tiso, N.; Rampoldi, L.; Pallavicini, A.; Zimbello, R.; Pan-
dolfo,
[44299] D.; Valle, G.; Lanfranchi, G.; Danieli, G. A.: Fine mapping

of five
[44300] human skeletal muscle genes: alpha-tropomyosin, beta-tropomyosin,
[44301] troponin-I slow-twitch, troponin-I fast-twitch, and troponin-C fast. Biochem.
[44302] Biophys. Res. Commun. 230: 347-350, 1997.

[44303] 6670.

[44304] Blanchard, E. M.; Iizuka, K.; Christe, M.; Conner, D. A.; Geisterfer-Lowrance,
[44305] A.; Schoen, F. J.; Maughan, D. W.; Seidman, C. E.; Seidman, J. G.
[44306] : Targeted ablation of the murine alpha-tropomyosin gene. Circulation
[44307] Res. 81: 1005-1010, 1997.

[44308] 6671.

[44309] Brown, J. H.; Kim, K.-H.; Jun, G.; Greenfield, N. J.; Dominguez,
[44310] R.; Volkmann, N.; Hitchcock-DeGregori, S. E.; Cohen, C.: Deciphering
[44311] the design of the tropomyosin molecule. Proc. Nat. Acad. Sci. 98:
[44312] 8496-8501, 2001.

[44313] 6672.

[44314] Eyre, H.; Akkari, P. A.; Wilton, S. D.; Callen, D. C.; Baker, E.;

[44315] Laing, N. G.: Assignment of the human skeletal muscle alpha-tropomyosin

[44316] gene (TPM1) to band 15q22 by fluorescence in situ hybridization. Cytogenet.

[44317] Cell Genet. 69: 15-17, 1995.

[44318] 6673.

[44319] Torrents, D.; Mykkanen, J.; Pineda, M.; Feliubadalo, L.; Estevez,

[44320] R.; de Cid, R.; Sanjurjo, P.; Zorzano, A.; Nunes, V.; Huoponen, K.;

[44321] Reinikainen, A.; Simell, O.; Savontaus, M.-L.; Aula, P.; Palacin,

[44322] M.: Identification of SLC7A7, encoding y(+)LAT-1, as the lysinuric

[44323] protein intolerance gene. Nature Genet. 21: 293-296, 1999.

[44324] 6674.

[44325] Tauchi, H.; Matsuura, S.; Isomura, M.; Kinjo, T.; Nakamura, A.;

[44326] Sakamoto, S.; Kondo, N.; Endo, S.; Komatsu, K.; Nakamura, Y.: Sequence
[44327] analysis of an 800-kb genomic DNA region on chromosome 8q21 that contains
[44328] the Nijmegen breakage syndrome gene, NBS1. Genomics 55: 242-247,
[44329] 1999.

[44330] 6675.

[44331] Hamajima, N.; Matsuda, K.; Sakata, S.; Tamaki, N.; Sasaki, M.;
[44332] Nonaka, M.: A novel gene family defined by human dihydropyrimidinase
[44333] and three related proteins with differential tissue distribution. Gene 180:
[44334] 157-163, 1996.

[44335] 6676.

[44336] Sia, C. L.; Traniello, S.; Pontremoli, S.; Horecker, B. L.: Studies
[44337] on the subunit structure of rabbit liver fructose diphosphatase. Arch.
[44338] Biochem. Biophys. 132: 325-330, 1969.

[44339] 6677.

[44340] Boesiger, P.; Buchli, R.; Meier, D.; Steinmann, B.; Gitzel-
mann,
[44341] R.: Changes of liver metabolite concentrations in adults
with disorders
[44342] of fructose metabolism after intravenous fructose by (31)P
magnetic
[44343] resonance spectroscopy. Pediat. Res. 36: 436-440, 1994.

[44344] 6678.

[44345] Bonthron, D. T.; Brady, N.; Donaldson, I. A.; Steinmann, B.:
Molecular
[44346] basis of essential fructosuria: molecular cloning and mu-
tational analysis
[44347] of human ketohexokinase (fructokinase). Hum. Molec.
Genet. 3: 1627-1631,
[44348] 1994.

[44349] 6679.

[44350] Czapek, F.: Eine seltene Form von Diabetes mellitus.
Prager Med.
[44351] Wochenschr. 1: 245-249, 1876.

[44352] 6680.

[44353] Gitzelmann, R.; Steinmann, B.; Van Den Berghe, G.: Disorders of
[44354] fructose metabolism.In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.;
[44355] Valle, D.: The Metabolic Basis of Inherited Disease. New York:
[44356] McGraw-Hill (pub.) (6th ed.): 1989. Pp. 399-424.

[44357] 6681.

[44358] Hayward, B. E.; Fantes, J. A.; Warner, J. P.; Intody, S.; Leek,
[44359] J. P.; Markham, A. F.; Bonthron, D. T.: Co-localization of the ketohexokinase
[44360] and glucokinase regulator genes to a 500-kb region of chromosome 2p23. Mammalian
[44361] Genome 7: 454-458, 1996.

[44362] 6682.

[44363] Khachadurian, A. K.: Nonalimentary fructosuria. (Letter) Pediatrics 32:
[44364] 455-457, 1963.

[44365] 6683.

[44366] Laron, Z.: Essential benign fructosuria. Arch. Dis. Child. 36:

[44367] 273-277, 1961.

[44368] 6684.

[44369] Lasker, M.: Essential fructosuria. Hum. Biol. 13: 51-63, 1941.

[44370] 6685.

[44371] Schapira, F.; Schapira, G.; Dreyfus, J.-C.: La lesion enzymatique

[44372] de la fructosurie benigne. Enzymol. Biol. Clin. 1: 170-175, 1961.

[44373] 6686.

[44374] Zimmer, K.: I. Levulose im Harn eines Diabetikers. Dtsch. Med.

[44375] Wochenschr. 2: 329 only, 1876.

[44376] 6687.

[44377] Richards, F. M.; McKee, S. A.; Rajpar, M. H.; Cole, T. R. P.;

[44378] Evans, D. G. R.; Jankowski, J. A.; McKeown, C.; Sanders, D. S. A.;

[44379] Maher, E. R.: Germline E-cadherin gene (CDH1) mutations predispose

[44380] to familial gastric cancer and colorectal cancer. Hum. Molec. Genet. 8:

[44381] 607-610, 1999.

[44382] 6688.

[44383] Riethmacher, D.; Brinkmann, V.; Birchmeier, C.: A targeted mutation

[44384] in the mouse E-cadherin gene results in defective preimplantation

[44385] development. Proc. Nat. Acad. Sci. 92: 855-859, 1995.

[44386] 6689.

[44387] Risinger, J. I.; Berchuck, A.; Kohler, M. F.; Boyd, J.: Mutations

[44388] of the E-cadherin gene in human gynecologic cancers. Nature Genet. 7:

[44389] 98-102, 1994.

[44390] 6690.

[44391] Scherer, G.; Bausch, E.; Gaa, A.; von Deimling, O.: Gene mapping

[44392] on mouse chromosome 8 by interspecific crosses: new data on a linkage

[44393] group conserved on human chromosome 16q. Genomics 5: 275-282, 1989.

[44394] 6691.

[44395] Sokoloff, B.: Predisposition to cancer in the Bonaparte family. Am.
[44396] J. Surg. 40: 673-678, 1938.

[44397] 6692.

[44398] Wheeler, J. M. D.; Warren, B. F.; Mortensen, N. J. M.; Kim, H.
[44399] C.; Biddolph, S. C.; Elia, G.; Beck, N. E.; Williams, G. T.; Shepherd,
[44400] N. A.; Bateman, A. C.; Bodmer, W. F.: An insight into the genetic
[44401] pathway of adenocarcinoma of the small intestine. Gut 50: 218-223,
[44402] 2002.

[44403] 6693.

[44404] Yoon, K.-A.; Ku, J.-L.; Yang, H.-K.; Kim, W. H.; Park, S. Y.;
[44405] Park, J.-G.: Germline mutations of E-cadherin gene in Ko-rean familial
[44406] gastric cancer patients. J. Hum. Genet. 44: 177-180, 1999.
[44407] 6694.

[44408] Cybulsky, M.; Fries, J. W.; Williams, A. J.; Sultan, P.; Eddy,

[44409] R. L.; Byers, M. G.; Shows, T. B.; Gimbrone, M. A., Jr.; Collins,

[44410] T.: The human VCAM1 gene is assigned to chromosome 1p31-p32. (Abstract) Cytogenet.

[44411] Cell Genet. 58: 1852, 1991.

[44412] 6695.

[44413] Cybulsky, M. I.; Fries, J. W. U.; Williams, A. J.; Sultan, P.;

[44414] Eddy, R.; Byers, M.; Shows, T.; Gimbrone, M. A., Jr.; Collins, T.

[44415] : Gene structure, chromosomal location, and basis for alternative

[44416] mRNA splicing of the human VCAM1 gene. Proc. Nat. Acad. Sci. 88:

[44417] 7859-7863, 1991.

[44418] 6696.

[44419] Kumar, A. G.; Dai, X. Y.; Kozak, C. A.; Mims, M. P.; Gotto, A.

[44420] M.; Ballantyne, C. M.: Murine VCAM-1: molecular cloning, mapping,

[44421] and analysis of a truncated form. J. Immun. 153: 4088-4098, 1994.

[44422] 6697.

[44423] Bardwick, P. A.; Zvaifler, N. J.; Gill, G. N.; Newman, D.; Greenway,

[44424] G. D.; Resnick, D. L.: Plasma cell dyscrasia with polyneuropathy,

[44425] organomegaly, endocrinopathy, M protein, and skin changes: the POEMS

[44426] syndrome. Report on two cases and a review of the literature. Medicine 59:

[44427] 311-322, 1980.

[44428] 6698.

[44429] Frangiskakis, J. M.; Ewart, A. K.; Morris, C. A.; Mervis, C. B.;

[44430] Bertrand, J.; Robinson, B. F.; Klein, B. P.; Ensing, G. J.; Everett,

[44431] L. A.; Green, E. D.; Proschel, C.; Gutowski, N. J.; Noble, M.; Atkinson,

[44432] D. L.; Odelberg, S. J.; Keating, M. T.: LIM-kinase1 hemizygosity

[44433] implicated in impaired visuospatial constructive cognition. Cell 86:

[44434] 59-69, 1996.

[44435] 6699.

[44436] Votruba, M.; Moore, A. T.; Bhattacharya, S. S.: Demonstra-
tion
[44437] of a founder effect and fine mapping of dominant optic
atrophy locus
[44438] on 3q28-qter by linkage disequilibrium method: a study
of 38 British
[44439] Isles pedigrees. Hum. Genet. 102: 79-86, 1998.

[44440] 6700.

[44441] Alhonen-Hongisto, L.; Leinonen, P.; Sinervirta, R.; Laine,
R.;
[44442] Winqvist, R.; Alitalo, K.; Janne, O. A.; Janne, J.: Mouse and
human
[44443] ornithine decarboxylase genes: methylation polymorphism
and amplification. Biochem.
[44444] J. 242: 205-210, 1987.

[44445] 6701.

[44446] Brabant, M.; McConlogue, L.; van Daalen Wetters, T.;
Coffino, P.
[44447] : Mouse ornithine decarboxylase gene: cloning, structure,
and expression. Proc.

[44448] Nat. Acad. Sci. 85: 2200-2204, 1988.

[44449] 6702.

[44450] Cox, D. R.; Trouillot, T.; Ashley, P. L.; Brabant, M.; Coffino,

[44451] P.: A functional mouse ornithine decarboxylase gene (Odc) maps to

[44452] chromosome 12: further evidence of homoeology between mouse chromosome

[44453] 12 and the short arm of human chromosome 2. Cytogenet. Cell Genet. 48:

[44454] 92-94, 1988.

[44455] 6703.

[44456] Kasid, U.; Pfeifer, A.; Weichselbaum, R. R.; Dritschilo, A.; Mark,

[44457] G. E.: The raf oncogene is associated with a radiation-resistant

[44458] human laryngeal cancer. Science 237: 1039-1041, 1987.

[44459] 6704.

[44460] Bandmann, O.; Davis, M. B.; Marsden, C. D.; Wood, N. W.: The human

[44461] homologue of the weaver mouse gene in familial and sporadic Parkinson's

[44462] disease. Neuroscience 72: 877-879, 1996.

[44463] 6705.

[44464] Cohn, J. A.; Friedman, K. J.; Noone, P. G.; Knowles, M. R.; Silverman,

[44465] L. M.; Jowell, P. S.: Relation between mutations of the cystic fibrosis

[44466] gene and idiopathic pancreatitis. New Eng. J. Med. 339: 653-658,

[44467] 1998.

[44468] 6706.

[44469] Trent, J. M.; Rosenfeld, S. B.; Meyskens, F. L.: Chromosome 6q

[44470] involvement in human malignant melanoma. Cancer Genet. Cytogenet. 9:

[44471] 177-180, 1983.

[44472] 6707.

[44473] Puig, S.; Ruiz, A.; Lazaro, C.; Castel, T.; Lynch, M.; Palou,

[44474] J.; Vilalta, A.; Weissenbach, J.; Mascaro, J.-M.; Estivill, X.: Chromosome

[44475] 9p deletions in cutaneous malignant melanoma tumors: the minimal deleted

[44476] region involves markers outside the p16 (CDKN2) gene. Am. J. Hum.

[44477] Genet. 57: 395-402, 1995.

[44478] 6708.

[44479] Briggs, R. C.; Briggs, J. A.; Ozer, J.; Sealy, L.; Dworkin, L.

[44480] L.; Kingsmore, S. F.; Seldin, M. F.; Kaur, G. P.; Athwal, R. S.; Dessypris,

[44481] E. N.: The human myeloid cell nuclear differentiation anti-gen gene

[44482] is one of at least two related interferon-inducible genes located

[44483] on chromosome 1q that are expressed specifically in hematopoietic

[44484] cells.. Blood 83: 2153-2162, 1994.

[44485] 6709.

[44486] Daheron, L.; Veinstein, A.; Brizard, F.; Drabkin, H.; Lacotte,

[44487] L.; Guilhot, F.; Larsen, C. J.; Brizard, A.; Roche, J.: Human LPP

[44488] gene is fused to MLL in a secondary acute leukemia with a t(3;11)(q28;q23). Genes

[44489] Chromosomes Cancer 31: 382-389, 2001.

[44490] 6710.

[44491] Domer, P. H.; Fakharzadeh, S. S.; Chen, C.-S.; Jockel, J.; Johansen,

[44492] L.; Silverman, G. A.; Kersey, J. H.; Korsmeyer, S. J.: Acute mixed-lineage

[44493] leukemia t(4;11)(q21;q23) generates an MLL-AF4 fusion product. Proc.

[44494] Nat. Acad. Sci. 90: 7884-7888, 1993.

[44495] 6711.

[44496] Gu, Y.; Nakamura, T.; Alder, H.; Prasad, R.; Canaani, O.; Cimino,

[44497] G.; Croce, C. M.; Canaani, E.: The t(4;11) chromosome translocation

[44498] of human acute leukemias fuses the ALL-1 gene, related to Drosophila

[44499] trithorax, to the AF-4 gene. Cell 71: 701-708, 1992.

[44500] 6712.

[44501] Megonigal, M. D.; Rappaport, E. F.; Jones, D. H.; Williams, T.

[44502] M.; Lovett, B. D.; Kelly, K. M.; Lerou, P. H.; Moulton, T.; Budarf,

[44503] M. L.; Felix, C. A.: t(11;22)(q23;q11.2) in acute myeloid leukemia

[44504] of infant twins fuses MLL with hCDCrel, a cell division cycle gene

[44505] in the genomic region of deletion in DiGeorge and velocardiofacial

[44506] syndromes. Proc. Nat. Acad. Sci. 95: 6413-6418, 1998.

[44507] 6713.

[44508] Nakamura, T.; Alder, H.; Gu, Y.; Prasad, R.; Canaani, O.; Kamada,

[44509] N.; Gale, R. P.; Lange, B.; Crist, W. M.; Nowell, P. C.; Croce, C.

[44510] M.; Canaani, E.: Genes on chromosomes 4, 9, and 19 involved in 11q23

[44511] abnormalities in acute leukemia share sequence homology and/or common

[44512] motifs. Proc. Nat. Acad. Sci. 90: 4631-4635, 1993.

[44513] 6714.

[44514] Raffini, L. J.; Slater, D. J.; Rappaport, E. F.; Lo Nigro, L.;

[44515] Cheung, N.-K. V.; Biegel, J. A.; Nowell, P. C.; Lange, B. J.; Felix,

[44516] C. A.: Panhandle and reverse-panhandle PCR enable cloning of der(11)

[44517] and der(other) genomic breakpoint junctions of MLL translocations

[44518] and identify complex translocation of MLL, AF-4, and CDK6. Proc.

[44519] Nat. Acad. Sci. 99: 4568-4573, 2002.

[44520] 6715.

[44521] Hickok, N. J.; Seppanen, P. J.; Gunsalus, G. L.; Janne, O. A.:

[44522] Complete amino acid sequence of human ornithine decarboxylase deduced

[44523] from complementary DNA. DNA 6: 179-187, 1987.

[44524] 6716.

[44525] Michalak, M.; Quackenbush, E. J.; Letarte, M.: Inhibition of

[44526] Na+/Ca2+ exchanger activity in cardiac and skeletal muscle sarcolemmal

[44527] vesicles by monoclonal antibody 44D7. J. Biol. Chem. 261: 92-95,

[44528] 1986.

[44529] 6717.

[44530] Posillico, J. T.; Srikanta, S.; Brown, E. M.; Eisenbarth, G. S.

[44531] : The 4F2 cell surface protein modulates intracellular calcium. (Abstract) Clin.

[44532] Res. 33: 385A only, 1985.

[44533] 6718.

[44534] Posillico, J. T.; Srikanta, S.; Eisenbarth, G.; Quaranta, V.;

[44535] Kajiji, S.; Brown, E. M.: Binding of monoclonal antibody (4F2) to

[44536] its cell surface antigen on dispersed adenomatous parathyroid cells

[44537] raises cytosolic calcium and inhibits parathyroid hormone secretion. J.

[44538] Clin. Endocr. Metab. 64: 43-50, 1987.

[44539] 6719.

[44540] Quackenbush, E.; Clabby, M.; Gottesdiener, K. M.; Barbosa, J.;

[44541] Jones, N. H.; Strominger, J. L.; Speck, S.; Leiden, J. M.: Molecular

[44542] cloning of complementary DNAs encoding the heavy chain of the human

[44543] 4F2 cell-surface antigen: a type II membrane glycoprotein

involved
[44544] in normal and neoplastic cell growth. Proc. Nat. Acad. Sci. 84:
[44545] 6526-6530, 1987. Note: Correction: Proc. Nat. Acad. Sci. 84: 6526-6530,
[44546] 1987.

[44547] 6720.

[44548] McQuibban, G. A.; Gong, J.-H.; Tam, E. M.; McCulloch, C. A. G.;
[44549] Clark-Lewis, I.; Overall, C. M.: Inflammation dampened by gelatinase
[44550] A cleavage of monocyte chemoattractant protein-3. Science 289: 1202-1206,
[44551] 2000.

[44552] 6721.

[44553] Fu, Y.-H.; Pizzuti, A.; Fenwick, R. G., Jr.; King, J.; Rajnarayan,
[44554] S.; Dunne, P. W.; Dubel, J.; Nasser, G. A.; Ashizawa, T.; de Jong,
[44555] P.; Wieringa, B.; Korneluk, R.; Perryman, M. B.; Epstein, H. F.; Caskey,
[44556] C. T.: An unstable triplet repeat in a gene related to my-

otonic muscular

[44557] dystrophy. Science 255: 1256-1258, 1992.

[44558] 6722.

[44559] Marsden, P. A.; Heng, H. H. Q.; Scherer, S. W.; Stewart, R. J.;

[44560] Hall, A. V.; Shi, X.-M.; Tsui, L.-C.; Schappert, K. T.: Structure

[44561] and chromosomal localization of the human constitutive endothelial

[44562] nitric oxide synthase gene. J. Biol. Chem. 268: 17478-17488, 1993.

[44563] 6723.

[44564] Miyamoto, Y.; Saito, Y.; Nakayama, M.; Shimasaki, Y.; Yoshimura,

[44565] T.; Yoshimura, M.; Harada, M.; Kajiyama, N.; Kishimoto, I.; Kuwahara,

[44566] K.; Hino, J.; Ogawa, E.; Hamanaka, I.; Kamitani, S.; Takahashi, N.;

[44567] Kawakami, R.; Kangawa, K.; Yasue, H.; Nakao, K.: Replication protein

[44568] A1 reduces transcription of the endothelial nitric oxide synthase

[44569] gene containing a -786T-C mutation associated with coronary spastic
[44570] angina. Hum. Molec. Genet. 9: 2629-2637, 2000.

[44571] 6724.

[44572] Nakayama, M.; Yasue, H.; Yoshimura, M.; Shimasaki, Y.; Kugiyama,
[44573] K.; Ogawa, H.; Motoyama, T.; Saito, Y.; Ogawa, Y.; Miyamoto, Y.; Nakao,
[44574] K.: T(-786)-C mutation in the 5-prime-flanking region of the endothelial
[44575] nitric oxide synthase gene is associated with coronary spasm. Circulation 99:
[44576] 2864-2870, 1999.

[44577] 6725.

[44578] Persu, A.; Stoenoiu, M. S.; Messiaen, T.; Davila, S.; Robino,
[44579] C.; El-Khattabi, O.; Mourad, M.; Horie, S.; Feron, O.; Balli-gand,
[44580] J.-L.; Wattiez, R.; Pirson, Y.; Chaveau, D.; Lens, X. M.; De-vuyst,
[44581] O.: Modifier effect of ENOS in autosomal dominant poly-cystic kidney

[44582] disease. Hum. Molec. Genet. 11: 229-241, 2002.

[44583] 6726.

[44584] Bloch, K. D.; Wolfram, J. R.; Brown, D. M.; Roberts, J. D., Jr.;

[44585] Zapol, D. G.; Lepore, J. J.; Filippov, G.; Thomas, J. E.; Jacob, H.

[44586] J.; Bloch, D. B.: Three members of the nitric oxide synthase II gene

[44587] family (NOS2A, NOS2B, and NOS2C) colocalize to human chromosome 17. Genomics 27:

[44588] 526-530, 1995.

[44589] 6727.

[44590] Chartrain, N. A.; Geller, D. A.; Koty, P. P.; Sitrin, N. F.; Nussler,

[44591] A. K.; Hoffman, E. P.; Billiar, T. R.; Hutchinson, N. I.; Mudgett,

[44592] J. S.: Molecular cloning, structure, and chromosomal localization

[44593] of the human inducible nitric oxide synthase gene. J. Biol. Chem. 269:

[44594] 6765-6772, 1994.

[44595] 6728.

[44596] Choi, H.-S.; Rai, P. R.; Chu, H. W.; Cool, C.; Chan, E. D.: Analysis

[44597] of nitric oxide synthase and nitrotyrosine expression in human pulmonary

[44598] tuberculosis. Am. J. Resp. Crit. Care Med. 166: 178-186, 2002.

[44599] 6729.

[44600] Facchetti, F.; Vermi, W.; Fiorentini, S.; Chilosi, M.; Caruso,

[44601] A.; Duse, M.; Notarangelo, L. D.; Badolato, R.: Expression of inducible

[44602] nitric oxide synthase in human granulomas and histiocytic reactions. Am.

[44603] J. Path. 154: 145-152, 1999.

[44604] 6730.

[44605] Pyronnet, S.; Pradayrol, L.; Sonenberg, N.: A cell cycle-dependent

[44606] internal ribosome entry site. Molec. Cell 5: 607-616, 2000.

[44607] 6731.

[44608] Radford, D. M.; Nakai, H.; Byers, M. G.; Eddy, R. L.; Haley,

L.

[44609] L.; Henry, W. M.; Shows, T. B.: Mapping the ornithine de-
carboxylase

[44610] gene (ODC1 and ODC2) to 2p25 and 7q31-qter, respec-
tively. (Abstract) Cytogenet.

[44611] Cell Genet. 46: 678 only, 1987.

[44612] 6732.

[44613] Villani, V.; Coffino, P.; D'Eustachio, P.: Linkage genetics of

[44614] mouse ornithine decarboxylase (Odc). Genomics 5:
636-638, 1989.

[44615] 6733.

[44616] Winqvist, R.; Makela, T. P.; Seppanen, P.; Janne, O. A.; Al-
honen-Hongisto,

[44617] L.; Janne, J.; Grzeschik, K.-H.; Alitalo, K.: Human ornithine
decarboxylase

[44618] sequences map to chromosome regions 2pter-p23 and
7cen-qter but are

[44619] not coamplified with the NMYC oncogene. Cytogenet. Cell
Genet. 42:

[44620] 133-140, 1986.

[44621] 6734.

[44622] Yang-Feng, T. L.; Barton, D. E.; Thelander, L.; Lewis, W. H.; Srinivasan,
[44623] P. R.; Francke, U.: Ribonucleotide reductase M2 subunit sequences
[44624] mapped to four different chromosomal sites in humans and mice: functional
[44625] locus identified by its amplification in hydroxyurea-resistant cell
[44626] lines. Genomics 1: 77-86, 1987.

[44627] 6735.

[44628] Clark, C. C.; Cohen, I.; Eichstetter, I.; Cannizzaro, L. A.; McPherson,
[44629] J. D.; Wasmuth, J. J.; Iozzo, R. V.: Molecular cloning of the human
[44630] proto-oncogene Wnt-5A and mapping of the gene (WNT5A) to chromosome
[44631] 3p14-p21. Genomics 18: 249-260, 1993.

[44632] 6736.

[44633] He, X.; Saint-Jeannet, J.-P.; Wang, Y.; Nathans, J.; Dawid, I.;
[44634] Varmus, H.: A member of the frizzled protein family mediating axis

[44635] induction by Wnt-5A. Science 275: 1652-1654, 1997.

[44636] 6737.

[44637] Adelaide, J.; Mattei, M.-G.; Marics, I.; Raybaud, F.; Planche,

[44638] J.; De Lapeyriere, O.; Birnbaum, D.: Chromosomal local-
ization of

[44639] the hst oncogene and its co-amplification with the int.2
oncogene

[44640] in a human melanoma. Oncogene 2: 413-416, 1988.

[44641] 6738.

[44642] Dudley, A. T.; Ros, M. A.; Tabin, C. J.: A re-examination of
proximodistal

[44643] patterning during vertebrate limb development. Nature
418: 539-544,

[44644] 2002.

[44645] 6739.

[44646] Feldman, B.; Poueymirou, W.; Papaionnou, V. E.; DeChiara,
T. M.;

[44647] Goldfarb, M.: Requirement of FGF-4 for postimplantation
mouse development. Science 267:

[44648] 246-249, 1995.

[44649] 6740.

[44650] Huebner, K.; Ferrari, A. C.; Delli Bovi, P.; Croce, C. M.; Basilico,

[44651] C.: The FGF-related oncogene, K-FGF, maps to human chromosome region

[44652] 11q13, possibly near int-2. Oncogene Res. 3: 263-270, 1988.

[44653] 6741.

[44654] Sakamoto, H.; Mori, M.; Taira, M.; Yoshida, T.; Matsukawa, S.;

[44655] Shimizu, K.; Sekiguchi, M.; Terada, M.; Sugimura, T.: Transforming

[44656] gene from human stomach cancers and a noncancerous portion of stomach

[44657] mucosa. Proc. Nat. Acad. Sci. 83: 3997-4001, 1986.

[44658] 6742.

[44659] Sun, X.; Lewandoski, M.; Meyers, E. N.; Liu, Y.-H.; Maxson, R.

[44660] E., Jr.; Martin, G. R.: Conditional inactivation of Fgf4 reveals

[44661] complexity of signalling during limb bud development. Nature Genet. 25:

[44662] 83-86, 2000.

[44663] 6743.

[44664] Sun, X.; Mariani, F. V.; Martin, G. R.: Functions of FGF sig-
nalling
[44665] from the apical ectodermal ridge in limb development.
Nature 418:
[44666] 501-508, 2002.

[44667] 6744.

[44668] Dusetti, N. J.; Frigerio, J.-M.; Fox, M. F.; Swallow, D. M.;
Dagorn,
[44669] J.-C.; Iovanna, J. L.: Molecular cloning, genomic organiza-
tion, and
[44670] chromosomal localization of the human pancreatitis-asso-
ciated protein
[44671] (PAP) gene. Genomics 19: 108-114, 1994.

[44672] 6745.

[44673] Dusetti, N. J.; Frigerio, J.-M.; Keim, V.; Dagorn, J.-C.; Io-
vanna,
[44674] J. L.: Structural organization of the gene encoding the rat
pancreatitis-associated
[44675] protein: analysis of its evolutionary history reveals an an-

cient divergence
[44676] from the other carbohydrate-recognition domain-containing genes. J.
[44677] Biol. Chem. 268: 14470-14475, 1993.

[44678] 6746.

[44679] Dusetti, N. J.; Frigerio, J. M.; Szpirer, C.; Dagorn, J. C.; Iovanna,
[44680] J. L.: Cloning, expression and chromosomal localization of the rat
[44681] pancreatitis-associated protein III gene. Biochem. J. 307: 9-16,
[44682] 1995.

[44683] 6747.

[44684] Frigerio, J.-M.; Dusetti, N. J.; Garrido, P.; Dagorn, J.-C.; Iovanna,
[44685] J. L.: The pancreatitis associated protein III (PAP III), a new member
[44686] of the PAP gene family. Biochim. Biophys. Acta 1216: 329-331, 1993.
[44687] 6748.

[44688] Frigerio, J.-M.; Dusetti, N. J.; Keim, V.; Dagorn, J.-C.; Io-

vanna,

[44689] J. L.: Identification of a second rat pancreatitis-associated protein:

[44690] messenger RNA cloning, gene structure, and expression during acute

[44691] pancreatitis. Biochemistry 32: 9236-9241, 1993.

[44692] 6749.

[44693] Itoh, T.; Sawabu, N.; Motoo, Y.; Funakoshi, A.; Teraoka, H.: The

[44694] human pancreatitis-associated protein (PAP)-encoding gene generates

[44695] multiple transcripts through alternative use of 5-prime exons. Gene 155:

[44696] 283-287, 1995.

[44697] 6750.

[44698] Lasserre, C.; Simon, M. T.; Ishikawa, H.; Diriong, S.; Nguyen,

[44699] V. C.; Christa, L.; Vernier, P.; Brechot, C.: Structural orga-nization

[44700] and chromosomal localization of a human gene (HIP/PAP) encoding a

[44701] C-type lectin overexpressed in primary liver cancer. Eu-

rop. J. Biochem. 224:

[44702] 29-38, 1994.

[44703] 6751.

[44704] Orelle, B.; Keim, V.; Masciotra, L.; Dagorn, J.-C.; Iovanna,
J.-L.

[44705] : Human pancreatitis-associated protein: messenger RNA
cloning and

[44706] expression in pancreatic diseases. J. Clin. Invest. 90:
2284-2291,

[44707] 1992.

[44708] 6752.

[44709] Vasseur, S.; Frigerio, J. M.; Dusetti, N. J.; Keim, V.; Dagorn,

[44710] J. C.; Iovanna, J. L.: Two transcripts are generated from the
pancreatitis

[44711] associated protein II gene by alternative splicing in the
5-prime

[44712] untranslated region. Biochim. Biophys. Acta 1261:
272-274, 1995.

[44713] 6753.

[44714] Julier, C.; Lathrop, M.; Lalouel, J. M.; Kaplan, J. C.: Use of

[44715] multilocus tests of gene order: example for chromosome

22. (Abstract) Cytogenet.

[44716] Cell Genet. 40: 663-664, 1985.

[44717] 6754.

[44718] Julier, C.; Lathrop, M.; Lalouel, J. M.; Reghis, A.; Szajnert,

[44719] M. F.; Kaplan, J. C.: New restriction fragment length poly-
morphisms

[44720] on human chromosome 22 at loci SIS, MB and IGLV.
(Abstract) Cytogenet.

[44721] Cell Genet. 40: 664 only, 1985.

[44722] 6755.

[44723] Julier, C.; Reghis, A.; Szajnert, M. F.; Kaplan, J. C.; Lathrop,

[44724] G. M.; Lalouel, J. M.: A preliminary linkage map of human
chromosome

[44725] 22. (Abstract) Cytogenet. Cell Genet. 40: 665 only, 1985.

[44726] 6756.

[44727] Nakamura, Y.; Leppert, M.; O'Connell, P.; Wolff, R.; Holm,
T.;

[44728] Culver, M.; Martin, C.; Fujimoto, E.; Hoff, M.; Kumlin, E.;
White,

[44729] R.: Variable number of tandem repeat (VNTR) markers for

human gene
[44730] mapping. Science 235: 1616-1622, 1987.

[44731] 6757.

[44732] Romero-Herrera, A. E.; Lehmann, H.: The amino acid se-
quence of
[44733] human myoglobin and its minor fractions. Proc. Roy. Soc.
London 186B:
[44734] 249-279, 1974.

[44735] 6758.

[44736] Romero-Herrera, A. E.; Lehmann, H.: Primary structure of
human
[44737] myoglobin. Nature N.B. 232: 149-152, 1971.

[44738] 6759.

[44739] Weller, P.; Jeffreys, A. J.; Wilson, V.; Blanchetot, A.: Organi-
zation
[44740] of the human myoglobin gene. EMBO J. 3: 439-446, 1984.

[44741] 6760.

[44742] Wyman, A. R.; White, R.: A highly polymorphic locus in hu-
man
[44743] DNA. Proc. Nat. Acad. Sci. 77: 6754-6758, 1980.

[44744] 6761.

[44745] Chamberlain, J. S.; vanTuinen, P.; Reeves, A. A.; Philip, B. A.;

[44746] Caskey, C. T.: Isolation of cDNA clones for the catalytic gamma subunit

[44747] of mouse muscle phosphorylase kinase: expression of mRNA in normal

[44748] and mutant Phk mice. Proc. Nat. Acad. Sci. 84: 2886-2890, 1987.

[44749] 6762.

[44750] Jones, T. A.; da Cruz e Silva, E. F.; Spurr, N. K.; Sheer, D.;

[44751] Cohen, P. T.: Localization of the gene encoding the cat- alytic gamma

[44752] subunit of phosphorylase kinase to human chromosome bands 7p12-q21. Biochim.

[44753] Biophys. Acta 1048: 24-29, 1990.

[44754] 6763.

[44755] Maichele, A. J.; Chamberlain, J. S.: The gamma phosphory- lase kinase

[44756] gene, Phkg, maps to mouse chromosome 5 near Gus. Mammalian Genome 5:

[44757] 15-18, 1994.

[44758] 6764.

[44759] Wehner, M.; Kilimann, M. W.: Human cDNA encoding the muscle isoform

[44760] of the phosphorylase kinase gamma subunit (PHKG1). Hum. Genet. 96:

[44761] 616-618, 1995.

[44762] 6765.

[44763] Burwinkel, B.; Shiomi, S.; Al Zaben, A.; Kilimann, M. W.: Liver

[44764] glycogenosis due to phosphorylase kinase deficiency: PHKG2 gene structure

[44765] and mutations associated with cirrhosis. Hum. Molec. Genet. 7: 149-154,

[44766] 1998.

[44767] 6766.

[44768] Burwinkel, B.; Tanner, M. S.; Kilimann, M. W.: Phosphory-lase kinase

[44769] deficient liver glycogenosis: progression to cirrhosis in in-fancy

[44770] associated with PHKG2 mutations (H144Y and L225R).

(Letter) J. Med.

[44771] Genet. 37: 376-377, 2000.

[44772] 6767.

[44773] Clark, D.; Haynes, D.: The glycogen storage disease (gsd/gsd)

[44774] rat. Curr. Top. Cell. Regul. 29: 217-263, 1988.

[44775] 6768.

[44776] Hanks, S. K.: Messenger ribonucleic acid encoding an apparent

[44777] isoform of phosphorylase kinase catalytic subunit is abundant in the

[44778] adult testis. Molec. Endocr. 3: 110-116, 1989.

[44779] 6769.

[44780] Maichele, A. J.; Burwinkel, B.; Maire, I.; Sovik, O.; Kilimann,

[44781] M. W.: Mutations in the testis/liver isoform of the phosphorylase

[44782] kinase gamma subunit (PHKG2) cause autosomal liver glycogenosis in

[44783] the gsd rat and in humans. Nature Genet. 14: 337-340, 1996.

[44784] 6770.

[44785] Sovik, O.; deBarsy, T.; Maehle, B.: Phosphorylase kinase deficiency:
[44786] severe glycogen storage disease with evidence of autosomal recessive
[44787] mode of inheritance. (Letter) Europ. J. Pediat. 139: 210 only, 1982.
[44788] 6771.

[44789] Almarri, A.; Batchelor, J. R.: HLA and hepatitis B infection. Lancet 344:
[44790] 1194-1195, 1994.

[44791] 6772.

[44792] Ayala, F. J.; Escalante, A.; O'hUigin, C.; Klein, J.: Molecular

[44793] genetics of speciation and human origins. Proc. Nat. Acad. Sci. 91:
[44794] 6787-6794, 1994.

[44795] 6773.

[44796] Bell, J. I.; Estess, P.; St. John, T.; Saiki, R.; Watling, D. L.;

[44797] Erlich, H. A.; McDevitt, H. O.: DNA sequence and characterization
[44798] of human class II major histocompatibility complex beta chains from

[44799] the DR1 haplotype. Proc. Nat. Acad. Sci. 82: 3405-3409, 1985.

[44800] 6774.

[44801] Bodmer, J. G.; Marsh, S. G. E.; Albert, E.: Nomenclature for factors

[44802] of the HLA system, 1989. Immun. Today 11: 3-10, 1990.

[44803] 6775.

[44804] Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.; Bontrop,

[44805] R. E.; Charron, D.; Dupont, B.; Erlich, H. A.; Fauchet, R.; Mach,

[44806] B.; Mayr, W. R.; Parham, P.; Sasazuki, T.; Schreuder, G. M. T.; Strominger,

[44807] J. L.; Svejgaard, A.; Terasaki, P. I.: Nomenclature for factors of

[44808] the HLA system, 1996. Europ. J. Immunogenet. 24: 105-151, 1997.

[44809] 6776.

[44810] Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.; Dupont,

[44811] B.; Erlich, H. A.; Mach, B.; Mayr, W. R.; Parham, P.; Sasazuki, T.;

[44812] Schreuder, G. M. T.; Strominger, J. L.; Svejgaard, A.; Terasaki, P.

[44813] I.: Nomenclature for factors of the HLA system, 1994. Tissue Antigens 44:

[44814] 1-18, 1994.

[44815] 6777.

[44816] Bodmer, W. F.; Bodmer, J. G.; Batchelor, J. R.; Festenstein, H.;

[44817] Morris, P. J.: Histocompatibility Testing 1977. Copenhagen: Munksgaard

[44818] (pub.) 1978.

[44819] 6778.

[44820] Boss, J. M.; Strominger, J. L.: Cloning and sequence analysis

[44821] of the human major histocompatibility complex gene DC-3-beta. Proc.

[44822] Nat. Acad. Sci. 81: 5199-5203, 1984.

[44823] 6779.

[44824] Charron, D. J.; McDevitt, H. O.: Analysis of HLA-D region-associated

[44825] molecules with monoclonal antibody. Proc. Nat. Acad. Sci.

76: 6567-6571,

[44826] 1979.

[44827] 6780.

[44828] Chelladurai, M.; Honn, K. V.; Walz, D. A.: HLA-DR is a pro-coagulant. Biochem.

[44829] Biophys. Res. Commun. 178: 467-473, 1991.

[44830] 6781.

[44831] Corte, G.; Damiani, G.; Calabi, F.; Fabbi, M.; Bargellesi, A.

[44832] : Analysis of HLA-DR polymorphism by two-dimensional peptide mapping. Proc.

[44833] Nat. Acad. Sci. 78: 534-538, 1981.

[44834] 6782.

[44835] Das, H. K.; Lawrance, S. K.; Weissman, S. M.: Structure and nucleotide

[44836] sequence of the heavy chain gene of HLA-DR. Proc. Nat. Acad. Sci. 80:

[44837] 3543-3547, 1983.

[44838] 6783.

[44839] Delovitch, T. L.; Falk, J. A.: Evidence for structural homology

[44840] between murine and human Ia antigens. Immunogenetics 8: 405-418,

[44841] 1979.

[44842] 6784.

[44843] Dunham, I.; Sargent, C. A.; Dawkins, R. L.; Campbell, R. D.:

[44844] An analysis of variation in the long-range genomic organization of

[44845] the human major histocompatibility complex class II region by pulsed-field

[44846] gel electrophoresis. Genomics 5: 787-796, 1989.

[44847] 6785.

[44848] Erlich, H. A.; Stetler, D.; Saiki, R.; Gladstone, P.; Pious, D.

[44849] : Mapping of the genes encoding the HLA-DR alpha chain and the HLA-related

[44850] antigens to a chromosome 6 deletion by using genomic blotting. Proc.

[44851] Nat. Acad. Sci. 80: 2300-2304, 1983.

[44852] 6786.

[44853] Ferber, K. M.; Keller, E.; Albert, E. D.; Ziegler, A.-G.: Predictive

[44854] value of human leukocyte antigen class II typing for the development

[44855] of islet autoantibodies and insulin-dependent diabetes postpartum

[44856] in women with gestational diabetes. J. Clin. Endocr. Metab. 84:

[44857] 2342-2348, 1999.

[44858] 6787.

[44859] Fuller, T. C.; Einarson, M.; Pinto, C.; Ahern, A.; Yunis, E. J.

[44860] : Genetic evidence that HLA-DR (Ia) specifications include multiple

[44861] HLA-D determinants on a single haplotype. Transplant. Proc. 10:

[44862] 781-784, 1978.

[44863] 6788.

[44864] Hardy, D. A.; Bell, J. I.; Long, E. O.; Lindsten, T.; McDevitt,

[44865] H. O.: Mapping of the class II region of the human major

histocompatibility
[44866] complex by pulsed-field gel electrophoresis. Nature 323: 453-455,
[44867] 1986.

[44868] 6789.

[44869] Hui, K.; Festenstein, H.; de Klein, A.; Grosveld, G.; Grosveld,
[44870] F.: HLA-DR genotyping by restriction fragment length polymorphism
[44871] analyses. Immunogenetics 22: 231-239, 1985.

[44872] 6790.

[44873] Kaufman, J. F.; Strominger, J. L.: HLA-DR light chain has a polymorphic
[44874] N-terminal region and a conserved immunoglobulin-like C-terminal region. Nature 297:
[44875] 694-697, 1982.

[44876] 6791.

[44877] Korman, A. J.; Auffray, C.; Schamboeck, A.; Strominger, J. L.
[44878] : The amino acid sequence and gene organization of the heavy chain

[44879] of the HLA-DR antigen: homology to immunoglobulins. Proc. Nat. Acad.
[44880] Sci. 79: 6013-6017, 1982.

[44881] 6792.

[44882] Korman, A. J.; Knudsen, P. J.; Kaufman, J. F.; Strominger, J.

[44883] L.: cDNA clones for the heavy chain of HLA-DR antigens obtained after
[44884] immunopurification of polysomes by monoclonal antibody. Proc. Nat.
[44885] Acad. Sci. 79: 1844-1848, 1982.

[44886] 6793.

[44887] Kratzin, H.; Yang, C.; Gotz, H.; Pauly, E.; Kolbel, S.; Egert,

[44888] G.; Thinnes, F. P.; Wernet, P.; Altevogt, P.; Hilschmann, N.: Primarstruktur
[44889] menschlicher Histokompatibilitatsantigene der Klasse II. 1. Mitteilung:
[44890] aminosauresequenz der N-terminalen 198 Reste der beta-Kette des HLA-Dw2,2;DR2,2-Alloantigens. Hoppe
[44891] Seylers Z. Physiol. Chem. 362: 1665-1669, 1981.

[44892] 6794.

[44893] Lamm, L. U.: Another segregant series, DR, in HLA. Cyto-
genet.
[44894] Cell Genet. 22: 309-312, 1978.

[44895] 6795.

[44896] Larhammar, D.; Schenning, L.; Gustafsson, K.; Wiman, K.;
Claesson,
[44897] L.; Rask, L.; Peterson, P. A.: Complete amino acid se-
quence of an
[44898] HLA-DR antigen-like beta chain as predicted from the nu-
cleotide sequence:
[44899] similarities with immunoglobulins and HLA-A, -B, and -C
antigens. Proc.
[44900] Nat. Acad. Sci. 79: 3687-3691, 1982.

[44901] 6796.

[44902] Lee, J. S.; Trowsdale, J.; Bodmer, W. F.: cDNA clones cod-
ing
[44903] for the heavy chain of human HLA-DR antigen. Proc. Nat.
Acad. Sci. 79:
[44904] 545-549, 1982.

[44905] 6797.

[44906] Levine, F.; Mach, B.; Long, E.; Erlich, H.; Pious, D.: Mapping

[44907] in the HLA-D region with deletion variants and cloned genes. (Abstract) Cytogenet.
[44908] Cell Genet. 37: 523 only, 1984.

[44909] 6798.

[44910] Mann, D. L.; Abelson, L.; Harris, S. D.; Amos, D. B.: Second

[44911] genetic locus in the HLA region for human B-cell alloantigens. Nature 259:
[44912] 145-146, 1976.

[44913] 6799.

[44914] Mann, D. L.; Abelson, L.; Henkart, P.; Harris, S. D.; Amos, D.
[44915] B.: Specific B-lymphocyte alloantigens linked to HLA. Proc. Nat.
[44916] Acad. Sci. 72: 5103-5106, 1975.

[44917] 6800.

[44918] Markert, M. L.; Cresswell, P.: Polymorphism of human B-cell alloantigens:
[44919] evidence for three loci within the HLA system. Proc. Nat. Acad. Sci. 77:
[44920] 6101-6104, 1980.

[44921] 6801.

[44922] Mayer, W. E.; O'hUigin, C.; Klein, J.: Resolution of the HLA-DRB6

[44923] puzzle: a case of grafting a de novo-generated exon on an existing

[44924] gene. Proc. Nat. Acad. Sci. 90: 10720-10724, 1993.

[44925] 6802.

[44926] McDevitt, H. O.; Bodmer, W. F.: HLA, immune-response genes, and

[44927] disease. Lancet I: 1269-1275, 1974.

[44928] 6803.

[44929] Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Mapping

[44930] of the genes for LYAM1, a new lymphocyte adhesion molecule, and for

[44931] LAR, a new receptor-linked protein tyrosine phosphatase, to human

[44932] chromosome 1. (Abstract) Cytogenet. Cell Genet. 51: 990 only, 1989.

[44933] 6804.

[44934] Pentchev, P. G.; Comly, M. E.; Kruth, H. S.; Patel, S.; Proes-

tel,
[44935] M.; Weintroub, H.: The cholesterol storage disorder of the mutant
[44936] BALB/c mouse: a primary genetic lesion closely linked to defective
[44937] esterification of exogenously derived cholesterol and its relationship
[44938] to human type C Niemann-Pick disease. J. Biol. Chem. 261: 2772-2777,
[44939] 1986.

[44940] 6805.

[44941] Pentchev, P. G.; Comly, M. E.; Kruth, H. S.; Vanier, M. T.; Wenger,
[44942] D. A.; Patel, S.; Brady, R. O.: A defect in cholesterol esterification
[44943] in Niemann-Pick disease (type C) patients. Proc. Nat. Acad. Sci. 82:
[44944] 8247-8251, 1985.

[44945] 6806.

[44946] Pentchev, P. G.; Comly, M. E.; Vanier, M. T.; Kruth, H. S.; Patel,
[44947] S.; Brady, R. O.: Type C Niemann-Pick disease: a defect in

intracellular

[44948] esterification. (Abstract) Am. J. Hum. Genet. 37: A15, 1985.

[44949] 6807.

[44950] Pentchev, P. G.; Kruth, H. S.; Comly, M. E.; Butler, J. D.; Vanier,

[44951] M. T.; Wenger, D. A.; Patel, S.: Type C Niemann-Pick disease: a parallel

[44952] loss of regulatory responses in both the uptake and esterification

[44953] of low-density lipoprotein-derived cholesterol in cultured fibroblasts. J.

[44954] Biol. Chem. 261: 16775-16780, 1986.

[44955] 6808.

[44956] Ribeiro, I.; Marcao, A.; Amaral, O.; Sa Miranda, M. C.; Vanier,

[44957] M. T.; Millat, G.: Niemann-Pick type C disease: NPC1 mutations associated

[44958] with severe and mild cellular cholesterol trafficking alterations. Hum.

[44959] Genet. 109: 24-32, 2001.

[44960] 6809.

[44961] Roff, C. F.; Goldin, E.; Comly, M. E.; Cooney, A.; Brown, A.;

[44962] Vanier, M. T.; Miller, S. P. F.; Brady, R. O.; Pentchev, P. G.:
Type

[44963] C Niemann-Pick disease: use of hydrophobic amines to
study defective

[44964] cholesterol transport. Dev. Neurosci. 13: 315-319, 1991.

[44965] 6810.

[44966] Sakai, Y.; Miyawaki, S.; Shimizu, A.; Ohno, K.; Watanabe,
T.:

[44967] A molecular genetic linkage map of mouse chromosome
18, including

[44968] spm, Grl-1, Fim-2/c-fms, and Mbp. Biochem. Genet. 29:
103-113, 1991.

[44969] 6811.

[44970] Schedin, S.; Sindelar, P. J.; Pentchev, P.; Brunk, U.; Dallner,

[44971] G.: Peroxisomal impairment in Niemann-Pick type C dis-
ease. J. Biol.

[44972] Chem. 272: 6245-6251, 1997.

[44973] 6812.

[44974] Sokol, J.; Blanchette-Mackie, E. J.; Kruth, H. S.; Dwyer, N.
K.;

[44975] Amende, L. M.; Butler, J. D.; Robinson, E.; Patel, S.; Brady, R. O.;

[44976] Comly, M. E.; Vanier, M. T.; Pentchev, P. G.: Type C Niemann-Pick

[44977] disease: lysosomal accumulation and defective intracellular mobilization

[44978] of low density lipoprotein cholesterol. J. Biol. Chem. 263: 3411-3417,

[44979] 1988.

[44980] 6813.

[44981] Steinberg, S. J.; Ward, C. P.; Fensom, A. H.: Complementation

[44982] studies in Niemann-Pick disease type C indicate the existence of a

[44983] second group. J. Med. Genet. 31: 317-320, 1994.

[44984] 6814.

[44985] Sun, X.; Marks, D. L.; Park, W. D.; Wheatley, C. L.; Puri, V.;

[44986] O'Brien, J. F.; Kraft, D. L.; Lundquist, P. A.; Patterson, M. C.;

[44987] Pagano, R. E.; Snow, K.: Niemann-Pick C variant detection by altered

[44988] sphingolipid trafficking and correlation with mutations within a specific

[44989] domain of NPC1. Am. J. Hum. Genet. 68: 1361-1372, 2001.

[44990] 6815.

[44991] Thomas, G. H.; Tuck-Muller, C. M.; Miller, C. S.; Reynolds, L.

[44992] W.: Correction of sphingomyelinase deficiency in Niemann-Pick type

[44993] C fibroblasts by removal of lipoprotein fraction from culture media. J.

[44994] Inherit. Metab. Dis. 12: 139-151, 1989.

[44995] 6816.

[44996] Turpin, J. C.; Masson, M.; Baumann, N.: Clinical aspects of Niemann-Pick

[44997] type C disease in the adult. Dev. Neurosci. 13: 304-306, 1991.

[44998] 6817.

[44999] Vanier, M. T.; Duthel, S.; Rodriguez-Lafrasse, C.; Pentchev, P.;

[45000] Carstea, E. D.: Genetic heterogeneity in Niemann-Pick C disease:

[45001] a study using somatic cell hybridization and linkage analysis. Am.

[45002] J. Hum. Genet. 58: 118-125, 1996.

[45003] 6818.

[45004] Vanier, M. T.; Rodriguez-Lafrasse, C.; Rousson, R.; Duthel, S.;

[45005] Harzer, K.; Pentchev, P. G.; Revol, A.; Louisot, P.: Type C Niemann-Pick

[45006] disease: biochemical aspects and phenotypic heterogeneity. Dev. Neurosci. 13:

[45007] 307-314, 1991.

[45008] 6819.

[45009] Vanier, M. T.; Wenger, D. A.; Comly, M. E.; Rousson, R.; Brady,

[45010] R. O.; Pentchev, P. G.: Niemann-Pick disease group C: clinical variability

[45011] and diagnosis based on defective cholesterol esterification: a collaborative

[45012] study on 70 patients. Clin. Genet. 33: 331-348, 1988.

[45013] 6820.

[45014] Watari, H.; Blanchette-Mackie, E. J.; Dwyer, N. K.; Glick, J.

[45015] M.; Patel, S.; Neufeld, E. B.; Brady, R. O.; Pentchev, P. G.; Strauss,

[45016] J. F., III: Niemann-Pick C1 protein: obligatory roles for N-terminal

[45017] domains and lysosomal targeting in cholesterol mobilization. Proc.

[45018] Nat. Acad. Sci. 96: 805-810, 1999.

[45019] 6821.

[45020] Xie, C.; Turley, S. D.; Dietschy, J. M.: Cholesterol accumulation

[45021] in tissues of the Niemann-Pick type C mouse is determined by the rate

[45022] of lipoprotein-cholesterol uptake through the coated-pit pathway in

[45023] each organ. Proc. Nat. Acad. Sci. 96: 11992-11997, 1999.

[45024] 6822.

[45025] Yamamoto, T.; Nanba, E.; Ninomiya, H.; Higaki, K.; Taniguchi,

[45026] M.; Zhang, H.; Akaboshi, S.; Watanabe, Y.; Takeshima, T.; Inui, K.;

[45027] Okada, S.; Tanaka, A.; Sakuragawa, N.; Millat, G.; Vanier, M. T.;

[45028] Morris, J. A.; Pentchev, P. G.; Ohno, K.: NPC1 gene mutations in

[45029] Japanese patients with Niemann-Pick disease type C. Hum. Genet. 105:

[45030] 10-16, 1999.

[45031] 6823.

[45032] Wijmenga, C.; Hansen, R. S.; Gimelli, G.; Bjorck, E. J.; Davies,

[45033] E. G.; Valentine, D.; Belohradsky, B. H.; van Dongen, J. J.; Smeets,

[45034] D. F. C. M.; van den Heuvel, L. P. W. J.; Luyten, J. A. F. M.; Strengman,

[45035] E.; Weemaes, C.; Pearson, P. L.: Genetic variation in ICF syndrome:

[45036] evidence for genetic heterogeneity. Hum. Mutat. 16: 509-517, 2000.

[45037] 6824.

[45038] Ehrlich, M.; Buchanan, K. L.; Tsien, F.; Jiang, G.; Sun, B.; Uicker,

[45039] W.; Weemaes, C. M. R.; Smeets, D.; Sperling, K.; Belohradsky, B. H.;

[45040] Tommerup, N.; Misek, D. E.; Rouillard, J.-M.; Kuick, R.;

Hanash, S.

[45041] M.: DNA methyltransferase 3B mutations linked to the ICF syndrome

[45042] cause dysregulation of lymphogenesis genes. Hum. Molec. Genet. 10:

[45043] 2917-2931, 2001.

[45044] 6825.

[45045] Barash, V.; Mandel, H.; Sella, S.; Geiger, R.: 3-Hydroxy-3-methylglutaryl-coenzyme

[45046] A lyase deficiency: biochemical studies and family investi-gation of

[45047] four generations. J. Inherit. Metab. Dis. 13: 156-164, 1990.

[45048] 6826.

[45049] Berry, H. K.; Suchy, F. J.; Norman, E. J.: HMG CoA lyase de-ficiency

[45050] in double first cousins: relation of leucine defect to fat metabolism.

[45051] (Abstract) Am. J. Hum. Genet. 33: 37A, 1981.

[45052] 6827.

[45053] Duran, M.; Shutgens, R. B. H.; Ketel, A.; Heymans, H.;

Berntssen,
[45054] M. W. J.; Ketting, D.; Wadman, S. K.:
3-Hydroxy-3-methylglutaryl
[45055] coenzyme A lyase deficiency: postnatal management fol-
lowing prenatal
[45056] diagnosis by analysis of maternal urine. J. Pediat. 95:
1004-1007,
[45057] 1979.

[45058] 6828.

[45059] Faull, K.; Bolton, P.; Halpern, B.; Hammond, J.; Danks, D.
M.;
[45060] Hahnel, R.; Wilkinson, S. P.; Wysocki, S. J.; Masters, P. L.:
Patient
[45061] with defect in leucine metabolism. (Letter) New Eng. J.
Med. 294:
[45062] 1013, 1976.

[45063] 6829.

[45064] Faull, K. F.; Bolton, P. D.; Halpern, B.; Hammond, J.; Danks,
D.
[45065] M.: The urinary organic acid profile associated with
3-hydroxy-3-methylglutaric
[45066] aciduria. Clin. Chim. Acta 73: 553-559, 1976.

[45067] 6830.

[45068] Gibson, K. M.; Breuer, J.; Kaiser, K.; Nyhan, W. L.; McCoy, E.

[45069] E.; Ferreira, P.; Greene, C. L.; Blitzer, M. G.; Shapira, E.; Reverte,

[45070] F.; Conde, C.; Bagnell, P.; Cole, D. E. C.: 3-Hydroxy-3-methylglutaryl-coenzyme

[45071] A lyase deficiency: report of five new patients. J. Inherit. Metab.

[45072] Dis. 11: 76-87, 1988.

[45073] 6831.

[45074] Gibson, K. M.; Breuer, J.; Nyhan, W. L.: 3-Hydroxy-3-methylglutaryl-coenzyme

[45075] A lyase deficiency: review of 18 reported patients. Europ. J. Pediat. 148:

[45076] 180-186, 1988.

[45077] 6832.

[45078] Leonard, J. V.; Seakins, J. W. T.; Griffin, N. K.: Beta-hy-droxy-methylglutaricaciduria

[45079] presenting as Reye's syndrome. (Letter) Lancet I: 680, 1979.

[45080] 6833.

[45081] Mitchell, G. A.; Ozand, P. T.; Robert, M.-F.; Ashmarina, L.; Roberts,

[45082] J.; Gibson, K. M.; Wanders, R. J.; Wang, S.; Chevalier, I.; Plochl,

[45083] R.; Miziorko, H.: HMG CoA lyase deficiency: identification of five

[45084] causal point mutations in codons 41 and 42, including a frequent Saudi

[45085] Arabian mutation, R41Q. Am. J. Hum. Genet. 62: 295-300, 1998.

[45086] 6834.

[45087] Mitchell, G. A.; Robert, M.-F.; Fontaine, G.; Wang, S.; Lam-bert,

[45088] M.; Cole, D.; Lee, C.; Gibson, M.; Miziorko, H.: HMG CoA lyase (HL)

[45089] deficiency: detection of a causal mutation in an affected French-Canadian

[45090] sibship. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A173, 1992.

[45091] 6835.

[45092] Neufeld, E. J.; Mandel, H.; Raz, T.; Szargel, R.; Yandava, C.

N.;

[45093] Stagg, A.; Faure, S.; Barrett, T.; Buist, N.; Cohen, N.: Localization

[45094] of the gene for thiamine-responsive megaloblastic anemia syndrome,

[45095] on the long arm of chromosome 1, by homozygosity mapping. Am. J.

[45096] Hum. Genet. 61: 1335-1341, 1997.

[45097] 6836.

[45098] Raz, T.; Barrett, T.; Szargel, R.; Mandel, H.; Neufeld, E. J.;

[45099] Nosaka, K.; Viana, M. B.; Cohen, N.: Refined mapping of the gene

[45100] for thiamine-responsive megaloblastic anemia syndrome and evidence

[45101] for genetic homogeneity. Hum. Genet. 103: 455-461, 1998.

[45102] 6837.

[45103] Fredrickson, D. S.; Levy, R. I.: Familial hyperlipoproteinemia.In:

[45104] Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic

[45105] Basis of Inherited Disease. New York: McGraw-Hill (pub.)

(3rd

[45106] ed.): 1972. Pp. 545-614.

[45107] 6838.

[45108] Ameis, D.; Kobayashi, J.; Davis, R. C.; Ben-Zeev, O.; Mal-

loy, M.

[45109] J.; Kane, J. P.; Lee, G.; Wong, H.; Havel, R. J.; Schotz, M. C.:

[45110] Familial chylomicronemia (type I hyperlipoproteinemia)

due to a single

[45111] missense mutation in the lipoprotein lipase gene. J. Clin.

Invest. 87:

[45112] 1165-1170, 1991.

[45113] 6839.

[45114] Auwerx, J. H.; Babirak, S. P.; Fujimoto, W. Y.; Iverius, P. H.;

[45115] Brunzell, J. D.: Defective enzyme protein in lipoprotein li-

pase deficiency. Europ.

[45116] J. Clin. Invest. 19: 433-437, 1989.

[45117] 6840.

[45118] Beg, O. U.; Meng, M. S.; Skarlatos, S. I.; Previato, L.; Brun-

zell,

[45119] J. D.; Brewer, H. B., Jr.; Fojo, S. S.: Lipoprotein li-

pase(Bethesda):

[45120] a single amino acid substitution (ala176-to-thr) leads to abnormal

[45121] heparin binding and loss of enzymic activity. Proc. Nat. Acad. Sci. 87:

[45122] 3474-3478, 1990.

[45123] 6841.

[45124] Benlian, P.; De Gennes, J. L.; Foubert, L.; Zhang, H.; Gagne, S.

[45125] E.; Hayden, M.: Premature atherosclerosis in patients with familial

[45126] chylomicronemia caused by mutations in the lipoprotein lipase gene. New

[45127] Eng. J. Med. 335: 848-854, 1996.

[45128] 6842.

[45129] Berger, G. M. B.: An incomplete form of familial lipopro-tein lipase

[45130] deficiency presenting with type I hyperlipoproteinemia. Am. J. Clin.

[45131] Path. 88: 369-373, 1987.

[45132] 6843.

[45133] Berger, H.; Richter, A.; Gilardi, A.; Wagner, H.: Essential familial
[45134] hyperlipaemia in a 2-year-old child. Ann. Paediat. 199: 445-466,
[45135] 1962.

[45136] 6844.

[45137] Bergeron, J.; Normand, T.; Bharucha, A.; Ven Murthy, M. R.; Julien,
[45138] P.; Gagne, C.; Dionne, C.; De Braekeleer, M.; Brun, D.; Hayden, M.
[45139] R.; Lupien, P. J.: Prevalence, geographical distribution and genealogical
[45140] investigations of mutation 188 of lipoprotein lipase gene in the French
[45141] Canadian population of Quebec. Clin. Genet. 41: 206-210, 1992.
[45142] 6845.

[45143] Bertolini, S.; Simone, M. L.; Pes, G. M.; Ghisellini, M.; Rol-
leri,
[45144] M.; Bellocchio, A.; Elicio, N.; Masturzo, P.; Calandra, S.: Pseudodominance
[45145] of lipoprotein lipase (LPL) deficiency due to a nonsense

mutation

[45146] (tyr302-to-term) in exon 6 of LPL gene in an Italian family from Sardinia

[45147] (LPL-Olbia). Clin. Genet. 57: 140-147, 2000.

[45148] 6846.

[45149] Boer, J. M. A.; Kuivenhoven, J. A.; Feskens, E. J. M.; Schouten,

[45150] E. G.; Havekes, L. M.; Seidell, J. C.; Kastelein, J. J. P.; Kromhout,

[45151] D.: Physical activity modulates the effect of a lipoprotein lipase

[45152] mutation (D9N) on plasma lipids and lipoproteins. Clin. Genet. 56:

[45153] 158-163, 1999.

[45154] 6847.

[45155] Boggs, J. D.; Hsia, D. Y.-Y.; Mais, R. F.; Bigler, J. A.: The

[45156] genetic mechanism of idiopathic hyperlipemia. New Eng. J. Med. 257:

[45157] 1101-1108, 1957.

[45158] 6848.

[45159] Breckenridge, W. C.; Little, A. C.; Steiner, G.; Chow, A.;

Poapst,

[45160] M.: Hypertriglyceridemia associated with deficiency of C-II apoprotein

[45161] in plasma lipoproteins. New Eng. J. Med. 298: 1265-1273, 1978.

[45162] 6849.

[45163] Brunzell, J. D.; Chait, A.; Nikkila, E. A.; Ehnholm, C.; Huttunen,

[45164] J. K.; Steiner, G.: Heterogeneity of primary lipoprotein lipase deficiency. Metabolism 29:

[45165] 624-629, 1980.

[45166] 6850.

[45167] Busca, R.; Martinez, M.; Vilella, E.; Pognonec, P.; Deeb, S.;

[45168] Auwerx, J.; Reina, M.; Vilaro, S.: The mutation gly142-to-glu in

[45169] human lipoprotein lipase produces a missorted protein that is diverted

[45170] to lysosomes. J. Biol. Chem. 271: 2139-2146, 1996.

[45171] 6851.

[45172] Cantin, B.; Boudriau, S.; Bertrand, M.; Brun, L.-D.; Gagne, C.;

[45173] Rogers, P. A.; Ven Murthy, M. R.; Lupien, P.-J.; Julien, P.: Hemolysis
[45174] in primary lipoprotein lipase deficiency. Metabolism 44: 652-658,
[45175] 1995.

[45176] 6852.

[45177] Chimienti, G.; Capurso, A.; Resta, F.; Pepe, G.: A G-to-C change
[45178] at the donor splice site of intron 1 causes lipoprotein lipase deficiency
[45179] in a Southern-Italian family. Biochem. Biophys. Res. Commun. 187:
[45180] 620-627, 1992.

[45181] 6853.

[45182] Clark, A. G.; Weiss, K. M.; Nickerson, D. A.; Taylor, S. L.; Buchanan,
[45183] A.; Stengard, J.; Salomaa, V.; Vartiainen, E.; Perola, M.; Boerwinkle,
[45184] E.; Sing, C. F.: Haplotype structure and population genetic inferences
[45185] from nucleotide-sequence variation in human lipoprotein lipase. Am.

[45186] J. Hum. Genet. 63: 595-612, 1998.

[45187] 6854.

[45188] Clee, S. M.; Loubser, O.; Collins, J.; Kastelein, J. J. P.; Hay-
den,
[45189] M. R.: The LPL S447X cSNP is associated with decreased
blood pressure
[45190] and plasma triglycerides, and reduced risk of coronary
artery disease. Clin.
[45191] Genet. 60: 293-300, 2001.

[45192] 6855.

[45193] De Braekeleer, M.; Dionne, C.; Gagne, C.; Julien, P.; Brun,
D.;
[45194] Ven Murthy, M. R.; Lupien, P.-J.: Founder effect in familial
hyperchylomicronemia
[45195] among French Canadians of Quebec. Hum. Hered. 41:
168-173, 1991.
[45196] 6856.

[45197] De Bruin, T. W. A.; Mailly, F.; Van Barlingen, H. H. J. J.;
Fisher,
[45198] R.; Castro Cabezas, M.; Talmud, P.; Dallinga-Thie, G. M.;
Humphries,

[45199] S. E.: Lipoprotein lipase gene mutations D9N and N291S in four pedigrees
[45200] with familial combined hyperlipidaemia. Europ. J. Clin. In-
vest. 26:
[45201] 631-639, 1996.

[45202] 6857.

[45203] Deeb, S. S.; Peng, R.: Structure of the human lipoprotein lipase
[45204] gene. Biochemistry 28: 4131-4135, 1989.

[45205] 6858.

[45206] Henderson, H. E.; Hassan, F.; Berger, G. M. B.; Hayden, M. R.
[45207] : The lipoprotein lipase gly188-to-glu mutation in South Africans
[45208] of Indian descent: evidence suggesting common origins and an increased
[45209] frequency. J. Med. Genet. 29: 119-122, 1992.

[45210] 6859.

[45211] Henderson, H. E.; Hassan, F.; Marais, D.; Hayden, M. R.: A new
[45212] mutation destroying disulphide bridging in the C-terminal

domain of
[45213] lipoprotein lipase. Biochem. Biophys. Res. Commun. 227: 189-194,
[45214] 1996.

[45215] 6860.

[45216] Henderson, H. E.; Ma, Y.; Hassan, M. F.; Monsalve, M. V.; Marais,
[45217] A. D.; Winkler, F.; Gubernator, K.; Peterson, J.; Brunzell, J. D.;
[45218] Hayden, M. R.: Amino acid substitution (ile194-to-thr) in exon 5
[45219] of the lipoprotein lipase gene causes lipoprotein lipase deficiency
[45220] in three unrelated probands: support for a multicentric origin. J.
[45221] Clin. Invest. 87: 2005-2011, 1991.

[45222] 6861.

[45223] Kousseff, B. G.; Matsuoka, L. Y.; Stenn, K. S.; Hobbins, J. C.;
[45224] Mahoney, M. J.; Hashimoto, K.: Prenatal diagnosis of Sjo-gren-Larsson
[45225] syndrome. J. Pediat. 101: 998-1001, 1982.

[45226] 6862.

[45227] Aoki, K.; Siegel, F. L.: Hyperphenylalaninemia: disaggrega-
[45228] tion of brain polyribosomes in young rats. Science 168:
129-130, 1970.

[45229] 6863.

[45230] Apold, J.; Eiken, H. G.; Odland, E.; Fredriksen, A.; Bakken,
A.;
[45231] Lorens, J. B.; Boman, H.: A termination mutation prevalent
in Norwegian
[45232] haplotype 7 phenylketonuria genes. Am. J. Hum. Genet.
47: 1002-1007,
[45233] 1990.

[45234] 6864.

[45235] Apold, J.; Eiken, H. G.; Svensson, E.; Kunert, E.; Kozak, L.;
Cechak,
[45236] P.; Guttler, F.; Giltay, J.; Lichter-Konecki, U.; Melle, D.;
Jaruzelska,
[45237] J. M.: The phenylketonuria G272X haplotype 7 mutation in
European
[45238] populations. Hum. Genet. 92: 107-109, 1993.

[45239] 6865.

[45240] Arthur, L. J. H.; Hulme, J. D.: Intelligent, small for dates baby

[45241] born to oligophrenic phenylketonuric mother after low phenylalanine

[45242] diet during pregnancy. Pediatrics 46: 235-239, 1970.

[45243] 6866.

[45244] Auerbach, V. H.; DiGeorge, A. M.; Carpenter, G. G.: Phenylalaninemia:

[45245] a study of the diversity of disorders which produce elevation of blood

[45246] concentrations of phenylalanine.In: Nyhan, W. L.: Amino Acid Metabolism

[45247] and Genetic Variation. New York: McGraw-Hill (pub.) 1967. Pp.

[45248] 11-68.

[45249] 6867.

[45250] Avigad, S.; Cohen, B. E.; Bauer, S.; Schwartz, G.; Frydman, M.;

[45251] Woo, S. L. C.; Niny, Y.; Shiloh, Y.: A single origin of phenylketonuria

[45252] in Yemenite Jews. Nature 344: 168-170, 1990.

[45253] 6868.

[45254] Avigad, S.; Cohen, B. E.; Woo, S. L. C.; Shiloh, Y.: A specific

[45255] deletion within the phenylalanine hydroxylase gene is common to most

[45256] Yemenite Jewish phenylketonuria patients. (Abstract) Am. J. Hum.

[45257] Genet. 41: A205 only, 1987.

[45258] 6869.

[45259] Baric, I.; Mardesic, D.; Sarnavoka, V.; Lichter-Konecki, U.; Konecki,

[45260] D. S.; Trefz, F. K.: Geographical distribution of the P281L mutation

[45261] at the phenylalanine hydroxylase locus: possible origin in southeastern

[45262] Europe. J. Inherit. Metab. Dis. 17: 376-377, 1994.

[45263] 6870.

[45264] Barranger, J. A.; Geiger, P. J.; Arezino, A.; Bessman, S. P.:

[45265] Isozymes of phenylalanine hydroxylase. Science 175: 903-905, 1972.

[45266] 6871.

[45267] Bartholome, K.; Olek, K.; Trefz, F.: Compound heterozy-
gotes in
[45268] hyperphenylalaninaemia. Hum. Genet. 65: 405-406,
1984.
[45269] 6872.

[45270] Benit, P.; Rey, F.; Blandin-Savoja, F.; Munnich, A.; Abadie,
V.;
[45271] Rey, J.: The mutant genotype is the main determinant of
the metabolic
[45272] phenotype in phenylalanine hydroxylase deficiency. Molec.
Genet.
[45273] Metab. 68: 43-47, 1999.

[45274] 6873.

[45275] Berg, K.; Saugstad, L. F.: A linkage study of phenylke-
tonuria. Clin.
[45276] Genet. 6: 147-152, 1974.

[45277] 6874.

[45278] Bickel, H.: The first treatment of phenylketonuria. Europ. J.

[45279] Pediat. 155 (suppl. 1): 2-3, 1996.

[45280] 6875.

[45281] Bickel, H.; Gerard, J.; Hickmans, E. M.: The influence of phenylalanine

[45282] intake on the chemistry and behavior of a phenylketonuric child. Acta

[45283] Paediat. 43: 64-77, 1954.

[45284] 6876.

[45285] Sekiguchi, K.; Klos, A. M.; Kurachi, K.; Yoshitake, S.; Hako-mori,

[45286] S.: Human liver fibronectin complementary DNAs: identifi-cation of

[45287] two different messenger RNAs possibly encoding the al-pha and beta

[45288] subunits of plasma fibronectin. Biochemistry 25: 4936-4941, 1986.

[45289] 6877.

[45290] Shirakami, A.; Shigekiyo, T.; Hirai, Y.; Takeichi, T.; Kawauchi,

[45291] S.; Saito, S.; Miyoshi, K.: Plasma fibronectin deficiency in eight

[45292] members of one family. Lancet I: 473-474, 1986.

[45293] 6878.

[45294] Shows, T. B.: Personal Communication. Buffalo, N. Y. 1982.

[45295] 6879.

[45296] Skow, L. C.; Adkison, L.; Womack, J. E.; Beamer, W. G.; Taylor,

[45297] B. A.: Mapping of the mouse fibronectin gene (Fn-1) to chromosome

[45298] 1: conservation of the Idh-1--Cryg--Fn-1 synteny group in mammals. Genomics 1:

[45299] 283-286, 1987.

[45300] 6880.

[45301] Smith, M.; Gold, L. I.; Pearlstein, E.; Krinsky, A.: Expression

[45302] of mouse and human fibronectin in hybrid cells. (Abstract) Cytogenet.

[45303] Cell Genet. 25: 205 only, 1979.

[45304] 6881.

[45305] Smith, M.; Krinsky, A. M.; Arredondo-Vega, F. X.; Pearl-stein,

[45306] E.: Production of soluble fibronectin by RAG x human fi-broblast hybrids.

[45307] (Abstract) Cytogenet. Cell Genet. 32: 318 only, 1982.

[45308] 6882.

[45309] Wu, B.-L.; Milunsky, A.; Wyandt, H.; Hoth, C.; Baldwin, C.; Skare,

[45310] J.: In situ hybridization applied to Waardenburg syndrome. Cytogenet.

[45311] Cell Genet. 63: 29-32, 1993.

[45312] 6883.

[45313] Zardi, L.; Cianfriglia, M.; Balza, E.; Carnemolla, B.; Siri, A.;

[45314] Croce, C. M.: Species-specific monoclonal antibodies in the assignment

[45315] of the gene for human fibronectin to chromosome 2. EMBO J. 1: 929-933,

[45316] 1982.

[45317] 6884.

[45318] Zardi, L.; Siri, A.; Carnemolla, B.; Santi, L.; Gardner, W. D.;

[45319] Hoch, S. O.: Fibronectin: a chromatin-associated protein? Cell 18:

[45320] 649-657, 1979.

[45321] 6885.

[45322] Adkison, L. R.; White, R. A.; Haney, D. M.; Lee, J. C.; Pusey,

[45323] K. T.; Gardner, J.: The fibronectin receptor, alpha subunit (Itga5)

[45324] maps to murine chromosome 15, distal to D15Mit16. Mammalian Genome 5:

[45325] 456-457, 1994.

[45326] 6886.

[45327] Argraves, W. S.; Pytela, R.; Suzuki, S.; Millan, J. L.; Pier-schbacher,

[45328] M. D.; Ruoslahti, E.: cDNA sequences from the alpha sub-unit of the

[45329] fibronectin receptor predict a transmembrane domain and a short cytoplasmic

[45330] peptide. J. Biol. Chem. 261: 12922-12924, 1986.

[45331] 6887.

[45332] Argraves, W. S.; Suzuki, S.; Arai, H.; Thompson, K.; Pier-schbacher,

[45333] M. D.; Ruoslahti, E.: Amino acid sequence of the human fi-bronectin

[45334] receptor. J. Cell Biol. 105: 1183-1190, 1987.

[45335] 6888.

[45336] Fitzgerald, L. A.; Poncz, M.; Steiner, B.; Rall, S. C., Jr.; Ben-
nett,
[45337] J. S.; Phillips, D. R.: Comparison of cDNA-derived protein
sequences
[45338] of the human fibronectin and vitronectin receptor alpha-
subunits and
[45339] platelet glycoprotein IIb. Biochemistry 26: 8158-8165,
1987.
[45340] 6889.

[45341] Krissansen, G. W.; Yuan, Q.; Jenkins, D.; Jiang, W.-M.;
Rooke,
[45342] L.; Spurr, N. K.; Eccles, M.; Leung, E.; Watson, J. D.: Chro-
mosomal
[45343] locations of the genes coding for the integrin beta-6 and
beta-7 subunits. Immunogenetics 35:
[45344] 58-61, 1992.

[45345] 6890.

[45346] Sosnoski, D.; Emanuel, B. S.; Hawkins, A. L.; van Tuinen,
P.; Ledbetter,
[45347] D. H.; Nussbaum, R. L.; Kaos, F.-T.; Schwartz, E.; Phillips,
D.; Bennett,
[45348] J. S.; Fitzgerald, L. A.; Poncz, M.: Chromosomal localiza-

tion of

[45349] the genes for the vitronectin and fibronectin receptors alpha-subunits

[45350] and for platelet glycoproteins IIb and IIIa. J. Clin. Invest. 81:

[45351] 1993-1998, 1988.

[45352] 6891.

[45353] Spurr, N. K.; Rooke, L.: Confirmation of the assignment of the

[45354] vitronectin (VNRA) and fibronectin (FNRA) receptor alpha-subunits. Ann.

[45355] Hum. Genet. 55: 217-223, 1991.

[45356] 6892.

[45357] Akula, S. M.; Pramod, N. P.; Wang, F.-Z.; Chandran, B.: Integrin

[45358] alpha-3/beta-1 (CD 49c/29) is a cellular receptor for Kaposi's sarcoma-associated

[45359] herpesvirus (KSHV/HHV-8) entry into the target cells. Cell 108:

[45360] 407-419, 2002.

[45361] 6893.

[45362] Arregui, C.; Pathre, P.; Lilien, J.; Balsamo, J.: The nonre-
ceptor
[45363] tyrosine kinase Fer mediates cross-talk between N-
cadherin and beta-1-integrins. J.
[45364] Cell Biol. 149: 1263-1273, 2000.

[45365] 6894.

[45366] Giuffra, L. A.; Lichter, P.; Wu, J.; Kennedy, J. L.; Pakstis, A.

[45367] J.; Rogers, J.; Kidd, J. R.; Harley, H.; Jenkins, T.; Ward, D.
C.;
[45368] Kidd, K. K.: Genetic and physical mapping and population
studies
[45369] of a fibronectin receptor beta-subunit-like sequence on
human chromosome
[45370] 19. Genomics 8: 340-346, 1990.

[45371] 6895.

[45372] Giuffra, L. A.; Wu, J.; Lichter, P.; Kennedy, J. L.; Castiglione,

[45373] C.; Pakstis, A. J.; Ward, D.; Kidd, K. K.: Mapping of a fi-
bronectin
[45374] receptor beta subunit-like sequence to chromosome 19.
(Abstract) Am.
[45375] J. Hum. Genet. 45 (suppl.): A141 only, 1989.

[45376] 6896.

[45377] Goodfellow, P. J.; Nevanlinna, H. A.; Gorman, P.; Sheer, D.; Lam,

[45378] G.; Goodfellow, P. N.: Assignment of the gene encoding the beta-subunit

[45379] of the human fibronectin receptor (beta-FNR) to chromosome 10p11.2. Ann.

[45380] Hum. Genet. 53: 15-22, 1989.

[45381] 6897.

[45382] Graus-Porta, D.; Blaess, S.; Senften, M.; Littlewood-Evans, A.;

[45383] Damsky, C.; Huang, Z.; Orban, P.; Klein, R.; Schittny, J. C.; Muller,

[45384] U.: Beta-1-class integrins regulate the development of laminae and

[45385] folia in the cerebral and cerebellar cortex. Neuron 31: 367-379,

[45386] 2001.

[45387] 6898.

[45388] Hynes, R. O.: Integrins: a family of cell surface receptors. Cell 48:

[45389] 549-554, 1987.

[45390] 6899.

[45391] Johansson, S.; Forsberg, E.; Lundgren, B.: Comparison of fibronectin

[45392] receptors from rat hepatocytes and fibroblasts. J. Biol. Chem. 262:

[45393] 7819-7824, 1987.

[45394] 6900.

[45395] Lu, T. T.; Cyster, J. G.: Integrin-mediated long-term B cell retention

[45396] in the splenic marginal zone. Science 297: 409-412, 2002.

[45397] 6901.

[45398] Messer Peters, P.; Kamarck, M. E.; Hemler, M. E.; Stro-minger,

[45399] J. L.; Ruddle, F. H.: Genetic and biochemical characteriza-tion of

[45400] human lymphocyte cell surface antigens: the A-1A5 and A-3A4 determinants. J.

[45401] Exp. Med. 159: 1441-1454, 1984.

[45402] 6902.

[45403] Pytela, R.; Pierschbacher, M. D.; Ginsberg, M. H.; Plow, E. F.;

[45404] Ruoslahti, E.: Platelet membrane glycoprotein IIb/IIIa: member of

[45405] a family of arg-gly-asp-specific adhesion receptors. Science 231:

[45406] 1159-1162, 1986.

[45407] 6903.

[45408] Woods, V. L., Jr.; Pischel, K. D.; Avery, E. D.; Bluestein, H.

[45409] G.: Antigenic polymorphism of human very late activation protein-2

[45410] (platelet glycoprotein Ia-IIa): platelet alloantigen Hc(a). J. Clin.

[45411] Invest. 83: 978-985, 1989.

[45412] 6904.

[45413] Wu, J. S.; Giuffra, L. A.; Goodfellow, P. J.; Myers, S.; Carson,

[45414] N. L.; Anderson, L.; Hoyle, L. S.; Simpson, N. E.; Kidd, K. K.: The

[45415] beta subunit locus of the human fibronectin receptor: DNA restriction

[45416] fragment length polymorphism and linkage mapping

studies. Hum. Genet. 83:
[45417] 383-390, 1989.

[45418] 6905.

[45419] Huang, F.; Shi, L. J.; Heng, H. H. Q.; Fei, J.; Guo, L.-H.: As-
signment
[45420] of the human GABA transporter gene (GABATHG) locus to
chromosome 3p24-p25. Genomics 29:
[45421] 302-304, 1995.

[45422] 6906.

[45423] Lam, D. M.-K.; Fei, J.; Zhang, X.-Y.; Tam, A. C. W.; Zhu,
L.-H.;
[45424] Huang, F.; King, S. C.; Guo, L.-H.: Molecular cloning and
structure
[45425] of the human (GABATHG) GABA transporter gene. Molec.
Brain Res. 19:
[45426] 227-232, 1993.

[45427] 6907.

[45428] Nelson, H.; Mandiyan, S.; Nelson, N.: Cloning of the hu-
man brain
[45429] GABA transporter. FEBS Lett. 269: 181-184, 1990.

[45430] 6908.

[45431] Brenner, B.; Sanchez-Vega, B.; Wu, S.-M.; Lanir, N.; Stafford,
[45432] D. W.; Solera, J.: A missense mutation in a gamma-glutamyl carboxylase
[45433] gene causes combined deficiency of all vitamin K-dependent blood coagulation
[45434] factors. Blood 92: 4554-4559, 1998.

[45435] 6909.

[45436] Brenner, B.; Tavori, S.; Zivelin, A.; Keller, C. B.; Suttie, J.

[45437] W.; Tatarsky, I.; Seligsohn, U.: Hereditary deficiency of all vitamin
[45438] K-dependent procoagulants and anticoagulants. Brit. J. Haemat. 75:
[45439] 537-542, 1990.

[45440] 6910.

[45441] Kuo, W.-L.; Stafford, D. W.; Cruces, J.; Gray, J.; Solera, J.:

[45442] Chromosomal localization of the gamma-glutamyl carboxylase gene at
[45443] 2p12. Genomics 25: 746-748, 1995.

[45444] 6911.

[45445] Mutucumarana, V. P.; Stafford, D. W.; Stanley, T. B.; Jin, D.-Y.;
[45446] Solera, J.; Brenner, B.; Azerad, R.; Wu, S.-M.: Expression and characterization
[45447] of the naturally occurring mutation L394R in human gamma-glutamyl
[45448] carboxylase. J. Biol. Chem. 275: 32572-32577, 2000.

[45449] 6912.

[45450] Soute, B. A. M.; Ulrich, M. M. W.; Watson, A. D. J.; Maddi-son,
[45451] J. E.; Ebberink, R. H. M.; Vermmer, C.: Congenital defi-ciency of
[45452] all vitamin K-dependent blood coagulation factors due to a defective
[45453] vitamin K-dependent carboxylase in Devon Rex cats. Thromb. Haemost. 68:
[45454] 521-525, 1992.

[45455] 6913.

[45456] Spronk, H. M. H.; Farah, R. A.; Buchanan, G. R.; Vermeer, C.; Soute,
[45457] B. A. M.: Novel mutation in the gamma-glutamyl carboxy-lase gene resulting

[45458] in congenital combined deficiency of all vitamin K-dependent blood

[45459] coagulation factors. Blood 96: 3650-3652, 2000.

[45460] 6914.

[45461] Wu, S.-M.; Cheung, W.-F.; Frazier, D.; Stafford, D. W.: Cloning

[45462] and expression of the cDNA for human gamma-glutamyl carboxylase. Science 254:

[45463] 1634-1636, 1991.

[45464] 6915.

[45465] Wu, S. M.; Stafford, D. W.; Frazier, L. D.; Fu, Y. Y.; High, K.

[45466] A.; Chu, K.; Sanchez-Vega, B.; Solera, J.: Genomic se-quence and transcription

[45467] start site for the human gamma-glutamyl carboxylase. Blood 89: 4058-4062,

[45468] 1997.

[45469] 6916.

[45470] Bibas, M.; Zampa, G.; Procopio, A.; Guaitolini, R.: High serum

[45471] gamma-glutamyltransferase concentrations in a family. (Letter) New

[45472] Eng. J. Med. 330: 1832-1833, 1994.

[45473] 6917.

[45474] Courtay, C.; Heisterkamp, N.; Siest, G.; Groffen, J.: Expres-
sion

[45475] of multiple gamma-glutamyltransferase genes in man.
Biochem. J. 297:

[45476] 503-508, 1994.

[45477] 6918.

[45478] Figlewicz, D. A.; Delattre, O.; Guellaen, G.; Krizus, A.;
Thomas,

[45479] G.; Zucman, J.; Rouleau, G. A.: Mapping of human
gamma-glutamyl transpeptidase

[45480] genes on chromosome 22 and other human autosomes.
Genomics 17: 299-305,

[45481] 1993.

[45482] 6919.

[45483] Grausz, J. D.; Bulle, F.; Hillion, J.; Bernheim, A.; Miglierina,

[45484] R.; Guellaen, G.; Berger, R.: Indication that a gamma glu-
tamyl transferase

[45485] (GGT) gene is located centromeric to Ig-lambda (IGL) on
chromosome

[45486] 22q11. (Abstract) Cytogenet. Cell Genet. 51: 1007 only, 1989.

[45487] 6920.

[45488] Pawlak, A.; Wu, S.-J.; Bulle, F.; Suzuki, A.; Chikhi, N.; Ferry,

[45489] N.; Baik, J.-H.; Siegrist, S.; Guellaen, G.: Different gamma-glutamyl

[45490] transpeptidase mRNAs are expressed in human liver and kidney. Biochem.

[45491] Biophys. Res. Commun. 164: 912-918, 1989.

[45492] 6921.

[45493] Rajpert-De Meyts, E.; Heisterkamp, N.; Groffen, J.: Cloning and

[45494] nucleotide sequence of human gamma-glutamyl transpeptidase. Proc.

[45495] Nat. Acad. Sci. 85: 8840-8844, 1988.

[45496] 6922.

[45497] Visvikis, A.: Personal Communication. Nancy, France 9/26/1997.

[45498] 6923.

[45499] Visvikis, A.; Thioudellet, C.; Oster, T.; Fournel-Gigleux, S.;

[45500] Wellman, M.; Siest, G.: High-level expression of enzymatically active
[45501] mature human gamma-glutamyltransferase in transgenic V79 Chinese hamster
[45502] cells. Proc. Nat. Acad. Sci. 88: 7361-7365, 1991.

[45503] 6924.

[45504] Dean, M.; Lucas-Derse, S.; Bolos, A.; O'Brien, S. J.; Kirkness,
[45505] E. F.; Fraser, C. M.; Goldman, D.: Genetic mapping of the beta-1
[45506] GABA receptor gene to human chromosome 4, using a tetranucleotide
[45507] repeat polymorphism. Am. J. Hum. Genet. 49: 621-626, 1991.
[45508] 6925.

[45509] Kirkness, E. F.; Kusiak, J. W.; Fleming, J. T.; Menninger, J.;

[45510] Gocayne, J. D.; Ward, D. C.; Venter, J. C.: Isolation, characterization,
[45511] and localization of human genomic DNA encoding the beta-1 subunit
[45512] of the GABA-A receptor (GABRB1). Genomics 10: 985-995, 1991.

[45513] 6926.

[45514] Lopez de Castro, J. A.; Strominger, J. L.; Strong, D. M.; Orr,

[45515] H. T.: Structure of crossreactive human histocompatibility antigens

[45516] HLA-A28 and HLA-A2: possible implications for the gen-eration of HLA

[45517] polymorphism. Proc. Nat. Acad. Sci. 79: 3813-3817, 1982.

[45518] 6927.

[45519] Malissen, M.; Malissen, B.; Jordan, B. R.: Exon/intron orga-nization

[45520] and complete nucleotide sequence of an HLA gene. Proc. Nat. Acad.

[45521] Sci. 79: 893-897, 1982.

[45522] 6928.

[45523] Mann, D. L.; Rogentine, G. N., Jr.; Fahey, J. L.; Nathenson, S.

[45524] G.: Molecular heterogeneity of human lymphoid (HL-A) al-loantigens. Science 163:

[45525] 1460-1462, 1969.

[45526] 6929.

[45527] MacDonald, K. S.; Fowke, K. R.; Kimani, J.; Dunand, V. A.; Nagelkerke,

[45528] N. J. D.; Ball, T. B.; Oyugi, J.; Njagi, E.; Gaur, L. K.; Brunham,

[45529] R. C.; Wade, J.; Luscher, M. A.; Krausa, P.; Rowland-Jones, S.; Ngugi,

[45530] E.; Bwayo, J. J.; Plummer, F. A.: Influence of HLA supertypes on

[45531] susceptibility and resistance to human immunodeficiency virus type

[45532] 1 infection. J. Infect. Dis. 181: 1581-1589, 2000.

[45533] 6930.

[45534] Mayr, W. R.; Mayr, D.: Analysis of the linkage between the HL-A

[45535] loci and the genes of other markers. Humangenetik 24: 129-133, 1974.

[45536] 6931.

[45537] Messer, G.; Zemmour, J.; Orr, H. T.; Parham, P.; Weiss, E. H.;

[45538] Girdlestone, J.: HLA-J, a second inactivated class I HLA gene related

[45539] to HLA-G and HLA-A. J. Immun. 148: 4043-4053, 1992.

[45540] 6932.

[45541] Morton, C. C.; Kirsch, I. R.; Nance, W. E.; Evans, G. A.; Kor-
man,
[45542] A. J.; Strominger, J. L.: Orientation of loci within the hu-
man major
[45543] histocompatibility complex by chromosomal in situ hy-
bridization. Proc.
[45544] Nat. Acad. Sci. 81: 2816-2820, 1984.

[45545] 6933.

[45546] Morton, N. E.; Rao, D. C.; Lindsten, J.; Hulten, M.; Yee, S.:

[45547] A chiasma map of man. Hum. Hered. 27: 38-51, 1977.

[45548] 6934.

[45549] Mulley, J. C.; Hay, J.; Sheffield, L. J.; Sutherland, G. R.:

[45550] Regional localization for HLA by recombination with a
fragile site
[45551] at 6p23. Am. J. Hum. Genet. 35: 1284-1288, 1983.

[45552] 6935.

[45553] Newell, W. R.; Trowsdale, J.; Beck, S.: MHCDB--database of
the
[45554] human MHC. Immunogenetics 40: 109-115, 1994.

[45555] 6936.

[45556] Orr, H. T.; Bach, F. H.; Ploegh, H. L.; Strominger, J. L.; Ka-
vathas,
[45557] P.; DeMars, R.: Use of HLA loss mutants to analyse the
structure
[45558] of the human major histocompatibility complex. Nature
296: 454-456,
[45559] 1982.

[45560] 6937.

[45561] Paabo, S.: Molecular cloning of ancient Egyptian mummy
DNA. Nature 314:
[45562] 644-645, 1985.

[45563] 6938.

[45564] Patel, R.; Mickey, M. R.; Terasaki, P. I.: Serotyping for ho-
motransplantation
[45565] of kidneys from unrelated donors. New Eng. J. Med. 279:
501-506,
[45566] 1968.

[45567] 6939.

[45568] Payne, R.; Tripp, M.; Weigle, J.; Bodmer, W. F.; Bodmer, J.
G.

[45569] : A new leukocyte isoantigen system in man. Cold Spring Harbor Symp.

[45570] Quant. Biol. 29: 285-295, 1964.

[45571] 6940.

[45572] Polacek, L. A.; Phillips, R. B.; Hackbarth, S. A.; Duquesnoy,

[45573] R. J.: A linkage study of the HLA region using C-band heteromorphisms. Clin.

[45574] Genet. 23: 177-185, 1983.

[45575] 6941.

[45576] Ragoussis, J.; Bloemer, K.; Pohla, H.; Messer, G.; Weiss, E. H.;

[45577] Ziegler, A.: A physical map including a new class I gene (cda12)

[45578] of the human major histocompatibility complex (A2/B13 haplotype) derived

[45579] from a monosomy 6 mutant cell line. Genomics 4: 301-308, 1989.

[45580] 6942.

[45581] Ragoussis, J.; van der Bliek, A.; Trowsdale, J.; Ziegler, A.:

[45582] Mapping of HLA genes using pulsed field gradient electrophoreses. FEBS

[45583] Lett. 204: 1-4, 1986.

[45584] 6943.

[45585] Salter, R. D.; Norment, A. M.; Chen, B. P.; Clayberger, C.; Krensky,

[45586] A. M.; Littman, D. R.; Parham, P.: Polymorphism in the alpha(3) domain

[45587] of HLA-A molecules affects binding to CD8. Nature 338: 345-347,

[45588] 1989.

[45589] 6944.

[45590] Schunter, F.; Wernet, P.; Kompf, J.; Bissbort, S.; Gohler, F.

[45591] : Mapping of the linkage group GLO-Bf-HLA-B, C, A-PGM. II. Segregation

[45592] analysis. Hum. Genet. 44: 321-331, 1978.

[45593] 6945.

[45594] Shivdasani, R. A.; Haluska, F. G.; Dock, N. L.; Dover, J. S.;

[45595] Kineke, E. J.; Anderson, K. C.: Graft-versus-host disease associated

[45596] with transfusion of blood from unrelated HLA-homozygous donors. New

[45597] Eng. J. Med. 328: 766-770, 1993.

[45598] 6946.

[45599] Shukla, H.; Gillespie, G. A.; Srivastava, R.; Collins, F.; Chorney,

[45600] M. J.: A class I jumping clone places the HLA-G gene ap-

proximately

[45601] 100 kilobases from HLA-H within the HLA-A subregion of the human MHC. Genomics 10:

[45602] 905-914, 1991.

[45603] 6947.

[45604] Snell, G. D.: Studies in histocompatibility. Science 213: 172-178,

[45605] 1981.

[45606] 6948.

[45607] Solheim, B. G.; Bratlie, A.; Sandberg, L.; Staub-Nielsen, L.;

[45608] Thorsby, E.: Further evidence of a third HL-A locus. Tissue Antigens 3:

[45609] 439-453, 1973.

[45610] 6949.

[45611] Strachan, T.: Molecular genetics and polymorphism of

class I

[45612] HLA antigens. Brit. Med. Bull. 43: 1-14, 1987.

[45613] 6950.

[45614] Szpak, Y.; Vieville, J.-C.; Tabary, T.; Naud, M.-C.; Chopin, M.;

[45615] Edelson, C.; Cohen, J. H. M.; Dausset, J.; de Kozak, Y.; Pla, M.:

[45616] Spontaneous retinopathy in HLA-A29 transgenic mice. Proc. Nat. Acad.

[45617] Sci. 98: 2572-2576, 2001.

[45618] 6951.

[45619] Teshima, T.; Ordemann, R.; Reddy, P.; Gagin, S.; Liu, C.; Cooke,

[45620] K. R.; Ferrara, J. L. M.: Acute graft-versus-host disease does not

[45621] require alloantigen expression on host epithelium. Nature Med. 8:

[45622] 575-581, 2002.

[45623] 6952.

[45624] Thorsby, E.; Sandberg, L.; Lindholm, A.; Kissmeyer-Nielsen, F.

[45625] : The HL-A system: evidence of a third sub-locus. Scand. J. Haemat. 7:
[45626] 195-200, 1970.

[45627] 6953.

[45628] Tragardh, L.; Rask, L.; Wiman, K.; Fohlman, J.; Peterson, P. A.
[45629] : Amino acid sequence of an immunoglobulin HLA antigen heavy chain
[45630] domain. Proc. Nat. Acad. Sci. 76: 5839-5842, 1979.

[45631] 6954.

[45632] Trowsdale, J.: Personal Communication. London, England 1/12/1983.
[45633] 6955.

[45634] Trowsdale, J.; Ragoussis, J.; Campbell, R. D.: Map of the human
[45635] MHC. Immun. Today 12: 443-446, 1991.

[45636] 6956.

[45637] Van Leeuwen, A.; Eernisse, J. G.; Van Rood, J. J.: A new leucocyte
[45638] group with two alleles: leucocyte group five. Vox Sang. 9: 431-446,

[45639] 1964.

[45640] 6957.

[45641] Van Rood, J. J.: Leucocyte grouping and organ transplantation. Brit.

[45642] J. Haemat. 16: 211–220, 1969.

[45643]

[45644] 100. Van Rood, J. J.: Tissue typing and organ transplantation. Lancet I:

[45645] 1142–1146, 1969.

[45646]

[45647] 101. Van Rood, J. J.; Amos, D. B.: In memoriam: Ruggero Ceppellini,

[45648] 1917–1988. Hum. Immun. 23: 1–3, 1988.

[45649]

[45650] 102. Van Rood, J. J.; Van Leeuwen, A.: Leukocyte grouping: a method

[45651] and its application. J. Clin. Invest. 42: 1382–1390, 1963.

[45652]

[45653] 103. Van Someren, H.; Westerveld, A.; Hagemeijer, A.;

Mees, J. R.;

[45654] Meera Khan, P.; Zaalberg, O. B.: Human antigen and enzyme markers

[45655] in man-Chinese hamster somatic cell hybrids: evidence for synteny

[45656] between the HL-A, PGM-3, ME-1, and IPO-B loci. Proc. Nat. Acad. Sci. 71:

[45657] 962-965, 1974.

[45658]

[45659] 104. Walford, R. L.; Finkelstein, S.; Hanna, C.; Collins, Z.: Third

[45660] sublocus in the HL-A human transplantation system. Nature 224: 74-75,

[45661] 1969.

[45662]

[45663] 105. Weitkamp, L. R.; Van Rood, J. J.; Thorsby, E.; Bias, W. B.; Fotino,

[45664] M.; Lawler, S. D.; Dausset, J.; Mayr, W. R.; Bodmer, J.; Ward, F.

[45665] S.; Seignalet, J.; Payne, R.; Kissmeyer-Nielsen, F.; Gatti, R. A.;

[45666] Sachs, J. A.; Lamm, L. U.: The relation of parental sex and

age to
[45667] recombination in the HL-A system. Hum. Hered. 23:
197-205, 1973.
[45668]

[45669] 106. Wolski, K. P.; Schmid, F. R.; Mittal, K. K.: Genetic
linkage
[45670] between the HL-A system and a deficit of the second
component (C2)
[45671] of complement. Science 188: 1020-1022, 1975.

[45672]

[45673] 107. Ziegler, A.; Ragoussis, J.; Fonatsch, C.; Weiss, E.: A
physical
[45674] map of the human major histocompatibility (HLA) complex
derived from
[45675] a monosomy 6 mutant cell line. (Abstract) Cytogenet. Cell
Genet. 51:
[45676] 1116 only, 1989.

[45677] 6958.

[45678] Wildhage, I.; Trusheim, H.; Goke, B.; Lankat-Buttgereit, B.:
Gene
[45679] expression of the human glucagon-like peptide-1 recep-

tor is regulated
[45680] by Sp1 and Sp3. Endocrinology 140: 624-631, 1999.

[45681] 6959.

[45682] Rucknagel, D. L.; Shreffler, D. C.; Halstead, S. B.: The Bangkok
[45683] variant of the serum group-specific component (Gc) and the frequency
[45684] of the Gc alleles in Thailand. Am. J. Hum. Genet. 20: 478-485, 1968.
[45685] 6960.

[45686] Ashraf, J.; Thompson, E. B.: Identification of the activa-tion-labile
[45687] gene: a single point mutation in the human glucocorticoid receptor
[45688] presents as two distinct receptor phenotypes. Molec. En-docr. 7:
[45689] 631-642, 1993.

[45690] 6961.

[45691] Bamberger, C. M.; Bamberger, A.-M.; de Castro, M.; Chrousos, G.
[45692] P.: Glucocorticoid receptor beta, a potential endogenous

inhibitor
[45693] of glucocorticoid action in humans. J. Clin. Invest. 95:
2435-2441,
[45694] 1995.

[45695] 6962.

[45696] Breslin, M. B.; Geng, C.-D.; Vedeckis, W. V.: Multiple pro-
moters
[45697] exist in the human GR gene, one of which is activated by
glucocorticoids. Molec.
[45698] Endocr. 15: 1381-1395, 2001.

[45699] 6963.

[45700] Bronnegard, M.; Werner, S.; Gustafsson, J.-A.: Primary
cortisol
[45701] resistance associated with a thermolabile glucocorticoid
receptor
[45702] in a patient with fatigue as the only symptom. J. Clin. In-
vest. 78:
[45703] 1270-1278, 1986.

[45704] 6964.

[45705] Bledsoe, R. K.; Montana, V. G.; Stanley, T. B.; Delves, C. J.;

[45706] Apolito, C. J.; McKee, D. D.; Consler, T. G.; Parks, D. J.;

Stewart,
[45707] E. L.; Willson, T. M.; Lambert, M. H.; Moore, J. T.; Pearce, K. H.;
[45708] Xu, H. E.: Crystal structure of the glucocorticoid receptor ligand
[45709] binding domain reveals a novel mode of receptor dimer-ization and coactivator
[45710] recognition. Cell 110: 93-105, 2002.

[45711] 6965.

[45712] Carlstedt-Duke, J.; Stromstedt, P.-E.; Wrange, O.; Bergman, T.;
[45713] Gustafsson, J.-A.; Jornvall, H.: Domain structure of the glucocorticoid
[45714] receptor protein. Proc. Nat. Acad. Sci. 84: 4437-4440, 1987.
[45715] 6966.

[45716] Chrousos, G. P.; Renquist, D.; Brandon, D.; Eil, C.; Pugeat, M.;
[45717] Vigersky, R.; Cutler, G. B., Jr.; Loriaux, D. L.; Lipsett, M. B.:

[45718] Glucocorticoid hormone resistance during primate evolu-tion: receptor-mediated
[45719] mechanisms. Proc. Nat. Acad. Sci. 79: 2036-2040, 1982.

[45720] 6967.

[45721] Chrousos, G. P.; Vingerhoeds, A.; Brandon, D.; Eil, C.; Pugeat,

[45722] M.; DeVroede, M.; Loriaux, D. L.; Lipsett, M. B.: Primary cortisol

[45723] resistance in man: a glucocorticoid receptor-mediated disease. J.

[45724] Clin. Invest. 69: 1261-1269, 1982.

[45725] 6968.

[45726] Chrousos, G. P.; Vingerhoeds, A. C. M.; Loriaux, D. L.; Lipsett,

[45727] M. B.: Primary cortisol resistance: a family study. J. Clin. Endocr.

[45728] Metab. 56: 1243-1245, 1983.

[45729] 6969.

[45730] Diamond, M. I.; Robinson, M. R.; Yamamoto, K. R.: Regula-tion

[45731] of expanded polyglutamine protein aggregation and nu-clear localization

[45732] by the glucocorticoid receptor. Proc. Nat. Acad. Sci. 97: 657-661,

[45733] 2000.

[45734] 6970.

[45735] Dobson, M. G.; Redfern, C. P. F.; Unwin, N.; Weaver, J. U.: The

[45736] N363S polymorphism of the glucocorticoid receptor: po-
tential contribution

[45737] to central obesity in men and lack of association with
other risk

[45738] factors for coronary heart disease and diabetes mellitus. J.
Clin.

[45739] Endocr. Metab. 86: 2270-2274, 2001.

[45740] 6971.

[45741] Encio, I. J.; Detera-Wadleigh, S. D.: The genomic structure
of

[45742] the human glucocorticoid receptor. J. Biol. Chem. 266:
7182-7188,

[45743] 1991.

[45744] 6972.

[45745] Francke, U.; Foellmer, B. E.: The glucocorticoid receptor
gene

[45746] is in 5q31-q32. Genomics 4: 610-612, 1989.

[45747] 6973.

[45748] Gehring, U.; Segnitz, B.; Foellmer, B.; Francke, U.: Chro-
mosome
[45749] assignment of the human gene for the glucocorticoid re-
ceptor (GRL).
[45750] (Abstract) Cytogenet. Cell Genet. 37: 476, 1984.

[45751] 6974.

[45752] Gehring, U.; Segnitz, B.; Foellmer, B.; Francke, U.: Assign-
ment
[45753] of the human gene for the glucocorticoid receptor to
chromosome 5. Proc.
[45754] Nat. Acad. Sci. 82: 3751-3755, 1985.

[45755] 6975.

[45756] Giuffra, L. A.; Kennedy, J. L.; Castiglione, C. M.; Evans, R.
[45757] M.; Wasmuth, J. J.; Kidd, K. K.: Glucocorticoid receptor
maps to
[45758] the distal long arm of chromosome 5. Cytogenet. Cell
Genet. 49:
[45759] 313-314, 1988.

[45760] 6976.

[45761] Hollenberg, S. M.; Weinberger, C.; Ong, E. S.; Cerelli, G.; Oro,
[45762] A.; Lebo, R.; Thompson, E. B.; Rosenfeld, M. G.; Evans, R. M.: Primary
[45763] structure and expression of a functional human glucocorticoid receptor
[45764] cDNA. Nature 318: 635-641, 1985.

[45765] 6977.

[45766] Huebner, K.; Nagarajan, L.; Besa, E.; Angert, E.; Lange, B. J.;
[45767] Cannizzaro, L. A.; van den Berghe, H.; Santoli, D.; Finan, J.; Croce,
[45768] C. M.; Nowell, P. C.: Order of genes on human chromosome 5q with
[45769] respect to 5q interstitial deletions. Am. J. Hum. Genet. 46: 26-36,
[45770] 1990.

[45771] 6978.

[45772] Huizenga, N. A. T. M.; de Lange, P.; Koper, J. W.; Clayton, R.
[45773] N.; Farrell, W. E.; van der Lely, A. J.; Brinkmann, A. O.; de Jong,

[45774] F. H.; Lamberts, S. W. J.: Human adrenocorticotropin-secreting pituitary
[45775] adenomas show frequent loss of heterozygosity at the glucocorticoid
[45776] receptor gene locus. J. Clin. Endocr. Metab. 83: 917-921, 1998.
[45777] 6979.

[45778] Huizenga, N. A. T. M.; De Lange, P.; Koper, J. W.; De Herder,
[45779] W. W.; Abs, R.; Kasteren, J. H. L. M. V.; De Jong, F. H.; Lamberts,
[45780] S. W. J.: Five patients with biochemical and/or clinical generalized
[45781] glucocorticoid resistance without alterations in the glucocorticoid
[45782] receptor gene. J. Clin. Endocr. Metab. 85: 2076-2081, 2000.
[45783] 6980.

[45784] Huizenga, N. A. T. M.; Koper, J. W.; de Lange, P.; Pols, H. A.
[45785] P.; Stolk, R. P.; Burger, H.; Grobbee, D. E.; Brinkmann, A. O.; de

[45786] Jong, F. H.; Lamberts, S. W. J.: A polymorphism in the glucocorticoid

[45787] receptor gene may be associated with an increased sensitivity to glucocorticoids

[45788] in vivo. J. Clin. Endocr. Metab. 83: 144-151, 1998.

[45789] 6981.

[45790] Hurley, D. M.; Accili, D.; Stratakis, C. A.; Karl, M.; Vamvakopoulos,

[45791] N.; Rorer, E.; Constantine, K.; Taylor, S. I.; Chrousos, G. P.: Point

[45792] mutation causing a single amino acid substitution in the hormone binding

[45793] domain of the glucocorticoid receptor in familial glucocorticoid resistance. J.

[45794] Clin. Invest. 87: 680-686, 1991.

[45795] 6982.

[45796] Iida, S.; Gomi, M.; Moriwaki, K.; Itoh, Y.; Hirobe, K.; Matsuzawa,

[45797] Y.; Katagiri, S.; Yonezawa, T.; Tarui, S.: Primary cortisol resistance

[45798] accompanied by a reduction in glucocorticoid receptors in two members

[45799] of the same family. J. Clin. Endocr. Metab. 60: 967-971, 1985.

[45800] 6983.

[45801] Pierce, E. A.; Dame, M. C.; Bouillon, R.; Van Baelen, H.; DeLuca,

[45802] H. F.: Monoclonal antibodies to human vitamin D-binding protein. Proc.

[45803] Nat. Acad. Sci. 82: 8429-8433, 1985.

[45804] 6984.

[45805] Schoentgen, F.; Metz-Boutigue, M.-H.; Jolles, J.; Constans, J.;

[45806] Jolles, P.: Homology between the human vitamin D-binding protein

[45807] (group specific component), alpha-fetoprotein and serum albumin. FEBS

[45808] Lett. 185: 47-50, 1985.

[45809] 6985.

[45810] Seppala, M.; Ruoslahti, E.; Makela, O.: Inheritance and genetic

[45811] linkage of Gc and TF groups. Acta Genet. Statist. Med. 17: 47-54,

[45812] 1967.

[45813] 6986.

[45814] Shibata, T.; Abe, T.: Linkage between the loci for serum albumin

[45815] and vitamin D binding protein (GC) in the Japanese quail. Animal

[45816] Genet. 27: 195-197, 1996.

[45817] 6987.

[45818] Svasti, J.; Kurosky, A.; Bennett, A.; Bowman, B. H.: Molecu-lar

[45819] basis for the three major forms of human serum vitamin D binding protein

[45820] (group-specific component). Biochemistry 18: 1611-1617, 1979.

[45821] 6988.

[45822] Szathmary, E. J. E.: The effect of Gc genotype on fasting insulin

[45823] level in Dogrib Indians. Hum. Genet. 75: 368-372, 1987.

[45824] 6989.

[45825] Thymann, M.; Hjalmarsson, K.; Svensson, M.: Five new Gc variants

[45826] detected by isoelectric focusing in agarose gel. Hum. Genet. 60:
[45827] 340-343, 1982.

[45828] 6990.

[45829] Vavrusa, B.; Cleve, H.; Constans, J.: A deficiency mutant of
[45830] the Gc system. Hum. Genet. 65: 102-107, 1983.

[45831] 6991.

[45832] Weitkamp, L. R.: Comparative gene mapping: linkage be-
tween the
[45833] albumin and Gc loci in the horse. (Abstract) Am. J. Hum. Genet. 30:
[45834] 128A only, 1978.

[45835] 6992.

[45836] Weitkamp, L. R.: Concerning the linkage relationships of
the
[45837] Gc and MNSs loci. Hum. Genet. 43: 215-220, 1978.

[45838] 6993.

[45839] Weitkamp, L. R.; Rucknagel, D. L.; Gershowitz, H.: Genetic
linkage
[45840] between structural loci for albumin and group specific

component (Gc). Am.
[45841] J. Hum. Genet. 18: 559-571, 1966.

[45842] 6994.

[45843] Witke, F. W.; Gibbs, P. E. M.; Zielinski, R.; Yang, F.; Bow-
man,
[45844] B. H.; Dugaiczyk, A.: Complete structure of the human Gc
gene: differences
[45845] and similarities between members of the albumin gene
family. Genomics 16:
[45846] 751-754, 1993.

[45847] 6995.

[45848] Yamamoto, N.; Homma, S.: Vitamin D-3 binding protein
(group-specific
[45849] component) is a precursor for the macrophage-activating
signal factor
[45850] from lysophosphatidylcholine-treated lymphocytes. Proc.
Nat. Acad.
[45851] Sci. 88: 8539-8543, 1991.

[45852] 6996.

[45853] Yamamoto, Y.; Nishimoto, H.; Ikemoto, S.: Interstitial
deletion

[45854] of the proximal long arm of chromosome 4 associated with father-child
[45855] incompatibility within the Gc-system: probable reduced gene dosage
[45856] effect and partial piebald trait. Am. J. Med. Genet. 32: 520-523,
[45857] 1989.

[45858] 6997.

[45859] Yang, F.; Bergeron, J. M.; Linehan, L. A.; Lalley, P. A.; Sak-aguchi,
[45860] A. Y.; Bowman, B. H.: Mapping and conservation of the group-specific
[45861] component gene in mouse. Genomics 7: 509-516, 1990.

[45862] 6998.

[45863] Yang, F.; Brune, J. L.; Naylor, S. L.; Cupples, R. L.; Naber-haus,
[45864] K. H.; Bowman, B. H.: Human group-specific component (Gc) is a member
[45865] of the albumin family. Proc. Nat. Acad. Sci. 82: 7994-7998, 1985.
[45866] 6999.

[45867] Yang, F.; Luna, V. J.; McAnelly, R. D.; Naberhaus, K. H.; Cupples,
[45868] R. L.; Bowman, B. H.: Evolutionary and structural relationships among
[45869] the group-specific component, albumin and alpha-fetoprotein. Nucleic
[45870] Acids Res. 13: 8007-8017, 1985.

[45871] 7000.

[45872] Yasuda, T.; Ikehara, Y.; Takagi, S.; Mizuta, K.; Kishi, K.: A

[45873] hereditary double double-banded variation in the vitamin D-binding

[45874] protein (GC) system analyzed by immunoblotting: duplication of the

[45875] 1F and 1A2 genes? Hum. Genet. 82: 89-91, 1989.

[45876] 7001.

[45877] Boguszewski, C. L.; Svensson, P.-A.; Jansson, T.; Clark, R.; Carlsson,

[45878] L. M. S.; Carlsson, B.: Cloning of two novel growth hormone transcripts

[45879] expressed in human placenta. J. Clin. Endocr. Metab. 83: 2878-2885,

[45880] 1998.

[45881] 7002.

[45882] Chen, E. Y.; Liao, Y.-C.; Smith, D. H.; Barrera-Saldana, H. A.;

[45883] Gelinas, R. E.; Seeburg, P. H.: The human growth hormone locus: nucleotide

[45884] sequence, biology, and evolution. Genomics 4: 479-497,

1989.

[45885] 7003.

[45886] Frankenne, F.; Rentier-Delrue, F.; Scippo, M.-L.; Martial, J.;

[45887] Hennen, G.: Expression of the growth hormone variant gene in human

[45888] placenta. J. Clin. Endocr. Metab. 64: 635-637, 1987.

[45889] 7004.

[45890] Lewis, U. J.; Dunn, J. T.; Bonewald, L. F.; Seavey, B. K.; VanderLaan,

[45891] W. P.: A naturally occurring structural variant of human growth hormone. J.

[45892] Biol. Chem. 253: 2679-2687, 1978.

[45893] 7005.

[45894] Liebhaber, S. A.; Urbanek, M.; Ray, J.; Tuan, R. S.; Cooke, N.

[45895] E.: Characterization and histologic localization of human growth

[45896] hormone-variant gene expression in the placenta. J. Clin. Invest. 83:

[45897] 1985-1991, 1989.

[45898] 7006.

[45899] MacLeod, J. N.; Liebhaber, S. A.; MacGillivray, M. H.; Cooke, N.
[45900] E.: Identification of a splice-site mutation in the human growth
[45901] hormone-variant gene. Am. J. Hum. Genet. 48: 1168-1174, 1991.
[45902] 7007.

[45903] MacLeod, J. N.; Worsley, I.; Ray, J.; Friesen, H. G.; Lieb-haber,
[45904] S. A.; Cooke, N. E.: Human growth hormone variant is a biologically
[45905] active somatogen and lactogen. Endocrinology 128: 1298-1302, 1991.
[45906] 7008.

[45907] Owerbach, D.; Rutter, W. J.; Martial, J. A.; Baxter, J. D.; Shows,
[45908] T. B.: Genes for growth hormone, chorionic somatomam-motropin and
[45909] growth hormone-like genes on chromosome 17 in hu-mans. Science 209:
[45910] 289-292, 1980.

[45911] 7009.

[45912] Harada, H.; Fujita, T.; Miyamoto, M.; Kimura, Y.; Maruyama, M.;

[45913] Furia, A.; Miyata, T.; Taniguchi, T.: Structurally similar but functionally

[45914] distinct factors, IRF-1 and IRF-2, bind to the same regulatory elements

[45915] of IFN and IFN-inducible genes. Cell 58: 729-739, 1989.

[45916] 7010.

[45917] Digilio, M. C.; Conti, E.; Sarkozy, A.; Mingarelli, R.; Dottorini,

[45918] T.; Marino, B.; Pizzuti, A.; Dallapiccola, B.: Grouping of multiple-lentigines/LEOPARD

[45919] and Noonan syndromes on the PTPN11 gene. Am. J. Hum. Genet. 71:

[45920] 389-394, 2002.

[45921] 7011.

[45922] Guo, N.; Mogues, T.; Weremowicz, S.; Morton, C. C.; Sastry, K.

[45923] N.: The human ortholog of rhesus mannose-binding protein-A gene is

[45924] an expressed pseudogene that localizes to chromosome 10. Mammalian

[45925] Genome 9: 246-249, 1998.

[45926] 7012.

[45927] Moreau, J.-F.; Donaldson, D. D.; Bennett, F.; Witek-Gi-
annotti,
[45928] J.; Clark, S. C.; Wong, G. C.: Leukaemia inhibitory factor is
identical
[45929] to the myeloid growth factor human interleukin for DA
cells. Nature 336:
[45930] 690-692, 1988.

[45931] 7013.

[45932] Nakashima, K.; Yanagisawa, M.; Arakawa, H.; Kimura, N.;
Hisatsune,
[45933] T.; Kawabata, M.; Miyazono, K.; Taga, T.: Synergistic sig-
naling in
[45934] fetal brain by STAT3-Smad1 complex bridged by p300.
Science 284:
[45935] 479-482, 1999.

[45936] 7014.

[45937] Patterson, P. H.: Leukemia inhibitory factor, a cytokine at
the
[45938] interface between neurobiology and immunology. Proc.

Nat. Acad. Sci. 91:
[45939] 7833-7835, 1994.

[45940] 7015.

[45941] Sutherland, G. R.; Baker, E.; Hyland, V. J.; Callen, D. F.; Stahl,
[45942] J.; Gough, N. M.: The gene for human leukemia inhibitory factor (LIF)
[45943] maps to 22q12. Leukemia 3: 9-13, 1989.

[45944] 7016.

[45945] Williams, R. L.; Hilton, D. J.; Pease, S.; Willson, T. A.; Stew-art,
[45946] C. L.; Gearing, D. P.; Wagner, E. F.; Metcalf, D.; Nicola, N. A.;
[45947] Gough, N. M.: Myeloid leukaemia inhibitory factor main-tains the developmental
[45948] potential of embryonic stem cells. Nature 336: 684-687, 1988.

[45949] 7017.

[45950] Bae, J.; Leo, C. P.; Hsu, S. Y.; Hsueh, A. J. W.: MCL-1S, a splicing
[45951] variant of the antiapoptotic BCL-2 family member MCL-1,

encodes a
[45952] proapoptotic protein possessing only the BH3 domain. J. Biol. Chem. 275:
[45953] 25255-25261, 2000.

[45954] 7018.

[45955] Craig, R. W.; Jabs, E. W.; Zhou, P.; Kozopas, K. M.; Hawkins, A.
[45956] L.; Rochelle, J. M.; Seldin, M. F.; Griffin, C. A.: Human and mouse
[45957] chromosomal mapping of the myeloid cell leukemia-1 gene: MCL1 maps
[45958] to human chromosome 1q21, a region that is frequently altered in preneoplastic
[45959] and neoplastic disease. Genomics 23: 457-463, 1994.

[45960] 7019.

[45961] Kozopas, K. M.; Yang, T.; Buchan, H. L.; Zhou, P.; Craig, R. W.
[45962] : MCL1, a gene expressed in programmed myeloid cell differentiation,
[45963] has sequence similarity to BCL2. Proc. Nat. Acad. Sci. 90: 3516-3520,
[45964] 1993.

[45965] 7020.

[45966] Rinkenberger, J. L.; Horning, S.; Klocke, B.; Roth, K.; Ko-
rsmeyer,

[45967] S. J.: Mcl-1 deficiency results in peri-implantation embry-
onic lethality. Genes

[45968] Dev. 14: 23-27, 2000.

[45969] 7021.

[45970] Hizawa, N.; Yamaguchi, E.; Furuya, K.; Ohnuma, N.; Ko-
dama, N.;

[45971] Kojima, J.; Ohe, M.; Kawakami, Y.: Association between
high serum

[45972] total IgE levels and D11S97 on chromosome 11q13 in
Japanese subjects. J.

[45973] Med. Genet. 32: 363-369, 1995.

[45974] 7022.

[45975] Sandford, A. J.; Shirakawa, T.; Moffatt, M. F.; Daniels, S. E.;

[45976] Ra, C.; Faux, J. A.; Young, R. P.; Nakamura, Y.; Lathrop, G.
M.; Cookson,

[45977] W. O. C. M.; Hopkin, J. M.: Localisation of atopy and beta
subunit

[45978] of high-affinity IgE receptor (FCER1) on chromosome 11q.

Lancet 341:
[45979] 332-334, 1993.

[45980] 7023.

[45981] Harada, H.; Kondo, T.; Ogawa, S.; Tamura, T.; Kitagawa, M.; Tanaka,
[45982] N.; Lamphier, M. S.; Hirai, H.; Taniguchi, T.: Accelerated exon skipping
[45983] of IRF-1 mRNA in human myelodysplasia/leukemia: a possible mechanism
[45984] of tumor suppressor inactivation Oncogene 9:
3313-3320, 1994.

[45985] 7024.

[45986] Harada, H.; Willison, K.; Sakakibara, J.; Miyamoto, M.; Fu-jita,
[45987] T.; Taniguchi, T.: Absence of the type I IFN system in EC cells:
[45988] transcriptional activator (IRF-1) and repressor (IRF-2) genes are
[45989] developmentally regulated. Cell 63: 303-312, 1990.

[45990] 7025.

[45991] Itoh, S.; Harada, H.; Nakamura, Y.; White, R.; Taniguchi, T.:

[45992] Assignment of the human interferon regulatory factor-1 (IRF1) gene

[45993] to chromosome 5q23-q31. Genomics 10: 1097-1099, 1991.

[45994] 7026.

[45995] Ko, J.; Gendron-Fitzpatrick, A.; Splitter, G. A.: Susceptibil-ity

[45996] of IFN regulatory factor-1 and IFN consensus sequence binding protein-deficient

[45997] mice to brucellosis. J. Immun. 168: 2433-2440, 2002.

[45998] 7027.

[45999] Miyamoto, M.; Fujita, T.; Kimura, Y.; Maruyama, M.; Harada, H.;

[46000] Sudo, Y.; Miyata, T.; Taniguchi, T.: Regulated expression of a gene

[46001] encoding a nuclear factor, IRF-1, that specifically binds to IFN-beta

[46002] gene regulatory elements. Cell 54: 903-913, 1988.

[46003] 7028.

[46004] Nozawa, H.; Oda, E.; Ueda, S.; Tamura, G.; Maesawa, C.; Muto, T.;

[46005] Taniguchi, T.; Tanaka, N.: Functionally inactivating point mutation

[46006] in the tumor-suppressor IRF-1 gene identified in human gastric cancer. Int.

[46007] J. Cancer 77: 522-527, 1998.

[46008] 7029.

[46009] Tamura, G.; Sakata, K.; Nishizuka, S.; Maesawa, C.; Suzuki, Y.;

[46010] Terashima, M.; Eda, Y.; Satodate, R.: Allelotype of adenoma and differentiated

[46011] adenocarcinoma of the stomach. J. Path. 180: 371-377, 1996.

[46012] 7030.

[46013] Willman, C. L.; Sever, C. E.; Pallavicini, M. G.; Harada, H.;

[46014] Tanaka, N.; Slovak, M. L.; Yamamoto, H.; Harada, K.; Meeker, T. C.;

[46015] List, A. F.; Taniguchi, T.: Deletion of IRF-1, mapping to chromosome

[46016] 5q31.1, in human leukemia and preleukemic myelodysplasia. Science 259:

[46017] 968-971, 1993.

[46018] 7031.

[46019] Yamada, G.; Ogawa, M.; Akagi, K.; Miyamoto, H.; Nakano, N.; Itoh,

[46020] S.; Miyazaki, J.; Nishikawa, S.; Yamamura, K.; Taniguchi, T.: Specific

[46021] depletion of the B-cell population induced by aberrant ex-pression

[46022] of human interferon regulatory factor 1 gene in transgenic mice. Proc.

[46023] Nat. Acad. Sci. 88: 532-536, 1991.

[46024] 7032.

[46025] Harada, H.; Kitagawa, M.; Tanaka, N.; Yamamoto, H.; Harada, K.;

[46026] Ishihara, M.; Taniguchi, T.: Anti-oncogenic and oncogenic potentials

[46027] of interferon regulatory factors-1 and -2. Science 259: 971-974,

[46028] 1993.

[46029] 7033.

[46030] Harada, H.; Takahashi, E.-I.; Itoh, S.; Harada, K.; Hori, T.-A.;

[46031] Taniguchi, T.: Structure and regulation of the human interferon regulatory
[46032] factor 1 (IRF-1) and IRF-2 genes: implications for a gene network
[46033] in the interferon system. Molec. Cell. Biol. 14: 1500-1509, 1994.
[46034] 7034.

[46035] Hida, S.; Ogasawara, K.; Sato, K.; Abe, M.; Takayanagi, H.; Yokochi,
[46036] T.; Sato, T.; Hirose, S.; Shirai, T.; Taki, S.; Taniguchi, T.: CD8+
[46037] T cell-mediated skin disease in mice lacking IRF-2, the transcriptional
[46038] attenuator of interferon-alpha/beta signaling. Immunity 13: 643-655,
[46039] 2000.

[46040] 7035.

[46041] Nishio, Y.; Noguchi, E.; Ito, S.; Ichikawa, E.; Umebayashi, Y.;
[46042] Otsuka, F.; Arinami, T.: Mutation and association analysis of the
[46043] interferon regulatory factor 2 gene (IRF2) with atopic der-

matitis. J.

[46044] Hum. Genet. 46: 664-667, 2001.

[46045] 7036.

[46046] Guo, B.; Yu, Y.; Leibold, E. A.: Iron regulates cytoplasmic levels

[46047] of a novel iron-responsive element-binding protein without aconitase

[46048] activity. J. Biol. Chem. 269: 24252-24260, 1994.

[46049] 7037.

[46050] Henderson, B. R.; Seiser, C.; Kuhn, L. C.: Characterization of

[46051] a second RNA-binding protein in rodents with specificity for iron-responsive

[46052] elements. J. Biol. Chem. 268: 27327-27334, 1993.

[46053] 7038.

[46054] Hentze, M. W.; Kuhn, L. C.: Molecular control of vertebrate iron

[46055] metabolism: mRNA-based regulatory circuits operated by iron, nitric

[46056] oxide, and oxidative stress. Proc. Nat. Acad. Sci. 93: 8175-8182,

[46057] 1996.

[46058] 7039.

[46059] Iwai, K.; Klausner, R. D.; Rouault, T. A.: Requirements for iron-regulated

[46060] degradation of the RNA binding protein, iron regulatory protein 2. EMBO

[46061] J. 14: 5350-5357, 1995.

[46062] 7040.

[46063] Rouault, T. A.; Tang, C. K.; Kaptain, S.; Burgess, W. H.; Haile,

[46064] D. J.; Samaniego, F.; McBride, O. W.; Harford, J. B.; Klausner, R.

[46065] D.: Cloning of the cDNA encoding an RNA regulatory protein: the human

[46066] iron-responsive element-binding protein. Proc. Nat. Acad. Sci. 87:

[46067] 7958-7962, 1990.

[46068] 7041.

[46069] LaVaute, T.; Smith, S.; Cooperman, S.; Iwai, K.; Land, W.; Meyron-Holtz,

[46070] E.; Drake, S. K.; Miller, G.; Abu-Asab, M.; Tsokos, M.;

Switzer, R.,
[46071] III; Grinberg, A.; Love, P.; Tresser, N.; Rouault, T. A.: Targeted
[46072] deletion of the gene encoding iron regulatory protein-2 causes misregulation
[46073] of iron metabolism and neurodegenerative disease in mice. Nature
[46074] Genet. 27: 209-214, 2001.

[46075] 7042.

[46076] Samaniego, F.; Chin, J.; Iwai, K.; Rouault, T. A.; Klausner, R.
[46077] D.: Molecular characterization of a second iron-responsive element
[46078] binding protein, iron regulatory protein 2: structure, function, and
[46079] post-translational regulation. J. Biol. Chem. 269: 30904-30910,
[46080] 1994.

[46081] 7043.

[46082] Kao, W. W.-Y.; Liu, C.-Y.; Converse, R. L.; Shiraishi, A.; Kao,
[46083] C. W.-C.; Ishizaki, M.; Doetschman, T.; Duffy, J.: Keratin

12-deficient
[46084] mice have fragile corneal epithelia. Invest. Ophthal. Vis. Sci. 37:
[46085] 2572-2584, 1996.

[46086] 7044.

[46087] Moll, R.; Franke, W. W.; Schiller, D. L.; Geiger, B.; Krepler,

[46088] R.: The catalog of human cytokeratins: patterns of ex-pression in
[46089] normal epithelia, tumors and cultured cells. Cell 31: 11-24, 1982.

[46090] 7045.

[46091] Raimondi, E.; Moralli, D.; De Carli, L.; Ceratto, N.; Balzaretti,
[46092] M.; Leube, R.; Collin, C.; Romano, V.: Assignment of the human cytokeratin
[46093] 3 gene (KRT3) to 12q12-q13 by FISH. Cytogenet. Cell Genet. 66: 162-163,
[46094] 1994.

[46095] 7046.

[46096] Schermer, A.; Galvin, S.; Sun, T.-T.: Differentiation-related

[46097] expression of a major 64K corneal keratin in vivo and in

culture suggests
[46098] limbal location of corneal epithelial stem cells. J. Cell Biol. 103:
[46099] 49-62, 1986.

[46100] 7047.

[46101] Tremblay, M.; Dube, I.: Meesmann's corneal dystrophy: ultrastructural
[46102] features. Canad. J. Ophthal. 17: 24-28, 1982.

[46103] 7048.

[46104] Casanova, M. L.; Bravo, A.; Ramirez, A.; Morreale de Escobar, G.;
[46105] Were, F.; Merlino, G.; Vidal, M.; Jorcano, J. L.: Exocrine pancreatic
[46106] disorders in transsgenic (sic) mice expressing human keratin 8. J.
[46107] Clin. Invest. 103: 1587-1595, 1999.

[46108] 7049.

[46109] Jackson, B. W.; Grund, C.; Schmid, E.; Burke, K.; Franke, W.; Illmensee,
[46110] K.: Formation of cytoskeletal elements during mouse embryogenesis:

[46111] intermediate filaments of the cytokeratin type and desmosomes in preimplantation
[46112] embryos. Differentiation 17: 161-179, 1980.

[46113] 7050.

[46114] Ku, N.-O.; Gish, R.; Wright, T. L.; Omary, M. B.: Keratin 8 mutations
[46115] in patients with cryptogenic liver disease. New Eng. J. Med. 344:
[46116] 1580-1587, 2001.

[46117] 7051.

[46118] Ku, N.-O.; Wright, T. L.; Terrault, N. A.; Gish, R.; Omary, M.
[46119] B.: Mutation of human keratin 18 in association with cryptogenic
[46120] cirrhosis. J. Clin. Invest. 99: 19-23, 1997.

[46121] 7052.

[46122] Waseem, A.; Alexander, C. M.; Steel, J. B.; Lane, E. B.: Embryonic
[46123] simple epithelial keratins 8 and 18: chromosomal location emphasizes
[46124] difference from other keratin pairs. New Biologist 2:

464-478, 1990.

[46125] 7053.

[46126] Yamamoto, R.; Kao, L.-C.; McKnight, C. E.; Strauss, J. F., III

[46127] : Cloning and sequence of cDNA for human placental cy-

tokeratin 8:

[46128] regulation of the mRNA in trophoblastic cells by cAMP.

Molec. Endocr. 4:

[46129] 370-374, 1990.

[46130] 7054.

[46131] Dunham, I.; Sargent, C. A.; Trowsdale, J.; Campbell, R. D.:

Molecular

[46132] mapping of the human major histocompatibility complex

by pulsed-field

[46133] gel electrophoresis. Proc. Nat. Acad. Sci. 84: 7237-7241,

1987.

[46134] 7055.

[46135] Susa, S.; Daimon, M.; Kondo, H.; Kondo, M.; Yamatani, K.;

Sasaki,

[46136] H.: Identification of a novel mutation of the CPO gene in a

Japanese

[46137] hereditary coproporphyria family. Am. J. Med. Genet. 80:

204-206,

[46138] 1998.

[46139] 7056.

[46140] Taketani, S.; Kohno, H.; Furukawa, T.; Yoshinaga, T.; Tokunaga,

[46141] R.: Molecular cloning, sequencing and expression of cDNA encoding

[46142] human coproporphyrinogen oxidase. Biochim. Biophys. Acta 1183:

[46143] 547-549, 1994.

[46144] 7057.

[46145] Wiman, A.; Floderus, Y.; Harper, P.: Two novel mutations and

[46146] coexistence of the 991C-T and the 1339C-T mutation on a single allele

[46147] in the coproporphyrinogen oxidase gene in Swedish pa- tients with hereditary

[46148] coproporphyria. J. Hum. Genet. 47: 407-412, 2002.

[46149] 7058.

[46150] Kogan, S. C.; Lagasse, E.; Atwater, S.; Bae, S.; Weissman, I.;

[46151] Ito, Y.; Bishop, J. M.: The PEBP2-beta-MYH11 fusion cre-

ated by inv(16)(p13;q22)

[46152] in myeloid leukemia impairs neutrophil maturation and contributes

[46153] to granulocytic dysplasia. Proc. Nat. Acad. Sci. 95: 11863-11868,

[46154] 1998.

[46155] 7059.

[46156] Liu, P.; Tarle, S. A.; Hajra, A.; Claxton, D. F.; Marlton, P.;

[46157] Freedman, M.; Siciliano, M. J.; Collins, F. S.: Fusion be-

[46158] tween transcription factor CBF-beta/PEBP2-beta and a myosin heavy chain in acute myeloid

[46159] leukemia. Science 261: 1041-1044, 1993.

[46160] 7060.

[46161] Liu, P. P.; Hajra, A.; Wijmenga, C.; Collins, F. S.: Molecular

[46162] pathogenesis of the chromosome 16 inversion in the M4Eo subtype of

[46163] acute myeloid leukemia. Blood 85: 2289-2302, 1995.

[46164] 7061.

[46165] Lutterbach, B.; Hou, Y.; Durst, K. L.; Hiebert, S. W.: The

inv(16)
[46166] encodes an acute myeloid leukemia 1 transcriptional corepressor. Proc.
[46167] Nat. Acad. Sci. 96: 12822-12827, 1999.

[46168] 7062.

[46169] O'Reilly, J.; Chipper, L.; Springall, F.; Herrmann, R.: A unique
[46170] structural abnormality of chromosome 16 resulting in a CBF-beta-MYH11
[46171] fusion transcript in a patient with acute myeloid leukemia, FAB M4. Cancer
[46172] Genet. Cytogenet. 121: 52-55, 2000.

[46173] 7063.

[46174] Ogawa, E.; Inuzuka, M.; Maruyama, M.; Satake, M.; Naito-Fujimoto,
[46175] M.; Ito, Y.; Shigesada, K.: Molecular cloning and character-ization
[46176] of PEBP2-beta, the heterodimeric partner of a novel Drosophila runt-related
[46177] DNA binding protein PEBP2-alpha. Virology 194: 314-331, 1993.
[46178] 7064.

[46179] Wang, S.; Wang, Q.; Crute, B. E.; Melnikova, I. N.; Keller, S.

[46180] R.; Speck, N. A.: Cloning and characterization of subunits of the

[46181] T-cell receptor and murine leukemia virus enhancer core-binding factor. Molec.

[46182] Cell. Biol. 13: 3324-3339, 1993.

[46183] 7065.

[46184] Pellegata, N. S.; Dieguez-Lucena, J. L.; Joensuu, T.; Lau, S.;

[46185] Montgomery, K. T.; Krahe, R.; Kivela, T.; Kucherlapati, R.; Forsius,

[46186] H.; de la Chapelle, A.: Mutations in KERA, encoding keratocan, cause

[46187] cornea plana. Nature Genet. 25: 91-95, 2000.

[46188] 7066.

[46189] Brooks, B. A.; McBride, O. W.; Dolphin, C. T.; Farrall, M.; Scambler,

[46190] P. J.; Gonzalez, F. J.; Idle, J. R.: The gene CYP3 encoding P450PCN1

[46191] (nifedipine oxidase) is tightly linked to the gene COL1A2 encoding

[46192] collagen type 1 alpha on 7q21-q22.1. Am. J. Hum. Genet.

43: 280-284,

[46193] 1988.

[46194] 7067.

[46195] Chen, H.; Sandler, D. P.; Taylor, J. A.; Shore, D. L.; Liu, E.;

[46196] Bloomfield, C. D.; Bell, D. A.: Increased risk for myelodys-
plastic

[46197] syndromes in individuals with glutathione transferase
theta 1 (GSTT1)

[46198] gene defect. Lancet 347: 295-297, 1996.

[46199] 7068.

[46200] Daly, A. K.; Salh, B. S.; Bilton, D.; Allen, J.; Knight, A. D.;

[46201] Webb, A. K.; Braganza, J. M.; Idle, J. R.: Deficient nifedipine
oxidation:

[46202] a rare inherited trait associated with cystic fibrosis kin-
dreds. Pharmacogenetics 2:

[46203] 19-24, 1992.

[46204] 7069.

[46205] Elshourbagy, N. A.; Guzelian, P. S.: Separation, purifica-
tion,

[46206] and characterization of a novel form of hepatic cy-

tochrome P-450 from
[46207] rats treated with pregnenolone-16-alpha-carbonitrile. J. Biol. Chem. 255:
[46208] 1279-1285, 1980.

[46209] 7070.

[46210] Felix, C. A.; Walker, A. H.; Lange, B. J.; Williams, T. M.; Winick,
[46211] N. J.; Cheung, N.-K. V.; Lovett, B. D.; Nowell, P. C.; Blair, I. A.;
[46212] Rebbeck, T. R.: Association of CYP3A4 genotype with treatment-related
[46213] leukemia. Proc. Nat. Acad. Sci. 95: 13176-13181, 1998.

[46214] 7071.

[46215] Forrester, L. M.; Neal, G. E.; Judah, D. J.; Glancey, M. J.; Wolf,
[46216] C. R.: Evidence for involvement of multiple forms of cytochrome P-450
[46217] in aflatoxin B(1) metabolism in human liver. Proc. Nat. Acad. Sci. 87:
[46218] 8306-8310, 1990.

[46219] 7072.

[46220] Gonzalez, F. J.; Schmid, B. J.; Umeno, M.; McBride, O. W.; Hardwick,

[46221] J. P.; Meyer, U. A.; Gelboin, H. V.; Idle, J. R.: Human P450PCN1:

[46222] sequence, chromosome localization, and direct evidence through cDNA

[46223] expression that P450PCN1 is nifedipine oxidase. DNA 7: 79-86, 1988.

[46224] 7073.

[46225] Hoyo-Vadillo, C.; Castaneda-Hernandez, G.; Herrera, J. E.; Vidal-Garate,

[46226] J.; Moreno-Ramos, A.; Chavez, F.; Hong, E.: Pharmacoki-netics of nifedipine

[46227] slow release tablet in Mexican subjects: further evidence for an oxidation

[46228] polymorphism. J. Clin. Pharm. 29: 816-820, 1989.

[46229] 7074.

[46230] Inoue, K.; Inazawa, J.; Nakagawa, H.; Shimada, T.; Ya-mazaki, H.;

[46231] Guengerich, F. P.; Abe, T.: Assignment of the human cy-tochrome P-450

[46232] nifedipine oxidase gene (CYP3A4) to chromosome 7 at

band q22.1 by
[46233] fluorescence in situ hybridization. Jpn. J. Hum. Genet. 37: 133-138,
[46234] 1992.

[46235] 7075.

[46236] Kittles, R. A.; Chen, W.; Panguluri, R. K.; Ahaghotu, C.; Jackson,
[46237] A.; Adebamowo, C. A.; Griffin, R.; Williams, T.; Ukoli, F.; Adams-Campbell,
[46238] L.; Kwagyan, J.; Isaacs, W.; Freeman, V.; Dunston, G. M.: CYP3A4-V
[46239] and prostate cancer in African Americans: causal or confounding association
[46240] because of population stratification? Hum. Genet. 110: 553-560,
[46241] 2002.

[46242] 7076.

[46243] Kleinbloesem, C. H.; van Brummelen, P.; Faber, H.; Danhof, M.;
[46244] Vermeulen, N. P. E.; Breimer, D. D.: Variability in nifedipine pharmacokinetics
[46245] and dynamics: a new oxidation polymorphism in man.

Biochem. Pharm. 33:
[46246] 3721-3724, 1984.

[46247] 7077.

[46248] Lehmann, J. M.; McKee, D. D.; Watson, M. A.; Willson, T. M.; Moore,
[46249] J. T.; Kliewer, S. A.: The human orphan nuclear receptor PXR is activated
[46250] by compounds that regulate CYP3A4 gene expression and cause drug interactions. J.
[46251] Clin. Invest. 102: 1016-1023, 1998.

[46252] 7078.

[46253] Lown, K. S.; Bailey, D. G.; Fontana, R. J.; Janardan, S. K.; Adair,
[46254] C. H.; Fortlage, L. A.; Brown, M. B.; Guo, W.; Watkins, P. B.: Grapefruit
[46255] juice increases felodipine oral availability in humans by decreasing
[46256] intestinal CYP3A protein expression. J. Clin. Invest. 99: 2545-2553,
[46257] 1997.

[46258] 7079.

[46259] Molowa, D. T.; Schuetz, E. G.; Wrighton, S. A.; Watkins, P. B.;

[46260] Kremers, P.; Mendez-Picon, G.; Parker, G. A.; Guzelian, P. S.: Complete

[46261] cDNA sequence of a cytochrome P-450 inducible by glu-

cocorticoids in

[46262] human liver. Proc. Nat. Acad. Sci. 83: 5311-5315, 1986.

[46263] 7080.

[46264] Paris, P. L.; Kupelian, P. A.; Hall, J. M.; Williams, T. L.; Levin,

[46265] H.; Klein, E. A.; Casey, G.; Witte, J. S.: Association between a

[46266] CYP3A4 genetic variant and clinical presentation in African-American

[46267] prostate cancer patients. Cancer Epidemiol. Biomarkers Prev. 8:

[46268] 901-905, 1999.

[46269] 7081.

[46270] Rebbeck, T. R.; Jaffe, J. M.; Walker, A. H.; Wein, A. J.; Malkowicz,

[46271] S. B.: Modification of clinical presentation of prostate tu-

mors by

[46272] a novel genetic variant in CYP3A4. J. Nat. Cancer Inst. 90: 1225-1229,

[46273] 1998.

[46274] 7082.

[46275] Renwick, A. G.; Robertson, D. R. C.; Macklin, B.; Challenor, V.;

[46276] Waller, D. G.; George, C. F.: The pharmacokinetics of oral nifedipine--a

[46277] population study. Brit. J. Clin. Pharm. 25: 701-708, 1988.

[46278] 7083.

[46279] Spielberg, S. P.: Personal Communication. Toronto, On-tario,

[46280] Canada 2/26/1988.

[46281] 7084.

[46282] Thum, T.; Borlak, J.: Gene expression in distinct regions of

[46283] the heart. Lancet 355: 979-983, 2000.

[46284] 7085.

[46285] Chen, Y.; Faraco, J.; Yin, W.; Germiller, J.; Francke, U.; Bonadio,

[46286] J.: Structure, chromosomal localization, and expression

pattern of
[46287] the murine Magp gene. J. Biol. Chem. 268: 27381-27389, 1993.
[46288] 7086.

[46289] Faraco, J.; Bashir, M.; Rosenbloom, J.; Francke, U.: Characterization
[46290] of the human gene for microfibril-associated glycoprotein (MFAP2),
[46291] assignment to chromosome 1p36.1-p35, and linkage to D1S170. Genomics 25:
[46292] 630-637, 1995.

[46293] 7087.

[46294] Gibson, M. A.; Hughes, J. L.; Fanning, J. C.; Cleary, E. G.: The
[46295] major antigen of elastin-associated microfibrils is a 31-kDa glycoprotein.
[46296] J. Biol. Chem. 261: 11429-11436, 1986.

[46297] 7088.

[46298] Low, F. N.: Microfibrils: fine filamentous components of the tissue
[46299] space. Anat. Rec. 142: 131-137, 1962.

[46300] 7089.

[46301] Bondurand, N.; Pingault, V.; Goerich, D. E.; Lemort, N.; Sock,

[46302] E.; Le Caignec, C.; Wegner, M.; Goossens, M.: Interaction among SOX10,

[46303] PAX3 and MITF, three genes altered in Waardenburg syndrome. Hum.

[46304] Molec. Genet. 9: 1907-1917, 2000.

[46305] 7090.

[46306] Kelavkar, U. P.; Badr, K. F.: Effects of mutant p53 expression

[46307] on human 15-lipoxygenase-promoter activity and murine 12/15-lipoxygenase

[46308] gene expression: evidence that 15-lipoxygenase is a mutator gene. Proc.

[46309] Nat. Acad. Sci. 96: 4378-4383, 1999.

[46310] 7091.

[46311] Sigal, E.; Craik, C. S.; Highland, E.; Grunberger, D.; Costello,

[46312] L. L.; Dixon, R. A. F.; Nadel, J. A.: Molecular cloning and primary

[46313] structure of human 15-lipoxygenase. Biochem. Biophys. Res. Commun. 157:

[46314] 457-464, 1988.

[46315] 7092.

[46316] Yoshimoto, T.; Suzuki, H.; Yamamoto, S.; Takai, T.; Yokoyama, C.;

[46317] Tanabe, T.: Cloning and sequence analysis of the cDNA for arachidonate

[46318] 12-lipoxygenase of porcine leukocytes. Proc. Nat. Acad. Sci. 87:

[46319] 2142-2146, 1990.

[46320] 7093.

[46321] Minoshima, S.; Fukuyama, R.; Yamamoto, T.; Shimizu, N.: Mapping

[46322] of human long-chain acyl-CoA synthetase to chromosome 4. (Abstract) Cytogenet.

[46323] Cell Genet. 58: 1888 only, 1991.

[46324] 7094.

[46325] Cantu, E. S.; Sprinkle, T. J.; Ghosh, B.; Singh, I.: The human

[46326] palmitoyl-CoA ligase (FACL2) gene maps to the chromosome 4q34-q35

[46327] region by fluorescence in situ hybridization (FISH) and somatic cell
[46328] hybrid panels. Genomics 28: 600-602, 1995.

[46329] 7095.

[46330] Adelman, J. P.; Bond, C. T.; Douglass, J.; Herbert, E.: Two mammalian
[46331] genes transcribed from opposite strands of the same DNA locus. Science 235:
[46332] 1514-1517, 1987.

[46333] 7096.

[46334] Adelman, J. P.; Mason, A. J.; Hayflick, J. S.; Seeburg, P. H.:

[46335] Isolation of the gene and hypothalamic cDNA for the common precursor
[46336] of gonadotropin-releasing hormone and prolactin release-inhibiting
[46337] factor in human and rat. Proc. Nat. Acad. Sci. 83:
179-183, 1986.

[46338] 7097.

[46339] Barni, T.; Maggi, M.; Fantoni, G.; Granchi, S.; Mancina, R.; Gulisano,
[46340] M.; Marra, F.; Macorsini, E.; Luconi, M.; Rotella, C.; Serio,

M.;

[46341] Balboni, G. C.; Vannelli, G. B.: Sex steroids and odorants modulate

[46342] gonadotropin-releasing hormone secretion in primary cultures of human

[46343] olfactory cells. J. Clin. Endocr. Metab. 84: 4266-4273, 1999.

[46344] 7098.

[46345] Casan, E. M.; Raga, F.; Bonilla-Musoles, F.; Polan, M. L.: Human

[46346] oviductal gonadotropin-releasing hormone: possible im-

[46347] plications in fertilization, early embryonic development, and implanta-

[46348] tion. J. Clin. Enodcr. Metab. 85: 1377-1381, 2000.

[46349] 7099.

[46350] Cattanach, B. M.; Iddon, C. A.; Charlton, H. M.; Chiappa, S. A.;

[46351] Fink, G.: Gonadotrophin-releasing hormone deficiency in a mutant

[46352] mouse with hypogonadism. Nature 269: 338-340, 1977.

[46353] 7100.

[46354] Dong, K.-W.; Yu, K.-L.; Chen, Z.-G.; Chen, Y.-D.; Roberts, J. L.

[46355] : Characterization of multiple promoters directing tissue-specific

[46356] expression of the human gonadotropin-releasing hormone gene. Endocrinology 138:

[46357] 2754-2762, 1997.

[46358] 7101.

[46359] Dong, K.-W.; Zheng, H.-M.; Wen, Z.-Y.; Chen, Z.-G.: The POU homeodomain

[46360] protein Oct-1 binds cis-regulatory element essential for the human

[46361] GnRH upstream promoter activity in the JEG-3 cells. J. Clin. Endocr.

[46362] Metab. 86: 2838-2844, 2001.

[46363] 7102.

[46364] Duan, W. R.; Ito, M.; Park, Y.; Maizels, E. T.; Hunzicker-Dunn,

[46365] M.; Jameson, J. L.: GnRH regulates early growth response protein

[46366] 1 transcription through multiple promoter elements. Molec. Endocr. 16:

[46367] 221-233, 2002.

[46368] 7103.

[46369] Eraly, S. A.; Nelson, S. B.; Huang, K. M.; Mellon, P. L.: Oct-1

[46370] binds promoter elements required for transcription of the GnRH gene. Molec.

[46371] Endocr. 12: 469-481, 1998.

[46372] 7104.

[46373] Ewer, R. W.: Familial monotropic pituitary gonadotropin insufficiency. J.

[46374] Clin. Endocr. 28: 783-788, 1968.

[46375] 7105.

[46376] Gibson, M. J.; Krieger, D. T.; Charlton, H. M.; Zimmerman, E.

[46377] A.; Silverman, A.-J.; Perlow, M. J.: Mating and pregnancy can occur

[46378] in genetically hypogonadal mice with preoptic area brain grafts. Science 225:

[46379] 949-951, 1984.

[46380] 7106.

[46381] Hayflick, J. S.; Adelman, J. P.; Seeburg, P. H.: The complete

[46382] nucleotide sequence of the human gonadotropin-releasing hormone gene. Nucleic

[46383] Acids Res. 17: 6403-6404, 1989.

[46384] 7107.

[46385] Kelley, C. G.; Lavorgna, G.; Clark, M. E.; Boncinelli, E.; Mellon,

[46386] P. L.: The Otx2 homeoprotein regulates expression from the gonadotropin-releasing

[46387] hormone proximal promoter. Molec. Endocr. 14: 1246-1256, 2000.

[46388] 7108.

[46389] Lawson, M. A.; Buhain, A. R.; Jovenal, J. C.; Mellon, P. L.:

[46390] Multiple factors interacting at the GATA sites of the gonadotropin-releasing

[46391] hormone neuron-specific enhancer regulate gene expression. Molec.

[46392] Endocr. 12: 364-377, 1998.

[46393] 7109.

[46394] Mason, A. J.; Pitts, S. L.; Nikolics, K.; Szonyi, E.; Wilcox,

[46395] J. N.; Seeburg, P. H.; Stewart, T. A.: The hypogonadal mouse: reproductive
[46396] functions restored by gene therapy. Science 234: 1372-1378, 1986.
[46397] 7110.

[46398] Nachtigall, L. B.; Boepple, P. A.; Pralong, F. P.; Crowley, W.

[46399] F., Jr.: Adult-onset idiopathic hypogonadotropic hypogo-nadism: a
[46400] treatable form of male infertility. New Eng. J. Med. 336: 410-415,
[46401] 1997.

[46402] 7111.

[46403] Nikolics, K.; Mason, A. J.; Szonyi, E.; Ramachandran, J.; Seeburg,
[46404] P. H.: A prolactin-inhibiting factor within the precursor for human
[46405] gonadotropin-releasing hormone. Nature 316: 511-517, 1985.
[46406] 7112.

[46407] Seeburg, P. H.; Adelman, J. P.: Characterization of cDNA for

[46408] precursor of human luteinizing hormone releasing hormone. Nature 311:
[46409] 666-668, 1984.

[46410] 7113.

[46411] Williamson, P.; Lang, J.; Boyd, Y.: The gonadotropin-releasing
[46412] hormone (Gnrh) gene maps to mouse chromosome 14 and identifies a homologous
[46413] region on human chromosome 8. Somat. Cell Molec. Genet. 17: 609-615,
[46414] 1991.

[46415] 7114.

[46416] Wolfe, A.; Kim, H. H.; Tobet, S.; Stafford, D. E. J.; Radovick,
[46417] S.: Identification of a discrete promoter region of the human GnRH
[46418] gene that is sufficient for directing neuron-specific expression:
[46419] a role for POU homeodomain transcription factors. Molec. Endocr. 16:
[46420] 435-449, 2002.

[46421] 7115.

[46422] Yang-Feng, T. L.; Seeburg, P. H.; Francke, U.: Human luteinizing
[46423] hormone-releasing hormone gene (LHRH) is located on short arm of chromosome
[46424] 8 (region 8p11.2-p21). Somat. Cell Molec. Genet. 12: 95-100, 1986.
[46425] 7116.

[46426] Axelrod, L.; Neer, R. M.; Kliman, B.: Hypogonadism in a male with
[46427] immunologically active, biologically inactive luteinizing hormone:
[46428] an exception to a venerable rule. J. Clin. Endocr. Metab. 48: 279-287,
[46429] 1979.

[46430] 7117.

[46431] Beitins, I. Z.; Axelrod, L.; Ostrea, T.; Little, R.; Badger, T.

[46432] M.: Hypogonadism in a male with an immunologically active, biologically
[46433] inactive luteinizing hormone: characterization of the abnormal hormone. J.
[46434] Clin. Endocr. Metab. 52: 1143-1149, 1981.

[46435] 7118.

[46436] Curtin, D.; Jenkins, S.; Farmer, N.; Anderson, A. C.; Haisenleder,

[46437] D. J.; Rissman, E.; Wilson, E. M.; Shupnik, M. A.: Androgen suppression

[46438] of GnRH-stimulated rat LH-beta gene transcription occurs through Sp1

[46439] sites in the distal GnRH-responsive promoter region. Molec. Endocr. 15:

[46440] 1906-1917, 2001.

[46441] 7119.

[46442] Furui, K.; Suganuma, N.; Tsukahara, S.-I.; Asada, Y.; Kikkawa,

[46443] F.; Tanaka, M.; Ozawa, T.; Tomoda, Y.: Identification of two point

[46444] mutations in the gene coding luteinizing hormone (LH) beta-subunit,

[46445] associated with immunologically anomalous LH variants. J. Clin. Endocr.

[46446] Metab. 78: 107-113, 1994.

[46447] 7120.

[46448] Hibberd, M. L.; Sumiya, M.; Summerfield, J. A.; Booy, R.; Levin,
[46449] M.; Meningococcal Research Group: Association of variants of the
[46450] gene for mannose-binding lectin with susceptibility to meningococcal
[46451] disease. Lancet 353: 1049-1053, 1999.

[46452] 7121.

[46453] Jack, D.; Bidwell, J.; Turner, M.; Wood, N.: Simultaneous genotyping
[46454] for all three known structural mutations in the human mannose-binding
[46455] lectin gene. Hum. Mutat. 9: 41-46, 1997.

[46456] 7122.

[46457] Lipscombe, R. J.; Sumiya, M.; Hill, A. V. S.; Lau, Y. L.; Levinsky,
[46458] R. J.; Summerfield, J. A.; Turner, M. W.: High frequencies in African
[46459] and non-African populations of independent mutations in the mannose
[46460] binding protein gene. Hum. Molec. Genet. 1: 709-715, 1992.

[46461] 7123.

[46462] Madsen, H. O.; Garred, P.; Kurtzhals, J. A. L.; Lamm, L. U.; Ryder,

[46463] L. P.; Thiel, S.; Svejgaard, A.: A new frequent allele is the missing

[46464] link in the structural polymorphism of the human man-nan-binding protein. Immunogenetics 40:

[46465] 37-44, 1994.

[46466] 7124.

[46467] Orren, A.; Potter, P. C.; Cooper, R. C.; du Toit, E.: Defi-ciency

[46468] of the sixth component of complement and susceptibility to Neisseria

[46469] meningitidis infections: studies in 10 families and five isolated

[46470] cases. Immunology 62: 249-253, 1987.

[46471] 7125.

[46472] Sastry, K.; Herman, G. A.; Day, L.; Deignan, E.; Bruns, G.; Morton,

[46473] C. C.; Ezekowitz, R. A. B.: The human mannose-binding protein gene:

[46474] exon structure reveals its evolutionary relationship to a human pulmonary
[46475] surfactant gene and localization to chromosome 10. J. Exp. Med. 170:
[46476] 1175-1189, 1989.

[46477] 7126.

[46478] Schuffenecker, I.; Narod, S. A.; Ezekowitz, R. A. B.; Sobol, H.;
[46479] Feunteun, J.; Lenoir, G. M.: The gene for mannose-binding protein
[46480] maps to chromosome 10 and is a marker for multiple endocrine neoplasia
[46481] type 2. Cytogenet. Cell Genet. 56: 99-102, 1991.

[46482] 7127.

[46483] Soothill, J. F.; Harvey, B. A.: Defective opsonization: a common
[46484] immunity deficiency. Arch. Dis. Child 51: 91-99, 1976.

[46485] 7128.

[46486] Sumiya, M.; Super, M.; Tabona, P.; Levinsky, R. J.; Arai, T.;

[46487] Turner, M. W.; Summerfield, J. A.: Molecular basis of opsonic defect

[46488] in immunodeficient children. Lancet 337: 1569-1570, 1991.

[46489] 7129.

[46490] Summerfield, J. A.; Ryder, S.; Sumiya, M.; Thursz, M.; Gorchein,

[46491] A.; Monteil, M. A.; Turner, M. W.: Mannose binding protein gene mutations

[46492] associated with unusual and severe infections in adults. Lancet 345:

[46493] 886-889, 1995.

[46494] 7130.

[46495] Summerfield, J. A.; Sumiya, M.; Levin, M.; Turner, M. W.: Association

[46496] of mutations in mannose binding protein gene with child-hood infection

[46497] in consecutive hospital series. Brit. Med. J. 314: 1229-1231, 1997.

[46498] 7131.

[46499] Super, M.; Gillies, S. D.; Foley, S.; Sastry, K.; Schweinle, J.-E.;

[46500] Silverman, V. J.; Ezekowitz, R. A. B.: Distinct and overlap-ping functions

[46501] of allelic forms of human mannose binding protein. Nature Genet. 2:
[46502] 50-55, 1992.

[46503] 7132.

[46504] Super, M.; Thiel, S.; Lu, J.; Levinsky, R. J.; Turner, M. W.:

[46505] Association of low levels of mannose-binding protein with a common
[46506] defect of opsonization. Lancet II: 1236-1239, 1989.

[46507] 7133.

[46508] Taylor, M. E.; Brickell, P. M.; Craig, R. K.; Summerfield, J.

[46509] A.: Structure and evolutionary origin of the gene encoding a human
[46510] serum mannose-binding protein. Biochem. J. 262:
763-771, 1989.
[46511] 7134.

[46512] Turner, M. W.: Deficiency of mannan binding protein--a new complement
[46513] deficiency syndrome. Clin. Exp. Immun. 86 (suppl. 1):
53-56, 1991.
[46514] 7135.

[46515] Turner, M. W.; Dinan, L.; Heatley, S.; Jack, D. L.; Boettcher,

[46516] B.; Lester, S.; McCluskey, J.; Roberton, D.: Restricted poly-
morphism

[46517] of the mannose-binding lectin gene of indigenous Aus-
tralians. Hum.

[46518] Molec. Genet. 9: 1481-1486, 2000.

[46519] 7136.

[46520] White, R. A.; Dowler, L. L.; Adkison, L. R.; Ezekowitz, R. A.

[46521] B.; Sastry, K. N.: The murine mannose-binding protein
genes (Mbl

[46522] 1 and Mbl 2) localize to chromosomes 14 and 19. Mam-
malian Genome 5:

[46523] 807-809, 1994.

[46524] 7137.

[46525] Chern, C. J.; Croce, C. M.: Confirmation of the synteny of
the

[46526] human genes for mannose phosphate isomerase and
pyruvate kinase and

[46527] of their assignment to chromosome 15. Cytogenet. Cell
Genet. 15:

[46528] 299-305, 1975.

[46529] 7138.

[46530] Brook, J. D.; Shaw, D. J.; Meredith, A. L.; Bruns, G. A. P.; Harper,

[46531] P. S.: Localisation of genetic markers and orientation of the linkage

[46532] group on chromosome 19. Hum. Genet. 68: 282-285, 1984.

[46533] 7139.

[46534] Bruns, G. A. P.; Regina, V. M.; Gerald, P. S.: Lysosomal DNase

[46535] and chromosome 19.(Abstract) J. Cell Biol. 83: 444a only, 1979.

[46536] 7140.

[46537] Kawane, K.; Fukuyama, H.; Kondoh, G.; Takeda, J.; Ohsawa, Y.; Uchiyama,

[46538] Y.; Nagata, S.: Requirement of DNase II for definitive erythropoiesis

[46539] in the mouse fetal liver. Science 292: 1546-1549, 2001.

[46540] 7141.

[46541] Yasuda, T.; Nadano, D.; Sawazaki, K.; Kishi, K.: Genetic polymorphism

[46542] of human deoxyribonuclease II (DNase II): low activity levels in urine

[46543] and leukocytes are due to an autosomal recessive allele. Ann. Hum.

[46544] Genet. 56: 1-10, 1992.

[46545] 7142.

[46546] Yasuda, T.; Takeshita, H.; Iida, R.; Nakajima, T.; Hosomi, O.;

[46547] Nakashima, Y.; Kishi, K.: Molecular cloning of the cDNA encoding

[46548] human deoxyribonuclease II. J. Biol. Chem. 273: 2610-2616, 1998.

[46549] 7143.

[46550] Yasuda, T.; Takeshita, H.; Iida, R.; Nakajima, T.; Hosomi, O.;

[46551] Nakashima, Y.; Mogi, K.; Kishi, K.: Chromosomal localization of a

[46552] human deoxyribonuclease II gene (DNASE2) to 19p13.2-p13.1 using both

[46553] the polymerase chain reaction and fluorescence in situ hybridization

[46554] analysis. Biochem. Biophys. Res. Commun. 244: 815-818,

1998.

[46555] 7144.

[46556] Baylin, S. B.: Tying it all together: epigenetics, genetics, cell

[46557] cycle, and cancer. Science 277: 1948-1949, 1997.

[46558] 7145.

[46559] Bestor, T.; Laudano, A.; Mattaliano, R.; Ingram, V.: Cloning and

[46560] sequencing of a cDNA encoding DNA methyltransferase of mouse cells:

[46561] the carboxyl-terminal domain of the mammalian enzymes is related to

[46562] bacterial restriction methyltransferases. J. Molec. Biol. 203: 971-983,

[46563] 1988.

[46564] 7146.

[46565] Bestor, T. H.: The DNA methyltransferases of mammals. Hum. Molec.

[46566] Genet. 9: 2395-2402, 2000.

[46567] 7147.

[46568] Bestor, T. H.: DNA methylation: evolution of a bacterial

immune

[46569] function into a regulator of gene expression and genome structure

[46570] in higher eukaryotes. Phil. Trans. Roy. Soc. London B 326: 179-187,

[46571] 1990.

[46572] 7148.

[46573] Chuang, L. S.-H.; Ian, H.-I.; Koh, T.-W.; Ng, H.-H.; Xu, G.; Li,

[46574] B. F. L.: Human DNA-(cytosine-5) methyltransferase-PCNA complex as

[46575] a target for p21(WAF1). Science 277: 1996-2000, 1997.

[46576] 7149.

[46577] El-Deiry, W. S.; Nelkin, B. D.; Celano, P.; Chiu Yen, R.-W.; Falco,

[46578] J. P.; Hamilton, S. R.; Baylin, S. B.: High expression of the DNA

[46579] methyltransferase gene characterizes human neoplastic cells and progression

[46580] stages of colon cancer. Proc. Nat. Acad. Sci. 88: 3470-3474, 1991.

[46581] 7150.

[46582] Fuks, F.; Burgers, W. A.; Brehm, A.; Hughes-Davies, L.; Kouzarides,
[46583] T.: DNA methyltransferase Dnmt1 associates with histone deacetylase
[46584] activity. Nature Genet. 24: 88-91, 2000.

[46585] 7151.

[46586] Howell, C. Y.; Bestor, T. H.; Ding, F.; Latham, K. E.; Mertineit,
[46587] C.; Trasler, J. M.; Chaillet, J. R.: Genomic imprinting dis-rupted
[46588] by a maternal effect mutation in the Dnmt1 gene. Cell 104: 829-838,
[46589] 2001.

[46590] 7152.

[46591] Hsu, D.-W.; Lin, M.-J.; Lee, T.-L.; Wen, S.-C.; Chen, X.; Shen,
[46592] C.-K. J.: Two major forms of DNA (cytosine-5) methyl-transferase in
[46593] human somatic tissues. Proc. Nat. Acad. Sci. 96: 9751-9756, 1999.

[46594] 7153.

[46595] Lee, P. P.; Fitzpatrick, D. R.; Beard, C.; Jessup, H. K.; Lehar,

[46596] S.; Makar, K. W.; Perez-Melgosa, M.; Sweetser, M. T.;
Schlissel, M.

[46597] S.; Nguyen, S.; Cherry, S. R.; Tsai, J. H.; Tucker, S. M.;
Weaver,

[46598] W. M.; Kelso, A.; Jaenisch, R.; Wilson, C. B.: A critical role
for

[46599] Dnmt1 and DNA methylation in T cell development, func-
tion, and survival. Immunity 15:

[46600] 763-774, 2001.

[46601] 7154.

[46602] Li, E.; Bestor, T. H.; Jaenisch, R.: Targeted mutation of the

[46603] DNA methyltransferase gene results in embryonic lethal-
ity. Cell 69:

[46604] 915-926, 1992.

[46605] 7155.

[46606] Lyko, F.; Ramsahoye, B. H.; Kashevsky, H.; Tudor, M.;
Mastrangelo,

[46607] M.-A.; Orr-Weaver, T. L.; Jaenisch, R.: Mammalian
(cytosine-5) methyltransferases

[46608] cause genomic DNA methylation and lethality in

Drosophila. Nature

[46609] Genet. 23: 363-366, 1999.

[46610] 7156.

[46611] Rhee, I.; Bachman, K. E.; Park, B. H.; Jair, K.-W.; Yen, R.-W.

[46612] C.; Schuebel, K. E.; Cui, H.; Feinberg, A. P.; Lengauer, C.; Kinzler,

[46613] K. W.; Baylin, S. B.; Vogelstein, B.: DNMT1 and DNMT3b cooperate

[46614] to silence genes in human cancer cells. Nature 416: 552-556, 2002.

[46615] 7157.

[46616] Rhee, I.; Jair, K.-W.; Yen, R-W. C.; Lengauer, C.; Herman, J.

[46617] G.; Kinzler, K. W.; Vogelstein, B.; Baylin, S. B.; Schuebel, K. E.

[46618] : CpG methylation is maintained in human cancer cells lacking DNMT1. Nature 404:

[46619] 1003-1007, 2000.

[46620] 7158.

[46621] Robertson, K. D.; Ait-Si-Ali, S.; Yokochi, T.; Wade, P. A.; Jones,

[46622] P. L.; Wolffe, A. P.: DNMT1 forms a complex with Rb, E2F1

and HDAC1
[46623] and represses transcription from E2F-responsive promoters. Nature
[46624] Genet. 25: 338-342, 2000.

[46625] 7159.

[46626] Rouleau, J.; Tanigawa, G.; Szyf, M.: The mouse DNA methyltransferase
[46627] 5-prime region: a unique housekeeping gene promoter. J. Biol. Chem. 267:
[46628] 7368-7377, 1992.

[46629] 7160.

[46630] Rountree, M. R.; Bachman, K. E.; Baylin, S. B.: DNMT1 binds HDAC2
[46631] and a new co-repressor, DMAP1, to form a complex at replication foci. Nature
[46632] Genet. 25: 269-277, 2000.

[46633] 7161.

[46634] Tucker, K. L.; Talbot, D.; Lee, M. A.; Leonhardt, H.; Jaenisch,
[46635] R.: Complementation of methylation deficiency in embryonic stem cells

[46636] by a DNA methyltransferase minigene. Proc. Nat. Acad. Sci. 93: 12920-12925,

[46637] 1996.

[46638] 7162.

[46639] Yen, R.-W. C.; Vertino, P. M.; Nelkin, B. D.; Yu, J. J.; El-Deiry,

[46640] W.; Cumaraswamy, A.; Lennon, G. G.; Trask, B. J.; Celano, P.; Baylin,

[46641] S. B.: Isolation and characterization of the cDNA encoding human

[46642] DNA methyltransferase. Nucleic Acids Res. 20: 2287-2291, 1992.

[46643] 7163.

[46644] Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification

[46645] of a novel FAS/ER-alpha fusion transcript expressed in human cancer

[46646] cells. Biochim. Biophys. Acta 1493: 373-377, 2000.

[46647] 7164.

[46648] Zuppan, P.; Hall, J. M.; Lee, M. K.; Ponglikitmongkol, M.; King,

[46649] M.-C.: Possible linkage of the estrogen receptor gene to

breast cancer
[46650] in a family with late-onset disease. Am. J. Hum. Genet. 48: 1065-1068,
[46651] 1991.

[46652] 7165.

[46653] Zuppan, P. J.; Hall, J. M.; Ponglikitmongkol, M.; Spielman, R.;
[46654] King, M. C.: Polymorphisms at the estrogen receptor (ESR) locus and
[46655] linkage relationships on chromosome 6q. (Abstract) Cyto-genet. Cell
[46656] Genet. 51: 1116 only, 1989.

[46657] 7166.

[46658] Calabi, F.; Cilli, V.: CBFA2T1, a gene rearranged in human leukemia,
[46659] is a member of a multigene family. Genomics 52: 332-341, 1998.
[46660] 7167.

[46661] Erickson, P.; Gao, J.; Chang, K.-S.; Look, T.; Whisenant, E.; Raimondo,
[46662] S.; Lasher, R.; Trujillo, J.; Rowley, J. D.; Drabkin, H. A.:

Identification

[46663] of breakpoints in t(8;21) acute myelogenous leukemia and isolation

[46664] of a fusion transcript, AML1/ETO, with similarity to Drosophila segmentation

[46665] gene, runt. Blood 80: 1825-1831, 1992.

[46666] 7168.

[46667] Miyamoto, T.; Weissman, I. L.; Akashi, K.: AML1/ETO-expressing

[46668] nonleukemic stem cells in acute myelogenous leukemia with 8;21 chromosomal

[46669] translocation. Proc. Nat. Acad. Sci. 97: 7521-7526, 2000.

[46670] 7169.

[46671] Miyoshi, H.; Kozu, T.; Shimizu, K.; Enomoto, K.; Maseki, N.; Kaneko,

[46672] Y.; Kamada, N.; Ohki, M.: The t(8;21) translocation in acute myeloid

[46673] leukemia results in production of an AML1-MTG8 fusion transcript. EMBO

[46674] J. 12: 2715-2721, 1993.

[46675] 7170.

[46676] Miyoshi, H.; Shimizu, K.; Kozu, T.; Maseki, N.; Kaneko, Y.; Ohki,

[46677] M.: t(8;21) breakpoints on chromosome 21 in acute myeloid leukemia

[46678] are clustered within a limited region of a single gene, AML1. Proc.

[46679] Nat. Acad. Sci. 88: 10431-10434, 1991.

[46680] 7171.

[46681] Linggi, B.; Muller-Tidow, C.; van de Locht, L.; Hu, M.; Nip, J.;

[46682] Serve, H.; Berdel, W. E.; van der Reijden, B.; Quelle, D. E.; Rowley,

[46683] J. D.; Cleveland, J.; Jansen, J. H.; Pandolfi, P. P.; Hiebert, S.

[46684] W.: The t(8;21) fusion protein, AML1-ETO, specifically represses

[46685] the transcription of the p14(ARF) tumor suppressor in acute myeloid

[46686] leukemia. Nature Med. 8: 743-750, 2002.

[46687] 7172.

[46688] Minucci, S.; Maccarana, M.; Cioce, M.; De Luca, P.; Gelmetti, V.;

[46689] Segalla, S.; Di Croce, L.; Giavara, S.; Matteucci, C.; Gobbi, A.;

[46690] Bianchini, A.; Colombo, E.; Schiavoni, I.; Badaracco, G.; Hu, X.;

[46691] Lazar, M. A.; Landsberger, N.; Nervi, C.; Pelicci, P. G.: Oligomerization

[46692] of RAR and AML1 transcription factors as a novel mechanism of oncogenic

[46693] activation. Molec. Cell 5: 811-820, 2000.

[46694] 7173.

[46695] Niwa-Kawakita, M.; Miyoshi, H.; Gotoh, O.; Matsushima, Y.; Nishimura,

[46696] M.; Shisa, H.; Ohki, M.: Cloning and gene mapping of the mouse homologue

[46697] of the CBFA2T1 gene associated with human acute myeloid leukemia. Genomics 29:

[46698] 755-759, 1995.

[46699] 7174.

[46700] Nucifora, G.; Rowley, J. D.: The AML1 and ETO genes in acute myeloid

[46701] leukemia with a t(8;21). Leukemia Lymphoma 14: 353-362, 1994.

[46702] 7175.

[46703] Schoch, C.; Kohlmann, A.; Schnittger, S.; Brors, B.; Dugas, M.;

[46704] Mergenthaler, S.; Kern, W.; Hiddemann, S.; Eils, R.; Haferlach, T.

[46705] : Acute myeloid leukemias with reciprocal rearrangements can be distinguished

[46706] by specific gene expression profiles. Proc. Nat. Acad. Sci. 99:

[46707] 10008-10013, 2002.

[46708] 7176.

[46709] Wolford, J. K.; Bogardus, C.; Prochazka, M.: Polymorphism in

[46710] the 3-prime untranslated region of MTG8 is associated with obesity

[46711] in Pima Indian males. Biochem. Biophys. Res. Commun. 246: 624-626,

[46712] 1998.

[46713] 7177.

[46714] Wolford, J. K.; Prochazka, M.: Structure and expression of the

[46715] human MTG8/ETO gene. Gene 212: 103-109, 1998.

[46716] 7178.

[46717] Yergeau, D. A.; Hetherington, C. J.; Wang, Q.; Zhang, P.; Sharpe,

[46718] A. H.; Binder, M.; Marin-Padilla, M.; Tenen, D. G.; Speck, N. A.;

[46719] Zhang, D.-E.: Embryonic lethality and impairment of haematopoiesis

[46720] in mice heterozygous for an AML1-ETO fusion gene. Na-ture Genet. 15:

[46721] 303-306, 1997.

[46722] 7179.

[46723] Pause, A.; Belsham, G. J.; Gingras, A.-C.; Donze, O.; Lin, T.-A.;

[46724] Lawrence, J. C., Jr.; Sonenberg, N.: Insulin-dependent stimulation

[46725] of protein synthesis by phosphorylation of a regulator of 5-prime-cap

[46726] function. Nature 371: 762-767, 1994.

[46727] 7180.

[46728] Pyronnet, S.; Imataka, H.; Gingras, A.-C.; Fukunaga, R.;

Hunter,
[46729] T.; Sonenberg, N.: Human eukaryotic translation initiation factor
[46730] 4G (eIF4G) recruits Mnk1 to phosphorylate eIF4E. EMBO J. 18: 270-279,
[46731] 1999.

[46732] 7181.

[46733] Waskiewicz, A. J.; Flynn, A.; Proud, C. G.; Cooper, J. A.: Mitogen-activated
[46734] protein kinases activate the serine/threonine kinases Mnk1 and Mnk2. EMBO
[46735] J. 16: 1909-1920, 1997.

[46736] 7182.

[46737] Zhang, Y.; Saison, M.; Spaepen, M.; De Strooper, B.; Van Leuven,
[46738] F.; David, G.; Van den Berghe, H.; Cassiman, J.-J.: Mapping of human
[46739] fibronectin receptor beta subunit gene to chromosome 10. Somat. Cell
[46740] Molec. Genet. 14: 99-104, 1988.

[46741] 7183.

[46742] Kondoh, S.; Sugawara, H.; Harada, N.; Matsumoto, N.; Ohashi, H.;

[46743] Sato, M.; Kantaputra, P. N.; Ogino, T.; Tomita, H.; Ohta, T.; Kishino,

[46744] T.; Fukushima, Y.; Niikawa, N.; Yoshiura, K.: A novel gene is disrupted

[46745] at a 14q13 breakpoint of t(2;14) in a patient with mirror-image polydactyly

[46746] of hands and feet. J. Hum. Genet. 47: 136-139, 2002.

[46747] 7184.

[46748] Argraves, W. S.; Dickerson, K.; Burgess, W. H.; Ruoslahti, E.:

[46749] Fibulin, a novel protein that interacts with the fibronectin receptor-beta

[46750] subunit cytoplasmic domain. Cell 58: 623-629, 1989.

[46751] 7185.

[46752] Korenberg, J. R.; Chen, X.-N.; Tran, H.; Argraves, W. S.: Localization

[46753] of the human gene for fibulin-1 (FBLN1) to chromosome band 22q13.3.

[46754] Cytogenet. Cell Genet. 68: 192-193, 1995.

[46755] 7186.

[46756] Mattei, M.-G.; Pan, T.-C.; Zhang, R.-Z.; Timpl, R.; Chu, M.-L.

[46757] : The fibulin-1 gene (FBLN1) is located on human chromosome 22 and

[46758] on mouse chromosome 15. Genomics 22: 437-438, 1994.

[46759] 7187.

[46760] Karl, M.; Lamberts, S. W.; Koper, J. W.; Katz, D. A.; Huizenga,

[46761] N. E.; Kino, T.; Haddad, B. R.; Hughes, M. R.; Chrousos, G. P.: Cushing's

[46762] disease preceded by generalized glucocorticoid resistance: clinical

[46763] consequences of a novel, dominant-negative glucocorticoid receptor

[46764] mutation. Proc. Assoc. Am. Phys. 108: 296-307, 1996.

[46765] 7188.

[46766] Pavlakis, G. N.; Hizuka, N.; Gorden, P.; Seeburg, P. H.; Hamber,

[46767] D. H.: Expression of two human growth hormone genes in monkey cells

[46768] infected by simian virus 40 recombinants. Proc. Nat. Acad. Sci. 78:

[46769] 7398-7402, 1981.

[46770] 7189.

[46771] Geijsen, N.; Uings, I. J.; Pals, C.; Armstrong, J.; McKinnon, M.;

[46772] Raaijmakers, J. A. M.; Lammers, J.-W. J.; Koenderman, L.; Coffer,

[46773] P. J.: Cytokine-specific transcriptional regulation through an IL-5R-alpha

[46774] interacting protein. Science 293: 1136-1138, 2001.

[46775] 7190.

[46776] Gough, N. M.; Rakar, S.: Localization of the IL-5 receptor gene

[46777] to the distal half of murine chromosome 6 using recombinant inbred

[46778] strains of mice. Genomics 12: 855-856, 1992.

[46779] 7191.

[46780] Isobe, M.; Kumura, Y.; Murata, Y.; Takaki, S.; Tominaga, A.; Takatsu,

[46781] K.; Ogita, Z.: Localization of the gene encoding the alpha

subunit
[46782] of human interleukin-5 receptor (IL5RA) to chromosome region 3p24-3p26. Genomics 14:
[46783] 755-758, 1992.

[46784] 7192.

[46785] Jacob, C. O.; Mykytyn, K.; Varcony, T.; Drabkin, H. A.: Mapping
[46786] of the interleukin 5 receptor gene to human chromosome 3p25-p26 and
[46787] to mouse chromosome 6 close to the Raf-1 locus with polymorphic tandem
[46788] repeat sequences. Mammalian Genome 4: 435-439, 1993.

[46789] 7193.

[46790] Sun, Z.; Yergeau, D. A.; Tuypens, T.; Tavernier, J.; Paul, C. C.;
[46791] Baumann, M. A.: Tenen, D. G.; Ackerman, S. J.: Identification and
[46792] characterization of a functional promoter region in the human eosinophil
[46793] IL-5 receptor alpha subunit gene. J. Biol. Chem. 270: 1462-1471,
[46794] 1995.

[46795] 7194.

[46796] Tavernier, J.; Tuypens, T.; Plaetinck, G.; Verhee, A.; Fiers, W.;

[46797] Devos, R.: Molecular basis of the membrane-anchored and two soluble

[46798] isoforms of the human interleukin 5 receptor alpha subunit. Proc.

[46799] Nat. Acad. Sci. 89: 7041-7045, 1992.

[46800] 7195.

[46801] Zhang, J.; Kuvelkar, R.; Cheewatrakoolpong, B.; Williams, S.; Egan,

[46802] R. W.; Billah, M. M.: Evidence for multiple promoters of the human

[46803] IL-5 receptor alpha subunit gene: a novel 6-base pair element determines

[46804] cell-specific promoter function. J. Immun. 15: 5412-5421, 1997.

[46805] 7196.

[46806] Povey, S.; Morton, N. E.; Sherman, S. L.: Report of the committee

[46807] on the genetic constitution of chromosomes 1 and 2. Cytogenet. Cell

[46808] Genet. 40: 67-106, 1985.

[46809] 7197.

[46810] Bird, A.; Taggart, M.; Frommer, M.; Miller, O. J.; Macleod, D.

[46811] : A fraction of the mouse genome that is derived from islands of nonmethylated,

[46812] CpG-rich DNA. Cell 40: 91-99, 1985.

[46813] 7198.

[46814] Pikkarainen, T.; Kallunki, T.; Tryggvason, K.: Human laminin B2

[46815] chain: comparison of the complete amino acid sequence with the B1

[46816] chain reveals variability in sequence homology between different structural

[46817] domains. J. Biol. Chem. 263: 6751-6758, 1988.

[46818] 7199.

[46819] Sasaki, M.; Yamada, Y.: The laminin B2 chain has a multidomain

[46820] structure homologous to the B1 chain. J. Biol. Chem. 262: 17111-17117,

[46821] 1987.

[46822] 7200.

[46823] Smyth, N.; Vatansever, H. S.; Murray, P.; Meyer, M.; Frie, C.;

[46824] Paulsson, M.; Edgar, D.: Absence of basement membranes after targeting

[46825] the LAMC1 gene results in embryonic lethality due to failure of endoderm

[46826] differentiation. J. Cell Biol. 144: 151-160, 1999.

[46827] 7201.

[46828] Aberdam, D.; Galliano, M.-F.; Vailly, J.; Pulkkinen, L.; Bonifas,

[46829] J.; Christiano, A. M.; Tryggvason, K.; Uitto, J.; Epstein, E. H.,

[46830] Jr.; Ortonne, J.-P.; Meneguzzi, G.: Herlitz's junctional epidermolysis

[46831] bullosa is linked to mutations in the gene (LAMC2) for the gamma-2

[46832] subunit of nicein/kalinin (laminin-5). Nature Genet. 6: 299-304,

[46833] 1994.

[46834] 7202.

[46835] Aberdam, D.; Galliano, M. F.; Mattei, M.-G.; Pisani-Spadafora,

[46836] A.; Ortonne, J. P.; Meneguzzi, G.: Assignment of mouse nicein genes

[46837] to chromosomes 1 and 18. Mammalian Genome 5: 229-233, 1994.

[46838] 7203.

[46839] Airenne, T.; Haakana, H.; Sainio, K.; Kallunki, T.; Kallunki, P.;

[46840] Sariola, H.; Tryggvason, K.: Structure of the human laminin gamma-2

[46841] chain gene (LAMC2): alternative splicing with different tissue distribution

[46842] of two transcripts. Genomics 32: 54-64, 1996.

[46843] 7204.

[46844] Baudoin, C.; Miquel, C.; Gagnoux-Palacios, L.; Pulkkinen, L.; Christiano,

[46845] A. M.; Uitto, J.; Tadini, G.; Ortonne, J.-P.; Meneguzzi, G.: A novel

[46846] homozygous nonsense mutation in the LAMC2 gene in patients with the

[46847] Herlitz junctional epidermolysis bullosa. Hum. Molec.

Genet. 3:
[46848] 1909-1910, 1994.

[46849] 7205.

[46850] Kallunki, P.; Sainio, K.; Eddy, R.; Byers, M.; Kallunki, T.; Sariola,
[46851] H.; Beck, K.; Hirvonen, H.; Shows, T. B.; Tryggvason, K.: A truncated
[46852] laminin chain homologous to the B2 chain: structure, spatial expression,
[46853] and chromosomal assignment. J. Cell Biol. 119: 679-693, 1992.

[46854] 7206.

[46855] Nakano, A.; Chao, S.-C.; Pulkkinen, L.; Murrell, D.; Bruckner-Tuderman,
[46856] L.; Pfendner, E.; Uitto, J.: Laminin 5 mutations in junctional epidermolysis
[46857] bullosa: molecular basis of Herlitz vs non-Herlitz phenotypes. Hum.
[46858] Genet. 110: 41-51, 2002.

[46859] 7207.

[46860] Pulkkinen, L.; Christiano, A. M.; Airenne, T.; Haakana, H.;

Tryggvason,
[46861] K.; Uitto, J.: Mutations in the gamma-2 chain gene (LAMC2) of kalinin/laminin
[46862] 5 in the junctional forms of epidermolysis bullosa. Nature Genet. 6:
[46863] 293-298, 1994.

[46864] 7208.

[46865] Takizawa, Y.; Pulkkinen, L.; Chao, S.-C.; Nakajima, H.; Nakano,
[46866] Y.; Shimizu, H.; Uitto, J.: Complete paternal uniparental isodisomy
[46867] of chromosome 1: a novel mechanism for Herlitz junctional epidermolysis
[46868] bullosa. J. Invest. Derm. 206: 307-311, 2000.

[46869] 7209.

[46870] Ashton, G. H. S.; Mellerio, J. E.; Dunnill, M. G. S.; Pulkkinen,
[46871] L.; Christiano, A. M.; Uitto, J.; Eady, R. A. J.; McGrath, J. A.:

[46872] A recurrent laminin 5 mutation in British patients with lethal (Herlitz)
[46873] junctional epidermolysis bullosa: evidence for a muta-

tional hotspot
[46874] rather than propagation of an ancestral allele. Brit. J. Derm. 136:
[46875] 674-677, 1997.

[46876] 7210.

[46877] Benlian, P.; Foubert, L.; Gagne, E.; Bernard, L.; De Gennes, J.
[46878] L.; Langlois, S.; Robinson, W.; Hayden, M.: Complete paternal isodisomy
[46879] for chromosome 8 unmasked by lipoprotein lipase deficiency. Am. J.
[46880] Hum. Genet. 59: 431-436, 1996.

[46881] 7211.

[46882] Engel, E.: Uniparental disomy revisited: the first twelve years. Am.
[46883] J. Med. Genet. 46: 670-674, 1993.

[46884] 7212.

[46885] Floeth, M.; Bruckner-Tuderman, L.: Digenic junctional epidermolysis
[46886] bullosa: mutations in COL17A1 and LAMB3 genes. Am. J. Hum. Genet. 65:

[46887] 1530-1537, 1999.

[46888] 7213.

[46889] Gache, Y.; Allegra, M.; Bodemer, C.; Pisani-Spadafora, A.; de Prost,

[46890] Y.; Ortonne, J. P.; Meneguzzi, G.: Genetic bases of severe junctional
[46891] epidermolysis bullosa presenting spontaneous amelioration with aging. Hum.
[46892] Molec. Genet. 10: 2453-2461, 2001.

[46893] 7214.

[46894] Jonkman, M. F.; Scheffer, H.; Stulp, R.; Pas, H. H.; Nijenhuis,
[46895] M.; Heeres, K.; Owaribe, K.; Pulkkinen, L.; Uitto, J.: Revertant
[46896] mosaicism in epidermolysis bullosa caused by mitotic gene conversion. Cell 88:
[46897] 543-551, 1997.

[46898] 7215.

[46899] Kivirikko, S.; McGrath, J. A.; Pulkkinen, L.; Uitto, J.; Christiano,
[46900] A. M.: Mutational hotspots in the LAMB3 gene in the lethal (Herlitz)
[46901] type of junctional epidermolysis bullosa. Hum. Molec. Genet. 5:
[46902] 231-237, 1996.

[46903] 7216.

[46904] Ledbetter, D. H.; Engel, E.: Uniparental disomy in humans: development

[46905] of an imprinting map and its implications for prenatal diagnosis. Hum.

[46906] Molec. Genet. 4: 1757-1764, 1995.

[46907] 7217.

[46908] McGrath, J. A.; Pulkkinen, L.; Christiano, A. M.; Leigh, I. M.;

[46909] Eady, R. A. J.; Uitto, J.: Altered laminin 5 expression due to mutations

[46910] in the gene encoding the beta-3 chain (LAMB3) in generalized atrophic

[46911] benign epidermolysis bullosa. J. Invest. Derm. 104: 467-474, 1995.

[46912] 7218.

[46913] Mellerio, J. E.; Eady, R. A. J.; Atherton, D. J.; Lake, B. D.;

[46914] McGrath, J. A.: E210K mutation in the gene encoding the beta-3 chain

[46915] of laminin-5 (LAMB3) is predictive of a phenotype of generalized atrophic

[46916] benign epidermolysis bullosa. Brit. J. Derm. 139:

325-331, 1998.

[46917] 7219.

[46918] Schoenmakers, E. P. P. M.; Wanschura, S.; Mols, R.;
Bullerdiek,

[46919] J.; Van den Berghe, H.; Van de Ven, W. J. M.: Recurrent re-
arrangements

[46920] in the high mobility group protein gene, HMGI-C, in be-
nign mesenchymal

[46921] tumours. Nature Genet. 10: 436-444, 1995.

[46922] 7220.

[46923] Turc-Carel, C.; Pietrzak, E.; Kakati, S.; Kinniburgh, A. J.;
Sandberg,

[46924] A. A.: The human int-1 gene is located at chromosome
region 12q12-12q13

[46925] and is not rearranged in myxoid liposarcoma with
t(12;16)(q13;p11). Oncogene

[46926] Res. 1: 397-405, 1987.

[46927] 7221.

[46928] Bochkov, V. N.; Kadl, A.; Huber, J.; Gruber, F.; Binder, B. R.;

[46929] Leitinger, N.: Protective role of phospholipid oxidation
products

[46930] in endotoxin-induced tissue damage. Nature 419: 77-81, 2002.

[46931] 7222.

[46932] Hassett, C.; Robinson, K. B.; Beck, N. B.; Omiecinski, C. J.:

[46933] The human microsomal epoxide hydrolase gene (EPHX1): complete nucleotide

[46934] sequence and structural characterization. Genomics 23: 433-442,

[46935] 1994.

[46936] 7223.

[46937] Jackson, M. R.; Craft, J. A.; Burchell, B.: Nucleotide and deduced

[46938] amino acid sequence of human liver microsomal epoxide hydrolase. Nucleic

[46939] Acids Res. 15: 7188 only, 1987.

[46940] 7224.

[46941] Kutt, H.; Brennan, R.; Dehejia, H.; Verebely, K.: Diphenylhydantoin

[46942] intoxication: a complication of isoniazid therapy. Am. Rev. Resp.

[46943] Dis. 101: 377-383, 1970.

[46944] 7225.

[46945] Kutt, H.; Verebely, K.; McDowell, F.: Inhibition of diphenyl-
hydantoin
[46946] metabolism in rat and in rat liver microsome by antituber-
cular drugs. Neurology 18:
[46947] 706-710, 1968.

[46948] 7226.

[46949] Kutt, H.; Wolk, M.; Scherman, R.; McDowell, F.: Insufficient

[46950] parahydroxylation as a cause of diphenylhydantoin toxic-
ity. Neurology 14:
[46951] 542-548, 1964.

[46952] 7227.

[46953] McGlynn, K. A.; Rosvold, E. A.; Lustbader, E. D.; Hu, Y.;
Clapper,
[46954] M. L.; Zhou, T.; Wild, C. P.; Xia, X.-L.; Baffoe-Bonnie, A.;
Ofori-Adjei,
[46955] D.; Chen, G.-C.; London, W. T.; Shen, F.-M.; Buetow, K. H.:
Susceptibility
[46956] to hepatocellular carcinoma is associated with genetic
variation in
[46957] the enzymatic detoxification of aflatoxin B1. Proc. Nat.

Acad. Sci. 92:
[46958] 2384-2387, 1995.

[46959] 7228.

[46960] Nadeau, J. H.: Personal Communication. Bar Harbor, Me. 6/22/1988.

[46961] 7229.

[46962] Phelan, M. C.; Pellock, J. M.; Nance, W. E.: Discordant ex-
pression
[46963] of fetal hydantoin syndrome in a pair of dizygotic twins with different
[46964] fathers. (Abstract) Am. J. Hum. Genet. 33: 67A only, 1981.

[46965] 7230.

[46966] Sabry, M. A.; Farag, T. I.: Hand anomalies in fetal-
hydantoin
[46967] syndrome: from nail/phalangeal hypoplasia to unilateral acheiria.
[46968] (Letter) Am. J. Med. Genet. 62: 410-412, 1996.

[46969] 7231.

[46970] Skoda, R. C.; Demierre, A.; McBride, O. W.; Gonzalez, F. J.; Meyer,
[46971] U. A.: Human microsomal xenobiotic epoxide hydrolase:

complementary

[46972] DNA sequence, complementary DNA-directed expression in COS-1 cells,

[46973] and chromosomal localization. J. Biol. Chem. 263: 1549-1554, 1988.

[46974] 7232.

[46975] Spielberg, S. P.; Gordon, G. B.; Blake, D. A.; Goldstein, D. A.;

[46976] Herlong, H. F.: Predisposition to phenytoin hepatotoxicity assessed

[46977] in vitro. New Eng. J. Med. 305: 722-727, 1981.

[46978] 7233.

[46979] Strickler, S. M.; Dansky, L. V.; Miller, M. A.; Seni, M. H.; Andermann,

[46980] E.; Spielberg, S. P.: Genetic predisposition to phenytoin-induced

[46981] birth defects. Lancet II: 746-749, 1985.

[46982] 7234.

[46983] Vasko, M. R.; Bell, R. D.; Daly, D. D.: Inheritance of diphenylhydantoin

[46984] hypometabolism: a pharmacokinetic study of one family.

(Abstract) Clin.

[46985] Pharm. Therap. 21: 120 only, 1977.

[46986] 7235.

[46987] Vasko, M. R.; Bell, R. D.; Daly, D. D.; Pippenger, C. E.: In-
heritance

[46988] of phenyltoin hypometabolism: a kinetic study of one
family. Clin.

[46989] Pharm. Therap. 27: 96-103, 1980.

[46990] 7236.

[46991] Vermeij, P.; Ferrari, M. D.; Buruma, O. J. S.; Veenema, H.;
de

[46992] Wolff, F. A.: Inheritance of poor phenytoin parahydroxyla-
tion capacity

[46993] in a Dutch family. Clin. Pharm. Therap. 44: 588-593,
1988.

[46994] 7237.

[46995] Vesell, E. S.: Pharmacogenetics: multiple interactions be-
tween

[46996] genes and environment as determinants of drug response.
(Editorial) Am.

[46997] J. Med. 66: 183-187, 1979.

[46998] 7238.

[46999] Zusterzeel, P. L. M.; Peters, W. H. M.; Visser, W.; Hermsen, K.

[47000] J. M.; Roelofs, H. M. J.; Steegers, E. A. P.: A polymorphism in the

[47001] gene for microsomal epoxide hydrolase is associated with pre-eclampsia. J.

[47002] Med. Genet. 38: 234-237, 2001.

[47003] 7239.

[47004] Fonatsch, C.; Latza, U.; Durkop, H.; Rieder, H.; Stein, H.: Assignment

[47005] of the human CD30 (Ki-1) gene to 1p36. Genomics 14: 825-826, 1992.

[47006] 7240.

[47007] Ma, T.; Yang, B.; Umenishi, F.; Verkman, A. S.: Closely spaced

[47008] tandem arrangement of AQP2, AQP5, and AQP6 genes in a 27-kilobase

[47009] segment at chromosome locus 12q13. Genomics 43: 387-389, 1997.

[47010] 7241.

[47011] Van Heyningen, V.; Bobrow, M.; Bodmer, W. F.; Gardiner, S. E.;

[47012] Povey, S.; Hopkinson, D. A.: Chromosome assignment of some human

[47013] enzyme loci: mitochondrial malate dehydrogenase to 7, mannosephosphate

[47014] isomerase and pyruvate kinase to 15 and probably, esterase D to 13. Ann.

[47015] Hum. Genet. 38: 295-303, 1975.

[47016] 7242.

[47017] Larson, L. M.; Bruce, A. W.; Saumur, J. H.; Wasdahl, W. A.: Further

[47018] evidence by gene dosage for the regional assignment of erythrocyte

[47019] acid phosphatase (ACP1) and malate dehydrogenase (MDH1) loci on chromosome

[47020] 2p. Clin. Genet. 22: 220-225, 1982.

[47021] 7243.

[47022] Bell, G. I.; Gerhard, D. S.; Fong, N. M.; Sanchez-Pescador, R.;

[47023] Rall, L. B.: Isolation of the human insulin-like growth factor genes:

[47024] insulin-like growth factor II and insulin genes are contiguous. Proc.

[47025] Nat. Acad. Sci. 82: 6450-6454, 1985.

[47026] 7244.

[47027] Bell, G. I.; Merryweather, J. P.; Sanchez-Pescador, R.; Stempien,

[47028] M. M.; Priestley, L.; Scott, J.; Rall, L. B.: Sequence of a cDNA

[47029] clone encoding human preproinsulin-like growth factor II. Nature 310:

[47030] 775-777, 1984.

[47031] 7245.

[47032] Brissenden, J. E.; Ullrich, A.; Francke, U.: Chromosomal mapping

[47033] of loci for insulin-like growth factors I and II and for epidermal

[47034] growth factor in man. (Abstract) Am. J. Hum. Genet. 36: 133S, 1984.

[47035] 7246.

[47036] Brown, K. W.; Villar, A. J.; Bickmore, W.; Clayton-Smith, J.; Catchpoole,

[47037] D.; Maher, E. R.; Reik, W.: Imprinting mutation in the Beckwith-Wiedemann
[47038] syndrome leads to biallelic IGF2 expression through an H19-independent
[47039] pathway. Hum. Molec. Genet. 5: 2027-2032, 1996.

[47040] 7247.

[47041] Burgisser, D. M.; Roth, B. V.; Giger, R.; Luthi, C.; Weigl, S.;

[47042] Zarn, J.; Humbel, R. E.: Mutants of human insulin-like growth factor
[47043] II with altered affinities for the type 1 and type 2 insulin-like
[47044] growth factor receptor. J. Biol. Chem. 266: 1029-1033, 1991.
[47045] 7248.

[47046] Comings, D. E.: A general theory of carcinogenesis. Proc. Nat.
[47047] Acad. Sci. 70: 3324-3328, 1973.

[47048] 7249.

[47049] Constancia, M.; Dean, W.; Lopes, S.; Moore, T.; Kelsey, G.; Reik,
[47050] W.: Deletion of a silencer element in Igf2 results in loss of

imprinting

[47051] independent of H19. Nature Genet. 26: 203-206, 2000.

[47052] 7250.

[47053] Constancia, M.; Hemberger, M.; Hughes, J.; Dean, W.; Ferguson-Smith,

[47054] A.; Fundele, R.; Stewart, F.; Kelsey, G.; Fowden, A.; Sibley, C.;

[47055] Reik, W.: Placental-specific IGF-II is a major modulator of placental

[47056] and fetal growth. Nature 417: 945-948, 2002.

[47057] 7251.

[47058] Daughaday, W. H.; Emanuele, M. A.; Brooks, M. H.; Barbato, A.

[47059] L.; Kapadia, M.; Rotwein, P.: Synthesis and secretion of insulin-like

[47060] growth factor II by a leiomyosarcoma with associated hypoglycemia. New

[47061] Eng. J. Med. 319: 1434-1440, 1988.

[47062] 7252.

[47063] de Pagter-Holthuizen, P.; Hoppener, J. W. M.; Jansen, M.; Geurts

[47064] van Kessel, A. H. M.; van Ommen, G. J. B.; Sussenbach, J. S.: Chromosomal

[47065] localization and preliminary characterization of the human gene encoding

[47066] insulin-like growth factor II. Hum. Genet. 69: 170-173, 1985.

[47067] 7253.

[47068] Drummond, I. A.; Madden, S. L.; Rohwer-Nutter, P.; Bell, G. I.;

[47069] Sukhatme, V. P.; Rauscher, F. J., III: Repression of the insulin-like

[47070] growth factor II gene by the Wilms tumor suppressor WT1. Science 257:

[47071] 674-678, 1992.

[47072] 7254.

[47073] Dull, T. J.; Gray, A.; Hayflick, J. S.; Ullrich, A.: Insulin-like

[47074] growth factor II precursor gene organization in relation to insulin

[47075] gene family. Nature 310: 777-781, 1984.

[47076] 7255.

[47077] Gaunt, T. R.; Cooper, J. A.; Miller, G. J.; Day, I. N. M.;

O'Dell,

[47078] S. D.: Positive associations between single nucleotide polymorphisms

[47079] in the IGF2 gene region and body mass index in adult males. Hum.

[47080] Molec. Genet. 10: 1491-1501, 2001.

[47081] 7256.

[47082] Haselbacher, G. K.; Irminger, J.-C.; Zapf, J.; Ziegler, W. H.;

[47083] Humbel, R. E.: Insulin-like growth factor II in human adrenal pheochromocytoma

[47084] and Wilms tumors: expression at the mRNA and protein level. Proc.

[47085] Nat. Acad. Sci. 84: 1104-1106, 1987.

[47086] 7257.

[47087] Jansen, M.; van Schaik, F. M. A.; van Tol, H.; Van den Brande,

[47088] J. L.; Sussenbach, J. S.: Nucleotide sequences of cDNAs encoding

[47089] precursors of human insulin-like growth factor II (IGF-II) and an

[47090] IGF-II variant. FEBS Lett. 179: 243-246, 1985.

[47091] 7258.

[47092] Jeon, J.-T.; Carlborg, O.; Tornsten, A.; Giuffra, E.; Amarger,
[47093] V.; Chardon, P.; Andersson-Eklund, L.; Andersson, K.; Hansson, I.;
[47094] Lundstrom, K.; Andersson, L.: A paternally expressed QTL affecting
[47095] skeletal and cardiac muscle mass in pigs maps to the IGF2 locus. (Letter) Nature
[47096] Genet. 21: 157-158, 1999.

[47097] 7259.

[47098] Le Stunff, C.; Fallin, D.; Bougneres, P.: Paternal transmis-
[47099] sion of the very common class I INS VNTR alleles predisposes to childhood
[47100] obesity. Nature Genet. 29: 96-99, 2001.

[47101] 7260.

[47102] Lighten, A. D.; Hardy, K.; Winston, R. M. L.; Moore, G. E.: IGF2
[47103] is parentally imprinted in human preimplantation em-
bryos. (Letter) Nature

[47104] Genet. 122-123, 1997.

[47105] 7261.

[47106] Moore, T.; Constancia, M.; Zubair, M.; Bailleul, B.; Feil, R.;

[47107] Sasaki, H.; Reik, W.: Multiple imprinted sense and anti-
sense transcripts,

[47108] differential methylation and tandem repeats in a putative
imprinting

[47109] control region upstream of mouse Igf2. Proc. Nat. Acad.
Sci. 94:

[47110] 12509-12514, 1997.

[47111] 7262.

[47112] Henschen, A.; Kehl, M.; Deutsch, E.: Novel structure eluci-
dation

[47113] strategy for genetically abnormal fibrinogens with incom-
plete fibrinopeptide

[47114] release as applied to fibrinogen Schwarzach. Hoppe
Seylers Z. Physiol.

[47115] Chem. 364: 1747-1751, 1983.

[47116] 7263.

[47117] Henschen, A.; Southan, C.; Soria, J.; Soria, C.; Samama, M.:

[47118] Structure abnormality of fibrinogen Metz and its relation to the clotting

[47119] defect. (Abstract) Thromb. Haemost. 45: 103 only, 1981.

[47120] 7264.

[47121] Higgins, D. L.; Shafer, J. A.: Fibrinogen Petoskey, a dysfibrinogenemia

[47122] characterized by replacement of arg-A(alpha)16 by a histidyl residue:

[47123] evidence for thrombin-catalyzed hydrolysis at a histidyl residue. J.

[47124] Biol. Chem. 256: 12013-12017, 1981.

[47125] 7265.

[47126] Humphries, S. E.; Imam, A. M. A.; Robbins, T. P.; Cook, M.; Carritt,

[47127] B.; Ingle, C.; Williamson, R.: The identification of a DNA polymorphism

[47128] of the alpha fibrinogen gene, and the regional assignment of the human

[47129] fibrinogen genes to 4q26-qter. Hum. Genet. 68: 148-153, 1984.

[47130] 7266.

[47131] Imperato, C.; Dettori, A. G.: Ipofibrinogenemia congenita con
[47132] fibrinoastenia. Helv. Paediat. Acta 13: 380-399, 1958.

[47133] 7267.

[47134] Jackson, D. P.; Beck, E. A.; Charache, P.: Congenital disor-ders
[47135] of fibrinogen. Fed. Proc. 24: 816-821, 1965.

[47136] 7268.

[47137] Jandrot-Perrus, M.; Aurousseau, M.-H.; Rabiet, M.-J.; Josso, F.
[47138] : Fibrinogen Bondy: a new case of dysfibrinogenemia. Iso-lation of
[47139] the abnormal fibrinogen molecules. Thromb. Res. 27: 659-670, 1982.

[47140] 7269.

[47141] Kant, J. A.; Crabtree, G. R.: The rat fibrinogen genes: link-age
[47142] of the A-alpha and gamma chain genes. J. Biol. Chem. 258: 4666-4667,
[47143] 1983.

[47144] 7270.

[47145] Kant, J. A.; Fornace, A. J., Jr.; Saxe, D.; Simon, M. I.; McBride,
[47146] O. W.; Crabtree, G. R.: Evolution and organization of the fibrinogen
[47147] locus on chromosome 4: gene duplication accompanied by transposition
[47148] and inversion. Proc. Nat. Acad. Sci. 82: 2344-2348, 1985.

[47149] 7271.

[47150] Kohn, P. H.; Cruz, A. C.; Kitchens, C. S.: Autosomal dominant
[47151] hypofibrinogenemia associated with a balanced 7p/12q translocation
[47152] in three generations of a family. (Abstract) Clin. Res. 31: 316A
[47153] only, 1983.

[47154] 7272.

[47155] Koopman, J.; Haverkate, F.; Grimbergen, J.; Egbring, R.; Lord,
[47156] S. T.: Fibrinogen Marburg: a homozygous case of dysfibrinogenemia,
[47157] lacking amino acids A-alpha 461-610 (lys461 AAA-to-stop TAA). Blood 80:

[47158] 1972-1979, 1992.

[47159] 7273.

[47160] Koopman, J.; Haverkate, F.; Grimbergen, J.; Lord, S. T.; Mosesson,

[47161] M. W.; DiOrio, J. P.; Siebenlist, K. S.; Legrand, C.; Soria, J.; Soria,

[47162] C.; Caen, J. P.: Molecular basis for fibrinogen Dusart (A-alpha 554

[47163] arg-to-cys) and its association with abnormal fibrin polymerization

[47164] and thrombophilia. J. Clin. Invest. 91: 1637-1643, 1993.

[47165] 7274.

[47166] Kudryk, B.; Blomback, B.; Blomback, M.: Fibrinogen Detroit--an

[47167] abnormal fibrinogen with nonfunctional NH(2)-terminal polymerization

[47168] domain. Thromb. Res. 9: 25-36, 1976.

[47169] 7275.

[47170] Lachmann, H. J.; Chir, B.; Booth, D. R.; Booth, S. E.; Bybee,

[47171] A.; Gilbertson, J. A.; Gillmore, J. D.; Pepys, M. B.; Hawkins, P.

[47172] N.: Misdiagnosis of hereditary amyloidosis as AL (primary) amyloidosis. New

[47173] Eng. J. Med. 346: 1786-1791, 2002.

[47174] 7276.

[47175] Lee, M. H.; Kaczmarek, E.; Chin, D. T.; Oda, A.; McIntosh, S.;

[47176] Bauer, K. A.; Clyne, L. P.; McDonagh, J.: Fibrinogen Ledyard (A-alpha

[47177] arg16-to-cys): biochemical and physiologic characterization. Blood 78:

[47178] 1744-1752, 1991.

[47179] 7277.

[47180] Maekawa, H.; Yamazumi, K.; Muramatsu, S.; Kaneko, M.; Hirata,

[47181] H.; Takahashi, N.; Arocha-Pinango, C. L.; Rodriguez, S.; Nagy, H.;

[47182] Perez-Requejo, J. L.; Matsuda, M.: Fibrinogen Lima: a homozygous

[47183] dysfibrinogen with an A-alpha-arginine141-to-serine substitution associated

[47184] with extra N-glycosylation at A-alpha-asparagine-139--impaired fibrin

[47185] gel formation but normal fibrin-facilitated plasminogen activation
[47186] catalyzed by tissue-type plasminogen activator. J. Clin. In-
vest. 90:
[47187] 67-76, 1992.

[47188] 7278.

[47189] Maekawa, H.; Yamazumi, K.; Muramatsu, S.; Kaneko, M.; Hirata,
[47190] H.; Takahashi, N.; de Bosch, N. B.; Carvajal, Z.; Ojeda, A.; Arocha-Pinango,
[47191] C. L.; Matsuda, M.: An A-alpha ser-434 to N-glycosylated asn substitution
[47192] in a dysfibrinogen, fibrinogen Caracas II, characterized by impaired
[47193] fibrin gel formation. J. Biol. Chem. 266: 11575-11581, 1991.
[47194] 7279.

[47195] Mammen, E. F.; Prasad, A. S.; Barnhart, M. I.; Au, C. C.: Congenital
[47196] dysfibrinogenemia: fibrinogen Detroit. J. Clin. Invest. 48: 235-249,
[47197] 1969.

[47198] 7280.

[47199] Marder, V. J.; Budzynski, A. Z.: Fibrinogen and its derivatives,

[47200] hereditary and acquired abnormalities. Schweiz. Med. Wschr. 104:

[47201] 1338-1342, 1974.

[47202] 7281.

[47203] Marino, M. W.; Fuller, G. M.; Elder, F. F. B.: Chromosomal localization

[47204] of human and rat A-alpha, B-beta, and gamma fibrinogen genes by in

[47205] situ hybridization. Cytogenet. Cell Genet. 42: 36-41, 1986.

[47206] 7282.

[47207] Martinez, J.; Holburn, R. R.; Shapiro, S.; Erslev, A. J.: A hereditary

[47208] hypodysfibrinogenemia characterized by fibrinogen hypercatabolism. J.

[47209] Clin. Invest. 53: 600-611, 1974.

[47210] 7283.

[47211] McDonagh, R. P.; Carrell, N. A.; Roberts, H. R.; Blatt, P. M.;

[47212] McDonagh, J.: Fibrinogen Chapel Hill: hypodysfibrinogenemia with

[47213] a tertiary polymerization defect. Am. J. Hemat. 9: 23-38, 1980.

[47214] 7284.

[47215] Menache, D.: Dysfibrinogenemie constitutionnelle et familiale. Proc.

[47216] IX Cong. Europ. Soc. Hemat. 1255-1259, 1963.

[47217] 7285.

[47218] Menache, D.: Constitutional and familial abnormal fibrinogen. Thromb.

[47219] Diath. Haemorrh. 10 (suppl. 13): 173-185, 1964.

[47220] 7286.

[47221] Miyashita, C.; Schwamborn, J.; von Blohn, G.; Wenzel, E.; Hellstern,

[47222] P.: Preliminary report concerning two new cases of congenital dysfibrinogenemia

[47223] (Homburg II and Homburg III).In: Henschen, A.; Hessel, B.; McDonagh,

[47224] J.; Saldeen, T.: Fibrinogen--Structural Variants and Interactions.

[47225] Berlin: Walter de Gruyter (pub.) 1985. Pp. 237-246.

[47226] 7287.

[47227] Miyata, T.; Terukina, S.; Matsuda, M.; Kasamatsu, A.; Takeda,

[47228] Y.; Murakami, T.; Iwanaga, S.: Fibrinogens Kawaguchi and Osaka: an

[47229] amino acid substitution of A alpha arginine-16 to cysteine which forms

[47230] an extra interchain disulfide bridge between the two A al-pha chains. J.

[47231] Biochem. 102: 93-101, 1987.

[47232] 7288.

[47233] Morris, S.; Denninger, M. H.; Finlayson, J. S.; Menache, D.:

[47234] Fibrinogen Lille: A (alpha) 7 asp-to-asn. (Abstract) Thromb. Haemost. 46:

[47235] 104 only, 1981.

[47236] 7289.

[47237] Mosesson, M. W.; Siebenlist, K. R.; Hainfeld, J. F.; Wall, J.

[47238] S.; Soria, J.; Soria, C.; Caen, J. P.: The relationship between the

[47239] fibrinogen D domain self-association/cross-linking site (gamma-XL)

[47240] and the fibrinogen Dusart abnormality (A-alpha R554C-albumin): clues

[47241] to thrombophilia in the 'Dusart syndrome.' J. Clin. Invest. 97:

[47242] 2342-2350, 1996.

[47243] 7290.

[47244] Neerman-Arbez, M.; Antonarakis, S. E.; Honsberger, A.; Morris,

[47245] M. A.: The 11 kb FGA deletion responsible for congenital afibrinogenaemia

[47246] is mediated by a short direct repeat in the fibrinogen gene cluster. Europ.

[47247] J. Hum. Genet. 7: 897-902, 1999.

[47248] 7291.

[47249] Klugbauer, S.; Rabes, H. M.: The transcription coactivator HTIF1

[47250] and a related protein are fused to the RET receptor tyrosine kinase

[47251] in childhood papillary thyroid carcinomas. Oncogene 18: 4388-4393,

[47252] 1999.

[47253] 7292.

[47254] Lairmore, T. C.; Dou, S.; Howe, J. R.; Chi, D.; Carlson, K.; Veile,

[47255] R.; Mishra, S. K.; Wells, S. A., Jr.; Donis-Keller, H.: A 1.5-megabase

[47256] yeast artificial chromosome contig from human chromosome 10q11.2 connecting

[47257] three genetic loci (RET, D10S94, and D10S102) closely linked to the

[47258] MEN2A locus. Proc. Nat. Acad. Sci. 90: 492-496, 1993.

[47259] 7293.

[47260] Lombardo, F.; Baudin, E.; Chiefari, E.; Arturi, F.; Bardet, S.;

[47261] Caillou, B.; Conte, C.; Dallapiccola, B.; Giuffrida, D.; Bidart, J.-M.;

[47262] Schlumberger, M.; Filetti, S.: Familial medullary thyroid carcinoma:

[47263] clinical variability and low aggressiveness associated with RET mutation

[47264] at codon 804. J. Clin. Endocr. Metab. 87: 1674-1680, 2002.

[47265] 7294.

[47266] Lore, F.; Di Cairano, G.; Talidis, F.: Unilateral renal agene-
sis
[47267] in a family with medullary thyroid carcinoma. (Letter) New
Eng. J.
[47268] Med. 342: 1218-1219, 2000.

[47269] 7295.

[47270] Machens, A.; Gimm, O.; Hinze, R.; Hoppner, W.; Boehm, B.
O.; Dralle,
[47271] H.: Genotype-phenotype correlations in hereditary
medullary thyroid
[47272] carcinoma: oncological features and biochemical proper-
ties. J. Clin.
[47273] Endocr. Metab. 86: 1104-1109, 2001.

[47274] 7296.

[47275] Mendelsohn, C.; et al; et al: Function of the retinoic acid
receptors
[47276] (RARs) during development (II). Multiple abnormalities at
various
[47277] stages of organogenesis in RAR double mutants. Develop-
ment 120:

[47278] 2749-2771, 1994.

[47279] 7297.

[47280] Menko, F. H.; van der Luijt, R. B.; de Valk, I. A. J.; Toorians,

[47281] A. W. F. T.; Sepers, J. M.; van Diest, P. J.; Lips, C. J. M.:
Atypical

[47282] MEN type 2B associated with two germline RET mutations
on the same

[47283] allele not involving codon 918. J. Clin. Endocr. Metab. 87:
393-397,

[47284] 2002.

[47285] 7298.

[47286] Mulligan, L. M.; Kwok, J. B. J.; Healey, C. S.; Elsdon, M. J.;

[47287] Eng, C.; Gardner, E.; Love, D. R.; Mole, S. E.; Moore, J. K.;
Papi,

[47288] L.; Ponder, M. A.; Telenius, H.; Tunnacliffe, A.; Ponder, B.
A. J.

[47289] : Germ-line mutations of the RET proto-oncogene in mul-
tiple endocrine

[47290] neoplasia type 2A. Nature 363: 458-460, 1993.

[47291] 7299.

[47292] Munnes, M.; Fanaei, S.; Schmitz, B.; Muiznieks, I.; Holschneider,

[47293] A. M.; Doerfler, W.: Familial form of Hirschsprung disease: nucleotide

[47294] sequence studies reveal point mutations in the RET proto-oncogene

[47295] in two of six families but not in other candidate genes. Am. J. Med.

[47296] Genet. 94: 19-27, 2000.

[47297] 7300.

[47298] Nakata, T.; Kitamura, Y.; Shimizu, K.; Tanaka, S.; Fujimori, M.;

[47299] Yokoyama, S.; Ito, K.; Emi, M.: Fusion of a novel gene, ELKS, to

[47300] RET due to translocation t(10;12)(q11;p13) in a papillary thyroid

[47301] carcinoma. Genes Chromosomes Cancer 25: 97-103, 1999.

[47302] 7301.

[47303] Niccoli-Sire, P.; Murat, A.; Rohmer, V.; Franc, S.; Chabrier,

[47304] G.; Baldet, L.; Maes, B.; Savagner, F.; Giraud, S.; Bezieau, S.; Kottler,

[47305] M.-L.; Morange, S.; Conte-Devolx, B.: The French Calcitonin Tumors

[47306] Study Group (GETC).: Familial medullary thyroid carcinoma with noncysteine

[47307] RET mutations: phenotype-genotype relationship in a large series of

[47308] patients. J. Clin. Endocr. Metab. 86: 3746-3753, 2001.

[47309] 7302.

[47310] Pachnis, V.; Mankoo, B.; Costantini, F.: Expression of the c-ret

[47311] proto-oncogene during mouse embryogenesis. Development 119: 1005-1017,

[47312] 1993.

[47313] 7303.

[47314] Pasini, B.; Hofstra, R. M. W.; Yin, L.; Bocciardi, R.; Santamaria,

[47315] G.; Grootscholten, P. M.; Ceccherini, I.; Patrone, G.; Priolo, M.;

[47316] Buys, C. H. C. M.; Romeo, G.: The physical map of the human RET proto-oncogene. Oncogene 11:

[47317] 1737-1743, 1995.

[47318] 7304.

[47319] Pelet, A.; Geneste, O.; Edery, P.; Pasini, A.; Chappuis, S.; Attie,

[47320] T.; Munnich, A.; Lenoir, G.; Lyonnet, S.; Billaud, M.: Various mechanisms

[47321] cause RET-mediated signaling defects in Hirschsprung's disease. J.

[47322] Clin. Invest. 101: 1415-1423, 1998.

[47323] 7305.

[47324] Pigny, P.; Bauters, C.; Wemeau, J.-L.; Houcke, M. L.; Crepin,

[47325] M.; Caron, P.; Giraud, S.; Calender, A.; Buisine, M.-P.; Kerckaert,

[47326] J.-P.; Porchet, N.: A novel 9-base pair duplication in RET exon 8

[47327] in familial medullary thyroid carcinoma. J. Clin. Endocr. Metab. 84:

[47328] 1700-1704, 1999.

[47329] 7306.

[47330] Pierotti, M. A.; Santoro, M.; Jenkins, R. B.; Sozzi, G.; Bongarzone,

[47331] I.; Grieco, M.; Monzini, N.; Miozzo, M.; Herrmann, M. A.; Fusco, A.;

[47332] Hay, I. D.; Della Porta, G.; Vecchio, G.: Characterization of an

[47333] inversion on the long arm of chromosome 10 juxtaposing D10S170 and

[47334] RET and creating the oncogenic sequence RET/PTC. Proc. Nat. Acad.

[47335] Sci. 89: 1616–1620, 1992.

[47336] 7307.

[47337] Rodrigues, G. A.; Park, M.: Dimerization mediated through a leucine

[47338] zipper activates the oncogenic potential of the met receptor tyrosine

[47339] kinase. Molec. Cell. Biol. 13: 6711–6722, 1993.

[47340] 7308.

[47341] Romeo, G.; Ronchetto, P.; Luo, Y.; Barone, V.; Seri, M.; Ceccherini,

[47342] I.; Pasini, B.; Bocciardi, R.; Lerone, M.; Kaariainen, H.; Martucciello,

[47343] G.: Point mutations affecting the tyrosine kinase domain of the RET

[47344] proto-oncogene in Hirschsprung's disease. Nature 367: 377-378, 1994.

[47345] 7309.

[47346] Salvatore, D.; Barone, M. V.; Salvatore, G.; Melillo, R. M.; Chiappetta,

[47347] G.; Mineo, A.; Fenzi, G.; Vecchio, G.; Fusco, A.; Santoro, M.: Tyrosines

[47348] 1015 and 1062 are in vivo autophosphorylation sites in Ret and Ret-derived

[47349] oncoproteins. J. Clin. Endocr. Metab. 85: 3898-3907, 2000.

[47350] 7310.

[47351] Robinson, M. F.; Cote, G. J.; Nunziata, V.; Brandi, M. L.; Ferrer,

[47352] J. P.; Martins Bugalho, M. J. G.; Almeida Ruas, M. M.; Chik, C.; Colantuoni,

[47353] V.; Gagel, R. F.: Mutation of a specific codon of the RET proto-oncogene

[47354] in the multiple endocrine neoplasia type 2A/cutaneous lichen amyloidosis

[47355] syndrome. (Abstract) Fifth International Workshop on Multiple Endocrine

[47356] Neoplasia, Stockholm, Archipelago , 1994.

[47357] 7311.

[47358] Boyer, S. H.: Human organ alkaline phosphatases: dis-
crimination
[47359] by several means including starch gel electrophoresis of
antienzyme-enzyme
[47360] supernatant fluids. Ann. N.Y. Acad. Sci. 103: 938-950,
1963.

[47361] 7312.

[47362] Thompson, E. M.; Young, I. D.; Hall, C. M.; Pembrey, M. E.:
Recurrence
[47363] risks and prognosis in severe sporadic osteogenesis im-
perfecta. J.
[47364] Med. Genet. 24: 390-405, 1987.

[47365] 7313.

[47366] Deutman, A. F.; van Blommestein, J. D. A.; Henkes, H. E.;
Waardenburg,
[47367] P. J.; Solleveld-van Driest, E.: Butterfly-shaped pigment
dystrophy
[47368] of the fovea. Arch. Ophthal. 83: 558-569, 1970.

[47369] 7314.

[47370] Okuda, T.; Cleveland, J. L.; Downing, J. R.: PCTAIRE-1 and PCTAIRE-3,
[47371] two members of a novel cdc2/CDC28-related protein kinase gene family. Oncogene 7:
[47372] 2249-2258, 1992.

[47373] 7315.

[47374] Okuda, T.; Valentine, V. A.; Shapiro, D. N.; Downing, J. R.: Cloning
[47375] of genomic loci and chromosomal localization of the human PCTAIRE-1
[47376] and -3 protein kinase genes. Genomics 21: 217-221, 1994.
[47377] 7316.

[47378] Green, M. C.; Shultz, L. D.; Nedzi, L. A.: Abnormal nuclear morphology
[47379] of leukocytes in the mouse mutant ichthyosis. Transplantation 20:
[47380] 172-175, 1975.

[47381] 7317.

[47382] Hoffmann, K.; Dreger, C. K.; Olins, A. L.; Olins, D. E.; Shultz,

[47383] L. D.; Lucke, B.; Karl, H.; Kaps, R.; Muller, D.; Vaya, A.; Az-
nar,
[47384] J.; Ware, R. E.; Cruz, N. S.; Lindner, T. H.; Herrmann, H.;
Reis,
[47385] A.; Sperling, K.: Mutations in the gene encoding the lamin
B receptor
[47386] produce an altered nuclear morphology in granulocytes
(Pelger-Huet
[47387] anomaly). Nature Genet. 31: 410-414, 2002.

[47388] 7318.

[47389] Donald, L. J.: The genetics of placental alkaline phos-
phatase:
[47390] a possible 'null' allele. Ann. Hum. Genet. 38: 7-18, 1974.

[47391] 7319.

[47392] Donald, L. J.; Robson, E. B.: Rare variants of placental al-
kaline
[47393] phosphatase. Ann. Hum. Genet. 37: 303-313, 1974.

[47394] 7320.

[47395] Edwards, J. H.; Wingham, J.: Data on linkage between the
locus
[47396] determining placental alkaline phosphatase and other

markers. Ann.

[47397] Hum. Genet. 30: 233-237, 1967.

[47398] 7321.

[47399] Garattini, E.; Margolis, J.; Heimer, E.; Felix, A.; Udenfriend,

[47400] S.: Human placental alkaline phosphatase in liver and intestine. Proc.

[47401] Nat. Acad. Sci. 82: 6080-6084, 1985.

[47402] 7322.

[47403] Gogolin, K. J.; Slaughter, C. A.; Harris, H.: Electrophoresis

[47404] of enzyme-monoclonal antibody complexes: studies of human placental

[47405] alkaline phosphatase polymorphism. Proc. Nat. Acad. Sci. 78: 5061-5065,

[47406] 1981.

[47407] 7323.

[47408] Henthorn, P. S.; Knoll, B. J.; Raducha, M.; Rothblum, K. N.; Slaughter,

[47409] C.; Weiss, M.; Lafferty, M. A.; Fischer, T.; Harris, H.: Products

[47410] of two common alleles at the locus for human placental

alkaline phosphatase
[47411] differ by seven amino acids. Proc. Nat. Acad. Sci. 83:
5597-5601,
[47412] 1986.

[47413] 7324.

[47414] Kam, W.; Clauser, E.; Kim, Y. S.; Kan, Y. W.; Rutter, W. J.:

[47415] Cloning, sequencing, and chromosomal localization of
human term placental
[47416] alkaline phosphatase cDNA. Proc. Nat. Acad. Sci. 82:
8715-8719,
[47417] 1985.

[47418] 7325.

[47419] Knoll, B. J.; Rothblum, K. N.; Longley, M.: Nucleotide se-
quence
[47420] of the human placental alkaline phosphatase gene: evolu-
tion of the
[47421] 5-prime flanking region by deletion/substitution. J. Biol.
Chem. 263:
[47422] 12020-12027, 1988.

[47423] 7326.

[47424] Lucarelli, P.; Scacchi, R.; Corbo, R. M.; Benincasa, A.; Pal-

marino,
[47425] R.: Human placental alkaline phosphatase electrophoretic alleles:
[47426] quantitative studies. Am. J. Hum. Genet. 34: 331-336, 1982.
[47427] 7327.

[47428] Martin, D.; Spurr, N. K.; Trowsdale, J.: RFLP of the human placental
[47429] alkaline phosphatase gene (PLAP). Nucleic Acids Res. 15: 9104 only,
[47430] 1987.

[47431] 7328.

[47432] Martin, D.; Tucker, D. F.; Gorman, P.; Sheer, D.; Spurr, N. K.;
[47433] Trowsdale, J.: The human placental alkaline phosphatase gene and
[47434] related sequences map to chromosome 2 band q37. Ann. Hum. Genet. 51:
[47435] 145-152, 1987.

[47436] 7329.

[47437] Millan, J. L.; Beckman, G.; Jeppsson, A.; Stigbrand, T.: Ge-

netic

[47438] variants of placental alkaline phosphatase as detected by a monoclonal

[47439] antibody. Hum. Genet. 60: 145-149, 1982.

[47440] 7330.

[47441] Millan, J. L.; Stigbrand, T.: Antigenic determinants of human

[47442] placental and testicular placental-like alkaline phosphatases as mapped

[47443] by monoclonal antibodies. Europ. J. Biochem. 136: 1-7, 1983.

[47444] 7331.

[47445] Palmarino, R.; Corbo, R. M.; Lucarelli, P.: Human placental alkaline

[47446] phosphatase: analysis of genetically determined rare variants. Hum.

[47447] Biol. 51: 341-352, 1979.

[47448] 7332.

[47449] Raimondi, E.; Talarico, D.; Moro, L.; Rutter, W. J.; Della Valle,

[47450] G.; De Carli, L.: Regional mapping of the human placental

alkaline
[47451] phosphatase gene (ALPP) to 2q37 by in situ hybridization. Cytogenet.
[47452] Cell Genet. 47: 98-99, 1988.

[47453] 7333.

[47454] Robinson, J. C.; Goldsmith, L. A.: Genetically determined variants
[47455] of serum alkaline phosphatase: a review. Vox Sang. 13: 289-307,
[47456] 1967.

[47457] 7334.

[47458] Robson, E. B.; Harris, H.: Genetics of the alkaline phos-phatase
[47459] polymorphism of the human placenta. Nature 207: 1257-1259, 1965.
[47460] 7335.

[47461] Slaughter, C. A.; Gogolin, K. J.; Coseo, M. C.; Meyer, L. J.;

[47462] Lesko, J.; Harris, H.: Discrimination of human placental al-kaline
[47463] phosphatase allelic variants by monoclonal antibodies. Am. J. Hum.

[47464] Genet. 35: 1-20, 1983.

[47465] 7336.

[47466] Beckman, G.; Beckman, L.; Kivela, A.; Millan, J. L.; Sik-
strom,
[47467] C.: A new PstI restriction fragment length polymorphism
(RFLP) of
[47468] placental alkaline phosphatase: RFLP haplotypes and cor-
relation with
[47469] electrophoretic types. Hum. Hered. 41: 122-128, 1991.

[47470] 7337.

[47471] Beckman, G.; Beckman, L.; Sikstrom, C.; Millan, J. L.: DNA
polymorphism
[47472] of alkaline phosphatase isozyme genes: linkage disequi-
libria between
[47473] placental and germ-cell alkaline phosphatase alleles. Am.
J. Hum.
[47474] Genet. 51: 1066-1070, 1992.

[47475] 7338.

[47476] Goldstein, D. J.; Gogolin, K. J.; Harris, H.: Possible new lo-
cus
[47477] for alkaline phosphatase in humans. (Abstract) Sixth Int.

Cong. Hum.

[47478] Genet., Jerusalem , 1981.

[47479] 7339.

[47480] Millan, J. L.; Eriksson, A.; Stigbrand, T.: A possible new locus

[47481] of alkaline phosphatase expressed in human testis. Hum. Genet. 62:

[47482] 293-295, 1982.

[47483] 7340.

[47484] Millan, J. L.; Manes, T.: Seminoma-derived Nagao isozyme is encoded

[47485] by a germ-cell alkaline phosphatase gene. Proc. Nat. Acad. Sci. 85:

[47486] 3024-3028, 1988.

[47487] 7341.

[47488] Abe, A.; Emi, N.; Tanimoto, M.; Terasaki, H.; Marunouchi, T.; Saito,

[47489] H.: Fusion of the platelet-derived growth factor receptor beta to

[47490] a novel gene CEV14 in acute myelogenous leukemia after clonal evolution. Blood 90:

[47491] 4271-4277, 1997.

[47492] 7342.

[47493] Buchberg, A. M.; Jenkins, N. A.; Copeland, N. G.: Localization

[47494] of the murine macrophage colony-stimulating factor gene to chromosome

[47495] 3 using interspecific backcross analysis. Genomics 5: 363-367, 1989.

[47496] 7343.

[47497] Claesson-Welsh, L.; Eriksson, A.; Moren, A.; Severinsson, L.; Ek,

[47498] B.; Ostman, A.; Betsholtz, C.; Heldin, C.-H.: cDNA cloning and expression

[47499] of a human platelet-derived growth factor (PDGF) receptor specific

[47500] for B-chain-containing PDGF molecules. Molec. Cell. Biol. 8: 3476-3486,

[47501] 1988.

[47502] 7344.

[47503] Escobedo, J. A.; Fried, V. A.; Daniel, T. O.; Williams, L. T.:

[47504] Primary structure of the platelet-derived growth factor.

(Abstract) Clin.

[47505] Res. 34: 544A, 1986.

[47506] 7345.

[47507] Gronwald, R. G. K.; Grant, F. J.; Haldeman, B. A.; Hart, C. E.;

[47508] O'Hara, P. J.; Hagen, F. S.; Ross, R.; Bowen-Pope, D. F.; Murray,

[47509] M. J.: Cloning and expression of a cDNA coding for the human platelet-derived

[47510] growth factor receptor: evidence for more than one receptor class. Proc.

[47511] Nat. Acad. Sci. 85: 3435-3439, 1988.

[47512] 7346.

[47513] Klinghoffer, R. A.; Mueting-Nelsen, P. F.; Faerman, A.; Shani,

[47514] M.; Soriano, P.: The two PDGF receptors maintain conserved signaling

[47515] in vivo despite divergent embryological functions. Molec. Cell 7:

[47516] 343-354, 2001.

[47517] 7347.

[47518] Kulkarni, S.; Heath, C.; Parker, S.; Chase, A.; Iqbal, S.; Pocock,

[47519] C. F.; Kaeda, J.; Cwynarski, K.; Goldman, J. M.; Cross, N. C. P.:

[47520] Fusion of H4/D10S170 to the platelet-derived growth factor receptor

[47521] beta in BCR-ABL-negative myeloproliferative disorders with a t(5;10)(q33;q21). Cancer

[47522] Res. 60: 3592-3598, 2000.

[47523] 7348.

[47524] Leal, F.; Williams, L. T.; Robbins, K. C.; Aaronson, S. A.: Evidence

[47525] that the v-sis gene product transforms by interaction with the receptor

[47526] for platelet-derived growth factor. Science 230: 327-330, 1985.

[47527] 7349.

[47528] Magnusson, M. K.; Meade, K. E.; Brown, K. E., Arthur, D. C.; Krueger,

[47529] L. A.; Barrett, A. J.; Dunbar, C. E.: Rabaptin-5 is a novel fusion

[47530] partner to platelet-derived growth factor beta receptor in

chronic
[47531] myelomonocytic leukemia. Blood 98: 2518-2525, 2001.

[47532] 7350.

[47533] Matsui, T.; Heidaran, M.; Miki, T.; Popescu, N.; La Rochelle,
[47534] W.; Kraus, M.; Pierce, J.; Aaronson, S.: Isolation of a novel receptor
[47535] cDNA establishes the existence of two PDGF receptor genes. Science 243:
[47536] 800-804, 1989.

[47537] 7351.

[47538] Ross, T. S.; Bernard, O. A.; Berger, R.; Gilliland, D. G.: Fu-
[47539] sion of Huntingtin interacting protein 1 to platelet-derived growth factor-beta
[47540] receptor (PDGF-beta-R) in chronic myelomonocytic leukemia with t(5;7)(q33;q11.2). Blood 91:
[47541] 4419-4426, 1998.

[47542] 7352.

[47543] Steer, E. J.; Cross, N. C. P.: Myeloproliferative disorders with

[47544] translocations of chromosome 5q31-35: role of the platelet-derived
[47545] growth factor receptor beta. Acta Haemat. 107: 113-122, 2002.
[47546] 7353.

[47547] Treacher Collins Syndrome Collaborative Group: Positional cloning
[47548] of a gene involved in the pathogenesis of Treacher Collins syndrome. Nature
[47549] Genet. 12: 130-136, 1996.

[47550] 7354.

[47551] Yarden, Y.; Escobedo, J. A.; Kuang, W.-J.; Yang-Feng, T. L.; Daniel,
[47552] T. O.; Tremble, P. M.; Chen, E. Y.; Ando, M. E.; Harkins, R. N.; Francke,
[47553] U.; Fried, V. A.; Ullrich, A.; Williams, L. T.: Structure of the

[47554] receptor for platelet-derived growth factor helps define a family
[47555] of closely related growth factor receptors. Nature 323: 226-232,
[47556] 1986.

[47557] 7355.

[47558] Allcock, R. J. N.; Williams, J. H.; Price, P.: The central MHC

[47559] gene, BAT1, may encode a protein that down-regulates cytokine production. Genes

[47560] Cells 6: 487-494, 2001.

[47561] 7356.

[47562] Nunes, M.; Peelman, L.; Vaiman, M.; Bourgeaux, N.; Chardon, P.

[47563] : Characterization of six new loci within the swine major histocompatibility

[47564] complex class III region. Mammalian Genome 5: 616-622, 1994.

[47565] 7357.

[47566] Peelman, L. J.; Chardon, P.; Nunes, M.; Renard, C.; Gef-

frotin,

[47567] C.; Vaiman, M.; Van Zeveren, A.; Coppieters, W.; van de Weghe, A.;

[47568] Bouquet, Y.; Choy, W. W.; Strominger, J. L.; Spies, T.: The BAT1

[47569] gene in the MHC encodes an evolutionarily conserved putative nuclear

[47570] RNA helicase of the DEAD family. Genomics 26: 210-218, 1995.

[47571] 7358.

[47572] Spies, T.; Blanck, G.; Bresnahan, M.; Sands, J.; Strominger, J.

[47573] L.: A new cluster of genes within the human major histocompatibility

[47574] complex. Science 243: 214-217, 1989.

[47575] 7359.

[47576] Spies, T.; Bresnahan, M.; Strominger, J. L.: Human major histocompatibility

[47577] complex contains a minimum of 19 genes between the complement cluster

[47578] and HLA-B. Proc. Nat. Acad. Sci. 86: 8955-8958, 1989.

[47579] 7360.

[47580] Colosimo, A.; Calabrese, G.; Gennarelli, M.; Ruzzo, A. M.; Sangiuolo,

[47581] F.; Magnani, M.; Palka, G.; Novelli, G.; Dallapiccola, B.: Assignment

[47582] of the hexokinase type 3 gene (HK3) to human chromosome band 5q35.3

[47583] by somatic cell hybrids and in situ hybridization. Cyto-
genet. Cell
[47584] Genet. 74: 187-188, 1996.

[47585] 7361.

[47586] Furuta, H.; Nishi, S.; Le Beau, M. M.; Fernald, A. A.; Yano,
H.;
[47587] Bell, G. I.: Sequence of human hexokinase III cDNA and
assignment
[47588] of the human hexokinase III gene (HK3) to chromosome
band 5q35.2 by
[47589] fluorescence in situ hybridization. Genomics 36: 206-209,
1996.
[47590] 7362.

[47591] Harris, H.; Hopkinson, D. A.: Average heterozygosity per
locus
[47592] in man: an estimate based on the incidence of enzyme
polymorphism. Ann.
[47593] Hum. Genet. 36: 9-20, 1972.

[47594] 7363.

[47595] Hopkinson, D. A.; Edwards, Y. H.; Harris, H.: The distribu-
tion

[47596] of subunit numbers and subunit sizes of enzymes: a study of the products
[47597] of 100 gene loci. Ann. Hum. Genet. 39: 383-411, 1976.

[47598] 7364.

[47599] Povey, S.; Corney, G.; Harris, H.: Genetically determined polymorphism
[47600] in a form of hexokinase, HK III, found in human leukocytes. Ann.
[47601] Hum. Genet. 38: 407-415, 1975.

[47602] 7365.

[47603] Altay, C.; Alper, C. A.; Nathan, D. G.: Normal and variant isoenzymes
[47604] of human blood cell hexokinase and the isoenzyme pattern in hemolytic
[47605] anemia. Blood 36: 219-227, 1970.

[47606] 7366.

[47607] Andreoni, F.; Ruzzo, A.; Magnani, M.: Structure of the 5-prime
[47608] region of the human hexokinase type I (HKI) gene and identification
[47609] of an additional testis-specific HKI mRNA. Biochim. Bio-

phys. Acta 1493:
[47610] 19-26, 2000.

[47611] 7367.

[47612] Bianchi, M.; Magnani, M.: Hexokinase mutations that pro-
duce nonspherocytic
[47613] hemolytic anemia. Blood Cells Molecules Dis. 21: 2-8,
1995.
[47614] 7368.

[47615] Chern, C. J.: Localization of the structural genes for hex-
okinase-1
[47616] and inorganic pyrophosphatase on region (pter-q24) of
human chromosome
[47617] 10. Cytogenet. Cell Genet. 17: 338-342, 1976.

[47618] 7369.

[47619] Dallapiccola, B.; Lungarotti, M. S.; Magnani, M.; Dacha, M.:
Evidence
[47620] of gene dosage effect for HK1 in the red cells of a patient
with trisomy
[47621] 10pter leads to p13. Ann. Genet. 24: 45-47, 1981.

[47622] 7370.

[47623] Dallapiccola, B.; Novelli, G.; Micara, G.; Delaroche, I.;

Moric-Petrovic,
[47624] S.; Magnani, M.: Regional mapping of hexokinase-1 within the short
[47625] arm of chromosome 10. Hum. Hered. 34: 156-160, 1984.

[47626] 7371.

[47627] Daniele, A.; Altruda, F.; Ferrone, M.; Silengo, L.; Romeo, G.;
[47628] Archidiacono, N.; Rocchi, M.: Mapping of human hexokinase 1 gene
[47629] to 10q11-qter. Hum. Hered. 42: 107-110, 1992.

[47630] 7372.

[47631] Gelb, B. D.; Worley, K. C.; Griffin, L. D.; Adams, V.; Chinault,
[47632] A. C.; McCabe, E. R. B.: Characterization of human genomic artificial
[47633] chromosome inserts containing hexokinase 1 coding information on chromosome
[47634] 10. Biochem. Med. Metab. Biol. 47: 267-269, 1992.

[47635] 7373.

[47636] Gitelman, B. J.; Simpson, N. E.: Regional mapping of the locus

[47637] for hexokinase-1 (HK1) to 10p11-q23 by gene dosage in human fibroblasts. Hum.
[47638] Genet. 60: 227-229, 1982.

[47639] 7374.

[47640] Griffin, L. D.; Gelb, B. D.; Wheeler, D. A.; Davison, D.; Adams,
[47641] V.; McCabe, E. R. B.: Mammalian hexokinase 1: evolution-ary conservation
[47642] and structure to function analysis. Genomics 11: 1014-1024, 1991.
[47643] 7375.

[47644] Mori, C.; Welch, J. E.; Fulcher, K. D.; O'Brien, D. A.; Eddy,
[47645] E. M.: Unique hexokinase messenger ribonucleic acids lacking the
[47646] porin-binding domain are developmentally expressed in mouse spermatogenic
[47647] cells. Biol. Reprod. 49: 191-203, 1993.

[47648] 7376.

[47649] Ollendorff, V.; Szepetowski, P.; Mattei, M.-G.; Gaudray, P.; Birnbaum,
[47650] D.: New gene in the homologous human 11q13-q14 and

mouse 7F chromosomal
[47651] regions. Mammalian Genome 2: 195-200, 1992.

[47652] 7377.

[47653] Guilford, P.; Hopkins, J.; Harraway, J.; McLeod, M.; McLeod, N.;
[47654] Harawira, P.; Taite, H.; Scoular, R.; Miller, A.; Reeve, A. E.: E-cadherin
[47655] germline mutations in familial gastric cancer. Nature 392: 402-405,
[47656] 1998.

[47657] 7378.

[47658] Huntsman, D. G.; Carneiro, F.; Lewis, F. R.; MacLeod, P. M.; Hayashi,
[47659] A.; Monaghan, K. G.; Maung, R.; Seruca, R.; Jackson, C. E.; Caldas,
[47660] C.: Early gastric cancer in young, asymptomatic carriers of germ-line
[47661] E-cadherin mutations. New Eng. J. Med. 344: 1904-1909, 2001.
[47662] 7379.

[47663] Karl, M.; Lamberts, S. W. J.; Detera-Wadleigh, S. D.; Encio,

I.

[47664] J.; Stratakis, C. A.; Hurley, D. M.; Accili, D.; Chrousos, G. P.:

[47665] Familial glucocorticoid resistance caused by a splice site deletion

[47666] in the human glucocorticoid receptor gene. J. Clin. Endocr. Metab. 76:

[47667] 683-689, 1993.

[47668] 7380.

[47669] Kino, T.; Stauber, R. H.; Resau, J. H.; Pavlakis, G. N.; Chrousos,

[47670] G. P.: Pathologic human GR mutant has a transdominant negative effect

[47671] on the wild-type GR by inhibiting its translocation into the nucleus:

[47672] importance of the ligand-binding domain for intracellular GR trafficking. J.

[47673] Clin. Endocr. Metab. 86: 5600-5608, 2001.

[47674] 7381.

[47675] Kontula, K.; Pelkonen, R.; Andersson, L.; Sivula, A.: Gluco-corticoid

[47676] receptors in adrenocorticoid disorders. J. Clin. Endocr.

Metab. 51:
[47677] 654-657, 1980.

[47678] 7382.

[47679] Koper, J. W.; Stolk, R. P.; de Lange, P.; Huizenga, N. A.; Molijn,
[47680] G. J.; Pols, H. A.; Grobbee, D. E.; Karl, M.; de Jong, F. H.; Brinkmann,
[47681] A. O.; Lamberts, S. W.: Lack of association between five polymorphisms
[47682] in the human glucocorticoid receptor gene and glucocor-ticoid resistance. Hum.
[47683] Genet. 99: 663-668, 1997.

[47684] 7383.

[47685] Lamberts, S. W. J.; Poldermans, D.; Zweens, M.; de Jong, F. H.
[47686] : Familial cortisol resistance: differential diagnostic and therapeutic
[47687] aspects. J. Clin. Endocr. Metab. 63: 1328-1333, 1986.

[47688] 7384.

[47689] Lind, U.; Carlstedt-Duke, J.; Gustafsson, J.-A.; Wright, A. P.

[47690] H.: Identification of single amino acid substitutions of cys-736

[47691] that affect the steroid-binding affinity and specificity of the glucocorticoid

[47692] receptor using phenotypic screening in yeast. Molec. Endocr. 10:

[47693] 1358-1370, 1996.

[47694] 7385.

[47695] Lipsett, M. B.; Chrousos, G. P.; Tomita, M.; Brandon, D. D.; Loriaux,

[47696] D. L.: The defective glucocorticoid receptor in man and nonhuman

[47697] primates. Recent Prog. Horm. Res. 41: 199-247, 1985.

[47698] 7386.

[47699] Lipsett, M. B.; Tomita, M.; Brandon, D. D.; De Vroede, M. M.;

[47700] Loriaux, D. L.; Chrousos, G. P.: Cortisol resistance in man.In: Chrousos,

[47701] G. P.; Loriaux, D. L.; Lipsett, M. B.: Steroid Hormone Resistance:

[47702] Mechanisms and Clinical Aspects. New York: Plenum Press (pub.)

[47703] 1986. Pp. 97-109.

[47704] 7387.

[47705] McNally, J. G.; Muller, W. G.; Walker, D.; Wolford, R.; Hager,

[47706] G. L.: The glucocorticoid receptor: rapid exchange with regulatory

[47707] sites in living cells. Science 287: 1262-1265, 2000.

[47708] 7388.

[47709] Montkowski, A.; Barden, N.; Wotjak, C.; Stec, I.; Ganster, J.;

[47710] Meaney, M.; Engelmann, M.; Reul, J. M. H. M.; Landgraf, R.; Holsboer,

[47711] F.: Long-term antidepressant treatment reduces be-

havioural deficits

[47712] in transgenic mice with impaired glucocorticoid receptor function. J.

[47713] Neuroendocrinology 7: 841-845, 1995.

[47714] 7389.

[47715] Muller, M.; Renkawitz, R.: Review: the glucocorticoid re-

ceptor. Biochim.

[47716] Biophys. Acta 1088: 171-182, 1991.

[47717] 7390.

[47718] Nawata, H.; Sekiya, K.; Higuchi, K.; Kato, K.-I.; Ibayashi, H.

[47719] : Decreased deoxyribonucleic acid binding of glucocorti-
coid-receptor

[47720] complex in cultured skin fibroblasts from a patient with
the glucocorticoid

[47721] resistance syndrome. J. Clin. Endocr. Metab. 65: 219-226,
1987.

[47722] 7391.

[47723] Oakley, R. H.; Sar, M.; Cidlowski, J. A.: The human gluco-
corticoid

[47724] receptor isoform: expression, biochemical properties, and
putative

[47725] function. J. Biol. Chem. 271: 9550-9559, 1996.

[47726] 7392.

[47727] Pepin, M.-C.; Pothier, F.; Barden, N.: Impaired type II glu-
cocorticoid-receptor

[47728] function in mice bearing antisense RNA transgene. Nature
355: 725-728,

[47729] 1992.

[47730] 7393.

[47731] Ray, D. W.; Davis, J. R. E.; White, A.; Clark, A. J. L.: Gluco-
corticoid
[47732] receptor structure and function in glucocorticoid-resistant
small
[47733] cell lung carcinoma cells. Cancer Res. 56: 3276-3280,
1996.
[47734] 7394.

[47735] Reichardt, H. M.; Kaestner, K. H.; Tuckermann, J.; Kretz,
O.;
[47736] Wessely, O.; Bock, R.; Gass, P.; Schmid, W.; Herrlich, P.;
Angel,
[47737] P.; Schutz, G.: DNA binding of the glucocorticoid receptor
is not
[47738] essential for survival. Cell 93: 531-541, 1998.

[47739] 7395.

[47740] Rivers, C.; Levy, A.; Hancock, J.; Lightman, S.; Norman, M.:

[47741] Insertion of an amino acid in the DNA-binding domain of
the glucocorticoid
[47742] receptor as a result of alternative splicing. J. Clin. Endocr.
Metab. 84:
[47743] 4283-4286, 1999.

[47744] 7396.

[47745] Roux, S.; Terouanne, B.; Balaguer, P.; Jausons-Loffreda, N.; Pons,

[47746] M.; Chambon, P.; Gronemeyer, H.; Nicolas, J.-C.: Mutation of isoleucine

[47747] 747 by a threonine alters the ligand responsiveness of the human glucocorticoid

[47748] receptor. Molec. Endocr. 10: 1214-1226, 1996.

[47749] 7397.

[47750] Strickland, I.; Kisich, K.; Hauk, P. J.; Vottero, A.; Chrousos,

[47751] G. P.; Klemm, D. J.; Leung, D. Y. M.: High constitutive glucocorticoid

[47752] receptor beta in human neutrophils enables them to reduce their spontaneous

[47753] rate of cell death in response to corticosteroids. J. Exp. Med. 193:

[47754] 585-593, 2001.

[47755] 7398.

[47756] Theriault, A.; Boyd, E.; Harrap, S. B.; Hollenberg, S. M.; Connor,

[47757] J. M.: Regional chromosomal assignment of the human

glucocorticoid
[47758] receptor gene to 5q31. Hum. Genet. 83: 289-291, 1989.

[47759] 7399.

[47760] Theriault, A.; Harrap, S. B.; Hollenberg, S. M.; Boyd, E.; Connor,
[47761] J. M.: Regional chromosomal assignment of the glucocorticoid receptor
[47762] gene to 5q31. (Abstract) Cytogenet. Cell Genet. 51: 1089, 1989.
[47763] 7400.

[47764] Tronche, F.; Kellendonk, C.; Kretz, O.; Gass, P.; Anlag, K.; Orban,
[47765] P. C.; Bock, R.; Klein, R.; Schutz, G.: Disruption of the glucocorticoid
[47766] receptor gene in the nervous system results in reduced anxiety. Nature
[47767] Genet. 23: 99-103, 1999.

[47768] 7401.

[47769] Vingerhoeds, A. C. M.; Thijssen, J. H. H.; Schwarz, F.: Spontaneous
[47770] hypercortisolism without Cushing's syndrome. J. Clin. En-

docr. Metab. 43:

[47771] 1128-1133, 1976.

[47772] 7402.

[47773] Webster, J. C.; Oakley, R. H.; Jewell, C. M.; Cidlowski, J. A.

[47774] : Proinflammatory cytokines regulate human glucocorti-
coid receptor

[47775] gene expression and lead to the accumulation of the
dominant negative

[47776] beta-isoform: a mechanism for the generation of gluco-
corticoid resistance. Proc.

[47777] Nat. Acad. Sci. 98: 6865-6870, 2001.

[47778] 7403.

[47779] Weinberger, C.; Evans, R.; Rosenfeld, M. G.; Hollenberg, S.
M.;

[47780] Skarecky, D.; Wasmuth, J. J.: Assignment of the human
gene encoding

[47781] the glucocorticoid receptor to the q11-q13 region on
chromosome 5.

[47782] (Abstract) Cytogenet. Cell Genet. 40: 776, 1985.

[47783] 7404.

[47784] Weinberger, C.; Giguere, V.; Hollenberg, S. M.; Thompson,

C.;

[47785] Arriza, J.; Evans, R. M.: Human steroid receptors and erb-A gene

[47786] products form a superfamily of enhancer-binding proteins. Clin. Physiol.

[47787] Biochem. 5: 179-189, 1987.

[47788] 7405.

[47789] Weinberger, C.; Hollenberg, S. M.; Ong, E. S.; Harmon, J. M.;

[47790] Brower, S. T.; Cidlowski, J.; Thompson, E. B.; Rosenfeld, M. G.; Evans,

[47791] R. M.: Identification of human glucocorticoid receptor complementary

[47792] DNA clones by epitope selection. Science 228: 740-742, 1985.

[47793] 7406.

[47794] Seeburg, P. H.: The human growth hormone gene family: nucleotide

[47795] sequences show recent divergence and predict a new polypeptide hormone. DNA 1:

[47796] 239-249, 1982.

[47797] 7407.

[47798] Abdul-Latif, H.; Leiberman, E.; Brown, M. R.; Carmi, R.; Parks,
[47799] J. S.: Growth hormone deficiency type IB caused by cryptic splicing
[47800] of the GH-1 gene. J. Pediat. Endocr. Metab. 13: 21-28, 2000.
[47801] 7408.

[47802] Akinci, A.; Kanaka, C.; Eble, A.; Akar, N.; Vidinlisan, S.; Mullis,
[47803] P. E.: Isolated growth hormone (GH) deficiency type IA associated
[47804] with a 45-kilobase gene deletion within the human GH gene cluster. J.
[47805] Clin. Endocr. Metab. 75: 437-441, 1992.

[47806] 7409.

[47807] Archer, J. S.; Kennan, W. S.; Gould, M. N.; Bremel, R. D.: Human
[47808] growth hormone (hGH) secretion in milk of goats after direct transfer
[47809] of the hGH gene into the mammary gland by using replication-defective
[47810] retrovirus vectors. Proc. Nat. Acad. Sci. 91: 6840-6844,

1994.

[47811] 7410.

[47812] Baum, H. B. A.; Katznelson, L.; Sherman, J. C.; Biller, B. M. K.;

[47813] Hayden, D. L.; Schoenfeld, D. A.; Cannistraro, K. E.; Klibanski, A.

[47814] : Effects of physiological growth hormone (GH) therapy on cognition

[47815] and quality of life in patients with adult-onset GH deficiency. J.

[47816] Clin. Endocr. Metab. 83: 3184-3189, 1998.

[47817] 7411.

[47818] Baxter, J. D.: Personal Communication. San Francisco, Calif.

[47819] 4/1981.

[47820] 7412.

[47821] Baxter, J. D.; Seeburg, P. H.; Shine, J.; Martial, J. A.; Goodman,

[47822] H. M.: The gene for growth hormone (GH): DNA sequence, expression,

[47823] regulation. (Abstract) Clin. Res. 25: 461A, 1977.

[47824] 7413.

[47825] Behncken, S. N.; Rowlinson, S. W.; Rowland, J. E.; Conway-Campbell,

[47826] B. L.; Monks, T. A.; Waters, M. J.: Aspartate 171 is the major primate-specific

[47827] determinant of human growth hormone. J. Biol. Chem. 272: 27077-27083,

[47828] 1997.

[47829] 7414.

[47830] Bianda, T.; Glatz, Y.; Bouillon, R.; Froesch, E. R.; Schmid, C.

[47831] : Effects of short-term insulin-like growth factor-I (IGF-I) or growth

[47832] hormone (GH) treatment on bone metabolism and on production of 1,25-dihydroxycholecalciferol

[47833] in GH-deficient adults. J. Clin. Endocr. Metab. 98: 81-87, 1998.

[47834] 7415.

[47835] Binder, G.; Brown, M.; Parks, J. S.: Mechanisms responsible for

[47836] dominant expression of human growth hormone gene mutations. J. Clin.

[47837] Endocr. Metab. 81: 4047-4050, 1996.

[47838] 7416.

[47839] Binder, G.; Ranke, M. B.: Screening for growth hormone (GH) gene

[47840] splice-site mutations in sporadic cases with severe isolated GH deficiency

[47841] using ectopic transcript analysis. J. Clin. Endocr. Metab. 80: 1247-1252,

[47842] 1995.

[47843] 7417.

[47844] Blethen, S. L.; Baptista, J.; Kuntze, J.; Foley, T.; LaFranchi,

[47845] S.; Johanson, A.: Adult height in growth hormone (GH)-deficient children

[47846] treated with biosynthetic GH. J. Clin. Endocr. Metab. 82: 418-420,

[47847] 1997.

[47848] 7418.

[47849] Boguszewski, C. L.; Jansson, C.; Boguszewski, M. C. S.; Rosberg,

[47850] S.; Carlsson, B.; Albertsson-Wikland, K.; Carlsson, L. M. S.: Increased

[47851] proportion of circulating non-22-kilodalton growth hormone isoforms

[47852] in short children: a possible mechanism for growth failure. J. Clin.

[47853] Endocr. Metab. 82: 2944-2949, 1997.

[47854] 7419.

[47855] Boguszewski, C. L.; Johannsson, G.; Bengtsson, B.-A.; Johansson,

[47856] A.; Carlsson, B.; Carlsson, L. M. S.: Circulating non-22-kilodalton

[47857] growth hormone isoforms in acromegalic men before and after transsphenoidal

[47858] surgery. J. Clin. Endocr. Metab. 82: 1516-1521, 1997.

[47859] 7420.

[47860] Braga, S.; Phillips, J. A., III; Joss, E.; Schwarz, H.; Zuppinger,

[47861] K.: Familial growth hormone deficiency resulting from a 7.6 kb deletion

[47862] within the growth hormone gene cluster. Am. J. Med. Genet. 25: 443-452,

[47863] 1986.

[47864] 7421.

[47865] Burman, P.; Johansson, A. G.; Siegbahn, A.; Vessby, B.; Karlsson,

[47866] F. A.: Growth hormone (GH)-deficient men are more responsive to GH

[47867] replacement therapy than women. J. Clin. Endocr. Metab. 82: 550-555,

[47868] 1997.

[47869] 7422.

[47870] Cassorla, F.; Mericq, V.; Eggers, M.; Avila, A.; Garcia, C.; Fuentes,

[47871] A.; Rose, S. R.; Cutler, G. B., Jr.: Effects of luteinizing hormone-releasing

[47872] hormone analog-induced pubertal delay in growth hormone (GH)-deficient

[47873] children treated with GH: preliminary results. J. Clin. Endocr. Metab. 82:

[47874] 3989-3992, 1997.

[47875] 7423.

[47876] Chakravarti, A.; Phillips, J. A., III; Mellits, K. H.; Buetow,

[47877] K. H.; Seeburg, P. H.: Patterns of polymorphism and link-

age disequilibrium

[47878] suggest independent origins of the human growth hormone gene cluster. Proc.

[47879] Nat. Acad. Sci. 81: 6085-6089, 1984.

[47880] 7424.

[47881] Chapman, I. M.; Pescovitz, O. H.; Murphy, G.; Treep, T.; Cerchio,

[47882] K. A.; Krupa, D.; Gertz, B.; Polvino, W. J.; Skiles, E. H.; Pezzoli,

[47883] S. S.; Thorner, M. O.: Oral administration of growth hormone (GH)

[47884] releasing peptide-mimetic MK-677 stimulates the GH/insulin-like growth

[47885] factor-I axis in selected GH-deficient adults. J. Clin. Endocr. Metab. 82:

[47886] 3455-3463, 1997.

[47887] 7425.

[47888] Chen, E.: Personal Communication. San Francisco, Calif. 10/25/1988.

[47889] 7426.

[47890] Chen, S.; Leger, J.; Garel, C.; Hassan, M.; Czernichow, P.:

Growth
[47891] hormone deficiency with ectopic neurohypophysis: anatomical variations
[47892] and relationship between the visibility of the pituitary stalk asserted
[47893] by magnetic resonance imaging and anterior pituitary function. J.
[47894] Clin. Endocr. Metab. 84: 2408-2413, 1999.

[47895] 7427.

[47896] Cogan, J. D.; Phillips, J. A., III; Sakati, N.; Frisch, H.; Schober,
[47897] E.; Milner, R. D. G.: Heterogeneous growth hormone (GH) gene mutations
[47898] in familial GH deficiency. J. Clin. Endocr. Metab. 76: 1224-1228,
[47899] 1993.

[47900] 7428.

[47901] Cogan, J. D.; Prince, M. A.; Lekhakula, S.; Bundey, S.; Fu-trakul,
[47902] A.; McCarthy, E. M. S.; Phillips, J. A., III: A novel mecha-nism of
[47903] aberrant pre-mRNA splicing in humans. Hum. Molec.

Genet. 6: 909-912,

[47904] 1997.

[47905] 7429.

[47906] Cogan, J. D.; Ramel, B.; Lehto, M.; Phillips, J., III; Prince,

[47907] M.; Blizzard, R. M.; de Ravel, T. J. L.; Brammert, M.; Groop,
L.:

[47908] A recurring dominant negative mutation causes autosomal
dominant growth

[47909] hormone deficiency: a clinical research center study. J. Clin. Endocr.

[47910] Metab. 80: 3591-3595, 1995.

[47911] 7430.

[47912] Colao, A.; Cuocolo, A.; Di Somma, C.; Cerbone, G.; Morte, A. M.

[47913] D.; Nicolai, E.; Lucci, R.; Salvatore, M.; Lombardi, G.: Im-paired

[47914] cardiac performance in elderly patients with growth hor-mone deficiency. J.

[47915] Clin. Endocr. Metab. 84: 3950-3955, 1999.

[47916] 7431.

[47917] Schneider, P. M.; Witzel-Schlomp, K.; Rittner, C.; Zhang, L.:

[47918] The endogenous retroviral insertion in the human com-plement C4 gene

[47919] modulates the expression of homologous genes by anti-sense inhibition. Immunogenetics 53:

[47920] 1-9, 2001.

[47921] 7432.

[47922] Suto, Y.; Tokunaga, K.; Watanabe, Y.; Hirai, M.: Visual

demonstration

[47923] of the organization of the human complement C4 and 21-hydroxylase

[47924] genes by high-resolution fluorescence in situ hybridization. Genomics 33:

[47925] 321-324, 1996.

[47926] 7433.

[47927] Teisberg, P.; Akesson, I.; Olaisen, B.; Gedde-Dahl, T., Jr.; Thorsby,

[47928] E.: Genetic polymorphism of C4 in man and localization of a structural

[47929] C4 locus to the HLA gene complex of chromosome 6. Nature 264: 253-254,

[47930] 1976.

[47931] 7434.

[47932] Boerkoel, C. F.; Takashima, H.; Stankiewicz, P.; Garcia, C. A.;

[47933] Leber, S. M.; Rhee-Morris, L.; Lupski, J. R.: Periaxin mutations

[47934] cause recessive Dejerine-Sottas neuropathy. Am. J. Hum. Genet. 68:

[47935] 325-333, 2001. Note: Erratum: Am. J. Hum. Genet. 68:

557 only, 2001.

[47936] 7435.

[47937] Pischel, K. D.; Marlin, S. D.; Springer, T. A.; Woods, V. L., Jr.;

[47938] Bluestein, H. G.: Polymorphism of lymphocyte function-associated

[47939] antigen-1 demonstrated by a lupus patient's alloantiserum. J. Clin.

[47940] Invest. 79: 1607-1614, 1987.

[47941] 7436.

[47942] Gantz, I.; Miwa, H.; Konda, Y.; Shimoto, Y.; Tashiro, T.; Watson,

[47943] S. J.; DelValle, J.; Yamada, T.: Molecular cloning, expression, and

[47944] gene localization of a fourth melanocortin receptor. J. Biol. Chem. 268:

[47945] 15174-15179, 1993.

[47946] 7437.

[47947] Akey, J. M.; Wang, H.; Xiong, M.; Wu, H.; Liu, W.; Shriver, M.

[47948] D.; Jin, L.: Interaction between the melanocortin-1 recep-

tor and
[47949] P genes contributes to inter-individual variation in skin pigmentation
[47950] phenotypes in a Tibetan population. Hum. Genet. 108: 516-520, 2001.
[47951] 7438.

[47952] Bastiaens, M.; ter Huurne, J.; Gruis, N.; Bergman, W.; Westendorp,
[47953] R.; Vermeer, B.-J.; Bouwes Bavinck, J.-N.: The melanocortin-1-receptor
[47954] gene is the major freckle gene. Hum. Molec. Genet. 10: 1701-1708,
[47955] 2001.

[47956] 7439.

[47957] Chottiner, E. G.; Shewach, D. S.; Datta, N. S.; Ashcraft, E.; Gribbin,
[47958] D.; Ginsburg, D.; Fox, I. H.; Mitchell, B. S.: Cloning and expression
[47959] of human deoxycytidine kinase cDNA. Proc. Nat. Acad. Sci. 88: 1531-1535,
[47960] 1991.

[47961] 7440.

[47962] Huang, S.-H.; Tomich, J. M.; Wu, H.; Jong, A.; Holcenberg, J.:
[47963] Human deoxycytidine kinase: sequence of cDNA clones and analysis of
[47964] expression in cell lines with and without enzyme activity. J. Biol.
[47965] Chem. 264: 14762-14768, 1989. Note: Correction: J. Biol. Chem. 266:
[47966] 5353 only, 1991.

[47967] 7441.

[47968] Johansson, M.; Brismar, S.; Karlsson, A.: Human deoxycy-tidine
[47969] kinase is located in the cell nucleus. Proc. Nat. Acad. Sci. 94:
[47970] 11941-11945, 1997.

[47971] 7442.

[47972] Song, J. J.; Walker, S.; Chen, E.; Johnson, E. E., II; Spychala,
[47973] J.; Gribbin, T.; Mitchell, B. S.: Genomic structure and chro-mosomal
[47974] localization of the human deoxycytidine kinase gene. Proc. Nat. Acad.

[47975] Sci. 90: 431-434, 1993.

[47976] 7443.

[47977] Nurnberg, P.; Thiele, H.; Chandler, D.; Hohne, W.; Cunningham,

[47978] M. L.; Ritter, H.; Leschik, G.; Uhlmann, K.; Mischung, C.; Harrop,

[47979] K.; Goldblatt, J.; Borochowitz, Z. U.; Kotzot, D.; Westermann, F.;

[47980] Mundlos, S.; Braun, H.-S.; Laing, N.; Tinschert, S.: Heterozygous

[47981] mutations in ANKH, the human ortholog of the mouse progressive ankylosis

[47982] gene, result in craniometaphyseal dysplasia. Nature Genet. 28: 37-41,

[47983] 2001.

[47984] 7444.

[47985] Nurnberg, P.; Tinschert, S.; Mrug, M.; Hampe, J.; Muller, C. R.;

[47986] Fuhrmann, E.; Braun, H.-S.; Reis, A.: The gene for autosomal dominant

[47987] craniometaphyseal dysplasia maps to chromosome 5q and is distinct

[47988] from the growth hormone-receptor gene. Am. J. Hum. Genet. 61: 918-923,
[47989] 1997.

[47990] 7445.

[47991] Soriano, P.; Montgomery, C.; Geske, R.; Bradley, A.: Targeted
[47992] disruption of the c-src proto-oncogene leads to osteopetrosis in mice. Cell 64:
[47993] 693-702, 1991.

[47994] 7446.

[47995] Zhang, S.; Xu, L.; Lee, J.; Xu, T.: Drosophila Atrophin homolog
[47996] functions as a transcriptional corepressor in multiple developmental
[47997] processes. Cell 108: 45-56, 2002.

[47998] 7447.

[47999] Stegmann, A. P. A.; Honders, M. W.; Bolk, M. W. J.; Wessels, J.;
[48000] Willemze, R.; Landegent, J. E.: Assignment of the human deoxycytidine
[48001] kinase (DCK) gene to chromosome 4 band q13.3-q21.1.

Genomics 17:
[48002] 528-529, 1993.

[48003] 7448.

[48004] Sanchez-Madrid, F.; Nagy, J.; Robbins, E.; Simon, P.; Springer,
[48005] T. A.: A human leukocyte differentiation antigen family with distinct
[48006] alpha subunits and a common beta subunit: the lympho-cyte function-associated
[48007] antigen (LFA-1), the C3bi complement receptor (OKM1/Mac-1), and the
[48008] p150,95 molecule. J. Exp. Med. 158: 1785-1803, 1983.

[48009] 7449.

[48010] Saint-Ruf, C.; Panigada, M.; Azogui, O.; Debey, P.; von Boehmer,
[48011] H.; Grassi, F.: Different initiation of pre-TCR and gamma-delta-TCR
[48012] signalling. Nature 406: 524-527, 2000.

[48013] 7450.

[48014] Delague, V.; Bareil, C.; Tuffery, S.; Bouvagnet, P.; Chouery, E.;

[48015] Koussa, S.; Maisonobe, T.; Loiselet, J.; Megarbane, A.; Claustres,

[48016] M.: Mapping of a new locus for autosomal recessive demyelinating

[48017] Charcot-Marie-Tooth disease to 19q13.1-13.3 in a large consanguineous

[48018] Lebanese family: exclusion of MAG as a candidate gene. Am. J. Hum.

[48019] Genet. 67: 236-243, 2000.

[48020] 7451.

[48021] Guilbot, A.; Williams, A.; Ravise, N.; Verny, C.; Brice, A.; Sherman,

[48022] D. L.; Brophy, P. J.; LeGuern, E.; Delague, V.; Bareil, C.; Megarbane,

[48023] A.; Claustres, M.: A mutation in periaxin is responsible for CMT4F,

[48024] an autosomal recessive form of Charcot-Marie-Tooth disease. Hum.

[48025] Molec. Genet. 10: 415-421, 2001.

[48026] 7452.

[48027] Roa, B. B.; Dyck, P. J.; Marks, H. G.; Chance, P. F.; Lupski,

[48028] J. R.: Dejerine-Sottas syndrome associated with point mutation in
[48029] the peripheral myelin protein 22 (PMP22) gene. Nature Genet. 5:
[48030] 269-273, 1993.

[48031] 7453.

[48032] Aida, K.; Koishi, S.; Inoue, M.; Nakazato, M.; Tawata, M.; Onaya,
[48033] T.: Familial hypocalciuric hypercalcemia associated with mutation
[48034] in the human Ca(2+)-sensing receptor gene. J. Clin. Endocr. Metab. 80:
[48035] 2594-2598, 1995.

[48036] 7454.

[48037] Bai, M.; Janicic, N.; Trivedi, S.; Quinn, S. J.; Cole, D. E. C.;

[48038] Brown, E. M.; Hendy, G. N.: Markedly reduced activity of mutant calcium-sensing
[48039] receptor with an inserted Alu element from a kindred with familial
[48040] hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. J.

[48041] Clin. Invest. 99: 1917-1925, 1997.

[48042] 7455.

[48043] Bai, M.; Pearce, S. H. S.; Kifor, O.; Trivedi, S.; Stauffer, U.

[48044] G.; Thakker, R. V.; Brown, E. M.; Steinmann, B.: In vivo and in vitro

[48045] characterization of neonatal hyperparathyroidism resulting from a

[48046] de novo, heterozygous mutation in the Ca(2+)-sensing receptor gene:

[48047] Normal maternal calcium homeostasis as a cause of secondary hyperparathyroidism

[48048] in familial benign hypocalciuric hypercalcemia. J. Clin. Invest. 99:

[48049] 88-96, 1997.

[48050] 7456.

[48051] Brown, E. M.; Gamba, G.; Riccardi, D.; Lombardi, M.; Butters, R.;

[48052] Kifor, O.; Sun, A.; Hediger, M.; Lytton, M. A.; Hebert, S. C.: Cloning

[48053] and characterization of an extracellular Ca(2+)-sensing receptor from

[48054] bovine parathyroid. Nature 366: 575-580, 1993.

[48055] 7457.

[48056] Chou, Y.-H. W.; Pollak, M. R.; Brandi, M. L.; Toss, G.; Arn-
qvist,
[48057] H.; Atkinson, A. B.; Papapoulos, S. E.; Marx, S.; Brown, E.
M.; Seidman,
[48058] J. G.; Seidman, C. E.: Mutations in the human
Ca(2+)-sensing-receptor
[48059] gene that cause familial hypocalciuric hypercalcemia. Am.
J. Hum.
[48060] Genet. 56: 1075-1079, 1995.

[48061] 7458.

[48062] Clapham, D. E.: Mutations in G protein-linked receptors:
novel
[48063] insights on disease. Cell 75: 1237-1239, 1993.

[48064] 7459.

[48065] Cole, D. E. C.; Forsythe, C. R.; Dooley, J. M.; Grantmyre, E.

[48066] B.; Salisbury, S. R.: Primary neonatal hyperparathyroidism:
a devastating
[48067] neurodevelopmental disorder if left untreated. J. Craniofa-
cial Genet.

[48068] Dev. Biol. 10: 205-214, 1990.

[48069] 7460.

[48070] Selden, J. R.; Emanuel, B. S.; Wang, E.; Cannizzaro, L.; Palumbo,

[48071] A.; Erikson, J.; Nowell, P. C.; Rovera, G.; Croce, C. M.: Am-

plified

[48072] C-lambda and c-abl genes are on the same marker chro-

mosome in K562

[48073] leukemia cells. Proc. Nat. Acad. Sci. 80: 7289-7292, 1983.

[48074] 7461.

[48075] Mufti, G. J.; Hamblin, T. J.; Oscier, D. G.; Johnson, S.: Common

[48076] ALL with pre-B-cell features showing (8;14) and (14;18) chromosome

[48077] translocations. Blood 62: 1141-1146, 1983.

[48078] 7462.

[48079] Nakayama, K.; Nakayama, K.; Negishi, I.; Kuida, K.; Sawa, H.;

[48080] Loh, D. Y.: Targeted disruption of Bcl-2-alpha-beta in mice: occurrence

[48081] of gray hair, polycystic kidney disease, and lymphocy-

topenia. Proc.
[48082] Nat. Acad. Sci. 91: 3700-3704, 1994.

[48083] 7463.

[48084] Negrini, M.; Silini, E.; Kozak, C.; Tsujimoto, Y.; Croce, C. M.
[48085] : Molecular analysis of mbcl-2: structure and expression of the murine
[48086] gene homologous to the human gene involved in follicular lymphoma. Cell 49:
[48087] 455-463, 1987.

[48088] 7464.

[48089] Ngan, B.-Y.; Chen-Levy, Z.; Weiss, L. M.; Warnke, R. A.; Cleary,
[48090] M. L.: Expression in non-Hodgkin's lymphoma of the BCL-2 protein
[48091] associated with the t(14;18) chromosomal translocation. New Eng.
[48092] J. Med. 318: 1638-1644, 1988.

[48093] 7465.

[48094] Nunez, G.; Hockenbery, D.; McDonnell, T. J.; Sorensen, C. M.;

[48095] Korsmeyer, S. J.: Bcl-2 maintains B cell memory. Nature 353: 71-73,

[48096] 1991.

[48097] 7466.

[48098] Nunez, G.; Seto, M.; Seremetis, S.; Ferrero, D.; Grignani, F.;

[48099] Korsmeyer, S. J.; Dalla-Favera, R.: Growth- and tumor-promoting effects

[48100] of deregulated BCL2 in human B-lymphoblastoid cells. Proc. Nat. Acad.

[48101] Sci. 86: 4589-4593, 1989.

[48102] 7467.

[48103] Pegoraro, L.; Palumbo, A.; Erikson, J.; Falda, M.; Giovanazzo,

[48104] B.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.; Croce, C. M.: A 14;18

[48105] and an 8;14 chromosome translocation in a cell line derived from an

[48106] acute B-cell leukemia. Proc. Nat. Acad. Sci. 81: 7166-7170, 1984.

[48107] 7468.

[48108] Reed, J. C.; Cuddy, M.; Slabiak, T.; Croce, C. M.; Nowell, P.

[48109] C.: Oncogenic potential of bcl-2 demonstrated by gene transfer. Nature 336:

[48110] 259-261, 1988.

[48111] 7469.

[48112] Sagot, Y.; Dubois-Dauphin, M.; Tan, S. A.; de Bilbao, F.; Aebischer,

[48113] P.; Martinou, J.-C.; Kato, A. C.: Bcl-2 overexpression pre-

vents motoneuron

[48114] cell body loss but not axonal degeneration in a mouse model of a neurodegenerative

[48115] disease. J. Neurosci. 15: 7727-7733, 1995.

[48116] 7470.

[48117] Sentman, C. L.; Shutter, J. R.; Hockenbery, D.; Kanagawa, O.;

[48118] Korsmeyer, S. J.: bcl-2 inhibits multiple forms of apopto-

sis but

[48119] not negative selection in thymocytes. Cell 67: 879-888, 1991.

[48120] 7471.

[48121] Silverman, G. A.; Green, E. D.; Young, R. L.; Jockel, J. I.; Domer,

[48122] P. H.; Korsmeyer, S. J.: Meiotic recombination between yeast artificial
[48123] chromosomes yields a single clone containing the entire BCL2 protooncogene. Proc.
[48124] Nat. Acad. Sci. 87: 9913-9917, 1990.

[48125] 7472.

[48126] Strasser, A.; Harris, A. W.; Cory, S.: bcl-2 transgene in-hibits
[48127] T cell death and perturbs thymic self-censorship. Cell 67: 889-899,
[48128] 1991.

[48129] 7473.

[48130] Sugino, N.; Suzuki, T.; Kashida, S.; Karube, A.; Takiguchi, S.;
[48131] Kato, H.: Expression of Bcl-2 and Bax in the human cor-pus luteum
[48132] during the menstrual cycle in early pregnancy: regulation by human
[48133] chorionic gonadotropin. J. Clin. Endocr. Metab. 85: 4379-4386, 2000.
[48134] 7474.

[48135] Tsujimoto, Y.: Overexpression of the human BCL-2 gene product
[48136] results in growth enhancement of Epstein-Barr virus-immortalized B
[48137] cells. Proc. Nat. Acad. Sci. 86: 1958-1962, 1989.

[48138] 7475.

[48139] Tsujimoto, Y.; Bashir, M. M.; Givol, I.; Cossman, J.; Jaffe, E.;
[48140] Croce, C. M.: DNA rearrangements in human follicular lymphoma can
[48141] involve the 5-prime or the 3-prime region of the bcl-2 gene. Proc.
[48142] Nat. Acad. Sci. 84: 1329-1331, 1987.

[48143] 7476.

[48144] Tsujimoto, Y.; Cossman, J.; Jaffe, E.; Croce, C. M.: Involve-ment
[48145] of the bcl-2 gene in human follicular lymphoma. Science 228: 1440-1443,
[48146] 1985.

[48147] 7477.

[48148] Tsujimoto, Y.; Croce, C. M.: Analysis of the structure,

transcripts,
[48149] and protein products of bcl-2, the gene involved in human follicular
[48150] lymphoma. Proc. Nat. Acad. Sci. 83: 5214-5218, 1986.

[48151] 7478.

[48152] Tsujimoto, Y.; Finger, L. R.; Yunis, J.; Nowell, P. C.; Croce,

[48153] C. M.: Cloning of the chromosome breakpoint of neoplastic B cells
[48154] with the t(14;18) chromosome translocation. Science 226: 1097-1099,
[48155] 1985.

[48156] 7479.

[48157] Tsujimoto, Y.; Gorham, J.; Cossman, J.; Jaffe, E.; Croce, C. M.
[48158] : The t(14;18) chromosome translocations involved in B-cell neoplasms
[48159] result from mistakes in VDJ joining. Science 229: 1390-1393, 1985.
[48160] 7480.

[48161] Vanasse, G. J.; Halbrook, J.; Thomas, S.; Burgess, A.; Hoekstra,

[48162] M. F.; Disteche, C. M.; Willerford, D. M.: Genetic pathway to recurrent
[48163] chromosome translocations in murine lymphoma involves V(D)J recombinase. J.
[48164] Clin. Invest. 103: 1669-1675, 1999.

[48165] 7481.

[48166] Vaskivuo, T. E.; Anttonen, M.; Herva, R.; Billig, H.; Dorland,
[48167] M.; Te Velde, E. R.; Stenback, F.; Heikinheimo, M.; Tapanainen, J.
[48168] S.: Survival of human ovarian follicles from fetal to adult life:
[48169] apoptosis, apoptosis-related proteins, and transcription factor GATA-4. J.
[48170] Clin. Endocr. Metab. 86: 3421-3429, 2001.

[48171] 7482.

[48172] Vaux, D. L.; Cory, S.; Adams, J. M.: Bcl-2 gene promotes haemopoietic
[48173] cell survival and cooperates with c-myc to immortalize pre-B cells. Nature 335:
[48174] 440-442, 1988.

[48175] 7483.

[48176] Wang, H.-G.; Rapp, U. R.; Reed, J. C.: Bcl-2 targets the protein
[48177] kinase Raf-1 to mitochondria. Cell 87: 629-638, 1996.

[48178] 7484.

[48179] Williams, G. T.: Programmed cell death: apoptosis and oncogenesis. Cell 65:
[48180] 1097-1098, 1991.

[48181] 7485.

[48182] Yunis, J. J.; Frizzera, G.; Oken, M. M.; McKenna, J.; Theologides,
[48183] A.; Arnesen, M.: Multiple recurrent genomic defects in follicular
[48184] lymphoma: a possible model for cancer. New Eng. J. Med. 316: 79-84,
[48185] 1987.

[48186] 7486.

[48187] Yunis, J. J.; Mayer, M. G.; Arnesen, M. A.; Aeppli, D. P.; Oken,
[48188] M. M.; Frizzera, G.: Bcl-2 and other genomic alterations in the prognosis
[48189] of large-cell lymphoma. New Eng. J. Med. 320:

1047-1054, 1989.

[48190] 7487.

[48191] Cleary, M. L.; Mellentin, J. D.; Spies, J.; Smith, S. D.: Chro-
mosomal

[48192] translocation involving the beta T cell receptor gene in
acute leukemia.

[48193] J. Exp. Med. 167: 682-687, 1988.

[48194] 7488.

[48195] Kuo, S. S.; Mellentin, J. D.; Copeland, N. G.; Gilbert, D. J.;

[48196] Jenkins, N. A.; Cleary, M. L.: Structure, chromosome map-
ping, and

[48197] expression of the mouse Lyl-1 gene. Oncogene 6:
961-968, 1991.

[48198] 7489.

[48199] Inoguchi, K.; Yoshioka, H.; Khaleduzzaman, M.; Ninomiya,
Y.: The

[48200] mRNA for alpha-1(XIX) collagen chain, a new member of
FACITs, contains

[48201] a long unusual 3-prime untranslated region and displays
many unique

[48202] splicing variants. J. Biochem. 117: 137-146, 1995.

[48203] 7490.

[48204] Khaleduzzaman, M.; Sumiyoshi, H.; Ueki, Y.; Inoguchi, K.; Ninomiya,

[48205] Y.; Yoshioka, H.: Structure of the human type XIX collagen (COL19A1)

[48206] gene, which suggests it has arisen from an ancestor gene of the FACIT

[48207] family. Genomics 45: 304-312, 1997.

[48208] 7491.

[48209] Myers, J. C.; Sun, M. J.; D'Ippolito, J. A.; Jabs, E. W.; Neil-son,

[48210] E. G.; Dion, A. S.: Human cDNA clones transcribed from an unusually

[48211] high molecular weight RNA encode a new collagen chain. Gene 123:

[48212] 211-217, 1993.

[48213] 7492.

[48214] Yoshioka, H.; Zhang, H.; Ramirez, F.; Mattei, M.-G.; Moradi-Ameli,

[48215] M.; van der Rest, M.; Gordon, M. K.: Synteny between the loci for

[48216] a novel FACIT-like collagen (D6S228E) and alpha 1(IX) collagen (COL9A1)

[48217] on 6q12-q14 in humans. Genomics 13: 884-886, 1992.

[48218] 7493.

[48219] Gerecke, D. R.; Olson, P. F.; Koch, M.; Knoll, J. H. M.; Taylor,

[48220] R.; Hudson, D. L.; Champliaud, M.-F.; Olsen, B. R.; Burgeson, R. E.

[48221] : Complete primary structure of two splice variants of collagen XII,

[48222] and assignment of alpha-1(XII) collagen (COL12A1), alpha-1(IX) collagen

[48223] (COL9A1), and alpha-1(XIX) collagen (COL19A1) to human chromosome

[48224] 6q12-q13. Genomics 41: 236-242, 1997.

[48225] 7494.

[48226] Olufemi, S.-E.; Green, J. S.; Manickam, P.; Guru, S. C.; Agarwal,

[48227] S. K.; Kester, M. B.; Dong, Q.; Burns, A. L.; Spiegel, A. M.; Marx,

[48228] S. J.; Collins, F. S.; Chandrasekharappa, S. C.: Common ancestral

[48229] mutation in the MEN1 gene is likely responsible for the prolactinoma
[48230] variant of MEN1 (MEN1-Burin) in four kindreds from New-foundland. Hum.
[48231] Mutat. 11: 264-269, 1998.

[48232] 7495.

[48233] Tong, Q.; Dalgin, G.; Xu, H.; Ting, C.-N.; Leiden, J. M.; Ho-tamisligil,
[48234] G. S.: Function of GATA transcription factors in preadipocyte-adipocyte
[48235] transition. Science 290: 134-138, 2000.

[48236] 7496.

[48237] Litt, M.; Buroker, N. E.; Kondoleon, S.; Douglass, J.; Liston,

[48238] D.; Sheehy, R.; Magenis, R. E.: Chromosomal localization of the human
[48239] proenkephalin and prodynorphin genes. Am. J. Hum. Genet. 42: 327-334,
[48240] 1988.

[48241] 7497.

[48242] Litt, M.; Buroker, N. E.; Kondoleon, S. K.; Liston, D.; Dou-glass,

[48243] J.; Sheehy, R.; Magenis, R. E.: Chromosomal localization of the human
[48244] proenkephalin and prodynorphin genes. (Abstract) Cyto-genet. Cell
[48245] Genet. 46: 651 only, 1987.

[48246] 7498.

[48247] Berkovic, S. F.; McIntosh, A.; Howell, R. A.; Mitchell, A.; Sheffield,
[48248] L. J.; Hopper, J. L.: Familial temporal lobe epilepsy: a com-mon disorder
[48249] identified in twins. Ann. Neurol. 40: 227-235, 1996.

[48250] 7499.

[48251] Cendes, F.; Lopes-Cendes, I.; Andermann, E.; Andermann, F.: Familial
[48252] temporal lobe epilepsy: a clinically heterogeneous syn-drome. Neurology 50:
[48253] 554-557, 1998.

[48254] 7500.

[48255] Horikawa, S.; Takai, T.; Toyosato, M.; Takahashi, H.; Noda, M.;
[48256] Kakidani, H.; Kubo, T.; Hirose, T.; Inayama, S.; Hayashida,

H.; Miyata,
[48257] T.; Numa, S.: Isolation and structural organization of the human
[48258] preproenkephalin B gene. Nature 306: 611-614, 1983.

[48259] 7501.

[48260] Grigorenko, E. L.; Wood, F. B.; Meyer, M. S.; Hart, L. A.; Speed,
[48261] W. C.; Shuster, A.; Pauls, D. L.: Susceptibility loci for dis-tinct
[48262] components of developmental dyslexia on chromosomes 6 and 15. Am.
[48263] J. Hum. Genet. 60: 27-39, 1997.

[48264] 7502.

[48265] Lorand, L.; Velasco, P. T.; Rinne, J. R.; Amare, M.; Miller, L.
[48266] K.; Zucker, M. L.: Autoimmune antibody (IgG Kansas) against the fibrin
[48267] stabilizing factor (factor XIII) system. Proc. Nat. Acad. Sci. 85:
[48268] 232-236, 1988.

[48269] 7503.

[48270] Takahashi, K.; Newburger, P. E.; Cohen, H. J.: Glutathione

peroxidase
[48271] protein: absence in selenium deficiency states and correlation with
[48272] enzymatic activity. J. Clin. Invest. 77: 1402-1404, 1986.

[48273] 7504.

[48274] Giuili, G.; Scholl, U.; Bulle, F.; Guellaen, G.: Molecular cloning
[48275] of the cDNAs coding for the two subunits of soluble guanylyl cyclase
[48276] from human brain. FEBS Lett. 304: 83-88, 1992.

[48277] 7505.

[48278] Yan, B.; Heus, J.; Lu, N.; Nichols, R. C.; Raben, N.; Plotz, P.
[48279] H.: Transcriptional regulation of the human acid alpha-glucosidase
[48280] gene: identification of a repressor element and its transcription
[48281] factors Hes-1 and YY1. J. Biol. Chem. 276: 1789-1793, 2001.
[48282] 7506.

[48283] Yan, B.; Raben, N.; Plotz, P. H.: Hes-1, a known transcriptional

[48284] repressor, acts as a transcriptional activator for the human acid

[48285] alpha-glucosidase gene in human fibroblast cells. Biochem. Biophys.

[48286] Res. Commun. 291: 582-587, 2002.

[48287] 7507.

[48288] Pannett, A. A. J.; Thakker, R. V.: Somatic mutations in MEN type

[48289] 1 tumors, consistent with the Knudson 'two-hit' hypothesis. J. Clin.

[48290] Endocr. Metab. 86: 4371-4374, 2001.

[48291] 7508.

[48292] Petty, E. M.; Green, J. S.; Marx, S. J.; Taggart, R. T.; Farid,

[48293] N.; Bale, A. E.: Mapping the gene for hereditary hyperparathyroidism

[48294] and prolactinoma (MEN1-Burin) to chromosome 11q: evidence for a founder

[48295] effect in patients from Newfoundland. Am. J. Hum. Genet. 54: 1060-1066,

[48296] 1994.

[48297] 7509.

[48298] Prezant, T. R.; Levine, J.; Melmed, S.: Molecular characterization
[48299] of the Men1 tumor suppressor gene in sporadic pituitary tumors. J.
[48300] Clin. Endocr. Metab. 83: 1388-1391, 1998.

[48301] 7510.

[48302] Prosser, P. R.; Karam, J. H.; Townsend, J. J.; Frosham, P. H.

[48303] : Prolactin-secreting pituitary adenomas in multiple endocrine adenomatosis,
[48304] type I. Ann. Intern. Med. 91: 41-44, 1979.

[48305] 7511.

[48306] Regan, P. T.; Malagelada, J.-R.: A reappraisal of clinical, roentgenographic,
[48307] and endoscopic features of the Zollinger-Ellison syndrome. Mayo Clin.
[48308] Proc. 53: 19-23, 1978.

[48309] 7512.

[48310] Reimer, D.; Singh, S. M.: A kindred with 5 cases of multiple
[48311] endocrine adenomatosis type I. Hum. Hered. 31: 84-88, 1981.

[48312] 7513.

[48313] Richard, C. W., III; Withers, D. A.; Meeker, T. C.; Maurer, S.;

[48314] Evans, G. A.; Myers, R. M.; Cox, D. R.: A radiation hybrid map of

[48315] the proximal long arm of human chromosome 11 containing the multiple

[48316] endocrine neoplasia type 1 (MEN-1) and bcl-1 disease loci. Am. J.

[48317] Hum. Genet. 49: 1189-1196, 1991.

[48318] 7514.

[48319] Rochelle, J. M.; Watson, M. L.; Oakey, R. J.; Seldin, M. F.:

[48320] A linkage map of mouse chromosome 19:definition of comparative mapping

[48321] relationships with human chromosomes 10 and 11 including the MEN1

[48322] locus. Genomics 14: 26-31, 1992.

[48323] 7515.

[48324] Sato, K.; Obara, T.; Yamazaki, K.; Kanbe, M.; Nakajima, K.; Yamada,

[48325] A.; Yanagisawa, T.; Kato, Y.; Nishikawa, T.; Takano, K.: Somatic

[48326] mutations of the MEN1 gene and microsatellite instability in a case
[48327] of tertiary hyperparathyroidism occurring during high phosphate therapy
[48328] for acquired, hypophosphatemic osteomalacia. J. Clin. Endocr. Metab. 86:
[48329] 5564-5571, 2001.

[48330] 7516.

[48331] Sato, M.; Matsubara, S.; Miyauchi, A.; Ohye, H.; Imachi, H.; Murao,
[48332] K.; Takahara, J.: Identification of five novel germline mutations
[48333] of the MEN1 gene in Japanese multiple endocrine neoplasia type 1 (MEN1)
[48334] families. J. Med. Genet. 35: 915-919, 1998.

[48335] 7517.

[48336] Sawicki, M. P.; Wan, Y.-J. Y.; Johnson, C. L.; Berenson, J.; Gatti,
[48337] R.; Passaro, E., Jr.: Loss of heterozygosity on chromosome 11 in
[48338] sporadic gastrinomas. Hum. Genet. 89: 445-449, 1992.

[48339] 7518.

[48340] Schimke, R. N.: Multiple endocrine adenomatosis syn-
dromes. Adv.
[48341] Intern. Med. 21: 249-265, 1976.

[48342] 7519.

[48343] Schimke, R. N.: Genetic aspects of multiple endocrine
neoplasia. Ann.
[48344] Rev. Med. 35: 25-31, 1984.

[48345] 7520.

[48346] Schimke, R. N.: Multiple endocrine neoplasia: search for
the
[48347] oncogenic trigger. (Editorial) New Eng. J. Med. 314:
1315-1316,
[48348] 1986.

[48349] 7521.

[48350] Schulte, K.-M.; Heinze, M.; Mengel, M.; Simon, D.; Scheur-
ing,
[48351] S.; Kohrer, K.; Roher, H.-D.: MEN I gene mutations in spo-
radic adrenal
[48352] adenomas. Hum. Genet. 105: 603-610, 1999.

[48353] 7522.

[48354] Schussheim, D. H.; Skarulis, M. C.; Agarwal, S. K.; Si-
monds, W.
[48355] F.; Burns, A. L.; Spiegel, A. M.; Marx, S. J.: Multiple en-
docrine
[48356] neoplasia type 1: new clinical and basic findings. Trends
Endocr.
[48357] Metab. 12: 173-178, 2001.

[48358] 7523.

[48359] Skogseid, B.; Larsson, C.; Lindgren, P.-G.; Kvanta, E.; Ras-
tad,
[48360] J.; Theodorsson, E.; Wide, L.; Wilander, E.; Oberg, K.: Clini-
cal
[48361] and genetic features of adrenocortical lesions in multiple
endocrine
[48362] neoplasia type 1. J. Clin. Endocr. Metab. 75: 76-81, 1992.

[48363] 7524.

[48364] Snyder, N.; Scurry, M.; Hughes, W.: Hypergastrinemia in
familial
[48365] multiple endocrine adenomatosis. Ann. Intern. Med. 80:
321-325,

[48366] 1974.

[48367] 7525.

[48368] Snyder, N., III; Scurry, M. T.; Deiss, W. P., Jr.: Five families

[48369] with multiple endocrine adenomatosis. Ann. Intern. Med. 76: 53-58,

[48370] 1972.

[48371] 7526.

[48372] Stacpoole, P. W.; Jaspan, J.; Kasselberg, A. G.; Halter, S. A.;

[48373] Polonsky, K.; Gluck, F. W.; Liljenquist, J. E.; Rabin, D.: A fa-milial

[48374] glucagonoma syndrome: genetic, clinical and biochemical features. Am.

[48375] J. Med. 70: 1017-1026, 1981.

[48376] 7527.

[48377] Stadil, F.; Stage, G.; Rehfeld, J. F.; Efsen, F.; Fischerman,

[48378] K.: Treatment of Zollinger-Ellison syndrome with strepto-zotocin. New

[48379] Eng. J. Med. 294: 1440-1442, 1976.

[48380] 7528.

[48381] Stratakis, C. A.; Schussheim, D. H.; Freedman, S. M.; Keil, M.
[48382] F.; Pack, S. D.; Agarwal, S. K.; Skarulis, M. C.; Weil, R. J.; Lubensky,
[48383] I. A.; Zhuang, Z.; Oldfield, E. H.; Marx, S. J.: Pituitary macroadenoma
[48384] in a 5-year-old: an early expression of multiple endocrine neoplasia
[48385] type 1. J. Clin. Endocr. Metab. 85: 4776-4780, 2000.

[48386] 7529.

[48387] Straus, E.; Johnson, G. F.; Yalow, R. S.: Canine Zollinger-Ellison
[48388] syndrome. Gastroenterology 72: 380-381, 1977.

[48389] 7530.

[48390] Tahara, H.; Imanishi, Y.; Yamada, T.; Tsujimoto, Y.; Tabata, T.;
[48391] Inoue, T.; Inaba, M.; Morii, H.; Nishizawa, Y.: Rare somatic inactivation
[48392] of the multiple endocrine neoplasia type 1 gene in secondary hyperparathyroidism
[48393] of uremia. J. Clin. Endocr. Metab. 85: 4113-4117, 2000.

[48394] 7531.

[48395] Tateishi, R.; Wada, A.; Ishiguro, S.; Ehara, M.; Sakamoto, H.;

[48396] Miki, T.; Mori, Y.; Matsui, Y.; Ishikawa, O.: Coexistence of bilateral

[48397] pheochromocytoma and pancreatic islet cell tumor: report of a case

[48398] and review of the literature. Cancer 42: 2928-2934, 1978.

[48399] 7532.

[48400] Teh, B. T.; Esapa, C. T.; Houlston, R.; Grandell, U.; Farnebo,

[48401] F.; Nordenskjold, M.; Pearce, C. J.; Carmichael, D.; Lars-son, C.;

[48402] Harris, P. E.: A family with isolated hyperparathyroidism segregating

[48403] a missense MEN1 mutation and showing loss of the wild-type alleles

[48404] in the parathyroid tumors. (Letter) Am. J. Hum. Genet. 63: 1554-1549,

[48405] 1998.

[48406]

[48407] 100. Teh, B. T.; Kytola, S.; Farnebo, F.; Bergman, L.; Wong, F. K.;
[48408] Weber, G.; Hayward, N.; Larsson, C.; Skogseid, B.; Beckers, A.; Phelan,
[48409] C.; Edwards, M.; and 29 others: Mutation analysis of the MEN1 gene
[48410] in multiple endocrine neoplasia type 1, familial acromegaly and familial
[48411] isolated hyperparathyroidism. J. Clin. Endocr. Metab. 83: 2621-2626,
[48412] 1998.

[48413]

[48414] 101. Thakker, R. V.: Multiple endocrine neoplasia: syn-dromes of the
[48415] twentieth century. (Editorial) J. Clin. Endocr. Metab. 83: 2617-2620,
[48416] 1998. 1. Aach, R.; Kissane, J. M.: Clinicopathologic confer-ence:
[48417] multiple endocrine adenomatosis. Am. J. Med. 47: 608-618, 1969.

[48418]

[48419] 102. Thakker, R. V.; Bouloux, P.; Wooding, C.; Chotai, K.;

Broad,
[48420] P. M.; Spurr, N. K.; Besser, G. M.; O'Riordan, J. L. H.: Association
[48421] of parathyroid tumors in multiple endocrine neoplasia type I with
[48422] loss of alleles on chromosome 11. New Eng. J. Med. 321: 218-224,
[48423] 1989.

[48424]

[48425] 103. The European Consortium on MEN1: Construction of a 1.2-Mb
[48426] sequence-ready contig of chromosome 11q13 encompassing the multiple
[48427] endocrine neoplasia type 1 (MEN1) gene. Genomics 44: 94-100, 1997.

[48428]

[48429] 104. Underwood, L. E.; Jacobs, N. M.: Familial endocrine adenomatosis. Am.
[48430] J. Dis. Child. 106: 218-223, 1963.

[48431]

[48432] 105. Vance, J. E.; Stoll, R. W.; Kitabchi, A. E.; Buchanan, K.

D.;
[48433] Hollander, D.; Williams, R. H.: Familial nesidioblastosis as the
[48434] predominant manifestation of multiple endocrine adeno-matosis. Am.
[48435] J. Med. 52: 211-227, 1972.

[48436]

[48437] 106. Vance, J. E.; Stoll, R. W.; Kitabchi, A. E.; Williams, R. H.;
[48438] Wood, F. C., Jr.: Nesidioblastosis in familial endocrine adenomatosis. J.A.M.A. 207:
[48439] 1679-1682, 1969.

[48440]

[48441] 107. Verges, B.; Boureille, F.; Goudet, P.; Murat, A.; Beck-ers, A.;
[48442] Sassolas, G.; Cougard, P.; Chambe, B.; Montvernay, C.; Calender, A.;
[48443] Groupe d'Etude des Neoplasies Endocriniennes Multiples: Pituitary
[48444] disease in MEN type 1 (MEN1): data from the France-Bel-gium MEN1 multicenter
[48445] study. J. Clin. Endocr. Metab. 87: 457-465, 2002.

[48446]

[48447] 108. Vortmeyer, A. O.; Boni, R.; Pak, E.; Pack, S.; Zhuang, Z.: Multiple

[48448] endocrine neoplasia 1 gene alterations in MEN1-associated and sporadic

[48449] lipomas. (Letter) J. Nat. Cancer Inst. 90: 398-399, 1998.

[48450]

[48451] 109. Wautot, V.; Vercherat, C.; Lespinasse, J.; Chambe, B.; Lenoir,

[48452] G. M.; Zhang, C. X.; Porchet, N.; Cordier, M.; Beroud, C.; Calender,

[48453] A.: Germline mutation profile of MEN1 in multiple endocrine neoplasia

[48454] type 1: search for correlation between phenotype and the functional

[48455] domains of the MEN1 protein. Hum. Mutat. 20: 35-47, 2002.

[48456]

[48457] 110. Way, L.; Goldman, L.; Dunphy, J. E.: Zollinger-Ellison syndrome:

[48458] an analysis of twenty-five cases. Am. J. Surg. 116: 293-304, 1968.

[48459]

[48460] 111. Wermer, P.: Genetic aspects of adenomatosis of en-
docrine glands. Am.
[48461] J. Med. 16: 363-371, 1954.

[48462]

[48463] 112. Williams, E. D.; Celestin, L. R.: The association of
bronchial
[48464] carcinoid and pluriglandular adenomatosis. Thorax 17:
120-127, 1962.
[48465]

[48466] 113. Wolfe, M. M.; Jensen, R. T.: Zollinger-Ellison syn-
drome: current
[48467] concepts in diagnosis and management. New Eng. J. Med.
317: 1200-1209,
[48468] 1987.

[48469]

[48470] 114. Yu, F.; Venzon, D. J.; Serrano, J.; Goebel, S. U.; Dopp-
man, J.
[48471] L.; Gibril, F.; Jensen, R. T.: Prospective study of the clinical
[48472] course, prognostic factors, causes of death, and survival
in patients

[48473] with long-standing Zollinger-Ellison syndrome. J. Clin. Oncol. 17:
[48474] 615-630, 1999.

[48475]

[48476] 115. Zollinger, R. M.; Ellison, E. H.: Primary peptic ulcerations
[48477] of the jejunum associated with islet cell tumors of the pancreas. Ann.
[48478] Surg. 142: 709-728, 1955.

[48479] 7533.

[48480] Losowsky, M. S.; Miloszewski, K. J. A.: Factor XIII. Brit. J.

[48481] Haemat. 37: 1-5, 1977.

[48482] 7534.

[48483] McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,
[48484] Canada 2/1/1989.

[48485] 7535.

[48486] McDonagh, J.; McDonagh, R. P.; Myllyla, G.; Ikkala, E.: Factor
[48487] XIII deficiency: a genetic study of two affected kindreds in

Finland. Blood 43:
[48488] 327-332, 1974.

[48489] 7536.

[48490] McDonagh, J.; McDonagh, R. P., Jr.; Duckert, F.: Genetic aspects
[48491] of factor XIII deficiency. Ann. Hum. Genet. 35: 197-206, 1971.

[48492] 7537.

[48493] Mikkola, H.; Syrjala, M.; Rasi, V.; Vahtera, E.; Hamalainen, E.;
[48494] Peltonen, L.; Palotie, A.: Deficiency in the A-subunit of coagulation
[48495] factor XIII: two novel point mutations demonstrate different effects
[48496] on transcript levels. Blood 84: 517-525, 1994.

[48497] 7538.

[48498] Mikkola, H.; Yee, V. C.; Syrjala, M.; Seitz, R.; Egbring, R.;

[48499] Petrini, P.; Ljung, R.; Ingerslev, J.; Teller, D. C.; Peltonen, L.;
[48500] Palotie, A.: Four novel mutations in deficiency of coagulation factor

[48501] XIII: consequences to expression and structure of the A subunit. Blood 87:
[48502] 141-151, 1996.

[48503] 7539.

[48504] Nishigaki, T.; Tokunaga, K.; Ishii, C.; Omoto, K.; Akaza, T.;

[48505] Akiyama, N.; Juji, T.; Murata, S.; Naito, S.; Saji, H.; Satoh, H.;
[48506] Shiraki, T.; Tokunaga, K.; Yasuda, N.: No evidence for linkage between
[48507] the loci for coagulation factor XIII-A and HLA. Hum. Genet. 72:
[48508] 266-267, 1986.

[48509] 7540.

[48510] Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg, P.; Thorsby, E.; Siverts,
[48511] A.; Jonassen, R.; Wilhelmy, M. C.: A structural locus for co-agulation
[48512] factor XIIIA (F13A) is located distal to the HLA region on chromosome
[48513] 6p in man. Am. J. Hum. Genet. 37: 215-220, 1985.

[48514] 7541.

[48515] Olaisen, B.; Siverts, A.; Gedde-Dahl, T., Jr.: Linkage data for
[48516] the coagulation factor genes FXIIIA and FXIIIB. (Abstract) Cytogenet.
[48517] Cell Genet. 37: 560 only, 1984.

[48518] 7542.

[48519] Ratnoff, O. D.; Steinberg, A. G.: Inheritance of fibrin-stabilizing-factor
[48520] deficiency. Lancet I: 25-26, 1968.

[48521] 7543.

[48522] Schwartz, M. L.; Pizzo, S. V.; Hill, R. L.; McKee, P. A.: The
[48523] subunit structures of human plasma and platelet factor XIII (fibrin-stabilizing
[48524] factor). J. Biol. Chem. 246: 5851-5854, 1971.

[48525] 7544.

[48526] Sebetan, I. M.: A new method for FXIIIA genetic variants determination
[48527] using isoelectric focusing in 1 M urea: evidence for three structural
[48528] gene loci, FXIIIA1, FXIIIA2, and FXIIIA3. Hum. Genet. 81: 95-97,

[48529] 1988.

[48530] 7545.

[48531] Shmerling, D. H.; Jung, E.; Duckert, F.: Eine neue familiaere

[48532] Koagulopathie infolge Mangels an fibrinstabilisierendem Faktor. Helv.

[48533] Paediat. Acta 15: 471-478, 1960.

[48534] 7546.

[48535] Souri, M.; Yee, V. C.; Kasai, K.; Kaneshiro, T.; Narasaki, K.;

[48536] Castaman, G.; Ichinose, A.: Novel Y283C mutation of the A subunit

[48537] for coagulation factor XIII: molecular modelling predicts its impaired

[48538] protein folding and dimer formation. Brit. J. Haemat. 113: 652-654,

[48539] 2001.

[48540] 7547.

[48541] Standen, G. R.; Bowen, D. J.: Factor XIII A-Bristol 1: detection

[48542] of a nonsense mutation (arg171-to-stop codon) in factor XIII A subunit

[48543] deficiency. Brit. J. Haemat. 85: 769-772, 1993.

[48544] 7548.

[48545] Steinberg, A. G.; Ratnoff, O. D.: Inheritance of factor XIII.

[48546] (Letter) Am. J. Hum. Genet. 22: 597-598, 1970.

[48547] 7549.

[48548] Suzuki, K.; Matsui, K.; Ito, S.; Fujita, K.; Matsumoto, H.: Polymorphism

[48549] of the A subunit of coagulation factor XIII: evidence for subtypes

[48550] of the FXIIIA*I and FXIIIA*2 alleles. Am. J. Hum. Genet. 43: 170-174,

[48551] 1988.

[48552] 7550.

[48553] Takahashi, N.; Takahashi, Y.; Putnam, F. W.: Primary structure

[48554] of blood coagulation factor XIIIa (fibrinoligase, transglu-

[48555] taminase) from human placenta. Proc. Nat. Acad. Sci. 83: 8019-8023, 1986.

[48556] 7551.

[48557] Takahashi, N.; Tsukamoto, H.; Umeyama, H.; Castaman, G.; Rodeghiero,
[48558] F.; Ichinose, A.: Molecular mechanisms of type II factor XIII deficiency:
[48559] novel gly562-to-arg mutation and C-terminal truncation of the A subunit
[48560] cause factor XIII deficiency as characterized in a mammalian expression
[48561] system. Blood 91: 2830-2838, 1998.

[48562] 7552.

[48563] Wong, P.; Komarnicki, L.; Schroeder, M. L.; Lewis, M.; Kaita,
[48564] H.; Philipps, S.; Stranc, L.; McAlpine, P. J.: Analysis for linkage
[48565] between F13A and three chromosome 6 marker loci: evidence for 6pter:F13A:HLA:GLO1:cen
[48566] gene order. Hum. Genet. 79: 228-230, 1988.

[48567] 7553.

[48568] Zahir, M.: Congenital deficiency of fibrin-stabilizing factor:
[48569] report of a case and family study. J.A.M.A. 207: 751-753, 1969.

[48570] 7554.

[48571] Zoghbi, H. Y.; Daiger, S. P.; McCall, A.; O'Brien, W. E.; Beaudet,

[48572] A. L.: Extensive DNA polymorphism at the factor XIIIa (F13A) locus

[48573] and linkage to HLA. Am. J. Hum. Genet. 42: 877-883, 1988.

[48574] 7555.

[48575] Ayyagari, R.; Griesinger, I. B.; Bingham, E.; Lark, K. K.; Mo-

[48576] roi, S. E.; Sieving, P. A.: Autosomal dominant hemorrhagic macular dystrophy

[48577] not associated with the TIMP3 gene. Arch. Ophthal. 118: 85-92, 2000.

[48578] 7556.

[48579] Felbor, U.; Suvanto, E. A.; Forsius, H. R.; Eriksson, A. W.; Weber,

[48580] B. H. F.: Autosomal recessive Sorsby fundus dystrophy re-

[48581] visited: molecular evidence for dominant inheritance. Am. J. Hum. Genet. 60:

[48582] 57-62, 1997.

[48583] 7557.

[48584] Forsius, H. R.; Eriksson, A. W.; Suvanto, E. A.; Alanko, H. I.

[48585] : Pseudoinflammatory fundus dystrophy with autosomal recessive inheritance. Am.

[48586] J. Ophthal. 94: 634-649, 1982.

[48587] 7558.

[48588] Weber, B. H. F.; Vogt, G.; Pruett, R. C.; Stohr, H.; Felbor, U.

[48589] : Mutations in the tissue inhibitor metalloproteinases-3 (TIMP3) in

[48590] patients with Sorsby's fundus dystrophy. Nature Genet. 8: 352-356,

[48591] 1994.

[48592] 7559.

[48593] Blanchette, F.; Day, R.; Dong, W.; Laprise, M.-H.; Dubois, C. M.

[48594] : TGF-beta-1 regulates gene expression of its own converting enzyme

[48595] furin. J. Clin. Invest. 99: 1974-1983, 1997.

[48596] 7560.

[48597] Copeland, N. G.; Gilbert, D. J.; Chretien, M.; Seidah, N. G.;

Jenkins,
[48598] N. A.: Regional localization of three convertases, PC1 (Nec-1), PC2
[48599] (Nec-2), and furin (Fur), on mouse chromosomes. Genomics 13: 1356-1358,
[48600] 1992.

[48601] 7561.

[48602] Dubois, C. M.; Laprise, M.-H.; Blanchette, F.; Gentry, L. E.; Leduc,
[48603] R.: Processing of transforming growth factor beta-1 precursor by
[48604] human furin convertase. J. Biol. Chem. 270: 10618-10624, 1995.
[48605] 7562.

[48606] Hendy, G. N.; Bennett, H. P. J.; Gibbs, B. F.; Lazure, C.; Day,
[48607] R.; Seidah, N. G.: Proparathyroid hormone is preferentially cleaved
[48608] to parathyroid hormone by the prohormone convertase furin: a mass
[48609] spectrometric study. J. Biol. Chem. 270: 9517-9525, 1995.

[48610] 7563.

[48611] Mbikay, M.; Seidah, N. G.; Chretien, M.; Simpson, E. M.: Chromosomal

[48612] assignment of the genes for proprotein convertases PC4, PC5, and PACE

[48613] 4 in mouse and human. Genomics 26: 123-129, 1995.

[48614] 7564.

[48615] Roebroek, A. J. M.; Schalken, J. A.; Leunissen, J. A. M.; On-nekink,

[48616] C.; Bloemers, H. P. J.; Van de Ven, W. J. M.: Evolutionary conserved

[48617] close linkage of the c-fes/fps proto-oncogene and ge-netic sequences

[48618] encoding a receptor-like protein. EMBO J. 5: 2197-2202, 1986.

[48619] 7565.

[48620] Schalken, J. A.; Roebroek, A. J. M.; Oomen, P. P. C. A.; Wa-genaar,

[48621] S. S.; Debruyne, F. M. J.; Bloemers, H. P. J.; Van de Ven, W. J. M.

[48622] : FUR gene expression as a discriminating marker for small cell and

[48623] nonsmall cell lung carcinomas. J. Clin. Invest. 80:
1545-1549, 1987.

[48624] 7566.

[48625] Seidah, N. G.; Mattei, M. G.; Gaspar, L.; Benjannet, S.;
Mbikay,

[48626] M.; Chretien, M.: Chromosomal assignments of the genes
for neuroendocrine

[48627] convertase PC1 (NEC1) to human 5q15-21, neuroen-
docrine convertase

[48628] PC2 (NEC2) to human 20p11.1-11.2, and furin (mouse
7[D1-E2] region). Genomics 11:

[48629] 103-107, 1991.

[48630] 7567.

[48631] Curtis, E. J.; Fraser, F. C.; Warburton, D.: Congenital cleft

[48632] lip and palate. Am. J. Dis. Child. 102: 853-857, 1961.

[48633] 7568.

[48634] Awdeh, Z. L.; Alper, C. A.: Inherited structural polymor-
phism

[48635] of the fourth component of human complement. Proc.
Nat. Acad. Sci. 77:

[48636] 3576-3580, 1980.

[48637] 7569.

[48638] Awdeh, Z. L.; Ochs, H. D.; Alper, C. A.: Genetic analysis of C4

[48639] deficiency. J. Clin. Invest. 67: 260-263, 1981.

[48640] 7570.

[48641] Brade, V.; Hall, R. E.; Colten, H. R.: Biosynthesis of pro-C3,

[48642] a precursor of the third component of complement. J. Exp. Med. 146:

[48643] 759-765, 1977.

[48644] 7571.

[48645] Carroll, M. C.; Campbell, R. D.; Bentley, D. R.; Porter, R. R.

[48646] : A molecular map of the human major histocompatibility complex class

[48647] III region linking complement genes C4, C2 and factor B. Nature 307:

[48648] 237-241, 1984.

[48649] 7572.

[48650] Carroll, M. C.; Porter, R. R.: Cloning of a human comple-ment component

[48651] C4 gene. Proc. Nat. Acad. Sci. 80: 264-267, 1983.

[48652] 7573.

[48653] Chan, A. C.; Mitchell, K. R.; Munns, T. W.; Karp, D. R.; Atkinson,

[48654] J. P.: Identification and partial characterization of the se-

creted

[48655] form of the fourth component of human complement: evi-

dence that it

[48656] is different from major plasma form. Proc. Nat. Acad. Sci. 80: 268-272,

[48657] 1983.

[48658] 7574.

[48659] Cream, J. J.; Olaisen, B.; Teisberg, P.; Soler, A. V.; Thomp-

son,

[48660] R. A.: Genetic basis of acquired C4 deficiency. Clin. Genet. 16:

[48661] 297-300, 1979.

[48662] 7575.

[48663] Cunningham-Rundles, C.; Dupont, B.; Jersild, C.; Tegoli, C.; Whitsett,

[48664] C.; Good, R. A.: Are HLA and Chido related antigenic groups? Transplant.

[48665] Proc. 9: 33-38, 1977.

[48666] 7576.

[48667] Cunningham-Rundles, C.; Tegoli, J.; Dupont, B.; Whitsett, C.; Good,

[48668] R. A.: Chemical studies on the Chido antigen. Transplant. Proc. 9:

[48669] 647-652, 1977.

[48670] 7577.

[48671] Curman, B.; Ostberg, L.; Sandberg, L.; Malmheden-Erikkson, I.;

[48672] Stalenheim, G.; Rask, L.; Peterson, P. A.: H-2 linked Ss protein

[48673] is C-4 component of complement. Nature 258: 243-245, 1975.

[48674] 7578.

[48675] Ellman, L.; Green, I.; Frank, M.: Genetically controlled total

[48676] deficiency of the fourth component of complement in the guinea pig. Science 170:

[48677] 74-75, 1970.

[48678] 7579.

[48679] Fontaine, M.; Daveau, M.; Lebreton, J. P.: A common antigenic
[48680] determinant on human C4b and C3b. Molec. Immun. 17: 1075-1078, 1980.
[48681] 7580.
[48682] Giles, C. M.: A new genetic variant for Chido. Vox Sang. 46:
[48683] 149-156, 1984.
[48684] 7581.
[48685] Hall, R. E.; Colten, H. R.: Genetic defect in biosynthesis of
[48686] the precursor form of the fourth component of complement. Science 199:
[48687] 69-70, 1978.
[48688] 7582.
[48689] Hall, R. E.; Colten, H. R.: Cell-free synthesis of the fourth
[48690] component of guinea pig complement (C4): identification of a precursor
[48691] of serum C4 (pro-C4). Proc. Nat. Acad. Sci. 74: 1707-1710, 1977.
[48692] 7583.

[48693] Harris, J. P.; Tegoli, J.; Swanson, J.; Fisher, N.; Gavin, J.;

[48694] Noades, J.: A nebulous antibody responsible for cross-matching difficulties

[48695] (Chido). Vox Sang. 12: 140-142, 1967.

[48696] 7584.

[48697] Hobart, M. J.; Lachmann, P. J.: Allotypes of complement components

[48698] in man. Transplant. Rev. 32: 26-42, 1976.

[48699] 7585.

[48700] Kramer, J.; Fulop, T.; Rajczy, K.; Ahn Tuan, N.; Fust, G.: A

[48701] marked drop in the incidence of the null allele of the B gene of the

[48702] fourth component of complement (C4B*Q0) in elderly subjects: C4B*Q0

[48703] as a probable negative selection factor for survival. Hum. Genet. 86:

[48704] 595-598, 1991.

[48705] 7586.

[48706] Kronke, M.; Geezy, A. F.; Hadding, U.; Bitter-Suermann, D.: Linkage

[48707] of C4 and C4 deficiency to Bf and GPLA. Immunogenetics 5: 461-466,

[48708] 1977.

[48709] 7587.

[48710] Lachmann, P. J.; Grennan, D.; Martin, A.; Demant, P.: Identification

[48711] of Ss protein as murine C4. Nature 258: 242-243, 1975.

[48712] 7588.

[48713] Lhotta, K.; Schlogl, A.; Uring-Lambert, B.; Kronenberg, F.; Konig,

[48714] P.: Complement C4 phenotypes in patients with end-stage renal disease. Nephron 72:

[48715] 442-446, 1996.

[48716] 7589.

[48717] Mascart-Lemone, F.; Hauptmann, G.; Goetz, J.; Duchateau, J.; Delespesse,

[48718] G.; Vray, B.; Dab, I.: Genetic deficiency of C4 presenting with recurrent

[48719] infections and a SLE-like disease: genetic and immunologic studies. Am.

[48720] J. Med. 75: 295-304, 1983.

[48721] 7590.

[48722] Meo, T.; Krasteff, T.; Shreffler, D. C.: Immunochemical characterization

[48723] of murine H-2 controlled Ss (serum substance) protein through identification

[48724] of its human homologue as the fourth component of complement. Proc.

[48725] Nat. Acad. Sci. 72: 4536-4540, 1975.

[48726] 7591.

[48727] Middleton, J.; Crookston, M. C.: Chido-substance in plasma. Vox

[48728] Sang. 23: 256-261, 1972.

[48729] 7592.

[48730] Middleton, J.; Crookston, M. C.; Falk, J. A.; Robson, E. B.; Cook,

[48731] P. J. L.; Batchelor, J. R.; Bodmer, J.; Ferrara, G. B.; Festen-stein,

[48732] J.; Harris, H.; Kissmeyer-Nielsen, F.; Lawler, S. D.; Sachs, J. A.;

[48733] Wolf, E.: Linkage of Chido and HL-A. Tissue Antigens 4: 366-373,

[48734] 1974.

[48735] 7593.

[48736] O'Neill, G. J.: The genetic control of Chido and Rodgers blood

[48737] group substances. Seminars Hemat. 18: 32-38, 1981.

[48738] 7594.

[48739] O'Neill, G. J.; Yang, S. Y.; Dupont, B.: Two HLA-linked loci

[48740] controlling the fourth component of human complement. Proc. Nat.

[48741] Acad. Sci. 75: 5165-5169, 1978.

[48742] 7595.

[48743] O'Neill, G. J.; Yang, S. Y.; Dupont, B.: Chido and Rodgers blood

[48744] groups: relationships to C4 and HLA. Transplant. Proc. 10: 749-751,

[48745] 1978.

[48746] 7596.

[48747] O'Neill, G. J.; Yang, S. Y.; Tegoli, J.; Berger, R.; Dupont, B.

[48748] : Chido and Rodgers blood groups are distinct antigenic

components

[48749] of human C4. Nature 273: 668-670, 1978.

[48750] 7597.

[48751] Ochs, H. D.; Rosenfeld, S. I.; Thomas, E. D.; Giblett, E. R.;

[48752] Alper, C. A.; Dupont, B.; Schaller, J. G.; Gilliland, B. C.; Hansen,

[48753] J. A.; Wedgwood, R. J.: Linkage between the gene (or genes) controlling

[48754] synthesis of the fourth component of complement and the major histocompatibility

[48755] complex. New Eng. J. Med. 296: 470-475, 1977.

[48756] 7598.

[48757] Olaisen, B.; Teisberg, P.; Nordhagen, R.; Michaelsen, T.; Gedde-Dahl,

[48758] T., Jr.: Human complement C4 locus is duplicated on some chromosomes. Nature 279:

[48759] 736-737, 1979.

[48760] 7599.

[48761] Partanen, J.; Kere, J.; Wessberg, S.; Koskimies, S.: Determi-nation

[48762] of deletion sizes in the MHC-linked complement C4 and

steroid 21-hydroxylase

[48763] genes by pulsed-field gel electrophoresis. Genomics 5: 345-349,

[48764] 1989.

[48765] 7600.

[48766] Partanen, J.; Koskimies, S.; Johansson, E.: C4 null phenotypes

[48767] among lupus erythematosus patients are predominantly the result of

[48768] deletions covering C4 and closely linked 21-hydroxylase A genes. J.

[48769] Med. Genet. 25: 387-391, 1988.

[48770] 7601.

[48771] Petersen, G. B.; Sorensen, I. J.; Buskjaer, L.; Lamm, L. U.:

[48772] Genetic studies of complement C4 in man. Hum. Genet. 53: 31-36,

[48773] 1979.

[48774] 7602.

[48775] Davidson, D.: The function and evolution of Msx genes: pointers

[48776] and paradoxes. Trends Genet. 11: 405-411, 1995.

[48777] 7603.

[48778] Jabs, E. W.; Muller, U.; Li, X.; Ma, L.; Luo, W.; Haworth, I. S.;

[48779] Klisak, I.; Sparkes, R.; Warman, M. L.; Mulliken, J. B.;
Snead, M.

[48780] L.; Maxson, R.: A mutation in the homeodomain of the
human MSX2 gene

[48781] in a family affected with autosomal dominant craniosyn-
ostosis. Cell 75:

[48782] 443-450, 1993.

[48783] 7604.

[48784] Li, X.; Ma, L.; Snead, M.; Haworth, I.; Sparkes, R.; Jackson,
C.;

[48785] Warman, M.; Mulliken, J.; Maxson, R.; Muller, U.; Jabs, E.: A
mutation

[48786] in the homeodomain of the MSX2 gene in a family affected
with craniosynostosis,

[48787] Boston type. (Abstract) Am. J. Hum. Genet. 53 (suppl.):
A213 only,

[48788] 1993.

[48789] 7605.

[48790] Liu, Y. H.; Kundu, R.; Wu, L.; Luo, W.; Ignelzi, M. A., Jr.;

Snead,
[48791] M. L.; Maxson, R. E., Jr.: Premature suture closure and ectopic cranial
[48792] bone in mice expressing Msx2 transgenes in the developing skull. Proc.
[48793] Nat. Acad. Sci. 92: 6137-6141, 1995.

[48794] 7606.

[48795] Ma, L.; Golden, S.; Wu, L.; Maxson, R.: The molecular basis of
[48796] Boston-type craniosynostosis: the pro148-to-his mutation in the N-terminal
[48797] arm of the MSX2 homeodomain stabilizes DNA binding without altering
[48798] nucleotide sequence preferences. Hum. Molec. Genet. 5: 1915-1920,
[48799] 1996.

[48800] 7607.

[48801] Muller, U.; Warman, M. L.; Mulliken, J. B.; Weber, J. L.: Assignment
[48802] of a gene locus involved in craniosynostosis to chromosome 5qter. Hum.
[48803] Molec. Genet. 2: 119-122, 1993.

[48804] 7608.

[48805] Satokata, I.; Ma, L.; Ohshima, H.; Bei, M.; Woo, I.; Nishizawa,

[48806] K.; Maeda, T.; Takano, Y.; Uchiyama, M.; Heaney, S.; Peters, H.; Tang,

[48807] Z.; Maxson, R.; Maas, R.: Msx2 deficiency in mice causes pleiotropic

[48808] defects in bone growth and ectodermal organ formation. Nature Genet. 24:

[48809] 391-395, 2000.

[48810] 7609.

[48811] Takahashi, C.; Akiyama, N.; Matsuzaki, T.; Takai, S.; Kitayama,

[48812] H.; Noda, M.: Characterization of a human MSX-2 cDNA and its fragment

[48813] isolated as a transformation suppressor gene against v-Ki-ras oncogene. Oncogene 12:

[48814] 2137-2146, 1996.

[48815] 7610.

[48816] Warman, M. L.; Mulliken, J. B.; Hayward, P. G.; Muller, U.: Newly

[48817] recognized autosomal dominant disorder with craniosyn-
ostosis. Am.
[48818] J. Med. Genet. 46: 444-449, 1993.
[48819] 7611.
[48820] Wilkie, A. O. M.; Tang, Z.; Elanko, N.; Walsh, S.; Twigg, S.
R.
[48821] F.; Hurst, J. A.; Wall, S. A.; Chrzanowska, K. H.; Maxson, R.
E.,
[48822] Jr.: Functional haploinsufficiency of the human homeobox
gene MSX2
[48823] causes defects in skull ossification. Nature Genet. 24:
387-390,
[48824] 2000.
[48825] 7612.
[48826] Rius, C.; Smith, J. D.; Almendro, N.; Langa, C.; Botella, L.
M.;
[48827] Marchuk, D. A.; Vary, C. P. H.; Bernabeu, C.: Cloning of the
promoter
[48828] region of human endoglin, the target gene for hereditary
hemorrhagic
[48829] telangiectasia type 1. Blood 92: 4677-4690, 1998.

[48830] 7613.

[48831] Shovlin, C. L.; Hughes, J. M. B.; Scott, J.; Seidman, C. E.; Seidman,

[48832] J. G.: Characterization of endoglin and identification of novel mutations

[48833] in hereditary hemorrhagic telangiectasia. Am. J. Hum. Genet. 61:

[48834] 68-79, 1997.

[48835] 7614.

[48836] Kanai, Y.; Hediger, M. A.: Primary structure and functional characterization

[48837] of a high-affinity glutamate transporter. Nature 360: 467-471, 1992.

[48838] 7615.

[48839] Lin, C. G.; Orlov, I.; Ruggiero, A. M.; Dykes-Hoberg, M.; Lee,

[48840] A.; Jackson, M.; Rothstein, J. D.: Modulation of the neu-ronal glutamate

[48841] transporter EAAC1 by the interacting protein GTRAP3-18. Nature 410:

[48842] 84-88, 2001.

[48843] 7616.

[48844] Smith, C. P.; Weremowicz, S.; Kanai, Y.; Stelzner, M.; Mor-
ton,
[48845] C. C.; Hediger, M. A.: Assignment of the gene coding for
the human
[48846] high-affinity glutamate transporter EAAC1 to 9p24: po-
tential role
[48847] in dicarboxylic aminoaciduria and neurodegenerative dis-
orders. Genomics 20:
[48848] 335-336, 1994.

[48849] 7617.

[48850] Ahn, J.; Ludecke, H.-J.; Lindow, S.; Horton, W. A.; Lee, B.;
Wagner,
[48851] M. J.; Horsthemke, B.; Wells, D. E.: Cloning of the putative
tumour
[48852] suppressor gene for hereditary multiple exostoses (EXT1).
Nature
[48853] Genet. 11: 137-143, 1995.

[48854] 7618.

[48855] Bakker, E.; Smoor, M. A.; Maat, A.; van Haeringen, A.; van
de Kamp,

[48856] J. J. P.; van Ommen, G. J. B.: Hereditary multiple exostoses in two

[48857] Dutch families, LGS region excluded. (Abstract) Human Genome Mapping

[48858] Workshop 93 15, 1993.

[48859] 7619.

[48860] Beals, R.; Weber, C.; Wright, A.; Rowe, S.; Lovrien, E. W.: Multiple

[48861] exostosis may be loosely linked to ABO. (Abstract) Cytogenet. Cell

[48862] Genet. 37: 416-417, 1984.

[48863] 7620.

[48864] Beighton, P.; Sujansky, E.; Patzak, B.; Portele, K. A.: Genetic

[48865] skeletal dysplasias in the Museum of Pathological Anatomy, Vienna. Am.

[48866] J. Med. Genet. 47: 843-847, 1993.

[48867] 7621.

[48868] Blanton, S. H.; Hogue, D.; Wagner, M.; Wells, D.; Young, I. D.;

[48869] Hecht, J. T.: Hereditary multiple exostoses: confirmation

of linkage
[48870] to chromosomes 8 and 11. Am. J. Med. Genet. 62:
150-159, 1996.
[48871] 7622.

[48872] Bovee, J. V. M. G.; Cleton-Jansen, A.-M.; Wuyts, W.;
Caethoven,
[48873] G.; Taminiau, A. H. M.; Bakker, E.; Van Hul, W.; Cornelisse,
C. J.;
[48874] Hogendoorn, P. C. W.: EXT-mutation analysis and loss of
heterozygosity
[48875] in sporadic and hereditary osteochondromas and sec-
ondary chondrosarcomas. Am.
[48876] J. Hum. Genet. 65: 689-698, 1999.

[48877] 7623.

[48878] Brooks, A. P.; Wynne-Davies, R.: A family with diaphyseal
aclasis
[48879] and peripheral dysostosis. J. Med. Genet. 17: 277-280,
1980.
[48880] 7624.

[48881] Buhler, E. M.; Malik, N. J.: The tricho-rhino-phalangeal
syndrome(s):
[48882] chromosome 8 long arm deletion: is there a shortest re-

gion of overlap
[48883] between reported cases? TRP I and TRP II syndromes: are they separate
[48884] entities? (Editorial) Am. J. Med. Genet. 19: 113-119, 1984.

[48885] 7625.

[48886] Buur, T.; Morch, M. M.: Hereditary multiple exostoses with spinal
[48887] cord compression. (Letter) J. Neurol. Neurosurg. Psychiat. 46: 96-98,
[48888] 1983.

[48889] 7626.

[48890] Cheung, P. K.; McCormick, C.; Crawford, B. E.; Esko, J. D.; Tufaro,
[48891] F.; Duncan, G.: Etiological point mutations in the heredi-tary multiple
[48892] exostoses gene EXT1: a functional analysis of heparan sulfate polymerase
[48893] activity. Am. J. Hum. Genet. 69: 55-66, 2001.

[48894] 7627.

[48895] Cook, A.; Raskind, W.; Blanton, S. H.; Pauli, R. M.; Gregg, R.

[48896] G.; Francomano, C. A.; Puffenberger, E.; Conrad, E. U.; Schmale, G.;

[48897] Schellenberg, G.; Wijsman, E.; Hecht, J. T.; Wells, D.; Wagner, M.

[48898] J.: Genetic heterogeneity in families with hereditary multiple exostoses. Am.

[48899] J. Hum. Genet. 53: 71-79, 1993.

[48900] 7628.

[48901] Del-Rio, R.; Navarra, E.; Ferrando, J.; Mascaro, J. M.: Multiple

[48902] exostoses syndrome presenting as nail malalignment and longitudinal

[48903] dystrophy of fingers. (Letter) Arch. Derm. 128: 1655-1656, 1992.

[48904] 7629.

[48905] Francannet, C.; Cohen-Tanugi, A.; Le Merrer, M.; Munnich, A.;

[48906] Bonaventure, J.; Legeai-Mallet, L.: Genotype-phenotype correlation

[48907] in hereditary multiple exostoses. J. Med. Genet. 38: 430-434, 2001.

[48908] 7630.

[48909] Gardner, E. J.; Shupe, J. L.; Leone, N. C.; Olson, A. E.: Hereditary
[48910] multiple exostosis: a comparative genetic evaluation in man and horses. J.
[48911] Hered. 66: 318-322, 1975.

[48912] 7631.

[48913] Giedion, A.: Personal Communication. Zurich, Switzerland 1972.
[48914] 7632.

[48915] Hall, C. R.; Cole, W. G.; Haynes, R.; Hecht, J. T.: Reevaluation
[48916] of a genetic model for the development of exostosis in hereditary
[48917] multiple exostosis. Am. J. Med. Genet. 112: 1-5, 2002.

[48918] 7633.

[48919] Hall, J. G.; Wilson, R. D.; Kalousek, D.; Beauchamp, R.: Familial
[48920] multiple exostoses--no chromosome 8 deletion observed. Am. J. Med.
[48921] Genet. 22: 639-640, 1985.

[48922] 7634.

[48923] Hamers, A.; Peeters, G.; Jongbloet, P.; Geraedts, J.: Micro-cytogenetics

[48924] of chromosome 8q. (Abstract) 8th Int. Chrom. Conf., Lubeck Abstract-8,

[48925] 9/1983.

[48926] 7635.

[48927] Hecht, J. T.; Hogue, D.; Strong, L. C.; Hansen, M. F.; Blan-ton,

[48928] S. H.; Wagner, M.: Hereditary multiple exostosis and chondrosarcoma:

[48929] linkage to chromosome 11 and loss of heterozygosity for EXT-linked

[48930] markers on chromosomes 11 and 8. Am. J. Hum. Genet. 56: 1125-1131,

[48931] 1995.

[48932] 7636.

[48933] Hecht, J. T.; Hogue, D.; Wang, Y.; Blanton, S. H.; Wagner, M.;

[48934] Strong, L. C.; Raskind, W.; Hansen, M. F.; Wells, D.: Hereditary

[48935] multiple exostoses (EXT): mutational studies of familial EXT1 cases

[48936] and EXT-associated malignancies. Am. J. Hum. Genet. 60: 80-86, 1997.

[48937] 7637.

[48938] Lacey, S. W.; Sanders, J. M.; Rothberg, K. G.; Anderson, R. G.

[48939] W.; Kamen, B. A.: Complementary DNA for the folate binding protein

[48940] correctly predicts anchoring to the membrane by glycosyl-

phosphatidylinositol. J.

[48941] Clin. Invest. 84: 715-720, 1989.

[48942] 7638.

[48943] Istvan, E. S.; Deisenhofer, J.: Structural mechanism for statin

[48944] inhibition of HMG-CoA reductase. Science 292: 1160-1164, 2001.

[48945] 7639.

[48946] Piedrahita, J. A.; Oetama, B.; Bennett, G. D.; van Waes, J.; Kamen,

[48947] B. A.; Richardson, J.; Lacey, S. W.; Anderson, R. G. W.; Finnell,

[48948] R. H.: Mice lacking the folic acid-binding protein Folbp1 are defective

[48949] in early embryonic development. Nature Genet. 23: 228-232, 1999.

[48950] 7640.

[48951] Shaw, G. M.; Jensvold, N. G.; Wasserman, C. R.; Lammer, E. J.

[48952] : Epidemiologic characteristics of phenotypically distinct neural

[48953] tube defects among 0.7 million California births,

1983-1987. Teratology 49:
[48954] 143-149, 1994.

[48955] 7641.

[48956] Werler, M. M.; Shapiro, S.; Mitchell, A. A.: Periconceptional

[48957] folic acid exposure and risk of occurrent neural tube de-
fects. J.A.M.A. 269:
[48958] 1257-1261, 1993.

[48959] 7642.

[48960] Sipila, K.; Aula, P.: Database for the mutations of the
Finnish
[48961] disease heritage. Hum. Mutat. 19: 16-22, 2002.

[48962] 7643.

[48963] Rimokh, R.; Gadoux, M.; Bertheas, M.-F.; Berger, F.;
Garoscio,
[48964] M.; Deleage, G.; Germain, D.; Magaud, J.-P.: FVT-1, a
novel human
[48965] transcription unit affected by variant translocation
t(2;18)(p11;q21)
[48966] of follicular lymphoma. Blood 81: 136-142, 1993.

[48967] 7644.

[48968] Tortoriello, D. V.; Sidis, Y.; Holtzman, D. A.; Holmes, W. E.;

[48969] Schneyer, A. L.: Human follistatin-related protein: a structural

[48970] homologue of follistatin with nuclear localization. Endocrinology 142:

[48971] 3426-3434, 2001.

[48972] 7645.

[48973] Lee, H.-K.; Barbarosie, M.; Kameyama, K.; Bear, M. F.; Huganir,

[48974] R. L.: Regulation of distinct AMPA receptor phosphorylation sites

[48975] during bidirectional synaptic plasticity. Nature 405: 955-959, 2000.

[48976] 7646.

[48977] Mack, V.; Burnashev, N.; Kaiser, K. M. M.; Rozov, A.; Jensen, V.;

[48978] Hvalby, O.; Seeburg, P. H.; Sakmann, B.; Sprengel, R.: Conditional

[48979] restoration of hippocampal synaptic potentiation in GluR-A-deficient

[48980] mice. Science 292: 2501-2504, 2001.

[48981] 7647.

[48982] Puckett, C.; Gomez, C. M.; Korenberg, J. R.; Tung, H.; Meier, T.

[48983] J.; Chen, X. N.; Hood, L.: Molecular cloning and chromosomal localization

[48984] of one of the human glutamate receptor genes. Proc. Nat. Acad. Sci. 88:

[48985] 7557-7561, 1991.

[48986] 7648.

[48987] Shi, S.-H.; Hayashi, Y.; Petralla, R. S.; Zaman, S. H.; Wenthold,

[48988] R. J.; Svoboda, K.; Malinow, R.: Rapid spine delivery and redistribution

[48989] of AMPA receptors after synaptic NMDA receptor activation. Science 284:

[48990] 1811-1816, 1999.

[48991] 7649.

[48992] Zamanillo, D.; Sprengel, R.; Hvalby, O.; Jensen, V.; Burnashev,

[48993] N.; Rozov, A.; Kaiser, K. M. M.; Koster, H. J.; Borchardt, T.; Worley,

[48994] P.; Lubke, J.; Frotscher, M.; Kelly, P. H.; Sommer, B.; Andersen,
[48995] P.; Seeburg, P. H.; Sakmann, B.: Importance of AMPA receptors for
[48996] hippocampal synaptic plasticity but not for spatial learning. Science 284:
[48997] 1805-1811, 1999.

[48998] 7650.

[48999] Hardingham, G. E.; Fukunaga, Y.; Bading, H.: Extrasynaptic NMDARs
[49000] oppose synaptic NMDARs by triggering CREB shut-off and cell death
[49001] pathways. Nature Neurosci. 5: 405-414, 2002.

[49002] 7651.

[49003] Takano, H.; Onodera, O.; Tanaka, H.; Mori, H.; Sakimura, K.; Hori,
[49004] T.; Kobayashi, H.; Mishina, M.; Tsuji, S.: Chromosomal localization
[49005] of the epsilon-1, epsilon-3, and zeta-1 subunit genes of the human
[49006] NMDA receptor channel. Biochem. Biophys. Res. Commun. 197: 922-926,

[49007] 1993.

[49008] 7652.

[49009] Aral, B.; Schlenzig, J.-S.; Liu, G.; Kamoun, P.: Database cloning

[49010] human delta-1-pyrroline-5-carboxylate synthetase (P5CS) cDNA: a bifunctional

[49011] enzyme catalyzing the first two steps in proline biosyn- thesis. C.

[49012] R. Acad. Sci. III 319: 171-178, 1996.

[49013] 7653.

[49014] Baumgartner, M. R.; Hu, C. A.; Almashanu, S.; Steel, G.; Obie,

[49015] C.; Aral, B.; Rabier, D.; Kamoun, P.; Saudubray, J.-M.; Valle, D.

[49016] : Hyperammonemia with reduced ornithine, citrulline, arginine and

[49017] proline: a new inborn error caused by a mutation in the gene encoding

[49018] delta-1-pyrroline-5-carboxylate synthase. Hum. Molec. Genet. 9:

[49019] 2853-2858, 2000.

[49020] 7654.

[49021] Hu, C. A.; Lin, W.-W.; Obie, C.; Valle, D.: Molecular enzymology

[49022] of mammalian delta-1-pyrroline-5-carboxylate synthase. J. Biol. Chem.

[49023] 274: 6754-6762, 1999.

[49024] 7655.

[49025] Meera Khan, P.; Doppert, B. A.; Hagemeijer, A.; Westerveld, A.

[49026] : The human loci for phosphopyruvate hydratase and guanylate kinase

[49027] are syntenic with the PGD-PGM1 linkage group in man-Chinese hamster

[49028] somatic cell hybrids. Cytogenet. Cell Genet. 13: 130-131, 1974.

[49029] 7656.

[49030] Berkovitz, G. D.; Guerami, A.; Brown, T. R.; MacDonald, P. C.;

[49031] Migeon, C. J.: Familial gynecomastia with increased extraglandular

[49032] aromatization of plasma carbon(19)-steroids. J. Clin. Invest. 75:

[49033] 1763-1769, 1985.

[49034] 7657.

[49035] Itoh, H.; Toyama, R.; Kozasa, T.; Tsukamoto, T.; Matsuoka, M.;

[49036] Kaziro, Y.: Presence of three distinct molecular species of G(i)

[49037] protein alpha subunit: structure of rat cDNAs and human genomic DNAs. J.

[49038] Biol. Chem. 263: 6656-6664, 1988.

[49039] 7658.

[49040] Jiang, M.; Gold, M. S.; Boulay, G.; Spicher, K.; Peyton, M.; Brabet,

[49041] P.; Srinivasan, Y.; Rudolph, U.; Ellison, G.; Birnbaumer, L.: Multiple

[49042] neurological abnormalities in mice deficient in the G protein G(o). Proc.

[49043] Nat. Acad. Sci. 95: 3269-3274, 1998.

[49044] 7659.

[49045] Kroll, S. D.; Chen, J.; De Vivo, M.; Carty, D. J.; Buku, A.; Premont,

[49046] R. T.; Iyengar, R.: The Q205LGo-alpha subunit expressed

in NIH-3T3
[49047] cells induces transformation. J. Biol. Chem. 267:
23183-23188, 1992.
[49048] 7660.

[49049] Murtagh, J. J., Jr.; Eddy, R.; Shows, T. B.; Moss, J.; Vaughan,

[49050] M.: Different forms of Go alpha mRNA arise by alternative
splicing
[49051] of transcripts from a single gene on human chromosome
16. Molec.
[49052] Cell. Biol. 11: 1146-1155, 1991.

[49053] 7661.

[49054] Ram, P. T.; Horvath, C. M.; Iyengar, R.: Stat3-mediated
transformation
[49055] of NIH-3T3 cells by the constitutively active Q205L G-
alpha(O) protein. Science 287:
[49056] 142-144, 2000.

[49057] 7662.

[49058] Strathmann, M.; Wilkie, T. M.; Simon, M. I.: Alternative
splicing
[49059] produces transcripts encoding two forms of the alpha
subunit of GTP-binding

[49060] protein G(o). Proc. Nat. Acad. Sci. 87: 6477-6481, 1990.

[49061] 7663.

[49062] Tsukamoto, T.; Toyama, R.; Itoh, H.; Kozasa, T.; Matsuoka, M.;

[49063] Kaziro, Y.: Structure of the human gene and two rat cD-NAs encoding

[49064] the alpha chain of GTP-binding regulatory protein G(o): two different

[49065] mRNAs are generated by alternative splicing. Proc. Nat. Acad. Sci. 88:

[49066] 2974-2978, 1991.

[49067] 7664.

[49068] Valenzuela, D.; Han, X.; Mende, U.; Fankhauser, C.; Mashimo, H.;

[49069] Huang, P.; Pfeffer, J.; Neer, E. J.; Fishman, M. C.: G-al-pha-o is

[49070] necessary for muscarinic regulation of Ca(2+) channels in mouse heart. Proc.

[49071] Nat. Acad. Sci. 94: 1727-2732, 1997.

[49072] 7665.

[49073] Belluscio, L.; Gold, G. H.; Nemes, A.; Axel, R.: Mice defi-

cient
[49074] in G(olf) are anosmic. Neuron 20: 69-81, 1998.

[49075] 7666.

[49076] Buck, L. B.: Information coding in the mammalian olfactory system. Cold
[49077] Spring Harbor Symp. Quant. Biol. 61: 147-155, 1996.

[49078] 7667.

[49079] Jones, D. T.; Reed, R. R.: Golf: an olfactory neuron specific
[49080] G protein involved in odorant signal transduction. Science 244:
[49081] 790-795, 1989.

[49082] 7668.

[49083] Lindgren, V.; Luskey, K. L.; Russell, D. W.; Francke, U.: Human
[49084] genes involved in cholesterol metabolism: chromosomal mapping of the
[49085] loci for the low density lipoprotein receptor and 3-hydroxy-3-methylglutaryl-coenzyme
[49086] A reductase with cDNA probes. Proc. Nat. Acad. Sci. 82: 8567-8571,
[49087] 1985.

[49088] 7669.

[49089] Luskey, K. L.: Conservation of promoter sequence but not complex

[49090] intron splicing pattern in human and hamster genes for 3-hydroxy-3-methylglutaryl

[49091] coenzyme A reductase. Molec. Cell. Biol. 7: 1881-1893, 1987.

[49092] 7670.

[49093] Mohandas, T.; Heinzmann, C.; Sparkes, R. S.; Wasmuth, J.; Edwards,

[49094] P.; Lusis, A. J.: Assignment of human 3-hydroxy-3-methylglutaryl

[49095] coenzyme A reductase gene to q13-q23 region of chromosome 5. Somat.

[49096] Cell Molec. Genet. 12: 89-94, 1986.

[49097] 7671.

[49098] Osborne, T. F.; Goldstein, J. L.; Brown, M. S.: 5-prime end of

[49099] HMG CoA reductase gene contains sequences responsible for cholesterol-mediated

[49100] inhibition of transcription. Cell 42: 203-212, 1985.

[49101] 7672.

[49102] Van Doren, M.; Broihier, H. T.; Moore, L. A.; Lehmann, R.: HMG-CoA

[49103] reductase guides migrating primordial germ cells. Nature 396: 466-469,

[49104] 1998.

[49105] 7673.

[49106] Ayte, J.; Gil-Gomez, G.; Haro, D.; Marrero, P. F.; Hegardt, F.

[49107] G.: Rat mitochondrial and cytosolic 3-hydroxy-3-methylglutaryl-CoA

[49108] synthases are encoded by two different genes. Proc. Nat. Acad. Sci. 87:

[49109] 3874-3878, 1990.

[49110] 7674.

[49111] Gil, G.; Smith, J. R.; Goldstein, J. L.; Brown, M. S.: Optional

[49112] exon in the 5-prime-untranslated region of 3-hydroxy-3-methylglutaryl

[49113] coenzyme A synthase gene: conserved sequence and splicing pattern

[49114] in humans and hamsters. Proc. Nat. Acad. Sci. 84:

1863-1866, 1987.

[49115] 7675.

[49116] Leonard, S.; Arbogast, D.; Geyer, D.; Jones, C.; Sinensky, M.:

[49117] Localization of the gene encoding 3-hydroxy-3-methylglutaryl-coenzyme

[49118] A synthase to human chromosome 5. Proc. Nat. Acad. Sci. 83: 2187-2189,

[49119] 1986.

[49120] 7676.

[49121] Mehrabian, M.; Callaway, K. A.; Clarke, C. F.; Tanaka, R. D.; Greenspan,

[49122] M.; Lusis, A. J.; Sparkes, R. S.; Mohandas, T.; Edmond, J.; Fogelman,

[49123] A. M.; Edwards, P. A.: Regulation of rat liver 3-hydroxy-3-methylglutaryl

[49124] coenzyme A synthase and the chromosomal localization of the human

[49125] gene. J. Biol. Chem. 261: 16249-16255, 1986.

[49126] 7677.

[49127] Rokosz, L. L.; Boulton, D. A.; Butkiewicz, E. A.; Sanyal, G.;

Cueto,
[49128] M. A.; Lachance, P. A.; Hermes, J. D.: Human cytoplasmic 3-hydroxy-3-methylglutaryl
[49129] coenzyme A synthase: expression, purification, and characterization
[49130] of recombinant wild-type and cys129 mutant enzymes. Arch. Biochem.
[49131] Biophys. 312: 1-13, 1994.

[49132] 7678.

[49133] Russ, A. P.; Ruzicka, V.; Maerz, W.; Appelhans, H.; Gross, W.:
[49134] Amplification and direct sequencing of a cDNA encoding human cytosolic
[49135] 3-hydroxy-3-methylglutaryl-coenzyme A synthase. Biochim. Biophys.
[49136] Acta 1132: 329-331, 1992.

[49137] 7679.

[49138] Hennies, H.-C.; Hagedorn, M.; Reis, A.: Palmoplantar keratoderma
[49139] in association with carcinoma of the esophagus maps to chromosome
[49140] 17q distal to the keratin gene cluster. Genomics 29:

537-540, 1995.

[49141] 7680.

[49142] Howel-Evans, W.; McConnell, R. B.; Clarke, C. A.; Shep-

pard, P.

[49143] M.: Carcinoma of the oesophagus with keratosis palmaris et plantaris

[49144] (tylosis): a study of two families. Quart. J. Med. 27: 413-429,

[49145] 1958.

[49146] 7681.

[49147] Hennekam, R. C. M.: Hereditary multiple exostoses. J. Med. Genet. 28:

[49148] 262-266, 1991.

[49149] 7682.

[49150] Ho, S. U.; Lipton, H. L.: Hereditary multiple exostoses with

[49151] myelopathy. Arch. Neurol. 36: 714, 1979.

[49152] 7683.

[49153] Hogue, D: Personal Communication. Houston, Tex 2/6/1998.

[49154] 7684.

[49155] Knudson, A. G., Jr.: Mutation and cancer: statistical study of
[49156] retinoblastoma. Proc. Nat. Acad. Sci. 68: 820-823, 1971.
[49157] 7685.
[49158] Krooth, R. S.; Macklin, M. T.; Hilbish, T. F.: Diaphysial aclasis
[49159] (multiple exostoses) on Guam. Am. J. Hum. Genet. 13: 340-347, 1961.
[49160] 7686.
[49161] Hou, J.; Parrish, J.; Ludecke, H.-J.; Sapru, M.; Wang, Y.; Chen,
[49162] W.; Hill, A.; Siegel-Bartelt, J.; Northrup, H.; Elder, F. F. B.; Chinault,
[49163] C.; Horsthemke, B.; Wagner, M. J.; Wells, D. E.: A 4-megabase YAC
[49164] contig that spans the Langer-Giedion syndrome region on human chromosome
[49165] 8q24.1: use in refining the location of the trichorhinopha-langeal
[49166] syndrome and multiple exostoses genes (TRPS1 and EXT1). Genomics 29:
[49167] 87-97, 1995.

[49168] 7687.

[49169] Legeai-Mallet, L.; Margaritte-Jeannin, P.; Lemdani, M.; Le Merrer,

[49170] M.; Plauchu, H.; Maroteaux, P.; Munnich, A.; Clerget-Darpoux, F.:

[49171] An extension of the admixture test for the study of genetic heterogeneity

[49172] in hereditary multiple exostoses. Hum. Genet. 99: 298-302, 1997.

[49173] 7688.

[49174] Legeai-Mallet, L.; Munnich, A.; Maroteaux, P.; Le Merrer, M.:

[49175] Incomplete penetrance and expressivity skewing in hereditary multiple

[49176] exostoses. Clin. Genet. 52: 12-16, 1997.

[49177] 7689.

[49178] Le Merrer, M.; Ben Othmane, K.; Stanescu, V.; Lyonnet, S.; Van

[49179] Maldergem, L.; Royer, G.; Munnich, A.; Maroteaux, P.: The gene for

[49180] hereditary multiple exostoses does not map to the Langer-Giedion region

[49181] (8q23-q24). J. Med. Genet. 29: 713-715, 1992.

[49182] 7690.

[49183] Lin, X.; Wells, D.: Isolation of the mouse cDNA homolo-
gous to
[49184] the human EXT1 gene responsible for hereditary multiple
exostoses. DNA
[49185] Seq. 7: 199-202, 1997.

[49186] 7691.

[49187] Lohmann, D. R.; Brandt, B.; Hopping, W.; Passarge, E.;
Horsthemke,
[49188] B.: Spectrum of small length germline mutations in the
RB1 gene. Hum.
[49189] Molec. Genet. 3: 2187-2193, 1994.

[49190] 7692.

[49191] Lohmann, D. R.; Buiting, K.; Ludecke, H.-J.; Horsthemke,
B.:
[49192] The murine Ext1 gene shows a high level of sequence
similarity with
[49193] its human homologue and is part of a conserved linkage
group on chromosome
[49194] 15. Cytogenet. Cell Genet. 76: 164-166, 1997.

[49195] 7693.

[49196] Ludecke, H.-J.; Ahn, J.; Lin, X.; Hill, A.; Wagner, M. J.; Schomburg,

[49197] L.; Horsthemke, B.; Wells, D. E.: Genomic organization and promoter

[49198] structure of the human EXT1 gene. Genomics 40: 351-354, 1997.

[49199] 7694.

[49200] Ludecke, H.-J.; Wagner, M. J.; Nardmann, J.; La Pillo, B.; Parrish,

[49201] J. E.; Willems, P. J.; Haan, E. A.; Frydman, M.; Hamers, G. J. H.;

[49202] Wells, D. E.; Horsthemke, B.: Molecular dissection of a contiguous

[49203] gene syndrome: localization of the genes involved in the Langer-Giedion

[49204] syndrome. Hum. Molec. Genet. 4: 31-36, 1995.

[49205] 7695.

[49206] Matsuno, T.; Ichioka, Y.; Yagi, T.; Ishii, S.: Spindle-cell sarcoma

[49207] in patients who have osteochondromatosis: a report of two cases. J.

[49208] Bone Joint Surg. 70A: 137-141, 1988.

[49209] 7696.

[49210] McCormick, C.; Duncan, G.; Goutsos, K. T.; Tufaro, F.: The putative

[49211] tumor suppressors EXT1 and EXT2 form a stable complex that accumulates

[49212] in the Golgi apparatus and catalyzes the synthesis of heparan sulfate. Proc.

[49213] Nat. Acad. Sci. 97: 668-673, 2000.

[49214] 7697.

[49215] McCormick, C.; Leduc, Y.; Martindale, D.; Mattison, K.; Esford,

[49216] L. E.; Dyer, A. P.; Tufaro, F.: The putative tumour suppressor EXT1

[49217] alters the expression of cell-surface heparan sulfate. Nature Genet. 19:

[49218] 158-161, 1998.

[49219] 7698.

[49220] Mertens, F.; Rydholm, A.; Kreicbergs, A.; Willen, H.; Jonsson,

[49221] K.; Heim, S.; Mitelman, F.; Mandahl, N.: Loss of chromo-

some band
[49222] 8q24 in sporadic osteocartilaginous exostoses. Genes Chromosomes
[49223] Cancer 9: 8-12, 1994.

[49224] 7699.

[49225] Morgan, J. P.; Carlson, W. D.; Adams, O. R.: Hereditary multiple
[49226] exostosis in the horse. J. Am. Vet. Med. Assoc. 140: 1320-1322,
[49227] 1962.

[49228] 7700.

[49229] Ochsner, P. E.: Zum Problem der neoplastischen Entartung bei
[49230] multipen kartilaginaren Exostosen. Z. Orthop. 116: 369-378, 1978.
[49231] 7701.

[49232] Ogle, R. F.; Dalzell, P.; Turner, G.; Wass, D.; Yip, M.-Y.: Multiple
[49233] exostoses in a patient with t(8;11)(q24.11;p15.5). J. Med. Genet. 28:
[49234] 881-883, 1991.

[49235] 7702.

[49236] Peterson, H. A.: Multiple hereditary osteochondromata. Clin.

[49237] Orthop. 239: 222-230, 1989.

[49238] 7703.

[49239] Philippe, C.; Porter, D. E.; Emerton, M. E.; Wells, D. E.; Simpson,

[49240] A. H. R. W.; Monaco, A. P.: Mutation screening of the EXT1 and EXT2

[49241] genes in patients with hereditary multiple exostoses. Am. J. Hum.

[49242] Genet. 61: 520-528, 1997.

[49243] 7704.

[49244] Raskind, W. H.; Conrad, E. U.; Chansky, H.; Matsushita, M.: Loss

[49245] of heterozygosity in chondrosarcomas for markers linked to hereditary

[49246] multiple exostoses loci on chromosomes 8 and 11. Am. J. Hum. Genet. 56:

[49247] 1132-1139, 1995.

[49248] 7705.

[49249] Raskind, W. H.; Conrad, E. U., III; Matsushita, M.; Wijsman, E.
[49250] M.; Wells, D. E.; Chapman, N.; Sandell, L. J.; Wagner, M.; Houck,
[49251] J.: Evaluation of locus heterogeneity and EXT1 mutations in 34 families
[49252] with hereditary multiple exostoses. Hum. Mutat. 11: 231-239, 1998.
[49253] 7706.

[49254] Scholz, W.; Murken, J. D.: Koppelungsuntersuchungen bei Familien
[49255] mit multiplen cartilaginaeren Exostosen. Z. Menschl. Vererb. Konstitutionsl. 37:
[49256] 178-192, 1963.

[49257] 7707.

[49258] Seki, H.; Kubota, T.; Ikegawa, S.; Haga, N.; Fujioka, F.; Ohzeki,
[49259] S.; Wakui, K.; Yoshikawa, H.; Takaoka, K.; Fukushima, Y.: Mutation
[49260] frequencies of EXT1 and EXT2 in 43 Japanese families with hereditary
[49261] multiple exostoses. Am. J. Med. Genet. 99: 59-62, 2001.

[49262] 7708.

[49263] Shapiro, F.; Simon, S.; Glimcher, M. J.: Hereditary multiple

[49264] exostoses: anthropometric, roentgenographic, and clinical aspects. J.

[49265] Bone Joint Surg. 61A: 815-824, 1979.

[49266] 7709.

[49267] Shupe, J. L.; Leone, N. C.; Olson, A. E.; Gardner, E. J.: Hereditary

[49268] multiple exostoses: clinicopathologic features of a comparative study

[49269] in horses and man. Am. J. Vet. Res. 40: 751-757, 1979.

[49270] 7710.

[49271] Silverman, F. N.: Caffey's Pediatric X-ray Diagnosis: An Integrated

[49272] Imaging Approach. Chicago: Year Book Med. Publ. (pub.) 1: 1985.

[49273] Pp. 433-434.

[49274] 7711.

[49275] Solomon, L.: Hereditary multiple exostosis. J. Bone Joint Surg. 45B:

[49276] 292-304, 1963.

[49277] 7712.

[49278] Solomon, L.: Hereditary multiple exostosis. Am. J. Hum. Genet. 16:

[49279] 351-363, 1964.

[49280] 7713.

[49281] Sugiura, Y.; Sugiura, I.; Iwata, H.: Hereditary multiple ex-

ostosis:

[49282] diaphyseal aclasis. Jpn. J. Hum. Genet. 21: 149-167, 1976.

[49283] 7714.

[49284] Vinstein, A. L.; Franken, E. A., Jr.: Hereditary multiple ex-

ostoses:

[49285] report of a case with spinal cord compression. Am. J. Roentgen. 112:

[49286] 405-407, 1971.

[49287] 7715.

[49288] Strathdee, C. A.; Gavish, H.; Shannon, W. R.; Buchwald, M.: Cloning

[49289] of cDNAs for Fanconi's anaemia by functional comple-

mentation. Nature 356:

[49290] 763-767, 1992.

[49291] 7716.

[49292] Koizumi, T.; Hendel, E.; Lalley, P. A.; Tchetgen, M.-B. N.; Nadeau,

[49293] J. H.: Homologs of genes and anonymous loci on human chromosome 13

[49294] map to mouse chromosomes 8 and 14. Mammalian Genome 6: 263-268,

[49295] 1995.

[49296] 7717.

[49297] Bu, X.; Yang, H.-Y.; Shohat, M.; Rotter, J. I.: Two-locus mi-tochondrial

[49298] and nuclear gene models for mitochondrial disorders. Genet. Epidemiol. 9:

[49299] 27-44, 1992.

[49300] 7718.

[49301] Bykhovskaya, Y.; Estivill, X.; Taylor, K.; Hang, T.; Hamon, M.;

[49302] Casano, R. A. M. S.; Yang, H.; Rotter, J. I.; Shohat, M.; Fis-chel-Ghodsian,

[49303] N.: Candidate locus for a nuclear modifier gene for mater-

nally inherited
[49304] deafness. Am. J. Hum. Genet. 66: 1905-1910, 2000.

[49305] 7719.

[49306] Bykhovskaya, Y.; Shohat, M.; Ehrenman, K.; Johnson, D.; Hamon,
[49307] M.; Cantor, R. M.; Aouizerat, B.; Bu, X.; Rotter, J. I.; Jaber, L.;
[49308] Fischel-Ghodsian, N.: Evidence for complex nuclear inher-itance in
[49309] a pedigree with nonsyndromic deafness due to a homo-plasmic mitochondrial
[49310] mutation. Am. J. Med. Genet. 77: 421-426, 1998.

[49311] 7720.

[49312] Bykhovskaya, Y.; Yang, H.; Taylor, K.; Hang, T.; Tun, R. Y. M.;
[49313] Estivill, X.; Casano, R. A. M. S.; Majamaa, K.; Shohat, M.; Fischel-Ghodsian,
[49314] N.: Modifier locus for mitochondrial DNA disease: linkage and linkage
[49315] disequilibrium mapping of a nuclear modifier gene for maternally inherited
[49316] deafness. Genet. Med. 3: 177-180, 2001.

[49317] 7721.

[49318] Guan, M.-X.; Fischel-Ghodsian, N.; Attardi, G.: Biochemical evidence

[49319] for nuclear gene involvement in phenotype of non-syndromic deafness

[49320] associated with mitochondrial 12S rRNA mutation. Hum. Molec. Genet. 5:

[49321] 963-971, 1996.

[49322] 7722.

[49323] Jaber, L.; Shohat, M.; Bu, X.; Fischel-Ghodsian, N.; Yang, H.-Y.;

[49324] Wang, S.-J.; Rotter, J. I.: Sensorineural deafness inherited as a

[49325] tissue specific mitochondrial disorder. J. Med. Genet. 29: 86-90,

[49326] 1992.

[49327] 7723.

[49328] Prezant, T. R.; Agapian, J. V.; Bohlman, M. C.; Bu, X.; Oztas,

[49329] S.; Qiu, W.-Q.; Arnos, K. S.; Cortopassi, G. A.; Jaber, L.; Rotter,

[49330] J. I.; Shohat, M.; Fischel-Ghodsian, N.: Mitochondrial ribo-
somal
[49331] RNA mutation associated with both antibiotic-induced and
non-syndromic
[49332] deafness. Nature Genet. 4: 289-294, 1993.

[49333] 7724.

[49334] Prezant, T. R.; Shohat, M.; Jaber, L.; Pressman, S.; Fischel-
Ghodsian,
[49335] N.: Biochemical characterization of a pedigree with mito-
chondrially
[49336] inherited deafness. Am. J. Med. Genet. 44: 465-472,
1992.
[49337] 7725.

[49338] Chen, M.; Tomkins, D. J.; Auerbach, W.; McKerlie, C.;
Youssoufian,
[49339] H.; Liu, L.; Gan, O.; Carreau, M.; Auerbach, A.; Groves, T.;
Guidos,
[49340] C. J.; Freedman, M. H.; Cross, J.; Percy, D. H.; Dick, J. E.;
Joyner,
[49341] A. L.; Buchwald, M.: Inactivation of Fac in mice produces
inducible
[49342] chromosomal instability and reduced fertility reminiscent

of Fanconi
[49343] anaemia. Nature Genet. 12: 448-451, 1996.

[49344] 7726.

[49345] Gibson, R. A.; Ford, D.; Jansen, S.; Savoia, A.; Havenga, C.; Milner,
[49346] R. D.; de Ravel, T. J.; Cohn, R. J.; Ball, S. E.; Roberts, I.; Llerena,
[49347] J. C.; Vorechovsky, I.; Pearson, T.; Birjandi, F.; Hussein, S. S.;
[49348] Murer-Orlando, M.; Easton, D. F.; Mathew, C. G.: Genetic mapping
[49349] of the FACC gene and linkage analysis in Fanconi anaemia families. J.
[49350] Med. Genet. 31: 868-871, 1994.

[49351] 7727.

[49352] Hoatlin, M. E.; Christianson, T. A.; Keeble, W. W.; Ham-
mond, A.
[49353] T.; Zhi, Y.; Heinrich, M. C.; Tower, P. A.; Bagby, G. C., Jr.: The
[49354] Fanconi anemia group C gene product is located in both the nucleus
[49355] and cytoplasm of human cells. Blood 91: 1418-1425,

1998.

[49356] 7728.

[49357] Montgomery, R. R.; Kunicki, T. J.; Taves, C.; Pidard, D.; Corcoran,

[49358] M.: Diagnosis of Bernard-Soulier syndrome and Glanz-mann's thrombasthenia

[49359] with a monoclonal assay on whole blood. J. Clin. Invest. 71: 385-389,

[49360] 1983.

[49361] 7729.

[49362] Steinmann, B.; Gitzelmann, R.: The diagnosis of hereditary fructose

[49363] intolerance. Helv. Paediat. Acta 36: 297-316, 1981.

[49364] 7730.

[49365] Baerlocher, K.; Gitzelmann, R.; Nussli, R.; Dumermuth, G.: Infantile

[49366] lactic acidosis due to hereditary fructose-1,6-diphosphatase deficiency. Helv.

[49367] Paediat. Acta 26: 489-506, 1971.

[49368] 7731.

[49369] Baker, L.; Winegrad, A. I.: Fasting hypoglycaemia and

metabolic
[49370] acidosis associated with deficiency of hepatic fructose-1,6-diphosphatase
[49371] activity. Lancet II: 13-16, 1970.

[49372] 7732.

[49373] Besley, G. T. N.; Walter, J. H.; Lewis, M. A.; Chard, C. R.; Addison,
[49374] G. M.: Fructose-1,6-bisphosphatase deficiency: severe phenotype with
[49375] normal leukocyte enzyme activity. J. Inherit. Metab. Dis. 17: 333-335,
[49376] 1994.

[49377] 7733.

[49378] Buhrdel, P.; Bohme, H.-J.; Didt, L.: Biochemical and clinical
[49379] observations in four patients with fructose-1,6-diphosphatase deficiency. Europ.
[49380] J. Pediat. 149: 574-576, 1990.

[49381] 7734.

[49382] El-Maghrabi, M. R.; Lange, A. J.; Jiang, W.; Yamagata, K.; Stoffel,
[49383] M.; Takeda, J.; Fernald, A. A.; Le Beau, M. M.; Bell, G. I.;

Baker,

[49384] L.; Pilkis, S. J.: Human fructose-1,6-bisphosphatase gene (FBP1):

[49385] exon-intron organization, localization to chromosome bands 9q22.2-q22.3,

[49386] and mutation screening in subjects with fructose-1,6-bisphosphatase

[49387] deficiency. Genomics 27: 520-525, 1995.

[49388] 7735.

[49389] Gitzelmann, R.; Steinmann, B.; van den Berghe, G.: Disorders of

[49390] fructose metabolism.In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.;

[49391] Valle, D.: The Metabolic Basis of Inherited Disease. New York:

[49392] McGraw-Hill (pub.) (6th ed.) 1: 1989. Pp. 413-417.

[49393] 7736.

[49394] Greene, H. L.; Stifel, F. B.; Herman, R. H.: 'Ketotic hypoglycemia'

[49395] due to hepatic fructose-1,6-diphosphatase deficiency. Am. J. Dis.

[49396] Child. 124: 415-420, 1972.

[49397] 7737.

[49398] Hulsmann, W. C.; Fernandez, J.: A child with lactacidemia and

[49399] fructose diphosphatase deficiency in the liver. Pediat. Res. 5:

[49400] 633-637, 1971.

[49401] 7738.

[49402] Kikawa, Y.; Inuzuka, M.; Jin, B. Y.; Kaji, S.; Koga, J.; Ya-mamoto,

[49403] Y.; Fujisawa, K.; Hata, I.; Nakai, A.; Shigematsu, Y.; Mizunuma, H.;

[49404] Taketo, A.; Mayumi, M.; Sudo, M.: Identification of genetic mutations

[49405] in Japanese patients with fructose-1,6-biphosphatase de-ficiency. Am.

[49406] J. Hum. Genet. 61: 852-861, 1997.

[49407] 7739.

[49408] Kikawa, Y.; Inuzuka, M.; Jin, B. Y.; Kaji, S.; Yamamoto, Y.; Shigematsu,

[49409] Y.; Nakai, A.; Taketo, A.; Ohura, T.; Mikami, H.; Mizunuma, H.; Suzuki,

[49410] Y.; Narisawa, K.; Sudo, M.: Identification of a genetic mutation

[49411] in a family with fructose-1,6-biphosphatase deficiency. Biochem.

[49412] Biophys. Res. Commun. 210: 797-804, 1995.

[49413] 7740.

[49414] Melancon, S. B.; Nadler, H. L.: Detection of fructose-1,6-diphosphatase

[49415] deficiency with use of white blood cells. (Letter) New Eng. J. Med. 286:

[49416] 731-732, 1972.

[49417] 7741.

[49418] Moses, S. W.; Bashan, N.; Flasterstein, B. F.; Rachmel, A.; Gutman,

[49419] A.: Fructose-1,6-diphosphatase deficiency in Israel. Israel J. Med.

[49420] Sci. 27: 1-4, 1991.

[49421] 7742.

[49422] Odievre, M.; Brivet, M.; Moatti, N.; Dreyfus, J.-C.; Beaufils,

[49423] F.; Lejeune, C.; Feffer, J.: Defict en fructose-1,6-diphosphatase

[49424] chez deux soeurs. Arch. Franc. Pediat. 32: 113-122, 1975.

[49425] 7743.

[49426] Pagliara, A. S.; Karl, I. E.; Keating, J. P.; Brown, B. I.; Kipnis,

[49427] D. M.: Hepatic fructose-1,6-diphosphatase deficiency. A cause of

[49428] lactic acidosis and hypoglycemia in infancy. J. Clin. Invest. 51:

[49429] 2115-2123, 1972.

[49430] 7744.

[49431] Holtzman, N. A.; Kronmal, R. A.; van Doorninck, W.; Azen, C.;

[49432] Koch, R.: Effect of age at loss of dietary control on intel-lectual

[49433] performance and behavior of children with phenylke-tonuria. New Eng.

[49434] J. Med. 314: 593-598, 1986.

[49435] 7745.

[49436] Amit, R.; Bashan, N.; Abarbanel, J. M.; Shapira, Y.; Sofer, S.;

[49437] Moses, S.: Fatal familial infantile glycogen storage disease:

multisystem
[49438] phosphofructokinase deficiency. Muscle Nerve 15:
455-458, 1992.
[49439] 7746.

[49440] Argov, Z.; Barash, V.; Soffer, D.; Sherman, J.; Raben, N.:
Late-onset
[49441] muscular weakness in phosphofructokinase deficiency due
to exon 5/intron
[49442] 5 junction point mutation: a unique disorder or the natu-
ral course
[49443] of this glycolytic disorder? Neurology 44: 1097-1100,
1994.
[49444] 7747.

[49445] Bruno, C.; Minetti, C.; Shanske, S.; Morreale, G.; Bado, M.;
Cordone,
[49446] G.; DiMauro, S.: Combined defects of muscle phospho-
fructokinase and
[49447] AMP deaminase in a child with myoglobinuria. Neurology
50: 296-298,
[49448] 1998.

[49449] 7748.

[49450] Danon, M. J.; Carpenter, S.; Manaligod, J. R.; Schliselfeld,

L.

[49451] H.: Fatal infantile glycogen storage disease: deficiency of phosphofructokinase

[49452] and phosphorylase b kinase. Neurology 31: 1303-1307, 1981.

[49453] 7749.

[49454] Danon, M. J.; Servidei, S.; DiMauro, S.; Vora, S.: Late-onset

[49455] muscle phosphofructokinase deficiency. Neurology 38: 956-960, 1988.

[49456] 7750.

[49457] Davidson, M.; Miranda, A. F.; Bender, A. N.; DiMauro, S.; Vora,

[49458] S.: Muscle phosphofructokinase deficiency: biochemical and immunological

[49459] studies of phosphofructokinase isozymes in muscle cul- ture. J. Clin.

[49460] Invest. 72: 545-550, 1983.

[49461] 7751.

[49462] Guibaud, P.; Carrier, H.; Mathieu, M.; Dorche, C.; Par- choux, B.;

[49463] Bethenod, M.; Larbre, F.: Observation familiale de dystro-

phie musculaire
[49464] congenitale par deficit en phosphofructokinase. Arch. Franc. Pediat. 35:
[49465] 1105-1115, 1978.

[49466] 7752.

[49467] Haller, R. G.; Lewis, S. F.: Glucose-induced exertional fa-
tigue
[49468] in muscle phosphofructokinase deficiency. New Eng. J. Med. 324:
[49469] 364-369, 1991.

[49470] 7753.

[49471] Hamaguchi, T.; Nakajima, H.; Noguchi, T.; Nakagawa, C.; Kuwajima,
[49472] M.; Kono, N.; Tarui, S.; Matsuzawa, Y.: Novel missense mutation (W686C)
[49473] of the phosphofructokinase-M gene in a Japanese patient with a mild
[49474] form of glycogenosis VII. Hum. Mutat. 8: 273-275, 1996.

[49475] 7754.

[49476] Hays, A. P.; Hallett, M.; Delfs, J.; Morris, J.; Sotrel, A.; Shevchuk,

[49477] M. M.; DiMauro, S.: Muscle phosphofructokinase defi-
ciency: abnormal
[49478] polysaccharide in a case of late-onset myopathy. Neurol-
ogy 31: 1077-1086,
[49479] 1981.

[49480] 7755.

[49481] Howard, T. D.; Akots, G.; Bowden, D. W.: Physical and ge-
netic
[49482] mapping of the muscle phosphofructokinase gene (PFKM):
reassignment
[49483] to human chromosome 12q. Genomics 34: 122-127,
1996.
[49484] 7756.

[49485] Layzer, R. B.; Rowland, L. P.; Bank, W. J.: Physical and ki-
netic
[49486] properties of human phosphofructokinase from skeletal
muscle and erythrocytes. J.
[49487] Biol. Chem. 244: 3823-3831, 1969.

[49488] 7757.

[49489] Layzer, R. B.; Rowland, L. P.; Ranney, H. M.: Muscle phos-
phofructokinase

[49490] deficiency. Arch. Neurol. 17: 512-523, 1967.

[49491] 7758.

[49492] Mineo, I.; Kono, N.; Hara, N.; Shimizu, T.; Yamada, Y.; Kawachi,

[49493] M.; Kiyokawa, H.; Wang, Y. L.; Tarui, S.: Myogenic hyper-uricemia:

[49494] a common pathophysiologic feature of glycogenosis types III, V, and

[49495] VII. New Eng. J. Med. 317: 75-80, 1987.

[49496] 7759.

[49497] Nakajima, H.: Personal Communication. Osaka, Japan 6/15/1997.

[49498] 7760.

[49499] Nakajima, H.; Kono, N.; Yamasaki, T.; Hotta, K.; Kawachi, M.;

[49500] Kuwajima, M.; Noguchi, T.; Tanaka, T.; Tarui, S.: Genetic defect

[49501] in muscle phosphofructokinase deficiency: abnormal splicing of the

[49502] muscle phosphofructokinase gene due to a point mutation at the 5-prime-splice

[49503] site. J. Biol. Chem. 265: 9392-9395, 1990.

[49504] 7761.

[49505] Nakajima, H.; Noguchi, T.; Yamasaki, T.; Kono, N.; Tanaka, T.;

[49506] Tarui, S.: Cloning of human muscle phosphofructokinase cDNA. FEBS

[49507] Lett. 223: 113-116, 1987.

[49508] 7762.

[49509] Nishikawa, M.; Tsukiyama, K.; Enomoto, T.; Tarui, S.; Okuno, G.;

[49510] Ueda, K.; Ikura, T.; Tsujii, T.; Sugase, T.; Suda, M.; Tanaka, T.

[49511] : A new type of skeletal muscle glycogenosis due to phos-phofructokinase

[49512] deficiency. Proc. Jpn. Acad. 41: 350-353, 1965.

[49513] 7763.

[49514] Raben, N.; Sherman, J.; Miller, F.; Mena, H.; Plotz, P.: A 5-prime

[49515] splice junction mutation leading to exon deletion in an Ashkenazic

[49516] Jewish family with phosphofructokinase deficiency (Tarui

disease). J.

[49517] Biol. Chem. 268: 4963-4967, 1993.

[49518] 7764.

[49519] Raben, N.; Sherman, J.; Nicastri, C.; Adams, E.; Argov, Z.; Nakajima,

[49520] H.; Plotz, P.: A limited number of mutations in the phosphofructokinase

[49521] gene in Ashkenazi Jewish patients with glycogenosis VII (Tarui disease).(Abstract) Am.

[49522] J. Hum. Genet. 53 (suppl.): A942, 1993.

[49523] 7765.

[49524] Raben, N.; Sherman, J. B.: Mutations in muscle phosphofructokinase

[49525] gene. Hum. Mutat. 6: 1-6, 1995.

[49526] 7766.

[49527] Ristow, M.; Vorgerd, M.; Mohlig, M.; Schatz, H.; Pfeiffer, A.

[49528] : Deficiency of phosphofructo-1-kinase/muscle subtype in humans impairs

[49529] insulin secretion and causes insulin resistance. J. Clin. Invest. 100:

[49530] 2833-2841, 1997.

[49531] 7767.

[49532] Bolliger-Stucki, B.; Lord, S. T.; Furlan, M.: Fibrinogen Milano

[49533] XII: a dysfunctional variant containing 2 amino acid substitutions,

[49534] A-alpha R16C and gamma G165R. Blood 98: 351-357, 2001.

[49535] 7768.

[49536] Li, Y.-M.; Xu, M.; Lai, M.-T.; Huang, Q.; Castro, J. L.; DiMuzio-Mower,

[49537] J.; Harrison, T.; Lellis, C.; Nadin, A.; Neduvelil, J. G.; Register,

[49538] R. B.; Sardana, M. K.; Shearman, M. S.; Smith, A. L.; Shi, X.-P.;

[49539] Yin, K.-C.; Shafer, J. A.; Gardell, S. J.: Photoactivated gamma-secretase

[49540] inhibitors directed to the active site covalently label presenilin

[49541] 1. Nature 405: 689-694, 2000.

[49542] 7769.

[49543] Lopera, F.; Ardilla, A.; Martinez, A.; Madrigal, L.; Arango-Viana,

[49544] J. C.; Lemere, C. A.; Arango-Lasprilla, J. C.; Hincapie, L.; Arcos-Burgos,

[49545] M.; Ossa, J. E.; Behrens, I. M.; Norton, J.; Lendon, C.; Goate, A.

[49546] M.; Ruiz-Linares, A.; Rosselli, M.; Kosik, K. S.: Clinical features

[49547] of early-onset Alzheimer disease in a large kindred with an E280A

[49548] presenilin-1 mutation. J.A.M.A. 277: 793-799, 1997.

[49549] 7770.

[49550] Branson, H. E.; Schmer, G.; Dillard, D. H.: Fibrinogen Seattle:

[49551] a qualitatively abnormal fibrinogen in a patient with tetralogy of

[49552] Fallot. Am. J. Clin. Path. 67: 236-240, 1977.

[49553] 7771.

[49554] Brennan, S. O.; Hammonds, B.; George, P. M.: Aberrant hepatic

[49555] processing causes removal of activation peptide and primary polymerisation

[49556] site from fibrinogen Canterbury (A-alpha 20 val-to-asp). J. Clin.

[49557] Invest. 96: 2854-2858, 1995.

[49558] 7772.

[49559] Carrell, N.; Gabriel, D. A.; Blatt, P. M.; Carr, M. E.; McDon-
agh,
[49560] J.: Hereditary dysfibrinogenemia in a patient with throm-
botic disease. Blood 62:
[49561] 432-447, 1983.

[49562] 7773.

[49563] Collen, A.; Maas, A.; Kooistra, T.; Lupu, F.; Grimbergen, J.;

[49564] Haas, F. J. L. M.; Biesma, D. H.; Koolwijk, P.; Koopman, J.;
van Hinsbergh,
[49565] V. W. M.: Aberrant fibrin formation and cross-linking of
fibrinogen
[49566] Nieuwegein, a variant with a shortened A-alpha-chain, al-
ters endothelial
[49567] capillary tube formation. Blood 97: 973-980, 2001.

[49568] 7774.

[49569] Crabtree, G. R.; Kant, J. A.: Molecular cloning of cDNA for
the
[49570] alpha, beta, and gamma chains of rat fibrinogen: a family
of coordinately

[49571] regulated genes. J. Biol. Chem. 256: 9718-9723, 1981.

[49572] 7775.

[49573] Crabtree, G. R.; Kant, J. A.: Coordinate accumulation of the

[49574] mRNAs for the alpha, beta, and gamma chains of rat fibrinogen following

[49575] defibrination. J. Biol. Chem. 257: 7277-7279, 1982.

[49576] 7776.

[49577] Crum, E. D.; Shainoff, J. R.; Graham, R. C.; Ratnoff, O. D.:

[49578] Fibrinogen Cleveland II: an abnormal fibrinogen with defective release

[49579] of fibrinopeptide A. J. Clin. Invest. 53: 1308-1319, 1974.

[49580] 7777.

[49581] Davison, A. M.: The United Kingdom Medical Research Council's

[49582] glomerulonephritis registry. Contrib. Nephrol. 48: 24-35, 1985.

[49583] 7778.

[49584] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Fibrinogen

[49585] and fibrinopeptides. Washington: National Biomedical Re-

search Foundation
[49586] (pub.) 5: 1972. Pp. D87-D97.

[49587] 7779.

[49588] Doolittle, R. F.; Chen, R.; Glasgow, C.; Mross, B.; Weinstein,
[49589] M.: The molecular constancy of fibrinopeptides A and B from 125 individual
[49590] humans. Humangenetik 10: 15-29, 1970.

[49591] 7780.

[49592] Doolittle, R. F.; Takagi, T.; Cottrell, B. A.: Platelet and plasma
[49593] fibrinogens are identical gene products. Science 185: 368-370, 1974.
[49594] 7781.

[49595] Drew, A. F.; Liu, H.; Davidson, J. M.; Daugherty, C. C.; Degen,
[49596] J. L.: Wound-healing defects in mice lacking fibrinogen. Blood 97:
[49597] 3691-3698, 2001.

[49598] 7782.

[49599] Ebert, R. F.: Index of Variant Human Fibrinogens.

Rockville,
[49600] Md.: Privately published (pub.) 1990.

[49601] 7783.

[49602] Ebert, R. F.; Schreiber, W. E.; Bell, W. R.: Fibrinogen Seattle

[49603] II: congenital dysfibrinogenemia with an arg (A alpha 16)-to-his substitution. Thromb.
[49604] Res. 43: 7-13, 1986.

[49605] 7784.

[49606] Forman, W. B.; Ratnoff, O. D.; Boyer, M. H.: An inherited qualitative
[49607] abnormality in plasma fibrinogen: fibrinogen Cleveland. J. Lab. Clin.
[49608] Med. 72: 455-472, 1968.

[49609] 7785.

[49610] Fu, Y.; Cao, Y.; Hertzberg, K. M.; Grieninger, G.: Fibrino-gen
[49611] alpha genes: conservation of bipartite transcripts and car-boxy-terminal-extended
[49612] alpha subunits in vertebrates. Genomics 30: 71-76, 1995.

[49613] 7786.

[49614] Fu, Y.; Weissbach, L.; Plant, P. W.; Oddoux, C.; Cao, Y.; Liang,
[49615] T. J.; Roy, S. N.; Redman, C. M.; Grieninger, G.: Carboxy-terminal-extended
[49616] variant of the human fibrinogen alpha subunit: a novel exon conferring
[49617] marked homology to beta and gamma subunits. Biochemistry 31: 11968-11972,
[49618] 1992.

[49619] 7787.

[49620] Fuchs, G.; Egbring, R.; Havemann, K.: Fibrinogen Marburg--new
[49621] genetic variant of fibrinogen. Blut 34: 107-118, 1977.

[49622] 7788.

[49623] Funk, C.; Straub, P. W.: Hereditary abnormality of fibrin monomer
[49624] aggregation ('fibrinogen Zurich II'). Europ. J. Clin. Invest. 1:
[49625] 131, 1970.

[49626] 7789.

[49627] Galanakis, D. K.: Inherited dysfibrinogenemia: emerging

abnormal
[49628] structure associations with pathologic and nonpathologic dysfunctions. Semin.
[49629] Thromb. Hemost. 19: 386-395, 1993.

[49630] 7790.

[49631] Galanakis, D. K.; Henschen, A.; Keeling, M.; Kehl, M.; Dismore,
[49632] R.; Peerschke, E. I.: Fibrinogen Louisville: an A-alpha-16-arg-to-his
[49633] defect that forms no hybrid molecules in heterozygous individuals
[49634] and inhibits aggregation of normal fibrin monomers. Ann. N.Y. Acad.
[49635] Sci. 408: 644-648, 1983.

[49636] 7791.

[49637] Galanakis, D. K.; Henschen, A.; Peerschke, E. I. B.; Kehl, M.

[49638] : Fibrinogen Stony Brook, a heterozygous A-alpha-16-arg-to-cys dysfibrinogenemia:
[49639] evaluation of diminished platelet aggregation support and of enhanced
[49640] inhibition of fibrin assembly. J. Clin. Invest. 84: 295-304,

1989.

[49641] 7792.

[49642] Godal, H. C.; Brosstad, F.; Kierulf, P.: Three new cases of an

[49643] inborn qualitative fibrinogen defect (fibrinogen Oslo II). Scand.

[49644] J. Haemat. 20: 57–62, 1978.

[49645] 7793.

[49646] Gralnick, H. R.; Coller, B. S.; Fratantoni, J. C.; Martinez, J.

[49647] : Fibrinogen Bethesda III: a hypodysfibrinogenemia. Blood 53: 28–46,

[49648] 1979.

[49649] 7794.

[49650] Gralnick, H. R.; Finlayson, J. S.: Congenital dysfibrinogen-emias. Ann.

[49651] Intern. Med. 77: 471–473, 1972.

[49652] 7795.

[49653] Gralnick, H. R.; Givelber, H. M.; Finlayson, J. S.: Congenital

[49654] dysfibrinogenemia: fibrinogen Bethesda II.In: Abstracts, International

[49655] Society on Thrombosis and Haemostasis. Oslo: Villco Trykkeri (pub.)

[49656] 1972.

[49657] 7796.

[49658] Gralnick, H. R.; Givelber, H. M.; Shainoff, J. R.; Finlayson,

[49659] J. S.: Fibrinogen Bethesda: a congenital dysfibrinogenemia with delayed

[49660] fibrinopeptide release. J. Clin. Invest. 50: 1819-1830, 1971.

[49661] 7797.

[49662] Hampton, J. W.; Garrison, M. S.: Fibrinogen and fibrin-stabilizing

[49663] factor. Med. Clin. N. Am. 56: 133-143, 1972.

[49664] 7798.

[49665] Hampton, J. W.; Morton, R. O.; Bannerjee, D.; Kalmaz, E.: Defective

[49666] fibrin cross-linkages: a genetic and biochemical study of three families.

[49667] (Abstract) J. Clin. Invest. 50: 42A only, 1971.

[49668] 7799.

[49669] Hamsten, A.; Iselius, L.; de Faire, U.; Blomback, M.: Genetic
[49670] and cultural inheritance of plasma fibrinogen concentration. Lancet II:
[49671] 988-990, 1987.

[49672] 7800.

[49673] Alzheimer's Disease Collaborative Group: The structure of the
[49674] presenilin 1 (S182) gene and identification of six novel mutations
[49675] in early onset AD families. Nature Genet. 11: 219-222, 1995.
[49676] 7801.

[49677] Athan, E. S.; Williamson, J.; Ciappa, A.; Santana, V.; Romas, S.
[49678] N.; Lee, J. H.; Rondon, H.; Lantigua, R. A.; Medrano, M.; Torres,
[49679] M.; Arawaka, S.; Rogaeva, E.; and 10 others: A founder mutation
[49680] in presenilin 1 causing early-onset Alzheimer disease in unrelated
[49681] Caribbean Hispanic families. J.A.M.A. 286: 2257-2263,

2001.

[49682] 7802.

[49683] Buckler, A. J.; Chang, D. D.; Graw, S. L.; Brook, J. D.;
Haber,

[49684] D. A.; Sharp, P. A.; Housman, D. E.: Exon amplification: a
strategy

[49685] to isolate mammalian genes based on RNA splicing. Proc.
Nat. Acad.

[49686] Sci. 88: 4005-4009, 1991.

[49687] 7803.

[49688] Campion, D.; Brice, A.; Hannequin, D.; Tardieu, S.; Dubois,
B.;

[49689] Calenda, A.; Brun, E.; Penet, C.; Tayot, J.; Martinez, M.;
Bellis,

[49690] M.; Mallet, J.; Agid, Y.; Clerget-Darpoux, F.: A large pedi-
gree with

[49691] early-onset Alzheimer's disease: clinical, neuropathologic,
and genetic

[49692] characterization. Neurology 45: 80-85, 1995.

[49693] 7804.

[49694] Clark, R. F.; Hutton, M.; Talbot, C.; Wragg, M.; Lendon, C.;

Busfield,
[49695] F.; Han, S. W.; Perez-Tur, J.; Adams, M.; Fuldner, R.; Roberts, G.;
[49696] Karran, E.; Hardy, J.; Goate, A.: The role of presenilin 1 in the
[49697] genetics of Alzheimer's disease. Cold Spring Harbor Symp. Quant.
[49698] Biol. 61: 551-558, 1996.

[49699] 7805.

[49700] Borchelt, D. R.; Thinakaran, G.; Eckman, C. B.; Lee, M. K.; Davenport,
[49701] F.; Ratovitsky, T.; Prada, C.-M.; Kim, G.; Seekins, S.; Yager, D.;
[49702] Slunt, H. H.; Wang, R.; Seeger, M.; Levey, A. I.; Gandy, S. E.; Copeland,
[49703] N. G.; Jenkins, N. A.; Price, D. L.; Younkin, S. G; Sisodia, S. S.
[49704] : Familial Alzheimer's disease-linked presenilin 1 variants elevate
[49705] A-beta-1-42/1-40 ratio in vitro and in vivo. Neuron 17: 1005-1013,
[49706] 1996.

[49707] 7806.

[49708] Citron, M.; Westaway, D.; Xia, W.; Carlson, G.; Diehl, T.; Levesque,

[49709] G.; Johnson-Wood, K.; Lee, M.; Seubert, P.; Davis, A.; Kholodenko,

[49710] D.; Motter, R.; Sherrington, R.; Perry, B.; Yao, H.; Strome, R.; Lieberburg,

[49711] I.; Rommens, J.; Kim. S.; Schenk, D.; Fraser, P.; St George Hyslop,

[49712] P.; Selkoe, D. J.: Mutant presenilins of Alzheimer's disease increase

[49713] production of 42-residue amyloid beta-protein in both transfected

[49714] cells and transgenic mice. Nature Med. 3: 67-72, 1997.

[49715] 7807.

[49716] Crook, R.; Verkkoniemi, A.; Perez-Tur, J.; Mehta N.; Baker, M.;

[49717] Houlden, H.; Farrer, M.; Hutton, M.; Lincoln, S.; Hardy, J.; Gwinn,

[49718] K.; Somer, M.; Paetau, A.; Kalimo, H.; Ylikoski, R.; Poyho-nen, M.;

[49719] Kucera, S.; Haltia, M.: A variant of Alzheimer's disease with

spastic
[49720] paraparesis and unusual plaques due to deletion of exon 9 of presenilin
[49721] 1. Nature Med. 4: 452-455, 1998.

[49722] 7808.

[49723] Cruts, M.; Van Broeckhoven, C.: Presenilin mutations in Alzheimer's
[49724] disease. Hum. Mutat. 11: 183-190, 1998.

[49725] 7809.

[49726] Cruts, M.; van Duijn, C. M.; Backhovens, H.; Van den Broeck, M.;
[49727] Wehnert, A.; Serneels, S.; Sherrington, R.; Hutton, M.; Hardy, J.;
[49728] St George-Hyslop, P. H.; Hofman, A.; Van Broeckhoven, C.: Estimation
[49729] of the genetic contribution of presenilin-1 and -2 muta- tions in a
[49730] population-based study of presenile Alzheimer disease. Hum. Molec.
[49731] Genet. 7: 43-51, 1998.

[49732] 7810.

[49733] Davis, J. A.; Naruse, S.; Chen, H.; Eckman, C.; Younkin, S.; Price,
[49734] D. L.; Borchelt, D. R.; Sisodia, S. S.; Wong, P. C.: An Alzheimer's
[49735] disease-linked PS1 variant rescues the developmental ab-normalities
[49736] of PS1-deficient embryos. Neuron 20: 603-609, 1998.

[49737] 7811.

[49738] De Jonghe, C.; Cruts, M.; Rogaeva, E. A.; Tysoe, C.; Single-ton,
[49739] A.; Vanderstichele, H.; Meschino, W.; Dermaut, B.; Vander-hoeven, I.;
[49740] Backhovens, H.; Vanmechelen, E.; Morris, C. M.; Hardy, J.; Rubinsztein,
[49741] D. C.; St George-Hyslop, P. H.; Van Broeckhoven, C.: Aberrant splicing
[49742] in the presenilin-1 intron 4 mutation causes presenile Alzheimer's
[49743] disease by increased A-beta-42 secretion. Hum. Molec. Genet. 8:
[49744] 1529-1540, 1999.

[49745] 7812.

[49746] Dermaut, B.; Cruts, M.; Slooter, A. J. C.; Van Gestel, S.; De

[49747] Jonghe, C.; Vanderstichele, H.; Vanmechelen, E.; Breteler, M. M.;

[49748] Hofman, A.; van Duijn, C. M.; Van Broeckhoven, C.: The glu318-to-gly

[49749] substitution in presenilin 1 is not causally related to Alzheimer

[49750] disease. (Letter) Am. J. Hum. Genet. 64: 290-292, 1999.

[49751] 7813.

[49752] De Strooper, B.; Annaert, W.; Cupers, P.; Saftig, P.; Craes-saerts,

[49753] K.; Mumm, J. S.; Schroeter, E. H.; Schrijvers, V.; Wolfe, M. S.; Ray,

[49754] W. J.; Goate, A.; Kopan, R.: A presenilin-1-dependent gamma-secretase-like

[49755] protease mediates release of Notch intracellular domain. Nature 398:

[49756] 518-522, 1999.

[49757] 7814.

[49758] De Strooper, B.; Saftig, P.; Craessaerts, K.; Vanderstichele,

[49759] H.; Guhde, G.; Annaert, W.; Von Figura, K.; Van Leuven, F.:

Deficiency
[49760] of presenilin-1 inhibits the normal cleavage of amyloid precursor
[49761] protein. Nature 391: 387-390, 1998.

[49762] 7815.

[49763] Devi, G.; Fotiou, A.; Jyrinji, D.; Tycko, B.; DeArmand, S.; Rogaeva,
[49764] E.; Song, Y.-Q.; Medieros, H.; Liang, Y.; Orlacchio, A.; Williamson,
[49765] J.; St George-Hyslop, P.; Mayeux, R.: Novel presenilin 1 mutations
[49766] associated with early onset of dementia in a family with both early-onset
[49767] and late-onset Alzheimer disease. Arch. Neurol. 57: 1454-1457, 2000.
[49768] 7816.

[49769] Dineley, K. T.; Xia, X.; Bui, D.; Sweatt, J. D.; Zheng, H.: Ac-celerated
[49770] plaque accumulation, associative learning deficits, and up-regulation
[49771] of alpha-7 nicotinic receptor protein in transgenic mice co-expressing

[49772] mutant human presenilin 1 and amyloid precursor proteins. J. Biol.

[49773] Chem. 277: 22768-22780, 2002.

[49774] 7817.

[49775] Donoviel, D. B.; Hadjantonakis, A.-K.; Ikeda, M.; Zheng, H.; St

[49776] George Hyslop, P.; Bernstein, A.: Mice lacking both presenilin genes

[49777] exhibit early embryonic patterning defects. Genes Dev. 13: 2801-2810,

[49778] 1999.

[49779] 7818.

[49780] Duff, K.; Eckman, C.; Zehr, C.; Yu, X; Prada, C.-M.; Perez-tur,

[49781] J.; Hutton, M.; Buee, L.; Harigaya, Y.; Yager, D.; Morgan, D.; Gordon,

[49782] M. N.; Holcomb, L.; Refolo, L.; Zenk, B.; Hardy, J.; Youndkin, S.

[49783] : Increased amyloid-beta-42(43) in brains of mice expressing mutant

[49784] presenilin 1. Nature 383: 710-713, 1996.

[49785] 7819.

[49786] Burrows, N. P.; Nicholls, A. C.; Yates, J. R. W.; Gatward, G.;

[49787] Sarathachandra, P.; Richards, A.; Pope, F. M.: The gene encoding

[49788] collagen alpha-1(V) (COL5A1) is linked to mixed Ehlers-Danlos syndrome

[49789] type I/II. J. Invest. Derm. 106: 1273-1276, 1996.

[49790] 7820.

[49791] Blair, I. P.; Hulme, D.; Dawkins, J. L.; Nicholson, G. A.: A YAC-based

[49792] transcript map of human chromosome 9q22.1-q22.3 encompassing the loci

[49793] for hereditary sensory neuropathy type I and multiple self-healing

[49794] squamous epithelioma. Genomics 51: 277-281, 1998.

[49795] 7821.

[49796] Brennan, R. W.; Dehejia, H.; Kutt, H.; Verebely, K.; McDowell,

[49797] F.: Diphenylhydantoin intoxication attendant to slow inactivation

[49798] of isoniazid. Neurology 20: 687-693, 1970.

[49799] 7822.

[49800] Brown, S.; Chalmers, D. E.: Microsomal epoxide hydrolase activity

[49801] in human x mouse hybrid cells. Biochem. Biophys. Res. Commun. 137:

[49802] 775-780, 1986.

[49803] 7823.

[49804] Buehler, B. A.; Delimont, D.; van Waes, M.; Finnell, R. H.: Prenatal

[49805] prediction of risk of the fetal hydantoin syndrome. New Eng. J. Med. 322:

[49806] 1567-1571, 1990.

[49807] 7824.

[49808] Chodirker, B. N.; Chudley, A. E.; Reed, M. H.; Persaud, T. V. N.

[49809] : Possible prenatal hydantoin effect in a child born to a nonepileptic

[49810] mother. Am. J. Med. Genet. 27: 373-378, 1987.

[49811] 7825.

[49812] De Smet, L.; Debeer, P.: Fetal hydantoin syndrome with unilateral

[49813] atypical cleft hand: additional evidence for vascular disruption. Genet.

[49814] Counsel. 13: 157-161, 2002.

[49815] 7826.

[49816] de Wolff, F. A.; Vermeij, P.; Ferrari, M. D.; Buruma, O. J. S.;

[49817] Breimer, D. D.: Impairment of phenytoin parahydroxylation as a cause

[49818] of severe intoxication. Ther. Drug Monit. 5: 213-215, 1983.

[49819] 7827.

[49820] Gennis, M. A.; Vemuri, R.; Burns, E. A.; Hill, J. V.; Miller, M.

[49821] A.; Spielberg, S. P.: Familial occurrence of hypersensitivity to

[49822] phenytoin. Am. J. Med. 91: 631-634, 1991.

[49823] 7828.

[49824] Goldman, A. S.; Van Dyke, D. C.; Gupta, C.; Katsumata, M.: Elevated

[49825] glucocorticoid receptor levels in lymphocytes of children with the

[49826] fetal hydantoin syndrome (FHS). Am. J. Med. Genet. 28: 607-618,

[49827] 1987.

[49828] 7829.

[49829] Hanson, J. W.; Myrianthopoulos, N. C.; Sedgwick, M. H. A.; Smith,

[49830] D. W.: Risks to the offspring of women treated with hydantoin anticonvulsants,

[49831] with emphasis on the fetal hydantoin syndrome. J. Pediat. 89: 662-668,

[49832] 1976.

[49833] 7830.

[49834] Hartsfield, J. K., Jr.; Sutcliffe, M. J.; Everett, E. T.; Hassett,

[49835] C.; Omiecinski, C. J.; Saari, J. A.: Assignment of microsomal epoxide

[49836] hydrolase (EPHX1) to human chromosome 1q42.1 by in situ hybridization. Cytogenet.

[49837] Cell Genet. 83: 44-45, 1998.

[49838] 7831.

[49839] Aoyama, T.; Francke, U.; Dietz, H. C.; Furthmayr, H.: Quantitative

[49840] differences in biosynthesis and extracellular deposition of fibrillin

[49841] in cultured fibroblasts distinguish five groups of Marfan syndrome
[49842] patients and suggest distinct pathogenetic mechanisms. J. Clin. Invest. 94:
[49843] 130-137, 1994.

[49844] 7832.

[49845] Biery, N. J.; Eldadah, Z. A.; Moore, C. S.; Stetten, G.; Spencer,
[49846] F.; Dietz, H. C.: Revised genomic organization of FBN1 and significance
[49847] for regulated gene expression. Genomics 56: 70-77, 1999.
[49848] 7833.

[49849] Caputi, M.; Kendzior, R. J., Jr.; Beemon, K. L.: A nonsense mutation
[49850] in the fibrillin-1 gene of a Marfan syndrome patient induces NMD and
[49851] disrupts an exonic splicing enhancer. Genes Dev. 16: 1754-1759,
[49852] 2002.

[49853] 7834.

[49854] Cech, T. R.; Bass, B. L.: Biological catalysis by RNA. Ann. Rev.

[49855] Biochem. 55: 599-629, 1986.

[49856] 7835.

[49857] Chikumi, H.; Yamamoto, T.; Ohta, Y.; Nanba, E.; Nagata, K.; Ninomiya,

[49858] H.; Narasaki, K.; Katoh, T.; Hisatome, I.; Ono, K.; Tanaka, Y.; Kuroda,

[49859] H.; Ohgi, S.: Fibrillin gene (FBN1) mutations in Japanese patients

[49860] with Marfan syndrome. Am. J. Hum. Genet. 45: 115-118, 2000.

[49861] 7836.

[49862] Collod, G.; Beroud, C.; Soussi, T.; Junien, C.; Boileau, C.: Software

[49863] and database for the analysis of mutations in the human FBN1 gene. Nucleic

[49864] Acids Res. 24: 137-140, 1996.

[49865] 7837.

[49866] Collod-Beroud, G.; Beroud, C.; Ades, L.; Black, C.; Boxer, M.;

[49867] Brock, D. J.; Godfrey, M.; Hayward, C.; Karttunen, L.; Milewicz, D.;

[49868] Peltonen, L.; Richards, R. I.; Wang, M.; Junien, C.; Boileau, C.:

[49869] Marfan database (second edition): software and database for the analysis

[49870] of mutations in the human FBN1 gene. Nucleic Acids Res. 25: 147-150,

[49871] 1997.

[49872] 7838.

[49873] Usala, A.-L.; Madigan, T.; Burguera, B.; Sinha, M. K.; Caro, J.

[49874] F.; Cunningham, P.; Powell, J. G.; Butler, P. C.: Treatment of insulin-resistant

[49875] diabetic ketoacidosis with insulin-like growth factor I in an adolescent

[49876] with insulin-dependent diabetes. New Eng. J. Med. 327: 853-857,

[49877] 1992.

[49878] 7839.

[49879] Vaessen, N.; Janssen, J. A.; Heutink, P.; Hofman, A.; Lamberts,

[49880] S. W. J.; Oostra, B. A.; Pols, H. A. P.; van Duijn, C. M.: Association
[49881] between genetic variation in the gene for insulin-like growth factor-1
[49882] and low birthweight. Lancet 359: 1036-1037, 2002.

[49883] 7840.

[49884] Van Wyk, J. J.; Svoboda, M. E.; Underwood, L. E.: Evidence from
[49885] radioligand assays that somatomedin-C and insulin-like growth factor-I
[49886] are similar to each other and different from other somatomedins. J.
[49887] Clin. Endocr. Metab. 50: 206-208, 1980.

[49888] 7841.

[49889] Zalin, A. M.; Jones, S.; Fitch, N. J. S.; Ramsden, D. B.: Familial
[49890] nephropathic non-neuropathic amyloidosis: clinical features, immunohistochemistry
[49891] and chemistry. Quart. J. Med. 81: 945-956, 1991.

[49892] 7842.

[49893] Ceccherini, I.; Romei, C.; Barone, V.; Pacini, F.; Martino, E.;

[49894] Loviselli, A.; Pinchera, A.; Romeo, G.: Identification of the cys634-to-tyr
[49895] mutation of the RET proto-oncogene in a pedigree with multiple endocrine
[49896] neoplasia type 2A and localized cutaneous lichen amyloi-dosis. J.
[49897] Endocr. Invest. 17: 201-204, 1994.

[49898] 7843.

[49899] Sarkar, F. H.; Gupta, S. L.: Receptors for human gamma interferon:
[49900] binding and crosslinking of 125-I-labeled recombinant human gamma
[49901] interferon to receptors on WISH cells. Proc. Nat. Acad. Sci. 81:
[49902] 5160-5164, 1984.

[49903] 7844.

[49904] Slate, D. L.; Ruddle, F. H.: Antibodies to chromosome 21 coded
[49905] cell surface components can block response to human in-terferon. Cytogenet.
[49906] Cell Genet. 22: 265-269, 1978.

[49907] 7845.

[49908] Slate, D. L.; Shulman, L.; Lawrence, J. B.; Revel, M.; Ruddle,

[49909] F. H.: Presence of human chromosome 21 alone is sufficient for hybrid

[49910] cell sensitivity to human interferon. J. Virol. 25: 319-325, 1978.

[49911] 7846.

[49912] Takaoka, A.; Mitani, Y.; Suemori, H.; Sato, M.; Yokochi, T.; Noguchi,

[49913] S.; Tanaka, N.; Taniguchi, T.: Cross talk between interferon-gamma

[49914] and -alpha/beta signaling components in caveolar membrane domains. Science 288:

[49915] 2357-2360, 2000.

[49916] 7847.

[49917] Tan, Y. H.: Chromosome 21 and the cell growth inhibitory effect

[49918] of human interferon preparations. Nature 260: 141-143, 1976.

[49919] 7848.

[49920] Tan, Y. H.; Schneider, E. L.; Tischfield, J.; Epstein, C. J.;

[49921] Ruddle, F. H.: Human chromosome 21 dosage: effect on the expression
[49922] of the interferon induced antiviral state. Science 186: 61-63, 1974.
[49923] 7849.

[49924] Tan, Y. H.; Tischfield, J.; Ruddle, F. H.: The linkage of genes
[49925] for the human interferon-induced antiviral protein and indophenoloxidase-B
[49926] traits to chromosome G-21. J. Exp. Med. 37: 317-330, 1973.
[49927] 7850.

[49928] Weil, J.; Tucker, G.; Epstein, L. B.; Epstein, C. J.: Interferon
[49929] induction of (2-prime-5-prime) oligoisoadenylate synthetase in diploid
[49930] and trisomy 21 human fibroblasts: relation to dosage of the interferon
[49931] receptor gene (IFRC). Hum. Genet. 65: 108-111, 1983.

[49932] 7851.

[49933] Wiranowska-Stewart, M.; Stewart, W. E., II: The role of human

[49934] chromosome 21 in sensitivity to interferons. J. Gen. Virol. 37:
[49935] 629-633, 1977.

[49936] 7852.

[49937] Aguet, M.; Dembic, Z.; Merlin, G.: Molecular cloning and expression
[49938] of the human interferon-gamma receptor. Cell 55: 273-280, 1988.

[49939] 7853.

[49940] Alcaide-Loridan, C.; Le Coniat, M.; Bono, R.; Benech, P.; Couillin,
[49941] P.; Van Cong, N.; Fisher, D. N.; Berger, R.; Fellous, M.: Mapping
[49942] of the human interferon gamma response. (Abstract) Cytogenet. Cell
[49943] Genet. 51: 949 only, 1989.

[49944] 7854.

[49945] Branca, A. A.; Baglioni, C.: Evidence that types I and II interferons
[49946] have different receptors. Nature 294: 768-770, 1981.

[49947] 7855.

[49948] Casanova, J.-L.; Jouanguy, E.; Lamhamedi, S.; Blanche, S.; Fischer,

[49949] A.: Immunological conditions of children with BCG disseminated infection.

[49950] (Letter) Lancet 346: 581 only, 1995.

[49951] 7856.

[49952] Celada, A.; Allen, R.; Esparza, I.; Gray, P. W.; Schreiber, R.

[49953] D.: Demonstration and partial characterization of the interferon-gamma

[49954] receptor on human mononuclear phagocytes. J. Clin. Invest. 76: 2196-2205,

[49955] 1985.

[49956] 7857.

[49957] Lee, W. M.; Galbraith, R. M.: The extracellular actin-scavenger

[49958] system and actin toxicity. New Eng. J. Med. 326: 1335-1341, 1992.

[49959] 7858.

[49960] Pilz, A.; Moseley, H.; Peters, J.; Abbott, C.: Comparative

mapping
[49961] of mouse chromosome 2 and human chromosome 9q: the genes for gelsolin
[49962] and dopamine beta-hydroxylase map to mouse chromosome 2. Genomics 12:
[49963] 715-719, 1992.

[49964] 7859.

[49965] de Lecea, L.; Kilduff, T. S.; Peyron, C.; Gao, X.-B.; Foye, P.

[49966] E.; Danielson, P. E.; Fukuhara, C.; Battenberg, E. L. F.; Gautvik,
[49967] V. T.; Bartlett, F. S., II; Frankel, W. N.; van den Pol, A. N.; Bloom,
[49968] F. E.; Gautvik, K. M.; Sutcliffe, J. G.: The hypocretins: hypothalamus-specific
[49969] peptides with neuroexcitatory activity. Proc. Nat. Acad. Sci. 95:
[49970] 322-327, 1998.

[49971] 7860.

[49972] Lin, L.; Faraco, J.; Li, R.; Kadotani, H.; Rogers, W.; Lin, X.;

[49973] Qiu, X.; de Jong, P. J.; Nishino, S.; Mignot, E.: The sleep disorder

[49974] canine narcolepsy is caused by a mutation in the hypocretin (orexin)
[49975] receptor 2 gene. Cell 98: 365-376, 1999.

[49976] 7861.

[49977] Ostrander, E. A.; Giniger, E.: Let sleeping dogs lie? Nature

[49978] Genet. 23: 3-4, 1999.

[49979] 7862.

[49980] Peyron, C.; Faraco, J.; Rogers, W.; Ripley, B.; Overeem, S.; Charnay,
[49981] Y.; Nevsimalova, S.; Aldrich, M.; Reynolds, D.; Albin, R.; Li, R.;
[49982] Hungs, M.; and 10 others: A mutation in a case of early onset narcolepsy
[49983] and a generalized absence of hypocretin peptides in human narcoleptic
[49984] brains. Nature Med. 6: 991-997, 2000.

[49985] 7863.

[49986] Siegel, J. M.: Narcolepsy: a key role for hypocretins (orexins). Cell 98:
[49987] 409-412, 1999.

[49988] 7864.

[49989] Thannickal, T. C.; Moore, R. Y.; Nienhuis, R.; Ramanathan, L.;

[49990] Gulyani, S.; Aldrich, M.; Cornford, M.; Siegel, J. M.: Reduced number

[49991] of hypocretin neurons in human narcolepsy. Neuron 27: 469-474, 2000.

[49992] 7865.

[49993] Chakrani, F.; Armand, J.-P.; Lenoir, G.; Ju, L.; Liang, J.-P.;

[49994] May, E.; May, P.: Mutations clustered in exon 5 of the p53 gene in

[49995] primary nasopharyngeal carcinomas from southeastern Asia. Int. J.

[49996] Cancer 61: 316-320, 1995.

[49997] 7866.

[49998] Koufos, A.; Hansen, M. F.; Copeland, N. G.; Jenkins, N. A.; Lampkin,

[49999] B. C.; Cavenee, W. K.: Loss of heterozygosity in three embryonal

[50000] tumours suggests a common pathogenetic mechanism. Nature 316: 330-334,

[50001] 1985.

[50002] 7867.

[50003] Lam, W. W. K.; Hatada, I.; Ohishi, S.; Mukai, T.; Joyce, J. A.;

[50004] Cole, T. R. P.; Donnai, D.; Reik, W.; Schofield, P. N.; Maher,
E.

[50005] R.: Analysis of germline CDKN1C (p57-KIP2) mutations in
familial

[50006] and sporadic Beckwith-Wiedemann syndrome (BWS) pro-
vides a novel genotype-phenotype

[50007] correlation. J. Med. Genet. 36: 518-523, 1999.

[50008] 7868.

[50009] Eglen, R. M.; Whiting, R. L.: Muscarinic receptor subtypes:
a

[50010] critique of the current classification and a proposal for a
working

[50011] nomenclature. J. Auton. Pharm. 6: 323-346, 1986.

[50012] 7869.

[50013] Goyal, R. K.: Muscarinic receptor subtypes: physiology and
clinical

[50014] implications. New Eng. J. Med. 321: 1022-1029, 1989.

[50015] 7870.

[50016] Liao, C. F.; Themmen, A. P.; Joho, R.; Barberis, C.; Birn-
baumer,
[50017] M.; Birnbaumer, L.: Molecular cloning and expression of a
fifth muscarinic
[50018] acetylcholine receptor. J. Biol. Chem. 264: 7328-7337,
1989.
[50019] 7871.

[50020] Peralta, E. G.; Ashkenazi, A.; Winslow, J. W.; Smith, D. H.;
Ramachandran,
[50021] J.; Capon, D. J.: Distinct primary structures, ligand-bind-
ing properties
[50022] and tissue-specific expression of four human muscarinic
acetylcholine
[50023] receptors. EMBO J. 6: 3923-3929, 1987.

[50024] 7872.

[50025] Chini, B.; Raimond, E.; Elgoyhen, A. B.; Moralli, D.;
Balzaretti,
[50026] M.; Heinemann, S.: Molecular cloning and chromosomal
localization
[50027] of the human alpha-7-nicotinic receptor subunit gene
(CHRNA7). Genomics 19:

[50028] 379-381, 1994.

[50029] 7873.

[50030] Doucette-Stamm, L.; Monteggia, L. M.; Donnelly-Roberts, D.; Wang,

[50031] M. T.; Lee, J.; Tian, J.; Giordano, T.: Cloning and sequence of the

[50032] human alpha-7 nicotinic acetylcholine receptor. Drug Dev. Res. 30:

[50033] 252-256, 1993.

[50034] 7874.

[50035] Elmslie, F.; Williamson, M. P.; Rees, M.; Kerr, M.; Kjeldsen, M.

[50036] J.; Pang, K. A.; Sundqvist, A.; Friis, M. L.; Chadwick, D.; Richens,

[50037] A.; Curtis, D.; Whitehouse, W. P.; Gardiner, R. M.: Genetic mapping

[50038] of a major susceptibility locus for juvenile myoclonic epilepsy on

[50039] chromosome 15q. Medizinische Genetik 9: 15 only, 1997.

[50040] 7875.

[50041] Freedman, R.; Coon, H.; Myles-Worsley, M.; Orr-Urtreger,

A.; Olincy,

[50042] A.; Davis, A.; Polymeropoulos, M.; Holik, J.; Hopkins, J.; Hoff, M.;

[50043] Rosenthal, J.; Waldo, M. C.; and 11 others: Linkage of a neurophysiological

[50044] deficit in schizophrenia to a chromosome 15 locus. Proc. Nat. Acad.

[50045] Sci. 94: 587-592, 1997.

[50046] 7876.

[50047] Orr-Urtreger, A.; Seldin, M. F.; Baldini, A.; Beaudet, A. L.:

[50048] Cloning and mapping of the mouse alpha-7-neuronal nicotinic acetylcholine

[50049] receptor. Genomics 26: 399-402, 1995.

[50050] 7877.

[50051] Peng, X.; Katz, M.; Gerzanich, V.; Anand, R.; Lindstrom, J.: Human

[50052] alpha-7 acetylcholine receptor: cloning of the alpha-7 subunit from

[50053] the SH-SY5Y cell line and determination of pharmacologi-cal properties

[50054] of native receptors and functional alpha-7 homomers ex-

pressed in Xenopus
[50055] oocytes. Molec. Pharm. 45: 546-554, 1994.

[50056] 7878.

[50057] Wisniewski, L.; Hassold, T.; Heffelfinger, J.; Higgins, J. V.:

[50058] Cytogenetic and clinical studies in five cases of inv dup (15). Hum.
[50059] Genet. 50: 259-270, 1979.

[50060] 7879.

[50061] Mortier, G. R.; Wilkin, D. J.; Wilcox, W. R.; Rimoin, D. L.; Lachman,
[50062] R. S.; Eyre, D. R.; Cohn, D. H.: A radiographic, morpho-logic, biochemical
[50063] and molecular analysis of a case of achondrogenesis type II resulting
[50064] from substitution for a glycine residue (gly691-to-arg) in the type
[50065] II collagen trimer. Hum. Molec. Genet. 4: 285-288, 1995.

[50066] 7880.

[50067] Nelson, F.; Dahlberg, L.; Laverty, S.; Reiner, A.; Pidoux, I.;

[50068] Ionescu, M.; Fraser, G. L.; Brooks, E.; Tanzer, M.; Rosen-

berg, L.

[50069] C.; Dieppe, P.; Poole, A. R.: Evidence for altered synthesis of type

[50070] II collagen in patients with osteoarthritis. J. Clin. Invest. 102:

[50071] 2115-2125, 1998.

[50072] 7881.

[50073] Nunez, A. M.; Francomano, C.; Young, M. F.; Martin, G. R.; Yamada,

[50074] Y.: Isolation and partial characterization of genomic clones coding

[50075] for a human pro-alpha-1(II) collagen chain and demonstration of restriction

[50076] fragment length polymorphism at the 3-prime end of the gene. Biochemistry 24:

[50077] 6343-6348, 1985.

[50078] 7882.

[50079] Pope, F. M.; Cheah, K. S. E.; Nicholls, A. C.; Price, A. B.; Grosveld,

[50080] F. G.: Lethal osteogenesis imperfecta congenita and a 300 base pair

[50081] gene deletion for alpha-1-like collagen. Brit. Med. J. 288:

431-434,

[50082] 1984.

[50083] 7883.

[50084] Reginato, A. J.; Passano, G. M.; Neumann, G.; Falasca, G. F.;

[50085] Diaz-Valdez, M.; Jimenez, S. A.; Williams, C. J.: Familial spondyloepiphyseal

[50086] dysplasia tarda, brachydactyly, and precocious osteoarthritis associated

[50087] with an arginine75-to-cysteine mutation in the procollagen type II

[50088] gene in a kindred of Chiloe Islanders. I. Clinical, radiographic,

[50089] and pathologic findings. Arthritis Rheum. 37: 1078-1086, 1994.

[50090] 7884.

[50091] Ritvaniemi, P.; Hyland, J.; Ignatius, J.; Kivirikko, K. I.; Prockop,

[50092] D. J.; Ala-Kokko, L.: A fourth example suggests that premature termination

[50093] codons in the COL2A1 gene are a common cause of the Stickler syndrome:

[50094] analysis of the COL2A1 gene by denaturing gradient gel electrophoresis. Genomics 17:
[50095] 218-221, 1993.

[50096] 7885.

[50097] Sangiorgi, F.; Huerre-Jeanpierre, C.; Weil, D.; Grzeschik, K.

[50098] H.; Junien, C.; Sobel, M.; Ramirez, F.: Chromosomal assignment of
[50099] the human type II collagen gene. (Abstract) Am. J. Hum. Genet. 36:
[50100] 208S only, 1984.

[50101] 7886.

[50102] Sangiorgi, F. O.; Benson-Chanda, V.; de Wet, W. J.; Sobel, M.
[50103] E.; Tsipouras, P.; Ramirez, F.: Isolation and partial characterization
[50104] of the entire human pro-alpha-1(II) collagen gene. Nucleic Acids
[50105] Res. 13: 2207-2225, 1985.

[50106] 7887.

[50107] Sher, C.; Ramesar, R.; Martell, R.; Learmonth, I.; Tsipouras,

[50108] P.; Beighton, P.: Mild spondyloepiphyseal dysplasia (Namaqualand

[50109] type): genetic linkage to the type II collagen gene (COL2A1). Am.

[50110] J. Hum. Genet. 48: 518-524, 1991.

[50111] 7888.

[50112] Siggers, D. C.: U. Kniest disease.:In: Bergsma, D. (ed.): Skeletal

[50113] Dysplasias. New York: American Elsevier 1974. Pg. 432-442.

[50114] 7889.

[50115] Solomon, E.; Hiorns, L. R.; Cheah, K. S. E.; Parkar, M.; Weiss,

[50116] E.; Flavell, R. A.: Assignment of a human alpha-1(I)-like collagen

[50117] gene to chromosome 12, by molecular hybridization. (Abstract) Cytogenet.

[50118] Cell Genet. 37: 588-589, 1984.

[50119] 7890.

[50120] Spector, T. D.; Cicuttini, F.; Baker, J.; Loughlin, J.; Hart,

[50121] D.: Genetic influences on osteoarthritis in women: a twin

study. Brit.

[50122] Med. J. 312: 940-944, 1996.

[50123] 7891.

[50124] Stoker, N. G.; Cheah, K. S. E.; Griffin, J. R.; Pope, F. M.; Solomon,

[50125] E.: A highly polymorphic region 3-prime to the human type II collagen

[50126] gene. Nucleic Acids Res. 13: 4613-4622, 1985.

[50127] 7892.

[50128] Strom, C. M.: Achondroplasia due to DNA insertion into the type

[50129] II collagen gene. (Abstract) Pediat. Res. 18: 226A only, 1984.

[50130] 7893.

[50131] Strom, C. M.; Eddy, R. L.; Shows, T. B.: Localization of human

[50132] type II procollagen gene (COL2A1) to chromosome 12. Somat. Cell Molec.

[50133] Genet. 10: 651-655, 1984.

[50134] 7894.

[50135] Strom, C. M.; Upholt, W. B.: Isolation and characterization

of

[50136] genomic clones corresponding to the human type II pro-collagen gene. Nucleic

[50137] Acids Res. 12: 1025-1038, 1984.

[50138] 7895.

[50139] Sykes, B.; Smith, R.; Vipond, S.; Paterson, C.; Cheah, K.; Solomon,

[50140] E.: Exclusion of the alpha-1(II) cartilage collagen gene as the mutant

[50141] locus in type IA osteogenesis imperfecta. J. Med. Genet. 22: 187-191,

[50142] 1985.

[50143] 7896.

[50144] Sykes, B. C.; Ogilvie, D. J.; Wordsworth, B. P.: Lethal osteo-genesis

[50145] imperfecta and a collagen gene deletion: length polymor-phism provides

[50146] an alternative explanation. Hum. Genet. 70: 35-37, 1985.

[50147] 7897.

[50148] Takahashi, E.; Hori, T.; Sutherland, G. R.: Mapping of the human

[50149] type II collagen gene (COL2A1) proximal to fra(12)(q13.1) by nonisotopic
[50150] in situ hybridization. Cytogenet. Cell Genet. 54: 84-85, 1990.
[50151] 7898.

[50152] Griffin, C. A.; McKeon, C.; Israel, M. A.; Gegonne, A.; Ghysdael,
[50153] J.; Stehelin, D.; Douglass, E. C.; Green, A. A.; Emanuel, B. S.:
[50154] Comparison of constitutional and tumor-associated 11;22 translocations:
[50155] nonidentical breakpoints on chromosomes 11 and 22. Proc. Nat. Acad.
[50156] Sci. 83: 6122-6126, 1986.

[50157] 7899.

[50158] Mastrangelo, T.; Modena, P.; Tornielli, S.; Bullrich, F.; Testi,
[50159] M. A.; Mezzelani, A.; Radice, P.; Azzarelli, A.; Pilotti, S.; Croce,
[50160] C. M.; Pierotti, M. A.; Sozzi, G.: A novel zinc finger gene is fused
[50161] to EWS in small round cell tumor. Oncogene 19:

3799-3804, 2000.

[50162] 7900.

[50163] Baden, H. P.; Roth, S. I.; Goldsmith, L. A.; Lee, L. D.: Keratohyalin

[50164] protein in disorders of keratinization. J. Invest. Derm. 62: 411-414,

[50165] 1974.

[50166] 7901.

[50167] Gan, S.-Q.; McBride, O. W.; Idler, W. W.; Markova, N.; Steinert,

[50168] P. M.: Organization, structure, and polymorphisms of the human profilaggrin

[50169] gene. Biochemistry 29: 9432-9440, 1990.

[50170] 7902.

[50171] Holbrook, K. A.; Dale, B. A.; Brown, K. S.: Abnormal epidermal

[50172] keratinization in the repeated epilation mutant mouse. J. Cell Biol. 92:

[50173] 387-397, 1982.

[50174] 7903.

[50175] McKinley-Grant, L. J.; Idler, W. W.; Bernstein, I. A.; Parry,

D.

[50176] A. D.; Cannizzaro, L.; Croce, C. M.; Huebner, K.; Lessin, S. R.; Steinert,

[50177] P. M.: Characterization of a cDNA clone encoding human filaggrin

[50178] and localization of the gene to chromosome region 1q21. Proc. Nat.

[50179] Acad. Sci. 86: 4848-4852, 1989.

[50180] 7904.

[50181] Rothnagel, J. A.; Longley, M. A.; Bundman, D. S.; Naylor, S. L.;

[50182] Lalley, P. A.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Roop,

[50183] D. R.: Characterization of the mouse loricrin gene: linkage with

[50184] profilaggrin and the flaky tail and soft coat mutant loci on chromosome

[50185] 3. Genomics 23: 450-456, 1994.

[50186] 7905.

[50187] Sybert, V. P.; Dale, B. A.; Holbrook, K. A.: Ichthyosis vul-garis:

[50188] identification of a defect in synthesis of filaggrin corre-

lated with
[50189] an absence of keratohyaline granules. J. Invest. Derm. 84: 191-194,
[50190] 1985.

[50191] 7906.

[50192] Volz, A.; Korge, B. P.; Compton, J. G.; Ziegler, A.; Steinert,

[50193] P. M.; Mischke, D.: Physical mapping of a functional clus-
ter of epidermal
[50194] differentiation genes on chromosome 1q21. Genomics 18: 92-99, 1993.
[50195] 7907.

[50196] Barbosa, C. A. A.; Koury, W. H.; Krieger, H.: Linkage data on
[50197] MN and the Hb beta locus. Am. J. Hum. Genet. 27: 797-801, 1975.
[50198] 7908.

[50199] Riddell, D. C.; Wang, H.; Umbenhauer, D. R.; Beaume, P. H.; Guengerich,
[50200] F. P.; Hamerton, J. L.: Regional assignment for the genes encoding
[50201] human P450IIIA3 (CYP3) and P450IIC9 (CYP2C). (Abstract)

Cytogenet.

[50202] Cell Genet. 46: 682, 1987.

[50203] 7909.

[50204] Yoder, J. A.; Yen, R.-W. C.; Vertino, P. M.; Bestor, T. H.; Baylin,

[50205] S. B.: New 5-prime regions of the murine and human genes for DNA

[50206] (cytosine-5)-methyltransferase. J. Biol. Chem. 271: 31092-31097,

[50207] 1996.

[50208] 7910.

[50209] Hofmann, S.; Lichtner, P.; Schuffenhauer, S.; Gerbitz, K.-D.; Meitinger,

[50210] T.: Assignment of the human genes coding for cytochrome c oxidase

[50211] subunits Va (COX5A), VIc (COX6C) and VIIc (COX7C) to chromosome bands

[50212] 15q25, 8q22-q23 and 5q14 and of three pseudogenes (COX5AP1, COX6CP1,

[50213] COX7CP1) to 14q22, 16p12 and 13q14-q21 by FISH and radiation hybrid

[50214] mapping. Cytogenet. Cell Genet. 83: 226-227, 1998.

[50215] 7911.

[50216] Martin-Gallardo, A.; McCombie, W. R.; Gocayne, J. D.; FitzGerald,

[50217] M. G.; Wallace, S.; Lee, B. M. B.; Lamerdin, J.; Trapp, S.; Kelley,

[50218] J. M.; Liu, L.-I.; Dubnick, M.; Johnston-Dow, L. A.; Kerlavage, A.

[50219] R.; de Jong, P.; Carrano, A.; Fields, C.; Venter, J. C.: Auto-mated

[50220] DNA sequencing and analysis of 106 kilobases from hu-man chromosome

[50221] 19q13.3. Nature Genet. 1: 34-39, 1992.

[50222] 7912.

[50223] Volker, M.; Mone, M. J.; Karmakar, P.; van Hoffen, A.; Schul,

[50224] W.; Vermeulen, W.; Hoeijmakers, J. H. J.; van Driel, R.; van Zeeland,

[50225] A. A.; Mullenders, L. H. F.: Sequential assembly of the nu-cleotide

[50226] excision repair factors in vivo. Molec. Cell 8: 213-224, 2001.

[50227] 7913.

[50228] Schellens, J. H. M.; Soons, P. A.; Breimer, D. D.: Lack of bi-
modality
[50229] in nifedipine plasma kinetics in a large population of
healthy subjects. Biochem.
[50230] Pharm. 37: 2507-2510, 1988.

[50231] 7914.

[50232] Shet, M. S.; Fisher, C. W.; Holmans, P. L.; Estabrook, R. W.:

[50233] Human cytochrome P450 3A4: enzymatic properties of a
purified recombinant
[50234] fusion protein containing NADPH-P450 reductase. Proc.
Nat. Acad.
[50235] Sci. 90: 11748-11752, 1993.

[50236] 7915.

[50237] Shimada, T.; Guengerich, F. P.: Evidence for cytochrome P-
450(NF),
[50238] the nifedipine oxidase, being the principal enzyme in-
volved in the
[50239] bioactivation of aflatoxins in human liver. Proc. Nat. Acad.
Sci. 86:
[50240] 462-465, 1989.

[50241] 7916.

[50242] Spurr, N. K.; Gough, A. C.; Stevenson, K.; Wolf, C. R.: The human
[50243] cytochrome P450 CYP3 locus: assignment to chromosome 7q22-qter. Hum.
[50244] Genet. 81: 171-174, 1989.

[50245] 7917.

[50246] Walker, A. H.; Jaffe, J. M.; Gunasegaram, S.; Cummings, S. A.;
[50247] Huang, C.-S.; Chern, H.-D.; Olopade, O. I.; Weber, B. L.; Rebbeck,
[50248] T. R.: Characterization of an allelic variant in the nifedip-ine-specific
[50249] element of CYP3A4: ethnic distribution and implications for prostate
[50250] cancer risk. Hum. Mutat. 12: 289-293, 1998.

[50251] 7918.

[50252] Watkins, P. B.; Wrighton, S. A.; Maurel, P.; Schuetz, E. G.; Mendez-Picon,
[50253] G.; Parker, G. A.; Guzelian, P. S.: Identification of an in-ducible
[50254] form of cytochrome P-450 in human liver. Proc. Nat. Acad. Sci. 82:

[50255] 6310-6314, 1985.

[50256] 7919.

[50257] Wolf, C. R.; Smith, C. A. D.; Gough, A. C.; Moss, J. E.; Val-
lis,

[50258] K. A.; Howard, G.; Carey, F. J.; Mills, K.; McNee, W.;
Carmichael,

[50259] J.; Spurr, N. K.: Relationship between the debrisoquine hy-
droxylase

[50260] polymorphism and cancer susceptibility. Carcinogenesis
13: 1035-1038,

[50261] 1992.

[50262] 7920.

[50263] Wrighton, S. A.; Stevens, J. C.: The human hepatic cy-
tochromes

[50264] P450 involved in drug metabolism. Crit. Rev. Toxicol. 22:
1-21,

[50265] 1992.

[50266] 7921.

[50267] Wrighton, S. A.; Vandenbranden, M.: Isolation and charac-
terization

[50268] of human fetal liver cytochrome P450HLp2: a third mem-

ber of the P450III
[50269] gene family. Arch. Biochem. Biophys. 268: 144-151, 1989.
[50270] 7922.
[50271] Xie, W.; Barwick, J. L.; Downes, M.; Blumberg, B.; Simon, C. M.;
[50272] Nelson, M. C.; Neuschwander-Tetri, B. A.; Brunt, E. M.; Guzelian,
[50273] P. S.; Evans, R. M.: Humanized xenobiotic response in mice expressing
[50274] nuclear receptor SXR. Nature 406: 435-439, 2000.
[50275] 7923.
[50276] Gray, I. C.; Nobile, C.; Muresu, R.; Ford, S.; Spurr, N. K.: A
[50277] 2.4-megabase physical map spanning the CYP2C gene cluster on chromosome
[50278] 10q24. Genomics 28: 328-332, 1995.
[50279] 7924.
[50280] Guengerich, F. P.; Distlerath, L. M.; Reilly, P. E. B.; Wolff,
[50281] T.; Shimada, T.; Umbenhauer, D. R.; Martin, M. V.: Hu-man-liver cytochromes
[50282] P-450 involved in polymorphisms of drug oxidation.

Xenobiotica 16:
[50283] 367-378, 1986.

[50284] 7925.

[50285] Weinberger, C.; Hollenberg, S. M.; Rosenfeld, M. G.; Evans, R.
[50286] M.: Domain structure of human glucocorticoid receptor and its relationship
[50287] to the v-erb-A oncogene product. Nature 318: 670-672, 1985.
[50288] 7926.

[50289] Colao, A.; Di Somma, C.; Pivonello, R.; Cuocolo, A.; Spinelli,
[50290] L.; Bonaduce, D.; Salvatore, M.; Lombardi, G.: The cardio-vascular
[50291] risk of adult GH deficiency (GHD) improved after GH re-placement and
[50292] worsened in untreated GHD: a 12-month prospective study. J. Clin.
[50293] Endocr. Metab. 87: 1088-1093, 2002.

[50294] 7927.

[50295] Lenka, N.; Vijayasarathy, C.; Mullick, J.; Avadhani, N. G.:

Structural
[50296] organization and transcription regulation of nuclear genes encoding
[50297] the mammalian cytochrome c oxidase complex. Prog. Nucleic Acid Res.
[50298] Molec. Biol. 61: 309-344, 1998.

[50299] 7928.

[50300] Otsuka, M.; Mizuno, Y.; Yoshida, M.; Kagawa, Y.; Ohta, S.: Nucleotide
[50301] sequence of cDNA encoding human cytochrome c oxidase subunit Vlc. Nucleic
[50302] Acids Res. 16: 10916 only, 1988.

[50303] 7929.

[50304] Akbari, O.; Freeman, G. J.; Meyer, E. H.; Greenfield, E. A.; Chang,
[50305] T. T.; Sharpe, A. H.; Berry, G.; DeKruyff, R. H.; Umetsu, D. T.:
[50306] Antigen-specific regulatory T cells develop via the ICOS-ICOS-ligand
[50307] pathway and inhibit allergen-induced airway hyperreactivity. Nature
[50308] Med. 8: 1024-1032, 2002.

[50309] 7930.

[50310] Cox, D. R.; Burmeister, M.; Price, E. R.; Kim, S.; Myers, R. M.

[50311] : Radiation hybrid mapping: a somatic cell genetic method for constructing

[50312] high-resolution maps of mammalian chromosomes. Science 250: 245-250,

[50313] 1990.

[50314] 7931.

[50315] Crawley, J. B.; Williams, L. M.; Mander, T.; Brennan, F. M.; Foxwell,

[50316] B. M.: Interleukin-10 stimulation of phosphatidylinositol 3-kinase

[50317] and p70 S6 kinase is required for the proliferative but not the antiinflammatory

[50318] effects of the cytokine. J. Biol. Chem. 271: 16357-16362, 1996.

[50319] 7932.

[50320] Eskdale, J.; Gallagher, G.; Verweij, C. L.; Keijsers, V.; Westendorp,

[50321] R. G. J.; Huizinga, T. W. J.: Interleukin 10 secretion in relation

[50322] to human IL-10 locus haplotypes. Proc. Nat. Acad. Sci. 95: 9465-9470,
[50323] 1998.

[50324] 7933.

[50325] Eskdale, J.; Kube, D.; Tesch, H.; Gallagher, G.: Mapping of the
[50326] human IL10 gene and further characterization of the 5-prime flanking
[50327] sequence. Immunogenetics 46: 120-128, 1997.

[50328] 7934.

[50329] Farmer, M. A.; Sundberg, J. P.; Bristol, I. J.; Churchill, G. A.;

[50330] Li, R.; Elson, C. O.; Leiter, E. H.: A major quantitative trait locus
[50331] on chromosome 3 controls colitis severity in IL-10-deficient mice. Proc.
[50332] Nat. Acad. Sci. 98: 13820-13825, 2001.

[50333] 7935.

[50334] Franchimont, D.; Martens, H.; Hagelstein, M.-T.; Louis, E.; Dewe,
[50335] W.; Chrousos, G. P.; Belaiche, J.; Geenen, V.: Tumor necrosis factor

[50336] alpha decreases, and interleukin-10 increases, the sensitivity of
[50337] human monocytes to dexamethasone: potential regulation of the glucocorticoid
[50338] receptor. J. Clin. Endocr. Metab. 84: 2834-2839, 1999.

[50339] 7936.

[50340] Gesser, B.; Leffers, H.; Jinquan, T.; Vestergaard, C.; Kirstein,
[50341] N.; Sindet-Pedersen, S.; Jensen, S. L.; Thestrup-Pedersen, K.; Larsen,
[50342] C. G.: Identification of functional domains on human interleukin
[50343] 10. Proc. Nat. Acad. Sci. 94: 14620-14625, 1997.

[50344] 7937.

[50345] Gibson, A. W.; Edberg, J. C.; Wu, J.; Westendorp, R. G. J.; Huizinga,
[50346] T. W. J.; Kimberly, R. P.: Novel single nucleotide polymorphisms
[50347] in the distal IL-10 promoter affect IL-10 production and enhance the
[50348] risk of systemic lupus erythematosus. J. Immun. 166: 3915-3922,

[50349] 2001.

[50350] 7938.

[50351] Goudy, K.; Song, S.; Wasserfall, C.; Zhang, Y. C.; Kapturczak,

[50352] M.; Muir, A.; Powers, M.; Scott-Jorgensen, M.; Campbell-Thompson,

[50353] M.; Crawford, J. M.; Ellis, T. M.; Flotte, T. R.; Atkinson, M. A.

[50354] : Adeno-associated virus vector-mediated IL-10 gene delivery prevents

[50355] type 1 diabetes in NOD mice. Proc. Nat. Acad. Sci. 98: 13913-13918,

[50356] 2001.

[50357] 7939.

[50358] Grove, J.; Daly, A. K.; Bassendine, M. F.; Gilvarry, E.; Day,

[50359] C. P.: Interleukin 10 promoter region polymorphisms and susceptibility

[50360] to advanced alcoholic liver disease. Gut 46: 540-545, 2000.

[50361] 7940.

[50362] Kim, J. M.; Brannan, C. I.; Copeland, N. G.; Jenkins, N. A.;

Khan,
[50363] T. A.; Moore, K. W.: Structure of the mouse Il-10 gene and chromosomal
[50364] localization of the mouse and human genes. J. Immun. 148: 3618-3623,
[50365] 1992.

[50366] 7941.

[50367] Kitagawa, N.; Goto, M.; Kurozumi, K.; Maruo, S.; Fukayama, M.;
[50368] Naoe, T.; Yasukawa, M.; Hino, K.; Suzuki, T.; Todo, S.; Takada, K.
[50369] : Epstein-Barr virus-encoded poly(A)- RNA supports Burkitt's lymphoma
[50370] growth through interleukin-10 induction. EMBO J. 19: 6742-6750,
[50371] 2000.

[50372] 7942.

[50373] Kuhn, R.; Lohler, J.; Rennick, D.; Rajewsky, K.; Muller, K.:
[50374] Interleukin-10-deficient mice develop chronic enterocoli-tis. Cell 75:
[50375] 263-274, 1993.

[50376] 7943.

[50377] Lee, T.-S.; Chau, L.-Y.: Heme oxygenase-1 mediates the anti-inflammatory

[50378] effect of interleukin-10 in mice. Nature Med. 8: 240-246, 2002.

[50379] 7944.

[50380] Meng, X.; Sawamura, D.; Tamai, K.; Hanada, K.; Ishida, H.; Hashimoto,

[50381] I.: Keratinocyte gene therapy for systemic diseases: circulating

[50382] interleukin 10 released from gene-transferred keratinocytes inhibits

[50383] contact hypersensitivity at distant areas of the skin. J. Clin. Invest. 101:

[50384] 1462-1467, 1998.

[50385] 7945.

[50386] Pinderski Oslund, L. J.; Hedrick, C. C.; Olvera, T.; Hagenbaugh,

[50387] A.; Territo, M.; Berliner, J. A.; Fyfe, A. I.: Interleukin-10 blocks

[50388] atherosclerotic events in vitro and in vivo. Arterioscler. Thromb.

[50389] Vasc. Biol. 19: 2847-2853, 1999.

[50390] 7946.

[50391] Rosenwasser, L. J.; Borish, L.: Genetics of atopy and asthma:

[50392] the rationale behind promoter-based candidate gene studies (IL-4 and

[50393] IL-10). Am. J. Resp. Crit. Care Med. 156: S152-S155, 1997.

[50394] 7947.

[50395] Shin, H. D.; Winkler, C.; Stephens, J. C.; Bream, J.; Young, H.;

[50396] Goedert, J. J.; O'Brien, T. R.; Vlahov, D.; Buchbinder, S.; Giorgi,

[50397] J.; Rinaldo, C.; Donfield, S.; Willoughby, A.; O'Brien, S. J.; Smith,

[50398] M. W.: Genetic restriction of HIV-1 pathogenesis to AIDS by promoter

[50399] alleles of IL10. Proc. Nat. Acad. Sci. 97: 14467-14472, 2000.

[50400] 7948.

[50401] Terkeltaub, R. A.: IL-10: an 'immunologic scalpel' for atherosclerosis?

[50402] (Editorial) Atheroscler. Thromb. Vasc. Biol. 19:
2823-2825, 1999.

[50403] 7949.

[50404] Turner, D. M.; Williams, D. M.; Sankaran, D.; Lazarus, M.;
Sinnott,

[50405] P. J.; Hutchinson, I. V.: An investigation of polymorphism
in the

[50406] interleukin-10 gene promoter. Europ. J. Immunogenet.
24: 1-8, 1997.

[50407] 7950.

[50408] Vieira, P.; de Waal-Malefyt, R.; Dang, M.-N.; Johnson, K.
E.;

[50409] Kastelein, R.; Fiorentino, D. F.; deVries, J. E.; Roncarolo,
M.-G.;

[50410] Mosmann, T. R.; Moore, K. W.: Isolation and expression of
human cytokine

[50411] synthesis inhibitory factor cDNA clones: homology to Ep-
stein-Barr

[50412] virus open reading frame BCRFI. Proc. Nat. Acad. Sci. 88:
1172-1176,

[50413] 1991.

[50414] 7951.

[50415] Westendorp, R. G. J.; Langermans, J. A. M.; Huizinga, T. W. G.;

[50416] Elouali, A. H.; Boomsma, D. I.; Verweij, C. L.; Vanden-broucke, J.

[50417] P.: Genetic influence on cytokine production and fatal meningococcal

[50418] disease. Lancet 349: 170-173, 1997.

[50419] 7952.

[50420] Armstrong, E.; Cannizzaro, L.; Bergman, M.; Huebner, K.; Alitalo,

[50421] K.: The c-src tyrosine kinase (CSK) gene, a potential an-tioncogene,

[50422] localizes to human chromosome region 15q23-q25. Cy-togenet. Cell Genet. 60:

[50423] 119-120, 1992.

[50424] 7953.

[50425] Cloutier, J.-F.; Veillette, A.: Association of inhibitory tyro-sine

[50426] protein kinase p50(csk) with protein tyrosine phosphatase PEP in T

[50427] cells and other hemopoietic cells. EMBO J. 15: 4909-4918, 1996.

[50428] 7954.

[50429] Partanen, J.; Armstrong, E.; Bergman, M.; Makela, T. P.; Hirvonen,

[50430] H.; Huebner, K.; Alitalo, K.: Cyl encodes a putative cyto-plasmic

[50431] tyrosine kinase lacking the conserved tyrosine autophos-phorylation

[50432] site (Y416-src). Oncogene 6: 2013-2018, 1991.

[50433] 7955.

[50434] Morison, I. M.; Becroft, D. M.; Taniguchi, T.; Woods, C. G.; Reeve,

[50435] A. E.: Somatic overgrowth associated with overexpression of insulin-like

[50436] growth factor II. Nature Med. 2: 311-316, 1996.

[50437] 7956.

[50438] Porteous, D. J.; Bickmore, W.; Christie, S.; Boyd, P. A.; Cranston,

[50439] G.; Fletcher, J. M.; Gosden, J. R.; Rout, D.; Seawright, A.; Simola,

[50440] K. O. J.; van Heyningen, V.; Hastie, N. D.: HRAS1-selected chromosome

[50441] transfer generates markers that colocalize aniridia- and genitourinary
[50442] dysplasia-associated translocation breakpoints and the Wilms tumor
[50443] gene within band 11p13. Proc. Nat. Acad. Sci. 84: 5355-5359, 1987.
[50444] 7957.

[50445] van Heyningen, V.; Bickmore, W. A.; Seawright, A.; Fletcher, J.
[50446] M.; Maule, J.; Fekete, G.; Gessler, M.; Bruns, G. A. P.; Huerre-Jeanpierre,
[50447] C.; Junien, C.; Williams, B. R. G.; Hastie, N. D.: Role for the Wilms
[50448] tumor gene in genital development? Proc. Nat. Acad. Sci. 87: 5383-5386,
[50449] 1990.

[50450] 7958.

[50451] Webster, A. D. B.; Barnes, D. E.; Arlett, C. F.; Lehmann, A. R.;
[50452] Lindahl, T.: Growth retardation and immunodeficiency in a patient
[50453] with mutations in the DNA ligase I gene. Lancet 339:

1508-1509,

[50454] 1992.

[50455] 7959.

[50456] Webster, D.; Arlett, C. F.; Harcourt, S. A.; Teo, I.; Henderson,

[50457] L.: A new syndrome of immunodeficiency and increased cellular sensitivity

[50458] to DNA damaging agents.In: Bridges, B. A.; Harnden, D. G.: Ataxia-telangiectasia:

[50459] A Cellular and Molecular Link between Cancer, Neuropathology, and

[50460] Immune Deficiency. New York: John Wiley (pub.) 1982. Pp. 379-386.

[50461] 7960.

[50462] Ahuja, H. G.; Felix, C. A.; Aplan, P. D.: The t(11;20)(p15;q11)

[50463] chromosomal translocation associated with therapy-related myelodysplastic

[50464] syndrome results in an NUP98-TOP1 fusion. Blood 94: 3258-3261, 1999.

[50465] 7961.

[50466] Baumgartner, B.; Klett, C.; Hameister, H.; Richter, A.;

Knippers,
[50467] R.: Mouse genes encoding DNA topoisomerase I. Mammalian Genome 5:
[50468] 19-25, 1994.

[50469] 7962.

[50470] Chang, J.-Y.; Liu, J.-F.; Juang, S.-H.; Liu, T.-W.; Chen, L.-T.
[50471] : Novel mutation of topoisomerase I in rendering cells resistant to
[50472] camptothecin. Cancer Res. 62: 3716-3721, 2002.

[50473] 7963.

[50474] D'Arpa, P.; Machlin, P. S.; Ratrie, H., III; Rothfield, N. F.;

[50475] Cleveland, D. W.; Earnshaw, W. C.: cDNA cloning of human DNA topoisomerase
[50476] I: catalytic activity of a 67.7-kDa carboxyl-terminal fragment. Proc.
[50477] Nat. Acad. Sci. 85: 2543-2547, 1988.

[50478] 7964.

[50479] Juan, C.-C.; Hwang, J.; Liu, A. A.; Whang-Peng, J.; Knutsen, T.;
[50480] Huebner, K.; Croce, C. M.; Zhang, H.; Wang, J. C.; Liu, L. F.:

Human
[50481] DNA topoisomerase I is encoded by a single-copy gene that maps to
[50482] chromosome region 20q12-13.2. Proc. Nat. Acad. Sci. 85: 8910-8913,
[50483] 1988.

[50484] 7965.

[50485] Kunze, N.; Yang, G.; Dolberg, M.; Sundarp, R.; Knippers, R.; Richter,
[50486] A.: Structure of the human type I DNA topoisomerase gene. J. Biol.
[50487] Chem. 266: 9610-9616, 1991.

[50488] 7966.

[50489] Kunze, N.; Yang, G. C.; Jiang, Z. Y.; Hameister, H.; Adolph, S.;
[50490] Wiedorn, K.-H.; Richter, A.; Knippers, R.: Localization of the active
[50491] type I DNA topoisomerase gene on human chromosome 20q11.2-13.1, and
[50492] two pseudogenes on chromosomes 1q23-24 and 22q11.2-13.1. Hum. Genet. 84:
[50493] 6-10, 1989.

[50494] 7967.

[50495] Tamura, H.; Kohchi, C.; Yamada, R.; Ikeda, T.; Koiwai, O.; Patterson,

[50496] E.; Keene, J. D.; Okada, K.; Kjeldsen, E.; Nishikawa, K.; Andoh, T.

[50497] : Molecular cloning of a cDNA of a camptothecin-resistant human DNA

[50498] topoisomerase I and identification of mutation sites. Nucleic Acids

[50499] Res. 19: 69-75, 1991.

[50500] 7968.

[50501] Siciliano, M. J.; Bachinski, L.; Dolf, G.; Carrano, A. V.; Thompson,

[50502] L. H.: Chromosomal assignments of human DNA repair genes that complement

[50503] Chinese hamster ovary (CHO) cell mutants. (Abstract) Cytogenet. Cell

[50504] Genet. 46: 691-692, 1987.

[50505] 7969.

[50506] Thompson, L. H.; Carrano, A. V.; Sato, K.; Salazar, E. P.; White,

[50507] B. F.; Stewart, S. A.; Minkler, J. L.; Siciliano, M. J.: Identification
[50508] of nucleotide-excision-repair genes on human chromosomes 2 and 13
[50509] by functional complementation in hamster-human hybrids. Somat. Cell
[50510] Molec. Genet. 13: 539-551, 1987.

[50511] 7970.

[50512] Brandi, M. L.; Gagel, R. F.; Angeli, A.; Bilezikian, J. P.; Beck-Peccoz,
[50513] P.; Bordi, C.; Conte-Devolx, B.; Falchetti, A.; Gheri, R. G.; Libroia,
[50514] A.; Lips, C. J. M.; Lombardi, G.; and 12 others: Guidelines for
[50515] diagnosis and therapy of MEN type 1 and type 2. J. Clin. Endocr.
[50516] Metab. 86: 5658-5671, 2001.

[50517] 7971.

[50518] Buchta, R. M.; Kaplan, J. M.: Zollinger-Ellison syndrome in a
[50519] nine-year-old child: a case report and review of the entity in childhood. Pediatrics 47:

[50520] 594-598, 1971.

[50521] 7972.

[50522] Bystrom, C.; Larsson, C.; Blomberg, C.; Sandelin, K.; Falk-
mer,
[50523] U.; Skogseid, B.; Oberg, K.; Werner, S.; Nordenskjold, M.:
Localization
[50524] of the MEN1 gene to a small region within chromosome
11q13 by deletion
[50525] mapping in tumors. Proc. Nat. Acad. Sci. 87: 1968-1972,
1990.

[50526] 7973.

[50527] Carling, T.; Correa, P.; Hessman, O.; Hedberg, J.; Skogseid,
B.;
[50528] Lindberg, D.; Rastad, J.; Westin, G.; Akerstrom, G.:
Parathyroid
[50529] MEN1 gene mutations in relation to clinical characteristics
of nonfamilial
[50530] primary hyperparathyroidism. J. Clin. Endocr. Metab. 83:
2960-2963,
[50531] 1998.

[50532] 7974.

[50533] Cebrian, A.; Herrera-Pombo, J. L.; Diez, J. J.; Sanchez-Vi-
lar,
[50534] O.; Lara, J. I.; Vazquez, C.; Pico, A.; Osorio, A.; Martinez-
Delgado,
[50535] B.; Benitez, J.; Robledo, M.: Genetic and clinical analysis in
10
[50536] Spanish patients with multiple endocrine neoplasia type 1.
Europ.
[50537] J. Hum. Genet. 7: 585-589, 1999.

[50538] 7975.

[50539] Chedid, A.; Hoversland, R. C.; Chejfec, G.; McCune, S. A.;
Jurin,
[50540] R. R.; Evans, E. E.: Hereditary pituitary prolactinomas: a
new rat
[50541] model. Acta Endocr. 119: 535-542, 1988.

[50542] 7976.

[50543] Cocco, A. E.; Conway, S. J.: Zollinger-Ellison syndrome as-
sociated
[50544] with ovarian mucinous cystadenocarcinoma. New Eng. J.
Med. 293:
[50545] 485-486, 1975.

[50546] 7977.

[50547] Crabtree, J. S.; Scacheri, P. C.; Ward, J. M.; Garrett-Beal, L.;

[50548] Emmert-Buck, M. R.; Edgemon, K. A.; Lorang, D.; Libutti,
S. K.; Chandrasekharappa,

[50549] S. C.; Marx, S. J.; Spiegel, A. M.; Collins, F. S.: A mouse
model

[50550] of multiple endocrine neoplasia, type 1, develops multiple
endocrine

[50551] tumors. Proc. Nat. Acad. Sci. 98: 1118-1123, 2001.

[50552] 7978.

[50553] Cutler, R. E.; Reiss, E.; Ackerman, L. V.: Familial hyper-
parathyroidism:

[50554] a kindred involving eleven cases, with a discussion of pri-
mary chief-cell

[50555] hyperplasia. New Eng. J. Med. 270: 859-865, 1964.

[50556] 7979.

[50557] Chandrasekharappa, S. C.; Guru, S. C.; Manickam, P.;
Olufemi,

[50558] S.-E.; Collins, F. S.; Emmert-Buck, M. R.; Debelenko, L. V.;
Zhuang,

[50559] Z.; Lubensky, I. A.; Liotta, L. A.; Crabtree, J. S.; Wang, Y.;

Roe,

[50560] B. A.; Weisemann, J.; Boguski, M. S.; Agarwal, S. K.; Kester, M. B.;

[50561] Kim, Y. S.; Heppner, C.; Dong, Q.; Spiegel, A. M.; Burns, A. L.; Marx,

[50562] S. J.: Positional cloning of the gene for multiple endocrine neoplasia-type

[50563] 1. Science 276: 404-406, 1997.

[50564] 7980.

[50565] Darling, T. N.; Skarulis, M. C.; Steinberg, S. M.; Marx, S. J.;

[50566] Spiegel, A. M.; Turner, M.: Multiple facial angiofibromas and collagenomas

[50567] in patients with multiple endocrine neoplasia type 1. Arch. Derm. 133:

[50568] 853-857, 1997.

[50569] 7981.

[50570] Debelenko, L. V.; Brambilla, E.; Agarwal, S. K.; Swalwell, J.

[50571] I.; Kester, M. B.; Lubensky, I. A.; Zhuang, Z.; Guru, S. C.; Manickam,

[50572] P.; Olufemi, S.-E.; Chandrasekharappa, S. C.; Crabtree, J. S.; Kim,

[50573] Y. S.; Heppner, C.; Burns, A. L.; Spiegel, A. M.; Marx, S. J.; Liotta,
[50574] L. A.; Collins, F. S.; Travis, W. D.; Emmert-Buck, M. R.: Identification
[50575] of MEN1 gene mutations in sporadic carcinoid tumors of the lung. Hum.
[50576] Molec. Genet. 6: 2285-2290, 1997.

[50577] 7982.

[50578] Deveney, C. W.; Deveney, K. S.; Way, L. W.: The Zollinger-El-lison
[50579] syndrome--23 years later. Ann. Surg. 188: 384-391, 1978.

[50580] 7983.

[50581] Ellison, E. H.; Wilson, S. D.: The Zollinger-Ellison syn-drome.
[50582] Re-appraisal and evaluation of 260 registered cases. Ann. Surg. 160:
[50583] 512-530, 1964.

[50584] 7984.

[50585] Ellison, E. H.; Wilson, S. D.: The Zollinger-Ellison syn-drome

[50586] updated. Surg. Clin. N. Am. 47: 1115-1124, 1967.

[50587] 7985.

[50588] Farid, N. R.: Personal Communication. St. John's, New-
foundland,
[50589] Canada 7/28/1980.

[50590] 7986.

[50591] Farid, N. R.: Personal Communication. Riyadh, Saudi Ara-
bia
[50592] 1/11/1994.

[50593] 7987.

[50594] Farid, N. R.; Buehler, S.; Russell, N. A.; Maroun, F. B.;
Allerdice,
[50595] P.; Smyth, H. S.: Prolactinomas in familial multiple en-
docrine neoplasia
[50596] syndrome type I: relationship to HLA and carcinoid tu-
mors. Am. J.
[50597] Med. 69: 874-880, 1980.

[50598] 7988.

[50599] Farnebo, F.; Teh, B. T.; Kytola, S.; Svensson, A.; Phelan, C.;

[50600] Sandelin, K.; Thompson, N. W.; Hoog, A.; Weber, G.;

Farnebo, L.-O.;

[50601] Larsson, C.: Alterations of the MEN1 gene in sporadic parathyroid

[50602] tumors. J. Clin. Endocr. Metab. 83: 2627-2630, 1998.

[50603] 7989.

[50604] Friesen, S. R.; Schimke, R. N.; Pearse, A. G. E.: Genetic as-pects

[50605] of the Z-E syndrome: prospective studies in two kindred: antral gastrin

[50606] cell hyperplasia. Ann. Surg. 176: 370-383, 1972.

[50607] 7990.

[50608] Fujimori, M.: Personal Communication. Matsumoto, Japan 1/5/1999.

[50609] 7991.

[50610] Fujimori, M.; Shirahama, S.; Sakurai, A.; Hashizume, K.; Hama,

[50611] Y.; Ito, K.; Shingu, K.; Kobayashi, S.; Amano, J.; Fukushima, Y.:

[50612] Novel V184E MEN1 germline mutation in a Japanese kin-dred with familial

[50613] hyperparathyroidism. Am. J. Med. Genet. 80: 221-222,

1998.

[50614] 7992.

[50615] Fujimori, M.; Wells, S. A., Jr.; Nakamura, Y.: Fine-scale mapping

[50616] of the gene responsible for multiple endocrine neoplasia type 1 (MEN1). Am.

[50617] J. Hum. Genet. 50: 399-403, 1992.

[50618] 7993.

[50619] Friedman, E.; Sakaguchi, K.; Bale, A. E.; Falchetti, A.; Streeten,

[50620] E.; Zimering, M. B.; Weinstein, L. S.; McBride, W. O.; Naka-mura, Y.;

[50621] Brandi, M.-L.; Norton, J. A.; Aurbach, G. D.; Spiegel, A. M.; Marx,

[50622] S. J.: Clonality of parathyroid tumors in familial multiple endocrine

[50623] neoplasia type 1. New Eng. J. Med. 321: 213-218, 1989. Note: Erratum:

[50624] New Eng. J. Med. 321: 1057 only, 1989...

[50625] 7994.

[50626] Gaitan, D.; Loosen, P. T.; Orth, D. N.: Two patients with

Cushing's
[50627] disease in a kindred with multiple endocrine neoplasia type I. J.
[50628] Clin. Endocr. Metab. 76: 1580-1582, 1993.

[50629] 7995.

[50630] Giraud, S.; Zhang, C. X.; Serova-Sinilnikova, O.; Wautot, V.;

[50631] Salandre, J.; Buisson, N.; Waterlot, C.; Bauters, C.; Porchet, N.;
[50632] Aubert, J.-P.; Emy, P.; Cadiot, G.; and 29 others: Germline mutation
[50633] analysis in patients with multiple endocrine neoplasia type 1 and
[50634] related disorders. Am. J. Hum. Genet. 63: 455-467, 1998.

[50635] 7996.

[50636] Gisselsson, D.; Hoglund, M.; Mertens, F.; Del Cin, P.; Mandahl,
[50637] N.: Hibernomas are characterized by homozygous deletions in the multiple
[50638] endocrine neoplasia type I region: metaphase fluorescence in situ
[50639] hybridization reveals complex rearrangements not de-

tected by conventional
[50640] cytogenetics. Am. J. Path. 155: 61-66, 1999.

[50641] 7997.

[50642] Olson, S.; Wang, M. G.; Carafoli, E.; Strehler, E. E.; McBride,

[50643] O. W.: Localization of two genes encoding plasma mem-
brane Ca(2+)-transporting
[50644] ATPases to human chromosomes 1q25-32 and
12q21-23. Genomics 9:
[50645] 629-641, 1991.

[50646] 7998.

[50647] Brandt, P.; Ibrahim, E.; Bruns, G. A. P.; Neve, R. L.: Deter-
mination
[50648] of the nucleotide sequence and chromosomal localization
of the ATP2B2
[50649] gene encoding human Ca(2+)-pumping ATPase isoform
PMCA2. Genomics 14:
[50650] 484-487, 1992.

[50651] 7999.

[50652] Kozel, P. J.; Friedman, R. A.; Erway, L. C.; Yamoah, E. N.;
Liu,
[50653] L. H.; Riddle, T.; Duffy, J. J.; Doetschman, T.; Miller, M. L.;

Cardell,
[50654] E. L.; Shull, G. E.: Balance and hearing deficits in mice with a
[50655] null mutation in the gene encoding plasma membrane Ca(2+)-ATPase isoform
[50656] 2. J. Biol. Chem. 273: 18693-18696, 1998.

[50657] 8000.

[50658] Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.; Crossey, P.

[50659] A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M. I.; Zbar,
[50660] B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von Hippel-Lindau
[50661] disease tumour suppressor gene: identification of germline deletions
[50662] by pulsed field gel electrophoresis. Hum. Molec. Genet. 2: 879-882,
[50663] 1993.

[50664] 8001.

[50665] Street, V. A.; McKee-Johnson, J. W.; Fonseca, R. C.; Tempel, B.
[50666] L.; Noben-Trauth, K.: Mutations in a plasma membrane

Ca(2+)-ATPase

[50667] gene cause deafness in deafwaddler mice. Nature Genet. 19: 390-394,

[50668] 1998.

[50669] 8002.

[50670] Wang, M. G.; Yi, H.; Hilfiker, H.; Carafoli, E.; Strehler, E. E.;

[50671] McBride, O. W.: Localization of two genes encoding plasma membrane

[50672] Ca(2+)-ATPases isoforms 2 (ATP2B2) and 3 (ATP2B3) to human chromosomes

[50673] 3p26-p25 and Xq28, respectively. Cytogenet. Cell Genet. 67: 41-45,

[50674] 1994.

[50675] 8003.

[50676] Shi, G.-P.; Munger, J. S.; Meara, J. P.; Rich, D. H.; Chapman,

[50677] H. A.: Molecular cloning and expression of human alveolar macrophage

[50678] cathepsin S, an elastinolytic cysteine protease. J. Biol. Chem. 267:

[50679] 7258-7262, 1992.

[50680] 8004.

[50681] Shi, G.-P.; Webb, A. C.; Foster, K. E.; Knoll, J. H. M.; Lemere,

[50682] C. A.; Munger, J. S.; Chapman, H. A.: Human cathepsin S: chromosomal

[50683] localization, gene structure, and tissue distribution. J. Biol. Chem. 269:

[50684] 11530-11536, 1994.

[50685] 8005.

[50686] Berube, D.; Luu-The, V.; Simard, J.; Gagne, R.; Labrie, F.: Localization

[50687] of the beta-estradiol dehydrogenase genes to q11-q12 of chromosome

[50688] 17. (Abstract) Cytogenet. Cell Genet. 51: 962 only, 1989.

[50689] 8006.

[50690] Geissler, W. M.; Davis, D. L.; Wu, L.; Bradshaw, K. D.; Patel,

[50691] S.; Mendonca, B. B.; Elliston, K. O.; Wilson, J. D.; Russell, D. W.;

[50692] Andersson, S.: Male pseudohermaphroditism caused by mutations of

[50693] testicular 17-beta-hydroxysteroid dehydrogenase 3. Na-

ture Genet. 7:
[50694] 34-39, 1994.

[50695] 8007.

[50696] Green, V. L.; Speirs, V.; Landolt, A. M., Foy, P. M.; Atkin, S.

[50697] L.: 17-beta-hydroxysteroid dehydrogenase type 1, 2, 3, and 4 expression
[50698] and enzyme activity in human anterior pituitary adenomas. J. Clin.
[50699] Endocr. Metab. 84: 1340-1345, 1999.

[50700] 8008.

[50701] Harkness, R. A.; Thistlethwaite, D.; Darling, J. A. B.; Skakkebaek,
[50702] N. E.; Corker, C. S.: Neutral 17-beta-hydroxysteroid oxidoreductase
[50703] deficiency in testes causing male pseudohermaphroditism in an infant. J.
[50704] Inherit. Metab. Dis. 2: 51-54, 1979.

[50705] 8009.

[50706] Huang, Y.-W.; Pineau, I.; Chang, H.-J.; Azzi, A.; Bellemare, V.;
[50707] Laberge, S.; Lin, S.-X.: Critical residues for the specificity

of

[50708] cofactors and substrates in human estrogenic 17-beta-hydroxysteroid

[50709] dehydrogenase 1: variants designed from the three-dimensional structure

[50710] of the enzyme. Molec. Endocr. 15: 2010-2020, 2001.

[50711] 8010.

[50712] Luu-The, V.; Labrie, C.; Simard, J.; Lachance, Y.; Zhao, H.-F.;

[50713] Couet, J.; Leblanc, G.; Labrie, F.: Structure of two in tan-dem human

[50714] 17-beta-hydroxysteroid dehydrogenase genes. Molec. Endocr. 4: 268-275,

[50715] 1990.

[50716] 8011.

[50717] Migeon, C. J.: Personal Communication. Baltimore, Md. 10/29/1990.

[50718] 8012.

[50719] Normand, T.; Narod, S.; Labrie, F.; Simard, J.: Detection of polymorphisms

[50720] in the estradiol 17-beta-hydroxysteroid dehydrogenase II

gene at the
[50721] EDH17B2 locus on 17q11-q21. Hum. Molec. Genet. 2: 479-483, 1993.
[50722] 8013.
[50723] Russell, D. W.: Personal Communication. Dallas, Texas 5/9/1994.
[50724] 8014.
[50725] Tremblay, Y.; Ringler, G. E.; Morel, Y.; Mohandas, T. K.; Labrie,
[50726] F.; Strauss, J. F., III; Miller, W. L.: Regulation of the gene for
[50727] estrogenic 17-ketosteroid reductase lying on chromosome 17cen-q25. J.
[50728] Biol. Chem. 264: 20458-20462, 1989.
[50729] 8015.
[50730] Winqvist, R.; Peltoketo, H.; Isomaa, V.; Grzeschik, K. H.; Mannermaa,
[50731] A.; Vihko, R.: The gene for 17-beta-hydroxysteroid dehydrogenase
[50732] maps to human chromosome 17, bands q12-q21, and shows an RFLP with
[50733] Scal. Hum. Genet. 85: 473-476, 1990.

[50734] 8016.

[50735] Casey, M. L.; MacDonald, P. C.; Andersson, S.: 17-beta-hydroxysteroid
[50736] dehydrogenase type 2: chromosomal assignment and progestin regulation
[50737] of gene expression in human endometrium. J. Clin. Invest. 94: 2135-2141,
[50738] 1994.

[50739] 8017.

[50740] Durocher, F.; Morissette, J.; Labrie, Y.; Labrie, F.; Simard, J.

[50741] : Mapping of the HSD17B2 gene encoding type II 17-beta-hydroxysteroid
[50742] dehydrogenase close to D16S422 on chromosome 16q24.1-q24.2. Genomics 25:
[50743] 724-726, 1995.

[50744] 8018.

[50745] Bouillaud, F.; Villarroya, F.; Hentz, E.; Raimbault, S.; Cas-sard,
[50746] A.-M.; Ricquier, D.: Detection of brown adipose tissue un-coupling
[50747] protein mRNA in adult patients by a human genomic

probe. Clin. Sci. 75:
[50748] 21-27, 1988.

[50749] 8019.

[50750] Bouillaud, F.; Weissenbach, J.; Ricquier, D.: Complete cDNA-derived
[50751] amino acid sequence of rat brown fat uncoupling protein. J. Biol.
[50752] Chem. 261: 1487-1491, 1986.

[50753] 8020.

[50754] Cassard, A. M.; Bouillaud, F.; Mattei, M. G.; Hentz, E.; Raimbault,
[50755] S.; Thomas, M.; Ricquier, D.: Human uncoupling protein gene: structure,
[50756] comparison with rat gene, and assignment to the long arm of chromosome 8021.
[50757] Echtay, K. S.; Roussel, D.; St-Pierre, J.; Jekabsons, M. B.; Cadenas,
[50758] S.; Stuart, J. A.; Harper, J. A.; Roebuck, S. J.; Morrison, A.; Pickering,
[50759] S.; Clapham, J. C.; Brand, M. D.: Superoxide activates mitochondrial
[50760] uncoupling proteins. Nature 415: 96-99, 2002.

[50761] 8022.

[50762] Echtay, K. S.; Winkler, E.; Klingenberg, M.: Coenzyme Q is an

[50763] obligatory cofactor for uncoupling protein function. Nature 408:

[50764] 609-613, 2000.

[50765] 8023.

[50766] J. Cell. Biochem. 43: 255-264, 1990.

[50767]

[50768] 4. Clement, K.; Ruiz, J.; Cassard-Doulcier, A. M.; Bouillaud, F.;

[50769] Ricquier, D.; Basdevant, A.; Guy-Grand, B.; Froguel, P.: Additive

[50770] effect of A-G (-3826) variant of the uncoupling protein gene and the

[50771] trp64arg mutation of the beta 3-adrenergic receptor gene on weight

[50772] gain in morbid obesity. Int. J. Obes. Relat. Metab. Disord. 20:

[50773] 1062-1066, 1996.

[50774] 8024.

[50775] Enerback, S.; Jacobsson, A.; Simpson, E. M.; Guerra, C.; Yamashita,
[50776] H.; Harper, M.-E.; Kozak, L. P.: Mice lacking mitochondrial uncoupling
[50777] protein are cold-sensitive but not obese. Nature 387: 90-93, 1997.
[50778] 8025.

[50779] Fletcher, C.; Norman, D. J.; Germond, E.; Heintz, N.: A multilocus
[50780] linkage map of mouse chromosome 8. Genomics 9: 737-741, 1991.
[50781] 8026.

[50782] Jacobson, A.; Stadler, U.; Glotzer, M. A.; Kozak, L. P.: Mi-tochondrial
[50783] uncoupling protein from mouse brown fat: molecular cloning, genetic
[50784] mapping and mRNA expression. J. Biol. Chem. 260: 16250-16254, 1985.
[50785] 8027.

[50786] Lowell, B. B.; S-Susulic, V.; Hamann, A.; Lawitts, J. A.; Himms-Hagen,
[50787] J.; Boyer, B. B.; Kozak, L. P.; Flier, J. S.: Development of

obesity
[50788] in transgenic mice after genetic ablation of brown adipose tissue. Nature 366:
[50789] 740-742, 1993.

[50790] 8028.

[50791] Oppert, J. M.; Vohl, M. C.; Chagnon, M.; Dionne, F. T.; Cassard-Doulcier,
[50792] A. M.; Ricquier, D.; Perusse, L.; Bouchard, C.: DNA poly-morphism
[50793] in the uncoupling protein (UCP) gene and human body fat. Int. J.
[50794] Obes. Relat. Metab. Disord. 18: 526-531, 1994.

[50795] 8029.

[50796] Ridley, R. G.; Patel, H. V.; Gerber, G. E.; Morton, R. C.; Freeman,
[50797] K. B.: Complete nucleotide and derived amino acid se-quence of cDNA
[50798] encoding the mitochondrial uncoupling protein of rat brown adipose
[50799] tissue: lack of mitochondrial targeting presequence. Nu-cleic Acids
[50800] Res. 14: 4025-4035, 1986.

[50801] 8030.

[50802] Urhammer, S. A.; Fridberg, M.; Sorensen, T. I.; Echwald, S. M.;

[50803] Andersen, T.; Tybjaerg-Hansen, A.; Clausen, J. O.; Pedersen, O.:

[50804] Studies of genetic variability of the uncoupling protein 1 gene in

[50805] Caucasian subjects with juvenile-onset obesity. J. Clin. Endocr.

[50806] Metab. 82: 4069-4074, 1997.

[50807] 8031.

[50808] Reich, C. B.; Franklin, E. C.: Familial Mediterranean fever in

[50809] an Italian family. Arch. Intern. Med. 125: 337-340, 1970.

[50810] 8032.

[50811] Di Marzo, V.; Goparaju, S. K.; Wang, L.; Liu, J.; Batkai, S.; Jarai,

[50812] Z.; Fezza, F.; Miura, G. I.; Palmiter, R. D.; Sugiura, T.; Kunos,

[50813] G.: Leptin-regulated endocannabinoids are involved in maintaining

[50814] food intake. Nature 410: 822-825, 2001.

[50815] 8033.

[50816] Gerard, C. M.; Mollereau, C.; Vassart, G.; Parmentier, M.: Molecular

[50817] cloning of a human cannabinoid receptor which is also expressed in

[50818] testis. Biochem. J. 279: 129-134, 1991.

[50819] 8034.

[50820] Hoehe, M. R.; Caenazzo, L.; Martinez, M. M.; Hsieh, W.-T.; Modi,

[50821] W. S.; Gershon, E. S.; Bonner, T. I.: Genetic and physical mapping

[50822] of the human cannabinoid receptor gene to chromosome 6q14-q15. New

[50823] Biologist 3: 880-885, 1991.

[50824] 8035.

[50825] Ledent, C.; Valverde, O.; Cossu, G.; Petitet, F.; Aubert, J.-F.;

[50826] Beslot, F.; Bohme, G. A.; Imperato, A.; Pedrazzini, T.; Roques, B.

[50827] P.; Vassart, G.; Fratta, W.; Parmentier, M.: Unresponsive-ness to

[50828] cannabinoids and reduced addictive effects of opiates in CB(1) receptor

[50829] knockout mice. Science 283 401-404, 1999.

[50830] 8036.

[50831] Marsicano, G.; Wotjak, C. T.; Azad, S. C.; Bisogno, T.; Rammes,

[50832] G.; Cascio, M. G.; Hermann, H.; Tang, J.; Hofmann, C.; Zieglgansberger,

[50833] W.; Di Marzo, V.; Lutz, B.: The endogenous cannabinoid system controls

[50834] extinction of aversive memories. Nature 418: 530-534, 2002.

[50835] 8037.

[50836] Matsuda, L. A.; Lolait, S. J.; Brownstein, M. J.; Young, A. C.;

[50837] Bonner, T. I.: Structure of a cannabinoid receptor and functional

[50838] expression of the cloned cDNA. Nature 346: 561-564, 1990.

[50839] 8038.

[50840] Modi, W. S.; Bonner, T. I.: Localization of the cannabanoid (sic)

[50841] receptor locus using non-isotopic in situ hybridization. (Abstract) Cytogenet.

[50842] Cell Genet. 58: 1915 only, 1991.

[50843] 8039.

[50844] Panikashvili, D.; Simeonidou, C.; Ben-Shabat, S.; Hanus, L.; Breuer,

[50845] A.; Mechoulam, R.; Shohami, E.: An endogenous cannabi-noid (2-AG)

[50846] is neuroprotective after brain injury. Nature 413: 527-531, 2001.

[50847] 8040.

[50848] Beechey, C.; Tweedie, S.; Spurr, N.; Ball, S.; Peters, J.; Edwards,

[50849] Y.: Mapping of mouse carbonic anhydrase-3, Car-3: another locus in

[50850] the homologous region of mouse chromosome 3 and human chromosome 8. Genomics 6:

[50851] 692-696, 1990.

[50852] 8041.

[50853] Carter, N.; Jeffery, S.; Shiels, A.; Edwards, Y.; Tipler, T.; Hopkinson,

[50854] D. A.: Characterization of human carbonic anhydrase III from skeletal
[50855] muscle. Biochem. Genet. 17: 837-854, 1979.

[50856] 8042.

[50857] Edwards, Y. H.; Lloyd, J.; Parkar, M.; Povey, S.: Human muscle
[50858] specific carbonic anhydrase, CA3, is on chromosome 8. (Abstract) Cytogenet.
[50859] Cell Genet. 40: 621 only, 1985.

[50860] 8043.

[50861] Edwards, Y. H.; Lloyd, J. C.; Parkar, M.; Povey, S.: The gene
[50862] for human muscle specific carbonic anhydrase (CAIII) is assigned to
[50863] chromosome 8. Ann. Hum. Genet. 50: 41-47, 1986.

[50864] 8044.

[50865] Heath, R.; Carter, N. D.; Jeffery, S.; Edwards, R. J.; Watts, D.
[50866] C.; Watts, R. L.: Evaluation of carrier detection of Duchenne muscular
[50867] dystrophy using carbonic anhydrase III and creatine ki- nase. Am. J.

[50868] Med. Genet. 21: 291-296, 1985.

[50869] 8045.

[50870] Lloyd, J.; Brownson, C.; Tweedie, S.; Charlton, J.; Edwards, Y.

[50871] H.: Human muscle carbonic anhydrase: gene structure and DNA methylation

[50872] patterns in fetal and adult tissues. Genes Dev. 1: 594-602, 1987.

[50873] 8046.

[50874] Lloyd, J.; McMillan, S.; Hopkinson, D.; Edwards, Y. H.: Nucleotide

[50875] sequence and derived amino acid sequence of a cDNA encoding human

[50876] muscle carbonic anhydrase. Gene 41: 233-239, 1986.

[50877] 8047.

[50878] Lloyd, J. C.; Isenberg, H.; Hopkinson, D. A.; Edwards, Y. H.:

[50879] Isolation of a cDNA clone for the human muscle specific carbonic anhydrase,

[50880] CA III. Ann. Hum. Genet. 49: 241-251, 1985.

[50881] 8048.

[50882] Wade, R.; Gunning, P.; Eddy, R.; Shows, T.; Kedes, L.: Nucleotide
[50883] sequence, tissue-specific expression, and chromosome location of human
[50884] carbonic anhydrase III: the human CAIII gene is located on the same
[50885] chromosome as the closely linked CAI and CAII genes. Proc. Nat. Acad.
[50886] Sci. 83: 9571-9575, 1986.

[50887] 8049.

[50888] Ueki, Y.; Tiziani, V.; Santanna, C.; Fukai, N.; Maulik, C.; Garfinkle,
[50889] J.; Ninomiya, C.; doAmaral, C.; Peters, H.; Habal, M.; Rhee-Morris,
[50890] L.; Doss, J. B.; Kreiborg, S.; Olsen, B. R.; Reichenberger, E.: Mutations
[50891] in the gene encoding c-Abl-binding protein SH3BP2 cause cherubism. Nature
[50892] Genet. 28: 125-126, 2001.

[50893] 8050.

[50894] Hall, C.; Monfries, C.; Smith, P.; Lim, H. H.; Kozma, R.; Ahmed,

[50895] S.; Vanniasingham, V.; Leung, T.; Lim, L.: Novel human brain cDNA

[50896] encoding a 34,000 M(r) protein n-chimaerin, related to both the regulatory

[50897] domain of protein kinase C and BCR, the product of the breakpoint

[50898] cluster region gene. J. Molec. Biol. 211: 11-16, 1990.

[50899] 8051.

[50900] Hall, C.; Sin, W. C.; Teo, M.; Michael, G. J.; Smith, P.; Dong,

[50901] J. M.; Lim, H. H.; Manser, E.; Spurr, N. K.; Jones, T. A.; Lim, L.

[50902] : Alpha-2-chimerin, an SH2-containing GTPase-activating protein for

[50903] the ras-related protein p21-rac derived by alternate splicing of the

[50904] human n-chimerin gene, is selectively expressed in brain regions and

[50905] testes. Molec. Cell. Biol. 13: 4986-4998, 1993.

[50906] 8052.

[50907] Brewer, C.; Holloway, S.; Zawalnyski, P.; Schinzel, A.; Fitz-Patrick,

[50908] D.: A chromosomal deletion map of human malformations. Am. J. Hum.

[50909] Genet. 63: 1153-1159, 1998.

[50910] 8053.

[50911] Brewer, C. M.; Leek, J. P.; Green, A. J.; Holloway, S.; Bonthron,

[50912] D. T.; Markham, A. F.; FitzPatrick, D. R.: A locus for isolated cleft

[50913] palate, located on human chromosome 2q32. Am. J. Hum. Genet. 65:

[50914] 387-396, 1999.

[50915] 8054.

[50916] Carter, C. O.; Evans, K.; Coffey, R.; Roberts, J. A. F.; Buck,

[50917] A.; Roberts, M. F.: A family study of isolated cleft palate. J.

[50918] Med. Genet. 19: 329-331, 1982.

[50919] 8055.

[50920] Christensen, K.; Holm, N. V.; Olsen, J.; Kock, K.; Fogh-Andersen,

[50921] P.: Selection bias in genetic-epidemiological studies of cleft lip

[50922] and palate. Am. J. Hum. Genet. 51: 654-659, 1992.

[50923] 8056.

[50924] Christensen, K.; Mitchell, L. E.: Familial recurrence-pattern

[50925] analysis of nonsyndromic isolated cleft palate: a Danish registry

[50926] study. Am. J. Hum. Genet. 58: 182-190, 1996.

[50927] 8057.

[50928] Jenkins, M.; Stady, C.: Dominant inheritance of cleft of the soft

[50929] palate. Hum. Genet. 53: 341-342, 1980.

[50930] 8058.

[50931] Shields, E. D.; Bixler, D.; Fogh-Andersen, P.: Cleft palate:

[50932] a genetic and epidemiologic investigation. Clin. Genet. 20: 13-24,

[50933] 1981.

[50934] 8059.

[50935] Hwang, S. J.; Beaty, T. H.; Panny, S. R.; Street, N. A.; Joseph,

[50936] J. M.; Gordon, S.; McIntosh, I.; Francomano, C. A.: Associ-

ation study
[50937] of transforming growth factor alpha (TGF-alpha) TaqI polymorphism
[50938] and oral clefts: indication of gene-environment interaction in a population-based
[50939] sample of infants with birth defects. Am. J. Epidemiol. 141: 629-636,
[50940] 1995.

[50941] 8060.

[50942] Van Dyke, D. C.; Goldman, A. S.; Spielman, R. S.; Zmijew-
[50943] ski, C.
M.: Segregation of HLA in families with oral clefts: evi-
[50944] dence against
linkage between isolated cleft palate and HLA. Am. J. Med. Genet. 15:
[50945] 85-88, 1983.

[50946] 8061.

[50947] Czarny-Ratajczak, M.; Lohiniva, J.; Rogala, P.; Kozlowski, K.;
[50948] Perala, M.; Carter, L.; Spector, T. D.; Kolodziej, L.; Seppa-
[50949] nen, U.;
Glazar, R.; Krolewski, J.; Latos-Bielenska, A.; Ala-Kokko,

L.: A
[50950] mutation in COL9A1 causes multiple epiphyseal dysplasia: further evidence
[50951] for locus heterogeneity. Am. J. Hum. Genet. 69: 969-980, 2001.
[50952] 8062.

[50953] Eyre, D. R.; Apon, S.; Wu, J.-J.; Ericsson, L. H.; Walsh, K. A.

[50954] : Collagen type IX: evidence for covalent linkages to type II collagen
[50955] in cartilage. FEBS Lett. 220: 337-341, 1987.

[50956] 8063.

[50957] Fassler, R.; Schnegelsberg, P. N. J.; Dausman, J.; Shinya, T.;
[50958] Muragaki, Y.; McCarthy, M. T.; Olsen, B. R.; Jaenisch, R.: Mice lacking
[50959] alpha 1 (IX) collagen develop noninflammatory degenera- tive joint disease Proc.
[50960] Nat. Acad. Sci. 91: 5070-5074, 1994.

[50961] 8064.

[50962] Hagg, R.; Hedbom, E.; Mollers, U.; Aszodi, A.; Fassler, R.; Bruckner,

[50963] P.: Absence of the alpha-1(IX) chain leads to a functional knock-out

[50964] of the entire collagen IX protein in mice. J. Biol. Chem. 272: 20650-20654,

[50965] 1997.

[50966] 8065.

[50967] Jacenko, O.; Olsen, B. R.; Warman, M. L.: Of mice and men: heritable

[50968] skeletal disorders. (Editorial) Am. J. Hum. Genet. 54: 163-168,

[50969] 1994.

[50970] 8066.

[50971] Kimura, T.; Mattei, M.-G.; Stevens, J. W.; Goldring, M. B.; Ninomiya,

[50972] Y.; Olsen, B. R.: Molecular cloning of rat and human type IX collagen

[50973] cDNA and localization of the alpha1(IX) gene on the human chromosome

[50974] 6. Europ. J. Biochem. 179: 71-78, 1989.

[50975] 8067.

[50976] Mayne, R.; van der Rest, M.; Ninomiya, Y.; Olsen, B. R.: The structure

[50977] of type IX collagen. Ann. N.Y. Acad. Sci. 460: 38-46, 1985.

[50978] 8068.

[50979] McCormick, D.; van der Rest, M.; Goodship, J.; Lozano, G.;

Ninomiya,
[50980] T.; Olsen, B. R.: Structure of the glycosaminoglycan do-
main in the
[50981] type IX collagen-proteoglycan. Proc. Nat. Acad. Sci. 84:
4044-4048,
[50982] 1987.

[50983] 8069.

[50984] Muragaki, Y.; Nishimura, I.; Henney, A.; Ninomiya, Y.;
Olsen,
[50985] B. R.: The alpha-1(IX) collagen gene gives rise to two dif-
ferent
[50986] transcripts in both mouse embryonic and human fetal
RNA. Proc. Nat.
[50987] Acad. Sci. 87: 2400-2404, 1990.

[50988] 8070.

[50989] Nakata, K.; Ono, K.; Miyazaki, J.; Olsen, B. R.; Muragaki, Y.;

[50990] Adachi, E.; Yamamura, K.; Kimura, T.: Osteoarthritis asso-
ciated with
[50991] mild chondrodysplasia in transgenic mice expressing al-
pha-1(IX) collagen
[50992] chains with a central deletion. Proc. Nat. Acad. Sci. 90:

2870-2874,

[50993] 1993.

[50994] 8071.

[50995] Warman, M. L.; Tiller, G. E.; Polumbo, P. A.; Seldin, M. F.; Rochelle,

[50996] J. M.; Knoll, J. H. M.; Cheng, S.-D.; Olsen, B. R.: Physical and

[50997] linkage mapping of the human and murine genes for the alpha-1 chain

[50998] of type IX collagen (COL9A1). Genomics 17: 694-698, 1993.

[50999] 8072.

[51000] Burrows, N. P.; Nicholls, A. C.; Richards, A. J.; Luccarini, C.;

[51001] Harrison, J. B.; Yates, J. R. W.; Pope, F. M.: A point muta-tion in

[51002] an intronic branch site results in aberrant splicing of COL5A1 and

[51003] in Ehlers-Danlos syndrome type II in two British families. Am. J.

[51004] Hum. Genet. 63: 390-398, 1998.

[51005] 8073.

[51006] Caridi, G.; Pezzolo, A.; Bertelli, R.; Gimelli, G.; Di Donato,
[51007] A.; Candiano, G.; Ghiggeri, G. M.: Mapping of the human COL5A1 gene
[51008] to chromosome 9q34.3. Hum. Genet. 90: 174-176, 1992.
[51009] 8074.
[51010] Fichard, A.; Kleman, J.-P.; Ruggiero, F.: Another look at collagen
[51011] V and XI molecules. Matrix Biol. 14: 515-531, 1994.
[51012] 8075.
[51013] Giunta, C.; Steinmann, B.: Compound heterozygosity for a disease-causing
[51014] G1489D and disease-modifying G530S substitution in COL5A1 of a patient
[51015] with the classical type of Ehlers-Danlos syndrome: an explanation
[51016] of intrafamilial variability? Am. J. Med. Genet. 90: 72-79, 2000.
[51017] 8076.
[51018] Greenspan, D. S.; Byers, M. G.; Eddy, R. L.; Cheng, W.; Jani-Sait,
[51019] S.; Shows, T. B.: Human collagen gene COL5A1 maps to

the q34.2-q34.3
[51020] region of chromosome 9, near the locus for nail-patella syndrome. Genomics 12:
[51021] 836-837, 1992.

[51022] 8077.

[51023] Hamann, K. J.; Ten, R. M.; Loegering, D. A.; Jenkins, R. B.; Heise,
[51024] M. T.; Schad, C. R.; Pease, L. R.; Gleich, G. J.; Barker, R. L.:
[51025] Structure and chromosome localization of the human eosinophil-derived
[51026] neurotoxin and eosinophil cationic protein genes: evi-dence for intronless
[51027] coding sequences in the ribonuclease gene superfamily. Genomics 7:
[51028] 535-546, 1990.

[51029] 8078.

[51030] Mastrianni, D. M.; Eddy, R. L.; Rosenberg, H. F.; Corrette, S.
[51031] E.; Shows, T. B.; Tenen, D. G.; Ackerman, S. J.: Localization of
[51032] the human eosinophil Charcot-Leyden crystal protein

(lysophospholipase)

[51033] gene (CLC) to chromosome 19 and the human ribonuclease 2 (eosinophil-derived

[51034] neurotoxin) and ribonuclease 3 (eosinophil cationic protein) genes

[51035] (RNS2 and RNS3) to chromosome 14. Genomics 13: 240-242, 1992.

[51036] 8079.

[51037] Benner, S. A.: The past as the key to the present: resurrection

[51038] of ancient proteins from eosinophils. (Commentary) Proc. Nat. Acad.

[51039] Sci. 99: 4760-4761, 2002.

[51040] 8080.

[51041] Rosenberg, H. F.; Tenen, D. G.; Ackerman, S. J.: Molecular cloning

[51042] of the human eosinophil-derived neurotoxin: a member of the ribonuclease

[51043] gene family. Proc. Nat. Acad. Sci. 86: 4460-4464, 1989.

[51044] 8081.

[51045] Zhang, J.; Rosenberg, H. F.: Complementary advantageous

substitutions

[51046] in the evolution of an antiviral RNase of higher primates. Proc.

[51047] Nat. Acad. Sci. 99: 5486-5491, 2002.

[51048] 8082.

[51049] Goetzl, E. J.: Personal Communication. Boston, Mass. 12/4/1978.

[51050] 8083.

[51051] Pearson, W. R.; Vorachek, W. R.; Xu, S.; Berger, R.; Hart, I.;

[51052] Vannais, D.; Patterson, D.: Identification of class-mu glutathione

[51053] transferase genes GSTM1-GSTM5 on human chromosome 1p13. Am. J. Hum.

[51054] Genet. 53: 220-233, 1993.

[51055] 8084.

[51056] Hillarp, A.; Dahlback, B.: Novel subunit in C4b-binding protein

[51057] required for protein S binding. J. Biol. Chem. 263: 12759-12764,

[51058] 1988.

[51059] 8085.

[51060] Hillarp, A.; Dahlback, B.: Cloning of cDNA coding for the beta-chain
[51061] of human complement component C4b-binding protein: sequence homology
[51062] with the alpha chain. Proc. Nat. Acad. Sci. 87: 1183-1187, 1990.
[51063] 8086.

[51064] Hillarp, A.; Pardo-Manuel, F.; Ramos Ruiz, R.; Rodriguez de Cordoba,
[51065] R.; Dahlback, B.: The human C4b-binding protein beta-chain gene. J.
[51066] Biol. Chem. 268: 15017-15023, 1993.

[51067] 8087.

[51068] Pardo-Manuel, F.; Rey-Campos, J.; Hillarp, A.; Dahlback, B.; Rodriguez
[51069] de Cordoba, S.: Human genes for the alpha and beta chains of complement
[51070] C4b-binding protein are closely linked in a head-to-tail arrangement. Proc.
[51071] Nat. Acad. Sci. 87: 4529-4532, 1990.

[51072] 8088.

[51073] Rodriguez de Cordoba, S.; Perez-Blas, M.; Ramos-Ruiz, R.; Sanchez-Corral,
[51074] P.; Pardo-Manuel de Villena, F.; Rey-Campos, J.: The gene coding
[51075] for the beta-chain of C4b-binding protein (C4BPB) has become a pseudogene
[51076] in the mouse. Genomics 21: 501-509, 1994.

[51077] 8089.

[51078] Asghar, S. S.; Venneker, G. T.; van Meegen, M.; Meinardi, M. M.
[51079] H. M.; Hulsmans, R.-F. H. J.; de Waal, L. P.: Hereditary deficiency
[51080] of C5 in association with discoid lupus erythematosus. J. Am. Acad.
[51081] Derm. 24: 376-378, 1991.

[51082] 8090.

[51083] Collaborative Study on the Genetics of Asthma: A genome-wide
[51084] search for asthma susceptibility loci in ethnically diverse populations. Nature
[51085] Genet. 15: 389-392, 1997.

[51086] 8091.

[51087] Gavett, S. H.; O'Hearn, D. J.; Li, X.; Huang, S. K.; Finkel-
man,
[51088] F. D.; Wills-Karp, M.: Interleukin 12 inhibits antigen-in-
duced airway
[51089] hyperresponsiveness, inflammation, and Th2 cytokine ex-
pression in
[51090] mice. J. Exp. Med. 182: 1527-1536, 1995.

[51091] 8092.

[51092] Heist, D. G.; Solis-Cohen, S.; Solis-Cohen, M.: A study of
the
[51093] virulence of meningococci for man and human suscepti-
bility to meningococcic
[51094] infection. J. Immun. 7: 1-33, 1922.

[51095] 8093.

[51096] Jacobs, J. C.; Miller, M. E.: Fatal familial Leiner's disease:

[51097] a deficiency of the opsonic activity of serum complement.
Pediatrics 49:
[51098] 225-232, 1972.

[51099] 8094.

[51100] Jeremiah, S. J.; West, L. F.; Davis, M.; Povey, S.; Carritt, B.;

[51101] Fey, G. H.: The assignment of the human gene coding for complement

[51102] C5 to chromosome 9q22-9q33. Ann. Hum. Genet. 52: 111-116, 1988.

[51103] 8095.

[51104] Jeremiah, S. J.; West, L. F.; Davis, M. B.; Povey, S.; Carritt,

[51105] B.; Fey, G.: Assignment of human complement component C5 to chromosome

[51106] 9. (Abstract) Cytogenet. Cell Genet. 46: 634 only, 1987.

[51107] 8096.

[51108] Karp, C. L.; Grupe, A.; Schadt, E.; Ewart, S. L.; Keane-Moore,

[51109] M.; Cuomo, P. J.; Kohl, J.; Wahl, L.; Kuperman, D.; Germer, S.; Aud,

[51110] D.; Peltz, G.; Wills-Karp, M.: Identification of complement factor

[51111] 5 as a susceptibility locus for experimental allergic asthma. Nature

[51112] Immun. 1: 221-226, 2000.

[51113] 8097.

[51114] McLean, R. H.; Peter, G.; Gold, R.; Guerra, L.; Yunis, E. J.; Kreutzer,

[51115] D. L.: Familial deficiency of C5 in humans: intact but defi-cient

[51116] alternative complement pathway activity. Clin. Immun. Im-munopath. 21:

[51117] 62-76, 1981.

[51118] 8098.

[51119] Miller, M. E.; Nilsson, U. R.: A familial deficiency of the phagocytosis-enhancing

[51120] activity of serum related to a dysfunction of the fifth com-ponent

[51121] of complement (C5). New Eng. J. Med. 282: 354-358, 1970.

[51122] 8099.

[51123] Miller, M. E.; Seals, J.; Kaye, R.; Levitsky, L. C.: A familial,

[51124] plasma-associated defect of phagocytosis: a new cause of recurrent

[51125] bacterial infections. Lancet II: 60-63, 1968.

[51126] 8100.

[51127] Ober, C.; Cox, N. J.; Abney, M.; Di Rienzo, A.; Lander, E. S.;

[51128] Changyaleket, B.; Gidley, H.; Kurtz, B.; Lee, J.; Nance, M.; Pettersson,

[51129] A.; Prescott, J.; Richardson, A.; Schlenker, E.; Summerhill, E.; Willadsen,

[51130] S.; Parry, R.; Collaborative Study on the Genetics of Asthma: Genome-wide

[51131] search for asthma susceptibility loci in a founder population. Hum.

[51132] Molec. Genet. 7: 1393-1398, 1998.

[51133] 8101.

[51134] Ooi, Y. M.; Colten, H. R.: Genetic defect in secretion of complement

[51135] C5 in mice. Nature 282: 207-208, 1979.

[51136] 8102.

[51137] Rosenfeld, S. I.; Baum, J.; Steigbigel, R. T.; Leddy, J. P.:

[51138] Hereditary deficiency of the fifth component of complement in man.

[51139] II. Biological properties of C5-deficient human serum. J. Clin. Invest. 57:

[51140] 1635-1643, 1976.

[51141] 8103.

[51142] Rosenfeld, S. I.; Kelly, M. E.; Leddy, J. P.: Hereditary deficiency
[51143] of the fifth component of complement in man. I. Clinical, immunochemical,
[51144] and family studies. J. Clin. Invest. 57: 1626-1634, 1976.

[51145] 8104.

[51146] Rosenfeld, S. I.; Leddy, J. P.: Hereditary deficiency of fifth
[51147] component of complement (C5) in man. (Abstract) J. Clin. Invest. 53:
[51148] 67A only, 1974.

[51149] 8105.

[51150] Rosenfeld, S. I.; Weitkamp, L. R.; Countryman, J. K.: Non-linkage
[51151] for a locus of human complement C5 deficiency to the complement C6
[51152] structural locus. Immunogenetics 7: 95-97, 1978.

[51153] 8106.

[51154] Schifferli, J. A.; Hirschel, B.: Meningococcal meningitis in
[51155] the first case of complement deficiency. (Letter) Lancet II: 1240

[51156] only, 1985.

[51157] 8107.

[51158] Snyderman, R.; Durack, D. T.; McCarty, G. A.; Ward, F. E.; Meadows,

[51159] L.: Deficiency of the fifth component of complement in human subjects:

[51160] clinical, genetic and immunologic studies in a large kindred. Am.

[51161] J. Med. 67: 638-645, 1979.

[51162] 8108.

[51163] Tack, B. F.; Morris, S. C.; Prahl, J. W.: Fifth component of

[51164] human complement: purification from plasma and polypeptide chain structure. Biochemistry 18:

[51165] 1490-1497, 1979.

[51166] 8109.

[51167] Weitkamp, L. R.; Rosenfeld, S.; Johnston, E.: Complement C5:

[51168] immunofixation electrophoresis, quantitative variants, and nonlinkage

[51169] to HLA. Cytogenet. Cell Genet. 22: 651-654, 1978.

[51170] 8110.

[51171] Wetsel, R. A.; Fleischer, D. T.; Haviland, D. L.: Deficiency

[51172] of the murine fifth complement component (C5): a 2-base pair gene

[51173] deletion in a 5-prime-exon. J. Biol. Chem. 265: 2435-2440, 1990.

[51174] 8111.

[51175] Wetsel, R. A.; Lemons, R. S.; Le Beau, M. M.; Barnum, S. R.; Noack,

[51176] D.; Tack, B. F.: Molecular analysis of human complement component

[51177] C5: localization of the structural gene to chromosome 9. Biochemistry 27:

[51178] 1474-1482, 1988.

[51179] 8112.

[51180] Carbone, I.; Bruno, C.; Sotgia, F.; Bado, M.; Broda, P.; Masetti,

[51181] E.; Panella, A.; Zara, F.; Bricarelli, F. D.; Cordone, G.; Lisanti,

[51182] M. P.; Minetti, C.: Mutation in the CAV3 gene causes partial caveolin-3

[51183] deficiency and hyperCKemia. Neurology 54: 1373-1376, 2000.

[51184] 8113.

[51185] Aguzzi, A.; Weissmann, C.: A suspicious signature. Nature 383:

[51186] 666-667, 1996.

[51187] 8114.

[51188] Collinge, J.; Sidle, K. C. L.; Heads, J.; Ironside, J.; Hill,

[51189] A. F.: Molecular analysis of prion strain variation and the aetiology

[51190] of 'new variant' CJD. Nature 383: 685-690, 1996.

[51191] 8115.

[51192] Kishi, K.; Yasuda, T.; Awazu, S.; Mizuta, K.: Genetic poly-morphism

[51193] of human urine deoxyribonuclease I. Hum. Genet. 81: 295-297, 1989.

[51194] 8116.

[51195] Kishi, K.; Yasuda, T.; Ikehara, Y.; Sawazaki, K.; Sato, W.; Iida,

[51196] R.: Human serum deoxyribonuclease I (DNase I) polymor-phism: pattern

[51197] similarities among isozymes from serum, urine, kidney, liver, and

[51198] pancreas. Am. J. Hum. Genet. 47: 121-126, 1990.

[51199] 8117.

[51200] Napirei, M.; Karsunky, H.; Zevnik, B.; Stephan, H.; Mannherz, H.

[51201] G.; Moroy, T.: Features of systemic lupus erythematosus in Dnase1-deficient

[51202] mice. Nature Genet. 25: 177-181, 2000.

[51203] 8118.

[51204] Yasuda, T.; Awazu, S.; Sato, W.; Iida, R.; Tanaka, Y.; Kishi, K.

[51205] : Human genetically polymorphic deoxyribonuclease: purification, characterization,

[51206] and multiplicity of urine deoxyribonuclease I. J. Biochem. 108:

[51207] 393-398, 1990.

[51208] 8119.

[51209] Yasuda, T.; Kishi, K.; Yanagawa, Y.; Yoshida, A.: Structure of

[51210] the human deoxyribonuclease I (DNase I) gene: identifica-

tion of the
[51211] nucleotide substitution that generates its classical genetic polymorphism. Ann.
[51212] Hum. Genet. 59: 1-15, 1995.

[51213] 8120.

[51214] Yasuda, T.; Nadano, D.; Iida, R.; Takeshita, H.; Lane, S. A.; Callen,
[51215] D. F.; Kishi, K.: Chromosomal assignment of the human deoxyribonuclease
[51216] I gene, DNASE1 (DNL1), to band 16p13.3 using the poly-merase chain
[51217] reaction. Cytogenet. Cell Genet. 70: 221-223, 1995.

[51218] 8121.

[51219] Yasutomo, K.; Horiuchi, T.; Kagami, S.; Tsukamoto, H.; Hashimura,
[51220] C.; Urushihara, M.; Kuroda, Y.: Mutation of DNASE1 in people with
[51221] systemic lupus erythematosus. Nature Genet. 28: 313-314, 2001.
[51222] 8122.

[51223] Muller, U.; Steinberger, D.; Nemeth, A. H.: Clinical and

molecular
[51224] genetics of primary dystonias. Neurogenetics 1: 165-177, 1998.
[51225] 8123.
[51226] Ozelius, L. J.; Hewett, J. W.; Page, C. E.; Bressman, S. B.; Kramer,
[51227] P. L.; Shalish, C.; de Leon, D.; Brin, M. F.; Raymond, D.; Corey,
[51228] D. P.; Fahn, S.; Risch, N. J.; Buckler, A. J.; Gusella, J. F.; Breakefield,
[51229] X. O.: The early-onset torsion dystonia gene (DYT1) en-codes an ATP-binding
[51230] protein. Nature Genet. 17: 40-48, 1997.
[51231] 8124.
[51232] Winograd, J.; Reilly, M. P.; Roe, R.; Lutz, J.; Laughner, E.;
[51233] Xu, X.; Hu, L.; Asakura, T.; vander Kolk, C.; Strandberg, J. D.; Semenza,
[51234] G. L.: Perinatal lethality and multiple craniofacial malfor-mations
[51235] in MSX2 transgenic mice. Hum. Molec. Genet. 6: 369-378, 1997.
[51236] 8125.

[51237] Wuyts, W.; Reardon, W.; Preis, S.; Homfray, T.; Rasore-Quartino,
[51238] A.; Christians, H.; Willems, P. J.; Van Hul, W.: Identification of
[51239] mutations in the MSX2 homeobox gene in families affected with foramina
[51240] parietalia permagna. Hum. Molec. Genet. 9: 1251-1255, 2000.
[51241] 8126.

[51242] Jabs, E. W.; Li, X.; Scott, A. F.; Meyers, G.; Chen, W.; Eccles,
[51243] M.; Mao, J.; Charnas, L. R.; Jackson, C. E.; Jaye, M.: Jackson-Weiss
[51244] and Crouzon syndromes are allelic with mutations in fibroblast growth
[51245] factor receptor 2. Nature Genet. 8: 275-279, 1994.

[51246] 8127.

[51247] Roscioli, T.; Flanagan, S.; Kumar, P.; Masel, J.; Gattas, M.;
[51248] Hyland, V. J.; Glass, I. A.: Clinical findings in a patient with
[51249] FGFR1 P252R mutation and comparison with the literature. Am. J. Med.
[51250] Genet. 93: 22-28, 2000.

[51251] 8128.

[51252] Vinton, N. E.; Dahlstrom, K. A.; Stobel, C. T.; Ament, M. E.:

[51253] Macrocytosis and pseudoalbinism: manifestations of sele-
nium deficiency. J.

[51254] Pediat. 111: 711-717, 1987.

[51255] 8129.

[51256] Wijnen, L. M.; Monteba-van Heuvel, M.; Pearson, P. L.;
Meera Khan,

[51257] P.: Assignment of a gene for glutathione peroxidase
(GPX-1) to human

[51258] chromosome 3. Cytogenet. Cell Genet. 22: 232-238,
1978.

[51259] 8130.

[51260] Yannicelli, S.; Hambidge, K. M.; Picciano, M. F.: Decreased
selenium

[51261] intake and low plasma selenium concentrations leading to
clinical

[51262] symptoms in a child with propionic acidaemia. J. Inherit.
Metab.

[51263] Dis. 15: 261-268, 1992.

[51264] 8131.

[51265] Chambers, I.; Frampton, J.; Goldfarb, P.; Affara, N.; McBain, W.;

[51266] Harrison, P. R.: The structure of the mouse glutathione peroxidase

[51267] gene: the selenocysteine in the active site is encoded by the 'termination'

[51268] codon, TGA. EMBO J. 5: 1221-1227, 1986.

[51269] 8132.

[51270] Chu, F.-F.; Esworthy, R. S.; Doroshow, J. H.; Doan, K.; Liu, X.-F.

[51271] : Expression of glutathione peroxidase in human liver in addition

[51272] to kidney, heart, lung, and breast in humans and rodents. Blood 79:

[51273] 3233-3238, 1992.

[51274] 8133.

[51275] Takahashi, K.; Akasaka, M.; Yamamoto, Y.; Kobayashi, C.; Mizoguchi,

[51276] J.; Koyama, J.: Primary structure of human plasma glutathione peroxidase

[51277] deduced from cDNA sequences. J. Biochem. 108: 145-148, 1990.

[51278] 8134.

[51279] Kelner, M. J.; Montoya, M. A.: Structural organization of the

[51280] human selenium-dependent phospholipid hydroperoxide glutathione peroxidase

[51281] gene (GPX4): chromosomal localization to 19p13.3. Biochem. Biophys.

[51282] Res. Commun. 249: 53-55, 1998.

[51283] 8135.

[51284] Roveri, A.; Casasco, A.; Maiorino, M.; Dalan, P.; Calligaro, A.;

[51285] Ursini, F.: Phospholipid hydroperoxide glutathione peroxi-dase of

[51286] rat testis. J. Biol. Chem. 267: 6142-6146, 1992.

[51287] 8136.

[51288] Ikeuchi, T.; Shimohata, T.; Nakano, R.; Koide, R.; Takano, H.;

[51289] Tsuji, S.: A case of primary torsion dystonia in Japan with the 3-bp

[51290] (GAG) deletion in the DYT1 gene with a unique clinical presentation.

[51291] (Letter) Neurogenetics 2: 189-190, 1999.

[51292] 8137.

[51293] Kamm, C.; Castelon-Konkiewitz, E.; Naumann, M.; Heinen, F.; Brack,

[51294] M.; Nebe, A.; Ceballos-Baumann, A.; Gasser, T.: GAG deletion in the

[51295] DYT1 gene in early limb-onset idiopathic torsion dystonia in Germany. Mov.

[51296] Disord. 14: 681-683, 1999.

[51297] 8138.

[51298] Arce, M. A.; Thompson, E. S.; Wagner, S.; Coyne, K. E.; Ferdman,

[51299] B. A.; Lublin, D. M.: Molecular cloning of RhD cDNA de-rived from

[51300] a gene present in RhD-positive, but not RhD-negative in-dividuals. Blood 82:

[51301] 651-655, 1993.

[51302] 8139.

[51303] Bennett, P. R.; Le Van Kim, C.; Colin, Y.; Warwick, R. M.; Cherif-Zahar,

[51304] B.; Fisk, N. M.; Cartron, J.-P.: Prenatal determination of fe-

tal

[51305] RhD type by DNA amplification. New Eng. J. Med. 329: 607-610, 1993.

[51306] 8140.

[51307] Bowman, J. M.: RhD hemolytic disease of the newborn. (Editorial) New

[51308] Eng. J. Med. 339: 1775-1777, 1998.

[51309] 8141.

[51310] Cartron, J.-P.: Defining the Rh blood group antigens: bio-chemistry

[51311] and molecular genetics. Blood Rev. 8: 199-212, 1994.

[51312] 8142.

[51313] Colin, Y.; Cherif-Zahar, B.; Le Van Kim, C.; Raynal, V.; Van Huffel,

[51314] V.; Cartron, J.-P.: Genetic basis of the RhD-positive and RhD-negative

[51315] blood group polymorphism as determined by Southern analysis. Blood 78:

[51316] 2747-2752, 1991.

[51317] 8143.

[51318] Diamond, L. K.; Blackfan, K. D.; Baty, J. M.: Erythroblasto-

sis
[51319] fetalis and its association with universal edema of the fetus, icterus
[51320] gravis neonatorum and anemia of the newborn. J. Pediat. 1: 269-309,
[51321] 1932.

[51322] 8144.

[51323] du Toit, E. D.; Martell, R. W.; Botha, I.; Kriel, C. J.: Anti-D
[51324] antibodies in the Rh-positive mothers. S. Afr. Med. J. 75: 452,
[51325] 1989.

[51326] 8145.

[51327] Garratty, G.: Severe reactions associated with transfusion of
[51328] patients with sickle cell disease. Transfusion 37: 357-361, 1997.
[51329] 8146.

[51330] Huang, C.-H.: Personal Communication. New York City, N. Y.
[51331] 10/11/1996.

[51332] 8147.

[51333] Huang, C.-H.; Chen, Y.; Reid, M.; Ghosh, S.: Genetic re-
combination
[51334] at the human RH locus: a family study of the red-cell
Evans phenotype
[51335] reveals a transfer of exons 2-6 from the RHD to the RHCE
gene. Am.
[51336] J. Hum. Genet. 59: 825-833, 1996.

[51337] 8148.

[51338] Huang, C.-H.; Reid, M. E.; Chen, Y.; Coghlan, G.; Okubo,
Y.:
[51339] Molecular definition of red cell Rh haplotypes by tightly
linked SphI
[51340] RFLPs. Am. J. Hum. Genet. 58: 133-142, 1996.

[51341] 8149.

[51342] Hyland, C. A.; Wolter, L. C.; Liew, Y. W.; Saul, A.: A South-
ern
[51343] analysis of Rh blood group genes: association between re-
striction
[51344] fragment length polymorphism patterns and Rh serotypes.
Blood 83:
[51345] 566-572, 1994.

[51346] 8150.

[51347] Issitt, P. D.: The Rh blood group system, 1988: eight new antigens
[51348] in nine years and some observations on the biochemistry and genetics
[51349] of the system. Transfusion Med. Rev. 3: 1-12, 1989.

[51350] 8151.

[51351] Kemp, T. J.; Poulter, M.; Carritt, B.: A recombination hot spot
[51352] in the Rh genes revealed by analysis of unrelated donors with the
[51353] rare D-- phenotype. Am. J. Hum. Genet. 59: 1066-1073, 1996.

[51354] 8152.

[51355] Legler, T. J.; Eber, S. W.; Lakomek, M.; Lynen, R.; Maas, J. H.;
[51356] Pekrun, A.; Repas-Humpe, M.; Schroter, W.; Kohler, M.: Application
[51357] of RHD and RHCE genotyping for correct blood group de-termination in
[51358] chronically transfused patients. Transfusion 39: 852-855, 1999.

[51359] 8153.

[51360] Le Van Kim, C.; Cherif-Zahar, B.; Raynal, V.; Mouro, I.; Lopez,

[51361] M.; Cartron, J. P.; Colin, Y.: Multiple Rh messenger RNA isoforms

[51362] are produced by alternative splicing. Blood 80: 1074-1078, 1992.

[51363] 8154.

[51364] Le Van Kim, C.; Mouro, I.; Cherif-Zahar, B.; Raynal, V.; Cherrier,

[51365] C.; Cartron, J.-P.; Colin, Y.: Molecular cloning and primary structure

[51366] of the human blood group RhD polypeptide. Proc. Nat. Acad. Sci. 89:

[51367] 10925-10929, 1992.

[51368] 8155.

[51369] Levine, P.; Katzin, E. M.; Burnham, L.: Isoimmunization in pregnancy:

[51370] its possible bearing on the etiology of erythroblastosis foetalis. J.A.M.A. 116:

[51371] 825-827, 1941.

[51372] 8156.

[51373] Lo, Y. M. D.; Hjelm, N. M.; Fidler, C.; Sargent, I. L.; Murphy,

[51374] M. F.; Chamberlain, P. F.; Poon, P. M. K.; Redman, C. W. G.; Wainscoat,

[51375] J. S.: Prenatal diagnosis of fetal RhD status by molecular analysis

[51376] of maternal plasma. New Eng. J. Med. 339: 1734-1738, 1998.

[51377] 8157.

[51378] Miyoshi, O.; Yabe, R.; Wakui, K.; Fukushima, Y.; Koizumi, S.;

[51379] Uchikawa, M.; Kajii, T.; Numakura, C.; Takahashi, S.; Hayasaka, K.;

[51380] Niikawa, N.: Two cases of mosaic RhD blood-group phe-notypes and paternal

[51381] isodisomy for chromosome 1. Am. J. Med. Genet. 104: 250-256, 2001.

[51382] 8158.

[51383] Okuda, H.; Kawano, M.; Iwamoto, S.; Tanaka, M.; Seno, T.; Okubo,

[51384] Y.; Kajii, E.: The RHD gene is highly detectable in RhD-negative

[51385] Japanese donors. J. Clin. Invest. 100: 373-379, 1997.

[51386] 8159.

[51387] Race, R. R.: An 'incomplete' antibody in human serum. (Letter) Nature 153:

[51388] 771-772, 1944.

[51389] 8160.

[51390] Race, R. R.; Sanger, R.: Blood Groups in Man. Oxford: Blackwell

[51391] (pub.) (6th ed.): 1975.

[51392] 8161.

[51393] Rouillac, C.; Le Van Kim, C.; Beolet, M.; Cartron, J.-P.; Colin,

[51394] Y.: Leu110-to-pro substitution in the RhD polypeptide is responsible

[51395] for the D(VII) category blood group phenotype. Am. J. Hemat. 49:

[51396] 87-88, 1995.

[51397] 8162.

[51398] Smythe, J. S.; Avent, N. D.; Judson, P. A.; Parsons, S. F.; Martin,

[51399] P. G.; Anstee, D. J.: Expression of RHD and RHCE gene products using

[51400] retroviral transduction of K562 cells establishes the molecular basis

[51401] of Rh blood group antigens. Blood 87: 2968-2973, 1996.

[51402] 8163.

[51403] Spanos, T.; Karageorga, M.; Ladis, V.; Peristeri, J.; Hatzil-iami,

[51404] A.; Kattamis, C.: Red cell alloantibodies in patients with thalassemia. Vox

[51405] Sang. 58: 50-55, 1990.

[51406] 8164.

[51407] Wagner, F. F.; Flegel, W. A.: RHD gene deletion occurred in the

[51408] Rhesus box. Blood 95: 3662-3668, 2000.

[51409] 8165.

[51410] Wagner, F. F.; Gassner, C.; Muller, T. H.; Schonitzer, D.; Schunter,

[51411] F.; Flegel, W. A.: Molecular basis of weak D phenotypes. Blood 93:

[51412] 385-393, 1999.

[51413] 8166.

[51414] Wagner, F. F.; Ladewig, B.; Angert, K. S.; Heymann, G. A.; Eicher,

[51415] N. I.; Flegel, W. A.: The DAU allele cluster of the RHD gene. Blood 100:

[51416] 306-311, 2002.

[51417] 8167.

[51418] Wiener, A. S.: The Rh series of allelic genes. Science 100:

[51419] 595-597, 1944.

[51420] 8168.

[51421] Wang, X.; Chan, S. J.; Eddy, R. L.; Byers, M. G.; Fukushima, Y.;

[51422] Henry, W. M.; Haley, L. L.; Steiner, D. F.; Shows, T. B.: Chromosome

[51423] assignment of cathepsin B (CTSB) to 8p22 and cathepsin H (CTSH) to

[51424] 15q24-q25. (Abstract) Cytogenet. Cell Genet. 46:

710-711, 1987.

[51425] 8169.

[51426] Gelb, B. D.; Shi, G.-P.; Heller, M.; Weremowicz, S.; Morton, C.;

[51427] Desnick, R. J.; Chapman, H. A.: Structure and chromosomal assignment
[51428] of the human cathepsin K gene. Genomics 41: 258-262, 1997.
[51429] 8170.
[51430] Shiang, R.; Lidral, A. C.; Ardinger, H. H.; Buetow, K. H.; Romitti,
[51431] P. A.; Munger, R. G.; Murray, J. C.: Association of transforming
[51432] growth-factor alpha gene polymorphisms with nonsyndromic cleft palate
[51433] only (CPO). Am. J. Hum. Genet. 53: 836-843, 1993.
[51434] 8171.
[51435] Shields, E. D.; Bixler, D.; Fogh-Andersen, P.: Facial clefts
[51436] in Danish twins. Cleft Palate J. 16: 1-6, 1979.
[51437] 8172.
[51438] Pollack, M. S.; Ochs, H. D.; Dupont, B.: HLA typing of cultured
[51439] amniotic cells for the prenatal diagnosis of complement C4 deficiency. Clin.
[51440] Genet. 18: 197-200, 1980.

[51441] 8173.

[51442] Porter, R. R.; Reid, K. B. M.: The biochemistry of comple-
ment. Nature 275:
[51443] 699-704, 1978.

[51444] 8174.

[51445] Rittner, C.; Bertrams, J.: On the significance of C2, C4, and
[51446] factor B polymorphisms in disease. Hum. Genet. 56:
235-247, 1981.
[51447] 8175.

[51448] Rittner, C.; Hauptmann, G.; Grosse-Wilde, H.; Grosshans,
E.; Tongio,
[51449] M. M.; Mayer, S.: Linkage between HL-A (major histocom-
patibility
[51450] complex) and genes controlling the fourth component of
complement.In:
[51451] Histocompatibility Testing 1975. Copenhagen: Munks-
gaard (pub.)
[51452] 1976. Pp. 945-953.

[51453] 8176.

[51454] Roos, M. H.; Mollenhauer, E.; Demant, P.; Rittner, C.: A
molecular

[51455] basis for the two locus model of human complement component C4. Nature 298:
[51456] 854-856, 1982.

[51457] 8177.

[51458] Rosenfeld, S. I.; Ruddy, S.; Austen, K. F.: Structural polymorphism
[51459] of the fourth component of human complement. J. Clin. Invest. 48:
[51460] 2283-2292, 1969.

[51461] 8178.

[51462] Roychoudhury, A. K.; Nei, M.: Human Polymorphic Genes: World
[51463] Distribution. New York: Oxford Univ. Press (pub.) 1988.

[51464] 8179.

[51465] Schaller, J. G.; Gilliland, B. G.; Ochs, H. D.; Leddy, J. P.;

[51466] Agodoa, L. C. Y.; Rosenfeld, S. I.: Severe systemic lupus erythematosus
[51467] with nephritis in a boy with deficiency of the fourth component of
[51468] complement. Arthritis Rheum. 20: 1519-1525, 1977.

[51469] 8180.

[51470] Shevach, E. M.; Frank, M. M.; Green, I.: Linkage of gene controlling

[51471] the synthesis of the fourth component of complement to the major histocompatibility

[51472] complex of the guinea pig. Immunogenetics 3: 595-602, 1976.

[51473] 8181.

[51474] Shreffler, D. C.: The S region of the mouse major histo-compatibility

[51475] complex (H-2): genetic variation and functional role in complement

[51476] system. Transplant. Rev. 32: 140-167, 1976.

[51477] 8182.

[51478] Torisu, M.; Sonozaki, H.; Inai, S.; Arata, M.: Deficiency of

[51479] the fourth component of complement in man. J. Immuno-genet. 104:

[51480] 728-737, 1970.

[51481] 8183.

[51482] Wank, R.; Schendel, D. J.; O'Neill, G. J.; Riethmuller, G.; Held,

[51483] E.; Feucht, H. E.: Rare variant of complement C4 is seen in high
[51484] frequency in patients with primary glomerulonephritis. Lancet I:
[51485] 872-874, 1984.

[51486] 8184.

[51487] Welch, T. R.; Beischel, L.: C4 uremic variant: an acquired C4
[51488] allotype. Immunogenetics 22: 553-562, 1985.

[51489] 8185.

[51490] Welch, T. R.; Beischel, L. S.; Choi, E.; Balakrishnan, K.; Bishof,
[51491] N. A.: Uniparental isodisomy 6 associated with deficiency of the
[51492] fourth component of complement. J. Clin. Invest. 86: 675-678, 1990.
[51493] 8186.

[51494] Whitehead, A. S.; Colten, H. R.; Chang, C. C.; Demars, R.: Localization
[51495] of the human MHC-linked complement genes between HLA-B and HLA-DR

[51496] by using HLA mutant cell lines. J. Immun. 134: 641-643, 1985.

[51497] 8187.

[51498] Wilson, W. A.; Perez, M. C.: Complete C4B deficiency in black

[51499] Americans with systemic lupus erythematosus. J. Rheum. 15: 1855-1858,

[51500] 1988.

[51501] 8188.

[51502] Andersson, A.; Dahlback, B.; Hanson, C.; Hillarp, A.; Levan, G.;

[51503] Szpirer, J.; Szpirer, C.: Genes for C4b-binding protein al-pha- and

[51504] beta-chains (C4BPA and C4BPB) are located on chromo-some 1, band 1q32,

[51505] in humans and on chromosome 13 in rats. Somat. Cell Molec. Genet. 16:

[51506] 493-500, 1990.

[51507] 8189.

[51508] Rodriguez de Cordoba, S.; Lublin, D. M.; Rubinstein, P.; Atkinson,

[51509] J. P.: Human genes for three complement components that regulate
[51510] the activation of C3 are tightly linked. J. Exp. Med. 161: 1189-1195,
[51511] 1985.

[51512] 8190.

[51513] Hofstra, R. M. W.; Osinga, J.; Tan-Sindhunata, G.; Wu, Y.; Kamsteeg,
[51514] E.-J.; Stulp, R. P.; van Ravenswaaij-Arts, C.; Majoor-Krakauer, D.;
[51515] Angrist, M.; Chakravarti, A.; Meijers, C.; Buys, C. H. C. M.: A homozygous
[51516] mutation in the endothelin-3 gene associated with a combined Hirschsprung
[51517] type 2 and Hirschsprung phenotype (Shah-Waardenburg syndrome). Nature
[51518] Genet. 12: 445-447, 1996.

[51519] 8191.

[51520] Lenzerini, L.; Benatti, U.; Morelli, A.; Pontremoli, S.; De Flora,
[51521] A.; Piazza, A.; Rinaldi, A.; Filippi, G.; Siniscalco, M.: Genetic

[51522] variation in the quantitative levels of an NADP(H)-binding protein

[51523] (FX) in human erythrocytes. Blood 57: 209-217, 1981.

[51524] 8192.

[51525] Morelli, A.; De Flora, A.: Isolation and partial characterization

[51526] of an NADP- and NADPH-binding protein from human erythrocytes. Arch.

[51527] Biochem. Biophys. 179: 698-705, 1977.

[51528] 8193.

[51529] Sullivan, F. X.; Kumar, R.; Kriz, R.; Stahl, M.; Xu, G.-Y.; Rouse,

[51530] J.; Chang, X.; Boodhoo, A.; Potvin, B.; Cumming, D. A.: Molecular

[51531] cloning of human GDP-mannose 4,6-dehydratase and reconstitution of

[51532] GDP-fucose biosynthesis in vitro. J. Biol. Chem. 273: 8193-8202,

[51533] 1998.

[51534] 8194.

[51535] Tonetti, M.; Sturla, L.; Bisso, A.; Benatti, U.; De Flora, A.:

[51536] Synthesis of GDP-L-fucose by the human FX protein. J. Biol. Chem. 271:
[51537] 27274-27279, 1996.

[51538] 8195.

[51539] Boyle, J. M.; Hey, Y.; Myers, K.; Stern, P. L.; Grzeschik, F.-H.;
[51540] Ikehara, Y.; Misumi, Y.; Fox, M.: Regional localization of a trophoblast
[51541] antigen-related sequence and 16 other sequences to human chromosome
[51542] 6q using somatic cell hybrids. Genomics 12: 693-698, 1992.
[51543] 8196.

[51544] Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp, N.;
[51545] Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins, N.
[51546] A.: A genetic linkage map of mouse chromosome 10: localization of
[51547] eighteen molecular markers using a single interspecific backcross. Genetics 125:
[51548] 855-866, 1990.

[51549] 8197.

[51550] Mouillet-Richard, S.; Ermonval, M.; Chebassier, C.; La-
planche,
[51551] J. L.; Lehmann, S.; Launay, J. M.; Kellermann, O.: Signal
transduction
[51552] through prion protein. Science 289: 1925-1928, 2000.

[51553] 8198.

[51554] Casari, G.; Barlassina, C.; Cusi, D.; Zagato, L.; Muirhead,
R.;
[51555] Righetti, M.; Nembri, P.; Amar, K.; Gatti, M.; Macciardi, F.;
Binelli,
[51556] G.; Bianchi, G.: Association of the alpha-adducin locus
with essential
[51557] hypertension. Hypertension 25: 320-326, 1995.

[51558] 8199.

[51559] Lee, R. T.; Peterson, C. L.; Calman, A. F.; Herskowitz, I.;
O'Donnell,
[51560] J. J.: Cloning of a human galactokinase gene (GK2) on
chromosome
[51561] 15 by complementation in yeast. Proc. Nat. Acad. Sci. 89:
10887-10891,

[51562] 1992.

[51563] 8200.

[51564] Pastuszak, I.; O'Donnell, J.; Elbein, A. D.: Identification of

[51565] the GalNAc kinase amino acid sequence. J. Biol. Chem.
271: 23653-23656,

[51566] 1996.

[51567] 8201.

[51568] Ball, S. P.; Cook, P. J. L.; Mars, M.; Buckton, K. E.: Linkage

[51569] between dentinogenesis imperfecta and Gc. Ann. Hum.
Genet. 46: 35-40,

[51570] 1982.

[51571] 8202.

[51572] Crosby, A. H.; Edwards, S. J.; Murray, J. C.; Dixon, M. J.:
Genomic

[51573] organization of the human osteopontin gene: exclusion of
the locus

[51574] from a causative role in the pathogenesis of dentinogene-
sis imperfecta

[51575] type II. Genomics 27: 155-160, 1995.

[51576] 8203.

[51577] Abernethy, T. J.; Avery, O. T.: The occurrence during acute infections
[51578] of a protein not normally present in the blood. I. Distribution of
[51579] the reactive protein in patients' sera and the effect of calcium on
[51580] the flocculation reaction with C polysaccharide of pneumococcus. J.
[51581] Exp. Med. 73: 173-182, 1941.

[51582] 8204.

[51583] Floyd-Smith, G. A.; Whitehead, A. S.; Colten, H. R.; Francke, U.
[51584] : The human C-reactive protein gene (CRP) and serum amyloid P component
[51585] gene (APCS) are located on the proximal long arm of chromosome 1. Immunogenetics 24:
[51586] 171-176, 1986.

[51587] 8205.

[51588] Tillett, W. S.; Francis, T., Jr.: Serological reactions in pneumonia
[51589] with a nonprotein somatic fraction of pneumococcus. J. Exp. Med. 52:

[51590] 561-585, 1930.

[51591] 8206.

[51592] Stallings, R. L.; Olson, E.; Strauss, A. W.; Thompson, L. H.; Bachinski,

[51593] L. L.; Siciliano, M. J.: Human creatine kinase genes on chromosomes

[51594] 15 and 19, and proximity of the gene for the muscle form to the genes

[51595] for apolipoprotein C2 and excision repair. Am. J. Hum. Genet. 43:

[51596] 144-151, 1988.

[51597] 8207.

[51598] Bark, C. J.: Mitochondrial creatine kinase: a poor prognostic

[51599] sign. J.A.M.A. 243: 2058-2060, 1980.

[51600] 8208.

[51601] Steeghs, K.; Merkx, G.; Wieringa, B.: The ubiquitous mitochondrial

[51602] creatine kinase gene maps to a conserved region on human chromosome

[51603] 15q15 and mouse chromosome 2 bands F1-F3. Genomics

24: 193-195,
[51604] 1994.

[51605] 8209.

[51606] Klein, S. C.; Haas, R. C.; Perryman, M. B.; Billadello, J. J.;

[51607] Strauss, A. W.: Regulatory element analysis and structural characterization
[51608] of the human sarcomeric mitochondrial creatine kinase gene. J. Biol.
[51609] Chem. 266: 18058-18065, 1991.

[51610] 8210.

[51611] Richard, I.; Devaud, C.; Cherif, D.; Cohen, D.; Beckmann, J. S.
[51612] : The gene for creatine kinase, mitochondrial 2 (sarcomeric; CKMT2),
[51613] maps to chromosome 5q13.3. Genomics 18: 134-136, 1993.
[51614] 8211.

[51615] Holst, L. S.; Langin, D.; Mulder, H.; Laurell, H.; Grober, J.;

[51616] Bergh, A.; Mohrenweiser, H. W.; Edgren, G.; Holm, C.: Molecular cloning,
[51617] genomic organization, and expression of a testicular iso-

form of hormone-sensitive
[51618] lipase. Genomics 35: 441-447, 1996.

[51619] 8212.

[51620] Levitt, R. C.; Liu, Z.; Nouri, N.; Meyers, D. A.; Brandriff, B.;

[51621] Mohrenweiser, H. M.: Mapping of the gene for hormone sensitive lipase
[51622] (LIPE) to chromosome 19q13.1-q13.2. Cytogenet. Cell Genet. 69: 211-214,
[51623] 1995.

[51624] 8213.

[51625] Li, Z.; Sumida, M.; Birchbauer, A.; Schotz, M. C.; Reue, K.: Isolation
[51626] and characterization of the gene for mouse hormone-sensitive lipase. Genomics 24:
[51627] 259-265, 1994.

[51628] 8214.

[51629] Osuga, J.; Ishibashi, S.; Oka, T.; Yagyu, H.; Tozawa, R.; Fujimoto,
[51630] A.; Shionoiri, F.; Yahagi, N.; Kraemer, F. B.; Tsutsumi, O.; Yamada,
[51631] N.: Targeted disruption of hormone-sensitive lipase re-

sults in male

[51632] sterility and adipocyte hypertrophy, but not in obesity. Proc. Nat.

[51633] Acad. Sci. 97: 787-792, 2000.

[51634] 8215.

[51635] Schonk, D.; van Dijk, P.; Riegmann, P.; Trapman, J.; Holm, C.;

[51636] Willcocks, T. C.; Sillekens, P.; van Venrooij, W.; Wimmer, E.; Geurts

[51637] van Kessel, A.; Ropers, H.-H.; Wieringa, B.: Assignment of seven

[51638] genes to distinct intervals on the midportion of human chromosome

[51639] 19q surrounding the myotonic dystrophy gene region. Cytogenet. Cell

[51640] Genet. 54: 15-19, 1990.

[51641] 8216.

[51642] Wang, S.; Lapierre, P.; Robert, M.-F.; Nadeau, J. H.; Mitchell,

[51643] G. A.: Hormone-sensitive lipase maps to proximal chromosome 7 in

[51644] mice and is genetically distinct from the Ad and Tub loci.

Genomics 24:
[51645] 416-417, 1994.

[51646] 8217.

[51647] Dal Cin, P.; Turc-Carel, C.; Sandberg, A. A.: Consistent in-
volvement
[51648] of band 12q14 in two different translocations in three
lipomas from
[51649] the same patient. Cancer Genet. Cytogenet. 31: 237-240,
1988.
[51650] 8218.

[51651] Levi-Strauss, M.; Carroll, M. C.; Steinmetz, M.; Meo, T.: A
previously
[51652] undetected MHC gene with an unusual periodic structure.
Science 240:
[51653] 201-204, 1988.

[51654] 8219.

[51655] Speiser, P. W.; White, P. C.: Structure of the human RD
gene:
[51656] a highly conserved gene in the class III region of the major
histocompatibility
[51657] complex. DNA 8: 745-751, 1989.

[51658] 8220.

[51659] Surowy, C. S.; Gosink, J. J.; Spritz, R. A.: cDNA cloning of the

[51660] human 'RD' protein, a possible novel RNA-binding protein. (Abstract) Am.

[51661] J. Hum. Genet. 43: A203 only, 1988.

[51662] 8221.

[51663] Yamaguchi, Y.; Takagi, T.; Wada, T.; Yano, K.; Furuya, A.; Sugimoto,

[51664] S.; Hasegawa, J.; Handa, H.: NELF, a multisubunit complex containing

[51665] RD, cooperates with DSIF to repress RNA polymerase II elongation. Cell 97:

[51666] 41-51, 1999.

[51667] 8222.

[51668] Gross, M.-S.; Guyonnet-Duperat, V.; Porchet, N.; Bern-

[51669] heim, A.; Aubert, J. P.; Nguyen, V. C.: Mucin 4 (MUC4) gene: re-

[51670] gional assignment (3q29) and RFLP analysis. Ann. Genet. 35: 21-26, 1992.

[51671] 8223.

[51672] Porchet, N.; Van Cong, N.; Dufosse, J.; Audie, J. P.; Guyon-
net-Duperat,
[51673] V.; Gross, M. S.; Denis, C.; Degand, P.; Bernheim, A.;
Aubert, J.
[51674] P.: Molecular cloning and chromosomal localization of a
novel human
[51675] tracheo-bronchial mucin cDNA containing tandemly re-
peated sequences
[51676] of 48 base pairs. Biochem. Biophys. Res. Commun. 175:
414-422,
[51677] 1991.

[51678] 8224.

[51679] Van Cong, N.; Aubert, J.-P.; Gross, M.-S.; Porchet, N.; De-
gand,
[51680] P.; Bernheim, A.: Tracheobronchial mucin 4 (MUC4) gene:
assignment
[51681] to 3q29 and polymorphism of VNTR type. (Abstract) Cyto-
genet. Cell
[51682] Genet. 58: 1879-1880, 1991.

[51683] 8225.

[51684] Pigny, P.; Pratt, W. S.; Laine, A.; Leclercq, A.; Swallow, D.
M.;

[51685] Nguyen, V. C.; Aubert, J. P.; Porchet, N.: The MUC5AC gene: RFLP

[51686] analysis with the Jer58 probe. Hum. Genet. 96: 367-368, 1995.

[51687] 8226.

[51688] Collins, F. S.: Personal Communication. Bethesda, Md. 11/17/1993.

[51689] 8227.

[51690] Blanck, O.; Perrin, C.; Mziaut, H.; Darbon, H.; Mattei, M. G.;

[51691] Miquelis, R.: Molecular cloning, cDNA analysis, and localization

[51692] of a monomer of the N-acetylglucosamine-specific receptor of the thyroid,

[51693] NAGR1, to chromosome 19p13.3-13.2. Genomics 21: 18-26, 1994. Note:

[51694] Erratum: Genomics 27: 561 only, 1995.

[51695] 8228.

[51696] Datar, K. V.; Dreyfuss, G.; Swanson, M. S.: The human hnRNP M

[51697] proteins: identification of a methionine/arginine-rich repeat motif

[51698] in ribonucleoproteins. Nucleic Acids Res. 21: 439-446, 1993.

[51699] 8229.

[51700] Hull, E.; Sarkar, M.; Spruijt, M. P. N.; Hoppener, J. W. M.; Dunn,

[51701] R.; Schachter, H.: Organization and localization to chromosome 5

[51702] of the human UDP-N-acetylglucosamine:alpha-3-D-mannoside beta-1,2-N-acetylglucosaminyltransferase

[51703] 1 gene. Biochem. Biophys. Res. Commun. 176: 608-615, 1991.

[51704] 8230.

[51705] Ioffe, E.; Stanley, P.: Mice lacking N-acetylglucosaminyltransferase

[51706] I activity die at mid-gestation, revealing an essential role for complex

[51707] or hybrid N-linked carbohydrates. Proc. Nat. Acad. Sci. 91: 728-732,

[51708] 1994.

[51709] 8231.

[51710] Kumar, R.; Stanley, P.: Transfection of a human gene that

corrects

[51711] the Lec1 glycosylation defect: evidence for transfer of the structural

[51712] gene for N-acetylglucosaminyltransferase I. Molec. Cell. Biol. 9:

[51713] 5713-5717, 1989. Note: Erratum: Molec. Cell. Biol. 10: 3857 only,

[51714] 1990.

[51715] 8232.

[51716] Kumar, R.; Yang, J.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Stanley,

[51717] P.: Cloning and expression of the murine gene and chromosomal location

[51718] of the human gene encoding N-acetylglucosaminyltransferase I. Glycobiology 2:

[51719] 383-393, 1992.

[51720] 8233.

[51721] Kumar, R.; Yang, J.; Larsen, R. D.; Stanley, P.: Cloning and expression

[51722] of N-acetylglucosaminyltransferase I, the medial Golgi transferase

[51723] that initiates complex N-linked carbohydrate formation.

Proc. Nat.
[51724] Acad. Sci. 87: 9948-9952, 1990.

[51725] 8234.

[51726] Metzler, M.; Gertz, A.; Sarkar, M.; Schachter, H.; Schrader, J.
[51727] W.; Marth, J. D.: Complex asparagine-linked oligosaccharides are
[51728] required for morphogenic events during post-implantation development. EMBO
[51729] J. 13: 2056-2065, 1994.

[51730] 8235.

[51731] Pownall, S.; Kozak, C. A.; Schappert, K.; Sarkar, M.; Hull, E.;
[51732] Schachter, H.; Marth, J. D.: Molecular cloning and characterization
[51733] of the mouse UDP-N-acetylglucosamine:alpha-3-D-mannoside beta-1,2-N-acetylglucosaminyltransferase
[51734] I gene. Genomics 12: 699-704, 1992.

[51735] 8236.

[51736] Puthalakath, H.; Burke, J.; Gleeson, P. A.: Glycosylation de-

fect
[51737] in lec1 Chinese hamster ovary mutant is due to a point mutation in
[51738] N-acetylglucosaminyltransferase 1 gene. J. Biol. Chem. 271: 27818-27822,
[51739] 1996.

[51740] 8237.

[51741] Shows, T. B.: Personal Communication. Buffalo, N.Y. 4/13/1999.
[51742] 8238.

[51743] Yip, B.; Chen, S.-H.; Mulder, H.; Hoppener, J. W. M.; Schachter,
[51744] H.: Organization of the human beta-1,2-N-acetylglucosaminyltransferase
[51745] I gene (MGAT1), which controls complex and hybrid N-glycan synthesis. Biochem.
[51746] J. 321: 465-474, 1997.

[51747] 8239.

[51748] Tan, J.; D'Agostaro, G. A. F.; Bendiak, B.; Reck, F.; Sarkar,
[51749] M.; Squire, J. A.; Leong, P.; Schachter, H.: The human UDP-N-acetylglu-

cosamine:alpha-6-D-mannoside-beta-1,2-N-acetylgluco saminyltransfe

[51750] rase II gene (MGAT2): cloning of genomic DNA, localiza- tion to chromosome

[51751] 14q21, expression in insect cells and purification of the recombinant

[51752] protein. Europ. J. Biochem. 231: 317-328, 1995.

[51753] 8240.

[51754] Kondo, M.; Scherer, D. C.; Miyamoto, T.; King, A. G.; Akashi,

[51755] K.; Sugamura, K.; Weissman, I. L.: Cell-fate conversion of lymphoid-committed

[51756] progenitors by instructive actions of cytokines. Nature 407: 383-386,

[51757] 2000.

[51758] 8241.

[51759] Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Larson, R. A.;

[51760] Rowley, J. D.; Gasson, J. C.; Golde, D. W.; Sherr, C. J.: Evi- dence

[51761] for the involvement of GM-CSF and FMS in the deletion (5q) in myeloid

[51762] disorders. Science 231: 984-987, 1986.

[51763] 8242.

[51764] Morgan, R.; Hecht, B. K.; Sandberg, A. A.; Hecht, F.; Smith, S.

[51765] D.: Chromosome 5q35 breakpoint in malignant histiocytosis. (Letter) New

[51766] Eng. J. Med. 314: 1322 only, 1986.

[51767] 8243.

[51768] Ridge, S. A.; Worwood, M.; Oscier, D.; Jacobs, A.; Padua, R. A.

[51769] : FMS mutations in myelodysplastic, leukemic, and normal subjects. Proc.

[51770] Nat. Acad. Sci. 87: 1377-1380, 1990.

[51771] 8244.

[51772] Roberts, W. M.; Look, A. T.; Roussel, M. F.; Sherr, C. J.: Tandem

[51773] linkage of human CSF-1 receptor (c-fms) and PDGF receptor genes. Cell 55:

[51774] 655-661, 1988.

[51775] 8245.

[51776] Sapi, E.; Flick, M. B.; Kacinski, B. M.: The first intron of

[51777] human c-fms proto-oncogene contains a processed pseudogene (RPL7P)

[51778] for ribosomal protein L7. Genomics 22: 641-645, 1994.

[51779] 8246.

[51780] Verbeek, J. S.; Roebroek, A. J. M.; van den Ouweland, A. M. W.;

[51781] Bloemers, H. P. J.; Van de Ven, W. J. M.: Human c-fms proto-oncogene:

[51782] comparative analysis with an abnormal allele. Molec. Cell. Biol. 5:

[51783] 422-426, 1985.

[51784] 8247.

[51785] Verbeek, J. S.; van Heerikhuizen, H.; de Pauw, B. E.; Haa-nen,

[51786] C.; Bloemers, H. P. J.; Van de Ven, W. J. M.: A hereditary abnormal

[51787] c-fms proto-oncogene in a patient with acute lymphocytic leukaemia

[51788] and congenital hypothyroidism. Brit. J. Haemat. 61: 135-138, 1985.

[51789] 8248.

[51790] Yarden, Y.; Ullrich, A.: Growth factor receptor tyrosine kinases. Ann.

[51791] Rev. Biochem. 57: 443-478, 1988.

[51792] 8249.

[51793] Brown, J. R.; Ye, H.; Bronson, R. T.; Dikkes, P.; Greenberg, M.

[51794] E.: A defect in nurturing in mice lacking the immediate early gene

[51795] fosB. Cell 86: 297-309, 1996.

[51796] 8250.

[51797] Forsdyke, D.: Personal Communication. Kingston, Ontario, Canada

[51798] 6/8/1992.

[51799] 8251.

[51800] Lazo, P. S.; Dorfman, K.; Noguchi, T.; Mattei, M. G.; Bravo, R.

[51801] : Structure and mapping of the fosB gene: FosB downregulates the activity

[51802] of the fosB promoter. Nucleic Acids Res. 20: 343-350, 1992.

[51803] 8252.

[51804] Nakabeppu, Y.; Nathans, D.: A naturally occurring truncated form

[51805] of FosB that inhibits Fos/Jun transcriptional activity. Cell 64:

[51806] 751-759, 1991.

[51807] 8253.

[51808] Siderovski, D. P.; Blum, S.; Forsdyke, R. E.; Forsdyke, D. R.:

[51809] A set of human putative lymphocyte G0/G1 switch genes includes genes

[51810] homologous to rodent cytokine and zinc finger protein-encoding genes. DNA

[51811] Cell Biol. 9: 579-587, 1990.

[51812] 8254.

[51813] Kelz, M. B.; Chen, J.; Carlezon, W. A., Jr.; Whisler, K.; Gilden,

[51814] L.; Beckmann, A. M.; Steffen, C.; Zhang, Y.-J.; Marotti, L.; Self,

[51815] D. W.; Tkatch, T.; Baranauskas, G.; Surmeier, D. J.; Neve, R. L.;

[51816] Duman, R. S.; Picciotto, M. R.; Nestler, E. J.: Expression of

the

[51817] transcription factor delta-FosB in the brain controls sensi-
tivity

[51818] to cocaine. Nature 401: 272-276, 1999.

[51819] 8255.

[51820] Amaravadi, L. S.; Neff, A. W.; Sleeman, J. P.; Smith, R. C.:
Autonomous

[51821] neural axis formation by ectopic expression of the pro-
tooncogene c-ski. Dev.

[51822] Biol. 192: 392-404, 1997.

[51823] 8256.

[51824] Balazs, I.; Grzeschik, K. H.; Stavnezer, E.: Assignment of
the

[51825] human homologue of a chicken oncogene to chromosome
1. (Abstract) Cytogenet.

[51826] Cell Genet. 37: 410-411, 1984.

[51827] 8257.

[51828] Berk, M.; Desai, S. Y.; Heyman, H. C.; Colmenares, C.: Mice
lacking

[51829] the ski proto-oncogene have defects in neurulation, cran-
iofacial,

[51830] patterning, and skeletal muscle development. Genes Dev. 11: 2029-2039,
[51831] 1997.

[51832] 8258.

[51833] Chaganti, R. S. K.; Balazs, I.; Jhanwar, S. C.; Murty, V. V. V.

[51834] S.; Koduru, P. R. K.; Grzeschik, K.-H.; Stavnezer, E.: The cellular
[51835] homologue of the transforming gene of SKV avian retro- virus maps to
[51836] human chromosome region 1q22-q24. Cytogenet. Cell Genet. 43: 181-186,
[51837] 1986.

[51838] 8259.

[51839] Colmenares, C.; Heilstedt, H. A.; Shaffer, L. G.; Schwartz, S.;
[51840] Berk, M.; Murray, J. C.; Stavnezer, E.: Loss of the SKI proto-oncogene
[51841] in individuals affected with 1p36 deletion syndrome is predicted by
[51842] strain-dependent defects in Ski -/- mice. Nature Genet. 30: 106-109,

[51843] 2002.

[51844] 8260.

[51845] Kaufman, C. D.; Martinez-Rodriguez, G.; Hackett, P. B., Jr.: Ectopic

[51846] expression of c-ski disrupts gastrulation and neural patterning in

[51847] zebrafish. Mech. Dev. 95: 147-162, 2000.

[51848] 8261.

[51849] Nomura, N.; Sasamoto, S.; Ishii, S.; Date, T.; Matsui, M.; Ishizaki,

[51850] R.: Isolation of human cDNA clones of SKI and the SKI-related gene,

[51851] SNO. Nucleic Acids Res. 17: 5489-5500, 1989.

[51852] 8262.

[51853] Shapira, S. K.; McCaskill, C.; Northrup, H.; Spikes, A. S.; Elder,

[51854] F. F. B.; Sutton, V. R.; Korenberg, J. R.; Greenberg, F.; Shaffer,

[51855] L. G.: Chromosome 1p36 deletions: the clinical phenotype and molecular

[51856] characterization of a common newly delineated syndrome.

Am. J. Hum.

[51857] Genet. 61: 642-650, 1997.

[51858] 8263.

[51859] Shinagawa, T.; Nomura, T.; Colmenares, C.; Ohira, M.; Nakagawara,

[51860] A.; Ishii, S.: Increased susceptibility to tumorigenesis of ski-deficient

[51861] heterozygous mice. Oncogene 20: 8100-8108, 2001.

[51862] 8264.

[51863] Slavotinek, A.; Shaffer, L. G.; Shapira, S. K.: Monosomy 1p36. J.

[51864] Med. Genet. 36: 657-663, 1999.

[51865] 8265.

[51866] Sun, Y.; Liu, X.; Eaton, E. N.; Lane, W. S.; Lodish, H. F.; Weinberg,

[51867] R. A.: Interaction of the Ski oncoprotein with Smad3 regulates TGF-beta

[51868] signaling. Molec. Cell 4: 499-509, 1999.

[51869] 8266.

[51870] Sutrave, P.; Kelly, A. M.; Hughes, S. H.: ski can cause se-

lective
[51871] growth of skeletal muscle in transgenic mice. Genes Dev. 4: 1462-1472,
[51872] 1990.

[51873] 8267.

[51874] Boyd, M. T.; Vlatkovic, N.; Haines, D. S.: A novel cellular protein
[51875] (MTBP) binds to MDM2 and induces a G1 arrest that is suppressed by
[51876] MDM2. J. Biol. Chem. 275: 31883-31890, 2000.

[51877] 8268.

[51878] Bueso-Ramos, C. E.; Yang, Y.; deLeon, E.; McCown, P.; Stass, S.
[51879] A.; Albitar, M.: The human MDM-2 oncogene is overex-pressed in leukemias. Blood 82:
[51880] 2617-2623, 1993.

[51881] 8269.

[51882] Buschmann, T.; Fuchs, S. Y.; Lee, C.-G.; Pan, Z.-Q.; Ronai, Z.
[51883] : SUMO-1 modification of Mdm2 prevents its self-ubiquitination and

[51884] increases Mdm2 ability to ubiquitinate p53. Cell 101: 753-762, 2000.

[51885] Note: Erratum: Cell 107: 549 only, 2001.

[51886] 8270.

[51887] Otsu, M.; Hiles, I.; Gout, I.; Fry, M. J.; Ruiz-Larrea, F.; Panayotou,

[51888] G.; Thompson, A.; Dhand, R.; Hsuan, J.; Totty, N.; Smith, A. D.; Morgan,

[51889] S. J.; Courtneidge, S. A.; Parker, P. J.; Waterfield, M. D.: Characterization

[51890] of two 85 kd proteins that associate with receptor tyrosine kinases,

[51891] middle-T/pp60(c-src) complexes, and PI3-kinase. Cell 65: 91-104,

[51892] 1991.

[51893] 8271.

[51894] Volinia, S.; Patracchini, P.; Otsu, M.; Hiles, I.; Gout, I.; Cal-zolari,

[51895] E.; Bernardi, F.; Rooke, L.; Waterfield, M. D.: Chromosomal localization

[51896] of human p85-alpha, a subunit of phosphatidylinositol 3-kinase, and

[51897] its homologue p85-beta. Oncogene 7: 789-793, 1992.

[51898] 8272.

[51899] Hilliker, C. E.; Darville, M. I.; Aly, M. S.; Chikri, M.; Szpirer,

[51900] C.; Marynen, P.; Rousseau, G. G.; Cassiman, J.-J.: Human and rat

[51901] chromosomal localization of two genes for 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase

[51902] by analysis of somatic cell hybrids and in situ hybridiza- tion. Genomics 10:

[51903] 867-873, 1991.

[51904] 8273.

[51905] Hu, Z.; Bonifas, J. M.; Beech, J.; Bench, G.; Shigihara, T.; Ogawa,

[51906] H.; Ikeda, S.; Mauro, T.; Epstein, E. H., Jr.: Mutations in ATP2C1,

[51907] encoding a calcium pump, cause Hailey-Hailey disease. Nature Genet. 24:

[51908] 61-65, 2000.

[51909] 8274.

[51910] Steinkasserer, A.; Cockburn, D. J.; Black, D. M.; Boyd, Y.; Solomon,

[51911] E.; Sim, R. B.: Assignment of apolipoprotein H (APOH: beta-2-glycoprotein
[51912] I) to human chromosome 17q23-qter; determination of the major expression
[51913] site. Cytogenet. Cell Genet. 60: 31-33, 1992.

[51914] 8275.

[51915] Steinkasserer, A.; Dorner, C.; Wurzner, R.; Sim, R. B.: Hu-man
[51916] beta-2-glycoprotein I: molecular analysis of DNA and amino acid polymorphism. Hum.
[51917] Genet. 91: 401-402, 1993.

[51918] 8276.

[51919] Walter, H.; Hilling, M.; Brachtel, R.; Hitzeroth, H. W.: On the
[51920] population genetics of beta-2-glycoprotein I. Hum. Hered. 29: 236-241,
[51921] 1979.

[51922] 8277.

[51923] Budarf, M. L.; Konkle, B. A.; Ludlow, L. B.; Michaud, D.; Li, M.;
[51924] Yamashiro, D. J.; McDonald-McGinn, D.; Zackai, E. H.;

Driscoll, D.

[51925] A.: Identification of a patient with Bernard-Soulier syndrome and

[51926] a deletion in the DiGeorge/velo-cardio-facial chromosomal region in

[51927] 22q11.2. Hum. Molec. Genet. 4: 763-766, 1995.

[51928] 8278.

[51929] Zieger, B.; Hashimoto, Y.; Ware, J.: Alternative expression of

[51930] platelet glycoprotein Ib-beta mRNA from an adjacent 5-prime gene with

[51931] an imperfect polyadenylation signal sequence. J. Clin. Invest. 99:

[51932] 520-525, 1997.

[51933] 8279.

[51934] Bakker, E.; Pearson, P. L.; Meera Khan, P.; Schreuder, G. M. T.;

[51935] Madan, K.: Orientation of major histocompatibility (MHC) genes relative

[51936] to the centromere of human chromosome 6. Clin. Genet. 15: 198-202,

[51937] 1979.

[51938] 8280.

[51939] Goldman, D.; O'Brien, S. J.; Lucas-Derse, S.; Dean, M.: Linkage

[51940] mapping of human polymorphic proteins identified by two-dimensional

[51941] electrophoresis. Genomics 11: 875-884, 1991.

[51942] 8281.

[51943] Ducros, A.; Denier, C.; Joutel, A.; Vahedi, K.; Michel, A.; Darcel,

[51944] F.; Madigand, M.; Guerouaou, D.; Tison, F.; Julien, J.; Hirsch, E.;

[51945] Chedru, F.; Bisgard, C.; Lucotte, G.; Despres, P.; Billard, C.; Barthez,

[51946] M. A.; Ponsot, G.; Bousser, M. G.; Tournier-Lasserve, E.: Recurrence

[51947] of the T666M calcium channel CACNA1A gene mutation in familial hemiplegic

[51948] migraine with progressive cerebellar ataxia. Am. J. Hum. Genet. 64:

[51949] 89-98, 1999.

[51950] 8282.

[51951] Andre, P.; Srinivasa Prasad, K. S.; Denis, C. V.; He, M.; Papalia,

[51952] J. M.; Hynes, R. O.; Phillips, D. R.; Wagner, D. D.: CD40L stabilizes

[51953] arterial thrombi by a beta(3) integrin-dependent mechanism. Nature

[51954] Med. 8: 247-252, 2002.

[51955] 8283.

[51956] Roninson, I. B.; Chin, J. E.; Choi, K.; Gros, P.; Housman, D.

[51957] E.; Fojo, A.; Shen, D.; Gottesman, M. M.; Pastan, I.: Isolation of

[51958] human mdr DNA sequences amplified in multidrug-resistant KB carcinoma

[51959] cells. Proc. Nat. Acad. Sci. 83: 4538-4542, 1986.

[51960] 8284.

[51961] Synold, T. W.; Dussault, I.; Forman, B. M.: The orphan nuclear

[51962] receptor SXR coordinately regulates drug metabolism and efflux. Nature

[51963] Med. 7: 584-590, 2001.

[51964] 8285.

[51965] Alonso, E. M.; Snover, D. C.; Montag, A.; Freese, D. K.; Whitington,
[51966] P. F.: Histologic pathology of the liver in progressive familial
[51967] intrahepatic cholestasis. J. Pediat. Gastroent. Nutr. 18: 128-133,
[51968] 1994.

[51969] 8286.

[51970] Deleuze, J. F.; Jacquemin, E.; Dubuisson, C.; Cresteil, D.; Dumont,
[51971] M.; Erlinger, S.; Bernard, O.; Hadchouel, M.: Defect of multidrug-resistance
[51972] 3 gene expression in a subtype of progressive familial intrahepatic
[51973] cholestasis. Hepatology 23: 904-908, 1996.

[51974] 8287.

[51975] Woods, K. A.; Camacho-Hubner, C.; Bergman, R. N.; Barter, D.;
[51976] Clark, A. J. L.; Savage, M. O.: Effects of insulin-like growth factor
[51977] I (IGF-I) therapy on body composition and insulin resistance in IGF-I

[51978] gene deletion. J. Clin. Endocr. Metab. 85: 1407-1411, 2000.

[51979] 8288.

[51980] Woods, K. A.; Camacho-Hubner, C.; Savage, M. O.; Clark, A. J.

[51981] L.: Intrauterine growth retardation and postnatal growth failure

[51982] associated with deletion of the insulin-like growth factor I gene. New

[51983] Eng. J. Med. 335: 1363-1367, 1996.

[51984] 8289.

[51985] Yang-Feng, T. L.; Brissenden, J. E.; Ullrich, A.; Francke, U.

[51986] : Sub-regional localization of human genes for insulin-like growth

[51987] factors I (IGF1) and II (IGF2) by in situ hybridization. (Abstract) Cytogenet.

[51988] Cell Genet. 40: 782 only, 1985.

[51989] 8290.

[51990] Yanovski, J. A.; Sovik, K. N.; Nguyen, T. T.; Sebring, N. G.:

[51991] Insulin-like growth factors and bone mineral density in African American

[51992] and white girls. J. Pediat. 137: 826-832, 2000.

[51993] 8291.

[51994] Zhu, J.; Kahn, C. R.: Analysis of a peptide hormone-receptor

[51995] interaction in the yeast two-hybrid system. Proc. Nat. Acad. Sci. 94:

[51996] 13063-13068, 1997.

[51997] 8292.

[51998] Nishi, S.; Seino, S.; Bell, G. I.: Human hexokinase: sequences

[51999] of amino- and carboxyl-terminal halves are homologous. Biochem. Biophys.

[52000] Res. Commun. 157: 937-943, 1988.

[52001] 8293.

[52002] Rijksen, G.; Akkerman, J. W. N.; van den Wall Bake, A. W. L.;

[52003] Pott Hofstede, D.; Staal, G. E. J.: Generalized hexokinase deficiency

[52004] in the blood cells of a patient with nonspherocytic hemolytic anemia. Blood 61:

[52005] 12-18, 1983.

[52006] 8294.

[52007] Ritter, H.; Friedrichson, U.; Schmitt, J.: Genetic polymor-phism

[52008] of hexokinase in primates. Humangenetik 22: 265-266, 1974.

[52009] 8295.

[52010] Rogers, P. A.; Fisher, R. A.; Harris, H.: An electrophoretic

[52011] study of the distribution and properties of human hexoki-nases. Biochem.

[52012] Genet. 13: 857-866, 1975.

[52013] 8296.

[52014] Ruzzo, A.; Andreoni, F.; Magnani, M.: Structure of the human

[52015] hexokinase type I gene and nucleotide sequence of the 5-prime flanking

[52016] region. Biochem. J. 331: 607-613, 1998.

[52017] 8297.

[52018] Schimke, R. T.; Grossbard, L.: Studies on isozymes of hexokinase

[52019] in animal tissues. Ann. N.Y. Acad. Sci. 151: 332-350, 1968.

[52020] 8298.

[52021] Shows, T. B.: Synteny of human genes for glutamic oxaloacetic

[52022] transaminase and hexokinase in somatic cell hybrids. Cytogenet. Cell

[52023] Genet. 13: 143-145, 1974.

[52024] 8299.

[52025] Shows, T. B.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry,

[52026] W. M.; Nishi, S.; Bell, G. I.: Localization of the human hexokinase

[52027] I gene (HK1) to chromosome 10q22. (Abstract) Cytogenet. Cell Genet. 51:

[52028] 1079 only, 1989.

[52029] 8300.

[52030] Ureta, T.: The comparative isozymology of vertebrate hexokinases. Comp.

[52031] Biochem. Physiol. 71B: 549-555, 1982.

[52032] 8301.

[52033] Kobayashi, M.; Takamatsu, K.; Saitoh, S.; Miura, M.; Noguchi, T.

[52034] : Molecular cloning of hippocalcin, a novel calcium-binding protein

[52035] of the recoverin family exclusively expressed in hippocampus. Biochem.

[52036] Biophys. Res. Commun. 189: 511-517, 1992.

[52037] 8302.

[52038] Takamatsu, K.; Kobayashi, M.; Saitoh, S.; Fujishiro, M.; Noguchi,

[52039] T.: Molecular cloning of human hippocalcin cDNA and chromosomal mapping

[52040] of its gene. Biochem. Biophys. Res. Commun. 200:

606-611, 1994.

[52041] 8303.

[52042] Borrego, S.; Ruiz, A.; Saez, M. E.; Gimm, O.; Gao, X.; Lopez-Alonso,

[52043] M.; Hernandez, A.; Wright, F. A.; Antinolo, G.; Eng, C.: RET genotypes

[52044] comprising specific haplotypes of polymorphic variants predispose

[52045] to isolated Hirschsprung disease. J. Med. Genet. 37: 572-578, 2000.

[52046] 8304.

[52047] Borrego, S.; Saez, M. E.; Ruiz, A.; Gimm, O.; Lopez-Alonso, M.;

[52048] Antinolo, G.; Eng, C.: Specific polymorphisms in the RET proto-oncogene

[52049] are over-represented in patients with Hirschsprung dis- ease and may

[52050] represent loci modifying phenotypic expression. J. Med. Genet. 36:

[52051] 771-774, 1999.

[52052] 8305.

[52053] Goldstein, J. L.; Brown, M. S.; Stone, N. J.: Genetics of the

[52054] LDL receptor: evidence that the mutations affecting binding and internalization
[52055] are allelic. Cell 12: 629-641, 1977.

[52056] 8306.

[52057] Goldstein, J. L.; Dana, S. E.; Brunschede, G. Y.; Brown, M. S.
[52058] : Genetic heterogeneity in familial hypercholesterolemia: evidence
[52059] for two different mutations affecting functions of low-density lipoprotein
[52060] receptor. Proc. Nat. Acad. Sci. 72: 1092-1096, 1975.

[52061] 8307.

[52062] Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.: Molecular genetics
[52063] of the LDL receptor gene in familial hypercholesterolemia. Hum. Mutat. 1:
[52064] 445-466, 1992.

[52065] 8308.

[52066] Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.; Russell, D. W.:

[52067] Deletion of exon encoding cysteine-rich repeat of low density lipoprotein

[52068] receptor alters its binding specificity in a subject with familial
[52069] hypercholesterolemia. J. Biol. Chem. 261: 13114-13120, 1986.
[52070] 8309.

[52071] Hobbs, H. H.; Russell, D. W.; Brown, M. S.; Goldstein, J. L.:
[52072] The LDL receptor locus in familial hypercholesterolemia: mutational
[52073] analysis of a membrane protein. Annu. Rev. Genet. 24: 133-170, 1990.
[52074] 8310.

[52075] Horsthemke, B.; Beisiegel, U.; Dunning, A.; Havinga, J. R.; Williamson,
[52076] R.; Humphries, S.: Unequal crossing-over between two Alu-repetitive
[52077] DNA sequences in the low-density-lipoprotein-receptor gene: a possible
[52078] mechanism for the defect in a patient with familial hypercholesterolaemia. Europ.
[52079] J. Biochem. 164: 77-81, 1987.

[52080] 8311.

[52081] Khachadurian, A. K.: The inheritance of essential familial hypercholesterolemia. Am.

[52082] J. Med. 37: 402-407, 1964.

[52083] 8312.

[52084] Kingsley, D. M.; Krieger, M.: Receptor-mediated endocytosis of

[52085] low density lipoprotein: somatic cell mutants define multiple genes

[52086] required for expression of surface-receptor activity. Proc. Nat.

[52087] Acad. Sci. 81: 5454-5458, 1984.

[52088] 8313.

[52089] Knight, B. L.; Gavigan, S. J. P.; Soutar, A. K.; Patel, D. D.

[52090] : Defective processing and binding of low-density lipoprotein receptors

[52091] in fibroblasts from a familial hypercholesterolaemic subject. Europ.

[52092] J. Biochem. 179: 693-698, 1989.

[52093] 8314.

[52094] Knoblauch, H.; Muller-Myhsok, B.; Busjahn, A.; Ben Avi, L.; Bahring,

[52095] S.; Baron, H.; Heath, S. C.; Uhlmann, R.; Faulhaber, H.-D.; Shpitzen,
[52096] S.; Aydin, A.; Reshef, A.; and 11 others: A cholesterol-lowering
[52097] gene maps to chromosome 13q. Am. J. Hum. Genet. 66: 157-166, 2000.
[52098] 8315.

[52099] Komuro, I.; Kato, H.; Nakagawa, T.; Takahashi, K.; Mimori, A.;
[52100] Takeuchi, F.; Nishida, Y.; Miyamoto, T.: Case report: the longest-lived
[52101] patient with homozygous familial hypercholesterolemia secondary to
[52102] a defect in internalization of the LDL receptor. Am. J. Med. Sci. 294:
[52103] 341-345, 1987.

[52104] 8316.

[52105] Kotze, M. J.; Langenhoven, E.; Retief, A. E.; Steyn, K.; Marais,
[52106] M. P.; Grobbelaar, J. J.; Oosthuizen, C. J. J.; Weich, H. F. H.; Benade,
[52107] A. J. S.: Haplotype associations of three DNA polymor-

phisms at the
[52108] human low density lipoprotein receptor gene locus in familial hypercholesterolemia. J.
[52109] Med. Genet. 24: 750-755, 1987.

[52110] 8317.

[52111] Grundy, H. O.; Peltz, G.; Moore, K. W.; Golbus, M. S.; Jackson,
[52112] L. G.; Lebo, R. V.: The polymorphic Fc-gamma receptor II gene maps
[52113] to human chromosome 1q. Immunogenetics 29: 331-339, 1989.
[52114] 8318.

[52115] Kudo, A.; Melchers, F.: A second gene, Vpre-B in the lambda 5
[52116] locus of the mouse, which appears to be selectively expressed in pre-B
[52117] lymphocytes. EMBO J. 6: 2267-2272, 1987.

[52118] 8319.

[52119] Mattei, M.-G.; Fumoux, F.; Roeckel, N.; Fougereau, M.; Schiff,
[52120] C.: The human pre-B-specific lambda-like cluster is lo-

cated in the
[52121] 22q11.2-22q12.3 region, distal to the IgC-lambda locus. Genomics 9:
[52122] 544-546, 1991.

[52123] 8320.

[52124] Minegishi, Y.; Coustan-Smith, E.; Wang, Y.-H.; Cooper, M. D.; Campana,
[52125] D.; Conley, M. E.: Mutations in the human lambda-5/14.1 gene result
[52126] in B cell deficiency and agammaglobulinemia. J. Exp. Med. 187: 71-77,
[52127] 1998.

[52128] 8321.

[52129] Pillai, S.; Baltimore, D.: The omega and iota surrogate im-munoglobulin
[52130] light chains. Curr. Top. Microbiol. Immun. 137: 136-139, 1988.
[52131] 8322.

[52132] Sakaguchi, N.; Melchers, F.: Lambda 5, a new light-chain-related
[52133] locus selectively expressed in pre-B lymphocytes. Nature

324: 579-582,

[52134] 1986.

[52135] 8323.

[52136] Schiff, C.; Milili, M.; Fougereau, M.: Isolation of early immunoglobulin

[52137] lambda-like gene transcripts in human fetal liver. Europ. J. Immun. 19:

[52138] 1873-1878, 1989.

[52139] 8324.

[52140] Lin, P.-F.; Slate, D. L.; Lawyer, F. C.; Ruddle, F. H.: Assignment

[52141] of the murine interferon sensitivity and cytoplasmic superoxide dismutase

[52142] genes to chromosome 16. Science 209: 285-287, 1980.

[52143] 8325.

[52144] Holmes, W. E.; Lee, J.; Kuang, W.-J.; Rice, G. C.; Wood, W. I.

[52145] : Structure and functional expression of a human interleukin-8 receptor. Science 253:

[52146] 1278-1280, 1991.

[52147] 8326.

[52148] Nishimura, D.; Buetow, K. H.; Yamada, Y.; Murray, J. C.: RFLPs
[52149] and linkage relationships of the human laminin B2 gene. Genomics 3:
[52150] 393-395, 1988.

[52151] 8327.

[52152] Juppner, H.: Personal Communication. Boston, Mass. 6/26/1995.
[52153] 8328.

[52154] Karaplis, A. C.; Luz, A.; Glowacki, J.; Bronson, R. T.; Ty-
bulewicz,
[52155] V. L. J.; Kronenberg, H. M.; Mulligan, R. C.: Lethal skeletal dysplasia
[52156] from targeted disruption of the parathyroid hormone-re-
lated peptide
[52157] gene. Genes Dev. 8: 277-289, 1994.

[52158] 8329.

[52159] Schipani, E.; Kruse, K.; Juppner, H.: A constitutively active
[52160] mutant PTH-PTHrP receptor in Jansen-type metaphyseal chondrodysplasia. Science 268:
[52161] 98-100, 1995.

[52162] 8330.

[52163] Drayna, D.; Fielding, C.; McLean, J.; Baer, B.; Castro, G.; Chen,

[52164] E.; Comstock, L.; Henzel, W.; Kohr, W.; Rhee, L.; Wion, K.; Lawn,

[52165] R.: Cloning and expression of human apolipoprotein D cDNA. J. Biol.

[52166] Chem. 261: 16535-16539, 1986.

[52167] 8331.

[52168] Drayna, D.; Scott, J. D.; Lawn, R.: Multiple RFLPs at the human

[52169] apolipoprotein D (APOD) locus. Nucleic Acids Res. 15: 9617 only,

[52170] 1987.

[52171] 8332.

[52172] Mattei, M. G.; Bruce, B.; Karsenty, G.: Mouse alpha-1 type V

[52173] collagen gene maps to the [A2-B] region of chromosome 2. Genomics 16:

[52174] 786-788, 1993.

[52175] 8333.

[52176] Nicholls, A. C.; McCarron, S.; Narcisi, P.; Pope, F. M.: Molecular
[52177] abnormalities of type V collagen in Ehlers Danlos syndrome (Abstract) Am.
[52178] J. Hum. Genet. 55 (suppl.): A233, 1994.

[52179] 8334.

[52180] Nicholls, A. C.; Oliver, J. E.; McCarron, S.; Harrison, J. B.;

[52181] Greenspan, D. S.; Pope, F. M.: An exon skipping mutation of a type
[52182] V collagen gene (COL5A1) in Ehlers-Danlos syndrome. J. Med. Genet. 33:
[52183] 940-946, 1996.

[52184] 8335.

[52185] Schwarze, U.; Atkinson, M.; Hoffman, G. G.; Greenspan, D. S.;
[52186] Byers, P. H.: Null alleles of the COL5A1 gene of type V collagen
[52187] are a cause of the classical forms of Ehlers-Danlos syndrome (types
[52188] I and II). Am. J. Hum. Genet. 66: 1757-1765, 2000.

[52189] 8336.

[52190] Takahara, K.; Hoffman, G. G.; Greenspan, D. S.: Complete structural
[52191] organization of the human alpha-1(V) collagen gene (COL5A1): divergence
[52192] from the conserved organization of other characterized fibrillar collagen
[52193] genes. Genomics 29: 588-597, 1995.

[52194] 8337.

[52195] Takahara, K.; Sato, Y.; Okazawa, K.; Okamoto, N.; Noda, A.; Yaoi,
[52196] Y.; Kato, I.: Complete primary structure of human collagen alpha-1(V)
[52197] chain. J. Biol. Chem. 266: 13124-13129, 1991.

[52198] 8338.

[52199] Baas, F.; Bikker, H.; Van Ommen, G.-J. B.; de Vijlder, J. J. M.
[52200] : Unusual scarcity of restriction site polymorphism in the human thyroglobulin
[52201] gene: a linkage study suggesting autosomal dominance of a defective
[52202] thyroglobulin allele. Hum. Genet. 67: 301-305, 1984.

[52203] 8339.

[52204] Takahara, K.; Schwarze, U.; Imamura, Y.; Hoffman, G. G.; Toriello,

[52205] H.; Smith, L. T.; Byers, P. H.; Greenspan, D. S.: Order of intron

[52206] removal influences multiple splice outcomes, including a two-exon

[52207] skip, in a COL5A1 acceptor-site mutation that results in abnormal

[52208] pro-alpha-1(V) N-propeptides and Ehlers-Danlos syndrome type I. Am.

[52209] J. Hum. Genet. 71: 451-465, 2002.

[52210] 8340.

[52211] Wenstrup, R. J.; Florer, J. B.; Willing, M. C.; Giunta, C.; Steinmann,

[52212] B.; Young, F.; Susic, M.; Cole, W. G.: COL5A1 haploinsufficiency

[52213] is a common molecular mechanism underlying the classical form of EDS. Am.

[52214] J. Hum. Genet. 66: 1766-1776, 2000.

[52215] 8341.

[52216] Imamura, Y.; Scott, I. C.; Greenspan, D. S.: The pro-alpha3(V)
[52217] collagen chain: complete primary structure, expression domains in
[52218] adult and developing tissues, and comparison to the structures and
[52219] expression domains of the other types V and XI procollagen chains. J.
[52220] Biol. Chem. 275: 8749-8759, 2000.

[52221] 8342.

[52222] Bonaldo, P.; Braghetta, P.; Zanetti, M.; Piccolo, S.; Volpin, D.;
[52223] Bressan, G. M.: Collagen VI deficiency induces early onset myopathy
[52224] in the mouse: an animal model for Bethlem myopathy. Hum. Molec. Genet. 7:
[52225] 2135-2140, 1998.

[52226] 8343.

[52227] Chu, M.-L.; Mann, K.; Deutzmann, R.; Pribula-Conway, D.; Hsu-Chen,
[52228] C.-C.; Bernard, M. P.; Timpl, R.: Characterization of three constituent

[52229] chains of collagen type VI by peptide sequences and cDNA clones. Europ.

[52230] J. Biochem. 168: 309-317, 1987.

[52231] 8344.

[52232] Chung, E.; Rhodes, R. K.; Miller, E. J.: Isolation of three collagenous

[52233] components of probable basement membrane origin from several tissues. Biochem.

[52234] Biophys. Res. Commun. 71: 1167-1174, 1976.

[52235] 8345.

[52236] Cutting, G.; Francomano, C. A.; Chu, M. L.; Timpl, R.; Mc-Cormick,

[52237] M. K.; Warren, A. C.; Hong, H. K.; Pyeritz, R. E.; An-tonarakis, S.

[52238] E.: Genetic linkage analysis and macrorestriction mapping of COL6A1

[52239] and COL6A2, structural genes of type VI collagen. (Abstract) Am.

[52240] J. Hum. Genet. 43: A141 only, 1988.

[52241] 8346.

[52242] Davies, G. E.; Howard, C. M.; Farrer, M. J.; Coleman, M. M.;

Bennett,
[52243] L. B.; Cullen, L. M.; Wyse, R. K. H.; Burn, J.; Williamson, R.; Kessling,
[52244] A. M.: Genetic variation in the COL6A1 region is associated with
[52245] congenital heart defects in trisomy 21 (Down's syndrome). Ann. Hum.
[52246] Genet. 59: 253-269, 1995.

[52247] 8347.

[52248] Delabar, J.-M.; Chettouh, Z.; Rahmani, Z.; Theophile, D.; Blouin,
[52249] J.-L.; Bono, R.; Kraus, J.; Barton, J.; Patterson, D.; Sinet, P.-M.
[52250] : Gene-dosage mapping of 30 DNA markers on chromosome 21. Genomics 13:
[52251] 887-889, 1992.

[52252] 8348.

[52253] Engel, J.; Furthmayr, H.; Odermatt, E.; Von der Mark, H.; Aumailley,
[52254] M.; Fleishmajer, R.; Timpl, R.: Structure and macromolecular organization
[52255] of type VI collagen. Ann. N.Y. Acad. Sci. 460: 25-37,

1985.

[52256] 8349.

[52257] Francomano, C. A.; Cutting, G. R.; McCormick, M. K.; Chu, M. L.;

[52258] Timpl, R.; Hong, H. K.; Antonarakis, S. E.: The COL6A1 and COL6A2

[52259] genes exist as a gene cluster and detect highly informa-tive DNA polymorphisms

[52260] in the telomeric region of human chromosome 21q. Hum. Genet. 87:

[52261] 162-166, 1991.

[52262] 8350.

[52263] Furuto, D. K.; Miller, E. J.: Characterization of a unique collagenous

[52264] fraction from limited pepsin digests of human placental tissue: molecular

[52265] organization of the native aggregate. Biochemistry 20: 1635-1640,

[52266] 1981.

[52267] 8351.

[52268] Heiskanen, M.; Saitta, B.; Palotie, A.; Chu, M.-L.: Head to

tail

[52269] organization of the human COL6A1 and COL6A2 genes by fiber-FISH. Genomics 29:

[52270] 801-803, 1995.

[52271] 8352.

[52272] Hessle, H.; Engvall, E.: Type VI collagen: studies on its lo-

[52273] calization, structure, and biosynthetic form with monoclonal anti-

[52274] bodies. J. Biol. Chem. 259: 3955-3961, 1984.

[52275] 8353.

[52276] Jander, R.; Rauterberg, J.; Voss, B.; von Bassewitz, D. B.: A

[52277] cysteine-rich collagenous protein from bovine placenta: isolation

[52278] of its constituent polypeptide chains and some properties of the non-denatured

[52279] protein. Europ. J. Biochem. 114: 17-25, 1981.

[52280] 8354.

[52281] Agar, N. S.; Board, P. G.; Bell, K.: Studies of erythrocyte glyoxalase

[52282] II in various domestic species: discovery of glyoxalase II

deficiency
[52283] in the horse. Animal Blood Groups Biochem. Genet. 15: 67-70, 1984.
[52284] 8355.
[52285] Ball, J. C.; Vander Jagt, D. L.: Purification of S-2-hydroxyacylglutathione
[52286] hydrolase (glyoxalase II) from rat erythrocytes. Anal. Biochem. 98:
[52287] 472-477, 1979.
[52288] 8356.
[52289] Board, P. G.: Genetic polymorphism of human erythrocyte glyoxalase
[52290] II. Am. J. Hum. Genet. 32: 690-694, 1980.
[52291] 8357.
[52292] Board, P. G.; Gibbs, C. J., Jr.; Gajdusek, D. C.: Polymor-phism
[52293] of erythrocyte glyoxalase II in anthropoid primates. Folia Primatol. 36:
[52294] 138-143, 1981.
[52295] 8358.
[52296] Honey, N. K.; Shows, T. B.: Assignment of the glyoxalase II

gene
[52297] (HAGH) to human chromosome 16. Hum. Genet. 58:
358-361, 1981.
[52298] 8359.

[52299] Mulley, J. C.; Barton, N.; Callen, D. F.: Localisation of hu-
man
[52300] PGP and HAGH genes to 16p13.3. Cytogenet. Cell Genet.
53: 175-176,
[52301] 1990.

[52302] 8360.

[52303] Mulley, J. C.; Callen, D. F.: New regional localisations for
HAGH
[52304] and PGP on human chromosome 16. Hum. Genet. 74:
423-424, 1986.
[52305] 8361.

[52306] Ridderstrom, M.; Saccucci, F.; Hellman, U.; Bergman, T.;
Principato,
[52307] G.; Mannervik, B.: Molecular cloning, heterologous ex-
pression, and
[52308] characterization of human glyoxalase II. J. Biol. Chem.
271: 319-323,
[52309] 1996.

[52310] 8362.

[52311] Valentine, W. N.; Paglia, D. E.; Neerhout, R. C.; Konrad, P. N.

[52312] : Erythrocyte glyoxalase II deficiency with coincidental hereditary

[52313] elliptocytosis. Blood 36: 797-808, 1970.

[52314] 8363.

[52315] Beranova, M.; Oliveira, L. M. B.; Bedecarrats, G. Y.; Schipani,

[52316] E.; Vallejo, M.; Ammini, A. C.; Quintos, J. B.; Hall, J. E.; Martin,

[52317] K. A.; Hayes, F. J.; Pitteloud, N.; Kaiser, U. B.; Crowley, W. F.,

[52318] Jr.; Seminara, S. B.: Prevalence, phenotypic spectrum, and modes

[52319] of inheritance of gonadotropin-releasing hormone receptor mutations

[52320] in idiopathic hypogonadotropic hypogonadism. J. Clin. Endocr. Metab. 86:

[52321] 1580-1588, 2001.

[52322] 8364.

[52323] Caron, P.; Chauvin, S.; Christin-Maitre, S.; Bennet, A.; Lahlou,
[52324] N.; Counis, R.; Bouchard, P.; Kottler, M.-L.: Resistance of hypogonadic
[52325] patients with mutated GnRH receptor genes to pulsatile GnRH administration. J.
[52326] Clin. Endocr. Metab. 84: 990-996, 1999.

[52327] 8365.

[52328] Costa, E. M. F.; Bedecarrats, G. Y.; Mendonca, B. B.; Arn-
hold,
[52329] I. J. P.; Kaiser, U. B.; Latronico, A. C.: Two novel mutations in
[52330] the gonadotropin- releasing hormone receptor gene in Brazilian patients
[52331] with hypogonadotropic hypogonadism and normal olfac-
tion. J. Clin.
[52332] Endocr. Metab. 86: 2680-2686, 2001.

[52333] 8366.

[52334] de Roux, N.; Young, J.; Brailly-Tabard, S.; Misrahi, M.; Mil-
grom,
[52335] E.; Schaison, G.: The same molecular defects of the go-
nadotropin-releasing

[52336] hormone receptor determine a variable degree of hypogonadism in affected
[52337] kindred. J. Clin. Endocr. Metab. 84: 567-572, 1999.

[52338] 8367.

[52339] de Roux, N.; Young, J.; Misrahi, M.; Genet, R.; Chanson, P.; Schaison,
[52340] G.; Milgrom, E.: A family with hypogonadotropic hypogonadism and
[52341] mutations in the gonadotropin-releasing hormone receptor. New Eng.
[52342] J. Med. 337: 1597-1602, 1997.

[52343] 8368.

[52344] Fan, N. C.; Jeung, E.-B.; Peng, C.; Olofsson, J. I.; Krisinger,
[52345] J.; Leung, P. C. K.: The human gonadotropin-releasing hormone (GnRH)
[52346] receptor gene: cloning, genomic organization and chromosomal assignment. Molec.
[52347] Cell. Endocr. 103: R1-R6, 1994.

[52348] 8369.

[52349] Hasselback, R.; Marion, R. B.; Thomas, J. W.: Congenital hypofibrinogenemia

[52350] in five members of a family. Canad. Med. Assoc. J. 88: 19-22, 1963.

[52351] 8370.

[52352] Iwashita, T.; Murakami, H.; Asai, N.; Takahashi, M.: Mech-
anism
[52353] of Ret dysfunction by Hirschsprung mutations affecting its extracellular
[52354] domain. Hum. Molec. Genet. 5: 1577-1580, 1996.

[52355] 8371.

[52356] Lehrman, M. A.; Goldstein, J. L.; Brown, M. S.; Russell, D. W.;
[52357] Schneider, W. J.: Internalization-defective LDL receptors produced
[52358] by genes with nonsense and frameshift mutations that truncate the
[52359] cytoplasmic domain. Cell 41: 735-743, 1985.

[52360] 8372.

[52361] Nezer, C.; Moreau, L.; Brouwers, B.; Coppieters, W.; De-
tilleux,
[52362] J.; Hanset, R.; Karim, L.; Kvasz, A.; Leroy, P.; Georges, M.: An

[52363] imprinted QTL with major effect on muscle mass and fat deposition

[52364] maps to the IGF2 locus in pigs. (Letter) Nature Genet. 21: 155-156,

[52365] 1999.

[52366] 8373.

[52367] Bird, A. P.: CpG islands as gene markers in the vertebrate nucleus. Trends

[52368] Genet. 3: 342-347, 1987.

[52369] 8374.

[52370] Maartmann-Moe, K.; Wang, H. S.; Donald, L. J.; Hamerton, J. L.;

[52371] Berg, K.: Data from hybrid cell lines raise the possibility that

[52372] factors controlling the low density lipoprotein receptor activity

[52373] may reside on human chromosome 21, 5 or both. (Abstract) Cytogenet.

[52374] Cell Genet. 32: 295-296, 1982.

[52375] 8375.

[52376] Miyake, Y.; Tajima, S.; Funahashi, T.; Yamamoto, A.: Anal-

ysis
[52377] of a recycling-impaired mutant of low density lipoprotein receptor
[52378] in familial hypercholesterolemia. J. Biol. Chem. 264: 16584-16590,
[52379] 1989.

[52380] 8376.

[52381] Ott, J.; Schrott, H. G.; Goldstein, J. L.; Hazzard, W. R.; Allen,
[52382] F. H.; Falk, C. T.; Motulsky, A. G.: Linkage studies in a large kindred
[52383] with familial hypercholesterolemia. Am. J. Hum. Genet. 26: 598-603,
[52384] 1974.

[52385] 8377.

[52386] Seftel, H. C.; Baker, S. G.; Sandler, M. P.; Forman, M. B.; Joffe,
[52387] B. I.; Mendelsohn, D.; Jenkins, T.; Mieny, C. J.: A host of hypercholesterolaemic
[52388] homozygotes in South Africa. Brit. Med. J. 281: 633-636, 1980.
[52389] 8378.

[52390] Grundy, H. O.; Peltz, G.; Barsh, G.; Moore, K.; Golbus, M. S.;
[52391] Lebo, R. V.: Immunoglobulin G Fc receptor II and Fc receptor III
[52392] genes map to chromosome 1 by spot-blot chromosome analysis. (Abstract) Am.
[52393] J. Hum. Genet. 43: A145, 1988.

[52394] 8379.

[52395] Hibbs, M. L.; Bonadonna, L.; Scott, B. M.; McKenzie, I. F. C.;
[52396] Hogarth, P. M.: Molecular cloning of a human immunoglobulin G Fc
[52397] receptor. Proc. Nat. Acad. Sci. 85: 2240-2244, 1988.

[52398] 8380.

[52399] Hibbs, M. L.; Hogarth, P. M.; McKenzie, I. F. C.: The mouse Ly-17
[52400] locus identifies a polymorphism of the Fc receptor. Immunogenetics 22:
[52401] 335-348, 1985.

[52402] 8381.

[52403] Lebo, R. V.; Lynch, E. D.; Wiegant, J.; Moore, K.; Troun-

stine,
[52404] M.; van der Ploeg, M.: Multicolor fluorescence in situ hybridization
[52405] and pulsed field electrophoresis dissect CMT1B gene region. Hum.
[52406] Genet. 88: 13-20, 1991.

[52407] 8382.

[52408] Moser, K. L.; Neas, B. R.; Salmon, J. E.; Yu, H.; Gray-McGuire,
[52409] C.; Asundi, N.; Bruner, G. R.; Fox, J.; Kelly, J.; Henshall, S.; Bacino,
[52410] D.; Dietz, M.; Hogue, R.; Koelsch, G.; Nightingale, L.; Shaver, T.;
[52411] Abdou, N. I.; Albert, D. A.; Carson, C.; Petri, M.; Treadwell, E.
[52412] L.; James, J. A.; Harley, J. B.: Genome scan of human systemic lupus
[52413] erythematosus: evidence for linkage on chromosome 1q in African-American
[52414] pedigrees. Proc. Nat. Acad. Sci. 95: 14869-14874, 1998.

[52415] 8383.

[52416] Oakey, R. J.; Watson, M. L.; Seldin, M. F.: Construction of a

[52417] physical map on mouse and human chromosome 1: comparison of 13 Mb
[52418] of mouse and 11 Mb of human DNA. Hum. Molec. Genet. 1: 613-620,
[52419] 1992.

[52420] 8384.

[52421] Peltz, G. A.; Grundy, H. O.; Lebo, R. V.; Yssel, H.; Barsh, G.
[52422] S.; Moore, K. W.: Human Fc-gamma-RIII: cloning, expression, and identification
[52423] of the chromosomal locus of two Fc receptors for IgG. Proc. Nat.
[52424] Acad. Sci. 86: 1013-1017, 1989.

[52425] 8385.

[52426] Qiu, W. Q.; de Bruin, D.; Brownstein, B. H.; Pearse, R.; Ravetch,
[52427] J. V.: Organization of the human and mouse low-affinity Fc-gamma-R
[52428] genes: duplication and recombination. Science 248: 732-735, 1990.
[52429] 8386.

[52430] Salmon, J. E.; Millard, S.; Schachter, L. A.; Arnett, F. C.;

Ginzler,
[52431] E. M.; Gourley, M. F.; Ramsey-Goldman, R.; Peterson, M. G. E.; Kimberly,
[52432] R. P.: Fc-gamma-RIIA alleles are heritable risk factors for lupus
[52433] nephritis in African Americans. J. Clin. Invest. 97: 1348-1354,
[52434] 1996.

[52435] 8387.

[52436] Sammartino, L.; Webber, L. M.; Hogarth, P. M.; McKenzie, I. F.
[52437] C.; Garson, O. M.: Assignment of the gene coding for human FcRII
[52438] (CD32) to bands q23q24 on chromosome 1. Immunogenetics 28: 380-381,
[52439] 1988.

[52440] 8388.

[52441] Stein, M.-P.; Edberg, J. C.; Kimberly, R. P.; Mangan, E. K.; Bharadwaj,
[52442] D.; Mold, C.; Du Clos, T. W.: C-reactive protein binding to Fc-gamma-RIIa
[52443] on human monocytes and neutrophils is allele-specific. J.

Clin. Invest. 105:
[52444] 369-376, 2000.

[52445] 8389.

[52446] Unkeless, J. C.: Function and heterogeneity of human Fc receptors
[52447] for immunoglobulin G. J. Clin. Invest. 83: 355-361, 1989.

[52448] 8390.

[52449] Ogawa, O.; Becroft, D. M.; Morison, I. M.; Eccles, M. R.; Skeen,
[52450] J. E.; Mauger, D. C.; Reeve, A. E.: Constitutional relaxation of
[52451] insulin-like growth factor II gene imprinting associated with Wilms'
[52452] tumour and gigantism. Nature Genet. 5: 408-412, 1993.

[52453] 8391.

[52454] Pedone, P. V.; Tirabosco, R.; Cavazzana, A. O.; Ungaro, P.; Basso,
[52455] G.; Luksch, R.; Carli, M.; Bruni, C. B.; Frunzio, R.; Riccio, A.:
[52456] Mono- and bi-allelic expression of insulin-like growth factor II gene

[52457] in human muscle tumors. Hum. Molec. Genet. 3:
1117-1121, 1994.
[52458] 8392.
[52459] Reeve, A. E.; Eccles, M. R.; Wilkins, R. J.; Bell, G. I.; Millow,
[52460] L. J.: Expression of insulin-like growth factor-II transcripts
in
[52461] Wilms' tumour. Nature 317: 258-260, 1985.
[52462] 8393.
[52463] Scott, J.; Cowell, J.; Robertson, M. E.; Priestley, L. M.;
Wadey,
[52464] R.; Hopkins, B.; Pritchard, J.; Bell, G. I.; Rall, L. B.; Graham,
[52465] C. F.; Knott, T. J.: Insulin-like growth factor-II gene ex-
pression
[52466] in Wilms' tumour and embryonic tissues. Nature 317:
260-262, 1985.
[52467] 8394.
[52468] Shen, S.-J.; Daimon, M.; Wang, C.-Y.; Jansen, M.; Ilan, J.:
Isolation
[52469] of an insulin-like growth factor II cDNA with a unique
5-prime untranslated
[52470] region from human placenta. Proc. Nat. Acad. Sci. 85:

1947-1951,

[52471] 1988.

[52472] 8395.

[52473] Sun, F.-L.; Dean, W. L.; Kelsey, G.; Allen, N. D.; Reik, W.:

[52474] Transactivation of IGF2 in a mouse model of Beckwith-Wiedemann syndrome. Nature 389:

[52475] 809-815, 1997.

[52476] 8396.

[52477] Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[52478] B.: Insulin-like growth factor genes: chromosome organi-zation and

[52479] association with disease. (Abstract) Am. J. Hum. Genet. 36: 121S,

[52480] 1984.

[52481] 8397.

[52482] Weksberg, R.; Shen, D. R.; Fei, Y.-L.; Song, Q. L.; Squire, J.

[52483] : Disruption of insulin-like growth factor 2 imprinting in Beckwith-Wiedemann

[52484] syndrome. Nature Genet. 5: 143-150, 1993.

[52485] 8398.

[52486] Zaina, S.; Pettersson, L.; Ahren, B.; Branen, L.; Bassim Has-
san,
[52487] A.; Lindholm, M.; Mattson, R.; Thyberg, J.; Nilsson, J.: In-
sulin-like
[52488] growth factor II plays a central role in atherosclerosis in a
mouse
[52489] model. J. Biol. Chem. 277: 4505-4511, 2002.

[52490] 8399.

[52491] Dixon, P. H.; Weerasekera, N.; Linton, K. J.; Donaldson, O.;
Chambers,
[52492] J.; Egginton, E.; Weaver, J.; Nelson-Piercy, C.; de Swiet, M.;
Warnes,
[52493] G.; Elias, E.; Higgins, C. F.; Johnston, D. G.; McCarthy, M.
I.; Williamson,
[52494] C.: Heterozygous MDR3 missense mutation associated
with intrahepatic
[52495] cholestasis of pregnancy: evidence for a defect in protein
trafficking. Hum.
[52496] Molec. Genet. 9: 1209-1217, 2000.

[52497] 8400.

[52498] Jacquemin, E.; Cresteil, D.; Manouvrier, S.; Boute, O.; Had-
chouel,
[52499] M.: Heterozygous non-sense mutation of the MDR3 gene
in familial
[52500] intrahepatic cholestasis of pregnancy. Lancet 353:
210-211, 1999.
[52501] 8401.

[52502] Chang-Yeh, A.; Jabs, E. W.; Li, X.; Dracopoli, N. C.; Huang,
R.
[52503] C. C.: The IPP gene is assigned to human chromosome
1p32-1p22. Genomics 15:
[52504] 239-241, 1993.

[52505] 8402.

[52506] Chang-Yeh, A.; Mold, D. E.; Brilliant, M. H.; Huang, R. C.
C.:
[52507] The mouse intracisternal A particle-promoted placental
gene retrotransposition
[52508] is mouse-strain-specific. Proc. Nat. Acad. Sci. 90:
292-296, 1993.
[52509] 8403.

[52510] Chang-Yeh, A.; Mold, D. E.; Huang, R. C. C.: Identification
of

[52511] a novel murine IAP-promoted placenta-expressed gene. Nucleic Acids
[52512] Res. 19: 3667-3672, 1991.

[52513] 8404.

[52514] Mold, D. E.; Chang-Yeh, A.; Huang, R. C. C.: Cell lineage-specific
[52515] expression of the MIPP gene. Biochem. Biophys. Res. Commun. 177:
[52516] 1062-1067, 1991.

[52517] 8405.

[52518] Cama, A.; de la Luz Sierra, M.; Quon, M. J.; Ottini, L.; Gor-den,
[52519] P.; Taylor, S. I.: Substitution of glutamic acid for alanine 1135
[52520] in the putative 'catalytic loop' of the tyrosine kinase do-main of
[52521] the human insulin receptor: a mutation that impairs pro-teolytic processing
[52522] into subunits and inhibits receptor tyrosine kinase activity. J.
[52523] Biol. Chem. 268: 8060-8069, 1993.

[52524] 8406.

[52525] Cama, A.; Quon, M. J.; de la Luz Sierra, M.; Taylor, S. I.: Substitution

[52526] of isoleucine for methionine at position 1153 in the beta-subunit

[52527] of the human insulin receptor. J. Biol. Chem. 267: 8383-8389, 1992.

[52528] 8407.

[52529] Caro, J. F.; Raju, S. M.; Sinha, M. K.; Goldfine, I. D.; Dohm,

[52530] G. L.: Heterogeneity of human liver, muscle, and adipose tissue insulin

[52531] receptor. Biochem. Biophys. Res. Commun. 151: 123-129, 1988.

[52532] 8408.

[52533] Christiansen, K.; Tranum-Jensen, J.; Carlsen, J.; Vinten, J.:

[52534] A model for the quaternary structure of human placental insulin receptor

[52535] deduced from electron microscopy. Proc. Nat. Acad. Sci. 88: 249-252,

[52536] 1991.

[52537] 8409.

[52538] Cocozza, S.; Porcellini, A.; Riccardi, G.; Monticelli, A.; Condorelli,
[52539] G.; Ferrara, A.; Pianese, L.; Miele, C.; Capaldo, B.; Beguinot, F.;
[52540] Varrone, S.: NIDDM associated with mutation in tyrosine kinase domain
[52541] of insulin receptor gene. Diabetes 41: 521-526, 1992.

[52542] 8410.

[52543] Due, C.; Simonsen, M.; Olsson, L.: The major histocompatibility
[52544] complex class I heavy chain as a structural subunit of the human cell
[52545] membrane insulin receptor: implications for the range of biological
[52546] functions of histocompatibility antigens. Proc. Nat. Acad. Sci. 83:
[52547] 6007-6011, 1986.

[52548] 8411.

[52549] Ebina, Y.; Ellis, L.; Jarnagin, K.; Edery, M.; Graf, L.; Clauser,
[52550] E.; Ou, J.-H.; Masiarz, F.; Kan, Y. W.; Goldfine, I. D.; Roth, R.

[52551] A.; Rutter, W. J.: The human insulin receptor cDNA: the structural
[52552] basis for hormone activated transmembrane signalling. Cell 40: 747-758,
[52553] 1985.

[52554] 8412.

[52555] Elbein, S. C.; Corsetti, L.; Ullrich, A.; Permutt, M. A.: Multi-ple
[52556] restriction fragment length polymorphisms at the insulin receptor
[52557] locus: a highly informative marker for linkage analysis. Proc. Nat.
[52558] Acad. Sci. 83: 5223-5227, 1986.

[52559] 8413.

[52560] Elbein, S. C.; Sorensen, L. K.; Schumacher, M. C.: Methion-ine
[52561] for valine substitution in exon 17 of the insulin receptor gene in
[52562] a pedigree with familial NIDDM. Diabetes 42: 429-434, 1993.
[52563] 8414.

[52564] Ferrannini, E.; Muggeo, M.; Navalesi, R.; Pilo, A.: Impaired

[52565] insulin degradation in a patient with insulin resistance and acanthosis

[52566] nigricans. Am. J. Med. 73: 148-154, 1982.

[52567] 8415.

[52568] Grigorescu, F.; Flier, J. S.; Kahn, C. R.: Characterization of

[52569] binding and phosphorylation defects of erythrocyte insulin receptors

[52570] in the type A syndrome of insulin resistance. Diabetes 35: 127-138,

[52571] 1986.

[52572] 8416.

[52573] Grigorescu, F.; Flier, J. S.; Kahn, C. R.: Defect in insulin

[52574] receptor phosphorylation in erythrocytes and fibroblasts associated

[52575] with severe insulin resistance. J. Biol. Chem. 259: 15003-15006,

[52576] 1984.

[52577] 8417.

[52578] Grunberger, G.; Zick, Y.; Gordon, G.: Defect in phosphory-

lation
[52579] of insulin receptors in cells from an insulin-resistant patient with
[52580] normal insulin binding. Science 223: 932-934, 1984.

[52581] 8418.

[52582] Hone, J.; Accili, D.; Al-Gazali, L. I.; Lestringant, G.; Orban,

[52583] T.; Taylor, S. I.: Homozygosity for a new mutation (ile119-to-met)
[52584] in the insulin receptor gene in five sibs with familial insulin resistance. J.
[52585] Med. Genet. 31: 715-716, 1994.

[52586] 8419.

[52587] Kadowaki, T.; Bevins, C. L.; Cama, A.; Ojamaa, K.; Marcus-Samuels,
[52588] B.; Kadowaki, H.; Beitz, L.; McKeon, C.; Taylor, S. I.: Two mutant
[52589] alleles of the insulin receptor gene in a patient with extreme insulin
[52590] resistance. Science 240: 787-790, 1988.

[52591] 8420.

[52592] Kadowaki, T.; Kadowaki, H.; Accili, D.; Taylor, S. I.: Substi-

tution
[52593] of lysine for asparagine at position 15 in the alpha-sub-
unit of the
[52594] human insulin receptor: a mutation that impairs transport
of receptors
[52595] to the cell surface and decreases the affinity of insulin
binding. J.
[52596] Biol. Chem. 265: 19143-19150, 1990.

[52597] 8421.

[52598] Kadowaki, T.; Kadowaki, H.; Accili, D.; Yazaki, Y.; Taylor,
S.
[52599] I.: Substitution of arginine for histidine at position 209 in
the
[52600] alpha-subunit of the human insulin receptor: a mutation
that impairs
[52601] receptor dimerization and transport of receptors to the
cell surface. J.
[52602] Biol. Chem. 266: 21224-21231, 1991.

[52603] 8422.

[52604] Kadowaki, T.; Kadowaki, H.; Rechler, M. M.; Serrano-Rios,
M.;
[52605] Roth, J.; Gorden, P.; Taylor, S. I.: Five mutant alleles of the

insulin
[52606] receptor gene in patients with genetic forms of insulin resistance. J.
[52607] Clin. Invest. 86: 254-264, 1990.

[52608] 8423.

[52609] Kadowaki, T.; Kadowaki, H.; Taylor, S. I.: A nonsense mutation
[52610] causing decreased levels of insulin receptor mRNA: detection by a
[52611] simplified technique for direct sequencing of genomic DNA amplified
[52612] by the polymerase chain reaction. Proc. Nat. Acad. Sci. 87: 658-662,
[52613] 1990.

[52614] 8424.

[52615] Kahn, C. R.; Flier, J. S.; Bar, R. S.; Archer, J. A.; Gorden,
[52616] P.; Martin, M. M.; Roth, J.: The syndromes of insulin resistance
[52617] and acanthosis nigricans: insulin receptor disorders in man. New
[52618] Eng. J. Med. 294: 739-745, 1976.

[52619] 8425.

[52620] Kahn, C. R.; Goldstein, B. J.: Molecular defects in insulin action. Science 245:

[52621] 13 only, 1989.

[52622] 8426.

[52623] Kahn, C. R.; White, M. F.: The insulin receptor and the molecular

[52624] mechanism of insulin action. J. Clin. Invest. 82: 1151-1156, 1988.

[52625] 8427.

[52626] Kakehi, T.; Hisatomi, A.; Kuzuya, H.; Yoshimasa, Y.; Okamoto,

[52627] M.; Yamada, K.; Nishimura, H.; Kosaki, A.; Nawata, H.; Umeda, F.;

[52628] Ibayashi, H.; Imura, H.: Defective processing of insulin-receptor

[52629] precursor in cultured lymphocytes from a patient with extreme insulin

[52630] resistance. J. Clin. Invest. 81: 2020-2022, 1988.

[52631] 8428.

[52632] Kittur, D.; Shimizu, Y.; DeMars, R.; Edidin, M.: Insulin

binding
[52633] to human B lymphoblasts is a function of HLA haplotype. Proc. Nat.
[52634] Acad. Sci. 84: 1351-1355, 1987.

[52635] 8429.

[52636] Klinkhamer, M. P.; Groen, N. A.; van der Zon, G. C. M.; Lindhout,
[52637] D.; Sandkuyl, L. A.; Krans, H. M. J.; Moller, W.; Maassen, J. A.:
[52638] A leucine-to-proline mutation in the insulin receptor in a family
[52639] with insulin resistance. EMBO J. 8: 2503-2507, 1989.

[52640] 8430.

[52641] Krook, A.; Brueton, L.; O'Rahilly, S.: Homozygous non-sense mutation
[52642] in the insulin receptor gene in infant with leprechaunism. Lancet 342:
[52643] 277-278, 1993.

[52644] 8431.

[52645] Kulkarni, R. N.; Bruning, J. C.; Winnay, J. N.; Postic, C.; Magnuson,

[52646] M. A.; Kahn, C. R.: Tissue-specific knockout of the insulin receptor
[52647] in pancreatic beta cells creates an insulin secretory defect similar
[52648] to that in type 2 diabetes. Cell 96: 329-339, 1999.

[52649] 8432.

[52650] Chan, A. M.-L.; Hilkens, J.; Kroezen, V.; Mitchell, P. J.; Scambler,
[52651] P.; Wainwright, B. J.; Williamson, R.; Cooper, C. S.: Molecular cloning
[52652] and localization to chromosome 6 of mouse INT1L1 gene. Somat. Cell
[52653] Molec. Genet. 15: 555-562, 1989.

[52654] 8433.

[52655] Huguet, E. L.; McMahon, J. A.; McMahon, A. P.; Bicknell, R.; Harris,
[52656] A. L.: Differential expression of human Wnt genes 2, 3, 4, and 7B
[52657] in human breast cell lines and normal and disease states of human
[52658] breast tissue. Cancer Res. 54: 2615-2621, 1994.

[52659] 8434.

[52660] McCoy, P. A.; Shao, Y.; Wolpert, C. M.; Donnelly, S. L.;
Ashley-Koch,
[52661] A.; Abel, H. L.; Ravan, S. A.; Abramson, R. K.; Wright, H.
H.; DeLong,
[52662] G. R.; Cuccaro, M. L.; Gilbert, J. R.; Pericak-Vance, M. A.:
No association
[52663] between the WNT2 gene and autistic disorder. Am. J. Med.
Genet. (Neuropsychiat.
[52664] Genet.) 114: 106-109, 2002.

[52665] 8435.

[52666] Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford,
G.;
[52667] McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature
for int-1 and
[52668] related genes: the Wnt gene family. Cell 64: 231-232,
1991.
[52669] 8436.

[52670] Wainwright, B. J.; Scambler, P. J.; Stanier, P.; Watson, E. K.;

[52671] Bell, G.; Wicking, C.; Estivill, X.; Courtney, M.; Bour, A.;
Pedersen,

[52672] P. S.; Williamson, R.; Farrall, M.: Isolation of a human gene with
[52673] protein sequence similarity to human and murine int-1 and the Drosophila
[52674] segment polarity mutant wingless. EMBO J. 7: 1743-1748, 1988.
[52675] 8437.

[52676] Wassink, T. H.; Piven, J.; Vieland, V. J.; Huang, J.; Swider-ski,
[52677] R. E.; Pietila, J.; Braun, T.; Beck, G.; Folstein, S. E.; Haines,
[52678] J. L.; Sheffield, V. C.: Evidence supporting WNT2 as an autism susceptibility
[52679] gene. Am. J. Med. Genet. 105: 406-413, 2001.

[52680] 8438.

[52681] Kluck, P. M. C.; Wiegant, J.; Jansen, R. P. M.; Bolk, M. W. J.;
[52682] Raap, A. K.; Willemze, R.; Landegent, J. E.: The human in-terleukin-6
[52683] receptor alpha-chain gene is localized on chromosome 1 band q21. Hum.
[52684] Genet. 90: 542-544, 1993.

[52685] 8439.

[52686] Szpirer, J.; Szpirer, C.; Riviere, M.; Houart, C.; Baumann, M.;

[52687] Fey, G. H.; Poli, V.; Cortese, R.; Islam, M. Q.; Levan, G.: The interleukin-6-dependent

[52688] DNA-binding protein gene (transcription factor 5: TCF5) maps to human

[52689] chromosome 20 and rat chromosome 3, the IL6 receptor locus (IL6R)

[52690] to human chromosome 1 and rat chromosome 2, and the rat IL6 gene to

[52691] rat chromosome 4. Genomics 10: 539-546, 1991.

[52692] 8440.

[52693] Yamasaki, K.; Taga, T.; Hirata, Y.; Yawata, H.; Kawanishi, Y.;

[52694] Seed, B.; Taniguchi, T.; Hirano, T.; Kishimoto, T.: Cloning and expression

[52695] of the human interleukin-6 (BSF-2/IFN-beta-2) receptor. Science 241:

[52696] 825-828, 1988.

[52697] 8441.

[52698] Melen, K.; Keskinen, P.; Ronni, T.; Sareneva, T.; Lounatmaa, K.;

[52699] Julkunen, I.: Human MxB protein, an interferon-alpha-inducible GTPase,
[52700] contains a nuclear targeting signal and is localized in the hemochromatin
[52701] region beneath the nuclear envelope. J. Biol. Chem. 271: 23478-23486,
[52702] 1996.

[52703] 8442.

[52704] Berry, M. J.; Banu, L.; Larsen, P. R.: Type I iodothyronine deiodinase
[52705] is a selenocysteine-containing enzyme. Nature 349: 438-440, 1991.
[52706] 8443.

[52707] Berry, M. J.; Grieco, D.; Taylor, B. A.; Maia, A. L.; Kieffer,

[52708] J. D.; Beamer, W.; Glover, E.; Poland, A.; Larsen, P. R.: Physiological
[52709] and genetic analyses of inbred mouse strains with a type I iodothyronine
[52710] 5-prime deiodinase deficiency. J. Clin. Invest. 92: 1517-1528, 1993.
[52711] 8444.

[52712] Hatfield, D.; Diamond, A.: UGA: a split personality in the universal
[52713] genetic code. (Letter) Trends Genet. 9: 69-70, 1993.

[52714] 8445.

[52715] Jakobs, T. C.; Koehler, M. R.; Schmutzler, C.; Glaser, F.; Schmid,
[52716] M.; Kohrle, J.: Structure of the human type I iodothyronine 5-prime-deiodinase
[52717] gene and localization to chromosome 1p32-p33. Genomics 42: 361-363,
[52718] 1997.

[52719] 8446.

[52720] Jansen, M.; Krenning, E. P.; Oostdijik, W.; Docter, R.; Kingma,
[52721] B. E.; Van den Brande, J. V. L.; Hennemann, G.: Hyperthyroxinaemia
[52722] due to decreased peripheral triiodothyronine production. Lancet II:
[52723] 849-851, 1982.

[52724] 8447.

[52725] Kleinhaus, N.; Faber, J.; Kahana, L.; Schneer, J.; Scheinfeld,

[52726] M.: Euthyroid hyperthyroxinemia due to a generalized 5-prime-deiodinase
[52727] defect. J. Clin. Endocr. Metab. 66: 684-688, 1988.

[52728] 8448.

[52729] Maia, A. L.; Berry, M. J.; Sabbag, R.; Harney, J. W.; Larsen, P.
[52730] R.: Structural and functional differences in the dio1 gene in mice
[52731] with inherited type 1 deiodinase deficiency. Molec. En- docr. 9: 969-980,
[52732] 1995.

[52733] 8449.

[52734] Mandel, S. J.; Berry, M. J.; Kieffer, J. D.; Harney, J. W.; Warne,
[52735] R. L.; Larsen, P. R.: Cloning and in vitro expression of the human
[52736] selenoprotein, type I iodothyronine deiodinase. J. Clin. En- docr.
[52737] Metab. 75: 1133-1139, 1992.

[52738] 8450.

[52739] Toyoda, N.; Kleinhaus, N.; Larsen, P. R.: The structure of

the

[52740] coding and 5-prime flanking region of the type 1 iodothyronine deiodinase

[52741] (dio1) gene is normal in a patient with suspected congenital dio1

[52742] deficiency. J. Clin. Endocr. Metab. 81: 2121-2124, 1996.

[52743] 8451.

[52744] Alli, C.; Consalez, G. G.: Linkage mapping of Csrp to proximal

[52745] mouse chromosome 3. Mammalian Genome 9: 172 only, 1998.

[52746] 8452.

[52747] Erdel, M.; Weiskirchen, R.: Assignment of CSRP1 encoding the LIM

[52748] domain protein CRP1, to human chromosome 1q32 by fluorescence in situ

[52749] hybridization. Cytogenet. Cell Genet. 83: 10-11, 1998.

[52750] 8453.

[52751] Liebhaber, S. A.; Emery, J. G.; Urbanek, M.; Wang, X.; Cooke, N.

[52752] E.: Characterization of a human cDNA encoding a widely

expressed
[52753] and highly conserved cysteine-rich protein with an un-
usual zinc-finger
[52754] motif. Nucleic Acids Res. 18: 3871-3879, 1990.

[52755] 8454.

[52756] Weiskirchen, R.; Pino, J. D.; Macalma, T.; Bister, K.; Beck-
erle,
[52757] M. C.: The cysteine-rich protein family of highly related
LIM domain
[52758] proteins. J. Biol. Chem. 270: 28946-28954, 1995.

[52759] 8455.

[52760] Asselta, R.; Duga, S.; Simonic, T.; Malcovati, M.; San-
tagostino,
[52761] E.; Giangrande, P. L. F.; Mannucci, P. M.; Tenchini, M. L.:
Afibrinogenemia:
[52762] first identification of a splicing mutation in the fibrinogen
gamma
[52763] chain gene leading to a major gamma chain truncation.
Blood 96:
[52764] 2496-2500, 2000.

[52765] 8456.

[52766] Bantia, S.; Bell, W. R.; Dang, C. V.: Polymerization defect of

[52767] fibrinogen Baltimore III due to a gamma-asn308-to-ile mutation. Blood 75:

[52768] 1659-1663, 1990.

[52769] 8457.

[52770] Bantia, S.; Mane, S. M.; Bell, W. R.; Dang, C. V.: Fibrinogen

[52771] Baltimore I: polymerization defect associated with a gamma(292)gly-to-val

[52772] (GGC-GTC) mutation. Blood 76: 2279-2283, 1990.

[52773] 8458.

[52774] Brook, J. G.; Tabori, S.; Tatarsky, I.; Hashmonai, M.; Schramek,

[52775] A.: Fibrinogen 'Haifa'--a new fibrinogen variant: a case report. Haemostasis 13:

[52776] 277-281, 1983.

[52777] 8459.

[52778] Brown, C. H., III; Crowe, M. F.: Defective alpha polymerization

[52779] in the conversion of fibrinogen Baltimore to fibrin. J. Clin. Invest. 55:

[52780] 1190-1194, 1975.

[52781] 8460.

[52782] Budzynski, A. Z.; Marder, V. J.; Menache, D.; Guillin, M.-C.:

[52783] Defect in the gamma polypeptide chain of a congenital abnormal fibrinogen

[52784] (Paris I). Nature 252: 66-68, 1974.

[52785] 8461.

[52786] Cote, H. C. F.; Lord, S. T.; Pratt, K. P.: Gamma-chain dysfibrinogenemias:

[52787] molecular structure-function relationships of naturally occurring

[52788] mutations in the gamma chain of human fibrinogen. J. Am. Soc. Hemat. 92:

[52789] 2195-2212, 1998.

[52790] 8462.

[52791] Crabtree, G. R.; Kant, J. A.: Organization of the rat gamma-fibrinogen

[52792] gene: alternative mRNA splice patterns produce the gamma-A and gamma-B

[52793] (gamma-prime) chains of fibrinogen. Cell 31: 159-166, 1982.

[52794] 8463.

[52795] Ebert, R. F.; Bell, W. R.: Fibrinogen Baltimore IV: congenital

[52796] dysfibrinogenemia with delayed fibrin monomer polymer-
ization. Thromb.

[52797] Res. 38: 121-128, 1985.

[52798] 8464.

[52799] Ebert, R. F.; Bell, W. R.: Fibrinogen Baltimore III: congenital

[52800] dysfibrinogenemia with a shortened gamma-subunit.
Thromb. Res. 51:

[52801] 251-258, 1988.

[52802] 8465.

[52803] Fernandez, F. J.; Noguerol, P.; Sosa, R.; Cuesta, B.; Paramo,

[52804] J. A.; Rocha, E.: Fibrinogen Sevilla, a congenital dysfibrino-
genemia

[52805] characterized by an abnormal monomer aggregation and
a defective plasmin

[52806] lysis. Clin. Chim. Acta 179: 239-250, 1989.

[52807] 8466.

[52808] Fornace, A. J., Jr.; Cummings, D. E.; Comeau, C. M.; Kant,

J.

[52809] A.; Crabtree, G. R.: Structure of the human gamma-fibrinogen gene:

[52810] alternate mRNA splicing near the 3-prime end of the gene produces

[52811] gamma-A and gamma-B forms of gamma-fibrinogen. J. Biol. Chem. 259:

[52812] 12826-12830, 1984.

[52813] 8467.

[52814] Haverkate, F.; Samama, M.: Familial dysfibrinogenemia and thrombophilia:

[52815] Report on a study of the SSC subcommittee on fibrinogen. Thromb.

[52816] Haemost. 73: 151-161, 1995.

[52817] 8468.

[52818] Bachman, N. J.; Wu, W.; Schmidt, T. R.; Grossman, L. I.; Lomax,

[52819] M. I.: The 5-prime region of the COX4 gene contains a novel overlapping

[52820] gene, NOC4. Mammalian Genome 10: 506-512, 1999.

[52821] 8469.

[52822] Brown, W. M.; Prager, E. M.; Wang, A.; Wilson, A. C.: Mitochondrial

[52823] DNA sequences of primates: tempo and mode of evolution. J. Molec.

[52824] Evol. 18: 225-239, 1982.

[52825] 8470.

[52826] Darras, B. T.; Zeviani, M.; Schon, E. A.; Francke, U.: Sequences

[52827] homologous to cytochrome c oxidase subunit IV are located on human

[52828] chromosomes 14q21-qter and 16q22-q24. (Abstract) Cytogenet. Cell

[52829] Genet. 46: 603 only, 1987.

[52830] 8471.

[52831] Lomax, M. I.; Hewett-Emmett, D.; Yang, T. L.; Grossman, L. I.:

[52832] Rapid evolution of the human gene for cytochrome c oxidase subunit

[52833] IV. Proc. Nat. Acad. Sci. 89: 5266-5270, 1992.

[52834] 8472.

[52835] Lomax, M. I.; Welch, M. D.; Darras, B. T.; Francke, U.;

Grossman,

[52836] L. I.: Novel use of a chimpanzee pseudogene for chromosomal mapping

[52837] of human cytochrome c oxidase subunit IV. Gene 86: 209-216, 1990.

[52838] 8473.

[52839] Makris, G. J.; Samuelson, L. C.; Lomax, M. I.: The gene encoding

[52840] subunit IV of cytochrome c oxidase maps to mouse chromosome 8. Mammalian

[52841] Genome 7: 789-790, 1996.

[52842] 8474.

[52843] Zeviani, M.; Nakagawa, M.; Herbert, J.; Lomax, M. I.; Grossman,

[52844] L. I.; Sherbany, A. A.; Miranda, A. F.; DiMauro, S.; Schon, E. A.

[52845] : Isolation of a cDNA clone encoding subunit IV of human cytochrome

[52846] c oxidase. Gene 55: 205-217, 1987.

[52847] 8475.

[52848] Wang, X.; Lee, G.; Liebhaber, S. A.; Cooke, N. E.: Human

cysteine-rich
[52849] protein: a member of the LIM/double-finger family displaying coordinate
[52850] serum induction with c-myc. J. Biol. Chem. 267: 9176-9184, 1992.
[52851] 8476.

[52852] Wang, X.; Ray, K.; Szpirer, J.; Levan, G.; Liebhaber, S. A.; Cooke,
[52853] N. E.: Analysis of the human cysteine-rich protein gene (CSRP), assignment
[52854] to chromosome 1q24-1q32, and identification of an associated MspI
[52855] polymorphism. Genomics 14: 391-397, 1992.

[52856] 8477.

[52857] Courseaux, A.; Grosgeorge, J.; Gaudray, P.; Pannett, A. A. J.;
[52858] Forbes, S. A.; Williamson, C.; Bassett, D.; Thakker, R. V.; Teh, B.
[52859] T.; Farnebo, F.; Shepherd, J.; Skogseid, B.; Larsson, C.; Giraud,
[52860] S.; Zhang, C. X.; Salandre, J.; Calender, A.: Definition of the minimal

[52861] MEN1 candidate area based on a 5-Mb integrated map of proximal 11q13. Genomics 37:
[52862] 354-365, 1996.

[52863] 8478.

[52864] Dorin, J. R.; Emslie, E.; van Heyningen, V.: Related cal-
cium-binding
[52865] proteins map to the same subregion of chromosome 1q and to an extended
[52866] region of synteny on mouse chromosome 3. Genomics 8: 420-426, 1990.
[52867] 8479.

[52868] Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Modi, W. S.;
[52869] Nienhuis, A. W.: Chromosomal localization and racial dis-
tribution
[52870] of the polymorphic human dihydrofolate reductase pseu-
dogene (DHFRPI). Am.
[52871] J. Hum. Genet. 42: 345-352, 1988.

[52872] 8480.

[52873] Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Nienhuis, A.

[52874] W.: Chromosomal localization and racial distribution of the polymorphic
[52875] hDHFR-psi-1 pseudogene. (Abstract) Clin. Res. 33: 328A only, 1985.
[52876] 8481.

[52877] Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Nienhuis, A.
[52878] W.: A novel form of human polymorphism involving the hDHFR-psi-1
[52879] pseudogene identifies three RFLPs. Nucleic Acids Res. 15: 5501 only,
[52880] 1987.

[52881] 8482.

[52882] Anagnou, N. P.; O'Brien, S. J.; Shimada, T.; Nash, W. G.; Chen,
[52883] M.-J.; Nienhuis, A. W.: Chromosomal organization of the human dihydrofolate
[52884] reductase genes: dispersion, selective amplification and a novel form
[52885] of polymorphism. Proc. Nat. Acad. Sci. 81: 5170-5174, 1984.
[52886] 8483.

[52887] Blakley, R. L.; Sorrentino, B. P.: In vitro mutations in dihydrofolate
[52888] reductase that confer resistance to methotrexate: potential for clinical
[52889] application. Hum. Mutat. 11: 259-263, 1998.

[52890] 8484.

[52891] Chen, M.-J.; Shimada, T.; Moulton, A. D.; Harrison, M.; Nienhuis,
[52892] A. W.: Intronless human dihydrofolate reductase genes are derived
[52893] from processed RNA molecules. Proc. Nat. Acad. Sci. 79: 7435-7439,
[52894] 1982.

[52895] 8485.

[52896] Craik, C. S.; Rutter, W. J.; Fletterick, R.: Splice junctions:

[52897] association with variation in protein structure. Science 220: 1125-1129,
[52898] 1983.

[52899] 8486.

[52900] Erbe, R. W.: Inborn errors of folate metabolism. New Eng. J.

[52901] Med. 293: 753-757 and 807-812, 1975.

[52902] 8487.

[52903] Funanage, V. L.; Myoda, T. T.; Moses, P. A.; Cowell, H. R.: Assignment

[52904] of the human dihydrofolate reductase gene to the q11-q22 region of

[52905] chromosome 5. Molec. Cell. Biol. 4: 2010-2016, 1984.

[52906] 8488.

[52907] Hoffbrand, A. V.; Tripp, E.; Jackson, B. F. A.; Luck, W. E.; Frater-Schroder,

[52908] M.: Hereditary abnormal transcobalamin II previously di-agnosed as

[52909] congenital dihydrofolate reductase deficiency. (Letter) New Eng.

[52910] J. Med. 310: 789-790, 1984.

[52911] 8489.

[52912] Killary, A. M.; Leach, R. J.; Moran, R. G.; Fournier, R. E. K.

[52913] : Assignment of genes encoding dihydrofolate reductase and hexosaminidase

[52914] B to Mus musculus chromosome 13. Somat. Cell Molec. Genet. 12: 641-648,

[52915] 1986.

[52916] 8490.

[52917] Maurer, B.; Barker, P. E.; Masters, J. N.; D'Eustachio, P.; Ruddle,

[52918] F. H.; Attardi, G.: Chromosomal location of the normal human DHFR

[52919] gene and of its amplified copies in methotrexate resistant cell variants.

[52920] (Abstract) Cytogenet. Cell Genet. 37: 534 only, 1984.

[52921] 8491.

[52922] Maurer, B. J.; Barker, P. E.; Masters, J. N.; Ruddle, F. H.; Attardi,

[52923] G.: Human dihydrofolate reductase gene is located in chromosome 5

[52924] and is unlinked to the related pseudogenes. Proc. Nat. Acad. Sci. 81:

[52925] 1484-1488, 1984.

[52926] 8492.

[52927] Maurer, B. J.; Carlock, L.; Wasmuth, J.; Attardi, G.: Assignment

[52928] of human dihydrofolate reductase gene to band q23 of

chromosome 5

[52929] and of related pseudogene psiHD1 to chromosome 3. So-
mat. Cell Molec.

[52930] Genet. 11: 79-85, 1985.

[52931] 8493.

[52932] Myoda, T. T.; Funanage, V. L.: Personal Communication.
Wilmington,

[52933] Del. 10/7/1983.

[52934] 8494.

[52935] Singer, M. J.; Mesner, L. D.; Friedman, C. L.; Trask, B. J.;
Hamlin,

[52936] J. L.: Amplification of the human dihydrofolate reductase
gene via

[52937] double minutes is initiated by chromosome breaks. Proc.
Nat. Acad.

[52938] Sci. 97: 7921-7926, 2000.

[52939] 8495.

[52940] Tauro, G. P.; Danks, D. M.; Rowe, P. B.; Van der Weyden,
M. B.;

[52941] Schwarz, M. A.; Collins, V. L.; Neal, B. W.: Dihydrofolate
reductase

[52942] deficiency causing megaloblastic anemia in two families. New Eng.
[52943] J. Med. 294: 466-470, 1976.

[52944] 8496.

[52945] Walters, T. R.: Congenital megaloblastic anemia responsive to
[52946] N(5)-formyltetrahydrofolic acid administration. J. Pediat. 70: 686-687,
[52947] 1967.

[52948] 8497.

[52949] Ali, G.; Wasco, W.; Cai, X.; Szabo, P.; Sheu, K.-F. R.; Cooper,
[52950] A. J. L.; Gaston, S. M.; Gusella, J. F.; Tanzi, R. E.; Blass, J. P.
[52951] : Isolation, characterization, and mapping of gene encoding dihydrolipoyl
[52952] succinyltransferase (E2k) of human alpha-ketoglutarate dehydrogenase
[52953] complex. Somat. Cell Molec. Genet. 20: 99-105, 1994.

[52954] 8498.

[52955] Nakano, K.; Matuda, S.; Sakamoto, T.; Takase, C.; Nakagawa, S.;

[52956] Ohta, S.; Ariyama, T.; Inazawa, J.; Abe, T.; Miyata, T.: Human dihydrolipoamide
[52957] succinyltransferase: cDNA cloning and localization on chromosome 14q24.2-q24.3.
[52958] Biochim. Biophys. Acta 1216: 360-368, 1993.

[52959] 8499.

[52960] Nakano, K.; Takase, C.; Sakamoto, T.; Ohta, S.; Nakagawa, S.; Ariyama,
[52961] T.; Inazawa, J.; Abe, T.; Matuda, S.: An unspliced cDNA for human
[52962] dihydrolipoamide succinyltransferase: characterization and mapping
[52963] of the gene to chromosome 14q24.2-q24.3. Biochem. Biophys. Res.
[52964] Commun. 196: 527-533, 1993.

[52965] 8500.

[52966] Patel, M. S.; Harris, R. A.: Mammalian alpha-keto acid dehydrogenase
[52967] complexes: gene regulation and genetic defects. FASEB J. 9: 1164-1172,
[52968] 1995.

[52969] 8501.

[52970] Wenstrup, R. J.; Langland, G. T.; Willing, M. C.; D'Souza, V.

[52971] N.; Cole, W. G.: A splice-junction mutation in the region of COL5A1

[52972] that codes for the carboxyl propeptide of pro-alpha-1(V) chains results

[52973] in the gravis form of the Ehlers-Danlos syndrome (type I). Hum. Molec.

[52974] Genet. 5: 1733-1736, 1996.

[52975] 8502.

[52976] De Paepe, A.; Nuytinck, L.; Hausser, I.; Anton-Lamprecht, I.; Naeyaert,

[52977] J.-M.: Mutations in the COL5A1 gene are causal in the Ehlers-Danlos

[52978] syndromes I and II. Am. J. Hum. Genet. 60: 547-554, 1997.

[52979] 8503.

[52980] Palmer, G.; Manen, D.; Bonjour, J.-P.; Caverzasio, J.: Char-

[52981] acterization of the human Glvr-1 phosphate transporter/retrovirus receptor gene

[52982] and promoter region. Gene 226: 25-33, 1999.

[52983] 8504.

[52984] Fan, N. C.; Peng, C.; Krisinger, J.; Leung, P. C. K.: The hu-
man
[52985] gonadotropin-releasing hormone receptor gene: complete
structure including
[52986] multiple promoters, transcription initiation sites, and
polyadenylation
[52987] signals. Molec. Cell. Endocr. 107: R1-R8, 1995.

[52988] 8505.

[52989] Nakano, A.; Pfendner, E.; Pulkkinen, L.; Hashimoto, I.;
Uitto,
[52990] J.: Herlitz junctional epidermolysis bullosa: novel and re-
current
[52991] mutations in the LAMB3 gene and the population carrier
frequency. J.
[52992] Invest. Derm. 115: 493-498, 2000.

[52993] 8506.

[52994] Boghosian-Sell, L.; Comings, D. E.; Overhauser, J.:
Tourette syndrome
[52995] in a pedigree with a 7;18 translocation: identification of a

YAC spanning
[52996] the translocation breakpoint at 18q22.3. Am. J. Hum. Genet. 59:
[52997] 999-1005, 1996.

[52998] 8507.

[52999] Collod-Beroud, G.; Lackmy-Port-Lys, M.; Jondeau, G.; Mathieu, M.;
[53000] Maingourd, Y.; Coulon, M.; Guillotel, M.; Junien, C.; Boileau, C.
[53001] : Demonstration of the recurrence of Marfan-like skeletal and cardiovascular
[53002] manifestations due to germline mosaicism for an FBN1 mutation. (Letter) Am.
[53003] J. Hum. Genet. 65: 917-921, 1999.

[53004] 8508.

[53005] Corson, G. M.; Chalberg, S. C.; Dietz, H. C.; Charbonneau, N.

[53006] L.; Sakai, L. Y.: Fibrillin binds calcium and is coded by cD-NAs that

[53007] reveal a multidomain structure and alternatively spliced exons at

[53008] the 5-prime end. Genomics 17: 476-484, 1993.

[53009] 8509.

[53010] De Paepe, A.; Devereux, R. B.; Dietz, H. C.; Hennekam, R. C. M.;

[53011] Pyeritz, R. E.: Revised diagnostic criteria for the Marfan syndrome. Am.

[53012] J. Med. Genet. 62: 417-426, 1996.

[53013] 8510.

[53014] Dietz, H. C.; Cutting, G. R.; Pyeritz, R. E.; Maslen, C. L.; Sakai,

[53015] L. Y.; Corson, G. M.; Puffenberger, E. G.; Hamosh, A.; Nan-thakumar,

[53016] E. J.; Curristin, S. M.; Stetten, G.; Meyers, D. A.; Franco-mano, C.

[53017] A.: Marfan syndrome caused by a recurrent de novo mis-
sense mutation
[53018] in the fibrillin gene. Nature 352: 337-339, 1991.

[53019] 8511.

[53020] Dietz, H. C.; Kendzior, R. J., Jr.: Maintenance of an open
reading
[53021] frame as an additional level of scrutiny during splice site
selection. Nature
[53022] Genet. 8: 183-188, 1994.

[53023] 8512.

[53024] Dietz, H. C.; McIntosh, I.; Sakai, L. Y.; Corson, G. M.; Chal-
berg,
[53025] S. C.; Pyeritz, R. E.; Francomano, C. A.: Four novel FBN1
mutations:
[53026] significance for mutant transcript level and EGF-like do-
main calcium
[53027] binding in the pathogenesis of Marfan syndrome. Ge-
nomics 17: 468-475,
[53028] 1993.

[53029] 8513.

[53030] Dietz, H. C.; Pyeritz, R. E.: Molecular biology: to the heart

[53031] of the matter. (Editorial) New Eng. J. Med. 330: 930-932, 1994.

[53032] 8514.

[53033] Dietz, H. C.; Pyeritz, R. E.: Mutations in the human gene for

[53034] fibrillin-1 (FBN1) in the Marfan syndrome and related disorders. Hum.

[53035] Molec. Genet. 4: 1799-1809, 1995.

[53036] 8515.

[53037] Dietz, H. C.; Pyeritz, R. E.; Puffenberger, E. G.; Kendzior, R.

[53038] J., Jr.; Corson, G. M.; Maslen, C. L.; Sakai, L. Y.; Francomano, C.

[53039] A.; Cutting, G. R.: Marfan phenotype variability in a family segregating

[53040] a missense mutation in the epidermal growth factor-like motif of the

[53041] fibrillin gene. J. Clin. Invest. 89: 1674-1680, 1992.

[53042] 8516.

[53043] Dietz, H. C.; Saraiva, J. M.; Pyeritz, R. E.; Cutting, G. R.;

[53044] Francomano, C. A.: Clustering of fibrillin (FBN1) missense mutations

[53045] in Marfan syndrome patients at cysteine residues in EGF-like domains. Hum.

[53046] Mutat. 1: 366-374, 1992.

[53047] 8517.

[53048] Dietz, H. C.; Sood, I.; McIntosh, I.: The phenotypic continuum

[53049] associated with FBN1 mutations includes the Shprintzen-Goldberg syndrome.

[53050] (Abstract) Am. J. Hum. Genet. 57: A211, 1995.

[53051] 8518.

[53052] Dietz, H. C.; Valle, D.; Francomano, C. A.; Kendzior, R. J., Jr.;

[53053] Pyeritz, R. E.; Cutting, G. R.: The skipping of constitutive exons

[53054] in vivo induced by nonsense mutations. Science 259: 680-683, 1993.

[53055] 8519.

[53056] Downing, A. K.; Knott, V.; Werner, J. M.; Cardy, C. M.; Campbell,

[53057] I. D.; Handford, P. A.: Solution structure of a pair of calcium-binding

[53058] epidermal growth factor-like domains: implications for the Marfan
[53059] syndrome and other genetic disorders. Cell 85: 597-605, 1996.
[53060] 8520.

[53061] Eldadah, Z. A.; Brenn, T.; Furthmayr, H.; Dietz, H. C.: Ex-pression
[53062] of a mutant human fibrillin allele upon a normal human or murine genetic
[53063] background recapitulates a Marfan cellular phenotype. J. Clin. Invest. 95:
[53064] 874-880, 1995.

[53065] 8521.

[53066] Eldadah, Z. A.; Grifo, J. A.; Dietz, H. C.: Marfan syndrome as
[53067] a paradigm for transcript-targeted preimplantation diag-nosis of heterozygous
[53068] mutations. Nature Med. 1: 798-803, 1995.

[53069] 8522.

[53070] Francke, U.; Berg, M. A.; Tynan, K.; Brenn, T.; Liu, W.; Aoyama,

[53071] T.; Gasner, C.; Miller, D. C.; Furthmayr, H.: A gly1127ser mutation
[53072] in an EGF-like domain of the fibrillin-1 gene is a risk factor for
[53073] ascending aortic aneurysm and dissection. Am. J. Hum. Genet. 56:
[53074] 1287-1296, 1995.

[53075] 8523.

[53076] Gayraud, B.; Keene, D. R.; Sakai, L. Y.; Ramirez, F.: New insights
[53077] into the assembly of extracellular microfibrils from the analysis
[53078] of the fibrillin 1 mutation in the tight skin mouse. J. Cell Biol. 150:
[53079] 667-679, 2000.

[53080] 8524.

[53081] Glesby, M. J.; Pyeritz, R. E.: Association of mitral valve prolapse
[53082] and systemic abnormalities of connective tissue: a phenotypic continuum. J.A.M.A. 262:
[53083] 523-528, 1989.

[53084] 8525.

[53085] Godfrey, M.; Vandemark, N.; Wang, M.; Velinov, M.; War-
gowski,
[53086] D.; Tsipouras, P.; Han, J.; Becker, J.; Robertson, W.; Droste,
S.;
[53087] Rao, V. H.: Prenatal diagnosis and a donor splice site mu-
tation in
[53088] fibrillin in a family with Marfan syndrome. Am. J. Hum.
Genet. 53:
[53089] 472-480, 1993.

[53090] 8526.

[53091] Hayward, C.; Brock, D. J. H.: Fibrillin-1 mutations in Mar-
fan
[53092] syndrome and other type-1 fibrillinopathies. Hum. Mutat.
10: 415-423,
[53093] 1997.

[53094] 8527.

[53095] Hayward, C.; Keston, M.; Brock, D. J. H.; Dietz, H. C.: Fib-
rillin
[53096] (FBN1) mutations in Marfan syndrome. (Letter) Hum. Mu-
tat. 1: 79,

[53097] 1992.

[53098] 8528.

[53099] Hayward, C.; Porteous, M. E.; Brock, D. J. H.: Mutation screening

[53100] of all 65 exons of the fibrillin-1 gene in 60 patients with Marfan

[53101] syndrome: report of 12 novel mutations. Hum. Mutat. 10: 280-289,

[53102] 1997.

[53103] 8529.

[53104] Hayward, C.; Porteous, M. E. M.; Brock, D. J. H.: Identification

[53105] of a novel nonsense mutation in the fibrillin gene (FBN1) using nonisotopic

[53106] techniques. Hum. Mutat. 3: 159-162, 1994.

[53107] 8530.

[53108] Hewett, D. R.; Lynch, J. R.; Smith, R.; Sykes, B. C.: A novel

[53109] fibrillin mutation in the Marfan syndrome which could disrupt calcium

[53110] binding of the epidermal growth factor-like module. Hum. Molec. Genet. 2:

[53111] 475-477, 1993.

[53112] 8531.

[53113] Hollister, D. W.; Godfrey, M.; Sakai, L. Y.; Pyeritz, R. E.:

[53114] Immunohistologic abnormalities of the microfibrillar-fiber system

[53115] in the Marfan syndrome. New Eng. J. Med. 323: 152-159, 1990.

[53116] 8532.

[53117] Hutchinson, S.; Wordsworth, B. P.; Handford, P. A.: Marfan syndrome

[53118] caused by a mutation in FBN1 that gives rise to cryptic splicing and

[53119] a 33 nucleotide insertion in the coding sequence. Hum. Genet. 109:

[53120] 416-420, 2001.

[53121] 8533.

[53122] Judge, D. P.; Biery, N. J.; Dietz, H. C.: Characterization and

[53123] microsatellite markers flanking FBN1: utility in the diagnostic evaluation

[53124] for Marfan syndrome. Am. J. Med. Genet. 99: 39-47, 2001.

[53125] 8534.

[53126] Kainulainen, K.; Sakai, L. Y.; Child, A.; Pope, F. M.;
Puhakka,
[53127] L.; Ryhanen, L.; Palotie, A.; Kaitila, I.; Peltonen, L.: Two
mutations
[53128] in Marfan syndrome resulting in truncated fibrillin
polypeptides. Proc.
[53129] Nat. Acad. Sci. 89: 5917-5921, 1992.

[53130] 8535.

[53131] Grosse, R.; Schoneberg, T.; Schultz, G.; Gudermann, T.:
Inhibition
[53132] of gonadotropin-releasing hormone receptor signaling by
expression
[53133] of a splice variant of the human receptor. Molec. Endocr.
11: 1305-1318,
[53134] 1997.

[53135] 8536.

[53136] Kaiser, U. B.; Dushkin, H.; Altherr, M. R.; Beier, D. R.; Chin,
[53137] W. W.: Chromosomal localization of the gonadotropin-releas-
ing hormone
[53138] receptor gene to human chromosome 4q13.1-q21.1 and

mouse chromosome
[53139] 5. Genomics 20: 506-508, 1994.

[53140] 8537.

[53141] Kakar, S. S.; Musgrove, L. C.; Devor, D. C.; Sellers, J. C.; Neill,
[53142] J. D.: Cloning, sequencing, and expression of human gonadotropin
[53143] releasing hormone (GnRH) receptor. Biochem. Biophys. Res. Commun. 189:
[53144] 289-295, 1992.

[53145] 8538.

[53146] Kakar, S. S.; Neill, J. D.: The human gonadotropin-releasing
[53147] hormone receptor gene (GNRHR) maps to chromosome band 4q13. Cytogenet.
[53148] Cell Genet. 70: 211-214, 1995.

[53149] 8539.

[53150] Kottler, M.-L.; Chauvin, S.; Lahlou, N.; Harris, C. E.; Johnston,
[53151] C. J.; Lagarde, J.-P.; Bouchard, P.; Farid, N. R.; Counis, R.: A

[53152] new compound heterozygous mutation of the go-
nadotropin-releasing hormone
[53153] receptor (L314X, Q106R) in a woman with complete hy-
pogonadotropic
[53154] hypogonadism: chronic estrogen administration amplifies
the gonadotropin
[53155] defect. J. Clin. Endocr. Metab. 85: 3002-3008, 2000.

[53156] 8540.

[53157] Kottler, M.-L.; Counis, R.; Bouchard, P.: Mutations of the
GnRH
[53158] receptor gene: a new cause of autosomal-recessive hy-
pogonadotropic
[53159] hypogonadism. Arch. Med. Res. 30: 481-485, 1999.

[53160] 8541.

[53161] Kottler, M. L.; Lorenzo, F.; Bergametti, F.; Commercon, P.;
Souchier,
[53162] C.; Counis, R.: Subregional mapping of the human go-
nadotropin-releasing
[53163] hormone receptor (GnRH-R) gene to 4q between the
markers D4S392 and
[53164] D4S409. Hum. Genet. 96: 477-480, 1995.

[53165] 8542.

[53166] Layman, L. C.; Cohen, D. P.; Jin, M.; Xie, J.; Li, Z.; Reindol-
lar,

[53167] R. H.; Bolbolan, S.; Bick, D. P.; Sherins, R. R.; Duck, L. W.;
Musgrove,

[53168] L. C.; Sellers, J. C.; Neill, J. D.: Mutations in gonadotropin-re-
leasing

[53169] hormone receptor gene cause hypogonadotropic hypogo-
nadism. (Letter) Nature

[53170] Genet. 18: 14-15, 1998.

[53171] 8543.

[53172] Leung, P. C. K.; Squire, J.; Peng, C.; Fan, N.; Hayden, M. R.;

[53173] Olofsson, J. I.: Mapping of the gonadotropin-releasing
hormone (GnRH)

[53174] receptor gene to human chromosome 4q21.2 by fluores-
cence in situ hybridization. Mammalian

[53175] Genome 6: 309-310, 1995.

[53176] 8544.

[53177] Mason, A. J.; Hayflick, J. S.; Zoeller, R. T.; Young, W. S., III;

[53178] Phillips, H. S.; Nikolics, K.; Seeburg, P. H.: A deletion trun-
cating

[53179] the gonadotropin-releasing hormone gene is responsible for hypogonadism

[53180] in the 'hpg' mouse. Science 234: 1366-1371, 1986.

[53181] 8545.

[53182] Morrison, N.; Sellar, R. E.; Boyd, E.; Eidne, K. A.; Connor, J.

[53183] M.: Assignment of the gene encoding the human gonadotropin-releasing

[53184] hormone receptor to 4q13.2-13.3 by fluorescence in situ hybridization. Hum.

[53185] Genet. 93: 714-715, 1994.

[53186] 8546.

[53187] Pitteloud, N.; Boepple, P. A.; DeCruz, S.; Valkenburgh, S. B.;

[53188] Crowley, W. F., Jr.; Hayes, F. J.: The fertile eunuch variant of

[53189] idiopathic hypogonadotropic hypogonadism: spontaneous reversal associated

[53190] with a homozygous mutation in the gonadotropin-releasing hormone receptor. J.

[53191] Clin. Endocr. Metab. 86: 2470-2475, 2001.

[53192] 8547.

[53193] Pralong, F. P.; Gomez, F.; Castillo, E.; Cotecchia, S.; Abuin,

[53194] L.; Aubert, M. L.; Portmann, L.; Gaillard, R. C.: Complete hypogonadotropic

[53195] hypogonadism associated with a novel inactivating mutation of the

[53196] gonadotropin-releasing hormone receptor. J. Clin. Endocr. Metab. 84:

[53197] 3811-3816, 1999.

[53198] 8548.

[53199] Szende, B.; Srkalovic, G.; Timar, J.; Mulchahey, J. J.; Neill,

[53200] J. D.; Lapis, K.; Csikos, A.; Szepeshazi, K.; Schally, A. V.: Localization

[53201] of receptors for luteinizing hormone-releasing hormone in pancreatic

[53202] and mammary cancer cells. Proc. Nat. Acad. Sci. 88: 4153-4156, 1991.

[53203] 8549.

[53204] Aganna, E.; Aksentijevich, I.; Hitman, G. A.; Kastner, D. L.; Hoepelman,

[53205] A. I. M.; Posma, F. D.; Zweers, E. J. K.; McDermott, M. F.: Tumor

[53206] necrosis factor receptor-associated periodic syndrome (TRAPS) in a

[53207] Dutch family: evidence for a TNFRSF1A mutation with reduced penetrance. Europ.

[53208] J. Hum. Genet. 9: 63-66, 2001.

[53209] 8550.

[53210] Mulley, J.; Saar, K.; Hewitt, G.; Ruschendorf, F.; Phillips, H.;

[53211] Colley, A.; Sillence, D.; Reis, A.; Wilson, M.: Gene localization

[53212] for an autosomal dominant familial periodic fever to 12p13. Am. J.

[53213] Hum. Genet. 62: 884-889, 1998.

[53214] 8551.

[53215] Williamson, L. M.; Hull, D.; Mehta, R.; Reeves, W. G.; Robinson,

[53216] B. H. B.; Toghill, P. J.: Familial hibernian fever. Quart. J. Med. 51:

[53217] 469-480, 1982.

[53218] 8552.

[53219] Bahler, M.; Kehrer, I.; Gordon, L.; Stoffler, H.-E.; Olsen, A.

[53220] S.: Physical mapping of human myosin-IXB (MYO9B), the human orthologue
[53221] of the rat myosin myr 5, to chromosome 19p13.1. Genomics 43: 107-109,
[53222] 1997.

[53223] 8553.

[53224] Miyajima, N.; Kadowaki, Y.; Fukushige, S.; Shimizu, S.; Semba,
[53225] K.; Yamanashi, Y.; Matsubara, K.; Toyoshima, K.; Yamamoto, T.: Identification
[53226] of two novel members of erbA superfamily by molecular cloning: the
[53227] gene products of the two are highly related to each other. Nucleic
[53228] Acids Res. 16: 11057-11073, 1988.

[53229] 8554.

[53230] Qiu, Y.; Krishnan, V.; Zeng, Z.; Gilbert, D. J.; Copeland, N. G.;
[53231] Gibson, L.; Yang-Feng, T.; Jenkins, N. A.; Tsai, M.-J.; Tsai, S. Y.
[53232] : Isolation, characterization, and chromosomal localization of mouse

[53233] and human COUP-TF I and II genes. Genomics 29: 240-246, 1995.

[53234] 8555.

[53235] Hawiger, J.; Timmons, S.; Kloczewiak, M.; Strong, D. D.; Doolittle,

[53236] R. F.: Gamma and alpha chains of human fibrinogen possess sites reactive

[53237] with human platelet receptors. Proc. Nat. Acad. Sci. 79: 2068-2071,

[53238] 1982.

[53239] 8556.

[53240] Henry, I.; Uzan, G.; Weil, D.; Nicolas, H.; Kaplan, J. C.; Marguerie,

[53241] G.; Kahn, A.; Junien, C.: The genes coding for the A-alpha, B-beta,

[53242] and gamma chains of fibrinogen are located on chromosome 4. (Abstract) Cytogenet.

[53243] Cell Genet. 37: 490-491, 1984.

[53244] 8557.

[53245] Koopman, J.; Haverkate, F.; Briet, E.; Lord, S. T.: A congenitally

[53246] abnormal fibrinogen (Vlissingen) with a 6-base deletion in the gamma-chain

[53247] gene, causing defective calcium binding and impaired fibrin polymerization. J.

[53248] Biol. Chem. 266: 13456-13461, 1991.

[53249] 8558.

[53250] Koopman, J.; Haverkate, F.; Lord, S.; Caekebeke-Peerlinck, K.;

[53251] Brommer, E.; Briet, E.: A six base deletion in the gamma-chain gene

[53252] of dysfibrinogen Vlissingen, coding for asn319 and asp320, resulting

[53253] in defective interaction with calcium. (Abstract) Thromb. Haemost. 62:

[53254] 158 only, 1989.

[53255] 8559.

[53256] Lounes, K. C.; Soria, C.; Mirshahi, S. S.; Desvignes, P.; Mirshahi,

[53257] M.; Bertrand, O.; Bonnet, P.; Koopman, J.; Soria, J.: Fibrinogen

[53258] Ales: a homozygous case of dysfibrinogenemia (gamma-asp330-to-val)

[53259] characterized by a defective fibrin polymerization site 'a'. Blood 96:
[53260] 3473-3479, 2000.

[53261] 8560.

[53262] Marder, V. J.: Personal Communication. Philadelphia, Pa. 12/8/1974.
[53263] 8561.

[53264] Margaglione, M.; Santacroce, R.; Colaizzo, D.; Seripa, D.; Vecchione,
[53265] G.; Lupone, M. R.; De Lucia, D.; Fortina, P.; Grandone, E.; Perricone,
[53266] C.; Di Minno, G.: A G-to-A mutation in IVS-3 of the human gamma fibrinogen
[53267] gene causing afibrinogenemia due to abnormal RNA splicing. Blood 96:
[53268] 2501-2505, 2000.

[53269] 8562.

[53270] Matsuda, M.; Baba, M.; Morimoto, K.; Nakamikawa, C.: 'Fibrinogen
[53271] Tokyo II': an abnormal fibrinogen with an impaired polymerization

[53272] site on the aligned DD domain of fibrin molecules. J. Clin. Invest. 72:

[53273] 1034-1041, 1983.

[53274] 8563.

[53275] Miyata, T.; Furukawa, K.; Iwanaga, S.; Takamatsu, J.; Saito, H.

[53276] : Fibrinogen Nagoya, a replacement of glutamine-329 by arginine in

[53277] the gamma-chain that impairs the polymerization of fibrin monomer. J.

[53278] Biochem. 105: 10-14, 1989.

[53279] 8564.

[53280] Mosesson, M. W.; Amrani, D. L.; Menache, D.: Studies on the structural

[53281] abnormality of fibrinogen Paris I. J. Clin. Invest. 57: 782-790,

[53282] 1976.

[53283] 8565.

[53284] Mosesson, M. W.; Siebenlist, K. R.; DiOrio, J. P.; Matsuda, M.;

[53285] Hainfeld, J. F.; Wall, J. S.: The role of fibrinogen D domain

intermolecular
[53286] association sites in the polymerization of fibrin and fib-
rinogen Tokyo
[53287] II (gamma-275 Arg-to-Cys). J. Clin. Invest. 96:
1053-1058, 1995.
[53288] 8566.

[53289] Mullin, J. L.; Brennan, S. O.; Ganly, P. S.; George, P. M.:
Fibrinogen
[53290] Hillsborough: a novel gamma-gly309asp dysfibrinogen
with impaired
[53291] clotting. Blood 99: 3597-3601, 2002.

[53292] 8567.

[53293] Okumura, N.; Furihata, K.; Terasawa, F.; Nagagoshi, R.;
Ueno,
[53294] I.; Katsuyama, T.: Fibrinogen Matsumoto I: a gamma 364
asp-to-his
[53295] (GAT-to-CAT) substitution associated with defective fibrin
polymerization. Thromb.
[53296] Haemost. 75: 887-891, 1996.

[53297] 8568.

[53298] Okumura, N.; Gorkun, O. V.; Lord, S. T.: Severely impaired

polymerization
[53299] of recombinant fibrinogen gamma-364 asp-to-his, the substitution discovered
[53300] in a heterozygous individual. J. Biol. Chem. 272: 29596-29601, 1997.
[53301] 8569.

[53302] Reber, P.; Furlan, M.; Henschen, A.; Kaudewitz, H.; Barbui, T.;
[53303] Hilgard, P.; Nenci, G. G.; Berrettini, M.; Beck, E. A.: Three abnormal
[53304] fibrinogen variants with the same amino acid substitution (gamma275
[53305] arg-to-his): fibrinogens Bergamo II, Essen and Perugia. Thromb. Haemost. 56:
[53306] 401-406, 1986.

[53307] 8570.

[53308] Reber, P.; Furlan, M.; Rupp, C.; Kehl, M.; Henschen, A.; Mannucci,
[53309] P. M.; Beck, E. A.: Characterization of fibrinogen Milano I: amino
[53310] acid exchange gamma-330 asp-to-val impairs fibrin polymerization. Blood 67:

[53311] 1751-1756, 1986.

[53312] 8571.

[53313] Rixon, M. W.; Chung, D. W.; Davie, E. W.: Nucleotide se-
quence
[53314] of the gene for the gamma chain of human fibrinogen.
Biochemistry 24:
[53315] 2077-2086, 1985.

[53316] 8572.

[53317] Rosenberg, J. B.; Newman, P. J.; Mosesson, M. W.; Guillin,
M.-C.;
[53318] Amrani, D. L.: Paris I dysfibrinogenemia: a point mutation
in intron
[53319] 8 results in insertion of a 15 amino acid sequence in fib-
rinogen gamma-chain. Thromb.
[53320] Haemost. 69: 217-220, 1993.

[53321] 8573.

[53322] Schmelzer, C. H.; Ebert, R. F.; Bell, W. R.: Fibrinogen Balti-
more
[53323] IV: congenital dysfibrinogenemia with a gamma-275
(arg-to-cys) substitution. Thromb.
[53324] Res. 56: 307-316, 1989.

[53325] 8574.

[53326] Siebenlist, K. R.; Mosesson, M. W.; Di Orio, J. P.; Tavori, S.;

[53327] Tatarsky, I.; Rimon, A.: The polymerization of fibrin pre-
pared from

[53328] fibrinogen Haifa (gamma-275-arg-to-his). Thromb.
Haemost. 62: 875-879,

[53329] 1989.

[53330] 8575.

[53331] Terukina, S.; Matsuda, M.; Hirata, H.; Takeda, Y.; Miyata,
T.;

[53332] Takao, T.; Shimonishi, Y.: Substitution of gamma-arg-275
by cys in

[53333] an abnormal fibrinogen, 'fibrinogen Osaka II': evidence for
a unique

[53334] solitary cystine structure at the mutation site. J. Biol.
Chem. 263:

[53335] 13579-13587, 1988.

[53336] 8576.

[53337] Terukina, S.; Yamazumi, K.; Okamoto, K.; Yamashita, H.;
Ito, Y.;

[53338] Matsuda, M.: Fibrinogen Kyoto III: a congenital dysfibrino-

gen with

[53339] a gamma aspartic acid-330 to tyrosine substitution. Blood 74: 2681-2687,

[53340] 1989.

[53341] 8577.

[53342] Yamazumi, K.; Shimura, K.; Terukina, S.; Takahashi, N.; Matsuda,

[53343] M.: A gamma methionine-310 to threonine substitution and consequent

[53344] N-glycosylation at gamma asparagine-308 identified in a congenital

[53345] dysfibrinogenemia associated with posttraumatic bleeding, fibrinogen

[53346] Asahi. J. Clin. Invest. 83: 1590-1597, 1989.

[53347] 8578.

[53348] Yamazumi, K.; Terukina, S.; Onohara, S.; Matsuda, M.: Normal

[53349] plasmic cleavage of the gamma-chain variant of 'fibrinogen Saga' with

[53350] an arg275-to-his substitution. Thromb. Haemost. 60: 476-480, 1988.

[53351] 8579.

[53352] Yoshida, N.; Hirata, H.; Morigami, Y.; Imaoka, S.; Matsuda, M.;

[53353] Yamazumi, K.; Asakura, S.: Characterization of an abnormal fibrinogen

[53354] Osaka V with the replacement of gamma-arginine 375 by glycine: the

[53355] lack of high affinity calcium binding to D-domains and the lack of

[53356] protective effect of calcium on fibrinolysis. J. Biol. Chem. 267:

[53357] 2753-2759, 1992.

[53358] 8580.

[53359] Petek, E.; Windpassinger, C.; Vincent, J. B.; Cheung, J.; Boright,

[53360] A. P.; Scherer, S. W.; Kroisel, P. M.; Wagner, K.: Disruption of

[53361] a novel gene (IMMP2L) by a breakpoint in 7q31 associated with Tourette

[53362] syndrome. Am. J. Hum. Genet. 848-858, 2001.

[53363] 8581.

[53364] Azen, E. A.; Leutenegger, W.; Peters, E. H.: Evolutionary and

[53365] dietary aspects of salivary basic (Pb) and post Pb (PPb) proteins

[53366] in anthropoid primates. Nature 273: 775-778, 1978.

[53367] 8582.

[53368] Peters, E. H.; Goodfriend, T.; Azen, E. A.: Human Pb, human post-Pb,

[53369] and nonhuman primate Pb proteins: immunological and biochemical relationships. Biochem.

[53370] Genet. 15: 947-962, 1977.

[53371] 8583.

[53372] Sabatini, L. M.; Azen, E. A.: Histatins, a family of salivary

[53373] histidine-rich proteins, are encoded by at least two loci (HIS1 and

[53374] HIS2). Biochem. Biophys. Res. Commun. 160: 495-502, 1989.

[53375] 8584.

[53376] vanderSpek, J. C.; Wyandt, H. E.; Skare, J. C.; Milunsky, A.; Oppenheim,

[53377] F. G.; Troxler, R. F.: Localization of the genes for histatins to

[53378] human chromosome 4q13 and tissue distribution of the

mRNAs. Am. J.

[53379] Hum. Genet. 45: 381-387, 1989.

[53380] 8585.

[53381] Azen, E. A.: Properties of salivary basic proteins showing polymorphism. Biochem.

[53382] Genet. 9: 69-86, 1973.

[53383] 8586.

[53384] Azen, E. A.: Personal Communication. Madison, Wis. 3/27/1990.

[53385] 8587.

[53386] Azen, E. A.: Genetic polymorphism of basic proteins from parotid

[53387] saliva. Science 176: 673-674, 1972.

[53388] 8588.

[53389] Helmerhorst, E. J.; Troxler, R. F.; Oppenheim, F. G.: The human

[53390] salivary peptide histatin 5 exerts its antifungal activity through

[53391] the formation of reactive oxygen species. Proc. Nat. Acad. Sci. 98:

[53392] 14637-14642, 2001.

[53393] 8589.

[53394] Oppenheim, F. G.; Xu, T.; McMillian, F. M.; Levitz, S. M.; Diamond,

[53395] R. D.; Offner, G. D.; Troxler, R. F.: Histatins, a novel family of

[53396] histidine-rich proteins in human parotid secretion. Isola-tion, characterization,

[53397] primary structure, and fungistatic effects on Candida albi-cans. J.

[53398] Biol. Chem. 263: 7472-7477, 1988.

[53399] 8590.

[53400] Peters, E. H.; Azen, E. A.: Isolation and partial characteri-zation

[53401] of human parotid basic proteins. Biochem. Genet. 15: 925-946, 1977.

[53402] 8591.

[53403] Kaelbling, M.; Eddy, R.; Shows, T. B.; Copeland, N. G.; Gilbert,

[53404] D. J.; Jenkins, N. A.; Klinger, H. P.; O'Hara, B.: Localization of

[53405] the human gene allowing infection by gibbon ape leukemia virus to

[53406] human chromosome region 2q11-q14 and to the homologous region on mouse

[53407] chromosome 2. J. Virol. 65: 1743-1747, 1991.

[53408] 8592.

[53409] Kavanaugh, M. P.; Miller, D. G.; Zhang, W.; Law, W.; Kozak, S.

[53410] L.; Kabat, D.; Miller, A. D.: Cell-surface receptors for gibbon ape

[53411] leukemia virus and amphotropic murine retrovirus are inducible sodium-dependent

[53412] phosphate symporters. Proc. Nat. Acad. Sci. 91: 7071-7075, 1994.

[53413] 8593.

[53414] O'Hara, B.; Johann, S. V.; Klinger, H. P.; Blair, D. G.; Rubinson,

[53415] H.; Dunn, K. J.; Sass, P.; Vitek, S. M.; Robins, T.: Characterization

[53416] of a human gene conferring sensitivity to infection by gibbon ape

[53417] leukemia virus. Cell Growth Differ. 1: 119-127, 1990.

[53418] 8594.

[53419] Palmer, G.; Manen, D.; Bonjour, J.-P.; Caverzasio, J.: Species-specific
[53420] mechanisms control the activity of the Pit1/PIT1 phosphate transporter
[53421] gene promoter in mouse and human. Gene 279: 49-62, 2001.
[53422] 8595.

[53423] Welch, W. J.; Diamond, M. I.: Glucocorticoid modulation of androgen
[53424] receptor nuclear aggregation and cellular toxicity is associated with
[53425] distinct forms of soluble expanded polyglutamine protein. Hum. Molec.
[53426] Genet. 10: 3063-3074, 2001.

[53427] 8596.

[53428] Aizawa, T.; Asanuma, N.; Terauchi, Y.; Suzuki, N.; Komatsu, M.;
[53429] Itoh, N.; Nakabayashi, T.; Hidaka, H.; Ohnota, H.; Yamauchi, K.; Yasuda,
[53430] K.; Yazaki, Y.; Kadowaki, T.; Hashizume, K.: Analysis of the pancreatic
[53431] beta cell in the mouse with targeted disruption of the

pancreatic
[53432] beta cell-specific glucokinase gene. Biochem. Biophys. Res. Commun. 229:
[53433] 460-465, 1996.

[53434] 8597.

[53435] Bali, D.; Svetlanov, A.; Lee, H.-W.; Fusco-DeMane, D.; Leiser,
[53436] M.; Li, B.; Barzilai, N.; Surana, M.; Hou, H.; Fleischer, N.; DePinho,
[53437] R.; Rossetti, L.; Efrat, S.: Animal model for maturity-onset diabetes
[53438] of the young generated by disruption of the mouse glucokinase gene. J.
[53439] Biol. Chem. 270: 21464-21467, 1995.

[53440] 8598.

[53441] Byrne, M. M.; Sturis, J.; Clement, K.; Vionnet, N.; Pueyo, M. E.;
[53442] Stoffel, M.; Takeda, J.; Passa, P.; Cohen, D.; Bell, G. I.; Velho,
[53443] G.; Froguel, P.; Polonsky, K. S.: Insulin secretory abnormalities
[53444] in subjects with hyperglycemia due to glucokinase muta-

tions. J. Clin.
[53445] Invest. 93: 1120-1130, 1994.

[53446] 8599.

[53447] Chiu, K. C.; Tanizawa, Y.; Permutt, M. A.: Non-sense mutation
[53448] of glucokinase gene. (Letter) Lancet 341: 385-386, 1993.

[53449] 8600.

[53450] Concannon, P.: Personal Communication. Seattle, Wash. 7/26/1995.
[53451] 8601.

[53452] Froguel, P.; Velho, G.: Non-sense mutation of glucokinase gene.
[53453] (Letter) Lancet 341: 385 only, 1993.

[53454] 8602.

[53455] Froguel, P.; Zouali, H.; Vionnet, N.; Velho, G.; Vaxillaire, M.;
[53456] Sun, F.; Lesage, S.; Stoffel, M.; Takeda, J.; Passa, P.; Permutt,
[53457] M. A.; Beckmann, J. S.; Bell, G. I.; Cohen, D.: Familial hyperglycemia
[53458] due to mutations in glucokinase: definition of a subtype of

diabetes
[53459] mellitus. New Eng. J. Med. 328: 697-702, 1993.

[53460] 8603.

[53461] Glaser, B.; Kesavan, P.; Heyman, M.; Davis, E.; Cuesta, A.; Buchs,
[53462] A.; Stanley, C. A.; Thornton, P. S.; Permutt, M. A.; Matschinsky,
[53463] F. M.; Herold, K. C.: Familial hyperinsulinism caused by an activating
[53464] glucokinase mutation. New Eng. J. Med. 338: 226-230, 1998.

[53465] 8604.

[53466] Glaser, B.; Thornton, P. S.; Herold, K.; Stanley, C. A.: Clini-cal
[53467] and molecular heterogeneity of familial hyperinsulinism. (Letter) J.
[53468] Pediat. 133: 801-802, 1998.

[53469] 8605.

[53470] Grupe, A.; Hultgren, B.; Ryan, A.; Ma, Y. H.; Bauer, M.; Stewart,
[53471] T. A.: Transgenic knockouts reveal a critical requirement

for pancreatic

[53472] beta cell glucokinase in maintaining glucose homeostasis. Cell 83:

[53473] 69-78, 1995.

[53474] 8606.

[53475] Hattersley, A. T.; Turner, R. C.; Permutt, M. A.; Patel, P.; Tanizawa,

[53476] Y.; Chiu, K. C.; O'Rahilly, S.; Watkins, P. J.; Wainscoat, J. S.:

[53477] Linkage of type 2 diabetes to the glucokinase gene. Lancet 339:

[53478] 1307-1310, 1992.

[53479] 8607.

[53480] Heimberg, H.; De Vos, A.; Moens, K.; Quartier, E.; Bouwens, L.;

[53481] Pipeleers, D.; Van Schaftingen, E.; Madsen, O.; Schuit, F.: The glucose

[53482] sensor protein glucokinase is expressed in glucagon-producing alpha-cells. Proc.

[53483] Nat. Acad. Sci. 93: 7036-7041, 1996.

[53484] 8608.

[53485] Katagiri, H.; Asano, T.; Ishihara, H.; Inukai, K.; Anai, M.;

Miyazaki,
[53486] J.; Tsukuda, K.; Kikuchi, M.; Yazaki, Y.; Oka, Y.: Nonsense mutation
[53487] of glucokinase gene in late-onset non-insulin-dependent diabetes mellitus. Lancet 340:
[53488] 1316-1317, 1992.

[53489] 8609.

[53490] Matschinsky, F.; Liang, Y.; Kesavan, P.; Wang, L.; Froguel, P.;
[53491] Velho, G.; Cohen, D.; Permutt, M. A.; Tanizawa, Y.; Jetton, T. L.;
[53492] Niswender, K.; Magnuson, M. A.: Glucokinase as pancreatic beta-cell
[53493] glucose sensor and diabetes gene. J. Clin. Invest. 92: 2092-2098,
[53494] 1993.

[53495] 8610.

[53496] Matschinsky, F. M.: Glucokinase as glucose sensor and metabolic
[53497] signal generator in pancreatic beta-cells and hepatocytes. Diabetes 39:
[53498] 647-652, 1990.

[53499] 8611.

[53500] Matsutani, A.; Janssen, R.; Donis-Keller, H.; Permutt, M. A.:

[53501] A polymorphic (CA)n repeat element maps the human glucokinase gene

[53502] (GCK) to chromosome 7p. Genomics 12: 319-325, 1992.

[53503] 8612.

[53504] Mishra, S. K.; Helms, C.; Dorsey, D.; Permutt, M. A.; Donis-Keller,

[53505] H.: A 2-cM genetic linkage map of human chromosome 7p that includes

[53506] 47 loci. Genomics 12: 326-334, 1992.

[53507] 8613.

[53508] Njolstad, P. R.; Sovik, O.; Cuesta-Munoz, A.; Bjorkhaug, L.; Massa,

[53509] O.; Barbetti, F.; Undlien, D. E.; Shiota, C.; Magnuson, M. A.; Molven,

[53510] A.; Matschinsky, F. M.; Bell, G. I.: Neonatal diabetes mellitus due

[53511] to complete glucokinase deficiency. New Eng. J. Med. 344: 1588-1592,

[53512] 2001.

[53513] 8614.

[53514] Permutt, M. A.; Chiu, K. C.; Tanizawa, Y.: Glucokinase and NIDDM:

[53515] a candidate gene that paid off. Diabetes 41: 1367-1372, 1992.

[53516] 8615.

[53517] Randle, P. J.: Glucokinase and candidate genes for type 2 (non-insulin-dependent)

[53518] diabetes mellitus. Diabetologia 36: 269-275, 1993.

[53519] 8616.

[53520] Rowe, R. E.; Wapelhorst, B.; Bell, G. I.; Risch, N.; Spielman,

[53521] R. S.; Concannon, P.: Linkage and association between insulin-dependent

[53522] diabetes mellitus (IDDM) susceptibility and markers near the glucokinase

[53523] gene on chromosome 7. Nature Genet. 10: 240-245, 1995.

[53524] 8617.

[53525] Stoffel, M.; Bell, K. L.; Blackburn, C. L.; Powell, K. L.; Seo,

[53526] T. S.; Takeda, J.; Vionnet, N.; Xiang, K.-S.; Gidh-Jain, M.; Pilkis,
[53527] S. J.; Ober, C.; Bell, G. I.: Identification of glucokinase mutations
[53528] in subjects with gestational diabetes mellitus. Diabetes 42: 937-940,
[53529] 1993.

[53530] 8618.

[53531] Hayashi, K.; Yano, H.; Hashida, T.; Takeuchi, R.; Takeda, O.; Asada,
[53532] K.; Takahashi, E.; Kato, I.; Sobue, K.: Genomic structure of the
[53533] human caldesmon gene. Proc. Nat. Acad. Sci. 89: 12122-12126, 1992.
[53534] 8619.

[53535] Humphrey, M. B.; Herrera-Sosa, H.; Gonzalez, G.; Lee, R.; Bryan,
[53536] J.: Cloning of cDNAs encoding human caldesmons. Gene 112: 197-204,
[53537] 1992.

[53538] 8620.

[53539] Dickson, K. M.; Bergeron, J. J. M.; Shames, I.; Colby, J.; Nguyen,
[53540] D. T.; Chevet, E.; Thomas, D. Y.; Snipes, G. J.: Association of calnexin
[53541] with mutant peripheral myelin protein-22 ex vivo: a basis for 'gain-of-function'
[53542] ER diseases. Proc. Nat. Acad. Sci. 99: 9852-9857, 2002.

[53543] 8621.

[53544] Gray, P. W.; Byers, M. G.; Eddy, R. L.; Shows, T. B.: The as-signment
[53545] of the calnexin gene to the q35 region of chromosome 5. (Abstract) Human
[53546] Genome Mapping Workshop 93 9 only, 1993.

[53547] 8622.

[53548] Schrag, J. D.; Bergeron, J. J. M.; Li, Y.; Borisova, S.; Hahn,
[53549] M.; Thomas, D. Y.; Cygler, M.: The structure of calnexin, an ER chaperone
[53550] involved in quality control of protein folding. Molec. Cell 8: 633-644,
[53551] 2001.

[53552] 8623.

[53553] Tjoelker, L. W.; Seyfried, C. E.; Eddy, R. L., Jr.; Byers, M. G.;

[53554] Shows, T. B.; Calderon, J.; Schreiber, R. B.; Gray, P. W.: Human,

[53555] mouse, and rat calnexin cDNA cloning: identification of potential

[53556] calcium binding motifs and gene localization to human chromosome 5. Biochemistry 33:

[53557] 3229-3236, 1994.

[53558] 8624.

[53559] Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Ohmi-Imajoh,

[53560] S.; Suzuki, K.; Shimizu, Y.; Shimizu, N.: Four genes for the calpain

[53561] family locate on four distinct human chromosomes. Cytogenet. Cell

[53562] Genet. 51: 1054-1055, 1989.

[53563] 8625.

[53564] Chen, C.-S.; Bejeck, B. E.; Kersey, J. H.: A mapping study of

[53565] 13 genes on human chromosome bands 4q11-q25. Cytogenet. Cell Genet. 69:

[53566] 260-265, 1995.

[53567] 8626.

[53568] Fujiwara, Y.; Miwa, M.; Nogami, M.; Okumura, K.; Nobori, T.; Suzuki,

[53569] T.; Ueda, M.: Genomic organization and chromosomal localization of

[53570] the human casein gene family. Hum. Genet. 99: 368-373, 1997.

[53571] 8627.

[53572] Rijnkels, M.; Meershoek, E.; de Boer, H. A.; Pieper, F. R.: Physical

[53573] map and localization of the human casein gene locus. Mammalian Genome 8:

[53574] 285-286, 1997.

[53575] 8628.

[53576] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Silver

[53577] Spring, Md.: National Biomedical Research Foundation (pub.) 1976.

[53578] 8629.

[53579] Ferretti, L.; Leone, P.; Sgaramella, V.: Long range restric-

tion

[53580] analysis of the bovine casein genes. Nucleic Acids Res. 18: 6829-6833,

[53581] 1990.

[53582] 8630.

[53583] Lonnerdal, B.; Bergstrom, S.; Andersson, Y.; Hjalmarsson, K.; Sundqvist,

[53584] A. K.; Hernell, O.: Cloning and sequencing of a cDNA encoding human

[53585] milk beta-casein. FEBS Lett. 269: 153-156, 1990.

[53586] 8631.

[53587] McConkey, E. H.; Menon, R.; Williams, G.; Baker, E.; Sutherland,

[53588] G. R.: Assignment of the gene for beta-casein (CSN2) to 4q13-q21

[53589] in humans and 3p13-p12 in chimpanzees. Cytogenet. Cell Genet. 72:

[53590] 60-62, 1996.

[53591] 8632.

[53592] Menon, R. S.; Chang, Y.-F.; Jeffers, K. F.; Ham, R. G.: Exon skipping

[53593] in human beta-casein. Genomics 12: 13-17, 1992.

[53594] 8633.

[53595] Menon, R. S.; Chang, Y.-F.; Jeffers, K. F.; Jones, C.; Ham, R.

[53596] G.: Regional localization of human beta-casein gene (CSN2) to 4pter-q21. Genomics 13:

[53597] 225-226, 1992.

[53598] 8634.

[53599] Menon, R. S.; Ham, R. G.: Human beta-casein: partial cDNA sequence

[53600] and apparent polymorphism. Nucleic Acids Res. 17: 2869, 1989.

[53601] 8635.

[53602] Threadgill, D. W.; Womack, J. E.: Genomic analysis of the major

[53603] bovine milk protein genes. Nucleic Acids Res. 18: 6935-6942, 1990.

[53604] 8636.

[53605] Wyszomierski, S. L.; Rosen, J. M.: Cooperative effects of STAT5

[53606] (signal transducer and activator of transcription 5) and C/

EBP beta
[53607] (CCAAT/enhancer-binding protein-beta) on beta-casein gene transcription
[53608] are mediated by the glucocorticoid receptor. Molec. En-
docr. 15:
[53609] 228-240, 2001.

[53610] 8637.

[53611] Kurotaki, N.; Imaizumi, K.; Harada, N.; Masuno, M.; Kon-
doh, T.;
[53612] Nagai, T.; Ohashi, H.; Naritomi, K.; Tsukahara, M.; Makita,
Y.; Sugimoto,
[53613] T.; Sonoda, T.; and 11 others: Haploinsufficiency of NSD1
causes
[53614] Sotos syndrome. Nature Genet. 30: 365-366, 2002.

[53615] 8638.

[53616] Vandenberg, P.; Khillan, J. S.; Prockop, D. J.; Helminen, H.;

[53617] Kontusaari, S.; Ala-Kokko, L.: Expression of a partially
deleted
[53618] gene of a human type II procollagen (COL2A1) in trans-
genic mice produces
[53619] a chondrodysplasia. Proc. Nat. Acad. Sci. 88: 7640-7644,

1991.

[53620] 8639.

[53621] van der Rest, M.; Rosenberg, L. C.; Olsen, B. R.; Poole, A. R.

[53622] : Chondrocalcin is identical with the C-propeptide of type II procollagen. Biochem.

[53623] J. 237: 923-925, 1986.

[53624] 8640.

[53625] Vikkula, M.; Ritvaniemi, P.; Vuorio, A. F.; Kaitila, I.; Ala-Kokko,

[53626] L.; Peltonen, L.: A mutation in the amino-terminal end of the triple

[53627] helix of type II collagen causing severe osteochondrodys-plasia. Genomics 16:

[53628] 282-285, 1993.

[53629] 8641.

[53630] Vissing, H.; D'Alessio, M.; Lee, B.; Ramirez, F.; Godfrey, M.;

[53631] Hollister, D. W.: Glycine to serine substitution in the triple helical

[53632] domain of pro-alpha-1(II) collagen results in a lethal peri-natal form

[53633] of short-limbed dwarfism. J. Biol. Chem. 264: 18265-18267, 1989.

[53634] 8642.

[53635] Wagner, H.: Ein bisher unbeknantes Erbleiden des Auges (degeneratio

[53636] hyaloideo-retinalis hereditaria), beobachtet im Kanton, Zurich. Klin.

[53637] Mbl. Augenheilk. 100: 840-857, 1938.

[53638] 8643.

[53639] Weis, M. A.; Wilkin, D. J.; Kim, H. J.; Wilcox, W. R.; Lachman,

[53640] R. S.; Rimoin, D. L.; Cohn, D. H.; Eyre, D. R.: Structurally abnormal

[53641] type II collagen in a severe form of Kniest dysplasia caused by an

[53642] exon 24 skipping mutation. J. Biol. Chem. 273: 4761-4768, 1998.

[53643] 8644.

[53644] Weiss, E. H.; Cheah, S. E.; Grosveld, F. G.; Dahl, H. H. M.; Solomon,

[53645] E.; Flavell, R. A.: Isolation and characterization of a human collagen

[53646] alpha1(I)-like gene from a cosmid library. Nucleic Acids Res. 10:
[53647] 1981-1992, 1982.

[53648] 8645.

[53649] Wilkin, D. J.; Bogaert, R.; Lachman, R. S.; Rimoin, D. L.; Eyre,
[53650] D. R.; Cohn, D. H.: A single amino acid substitution (G103D) in the
[53651] type II collagen triple helix produces Kniest dysplasia. Hum. Molec.
[53652] Genet. 3: 1999-2003, 1994.

[53653] 8646.

[53654] Williams, C. J.; Considine, E. L.; Knowlton, R. G.; Reginato,
[53655] A.; Neumann, G.; Harrison, D.; Buxton, P.; Jimenez, S.; Prockop, D.
[53656] J.: Spondyloepiphyseal dysplasia and precocious os-teoarthritis in
[53657] a family with an arg75-to-cys mutation in the procollagen type II
[53658] gene (COL2A1). Hum. Genet. 92: 499-505, 1993.

[53659] 8647.

[53660] Williams, C. J.; Rock, M.; Considine, E.; McCarron, S.; Gow, P.;

[53661] Ladda, R.; McLain, D.; Michels, V. M.; Murphy, W.; Prockop, D. J.;

[53662] Ganguly, A.: Three new point mutations in type II procollagen (COL2A1)

[53663] and identification of a fourth family with the COL2A1 arg519-to-cys

[53664] base substitution using conformation sensitive gel electrophoresis. Hum.

[53665] Molec. Genet. 4: 309-312, 1995.

[53666] 8648.

[53667] Winterpacht, A.; Hilbert, M.; Schwarze, U.; Mundlos, S.; Spranger,

[53668] J.; Zabel, B. U.: Kniest and Stickler dysplasia phenotypes caused

[53669] by collagen type II gene (COL2A1) defect. Nature Genet. 3: 323-326,

[53670] 1993.

[53671] 8649.

[53672] Winterpacht, A.; Schwarze, U.; Mundlos, S.; Menger, H.; Spranger,

[53673] J.; Zabel, B.: Alternative splicing as the result of a type II procollagen
[53674] gene (COL2A1) mutation in a patient with Kniest dysplasia. Hum. Molec.
[53675] Genet. 3: 1891-1893, 1994.

[53676] 8650.

[53677] Wu, J.-J.; Eyre, D. R.: Structural analysis of cross-linking
[53678] domains in cartilage type XI collagen: insights on polymeric assembly. J.
[53679] Biol. Chem. 270: 18865-18870, 1995.

[53680] 8651.

[53681] Yoo, T. J.; Tomoda, K.; Stuart, J. M.; Kang, A. H.; Townes, A.
[53682] S.: Type II collagen-induced autoimmune otospongiosis: a preliminary
[53683] report. Ann. Otol. Rhinol. Laryng. 92: 103-108, 1983.

[53684] 8652.

[53685] Young, M. F.; Vogeli, G.; Nunez, A. M.; Fernandez, M. P.; Sullivan,
[53686] M.; Sobel, M. E.: Isolation of cDNA and genomic DNA clones encoding

[53687] type II collagen. Nucleic Acids Res. 12: 4207-4228, 1984.

[53688] 8653.

[53689] Bateman, J. F.; Lamande, S. R.; Dahl, H.-H. M.; Chan, D.; Cole,

[53690] W. G.: Substitution of arginine for glycine 664 in the collagen alpha-1(I)

[53691] chain in lethal perinatal osteogenesis imperfecta. J. Biol. Chem. 263:

[53692] 11627-11630, 1988.

[53693] 8654.

[53694] Brandi, M. L.; Aurbach, G. D.; Fitzpatrick, L. A.; Quarto, R.;

[53695] Spiegel, A. M.; Bliziotes, M. M.; Norton, J. A.; Doppman, J. L.; Marx,

[53696] S. J.: Parathyroid mitogenic activity in plasma from patients with

[53697] familial multiple endocrine neoplasia type 1. New Eng. J. Med. 314:

[53698] 1287-1293, 1986.

[53699] 8655.

[53700] Colao, A.; Di Somma, C.; Pivonello, R.; Loche, S.; Aimaretti,

[53701] G.; Cerbone, G.; Faggiano, A.; Corneli, G.; Ghigo, E.; Lombardi, G.

[53702] : Bone loss is correlated to the severity of growth hormone deficiency

[53703] in adult patients with hypopituitarism. J. Clin. Endocr. Metab. 84:

[53704] 1919-1924, 1999.

[53705] 8656.

[53706] Lee, S.; Lin, M.; Mele, A.; Cao, Y.; Farmar, J.; Russo, D.; Redman,

[53707] C.: Proteolytic processing of big endothelin-3 by the Kell blood

[53708] group protein. Blood 94: 1440-1450, 1999.

[53709] 8657.

[53710] Pingault, V.; Bondurand, N.; Lemort, N.; Sancandi, M.; Ceccherini,

[53711] I.; Hugot, J.-P.; Jouk, P.-S.; Goossens, M.: A heterozygous endothelin

[53712] 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage

[53713] effect of EDN3/EDNRB gene mutations on neurocristopathy phenotypes?

[53714] (Letter) J. Med. Genet. 38: 205-208, 2001.

[53715] 8658.

[53716] Svensson, P.-J.; Von Tell, D.; Molander, M.-L.; Anvret, M.; Nordenskjold,

[53717] A.: A heterozygous frameshift mutation in the endothelin-3 (EDN-3)

[53718] gene in isolated Hirschsprung's disease. Pediat. Res. 45: 714-717,

[53719] 1999.

[53720] 8659.

[53721] Cyr, C.; Huebner, K.; Druck, T.; Kris, R.: Cloning and chromosomal

[53722] localization of a human endothelin ETA receptor. Biochem. Biophys.

[53723] Res. Commun. 181: 184-190, 1991.

[53724] 8660.

[53725] Hosoda, K.; Nakao, K.; Tamura, N.; Arai, H.; Ogawa, Y.; Suga, S.;

[53726] Nakanishi, S.; Imura, H.: Organization, structure, chromosomal assignment,

[53727] and expression of the gene encoding the human endothe-

lin-A receptor. J.

[53728] Biol. Chem. 267: 18797-18804, 1992.

[53729] 8661.

[53730] Tzourio, C.; El Amrani, M.; Poirier, O.; Nicaud, V.; Bousser, M.-G.;

[53731] Alperovitch, A.: Association between migraine and endothelin type

[53732] A receptor (ETA -231 A/G) gene polymorphism. Neurology 56: 1273-1277,

[53733] 2001.

[53734] 8662.

[53735] Carrasquillo, M. M.; McCallion, A. S.; Puffenberger, E. G.; Kashuk,

[53736] C. S.; Nouri, N.; Chakravarti, A.: Genome-wide association study

[53737] and mouse model identify interaction between RET and EDNRB pathways

[53738] in Hirschsprung disease. Nature Genet. 32: 237-244, 2002.

[53739] 8663.

[53740] Charlton, P.; Guida, L.; Copeland, N.; Jenkins, N.; Munroe,

D.;

[53741] Greenberg, F.; Fiedorek, F. T.; Nicholls, R. D.: Genetic approach

[53742] to function of the neuropeptide galanin. (Abstract) Am. J. Hum. Genet. 53

[53743] (suppl.): A1137, 1993.

[53744] 8664.

[53745] Evans, H.; Baumgartner, M.; Shine, J.; Herzog, H.: Genomic organization

[53746] and localization of the gene encoding human prepro-galanin. Genomics 18:

[53747] 473-477, 1993.

[53748] 8665.

[53749] Evans, H. F.; Shine, J.: Human galanin: molecular cloning reveals

[53750] a unique structure. Endocrinology 129: 1682-1684, 1991.

[53751] 8666.

[53752] Harris, G. W.: Neural control of the pituitary gland. Physiol.

[53753] Rev. 28: 139-179, 1948.

[53754] 8667.

[53755] Holmes, F. E.; Mahoney, S.; King, V. R.; Bacon, A.; Kerr, N. C.

[53756] H.; Pachnis, V.; Curtis, R.; Priestley, J. V.; Wynick, D.: Targeted

[53757] disruption of the galanin gene reduces the number of sensory neurons

[53758] and their regenerative capacity. Proc. Nat. Acad. Sci. 97: 11563-11568,

[53759] 2000.

[53760] 8668.

[53761] Lopez, F. J.; Merchenthaler, I.; Ching, M.; Wisniewski, M. G.;

[53762] Negro-Vilar, A.: Galanin: a hypothalamic-hypophysiotropic hormone

[53763] modulating reproductive functions. Proc. Nat. Acad. Sci. 88: 4508-4512,

[53764] 1991.

[53765] 8669.

[53766] Lundkvist, J.; Land, T.; Kahl, U.; Bedecs, K.; Bartfai, T.: cDNA

[53767] sequence, ligand binding, and regulation of galanin/GMAP in mouse

[53768] brain. Neurosci. Lett. 200: 121-124, 1995.

[53769] 8670.

[53770] O'Meara, G.; Coumis, U.; Ma, S. Y.; Kehr, J.; Mahoney, S.; Bacon,

[53771] A.; Allen, S. J.; Holmes, F.; Kahl, U.; Wang, F. H.; Kearns, I. R.;

[53772] Ove-Ogren, S.; Dawbarn, D.; Mufson, E. J.; Davies, C.; Dawson, G.;

[53773] Wynick, D.: Galanin regulates the postnatal survival of a subset

[53774] of basal forebrain cholinergic neurons. Proc. Nat. Acad. Sci. 97:

[53775] 11569-11574, 2000.

[53776] 8671.

[53777] Rattan, S.: Role of galanin in the gut. Gastroenterology 100:

[53778] 1762-1768, 1991.

[53779] 8672.

[53780] Schmidt, W. E.; Kratzin, H.; Eckart, K.; Drevs, D.; Mund-kowski,

[53781] G.; Clemens, A.; Katsoulis, S.; Schafer, H.; Gallwitz, B.;

Creutzfeldt,
[53782] W.: Isolation and primary structure of pituitary human galanin, a
[53783] 30-residue nonamidated neuropeptide. Proc. Nat. Acad. Sci. 88: 11435-11439,
[53784] 1991.

[53785] 8673.

[53786] Steiner, R. A.; Hohmann, J. G.; Holmes, A.; Wrenn, C. C.; Cadd,
[53787] G.; Jureus, A.; Clifton, D. K.; Luo, M.; Gutshall, M.; Ma, S. Y.;
[53788] Mufson, E. J.; Crawley, J. N.: Galanin transgenic mice dis-play cognitive
[53789] and neurochemical deficits characteristic of Alzheimer's disease. Proc.
[53790] Nat. Acad. Sci. 98: 4184-4189, 2001.

[53791] 8674.

[53792] Wynick, D.; Small, C. J.; Bacon, A.; Holmes, F. E.; Norman, M.;
[53793] Ormandy, C. J.; Kilic, E.; Kerr, N. C. H.; Ghatei, M.; Tala-mantes,
[53794] F.; Bloom, S. R.; Pachnis, V.: Galanin regulates prolactin

release
[53795] and lactotroph proliferation. Proc. Nat. Acad. Sci. 95:
12671-12676,
[53796] 1998.

[53797] 8675.

[53798] Appert, H.; Rutherford, T.; Tarr, G.; Wiest, J.; Thomford,
N.;
[53799] McCorquodale, M.; McCorquodale, D. J.: Isolation of a
cDNA for human
[53800] galactosyltransferase. (Abstract) Am. J. Hum. Genet. 39:
A186 only,
[53801] 1986.

[53802] 8676.

[53803] Appert, H. E.; Rutherford, T. J.; Tarr, G. E.; Wiest, J. S.;
Thomford,
[53804] N. R.; McCorquodale, D. J.: Isolation of a cDNA coding for
human
[53805] galactosyltransferase. Biochem. Biophys. Res. Commun.
139: 163-168,
[53806] 1986.

[53807] 8677.

[53808] Asano, M.; Furukawa, K.; Kido, M.; Matsumoto, S.; Umesaki, Y.;

[53809] Kochibe, N.; Iwakura, Y.: Growth retardation and early death of beta-1,4-galactosyltransferase

[53810] knockout mice with augmented proliferation and abnormal differentiation

[53811] of epithelial cells. EMBO J. 16: 1850-1857, 1997.

[53812] 8678.

[53813] Duncan, A. M. V.; McCorquodale, M. M.; Morgan, C.; Rutherford,

[53814] T. J.; Appert, H. E.; McCorquodale, D. J.: Chromosomal localization

[53815] of the gene for a human galactosyltransferase (GT-1). Biochem. Biophys.

[53816] Res. Commun. 141: 1185-1188, 1986.

[53817] 8679.

[53818] Furukawa, K.; Roth, S.; Sawicki, J.: Several galactosyltransferase

[53819] activities are associated with mouse chromosome 17. Genetics 114:

[53820] 983-991, 1986.

[53821] 8680.

[53822] Hansske, B.; Thiel, C.; Lubke, T.; Hasilik, M.; Honing, S.; Peters,

[53823] V.; Heidemann, P. H.; Hoffmann, G. F.; Berger, E. G.; von Figura,

[53824] K.; Korner, C.: Deficiency of UDP-galac-tose:N-acetylglucosamine beta-1,4-galactosyltransferase

[53825] I causes the congenital disorder of glycosylation type IId. J. Clin.

[53826] Invest. 109: 725-733, 2002.

[53827] 8681.

[53828] Kotani, N.; Asano, M.; Iwakura, Y.; Takasaki, S.: Knockout of

[53829] mouse beta-1,4-galactosyltransferase-1 gene results in a dramatic

[53830] shift of outer chain moieties of N-glycans from type 2 to type 1 chains

[53831] in hepatic membrane and plasma glycoproteins. Biochem. J. 357: 827-834,

[53832] 2001.

[53833] 8682.

[53834] Lo, N.-W.; Shaper, J. H.; Pevsner, J.; Shaper, N. L.: The expanding

[53835] beta-4-galactosyltransferase gene family: messages from the databanks. Glycobiology 8:

[53836] 517-526, 1998.

[53837] 8683.

[53838] Masri, K. A.; Appert, H. E.; Fukuda, M. N.: Identification of

[53839] the full-length coding sequence for human galactosyltransferase (beta-N-acetylglucosaminide:

[53840] beta-1,4-galactosyltransferase). Biochem. Biophys. Res. Commun. 157:

[53841] 657-663, 1988.

[53842] 8684.

[53843] Mengle-Gaw, L.; McCoy-Haman, M. F.; Tiemeier, D. C.: Genomic

[53844] structure and expression of human beta-1,4-galactosyltransferase. Biochem.

[53845] Biophys. Res. Commun. 176: 1269-1276, 1991.

[53846] 8685.

[53847] Schnyder-Candrian, S.; Borsig, L.; Moser, R.; Berger, E. G.:

[53848] Localization of alpha-1,3-fucosyltransferase VI in Weibel-Palade bodies

[53849] of human endothelial cells. Proc. Nat. Acad. Sci. 97: 8369-8374,

[53850] 2000.

[53851] 8686.

[53852] Shaper, N. L.; Shaper, J. H.; Bertness, V.; Chang, H.; Kirsch,

[53853] I. R.; Hollis, G. F.: The human galactosyltransferase gene is on

[53854] chromosome 9 at band p13. Somat. Cell Molec. Genet. 12: 633-636,

[53855] 1986.

[53856] 8687.

[53857] Shaper, N. L.; Shaper, J. H.; Meuth, J. L.; Fox, J. L.; Chang,

[53858] H.; Kirsch, I. R.; Hollis, G. F.: Bovine galactosyltransferase: identification

[53859] of a clone by direct immunological screening of a cDNA expression

[53860] library. Proc. Nat. Acad. Sci. 83: 1573-1577, 1986.

[53861] 8688.

[53862] Puech, A.; Saint-Jore, B.; Funke, B.; Gilbert, D. J.; Sirotkin,

[53863] H.; Copeland, N. G.; Jenkins, N. A.; Kucherlapati, R.; Morrow, B.;

[53864] Skoultchi, A. I.: Comparative mapping of the human 22q11 chromosomal

[53865] region and the orthologous region in mice reveals complex changes

[53866] in gene organization. Proc. Nat. Acad. Sci. 94: 14608-14613, 1997.

[53867] 8689.

[53868] Janus, C.; Pearson, J.; McLaurin, J.; Mathews, P. M.; Jiang, Y.;

[53869] Schmidt, S. D.; Chishti, M. A.; Horne, P.; Heslin, D.; French, J.;

[53870] Mount, H. T. J.; Nixon, R. A.; Mercken, M.; Bergeron, C.; Fraser,

[53871] P. E.; St George-Hyslop, P.; Westaway, D.: A-beta peptide immunization

[53872] reduces behavioural impairment and plaques in a model of Alzheimer's

[53873] disease. Nature 408: 979-982, 2000.

[53874] 8690.

[53875] Jenkins, E. C.; Devine-Gage, E. A.; Robakis, N. K.; Yao, X.-L.;

[53876] Brown, W. T.; Houck, G. E., Jr.; Wolfe, G.; Ramakrishna, N.; Silverman,

[53877] W. P.; Wisniewski, H. M.: Fine mapping of an Alzheimer disease-associated

[53878] gene encoding beta-amyloid protein. Biochem. Biophys. Res. Commun. 151:

[53879] 1-8, 1988.

[53880] 8691.

[53881] Jones, C. T.; Morris, S.; Yates, C. M.; Moffoot, A.; Sharpe, C.;

[53882] Brock, D. J. H.; St. Clair, D.: Mutation in codon 713 of the beta

[53883] amyloid precursor protein gene presenting with schizophrenia. Nature

[53884] Genet. 1: 306-309, 1992.

[53885] 8692.

[53886] Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.-H.; Goldstein, L.

[53887] S.: Axonal transport of amyloid precursor protein is mediated by

[53888] direct binding to the kinesin light chain subunit of kinesin-I. Neuron 28:

[53889] 449-459, 2000.

[53890] 8693.

[53891] Kamino, K.; Orr, H. T.; Payami, H.; Wijsman, E. M.; Alonso, M.

[53892] E.; Pulst, S. M.; Anderson, L.; O'dahl, S.; Nemens, E.; White, J.

[53893] A.; Sadovnick, A. D.; Ball, M. J.; Kaye, J.; Warren, A.; McInnis,

[53894] M.; Antonarakis, S. E.; Korenberg, J. R.; Sharma, V.; Kukull, W.;

[53895] Larson, E.; Heston, L. L.; Martin, G. M.; Bird, T. D.; Schellenberg,

[53896] G. D.: Linkage and mutational analysis of familial Alzheimer disease

[53897] kindreds for the APP gene region. Am. J. Hum. Genet. 51: 998-1014,

[53898] 1992.

[53899] 8694.

[53900] Kaneko, I.; Yamada, N.; Sakuraba, Y.; Kamenosono, M.; Tutumi,

[53901] S.: Suppression of mitochondrial succinate dehydroge-
nase, a primary
[53902] target of beta-amyloid, and its derivative racemized at ser
residue. J.
[53903] Neurochem. 65: 2585-2593, 1995.

[53904] 8695.

[53905] Karlinsky, H.; Vaula, G.; Haines, J. L.; Ridgley, J.; Bergeron,

[53906] C.; Mortilla, M.; Tupler, R. G.; Percy, M. E.; Robitaille, Y.;
Noldy,
[53907] N. E.; Yip, T. C. K.; Tanzi, R. E.; Gusella, J. F.; Becker, R.;
Berg,
[53908] J. M.; Crapper McLachlan, D. R.; St. George-Hyslop, P. H.:
Molecular
[53909] and prospective phenotypic characterization of a pedigree
with familial
[53910] Alzheimer's disease and a missense mutation in codon
717 of the beta-amyloid
[53911] precursor protein gene. Neurology 42: 1445-1453, 1992.

[53912] 8696.

[53913] Korenberg, J. R.; Pulst, S.-M.; Neve, R. L.; West, R.: The
Alzheimer

[53914] amyloid precursor protein maps to human chromosome 21 bands q21.105-q21.05. Genomics 5:
[53915] 124-127, 1989.

[53916] 8697.

[53917] Kumar-Singh, S.; De Jonghe, C.; Cruts, M.; Kleinert, R.; Wang,
[53918] R.; Mercken, M.; De Strooper, B.; Vanderstichele, H.; Lofgren, A.;
[53919] Vanderhoeven, I.; Backhovens, H.; Vanmechelen, E.; Kroisel, P. M.;
[53920] Van Broeckhoven, C.: Nonfibrillar diffuse amyloid deposition due
[53921] to a gamma-42-secretase site mutation points to an essential role
[53922] for N-truncated A-beta-42 in Alzheimer's disease. Hum. Molec. Genet. 9:
[53923] 2589-2598, 2000.

[53924] 8698.

[53925] Lashuel, H. A.; Hartley, D.; Petre, B. M.; Walz, T.; Lansbury,
[53926] P. T., Jr.: Amyloid pores from pathogenic mutations. Nature 418:

[53927] 291 only, 2002.

[53928] 8699.

[53929] Levy, E.; Carman, M. D.; Fernandez-Madrid, I. J.; Power, M. D.;

[53930] Lieberburg, I.; van Duinen, S. G.; Bots, G. T. A. M.; Luyendijk, W.;

[53931] Frangione, B.: Mutation of the Alzheimer's disease amyloid gene in

[53932] hereditary cerebral hemorrhage, Dutch type. Science 248: 1124-1126,

[53933] 1990.

[53934] 8700.

[53935] Lewis, J.; Dickson, D. W.; Lin, W.-L.; Chisholm, L.; Corral, A.;

[53936] Jones, G.; Yen, S.-H.; Sahara, N.; Skipper, L.; Yager, D.; Eckman,

[53937] C.; Hardy, J.; Hutton, M.; McGowan, E.: Enhanced neu-

rofibrillary

[53938] degeneration in transgenic mice expressing mutant tau and APP. Science 293:

[53939] 1487-1491, 2001.

[53940] 8701.

[53941] Lorenzo, A.; Yuan, M.; Zhang, Z.; Paganetti, P. A.; Sturch-
ler-Pierrat,
[53942] C.; Staufenbiel, M.; Mautino, J.; Vigo, F. S.; Sommer, B.;
Yankner,
[53943] B. A.: Amyloid beta interacts with the amyloid precursor
protein:
[53944] a potential toxic mechanism in Alzheimer's disease. Na-
ture Neurosci. 3:
[53945] 460-464, 2000.

[53946] 8702.

[53947] Lovett, M.; Goldgaber, D.; Ashley, P.; Cox, D. R.; Gajdusek,
D.
[53948] C.; Epstein, C. J.: The mouse homolog of the human amy-
loid beta protein
[53949] (AD-AP) gene is located on the distal end of mouse chro-
mosome 16:
[53950] further extension of the homology between human chro-
mosome 21 and
[53951] mouse chromosome 16. Biochem. Biophys. Res. Commun.
144: 1069-1075,
[53952] 1987.

[53953] 8703.

[53954] Luyendijk, W.; Bots, G. T. A. M.: Hereditary cerebral haem-
orrhage.(Letter) Scand.
[53955] J. Clin. Lab. Invest. 46: 391 only, 1986.

[53956] 8704.

[53957] Luyendijk, W.; Bots, G. T. A. M.; Vegter-van der Vlis, M.;
Went,
[53958] L. N.; Frangione, B.: Hereditary cerebral haemorrhage
caused by cortical
[53959] amyloid angiopathy. J. Neurol. Sci. 85: 267-280, 1988.

[53960] 8705.

[53961] Majocha, R. E.; Benes, F. M.; Reifel, J. L.; Rodenrys, A. M.;

[53962] Marotta, C. A.: Laminar-specific distribution and infras-
tructural
[53963] detail of amyloid in the Alzheimer disease cortex visual-
ized by computer-enhanced
[53964] imaging of epitopes recognized by monoclonal antibod-
ies. Proc. Nat.
[53965] Acad. Sci. 85: 6182-6186, 1988.

[53966] 8706.

[53967] Mant, R.; Asherson, P.; Gill, M.; McGuffin, P.; Owen, M.: Schizophrenia

[53968] scepticism.(Letter) Nature Genet. 2: 12 only, 1992.

[53969] 8707.

[53970] Maruyama, K.; Tomita, T.; Shinozaki, K.; Kume, H.; Asada, H.;

[53971] Saido, T. C.; Ishiura, S.; Iwatsubo, T.; Obata, K.: Familial Alzheimer's

[53972] disease-linked mutations at val717 of amyloid precursor protein are

[53973] specific for the increased secretion of A-beta-42(43). Biochem. Biophys.

[53974] Res. Commun. 227: 730-735, 1996.

[53975] 8708.

[53976] Masters, C. L.; Simms, G.; Weinman, N. A.; Multhaup, G.; McDonald,

[53977] B. L.; Beyreuther, K.: Amyloid plaque core protein in Alzheimer disease

[53978] and Down syndrome. Proc. Nat. Acad. Sci. 82: 4245-4249, 1985.

[53979] 8709.

[53980] Meziane, H.; Dodart, J.-C.; Mathis, C.; Little, S.; Clemens, J.;

[53981] Paul, S. M.; Ungerer, A.: Memory-enhancing effects of se-creted forms

[53982] of the beta-amyloid precursor protein in normal and amnestic mice. Proc.

[53983] Nat. Acad. Sci. 95: 12683-12688, 1998.

[53984] 8710.

[53985] Miravalle, L.; Tokuda, T.; Chiarle, R.; Giaccone, G.; Bugiani,

[53986] O.; Tagliavini, F.; Frangione, B.; Ghiso, J.: Substitutions at codon

[53987] 22 of Alzheimer's A-beta peptide induce diverse confor-mational changes

[53988] and apoptotic effects in human cerebral endothelial cells. J. Biol.

[53989] Chem. 275: 27110-27116, 2000.

[53990] 8711.

[53991] Denker, S. P.; Huang, D. C.; Orlowski, J.; Furthmayr, H.; Barber,

[53992] D. L.: Direct binding of the Na-H exchanger NHE1 to ERM proteins

[53993] regulates the cortical cytoskeleton and cell shape independently of

[53994] H(+) translocation. Molec. Cell 6: 1425-1436, 2000.

[53995] 8712.

[53996] Dudley, C. R. K.; Giuffra, L. A.; Tippett, P.; Kidd, K. K.; Reeders,

[53997] S. T.: The Na+/H+ antiporter: a 'melt' polymorphism allows regional

[53998] mapping to the short arm of chromosome 1. Hum. Genet. 86: 79-83,

[53999] 1990.

[54000] 8713.

[54001] Franchi, A.; Perucca-Lostanlen, D.; Pouyssegur, J.: Functional

[54002] expression of a human Na+/H+ antiporter gene transfected into antiporter-deficient

[54003] mouse L cells. Proc. Nat. Acad. Sci. 83: 9388-9392, 1986.

[54004] 8714.

[54005] Lifton, R. P.; Sardet, C.; Pouyssegur, J.; Lalouel, J.-M.: Cloning

[54006] of the human genomic amiloride-sensitive Na+/H+ an-

tiporter gene, identification
[54007] of genetic polymorphisms, and localization on the genetic map of chromosome
[54008] 1p. Genomics 7: 131-135, 1990.

[54009] 8715.

[54010] Mattei, M.-G.; Galloni, M.; Sardet, C.; Franchi, A.; Counil-lon,
[54011] L.; Passage, E.; Pouyssegur, J.: Localization of the an-tiporter gene
[54012] (APNH) and chromosomal homology between human 1p, mouse 4 and Chinese
[54013] hamster 2q. (Abstract) Cytogenet. Cell Genet. 51: 1041, 1989.

[54014] 8716.

[54015] Mattei, M.-G.; Sardet, C.; Franchi, A.; Pouyssegur, J.: Chromosomal
[54016] mapping of the amiloride-sensitive Na+/H+ antiporter gene. (Abstract) Cytogenet.
[54017] Cell Genet. 46: 658-659, 1987.

[54018] 8717.

[54019] Mattei, M.-G.; Sardet, C.; Franchi, A.; Pouyssegur, J.: The

human

[54020] amiloride-sensitive Na+/H+ antiporter: localization to chromosome

[54021] 1 by in situ hybridization. Cytogenet. Cell Genet. 48: 6-8, 1988.

[54022] 8718.

[54023] Mendoza, S. A.: The Na+/H+ antiport is a mediator of cell proliferation. Acta

[54024] Paediat. Scand. 76: 545-547, 1987.

[54025] 8719.

[54026] Morahan, G.; Rakar, S.: Localization of the mouse Na+/H+ exchanger

[54027] gene on distal chromosome 4. Genomics 15: 231-232, 1993.

[54028] 8720.

[54029] Sardet, C.; Franchi, A.; Pouyssegur, J.: Molecular cloning, primary

[54030] structure, and expression of the human growth factor-activatable Na(+)/H(+)

[54031] antiporter. Cell 56: 271-280, 1989.

[54032] 8721.

[54033] Kitawaki, J.; Koshiba, H.; Ishihara, H.; Kusuki, I.; Tsukamoto,

[54034] K.; Honjo, H.: Progesterone induction of 17-beta-hydroxysteroid dehydrogenase

[54035] type 2 during the secretory phase occurs in the en-dometrium of estrogen-dependent

[54036] benign diseases but not in normal endometrium. J. Clin. Endocr. Metab. 85:

[54037] 3292-3296, 2000.

[54038] 8722.

[54039] Suzuki, T.; Sasano, H.; Andersson, S.; Mason, J. I.: 3-beta-hydroxysteroid

[54040] dehydrogenase/delta(5-4)-isomerase activity associated with the human

[54041] 17-beta-hydroxysteroid dehydrogenase type 2 isoform. J. Clin. Endocr.

[54042] Metab. 85: 3669-3672, 2000.

[54043] 8723.

[54044] Wu, L.; Einstein, M.; Geissler, W. M.; Chan, H. K.; Elliston, K.

[54045] O.; Andersson, S.: Expression cloning and characterization of human

[54046] 17-beta-hydroxysteroid dehydrogenase type 2, a micro-somal enzyme possessing

[54047] 20-alpha-hydroxysteroid dehydrogenase activity. J. Biol. Chem. 268:

[54048] 12964-12969, 1993.

[54049] 8724.

[54050] Davare, M. A.; Avdonin, V.; Hall, D. D.; Peden, E. M.; Burette,

[54051] A.; Weinberg, R. J.; Horne, M. C.; Hoshi, T.; Hell, J. W.: A beta-2

[54052] adrenergic receptor signaling complex assembled with the Ca(2+) channel

[54053] Ca(V)1.2. Science 293: 98-101, 2001.

[54054] 8725.

[54055] Jacobs, P. A.; Brunton, M.; Frackiewicz, A.; Newton, M.; Cook,

[54056] P. J. L.; Robson, E. B.: Studies on a family with three cytogenetic

[54057] markers. Ann. Hum. Genet. 33: 325-336, 1970.

[54058] 8726.

[54059] Brown, A.; Bernier, G.; Mathieu, M.; Rossant, J.; Kothary, R.:

[54060] The mouse dystonia musculorum gene is a neural isoform of bullous

[54061] pemphigoid antigen 1. Nature Genet. 10: 301-306, 1995.

[54062] 8727.

[54063] Brown, A.; Dalpe, G.; Mathieu, M.; Kothary, R.: Cloning and characterization

[54064] of the neural isoforms of human dystonin. Genomics 29: 777-780,

[54065] 1995.

[54066] 8728.

[54067] Copeland, N. G.; Gilbert, D. J.; Li, K.; Sawamura, D.; Giu-dice,

[54068] G. J.; Chu, M.-L.; Jenkins, N. A.; Uitto, J.: Chromosomal lo-calization

[54069] of mouse bullous pemphigoid antigens, BPAG1 and BPAG2: identification

[54070] of a new region of homology between mouse and human chromosomes. Genomics 15:

[54071] 180-181, 1993.

[54072] 8729.

[54073] Diaz, L. A.; Ratrie, H., III; Saunders, W. S.; Futamura, S.; Squiquera,

[54074] H. L.; Anhalt, G. J.; Giudice, G. J.: Isolation of a human epi-dermal

[54075] cDNA corresponding to the 180-kD autoantigen recog-nized by bullous

[54076] pemphigoid and herpes gestationis sera: immunolocaliza-
tion of this
[54077] protein to the hemidesmosome. J. Clin. Invest. 86:
1088-1094, 1990.
[54078] 8730.

[54079] Guo, L.; Degenstein, L.; Dowling, J.; Yu, Q.-C.; Wollmann,
R.;
[54080] Perman, B.; Fuchs, E.: Gene targeting of BPAG1: abnormal-
ities in
[54081] mechanical strength and cell migration in stratified ep-
ithelia and
[54082] neurologic degeneration. Cell 81: 233-243, 1995.

[54083] 8731.

[54084] Minoshima, S.; Amagai, M.; Kudoh, J.; Fukuyama, R.;
Hashimoto,
[54085] T.; Nishikawa, T.; Shimizu, N.: Localization of the human
gene for
[54086] 230-kDa bullous pemphigoid autoantigen to the pter-q15
region of chromosome
[54087] 6. (Abstract) Cytogenet. Cell Genet. 58: 1914-1915, 1991.

[54088] 8732.

[54089] Minoshima, S.; Amagai, M.; Kudoh, J.; Fukuyama, R.; Hashimoto,
[54090] T.; Nishikawa, T.; Shimizu, N.: Localization of the human gene for
[54091] 230-kDal bullous pemphigoid autoantigen (BPAG1) to chromosome 6pter-q15. Cytogenet.
[54092] Cell Genet. 57: 30-32, 1991.

[54093] 8733.

[54094] Ryynanen, M.; Knowlton, R. G.; Kero, M.; Sawamura, D.; Li, K.;
[54095] Giudice, G. J.; Diaz, L. A.; Uitto, J.: Bullous pemphigoid antigens
[54096] (BPAGs): identification of RFLPs in human BPAG1 and BPAG2, and exclusion
[54097] as candidate genes in a large kindred with dominant epi-dermolysis
[54098] bullosa simplex. Genomics 11: 1025-1029, 1991.

[54099] 8734.

[54100] Sawamura, D.; Nomura, K.; Sugita, Y.; Mattei, M.-G.; Chu, M.-L.;
[54101] Knowlton, R.; Uitto, J.: Bullous pemphigoid antigen (BPAG1): cDNA

[54102] cloning and mapping of the gene to the short arm of human chromosome

[54103] 6. Genomics 8: 722-726, 1990.

[54104] 8735.

[54105] Stanley, J. R.; Tanaka, T.; Mueller, S.; Klaus-Kovtun, V.; Roop,

[54106] D.: Isolation of complementary DNA for bullous pemphigoid antigen

[54107] by use of patients' autoantibodies. J. Clin. Invest. 82: 1864-1870,

[54108] 1988.

[54109] 8736.

[54110] Tamai, K.; Sawamura, D.; Do, H. C.; Tamai, Y.; Li, K.; Uitto,

[54111] J.: The human 230-kD bullous pemphigoid antigen gene (BPAG1): exon-intron

[54112] organization and identification of regulatory tissue specific elements

[54113] in the promoter region. J. Clin. Invest. 92: 814-822, 1993.

[54114] 8737.

[54115] Yang, Y.; Bauer, C.; Strasser, G.; Wollman, R.; Julien, J.-P.;

[54116] Fuchs, E.: Integrators of the cytoskeleton that stabilize microtubules. Cell 98:

[54117] 229-238, 1999.

[54118] 8738.

[54119] Chavanas, S.; Gache, Y.; Tadini, G.; Pulkkinen, L.; Uitto, J.;

[54120] Ortonne, J. P.; Meneguzzi, G.: A homozygous in-frame deletion in

[54121] the collagenous domain of bullous pemphigoid antigen BP180 (type XVII

[54122] collagen) causes generalized atrophic benign epidermolysis bullosa. J.

[54123] Invest. Derm. 109: 74-78, 1997.

[54124] 8739.

[54125] Darling, T. N.; Koh, B. B.; Bale, S. J.; Compton, J. G.; Bauer,

[54126] J. W.; Hintner, H.; Yancey, K. B.: A deletion mutation in COL17A1

[54127] in five Austrian families with generalized atrophic benign epidermolysis

[54128] bullosa represents propagation of an ancestral allele. J. Invest.

[54129] Derm. 110: 170-173, 1998.

[54130] 8740.

[54131] Darling, T. N.; Yee, C.; Koh, B.; McGrath, J. A.; Bauer, J. W.;

[54132] Uitto, J.; Hintner, H.; Yancey, K. B.: Cycloheximide facili-
tates

[54133] the identification of aberrant transcripts resulting from a
novel

[54134] splice-site mutation in COL17A1 in a patient with gener-
alized atrophic

[54135] benign epidermolysis bullosa. J. Invest. Derm. 110:
165-169, 1998.

[54136] 8741.

[54137] Floeth, M.; Fiedorowicz, J.; Schacke, H.; Hammami-
Hauasli, N.;

[54138] Owaribe, K.; Trueb, R. M.; Bruckner-Tuderman, L.: Novel
homozygous

[54139] and compound heterozygous COL17A1 mutations associ-
ated with junctional

[54140] epidermolysis bullosa. J. Invest. Derm. 111: 528-533,
1998.

[54141] 8742.

[54142] Montgomery, J. C.; Shows, T. B.; Venta, P. J.; Tashian, R. E.:

[54143] Gene for novel human carbonic anhydrase (CA) isozyme on chromosome

[54144] 16 is unlinked to the CA1/CA2/CA3 gene cluster. (Abstract) Am. J.

[54145] Hum. Genet. 41: A229 only, 1987.

[54146] 8743.

[54147] Montgomery, J. C.; Venta, P. J.; Eddy, R. L.; Fukushima, Y.-S.;

[54148] Shows, T. B.; Tashian, R. E.: Characterization of the human gene

[54149] for a newly discovered carbonic anhydrase, CA VII, and its localization

[54150] to chromosome 16. Genomics 11: 835-848, 1991.

[54151] 8744.

[54152] Venta, P. J.; Montgomery, J. C.; Tashian, R. E.: Molecular genetics

[54153] of carbonic anhydrase isozymes. Isozymes: Curr. Top. Biol. Med. Res. 14:

[54154] 59-72, 1987.

[54155] 8745.

[54156] Aldred, P.; Fu, P.; Barrett, G.; Penschow, J. D.; Wright, R.

D.;

[54157] Coghlan, J. P.; Fernley, R. T.: Human secreted carbonic anhydrase:

[54158] cDNA cloning, nucleotide sequence, and hybridization histochemistry. Biochemistry 30:

[54159] 569-575, 1991.

[54160] 8746.

[54161] Karhumaa, P.; Leinonen, J.; Parkkila, S.; Kaunisto, K.; Tapanainen,

[54162] J.; Rajaniemi, H.: The identification of secreted carbonic anhydrase

[54163] VI as a constitutive glycoprotein of human and rat milk. Proc. Nat.

[54164] Acad. Sci. 98: 11604-11608, 2001.

[54165] 8747.

[54166] Murakami, H.; Sly, W. S.: Purification and characterization of

[54167] human salivary carbonic anhydrase. J. Biol. Chem. 262: 1382-1388,

[54168] 1987.

[54169] 8748.

[54170] Sutherland, G. R.; Baker, E.; Fernandez, K. E. W.; Callen, D. F.;
[54171] Aldred, P.; Coghlan, J. P.; Wright, R. D.; Fernley, R. T.: The gene
[54172] for human carbonic anhydrase VI (CA6) is on the tip of the short arm
[54173] of chromosome 1. Cytogenet. Cell Genet. 50: 149-150, 1989.
[54174] 8749.
[54175] Thatcher, B. J.; Doherty, A. E.; Orvisky, E.; Martin, B. M.; Henkin,
[54176] R. I.: Gustin from human parotid saliva is carbonic anhydrase VI. Biochem.
[54177] Biophys. Res. Commun. 250: 635-641, 1998.
[54178] 8750.
[54179] Chauhan, S. S.; Popescu, N. C.; Ray, D.; Fleischmann, R.; Gottesman,
[54180] M. M.; Troen, B. R.: Cloning, genomic organization, and chromosomal
[54181] localization of human cathepsin L. J. Biol. Chem. 268: 1039-1045,
[54182] 1993.

[54183] 8751.

[54184] Fan, Y.-S.; Byers, M. G.; Eddy, R. L.; Joseph, L.; Sukhatme, V.;

[54185] Chan, S.-J.; Shows, T. B.: Cathepsin L (CTSL) is located in the chromosome

[54186] 9q21-q22 region: a related sequence is located on chromosome 10. (Abstract) Cytogenet.

[54187] Cell Genet. 51: 996 only, 1989.

[54188] 8752.

[54189] Gal, S.; Gottesman, M. M.: Isolation and sequence of a cDNA for

[54190] human pro-(cathepsin L). Biochem. J. 253: 303-306, 1988.

[54191] 8753.

[54192] Joseph, L. J.; Chang, L. C.; Stamenkovich, D.; Sukhatme, V. P.

[54193] : Complete nucleotide and deduced amino acid sequences of human and

[54194] murine preprocathepsin L: an abundant transcript induced by transformation

[54195] of fibroblasts. J. Clin. Invest. 81: 1621-1629, 1988.

[54196] 8754.

[54197] Stypmann, J.; Glaser, K.; Roth, W.; Tobin, D. J.; Petermann, I.;

[54198] Matthias, R.; Monnig, G.; Haverkamp, W.; Breithardt, G.; Schmahl,

[54199] W.; Peters, C.; Reinheckel, T.: Dilated cardiomyopathy in mice deficient

[54200] for the lysosomal cysteine peptidase cathepsin L. Proc. Nat. Acad.

[54201] Sci. 99: 6234-6239, 2002.

[54202] 8755.

[54203] Kere, J.; Ruutu, T.; Davies, K. A.; Roninson, I. B.; Watkins, P.

[54204] C.; Winqvist, R.; de la Chapelle, A.: Chromosome 7 long arm deletion

[54205] in myeloid disorders: a narrow breakpoint region in 7q22 defined by

[54206] molecular mapping. Blood 73: 230-234, 1989.

[54207] 8756.

[54208] Neufeld, E. J.: Personal Communication. Boston, Mass. 2/21/1995.

[54209] 8757.

[54210] Neufeld, E. J.; Skalnik, D. G.; Lievens, P. M.-J.; Orkin, S. H.

[54211] : Human CCAAT displacement protein is homologous to the Drosophila

[54212] homeoprotein, cut. Nature Genet. 1: 50-55, 1992.

[54213] 8758.

[54214] Scherer, S. W.; Neufeld, E. J.; Lievens, P. M.-J.; Orkin, S. H.;

[54215] Kim, J.; Tsui, L.-C.: Regional localization of the CCAAT displacement

[54216] protein gene (CUTL1) to 7q22 by analysis of somatic cell hybrids. Genomics 15:

[54217] 695-696, 1993.

[54218] 8759.

[54219] Snyder, S. R.; Wang, J.; Waring, J. F.; Ginder, G. D.: Identification

[54220] of CCAAT displacement protein (CDP/cut) as a locus-specific repressor

[54221] of major histocompatibility complex gene expression in human tumor

[54222] cells. J. Biol. Chem. 276: 5323-5330, 2001.

[54223] 8760.

[54224] Zeng, W. R.; Scherer, S. W.; Koutsilieris, M.; Huizenga, J. J.;

[54225] Filteau, F.; Tsui, L.-C.; Nepveu, A.: Loss of heterozygosity and

[54226] reduced expression of the CUTL1 gene in uterine leiomyomas. Oncogene 14:

[54227] 2355-2365, 1997.

[54228] 8761.

[54229] Birkenmeier, E. H.; Gwynn, B.; Howard, S.; Jerry, J.; Gordon, J.

[54230] I.; Landschulz, W. H.; McKnight, S. L.: Tissue-specific expression,

[54231] developmental regulation, and genetic mapping of the gene encoding

[54232] CCAAT/enhancer binding protein. Genes Dev. 3: 1146-1156, 1989.

[54233] 8762.

[54234] Flodby, P.; Barlow, C.; Kylefjord, H.; Ahrlund-Richter, L.; Xanthopoulos,

[54235] K. G.: Increased hepatic cell proliferation and lung abnormalities

[54236] in mice deficient in CCAAT/enhancer binding protein alpha. J. Biol.

[54237] Chem. 271: 24753-24760, 1996.

[54238] 8763.

[54239] Landschulz, W. H.; Johnson, P. F.; McKnight, S. L.: The DNA binding

[54240] domain of the rat liver nuclear protein C/EBP is bipartite. Science 243:

[54241] 1681-1688, 1989.

[54242] 8764.

[54243] Blackburn, M. R.; Datta, S. K.; Kellems, R. E.: Adenosine deaminase-deficient

[54244] mice generated using a two-stage genetic engineering strategy exhibit

[54245] a combined immunodeficiency. J. Biol. Chem. 273: 5093-5100, 1998.

[54246] 8765.

[54247] Blaese, R. M.; Culver, K. W.; Miller, A. D.; Carter, C. S.; Fleisher,

[54248] T.; Clerici, M.; Shearer, G.; Chang, L., Chiang, Y.; Tolstoshev, P.;

[54249] Greenblatt, J. J.; Rosenberg, S. A.; Klein, H.; Berger, M.; Mullen,
[54250] C. A.; Ramsey, W. J.; Muul, L.; Morgan, R. A.; Anderson, W. F.: T
[54251] lymphocyte-directed gene therapy for ADA-SCID: initial trial results
[54252] after 4 years. Science 270: 475-480, 1995.

[54253] 8766.

[54254] Bollinger, M. E.; Arredondo-Vega, F. X.; Santisteban, I.; Schwarz,
[54255] K.; Hershfield, M. S.; Lederman, H. M.: Hepatic dysfunction as a
[54256] complication of adenosine deaminase deficiency. New Eng. J. Med. 334:
[54257] 1367-1371, 1996.

[54258] 8767.

[54259] Bonthron, D. T.; Markham, A. F.; Ginsburg, D.; Orkin, S. H.:
[54260] Identification of a point mutation in the adenosine deaminase gene
[54261] responsible for immunodeficiency. J. Clin. Invest. 76: 894-897,

[54262] 1985.

[54263] 8768.

[54264] Bordignon, C.; Notarangelo, L. D.; Nobili, N.; Ferrari, G.; Casorati,

[54265] G.; Panina, P.; Mazzolari, E.; Maggioni, D.; Rossi, C.; Servida, P.;

[54266] Ugazio, A. G.; Mavilio, F.: Gene therapy in peripheral blood lymphocytes

[54267] and bone marrow for ADA: immunodeficient patients. Science 270:

[54268] 470-475, 1995.

[54269] 8769.

[54270] Bortin, M. M.; Rimm, A. A. (eds.): Severe combined immunodeficiency

[54271] disease: characterization of the disease and results of transplantation. J.A.M.A. 238:

[54272] 591-600, 1977.

[54273] 8770.

[54274] Boss, G. R.; Thompson, L. F.; O'Connor, R. D.; Ziering, R. W.;

[54275] Seegmiller, J. E.: Ecto-5-prime-nucleotidase deficiency:

association
[54276] with adenosine deaminase deficiency and nonassociation with deoxyadenosine
[54277] toxicity. Clin. Immun. Immunopath. 19: 1-7, 1981.

[54278] 8771.

[54279] Bottini, N.; De Luca, D.; Saccucci, P.; Fiumara, A.; Elia, M.;

[54280] Porfirio, M. C.; Lucarelli, P.; Curatolo, P.: Autism: evidence of
[54281] association with adenosine deaminase genetic polymorphism. Neurogenetics 3:
[54282] 111-113, 2001.

[54283] 8772.

[54284] Chen, S.-H.; Ochs, H. D.; Scott, C. R.: Adenosine deaminase deficiency:
[54285] disappearance of adenine deoxynucleotides from a patient's erythrocytes
[54286] after successful marrow transplantation. J. Clin. Invest. 62: 1386-1389,
[54287] 1978.

[54288] 8773.

[54289] Chen, S.-H.; Ochs, H. D.; Scott, C. R.; Giblett, E. R.:

Adenosine
[54290] deaminase and nucleoside phosphorylase activity in patients with immunodeficiency
[54291] syndromes. Clin. Immun. Immunopath. 13: 156-160, 1979.
[54292] 8774.

[54293] Chen, S.-H.; Scott, C. R.; Giblett, E. R.: Adenosine deaminase:
[54294] demonstration of a 'silent' gene associated with combined immunodeficiency
[54295] disease. Am. J. Hum. Genet. 26: 103-107, 1974.

[54296] 8775.

[54297] Cohen, A.; Hirschhorn, R.; Horowitz, S. D.; Rubinstein, A.; Polmar,
[54298] S. H.; Hong, R.; Martin, D. W., Jr.: Deoxyadenosine triphosphate
[54299] as a potentially toxic metabolite in adenosine deaminase deficiency. Proc.
[54300] Nat. Acad. Sci. 75: 472-476, 1978.

[54301] 8776.

[54302] Cook, P. J. L.; Hopkinson, D. A.; Robson, E. B.: The linkage

[54303] relationships of adenosine deaminase. Ann. Hum. Genet. 34: 187-188,

[54304] 1970.

[54305] 8777.

[54306] Creagan, R. P.; Tischfield, J. A.; Nichols, E. A.; Ruddle, F.

[54307] H.: Autosomal assignment of the gene for the form of adenosine deaminase

[54308] which is deficient in patients with combined immunodeficiency syndrome.

[54309] (Letter) Lancet II: 1449 only, 1973.

[54310] 8778.

[54311] Culver, K. W.: Gene Therapy: A Primer for Physicians. Larchmont,

[54312] New York: Mary Ann Liebert, Inc. , 1996.

[54313] 8779.

[54314] Daddona, P. E.; Kelley, W. N.: Human adenosine deaminase: stoichiometry

[54315] of the adenosine deaminase-binding protein complex. Biochim. Biophys.

[54316] Acta 580: 302-311, 1979.

[54317] 8780.

[54318] Daddona, P. E.; Mitchell, B. S.; Meuwissen, H. J.; Davidson, B.

[54319] L.; Wilson, J. M.; Koller, C. A.: Adenosine deaminase deficiency

[54320] with normal immune function: an acidic enzyme mutation. J. Clin.

[54321] Invest. 72: 483-492, 1983.

[54322] 8781.

[54323] Detter, J. C.; Stamatoyannopoulos, G.; Giblett, E. R.; Motulsky,

[54324] A. G.: Adenosine deaminase: racial distribution and report of a new

[54325] phenotype. J. Med. Genet. 7: 356-357, 1970.

[54326] 8782.

[54327] Dissing, J.; Knudsen, B.: Adenosine-deaminase deficiency and

[54328] combined immunodeficiency syndrome. (Letter) Lancet II: 1316, 1972.

[54329] 8783.

[54330] Dissing, J.; Knudsen, J. B.: A new red cell adenosine deam-

inase
[54331] phenotype in man. Hum. Hered. 19: 375-377, 1969.

[54332] 8784.

[54333] Dyminski, J. W.; Daoud, A.; Lampkin, B. C.; Limouze, S.; Donofrio,
[54334] J.; Coleman, M. S.; Hutton, J. J.: Immunological and biochemical
[54335] profiles in response to transfusion therapy in adenosine deaminase-deficient
[54336] patient with severe combined immunodeficiency disease. Clin. Immun.
[54337] Immunopath. 14: 307-326, 1979.

[54338] 8785.

[54339] Egashira, M.; Ariga, T.; Kawamura, N.; Miyoshi, O.; Niikawa, N.;
[54340] Sakiyama, Y.: Visible integration of the adenosine deaminase (ADA)
[54341] gene into the recipient genome after gene therapy. Am. J. Med. Genet. 75:
[54342] 314-317, 1998.

[54343] 8786.

[54344] Ellis, N. A.; Lennon, D. J.; Proytcheva, M.; Alhadeff, B.; Henderson,

[54345] E. E.; German, J.: Somatic intragenic recombination within the mutated

[54346] locus BLM can correct the high SCE phenotype of Bloom syndrome cells. Am.

[54347] J. Hum. Genet. 57: 1019-1027, 1995.

[54348] 8787.

[54349] Ferrari, G.; Rossini, S.; Giavazzi, R.; Maggioni, D.; Nobili,

[54350] N.; Soldati, M.; Ungers, G.; Mavilio, F.; Gilboa, E.; Bor-dignon, C.

[54351] : An in vivo model of somatic cell gene therapy for human severe combined

[54352] immunodeficiency. Science 251: 1363-1366, 1991.

[54353] 8788.

[54354] Giblett, E. R.; Anderson, J. E.; Cohen, F.; Pollara, B.; Meuwissen,

[54355] H. J.: Adenosine-deaminase deficiency in two patients with severely

[54356] impaired cellular immunity. Lancet I: 1067-1069, 1972.

[54357] 8789.

[54358] Hart, S. L.; Lane, A. B.; Jenkins, T.: Partial adenosine deaminase
[54359] deficiency: another family from southern Africa. Hum. Genet. 74:
[54360] 307-312, 1986.

[54361] 8790.

[54362] Herbschleb-Voogt, E.; Scholten, J.-W.; Meera Khan, P.: Basic
[54363] defect in the expression of adenosine deaminase in ADA SCID disease.
[54364] II. Deficiency of ADA-CRM detected in heterozygote human-Chinese hamster
[54365] cell hybrids. Hum. Genet. 63: 121-125, 1983.

[54366] 8791.

[54367] Hershfield, M. S.: PEG-ADA: an alternative to haploidenti-cal
[54368] bone marrow transplantation and an adjunct to gene therapy for adenosine
[54369] deaminase deficiency. Hum. Mutat. 5: 107-112, 1995.

[54370] 8792.

[54371] Hershfield, M. S.; Buckley, R. H.; Greenberg, M. L.; Melton,

A.

[54372] L.; Schiff, R.; Hatem, C.; Kurtzberg, J.; Markert, M. L.; Kobayashi,

[54373] R. H.; Kobayashi, A. L.; Abuchowski, A.: Treatment of adenosine deaminase

[54374] deficiency with polyethylene glycol-modified adenosine deaminase. New

[54375] Eng. J. Med. 316: 589-596, 1987.

[54376] 8793.

[54377] Hershfield, M. S.; Kredich, N. M.: S-adenosylhomocysteine hydrolase

[54378] is an adenosine-binding protein: a target for adenosine toxicity. Science 202:

[54379] 757-760, 1978.

[54380] 8794.

[54381] Presley, J. F.; Ward, T. H.; Pfeifer, A. C.; Siggia, E. D.; Phair,

[54382] R. D.; Lippincott-Schwartz, J.: Dissection of COPI and Arf1 dynamics

[54383] in vivo and role in Golgi membrane transport. Nature 417: 187-193,

[54384] 2002.

[54385] 8795.

[54386] Tsai, S.-C.; Haun, R. S.; Tsuchiya, M.; Moss, J.; Vaughan, M.:

[54387] Isolation and characterization of the human gene for ADP-ribosylation

[54388] factor 3, a 20-kDa guanine nucleotide-binding protein activator of

[54389] cholera toxin. J. Biol. Chem. 266: 23053-23059, 1991.

[54390] 8796.

[54391] Lachman, H. M.; Papolos, D. F.; Saito, T.; Yu, Y. M.; Szum-lanski,

[54392] C. L.; Weinshilboum, R. M.: Human catechol-O-methyltransferase pharmacogenetics:

[54393] description of a functional polymorphism and its potential application

[54394] to neuropsychiatric disorders. Pharmacogenetics 6: 243-250, 1996.

[54395] 8797.

[54396] Tiihonen, J.; Hallikainen, T.; Lachman, H.; Saito, T.; Volavka,

[54397] J.; Kauhanen, J.; Salonen, J. T.; Ryynanen, O.-P.; Koulu, M.; Karvonen,

[54398] M. K.; Pohjalainen, T.; Syvalahti, E.; Hietala, J.: Association between
[54399] the functional variant of the catechol-O-methyltransferase (COMT)
[54400] gene and type 1 alcoholism. Molec. Psychiat. 4: 286-289, 1999.
[54401] 8798.

[54402] Alder, A.: Ueber konstitutionell bedingte Granulationsver-aenderungen
[54403] der Leukocyten. Dtsch. Arch. Klin. Med. 183: 372-378, 1939.
[54404] 8799.

[54405] Bohren, K. M.; Bullock, B.; Wermuth, B.; Gabbay, K. H.: The aldo-keto
[54406] reductase superfamily: cDNAs and deduced amino acid sequences of human
[54407] aldehyde and aldose reductases. J. Biol. Chem. 264: 9547-9551, 1989.
[54408] 8800.

[54409] Charlesworth, D.: Starch-gel electrophoresis of four en-zymes from
[54410] human red blood cells: glyceraldehyde-3-phosphate de-

hydrogenase, fructoaldolase,
[54411] glyoxalase II and sorbitol dehydrogenase. Ann. Hum. Genet. 35: 477-484,
[54412] 1972.

[54413] 8801.

[54414] Frade, R.; Balbo, M.; Barel, M.: RB18A, whose gene is localized
[54415] on chromosome 17q12-q21.1, regulates in vivo p53 transactivating activity. Cancer
[54416] Res. 60: 6585-6589, 2000.

[54417] 8802.

[54418] Wolffe, A. P.: Transcriptional control: sinful repression. Nature 387:
[54419] 16-17, 1997.

[54420] 8803.

[54421] Wu, L.; Aster, J. C.; Blacklow, S. C.; Lake, R.; Artavanis-Tsakonas,
[54422] S.; Griffin, J. D.: MAML1, a human homologue of Drosophila mastermind,
[54423] is a transcriptional co-activator for NOTCH receptors. Nature Genet. 26:

[54424] 484-489, 2000.

[54425] 8804.

[54426] Inohara, N.; Koseki, T.; del Peso, L.; Hu, Y.; Yee, C.; Chen, S.;

[54427] Carrio, R.; Merino, J.; Liu, D.; Ni, J.; Nunez, G.: Nod1, an Apaf-1-like

[54428] activator of caspase-9 and nuclear factor-kappa-B. J. Biol. Chem. 274:

[54429] 14560-14567, 1999.

[54430] 8805.

[54431] Weng, Z.; Fluckiger, A.-C.; Nisitani, S.; Wahl, M. I.; Le, L. Q.;

[54432] Hunter, C. A.; Fernal, A. A.; le Beau, M. M.; Witte, O. N.: A DNA

[54433] damage and stress inducible G protein-coupled receptor blocks cells

[54434] in G2/M. Proc. Nat. Acad. Sci. 95: 12334-12339, 1998.

[54435] 8806.

[54436] Charroux, B.; Pellizzoni, L.; Perkinson, R. A.; Shevchenko, A.;

[54437] Mann, M.; Dreyfuss, G.: Gemin3: a novel DEAD box protein

that interacts

[54438] with SMN, the spinal muscular atrophy gene product, and is a component

[54439] of gems. J. Cell Biol. 147: 1181-1193, 1999.

[54440] 8807.

[54441] Grundhoff, A. T.; Kremmer, E.; Tureci, O.; Glieden, A.; Gindorf,

[54442] C.; Atz, J.; Mueller-Lantzsch, N.; Schubach, W. H.; Grasser, F. A.

[54443] : Characterization of DP103, a novel DEAD box protein that binds to

[54444] the Epstein-Barr virus nuclear proteins EBNA2 and EBNA3C. J. Biol.

[54445] Chem. 274: 19136-19144, 1999.

[54446] 8808.

[54447] Hunt, S. L.; Hsuan, J. J.; Totty, N.; Jackson, R. J.: unr, a cel-lular

[54448] cytoplasmic RNA-binding protein with five cold-shock domains, is required

[54449] for internal initiation of translation of human rhinovirus RNA. Genes

[54450] Dev. 13: 437-448, 1999.

[54451] 8809.

[54452] Matsuda, S.; Katsumata, R.; Okuda, T.; Yamamoto, T.; Miyazaki,

[54453] K.; Senga, T.; Machida, K.; Thant, A. A.; Nakatsugawa, S.; Hamaguchi,

[54454] M.: Molecular cloning and characterization of human MAWD, a novel

[54455] protein containing WD-40 repeats frequently overex- pressed in breast

[54456] cancer. Cancer Res. 60: 13-17, 2000.

[54457] 8810.

[54458] Pasteris, N. G.; Trask, B. J.; Sheldon, S.; Gorski, J. L.: Dis- cordant

[54459] phenotype of two overlapping deletions involving the PAX3 gene in

[54460] chromosome 2q35. Hum. Molec. Genet. 2: 953-959, 1993.

[54461] 8811.

[54462] Beeson, D.; Brydson, M.; Betty, M.; Jeremiah, S.; Povey, S.; Vincent,

[54463] A.; Newsom-Davis, J.: Primary structure of the human muscle acetylcholine

[54464] receptor cDNA cloning of the gamma and epsilon subunits. Europ. J.

[54465] Biochem. 215: 229-238, 1993.

[54466] 8812.

[54467] Ohno, K.; Hutchinson, D. O.; Milone, M.; Brengman, J. M.; Bouzat,

[54468] C.; Sine, S. M.; Engel, A. G.: Congenital myasthenic syndrome caused

[54469] by prolonged acetylcholine receptor channel openings due to a mutation

[54470] in the M2 domain of the epsilon subunit. Proc. Nat. Acad. Sci. 92:

[54471] 758-762, 1995.

[54472] 8813.

[54473] Ohno, K.; Quiram, P. A.; Milone, M.; Wang, H.-L.; Harper, M. C.;

[54474] Pruitt, J. N., II; Brengman, J. M.; Pao, L.; Fischbeck, K. H.; Crawford,

[54475] T. O.; Sine, S. M.; Engel, A. G.: Congenital myasthenic syndromes

[54476] due to heteroallelic nonsense/missense mutations in the acetylcholine

[54477] receptor epsilon subunit gene: identification and functional characterization
[54478] of six new mutations. Hum. Molec. Genet. 6: 753-766, 1997.
[54479] 8814.

[54480] Ohno, K.; Wang, H.-L.; Milone, M.; Bren, N.; Brengman, J. M.; Nakano,
[54481] S.; Quiram, P.; Pruitt, J. N.; Sine, S. M.; Engel, A. G.: Congenital
[54482] myasthenic syndrome caused by decreased agonist binding affinity due
[54483] to a mutation in the acetylcholine receptor epsilon subunit. Neuron 17:
[54484] 157-170, 1996.

[54485] 8815.

[54486] Sieb, J. P.; Dorfler, P.; Tzartos, S.; Wewer, U. M.; Ruegg, M.

[54487] A.; Meyer, D.; Baumann, I.; Lindemuth, R.; Jakschik, J.; Ries, F.
[54488] : Congenital myasthenic syndromes in two kinships with end-plate acetylcholine
[54489] receptor and utrophin deficiency. Neurology 50: 54-61, 1998.

[54490] 8816.

[54491] Sieb, J. P.; Kraner, S.; Rauch, M.; Steinlein, O. K.: Immature

[54492] end-plates and utrophin deficiency in congenital myas-
thenic syndrome

[54493] caused by epsilon-AChR subunit truncating mutations.
Hum. Genet. 107:

[54494] 160-164, 2000.

[54495] 8817.

[54496] Witzemann, V.; Schwarz, H.; Koenen, M.; Berberich, C.;
Villarroel,

[54497] A.; Wernig, A.; Brenner, H. R.; Sakmann, B.: Acetylcholine
receptor

[54498] epsilon-subunit deletion causes muscle weakness and at-
rophy in juvenile

[54499] and adult mice. Proc. Nat. Acad. Sci. 93: 13286-13291,
1996.

[54500] 8818.

[54501] Bartels, C. F.; Zelinski, T.; Lockridge, O.: Mutation at
codon

[54502] 322 in the human acetylcholinesterase (ACHE) gene ac-
counts for YT

[54503] blood group polymorphism. Am. J. Hum. Genet. 52: 928-936, 1993.

[54504] 8819.

[54505] Coates, P. M.; Simpson, N. E.: Genetic variation in human erythrocyte

[54506] acetylcholinesterase. Science 175: 1466-1467, 1972.

[54507] 8820.

[54508] Ehrlich, G.; Viegas-Pequignot, E.; Ginzberg, D.; Sindel, L.; Soreq,

[54509] H.; Zakut, H.: Mapping the human acetylcholinesterase gene to chromosome

[54510] 7q22 by fluorescent in situ hybridization coupled with selective PCR

[54511] amplification from a somatic hybrid cell panel and chromosome-sorted

[54512] DNA libraries. Genomics 13: 1192-1197, 1992.

[54513] 8821.

[54514] Feng, G.; Krejci, E.; Molgo, J.; Cunningham, J. M.; Massoulie,

[54515] J.; Sanes, J. R.: Genetic analysis of collagen Q: roles in acetylcholinesterase

[54516] and butyrylcholinesterase assembly and in synaptic structure and function. J.

[54517] Cell Biol. 144: 1349-1360, 1999.

[54518] 8822.

[54519] Getman, D. K.; Eubanks, J. H.; Camp, S.; Evans, G. A.; Taylor,

[54520] P.: The human gene encoding acetylcholinesterase is located on the

[54521] long arm of chromosome 7. Am. J. Hum. Genet. 51: 170-177, 1992.

[54522] 8823.

[54523] Lapidot-Lifson, Y.; Prody, C. A.; Ginzberg, D.; Meytes, D.; Zakut,

[54524] H.; Soreq, H.: Coamplification of human acetylcholinesterase and

[54525] butyrylcholinesterase genes in blood cells: correlation with various

[54526] leukemias and abnormal megakaryocytopoiesis. Proc. Nat. Acad. Sci. 86:

[54527] 4715-4719, 1989.

[54528] 8824.

[54529] Meshorer, E.; Erb, C.; Gazit, R.; Pavlovsky, L.; Kaufer, D.; Friedman,
[54530] A.; Glick, D.; Ben-Arie, N.; Soreq, H.: Alternative splicing and
[54531] neuritic mRNA translocation under long-term neuronal hypersensitivity. Science 295:
[54532] 508-512, 2002.

[54533] 8825.

[54534] Rachinsky, T. L.; Crenshaw, E. B., III; Taylor, P.: Assign-
ment
[54535] of the gene for acetylcholinesterase to distal mouse chro-
mosome 5. Genomics 14:
[54536] 511-514, 1992.

[54537] 8826.

[54538] Rotundo, R. L.; Gomez, A. M.; Fernandez-Valle, C.; Ran-
dall, W.
[54539] R.: Allelic variants of acetylcholinesterase: genetic evi-
dence that
[54540] all acetylcholinesterase forms in avian nerves and muscles are encoded
[54541] by a single gene. Proc. Nat. Acad. Sci. 85: 7805-7809, 1988.

[54542] 8827.

[54543] Shapira, M.; Tur-Kaspa, I.; Bosgraaf, L.; Livni, N.; Grant, A.

[54544] D.; Grisaru, D.; Korner, M.; Ebstein, R. P.; Soreq, H.: A transcription-activating

[54545] polymorphism in the ACHE promoter associated with acute sensitivity

[54546] to anti-acetylcholinesterases. Hum. Molec. Genet. 9: 1273-1281,

[54547] 2000.

[54548] 8828.

[54549] Drayna, D. T.; McLean, J. W.; Wion, K. L.; Trent, J. M.; Drabkin,

[54550] H. A.; Lawn, R. M.: Human apolipoprotein D gene: gene sequence, chromosome

[54551] localization, and homology to the alpha-2mu-globulin superfamily. DNA 6:

[54552] 199-204, 1987.

[54553] 8829.

[54554] Fielding, P. E.; Fielding, C. J.: A cholesteryl ester transfer

[54555] complex in human plasma. Proc. Nat. Acad. Sci. 77: 3327-3330, 1980.

[54556] 8830.

[54557] Graham, C. A.; McGrew, W. C.: Menstrual synchrony in female undergaduates

[54558] living on a coeducational campus. Psychoneuroendocrinology 5: 245-252,

[54559] 1980.

[54560] 8831.

[54561] Kamboh, M. I.; Albers, J. J.; Majumder, P. P.; Ferrell, R. E.:

[54562] Genetic studies of human apolipoproteins. IX. Apolipoprotein D polymorphism

[54563] and its relation to serum lipoprotein lipid levels. Am. J. Hum. Genet. 45:

[54564] 147-154, 1989.

[54565] 8832.

[54566] McClintock, M. K.: Menstrual synchrony and suppression. Nature 229:

[54567] 244-245, 1971.

[54568] 8833.

[54569] McClintock, M. K.: Estrous synchrony and its mediation by airborn

[54570] chemical communication (Rattus norvegicus). Horm. Behav. 10: 264-276,
[54571] 1978.

[54572] 8834.

[54573] Quadagno, D. M.; Shubeita, H. E.; Deck, J.; Francoeur, D.: Influence
[54574] of male social contacts, exercise and all-female living conditions
[54575] on the menstrual cycle. Psychoneuroendocrinology 6: 239-244, 1981.
[54576] 8835.

[54577] Warden, C. H.; Diep, A.; Taylor, B. A.; Lusis, A. J.: Localization
[54578] of the gene for apolipoprotein D on mouse chromosome 16. Genomics 12:
[54579] 851-852, 1992.

[54580] 8836.

[54581] Zeng, C.; Spielman, A. I.; Vowels, B. R.; Leyden, J. J.; Biemann,
[54582] K.; Preti, G.: A human axillary odorant is carried by apolipoprotein

[54583] D. Proc. Nat. Acad. Sci. 93: 6626-6630, 1996.

[54584] 8837.

[54585] Ruttledge, M. H.; Narod, S. A.; Dumanski, J. P.; Parry, D. M.;

[54586] Eldridge, R.; Wertelecki, W.; Parboosingh, J.; Faucher, M.-C.; Lenoir,

[54587] G. M.; Collins, V. P.; Nordenskjold, M.; Rouleau, G. A.: Presymptomatic

[54588] diagnosis for neurofibromatosis 2 with chromosome 22 markers. Neurology 43:

[54589] 1753-1760, 1993.

[54590] 8838.

[54591] Ruttledge, M. H.; Sarrazin, J.; Rangaratnam, S.; Phelan, C. M.;

[54592] Twist, E.; Merel, P.; Delattre, O.; Thomas, G.; Norden-skjold, M.;

[54593] Collins, V. P.; Dumanski, J. P.; Rouleau, G. A.: Evidence for the

[54594] complete inactivation of the NF2 gene in the majority of sporadic

[54595] meningiomas. Nature Genet. 6: 180-184, 1994.

[54596] 8839.

[54597] Sainz, J.; Huynh, D. P.; Figueroa, K.; Ragge, N. K.; Baser, M.

[54598] E.; Pulst, S.-M.: Mutations of the neurofibromatosis type 2 gene

[54599] and lack of the gene product in vestibular schwannomas. Hum. Molec.

[54600] Genet. 3: 885-891, 1994.

[54601] 8840.

[54602] Schulze, K. M. M.; Hanemann, C. O.; Muller, H. W.; Hanen-burg,

[54603] H.: Transduction of wild-type merlin into human schwan-noma cells

[54604] decreases schwannoma cell growth and induces apopto-sis. Hum. Molec.

[54605] Genet. 11: 69-76, 2002.

[54606] 8841.

[54607] Scoles, D. R.; Baser, M. E.; Pulst, S.-M.: A missense muta-tion

[54608] in the neurofibromatosis 2 gene occurs in patients with mild and severe

[54609] phenotypes. Neurology 47: 544-546, 1996.

[54610] 8842.

[54611] Scoles, D. R.; Huynh, D. P.; Chen, M. S.; Burke, S. P.; Gut-
mann,
[54612] D. H.; Pulst, S.-M.: The neurofibromatosis 2 tumor sup-
pressor protein
[54613] interacts with hepatocyte growth factor-regulated tyrosine
kinase
[54614] substrate. Hum. Molec. Genet. 9: 1567-1574, 2000.

[54615] 8843.

[54616] Seizinger, B. R.; Martuza, R. L.; Gusella, J. F.: Loss of genes

[54617] on chromosome 22 in tumorigenesis of human acoustic
neuroma. Nature 322:
[54618] 644-647, 1986.

[54619] 8844.

[54620] Seizinger, B. R.; Rouleau, G.; Ozelius, L. J.; Lane, A. H.; St.

[54621] George-Hyslop, P.; Huson, S.; Gusella, J. F.; Martuza, R. L.:
Common
[54622] pathogenetic mechanism for three tumor types in bilateral
acoustic

[54623] neurofibromatosis. Science 236: 317-319, 1987.

[54624] 8845.

[54625] Sestini, R.; Vivarelli, R.; Balestri, P.; Ammannati, F.; Mon-
tali,
[54626] E.; Papi, L.: Neurofibromatosis type 2 attributable to
gonosomal
[54627] mosaicism in a clinically normal mother, and identification
of seven
[54628] novel mutations in the NF2 gene. Hum. Genet. 107:
366-371, 2000.

[54629] 8846.

[54630] Siggers, D. C.; Boyer, S. H.; Eldridge, R.: Nerve-growth
factor
[54631] in disseminated neurofibromatosis. (Letter) New Eng. J.
Med. 292:
[54632] 1134, 1975.

[54633] 8847.

[54634] Stokowski, R. P.; Cox, D. R.: Functional analysis of the
neurofibromatosis
[54635] type 2 protein by means of disease-causing point muta-
tions. Am. J.

[54636] Hum. Genet. 66: 873-891, 2000.

[54637] 8848.

[54638] Thomas, G.: Personal Communication. Paris, France 11/17/1993.

[54639] 8849.

[54640] Trofatter, J. A.; MacCollin, M. M.; Rutter, J. L.; Murrell, J.

[54641] R.; Duyao, M. P.; Parry, D. M.; Eldridge, R.; Kley, N.; Menon, A.

[54642] G.; Pulaski, K.; Haase, V. H.; Ambrose, C. M.; Munroe, D.; Bove, C.;

[54643] Haines, J. L.; Martuza, R. L.; MacDonald, M. E.; Seizinger, B. R.;

[54644] Short, M. P.; Buckler, A. J.; Gusella, J. F.: A novel moesin-, ezrin-,

[54645] radixin-like gene is a candidate for the neurofibromatosis 2 tumor

[54646] suppressor. Cell 72: 791-800, 1993.

[54647] 8850.

[54648] Wellenreuther, R.; Kraus, J. A.; Lenartz, D.; Menon, A. G.; Schramm,

[54649] J.; Louis, D. N.; Ramesh, V.; Gusella, J. F.; Wiestler, O. D.;

von

[54650] Deimling, A.: Analysis of the neurofibromatosis 2 gene reveals molecular

[54651] variants of meningioma. Am. J. Path. 146: 827-832, 1995.

[54652] 8851.

[54653] Wertelecki, W.; Rouleau, G. A.; Superneau, D. W.; Forehand, L.

[54654] W.; Williams, J. P.; Haines, J. L.; Gusella, J. F.: Neurofibromatosis

[54655] 2: clinical and DNA linkage studies of a large kindred. New Eng.

[54656] J. Med. 319: 278-283, 1988.

[54657] 8852.

[54658] Wishart, J. H.: Case of tumours in the skull, dura mater, and

[54659] brain. Edinburgh Med. Surg. J. 18: 393-397, 1822.

[54660] 8853.

[54661] Wolff, R. K.; Frazer, K. A.; Jackler, R. K.; Lanser, M. J.; Pitts,

[54662] L. H.; Cox, D. R.: Analysis of chromosome 22 deletions in neurofibromatosis

[54663] type 2-related tumors. Am. J. Hum. Genet. 51: 478-485,

1992.

[54664] 8854.

[54665] Worster-Drought, C.; Dickson, W. E. C.; McMenemey, W. H.: Multiple

[54666] meningeal and perineural tumors with analogous changes in the glia

[54667] and ependyma (neurofibroblastomatosis). Brain 60: 85-117, 1937.

[54668] 8855.

[54669] Wu, C. L.; Thakker, N.; Neary, W.; Black, G.; Lye, R.; Ramsden,

[54670] R. T.; Read, A. P.; Evans, D. G. R.: Differential diagnosis of type

[54671] 2 neurofibromatosis: molecular discrimination of NF2 and sporadic

[54672] vestibular schwannomas. J. Med. Genet. 35: 973-977, 1998.

[54673] 8856.

[54674] Young, D. F.; Eldridge, R.; Gardner, W. J.: Bilateral acoustic

[54675] neuroma in a large kindred. J.A.M.A. 214: 347-353, 1970.

[54676] 8857.

[54677] Zucman-Rossi, J.; Legoix, P.; Der Sarkissian, H.; Cheret, G.;
[54678] Sor, F.; Bernardi, A.; Cazes, L.; Giraud, S.; Ollagnon, E.; Lenoir,
[54679] G.; Thomas, G.: NF2 gene in neurofibromatosis type 2 patients. Hum.
[54680] Molec. Genet. 7: 2095-2101, 1998.

[54681] 8858.

[54682] Gatalica, B.; Pulkkinen, L.; Li, K.; Kuokkanen, K.; Ryynanen, M.;
[54683] McGrath, J. A.; Uitto, J.: Cloning of the human type XVII collagen
[54684] gene (COL17A1), and detection of novel mutations in generalized atrophic
[54685] benign epidermolysis bullosa. Am. J. Hum. Genet. 60: 352-365, 1997.
[54686] 8859.

[54687] Wang, H.; Iakova, P.; Wilde, M.; Welm, A.; Goode, T.; Roesler,
[54688] W. J.; Timchenko, N. A.: C/EBP-alpha arrests cell proliferation through
[54689] direct inhibition of Cdk2 and Cdk4. Molec. Cell 8:

817-828, 2001.

[54690] 8860.

[54691] Wang, N.; Finegold, M. J.; Bradley, A.; Ou, C. N.; Abdel-sayed,

[54692] S. V.; Wilde, M. D.; Taylor, L. R.; Wilson, D. R.; Darlington, G.

[54693] J.: Impaired energy homeostasis in C/EBP-alpha knockout mice. Science 269:

[54694] 1108-1112, 1995.

[54695] 8861.

[54696] Wu, Z.; Rosen, E. D.; Brun, R.; Hauser, S.; Adelmant, G.; Troy,

[54697] A. E.; McKeon, C.; Darlington, G. J.; Spiegelman, B. M.: Cross-regulation

[54698] of C/EBP-alpha and PPAR-gamma controls the transcrip-tional pathway

[54699] of adipogenesis and insulin sensitivity. Molec. Cell 3: 151-158,

[54700] 1999.

[54701] 8862.

[54702] Cleutjens, C. B. J. M.; van Eekelen, C. C. E. M.; van Dekken,

H.;

[54703] Smit, E. M. E.; Hagemeijer, A.; Wagner, M. J.; Wells, D. E.; Trapman,

[54704] J.: The human C/EBP-delta (CRP3/CELF) gene: structure and chromosomal

[54705] localization. Genomics 16: 520-523, 1993.

[54706] 8863.

[54707] Williams, S. C.; Cantwell, C. A.; Johnson, P. F.: A family of

[54708] C/EBP-related proteins capable of forming covalently linked leucine

[54709] zipper dimers in vitro. Genes Dev. 5: 1553-1567, 1991.

[54710] 8864.

[54711] Al-Awqati, Q.; Preisig, P. A.: Size does matter: will knock-out

[54712] of p21(WAF1/CIP1) save the kidney by limiting compen-satory renal growth?

[54713] (Commentary) Proc. Nat. Acad. Sci. 96: 10551-10553, 1999.

[54714] 8865.

[54715] Chedid, M.; Michieli, P.; Lengel, C.; Huppi, K.; Givol, D.: A

[54716] single nucleotide substitution at codon 31 (ser/arg) de-

fines a polymorphism

[54717] in a highly conserved region of the p53-inducible gene WAF1/CIP1. Oncogene 9:

[54718] 3021-3024, 1994.

[54719] 8866.

[54720] Cheng, T.; Rodrigues, N.; Shen, H.; Yang, Y.; Dombkowski, D.; Sykes,

[54721] M.; Scadden, D. T.: Hematopoietic stem cell quiescence maintained

[54722] by p21(cip1/waf1). Science 287: 1804-1808, 2000.

[54723] 8867.

[54724] Harper, J. W.; Adami, G. R.; Wei, N.; Keyomarsi, K.; Elledge, S.

[54725] J.: The p21 Cdk-interacting protein Cip1 is a potent inhibitor of

[54726] G1 cyclin-dependent kinases. Cell 75: 805-816, 1993.

[54727] 8868.

[54728] Huppi, K.; Siwarski, D.; Dosik, J.; Michieli, P.; Chedid, M.; Reed,

[54729] S.; Mock, B.; Givol, D.; Mushinski, J. F.: Molecular cloning, sequencing,

[54730] chromosomal localization and expression of mouse p21 (Waf1). Oncogene 9:
[54731] 3017-3020, 1994.

[54732] 8869.

[54733] Megyesi, J.; Price, P. M.; Tamayo, E.; Safirstein, R. L.: The

[54734] lack of a functional p21(WAF1/CIP1) gene ameliorates progression to
[54735] chronic renal failure. Proc. Nat. Acad. Sci. 96: 10830-10835, 1999.
[54736] 8870.

[54737] Mousses, S.; Ozcelik, H.; Lee, P. D.; Malkin, D.; Bull, S. B.;

[54738] Andrulis, I. L.: Two variants of the CIP1/WAF1 gene occur together
[54739] and are associated with human cancer. Hum. Molec. Genet. 4: 1089-1092,
[54740] 1995.

[54741] 8871.

[54742] Heintz, N.; Zoghbi, H.: Alpha-synuclein--a link between Parkinson
[54743] and Alzheimer diseases? Nature Genet. 16: 325-327, 1997.

[54744] 8872.

[54745] Iwata, N.; Tsubuki, S.; Takaki, Y.; Shirotani, K.; Lu, B.; Gerard,

[54746] N. P.; Gerard, C.; Hama, E.; Lee, H.-J.; Saido, T. C.: Metabolic

[54747] regulation of brain A-beta by neprilysin. Science 292: 1550-1552,

[54748] 2001.

[54749] 8873.

[54750] Kang, J.; Lemaire, H.-G.; Unterbeck, A.; Salbaum, J. M.; Masters,

[54751] C. L.; Grzeschik, K.-H.; Multhaup, G.; Beyreuther, K.; Muller-Hill,

[54752] B.: The precursor of Alzheimer's disease amyloid A4 protein resembles

[54753] a cell-surface receptor. Nature 325: 733-736, 1987.

[54754] 8874.

[54755] Lu, P.-J.; Wulf, G.; Zhou, X. Z.; Davies, P.; Lu, K. P.: The

[54756] prolyl isomerase Pin1 restores the function of Alzheimer-associated

[54757] phosphorylated tau protein. Nature 399: 784-788, 1999.

[54758] 8875.

[54759] Helmerhorst, E. J.; van't Hof, W.; Breeuwer, P.; Veerman, E. C.

[54760] I.; Abee, T.; Troxler, R. F.; Nieuw Amerongen, A. V.; Oppenheim, F.

[54761] G.: Characterization of histatin 5 with respect to amphipathicity,

[54762] hydrophobicity, and effects on cell and mitochondrial membrane integrity

[54763] excludes a candidacidal mechanism of pore formation. J. Biol. Chem. 276:

[54764] 5643-5649, 2001.

[54765] 8876.

[54766] Sabatini, L. M.; Azen, E. A.: Two coding change mutations in the

[54767] HIS2(2) allele characterize the salivary histatin 3-2 protein variant. Hum.

[54768] Mutat. 4: 12-19, 1994.

[54769] 8877.

[54770] Troxler, R. F.; Offner, G. D.; Xu, T.; Vanderspek, J. C.; Oppenheim,

[54771] F. G.: Structural relationship between human salivary histatins. J.

[54772] Dent. Res. 69: 2-6, 1990.

[54773] 8878.

[54774] vanderSpek, J. C.; Wyandt, H. E.; Skare, J. C.; Milunsky, A.;

[54775] Oppenheim, F. G.; Troxler, R. F.: Localization of the genes for histatins

[54776] to human chromosome 4q13 and tissue distribution of the mRNAs. Am.

[54777] J. Hum. Genet. 45: 381-387, 1989.

[54778] 8879.

[54779] Stamey, T. A.; Yang, N.; Hay, A. R.; McNeal, J. E.; Freiha, F.

[54780] S.; Redwine, E.: Prostate-specific antigen as a serum marker for

[54781] adenocarcinoma of the prostate. New Eng. J. Med. 317: 909-916, 1987.

[54782] 8880.

[54783] Seldin, M. F.: Personal Communication. Durham, N. C. 9/19/1990.

[54784] 8881.

[54785] Sherman, L.; Wainwright, D.; Ponta, H.; Herrlich, P.: A splice
[54786] variant of CD44 expressed in the apical ectodermal ridge presents
[54787] fibroblast growth factors to limb mesenchyme and is required for limb
[54788] outgrowth. Genes Dev. 12: 1058-1071, 1998.

[54789] 8882.

[54790] Stefanova, I.; Hilgert, I.; Bazil, V.; Kristofova, H.; Horejsi,

[54791] V.: Human leucocyte surface glycoprotein CDw44 and lymphocyte homing
[54792] receptor are identical molecules. Immunogenetics 29: 402-404, 1989.

[54793] 8883.

[54794] Teder, P.; Vandivier, R. W.; Jiang, D.; Liang, J.; Cohn, L.; Pure,
[54795] E.; Henson, P. M.; Noble, P. W.: Resolution of lung inflammation
[54796] by CD44. Science 296: 155-158, 2002.

[54797] 8884.

[54798] Telen, M. J.: Personal Communication. Durham, N. C.

12/30/1992.

[54799] 8885.

[54800] Telen, M. J.; Eisenbarth, G. S.; Haynes, B. F.: Human erythrocyte

[54801] antigens: regulation of expression of a novel erythrocyte surface

[54802] antigen by the inhibitor Lutheran In(Lu) gene. J. Clin. Invest. 71:

[54803] 1878-1886, 1983.

[54804] 8886.

[54805] Mueller, T. J.; Morrison, M.: Detection of a variant of protein

[54806] 3, the major transmembrane protein of the human erythrocyte. J. Biol.

[54807] Chem. 252: 6573-6576, 1977.

[54808] 8887.

[54809] Mueller, T. J.; Morrison, M.: Heterogeneity of the major protein

[54810] in the human erythrocyte membrane. (Abstract) Fed. Proc. 36: 747,

[54811] 1977.

[54812] 8888.

[54813] Palatnik, M.; da Silva Simoes, M. L. M.; Alves, Z. M. S.; Laranjeira,

[54814] N. S. M.: The 60 and 63 kDa proteolytic peptides of the red cell

[54815] membrane band-3 protein: their prevalence in human and non-human primates. Hum.

[54816] Genet. 86: 126-130, 1990.

[54817] 8889.

[54818] Palumbo, A. P.; Isobe, M.; Huebner, K.; Shane, S.; Rovera, G.;

[54819] Demuth, D.; Curtis, P. J.; Ballantine, M.; Croce, C. M.; Showe, L.

[54820] C.: Chromosomal localization of a human band 3-like gene to region

[54821] 7q35-7q36. Am. J. Hum. Genet. 39: 307-316, 1986.

[54822] 8890.

[54823] Pawloski, J. R.; Hess, D. T.; Stamler, J. S.: Export by red blood

[54824] cells of nitric oxide bioactivity. Nature 409: 622-626, 2001.

[54825] 8891.

[54826] Peters, L. L.; Shivdasani, R. A.; Liu, S.-C.; Hanspal, M.; John,

[54827] K. M.; Gonzalez, J. M.; Brugnara, C.; Gwynn, B.; Mohandas, N.; Alper,

[54828] S. L.; Orkin, S. H.; Lux, S. E.: Anion exchanger 1 (band 3) is required

[54829] to prevent erythrocyte membrane surface loss but not to form the membrane

[54830] skeleton. Cell 86: 917-927, 1996.

[54831] 8892.

[54832] Prchal, J. T.; Guan, Y.; Jarolim, P.; Palek, J.; Showe, L.; Bertoli,

[54833] L.: Hereditary spherocytosis in a large family is linked with the

[54834] band 3 gene and not with alpha-spectrin, beta-spectrin or ankyrin.

[54835] (Abstract) Blood 78 (suppl.): 81a, 1991.

[54836] 8893.

[54837] Ranney, H. M.; Rosenberg, G. H.; Morrison, M.; Mueller, T. J.

[54838] : Frequencies of band 3 variants of human red cell membranes in some
[54839] different populations. Brit. J. Haemat. 76: 262-267, 1990.

[54840] 8894.

[54841] Ravindranath, Y.; Goyette, G., Jr.; Johnson, R. M.: South-east
[54842] Asian ovalocytosis in an African-American family. (Letter) Blood 84:
[54843] 2823-2824, 1994.

[54844] 8895.

[54845] Reinhart, W. H.; Wyss, E. J.; Arnold, D.; Ott, P.: Hereditary
[54846] spherocytosis associated with protein band 3 defect in a Swiss kindred. Brit.
[54847] J. Haemat. 86: 147-155, 1994.

[54848] 8896.

[54849] Ribeiro, M. L.; Alloisio, N.; Almeida, H.; Gomes, C.; Texier,
[54850] P.; Lemos, C.; Mimoso, G.; Morle, L.; Bey-Cabet, F.; Rudigoz, R.-C.;
[54851] Delaunay, J.; Tamagnini, G.: Severe hereditary spherocytosis and

[54852] distal renal tubular acidosis associated with the total absence of

[54853] band 3. Blood 96: 1602-1604, 2000.

[54854] 8897.

[54855] Rybicki, A. C.; Qiu, J. J. H.; Musto, S.; Rosen, N. L.; Nagel,

[54856] R. L.; Schwartz, R. S.: Human erythrocyte protein 4.2 deficiency

[54857] associated with hemolytic anemia and a homozygous 40 glutamic acid-to-lysine

[54858] substitution in the cytoplasmic domain of band 3 (band 3 Montefiore). Blood 81:

[54859] 2155-2165, 1993.

[54860] 8898.

[54861] Saad, S. T. O.; Liu, S. C.; Golan, D.; Corbett, J. B.; Thatte,

[54862] H. S.; Derick, L.; Hanspal, M.; Jarolim, P.; Fibach, E.; Palek, J.

[54863] : Mechanism underlying band 3 deficiency in a subset of patients with

[54864] hereditary spherocytosis (HS). (Abstract) Blood 78 (suppl.): 81a,

[54865] 1991.

[54866] 8899.

[54867] Sahr, K. E.; Taylor, W. M.; Daniels, B. P.; Rubin, H. L.; Jarolim,

[54868] P.: The structure and organization of the human erythroid anion exchanger

[54869] (AE1) gene. Genomics 24: 491-501, 1994.

[54870] 8900.

[54871] Schofield, A. E.; Martin, P. G.; Spillett, D.; Tanner, M. J. A.

[54872] : The structure of the human red blood cell anion exchanger (EPB3,

[54873] AE1, Band 3) gene. Blood 84: 2000-2012, 1994.

[54874] 8901.

[54875] Schulman, S.; Roth, E. F., Jr.; Cheng, B.; Rybicki, A. C.; Sussman,

[54876] I. I.; Wong, M.; Wang, W.; Ranney, H. M.; Nagel, R. L.; Schwartz,

[54877] R. S.: Growth of Plasmodium falciparum in human erythrocytes containing

[54878] abnormal membrane proteins. Proc. Nat. Acad. Sci. 87: 7339-7343,

[54879] 1990.

[54880] 8902.

[54881] Serjeantson, S.; Bryson, K.; Amato, D.; Babona, D.: Malaria and

[54882] hereditary ovalocytosis. Hum. Genet. 37: 161-167, 1977.

[54883] 8903.

[54884] Showe, L. C.; Ballantine, M.; Huebner, K.: Localization of the

[54885] gene for the erythroid anion exchange protein, band 3 (EMPB3), to

[54886] human chromosome 17. Genomics 1: 71-76, 1987.

[54887] 8904.

[54888] Southgate, C. D.; Chisti, A. H.; Mitchell, B.; Yi, S. J.; Palek,

[54889] J.: Targeted disruption of the murine erythroid band 3 gene results

[54890] in spherocytosis and severe haemolytic anaemia despite a normal membrane

[54891] skeleton. Nature Genet. 14: 227-230, 1996.

[54892] 8905.

[54893] Stewart, E. A.; Kopito, R.; Bowcock, A. M.: A PstI polymor-
phism

[54894] for the human erythrocyte surface protein band 3 (EPB3) demonstrates

[54895] close linkage of EPB3 to the nerve growth factor receptor. Genomics 5:

[54896] 633-635, 1989.

[54897] 8906.

[54898] Tanner, M. J. A.: Molecular and cellular biology of the erythrocyte

[54899] anion exchanger (AE1). Seminars Hemat. 30: 34-57, 1993.

[54900] 8907.

[54901] Tanner, M. J. A.; Bruce, L.; Martin, P. G.; Rearden, D. M.; Jones,

[54902] G. L.: Melanesian hereditary ovalocytes have a deletion in red cell

[54903] band 3. (Letter) Blood 78: 2785-2786, 1991.

[54904] 8908.

[54905] Tanphaichitr, V. S.; Sumboonnanonda, A.; Ideguchi, H.; Shayakul,

[54906] C.; Brugnara, C.; Takao, M.; Veerakul, G.; Alper, S. L.: Novel AE1

[54907] mutations in recessive distal renal tubular acidosis: loss-

of-function
[54908] is rescued by glycophorin A. J. Clin. Invest. 102: 2173-2179, 1998.
[54909] 8909.
[54910] Toye, A. M.; Bruce, L. J.; Unwin, R. J.; Wrong, O.; Tanner, M.
[54911] J. A.: Band 3 Walton, a C-terminal deletion associated with distal
[54912] renal tubular acidosis, is expressed in the red cell membrane but
[54913] retained internally in kidney cells. Blood 99: 342-347, 2002.
[54914] 8910.
[54915] Wrong, O. M.; Feest, T. G.; MacIver, A. G.: Immune-related potassium-losing
[54916] interstitial nephritis: a comparison with distal renal tubular acidosis. Quart.
[54917] J. Med. 86: 513-534, 1993.
[54918] 8911.
[54919] Yannoukakos, D.; Vasseur, C.; Driancourt, C.; Blouquit, Y.; Delaunay,
[54920] J.; Wajcman, H.; Bursaux, E.: Human erythrocyte band 3

polymorphism
[54921] (band 3 Memphis): characterization of the structural modification
[54922] (lys56-to-glu) by protein chemistry methods. Blood 78: 1117-1120,
[54923] 1991.

[54924] 8912.

[54925] Mallinson, G.; Soo, K. S.; Schall, T. J.; Pisacka, M.; Anstee,

[54926] D. J.: Mutations in the erythrocyte chemokine receptor (Duffy) gene:
[54927] the molecular basis of the Fy(a)/Fy(b) antigens and identification
[54928] of a deletion in the Duffy gene of an apparently healthy individual
[54929] with the Fy(a-b-) phenotype. Brit. J. Haemat. 90: 823-829, 1995.
[54930] 8913.

[54931] Miller, L. H.; Mason, S. J.; Clyde, D. F.; McGinnis, M. H.: The
[54932] resistance factor to Plasmodium vivax in blacks: the Duffy blood group
[54933] genotype, FyFy. New Eng. J. Med. 295: 302-304, 1976.

[54934] 8914.

[54935] Miller, L. H.; Mason, S. J.; Dvorak, J. A.: Erythrocyte recep-
tors
[54936] of Plasmodium knowlesi malaria: Duffy blood group de-
terminants. Science 189:
[54937] 561-562, 1975.

[54938] 8915.

[54939] Nance, W. E.; Conneally, M.; Kang, K. W.; Reed, T. E.;
Schroder,
[54940] J.; Rose, S.: Genetic linkage analysis of human hemoglobin
variants. Am.
[54941] J. Hum. Genet. 22: 453-459, 1970.

[54942] 8916.

[54943] Nichols, M. E.; Rubinstein, P.; Barnwell, J.; Rodriguez de
Cordoba,
[54944] S.; Rosenfield, R. E.: A new human Duffy blood group
specificity
[54945] defined by a murine monoclonal antibody: immunogenet-
ics and association
[54946] with susceptibility to Plasmodium vivax. J. Exp. Med. 166:
776-785,

[54947] 1987.

[54948] 8917.

[54949] Olsson, M. L.; Smythe, J. S.; Hansson, C.; Poole, J.; Mallinson,

[54950] G.; Jones, J.; Avent, N. D.; Daniels, G.: The Fy(x) phenotype is

[54951] associated with a missense mutation in the Fy(b) allele predicting

[54952] Arg89Cys in the Duffy glycoprotein. Brit. J. Haemat. 103: 1184-1191,

[54953] 1998.

[54954] 8918.

[54955] Palmer, C. G.; Christian, J. C.; Merritt, A. D.: Partial trisomy

[54956] 1 due to a 'shift' and probable location of the Duffy (Fy) locus. Am.

[54957] J. Hum. Genet. 29: 371-377, 1977.

[54958] 8919.

[54959] Parasol, N.; Reid, M.; Rios, M.; Castilho, L.; Harari, I.; Kosower,

[54960] N. S.: A novel mutation in the coding sequence of the FY*B allele

[54961] of the Duffy chemokine receptor gene is associated with an altered
[54962] erythrocyte phenotype. Blood 92: 2237-2243, 1998.

[54963] 8920.

[54964] Pasvol, G.; Wilson, R. J. M.: The interaction of malaria parasites
[54965] with red blood cells. Brit. Med. Bull. 38: 133-140, 1982.

[54966] 8921.

[54967] Peiper, S. C.; Wang, Z.; Neote, K.; Martin, A. W.; Showell, H.
[54968] J.; Conklyn, M. J.; Ogborne, K.; Hadley, T. J.; Lu, Z.; Hesselgesser,
[54969] J.; Horuk, R.: The Duffy antigen/receptor for chemokines (DARC) is
[54970] expressed in endothelial cells of Duffy negative individuals who lack
[54971] the erythrocyte receptor. J. Exp. Med. 181: 1311-1317, 1995.

[54972] 8922.

[54973] Ritter, H.: Zur formalen Genetik des Duffy-systems. Untersuchung

[54974] von 247 Familien. Humangenetik 4: 59-61, 1967.

[54975] 8923.

[54976] Robson, E. B.; Cook, P. J. L.; Corney, G.; Hopkinson, D. A.; Noades,
[54977] J.; Cleghorn, T. E.: Linkage data on Rh, PGM, PGD, peptidase C and
[54978] Fy from family studies. Ann. Hum. Genet. 36: 393-399, 1973.

[54979] 8924.

[54980] Szabo, M. C.; Soo, K. S.; Zlotnik, A.; Schall, T. J.: Chemokine
[54981] class differences in binding to the Duffy antigen-erythrocyte chemokine
[54982] receptor. J. Biol. Chem. 270: 25348-25351, 1995.

[54983] 8925.

[54984] Tamasauskas, D.; Powell, V.; Saksela, K.; Yazdanbakhsh, K.: A
[54985] homologous naturally occurring mutation in Duffy and CCR5 leading
[54986] to reduced receptor expression. Blood 97: 3651-3654, 2001.

[54987] 8926.

[54988] Tang, T.; Owen, J. D.; Du, J.; Walker, C. L.; Richmond, A.: Molecular

[54989] cloning and characterization of a mouse gene with homology to the

[54990] Duffy-antigen receptor for chemokines. DNA Seq. 9: 129-143, 1999.

[54991] 8927.

[54992] Tournamille, C.; Kim Le Van, C.; Gane, P.; Le Pennec, P. Y.; Roubinet,

[54993] F.; Babinet, J.; Cartron, J. P.; Colin, Y.: Arg89Cys substitution

[54994] results in very low membrane expression of the Duffy antigen/receptor

[54995] for chemokines in Fy(x) individuals. Blood 92: 2147-2156, 1998.

[54996] Note: Erratum. Blood 95: 2753 only, 2000.

[54997] 8928.

[54998] Tournamille, C.; Le Van Kim, C.; Gane, P.; Cartron, J.-P.; Colin,

[54999] Y.: Molecular basis and PCR-DNA typing of the Fya/fyb blood group

[55000] polymorphism. Hum. Genet. 95: 407-410, 1995.

[55001] 8929.

[55002] Weitkamp, L. R.: Personal Communication. Rochester, N. Y.

[55003] 1972.

[55004] 8930.

[55005] Zimmerman, P. A.; Woolley, I.; Masinde, G. L.; Miller, S. M.;

[55006] McNamara, D. T.; Hazlett, F.; Mgone, C. S.; Alpers, M. P.; Genton,

[55007] B.; Boatin, B. A.; Kazura, J. W.: Emergence of FY*A(null) in a Plasmodium

[55008] vivax-endemic region of Papua New Guinea. Proc. Nat. Acad. Sci. 96:

[55009] 13973-13977, 1999.

[55010] 8931.

[55011] Anderson, S. E.; McKenzie, J. L.; McLoughlin, K.; Beard, M. E.

[55012] J.; Hart, D. N. J.: The inheritance of abnormal sialoglyco-proteins

[55013] found in a Gerbich negative individual. Pathology 18: 407-412, 1986.

[55014] 8932.

[55015] Anstee, D. J.; Parsons, S. F.; Ridgwell, K.; Tanner, M. J. A.;

[55016] Merry, A. H.; Thomson, E. E.; Judson, P. A.; Johnson, P.;
Bates, S.;

[55017] Fraser, I. D.: Two individuals with elliptocytic red cells ap-
parently

[55018] lack three minor erythrocyte membrane sialoglycopro-
teins. Biochem.

[55019] J. 218: 615-619, 1984.

[55020] 8933.

[55021] Barnes, R.; Lewis, T. L. T.: A rare antibody (anti-Ge) caus-
ing

[55022] hemolytic disease of the newborn. Lancet II: 1285-1286,
1961.

[55023] 8934.

[55024] Bennett, V.: The membrane skeleton of human erythro-
cytes and its

[55025] implications for more complex cells. Annu. Rev. Biochem.
54: 273-304,

[55026] 1985.

[55027] 8935.

[55028] Bloomfield, L.; Rowe, G. P.; Green, C.: The Webb (Wb) antigen
[55029] in South Wales donors. Hum. Hered. 36: 352-356, 1986.

[55030] 8936.

[55031] Booth, P. B.; McLoughlin, K.: The Gerbich blood group system,
[55032] especially in Melanesians. Vox Sang. 22: 73-84, 1972.

[55033] 8937.

[55034] Cartron, J.-P.; Colin, Y.; Kudo, S.; Fukuda, M.: Molecular genetics
[55035] of human erythrocyte sialoglycoproteins A, B, C, and D.In: Harris,
[55036] J. R.: Erythroid Cells. Blood Cell Biochemistry. New York: Plenum
[55037] Press (pub.) 1: 1990. Pp. 299-335.

[55038] 8938.

[55039] Chang, S.; Reid, M. E.; Conboy, J.; Kan, Y. W.; Mohandas, N.:
[55040] Molecular characterization of erythrocyte glycophorin C variants. Blood 77:
[55041] 644-648, 1991.

[55042] 8939.

[55043] Tommerup, N.; Leffers, H.: Assignment of the human genes encoding

[55044] 14-3-3 eta (YWHAH) to 22q12, 14-3-3 zeta (YWHAZ) to 2p25.1-p25.2,

[55045] and 14-3-3 beta (YWHAB) to 20q13.1 by in situ hybridization. Genomics 33:

[55046] 149-150, 1996.

[55047] 8940.

[55048] Yaffe, M. B.; Rittinger, K.; Volinia, S.; Caron, P. R.; Aitken,

[55049] A.; Leffers, H.; Gamblin, S. J.; Smerdon, S. J.; Cantley, L. C.:

[55050] The structural basis for 14-3-3:phosphopeptide binding specificity. Cell 91:

[55051] 961-971, 1997.

[55052] 8941.

[55053] Ben-Yosef, T.; Eden, A.; Benvenisty, N.: Characterization of murine

[55054] BCAT genes: Bcat1, a c-Myc target, and its homolog, Bcat2. Mammalian

[55055] Genome 9: 595-597, 1998.

[55056] 8942.

[55057] Benvenisty, N.; Leder, A.; Kuo, A.; Leder, P.: An embryonically

[55058] expressed gene is a target for c-Myc regulation via the c-Myc-binding

[55059] sequence. Genes Dev. 6: 2513-2523, 1992.

[55060] 8943.

[55061] Eden, A.; Simchen, G.; Benvenisty, N.: Two yeast homologs of ECA39,

[55062] a target for c-Myc regulation, code for cytosolic and mitochondrial

[55063] branched-chain amino acid aminotransferases. J. Biol. Chem. 271:

[55064] 20242-20245, 1996.

[55065] 8944.

[55066] Jones, C.; Moore, E. E.: Isolation of mutants lacking branched-chain

[55067] amino acid transaminase. Somat. Cell Genet. 2: 235-243, 1976.

[55068] 8945.

[55069] Jones, C.; Moore, E. E.: Assignment of the human gene

complementing

[55070] the auxotrophic marker TRANS-minus (BCT1) to chromosome 12. (Abstract) Cytogenet.

[55071] Cell Genet. 25: 168 only, 1979.

[55072] 8946.

[55073] Jones, C.; Moore, E. E.: Localization of a gene which complements

[55074] branched-chain amino acid transaminase deficiency to the short arm

[55075] of human chromosome 12. Hum. Genet. 66: 206-211, 1984.

[55076] 8947.

[55077] Naylor, S. L.; Shows, T. B.: Branched-chain aminotransferase genes

[55078] (BCT-1 and BCT-2) assigned to human chromosomes 12 and 19 using alpha-keto

[55079] acid selection media. (Abstract) Cytogenet. Cell Genet. 25: 191-192,

[55080] 1979.

[55081] 8948.

[55082] Naylor, S. L.; Shows, T. B.: Branched-chain aminotrans-

ferase deficiency
[55083] in Chinese hamster cells complemented by two independent genes on
[55084] human chromosomes 12 and 19. Somat. Cell Genet. 6: 641-652, 1980.
[55085] 8949.

[55086] Schuldiner, O.; Eden, A.; Ben-Yosef, T.; Yanuka, O.; Simchen, G.;
[55087] Benvenisty, N.: ECA39, a conserved gene regulated by c-Myc in mice,
[55088] is involved in G1/S cell cycle regulation in yeast. Proc. Nat. Acad.
[55089] Sci. 93: 7143-7148, 1996.

[55090] 8950.

[55091] Tanaka, K.; Rosenberg, L. E.: Disorders of branched chain amino
[55092] acid and organic acid metabolism.In: Stanbury, J. B.; Wyngaarden,
[55093] J. B.; Fredrickson, D. S.; Goldstein, J. L.; Brown, M. S.: The Metabolic
[55094] Basis of Inherited Disease. New York: McGraw-Hill (pub.) (5th

[55095] ed.): 1983. Pp. 450-451.

[55096] 8951.

[55097] Bassett, D. E., Jr.; Boguski, M. S.; Spencer, F.; Reeves, R.; Kim,

[55098] S.; Weaver, T.; Hieter, P.: Genome cross-referencing and XREFdb:

[55099] implications for the identification and analysis of genes mutated

[55100] in human disease. Nature Genet. 15: 339-344, 1997.

[55101] 8952.

[55102] Bijlsma, E. K.; Merel, P.; Bosch, D. A.; Westerveld, A.; De-lattre,

[55103] O.; Thomas, G.; Hulsebos, T. J. M.: Analysis of mutations in the

[55104] SCH gene in schwannomas. Genes Chromosomes Cancer 11: 7-14, 1994.

[55105] 8953.

[55106] Nagase, S.; Shimamune, K.; Shumiya, S.: Albumin-defi-cient rat

[55107] mutant. Science 205: 590-591, 1979.

[55108] 8954.

[55109] Cattanach, B. M.; Barr, J. A.; Evans, E. P.; Burtenshaw, M.; Beechey,
[55110] C. V.; Leff, S. E.; Brannan, C. I.; Copeland, N. G.; Jenkins, N. A.;
[55111] Jones, J.: A candidate mouse model for Prader-Willi syndrome which
[55112] shows an absence of Snrpn expression. Nature Genet. 2: 270-274,
[55113] 1992.

[55114] 8955.

[55115] Telen, M. J.; Palker, T. J.; Haynes, B. F.: Human erythrocyte
[55116] antigens: II. The In(Lu) gene regulates expression of an antigen on
[55117] an 80-kilodalton protein of human erythrocytes. Blood 64: 599-606,
[55118] 1984.

[55119] 8956.

[55120] Colin, Y.; Rahuel, C.; London, J.; Romeo, P. H.; d'Auriol, L.;
[55121] Galibert, F.; Cartron, J.-P.: Isolation of cDNA clones and complete
[55122] amino acid sequence of human erythrocyte glycophorin C.

J. Biol.

[55123] Chem. 261: 229-233, 1986.

[55124] 8957.

[55125] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Collagen

[55126] alpha-1 chain . Washington: National Biomedical Research Foundation

[55127] (pub.) 5: 1972. Pp. D297-D300.

[55128] 8958.

[55129] Groussin, P.; Renard, J.-P.; Chavanne, D.; Bertrand, G.; Avril,

[55130] J.; Simier, J.-L.: Antigens HLA d'une famille ayant une ade-nomatose

[55131] polyendocrienne de type I. (Letter) Presse Med. 9: 1033-1034, 1980.

[55132] 8959.

[55133] Jung, K. Y.; Warter, S.; Rumpler, Y.: Assignment of the GDH loci

[55134] to human chromosomes 10q23 and Xq24 by in situ hy-bridization. Ann.

[55135] Genet. 32: 109-110, 1989.

[55136] 8960.

[55137] Ho, Y.; Elefant, F.; Cooke, N.; Liebhaber, S.: A defined locus

[55138] control region determinant links chromatin domain acetylation with

[55139] long-range gene activation. Molec. Cell 9: 291-302, 2002.

[55140] 8961.

[55141] Accolla, R. S.; Gross, N.; Carrel, S.; Corte, G.: Distinct forms

[55142] of both alpha and beta subunits are present in the human Ia molecular

[55143] pool. Proc. Nat. Acad. Sci. 78: 4549-4551, 1981.

[55144] 8962.

[55145] Haavisto, A.-M.; Pettersson, K.; Bergendahl, M.; Virkamaki, A.;

[55146] Huhtaniemi, I.: Occurrence and biological properties of a common

[55147] genetic variant of luteinizing hormone. J. Clin. Endocr. Metab. 80:

[55148] 1257-1263, 1995.

[55149] 8963.

[55150] Motyckova, G.; Weilbaecher, K. N.; Horstmann, M.; Rie-
man, D. J.;

[55151] Fisher, D. Z.; Fisher, D. E.: Linking osteopetrosis and pyc-
nodysostosis:

[55152] regulation of cathepsin K expression by the microph-
thalmia transcription

[55153] factor family. Proc. Nat. Acad. Sci. 98: 5798-5803, 2001.

[55154] 8964.

[55155] Iida, M.; Murakami, T.; Ishida, K.; Mizuno, A.; Kuwajima,
M.;

[55156] Shima, K.: Substitution at codon 269
(glutamine-to-proline) of the

[55157] leptin receptor (OB-R) cDNA is the only mutation found in
the Zucker

[55158] fatty (fa/fa) rat. Biochem. Biophys. Res. Commun. 224:
597-604,

[55159] 1996.

[55160] 8965.

[55161] Gonnaud, P. M.; Sturtz, F.; Fourbil, Y.; Bonnebouche, C.;
Tranchant,

[55162] C.; Warter, J. M.; Chazot, G.; Bady, B.; Vial, C.; Brechard, A.
S.;

[55163] Vandenberghe, A.: DNA analysis as a tool to confirm the diagnosis

[55164] of asymptomatic hereditary neuropathy with liability to pressure palsies

[55165] (HNPP) with further evidence for the occurrence of de novo mutations. Acta.

[55166] Neurol. Scand. 92: 313-318, 1995.

[55167] 8966.

[55168] Thomas, K. R.; Folger, K. R.; Capecchi, M. R. :Cell 44: 419-428,

[55169] 1986.

[55170] 8967.

[55171] Peters, D. J. M.; Spruit, L.; Saris, J. J.; Ravine, D.; Sandkuijl,

[55172] L. A.; Fossdal, R.; Boersma, J.; van Eijk, R.; Norby, S.; Constantinou-Deltas,

[55173] C. D.; Pierides, A.; Brissenden, J. E.; Frants, R. R.; van Ommen,

[55174] G.-J. B.; Breuning, M. H.: Chromosome 4 localization of a second

[55175] gene for autosomal dominant polycystic kidney disease. Nature Genet. 5:

[55176] 359-362, 1993.

[55177] 8968.

[55178] Watanabe, K.; Kessler, C. A.; Bachurski, C. J.; Kanda, Y.; Richardson,

[55179] B. D.; Stanek, J.; Handwerger, S.; Brar, A. K.: Identification of

[55180] a decidua-specific enhancer on the human prolactin gene with two critical

[55181] activator protein 1 (AP-1) binding sites. Molec. Endocr. 15: 638-653,

[55182] 2001.

[55183] 8969.

[55184] Guillon, H.; de Massy, B.: An initiation site for meiotic crossing-over

[55185] and gene conversion in the mouse. Nature Genet. 32: 296-299, 2002.

[55186] 8970.

[55187] Berx, G.; Cleton-Jansen, A.-M.; Nollet, F.; de Leeuw, W. J. F.;

[55188] van de Vijver, M. J.; Cornelisse, C.; van Roy, F.: E-cadherin is

[55189] a tumour/invasion suppressor gene mutated in human lobular breast
[55190] cancers. EMBO J. 14: 6107-6115, 1995.

[55191] 8971.

[55192] Haines, J. L.; Boustany, R.-M. N.; Alroy, J.; Auger, K. J.; Shook,
[55193] K. S.; Terwedow, H.; Lerner, T. J.: Chromosomal localization of two
[55194] genes underlying late-infantile neuronal ceroid lipofuscinosis. Neurogenetics 1:
[55195] 217-222, 1998.

[55196] 8972.

[55197] Hari, K. L.; Santerre, A.; Sekelsky, J. J.; McKim, K. S.; Boyd,
[55198] J. B.; Hawley, R. S.: The mei-41 gene of D. melanogaster is a structural
[55199] and functional homolog of the human ataxia telangiectasia gene. Cell 82:
[55200] 815-821, 1995.

[55201] 8973.

[55202] Krasnoshtein, F.; Buchwald, M.: Developmental expression of the

[55203] Fac gene correlates with congenital defects in Fanconi anemia patients. Hum.
[55204] Molec. Genet. 5: 85-93, 1996.

[55205] 8974.

[55206] Hoeg, J. M.; Osborne, J. C., Jr.; Gregg, R. E.; Brewer, H. B.,

[55207] Jr.: Initial diagnosis of lipoprotein lipase deficiency in a 75-year-old
[55208] man. Am. J. Med. 75: 889-892, 1983.

[55209] 8975.

[55210] Barth, M. L.; Ward, C.; Harris, A.; Saad, A.; Fensom, A.: Frequency
[55211] of arylsulphatase A pseudodeficiency associated mutations in a healthy
[55212] population. J. Med. Genet. 31: 667-671, 1994.

[55213] 8976.

[55214] Sander, J. E.; Malamud, N.; Cowan, M. J.; Packman, S.; Amman,
[55215] A. J.; Wara, D. W.: Intermittent ataxia and immunodeficiency with
[55216] multiple carboxylase deficiencies: a biotin-responsive disorder. Ann.

[55217] Neurol. 8: 544-547, 1980.

[55218] 8977.

[55219] Lyon, M. F.; Hulse, E. V.: An inherited kidney disease of mice

[55220] resembling human nephronophthisis. J. Med. Genet. 8: 41-48, 1971.

[55221] 8978.

[55222] Mitchell, G. A.; Robert, M.-F.; Hruz, P. W.; Wang, S.; Fontaine,

[55223] G.; Behnke, C. E.; Mende-Mueller, L. M.; Schappert, K.; Lee, C.; Gibson,

[55224] K. M.; Miziorko, H. M.: 3-Hydroxy-3-methylglutaryl coen-zyme A lyase

[55225] (HL): cloning of human and chicken liver HL cDNAs and characterization

[55226] of a mutation causing human HL deficiency. J. Biol. Chem. 268: 4376-4381,

[55227] 1993.

[55228] 8979.

[55229] Papayannis, A. G.; Israels, M. C. G.: Glanzmann's disease and

[55230] trait. (Letter) Lancet II: 44 only, 1970.

[55231] 8980.

[55232] Dreyer, B.; Tranebjaerg, L.; Brox, V.; Rosenberg, T.; Moller, C.;

[55233] Beneyto, M.; Weston, M. D.; Kimberling, W. J.; Nilssen, O.: A common

[55234] ancestral origin of the frequent and widespread 2299delG USH2A mutation. Am.

[55235] J. Hum. Genet. 69: 228-234, 2001. Note: Erratum: Am. J. Hum. Genet.

[55236] 69: 922 only, 2001.

[55237] 8981.

[55238] Moser, H. W.: Adrenoleukodystrophy: phenotype, genetics, pathogenesis

[55239] and therapy. Brain 120: 1485-1508, 1997.

[55240] 8982.

[55241] Batischev, A. I.; Chernyak, N. B.; Torakev, Y. N.: Detection

[55242] of a new abnormal variant of glucose-6-phosphate dehy-drogenase in

[55243] human red cells. Bulleten Eksperimental Noi Biologii I Meditsiny 84:

[55244] 728-731, 1977.

[55245] 8983.

[55246] Parolini, S.; Bottino, C.; Falco, M.; Augugliaro, R.; Giliani,

[55247] S.; Franceschini, R.; Ochs, H. D.; Wolf, H.; Bonnefoy, J.-Y.; Biassoni,

[55248] R.; Moretta, L.; Notarangelo, L. D.; Moretta, A.: X-linked lymphoproliferative

[55249] disease: 2B4 molecules displaying inhibitory rather than activating

[55250] function are responsible for the inability of natural killer cells

[55251] to kill Epstein-Barr virus-infected cells. J. Exp. Med. 192: 337-346,

[55252] 2000.

[55253] 8984.

[55254] Rosse, W. F.: Epidemiology of PNH. Lancet 348: 560 only, 1996.

[55255] 8985.

[55256] Hawkins, J. R.; Taylor, A.; Goodfellow, P. N.; Migeon, C. J.;

[55257] Smith, K. D.; Berkovitz, G. D.: Evidence for increased prevalence

[55258] of SRY mutations in XY females with complete rather than partial gonadal
[55259] dysgenesis. Am. J. Hum. Genet. 51: 979-984, 1992.

[55260] 8986.

[55261] Sato, T. N.; Qin, Y.; Kozak, C. A.; Audus, K. L.: tie-1 and tie-2
[55262] define another class of putative receptor tyrosine kinase genes expressed
[55263] in early embryonic vascular system. Proc. Nat. Acad. Sci. 90: 9355-9358,
[55264] 1993.

[55265] 8987.

[55266] Phelan, C. M.; Lancaster, J. M.; Tonin, P.; Gumbs, C.; Cochran,
[55267] C.; Carter, R.; Ghadirian, P.; Perret, C.; Moslehi, R.; Dion, F.;
[55268] Faucher, M. C.; Dole, K.; Karimi, S.; Foulkes, W.; Lounis, H.; Warner,
[55269] E.; Goss, P.; Anderson, D.; Larsson, C.; Narod, S. A.; Futreal, P.
[55270] A.: Mutation analysis of the BRCA2 gene in 49 site-specific breast

[55271] cancer families. Nature Genet. 13: 120-122, 1996.

[55272] 8988.

[55273] Harrington, J. J.; Lieber, M. R.: The characterization of a mammalian

[55274] DNA structure-specific endonuclease. EMBO J. 13: 1235-1246, 1994.

[55275] 8989.

[55276] Lee, M. P.; DeBaun, M.; Randhawa, G.; Reichard, B. A.; Elledge,

[55277] S. J.; Feinberg, A. P.: Low frequency of p57(KIP2) mutation in Beckwith-Wiedemann

[55278] syndrome. Am. J. Hum. Genet. 61: 304-309, 1997.

[55279] 8990.

[55280] Yi, T.; Cleveland, J. L.; Ihle, J. N.: Identification of novel

[55281] protein tyrosine phosphatases of hematopoietic cells by polymerase

[55282] chain reaction amplification. Blood 78: 2222-2228, 1991.

[55283] 8991.

[55284] Amano, M.; Mukai, H.; Ono, Y.; Chihara, K.; Matsui, T.; Hamajima,

[55285] Y.; Okawa, K.; Iwamatsu, A.; Kaibuchi, K.: Identification of a putative

[55286] target for rho as the serine-threonine kinase protein kinase N. Science 271:

[55287] 648-651, 1996.

[55288] 8992.

[55289] Gorlatov, S. N.; Stadtman, T. C.: Human thioredoxin reductase

[55290] from HeLa cells: selective alkylation of selenocysteine in the protein

[55291] inhibits enzyme activity and reduction with NADPH influences affinity

[55292] to heparin. Proc. Nat. Acad. Sci. 95: 8520-8525, 1998.

[55293] 8993.

[55294] Ibraghimov-Beskrovnaya, O.; Dackowski, W. R.; Foggensteiner, L.;

[55295] Coleman, N.; Thiru, S.; Petry, L. R.; Burn, T. C.; Connors, T. D.;

[55296] Van Raay, T.; Bradley, J.; Qian, F.; Onuchic, L. F.; Watnick, T. J.;

[55297] Piontek, K.; Hakim, R. M.; Landes, G. M.; Germino, G. G.; Sandford,

[55298] R.; Klinger, K. W.: Polycystin: in vitro synthesis, in vivo tissue
[55299] expression, and subcellular localization identifies a large membrane-associated
[55300] protein. Proc. Nat. Acad. Sci. 94: 6397-6402, 1997.

[55301] 8994.

[55302] McPherron, A. C.; Lee, S.-J.: Suppression of body fat accumulation
[55303] in myostatin-deficient mice. J. Clin. Invest. 109: 595-601, 2002.
[55304] 8995.

[55305] Abdelhak, S.; Kalatzis, V.; Heilig, R.; Compain, S.; Samson, D.;
[55306] Vincent, C.; Levi-Acobas, F.; Cruaud, C.; Le Merrer, M.; Mathieu,
[55307] M.; Konig, R.; Vigneron, J.; Weissenbach, J.; Petit, C.; Weil, D.
[55308] : Clustering of mutations responsible for branchio-oto-renal (BOR)
[55309] syndrome in the eyes absent homologous region (eyaHR) of EYA1. Hum.
[55310] Molec. Genet. 6: 2247-2255, 1997.

[55311] 8996.

[55312] Masuda, H.; Tanaka, K.; Takagi, M.; Ohgami, K.; Sakamaki, T.; Shibata,

[55313] N.; Takahashi, K.: Molecular cloning and characterization of human

[55314] non-smooth muscle calponin. J. Biochem. 120: 415-424, 1996.

[55315] 8997.

[55316] Sarfarazi, M.; Child, A.; Stoilova, D.; Brice, G.; Desai, T.; Trifan,

[55317] O. C.; Poinoosawmy, D.; Crick, R. P.: Localization of the fourth

[55318] locus (GLC1E) for adult-onset primary open-angle glaucoma to the 10p15-p14

[55319] region. Am. J. Hum. Genet. 62: 641-652, 1998.

[55320] 8998.

[55321] Vlangos, C. N.; Das, P.; Patel, P. I.; Elsea, S. H.: Assignment

[55322] of developmentally regulated GTP-binding protein (DRG2) to human chromosome

[55323] band 17p11.2 with somatic cell hybrids and localization to the Smith-Magenis

[55324] syndrome critical interval. Cytogenet. Cell Genet. 88: 283-285,
[55325] 2000.

[55326] 8999.

[55327] Augustin, M.; Bamberger, C.; Paul, D.; Schmale, H.: Cloning and
[55328] chromosomal mapping of the human p53-related KET gene to chromosome
[55329] 3q27 and its murine homolog Ket to mouse chromosome 16. Mammalian
[55330] Genome 9: 899-902, 1998.

[55331] 9000.

[55332] Ohmori, M.; Ohta, M.; Shimura, H.; Shimura, Y.; Suzuki, K.; Kohn,
[55333] L. D.: Cloning of the single strand DNA-binding protein important
[55334] for maximal expression and thyrotropin (TSH)-induced negative regulation
[55335] of the TSH receptor. Molec. Endocr. 10: 1407-1424, 1996.

[55336] 9001.

[55337] Peyrard, M.: Personal Communication. Stockholm, Sweden

1/7/1999.

[55338] 9002.

[55339] Ware, F. E.; Lehrman, M. A.: Expression cloning of a novel suppressor

[55340] of the Lec15 and Lec35 glycosylation mutations of Chinese hamster

[55341] ovary cells. J. Biol. Chem. 271: 13935-13938, 1996.

[55342] 9003.

[55343] Boehm, J.; Orth, T.; Van Nguyen, P.; Soling, H. D.: Systemic lupus

[55344] erythematosus is associated with increased auto-antibody titers against

[55345] calreticulin and grp94, but calreticulin is not the Ro/SS-A antigen. Europ.

[55346] J. Clin. Invest. 24: 248-257, 1994.

[55347] 9004.

[55348] Burns, K.; Duggan, B.; Atkinson, E. A.; Famulski, K. S.; Nemer,

[55349] M.; Bleackley, R. C.; Michalak, M.: Modulation of gene expression

[55350] by calreticulin binding to the glucocorticoid receptor. Na-

ture 367:

[55351] 476-480, 1994.

[55352] 9005.

[55353] Telen, M. J.; Whitsett, C. F.: Erythrocyte acetyl-
cholinesterase

[55354] bears the Cartwright blood group antigens. (Abstract)
Clin. Res. 40:

[55355] 170A only, 1992.

[55356] 9006.

[55357] David, G.; Terris, B.; Marchio, A.; Lavau, C.; Dejean, A.:
The

[55358] acute promyelocytic leukemia PML-RAR-alpha protein in-
duces hepatic

[55359] preneoplastic and neoplastic lesions in transgenic mice.
Oncogene 14:

[55360] 1547-1554, 1997.

[55361] 9007.

[55362] Wjst, M.; Fischer, G.; Immervoll, T.; Jung, M.; Saar, K.;
Rueschendorf,

[55363] F.; Reis, A.; Ulbrecht, M.; Gomolka, M.; Weiss, E. H.;
Jaeger, L.;

[55364] Nickel, R.; and 14 others: A genome-wide search for linkage to
[55365] asthma. Genomics 58: 1-8, 1999.
[55366] 9008.
[55367] Kalachikov, S.; Evgrafov, O.; Ross, B.; Winawer, M.; Barker-Cummings,
[55368] C.; Boneschi, F. M.; Choi, C.; Morozov, P.; Das, K.; Teplitskaya,
[55369] E.; Yu, A.; Cayanis, E.; Penchaszadeh, G.; Kottmann, A. H.; Pedley,
[55370] T. A.; Hauser, W. A.; Ottman, R.; Gilliam, T. C.: Mutations in LGI1
[55371] cause autosomal-dominant partial epilepsy with auditory features. Nature
[55372] Genet. 30: 335-341, 2002.
[55373] 9009.
[55374] Dezateux, C. A.; Hyde, J. C.; Hoey, H. M. C. V.; O'Riordan, J.
[55375] L. H.; Spitz, L.; Taylor, G. W.; Grant, D. B.: Neonatal hyperparathyroidism. Europ.
[55376] J. Pediat. 142: 135-136, 1984.

[55377] 9010.

[55378] Gordon, M. S.; Kato, R. M.; Lansigan, F.; Thompson, A. A.; Wall,

[55379] R.; Rawlings, D. J.: Aberrant B cell receptor signaling from B29

[55380] (Ig-beta, CD79b) gene mutations of chronic lymphocytic leukemia B

[55381] cells. Proc. Nat. Acad. Sci. 97: 5504-5509, 2000.

[55382] 9011.

[55383] de Sousa, M.; Reimao, R.; Lacerda, R.; Hugo, P.; Kaufmann, S.

[55384] H. E.; Porto, G.: Iron overload in beta-2-microglobulin-deficient

[55385] mice. Immun. Lett. 39: 105-111, 1994.

[55386] 9012.

[55387] Morissette, J.; Rheaume, E.; Leblanc, J.-F.; Luu-The, V.; Labrie,

[55388] F.; Simard, J.: Genetic linkage mapping of HSD3B1 and HSD3B2 encoding

[55389] human types I and II 3-beta-hydroxysteroid dehydroge-nase/delta-5-delta-4-isomerase

[55390] close to D1S514 and the centromeric D1Z5 locus. Cyto-
genet. Cell
[55391] Genet. 69: 59-62, 1995.

[55392] 9013.

[55393] Morrison, N.; Nickson, D. A.; McBride, M. W.; Mueller, U.
W.; Boyd,
[55394] E.; Sutcliffe, R. G.: Regional chromosomal assignment of
human 3-beta-hydroxy-5-ene
[55395] steroid dehydrogenase to 1p13.1 by non-isotopic in situ
hybridisation.
[55396] Hum. Genet. 87: 223-225, 1991.

[55397] 9014.

[55398] Gallione, C. J.; Pasyk, K. A.; Boon, L. M.; Lennon, F.; John-
son,
[55399] D. W.; Helmbold, E. A.; Markel, D. S.; Vikkula, M.; Mulliken,
J. B.;
[55400] Warman, M. L.; Pericak-Vance, M. A.; Marchuk, D. A.: A
gene for familial
[55401] venous malformations maps to chromosome 9p in a sec-
ond large kindred. J.
[55402] Med. Genet. 32: 197-199, 1995.

[55403] 9015.

[55404] Chini, B.; Clementi, F.; Hukovic, N.; Sher, E.: Neuronal-type

[55405] alpha-bungarotoxin receptors and the alpha-5-nicotinic receptor subunit

[55406] gene are expressed in neuronal and nonneuronal human cell lines. Hum.

[55407] Genet. 89: 1572-1576, 1992.

[55408] 9016.

[55409] Kumar, B. V.; Aleman-Gomez, J. A.; Colwell, N.; Lopez-Candales,

[55410] A.; Bosner, M. S.; Spilburg, C. A.; Lowe, M.; Lange, L. G.: Structure

[55411] of the human pancreatic cholesterol esterase gene. Bio-chemistry 31:

[55412] 6077-6081, 1992.

[55413] 9017.

[55414] Lidberg, U.; Nilsson, J.; Stromberg, K.; Stenman, G.; Sahlin, P.;

[55415] Enerback, S.; Bjursell, G.: Genomic organization, sequence analysis,

[55416] and chromosomal localization of the human carboxyl ester lipase (CEL)
[55417] gene and a CEL-like (CELL) gene. Genomics 13: 630-640, 1992.
[55418] 9018.
[55419] Galland, F.; Stefanova, M.; Pirisi, V.; Birnbaum, D.: Characterization
[55420] of a murine glyceraldehyde-3-phosphate dehydrogenase pseudogene. Biochimie 72:
[55421] 759-762, 1990.
[55422] 9019.
[55423] Lidmer, A.-S.; Kannius, M.; Lundberg, L.; Bjursell, G.; Nilsson,
[55424] J.: Molecular cloning and characterization of the mouse carboxyl
[55425] ester lipase gene and evidence for expression in the lactating mammary
[55426] gland. Genomics 29: 115-122, 1995.
[55427] 9020.
[55428] Nilsson, J.; Hellquist, M.; Bjursell, G.: The human carboxyl ester

[55429] lipase-like (CELL) gene is ubiquitously expressed and contains a hypervariable
[55430] region. Genomics 17: 416-422, 1993.

[55431] 9021.

[55432] Taylor, A. K.; Zambaux, J. L.; Klisak, I.; Mohandas, T.; Sparkes,
[55433] R. S.; Schotz, M. C.; Lusis, A. J.: Carboxyl-ester lipase: a highly
[55434] polymorphic locus on chromosome 9qter. Genomics 10: 425-431, 1991.

[55435] 9022.

[55436] Vanin, E. F.: Processed pseudogenes: characteristics and evolution.
[55437] Annu. Rev. Genet. 19: 253-272, 1985.

[55438] 9023.

[55439] Klewer, S. E.; Krob, S. L.; Kolker, S. J.; Kitten, G. T.: Ex-pression
[55440] of type VI collagen in the developing mouse heart. :Dev. Dyn. 211:
[55441] 248-255, 1998.

[55442] 9024.

[55443] Lamande, S. R.; Bateman, J. F.; Hutchison, W.; McKinlay Gardner,
[55444] R. J.; Bower, S. P.; Byrne, E.; Dahl, H.-H. M.: Reduced collagen
[55445] VI causes Bethlem myopathy: a heterozygous COL6A1 nonsense mutation
[55446] results in mRNA decay and functional haploinsufficiency. Hum. Molec.
[55447] Genet. 7: 981-989, 1998.

[55448] 9025.

[55449] Murata, K.; Motoyama, T.; Suka, M.; Ohno, M.; Kuboki, Y.: High
[55450] production of type VI collagen in multiple fibromatosis with multiple
[55451] articular dysplasia. Biochem. Biophys. Res. Commun. 147: 275-281,
[55452] 1987.

[55453] 9026.

[55454] Pepe, G.; Giusti, B.; Bertini, E.; Brunelli, T.; Saitta, B.; Comeglio,
[55455] P.; Bolognese, A.; Merlini, L.; Federici, G.; Abbate, R.; Chu, M.-L.

[55456] : A heterozygous splice site mutation in COL6A1 leading to an in-frame
[55457] deletion of the alpha-1(VI) collagen chain in an Italian family affected
[55458] by Bethlem myopathy. Biochem. Biophys. Res. Commun. 258: 802-807,
[55459] 1999.

[55460] 9027.

[55461] Roest, P. A. M.; Roberts, R. G.; Sugino, S.; van Ommen, G.-J.
[55462] B.; den Dunnen, J. T.: Protein truncation test (PTT) for rapid detection
[55463] of translation-terminating mutations. Hum. Molec. Genet. 2: 1719-1721,
[55464] 1993.

[55465] 9028.

[55466] Scacheri, P. C.; Gillanders, E. M.; Subramony, S. H.; Veda-narayanan,
[55467] V.; Crowe, C. A.; Thakore, N.; Bingler, M.; Hoffman, E. P.: Novel
[55468] mutations in collagen VI genes: expansion of the Bethlem myopathy

[55469] phenotype. Neurology 58: 593-602, 2002.

[55470] 9029.

[55471] Avramopoulos, D.; Cox, T.; Forrest, G. L.; Chakravarti, A.; Antonarakis,

[55472] S. E.: Linkage mapping of the carbonyl reductase (CBR) gene on human

[55473] chromosome 21 using a DNA polymorphism in the 3-prime untranslated

[55474] region. Genomics 13: 447-448, 1992.

[55475] 9030.

[55476] Forrest, G. L.; Akman, S.; Krutzik, S.; Paxton, R. J.; Sparkes,

[55477] R. S.; Doroshow, J.; Felsted, R. L.; Glover, C. J.; Mohandas, T.;

[55478] Bachur, N. R.: Induction of a human carbonyl reductase gene located

[55479] on chromosome 21. Biochim. Biophys. Acta 1048: 149-155, 1990.

[55480] 9031.

[55481] Lemieux, N.; Malfoy, B.; Forrest, G. L.: Human carbonyl reductase

[55482] (CBR) localized to band 21q22.1 by high-resolution fluo-
rescence in
[55483] situ hybridization displays gene dosage effects in trisomy
21 cells. Genomics 15:
[55484] 169-172, 1993.

[55485] 9032.

[55486] Watanabe, K.; Sugawara, C.; Ono, A.; Fukuzumi, Y.;
Itakura, S.;
[55487] Yamazaki, M.; Tashiro, H.; Osoegawa, K.; Soeda, E.; No-
mura, T.: Mapping
[55488] of a novel human carbonyl reductase, CBR3, and riboso-
mal pseudogenes
[55489] to human chromosome 21q22.2. Genomics 52: 95-100,
1998.
[55490] 9033.

[55491] Wei, J.; Dlouhy, S. R.; Hara, A.; Ghetti, B.; Hodes, M. E.:
Cloning
[55492] a cDNA for carbonyl reductase (Cbr) from mouse cerebel-
lum: murine
[55493] genes that express Cbr map to chromosomes 16 and 11.
Genomics 34:
[55494] 147-148, 1996.

[55495] 9034.

[55496] Wermuth, B.; Bohren, K. M.; Heinemann, G.; von Wartburg, J.-P.;

[55497] Gabbay, K. H.: Human carbonyl reductase: nucleotide se-

quence analysis

[55498] of a cDNA and amino acid sequence of the encoded pro-

tein. J. Biol.

[55499] Chem. 263: 16185-16188, 1988.

[55500] 9035.

[55501] Lowe, D. G.; Klisak, I.; Sparkes, R. S.; Mohandas, T.; Goed-

del,

[55502] D. V.: Chromosomal distribution of three members of the human natriuretic

[55503] peptide receptor/guanylyl cyclase gene family. Genomics 8: 304-312,

[55504] 1990.

[55505] 9036.

[55506] Morgan, D.; Diamond, D. M.; Gottschall, P. E.; Ugen, K. E.; Dickey,

[55507] C.; Hardy, J.; Duff, K.; Jantzen, P.; DiCarlo, G.; Wilcock, D.; Connor,

[55508] K.; Hatcher, J.; Hope, C.; Gordon, M.; Arendash, G. W.: A-beta peptide

[55509] vaccination prevents memory loss in an animal model of Alzheimer's

[55510] disease. Nature 408: 982-985, 2000.

[55511] 9037.

[55512] Mullan, M.; Crawford, F.; Axelman, K.; Houlden, H.; Lilius, L.;

[55513] Winblad, B.; Lannfelt, L. :Nature Genet. 1: 345-349, 1992.

[55514] 9038.

[55515] Mullan, M.; Crawford, F.; Axelman, K.; Houlden, H.; Lilius, L.;

[55516] Winblad, B.; Lannfelt, L.: A pathogenic mutation for probable Alzheimer's

[55517] disease in the APP gene at the N-terminus of beta-amyloid. Nature

[55518] Genet. 1: 345-347, 1992.

[55519] 9039.

[55520] Multhaup, G.; Schlicksupp, A.; Hesse, L.; Beher, D.; Ruppert,

[55521] T.; Masters, C. L.; Beyreuther, K.: The amyloid precursor

protein

[55522] of Alzheimer's disease in the reduction of copper(II) to copper (I). Science 271:

[55523] 1406-1409, 1996.

[55524] 9040.

[55525] Murrell, J.; Farlow, M.; Ghetti, B.; Benson, M. D.: A mutation

[55526] in the amyloid precursor protein associated with hereditary Alzheimer's

[55527] disease. Science 254: 97-99, 1991.

[55528] 9041.

[55529] Naruse, S.; Igarashi, S.; Kobayashi, H.; Aoki, K.; Inuzuka, T.;

[55530] Kaneko, K.; Shimizu, T.; Iihara, K.; Kojima, T.; Miyatake, T.; Tsuji,

[55531] S.: Mis-sense mutation val-to-ile in exon 17 of amyloid precursor

[55532] protein gene in Japanese familial Alzheimer's disease.(Letter) Lancet 337:

[55533] 978-979, 1991.

[55534] 9042.

[55535] Natte, R.; Maat-Schieman, M. L. C.; Haan, J.; Bornebroek, M.;
[55536] Roos, R. A. C.; van Duinen, S. G.: Dementia in hereditary cerebral
[55537] hemorrhage with amyloidosis-Dutch type is associated with cerebral
[55538] amyloid angiopathy but is independent of plaques and neurofibrillary
[55539] tangles. Ann. Neurol. 50: 765-772, 2001.

[55540] 9043.

[55541] Olson, J. M.; Goddard, K. A. B.; Dudek, D. M.: A second lo-cus
[55542] for very-late-onset Alzheimer disease: a genome scan re-veals linkage
[55543] to 20p and epistasis between 20p and the amyloid pre-cursor protein
[55544] region. Am. J. Hum. Genet. 71: 154-161, 2002.

[55545] 9044.

[55546] Olson, J. M.; Goddard, K. A. B.; Dudek, D. M.: The amyloid precursor
[55547] protein locus and very-late-onset Alzheimer disease. Am. J. Hum.

[55548] Genet. 69: 895-899, 2001.

[55549] 9045.

[55550] Patterson, D.; Gardiner, K.; Kao, F.-T.; Tanzi, R.; Watkins, P.;

[55551] Gusella, J. F.: Mapping of the gene encoding the beta-amyloid precursor

[55552] protein and its relationship to the Down syndrome region of chromosome

[55553] 21. Proc. Nat. Acad. Sci. 85: 8266-8270, 1988.

[55554] 9046.

[55555] Peacock, M. L.; Murman, D. L.; Sima, A. A. F.; Warren, J. T.,

[55556] Jr.; Roses, A. D.; Fink, J. K.: Novel amyloid precursor protein gene

[55557] mutation (codon 665asp) in a patient with late-onset Alzheimer disease. Ann.

[55558] Neurol. 35: 432-438, 1994.

[55559] 9047.

[55560] Prelli, F.; Levy, E.; van Duinen, S. G.; Bots, G. T. A. M.; Luyendijk,

[55561] W.; Frangione, B.: Expression of a normal and variant Alzheimer's

[55562] beta-protein gene in amyloid of hereditary cerebral hemorrhage, Dutch
[55563] type: DNA and protein diagnostic assays. Biochem. Biophys. Res. Commun. 170:
[55564] 301-307, 1990.

[55565] 9048.

[55566] Robakis, N. K.; Ramakrishna, N.; Wolfe, G.; Wisniewski, H. M.
[55567] : Molecular cloning and characterization of a cDNA encoding the cerebrovascular
[55568] and the neuritic plaque amyloid peptides. Proc. Nat. Acad. Sci. 84:
[55569] 4190-4194, 1987.

[55570] 9049.

[55571] Robakis, N. K.; Wisniewski, H. M.; Jenkins, E. C.; Devine-Gage,
[55572] E. A.; Houck, G. E.; Yao, X.-L.; Ramakrishna, N.; Wolfe, G.; Silverman,
[55573] W. P.; Brown, W. T.: Chromosome 21q21 sublocalisation of gene encoding
[55574] beta-amyloid peptide in cerebral vessels and neuritic (senile) plaques

[55575] of people with Alzheimer disease and Down syndrome.(Letter) Lancet I:
[55576] 384-385, 1987.

[55577] 9050.

[55578] Sandbrink, R.; Masters, C. L.; Beyreuther, K.: Beta A4-amyloid
[55579] protein precursor mRNA isoforms without exon 15 are ubiquitously expressed
[55580] in rat tissues including brain, but not in neurons. J. Biol. Chem. 269:
[55581] 1510-1517, 1994.

[55582] 9051.

[55583] Schellenberg, G. D.; Anderson, L.; O'dahl, S.; Wisjman, E. M.;
[55584] Sadovnick, A. D.; Ball, M. J.; Larson, E. B.; Kukull, W. A.; Martin,
[55585] G. M.; Roses, A. D.; Bird, T. D.: APP(717), APP(693), and PRIP gene
[55586] mutations are rare in Alzheimer disease. Am. J. Hum. Genet. 49:
[55587] 511-517, 1991.

[55588] 9052.

[55589] Schenk, D.; Barbour, R.; Dunn, W.; Gordon, G.; Grajeda, H.; Guido,

[55590] T.; Hu, K.; Huang, J.; Johnson-Wood, K.; Khan, K.; Kholo-denko, D.;

[55591] Lee, M.; and 13 others: Immunization with amyloid-beta attenuates

[55592] Alzheimer-disease-like pathology in the PDAPP mouse. Nature 400:

[55593] 173-177, 1999.

[55594] 9053.

[55595] Schmaier, A. H.; Dahl, L. D.; Rozemuller, A. J. M.; Roos, R. A.

[55596] C.; Wagner, S. L.; Chung, R.; Van Nostrand, W. E.: Protease nexin-2/amyloid-beta

[55597] protein precursor: a tight-binding inhibitor of coagulation factor

[55598] IXa. J. Clin. Invest. 92: 2540-2545, 1993.

[55599] 9054.

[55600] Senior, K.: Alzheimer's and prion disease: do they share a pathogenic

[55601] mechanism? Lancet 355: 1434 only, 2000.

[55602] 9055.

[55603] Sisodia, S. S.; Gallagher, M.: A role for the beta-amyloid precursor

[55604] protein in memory? Proc. Nat. Acad. Sci. 95: 12074-12076, 1998.

[55605] 9056.

[55606] Smith, R. P.; Higuchi, D. A.; Broze, G. J., Jr.: Platelet coagulation

[55607] factor XI(a)-inhibitor: a form of Alzheimer amyloid precursor protein. Science 248:

[55608] 1126-1128, 1990.

[55609] 9057.

[55610] St. George-Hyslop, P.; Crapper McLachlan, D.; Tuda, T.; Rogaev,

[55611] E.: Alzheimer's disease and possible gene interaction.(Letter) Science 263:

[55612] 537 only, 1994.

[55613] 9058.

[55614] Sturchler-Pierrat, C.; Abramowski, D.; Duke, M.; Wiederhold, K.-H.;

[55615] Mistl, C.; Rothacher, S.; Ledermann, B.; Burki, K.; Frey, P.; Paganetti,

[55616] P. A.; Waridel, C.; Calhoun, M. E.; Jucker, M.; Probst, A.; Staufenbiel,

[55617] M.; Sommer, B.: Two amyloid precursor protein transgenic mouse models

[55618] with Alzheimer disease-like pathology. Proc. Nat. Acad. Sci. 94:

[55619] 13287-13292, 1997.

[55620] 9059.

[55621] Sugarman, M. C.; Yamasaki, T. R.; Oddo, S.; Echegoyen, J. C.;

[55622] Murphy, M. P.; Golde, T. E.; Jannatipour, M.; Leissring, M. A.; LaFerla,

[55623] F. M.: Inclusion body myositis-like phenotype induced by transgenic

[55624] overexpression of beta-APP in skeletal muscle. Proc. Nat. Acad. Sci. 99:

[55625] 6334-6339, 2002.

[55626] 9060.

[55627] Suzuki, N.; Cheung, T. T.; Cai, X.-D.; Odaka, A.; Otvos, L., Jr.;

[55628] Eckman, C.; Golde, T. E.; Younkin, S. G.: An increased percentage

[55629] of long amyloid beta protein secreted by familial amyloid beta protein

[55630] precursor (beta-APP-717) mutants. Science 264: 1336-1340, 1994.

[55631] 9061.

[55632] Tang, M. X.; Jacobs, D.; Stern, Y.; Marder, K.; Schofield, P.;

[55633] Gurland, B.; Andrews, H.; Mayeux, R.: Effect of oestrogen during

[55634] menopause on risk and age of onset of Alzheimer's disease. Lancet 348:

[55635] 429-432, 1996.

[55636] 9062.

[55637] AbdAlla, S.; Lother, H.; Quitterer, U.: AT(1)-receptor heterodimers

[55638] show enhanced G-protein activation and altered receptor sequestration. Nature 407:

[55639] 94-98, 2000.

[55640] 9063.

[55641] Bergsma, D. J.; Ellis, C.; Kumar, C.; Nuthulaganti, P.; Ker-

sten,

[55642] H.; Elshourbagy, N.; Griffin, E.; Stadel, J. M.; Aiyar, N.: Cloning

[55643] and characterization of a human angiotensin II type 1 receptor. Biochem.

[55644] Biophys. Res. Commun. 183: 989-995, 1992.

[55645] 9064.

[55646] Bonnardeaux, A.; Davies, E.; Jeunemaitre, X.; Fery, I.; Charru,

[55647] A.; Clauser, E.; Tiret, L.; Cambien, F.; Corvol, P.; Soubrier, F.

[55648] : Angiotensin II type 1 receptor gene polymorphisms in human essential

[55649] hypertension. Hypertension 24: 63-69, 1994.

[55650] 9065.

[55651] Curnow, K. M.; Pascoe, L.; White, P. C.: Genetic analysis of the

[55652] human type-1 angiotensin II receptor. Molec. Endocr. 6: 1113-1118,

[55653] 1992.

[55654] 9066.

[55655] Elton, T. S.; Stephan, C. C.; Taylor, G. R.; Kimball, M. G.; Martin,
[55656] M. M.; Durand, J. N.; Oparil, S.: Isolation of two distinct type
[55657] I angiotensin II receptor genes. Biochem. Biophys. Res. Commun. 184:
[55658] 1067-1073, 1992.

[55659] 9067.

[55660] Furuta, H.; Guo, D.-F.; Inagami, T.: Molecular cloning and sequencing
[55661] of the gene encoding human angiotensin II type 1 receptor. Biochem.
[55662] Biophys. Res. Commun. 183: 8-13, 1992.

[55663] 9068.

[55664] Gemmill, R. M.; Drabkin, H. A.: Report of The Second International
[55665] Workshop on Human Chromosome 3 Mapping. Cytogenet. Cell Genet. 57:
[55666] 162-166, 1991.

[55667] 9069.

[55668] Guo, D.-F.; Furuta, H.; Mizukoshi, M.; Inagami, T.: The ge-

nomic
[55669] organization of human angiotensin II type 1 receptor. Biochem. Biophys.
[55670] Res. Commun. 200: 313-319, 1994.
[55671] 9070.
[55672] Harada, K.; Komuro, I.; Hayashi, D.; Sugaya, T.; Murakami, K.;
[55673] Yazaki, Y.: Angiotensin II type 1a receptor is involved in the occurrence
[55674] of reperfusion arrhythmias. Circulation 97: 315-317, 1998.
[55675] 9071.
[55676] Haywood, G. A.; Gullestad, L.; Katsuya, T.; Hutchinson, H. G.;
[55677] Pratt, R. E.; Horiuchi, M.; Fowler, M. B.: AT(1) and AT(2) angiotensin
[55678] receptor gene expression in human heart failure. Circulation 95:
[55679] 1201-1206, 1997.
[55680] 9072.
[55681] Herzig, T. C.; Jobe, S. M.; Aoki, H.; Molkentin, J. D.; Cow-

ley,
[55682] A. W., Jr.; Izumo, S.; Markham, B. E.: Angiotensin II type-1a receptor
[55683] gene expression in the heart: AP-1 and GATA-4 participate in the response
[55684] to pressure overload. Proc. Nat. Acad. Sci. 94: 7543-7548, 1997.
[55685] 9073.

[55686] Ito, M.; Oliverio, M. I.; Mannon, P. J.; Best, C. F.; Maeda, N.;

[55687] Smithies, O.; Coffman, T. M.: Regulation of blood pressure by type
[55688] 1A angiotensin II receptor gene. Proc. Nat. Acad. Sci. 92: 3521-3525,
[55689] 1995.

[55690] 9074.

[55691] Iyer, S. N.; Lu, D.; Katovich, M. J.; Raizada, M. K.: Chronic

[55692] control of high blood pressure in the spontaneously hypertensive rat
[55693] by delivery of angiotensin type 1 receptor antisense. Proc. Nat.
[55694] Acad. Sci. 93: 9960-9965, 1996.

[55695] 9075.

[55696] Konishi, H.; Kuroda, S.; Inada, Y.; Fujisawa, Y.: Novel sub-
type
[55697] of human angiotensin II type 1 receptor: cDNA cloning
and expression. Biochem.
[55698] Biophys. Res. Commun. 199: 467-474, 1994.

[55699] 9076.

[55700] Martens, J. R.; Reaves, P. Y.; Lu, D.; Katovich, M. J.; Bere-
cek,
[55701] K. H.; Bishop, S. P.; Raizada, M. K.; Gelband, C. H.: Preven-
tion
[55702] of renovascular and cardiac pathophysiological changes in
hypertension
[55703] by angiotensin II type 1 receptor antisense gene therapy.
Proc. Nat.
[55704] Acad. Sci. 95: 2664-2669, 1998.

[55705] 9077.

[55706] Martin, M. M.; Willardson, B. M.; Burton, G. F.; White, C. R.;

[55707] McLaughlin, J. N.; Bray, S. M.; Ogilvie, J. W., Jr.; Elton, T. S.

[55708] : Human angiotensin II type 1 receptor isoforms encoded
by messenger

[55709] RNA splice variants are functionally distinct. Molec. Endocr. 15:
[55710] 281-293, 2001.

[55711] 9078.

[55712] Mauzy, C. A.; Hwang, O.; Egloff, A. M.; Wu, L.-H.; Chung, F.-Z.
[55713] : Cloning, expression, and characterization of a gene encoding the
[55714] human angiotensin II type 1A receptor. Biochem. Biophys. Res. Commun. 186:
[55715] 277-284, 1992.

[55716] 9079.

[55717] Murphy, T. J.; Alexander, R. W.; Griendling, K. K.; Runge, M.
[55718] S.; Bernstein, K. E.: Isolation of a cDNA encoding the vascular type-1
[55719] angiotensin II receptor. Nature 351: 233-236, 1991.

[55720] 9080.

[55721] Paradis, P.; Dali-Youcef, N.; Paradis, F. W.; Thibault, G.; Nemer,
[55722] M.: Overexpression of angiotensin II type I receptor in car-

diomyocytes

[55723] induces cardiac hypertrophy and remodeling. Proc. Nat. Acad. Sci. 97:

[55724] 931-936, 2000.

[55725] 9081.

[55726] Sasaki, K.; Murohara, T.; Ikeda, H.; Sugaya, T.; Shimada, T.;

[55727] Shintani, S.; Imaizumi, T.: Evidence for the importance of angiotensin

[55728] II type 1 receptor in ischemia-induced angiogenesis. J. Clin. Invest. 109:

[55729] 603-611, 2002.

[55730] 9082.

[55731] Oliverio, M. I.; Kim, H-S.; Ito, M.; Le, T.; Audoly, L.; Best,

[55732] C. F.; Hiller, S.; Kluckman, K.; Maeda, N.; Smithies, O.; Coffman,

[55733] T. M.: Reduced growth, abnormal kidney structure, and type 2 (AT2)

[55734] angiotensin receptor-mediated blood pressure regulation in mice lacking

[55735] both AT1A and AT1B receptors for angiotensin II. Proc.

Nat. Acad.

[55736] Sci. 95: 15496-15501, 1998.

[55737] 9083.

[55738] Sasaki, K.; Yamano, Y.; Bardhan, S.; Iwai, N.; Murray, J. J.;

[55739] Hasegawa, M.; Matsuda, Y.; Inagami, T.: Cloning and ex-
pression of

[55740] a complementary DNA encoding a bovine adrenal an-
giotensin II type-1

[55741] receptor. Nature 351: 230-233, 1991.

[55742] 9084.

[55743] Scott, A. F.: Personal Communication. Baltimore, Md.
3/20/2001.

[55744] 9085.

[55745] Szpirer, C.; Riviere, M.; Szpirer, J.; Levan, G.; Guo, D. F.;

[55746] Iwai, N.; Inagami, T.: Chromosomal assignment of human
and rat hypertension

[55747] candidate genes: type 1 angiotensin II receptor genes and
the SA gene. J.

[55748] Hypertension 11: 919-925, 1993.

[55749] 9086.

[55750] Takayanagi, R.; Ohnaka, K.; Sakai, Y.; Nakao, R.; Yanase, T.;

[55751] Haji, M.; Inagami, T.; Furuta, H.; Gou, D.-F.; Nakamuta, M.; Nawata,

[55752] H.: Molecular cloning, sequence analysis and expression of a cDNA

[55753] encoding human type-1 angiotensin II receptor. Biochem. Biophys.

[55754] Res. Commun. 183: 910-916, 1992.

[55755] 9087.

[55756] Tsuchida, S.; Matsusaka, T.; Chen, X.; Okubo, S.; Niimura, F.;

[55757] Nishimura, H.; Fogo, A.; Utsunomiya, H.; Inagami, T.; Ichikawa, I.

[55758] : Murine double nullizygotes of the angiotensin type 1A and 1B receptor

[55759] genes duplicate severe abnormal phenotypes of angiotensinogen nullizygotes. J.

[55760] Clin. Invest. 101: 755-760, 1998.

[55761] 9088.

[55762] Erba, H. P.; Eddy, R.; Shows, T.; Kedes, L.; Gunning, P.: Structure,

[55763] chromosome location, and expression of the human gamma-actin gene:

[55764] differential evolution, location, and expression of the cytoskeletal

[55765] beta- and gamma-actin genes. Molec. Cell. Biol. 8: 1775-1789, 1988.

[55766] 9089.

[55767] Ueyama, H.; Inazawa, J.; Nishino, H.; Ohkubo, I.; Miwa, T.: FISH

[55768] localization of human cytoplasmic actin genes ACTB to 7p22 and ACTG1

[55769] to 17q25 and characterization of related pseudogenes. Cytogenet.

[55770] Cell Genet. 74: 221-224, 1996.

[55771] 9090.

[55772] Beggs, A. H.; Byers, T. J.; Knoll, J. H. M.; Boyce, F. M.; Bruns,

[55773] G. A. P.; Kunkel, L. M.: Cloning and characterization of two human

[55774] skeletal muscle alpha-actinin genes located on chromosomes 1 and 11. J.

[55775] Biol. Chem. 267: 9281-9288, 1992.

[55776] 9091.

[55777] Beggs, A. H.; Phillips, H. A.; Kozman, H.; Mulley, J. C.; Wilton,

[55778] S. D.; Kunkel, L. M.; Laing, N. G.: A (CA)n repeat polymor-phism for

[55779] the human skeletal muscle alpha-actinin gene ACTN2 and its localization

[55780] on the linkage map of chromosome 1. Genomics 13: 1314-1315, 1992.

[55781] 9092.

[55782] Mills, M. A.; Yang, N.; Weinberger, R. P.; Vander Woude, D. L.;

[55783] Beggs, A. H.; Easteal, S.; North, K. N.: Differential expres-sion

[55784] of the actin-binding proteins, alpha-actinin-2 and -3, in different

[55785] species: implications for the evolution of functional re-dundancy. Hum.

[55786] Molec. Genet. 10: 1335-1346, 2001.

[55787] 9093.

[55788] Youssoufian, H.; McAfee, M.; Kwiatkowski, D. J.: Cloning and chromosomal

[55789] localization of the human cytoskeletal alpha-actinin gene reveals
[55790] linkage to the beta-spectrin gene. Am. J. Hum. Genet. 47: 62-72,
[55791] 1990.

[55792] 9094.

[55793] Attisano, L.; Carcamo, J.; Ventura, F.; Weis, F. M. B.; Massague,
[55794] J.; Wrana, J. L.: Identification of human activin and TGF-beta type
[55795] I receptors that form heteromeric kinase complexes with type II receptors. Cell 75:
[55796] 671-680, 1993.

[55797] 9095.

[55798] Mathews, L. S.; Vale, W. W.: Expression cloning of an activin
[55799] receptor, a predicted transmembrane serine kinase. Cell 65: 973-982,
[55800] 1991.

[55801] 9096.

[55802] Matsuzaki, K.; Xu, J.; Wang, F.; McKeehan, W. L.; Krum-

men, L.;

[55803] Kan, M.: A widely expressed transmembrane serine/ threonine kinase

[55804] that does not bind activin, inhibin, transforming growth factor beta,

[55805] or bone morphogenic factor. J. Biol. Chem. 268: 12719-12723, 1993.

[55806] 9097.

[55807] Roijer, E.; Miyazono, K.; Astrom, A.-K.; Geurts van Kessel, A.;

[55808] ten Dijke, P.; Stenman, G.: Chromosomal localization of three human

[55809] genes encoding members of the TGF-beta superfamily of type I serine/threonine

[55810] kinase receptors. Mammalian Genome 9: 266-268, 1998.

[55811] 9098.

[55812] ten Dijke, P.; Ichijo, H.; Franzen, P.; Schulz, P.; Saras, J.;

[55813] Toyoshima, H.; Heldin, C.-H.; Miyazono, K.: Activin recep- tor-like

[55814] kinases: a novel subclass of cell-surface receptors with predicted

[55815] serine/threonine kinase activity. Oncogene 8: 2879-2887,

1993.

[55816] 9099.

[55817] Chen, M.; Pan, Z.-Q.; Hurwitz, J.: Studies of the cloned 37-kDa

[55818] subunit of activator 1 (replication factor C) of HeLa cells. Proc.

[55819] Nat. Acad. Sci. 89: 5211-5215, 1992.

[55820] 9100.

[55821] Alcalay, M.; Zangrilli, D.; Fagioli, M.; Pandolfi, P. P.; Mencarelli,

[55822] A.; Lo Coco, F.; Biondi, A.; Grignani, F.; Pelicci, P. G.: Expression

[55823] pattern of the RAR-alpha-PML fusion gene in acute promyelocytic leukemia. Proc.

[55824] Nat. Acad. Sci. 89: 4840-4844, 1992.

[55825] 9101.

[55826] Brown, D.; Kogan, S.; Lagasse, E.; Weissman, I.; Alcalay, M.; Pelicci,

[55827] P. G.; Atwater, S.; Bishop, J. M.: A PMLRAR-alpha transgene initiates

[55828] murine acute promyelocytic leukemia. Proc. Nat. Acad. Sci.

94: 2551-2556,

[55829] 1997.

[55830] 9102.

[55831] Chang, K.-S.; Fan, Y.-H.; Andreeff, M.; Liu, J.; Mu, Z.-M.: The

[55832] PML gene encodes a phosphoprotein associated with the nuclear matrix. Blood 85:

[55833] 3646-3653, 1995.

[55834] 9103.

[55835] Cleary, M. L.: Oncogenic conversion of transcription fac-tors by

[55836] chromosomal translocations. Cell 66: 619-622, 1991.

[55837] 9104.

[55838] Jonkman, M. F.: Revertant mosaicism in human genetic disorders. Am.

[55839] J. Med. Genet. 85: 361-364, 1999.

[55840] 9105.

[55841] Jonkman, M. F.; de Jong, M. C. J. M.; Heeres, K.; Pas, H. H.;

[55842] van der Meer, J. B.; Owaribe, K.; Martinez de Velasco, A. M.; Niessen,

[55843] C. M.; Sonnenberg, A.: 180-kD bullous pemphigoid antigen (BP180)

[55844] is deficient in generalized atrophic benign epidermolysis bullosa. J.

[55845] Clin. Invest. 95: 1345-1352, 1995.

[55846] 9106.

[55847] Jonkman, M. F.; De Jong, M. C. J. M.; Heeres, K.; Steijlen, P.

[55848] M.; Owaribe, K.; Kuster, W.; Meurer, M.; Gedde-Dahl, T., Jr.; Sonnenberg,

[55849] A.; Bruckner-Tuderman, L.: Generalized atrophic benign epidermolysis

[55850] bullosa: either 180-kd bullous pemphigoid antigen or laminin-5 deficiency. Arch.

[55851] Derm. 132: 145-150, 1996.

[55852] 9107.

[55853] Li, K.; Sawamura, D.; Giudice, G. J.; Diaz, L. A.; Mattei, M.-G.;

[55854] Chu, M.-L.; Uitto, J.: Genomic organization of collagenous domains

[55855] and chromosomal assignment of human 180-kDa bullous pemphigoid antigen-2,

[55856] a novel collagen of stratified squamous epithelium. J. Biol. Chem. 266:
[55857] 24064-24069, 1991.

[55858] 9108.

[55859] Li, K.; Tamai, K.; Tan, E. M. L.; Uitto, J.: Cloning of type

[55860] XVII collagen: complementary and genomic sequences of mouse 180-kDa
[55861] bullous pemphigoid antigen (BPAG2) predict an interrupted collagenous
[55862] domain, a transmembranous segment, and unusual features in the 5-prime
[55863] end of the gene and the 3-prime-untranslated region of the mRNA. J.
[55864] Biol. Chem. 268: 8825-8834, 1993.

[55865] 9109.

[55866] Sawamura, D.; Li, K.; Nomura, K.; Sugita, Y.; Christiano, A. M.;
[55867] Uitto, J.: Bullous pemphigoid antigen: cDNA cloning, cellular expression,
[55868] and evidence for polymorphism of the human gene. J. Invest. Derm. 96:

[55869] 908-915, 1991.

[55870] 9110.

[55871] Sawamura, D.; Li, K.; Uitto, J.: 230-kD and 180-kD bullous pemphigoid

[55872] antigens are distinct gene products. (Letter) J. Invest. Derm. 98:

[55873] 942-943, 1992.

[55874] 9111.

[55875] Schacke, H.; Schumann, H.; Hammami-Hauasli, N.; Raghu-

[55876] nath, M.;

[55876] Bruckner-Tuderman, L.: Two forms of collagen XVII in keratinocytes.

[55877] A full-length transmembrane protein and a soluble ectodomain. J.

[55878] Biol. Chem. 273: 25937-25943, 1998.

[55879] 9112.

[55880] Schumann, H.; Hammami-Hauasli, N.; Pulkkinen, L.; Mau-

[55881] viel, A.;

[55881] Kuster, W.; Luthi, U.; Owaribe, K.; Uitto, J.; Bruckner-Tu-

[55882] derman,

[55882] L.: Three novel homozygous point mutations and a new

polymorphism
[55883] in the COL17A1 gene: relation to biological and clinical phenotypes
[55884] of junctional epidermolysis bullosa. Am. J. Hum. Genet. 60: 1344-1353,
[55885] 1997.

[55886] 9113.

[55887] Tasanen, K.; Eble, J. A.; Aumailley, M.; Schumann, H.; Baetge,
[55888] J.; Tu, H.; Bruckner, P.; Bruckner-Tuderman, L.: Collagen XVII is
[55889] destabilized by a glycine substitution mutation in the cell adhesion
[55890] domain Col15. J. Biol. Chem. 275: 3093-3099, 2000.

[55891] 9114.

[55892] Tasanen, K.; Floeth, M.; Schumann, H.; Bruckner-Tuder-
man, L.:
[55893] Hemizygosity for a glycine substitution in collagen XVII: unfolding
[55894] and degradation of the ectodomain. J. Invest. Derm. 115: 207-212,
[55895] 2000.

[55896] 9115.

[55897] Wahn, V.; Stephan, V.; Hirschhorn, R.: Reverse mutations--spontaneous

[55898] amelioration or cure of inherited disorders? Europ. J. Pediat. 157:

[55899] 613-617, 1998.

[55900] 9116.

[55901] Youssoufian, H.: Natural gene therapy and the Darwinian legacy. Nature

[55902] Genet. 13: 255-256, 1996.

[55903] 9117.

[55904] Buettner, R.; Schaffler, A.; Arndt, H.; Rogler, G.; Nusser, J.;

[55905] Zietz, B.; Enger, I.; Hugl, S.; Cuk, A.; Scholmerich, J.; Palitzsch,

[55906] K.-D.: The trp64arg polymorphism of the beta-3-adrenergic receptor

[55907] gene is not associated with obesity or type 2 diabetes mellitus in

[55908] a large population-based Caucasian cohort. J. Clin. Endocr. Metab. 83:

[55909] 2892-2897, 1998.

[55910] 9118.

[55911] Clement, K.; Vaisse, C.; Manning, B. S. J.; Basdevant, A.;
Guy-Grand,
[55912] B.; Ruiz, J.; Silver, K. D.; Shuldiner, A. R.; Froguel, P.;
Strosberg,
[55913] A. D.: Genetic variation in the beta-3-adrenergic receptor
and an
[55914] increased capacity to gain weight in patients with morbid
obesity. New
[55915] Eng. J. Med. 333: 352-354, 1995.

[55916] 9119.

[55917] Elbein, S. C.; Hoffman, M.; Barrett, K.; Wegner, K.; Miles,
C.;
[55918] Bachman, K.; Berkowitz, D.; Shuldiner, A. R.; Leppert, M.
F.; Hasstedt,
[55919] S.: Role of the beta-3-adrenergic receptor locus in obesity
and noninsulin-dependent
[55920] diabetes among members of Caucasian families with a di-
abetic sibling
[55921] pair. J. Clin. Endocr. Metab. 81: 4422-4427, 1996.

[55922] 9120.

[55923] Emorine, L. J.; Marullo, S.; Briend-Sutren, M.-M.; Patey, G.; Tate,
[55924] K.; Delavier-Klutchko, C.; Strosberg, A. D.: Molecular characterization
[55925] of the human beta-3-adrenergic receptor. Science 245: 1118-1121,
[55926] 1989.

[55927] 9121.

[55928] Festa, A.; Krugluger, W.; Shnawa, N.; Hopmeier, P.; Haffner, S.
[55929] M.; Schernthaner, G.: Trp64Arg polymorphism of the beta-3-adrenergic
[55930] receptor gene in pregnancy: association with mild gestational diabetes
[55931] mellitus. J. Clin. Endocr. Metab. 84: 1695-1699, 1999.

[55932] 9122.

[55933] Gagnon, J.; Mauriege, P.; Roy, S.; Sjostrom, D.; Chagnon, Y. C.;
[55934] Dionne, F. T.; Oppert, J.-M.; Perusse, L.; Sjostrom, L.; Bouchard,
[55935] C.: The trp64arg mutation of the beta-3 adrenergic receptor gene

[55936] has no effect on obesity phenotypes in the Quebec Family Study and
[55937] Swedish Obese Subjects cohorts. J. Clin. Invest. 98: 2086-2093,
[55938] 1996.

[55939] 9123.

[55940] Garcia-Rubi, E.; Starling, R. D.; Tchernof, A.; Matthews, D. E.;
[55941] Walston, J. D.; Shuldiner, A. R.; Silver, K.; Poehlman, E. T.; Calles-Escandon,
[55942] J.: Trp64Arg variant of the beta-3-adrenoceptor and in-
sulin resistance
[55943] in obese postmenopausal women. J. Clin. Endocr. Metab. 83: 4002-4005,
[55944] 1998.

[55945] 9124.

[55946] Hoffstedt, J.; Poirier, O.; Thorne, A.; Lonnqvist, F.; Her-
rmann,
[55947] S. M.; Cambien, F.; Arner, P.: Polymorphism of the human beta-3-adrenoceptor
[55948] gene forms a well-conserved haplotype that is associated with moderate

[55949] obesity and altered receptor function. Diabetes 48: 203-205, 1999.

[55950] 9125.

[55951] Kim-Motoyama, H.; Yasuda, K.; Yamaguchi, T.; Yamada, N.; Katakura,

[55952] T.; Shuldiner, A. R.; Akanuma, Y.; Ohashi, Y.; Yazaki, Y.; Kadowaki,

[55953] T.: A mutation of the beta-3-adrenergic receptor is associated with

[55954] visceral obesity but decreased serum triglyceride. Diabetologia 40:

[55955] 469-472, 1997.

[55956] 9126.

[55957] Mitchell, B. D.; Blangero, J.; Comuzzie, A. G.; Almasy, L. A.;

[55958] Shuldiner, A. R.; Silver, K.; Stern, M. P.; MacCluer, J. W.; Hixson,

[55959] J. E.: A paired sibling analysis of the beta-3 adrenergic receptor

[55960] and obesity in Mexican Americans. J. Clin. Invest. 101: 584-587,

[55961] 1998.

[55962] 9127.

[55963] Nagase, T.; Aoki, A.; Yamamoto, M.; Yasuda, H.; Kado, S.; Nishikawa,

[55964] M.; Kugai, N.; Akatsu, T.; Nagata, N.: Lack of association between

[55965] the trp64arg mutation in the beta-3-adrenergic receptor gene and obesity

[55966] in Japanese men: a longitudinal analysis. J. Clin. Endocr. Metab. 82:

[55967] 1284-1287, 1997.

[55968] 9128.

[55969] Shihara, N.; Yasuda, K.; Moritani, T.; Ue, H.; Adachi, T.; Tanaka,

[55970] H.; Tsuda, K.; Seino, Y.: The association between Trp64Arg polymorphism

[55971] of the beta-3-adrenergic receptor and autonomic nervous system activity. J.

[55972] Clin. Endocr. Metab. 84: 1623-1627, 1999.

[55973] 9129.

[55974] Van Spronsen, A.; Nahmias, C.; Krief, S.; Briend-Sutren, M.-M.;

[55975] Strosberg, A. D.; Emorine, L. J.: The promoter and intron/ exon structure

[55976] of the human and mouse beta-3-adrenergic-receptor genes. Europ. J.

[55977] Biochem. 213: 1117-1124, 1993.

[55978] 9130.

[55979] Urhammer, S. A.; Hansen, T.; Borch-Johnsen, K.; Pedersen, O.:

[55980] Studies of the synergistic effect of the trp/arg64 polymor- phism of

[55981] the beta-3-adrenergic receptor gene and the -3826 A-G variant of the

[55982] uncoupling protein-1 gene on features of obesity and in- sulin resistance

[55983] in a population-based sample of 379 young Danish sub- jects. J. Clin.

[55984] Endocr. Metab. 85: 3151-3154, 2000.

[55985] 9131.

[55986] Walston, J.; Silver, K.; Bogardus, C.; Knowler, W. C.; Celi, F.

[55987] S.; Austin, S.; Manning, B.; Strosberg, A. D.; Stern, M. P.; Raben,

[55988] N.; Sorkin, J. D.; Roth, J.; Shuldiner, A. R.: Time of onset of non-insulin-dependent
[55989] diabetes mellitus and genetic variation in the beta-3-adrenergic-receptor
[55990] gene. New Eng. J. Med. 333: 343-347, 1995.

[55991] 9132.

[55992] Walston, J.; Silver, K.; Hilfiker, H.; Andersen, R. E.; Seibert,

[55993] M.; Beamer, B.; Roth, J.; Poehlman, E.; Shuldiner, A. R.: Insulin
[55994] response to glucose is lower in individuals homozygous for the arg64
[55995] variant of the beta-3-andrenergic receptor. J. Clin. Endocr. Metab. 85:
[55996] 4019-4022, 2000.

[55997] 9133.

[55998] Hershfield, M. S.; Mitchell, B. S.: Immunodeficiency diseases
[55999] caused by adenosine deaminase deficiency and purine nucleoside phosphorylase
[56000] deficiency.In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle,

[56001] D. (eds.): The Metabolic and Molecular Bases of Inherited

Disease.

[56002] Vol. 2. New York: McGraw-Hill (7th ed.): 1995. Pp. 1725-1768.

[56003] 9134.

[56004] Hirschhorn, R.: Identification of two new missense muta-tions

[56005] (R156C and S291L) in two ADA(-) SCID patients unusual for response

[56006] to therapy with partial exchange transfusions. Hum. Mu-tat. 1: 166-168,

[56007] 1992.

[56008] 9135.

[56009] Hirschhorn, R.: Personal Communication. New York, N.Y. 1976.

[56010] 9136.

[56011] Hirschhorn, R.; Borkowsky, W.; Jiang, C.-K.; Yang, D. R.; Jenkins,

[56012] T.: Two newly identified mutations (Thr233Ile and Leu152Met) in partially

[56013] adenosine deaminase-deficient (ADA-) individuals that result in differing

[56014] biochemical and metabolic phenotypes. Hum. Genet. 100:

22-29, 1997.

[56015] 9137.

[56016] Hirschhorn, R.; Chakravarti, V.; Puck, J.; Douglas, S. D.: Homozygosity

[56017] for a newly identified missense mutation in a patient with very severe

[56018] combined immunodeficiency due to adenosine deaminase deficiency (ADA-SCID). Am.

[56019] J. Hum. Genet. 49: 878-885, 1991.

[56020] 9138.

[56021] Hirschhorn, R.; Ellenbogen, A.: Genetic heterogeneity in adenosine

[56022] deaminase (ADA) deficiency: five different mutations in five new patients

[56023] with partial ADA deficiency. Am. J. Hum. Genet. 38: 13-25, 1986.

[56024] 9139.

[56025] Hirschhorn, R.; Ellenbogen, A.; Tzall, S.: Five missense mutations

[56026] at the adenosine deaminase locus (ADA) detected by altered restriction

[56027] fragments and their frequency in ADA-patients with se-

vere combined
[56028] immunodeficiency (ADA-SCID). Am. J. Med. Genet. 42: 201-207, 1992.
[56029] 9140.

[56030] Hirschhorn, R.; Levytska, V.; Parkman, R.: A mutant form of adenosine
[56031] deaminase in severe combined immunodeficiency. (Abstract) J. Clin.
[56032] Invest. 53: 33A, 1974.

[56033] 9141.

[56034] Hirschhorn, R.; Martiniuk, F.; Roegner-Maniscalco, V.; El-lenbogen,
[56035] A.; Perignon, J.-L.; Jenkins, T.: Genetic heterogeneity in partial
[56036] adenosine deaminase deficiency. J. Clin. Invest. 71: 1887-1892,
[56037] 1983.

[56038] 9142.

[56039] Hirschhorn, R.; Papageorgiou, P. S.; Kesarwala, H. H.; Taft, L.
[56040] T.: Amelioration of neurologic abnormalities after 'enzyme

replacement'
[56041] in adenosine deaminase deficiency. New Eng. J. Med. 303: 377-380,
[56042] 1980.

[56043] 9143.

[56044] Hirschhorn, R.; Roegner, V.; Jenkins, T.; Seaman, C.; Piomelli,
[56045] S.; Borkowsky, W.: Erythrocyte adenosine deaminase deficiency without
[56046] immunodeficiency: evidence for an unstable mutant enzyme. J. Clin.
[56047] Invest. 64: 1130-1139, 1979.

[56048] 9144.

[56049] Hirschhorn, R.; Tzall, S.; Ellenbogen, A.: Hot spot mutations
[56050] in adenosine deaminase deficiency. Proc. Nat. Acad. Sci. 87: 6171-6175,
[56051] 1990.

[56052] 9145.

[56053] Hirschhorn, R.; Tzall, S.; Ellenbogen, A.; Orkin, S. H.: Identification

[56054] of a point mutation resulting in a heat-labile adenosine deaminase

[56055] (ADA) in two unrelated children with partial ADA deficiency. J. Clin.

[56056] Invest. 83: 497-501, 1989.

[56057] 9146.

[56058] Hirschhorn, R.; Vawter, G. F.; Kirkpatrick, J. A., Jr.; Rosen,

[56059] F. S.: Adenosine deaminase deficiency: frequency and comparative

[56060] pathology in autosomally recessive severe combined immunodeficiency. Clin.

[56061] Immun. Immunopath. 14: 107-120, 1979.

[56062] 9147.

[56063] Hirschhorn, R.; Yang, D. R.; Israni, A.: An asp8-to-asn substitution

[56064] results in the adenosine deaminase (ADA) genetic polymorphism (ADA

[56065] 2 allozyme): occurrence on different chromosomal backgrounds and apparent

[56066] intragenic crossover. Ann. Hum. Genet. 58: 1-9, 1994.

[56067] 9148.

[56068] Hirschhorn, R.; Yang, D. R.; Israni, A.; Huie, M. L.; Ownby, D.

[56069] R.: Somatic mosaicism for a newly identified splice-site mutation

[56070] in a patient with adenosine deaminase-deficient immunodeficiency and

[56071] spontaneous clinical recovery. Am. J. Hum. Genet. 55: 59-68, 1994.

[56072] 9149.

[56073] Hirschhorn, R.; Yang, D. R.; Puck, J. M.; Huie, M. L.; Jiang,

[56074] C.-K.; Kurlandsky, L. E.: Spontaneous in vivo reversion to

normal

[56075] of an inherited mutation in a patient with adenosine deaminase deficiency. Nature

[56076] Genet. 13: 290-295, 1996.

[56077] 9150.

[56078] Hong, R.; Galti, R.; Rathbun, J. C.; Good, R. A.: Thymic hypoplasia

[56079] and thyroid dysfunction. New Eng. J. Med. 282: 470-474, 1970.

[56080] 9151.

[56081] Honig, J.; Martiniuk, F.; D'Eustachio, P.; Zamfirescu, C.; Desnick,

[56082] R.; Hirschhorn, K.; Hirschhorn, L. R.; Hirschhorn, R.: Confirmation

[56083] of the regional localization of the genes for human acid alpha-glucosidase

[56084] (GAA) and adenosine deaminase (ADA) by somatic cell hybridization. Ann.

[56085] Hum. Genet. 48: 49-56, 1984.

[56086] 9152.

[56087] Hopkinson, D. A.; Cook, P. J. L.; Harris, H.: Further data on

[56088] the adenosine deaminase (ADA) polymorphism and a report of a new phenotype. Ann.

[56089] Hum. Genet. 32: 361-368, 1969.

[56090] 9153.

[56091] Hutton, J. J.; Wiginton, D. A.; Coleman, M. S.; Fuller, S. A.;

[56092] Limouze, S.; Lampkin, B. C.: Biochemical and functional abnormalities

[56093] in lymphocytes from an adenosine deaminase-deficient patient during

[56094] enzyme replacement therapy. J. Clin. Invest. 68: 413-421, 1981.

[56095] 9154.

[56096] Jenkins, T.; Rabson, A. R.; Nurse, G. T.; Lane, A. B.; Hopkinson,

[56097] D. A.: Deficiency of adenosine deaminase not associated with severe

[56098] combined immunodeficiency. J. Pediat. 89: 732-736, 1976.

[56099] 9155.

[56100] Jhanwar, S. C.; Berkvens, T. M.; Breukel, C.; van Ormondt, H.;

[56101] van der Eb, A. J.; Meera Khan, P.: Localization of human adenosine
[56102] deaminase (ADA) gene sequences to the q12-q13.11 region of chromosome
[56103] 20 by in situ hybridization. Cytogenet. Cell Genet. 50: 168-171,
[56104] 1989.

[56105] 9156.

[56106] Jiang, C.-K.; Hong, R.; Horowitz, S. D.; Kong, X.-P.; Hirschhorn,
[56107] R.: An adenosine deaminase (ADA) allele contains two newly identified
[56108] deleterious mutations (Y97C and L106V) that interact to abolish enzyme
[56109] activity. Hum. Molec. Genet. 6: 2271-2278, 1997.

[56110] 9157.

[56111] Kaitila, I.; Rimoin, D. L.; Cederbaum, S. D.; Stiehm, E. R.; Lechman,
[56112] R. S.: Chondroosseous histopathology in adenosine deaminase deficient
[56113] combined immunodeficiency disease. Birth Defects Orig. Art. Ser. XII(6):

[56114] 115-121, 1976.

[56115] 9158.

[56116] Kellems, R. E.; Yeung, C.-Y.; Ingolia, D. E.: Adenosine deaminase

[56117] deficiency and severe combined immunodeficiencies. Trends Genet. 1:

[56118] 278-283, 1985.

[56119] 9159.

[56120] Kenny, A. B.; Hitzig, W. H.: Bone marrow transplantation for

[56121] severe combined immunodeficiency disease: reported from 1968-1977. Europ.

[56122] J. Pediat. 131: 155-176, 1979.

[56123] 9160.

[56124] Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine deaminase

[56125] expression and severe combined immunodeficiency dis- ease in humans. Proc.

[56126] Nat. Acad. Sci. 77: 4211-4215, 1980.

[56127] 9161.

[56128] Kredich, N. M.; Martin, D. W., Jr.: Role of
5-adenosylhomocysteine
[56129] in adenosine-mediated toxicity in cultured mouse T-
lymphoma cells. Cell 12:
[56130] 931-938, 1977.

[56131] 9162.

[56132] Levy, Y.; Hershfield, M. S.; Fernandez-Mejia, C.; Polmar, S.
H.;
[56133] Scudiery, D.; Berger, M.; Sorensen, R. U.: Adenosine
deaminase deficiency
[56134] with late onset of recurrent infections: response to treat-
ment with
[56135] polyethylene glycol-modified adenosine deaminase. J. Pe-
diat. 113:
[56136] 312-317, 1988.

[56137] 9163.

[56138] Sparkes, R. S.; Klisak, I.; Miller, W. L.: Regional mapping of

[56139] genes encoding human steroidogenic enzymes: P450scc
to 15q23-q24;
[56140] adrenodoxin to 11q22; adrenodoxin reductase to
17q24-q25; and P450c17

[56141] to 10q24-q25. DNA Cell Biol. 10: 359-365, 1991.

[56142] 9164.

[56143] Makino, Y.; Shibata, K.; Makino, I.; Kangawa, K.;
Kawarabayashi,
[56144] T.: Alteration of the adrenomedullin receptor components
gene expression
[56145] associated with the blood pressure in pregnancy-induced
hypertension. J.
[56146] Clin. Endocr. Metab. 86: 5079-5082, 2001.

[56147] 9165.

[56148] McLatchie, L. M.; Fraser, N. J.; Main, M. J.; Wise, A.; Brown,
[56149] J.; Thompson, N.; Solari, R.; Lee, M. G.; Foord, S. M.:
RAMPs regulate
[56150] the transport and ligand specificity of the calcitonin-recep-
tor-like
[56151] receptor. Nature 393: 333-339, 1998.

[56152] 9166.

[56153] Bell, A. C.; Felsenfeld, G.: Methylation of a CTCF-de-
pendent boundary
[56154] controls imprinted expression of the Igf2 gene. Nature
405: 482-485,

[56155] 2000.

[56156] 9167.

[56157] Ferguson-Smith, A. C.; Cattanach, B. M.; Barton, S. C.; Beechey,

[56158] C. V.; Surani, M. A.: Embryological and molecular investi-gations

[56159] of parental imprinting on mouse chromosome 7. Nature 351: 667-670,

[56160] 1991.

[56161] 9168.

[56162] Ohno, S.: Ancient linkage groups and frozen accidents. Nature 244:

[56163] 259-262, 1973.

[56164] 9169.

[56165] Bateman, J. B.; Kojis, T.; Heinzmann, C.; Klisak, I.; Diep, A.;

[56166] Carper, D.; Nishimura, C.; Mohandas, T.; Sparkes, R. S.: Mapping

[56167] of aldose reductase gene sequences to human chromo-somes 1, 3, 7, 9,

[56168] 11, and 13. Genomics 17: 560-565, 1993.

[56169] 9170.

[56170] Brown, L.; Hedge, P. J.; Markham, A. F.; Graham, A.: A human aldehyde

[56171] dehydrogenase (aldose reductase) pseudogene: nucleotide sequence analysis

[56172] and assignment to chromosome 3. Genomics 13: 465-468, 1992.

[56173] 9171.

[56174] Chung, S.; LaMendola, J.: Cloning and sequence determination of

[56175] human placental aldose reductase gene. J. Biol. Chem. 264: 14775-14777,

[56176] 1989.

[56177] 9172.

[56178] Graham, A.; Brown, L.; Hedge, P. J.; Gammack, A. J.; Markham, A.

[56179] F.: Structure of the human aldose reductase gene. J. Biol. Chem. 266:

[56180] 6872-6877, 1991.

[56181] 9173.

[56182] Graham, A.; Heath, P.; Morten, J. E. N.; Markham, A. F.:

The human

[56183] aldose reductase gene maps to chromosome region 7q35. Hum. Genet. 86:

[56184] 509-514, 1991.

[56185] 9174.

[56186] Nishimura, C.; Matsuura, Y.; Kokai, Y.; Akera, T.; Carper, D.;

[56187] Morjana, N.; Lyons, C.; Flynn, T. G.: Cloning and expres-sion of human

[56188] aldose reductase. J. Biol. Chem. 265: 9788-9792, 1990.

[56189] 9175.

[56190] Shah, V. O.; Dorin, R. I.; Sun, Y.; Braun, M.; Zager, P. G.: Aldose

[56191] reductase gene expression is increased in diabetic nephropathy. J.

[56192] Clin. Endocr. Metab. 82: 2294-2298, 1997.

[56193] 9176.

[56194] Pericak-Vance, M. A.; Bebout, J. L.; Gaskell, P. C., Jr.; Ya-maoka,

[56195] L. H.; Hung, W.-Y.; Alberts, M. J.; Walker, A. P.; Bartlett, R. J.;

[56196] Haynes, C. A.; Welsh, K. A.; Earl, N. L.; Heyman, A.; Clark, C. M.;
[56197] Roses, A. D.: Linkage studies in familial Alzheimer dis-
ease: evidence
[56198] for chromosome 19 linkage. Am. J. Hum. Genet. 48: 1034-1050, 1991.
[56199] 9177.
[56200] Roberds, S. L.; Anderson, J.; Basi, G.; Bienkowski, M. J.; Branstetter,
[56201] D. G.; Chen, K. S.; Freedman, S. B.; Frigon, N. L.; Games, D.; Hu,
[56202] K.; Johnson-Wood, K.; Kappenman, K. E.; and 20 others: BACE knockout
[56203] mice are healthy despite lacking the primary beta-
secretase activity
[56204] in brain: implications for Alzheimer's disease therapeutics. Hum.
[56205] Molec. Genet. 10: 1317-1324, 2001.
[56206] 9178.
[56207] Tanzi, R.; St. George-Hyslop, P.; Haines, J.; Neve, R.; Polinsky,
[56208] R.; Conneally, P. M.; Gusella, J. F.: Genetic linkage analysis

of
[56209] the Alzheimer's associated amyloid beta protein gene with familial
[56210] Alzheimer's disease and chromosome 21.(Abstract) Cyto-genet. Cell
[56211] Genet. 46: 703 only, 1987.

[56212] 9179.

[56213] Tanzi, R. E.; Bird, E. D.; Latt, S. A.; Neve, R. L.: The amyloid

[56214] beta protein gene is not duplicated in brains from patients with Alzheimer's
[56215] disease. Science 238: 666-669, 1987.

[56216] 9180.

[56217] Tanzi, R. E.; St. George-Hyslop, P. H.; Haines, J. L.; Polin-sky,
[56218] R. J.; Nee, L.; Foncin, J.-F.; Neve, R. L.; McClatchey, A. I.; Conneally,
[56219] P. M.; Gusella, J. F.: The genetic defect in familial Alzheimer's
[56220] disease is not tightly linked to the amyloid beta-protein gene. Nature 329:
[56221] 156-157, 1987.

[56222] 9181.

[56223] Van Broeckhoven, C.; Backhovens, H.; Raeymaekers, P.; Wehnert,

[56224] A.; Horsthemke, B.; Beyreuther, K.; Genthe, A.; Barton, A.; Hardy,

[56225] J.; Irving, N.; Williamson, R.; Vandenberghe, A.: Linkage study between

[56226] the amyloid gene and familial Alzheimer dis-ease.(Abstract) Cytogenet.

[56227] Cell Genet. 46: 708 only, 1987.

[56228] 9182.

[56229] Botto, M.; Hawkins, P. N.; Bickerstaff, M. C.; Herbert, J.; Bygrave,

[56230] A. E.; McBride, A.; Hutchinson, W. L.; Tennent, G. A.; Wal-port, M.

[56231] J.; Pepys, M. B.: Amyloid deposition is delayed in mice with targeted

[56232] deletion of the serum amyloid P component gene. Nature Med. 3: 855-859,

[56233] 1997.

[56234] 9183.

[56235] Tanzi, R. E.; Vaula, G.; Romano, D. M.; Mortilla, M.; Huang, T.

[56236] L.; Tupler, R. G.; Wasco, W.; Hyman, B. T.; Haines, J. L.; Jenkins,

[56237] B. J.; Kalaitsidaki, M.; Warren, A. C.; McInnis, M. C.; Antonarakis,

[56238] S. E.; Karlinsky, H.; Percy, M. E.; Connor, L.; Growdon, J.; Crapper-McLachlan,

[56239] D. R.; Gusella, J. F.; St. George-Hyslop, P. H.: Assessment of amyloid

[56240] beta-protein precursor gene mutations in a large set of familial and

[56241] sporadic Alzheimer disease cases. Am. J. Hum. Genet. 51: 273-282,

[56242] 1992.

[56243] 9184.

[56244] Van Broeckhoven, C.; Haan, J.; Bakker, E.; Hardy, J. A.; Van Hul,

[56245] W.; Wehnert, A.; Vegter-Van der Vlis, M.; Roos, R. A. C.: Amyloid

[56246] beta protein precursor gene and hereditary cerebral hemorrhage with

[56247] amyloidosis (Dutch). Science 248: 1120-1122, 1990.

[56248]

[56249] 100. van Duijn, C. M.; Hendriks, L.; Cruts, M.; Hardy, J. A.; Hofman,

[56250] A.; Van Broeckhoven, C.: Amyloid precursor protein gene mutation

[56251] in early-onset Alzheimer's disease.(Letter) Lancet 337: 978 only,

[56252] 1991.

[56253]

[56254] 101. van Duinen, S. G.; Castano, E. M.; Prelli, F.; Bots, G. T. A.

[56255] B.; Luyendijk, W.; Frangione, B.: Hereditary cerebral hem-

[56256] with amyloidosis in patients of Dutch origin is related to Alzheimer

[56257] disease. Proc. Nat. Acad. Sci. 84: 5991-5994, 1987.

[56258]

[56259] 102. van Leeuwen, F. W.; de Kleijn, D. P. V.; van den Hurk, H. H.;

[56260] Neubauer, A.; Sonnemans, M. A. F.; Sluijs, J. A.; Koycu, S.;

Ramdjielal,
[56261] R. D. J.; Salehi, A.; Martens, G. J. M.; Grosveld, F. G.; Burbach,
[56262] J. P. H.; Hol, E. M.: Frameshift mutants of beta-amyloid precursor
[56263] protein and ubiquitin-B in Alzheimer's and Down patients. Science 279:
[56264] 242-247, 1998.

[56265]

[56266] 103. Van Nostrand, W. E.; Wanger, S. L.; Suzuki, M.; Choi, B. H.;
[56267] Farrow, J. S.; Geddes, J. W.; Cotman, C. W.; Cunningham, D. D.: Protease
[56268] nexin-II, a potent antichymotrypsin, shows identity to amyloid beta-protein
[56269] precursor. Nature 341: 546-549, 1989.

[56270]

[56271] 104. Walsh, D. M.; Klyubin, I.; Fadeeva, J. V.; Cullen, W. K.; Anwyl,
[56272] R.; Wolfe, M. S.; Rowan, M. J.; Selkoe, D. J.: Naturally secreted
[56273] oligomers of amyloid beta protein potently inhibit hip-

pocampal long-term
potentiation in vivo. Nature 416: 535-539, 2002.

105. Wattendorff, A. R.; Bots, G. T. A. M.; Went, L. N.; Endtz, L.
J.: Familial cerebral amyloid angiopathy presenting as re-current
cerebral haemorrhage. J. Neurol. Sci. 55: 121-135, 1982.

106. Weggen, S.; Eriksen, J. L.; Das, P.; Sagi, S. A.; Wang, R.; Pietrzik,
C. U.; Findlay, K. A.; Smith, T. E.; Murphy, M. P.; Bulter, T.; Kang,
D. E.; Marquez-Sterling, N.; Golde, T. E.; Koo, E. H.: A subset of
NSAIDs lower amyloidogenic A-beta-42 independently of cyclooxygenase
activity. Nature 414: 212-216, 2001.

107. Wilson, C. A.; Doms, R. W.; Zheng, H.; Lee, V. M.-Y.: Presenilins

[56287] are not required for A-beta-42 production in the early secretory pathway. Nature

[56288] Neurosci. 5: 849-855, 2002.

[56289]

[56290] 108. Wyss-Coray, T.; Lin, C.; Yan, F.; Yu, G.; Rohde, M.; McConlogue,

[56291] L.; Masliah, E.; Mucke, L.: TGF-beta-1 promotes microglial amyloid-beta

[56292] clearance and reduces plaque burden in transgenic mice. Nature Med. 7:

[56293] 612-618, 2001.

[56294]

[56295] 109. Xu, H.; Gouras, G. K.; Greenfield, J. P.; Vincent, B.; Naslund,

[56296] J.; Mazzarelli, L.; Fried, G.; Jovanovic, J. N.; Seeger, M.; Relkin,

[56297] N. R.; Liao, F.; Checler, F.; Buxbaum, J. D.; Chait, B. T.; Thinakaran,

[56298] G.; Sisodia, S. S.; Wang, R.; Greengard, P.; Gandy, S.: Estrogen

[56299] reduces neuronal generation of Alzheimer beta-amyloid peptides. Nature

[56300] Med. 4: 447-451, 1998.

[56301]

[56302] 110. Yamatsuji, T.; Matsui, T.; Okamoto, T.; Komatsuzaki, K.; Takeda,

[56303] S.; Fukumoto, H.; Iwatsubo, T.; Suzuki, N.; Asami-Odaka, A.; Ireland,

[56304] S.; Kinane, T. B.; Giambarella, U.; Nishimoto, I.: G protein-mediated

[56305] neuronal DNA fragmentation induced by familial Alzheimer's disease-associated

[56306] mutants of APP. Science 272: 1349-1352, 1996.

[56307]

[56308] 111. Yan, S. D.; Chen, X.; Fu, J.; Chen, M.; Zhu, H.; Roher, A.; Slattery,

[56309] T.; Zhao, L.; Nagashima, M.; Morser, J.; Migheli, A.; Nawroth, P.;

[56310] Stern, D.; Schmidt, A. M.: RAGE and amyloid-beta peptide neurotoxicity

[56311] in Alzheimer's disease. Nature 382: 685-691, 1996.

[56312]

[56313] 112. Yoshikai, S.; Sasaki, H.; Doh-ura, K.; Furuya, H.;

Sakaki, Y.

[56314] : Genomic organization of the human amyloid beta-protein precursor

[56315] gene. Gene 87: 257-263, 1990.

[56316]

[56317] 113. Yoshioka, K.; Miki, T.; Katsuya, T.; Ogihara, T.; Sakaki, Y.

[56318] : The 717-val-to-ile substitution in amyloid precursor protein is

[56319] associated with familial Alzheimer's disease regardless of ethnic

[56320] groups. Biochem. Biophys. Res. Commun. 178: 1141-1146, 1991.

[56321]

[56322] 114. Zabel, B. U.; Salbaum, J. M.; Multhaup, G.; Master, C. L.; Bohl,

[56323] J.; Beyreuther, K.: Sublocalization of the gene for the pre-cursor

[56324] of Alzheimer's disease amyloid A4 protein on chromo-some 21.(Abstract) Cytogenet.

[56325] Cell Genet. 46: 725-726, 1987.

[56326]

[56327] 115. Zeldenrust, S. R.; Murrell, J.; Farlow, M.; Ghetti, B.; Roses,

[56328] A. D.; Benson, M. D.: RFLP analysis for APP 717 mutations associated

[56329] with Alzheimer's disease. J. Med. Genet. 30: 476-478, 1993.

[56330] 9185.

[56331] Floyd-Smith, G. A.; Whitehead, A. S.; Colten, H. R.; Francke, U.

[56332] : Human serum amyloid P component (SAP) is located on the proximal

[56333] long arm of chromosome 1. (Abstract) Cytogenet. Cell Genet. 40:

[56334] 631 only, 1985.

[56335] 9186.

[56336] Ionasescu, V.; Burns, T.; Searby, C.; Ionasescu, R.: Linkage between

[56337] the loci for Duffy (FY) and serum amyloid P component (APCS) on human

[56338] chromosome 1. Cytogenet. Cell Genet. 45: 240-241, 1987.

[56339] 9187.

[56340] Mantzouranis, E. C.; Dowton, S. B.; Whitehead, A. S.; Edge, M.
[56341] D.; Bruns, G. A. P.; Colten, H. R.: Human serum amyloid P component:
[56342] cDNA isolation, complete sequence of pre-serum amyloid P component,
[56343] and localization of the gene to chromosome 1. J. Biol. Chem. 260:
[56344] 7752-7756, 1985.

[56345] 9188.

[56346] Mortensen, R. F.; Le, P. T.; Taylor, B. A.: Mouse serum amyloid
[56347] P-component (SAP) levels controlled by a locus on chromosome 1. Immunogenetics 22:
[56348] 367-375, 1985.

[56349] 9189.

[56350] Pepys, M. B.; Herbert, J.; Hutchinson, W. L.; Tennent, G. A.; Lachmann,
[56351] H. J.; Gallimore, J. R.; Lovat, L. B.; Bartfai, T.; Alanine, A.; Hertel,
[56352] C.; Hoffmann, T.; Jakob-Roetne, R.; and 11 others: Targeted pharmacological

[56353] depletion of serum amyloid P component for treatment of human amyloidosis. Nature 417:
[56354] 254-259, 2002.

[56355] 9190.

[56356] Prelli, F.; Pras, M.; Frangione, B.: The primary structure of

[56357] human tissue amyloid P component from a patient with primary idiopathic
[56358] amyloidosis. J. Biol. Chem. 260: 12895-12898, 1985.

[56359] 9191.

[56360] Whitehead, A. S.; Rits, M.; Michaelson, J.: Molecular genet-ics
[56361] of mouse serum amyloid P component (SAP): cloning and gene mapping. Immunogenetics 28:
[56362] 388-390, 1988.

[56363] 9192.

[56364] Woo, P.; O'Brien, J.; Robson, M.; Ansell, B. M.: A genetic marker
[56365] for systemic amyloidosis in juvenile arthritis. Lancet I: 767-769,
[56366] 1987.

[56367] 9193.

[56368] Wasco, W.; Brook, J. D.; Tanzi, R. E.: The amyloid precursor-like

[56369] protein (APLP) gene maps to the long arm of human chromosome 19. Genomics 15:

[56370] 237-239, 1993.

[56371] 9194.

[56372] Leach, R.; Ko, M.; Krawetz, S. A.: Assignment of amyloid-precursor-like

[56373] protein 2 gene (APLP2) to 11q24 by fluorescent in situ hybridization. Cytogenet.

[56374] Cell Genet. 87: 215-216, 1999.

[56375] 9195.

[56376] von der Kammer, H.; Loffler, C.; Hanes, J.; Klaudiny, J.; Scheit,

[56377] K. H.; Hansmann, I.: The gene for the amyloid precursor-like protein

[56378] APLP2 is assigned to human chromosome 11q23-25. Genomics 10: 308-311,

[56379] 1994.

[56380] 9196.

[56381] von Koch, C. S.; Lahiri, D. K.; Mammen, A. L.; Copeland, N. G.;
[56382] Gilbert, D. J.; Jenkins, N. A.; Sisodia, S. S.: The mouse APLP2 gene:
[56383] chromosomal localization and promoter characterization. J. Biol.
[56384] Chem. 270: 25475-25480, 1995.

[56385] 9197.

[56386] Yan, Y. C.; Bai, Y.; Wang, L.; Miao, S.; Koide, S. S.: Characterization
[56387] of cDNA encoding a human sperm membrane protein related to A4 amyloid
[56388] protein. Proc. Nat. Acad. Sci. 87: 2405-2408, 1990.

[56389] 9198.

[56390] Yang, Y.; Martin, L.; Cuzin, F.; Mattei, M.-G.; Rassoulzadegan,
[56391] M.: Genomic structure and chromosomal localization of the mouse CDEI-binding
[56392] protein CDEBP (APLP2) gene and promoter sequences. Genomics 35:
[56393] 24-29, 1996.

[56394] 9199.

[56395] Wallukat, G.; Homuth, V.; Fischer, T.; Lindschau, C.; Horstkamp,

[56396] B.; Jupner, A.; Baur, E.; Nissen, E.; Vetter, K.; Neichel, D.; Dudenhausen,

[56397] J. W.; Haller, H.; Luft, F. C.: Patients with preeclampsia de-velop

[56398] agonistic autoantibodies against the angiotensin AT-1 re-ceptor. J.

[56399] Clin. Invest. 103: 945-952, 1999.

[56400] 9200.

[56401] Wang, W. Y. S.; Zee, R. Y. L.; Morris, B. J.: Association of

[56402] angiotensin II type 1 receptor gene polymorphism with essential hypertension. Clin.

[56403] Genet. 51: 31-34, 1997.

[56404] 9201.

[56405] Wilson, G. N.; Dasouki, M.; Barr, M., Jr.: Further delineation

[56406] of the dup(3q) syndrome. Am. J. Med. Genet. 22: 117-123, 1985.

[56407] 9202.

[56408] Tweardy, D. J.; Cannizzaro, L. A.; Palumbo, A. P.; Shane, S.;

[56409] Huebner, K.; Vantuinen, P.; Ledbetter, D. H.; Finan, J. B.; Nowell,

[56410] P. C.; Rovera, G.: Molecular cloning and characterization of a cDNA

[56411] for human granulocyte colony-stimulating factor (G-CSF) from a glioblastoma

[56412] multiforme cell line and localization of the G-CSF gene to chromosome

[56413] band 17q21. Oncogene Res. 1: 209-220, 1987.

[56414] 9203.

[56415] Wada, K.; Yokotani, N.; Hunter, C.; Doi, K.; Wenthold, R. J.; Shimasaki,

[56416] S.: Differential expression of two distinct forms of mRNA encoding

[56417] members of a dipeptidyl aminopeptidase family. Proc. Nat. Acad. Sci. 89:

[56418] 197-201, 1992.

[56419] 9204.

[56420] Wada, K.; Zimmerman, K. L.; Adamson, M. C.; Yokotani, N.; Wenthold,

[56421] R. J.; Kozak, C. A.: Genetic mapping of the mouse gene encoding dipeptidyl
[56422] aminopeptidase-like proteins. Mammalian Genome 4: 234-237, 1993.
[56423] 9205.

[56424] Yokotani, N.; Doi, K.; Wenthold, R. J.; Wada, K.: Non-conservation
[56425] of a catalytic residue in a depeptidyl aminopeptidase IV-related protein
[56426] encoded by a gene on human chromosome 7. Hum. Molec. Genet. 2: 1037-1039,
[56427] 1993.

[56428] 9206.

[56429] Chang, T.-M.; Neville, D. M., Jr.: Demonstration of diph-theria
[56430] toxin receptors on surface membranes from both toxin-sensitive and
[56431] toxin-resistant species. J. Biol. Chem. 253: 6866-6871, 1978.
[56432] 9207.

[56433] Creagan, R. P.; Chen, S.-H.; Ruddle, F. H.: Genetic analysis of

[56434] the cell surface: association of human chromosome 5 with sensitivity
[56435] to diphtheria toxin in mouse-human somatic cell hybrids. Proc. Nat.
[56436] Acad. Sci. 72: 2237-2241, 1975.

[56437] 9208.

[56438] Fen, Z.; Dhadly, M. S.; Yoshizumi, M.; Hilkert, R. J.; Quert-ermous,
[56439] T.; Eddy, R. L.; Shows, T. B.; Lee, M.-E.: Structural organi-zation
[56440] and chromosomal assignment of the gene encoding the human heparin-binding
[56441] epidermal growth factor-like growth factor/diphtheria toxin receptor. Biochemistry 32:
[56442] 7932-7938, 1993.

[56443] 9209.

[56444] George, D. L.; Francke, U.: Regional mapping of human genes for
[56445] hexosaminidase B and diphtheria toxin sensitivity on chromosome 5
[56446] using mouse X human hybrid cells. Somat. Cell Genet. 3: 629-638,

[56447] 1977.

[56448] 9210.

[56449] Gupta, R. S.; Siminovitch, L.: Isolation and characterization

[56450] of mutants of human diploid fibroblasts resistant to diphtheria toxin. Proc.

[56451] Nat. Acad. Sci. 75: 3337-3340, 1978.

[56452] 9211.

[56453] Hayes, H.; Kaneda, Y.; Uchida, T.; Okada, Y.: Regional assignment

[56454] of the gene for diphtheria toxin sensitivity using subchromosomal

[56455] fragments in microcell hybrids. Chromosoma 96: 26-32, 1987.

[56456] 9212.

[56457] Higashiyama, S.; Abraham, J. A.; Miller, J.; Fiddes, J. C.; Klagsbrun,

[56458] M.: A heparin-binding growth factor secreted by macrophage-like cells

[56459] that is related to EGF. Science 251: 936-939, 1991.

[56460] 9213.

[56461] Wang, Q.; Curren, M. E.; Splawski, I.; Burn, T. C.; Millholland,
[56462] J. M.; VanRaay, T. J.; Shen, J.; Timothy, K. W.; Vincent, G. M.; de
[56463] Jager, T.; Schwartz, P. J.; Towbin, J. A.; Moss, A. J.; Atkinson,
[56464] D. L.; Landes, G. M.; Connors, T. D.; Keating, M. T.: Positional
[56465] cloning of a novel potassium channel gene: KVLQT1 mutations cause
[56466] cardiac arrhythmias. Nature Genet. 12: 17-23, 1996.

[56467] 9214.

[56468] Kresse, H.; Rosthoj, S.; Quentin, E.; Hollmann, J.; Glossl, J.;

[56469] Okada, S.; Tonnesen, T.: Glycosaminoglycan-free small proteoglycan
[56470] core protein is secreted by fibroblasts from a patient with a syndrome
[56471] resembling progeroid. Am. J. Hum. Genet. 41: 436-453, 1987.

[56472] 9215.

[56473] Bellus, G. A.; Spector, E. B.; Speiser, P. W.; Weaver, C. A.; Garber,

[56474] A. T.; Bryke, C. R.; Israel, J.; Rosengren, S. S.; Webster, M. K.;
[56475] Donoghue, D. J.; Francomano, C. A.: Distinct missense mutations of
[56476] the FGFR3 lys650 codon modulate receptor kinase activation and the
[56477] severity of the skeletal dysplasia phenotype. Am. J. Hum. Genet. 67:
[56478] 1411-1421, 2000.

[56479] 9216.

[56480] Kondo, M.; Scherer, D. C.; Miyamoto, T.; King, A. G.; Akashi, K.;
[56481] Sugamura, K.; Weissman, I. L.: Cell-fate conversion of lymphoid-committed
[56482] progenitors by instructive actions of cytokines. Nature 407: 383-386,
[56483] 2000.

[56484] 9217.

[56485] Tavernier, J.; Devos, R.; Cornelis, S.; Tuypens, T.; Van der Heyden,
[56486] J.; Fiers, W.; Plaetinck, G.: A human high affinity interleukin-5

[56487] receptor (IL5R) is composed of an IL5-specific alpha chain and a beta

[56488] chain shared with the receptor for GM-CSF. Cell 66: 1175-1184, 1991.

[56489] 9218.

[56490] Prohl, C.; Pelzer, W.; Diekert, K.; Kmita, H.; Bedekovics, T.;

[56491] Kispal, G.; Lill, R.: The yeast mitochondrial carrier Leu5p and its

[56492] human homologue Graves' disease protein are required for accumulation

[56493] of coenzyme A in the matrix. Molec. Cell Biol. 21: 1089-1097, 2001.

[56494] 9219.

[56495] Rossi, E.; Zarrilli, R.; Zuffardi, O.: Regional assignment of

[56496] the gene coding for a human Graves' disease autoantigen to 10q21.3-q22.1. Hum.

[56497] Genet. 90: 653-654, 1993.

[56498] 9220.

[56499] Zarrilli, R.; Oates, E. L.; McBride, O. W.; Lerman, M. I.; Chan,

[56500] J. Y.; Santisteban, P.; Ursini, M. V.; Notkins, A. L.; Kohn, L.

D.

[56501] : Sequence and chromosomal assignment of a novel cDNA identified by

[56502] immunoscreening of a thyroid expression library: similarity to a family

[56503] of mitochondrial solute carrier proteins. Molec. Endocr. 3: 1498-1508,

[56504] 1989.

[56505] 9221.

[56506] Douglas, A. J.; Fox, M. F.; Abbott, C. M.; Hinks, L. J.; Sharpe,

[56507] G.; Povey, S.; Thompson, R. J.: Structure and chromosomal localization

[56508] of the human 2-prime,3-prime-cyclic nucleotide 3-prime-phosphodiesterase

[56509] gene. Ann. Hum. Genet. 56: 243-254, 1992.

[56510] 9222.

[56511] Douglas, A. J.; Fox, M. F.; Hinks, L. J.; Povey, S.; Thompson,

[56512] R. J.: Localization of the myelin specific enzyme 2-prime,3-prime-cyclic

[56513] nucleotide-3-prime-phosphohydrolase to 17q21.

(Abstract) Cytogenet.

[56514] Cell Genet. 58: 2004 only, 1991.

[56515] 9223.

[56516] Kurihara, T.; Monoh, K.; Sakimura, K.; Takahashi, Y.: Al-
ternative

[56517] splicing of mouse brain 2-prime,3-prime-cyclic nu-
cleotide 3-prime-phosphodiesterase

[56518] mRNA. Biochem. Biophys. Res. Commun. 170:
1074-1081, 1990.

[56519] 9224.

[56520] Monoh, K.; Kurihara, T.; Takahashi, Y.; Ichikawa, T.; Ku-
manishi,

[56521] T.; Hayashi, S.; Minoshima, S.; Shimizu, N.: Structure, ex-
pression

[56522] and chromosomal localization of the gene encoding hu-
man 2-prime,3-prime-cyclic-nucleotide

[56523] 3-prime-phosphodiesterase. Gene 129: 297-301, 1993.

[56524] 9225.

[56525] Stepanova, L.; Leng, X.; Parker, S. B.; Harper, J. W.: Mam-
malian

[56526] p50(Cdc37) is a protein kinase-targeting subunit of

Hsp90 that binds
[56527] and stabilizes Cdk4. Genes Dev. 10: 1491-1502, 1996.

[56528] 9226.

[56529] Wolfel, T.; Hauer, M.; Schneider, J.; Serrano, M.; Wolfel, C.;

[56530] Klehmann-Hieb, E.; De Plaen, E.; Hankeln, T.; Meyer zum Buschenfelde,
[56531] K.-H.; Beach, D.: A p16(INK4a)-insensitive CDK4 mutant targeted by
[56532] cytolytic T lymphocytes in a human melanoma. Science 269: 1281-1284,
[56533] 1995.

[56534] 9227.

[56535] Zuo, L.; Weger, J.; Yang, Q.; Goldstein, A. M.; Tucker, M. A.;
[56536] Walker, G. J.; Hayward, N.; Dracopoli, N. C.: Germline mu-tations
[56537] in the p16(INK4a) binding domain of CDK4 in familial melanoma. Nature
[56538] Genet. 12: 97-99, 1996.

[56539] 9228.

[56540] Bernier, L.; Colman, D. R.; D'Eustachio, P.: Chromosomal

locations
[56541] of genes encoding 2-prime,3-prime cyclic nucleotide 3-prime-phosphodiesterase
[56542] and glial fibrillary acidic protein in the mouse. J. Neurosci. Res. 20:
[56543] 497-504, 1988.

[56544] 9229.

[56545] Bifulco, M.; Laezza, C.; Stingo, S.; Wolff, J.: 2-prime,3-prime-cyclic
[56546] nucleotide 3-prime-phosphodiesterase: a membrane-bound, microtubule-associated
[56547] protein and membrane anchor for tubulin. Proc. Nat. Acad. Sci. 99:
[56548] 1807-1812, 2002.

[56549] 9230.

[56550] Sprinkle, T. J.; Kouri, R. E.; Fain, P. D.; Stoming, T. A.; Whitney,
[56551] J. B., III: Chromosomal mapping of the human CNP gene using a meiotic
[56552] crossover DNA panel, PCR, and allele-specific probes. Ge-nomics 16:
[56553] 542-545, 1993.

[56554] 9231.

[56555] Sprinkle, T. J.; Lanclos, K. D.; Lapp, D. F.: Assignment of the

[56556] human 2-prime,3-prime-cyclic nucleotide 3-prime-phosphohydrolase gene

[56557] to chromosome 17. Genomics 13: 877-880, 1992.

[56558] 9232.

[56559] Vogel, U. S.; Thompson, R. J.: Molecular structure, local-ization,

[56560] and possible functions of the myelin-associated enzyme 2-prime,3-prime-cyclic

[56561] nucleotide 3-prime-phosphodiesterase. J. Neurochem. 50: 1667-1677,

[56562] 1988.

[56563] 9233.

[56564] Bibb, J. A.; Chen, J.; Taylor, J. R.; Svenningsson, P.; Nishi,

[56565] A.; Snyder, G. L.; Yan, Z.; Sagawa, Z. K.; Ouimet, C. C.; Nairn, A.

[56566] C.; Nestler, E. J.; Greengard, P.: Effects of chronic expo-sure to

[56567] cocaine are regulated by the neuronal protein Cdk5. Na-

ture 410:
[56568] 376-380, 2001.

[56569] 9234.

[56570] Demetrick, D. J.; Matsumoto, S.; Hannon, G. J.; Okamoto, K.; Xiong,
[56571] Y.; Zhang, H.; Beach, D. H.: Chromosomal mapping of the genes for
[56572] the human cell cycle proteins cyclin C (CCNC), cyclin E (CCNE), p21
[56573] (CDKN1) and KAP (CDKN3). Cytogenet. Cell Genet. 69: 190-192, 1995.

[56574] 9235.

[56575] Gyuris, J.; Golemis, E.; Chertkov, H.; Brent, R.: Cdi1, a human
[56576] G1 and S phase protein phosphatase that associates with Cdk2. Cell 75:
[56577] 791-803, 1993.

[56578] 9236.

[56579] Hannon, G. J.; Casso, D.; Beach, D.: KAP: a dual specificity phosphatase
[56580] that interacts with cyclin-dependent kinases. Proc. Nat.

Acad. Sci. 91:
[56581] 1731-1735, 1994.

[56582] 9237.

[56583] Inaba, T.; Matsushime, H.; Valentine, M.; Roussel, M. F.; Sherr,
[56584] C. J.; Look, A. T.: Genomic organization, chromosomal localization,
[56585] and independent expression of human cyclin D genes. Genomics 13:
[56586] 565-574, 1992.

[56587] 9238.

[56588] Kim, H. A.; Pomeroy, S. L.; Whoriskey, W.; Pawlitzky, I.; Benowitz,
[56589] L. I.; Sicinski, P.; Stiles, C. D.; Roberts, T. M.: A developmentally
[56590] regulated switch directs regenerative growth of Schwann cells through
[56591] cyclin D1. Neuron 26: 405-416, 2000.

[56592] 9239.

[56593] Xiong, Y.; Menninger, J.; Beach, D.; Ward, D. C.: Molecular cloning

[56594] and chromosomal mapping of CCND genes encoding human D-type cyclins. Genomics 13:

[56595] 575-584, 1992.

[56596] 9240.

[56597] Stamper, C. C.; Zhang, Y.; Tobin, J. F.; Erbe, D. V.; Ikemizu,

[56598] S.; Davis, S. J.; Stahl, M. L.; Seehra, J.; Somers, W. S.; Mosyak,

[56599] L.: Crystal structure of the B7-1/CTLA-4 complex that inhibits human

[56600] immune responses. Nature 410: 608-611, 2001. Note: Erratum: Nature

[56601] 411: 617 only, 2001.

[56602] 9241.

[56603] Allenspach, E. J.; Cullinan, P.; Tong, J.; Tang, Q.; Tesciuba,

[56604] A. G.; Cannon, J. L.; Takahashi, S. M.; Morgan, R.; Burkhardt, J.

[56605] K.; Sperling, A. I.: ERM-dependent movement of CD43 defines a novel

[56606] protein complex distal to the immunological synapse. Immunity 15:

[56607] 739-750, 2001.

[56608] 9242.

[56609] Bonilha, V. L.; Rodriguez-Boulan, E.: Polarity and develop-
mental

[56610] regulation of two PDZ proteins in the retinal pigment ep-
ithelium. Invest.

[56611] Ophthal. Vis. Sci. 42: 3274-3282, 2001.

[56612] 9243.

[56613] Brunet, J.-F.; Dosseto, M.; Denizot, F.; Mattei, M.-G.;
Clark,

[56614] W. R.; Haqqi, T. M.; Ferrier, P.; Nabholz, M.; Schmitt-Ver-
hulst, A.-M.;

[56615] Luciani, M.-F.; Golstein, P.: The inducible cytotoxic T-
lymphocyte-associated

[56616] gene transcript CTLA-1 sequence and gene localization to
mouse chromosome

[56617] 14. Nature 322: 268-271, 1986.

[56618] 9244.

[56619] Crosby, J. L.; Bleackley, R. C.; Nadeau, J. H.: A complex of
serine

[56620] protease genes expressed preferentially in cytotoxic T-

lymphocytes

[56621] is closely linked to the T-cell receptor alpha- and delta-chain genes

[56622] on mouse chromosome 14. Genomics 6: 252-259, 1990.

[56623] 9245.

[56624] Dahl, C. A.; Bach, F. H.; Chan, W.; Huebner, K.; Russo, G.; Croce,

[56625] C. M.; Herfurth, T.; Cairns, J. S.: Isolation of a cDNA clone encoding

[56626] a novel form of granzyme B from human NK cells and mapping to chromosome

[56627] 14. Hum. Genet. 84: 465-470, 1990.

[56628] 9246.

[56629] Okajima, T.; Fukumoto, S.; Furukawa, K.; Urano, T.; Fu-

rukawa, K.

[56630] : Molecular basis for the progeroid variant of Ehlers-Dan-los syndrome:

[56631] identification and characterization of two mutations in galactosyltransferase

[56632] I gene. J. Biol. Chem. 274: 28841-28844, 1999.

[56633] 9247.

[56634] Okajima, T.; Yoshida, K.; Kondo, T.; Furukawa, K.: Human homolog
[56635] of Caenorhabditis elegans sqv-3 gene is galactosyltransferase I involved
[56636] in the biosynthesis of the glycosaminoglycan-protein linkage region
[56637] of proteoglycans. J. Biol. Chem. 274: 22915-22918, 1999.

[56638] 9248.

[56639] Trueb, B.; Winterhalter, K. H.: Type VI collagen is composed
[56640] of a 200 kD subunit and two 140 kD subunits. EMBO J. 5: 2815-2819,
[56641] 1986.

[56642] 9249.

[56643] Weil, D.; Mattei, M.-G.; Passage, E.; Van Cong, N.; Pribula-Conway,
[56644] D.; Mann, K.; Deutzmann, R.; Timpl, R.; Chu, M.-L.: Cloning and chromosomal
[56645] localization of human genes encoding the three chains of type VI collagen. Am.
[56646] J. Hum. Genet. 42: 435-445, 1988.

[56647] 9250.

[56648] Muragaki, Y.; Mattei, M.-G.; Yamaguchi, N.; Olsen, B. R.; Ninomiya,

[56649] Y.: The complete primary structure of the human alpha-1(VIII) chain

[56650] and assignment of its gene (COL8A1) to chromosome 3. Europ. J. Biochem. 197:

[56651] 615-622, 1991.

[56652] 9251.

[56653] Muragaki, Y.; Shiota, C.; Inoue, M.; Ooshima, A.; Olsen, B. R.;

[56654] Ninomiya, Y.: Alpha-1(VIII)-collagen gene transcripts encode a short-chain

[56655] collagen polypeptide and are expressed by various epithelial, endothelial

[56656] and mesenchymal cells in newborn mouse tissues. Europ. J. Biochem. 207:

[56657] 895-902, 1992.

[56658] 9252.

[56659] Schafer, B. W.; Wicki, R.; Engelkamp, D.; Mattei, M.-G.; Heizmann,

[56660] C. W.: Isolation of a YAC clone covering a cluster of nine S100 genes
[56661] on human chromosome 1q21: rationale for a new nomen-clature of the
[56662] S100 calcium-binding protein family. Genomics 25: 638-643, 1995.
[56663] 9253.

[56664] van Heyningen, V.; Emslie, E.; Dorin, J. R.: Related calcium binding
[56665] proteins map to the same sub-region of chromosome 1q and to an extended
[56666] region of synteny on mouse chromosome 3. (Abstract) Cytogenet.
[56667] Cell Genet. 51: 1095, 1989.

[56668] 9254.

[56669] van Heyningen, V.; Emslie, E.; Dorin, J. R.: Related calcium binding
[56670] proteins map to the same sub-region of chromosome 1q and to an extended
[56671] region of synteny on mouse chromosome 3. (Abstract) Cytogenet. Cell
[56672] Genet. 51: 1095 only, 1989.

[56673] 9255.

[56674] Buonavista, N.; Balzano, C.; Pontarotti, P.; Le Paslier, D.; Golstein,

[56675] P.: Molecular linkage of the human CTLA4 and CD28 Ig-superfamily

[56676] genes in yeast artificial chromosomes. Genomics 13: 856-861, 1992.

[56677] 9256.

[56678] Howard, T. A.; Rochelle, J. M.; Seldin, M. F.: Cd28 and Ctla-4,

[56679] two related members of the Ig supergene family, are tightly linked

[56680] on proximal mouse chromosome 1. Immunogenetics 33: 74-76, 1991.

[56681] 9257.

[56682] Albertson, D. G.; Ylstra, B.; Segraves, R.; Collins, C.; Dair-kee,

[56683] S. H.; Kowbel, D.; Kuo, W.- L.; Gray, J. W.; Pinkel, D.: Quantitative

[56684] mapping of amplicon structure by array CGH identifies CYP24 as a candidate

[56685] oncogene. Nature Genet. 25: 144-146, 2000.

[56686] 9258.

[56687] Chen, K.-S.; Prahl, J. M.; DeLuca, H. F.: Isolation and ex-
pression

[56688] of human 1,25-dihydroxyvitamin D3 24-hydroxylase
cDNA. Proc. Nat.

[56689] Acad. Sci. 90: 4543-4547, 1993.

[56690] 9259.

[56691] Hahn, C. N.; Baker, E.; Laslo, P.; May, B. K.; Omdahl, J. L.;
Sutherland,

[56692] G. R.: Localization of the human vitamin D 24-hydroxylase
gene (CYP24)

[56693] to chromosome 20q13.2-q13.3. Cytogenet. Cell Genet.
62: 192-193,

[56694] 1993.

[56695] 9260.

[56696] Labuda, M.; Lemieux, N.; Tihy, F.; Prinster, C.; Glorieux, F.
H.

[56697] : Human 25-hydroxyvitamin D 24-hydroxylase cy-
tochrome P450 subunit

[56698] maps to a different chromosomal location than that of
pseudovitamin

[56699] D-deficient rickets. J. Bone Miner. Res. 8: 1397-1406, 1993.

[56700] 9261.

[56701] Malas, S.; Peters, J.; Abbott, C.: The genes for endothelin 3,

[56702] vitamin D 24-hydroxylase, and melanocortin 3 receptor map to distal

[56703] mouse chromosome 2, in the region of conserved synteny with human

[56704] chromosome 20. Mammalian Genome 5: 577-579, 1994.

[56705] 9262.

[56706] Ohyama, Y.; Noshiro, M.; Okuda, K.: Cloning and expres-sion of

[56707] cDNA encoding 25-hydroxyvitamin D(3) 24-hydroxylase. FEBS Lett. 278:

[56708] 195-198, 1991.

[56709] 9263.

[56710] Mendel, D. B.; Khavari, P. A.; Conley, P. B.; Graves, M. K.; Hansen,

[56711] L. P.; Admon, A.; Crabtree, G. R.: Characterization of a co-factor

[56712] that regulates dimerization of a mammalian homeodomain protein. Science 254:
[56713] 1762-1767, 1991.

[56714] 9264.

[56715] Hogenesch, J. B.; Chan, W. K.; Jackiw, V. H.; Brown, R. C.; Gu,
[56716] Y.-Z.; Pray-Grant, M.; Perdew, G. H.; Bradfield, C. A.: Characterization
[56717] of a subset of the basic-helix-loop-helix-PAS superfamily that interacts
[56718] with components of the dioxin signaling pathway. J. Biol. Chem. 272:
[56719] 8581-8593, 1997.

[56720] 9265.

[56721] Wang, G. L.; Jiang, B.-H.; Rue, E. A.; Semenza, G. L.: Hypoxia-inducible
[56722] factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by
[56723] cellular O(2) tension. Proc. Nat. Acad. Sci. 92: 5510-5514, 1995.

[56724] 9266.

[56725] Neerman-Arbez, M.; de Moerloose, P.; Bridel, C.; Hons-
berger, A.;
[56726] Schonborner, A.; Rossier, C.; Peerlinck, K.; Claeyssens, S.;
Di Michele,
[56727] D.; d'Oiron, R.; Dreyfus, M.; Laubriat-Bianchin, M.; Dieval,
J.; Antonarakis,
[56728] S. E.; Morris, M. A.: Mutations in the fibrinogen A-alpha
gene account
[56729] for the majority of cases of congenital afibrinogenemia.
Blood 96:
[56730] 149-152, 2000.

[56731] 9267.

[56732] Le Beau, M. M.; Lemons, R. S.; Carrino, J. J.; Pettenati, M.
J.;
[56733] Souza, L. M.; Diaz, M. O.; Rowley, J. D.: Chromosomal lo-
calization
[56734] of the human G-CSF gene to 17q11 proximal to the
breakpoint of the
[56735] t(15;17) in acute promyelocytic leukemia. Leukemia 1:
795-799, 1987.
[56736] 9268.

[56737] Posteraro, P.; Sorvillo, S.; Gagnoux-Palacios, L.; Angelo,

C.;

[56738] Paradisi, M.; Meneguzzi, G.; Castiglia, D.; Zambruno, G.: Compound

[56739] heterozygosity for an out-of-frame deletion and a splice site mutation

[56740] in the LAMB3 gene causes nonlethal junctional epidermolysis bullosa. Biochem.

[56741] Biophys. Res. Commun. 243: 758-764, 1998.

[56742] 9269.

[56743] Pulkkinen, L.; Christiano, A. M.; Gerecke, D.; Wagman, D. W.;

[56744] Burgeson, R. E.; Pittelkow, M. R.; Uitto, J.: A homozygous nonsense

[56745] mutation in the beta-3 chain gene of laminin 5 (LAMB3) in Herlitz

[56746] junctional epidermolysis bullosa. Genomics 24: 357-360, 1994.

[56747] 9270.

[56748] Pulkkinen, L.; Gerecke, D. R.; Christiano, A. M.; Wagman, D. W.;

[56749] Burgeson, R. E.; Uitto, J.: Cloning of the beta-3 chain gene (LAMB3)

[56750] of human laminin 5, a candidate gene in junctional epidermolysis bullosa. Genomics 25:
[56751] 192-198, 1995.

[56752] 9271.

[56753] Pulkkinen, L.; Meneguzzi, G.; McGrath, J. A.; Xu, Y.; Blanchet-Bardon,
[56754] C.; Ortonne, J.-P.; Christiano, A. M.; Uitto, J.: Predominance of
[56755] the recurrent mutation R635X in the LAMB3 gene in European patients
[56756] with Herlitz junctional epidermolysis bullosa has implications for
[56757] mutation detection strategy. J. Invest. Derm. 109:
232-237, 1997.
[56758] 9272.

[56759] Pulkkinen, L.; Uitto, J.: Heterozygosity for premature termination
[56760] codon mutations in LAMB3 in siblings with non-lethal junctional epidermolysis
[56761] bullosa. (Letter) J. Invest. Derm. 111: 1244-1245, 1998.

[56762] 9273.

[56763] Robbins, P. B.; Lin, Q.; Goodnough, J. B.; Tian, H.; Chen, X.;

[56764] Khavari, P. A.: In vivo restoration of laminin 5 beta-3 ex-

[56765] pression and function in junctional epidermolysis bullosa. Proc. Nat. Acad.

[56766] Sci. 98: 5193-5198, 2001.

[56767] 9274.

[56768] Takizawa, Y.; Pulkkinen, L.; Shimizu, H.; Lin, H.; Hagiwara, S.;

[56769] Nishikawa, T.; Uitto, J.: Maternal uniparental meroisodis-

[56770] omy in the LAMB3 region of chromosome 1 results in lethal junctional epidermolysis

[56771] bullosa. J. Invest. Derm. 110: 828-831, 1998.

[56772] 9275.

[56773] Takizawa, Y.; Shimizu, H.; Pulkkinen, L.; Hiraoka, Y.; Mc-

[56774] Grath, J. A.; Suzumori, K.; Aiso, S.; Uitto, J.; Nishikawa, T.: Novel mutations

[56775] in the LAMB3 gene shared by two Japanese unrelated families with Herlitz

[56776] junctional epidermolysis bullosa, and their application for prenatal

[56777] testing. J. Invest. Derm. 110: 174-178, 1998.

[56778] 9276.

[56779] Haaparanta, T.; Uitto, J.; Ruoslahti, E.; Engvall, E.: Molecular

[56780] cloning of the cDNA encoding human laminin A chain. Matrix 11: 151-160,

[56781] 1991.

[56782] 9277.

[56783] Chu, M.-L.; de Wet, W.; Bernard, M.; Ding, J.-F.; Morabito, M.;

[56784] Myers, J.; Williams, C.; Ramirez, F.: Human pro-alpha-1(I) collagen

[56785] gene structure reveals evolutionary conservation of a pattern of introns

[56786] and exons. Nature 310: 337-340, 1984.

[56787] 9278.

[56788] Cole, W. G.; Chan, D.; Chambers, G. W.; Walker, I. D.; Bateman,

[56789] J. F.: Deletion of 24 amino acids from the pro-alpha-1(I)

chain of

[56790] type I procollagen in a patient with the Ehlers-Danlos syn-
drome type

[56791] VII. J. Biol. Chem. 261: 5496-5503, 1986.

[56792] 9279.

[56793] Gallagher, P. G.; Forget, B. G.: Structure, organization, and

[56794] expression of the band 7.2b gene, a candidate gene for
hereditary

[56795] hydrocytosis. J. Biol. Chem. 270: 26358-26363, 1995.

[56796] 9280.

[56797] Gallagher, P. G.; Romana, M.; Lieman, J. H.; Ward, D. C.:
cDNA

[56798] structure, tissue-specific expression, and chromosomal
localization

[56799] of the murine band 7.2b gene. Blood 86: 359-365, 1995.

[56800] 9281.

[56801] Gallagher, P. G.; Upender, M.; Ward, D. C.; Forget, B. G.:
The

[56802] gene for human erythrocyte membrane protein band 7.2
(EPB72) maps

[56803] to 9q33-q34 centromeric to the Philadelphia chromosome

translocation

[56804] breakpoint region. Genomics 18: 167-169, 1993.

[56805] 9282.

[56806] Hiebl-Dirschmied, C. M.; Entler, B.; Glotzmann, C.; Mau-
rer-Fogy,

[56807] I.; Stratowa, C.; Prohaska, R.: Cloning and nucleotide se-
quence of

[56808] cDNA encoding human erythrocyte band 7 integral mem-
brane protein. Biochim.

[56809] Biophys. Acta 1090: 123-124, 1991.

[56810] 9283.

[56811] Pilz, A.; Prohaska, R.; Peters, J.; Abbott, C.: Genetic linkage

[56812] analysis of the Ak1, Col5a1, Epb7.2, Fpgs, Grp78, Pbx3,
and Notch1

[56813] genes in the region of mouse chromosome 2 homologous
to human chromosome

[56814] 9q. Genomics 21: 104-109, 1994.

[56815] 9284.

[56816] Unfried, I.; Entler, B.; Prohaska, R.: The organization of the

[56817] gene (EPB72) encoding the human erythrocyte band 7 in-

tegral membrane
[56818] protein (protein 7.2b). Genomics 30: 521-528, 1995.

[56819] 9285.

[56820] Westberg, J. A.; Entler, B.; Prohaska, R.; Schroder, J. P.: The

[56821] gene coding for erythrocyte protein band 7.2b (EPB72) is located in
[56822] band q34.1 of human chromosome 9. Cytogenet. Cell Genet. 63: 241-243,
[56823] 1993.

[56824] 9286.

[56825] Zhu, Y.; Paszty, C.; Turetsky, T.; Tsai, S.; Kuypers, F. A.; Lee,
[56826] G.; Cooper, P.; Gallagher, P. G.; Stevens, M. E.; Rubin, E.; Mohandas,
[56827] N.; Mentzer, W. C.: Stomatocytosis is absent in 'stom-atin'-deficient
[56828] murine red blood cells. Blood 93: 2404-2410, 1999.

[56829] 9287.

[56830] de la Chapelle, A.; Sistonen, P.; Lehvaslaiho, H.; Ikkala, E.;

[56831] Juvonen, E.: Familial erythrocytosis genetically linked to

erythropoietin
[56832] receptor gene. Lancet 341: 82-84, 1993.

[56833] 9288.

[56834] Juvonen, E.; Ikkala, E.; Fyhrquist, F.; Ruutu, T.: Autosomal

[56835] dominant erythrocytosis caused by increased sensitivity to erythropoietin. Blood 78:
[56836] 3066-3069, 1991.

[56837] 9289.

[56838] Neerman-Arbez, M.; Honsberger, A.; Antonarakis, S. E.; Morris,
[56839] M. A.: Deletion of the fibrogen (sic) alpha-chain gene (FGA) causes
[56840] congenital afibrogenemia (sic). J. Clin. Invest. 103: 215-218, 1999.
[56841] 9290.

[56842] Olaisen, B.; Teisberg, P.; Gedde-Dahl, T., Jr.: Fibrinogen gamma
[56843] chain locus is on chromosome 4 in man. Hum. Genet. 61: 24-26, 1982.
[56844] 9291.

[56845] Qureshi, G. D.; Evans, H. J.; Vennart, R. M.; Magnant, J. P.;

[56846] Sabau, J. M.; Willoughby, J. B.; Koehn, J. A.: Fibrinogen White Marsh--a
[56847] new human fibrinogen variant with alpha chain defect. (Abstract) Clin.
[56848] Res. 31: 321A only, 1983.

[56849] 9292.

[56850] Ratnoff, O. D.; Bennett, B.: The genetics of hereditary dis-orders
[56851] of blood coagulation. Science 179: 1291-1298, 1973.

[56852] 9293.

[56853] Reber, P.; Furlan, M.; Beck, E. A.; Barbui, T.: Fibrinogen Bergamo
[56854] III and fibrinogen Torino: two further variants with heredi-tary molecular
[56855] defects in fibrinopeptide A. Thromb. Res. 46: 163-167, 1987.
[56856] 9294.

[56857] Reber, P.; Furlan, M.; Beck, E. A.; Finazzi, G.; Buelli, M.; Barbui,
[56858] T.: Fibrinogen Bergamo I (A alpha 16 arg-to-cys): suscep-tibility

[56859] towards thrombin following aminoethylation, methylation or carboxamidomethylation

[56860] of cysteine residues. Thromb. Haemost. 54: 390-393, 1985.

[56861] 9295.

[56862] Rupp, C.; Beck, E. A.: Congenital dysfibrinogenemia.In: Beck,

[56863] E. A.; Furlan, M.: Variants of Human Fibrinogen. Berne: Hans Huber

[56864] (pub.) 1984. Pp. 65-130.

[56865] 9296.

[56866] Samama, M.; Soria, J.; Soria, C.; Bousser, J.: Dysfibrinogenemie

[56867] congenitale et familiale sans tendance hemorragique. Nouv. Rev. Franc.

[56868] Hemat. 9: 817-832, 1969.

[56869] 9297.

[56870] Siebenlist, K. R.; Prchal, J. T.; Mosesson, M. W.: Fibrinogen

[56871] Birmingham: a heterozygous dysfibrinogenemia (A alpha 16 arg-to-his)

[56872] containing heterodimeric molecules. Blood 71: 613-618,

1988.

[56873] 9298.

[56874] Soria, J.; Soria, C.; Bertrand, O.; Dunn, F.; Drouet, L.; Caen,

[56875] J. P.: Plasminogen Paris I: congenital abnormal plasmino-
gen and its

[56876] incidence in thrombosis. Thromb. Res. 32: 229-238,
1983.

[56877] 9299.

[56878] Soria, J.; Soria, C.; Hedner, U.; Nilsson, I. M.; Bergqvist, D.;

[56879] Samama, M.: Episodes of increased fibronectin level ob-
served in a

[56880] patient suffering from recurrent thrombosis related to
congenital

[56881] hypodysfibrinogenaemia (fibrinogen Malmoe). Brit. J.
Haemat. 61:

[56882] 727-738, 1985.

[56883] 9300.

[56884] Soria, J.; Soria, C.; Samama, M.; Poirot, E.; Kling, C.: Fib-
rinogen

[56885] Troyes-fibrinogen Metz: two new cases of congenital dys-
fibrinogenemia. Thromb.

[56886] Diath. Haemorrh. 27: 619-633, 1972.

[56887] 9301.

[56888] Southan, C.; Henschen, A.; Lottspeich, F.: The search for molecular

[56889] defects in abnormal fibrinogens.In: Henschen, A.; Graeff, H.; Lottspeich,

[56890] F.: Fibrinogen--Recent Biochemical and Medical Aspects. Berlin:

[56891] W. de Gruyter (pub.) 1982. Pp. 153-166.

[56892] 9302.

[56893] Southan, C.; Lane, D. A.; Knight, I.; Ireland, H.; Bottomley,

[56894] J.: Fibrinogen Manchester: detection of a heterozygous phenotype

[56895] in the intraplatelet pool. Biochem. J. 229: 723-730, 1985.

[56896] 9303.

[56897] Streiff, F.; Alexandre, P.; Vigneron, C.; Soria, J.; Soria, C.;

[56898] Mester, L.: Un nouveau cas d'anomalie constitutionnelle et familiale

[56899] du fibrinogene sans diathese hemorragique. Thromb. Diath. Haemorrh. 26:

[56900] 565-576, 1971.

[56901] 9304.

[56902] Suh, T. T.; Holmback, K.; Jensen, N. J.; Daugherty, C. C.; Small,

[56903] K.; Simon, D. I.; Potter, S.; Degen, J. L.: Resolution of spontaneous

[56904] bleeding events but failure of pregnancy in fibrinogen-deficient mice. Genes

[56905] Dev. 9: 2020-2033, 1995.

[56906] 9305.

[56907] Thomas, A.; Lamlum, H.; Humphries, S.; Green, F.: Linkage disequilibrium

[56908] across the fibrinogen locus as shown by five genetic polymorphisms,

[56909] G/A(-455) (HaeIII), C/T(-148) (HindIII/AluI), T/G(+1689) (AvaII),

[56910] and BclI (beta-fibrinogen) and TaqI (alpha-fibrinogen), and their

[56911] detection by PCR. Hum. Mutat. 3: 79-81, 1994.

[56912] 9306.

[56913] Uemichi, T.; Liepnieks, J. J.; Alexander, F.; Benson, M. D.:

[56914] The molecular basis of renal amyloidosis in Irish-
American and Polish-Canadian
[56915] kindreds. Quart. J. Med. 89: 745-750, 1996.

[56916] 9307.

[56917] Uemichi, T.; Liepnieks, J. J.; Benson, M. D.: Fibrinogen In-
dianapolis:
[56918] a fibrinogen A-alpha chain associated with hereditary
amyloidosis.
[56919] (Abstract) Clin. Res. 41: 133 only, 1993.

[56920] 9308.

[56921] Uemichi, T.; Liepnieks, J. J.; Benson, M. D.: Hereditary re-
nal
[56922] amyloidosis with a novel variant fibrinogen. J. Clin. Invest.
93:
[56923] 731-736, 1994.

[56924] 9309.

[56925] Uemichi, T.; Liepnieks, J. J.; Yamada, T.; Gertz, M. A.;
Bang,
[56926] N.; Benson, M. D.: A frame shift mutation in the fibrinogen
A-alpha
[56927] chain gene in a kindred with renal amyloidosis. Blood 87:

4197-4203,
[56928] 1996.

[56929] 9310.

[56930] Uemichi, T.; Liepnieks, J. J.; Yamada, T.; Gertz, M. A. Bang,

[56931] N.; Benson, M. D.: A frame shift mutation in the fibrinogen A-alpha

[56932] chain gene in a kindred with renal amyloidosis. Blood 87: 4197-4203,

[56933] 1996.

[56934] 9311.

[56935] Uzan, G.; Besmond, C.; Kahn, A.; Marguerie, G.: Cell free translation

[56936] of messenger RNA for human fibrinogen. Biochem. Int. 4: 271-278,

[56937] 1982.

[56938]

[56939] 100. Uzan, G.; Courtois, G.; Besmond, C.; Frain, M.; Sala-Trepat,

[56940] J.; Kahn, A.; Marguerie, G.: Analysis of fibrinogen genes in patients

[56941] with congenital afibrinogenemia. Biochem. Biophys. Res.

Commun. 120:
[56942] 376-383, 1984.

[56943]

[56944] 101. Verhaeghe, R.; Verstraete, M.; Vermylen, J.;
Vermylen, C.: Fibrinogen
[56945] 'Leuven', another genetic variant. Brit. J. Haemat. 26:
421-434,
[56946] 1974.

[56947]

[56948] 102. Verstraete, M.: Discussion. Thromb. Diath. Haem-
orrh. 39 (suppl.):
[56949] 334-337, 1970.

[56950]

[56951] 103. Vogel, F.; Motulsky, A. G.: Human Genetics: Problems
and Approaches.
[56952] New York: Springer-Verlag (pub.) 1979. Pp. 262 only.

[56953]

[56954] 104. von Felton, A.; Duckert, F.; Frick, P. G.: Familial dis-
turbance
[56955] of fibrin monomer aggregation. Brit. J. Haemat. 12:

667-677, 1966.

[56956]

[56957] 105. Wehinger, H.; Klinge, O.; Alexandrakis, E.; Schur-
mann, J.; Witt,
[56958] J.; Seydewitz, H. H.: Hereditary hypofibrinogenemia with
fibrinogen
[56959] storage in the liver. Europ. J. Pediat. 141: 109-112, 1983.

[56960]

[56961] 106. Winckelmann, G.; Augustin, R.; Bandilla, K.: Congeni-
tal dysfibrinogenemia:
[56962] report of a new family (fibrinogen Wiesbaden).In: Ab-
stracts, International
[56963] Society on Thrombosis and Haemostasis. Oslo: Villco
Trykkeri (pub.)
[56964] 1971. Pp. 64 only.

[56965]

[56966] 107. Xiao, Q.; Danton, M. J. S.; Witte, D. P.; Kowala, M. C.;
Valentine,
[56967] M. T.; Degen, J. L.: Fibrinogen deficiency is compatible
with the
[56968] development of atherosclerosis in mice. J. Clin. Invest.

101: 1184-1194,
[56969] 1998.

[56970]

[56971] 108. Yoshida, N.; Okuma, M.; Hirata, H.; Matsuda, M.; Ya-
mazumi, K.;
[56972] Asakura, S.: Fibrinogen Kyoto II, a new congenitally ab-
normal molecule,
[56973] characterized by the replacement of A-alpha proline-18
by leucine. Blood 78:
[56974] 149-153, 1991.

[56975]

[56976] 109. Zietz, B. H.; Scott, J. L.: An inherited defect in fib-
rinogen
[56977] polymerization: fibrinogen Los Angeles. (Abstract) Clin.
Res. 18:
[56978] 179 only, 1970.

[56979] 9312.

[56980] Alward, W. L. M.; Semina, E. V.; Kalenak, J. W.; Heon, E.;
Sheth,
[56981] B. P.; Stone, E. M.; Murray, J. C.: Autosomal dominant iris
hypoplasia

[56982] is caused by a mutation in the Rieger syndrome (RIEG/PITX2) gene. Am.
[56983] J. Ophthal. 125: 98-100, 1998.

[56984] 9313.

[56985] Heon, E.; Sheth, B. P.; Kalenak, J. W.; Sunden, S. L. F.; Streb,
[56986] L. M.; Taylor, C. M.; Alward, W. L. M.; Sheffield, V. C.; Stone, E.
[56987] M.: Linkage of autosomal dominant iris hypoplasia to the region of
[56988] the Rieger syndrome locus (4q25). Hum. Molec. Genet. 4: 1435-1439,
[56989] 1995.

[56990] 9314.

[56991] Kulak, S. C.; Kozlowski, K.; Semina, E. V.; Pearce, W. G.; Walter,
[56992] M. A.: Mutation in the RIEG1 gene in patients with iridogo-niodysgenesis
[56993] syndrome. Hum. Molec. Genet. 7: 1113-1117, 1998.

[56994] 9315.

[56995] Walter, M. A.; Mirzayans, F.; Mears, A. J.; Hickey, K.;

Pearce,
[56996] W. G.: Autosomal-dominant iridogoniodysgenesis and Axenfeld-Rieger
[56997] syndrome are genetically distinct. Ophthalmology 103: 1907-1915,
[56998] 1996.

[56999] 9316.

[57000] Metcalf, D.: The granulocyte-macrophage colony-stimulating factors. Science 229:
[57001] 16-22, 1985.

[57002] 9317.

[57003] Nagata, S.; Tsuchiya, M.; Asano, S.; Kaziro, Y.; Yamazaki, T.;
[57004] Yamamoto, O.; Hirata, Y.; Kubota, N.; Oheda, M.; Nomura, H.; Ono,
[57005] M.: Molecular cloning and expression of cDNA for the human granulocyte
[57006] colony-stimulating factor. Nature 319: 415-418, 1986.

[57007] 9318.

[57008] Nagata, S.; Tsuchiya, M.; Asano, S.; Yamamoto, O.; Hirata, Y.;

[57009] Kubota, N.; Oheda, M.; Nomura, H.; Yamazaki, T.: The chromosomal
[57010] gene structure and two mRNAs for human granulocyte colony-stimulating
[57011] factor. EMBO J. 5: 575-581, 1986.

[57012] 9319.

[57013] Petit, I.; Szyper-Kravitz, M.; Nagler, A.; Lahav, M.; Peled, A.;
[57014] Habler, L.; Ponomaryov, T.; Taichman, R. S.; Arenzana-Seisdedos, F.;
[57015] Fujii, N.; Sandbank, J.; Zipori, D.; Lapidot, T.: G-CSF in-duces stem
[57016] cell mobilization by decreasing bone marrow SDF-1 and up-regulating
[57017] CXCR4. Nature Immun. 3: 687-694, 2002.

[57018] 9320.

[57019] Simmers, R. N.; Smith, J.; Shannon, M. F.; Wong, G.; Lopez, A.
[57020] F.; Baker, E.; Sutherland, G. R.; Vadas, M. A.: Localization of the
[57021] human G-CSF gene to the region of a breakpoint in the translocation

[57022] typical of acute promyelocytic leukemia. Hum. Genet. 78: 134-136,

[57023] 1988.

[57024] 9321.

[57025] Souza, L. M.; Boone, T. C.; Gabrilove, J.; Lai, P. H.; Zsebo, K.

[57026] M.; Murdock, D. C.; Chazin, V. R.; Bruszewski, J.; Lee, H.; Chen,

[57027] K. K.; Barendt, J.; Platzer, E.; Moore, M. A. S.; Mertelsmann, R.;

[57028] Welte, K.: Recombinant human granulocyte colony-stimulating factor:

[57029] effects on normal and leukemic myeloid cells. Science 232: 61-65,

[57030] 1986.

[57031] 9322.

[57032] Sudo, S.; Yamada, H.; Kikuchi, K.; Sumie, A.; Yamashita, Y.; Tumura,

[57033] N.; Kawaguchi, I.; Fujimoto, S.; Kato, A.; Yamaguchi, J.: A case

[57034] of ovarian carcinoma with production of granulocyte colony-stimulating

[57035] factor. Brit. J. Haemat. 92: 137-139, 1996.

[57036] 9323.

[57037] Tse, W.; Zhu, W.; Chen, H. S.; Cohen, A.: A novel gene, AF1q,

[57038] fused to MLL in t(1;11)(q21;q23), is specifically expressed in leukemic

[57039] and immature hematopoietic cells. Blood 85: 650-656, 1995.

[57040] 9324.

[57041] Scott, M. P.: Vertebrate homeobox gene nomenclature. Cell 71:

[57042] 551-553, 1992.

[57043] 9325.

[57044] Carr, D. W.; Hausken, Z. E.; Fraser, I. D.; Stofko-Hahn, R. E.;

[57045] Scott, J. D: Association of the type II cAMP-dependent protein kinase

[57046] with a human thyroid RII-anchoring protein: cloning and characterization

[57047] of the RII-binding domain. J. Biol. Chem. 267: 13376-13382, 1992.

[57048] 9326.

[57049] Carr, D. W.; Stofko-Hahn, R. E.; Fraser, I. D.; Bishop, S. M.;

[57050] Acott, T. S.; Brennan, R. G.; Scott, J. D.: Interaction of the regulatory

[57051] subunit (RII) of cAMP-dependent protein kinase with RII-anchoring

[57052] proteins occurs through an amphipathic helix binding motif. J. Biol.

[57053] Chem. 266: 14188-14192, 1991.

[57054] 9327.

[57055] Rubino, D.; Driggers, P.; Arbit, D.; Kemp, L.; Miller, B.; Coso,

[57056] O.; Pagliai, K.; Gray, K.; Gutkind, S.; Segars, J.: Characteri-zation

[57057] of Brx, a novel Dbl family member that modulates estro-gen receptor

[57058] action. Oncogene 16: 2513-2526, 1998.

[57059] 9328.

[57060] Sterpetti, P.; Hack, A. A.; Bashar, M. P.; Park, B.; Cheng, S.-D.;

[57061] Knoll, J. H. M.; Urano, T.; Feig, L. A.; Toksoz, D.: Activation

of

[57062] the Lbc Rho exchange factor proto-oncogene by truncation of an extended

[57063] C terminus that regulates transformation and targeting. Molec. Cell.

[57064] Biol. 19: 1334-1345, 1999.

[57065] 9329.

[57066] Toksoz, D.; Williams, D. A.: Novel human oncogene lbc detected

[57067] by transfection with distinct homology regions to signal transduction

[57068] products. Oncogene 9: 621-628, 1994.

[57069] 9330.

[57070] Hirai, H.; Tanaka, K.; Yoshie, O.; Ogawa, K.; Kenmotsu, K.; Takamori,

[57071] Y.; Ichimasa, M.; Sugamura, K.; Nakamura, M.; Takano, S.; Nagata,

[57072] K.: Prostaglandin D2 selectively induces chemotaxis in T helper type

[57073] 2 cells, eosinophils, and basophils via seven-transmembrane receptor

[57074] CRTH2. J. Exp. Med. 193: 255-261, 2001.

[57075] 9331.

[57076] Margolis, R. L.; O'Hearn, E.; Rosenblatt, A.; Willour, V.; Holmes,

[57077] S. E.; Franz, M. L.; Callahan, C.; Hwang, H. S.; Troncoso, J. C.;

[57078] Ross, C. A.: A disorder similar to Huntington's disease is associated

[57079] with a novel CAG repeat expansion. Ann. Neurol. 50: 373-380, 2001.

[57080] 9332.

[57081] Lacy, S. E.; Bonnemann, C. G.; Buzney, E. A.; Kunkel, L. M.: Identification

[57082] of FLRT1, FLRT2, and FLRT3: a novel family of transmembrane leucine-rich

[57083] repeat proteins. Genomics 62: 417-426, 1999.

[57084] 9333.

[57085] Borges, L.; Hsu, M.-L.; Fanger, N.; Kubin, M.; Cosman, D.: A family

[57086] of human lymphoid and myeloid Ig-like receptors, some of which bind

[57087] to MHC class I molecules. J. Immun. 159: 5192-5196, 1997.

[57088] 9334.

[57089] Andre, P.; Biassoni, R.; Colonna, M.; Cosman, D.; Lanier, L. L.;

[57090] Long, E. O.; Lopez-Botet, M.; Moretta, A.; Moretta, L.; Parham, P.;

[57091] Trowsdale, J.; Vivier, E.; Wagtmann, N.; Wilson, M. J.: New nomenclature

[57092] for MHC receptors. Nature Immun. 2: 661 only, 2001.

[57093] 9335.

[57094] Banham, A. H.; Colonna, M.; Cella, M.; Micklem, K. J.; Pulford,

[57095] K.; Willis, A. C.; Mason, D. Y.: Identification of the CD85 antigen

[57096] as ILT2, an inhibitory MHC class I receptor of the immunoglobulin

[57097] superfamily. J. Leukoc. Biol. 65: 841-845, 1999.

[57098] 9336.

[57099] Colonna, M.; Navarro, F.; Bellon, T.; Llano, M.; Garcia, P.; Samaridis,

[57100] J.; Angman, L.; Cella, M.; Lopez-Botet, M.: A common inhibitory receptor

[57101] for major histocompatibility complex class I molecules on human lymphoid

[57102] and myelomonocytic cells. J. Exp. Med. 186: 1809-1818, 1997.

[57103] 9337.

[57104] Fenster, S. D.; Chung, W. J.; Zhai, R.; Cases-Langhoff, C.; Voss,

[57105] B.; Garner, A. M.; Kaempf, U.; Kindler, S.; Gundelfinger, E. D.; Garner,

[57106] C. C.: Piccolo, a presynaptic zinc finger protein structurally related
[57107] to Bassoon. Neuron 25: 203-214, 2000.

[57108] 9338.

[57109] Tung, R. M.; Blenis, J.: A novel human SPS1/STE20 homologue, KHS,
[57110] activates Jun N-terminal kinase. Oncogene 14: 653-659, 1997.
[57111] 9339.

[57112] Redeker, E.; Hoovers, J. M.; Alders, M.; van Moorsel, C. J.; Ivens,
[57113] A. C.; Gregory, S.; Kalikin, L.; Bliek, J.; de Galan, L.; van den
[57114] Bogaard, R.; Visser, J.; van der Voort, R.; Feinberg, A. P.; Little,
[57115] P. F. R.; Westerveld, A.; Mannens, M.: An integrated physical map
[57116] of 210 markers assigned to the short arm of human chromosome 11. Genomics 21:
[57117] 538-550, 1994.

[57118] 9340.

[57119] Rosnet, O.; Mattei, M.-G.; Delattre, O.; Schiff, C.: VPREB3: cDNA
[57120] characterization and expression in human and chromo-some mapping in
[57121] human and mouse. Cytogenet. Cell Genet. 87: 205-208, 1999.
[57122] 9341.

[57123] Hayashi, T.; Huang, J.; Deeb, S. S.: RINX(VSX1), a novel homeobox
[57124] gene expressed in the inner nuclear layer of the adult retina. Genomics 67:
[57125] 128-139, 2000.

[57126] 9342.

[57127] Ohtoshi, A.; Justice, M. J.; Behringer, R. R.: Isolation and characterization
[57128] of Vsx1, a novel mouse CVC paired-like homeobox gene expressed during
[57129] embryogenesis and in the retina. Biochem. Biophys. Res. Commun. 286:
[57130] 133-140, 2001.

[57131] 9343.

[57132] Semina, E. V.; Mintz-Hittner, H. A.; Murray, J. C.: Isolation

[57133] and characterization of a novel human paired-like homeodomain-containing

[57134] transcription factor gene, VSX1, expressed in ocular tissues. Genomics 63:

[57135] 289-293, 2000.

[57136] 9344.

[57137] Jones, J. M.; Morrell, J. C.; Gould, S. J.: Identification and

[57138] characterization of HAOX1, HAOX2, and HAOX3, three human peroxisomal

[57139] 2-hydroxy acid oxidases. J. Biol. Chem. 275:
12590-12597, 2000.

[57140] 9345.

[57141] Martignetti, J. A.; Al Aqeel, A.; Al Sewairi, W.; Boumah, C. E.;

[57142] Kambouris, M.; Al Mayouf, S.; Sheth, K. V.; Al Eid, W.; Dowling, O.;

[57143] Harris, J.; Glucksman, M. J.; Bahabri, S.; Meyer, B. F.; Desnick,

[57144] R. J.: Mutation of the matrix metalloproteinase 2 gene (MMP2) causes

[57145] a multicentric osteolysis and arthritis syndrome. Nature Genet. 28:

[57146] 261-265, 2001.

[57147] 9346.

[57148] Horikoshi, N.; Cong, J.; Kley, N.; Shenk, T.: Isolation of differentially

[57149] expressed cDNAs from p53-dependent apoptotic cells: activation of

[57150] the human homologue of the Drosophila peroxidasin gene. Biochem.

[57151] Biophys. Res. Commun. 261: 864-869, 1999.

[57152] 9347.

[57153] Weiler, S. R.; Taylor, S. M.; Deans, R. J.; Kan-Mitchell, J.; Mitchell,

[57154] M. S.; Trent, J. M.: Assignment of a human melanoma associated gene

[57155] MG50 (D2S448) to chromosome 2p25.3 by fluorescence in situ hybridization. Genomics 22:

[57156] 243-244, 1994.

[57157] 9348.

[57158] Matloubian, M.; David, A.; Engel, S.; Ryan, J. E.; Cyster, J.

G.

[57159] : A transmembrane CXC chemokine is a ligand for HIV-coreceptor Bonzo. Nature

[57160] Immun. 1: 298-304, 2000.

[57161] 9349.

[57162] Fedele, M.; Benvenuto, G.; Pero, R.; Majello, B.; Battista, S.;

[57163] Lembo, F.; Vollono, E.; Day, P. M.; Santoro, M.; Lania, L.; Bruni,

[57164] C. B.; Fusco, A.; Chiariotti, L.: A novel member of the BTB/POZ family,

[57165] PATZ, associates with the RNF4 RING finger protein and acts as a transcriptional

[57166] repressor. J. Biol. Chem. 275: 7894-7901, 2000.

[57167] 9350.

[57168] Kobayashi, A.; Yamagiwa, H.; Hoshino, H.; Muto, A.; Sato, K.; Morita,

[57169] M.; Hayashi, N.; Yamamoto, M.; Igarashi, K.: A combinato-rial code

[57170] for gene expression generated by transcription factor Bach2 and MAZR

[57171] (MAZ-related factor) through the BTB/POZ domain. Molec.

Cell. Biol. 20:

[57172] 1733-1746, 2000.

[57173] 9351.

[57174] McAleer, M. A.; Breen, M. A.; White, N. L.; Matthews, N.: pABC11

[57175] (also known as MOAT-C and MRP5), a member of the ABC family of proteins,

[57176] has anion transporter activity but does not confer multidrug resistance

[57177] when overexpressed in human embryonic kidney 293 cells. J. Biol.

[57178] Chem. 274: 23541-23548, 1999.

[57179] 9352.

[57180] Oguri, T.; Isobe, T.; Suzuki, T.; Nishio, K.; Fujiwara, Y.; Katoh,

[57181] O.; Yamakido, M.: Increased expression of the MRP5 gene is associated

[57182] with exposure to platinum drugs in lung cancer. Int. J. Cancer 86:

[57183] 95-100, 2000.

[57184] 9353.

[57185] Suzuki, T.; Nishio, K.; Sasaki, H.; Kurokawa, H.; Saito-Ohara,
[57186] F.; Ikeuchi, T.; Tanabe, S.; Terada, M.; Saijo, N.: cDNA cloning
[57187] of a short type of multidrug resistance protein homologue, SMRP, from
[57188] a human lung cancer cell line. Biochem. Biophys. Res. Commun. 238:
[57189] 790-794, 1997.

[57190] 9354.

[57191] Suzuki, T.; Sasaki, H.; Kuh, H.-J.; Agui, M.; Tatsumi, Y.; Tanabe,
[57192] S.; Terada, M.; Saijo, N.; Nishio, K.: Detailed structural analysis
[57193] on both human MRP5 and mouse mrp5 transcripts. Gene 242: 167-173,
[57194] 2000.

[57195] 9355.

[57196] Wijnholds, J.; Mol, C. A. A. M.; van Deemter, L.; de Haas, M.;
[57197] Scheffer, G. L.; Baas, F.; Beijnen, J. H.; Scheper, R. J.; Hatse,

[57198] S.; De Clercq, E.; Balzarini, J.; Borst, P.: Multidrug-resistance
[57199] protein 5 is a multispecific organic anion transporter able to transport
[57200] nucleotide analogs. Proc. Nat. Acad. Sci. 97: 7476-7481, 2000.
[57201] 9356.

[57202] Tateishi, S.; Sakuraba, Y.; Masuyama, S.; Inoue, H.; Yamaizumi,
[57203] M.: Dysfunction of human Rad18 results in defective postreplication
[57204] repair and hypersensitivity to multiple mutagens. Proc. Nat. Acad.
[57205] Sci. 97: 7927-7932, 2000.

[57206] 9357.

[57207] Arakawa, H.; Hauschild, J.; Buerstedde, J.-M.: Requirement of
[57208] the activation-induced deaminase (AID) gene for immunoglobulin gene
[57209] conversion. Science 295: 1301-1306, 2002.

[57210] 9358.

[57211] Fagarasan, S.; Kinoshita, K.; Muramatsu, M.; Ikuta, K.; Honjo,
[57212] T.: In situ class switching and differentiation to IgA producing
[57213] cells in the gut lamina propria. Nature 413: 639-643, 2001.
[57214] 9359.

[57215] Muramatsu, M.; Kinoshita, K.; Fagarasan, S.; Yamada, S.; Shinkai,
[57216] Y.; Honjo, T.: Class switch recombination and hypermutation require
[57217] activation-induced cytidine deaminase (AID), a potential RNA editing
[57218] enzyme. Cell 102: 553-563, 2000.

[57219] 9360.

[57220] Muramatsu, M.; Sankaranand, V. S.; Anant, S.; Sugai, M.; Kinoshita,
[57221] K.; Davidson, N. O.; Honjo, T.: Specific expression of activation-induced
[57222] cytidine deaminase (AID), a novel member of the RNA-editing deaminase
[57223] family in germinal center B cells. J. Biol. Chem. 274:

18470-18476,

[57224] 1999.

[57225] 9361.

[57226] Rowley, D. R.; Tindall, D. J.: Responses of NBT-II bladder carcinoma

[57227] cells to conditioned medium from normal fetal urogenital sinus. Cancer

[57228] Res. 47: 2955-2960, 1987.

[57229] 9362.

[57230] Kasaian, M. T.; Whitters, M. J.; Carter, L. L.; Lowe, L. D.; Jussif,

[57231] J. M.; Deng, B.; Johnson, K. A.; Witek, J. S.; Senices, M.; Konz,

[57232] R. F.; Wurster, A. L.; Donaldson, D. D.; Collins, M.; Young, D. A.;

[57233] Grusby, M. J.: IL-21 limits NK cell responses and promotes antigen-specific

[57234] T cell activation: a mediator of the transition from innate to adaptive

[57235] immunity. Immunity 16: 559-569, 2002.

[57236] 9363.

[57237] Brattsand, M.; Egelrud, T.: Purification, molecular cloning, and
[57238] expression of a human stratum corneum trypsin-like serine protease
[57239] with possible function in desquamation. J. Biol. Chem. 274: 30033-30040,
[57240] 1999.

[57241] 9364.

[57242] Choglay, A. A.; Chapple, J. P.; Blatch, G. L.; Cheetham, M. E.
[57243] : Identification and characterization of a human mitochondrial homologue
[57244] of the bacterial co-chaperone GrpE. Gene 267: 125-134, 2001.
[57245] 9365.

[57246] Zheng, P.; Eastman, J.; Vande Pol, S.; Pimplikar, S. W.: PAT1,
[57247] a microtubule-interacting protein, recognizes the basolateral sorting
[57248] signal of amyloid precursor protein. Proc. Nat. Acad. Sci. 95: 14745-14750,
[57249] 1998.

[57250] 9366.

[57251] De Valck, D.; Jin, D. Y.; Heyninck, K.; Van de Craen, M.; Contreras,

[57252] R.; Fiers, W.; Jeang, K. T.; Beyaert, R.: The zinc finger protein

[57253] A20 interacts with a novel anti-apoptotic protein which is cleaved

[57254] by specific caspases. Oncogene 18: 4182-4190, 1999.

[57255] 9367.

[57256] Gachon, F.; Peleraux, A.; Thebault, S.; Dick, J.; Lemasson, I.;

[57257] Devaux, C.; Mesnard, J.-M.: CREB-2, a cellular CRE-dependent transcription

[57258] repressor, functions in association with tax as an activator of the

[57259] human T-cell leukemia virus type 1 promoter. J. Virol. 72: 8332-8337,

[57260] 1998.

[57261] 9368.

[57262] Ling, L.; Goeddel, D. V.: T6BP, a TRAF6-interacting protein involved

[57263] in IL-1 signaling. Proc. Nat. Acad. Sci. 97: 9567-9572, 2000.

[57264] 9369.

[57265] Cowell, I. G.; Skinner, A.; Hurst, H. C.: Transcriptional repression

[57266] by a novel member of the bZIP family of transcription factors. Molec.

[57267] Cell. Biol. 12: 3070-3077, 1992.

[57268] 9370.

[57269] Hulme, D. J.; Blair, I. P.; Dawkins, J. L.; Nicholson, G. A.:

[57270] Exclusion of NFIL3 as the gene causing hereditary sensory neuropathy

[57271] type I by mutation analysis. Hum. Genet. 106: 594-596, 2000.

[57272] 9371.

[57273] Ikushima, S.; Inukai, T.; Inaba, T.; Nimer, S. D.; Cleveland, J.

[57274] L.; Look, A. T.: Pivotal role for the NFIL3/E4BP4 transcription factor

[57275] in interleukin 3-mediated survival of pro-B lymphocytes. Proc. Nat.

[57276] Acad. Sci. 94: 2609-2614, 1997.

[57277] 9372.

[57278] Zhang, W.; Zhang, J.; Kornuc, M.; Kwan, K.; Frank, R.;
Nimer, S.
[57279] D.: Molecular cloning and characterization of NF-IL3A, a
transcriptional
[57280] activator of the human interleukin-3 promoter. Molec.
Cell. Biol. 15:
[57281] 6055-6063, 1995.

[57282] 9373.

[57283] Ozaki, K.; Kikly, K.; Michalovich, D.; Young, P. R.; Leonard,
W.
[57284] J.: Cloning of a type I cytokine receptor most related to
the IL-2
[57285] receptor beta chain. Proc. Nat. Acad. Sci. 97:
11439-11444, 2000.
[57286] 9374.

[57287] Parrish-Novak, J.; Dillon, S. R.; Nelson, A.; Hammond, A.;
Sprecher,
[57288] C.; Gross, J. A.; Johnston, J.; Madden, K.; Xu, W.; West, J.;
Schrader,

[57289] S.; Burkhead, S.; and 26 others: Interleukin 21 and its receptor

[57290] are involved in NK cell expansion and regulation of lymphocyte function. Nature 408:

[57291] 57-63, 2000.

[57292] 9375.

[57293] Reither, A.; Hehlmann, R.; Goldman, J. M.; Cross, N. C.: The 8p11

[57294] myeloproliferative syndrome. Med. Klin. 94: 207-210, 1999.

[57295] 9376.

[57296] Bartholin, L.; Maguer-Satta, V.; Hayette, S.; Martel, S.; Gadoux,

[57297] M.; Bertrand, S.; Corbo, L.; Lamadon, C.; Morera, A.-M.; Magaud, J.-P.;

[57298] Rimokh, R.: FLRG, an activin-binding protein, is a new target of

[57299] TGF-beta transcription activation through Smad proteins. Oncogene 20:

[57300] 5409-5419, 2001.

[57301] 9377.

[57302] Rimokh, R.; Berger, F.; Delsol, G.; Charrin, C.; Bertheas, M. F.;

[57303] Ffrench, M.; Garoscio, M.; Felman, P.; Coiffier, C.; Bryon, P. A.

[57304] : Rearrangement and overexpression of the BCL-1/PRAD-1 gene in intermediate

[57305] lymphocytic lymphomas and in t(11q13)-bearing leukemias. Blood 81:

[57306] 3063-3067, 1993.

[57307] 9378.

[57308] Bahou, W. F.; Campbell, A. D.; Wicha, M. S.: cDNA cloning and

[57309] molecular characterization of MSE55, a novel human serum constituent

[57310] protein that displays bone marrow stromal/endothelial cell-specific

[57311] expression. J. Biol. Chem. 267: 13986-13992, 1992.

[57312] 9379.

[57313] Burbelo, P. D.; Drechsel, D.; Hall, A.: A conserved binding motif

[57314] defines numerous candidate target proteins for both Cdc42 and Rac

[57315] GTPases. J. Biol. Chem. 270: 29071-29074, 1995.

[57316] 9380.

[57317] Hermey, G.; Riedel, I. B.; Hampe, W.; Schaller, H. C.; Hermans-Borgmeyer,

[57318] I.: Identification and characterization of SorCS, a third member

[57319] of a novel receptor family. Biochem. Biophys. Res. Commun. 266:

[57320] 347-351, 1999.

[57321] 9381.

[57322] Rezgaoui, M.; Hermey, G.; Riedel, I. B.; Hampe, W.; Schaller, H.

[57323] C.; Hermans-Borgmeyer, I.: Identification of SorCS2, a novel member

[57324] of the VPS10 domain containing receptor family, prominently expressed

[57325] in the developing mouse brain. Mech. Dev. 100: 335-338, 2001.

[57326] 9382.

[57327] Oda, K.; Arakawa, H.; Tanaka, T.; Matsuda, K.; Tanikawa, C.; Mori,

[57328] T.; Nishimori, H.; Tamai, K.; Tokino, T.; Nakamura, Y.; Taya, Y.:

[57329] p53AIP1, a potential mediator of p53-dependent apoptosis, and its

[57330] regulation by Ser-46-phosphorylated p53. Cell 102: 849-862, 2000.

[57331] 9383.

[57332] Strotmann, R.; Harteneck, C.; Nunnenmacher, K.; Schultz, G.; Plant,

[57333] T. D: OTRPC4, a nonselective cation channel that confers sensitivity

[57334] to extracellular osmolarity Nature Cell. Biol. 2: 695-702, 2000.

[57335] 9384.

[57336] Wissenbach, U.; Bodding, M.; Freichel, M.; Flockerzi, V.: Trp12,

[57337] a novel Trp related protein from kidney FEBS Lett. 485: 127-134,

[57338] 2000.

[57339] 9385.

[57340] Baron, C. L.; Malhotra, V.: Role of diacylglycerol in PKD recruitment

[57341] to the TGN and protein transport to the plasma membrane. Science 295:
[57342] 325-328, 2002.

[57343] 9386.

[57344] Jamora, C.; Yamanouye, N.; Van Lint, J.; Laudenslager, J.; Vandenheede,
[57345] J. R.; Faulkner, D. J.; Malhotra, V.: G-beta-gamma-mediated regulation
[57346] of Golgi organization is through the direct activation of protein
[57347] kinase D. Cell 98: 59-68, 1999.

[57348] 9387.

[57349] Johannes, F.-J.; Prestle, J.; Eis, S.; Oberhagemann, P.; Pfizenmaier,
[57350] K.: PKC-mu is a novel, atypical member of the protein kinase C family. J.
[57351] Biol. Chem. 269: 6140-6148, 1994.

[57352] 9388.

[57353] Maeda, Y.; Beznoussenko, G. V.; Van Lint, J.; Mironov, A. A.; Malhotra,
[57354] V.: Recruitment of protein kinase D to the trans-Golgi

network via
[57355] the first cysteine-rich domain. EMBO J. 20: 5982-5990, 2001.
[57356] 9389.

[57357] Owczarek, C. M.; Portbury, K. J.; Kola, I.; Hertzog, P. J.: Assignment
[57358] of protein kinase C mu (PRKCM) to human chromosome band 14q11 with
[57359] somatic cell hybrids and radiation hybrids. Cytogenet. Cell Genet. 89:
[57360] 240-241, 2000.

[57361] 9390.

[57362] Ouchi, N.; Kihara, S.; Arita, Y.; Okamoto, Y.; Maeda, K.; Kuriyama,
[57363] H.; Hotta, K.; Nishida, M.; Takahashi, M.; Muraguchi, M.; Ohmoto,
[57364] Y.; Nakamura, T.; Yamashita, S.; Funahashi, T.; Matsuzawa, Y.: Adiponectin,
[57365] an adipocyte-derived plasma protein, inhibits endothelial NF-kappa-B
[57366] signaling through a cAMP-dependent pathway. Circulation 102: 1296-1301,

[57367] 2000.

[57368] 9391.

[57369] Schaffler, A.; Orso, E.; Palitzsch, K.-D.; Buchler, C.; Drob-
nik,

[57370] W.; Furst, A.; Scholmerich, J.; Schmitz, G.: The human
apM-1, an

[57371] adipocyte-specific gene linked to the family of TNF's and
to genes

[57372] expressed in activated T cells, is mapped to chromosome
1q21.3-q23,

[57373] a susceptibility locus identified for familial combined hy-
perlipidaemia

[57374] (FCH). Biochem. Biophys. Res. Commun. 260: 416-425,
1999.

[57375] 9392.

[57376] Gong, T.-W. L.; Burmeister, M.; Lomax, M. I.: The novel
gene D4Mil1e

[57377] maps to mouse chromosome 4 and human chromosome
1p36. Mammalian Genome 7:

[57378] 790-791, 1996.

[57379] 9393.

[57380] Nangaku, M.; Sato-Yoshitake, R.; Okada, Y.; Noda, Y.; Takemura,
[57381] R.; Yamazaki, H.; Hirokawa, N.: KIF1B, a novel microtubule plus end-directed
[57382] monomeric motor protein for transport of mitochondria. Cell 79:
[57383] 1209-1220, 1994.

[57384] 9394.

[57385] Saito, M.; Hayashi, Y.; Suzuki, T.; Tanaka, H.; Hozumi, I.; Tsuji,
[57386] S.: Linkage mapping of the gene for Charcot-Marie-Tooth disease type
[57387] 2 to chromosome 1p (CMT2A) and the clinical features of CMT2A. Neurology 49:
[57388] 1630-1635, 1997.

[57389] 9395.

[57390] Yang, H. W.; Chen, Y. Z.; Takita, J.; Soeda, E.; Piao, H. Y.; Hayashi,
[57391] Y.: Genomic structure and mutational analysis of the human KIF1B
[57392] gene which is homozygously deleted in neuroblastoma at chromosome

[57393] 1p36.2. Oncogene 20: 5075-5083, 2001.

[57394] 9396.

[57395] Hu, P.; Yu, L.; Zhang, M.; Zheng, L.; Zhao, Y.; Fu, Q.; Zhao, S.

[57396] : Molecular cloning and mapping of the brain-abundant B1-gamma subunit

[57397] of protein phosphatase 2A, PPP2R2C, to human chromosome 4p16. Genomics 67:

[57398] 83-86, 2000.

[57399] 9397.

[57400] Clark, R. M.; Marker, P. C.; Kingsley, D. M.: A novel candidate

[57401] gene for mouse and human preaxial polydactyly with altered expression

[57402] in limbs of hemimelic extra-toes mutant mice. Genomics 67: 19-27,

[57403] 2000.

[57404] 9398.

[57405] Knudsen, T. B.; Kochhar, D. M.: The role of morphogenetic cell

[57406] death during abnormal limb-bud outgrowth in mice het-

erozygous for
[57407] the dominant mutation hemimelia-extra toe (Hmx). J. Embryol. Exp.
[57408] Morphol. 65 (suppl.): 289-307, 1981.

[57409] 9399.

[57410] Matsuda, S.; Kawamura-Tsuzuku, J.; Ohsugi, M.; Yoshida, M.; Emi,
[57411] M.; Nakamura, Y.; Onda, M.; Yoshida, Y.; Nishiyama, A.; Yamamoto,
[57412] T.: Tob, a novel protein that interacts with p185erbB2, is associated
[57413] with anti-proliferative activity. Oncogene 12: 705-713, 1996.
[57414] 9400.

[57415] Tzachanis, D.; Freeman, G. J.; Hirano, N.; van Puijenbroek, A.
[57416] A. F. L.; Delfs, M. W.; Berezovskaya, A.; Nadler, L. M.; Boussiotis,
[57417] V. A.: Tob is a negative regulator of activation that is expressed
[57418] in anergic and quiescent T cells. Nature Immun. 2: 1174-1182, 2001.

[57419] 9401.

[57420] Yoshida, Y.; Tanaka, S.; Umemori, H.; Minowa, O.; Usui, M.; Ikematsu,

[57421] N.; Hosoda, E.; Imamura, T.; Kuno, J.; Yamashita, T.; Miya-zono, K.;

[57422] Noda, M.; Noda, T.; Yamamoto, T.: Negative regulation of BMP/Smad

[57423] signaling by Tob in osteoblasts. Cell 103: 1085-1097, 2000.

[57424] 9402.

[57425] Dell'Angelica, E. C.; Puertollano, R.; Mullins, C.; Aguilar, R.

[57426] C.; Vargas, J. D.; Hartnell, L. M.; Bonifacino, J. S.: GGAs: a family

[57427] of ADP ribosylation factor-binding proteins related to adaptors and

[57428] associated with the Golgi complex. J. Cell Biol. 149: 81-93, 2000.

[57429] 9403.

[57430] Hirst, J.; Lui, W. W. Y.; Bright, N. A.; Totty, N.; Seaman, M.

[57431] N. J.; Robinson, M. S.: A family of proteins with gamma-adaptin and

[57432] VHS domains that facilitate trafficking between the trans-Golgi network

[57433] and the vacuole/lysosome. J. Cell Biol. 149: 67-69, 2000.

[57434] 9404.

[57435] Blumberg, H.; Conklin, D.; Xu, W.; Grossmann, A.; Brender, T.;

[57436] Carollo, S.; Eagan, M.; Foster, D.; Haldeman, B. A.; Ham-mond, A.;

[57437] Haugen, H.; Jelinek, L.; and 14 others: Interleukin 20: dis-covery,

[57438] receptor identification, and role in epidermal function. Cell 104:

[57439] 9-19, 2001.

[57440] 9405.

[57441] Huang, E. Y.; Madireddi, M. T.; Gopalkrishnan, R. V.; Leszczyniecka,

[57442] M.; Su, Z.; Lebedeva, I. V.; Kang, D.; Jiang, H.; Lin, J. J.; Alexandre,

[57443] D.; Chen, Y.; Vozhilla, N.: and 9 others: Genomic struc-ture, chromosomal

[57444] localization and expression profile of a novel melanoma differentiation

[57445] associated (mda-7) gene with cancer specific growth suppressing and
[57446] apoptosis inducing properties. Oncogene 20: 7051-7063, 2001.
[57447] 9406.

[57448] Jiang, H.; Lin, J. J.; Su, Z.-Z.; Goldstein, N. I.; Fisher, P.

[57449] B.: Subtraction hybridization identifies a novel melanoma differentiation
[57450] associated gene, mda-7, modulated during human melanoma differentiation,
[57451] growth and progression. Oncogene 11: 2477-2486, 1995.

[57452] 9407.

[57453] Jiang, H.; Su, Z.-Z.; Lin, J. J.; Goldstein, N. I.; Young, C. S.

[57454] H.; Fisher, P. B.: The melanoma differentiation associated gene mda-7
[57455] suppresses cancer cell growth. Proc. Nat. Acad. Sci. 93: 9160-9165,
[57456] 1996.

[57457] 9408.

[57458] Su, Z.; Madireddi, M. T.; Lin, J. J.; Young, C. S. H.; Kitada,

[57459] S.; Reed, J. C.; Goldstein, N. I.; Fisher, P. B.: The cancer growth
[57460] suppressor gene mda-7 selectively induces apoptosis in human breast
[57461] cancer cells and inhibits tumor growth in nude mice. Proc. Nat. Acad.
[57462] Sci. 95: 14400-14405, 1998.

[57463] 9409.

[57464] Cheng, Y.; Lotan, R.: Molecular cloning and characterization of
[57465] a novel retinoic acid-inducible gene that encodes a putative G protein-coupled
[57466] receptor. J. Biol. Chem. 273: 35008-35015, 1998.

[57467] 9410.

[57468] Mansour, S. J.; Skaug, J.; Zhao, X.-H.; Giordano, J.; Scherer,
[57469] S. W.; Melancon, P.: p200 ARF-GEP1: a Golgi-localized guanine nucleotide
[57470] exchange protein whose Sec7 domain is targeted by the drug brefeldin
[57471] A. Proc. Nat. Acad. Sci. 96: 7968-7973, 1999.

[57472] 9411.

[57473] Morinaga, N.; Moss, J.; Vaughan, M.: Cloning and expres-
sion of

[57474] a cDNA encoding a bovine brain brefeldin A-sensitive
guanine nucleotide-exchange

[57475] protein for ADP-ribosylation factor. Proc. Nat. Acad. Sci.
94: 12926-12931,

[57476] 1997.

[57477] 9412.

[57478] Morinaga, N.; Tsai, S.-C.; Moss, J.; Vaughan, M.: Isolation
of

[57479] a brefeldin A-inhibited guanine nucleotide-exchange pro-
tein for ADP

[57480] ribosylation factor (ARF) 1 and ARF3 that contains a
Sec7-like domain. Proc.

[57481] Nat. Acad. Sci. 93: 12856-12860, 1996.

[57482] 9413.

[57483] Togawa, A.; Morinaga, N.; Ogasawara, M.; Moss, J.;
Vaughan, M.

[57484] : Purification and cloning of a brefeldin A-inhibited gua-
nine nucleotide-exchange

[57485] protein for ADP-ribosylation factors. J. Biol. Chem. 274:
12308-12315,
[57486] 1999.

[57487] 9414.

[57488] Zou, H.; McGarry, T. J.; Bernal, T.; Kirschner, M. W.: Identi-
fication
[57489] of a vertebrate sister-chromatid separation inhibitor in-
volved in
[57490] transformation and tumorigenesis. Science 285: 418-422,
1999.
[57491] 9415.

[57492] Colombo, R.: Dating the origin of the V170M mutation
causing non-type
[57493] I cystinuria in Libyan Jews by linkage disequilibrium and
physical
[57494] mapping of the SLC7A9 gene. Genomics 69: 131-134,
2000.
[57495] 9416.

[57496] Kim, H.; Scorillas, A.; Katsaros, D.; Yousef, G. M.; Masso-
brio,
[57497] M.; Fracchioli, S.; Piccinno, R.; Gordini, G.; Diamandis, E.
P.:

[57498] Human kallikrein gene 5 (KLK5) expression is an indicator of poor
[57499] prognosis in ovarian cancer. Brit. J. Cancer 84: 643-650, 2001.
[57500] 9417.

[57501] Yousef, G. M.; Diamandis, E. P.: The new kallikrein-like gene,
[57502] KLK-L2: molecular characterization, mapping, tissue expression, and
[57503] hormonal regulation. J. Biol. Chem. 274: 37511-37516, 1999.
[57504] 9418.

[57505] Bidart, J.-M.; Mian, C.; Lazar, V.; Russo, D.; Filetti, S.; Caillou,
[57506] B.; Schlumberger, M.: Expression of pendrin and the Pendred syndrome
[57507] (PDS) gene in human thyroid tissues. J. Clin. Endocr. Metab. 85:
[57508] 2028-2033, 2000.

[57509] 9419.

[57510] Coyle, B.; Reardon, W.; Herbrick, J.-A.; Tsui, L.-C.; Gausden,

[57511] E.; Lee, J.; Coffey, R.; Grueters, A.; Grossman, A.; Phelps, P. D.;
[57512] Luxon, L.; Kendall-Taylor, P.; Scherer, S. W.; Trembath, R. C.: Molecular
[57513] analysis of the PDS gene in Pendred syndrome (sensorineural hearing
[57514] loss and goitre). Hum. Molec. Genet. 7: 1105-1112, 1998.

[57515] 9420.

[57516] Everett, L. A.; Belyantseva, I. A.; Noben-Trauth, K.; Cantos, R.;
[57517] Chen, A.; Thakkar, S. I.; Hoogstraten-Miller, S. L.; Kachar, B.; Wu,
[57518] D. K.; Green, E. D.: Targeted disruption of mouse Pds provides insight
[57519] about the inner-ear defects encountered in Pendred syndrome. Hum.
[57520] Molec. Genet. 10: 153-161, 2001.

[57521] 9421.

[57522] Everett, L. A.; Morsli, H.; Wu, D. K.; Green, E. D.: Expression
[57523] pattern of the mouse ortholog of the Pendred's syndrome gene (Pds)

[57524] suggests a key role for pendrin in the inner ear. Proc. Nat. Acad.
[57525] Sci. 96: 9727-9732, 1999.

[57526] 9422.

[57527] Kraiem, Z.; Heinrich, R.; Sadeh, O.; Shiloni, E.; Nassir, E.;

[57528] Hazani, E.; Glaser, B.: Sulfate transport is not impaired in Pendred
[57529] syndrome thyrocytes. J. Clin. Endocr. Metab. 84: 2574-2576, 1999.

[57530] 9423.

[57531] Lopez-Bigas, N.; Rabionet, R.; de Cid, R.; Govea, N.; Gas-parini,
[57532] P.; Zelante, L.; Arbones, M. L.; Estivill, X.: Splice-site mu-tation
[57533] in the PDS gene may result in intrafamilial variability for deafness
[57534] in Pendred syndrome. Hum. Mutat. 14: 520-526, 1999.

[57535] 9424.

[57536] Scott, D. A.; Wang, R.; Kreman, T. M.; Andrews, M.; Mc-Donald,
[57537] J. M.; Bishop, J. R.; Smith, R. J. H.; Karniski, L. P.; Sheffield,

[57538] V. C.: Functional differences of the PDS gene product are associated
[57539] with phenotypic variation in patients with Pendred syndrome and non-syndromic
[57540] hearing loss (DFNB4). Hum. Molec. Genet. 9: 1709-1715, 2000.
[57541] 9425.

[57542] Markert, M. L.; Hershfield, M. S.; Schiff, R. I.; Buckley, R.

[57543] H.: Adenosine deaminase and purine nucleoside phosphorylase deficiencies:
[57544] evaluation of therapeutic interventions in eight patients. J. Clin.
[57545] Immun. 7: 389-399, 1987.

[57546] 9426.

[57547] Liquori, C. L.; Ricker, K.; Moseley, M. L.; Jacobsen, J. F.; Kress,
[57548] W.; Naylor, S. L.; Day, J. W.; Ranum, L. P. W.: Myotonic dystrophy
[57549] type 2 caused by a CCTG expansion in intron 1 of ZNF9. Science 293:
[57550] 864-867, 2001.

[57551] 9427.

[57552] Lusis, A. J.; Rajavashisth, T. B.; Klisak, I.; Heinzmann, C.; Mohandas,

[57553] T.; Sparkes, R. S.: Mapping of the gene for CNBP, a finger protein,

[57554] to human chromosome 3q13.3-q24. Genomics 8: 411-414, 1990. Note:

[57555] Erratum: Genomics 9: 564 only, 1991.

[57556] 9428.

[57557] Rajavashisth, T. B.; Taylor, A. K.; Andalibi, A.; Svenson, K. L.;

[57558] Lusis, A. J.: Identification of a zinc finger protein that binds

[57559] to the sterol regulatory element. Science 245: 640-643, 1989.

[57560] 9429.

[57561] Ewen, M. E.; Xing, Y.; Lawrence, J. B.; Livingston, D. M.: Molecular

[57562] cloning, chromosomal mapping, and expression of the cDNA for p107,

[57563] a retinoblastoma gene product-related protein. Cell 66: 1155-1164,

[57564] 1991.

[57565] 9430.

[57566] Huppi, K.; Siwarski, D.; Mock, B. A.; Dosik, J.; Hamel, P. A.:

[57567] Molecular cloning, chromosomal mapping, and expression of the mouse

[57568] p107 gene. Mammalian Genome 7: 353-355, 1996.

[57569] 9431.

[57570] Kim, K. K.; Soonpaa, M. H.; Wang, H.; Field, L. J.: Developmental

[57571] expression of p107 mRNA and evidence for alternative splicing of the

[57572] p107 (RBL1) gene product. Genomics 28: 520-529, 1995.

[57573] 9432.

[57574] LeCouter, J. E.; Kablar, B.; Hardy, W. R.; Ying, C.; Megeney, L.

[57575] A.; May, L. L.; Rudnicki, M. A.: Strain-dependent myeloid hyperplasia,

[57576] growth deficiency, and accelerated cell cycle in mice lacking the

[57577] Rb-related p107 gene. Molec. Cell. Biol. 18: 7455-7465, 1998.

[57578] 9433.

[57579] Bonner, T. I.; Modi, W. S.; Seuanez, H. N.; O'Brien, S. J.: Chromosomal

[57580] mapping of five human genes encoding muscarinic acetyl-choline receptors.

[57581] (Abstract) Cytogenet. Cell Genet. 58: 1850-1851, 1991.

[57582] 9434.

[57583] Anand, R.; Lindstrom, J.: Chromosomal localization of seven neuronal

[57584] nicotinic acetylcholine receptor subunit genes in humans. Genomics 13:

[57585] 962-967, 1992.

[57586] 9435.

[57587] Bessis, A.; Simon-Chazottes, D.; Devillers-Thiery, A.; Guenet,

[57588] J.-L.; Changeux, J.-P.: Chromosomal localization of the mouse genes

[57589] coding for alpha-2, alpha-3, alpha-4 and beta-2 subunits of neuronal

[57590] nicotinic acetylcholine receptor. FEBS Lett. 264: 48-52, 1990.

[57591] 9436.

[57592] Wilcox, W. R.; Tavormina, P. L.; Krakow, D.; Kitoh, H.; Lachman,

[57593] R. S.; Wasmuth, J. J.; Thompson, L. M.; Rimoin, D. L.: Molecular,

[57594] radiologic, and histopathologic correlations in thanatophoric dysplasia. Am.

[57595] J. Med. Genet. 78: 274-281, 1998.

[57596] 9437.

[57597] Wilkie, A. O. M.: Craniosynostosis: genes and mecha-nisms. Hum.

[57598] Molec. Genet. 6: 1647-1656, 1997.

[57599] 9438.

[57600] Armstrong, E.; Partanen, J.; Cannizzaro, L.; Huebner, K.; Alitalo,

[57601] K.: Localization of the fibroblast growth factor receptor-4 gene

[57602] to chromosome region 5q33-qter. Genes Chromosomes Cancer 4: 94-98,

[57603] 1992.

[57604] 9439.

[57605] Bange, J.; Prechtl, D.; Cheburkin, Y.; Specht, K.; Harbeck, N.;
[57606] Schmitt, M.; Knyazeva, T.; Muller, S.; Gartner, S.; Sures, I.; Wang,
[57607] H.; Imyanitov, E.; Haring, H.-U.; Knayzev, P.; Iacobelli, S.; Hofler,
[57608] H.; Ullrich, A.: Cancer progression and tumor cell motility are associated
[57609] with the FGFR4 Arg388 allele. Cancer Res. 62: 840-847, 2002.
[57610] 9440.

[57611] Holtrich, U.; Brauninger, A.; Strebhardt, K.; Rubsamen-Waigmann,
[57612] H.: Two additional protein-tyrosine kinases expressed in human lung:
[57613] fourth member of the fibroblast growth factor receptor family and
[57614] an intracellular protein-tyrosine kinase. Proc. Nat. Acad. Sci. 88:
[57615] 10411-10415, 1991.

[57616] 9441.

[57617] Kostrzewa, M.; Muller, U.: Genomic structure and com-

plete sequence
[57618] of the human FGFR4 gene. Mammalian Genome 9:
131-135, 1998.
[57619] 9442.

[57620] Partanen, J.; Makela, T. P.; Eerola, E.; Korhonen, J.; Hirvo-
nen,
[57621] H.; Claesson-Welsh, L.; Alitalo, K.: FGFR-4, a novel acidic
fibroblast
[57622] growth factor receptor with a distinct expression pattern.
EMBO J. 10:
[57623] 1347-1354, 1991.

[57624] 9443.

[57625] Scott, A. F.: Personal Communication. Baltimore, Md.
10/12/1999.
[57626] 9444.

[57627] Vainikka, S.; Partanen, J.; Bellosta, P.; Coulier, F.; Basilico,

[57628] C.; Jaye, M.; Alitalo, K.: Fibroblast growth factor receptor-
4 shows
[57629] novel features in genomic structure, ligand binding and
signal transduction. EMBO
[57630] J. 11: 4273-4280, 1992.

[57631] 9445.

[57632] Warrington, J. A.; Bailey, S. K.; Armstrong, E.; Aprelikova, O.;

[57633] Alitalo, K.; Dolganov, G. M.; Wilcox, A. S.; Sikela, J. M.; Wolfe,

[57634] S. F.; Lovett, M.; Wasmuth, J. J.: A radiation hybrid map of 18 growth

[57635] factor, growth factor receptor, hormone receptor, or neu-rotransmitter

[57636] receptor genes on the distal region of the long arm of chromosome

[57637] 5. Genomics 13: 803-808, 1992.

[57638] 9446.

[57639] Celik, M.; Gokmen, N.; Erbayraktar, S.; Akhisaroglu, M.; Konakc,

[57640] S.; Ulukus, C.; Genc, S.; Genc, K.; Sagiroglu, E.; Cerami, A.; Brines,

[57641] M.: Erythropoietin prevents motor neuron apoptosis and neurologic

[57642] disability in experimental spinal cord ischemic injury. Proc. Nat.

[57643] Acad. Sci. 99: 2258-2263, 2002.

[57644] 9447.

[57645] Digicaylioglu, M.; Lipton, S. A.: Erythropoietin-mediated neuroprotection

[57646] involves cross-talk between Jak2 and NF-kappa-B signalling cascades. Nature 412:

[57647] 641-647, 2001.

[57648] 9448.

[57649] Eschbach, J. W.; Egrie, J. C.; Downing, M. R.; Browne, J. K.; Adamson,

[57650] J. W.: Correction of the anemia of end-stage renal disease with recombinant

[57651] human erythropoietin: results of a combined phase I and II clinical

[57652] trial. New Eng. J. Med. 316: 73-78, 1987.

[57653] 9449.

[57654] Goldberg, M. A.; Dunning, S. P.; Bunn, H. F.: Regulation of the

[57655] erythropoietin gene: evidence that the oxygen sensor is a heme protein. Science 242:

[57656] 1412-1415, 1988.

[57657] 9450.

[57658] Gorio, A.; Gokmen, N.; Erbayraktar, S.; Yilmaz, O.; Madaschi, L.;

[57659] Cichetti, C.; Di Giulio, A. M.; Vardar, E.; Cerami, A.; Brines, M.

[57660] : Recombinant human erythropoietin counteracts secondary injury and

[57661] markedly enhances neurological recovery from experimental spinal cord

[57662] trauma. Proc. Nat. Acad. Sci. 99: 9450-9455, 2002.

[57663] 9451.

[57664] Grimm, C.; Wenzel, A.; Groszer, M.; Mayser, H.; Seeliger, M.; Samardzija,

[57665] M.; Bauer, C.; Gassmann, Z.; Reme, C. E.: HIF-1-induced erythropoietin

[57666] in the hypoxic retina protects against light-induced retinal degeneration. Nature

[57667] Med. 8: 718-724, 2002.

[57668] 9452.

[57669] Jacobs, K.; Shoemaker, C.; Rudersdorf, R.; Neill, S. D.; Kaufman,

[57670] R. J.; Mufson, A.; Seehra, J.; Jones, S. S.; Hewick, R.; Fritsch,

[57671] E. F.; Kawakita, M.; Shimizu, T.; Miyake, T.: Isolation and characterization

[57672] of genomic and cDNA clones of human erythropoietin. Nature 313:

[57673] 806-810, 1985.

[57674] 9453.

[57675] Junk, A. K.; Mammis, A.; Savitz, S. I.; Singh, M.; Roth, S.; Malhotra,

[57676] S.; Rosenbaum, P. S.; Cerami, A.; Brines, M.; Rosenbaum, D. M.: Erythropoietin

[57677] administration protects retinal neurons from acute ischemia-reperfusion

[57678] injury. Proc. Nat. Acad. Sci. 99: 10659-10664, 2002.

[57679] 9454.

[57680] Kessler, P. D.; Podsakoff, G. M.; Chen, X.; McQuiston, S. A.; Colosi,

[57681] P. C.; Matelis, L. A.; Kurtzman, G. J.; Byrne, B. J.: Gene delivery

[57682] to skeletal muscle results in sustained expression and systemic delivery

[57683] of a therapeutic protein. Proc. Nat. Acad. Sci. 93: 14082-14087,

[57684] 1996.

[57685] 9455.

[57686] Lacombe, C.; Tambourin, P.; Mattei, M. G.; Simon, D.; Guenet,

[57687] J. L.: The murine erythropoietin gene is localized on chromosome

[57688] 5. (Letter) Blood 72: 1440-1442, 1988.

[57689] 9456.

[57690] Law, M. L.; Cai, G.-Y.; Lin, F.-K.; Wei, Q.; Huang, S.-Z.; Hartz,

[57691] J. H.; Morse, H.; Lin, C.-H.; Jones, C.; Kao, F.-T.: Chromosomal

[57692] assignment of the human erythropoietin gene and its DNA polymorphism. Proc.

[57693] Nat. Acad. Sci. 83: 6920-6924, 1986.

[57694] 9457.

[57695] Lee, R.; Kertesz, N.; Joseph, S. B.; Jegalian, A. Wu, H.: Erythropoietin

[57696] (Epo) and EpoR expression and 2 waves of erythropoiesis. Blood 98:

[57697] 1408-1415, 2001.

[57698] 9458.

[57699] Lee-Huang, S.: Cloning and expression of human erythro-
poietin
[57700] cDNA in Escherichia coli. Proc. Nat. Acad. Sci. 81:
2708-2712, 1984.
[57701] 9459.

[57702] Lin, F.-K.; Suggs, S.; Lin, C.-H.; Browne, J. K.; Smalling, R.;

[57703] Egrie, J. C.; Chen, K. K.; Fox, G. M.; Martin, F.; Stabinsky,
Z.;
[57704] Badrawi, S. M.; Lai, P.-H.; Goldwasser, E.: Cloning and ex-
pression
[57705] of the human erythropoietin gene. Proc. Nat. Acad. Sci.
82: 7580-7584,
[57706] 1985.

[57707] 9460.

[57708] Macdougall, I. C.; Gray, S. J.; Elston, O.; Breen, C.; Jenkins,

[57709] B.; Browne, J.; Egrie, J.: Pharmacokinetics of novel erythro-
poiesis
[57710] stimulating protein compared with Epoetin alfa in dialysis
patients. J.
[57711] Am. Soc. Nephrol. 10: 2392-2395, 1999.

[57712] 9461.

[57713] McDonald, J. D.; Lin, F.-K.; Goldwasser, E.: Cloning, se-
quencing,
[57714] and evolutionary analysis of the mouse erythropoietin
gene. Molec.
[57715] Cell. Biol. 6: 842-848, 1986.

[57716] 9462.

[57717] Naffakh, N.; Henri, A.; Villeval, J. L.; Rouyer-Fessard, P.;
Moullier,
[57718] P.; Blumenfeld, N.; Danos, O.; Vainchenker, W.; Heard, J.
M.; Beuzard,
[57719] Y.: Sustained delivery of erythropoietin in mice by geneti-
cally modified
[57720] skin fibroblasts. Proc. Nat. Acad. Sci. 92: 3194-3198,
1995.
[57721] 9463.

[57722] Osborne, W. R. A.; Ramesh, N.; Lau, S.; Clowes, M. M.;
Dale, D.
[57723] C.; Clowes, A. W.: Gene therapy for long-term expression
of erythropoietin
[57724] in rats. Proc. Nat. Acad. Sci. 92: 8055-8058, 1995.

[57725] 9464.

[57726] Powell, J. S.; Berkner, K. L.; Lebo, R. V.; Adamson, J. W.: Human

[57727] erythropoietin gene: high level expression in stably transfected mammalian

[57728] cells and chromosome localization. Proc. Nat. Acad. Sci. 83: 6465-6469,

[57729] 1986.

[57730] 9465.

[57731] Romanowski, R. R.; Sytkowski, A. J.: The molecular structure

[57732] of human erythropoietin. Hemat. Oncol. Clin. North Am. 8: 885-894,

[57733] 1994.

[57734] 9466.

[57735] Sakanaka, M.; Wen, T.-C.; Matsuda, S.; Masuda, S.; Morishita,

[57736] E.; Nagao, M.; Sasaki, R.: In vivo evidence that erythropoietin protects

[57737] neurons from ischemic damage. Proc. Nat. Acad. Sci. 95: 4635-4640,

[57738] 1998.

[57739] 9467.

[57740] Semenza, G. L.; Dureza, R. C.; Traystman, M. D.; Gearhart,
J.

[57741] D.; Antonarakis, S. E.: Different DNA sequences control
liver versus

[57742] kidney expression of the human erythropoietin gene in
transgenic mice.

[57743] (Abstract) Am. J. Hum. Genet. 45 (suppl.): A116 only,
1989.

[57744] 9468.

[57745] Semenza, G. L.; Traystman, M. D.; Gearhart, J. D.; An-
tonarakis,

[57746] S. E.: Polycythemia in transgenic mice expressing the hu-
man erythropoietin

[57747] gene. Proc. Nat. Acad. Sci. 86: 2301-2305, 1989.

[57748] 9469.

[57749] Sherwood, J. B.; Shouval, D.: Continuous production of
erythropoietin

[57750] by an established human renal carcinoma cell line: devel-
opment of

[57751] the cell line. Proc. Nat. Acad. Sci. 83: 165-169, 1986.

[57752] 9470.

[57753] Shoemaker, C. B.; Mitsock, L. D.: Murine erythropoietin gene:

[57754] cloning, expression, and human gene homology. Molec. Cell. Biol. 6:

[57755] 849-858, 1986.

[57756] 9471.

[57757] Siren, A.-L.; Fratelli, M.; Brines, M.; Goemans, C.; Casagrande,

[57758] S.; Lewczuk, P.; Keenan, S.; Gleiter, C.; Pasquali, C.; Capo-

bianco,

[57759] A.; Mennini, T.; Heumann, R.; Cerami, A.; Ehrenreich, H.; Ghezzi,

[57760] P.: Erythropoietin prevents neuronal apoptosis after cere-

bral ischemia

[57761] and metabolic stress. Proc. Nat. Acad. Sci. 98: 4044-4049, 2001.

[57762] 9472.

[57763] Clancy, D. J.; Gems, D.; Harshman, L. G.; Oldham, S.; Stocker,

[57764] H.; Hafen, E.; Leevers, S. J.; Partridge, L.: Extension of life-span
[57765] by loss of CHICO, a Drosophila insulin receptor substrate protein. Science 292:
[57766] 104-106, 2001.

[57767] 9473.

[57768] Chai, K. X.; Chen, L.-M.; Chao, J.; Chao, L.: Kallistatin: a novel
[57769] human serine proteinase inhibitor: molecular cloning, tis-sue distribution,
[57770] and expression in Escherichia coli. J. Biol. Chem. 268: 24498-24505,
[57771] 1993.

[57772] 9474.

[57773] Chai, K. X.; Ward, D. C.; Chao, J.; Chao, L.: Molecular cloning,
[57774] sequence analysis, and chromosomal localization of the human protease
[57775] inhibitor 4 (kallistatin) gene (PI4). Genomics 23: 370-378, 1994.
[57776] 9475.

[57777] Zhou, G. X.; Chao, L.; Chao, J.: Kallistatin: a novel human tissue
[57778] kallikrein inhibitor; purification, characterization, and reactive
[57779] center sequence. J. Biol. Chem. 267: 25873-25880, 1992.

[57780] 9476.

[57781] Christmanson, L.; Rorsman, F.; Stenman, G.; Westermark, P.; Betsholtz,
[57782] C.: The human islet amyloid polypeptide (IAPP) gene: organization,
[57783] chromosomal localization and functional identification of a promoter
[57784] region. FEBS Lett. 267: 160-166, 1990.

[57785] 9477.

[57786] Cockburn, D. C.; Holt, S. M.; Roberts, A. N.; Cooper, G. J. S.;
[57787] Reid, K. B. M.; Boyd, Y.: Localization of the amylin locus to chromosome
[57788] 12. (Abstract) Cytogenet. Cell Genet. 51: 977, 1989.

[57789] 9478.

[57790] Cooper, G. J.; Willis, A. C.; Clark, A.; Turner, R. C.; Sim, R.

[57791] B.; Reid, K. B.: Purification and characterization of a pep-
tide from
[57792] amyloid-rich pancreases of type 2 diabetic patients. Proc.
Nat. Acad.
[57793] Sci. 84: 8628-8632, 1987.

[57794] 9479.

[57795] Cooper, G. J. S.; Leighton, B.; Dimitriadis, G. D.; Parry-
Billings,
[57796] M.; Kowalchuk, J. M.; Howland, K.; Rothbard, J. B.; Willis,
A. C.;
[57797] Reid, K. B. M.: Amylin found in amyloid deposits in human
type 2
[57798] diabetes mellitus may be a hormone that regulates glyco-
gen metabolism
[57799] in skeletal muscle. Proc. Nat. Acad. Sci. 85: 7763-7766,
1988.
[57800] 9480.

[57801] Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[57802] M.; Sanke, T.; Steiner, D. F.; Shows, T. B.; Bell, G. I.: Local-
ization
[57803] of the human islet amyloid polypeptide gene (IAP) to
chromosome 12p12.3.

[57804] (Abstract) Cytogenet. Cell Genet. 51: 997, 1989.

[57805] 9481.

[57806] Harvey, A. M.: Medical students on the march: Brown, MacCallum

[57807] and Opie. Johns Hopkins Med. J. 134: 330-345, 1974.

[57808] 9482.

[57809] Hoovers, J. M. N.; Redeker, E.; Speleman, F.; Hoppener, J. W. M.;

[57810] Bhola, S.; Bliek, J.; van Roy, N.; Leschot, N. J.; Westerveld, A.;

[57811] Mannens, M.: High-resolution chromosomal localization of the human

[57812] calcitonin/CGRP/IAPP gene family members. Genomics 15: 525-529,

[57813] 1993.

[57814] 9483.

[57815] Hoppener, J. W. M.; Oosterwijk, C.; van Hulst, K. L.; Ver-beek,

[57816] J. S.; Capel, P. J. A.; de Koning, E. J. P.; Clark, A.; Jansz, H.

[57817] S.; Lips, C. J. M.: Molecular physiology of the islet amyloid polypeptide

[57818] (IAPP)/amylin gene in man, rat, and transgenic mice. J. Cell. Biochem. 55S:

[57819] 39-53, 1994.

[57820] 9484.

[57821] Jaikaran, E. T. A. S.; Clark, A.: Islet amyloid and type 2 diabetes:

[57822] from molecular misfolding to islet pathophysiology. Biochim. Biophys.

[57823] Acta 1537: 179-203, 2001.

[57824] 9485.

[57825] Janson, J.; Soeller, W. C.; Roche, P. C.; Nelson, R. T.; Torchia,

[57826] A. J.; Kreutter, D. K.; Butler, P. C.: Spontaneous diabetes mellitus

[57827] in transgenic mice expressing human islet amyloid polypeptide. Proc.

[57828] Nat. Acad. Sci. 93: 7283-7288, 1996.

[57829] 9486.

[57830] Johnson, K. H.; O'Brien, T. D.; Betsholtz, C.; Westermark, P.

[57831] : Islet amyloid, islet-amyloid polypeptide, and diabetes

mellitus. New

[57832] Eng. J. Med. 321: 513-518, 1989.

[57833] 9487.

[57834] Lorenzo, A.; Razzaboni, B.; Weir, G. C.; Yankner, B. A.: Pancreatic

[57835] islet cell toxicity of amylin associated with type-2 diabetes mellitus. Nature 368:

[57836] 756-760, 1994.

[57837] 9488.

[57838] Mosselman, S.; Hoppener, J. W. M.; Lips, C. J. M.; Jansz, H. S.

[57839] : The complete islet amyloid polypeptide precursor is encoded by two

[57840] exons. FEBS Lett. 247: 154-158, 1989.

[57841] 9489.

[57842] Mosselman, S.; Hoppener, J. W. M.; Zandberg, J.; van Mansfeld,

[57843] A. D. M.; Geurts van Kessel, A. H. M.; Lips, C. J. M.; Jansz, H. S.

[57844] : Islet amyloid polypeptide: identification and chromosomal localization

[57845] of the human gene. FEBS Lett. 239: 227-232, 1988.

[57846] 9490.

[57847] Nishi, M.; Sanke, T.; Seino, S.; Eddy, R. L.; Fan, Y.-S.; By-
ers,
[57848] M. G.; Shows, T. B.; Bell, G. I.; Steiner, D. F.: Human islet
amyloid
[57849] polypeptide gene: complete nucleotide sequence, chro-
mosomal localization,
[57850] and evolutionary history. Molec. Endocr. 3: 1775-1781,
1989.
[57851] 9491.

[57852] Opie, E. L.: On the relation of the chronic interstitial pan-
creatitis
[57853] to the islands of Langerhans and to diabetes mellitus. J.
Exp. Med. 5:
[57854] 397-428, 1900.

[57855] 9492.

[57856] Bergsagel, P. L.; Chesi, M.; Nardini, E.; Brents, L. A.; Kirby,

[57857] S. L.; Kuehl, W. M.: Promiscuous translocations into im-
munoglobulin
[57858] heavy chain switch regions in multiple myeloma. Proc.

Nat. Acad.
[57859] Sci. 93: 13931-13936, 1996.

[57860] 9493.

[57861] Breitbart, A. S.; Eaton, C.; McCarthy, J. G.: Crouzon's syn-
drome
[57862] associated with acanthosis nigricans: ramifications for the
craniofacial
[57863] surgeon. Ann. Plastic Surg. 22: 310-315, 1989.

[57864] 9494.

[57865] Brodie, S. G.; Kitoh, H.; Lipson, M.; Sifry-Platt, M.; Wilcox,

[57866] W. R.: Thanatophoric dysplasia type I with syndactyly. Am.
J. Med.
[57867] Genet. 80: 260-262, 1998.

[57868] 9495.

[57869] Cappellen, D.; De Oliveira, C.; Ricol, D.; Gil Diez de Med-
ina,
[57870] S.; Bourdin, J.; Sastre-Garau, X.; Chopin, D.; Thiery, J. P.;
Radvanyi,
[57871] F.: Frequent activating mutations of FGFR3 in human
bladder and cervix
[57872] carcinomas. (Letter) Nature Genet. 23: 18-20, 1999.

[57873] 9496.

[57874] Chen, L.; Adar, R.; Yang, X.; Monsonego, E. O.; Li, C.; Hauschka,

[57875] P. V.; Yayon, A.; Deng, C.-X.: Gly369-to-cys mutation in mouse FGFR3

[57876] causes achondroplasia by affecting both chondrogenesis and osteogenesis. J.

[57877] Clin. Invest. 104: 1517-1525, 1999.

[57878] 9497.

[57879] Chen, L.; Li, C.; Qiao, W.; Xu, X.; Deng, C.: A ser365-cys mutation

[57880] of fibroblast growth factor receptor 3 in mouse downregulates Ihh/PTHrP

[57881] signals and causes severe achondroplasia. Hum. Molec. Genet. 10:

[57882] 457-465, 2001.

[57883] 9498.

[57884] Chesi, M.; Nardini, E.; Brents, L. A.; Schrock, E.; Ried, T.;

[57885] Kuehl, W. M.; Bergsagel, P. L.: Frequent translocation t(4;14)(p16.3;q32.3)

[57886] in multiple myeloma is associated with increased expres-

sion and activating
[57887] mutations of fibroblast growth factor receptor 3. Nature Genet. 16:
[57888] 260-264, 1997.

[57889] 9499.

[57890] Cohen, M. M., Jr.: Let's call it 'Crouzonodermoskeletal syndrome'
[57891] so we won't be prisoners of our own conventional termi-nology. (Letter) Am.
[57892] J. Med. Genet. 84: 74 only, 1999.

[57893] 9500.

[57894] Colvin, J. S.; Bohne, B. A.; Harding, G. W.; McEwen, D. G.; Ornitz,
[57895] D. M.: Skeletal overgrowth and deafness in mice lacking fibroblast
[57896] growth factor receptor 3. Nature Genet. 12: 390-397, 1996.
[57897] 9501.

[57898] Deng, C.; Wynshaw-Boris, A.; Zhou, F.; Kuo, A.; Leder, P.: Fibroblast
[57899] growth factor receptor 3 is a negative regulator of bone

growth. Cell 84:
[57900] 911-921, 1996.

[57901] 9502.

[57902] Deutz-Terlouw, P. P.; Losekoot, M.; Aalfs, C. M.; Hen-
nekam, R.
[57903] C. M.; Bakker, E.: Asn540thr substitution in the fibroblast
growth
[57904] factor receptor 3 tyrosine kinase domain causing
hypochondroplasia. Hum.
[57905] Mutat. Suppl. 1: S62-S65, 1998.

[57906] 9503.

[57907] El Ghouzzi, V.; Lajeunie, E.; Le Merrer, M.; Cormier-Daire,
V.;
[57908] Renier, D.; Munnich, A.; Bonaventure, J.: Mutations within
or upstream
[57909] of the basic helix-loop-helix domain of the TWIST gene
are specific
[57910] to Saethre-Chotzen syndrome. Europ. J. Hum. Genet. 7:
27-33, 1999.
[57911] 9504.

[57912] Fofanova, O. V.; Takamura, N.; Kinoshita, E.; Meerson, E.

M.;

[57913] Iljina, V. K.; Nechvolodova, O. L.; Evgrafov, O. V.; Peterkova, V.

[57914] A.; Yamashita, S.: A missense mutation of C1659 in the fibroblast

[57915] growth factor receptor 3 gene in Russian patients with hypochondroplasia. Endocr.

[57916] J. 45: 791-795, 1998.

[57917] 9505.

[57918] Francomano, C. A.: Personal Communication. Bethesda, Md. 1/19/1999.

[57919] 9506.

[57920] Francomano, C. A.: Personal Communication. Baltimore, Md.

[57921] 12/1/1994.

[57922] 9507.

[57923] Golla, A.; Lichtner, P.; von Gernet, S.; Winterpacht, A.; Fairley,

[57924] J.; Murken, J.; Schuffenhauer, S.: Phenotypic expression of the fibroblast

[57925] growth factor receptor 3 (FGFR3) mutation P250R in a

large craniosynostosis
[57926] family. J. Med. Genet. 34: 683-684, 1997.

[57927] 9508.

[57928] Graham, J. M., Jr.; Braddock, S. R.; Mortier, G. R.; Lachman,

[57929] R.; Van Dop, C.; Jabs, E. W.: Syndrome of coronal cran-
iosynostosis
[57930] with brachydactyly and carpal tarsal coalition due to
Pro250Arg mutation
[57931] in FGFR3 gene. Am. J. Med. Genet. 77: 322-329, 1998.

[57932] 9509.

[57933] Grigelioniene, G.; Hagenas, L.; Eklof, O.; Neumeyer, L.;
Haereid,
[57934] P. E.; Anvret, M.: A novel missense mutation ile538-to-val
in the
[57935] fibroblast growth factor receptor 3 in hypochondroplasia.
(Abstract) Hum.
[57936] Mutat. 11: 333 only, 1998.

[57937] 9510.

[57938] Gripp, K. W.; McDonald-McGinn, D. M.; Gaudenz, K.;
Whitaker, L.
[57939] A.; Bartlett, S. P.; Glat, P. M.; Cassileth, L. B.; Mayro, R.;

Zackai,
[57940] E. H.; Muenke, M.: Identification of a genetic cause for isolated
[57941] unilateral coronal synostosis: a unique mutation in the fibroblast
[57942] growth factor receptor 3. J. Pediat. 132: 714-716, 1998.

[57943] 9511.

[57944] Francomano, C. A.; Bellus, G. A.; Szabo, J.; McIntosh, I.; Dorst,
[57945] J.; Lee, R.; Hurko, O.; Fraley, A. E.; Bamshad, M. J.: A new skeletal
[57946] dysplasia with severe tibial bowing, profound developmental delay
[57947] and acanthosis nigricans is caused by a Lys 650 Met mutation in fibroblast
[57948] growth factor receptor 3 (FGFR3). (Abstract) Am. J. Hum. Genet. 59
[57949] (suppl.): A25 only, 1996.

[57950] 9512.

[57951] Henderson, J. E.; Naski, M. C.; Aarts, M. M.; Wang, D.; Cheng,
[57952] L.; Goltzman, D.; Ornitz, D. M.: Expression of FGFR3 with

the G380R

[57953] achondroplasia mutation inhibits proliferation and matu-
ration of CFK2

[57954] chondrocytic cells. J. Bone Miner. Res. 15: 155-165, 2000.

[57955] 9513.

[57956] Hollway, G. E.; Suthers, G. K.; Battese, K. M.; Turner, A. M.;

[57957] David, D. J.; Mulley, J. C.: Deafness due to pro250-to-arg
mutation

[57958] of FGFR3. (Letter) Lancet 351: 877-878, 1998.

[57959] 9514.

[57960] Ikegawa, S.; Fukushima, Y.; Isomura, M.; Takada, F.; Naka-
mura,

[57961] Y.: Mutations of the fibroblast growth factor receptor-3
gene in

[57962] one familial and six sporadic cases of achondroplasia in
Japanese

[57963] patients. Hum. Genet. 96: 309-311, 1995.

[57964] 9515.

[57965] Intini, D.; Baldini, L.; Fabris, S.; Lombardi, L.; Ciceri, G.;

[57966] Maiolo, A. T.; Neri, A.: Analysis of FGFR3 gene mutations

in multiple
[57967] myeloma patients with t(4;14). Brit. J. Haemat. 114: 362-364, 2001.
[57968] 9516.

[57969] Iwata, T.; Chen, L.; Li, C.; Ovchinnikov, D. A.; Behringer, R.

[57970] R.; Francomano, C. A.; Deng, C.-X.: A neonatal lethal mutation in
[57971] FGFR3 uncouples proliferation and differentiation of growth plate
[57972] chondrocytes in embryos. Hum. Molec. Genet. 9: 1603-1613, 2000.
[57973] 9517.

[57974] Iwata, T.; Li, C.-L.; Deng, C.-X.; Francomano, C. A.: Highly

[57975] activated Fgfr3 with the K644M mutation causes prolonged survival
[57976] in severe dwarf mice. Hum. Molec. Genet. 10: 1255-1264, 2001.
[57977] 9518.

[57978] Jabs, E. W.; Li, X.; Scott, A. F.; Meyers, G.; Chen, W.; Eccles,

[57979] M.; Mao, J. I.; Charnas, L. R.; Jackson, C. E.; Jaye, M.: Jackson-Weiss

[57980] and Crouzon syndromes are allelic with mutations in fibroblast growth
[57981] factor receptor 2. Nature Genet. 8: 275-279, 1994.

[57982] 9519.

[57983] Jang, J.-H.: Identification and characterization of soluble isoform
[57984] of fibroblast growth factor receptor 3 in human SaOS-2 osteosarcoma
[57985] cells. Biochem. Biophys. Res. Commun. 292: 378-382, 2002.
[57986] 9520.

[57987] Jang, J.-H.; Shin, K.-H.; Park, J.-G.: Mutations in fibroblast
[57988] growth factor receptor 2 and fibroblast growth factor receptor 3 genes
[57989] associated with human gastric and colorectal cancers. Cancer Res. 61:
[57990] 3541-3543, 2001.

[57991] 9521.

[57992] Clausen, J. O.; Hansen, T.; Bjorbaek, C.; Echwald, S. M.; Urhammer,
[57993] S. A.; Rasmussen, S.; Andersen, C. B.; Hansen, L.; Almind,

K.; Winther,
[57994] K.; Haraldsdottir, J.; Borch-Johnsen, K.; Pedersen, O.: In-
sulin resistance:
[57995] interactions between obesity and a common variant of in-
sulin receptor
[57996] substrate-1. Lancet 346: 397-402, 1995.

[57997] 9522.

[57998] Esposito, D. L.; Mammarella, S.; Ranieri, A.; Loggia, F. D.;
Capani,
[57999] F.; Consoli, A.; Mariani-Costantini, R.; Caramia, F. G.;
Cama, A.;
[58000] Battista, P.: Deletion of gly723 in the insulin receptor sub-
strate-1
[58001] of a patient with noninsulin-dependent diabetes mellitus.
Hum. Mutat. 7:
[58002] 364-366, 1996.

[58003] 9523.

[58004] Hribal, M. L.; Federici, M.; Porzio, O.; Lauro, D.; Borboni,
P.;
[58005] Accili, D.; Lauro, R.; Sesti, G.: The gly-to-arg(972) amino
acid
[58006] polymorphism in insulin receptor substrate-1 affects glu-

cose metabolism
[58007] in skeletal muscle cells. J. Clin. Endocr. Metab. 85: 2004-2013,
[58008] 2000.

[58009] 9524.

[58010] Kido, Y.; Burks, D. J.; Withers, D.; Bruning, J. C.; Kahn, C.
[58011] R.; White, M. F.; Accili, D.: Tissue-specific insulin resistance
[58012] in mice with mutations in the insulin receptor, IRS-1, and IRS-2. J.
[58013] Clin. Invest. 105: 199-205, 2000.

[58014] 9525.

[58015] Kulkarni, R. N.; Winnay, J. N.; Daniels, M.; Bruning, J. C.; Flier,
[58016] S. N.; Hanahan, D.; Kahn, C. R.: Altered function of insulin receptor
[58017] substrate-1-deficient mouse islets and cultured beta-cell lines. J.
[58018] Clin. Invest. 104: R69-R75, 1999.

[58019] 9526.

[58020] Laakso, M.; Malkki, M.; Kekalainen, P.; Kuusisto, J.; Deeb,

S.

[58021] S.: Insulin receptor substrate-1 variants in non-in-
sulin-dependent

[58022] diabetes. J. Clin. Invest. 94: 1141-1146, 1994.

[58023] 9527.

[58024] Myers, M. G., Jr.; Sun, X. J.; White, M. F.: The IRS-1 signal-
ing

[58025] system. Trends Biochem. Sci. 19: 289-293, 1994.

[58026] 9528.

[58027] Nishiyama, M.; Inazawa, J.; Ariyama, T.; Nakamura, Y.;
Matsufuji,

[58028] S.; Furusaka, A.; Tanaka, T.; Hayashi, S.; Wands, J. R.: The
human

[58029] insulin receptor substrate-1 gene (IRS1) is localized on
2q36. Genomics 20:

[58030] 139-141, 1994.

[58031] 9529.

[58032] Nishiyama, M.; Wands, J. R.: Cloning and increased ex-
pression

[58033] of an insulin receptor substrate-1-like gene in human
hepatocellular

[58034] carcinoma. Biochem. Biophys. Res. Commun. 183: 280-285, 1992.

[58035] 9530.

[58036] Ogata, N.; Chikazu, D.; Kubota, N.; Terauchi, Y.; Tobe, K.; Azuma,

[58037] Y.; Ohta, T.; Kadowaki, T.; Nakamura, K.; Kawaguchi, H.: Insulin

[58038] receptor substrate-1 in osteoblast is indispensable for maintaining

[58039] bone turnover. J. Clin. Invest. 105: 935-943, 2000.

[58040] 9531.

[58041] Ogihara, T.; Isobe, T.; Ichimura, T.; Taoka, M.; Funaki, M.; Sakoda,

[58042] H.; Onishi, Y.; Inukai, K.; Anai, M.; Fukushima, Y.; Kikuchi, M.;

[58043] Yazaki, Y.; Oka, Y.; Asano, T.: 14-3-3 protein binds to in-sulin receptor

[58044] substrate-1, one of the binding sites of which is in the phosphotyrosine

[58045] binding domain. J. Biol. Chem. 272: 25267-25274, 1997.

[58046] 9532.

[58047] Porzio, O.; Federici, M.; Hribal, M. L.; Lauro, D.; Accili, D.;

[58048] Lauro, R.; Borboni, P.; Sesti, G.: The gly972-to-arg amino acid polymorphism

[58049] in IRS-1 impairs insulin secretion in pancreatic beta cells. J. Clin.

[58050] Invest. 104: 357-364, 1999.

[58051] 9533.

[58052] Stoffel, M.; Espinosa, R., III; Keller, S. R.; Lienhard, G. E.;

[58053] Le Beau, M. M.; Bell, G. I.: Human insulin receptor substrate-1 gene

[58054] (IRS1): chromosomal localization to 2q35-q36.1 and identification

[58055] of a simple tandem repeat DNA polymorphism. Diabetologia 36: 335-337,

[58056] 1993.

[58057] 9534.

[58058] Sun, X. J.; Rothenberg, P.; Kahn, C. R.; Backer, J. M.; Araki,

[58059] E.; Wilden, P. A.; Cahill, D. A.; Goldstein, B. J.; White, M. F.:

[58060] Structure of the insulin receptor substrate IRS-1 defines a unique

[58061] signal transduction protein. Nature 352: 73-77, 1991.

[58062] 9535.

[58063] Diaz, M. O.; Bohlander, S.: Nomenclature of the human interferon

[58064] genes. J. Interferon Res. 13: 443-444, 1993.

[58065] 9536.

[58066] Henco, K.; Brosius, J.; Fujisawa, A.; Fujisawa, J.-I.; Haynes,

[58067] J. R.; Hochstadt, J.; Kovacic, T.; Pasek, M.; Schambock, A.; Schmid,

[58068] J.; Todokoro, K.; Walchli, M.; Nagata, S.; Weissmann, C.: Structural

[58069] relationship of human interferon alpha genes and pseudogenes. J.

[58070] Molec. Biol. 185: 227-260, 1985.

[58071] 9537.

[58072] Olopade, O. I.; Bohlander, S. K.; Pomykala, H.; Maltepe, E.; Van

[58073] Melle, E.; Le Beau, M. M.; Diaz, M. O.: Mapping of the shortest region

[58074] of overlap of deletions of the short arm of chromosome 9 associated

[58075] with human neoplasia. Genomics 14: 437-443, 1992.

[58076] 9538.

[58077] Georges-Labouesse, E.; Messaddeq, N.; Yehia, G.; Cadal-
bert, L.;

[58078] Dierich, A.; Le Meur, M.: Absence of integrin alpha-6 leads
to epidermolysis

[58079] bullosa and neonatal death in mice. Nature Genet. 13:
370-373, 1996.

[58080] 9539.

[58081] Hogervorst, F.; Kuikman, I.; Geurts van Kessel, A.; Son-
nenberg,

[58082] A.: Molecular cloning of the human alpha-6 integrin sub-
unit: alternative

[58083] splicing of alpha-6 mRNA and chromosomal localization
of the alpha-6

[58084] and beta-4 genes. Europ. J. Biochem. 199: 425-433,
1991.

[58085] 9540.

[58086] Ruzzi, L.; Gagnoux-Palacios, L.; Pinola, M.; Belli, S.;
Meneguzzi,

[58087] G.; D'Alessio, M.; Zambruno, G.: A homozygous mutation
in the integrin

[58088] alpha-6 gene in junctional epidermolysis bullosa with pyloric atresia. J.

[58089] Clin. Invest. 99: 2826-2831, 1997.

[58090] 9541.

[58091] Ruzzi, L.; Gagnoux-Palacios, L.; Pinola, M.; Belli, S.; Meneguzzi,

[58092] G.; Zambruno, G.; D'Alessio, M.: A homozygous mutation in the gene

[58093] encoding integrin alpha6 in junctional epidermolysis bullosa with

[58094] pyloric atresia. (Abstract) Medizinische Genetik 9: 13-14, 1997.

[58095] 9542.

[58096] Tamura, R. N.; Rozzo, C.; Starr, L.; Chambers, J.; Reichardt, L.

[58097] F.; Cooper, H. M.; Quaranta, V.: Epithelial integrin alpha-6/beta-4:

[58098] complete primary structure of alpha-6 and variant forms of beta-4. J.

[58099] Cell Biol. 111: 1593-1604, 1990.

[58100] 9543.

[58101] Chavanas, S.; Gache, Y.; Vailly, J.; Kanitakis, J.; Pulkkinen,

[58102] L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Splicing modu-
lation

[58103] of integrin beta-4 pre-mRNA carrying a branch point mu-
tation underlies

[58104] epidermolysis bullosa with pyloric atresia undergoing
spontaneous

[58105] amelioration with ageing. Hum. Molec. Genet. 8:
2097-2105, 1999.

[58106] 9544.

[58107] Inoue, M.; Tamai, K.; Shimizu, H.; Owaribe, K.; Nakama,
T.; Hashimoto,

[58108] T.; McGrath, J. A.: A homozygous missense mutation in
the cytoplasmic

[58109] tail of beta-4 integrin, G931D, that disrupts hemidesmo-
some assembly

[58110] and underlies non-Herlitz junctional epidermolysis bul-
losa without

[58111] pyloric atresia? J. Invest. Derm. 114: 1061-1064, 2000.

[58112] 9545.

[58113] Mellerio, J. E.; Pulkkinen, L.; McMillan, J. R.; Lake, B. D.; Horn,

[58114] H. M.; Tidman, M. J.; Harper, J. I.; McGrath, J. A.; Uitto, J.; Eady,

[58115] R. A. J.: Pyloric atresia-junctional epidermolysis bullosa syndrome:

[58116] mutations in the integrin beta-4 gene (ITGB4) in two un-related patients

[58117] with mild disease. Brit. J. Derm. 139: 862-871, 1998.

[58118] 9546.

[58119] Carayannopoulos, M. O.; Chi, M. M.-Y.; Cui, Y.; Pingster-haus, J.

[58120] M.; McKnight, R. A.; Mueckler, M.; Devaskar, S. U.; Moley, K. H.:

[58121] GLUT8 is a glucose transporter responsible for insulin-stimulated

[58122] glucose uptake in the blastocyst. Proc. Nat. Acad. Sci. 97: 7313-7318,

[58123] 2000.

[58124] 9547.

[58125] Doege, H.; Schurmann, A.; Bahrenberg, G.; Brauers, A.; Joost, H.

[58126] G.: GLUT8, a novel member of the sugar transport facilitator family

[58127] with glucose transport activity. J. Biol. Chem. 275: 16275-16280,

[58128] 2000.

[58129] 9548.

[58130] Ibberson, M.; Uldry, M.; Thorens, B.: GLUTX1, a novel mammalian

[58131] glucose transporter expressed in the central nervous system and insulin-sensitive

[58132] tissues. J. Biol. Chem. 275: 4607-4612, 2000.

[58133] 9549.

[58134] Kool, M.; de Haas, M.; Scheffer, G. L.; Scheper, R. J.; van Eijk,

[58135] M. J.; Juijn, J. A.; Baas, F.; Borst, P.: Analysis of expression

[58136] of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the multidrug

[58137] resistance-associated protein gene (MRP1), in human cancer cell lines. Cancer

[58138] Res. 57: 3537-3547, 1997.

[58139] 9550.

[58140] Mahlknecht, U.; Schnittger, S.; Ottmann, O. G.; Schoch, C.; Mosebach,

[58141] M.; Hiddemann, W.; Hoelzer, D.: Chromosomal organization and localization

[58142] of the human histone deacetylase 5 gene (HDAC5). Biochim. Biophys.

[58143] Acta 1493: 342-348, 2000.

[58144] 9551.

[58145] McKinsey, T. A.; Zhang, C.-L.; Lu, J.; Olson, E. N.: Signal-dependent

[58146] nuclear export of a histone deacetylase regulates muscle differentiation. Nature 408:

[58147] 106-111, 2000.

[58148] 9552.

[58149] Chesney, J.; Mitchell, R.; Benigni, F.; Bacher, M.; Spiegel, L.;

[58150] Al-Abed, Y.; Han, J. H.; Metz, C.; Bucala, R.: An inducible gene

[58151] product for 6-phosphofructo-2-kinase with an AU-rich

instability element:
[58152] role in tumor cell glycolysis and the Warburg effect. Proc. Nat.
[58153] Acad. Sci. 96: 3047-3052, 1999.

[58154] 9553.

[58155] Manzano, A.; Rosa, J. L.; Ventura, F.; Perez, J. X.; Nadal, M.;
[58156] Estivill, X.; Ambrosio, S.; Gil, J.; Bartrons, R.: Molecular cloning,
[58157] expression, and chromosomal localization of a ubiqui-tously expressed
[58158] human 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene (PFKFB3). Cytogenet.
[58159] Cell Genet. 83: 214-217, 1998.

[58160] 9554.

[58161] Nicholl, J.; Hamilton, J. A.; Sutherland, G. R.; Sutherland, R.
[58162] L.; Watts, C. K. W.: The third human isoform of 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase
[58163] (PFKFB3) map position 10p14-p15. Chromosome Res. 5: 150 only, 1997.

[58164] 9555.

[58165] Sakai, A.; Kato, M.; Fukasawa, M.; Ishiguro, M.; Furuya, E.; Sakakibara,

[58166] R.: Cloning of cDNA encoding for a novel isozyme of fructose 6-phosphate,2-kinase/fructose

[58167] 2,6-bisphosphatase from human placenta. J. Biochem. 119: 506-511,

[58168] 1996.

[58169] 9556.

[58170] Sakakibara, R.; Kato, M.; Okamura, N.; Nakagawa, T.; Komada, Y.;

[58171] Tominaga, N.; Shimojo, M.; Fukasawa, M.: Characterization of a human

[58172] placental fructose-6-phosphate,2-kinase/fructose-2,6-bisphosphatase. J.

[58173] Biochem. 122: 122-128, 1997.

[58174] 9557.

[58175] Manzano, A.; Perez, J. X.; Nadal, M.; Estivill, X.; Lange, A.;

[58176] Bartrons, R.: Cloning, expression and chromosomal localization of

[58177] a human testis

6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase
[58178] gene. Gene 229: 83-89, 1999.

[58179] 9558.

[58180] Larsen, M.; Ressler, S. J.; Gerdes, M. J.; Lu, B.; Byron, M.; Lawrence,
[58181] J. B.; Rowley, D. R.: The WFDC1 gene encoding ps20 local-izes to 16q24,
[58182] a region of LOH in multiple cancers. Mammalian Genome 11: 767-773,
[58183] 2000.

[58184] 9559.

[58185] Larsen, M.; Ressler, S. J.; Lu, B.; Gerdes, M. J.; McBride, L.;

[58186] Dang, T. D.; Rowley, D. R.: Molecular cloning and expres-sion of ps20
[58187] growth inhibitor: a novel WAP-type 'four-disulfide core' domain protein
[58188] expressed in smooth muscle. J. Biol. Chem. 273: 4574-4584, 1998.
[58189] 9560.

[58190] Rowley, D. R.: Characterization of a fetal urogenital sinus mesenchymal

[58191] cell line U4F: secretion of a negative growth factor regulatory activity. In

[58192] Vitro Cell. Dev. Biol. 28A: 29-38, 1992.

[58193] 9561.

[58194] Rowley, D. R.; Dang, T. D.; Larsen, M.; Gerdes, M. J.; McBride,

[58195] L.; Lu, B.: Purification of a novel protein (ps20) from urogenital

[58196] sinus mesenchymal cells with growth inhibitory properties in vitro. J.

[58197] Biol. Chem. 270: 22058-22065, 1995.

[58198] 9562.

[58199] Hildebrand, J. D.; Taylor, J. M.; Parsons, T. J.: An SH3 domain-containing

[58200] GTPase-activating protein for Rho and Cdc42 associates with focal

[58201] adhesion kinase. Molec. Cell. Biol. 16: 3169-3178, 1996.

[58202] 9563.

[58203] Smith, D. E.; Renshaw, B. R.; Ketchem, R. R.; Kubin, M.; Garka,

[58204] K. E.; Sims, J. E.: Four new members expand the inter-

leukin-1 superfamily. J.

[58205] Biol. Chem. 275: 1169-1175, 2000.

[58206] 9564.

[58207] Kumar, S.; McDonnell, P. C.; Lehr, R.; Tierney, L.; Tzimas, M.

[58208] N.; Griswold, D. E.; Capper, E. A.; Tal-Singer, R.; Wells, G. I.;

[58209] Doyle, M. L.; Young, P. R.: Identification and initial characterization

[58210] of four novel members of the interleukin-1 family. J. Biol. Chem. 275:

[58211] 10308-10314, 2000.

[58212] 9565.

[58213] Masmoudi, S.; Antonarakis, S. E.; Schwede, T.; Ghorbel, A. M.;

[58214] Gratri, M.; Pappasavas, M.-P.; Drira, M.; Elgaied-Boulila, A.; Wattenhofer,

[58215] M.; Rossier, C.; Scott, H. S.; Ayadi, H.; Guipponi, M.: Novel missense

[58216] mutations of TMPRSS3 in two consanguineous Tunisian families with

[58217] nonsyndromic autosomal recessive deafness. Hum. Mutat.

18: 101-108,

[58218] 2001.

[58219] 9566.

[58220] Borkhardt, A.; Bojesen, S.; Haas, O. A.; Fuchs, U.; Bartel-
heimer,

[58221] D.; Loncarevic, I. F.; Bohle, R. M.; Harbott, J.; Repp, R.;
Jaeger,

[58222] U.; Viehmann, S.; Henn, T.; Korth, P.; Scharr, D.; Lampert,
F.: The

[58223] human GRAF gene is fused to MLL in a unique
t(5;11)(q31;q23) and both

[58224] alleles are disrupted in three cases of myelodysplastic
syndrome/acute

[58225] myeloid leukemia with a deletion 5q. Proc. Nat. Acad. Sci.
97: 9168-9173,

[58226] 2000.

[58227] 9567.

[58228] Krystal, G. W.; Armstrong, B. C.; Battey, J. F.: N-myc mRNA
forms

[58229] an RNA-RNA duplex with endogenous antisense tran-
scripts. Molec. Cell.

[58230] Biol. 10: 4180-4191, 1990.

[58231] 9568.

[58232] Torrents, D.; Estevez, R.; Pineda, M.; Fernandez, E.; Lloberas,

[58233] J.; Shi, Y.-B.; Zorzano, A.; Palacin, M.: Identification and characterization

[58234] of a membrane protein (y(+)L amino acid transporter-1) that associates

[58235] with 4F2hc to encode the amino acid transport activity y(+)L: a candidate

[58236] gene for lysinuric protein intolerance. J. Biol. Chem. 273: 32437-32445,

[58237] 1998.

[58238] 9569.

[58239] Yousef, G. M.; Chang, A.; Diamandis, E. P.: Identification and

[58240] characterization of KLK-L4, a new kallikrein-like gene that appears

[58241] to be down-regulated in breast cancer tissues. J. Biol. Chem. 275:

[58242] 11891-11898, 2000.

[58243] 9570.

[58244] Haft, C. R.; de la Luz Sierra, M.; Bafford, R.; Lesniak, M. A.;

[58245] Barr, V. A.; Taylor, S. I.: Human orthologs of yeast vacuolar protein

[58246] sorting proteins Vps26, 29, and 35: assembly into multi-meric complexes. Molec.

[58247] Biol. Cell 11: 4105-4116, 2000.

[58248] 9571.

[58249] Busfield, S. J.; Comrack, C. A.; Yu, G.; Chickering, T. W.; Smutko,

[58250] J. S.; Zhou, H.; Leiby, K. R.; Holmgren, L. M.; Gearing, D. P.; Pan,

[58251] Y.: Identification and gene organization of three novel members of

[58252] the IL-1 family on human chromosome 2. Genomics 66: 213-216, 2000.

[58253] 9572.

[58254] Mulero, J. J.; Pace, A. M.; Nelken, S. T.; Loeb, D. B.; Correa,

[58255] T. R.; Drmanac, R.; Ford, J. E.: IL1HY1: a novel interleukin-1 receptor

[58256] antagonist gene. Biochem. Biophys. Res. Commun. 263: 702-706, 1999.

[58257] 9573.

[58258] Ben-Yosef, T.; Wattenhofer, M.; Riazuddin, S.; Ahmed, Z. M.; Scott,

[58259] H. S.; Kudoh, J.; Shibuya, K.; Antonarakis, S. E.; Bonne-Tamir, B.;

[58260] Radhakrishna, U.; Naz, S.; Ahmed, Z.; Riazuddin, S.; Pandya, A.; Nance,

[58261] W. E.; Wilcox, E. R.; Friedman, T. B.; Morell, R. J.: Novel mutations

[58262] of TMPRSS3 in four DFNB8/B10 families segregating con-genital autosomal

[58263] recessive deafness. (Letter) J. Med. Genet. 38: 396-400, 2001.

[58264] 9574.

[58265] Pratt, W. S.; Crawley, S.; Hicks, J.; Ho, J.; Nash, M.; Kim, Y.

[58266] S.; Gum, J. R.; Swallow, D. M.: Multiple transcripts of MUC3: evidence

[58267] for two genes, MUC3A and MUC3B. Biochem. Biophys. Res. Commun. 275:

[58268] 916-923, 2000.

[58269] 9575.

[58270] Ishida, N.; Ito, M.; Yoshioka, S.; Sun-Wada, G.-H.; Kawakita, M.

[58271] : Functional expression of human Golgi CMP-sialic acid transporter

[58272] in the Golgi complex of a transporter-deficient Chinese hamster ovary

[58273] cell mutant. J. Biochem. 124: 171-178, 1998.

[58274] 9576.

[58275] Angata, T.; Varki, A.: Cloning, characterization, and phy-logenetic

[58276] analysis of Siglec-9, a new member of the CD33-related group of Siglecs:

[58277] evidence for co-evolution with sialic acid synthesis path-ways. J.

[58278] Biol. Chem. 275: 22127-22135, 2000.

[58279] 9577.

[58280] Foussias, G.; Yousef, G. M.; Diamandis, E. P.: Identification

[58281] and molecular characterization of a novel member of the Siglec family

[58282] (SIGLEC9). Genomics 67: 171-178, 2000.

[58283] 9578.

[58284] Zhang, J. Q.; Nicoll, G.; Jones, C.; Crocker, P. R.: Siglec-9,

[58285] a novel sialic acid binding member of the immunoglobulin superfamily

[58286] expressed broadly on human blood leukocytes. J. Biol. Chem. 275:

[58287] 22121-22126, 2000.

[58288] 9579.

[58289] Jinks-Robertson, S.: The genome's best friend. Nature Genet. 31:

[58290] 331-332, 2002.

[58291] 9580.

[58292] Wattenhofer, M.; Shibuya, K.; Kudoh, J.; Lyle, R.; Michaud, J.;

[58293] Rossier, C.; Kawasaki, K.; Asakawa, S.; Minoshima, S.; Berry, A.;

[58294] Bonne-Tamir, B.; Shimizu, N.; Antonarakis, S. E.; Scott, H. S.: Isolation

[58295] and characterization of the UBASH3A gene on 21q22.3 encoding a potential

[58296] nuclear protein with a novel combination of domains. Hum. Genet. 108:

[58297] 140-147, 2001.

[58298] 9581.

[58299] Kasof, G. M.; Gomes, B. C.: Livin, a novel inhibitor of apoptosis

[58300] protein family member. J. Biol. Chem. 276: 3238-3246, 2001.

[58301] 9582.

[58302] Lin, J.-H.; Deng, G.; Huang, Q.; Morser, J.: KIAP, a novel member

[58303] of the inhibitor of apoptosis protein family. Biochem. Bio-phys. Res.

[58304] Commun. 279: 820-831, 2000.

[58305] 9583.

[58306] Vucic, D.; Stennicke, H. R.; Pisabarro, M. T.; Salvesen, G. S.;

[58307] Dixit, V. M.: ML-IAP, a novel inhibitor of apoptosis that is preferentially

[58308] expressed in human melanomas. Curr. Biol. 10: 1359-1366, 2000.

[58309] 9584.

[58310] Balemans, W.; Ebeling, M.; Patel, N.; Van Hul, E.; Olson, P.;

Dioszegi,
[58311] M.; Lacza, C.; Wuyts, W.; Van Den Ende, J.; Willems, P.; Paes-Alves,
[58312] A. F.; Hill, S.; and 9 others: Increased bone density in sclerosteosis
[58313] is due to the deficiency of a novel secreted protein (SOST). Hum.
[58314] Molec. Genet. 10: 537-543, 2001.

[58315] 9585.

[58316] Kobayashi, S.; Akiyama, T.; Nata, K.; Abe, M.; Tajima, M.; Shervani,
[58317] N. J.; Unno, M.; Matsuno, S.; Sasaki, H.; Takasawa, S.; Okamoto, H.
[58318] : Identification of a receptor for Reg (regenerating gene) protein,
[58319] a pancreatic beta-cell regeneration factor. J. Biol. Chem. 275:
[58320] 10723-10726, 2000.

[58321] 9586.

[58322] Van Hul, W.; Wuyts, W.; Hendrickx, J.; Speleman, F.; Wauters, J.;
[58323] De Boulle, K.; Van Roy, N.; Bossuyt, P.; Willems, P. J.: Iden-

tification

[58324] of a third EXT-like gene (EXTL3) belonging to the EXT gene family. Genomics 47:

[58325] 230-237, 1998.

[58326] 9587.

[58327] Brdicka, T.; Pavlistova, D.; Leo, A.; Bruyns, E.; Korinek, V.;

[58328] Angelisova, P.; Scherer, J.; Shevchenko, A.; Shevchenko, A.; Hilgert,

[58329] I.; Cerny, J.; Drbal, K.; Kuramitsu, Y.; Kornacker, B.; Horejsi, V.;

[58330] Schraven, B.: Phosphoprotein associated with glycosphingolipid-enriched

[58331] microdomains (PAG), a novel ubiquitously expressed transmembrane adaptor

[58332] protein, binds the protein tyrosine kinase Csk and is involved in

[58333] regulation of T cell activation. J. Exp. Med. 191: 1591-1604, 2000.

[58334] 9588.

[58335] Dowler, S.; Currie, R. A.; Downes, C. P.; Alessi, D. R.: DAPP1:

[58336] a dual adaptor for phosphotyrosine and

3-phosphoinositides. Biochem.
[58337] J. 342: 7-12, 1999.

[58338] 9589.

[58339] Marshall, A. J.; Niiro, H.; Lerner, C. G.; Yun, T. J.; Thomas,

[58340] S.; Disteche, C. M.; Clark, E. A.: A novel B lymphocyte-associated
[58341] adaptor protein, Bam32, regulates antigen receptor signaling downstream
[58342] of phosphatidylinositol 3-kinase. J. Exp. Med. 191: 1319-1331, 2000.
[58343] 9590.

[58344] Cantor, S. B.; Bell, D. W.; Ganesan, S.; Kass, E. M.; Drapkin,

[58345] R.; Grossman, S.; Wahrer, D. C. R.; Sgroi, D. C.; Lane, W. S.; Haber,
[58346] D. A.; Livingston, D. M.: BACH1, a novel helicase-like protein, interacts
[58347] directly with BRCA1 and contributes to its DNA repair function. Cell 105:
[58348] 149-160, 2001.

[58349] 9591.

[58350] Cheung, I.; Schertzer, M.; Rose, A.; Lansdorp, P. M.: Dis-

ruption

[58351] of dog-1 in Caenorhabditis elegans triggers deletions upstream of

[58352] guanine-rich DNA. Nature Genet. 31: 405-409, 2002.

[58353] 9592.

[58354] Kinzler, K. W.; Vogelstein, B. :Nature 386: 761-763, 1997.

[58355] 9593.

[58356] Schmitt, C.; von Kobbe, C.; Bachi, A.; Pante, N.; Rodrigues, J.

[58357] P.; Boscheron, C.; Rigaut, G.; Wilm, M.; Seraphin, B.; Carmo-Fonseca,

[58358] M.; Izaurralde, E.: Dbp5, a DEAD-box protein required for mRNA export,

[58359] is recruited to the cytoplasmic fibrils of nuclear pore complex via

[58360] a conserved interaction with CAN/Nup159p. EMBO J. 18: 4332-4347,

[58361] 1999.

[58362] 9594.

[58363] Pati, D.; Meistrich, M. L.; Plon, S. E.: Human Cdc34 and Rad6B

[58364] ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced
[58365] transcription for proteolysis. Molec. Cell. Biol. 19: 5001-5013,
[58366] 1999.

[58367] 9595.

[58368] Courjal, F.; Louason, G.; Speiser, P.; Katsaros, D.; Zeillinger,
[58369] R.; Theillet, C.: Cyclin gene amplification and overexpres-sion in
[58370] breast and ovarian cancers: evidence for the selection of cyclin D1
[58371] in breast and cyclin E in ovarian tumors. Int. J. Cancer 69: 247-253,
[58372] 1996.

[58373] 9596.

[58374] Strohmaier, H.; Spruck, C. H.; Kaiser, P.; Won, K.-A.; Sangfelt,
[58375] O.; Reed, S. I.: Human F-box protein hCdc4 targets cyclin E for proteolysis
[58376] and is mutated in a breast cancer cell line. Nature 413: 316-322,

[58377] 2001.

[58378] 9597.

[58379] Matsuda, S.; Iriyama, C.; Yokozaki, S.; Ichigotani, Y.; Shi-
rafuji,

[58380] N.; Yamaki, K.; Hayakawa, T.; Hamaguchi, M.: Cloning and
sequencing

[58381] of a novel human gene that encodes a putative target pro-
tein of Nesh-SH3. J.

[58382] Hum. Genet. 46: 483-486, 2001.

[58383] 9598.

[58384] Jager, D.; Stockert, E.; Jager, E.; Gure, A. O.; Scanlan, M. J.;

[58385] Knuth, A.; Old, L. J.; Chen, Y.-T.: Serological cloning of a
melanocyte

[58386] rab guanosine 5-prime-triphosphate-binding protein and
a chromosome

[58387] condensation protein from a melanoma complementary
DNA library. Cancer

[58388] Res. 60: 3584-3591, 2000.

[58389] 9599.

[58390] Loftus, S. K.; Larson, D. M.; Baxter, L. L.; Antonellis, A.;
Chen,

[58391] Y.; Wu, X.; Jiang, Y.; Bittner, M.; Hammer, J. A., III; Pavan, W.

[58392] J.: Mutation of melanosome protein RAB38 in chocolate mice. Proc.

[58393] Nat. Acad. Sci. 99: 4471-4476, 2002.

[58394] 9600.

[58395] Matsuyoshi, N.; Tanaka, T.; Toda, K.; Imamura, S.: Identification

[58396] of novel cadherins expressed in human melanoma cells. J. Invest.

[58397] Derm. 108: 908-913, 1997.

[58398] 9601.

[58399] Gumienny, T. L.; Brugnera, E.; Tosello-Trampont, A.-C.; Kinchen,

[58400] J. M.; Haney, L. B.; Nishiwaki, K.; Walk, S. F.; Nemergut, M. E.;

[58401] Macara, I. G.; Francis, R.; Schedl, T.; Qin, Y.; Van Aelst, L.; Hengartner,

[58402] M. O.; Ravichandran, K. S.: CED-12/ELMO, a novel member of the CrkII/Dock180/Rac

[58403] pathway, is required for phagocytosis and cell migration. Cell 107:

[58404] 27-41, 2001.

[58405] 9602.

[58406] Tanner, S. M.; Austin, J. L.; Leone, G.; Rush, L. J.; Plass, C.;

[58407] Heinonen, K.; Mrozek, K.; Sill, H.; Knuutila, S.; Kolitz, J. E.; Archer,

[58408] K. J.; Caligiuri, M. A.; Bloomfield, C. D.; de la Chapelle, A.: BAALC,

[58409] the human member of a novel mammalian neuroectoderm gene lineage,

[58410] is implicated in hematopoiesis and acute leukemia. Proc. Nat. Acad.

[58411] Sci. 98: 13901-13906, 2001.

[58412] 9603.

[58413] Wang, H.; Fei, Y.-J.; Kekuda, R.; Yang-Feng, T. L.; Devoe, L. D.;

[58414] Leibach, F. H.; Prasad, P. D.; Ganapathy, V.: Structure, function,

[58415] and genomic organization of human Na(+)-dependent high-affinity dicarboxylate

[58416] transporter. Am. J. Physiol. Cell Physiol. 278: C1019-C1030, 2000.

[58417] 9604.

[58418] Yun, C. H. C.; Oh, S.; Zizak, M.; Steplock, D.; Tsao, S.; Tse,

[58419] C.-M.; Weinman, E. J.; Donowitz, M.: cAMP-mediated inhi-
bition of

[58420] the epithelial brush border Na(+)/H(+) exchanger, NHE3,
requires an

[58421] associated regulatory protein. Proc. Nat. Acad. Sci. 94:
3010-3015,

[58422] 1997.

[58423] 9605.

[58424] Li, Y.; Chin, L.-S.; Weigel, C.; Li, L.: Spring, a novel RING

[58425] finger protein that regulates synaptic vesicle exocytosis. J.
Biol.

[58426] Chem. 276: 40824-40833, 2001.

[58427] 9606.

[58428] Reymond, A.; Meroni, G.; Fantozzi, A.; Merla, G.; Cairo, S.;
Luzi,

[58429] L.; Riganelli, D.; Zanaria, E.; Messali, S.; Cainarca, S.; Guf-
fanti,

[58430] A.; Minucci, S.; Pelicci, P. G.; Ballabio, A.: The tripartite
motif

[58431] family identifies cell compartments. EMBO J. 20:
2140-2151, 2001.
[58432] 9607.

[58433] Boles, K. S.; Mathew, P. A.: Molecular cloning of CS1, a
novel
[58434] human natural killer cell receptor belonging to the CD2
subset of
[58435] the immunoglobulin superfamily. Immunogenetics 52:
302-307, 2001.
[58436] 9608.

[58437] Bouchon, A.; Cella, M.; Grierson, H. L.; Cohen, J. I.;
Colonna,
[58438] M.: Cutting edge: activation of NK cell-mediated cytotoxi-
city by
[58439] a SAP-independent receptor of the CD2 family. J. Immun.
167: 5517-5521,
[58440] 2001.

[58441] 9609.

[58442] Habas, R.; Kato, Y.; He, X.: Wnt/Frizzled activation of Rho
regulates
[58443] vertebrate gastrulation and requires a novel Formin ho-
mology protein

[58444] Daam1. Cell 107: 843-854, 2001.

[58445] 9610.

[58446] Peters, C. S.; Liang, X.; Li, S.; Kannan, S.; Peng, Y.; Taub, R.;

[58447] Diamond, R. H.: ATF-7, a novel bZIP protein, interacts with the PRL-1

[58448] protein-tyrosine phosphatase. J. Biol. Chem. 276: 13718-13726, 2001.

[58449] 9611.

[58450] White, J. H.; McIllhinney, R. A. J.; Wise, A.; Ciruela, F.; Chan,

[58451] W.-Y.; Emson, P. C.; Billinton, A.; Marshall, F. H.: The GABA-B receptor

[58452] interacts directly with the related transcription factors CREB2 and

[58453] ATFx. Proc. Nat. Acad. Sci. 97: 13967-13972, 2000.

[58454] 9612.

[58455] Futai, E.; Kubo, T.; Sorimachi, H.; Suzuki, K.; Maeda, T.: Molecular

[58456] cloning of PalBH, a mammalian homologue of the As-pergillus atypical

[58457] calpain PalB. Biochim. Biophys. Acta 1517: 316-319, 2001.

[58458] 9613.

[58459] Huang, Y.; Wang, K. K. W.: The calpain family and human disease. Trends

[58460] Molec. Med. 7: 355-362, 2001.

[58461] 9614.

[58462] Lee, H.-J.; Sorimachi, H.; Jeong, S.-Y.; Ishiura, S.; Suzuki, K.

[58463] : Molecular cloning and characterization of a novel tissue-specific

[58464] calpain predominantly expressed in digestive tract. Biol. Chem. 379:

[58465] 175-183, 1998.

[58466] 9615.

[58467] Yoshikawa, Y.; Mukai, H.; Hino, F.; Asada, K.; Kato, I.: Isolation

[58468] of two novel genes, down-regulated in gastric cancer. Jpn. J. Cancer

[58469] Res. 91: 459-463, 2000.

[58470] 9616.

[58471] Burgess, D. L.; Gefrides, L. A.; Foreman, P. J.; Noebels, J. L.

[58472] : A cluster of three novel Ca(2+) channel gamma subunit genes on chromosome

[58473] 19q13.4: evolution and expression profile of the gamma subunit gene

[58474] family. Genomics 71: 339-350, 2001.

[58475] 9617.

[58476] Chu, P.-J.; Robertson, H. M.; Best, P. M.: Calcium channel gamma

[58477] subunits provide insights into the evolution of this gene family. Gene 280:

[58478] 37-48, 2001.

[58479] 9618.

[58480] Chen, X.; Wen, S.; Fukuda, M. N.; Gavva, N. R.; Hsu, D.; Akama,

[58481] T. O.; Yang-Feng, T.; Shen, C. K. J.: Human ITCH is a coregulator

[58482] of the hematopoietic transcription factor NF-E2. Genomics 73: 238-241,

[58483] 2001.

[58484] 9619.

[58485] D'Andrea, A. D.; Serhan, C. N.: Relieving the Itch. Nature Genet. 18:
[58486] 97-99, 1998.

[58487] 9620.

[58488] Perry, W. L.; Hustad, C. M.; Swing, D. A.; O'Sullivan, T. N.; Jenkins,
[58489] N. A.; Copeland, N. G.: The itchy locus encodes a novel ubiquitin
[58490] protein ligase that is disrupted in a18H mice. Nature Genet. 18:
[58491] 143-146, 1998.

[58492] 9621.

[58493] Qiu, L.; Joazeiro, C.; Fang, N.; Wang, H.-Y.; Elly, C.; Alt-man,
[58494] Y.; Fang, D.; Hunter, T.; Liu, Y.-C.: Recognition and ubiq-uitination
[58495] of Notch by Itch, a Hect-type E3 ubiquitin ligase. J. Biol. Chem. 275:
[58496] 35734-35737, 2000.

[58497] 9622.

[58498] Winberg, G.; Matskova, L.; Chen, F.; Plant, P.; Rotin, D.;

Gish,
[58499] G.; Ingham, R.; Ernberg, I.; Pawson, T.: Latent membrane protein
[58500] 2A of Epstein-Barr virus binds WW domain E3 protein-ubiquitin ligases
[58501] that ubiquitinate B-cell tyrosine kinases. Molec. Cell. Biol. 20:
[58502] 8526-8535, 2000.

[58503] 9623.

[58504] Bradley, K. A.; Mogridge, J.; Mourez, M.; Collier, R. J.; Young,
[58505] J. A. T.: Identification of the cellular receptor for anthrax toxin. Nature 414:
[58506] 160-161, 2001.

[58507] 9624.

[58508] Krause, A.; Neitz, S.; Magert, H.-J.; Schulz, A.; Forssmann, W.-G.;
[58509] Schulz-Knappe, P.; Adermann, K.: LEAP-1, a novel highly disulfide-bonded
[58510] human peptide, exhibits antimicrobial activity. FEBS Lett. 480:
[58511] 147-150, 2000.

[58512] 9625.

[58513] Nicolas, G.; Bennoun, M.; Devaux, I.; Beaumont, C.; Grandchamp,

[58514] B.; Kahn, A.; Vaulont, S.: Lack of hepcidin gene expression and severe

[58515] tissue iron overlad in upstream stimulatory factor 2 (USF2) knockout

[58516] mice. Proc. Natl. Acad. Sci. 98: 8780-8785, 2001.

[58517] 9626.

[58518] Nicolas, G.; Bennoun, M.; Porteu, A.; Mativet, S.; Beaumont, C.;

[58519] Grandchamp, B.; Sirito, M.; Sawadogo, M.; Kahn, A.; Vaulont, S.:

[58520] Severe iron deficiency anemia in transgenic mice expressing liver

[58521] hepcidin. Proc. Nat. Acad. Sci. 99: 4596-4601, 2002.

[58522] 9627.

[58523] Park, C. H.; Valore, E. V.; Waring, A. J.; Ganz, T.: Hepcidin,

[58524] a urinary antimicrobial peptide synthesized in the liver. J. Biol.

[58525] Chem. 276: 7806-7810, 2001.

[58526] 9628.

[58527] Pigeon, C.; Ilyin, G.; Courselaud, B.; Leroyer, P.; Turlin, B.;

[58528] Brissot, P.; Loreal, O.: A new mouse liver-specific gene, encoding

[58529] a protein homologous to human antimicrobial peptide hepcidin, is overexpressed

[58530] during iron overload. J. Biol. Chem. 276: 7811-7819, 2001.

[58531] 9629.

[58532] Lin, M.; Napoli, J. L.: cDNA cloning and expression of a human

[58533] aldehyde dehydrogenase (ALDH) active with 9-cis-retinal and identification

[58534] of a rat ortholog, ALDH12. J. Biol. Chem. 275: 40106-40112, 2000.

[58535] 9630.

[58536] Dragon, F.; Pogacic, V.; Filipowicz, W.: In vitro assembly of

[58537] human H/ACA small nucleolar RNPs reveals unique features of U17 and

[58538] telomerase RNAs. Molec. Cell. Biol. 20: 3037-3048, 2000.

[58539] 9631.

[58540] Tollervey, D.; Kiss, T.: Function and synthesis of small nucleolar

[58541] RNAs. Curr. Opin. Cell Biol. 9: 337-342, 1997.

[58542] 9632.

[58543] Pogacic, V.; Dragon, F.; Filipowicz, W.: Human H/ACA small nucleolar

[58544] RNPs and telomerase share evolutionarily conserved proteins NHP2 and

[58545] NOP10. Molec. Cell. Biol. 20: 9028-9040, 2000.

[58546] 9633.

[58547] Ichtchenko, K.; Nguyen, T.; Sudhof, T. C.: Structures, alternative

[58548] splicing, and neurexin binding of multiple neuroligins. J. Biol.

[58549] Chem. 271: 2676-2682, 1996.

[58550] 9634.

[58551] Hammarsund, M.; Wilson, W.; Corcoran, M.; Merup, M.; Einhorn, S.;

[58552] Grander, D.; Sangfelt, O.: Identification and characterization of

[58553] two novel human mitochondrial elongation factor genes,

hEFG2 and hEFG1,

[58554] phylogenetically conserved through evolution. Hum. Genet. 109: 542-550,

[58555] 2001.

[58556] 9635.

[58557] Ishibashi, K.; Suzuki, M.; Sasaki, S.; Imai, M.: Identification

[58558] of a new multigene four-transmembrane family (MS4A) related to CD20,

[58559] HTm4 and beta subunit of the high-affinity IgE receptor. Gene 264:

[58560] 87-93, 2001.

[58561] 9636.

[58562] Liang, Y.; Tedder, T. F.: Identification of a CD20-, Fc-epsilon-RI-beta-related

[58563] gene family: sixteen new MS4A family members expressed in human and

[58564] mouse. Genomics 72: 119-127, 2001.

[58565] 9637.

[58566] Ishii, H.; Vecchione, A.; Murakumo, Y.; Baldassarre, G.; Numata,

[58567] S.; Trapasso, F.; Alder, H.; Baffa, R.; Croce, C. M.:

FEZ1/LZTS1
[58568] gene at 8p22 suppresses cancer cell growth and regulates mitosis. Proc.
[58569] Nat. Acad. Sci. 98: 10374-10379, 2001.

[58570] 9638.

[58571] Mahon, M. J.; Donowitz, M.; Yun, C. C.; Segre, G. V.: Na+/H+ exchanger
[58572] regulatory factor 2 directs parathyroid hormone 1 receptor signalling. Nature 417:
[58573] 858-861, 2002.

[58574] 9639.

[58575] Poulat, F.; de Santa Barbara, P.; Desclozeaux, M.; Soullier, S.;
[58576] Moniot, B.; Bonneaud, N.; Boizet, B.; Berta, P. The human testis
[58577] determining factor SRY binds a nuclear factor containing PDZ protein
[58578] interaction domains. J. Biol. Chem. 272: 7167-7172, 1997.
[58579] 9640.

[58580] Yun, C.-H. C.; Lamprecht, G.; Forster, D. V.; Sidor, A.:

NHE3

[58581] kinase A regulatory protein E3KARP binds the epithelial brush border

[58582] Na(+)/H(+) exchanger NHE3 and the cytoskeletal protein ezrin. J.

[58583] Biol. Chem. 273: 25856-25863, 1998.

[58584] 9641.

[58585] Hall, R. A.; Ostedgaard, L. S.; Premont, R. T.; Blitzer, J. T.;

[58586] Rahman, N.; Welsh, M. J.; Lefkowitz, R. J.: A C-terminal motif found

[58587] in the beta-2-adrenergic receptor, P2Y1 receptor and cystic fibrosis

[58588] transmembrane conductance regulator determines binding to the Na(+)/H(+)

[58589] exchanger regulatory factor family of PDZ proteins. Proc. Nat. Acad.

[58590] Sci. 95: 8496-8501, 1998.

[58591] 9642.

[58592] Druck, T.; Podolski, J.; Byrski, T.; Wyrwicz, L.; Zajaczek, S.;

[58593] Kata, G.; Borowka, A.; Lubinski, J.; Huebner, K.: The DIRC1 gene

[58594] at chromosome 2q33 spans a familial RCC-associated t(2;3)(q33;q21)
[58595] chromosome translocation. J. Hum. Genet. 46: 583-589, 2001.
[58596] 9643.
[58597] Podolski, J.; Zajaczek, S.; Byrski, T.; Druck, T.; Zimonjic, D.
[58598] B.; Popescu, N. C.; Lubinski, J.; Huebner, K.: Characterization of
[58599] a familial RCC-associated t(2;3)(q33;q21) chromosome translocation. J.
[58600] Hum. Genet. in-press, 2001.
[58601] 9644.
[58602] Bruick, R. K.; McKnight, S. L.: A conserved family of prolyl-4-hydroxylases
[58603] that modify HIF. Science 294: 1337-1340, 2001.
[58604] 9645.
[58605] Epstein, A. C. R.; Gleadle, J. M.; McNeill, L. A.; Hewitson, K.
[58606] S.; O'Rourke, J.; Mole, D. R.; Mukherji, M.; Metzen, E.; Wilson, M.
[58607] I.; Dhanda, A.; Tian, Y.-M.; Masson, N.; Hamilton, D. L.; Jaakkola,

[58608] P.; Barstead, R.; Hodgkin, J.; Maxwell, P. H.; Pugh, C. W.; Schofield,

[58609] C. J.; Ratcliffe, P. J.: C. elegans EGL-9 and mammalian homologs

[58610] define a family of dioxygenases that regulate HIF by prolyl hydroxylation. Cell 107:

[58611] 43-54, 2001.

[58612] 9646.

[58613] Dupuy, D.; Aubert, I.; Duperat, V. G.; Petit, J.; Taine, L.; Stef,

[58614] M.; Bloch, B.; Arveiler, B.: Mapping, characterization, and expression

[58615] analysis of the SM-20 human homologue, C1orf12, and identification

[58616] of a novel related gene, SCAND2. Genomics 69: 348-354, 2000.

[58617] 9647.

[58618] Ritter, J. K.; Crawford, J. M.; Owens, I. S.: Cloning of two human

[58619] liver bilirubin UDP-glucuronosyltransferase cDNAs with expression

[58620] in COS-1 cells. J. Biol. Chem. 266: 1043-1047, 1991.

[58621] 9648.

[58622] Rheaume, E.; Leblanc, J. F.; Lachance, Y.; Labrie, F.; Simard,
[58623] J.: Detection of frequent BglII polymorphism by polymerase chain
[58624] reaction and TaqI restriction fragment length polymorphism for 3-beta-hydroxysteroid
[58625] dehydrogenase/delta-5-delta-4 isomerase at the human HSDB3 locus (1p11-p13).
[58626] Hum. Genet. 87: 753-754, 1991.

[58627] 9649.

[58628] Yamaguchi, N.; Benya, P. D.; van der Rest, M.; Ninomiya, Y.: The
[58629] cloning and sequencing of alpha-1(VIII) collagen cDNAs demonstrate
[58630] that type VIII collagen is a short chain collagen and contains triple-helical
[58631] and carboxyl-terminal non-triple-helical domains similar to those
[58632] of type X collagen. J. Biol. Chem. 264: 16022-16029, 1989.

[58633] 9650.

[58634] Bleul, C. C.; Farzan, M.; Choe, H.; Parolin, C.; Clark-Lewis, I.;

[58635] Sodroski, J.; Springer, T. A.: The lymphocyte chemoattractant SDF-1

[58636] is a ligand for LESTR/fusin and blocks HIV-1 entry. Nature 382:

[58637] 829-833, 1996.

[58638] 9651.

[58639] Caruz, A.; Samsom, M.; Alonso, J. M.; Alcami, J.; Baleux, F.; Virelizier,

[58640] J. L.; Parmentier, M.; Arenzana-Seisdedos, F.: Genomic organization

[58641] and promoter characterization of human CXCR4 gene. FEBS Lett. 426:

[58642] 271-278, 1998.

[58643] 9652.

[58644] Clapham, P. R.; Blanc, D.; Weiss, R. A.: Specific cell surface

[58645] requirements for the infection of CD4-positive cells by human immunodeficiency

[58646] virus types 1 and 2 and by Simian immunodeficiency virus. Virology 181:

[58647] 703-715, 1991.

[58648] 9653.

[58649] Dragic, T.; Alizon, M.: Different requirements for mem-
brane fusion
[58650] mediated by the envelopes of human immunodeficiency
virus types 1
[58651] and 2. J. Virol. 67: 2355-2359, 1993.

[58652] 9654.

[58653] Federsppiel, B.; Melhado, I. G.; Duncan, A. M. V.; Delaney,
A.;
[58654] Schappert, K.; Clark-Lewis, I.; Jirik, F. R.: Molecular
cloning of
[58655] the cDNA and chromosomal localization of the gene for a
putative seven-transmembrane
[58656] segment (7-TMS) receptor isolated from human spleen.
Genomics 16:
[58657] 707-712, 1993.

[58658] 9655.

[58659] Feng, Y.; Broder, C. C.; Kennedy, P. E.; Berger, E. A.: HIV-1
[58660] entry cofactor: functional cDNA cloning of a seven-
transmembrane,

[58661] G protein-coupled receptor. Science 272: 872-876, 1996.

[58662] 9656.

[58663] Hendrix, C. W.; Flexner, C.; MacFarland, R. T.; Gian-
domenico, C.;

[58664] Fuchs, E. J.; Redpath, E.; Bridger, G.; Henson, G. W.: Phar-
macokinetics

[58665] and safety of AMD-3100, a novel antagonist of the CXCR-
4 chemokine

[58666] receptor, in human volunteers. Antimicrob. Agents
Chemother. 44:

[58667] 1667-1673, 2000.

[58668] 9657.

[58669] Herzog, H.; Hort, Y. J.; Shine, J.; Selbie, L. A.: Molecular
cloning,

[58670] characterization, and localization of the human homolog
to the reported

[58671] bovine NPY Y3 receptor: lack of NPY binding and activa-
tion. DNA Cell

[58672] Biol. 12: 465-471, 1993.

[58673] 9658.

[58674] Jazin, E. E.; Yoo, H.; Blomqvist, A. G.; Yee, F.; Weng, G.;

Walker,
[58675] M. W.; Salon, J.; Larhammar, D.; Wahlestedt, C.: A pro-
posed bovine
[58676] neuropeptide Y (NPY) receptor cDNA clone, or its human
homologue,
[58677] confers neither NPY binding sites nor NPY responsiveness
on transfected
[58678] cells. Regul. Pept. 47: 247-258, 1993.

[58679] 9659.

[58680] Liotta, L. A.: An attractive force in metastasis. Nature 410:

[58681] 24-25, 2001.

[58682] 9660.

[58683] Loetscher, M.; Geiser, T.; O'Reilly, T.; Zwahlen, R.; Baggi-
olini,
[58684] M.; Moser, B.: Cloning of a human seven-transmembrane
domain receptor,
[58685] LESTR, that is highly expressed in leukocytes. J. Biol.
Chem. 269:
[58686] 232-237, 1994.

[58687] 9661.

[58688] Lu, M.; Grove, E. A.; Miller, R. J.: Abnormal development of

[58689] the hippocampal dentate gyrus in mice lacking the CXCR4 chemokine
[58690] receptor. Proc. Nat. Acad. Sci. 99: 7090-7095, 2002.

[58691] 9662.

[58692] Ma, Q.; Jones, D.; Borghesani, P. R.; Segal, R. A.; Naga-sawa,
[58693] T.; Kishimoto, T.; Bronson, R. T.; Springer, T. A.: Impaired B-lymphopoiesis,
[58694] myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and
[58695] SDF-1-deficient mice. Proc. Nat. Acad. Sci. 95: 9448-9453, 1998.

[58696] 9663.

[58697] Muller, A.; Homey, B.; Soto, H.; Ge, N.; Catron, D.; Buchanan,
[58698] M. E.; McClanahan, T.; Murphy, E.; Yuan, W.; Wagner, S. N.; Barrera,
[58699] J. L.; Mohar, A.; Verastegui, E.; Zlotnik, A.: Involvement of chemokine
[58700] receptors in breast cancer metastasis. Nature 410: 50-56, 2001.

[58701] 9664.

[58702] Nagasawa, T.; Hirota, S.; Tachibana, K.; Takakura, N.; Nishikawa,
[58703] S.; Kitamura, Y.; Yoshida, N.; Kikutani, H.; Kishimoto, T.: Defects
[58704] of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking
[58705] the CXC chemokine PBSF/SDF-1. Nature 382: 635-638, 1996.
[58706] 9665.

[58707] Oberlin, E.; Amara, A.; Bachelerie, F.; Bessia, C.; Virelizier,
[58708] J.-L.; Arenzana-Seisdedos, F.; Schwartz, O.; Heard, J.-M.; Clark-Lewis,
[58709] I.; Legler, D. F.; Loetscher, M.; Baggiollini, M.; Moser, B.: The
[58710] CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection
[58711] by T-cell-line-adapted HIV-1. Nature 382: 833-835, 1996.
[58712] 9666.

[58713] Peled, A.; Petit, I.; Kollet, O.; Magid, M.; Ponomaryov, T.; Byk,
[58714] T.; Nagler, A.; Ben-Hur, H.; Many, A.; Shultz, L.; Lider, O.;

Alon,

[58715] R.; Zipori, D.; Lapidot, T.: Dependence of human stem cell engraftment

[58716] and repopulation of NOD/SCID mice on CXCR4. Science 283: 845-848,

[58717] 1999.

[58718] 9667.

[58719] Rimland, J.; Xin, W.; Sweetnam, P.; Saijoh, K.; Nestler, E. J.;

[58720] Duman, R. S.: Sequence and expression of a neuropeptide Y receptor

[58721] cDNA. Molec. Pharm. 40: 869-875, 1991.

[58722] 9668.

[58723] Tachibana, K.; Hirota, S.; Iizasa, H.; Yoshida, H.; Kawabata,

[58724] K.; Kataoka, Y.; Kitamura, Y.; Matsushima, K.; Yoshida, N.; Nishikawa,

[58725] S.; Kishimoto, T.; Nagasawa, T.: The chemokine receptor CXCR4 is

[58726] essential for vascularization of the gastrointestinal tract. Nature 393:

[58727] 591-594, 1998.

[58728] 9669.

[58729] Wegner, S. A.; Ehrenberg, P. K.; Chang, G.; Dayhoff, D. E.; Sleeker,
[58730] A. L.; Michael, N. L.: Genomic organization and functional characterization
[58731] of the chemokine receptor CXCR4, a major entry co-receptor for human
[58732] immunodeficiency virus type 1. J. Biol. Chem. 273: 4754-4760, 1998.
[58733] 9670.

[58734] Weiner, D. B.; Huebner, K.; Williams, W. V.; Greene, M. I.: Human
[58735] genes other than CD4 facilitate HIV-1 infection of murine cells. Pathobiology 59:
[58736] 361-371, 1991.

[58737] 9671.

[58738] Szostecki, C.; Guldner, H. H.; Netter, H. J.; Will, H.: Isola-tion
[58739] and characterization of cDNA encoding a human nuclear antigen predominantly
[58740] recognized by autoantibodies from patients with primary biliary cirrhosis. J.
[58741] Immun. 145: 4338-4347, 1990.

[58742] 9672.

[58743] Freeman, G. J.; Disteche, C. M.; Gribben, J. G.; Adler, D. A.;

[58744] Freedman, A. S.; Dougery, J.; Nadler, L. M.: The gene for B7, a costimulatory

[58745] signal for T-cell activation, maps to chromosomal region 3q13.3-3q21. Blood 79:

[58746] 489-494, 1992.

[58747] 9673.

[58748] Reeves, R. H.; Patch, D.; Sharpe, A. H.; Borriello, F.; Free-man,

[58749] G. J.; Edelhoff, S.; Disteche, C.: The costimulatory genes Cd80 and

[58750] Cd86 are linked on mouse chromosome 16 and human chromosome 3. Mammalian

[58751] Genome 8: 581-582, 1997.

[58752] 9674.

[58753] Selvakumar, A.; Mohanraj, B. K.; Eddy, R. L.; Shows, T. B.; White,

[58754] P. C.; Dupont, B.: Genomic organization and chromosomal location

[58755] of the human gene encoding the B-lymphocyte activation

antigen B7. Immunogenetics 36:
[58756] 175-181, 1992.

[58757] 9675.

[58758] Ha, H.; Kubagawa, H.; Burrows, P. D.: Molecular cloning and expression
[58759] pattern of a human gene homologous to the murine mb-1 gene. J. Immun. 148:
[58760] 1526-1531, 1992.

[58761] 9676.

[58762] Hashimoto, S.; Mohrenweiser, H. W.; Gregersen, P. K.; Chiorazzi,
[58763] N.: Chromosomal localization, genomic structure, and allelic polymorphism
[58764] of the human CD79a (Ig-alpha/mb-1) gene. Immunogenetics 40: 287-295,
[58765] 1994.

[58766] 9677.

[58767] Reth, M.: Antigen receptors on B lymphocytes. Annu. Rev. Immun. 10:
[58768] 97-121, 1992.

[58769] 9678.

[58770] Liang, Y.; Buckley, T. R.; Tu, L.; Langdon, S. D.; Tedder, T. F.

[58771] : Structural organization of the human MS4A gene cluster on chromosome

[58772] 11q12. Immunogenetics 53: 357-368, 2001.

[58773] 9679.

[58774] Tedder, T. F.; Disteche, C. M.; Louie, E.; Adler, D. A.; Croce,

[58775] C. M.; Schlossman, S. F.; Saito, H.: The gene that encodes the human

[58776] CD20 (B1) differentiation antigen is located on chromosome 11 near

[58777] the t(11;14)(q13;q32) translocation site. J. Immun. 142: 2555-2559,

[58778] 1989.

[58779] 9680.

[58780] Tedder, T. F.; Klejman, G.; Schlossman, S. F.; Saito, H.: Structure

[58781] of the gene encoding the human B lymphocyte differentiation antigen

[58782] CD20 (B1). J. Immun. 142: 2560-2568, 1989.

[58783] 9681.

[58784] Tedder, T. F.; Streuli, M.; Schlossman, S. F.; Saito, H.: Isolation

[58785] and structure of a cDNA encoding the B1 (CD20) cell-surface antigen

[58786] of human B lymphocytes. Proc. Nat. Acad. Sci. 85: 208-212, 1988.

[58787] 9682.

[58788] Celeste, A. J.; Rosen, V.; Buecker, J. L.; Kriz, R.; Wang, E. A.;

[58789] Wozney, J. M.: Isolation of the human gene for bone gla protein utilizing

[58790] mouse and rat cDNA clones. EMBO J. 5: 1885-1890, 1986.

[58791] 9683.

[58792] Desbois, C.; Seldin, M. F.; Karsenty, G.: Localization of the

[58793] osteocalcin gene cluster on mouse chromosome 3. Mammalian Genome 5:

[58794] 321-322, 1994.

[58795] 9684.

[58796] Ducy, P.; Karsenty, G.: Two distinct osteoblast-specific cis-acting

[58797] elements control expression of a mouse osteocalcin gene. Molec. Cell.

[58798] Biol. 15: 1858-1869, 1995.

[58799] 9685.

[58800] Ducy, P.; Zhang, R.; Geoffroy, V.; Ridall, A. L.; Karsenty, G.

[58801] : Osf2/Cbfa1: a transcriptional activator of osteoblast differentiation. Cell 89:

[58802] 747-754, 1997.

[58803] 9686.

[58804] Kaplan, F. S.; Tabas, J. A.; Zasloff, M. A.: Fibrodysplasia ossificans

[58805] progressiva: a clue from the fly? Calcif. Tissue Int. 47: 117-125,

[58806] 1990.

[58807] 9687.

[58808] Kerner, S. A.; Scott, R. A.; Pike, J. W.: Sequence elements in

[58809] the human osteocalcin gene confer basal activation and inducible response

[58810] to hormonal vitamin D(3). Proc. Nat. Acad. Sci. 86: 4455-4459, 1989.

[58811] 9688.

[58812] Hopken, U. E.; Lu, D.; Gerard, N. P.; Gerard, C.: The C5a chemoattractant

[58813] receptor mediates mucosal defense to infection. Nature 383: 86-89,

[58814] 1996.

[58815] 9689.

[58816] Chen, H.; Jawahar, S.; Qian, Y.; Duong, Q.; Chan, G.; Parker, A.;

[58817] Meyer, J. M.; Moore, K. J.; Chayen, S.; Gross, D. J.; Glaser, B.;

[58818] Permutt, M. A.; Fricker, L. D.: Missense polymorphism in the human

[58819] carboxypeptidase E gene alters enzymatic activity. Hum. Mutat. 18:

[58820] 120-131, 2001.

[58821] 9690.

[58822] Hall, C.; Manser, E.; Spurr, N. K.; Lim, L.: Assignment of the

[58823] human carboxypeptidase E (CPE) gene to chromosome 4. Genomics 15:

[58824] 461-463, 1993.

[58825] 9691.

[58826] Manser, E.; Fernandez, D.; Loo, L.; Goh, P. Y.; Monfries, C.; Hall,

[58827] C.; Lim, L.: Human carboxypeptidase E: isolation and characterization

[58828] of the cDNA, sequence conservation, expression and processing in vitro. Biochem.

[58829] J. 267: 517-525, 1990.

[58830] 9692.

[58831] Kas, K.; Schoenmakers, E. F. P. M.; Van de Ven, W. J. M.: Physical

[58832] map location of the human carboxypeptidase M gene (CPM) distal to

[58833] D12S375 and proximal to D12S8 at chromosome 12q15. Genomics 30:

[58834] 403-405, 1995.

[58835] 9693.

[58836] Rehli, M.; Krause, S. W.; Kreutz, M.; Andreesen, R.: Carboxypeptidase

[58837] M is identical to the MAX.1 antigen and its expression is

associated
[58838] with monocyte to macrophage differentiation. J. Biol. Chem. 270:
[58839] 15644-15649, 1995.

[58840] 9694.

[58841] Tan, F.; Chan, S. J.; Steiner, D. F.; Schilling, J. W.; Skidgel,

[58842] R. A.: Molecular cloning and sequencing of the cDNA for human membrane-bound
[58843] carboxypeptidase M: comparison with carboxypeptidases A, B, H, and
[58844] N. J. Biol. Chem. 264: 13165-13170, 1989.

[58845] 9695.

[58846] Gold, P.; Freedman, S. O.: Demonstration of tumor-specific antigens
[58847] in human colonic carcinomata by immunological tolerance and absorption
[58848] techniques. J. Exp. Med. 121: 439-462, 1965.

[58849] 9696.

[58850] Kamarck, M. E.; Elting, J. J.; Hart, J. T.; Goebel, S. J.; Rae,

[58851] P. M. M.; Nothdurft, M. A.; Nedwin, J. J.; Barnett, T. R.:

Carcinoembryonic
[58852] antigen family: expression in a mouse L-cell transfectant and characterization
[58853] of a partial cDNA in bacteriophage lambda-gt11. Proc. Nat. Acad.
[58854] Sci. 84: 5350-5354, 1987.

[58855] 9697.

[58856] Nishi, M.; Inazawa, J.; Inoue, K.; Nakagawa, H.; Taniwaki, M.;
[58857] Misawa, S.; Oikawa, S.; Nakazato, H.; Abe, T.: Regional chromosomal
[58858] assignment of carcinoembryonic antigen gene (CEA) to chromosome 19
[58859] at band q13.2. Cancer Genet. Cytogenet. 54: 77-81, 1991.
[58860] 9698.

[58861] Oikawa, S.; Nakazato, H.; Kosaki, G.: Primary structure of human
[58862] carcinoembryonic antigen (CEA) deduced from cDNA se-quence. Biochem.
[58863] Biophys. Res. Commun. 142: 511-528, 1987.

[58864] 9699.

[58865] Thompson, J.; Zimmermann, W.: The carcinoembryonic antigen gene
[58866] family: structure, expression and evolution. Tumor Biol. 9: 63-83,
[58867] 1988.

[58868] 9700.

[58869] Thompson, J. A.; Pande, H.; Paxton, R. J.; Shively, L.; Padma,
[58870] A.; Simmer, R. L.; Todd, C. W.; Riggs, A. D.; Shively, J. E.: Molecular
[58871] cloning of a gene belonging to the carcinoembryonic antigen gene family
[58872] and discussion of a domain model. Proc. Nat. Acad. Sci. 84: 2965-2969,
[58873] 1987.

[58874] 9701.

[58875] Willcocks, T. C.; Craig, I. W.: Characterization of the genomic
[58876] organization of human carcinoembryonic antigen (CEA): comparison with
[58877] other family members and sequence analysis of 5-prime controlling

[58878] region. Genomics 8: 492-500, 1990.

[58879] 9702.

[58880] Willcocks, T. C.; Craig, S. P.; Coates, D.; Craig, I. W.: Cod-
ing
[58881] sequences for carcinoembryonic antigen (CEA) assigned to
human chromosome
[58882] 19q13. (Abstract) Cytogenet. Cell Genet. 46: 716 only,
1987.
[58883] 9703.

[58884] Zimmer, R.; Thomas, P.: Mutations in the carcinoembroy-
onic antigen
[58885] gene in colorectal cancer patients: implications on liver
metastasis. Cancer
[58886] Res. 61: 2822-2826, 2001.

[58887] 9704.

[58888] Zimmermann, W.; Ortlieb, B.; Friedrich, R.; von Kleist, S.:
Isolation
[58889] and characterization of cDNA clones encoding the human
carcinoembryonic
[58890] antigen reveal a highly conserved repeating structure.
Proc. Nat.

[58891] Acad. Sci. 84: 2960-2964, 1987.

[58892] 9705.

[58893] Zimmermann, W.; Weber, B.; Ortlieb, B.; Rudert, F.;
Schempp, W.;
[58894] Fiebig, H.-H.; Shively, J. E.; von Kleist, S.; Thompson, J. A.:
Chromosomal
[58895] localization of the carcinoembryonic antigen gene family
and differential
[58896] expression in various tumors. Cancer Res. 48:
2550-2554, 1988.
[58897] 9706.

[58898] Brewton, R. G.; Wood, B. M.; Ren, Z.-X.; Gong, Y.; Tiller, G.
E.;
[58899] Warman, M. L.; Lee, B.; Horton, W. A.; Olsen, B. R.; Baker,
J. R.;
[58900] Mayne, R.: Molecular cloning of the alpha-3 chain of hu-
man type IX
[58901] collagen: linkage of the gene COL9A3 to chromosome
20q13.3. Genomics 30:
[58902] 329-336, 1995.

[58903] 9707.

[58904] Lohiniva, J.; Paassilta, P.; Seppanen, U.; Vierimaa, O.; Kivirikko,
[58905] S.; Ala-Kokko, L.: Splicing mutations in the COL3 domain of collagen
[58906] IX cause multiple epiphyseal dysplasia. Am. J. Med. Genet. 90: 216-222,
[58907] 2000.

[58908] 9708.

[58909] Paassilta, P.; Lohiniva, J.; Goring, H. H. H.; Perala, M.; Raina,
[58910] S. S.; Karppinen, J.; Hakala, M.; Palm, T.; Kroger, H.; Kaitila, I.;
[58911] Vanharanta, H.; Ott, J.; Ala-Kokko, L.: Identification of a novel
[58912] common genetic risk factor for lumbar disk disease. J.A.M.A. 285:
[58913] 1843-1849, 2001.

[58914] 9709.

[58915] Tiller, G. E.; Warman, M. L.; Gong, Y.; Knoll, J. H. M.; Mayne,
[58916] R.; and Brewton, R. G.: Physical and linkage mapping of the gene

[58917] for the alpha-3 chain of type IX collagen, COL9A3, to human chromosome
[58918] 20q13.3. Cytogenet. Cell Genet. 81: 205-207, 1998.

[58919] 9710.

[58920] Henry, I.; Bernheim, A.; Bernard, M.; van der Rest, M.; Kimura,
[58921] T.; Jeanpierre, C.; Barichard, F.; Berger, R.; Olsen, B. R.; Ramirez,
[58922] F.; Junien, C.: Mapping of a human fibrillar collagen gene, pro alpha-1(XI)(COL11A1),
[58923] to the p21 region of chromosome 1. Genomics 3: 87-90, 1988.
[58924] 9711.

[58925] Bernard, M.; Yoshioka, H.; Rodriguez, E.; van der Rest, M.; Kimura,
[58926] T.; Ninomiya, Y.; Olsen, B. R.; Ramirez, F.: Cloning and sequencing
[58927] of pro-alpha1(XI) collagen cDNA demonstrated that type XI belongs
[58928] to the fibrillar class of collagens and reveals that the expression
[58929] of the gene is not restricted to cartilaginous tissue. J. Biol.

Chem. 263:
[58930] 17159-17166, 1988.

[58931] 9712.

[58932] Li, Y.; Lacerda, D. A.; Warman, M.; Beier, D.; Oxford, J. T.; Morris,
[58933] N.; Andrikopoulos, K.; Ramirez, F.; Taylor, B.; Seegmiller, R.; Olsen,
[58934] B. R.: An abnormality in alpha-1(XI) collagen causes autosomal recessive
[58935] chondrodysplasia (cho) in mice. (Abstract) Molec. Biol. Cell 4 (suppl.
[58936] 1): 7a, 1993.

[58937] 9713.

[58938] Li, Y.; Lacerda, D. A.; Warman, M. L.; Beier, D. R.; Yoshioka,
[58939] H.; Ninomiya, Y.; Oxford, J. T.; Morris, N. P.; Andrikopoulos, K.;
[58940] Ramirez, F.; Wardell, B. B.; Lifferth, G. D.; Teuscher, C.; Woodward,
[58941] S. R.; Taylor, B. A.; Seegmiller, R. E.; Olsen, B. R.: A fibrillar

[58942] collagen gene, Col11a1, is essential for skeletal morpho-

genesis. Cell 80:

[58943] 423-430, 1995.

[58944] 9714.

[58945] Morris, N. P.; Bachinger, H. P.: Type XI collagen is a het-
erotrimer

[58946] with the composition (1alpha,2alpha,3alpha) retaining
non-triple-helical

[58947] domains. J. Biol. Chem. 262: 11345-11350, 1987.

[58948] 9715.

[58949] Ichinose, H.; Ohye, T.; Takahashi, E.; Seki, N.; Hori, T.;
Segawa,

[58950] M.; Nomura, Y.; Endo, K.; Tanaka, H.; Tsuji, S.; Fujita, K.;
Nagatsu,

[58951] T.: Hereditary progressive dystonia with marked diurnal
fluctuation

[58952] caused by mutations in the GTP cyclohydrolase I gene.
Nature Genet. 8:

[58953] 236-242, 1994.

[58954] 9716.

[58955] Schuback, D.; Kramer, P.; Ozelius, L.; Holmgren, G.; Fors-
gren,

[58956] L.; Kyllerman, M.; Wahlstrom, J.; Craft, C. M.; Nygaard, T.; Brin,

[58957] M.; de Leon, D.; Bressman, S.; Moskowitz, C. B.; Burke, R. E.; Sanner,

[58958] G.; Drugge, U.; Gusella, J. F.; Fahn, S.; Breakefield, X. O.: Dopamine

[58959] beta-hydroxylase gene excluded in four subtypes of hereditary dystonia. Hum.

[58960] Genet. 87: 311-316, 1991.

[58961] 9717.

[58962] Steinberger, D.; Korinthenberg, R.; Topka, H.; Berghauser, M.;

[58963] Wedde, R.; Muller, U.: Dopa-responsive dystonia: mutation analysis

[58964] of GCH1 and analysis of therapeutic doses of L-dopa. Neurology 55:

[58965] 1735-1737, 2000.

[58966] 9718.

[58967] Arahata, K.; Hayashi, Y. K.; Mizuno, Y.; Yoshida, M.; Ozawa, E.

[58968] : Dystrophin-associated glycoprotein and dystrophin co-localisation

[58969] at sarcolemma in Fukuyama congenital muscular dystrophy. (Letter) Lancet 342:
[58970] 623-624, 1993.

[58971] 9719.

[58972] Campanelli, J. T.; Roberds, S. L.; Campbell, K. P.; Scheller, R.
[58973] H.: A role for dystrophin-associated glycoproteins and utrophin in
[58974] agrin-induced AChR clustering. Cell 77: 663-674, 1994.

[58975] 9720.

[58976] Cao, W.; Henry, M. D.; Borrow, P.; Yamada, H.; Elder, J. H.; Ravkov,
[58977] E. V.; Nichol, S. T.; Compans, R. W.; Campbell, K. P.; Oldstone, M.
[58978] B. A.: Identification of alpha-dystroglycan as a receptor for lymphocytic
[58979] choriomeningitis virus and Lassa fever virus. Science 282: 2079-2081,
[58980] 1998.

[58981] 9721.

[58982] Ottman, R.; Risch, N.; Hauser, W. A.; Pedley, T. A.; Lee, J.

H.;

[58983] Barker-Cummings, C.; Lustenberger, A.; Nagle, K. J.; Lee, K. S.; Scheuer,

[58984] M. L.; Neystat, M.; Susser, M.; Wilhelmsen, K. C.: Localization of

[58985] a gene for partial epilepsy to chromosome 10q. Nature Genet. 10:

[58986] 56-60, 1995.

[58987] 9722.

[58988] Stogmann, E.; Zimprich, A.; Baumgartner, C.; Aull-Watschinger,

[58989] S.; Hollt, V.; Zimprich, F.: A functional polymorphism in the prodynorphin

[58990] gene promoter is associated with temporal lobe epilespy. Ann. Neurol. 51:

[58991] 260-263, 2002.

[58992] 9723.

[58993] Summar, M. L.; Phillips, J. A., III; Battey, J.; Castiglione, C.

[58994] M.; Kidd, K. K.; Maness, K. J.; Weiffenbach, B.; Gravius, T. C.:

[58995] Linkage relationships of human arginine vasopressin-

neurophysin-II
[58996] and oxytocin-neurophysin-I to prodynorphin and other loci on chromosome
[58997] 20. Molec. Endocr. 4: 947-950, 1990.

[58998] 9724.

[58999] Zimprich, A.; Kraus, J.; Woltje, M.; Mayer, P.; Rauch, E.; Hollt,
[59000] V.: An allelic variation in the human prodynorphin gene promoter
[59001] alters stimulus-induced expression. J. Neurochem. 74: 472-477, 2000.
[59002] 9725.

[59003] Craig, S. P.; Day, I. N. M.; Thompson, R. J.; Craig, I. W.: Localization
[59004] of human neurone-specific enolase to chromosome 12p13. (Abstract) Cytogenet.
[59005] Cell Genet. 51: 980 only, 1989.

[59006] 9726.

[59007] Craig, S. P.; Day, I. N. M.; Thompson, R. J.; Craig, I. W.: Localisation
[59008] of neurone-specific enolase (ENO2) to 12p13. Cytogenet.

Cell Genet. 54:
[59009] 71-73, 1990.

[59010] 9727.

[59011] Feo, S.; Oliva, D.; Barbieri, G.; Xu, W.; Fried, M.; Giallongo,

[59012] A.: The gene for the muscle-specific enolase is on the short arm
[59013] of human chromosome 17. Genomics 6: 192-194, 1990.

[59014] 9728.

[59015] Grzeschik, K.-H.: Assignment of human genes: beta-glucuronidase
[59016] to chromosome 7, adenylate kinase-1 to 9, a second enzyme with enolase
[59017] activity to 12, and mitochondrial IDH to 15. Cytogenet. Cell Genet. 16:
[59018] 142-148, 1976. Note: Alternate: Birth Defects Orig. Art. Ser. 12(7):
[59019] 142-148, 1976.

[59020] 9729.

[59021] Herbschleb-Voogt, E.; Monteba-van Heuvel, M.; Wijnen, L. M. M.;
[59022] Westerveld, A.; Pearson, P. L.; Meera Khan, P.: Chromoso-

mal assignment
[59023] and regional localization of CS, ENO-2, GAPDH, LDH-B, PEP-B and TPI
[59024] in man-rodent cell hybrids. Cytogenet. Cell Genet. 22: 482-486,
[59025] 1978.

[59026] 9730.

[59027] Hinks, L. J.; Day, I. N. M.: Further studies of enolase loci.

[59028] (Abstract) Cytogenet. Cell Genet. 58: 1854 only, 1991.

[59029] 9731.

[59030] Law, M. L.; Kao, F.: Regional mapping of the gene coding for enolase-2
[59031] on human chromosome 12. J. Cell Sci. 53: 245-254, 1982.
[59032] 9732.

[59033] Mattei, J. F.; Baeteman, M. A.; Mattei, M. G.; Ardissonne, J. P.;
[59034] Giraud, F.: Regional assignments of CS and ENO2 on chromosome 12.
[59035] (Abstract) Cytogenet. Cell Genet. 32: 297 only, 1982.

[59036] 9733.

[59037] Oliva, D.; Cali, L.; Feo, S.; Giallongo, A.: Complete structure
[59038] of the human gene encoding neuron-specific enolase. Genomics 10:
[59039] 157-165, 1991.

[59040] 9734.

[59041] Jenkins, N. A.; Justice, M. J.; Gilbert, D. J.; Chu, M.-L.; Copeland,
[59042] N. G.: Nidogen/entactin (Nid) maps to the proximal end of mouse chromosome
[59043] 13 linked to beige (bg) and identifies a new region of homology between
[59044] mouse and human chromosomes. Genomics 9: 401-403, 1991.
[59045] 9735.

[59046] Olavesen, M. G.; Bentley, E.; Mason, R. V. F.; Stephens, R. J.;
[59047] Ragoussis, J.: Fine mapping of 39 ESTs on human chromosome 6p23-p25. Genomics 46:
[59048] 303-306, 1997.

[59049] 9736.

[59050] Boyd, C. D.; Toth-Fejel, S.; Gadi, I. K.; Litt, M.; Condon, M.

[59051] R.; Kolbe, M.; Hagen, I. K.; Kurkinen, M.; Mackenzie, J. W.; Magenis,

[59052] E.: The genes coding for human pro alpha-1(IV) collagen and pro alpha-2(IV)

[59053] collagen are both located at the end of the long arm of chromosome

[59054] 13. Am. J. Hum. Genet. 42: 309-314, 1988.

[59055] 9737.

[59056] Brazel, D.; Pollner, R.; Oberbaumer, I.; Kuhn, K.: Human basement

[59057] membrane collagen (type IV): the amino acid sequence of the alpha-2(IV)

[59058] chain and its comparison with the alpha-1(IV) chain reveals deletions

[59059] in the alpha-1(IV) chain. Europ. J. Biochem. 172: 35-42, 1988.

[59060] 9738.

[59061] Griffin, C. A.; Emanuel, B. S.; Hansen, J. R.; Cavenee, W. K.;

[59062] Myers, J. C.: Human collagen genes encoding basement membrane alpha-1(IV)

[59063] and alpha-2(IV) chains map to the distal long arm of chromosome 13. Proc.

[59064] Nat. Acad. Sci. 84: 512-516, 1987.

[59065] 9739.

[59066] Haniel, A.; Welge-Lussen, U.; Kuhn, K.; Poschl, E.: Identification

[59067] and characterization of a novel transcriptional silencer in the human

[59068] collagen type IV gene COL4A2. J. Biol. Chem. 270: 11209-11215, 1995.

[59069] 9740.

[59070] Killen, P. D.; Francomano, C. A.; Yamada, Y.; Modi, W. S.; O'Brien,

[59071] S. J.: Partial structure of the human alpha-2(IV) collagen chain

[59072] and chromosomal localization of the gene (COL4A2). Hum. Genet. 77:

[59073] 318-324, 1987.

[59074] 9741.

[59075] Solomon, E.; Hall, V.; Kurkinen, M.: The human alpha-2(IV) collagen

[59076] gene, COL4A2, is syntenic with the alpha-1(IV) gene, COL4A1, on chromosome
[59077] 13. Ann. Hum. Genet. 51: 125-127, 1987.

[59078] 9742.

[59079] Apte, S.; Mattei, M.-G.; Olsen, B. R.: Cloning of human alpha-1(X)
[59080] collagen DNA and localization of the COL10A1 gene to the q21-q22 region
[59081] of human chromosome 6. FEBS Lett. 282: 393-396, 1991.

[59082] 9743.

[59083] Apte, S. S.; Seldin, M. F.; Hayashi, M.; Olsen, B. R.: Cloning

[59084] of the human and mouse type X collagen genes and mapping of the mouse
[59085] type X collagen gene to chromosome 10. Europ. J. Biochem. 206: 217-224,
[59086] 1992.

[59087] 9744.

[59088] Chan, D.; Weng, Y. M.; Graham, H. K.; Sillence, D. O.; Bateman,
[59089] J. F.: A nonsense mutation in the carboxyl-terminal domain of type

[59090] X collagen causes haploinsufficiency in Schmid metaphyseal chondrodysplasia. J.

[59091] Clin. Invest. 101: 1490-1499, 1998.

[59092] 9745.

[59093] Dharmavaram, R. M.; Elberson, M. A.; Peng, M.; Kirson, L. A.; Kelley,

[59094] T. E.; Jimenez, S. A.: Identification of a mutation in type X collagen

[59095] in a family with Schmid metaphyseal chondrodysplasia. Hum. Molec.

[59096] Genet. 3: 507-509, 1994.

[59097] 9746.

[59098] Alper, C. A.; Marcus, D.; Raum, D.; Petersen, B. H.; Spira, T.

[59099] J.: Genetic polymorphism in C8 beta-chains: evidence for two unlinked

[59100] genetic loci for the eighth component of human complement (C8). J.

[59101] Clin. Invest. 72: 1526-1531, 1983.

[59102] 9747.

[59103] Densen, P.; Brown, E. J.; O'Neill, G. J.; Tedesco, F.; Clark, R.

[59104] A.; Frank, M. M.; Webb, D.; Myers, J.: Inherited deficiency of C8

[59105] in a patient with recurrent meningococcal infections: further evidence

[59106] for a dysfunctional C8 molecule and nonlinkage to the HLA system. J.

[59107] Clin. Immun. 3: 90-99, 1983.

[59108] 9748.

[59109] Giraldo, G.; Degos, L.; Beth, E.; Sasportes, M.; Marcelli, A.;

[59110] Gharbi, R.; Day, N. K.: C8 deficiency in a family with xeroderma

[59111] pigmentosum: lack of linkage to HLA region. Clin. Immun. Immunopath. 8:

[59112] 377-384, 1977.

[59113] 9749.

[59114] Jasin, H. E.: Absence of the eighth component of complement in

[59115] association with systemic lupus erythematosus-like disease. J. Clin.

[59116] Invest. 60: 709-715, 1977.

[59117] 9750.

[59118] Kolb, W. P.; Muller-Eberhard, H. J.: The membrane attack mechanism
[59119] of complement: the three polypeptide chain structure of the eighth
[59120] component (C8). J. Exp. Med. 143: 1131-1139, 1976.

[59121] 9751.

[59122] Komatsu, M.; Imaoka, K.; Satoh, M.; Mikami, H.: Hereditary C8-alpha-gamma
[59123] deficiency associated with dwarfism in the rabbit. J. Hered. 81:
[59124] 413-417, 1990.

[59125] 9752.

[59126] Komatsu, M.; Yamamoto, K.; Kawashima, T.; Migita, S.: Genetic
[59127] deficiency of the alpha-gamma-subunit of the eighth complement component
[59128] in the rabbit. J. Immun. 134: 2607-2609, 1985.

[59129] 9753.

[59130] Marcus, D.; Spira, T. J.; Petersen, B. H.; Raum, D.; Alper, C.

[59131] A.: There are two unlinked genetic loci for human C8.

(Abstract) Molec.
[59132] Immun. 19: 1385 only, 1982.

[59133] 9754.

[59134] Matthews, N.; Stark, J. M.; Harper, P. S.; Doran, J.; Jones, D.

[59135] M.: Recurrent meningococcal infections associated with a functional
[59136] deficiency of the C8 component of human complement. Clin. Exp. Immun. 39:
[59137] 53-59, 1980.

[59138] 9755.

[59139] Merritt, A. D.; Petersen, B. H.; Biegel, A. A.; Meyers, D. A.;

[59140] Brooks, G. F.; Hodes, M. E.: Chromosome 6: linkage of the eighth
[59141] component of complement (C8) to the histocompatibility region (HLA). Cytogenet.
[59142] Cell Genet. 16: 331-334, 1976.

[59143] 9756.

[59144] Michelotti, G. A.; Snider, J. V.; Sodetz, J. M.: Genomic or-
[59145] ganization of human complement protein C8-alpha and further ex-

amination of its

[59146] linkage to C8-beta. Hum. Genet. 95: 513-518, 1995.

[59147] 9757.

[59148] Nakamura, S.; Ohue, O.; Abe, K.: Genetic polymorphism of human

[59149] complement component C81 in the Japanese population. Hum. Genet. 72:

[59150] 344-347, 1986.

[59151] 9758.

[59152] Pericak-Vance, M. A.; Elston, R. C.; Spira, T. J.; Band, J.:

[59153] Segregation and linkage analysis of immunochemical C8 levels in a

[59154] family with C8 beta-chain deficiency. (Abstract) Am. J. Hum. Genet. 34:

[59155] 109A only, 1982.

[59156] 9759.

[59157] Petersen, B. H.; Graham, J. A.; Brooks, G. F.: Human defi-ciency

[59158] of the eighth component of complement: the requirement of C8 for serum

[59159] Neisseria gonorrhoeae bactericidal activity. J. Clin. Invest.

57:

[59160] 283-290, 1976.

[59161] 9760.

[59162] Pickering, R. J.; Rynes, R. I.; LoCascio, N.; Monahan, J. B.;

[59163] Sodetz, J. M.: Identification of the alpha-gamma subunit of the eighth

[59164] component of complement (C8) in a patient with systemic lupus erythematosus

[59165] and absent C8 activity: patient and family studies. Clin. Immun.

[59166] Immunopath. 23: 323-334, 1982.

[59167] 9761.

[59168] Rittner, C.; Hargesheimer, W.; Mollenhauer, E.: Population and

[59169] formal genetics of the human C81(alpha-gamma) poly-morphism. Hum.

[59170] Genet. 67: 166-169, 1984.

[59171] 9762.

[59172] Rittner, C.; Hargesheimer, W.; Stradmann, B.; Bertrams, J.; Baur,

[59173] M. P.; Petersen, B. H.: Human C81 (alpha-gamma) poly-

morphism: detection

[59174] in the alpha-gamma subunit on SDS-PAGE, formal genetics and linkage

[59175] relationship. Am. J. Hum. Genet. 38: 482-491, 1986.

[59176] 9763.

[59177] Rogde, S.; Gedde-Dahl, T., Jr.; Teisberg, P.; Jonassen, R.; Hoyheim,

[59178] B.; Olaisen, B.: Linkage and association studies with C8A and C8B

[59179] RFLPs on chromosome 1. Ann. Hum. Genet. 56: 233-242, 1992.

[59180] 9764.

[59181] Rogde, S.; Mevag, B.; Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg,

[59182] P.: Structural genes for complement factor C8 on chromosome 1. (Abstract) Cytogenet.

[59183] Cell Genet. 37: 571 only, 1984.

[59184] 9765.

[59185] Rogde, S.; Mevag, B.; Teisberg, P.; Gedde-Dahl, T., Jr.; Tedesco,

[59186] F.; Olaisen, B.: Genetic polymorphism of complement

component C8. Hum.

[59187] Genet. 70: 211-216, 1985.

[59188] 9766.

[59189] Rogde, S.; Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg, P.: Two

[59190] complement component C8 loci are localized between PGM1 and Rh on

[59191] chromosome 1. (Abstract) Cytogenet. Cell Genet. 40: 734-735, 1985.

[59192] 9767.

[59193] Rogde, S.; Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg, P.: The

[59194] C8A and C8B loci are closely linked on chromosome 1. Ann. Hum. Genet. 50:

[59195] 139-144, 1986.

[59196] 9768.

[59197] Tedesco, F.; Densen, P.; Villa, M. A.; Petersen, B. H.; Sirchia,

[59198] G.: Two types of dysfunctional eighth component of com-plement (C8)

[59199] molecules in C8 deficiency in man: reconstitution of nor-

mal C8 from
[59200] the mixture of two abnormal C8 molecules. J. Clin. Invest. 71: 183-191,
[59201] 1983.

[59202] 9769.

[59203] Glaser, R. L.; Jiang, W.; Boyadjiev, S. A.; Tran, A. K.; Zachary,
[59204] A. A.; Van Maldergem, L.; Johnson, D.; Walsh, S.; Oldridge, M.; Wall,
[59205] S. A.; Wilkie, A. O. M.; Jabs, E. W.: Paternal origin of FGFR2 mutations
[59206] in sporadic cases of Crouzon syndrome and Pfeiffer syndrome. Am.
[59207] J. Hum. Genet. 66: 768-777, 2000.

[59208] 9770.

[59209] Jones, K. L.; Smith, D. W.; Harvey, M. A. S.; Hall, B. D.; Quan,
[59210] L.: Older paternal age and fresh gene mutation: data on additional
[59211] disorders. J. Pediat. 86: 84-88, 1975.

[59212] 9771.

[59213] Meyers, G. A.; Orlow, S. J.; Munro, I. R.; Przylepa, K. A.; Jabs,
[59214] E. W.: Fibroblast growth factor receptor 3 (FGFR3) transmembrane
[59215] mutation in Crouzon syndrome with acanthosis nigricans. Nature Genet. 11:
[59216] 462-464, 1995.

[59217] 9772.

[59218] Preston, R. A.; Post, J. C.; Keats, B. J. B.; Aston, C. E.; Ferrell,
[59219] R. E.; Priest, J.; Nouri, N.; Losken, H. W.; Morris, C. A.; Hurtt,
[59220] M. R.; Mulvihill, J. J.; Ehrlich, G. D.: A gene for Crouzon craniofacial
[59221] dysostosis maps to the long arm of chromosome 10. Nature Genet. 7:
[59222] 149-153, 1994.

[59223] 9773.

[59224] Reardon, W.; Winter, R. M.; Rutland, P.; Pulleyn, L. J.; Jones,
[59225] B. M.; Malcolm, S.: Mutations in the fibroblast growth factor receptor

[59226] 2 gene cause Crouzon syndrome. Nature Genet. 8:
98-103, 1994.
[59227] 9774.

[59228] Jankowski, S. A.; Gumucio, D. L.: Genes for tensin, villin
and
[59229] desmin are linked on mouse chromosome 1. Mammalian
Genome 6: 744-745,
[59230] 1995.

[59231] 9775.

[59232] Quax, W.; Meera Khan, P.; Quax-Jeuken, Y.; Bloemendal,
H.: The
[59233] human desmin and vimentin genes are located on differ-
ent chromosomes. Gene 38:
[59234] 189-196, 1985.

[59235] 9776.

[59236] Sibilia, M.; Fleischmann, A.; Behrens, A.; Stingl, L.; Carroll,
[59237] J.; Watt, F. M.; Schlessinger, J.; Wagner, E. F.: The EGF re-
ceptor
[59238] provides an essential survival signal for SOS-dependent
skin tumor
[59239] development. Cell 102: 211-220, 2000.

[59240] 9777.

[59241] Spurr, N. K.; Solomon, E.; Jansson, M.; Sheer, D.; Goodfel-
low,
[59242] P. N.; Bodmer, W. F.; Vennstrom, B.: Chromosomal locali-
sation of
[59243] the human homologues to the oncogenes erbA and B.
EMBO J. 3: 159-163,
[59244] 1984.

[59245] 9778.

[59246] Thaung, C.; West, K.; Clark, B. J.; McKie, L.; Morgan, J. E.;

[59247] Arnold, K.; Nolan, P. M.; Peters, J.; Hunter, A. J.; Brown, S.
D.
[59248] M.; Jackson, I. J.; Cross, S. H.: Novel ENU-induced eye mu-
tations
[59249] in the mouse: models for human eye disease. Hum. Molec.
Genet. 11:
[59250] 755-767, 2002.

[59251] 9779.

[59252] Baumann, C. A.; Ribon, V.; Kanzaki, M.; Thurmond, D. C.;
Mora,
[59253] S.; Shigematsu, S.; Bickel, P. E.; Pessin, J. E.; Saltiel, A. R.:

[59254] CAP defines a second signalling pathway required for in-
sulin-stimulated
[59255] glucose transport. Nature 407: 202-207, 2000.

[59256] 9780.

[59257] Bickel, P. E.; Scherer, P. E.; Schnitzer, J. E.; Oh, P.; Lisanti,

[59258] M. P.; Lodish, H. F.: Flotillin and epidermal surface antigen
define
[59259] a new family of caveolae-associated integral membrane
proteins. J.
[59260] Biol. Chem. 272: 13793-13802, 1997.

[59261] 9781.

[59262] Cho, Y.-J.; Chema, D.; Moskow, J. J.; Cho, M.; Schroeder,
W. T.;
[59263] Overbeek, P.; Buchberg, A. M.; Duvic, M.: Epidermal sur-
face antigen
[59264] (MS17S1) is highly conserved between mouse and human.
Genomics 27:
[59265] 251-258, 1995.

[59266] 9782.

[59267] Kayes, L. M.; Schroeder, W. T.; Marchuk, D. A.; Collins, F.
S.;

[59268] Riccardi, V. M.; Duvic, M.; Stephens, K.: The gene for a novel epidermal
[59269] antigen maps near the neurofibromatosis 1 gene. Genomics 14: 369-376,
[59270] 1992.

[59271] 9783.

[59272] Schroeder, W. T.; Siciliano, M. J.; Stewart-Galetka, S. L.; Duvic,
[59273] M.: The human gene for an epidermal surface antigen (M17S1) is located
[59274] at 17q11-12. Genomics 11: 481-482, 1991.

[59275] 9784.

[59276] Schroeder, W. T.; Stephens, K.; Stewart-Galetka, S.; Riccardi,
[59277] V.; Duvic, M.: The gene for an epidermal surface antigen is in close
[59278] proximity to the locus for von Recklinghausen neurofibromatosis. (Abstract) Clin.
[59279] Res. 39: 323A only, 1991.

[59280] 9785.

[59281] Mitchell, B. D.; Cole, S. A.; Bauer, R. L.; Iturria, S. J.; Ro-

driguez,
[59282] E. A.; Blangero, J.; MacCluer, J. W.; Hixson, J. E.: Genes influencing
[59283] variation in serum osteocalcin concentrations are linked to markers
[59284] on chromosomes 16q and 20q. J. Clin. Endocr. Metab. 85: 1362-1366,
[59285] 2000.

[59286] 9786.

[59287] Richards, A. J.; Yates, J. R. W.; Williams, R.; Payne, S. J.;

[59288] Pope, F. M.; Scott, J. D.; Snead, M. P.: A family with Stickler syndrome
[59289] type 2 has a mutation in the COL11A1 gene resulting in the substitution
[59290] of glycine 97 by valine in alpha-1(XI) collagen. Hum. Molec. Genet. 5:
[59291] 1339-1343, 1996.

[59292] 9787.

[59293] Cohn, R. D.; Henry, M. D.; Michele, D. E.; Barresi, R.; Saito,

[59294] F.; Moore, S. A.; Flanagan, J. D.; Skwarchuk, M. W.; Robbins, M. E.;

[59295] Mendell, J. R.; Williamson, R. A.; Campbell, K. P.: Disruption of
[59296] Dag1 in differentiated skeletal muscle reveals a role for dystroglycan
[59297] in muscle regeneration. Cell 110: 639-648, 2002.

[59298] 9788.

[59299] Apperley, J. F.; Gardembas, M.; Melo, J. V.; Russell-Jones, R.;
[59300] Bain, B. J.; Baxter, E. J.; Chase, A.; Chessells, J. M.; Colombat,
[59301] M.; Dearden, C. E.; Dimitrijevic, S.; Mahon, F.-X.; Marin, D.; Nikolova,
[59302] Z.; Olavarria, E.; Silberman, S.; Schultheis, B.; Cross, N. C. P.;
[59303] Goldman, J. M.: Response to imatinib mesylate in patients with chronic
[59304] myeloproliferative diseases with rearrangements of the platelet-derived
[59305] growth factor receptor beta. New Eng. J. Med. 347: 481-487, 2002.
[59306] 9789.

[59307] Shanske, A.; Bogdanow, A.; Shprintzen, R. J.; Marion, R.

W.:

[59308] Marshall syndrome and a defect at the COL11A1 locus. (Letter) Am.

[59309] J. Hum. Genet. 63: 1558-1559, 1998.

[59310] 9790.

[59311] Yoshioka, H.; Inoguchi, K.; Khaleduzzaman, M.; Ninomiya, Y.; Andrikopoulos,

[59312] K.; Ramirez, F.: Coding sequence and alternative splicing of the

[59313] mouse alpha-1(XI) collagen gene (Col11a1). Genomics 28: 337-340,

[59314] 1995.

[59315] 9791.

[59316] Zhidkova, N. I.; Justice, S. K.; Mayne, R.: Alternative mRNA

[59317] processing occurs in the variable region of the pro-alpha-1(XI) and

[59318] pro-alpha-2(XI) collagen chains. J. Biol. Chem. 270: 9486-9493,

[59319] 1995.

[59320] 9792.

[59321] Spear, P. G.: A welcome mat for leprosy and Lassa fever.

Science 282:
[59322] 1999-2000, 1998.

[59323] 9793.

[59324] Tinsley, J. M.; Blake, D. J.; Zuellig, R. A.; Davies, K. E.:

[59325] Increasing complexity of the dystrophin-associated protein complex. Proc.
[59326] Nat. Acad. Sci. 91: 8307-8313, 1994.

[59327] 9794.

[59328] Williamson, R. A.; Henry, M. D.; Daniels, K. J.; Hrstka, R. F.;

[59329] Lee, J. C.; Sunada, Y.; Ibraghimov-Beskrovnaya, O.; Campbell, K. P.
[59330] : Dystroglycan is essential for early embryonic development: disruption
[59331] of Reichert's membrane in Dag1-null mice. Hum. Molec. Genet. 6:
[59332] 831-841, 1997.

[59333] 9795.

[59334] Yamada, H.; Denzer, A. J.; Hori, H.; Tanaka, T.; Anderson, L.
[59335] V. B.; Fujita, S.; Fukuta-Ohi, H.; Shimizu, T.; Ruegg, M. A.;

Matsumura,
[59336] K.: Dystroglycan is a dual receptor for agrin and laminin-2 in Schwann
[59337] cell membrane. J. Biol. Chem. 271: 23418-23423, 1996.

[59338] 9796.

[59339] Cote, P. D.; Moukhles, H.; Lindenbaum, M.; Carbonetto, S.: Chimaeric
[59340] mice deficient in dystroglycans develop muscular dystro-phy and have
[59341] disrupted myoneural synapses. Nature Genet. 23: 338-342, 1999.

[59342] 9797.

[59343] Gee, S. H.; Montanaro, F.; Lindenbaum, M. H.; Carbonetto, S.:
[59344] Dystroglycan-alpha, a dystrophin-associated glycopro-tein, is a functional
[59345] agrin receptor. Cell 77: 675-686, 1994.

[59346] 9798.

[59347] Gorecki, D. C.; Derry, J. M. J.; Barnard, E. A.: Dystroglycan:

[59348] brain localisation and chromosome mapping in the mouse. Hum. Molec.

[59349] Genet. 3: 1589-1597, 1994.

[59350] 9799.

[59351] Hayashi, Y. K.; Ogawa, M.; Tagawa, K.; Noguchi, S.; Ishi-
hara, T.;
[59352] Nonaka, I.; Arahata, K.: Selective deficiency of alpha-
dystroglycan
[59353] in Fukuyama-type congenital muscular dystrophy. Neurol-
ogy 57: 115-121,
[59354] 2001.

[59355] 9800.

[59356] Henry, M. D.; Campbell, K. P.: A role for dystroglycan in
basement
[59357] membrane assembly. Cell 95: 859-970, 1998.

[59358] 9801.

[59359] Ibraghimov-Beskrovnaya, O.; Ervasti, J. M.; Leveille, C. J.;
Slaughter,
[59360] C. A.; Sernett, S. W.; Campbell, K. P.: Primary structure of
dystrophin-associated
[59361] glycoproteins linking dystrophin to the extracellular ma-
trix. Nature 355:
[59362] 696-702, 1992.

[59363] 9802.

[59364] Ibraghimov-Beskrovnaya, O.; Milatovich, A.; Ozcelik, T.; Yang,

[59365] B.; Francke, U.; Campbell, K. P.: Dystroglycan: tissue distribution,

[59366] human muscle cDNA, genomic structure and chromosome mapping. (Abstract) Am.

[59367] J. Hum. Genet. 51 (suppl.): A130 only, 1992.

[59368] 9803.

[59369] Ibraghimov-Beskrovnaya, O.; Milatovich, A.; Ozcelik, T.; Yang,

[59370] B.; Koepnick, K.; Francke, U.; Campbell, K. P.: Human dystroglycan:

[59371] skeletal muscle cDNA, genomic structure, origin of tissue specific

[59372] isoforms and chromosomal localization. Hum. Molec. Genet. 2: 1651-1657,

[59373] 1993.

[59374] 9804.

[59375] Ma, J.; Nastuk, M. A.; McKechnie, B. A.; Fallon, J. R.: The agrin

[59376] receptor: localization in the postsynaptic membrane, interaction with
[59377] agrin, and relationship to the acetylcholine receptor. J. Biol. Chem. 268:
[59378] 25108-25117, 1993.

[59379] 9805.

[59380] Matsumura, K.; Nonaka, I.; Campbell, K. P.: Abnormal expression
[59381] of dystrophin-associated proteins in Fukuyama-type congenital muscular
[59382] dystrophy. Lancet 341: 521-522, 1993.

[59383] 9806.

[59384] Matsumura, K.; Tome, F. M. S.; Collin, H.; Azibi, K.; Chaouch,
[59385] M.; Kaplan, J.-C.; Fardeau, M.; Campbell, K. P.: Deficiency of the
[59386] 50K dystrophin-associated glycoprotein in severe childhood autosomal
[59387] recessive muscular dystrophy. Nature 359: 320-322, 1992.
[59388] 9807.

[59389] Matsumura, K.; Tome, F. M. S.; Ionasescu, V.; Ervasti, J. M.;

[59390] Anderson, R. D.; Romero, N. B.; Simon, D.; Recan, D.; Ka-
plan, J.-C.;

[59391] Fardeau, M.; Campbell, K. P.: Deficiency of dystrophin-as-
sociated

[59392] proteins in Duchenne muscular dystrophy patients lacking
COOH-terminal

[59393] domains of dystrophin. J. Clin. Invest. 92: 866-871, 1993.

[59394] 9808.

[59395] Michele, D. E.; Barresi, R.; Kanagawa, M.; Saito, F.; Cohn,
R.

[59396] D.; Satz, J. S.; Dollar, J.; Nishino, I.; Kelley, R. I.; Somer, H.;

[59397] Straub, V.; Mathews, K. D.; Moore, S. A.; Campbell, K. P.:
Post-translational

[59398] disruption of dystroglycan-ligand interactions in congeni-
tal muscular

[59399] dystrophies. Nature 418: 417-422, 2002.

[59400] 9809.

[59401] Moore, S. A.; Saito, F.; Chen, J.; Michele, D. E.; Henry, M.
D.;

[59402] Messing, A.; Cohn, R. D.; Ross-Barta, S. E.; Westra, S.;

Williamson,
[59403] R. A.; Hoshi, T.; Campbell, K. P.: Deletion of brain dystro-
glycan
[59404] recapitulates aspects of congenital muscular dystrophy.
Nature 418:
[59405] 422-425, 2002.

[59406] 9810.

[59407] Rambukkana, A.; Yamada, H.; Zanazzi, G.; Mathus, T.;
Salzer, J.
[59408] L.; Yurchenco, P. D.; Campbell, K. P.; Fischetti, V. A.: Role
of
[59409] alpha-dystroglycan as a Schwann cell receptor for My-
cobacterium leprae. Science 282:
[59410] 2076-2078, 1998.

[59411] 9811.

[59412] Sealock, R.; Froehner, S. C.: Dystrophin-associated pro-
teins
[59413] and synapse formation: is alpha-dystroglycan the agrin
receptor? Cell 77:
[59414] 617-619, 1994.

[59415] 9812.

[59416] Yamada, H.; Saito, F.; Fukuta-Ohi, H.; Zhong, D.; Hase, A.; Arai,
[59417] K.; Okuyama, A.; Maekawa, R.; Shimizu, T.; Matsumura, K.: Processing
[59418] of beta-dystroglycan by matrix metalloproteinase disrupts the link
[59419] between the extracellular matrix and cell membrane via the dystroglycan
[59420] complex. Hum. Molec. Genet. 10: 1563-1569, 2001.

[59421] 9813.

[59422] Blake, D. J.; Love, D. R.; Tinsley, J.; Morris, G. E.; Turley,
[59423] H.; Gatter, K.; Dickson, G.; Edwards, Y. H.; Davies, K. E.: Characterization
[59424] of a 4.8kb transcript from the Duchenne muscular dystrophy locus expressed
[59425] in schwannoma cells. Hum. Molec. Genet. 1: 103-109, 1992.
[59426] 9814.

[59427] Golub, T. R.; Barker, G. F.; Lovett, M.; Gilliland, D. G.: Fusion
[59428] of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic

[59429] leukemia with t(5;12) chromosomal translocation. Cell 77: 307-316,

[59430] 1994.

[59431] 9815.

[59432] Brissenden, J. E.; Ullrich, A.; Francke, U.: Chromosomal mapping

[59433] of loci for insulin-like growth factors I and II and for epidermal

[59434] growth factor in man. (Abstract) Am. J. Hum. Genet. 36: 133S only,

[59435] 1984.

[59436] 9816.

[59437] Brissenden, J. E.; Ullrich, A.; Francke, U.: Human chromosomal

[59438] mapping of genes for insulin-like growth factors I and II and epidermal

[59439] growth factor. Nature 310: 781-784, 1984.

[59440] 9817.

[59441] Morton, C. C.; Byers, M. G.; Nakai, H.; Bell, G. I.; Shows, T.

[59442] B.: Human genes for insulin-like growth factors I and II and epidermal

[59443] growth factor are located on 12q22-q24.1, 11p15, and 4q25-q27, respectively. Cytogenet.

[59444] Cell Genet. 41: 245-249, 1986.

[59445] 9818.

[59446] Chen, B.; Bronson, R. T.; Klaman, L. D.; Hampton, T. G.; Wang,

[59447] J.; Green, P. J.; Magnuson, T.; Douglas, P. S.; Morgan, J. P.; Neel,

[59448] B. G.: Mice mutant for Egfr and Shp2 have defective cardiac semilunar

[59449] valvulogenesis. Nature Genet. 24: 296-299, 2000.

[59450] 9819.

[59451] Ishii, H.; Baffa, R.; Numata, S.-I.; Murakumo, Y.; Rattan, S.;

[59452] Inoue, H.; Mori, M.; Fidanza, V.; Alder, H.; Croce, C. M.: The FEZ1

[59453] gene at chromosome 8p22 encodes a leucine-zipper protein, and its

[59454] expression is altered in multiple human tumors. Proc. Nat. Acad.

[59455] Sci. 96: 3928-3933, 1999.

[59456] 9820.

[59457] Li, G.; Hu, N.; Goldstein, A. M.; Tang, Z.-Z.; Roth, M. J.; Wang,
[59458] Q.-H.; Dawsey, S. M.; Han, X.-Y.; Ding, T.; Huang, J.; Giffen, C.;
[59459] Taylor, P. R.; Emmert-Buck, M. R.: Allelic loss on chromosome bands
[59460] 13q11-q13 in esophageal squamous cell carcinoma. Genes Chromosomes
[59461] Cancer 31: 390-397, 2001.

[59462] 9821.

[59463] Lo, H. S.; Hu, N.; Gere, S.; Lu, N.; Su, H.; Goldstein, A. M.;

[59464] Taylor, P. R.; Lee, M. P.: Identification of somatic mutations of
[59465] the RNF6 gene in human esophageal squamous cell carcinoma. Cancer
[59466] Res. 62: 4191-4193, 2002.

[59467] 9822.

[59468] Wang, M.; Price, C. E.; Han, J.; Cisler, J.; Imaizumi, K.; Van

[59469] Thienen, M. N.; DePaepe, A.; Godfrey, M.: Recurrent misplicing
[59470] of fibrillin exon 32 in two patients with neonatal Marfan

syndrome. Hum.
[59471] Molec. Genet. 4: 607-613, 1995.

[59472] 9823.

[59473] Beck, C.; Moulard, B.; Steinlein, O.; Guipponi, M.; Vallee, L.;
[59474] Montpied, P.; Baldy-Moulnier, M.; Malafosse, A.: A non-sense mutation
[59475] in the alpha-4 subunit of the nicotinic acetylcholine re-ceptor (CHRNA4)
[59476] cosegregates with 20q-linked benign neonatal familial convulsions
[59477] (EBN1). Neurobiol. Dis. 1: 95-99, 1994.

[59478] 9824.

[59479] Singh, N. A.; Charlier, C.; Stauffer, D.; DuPont, B. R.; Leach,
[59480] R. J.; Melis, R.; Ronen, G. M.; Bjerre, I.; Quattlebaum, T.; Murphy,
[59481] J. V.; McHarg, M. L.; Gagnon, D.; Rosales, T. O.; Peiffer, A.; Anderson,
[59482] V. E.; Leppert, M.: A novel potassium channel gene, KCNQ2, is mutated
[59483] in an inherited epilepsy of newborns. Nature Genet. 18:

25-29, 1998.

[59484] 9825.

[59485] Earnshaw, W. C.; Sullivan, K. F.; Machlin, P. S.; Cooke, C. A.;

[59486] Kaiser, D. A.; Pollard, T. D.; Rothfield, N. F.; Cleveland, D. W.

[59487] : Molecular cloning of cDNA for CENP-B, the major human centromere

[59488] autoantigen. J. Cell Biol. 104: 817-829, 1987.

[59489] 9826.

[59490] Seki, N.; Saito, T.; Kitagawa, K.; Masumoto, H.; Okazaki, T.; Hori,

[59491] T.-A.: Mapping of the human centromere protein B gene (CENPB) to

[59492] chromosome 20p13 by fluorescence in situ hybridization. Genomics 24:

[59493] 187-188, 1994.

[59494] 9827.

[59495] Sugimoto, K.; Yata, H.; Himeno, M.: Mapping of the human CENP-B

[59496] gene to chromosome 20 and the CENP-C gene to chro-

mosome 12 by a rapid
[59497] cycle DNA amplification procedure. Genomics 17:
240-242, 1993.
[59498] 9828.

[59499] Rothstein, J. D.; Martin, L. J.; Kuncl, R. W.: Decreased glu-
tamate
[59500] transport by the brain and spinal cord in amyotrophic lat-
eral sclerosis. New
[59501] Eng. J. Med. 326: 1464-1468, 1992.

[59502] 9829.

[59503] Gaynor, R. B.; Muchardt, C.; Xia, Y.; Klisak, I.; Mohandas,
T.;
[59504] Sparkes, R. S.; Lusis, A. J.: Localization of the gene for the
DNA-binding
[59505] protein AP-2 to human chromosome 6p22.3-pter. Ge-
nomics 10: 1100-1102,
[59506] 1991.

[59507] 9830.

[59508] Schorle, H.; Meier, P.; Buchert, M.; Jaenisch, R.; Mitchell, P.

[59509] J.: Transcription factor AP-2 essential for cranial closure
and craniofacial

[59510] development. Nature 381: 235-238, 1996.

[59511] 9831.

[59512] Warren, G.; Gordon, M.; Siracusa, L. D.; Buchberg, A. M.; Williams,

[59513] T.: Physical and genetic localization of the gene encoding the AP-2

[59514] transcription factor to mouse chromosome 13. Genomics 31: 234-237,

[59515] 1996.

[59516] 9832.

[59517] Williams, T.; Admon, A.; Luscher, B.; Tjian, R.: Cloning and expression

[59518] of AP-2, a cell-type-specific transcription factor that acti- vates

[59519] inducible enhancer elements. Genes Dev. 2: 1557-1569, 1988.

[59520] 9833.

[59521] Williamson, J. A.; Bosher, J. M.; Skinner, A.; Sheer, D.; Williams,

[59522] T.; Hurst, H. C.: Chromosomal mapping of the human and mouse homologues

[59523] of two new members of the AP-2 family of transcription factors. Genomics 35:
[59524] 262-264, 1996.

[59525] 9834.

[59526] Zhang, J.; Hagopian-Donaldson, S.; Serbedzija, G.; Elsemore, J.;
[59527] Plehn-Dujowich, D.; McMahon, A. P.; Flavell, R. A.; Williams, T.:
[59528] Neural tube, skeletal and body wall defects in mice lacking transcription
[59529] factor AP-2. Nature 381: 238-241, 1996.

[59530] 9835.

[59531] McGrath, J. A.; Duijf, P. H. G.; Doetsch, V.; Irvine, A. D.; de
[59532] Waal, R.; Vanmolkot, K. R. J.; Wessagowit, V.; Kelly, A.; Atherton,
[59533] D. J.; Griffiths, W. A. D.; Orlow, S. J.; van Haeringen, A.; Ausems,
[59534] M. G. E. M.; Yang, A.; McKeon, F.; Bamshad, M. A.; Brunner, H. G.;
[59535] Hamel, B. C. J.; van Bokhoven, H.: Hay-Wells syndrome is caused by

[59536] heterozygous missense mutations in the SAM domain of p63. Hum. Molec.
[59537] Genet. 10: 221-229, 2001.

[59538] 9836.

[59539] Gotoh, T.; Araki, M.; Mori, M.: Chromosomal localization of the
[59540] human arginase II gene and tissue distribution of its mRNA. Biochem.
[59541] Biophys. Res. Commun. 233: 487-491, 1997.

[59542] 9837.

[59543] Gotoh, T.; Sonoki, T.; Nagasaki, A.; Terada, K.; Takiguchi, M.;
[59544] Mori, M.: Molecular cloning of cDNA for nonhepatic mitochondrial
[59545] arginase (arginase II) and comparison of its induction with nitric
[59546] oxide synthase in a murine macrophage-like cell line. FEBS Lett. 395:
[59547] 119-122, 1996.

[59548] 9838.

[59549] Iyer, R.; Jenkinson, C. P.; Vockley, J. G.; Kern, R. M.; Grody,

[59550] W. W.; Cederbaum, S.: The human arginases and arginase deficiency. J.
[59551] Inherit. Metab. Dis. 21 (suppl. 1): 86-100, 1998.

[59552] 9839.

[59553] Morris, S. M., Jr.; Bhamidipati, D.; Kepka-Lenhart, D.: Human
[59554] type II arginase: sequence analysis and tissue-specific expression. Gene 193:
[59555] 157-161, 1997.

[59556] 9840.

[59557] Spector, E. B.; Rice, S. C. H.; Cederbaum, S. D.: Evidence for
[59558] two genes encoding human arginase. (Abstract) Am. J. Hum. Genet. 32:
[59559] 55A only, 1980.

[59560] 9841.

[59561] Spector, E. B.; Rice, S. C. H.; Cederbaum, S. D.: Immunologic
[59562] studies of arginase in tissues of normal human adult and arginase-deficient
[59563] patients. Pediat. Res. 17: 941-944, 1983.

[59564] 9842.

[59565] Vockley, J. G.; Jenkinson, C. P.; Shukla, H.; Kern, R. M.; Grody,

[59566] W. W.; Cederbaum, S. D.: Cloning and characterization of the human

[59567] type II arginase gene. Genomics 38: 118-123, 1996.

[59568] 9843.

[59569] Agarwal, V. R.; Ashanullah, C. I.; Simpson, E. R.; Bulun, S. E.

[59570] : Alternatively spliced transcripts of the aromatase cytochrome P450

[59571] (CYP19) gene in adipose tissue of women. J. Clin. Endocr. Metab. 82:

[59572] 70-74, 1997.

[59573] 9844.

[59574] Nalabolu, S. R.; Shukla, H.; Nallur, G.; Parimoo, S.; Weissman,

[59575] S. M.: Genes in a 220-kb region spanning the TNF cluster in human

[59576] MHC. Genomics 31: 215-222, 1996.

[59577] 9845.

[59578] DeChiara, T. M.; Kimble, R. B.; Poueymirou, W. T.; Rojas, J.; Masiakowski,

[59579] P.; Valenzuela, D. M.; Yancopoulos, G. D.: Ror2, encoding a receptor-like

[59580] tyrosine kinase, is required for cartilage and growth plate development. Nature

[59581] Genet. 24: 271-274, 2000.

[59582] 9846.

[59583] Oldridge, M.; Fortuna, A. M.; Maringa, M.; Propping, P.; Mansour,

[59584] S.; Pollitt, C.; DeChiara, T. M.; Kimble, R. B.; Valenzuela, D. M.;

[59585] Yancopoulos, G. D.; Wilkie, A. O. M.: Dominant mutations in ROR2,

[59586] encoding an orphan receptor tyrosine kinase, cause brachydactyly type

[59587] B. Nature Genet. 24: 375-378, 2000.

[59588] 9847.

[59589] De Windt, L. J.; Lim, H. W.; Bueno, O. F.; Liang, Q.; Delling,

[59590] U.; Braz, J. C.; Glascock, B. J.; Kimball, T. F.; del Monte, F.; Hajjar,

[59591] R. J.; Molkentin, J. D.: Targeted inhibition of calcineurin attenuates
[59592] cardiac hypertrophy in vivo. Proc. Nat. Acad. Sci. 98: 3322-3327,
[59593] 2001.

[59594] 9848.

[59595] Fuentes, J. J.; Genesca, L.; Kingsbury, T. J.; Cunningham, K. W.;
[59596] Perez-Riba, M.; Estivill, X.; de la Luna, S.: DSCR1, overex-pressed
[59597] in Down syndrome, is an inhibitor of calcineurin-mediated signaling
[59598] pathways. Hum. Molec. Genet. 9: 1681-1690, 2000.

[59599] 9849.

[59600] Giri, P.; Higuchi, S.; Kincaid, R. L.: Chromosomal mapping of
[59601] the human genes for the calmodulin-dependent protein phosphatase (calcineurin)
[59602] catalytic subunit. Biochem. Biophys. Res. Commun. 181: 252-258,
[59603] 1991.

[59604] 9850.

[59605] Guerini, D.; Klee, C. B.: Cloning of human calcineurin A: evidence

[59606] for two isozymes and identification of a polyproline structural domain. Proc.

[59607] Nat. Acad. Sci. 86: 9183-9187, 1989.

[59608] 9851.

[59609] Leinwand, L. A.: Calcineurin inhibition and cardiac hypertrophy:

[59610] a matter of balance. Proc. Nat. Acad. Sci. 98: 2947-2949, 2001.

[59611] 9852.

[59612] Malleret, G.; Haditsch, U.; Genoux, D.; Jones, M. W.; Bliss, T.

[59613] V. P.; Vanhoose, A. M.; Weitlauf, C.; Kandel, E. R.; Winder, D. G.;

[59614] Mansuy, I. M.: Inducible and reversible enhancement of learning,

[59615] memory, and long-term potentiation by genetic inhibition of calcineurin. Cell 104:

[59616] 675-686, 2001.

[59617] 9853.

[59618] Mansuy, I. M.; Mayford, M.; Jacob, B.; Kandel, E. R.; Bach, M.

[59619] E.: Restricted and regulated overexpression reveals cal-cineurin as

[59620] a key component in the transition from short-term to long-term memory. Cell 92:

[59621] 39-49, 1998.

[59622] 9854.

[59623] Rothermel, B. A.; McKinsey, T. A.; Vega, R. B.; Nicol, R. L.; Mammen,

[59624] P.; Yang, J.; Antos, C. L.; Shelton, J. M.; Bassel-Duby, R.; Olson,

[59625] E. N.; Williams, R. S.: Myocyte-enriched calcineurin-inter-acting

[59626] protein, MCIP1, inhibits cardiac hypertrophy in vivo. Proc. Nat.

[59627] Acad. Sci. 98: 3328-3333, 2001.

[59628] 9855.

[59629] Seitz, D. P.; Pasha, M. K.; Singh, B.; Chu, A.; Sharma, R. K.

[59630] : Localization and characterization of calcineurin in bovine

eye. Invest.
[59631] Ophthal. Vis. Sci. 43: 15–21, 2002.

[59632] 9856.

[59633] Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.

[59634] : Calcineurin A alpha (PPP3CA), calcineurin A beta
(PPP3CB) and calcineurin
[59635] B (PPP3R1) are located on human chromosomes 4,
10q21–q22 and 2p16–p15
[59636] respectively. Cytogenet. Cell Genet. 72: 236–241, 1996.

[59637] 9857.

[59638] Winder, D. G.; Mansuy, I. M.; Osman, M.; Moallem, T. M.;
Kandel,
[59639] E. R.: Genetic and pharmacological evidence for a novel,
intermediate
[59640] phase of long-term potentiation suppressed by cal-
cineurin. Cell 92:
[59641] 25–37, 1998.

[59642] 9858.

[59643] Bueno, O. F.; Wilkins, B. J.; Tymitz, K. M.; Glascock, B. J.;
Kimball,
[59644] T. F.; Lorenz, J. N.; Molkentin, J. D.: Impaired cardiac hy-

pertrophic
[59645] response in calcineurin A-beta-deficient mice. Proc. Nat. Acad. Sci. 99:
[59646] 4586-4591, 2002.

[59647] 9859.

[59648] Guerini, D.; Krinks, M. H.; Sikela, J. M.; Hahn, W. E.; Klee, C.
[59649] B.: Isolation and sequence of a cDNA clone for human cal-cineurin
[59650] B, the Ca(2+)-binding subunit of the Ca(2+)/calmodulin-stimulated
[59651] protein phosphatase. DNA 8: 675-682, 1989.

[59652] 9860.

[59653] Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.:
[59654] Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB) and calcineurin
[59655] B (PPP3R1) are located on human chromosomes 4, 10q21-q22 and 2p16-p15
[59656] respectively. Cytogenet. Cell Genet. 72: 236-241, 1996.

[59657] 9861.

[59658] Ferrari, S.; Calabretta, B.; deRiel, J. K.; Battini, R.; Ghezzo,

[59659] F.; Lauret, E.; Griffin, C.; Emanuel, B. S.; Gurrieri, F.; Baserga,

[59660] R.: Structural and functional analysis of a growth-regulated gene,

[59661] the human calcyclin. J. Biol. Chem. 262: 8325-8332, 1987.

[59662] 9862.

[59663] Alevizaki, M.; Stevenson, J. C.; Girgis, S. I.; MacIntyre, I.;

[59664] Legon, S.: Altered calcitonin gene in a young patient with osteoporosis. Brit.

[59665] Med. J. 298: 1215-1216, 1989.

[59666] 9863.

[59667] Amara, S. G.; Jonas, V.; Rosenfeld, M. G.; Ong, E. S.; Evans, R.

[59668] M.: Alternative RNA processing in calcitonin gene expression generates

[59669] mRNAs encoding different polypeptide products. Nature 298: 240-244,

[59670] 1982.

[59671] 9864.

[59672] Aiyar, N.; Rand, K.; Elshourbagy, N. A.; Zeng, Z.; Adamou,

J. E.;

[59673] Bergsma, D. J.; Li, Y.: A cDNA encoding the calcitonin gene-related

[59674] peptide type 1 receptor. J. Biol. Chem. 271: 11325-11329, 1996.

[59675] 9865.

[59676] Fluhmann, B.; Muff, R.; Hunziker, W.; Fischer, J. A.; Born, W.

[59677] : A human orphan calcitonin receptor-like structure. Biochem. Biophys.

[59678] Res. Commun. 206: 341-347, 1995.

[59679] 9866.

[59680] Foord, S. M.; Craig, R. K.: Isolation and characterisation of

[59681] a human calcitonin-gene-related-peptide receptor. Europ. J. Biochem. 170:

[59682] 373-379, 1987.

[59683] 9867.

[59684] Foord, S. M.; Marshall, F. H.: RAMPs: accessory proteins for seven

[59685] transmembrane domain receptors. Trends Pharm. Sci. 20: 184-187,

[59686] 1999.

[59687] 9868.

[59688] Kamitani, S.; Asakawa, M.; Shimekake, Y.; Kuwasako, K.; Nakahara,

[59689] K.; Sakata, T.: The RAMP2/CRLR complex is a functional adrenomedullin

[59690] receptor in human endothelial and vascular smooth muscle cells. FEBS

[59691] Lett. 448: 111-114, 1999.

[59692] 9869.

[59693] Nakazawa, I.; Nakajima, T.; Harada, H.; Ishigami, T.; Umemura,

[59694] S.; Emi, M.: Human calcitonin receptor-like receptor for adrenomedullin:

[59695] genomic structure, eight single-nucleotide polymorphisms, and haplotype

[59696] analysis. J. Hum. Genet. 46: 132-136, 2001.

[59697] 9870.

[59698] Dolmetsch, R. E.; Pajvani, U.; Fife, K.; Spotts, J. M.; Greenberg,

[59699] M. E.: Signaling to the nucleus by an L-type calcium chan-

nel-calmodulin
[59700] complex through the MAP kinase pathway. Science 294: 333-339, 2001.
[59701] 9871.
[59702] McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,
[59703] Canada 2/14/1992.
[59704] 9872.
[59705] Powers, P. A.; Gregg, R. G.; Hogan, K.: Linkage mapping of the
[59706] human gene for the alpha-1 subunit of the cardiac DHP-sensitive Ca(2+)
[59707] channel (CACNL1A1) to chromosome 12p13.2-pter using a dinucleotide
[59708] repeat. Genomics 14: 206-207, 1992.
[59709] 9873.
[59710] Powers, P. A.; Gregg, R. G.; Lalley, P. A.; Liao, M.; Hogan, K.
[59711] : Assignment of the human gene for the alpha-1 subunit of the cardiac
[59712] DHP-sensitive Ca(2+) channel (CCHL1A1) to chromosome

12p12-pter. Genomics 10:
[59713] 835-839, 1991.

[59714] 9874.

[59715] Sartor, H.; Ehlert, F.; Grzeschik, K.-H.; Muller, R.; Adolph, S.
[59716] : Assignment of two human cell cycle genes, CDC25C and CCNB1, to 5q31
[59717] and 5q12, respectively. Genomics 13: 911-912, 1992.

[59718] 9875.

[59719] Geng, Y.; Whoriskey, W.; Park, M. Y.; Bronson, R. T.; Medema, R.
[59720] H.; Li, T.; Weinberg, R. A.; Sicinski, P.: Rescue of cyclin D1 deficiency
[59721] by knockin cyclin E. Cell 97: 767-777, 1999.

[59722] 9876.

[59723] Moberg, K. H.; Bell, D. W.; Wahrer, D. C. R.; Haber, D. A.; Hariharan,
[59724] I. K.: Archipelago regulates cyclin E levels in Drosophila and is
[59725] mutated in human cancer cell lines. Nature 413: 311-316, 2001.

[59726] 9877.

[59727] Sheaff, R. J.; Groudine, M.; Gordon, M.; Roberts, J. M.; Clurman,

[59728] B. E.: Cyclin E-CDK2 is a regulator of p27(Kip1). Genes Dev. 11:

[59729] 1464-1478, 1997.

[59730] 9878.

[59731] Akoulitchev, S.; Chuikov, S.; Reinberg, D.: TFIIH is nega-tively

[59732] regulated by cdk8-containing mediator complexes. Na-ture 407: 102-106,

[59733] 2000.

[59734] 9879.

[59735] Hanson, R. D.; Hohn, P. A.; Popescu, N. C.; Ley, T. J.: A cluster

[59736] of hematopoietic serine protease genes is found on the same chromosomal

[59737] band as the human alpha/delta T-cell receptor locus. Proc. Nat. Acad.

[59738] Sci. 87: 960-963, 1990.

[59739] 9880.

[59740] Harper, K.; Mattei, M.-G.; Simon, D.; Suzan, M.; Guenet, J.-L.;
[59741] Haddad, P.; Sasportes, M.; Golstein, P.: Proximity of the CTLA-1
[59742] serine esterase and Tcr(alpha) loci in mouse and man. Im-
munogenetics 28:
[59743] 439-444, 1988.

[59744] 9881.

[59745] Klein, J. L.; Shows, T. B.; Dupont, B.; Trapani, J. A.: Ge-
nomic
[59746] organization and chromosomal assignment for a serine protease gene
[59747] (CSP-B), expressed by human cytotoxic lymphocytes. Ge-
nomics 5: 110-117,
[59748] 1989.

[59749] 9882.

[59750] Lin, C. C.; Meier, M.; Sorensen, O.; Sasi, R.; Tainaka, T.; Bleackley,
[59751] R. C.: Chromosome localization of two human serine pro-
tease genes
[59752] to region 14q11.2-q12 by in situ hybridization. Cyto-
genet. Cell Genet. 53:

[59753] 169-171, 1990.

[59754] 9883.

[59755] Motyka, B.; Korbutt, G.; Pinkoski, M. J.; Heibein, J. A.; Ca-
puto,
[59756] A.; Hobman, M.; Barry, M.; Shostak, I.; Sawchuk, T.;
Holmes, C. F.
[59757] B.; Gauldie, J.; Bleackley, R. C.: Mannose
6-phosphate/insulin-like
[59758] growth factor II receptor is a death receptor for granzyme
B during
[59759] cytotoxic T cell-induced apoptosis. Cell 103: 491-500,
2000.
[59760] 9884.

[59761] Miles, J. S.; Spurr, N. K.; Gough, A. C.; Jowett, T.; McLaren,

[59762] A. W.; Brook, J. D.; Wolf, C. R.: A novel human cytochrome
P450 gene
[59763] (P450IIB): chromosomal localization and evidence for al-
ternative splicing. Nucleic
[59764] Acids Res. 16: 5783-5795, 1988.

[59765] 9885.

[59766] Santisteban, I.; Povey, S.; Shephard, E. A.; Phillips, I. R.:

[59767] The major phenobarbital-inducible cytochrome P-450 gene subfamily
[59768] (P450IIB) mapped to the long arm of human chromosome 19. Ann. Hum.
[59769] Genet. 52: 129-135, 1988.

[59770] 9886.

[59771] Trask, B.; Fertitta, A.; Christensen, M.; Youngblom, J.; Bergmann,
[59772] A.; Copeland, A.; de Jong, P.; Mohrenweiser, H.; Olsen, A.; Carrano,
[59773] A.; Tynan, K.: Fluorescence in situ hybridization mapping of human
[59774] chromosome 19: cytogenetic band location of 540 cosmids and 70 genes
[59775] or DNA markers. Genomics 15: 133-145, 1993.

[59776] 9887.

[59777] Barletta, C.; Batticane, N.; Ragusa, R. M.; Leube, R.; Peschle,
[59778] C.; Romano, V.: Subchromosomal localization of two human cytokeratin
[59779] genes (KRT4 and KRT15) by in situ hybridization. Cytogenet. Cell

[59780] Genet. 54: 148-150, 1990.

[59781] 9888.

[59782] Barletta, C.; Batticane, N.; Ragusa, R. M.; Leube, R. E.; Franke,

[59783] W. W.; Peschle, C.; Romano, V.: Sub-chromosomal local-

ization of human

[59784] cytokeratin 4, 15, and 19 genes. (Abstract) Cytogenet. Cell Genet. 51:

[59785] 958 only, 1989.

[59786] 9889.

[59787] Fuchs, E.: Keratins as biochemical markers of epithelial differentiation. Trends

[59788] Genet. 4: 277-281, 1988.

[59789] 9890.

[59790] Mischke, D.; Wild, G.: Polymorphic keratins in human epi-

dermis. J.

[59791] Invest. Derm. 88: 191-197, 1987.

[59792] 9891.

[59793] Mischke, D.; Wille, G.; Wild, A. G.: Allele frequencies and segregation

[59794] of human polymorphic keratins K4 and K5. Am. J. Hum. Genet. 46:
[59795] 548-552, 1990.

[59796] 9892.

[59797] Romano, V.; Bosco, P.; Costa, G.; Leube, R.; Franke, W. W.; Rocchi,
[59798] M.; Romeo, G.: Chromosomal assignment of cytokeratin genes. (Abstract) Cytogenet.
[59799] Cell Genet. 46: 683 only, 1987.

[59800] 9893.

[59801] Romano, V.; Bosco, P.; Rocchi, M.; Costa, G.; Leube, R. E.; Franke,
[59802] W. W.; Romeo, G.: Chromosomal assignments of human type I and type
[59803] II cytokeratin genes to different chromosomes. Cytogenet. Cell Genet. 48:
[59804] 148-151, 1988.

[59805] 9894.

[59806] Rugg, E. L.; McLean, W. H. I.; Allison, W. E.; Lunny, D. P.; Lane,
[59807] E. B.; Macleod, R. I.; Felix, D. H.; Munro, C. S.: White

sponge nevus
[59808] (WSN) is caused by mutations in mucosal keratin K4. (Abstract) Am.
[59809] J. Hum. Genet. 57: A47 only, 1995.

[59810] 9895.

[59811] Rugg, E. L.; McLean, W. H. I.; Allison, W. E.; Lunny, D. P.; Macleod,
[59812] R. I.; Felix, D. H.; Lane, E. B.; Munro, C. S.: A mutation in the
[59813] mucosal keratin K4 is associated with oral white sponge nevus. Nature
[59814] Genet. 11: 450-452, 1995.

[59815] 9896.

[59816] Terrinoni, A.; Candi, E.; Oddi, S.; Gobello, T.; Camaione, D.
[59817] B.; Mazzanti, C.; Zambruno, G.; Knight, R.; Melino, G.: A glutamine
[59818] insertion in the 1A alpha helical domain of the keratin 4 gene in
[59819] a familial case of white sponge nevus. J. Invest. Derm. 114: 388-391,
[59820] 2000.

[59821] 9897.

[59822] Schrijver, H. M.; Crusius, J. B. A.; Uitdehaag, B. M. J.; Gar-
cia
[59823] Gonzalez, M. A.; Kostense, P. J.; Polman, C. H.; Pena, A. S.:
Association
[59824] of interleukin-1-beta and interleukin-1 receptor antago-
nist genes
[59825] with disease severity in MS. Neurology 52: 595-599,
1999.

[59826] 9898.

[59827] Terasaki, P. I.; Park, M. S.; Opelz, G.; Ting, A.: Multiple
sclerosis
[59828] and high incidence of a B-lymphocyte antigen. Science
193: 1245-1247,
[59829] 1976.

[59830] 9899.

[59831] Reardon, W.; Wilkes, D.; Rutland, P.; Pulleyn, L. J.; Mal-
colm,
[59832] S.; Dean, J. C. S.; Evans, R. D.; Jones, B. M.; Hayward, R.;
Hall,
[59833] C. M.; Nevin, N. C.; Baraitser, M.; Winter, R. M.: Craniosyn-
ostosis

[59834] associated with FGFR3 pro250-to-arg mutation results in a range of
[59835] clinical presentations including unisutural sporadic craniosynostosis. J.
[59836] Med. Genet. 34: 632-636, 1997.

[59837] 9900.

[59838] Reddy, B. S. N.; Garg, B. R.; Padiyar, N. V.; Krishnaram, A. S.
[59839] : An unusual association of acanthosis nigricans and Crouzon's disease:
[59840] a case report. J. Derm. 12: 85-90, 1985.

[59841] 9901.

[59842] Robin, N. H.; Scott, J. A.; Cohen, A. R.; Goldstein, J. A.: Nonpenetrance
[59843] in FGFR3-associated coronal synostosis syndrome. (Letter) Am. J.
[59844] Med. Genet. 80: 296-297, 1998.

[59845] 9902.

[59846] Roscioli, T.; Flanagan, S.; Mortimore, R. J.; Kumar, P.; Weedon,
[59847] D.; Masel, J.; Lewandowski, R.; Hyland, V.; Glass, I. A.: Pre-

mature
[59848] calvarial synostosis and epidermal hyperplasia
(Beare-Stevenson syndrome-like
[59849] anomalies) resulting from a P250R missense mutation in
the gene encoding
[59850] fibroblast growth factor receptor 3. Am. J. Med. Genet.
101: 187-194,
[59851] 2001.

[59852] 9903.

[59853] Rousseau, F.; Bonaventure, J.; Legeai-Mallet, L.; Pelet, A.;
Rozet,
[59854] J.-M.; Maroteaux, P.; Le Merrer, M.; Munnich, A.: Muta-
tions in the
[59855] gene encoding fibroblast growth factor receptor-3 in
achondroplasia. Nature 371:
[59856] 252-254, 1994.

[59857] 9904.

[59858] Rousseau, F.; El Ghouzzi, V.; Delezoide, A. L.; Legeai-Mal-
let,
[59859] L.; Le Merrer, M.; Munnich, A.; Bonaventure, J.: Missense
FGFR3 mutations
[59860] create cysteine residues in thanatophoric dwarfism type I

(TD1). Hum.
[59861] Molec. Genet. 5: 509-512, 1996.

[59862] 9905.

[59863] Rousseau, F.; Saugier, P.; Le Merrer, M.; Munnich, A.;
Delezoide,
[59864] A.-L.; Maroteaux, P.; Bonaventure, J.; Narcy, F.; Sanak, M.:
Stop
[59865] codon FGFR3 mutations in thanatophoric dwarfism type 1.
Nature Genet. 10:
[59866] 11-12, 1995.

[59867] 9906.

[59868] Sawai, H.; Komori, S.; Ida, A.; Henmi, T.; Bessho, T.;
Koyama,
[59869] K.: Prenatal diagnosis of thanatophoric dysplasia by muta-
tional analysis
[59870] of the fibroblast growth factor receptor 3 gene and a pro-
posed correction
[59871] of previously published PCR results. Prenatal Diag. 19:
21-24, 1999.
[59872] 9907.

[59873] Schweitzer, D. N.; Graham, J. M., Jr.; Lachman, R. S.; Jabs,

E.

[59874] W.; Okajima, K.; Przylepa, K. A.; Shanske, A.; Chen, K.; Neidich,

[59875] J. A.; Wilcox, W. R.: Subtle radiographic findings of achondroplasia

[59876] in patients with Crouzon syndrome with acanthosis nigricans due to

[59877] an ala391-to-glu substitution in FGFR3. Am. J. Med. Genet. 98: 75-91,

[59878] 2001.

[59879] 9908.

[59880] Scotet, E.; Houssaint, E.: The choice between alternative IIIb

[59881] and IIIc exons of the FGFR-3 gene is not strictly tissue-specific. Biochim.

[59882] Biophys. Acta 1264: 238-242, 1995.

[59883] 9909.

[59884] Shiang, R.; Thompson, L. M.; Zhu, Y.-Z.; Church, D. M.; Fielder,

[59885] T. J.; Bocian, M.; Winokur, S. T.; Wasmuth, J. J.: Mutations in the

[59886] transmembrane domain of FGFR3 cause the most common

genetic form of
[59887] dwarfism, achondroplasia. Cell 78: 335-342, 1994.

[59888] 9910.

[59889] Shimizu, A.; Tada, K.; Shukunami, C.; Hiraki, Y.; Kurokawa, T.;
[59890] Magane, N.; Kurokawa-Seo, M.: A novel alternatively spliced fibroblast
[59891] growth factor receptor 3 isoform lacking the acid box domain is expressed
[59892] during chondrogenic differentiation of ATDC5 cells. J. Biol. Chem. 276:
[59893] 11031-11040, 2001.

[59894] 9911.

[59895] Shimizu, A.; Takashima, Y.; Kurokawa-Seo, M.: FGFR3 isoforms
[59896] have distinct functions in the regulation of growth and cell morphology. Biochem.
[59897] Biophys. Res. Commun. 290: 113-120, 2002.

[59898] 9912.

[59899] Sibley, K.; Cuthbert-Heavens, D.; Knowles, M. A.: Loss of heterozygosity

[59900] at 4p16.3 and mutation of FGFR3 in transitional cell carcinoma. Oncogene 20:
[59901] 686-691, 2001.

[59902] 9913.

[59903] Spranger, J.: Bone dysplasia families. Path. Immunopath. Res. 7:
[59904] 76-80, 1988.

[59905] 9914.

[59906] Su, W.-C. S.; Kitagawa, M.; Xue, N.; Xie, B.; Garofalo, S.; Cho,
[59907] J.; Deng, C.; Horton, W. A.; Fu, X.-Y.: Activation of Stat1 by mutant
[59908] fibroblast growth-factor receptor in thanatophoric dysplasia type
[59909] II dwarfism. Nature 386: 288-292, 1997.

[59910] 9915.

[59911] Superti-Furga, A.; Eich, G.; Bucher, H. U.; Wisser, J.; Giedion,
[59912] A.; Gitzelmann, R.; Steinmann, B.: A glycine
375-to-cysteine substitution
[59913] in the transmembrane domain of the fibroblast growth

factor receptor-3
[59914] in a newborn with achondroplasia. Europ. J. Pediat. 154: 215-219,
[59915] 1995.

[59916] 9916.

[59917] Superti-Furga, A.; Locher, M. L.; Steinlin, M.; Elch, G.; Huisman,
[59918] T.; Steinmann, B.; Sailer, H. F.; Boltshauser, E.: Crouzon syndrome
[59919] with acanthosis nigricans, spinal stenosis and desmo-osteoblastomas:
[59920] pleiotropic effects of the FGFR-3 ala391glu mutation. J. Craniomaxillofac.
[59921] Surg. (Suppl.) 24: 112 only, 1996.

[59922] 9917.

[59923] Szabo, J.; Bellus, G. A.; Kaitila, I.; Francomano, C. A.: Fi-broblast
[59924] growth factor receptor 3 (FGFR3) mutations in sporadic cases of achondroplasia
[59925] occur exclusively on the paternally derived chromosome. Am. J. Hum.
[59926] Genet.(Abstract) 59 (suppl.): A287 only, 1996.

[59927] 9918.

[59928] Tavormina, P. L.; Bellus, G. A.; Webster, M. K.; Bamshad, M. J.;

[59929] Fraley, A. E.; McIntosh, I.; Szabo, J.; Jiang, W.; Jabs, E. W.; Wilcox,

[59930] W. R.; Wasmuth, J. J.; Donoghue, D. J.; Thompson, L. M.; Francomano,

[59931] C. A.: A novel skeletal dysplasia with developmental delay and acanthosis

[59932] nigricans is caused by a lys650-to-met mutation in the fi-broblast

[59933] growth factor receptor 3 gene. Am. J. Hum. Genet. 64: 722-731, 1999.

[59934] 9919.

[59935] Tavormina, P. L.; Rimoin, D. L.; Cohn, D. H.; Zhu, Y.-Z.; Shiang,

[59936] R.; Wasmuth, J. J.: Another mutation that results in the substitution

[59937] of an unpaired cysteine residue in the extracellular do-main of FGFR3

[59938] in thanatophoric dysplasia type I. Hum. Molec. Genet. 4: 2175-2177,

[59939] 1995.

[59940] 9920.

[59941] Tavormina, P. L.; Shiang, R.; Thompson, L. M.; Zhu, Y.-Z.; Wilkin,

[59942] D. J.; Lachman, R. S.; Wilcox, W. R.; Rimoin, D. L.; Cohn, D. H.;

[59943] Wasmuth, J. J.: Thanatophoric dysplasia (types I and II) caused by

[59944] distinct mutations in fibroblast growth factor receptor 3. Nature

[59945] Genet. 9: 321-328, 1995.

[59946] 9921.

[59947] Thompson, L. M.; Plummer, S.; Schalling, M.; Altherr, M. R.; Gusella,

[59948] J. F.; Housman, D. E.; Wasmuth, J. J.: A gene encoding a fi-broblast

[59949] growth factor receptor isolated from the Huntington dis-ease gene region

[59950] of human chromosome 4. Genomics 11: 1133-1142, 1991.

[59951] 9922.

[59952] Tsai, F.-J.; Tsai, C.-H.; Chang, J.-G.; Wu, J.-Y.: Mutations

[59953] in the fibroblast growth factor receptor 3 (FGFR3) cause achondroplasia,

[59954] hypochondroplasia, and thanatophoric dysplasia: Taiwanese data. (Letter) Am.

[59955] J. Med. Genet. 86: 300-301, 1999.

[59956] 9923.

[59957] Tsai, F.-J.; Wu, J.-Y.; Tsai, C.-H.; Chang, J.-G.: Identification

[59958] of a common N540K mutation in 8/18 Taiwanese hypochondroplasia patients:

[59959] further evidence for genetic heterogeneity. (Letter) Clin. Genet. 55:

[59960] 279-280, 1999.

[59961] 9924.

[59962] von Gernet, S.; Schuffenhauer, S.; Golla, A.; Lichtner, P.; Balg,

[59963] S.; Muhlbauer, W.; Murken, J.; Fairley, J.; Meitinger, T.: Craniosynostosis

[59964] suggestive of Saethre-Chotzen syndrome: clinical description of a

[59965] large kindred and exclusion of candidate regions on 7p. Am. J. Med.

[59966] Genet. 63: 177-184, 1996.

[59967] 9925.

[59968] Frydman, M.; Bonne-Tamir, B.; Farrer, L. A.; Conneally, P. M.;

[59969] Magazanik, A.; Ashbel, S.; Goldwitch, Z.: Assignment of the gene

[59970] for Wilson disease to chromosome 13: linkage to the esterase D locus. Proc.

[59971] Nat. Acad. Sci. 82: 1819-1821, 1985.

[59972] 9926.

[59973] Scheuner, D.; Eckman, C.; Jensen, M.; Song, X.; Citron, M.; Suzuki,

[59974] N.; Bird, T. D.; Hardy, J.; Hutton, M.; Kukull, W.; Larson, E.; Levy-Lahad,

[59975] E.: Secreted amyloid beta-protein similar to that in the senile plaques

[59976] of Alzheimer's disease is increased in vivo by the presenilin 1 and

[59977] 2 and APP mutations linked to familial Alzheimer's disease. Nature

[59978] Med. 2: 864-870, 1996.

[59979] 9927.

[59980] Sherrington, R.; Rogaev, E. I.; Liang, Y.; Rogaeva, E. A.; Levesque,

[59981] G.; Ikeda, M.; Chi, H.; Lin, C.; Li, G.; Holman, K.; Tsuda, T.; Mar,

[59982] L.; Foncin, J.-F.; Bruni, A. C.; Montesi, M. P.; Sorbi, S.; Rainero,

[59983] I.; Pinessi, L.; Nee, L.; Chumakov, I.; Pollen, D.; Brookes, A.; Sanseau,

[59984] P.; Polinsky, R. J.; Wasco, W.; Da Silva, H. A. R.; Haines, J. L.;

[59985] Pericak-Vance, M. A.; Tanzi, R. E.; Roses, A. D.; Fraser, P. E.; Rommens,

[59986] J. M.; St George-Hyslop, P. H.: Cloning of a gene bearing mis-sense

[59987] mutations in early-onset familial Alzheimer's disease. Nature 375:

[59988] 754-760, 1995.

[59989] 9928.

[59990] St. George-Hyslop, P.; Haines, J.; Rogaev, E.; Mortilla, M.; Vaula,

[59991] G.; Pericak-Vance, M.; Foncin, J.-F.; Montesi, M.; Bruni, A.; Sorbi,

[59992] S.; Rainero, I.; Pinessi, L.; Pollen, D.; Polinsky, R.; Nee, L.; Kennedy,

[59993] J.; Macciardi, F.; Rogaeva, E.; Liang, Y.; Alexandrova, N.; Lukiw,

[59994] W.; Schlumpf, K.; Tanzi, R.; Tsuda, T.; Farrer, L.; Cantu, J.-M.;

[59995] Duara, R.; Amaducci, L.; Bergamini, L.; Gusella, J.; Roses, A.; Crapper

[59996] McLachlan, D.: Genetic evidence for a novel familial Alzheimer's

[59997] disease locus on chromosome 14. Nature Genet. 2: 330-334, 1992.

[59998] 9929.

[59999] St. George-Hyslop, P. H.; Haines, J. L.; Farrer, L. A.; Polinsky,

[60000] R.; Van Broeckhoven, C.; Goate, A.; Crapper McLachlan, D. R.; Orr,

[60001] H.; Bruni, A. C.; Sorbi, S.; Rainero, I.; Foncin, J.-F.; Pollen, D.;

[60002] Cantu, J. M.; Tupler, R.; Voskresenskaya, N.; Mayeux, R.; Growdon,

[60003] J.; Fried, V. A.; Myers, R. H.; Nee, L.; Backhovens, H.; Martin, J.

[60004] J.; Rossor, M.; Owen, M. J.; Mullan, M.; Percy, M. E.; Karlinsky,

[60005] H.; Rich, S.; Heston, L.; Montesi, M.; Mortilla, M.; Nacmias, N.;

[60006] Gusella, J. F.; Hardy, J. A.; other members of the FAD Collaborative

[60007] Study Group: Genetic linkage studies suggest that Alzheimer's disease

[60008] is not a single homogeneous disorder. Nature 347: 194-197, 1990.

[60009] 9930.

[60010] Moehlmann, T.; Winkler, E.; Xia, X.; Edbauer, D.; Murrell, J.;

[60011] Capell, A.; Kaether, C.; Zheng, H.; Ghetti, B.; Haass, C.; Steiner,

[60012] H.: Presenilin-1 mutations of leucine 166 equally affect the generation

[60013] of the Notch and APP intracellular domains independent of their effect

[60014] on A-beta(42) production. Proc. Nat. Acad. Sci. 99: 8025-8030, 2002.

[60015] 9931.

[60016] Morelli, L.; Prat, M. I.; Levy, E.; Mangone, C. A.; Castano, E.

[60017] M.: Presenilin 1 met146leu variant due to an A-T transversion in

[60018] an early-onset familial Alzheimer's disease pedigree from Argentina. Clin.

[60019] Genet. 53: 469-473, 1998.

[60020] 9932.

[60021] Morgan, D.; Diamond, D. M.; Gottschall, P. E.; Ugen, K. E.; Dickey,

[60022] C.; Hardy, J.; Duff, K.; Jantzen, P.; DiCarlo, G.; Wilcock, D.; Connor,

[60023] K.; Hatcher, J.; Hope, C.; Gordon, M.; Arendash, G. W.: A-beta peptide

[60024] vaccination prevents memory loss in an animal model of Alzheimer's

[60025] disease. Nature 408: 982-985, 2000. Note: Erratum Na-ture 412:

[60026] 660 only, 2001.

[60027] 9933.

[60028] Mullan, M.; Houlden, H.; Windelspecht, M.; Fidani, L.;

Lombardi,
[60029] C.; Diaz, P.; Rossor, M.; Crook, R.; Hardy, J.; Duff, K.; Crawford,
[60030] F.: A locus for familial early-onset Alzheimer's disease on the long
[60031] arm of chromosome 14, proximal to the alpha-1-antichymotrypsin gene. Nature
[60032] Genet. 2: 340-342, 1992.

[60033] 9934.

[60034] Nechiporuk, A.; Fain, P.; Kort, E.; Nee, L. E.; Frommelt, E.;

[60035] Polinsky, R. J.; Korenberg, J. R.; Pulst, S.-M.: Linkage of fa-milial
[60036] Alzheimer disease to chromosome 14 in two large early-onset pedigrees:
[60037] effects of marker allele frequencies on lod scores. Am. J. Med. Genet. 48:
[60038] 63-66, 1993.

[60039] 9935.

[60040] Page, K.; Hollister, R.; Tanzi, R. E.; Hyman, B. T.: In situ

[60041] hybridization analysis of presenilin 1 mRNA in Alzheimer disease and

[60042] in lesioned rat brain. Proc. Nat. Acad. Sci. 93: 14020-14024, 1996.

[60043] 9936.

[60044] Perez-Tur, J.; Froelich, S.; Prihar, G.; Crook, R.; Baker, M.;

[60045] Duff, K.; Wragg, M.; Busfield, F.; Lendon, C.; Clark, R. F.; Roques,

[60046] P.; Fuldner, R. A.; Johnston, J.; Cowburn, R.; Forsell, C.; Axelman,

[60047] K.; Lilius, L.; Houlden, H.; Karran, E.; Roberts, G. W.; Rossor, M.;

[60048] Adams, M. D.; Hardy, J.; Goate, A.; Lannfelt, L.; Hutton, M.: A mutation

[60049] in Alzheimer's disease destroying a splice acceptor site in the presenilin-1

[60050] gene. Neuroreport 7: 297-301, 1995.

[60051] 9937.

[60052] Poorkaj, P.; Sharma, V.; Anderson, L.; Nemens, E.; Alonso, M.

[60053] E.; Orr, H.; White, J.; Heston, L.; Bird, T. D.; Schellenberg, G.

[60054] D.: Missense mutations in the chromosome 14 familial Alzheimer's

[60055] disease presenilin 1 gene. Hum. Mutat. 11: 216-221, 1998.

[60056] 9938.

[60057] Qian, S.; Jiang, P.; Guan, X.-M.; Singh, G.; Trumbauer, M. E.;

[60058] Yu, H.; Chen, H. Y.; Van der Ploeg, L. H. T.; Zheng, H.: Mu-tant human

[60059] presenilin 1 protects presenilin 1 null mouse against em-bryonic lethality

[60060] and elevates A-beta-1-42/43 expression. Neuron 20: 611-617, 1998.

[60061] 9939.

[60062] Raux, G.; Gantier, R.; Thomas-Anterion, C.; Boulliat, J.; Verpillat,

[60063] P.; Hannequin, D.; Brice, A.; Frebourg, T.; Campion, D.: Dementia

[60064] with prominent frontotemporal features associated with L113P presenilin

[60065] 1 mutation. Neurology 55: 1577-1578, 2000.

[60066] 9940.

[60067] Reznik-Wolf, H; Treves, T. A.; Davidson, M.; Aharon-Peretz, J.;

[60068] St. George Hyslop, P. H.; Chapman, J.; Korczyn, A. D.; Goldman, B.;

[60069] Friedman, E.: A novel mutation of presenilin 1 in familial Alzheimer's

[60070] disease in Israel detected by denaturing gradient gel electrophoresis. Hum.

[60071] Genet. 98: 700-702, 1996.

[60072] 9941.

[60073] Rozen, R.; Barton, D.; Du, J.; Hum, D. W.; MacKenzie, R. E.; Francke,

[60074] U.: Chromosomal localization of the gene for the human trifunctional

[60075] enzyme, methylenetetrahydrofolate dehydrogenase-methenyltetrahydrofolate

[60076] cyclohydrolase-formyltetrahydrofolate synthetase. Am. J. Hum. Genet. 44:

[60077] 781-786, 1989.

[60078] 9942.

[60079] Russo, C.; Schettini, G.; Saido, T. C.; Hulette, C.; Lippa, C.;

[60080] Lannfelt, L.; Ghetti, B.; Gambetti, P.; Tabaton, M.; Teller, J. K.

[60081] : Presenilin-1 mutations in Alzheimer's disease. Nature 405: 531-532,

[60082] 2000.

[60083] 9943.

[60084] Schellenberg, G. D.; Bird, T. D.; Wijsman, E. M.; Orr, H. T.;

[60085] Anderson, L.; Nemens, E.; White, J. A.; Bonnycastle, L.; Weber, J.

[60086] L.; Alonso, M. E.; Potter, H.; Heston, L. L.; Martin, G. M.: Genetic

[60087] linkage evidence for a familial Alzheimer's disease locus on chromosome

[60088] 14. Science 258: 668-671, 1992.

[60089] 9944.

[60090] Rogaev, E. I.; Sherrington, R.; Wu, C.; Levesque, G.; Liang, Y.;

[60091] Rogaeva, E. A.; Ikeda, M.; Holman, K.; Lin, C.; Lukiw, W. J.; de Jong,

[60092] P. J.; Fraser, P. E.; Rommens, J. M.; St. George-Hyslop, P.: Analysis

[60093] of the 5-prime sequence, genomic structure, and alternative splicing

[60094] of the presenilin-1 gene (PSEN1) associated with early onset Alzheimer
[60095] disease. Genomics 40: 415-424, 1997.

[60096] 9945.

[60097] Schellenberg, G. D.; Boehnke, M.; Wijsman, E. M.; Moore, D. K.;
[60098] Martin, G. M.; Bird, T. D.: Genetic association and linkage analysis
[60099] of the apolipoprotein CII locus and familial Alzheimer's disease. Ann.
[60100] Neurol. 31: 223-227, 1992.

[60101] 9946.

[60102] Schellenberg, G. D.; Payami, H.; Wijsman, E. M.; Orr, H. T.; Goddard,
[60103] K. A. B.; Anderson, L.; Nemens, E.; White, J. A.; Alonso, M. E.; Ball,
[60104] M. J.; Kaye, J.; Morris, J. C.; Chui, H.; Sadovnick, A. D.; Heston,
[60105] L. L.; Martin, G. M.; Bird, T. D.: Chromosome 14 and late-onset familial
[60106] Alzheimer disease (FAD). Am. J. Hum. Genet. 53: 619-628, 1993.

[60107] 9947.

[60108] Shen, J.; Bronson, R. T.; Chen, D. F.; Xia, W.; Selkoe, D. J.;

[60109] Tonegawa, S.: Skeletal and CNS defects in presenilin-
1-deficient

[60110] mice. Cell 89: 629-639, 1997.

[60111] 9948.

[60112] Smith, M. J.; Kwok, J. B. J.; McLean, C. A.; Kril, J. J.; Broe,

[60113] G. A.; Nicholson, G. A.; Cappai, R.; Hallupp, M.; Cotton, R.
G. H.;

[60114] Masters, C. L.; Schofield, P. R.; Brooks, W. S.: Variable
phenotype

[60115] of Alzheimer's disease with spastic paraparesis. Ann.
Neurol. 49:

[60116] 125-129, 2001.

[60117] 9949.

[60118] Sorbi, S.; Nacmias, B.; Forleo, P.; Piacentini, S.; Sherring-
ton,

[60119] R.; Rogaev, E.; St George Hyslop, P.; Amaducci, L.: Mis-
sense mutation

[60120] of S182 gene in Italian families with early-onset
Alzheimer's disease.

[60121] (Letter) Lancet 346: 439-440, 1995.

[60122] 9950.

[60123] St. George-Hyslop, P. H.; Levesque, G.; Levesque, L.; Rommens,

[60124] J.; Westaway, D.; Fraser, P. E.: Two homologous genes causing early-onset

[60125] familial Alzheimer's disease. Cold Spring Harbor Symp. Quant. Biol. 61:

[60126] 559-564, 1996.

[60127] 9951.

[60128] Shitoh, K.; Konishi, F.; Iijima, T.; Ohdaira, T.; Sakai, K.; Kanazawa,

[60129] K.; Miyaki, M.: A novel case of a sporadic desmoid tumour with mutation

[60130] of the beta catenin gene. J. Clin. Path. 52: 695-696, 1999.

[60131] 9952.

[60132] Hart, T. C.; Zhang, Y.; Gorry, M. C.; Hart, P. S.; Cooper, M.;

[60133] Marazita, M. L.; Marks, J. M.; Cortelli, J. R.; Pallos, D.: A mutation

[60134] in the SOS1 gene causes hereditary gingival fibromatosis type 1. Am.

[60135] J. Hum. Genet. 70: 943-954, 2002.

[60136] 9953.

[60137] Chernova, O. B.; Somerville, R. P. T.; Cowell, J. K.: A novel

[60138] gene, LGI1, from 10q24 is rearranged and downregulated in malignant

[60139] brain tumors. Oncogene 17: 2873-2881, 1998.

[60140] 9954.

[60141] Hamilton, S. R.; Liu, B.; Parsons, R. E.; Papadopoulos, N.; Jen,

[60142] J.; Powell, S. M.; Krush, A. J.; Berk, T.; Cohen, Z.; Tetu, B.; Burger,

[60143] P. C.; Wood, P. A.; Taqi, F.; Booker, S. V.; Petersen, G. M.; Offerhaus,

[60144] G. J. A.; Tersmette, A. C.; Giardiello, F. M.; Vogelstein, B.; Kinzler,

[60145] K. W.: The molecular basis of Turcot's syndrome. New Eng. J. Med. 332:

[60146] 839-847, 1995.

[60147] 9955.

[60148] Holland, E. C.; Celestino, J.; Dai, C.; Schaefer, L.; Sawaya,

[60149] R. E.; Fuller, G. N.: Combined activation of Ras and Akt in neural

[60150] progenitors induces glioblastoma formation in mice. Na-

ture Genet. 25:
[60151] 55-57, 2000.

[60152] 9956.

[60153] Efstratiadis, A.; Posakony, J. W.; Maniatis, T.; Lawn, R. M.;

[60154] O'Connell, C.; Spritz, R. A.; DeRiel, J. K.; Forget, B. G.; Weissman,
[60155] S. M.; Slightom, J. L.; Blechl, A. E.; Smithies, O.; Baralle, F. E.;
[60156] Shoulders, C. C.; Proudfoot, N. J.: The structure and evolution of
[60157] the human beta-globin gene family. Cell 21: 653-668, 1980.
[60158] 9957.

[60159] Jin, Y.; Dietz, H. C.; Montgomery, R. A.; Bell, W. R.; McIntosh,
[60160] I.; Coller, B.; Bray, P. F.: Glanzmann thrombasthenia: cooperation
[60161] between sequence variants in Cis during splice site selection. J.
[60162] Clin. Invest. 98: 1745-1754, 1996.

[60163] 9958.

[60164] Levine, M. A.; Modi, W. S.; O'Brien, S. J.: Chromosomal localization
[60165] of the genes encoding two forms of the G-protein beta polypeptide,
[60166] beta-1 and beta-3, in man. Genomics 8: 380-386, 1990.

[60167] 9959.

[60168] Stallings-Mann, M. L.; Ludwiczak, R. L.; Klinger, K. W.; Rottman,
[60169] F.: Alternative splicing of exon 3 of the human growth hormone receptor
[60170] is the result of an unusual genetic polymorphism. Proc. Nat. Acad.
[60171] Sci. 93: 12394-12399, 1996.

[60172] 9960.

[60173] Friedman, E.; Gejman, P. V.; Martin, G. A.; McCormick, F.: Nonsense
[60174] mutations in the C-terminal SH2 region of the GTPase activating protein
[60175] (GAP) gene in human tumours. Nature Genet. 5: 242-247, 1993.
[60176] 9961.

[60177] Hsieh, C. L.; Francke, U.: The gene for GTPase activating protein

[60178] (GAP) is on human chromosome 5q and mouse chromosome 13. (Abstract) Cytogenet.

[60179] Cell Genet. 51: 1016 only, 1989.

[60180] 9962.

[60181] Hsieh, C. L.; Vogel, U. S.; Dixon, R. A.; Francke, U.: Chromosome

[60182] localization and cDNA sequence of murine and human genes for ras p21

[60183] GTPase activating protein (GAP). Somat. Cell Molec. Genet. 15: 579-590,

[60184] 1989.

[60185] 9963.

[60186] Lemons, R. S.; Espinosa, R., III; Rebentisch, M.; McCormick, F.;

[60187] Ladner, M.; Le Beau, M. M.: Chromosomal localization of the gene

[60188] encoding GTPase-activating protein (RASA) to human chromosome 5, bands

[60189] q13-q15. Genomics 6: 383-385, 1990.

[60190] 9964.

[60191] Mitsudomi, T.; Friedman, E.; Gejman, P. V.; McCormick, F.; Gazdar,

[60192] A. F.: Genetic analysis of the catalytic domain of the GAP gene in

[60193] human lung cancer cell lines. Hum. Genet. 93: 27-31, 1994.

[60194] 9965.

[60195] Trahey, M.; Wong, G.; Halenbeck, R.; Rubinfeld, B.; Martin, G.

[60196] A.; Ladner, M.; Long, C. M.; Crosier, W. J.; Watt, K.; Koths, K.;

[60197] McCormick, F.: Molecular cloning of two types of GAP complementary

[60198] DNA from human placenta. Science 242: 1697-1700, 1988.

[60199] 9966.

[60200] Blatt, C.; Eversole-Cire, P.; Cohn, V. H.; Zollman, S.; Fournier,

[60201] R. E. K.; Mohandas, L. T.; Nesbitt, M.; Lugo, T.; Jones, D. T.; Reed,

[60202] R. R.; Weiner, L. P.; Sparkes, R. S.; Simon, M. I.: Chromo-

somal localization
[60203] of genes encoding guanine nucleotide-binding protein subunits in mouse
[60204] and human. Proc. Nat. Acad. Sci. 85: 7642-7646, 1988.

[60205] 9967.

[60206] Allen, B.; Ostrer, H.; Stein, J.; Stein, G.: Histone gene clusters
[60207] map to chromosomes 1 and 6. (Abstract) Cytogenet. Cell Genet. 51:
[60208] 950 only, 1989.

[60209] 9968.

[60210] Carozzi, N.; Marashi, F.; Plumb, M.; Zimmerman, S.; Zimmerman,
[60211] A.; Coles, L. S.; Wells, J. R. E.; Stein, G.; Stein, J.: Clustering
[60212] of human H1 and core histone genes. Science 224: 1115-1117, 1984.
[60213] 9969.

[60214] Chandler, M. E.; Kedes, L. H.; Cohn, R. H.; Yunis, J. J.: Genes
[60215] coding for histone proteins in man are located on the distal end of

[60216] chromosome 7. Science 205: 908-910, 1979.

[60217] 9970.

[60218] Clark, S. J.; Krieg, P. A.; Wells, J. R. E.: Isolation of a clone

[60219] containing human histone genes. Nucleic Acids Res. 9:
1583-1597,
[60220] 1981.

[60221] 9971.

[60222] Delange, R. J.; Smith, E. L.: Histones: structure and func-
tion. Ann.
[60223] Rev. Biochem. 40: 279-314, 1971.

[60224] 9972.

[60225] Felsenfeld, G.: Chromatin as an essential part of the tran-
scriptional
[60226] mechanism. Nature 355: 219-224, 1992.

[60227] 9973.

[60228] Green, L.; Van Antwerpen, R.; Stein, J.; Stein, G.; Tripputi,
P.;
[60229] Emanuel, B.; Selden, J.; Croce, C.: A major human histone
gene cluster
[60230] on the long arm of chromosome 1. Science 226: 838-840,

1984.

[60231] 9974.

[60232] Heintz, N.; Zernik, M.; Roeder, R. G.: The structure of the human

[60233] histone genes: clustered but not tandemly repeated. Cell 24: 661-668,

[60234] 1981.

[60235] 9975.

[60236] Hentschel, C. C.; Birnstiel, M. L.: The organization and expression

[60237] of histone gene families. Cell 25: 301-313, 1981.

[60238] 9976.

[60239] Kedes, L.; Maxson, R.: Histone gene organization: paradigm lost. Nature 294:

[60240] 11-12, 1981.

[60241] 9977.

[60242] Lichtler, A. C.; Sierra, F.; Clark, S.; Wells, J. R. E.; Stein,

[60243] J. L.; Stein, G. S.: Multiple H4 histone mRNAs of HeLa cells are

[60244] encoded in different genes. Nature 298: 195-198, 1982.

[60245] 9978.

[60246] McAlpine, P. J.: Personal Communication. Winnipeg, Mani-
toba,
[60247] Canada 1989.

[60248] 9979.

[60249] Sierra, F.; Lichtler, A.; Marashi, F.; Rickles, R.; Van Dyke,

[60250] T.; Clark, S.; Wells, J.; Stein, G.; Stein, J.: Organization of
human
[60251] histone genes. Proc. Nat. Acad. Sci. 79: 1795-1799, 1982.

[60252] 9980.

[60253] Steffensen, D. M.: Human histone genes mapped to chro-
mosome 7.
[60254] (Abstract) Cytogenet. Cell Genet. 25: 211 only, 1979.

[60255] 9981.

[60256] Szabo, P.; Yu, L. C.; Borun, T.; Varicchio, F.; Siniscalco, M.;

[60257] Prensky, W.: Localization of the histone genes in man. Cy-
togenet.
[60258] Cell Genet. 22: 359-363, 1978.

[60259] 9982.

[60260] Tripputi, P.; Emanuel, B. S.; Croce, C. M.; Green, L. G.; Stein,
[60261] G. S.; Stein, J. L.: Human histone genes map to multiple chromosomes. Proc.
[60262] Nat. Acad. Sci. 83: 3185-3188, 1986.

[60263] 9983.

[60264] Yunis, J. J.; Chandler, M. E.: Localization of histone genes
[60265] to bands 7q32-36 in man and the homologous chromosome segments in
[60266] chimpanzee, gorilla, and orangutan. (Abstract) Cytogenet. Cell Genet. 25:
[60267] 220 only, 1979.

[60268] 9984.

[60269] Doyle, J.; Hoffman, S.; Ucla, C.; Reith, W.; Mach, B.; Stubbs,
[60270] L.: Locations of human and mouse genes encoding the RFX1 and RFX2
[60271] transcription factor proteins. Genomics 35: 227-230, 1996.
[60272] 9985.

[60273] Pugliatti, L.; Derre, J.; Berger, R.; Ucla, C.; Reith, W.; Mach,

[60274] B.: The genes for MHC class II regulatory factors RFX1 and RFX2 are
[60275] located on the short arm of chromosome 19. Genomics 13: 1307-1310,
[60276] 1992.

[60277] 9986.

[60278] Reith, W.; Ucla, C.; Barras, E.; Gaud, A.; Durand, B.; Herrero-Sanchez,
[60279] C.; Kobr, M.; Mach, B.: RFX1, a transactivator of hepatitis B virus
[60280] enhancer I, belongs to a novel family of homodimeric and heterodimeric
[60281] DNA-binding proteins. Molec. Cell. Biol. 14: 1230-1244, 1994.
[60282] 9987.

[60283] Claesson, L.; Larhammar, D.; Rask, L.; Peterson, P. A.: cDNA clone
[60284] for the human invariant gamma chain of class II histocompatibility
[60285] antigens and its implications for the protein structure. Proc. Nat.
[60286] Acad. Sci. 80: 7395-7399, 1983.

[60287] 9988.

[60288] Claesson-Welsh, L.; Barker, P. E.; Larhammar, D.; Rask, L.; Ruddle,

[60289] F. H.; Peterson, P. A.: The gene encoding the human class II antigen-associated

[60290] gamma chain is located on chromosome 5. Immuno-genetics 20: 89-93,

[60291] 1984.

[60292] 9989.

[60293] Driessen, C.; Bryant, R. A.; Lennon-Dumenil, A. M.; Vil-ladangos,

[60294] J. A.; Bryant, P. W.; Shi, G. P.; Chapman, H. A.; Ploegh, H. L.:

[60295] Cathepsin S controls the trafficking and maturation of MHC class II

[60296] molecules in dendritic cells. J. Cell Biol. 147: 775-790, 1999.

[60297] 9990.

[60298] Genuardi, M.; Saunders, G. F.: Localization of the HLA class II-associated

[60299] invariant chain gene to human chromosome band 5q32. Immunogenetics 28:

[60300] 53-56, 1988.

[60301] 9991.

[60302] Ferdinandusse, S.; Denis, S.; Clayton, P. T.; Graham, A.; Rees,

[60303] J. E.; Allen, J. T.; McLean, B. N.; Brown, A. Y.; Vreken, P.; Waterham,

[60304] H. R.; Wanders, R. J. A.: Mutations in the gene encoding peroxisomal

[60305] alpha-methylacyl-CoA racemase cause adult-onset sensory motor neuropathy. Nature

[60306] Genet. 24: 188-191, 2000.

[60307] 9992.

[60308] Bathori, G.; Parolini, I.; Tombola, F.; Szabo, I.; Messina, A.;

[60309] Oliva, M.; De Pinto, V.; Lisanti, M.; Sargiacomo, M.; Zoratti, M.

[60310] : Porin is present in the plasma membrane where it is concentrated

[60311] in caveolae and caveolae-related domains. J. Biol. Chem. 274: 29607-29612,

[60312] 1999.

[60313] 9993.

[60314] Huizing, M.; Ruitenbeek, W.; Thinnes, F. P.; DePinto, V.; Wendel,
[60315] U.; Trijbels, F. J. M.; Smit, L. M. E.; Ter Laak, H. J.; Van Den Heuvel,
[60316] L. P.: Deficiency of the voltage-dependent anion channel: a novel
[60317] cause of mitochondriopathy. Pediat. Res. 39: 760-765, 1996.
[60318] 9994.

[60319] Cornish, A. L.; Freeman, S.; Forbes, G.; Ni, J.; Zhang, M.; Cepeda,
[60320] M.; Gentz, R.; Augustus, M.; Carter, K. C.; Crocker, P. R.: Characterization
[60321] of siglec-5, a novel glycoprotein expressed on myeloid cells related
[60322] to CD33. Blood 92: 2123-2132, 1998.

[60323] 9995.

[60324] Kim, H.-S.: Assignment of the human OB binding protein-2 gene
[60325] (CD33L2) to chromosome 19q13.3 by radiation hybrid mapping. Cytogenet.
[60326] Cell Genet. 84: 96 only, 1999.

[60327] 9996.

[60328] Patel, N.; Brinkman-Van der Linden, E. C. M.; Altmann, S. W.; Gish,

[60329] K.; Balasubramanian, S.; Timans, J. C.; Peterson, D.; Bell, M. P.;

[60330] Bazan, J. F.; Varki, A.; Kastelein, R. A.: OB-BP1/Siglec-6: a leptin-

[60331] and sialic acid-binding protein of the immunoglobulin superfamily. J.

[60332] Biol. Chem. 274: 22729-22738, 1999.

[60333] 9997.

[60334] Creutz, C. E.; Tomsig, J. L.; Snyder, S. L.; Gautier, M.-C.; Skouri,

[60335] F.; Beisson, J.; Cohen, J.: The copines, a novel class of C2 domain-containing,

[60336] calcium-dependent, phospholipid-binding proteins conserved from Paramecium

[60337] to humans. J. Biol. Chem. 273: 1393-1402, 1998.

[60338] 9998.

[60339] Caudell, E. G.; Caudell, J. J.; Tang, C.-H.; Yu, T.-K.; Frederick,

[60340] M. J.; Grimm, E. A.: Characterization of human copine III as a phosphoprotein

[60341] with associated kinase activity. Biochemistry 39: 13034-13043, 2000.

[60342] 9999.

[60343] Ito, M.; Yuan, C.-X.; Okano, H. J.; Darnell, R. B.; Roeder, R.

[60344] G.: Involvement of the TRAP220 component of the TRAP/SMCC coactivator

[60345] complex in embryonic development and thyroid hormone action. Molec.

[60346] Cell 5: 683-693, 2000.

[60347] 10000.

[60348] Yuan, C.-X.; Ito, M.; Fondell, J. D.; Fu, Z.-Y.; Roeder, R. G.

[60349] : The TRAP220 component of a thyroid hormone receptor-associated protein

[60350] (TRAP) coactivator complex interacts directly with nuclear receptors

[60351] in a ligand-dependent fashion. Proc. Nat. Acad. Sci. 95: 7939-7944,

[60352] 1998. Note: Erratum: Proc. Nat. Acad. Sci. 95: 14584 only, 1998.

[60353] 10001.

[60354] Zhu, Y.; Qi, C.; Jain, S.; Le Beau, M. M.; Espinosa, R., III; Atkins,

[60355] G. B.; Lazar, M. A.; Yeldandi, A. V.; Rao, M. S.; Reddy, J. K.: Amplification

[60356] and overexpression of peroxisome proliferator-activated receptor binding

[60357] protein (PBP/PPARBP) gene in breast cancer. Proc. Nat. Acad. Sci. 96:

[60358] 10848-10853, 1999.

[60359] 10002.

[60360] Zhu, Y.; Qi, C.; Jain, S.; Rao, M. S.; Reddy, J. K.: Isolation

[60361] and characterization of PBP, a protein that interacts with peroxisome

[60362] proliferator-activated receptor. J. Biol. Chem. 272: 25500-25506,

[60363] 1997.

[60364] 10003.

[60365] Kadereit, S.; Gewert, D. R.; Galabru, J.; Hovanessian, A. G.; Meurs,

[60366] E. F.: Molecular cloning of two new interferon-induced,

highly related
[60367] nuclear phosphoproteins. J. Biol. Chem. 268:
24432-24441, 1993.
[60368] 10004.

[60369] Welsh, G. I.; Kadereit, S.; Coccia, E. M.; Hovanessian, A. G.;

[60370] Meurs, E. F.: Colocalization within the nucleolus of two
highly related
[60371] IFN-induced human nuclear phosphoproteins with nucle-
olin. Exp. Cell
[60372] Res. 250: 62-74, 1999.

[60373] 10005.

[60374] Pohl, U.; Smith, J. S.; Tachibana, I.; Ueki, K.; Lee, H. K.; Ra-
maswamy,
[60375] S.; Wu, Q.; Mohrenweiser, H. W.; Jenkins, R. B.; Louis, D.
N.: EHD2,
[60376] EHD3, and EHD4 encode novel members of a highly con-
served family of
[60377] EH domain-containing proteins. Genomics 63: 255-262,
2000.
[60378] 10006.

[60379] Schweizer, A.; Valdenaire, O.; Koster, A.; Lang, Y.; Schmitt,

G.;

[60380] Lenz, B.; Bluethmann, H.; Rohrer, J.: Neonatal lethality in mice

[60381] deficient in XCE, a novel member of the endothelin-converting enzyme

[60382] and neutral endopeptidase family. J. Biol. Chem. 274: 20450-20456,

[60383] 1999.

[60384] 10007.

[60385] Valdenaire, O.; Richards, J. G.; Faull, R. L. M.; Schweizer, A.

[60386] : XCE, a new member of the endothelin-converting enzyme and neutral

[60387] endopeptidase family, is preferentially expressed in the CNS. Molec.

[60388] Brain Res. 64: 211-221, 1999.

[60389] 10008.

[60390] Valdenaire, O.; Rohrbacher, E.; Langeveld, A.; Schweizer, A.; Meijers,

[60391] C.: Organization and chromosomal localization of the human ECEL1

[60392] (XCE) gene encoding a zinc metallopeptidase involved in

the nervous
[60393] control of respiration. Biochem. J. 346: 611-616, 2000.

[60394] 10009.

[60395] Hengst, U.; Albrecht, H.; Hess, D.; Monard, D.: The phosphatidylethanolamine-binding
[60396] protein is the prototype of a novel family of serine protease inhibitors. J.
[60397] Biol. Chem. 276: 535-540, 2001.

[60398] 10010.

[60399] Hori, N.; Chae, K.; Murakawa, K.; Matoba, R.; Fukushima, A.; Okubo,
[60400] K.; Matsubara, K.: A human cDNA sequence homologue of bovine phosphatidylethanolamine-binding
[60401] protein. Gene 140: 293-294, 1994.

[60402] 10011.

[60403] Moore, C.; Perry, A. C. F.; Love, S.; Hall, L.: Sequence analysis
[60404] and immunolocalisation of phosphatidylethanolamine binding protein
[60405] (PBP) in human brain tissue. Molec. Brain Res. 37: 74-78, 1996.

[60406] 10012.

[60407] Schoentgen, F.; Jolles, P.: From structure to function: pos-
sible
[60408] biological roles of a new widespread protein family bind-
ing hydrophobic
[60409] ligands and displaying a nucleotide binding site. FEBS Lett.
369:
[60410] 22-26, 1995.

[60411] 10013.

[60412] Seddiqi, N.; Bollengier, F.; Alliel, P. M.; Perin, J.-P.; Bonnet,
[60413] F.; Bucquoy, S.; Jolles, P.; Schoentgen, F.: Amino acid se-
quence
[60414] of the Homo sapiens brain 21-23-kDa protein
(neuropolypeptide h3),
[60415] comparison with its counterparts from Rattus norvegicus
and Bos tauras
[60416] species, and expression of its mRNA in different tissues. J.
Molec.
[60417] Evol. 39: 655-660, 1994.

[60418] 10014.

[60419] Tohdoh, N.; Tojo, S.; Agui, H.; Ojika, K.: Sequence homol-

ogy of
[60420] rat and human HCNP precursor proteins, bovine phosphatidylethanolamine-binding
[60421] protein and rat 23-kDa protein associated with the opioid-binding
[60422] protein. Molec. Brain Res. 30: 381-384, 1995.

[60423] 10015.

[60424] Yeung, K.; Seitz, T.; Li, S.; Janosch, P.; McFerran, B.; Kaiser,
[60425] C.; Fee, F.; Katsanakis, K. D.; Rose, D. W.; Mischak, H.; Sedivy,
[60426] J. M.; Kolch, W.: Suppression of Raf-1 kinase activity and MAP kinase
[60427] signalling by RKIP. Nature 401: 173-177, 1999.

[60428] 10016.

[60429] Heinemann, T.; Bulwin, G. C.; Randall, J.; Schnieders, B.; Sandhoff,
[60430] K.; Volk, H. D.; Milford, E.; Gullans, S. R.; Utku, N.: Genomic organization
[60431] of the gene coding for TIRC7, a novel membrane protein essential for
[60432] T cell activation. Genomics 57: 398-406, 1999.

[60433] 10017.

[60434] Li, Y.-P.; Chen, W.; Liang, Y.; Li, E.; Stashenko, P.:
ATP6i-deficient
[60435] mice exhibit severe osteopetrosis due to loss of osteo-
clast-mediated
[60436] extracellular acidification. Nature Genet. 23: 447-451,
1999.
[60437] 10018.

[60438] Li, Y.-P.; Chen, W.; Stashenko, P.: Molecular cloning and
characterization
[60439] of a putative novel human osteoclast-specific 116-kDa
vacuolar proton
[60440] pump subunit. Biochem. Biophys. Res. Commun. 218:
813-821, 1996.
[60441] 10019.

[60442] Scimeca, J.-C.; Franchi, A.; Trojani, C.; Parrinello, H.;
Grosgeorge,
[60443] J.; Robert, C.; Jaillon, O.; Poirier, C.; Gaudray, P.; Carle, G.
F.
[60444] : The gene encoding the mouse homologue of the human
osteoclast-specific
[60445] 116-kDa V-ATPase subunit bears a deletion in osteoscle-

rotic (oc/oc)

[60446] mutants. Bone 26: 207-213, 2000.

[60447] 10020.

[60448] Sobacchi, C.; Frattini, A.; Orchard, P.; Porras, O.; Tezcan, I.;

[60449] Andolina, M.; Babul-Hirji, R.; Baric, I.; Canham, N.; Chitayat, D.;

[60450] Dupuis-Girod, S.; Ellis, I.; and 21 others: The mutational spectrum

[60451] of human malignant autosomal recessive osteopetrosis. Hum. Molec.

[60452] Genet. 10: 1767-1773, 2001.

[60453] 10021.

[60454] Utku, N.; Heinemann, T.; Tullius, S. G.; Bulwin, G. C.; Beinke,

[60455] S.; Blumberg, R. S.; Beato, F.; Randall, J.; Kojima, R.; Busconi,

[60456] L.; Robertson, E. S.; Schulein, R.; Volk, H. D.; Milford, E. L.; Gullans,

[60457] S. R.: Prevention of acute allograft rejection by antibody targeting

[60458] of TIRC7, a novel T cell membrane protein. Immunity 9:

509-518,

[60459] 1998.

[60460] 10022.

[60461] Blum, R.; Feick, P.; Puype, M.; Vandekerckhove, J.; Klengel, R.;

[60462] Nastainczyk, W.; Schulz, I.: Tmp21 and p24A, two type I proteins

[60463] enriched in pancreatic microsomal membranes, are members of a protein

[60464] family involved in vesicular trafficking. J. Biol. Chem. 271: 17183-17189,

[60465] 1996.

[60466] 10023.

[60467] Horer, J.; Blum, R.; Feick, P.; Nastainczyk, W.; Schulz, I.: A

[60468] comparative study of rat and human Tmp21 (p23) reveals the pseudogene-like

[60469] features of human Tmp21-II. DNA Seq. 10: 121-126, 1999.

[60470] 10024.

[60471] Sherrington, R.; Rogaev, E. I.; Liang, Y.; Rogaeva, E. A.; Levesque,

[60472] G.; Ikeda, M.; Chi, H.; Lin, C.; Li, G.; Holman, K.; Tsuda, T.; Mar,
[60473] L.; Foncin, J.-F.; Bruni, A. C.; Montesi, M. P.; Sorbi, S.; Rainero,
[60474] I.; Pinessi, L.; Nee, L.; Chumakov, I.; Pollen, D.; Brookes, A.; Sanseau,
[60475] P.; Polinsky, R. J.; Wasco, W.; Da Silva, H. A. R.; Haines, J. L.;
[60476] Pericak-Vance, M. A.; Tanzi, R. E.; Roses, A. D.; Fraser, P. E.; Rommens,
[60477] J. M.; St George-Hyslop, P. H.: Cloning of a gene bearing missense
[60478] mutations in early-onset familial Alzheimer's disease. Nature 375:
[60479] 754-760, 1995.

[60480] 10025.

[60481] Postma, A. V.; Bezzina, C. R.; de Vries, J. F.; Wilde, A. A. M.;
[60482] Moorman, A. F. M.; Mannens, M. M. A. M.: Genomic organisation and
[60483] chromosomal localisation of two members of the KCND ion channel family,

[60484] KCND2 and KCND3. Hum. Genet. 106: 614-619, 2000.

[60485] 10026.

[60486] Zhu, X.-R.; Wulf, A.; Schwarz, M.; Isbrandt, D.; Pongs, O.: Characterization

[60487] of human Kv4.2 mediating a rapidly-inactivating transient voltage-sensitive

[60488] K+ current. Receptors Channels 6: 387-400, 1999.

[60489] 10027.

[60490] Wong, A. C. C.; Shkolny, D.; Dorman, A.; Willingham, D.; Roe, B.

[60491] A.; McDermid, H. E.: Two novel human RAB genes with near identical

[60492] sequence each map to a telomere-associated region: the subtelomeric

[60493] region of 22q13.3 and the ancestral telomere band 2q13. Genomics 59:

[60494] 326-334, 1999.

[60495] 10028.

[60496] Harada, H.; Nagai, H.; Tsuneizumi, M.; Mikami, I.; Sugano, S.;

[60497] Emi, M.: Identification of DMC1, a novel gene in the TOC

region on

[60498] 17q25.1 that shows loss of expression in multiple human cancers. J.

[60499] Hum. Genet. 46: 90-95, 2001.

[60500] 10029.

[60501] van den Eynde, B. J.; Gaugler, B.; Probst-Kepper, M.; Michaux,

[60502] L.; Devuyst, O.; Lorge, F.; Weynants, P.; Boon, T.: A new antigen

[60503] recognized by cytolytic T lymphocytes on a human kidney tumor results

[60504] from reverse strand transcription. J. Exp. Med. 190: 1793-1799,

[60505] 1999.

[60506] 10030.

[60507] Kile, B. T.; Viney, E. M.; Willson, T. A.; Brodnicki, T. C.; Cancilla,

[60508] M. R.; Herlihy, A. S.; Croker, B. A.; Baca, M.; Nicola, N. A.; Hilton,

[60509] D. J.; Alexander, W. S.: Cloning and characterization of the genes

[60510] encoding the ankyrin repeat and SOCS box-containing

proteins Asb-1,
[60511] Asb-2, Asb-3, and Asb-4. Gene 258: 31-41, 2000.

[60512] 10031.

[60513] Liu, Y.; Corcoran, M.; Rasool, O.; Ivanova, G.; Ibbotson, R.; Grander,
[60514] D.; Iyengar, A.; Baranova, A.; Kashuba, V.; Merup, M.; Wu, X.; Gardiner,
[60515] A.; and 12 others: Cloning of two candidate tumor suppressor genes
[60516] within a 10 kb region on chromosome 13q14, frequently deleted in chronic
[60517] lymphocytic leukemia. Oncogene 15: 2463-2473, 1997.

[60518] 10032.

[60519] Pappu, R.; Cheng, A. M.; Li, B.; Gong, Q.; Chiu, C.; Griffin, N.;
[60520] White, M.; Sleckman, B. P.; Chan, A. C.: Requirement for B cell linker
[60521] protein (BLNK) in B cell development. Science 286: 1949-1954, 1999.

[60522] 10033.

[60523] Wienands, J.; Schweikert, J.; Wollscheid, B.; Jumaa, H.;

Nielsen,
[60524] P. J.; Reth, M.: SLP-65: a new signaling component in B lymphocytes
[60525] which requires expression of the antigen receptor for phosphorylation. J.
[60526] Exp. Med. 188: 791-795, 1998.

[60527] 10034.

[60528] Agarwal, A. K.; Garg, A.: A novel heterozygous mutation in peroxisome
[60529] proliferator-activated receptor-gamma gene in a patient with familial
[60530] partial lipodystrophy. J. Clin. Endocr. Metab. 87: 408-411, 2002.
[60531] 10035.

[60532] Cosman, D.; Fanger, N.; Borges, L.; Kubin, M.; Chin, W.; Peterson,
[60533] L.; Hsu, M.-L.: A novel immunoglobulin superfamily receptor for cellular
[60534] and viral MHC class I molecules. Immunity 7: 273-282, 1997.
[60535] 10036.

[60536] Muto, T.; Muramatsu, M.; Taniwaki, M.; Kinoshita, K.;

Honjo, T.

[60537] : Isolation, tissue distribution, and chromosomal localization of

[60538] the human activation-induced cytidine deaminase (AID) gene. Genomics 68:

[60539] 85-88, 2000.

[60540] 10037.

[60541] Geck, P.; Maffini, M. V.; Szelei, J.; Sonnenschein, C.; Soto, A.

[60542] M.: Androgen-induced proliferative quiescence in prostate cancer

[60543] cells: the role of AS3 as its mediator. Proc. Nat. Acad. Sci. 97:

[60544] 10185-10190, 2000.

[60545] 10038.

[60546] Kas, K.; Finger, E.; Grall, F.; Gu, X.; Akbarali, Y.; Boltax, J.;

[60547] Weiss, A.; Oettgen, P.; Kapeller, R.; Libermann, T. A.: ESE-3, a

[60548] novel member of an epithelium-specific Ets transcription factor subfamily,

[60549] demonstrates different target gene specificity from ESE-1.

J. Biol.

[60550] Chem. 275: 2986-2998, 2000.

[60551] 10039.

[60552] Kleinbaum, L. A.; Duggan, C.; Ferreira, E.; Coffey, G. P.; Buttice,

[60553] G.; Burton, F. H.: Human chromosomal localization, tissue/tumor expression,

[60554] and regulatory function of the ets family gene EHF. Biochem. Biophys.

[60555] Res. Commun. 264: 119-126, 1999.

[60556] 10040.

[60557] Davidson, J. D.; Riley, B.; Burright, E.N.; Duvick, L.A.; Zoghbi,

[60558] H.Y.; Orr, H. T.: Identification and characterization of an ataxin-1-interacting

[60559] protein: A1Up, a ubiquitin-like nuclear protein. Hum. Molec. Genet. 9:

[60560] 2305-2312, 2000.

[60561] 10041.

[60562] Arita, Y.; Kihara, S.; Ouchi, N.; Takahashi, M.; Maeda, K.; Miyagawa,

[60563] J.; Hotta, K.; Shimomura, I.; Nakamura, T.; Miyaoka, K.; Kuriyama,

[60564] H.; Nishida, M.; Yamashita, S.; Okubo, K.; Matsubara, K.; Muraguchi,

[60565] M.; Ohmoto, Y.; Funahashi, T.; Matsuzawa, Y.: Paradoxical decrease

[60566] of an adipose-specific protein, adiponectin, in obesity. Biochem.

[60567] Biophys. Res. Commun. 257: 79-83, 1999.

[60568] 10042.

[60569] Comuzzie, A. G.; Funahashi, T.; Sonnenberg, G.; Martin, L. J.;

[60570] Jacob, H. J.; Kwitek Black, A. E.; Maas, D.; Takahashi, M.; Kihara,

[60571] S.; Tanaka, S.; Matsuzawa, Y.; Blangero, J.; Cohen, D.; Kissebah,

[60572] A.: The genetic basis of plasma variation in adiponectin, a global

[60573] endophenotype for obesity and the metabolic syndrome. J. Clin. Endocr.

[60574] Metab. 86: 4321-4325, 2001.

[60575] 10043.

[60576] Das, K.; Lin, Y.; Widen, E.; Zhang, Y.; Scherer, P. E.: Chromosomal

[60577] localization, expression pattern, and promoter analysis of the mouse

[60578] gene encoding adipocyte-specific secretory protein Acrp30. Biochem.

[60579] Biophys. Res. Commun. 280: 1120-1129, 2001.

[60580] 10044.

[60581] Maeda, K.; Okubo, K.; Shimomura, I.; Funahashi, T.; Matsuzawa,

[60582] Y.; Matsubara, K.: cDNA cloning and expression of a novel adipose

[60583] specific collagen-like factor, apM1 (adipose most abundant gene transcript

[60584] 1). Biochem. Biophys. Res. Commun. 221: 286-289, 1996.

[60585] 10045.

[60586] Maeda, N.; Shimomura, I.; Kishida, K.; Nishizawa, H.; Matsuda,

[60587] M.; Nagaretani, H.; Furuyama, N.; Kondo, H.; Takahashi, M.; Arita,

[60588] Y.; Komuro, R.; Ouchi, N.; Kihara, S.; Tochino, Y.; Okutomi, K.; Horie,

[60589] M.; Takeda, S.; Aoyama, T.; Funahashi, T.; Matsuzawa, Y.: Diet-induced

[60590] insulin resistance in mice lacking adiponectin/ACRP30. Nature Med. 8:

[60591] 731-737, 2002.

[60592] 10046.

[60593] Ouchi, N.; Kihara, S.; Arita, Y.; Maeda, K.; Kuriyama, H.; Okamoto,

[60594] Y.; Hotta, K.; Nishida, M.; Takahashi, M.; Nakamura, T.; Yamashita,

[60595] S.; Funahashi, T.; Matsuzawa, Y.: Novel modulator for en-dothelial

[60596] adhesion molecules: adipocyte-derived plasma protein adiponectin. Circulation 100:

[60597] 2473-2476, 1999.

[60598] 10047.

[60599] Saito, K.; Tobe, T.; Minoshima, S.; Asakawa, S.; Sumiya, J.; Yoda,

[60600] M.; Nakano, Y.; Shimizu, N.; Tomita, M.: Organization of the gene

[60601] for gelatin-binding protein (GBP28). Gene 229: 67-73, 1999.

[60602] 10048.

[60603] Zhao, L.; Gregoire, F.; Sul, H. S.: Transient induction of ENC-1,

[60604] a kelch-related actin-binding protein, is required for adipocyte differentiation. J.

[60605] Biol. Chem. 275: 16845-16850, 2000.

[60606] 10049.

[60607] Kuno, K.; Kanada, N.; Nakashima, E.; Fujiki, F.; Ichimura, F.;

[60608] Matsushima, K.: Molecular cloning of a gene encoding a new type of

[60609] metalloproteinase-disintegrin family protein with thrombospondin motifs

[60610] as an inflammation associated gene. J. Biol. Chem. 272: 556-562,

[60611] 1997.

[60612] 10050.

[60613] Hansen, L. L.; Jorgensen, R.; Justesen, J.: Assignment of the

[60614] human mitochondrial translational release factor 1 (MTRF1) to chromosome

[60615] 13q14.1-q14.3 and of the human mitochondrial ribosome recycling factor

[60616] (MRRF) to chromosome 9q32-q34.1 with radiation hybrid mapping. Cytogenet.

[60617] Cell Genet. 88: 91-92, 2000.

[60618] 10051.

[60619] Zhang, Y.; Spremulli, L. L.: Identification and cloning of human

[60620] mitochondrial translational release factor 1 and the ribo- some recycling

[60621] factor. Biochim. Biophys. Acta 1443: 245-250, 1998.

[60622] 10052.

[60623] Sim, D. L.; Chow, V. T.: The novel human HUEL (C4orf1) gene maps

[60624] to chromosome 4p12-p13 and encodes a nuclear protein containing the

[60625] nuclear receptor interaction motif. Genomics 59: 224-233, 1999.

[60626] 10053.

[60627] Stelnicki, E. J.; Arbeit, J.; Cass, D. L.; Saner, C.; Harrison,

[60628] M.; Largman, C.: Modulation of the human homeobox

genes PRX-2 and
[60629] HOXB13 in scarless fetal wounds. J. Invest. Derm. 111: 57-63, 1998.
[60630] 10054.

[60631] Beadling, C.; Johnson, K. W.; Smith, K. A.: Isolation of interleukin
[60632] 2-induced immediate-early genes. Proc. Nat. Acad. Sci. 90: 2719-2723,
[60633] 1993.

[60634] 10055.

[60635] Gong, R.; Yu, L.; Zhang, H.; Tu, Q.; Zhao, Y.; Yang, J.; Xu, Y.;
[60636] Zhao, S.: Assignment of human GADD45G to chromosome 9q22.1-q22.3
[60637] by radiation hybrid mapping. Cytogenet. Cell Genet. 88: 95-96, 2000.
[60638] 10056.

[60639] Nakayama, K.; Hara, T.; Hibi, M.; Hirano, T.; Miyajima, A.: A
[60640] novel oncostatin M-inducible gene OIG37 forms a gene family with MyD118
[60641] and GADD45 and negatively regulates cell growth. J. Biol.

Chem. 274:
[60642] 24766-24772, 1999.

[60643] 10057.

[60644] Suzuki, M.; Watanabe, T. K.; Fujiwara, T.; Nakamura, Y.; Takahashi,
[60645] E.; Tanigami, A.: Molecular cloning, expression, and mapping of a
[60646] novel human cDNA, GRP17, highly homologous to human gadd45 and murine
[60647] MyD118. J. Hum. Genet. 44: 300-303, 1999.

[60648] 10058.

[60649] Zhang, X.; Sun, H.; Danila, D. C.; Johnson, S. R.; Zhou, Y.; Swearingen,
[60650] B.; Klibanski, A.: Loss of expression of GADD45-gamma, a growth inhibitory
[60651] gene, in human pituitary adenomas: implications for tumorigenesis. J.
[60652] Clin. Endocr. Metab. 87: 1262-1267, 2002.

[60653] 10059.

[60654] Donzeau, M.; Kaldi, K.; Adam, A.; Paschen, S.; Wanner, G.; Guiard,

[60655] B.; Bauer, M. F.; Neupert, W.; Brunner, M.: Tim23 links the inner
[60656] and outer mitochondrial membranes. Cell 101: 401-412, 2000.
[60657] 10060.
[60658] Scott, D. A.; Wang, R.; Kreman, T. M.; Sheffield, V. C.; Karniski,
[60659] L. P.: The Pendred syndrome gene encodes a chloride-iodide transport
[60660] protein. Nature Genet. 21: 440-443, 1999.
[60661] 10061.
[60662] Scott, A. F.: Personal Communication. Baltimore, Md. 10/23/2001.
[60663] 10062.
[60664] Pajor, A. M.: Sodium-coupled transporters for Krebs cycle intermediates. Annu.
[60665] Rev. Physiol. 61: 663-682, 1999.
[60666] 10063.
[60667] McGaughran, J. M.; Ward, H. B.; Evans, D. G.: WAGR syndrome and
[60668] multiple exostoses in a patient with del(11)(p11.2p14.2).

J. Med.

[60669] Genet. 32: 823-824, 1995.

[60670] 10064.

[60671] Wuyts, W.; Van Hul, W.; Wauters, J.; Nemtsova, M.;
Reyniers, E.;

[60672] Van Hul, E. V.; De Boulle, K.; de Vries, B. B.; Hendrickx, J.;
Herrygers,

[60673] I.; Bossuyt, P.; Balemans, W.; Fransen, E.; Vits, L.; Coucke,
P.;

[60674] Nowak, N. J.; Shows, T. B.; Mallet, L.; van den Ouweland,
A. M.; McGaughran,

[60675] J.; Halley, D. J.; Willems, P. J.: Positional cloning of a gene
involved

[60676] in hereditary multiple exostoses. Hum. Molec. Genet. 5:
1547-1557,

[60677] 1996.

[60678] 10065.

[60679] Creasy, C. L.; Chernoff, J.: Cloning and characterization of
a

[60680] human protein kinase with homology to Ste20. J. Biol.
Chem. 270:

[60681] 21695-21700, 1995.

[60682] 10066.

[60683] Taylor, L. K.; Wang, H.-C. R.; Erikson, R. L.: Newly identified

[60684] stress-responsive protein kinases, Krs-1 and Krs-2. Proc. Nat. Acad.

[60685] Sci. 93: 10099-10104, 1996.

[60686] 10067.

[60687] Carmo-Fonseca, M.; Kern, H.; Hurt, E. C.: Human nucleoporin p62

[60688] and the essential yeast nuclear pore protein NSP1 show sequence homology

[60689] and a similar domain organization. Europ. J. Cell Biol. 55: 17-30,

[60690] 1991.

[60691] 10068.

[60692] Lennard, A.; Gorman, P.; Carrier, M.; Griffiths, S.; Scotney, H.;

[60693] Sheer, D.; Solari, R.: Cloning and chromosome mapping of the human

[60694] interleukin-1 receptor antagonist gene. Cytokine 4: 83-89, 1992.

[60695] 10069.

[60696] Mansfield, J. C.; Holden, H.; Tarlow, J. K.; Di Giovine, F. S.;

[60697] McDowell, T. L.; Wilson, A. G.; Holdsworth, C. D.; Duff, G. W.: Novel

[60698] genetic association between ulcerative colitis and the anti-inflammatory

[60699] cytokine interleukin-1 receptor antagonist. Gastroenterol-ogy 106:

[60700] 637-642, 1994.

[60701] 10070.

[60702] Patterson, D.; Jones, C.; Hart, I.; Bleskan, J.; Berger, R.; Geyer,

[60703] D.; Eisenberg, S. P.; Smith, M. F., Jr.; Arend, W. P.: The hu-man

[60704] interleukin-1 receptor antagonist (IL1RN) gene is located in the chromosome

[60705] 2q14 region. Genomics 15: 173-176, 1993.

[60706] 10071.

[60707] Steinkasserer, A.; Spurr, N. K.; Cox, S.; Jeggo, P.; Sim, R. B.

[60708] : The human IL-1 receptor antagonist gene (IL1RN) maps to chromosome

[60709] 2q14-q21, in the region of the IL-1-alpha and IL-1-beta loci. Genomics 13:
[60710] 654-657, 1992.

[60711] 10072.

[60712] Tarlow, J. K.; Blakemore, A. I. F.; Lennard, A.; Solari, R.; Hughes,
[60713] H. N.; Steinkasserer, A.; Duff, G. W.: Polymorphism in human IL-1
[60714] receptor antagonist gene intron 2 is caused by variable numbers of
[60715] an 86-bp tandem repeat. Hum. Genet. 91: 403-404, 1993.
[60716] 10073.

[60717] Tarlow, J. K.; Clay, F. E.; Cork, M. J.; Blakemore, A. I. F.;

[60718] McDonagh, A. J. G.; Messenger, A. G.; Duff, G. W.: Severity of alopecia
[60719] areata is associated with a polymorphism in the interleukin-1 receptor
[60720] antagonist gene. J. Invest. Dermatol. 103: 387-390, 1994.

[60721] 10074.

[60722] Wuyts, W.; Ramlakhan, S.; Van Hul, W.; Hecht, J. T.; van

den Ouweland,
[60723] A. M. W.; Raskind, W. H.; Hofstede, F. C.; Reyniers, E.; Wells, D.
[60724] E.; de Vries, B.; Conrad, E. U.; Hill, A.; Zalatayev, D.; Weissenbach,
[60725] J.; Wagner, M. J.; Bakker, E.; Halley, D. J. J.; Willems, P. J.:
[60726] Refinement of the multiple exostoses locus (EXT2) to a 3-cM interval
[60727] on chromosome 11. Am. J. Hum. Genet. 57: 382-387, 1995.
[60728] 10075.
[60729] Schild, D.; Brake, A. J.; Kiefer, M. C.; Young, D.; Barr, P. J.
[60730] : Cloning of three human multifunctional de novo purine biosynthetic
[60731] genes by functional complementation of yeast mutations. Proc. Nat.
[60732] Acad. Sci. 87: 2916-2920, 1990.
[60733] 10076.
[60734] Garrow, T. A.; Brenner, A. A.; Whitehead, V. M.; Chen, X.-N.; Duncan,
[60735] R. G.; Korenberg, J. R.; Shane, B.: Cloning of human cDNAs

encoding

[60736] mitochondrial and cytosolic serine hydroxymethyltrans-
ferases and chromosomal

[60737] localization. J. Biol. Chem. 268: 11910-11916, 1993.

[60738] 10077.

[60739] Kao, F.-T.; Chasin, L. A.; Puck, T. T.: Genetics of somatic
mammalian

[60740] cells. X. Complementation analysis of glycine-requiring
mutants. Proc.

[60741] Nat. Acad. Sci. 64: 1284-1291, 1969.

[60742] 10078.

[60743] Stover, P. J.; Chen, L. H.; Suh, J. R.; Stover, D. M.; Key-
omarsi,

[60744] K.; Shane, B.: Molecular cloning, characterization, and
regulation

[60745] of the human mitochondrial serine hydroxymethyltrans-
ferase gene. J.

[60746] Biol. Chem. 272: 1842-1848, 1997.

[60747] 10079.

[60748] Alper, C. A.: Inherited structural polymorphism in human
C2: evidence

[60749] for genetic linkage between C2 and Bf. J. Exp. Med. 144: 1111-1115,

[60750] 1976.

[60751] 10080.

[60752] Wuyts, W.; Van Hul, W.; Wauters, J.; Nemtsova, M.; Reyniers, E.;

[60753] Van Hul, E.; De Boulle, K.; de Vries, B. B. A.; Hendrickx, J.; Herrygers,

[60754] I.; Bossuyt, P.; Balemans, W.; Fransen, E.; Vits, L.; Coucke, P.;

[60755] Nowak, N. J.; Shows, T. B.; Mallet, L.; van den Ouweland, A. M. W.;

[60756] McGaughran, J.; Halley, D. J. J.; Willems, P. J.: Positional cloning

[60757] of a gene involved in hereditary multiple exostoses. Hum. Molec.

[60758] Genet. 5: 1547-1557, 1996.

[60759] 10081.

[60760] Fulton, A. B.; Hansen, R. M.; Petersen, R. A.; Vanderveen, D. K.

[60761] : The rod photoreceptors in retinopathy of prematurity: an electroretinographic

[60762] study. Arch. Ophthal. 119: 499-505, 2001.

[60763] 10082.

[60764] Rochelle, J. M.; Watson, M. L.; Oakey, R. J.; Seldin, M. F.:

[60765] A linkage map of mouse chromosome 19: definition of comparative mapping

[60766] relationships with human chromosomes 10 and 11 in-cluding the MEN1

[60767] locus. Genomics 14: 26-31, 1992.

[60768] 10083.

[60769] Seldin, M. F.; Saunders, A. M.; Rochelle, J. M.; Howard, T. A.

[60770] : A proximal mouse chromosome 9 linkage map that fur-ther defines linkage

[60771] groups homologous with segments of human chromo-somes 11, 15, and 19. Genomics 9:

[60772] 678-685, 1991.

[60773] 10084.

[60774] Buckwalter, M. S.; Lossie, A. C.; Scarlett, L. M.; Camper, S. A.

[60775] : Localization of the human chromosome 5q genes Gabra-1, Gabrg-2,

[60776] I1-4, I1-5, and Irf-1 on mouse chromosome 11. Mam-
malian Genome 3:
[60777] 604-607, 1992.

[60778] 10085.

[60779] Jensen, K. B.; Dredge, B. K.; Stefani, G.; Zhong, R.; Buck-
anovich,
[60780] R. J.; Okano, H. J.; Yang, Y. Y. L.; Darnell, R. B.: Nova-1
regulates
[60781] neuron-specific alternative splicing and is essential for
neuronal
[60782] viability. Neuron 25: 359-371, 2000.

[60783] 10086.

[60784] Ozcelik, T.; Suedhof, T. C.; Francke, U.: The genes for in-
ositol
[60785] 1,4,5-triphosphate receptors 1 (ITPR1) and 3 (ITPR3) are
localized
[60786] on human chromosomes 3p and 6pter-p21, respectively.
(Abstract) Cytogenet.
[60787] Cell Genet. 58: 1880, 1991.

[60788] 10087.

[60789] Woodroofe, M. N.; Tunnacliffe, A.; Pym, B.; Goodfellow, P.

N.;

[60790] Walsh, F. S.: Human muscle cell surface antigen 16-3A5 is encoded

[60791] by a gene on chromosome 11. Somat. Cell Molec. Genet. 10: 535-540,

[60792] 1984.

[60793] 10088.

[60794] Sudo, T.; Ozawa, K.; Soeda, E.-I.; Nomura, N.; Ishii, S.: Mapping

[60795] of the human gene for the human immunodeficiency virus type 1 enhancer

[60796] binding protein HIV-EP2 to chromosome 6q23-q24. Genomics 12: 167-170,

[60797] 1992.

[60798] 10089.

[60799] Li, C.; Lusis, A. J.; Sparkes, R.; Tran, S.-M.; Gaynor, R.: Characterization

[60800] and chromosomal mapping of the gene encoding the cellular DNA binding

[60801] protein HTLF. Genomics 13: 658-664, 1992.

[60802] 10090.

[60803] Adams, P.; Falek, A.; Arnold, J.: Huntington disease in Georgia:
[60804] age at onset. Am. J. Hum. Genet. 43: 695-704, 1988.

[60805] 10091.

[60806] Almqvist, E.; Adam, S.; Bloch, M.; Fuller, A.; Welch, P.; Eisenberg,
[60807] D.; Whelan, D.; Macgregor, D.; Meschino, W.; Hayden, M. R.: Risk
[60808] reversals in predictive testing for Huntington disease. Am. J. Hum.
[60809] Genet. 61: 945-952, 1997.

[60810] 10092.

[60811] Almqvist, E.; Andrew, S.; Theilmann, J.; Goldberg, P.; Zeisler,
[60812] J.; Drugge, U.; Grandell, U.; Tapper-Persson, M.; Winblad, B.; Hayden,
[60813] M.; Anvret, M.: Geographical distribution of haplotypes in Swedish
[60814] families with Huntington's disease. Hum. Genet. 94: 124-128, 1994.

[60815] 10093.

[60816] Almqvist, E. W.; Elterman, D. S.; MacLeod, P. M.; Hayden, M. R.

[60817] : High incidence rate and absent family histories in one quarter of

[60818] patients newly diagnosed with Huntington disease in British Columbia. Clin.

[60819] Genet. 60: 198-205, 2001.

[60820] 10094.

[60821] Altherr, M. R.; Wasmuth, J. J.; Seldin, M. F.; Nadeau, J. H.; Baehr,

[60822] W.; Pittler, S. J.: Chromosome mapping of the rod pho-

toreceptor cGMP

[60823] phosphodiesterase beta-subunit gene in mouse and hu-

man: tight linkage

[60824] to the Huntington disease region (4p16.3). Genomics 12: 750-754,

[60825] 1992.

[60826] 10095.

[60827] de Vree, J. M. L.; Jacquemin, E.; Sturm, E.; Cresteil, D.; Bosma,

[60828] P. J.; Aten, J.; Deleuze, J.-F.; Desrochers, M.; Burdelski, M.; Bernard,

[60829] O.; Oude Elferink, R. P. J.; Hadchouel, M.: Mutations in the MDR3
[60830] gene cause progressive familial intrahepatic cholestasis. Proc. Nat.
[60831] Acad. Sci. 95: 282-287, 1998.

[60832] 10096.

[60833] Watanabe, T.; Bai, M.; Lane, C. R.; Matsumoto, S.; Minami-tani,
[60834] K.; Minagawa, M.; Niimi, H.; Brown, E. M.; Yasuda, T.: Fa-milial hypoparathyroidism:
[60835] identification of a novel gain of function mutation in transmembrane
[60836] domain 5 of the calcium-sensing receptor. J. Clin. Endocr. Metab. 83:
[60837] 2497-2502, 1998.

[60838] 10097.

[60839] Hasegawa, T.; Hasegawa, Y.; Aso, T.; Koto, S.; Nagai, T.; Tsuchiya,
[60840] Y.; Kim, K. C.; Ohashi, H.; Wakui, K.; Fukushima, Y.: HDR syndrome
[60841] (hypoparathyroidism, sensorineural deafness, renal dys-plasia) associated

[60842] with del(10)(p13). Am. J. Med. Genet. 73: 416-418, 1997.

[60843] 10098.

[60844] Henthorn, P. S.; Raducha, M.; Fedde, K. N.; Lafferty, M. A.; Whyte,

[60845] M. P.: Different missense mutations at the tissue-nonspecific alkaline

[60846] phosphatase gene locus in autosomal recessively inherited forms of

[60847] mild and severe hypophosphatasia. Proc. Nat. Acad. Sci. 89: 9924-9928,

[60848] 1992.

[60849] 10099.

[60850] Macfarlane, J. D.; Kroon, H. M.; van der Harten, J. J.: Phe-notypically

[60851] dissimilar hypophosphatasia in two sibships. Am. J. Med. Genet. 42:

[60852] 117-121, 1992.

[60853] 10100.

[60854] Moore, C. A.; Curry, C. J. R.; Henthorn, P. S.; Smith, J. A.;

[60855] Smith, J. C.; O'Lague, P.; Coburn, S. P.; Weaver, D. D.; Whyte, M.

[60856] P.: Mild autosomal dominant hypophosphatasia: in utero presentation
[60857] in two families. Am. J. Med. Genet. 86: 410-415, 1999.

[60858] 10101.

[60859] Mornet, E.: Personal Communication. Paris, France 6/3/1999.

[60860] 10102.

[60861] Yamada, N.; Makino, Y.; Clark, R. A.; Pearson, D. W.; Mat-
[60862] tei, M.-G.; Guenet, J.-L.; Ohama, E.; Fujino, I.; Miyawaki, A.; Furuichi, T.;
[60863] Mikoshiba, K.: Human inositol 1,4,5-triphosphate type-1 receptor,
[60864] InsP3R1: structure, function, regulation of expression and chromosomal
[60865] localization. Biochem. J. 302: 781-790, 1994.

[60866] 10103.

[60867] Maranto, A. R.: Primary structure, ligand binding, and lo-
[60868] calization of the human type 3 inositol 1,4,5-trisphosphate receptor expressed

[60869] in intestinal epithelium. J. Biol. Chem. 269: 1222-1230, 1994.

[60870] 10104.

[60871] Ozcelik, T.; Suedhof, T. C.; Francke, U.: The genes for inositol

[60872] 1,4,5-triphosphate receptors 1 (ITPR1) and 3 (ITPR3) are localized

[60873] on human chromosomes 3p and 6pter-p21, respectively. (Abstract) Cytogenet.

[60874] Cell Genet. 58: 1880 only, 1991.

[60875] 10105.

[60876] Yamamoto-Hino, M.; Sugiyama, T.; Hikichi, K.; Mattei, M. G.; Hasegawa,

[60877] K.; Sekine, S.; Sakurada, K.; Miyawaki, A.; Furuichi, T.; Hasegawa,

[60878] M.; Mikoshiba, K.: Cloning and characterization of human type 2 and

[60879] type 3 inositol 1,4,5-triphosphate receptors. Receptors Channels 2:

[60880] 9-22, 1994.

[60881] 10106.

[60882] Acquati, F.; Malgaretti, N.; Hauptschein, R.; Rao, P.; Gaidano,
[60883] G.; Taramelli, R.: A 2-Mb YAC contig linking the plas-minogen-apoprotein(a)
[60884] gene family to the insulin-like growth factor 2 receptor (IGF2R) gene
[60885] on the telomeric region of chromosome 6 (6q26-q27). Genomics 22:
[60886] 664-666, 1994.

[60887] 10107.

[60888] Barlow, D. P.; Stoger, R.; Herrmann, B. G.; Saito, K.; Schweifer,
[60889] N.: The mouse insulin-like growth factor type-2 receptor is imprinted
[60890] and closely linked to the Tme locus. Nature 349: 84-87, 1991.
[60891] 10108.

[60892] DeChiara, T. M.; Robertson, E. J.; Efstratiadis, A.: Parental
[60893] imprinting of the mouse insulin-like growth factor II gene. Cell 64:
[60894] 849-859, 1991.

[60895] 10109.

[60896] De Souza, A. T.; Hankins, G. R.; Washington, M. K.; Fine, R. L.;

[60897] Orton, T. C.; Jirtle, R. L.: Frequent loss of heterozygosity on 6q

[60898] at the mannose 6-phosphate/insulin-like growth factor II receptor

[60899] locus in human hepatocellular tumors. Oncogene 10: 1725-1729, 1995.

[60900] 10110.

[60901] De Souza, A. T.; Hankins, G. R.; Washington, M. K.; Orton, T. C.;

[60902] Jirtle, R. L.: M6P/IGF2R gene is mutated in human hepato-cellular

[60903] carcinomas with loss of heterozygosity. Nature Genet. 11: 447-449,

[60904] 1995.

[60905] 10111.

[60906] Feinberg, A. P.: Genomic imprinting and gene activation in cancer. Nature

[60907] Genet. 4: 110-113, 1993.

[60908] 10112.

[60909] Haig, D.; Graham, C.: Genomic imprinting and the strange case

[60910] of the insulin-like growth factor II receptor. Cell 64: 1045-1046,

[60911] 1991.

[60912] 10113.

[60913] Haig, D.; Westoby, M.: Parent-specific gene expression and the

[60914] triploid endosperm. Am. Nat. 134: 147-155, 1989.

[60915] 10114.

[60916] Kalscheuer, V. M.; Mariman, E. C.; Schepens, M. T.; Re-hder, H.;

[60917] Ropers, H.-H.: The insulin-like-growth factor type-2 re-ceptor gene

[60918] is imprinted in the mouse but not in humans. Nature Genet. 5: 74-78,

[60919] 1993.

[60920] 10115.

[60921] Kiess, W.; Blickenstaff, G. D.; Sklar, M. M.; Thomas, C. L.; Nissley,

[60922] S. P.; Sahagian, G. G.: Biochemical evidence that the type II insulin-like
[60923] growth factor receptor is identical to the cation-in-dependent mannose
[60924] 6-phosphate receptor. J. Biol. Chem. 263: 9339-9344, 1988.
[60925] 10116.
[60926] Killian, J. K.; Oka, Y.; Jang, H.-S.; Fu, X.; Waterland, R. A.;
[60927] Sohda, T.; Sakaguchi, S.; Jirtle, R. L.: Mannose 6-phosphate/insulin-like
[60928] growth factor 2 receptor (M6P/IGF2R) variants in American and Japanese
[60929] populations. Hum. Mutat. 18: 25-31, 2001.
[60930] 10117.
[60931] Kornfeld, S.; Mellman, I.: The biogenesis of lysosomes. Annu.
[60932] Rev. Cell Biol. 5: 483-525, 1989.
[60933] 10118.
[60934] Lau, M. M. H.; Stewart, C. E. H.; Liu, Z.; Bhatt, H.; Rotwein,
[60935] P.; Stewart, C. L.: Loss of the imprinted IGF2/cation-independent

[60936] mannose 6-phosphate receptor results in fetal overgrowth and perinatal
[60937] lethality. Genes Dev. 8: 2953-2963, 1994.

[60938] 10119.

[60939] Laureys, G.; Barton, D. E.; Ullrich, A.; Francke, U.: Chromosomal
[60940] mapping of the gene for the type II insulin-like growth factor receptor/cation-independent
[60941] mannose 6-phosphate receptor in man and mouse. Genomics 3: 224-229,
[60942] 1988.

[60943] 10120.

[60944] MacDonald, R. G.; Pfeffer, S. R.; Coussens, L.; Tepper, M. A.;
[60945] Brocklebank, C. M.; Mole, J. E.; Anderson, J. K.: A single receptor
[60946] binds both insulin-like growth factor II and mannose-6-phosphate. Science 239:
[60947] 1134-1137, 1988.

[60948] 10121.

[60949] Morgan, D. O.; Edman, J. D.; Standring, D. N.; Fried, V. A.;

Smith,
[60950] M. C.; Roth, R. A.; Rutter, W. J.: Insulin-like growth factor II
[60951] receptor as a multifunctional binding protein. Nature 329: 301-307,
[60952] 1987.

[60953] 10122.

[60954] Benharroch, D.; Meguerian-Bedoyan, Z.; Lamant, L.; Amin, C.; Brugieres,
[60955] L.; Terrier-Lacombe, M.-J.; Haralambieva, E.; Pulford, K.; Pileri,
[60956] S.; Morris, S. W.; Mason, D. Y.; Delsol, G.: ALK-positive lymphoma:
[60957] a single disease with a broad spectrum of morphology. Blood 91:
[60958] 2076-2084, 1998.

[60959] 10123.

[60960] Mathew, P.; Morris, S. W.; Kane, J. R.; Shurtleff, S. A.; Pasquini,
[60961] M.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.: Localization
[60962] of the murine homolog of the anaplastic lymphoma kinase

(Alk) gene
[60963] on mouse chromosome 17. Cytogenet. Cell Genet. 70: 143-144, 1995.
[60964] 10124.
[60965] Morris, S. W.; Kirstein, M. N.; Valentine, M. B.; Dittmer, K. G.;
[60966] Shapiro, D. N.; Saltman, D. L.; Look, A. T.: Fusion of a ki-
nase gene,
[60967] ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. Science 263:
[60968] 1281-1284, 1994.
[60969] 10125.
[60970] Otto, E.; Kunimoto, M.; McLaughlin, T.; Bennett, V.: Isola-
tion
[60971] and characterization of cDNAs encoding human brain ankyrins reveal
[60972] a family of alternatively spliced genes. J. Cell Biol. 114: 241-253,
[60973] 1991.
[60974] 10126.
[60975] Tse, W. T.; Menninger, J. C.; Yang-Feng, T. L.; Francke, U.;

Sahr,
[60976] K. E.; Lux, S. E.; Ward, D. C.; Forget, B. G.: Isolation and chromosomal
[60977] localization of a novel non-erythroid ankyrin gene. Genomics 10:
[60978] 858-866, 1991.

[60979] 10127.

[60980] Grundmann, U.; Amann, E.; Abel, K.-J.; Kupper, H. A.: Isolation
[60981] and expression of cDNA coding for a new member of the phospholipase
[60982] A2 inhibitor family. Behring Inst. Mitt. 82: 59-67, 1988.

[60983] 10128.

[60984] Hauptmann, R.; Maurer-Fogy, I.; Krystek, E.; Bodo, G.; Andree,
[60985] H.; Reutelingsperger, C. P. M.: Vascular anticoagulant beta, a novel
[60986] human Ca(2+)/phospholipid binding protein that inhibits coagulation
[60987] and phospholipase A-2 activity: its molecular cloning, expression
[60988] and comparison with VAC-alpha. Europ. J. Biochem. 185:

63-71, 1989.

[60989] 10129.

[60990] Tait, J. F.; Smith, C.; Frankenberry, D. A.; Miao, C. H.; Adler,

[60991] D. A.; Disteche, C. M.: Chromosomal mapping of the human annexin

[60992] IV (ANX4) gene. Genomics 12: 313-318, 1992.

[60993] 10130.

[60994] Bilezikian, J. P.; Morishima, A.; Bell, J.; Grumbach, M. M.: Increased

[60995] bone mass as a result of estrogen therapy in a man with aromatase

[60996] deficiency. New Eng. J. Med. 339: 599-603, 1998.

[60997] 10131.

[60998] Oldridge, M.; Temple, I. K. Santos, H. G.; Gibbons, R. J.; Mustafa,

[60999] Z.; Chapman, K. E.; Loughlin, J.; Wilkie, A. O. M.: Brachydactyly

[61000] type B: linkage to chromosome 9q22 and evidence for genetic heterogeneity. Am.

[61001] J. Hum. Genet. 64: 578-585, 1999.

[61002] 10132.

[61003] Breimer, L. H.; MacIntyre, I.; Zaidi, M.: Peptides from the calcitonin

[61004] genes: molecular genetics, structure and function. Biochem. J. 255:

[61005] 377-390, 1988.

[61006] 10133.

[61007] Schultz, D.; Mikala, G.; Yatani, A.; Engle, D. B.; Iles, D. E.;

[61008] Segers, B.; Sinke, R. J.; Weghuis, D. O.; Klockner, U.; Wakamori,

[61009] M.; Wang, J.-J.; Melvin, D.; Varadi, G.; Schwartz, A.: Cloning, chromosomal

[61010] localization, and functional expression of the alpha-1 subunit of

[61011] the L-type voltage-dependent calcium channel from nor- mal human heart. Proc.

[61012] Nat. Acad. Sci. 90: 6228-6232, 1993.

[61013] 10134.

[61014] Soldatov, N. M.: Genomic structure of human L-type Ca(2+) channel. Genomics 22:

[61015] 77-87, 1994.

[61016] 10135.

[61017] Sun, W.; McPherson, J. D.; Hoang, D. Q.; Wasmuth, J. J.; Evans,
[61018] G. A.; Montal, M.: Mapping of a human brain voltage-gated calcium
[61019] channel to human chromosome 12p13-pter. Genomics 14: 1092-1094,
[61020] 1992.

[61021] 10136.

[61022] Tsien, R. W.; Ellinor, P. T.; Horne, W. A.: Molecular diver-sity
[61023] of voltage-dependent Ca(2+) channels. Trends Pharm. Sci. 12: 349-354,
[61024] 1991.

[61025] 10137.

[61026] Chin, H.; Kozak, C. A.; Kim, H.-L.; Mock, B.; McBride, O. W.:
[61027] A brain L-type calcium channel alpha-1 subunit gene (CCHL1A2) maps
[61028] to mouse chromosome 14 and human chromosome 3. Genomics 11: 914-919,

[61029] 1991.

[61030] 10138.

[61031] Jinnah, H. A.; Yitta, S.; Drew, T.; Kim, B. S.; Visser, J. E.;

[61032] Rothstein, J. D.: Calcium channel activation and self-biting in mice. Proc.

[61033] Nat. Acad. Sci. 96: 15228-15232, 1999.

[61034] 10139.

[61035] Mori, Y.; Friedrich, T.; Kim, M.-S.; Mikami, A.; Nakai, J.; Ruth,

[61036] P.; Bosse, E.; Hofmann, F.; Flockerzi, V.; Furuichi, T.; Mikoshiba,

[61037] K.; Imoto, K.; Tanabe, T.; Numa, S.: Primary structure and functional

[61038] expression from complementary DNA of a brain calcium channel. Nature 350:

[61039] 398-402, 1991.

[61040] 10140.

[61041] Pennartz, C. M. A.; de Jeu, M. T. G.; Bos, N. P. A.; Schaap, J.;

[61042] Geurtsen, A. M. S.: Diurnal modulation of pacemaker potentials and

[61043] calcium current in the mammalian circadian clock. Nature 416: 286-290,

[61044] 2002.

[61045] 10141.

[61046] Platzer, J.; Engel, J.; Schrott-Fischer, A.; Stephan, K.; Bova,

[61047] S.; Chen, H.; Zheng, H.; Striessnig, J.: Congenital deafness and

[61048] sinoatrial node dysfunction in mice lacking class D L-type Ca(2+)

[61049] channels. Cell 102: 89-97, 2000.

[61050] 10142.

[61051] Seino, S.; Chen, L.; Seino, M.; Blondel, O.; Takeda, J.; John-son,

[61052] J. H.; Bell, G. I.: Cloning of the alpha-1 subunit of a volt-age-dependent

[61053] calcium channel expressed in pancreatic beta-cells. Proc. Nat. Acad.

[61054] Sci. 89: 584-588, 1992.

[61055] 10143.

[61056] Seino, S.; Yamada, Y.; Espinosa, R., III; Le Beau, M. M.; Bell,

[61057] G. I.: Assignment of the gene encoding the alpha-1 sub-
unit of the
[61058] neuroendocrine/brain-type calcium channel (CACNL1A2)
to human chromosome
[61059] 3, band p14.3. Genomics 13: 1375-1377, 1992.

[61060] 10144.

[61061] Powers, P. A.; Liu, S.; Hogan, K.; Gregg, R. G.: Skeletal
muscle
[61062] and brain isoforms of a beta-subunit of human voltage-
dependent calcium
[61063] channels are encoded by a single gene. J. Biol. Chem. 267:
22967-22972,
[61064] 1992.

[61065] 10145.

[61066] Gregg, R. G.; Messing, A.; Strube, C.; Beurg, M.; Moss, R.;
Behan,
[61067] M.; Sukhareva, M.; Haynes, S.; Powell, J. A.; Coronado, R.;
Powers,
[61068] P. A.: Absence of the beta subunit (cchb1) of the skeletal
muscle
[61069] dihydropyridine receptor alters expression of the alpha-1
subunit

[61070] and eliminates excitation-contraction coupling. Proc. Nat. Acad.

[61071] Sci. 93: 13961-13966, 1996.

[61072] 10146.

[61073] Iles, D. E.; Segers, B.; Sengers, R. C. A.; Monsieurs, K.; Heytens,

[61074] L.; Halsall, P. J.; Hopkins, P. M.; Ellis, F. R.; Hall-Curran, J.

[61075] L.; Stewart, A. D.; Wieringa, B.: Genetic mapping of the beta-1-

[61076] and gamma-subunits of the human skeletal muscle L-type voltage-dependent

[61077] calcium channel on chromosome 17q and exclusion as candidate genes

[61078] for malignant hyperthermia susceptibility. Hum. Molec. Genet. 2:

[61079] 863-868, 1993.

[61080] 10147.

[61081] Pragnell, M.; Sakamoto, J.; Jay, S. D.; Campbell, K. P.: Cloning

[61082] and tissue-specific expression of the brain calcium channel beta-subunit. FEBS

[61083] Lett. 291: 253-258, 1991.

[61084] 10148.

[61085] Boerman, R. H.; Ophoff, R. A.; Links, T. P.; van Eijk, R.; Sandkuijl,

[61086] L. A.; Elbaz, A.; Vale-Santos, J. E.; Wintzen, A. R.; van Deutekom,

[61087] J. C.; Isles, D. E.; Fontaine, B.; Padberg, G. W.; Frants, R. R.:

[61088] Mutation in DHP receptor alpha-1 subunit (CACLN1A3) (sic) gene in

[61089] a Dutch family with hypokalaemic periodic paralysis. J. Med. Genet. 32:

[61090] 44-47, 1995.

[61091] 10149.

[61092] Chaudhari, N.: A single nucleotide deletion in the skeletal muscle-specific

[61093] calcium channel transcript of muscular dysgenesis (mdg) mice. J.

[61094] Biol. Chem. 267: 25636-25639, 1992.

[61095] 10150.

[61096] Chin, H.; Krall, M.; Kim, H.-L.; Kozak, C. A.; Mock, B.: The gene

[61097] for the alpha-1 subunit of the skeletal muscle dihydropy-
ridine-sensitive
[61098] calcium channel (Cchl1a3) maps to mouse chromosome 1.
Genomics 14:
[61099] 1089-1091, 1992.

[61100] 10151.

[61101] Bulun, S. E.: Aromatase deficiency in women and men:
would you
[61102] have predicted the phenotypes? J. Clin. Endocr. Metab. 81:
867-871,
[61103] 1996.

[61104] 10152.

[61105] Carani, C.; Qin, K.; Simoni, M.; Faustini-Fustini, M.; Ser-
pente,
[61106] S.; Boyd, J.; Korach, K. S.; Simpson, E. R.: Effect of testos-
terone
[61107] and estradiol in a man with aromatase deficiency. New
Eng. J. Med. 337:
[61108] 91-95, 1997.

[61109] 10153.

[61110] Chen, S.; Besman, M. J.; Sparkes, R. S.; Zollman, S.; Klisak,

I.;

[61111] Mohandas, T.; Hall, P. F.; Shively, J. E.: Human aromatase: cDNA

[61112] cloning, Southern blot analysis, and assignment of the gene to chromosome

[61113] 15. DNA 7: 27-38, 1988.

[61114] 10154.

[61115] Chen, S.; Shively, J. E.; Nakajin, S.; Shinoda, M.; Hall, P. F.

[61116] : Amino terminal sequence analysis of human placenta aromatase. Biochem.

[61117] Biophys. Res. Commun. 135: 713-719, 1986.

[61118] 10155.

[61119] Conte, F. A.; Grumbach, M. M.; Ito, Y.; Fisher, C. R.; Simpson,

[61120] E. R.: A syndrome of female pseudohermaphrodism, hypergonadotropic

[61121] hypogonadism, and multicystic ovaries associated with missense mutations

[61122] in the gene encoding aromatase (P450arom). J. Clin. Endocr. Metab. 78:

[61123] 1287-1292, 1994.

[61124] 10156.

[61125] Corbin, C. J.; Graham-Lorence, S.; McPhaul, M.; Mason, J. I.; Mendelson,

[61126] C. R.; Simpson, E. R.: Isolation of a full-length cDNA insert encoding

[61127] human aromatase system cytochrome P-450 and its expression in nonsteroidogenic

[61128] cells. Proc. Nat. Acad. Sci. 85: 8948-8952, 1988.

[61129] 10157.

[61130] Deladoey, J.; Fluck, C.; Bex, M.; Yoshimura, N.; Harada, N.; Mullis,

[61131] P. E.: Aromatase deficiency caused by a novel P450(arom) gene mutation:

[61132] impact of absent estrogen production on serum gonadotropin concentration

[61133] in a boy. J. Clin. Endocr. Metab. 84: 4050-4054, 1999.

[61134] 10158.

[61135] Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Significant population

[61136] variation in adult male height associated with the Y chromosome and

[61137] the aromatase gene. J. Clin. Endocr. Metab. 86:
4147-4150, 2001.

[61138] 10159.

[61139] Evans, C. T.; Ledesma, D. B.; Schulz, T. Z.; Simpson, E. R.;
Mendelson,

[61140] C. R.: Isolation and characterization of a complementary
DNA specific

[61141] for human aromatase-system cytochrome P-450 mRNA.
Proc. Nat. Acad.

[61142] Sci. 83: 6387-6391, 1986.

[61143] 10160.

[61144] Fisher, C. R.; Graves, K. H.; Parlow, A. F.; Simpson, E. R.:

[61145] Characterization of mice deficient in aromatase (ArKO)
because of

[61146] targeted disruption of the cyp19 gene. Proc. Nat. Acad.
Sci. 95:

[61147] 6965-6970, 1998.

[61148] 10161.

[61149] George, F. W.; Matsumine, H.; McPhaul, M. J.; Somes, R. G.,
Jr.;

[61150] Wilson, J. D.: Inheritance of the henny feathering trait in

the Golden

[61151] Campine chicken: evidence for allelism with the gene that causes henny

[61152] feathering in the Sebright Bantam. J. Hered. 81: 107-110, 1990.

[61153] 10162.

[61154] George, F. W.; Wilson, J. D.: Pathogenesis of the henny feathering

[61155] trait in the Sebright Bantam chicken. J. Clin. Invest. 66: 57-65,

[61156] 1980.

[61157] 10163.

[61158] Harada, N.: Cloning of a complete cDNA encoding human aromatase:

[61159] immunochemical identification and sequence analysis. Biochem. Biophys.

[61160] Res. Commun. 156: 725-732, 1988.

[61161] 10164.

[61162] Harada, N.; Ogawa, H.; Shozu, M.; Yamada, K.: Genetic studies

[61163] to characterize the origin of the mutation in placental aromatase

[61164] deficiency. Am. J. Hum. Genet. 51: 666-672, 1992.

[61165] 10165.

[61166] Harada, N.; Ogawa, H.; Shozu, M.; Yamada, K.; Suhara, K.; Nishida,

[61167] E.; Takagi, Y.: Biochemical and molecular genetic analyses on placental

[61168] aromatase (P-450-AROM) deficiency. J. Biol. Chem. 267: 4781-4785,

[61169] 1992.

[61170] 10166.

[61171] Hemsell, D. L.; Edman, C. D.; Marks, J. F.; Siiteri, P. K.; MacDonald,

[61172] P. C.: Massive extraglandular aromatization of plasma androstenedione

[61173] resulting in feminization of a prepubertal boy. J. Clin. Invest. 60:

[61174] 455-464, 1977.

[61175] 10167.

[61176] Ito, Y.; Fisher, C. R.; Conte, F. A.; Grumbach, M. M.; Simpson,

[61177] E. R.: Molecular basis of aromatase deficiency in an adult female

[61178] with sexual infantilism and polycystic ovaries. Proc. Nat. Acad.

[61179] Sci. 90: 11673-11677, 1993.

[61180] 10168.

[61181] Leiberman, E.; Zachmann, M.: Familial adrenal feminization probably

[61182] due to increased steroid aromatization. Hormone Res. 37: 96-102,

[61183] 1992.

[61184] 10169.

[61185] Leshin, M.; Baron, J.; George, F. W.; Wilson, J. D.: Increased

[61186] estrogen formation and aromatase activity in fibroblasts cultured

[61187] from the skin of chickens with the Henny feathering trait. J. Biol.

[61188] Chem. 256: 4341-4344, 1981.

[61189] 10170.

[61190] Leshin, M.; George, F. W.; Wilson, J. D.: Increased estrogen

[61191] synthesis in the Sebright bantam is due to a mutation that causes

[61192] increased aromatase activity. Trans. Assoc. Am. Phys. 94: 97-105,

[61193] 1981.

[61194] 10171.

[61195] Mango, D.; Montemurro, A.; Scirpa, P.; Bompiani, A.; Menini, E.

[61196] : Four cases of pregnancy with low estrogen production due to placental

[61197] enzymatic deficiency. Europ. J. Obstet. Gynec. Reprod. Biol. 8:
[61198] 65-71, 1978.

[61199] 10172.

[61200] McTernan, P. G.; Anderson. L. A.; Anwar, A. J.; Eggo, M. C.; Crocker,
[61201] J.; Barnett, A. H.; Stewart, P. M.; Kumar, S.: Glucocorticoid regulation
[61202] of P450 aromatase activity in human adipose tissue: gen- der and site
[61203] differences. J. Clin. Endocr. Metab. 87: 1327-1336, 2002.

[61204] 10173.

[61205] Morishima, A.; Grumbach, M. M.; Simpson, E. R.; Fisher, C.; Qin,
[61206] K.: Aromatase deficiency in male and female siblings caused by a
[61207] novel mutation and the physiological role of estrogens. J. Clin.
[61208] Endocr. Metab. 80: 3689-3698, 1995.

[61209] 10174.

[61210] Mullis, P. E.; Yoshimura, N.; Kuhlmann, B.; Lippuner, K.;

Jaeger,
[61211] P.; Harada, H.: Aromatase deficiency in a female who is compound
[61212] heterozygote for two new point mutations in the P450(arom) gene: impact
[61213] of estrogens on hypergonadotropic hypogonadism, multi-cystic ovaries,
[61214] and bone densitometry in childhood. J. Clin. Endocr. Metab. 82:
[61215] 1739-1745, 1997.

[61216] 10175.

[61217] Phornphutkul, C.; Okubo, T.; Wu, K.; Harel, Z.; Tracy, T. F.,

[61218] Jr.; Pinar, H.; Chen, S.; Gruppuso, P. A.; Goodwin, G.: Aro-matase
[61219] P450 expression in a feminizing adrenal adenoma pre-senting as isosexual
[61220] precocious puberty. J. Clin. Endocr. Metab. 86: 649-652, 2001.
[61221] 10176.

[61222] Robertson, K. M.; O'Donnell, L.; Jones, M. E. E.; Meachem, S.
[61223] J.; Boon, W. C.; Fisher, C. R.; Graves, K. H.; McLachlan, R.

I.; Simpson,
[61224] E. R.: Impairment of spermatogenesis in mice lacking a functional
[61225] aromatase (cyp 19) gene. Proc. Nat. Acad. Sci. 96: 7986-7991, 1999.
[61226] 10177.

[61227] Sebastian, S.; Bulun, S. E.: A highly complex organization of
[61228] the regulatory region of the human CYP19 (aromatase) gene revealed
[61229] by the Human Genome Project. J. Clin. Endocr. Metab. 86: 4600-4602,
[61230] 2001.

[61231] 10178.

[61232] Allen, S. J.; O'Donnell, A.; Alexander, N. D. E.; Mgone, C. S.;
[61233] Peto, T. E. A.; Clegg, J. B.; Alpers, M. P.; Weatherall, D. J.: Prevention
[61234] of cerebral malaria in children in Papua New Guinea by southeast Asian
[61235] ovalocytosis band 3. Am. J. Trop. Med. Hyg. 60: 1056-1060, 1999.

[61236] 10179.

[61237] Alloisio, N.; Maillet, P.; Carre, G.; Texier, P.; Vallier, A.;

[61238] Baklouti, F.; Philippe, N.; Delaunay, J.: Hereditary spherocytosis

[61239] with band 3 deficiency: association with a nonsense mutation of the

[61240] band 3 gene (allele Lyon), and aggravation by a low-expression allele

[61241] occurring in trans (allele Genas). Blood 88: 1062-1069, 1996.

[61242] 10180.

[61243] Alloisio, N.; Texier, P.; Vallier, A.; Ribeiro, M. L.; Morle, L.;

[61244] Bozon, M.; Bursaux, E.; Maillet, P.; Goncalves, P.; Tanner, M. J.

[61245] A.; Tamagnini, G.; Delaunay, J.: Modulation of clinical expression

[61246] and band 3 deficiency in hereditary spherocytosis. Blood 90: 414-420,

[61247] 1997.

[61248] 10181.

[61249] Amato, D.; Booth, P. B.: Hereditary ovalocytosis in Melane-

sians. Papua

[61250] New Guinea Med. J. 20: 26-32, 1977.

[61251] 10182.

[61252] Baehner, R. L.; Gilchrist, G. S.; Anderson, E. J.: Hereditary

[61253] elliptocytosis and primary renal tubular acidosis in a sin-
gle family. Am.

[61254] J. Dis. Child. 115: 414-419, 1968.

[61255] 10183.

[61256] Baer, A.: Elliptocytosis, malaria, and fertility in Malaysia.
Hum.

[61257] Biol. 60: 909-915, 1988.

[61258] 10184.

[61259] Baer, A.; Lie-Injo, L. E.; Welch, Q. B.; Lewis, A. N.: Genetic

[61260] factors and malaria in the Temuan. Am. J. Hum. Genet. 28:
179-188,

[61261] 1976.

[61262] 10185.

[61263] Booth, P. B.: The occurrence of weak I(T) red cell antigen
among

[61264] Melanesians. Vox Sang. 22: 64-72, 1972.

[61265] 10186.

[61266] Booth, P. B.; Serjeantson, S.; Woodfield, D. G.; Amato, D.: Selective

[61267] depression of blood group antigens associated with hereditary ovalocytosis

[61268] among Melanesians. Vox Sang. 32: 99-110, 1977.

[61269] 10187.

[61270] Bracher, N. A.; Lyons, C. A.; Wessels, G.; Mansvelt, E.; Co-etzer,

[61271] T. L.: Band 3 Cape Town (E90K) causes severe hereditary spherocytosis

[61272] in combination with band 3 Prague III. Brit. J. Haemat. 113: 689-693,

[61273] 2001.

[61274] 10188.

[61275] Bruce, L. J.; Anstee, D. J.; Spring, F. A.; Tanner, M. J. A.:

[61276] Band 3 Memphis variant. II. Altered stilbene disulfonate binding and

[61277] the Diego (Di(a)) blood group antigen are associated with the human

[61278] erythrocyte band 3 mutation pro854-to-leu. J. Biol. Chem. 269: 16155-16158,

[61279] 1994.

[61280] 10189.

[61281] Bruce, L. J.; Ring, S. M.; Anstee, D. J.; Reid, M. E.; Wilkinson,

[61282] S.; Tanner, M. J. A.: Changes in the blood group Wright antigens

[61283] are associated with a mutation at amino acid 658 in human erythrocyte

[61284] band 3: a site of interaction between band 3 and glycophorin A under

[61285] certain conditions. Blood 85: 541-547, 1995.

[61286] 10190.

[61287] Bruce, L. J.; Tanner, M. J.; Zelinski, T.: The low incidence

[61288] blood group antigen, Wd(a), is associated with the substitution val557-to-met

[61289] in human erythrocyte band 3. (Abstract) Transfusion 35 (suppl.):

[61290] 52S only, 1995.

[61291] 10191.

[61292] Bruce, L. J.; Wrong, O.; Toye, A. M.; Young, M. T.; Ogle, G.;

[61293] Ismail, Z.; Sinha, A. K.; McMaster, P.; Hwaihwanje, I.; Nash, G. B.;

[61294] Hart, S.; Lavu, E.; Palmer, R.; Othman, A.; Unwin, R. J.; Tanner,

[61295] M. J. A.: Band 3 mutations, renal tubular acidosis and South-East

[61296] Asian ovalocytosis in Malaysia and Papua New Guinea: loss of up to

[61297] 95% band 3 transport in red cells. Biochem. J. 350: 41-51, 2000.

[61298] 10192.

[61299] Coetzer, T. L.; Beeton, L.; van Zyl, D.; Field, S. P.; Smart,

[61300] E.; Daniels, G. L.: Southeast Asian ovalocytosis in a South African

[61301] kindred with hemolytic anemia. Blood 87: 1656-1658, 1996.

[61302] 10193.

[61303] del Giudice, E.; Perrotta, S.; Pinto, L.; Cappellini, M. D.; Fiorelli,

[61304] G.; Cutillo, S.; Iolascon, A.: Hereditary spherocytosis characterized

[61305] by increased spectrin/band 3 ratio. Brit. J. Haemat. 80: 133-136,
[61306] 1992.

[61307] 10194.

[61308] del Giudice, E. M.; Perrotta, S.; Nobili, B.; Pinto, L.; Cutillo,
[61309] L.; Iolascon, A.: Coexistence of hereditary spherocytosis (HS) due
[61310] to band 3 deficiency and beta-thalassaemia trait: partial correction
[61311] of HS phenotype. Brit. J. Haemat. 85: 553-557, 1993.

[61312] 10195.

[61313] Dhermy, D.; Galand, C.; Bournier, O.; Boulanger, L.; Cynober,
[61314] T.; Schismanoff, P. O.; Bursaux, E.; Tchernia, G.; Boivin, P.; Garbarz,
[61315] M.: Heterogenous band 3 deficiency in hereditary spherocytosis related
[61316] to different band 3 gene defects. Brit. J. Haemat. 98: 32-40, 1997.

[61317] 10196.

[61318] Feest, T. G.; Proctor, S.; Brown, R.; Wrong, O. M.: Nephro-

calcinosis:
[61319] another cause of renal erythrocytosis. Brit. Med. J. 2: 605 only,
[61320] 1978.

[61321] 10197.

[61322] Fix, A. G.; Baer, A. S.; Lie-Injo, L. E.: The mode of inheri-tance
[61323] of ovalocytosis/elliptocytosis in Malaysian Orang Asli families. Hum.
[61324] Genet. 61: 250-253, 1982.

[61325] 10198.

[61326] Ganesan, J.; Lie-Injo, L. E.; Ong, B. P.: Abnormal hemoglobins,
[61327] glucose-6-phosphate dehydrogenase deficiency and hereditary ovalocytosis
[61328] in the Dayaks of Sarawak. Hum. Hered. 25: 258-262, 1975.
[61329] 10199.

[61330] Groves, J. D.; Ring, S. M.; Schofield, A. E.; Tanner, M. J. A.

[61331] : The expression of the abnormal human red cell anion transporter

[61332] from South-East Asian ovalocytes (band 3 SAO) in Xenopus oocytes. FEBS

[61333] Lett. 330: 186-190, 1993.

[61334] 10200.

[61335] Mohr, J.; Eiberg, H.: Colton blood groups: indication of linkage

[61336] with the Kidd (Jk) system as support for assignment to chromosome

[61337] 7. Clin. Genet. 11: 372-374, 1977.

[61338] 10201.

[61339] Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Hormones,

[61340] active peptides and toxins. Washington: National Biomedical Research

[61341] Foundation (pub.) 5: 1972. Pp. D205 only.

[61342] 10202.

[61343] Edbrooke, M. R.; Parker, D.; McVey, J. H.; Riley, J. H.; Sorenson,

[61344] G. D.; Pettengill, O. S.; Craig, R. K.: Expression of the human calcitonin/CGRP

[61345] gene in lung and thyroid carcinoma. EMBO J. 4: 715-724,

1985.

[61346] 10203.

[61347] Girgis, S. I.; Macdonald, D. W. R.; Stevenson, J. C.; Bevis, P.

[61348] J. R.; Lynch, C.; Wimalawansa, S. J.; Self, C. H.; Morris, H. R.;

[61349] MacIntyre, I.: Calcitonin gene-related peptide: potent va-

[61350] sodilator and major product of calcitonin gene. Lancet II: 14-16,

[61351] 1985.

[61352] 10204.

[61353] Goltzman, D.; Mitchell, J.: Interaction of calcitonin and calcitonin gene-related peptide at receptor sites in target tissues.

[61354] Science 227: 1343-1345, 1985.

[61355] 10205.

[61356] Henry, I.; Grandjouan, S.; Barichard, F.; Huerre-Jeanpierre, C.;

[61357] Junien, C.: Mitotic deletions of 11p15.5 in two different tumors

[61358] indicate that the CALCA locus is distal to the PTH locus.

Cytogenet.

[61359] Cell Genet. 50: 155-157, 1989.

[61360] 10206.

[61361] Hillyard, C. J.; Myers, C.; Abeyasekera, G.; Stevenson, J. C.;

[61362] Craig, R. K.; MacIntyre, I.: Katacalcin: a new plasma cal-
cium-lowering

[61363] hormone. Lancet I: 846-848, 1983.

[61364] 10207.

[61365] Hoovers, J. M. N.; Redeker, E.; Speleman, F.; Hoppener, J.
W.

[61366] M.; Bhola, S.; Bliek, J.; van Roy, N.; Leschot, N. J.; Wester-
veld,

[61367] A.; Mannens, M.: High-resolution chromosomal localiza-
tion of the

[61368] human calcitonin/CGRP/IAPP gene family members. Ge-
nomics 15: 525-529,

[61369] 1993.

[61370] 10208.

[61371] Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.;
Adema, G.

[61372] J.; Geurts van Kessel, A. H. M.; Lips, C. J. M.; Jansz, H. S.: A

[61373] third human CALC (pseudo)gene on chromosome 11. FEBS Lett. 233:

[61374] 57-63, 1988.

[61375] 10209.

[61376] Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Bakker, E.;

[61377] Pearson, P. L.; Geurts van Kessel, A. H. M.; Jansz, H. S.; Lips, C.

[61378] J. M.: Localization of the polymorphic human calcitonin gene on chromosome

[61379] 11. Hum. Genet. 66: 309-312, 1984.

[61380] 10210.

[61381] Jacobs, J. W.; Goodman, R. H.; Chin, W. W.; Dee, P. C.; Habener,

[61382] J. F.; Bell, N. H.; Potts, J. T., Jr.: Calcitonin messenger RNA encodes

[61383] multiple polypeptides in a single precursor. Science 213: 457-459,

[61384] 1981.

[61385] 10211.

[61386] Jonas, V.; Lin, C. R.; Kawashima, E.; Semon, D.; Swanson,

L. W.;

[61387] Mermod, J.-J.; Evans, R. M.; Rosenfeld, M. G.: Alternative RNA processing

[61388] events in human calcitonin/calcitonin gene-related peptide gene expression. Proc.

[61389] Nat. Acad. Sci. 82: 1994-1998, 1985.

[61390] 10212.

[61391] Kittur, S. D.; Hoppener, J. W. M.; Antonarakis, S. E.; Daniels,

[61392] J. D. J.; Meyers, D. A.; Maestri, N. E.; Jansen, M.; Korneluk, R.

[61393] G.; Nelkin, B. D.; Kazazian, H. H., Jr.: Linkage map of the short

[61394] arm of human chromosome 11: location of the genes for catalase, calcitonin,

[61395] and insulin-like growth factor II. Proc. Nat. Acad. Sci. 82: 5064-5067,

[61396] 1985.

[61397] 10213.

[61398] MacIntyre, I.; Hillyard, C. J.; Murphy, P. K.; Reynolds, J. J.;

[61399] Gaines-Das, R. E.; Craig, R. K.: A second plasma calcium-

lowering
[61400] peptide from the human calcitonin precursor. Nature 300: 460-462,
[61401] 1982.

[61402] 10214.

[61403] Mathe, A. A.; Agren, H.; Lindstrom, L.; Theodorsson, E.: Increased
[61404] concentration of calcitonin gene-related peptide in cerebrospinal
[61405] fluid of depressed patients: a possible trait marker of major depressive
[61406] disorder. Neurosci. Lett. 182: 138-142, 1994.

[61407] 10215.

[61408] Neher, R.; Riniker, B.; Rittel, W.; Zuber, H.: Thyrocalcitonin.

[61409] II. Struktur von alpha-Thyrocalcitonin. Helv. Chim. Acta 51: 917-924,
[61410] 1968.

[61411] 10216.

[61412] New, H. V.; Mudge, A. W.: Calcitonin gene-related peptide regulates
[61413] muscle acetylcholine receptor synthesis. Nature 323:

809-811, 1986.

[61414] 10217.

[61415] Przepiorka, D.; Baylin, S. B.; McBride, D. W.; Testa, J. R.; de

[61416] Bustros, A.; Nelkin, B. D.: The human calcitonin gene is located

[61417] on the short arm of chromosome 11. Biochem. Biophys. Res. Commun. 120:

[61418] 493-499, 1984.

[61419] 10218.

[61420] Rosenfeld, M. G.; Lin, C. R.; Amara, S. G.; Stolarsky, L.; Roos,

[61421] B. A.; Ong, E. S.; Evans, R. M.: Calcitonin mRNA polymor-phism: peptide

[61422] switching associated with alternative RNA splicing events. Proc.

[61423] Nat. Acad. Sci. 79: 1717-1721, 1982.

[61424] 10219.

[61425] Vora, S.; Seaman, C.; Durham, S.; Piomelli, S.: Isozymes of human

[61426] phosphofructokinase: identification and subunit structural characterization

[61427] of a new system. Proc. Nat. Acad. Sci. 77: 62-66, 1980.

[61428] 10220.

[61429] Mimmack, M. L.; Ryan, M.; Baba, H.; Navarro-Ruiz, J.; Iri-
tani,
[61430] S.; Faull, R. L. M.; McKenna, P. J.; Jones, P. B.; Arai, H.;
Starkey,
[61431] M.; Emson, P. C.; Bahn, S.: Gene expression analysis in
schizophrenia:
[61432] reproducible up-regulation of several members of the
apolipoprotein
[61433] L family located in a high-susceptibility locus for
schizophrenia
[61434] on chromosome 22. Proc. Nat. Acad. Sci. 99: 4680-4685,
2002.

[61435] 10221.

[61436] Schmauss, C.; McAllister, G.; Ohosone, Y.; Hardin, J. A.;
Lerner,
[61437] M. R.: A comparison of snRNP-associated Sm-
autoantigens: human N,
[61438] rat N and human B/B-prime. Nucleic Acids Res. 17:
1733-1743, 1989.

[61439] 10222.

[61440] Kovacs, G.; Kung, H.: Nonhomologous chromatid exchange in hereditary
[61441] and sporadic renal cell carcinomas. Proc. Nat. Acad. Sci. 88: 194-198,
[61442] 1991.

[61443] 10223.

[61444] Pompe, J. C.: Over Idiopathische hypertrophie van het hart. Nederl.
[61445] Tijdschr. Geneesk. 76: 304-312, 1932.

[61446] 10224.

[61447] Brunkow, M. E.; Gardner, J. C.; Van Ness, J.; Paeper, B. W.; Kovacevich,
[61448] B. R.; Proll, S.; Skonier, J. E.; Zhao, L.; Sabo, P. J.; Fu, Y.-H.;
[61449] Alisch, R. S.; Gillett, L.; Colbert, T.; Tacconi, P.; Galas, D.; Hamersma,
[61450] H.; Beighton, P.; Mulligan, J. T.: Bone dysplasia sclerosteosis results
[61451] from loss of the SOST gene product, a novel cystine knot-containing
[61452] protein. Am. J. Hum. Genet. 68: 577-589, 2001.

[61453] 10225.

[61454] Mansfield, E.; Chae, J. J.; Komarow, H. D.; Brotz, T. M.; Frucht,

[61455] D. M.; Aksentijevich, I.; Kastner, D. L.: The familial Mediterranean

[61456] fever protein, pyrin, associates with microtubules and colocalizes

[61457] with actin filaments. Blood 98: 851-859, 2001.

[61458] 10226.

[61459] Auer, I. A.; Schmidt, M. L.; Lee, V. M.-Y.; Curry, B.; Suzuki,

[61460] K.; Shin, R.-W.; Pentchev, P. G.; Carstea, E. D.; Tro-janowski, J.

[61461] Q.: Paired helical filament tau (PHFtau) in Niemann-Pick type C disease

[61462] is similar to PHFtau in Alzheimer's disease. Acta Neu-ropath. 90:

[61463] 547-551, 1995.

[61464] 10227.

[61465] Mules, E. H.; Dowling, C. E.; Petersen, M. B.; Kazazian, H. H.,

[61466] Jr.; Thomas, G. H.: A novel mutation in the invariant AG of the acceptor

[61467] splice site of intron 4 of the beta-hexosaminidase alpha-subunit gene

[61468] in two unrelated American black G(M2)-gangliosidosis (Tay-Sachs disease)

[61469] patients. Am. J. Hum. Genet. 48: 1181-1185, 1991.

[61470] 10228.

[61471] Montini, E.; Rugarli, E. I.; Van de Vosse, E.; Andolfi, G.; Mariani,

[61472] M.; Puca, A. A.; Consalez, G. G.; den Dunnen, J. T.; Ballabio, A.;

[61473] Franco, B.: A novel human serine-threonine phosphatase related to

[61474] the Drosophila retinal degeneration C (rdgC) gene is selectively expressed

[61475] in sensory neurons of neural crest origin. Hum. Molec. Genet. 6:

[61476] 1137-1145, 1997.

[61477] 10229.

[61478] Nishino, I.; Fu, J.; Tanji, K.; Yamada, T.; Shimojo, S.; Koori,

[61479] T.; Mora, M.; Riggs, J. E.; Oh, S. J.; Koga, Y.; Sue, C. M.; Yamamoto,

[61480] A.; Murakami, N.; Shanske, S.; Byrne, E.; Bonilla, E.; Non-aka, I.;

[61481] DiMauro, S.; Hirano, M.: Primary LAMP-2 deficiency causes X-linked

[61482] vacuolar cardiomyopathy and myopathy (Danon disease). Nature 406:

[61483] 906-910, 2000.

[61484] 10230.

[61485] Zonana, J.; Gault, J.; Davies, K. J. P.; Jones, M.; Browne, D.;

[61486] Litt, M.; Brockdorff, N.; Rastan, S.; Clarke, A.; Thomas, N. S. T.

[61487] : Detection of a molecular deletion at the DXS732 locus in a patient

[61488] with X-linked hypohidrotic ectodermal dysplasia (EDA), with the identification

[61489] of a unique junctional fragment. Am. J. Hum. Genet. 52: 78-84, 1993.

[61490] 10231.

[61491] Ueki, Y.; Naito, I.; Oohashi, T.; Sugimoto, M.; Seki, T.; Yoshioka,

[61492] H.; Sado, Y.; Sato, H.; Sawai, T.; Sasaki, F.; Matsuoka, M.; Fukuda,

[61493] S.; Ninomiya, Y.: Topoisomerase I and II consensus sequences in a
[61494] 17-kb deletion junction of the COL4A5 and COL4A6 genes and immunohistochemical
[61495] analysis of esophageal leiomyomatosis associated with Alport syndrome. Am.
[61496] J. Hum. Genet. 62: 253-261, 1998.

[61497] 10232.

[61498] Rowe, R. C.; Schroeder, M.-L.; Faiman, C.: Testosterone-induced
[61499] fertility in a patient with previously untreated Kallmann's syndrome. Fertil.
[61500] Steril. 40: 400-401, 1983.

[61501] 10233.

[61502] Hata, A.; Setoyama, C.; Shimada, K.; Takeda, E.; Kuroda, Y.; Akaboshi,
[61503] I.; Matsuda, I.: Ornithine transcarbamylase deficiency resulting
[61504] from a C-to-T substitution in exon 5 of the ornithine transcarbamylase
[61505] gene. Am. J. Hum. Genet. 45: 123-127, 1989.

[61506] 10234.

[61507] Huopaniemi, L.; Fellman, J.; Rantala, A.; Eriksson, A.; Forsius,

[61508] H.; de la Chapelle, A.; Alitalo, T.: Skewed secondary sex ratio in

[61509] the offspring of carriers of the 214G-A mutation of the RS1 gene. Ann.

[61510] Hum. Genet. 63: 521-533, 1999.

[61511] 10235.

[61512] Johnson, M. D.; Tho, S. P. T.; Behzadian, A.; McDonough, P. G.

[61513] : Molecular scanning of Yq11 (interval 6) in men with Sertoli-cell-only

[61514] syndrome. Am. J. Obstet. Gynec. 161: 1732-1737, 1989.

[61515] 10236.

[61516] Bolger, G.; Michaeli, T.; Martins, T.; St. John, T.; Steiner, B.;

[61517] Rodgers, L.; Riggs, M.; Wigler, M.; Ferguson, K.: A family of human

[61518] phosphodiesterases homologous to the dunce learning and memory gene

[61519] product of Drosophila melanogaster are potential targets

for antidepressant
[61520] drugs. Molec. Cell. Biol. 13: 6558-6571, 1993.

[61521] 10237.

[61522] Welcsh, P. L.; King, M.-C.: BRCA1 and BRCA2 and the genetics
[61523] of breast and ovarian cancer. Hum. Molec. Genet. 10: 705-713, 2001.
[61524] 10238.

[61525] Argyrokastritis, A.; Kamakari, S.; Kapsetaki, M.; Kritis, A.; Talianidis,
[61526] I.; Moschonas, N. K.: Human hepatocyte nuclear factor-4 (hHNF-4)
[61527] gene maps to 20q12-q13.1 between PLCG1 and D20S17. Hum. Genet. 99:
[61528] 233-236, 1997.

[61529] 10239.

[61530] Weaver, D. R.; Rivkees, S. A.; Carlson, L. L.; Reppert, S. M.:

[61531] Localization of melatonin receptors in mammalian brain.In: Klein,
[61532] D. C.; Moore, R. Y.; Reppert, S. M.: Suprachiasmatic Nucleus: The

[61533] Mind's Clock. New York: Oxford Press (pub.) 1991.

[61534] 10240.

[61535] Pickard, R. T.; Strifler, B. A.; Kramer, R. M.; Sharp, J. D.:

[61536] Molecular cloning of two new human paralogs of 85-kDa cytosolic phospholipase

[61537] A2. J. Biol. Chem. 274: 8823-8831, 1999.

[61538] 10241.

[61539] Lopes, J.; LeGuern, E.; Gouider, R.; Tardieu, S.; Abbas, N.; Birouk,

[61540] N.; Gugenheim, M.; Bouche, P.; Agid, Y.; Brice, A.; French CMT Collaborative

[61541] Research Group: Recombination hot spot in a 3.2-kb region of the

[61542] Charcot-Marie-Tooth type 1A repeat sequences: new tools for molecular

[61543] diagnosis of hereditary neuropathy with liability to pressure palsies

[61544] and of Charcot-Marie-Tooth type 1A. Am. J. Hum. Genet. 58: 1223-1230,

[61545] 1996.

[61546] 10242.

[61547] Hegele, R. A.; Cao, H.; Harris, S. B.; Zinman, B.; Hanley, A.

[61548] J. G.; Anderson, C. M.: Peroxisome proliferator-activated receptor-gamma2

[61549] P12A and type 2 diabetes in Canadian Oji-Cree. J. Clin. Endocr. Metab. 85:

[61550] 2014-2019, 2000.

[61551] 10243.

[61552] Chua, A. O.; Chizzonite, R.; Desai, B. B.; Truitt, T. P.; Nunes,

[61553] P.; Minetti, L. J.; Warrier, R. R.; Presky, D. H.; Levine, J. F.;

[61554] Gately, M. K.; Gubler, U.: Expression cloning of a human IL-12 receptor

[61555] component: a new member of the cytokine receptor superfamily with

[61556] strong homology to gp130. J. Immun. 153: 128-136, 1994.

[61557] 10244.

[61558] Warburg, M.; Tommerup, N.; Vestermark, S.; Parving, A.; Weismann,

[61559] K.; Russell, B.; Thomsen, H. K.: The Yemenite deaf-blind hypopigmentation

[61560] syndrome: a new oculo-dermato-auditory syndrome. Ophthal. Paediat.

[61561] Genet. 11: 201-207, 1990.

[61562] 10245.

[61563] Wedemeyer, N.; Peoples, R.; Himmelbauer, H.; Lehrach, H.; Francke,

[61564] U.; Wanker, E. E.: Localization of the human HIP1 gene close to the

[61565] elastin (ELN) locus on 7q11.23. Genomics 46: 313-315, 1997.

[61566] 10246.

[61567] Zetterstrom, R. H.; Solomin, L.; Jansson, L.; Hoffer, B. J.; Olson,

[61568] L.; Perlmann, T.: Dopamine neuron agenesis in Nurr1-deficient mice. Science 276:

[61569] 248-250, 1997.

[61570] 10247.

[61571] Ye, H.; Kelly, T. F.; Samadani, U.; Lim, L.; Rubio, S.; Overdier,

[61572] D. G.; Roebuck, K. A.; Costa, R. H.: Hepatic nuclear factor 3/fork

[61573] head homolog 11 is expressed in proliferating epithelial and mesenchymal
[61574] cells of embryonic and adult tissues. Molec. Cell. Biol. 17: 1626-1641,
[61575] 1997.

[61576] 10248.

[61577] Chu, C.-S.; Trapnell, B. C.; Murtagh, J. J., Jr.; Moss, J.; Dalemans,
[61578] W.; Jallat, S.; Mercenier, A.; Pavirani, A.; Lecocq, J.-P.; Cutting,
[61579] G. R.; Guggino, W. B.; Crystal, R. G.: Variable detection of exon
[61580] 9 coding sequences in cystic fibrosis transmembrane conductance regulator
[61581] gene mRNA transcripts in normal bronchial epithelium. EMBO J. 10:
[61582] 1355-1363, 1991.

[61583] 10249.

[61584] Bryan, J.; Kane, R. E.: Actin gelation in sea urchin egg extracts. Methods
[61585] Cell Biol. 25: 175-199, 1982.

[61586] 10250.

[61587] Bittner, R. E.; Anderson, L. V. B.; Burkhardt, E.; Bashir, R.;

[61588] Vafiadaki, E.; Ivanova, S.; Raffelsberger, T.; Maerk, I.;
Hoger, H.;

[61589] Jung, M.; Karbasiyan, M.; Storch, M.; Lassmann, H.; Moss,
J. A.; Davison,

[61590] K.; Harrison, R.; Bushby, K. M. D.; Reis, A.: Dysferlin dele-
tion

[61591] in SJL mice (SJL-Dysf) defines a natural model for limb gir-
dle muscular

[61592] dystrophy 2B. (Letter) Nature Genet. 23: 141-142, 1999.

[61593] 10251.

[61594] Wilgoss, A.; Leigh, I. M.; Barnes, M. R.; Dopping-Hepen-
stal, P.;

[61595] Eady, R. A. J.; Walter, J. M.; Kennedy, C. T. C.; Kelsell, D. P.:

[61596] Identification of a novel mutation R42P in the gap junction
protein

[61597] beta-3 associated with autosomal dominant erythrokera-
toderma variabilis. J.

[61598] Invest. Derm. 113: 1119-1122, 1999.

[61599] 10252.

[61600] Lamhonwah, A.-M.; Olpin, S. E.; Pollitt, R. J.; Vianey-Saban, C.;
[61601] Divry, P.; Guffon, N.; Besley, G. T. N.; Onizuka, R.; De Meirleir,
[61602] L. J.; Cvitanovic-Sojat, L.; Baric, I.; Dionisi-Vici, C.; Fumic, K.;
[61603] Maradin, M.; Tein, I.: Novel OCTN2 mutations: no genotype-phenotype
[61604] correlations: early carnitine therapy prevents cardiomyopathy. Am.
[61605] J. Med. Genet. 111: 271-284, 2002.

[61606] 10253.

[61607] Koi, M.; Johnson, L. A.; Kalikin, L. M.; Little, P. F. R.; Nakamura,
[61608] Y.; Feinberg, A. P.: Tumor cell growth arrest caused by subchromosomal
[61609] transferable DNA fragments from chromosome 11. Science 260: 361-364,
[61610] 1993.

[61611] 10254.

[61612] Yamada, T.; Ochi, H.; Hara, H.; Yoshimura, T.; Kobayashi T.:

[61613] A skeletal muscle sodium channel mutation in a Japanese family with
[61614] paramyotonia congenita. J. Neurol. Sci. 133: 192-193, 1995.
[61615] 10255.

[61616] Makoff, A.; Pilling, C.; Harrington, K.; Emson, P.: Human metabotropic
[61617] glutamate receptor type 7: molecular cloning and mRNA distribution
[61618] in the CNS. Molec. Brain Res. 40: 165-170, 1996.

[61619] 10256.

[61620] Esterbauer, H.; Oberkofler, H.; Krempler, F.; Patsch, W.: Human
[61621] peroxisome proliferator activated receptor gamma coactivator 1 (PPARGC1)
[61622] gene: cDNA sequence, genomic organization, chromosomal localization,
[61623] and tissue expression. Genomics 62: 98-102, 1999.

[61624] 10257.

[61625] Tsuchida, K.; Arai, K. Y.; Kuramoto, Y.; Yamakawa, N.; Hasegawa,

[61626] Y.; Sugino, H.: Identification and characterization of a novel follistatin-like
[61627] protein as a binding protein for the TGF-beta family. J. Biol. Chem. 275:
[61628] 40788-40796, 2000.

[61629] 10258.

[61630] Takatsu, H.; Yoshino, K.; Nakayama, K.: Adaptor gamma-ear homology
[61631] domain conserved in gamma-adaptin and GGA proteins that interact with
[61632] gamma-synergin. Biochem. Biophys. Res. Commun. 271: 719-725, 2000.
[61633] 10259.

[61634] Burbelo, P. D.; Snow, D. M.; Bahou, W.; Spiegel, S.: MSE55, a
[61635] Cdc42 effector protein, induces long cellular extensions in fibroblasts. Proc.
[61636] Nat. Acad. Sci. 96: 9083-9088, 1999.

[61637] 10260.

[61638] Tsavaler, L.; Shapero, M. H.; Morkowski, S.; Laus, R.: Trp-p8,

[61639] a novel prostate-specific gene, is up-regulated in prostate cancer

[61640] and other malignancies and shares high homology with transient receptor

[61641] potential calcium channel proteins. Cancer Res. 61: 3760-3769, 2001.

[61642] 10261.

[61643] Tomczak, J.; Boogen, C.; Grebner, E. E.: Distribution of a pseudodeficiency

[61644] allele among Tay-Sachs carriers. (Letter) Am. J. Hum. Genet. 53:

[61645] 537-539, 1993.

[61646] 10262.

[61647] Rafi, M. A.; Luzi, P.; Chen, Y. Q.; Wenger, D. A.: A large deletion

[61648] together with a point mutation in the GALC gene is a common mutant

[61649] allele in patients with infantile Krabbe disease. Hum. Molec. Genet. 4:

[61650] 1285-1289, 1995.

[61651] 10263.

[61652] Hazarika, P.; Dham, N.; Patel, P.; Cho, M.; Weidner, D.; Goldsmith,

[61653] L.; Duvic, M.: Flotillin 2 is distinct from epidermal surface antigen

[61654] (ESA) and is associated with filopodia formation. J. Cell. Biochem. 75:

[61655] 147-159, 1999.

[61656] 10264.

[61657] Satoh, S.; Daigo, Y.; Furukawa, Y.; Kato, T.; Miwa, N.; Nishiwaki,

[61658] T.; Kawasoe, T.; Ishiguro, H.; Fujita, M.; Tokino, T.; Sasaki, Y.;

[61659] Imaoka, S.; Murata, M.; Shimano, T.; Yamaoka, Y.; Nakamura, Y.: AXIN1

[61660] mutations in hepatocellular carcinomas, and growth suppression in

[61661] cancer cells by virus-mediated transfer of AXIN1. Nature Genet. 24:

[61662] 245-250, 2000.

[61663] 10265.

[61664] Zeng, L.; Fagotto, F.; Zhang, T.; Hsu, W.; Vasicek, T. J.; Perry,

[61665] W. L., III; Lee, J. J.; Tilghman, S. M.; Gumbiner, B. M.; Costantini,
[61666] F.: The mouse fused locus encodes axin, an inhibitor of the Wnt signaling
[61667] pathway that regulates embryonic axis formation. Cell 90: 181-192,
[61668] 1997.

[61669] 10266.

[61670] Andreev, J.; Simon, J.-P.; Sabatini, D. D.; Kam, J.; Plowman, G.;
[61671] Randazzo, P. A.; Schlessinger, J.: Identification of a new Pyk2 target
[61672] protein with Arf-GAP activity. Molec. Cell. Biol. 19: 2338-2350,
[61673] 1999.

[61674] 10267.

[61675] Burden, S.; Yarden, Y.: Neuregulins and their receptors: a versatile
[61676] signaling module in organogenesis and oncogenesis. Neuron 18: 847-855,
[61677] 1997.

[61678] 10268.

[61679] Busfield, S. J.; Michnick, D. A.; Chickering, T. W.; Revett, T.

[61680] L.; Ma, J.; Woolf, E. A.; Comrack, C. A.; Dussault, B. J.;
Woolf,

[61681] J.; Goodearl, A. D. J.; Gearing, D. P.: Characterization of a
neuregulin-related

[61682] gene, Don-1, that is highly expressed in restricted regions
of the

[61683] cerebellum and hippocampus. Molec. Cell. Biol. 17:
4007-4014, 1997.

[61684] 10269.

[61685] Carraway, K. L., III; Weber, J. L.; Unger, M. J.; Ledesma, J.;

[61686] Yu, N.; Gassmann, M.; Lai, C.: Neuregulin-2, a new ligand
of ErbB3/ErbB4-receptor

[61687] tyrosine kinases. Nature 387: 512-516, 1997.

[61688] 10270.

[61689] Chang, H.; Riese, D. J., II; Gilbert, W.; Stern, D. F.; McMa-
han,

[61690] U. J.: Ligands for ErbB-family receptors encoded by a
neuregulin-like

[61691] gene. Nature 387: 509-512, 1997.

[61692] 10271.

[61693] Ring, H. Z.; Chang, H.; Guilbot, A.; Brice, A.; LeGuern, E.; Francke,

[61694] U.: The human neuregulin-2 (NRG2) gene: cloning, mapping and evaluation

[61695] as a candidate for the autosomal recessive form of Charcot-Marie-Tooth

[61696] disease linked to 5q. Hum. Genet. 104: 326-332, 1999.

[61697] 10272.

[61698] Yamada, K.; Ichino, N.; Nishii, K.; Sawada, H.; Higashiyama, S.;

[61699] Ishiguro, H.; Nagatsu, T.: Characterization of the human NTAK gene

[61700] structure and distribution of the isoforms for rat NTAK mRNA. Gene 255:

[61701] 15-24, 2000.

[61702] 10273.

[61703] Yokoyama, M.; Nishi, Y.; Yoshii, J.; Okubo, K.; Matsubara, K.:

[61704] Identification and cloning of neuroblastoma-specific and nerve tissue-specific

[61705] genes through compiled expression profiles. DNA Res. 3: 311-320,

[61706] 1996.

[61707] 10274.

[61708] Hu, R.-J.; Lee, M. P.; Connors, T. D.; Johnson, L. A.; Burn, T.

[61709] C.; Su, K.; Landes, G. M.; Feinberg, A. P.: A 2.5-Mb tran-script map

[61710] of a tumor-suppressing subchromosomal transferable fragment from 11p15.5,

[61711] and isolation and sequence analysis of three novel genes. Genomics 46:

[61712] 9-17, 1997.

[61713] 10275.

[61714] Stephan, D.; Bon, C.; Holzwarth, J. A.; Galvan, M.; Pruss, R.

[61715] M.: Human metabotropic glutamate receptor 1: mRNA distribution, chromosome

[61716] localization and functional expression of two splice vari-ants. Neuropharmacology 35:

[61717] 1649-1660, 1996.

[61718] 10276.

[61719] Chen, M. S.; Huber, A. B.; van der Haar, M. E.; Frank, M.; Schnell,

[61720] L.; Spillmann, A. A.; Christ, F.; Schwab, M. E.: Nogo-A is a myelin-associated

[61721] neurite outgrowth inhibitor and an antigen for monoclonal antibody

[61722] IN-1. Nature 403: 434-439, 2000.

[61723] 10277.

[61724] Fournier, A. E.; GrandPre, T.; Strittmatter, S. M.: Identification

[61725] of a receptor mediating Nogo-66 inhibition of axonal regeneration. Nature 409:

[61726] 341-346, 2001.

[61727] 10278.

[61728] GrandPre, T.; Li, S.; Strittmatter, S. M.: Nogo-66 receptor antagonist

[61729] peptide promotes axonal regeneration. Nature 417: 547-551, 2002.

[61730] 10279.

[61731] GrandPre, T.; Nakamura, F.; Vartanian, T.; Strittmatter, S. M.

[61732] : Identification of the Nogo inhibitor of axon regeneration as a reticulon
[61733] protein. Nature 403: 439-444, 2000.

[61734] 10280.

[61735] Prinjha, R.; Moore, S. E.; Vinson, M.; Blake, S.; Morrow, R.; Christie,
[61736] G.; Michalovich, D.; Simmons, D. L.; Walsh, F. S.: Inhibitor of neurite
[61737] outgrowth in humans. (Letter) Nature 403: 383-384, 2000.
[61738] 10281.

[61739] Spillmann, A. A.; Bandtlow, C. E.; Lottspeich, F.; Keller, F.;

[61740] Schwab, M. E.: Identification and characterization of a bovine neurite
[61741] growth inhibitor (bNI-220). J. Biol. Chem. 273: 19283-19293, 1998.
[61742] 10282.

[61743] Yang, J.; Yu, L.; Bi, A. D.; Zhao, S.-Y.: Assignment of the human
[61744] reticulon 4 gene (RTN4) to chromosome 2p14-2p13 by radiation hybrid

[61745] mapping. Cytogenet. Cell Genet. 88: 101-102, 2000.

[61746] 10283.

[61747] Korner, C. G.; Wahle, E.: Poly(A) tail shortening by a mam-
malian
[61748] poly(A)-specific 3-prime-exoribonuclease. J. Biol. Chem.
272: 10448-10456,
[61749] 1997.

[61750] 10284.

[61751] Korner, C. G.; Wormington, M.; Muckenthaler, M.; Schnei-
der, S.;
[61752] Dehlin, E.; Wahle, E.: The deadenylating nuclease (DAN) is
involved
[61753] in poly(A) tail removal during the meiotic maturation of
Xenopus oocytes. EMBO
[61754] J. 17: 5427-5437, 1998.

[61755] 10285.

[61756] Dear, T. N.; Moller, A.; Boehm, T.: CAPN11: a calpain with
high
[61757] mRNA levels in testis and located on chromosome 6. Ge-
nomics 59:
[61758] 243-247, 1999.

[61759] 10286.

[61760] Ai, Y.; Jenkins, N. A.; Copeland, N. G.; Gilbert, D. J.; Bergsma,

[61761] D. J.; Stambolian, D.: Mouse galactokinase: isolation, characterization,

[61762] and location on chromosome 11. Genome Res. 5: 53-59, 1995.

[61763] 10287.

[61764] Ai, Y.; Zheng, Z.; O'Brien-Jenkins, A.; Bernard, D. J.; Wyn-shaw-Boris,

[61765] T.; Ning, C.; Reynolds, R.; Segal, S.; Huang, K.; Stambolian, D.:

[61766] A mouse model of galactose-induced cataracts. Hum. Molec. Genet. 9:

[61767] 1821-1827, 2000.

[61768] 10288.

[61769] Bergsma, D. J.; Ai, Y.; Skach, W. R.; Nesburn, K.; Anoia, E.; Van

[61770] Horn, S.; Stambolian, D.: Fine structure of the human galactokinase

[61771] GALK1 gene. Genome Res. 6: 980-985, 1996.

[61772] 10289.

[61773] Croce, C. M.; Huebner, K.; Koprowski, H.: Chromosome assignment

[61774] of the T-antigen gene of simian virus 40 in African green monkey cells

[61775] transformed by adeno 7-SV 40 hybrid. Proc. Nat. Acad. Sci. 71: 4116-4119,

[61776] 1974.

[61777] 10290.

[61778] de Jonge, A. J. R.; de Smit, S.; Kroos, M. A.; Reuser, A. J. J.

[61779] : Cotransfer of syntenic human genes into mouse cells using isolated

[61780] metaphase chromosomes or cellular DNA. Hum. Genet. 69: 32-38, 1985.

[61781] 10291.

[61782] Elsevier, S. M.; Kucherlapati, R. S.; Nichols, E. A.; Willecke,

[61783] K.; Creagan, R. P.; Giles, R. E.; McDougall, J. K.; Ruddle, F. H.

[61784] : Assignment and regional localization of a gene coding for galactokinase

[61785] to human chromosome 17q21-22. Birth Defects Orig. Art.

Ser. 11(3):
[61786] 117-119, 1975. Note: Alternate: Cytogenet. Cell Genet. 14: 287-289,
[61787] 1975.

[61788] 10292.

[61789] Klobutcher, L. A.; Ruddle, F. H.: Phenotype stabilisation and
[61790] integration of transferred material in chromosome-mediated gene transfer. Natur
[61791] e 280: 657-660, 1979.

[61792] 10293.

[61793] Kolosha, V.; Anoia, E.; de Cespedes, C.; Gitzelmann, R.; Shih,
[61794] L.; Casco, T.; Saborio, M.; Trejos, R.; Buist, N.; Tedesco, T.; Skach,
[61795] W.; Mitelmann, O.; Ledee, D.; Huang, K.; Stambolian, D.: Novel mutations
[61796] in 13 probands with galactokinase deficiency. Hum. Mutat. 15: 447-453,
[61797] 2000.

[61798] 10294.

[61799] Okajima, K.; Yazaki, M.; Wada, Y.: Thymidine kinase activity

[61800] in individuals with galactokinase deficiency. (Letter) Am. J. Hum.

[61801] Genet. 41: 503-504, 1987.

[61802] 10295.

[61803] Okano, Y.; Asada, M.; Fujimoto, A.; Ohtake, A.; Murayama, K.;

[61804] Hsiao, K.-J.; Choeh, K.; Yang, Y.; Cao, Q.; Reichardt, J. K. V.; Niihira,

[61805] S.; Imamura, T.; Yamano, T.: A genetic factor for age-related cataract:

[61806] identification and characterization of a novel galactokinase variant,

[61807] 'Osaka,' in Asians. Am. J. Hum. Genet. 68: 1036-1042, 2001.

[61808] 10296.

[61809] Ruddle, F. H.: Personal Communication. New Haven, Conn. 5/4/1982.

[61810] 10297.

[61811] Schoen, R. C.; Cox, S. H.; Wagner, R. P.: Thymidine-kinase activity

[61812] of cultured cells from individuals with inherited galactoki-
nase deficiency. Am.
[61813] J. Hum. Genet. 36: 815-822, 1984.

[61814] 10298.

[61815] O'Mahoney, J. V.; Guven, K. L.; Lin, J.; Joya, J. E.; Robinson,

[61816] C. S.; Wade, R. P.; Hardeman, E. C.: Identification of a
novel slow-muscle-fiber
[61817] enhancer binding protein, MusTRD1. Molec. Cell. Biol. 18:
6641-6652,
[61818] 1998.

[61819] 10299.

[61820] Osborne, L. R.; Campbell, T.; Daradich, A.; Scherer, S. W.;
Tsui,
[61821] L.-C.: Identification of a putative transcription factor gene
(WBSCR11)
[61822] that is commonly deleted in Williams-Beuren syndrome.
Genomics 57:
[61823] 279-284, 1999.

[61824] 10300.

[61825] Tassabehji, M.; Carette, M.; Wilmot, C.; Donnai, D.; Read,
A. P.;

[61826] Metcalfe, K.: A transcription factor involved in skeletal muscle

[61827] gene expression is deleted in patients with Williams syndrome. Europ.

[61828] J. Hum. Genet. 7: 737-747, 1999.

[61829] 10301.

[61830] Schmidt, A.; Wolde, M.; Thiele, C.; Fest, W.; Kratzin, H.; Podtelejnikov,

[61831] A. V.; Witke, W.; Huttner, W. B.; Soling, H.-D.: Endophilin I mediates

[61832] synaptic vesicle formation by transfer of arachidonate to lysophosphatidic

[61833] acid. Nature 401: 133-141, 1999.

[61834] 10302.

[61835] Bahr, A.; Hankeln, T.; Fiedler, T.; Hegemann, J.; Schmidt, E. R.

[61836] : Molecular analysis of METTL1, a novel human methyltransferase-like

[61837] gene with a high degree of phylogenetic conservation. Genomics 57:

[61838] 424-428, 1999.

[61839] 10303.

[61840] Aiba, A.; Chen, C.; Herrup, K.; Rosenmund, C.; Stevens, C. F.;

[61841] Tonegawa, S.: Reduced hippocampal long-term potentiation and context-specific

[61842] deficit in associative learning in mGluR1 mutant mice. Cell 79:

[61843] 365-375, 1994.

[61844] 10304.

[61845] Aiba, A.; Kano, M.; Chen, C.; Stanton, M. E.; Fox, G. D.; Herrup,

[61846] K.; Zwingman, T. A.; Tonegawa, S.: Deficient cerebellar long-term

[61847] depression and impaired motor learning in mGluR1 mutant mice. Cell 79:

[61848] 377-388, 1994.

[61849] 10305.

[61850] Conquet, F.; Bashir, Z. I.; Davies, C. H.; Daniel, H.; Ferraguti,

[61851] F.; Bordi, F.; Franz-Bacon, K.; Reggiani, A.; Matarese, V.; Conde,

[61852] F.; Collingridge, G. L.; Crepel, F.: Motor deficit and impair-
ment
[61853] of synaptic plasticity in mice lacking mGluR1. Nature 372:
237-243,
[61854] 1994.

[61855] 10306.

[61856] Ganesh, S.; Amano, K.; Yamakawa, K.: Assignment of the
gene GRM1
[61857] coding for metabotropic glutamate receptor 1 to human
chromosome band
[61858] 6q24 by in situ hybridization. Cytogenet. Cell Genet. 88:
314-315,
[61859] 2000.

[61860] 10307.

[61861] Ichise, T.; Kano, M.; Hashimoto, K.; Yanagihara, D.; Nakao,
K.;
[61862] Shigamoto, R.; Katsuki, M.; Alba, A.: mGluR1 in cerebellar
Purkinje
[61863] cells essential for long-term depression, synapse elimina-
tion, and
[61864] motor coordination. Science 288: 1832-1835, 2000.

[61865] 10308.

[61866] Kunishima, N.; Shimada, Y.; Tsuji, Y.; Sato, T.; Yamamoto, M.;

[61867] Kumasaka, T.; Nakanishi, S.; Jingami, H.; Morikawa, K.: Structural

[61868] basis of glutamate recognition by a dimeric metabotropic glutamate

[61869] receptor. Nature 407: 971-977, 2000.

[61870] 10309.

[61871] Okamoto, T.; Sekiyama, N.; Otsu, M.; Shimada, Y.; Sato, A.; Nakanishi,

[61872] S.; Jingami, H.: Expression and purification of the extra-cellular

[61873] ligand binding region of metabotropic glutamate receptor subtype 1. J.

[61874] Biol. Chem. 273: 13089-13096, 1998.

[61875] 10310.

[61876] Smitt, P. S.; Kinoshita, A.; De Leeuw, B.; Moll, W.; Coes-mans,

[61877] M.; Jaarsma, D.; Henzen-Logmans, S.; Vecht, C.; De Zeeuw, C.; Sekiyama,

[61878] N.; Nakanishi, S.; Shigemoto, R.: Paraneoplastic cerebellar ataxia
[61879] due to autoantibodies against a glutamate receptor. New Eng. J. Med. 342:
[61880] 21-27, 2000.

[61881] 10311.

[61882] Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hiro-sawa, M.;
[61883] Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction
[61884] of the coding sequences of unidentified human genes. XII. The complete
[61885] sequences of 100 new cDNA clones from brain which code for large proteins
[61886] in vitro. DNA Res. 5: 355-364, 1998.

[61887] 10312.

[61888] Chadwick, B. P.; Kidd, T.; Sgouros, J.; Ish-Horowicz, D.; Frischauf,
[61889] A.-M.: Cloning, mapping and expression of UBL3, a novel ubiquitin-like
[61890] gene. Gene 233: 189-195, 1999.

[61891] 10313.

[61892] Hodges, M.; Tissot, C.; Freemont, P. S. Protein regulation:

[61893] tag wrestling with relatives of ubiquitin. Curr. Biol. 8:
R749-R752,

[61894] 1998.

[61895] 10314.

[61896] Tanaka, H.; Arakawa, H.; Yamaguchi, T.; Shiraishi, K.;
Fukuda,

[61897] S.; Matsui, K.; Takei, Y.; Nakamura, Y.: A ribonucleotide
reductase

[61898] gene involved in a p53-dependent cell-cycle checkpoint
for DNA damage. Nature 404:

[61899] 42-49, 2000.

[61900] 10315.

[61901] Bora, R. S.; Kanamori, A.; Hirabayashi, Y.: Assignment of a
putative

[61902] acetyl-CoA transporter gene (Acatn) to mouse chromo-
some band 3E1-E3

[61903] by in situ hybridization. Cytogenet. Cell Genet. 83: 78-79,
1998.

[61904] 10316.

[61905] Veitia, R. A.; Ottolenghi, C.; Bissery, M.-C.; Fellous, A.: A

[61906] novel human gene, encoding a potential membrane pro-
tein conserved

[61907] from yeast to man, is strongly expressed in testis and
cancer cell

[61908] lines. Cytogenet. Cell Genet. 85: 217-220, 1999.

[61909] 10317.

[61910] Andree, B.; Hillemann, T.; Kessler-Ieckson, G.; Schmitt-
John, T.;

[61911] Jockusch, H.; Arnold, H.-H.; Brand, T.: Isolation and char-
acterization

[61912] of the novel Popeye gene family expressed in skeletal
muscle and heart. Dev.

[61913] Biol. 223: 371-382, 2000.

[61914] 10318.

[61915] DeYoung, K. L.; Ray, M. E.; Su, Y. A.; Anzick, S. L.; John-
stone,

[61916] R. W.; Trapani, J. A.; Meltzer, P. S.; Trent, J. M.: Cloning a
novel

[61917] member of the human interferon-inducible gene family
associated with

[61918] control of tumorigenicity in a model of human melanoma. Oncogene 15:
[61919] 453-457, 1997.

[61920] 10319.

[61921] Kirikoshi, H.; Sagara, N.; Koike, J.; Tanaka, K.; Sekihara, H.;

[61922] Hirai, M.; Katoh, M.: Molecular cloning and characterization of human
[61923] frizzled-4 on chromosome 11q14-q21. Biochem. Biophys. Res. Commun. 264:
[61924] 955-961, 1999.

[61925] 10320.

[61926] Kowal, R. C.; Jolsin, J. M.; Olson, E. N.; Schultz, R. A.: Assignment
[61927] of fibulin-5 (FBLN5) to human chromosome 14q31 by in situ hybridization
[61928] and radiation hybrid mapping. Cytogenet. Cell Genet. 87: 2-3, 1999.
[61929] 10321.

[61930] Nakamura, T.; Lozano, P. R.; Ikeda, Y.; Iwanaga, Y.; Hinek, A.;
[61931] Minamisawa, S.; Cheng, C.-F.; Kobuke, K.; Dalton, N.;

Takada, Y.;

[61932] Tashiro, K.; Ross, J., Jr.; Honjo, T.; Chien, K. R.: Fibulin-5/DANCE

[61933] is essential for elastogenesis in vivo. Nature 415: 171-175, 2002.

[61934] 10322.

[61935] Nakamura, T.; Ruiz-Lozano, P.; Lindner, V.; Yabe, D.; Taniwaki,

[61936] M.; Furukawa, Y.; Kobuke, K.; Tashiro, K.; Lu, Z.; Andon, N. L.; Schaub,

[61937] R.; Matsumori, A.; Sasayama, S.; Chien, K. R.; Honjo, T.: DANCE,

[61938] a novel secreted RGD protein expressed in developing, atherosclerotic,

[61939] and balloon-injured arteries. J. Biol. Chem. 274: 22476-22483, 1999.

[61940] 10323.

[61941] Yanagisawa, H.; Davis, E. C.; Starcher, B. C.; Ouchi, T.; Yanagisawa,

[61942] M.; Richardson, J. A.; Olson, E. N.: Fibulin-5 is an elastin-binding

[61943] protein essential for elastic fibre development in vivo. Na-

ture 415:

[61944] 168-171, 2002.

[61945] 10324.

[61946] Robitaille, J.; MacDonald, M. L. E.; Kaykas, A.; Sheldahl, L. C.;

[61947] Zeisler, J.; Dube, M.-P.; Zhang, L.-H.; Singaraja, R. R.; Guernsey,

[61948] D. L.; Zhang, B.; Siebert, L. F.; Hoskin-Mott, A.; Trese, M. T.; Pimstone,

[61949] S. N.; Shastry, B. S.; Moon, R. T.; Hayden, M. R.; Goldberg, Y. P.;

[61950] Samuels, M. E.: Mutant frizzled-4 disrupts retinal angio- genesis in

[61951] familial exudative vitreoretinopathy. Nature Genet. 32: 326-330,

[61952] 2002.

[61953] 10325.

[61954] Banfi, S; . Bassi, M. T.; Andolfi, G.; Marchitiello, A.; Zan- otta,

[61955] S.; Ballabio, A.; Casari, G.; Franco, B.: Identification and characterization

[61956] of AFG3L2, a novel paraplegin-related gene. Genomics 59:

51-58,

[61957] 1999.

[61958] 10326.

[61959] Dong, F.; Feldmesser, M.; Casadevall, A.; Rubin, C. S.: Molecular

[61960] characterization of a cDNA that encodes six isoforms of a novel murine

[61961] A kinase anchor protein. J. Biol. Chem. 273: 6533-6541, 1998.

[61962] 10327.

[61963] Dent, A. L.; Yewdell, J.; Puvion-Dutilleul, F.; Koken, M. H.; de

[61964] The, H.; Staudt, L. M.: LYSP100 associated nuclear do-

[61965] mains (LANDs): description of a new class of subnuclear structures and their relationship

[61966] to PML nuclear bodies. Blood 88: 1423-1426, 1996.

[61967] 10328.

[61968] Seeler, J. S.; Marchio, A.; Sitterlin, D.; Transy, C.; Dejean,

[61969] A.: Interaction of SP100 with HP1 proteins: a link between the promyelocytic

[61970] leukemia-associated nuclear bodies and the chromatin compartment. Proc.

[61971] Nat. Acad. Sci. 95: 7316-7321, 1998.

[61972] 10329.

[61973] Edwards, M. C.; Wong, C.; Elledge, S. J.: Human cyclin K, a novel

[61974] RNA polymerase II-associated cyclin possessing both car-boxy-terminal

[61975] domain kinase and Cdk-activating kinase activity. Molec. Cell. Biol. 18:

[61976] 4291-4300, 1998.

[61977] 10330.

[61978] Usami, S.; Abe, S.; Weston, M. D.; Shinkawa, H.; Van Camp, G.;

[61979] Kimberling, W. J.: Non-syndromic hearing loss associated with enlarged

[61980] vestibular aqueduct is caused by PDS mutations. Hum. Genet. 104:

[61981] 188-192, 1999.

[61982] 10331.

[61983] Scanlan, M. J.; Chen, Y.-T.; Williamson, B.; Gure, A. O.;

Stockert,
[61984] E.; Gordan, J. D.; Tureci, O.; Sahin, U.; Pfreundschuh, M.; Old, L.
[61985] J.: Characterization of human colon cancer antigens recognized by
[61986] autologous antibodies. Int. J. Cancer 76: 652-658, 1998.
[61987] 10332.
[61988] Halford, S.; Dulai, K. S.; Daw, S. C.; Fitzgibbon, J.; Hunt, D.
[61989] M.: Isolation and chromosomal localization of two human CDP-diacylglycerol
[61990] synthase (CDS) genes. Genomics 54: 140-144, 1998.
[61991] 10333.
[61992] Heacock, A. M.; Uhler, M. D.; Agranoff, B. W.: Cloning of CDP-diacylglycerol
[61993] synthase from a human neuronal cell line. J. Neurochem. 67: 2200-2203,
[61994] 1996.
[61995] 10334.
[61996] Weeks, R.; Dowhan, W.; Shen, H.; Balantac, N.; Meengs, B.; Nudelman,
[61997] E.; Leung, D. W.: Isolation and expression of an isoform of

human

[61998] CDP-diacylglycerol synthase cDNA. DNA Cell Biol. 16: 281-289, 1997.

[61999] 10335.

[62000] Volta, M.; Bulfone, A.; Gattuso, C.; Rossi, E.; Mariani, M.; Consalez,

[62001] G. G.; Zuffardi, O.; Ballabio, A.; Banfi, S.; Franco, B.: Iden-tification

[62002] and characterization of CDS2, a mammalian homolog of the Drosophila

[62003] CDP-diacylglycerol synthase gene. Genomics 55: 68-77, 1999.

[62004] 10336.

[62005] Borsani, G.; DeGrandi, A.; Ballabio, A.; Bulfone, A.; Bernard,

[62006] L.; Banfi, S.; Gattuso, C.; Mariani, M.; Dixon, M.; Donnai, D.; Metcalfe,

[62007] K.; Winter, R.; Robertson, M.; Axton, R.; Brown, A.; van Heyningen,

[62008] V.; Hanson, I.: EYA4, a novel vertebrate gene related to Drosophila

[62009] eyes absent. Hum. Molec. Genet. 8: 11-23, 1999.

[62010] 10337.

[62011] Kanamori, A.; Nakayama, J.; Fukuda, M. N.; Stallcup, W. B.; Sasaki,

[62012] K.; Fukuda, M.; Hirabayashi, Y.: Expression cloning and characterization

[62013] of a cDNA encoding a novel membrane protein required for the formation

[62014] of O-acetylated ganglioside: a putative acetyl-CoA transporter. Proc.

[62015] Nat. Acad. Sci. 94: 2897-2902, 1997.

[62016] 10338.

[62017] Paavola, P.; Horelli-Kuitunen, N.; Palotie, A.; Peltonen, L.:

[62018] Characterization of a novel gene, PNUTL2, on human chromosome 17q22-q23

[62019] and its exclusion as the Meckel syndrome gene. Genomics 55: 122-125,

[62020] 1999.

[62021] 10339.

[62022] Gregory, C. A.; Zabel, B.; Grant, M. E.; Boot-Handford, R. P.;

[62023] Wallis, G. A.: Equal expression of type X collagen mRNA

from mutant
[62024] and wild type COL10A1 alleles in growth plate cartilage from a patient
[62025] with metaphyseal chondrodysplasia type Schmid. (Letter) J. Med. Genet. 37:
[62026] 627-629, 2000.

[62027] 10340.

[62028] Tedesco, F.; Roncelli, L.; Petersen, B. H.; Agnello, V.; Sodetz,
[62029] J. M.: Two distinct abnormalities in patients with C8-alpha-gamma
[62030] deficiency: low level of C8-beta chain and presence of dysfunctional
[62031] C8-alpha-gamma subunit. J. Clin. Invest. 86: 884-888, 1990.
[62032] 10341.

[62033] Box, N. F.; Duffy, D. L.; Chen, W.; Stark, M.; Martin, N. G.; Sturm,
[62034] R. A.; Hayward, N. K.: MC1R genotype modifies risk of melanoma in
[62035] families segregating CDKN2A mutations. Am. J. Hum. Genet. 69: 765-773,

[62036] 2001.

[62037] 10342.

[62038] Box, N. F.; Wyeth, J. R.; O'Gorman, L. E.; Martin, N. G.; Sturm,

[62039] R. A.: Characterization of melanocyte stimulating hormone receptor

[62040] variant alleles in twins with red hair. Hum. Molec. Genet. 6: 1891-1897,

[62041] 1997.

[62042] 10343.

[62043] Chhajlani, V.; Wikberg, J. E. S.: Molecular cloning and expression

[62044] of the human melanocyte stimulating hormone receptor cDNA. FEBS Lett. 309:

[62045] 417-420, 1992.

[62046] 10344.

[62047] Flanagan, N.; Healy, E.; Ray, A.; Philips, S.; Todd, C.; Jackson,

[62048] I. J.; Birch-Machin, M. A.; Rees, J. L.: Pleiotropic effects of the

[62049] melanocortin 1 receptor (MC1R) gene on human pigmen-

tation. Hum. Molec.
[62050] Genet. 9: 2531-2537, 2000.

[62051] 10345.

[62052] Frandberg, P.-A.; Doufexis, M.; Kapas, S.; Chhajlani, V.: Human
[62053] pigmentation phenotype: a point mutation generates non-functional MSH
[62054] receptor. Biochem. Biophys. Res. Commun. 245: 490-492, 1998.
[62055] 10346.

[62056] Gantz, I.; Yamada, T.; Tashiro, T.; Konda, Y.; Shimoto, Y.; Miwa,
[62057] H.; Trent, J. M.: Mapping of the gene encoding the melanocortin-1
[62058] (alpha-melanocyte stimulating hormone) receptor (MC1R) to human chromosome
[62059] 16q24.3 by fluorescence in situ hybridization. Genomics 19: 394-395,
[62060] 1994.

[62061] 10347.

[62062] Healy, E.; Flannagan, N.; Ray, A.; Todd, C.; Jackson, I. J.;

Matthews,
[62063] J. N. S.; Birch-Machin, M. A.; Rees, J. L.: Melanocortin-
1-receptor
[62064] gene and sun sensitivity in individuals without red hair.
Lancet 355:
[62065] 1072-1073, 2000.

[62066] 10348.

[62067] Healy, E.; Jordan, S. A.; Budd, P. S.; Suffolk, R.; Rees, J. L.;

[62068] Jackson, I. J.: Functional variation of MC1R alleles from
red-haired
[62069] individuals. Hum. Molec. Genet. 10: 2397-2402, 2001.

[62070] 10349.

[62071] Jackson, I. J.: Colour-coded switches. Nature 362:
587-588,
[62072] 1993.

[62073] 10350.

[62074] Joerg, H.; Fries, H. R.; Meijerink, E.; Stranzinger, G. F.: Red

[62075] coat color in Holstein cattle is associated with a deletion
in the
[62076] MSHR gene. Mammalian Genome 7: 317-318, 1996.

[62077] 10351.

[62078] Johansson, M.; Marklund, L.; Sandberg, K.; Andersson, L.: Cosegregation

[62079] between the chestnut coat colour in horses and polymorphisms at the

[62080] melanocyte stimulating hormone (MSH) receptor locus. (Abstract) Animal

[62081] Genet. 25 (Suppl. 2): 35 only, 1994.

[62082] 10352.

[62083] John, P. R.; Ramsay, M.: Four novel variants in MC1R in red-haired

[62084] South African individuals of European descent: S83P, Y152X, A171D,

[62085] P256S. (Abstract) Hum. Mutat. 19: 461-462, 2002.

[62086] 10353.

[62087] Klungland, H.; Vage, D. I.; Gomez-Raya, L.; Adalsteinsson, S.;

[62088] Lien, S.: The role of melanocyte-stimulating hormone (MSH) receptor

[62089] in bovine coat color determination. Mammalian Genome 6: 636-639,

[62090] 1995.

[62091] 10354.

[62092] Koppula, S. V.; Robbins, L. S.; Lu, D.; Baack, E.; White, C. R.,

[62093] Jr.; Swanson, N. A.; Cone, R. D.: Identification of common polymorphisms

[62094] in the coding sequence of the human MSH receptor (MCIR) with possible

[62095] biological effects. Hum. Mutat. 9: 30-36, 1997.

[62096] 10355.

[62097] Marklund, L.; Johansson Moller, M.; Sandberg, K.; Anders-

[62098] son, L.

[62098] : A missense mutation in the gene for melanocyte-stimu-

[62099] lating hormone receptor (MC1R) is associated with the chestnut coat color in horses. Mammalian

[62100] Genome 7: 895-899, 1996.

[62101] 10356.

[62102] Mountjoy, K. G.; Robbins, L. S.; Mortrud, M. T.; Cone, R. D.:

[62103] The cloning of a family of genes that encode the

melanocortin receptors. Science 257:
[62104] 1248-1251, 1992.

[62105] 10357.

[62106] Palmer, J. S.; Duffy, D. L.; Box, N. F.; Aitken, J. F.;
O'Gorman,
[62107] L. E.; Green, A. C.; Hayward, N. K.; Martin, N. G.; Sturm, R.
A.:
[62108] Melanocortin-1 receptor polymorphisms and risk of
melanoma: is the
[62109] association explained solely by pigmentation phenotype?
Am. J. Hum.
[62110] Genet. 66: 176-186, 2000.

[62111] 10358.

[62112] Robbins, L. S.; Nadeau, J. H.; Johnson, K. R.; Kelly, M. A.;
Roselli-Rehfuss,
[62113] L.; Baack, E.; Mountjoy, K. G.; Cone, R. D.: Pigmentation
phenotypes
[62114] of variant extension locus alleles result from point muta-
tions that
[62115] alter MSH receptor function. Cell 72: 827-834, 1993.

[62116] 10359.

[62117] Savov, A.; Angelicheva, D.; Balassopoulou, A.; Jordanova, A.;

[62118] Noussia-Arvanitakis, S.; Kalaydjieva, L.: Double mutant alleles:

[62119] are they rare? Hum. Molec. Genet. 4: 1169-1171, 1995.

[62120] 10360.

[62121] Schioth, H. B.; Phillips, S. R.; Rudzish, R.; Birch-Machin, M.

[62122] A.; Wikberg, J. E. S.; Rees, J. L.: Loss of function mutations of

[62123] the human melanocortin 1 receptor are common and are associated with

[62124] red hair. Biochem. Biophys. Res. Commun. 260: 488-491, 1999.

[62125] 10361.

[62126] Smith, R.; Healy, E.; Siddiqui, S.; Flanagan, N.; Steijlen, P.

[62127] M.; Rosdahl, I.; Jacques, J. P.; Rogers, S.; Turner, R.; Jackson,

[62128] I. J.; Birch-Machin, M. A.; Rees, J. L.: Melanocortin 1 receptor

[62129] variants in an Irish population. J. Invest. Derm. 111: 119-122,

[62130] 1998.

[62131] 10362.

[62132] Theriault, A.; Boyd, E.; Whaley, K.; Sodetz, J. M.; Connor, J.

[62133] M.: Regional chromosomal assignment of genes encoding the alpha and

[62134] beta subunits of human complement protein c8 to 1p32. (Abstract) Cytogenet.

[62135] Cell Genet. 58: 1864 only, 1991.

[62136] 10363.

[62137] Theriault, A.; Boyd, E.; Whaley, K.; Sodetz, J. M.; Connor, J.

[62138] M.: Regional chromosomal assignment of genes encoding the alpha and

[62139] beta subunits of human complement protein C8 to 1p32. Hum. Genet. 88:

[62140] 703-704, 1992.

[62141] 10364.

[62142] Zhang, L.; Rittner, C.; Sodetz, J. M.; Schneider, P. M.; Kaufmann,

[62143] T.: The eighth component of human complement: molecular basis of

[62144] C8A (C81) polymorphism. Hum. Genet. 96: 281-284, 1995.

[62145] 10365.

[62146] Bahary, N.; Zorich, G.; Pachter, J. E.; Leibel, R. L.; Friedman,

[62147] J. M.: Molecular genetic linkage maps of mouse chromosomes 4 and

[62148] 6. Genomics 11: 33-47, 1991.

[62149] 10366.

[62150] Ross, S. C.; Densen, P.: Complement deficiency states and infection:

[62151] epidemiology, pathogenesis and consequences of neisserial and other

[62152] infections in an immune deficiency. Medicine 63: 243-273, 1984.

[62153] 10367.

[62154] Arnaout, M. A.; Gupta, S. K.; Pierce, M. W.; Tenen, D. G.: Amino

[62155] acid sequence of the alpha subunit of human leukocyte adhesion receptor

[62156] Mo1 (complement receptor type 3). J. Cell Biol. 106: 2153-2158,

[62157] 1988.

[62158] 10368.

[62159] Arnaout, M. A.; Remold-O'Donnell, E.; Pierce, M. W.; Harris, P.;

[62160] Tenen, D. G.: Molecular cloning of the alpha-subunit of human and

[62161] guinea pig leukocyte adhesion glycoprotein Mo1: chromosomal localization

[62162] and homology to the alpha-subunits of integrins. Proc. Nat. Acad.

[62163] Sci. 85: 2776-2780, 1988.

[62164] 10369.

[62165] Callen, D. F.; Chen, L. Z.; Nancarrow, J.; Whitmore, S. A.; Apostolou,

[62166] S.; Thompson, A. D.; Lane, S. A.; Stallings, R. L.; Hildebrand, C.

[62167] E.; Harris, P. G.; Sutherland, G. R.: Current state of the physical

[62168] map of human chromosome 16. (Abstract) Cytogenet. Cell Genet. 58:

[62169] 1998 only, 1991.

[62170] 10370.

[62171] Corbi, A. L.; Kishimoto, T. K.; Miller, L. J.; Springer, T. A.

[62172] : The human leukocyte adhesion glycoprotein Mac-1 (complement receptor

[62173] type 3, CD11b) alpha subunit: cloning, primary structure, and relation

[62174] to the integrins, von Willebrand factor and factor B. J. Biol. Chem. 263:

[62175] 12403-12411, 1988.

[62176] 10371.

[62177] Corbi, A. L.; Larson, R. S.; Kishimoto, T. K.; Springer, T. A.;

[62178] Morton, C. C.: Chromosomal location of the genes encod-
ing the leukocyte
[62179] adhesion receptors LFA-1, Mac-1 and p150,95: identifi-
cation of a gene
[62180] cluster involved in cell adhesion. J. Exp. Med. 167:
1597-1607,
[62181] 1988.

[62182] 10372.

[62183] Pierce, M. W.; Remold-O'Donnell, E.; Todd, R. F., III; Ar-
naout,
[62184] M. A.: N-terminal sequence of human leukocyte glycopro-
tein Mo1: conservation
[62185] across species and homology to platelet IIb/IIIa. Biochim.
Biophys.
[62186] Acta 874: 368-371, 1986.

[62187] 10373.

[62188] Simon, D. I.; Chen, Z.; Seifert, P.; Edelman, E. R.; Ballan-

tyne,
[62189] C. M.; Rogers, C.: Decreased neointimal formation in Mac-1 -/- mice
[62190] reveals a role for inflammation in vascular repair after angioplasty. J.
[62191] Clin. Invest. 105: 293-300, 2000.

[62192] 10374.

[62193] Springer, T. A.; Teplow, D. B.; Dreyer, W. J.: Sequence homology
[62194] of the LFA-1 and Mac-1 leukocyte adhesion glycoproteins and unexpected
[62195] relation to leukocyte interferon. Nature 314: 540-542, 1985.
[62196] 10375.

[62197] Brady, J. P.; Garland, D.; Duglas-Tabor, Y.; Robison, W. G., Jr.;
[62198] Groome, A.; Wawrousek, E. F.: Targeted disruption of the mouse alpha
[62199] A-crystallin gene induces cataract and cytoplasmic inclusion bodies
[62200] containing the small heat shock protein alpha B-crystallin. Proc.

[62201] Nat. Acad. Sci. 94: 884-889, 1997.

[62202] 10376.

[62203] Brakenhoff, R. H.; Henskens, H. A. M.; van Rossum, M. W. P. C.;

[62204] Lubsen, N. H.; Schoenmakers, J. G. G.: Activation of the gamma E-crystallin

[62205] pseudogene in the human hereditary Coppock-like cataract. Hum. Molec.

[62206] Genet. 3: 279-283, 1994.

[62207] 10377.

[62208] Cartier, M.; Breitman, M. L.; Tsui, L.-C.: A frameshift mutation

[62209] in the gamma-E-crystallin gene of the Elo mouse. Nature Genet. 2:

[62210] 42-45, 1992.

[62211] 10378.

[62212] Cobb, B. A.; Petrash, J. M.: Structural and functional changes

[62213] in the alpha-A-crystallin R116C mutant in hereditary cataracts. Biochemistry 39:

[62214] 15791-15798, 2000.

[62215] 10379.

[62216] de Jong, W. W.; Hendriks, W.: The eye lens crystallins: am-
biguity
[62217] as evolutionary strategy. J. Molec. Evol. 24: 121-129,
1986.
[62218] 10380.

[62219] Hawkins, J. W.; Van Keuren, M. L.; Piatigorsky, J.; Law, M.
L.;
[62220] Patterson, D.; Kao, F.-T.: Confirmation of assignment of
the human
[62221] alpha-1-crystallin gene (CRYA1) to chromosome 21 with
regional localization
[62222] to q22.3. Hum. Genet. 76: 375-380, 1987.

[62223] 10381.

[62224] Jaworski, C. J.; Piatigorsky, J.: A pseudo-exon in the func-
tional
[62225] human alpha-A-crystallin gene. Nature 337: 752-754,
1989.
[62226] 10382.

[62227] Kaye, N. W.; Lalley, P. A.; Petrash, J. M.; Church, R. L.: Re-
gional

[62228] assignment of the mouse alpha-A2-crystallin gene (Crya-1) to chromosome
[62229] 17A3-B by in situ hybridization. Cytogenet. Cell Genet. 53: 95-96,
[62230] 1990.

[62231] 10383.

[62232] Litt, M.; Kramer, P.; LaMorticella, D. M.; Murphey, W.; Lovrien,
[62233] E. W.; Weleber, R. G.: Autosomal dominant congenital cataract associated
[62234] with a missense mutation in the human alpha crystallin gene CRYAA. Hum.
[62235] Molec. Genet. 7: 471-474, 1998.

[62236] 10384.

[62237] Moormann, R. J. M.; den Dunnen, J. T.; Bloemendal, H.; Schoenmakers,
[62238] J. G. G.: Extensive intragenic sequence homology in two distinct
[62239] rat lens gamma-crystallin cDNAs suggests duplications of a primordial
[62240] gene. Proc. Nat. Acad. Sci. 79: 6876-6880, 1982.

[62241] 10385.

[62242] Petersen, M. B.; Slaugenhaupt, S. A.; Lewis, J. G.; Warren, A.

[62243] C.; Chakravarti, A.; Antonarakis, S. E.: A genetic linkage map of

[62244] 27 markers on human chromosome 21. Genomics 9: 407-419, 1991.

[62245] 10386.

[62246] Pras, E.; Frydman, M.; Levy-Nissenbaum, E.; Bakhan, T.; Raz, J.;

[62247] Assia, E. I.; Goldman, B.; Pras, E.: A nonsense mutation (W9X) in

[62248] CRYAA causes autosomal recessive cataract in an inbred Jewish Persian

[62249] family. Invest. Ophthal. Vis. Sci. 41: 3511-3515, 2000.

[62250] 10387.

[62251] Quax-Jeuken, Y.; Quax, W.; van Rens, G.; Meera Khan, P.; Bloemendal,

[62252] H.: Complete structure of the alpha-B-crystallin: conservation of

[62253] the exon-intron distribution in the two nonlinked alpha-crystallin

[62254] genes. Proc. Nat. Acad. Sci. 82: 5819-5823, 1985.

[62255] 10388.

[62256] Quax-Jeuken, Y.; Quax, W.; van Rens, G.; Meera Khan, P.; Bloemendal,

[62257] H.: Assignment of the human alpha-A-crystallin gene (CRYA1) to chromosome

[62258] 21. (Abstract) Cytogenet. Cell Genet. 40: 727-728, 1985.

[62259] 10389.

[62260] Schoffl, F.; Rascke, E.; Nagao, R. T.: The DNA sequence analysis

[62261] of soybean heat-shock genes and identification of possible regulatory

[62262] promoter elements. EMBO J. 3: 2491-2497, 1984.

[62263] 10390.

[62264] Skow, L. C.; Donner, M. E.: The locus encoding alpha-A-crystallin

[62265] is closely linked to H-2K on mouse chromosome 17. Genetics 110:

[62266] 723-732, 1985.

[62267] 10391.

[62268] Skow, L. C.; Kunz, H. W.; Gill, T. J., III: Linkage of the locus

[62269] encoding the A chain of alpha-crystallin (Acry-1) to the major histocompatibility

[62270] complex in the rat. Immunogenetics 22: 291-293, 1985.

[62271] 10392.

[62272] Wistow, G.: Domain structure and evolution in alpha-crystallins

[62273] and small heat shock proteins. FEBS Lett. 181: 1-6, 1985.

[62274] 10393.

[62275] Allen, E.; Yu, Q.-C.; Fuchs, E.: Mice expressing a mutant desmosomal

[62276] cadherin exhibit abnormalities in desmosomes, proliferation, and epidermal

[62277] differentiation. J. Cell Biol. 133: 1367-1382, 1996.

[62278] 10394.

[62279] Amagai, M.; Klaus-Kovtun, V.; Stanley, J. R.: Autoantibodies against

[62280] a novel epithelial cadherin in pemphigus vulgaris, a disease of cell

[62281] adhesion. Cell 67: 869-877, 1991.

[62282] 10395.

[62283] Koch, P. J.; Mahoney, M. G.; Ishikawa, H.; Pulkkinen, L.; Uitto,

[62284] J.; Schultz, L.; Murphy, G. F.; Whitaker-Menezes, D.; Stanley, J.

[62285] R.: Targeted disruption of the pemphigus vulgaris antigen (desmoglein

[62286] 3) gene in mice causes loss of keratinocyte cell adhesion with a phenotype

[62287] similar to pemphigus vulgaris. J. Cell Biol. 137: 1091-1102, 1997.

[62288] 10396.

[62289] Rickman, L.; Simrak, D.; Stevens, H. P.; Hunt, D. M.; King, I.

[62290] A.; Bryant, S. P.; Eady, R. A. J.; Leigh, I. M.; Arnemann, J.; Magee,

[62291] A. I.; Kelsell, D. P.; Buxton, R. S.: N-terminal deletion in a desmosomal

[62292] cadherin causes the autosomal dominant skin disease striate palmoplantar

[62293] keratoderma. Hum. Molec. Genet. 8: 971-976, 1999.

[62294] 10397.

[62295] Simrak, D.; Cowley, C. M. E.; Buxton, R. S.; Arnemann, J.: Tandem
[62296] arrangement of the closely linked desmoglein genes on human chromosome
[62297] 18. Genomics 25: 591-594, 1995.

[62298] 10398.

[62299] Wang, Y.; Amagai, M.; Minoshima, S.; Sakai, K.; Green, K. J.; Nishikawa,
[62300] T.; Shimizu, N.: The human genes for desmogleins (DSG1 and DSG3)
[62301] are located in a small region on chromosome 18q12. Genomics 20:
[62302] 492-495, 1994.

[62303] 10399.

[62304] Hunt, D. M.; Rickman, L.; Whittock, N. V.; Eady, R. A. J.; Simrak,
[62305] D.; Dopping-Hepenstal, P. J. C.; Stevens, H. P.; Armstrong, D. K.
[62306] B.; Hennies, H. C.; Kuster, W.; Hughes, A. E.; Arnemann, J.; Leigh,
[62307] I. M.; McGrath, J. A.; Kelsell, D. P.; Buxton, R. S.: Spectrum of

[62308] dominant mutations in the desmosomal cadherin desmoglein 1, causing

[62309] the skin disease striate palmoplantar keratoderma. Europ. J. Hum.

[62310] Genet. 9: 197-203, 2001.

[62311] 10400.

[62312] Warren, S. J. P.; Lin, M.-S.; Giudice, G. J.; Hoffmann, R. G.;

[62313] Hans-Filho, G.; Aoki, V.; Rivitti, E. A.; dos Santos, V.; Diaz, L.

[62314] A.: The prevalence of antibodies against desmoglein 1 in endemic

[62315] pemphigus foliaceus in Brazil. New Eng. J. Med. 343: 23-30, 2000.

[62316] 10401.

[62317] Wu, H.; Wang, Z. H.; Yan, A.; Lyle, S.; Fakharzadeh, S.; Wahl,

[62318] J. K.; Wheelock, M. J.; Ishikawa, H.; Uitto, J.; Amagai, M.; Stanley,

[62319] J. R.: Protection against pemphigus foliaceus by desmoglein 3 in

[62320] neonates. New Eng. J. Med. 343: 31-35, 2000.

[62321] 10402.

[62322] Arnemann, J.; Spurr, N. K.; Magee, A. I.; Buxton, R. S.: The human

[62323] gene (DSG2) coding for HDGC, a second member of the desmoglein subfamily

[62324] of the desmosomal cadherins, is, like DSG1 coding for desmoglein DGI,

[62325] assigned to chromosome 18. Genomics 13: 484-486, 1992.

[62326] 10403.

[62327] Koch, P. J.; Goldschmidt, M. D.; Walsh, M. J.; Zimbelmann, R.;

[62328] Franke, W. W.: Complete amino acid sequence of the epi-dermal desmoglein

[62329] precursor polypeptide and identification of a second type of desmoglein

[62330] gene. Europ. J. Cell Biol. 55: 200-208, 1991.

[62331] 10404.

[62332] Inskip, A.; Elexperu-Camiruaga, J.; Buxton, N.; Dias, P. S.; MacIntosh,

[62333] J.; Campbell, D.; Jones, P. W.; Yengi, L.; Talbot, J. A.; Strange,

[62334] R. C.; Fryer, A. A.: Identification of polymorphism at the glutathione
[62335] S-transferase, GSTM3 locus: evidence for linkage with GSTM1*A. Biochem.
[62336] J. 312: 713-716, 1995.

[62337] 10405.

[62338] Van Cong, N.; Laisney, V.; Gross, M. S.; Frezal, J.: Glutathione-S-transferases--tissues
[62339] distribution, number of loci, polymorphism, chromosome localization.
[62340] (Abstract) Cytogenet. Cell Genet. 37: 554, 1984.

[62341] 10406.

[62342] Law, M. L.; Kao, F.-T.: Induced segregation of human syntenic
[62343] genes by 5-bromodeoxyuridine plus near-visible light. Somat. Cell
[62344] Genet. 4: 465-476, 1978.

[62345] 10407.

[62346] Hopkinson, D. A.; Peters, J.; Harris, H.: Rare electrophoretic
[62347] variants of glycerol-3-phosphate dehydrogenase: evi-

dence for two structural
[62348] gene loci (GPD-1 and GPD-2). Ann. Hum. Genet. 37:
477-484, 1974.
[62349] 10408.

[62350] Kielty, C.; Povey, S.: Mapping of liver-specific enzymes:
alpha-glycerophosphate
[62351] dehydrogenase (GPD1) and others. (Abstract) Cytogenet.
Cell Genet. 32:
[62352] 290 only, 1982.

[62353] 10409.

[62354] Menaya, J.; Gonzalez-Manchon, C.; Parrilla, R.; Ayuso, M.
S.:
[62355] Molecular cloning, sequencing and expression of a cDNA
encoding a
[62356] human liver NAD-dependent alpha-glycerol-3-phosphate
dehydrogenase. Biochim.
[62357] Biophys. Acta 1262: 91-94, 1995.

[62358] 10410.

[62359] Prasad, R.; Zhadanov, A. B.; Sedkov, Y.; Bullrich, F.; Druck,
T.;
[62360] Rallapalli, R.; Yano, T.; Alder, H.; Croce, C. M.; Huebner,

K.; Mazo,
[62361] A.; Canaani, E.: Structure and expression pattern of human ALR, a
[62362] novel gene with strong homology to ALL-1 involved in acute leukemia
[62363] and to Drosophila trithorax. Oncogene 15: 549-560, 1997.
[62364] 10411.

[62365] Brown, L. J.; Stoffel, M.; Moran, S.; Lehn, D. A.; Pomije, C.;

[62366] Le Beau, M. M.; MacDonald, M. J.: Structure of the mitochondrial
[62367] glycerol-3-phosphate dehydrogenase gene and mapping to chromosome 10412.
[62368] (Abstract) Am. J. Hum. Genet. 57: A141 only, 1995.

[62369]

[62370] 2. Ferrer, J.; Aoki, M.; Behn, P.; Nestorowicz, A.; Riggs, A.; Permutt,
[62371] M. A.: Mitochondrial glycerol-3-phosphate dehydrogenase: cloning
[62372] of an alternatively spliced human islet-cell cDNA, tissue distribution,
[62373] physical mapping, and identification of a polymorphic ge-

netic marker. Diabetes 45:
[62374] 262-266, 1996.

[62375] 10413.

[62376] Novials, A.; Vidal, J.; Franco, C.; Ribera, F.; Sener, A.; Malaisse,
[62377] W. J.; Gomis, R.: Mutation in the calcium-binding domain of the mitochondrial
[62378] glycerophosphate dehydrogenase gene in a family of diabetic subjects. Biochem.
[62379] Biophys. Res. Comm. 231: 570-572, 1997.

[62380] 10414.

[62381] Shaw, M.-A.; Edwards, Y. H.; Hopkinson, D. A.: Human mitochondrial
[62382] glycerol phosphate dehydrogenase (GPD-M) isozymes. Ann. Hum. Genet. 46:
[62383] 11-23, 1982.

[62384] 10415.

[62385] Hashimoto, S.; Chiorazzi, N.; Gregersen, P. K.: The complete sequence
[62386] of the human CD79b (Ig-beta/B29) gene: identification of a conserved

[62387] exon/intron organization, immunoglobulin-like regulatory regions,

[62388] and allelic polymorphism. Immunogenetics 40: 145-149, 1994.

[62389] 10416.

[62390] Hashimoto, S.; Gregersen, P. K.; Chiorazzi, N.: The human Ig-beta

[62391] cDNA sequence, a homologue of murine B29, is identical in B cell and

[62392] plasma cell lines producing all the human Ig isotypes. J. Immun. 150:

[62393] 491-498, 1993.

[62394] 10417.

[62395] Wood, W. J., Jr.; Thompson, A. A.; Korenberg, J.; Chen, X.; May,

[62396] W.; Wall, R.; Denny, C. T.: Isolation and chromosomal mapping of

[62397] the human immunoglobulin-associated B29 gene (IGB). Genomics 16:

[62398] 187-192, 1993.

[62399] 10418.

[62400] Okabe, I.; Nussbaum, R. L.: Identification and chromoso-
mal mapping
[62401] of the mouse inositol polyphosphate 1-phosphatase gene.
Genomics 30:
[62402] 358-360, 1995.

[62403] 10419.

[62404] Woodcock, E. A.; Wang, B. H.; Arthur, J. F.; Lennard, A.;
Matkovich,
[62405] S. J.; Du, X.-J.; Brown, J. H.; Hannan, R. D.: Inositol
polyphosphate
[62406] 1-phosphatase is a novel antihypertrophic factor. J. Biol.
Chem. 277:
[62407] 22734-22742, 2002.

[62408] 10420.

[62409] York, J. D.; Veile, R. A.; Donis-Keller, H.; Majerus, P. W.:
Cloning,
[62410] heterologous expression, and chromosomal localization
of human inositol
[62411] polyphosphate 1-phosphatase. Proc. Nat. Acad. Sci. 90:
5833-5837,
[62412] 1993.

[62413] 10421.

[62414] Matsumoto, M.; Nakagawa, T.; Inoue, T.; Nagata, E.; Tanaka, K.;

[62415] Takano, H.; Minowa, O.; Kuno, J.; Sakakibara, S.; Yamada, M.; Yoneshima,

[62416] H.; Miyawaki, A; Fukuichi, T.; Furuichi, T.; Okano, H.; Mikoshiba,

[62417] K.; Noda, T.: Ataxia and epileptic seizures in mice lacking type

[62418] 1 inositol 1,4,5-triphosphate receptor. Nature 379: 168-171, 1996.

[62419] 10422.

[62420] Nucifora, F. C., Jr.; Li, S.-H.; Danoff, S.; Ullrich, A.; Ross,

[62421] C. A.: Molecular cloning of a cDNA for the human inositol 1,4,5-trisphosphate

[62422] receptor type 1, and the identification of a third alternatively spliced

[62423] variant. Molec. Brain Res. 32: 291-296, 1995.

[62424] 10423.

[62425] Bradham, D. M.; Igarashi, A.; Potter, R. L.; Grotendorst, G. R.

[62426] : Connective tissue growth factor: a cysteine-rich mitogen secreted
[62427] by human vascular endothelial cells is related to the SRC-induced
[62428] immediate early gene product CEF-10. J. Cell Biol. 114: 1285-1294,
[62429] 1991.

[62430] 10424.

[62431] Aguilar-Salinas, C. A.; Reyes-Rodriguez, E.; Ordonez-Sanchez, M.
[62432] L.; Torres, M. A.; Ramirez-Jimenez, S.; Dominguez-Lopez, A.; Martinez-Francois,
[62433] J. R.; Velasco-Perez, M. L.; Alpizar, M.; Garcia-Garcia, E.; Gomez-Perez,
[62434] F.; Rull, J.; Tusie-Luna, M. T.: Early-onset type 2 diabetes: metabolic
[62435] and genetic characterization in the Mexican population. J. Clin.
[62436] Endocr. Metab. 86: 220-226, 2001.

[62437] 10425.

[62438] DerKinderen, D. J.; Koten, J. W.; Tan, K. E. W. P.; Beemer, F.

[62439] A.; Van Romunde, L. K. J.; Den Otter, W.: Parental age in sporadic
[62440] hereditary retinoblastoma. Am. J. Ophthal. 110: 605-609, 1990.
[62441] 10426.

[62442] Orphanides, G.; Wu, W.-H.; Lane, W. S.; Hampsey, M.; Reinberg,
[62443] D.: The chromatin-specific transcription elongation factor FACT comprises
[62444] human SPT16 and SSRP1 proteins. Nature 400: 284-288, 1999.
[62445] 10427.

[62446] Shi, Y.; Ullrich, S. J.; Zhang, J.; Connolly, K.; Grze-
gorzewski,
[62447] K. J.; Barber, M. C.; Wang, W.; Wathen, K.; Hodge, V.; Fisher, C.
[62448] L.; Olsen, H.; Ruben, S. M.; Knyazev, I.; Cho, Y. H.; Kao, V.; Wilkinson,
[62449] K. A.; Carrell, J. A.; Ebner, R.: A novel cytokine receptor-lig-
and
[62450] pair: identification, molecular characterization, and in vivo immunomodulatory

[62451] activity. J. Biol. Chem. 275: 19167-19176, 2000.

[62452] 10428.

[62453] Lee, H.-K.; Lee, Y.-K.; Park, S.-H.; Kim, Y.-S.; Park, S. H.; Lee,

[62454] J. W.; Kwon, H.-B.; Soh, J.; Moore, D. D.; Choi, H.-S.: Structure

[62455] and expression of the orphan nuclear receptor SHP gene. J. Biol.

[62456] Chem. 273: 14398-14402, 1998.

[62457] 10429.

[62458] Nishigori, H.; Tomura, H.; Tonooka, N.; Kanamori, M.; Ya-mada, S.;

[62459] Sho, K.; Inoue, I.; Kikuchi, N.; Onigata, K.; Kojima, I.; Ko-hama,

[62460] T.; Yamagata, K.; and 9 others: Mutations in the small heterodimer

[62461] partner gene are associated with mild obesity in Japanese subjects. Proc.

[62462] Nat. Acad. Sci. 98: 575-580, 2001.

[62463] 10430.

[62464] Seol, W.; Choi, H.-S.; Moore, D. D.: An orphan nuclear

hormone

[62465] receptor that lacks a DNA binding domain and het-
erodimerizes with

[62466] other receptors. Science 272: 1336-1339, 1996.

[62467] 10431.

[62468] Friedman, L. S.; Ostermeyer, E. A.; Lynch, E. D.; Welcsh, P.;
Szabo,

[62469] C. I.; Meza, J. E.; Anderson, L. A.; Dowd, P.; Lee, M. K.;
Rowell,

[62470] S. E.; Ellison, J.; Boyd, J.; King, M.-C.: 22 genes from chro-
mosome

[62471] 17q21: cloning, sequencing, and characterization of mu-
tations in breast

[62472] cancer families and tumors. Genomics 25: 256-163,
1995.

[62473] 10432.

[62474] Linial, M.; Miller, K.; Scheller, R. H.: VAT-1: an abundant
membrane

[62475] protein from Torpedo cholinergic synaptic vesicles. Neu-
ron 2: 1265-1273,

[62476] 1989.

[62477] 10433.

[62478] Smith, T. M.; Lee, M. K.; Szabo, C. I.; Jerome, N.; McEuen, M.;
[62479] Taylor, M.; Hood, L.; King, M.-C.: Complete genomic sequence and
[62480] analysis of 117 kb of human DNA containing the gene BRCA1. Genome
[62481] Res. 6: 1029-1049, 1996.

[62482] 10434.

[62483] Echchakir, H.; Mami-Chouaib, F.; Vergnon, I.; Baurain, J.-F.; Karanikas,
[62484] V.; Chouaib, S.; Coulie, P. G.: A point mutation in the alpha-actinin-4
[62485] gene generates an antigenic peptide recognized by autologous cytolytic
[62486] T lymphocytes on a human lung carcinoma. Cancer Res. 61: 4078-4083,
[62487] 2001.

[62488] 10435.

[62489] Wieland, I.; Arden, K. C.; Michels, D.; Klein-Hitpass, L.; Bohm,
[62490] M.; Viars, C. S.; Weidle, U. H.: Isolation of DICE1: a gene frequently

[62491] affected by LOH and downregulated in lung carcinomas. Oncogene 18:
[62492] 4530-4537, 1999.

[62493] 10436.

[62494] Huebner, K.; Cannizzaro, L. A.; Nakamura, T.; Hillova, J.; Mariage-Samson,
[62495] R.; Hecht, F.; Hill, M.; Croce, C. M.: A rearranged trans-forming
[62496] gene, tre, is made up of human sequences derived from chromosome regions
[62497] 5q, 17q and 18q. Oncogene 3: 449-455, 1988.

[62498] 10437.

[62499] Nakamura, T.; Hillova, J.; Mariage-Samson, R.; Hill, M.: Molecular
[62500] cloning of a novel oncogene generated by DNA recombi-nation during
[62501] transfection. Oncogene Res. 2: 357-370, 1988.

[62502] 10438.

[62503] Nakamura, T.; Hillova, J.; Mariage-Samson, R.; Onno, M.; Huebner,
[62504] K.; Cannizzaro, L. A.; Boghosian-Sell, L.; Croce, C. M.; Hill,

M.

[62505] : A novel transcriptional unit of the tre oncogene widely expressed

[62506] in human cancer cells. Oncogene 7: 733-741, 1992.

[62507] 10439.

[62508] Chen, H.; Rossier, C.; Morris, M. A.; Scott, H. S.; Gos, A.; Bairoch,

[62509] A.; Antonarakis, S. E.: A testis-specific gene, TPTE, encodes a putative

[62510] transmembrane tyrosine phosphatase and maps to the pericentromeric

[62511] region of human chromosomes 21 and 13, and to chromosomes 15, 22,

[62512] and Y. Hum. Genet. 105: 399-409, 1999.

[62513] 10440.

[62514] Guipponi, M.; Tapparel, C.; Jousson, O.; Scamuffa, N.; Mas, C.;

[62515] Rossier, C.; Hutter, P.; Meda, P.; Lyle, R.; Reymond, A.; Antonarakis,

[62516] S. E.: The murine orthologue of the Golgi-localized TPTE protein

[62517] provides clues to the evolutionary history of the human

TPTE gene

[62518] family. Hum. Genet. 109: 569-575, 2001.

[62519] 10441.

[62520] Guipponi, M.; Yaspo, M.-L.; Riesselman, L.; Chen, H.; De Sario,

[62521] A.; Roizes, G.; Antonarakis, S. E.: Genomic structure of a copy of

[62522] the human TPTE gene which encompasses 87 kb on the short arm of chromosome

[62523] 21. Hum. Genet. 107: 127-131, 2000.

[62524] 10442.

[62525] Gruber, A. D.; Pauli, B. U.: Molecular cloning and bio-chemical

[62526] characterization of a truncated, secreted member of the human family

[62527] of Ca(2+)-activated Cl- channels. Biochim. Biophys. Acta 1444: 418-423,

[62528] 1999.

[62529] 10443.

[62530] Bause, E.; Bieberich, E.; Rolfs, A.; Volker, C.; Schmidt, B.:

[62531] Molecular cloning and primary structure of

Man(9)-mannosidase from
[62532] human kidney. Eur. J. Biochem. 217: 535-540, 1993.

[62533] 10444.

[62534] Tremblay, L. O; Campbell Dyke, N.; Herscovics, A.: Molec-
ular cloning,
[62535] chromosomal mapping and tissue-specific expression of a
novel human
[62536] alpha-1,2-mannosidase gene involved in N-glycan matu-
ration. Glycobiology 8:
[62537] 585-595, 1998.

[62538] 10445.

[62539] Nakano, M.; Yoshiura, K.; Oikawa, M.; Miyoshi, O.; Ya-
mada, K.;
[62540] Kondo, S.; Miwa, N.; Soeda, E.; Jinno, Y.; Fujii, T.; Niikawa,
N.
[62541] : Identification, characterization and mapping of the hu-
man ZIS (zinc-finger,
[62542] splicing) gene. Gene 225: 59-65, 1998.

[62543] 10446.

[62544] Koch, M.; Olson, P. F.; Albus, A.; Jin, W.; Hunter, D. D.;
Brunken,

[62545] W. J.; Burgeson, R. E.; Champliaud, M. F.: Characterization and expression

[62546] of the laminin gamma-3 chain: a novel, non-basement membrane-associated,

[62547] laminin chain. J. Cell Biol. 145: 605-617, 1999.

[62548] 10447.

[62549] Szekely, A. M.; Chen, Y.-H.; Zhang, C.; Oshima, J.; Weissman,

[62550] S. M.: Werner protein recruits DNA polymerase delta to the nucleolus. Proc.

[62551] Nat. Acad. Sci. 97: 11365-11370, 2000.

[62552] 10448.

[62553] Jedlitschky, G.; Cassidy, A. J.; Sales, M.; Pratt, N.; Burchell,

[62554] B.: Cloning and characterization of a novel human olfactory UDP-glucuronosyltransferase. Biochem.

[62555] J. 340: 837-843, 1999.

[62556] 10449.

[62557] Lazard, D.; Zupko, K.; Poria, Y.; Nef, P.; Lazarovits, J.; Horn,

[62558] S.; Khen, M.; Lancet, D.: Odorant signal termination by olfactory

[62559] UDP glucuronosyl transferase. Nature 349: 790-793, 1991.

[62560] 10450.

[62561] Baker, E.; Crawford, J.; Sutherland, G. R.; Freeman, C.; Parish,

[62562] C. R.; Hulett, M. D.: Human HPA endoglycosidase hep-aranase. Chromosome

[62563] Res. 7: 319 only, 1999.

[62564] 10451.

[62565] Hulett, M. D.; Freeman, C.; Hamdorf, B. J.; Baker, R. T.; Harris,

[62566] M. J.; Parish, C. R.: Cloning of mammalian heparanase, an important

[62567] enzyme in tumor invasion and metastasis. Nature Med. 5: 803-809,

[62568] 1999.

[62569] 10452.

[62570] Kussie, P. H.; Hulmes, J. D.; Ludwig, D. L.; Patel, S.; Navarro,

[62571] E. C.; Seddon, A. P.; Giorgio, N. A.; Bohlen, P: Cloning and functional

[62572] expression of a human heparanase gene. Biochem. Biophys. Res. Commun. 261:
[62573] 183-187, 1999.

[62574] 10453.

[62575] Toyoshima, M.; Nakajima, M.: Human heparanase: purification, characterization,
[62576] cloning, and expression. J. Biol. Chem. 274:
24153-24160, 1999.
[62577] 10454.

[62578] Vlodavsky, I.; Friedmann, Y.; Elkin, M.; Aingorn, H.; Atzmon, R.;
[62579] Ishai-Michaeli, R.; Bitan, M.; Pappo, O.; Peretz, T.; Michal, I.;
[62580] Spector, L.; Pecker, I.: Mammalian heparanase: gene cloning, expression
[62581] and function in tumor progression and metastasis. Nature Med. 5:
[62582] 793-802, 1999.

[62583] 10455.

[62584] Ikeda, A.; Ikeda, S.; Gridley, T.; Nishina, P. M.; Naggert, J.

[62585] K.: Neural tube defects and neuroepithelial cell death in

Tulp3 knockout
[62586] mice. Hum. Molec. Genet. 10: 1325-1334, 2001.

[62587] 10456.

[62588] Nishina, P. M.; North, M. A.; Ikeda, A.; Yan, Y.; Naggert, J. K.
[62589] : Molecular characterization of a novel tubby gene family member,
[62590] TULP3, in mouse and humans. Genomics 54: 215-220, 1998.
[62591] 10457.

[62592] Wu, X.; Kekuda, R.; Huang, W.; Fei, Y.-J.; Leibach, F. H.; Chen,
[62593] J.; Conway, S. J.; Ganapathy, V.: Identity of the organic cation
[62594] transporter OCT3 as the extraneuronal monoamine transporter (uptake-2)
[62595] and evidence for the expression of the transporter in the brain. J.
[62596] Biol. Chem. 273: 32776-32786, 1998.

[62597] 10458.

[62598] Seki, N.; Ohira, M.; Nagase, T.; Ishikawa, K.; Miyajima, N.;

Nakajima,

[62599] D.; Nomura, N.; Ohara, O.: Characterization of cDNA clones in size-fractionated

[62600] cDNA libraries from human brain. DNA Res. 4: 345-349, 1997.

[62601] 10459.

[62602] Sumoy, L.; Pluvinet, R.; Andreu, N.; Estivill, X.; Escarceller,

[62603] M.: PACSIN 3 is a novel SH3 domain cytoplasmic adapter protein of

[62604] the pacsin-syndapin-FAP52 gene family. Gene 262: 199-205, 2001.

[62605] 10460.

[62606] Hubener, C.; Mincheva, A.; Lichter, P.; Schraven, B.; Bruyns, E.

[62607] : Genomic organization and chromosomal localization of the human gene

[62608] encoding the T-cell receptor-interacting molecule (TRIM). Immunogenetics 51:

[62609] 154-158, 2000.

[62610] 10461.

[62611] Bruyns, E.; Marie-Cardine, A.; Kirchgessner, H.; Sagolla,

K.; Shevchenko,
[62612] A.; Mann, M.; Autschbach, F.; Bensussan, A.; Meuer, S.; Schraven,
[62613] B.: T cell receptor (TCR) interacting molecule (TRIM), a novel disulfide-linked
[62614] dimer associated with the TCR-CD3-zeta complex, recruits intracellular
[62615] signaling proteins to the plasma membrane. J. Exp. Med. 188: 561-575,
[62616] 1998.

[62617] 10462.

[62618] Zhao, R.; Qi, Y.; Chen, J.; Zhao, Z. J.: FYVE-DSP2, a FYVE domain-containing
[62619] dual specificity protein phosphatase that dephosphorylates phosphotidylinositol
[62620] (sic) 3-phosphate. Exp. Cell Res. 265: 329-338, 2001.

[62621] 10463.

[62622] Cui, X.; De Vivo, I.; Slany, R.; Miyamoto, A.; Firestein, R.; Cleary,
[62623] M. L.: Association of SET domain and myotubularin-related proteins
[62624] modulates growth control. Nature Genet. 18: 331-337,

1998.

[62625] 10464.

[62626] Mansour, S. J.; Herbrick, J.-A.; Scherer, S. W.; Melancon, P.:

[62627] Human GBF1 is a ubiquitously expressed gene of the Sec7 domain family

[62628] mapping to 10q24. Genomics 54: 323-327, 1998.

[62629] 10465.

[62630] Frye, R. A.: Characterization of five human cDNAs with homology

[62631] to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metab-olize NAD

[62632] and may have protein ADP-ribosyltransferase activity. Biochem. Biophys.

[62633] Res. Commun. 260: 273-279, 1999.

[62634] 10466.

[62635] Aula, N.; Salomaki, P.; Timonen, R.; Verheijen, F.; Mancini, G.;

[62636] Mansson, J.-E.; Aula, P.; Peltonen, L.: The spectrum of SLC17A5-gene

[62637] mutations resulting in free sialic acid-storage diseases in-dicates

[62638] some genotype-phenotype correlation. Am. J. Hum. Genet. 67: 832-840,

[62639] 2000.

[62640] 10467.

[62641] Biancheri, R.; Verbeek, E.; Rossi, A.; Gaggero, R.; Roc-catagliata,

[62642] L.; Gatti, R.; van Diggelen, O.; Verheijen, F. W.; Mancini, G. M.

[62643] S.: An Italian severe Salla disease variant associated with a SLC17A5

[62644] mutation earlier described in infantile sialic acid storage disease. Clin.

[62645] Genet. 61: 443-447, 2002.

[62646] 10468.

[62647] Belinsky, M. G.; Bain, L. J.; Balsara, B. B.; Testa, J. R.; Kruh,

[62648] G. D.: Characterization of MOAT-C and MOAT-D, new members of the

[62649] MRP/cMOAT subfamily of transporter proteins. J. Nat. Cancer Inst. 90:

[62650] 1735-1741, 1998.

[62651] 10469.

[62652] Fromm, M. F.; Leake, B.; Roden, D. M.; Wilkinson, G. R.; Kim, R.

[62653] B.: Human MRP3 transporter: identification of the 5-prime flanking

[62654] region, genomic organization and alternative splice variants. Biochim.

[62655] Biophys. Acta 1415: 369-374, 1999.

[62656] 10470.

[62657] Kiuchi, Y.; Suzuki, H.; Hirohashi, T.; Tyson, C. A.; Sugiyama,

[62658] Y.: cDNA cloning and inducible expression of human multidrug resistance

[62659] associated protein 3 (MRP3). FEBS Lett. 433: 149-152, 1998.

[62660] 10471.

[62661] Konig, J.; Rost, D.; Cui, Y.; Keppler, D.: Characterization of

[62662] the human multidrug resistance protein isoform MRP3 localized to the

[62663] basolateral hepatocyte membrane. Hepatology 29: 1156-1163, 1999.

[62664] 10472.

[62665] Kool, M.; van der Linden, M.; de Haas, M.; Scheffer, G. L.; de

[62666] Vree, J. M. L.; Smith, A. J.; Jansen, G.; Peters, G. J.; Ponne, N.;

[62667] Scheper, R. J.; Oude Elferink, R. P. J.; Baas, F.; Borst, P.: MRP3,

[62668] an organic anion transporter able to transport anti-cancer drugs. Proc.

[62669] Nat. Acad. Sci. 96: 6914-6919, 1999.

[62670] 10473.

[62671] Ortiz, D. F.; Li, S.; Iyer, R.; Zhang, X.; Novikoff, P.; Arias,

[62672] I. M.: MRP3, a new ATP-binding cassette protein localized to the

[62673] canalicular domain of the hepatocyte. Am. J. Physiol. 276: G1493-G1500,

[62674] 1999.

[62675] 10474.

[62676] Scott, A. F.: Personal Communication. Baltimore, Md. 9/18/2000.

[62677] 10475.

[62678] Cooper, M. P.; Machwe, A.; Orren, D. K.; Brosh, R. M.;

Ramsden,
[62679] D.; Bohr, V. A.: Ku complex interacts with and stimulates the Werner
[62680] protein. Genes Dev. 14: 907-912, 2000.

[62681] 10476.

[62682] Gangloff, S.; Soustelle, C.; Fabre, F.: Homologous recombination
[62683] is responsible for cell death in the absence of the Sgs1 and Srs2
[62684] helicases. Nature Genet. 25: 192-194, 2000.

[62685] 10477.

[62686] Huang, S.; Li, B.; Gray, M. D.; Oshima, J.; Mian, I. S.; Campisi,
[62687] J.: The premature ageing syndrome protein, WRN, is a 3-prime-5-prime
[62688] exonuclease. Nature Genet. 20: 114-115, 1998.

[62689] 10478.

[62690] Ketting, R. F.; Haverkamp, T. H. A.; van Luenen, H. G. A. M.; Plasterk,
[62691] R. H. A.: mut-7 of C. elegans, required for transposon silencing

[62692] and RNA interference, is a homolog of Werner syndrome helicase and
[62693] RNaseD. Cell 99: 133-141, 1999.

[62694] 10479.

[62695] Yokota, T.; Oritani, K.; Takahashi, I.; Ishikawa, J.; Mat-
[62696] suyama, A.; Ouchi, N.; Kihara, S.; Funahashi, T.; Tenner, A. J.; Tomiyama,
[62697] Y.; Matsuzawa, Y.: Adiponectin, a new member of the family of soluble
[62698] defense collagens, negatively regulates the growth of myelomonocytic
[62699] progenitors and the functions of macrophages. Blood 96: 1723-1732,
[62700] 2000.

[62701] 10480.

[62702] Hjalt, T. A.; Murray, J. C.: The human BARX2 gene: ge-
[62703] nomic structure, chromosomal localization, and single nucleotide polymor-
[62704] phisms. Genomics 62: 456-459, 1999.

[62705] 10481.

[62706] Jones, F. S.; Kioussi, C.; Copertino, D. W.; Kallunki, P.; Holst,

[62707] B. D.; Edelman, G. M.: Barx2, a new homeobox gene of the Bar class,

[62708] is expressed in neural and craniofacial structures during development. Proc.

[62709] Nat. Acad. Sci. 94: 2632-2637, 1997.

[62710] 10482.

[62711] Arking, D. E.; Krebsova, A.; Macek, M., Sr.; Macek, M., Jr.; Arking,

[62712] A.; Mian, I. S.; Fried, L.; Hamosh, A.; Dey, S.; McIntosh, I.; Dietz,

[62713] H. C.: Association of human aging with a functional variant of klotho. Proc.

[62714] Nat. Acad. Sci. 99: 856-861, 2002.

[62715] 10483.

[62716] Koh, N.; Fujimori, T.; Nishiguchi, S.; Tamori, A.; Shiomi, S.;

[62717] Nakatani, T.; Sugimura, K.; Kishimoto, T.; Kinoshita, S.; Kuroki,

[62718] T.; Nabeshima, Y.: Severely reduced production of Klotho

in human
[62719] chronic renal failure kidney. Biochem. Biophys. Res. Commun. 280:
[62720] 1015-1020, 2001.

[62721] 10484.

[62722] Kuro-o, M.; Matsumura, Y.; Aizawa, H.; Kawaguchi, H.; Suga, T.;
[62723] Utsugi, T.; Ohyama, Y.; Kurabayashi, M.; Kaname, T.; Kume, E.; Iwasaki,
[62724] H.; Iida, A.; Shiraki-Iida, T.; Nishikawa, S.; Nagai, R.; Nabeshima,
[62725] Y.: Mutation of the mouse klotho gene leads to a syndrome resembling
[62726] ageing. Nature 390: 45-51, 1997.

[62727] 10485.

[62728] Matsumura, Y.; Aizawa, H.; Shiraki-Iida, T.; Nagai, R.; Kuro-o,
[62729] M.; Nabeshima, Y.: Identification of the human klotho gene and its
[62730] two transcripts encoding membrane and secreted klotho protein. Biochem.
[62731] Biophys. Res. Commun. 242: 626-630, 1998.

[62732] 10486.

[62733] Mori, K.; Yahata, K.; Mukoyama, M.; Suganami, T.; Makino, H.; Nagae,

[62734] T.; Masuzaki, H.; Ogawa, Y.; Sugawara, A.; Nabeshima, Y.; Nakao, K.

[62735] : Disruption of klotho gene causes an abnormal energy homeostasis

[62736] in mice. Biochem. Biophys. Res. Commun. 278: 665-670, 2000.

[62737] 10487.

[62738] Saito, Y.; Nakamura, T.; Ohyama, Y; Suzuki, T.; Iida, A.; Shiraki-Iida,

[62739] T.; Kuro-o, M.; Nabeshima, Y.; Kurabayashi, M.; Nagai, R.: In vivo

[62740] klotho gene delivery protects against endothelial dysfunction in multiple

[62741] risk factor syndrome. Biochem. Biophys. Res. Commun. 276: 767-772,

[62742] 2000.

[62743] 10488.

[62744] Bloom, L.; Horvitz, H. R.: The Caenorhabditis elegans gene unc-76

[62745] and its human homologs define a new gene family involved in axonal
[62746] outgrowth and fasciculation. Proc. Nat. Acad. Sci. 94: 3414-3419,
[62747] 1997.

[62748] 10489.

[62749] Digilio, M. C.; Marino, B.; Giannotti, A.; Dallapiccola, B.: Single
[62750] atrium, atrioventricular canal/postaxial hexodactyly indicating Ellis-van
[62751] Creveld syndrome. Hum. Genet. 96: 251-253, 1996.

[62752] 10490.

[62753] Krawczak, M.; Cooper, D. N.: The human gene mutation database. Trends
[62754] Genet. 13: 121-122, 1997.

[62755] 10491.

[62756] Fujikawa-Adachi, K.; Nishimori, I.; Taguchi, T.; Onishi, S.: Human
[62757] carbonic anhydrase XIV (CA14): cDNA cloning, mRNA expression, and
[62758] mapping to chromosome 1. Genomics 61: 74-81, 1999.

[62759] 10492.

[62760] Vazquez, F.; Hastings, G.; Ortega, M.-A.; Lane, T. F.;
Oikemus,
[62761] S.; Lombardo, M.; Iruela-Arispe, M. L.: METH-1, a human
ortholog
[62762] of ADAMTS-1, and METH-2 are members of a new family
of proteins with
[62763] angio-inhibitory activity. J. Biol. Chem. 274:
23349-23357, 1999.

[62764] 10493.

[62765] Georgiadis, K. E.; Hirohata, S.; Seldin, M. F.; Apte, S. S.:
ADAM-TS8,
[62766] a novel metalloprotease of the ADAM-TS family located on
mouse chromosome
[62767] 9 and human chromosome 11. Genomics 62: 312-315,
1999.

[62768] 10494.

[62769] Whitmore, S. A.; Settasatian, C.; Crawford, J.; Lower, K. M.;
McCallum,
[62770] B.; Seshadri, R.; Cornelisse, C. J.; Moerland, E. W.; Cleton-
Jansen,
[62771] A.-M.; Tipping, A. J.; Mathew, C. G.; Savnio, M.; Savoia, A.;

Verlander,

[62772] P.; Auerbach, A. D.; Van Berkel, C.; Pronk, J. C.; Doggett, N. A.;

[62773] Callen, D. F.: Characterization and screening for mutations of the

[62774] growth arrest-specific 11 (GAS11) and C16orf3 genes at 16q24.3 in

[62775] breast cancer. Genomics 52: 325-331, 1998.

[62776] 10495.

[62777] Sugawara, M.; Nakanishi, T.; Fei, Y.-J.; Huang, W.; Ganap-athy,

[62778] M. E.; Leibach, F. H.; Ganapathy, V.: Cloning of an amino acid transporter

[62779] with functional characteristics and tissue expression pat-tern identical

[62780] to that of system A. J. Biol. Chem. 275: 16473-16477, 2000.

[62781] 10496.

[62782] Akashi, H.; Han, H.-J.; Iizaka, M.; Nakajima, Y.; Furukawa, Y.;

[62783] Sugano, S.; Imai, K.; Nakamura, Y.: Isolation and charac-terization

[62784] of a novel gene encoding a putative seven-span trans-
membrane protein,
[62785] TM7SF3. Cytogenet. Cell Genet. 88: 305-309, 2000.

[62786] 10497.

[62787] Hsieh, J.-C.; Kodjabachian, L.; Rebbert, M. L.; Rattner, A.;
Smallwood,
[62788] P. M.; Samos, C. H.; Nusse, R.; Dawid, I. B.; Nathans, J.: A
new
[62789] secreted protein that binds to Wnt proteins and inhibits
their activites
[62790] (sic). Nature 398: 431-436, 1999.

[62791] 10498.

[62792] Fedi, P.; Bafico, A.; Soria, A. N.; Burgess, W. H.; Miki, T.;
Bottaro,
[62793] D. P.; Kraus, M. H.; Aaronson, S. A.: Isolation and bio-
chemical characterization
[62794] of the human Dkk-1 homologue, a novel inhibitor of
mammalian Wnt signaling. J.
[62795] Biol. Chem. 274: 19465-19472, 1999.

[62796] 10499.

[62797] Krupnik, V. E.; Sharp, J. D.; Jiang, C.; Robison, K.; Chicker-

ing,

[62798] T. W.; Amaravadi, L.; Brown, D. E.; Guyot, D.; Mays, G.; Leiby, K.;

[62799] Chang, B.; Duong, T.; Goodearl, A. D. J.; Gearing, D. P.; Sokol, S.

[62800] Y.; McCarthy, S. A.: Functional and structural diversity of the human

[62801] Dickkopf gene family. Gene 238: 301-313, 1999.

[62802] 10500.

[62803] Roessler, E.; Du, Y.; Glinka, A.; Dutra, A.; Niehrs, C.; Muenke,

[62804] M.: The genomic structure, chromosome location, and analysis of the

[62805] human DKK1 head inducer gene as a candidate for holo-prosencephaly. Cytogenet.

[62806] Cell Genet. 89: 220-224, 2000.

[62807] 10501.

[62808] Chang, C.; Holtzman, D. A.; Chau, S.; Chickering, T.; Woolf, E.

[62809] A.; Holmgren, L. M.; Bodorova, J.; Gearing, D. P.; Holmes, W. E.;

[62810] Brivanlou, A. H.: Twisted gastrulation can function as a

BMP antagonist. Nature 410:
[62811] 483-487, 2001.

[62812] 10502.

[62813] Oelgeschlager, M.; Larrain, J.; Geissert, D.; De Robertis, E. M.
[62814] : The evolutionarily conserved BMP-binding protein Twisted gastrulation
[62815] promotes BMP signalling. Nature 405: 757-763, 2000.

[62816] 10503.

[62817] Ross, J. J.; Shimmi, O.; Vilmos, P.; Petryk, A.; Kim, H.; Gaudenz,
[62818] K.; Hermanson, S.; Ekker, S. C.; O'Connor, M. B.; Marsh, J. L.: Twisted
[62819] gastrulation is a conserved extracellular BMP antagonist. Nature 410:
[62820] 479-483, 2001.

[62821] 10504.

[62822] Scott, I. C.; Blitz, I. L.; Pappano, W. N.; Maas, S. A.; Cho, K.
[62823] W. Y.; Greenspan, D. S.: Homologues of twisted gastrulation are extracellular
[62824] cofactors in antagonism of BMP signalling. Nature 410:

475-478,

[62825] 2001.

[62826] 10505.

[62827] Gatignol, A.; Buckler-White, A.; Berkhout, B.; Jeang, K. T.: Characterization

[62828] of a human TAR RNA-binding protein that activates the HIV-1 LTR. Science 251:

[62829] 1597-1600, 1991.

[62830] 10506.

[62831] Gatignol, A.; Duarte, M.; Daviet, L.; Chang, Y.-N.; Jeang, K.-T.

[62832] : Sequential steps in Tat trans-activation of HIV-1 mediated through

[62833] cellular DNA, RNA, and protein binding factors. Gene Expr. 5: 217-228,

[62834] 1996.

[62835] 10507.

[62836] Kozak, C. A.; Gatignol, A.; Graham, K.; Jeang, K. T.; McBride,

[62837] O. W.: Genetic mapping in human and mouse of the locus encoding TRBP,

[62838] a protein that binds the TAR region of the human immun-
odeficiency
[62839] virus (HIV-1). Genomics 25: 66-72, 1995.

[62840] 10508.

[62841] Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.;
Lichtner,
[62842] P.; Gerbitz, K. D.; Neupert, W.; Brunner, M.; Hofmann, S.:
Genetic
[62843] and structural characterization of the human mitochon-
drial inner membrane
[62844] translocase. J. Molec. Biol. 289: 69-82, 1999.

[62845] 10509.

[62846] Phornphutkul, C.; Anikster, Y.; Huizing, M.; Braun, P.;
Brodie,
[62847] C.; Chou, J. Y.; Gahl, W. A.: The promoter of a lysosomal
membrane
[62848] transporter gene, CTNS, binds Sp-1, shares sequences
with the promoter
[62849] of an adjacent gene, CARKL, and causes cystinosis if mu-
tated in a
[62850] critical region. Am. J. Hum. Genet. 69: 712-721, 2001.

[62851] 10510.

[62852] Lebre, A.-S.; Jamot, L.; Takahashi, J.; Spasskey, N.; Leprince,

[62853] C.; Ravise, N.; Zander, Fujigasaki, H.; Kussel-Andermann, P.; Duyckaerts,

[62854] C.; Camonis, J. H.; Brice, A.: Ataxin-7 interacts with a Cbl-associated

[62855] protein that it recruits into neuronal intranuclear inclusions. Hum.

[62856] Molec. Genet. 10: 10:-1201-1213, 2001.

[62857] 10511.

[62858] Lin, W.-H.; Chiu, K. C.; Chang, H.-M.; Lee, K.-C.; Tai, T.-Y.;

[62859] Chuang, L.-M.: Molecular scanning of the human sorbin and SH3-domain-containing-1

[62860] (SORBS1) gene: positive association of the T228A polymorphism with

[62861] obesity and type 2 diabetes. Hum. Molec. Genet. 10: 1753-1760, 2001.

[62862] 10512.

[62863] Scott, A. F.: Personal Communication. Baltimore, Md. 9/13/2000.

[62864] 10513.

[62865] Nishi, M.; Mizushima, A.; Nakagawara, K.; Takeshima, H.: Characterization

[62866] of human junctophilin subtype genes. Biochem. Biophys. Res. Commun. 273:

[62867] 920-927, 2000.

[62868] 10514.

[62869] Takeshima, H.; Komazaki, S.; Nishi, M.; Iino, M.; Kangawa, K.:

[62870] Junctophilins: a novel family of junctional membrane complex proteins. Molec.

[62871] Cell 6: 11-22, 2000.

[62872] 10515.

[62873] Holmes, S. E.; O'Hearn, E.; Rosenblatt, A.; Callahan, C.; Hwang,

[62874] H. S.; Ingersoll-Ashworth, R. G.; Fleisher, A.; Stevanin, G.; Brice,

[62875] A.; Potter, N. T.; Ross, C. A.; Margolis, R. L.: A repeat ex-pansion

[62876] in the gene encoding junctophilin-3 is associated with Huntington

[62877] disease-like 2. Nature Genet. 29: 377-378, 2001. Note: Erratum:

[62878] Nature Genet. 30: 123 only, 2002.

[62879] 10516.

[62880] Iizaka, M.; Han, H.-J.; Akashi, H.; Furukawa, Y.; Nakajima, Y.;

[62881] Sugano, S.; Ogawa, M.; Nakamura, Y.: Isolation and chromosomal assignment

[62882] of a novel human gene, CORO1C, homologous to coronin-like actin-binding

[62883] proteins. Cytogenet. Cell Genet. 88: 221-224, 2000.

[62884] 10517.

[62885] Bowne, S. J.; Daiger, S. P.; Hims, M. M.; Sohocki, M. M.; Malone,

[62886] K. A.; McKie, A. B.; Heckenlively, J. R.; Birch, D. G.; Inglehearn,

[62887] C. F.; Bhattacharya, S. S.; Bird, A.; Sullivan, L. S.: Mutations

[62888] in the RP1 gene causing autosomal dominant retinitis pigmentosa. Hum.

[62889] Molec. Genet. 8: 2121-2128, 1999.

[62890] 10518.

[62891] Guillonneau, X.; Piriev, N. I.; Danciger, M.; Kozak, C. A.; Cideciyan,
[62892] A. V.; Jacobson, S. G.; Farber, D. B.: A nonsense mutation in a novel
[62893] gene is associated with retinitis pigmentosa in a family linked to
[62894] the RP1 locus. Hum. Molec. Genet. 8: 1541-1546, 1999.

[62895] 10519.

[62896] Iannaccone, A.; Cideciyan, A. V.; Sheffield, V. C.; Stone, E. M.;
[62897] Jacobson, S. G.: Phenotype of chromosome 8q-linked autosomal dominant
[62898] retinitis pigmentosa. (Abstract) Invest. Ophthal. Vis. Sci. 37:
[62899] S345 only, 1996.

[62900] 10520.

[62901] Liu, Q.; Zhou, J.; Daiger, S. P.; Farber, D. B.; Heckenlively,
[62902] J. R.; Smith, J. E.; Sullivan, L. S.; Zuo, J.; Milam, A. H.; Pierce,
[62903] E. A.: Identification and subcellular localization of the RP1 protein

[62904] in human and mouse photoreceptors. Invest. Ophthal. Vis. Sci. 43:
[62905] 22-32, 2002.

[62906] 10521.

[62907] Roderick, T. H.; Chang, B.; Hawes, N. L.; Heckenlively, J. R.:
[62908] A new dominant retinal degeneration (Rd4) associated with a chromosomal
[62909] inversion in the mouse. Genomics 42: 393-396, 1997.

[62910] 10522.

[62911] Chavez, R. A.; Gray, A. T.; Zhao, B. B.; Kindler, C. H.; Mazurek,
[62912] M. J.; Mehta, Y.; Forsayeth, J. R.; Yost, C. S.: TWIK-2, a new weak
[62913] inward rectifying member of the tandem pore domain potassium channel
[62914] family. J. Biol. Chem. 274: 7887-7892, 1999.

[62915] 10523.

[62916] Gray, A. T.; Kindler, C. H.; Sampson, E. R.; Yost, C. S.: As-
[62917] signment of KCNK6 encoding the human weak inward rectifier

potassium channel
[62918] TWIK-2 to chromosome band 19q13.1 by radiation hybrid mapping. Cytogenet.
[62919] Cell Genet. 84: 190-191, 1999.

[62920] 10524.

[62921] Pountney, D. J.; Gulkarov, I.; Vega-Saenz de Miera, E.; Holmes,
[62922] D.; Saganich, M.; Rudy, B.; Artman, M.; Coetzee, W. A.: Identification
[62923] and cloning of TWIK-originated similarity sequence (TOSS): a novel
[62924] human 2-pore K(+) channel principal subunit. FEBS Lett. 450: 191-196,
[62925] 1999.

[62926] 10525.

[62927] Salinas, M.; Reyes, R.; Lesage, F.; Fosset, M.; Heurteaux, C.;
[62928] Romey, G.; Lazdunski, M.: Cloning of a new mouse two-P domain channel
[62929] subunit and a human homologue with a unique pore structure. J. Biol.
[62930] Chem. 274: 11751-11760, 1999.

[62931] 10526.

[62932] Banikazemi, M.; Diaz, G. A.; Voussough, P.; Jalali, M.; Desnick,
[62933] R. J.; Gelb, B. D.: Localization of the thiamine-responsive megaloblastic
[62934] anemia syndrome locus to a 1.4-cM region of 1q23. Molec. Genet. Metab. 66:
[62935] 193-198, 1999.

[62936] 10527.

[62937] Diaz, G. A.; Banikazemi, M.; Oishi, K.; Desnick, R. J.; Gelb, B.
[62938] D.: Mutations in a new gene encoding a thiamine transporter cause
[62939] thiamine-responsive megaloblastic anaemia syndrome. Nature Genet. 22:
[62940] 309-312, 1999.

[62941] 10528.

[62942] Fleming, J. C.; Tartaglini, E.; Steinkamp, M. P.; Schorderet, D.
[62943] F.; Cohen, N.; Neufeld, E. J.: The gene mutated in thiamine-responsive

[62944] anaemia with diabetes and deafness (TRMA) encodes a functional thiamine
[62945] transporter. Nature Genet. 22: 305-308, 1999.

[62946] 10529.

[62947] Raz, T.; Labay, V.; Baron, D.; Szargel, R.; Anbinder, Y.; Barrett,
[62948] T.; Rabl, W.; Viana, M. B.; Mandel, H.; Baruchel, A.; Cayuela, J.-M.;
[62949] Cohen, N.: The spectrum of mutations, including four novel ones,
[62950] in the thiamine-responsive megaloblastic anemia gene SLC19A2 of eight
[62951] families. Hum. Mutat. 16: 37-43, 2000.

[62952] 10530.

[62953] Rindi, G.; Patrini, C.; Laforenza, U.; Mandel, H.; Berant, M.;

[62954] Viana, M. B.; Poggi, V.; Zarra, A. N.: Further studies on erythrocyte
[62955] thiamin transport and phosphorylation in seven patients with thiamin-responsive
[62956] megaloblastic anaemia. J. Inherit. Metab. Dis. 17: 667-677, 1994.

[62957] 10531.

[62958] Scharfe, C.; Hauschild, M.; Klopstock, T.; Janssen, A. J. M.; Heidemann,

[62959] P. H.; Meitinger, T.; Jaksch, M.: A novel mutation in the thiamine

[62960] responsive megaloblastic anaemia gene SLC19A2 in a patient with deficiency

[62961] of respiratory chain complex I. J. Med. Genet. 37: 669-673, 2000.

[62962] 10532.

[62963] Stagg, A. R.; Fleming, J. C.; Baker, M. A.; Sakamoto, M.; Cohen,

[62964] N.; Neufeld, E. J.: Defective high-affinity thiamine transporter

[62965] leads to cell death in thiamine-responsive megaloblastic anemia syndrome

[62966] fibroblasts. J. Clin. Invest. 103: 723-729, 1999.

[62967] 10533.

[62968] Barbetti, F.; Rocchi, M.; Bossolasco, M.; Cordera, R.; Sbraccia,

[62969] P.; Finelli, P.; Consalez, G. G.: The human skeletal muscle glycogenin

[62970] gene: cDNA, tissue expression, and chromosomal localization. Biochem.

[62971] Biophys. Res. Commun. 220: 72-77, 1996.

[62972] 10534.

[62973] Wang, D. W.; VanDeCarr, D.; Ruben, P. C.; George, A. L., Jr.;

[62974] Bennett, P. B.: Functional consequences of a domain 1/S6 segment

[62975] sodium channel mutation associated with painful congenital myotonia. FEBS

[62976] Lett. 448: 231-234, 1999.

[62977] 10535.

[62978] Wang, J.; Rojas, C. V.; Zhou, J.; Schwartz, L. S.; Nicholas, H.;

[62979] Hoffman, E. P.: Sequence and genomics structure of the human adult

[62980] skeletal muscle sodium channel alpha subunit gene on 17q. Biochem.

[62981] Biophys. Res. Commun. 182: 794-801, 1992.

[62982] 10536.

[62983] Machesky, L. M.; Reeves, E.; Wientjes, F.; Mattheyse, F. J.;

Grogan,
[62984] A.; Totty, N. F.; Burlingame, A. L.; Hsuan, J. J.; Segal, A. W.:
[62985] Mammalian actin-related protein 2/3 complex localizes to regions of
[62986] lamellipodial protrusion and is composed of evolutionarily conserved
[62987] proteins. Biochem. J. 328: 105-112, 1997.
[62988] 10537.
[62989] Wada, T.; Yoshikawa, Y.; Tokuyama, S.; Kuwabara, M.; Akita, H.;
[62990] Miyagi, T.: Cloning, expression, and chromosomal mapping of a human
[62991] ganglioside sialidase. Biochem. Biophys. Res. Commun. 261: 21-27,
[62992] 1999.
[62993] 10538.
[62994] Bigner, S. H.; Vogelstein, B.: Cytogenetics and molecular genetics
[62995] of malignant gliomas and medulloblastoma. Brain Path. 1: 12-18,
[62996] 1990.

[62997] 10539.

[62998] Yoshida, A.; Minowa, M. T.; Takamatsu, S.; Hara, T.; Oguri, S.;

[62999] Ikenaga, H.; Takeuchi, M.: Tissue specific expression and chromosomal

[63000] mapping of a human UDP-N-acetylglucosamine:alpha-1,3-D-mannoside beta-1,4-N-acetylglucosaminyltransferase. Glycobiology 9:

[63001] 303-310, 1999.

[63002] 10540.

[63003] Boelens, W. C.; Van Boekel, M. A.; De Jong, W. W.: HspB3, the

[63004] most deviating of the six known human small heat shock proteins. Biochim.

[63005] Biophys. Acta 1388: 513-516, 1998.

[63006] 10541.

[63007] Lam, W. Y.; Tsui, S. K. W.; Law, P. T. W.; Luk, S. C.; Fung, K.

[63008] P.; Lee, C. Y.; Waye, M. M. Y.: Isolation and characterization of

[63009] a human heart cDNA encoding a new member of the small

heat shock protein

[63010] family--HSPL27. Biochim. Biophys. Acta 1314: 120-124, 1996.

[63011] 10542.

[63012] Makoff, A.; Lelchuk, R.; Oxer, M.; Harrington, K.; Emson, P.:

[63013] Molecular characterization and localization of human metabotropic

[63014] glutamate receptor type 4. Molec. Brain Res. 37: 239-248, 1996.

[63015] 10543.

[63016] Pekhletski, R.; Gerlai, R.; Overstreet, L. S.; Huang, X. P.; Agopyan,

[63017] N.; Slater, N. T.; Abramow-Newerly, W.; Roder, J. C.; Hampson, D.

[63018] R.: Impaired cerebellar synaptic plasticity and motor per-formance

[63019] in mice lacking the mGluR4 subtype of metabotropic glu-tamate receptor. J.

[63020] Neurosci. 16: 6364-6373, 1996.

[63021] 10544.

[63022] Barbon, A.; Ferraboli, S.; Barlati, S.: Assignment of the hu-

man

[63023] metabotropic glutamate receptor gene GRM7 to chromo-
some 3p26.1-p25.2

[63024] by radiation hybrid mapping. Cytogenet. Cell Genet. 88:
288 only,

[63025] 2000.

[63026] 10545.

[63027] Shore, D.; Squire, M.; Nasmyth, K. A.: Characterization of
two

[63028] genes required for the position-effect control of yeast
mating-type

[63029] genes. EMBO J. 3: 2817-2823, 1984.

[63030] 10546.

[63031] Tanny, J. C.; Dowd, G. J.; Huang, J.; Hilz, H.; Moazed, D.:
An

[63032] enzymatic activity in the yeast Sir2 protein that is essential
for

[63033] gene silencing. Cell 99: 735-745, 1999.

[63034] 10547.

[63035] Verhaagh, S.; Schweifer, N.; Barlow, D. P.; Zwart, R.:
Cloning

[63036] of the mouse and human solute carrier 22a3 (Slc22a3/SLC22A3) identifies
[63037] a conserved cluster of three organic cation transporters on mouse
[63038] chromosome 17 and human 6q26-q27. Genomics 55: 209-218, 1999.
[63039] 10548.

[63040] Gorlich, D.; Hartmann, E.; Prehn, S.; Rapoport, T. A.: A protein
[63041] of the endoplasmic reticulum involved early in polypeptide translocation. Nature 357:
[63042] 47-52, 1992.

[63043] 10549.

[63044] Liu, J.; Shworak, N. W.; Sinay, P.; Schwartz, J. J.; Zhang, L.;

[63045] Fritze, L. M.; Rosenberg, R. D.: Expression of heparan sulfate D-glucosaminyl
[63046] 3-O-sulfotransferase isoforms reveals novel substrate specificities. J.
[63047] Biol. Chem. 274: 5185-5192, 1999.

[63048] 10550.

[63049] Shworak, N. W.; Liu, J.; Petros, L. M.; Zhang, L.; Kobayashi,

M.;

[63050] Copeland, N. G.; Jenkins, N. A.; Rosenberg, R. D.: Multiple isoforms

[63051] of heparan sulfate D-glucosaminyl 3-O-sulfotransferase: isolation,

[63052] characterization, and expression of human cDNAs and identification

[63053] of distinct genomic loci. J. Biol. Chem. 274: 5170-5184, 1999.

[63054] 10551.

[63055] McIlhatton, M. A.; Burrows, J. F.; Donaghy, P. G.; Chandu-loy, S.;

[63056] Johnston, P. G.; Russell, S. E. H.: Genomic organization, complex

[63057] splicing pattern and expression of a human septin gene on chromosome

[63058] 17q25.3. Oncogene 20: 5930-5939, 2001.

[63059] 10552.

[63060] Osaka, M.; Rowley, J. D.; Zeleznik-Le, N. J.: MSF (MLL septin-like

[63061] fusion), a fusion partner gene of MLL, in a therapy-related acute

[63062] myeloid leukemia with a t(11;17)(q23;q25). Proc. Nat. Acad. Sci. 96:
[63063] 6428-6433, 1999.

[63064] 10553.

[63065] Russell, S. E. H.; McIlhatton, M. A.; Burrows, J. F.; Donaghy,

[63066] P. G.; Chanduloy, S.; Petty, E. M.; Kalikin, L. M.; Church, S. W.;
[63067] McIlroy, S.; Harkin, D. P.; Keilty, G. W.; Cranston, A. N.; Weissenbach,
[63068] J.; Hickey, I.; Johnston, P. G.: Isolation and mapping of a human
[63069] septin gene to a region on chromosome 17q, commonly deleted in sporadic
[63070] epithelial ovarian tumors. Cancer Res. 60: 4729-4734, 2000.
[63071] 10554.

[63072] Karpinski, B. A.; Morle, G. D.; Huggenvik, J.; Uhler, M. D.; Leiden,
[63073] J. M.: Molecular cloning of human CREB-2: an ATF/CREB transcription
[63074] factor that can negatively regulate transcription from the cAMP response

[63075] element. Proc. Nat. Acad. Sci. 89: 4820-4824, 1992.

[63076] 10555.

[63077] Kasukabe, T.; Kobayashi, H.; Kaneko, Y.; Okabe-Kado, J.; Honma,

[63078] Y.: Identity of human normal counterpart (MmTRA1b) of mouse leukemogenesis-associated

[63079] gene (MmTRA1a) product as a plasma membrane phospholipid scramblase

[63080] and chromosome mapping of the human MmTRA1b/phospholipid scramblase

[63081] gene. Biochem. Biophys. Res. Commun. 249: 449-455, 1998.

[63082] 10556.

[63083] Zhou, Q.; Sims, P. J.; Wiedmer, T.: Identity of a conserved motif

[63084] in phospholipid scramblase that is required for Ca2+-accelerated transbilayer

[63085] movement of membrane phospholipids. Biochemistry 37: 2356-2360,

[63086] 1998.

[63087] 10557.

[63088] Zhou, Q.; Zhao, J.; Stout, J. G.; Luhm, R. A.; Wiedmer, T.; Sims,

[63089] P. J.: Molecular cloning of human plasma membrane phospholipid scramblase:

[63090] a protein mediating transbilayer movement of plasma membrane phospholipids. J.

[63091] Biol. Chem. 272: 18240-18244, 1997.

[63092] 10558.

[63093] Robinson, R. C.; Turbedsky, K.; Kaiser, D. A.; Marchand, J.-B.;

[63094] Higgs, H. N.; Choe, S.; Pollard, T. D.: Crystal structure of Arp2/3

[63095] complex. Science 294: 1679-1684, 2001.

[63096] 10559.

[63097] Volkmann, N.; Amann, K. J.; Stoilova-McPhie, S.; Egile, C.; Winter,

[63098] D. C.; Hazelwood, L.; Heuser, J. E.; Li, R.; Pollard, T. D.; Hanein,

[63099] D.: Structure of Arp2/3 complex in its activated state and in actin

[63100] filament branch junctions. Science 293: 2456-2459, 2001.

[63101] 10560.

[63102] Welch, M. D.; DePace, A. H.; Verma, S.; Iwamatsu, A.; Mitchison,

[63103] T. J.: The human Arp2/3 complex is composed of evolutionarily conserved

[63104] subunits and is localized to cellular regions of dynamic actin filament

[63105] assembly. J. Cell Biol. 138: 375-384, 1997.

[63106] 10561.

[63107] Odeberg, J.; Rosok, O.; Gudmundsson, G. H.; Ahmadian, A.; Roshani,

[63108] L.; Williams, C.; Larsson, C.; Ponten, F.; Uhlen, M.; Asheim, H.-C.;

[63109] Lundeberg, J.: Cloning and characterization of ZNF189, a novel human

[63110] Kruppel-like zinc finger gene localized to chromosome 9q22-q31. Genomics 50:

[63111] 213-221, 1998.

[63112] 10562.

[63113] Michel, J. J.; Xiong, Y.: Human CUL-1, but not other cullin family

[63114] members, selectively interacts with SKP1 to form a complex with SKP2
[63115] and cyclin A. Cell Growth Differ. 9: 435-449, 1998.

[63116] 10563.

[63117] Du, M.; Sansores-Garcia, L.; Zu, Z.; Wu, K. K.: Cloning and expression
[63118] analysis of a novel salicylate suppressible gene, Hs-CUL-3, a member
[63119] of cullin/Cdc53 family. J. Biol. Chem. 273: 24289-24292, 1998.
[63120] 10564.

[63121] Bao, S.; Shen, X.; Shen, K.; Liu, Y.; Wang, X.-F.: The mammalian
[63122] Rad24 homologous to yeast Saccharomyces cerevisiae Rad24 and Schizosaccharomyces
[63123] pombe Rad17 is involved in DNA damage checkpoint. Cell Growth Diff. 9:
[63124] 961-967, 1998.

[63125] 10565.

[63126] Bao, S.; Chang, M.-S.; Auclair, D.; Sun, Y.; Wang, Y.; Wong, W.-K.;

[63127] Zhang, J.; Liu, Y.; Qian, X.; Sutherland, R.; Magi-Galluzi, C.; Weisberg,
[63128] E.; Cheng, E. Y. S.; Hao, L.; Sasaki, H.; Campbell, M. S.; Kraeft,
[63129] S.-K.; Loda, M.; Lo, K.-M.; Chen, L. B.: Hrad17, a human homologue
[63130] of the Schizosaccharomyces pombe checkpoint gene rad17, is overexpressed
[63131] in colon carcinoma. Cancer Res. 59: 2023-2028, 1999.

[63132] 10566.

[63133] Bluyssen, H. A. R.; Naus, N. C.; van Os, R. I.; Jaspers, I.; Hoeijmakers,
[63134] J. H. J.; de Klein, A.: Human and mouse homologues of the Schizosaccharomyces
[63135] pombe rad17+ cell cycle checkpoint control gene. Genomics 55: 219-228,
[63136] 1999.

[63137] 10567.

[63138] Brunner, H. G.; Hamel, B. C. J.; van Bokhoven, H.: P63 gene mutations
[63139] and human developmental syndromes. Am. J. Med. Genet. 112: 284-290,

[63140] 2002.

[63141] 10568.

[63142] Crackower, M. A.; Motoyama, J.; Tsui, L.-C.: Defect in the maintenance

[63143] of the apical ectodermal ridge in the Dactylaplasia mouse. Dev. Biol. 201:

[63144] 78-79, 1998.

[63145] 10569.

[63146] Duijf, P. H. G.; Vanmolkot, K. R. J.; Propping, P.; Friedl, W.;

[63147] Krieger, E.; McKeon, F.; Dotsch, V.; Brunner, H. G.; van Bokhoven,

[63148] H.: Gain-of-function mutation in ADULT syndrome reveals the presence

[63149] of a second transactivation domain in p63. Hum. Molec. Genet. 11:

[63150] 799-804, 2002.

[63151] 10570.

[63152] Hibi, K.; Trink, B.; Patturajan, M.; Westra, W. H.; Caballero,

[63153] O. L.; Hill, D. E.; Ratovitski, E. A.; Jen, J.; Sidransky, D.: AIS

[63154] is an oncogene amplified in squamous cell carcinoma.

Proc. Nat. Acad.

[63155] Sci. 97: 5462-5467, 2000.

[63156] 10571.

[63157] Mills, A. A.; Zheng, B.; Wang, X. J.; Vogel, H.; Roop, D. R.;

[63158] Bradley, A.: p63 is a p53 homologue required for limb and epidermal

[63159] morphogenesis. Nature 398: 708-713, 1999.

[63160] 10572.

[63161] Osada, M.; Ohba, M.; Kawahara, C.; Ishioka, C.; Kanamaru, R.;

[63162] Katoh, I.; Ikawa, Y.; Nimura, Y.; Nakagawara, A.; Obinata, M.; Ikawa,

[63163] S.: Cloning and functional analysis of human p51, which structurally

[63164] and functionally resembles p53. Nature Med. 4: 839-843, 1998.

[63165] 10573.

[63166] Senoo, M.; Seki, N.; Ohira, M.; Sugano, S.; Watanabe, M.; Tachibana,

[63167] M.; Tanaka, T.; Shinkai, Y.; Kato, H.: A second p53-related protein,

[63168] p73L, with high homology to p73. Biochem. Biophys. Res. Commun. 248:

[63169] 603-607, 1998.

[63170] 10574.

[63171] Trink, B.; Okami, K.; Wu, L.; Sriuranpong, V.; Jen, J.; Sidransky,

[63172] D. :Nature Med. 4: 747-748, 1998.

[63173] 10575.

[63174] Yang, A.; Kaghad, M.; Wang, Y.; Gillett, E.; Fleming, M. D.; Dotsch,

[63175] V.; Andrews, N. C.; Caput, D.; McKeon, F.: p63, a p53 ho-

molog at

[63176] 3q27-29, encodes multiple products with transactivating, death-inducing,

[63177] and dominant-negative activities. Molec. Cell 2: 305-316, 1998.

[63178] 10576.

[63179] Yang, A.; Schweitzer, R.; Sun, D.; Kaghad, M.; Walker, N.; Bronson,

[63180] R. T.; Tabin, C.; Sharpe, A.; Caput, D.; Crum, C.; McKeon, F.: p63

[63181] is essential for regenerative proliferation in limb, cranio-facial
[63182] and epithelial development. Nature 398: 714-718, 1999.

[63183] 10577.

[63184] van Bokhoven, H.; Hamel, B. C.; Bamshad, M.; Sangiorgi, E.; Gurrieri,
[63185] F.; Duijf, P. H.; Vanmolkot, K. R.; van Beusekom, E.; van Beersum,
[63186] S. E.; Celli, J.; Merkx, G. F.; Tenconi, R.; and 13 others: p63
[63187] gene mutations in EEC syndrome, limb-mammary syn-drome, and isolated
[63188] split hand-foot malformation suggest a genotype-phe-notype correlation. Am.
[63189] J. Hum. Genet. 69: 481-492, 2001.

[63190] 10578.

[63191] Loijens, J. C.; Anderson, R. A.: Type I phosphatidylinosi-tol-4-phosphate
[63192] 5-kinases are distinct members of this novel lipid kinase family. J.
[63193] Biol. Chem. 271: 32937-32943, 1996.

[63194] 10579.

[63195] Xie, Y.; Zhu, L.; Zhao, G.: Assignment of type I phos-
phatidylinositol-4-phosphate
[63196] 5-kinase (PIP5K1A) to human chromosome bands
1q22-q24 by in situ hybridization. Cytogenet.
[63197] Cell Genet. 88: 197-199, 2000.

[63198] 10580.

[63199] Seki, N.; Sugano, S.; Suzuki, Y.; Nakagawara, A.; Ohira, M.;
Muramatsu,
[63200] M.; Saito, T.; Hori, T.: Isolation, tissue expression, and
chromosomal
[63201] assignment of human RGS5, a novel G-protein signaling
regulator gene. Hum.
[63202] Genet. 43: 202-205, 1998.

[63203] 10581.

[63204] Seelig, H. P.; Moosbrugger, I.; Ehrfeld, H.; Fink, T.; Renz,
M.;
[63205] Genth, E.: The major dermatomyositis-specific Mi-2 au-
toantigen is
[63206] a presumed helicase involved in transcriptional activation.
Arthritis

[63207] Rheum. 38: 1389-1399, 1995.

[63208] 10582.

[63209] Wade, P. A.; Gegonne, A.; Jones, P. L.; Ballestar, E.; Aubry, F.;

[63210] Wolffe, A. P.: Mi-2 complex couples DNA methylation to chromatin

[63211] remodelling and histone deacetylation. Nature Genet. 23: 62-66,

[63212] 1999.

[63213] 10583.

[63214] Zhang, Y.; LeRoy, G.; Seelig, H.-P.; Lane, W. S.; Reinberg, D.

[63215] : The dermatomyositis-specific autoantigen Mi2 is a component of a

[63216] complex containing histone deacetylase and nucleosome remodeling activities. Cell 95:

[63217] 279-289, 1998.

[63218] 10584.

[63219] Parma, J.; Duprez, L.; Van Sande, J.; Cochaux, P.; Gervy, C.;

[63220] Mockel, J.; Dumont, J.; Vassart, G.: Somatic mutations in the thyrotropin

[63221] receptor gene cause hyperfunctioning thyroid adenomas. Nature 365:
[63222] 649-651, 1993.

[63223] 10585.

[63224] Parma, J.; Duprez, L.; Van Sande, J.; Hermans, J.; Rocmans, P.;
[63225] Van Vliet, G.; Costagliola, S.; Rodien, P.; Dumont, J. E.; Vassart,
[63226] G.: Diversity and prevalence of somatic mutations in the thyrotropin
[63227] receptor and Gs-alpha genes as a cause of toxic thyroid adenomas. J.
[63228] Clin. Endocr. Metab. 82: 2695-2701, 1997.

[63229] 10586.

[63230] Rodien, P.; Bremont, C.; Luton, J.-P.; Raffin-Sanson, M.-L.; Parma,
[63231] J.; Duprez, L.; Vassart, G.: De la promiscuite chez les hormones
[63232] glycoproteiques: hyperthyroidie gestationnelle familiale par mutation
[63233] du recepteur de la TSH. Med./Sci. 15: 713-717, 1999.

[63234] 10587.

[63235] Rodien, P.; Bremont, C.; Raffin Sanson, M.-L.; Parma, J.; Van

[63236] Sande, J.; Costagliola, S.; Luton, J.-P.; Vassart, G.; Duprez, L.

[63237] : Familial gestational hyperthyroidism caused by a mutant thyrotropin

[63238] receptor hypersensitive to human chorionic gonadotropin. New Eng.

[63239] J. Med. 339: 1823-1826, 1998.

[63240] 10588.

[63241] Rousseau-Merck, M. F.; Misrahi, M.; Loosfelt, H.; Atger, M.; Milgrom,

[63242] E.; Berger, R.: Assignment of the human thyroid stimulating hormone

[63243] receptor (TSHR) gene to chromosome 14q31. Genomics 8: 233-236, 1990.

[63244] 10589.

[63245] Russo, D.; Betterle, C.; Arturi, F.; Chiefari, E.; Girelli, M.

[63246] E.; Filetti, S.: A novel mutation in the thyrotropin (TSH) receptor

[63247] gene causing loss of TSH binding but constitutive receptor activation

[63248] in a family with resistance to TSH. J. Clin. Endocr. Metab. 85:

[63249] 4238-4242, 2000.

[63250] 10590.

[63251] Russo, D.; Tumino, S.; Arturi, F.; Vigneri, P.; Grasso, G.; Pontecorvi,

[63252] A.; Filetti, S.; Belfiore, A.: Detection of an activating mutation

[63253] of the thyrotropin receptor in a case of an autonomously hyperfunctioning

[63254] thyroid insular carcinoma. J. Clin. Endocr. Metab. 82: 735-738,

[63255] 1997.

[63256] 10591.

[63257] Tonacchera, M.; Agretti, P.; Chiovato, L.; Rosellini, V.; Ceccarini,

[63258] G.; Perri, A.; Viacava, P.; Naccarato, A. G.; Miccoli, P.; Pinchera,

[63259] A.; Vitti, P.: Activating thyrotropin receptor mutations are present

[63260] in nonadenomatous hyperfunctioning nodules of toxic or autonomous
[63261] multinodular goiter. J. Clin. Endocr. Metab. 85: 2270-2274, 2000.
[63262] 10592.

[63263] Tonacchera, M.; Agretti, P.; Pinchera, A.; Rosellini, V.; Perri,
[63264] A.; Collecchi, P.; Vitti, P.; Chiovato, L.: Congenital hypothyroidism
[63265] with impaired thyroid response to thyrotropin (TSH) and absent circulating
[63266] thyroglobulin: evidence for a new inactivating mutation of the TSH
[63267] receptor gene. J. Clin. Endocr. Metab. 85: 1001-1008, 2000.
[63268] 10593.

[63269] Trulzsch, B.; Nebel, T.; Paschke, R.: The thyrotropin receptor
[63270] mutation database. (Editorial) Thyroid 9: 521-522, 1999.

[63271] 10594.

[63272] Van Sande, J.; Parma, J.; Tonacchera, M.; Swillens, S.; Dumont,

[63273] J.; Vassart, G.: Genetic basis of endocrine disease: Somatic and
[63274] germline mutations of the TSH receptor gene in thyroid diseases. J.
[63275] Clin. Endocr. Metab. 80: 2577-2585, 1995.

[63276] 10595.

[63277] Vassart, G.; Parmentier, M.; Libert, F.; Dumont, J.: Molecu-lar
[63278] genetics of the thyrotropin receptor. Trends Endocr. Metab. 2: 151-156,
[63279] 1991.

[63280] 10596.

[63281] Wadsworth, H. L.; Chazenbalk, G. D.; Nagayama, Y.; Russo, D.;
[63282] Rapoport, B.: An insertion in the human thyrotropin re-ceptor critical
[63283] for high affinity hormone binding. Science 249: 1423-1425, 1990.
[63284] 10597.

[63285] Aihara, T.; Miyoshi, Y.; Koyama, K.; Suzuki, M.; Takahashi, E.;

[63286] Monden, M.; Nakamura, Y.: Cloning and mapping of SMARCA5 encoding

[63287] hSNF2H, a novel human homologue of Drosophila ISWI. Cytogenet. Cell

[63288] Genet. 81: 191-193, 1998.

[63289] 10598.

[63290] Bochar, D. A.; Savard, J.; Wang, W.; Lafleur, D. W.; Moore, P.;

[63291] Cote, J.; Shiekhattar, R.: A family of chromatin remodeling factors

[63292] related to Williams syndrome transcription factor. Proc. Nat. Acad.

[63293] Sci. 97: 1038-1043, 2000.

[63294] 10599.

[63295] Bozhenok, L.; Wade, P. A.; Varga-Weisz, P.: WSTF-ISWI chromatin

[63296] remodeling complex targets heterochromatic replication foci. EMBO

[63297] J. 21: 2231-2241, 2002.

[63298] 10600.

[63299] LeRoy, G.; Orphanides, G.; Lane, W. S.; Reinberg, D.: Re-

quirement

[63300] of RSF and FACT for transcription of chromatin templates in vitro. Science 282:

[63301] 1900-1904, 1998.

[63302] 10601.

[63303] Poot, R. A.; Dellaire, G.; Hulsmann, B. B.; Grimaldi, M. A.; Corona,

[63304] D. F. V.; Becker, P. B.; Bickmore, W. A.; Varga-Weisz, P. D.: HuCHRAC,

[63305] a human ISWI chromatin remodelling complex contains hACF1 and two

[63306] novel histone-fold proteins. EMBO J. 19: 3377-3387, 2000.

[63307] 10602.

[63308] Burwinkel, B.; Kreuder, J.; Schweitzer, S.; Vorgerd, M.; Gempel,

[63309] K.; Gerbitz, K.-D.; Kilimann, M. W.: Carnitine transporter OCTN2

[63310] mutations in systemic primary carnitine deficiency: a novel Arg169Gln

[63311] mutation and a recurrent Arg282ter mutation associated with an unconventional

[63312] splicing abnormality. Biochem. Biophys. Res. Commun. 261: 484-487,
[63313] 1999.

[63314] 10603.

[63315] Alberati-Giani, D.; Cesura, A. M.; Broger, C.; Warren, W. D.; Rover,
[63316] S.; Malherbe, P.: Cloning and functional expression of human kynurenine
[63317] 3-monooxygenase. FEBS Lett. 410: 407-412, 1997.

[63318] 10604.

[63319] Nagler, D. K.; Sulea, T.; Menard, R.: Full-length cDNA of human
[63320] cathepsin F predicts the presence of a cystatin domain at the N-terminus
[63321] of the cysteine protease zymogen. Biochem. Biophys. Res. Commun. 257:
[63322] 313-318, 1999.

[63323] 10605.

[63324] Santamaria, I.; Velasco, G.; Pendas, A. M.; Paz, A.; Lopez-Otin,
[63325] C.: Molecular cloning and structural and functional char-

acterization
[63326] of human cathepsin F, a new cysteine proteinase of the papain family
[63327] with a long propeptide domain. J. Biol. Chem. 274: 13800-13809,
[63328] 1999.

[63329] 10606.

[63330] Wang, B.; Shi, G.-P.; Yao, P. M.; Li, Z.; Chapman, H. A.; Bromme,
[63331] D.: Human cathepsin F: molecular cloning, functional expression,
[63332] tissue localization, and enzymatic characterization. J. Biol. Chem. 273:
[63333] 32000-32008, 1998.

[63334] 10607.

[63335] Boyer, S. H.; Fainer, D. C.; Watson-Williams, E. J.: Lactate dehydrogenase
[63336] variant from human blood: evidence for molecular subunits. Science 141:
[63337] 642-643, 1963.

[63338] 10608.

[63339] Krolewski, J. J.; Lee, R.; Eddy, R.; Shows, T. B.; Dalla-Favera,
[63340] R.: Identification and chromosomal mapping of new human tyrosine
[63341] kinase genes. Oncogene 5: 277-282, 1990.

[63342] 10609.

[63343] Virtaneva, K. I.; Angelisova, P.; Baumruker, T.; Horejsi, V.; Nevanlinna,
[63344] H.; Schroder, J.: The genes for CD37, CD53, and R2, all members of
[63345] a novel gene family, are located on different chromosomes. Immunogenetics 37:
[63346] 461-465, 1993.

[63347] 10610.

[63348] Amiot, M.: Identification and analysis of cDNA clones encoding
[63349] CD53: a pan-leukocyte antigen related to membrane transport proteins.
[63350] J. Immun. 145: 4322-4325, 1990.

[63351] 10611.

[63352] Angelisova, P.; Vlcek, C.; Stefanova, I.; Lipoldova, M.;

Horejsi,
[63353] V.: The human leucocyte surface antigen CD53 is a protein structurally
[63354] similar to the CD37 and MRC OX-44 antigens. Immunogenetics 32:
[63355] 281-285, 1990.

[63356] 10612.

[63357] Gonzalez, M. E.; Pardo-Manuel de Villena, F.; Fernandez-Ruiz, E.;
[63358] Rodriguez de Cordoba, S.; Lazo, P. A.: The human CD53 gene, coding
[63359] for a four transmembrane domain protein, maps to chromosomal region
[63360] 1p13. Genomics 18: 725-728, 1993.

[63361] 10613.

[63362] Hadam, M. R.: Cluster report: CD53. In: Knapp, W.; Dorken, B.;
[63363] Gilks, W. R.; Rieber, E. P.; Schmidt, R. E.; Stein, H.; von dem Borne,
[63364] A. E. G. K.: Leucocyte Typing IV. White Cell Differentiation Antigens.
[63365] Oxford: Oxford Univ. Press (pub.) 1989. Pp. 674-678.

[63366] 10614.

[63367] Taguchi, T.; Bellacosa, A.; Zhou, J.-Y.; Gilbert, D. J.; Lazo,

[63368] P. A.; Copeland, N. G.; Jenkins, N. A.; Tsichlis, P. N.; Testa,
J.

[63369] R.: Chromosomal localization of the Ox-44 (CD53) leuko-
cyte antigen

[63370] gene in man and rodents. Cytogenet. Cell Genet. 64:
217-221, 1993.

[63371] 10615.

[63372] Yoshimura, K.; Toibana, A.; Nakahama, K.: Human
lysozyme: sequencing

[63373] of a cDNA, and expression and secretion by Saccha-
romyces cerevisiae. Biochem.

[63374] Biophys. Res. Commun. 150: 794-801, 1988.

[63375] 10616.

[63376] Salmikangas, P.; Mykkanen, O.-M.; Gronholm, M.; Heiska,
L.; Kere,

[63377] J.; Carpen, O.: Myotilin, a novel sarcomeric protein with
two Ig-like

[63378] domains, is encoded by a candidate gene for limb-girdle
muscular dystrophy. Hum.

[63379] Molec. Genet. 8: 1329-1336, 1999.

[63380] 10617.

[63381] Hauser, M. A.; Horrigan, S. K.; Salmikangas, P.; Torian, U. M.;

[63382] Viles, K. D.; Dancel, R.; Tim, R. W.; Taivainen, A.; Bartoloni, L.;

[63383] Gilchrist, J. M.; Stajich, J. M.; Gaskell, P. C.; Gilbert, J. R.;

[63384] Vance, J. M.; Pericak-Vance, M. A.; Carpen, O.; Westbrook, C. A.;

[63385] Speer, M. C.: Myotilin is mutated in limb girdle muscular dystrophy

[63386] 1A. Hum. Molec. Genet. 9: 2141-2147, 2000.

[63387] 10618.

[63388] Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G. M.;

[63389] Romijn, J. C.; Trapman, J.: Characterization of the prostate-specific

[63390] antigen gene: a novel human kallikrein-like gene. Biochem. Biophys.

[63391] Res. Commun. 159: 95-102, 1989.

[63392] 10619.

[63393] Santoro, M.; Carlomagno, F.; Hay, I. D.; Herrmann, M. A.; Grieco,
[63394] M.; Melillo, R.; Pierotti, M. A.; Bongarzone, I.; Della Porta, G.;
[63395] Berger, N.; Peix, J. L.; Paulin, C.; Fabien, N.; Vecchio, G.; Jenkins,
[63396] R. B.; Fusco, A.: Ret oncogene activation in human thyroid neoplasms
[63397] is restricted to the papillary cancer subtype. J. Clin. Invest. 89:
[63398] 1517-1522, 1992.

[63399] 10620.

[63400] Santoro, M.; Carlomagno, F.; Romano, A.; Bottaro, D. P.; Dathan,
[63401] N. A.; Grieco, M.; Fusco, A.; Vecchio, G.; Matoskova, B.; Kraus, M.
[63402] H.; Di Fiore, P. P.: Activation of RET as a dominant trans-forming
[63403] gene by germline mutations of MEN2A and MEN2B. Sci-ence 267: 381-383,
[63404] 1995.

[63405] 10621.

[63406] Schuchardt, A.; D'Agati, V.; Larsson-Blomberg, L.; Costantini,
[63407] F.; Pachnis, V.: Defects in the kidney and enteric nervous system
[63408] of mice lacking the tyrosine kinase receptor Ret. Nature 367: 380-383,
[63409] 1994.

[63410] 10622.

[63411] Shirahama, S.; Ogura, K.; Takami, H.; Ito, K.; Tohsen, T.; Miyauchi,
[63412] A.; Nakamura, Y.: Mutational analysis of the RET proto-oncogene in
[63413] 71 Japanese patients with medullary thyroid carcinoma. J. Hum. Genet. 43:
[63414] 101-106, 1998.

[63415] 10623.

[63416] Seri, M.; Yin, L.; Barone, A.; Bolino, A.; Celli, I.; Bocciardi,
[63417] R.; Pasini, B.; Ceccherini, I.; Lerone, M.; Kristoffersson, U.; Larsson,
[63418] L. T.; Casasa, J. M.; Cass, D. T.; Abramowicz, M. J.; Vanderwinden,

[63419] J.-M.; Kravcenkiene, I.; Baric, I.; Silengo, M.; Martucciello, G.;

[63420] Romeo, G.: Frequency of RET mutations in long- and short-segment

[63421] Hirschsprung disease. Hum. Mutat. 9: 243-249, 1997.

[63422] 10624.

[63423] Takahashi, M.; Buma, Y.; Hiai, H.: Isolation of ret proto-oncogene

[63424] cDNA with an amino-terminal signal sequence. Oncogene 4: 805-806,

[63425] 1989.

[63426] 10625.

[63427] Takahashi, M.; Buma, Y.; Iwamoto, T.; Inaguma, Y.; Ikeda, H.;

[63428] Hiai, H.: Cloning and expression of the ret proto-oncogene encoding

[63429] a tyrosine kinase with two potential transmembrane do-mains. Oncogene 3:

[63430] 571-578, 1988.

[63431] 10626.

[63432] Takahashi, M.; Ritz, J.; Cooper, G. M.: Activation of a novel

[63433] human transforming gene, ret, by DNA rearrangement. Cell 42: 581-588,
[63434] 1985.

[63435] 10627.

[63436] Tessitore, A.; Sinisi, A. A.; Pasquali, D.; Cardone, M.; Vi-
tale,
[63437] D.; Bellastella, A.; Colantuoni, V.: A novel case of multiple endocrine
[63438] neoplasia type 2A associated with two de novo mutations of the RET
[63439] protooncogene. J. Clin. Endocr. Metab. 84: 3522-3527, 1999.
[63440] 10628.

[63441] van Heyningen, V.: One gene--four syndromes. Nature 367: 319-320,
[63442] 1994.

[63443] 10629.

[63444] Xue, F.; Yu, H.; Maurer, L. H.; Memoli, V. A.; Nutile-
McMenemey,
[63445] N.; Schuster, M. K.; Bowden, D. W.; Mao, J.; Noll, W. W.: Germline

[63446] RET mutations in MEN 2A and FMTC and their detection by simple DNA
[63447] diagnostic tests. Hum. Molec. Genet. 3: 635-638, 1994.

[63448] 10630.

[63449] Yin, L.; Ceccherini, I.; Pasini, B.; Matera, I.; Bicocchi, M.

[63450] P.; Barone, V.; Bocciardi, R.; Kaariainen, H.; Weber, D.; De-voto,
[63451] M.; Romeo, G.: Close linkage with the RET protooncogene and boundaries
[63452] of deletion mutations in autosomal dominant Hirschsprung disease. Hum.
[63453] Molec. Genet. 2: 1803-1808, 1993.

[63454] 10631.

[63455] Feller, S. M.; Ren, R. B.; Hanafusa, H.; Baltimore, D.: SH2 and
[63456] SH3 domains as molecular adhesives: the interactions of crk and abl. Trends
[63457] Biochem. Sci. 19: 453-458, 1994.

[63458] 10632.

[63459] Benz-Lemoine, E.; Brizard, A.; Huret, J.-L.; Babin, P.; Guil-hot,

[63460] F.; Couet, D.; Tanzer, J.: Malignant histiocytosis: a specific t(2;5)(p23;q35)
[63461] translocation? Review of the literature. Blood 72: 1045-1047, 1988.
[63462] 10633.

[63463] Boultwood, J.; Rack, K.; Kelly, S.; Madden, J.; Sakaguchi, A. Y.;
[63464] Wang, L.-M.; Oscier, D. G.; Buckle, V. J.; Wainscoat, J. S.: Loss
[63465] of both CSF1R (FMS) alleles in patients with myelodysplasia and a
[63466] chromosome 5 deletion. Proc. Nat. Acad. Sci. 88: 6176-6180, 1991.
[63467] 10634.

[63468] De Qi Xu; Guilhot, S.; Galibert, F.: Restriction fragment length
[63469] polymorphism of the human c-fms gene. Proc. Nat. Acad. Sci. 82:
[63470] 2862-2865, 1985.

[63471] 10635.

[63472] Eccles, M. R.: Genes encoding the platelet-derived growth factor

[63473] (PDGF) receptor and colony-stimulating factor 1 (CSF-1) receptor are
[63474] physically associated in mice as in humans. Gene 108: 285-288, 1991.
[63475] 10636.

[63476] Gisselbrecht, S.; Fichelson, S.; Sola, B.; Bordereaux, D.; Hampe,
[63477] A.; Andre, C.; Galibert, F.; Tambourin, P. E.: Frequent c-fms activation
[63478] by proviral insertion in mouse myeloblastic leukaemias. Nature 329:
[63479] 259-261, 1987.

[63480] 10637.

[63481] Groffen, J.; Heisterkamp, N.; Spurr, N. K.; Dana, S. L.; Was-muth,
[63482] J. J.; Stephenson, J. R.: Regional assignment of the human c-fms
[63483] oncogene to band q34 of chromosome 5. (Abstract) Cyto-genet. Cell
[63484] Genet. 37: 484 only, 1984.

[63485] 10638.

[63486] Hampe, A.; Gobet, M.; Sherr, C. J.; Galibert, F.: Nucleotide sequence
[63487] of the feline retroviral oncogene v-fms shows unexpected homology
[63488] with oncogenes encoding tyrosine-specific protein kinases. Proc.
[63489] Nat. Acad. Sci. 81: 85-89, 1984.

[63490] 10639.

[63491] Hampe, A.; Shamoon, B.-M.; Gobet, M.; Sherr, C. J.; Galibert, F.
[63492] : Nucleotide sequence and structural organization of the human FMS
[63493] proto-oncogene. Oncogene Res. 4: 9-17, 1989.

[63494] 10640.

[63495] How, G.-F.; Venkatesh, B.; Brenner, S.: Conserved linkage between
[63496] the puffer fish (Fugu rubripes) and human genes for platelet-derived
[63497] growth factor receptor and macrophage colony-stimulating factor receptor. Genome
[63498] Res. 6: 1185-1191, 1996.

[63499] 10641.

[63500] Graham, M.; Adams, J. M.: Chromosome 8 breakpoint far 3-prime

[63501] of the c-myc oncogene in a Burkitt's lymphoma 2;8 variant translocation

[63502] is equivalent to the murine pvt-1 locus. EMBO J. 5: 2845-2851,

[63503] 1986.

[63504] 10642.

[63505] Berger, A.; Rosenthal, D.; Spiegel, S.: Sphingosylphosphocholine,

[63506] a signaling molecule which accumulates in Niemann-Pick disease type

[63507] A, stimulates DNA-binding activity of the transcription activator

[63508] protein AP-1. Proc. Nat. Acad. Sci. 92: 5885-5889, 1995.

[63509] 10643.

[63510] Bohmann, D.; Bos, T. J.; Admon, A.; Nishimura, T.; Vogt, P. K.;

[63511] Tjian, R.: Human proto-oncogene c-jun encodes a DNA binding protein

[63512] with structural and functional properties of transcription factor
[63513] AP-1. Science 238: 1386-1392, 1987.

[63514] 10644.

[63515] Bos, T. J.; Bohmann, D.; Tsuchie, H.; Tjian, R.; Vogt, P. K.:

[63516] v-jun encodes a nuclear protein with enhancer binding properties of
[63517] AP-1. Cell 52: 705-712, 1988.

[63518] 10645.

[63519] Haluska, F. G.; Huebner, K.; Isobe, M.; Nishimura, T.; Croce, C.
[63520] M.; Vogt, P. K.: Localization of the human JUN protooncogene to chromosome
[63521] region 1p31-32. Proc. Nat. Acad. Sci. 85: 2215-2218, 1988.
[63522] 10646.

[63523] Hattori, K.; Angel, P.; Le Beau, M. M.; Karin, M.: Structure and
[63524] chromosomal localization of the functional intronless human JUN protooncogene. Proc.
[63525] Nat. Acad. Sci. 85: 9148-9152, 1988.

[63526] 10647.

[63527] Lamph, W. W.; Wamsley, P.; Sassone-Corsi, P.; Verma, I. M.: Induction

[63528] of proto-oncogene JUN/AP-1 by serum and TPA. Nature 334: 629-631,

[63529] 1988.

[63530] 10648.

[63531] Marx, J. L.: 'Jun' is bustin' out all over. (Research News). Science 242:

[63532] 1377-1378, 1988.

[63533] 10649.

[63534] Mattei, M. G.; Simon-Chazottes, D.; Hirai, S.-I.; Ryseck, R.-P.;

[63535] Galcheva-Gargova, Z.; Guenet, J. L.; Mattei, J. F.; Bravo, R.; Yaniv,

[63536] M.: Chromosomal localization of the three members of the jun proto-oncogene

[63537] family in mouse and man. Oncogene 5: 151-156, 1990.

[63538] 10650.

[63539] Shaulian, E.; Karin, M.: AP-1 as a regulator of cell life and

[63540] death. Nature Cell Biol. 4: E131-E136, 2002.

[63541] 10651.

[63542] Shaulian, E.; Schreiber, M.; Piu, F.; Beeche, M.; Wagner, E. F.;

[63543] Karin, M.: The mammalian UV response: c-Jun induction is required

[63544] for exit from p53-imposed growth arrest. Cell 103: 897-907, 2000.

[63545] 10652.

[63546] Mattei, M. G.; Simon-Chazottes, D.; Hirai, S.; Ryseck, R. P.; Galcheva-Gargova,

[63547] Z.; Guenet, J. L.; Mattei, J. F.; Bravo, R.; Yaniv, M.: Chromo-somal

[63548] localization of the three members of the jun proto-oncogene family

[63549] in mouse and man. Oncogene 5: 151-156, 1990.

[63550] 10653.

[63551] Passegue, E.; Jochum, W.; Behrens, A.; Ricci, R.; Wagner, E. F.

[63552] : JunB can substitute for Jun in mouse development and cell proliferation. Nature

[63553] Genet. 30: 158-166, 2002.

[63554] 10654.

[63555] Passegue, E.; Jochum, W.; Schorpp-Kistner, M.; Mohle-Steinlein,

[63556] U.; Wagner, E. F.: Chronic myeloid leukemia with increased granulocyte

[63557] progenitors in mice lacking JunB expression in the myeloid lineage. Cell 104:

[63558] 21-32, 2001.

[63559] 10655.

[63560] Phinney, D. G.; Tseng, S. W.; Ryder, K.: Complex genetic organization

[63561] of junB: multiple blocks of flanking evolutionarily conserved sequence

[63562] at the murine and human junB loci. Genomics 28: 228-234, 1995.

[63563] 10656.

[63564] Sullivan, M.; Olsen, A. S.; Houslay, M. D.: Genomic organisation

[63565] of the human cyclic AMP-specific phosphodiesterase PDE4C gene and

[63566] its chromosomal localisation to 19p13.1, between RAB3A and JUND. Cell.

[63567] Signal. 11: 735-742, 1999.

[63568] 10657.

[63569] Weitzman, J. B.; Fiette, L.; Matsuo, K.; Yaniv, M.: JunD pro-tects

[63570] cells from p53-dependent senescence and apoptosis. Molec. Cell 6:

[63571] 1109-1119, 2000.

[63572] 10658.

[63573] Falush, D.; Almqvist, E. W.; Brinkmann, R. R.; Iwasa, Y.; Hayden,

[63574] M. R.: Measurement of mutational flow implies both a high new-mutation

[63575] rate for Huntington disease and substantial underascer-tainment of

[63576] late-onset cases. Am. J. Hum. Genet. 68: 373-385, 2001.

[63577] 10659.

[63578] Blin-Wakkach, C.; Lezot, F.; Ghoul-Mazgar, S.; Hotton, D.; Monteiro,

[63579] S.; Teillaud, C.; Pibouin, L.; Orestes-Cardoso, S.; Papager-

akis, P.;

[63580] Macdougall, M.; Robert, B.; Berdal, A.: Endogenous Msx1 antisense

[63581] transcript: in vivo and in vitro evidences, structure, and potential

[63582] involvement in skeleton development in mammals. Proc. Nat. Acad.

[63583] Sci. 98: 7336-7341, 2001.

[63584] 10660.

[63585] Goldfarb, L. G.; Vasconcelos, O.; Platonov, F. A.; Lunkes, A.;

[63586] Kipnis, V.; Kononova, S.; Chabrashvili, T.; Vladimirtsev, V. A.; Alexeev,

[63587] V. P.; Gajdusek, D. C.: Unstable triplet repeat and phenotypic variability

[63588] of spinocerebellar ataxia type 1. Ann. Neurol. 39: 500-506, 1996.

[63589] 10661.

[63590] Pellegris, G.; Illeni, M. T.; Rovini, D.; Vaglini, M.; Cascinelli,

[63591] N.; Ghidoni, A.: HLA complex and familial malignant melanoma. Int.

[63592] J. Cancer 29: 621-623, 1982.

[63593] 10662.

[63594] Steiglitz, B. M.; Greenspan, D. S.: Assignment of the mouse Pcolce2

[63595] gene, which encodes procollagen C-proteinase enhancer protein 2, to

[63596] chromosome 9 and localization of PCOLCE2 to human chromosome 3q23. Cytogenet.

[63597] Cell Genet. 95: 244-245, 2001.

[63598] 10663.

[63599] Xu, H.; Acott, T. S.; Wirtz, M. K.: Identification and expression

[63600] of a novel type I procollagen C-proteinase enhancer protein gene from

[63601] the glaucoma candidate region on 3q21-q24. Genomics 66: 264-273,

[63602] 2000.

[63603] 10664.

[63604] Peier, A. M.; Reeve, A. J.; Andersson, D. A.; Moqrich, A.; Earley,

[63605] T. J.; Hergarden, A. C.; Story, G. M.; Colley, S.; Hogenesch, J. B.;

[63606] McIntyre, P.; Bevan, S.; Patapoutian, A.: A heat-sensitive TRP channel

[63607] expressed in keratinocytes. Science 296: 2046-2049, 2002.

[63608] 10665.

[63609] Smith, G. D.; Gunthorpe, M. J.; Kelsell, R. E.; Hayes, P. D.; Reilly,

[63610] P.; Facer, P.; Wright, J. E.; Jerman, J. C.; Walhin, J.-P.; Ooi, L.;

[63611] Egerton, J.; Charles, K. J.; Smart, D.; Randall, A. D.; Anand, P.;

[63612] Davis, J. B.: TRPV3 is a temperature-sensitive vanilloid receptor-like

[63613] protein. Nature 418: 186-190, 2002.

[63614] 10666.

[63615] Xu, H.; Ramsey, I. S.; Kotecha, S. A.; Moran, M. M.; Chong, J.

[63616] A.; Lawson, D.; Ge, P.; Lilly, J.; Silos-Santiago, I.; Xie, Y.; DiStefano,

[63617] P. S.; Curtis, R.; Clapham, D. E.: TRPV3 is a calcium-permeable temperature-sensitive

[63618] cation channel. Nature 418: 181-186, 2002.

[63619] 10667.

[63620] Ching, Y.-P.; Zhou, H.-J.; Yuan, J.-G.; Qiang, B.-Q.; Kung, H.;

[63621] Jin, D.-Y.: Identification and characterization of FTSJ2, a novel

[63622] human nucleolar protein homologous to bacterial ribosomal RNA methyltransferase. Genomics 79:

[63623] 2-6, 2002.

[63624] 10668.

[63625] Brand, S. H.; Castle, J. D.: SCAMP-37, a new marker within the

[63626] general cell surface recycling system. EMBO J. 12: 3753-3761, 1993.

[63627] 10669.

[63628] Singleton, D. R.; Wu, T. T.; Castle, J. D.: Three mammalian SCAMPs

[63629] (secretory carrier membrane proteins) are highly related products

[63630] of distinct genes having similar subcellular distributions. J. Cell

[63631] Sci. 110: 2099-2107, 1997.

[63632] 10670.

[63633] Yamashita, A.; Ohnishi, T.; Kashima, I.; Taya, Y.; Ohno, S.: Human

[63634] SMG-1, a novel phosphatidylinositol 3-kinase-related protein kinase,

[63635] associates with components of the mRNA surveillance complex and is

[63636] involved in the regulation of nonsense-mediated mRNA decay. Genes

[63637] Dev. 15: 2215-2228, 2001.

[63638] 10671.

[63639] Bossolasco, M.; Lebel, M.; Lemieux, N.; Mes-Masson, A.-M.: The

[63640] human TDE gene homologue: localization to 20q13.1-13.3 and variable

[63641] expression in human tumor cell lines and tissue. Molec. Carcinogen. 26:

[63642] 189-200, 1999.

[63643] 10672.

[63644] Modarressi, M. H.; Cameron, J.; Taylor, K. E.; Wolfe, J.: Identification

[63645] and characterisation of a novel gene, TSGA10, expressed in testis. Gene 262:

[63646] 249-255, 2001.

[63647] 10673.

[63648] Bowlus, C. L.; Ahn, J.; Chu, T.; Gruen, J. R.: Cloning of a novel

[63649] MHC-encoded serine peptidase highly expressed by cortical epithelial

[63650] cells of the thymus. Cell. Immun. 196: 80-86, 1999.

[63651] 10674.

[63652] Cheunsuk, S.; Sparks, R.; Noveroske, J. K.; Hsu, T.; Justice, M.

[63653] J.; Gershwin, M. E.; Gruen, J. R.; Bowlus, C. L.: Expression, genomic

[63654] structure and mapping of the thymus specific protease prss16: a candidate

[63655] gene for insulin dependent diabetes mellitus susceptibility. J. Autoimmun. 18:

[63656] 311-316, 2002.

[63657] 10675.

[63658] Kamberov, E.; Makarova, O.; Roh, M.; Liu, A.; Karnak, D.;

Straight,
[63659] S.; Margolis, B.: Molecular cloning and characterization of Pals,
[63660] proteins associated with mLin-7. J. Biol. Chem. 275: 11425-11431,
[63661] 2000.

[63662] 10676.

[63663] Tseng, T.-C.; Marfatia, S. M.; Bryant, P. J.; Pack, S.; Zhuang,
[63664] A.; O'Brien, J. E.; Lin, L.; Hanada, T.; Chishti, A. H.: VAM-1: a
[63665] new member of the MAGUK family binds to human Veli-1 through a conserved
[63666] domain. Biochim. Biophys. Acta 1518: 249-259, 2001.

[63667] 10677.

[63668] Wei, X.; Malicki, J. M.: nagie oko, encoding a MAGUK-family protein,
[63669] is essential for cellular patterning of the retina. Nature Genet. 31:
[63670] 150-157, 2002. Note: Erratum: Nature Genet. 31: 439 only, 2002.
[63671] 10678.

[63672] Esser, V.; Russell, D. W.: Transport-deficient mutations in the
[63673] low density lipoprotein receptor: alterations in the cysteine-rich
[63674] and cysteine-poor regions of the protein block intracellular transport. J.
[63675] Biol. Chem. 263: 13276-13281, 1988.

[63676] 10679.

[63677] Fumeron, F.; Grandchamp, B.; Fricker, J.; Krempf, M.; Wolf, L.-M.;
[63678] Khayat, M.-C.; Boiffard, O.; Apfelbaum, M.: Presence of the French
[63679] Canadian deletion in a French patient with familial hypercholesterolemia.
[63680] (Letter) New Eng. J. Med. 326: 69 only, 1992.

[63681] 10680.

[63682] Gilbert, W.: Genes-in-pieces revisited. Science 228: 823-824,
[63683] 1985.

[63684] 10681.

[63685] Goldfarb, L. G.; Petersen, R. B.; Tabaton, M.; Brown, P.;

LeBlanc,
[63686] A. C.; Montagna, P.; Cortelli, P.; Julien, J.; Vital, C.; Pendel-
bury,
[63687] W. W.; Haltia, M.; Wills, P. R.; and 9 others: Fatal familial
insomnia
[63688] and familial Creutzfeldt-Jakob disease: disease phenotype
determined
[63689] by a DNA polymorphism. Science 258: 806-808, 1992.

[63690] 10682.

[63691] Graadt van Roggen, F.; van der Westhuyzen, D. R.; Marais,
A. D.;
[63692] Gevers, W.; Coetzee, G. A.: Low density lipoprotein recep-
tor founder
[63693] mutations in Afrikaner familial hypercholesterolaemic pa-
tients: a
[63694] comparison of two geographical areas. Hum. Genet. 88:
204-208, 1991.
[63695] 10683.

[63696] Gudnason, V.; King-Underwood, L.; Seed, M.; Sun, X.-M.;
Soutar,
[63697] A. K.; Humphries, S. E.: Identification of recurrent and
novel mutations

[63698] in exon 4 of the LDL receptor gene in patients with familial hypercholesterolemia

[63699] in the United Kingdom. Arteriosclerosis Thromb. 13: 56-63, 1993.

[63700] 10684.

[63701] Gudnason, V.; Sigurdsson, G.; Nissen, H.; Humphries, S. E.: Common

[63702] founder mutation in the LDL receptor gene causing familial hypercholesterolemia

[63703] in the Icelandic population. Hum. Mutat. 10: 36-44, 1997.

[63704] 10685.

[63705] Henderson, H. E.; Berger, G. M. B.; Marais, A. D.: A new LDL

[63706] receptor gene deletion mutation in the South African population. Hum.

[63707] Genet. 80: 371-374, 1988.

[63708] 10686.

[63709] Hobbs, H. H.: Personal Communication. Dallas, Tex. 12/1/1990.

[63710] 10687.

[63711] Hobbs, H. H.; Brown, M. S.; Russell, D. W.; Davignon, J.;

Goldstein,
[63712] J. L.: Deletion in the gene for the low-density-lipoprotein receptor
[63713] in a majority of French Canadians with familial hyperc-holesterolemia. New
[63714] Eng. J. Med. 317: 734-737, 1987.

[63715] 10688.

[63716] Hobbs, H. H.; Lehrman, M. A.; Yamamoto, T.; Russell, D. W.: Polymorphism
[63717] and evolution of Alu sequences in the human low density lipoprotein
[63718] receptor gene. Proc. Nat. Acad. Sci. 82: 7651-7655, 1985.

[63719] 10689.

[63720] Hobbs, H. H.; Leitersdorf, E.; Goldstein, J. L.; Brown, M. S.;

[63721] Russell, D. W.: Multiple CRM- mutations in familial hyper-cholesterolemia:
[63722] evidence for 13 alleles, including four deletions. J. Clin. Invest. 81:
[63723] 909-917, 1988.

[63724] 10690.

[63725] Horsthemke, B.; Dunning, A.; Humphries, S.: Identification

of
[63726] deletions in the human low density lipoprotein receptor gene. J.
[63727] Med. Genet. 24: 144-147, 1987.

[63728] 10691.

[63729] Horsthemke, B.; Kessling, A. M.; Seed, M.; Wynn, V.; Williamson,
[63730] R.; Humphries, S. E.: Identification of a deletion in the low density
[63731] lipoprotein (LDL) receptor gene in a patient with familial hypercholesterolaemia. Hum.
[63732] Genet. 71: 75-78, 1985.

[63733] 10692.

[63734] Jensen, H. K.; Jensen, T. G.; Faergeman, O.; Jensen, L. G.; Andresen,
[63735] B. S.; Corydon, M. J.; Andreasen, P. H.; Hansen, P. S.; Heath, F.;
[63736] Bolund, L.; Gregersen, N.: Two mutations in the same low-density
[63737] lipoprotein receptor allele act in synergy to reduce recep-tor function
[63738] in heterozygous familial hypercholesterolemia. Hum. Mu-

tat. 9: 437-444,

[63739] 1997.

[63740] 10693.

[63741] Jensen, J. M.; Kruse, T. A.; Brorholt-Petersen, J. U.; Chris-
tiansen,

[63742] T. M.; Jensen, H. K.; Kolvraa, S.; Faergeman, O.: Linking
genotype

[63743] to aorto-coronary atherosclerosis: a model using familial
hypercholesterolemia

[63744] and aorto-coronary calcification. Ann. Hum. Genet. 63:
511-520,

[63745] 1999.

[63746] 10694.

[63747] Kajinami, K.; Fujita, H.; Koizumi, J.; Mabuchi, H.; Takeda,
R.;

[63748] Ohta, M.: Genetically determined mild type of familial hy-
percholesterolemia

[63749] including normocholesterolemic patients: FH-Tonami-2.
Circulation 80

[63750] (suppl. 2): 278 only, 1989.

[63751] 10695.

[63752] Kajinami, K.; Mabuchi, H.; Itoh, H.; Michishita, I.; Takeda, M.;

[63753] Wakasugi, T.; Koizumi, J.; Takeda, R.: New variant of low density

[63754] lipoprotein receptor gene FH-Tonami. Arteriosclerosis 8: 187-192,

[63755] 1988.

[63756] 10696.

[63757] Suvorova, E. S.; Kurten, R. C.; Lupashin, V. V.: Identification

[63758] of a human orthologue of Sec34p as a component of the cis-Golgi vesicle

[63759] tethering machinery. J. Biol. Chem. 276: 22810-22818, 2001.

[63760] 10697.

[63761] Whyte, J. R. C.; Munro, S.: The Sec34/35 Golgi transport complex

[63762] is related to the exocyst, defining a family of complexes involved

[63763] in multiple steps of membrane traffic. Dev. Cell 1: 527-537, 2001.

[63764] 10698.

[63765] Haataja, L.; Groffen, J.; Heisterkamp, N.: Identification of a

[63766] novel Rac3-interacting protein C1D. Int. J. Molec. Med. 1: 665-670,

[63767] 1998.

[63768] 10699.

[63769] Nehls, P.; Keck, T.; Greferath, R.; Spiess, E.; Glaser, T.; Rothbarth,

[63770] K.; Stammer, H.; Werner, D.: cDNA cloning, recombinant expression

[63771] and characterization of polypeptides with exceptional DNA affinity. Nucleic

[63772] Acids Res. 26: 1160-1166, 1998.

[63773] 10700.

[63774] Rothbarth, K.; Hunziker, A.; Stammer, H.; Werner, D.: Pro-moter

[63775] of the gene encoding the 16 kDa DNA-binding and apop-tosis-inducing

[63776] C1D protein. Biochim. Biophys. Acta 1518: 271-275, 2001.

[63777] 10701.

[63778] Zamir, I.; Dawson, J.; Lavinsky, R. M.; Glass, C. K.; Rosen-

feld,

[63779] M. G.; Lazar, M. A.: Cloning and characterization of a corepressor

[63780] and potential component of the nuclear hormone receptor repression

[63781] complex. Proc. Nat. Acad. Sci. 94: 14400-14405, 1997.

[63782] 10702.

[63783] Nguyen, L. H.; Erzberger, J. P.; Root, J.; Wilson, D. M., III:

[63784] The human homolog of Escherichia coli Orn degrades small single-stranded

[63785] RNA and DNA oligomers. J. Biol. Chem. 275: 25900-25906, 2000.

[63786] 10703.

[63787] Bayani, J.; et al.; et al: Molecular cytogenetic analysis of medulloblastomas

[63788] and supratentorial primitive neuroectodermal tumors by using conventional

[63789] banding, comparative genomic hybridization, and spectral karyotyping. J.

[63790] Neurosurg. 93: 437-448, 2000.

[63791] 10704.

[63792] Grimm, T.; Teglund, S.; Tackels, D.; Sangiorgi, E.; Gurrieri, F.;

[63793] Schwartz, C.; Toftgard, R.: Genomic organization and embryonic expression

[63794] of Suppressor of Fused, a candidate gene for the split-hand/split-foot

[63795] malformation type 3. FEBS Lett. 505: 13-17, 2001.

[63796] 10705.

[63797] Stone, D. M.; Murone, M.; Luoh, S.-M.; Ye, W.; Armanini, M. P.;

[63798] Gurney, A.; Phillips, H.; Brush, J.; Goddard, A.; de Sauvage, F. J.;

[63799] Rosenthal, A.: Characterization of the human suppressor of fused,

[63800] a negative regulator of the zinc-finger transcription factor Gli. J.

[63801] Cell Sci. 112: 4437-4448, 1999.

[63802] 10706.

[63803] Nagle, D. L.; McGrail, S. H.; Vitale, J.; Woolf, E. A.; Dussault,

[63804] B. J., Jr.; DiRocco, L.; Holmgren, L.; Montagno, J.; Bork, P.; Huszar,

[63805] D.; Fairchild-Huntress, V.; Ge, P.; Keilty, J.; Ebelling, C.; Baldini,
[63806] L.; Gilchrist, J.; Burr, P.; Carlson, G. A.; Moore, K. J.: The mahogany
[63807] protein is a receptor involved in suppression of obesity. Nature 398:
[63808] 148-151, 1999.

[63809] 10707.

[63810] Zhang, Y.; Ng, H.-H.; Erdjument-Bromage, H.; Tempst, P.; Bird,
[63811] A.; Reinberg, D.: Analysis of the NuRD subunits reveals a histone
[63812] deacetylase core complex and a connection with DNA methylation. Genes
[63813] Dev. 13: 1924-1935, 1999.

[63814] 10708.

[63815] Takafuta, T.; Wu, G.; Murphy, G. F.; Shapiro, S. S.: Human beta-filamin
[63816] is a new protein that interacts with the cytoplasmic tail of glycoprotein
[63817] Ib-alpha. J. Biol. Chem. 273: 17531-17538, 1998.

[63818] 10709.

[63819] Van Gennip, A. H.; Mandel, H.; Stroomer, L. E.; van Cruchten, A.

[63820] G.: Effect of allopurinol on the xanthinuria in a patient with molybdenum

[63821] cofactor deficiency. Adv. Exp. Med. Biol. 370: 375-378, 1994.

[63822] 10710.

[63823] Margolin, J. F.; Friedman, J. R.; Meyer, W. K.; Vissing, H.; Thiesen,

[63824] H.-J.; Rauscher, F. J., III: Kruppel-associated boxes are po-tent

[63825] transcriptional repression domains. Proc. Nat. Acad. Sci. 91: 4509-4513,

[63826] 1994.

[63827] 10711.

[63828] Pineda, M.; Fernandez, E.; Torrents, D.; Estevez, R.; Lopez, C.;

[63829] Camps, M.; Lloberas, J.; Zorzano, A.; Palacin, M.: Identifi-cation

[63830] of a membrane protein, LAT-2, that co-expresses with 4F2 heavy chain,

[63831] an L-type amino acid transport activity with broad specificity for
[63832] small and large zwitterionic amino acids. J. Biol. Chem. 274: 19738-19744,
[63833] 1999.

[63834] 10712.

[63835] Alexander, W. S.; Rakar, S.; Robb, L.; Farley, A.; Willson, T.

[63836] A.; Zhang, J.-G.; Hartley, L.; Kikuchi, Y.; Kojima, T.; Nomura, H.;
[63837] Hasegawa, M.; Maeda, M.; Fabri, L.; Jachno, K.; Nash, A.; Metcalf,
[63838] D.; Nicola, N. A.; Hilton, D. J.: Suckling defect in mice lacking
[63839] the soluble haemopoietin receptor NR6. Curr. Biol. 9: 605-608, 1999.
[63840] 10713.

[63841] Elson, G. C. A.; Graber, P.; Losberger, C.; Herren, S.; Gretener,
[63842] D.; Menoud, L. N.; Wells, T. N. C.; Kosco-Vilbois, M. H.; Gauchat,
[63843] J.-F.: Cytokine-like factor-1, a novel soluble protein, shares homology

[63844] with members of the cytokine type I receptor family. J. Immun. 161:

[63845] 1371-1379, 1998.

[63846] 10714.

[63847] Grau, Y.; Carteret, C.; Simpson, P.: Mutations and chromosomal

[63848] rearrangements affecting the expression of snail, a gene involved

[63849] in embryonic patterning in Drosophila melanogaster. Genetics 108:

[63850] 347-360, 1984.

[63851] 10715.

[63852] Okubo, T.; Truong, T. K.; Yu, B.; Itoh, T.; Zhao, J.; Grube, B.;

[63853] Zhou, D.; Chen, S.: Down-regulation of promoter I.3 activity of the

[63854] human aromatase gene in breast tissue by zinc-finger protein, snail

[63855] (SnaH). Cancer Res. 15: 1338-1346, 2001.

[63856] 10716.

[63857] Paznekas, W. A.; Okajima, K.; Schertzer, M.; Wood, S.;

Jabs, E.

[63858] W.: Genomic organization, expression, and chromosome location of

[63859] the human SNAIL gene (SNAI1) and a related processed pseudogene (SNAI1P). Genomics 62:

[63860] 42-49, 1999.

[63861] 10717.

[63862] Sefton, M.; Sanchez, S.; Nieto, M. A.: Conserved and diver-gent

[63863] roles for members of the Snail family of transcription fac-tors in

[63864] the chick and mouse embryo. Development 124: 125:-3111-3121, 1998.

[63865] 10718.

[63866] Twigg, S. R. F.; Wilkie, A. O. M.: Characterisation of the human

[63867] snail (SNAI1) gene and exclusion as a major disease gene in craniosynostosis. Hum.

[63868] Genet. 105: 320-326, 1999.

[63869] 10719.

[63870] Shalata, A.; Mandel, H.; Reiss, J.; Szargel, R.; Cohen-Ak-

enine,
[63871] A.; Dorche, C.; Zabot, M.-T.; Van Gennip, A.; Abeling, N.; Berant,
[63872] M.; Cohen, N.: Localization of a gene for molybdenum co-factor deficiency,
[63873] on the short arm of chromosome 6, by homozygosity mapping. Am. J.
[63874] Hum. Genet. 63: 148-154, 1998.

[63875] 10720.

[63876] Hinderlich, S.; Stasche, R.; Zeitler, R.; Reutter, W.: A bi-functional
[63877] enzyme catalyzes the first two steps in N-acetylneuraminic acid biosynthesis
[63878] of rat liver: purification and characterization of UDP-N-acetylglucosamine
[63879] 2-epimerase/N-acetylmannosamine kinase. J. Biol. Chem. 272: 24313-24318,
[63880] 1997.

[63881] 10721.

[63882] Huizing, M.; Anikster, Y.: Personal Communication. Bethesda,
[63883] Md. 1/10/2000.

[63884] 10722.

[63885] Kayashima, T.; Matsuo, H.; Satoh, A.; Ohta, T.; Yoshiura, K.; Matsumoto,
[63886] N.; Nakane, Y.; Niikawa, N.; Kishino, T.: Nonaka myopathy is caused
[63887] by mutations in the UDP-N-acetylglucosamine-2-epimerase/N-acetylmannosamine
[63888] kinase gene (GNE). J. Hum. Genet. 47: 77-79, 2002.

[63889] 10723.

[63890] Keppler, O. T.; Hinderlich, S.; Langner, J.; Schwartz-Albiez, R.;
[63891] Reutter, W.; Pawlita, M.: UDP-GlcNAc 2-epimerase: a regulator of
[63892] cell surface sialylation. Science 284: 1372-1376, 1999.

[63893] 10724.

[63894] Nonaka, I.; Sunohara, N.; Ishiura, S.; Satoyoshi, E.: Familial
[63895] distal myopathy with rimmed vacuole and lamellar (myeloid) body formation. J.
[63896] Neurol. Sci. 51: 141-155, 1981.

[63897] 10725.

[63898] Schwarzkopf, M.; Knobeloch, K.-P.; Rohde, E.; Hinderlich, S.;

[63899] Wiechens, N.; Lucka, L.; Horak, I.; Reutter, W.; Horstkorte, R.:

[63900] Sialylation is essential for early development in mice. Proc. Nat.

[63901] Acad. Sci. 99: 5267-5270, 2002.

[63902] 10726.

[63903] Stasche, R.; Hinderlich, S.; Weise, C.; Effertz, K.; Lucka, L.;

[63904] Moormann, P.; Reutter, W.: A bifunctional enzyme cat-alyzes the first

[63905] two steps in N-acetylneuraminic acid biosynthesis of rat liver: molecular

[63906] cloning and functional expression of UDP-N-acetyl-glucosamine 2-epimerase/N-acetylmannosamine

[63907] kinase. J. Biol. Chem. 272: 24319-24324, 1997.

[63908] 10727.

[63909] Carter, M. G.; Johns, M. A.; Zeng, X.; Zhou, L.; Zink, M. C.; Mankowski,

[63910] J. L.; Donovan, D. M.; Baylin, S. B.: Mice deficient in the

candidate

[63911] tumor suppressor gene Hic1 exhibit developmental de-
fects of structures

[63912] affected in the Miller-Dieker syndrome. Hum. Molec.
Genet. 9: 413-419,

[63913] 2000.

[63914] 10728.

[63915] Chong, S. S.; Tanigami, A.; Roschke, A. V.; Ledbetter, D.
H.:

[63916] 14-3-3epsilon has no homology to LIS1 and lies telomeric
to it on

[63917] chromosome 17p13.3 outside the Miller-Dieker syndrome
chromosome region. Genome

[63918] Res. 6: 735-741, 1996.

[63919] 10729.

[63920] Grimm, C.; Sporle, R.; Schmid, T. E.; Adler, I.-D.; Adamski,
J.;

[63921] Schughart, K.; Graw, J.: Isolation and embryonic expres-
sion of the

[63922] novel mouse gene Hic1, the homologue of HIC1, a candi-
date gene for

[63923] the Miller-Dieker syndrome. Hum. Molec. Genet. 8:

697-710, 1999.

[63924] 10730.

[63925] Makos Wales, M.; Biel, M. A.; El Deiry, W.; Nelkin, B. D.; Issa,

[63926] J.-P.; Cavenee, W. K.; Kuerbitz, S. J.; Baylin, S. B.: p53 activates

[63927] expression of HIC-1, a new candidate tumour suppressor gene on 17p13.3. Nature

[63928] Med. 1: 570-577, 1995.

[63929] 10731.

[63930] Smith, D. J.; Salmi, M.; Bono, P.; Hellman, J.; Leu, T.; Jalkanen,

[63931] S.: Cloning of vascular adhesion protein 1 reveals a novel multifunctional

[63932] adhesion molecule. J. Exp. Med. 188: 17-27, 1998.

[63933] 10732.

[63934] Zhang, X.; McIntire, W. S.: Cloning and sequencing of a copper-containing,

[63935] topa quinone-containing monoamine oxidase from human placenta. Gene 179:

[63936] 279-286, 1996.

[63937] 10733.

[63938] Lomako, J.; Mazuruk, K.; Lomako, W. M.; Alonso, M. D.; Whelan,

[63939] W. J.; Rodriguez, I. R.: The human intron-containing gene for glycogenin

[63940] maps to chromosome 3, band q24. Genomics 33: 519-522, 1996.

[63941] 10734.

[63942] Tanaka, T.; Tsujimura, T.; Takeda, K.; Sugihara, A.; Maekawa, A.;

[63943] Terada, N.; Yoshida, N.; Akira, S.: Targeted disruption of ATF4 discloses

[63944] its essential role in the formation of eye lens fibres. Genes Cells 3:

[63945] 801-810, 1998.

[63946] 10735.

[63947] Maas, S.; Gerber, A. P.; Rich, A.: Identification and charac-terization

[63948] of a human tRNA-specific adenosine deaminase related to the ADAR family

[63949] of pre-mRNA editing enzymes. Proc. Nat. Acad. Sci. 96: 8895-8900,

[63950] 1999.

[63951] 10736.

[63952] Viskupic, E.; Cao, Y.; Zhang, W.; Cheng, C.; DePaoli-Roach, A.

[63953] A.; Roach, P. J.: Rabbit skeletal muscle glycogenin. Molecular cloning

[63954] and production of fully functional protein in Escherichia coli. J.

[63955] Biol. Chem. 267: 25759-25763, 1992.

[63956] 10737.

[63957] Hosokawa, Y.; Matsumoto, A.; Oka, J.; Itakura, H.; Yamaguchi, K.

[63958] : Isolation and characterization of a cDNA for rat liver cysteine

[63959] dioxygenase. Biochem. Biophys. Res. Commun. 168: 473-478, 1990.

[63960] 10738.

[63961] Jeremiah, S.; McCann, K. P.; Williams, A. C.; Ramsden, D. B.; Pilz,

[63962] A. J.; Fox, M. F.; Povey, S.: Chromosomal localisation of genes coding

[63963] for human and mouse liver cytosolic cysteine dioxygenase. Ann. Hum.

[63964] Genet. 60: 29-33, 1996.

[63965] 10739.

[63966] McCann, K. P.; Akbari, M. T.; Williams, A. C.; Ramsden, D. B.:

[63967] Human cysteine dioxygenase type I: primary structure derived from

[63968] base sequencing of cDNA. Biochim. Biophys. Acta 1209: 107-110, 1994.

[63969] 10740.

[63970] Ramsden, D. B.; Kapadi, A.; Fitch, N. J. S.; Farmer, M. J.; Bennett,

[63971] P.; Williams, A. C.: Human cysteine dioxygenase type I (CDO-I; EC

[63972] 1.13.11.20): 5-prime flanking region and intron-exon structure of

[63973] the gene. Molec. Path. 50: 269-271, 1997.

[63974] 10741.

[63975] Nelen, M. R.; van Staveren, W. C. G.; Peeters, E. A. J.; Ben Hassel,

[63976] M.; Gorlin, R. J.; Hamm, H.; Lindboe, C. F.; Fryns, J.-P.; Sijmons,
[63977] R. H.; Woods, D. G.; Mariman, E. C. M.; Padberg, G. W.; Kremer, H.
[63978] : Germline mutations in the PTEN/MMAC1 gene in patients with Cowden
[63979] disease. Hum. Molec. Genet. 6: 1383-1387, 1997.

[63980] 10742.

[63981] Fox, M. F.; Lahbib, F.; Pratt, W.; Attwood, J.; Gum, J.; Kim, Y.;
[63982] Swallow, D. M.: Regional localization of the intestinal mucin gene
[63983] MUC3 to chromosome 7q22. Ann. Hum. Genet. 56: 281-287, 1992.
[63984] 10743.

[63985] Kyo, K.; Muto, T.; Nagawa, H.; Lathrop, G. M.; Nakamura, Y.: Associations
[63986] of distinct variants of the intestinal mucin gene MUC3A with ulcerative
[63987] colitis and Crohn's disease. J. Hum. Genet. 46: 5-20, 2001.
[63988] 10744.

[63989] Caro, A.; Haines, S.: The history of Huntington's chorea. Update 91-95,

[63990] 7/1975.

[63991] 10745.

[63992] Swallow, D. M.; Povey, S.; Goodfellow, P. N. G.; Andrews, P.;

[63993] Harris, H.: The liver/bone/kidney isozyme of alkaline phosphatase

[63994] (ALPL) is coded by a gene on chromosome 1. (Abstract) Cytogenet.

[63995] Cell Genet. 40: 756 only, 1985.

[63996] 10746.

[63997] Zou, Y.-R.; Kottmann, A. H.; Kuroda, M.; Taniuchi, I.; Littman,

[63998] D. R.: Function of the chemokine receptor CXCR4 in haematopoiesis

[63999] and in cerebellar development. Nature 393: 595-599, 1998.

[64000] 10747.

[64001] Bean, A. J.; Dagerlind, A.; Hokfelt, T.; Dobner P. R.: Cloning

[64002] of human neurotensin/neuromedin N genomic sequences and expression
[64003] in the ventral mesencephalon of schizophrenics and age/ sex matched
[64004] controls. Neuroscience 50: 259-268, 1992.

[64005] 10748.

[64006] Dong, Z.; Wang, X.; Zhao, Q.; Townsend, C. M., Jr.; Evers, B. M.
[64007] : DNA methylation contributes to expression of the human neurotensin/neuromedin
[64008] N gene. Am. J. Physiol. 274: G535-G543, 1998.

[64009] 10749.

[64010] Gerhard, D. S.; Dobner, P. R.; Bruns, G. A. P.: Localization of
[64011] the neurotensin gene to human chromosome 12. (Abstract) Cytogenet.
[64012] Cell Genet. 51: 1003 only, 1989.

[64013] 10750.

[64014] Mai, J. K.; Triepel, J.; Metz, J.: Neurotensin in human brain. Neuroscience 22:
[64015] 499-524, 1987.

[64016] 10751.

[64017] Marondel, I; Renault, B.; Lieman, J.; Ward, D.; Kucherlapati, R.

[64018] : Physical mapping of the human neurotensin gene (NTS) between markers

[64019] D12S1444 and D12S81 on chromosome 12q21. Genomics 38: 243-245, 1996.

[64020] 10752.

[64021] Laurent, P.; Clerc, P.; Mattei, M.-G.; Forgez, P.; Dumont, X.;

[64022] Ferrara, P.; Caput, D.; Rostene, W.: Chromosomal localization of

[64023] mouse and human neurotensin receptor genes. Mammalian Genome 5:

[64024] 303-306, 1994.

[64025] 10753.

[64026] Le, F.; Groshan, K.; Zeng, X. P.; Richelson, E.: Characterization

[64027] of the genomic structure, promoter region, and a tetranucleotide repeat

[64028] polymorphism of the human neurotensin receptor gene. J. Biol. Chem. 272:

[64029] 1315-1322, 1997.

[64030] 10754.

[64031] Vincent, J.-P.: Neurotensin receptors: binding properties, transduction

[64032] pathways, and structure. Cell. Molec. Neurobiol. 15: 501-512, 1995.

[64033] 10755.

[64034] Vita, N.; Laurent, P.; Lefort, S.; Chalon, P.; Dumont, X.; Kaghad,

[64035] M.; Gully, D.; Le Fur, G.; Ferrara, P.; Caput, D.: Cloning and expression

[64036] of a complementary DNA encoding a high affinity human neurotensin

[64037] receptor. FEBS Lett. 317: 139-142, 1993.

[64038] 10756.

[64039] Bluteau, O.; Jeannot, E.; Bioulac-Sage, P.; Marques, J. M.; Blanc,

[64040] J.-F.; Bui, H.; Beaudoin, J.-C.; Franco, D.; Balabaud, C.; Laurent-Puig,

[64041] P.; Zucman-Rossi, J.: Bi-allelic inactivation of TCF1 in hepatic

[64042] adenomas. Nature Genet. 32: 312-315, 2002.

[64043] 10757.

[64044] Foster, J. H.; Donohue, T. A.; Berman, M. M.: Familial liver-cell

[64045] adenomas and diabetes mellitus. New Eng. J. Med. 299: 239-241, 1978.

[64046] 10758.

[64047] Blinder, M. A.; Andersson, T. R.; Abildgaard, U.; Tollefsen, D.

[64048] M.: Heparin cofactor II(Oslo): mutation of arg-189-to-his decreases

[64049] the affinity for dermatan sulfate. J. Biol. Chem. 264: 5128-5133,

[64050] 1989.

[64051] 10759.

[64052] Blinder, M. A.; Marasa, J. C.; Reynolds, C. H.; Deaven, L. L.;

[64053] Tollefsen, D. M.: Heparin cofactor II: cDNA sequence, chromosome

[64054] localization, restriction fragment length polymorphism, and expression

[64055] in Escherichia coli. Biochemistry 27: 752-759, 1988.

[64056] 10760.

[64057] Herzog, R.; Lutz, S.; Blin, N.; Marasa, J. C.; Blinder, M. A.;

[64058] Tollefsen, D. M.: Complete nucleotide sequence of the gene for human

[64059] heparin cofactor II and mapping to chromosomal band 22q11. Biochemistry 30:

[64060] 1350-1357, 1991.

[64061] 10761.

[64062] Kondo, S.; Tokunaga, F.; Kario, K.; Matsuo, T.; Koide, T.: Molecular

[64063] and cellular basis for type I heparin cofactor II deficiency (heparin

[64064] cofactor II Awaji). Blood 87: 1006-1012, 1996.

[64065] 10762.

[64066] Matsuo, T.; Kario, K.; Sakamoto, S.; Yamada, T.; Miki, T.; Hirase,

[64067] T.; Kobayashi, H.: Hereditary heparin cofactor II deficiency and

[64068] coronary artery disease. Thromb. Res. 65: 495-505, 1992.

[64069] 10763.

[64070] Ragg, H.: A new member of the plasma protease inhibitor gene family. Nucleic
[64071] Acids Res. 14: 1073-1088, 1986.

[64072] 10764.

[64073] Sie, P.; Dupouy, D.; Pichon, J.; Boneu, B.: Constitutional heparin
[64074] co-factor II deficiency associated with recurrent thrombo-sis. Lancet II:
[64075] 414-416, 1985.

[64076] 10765.

[64077] Tran, T. H.; Marbet, G. A.; Duckert, F.: Association of hereditary
[64078] heparin co-factor II deficiency with thrombosis. Lancet II: 413-414,
[64079] 1985.

[64080] 10766.

[64081] Villa, P.; Aznar, J.; Vaya, A.; Espana, F.; Ferrando, F.; Mira,

[64082] Y.; Estelles, A.: Hereditary homozygous heparin cofactor II deficiency
[64083] and the risk of developing venous thrombosis. Thromb. Haemost. 82:

[64084] 1011-1014, 1999.

[64085] 10767.

[64086] Quintrell, N.; Lebo, R.; Varmus, H.; Bishop, J. M.; Pettenati,

[64087] M. J.; Le Beau, M. M.; Diaz, M. O.; Rowley, J. D.: Identifica-
tion

[64088] of a human gene (HCK) that encodes a protein-tyrosine
kinase and is

[64089] expressed in hemopoietic cells. Molec. Cell. Biol. 7:
2267-2275,

[64090] 1987.

[64091] 10768.

[64092] Ziegler, S. F.; Marth, J. D.; Lewis, D. B.; Perlmutter, R. M.:

[64093] Novel protein-tyrosine kinase gene (hck) preferentially
expressed

[64094] in cells of hematopoietic origin. Molec. Cell. Biol. 7:
2276-2285,

[64095] 1987.

[64096] 10769.

[64097] Hitzler, J. K.; Soares, H. D.; Drolet, D. W.; Inaba, T.;
O'Connel,

[64098] S.; Rosenfeld, M. G.; Morgan, J. I.; Look, A. T.: Expression patterns
[64099] of the hepatic leukemia factor gene in the nervous system of developing
[64100] and adult mice. Brain Res. 820: 1-11, 1999.

[64101] 10770.

[64102] Honda, H.; Inaba, T.; Suzuki, T.; Oda, H.; Ebihara, Y.; Tsuiji,
[64103] K.; Nakahata, T.; Ishikawa, T.; Yazaki, Y.; Hirai, H.: Expres-
sion
[64104] of E2A-HLF chimeric protein induced T-cell apoptosis, B-cell maturation
[64105] arrest, and development of acute lymphoblastic leukemia. Blood 93:
[64106] 2780-2790, 1999.

[64107] 10771.

[64108] Hunger, S. P.: Chromosomal translocations involving the E2A gene
[64109] in acute lymphoblastic leukemia: clinical features and molecular pathogenesis. Blood 87:
[64110] 1211-1224, 1996.

[64111] 10772.

[64112] Inaba, T.; Roberts, W. M.; Shapiro, L. H.; Jolly, K. W.; Rai-
mondi,
[64113] S. C.; Smith, S. D.; Look, A. T.: Fusion of the leucine zipper
gene
[64114] HLF to the E2A gene in human acute B-lineage leukemia.
Science 257:
[64115] 531-534, 1992.

[64116] 10773.

[64117] Smith, K. S.; Rhee, J. W.; Naumovski, L.; Cleary, M. L.: Dis-
rupted
[64118] differentiation and oncogenic transformation of lymphoid
progenitors
[64119] in E2A-HLF transgenic mice. Molec. Cell. Biol. 19:
4443-4451, 1999.
[64120] 10774.

[64121] Gripp, K. W.; Wotton, D.; Edwards, M. C.; Roessler, E.;
Ades, L.;
[64122] Meinecke, P.; Richieri-Costa, A.; Zackai, E. H.; Massague,
J.; Muenke,
[64123] M.; Elledge, S. J.: Mutations in TGIF cause holoprosen-
cephaly and

[64124] link NODAL signalling to human neural axis determination. Nature

[64125] Genet. 25: 205-208, 2000.

[64126] 10775.

[64127] Nanni, L.; Ming, J. E.; Bocian, M.; Steinhaus, K.; Bianchi, D.

[64128] W.; de Die-Smulders, C.; Giannotti, A.; Imaizumi, K.; Jones, K. L.;

[64129] Del Campo, M.; Martin, R. A.; Meinecke, P.; Pierpont, M. E. M.; Robin,

[64130] N. H.; Young, I. D.; Roessler, E.; Muenke, M.: The mutational spectrum

[64131] of the Sonic hedgehog gene in holoprosencephaly: SHH mutations cause

[64132] a significant proportion of autosomal dominant holoprosencephaly. Hum.

[64133] Molec. Genet. 8: 2479-2488, 1999.

[64134] 10776.

[64135] Acampora, D.; D'Esposito, M.; Faiella, A.; Pannese, M.; Migliaccio,

[64136] E.; Morelli, F.; Stornaiuolo, A.; Nigro, V.; Simeone, A.; Boncinelli,

[64137] E.: The human HOX gene family. Nucleic Acids Res. 17: 10385-10402,

[64138] 1989.

[64139] 10777.

[64140] Balling, R.; Mutter, G.; Gruss, P.; Kessel, M.: Craniofacial abnormalities

[64141] induced by ectopic expression of the homeobox gene Hox-1.1 in transgenic

[64142] mice. Cell 58: 337-347, 1989.

[64143] 10778.

[64144] Bucan, M.; Yang-Feng, T.; Colberg-Poley, A. M.; Wolge-muth, D. J.;

[64145] Guenet, J.-L.; Francke, U.; Lehrach, H.: Genetic and cyto-genetic

[64146] localisation of the homeo box containing genes on mouse chromosome

[64147] 6 and human chromosome 7. EMBO J. 5: 2899-2905, 1986.

[64148] 10779.

[64149] Ferguson-Smith, A. C.; Fienberg, A.; Ruddle, F. H.: Isola-tion,

[64150] chromosomal localization, and nucleotide sequence of the human HOX

[64151] 1.4 homeobox. Genomics 5: 250-258, 1989.

[64152] 10780.

[64153] Gaunt, S. J.; Singh, P. B.: Homeogene expression patterns and

[64154] chromosomal imprinting. Trends Genet. 6: 208-212, 1990.

[64155] 10781.

[64156] Gehring, W. J.: The homeo box: a key to the understand-ing of development? Cell 40:

[64157] 3-5, 1985.

[64158] 10782.

[64159] Joyner, A. L.; Lebo, R. V.; Kan, Y. W.; Tjian, R.; Cox, D. R.;

[64160] Martin, G. R.: Comparative chromosome mapping of a conserved homoeo

[64161] box region in mouse and human. Nature 314: 173-175, 1985.

[64162] 10783.

[64163] McGinnis, W.; Garber, R. L.; Wirz, J.; Kuroiwa, A.; Gehring, W.

[64164] J.: A homologous protein-coding sequence in Drosophila homeotic genes
[64165] and its conservation in other metazoans. Cell 37: 403-408, 1984.
[64166] 10784.

[64167] McGinnis, W.; Levine, M. S.; Hafen, E.; Kuroiwa, A.; Gehring, W.
[64168] J.: A conserved DNA sequence in homoeotic genes of the Drosophila
[64169] Antennapedia and bithorax complexes. Nature 308: 428-433, 1984.
[64170] 10785.

[64171] Ohno, S.: Evolution by Gene Duplication. Heidelberg: Springer
[64172] (pub.) 1970.

[64173] 10786.

[64174] Rabin, M.; Ferguson-Smith, A.; Hart, C. P.; Ruddle, F. H.: Cognate
[64175] homeo-box loci mapped on homologous human and mouse chromosomes. Proc.
[64176] Nat. Acad. Sci. 83: 9104-9108, 1986.

[64177] 10787.

[64178] Rabin, M.; Hart, C. P.; Ferguson-Smith, A.; McGinnis, W.; Levine,

[64179] M.; Ruddle, F. H.: Two homoeo box loci mapped in evolutionarily related

[64180] mouse and human chromosomes. Nature 314: 175-178, 1985.

[64181] 10788.

[64182] Rieger, R.; Michaelis, A.; Green, M. M.: Glossary of Genetics

[64183] and Cytogenetics. New York: Springer-Verlag (pub.) 1976. Pp.

[64184] 281 only.

[64185] 10789.

[64186] Schughart, K.; Kappen, C.; Ruddle, F. H.: Duplication of large

[64187] genomic regions during the evolution of vertebrate homeobox genes. Proc.

[64188] Nat. Acad. Sci. 86: 7067-7071, 1989.

[64189] 10790.

[64190] Scott, M. P.: Vertebrate homeobox gene nomenclature.

(Letter) Cell 71:
[64191] 551-553, 1992.

[64192] 10791.

[64193] Greer, J. M.; Puetz, J.; Thomas, K. R.; Capecchi, M. R.: Maintenance
[64194] of functional equivalence during paralogous Hox gene evolution. Nature 403:
[64195] 661-665, 2000.

[64196] 10792.

[64197] Apiou, F.; Flagiello, D.; Cillo, C.; Malfoy, B.; Poupon, M.-F.;

[64198] Dutrillaux, B.: Fine mapping of human HOX gene clusters. Cytogenet.
[64199] Cell Genet. 73: 114-115, 1996.

[64200] 10793.

[64201] Chan, H. Y. E.; Warrick, J. M.; Gray-Board, G. L.; Paulson, H.
[64202] L.; Bonini, N. M.: Mechanisms of chaperone suppression of polyglutamine
[64203] disease: selectivity, synergy and modulation of protein solubility
[64204] in Drosophila. Hum. Molec. Genet. 9: 2811-2820, 2000.

[64205] 10794.

[64206] Chandler, J. H.; Reed, T. E.; Dejong, R. N.: Huntington's chorea

[64207] in Michigan. Neurology 10: 148-153, 1960.

[64208] 10795.

[64209] Chase, T. N.; Wexler, N. S.; Barbeau, A.: Huntington's Disease.

[64210] Advances in Neurology. New York: Raven Press (pub.) 23: 1979.

[64211] 10796.

[64212] Chen, S.; Ferrone, F. A.; Wetzel, R.: Huntington's disease age-of-onset

[64213] linked to polyglutamine aggregation nucleation. Proc. Nat. Acad.

[64214] Sci. 99: 11884-11889, 2002.

[64215] 10797.

[64216] Cheng, S. V.; Martin, G. R.; Nadeau, J. H.; Haines, J. L.; Bu-

can,

[64217] M.; Kozak, C. A.; MacDonald, M. E.; Lockyer, J. L.; Ledley, F. D.;

[64218] Woo, S. L. C.; Lehrach, H.; Gilliam, T. C.; Gusella, J. F.: Synteny

[64219] on mouse chromosome 5 of homologs for human DNA loci linked to the

[64220] Huntington disease gene. Genomics 4: 419-426, 1989.

[64221] 10798.

[64222] Chong, S. S.; Almqvist, E.; Telenius, H.; LaTray, L.; Nichol,

[64223] K.; Bourdelat-Parks, B.; Goldberg, Y. P.; Haddad, B. R.; Richards,

[64224] F.; Sillence, D.; Greenberg, C. R.; Ives, E.; Van den Engh, G.; Hughes,

[64225] M. R.; Hayden, M. R.: Contribution of DNA sequence and CAG size to

[64226] mutation frequencies of intermediate alleles for Hunting-
ton disease:
[64227] evidence from single sperm analyses. Hum. Molec. Genet.
6: 301-309,
[64228] 1997.

[64229] 10799.

[64230] Clarke, G.; Collins, R. A.; Leavitt, B. R.; Andrews, D. F.;
Hayden,
[64231] M. R.; Lumsden, C. J.; McInnes, R. R.: A one-hit model of
cell death
[64232] in inherited neuronal degenerations. Nature 406:
195-199, 2000.
[64233] 10800.

[64234] Coles, R.; Leggo, J.; Rubinsztein, D. C.: Analysis of the
5-prime
[64235] upstream sequence of the Huntington's disease (HD) gene
shows six
[64236] new rare alleles which are unrelated to the age at onset of
HD. J.
[64237] Med. Genet. 34: 371-374, 1997.

[64238] 10801.

[64239] Collins, F. S.; Richards, J. E.; Cole, J. L.; Gilliam, T. C.;

[64240] Gusella, J. F.: Chromosome jumping from D4S10 (G8) toward the Huntington

[64241] disease gene. (Abstract) Cytogenet. Cell Genet. 46: 597, 1987.

[64242] 10802.

[64243] Conneally, P. M.: Huntington disease: genetics and epidemiology. Am.

[64244] J. Hum. Genet. 36: 506-526, 1984.

[64245] 10803.

[64246] Conneally, P. M.; Haines, J. L.; Tanzi, R. E.; Wexler, N. S.;

[64247] Penchaszadeh, G. K.; Harper, P. S.; Folstein, S. E.; Cassiman, J.

[64248] J.; Myers, R. H.; Young, A. B.; Hayden, M. R.; Falek, A.; Tolosa,

[64249] E. S.; Crespi, S.; Di Maio, L.; Holmgren, G.; Anvret, M.; Kanazawa,

[64250] I.; Gusella, J. F.: Huntington disease: no evidence for locus heterogeneity. Genomics 5:

[64251] 304-308, 1989.

[64252] 10804.

[64253] Craufurd, D.; Dodge, A.; Kerzin-Storrar, L.; Harris, R.: Uptake
[64254] of presymptomatic predictive testing for Huntington's disease. Lancet II:
[64255] 603-605, 1989.

[64256] 10805.

[64257] Critchley, M.: Great Britain and the early history of Huntington's
[64258] chorea. Advances in Neurology. New York: Raven Press (pub.) 1:
[64259] 1973.

[64260] 10806.

[64261] Critchley, M.: The history of Huntington's chorea. (Editorial) Psych.
[64262] Med. 14: 725-727, 1984.

[64263] 10807.

[64264] Davies, S. W.; Turmaine, M.; Cozens, B. A.; DiFiglia, M.; Sharp,
[64265] A. H.; Ross, C. A.; Scherzinger, E.; Wanker, E. E.; Mangiarini, L.;
[64266] Bates, G. P.: Formation of neuronal intranuclear inclusions

underlies

[64267] the neurological dysfunction in mice transgenic for the HD mutation. Cell 90:

[64268] 537-548, 1997.

[64269] 10808.

[64270] Decruyenaere, M.; Evers-Kiebooms, G.; Boogaerts, A.; Cassiman,

[64271] J.-J.; Cloostermans, T.; Demyttenaere, K.; Dom, R.; Fryns, J.-P.;

[64272] Van den Berghe, H.: Prediction of psychological functioning one year

[64273] after the predictive test for Huntington's disease and impact of the

[64274] test result on reproductive decision making. J. Med. Genet. 33:

[64275] 737-743, 1996.

[64276] 10809.

[64277] De Rooij, K. E.; Dorsman, J. C.; Smoor, M. A.; den Dunnen, J.

[64278] T.; Van Ommen, G.-J. B.: Subcellular localization of the Huntington's

[64279] disease gene product in cell lines by immunofluorescence

and biochemical
[64280] subcellular fractionation. Hum. Molec. Genet. 5:
1093-1099, 1996.
[64281] 10810.

[64282] DiFiglia, M.; Sapp, E.; Chase, K. O.; Davies, S. W.; Bates, G.

[64283] P.; Vonsattel, J. P.; Aronin, N.: Aggregation of huntingtin
in neuronal
[64284] intranuclear inclusions and dystrophic neurites in brain.
Science 277:
[64285] 1990-1993, 1997.

[64286] 10811.

[64287] Doggett, N. A.; Cheng, J.-F.; Smith, C. L.; Cantor, C. R.:
The
[64288] Huntington disease locus is most likely within 325 kilo-
bases of the
[64289] chromosome 4p telomere. Proc. Nat. Acad. Sci. 86:
10011-10014, 1989.
[64290] 10812.

[64291] Dunah, A. W.; Jeong, H.; Griffin, A.; Kim, Y.-M.; Standaert,
D.
[64292] G.; Hersch, S. M.; Mouradian, M. M.; Young, A. B.; Tanese,

N.; Krainc,

[64293] D.: Sp1 and TAFII130 transcriptional activity disrupted in early

[64294] Huntington's disease. Science 296: 2238-2243, 2002.

[64295] 10813.

[64296] Durbach, N.; Hayden, M. R.: George Huntington: the man behind

[64297] the eponym. J. Med. Genet. 30: 406-409, 1993.

[64298] 10814.

[64299] Dure, L. S., IV; Landwehrmeyer, G. B.; Golden, J.; McNeil, S.

[64300] M.; Ge, P.; Aizawa, H.; Huang, Q.; Ambrose, C. M.; Duyao, M. P.; Bird,

[64301] E. D.; DiFiglia, M.; Gusella, J. F.; MacDonald, M. E.; Penney, J.

[64302] B.; Young, A. B.; Vonsattel, J.-P.: IT15 gene expression in fetal

[64303] human brain. Brain Res. 659: 33-41, 1994.

[64304] 10815.

[64305] Duyao, M.; Ambrose, C.; Myers, R.; Novelletto, A.; Per-sichetti,

[64306] F.; Frontali, M.; Folstein, S.; Ross, C.; Franz, M.; Abbott,

M.; Gray,
[64307] J.; Conneally, P.; and 30 others: Trinucleotide repeat length instability
[64308] and age of onset in Huntington's disease. Nature Genet. 4: 387-392,
[64309] 1993.

[64310] 10816.

[64311] Duyao, M. P.; Auerbach, A. B.; Ryan, A.; Persichetti, F.; Barnes,
[64312] G. T.; McNeil, S. M.; Ge, P.; Vonsattel, J.-P.; Gusella, J. F.; Joyner,
[64313] A. L.; MacDonald, M. E.: Inactivation of the mouse Hunt-ington's disease
[64314] gene homolog Hdh. Science 269: 407-410, 1995.

[64315] 10817.

[64316] Dyer, R. B.; McMurray, C. T.: Mutant protein in Huntington disease
[64317] is resistant to proteolysis in affected brain. Nature Genet. 29:
[64318] 270-278, 2001.

[64319] 10818.

[64320] Enna, S. J.; Bird, E. D.; Bennett, J. P., Jr.; Bylund, D. B.;

[64321] Yamamura, H. I.; Iversen, L. L.; Snyder, S. H.: Huntington's chorea:

[64322] changes in neurotransmitter receptors in the brain. New Eng. J. Med. 294:

[64323] 1305-1309, 1976.

[64324] 10819.

[64325] Erickson, R. P.: Chromosomal imprinting and the parent transmission

[64326] specific variation in expressivity of Huntington disease. (Letter) Am.

[64327] J. Hum. Genet. 37: 827-829, 1985.

[64328] 10820.

[64329] Faber, P. W.; Barnes, G. T.; Srinidhi, J.; Chen, J.; Gusella,

[64330] J. F.; MacDonald, M. E.: Huntingtin interacts with a family of WW

[64331] domain proteins. Hum. Molec. Genet. 7: 1463-1474, 1998.

[64332] 10821.

[64333] Man in't Veld, A. J.; Boomsma, F.; Moleman, P.; Schalekamp, M.

[64334] A. D. H.: Congenital dopamine-beta-hydroxylase deficiency: a novel
[64335] orthostatic syndrome. Lancet I: 183-188, 1987.

[64336] 10822.

[64337] Robertson, D.; Goldberg, M. R.; Onrot, J.; Hollister, A. S.; Wiley,
[64338] R.; Thompson, J. G., Jr.; Robertson, R. M.: Isolated failure of autonomic
[64339] noradrenergic neurotransmission: evidence for impaired beta-hydroxylation
[64340] of dopamine. New Eng. J. Med. 314: 1494-1497, 1986.

[64341] 10823.

[64342] Ogawa, O.; McNoe, L. A.; Eccles, M. R.; Morison, I. M.; Reeve,
[64343] A. E.: Human insulin-like growth factor type I and type II receptors
[64344] are not imprinted. Hum. Molec. Genet. 2: 2163-2165, 1993.

[64345] 10824.

[64346] Oshima, A.; Nolan, C. M.; Kyle, J. W.; Grubb, J. H.; Sly, W. S.

[64347] : The human cation-independent mannose 6-phosphate

receptor: cloning
[64348] and sequence of the full-length cDNA and expression of functional
[64349] receptor in COS cells. J. Biol. Chem. 263: 2553-2562, 1988.
[64350] 10825.

[64351] Puertollano, R.; Aguilar, R. C.; Gorshkova, I.; Crouch, R. J.;

[64352] Bonifacino, J. S.: Sorting of mannose 6-phosphate receptors mediated
[64353] by the GGAs. Science 292: 1712-1716, 2001.

[64354] 10826.

[64355] Rao, P. H.; Murty, V. V. V. S.; Gaidano, G.; Hauptschein, R.;

[64356] Dalla-Favera, R.; Chaganti, R. S. K.: Subregional mapping of 8 single
[64357] copy loci to chromosome 6 by fluorescence in situ hybridization. Cytogenet.
[64358] Cell Genet. 66: 272-273, 1994.

[64359] 10827.

[64360] Roth, R. A.: Structure of the receptor for insulin-like growth
[64361] factor II: the puzzle amplified. Science 239: 1269-1271,

1988.

[64362] 10828.

[64363] Sleutels, F.; Zwart, R.; Barlow, D. P.: The non-coding Air RNA

[64364] is required for silencing autosomal imprinted genes. Nature 415:

[64365] 810-813, 2002.

[64366] 10829.

[64367] Souza, R. F.; Appel, R.; Yin, J.; Wang, S.; Smolinski, K. N.;

[64368] Abraham, J. M.; Zou, T.-T.; Shi, Y.-Q.; Lei, J.; Cottrell, J.; Cymes,

[64369] K.; Biden, K.; Simms, L.; Leggett, B.; Lynch, P. M.; Frazier, M.;

[64370] Powell, S. M.; Harpaz, N.; Sugimura, H.; Young, J.; Meltzer, S. J.

[64371] : Microsatellite instability in the insulin-like growth factor II

[64372] receptor gene in gastrointestinal tumours. (Letter) Nature Genet. 14:

[64373] 255-257, 1996.

[64374] 10830.

[64375] Szebeny, G.; Rotwein, P.: The mouse insulin-like growth factor
[64376] II/cation-independent mannose 6-phosphate (IGF-II/MPR) receptor gene:
[64377] molecular cloning and genomic organization. Genomics 19: 120-129,
[64378] 1994.

[64379] 10831.

[64380] Tong, P. Y.; Tollefsen, S. E.; Kornfeld, S.: The cation-independent
[64381] mannose 6-phosphate receptor binds insulin-like growth factor II. J.
[64382] Biol. Chem. 263: 2585-2588, 1988.

[64383] 10832.

[64384] Waheed, A.; Braulke, T.; Junghans, U.; von Figura, K.: Man-nose
[64385] 6-phosphate/insulin like growth factor II receptor: the two types
[64386] of ligands bind simultaneously to one receptor at different sites. Biochem.
[64387] Biophys. Res. Commun. 152: 1248-1254, 1988.

[64388] 10833.

[64389] Xu, Y.; Goodyer, C. G.; Deal, C.; Polychronakos, C.: Functional

[64390] polymorphism in the parental imprinting of the human IGF2R gene. Biochem.

[64391] Biophys. Res. Commun. 197: 747-754, 1993.

[64392] 10834.

[64393] Young, L. E.; Fernandes, K.; McEvoy, T. G.; Butterwith, S. C.;

[64394] Gutierrez, C. G.; Carolan, C.; Broadbent, P. J.; Robinson, J. J.;

[64395] Wilmut, I.; Sinclair, K. D.: Epigenetic change in IGF2R is associated

[64396] with fetal overgrowth after sheep embryo culture. Nature Genet. 27:

[64397] 153-154, 2001.

[64398] 10835.

[64399] Zhu, Y.; Doray, B.; Poussu, A.; Lehto, V.-P.; Kornfeld, S.: Binding

[64400] of GGA2 to the lysosomal enzyme sorting motif of the mannose 6-phosphate

[64401] receptor. Science 292: 1716-1718, 2001.

[64402] 10836.

[64403] Brown, C. W.; Houston-Hawkins, D. E.; Woodruff, T. K.; Matzuk,

[64404] M. M.: Insertion of Inhbb into the Inhba locus rescues the Inhba-null

[64405] phenotype and reveals new activin functions. Nature Genet. 25: 453-457,

[64406] 2000.

[64407] 10837.

[64408] Matzuk, M. M.; Kumar, T. R.; Vassalli, A.; Bickenbach, J. R.; Roop,

[64409] D. R.; Jaenisch, R.; Bradley, A.: Functional analysis of activins

[64410] during mammalian development. Nature 374: 354-356, 1995.

[64411] 10838.

[64412] Mellor, S. L.; Cranfield, M.; Ries, R.; Pedersen, J.; Cancilla,

[64413] B.; de Kretser, D.; Groome, N. P.; Mason, A. J.; Risbridger, G. P.

[64414] : Localization of activin beta(A)-, beta(B)-, and

beta(C)-subunits

[64415] in human prostate and evidence for formation of new activin heterodimers

[64416] of beta(C)-subunit. J. Clin. Endocr. Metab. 85: 4851-4858, 2000.

[64417] 10839.

[64418] You, L.; Kruse, F. E.: Differential effect of activin A and BMP-7

[64419] on myofibroblast differentiation and the role of the Smad signaling

[64420] pathway. Invest. Ophthal. Vis. Sci. 43: 72-81, 2002.

[64421] 10840.

[64422] Valverde, P.; Healy, E.; Jackson, I.; Rees, J. L.; Thody, A. J.

[64423] : Variants of the melanocyte-stimulating hormone receptor gene are

[64424] associated with red hair and fair skin in humans. Nature Genet. 11:

[64425] 328-330, 1995.

[64426] 10841.

[64427] Valverde, P.; Healy, E.; Sikkink, S.; Haldane, F.; Thody, A. J.;

[64428] Carothers, A.; Jackson, I. J.; Rees, J. L.: The Asp84Glu variant

[64429] of the melanocortin 1 receptor (MC1R) is associated with melanoma. Hum.

[64430] Molec. Genet. 5: 1663-1666, 1996.

[64431] 10842.

[64432] Valverde, P.; Healy, F.; Jackson, I.; Rees, J. L.; Thody, A. J.

[64433] : Variants of the melanocyte-stimulating hormone receptor gene are

[64434] associated with red hair and fair skin in humans. Nature Genet. 11:

[64435] 328-330, 1995.

[64436] 10843.

[64437] van der Velden, P. A.; Sandkuijl, L. A.; Bergman, W.; Pavel, S.;

[64438] van Mourik, L.; Frants, R. R.; Gruis, N. A.: Melanocortin-1 receptor

[64439] variant R151C modifies melanoma risk in Dutch families with melanoma. Am.

[64440] J. Hum. Genet. 69: 774-779, 2001.

[64441] 10844.

[64442] Xu, X.; Thornwall, M.; Lundin, L.-G.; Chhajlani, V.: Val92Met
[64443] variant of the melanocyte stimulating hormone receptor gene. (Letter) Nature
[64444] Genet. 14: 384 only, 1996.

[64445] 10845.

[64446] Becher, R.; Gibas, Z.; Sandberg, A. A.: Chromosome 6 in malignant
[64447] melanoma. Cancer Genet. Cytogenet. 9: 173-175, 1983.

[64448] 10846.

[64449] Fountain, J. W.; Karayiorgou, M.; Ernstoff, M. S.; Kirkwood, J.
[64450] M.; Vlock, D. R.; Titus-Ernstoff, L.; Bouchard, B.; Vi-jayasaradhi,
[64451] S.; Houghton, A. N.; Lahti, J.; Kidd, V. J.; Housman, D. E.; Dracopoli,
[64452] N. C.: Homozygous deletions within human chromosome band 9p21 in
[64453] melanoma. Proc. Nat. Acad. Sci. 89: 10557-10561, 1992.

[64454] 10847.

[64455] Lacy, D. A.; Wang, Z.-E.; Symula, D. J.; McArthur, C. J.; Ru-

bin,

[64456] E. M.; Frazer, K. A.; Locksley, R. M.: Faithful expression of the

[64457] human 5q31 cytokine cluster in transgenic mice. J. Immun. 164: 4569-4574,

[64458] 2000.

[64459] 10848.

[64460] Taniyama, Y.; Kuroki, R.; Omura, F.; Seko, C.; Kikuchi, M.: Evidence

[64461] for intramolecular disulfide bond shuffling in the folding of mutant

[64462] human lysozyme. J. Biol. Chem. 266: 6456-6461, 1991.

[64463] 10849.

[64464] Loots, G. G.; Locksley, R. M.; Blankespoor, C. M.; Wang, Z. E.;

[64465] Miller, W.; Rubin, E. M.; Frazer, K. A.: Identification of a coordinate

[64466] regulator of interleukins 4, 13, and 5 by cross-species sequence comparisons. Science 288:

[64467] 136-140, 2000.

[64468] 10850.

[64469] McKenzie, A. N. J.; Culpepper, J. A.; de Waal Malefyt, R.; Briere,

[64470] F.; Punnonen, J.; Aversa, G.; Sato, A.; Dang, W.; Cocks, B. G.; Menon,

[64471] S.; de Vries, J. E.; Banchereau, J.; Zurawski, G.: Interleukin 13,

[64472] a T-cell-derived cytokine that regulates human monocyte and B-cell

[64473] function. Proc. Nat. Acad. Sci. 90: 3735-3739, 1993.

[64474] 10851.

[64475] Minty, A.; Chalon, P.; Derocq, J.-M.; Dumont, X.; Guille-mot, J.-C.;

[64476] Kaghad, M.; Labit, C.; Leplatois, P.; Liauzun, P.; Miloux, B.; Minty,

[64477] C.; Casellas, P.; Loison, G.; Lupker, J.; Shire, D.; Ferrara, P.;

[64478] Caput, D.: Interleukin-13 is a new human lymphokine regulating inflammatory

[64479] and immune responses. Nature 362: 248-250, 1993.

[64480] 10852.

[64481] Morgan, J. G.; Dolganov, G. M.; Robbins, S. E.; Hinton, L. M.;

[64482] Lovett, M.: The selective isolation of novel cDNAs encoded by the
[64483] regions surrounding the human interleukin 4 and 5 genes. Nucleic
[64484] Acids Res. 20: 5173-5179, 1992.

[64485] 10853.

[64486] Paul, W. E.: Personal Communication. Bethesda, Md. 3/6/1993.
[64487] 10854.

[64488] Punnonen, J.; Aversa, G.; Cocks, B. G.; McKenzie, A. N. J.; Menon,
[64489] S.; Zurawski, G.; de Waal Malefyt, R.; de Vries, J. E.: Inter-
leukin
[64490] 13 induces interleukin 4-independent IgG4 and IgE syn-
thesis and CD23
[64491] expression by human B cells. Proc. Nat. Acad. Sci. 90: 3730-3734,
[64492] 1993.

[64493] 10855.

[64494] Smirnov, D. V.; Smirnova, M. G.; Korobko, V. G.; Frolova, E. I.

[64495] : Tandem arrangement of human genes for interleukin-4 and interleukin-13:
[64496] resemblance in their organization. Gene 155: 277-281, 1995.
[64497] 10856.

[64498] Wills-Karp, M.; Luyimbazi, J.; Xu, X.; Schofield, B.; Neben, T.
[64499] Y.; Karp, C. L.; Donaldson, D. D.: Interleukin-13: central mediator
[64500] of allergic asthma. Science 282: 2258-2261, 1998.

[64501] 10857.

[64502] Zhu, Z.; Homer, R. J.; Wang, Z.; Chen, Q.; Geba, G. P.; Wang,
[64503] J.; Zhang, Y.; Elias, J. A.: Pulmonary expression of interleukin-13
[64504] causes inflammation, mucus hypersecretion, subepithelial fibrosis,
[64505] physiologic abnormalities, and eotaxin production. J. Clin. Invest. 103:
[64506] 779-788, 1999.

[64507] 10858.

[64508] Zurawski, G.; de Vries, J. E.: Interleukin 13 elicits a subset

[64509] of the activities of its close relative interleukin 4. Stem Cells 12:

[64510] 169-174, 1994.

[64511] 10859.

[64512] Li, C.; Lai, C.; Sigman, D. S.; Gaynor, R. B.: Cloning of a cellular

[64513] factor, interleukin binding factor, that binds to NFAT-like motifs

[64514] in the human immunodeficiency virus long terminal repeat. Proc. Nat.

[64515] Acad. Sci. 88: 7739-7743, 1991.

[64516] 10860.

[64517] Li, C.; Lusis, A. J.; Sparkes, R.; Nirula, A.; Gaynor, R.: Characterization

[64518] and chromosomal mapping of the gene encoding the cellular DNA binding

[64519] protein ILF. Genomics 13: 665-671, 1992.

[64520] 10861.

[64521] Chebath, J.; Merlin, G.; Metz, R.; Benech, P.; Revel, M.: Interferon-induced

[64522] 56,000 Mr protein and its mRNA in human cells: molecu-
lar cloning and
[64523] partial sequence of the cDNA. Nucleic Acids Res. 11:
1213-1226,
[64524] 1983.

[64525] 10862.

[64526] Kusari, J.; Szabo, P.; Grzeschik, K.-H.; Sen, G. C.: Chromo-
somal
[64527] localization of the interferon-inducible human gene en-
coding mRNA
[64528] 561. J. Interferon Res. 7: 53-59, 1987.

[64529] 10863.

[64530] Boone, C.; Chen, T.-R.; Ruddle, F. H.: Assignment of three
human
[64531] genes to chromosomes (LDH-A to 11, TK to 17 and IDH to
20) and evidence
[64532] for translocation between human and mouse chromo-
somes in somatic cell
[64533] hybrids. Proc. Nat. Acad. Sci. 68: 510-514, 1972.

[64534] 10864.

[64535] Chen, S.-H.; Fossum, B. L. G.; Giblett, E. R.: Genetic varia-

tion

[64536] of the soluble form of NADP-dependent isocitric dehydro-genase in man. Am.

[64537] J. Hum. Genet. 24: 325-329, 1972.

[64538] 10865.

[64539] Creagan, R. P.; Carritt, B.; Chen, S.-H.; Kucherlapati, R. S.;

[64540] McMorris, F. A.; Ricciuti, F.; Tan, Y. H.; Tischfield, J. A.; Ruddle,

[64541] F. H.: Chromosome assignments of genes in man using mouse-human somatic

[64542] cell hybrids: cytoplasmic isocitrate dehydrogenase (IDH 1) and malate

[64543] dehydrogenase (MDH 1) to chromosome 2. Am. J. Hum. Genet. 26: 604-613,

[64544] 1974.

[64545] 10866.

[64546] Glass, I. A.; Swindlehurst, C. A.; Aitken, D. A.; McCrea, W.; Boyd,

[64547] E.: Interstitial deletion of the long arm of chromosome 2 with normal

[64548] levels of isocitrate dehydrogenase. J. Med. Genet. 26:

127-130,
[64549] 1989.

[64550] 10867.

[64551] Henderson, N. S.: Isozymes of isocitrate dehydrogenase: subunit
[64552] structure and intracellular location. J. Exp. Zool. 158: 263-273,
[64553] 1965.

[64554] 10868.

[64555] Henderson, N. S.: Intracellular location and genetic control of
[64556] isozymes of NADP-dependent isocitrate dehydrogenase and malate dehydrogenase. Ann.
[64557] N.Y. Acad. Sci. 151: 429-440, 1968.

[64558] 10869.

[64559] Narahara, K.; Kimura, S.; Kikkawa, K.; Takahashi, Y.; Wakita, Y.;
[64560] Kasai, R.; Nagai, S.; Nishibayashi, Y.; Kimoto, H.: Probable assignment
[64561] of soluble isocitrate dehydrogenase (IDH-1) to 2q33.3. Hum. Genet. 71:

[64562] 37-40, 1985.

[64563] 10870.

[64564] Ruddle, F. H.: Linkage analysis in man by somatic cell genetics. Nature 242:

[64565] 165-169, 1973.

[64566] 10871.

[64567] Shows, T. B.: . (Abstract) 4th Int. Cong. Hum. Genet., Paris 165,

[64568] 1971.

[64569] 10872.

[64570] Shows, T. B.: Genetics of human-mouse somatic cell hybrids: linkage

[64571] of human genes for isocitrate dehydrogenase and malate dehydrogenase. Biochem.

[64572] Genet. 7: 193-204, 1972.

[64573] 10873.

[64574] Turner, B. M.; Fisher, R. A.; Garthwaite, E.; Whale, R. J.; Harris,

[64575] H.: An account of two new ICD-S variants not detectable in red blood

[64576] cells. Ann. Hum. Genet. 37: 469-476, 1974.

[64577] 10874.

[64578] Van Cong, N.: Personal Communication. Paris, France 1976.

[64579] 10875.

[64580] Weil, D.; Van Cong, N.; Finaz, C.; Rebourcet, R.; Cochet, C.;

[64581] de Grouchy, J.; Frezal, J.: Localisation regionale des genes humains

[64582] IDH-S, MDH-S, PGK, alpha-GAL, G6PD par l'hybridation cellulaire interspecifique. Hum.

[64583] Genet. 36: 205-211, 1977.

[64584] 10876.

[64585] Auron, P. E.; Webb, A. C.; Rosenwasser, L. J.; Mucci, S. F.; Rich,

[64586] A.; Wolff, S. M.; Dinarello, C. A.: Nucleotide sequence of human

[64587] monocyte interleukin 1 precursor cDNA. Proc. Nat. Acad. Sci. 81:

[64588] 7907-7911, 1984.

[64589] 10877.

[64590] Winter, H.; Labreze, C.; Chapalain, V.; Surleve-Bazeille, J. E.;

[64591] Mercier, M.; Rogers, M. A.; Taieb, A.; Schweizer, J.: A variable

[64592] monilethrix phenotype associated with a novel mutation, glu402lys,

[64593] in the helix termination motif of the type II hair keratin hHb1. J.

[64594] Invest. Derm. 111: 169-172, 1998.

[64595] 10878.

[64596] Francke, U.; Foellmer, B. E.; Haynes, B. F.: Chromosome mapping

[64597] of human cell surface molecules: monoclonal anti-human lymphocyte

[64598] antibodies 4F2, A3D8, and A1G3 define antigens controlled by different

[64599] regions of chromosome 11. Somat. Cell Genet. 9: 333-344, 1983.

[64600] 10879.

[64601] Gottesdiener, K. M.; Karpinski, B. A.; Lindsten, T.; Strominger,

[64602] J. L.; Jones, N. H.; Thompson, C. B.; Leiden, J. M.: Isolation

and
[64603] structural characterization of the human 4F2 heavy-chain gene, an
[64604] inducible gene involved in T-lymphocyte activation. Molec. Cell.
[64605] Biol. 8: 3809-3819, 1988.

[64606] 10880.

[64607] Haynes, B. F.; Hemler, M. E.; Mann, D. L.; Eisenbarth, G. S.; Shelhamer,
[64608] J.; Mostowski, H. S.; Thomas, C. A.; Strominger, J. L.; Fauci, A.
[64609] S.: Characterization of a monoclonal antibody (4F2) that binds to
[64610] human monocytes and to a subset of activated lympho- cytes. J. Immun. 126:
[64611] 1409-1414, 1981.

[64612] 10881.

[64613] Hemler, M. E.; Strominger, J. L.: Characterization of anti- gen
[64614] recognized by the monoclonal antibody (4F2): different molecular forms
[64615] on human T and B lymphoblastoid cell lines. J. Immun.

129: 623-628,
[64616] 1982.

[64617] 10882.

[64618] Lindsten, T.; June, C. H.; Thompson, C. B.; Leiden, J. M.: Regulation
[64619] of 4F2 heavy-chain gene expression during normal human T-cell activation
[64620] can be mediated by multiple distinct molecular mechanisms. Molec.
[64621] Cell. Biol. 8: 3820-3826, 1988.

[64622] 10883.

[64623] Lumadue, J. A.; Glick, A. B.; Ruddle, F. H.: Cloning, sequence
[64624] analysis, and expression of the large subunit of the human lymphocyte
[64625] activation antigen 4F2. Proc. Nat. Acad. Sci. 84: 9204-9208, 1987.
[64626] 10884.

[64627] Mastroberardino, L.; Spindler, B.; Pfeiffer, R.; Skelly, P. J.;

[64628] Loffing, J.; Shoemaker, C. B.; Verrey, F.: Amino-acid transport by

[64629] heterodimers of 4F2hc/CD98 and members of a permease family. Nature 395:

[64630] 288-291, 1998.

[64631] 10885.

[64632] Cohen-Haguenauer, O.; Barton, P. J. R.; Van Cong, N.; Cohen, A.;

[64633] Masset, M.; Buckingham, M.; Frezal, J.: Chromosomal assignment of

[64634] two myosin alkali light-chain genes encoding the ventricular/slow

[64635] skeletal muscle isoform and the atrial/fetal muscle isoform (MYL3,

[64636] MYL4). Hum. Genet. 81: 278-282, 1989.

[64637] 10886.

[64638] Seharaseyon, J.; Bober, E.; Hsieh, C.-L.; Fodor, W. L.; Francke,

[64639] U.; Arnold, H.-H.; Vanin, E. F.: Human embryonic/atrial myosin alkali

[64640] light chain gene: characterization, sequence and chromosomal location. Genomics 7:

[64641] 289-293, 1990.

[64642] 10887.

[64643] Strohman, R. C.; Micou-Eastwood, J.; Glass, C. A.; Mat-
suda, R.

[64644] : Human fetal muscle and cultured myotubes derived from
it contain

[64645] a fetal-specific myosin light chain. Science 221: 955-957,
1983.

[64646] 10888.

[64647] Copeland, N. G.; Hutchison, K. W.; Jenkins, N. A.: Excision
of

[64648] the DBA ecotropic provirus in dilute coat-color revertants
of mice

[64649] occurs by homologous recombination involving the viral
LTRs. Cell 33:

[64650] 379-387, 1983.

[64651] 10889.

[64652] El-Husseini, A. E.; Vincent, S. R.: Cloning and characteri-
zation

[64653] of a novel RING finger protein that interacts with class V
myosins. J.

[64654] Biol. Chem. 274: 19771-19777, 1999.

[64655] 10890.

[64656] Engle, L. J.; Kennett, R. H.: Cloning, analysis, and chromo-
somal
[64657] localization of myoxin (MYH12): the human homologue to
the mouse dilute
[64658] gene. Genomics 19: 407-416, 1994.

[64659] 10891.

[64660] Espreafico, E. M.; Coling, D. E.; Tsakraklides, V.; Krogh, K.;

[64661] Wolenski, J. S.; Kalinec, G.; Kachar, B.: Localization of
myosin-V
[64662] in the centrosome. Proc. Nat. Acad. Sci. 95: 8636-8641,
1998.
[64663] 10892.

[64664] Jenkins, N. A.; Copeland, N. G.; Taylor, B. A.; Lee, B. K.:
Dilute
[64665] (d) coat colour mutation of DNA/2J mice is associated with
the site
[64666] of integration of an ecotropic MuLV genome. Nature 293:
370-374,
[64667] 1981.

[64668] 10893.

[64669] Jones, J. M.; Huang, J.-D.; Mermall, V.; Hamilton, B. A.; Mooseker,
[64670] M. S.; Escayg, A.; Copeland, N. G.; Jenkins, N. A.; Meisler, M. H.
[64671] : The mouse neurological mutant flailer expresses a novel hybrid gene
[64672] derived by exon shuffling between Gnb5 and Myo5a. Hum. Molec. Genet. 9:
[64673] 821-828, 2000.

[64674] 10894.

[64675] Karcher, R. L.; Roland, J. T.; Zappacosta, F.; Huddleston, M. J.;
[64676] Annan, R. S.; Carr, S. A.; Gelfand, V. I.: Cell cycle regula-tion
[64677] of myosin-V by calcium/calmodulin-dependent protein kinase II. Science 293:
[64678] 1317-1320, 2001.

[64679] 10895.

[64680] Ptacek, L. J.; Trimmer, J. S.; Agnew, W. S.; Roberts, J. W.; Petajan,
[64681] J. H.; Leppert, M.: Paramyotonia congenita and hyper-kalemic periodic

[64682] paralysis map to the same sodium-channel gene locus. Am. J. Hum.

[64683] Genet. 49: 851-854, 1991.

[64684] 10896.

[64685] Jodice, C.; Malaspina, P.; Persichetti, F.; Novelletto, A.; Spadaro,

[64686] M.; Giunti, P.; Morocutti, C.; Terrenato, L.; Harding, A. E.; Frontali,

[64687] M.: Effect of trinucleotide repeat length and parental sex on phenotypic

[64688] variation in spinocerebellar ataxia 1. Am. J. Hum. Genet. 54: 959-965,

[64689] 1994.

[64690] 10897.

[64691] Klement, I. A.; Skinner, P. J.; Kaytor, M. D.; Yi, H.; Hersch,

[64692] S. M.; Clark, H. B.; Zoghbi, H. Y.; Orr, H. T.: Ataxin-1 nu-clear

[64693] localization and aggregation: role in polyglutamine-in-duced disease

[64694] in SCA1 transgenic mice. Cell 95: 41-53, 1998.

[64695] 10898.

[64696] Klockgether, T.; Burk, K.; Schulz, J. B.; Dichgans, J.; Wes-
sel,
[64697] K.; Auburger, G.: Absence of SCA1 mutation in idiopathic
cerebellar
[64698] ataxia. (Letter) J. Neurol. Neurosurg. Psychiat. 57:
1439-1440,
[64699] 1994.

[64700] 10899.

[64701] Lunkes, A.; Goldfarb, L. G.; Platonov, F. A.; Alexeev, V. P.;

[64702] Duenas-Barajas, E.; Gajdusek, D. C.; Auburger, G.: Auto-
somal dominant
[64703] spinocerebellar ataxia (SCA) in a Siberian founder popula-
tion: assignment
[64704] to the SCA1 locus. Exp. Neurol. 126: 310-312, 1994.

[64705] 10900.

[64706] Matilla, T.; Volpini, V.; Genis, D.; Rosell, J.; Corral, J.;
Davalos,
[64707] A.; Molins, A.; Estivill, X.: Presymptomatic analysis of
spinocerebellar
[64708] ataxia type 1 (SCA1) via the expansion of the SCA1 CAG-
repeat in a

[64709] large pedigree displaying anticipation and parental male bias. Hum.

[64710] Molec. Genet. 2: 2123-2128, 1993.

[64711] 10901.

[64712] Orr, H. T.; Chung, M.; Banfi, S.; Kwiatkowski, T. J., Jr.; Ser-

[64713] vadio, A.; Beaudet, A. L.; McCall, A. E.; Duvick, L. A.; Ranum, L. P. W.;

[64714] Zoghbi, H. Y.: Expansion of an unstable trinucleotide CAG repeat

[64715] in spinocerebellar ataxia type 1. Nature Genet. 4: 221-226, 1993.

[64716] 10902.

[64717] Juliano, R. L.; Ling, V.: A surface glycoprotein modulating drug

[64718] permeability in Chinese hamster ovary cell mutants. Biochim. Biophys.

[64719] Acta 455: 152-162, 1976.

[64720] 10903.

[64721] Lincke, C. R.; Smit, J. J.; van der Velde Koerts, T.; Borst, P.

[64722] : Structure of the human MDR3 gene and physical map-

ping of the human
[64723] MDR locus. J. Biol. Chem. 266: 5303-5310, 1991.

[64724] 10904.

[64725] Maggiore, G.; Bernard, O.; Hadchouel, M.; Lemonnier, A.; Alagille,
[64726] D.: Diagnostic value of serum gamma-glutamyl transpep-tidase activity
[64727] in liver diseases in children. J. Pediat. Gastroent. Nutr. 12: 21-26,
[64728] 1991.

[64729] 10905.

[64730] Maggiore, G.; Bernard, O.; Riely, C. A.; Hadchouel, M.; Lemonnier,
[64731] A.; Alagille, D.: Normal serum gamma-glu-tamyl-transpeptidase activity
[64732] identifies groups of infants with idiopathic cholestasis with poor
[64733] prognosis. J. Pediat. 111: 251-252, 1987.

[64734] 10906.

[64735] Offerhaus, J.: Personal Communication. Amsterdam, The Netherlands

[64736] 1/24/1994.

[64737] 10907.

[64738] Ruetz, S.; Gros, P.: Phosphatidylcholine translocase: a physiological

[64739] role for the mdr2 gene. Cell 77: 1071-1081, 1994.

[64740] 10908.

[64741] Smit, J. J. M.; Schinkel, A. H.; Oude Elferink, R. P. J.; Groen,

[64742] A. K.; Wagenaar, E.; van Deemter, L.; Mol, C. A. A. M.; Ottenhoff,

[64743] R.; van der Lugt, N. M. T.; van Roon, M. A.; van der Valk, M. A.;

[64744] Offerhaus, G. J. A.; Berns, A. J. M.; Borst, P.: Homozygous disruption

[64745] of the murine mdr2 P-glycoprotein gene leads to a complete absence

[64746] of phospholipid from bile and to liver disease. Cell 75: 451-462,

[64747] 1993.

[64748] 10909.

[64749] Van der Bliek, A. M.; Baas, F.; Ten Houte de Lange, T.; Koolman,

[64750] P. M.; Van der Velde-Koerts, T.; Borst, P.: The human mdr3 gene encodes
[64751] a novel P-glycoprotein homologue and gives rise to alternatively spliced
[64752] mRNAs in liver. EMBO J. 6: 3325-3331, 1987.

[64753] 10910.

[64754] Whitington, P. F.; Freese, D. K.; Alonso, E. M.; Schwarzenberg,
[64755] S. J.; Sharp, H. L.: Clinical and biochemical findings in progressive
[64756] familial intrahepatic cholestasis. J. Pediat. Gastroent. Nutr. 18:
[64757] 134-141, 1994.

[64758] 10911.

[64759] Barbosa, M. D. F. S.; Nguyen, Q. A.; Tchernev, V. T.; Ashley, J.
[64760] A.; Detter, J. C.; Blaydes, S. M.; Brandt, S. J.; Chotai, D.; Hodgman,
[64761] C.; Solari, R. C. E.; Lovett, M.; Kingsmore, S. F.: Identification
[64762] of the homologous beige and Chediak-Higashi syndrome genes. Nature 382:

[64763] 262-265, 1996.

[64764] 10912.

[64765] Fukai, K.; Oh, J.; Karim, M. A.; Moore, K. J.; Kandil, H. H.; Ito,

[64766] H.; Burger, J.; Spritz, R. A.: Homozygosity mapping of the gene for

[64767] Chediak-Higashi syndrome to chromosome 1q42-q44 in a segment of conserved

[64768] synteny that includes the mouse beige locus (bg). Am. J. Hum. Genet. 59:

[64769] 620-624, 1996.

[64770] 10913.

[64771] Jackson, I. J.: Homologous pigmentation mutations in human, mouse

[64772] and other model organisms. Hum. Molec. Genet. 6: 1613-1624, 1997.

[64773] 10914.

[64774] Karim, M. A.; Nagle, D. L.; Kandil, H. H.; Burger, J.; Moore, K.

[64775] J.; Spritz, R. A.: Mutations in the Chediak-Higashi syndrome gene

[64776] (CHS1) indicate requirement for the complete 3801 amino acid CHS protein. Hum.

[64777] Molec. Genet. 6: 1087-1089, 1997.

[64778] 10915.

[64779] Nagle, D. L.; Karim, M. A.; Woolf, E. A.; Holmgren, L.; Bork,

[64780] P.; Misumi, D. J.; McGrail, S. H.; Dussault, B. J., Jr.; Perou, C.

[64781] M.; Boissy, R. E.; Duyk, G. M.; Spritz, R. A.; Moore, K. J.: Identification

[64782] and mutation analysis of the complete gene for Chediak-Higashi syndrome. Nature

[64783] Genet. 14: 307-311, 1996.

[64784] 10916.

[64785] Perou, C. M.; Moore, K. J.; Nagle, D. L.; Misumi, D. J.; Woolf,

[64786] E. A.; McGrail, S. H.; Holmgren, L.; Brody, T. H.; Dussault, B. J.,

[64787] Jr.; Monroe, C. A.; Duyk, G. M.; Pryor, R. J.; Li, L.; Justice, M.

[64788] J.; Kaplan, J.: Identification of the murine beige gene by YAC complementation

[64789] and positional cloning. Nature Genet. 13: 303-308, 1996.

[64790] 10917.

[64791] Baysal, B. E.; Potkin, S. G.; Farr, J. E.; Higgins, M. J.; Korcz,

[64792] J.; Gollin, S. M.; James, M. R.; Evans, G. A.; Richard, C. W.,
III

[64793] : Bipolar affective disroder partially cosegregates with a
balanced

[64794] t(9;11)(p24;q23.1) chromosomal translocation in a small
pedigree. Am.

[64795] J. Med. Genet. 81: 81-91, 1998.

[64796] 10918.

[64797] Baysal, B. E.; Willett-Brozick, J. E.; Badner, J. A.; Corona,
W.;

[64798] Ferrell, R. E.; Nimgaonkar, V. L.; Detera-Wadleigh, S. D.: A
mannosyltransferase

[64799] gene at 11q23 is disrupted by a translocation breakpoint
that co-segregates

[64800] with bipolar affective disorder in a small family. Neuroge-
netics 4:

[64801] 43-53, 2002.

[64802] 10919.

[64803] Shima, D. T.; Scales, S. J.; Kreis, T. E.; Pepperkok, R.: Segregation

[64804] of COPI-rich and anterograde-cargo-rich domains in endoplasmic-reticulum-to-Golgi

[64805] transport complexes. Curr. Biol. 9: 821-824, 1999.

[64806] 10920.

[64807] Borg, J.-P.; Marchetto, S.; Le Bivic, A.; Ollendorff, V.; Jaulin-Bastard,

[64808] F.; Saito, H.; Fournier, E.; Adelaide, J.; Margolis, B.; Birnbaum,

[64809] D.: ERBIN: a basolateral PDZ protein that interacts with the mammalian

[64810] ERBB2/HER2 receptor. Nature Cell Biol. 2: 407-414, 2000.

[64811] 10921.

[64812] Dustin, M. L.; Olszowy, M. W.; Holdorf, A. D.; Li, J.; Bromley,

[64813] S.; Desai, N.; Widder, P.; Rosenberger, F.; van der Merwe, P. A.;

[64814] Allen, P. M.; Shaw, A. S.: A novel adaptor protein orchestrates receptor

[64815] patterning and cytoskeletal polarity in T-cell contacts. Cell 94:

[64816] 667-677, 1998.

[64817] 10922.

[64818] Kirsch, K. H.; Georgescu, M.-M.; Ishimaru, S.; Hanafusa,
H.: CMS:

[64819] an adapter molecule involved in cytoskeletal rearrange-
ments. Proc.

[64820] Nat. Acad. Sci. 96: 6211-6216, 1999.

[64821] 10923.

[64822] Shih, N.-Y.; Li, J.; Karpitskii, V.; Nguyen, A.; Dustin, M. L.;

[64823] Kanagawa, O.; Miner, J. H.; Shaw, A. S.: Congenital
nephrotic syndrome

[64824] in mice lacking CD2-associated protein. Science 286:
312-315, 1999.

[64825] 10924.

[64826] Kaplan, J. M.; Kim. S. H.; North, K. N.; Rennke, H.; Correia,
L.

[64827] A.; Tong, H.-Q.; Mathis, B. J.; Rodriguez-Perez, J.-C.;
Allen, P.

[64828] G.; Beggs, A. H.; Pollak, M. R.: Mutations in ACTN4, en-
coding alpha-actinin-4,

[64829] cause familial focal segmental glomerulosclerosis. Nature

Genet. 24:
[64830] 251-256, 2000.

[64831] 10925.

[64832] Schalling, M.; Hudson, T. J.; Buetow, K. H.; Housman, D. E.: Direct
[64833] detection of novel expanded trinucleotide repeats in the human genome. Nature
[64834] Genet. 4: 135-139, 1993.

[64835] 10926.

[64836] Xu, W.; Xie, Z.; Chung, D. W.; Davie, E. W.: A novel human actin-binding
[64837] protein homologue that binds to platelet glycoprotein Ib-alpha. Blood 92:
[64838] 1268-1276, 1998.

[64839] 10927.

[64840] Zhang, W.; Han, S. W.; McKeel, D. W.; Goate, A.; Wu, J. Y.: Interaction
[64841] of presenilins with the filamin family of actin-binding proteins. J.
[64842] Neurosci. 18: 914-922, 1998.

[64843] 10928.

[64844] Loughney, K.; Hill, T. R.; Florio, V. A.; Uher, L.; Rosman, G.

[64845] J.; Wolda, S. L.; Jones, B. A.; Howard, M. L.; McAllister-Lu-
cas, L.

[64846] M.; Sonnenburg, W. K.; Francis, S. H.; Corbin, J. D.; Beavo,
J. A.;

[64847] Ferguson, K.: Isolation and characterization of cDNAs en-
coding PDE5A,

[64848] a human cGMP-binding, cGMP-specific
3-prime,5-prime-cyclic nucleotide

[64849] phosphodiesterase. Gene 216: 139-147, 1998.

[64850] 10929.

[64851] Vissing, H.; Meyer, W. K.-H.; Aagaard, L.; Tommerup, N.;
Thiesen,

[64852] H.-J.: Repression of transcriptional activity by heterolo-
gous KRAB

[64853] domains present in zinc finger proteins. FEBS Lett. 369:
153-157,

[64854] 1995.

[64855] 10930.

[64856] Ziegelbauer, J.; Shan, B.; Yager, D.; Larabell, C.; Hoffmann,
B.;

[64857] Tjian, R.: Transcription factor MIZ-1 is regulated via microtubule
[64858] association. Molec. Cell 8: 339-349, 2001.

[64859] 10931.

[64860] Bolino, A.; Lonie, L. J.; Zimmer, M.; Boerkoel, C. F.; Takashima,
[64861] H.; Monaco, A. P.; Lupski, J. R.: Denaturing high-performance liquid
[64862] chromatography of the myotubularin-related 2 gene (MTMR2) in unrelated
[64863] patients with Charcot-Marie-Tooth disease suggests a low frequency
[64864] of mutation in inherited neuropathy. Neurogenetics 3: 107-109, 2001.

[64865] 10932.

[64866] Dolphin, C. T.; Beckett, D. J.; Janmohamed, A.; Cullingford, T.
[64867] E.; Smith, R. L.; Shephard, E. A.; Phillips, I. R.: The flavin-containing
[64868] monooxygenase 2 gene (FMO2) of humans, but not of other primates,
[64869] encodes a truncated, nonfunctional protein. J. Biol. Chem.

273:

[64870] 30599-30607, 1998.

[64871] 10933.

[64872] Williams, D. E.; Kelly, J.; Dutchuk, M.: Flavin-containing monooxygenase

[64873] in pulmonary microsomes of primates. In: Ingelman-Sundberg, M.; Gustafsson,

[64874] J.-A.; Orrenius, S. (eds.): Drug Metabolizing Enzymes: Genetics,

[64875] Regulation and Toxicology: Proceedings of the 8th International Symposium

[64876] on Microsomes and Drug Oxidations. Stockholm: Karolinska Institutet

[64877] 1990. P. 173.

[64878] 10934.

[64879] Johnson, J. L.; Wadman, S. K.: Molybdenum cofactor deficiency

[64880] and isolated sulfite oxidase deficiency.In: Scriver, C. R.; Beaudet,

[64881] A. L.; Sly, W. S.; Valle, D. (eds.): The Metabolic and Molecular

[64882] Bases of Inherited Disease. New York: McGraw-Hill 1995.

Pp. 2271-2283.

[64883] 10935.

[64884] Stallmeyer, B.; Drugeon, G.; Reiss, J.; Haenni, A. L.; Mendel,

[64885] R. R.: Human molybdopterin synthase gene: identification of a bicistronic

[64886] transcript with overlapping reading frames. Am. J. Hum. Genet. 64:

[64887] 698-705, 1999.

[64888] 10936.

[64889] Li-Hawkins, J.; Lund, E. G.; Turley, S. D.; Russell, D. W.: Disruption

[64890] of the oxysterol 7-alpha-hydroxylase gene in mice. J. Biol. Chem. 275:

[64891] 16536-16542, 2000.

[64892] 10937.

[64893] Setchell, K. D. R.; Schwarz, M.; O'Connell, N. C.; Lund, E. G.;

[64894] Davis, D. L.; Lathe, R.; Thompson, H. R.; Tyson, R. W.; Sokol, R.

[64895] J.; Russell, D. W.: Identification of a new inborn error in

bile

[64896] acid synthesis: mutation of the oxysterol 7-alpha-hydroxylase gene

[64897] causes severe neonatal liver disease. J. Clin. Invest. 102: 1690-1703,

[64898] 1998.

[64899] 10938.

[64900] Gunther, T.; Chen, Z.-F.; Kim, J.; Priemel, M.; Rueger, J. M.;

[64901] Amling, M.; Moseley, J. M.; Martin, T. J.; Anderson, D. J.; Karsenty,

[64902] G.: Genetic ablation of parathyroid glands reveals another source

[64903] of parathyroid hormone. Nature 406: 199-203, 2000.

[64904] 10939.

[64905] Kanemura, Y.; Hiraga, S.; Arita, N.; Ohnishi, T.; Izumoto, S.;

[64906] Mori, K.; Matsumura, H.; Yamasaki, M.; Fushiki, S.; Yoshimine, T.

[64907] : Isolation and expression analysis of a novel human homologue of

[64908] the Drosophila glial cells missing (gcm) gene. FEBS Lett.

442: 151-156,

[64909] 1999.

[64910] 10940.

[64911] Matsumoto-Taniura, N.; Pirollet, F.; Monroe, R.; Gerace, L.; Westendorf,

[64912] J. M.: Identification of novel M phase phosphoproteins by expression

[64913] cloning. Molec. Biol. Cell 7: 1455-1469, 1996.

[64914] 10941.

[64915] Doi, T.; Minami, T.; Itoh, M.; Aburatani, H.; Kawabe, Y.; Kodama,

[64916] T.; Kondo, N.; Satoh, Y.; Asayama, T.; Imanishi, T.: An al-ternative

[64917] form of nucleolysin binds to a T-cluster DNA in the si-lencer element

[64918] of platelet factor 4 gene. Biochem. Biophys. Res. Commun. 235: 625-630,

[64919] 1997.

[64920] 10942.

[64921] Kawakami, A.; Tian, Q.; Duan, X.; Streuli, M.; Schlossman, S. F.;

[64922] Anderson, P.: Identification and functional characterization of a
[64923] TIA-1-related nucleolysin. Proc. Nat. Acad. Sci. 89: 8681-8685,
[64924] 1992.

[64925] 10943.

[64926] Lemmens, I. H.; Kas, K.; Merregaert, J.; Van de Ven, W. J. M.:
[64927] Identification and molecular characterization of TM7SF2 in the FAUNA
[64928] gene cluster on human chromosome 11q13. Genomics 49: 437-442, 1998.
[64929] 10944.

[64930] Roebroek, A. J. M.; Contreras, B.; Pauli, I. G. L.; Van de Ven,
[64931] W. J. M.: cDNA cloning, genomic organization, and expression of the
[64932] human RTN2 gene, a member of a gene family encoding reticulons. Genomics 51:
[64933] 98-106, 1998.

[64934] 10945.

[64935] Kiriakidou, M.; Driscoll, D. A.; Lopez-Guisa, J. M.; Strauss, J.
[64936] F., III: Cloning and expression of primate Daxx cDNAs and mapping
[64937] of the human gene to chromosome 6p21.3 in the MHC region. DNA Cell
[64938] Biol. 16: 1289-1298, 1997.

[64939] 10946.

[64940] Yang, X.; Khosravi-Far, R.; Chang, H. Y.; Baltimore, D.: Daxx,
[64941] a novel Fas-binding protein that activates JNK and apoptosis. Cell 89:
[64942] 1067-1076, 1997.

[64943] 10947.

[64944] Errabolu, R.; Sanders, M. A.; Salisbury, J. L.: Cloning of a cDNA
[64945] encoding human centrin, an EF-hand protein of centrosomes and mitotic
[64946] spindle poles. J. Cell Sci. 107: 9-16, 1994.

[64947] 10948.

[64948] Parlati, F.; McNew, J. A.; Fukuda, R.; Miller, R.; Sollner, T.

[64949] H.; Rothman, J. E.: Topological restriction of SNARE-dependent membrane
[64950] fusion. Nature 407: 194-198, 2000.

[64951] 10949.

[64952] Scott, H. S.; Kyriakou, D. S.; Peterson, P.; Heino, M.; Tahti-nen,
[64953] M.; Krohn, K.; Chen, H.; Rossier, C.; Lalioti, M. D.; An-tonarakis,
[64954] S. E.: Characterization of a novel gene, C21orf2, on hu-man chromosome
[64955] 21q22.3 and its exclusion as the APECED gene by muta-tion analysis. Genomics 47:
[64956] 64-70, 1998.

[64957] 10950.

[64958] Dyer, M. R.; Walker, J. E.: Sequences of members of the human
[64959] gene family for the c subunit of mitochondrial ATP syn-thase. Biochem.
[64960] J. 293: 51-64, 1993.

[64961] 10951.

[64962] Beckstead, R.; Ortiz, J. A.; Sanchez, C.; Prokopenko, S. N.;

Chambon,
[64963] P.; Losson, R.; Bellen, H. J.: Bonus, a Drosophila homolog of TIF1
[64964] proteins, interacts with nuclear receptors and can inhibit beta-FTZ-F1-dependent
[64965] transcription. Molec. Cell 7: 753-765, 2001.

[64966] 10952.

[64967] Le Douarin, B.; Zechel, C.; Garnier, J.-M.; Lutz, Y.; Tora, L.;

[64968] Pierrat, B.; Heery, D.; Gronemeyer, H.; Chambon, P.; Los-
son, R.:
[64969] The N-terminal part of TIF1, a putative mediator of the ligand-dependent
[64970] activation function (AF-2) of nuclear receptors, is fused to B-raf
[64971] in the oncogenic protein T18. EMBO J. 14: 2020-2033, 1995.

[64972] 10953.

[64973] Thenot, S.; Henriquet, C.; Rochefort, H.; Cavailles, V.: Dif-
ferential
[64974] interaction of nuclear receptors with the putative human transcriptional
[64975] coactivator hTIF1. J. Biol. Chem. 272: 12062-12068,

1997.

[64976] 10954.

[64977] Feral, C.; Mattei, M. G.; Pawlak, A.; Guellaen, G.: Chromo-
somal

[64978] localization of three human poly(A)-binding protein genes
and four

[64979] related pseudogenes. Hum. Genet. 105: 347-353, 1999.

[64980] 10955.

[64981] Houng, A. K.; Maggini, L.; Clement, C. Y.; Reed, G. L.:
Identification

[64982] and structure of activated-platelet protein-1, a protein
with RNA-binding

[64983] domain motifs that is expressed by activated platelets.
Europ. J.

[64984] Biochem. 243: 209-218, 1997.

[64985] 10956.

[64986] Yang, H.; Duckett, C. S.; Lindsten, T.: iPABP, an inducible
poly(A)-binding

[64987] protein detected in activated human T cells. Molec. Cell.
Biol. 15:

[64988] 6770-6776, 1995.

[64989] 10957.

[64990] Liu, T.; DeCostanzo, A. J.; Liu, X.; Wang, H.; Hallagan, S.; Moon,

[64991] R. T.; Malbon, C. C.: G protein signaling from activated rat frizzled-1

[64992] to the beta-catenin-Lef-Tcf pathway. Science 292: 1718-1722, 2001.

[64993] 10958.

[64994] Sagara, N.; Toda, G.; Hirai, M.; Terada, M.; Katoh, M.: Molecular

[64995] cloning, differential expression, and chromosomal local-ization of

[64996] human frizzled-1, frizzled-2, and frizzled-7. Biochem. Biophys. Res.

[64997] Commun. 252: 117-122, 1998.

[64998] 10959.

[64999] Tokuhara, M.; Hirai, M.; Atomi, Y.; Terada, M.; Katoh, M.: Molecular

[65000] cloning of human frizzled-6. Biochem. Biophys. Res. Commun. 243:

[65001] 622-627, 1998.

[65002] 10960.

[65003] Tanaka, S.; Akiyoshi, T.; Mori, M.; Wands, J. R.; Sugimachi, K.

[65004] : A novel frizzled gene identified in human esophageal carcinoma mediates

[65005] APC/beta-catenin signals. Proc. Nat. Acad. Sci. 95: 10164-10169,

[65006] 1998.

[65007] 10961.

[65008] Winklbauer, R.; Medina, A.; Swain, R. K.; Steinbeisser, H.: Frizzled-7

[65009] signalling controls tissue separation during Xenopus gastrulation. Nature 413:

[65010] 856-860, 2001.

[65011] 10962.

[65012] Zhao, Y.; Loyer, P.; Li, H.; Valentine, V.; Kidd, V.; Kraft, A.

[65013] S.: Cloning and chromosomal location of a novel member of the myotonic

[65014] dystrophy family of protein kinases. J. Biol. Chem. 272: 10013-10020,

[65015] 1997.

[65016] 10963.

[65017] The European Consortium on MEN1: Construction of a 1.2-Mb sequence-ready

[65018] contig of chromosome 11q13 encompassing the multiple endocrine neoplasia

[65019] type 1 (MEN1) gene. Genomics 44: 94-100, 1997.

[65020] 10964.

[65021] Kohl, S.; Baumann, B.; Broghammer, M.; Jagle, H.; Sieving, P.;

[65022] Kellner, U.; Spegal, R.; Anastasi, M.; Zrenner, E.; Sharpe, L. T.;

[65023] Wissinger, B.: Mutations in the CNGB3 gene encoding the beta-subunit

[65024] of the cone photoreceptor cGMP-gated channel are responsible for achromatopsia

[65025] (ACHM3) linked to chromosome 8q21. Hum. Molec. Genet. 9: 2107-2116,

[65026] 2000.

[65027] 10965.

[65028] Amiel, J.; Bougeard, G.; Francannet, C.; Raclin, V.; Munnich, A.;

[65029] Lyonnet, S.; Frebourg, T.: TP63 gene mutation in ADULT syndrome. Europ.

[65030] J. Hum. Genet. 9: 642-645, 2001.

[65031] 10966.

[65032] Sperandeo, M. P.; Bassi, M. T.; Riboni, M.; Parenti, G.; Buoninconti,

[65033] A.; Manzoni, M.; Incerti, B.; Larocca, M. R.; Di Rocco, M.; Strisciuglio,

[65034] P.; Dianzani, I.; Parini, R.; Candito, M.; Endo, F.; Ballabio, A.;

[65035] Andria, G.; Sebastio, G.; Borsani, G.: Structure of the SLC7A7 gene

[65036] and mutational analysis of patients affected by lysinuric protein

[65037] intolerance. Am. J. Hum. Genet. 66: 92-99, 2000.

[65038] 10967.

[65039] Messina, A.; Oliva, M.; Rosato, C.; Huizing, M.; Ruitenbeek, W.;

[65040] van den Heuvel, L. P.; Forte, M.; Rocchi, M.; De Pinto, V.: Mapping

[65041] of the human voltage-dependent action channel isoforms 1 and 2 reconsidered. Bio

[65042] chem. Biophys. Res. Commun. 255: 707-710, 1999.

[65043] 10968.

[65044] Parnet, P.; Garka, K. E.; Bonnert, T. P.; Dower, S. K.; Sims, J.

[65045] E.: IL-1Rrp is a novel receptor-like molecule similar to the type

[65046] I interleukin-1 receptor and its homologues T1/ST2 and IL-1R AcP. J.

[65047] Biol. Chem. 271: 3967-3970, 1996.

[65048] 10969.

[65049] Torigoe, K.; Ushio, S.; Okura, T.; Kobayashi, S.; Taniai, M.; Kunikata,

[65050] T.; Murakami, T.; Sanou, O.; Kojima, H.; Fujii, M.; Ohta, T.; Ikeda,

[65051] M.; Ikegami, H.; Kurimoto, M.: Purification and characterization

[65052] of the human interleukin-18 receptor. J. Biol. Chem. 272: 25737-25742,

[65053] 1997.

[65054] 10970.

[65055] Torrents, D.; Estevez, R.; Pineda, M.; Fernandez, E.;

Lloberas,
[65056] J.; Shi, Y.-B.; Zorzano, A.; Palacin, M.: Identification and characterization
[65057] of a membrane protein ((+)L amino acid transporter-1) that associates
[65058] with 4F2hc to encode the amino acid transport activity y(+)L: a candidate
[65059] gene for lysinuric protein intolerance. J. Biol. Chem. 273: 32437-32445,
[65060] 1998.

[65061] 10971.

[65062] Akiba, H.; Atsuta, M.; Yagita, H.; Okumura, K.: Identifica-
tion
[65063] of rat OX40 ligand by molecular cloning. Biochem. Bio-
phys. Res. Commun. 251:
[65064] 131-136, 1998.

[65065] 10972.

[65066] Baum, P. R.; Gayle, R. B., III; Ramsdell, F.; Srinivasan, S.; Sorensen,
[65067] R. A.; Watson, M. L.; Seldin, M. F.; Baker, E.; Sutherland, G. R.;
[65068] Clifford, K. N.; Alderson, M. R.; Goodwin, R. G.; Fanslow,

W. C.:

[65069] Molecular characterization of murine and human OX40/OX40 ligand systems:

[65070] identification of a human OX40 ligand as the HTLV-1-regulated protein

[65071] gp34. EMBO J. 13: 3992-4001, 1994.

[65072] 10973.

[65073] Godfrey, W. R.; Fagnoni, F. F.; Harara, M. A.; Buck, D.; Engleman,

[65074] E. G.: Identification of a human OX-40 ligand, a costimulator of

[65075] CD4+ T cells with homology to tumor necrosis factor. J. Exp. Med. 180:

[65076] 757-762, 1994.

[65077] 10974.

[65078] Malmstrom, V.; Shipton, D.; Singh, B.; Al-Shamkhani, A.; Puklavec,

[65079] M. J.; Barclay, A. N.; Powrie, F.: CD134L expression on dendritic

[65080] cells in the mesenteric lymph nodes drives colitis in T cell-restored

[65081] SCID mice. J. Immun. 166: 6972-6981, 2001.

[65082] 10975.

[65083] Miura, S.; Ohtani, K.; Numata, N.; Niki, M.; Ohbo, K.; Ina, Y.;

[65084] Gojobori, T.; Tanaka, Y.; Tozawa, H.; Nakamura, M.; Suga-

mura, K.:

[65085] Molecular cloning and characterization of a novel glyco-

protein, gp34,

[65086] that is specifically induced by the human T-cell leukemia virus type

[65087] I transactivator p40-tax. Molec. Cell. Biol. 11: 1313-1325, 1991.

[65088] 10976.

[65089] Hilton, D. J.; Richardson, R. T.; Alexander, W. S.; Viney, E. M.;

[65090] Willson, T. A.; Sprigg, N. S.; Starr, R.; Nicholson, S. E.; Metcalf,

[65091] D.; Nicola, N. A.: Twenty proteins containing a C-terminal SOCS box

[65092] form five structural classes. Proc. Nat. Acad. Sci. 95: 114-119,

[65093] 1998.

[65094] 10977.

[65095] Cretney, E.; Takeda, K.; Yagita, H.; Glaccum, M.; Peschon, J. J.;
[65096] Smyth, M. J.: Increased susceptibility to tumor initiation and metastasis
[65097] in TNF-related apoptosis-inducing ligand-deficient mice. J. Immun. 168:
[65098] 1356-1361, 2002.

[65099] 10978.

[65100] Degli-Esposti, M. A.; Dougall, W. C.; Smolak, P. J.; Waugh, J.
[65101] Y.; Smith, C. A.; Goodwin, R. G.: The novel receptor TRAIL-R4 induces
[65102] NF-kappa-B and protects against TRAIL-mediated apoptosis, yet retains
[65103] an incomplete death domain. Immunity 7: 813-820, 1997.

[65104] 10979.

[65105] Shibui, A.; Tsunoda, T.; Seki, N.; Suzuki, Y.; Sugane, K.; Sugano,
[65106] S.: Isolation and chromosomal mapping of a novel human gene showing
[65107] homology to Na+/PO4 cotransporter. J. Hum. Genet. 44: 190-192, 1999.

[65108] 10980.

[65109] Bardoni, B.; Giglio, S.; Schenck, A.; Rocchi, M.; Mandel, J. L.

[65110] : Assignment of NUFIP1 (nuclear FMRP interacting protein 1) gene to

[65111] chromosome 13q14 and assignment of a pseudogene to chromosome 6q12. Cytogenet.

[65112] Cell Genet. 89: 11-13, 2000.

[65113] 10981.

[65114] Bardoni, B.; Schenck, A.; Mandel, J. L.: A novel RNA-binding nuclear

[65115] protein that interacts with the fragile X mental retardation (FMR1)

[65116] protein. Hum. Molec. Genet. 8: 2557-2566, 1999.

[65117] 10982.

[65118] Blagitko, N.; Schulz, U.; Schinzel, A. A.; Ropers, H.-H.; Kalscheuer,

[65119] V. M.: Gamma-2-COP, a novel imprinted gene on chro-mosome 7q32, defines

[65120] a new imprinting cluster in the human genome. Hum. Molec. Genet. 8:

[65121] 2387-2396, 1999.

[65122] 10983.

[65123] Putilina, T.; Jaworski, C.; Gentleman, S.; McDonald, B.; Kadiri,

[65124] M.; Wong, P.: Analysis of a human cDNA containing a tis-sue-specific

[65125] alternatively spliced LIM domain. Biochem. Biophys. Res. Commun. 252:

[65126] 433-439, 1998.

[65127] 10984.

[65128] Davenport, J. W.; Fernandes, E. R.; Harris, L. D.; Neale, G. A.

[65129] M.; Goorha, R.: The mouse mitotic checkpoint gene Bub1b, a novel

[65130] Bub1 family member, is expressed in a cell cycle-dependent manner. Genomics 55:

[65131] 113-117, 1999.

[65132] 10985.

[65133] Taylor, S. S.; Ha, E.; McKeon, F.: The human homologue of Bub3

[65134] is required for kinetochore localization of Bub1 and a

Mad3/Bub1-related
[65135] protein kinase. J. Cell Biol. 142: 1-11, 1998.

[65136] 10986.

[65137] Bonne, S.; van Hengel, J; van Roy, F.: Assignment of the plakophilin-2
[65138] gene (PKP2) and a plakophilin-2 pseudogene (PKP2P1) to human chromosome
[65139] bands 12p11 and 12p13, respectively, by in situ hybridization. Cytogenet.
[65140] Cell Genet. 88: 286-287, 2000.

[65141] 10987.

[65142] Mertens, C.; Kuhn, C.; Franke, W. W.: Plakophilins 2a and 2b:
[65143] constitutive proteins of dual location in the karyoplasm and the desmosomal
[65144] plaque. J. Cell Biol. 135: 1009-1025, 1996.

[65145] 10988.

[65146] Schmidt, A; Langbein, L.; Pratzel, S.; Rode, M.; Rackwitz, H.-R.;
[65147] Franke, W. W.: Plakophilin 3--a novel cell-type-specific desmosomal

[65148] plaque protein. Differentiation 64: 291-306, 1999.

[65149] 10989.

[65150] Bergstein, I.; Eisenberg, L. M.; Bhalerao, J.; Jenkins, N. A.;

[65151] Copeland, N. G.; Osborne, M. P.; Bowcock, A. M.; Brown,
A. M. C.:

[65152] Isolation of two novel WNT genes, WNT14 and WNT15,
one of which (WNT15)

[65153] is closely linked to WNT3 on human chromosome 17q21.
Genomics 46:

[65154] 450-458, 1997.

[65155] 10990.

[65156] Hartmann, C.; Tabin, C. J.: Wnt-14 plays a pivotal role in
inducing

[65157] synovial joint formation in the developing appendicular
skeleton. Cell 104:

[65158] 341-351, 2001.

[65159] 10991.

[65160] Saitoh, T.; Hirai, M.; Katoh, M.: Molecular cloning and
characterization

[65161] of WNT3A and WNT14 clustered in human chromosome
1q42 region. Biochem.

[65162] Biophys. Res. Commun. 284: 1168-1175, 2001.

[65163] 10992.

[65164] Habuchi, T.; Luscombe, M.; Elder, P. A.; Knowles, M. A.: Structure

[65165] and methylation-based silencing of a gene (DBCCR1) within a candidate

[65166] bladder cancer tumor suppressor region at 9q32-q33. Genomics 48:

[65167] 277-288, 1998.

[65168] 10993.

[65169] Habuchi, T.; Yoshida, O.; Knowles, M. A.: A novel candidate tumour

[65170] suppressor locus at 9q32-33 in bladder cancer: localization of the

[65171] candidate region within a single 840 kb YAC. Hum. Molec. Genet. 6:

[65172] 913-919, 1997.

[65173] 10994.

[65174] Nishiyama, H.; Takahashi, T.; Kakehi, Y.; Habuchi, T.; Knowles,

[65175] M. A.: Homozygous deletion at the 9q32-33 candidate tu-

mor suppressor
[65176] locus in primary human bladder cancer. Genes Chromo-
somes Cancer 26:
[65177] 171-175, 1999.

[65178] 10995.

[65179] Ayala-Madrigal, M. L.; Doerr, S.; Ramirez-Duenas, M. L.;
Hansmann,
[65180] I.: Assignment of KPNA4 and KPNB1 encoding karyopherin
alpha 4 and
[65181] beta 1 to human chromosome bands 11q22 and 17q21
respectively, by
[65182] in situ hybridization. Cytogenet. Cell Genet. 89: 258-259,
2000.
[65183] 10996.

[65184] Bayliss, R.; Littlewood, T.; Stewart, M.: Structural basis for

[65185] the interaction between FxFG nucleoporin repeats and im-
portin-beta
[65186] in nuclear trafficking. Cell 102: 99-108, 2000.

[65187] 10997.

[65188] Chi, N. C.; Adam, E. J. H.; Adam, S. A.: Sequence and char-
acterization

[65189] of cytoplasmic nuclear protein import factor p97. J. Cell Biol. 130:

[65190] 265-274, 1995.

[65191] 10998.

[65192] Gorlich, D.; Kostka, S.; Kraft, R.; Dingwall, C.; Laskey, R. A.;

[65193] Hartmann, E.; Prehn, S.: Two different subunits of im-portin cooperate

[65194] to recognize nuclear localization signals and bind them to the nuclear

[65195] envelope. Curr. Biol. 5: 383-392, 1995.

[65196] 10999.

[65197] Kutay, U.; Izaurralde, E.; Bischoff, F. R.; Mattaj, I. W.; Gor-lich,

[65198] D.: Dominant-negative mutants of importin-beta block multiple pathways

[65199] of import and export through the nuclear pore complex. EMBO J. 16:

[65200] 1153-1163, 1997.

[65201] 11000.

[65202] Matsuda, Y.; Hamatani, K.; Itoh, M.; Takahashi, E.; Araki,

R.;

[65203] Abe, M.: Localization of the importin-beta gene to mouse chromosome

[65204] 11D and rat chromosome 10q32.1. Genomics 36: 213-215, 1996.

[65205] 11001.

[65206] Thornton, C.; Snowden, M. A.; Carling, D.: Identification of a

[65207] novel AMP-activated protein kinase beta subunit isoform that is highly

[65208] expressed in skeletal muscle. J. Biol. Chem. 273: 12443-12450, 1998.

[65209] 11002.

[65210] Woods, A.; Cheung, P. C. F.; Smith, F. C.; Davison, M. D.; Scott,

[65211] J.; Beri, R. K.; Carling, D.: Characterization of AMP-activated protein

[65212] kinase beta and gamma subunits: assembly of the het-erotrimeric complex

[65213] in vitro. J. Biol. Chem. 271: 10282-10290, 1996.

[65214] 11003.

[65215] Lee, S.-J.: Expression of growth/differentiation factor 1 in

the
[65216] nervous system: conservation of a bicistronic structure. Proc. Nat.
[65217] Acad. Sci. 88: 4250-4254, 1991.

[65218] 11004.

[65219] Yamagoe, S.; Akasaka, T.; Uchida, T.; Hachiya, T.; Okabe, T.; Yamakawa,
[65220] Y.; Arai, T.; Mizuno, S.; Suzuki, K.: Expression of a neu-trophil
[65221] chemotactic protein LECT2 in human hepatocytes revealed by immunochemical
[65222] studies using polyclonal and monoclonal antibodies to a recombinant
[65223] LECT2. Biochem. Biophys. Res. Commun. 237: 116-120, 1997.
[65224] 11005.

[65225] Yamagoe, S.; Kameoka, Y.; Hashimoto, K.; Mizuno, S.; Suzuki, K.
[65226] : Molecular cloning, structural characterization, and chro-mosomal
[65227] mapping of the human LECT2 gene. Genomics 48: 324-329, 1998.

[65228] 11006.

[65229] Yamagoe, S.; Mizuno, S.; Suzuki, K.: Molecular cloning of human

[65230] and bovine LECT2 having a neutrophil chemotactic activity and its

[65231] specific expression in the liver. Biochim. Biophys. Acta 1396: 105-113,

[65232] 1998.

[65233] 11007.

[65234] Yamagoe, S.; Watanabe, T.; Mizuno, S.; Suzuki, K.: The mouse Lect2

[65235] gene: cloning of cDNA and genomic DNA, structural characterization

[65236] and chromosomal localization. Gene 216: 171-178, 1998.

[65237] 11008.

[65238] Yamagoe, S.; Yamakawa, Y.; Matsuo, Y.; Minowada, J.; Mizuno, S.;

[65239] Suzuki, K.: Purification and primary amino acid sequence of a novel

[65240] neutrophil chemotactic factor LECT2. Immun. Lett. 52: 9-13, 1996.

[65241] 11009.

[65242] Kullak-Ublick, G.-A.; Beuers, U.; Meier, P. J.; Domdey, H.; Paumgartner,

[65243] G.: Assignment of the human organic anion transporting polypeptide

[65244] (OATP) gene to chromosome 12p12 by fluorescence in situ hybridization. J.

[65245] Hepatol. 25: 985-987, 1996.

[65246] 11010.

[65247] Kullak-Ublick, G. A.; Hagenbuch, B.; Stieger, B.; Schtein-
gart,
[65248] C. D.; Hofmann, A. F.; Wolkoff, A. W.; Meier, P. J.: Molecu-
lar and
[65249] functional characterization of an organic anion transport-
ing polypeptide
[65250] cloned from human liver. :Gastroenterology 109:
1274-1282, 1995.
[65251] 11011.

[65252] El-Husseini, A. E.-D.; Schnell, E.; Chetkovich, D. M.; Nicoll,

[65253] R. A.; Bredt, D. S.: PSD-95 involvement in maturation of
excitatory
[65254] synapses. Science 290: 1364-1368, 2000.

[65255] 11012.

[65256] El-Husseini, A. E.-D.; Schnell, E.; Dakoji, S.; Sweeney, N.;
Zhou,
[65257] Q.; Prange, O.; Gauthier-Campbell, C.; Aguilera-Moreno,
A.; Nicoll,
[65258] R. A.; Bredt, D. S.: Synaptic strength regulated by palmi-

tate cycling
[65259] on PSD-95. Cell 108: 849-863, 2002.

[65260] 11013.

[65261] Kim, E.; Cho, K.-O.; Rothschild, A.; Sheng, M.: Heteromul-
timerization
[65262] and NMDA receptor-clustering activity of Chapsyn-110, a
member of
[65263] the PSD-95 family of proteins. Neuron 17: 103-113,
1996.
[65264] 11014.

[65265] Kim, E.; Niethammer, M.; Rothschild, A.; Jan, Y. N.; Sheng,
M.
[65266] : Clustering of Shaker-type K+ channels by interaction
with a family
[65267] of membrane-associated guanylate kinases. Nature 378:
85-88, 1995.
[65268] 11015.

[65269] Kistner, U.; Wenzel, B. M.; Veh, R. W.; Cases-Langhoff, C.;
Garner,
[65270] A. M.; Appeltauer, U.; Voss, B.; Gundelfinger, E. D.; Gar-
ner, C. C.
[65271] : SAP90, a rat presynaptic protein related to the product of

the Drosophila
[65272] tumor suppressor gene, dLg-A. J. Biol. Chem. 268: 4580-4583, 1993.
[65273] 11016.

[65274] Migaud, M.; Charlesworth, P.; Dempster, M.; Webster, L. C.; Watabe,
[65275] A. M.; Makhinson, M.; He, Y.; Ramsay, M. F.; Morris, R. G. M.; Morrison,
[65276] J. H.; O'Dell, T. J.; Grant, S. G. N.: Enhanced long-term potentiation
[65277] and impaired learning in mice with mutant postsynaptic density-95
[65278] protein. Nature 396: 433-439, 1998.

[65279] 11017.

[65280] Sattler, R.; Xiong, Z.; Lu, W.-Y.; Hafner, M.; MacDonald, J. F.;
[65281] Tymianski, M.: Specific coupling of NMDA receptor activation to nitric
[65282] oxide neurotoxicity by PSD-95 protein. Science 284: 1845-1848, 1999.
[65283] 11018.

[65284] Stathakis, D. G.; Hoover, K. B.; You, Z.; Bryant, P. J.: Hu-

man

[65285] postsynaptic density-95 (PSD95): location of the gene (DLG4) and possible

[65286] function in nonneural as well as in neural tissues. Genomics 44:

[65287] 71-82, 1997.

[65288] 11019.

[65289] Strippoli, P.; Petrini, M.; Lenzi, L.; Carinci, P.; Zannotti, M.

[65290] : The murine DSCR1-like (Down syndrome candidate region 1) gene family:

[65291] conserved synteny with the human orthologous genes. Gene 257: 223-232,

[65292] 2000.

[65293] 11020.

[65294] Yang, J.; Rothermel, B.; Vega, R. B.; Frey, N.; McKinsey, T. A.;

[65295] Olson, E. N.; Bassel-Duby, R.; Williams, R. S.: Independent signals

[65296] control expression of the calcineurin inhibitory proteins MCIP1 and

[65297] MCIP2 in striated muscles. Circ. Res. 87: 61e-68e, 2000.

[65298] 11021.

[65299] Zheng, B.; Larkin, D. W.; Albrecht, U.; Sun, Z. S.; Sage, M.; Eichele,

[65300] G.; Lee, C. C.; Bradley, A.: The mPer2 gene encodes a functional

[65301] component of the mammalian circadian clock. Nature 400: 169-173,

[65302] 1999.

[65303] 11022.

[65304] Lund, E. G.; Guileyardo, J. M.; Russell, D. W.: cDNA cloning of

[65305] cholesterol 24-hydroxylase, a mediator of cholesterol homeostasis

[65306] in the brain. Proc. Nat. Acad. Sci. 96: 7238-7243, 1999.

[65307] 11023.

[65308] Endo, S.; Suzuki, M.; Sumi, M.; Nairn, A. C.; Morita, R.; Ya-makawa,

[65309] K.; Greengard, P.; Ito, M.: Molecular identification of hu-man G-substrate,

[65310] a possible downstream component of the cGMP-de-pendent protein kinase

[65311] cascade in cerebellar Purkinje cells. Proc. Nat. Acad. Sci. 96:
[65312] 2467-2472, 1999.

[65313] 11024.

[65314] Nakamura, H.; Sudo, T.; Tsuiki, H.; Miyake, H.; Morisaki, T.; Sasaki,
[65315] J.; Masuko, N.; Kochi, M.; Ushio, Y.; Saya, H.: Identification of
[65316] a novel human homolog of the Drosophila dlg, P-dlg, specifically expressed
[65317] in the gland tissues and interacting with p55. FEBS Lett. 433: 63-67,
[65318] 1998.

[65319] 11025.

[65320] Funderburgh, J. L.; Perchellet, A. L.; Swiergiel, J.; Conrad, G.
[65321] W.; Justice, M. J.: Keratocan (Kera), a corneal keratan sulfate proteoglycan,
[65322] maps to the distal end of mouse chromosome 10. Genomics 52: 110-111,
[65323] 1998.

[65324] 11026.

[65325] Liu, C.-Y.; Shiraishi, A.; Kao, C. W.-C.; Converse, R. L.; Funderburgh,

[65326] J. L.; Corpuz, L. M.; Conrad, G. W.; Kao, W. W.-Y.: The cloning of

[65327] mouse keratocan cDNA and genomic DNA and the characterization of its

[65328] expression during eye development. J. Biol. Chem. 273: 22584-22588,

[65329] 1998.

[65330] 11027.

[65331] Tasheva, E. S.; Funderburgh, J. L.; Funderburgh, M. L.; Corpuz,

[65332] L. M.; Conrad, G. W.: Structure and sequence of the gene encoding

[65333] human keratocan. DNA Seq. 10: 67-74, 1999.

[65334] 11028.

[65335] Tasheva, E. S.; Pettenati, M.; Von Kap-Her, C.; Conrad, G. W.:

[65336] Assignment of keratocan gene (KERA) to human chromosome band 12q22

[65337] by in situ hybridization. Cytogenet. Cell Genet. 88: 244-245, 2000.

[65338] 11029.

[65339] Du, Y.; Weed, S. A.; Xiong, W.-C.; Marshall, T. D.; Parsons, J.

[65340] T.: Identification of a novel cortactin SH3 domain-binding protein

[65341] and its localization to growth cones of cultured neurons. Molec.

[65342] Cell. Biol. 18: 5838-5851, 1998.

[65343] 11030.

[65344] Bruick, R. K.: Expression of the gene encoding the proapoptotic

[65345] Nip3 protein is induced by hypoxia. Proc. Nat. Acad. Sci. 97: 9082-9087,

[65346] 2000.

[65347] 11031.

[65348] Soyombo, A. A.; Hofmann, S. L.: Molecular cloning and expression

[65349] of palmitoyl-protein thioesterase 2 (PPT2), a homolog of lysosomal

[65350] palmitoyl-protein thioesterase with a distinct substrate specificity. J.

[65351] Biol. Chem. 272: 27456-27463, 1997.

[65352] 11032.

[65353] Sarma, V.; Wolf, F. W.; Marks, R. M.; Shows, T. B.; Dixit, V. M.

[65354] : Cloning of a novel tumor necrosis factor-alpha-inducible primary

[65355] response gene that is differentially expressed in development and

[65356] capillary tube-like formation in vitro. J. Immun. 148: 3302-3312,

[65357] 1992.

[65358] 11033.

[65359] Richard, M.; Drouin, R.; Beaulieu, A. D.: ABC50, a novel human

[65360] ATP-binding cassette protein found in tumor necrosis factor-alpha-stimulated

[65361] synoviocytes. Genomics 53: 137-145, 1998.

[65362] 11034.

[65363] Richter-Cook, N. J.; Dever, T. E.; Hensold, J. O.; Merrick,

W.

[65364] C.: Purification and characterization of a new eukaryotic protein

[65365] translation factor: eukaryotic initiation factor 4H. J. Biol. Chem. 273:

[65366] 7579-7587, 1998.

[65367] 11035.

[65368] Hoogenraad, C. C.; Eussen, B. H. J.; Langeveld, A.; van Haperen,

[65369] R.; Winterberg, S.; Wouters, C. H.; Grosveld, F.; De Zeeuw, C. I.;

[65370] Galjart, N.: The murine CYLN2 gene: genomic organization, chromosome

[65371] localization, and comparison to the human gene that is located within

[65372] the 7q11.23 Williams syndrome critical region. Genomics 53: 348-358,

[65373] 1998.

[65374] 11036.

[65375] Araujo, H.; Danziger, N.; Cordier, J.; Glowinski, J.; Chneiweiss,

[65376] H.: Characterization of PEA-15, a major substrate for pro-

tein kinase
[65377] C in astrocytes. J. Biol. Chem. 268: 5911-5920, 1993.

[65378] 11037.

[65379] Bera, T. K.; Guzman, R. C.; Miyamoto, S.; Panda, D. K.; Sasaki,
[65380] M.; Hanyu, K.; Enami, J.; Nandi, S.: Identification of a mammary
[65381] transforming gene (MAT1) associated with mouse mammary carcinogenesis. Proc.
[65382] Nat. Acad. Sci. 91: 9789-9793, 1994.

[65383] 11038.

[65384] Condorelli, G.; Vigliotta, G.; Iavarone, C.; Caruso, M.; Tocchetti,
[65385] C. G.; Andreozzi, F.; Cafieri, A.; Tecce, M. F.; Formisano, P.; Beguinot,
[65386] L.; Beguinot, F.: PED/PEA-15 gene controls glucose transport and
[65387] is overexpressed in type 2 diabetes mellitus. EMBO J. 17: 3858-3866,
[65388] 1998.

[65389] 11039.

[65390] Danziger, N.; Yokoyama, M.; Jay, T.; Cordier, J.; Glowinski, J.;
[65391] Chneiweiss, H.: Cellular expression, developmental regulation, and
[65392] phylogenic conservation of PEA-15, the astrocytic major phosphoprotein
[65393] and protein kinase C substrate. J. Neurochem. 64: 1016-1025, 1995.
[65394] 11040.

[65395] Estelles, A.; Yokoyama, M.; Nothias, F.; Vincent, J.-D.; Glowinski,
[65396] J.; Vernier, P.; Chneiweiss, H.: The major astrocytic phosphoprotein
[65397] PEA-15 is encoded by two mRNAs conserved on their full length in mouse
[65398] and human. J. Biol. Chem. 271: 14800-14806, 1996.

[65399] 11041.

[65400] Hwang, S.; Kuo, W.-L.; Cochran, J. F.; Guzman, R. C.; Tsukamoto,
[65401] T.; Bandyopadhyay, G.; Myambo, K.; Collins, C. C.: Assignment of
[65402] HMAT1, the human homolog of the murine mammary

transforming gene (MAT1)
[65403] associated with tumorigenesis, to 1q21.1, a region frequently gained
[65404] in human breast cancers. Genomics 42: 540-542, 1997.

[65405] 11042.

[65406] Wolford, J. K.; Bogardus, C.; Ossowski, V.; Prochazka, M.: Molecular
[65407] characterization of the human PEA15 gene on 1q21-q22 and association
[65408] with type 2 diabetes mellitus in Pima Indians. Gene 241: 143-148,
[65409] 2000.

[65410] 11043.

[65411] Martin, F.; Malergue, F.; Pitari, G.; Philippe, J. M.; Philips,

[65412] S.; Chabret, C.; Granjeaud, S.; Mattei, M. G.; Mungall, A. J.; Naquet,
[65413] P.; Galland, F.: Vanin genes are clustered (human 6q22-24 and mouse
[65414] 10A2B1) and encode isoforms of pantetheinase ectoenzymes. Immunogenetics 53:
[65415] 296-306, 2001.

[65416] 11044.

[65417] Hendrich, B.; Bird, A.: Identification and characterization of

[65418] a family of mammalian methyl-CpG binding proteins. Molec. Cell. Biol. 18:

[65419] 6538-6547, 1998.

[65420] 11045.

[65421] Rasooly, R. S.: Personal Communication. Baltimore, Md. 2/23/1999.

[65422] 11046.

[65423] Zhang, Y.; Ng, H.-H.; Erdjument-Bromage, H; Tempst, P.; Bird, A.;

[65424] Reinberg, D.: Analysis of the NuRD subunits reveals a histone deacetylase

[65425] core complex and a connection with DNA methylation. Genes Dev. 13:

[65426] 1924-1935, 1999.

[65427] 11047.

[65428] Bellacosa, A.; Cicchillitti, L.; Schepis, F.; Riccio, A.; Yeung,

[65429] A. T.; Matsumoto, Y.; Golemis, E. A.; Genuardi, M.; Neri, G.: MED1,

[65430] a novel human methyl-CpG-binding endonuclease, interacts with DNA

[65431] mismatch repair protein MLH1. Proc. Nat. Acad. Sci. 96: 3969-3974,

[65432] 1999.

[65433] 11048.

[65434] Boland, C. R.; Thibodeau, S. N.; Hamilton, S. R.; Sidransky, D.;

[65435] Eshleman, J. R.; Burt, R. W.; Meltzer, S. J.; Rodriguez-Bigas, M.

[65436] A.; Fodde, R.; Ranzani, G. N.; Srivastava, S.: A National Cancer

[65437] Institute workshop on microsatellite instability for cancer detection

[65438] and familial predisposition: development of international criteria

[65439] for the determination of microsatellite instability in colorectal

[65440] cancer. Cancer Res. 58: 5248-5257, 1998.

[65441] 11049.

[65442] Hendrich, B.; Hardeland, U.; Ng, H.-H.; Jiricny, J.; Bird, A.:

[65443] The thymine glycosylase MBD4 can bind to the product of deamination
[65444] at methylated CpG sites. Nature 401: 301-304, 1999.

[65445] 11050.

[65446] Millar, C. B.; Guy, J.; Sansom, O. J.; Selfridge, J.; Mac-Dougall,
[65447] E.; Hendrich, B.; Keightley, P. D.; Bishop, S. M.; Clarke, A. R.;
[65448] Bird, A.: Enhanced CpG mutability and tumorigenesis in MBD4-deficient
[65449] mice. Science 297: 403-405, 2002.

[65450] 11051.

[65451] Riccio, A.; Aaltonen, L. A.; Godwin, A. K.; Loukola, A.; Percesepe,
[65452] A.; Salovaara, R.; Masciullo, V.; Genuardi M.; Paravatou-Pet-sotas,
[65453] M.; Bassi, D. E.; Ruggeri, B. A.; Klein-Szanto, A. J. P.; Testa, J.
[65454] R.; Neri, G.; Bellacosa, A.: The DNA repair gene MBD4 (MED1) is mutated
[65455] in human carcinomas with microsatellite instability. (Letter) Nature

[65456] Genet. 23: 266-268, 1999.

[65457] 11052.

[65458] Duncan, L. M.; Deeds, J.; Hunter, J.; Shao, J.; Holmgren, L. M.;

[65459] Woolf, E. A.; Tepper, R. I.; Shyjan, A. W.: Down-regulation of the

[65460] novel gene melastatin correlates with potential for melanoma metastasis. Cancer

[65461] Res. 58: 1515-1520, 1998.

[65462] 11053.

[65463] Fang, D.; Setaluri, V.: Expression and up-regulation of alternatively

[65464] spliced transcripts of melastatin, a melanoma metastasis-related gene,

[65465] in human melanoma cells. Biochem. Biophys. Res. Commun. 279: 53-61,

[65466] 2000.

[65467] 11054.

[65468] Hunter, J. J.; Shao, J.; Smutko, J. S.; Dussault, B. J.; Nagle,

[65469] D. L.; Woolf, E. A.; Holmgren, L. M.; Moore, K. J.; Shyjan, A. W.

[65470] : Chromosomal localization and genomic characterization of the mouse
[65471] melastatin gene (Mlsn1). Genomics 54: 116-123, 1998.

[65472] 11055.

[65473] Xu, X. Z.; Moebius, F.; Gill, D. L.; Montell, C.: Regulation of
[65474] melastatin, a TRP-related protein, through interaction with a cytoplasmic
[65475] isoform. Proc. Nat. Acad. Sci. 98: 10692-10697, 2001.

[65476] 11056.

[65477] Anaya, P.; Evans, S. C.; Dai, C.; Lozano, G.; May, G. S.: Isolation
[65478] of the Aspergillus nidulans sudD gene and its human homologue. Gene 211:
[65479] 323-329, 1998.

[65480] 11057.

[65481] Kalitsis, P.; Earle, E.; Fowler, K. J.; Choo, K. H. A.: Bub3 gene
[65482] disruption in mice reveals essential mitotic spindle checkpoint function
[65483] during early embryogenesis. Genes Dev. 14: 2277-2282, 2000.

[65484] 11058.

[65485] Ardley, H. C.; Moynihan, T. P.; Thompson, J.; Leek, J. P.; Markham,

[65486] A. F.; Robinson, P. A.: Rapid isolation of genomic clones for individual

[65487] members of human multigene families: identification and localisation

[65488] of UBE2L4, a novel member of a ubiquitin conjugating enzyme dispersed

[65489] gene family. Cytogenet. Cell Genet. 79: 188-192, 1997.

[65490] 11059.

[65491] Moynihan, T. P.; Cole, C. G.; Dunham, I.; O'Neil, L.; Markham,

[65492] A. F.; Robinson, P. A.: Fine-mapping, genomic organization, and transcript

[65493] analysis of the human ubiquitin-conjugating enzyme gene UBE2L3. Genomics 51:

[65494] 124-127, 1998.

[65495] 11060.

[65496] Cohen, L.; Henzel, W. J.; Baeuerle, P. A.: IKAP is a scaffold

[65497] protein of the I-kappa-B kinase complex. Nature 395:

292-296, 1998.

[65498] 11061.

[65499] Dong, J.; Edelmann, L.; Bajwa, A. M.; Kornreich, R.;
Desnick, R.

[65500] J.: Familial dysautonomia: detection of the IKBKAP
IVS20+6T-C and

[65501] R696P mutations and frequencies among Ashkenazi Jews.
Am. J. Med.

[65502] Genet. 110: 253-257, 2002.

[65503] 11062.

[65504] Hu, M. C.-T.; Qiu, W. R.; Wang, Y.-P.; Hill, D.; Ring, B. D.;
Scully,

[65505] S.; Bolon, B.; DeRose, M.; Luethy, R.; Simonet, W. S.;
Arakawa, T.;

[65506] Danilenko, D. M.: FGF-18, a novel member of the fibrob-
last growth

[65507] factor family, stimulates hepatic and intestinal prolifera-
tion. Molec.

[65508] Cell. Biol. 18: 6063-6074, 1998.

[65509] 11063.

[65510] Baens, M.; Chaffanet, M.; Cassiman, J.-J.; van den Berghe,

H.;

[65511] Marynen, P.: Construction and evaluation of a hncDNA library of human

[65512] 12 transcribed sequences derived from a somatic cell hybrid. Genomics 16:

[65513] 214-218, 1993.

[65514] 11064.

[65515] Fearnley, I. M.; Finel, M.; Skehel, J. M.; Walker, J. E.: NADH:ubiquinone

[65516] oxidoreductase from bovine heart mitochondria. Biochem. J. 278:

[65517] 821-829, 1991.

[65518] 11065.

[65519] Runswick, M. J.; Fearnley, I. M.; Skehel, J. M.; Walker, J. E.

[65520] : Presence of an acyl carrier protein in NADH:ubiquinone oxidoreductase

[65521] from bovine heart mitochondria. FEBS Lett. 286: 121-124, 1991.

[65522] 11066.

[65523] Bottorff, D.; Ebinu, J.; Stone, J. C.: RasGRP, a Ras activator:

[65524] mouse and human cDNA sequences and chromosomal

positions. Mammalian
[65525] Genome 10: 358-361, 1999.

[65526] 11067.

[65527] Ebinu, J. O.; Bottorff, D. A.; Chan, E. Y. W.; Stang, S. L.; Dunn,
[65528] R. J.; Stone, J. C.: RasGRP, a Ras guanyl nucleotide-releasing protein
[65529] with calcium- and diacylglycerol-binding motifs. Science 280: 1082-1086,
[65530] 1998.

[65531] 11068.

[65532] Kawasaki, H.; Springett, G. M.; Toki, S.; Canales, J. J.; Harlan,
[65533] P.; Blumenstiel, J. P.; Chen, E. J.; Bany, I. A.; Mochizuki, N.; Ashbacher,
[65534] A.; Matsuda, M.; Housman, D. E.; Graybiel, A. M.: A Rap guanine nucleotide
[65535] exchange factor enriched highly in the basal ganglia. Proc. Nat.
[65536] Acad. Sci. 95: 13278-13283, 1998.

[65537] 11069.

[65538] Bulman, D. E.; Scoggan, K. A.; van Oene, M. D.; Nicolle, M. W.;

[65539] Hahn, A. F.; Tollar, L. L.; Ebers, G. C.: A novel sodium channel

[65540] mutation in a family with hypokalemic periodic paralysis. Neurology 53:

[65541] 1932-1936, 1999.

[65542] 11070.

[65543] La Starza, R.; Wlodarska, I.; Aventin, A.; Falzetti, D.; Crescenzi,

[65544] B.; Martelli, M. F.; Van den Berghe, H.; Mecucci, C.: Molec-ular delineation

[65545] of 13q deletion boundaries in 20 patients with myeloid malignancies. Blood 91:

[65546] 231-237, 1998.

[65547] 11071.

[65548] El-Husseini, A. E.; Kwasnicka, D.; Yamada, T.; Hirohashi, S.; Vincent,

[65549] S. R.: BERP, a novel ring finger protein, binds to alpha-actinin-4. Biochem.

[65550] Biophys. Res. Commun. 267: 906-911, 2000.

[65551] 11072.

[65552] Favre, B.; Fontao, L.; Koster, J.; Shafaatian, R.; Jaunin, F.;

[65553] Saurat, J.-H.; Sonnenberg, A.; Borradori, L.: The hemidesmosomal

[65554] protein bullous pemphigoid antigen 1 and the integrin beta-4 subunit

[65555] bind to ERBIN: molecular cloning of multiple alternative splice variants

[65556] of ERBIN and analysis of their tissue expression. J. Biol. Chem. 276:

[65557] 32427-32436, 2001.

[65558] 11073.

[65559] Agnello, V.; Abel, G.; Elfahal, M.; Knight, G. B.; Zhang, Q.-X.

[65560] : Hepatitis C virus and other flaviviridae viruses enter cells via

[65561] low density lipoprotein receptor. Proc. Nat. Acad. Sci. 96: 12766-12771,

[65562] 1999.

[65563] 11074.

[65564] Allen, J. M.; Thompson, G. R.; Myant, N. B.; Steiner, R.;

Oakley,

[65565] C. M.: Cardiovascular complications of homozygous familial hypercholesterolaemia. Brit.

[65566] Heart J. 44: 361-368, 1980.

[65567] 11075.

[65568] Benlian, P.; Amselem, S.; Loux, N.; Pastier, D.; Giraud, G.; de

[65569] Gennes, J. L.; Turpin, G.; Monnier, L.; Rieu, D.; Douste-Blazy, P.;

[65570] Dastugue, B.; Goossens, M.; Junien, C.: A LDL receptor gene homozygous

[65571] mutation: PCR amplification, direct genomic sequencing, associated

[65572] haplotype, rapid screening for frequency. Ann. Genet. 33: 65-69,

[65573] 1990.

[65574] 11076.

[65575] Bertolini, S.; Lelli, N.; Coviello, D. A.; Ghisellini, M.; Masturzo,

[65576] P.; Tiozzo, R.; Elicio, N.; Gaddi, A.; Calandra, S.: A large deletion

[65577] in the LDL receptor gene--the cause of familial hyperc-

holesterolemia

[65578] in three Italian families: a study that dates back to the 17th century

[65579] (FH-Pavia). Am. J. Hum. Genet. 51: 123-134, 1992.

[65580] 11077.

[65581] Betard, C.; Kessling, A. M.; Roy, M.; Chamberland, A.; Lussier-Cacan,

[65582] S.; Davignon, J.: Molecular genetic evidence for a founder effect

[65583] in familial hypercholesterolemia among French Canadians. Hum. Genet. 88:

[65584] 529-536, 1992.

[65585] 11078.

[65586] Boehnke, M.; Arnheim, N.; Li, H.; Collins, F. S.: Fine-structure

[65587] genetic mapping of human chromosomes using the poly-merase chain reaction

[65588] on single sperm: experimental design considerations. Am. J. Hum.

[65589] Genet. 45: 21-32, 1989.

[65590] 11079.

[65591] Brown, M. S.; Goldstein, J. L.: Receptor-mediated endocy-
tosis:
[65592] insights from the lipoprotein receptor system. Proc. Nat.
Acad. Sci. 76:
[65593] 3330-3337, 1979.

[65594] 11080.

[65595] Defesche, J. C.; van Diermen, D. E.; Lansberg, P. J.; Lamp-
ing,
[65596] R. J.; Reymer, P. W. A.; Hayden, M. R.; Kastelein, J. J. P.:
South
[65597] African founder mutations in the low-density lipoprotein
receptor
[65598] gene causing familial hypercholesterolemia in the Dutch
population. Hum.
[65599] Genet. 92: 567-570, 1993.

[65600] 11081.

[65601] Defesche, J. C.; van de Ree, M. A.; Kastelein, J. J. P.; van
Diermen,
[65602] D. E.; Janssens, N. W. E.; van Doormaal, J. J.; Hayden, M.
R.: Detection
[65603] of the pro664-to-leu mutation in the low-density
lipoprotein receptor

[65604] and its relation to lipoprotein(a) levels in patients with familial

[65605] hypercholesterolemia of Dutch ancestry from The Netherlands and Canada. Clin.

[65606] Genet. 42: 273-280, 1992.

[65607] 11082.

[65608] Chen, H. I.; Einbond, A.; Kwak, S.-J.; Linn, H.; Koepf, E.; Peterson,

[65609] S.; Kelly, J. W.; Sudol, M.: Characterization of the WW domain of

[65610] human Yes-associated protein and its polyproline containing ligands. J.

[65611] Biol. Chem. 272: 17070-17077, 1997.

[65612] 11083.

[65613] Chen, H. I.; Sudol, M.: The WW domain of Yes-associated protein

[65614] binds a proline-rich ligand that differs from the consensus established

[65615] for Src homology 3-binding modules. Proc. Nat. Acad. Sci. 92: 7819-7823,

[65616] 1995.

[65617] 11084.

[65618] Millward, T.; Cron, P.; Hemmings, B. A.: Molecular cloning and

[65619] characterization of a conserved nuclear serine(threonine) protein

[65620] kinase. Proc. Nat. Acad. Sci. 92: 5022-5026, 1995.

[65621] 11085.

[65622] Hoggard, N.; Brintell, B.; Howell, A.; Weissenbach, J.; Var-ley,

[65623] J.: Allelic imbalance on chromosome 1 in human breast cancer. II.

[65624] Microsatellite repeat analysis. Genes Chromosomes Can-cer 12: 24-31,

[65625] 1995.

[65626] 11086.

[65627] White, G. R. M.; Varley, J. M.; Heighway, J.: Isolation and characterization

[65628] of a human homologue of the latrophilin gene from a re-gion of 1p31.1

[65629] implicated in breast cancer. Oncogene 17: 3513-3519, 1998.

[65630] 11087.

[65631] Heyer, B. S.; Warsowe, J.; Solter, D.; Knowles, B. B.; Acker-
man,
[65632] S. L.: New member of the Snf1/AMPK kinase family, Melk,
is expressed
[65633] in the mouse egg and preimplantation embryo. Molec. Re-
prod. Dev. 47:
[65634] 148-156, 1997.

[65635] 11088.

[65636] Merrill, R. A.; Plum, L. A.; Kaiser, M. E.; Clagett-Dame, M.:

[65637] A mammalian homolog of unc-53 is regulated by all-trans
retinoic acid
[65638] in neuroblastoma cells and embryos. Proc. Nat. Acad. Sci.
99: 3422-3427,
[65639] 2002.

[65640] 11089.

[65641] van Hille, B.; Richener, H.; Evans, D. B.; Green, J. R.; Bilbe,

[65642] G.: Identification of two subunit A isoforms of the vacuolar
H(+)-ATPase
[65643] in human osteoclastoma. J. Biol. Chem. 268: 7075-7080,
1993.

[65644] 11090.

[65645] van Hille, B.; Richener, H.; Green, J. R.; Bilbe, G.: The ubiq-
uitous

[65646] VA68 isoform of subunit A of the vacuolar H(+)-ATPase is
highly expressed

[65647] in human osteoclasts. Biochem. Biophys. Res. Commun.
214: 1108-1113,

[65648] 1995.

[65649] 11091.

[65650] Denning, G.; Jamieson, L.; Maquat, L. E.; Thompson, E. A.;
Fields,

[65651] A. P.: Cloning of a novel phosphatidylinositol kinase-re-
lated kinase:

[65652] characterization of the human SMG-1 RNA surveillance
protein. J.

[65653] Biol. Chem. 276: 22709-22714, 2001.

[65654] 11092.

[65655] Diaz-Meco, M. T.; Municio, M. M.; Sanchez, P.; Lozano, J.;
Moscat,

[65656] J.: Lambda-interacting protein, a novel protein that
specifically

[65657] interacts with the zinc finger domain of the atypical protein kinase

[65658] C isotype lambda/iota and stimulates its kinase activity in vitro

[65659] and in vivo. Molec. Cell. Biol. 16: 105-114, 1996.

[65660] 11093.

[65661] Kondo, M.; Ji, L.; Kamibayashi, C.; Tomizawa, Y.; Randle, D.; Sekido,

[65662] Y.; Yokota, J.; Kashuba, V.; Zabarovsky, E.; Kuzmin, I.; Lerman, M.;

[65663] Roth, J.; Minna, J. D.: Overexpression of candidate tumor suppressor

[65664] gene FUS1 isolated from the 3p21.3 homozygous deletion region leads

[65665] to G1 arrest and growth inhibition of lung cancer cells. Oncogene 20:

[65666] 6258-6262, 2001.

[65667] 11094.

[65668] Frank, D.; Mendelsohn, C. L.; Ciccone, E.; Svensson, K.; Ohlsson,

[65669] R.; Tycko, B.: A novel pleckstrin homology-related gene family defined

[65670] by Ipl/Tssc3, TDAG51, and Tih1: tissue-specific expression, chromosomal
[65671] location, and parental imprinting. Mammalian Genome 10: 1150-1159,
[65672] 1999.

[65673] 11095.

[65674] Sultana, R.; Yu, C.-E.; Yu, J.; Munson, J.; Chen, D.; Hua, W.;

[65675] Estes, A.; Cortes, F.; de la Barra, F.; Yu, D.; Haider, S. T.; Trask,
[65676] B. J.; Green, E. D.; Raskind, W. H.; Disteche, C. M.; Wijsman, E.;
[65677] Dawson, G.; Storm, D. R.; Schellenberg, G. D.; Villacres, E. C.:
[65678] Identification of a novel gene on chromosome 7q11.2 interrupted by
[65679] a translocation breakpoint in a pair of autistic twins. Genomics 80:
[65680] 129-134, 2002.

[65681] 11096.

[65682] Deshpande, K. L.; Seubert, P. H.; Tillman, D. M.; Farkas, W. R.;

[65683] Katze, J. R.: Cloning and characterization of cDNA encoding the rabbit
[65684] tRNA-guanine transglycosylase 60-kilodalton subunit. Arch. Biochem.
[65685] Biophys. 326: 1-7, 1996.

[65686] 11097.

[65687] Wilson, S. M.; Bhattacharyya, B.; Rachel, R. A.; Coppola, V.; Tessarollo,
[65688] L.; Householder, D. B.; Fletcher, C. F.; Miller, R. J.; Copeland,
[65689] N. G.; Jenkins, N. A.: Synaptic defects in ataxia mice result from
[65690] a mutation in Usp14, encoding a ubiquitin-specific protease. Nature
[65691] Genet. 7Oct, 2002. Note: Advance Electronic Publication.

[65692] 11098.

[65693] Triggs-Raine, B. L.; Akerman, B. R.; Clarke, J. T. R.; Gravel,
[65694] R. A.: Sequence of DNA flanking the exons of the HEXA gene, and identification
[65695] of mutations in Tay-Sachs disease. Am. J. Hum. Genet. 49: 1041-1054,

[65696] 1991.

[65697] 11099.

[65698] Triggs-Raine, B. L.; Gravel, R. A.: Diagnostic heteroduplexes:

[65699] simple detection of carriers of a 4-bp insertion mutation in Tay-Sachs

[65700] disease. Am. J. Hum. Genet. 46: 183-184, 1990.

[65701] 11100.

[65702] Triggs-Raine, B. L.; Mules, E. H.; Kaback, M. M.; Lim-Steele,

[65703] J. S. T.; Dowling, C. E.; Akerman, B. R.; Natowicz, M. R.; Grebner,

[65704] E. E.; Navon, R.; Welch, J. P.; Greenberg, C. R.; Thomas, G. H.; Gravel,

[65705] R. A.: A pseudodeficiency allele common in non-Jewish Tay-Sachs carriers:

[65706] implications for carrier screening. Am. J. Hum. Genet. 51: 793-801,

[65707] 1992.

[65708] 11101.

[65709] Trop, I.; Kaplan, F.; Brown, C.; Mahuran, D.; Hechtman, P.:

A

[65710] gly250-to-asp substitution in the alpha-subunit of hex-
osaminidase

[65711] A causes juvenile-onset Tay-Sachs disease in a Lebanese-
Canadian family. Hum.

[65712] Mutat. 1: 35-39, 1992.

[65713] 11102.

[65714] Trop, I.; Kaplan, F.; Hechtman, P.: Juvenile-onset Tay-
Sachs

[65715] disease in a Lebanese proband is caused by
gly(250)-to-asp substitution

[65716] in the alpha subunit of hexosaminidase A. (Abstract) Am.
J. Hum.

[65717] Genet. 47 (suppl.): A168, 1990.

[65718] 11103.

[65719] Whitley, C. B.; Anderson, R. A.; McIvor, R. S.: Heterozygos-
ity

[65720] for the 'DN allele' (G533-to-A) of the beta-
hexosaminidase alpha subunit

[65721] gene identified by direct DNA sequencing in a family with
the B1 variant

[65722] of G(M2)-gangliosidosis. Neuropediatrics 23: 96-101,

1992.

[65723] 11104.

[65724] Zlotogora, J.: Is the presence of two different Tay-Sachs disease

[65725] mutations in a Cajun population an unexpected observa-tion? (Letter) Am.

[65726] J. Hum. Genet. 52: 1014-1015, 1993.

[65727] 11105.

[65728] Zokaeem, G.; Bayleran, J.; Kaplan, P.; Hechtman, P.; Neufeld,

[65729] E. F.: A shortened beta-hexosaminidase alpha-chain in an Italian

[65730] patient with infantile Tay-Sachs disease. Am. J. Hum. Genet. 40:

[65731] 537-547, 1987.

[65732] 11106.

[65733] Cunningham, S. A.; Arrate, M. P.; Rodriguez, J. M.; Bjercke, R.

[65734] J.; Vanderslice, P.; Morris, A. P.; Brock, T. A.: A novel pro-tein

[65735] with homology to the junctional adhesion molecule: char-

acterization

[65736] of leukocyte interactions. J. Biol. Chem. 275: 34750-34756, 2000.

[65737] 11107.

[65738] Liang, T. W.; Chiu, H. H.; Gurney, A.; Sidle, A.; Tumas, D. B.;

[65739] Schow, P.; Foster, J.; Klassen, T.; Dennis, K.; DeMarco, R. A.; Pham,

[65740] T.; Frantz, G.; Fong, S.: Vascular endothelial-junctional adhesion

[65741] molecule (VE-JAM)/JAM 2 interacts with T, NK, and dendritic cells

[65742] through JAM 3. J. Immun. 168: 1618-1626, 2002.

[65743] 11108.

[65744] Palmeri, D.; van Zante, A.; Huang, C.-C.; Hemmerich, S.; Rosen,

[65745] S. D.: Vascular endothelial junction-associated molecule, a novel

[65746] member of the immunoglobulin superfamily, is localized to intercellular

[65747] boundaries of endothelial cells. J. Biol. Chem. 275: 19139-19145,

[65748] 2000.

[65749] 11109.

[65750] Arrate, M. P.; Rodriguez, J. M.; Tran, T. T.; Brock, T. A.; Cunningham,

[65751] S. A.: Cloning of human junctional adhesion molecule 3 (JAM3) and

[65752] its identification as the JAM2 counter-receptor. J. Biol. Chem. 276:

[65753] 45826-45832, 2001.

[65754] 11110.

[65755] Gregersen, N.; Winter, V. S.; Corydon, M. J.; Corydon, T. J.; Rinaldo,

[65756] P.; Ribes, A.; Martinez, G.; Bennett, M. J.; Vianey-Saban, C.; Bhala,

[65757] A.; Hale, D. E.; Lehnert, W.; Kmoch, S.; Roig, M.; Riudor, E.; Eiberg,

[65758] H.; Andresen, B. S.; Bross, P.; Bolund, L. A.; Kolvraa, S.: Identification

[65759] of four new mutations in the short-chain acyl-CoA dehydrogenase (SCAD)

[65760] gene in two patients: one of the variant alleles, 511C-T, is present

[65761] at an unexpectedly high frequency in the general popula-
tion, as was

[65762] the case for 625G-A, together conferring susceptibility to
ethylmalonic

[65763] aciduria. Hum. Molec. Genet. 7: 619-627, 1998.

[65764] 11111.

[65765] Kelly, C. L.; Wood, P. A.: Cloning and characterization of
the

[65766] mouse short-chain acyl-CoA dehydrogenase gene. Mam-
malian Genome 7:

[65767] 262-264, 1996.

[65768] 11112.

[65769] Naito, E.; Indo, Y.; Tanaka, K.: Short chain acyl-coenzyme
A dehydrogenase

[65770] (SCAD) deficiency: immunochemical demonstration of
molecular heterogeneity

[65771] due to variant SCAD with differing stability. J. Clin. Invest.
84:

[65772] 1671-1674, 1989.

[65773] 11113.

[65774] Naito, E.; Indo, Y.; Tanaka, K.: Short chain acyl-CoA dehy-

drogenase
[65775] (SCAD) deficiency: demonstration of molecular heterogeneity and identification
[65776] of point mutations. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A208,
[65777] 1989.

[65778] 11114.

[65779] Naito, E.; Indo, Y.; Tanaka, K.: Identification of two variant

[65780] short chain acyl-coenzyme A dehydrogenase alleles, each containing
[65781] a different point mutation in a patient with short chain acyl-coenzyme
[65782] A dehydrogenase deficiency. J. Clin. Invest. 85: 1575-1582, 1990.
[65783] 11115.

[65784] Naito, E.; Ozasa, H.; Ikeda, Y.; Tanaka, K.: Molecular cloning
[65785] and nucleotide sequence of cDNA encoding human short chain acyl-CoA
[65786] dehydrogenase (SCAD) and a study of its genetic deficiency. (Abstract) Am.
[65787] J. Hum. Genet. 43: A197, 1988.

[65788] 11116.

[65789] Naito, E.; Ozasa, H.; Ikeda, Y.; Tanaka, K.: Molecular cloning

[65790] and nucleotide sequence of complementary DNAs encoding human short

[65791] chain acyl-coenzyme A dehydrogenase and the study of the molecular

[65792] basis of human short chain acyl-coenzyme A dehydrogenase deficiency. J.

[65793] Clin. Invest. 83: 1605-1613, 1989.

[65794] 11117.

[65795] Garrett, R. M.; Bellissimo, D. B.; Rajagopalan, K. V.: Molecular

[65796] cloning of human liver sulfite oxidase. Biochim. Biophys. Acta 1262:

[65797] 147-149, 1995.

[65798] 11118.

[65799] Johnson, J. L.; Coyne, K. E.; Garrett, R. M.; Zabot, M.-T.; Dorche,

[65800] C.; Kisker, C.; Rajagopalan, K. V.: Isolated sulfite oxidase deficiency:

[65801] identification of 12 novel SUOX mutations in 10 patients.(Abstract) Hum.

[65802] Mutat. 20: 74 only, 2002.

[65803] 11119.

[65804] Chen, Y. Q.; Rafi, M. A.; de Gala, G.; Wenger, D. A.: Cloning

[65805] and expression of cDNA encoding human galactocerebrosidase, the enzyme

[65806] deficient in globoid cell leukodystrophy. Hum. Molec. Genet. 2:

[65807] 1841-1845, 1993.

[65808] 11120.

[65809] Chen, Y. Q.; Wenger, D. A.: Galactocerebrosidase from human urine:

[65810] purification and partial characterization. Biochim. Biophys. Acta 1170:

[65811] 53-61, 1993.

[65812] 11121.

[65813] Fiumara, A.; Pavone, L.; Siciliano, L.; Tine, A.; Parano, E.; Innico,

[65814] G.: Late-onset globoid cell leukodystrophy: report on 7

new patients. Child's

[65815] Nerv. Syst. 6: 194-197, 1990.

[65816] 11122.

[65817] Furuya, H.; Kukita, Y.; Nagano, S.; Sakai, Y.; Yamashita, Y.; Fukuyama,

[65818] H.; Inatomi, Y.; Saito, Y.; Koike, R.; Tsuji, S.; Fukumaki, Y.; Hayashi,

[65819] K.; Kobayashi, T.: Adult onset globoid cell leukodystrophy (Krabbe

[65820] disease): analysis of galactosylceramidase cDNA from four Japanese

[65821] patients. Hum. Genet. 100: 450-456, 1997.

[65822] 11123.

[65823] Kodama, S.; Igisu, H.; Siegel, D. A.; Suzuki, K.: Glycosylce-ramide

[65824] synthesis in the developing spinal cord and kidney of the twitcher

[65825] mouse, an enzymatically authentic model of human Krabbe disease. J.

[65826] Neurochem. 39: 1314-1318, 1982.

[65827] 11124.

[65828] Luzi, P.; Rafi, M. A.; Victoria, T.; Baskin, G. B.; Wenger, D.

[65829] A.: Characterization of the rhesus monkey galactocere-
brosidase (GALC)

[65830] cDNA and gene and identification of the mutation causing
globoid cell

[65831] leukodystrophy (Krabbe disease) in this primate. Ge-
nomics 42: 319-324,

[65832] 1997.

[65833] 11125.

[65834] Luzi, P.; Rafi, M. A.; Wenger, D. A.: Structure and organi-
zation

[65835] of the human galactocerebrosidase (GALC) gene. Ge-
nomics 26: 407-409,

[65836] 1995.

[65837] 11126.

[65838] Acharya, S.; Foletta, V. C.; Lee, J. W.; Rayborn, M. E.; Ro-
driguez,

[65839] I. R.; Young, W. S., III; Hollyfield, J. G.: SPACRCAN, a novel
human

[65840] interphotoreceptor matrix hyaluronan-binding proteogly-
can synthesized

[65841] by photoreceptors and pinealocytes. J. Biol. Chem. 275: 6945-6955,

[65842] 2000.

[65843] 11127.

[65844] Kuehn, M. H.; Hageman, G. S.: Molecular characterization and genomic

[65845] mapping of human IPM 200, a second member of a novel family of proteoglycans. Molec.

[65846] Cell Biol. Res. Commun. 2: 103-110, 1999.

[65847] 11128.

[65848] Kuehn, M. H.; Stone, E. M.; Hageman, G. S.: Organization of the

[65849] human IMPG2 gene and its evaluation as a candidate gene in age-related

[65850] macular degeneration and other retinal degenerative dis- orders. Invest.

[65851] Ophthal. Vis. Sci. 42: 3123-3129, 2001.

[65852] 11129.

[65853] Kang, D.; Jiang, H.; Wu, Q.; Pestka, S.; Fisher, P. B.: Cloning

[65854] and characterization of human ubiquitin-processing pro-

tease-43 from
[65855] terminally differentiated human melanoma cells using a rapid subtraction
[65856] hybridization protocol RaSH. Gene 267: 233-242, 2001.

[65857] 11130.

[65858] Li, X.-L.; Blackford, J. A.; Judge, C. S.; Liu, M.; Xiao, W.; Kalvakolanu,
[65859] D. V.; Hassel, B. A.: RNase-L-dependent destabilization of interferon-induced
[65860] mRNAs: a role for the 2-5A system in attenuation of the interferon
[65861] response. J. Biol. Chem. 275: 8880-8888, 2000.

[65862] 11131.

[65863] Liu, L.-Q.; Ilaria, R., Jr.; Kingsley, P. D.; Iwama, A.; van Et-ten,
[65864] R.; Palis, J.; Zhang, D. E.: A novel ubiquitin-specific pro-tease,
[65865] UBP43, cloned from leukemia fusion protein AML1-ETO-expressing mice,
[65866] functions in hematopoietic cell differentiation. Molec. Cell. Biol. 19:
[65867] 3029-3038, 1999.

[65868] 11132.

[65869] Schwer, H.; Liu, L.-Q.; Zhou, L.; Little, M.-T.; Pan, Z.; Het-
herington,
[65870] C. J.; Zhang, D.-E.: Cloning and characterization of a novel
human
[65871] ubiquitin-specific protease, a homologue of murine
UBP43 (Usp18). Genomics 65:
[65872] 44-52, 2000.

[65873] 11133.

[65874] Wang, K.; Zhou, B.; Kuo, Y.-M.; Zemansky, J.; Gitschier, J.:
A
[65875] novel member of a zinc transporter family is defective in
acrodermatitis
[65876] enteropathica. Am. J. Hum. Genet. 71: 66-73, 2002.

[65877] 11134.

[65878] Freeman, B. C.; Yamamoto, K. R.: Disassembly of tran-
scriptional
[65879] regulatory complexes by molecular chaperones. Science
296: 2232-2235,
[65880] 2002.

[65881] 11135.

[65882] Johnson, J. L.; Beito, T. G.; Krco, C. J.; Toft, D. O.: Characterization
[65883] of a novel 23-kilodalton protein of unactive progesterone receptor
[65884] complexes. Molec. Cell. Biol. 14: 1956-1963, 1994.

[65885] 11136.

[65886] Smith, D. F.; Faber, L. E.; Toft, D. O.: Purification of unactivated
[65887] progesterone receptor and identification of novel receptor-associated
[65888] proteins. J. Biol. Chem. 265: 3996-4003, 1990.

[65889] 11137.

[65890] Patterson, C. E.; Gao, J.; Rooney, A. P.; Davis, E. C.: Genomic
[65891] organization of mouse and human 65 kDa FK506-binding protein genes
[65892] and evolution of the FKBP multigene family. Genomics 79: 881-889,
[65893] 2002.

[65894] 11138.

[65895] Patterson, C. E.; Schaub, T.; Coleman, E. J.; Davis, E. C.:

Developmental
[65896] regulation of FKBP65: an ER localized extracellular matrix-binding
[65897] protein. Molec. Biol. Cell 11: 3925-3935, 2000.

[65898] 11139.

[65899] Cai, H.; Wang, Y.; McCarthy, D.; Wen, H.; Borchelt, D. R.; Price,
[65900] D. L.; Wong, P. C.: BACE1 is the major beta-secretase for generation
[65901] of A-beta peptides by neurons. Nature Neurosci. 4: 233-234, 2001.
[65902] 11140.

[65903] Meyaard, L.; Adema, G. J.; Chang, C.; Woollatt, E.; Suther-land,
[65904] G. R.; Lanier, L. L.; Phillips, J. H.: LAIR-1, a novel inhibitory
[65905] receptor expressed on human mononuclear leukocytes. Immunity 7:
[65906] 283-290, 1997.

[65907] 11141.

[65908] Sathish, J. G.; Johnson, K. G.; Fuller, K. J.; LeRoy, F. G.; Meyaard,

[65909] L.; Sims, M. J.; Matthews, R. J.: Constitutive association of SHP-1
[65910] with leukocyte-associated Ig-like receptor-1 in human T cells. J.
[65911] Immun. 166: 1763-1770, 2001.

[65912] 11142.

[65913] Xu, M.; Zhao, R.; Zhao, Z. J.: Identification and characterization
[65914] of leukocyte-associated Ig-like receptor-1 as a major anchor protein
[65915] of tyrosine phosphatase SHP-1 in hematopoietic cells. J. Biol. Chem. 275:
[65916] 17440-17446, 2000.

[65917] 11143.

[65918] Hofmann, R. M.; Pickart, C. M.: Noncanonical MMS2-encoded ubiquitin-conjugating
[65919] enzyme functions in assembly of novel polyubiquitin chains for DNA
[65920] repair. Cell 96: 645-653, 1999.

[65921] 11144.

[65922] Rothofsky, M. L.; Lin, S. L.: CROC-1 encodes a protein

which mediates
[65923] transcriptional activation of the human FOS promoter. Gene 195:
[65924] 141-149, 1997.

[65925] 11145.

[65926] Sancho, E.; Vila, M. R.; Sanchez-Pulido, L.; Lozano, J. J.; Paciucci,
[65927] R.; Nadal, M.; Fox, M.; Harvey, C.; Bercovich, B.; Loukili, N.; Ciechanover,
[65928] A.; Lin, S. L.; Sanz, F.; Estivill, X.; Valencia, A.; Thomson, T.

[65929] M.: Role of UEV-1, an inactive variant of the E2 ubiquitin-conjugating
[65930] enzymes, in in vitro differentiation and cell cycle behavior of HT-29-M6
[65931] intestinal mucosecretory cells. Molec. Cell. Biol. 18: 576-589,
[65932] 1998.

[65933] 11146.

[65934] Birn, H.; Verroust, P. J.; Nexo, E.; Hager, H.; Jacobsen, C.; Christensen,
[65935] E. I.; Moestrup, S. K.: Characterization of an epithelial ap-

proximately
[65936] 460-kDa protein that facilitates endocytosis of intrinsic factor-vitamin
[65937] B12 and binds receptor-associated protein. J. Biol. Chem. 272: 26497-26504,
[65938] 1997.

[65939] 11147.

[65940] Bork, P.; Beckmann, G.: The CUB domain: a widespread module in
[65941] developmentally regulated proteins. J. Molec. Biol. 231: 539-545,
[65942] 1993.

[65943] 11148.

[65944] Fyfe, J. C.; Giger, U.; Hall, C. A.; Jezyk, P. F.; Klumpp, S. A.;

[65945] Levine, J. S.; Patterson, D. F.: Inherited selective intestinal cobalamin
[65946] malabsorption and cobalamin deficiency in dogs. Pediat. Res. 29:
[65947] 24-31, 1991.

[65948] 11149.

[65949] Kristiansen, M.; Aminoff, M.; Jacobsen, C.; de la Chapelle,

A.;

[65950] Krahe, R.; Verroust, P. J.; Moestrup, S. K.: Cubilin P1297L mutation

[65951] associated with hereditary megaloblastic anemia 1 causes impaired

[65952] recognition of intrinsic factor-vitamin B12 by cubilin. Blood 96:

[65953] 405-409, 2000.

[65954] 11150.

[65955] Kozyraki, R.; Fyfe, J.; Kristiansen, M.; Gerdes, C.; Jacobsen,

[65956] C.; Cui, S.; Christensen, E. I.; Aminoff, M.; de la Chapelle, A.;

[65957] Krahe, R.; Verroust, P. J.; Moestrup, S. K.: The intrinsic factor-vitamin

[65958] B12 receptor, cubilin, is a high-affinity apolipoprotein A-I receptor

[65959] facilitating endocytosis of high-density lipoprotein. Na- ture Med. 5:

[65960] 656-661, 1999.

[65961] 11151.

[65962] De Strooper, B.; Konig, G.: A firm base for drug develop-

ment. Nature 402:

[65963] 471-472, 1999.

[65964] 11152.

[65965] Fan, W.; Bennett, B. D.; Babu-Khan, S.; Luo, Y.; Louis,

J.-C.;

[65966] McCaleb, M.; Citron, M.; Vassar, R.; Richards, W. G.: Re-

sponse to

[65967] Saunders et al. (1999). Science 286: 1255a, 1999. Note:

Electronic

[65968] Publication.

[65969] 11153.

[65970] Haniu, M.; Denis, P.; Young, Y.; Mendiaz, E. A.; Fuller, J.;

Hui,

[65971] J. O.; Bennett, B. D.; Kahn, S.; Ross, S.; Burgess, T.; Katta,

V.;

[65972] Rogers, G.; Vassar, R.; Citron, M.: Characterization of

Alzheimer's

[65973] beta-secretase protein BACE: a pepsin family member

with unusual properties. J.

[65974] Biol. Chem. 275: 21099-21106, 2000.

[65975] 11154.

[65976] Moestrup, S. K.; Kozyraki, R.; Kristiansen, M.; Kaysen, J. H.;

[65977] Rasmussen, H. H.; Brault, D.; Pontillon, F.; Goda, F. O.; Christensen,

[65978] E. I.; Hammond, T. G.; Verroust, P. J.: The intrinsic factor-vitamin

[65979] B12 receptor and target of teratogenic antibodies is a me-galin-binding

[65980] peripheral membrane protein with homology to develop-mental proteins. J.

[65981] Biol. Chem. 273: 5235-5242, 1998.

[65982] 11155.

[65983] Hong, L.; Koelsch, G.; Lin, X.; Wu, S.; Terzyan, S.; Ghosh, A.

[65984] K.; Zhang, X. C.; Tang, J.: Structure of the protease domain of memapsin

[65985] 2 (beta-secretase) complexed with inhibitor. Science 290: 150-153,

[65986] 2000.

[65987] 11156.

[65988] Hussain, I.; Powell, D.; Howlett, D. R.; Tew, D. G.; Meek, T. D.;

[65989] Chapman, C.; Gloger, I. S.; Murphy, K. E.; Southan, C. D.; Ryan, D.

[65990] M.; Smith, T. S.; Simmons, D. L.; Walsh, F. S.; Dingwall, C.; Christie,

[65991] G.: Identification of a novel aspartic protease (asp2) as beta-secretase Molec.

[65992] Cell Neurosci. 14: 419-427, 1999.

[65993] 11157.

[65994] Parker, A. E.; Van de Weyer, I.; Laus, M. C.; Verhasselt, P.; Luyten,

[65995] W. H. M. L.: Identification of a human homologue of the Schizosaccharomyces

[65996] pombe rad17+ checkpoint gene. J. Biol. Chem. 273: 18340-18346, 1998.

[65997] Note: Erratum: J. Biol. Chem. 274: 24438-24439, 1999.

[65998] 11158.

[65999] von Deimling, F.; Scharf, J. M.; Liehr, T.; Rothe, M.; Kelter,

[66000] A.-R.; Albers, P.; Dietrich, W. F.; Kunkel, L. M.; Wernert, N.; Wirth,

[66001] B.: Human and mouse RAD17 genes: identification, localization, genomic

[66002] structure and histological expression pattern in normal testis and

[66003] seminoma. Hum. Genet. 105: 17-27, 1999.

[66004] 11159.

[66005] Serra-Pages, C.; Medley, Q. G.; Tang, M.; Hart, A.; Streuli, M.

[66006] : Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting

[66007] proteins. J. Biol. Chem. 273: 15611-15620, 1998.

[66008] 11160.

[66009] Thomas, M. K.; Yao, K.-M.; Tenser, M. S.; Wong, G. G.; Habener,

[66010] J. F.: Bridge-1, a novel PDZ-domain coactivator of E2A-mediated regulation

[66011] of insulin gene transcription. Molec. Cell. Biol. 19: 8492-8504,

[66012] 1999.

[66013] 11161.

[66014] Watanabe, T. K.; Saito, A.; Suzuki, M.; Fujiwara, T.; Taka-hashi,

[66015] E.; Slaughter, C. A.; DeMartino, G. N.; Hendil, K. B.; Chung,

C. H.;

[66016] Tanahashi, N.; Tanaka, K.: cDNA cloning and characterization of a

[66017] human proteasomal modulator subunit, p27 (PSMD9). Genomics 50: 241-250,

[66018] 1998.

[66019] 11162.

[66020] Kennedy, J.; Rossi, D. L.; Zurawski, S. M.; Vega, F., Jr.: Kastelein,

[66021] R. A.; Wagner, J. L.; Hannum, C. H.; Zlotnick, A.: Mouse IL-17: a

[66022] cytokine preferentially expressed by alpha beta TCR + CD4-CD8-T cells. J.

[66023] Interferon Cytokine Res. 16: 611-617, 1996.

[66024] 11163.

[66025] Kotake, S.; Udagawa, N.; Takahashi, N.; Matsuzaki, K.; Itoh, K.;

[66026] Ishiyama, S.; Saito, S.; Inoue, K.; Kamatani, N.; Gillespie, M. T.;

[66027] Martin, T. J.; Suda, T.: IL-17 in synovial fluids from patients with

[66028] rheumatoid arthritis is a potent stimulator of osteoclasto-

genesis. J.

[66029] Clin. Invest. 103: 1345-1352, 1999.

[66030] 11164.

[66031] Rouvier, E.; Luciani, M.-F.; Mattei, M.-G.; Denizot, F.; Golstein,

[66032] P.: CTLA-8, cloned from an activated T cell, bearing AU-rich messenger

[66033] RNA instability sequences, and homologous to a Herpesvirus saimiri

[66034] gene. J. Immun. 150: 5445-5456, 1993.

[66035] 11165.

[66036] Yao, Z.; Painter, S. L.; Fanslow, W. C.; Ulrich, D.; Macduff, B.

[66037] M.; Spriggs, M. K.; Armitage, R. J.: Human IL-17: a novel cytokine

[66038] derived from T cells. J. Immun. 155: 5483-5486, 1995.

[66039] 11166.

[66040] Higuti, T.; Tsurumi, C.; Kawamura, Y.; Tsujita, H.; Osaka, F.;

[66041] Yoshihara, Y.; Tani, I.; Tanaka, K.; Ichihara, A.: Molecular cloning

[66042] of cDNA for the import precursor of human coupling factor 6 of H(+)-ATP

[66043] synthase in mitochondria. Biochem. Biophys. Res. Commun. 178: 793-799,

[66044] 1991.

[66045] 11167.

[66046] Javed, A. A.; Ogata, K.; Sanadi, D. R.: Human mitochondrial ATP

[66047] synthase: cloning cDNA for the nuclear-encoded precursor of coupling

[66048] factor 6. Gene 97: 307-310, 1991.

[66049] 11168.

[66050] Webster, K. A.; Oliver, N. A.; Wallace, D. C.: Assignment of an

[66051] oligomycin-resistance locus to human chromosome 10. Somat. Cell Genet. 8:

[66052] 223-244, 1982.

[66053] 11169.

[66054] Parker, A. E.; Van de Weyer, I.; Laus, M. C.; Oostveen, I.; Yon,

[66055] J.; Verhasselt, P.; Luyten, W. H. M. L.: A human homologue

of Schizosaccharomyces
[66056] pombe rad1+ checkpoint gene encodes an exonuclease. J. Biol. Chem. 273:
[66057] 18332-18339, 1998.

[66058] 11170.

[66059] Headon, D. J.; Overbeek, P. A.: Involvement of a novel Tnf receptor
[66060] homologue in hair follicle induction. Nature Genet. 22: 370-374,
[66061] 1999.

[66062] 11171.

[66063] Hashimoto, T.; Inazawa, J.; Okamoto, N.; Tagawa, Y.; Bessho, Y.;
[66064] Honda, Y.; Nakanishi, S.: The whole nucleotide sequence and chromosomal
[66065] localization of the gene for human metabotropic gluta- mate receptor
[66066] subtype 6. Europ. J. Neurosci. 9: 1226-1235, 1997.

[66067] 11172.

[66068] Masu, M.; Iwakabe, H.; Tagawa, Y.; Miyoshi, T.; Yamashita, M.;

[66069] Fokuda, Y.; Sasaki, H.; Hiroi, K.; Nakamura, Y.; Shigemoto, R.; Takada,
[66070] M.; Nakamura, K.; Nakao, K.; Katsuki, M.; Nakanishi, S.: Specific
[66071] deficit of the ON response in visual transmission by targeted disruption
[66072] of the mGluR6 gene. Cell 80: 757-765, 1995.

[66073] 11173.

[66074] Nakajima, Y.; Iwakabe, H.; Akazawa, C.; Nawa, H.; Shigemoto, R.;
[66075] Mizuno, N.; Nakanishi, S.: Molecular characterization of a novel
[66076] retinal metabotropic glutamate receptor mGluR6 with a high agonist
[66077] selectivity for L-2-amino-4-phosphonobutyrate. J. Biol. Chem. 268:
[66078] 11868-11873, 1993.

[66079] 11174.

[66080] Barbon, A.; Ferraboli, S.; Barlati, S.: Assignment of the human
[66081] metabotropic glutamate receptor gene GRM4 to chromosome 6 band p21.3

[66082] by radiation hybrid mapping. Cytogenet. Cell Genet. 88: 210 only, 2000.

[66083] 11175.

[66084] Inoue, N.; Hess, K. D.; Moreadith, R. W.; Richardson, L. L.; Handel, M. A.; Watson, M. L.; Zinn, A. R.: New gene family defined by MORC, a nuclear protein required for mouse spermatogenesis. Hum. Molec. Genet. 8: 1201-1207, 1999.

[66085] 11176.

[66086] Watson, M. L.; Zinn, A. R.; Inoue, N.; Hess, K. D.; Cobb, J.; Handel, M. A.; Halaban, R.; Duchene, C. C.; Albright, G. M.; Moreadith, R. W.: Identification of morc (microrchidia), a mutation that results in arrest of spermatogenesis at an early meiotic stage in the the mouse. Proc. Nat. Acad. Sci. 95: 14361-14366, 1998.

[66095] 11177.

[66096] Morita, R.; Miyazaki, E.; Fong, C. G.; Chen, X.-N.; Koren-
berg,
[66097] J. R.; Delgado-Escueta, A. V.; Yamakawa, K.: JH8, a gene
highly homologous
[66098] to the mouse jerky gene, maps to the region for childhood
absence
[66099] epilepsy on 8q24. Biochem. Biophys. Res. Commun. 248:
307-314, 1998.

[66100] 11178.

[66101] Toth, M.; Grimsby, J.; Buzsaki, G.; Donovan, G. P.: Epileptic
[66102] seizures caused by inactivation of a novel gene, jerky, re-
lated to
[66103] centromere binding protein-B in transgenic mice. Nature
Genet. 11:
[66104] 71-75, 1995. Note: Erratum: Nature Genet. 12: 110 only,
1996.

[66105] 11179.

[66106] Zeng, Z.; Kyaw, H.; Gakenheimer, K. R.; Augustus, M.; Fan,
P.;
[66107] Zhang, X.; Su, K.; Carter, K. C.; Li, Y.: Cloning, mapping,
and tissue

[66108] distribution of a human homologue of the mouse jerky gene product. Biochem.

[66109] Biophys. Res. Commun. 236: 389-395, 1997.

[66110] 11180.

[66111] Honda, K.; Yamada, T.; Endo, R.; Ino, Y.; Gotoh, M.; Tsuda, H.;

[66112] Yamada, Y.; Chiba, H.; Hirohashi, S.: Actinin-4, a novel actin-bundling

[66113] protein associated with cell motility and cancer invasion. J. Cell

[66114] Biol. 140: 1383-1393, 1998. Note: Erratum: J. Cell Biol. 143: 277

[66115] only, 1998.

[66116] 11181.

[66117] Kimura, M.; Matsuda, Y.; Eki, T.; Yoshioka, T.; Okumura, K.; Hanaoka,

[66118] F.; Okano, Y.: Assignment of STK6 to human chromosome 20q13.2-q13.3

[66119] and a pseudogene STK6P to 1q41-q42. Cytogenet. Cell Genet. 79: 201-203,

[66120] 1997.

[66121] 11182.

[66122] Bryan, J.; Edwards, R.; Matsudaira, P.; Otto, J.; Wulfkuhle, J.

[66123] : Fascin, an echinoid actin-bundling protein, is a homolog of the

[66124] Drosophila singed gene product. Proc. Nat. Acad. Sci. 90: 9115-9119,

[66125] 1993.

[66126] 11183.

[66127] Dressel, U.; Thormeyer, D.; Altincicek, B.; Paululat, A.; Eg-

[66128] M.; Schneider, S.; Tenbaum, S. P.; Renkawitz, R.; Baniah-

mad, A.:

[66129] Alien, a highly conserved protein with characteristics of a corepressor

[66130] for members of the nuclear hormone receptor superfam-

ily. Molec. Cell.

[66131] Biol. 19: 3383-3394, 1999.

[66132] 11184.

[66133] Schaefer, L.; Beermann, M. L.; Miller, J. B.: Coding se-

quence,

[66134] genomic organization, chromosomal localization, and ex-

pression pattern

[66135] of the signalosome component Cops2: the mouse homologue of Drosophila

[66136] alien. Genomics 56: 310-316, 1999.

[66137] 11185.

[66138] Scott, A. F.: Personal Communication. Baltimore, Md. 11/8/2000.

[66139] 11186.

[66140] Seeger, M.; Kraft, R.; Ferrell, K.; Dawadschargal, B.-O.; Dumdey,

[66141] R.; Schade, R.; Gordon, C.; Naumann, M.; Dubiel, W.: A novel protein

[66142] complex involved in signal transduction possessing similarities to

[66143] 26S proteasome subunits. FASEB J. 12: 469-478, 1998.

[66144] 11187.

[66145] Born, T. L.; Thomassen, E.; Bird, T. A.; Sims, J. E.: Cloning

[66146] of a novel receptor subunit, AcPL, required for interleukin-18 signaling. J.

[66147] Biol. Chem. 273: 29445-29450, 1998.

[66148] 11188.

[66149] Csoka, A. B.; Scherer, S. W.; Stern, R.: Expression analysis of
[66150] six paralogous human hyaluronidase genes clustered on chromosomes
[66151] 3p21 and 17q31. Genomics 60: 356-361, 1999.

[66152] 11189.

[66153] Spurkland, A.; Brinchmann, J. E.; Markussen, G.; Pedeu-tour, F.;
[66154] Munthe, E.; Lea, T.; Vartdal, F.; Aasheim, H.-C.: Molecular cloning
[66155] of a T cell-specific adapter protein (TSAd) containing an Src homology
[66156] (SH) 2 domain and putative SH3 and phosphotyrosine binding sites. J.
[66157] Biol. Chem. 273: 4539-4546, 1998.

[66158] 11190.

[66159] Fu, C.; Turck, C. W.; Kurosaki, T.; Chan, A. C.: BLNK: a central
[66160] linker protein in B cell activation. Immunity 9: 93-103, 1998.
[66161] 11191.

[66162] Ishiai, M.; Kurosaki, M.; Pappu, R.; Okawa, K.; Ronko, I.; Fu,
[66163] C.; Shibata, M.; Iwamatsu, A.; Chan, A. C.; Kurosaki, T.: BLNK required
[66164] for cloning Syk to PLC-gamma-2 and Rac1-JNK in B cells. Immunity 10:
[66165] 117-125, 1999.

[66166] 11192.

[66167] Minegishi, Y.; Rohrer, J.; Coustan-Smith, E.; Lederman, H. M.;
[66168] Pappu, R.; Campana, D.; Chan, A. C.; Conley, M. E.: An es-sential
[66169] role for BLNK in human B cell development. Science 286: 1954-1957,
[66170] 1999.

[66171] 11193.

[66172] Wang, L.; Mizzen, C.; Ying, C.; Candau, R.; Barlev, N.; Brownell,
[66173] J.; Allis, C. D.; Berger, S. L.: Histone acetyltransferase ac-tivity
[66174] is conserved between yeast and human GCN5 and is re-quired for complementation

[66175] of growth and transcriptional activation. Molec. Cell. Biol. 17:
[66176] 519-527, 1997.

[66177] 11194.

[66178] Xu, W.; Edmondson, D. G.; Evrard, Y. A.; Wakamiya, M.; Behringer,
[66179] R. R.; Roth, S. Y.: Loss of Gcn5l2 leads to increased apoptosis and
[66180] mesodermal defects during mouse development. Nature Genet. 26: 229-232,
[66181] 2000.

[66182] 11195.

[66183] Xu, W.; Edmondson, D. G.; Roth, S. Y.: Mammalian GCN5 and P/CAF
[66184] acetyltransferases have homologous amino-terminal domains important
[66185] for recognition of nucleosomal substrates. Molec. Cell. Biol. 18:
[66186] 5659-5669, 1998.

[66187] 11196.

[66188] Yang, X.-J.; Ogryzko, V. V.; Nishikawa, J.; Howard, B. H.;

Nakatani,
[66189] Y.: A p300/CBP-associated factor that competes with the adenoviral
[66190] oncoprotein E1A. Nature 382: 319-324, 1996.

[66191] 11197.

[66192] Ahmad, W.; Nomura, K.; McGrath, J. A.; Hashimoto, I.; Christiano,
[66193] A. M.: A homozygous nonsense mutation in the zinc-finger domain of
[66194] the human hairless gene underlies congenital atrichia. (Letter) J.
[66195] Invest. Derm. 113: 281-283, 1999.

[66196] 11198.

[66197] Ahmad, W.; Zlotogorski, A.; Panteleyev, A. A.; Lam, H.; Ahmad,
[66198] M.; ul Haque, M. F.; Abdallah, H. M.; Dragan, L.; Chris-tiano, A. M.
[66199] : Genomic organization of the human hairless gene (HR) and identification
[66200] of a mutation underlying congenital atrichia in an Arab Palestinian
[66201] family. Genomics 56: 141-148, 1999.

[66202] 11199.

[66203] Brooke, H. C.: Hairless mice. J. Hered. 17: 173-174, 1926.

[66204] 11200.

[66205] Cichon, S.; Anker, M.; Vogt, I. R.; Rohleder, H.; Putzstuck, M.;

[66206] Hillmer, A.; Farooq, S. A.; Al-Dhafri, K. S.; Ahmad, M.; Haque, S.;

[66207] Rietschel, M.; Propping, P.; Kruse, R.; Nothen, M. M.: Cloning, genomic

[66208] organization, alternative transcripts and mutational analysis of the

[66209] gene responsible for autosomal recessive universal congenital alopecia. Hum.

[66210] Molec. Genet. 7: 1671-1679, 1998.

[66211] 11201.

[66212] Davies, K.: Hair apparent. Nature 391: 537-539, 1998.

[66213] 11202.

[66214] Gallagher, C. H.; Canfield, P. J.; Greenoak, G. E; Reeve, V. E.

[66215] : Characterization and histogenesis of tumors in the hairless mouse

[66216] produced by low-dosage incremental ultraviolet radiation. J. Invest.

[66217] Derm. 83: 169-174, 1984.

[66218] 11203.

[66219] Morrissey, P. J.; Parkinson, D. R.; Schwartz, R. S.; Waksal, S.

[66220] D.: Immunologic abnormalities in HRS/J mice.: I. specific deficit

[66221] in T lymphocyte helper function in a mutant mouse. J. Immun. 125:

[66222] 1558-1562, 1980.

[66223] 11204.

[66224] Nomura, K.; Hashimoto, I.: Atrichia with papular lesions: successful

[66225] genetic counselling about having a child. (Letter) Brit. J. Derm. 139:

[66226] 742-743, 1998.

[66227] 11205.

[66228] Potter, G. B.; Beaudoin, G. M. J., III; DeRenzo, C. L.; Zarach,

[66229] J. M.; Chen, S. H.; Thompson, C. C.: The hairless gene mutated in

[66230] congenital hair loss disorders encodes a novel nuclear receptor corepressor. Genes

[66231] Dev. 15: 2687-2701, 2001.

[66232] 11206.

[66233] Schuddekopf, K.; Schorpp, M.; Boehm, T.: The whn transcription

[66234] factor encoded by the nude locus contains an evolutionarily conserved

[66235] and functionally indispensable activation domain. Proc. Nat. Acad.

[66236] Sci. 93: 9661-9664, 1996.

[66237] 11207.

[66238] Stoye, J. P.; Fenner, S.; Greenoak, G. E.; Moran, C.; Coffin,

[66239] J. M.: Role of endogenous retroviruses as mutagens: the hairless

[66240] mutation of mice. Cell 54: 383-391, 1988.

[66241] 11208.

[66242] Thompson, C. C.: Thyroid hormone-responsive genes in developing

[66243] cerebellum include a novel synaptotagmin and a hairless homolog. J.

[66244] Neurosci. 16: 7832-7840, 1996.

[66245] 11209.

[66246] Sillence, D. O.; Barlow, K. K.; Garber, A. P.; Hall, J. G.; Ri-
moin,

[66247] D. L.: Osteogenesis imperfecta type II: delineation of the
phenotype

[66248] with reference to genetic heterogeneity. Am. J. Med.
Genet. 17:

[66249] 407-423, 1984.

[66250] 11210.

[66251] Miao, D.; He, B.; Karaplis, A. C.; Goltzman, D.: Parathyroid

[66252] hormone is essential for normal fetal bone formation. J.
Clin. Invest. 109:

[66253] 1173-1182, 2002.

[66254] 11211.

[66255] Raizis, A. M.; Becroft, D. M.; Shaw, R. L.; Reeve, A. E.: A
mitotic

[66256] recombination in Wilms tumor occurs between the
parathyroid hormone

[66257] locus and 11p13. Hum. Genet. 70: 344-346, 1985.

[66258] 11212.

[66259] Betticher, D. C.; Thatcher, N.; Altermatt, H. J.; Hoban, P.;
Ryder,

[66260] W. D. J.; Heighway, J.: Alternate splicing produces a novel
cyclin

[66261] D1 transcript. Oncogene 11: 1005-1011, 1995.

[66262] 11213.

[66263] Chesi, M.; Bergsagel, P. L.; Brents, L. A.; Smith, C. M.; Ger-
hard,

[66264] D. S.; Kuehl, W. M.: Dysregulation of cyclin D1 by translo-

cation
[66265] into an IgH gamma switch region in two multiple myeloma cell lines. Blood 88:
[66266] 674-681, 1996.

[66267] 11214.

[66268] Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold, A.; Marx,
[66269] S. J.; Bale, A. E.: Physical mapping of chromosome 11q12-13 by pulsed
[66270] field gel electrophoresis (PFGE). (Abstract) Cytogenet. Cell Genet. 58:
[66271] 1959, 1991.

[66272] 11215.

[66273] Geng, Y.; Yu, Q.; Sicinska, E.; Das, M.; Bronson, R. T.; Sicinski,
[66274] P.: Deletion of the p27(Kip1) gene restores normal development in
[66275] cyclin D1-deficient mice. Proc. Nat. Acad. Sci. 98:
194-199, 2001.
[66276] 11216.

[66277] Hayette, S.; Gadoux, M.; Martel, S.; Bertrand, S.; Tigaud, I.;

[66278] Magaud, J.-P.; Rimokh, R.: FLRG (follistatin-related gene), a new
[66279] target of chromosomal rearrangement in malignant blood disorders. Oncogene 16:
[66280] 2949-2954, 1998.

[66281] 11217.

[66282] Chang, S. S.; Grunder, S.; Hanukoglu, A.; Rosler, A.; Mathew, P.
[66283] M.; Hanukoglu, I.; Schild, L.; Lu, Y.; Shimkets, R. A.; Nel-son-Williams,
[66284] C.; Rossier, B. C.; Lifton, R. P.: Mutations in subunits of the epithelial
[66285] sodium channel cause salt wasting with hyperkalaemic acidosis, pseudohypoaldosteronism
[66286] type 1. Nature Genet. 12: 248-253, 1996.

[66287] 11218.

[66288] Tajima, T.; Kitagawa, H.; Yokoya, S.; Tachibana, K.; Adachi, M.;
[66289] Nakae, J.; Suwa, S; Katoh, S.; Fujieda, K.: A novel missense mutation
[66290] of mineralocorticoid receptor gene in one Japanese family with a renal

[66291] form of pseudohypoaldosteronism type 1. J. Clin. Endocr. Metab. 85:
[66292] 4690-4694, 2000.

[66293] 11219.

[66294] Viemann, M.; Peter, M.; Lopez-Siguero, J. P.; Simic-Schleicher,
[66295] G.; Sippell, W. G.: Evidence for genetic heterogeneity of pseudohypoaldosteronism
[66296] type 1: identification of a novel mutation in the human mineralocorticoid
[66297] receptor in one sporadic case and no mutations in two autosomal dominant
[66298] kindreds. J. Clin. Endocr. Metab. 86: 2056-2059, 2001.

[66299] 11220.

[66300] Hoyer, L. W.: The factor VIII complex: structure and function. Blood 58:
[66301] 1-13, 1981.

[66302] 11221.

[66303] Lee, Y. L.; Helman, L.; Hoffman, T.; Laborda, J.: dlk, pG2 and
[66304] Pref-1 mRNAs encode similar proteins belonging to the

EGF-like superfamily.

[66305] Identification of polymorphic variants of this RNA. Biochim. Biophys.

[66306] Acta 1261: 223-232, 1995.

[66307] 11222.

[66308] Smas, C. M.; Sul, H. S.: Pref-1, a protein containing EGF-like

[66309] repeats, inhibits adipocyte differentiation. Cell 73: 725-734, 1993.

[66310] 11223.

[66311] Takada, S.; Paulsen, M.; Tevendale, M.; Tsai, C.-E.; Kelsey, G.;

[66312] Cattanach, B. M.; Ferguson-Smith, A. C.: Epigenetic analysis of the

[66313] Dlk1-Gtl2 imprinted domain on mouse chromosome 12: implications for

[66314] imprinting control from comparison with Igf2-H19. Hum. Molec. Genet. 11:

[66315] 77-86, 2002.

[66316] 11224.

[66317] Wylie, A. A.; Murphy, S. K.; Orton, T. C.; Jirtle, R. L.: Novel

[66318] imprinted DLK1/GTL2 domain on human chromosome 14 contains motifs
[66319] that mimic those implicated in IGF2/H19 regulation. Genome Res. 10:
[66320] 1711-1718, 2000.

[66321] 11225.

[66322] Okumura, K.; Nogami, M.; Taguchi, H.; Hisamatsu, H.; Tanaka, K.
[66323] : The genes for the alpha-type HC3 (PMSA2) and beta-type HC5 (PMSB1)
[66324] subunits of human proteasomes map to chromosomes 6q27 and 7p12-p13
[66325] by fluorescence in situ hybridization. Genomics 27: 377-379, 1995.
[66326] 11226.

[66327] Tamura, T.; Lee, D. H.; Osaka, F.; Fujiwara, T.; Shin, S.; Chung,
[66328] C. H.; Tanaka, K.; Ichihara, A.: Molecular cloning and se-quence analysis
[66329] of cDNAs for five major subunits of human proteasomes (multi-catalytic
[66330] proteinase complexes). Biochim. Biophys. Acta 1089:

95-102, 1991.

[66331] 11227.

[66332] Tamura, T.; Osaka, F.; Kawamura, Y.; Higuchi, T.; Ishida, N.; Nothwang,

[66333] H. G.; Tsurumi, C.; Tanaka, K.; Ichihara, A. :J. Molec. Biol. 244:

[66334] 1117-1124, 1994.

[66335] 11228.

[66336] DeVry, C. G.; Clarke, S.: Assignment of the protein L-isoaspartate

[66337] (D-aspartate) O-methyltransferase gene (PCMT1) to human chromosome

[66338] bands 6q24-q25 with radiation hybrid mapping. Cytogenet. Cell Genet. 84:

[66339] 130-131, 1999.

[66340] 11229.

[66341] Farrar, C.; Clarke, S.: Altered levels of S-adenosylmethionine

[66342] and S-adenosylhomocysteine in the brains of L-isoaspartyl (D-aspartyl)

[66343] O-methyltransferase-deficient mice. J. Biol. Chem. 277:

27856-27863,
[66344] 2002.

[66345] 11230.

[66346] Ingrosso, D.; Kagan, R. M.; Clarke, S.: Distinct C-terminal sequences
[66347] of isozymes I and II of the human erythrocyte L-isoaspartyl/D-aspartyl
[66348] protein methyltransferase. Biochem. Biophys. Res. Commun. 175: 351-358,
[66349] 1991.

[66350] 11231.

[66351] Kim, E.; Lowenson, J. D.; Clarke, S.; Young, S. G.: Phenotypic
[66352] analysis of seizure-prone mice lacking L-isoaspartate (D-aspartate)
[66353] O-methyltransferase. J. Biol. Chem. 274: 20671-20678, 1999.
[66354] 11232.

[66355] Lowenson, J. D.; Kim, E.; Young, S. G.; Clarke, S.: Limited accumulation
[66356] of damaged proteins in L-isoaspartyl (D-aspartyl) O-

methyltransferase-deficient
[66357] mice. J. Biol. Chem. 276: 20695-20702, 2001.

[66358] 11233.

[66359] MacLaren, D. C.; Kagan, R. M.; Clarke, S.: Alternative splic-
ing
[66360] of the human isoaspartyl protein carboxyl methyltrans-
ferase leads
[66361] to the generation of a C-terminal -RDEL sequence in
isozyme II. Biochem.
[66362] Biophys. Res. Commun. 185: 277-283, 1992.

[66363] 11234.

[66364] MacLaren, D. C.; O'Connor, C. M.; Xia, Y.-R.; Mehrabian,
M.; Klisak,
[66365] I.; Sparkes, R. S.; Clarke, S.; Lusis, A. J.: The L-
isoaspartyl/D-aspartyl
[66366] protein methyltransferase gene (PCMT1) maps to human
chromosome 6q22.3-6q24
[66367] and the syntenic region of mouse chromosome 10. Ge-
nomics 14: 852-856,
[66368] 1992.

[66369] 11235.

[66370] Ota, I. M.; Gilbert, J. M.; Clarke, S.: Two major isozymes of
[66371] the protein D-aspartyl/L-isoaspartyl methyltransferase from human
[66372] erythrocytes. Biochem. Biophys. Res. Commun. 151: 1136-1143, 1988.
[66373] 11236.
[66374] Ton, C. C. T.; Huff, V.; Call, K. M.; Cohn, S.; Strong, L. C.;
[66375] Housman, D. E.; Saunders, G. F.: Smallest region of over-lap in Wilms
[66376] tumor deletions uniquely implicates an 11p13 zinc finger gene as the
[66377] disease locus. Genomics 10: 293-297, 1991.
[66378] 11237.
[66379] Astrom, A.-K.; Voz, M. L.; Kas, K.; Roijer, E.; Wedell, B.; Mandahl,
[66380] N.; Van de Ven, W.; Mark, J.; Stenman, G.: Conserved mechanism of
[66381] PLAG1 activation in salivary gland tumors with and with-out chromosome
[66382] 8q12 abnormalities: identification of SII as a new fusion partner

[66383] gene. Cancer Res. 59: 918-923, 1999.

[66384] 11238.

[66385] Bullerdiek, J.; Bartnitzke, S.; Weinberg, M.; Chilla, R.; Haubrich,

[66386] J.; Schloot, W.: Rearrangements of chromosome region 12q13-q15 in

[66387] pleomorphic adenomas of the human salivary gland (PSA). Cytogenet.

[66388] Cell Genet. 45: 187-190, 1987.

[66389] 11239.

[66390] Bullerdiek, J.; Raabe, G.; Bartnitzke, S.; Boschen, C.; Schloot,

[66391] W.: Structural rearrangements of chromosome #8 involving 8q12--a

[66392] primary event in pleomorphic adenomas of the parotid gland. Genetica 72:

[66393] 85-92, 1987.

[66394] 11240.

[66395] Mark, J.; Dahlenfors, R.: Cytogenetical observations in 100 human

[66396] benign pleomorphic adenomas: specificity of the chromo-

somal aberrations
[66397] and their relationship to sites of localized oncogenes. Anticancer
[66398] Res. 6: 299-308, 1986.

[66399] 11241.

[66400] Stenman, G.; Sandros, J.; Mark, J.; Nordkvist, A.: High p21(RAS)
[66401] expression levels correlate with chromosome 8 rearrangements in benign
[66402] human mixed salivary gland tumors. Genes Chromosomes Cancer 1: 59-66,
[66403] 1989.

[66404] 11242.

[66405] Aksoy, M.; Erdem, S.; Dincol, G.; Erdogan, G.; Cilingiroglu, K.;
[66406] Dincol, K.: Combination of hereditary elliptocytosis and hereditary
[66407] spherocytosis. Clin. Genet. 6: 46-50, 1974.

[66408] 11243.

[66409] Goodman, S. R.; Shiffer, K. A.; Casoria, L. A.; Eyster, M. E.

[66410] : Identification of the molecular defect in the erythrocyte

membrane
[66411] skeleton of some kindreds with hereditary spherocytosis. Blood 60:
[66412] 772-784, 1982.

[66413] 11244.

[66414] Kimberling, W. J.; Taylor, R. A.; Chapman, R. G.; Lubs, H. A.
[66415] : Linkage and gene localization of hereditary spherocytosis (HS). Blood 52:
[66416] 859-867, 1978.

[66417] 11245.

[66418] Hinds, P. W.; Dowdy, S. F.; Eaton, E. N.; Arnold, A.; Weinberg,
[66419] R. A.: Function of a human cyclin gene as an oncogene. Proc. Nat.
[66420] Acad. Sci. 91: 709-713, 1994.

[66421] 11246.

[66422] Goldfarb, L. G.; Petersen, R. B.; Tabaton, M.; Brown, P.; LeBlanc,
[66423] A. C.; Montagna, P.; Cortelli, P.; Julien, J.; Vital, C.; Pendelbury,

[66424] W. W.; Haltia, M.; Wills, P. R.; Hauw, J. J.; McKeever, P. E.; Monari,

[66425] L.; Schrank, B.; Swergold, G. D.; Autilio-Gambetti, L.; Gaj-dusek,

[66426] D. C.; Lugaresi, E.; Gambetti, P.: Fatal familial insomnia and familial

[66427] Creutzfeldt-Jakob disease: disease phenotype determined by a DNA polymorphism. Science 258:

[66428] 806-808, 1992.

[66429] 11247.

[66430] Komatsu, H.; Iida, S.; Yamamoto, K.; Mikuni, C.; Nitta, M.; Takahashi,

[66431] T.; Ueda, R.; Seto, M.: A variant chromosome translocation at 11q13

[66432] identifying PRAD1/cyclin D1 as the BCL-1 gene . Blood 84: 1226-1231,

[66433] 1994.

[66434] 11248.

[66435] Kong, S.; Amos, C. I.; Luthra, R.; Lynch, P. M.; Levin, B.; Frazier,

[66436] M. L.: Effects of cyclin D1 polymorphism on age of onset of hereditary

[66437] nonpolyposis colorectal cancer. Cancer Res. 60: 249-252, 2000.

[66438] 11249.

[66439] Kong, S.; Wei, Q.; Amos, C. I.; Lynch, P. M.; Zong, J.; Frazier,

[66440] M. L.: Cyclin D1 polymorphism and increased risk of colorectal cancer

[66441] at young age. J. Nat. Cancer Inst. 93: 1106-1108, 2001.

[66442] 11250.

[66443] Ma, C.; Papermaster, D.; Cepko, C. L.: A unique pattern of photoreceptor

[66444] degeneration in cyclin D1 mutant mice. Proc. Nat. Acad. Sci. 95:

[66445] 9938-9943, 1998.

[66446] 11251.

[66447] Motokura, T.; Bloom, T.; Kim, H. G.; Juppner, H.; Ruderman, J.

[66448] V.; Kronenberg, H. M.; Arnold, A.: A novel cyclin encoded by a bcl1-linked

[66449] candidate oncogene. Nature 350: 512-515, 1991.

[66450] 11252.

[66451] Muller, H.; Lukas, J.; Schneider, A.; Warthoe, P.; Bartek, J.;

[66452] Eilers, M.; Strauss, M.: Cyclin D1 expression is regulated by the

[66453] retinoblastoma protein. Proc. Nat. Acad. Sci. 91: 2945-2949, 1994.

[66454] 11253.

[66455] Richard, C. W., III; Withers, D. A.; Meeker, T. C.; Maurer, S.;

[66456] Evans, G. A.; Myers, R. M.; Cox, D. R.: A radiation hybrid map of

[66457] the proximal long arm of human chromosome 11 containing the multiple

[66458] endocrine neoplasia type I (MEN-1) and bcl-1 disease loci. Am. J.

[66459] Hum. Genet. 49: 1189-1196, 1991.

[66460] 11254.

[66461] Rimokh, R.; Berger, F.; Bastard, C.; Klein, B.; French, M.; Archimbaud,

[66462] E.; Rouault, J. P.; Santa Lucia, B.; Duret, L.; Vuillaume, M.; et

[66463] al.: Rearrangement of CCND1 (BCL1/PRAD1) 3-prime untranslated region

[66464] in mantle-cell lymphomas and t(11q13)-associated leukemias. Blood 83:

[66465] 3689-3696, 1994.

[66466] 11255.

[66467] Rimokh, R.; Berger, F.; Delsol, G.; Charrin, C.; Bertheas, M.

[66468] F.; Ffrench, M.; Garoscio, M.; Felman, P.; Coiffier, C.; Bryon, P.

[66469] A.: Rearrangement and overexpression of the BCL-1/PRAD-1 gene in

[66470] intermediate lymphocytic lymphomas and in t(11q13)-bearing leukemias. Blood 81:

[66471] 3063-3067, 1993.

[66472] 11256.

[66473] Rosenberg, C. L.; Kim, H. G.; Shows, T. B.; Kronenberg, H. M.;

[66474] Arnold, A.: Rearrangement and overexpression of D11S287E, a candidate

[66475] oncogene on chromosome 11q13 in benign parathyroid tumors. Oncogene 6:

[66476] 449-453, 1991.

[66477] 11257.

[66478] Sicinski, P.; Donaher, J. L.; Parker, S. B.; Li, T.; Fazeli, A.;

[66479] Gardner, H.; Haslam, S. Z.; Bronson, R. T.; Elledge, S. J.; Weinberg,

[66480] R. A.: Cyclin D1 provides a link between development and oncogenesis

[66481] in the retina and breast. Cell 82: 621-630, 1995.

[66482] 11258.

[66483] Szepetowski, P.; Perucca-Lostanlen, D.; Gaudray, P.: Map-ping

[66484] genes according to their amplification status in tumor cells: contribution

[66485] to the map of 11q13. Genomics 16: 745-750, 1993.

[66486] 11259.

[66487] Converse, P. J.: Personal Communication. Baltimore, Md. 2/12/2002.

[66488] 11260.

[66489] Flowers, K. M.; Mellor, H.; Kimball, S. R.; Jefferson, L. S.:

[66490] Structure and sequence of the gene encoding the alpha-subunit of rat

[66491] translation initiation factor-2B. Biochim. Biophys. Acta 1264: 163-167,

[66492] 1995.

[66493] 11261.

[66494] Wang, R.; Cukerman, E.; Heng, H. H. Q.; Liew, C.-C.: Identification

[66495] of a locus of zinc finger genes in human chromosome 19q13.1-q13.3

[66496] region by fluorescence in situ hybridization. Somat. Cell Molec.

[66497] Genet. 22: 245-248, 1996.

[66498] 11262.

[66499] Clark, T. G.; Conway, S. J.; Scott, I. C.; Labosky, P. A.; Winnier,

[66500] G.; Bundy, J.; Hogan, B. L. M.; Greenspan, D. S.: The mammalian Tolloid-like

[66501] 1 gene, Tll1, is necessary for normal septation and positioning of

[66502] the heart. Development 126: 126:-2631-2642, 1999.

[66503] 11263.

[66504] Scott, I. C.; Clark, T. G.; Takahara, K.; Hoffman, G. G.; Eddy,

[66505] R. L.; Haley, L. L.; Shows, T. B.; Greenspan, D. S.: Assign-

ment of
[66506] TLL1 and TLL2, which encode human BMP-
1/tolloid-related metalloproteases,
[66507] to chromosomes 4q32-q33 and 10q23-q24 and assign-
ment of murine Tll2
[66508] to chromosome 19. Cytogenet. Cell Genet. 86: 64-65,
1999.
[66509] 11264.

[66510] Takahara, K.; Brevard, R.; Hoffman, G. G.; Suzuki, N.;
Greenspan,
[66511] D. S.: Characterization of a novel gene product
(mammalian tolloid-like)
[66512] with high sequence similarity to mammalian tolloid/bone
morphogenetic
[66513] protein-1. Genomics 34: 157-165, 1996.

[66514] 11265.

[66515] Bauer, W. O.; Nanda, I.; Beck, G.; Schmid, M.; Jakob, F.:
Human
[66516] puromycin-sensitive aminopeptidase: cloning of 3-prime
UTR, evidence
[66517] for a polymorphism at aa 140 and refined chromosomal
localization

[66518] to 17q21. Cytogenet. Cell Genet. 92: 221-224, 2001.

[66519] 11266.

[66520] Huber, G.; Thompson, A.; Gruninger, F.; Mechler, H.; Hochstrasser,

[66521] R.; Hauri, H.-P.; Malherbe, P.: cDNA cloning and molecular characterization

[66522] of human brain metalloprotease MP100: a beta-secretase candidate? J.

[66523] Neurochem. 72: 1215-1223, 1999.

[66524] 11267.

[66525] Osada, T.; Sakaki, Y.; Takeuchi, T.: Puromycin-sensitive aminopeptidase

[66526] gene (Psa) maps to mouse chromosome 11. Genomics 56: 361-362, 1999.

[66527] 11268.

[66528] Schonlein, C.; Loffler, J.; Huber, G.: Purification and char-acterization

[66529] of a novel metalloprotease from human brain with the ability to cleave

[66530] substrates derived from the N-terminus of beta-amyloid protein. Biochem.

[66531] Biophys. Res. Commun. 201: 45-53, 1994.

[66532] 11269.

[66533] Thompson, M. W.; Tobler, A.; Fontana, A.; Hersh, L. B.: Cloning

[66534] and analysis of the gene for the human puromycin-sensitive aminopeptidase. Biochem.

[66535] Biophys. Res. Commun. 258: 234-240, 1999.

[66536] 11270.

[66537] Tobler, A. R.; Constam, D. B.; Schmitt-Graff, A.; Malipiero, U.;

[66538] Schlapbach, R.; Fontana, A.: Cloning of the human puromycin-sensitive

[66539] aminopeptidase and evidence for expression in neurons. J. Neurochem. 68:

[66540] 889-897, 1997.

[66541] 11271.

[66542] Zhang, Y.; Cai, X.; Schlegelberger, B.; Zheng, S.: Assignment

[66543] of human putative tumor suppressor genes ST13 (alias SNC6) and ST14

[66544] (alias SNC19) to human chromosome bands 22q13 and

11q24-q25 by in
[66545] situ hybridization. Cytogenet. Cell Genet. 83: 56-57, 1998.
[66546] 11272.

[66547] Friedrich, R.; Fuentes-Prior, P.; Ong, E.; Coombs, G.; Hunter,
[66548] M.; Oehler, R.; Pierson, D.; Gonzalez, R.; Huber, R.; Bode, W.; Madison,
[66549] E. L.: Catalytic domain structures of MT-SP1/matriptase, a matrix-
[66550] degrading transmembrane serine proteinase. J. Biol. Chem. 277: 2160-2168,
[66551] 2002.

[66552] 11273.

[66553] Lin, C.-Y.; Wang, J.-K.; Torri, J.; Dou, L.; Sang, Q. A.; Dick-son,
[66554] R. B.: Characterization of a novel, membrane-bound, 80-kDa matrix-degrading
[66555] protease from human breast cancer cells: monoclonal an-tibody production,
[66556] isolation, and localization. J. Biol. Chem. 272: 9147-9152, 1997.

[66557] 11274.

[66558] Gacy, A. M.; Goellner, G. M.; Spiro, C.; Chen, X.; Gupta, G.;

[66559] Bradbury, E. M.; Dyer, R. B.; Mikesell, M. J.; Yao, J. Z.;
Johnson,

[66560] A. J.; Richter, A.; Melancon, S. B.; McMurray, C. T.: GAA in-
stability

[66561] in Friedreich's ataxia shares a common, DNA-directed and
intraallelic

[66562] mechanism with other trinucleotide diseases. Molec. Cell
1: 583-593,

[66563] 1998.

[66564] 11275.

[66565] Gordon, D. M.; Kogan, M.; Knight, S. A. B.; Dancis, A.;
Pain,

[66566] D.: Distinct roles for two N-terminal cleaved domains in
mitochondrial

[66567] import of the yeast frataxin homolog, Yfh1p. Hum. Molec.
Genet. 10:

[66568] 259-269, 2001.

[66569] 11276.

[66570] Koenig, M.; Mandel, J.-L.: Deciphering the cause of

Friedreich
[66571] ataxia. Curr. Opin. Neurobiol. 7: 689-694, 1997.

[66572] 11277.

[66573] Montermini, L.; Rodius, F.; Pianese, L.; Molto, M. D.; Cossee,
[66574] M.; Campuzano, V.; Cavalcanti, F.; Monticelli, A.; Palau, F.; Gyapay,
[66575] G.; Wenhert, M.; Zara, F.; Patel, P. I.; Cocozza, S.; Koenig, M.;
[66576] Pandolfo, M.: The Friedreich ataxia critical region spans a 150-kb
[66577] interval on chromosome 9q13. Am. J. Hum. Genet. 57: 1061-1067, 1995.

[66578] 11278.

[66579] Pandolfo, M.: Personal Communication. Montreal, Canada 4/28/1997.

[66580] 11279.

[66581] Ristow, M.; Pfister, M. F.; Yee, A. J.; Schubert, M.; Michael,
[66582] L.; Zhang, C.-Y.; Ueki, K.; Michael, M. D., II; Lowell, B. B.; Kahn,
[66583] C. R.: Frataxin activates mitochondrial energy conversion

and oxidative
[66584] phosphorylation. Proc. Nat. Acad. Sci. 97: 12239-12243, 2000.
[66585] 11280.
[66586] Sakamoto, N.; Chastain, P. D.; Parniewski, P.; Ohshima, K.; Pandolfo,
[66587] M.; Griffith, J. D.; Wells, R. D.: Sticky DNA: self-association properties
[66588] of long GAA-TTC repeats in R-R-Y triplex structures from Friedrich's
[66589] ataxia. Molec. Cell 3: 465-475, 1999.
[66590] 11281.
[66591] Santos, M. M.; Ohshima, K.; Pandolfo, M.: Frataxin defi-ciency
[66592] enhances apoptosis in cells differentiating into neuroecto-derm. Hum.
[66593] Molec. Genet. 10: 1935-1944, 2001.
[66594] 11282.
[66595] Shoichet, S. A.; Baumer, A. T.; Stamenkovic, D.; Sauer, H.; Pfeiffer,
[66596] A. F. H.; Kahn, C. R.; Muller-Wieland, D.; Richter, C.; Ris-

tow, M.

[66597] : Frataxin promotes antioxidant defense in a thiol-dependent manner

[66598] resulting in diminished malignant transformation in vitro. Hum. Molec.

[66599] Genet. 11: 815-821, 2002.

[66600] 11283.

[66601] Wilson, R. B.; Roof, D. M.: Respiratory deficiency due to loss

[66602] of mitochondrial DNA in yeast lacking the frataxin homo-logue. Nature

[66603] Genet. 16: 352-357, 1997.

[66604] 11284.

[66605] Zuhlke, C.; Laccone, F.; Cossee, M.; Kohlschutter, A.; Koenig,

[66606] M.; Schwinger, E.: Mutation of the start codon in the FRDA1 gene:

[66607] linkage analysis of three pedigrees with the ATG to ATT transversion

[66608] points to a unique common ancestor. Hum. Genet. 103: 102-105, 1998.

[66609] 11285.

[66610] Yuan, C.-X.; Ito, M.; Fondell, J. D.; Fu, Z.-Y.; Roeder, R. G.

[66611] : The TRAP220 component of a thyroid hormone receptor-
associated protein
[66612] (TRAP) coactivator complex interacts directly with nuclear
receptors
[66613] in a ligand-dependent fashion. Proc. Nat. Acad. Sci. 95:
7939-7944,
[66614] 1998.

[66615] 11286.

[66616] Zhang, J.; Fondell, J. D.: Identification of mouse TRAP100:
a
[66617] transcriptional coregulatory factor for thyroid hormone
and vitamin
[66618] D receptors. Molec. Endocr. 13: 1130-1140, 1999.

[66619] 11287.

[66620] Csoka, T. B.; Frost, G. I.; Heng, H. H. Q.; Scherer, S. W.;
Mohapatra,
[66621] G.; Stern, R.: The hyaluronidase gene HYAL1 maps to
chromosome 3p21.2-p21.3
[66622] in human and 9F1-F2 in mouse, a conserved candidate
tumor suppressor

[66623] locus. Genomics 48: 63-70, 1998.

[66624] 11288.

[66625] Frost, G. I.; Csoka, T. B.; Wong, T.; Stern, R.: Purification,

[66626] cloning, and expression of human plasma hyaluronidase. Biochem. Biophys.

[66627] Res. Commun. 236: 10-15, 1997.

[66628] 11289.

[66629] Laurent, T. C.; Fraser, J. R. E.: Hyaluronan. FASEB J. 6: 2397-2404,

[66630] 1992.

[66631] 11290.

[66632] Sturany, S.; Van Lint, J.; Muller, F.; Wilda, M.; Hameister, H.;

[66633] Hocker, M.; Brey, A.; Gern, U.; Vandenheede, J.; Gress, T.; Adler,

[66634] G.; Seufferlein, T.: Molecular cloning and characterization of the

[66635] human protein kinase D2: a novel member of the protein kinase D family

[66636] of serine threonine kinases. J. Biol. Chem. 276: 3310-3318, 2001.

[66637] 11291.

[66638] Fischer, W. H.; Schubert, D.: Characterization of a novel platelet-derived

[66639] growth factor-associated protein. J. Neurochem. 66: 2213-2216, 1996.

[66640] 11292.

[66641] Hayashi, A.; Seki, N.; Hattori, A.; Kozuma, S.; Saito, T.: PKC-nu,

[66642] a new member of the protein kinase C family, composes a fourth subfamily

[66643] with PKC-mu. Biochim. Biophys. Acta 1450: 99-106, 1999.

[66644] 11293.

[66645] Brodbeck, J.; Davies, A.; Courtney, J.-M.; Meir, A.; Bala-guero,

[66646] N.; Canti, C.; Moss, F. J.; Page, K. M.; Pratt, W. S.; Hunt, S. P.;

[66647] Barclay, J.; Rees, M.; Dolphin, A. C.: The ducky mutation in Cacna2d2

[66648] results in altered Purkinje cell morphology and is associ-ated with

[66649] the expression of a truncated alpha-2/delta-2 protein

with abnormal
[66650] function. J. Biol. Chem. 277: 7684-7693, 2002.

[66651] 11294.

[66652] Gao, B.; Sekido, Y.; Maximov, A.; Saad, M.; Forgacs, E.; Latif,
[66653] F.; Wei, M. H.; Lerman, M.; Lee, J.-H.; Perez-Reyes, E.; Bezprozvanny,
[66654] I.; Minna, J. D.: Functional properties of a new voltage-dependent
[66655] calcium channel alpha-2/delta auxiliary subunit gene (CACNA2D2). J.
[66656] Biol. Chem. 275: 12237-12242, 2000.

[66657] 11295.

[66658] Angrand, P.-O.; Apiou, F.; Stewart, A. F.; Dutrillaux, B.; Losson,
[66659] R.; Chambon, P.: NSD3, a new SET domain-containing gene, maps to
[66660] 8p12 and is amplified in human breast cancer cell lines. Genomics 74:
[66661] 79-88, 2001.

[66662] 11296.

[66663] Stec, I.; van Ommen, G.-J. B.; den Dunnen, J. T.: WHSC1L1, on
[66664] human chromosome 8p11.2, closely resembles WHSC1 and maps to a duplicated
[66665] region shared with 4p16.3. Genomics 76: 5-8, 2001.

[66666] 11297.

[66667] Corydon, M. J.; Andresen, B. S.; Bross, P.; Kjeldsen, M.; Andreasen,
[66668] P. H.; Eiberg, H.; Kolvraa, S.; Gregersen, N.: Structural or-
ganization
[66669] of the human short-chain acyl-CoA dehydrogenase gene. Mammalian Genome 8:
[66670] 922-926, 1997.

[66671] 11298.

[66672] Van Patten, S. M.; Ng, D. C.; Th'ng, J. P. H.; Angelos, K. L.;

[66673] Smith, A. J.; Walsh, D. A.: Molecular cloning of a rat testis form
[66674] of the inhibitor protein of cAMP-dependent protein ki-
nase. Proc.
[66675] Nat. Acad. Sci. 88: 5383-5387, 1991.

[66676] 11299.

[66677] Lee, D. K.; George, S. R.; Cheng, R.; Nguyen, T.; Liu, Y.; Brown,
[66678] M.; Lynch, K. R.; O'Dowd, B. F.: Identification of four novel human
[66679] G protein-coupled receptors expressed in the brain. Molec. Brain
[66680] Res. 86: 13-22, 2001.

[66681] 11300.

[66682] Cikos, S.; Gregor, P.; Koppel, J.: Cloning of a novel bio-genic
[66683] amine receptor-like G protein-coupled receptor ex-pressed in human
[66684] brain. Biochim. Biophys. Acta 1521: 66-72, 2001.

[66685] 11301.

[66686] Bascom, R. A.; Srinivasan, S.; Nussbaum, R. L.: Identifica-tion
[66687] and characterization of golgin-84, a novel Golgi integral membrane
[66688] protein with a cytoplasmic coiled-coil domain. J. Biol. Chem. 274:
[66689] 2953-2962, 1999. Note: Erratum: J. Biol. Chem 274: 12950 only, 1999.

[66690] 11302.

[66691] Ishizaka, Y.; Ochiai, M.; Tahira, T.; Sugimura, T.; Nagao, M.:

[66692] Activation of the ret-II oncogene without a sequence encoding a transmembrane

[66693] domain and transforming activity of two ret-II oncogene products differing

[66694] in carboxy-termini due to alternative splicing. Oncogene 4: 789-794,

[66695] 1989.

[66696] 11303.

[66697] Klugbauer, S.; Demidchik, E. P.; Lengfelder, E.; Rabes, H. M.:

[66698] Detection of a novel type of RET rearrangement (PTC5) in thyroid carcinomas

[66699] after Chernobyl and analysis of the involved RET-fused gene RFG5. Cancer

[66700] Res. 58: 198-203, 1998.

[66701] 11304.

[66702] Jiang, J. C.; Kirchman, P. A.; Zagulski, M.; Hunt, J.; Jazwinski,

[66703] S. M.: Homologs of the yeast longevity gene LAG1 in Caenorhabditis
[66704] elegans and human. Genome Res. 8: 1259-1272, 1998.

[66705] 11305.

[66706] Pan, H.; Qin, W.-X.; Huo, K.-K.; Wan, D.-F.; Yu, Y.; Xu, Z.-G.;
[66707] Hu, Q.-D.; Gu, K. T.; Zhou, X.-M.; Jiang, H.-Q.; Zhang, P.-P.; Huang,
[66708] Y.; Li, Y.-Y.; Gu, J.-R.: Cloning, mapping, and characterization
[66709] of a human homologue of the yeast longevity assurance gene LAG1. Genomics 77:
[66710] 58-64, 2001.

[66711] 11306.

[66712] Koivisto, U.-M.; Hubbard, A. L.; Mellman, I.: A novel cellular
[66713] phenotype for familial hypercholesterolemia due to a defect in polarized
[66714] targeting of LDL receptor. Cell 105: 575-585, 2001.

[66715] 11307.

[66716] Koivisto, U.-M.; Turtola, H.; Aalto-Setala, K.; Top, B.;

Frants,
[66717] R. R.; Kovanen, P. T.; Syvanen, A.-C.; Kontula, K.: The familial
[66718] hypercholesterolemia (FH)-North Karelia mutation of the low density
[66719] lipoprotein receptor gene deletes seven nucleotides of exon 6 and
[66720] is a common cause of FH in Finland. J. Clin. Invest. 90: 219-228,
[66721] 1992.

[66722] 11308.

[66723] Koivisto, U.-M.; Viikari, J. S.; Kontula, K.: Molecular characterization
[66724] of minor gene rearrangements in Finnish patients with heterozygous
[66725] familial hypercholesterolemia: identification of two common missense
[66726] mutations (Gly823-to-Asp and Leu380-to-His) and eight rare mutations
[66727] of the LDL receptor gene. Am. J. Hum. Genet. 57: 789-797, 1995.
[66728] 11309.

[66729] Kotze, M. J.; Langenhoven, E.; Warnich, L.; du Plessis, L.; Retief,
[66730] A. E.: The molecular basis and diagnosis of familial hypercholesterolaemia
[66731] in South African Afrikaners. Ann. Hum. Genet. 55: 115-121, 1991.
[66732] 11310.
[66733] Kotze, M. J.; Theart, L.; Peeters, A.; Langenhoven, E.: A de
[66734] novo duplication in the low density lipoprotein receptor gene. Hum.
[66735] Mutat. 6: 181-183, 1995.
[66736] 11311.
[66737] Kotze, M. J.; Warnich, L.; Langenhoven, E.; du Plessis, L.; Retief,
[66738] A. E.: An exon 4 mutation identified in the majority of South African
[66739] familial hypercholesterolaemics. J. Med. Genet. 27: 298-302, 1990.
[66740] 11312.
[66741] Landsberger, D.; Meiner, V.; Reshef, A.; Levy, Y.; van der Westhuyzen,

[66742] D. R.; Coetzee, G. A.; Leitersdorf, E.: A nonsense mutation in the
[66743] LDL receptor gene leads to familial hypercholesterolemia in the Druze
[66744] sect. Am. J. Hum. Genet. 50: 427-433, 1992.

[66745] 11313.

[66746] Langlois, S.: Personal Communication. Vancouver, British Columbia,
[66747] Canada 1989.

[66748] 11314.

[66749] Langlois, S.; Kastelein, J. J. P.; Hayden, M. R.: Characterization
[66750] of six partial deletions in the low-density-lipoprotein (LDL) receptor
[66751] gene causing familial hypercholesterolemia (FH). Am. J. Hum. Genet. 43:
[66752] 60-68, 1988.

[66753] 11315.

[66754] Lee, W. K.; Haddad, L.; Macleod, M. J.; Dorrance, A. M.; Wilson,
[66755] D. J.; Gaffney, D.; Dominiczak, M. H.; Packard, C. J.; Day, I.

N.;

[66756] Humphries, S. E.; Dominiczak, A. F.: Identification of a common low

[66757] density lipoprotein receptor mutation (C163Y) in the West of Scotland. J.

[66758] Med. Genet. 35: 573-578, 1998.

[66759] 11316.

[66760] Lehrman, M. A.; Goldstein, J. L.; Russell, D. W.; Brown, M. S.

[66761] : Duplication of seven exons in LDL receptor gene caused by Alu-Alu

[66762] recombination in a subject with familial hypercholes-terolemia. Cell 48:

[66763] 827-835, 1987.

[66764] 11317.

[66765] Lehrman, M. A.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.

[66766] : Alu-Alu recombination deletes splice acceptor sites and produces

[66767] secreted low density lipoprotein receptor in a subject with familial

[66768] hypercholesterolemia. J. Biol. Chem. 262: 3354-3361,

1987.

[66769] 11318.

[66770] Lehrman, M. A.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.

[66771] : Exon-Alu recombination deletes 5 kilobases from the low density

[66772] lipoprotein receptor gene, producing a null phenotype in familial

[66773] hypercholesterolemia. Proc. Nat. Acad. Sci. 83: 3679-3683, 1986.

[66774] 11319.

[66775] Lehrman, M. A.; Schneider, W. J.; Sudhof, T. C.; Brown, M. S.;

[66776] Goldstein, J. L.; Russell, D. W.: Mutation in LDL receptor: Alu-Alu

[66777] recombination deletes exons encoding transmembrane and cytoplasmic

[66778] domains. Science 227: 140-146, 1985.

[66779] 11320.

[66780] Leitersdorf, E.; Hobbs, H. H.: Personal Communication. Dallas,

[66781] Tex. 12/1990.

[66782] 11321.

[66783] Leitersdorf, E.; Hobbs, H. H.; Fourie, A. M.; Jacobs, M.; van

[66784] der Westhuyzen, D. R.; Coetzee, G. A.: Deletion in the first cysteine-rich

[66785] repeat of low-density lipoprotein receptor impairs its transport but

[66786] not lipoprotein binding in fibroblasts from a subject with familial

[66787] hypercholesterolemia. Proc. Nat. Acad. Sci. 85: 7912-7916, 1988.

[66788] 11322.

[66789] Leitersdorf, E.; Tobin, E. J.; Davignon, J.; Hobbs, H. H.: Common

[66790] low-density lipoprotein receptor mutations in the French Canadian

[66791] population. J. Clin. Invest. 85: 1014-1023, 1990.

[66792] 11323.

[66793] Leitersdorf, E.; van der Westhuyzen, D. R.; Coetzee, G. A.; Hobbs,

[66794] H. H.: Two common low density lipoprotein receptor gene mutations

[66795] cause familial hypercholesterolemia in Afrikaners. J. Clin. Invest. 84:
[66796] 954-961, 1989.

[66797] 11324.

[66798] Lelli, N.; Ghisellini, M.; Calandra, S.; Gaddi, A.; Ciarrocchi,

[66799] A.; Coviello, D. A.; Bertolini, S.: Duplication of exons 13, 14 and
[66800] 15 of the LDL-receptor gene in a patient with heterozygous familial
[66801] hypercholesterolemia. Hum. Genet. 86: 359-362, 1991.

[66802] 11325.

[66803] Leren, T. P.; Solberg, K.; Rodningen, O. K.; Tonstad, S.; Ose,
[66804] L.: Two founder mutations in the LDL receptor gene in Norwegian familial
[66805] hypercholesterolemia subjects. Atherosclerosis 111: 175-182, 1994.

[66806] 11326.

[66807] Li, H.; Gyllensten, U. B.; Cui, X.; Saiki, R. K.; Erlich, H. A.;

[66808] Arnheim, N.: Amplification and analysis of DNA sequences in single

[66809] human sperm and diploid cells. Nature 335: 414-417, 1988.

[66810] 11327.

[66811] Iiizumi, M.; Arakawa, H.; Mori, T.; Ando, A.; Nakamura, Y.: Isolation

[66812] of a novel gene, CABC1, encoding a mitochondrial protein that is highly

[66813] homologous to yeast activity of bc1 complex. Cancer Res. 62: 1246-1250,

[66814] 2002.

[66815] 11328.

[66816] Downes, G. B.; Gilbert, D. J.; Copeland, N. G.; Gautam, N.; Jenkins,

[66817] N. A.: Chromosomal mapping of five mouse G protein gamma subunits. Genomics 57:

[66818] 173-176, 1999.

[66819] 11329.

[66820] Modarressi, M. H.; Taylor, K. E.; Wolfe, J.: Cloning, charac-terization,

[66821] and mapping of the gene encoding the human G protein gamma-2 subunit. Biochem.

[66822] Biophys. Res. Commun. 272: 610-615, 2000.

[66823] 11330.

[66824] Yu, Y.; Zhang, C.; Zhou, G.; Wu, S.; Qu, X.; Wei, H.; Xing, G.;

[66825] Dong, C.; Zhai, Y.; Wan, J.; Ouyang, S.; Li, L., Zhang, S.; Zhou,

[66826] K.; Zhang, Y.; Wu, C.; He, F.: Gene expression profiling in human

[66827] fetal liver and identification of tissue- and developmental-stage-specific

[66828] genes through compiled expression profiles and efficient cloning of

[66829] full-length cDNAs. Genome Res. 11: 1392-1403, 2001.

[66830] 11331.

[66831] Ericsson, J.; Greene, J. M.; Carter, K. C.; Shell, B. K.; Duan,

[66832] D. R.; Florence, C.; Edwards, P. A.: Human geranylgeranyl diphosphate

[66833] synthase: isolation of the cDNA, chromosomal mapping and tissue expression. J.

[66834] Lipid Res. 39: 1731-1739, 1998.

[66835] 11332.

[66836] Kainou, T.; Kawamura, K.; Tanaka, K.; Matsuda, H.; Kawa-
mukai, M.
[66837] : Identification of the GGPS1 genes encoding geranylger-
anyl diphosphate
[66838] synthases from mouse and human. Biochim. Biophys. Acta
1437: 333-340,
[66839] 1999.

[66840] 11333.

[66841] Kuzuguchi, T.; Morita, Y.; Sagami, I.; Sagami, H.; Ogura,
K.:
[66842] Human geranylgeranyl diphosphate synthase: cDNA
cloning and expression. J.
[66843] Biol. Chem. 274: 5888-5894, 1999.

[66844] 11334.

[66845] Vicent, D.; Maratos-Flier, E.; Kahn, C. R.: The branch point
enzyme
[66846] of the mevalonate pathway for protein prenylation is over-
expressed
[66847] in the ob/ob mouse and induced by adipogenesis. Molec.
Cell. Biol. 20:
[66848] 2158-2166, 2000.

[66849] 11335.

[66850] Cases, S.; Stone, S. J.; Zhou, P.; Yen, E.; Tow, B.; Lardiza-
bal,

[66851] K. D.; Voelker, T.; Farese, R. V., Jr.: Cloning of DGAT2, a
second

[66852] mammalian diacylglycerol acyltransferase, and related
family members. J.

[66853] Biol. Chem. 276: 38870-38876, 2001.

[66854] 11336.

[66855] Chen, C.-K.; Zhang, K.; Church-Kopish, J.; Huang, W.;
Zhang, H.;

[66856] Chen, Y.-J.; Frederick, J. M.; Baehr, W.: Characterization of
human

[66857] GRK7 as a potential cone opsin kinase. Molec. Vision 7:
305-313,

[66858] 2001.

[66859] 11337.

[66860] Weiss, E. R.; Ducceschi, M. H.; Horner, T. J.; Li, A.; Craft, C.

[66861] M.; Osawa, S.: Species-specific differences in expression
of G-protein-coupled

[66862] receptor kinase (GRK) 7 and GRK1 in mammalian cone

photoreceptor cells:
[66863] implications for cone cell phototransduction. J. Neurosci. 21: 9175-9184,
[66864] 2001.

[66865] 11338.

[66866] Borregaard, N.; Cowland, J. B.: Granules of the human neutrophilic
[66867] polymorphonuclear leukocyte. Blood 89: 3503-3521, 1997.
[66868] 11339.

[66869] Chang, K. S.; Schroeder, W.; Siciliano, M. J.; Thompson, L. H.;
[66870] McCredie, K.; Beran, M.; Freireich, E. J.; Liang, J. C.; Trujillo,
[66871] J. M.; Stass, S. A.: The localization of the human myeloperoxidase
[66872] gene is in close proximity to the translocation breakpoint in acute
[66873] promyelocytic leukemia. Leukemia 1: 458-462, 1987.

[66874] 11340.

[66875] DeLeo, F. R.; Goedken, M.; McCormick, S. J.; Nauseef, W.

M.: A

[66876] novel form of hereditary myeloperoxidase deficiency linked to endoplasmic

[66877] reticulum/proteasome degradation. J. Clin. Invest. 101: 2900-2909,

[66878] 1998.

[66879] 11341.

[66880] Eiserich, J. P.; Baldus, S.; Brennan, M.-L.; Ma, W.; Zhang, C.;

[66881] Tousson, A.; Castro, L.; Lusis, A. J.; Nauseef, W. M.; White, C. R.;

[66882] Freeman, B. A.: Myeloperoxidase, a leukocyte-derived vascular NO

[66883] oxidase. Science 296: 2391-2394, 2002.

[66884] 11342.

[66885] Inazawa, J.; Inoue, K.; Nishigaki, H.; Tsuda, S.; Taniwaki, M.;

[66886] Misawa, S.; Abe, T.: Assignment of the human myeloper- oxidase gene

[66887] (MPO) to bands q21.3-q23 of chromosome 17. Cytogenet. Cell Genet. 50:

[66888] 135-136, 1989.

[66889] 11343.

[66890] Johnson, K.; Gemperlein, I.; Hudson, S.; Shane, S.; Rovera, G.

[66891] : Complete nucleotide sequence of the human myeloperoxidase gene. Nucleic

[66892] Acids Res. 17: 7985-7986, 1989.

[66893] 11344.

[66894] Kizaki, M.; Miller, C. W.; Selsted, M. E.; Koeffler, H. P.: Myeloperoxidase

[66895] (MPO) gene mutation in hereditary MPO deficiency. Blood 83: 1935-1940,

[66896] 1994.

[66897] 11345.

[66898] Klebanoff, S. J.: Myeloperoxidase. Proc. Assoc. Am. Physicians 111:

[66899] 383-389, 1999.

[66900] 11346.

[66901] Kudoh, J.; Minoshima, S.; Hashinaka, K.; Nishio, C.; Yamada, M.;

[66902] Shimizu, Y.; Shimizu, N.: Assignment of the myeloperoxidase gene

[66903] MPO to human chromosome 17 using somatic cell hybrids and flow-sorted

[66904] chromosomes. Jpn. J. Hum. Genet. 33: 315-324, 1988.

[66905] 11347.

[66906] Kudoh, J.; Minoshima, S.; Hashinaka, K.; Nishio, C.; Ya-mada, M.;

[66907] Shimizu, Y.; Shimizu, N.: Assignment of the myeloperoxi-dase (MPO)

[66908] gene to human chromosome 17. (Abstract) Cytogenet. Cell Genet. 46:

[66909] 641-642, 1987.

[66910] 11348.

[66911] Law, D. J.; Prasad, M. A.; King, S. E.; Spranger, K. D.; Lee,

[66912] Y. H.; Fox, R. E.; Collins, E. E.; Gebuhr, T. C.; Miller, D. E.; Petty,

[66913] E. M.: Localization of the human estrogen-responsive fin-ger protein

[66914] (EFP) gene (ZNF147) within a YAC contig containing the myeloperoxidase

[66915] (MPO) gene. Genomics 28: 361-363, 1995.

[66916] 11349.

[66917] Williams, M.; Lyu, M.-S.; Yang, Y.-L.; Lin, E. P.; Dunbrack, R.;
[66918] Birren, B.; Cunningham, J.; Hunter, K.: Ier5, a novel member of the
[66919] slow-kinetics immediate-early genes. Genomics 55: 327-334, 1999.
[66920] 11350.
[66921] Kiss, H.; Yang, Y.; Kiss, C.; Andersson, K.; Klein, G.; Imreh,
[66922] S.; Dumanski, J. P.: The transcriptional map of the common eliminated
[66923] region 1 (C3CER1) in 3p21.3. Europ. J. Hum. Genet. 10: 52-61, 2002.
[66924] 11351.
[66925] Arce, I.; Roda-Navarro, P.; Montoya, M. C.; Hernanz-Falcon, P.;
[66926] Puig-Kroger, A.; Fernandez-Ruiz, E.: Molecular and genomic characterization
[66927] of human DLEC, a novel member of the C-type lectin receptor gene family
[66928] preferentially expressed on monocyte-derived dendritic cells. Europ.
[66929] J. Immun. 31: 2733-2740, 2001.

[66930] 11352.

[66931] Dzionek, A.; Sohma, Y.; Nagafune, J.; Cella, M.; Colonna, M.; Facchetti,

[66932] F.; Gunther, G.; Johnston, I.; Lanzavecchia, A.; Nagasaka, T.; Okada,

[66933] T.; Vermi, W.; Winkels, G.; Yamamoto, T.; Zysk, M.; Yamaguchi, Y.;

[66934] Schmitz, J.: BDCA-2, a novel plasmacytoid dendritic cell-specific

[66935] type II C-type lectin, mediates antigen capture and is a potent inhibitor

[66936] of interferon alpha/beta induction. J. Exp. Med. 194: 1823-1834,

[66937] 2001.

[66938] 11353.

[66939] Antonarakis, S. E.: Personal Communication. Baltimore, Md.

[66940] 3/25/2002.

[66941] 11354.

[66942] McKemy, D. D.; Neuhausser, W. M.; Julius, D.: Identification of

[66943] a cold receptor reveals a general role for TRP channels in thermosensation. Nature 416:
[66944] 52-58, 2002.

[66945] 11355.

[66946] Peier, A. M.; Moqrich, A.; Hergarden, A. C.; Reeve, A. J.; Andersson,
[66947] D. A.; Story, G. M.; Earley, T. J.; Dragoni, I.; McIntyre, P.; Bevan,
[66948] S.; Patapoutian, A.: A TRP channel that senses cold stimuli and menthol. Cell 108:
[66949] 705-715, 2002.

[66950] 11356.

[66951] Feng, B.-J.; Huang, W.; Shugart, Y. Y.; Lee, M. K.; Zhang, F.;
[66952] Xia, J.-C.; Wang, H.-Y.; Huang, T.-B.; Jian, S.-W.; Huang, P.; Feng,
[66953] Q.-S.; Huang, L.-X.; and 18 others: Genome-wide scan for familial
[66954] nasopharyngeal carcinoma reveals evidence of linkage to chromosome
[66955] 4. Nature Genet. 31: 395-399, 2002.

[66956] 11357.

[66957] Ooi, E. E.; Ren, E. C.; Chan, S. H.: Association between microsatellites

[66958] within the human MHC and nasopharyngeal carcinoma. Int. J. Cancer 74:

[66959] 229-232, 1997.

[66960] 11358.

[66961] Roginski, R. S.; Mohan Raj, B. K.; Finkernagel, S. W.; Sciorra,

[66962] L. J.: Assignment of an ionotropic glutamate receptor-like gene (GRINL1A)

[66963] to human chromosome 15q22.1 by in situ hybridization. Cytogenet.

[66964] Cell Genet. 93: 143-144, 2001.

[66965] 11359.

[66966] Wydner, K. S.; Mohan Raj, B. K.; Sciorra, L. J.; Roginski, R. S.

[66967] : The mouse orthologue of the human ionotropic glutamate receptor-like

[66968] gene (GRINL1A) maps to mouse chromosome 9. Cytogenet. Cell Genet. 95:

[66969] 240-241, 2001.

[66970] 11360.

[66971] Vuoristo, J. T.; Berrettini, W. H.; Ala-Kokko, L.: C18orf2, a

[66972] novel, highly conserved intronless gene within intron 5 of the GNAL

[66973] gene on chromosome 18p11. Cytogenet. Cell Genet. 93: 19-22, 2001.

[66974] 11361.

[66975] Akiyama, H.; Hiraki, Y.; Noda, M.; Shigeno, C.; Ito, H.; Nakamura,

[66976] T.: Molecular cloning and biological activity of a novel Ha-Ras suppressor

[66977] gene predominantly expressed in skeletal muscle, heart, brain, and

[66978] bone marrow by differential display using clonal mouse EC cells, ATDC5. J.

[66979] Biol. Chem. 274: 32192-32197, 1999.

[66980] 11362.

[66981] Ito, H.; Akiyama, H.; Shigeno, C.; Nakamura, T.: Isolation, characterization,

[66982] and chromosome mapping of a human A-C1 Ha-Ras sup-

pressor gene (HRASLS). Cytogenet.
[66983] Cell Genet. 93: 36-39, 2001.

[66984] 11363.

[66985] Nakamura, T.; Yamazaki, Y.; Saiki, Y.; Moriyama, M.;
Largaespada,
[66986] D. A.; Jenkins, N. A.; Copeland, N. G.: Evi9 encodes a novel
zinc
[66987] finger protein that physically interacts with BCL6, a known
human
[66988] B-cell proto-oncogene product. Molec. Cell Biol. 20:
3178-3186,
[66989] 2000.

[66990] 11364.

[66991] Saiki, Y.; Yamazaki, Y.; Yoshida, M.; Katoh, O.; Nakamura,
T.:
[66992] Human EVI9, a homologue of the mouse myeloid leukemia
gene, is expressed
[66993] in the hematopoietic progenitors and down-regulated
during myeloid
[66994] differentiation of HL60 cells. Genomics 70: 387-391,
2000.
[66995] 11365.

[66996] Satterwhite, E.; Sonoki, T.; Willis, T. G.; Harder, L.; Nowak,

[66997] R.; Arriola, E. L.; Liu, H.; Price, H. P.; Gesk, S.; Steinemann, D.;

[66998] Schlegelberger, B.; Oscier, D. G.; Siebert, R.; Tucker, P. W.; Dyer,

[66999] M. J. S.: The BCL11 gene family: involvement of BCL11A in lymphoid

[67000] malignancies. Blood 98: 3413-3420, 2001.

[67001] 11366.

[67002] Frohlich, O.; Po, C.; Young, L. G.: Organization of the human

[67003] gene encoding the epididymis-specific EP2 protein variants and its

[67004] relationship to defensin genes. Biol. Reprod. 64: 1072-1079, 2001.

[67005] 11367.

[67006] Hamil, K. G.; Sivashanmugam, P.; Richardson, R. T.; Grossman, G.;

[67007] Ruben, S. M.; Mohler, J. L.; Petrusz, P.; O'rand, M. G.; French, F.

[67008] S.; Hall, S. H.: HE2-beta and HE2-gamma, new members of an epididymis-specific

[67009] family of androgen-regulated proteins in the human. Endocrinology 141:

[67010] 1245-1253, 2000.

[67011] 11368.

[67012] Jia, H. P.; Schutte, B. C.; Schudy, A.; Linzmeier, R.; Guthmiller,

[67013] J. M.; Johnson, G. K.; Tack, B. F.; Mitros, J. P.; Rosenthal, A.;

[67014] Ganz, T.; McCray, P. B., Jr.: Discovery of new human beta-defensins

[67015] using a genomics-based approach. Gene 263: 211-218, 2001.

[67016] 11369.

[67017] Li, P.; Chan, H. C.; He, B.; So, S. C.; Chung, Y. W.; Shang, Q.;

[67018] Zhang, Y.-D.; Zhang, Y.-L.: An antimicrobial peptide gene found in

[67019] the male reproductive system of rats. Science 291: 1783-1785, 2001.

[67020] 11370.

[67021] Osterhoff, C.; Kirchhoff, C.; Krull, N.; Ivell, R.: Molecular

[67022] cloning and characterization of a novel human sperm antigen (HE2)

[67023] specifically expressed in the proximal epididymis. Biol. Reprod. 50:

[67024] 516-525, 1994.

[67025] 11371.

[67026] Xu, L. L.; Shanmugam, N.; Segawa, T.; Sesterhenn, I. A.; McLeod,

[67027] D. G.; Moul, J. W.; Srivastava, S.: A novel androgen-regulated gene,

[67028] PMEPA1, located on chromosome 20q13 exhibits high level expression

[67029] in prostate. Genomics 66: 257-263, 2000.

[67030] 11372.

[67031] Wice, B. M.; Gordon, J. I.: A tetraspan membrane glycoprotein

[67032] produced in the human intestinal epithelium and liver that can regulate

[67033] cell density-dependent proliferation. J. Biol. Chem. 270: 21907-21918,

[67034] 1995.

[67035] 11373.

[67036] Kiss, H.; Kedra, D.; Kiss, C.; Kost-Alimova, M.; Yang, Y.; Klein,

[67037] G.; Imreh, S.; Dumanski, J. P.: The LZTFL1 gene is a part of a transcriptional

[67038] map covering 250 kb within the common eliminated region 1 (C3CER1)

[67039] in 3p21.3. Genomics 73: 10-19, 2001.

[67040] 11374.

[67041] Kingsbury, G. A.; Feeney, L. A.; Nong, Y.; Calandra, S. A.; Murphy,

[67042] C. J.; Corcoran, J. M.; Wang, Y.; Das, M. R. P.; Busfield, S. J.;

[67043] Fraser, C. C.; Villeval, J. L.: Cloning, expression, and function

[67044] of BLAME, a novel member of the CD2 family. J. Immun. 166: 5675-5680,

[67045] 2001.

[67046] 11375.

[67047] Quentmeier, H.; Drexler, H. G.; Fleckenstein, D.; Zaborski, M.;

[67048] Armstrong, A.; Sims, J. E.; Lyman, S. D.: Cloning of human

thymic

[67049] stromal lymphopoietin (TSLP) and signaling mechanisms leading to proliferation. Leukemia 15:

[67050] 1286-1292, 2001.

[67051] 11376.

[67052] Soumelis, V.; Reche, P. A.; Kanzler, H.; Yuan, W.; Edward, G.;

[67053] Homey, B.; Gilliet, M.; Ho, S.; Antonenko, S.; Lauerma, A.; Smith,

[67054] K.; Gorman, D.; Zurawski, S.; Abrams, J.; Menon, S.; Mc-Clanahan, T.;

[67055] de Waal-Malefyt, R.; Bazan, F.; Kastelein, R. A.; Liu, Y.-J.: Human

[67056] epithelial cells trigger dendritic cell-mediated allergic inflammation

[67057] by producing TSLP. Nature Immun. 3: 673-680, 2002.

[67058] 11377.

[67059] Gubitz, A. K.; Mourelatos, Z.; Abel, L.; Rappsilber, J.; Mann,

[67060] M.; Dreyfuss, G.: Gemin5, a novel WD repeat protein component of

[67061] the SMN complex that binds Sm proteins. J. Biol. Chem.

277: 5631-5636,

[67062] 2002.

[67063] 11378.

[67064] Guo, D.; Hasham, S.; Kuang, S.-Q.; Vaughan, C. J.; Boer-
winkle,

[67065] E.; Chen, H.; Abuelo, D.; Dietz, H. C.; Basson, C. T.; Shete,
S. S.;

[67066] Milewicz, D. M.: Familial thoracic aortic aneurysms and
dissections:

[67067] genetic heterogeneity with a major locus mapping to
5q13-14. Circulation 103:

[67068] 2461-2468, 2001.

[67069] 11379.

[67070] Vaughan, C. J.; Casey, M.; He, J.; Veuglers, M.; Henderson,
K.;

[67071] Guo, D.; Campagna, R.; Roman, M. J.; Milewicz, D. M.; De-
vereux, R.

[67072] B.; Basson, C. T.: Identification of a chromosome
11q23.2-q24 locus

[67073] for familial aortic aneurysm disease, a genetically hetero-
geneous

[67074] disorder. Circulation 103: 2469-2475, 2001.

[67075] 11380.

[67076] Xia, C.; Bao, Z.; Tabassam, F.; Ma, W.; Qiu, M.; Hua, S.; Liu,

[67077] M.: GCIP, a novel human Grap2 and cyclin D interacting protein, regulates

[67078] E2F-mediated transcriptional activity. J. Biol. Chem. 275: 20942-20948,

[67079] 2000.

[67080] 11381.

[67081] Liu, H.; Sugiura, M.; Nava, V. E.; Edsall, L. C.; Kono, K.; Poulton,

[67082] S.; Milstien, S.; Kohama, T.; Spiegel, S.: Molecular cloning and

[67083] functional characterization of a novel mammalian sphingosine kinase

[67084] type 2 isoform. J. Biol. Chem. 275: 19513-19520, 2000.

[67085] 11382.

[67086] Ballinger, C. A.; Connell, P.; Wu, Y.; Hu, Z.; Thompson, L. J.;

[67087] Yin, L.-Y.; Patterson, C.: Identification of CHIP, a novel tetratricopeptide

[67088] repeat-containing protein that interacts with heat shock

proteins

[67089] and negatively regulates chaperone functions. Molec. Cell. Biol. 19:

[67090] 4535-4545, 1999.

[67091] 11383.

[67092] Jiang, J.; Ballinger, C. A.; Wu, Y.; Dai, Q.; Cyr, D. M.; Ho-hfeld,

[67093] J.; Patterson, C.: CHIP is a U-box-dependent E3 ubiquitin ligase:

[67094] identification of Hsc70 as a target for ubiquitylation. J. Biol.

[67095] Chem. 276: 42938-42944, 2001.

[67096] 11384.

[67097] Gaide, O.; Martinon, F.; Micheau, O.; Bonnet, D.; Thome, M.; Tschopp,

[67098] J.: Carma1, a CARD-containing binding partner of Bcl10, induces Bcl10

[67099] phosphorylation and NF-kappa-B activation. FEBS Lett. 496: 121-127,

[67100] 2001.

[67101] 11385.

[67102] McAllister-Lucas, L. M.; Inohara, N.; Lucas, P. C.; Ruland, J.;

[67103] Benito, A.; Li, Q.; Chen, S.; Chen, F. F.; Yamaoka, S.; Verma, I.

[67104] M.; Mak, T. W.; Nunez, G.: Bimp1, a MAGUK family member linking protein

[67105] kinase C activation to Bcl10-mediated NF-kappa-B induction. J. Biol.

[67106] Chem. 276: 30589-30597, 2001.

[67107] 11386.

[67108] Wang, L.; Guo, Y.; Huang, W.-J.; Ke, X.; Poyet, J.-L.; Manji, G.

[67109] A.; Merriam, S.; Glucksmann, M. A.; DiStefano, P. S.; Alnemri, E.

[67110] S.; Bertin, J.: CARD10 is a novel caspase recruitment domain/membrane-associated

[67111] guanylate kinase family member that interacts with BCL10 and activates

[67112] NF-kappa-B. J. Biol. Chem. 276: 21405-21409, 2001.

[67113] 11387.

[67114] Gorlich, D.; Mattaj, I. W.: Nucleocytoplasmic transport. Science 271:

[67115] 1513-1518, 1996.

[67116] 11388.

[67117] Zhang, X.; Yang, H.; Corydon, M. J.; Zhang, X.; Pedersen, S.; Korenberg,

[67118] J. R.; Chen, X.-N.; Laporte, J.; Gregersen, N.; Niebuhr, E.; Liu,

[67119] G.; Bolund, L.: Localization of a human nucleoporin 155 gene (NUP155)

[67120] to the 5p13 region and cloning of its cDNA. Genomics 57: 144-151,

[67121] 1999.

[67122] 11389.

[67123] Lager, D. J.; Qian, Q.; Bengal, R. J.; Ishibashi, M.; Torres, V.

[67124] E.: The pck rat: a new model that resembles human autosomal dominant

[67125] polycystic kidney and liver disease. Kidney Int. 59: 126-136, 2001.

[67126] 11390.

[67127] Sanzen, T.; Harada, K.; Yasoshima, M.; Kawamura, Y.; Ishibashi,

[67128] M.; Nakanuma, Y.: Polycystic kidney rat is a novel animal

model of
[67129] Caroli's disease associated with congenital hepatic fibrosis. Am.
[67130] J. Path. 158: 1605-1612, 2001.

[67131] 11391.

[67132] Ward, C. J.; Hogan, M. C.; Rossetti, S.; Walker, D.; Sneddon, T.;
[67133] Wang, X.; Kubly, V.; Cunningham, J. M.; Bacallao, R.; Ishibashi, M.;
[67134] Milliner, D. S.; Torres, V. E.; Harris, P. C.: The gene mutated in
[67135] autosomal recessive polycystic kidney disease encodes a large, receptor-like
[67136] protein. Nature Genet. 30: 259-269, 2002.

[67137] 11392.

[67138] Takeuchi, T.; Shuman, M. A.; Craik, C. S.: Reverse biochemistry:
[67139] use of macromolecular protease inhibitors to dissect complex biological
[67140] processes and identify a membrane-type serine protease in epithelial
[67141] cancer and normal tissue. Proc. Nat. Acad. Sci. 96:

11054-11061,

[67142] 1999.

[67143] 11393.

[67144] Damiano, J. S.; Stehlik, C.; Pio, F.; Godzik, A.; Reed, J. C.:

[67145] CLAN, a novel human CED-4-like gene. Genomics 75:
77-83, 2001.

[67146] 11394.

[67147] Loux, N.; Benlian, P.; Pastier, D.; Boileau, C.; Cambou, J. P.;

[67148] Monnier, L.; Percheron, C.; Junien, C.: Recurrent mutation
at aa

[67149] 792 in the LDL receptor gene in a French patient. Hum.
Genet. 87:

[67150] 373-375, 1991.

[67151] 11395.

[67152] Le Beau, M. M.; Lemons, R. S.; Rosner, G. L.; Carrino, J. C.;

[67153] Reid, M. S.; Chisholm, R. L.; Diaz, M. O.; Weil, S. C.: Chro-
mosomal

[67154] localization of the gene encoding myeloperoxidase.
(Abstract) Cytogenet.

[67155] Cell Genet. 46: 645, 1987.

[67156] 11396.

[67157] De Deken, X.; Wang, D.; Many, M.-C.; Costagliola, S.; Libert, F.;

[67158] Vassart, G.; Dumont, J. E.; Miot, F.: Cloning of two human thyroid

[67159] cDNAs encoding new members of the NADPH oxidase family. J. Biol.

[67160] Chem. 275: 23227-23233, 2000.

[67161] 11397.

[67162] Lacroix, L.; Nocera, M.; Mian, C.; Caillou, B.; Virion, A.; Dupuy,

[67163] C.; Filetti, S.; Bidart, J. M.; Schlumberger, M.: Expression of nicotinamide

[67164] adenine dinucleotide phosphate oxidase flavoprotein DUOX genes and

[67165] proteins in human papillary and follicular thyroid carcinomas. Thyroid 11:

[67166] 1017-1023, 2001.

[67167] 11398.

[67168] Dupuy, C.; Ohayon, R.; Valent, A.; Noel-Hudson, M.-S.; Deme, D.;

[67169] Virion, A.: Purification of a novel flavoprotein involved in the

[67170] thyroid NADPH oxidase: cloning of the porcine and human cDNAs. J.

[67171] Biol. Chem. 274: 37265-37269, 1999.

[67172] 11399.

[67173] Moreno, J. C.; Bikker, H.; Kempers, M. J. E.; van Trotsen-burg,

[67174] A. S. P.; Baas, F.; de Vijlder, J. J. M.; Vulsma, T.; Ris-Stalpers,

[67175] C.: Inactivating mutations in the gene for thyroid oxidase 2 (THOX2)

[67176] and congenital hypothyroidism. New Eng. J. Med. 347: 95-102, 2002.

[67177] 11400.

[67178] Beratis, N. G.; LaBadie, G. U.; Hirschhorn, K.: An isozyme of

[67179] acid alpha-glucosidase with reduced catalytic activity for glycogen. Am.

[67180] J. Hum. Genet. 32: 137-149, 1980.

[67181] 11401.

[67182] D'Ancona, G. G.; Wurm, J.; Croce, C. M.: Genetics of type II glycogenosis:
[67183] assignment of the human gene for acid alpha-glucosidase to chromosome
[67184] 17. Proc. Nat. Acad. Sci. 76: 4526-4529, 1979.

[67185] 11402.

[67186] Dennis, J. A.; Moran, C.; Healy, P. J.: The bovine alpha-glucosidase
[67187] gene: coding region, genomic structure, and mutations that cause bovine
[67188] generalized glycogenosis. Mammalian Genome 11: 206-212, 2000.
[67189] 11403.

[67190] Halley, D. J. J.; Konings, A.; Hupkes, P.; Galjaard, H.: Re-gional
[67191] mapping of the human gene for lysosomal alpha-glucosidase by in situ
[67192] hybridization. Hum. Genet. 67: 326-328, 1984.

[67193] 11404.

[67194] Hermans, M. M. P.; De Graaff, E.; Kroos, M. A.; Mohkams-ing, S.;

[67195] Eussen, B. J.; Joosse, M.; Willemsen, R.; Kleijer, W. J.; Oostra,
[67196] B. A.; Reuser, A. J. J.: The effect of a single base pair deletion
[67197] (delta-T525) and a C1634T missense mutation (pro545leu) on the expression
[67198] of lysosomal alpha-glucosidase in patients with glycogen storage disease
[67199] type II. Hum. Molec. Genet. 3: 2213-2218, 1994.

[67200] 11405.

[67201] Hermans, M. M. P.; de Graaff, E.; Kroos, M. A.; Wisselaar, H. A.;
[67202] Oostra, B. A.; Reuser, A. J. J.: Identification of a point mutation
[67203] in the human lysosomal alpha-glucosidase gene causing infantile glycogenosis
[67204] type II. Biochem. Biophys. Res. Commun. 179: 919-926, 1991.
[67205] 11406.

[67206] Hermans, M. M. P.; Kroos, M. A.; de Graaff, E.; Oostra, B. A.;
[67207] Reuser, A. J. J.: Two mutations affecting the transport and

maturation
[67208] of lysosomal alpha-glucosidase in an adult case of glycogen storage
[67209] disease type II. Hum. Mutat. 2: 268-273, 1993.

[67210] 11407.

[67211] Hoefsloot, L. H.; Hoogeveen-Westerveld, M.; Kroos, M. A.; van
[67212] Beeumen, J.; Reuser, A. J. J.; Oostra, B. A.: Primary structure and
[67213] processing of lysosomal alpha-glucosidase; homology with the intestinal
[67214] sucrase-isomaltase complex. EMBO J. 7: 1697-1704, 1988.

[67215] 11408.

[67216] Hoefsloot, L. H.; Hoogeveen-Westerveld, M.; Reuser, A. J. J.;
[67217] Oostra, B. A.: Characterization of the human lysosomal alpha-glucosidase
[67218] gene. Biochem. J. 272: 493-497, 1990.

[67219] 11409.

[67220] Huie, M. L.; Chen, A. S.; Tsujino, S.; Shanske, S.; DiMauro,

S.;

[67221] Engel, A. G.; Hirschhorn, R.: Aberrant splicing in adult onset glycogen

[67222] storage disease type II (GSDII): molecular identification of an IVS1

[67223] (-13T to G) mutation in a majority of patients and a novel IVS10 (+GT

[67224] to CT) mutation. Hum. Molec. Genet. 3: 2231-2236, 1994.

[67225] 11410.

[67226] Huie, M. L.; Menaker, M.; McAlpine, P. J.; Hirschhorn, R.: Identification

[67227] of an E689K substitution as the molecular basis of the human acid

[67228] alpha-glucosidase type 4 allozyme (GAA*4). Ann. Hum. Genet. 60:

[67229] 365-368, 1996.

[67230] 11411.

[67231] Ko, T.-M.; Hwu, W.-L.; Lin, Y.-W.; Tseng, L.-H.; Hwa, H.-L.; Wang,

[67232] T.-R.; Chuang, S.-M.: Molecular genetic study of Pompe disease in

[67233] Chinese patients in Taiwan. Hum. Mutat. 13: 380-384,

1999.

[67234] 11412.

[67235] Okajima, T.; Fukumoto, S.; Miyazaki, H.; Ishida, H.; Kiso, M.;

[67236] Furukawa, K.; Urano, T.; Furukawa, K.: Molecular cloning of a novel

[67237] alpha-2,3-sialyltransferase (ST3Gal VI) that sialylates type II lactosamine

[67238] structures on glycoproteins and glycolipids. J. Biol. Chem. 274:

[67239] 11479-11486, 1999.

[67240] 11413.

[67241] Lin, C.-Y.; Shieh, J.-J.: Molecular study on the infantile form

[67242] of Pompe disease in Chinese in Taiwan. Acta Paediat. Sinica 37:

[67243] 115-121, 1996.

[67244] 11414.

[67245] Taniguchi, A.; Kaneta, R.; Morishita, K.; Matsumoto, K.: Gene

[67246] structure and transcriptional regulation of human Gal

beta-1,4(3)

[67247] GlcNac alpha-2,3-sialyltransferase VI (hST3Gal VI) gene in prostate

[67248] cancer cell line. Biochem. Biophys. Res. Commun. 287: 1148-1156,

[67249] 2001.

[67250] 11415.

[67251] Angata, T.; Kerr, S. C.; Greaves, D. R.; Varki, N. M.; Crocker,

[67252] P. R.; Varki, A.: Cloning and characterization of human Siglec-11:

[67253] a recently evolved signaling molecule that can interact with SHP-1

[67254] and SHP-2 and is expressed by tissue macrophages, including brain

[67255] microglia. J. Biol. Chem. 277: 24466-24474, 2002.

[67256] 11416.

[67257] Geddes, B. J.; Wang, L.; Huang, W.-J.; Lavellee, M.; Manji, G.

[67258] A.; Brown, M.; Jurman, M.; Cao, J.; Morgenstern, J.; Merriam, S.;

[67259] Glucksmann, M. A.; DiStefano, P. S.; Bertin, J.: Human

CARD12 is
[67260] a novel CED4/Apaf-1 family member that induces apoptosis. Biochem.
[67261] Biophys. Res. Commun. 284: 77-82, 2001.

[67262] 11417.

[67263] Poyet, J.-L.; Srinivasula, S. M.; Tnani, M.; Razmara, M.; Fernandes-Alnemri,
[67264] T.; Alnemri, E. S.: Identification of Ipaf, a human caspase-1-activating
[67265] protein related to Apaf-1. J. Biol. Chem. 276: 28309-28313, 2001.
[67266] 11418.

[67267] Chano, T.; et al; et al: Isolation, characterization and mapping
[67268] of the mouse and human RB1CC1 genes. Gene (in-press), 2002.
[67269] 11419.

[67270] Chano, T.; Kontani, K.; Teramoto, K.; Okabe, H.; Ikegawa, S.:
[67271] Truncating mutations of RB1CC1 in human breast cancers. Nature Genet. 31:
[67272] 285-288, 2002.

[67273] 11420.

[67274] Paris, M. J.; Williams, B. R. G.: Characterization of a 500-kb
[67275] contig spanning the region between c-Ha-Ras and MUC2 on chromosome
[67276] 11p15.5. Genomics 69: 196-202, 2000.

[67277] 11421.

[67278] Kim, N.; Takami, M.; Rho, J.; Josien, R.; Choi, Y.: A novel member
[67279] of the leukocyte receptor complex regulates osteoclast differentiation. J.
[67280] Exp. Med. 195: 201-209, 2002.

[67281] 11422.

[67282] Wilkinson, B.; Chen, J. Y.-F.; Han, P.; Rufner, K. M.; Goularte,
[67283] O. D.; Kaye, J.: TOX: an HMG box protein implicated in the

regulation
[67284] of thymocyte selection. Nature Immun. 3: 272-280, 2002.

[67285] 11423.

[67286] Kaneko, M.; Kudo, T.; Iwasaki, H.; Ikehara, Y.; Nishihara, S.;
[67287] Nakagawa, S.; Sasaki, K.; Shiina, T.; Inoko, H.; Saitou, N.; Narimatsu,
[67288] H.: Alpha-1,3-fucoslytransferase (sic) IX (Fuc-TIX) is very highly
[67289] conserved between human and mouse; molecular cloning, characterization
[67290] and tissue distribution of human Fuc-TIX. FEBS Lett. 452: 237-242,
[67291] 1999.

[67292] 11424.

[67293] Kaneko, M.; Kudo, T.; Iwasaki, H.; Shiina, T.; Inoko, H.; Kozaki,
[67294] T.; Saitou, N.; Narimatsu, H.: Assignment of the human alpha-1,3-fucosyltransferase
[67295] IX gene (FUT9) to chromosome band 6q16 by in situ hybridization. Cytogenet.
[67296] Cell Genet. 86: 329-330, 1999.

[67297] 11425.

[67298] Kas, K.; Lemahieum V.; Meyen, E.; Van de Ven, W. J.; Mer-
regaert,

[67299] J.: Isolation, cDNA, and genomic structure of a conserved
gene (NOF)

[67300] at chromosome 11q13 next to FAU and oriented in the
opposite transcriptional

[67301] orientation. Genomics 34: 433-436, 1996.

[67302] 11426.

[67303] Lane, J. D; Lucocq, J.; Pryde, J.; Barr, F. A.; Woodman, P. G.;

[67304] Allan, V. J.; Lowe, M.: Caspase-mediated cleavage of the
stacking

[67305] protein GRASP65 is required for Golgi fragmentation dur-
ing apoptosis. J.

[67306] Cell Biol. 156: 495-509, 2002.

[67307] 11427.

[67308] Sutterlin, C.; Hsu, P.; Mallabiabarrena, A.; Malhotra, V.:
Fragmentation

[67309] and dispersal of the pericentriolar Golgi complex is re-
quired for

[67310] entry into mitosis in mammalian cells. Cell 109: 359-369,

2002.

[67311] 11428.

[67312] D'Orazi, G.; Cecchinelli, B.; Bruno, T.; Manni, I.; Hi-
gashimoto,

[67313] Y.; Saito, S.; Gostissa, M.; Coen, S.; Marchetti, A.; Del Sal,
G.;

[67314] Piaggio, G.; Fanciulli, M.; Appella, E.; Soddu, S.: Home-
odomain-interacting

[67315] protein kinase-2 phosphorylates p53 at Ser 46 and medi-
ates apoptosis. Nature

[67316] Cell Biol. 4: 11-19, 2002.

[67317] 11429.

[67318] Hofmann, T. G.; Mincheva, A.; Lichter, P.; Droge, W.;
Schmitz,

[67319] M. L.: Human homeodomain-interacting protein kinase-2
(HIPK2) is

[67320] a member of the DYRK family of protein kinases and maps
to chromosome

[67321] 7q32-q34. Biochimie 82: 1123-1127, 2000.

[67322] 11430.

[67323] Hofmann, T. G.; Moller, A.; Sirma, H.; Zentgraf, H.; Taya,

Y.;

[67324] Droge, W.; Will, H.; Schmitz, M. L.: Regulation of p53 activity by

[67325] its interaction with homeodomain-interacting protein kinase-2. Nature

[67326] Cell Biol. 4: 1-10, 2002.

[67327] 11431.

[67328] Pierantoni, G. M.; Bulfone, A.; Pentimalli, F.; Fedele, M.; Iuliano,

[67329] R.; Santoro, M.; Chiariotti, L.; Ballabio, A.; Fusco, A.: The homeodomain-interacting

[67330] protein kinase 2 gene is expressed late in embryogenesis and preferentially

[67331] in retina, muscle, and neural tissues. Biochem. Biophys. Res. Commun. 290:

[67332] 942-947, 2002.

[67333] 11432.

[67334] Wang, Y.; Hofmann, T. G.; Runkel, L.; Haaf, T.; Schaller, H.; Debatin,

[67335] K.-M.; Hug, H.: Isolation and characterization of cDNAs for the protein

[67336] kinase HIPK2. Biochim. Biophys. Acta 1518: 168-172,

2001.

[67337] 11433.

[67338] Arpaia, E.; Dumbrille-Ross, A.; Maler, T.; Neote, K.; Tropak, M.;

[67339] Troxel, C.; Stirling, J. L.; Pitts, J. S.; Bapat, B.; Lamhonwah, A.

[67340] M.; Mahuran, D. J.; Schuster, S. M.; Clarke, J. T. R.; Lowden, J.

[67341] A.; Gravel, R. A.: Identification of an altered splice site in Ashkenazi

[67342] Tay-Sachs disease. Nature 333: 85-86, 1988.

[67343] 11434.

[67344] Takeda, K.; Nakai, H.; Hagiwara, H.; Tada, K.; Shows, T. B.; Byers,

[67345] M. G.; Myerowitz, R.: Fine assignment of beta-hexosaminidase A alpha-subunit

[67346] on 15q23-q24 by high resolution in situ hybridization. Tohoku J.

[67347] Exp. Med. 160: 203-211, 1990.

[67348] 11435.

[67349] Tanaka, A.; Ohno, K.; Sandhoff, K.; Maire, I.; Kolodny, E.

H.;

[67350] Brown, A.; Suzuki, K.: GM2-gangliosidosis B1 variant: analysis of

[67351] beta-hexosaminidase alpha gene abnormalities in seven patients. Am.

[67352] J. Hum. Genet. 46: 329-339, 1990.

[67353] 11436.

[67354] Tanaka, A.; Ohno, K.; Suzuki, K.: GM2-gangliosidosis B1 variant:

[67355] a wide geographic and ethnic distribution of the specific beta-hexosaminidase

[67356] alpha chain mutation originally identified in a Puerto Rican patient. Biochem.

[67357] Biophys. Res. Commun. 156: 1015-1019, 1988.

[67358] 11437.

[67359] Tanaka, A.; Punnett, H. H.; Suzuki, K.: A new point muta-tion

[67360] in the beta-hexosaminidase alpha subunit gene responsi-ble for infantile

[67361] Tay-Sachs disease in a non-Jewish Caucasian patient (a Kpn mutant). Am.

[67362] J. Hum. Genet. 47: 567-574, 1990.

[67363] 11438.

[67364] Thomas, G. H.; Raghavan, S.; Kolodny, E. H.; Frisch, A.; Neufeld,

[67365] E. F.; O'Brien, J. S.; Reynolds, L. W.; Miller, C. S.; Shapiro, J.;

[67366] Kazazian, H. H., Jr.; Heller, R. H.: Nonuniform deficiency of hexosaminidase

[67367] A in tissues and fluids of two unrelated individuals. Pediat. Res. 16:

[67368] 232-237, 1982.

[67369] 11439.

[67370] Thurmon, T. F.: Tay-Sachs genes in Acadians. (Letter) Am. J.

[67371] Hum. Genet. 53: 781-782, 1993.

[67372] 11440.

[67373] Croze, E.; Russell-Harde, D.; Wagner, T. C.; Pu, H.; Pfeffer, L.

[67374] M.; Perez, H. D.: The human type I interferon receptor: identification

[67375] of the interferon beta-specific receptor-associated phos-phoprotein. J.

[67376] Biol. Chem. 271: 33165-33168, 1996.

[67377] 11441.

[67378] Croze, E.; Usacheva, A.; Asarnow, D.; Minshall, R.D.; Perez, H.D.;

[67379] Colamonici, O.: Receptor for activated C-kinase (RACK-1), a WD motif-containing

[67380] protein, specifically associates with the human type I IFN receptor. J.

[67381] Immun. 165: 5127-5132, 2000.

[67382] 11442.

[67383] Domanski, P.; Witte, M.; Kellum, M.; Rubinstein, M.; Hackett, R.;

[67384] Pitha, P.; Colamonici, O. R.: Cloning and expression of a long form

[67385] of the beta subunit of the interferon alpha/beta receptor that is

[67386] required for signaling. J. Biol. Chem. 270: 21606-21611, 1995.

[67387] 11443.

[67388] Lutfalla, G.; Holland, S. J.; Cinato, E.; Monneron, D.; Reboul,

[67389] J.; Rogers, N. C.; Smith, J. M.; Stark, G. R.; Gardiner, K.; Mogensen,

[67390] K. E.; Kerr, I. M.; Uze, G.: Mutant U5A cells are complemented by
[67391] an interferon-alpha/beta receptor subunit generated by alternative
[67392] processing of a new member of a cytokine receptor gene cluster. EMBO
[67393] J. 14: 5100-5108, 1995.

[67394] 11444.

[67395] Platanias, L. C.; Uddin, S.; Domanski, P.; Colamonici, O. R.:

[67396] Differences in interferon alpha and beta signaling: interferon beta
[67397] selectively induces the interaction of the alpha and beta(L) subunits
[67398] of the type I interferon receptor. J. Biol. Chem. 271: 23630-23633,
[67399] 1996.

[67400] 11445.

[67401] Raz, R.; Cheung, K.; Ling, L.; Levy, D. E.: Three distinct loci

[67402] on human chromosome 21 contribute to interferon-alpha/beta responsiveness. Somat.
[67403] Cell Molec. Genet. 21: 139-145, 1995.

[67404] 11446.

[67405] Finch, J. L.; Webb, G. C.; Evdokiou, A; Cowled, P. A.: Chro-
mosomal

[67406] localization of the human urothelial tetraspan gene,
UPK1B, to 3q13.3-q21

[67407] and detection of a Taq1 polymorphism. Genomics 40:
501-503, 1997.

[67408] 11447.

[67409] Yu, J.; Lin, J.-H.; Wu, X.-R.; Sun, T.-T.: Uroplakins Ia and
Ib,

[67410] two major differentiation products of bladder epithelium,
belong to

[67411] a family of four transmembrane domain (4TM) proteins. J.
Cell Biol. 125:

[67412] 171-182, 1994.

[67413] 11448.

[67414] Colley, W. C.; Altshuller, Y. M.; Sue-Ling, C. K.; Copeland,
N.

[67415] G.; Gilbert, D. J.; Jenkins, N. A.; Branch, K. D.; Tsirka, S. E.;

[67416] Bollag, R. J.; Bollag, W. B.; Frohman, M. A.: Cloning and
expression

[67417] analysis of murine phospholipase D1. Biochem. J. 326: 745-753, 1997.

[67418] 11449.

[67419] Schwamborn, K.; Weil, R.; Courtois, G.; Whiteside, S. T.; Israel,

[67420] A.: Phorbol esters and cytokines regulate the expression of the NEMO-related

[67421] protein, a molecule involved in a NF-kappa-B-independent pathway. J.

[67422] Biol. Chem. 275: 22780-22789, 2000.

[67423] 11450.

[67424] Fischer, G.; Perez-Rodriguez, M.; Arguello, J. R.; Cox, S. T.;

[67425] McWhinnie, A.; Travers, P. J.; Madrigal, J. A.: Three novel MICB

[67426] alleles. Tissue Antigens 55: 166-170, 2000.

[67427] 11451.

[67428] Groh, V.; Bahram, S.; Bauer, S.; Herman, A.; Beauchamp, M.; Spies,

[67429] T.: Cell stress-regulated human major histocompatibility complex

[67430] class I gene expressed in gastrointestinal epithelium. Proc. Nat.
[67431] Acad. Sci. 93: 12445-12450, 1996.

[67432] 11452.

[67433] Nakai, A.; Tanabe, M.; Kawazoe, Y.; Inazawa, J.; Morimoto, R. I.;
[67434] Nagata, K.: HSF4, a new member of the human heat shock factor family
[67435] which lacks properties of a transcriptional activator. Molec. Cell.
[67436] Biol. 17: 469-481, 1997.

[67437] 11453.

[67438] Uchida, K.; Yoshimura, A.; Inazawa, J.; Yanagisawa, K.; Os-
ada,
[67439] H.; Masuda, A.; Saito, T.; Takahashi, T.; Miyajima, A.; Takahashi,
[67440] T.: Molecular cloning of CISH, chromosome assignment to 3p21.3, and
[67441] analysis of expression in fetal and adult tissues. Cyto-
genet. Cell
[67442] Genet. 78: 209-212, 1997.

[67443] 11454.

[67444] Yoshimura, A.; Ohkubo, T.; Kiguchi, T.; Jenkins, N. A.; Gilbert,

[67445] D. J.; Copeland, N. G.; Hara, T.; Miyajima, A.: A novel cytokine-inducible

[67446] gene CIS, encodes an SH2-containing protein that binds to tyrosine-phosphorylated

[67447] interleukin 3 and erythropoietin receptors. EMBO J. 14: 2816-2826,

[67448] 1995.

[67449] 11455.

[67450] Chen, H.; Antonarakis, S. E.: The SH3D1A gene maps to human chromosome

[67451] 21q22.1-q22.2. Cytogenet. Cell Genet. 78: 213-215, 1997.

[67452] 11456.

[67453] Pucharcos, C.; Estivill, X.; de la Luna, S.: Intersectin 2, a

[67454] new multimodular protein involved in clathrin-mediated endocytosis. FEBS

[67455] Lett. 478: 43-51, 2000.

[67456] 11457.

[67457] Abbas, N.; Lucking, C. B.; Ricard, S.; Durr, A.; Bonifati, V.;

[67458] De Michele, G.; Bouley, S.; Vaughan, J. R.; Gasser, T.; Marconi, R.;

[67459] Broussolle, E.; Brefel-Courbon, C.; and 13 others: A wide variety

[67460] of mutations in the parkin gene are responsible for autosomal recessive

[67461] parkinsonism in Europe. Hum. Molec. Genet. 8: 567-574, 1999.

[67462] 11458.

[67463] Farrer, M.; Chan, P.; Chen, R.; Tan, L.; Lincoln, S.; Hernandez,

[67464] D.; Forno, L.; Gwinn-Hardy, K.; Petrucelli, L.; Hussey, J.; Singleton,

[67465] A.; Tanner, C.; Hardy, J.; Langston, J. W.: Lewy bodies and parkinsonism

[67466] in families with parkin mutations. Ann. Neurol. 50: 293-300, 2001.

[67467] 11459.

[67468] Hattori, N.; Matsumine, H.; Asakawa, S.; Kitada, T.; Yoshino, H.;

[67469] Elibol, B.; Brookes, A. J.; Yamamura, Y.; Kobayashi, T.;

Wang, M.;
[67470] Yoritaka, A.; Minoshima, S.; Shimizu, N.; Mizuno, Y.: Point mutations
[67471] (thr240arg and gln311stop) in the parkin gene. Biochem. Biophys.
[67472] Res. Commun. 249: 754-758, 1998. Note: Erratum: Biochem. Biophys.
[67473] Res. Commun. 251: 666 only, 1998.

[67474] 11460.

[67475] Hedrich, K.; Kann, M.; Lanthaler, A. J.; Dalski, A.; Eskelson,
[67476] C.; Landt, O.; Schwinger, E.; Vieregge, P.; Lang, A. E.; Breakefield,
[67477] X. O.; Ozelius, L. J.; Pramstaller, P. P.; Klein, C.: The im-
portance
[67478] of gene dosage studies: mutational analysis of the parkin gene in
[67479] early-onset parkinsonism. Hum. Molec. Genet. 10: 1649-1656, 2001.
[67480] 11461.

[67481] Hedrich, K.; Marder, K.; Harris, J.; Kann, M.; Lynch, T.; Meija-Santana,
[67482] H.; Pramstaller, P. P.; Schwinger, E.; Bressman, S. B.; Fahn,

S.;

[67483] Klein, C.: Evaluation of 50 probands with early-onset Parkinson's

[67484] disease for parkin mutations. Neurology 58: 1239-1246, 2002.

[67485] 11462.

[67486] Huynh, D. P.; Scoles, D. R.; Ho, T. H.; Del Bigio, M. R.; Pulst,

[67487] S.-M.: Parkin is associated with actin filaments in neuronal and

[67488] nonneural cells. Ann. Neurol. 48: 737-744, 2000.

[67489] 11463.

[67490] Imai, Y.; Soda, M.; Inoue, H.; Hattori, N.; Mizuno, Y.; Taka-hashi,

[67491] R.: An unfolded putative transmembrane polypeptide, which can lead

[67492] to endoplasmic reticulum stress, is a substrate of parkin. Cell 105:

[67493] 891-902, 2001.

[67494] 11464.

[67495] Jeon, B. S.; Kim, J.-M.; Lee, D.-S.; Hattori, N.; Mizuno, Y.:

[67496] An apparently sporadic case with parkin gene mutation in a Korean
[67497] woman. Arch. Neurol. 58: 988-989, 2001.

[67498] 11465.

[67499] Szepetowski, P.; Simon, M.-P.; Grosgeorge, J.; Huebner, K.; Bastard,
[67500] C.; Evans, G. A.; Tsujimoto, Y.; Birnbaum, D.; Theillet, C.; Gaudray,
[67501] P.: Localization of 11q13 loci with respect to regional chromosomal
[67502] breakpoints. Genomics 12: 738-744, 1992.

[67503] 11466.

[67504] Wang, T. C.; Cardiff, R. D.; Zukerberg, L.; Lees, E.; Arnold,
[67505] A.; Schmidt, E. V.: Mammary hyperplasia and carcinoma in MMTV-cyclin
[67506] D1 transgenic mice. Nature 369: 669-671, 1994.

[67507] 11467.

[67508] Xiong, Y.; Connelly, T.; Futcher, B.; Beach, D.: Human D-type
[67509] cyclin. Cell 65: 691-699, 1991.

[67510] 11468.

[67511] Yu, Q.; Geng, Y.; Sicinski, P.: Specific protection against breast

[67512] cancers by cyclin D1 ablation. Nature 411: 1017-1021, 2001.

[67513] 11469.

[67514] Fukumoto, S.; Suzawa, M.; Takeuchi, Y.; Kodama, Y.; Nakayama, K.;

[67515] Ogata, E.; Matsumoto, T.; Fujita, T.: Absence of mutations in parathyroid

[67516] hormone (PTH)/PTH-related protein receptor complementary deoxyribonucleic

[67517] acid in patients with pseudohypoparathyroidism type Ib. J. Clin.

[67518] Endocr. Metab. 81: 2554-2558, 1996.

[67519] 11470.

[67520] Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M. R.; Wiesener,

[67521] M. S.; Eckardt, K.-U.; Houlston, R. S.; Richards, F. M.; Latif, F.;

[67522] Maher, E. R.: Identification of cyclin D1 and other novel targets

[67523] for the von Hippel-Lindau tumor suppressor gene by ex-
pression array

[67524] analysis and investigation of cyclin D1 genotype as a
modifier in

[67525] von Hippel-Lindau disease. Cancer Res. 62: 3803-3811,
2002.

[67526] 11471.

[67527] Bettoun, J. D.; Minagawa, M.; Kwan, M. Y.; Lee, H. S.; Ya-
suda,

[67528] T.; Hendy, G. N.; Goltzman, D.; White, J. H.: Cloning and
characterization

[67529] of the promoter regions of the human parathyroid hor-
mone (PTH)/PTH-related

[67530] peptide receptor gene: analysis of deoxyribonucleic acid
from normal

[67531] subjects and patients with pseudohypoparathyroidism
type 1b. J. Clin.

[67532] Endocr. Metab. 82: 1031-1040, 1997.

[67533] 11472.

[67534] Gelbert, L.; Schipani, E.; Juppner, H.; Abou-Samra, A.-B.;
Segre,

[67535] G. V.; Naylor, S.; Drabkin, H.; Heath, H., III: Chromosomal

localization
[67536] of the parathyroid hormone/parathyroid hormone-related protein receptor
[67537] gene to human chromosome 3p21.1-p24.2. J. Clin. En-
docr. Metab. 79:
[67538] 1046-1048, 1994.

[67539] 11473.

[67540] Jobert, A.-S.; Zhang, P.; Couvineau, A.; Bonaventure, J.; Roume,
[67541] J.; Le Merrer, M.; Silve, C.: Absence of functional receptors for
[67542] parathyroid hormone and parathyroid hormone-related peptide in Blomstrand
[67543] chondrodysplasia. J. Clin. Invest. 102: 34-40, 1998.

[67544] 11474.

[67545] Fearon, E. R.; Vogelstein, B.: A genetic model for colorec-
tal
[67546] tumorigenesis. Cell 61: 759-767, 1990.

[67547] 11475.

[67548] Goldgaber, D.; Goldfarb, L. G.; Brown, P.; Asher, D. M.; Brown,

[67549] W. T.; Lin, S.; Teener, J. W.; Feinstone, S. M.; Rubenstein, R.; Kascsak,

[67550] R. J.; Boellaard, J. W.; Gajdusek, D. C.: Mutations in familial Creutzfeldt-Jakob

[67551] disease and Gerstmann-Straussler-Scheinker's syndrome. Exp. Neurol. 106:

[67552] 204-206, 1989.

[67553] 11476.

[67554] Goldhammer, Y.; Gabizon, R.; Meiner, Z.; Sadeh, M.: An Is-raeli

[67555] family with Gerstmann-Straussler-Scheinker disease man-ifesting the

[67556] codon 102 mutation in the prion protein gene. Neurology 43: 2718-2719,

[67557] 1993.

[67558] 11477.

[67559] Griffith, J. S.: Self-replication and scrapie. Nature 215:

[67560] 1043-1044, 1967.

[67561] 11478.

[67562] Haltia, M.; Kovanen, J.; Goldfarb, L. G.; Brown, P.; Gaj-dusek,

[67563] D. C.: Familial Creutzfeldt-Jakob disease in Finland: epi-
demiological,
[67564] clinical, pathological and molecular genetic studies. Eu-
rop. J. Epidemiol. 7:
[67565] 494-500, 1991.

[67566] 11479.

[67567] Head, M. W.; Tissingh, G.; Uitdehaag, B. M. J.; Barkhof, F.;
Bunn,
[67568] T. J. R.; Ironside, J. W.; Kamphorst, W.; Scheltens, P.: Spo-
radic
[67569] Creutzfeldt-Jakob disease in a young Dutch valine ho-
mozygote: atypical
[67570] molecular phenotype. Ann. Neurol. 50: 258-261, 2001.

[67571] 11480.

[67572] Hegde, R. S.; Mastrianni, J. A.; Scott, M. R.; DeFea, K. A.;
Tremblay,
[67573] P.; Torchia, M.; DeArmond, S. J.; Prusiner, S. B.; Lingappa,
V. R.
[67574] : A transmembrane form of the prion protein in neurode-
generative disease. Science 279:
[67575] 827-834, 1998.

[67576] 11481.

[67577] Hegde, R. S.; Tremblay, P.; Groth, D.; DeArmond, S. J.; Prusiner,

[67578] S. B.; Lingappa, V. R.: Transmissible and genetic prion diseases

[67579] share a common pathway of neurodegeneration. Nature 402: 732-736,

[67580] 1999.

[67581] 11482.

[67582] Horwich, A. L.; Weissman, J. S.: Deadly conformations--protein

[67583] misfolding in prion disease. Cell 89: 499-510, 1997.

[67584] 11483.

[67585] Hsiao, K.; Baker, H. F.; Crow, T. J.; Poulter, M.; Owen, F.; Terwilliger,

[67586] J. D.; Westaway, D.; Ott, J.; Prusiner, S. B.: Linkage of a prion

[67587] protein missense variant to Gerstmann-Straussler syndrome. Nature 338:

[67588] 342-345, 1989.

[67589] 11484.

[67590] Hsiao, K.; Cass, C.; Conneally, P. M.; Dlouhy, S. R.; Hodes, M.

[67591] E.; Farlow, M. R.; Ghetti, B.; Prusiner, S. B.: Atypical Gerstmann-Straussler-Scheinker

[67592] syndrome with neurofibrillary tangles: no mutation in the prion protein

[67593] open-reading-frame in a patient of the Indiana kindred. (Abstract) Neurobiol.

[67594] Aging 11: 302, 1990.

[67595] 11485.

[67596] Hsiao, K.; Dlouhy, S. R.; Farlow, M. R.; Cass, C.; Da Costa, M.;

[67597] Conneally, P. M.; Hodes, M. E.; Ghetti, B.; Prusiner, S. B.: Mutant

[67598] prion proteins in Gerstmann-Straussler-Scheinker disease with neurofibrillary

[67599] tangles. Nature Genet. 1: 68-71, 1992.

[67600] 11486.

[67601] Hsiao, K.; Meiner, Z.; Kahana, E.; Cass, C.; Kahana, I.; Avrahami,

[67602] D.; Scarlato, G.; Abramsky, O.; Prusiner, S. B.; Gabizon, R.: Mutation

[67603] of the prion protein in Libyan Jews with Creutzfeldt-Jakob disease. New
[67604] Eng. J. Med. 324: 1091-1097, 1991.

[67605] 11487.

[67606] Hsiao, K. K.; Groth, D.; Scott, M.; Yang, S.-L.; Serban, H.; Rapp,
[67607] D.; Foster, D.; Torchia, M.; DeArmond, S. J.; Prusiner, S. B.: Serial
[67608] transmission in rodents of neurodegeneration from trans-genic mice
[67609] expressing mutant prion protein. Proc. Nat. Acad. Sci. 91: 9126-9130,
[67610] 1994.

[67611] 11488.

[67612] Ironside, J. W.; Sutherland, K.; Bell, J. E.; McCardle, L.; Bar-rie,
[67613] C.; Estebeiro, K.; Zeidler, M.; Will, R. G.: A new variant of Creutzfeldt-Jakob
[67614] disease: neuropathological and clinical features. Cold Spring Harbor
[67615] Symp. Quant. Biol. 61: 523-530, 1996.

[67616] 11489.

[67617] Jendroska, K.; Hoffmann, O.; Schelosky, L.; Lees, A. J.; Poewe,

[67618] W.; Daniel, S. E.: Absence of disease related prion protein in neurodegenerative

[67619] disorders presenting with Parkinson's syndrome. J. Neurol. Neurosurg.

[67620] Psychiat. 57: 1249-1251, 1994.

[67621] 11490.

[67622] Kaneko, K.; Zulianello, L.; Scott, M.; Cooper, C. M.; Wal-lace,

[67623] A. C.; James, T. L.; Cohen, F. E.; Prusiner, S. B.: Evidence for

[67624] protein X binding to a discontinuous epitope on the cellu-lar prion

[67625] protein during scrapie prion propagation. Proc. Nat. Acad. Sci. 94:

[67626] 10069-10074, 1997.

[67627] 11491.

[67628] Kitamoto, T.; Ohta, M.; Doh-ura, K.; Hitoshi, S.; Terao, Y.; Tateishi,

[67629] J.: Novel missense variants of prion protein in Creutzfeldt-Jakob
[67630] disease or Gerstmann-Straussler syndrome. Biochem. Biophys. Res.
[67631] Commun. 191: 709-714, 1993.

[67632] 11492.

[67633] Kocisko, D. A.; Come, J. H.; Priola, S. A.; Chesebro, B.; Raymond,
[67634] G. J.; Lansbury, P. T.; Caughey, B.: Cell-free formation of protease-resistant
[67635] prion protein. Nature 370: 471-474, 1994.

[67636] 11493.

[67637] Krasemann, S.; Zerr, I.; Weber, T.; Poser, S.; Kretzschmar, H.;
[67638] Hunsmann, G.; Bodemer, W.: Prion disease associated with a novel
[67639] nine octapeptide repeat insertion in the PRNP gene. Molec. Brain
[67640] Res. 34: 173-176, 1995.

[67641] 11494.

[67642] Kretzschmar, H. A.; Neumann, M.; Stavrou, D.: Codon 178

mutation

[67643] of the human prion protein gene in a German family (Backer family):

[67644] sequencing data from 72-year-old celloidin-embedded brain tissue. Acta

[67645] Neuropath. 89: 96-98, 1995.

[67646] 11495.

[67647] Kretzschmar, H. A.; Stowring, L. E.; Westaway, D.; Stub-

blebine,

[67648] W. H.; Prusiner, S. B.; DeArmond, S. J.: Molecular cloning of a human

[67649] prion protein cDNA. DNA 5: 315-324, 1986.

[67650] 11496.

[67651] Kuwahara, C.; Takeuchi, A. M.; Nishimura, T.; Haraguchi, K.; Kubosaki,

[67652] A.; Matsumoto, Y.; Saeki, K.; Matsumoto, Y.; Yokoyama, T.; Itohara,

[67653] S.; Onodera, T.: Prions prevent neuronal cell-line death. (Letter) Nature 400:

[67654] 225-226, 1999.

[67655] 11497.

[67656] Laplanche, J.-L.; El Hachimi, K. H.; Durieux, I.; Thuillet, P.;

[67657] Defebvre, L.; Delasnerie-Laupretre, N.; Peoc'h, K.; Foncin, J.-F.;

[67658] Destee, A.: Prominent psychiatric features and early onset in an

[67659] inherited prion disease with a new insertional mutation in the prion

[67660] protein gene. Brain 122: 2375-2386, 1999.

[67661] 11498.

[67662] Laplanche, J. L.; Chatelain, J.; Thomas, S.; Launay, J. M.; Gaultier,

[67663] C.; Derouesne, C.: Uncommon phenotype for a codon 178 mutation of

[67664] the human PrP gene. (Letter) Ann. Neurol. 31: 345, 1992.

[67665] 11499.

[67666] Le, Y.; Yazawa, H.; Gong, W.; Yu, Z.; Ferrans, V. J.; Murphy,

[67667] P. M.; Wang, J. M.: Cutting edge: the neurotoxic prion peptide fragment

[67668] PrP(106-126) is a chemotactic agonist for the G protein-coupled receptor

[67669] formyl peptide receptor-like 1. J. Immun. 166:

1448-1451, 2001.

[67670] 11500.

[67671] Gibbons, B.; Scott, D.; Hungerford, J. L.; Cheung, K. L.; Harrison,

[67672] C.; Attard-Montalto, S.; Evans, M.; Birch, J. M.; Kingston, J. E.

[67673] : Retinoblastoma in association with the chromosome breakage syndromes

[67674] Fanconi's anaemia and Bloom's syndrome: clinical and cytogenetic findings. Clin.

[67675] Genet. 47: 311-317, 1995.

[67676] 11501.

[67677] Girardet, A.; McPeek, M. S.; Leeflang, E. P.; Munier, F.; Arnheim,

[67678] N.; Claustres, M.; Pellestor, F.: Meiotic segregation analysis of

[67679] RB1 alleles in retinoblastoma pedigrees by use of single-sperm typing. Am.

[67680] J. Hum. Genet. 66: 167-175, 2000.

[67681] 11502.

[67682] Godbout, R.; Dryja, T. P.; Squire, J.; Gallie, B. L.; Phillips,

[67683] R. A.: Somatic inactivation of genes on chromosome 13 is a common
[67684] event in retinoblastoma. Nature 304: 451-453, 1983.

[67685] 11503.

[67686] Goodrich, D. W.; Wang, N. P.; Qian, Y.-W.; Lee, E. Y.-H. P.; Lee,
[67687] W.-H.: The retinoblastoma gene product regulates progression through
[67688] the G1 phase of the cell cycle. Cell 67: 293-302, 1991.

[67689] 11504.

[67690] Grace, E.; Drennan, J.; Colver, D.; Gordon, R. R.: The 13q deletion
[67691] syndrome. J. Med. Genet. 8: 351-357, 1971.

[67692] 11505.

[67693] Green, A. R.; Wyke, J. A.: Anti-oncogenes: a subset of regulatory
[67694] genes involved in carcinogenesis? Lancet II: 475-477, 1985.

[67695] 11506.

[67696] Greger, V.; Kerst, S.; Messmer, E.; Hopping, W.; Passarge, E.;

[67697] Horsthemke, B.: Application of linkage analysis to genetic counselling
[67698] in families with hereditary retinoblastoma. J. Med. Genet. 25: 217-221,
[67699] 1988.

[67700] 11507.

[67701] Greger, V.; Passarge, E.; Horsthemke, B.: Somatic mosaicism in
[67702] a patient with bilateral retinoblastoma. Am. J. Hum. Genet. 46:
[67703] 1187-1193, 1990.

[67704] 11508.

[67705] Hagstrom, S. A.; Dryja, T. P.: Mitotic recombination map of 13cen-13q14
[67706] derived from an investigation of loss of heterozygosity in retinoblastomas. Proc.
[67707] Nat. Acad. Sci. 96: 2952-2957, 1999.

[67708] 11509.

[67709] Hall, J. G.: Personal Communication. Vancouver, British Columbia,
[67710] Canada 5/29/1993.

[67711] 11510.

[67712] Hanahan, D.; Weinberg, R. A.: The hallmarks of cancer. Cell 100:

[67713] 57-70, 2000.

[67714] 11511.

[67715] Harbour, J. W.: Molecular basis of low-penetrance retinoblastoma. Arch.

[67716] Ophthal. 119: 1699-1704, 2001.

[67717] 11512.

[67718] Harbour, J. W.; Lai, S.-L.; Whang-Peng, J.; Gazdar, A. F.; Minna,

[67719] J. D.; Kaye, F. J.: Abnormalities in structure and expression of

[67720] the human retinoblastoma gene in SCLC. Science 241: 353-357, 1988.

[67721] 11513.

[67722] Hensel, C.; Hsieh, C.-L.; Lee, W.-H.; Pam-Lee, E.; Gazdar, A.;

[67723] Sakaguchi, A. Y.; Naylor, S. L.: Allele loss and lack of expression

[67724] of the RB-1 locus in small cell lung cancer. (Abstract) Am.

J. Hum.
[67725] Genet. 43: A25, 1988.

[67726] 11514.

[67727] Henson, J. W.; Schnitker, B. L.; Correa, K. M.; von Diem-
ling,
[67728] A.; Fassbender, F.; Xu, H.-J.; Benedict, W. F.; Yandell, D.
W.; Louis,
[67729] D. N.: The retinoblastoma gene is involved in malignant
progression
[67730] of astrocytomas. Ann. Neurol. 36: 714-721, 1994.

[67731] 11515.

[67732] Higgins, M. J.; Hansen, M. F.; Cavenee, W. K.; Lalande, M.:
Molecular
[67733] detection of chromosomal translocations that disrupt the
putative
[67734] retinoblastoma susceptibility locus. Molec. Cell. Biol. 9:
1-5,
[67735] 1989.

[67736] 11516.

[67737] Hoegerman, S. F.: Chromosome 13 long arm interstitial
deletion

[67738] may result from maternal inverted insertion. Science 205: 1035-1036,
[67739] 1979.

[67740] 11517.

[67741] Hogg, A.; Bia, B.; Onadim, Z.; Cowell, J. K.: Molecular mechanisms
[67742] of oncogenic mutations in tumours from patients with bilateral and
[67743] unilateral retinoblastoma. Proc. Nat. Acad. Sci. 90: 7351-7355,
[67744] 1993.

[67745] 11518.

[67746] Honavar, S. G.; Shields, C. L.; Shields, J. A.; Demirci, H.; Naduvilath,
[67747] T. J.: Intraocular surgery after treatment of retinoblastoma. Arch.
[67748] Ophthal. 119: 1613-1621, 2001.

[67749] 11519.

[67750] Honavar, S. G.; Singh, A. D.; Shields, C. L.; Meadows, A. T.;

[67751] Demirci, H.; Cater, J.; Shields, J. A.: Postenucleation adjuvant

[67752] therapy in high-risk retinoblastoma. Arch. Ophthal. 120: 923-931,

[67753] 2002.

[67754] 11520.

[67755] Hong, F. D.; Huang, H.-J. S.; To, H.; Young, L.-J. S.; Oro, A.;

[67756] Bookstein, R.; Lee, E. Y.-H. P.; Lee, W.-H.: Structure of the human

[67757] retinoblastoma gene. Proc. Nat. Acad. Sci. 86: 5502-5506, 1989.

[67758]

[67759] 100. Horowitz, J. M.; Park, S.-H.; Bogenmann, E.; Cheng, J.-C.; Yandell,

[67760] D. W.; Kaye, F. J.; Minna, J. D.; Dryja, T. P.; Weinberg, R. A.:

[67761] Frequent inactivation of the retinoblastoma anti-oncogene is restricted

[67762] to a subset of human tumor cells. Proc. Nat. Acad. Sci. 87: 2775-2779,

[67763] 1990.

[67764]

[67765] 101. Horowitz, J. M.; Park, S. H.; Yandell, D. W.; Weinberg,

R. A.
[67766] : Involvement of the retinoblastoma gene in the genesis of various
[67767] human tumors:.In: Kavenee, W.; Hastie, N.; Stanbridge, E.: Recessive
[67768] Oncogenes and Tumor Suppression: Current Communications in Molecular
[67769] Biology. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory
[67770] Press (pub.) 1989. Pp. 101-108.

[67771]

[67772] 102. Horsthemke, B.; Greger, V.; Barnert, H. J.; Hopping, W.; Passarge,
[67773] E.: Detection of submicroscopic deletions and a DNA polymorphism
[67774] at the retinoblastoma locus. Hum. Genet. 76: 257-261, 1987.

[67775]

[67776] 103. Horsthemke, B.; Greger, V.; Becher, R.; Passarge, E.: Mechanism
[67777] of i(6p) formation in retinoblastoma tumor cells. Cancer Genet. Cytogenet. 37:

[67778] 95-102, 1989.

[67779]

[67780] 104. Hsieh, J.-K.; Chan, F. S. G.; O'Connor, D. J.; Mit-
tnacht, S.;
[67781] Zhong, S.; Lu, X.: RB regulates the stability and the apop-
totic function
[67782] of p53 via MDM2. Molec. Cell 3: 181-193, 1999.

[67783]

[67784] 105. Huang, H.-J. S.; Yee, J.-K.; Shew, J.-Y.; Chen, P.-L.;
Bookstein,
[67785] R.; Friedmann, T.; Lee, E. Y.-H. P.; Lee, W.-H.: Suppression
of the
[67786] neoplastic phenotype by replacement of the RB gene in
human cancer
[67787] cells. Science 242: 1563-1566, 1988.

[67788]

[67789] 106. Janson, M.; Nordenskjold, M.: A constitutional muta-
tion within
[67790] the retinoblastoma gene detected by PFGE. Clin. Genet.
45: 5-10,
[67791] 1994.

[67792]

[67793] 107. Jensen, R. D.; Miller, R. W.: Retinoblastoma: epidemi-
ologic
[67794] characteristics. New Eng. J. Med. 285: 307-311, 1971.

[67795]

[67796] 108. Kimchi, A.; Wang, X.-F.; Weinberg, R. A.; Cheifetz, S.;
Massague,
[67797] J.: Absence of TGF-beta receptors and growth inhibitory
responses
[67798] in retinoblastoma cells. Science 240: 196-199, 1988.

[67799]

[67800] 109. Kitchin, F. D.; Ellsworth, R. M.: Pleiotropic effects of
the
[67801] gene for retinoblastoma. J. Med. Genet. 11: 244-246,
1974.

[67802]

[67803] 110. Kivela, T.: Trilateral retinoblastoma: a meta-analysis
of hereditary
[67804] retinoblastoma associated with primary ectopic intracra-
nial retinoblastoma. J.
[67805] Clin. Oncol. 17: 1829-1837, 1999.

[67806]

[67807] 111. Kivela, T.; Asko-Seljavaara, S.; Pihkala, U.; Hovi, L.; Heikkonen,

[67808] J.: Sebaceous carcinoma of the eyelid associated with retinoblastoma. Ophthalmology 108:

[67809] 1124-1128, 2001.

[67810]

[67811] 112. Klutz, M.; Brockmann, D.; Lohmann, D. R.: A parent-of-origin

[67812] effect in two families with retinoblastoma is associated with a distinct

[67813] splice mutation in the RB1 gene. Am. J. Hum. Genet. 71: 174-179,

[67814] 2002.

[67815]

[67816] 113. Knight, L. A.; Gardner, H. A.; Gallie, B. L.: Familial retinoblastoma:

[67817] segregation of chromosome 13 in four families. Am. J. Hum. Genet. 32:

[67818] 194-201, 1980.

[67819]

[67820] 114. Knudson, A. G.: Hereditary cancer, oncogenes and anti-oncogenes. Cancer
[67821] Res. 45: 1437-1443, 1985.

[67822]

[67823] 115. Knudson, A. G., Jr.: Mutation and cancer: statistical study
[67824] of retinoblastoma. Proc. Nat. Acad. Sci. 68: 820-823, 1971.

[67825]

[67826] 116. Knudson, A. G., Jr.: Genetics of human cancer. Annu. Rev. Genet. 20:
[67827] 231-251, 1986.

[67828]

[67829] 117. Knudson, A. G., Jr.; Hethcote, H. W.; Brown, B. W.: Mutation
[67830] and childhood cancer: a probabilistic model for the incidence of retinoblastoma. Proc.
[67831] Nat. Acad. Sci. 72: 5116-5120, 1975.

[67832]

[67833] 118. Knudson, A. G., Jr.; Meadows, A. T.; Nichols, W. W.; Hill, R.

[67834] : Chromosomal deletion and retinoblastoma. New Eng. J. Med. 295:
[67835] 1120-1123, 1976.

[67836]

[67837] 119. Laquis, S. J.; Rodriguez-Galindo, C.; Wilson, M. W.; Fleming,
[67838] J. C.; Haik, B. G.: Retinoblastoma in a patient with an X;13 translocation
[67839] and facial abnormalities consistent with 13q- syndrome. Am. J. Ophthal. 133:
[67840] 285-287, 2002.

[67841]

[67842] 120. Lee, W.-H.; Bookstein, R.; Hong, F.; Young, L.-J.; Shew, J.-Y.;
[67843] Lee, E. Y.-H. P.: Human retinoblastoma susceptibility gene: cloning,
[67844] identification, and sequence. Science 235: 1394-1399, 1987.

[67845]

[67846] 121. Lee, W.-H.; Shew, J.-Y.; Hong, F. D.; Sery, T. W.; Donoso, L.

[67847] A.; Young, L.-J.; Bookstein, R.; Lee, E. Y.-H. P.: The retinoblastoma
[67848] susceptibility gene encodes a nuclear phosphoprotein associated with
[67849] DNA binding activity. Nature 329: 642-645, 1987.

[67850]

[67851] 122. Lele, K. P.; Penrose, L. S.; Stallard, H. B.: Chromosome deletion
[67852] in a case of retinoblastoma. Ann. Hum. Genet. 27: 171-174, 1963.

[67853]

[67854] 123. Lemieux, N.; Messier, P. E.; Jacob, J. L.; Milot, J.; Richer,
[67855] C. L.: Precise cytogenetic localization of the Rb locus at subband
[67856] 13q14.11 by ultrastructural detection after immunochemical chromosome
[67857] banding. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A27, 1989.

[67858]

[67859] 124. Liu, Z.; Song, Y.; Bia, B.; Cowell, J. K.: Germline mutations

[67860] in the RB1 gene in patients with hereditary retinoblas-
toma. Genes
[67861] Chromosomes Cancer 14: 277-284, 1995.

[67862]

[67863] 125. Lohmann, D. R.; Brandt, B.; Hopping, W.; Passarge,
E.; Horsthemke,
[67864] B.: Spectrum of small length germline mutations in the
RB1 gene. Hum.
[67865] Molec. Genet. 3: 2187-2193, 1994.

[67866]

[67867] 126. Lohmann, D. R.; Brandt, B.; Hopping, W.; Passarge,
E.; Horsthemke,
[67868] B.: The spectrum of RB1 germ-line mutations in hereditary
retinoblastoma. Am.
[67869] J. Hum. Genet. 58: 940-949, 1996.

[67870]

[67871] 127. Lohmann, D. R.; Gerick, M.; Brandt, B.; Oelschlager,
U.; Lorenz,
[67872] B.; Passarge, E.; Horsthemke, B.: Constitutional RB1-gene
mutations
[67873] in patients with isolated unilateral retinoblastoma. Am. J.

Hum.
[67874] Genet. 61: 282-294, 1997.

[67875]

[67876] 128. Lomazzi, M.; Moroni, M. C.; Jensen, M. R.; Frittoli, E.; Helin,
[67877] K.: Suppression of the p53- or pRB-mediated G1 check-point is required
[67878] for E2F-induced S-phase entry. Nature Genet 31: 190-194, 2002.

[67879]

[67880] 129. Lueder, G. T.; Judisch, G. F.; Wen, B.-C.: Heritable retinoblastoma
[67881] and pinealoma. Arch. Ophthal. 109: 1707-1709, 1991.

[67882]

[67883] 130. Luo, R. X.; Postigo, A. A.; Dean, D. C.: Rb interacts with histone
[67884] deacetylase to repress transcription. Cell 92: 463-473, 1998.

[67885]

[67886] 131. Maat-Kievit, J. A.; Oepkes, D.; Hartwig, N. G.; Ver-meij-Keers,

[67887] C.; van Kamp, I. L.; van de Kamp, J. J. P.: A large retinoblastoma
[67888] detected in a fetus at 21 weeks of gestation. Prenatal Diag. 13:
[67889] 377-384, 1993.

[67890]

[67891] 132. MacKay, C. J.; Abramson, D. H.; Ellsworth, R. M.: Metastatic
[67892] patterns of retinoblastoma. Arch. Ophthal. 102: 391-396, 1984.

[67893]

[67894] 133. Macklin, M. T.: A study of retinoblastoma in Ohio. Am. J. Hum.
[67895] Genet. 12: 1-43, 1960.

[67896]

[67897] 134. Macklin, M. T.: Inheritance of retinoblastoma in Ohio. Arch.
[67898] Ophthal. 62: 842-851, 1959.

[67899]

[67900] 135. Manchester, P. T., Jr.: Retinoblastoma among off-spring of adult

[67901] survivors. Arch. Ophthal. 65: 546-549, 1961.

[67902]

[67903] 136. Mancini, D.; Singh, S.; Ainsworth, P.; Rodenhiser, D.: Constitutively

[67904] methylated CpG dinucleotides as mutation hot spots in the retinoblastoma

[67905] gene (RB1). Am. J. Hum. Genet. 61: 80-87, 1997.

[67906]

[67907] 137. Mancini, M. A.; Shan, B.; Nickerson, J. A.; Penman, S.; Lee,

[67908] W.-H.: The retinoblastoma gene product is a cell cycle-dependent,

[67909] nuclear matrix-associated protein. Proc. Nat. Acad. Sci. 91: 418-422,

[67910] 1994.

[67911]

[67912] 138. Marino, S.; Vooijs, M.; van der Gulden, H.; Jonker, J.; Berns,

[67913] A.: Induction of medulloblastomas in p53-null mutant mice by somatic

[67914] inactivation of Rb in the external granular layer cells of

the cerebellum. Gene
[67915] s Dev. 14: 994-1004, 2000.

[67916]

[67917] 139. Matsunaga, E.: Recurrence risks to relatives of patients with
[67918] retinoblastoma. Jpn. J. Ophthal. 22: 313-319, 1978.

[67919]

[67920] 140. Matsunaga, E.: Almost synchronous appearance of bilateral retinoblastomas.
[67921] (Letter) Am. J. Med. Genet. 11: 485-487, 1982.

[67922]

[67923] 141. Matsunaga, E.: Retinoblastoma: mutational mosaicism or host
[67924] resistance? Am. J. Med. Genet. 8: 375-387, 1981.

[67925]

[67926] 142. Matsunaga, E.: Hereditary retinoblastoma: host resistance and
[67927] second primary tumors. J. Nat. Cancer Inst. 65: 47-51, 1980.

[67928]

[67929] 143. Matsunaga, E.: Hereditary retinoblastoma: delayed mutation or
[67930] host resistance? Am. J. Hum. Genet. 30: 406-425, 1978.

[67931]

[67932] 144. Matsunaga, E.; Minoda, K.; Sasaki, M. S.: Parental age and seasonal
[67933] variation in the births of children with sporadic retinoblastoma:
[67934] a mutation-epidemiologic study. Hum. Genet. 84: 155-158, 1990.

[67935]

[67936] 145. Michalova, K.; Kloucek, F.; Musilova, J.: Deletion of 13q in
[67937] two patients with retinoblastoma, one probably due to 13q- mosaicism
[67938] in the mother. Hum. Genet. 61: 264-266, 1982.

[67939]

[67940] 146. Moll, A. C.; Imhof, S. M.; Schouten-Van Meeteren, A. Y. N.; Kuik,
[67941] D. J.; Hofman, P.; Boers, M.: Second primary tumors in hereditary

[67942] retinoblastoma: a register-based study, 1945-1997. Is there an age

[67943] effect on radiation-related risk? Ophthalmology 108: 1109-1114,

[67944] 2001.

[67945]

[67946] 147. Motegi, T.: Lymphocyte chromosome survey in 42 patients with

[67947] retinoblastoma: effort to detect 13q14 deletion mosaicism. Hum. Genet. 58:

[67948] 168-173, 1981.

[67949]

[67950] 148. Motegi, T.: High rate of detection of 13q14 deletion mosaicism

[67951] among retinoblastoma patients (using more extensive methods). Hum.

[67952] Genet. 61: 95-97, 1982.

[67953]

[67954] 149. Motegi, T.; Kaga, M.; Yanagawa, Y.; Kadowaki, H.; Watanabe, K.;

[67955] Inoue, A.; Komatsu, M.; Minoda, K.: A recognizable pattern

of the

[67956] midface of retinoblastoma patients with interstitial deletion of 13q. Hum.

[67957] Genet. 64: 160-162, 1983.

[67958]

[67959] 150. Motegi, T.; Komatsu, M.; Minoda, K.: Is the interstitial deletion

[67960] of 13q in retinoblastoma patients not transmissible? (Letter) Hum.

[67961] Genet. 64: 205, 1983.

[67962]

[67963] 151. Motegi, T.; Komatsu, M.; Nakazato, Y.; Ohuchi, M.; Minoda, K.

[67964] : Retinoblastoma in a boy with a de novo mutation of a 13/18 translocation:

[67965] the assumption that the retinoblastoma locus is at 13q141, particularly

[67966] at the distal portion of it. Hum. Genet. 60: 193-195, 1982.

[67967]

[67968] 152. Munier, F.; Spence, M. A.; Pescia, G.; Balmer, A.; Gail-

loud,
[67969] C.; Thonney, F.; van Melle, G.; Rutz, H. P.: Paternal selec-
tion favoring
[67970] mutant alleles of the retinoblastoma susceptibility gene.
Hum. Genet. 89:
[67971] 508-512, 1992.

[67972]

[67973] 153. Munier, F. L.; Wang, M. X.; Spence, M. A.; Thonney,
F.; Balmer,
[67974] A.; Pescia, G.; Donoso, L. A.; Murphree, A. L.: Pseudo low
penetrance
[67975] in retinoblastoma: fortuitous familial aggregation of spo-
radic cases
[67976] caused by independently derived mutations in two large
pedigrees. Arch.
[67977] Ophthal. 111: 1507-1511, 1993.

[67978]

[67979] 154. Murphree, A. L.; Benedict, W. F.: Retinoblastoma:
clues to human
[67980] oncogenesis. Science 223: 1028-1033, 1984.

[67981]

[67982] 155. Naumova, A.; Hansen, M.; Strong, L.; Jones, P. A.; Hadjistilianou,
[67983] D.; Mastrangelo, D.; Griegel, S.; Rajewsky, M. F.; Shields, J.; Donoso,
[67984] L.; Wang, M.; Sapienza, C.: Concordance between parental origin of
[67985] chromosome 13q loss and chromosome 6p duplication in sporadic retinoblastoma. Am.
[67986] J. Hum. Genet. 54: 274-281, 1994.

[67987]

[67988] 156. Naumova, A.; Sapienza, C.: The genetics of retinoblastoma, revisited. Am.
[67989] J. Hum. Genet. 54: 264-273, 1994.

[67990]

[67991] 157. Nevins, J. R.: The Rb/E2F pathway and cancer. Hum. Molec. Genet. 10:
[67992] 699-703, 2001.

[67993]

[67994] 158. Nichols, W. W.; Miller, R. C.; Sobel, M.; Hoffman, E.; Sparkes,
[67995] R. S.; Mohandas, T.; Veomett, I.; Davis, J. R.: Further ob-

servations
[67996] on a 13qXp translocation associated with retinoblastoma. Am. J. Ophthal. 89:
[67997] 621-627, 1980.

[67998]

[67999] 159. Nielsen, S. J.; Schneider, R.; Bauer, U.-M.; Bannister, A. J.;
[68000] Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein,
[68001] T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation
[68002] and HP1 to promoters. Nature 412: 561-565, 2001.

[68003]

[68004] 160. Nirankari, M. S.; Gulati, G. C.; Chaddah, M. R.: Retinoblastoma:
[68005] genetics and report of a family. Am. J. Ophthal. 53: 523-532, 1962.

[68006]

[68007] 161. Noorani, H. Z.; Khan, H. N.; Gallie, B. L.; Detsky, A. S.: Cost
[68008] comparison of molecular versus conventional screening of

relatives
[68009] at risk for retinoblastoma. Am. J. Hum. Genet. 59:
301-307, 1996.
[68010]

[68011] 162. Nussbaum, R.; Puck, J.: Recurrence risks for
retinoblastoma:
[68012] a model for autosomal dominant disorders with complex
inheritance. J.
[68013] Pediat. Ophthal. 13: 89-98, 1976.

[68014]

[68015] 163. Onadim, Z.; Hogg, A.; Baird, P. N.; Cowell, J. K.:
Oncogenic
[68016] point mutations in exon 20 of the RB1 gene in families
showing incomplete
[68017] penetrance and mild expression of the retinoblastoma
phenotype. Proc.
[68018] Nat. Acad. Sci. 89: 6177-6181, 1992.

[68019]

[68020] 164. Onadim, Z.; Woolford, A. J.; Kingston, J. E.; Hunger-
ford, J.
[68021] L.: The RB1 gene mutation in a child with ectopic intracra-

nial retinoblastoma. Brit.
[68022] J. Cancer 76: 1405-1409, 1997.

[68023]

[68024] 165. Ono, T.; Yoshida, M. C.: Chromosomal assignment of retinoblastoma
[68025] 1 gene (RB1) to mouse 14D3 and rat 15q12 by fluorescence in situ hybridization. Jpn.
[68026] J. Genet. 68: 617-621, 1993.

[68027]

[68028] 166. Orye, E.; Benoit, Y.; Coppieters, R.; Jeannin, P.; Vercruysse,
[68029] C.; Delaey, J.; Delbeke, M.-J.: A case of retinoblastoma, associated
[68030] with histiocytosis-X and mosaicism of a deleted D-group chromosome
[68031] (13q14-q31). Clin. Genet. 22: 37-39, 1982.

[68032]

[68033] 167. Orye, E.; Delbeke, M. J.; Vandenabeele, B.: Retinoblastoma and
[68034] long arm deletion of chromosome 13. Attempts to define the deleted

[68035] segment. Clin. Genet. 5: 457-464, 1974.

[68036]

[68037] 168. Orye, E.; Delbeke, M. J.; Vandenabeele, B.: Retinoblastoma and

[68038] D-chromosome deletions. (Letter) Lancet II: 1376, 1971.

[68039]

[68040] 169. Otterson, G. A.; Modi, S.; Nguyen, K.; Coxon, A. B.; Kaye, F.

[68041] J.: Temperature-sensitive RB mutations linked to incom- plete penetrance

[68042] of familial retinoblastoma in 12 families. Am. J. Hum. Genet. 65:

[68043] 1040-1046, 1999.

[68044]

[68045] 170. Otterson, G. W.; Chen, W.; Coxon, A. B.; Khleif, S. N.; Kaye,

[68046] F. J.: Incomplete penetrance of familial retinoblastoma linked to

[68047] germ-line mutations that result in partial loss of RB func- tion. Proc.

[68048] Nat. Acad. Sci. 94: 12036-12040, 1997.

[68049]

[68050] 171. Pendergrass, T. W.; Davis, S.: Incidence of retinoblas-
toma in
[68051] the United States. Arch. Ophthal. 98: 1204-1210, 1980.

[68052]

[68053] 172. Pennaneach, V.; Salles-Passador, I.; Munshi, A.;
Brickner, H.;
[68054] Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.-T.; Wang, J. Y.
J.; Fotedar,
[68055] R.; Fotedar, A.: The large subunit of replication factor C
promotes
[68056] cell survival after DNA damage in an LxCxE motif- and
Rb-dependent
[68057] manner. Molec. Cell 7: 715-727, 2001.

[68058]

[68059] 173. Riccardi, V. M.; Hittner, H. M.; Francke, U.; Pippin, S.;
Holmquist,
[68060] G. P.; Kretzer, F. L.; Ferrell, R.: Partial triplication and
deletion
[68061] of 13q: study of a family presenting with bilateral
retinoblastomas. Clin.

[68062] Genet. 15: 332-345, 1979.

[68063]

[68064] 174. Rivera, H.; Turleau, C.; de Grouchy, J.; Junien, C.; De-
spoisse,
[68065] S.; Zucker, J.-M.: Retinoblastoma-del(13q14): report of
two patients,
[68066] one with a trisomic sib due to maternal insertion; gene-
dosage effect
[68067] for esterase D. Hum. Genet. 59: 211-214, 1981.

[68068]

[68069] 175. Sakai, T.; Ohtani, N.; McGee, T. L.; Robbins, P. D.;
Dryja, T.
[68070] P.: Oncogenic germ-line mutations in Sp1 and ATF sites in
the human
[68071] retinoblastoma gene. Nature 353: 83-86, 1991.

[68072]

[68073] 176. Sakai, T.; Ohtani, N.; McGee, T. L.; Robbins, P. D.;
Dryja, T.
[68074] P.: Oncogenic germ-line mutations in Sp1 and ATF sites in
the human
[68075] retinoblastoma gene. Nature 353: 83-86, 1991.

[68076]

[68077] 177. Sakai, T.; Toguchida, J.; Ohtani, N.; Yandell, D. W.; Rapaport,
[68078] J. M.; Dryja, T. P.: Allele-specific hypermethylation of the retinoblastoma
[68079] tumor-suppressor gene. Am. J. Hum. Genet. 48: 880-888, 1991.
[68080]

[68081] 178. Schappert-Kimmijser, J.; Hemmes, G. D.; Nijland, R.: The heredity
[68082] of retinoblastoma. Ophthalmologica 151: 197-213, 1966.
[68083]

[68084] 179. Scheffer, H.; te Meerman, G. J.; Kruize, Y. C. M.; van den Berg,
[68085] A. H. M.; Penninga, D. P.; Tan, K. E. W. P.; der Kinderen, D. J.;
[68086] Buys, C. H. C. M.: Linkage analysis of families with heredi-tary retinoblastoma:
[68087] nonpenetrance of mutation, revealed by combined use of markers within
[68088] and flanking the RB1 gene. Am. J. Hum. Genet. 45: 252-260, 1989.

[68089]

[68090] 180. Schimke, R. N.; Lowman, J.; Cowan, G.: Retinoblas-
toma and osteogenic
[68091] sarcoma in sibs. Cancer 34: 2077-2079, 1974.

[68092]

[68093] 181. Schubert, E. L.; Strong, L. C.; Hansen, M. F.: A splic-
ing mutation
[68094] in RB1 in low penetrance retinoblastoma. Hum. Genet.
100: 557-563,
[68095] 1997.

[68096]

[68097] 182. Shields, C. L.; Honavar, S.; Shields, J. A.; Demirci, H.;
Meadows,
[68098] A. T.: Vitrectomy in eyes with unsuspected retinoblas-
toma. Ophthalmology 107:
[68099] 2250-2255, 2000.

[68100]

[68101] 183. Shiio, Y.; Yamamoto, T.; Yamaguchi, N.: Negative
regulation
[68102] of Rb expression by the p53 gene product. Proc. Nat.
Acad. Sci. 89:

[68103] 5206-5210, 1992.

[68104]

[68105] 184. Shroeder, W. T.; Chao, L.-Y.; Dao, D. D.; Strong, L. C.; Pathak,

[68106] S.; Riccardi, V.; Lewis, W. H.; Saunders, G. F.: Nonrandom loss of

[68107] maternal chromosome 11 alleles in Wilms tumors. Am. J. Hum. Genet. 40:

[68108] 413-420, 1987.

[68109]

[68110] 185. Sippel, K. C.; Fraioli, R. E.; Smith, G. D.; Schalkoff, M. E.;

[68111] Sutherland, J.; Gallie, B. L.; Dryja, T. P.: Frequency of so-matic

[68112] and germ-line mosaicism in retinoblastoma: implications for genetic

[68113] counseling. Am. J. Hum. Genet. 62: 610-619, 1998.

[68114]

[68115] 186. Smith, S. M.; Sorsby, A.: Retinoblastoma: some ge-netic aspects. Ann.

[68116] Hum. Genet. 23: 50-58, 1958.

[68117]

[68118] 187. Sparkes, R. S.: The genetics of retinoblastoma. Biochim. Biophys.

[68119] Acta 780: 95-118, 1985.

[68120]

[68121] 188. Sparkes, R. S.; Muller, H.; Klisak, I.; Abram, J. A.: Retinoblastoma

[68122] with 13q; chromosomal deletion associated with maternal paracentric

[68123] inversion of 13q. Science 203: 1027-1029, 1979.

[68124]

[68125] 189. Sparkes, R. S.; Murphree, A. L.; Lingua, R. W.; Sparkes, M. C.;

[68126] Field, L. L.; Funderburk, S. J.; Benedict, W. F.: Gene for hereditary

[68127] retinoblastoma assigned to human chromosome 13 by linkage to esterase

[68128] D. Science 219: 971-973, 1983.

[68129]

[68130] 190. Sparkes, R. S.; Sparkes, M. C.; Wilson, M. G.; Towner, J. W.;

[68131] Benedict, W.; Murphree, A. L.; Yunis, J. J.: Regional assign-
ment
[68132] of genes for human esterase D and retinoblastoma to
chromosome band
[68133] 13q14. Science 208: 1042-1044, 1980.

[68134]

[68135] 191. Sparkes, R. S.; Sparkes, M. C.; Wilson, M. G.; Towner,
J. W.;
[68136] Benedict, W.; Murphree, A. L.; Yunis, J. J.: Regional assign-
ment
[68137] of genes for human esterase D and retinoblastoma to
chromosome band
[68138] 13q14. (Abstract) Cytogenet. Cell Genet. 25: 209, 1979.

[68139]

[68140] 192. Squire, J.; Gallie, B. L.; Phillips, R. A.: A detailed anal-
ysis
[68141] of chromosomal changes in heritable and non-heritable
retinoblastoma. Hum.
[68142] Genet. 70: 291-301, 1985.

[68143]

[68144] 193. Squire, J.; Phillips, R. A.; Boyce, S.; Godbout, R.;

Rogers,
[68145] B.; Gallie, B. L.: Isochromosome 6p, a unique chromosomal abnormality
[68146] in retinoblastoma: verification by standard staining techniques, new
[68147] densitometric methods, and somatic cell hybridization. Hum. Genet. 66:
[68148] 46-53, 1984.

[68149]

[68150] 194. Stallard, H. B.: The conservation treatment of retinoblastoma. Trans.
[68151] Ophthal. Soc. 82: 473, 1962.

[68152]

[68153] 195. Stone, J. C.; Crosby, J. L.; Kozak, C. A.; Schievella, A. R.;
[68154] Bernards, R.; Nadeau, J. H.: The murine retinoblastoma homolog maps
[68155] to chromosome 14 near Es-10. Genomics 5: 70-75, 1989.

[68156]

[68157] 196. Strong, L. C.; Riccardi, V. M.; Ferrell, R. E.; Sparkes, R. S.

[68158] : Familial retinoblastoma and chromosome 13 deletion transmitted via

[68159] an insertional translocation. Science 213: 1501-1503, 1981.

[68160]

[68161] 197. Taylor, A. I.: Dq-, Dr and retinoblastoma. Human-genetik 10:

[68162] 209-217, 1970.

[68163]

[68164] 198. Thomas, D. M.; Carty, S. A.; Piscopo, D. M.; Lee, J.-S.; Wang,

[68165] W.-F.; Forrester, W. C.; Hinds, P. W.: The retinoblastoma protein

[68166] acts as a transcriptional coactivator required for os-teogenic differentiation. Molec.

[68167] Cell 8: 303-316, 2001.

[68168]

[68169] 199. Toguchida, J.; Ishizaki, K.; Sasaki, M. S.; Nakamura, Y.; Ikenaga,

[68170] M.; Kato, M.; Sugimot, M.; Kotoura, Y.; Yamamuro, T.: Preferential

[68171] mutation of paternally derived RB gene as the initial event in sporadic

[68172] osteosarcoma. Nature 338: 156-158, 1989.

[68173]

[68174] 200. Toguchida, J.; McGee, T. L.; Paterson, J. C.; Eagle, J. R.; Tucker,

[68175] S.; Yandell, D. W.; Dryja, T. P.: Complete genomic se-quence of the

[68176] human retinoblastoma susceptibility gene. Genomics 17: 535-543,

[68177] 1993.

[68178]

[68179] 201. Turleau, C.; de Grouchy, J.; Chavin-Colin, F.; De-spoisses, S.;

[68180] Leblanc, A.: Two cases of del(13q)-retinoblastoma and two cases of

[68181] partial trisomy due to a familial insertion. Ann. Genet. 26: 158-160,

[68182] 1983.

[68183]

[68184] 202. Turleau, C.; de Grouchy, J.; Chavin-Colin, F.; Junien,

C.; Seger,
[68185] J.; Schlienger, P.; Leblanc, A.; Haye, C.: Cytogenetic forms of retinoblastoma:
[68186] their incidence in a survey of 66 patients. Cancer Genet. Cytogenet. 16:
[68187] 321-334, 1985.

[68188]

[68189] 203. Verma, R. S.; Ramesh, K. H.; Samonte, R. V.; Conte, R. A.: Mapping
[68190] the homolog of the human Rb1 gene to chromosome 14 of higher primates. Mammalian
[68191] Genome 7: 591-592, 1996.

[68192]

[68193] 204. Vogel, F.: Genetics of retinoblastoma. Modern Trends in Ophthalmology.
[68194] (pub.) 1968.

[68195]

[68196] 205. Vogel, F.: Genetics of retinoblastoma.In: Genetic Counseling.
[68197] Heidelberg University, Science Library. Trans. by Sabine Kurth.

[68198] New York: Springer Verlag (pub.) 1969.

[68199]

[68200] 206. Vogel, F.: The genetics of retinoblastoma. Hum. Genet. 52:

[68201] 1-54, 1979.

[68202]

[68203] 207. Warburg, M.: Retinoblastoma.In: Goldberg, M. F.: Genetic and

[68204] Metabolic Eye Disease. Boston: Little, Brown and Co. (pub.) 1974.

[68205] Pp. 447-461.

[68206]

[68207] 208. Weichselbaum, R. R.; Beckett, M.; Diamond, A.: Some retinoblastomas,

[68208] osteosarcomas, and soft tissue sarcomas may share a common etiology. Proc.

[68209] Nat. Acad. Sci. 85: 2106-2109, 1988.

[68210]

[68211] 209. Weichselbaum, R. R.; Nove, J.; Little, J. B.: Fibroblasts from

[68212] a D-deletion type retinoblastoma patient are abnormally x-ray sensitive. Nature 266:
[68213] 726-727, 1977.

[68214]

[68215] 210. Weinberg, R. A.: The retinoblastoma protein and cell cycle control. Cell 81:
[68216] 323-330, 1995.

[68217]

[68218] 211. Whyte, P.; Buchkovich, K. J.; Horowitz, J. M.; Friend, S. H.;
[68219] Raybuck, M.; Weinberg, R. A.; Harlow, E.: Association be-tween an
[68220] oncogene and an anti-oncogene: the adenovirus E1A pro-teins bind to
[68221] the retinoblastoma gene product. Nature 334: 124-129, 1988.

[68222]

[68223] 212. Wiggs, J.; Nordenskjold, M.; Yandell, D.; Rapaport, J.; Grondin,
[68224] V.; Janson, M.; Werelius, B.; Petersen, R.; Craft, A.; Riedel, K.;

[68225] Liberfarb, R.; Walton, D.; Wilson, W.; Dryja, T. P.: Prediction of
[68226] the risk of hereditary retinoblastoma, using DNA polymorphisms within
[68227] the retinoblastoma gene. New Eng. J. Med. 318: 151-157, 1988.
[68228]

[68229] 213. Wilson, M. G.; Ebbin, A. J.; Towner, J. W.; Spencer, W. H.:
[68230] Chromosomal anomalies in patients with retinoblastoma. Clin. Genet. 12:
[68231] 1-8, 1977.
[68232]

[68233] 214. Wilson, M. G.; Melnyk, J.; Towner, J. W. J.: Retinoblastoma
[68234] and deletion D(14) syndrome. J. Med. Genet. 6: 322-327, 1969.
[68235]

[68236] 215. Wilson, M. G.; Towner, J. W.; Fujimoto, A.: Retinoblastoma and
[68237] D-chromosome deletions. Am. J. Hum. Genet. 25: 57-61, 1973.

[68238]

[68239] 216. Windle, J. J.; Albert, D. M.; O'Brien, J. M.; Marcus, D. M.;

[68240] Disteche, C. M.; Bernards, R.; Mellon, P. L.: Retinoblastoma in transgenic

[68241] mice. Nature 343: 665-669, 1990.

[68242]

[68243] 217. Yandell, D. W.; Campbell, T. A.; Dayton, S. H.; Pe-tersen, R.;

[68244] Walton, D.; Little, J. B.; McConkie-Rosell, A.; Buckley, E.; Dryja,

[68245] T.: Oncogenic point mutations in the human retinoblas-toma gene: their

[68246] application to genetic counseling. New Eng. J. Med. 321: 1689-1695,

[68247] 1989.

[68248]

[68249] 218. Yokota, J.; Akiyama, T.; Fung, Y.-K. T.; Benedict, W. F.; Namba,

[68250] Y.; Hanaoka, M.; Wada, M.; Terasaki, T.; Shimosato, Y.; Sugimura,

[68251] T.; Terada, M.: Altered expression of the retinoblastoma (RB) gene
[68252] in small-cell carcinoma of the lung. Oncogene 3: 471-475, 1988.
[68253]
[68254] 219. Zeschnigk, M.; Lohmann, D.; Horsthemke, B.: A PCR test for the
[68255] detection of hypermethylated alleles at the retinoblastoma locus. J.
[68256] Med. Genet. 36: 793-794, 1999.
[68257]
[68258] 220. Zhang, H. S.; Postigo, A. A.; Dean, D. C.: Active transcriptional
[68259] repression by the Rb-E2F complex mediates G1 arrest triggered by p16(INK4a),
[68260] TGF-beta, and contact inhibition. Cell 97: 53-61, 1999.
[68261]
[68262] 221. Zhu, X.; Dunn, J. M.; Phillips, R. A.; Goddard, A. D.; Paton,
[68263] K. E.; Becker, A.; Gallie, B. L.: Preferential germline mutation

[68264] of the paternal allele in retinoblastoma. Nature 340:
312-313, 1989.

[68265] 11521.

[68266] Baens, M.; Aerssens, J.; Van Zand, K.; Van den Berghe, H.;
Marynen,

[68267] P.: Isolation and regional assignment of human chromo-
some 12p cDNAs. Genomics 29:

[68268] 44-52, 1995.

[68269] 11522.

[68270] Mao, S.; Neale, G. A. M.; Goorha, R. M.: T-cell oncogene
rhombotin-2

[68271] interacts with retinoblastoma-binding protein 2. Onco-
gene 14: 1531-1539,

[68272] 1997.

[68273] 11523.

[68274] Benbrook, D.; Lernhardt, E.; Pfahl, M.: A new retinoic acid
receptor

[68275] identified from a hepatocellular carcinoma. (Letter) Nature
333:

[68276] 669-672, 1988.

[68277] 11524.

[68278] Brand, N.; Petkovich, M.; Krust, A.; Chambon, P.; de The, H.; Marchio,

[68279] A.; Tiollais, P.; Dejean, A.: Identification of a second human retinoic

[68280] acid receptor. (Letter) Nature 332: 850-853, 1988.

[68281] 11525.

[68282] Dejean, A.; Bougueleret, L.; Grzeschik, K.-H.; Tiollais, P.: Hepatitis

[68283] B virus DNA integration in a sequence homologous to v-erb-A and steroid

[68284] receptor genes in a hepatocellular carcinoma. Nature 322: 70-72,

[68285] 1986.

[68286] 11526.

[68287] de The, H.; del Mar Vivanco-Ruiz, M.; Tiollais, P.; Stunnenberg,

[68288] H.; Dejean, A.: Identification of a retinoic acid responsive element

[68289] in the retinoic acid receptor beta gene. Nature 343: 177-180, 1990.

[68290] 11527.

[68291] Aruffo, A.; Seed, B.: Molecular cloning of a CD28 cDNA by a high-efficiency

[68292] COS cell expression system. Proc. Nat. Acad. Sci. 84: 8573-8577,

[68293] 1987.

[68294] 11528.

[68295] Lafage-Pochitaloff, M.; Costello, R.; Couez, D.; Simonetti, J.;

[68296] Mannoni, P.; Mawas, C.; Olive, D.: Human CD28 and CTLA-4 Ig superfamily

[68297] genes are located on chromosome 2 at bands q33-q34. Immunogenetics 31:

[68298] 198-201, 1990.

[68299] 11529.

[68300] Lee, K. P.; Taylor, C.; Petryniak, B.; Turka, L. A.; June, C. H.;

[68301] Thompson, C. B.: The genomic organization of the CD28 gene: implications

[68302] for the regulation of CD28 mRNA expression and hetero-
geneity. J.
[68303] Immun. 145: 344-352, 1990.

[68304] 11530.

[68305] Lesslauer, W.; Gmunder, H.; Bohlen, P.: Purification and N-
terminal
[68306] amino acid sequence of the human T90/44 (CD28) anti-
gen. Immunogenetics 27:
[68307] 388-391, 1988.

[68308] 11531.

[68309] Okkenhaug, K.; Wu, L.; Garza, K. M.; La Rose, J.; Khoo, W.;
Odermatt,
[68310] B.; Mak, T. W.; Ohashi, P. S.; Rottapel, R.: A point mutation
in
[68311] CD28 distinguishes proliferative signals from survival sig-
nals. Nature
[68312] Immun. 2: 325-332, 2001.

[68313] 11532.

[68314] Dear, T. N.; Colledge, W. H.; Carlton, M. B. L.; Lavenir, I.;
Larson,
[68315] T.; Smith, A. J. H.; Warren, A. J.; Evans, M. J.; Sofroniew, M.

V.;

[68316] Rabbitts, T. H.: The Hox11 gene is essential for cell survival during

[68317] spleen development. Development 121: 2909-2915, 1995.

[68318] 11533.

[68319] Dear, T. N.; Sanchez-Garcia, I.; Rabbitts, T. H.: The HOX11 gene

[68320] encodes a DNA-binding nuclear transcription factor belonging to a

[68321] distinct family of homeobox genes. Proc. Nat. Acad. Sci. 90: 4431-4435,

[68322] 1993.

[68323] 11534.

[68324] Dube, I. D.; Kamel-Reid, S.; Yuan, C. C.; Lu, M.; Wu, X.; Corpus,

[68325] G.; Raimondi, S. C.; Crist, W. M.; Carroll, A. J.; Minowada, J.; Baker,

[68326] J. B.: A novel human homeobox gene lies at the chromosome 10 breakpoint

[68327] in lymphoid neoplasias with chromosomal translocation t(10;14). Blood 78:

[68328] 2996-3003, 1991.

[68329] 11535.

[68330] Hatano, M.; Roberts, C. W. M.; Minden, M.; Crist, W. M.; Korsmeyer,

[68331] S. J.: Deregulation of a homeobox gene, HOX11, by the t(10;14) in

[68332] T cell leukemia. Science 253: 79-82, 1991.

[68333] 11536.

[68334] Kagan, J.; Finan, J.; Letofsky, J.; Besa, E. C.; Nowell, P. C.;

[68335] Croce, C. M.: Alpha-chain locus of the T-cell antigen re-ceptor is

[68336] involved in the t(10;14) chromosome translocation of T-cell acute

[68337] lymphocytic leukemia. Proc. Nat. Acad. Sci. 84: 4543-4546, 1987.

[68338] 11537.

[68339] Kennedy, M. A.; Gonzalez-Sarmiento, R.; Kees, U. R.; Lam-pert, F.;

[68340] Dear, N.; Boehm, T.; Rabbitts, T. H.: HOX11, a homeobox-containing

[68341] T-cell oncogene on human chromosome 10q24. Proc.

Nat. Acad. Sci. 88:
[68342] 8900-8904, 1991.

[68343] 11538.

[68344] Lu, M.; Gong, Z. Y.; Shen, W. F.; Ho, A. D.: The TCL-3 proto-oncogene
[68345] altered by chromosomal translocation in T-cell leukemia codes for
[68346] a homeobox protein. EMBO J. 10: 2905-2910, 1991.

[68347] 11539.

[68348] Roberts, C. W. M.; Shutter, J. R.; Korsmeyer, S. J.: Hox11 controls
[68349] the genesis of the spleen. Nature 368: 747-750, 1994.

[68350] 11540.

[68351] Zutter, M.; Hockett, R. D.; Roberts, C. W. M.; McGuire, E. A.;
[68352] Bloomstone, J.; Morton, C. C.; Deaven, L. L.; Crist, W. M.; Carroll,
[68353] A. J.; Korsmeyer, S. J.: The t(10;14)(q24;q11) of T-cell acute lymphoblastic
[68354] leukemia juxtaposes the delta T-cell receptor with TCL3, a conserved

[68355] and activated locus at 10q24. Proc. Nat. Acad. Sci. 87:
3161-3165,
[68356] 1990.

[68357] 11541.

[68358] Alarcon, B.; Regueiro, J. R.; Arnaiz-Villena, A.; Terhorst, C.

[68359] : Familial defect in the surface expression of the T-cell re-
ceptor-CD3
[68360] complex. New Eng. J. Med. 319: 1203-1208, 1988.

[68361] 11542.

[68362] Caplan, S.; Zeliger, S.; Wang, L.; Baniyash, M.: Cell-
surface-expressed
[68363] T-cell antigen-receptor epsilon chain is associated with
the cytoskeleton. Proc.
[68364] Nat. Acad. Sci. 92: 4768-4772, 1995.

[68365] 11543.

[68366] Clevers, H.; Alarcon, B.; Wileman, T.; Terhorst, C.: The T
cell
[68367] receptor/CD3 complex: a dynamic protein ensemble.
Annu. Rev. Immun. 6:
[68368] 629-662, 1988.

[68369] 11544.

[68370] Grakoui, A.; Bromley, S. K.; Sumen, C.; Davis, M. M.; Shaw, A.

[68371] S.; Allen, P. M.; Dustin, M. L.: The immunological synapse: a molecular

[68372] machine controlling T cell activation. Science 285: 221-227, 1999.

[68373] 11545.

[68374] Krummel, M. F.; Sjaastad, M. D.; Wulfing, C.; Davis, M. M.: Differential

[68375] clustering of CD4 and CD3-zeta during T cell recognition. Science 289:

[68376] 1349-1352, 2000.

[68377] 11546.

[68378] Weissman, A. M.; Baniyash, M.; Hou, D.; Samelson, L. E.; Burgess,

[68379] W. H.; Klausner, R. D.: Molecular cloning of the zeta chain of the

[68380] T cell antigen receptor. Science 239: 1018-1021, 1988.

[68381] 11547.

[68382] Weissman, A. M.; Hou, D.; Orloff, D. G.; Modi, W. S.;

Seuanez,
[68383] H.; O'Brien, S. J.; Klausner, R. D.: Molecular cloning and chromosomal
[68384] localization of the human T-cell receptor zeta chain: distinction
[68385] from the molecular CD3 complex. Proc. Nat. Acad. Sci. 85: 9709-9713,
[68386] 1988.

[68387] 11548.

[68388] Weissman, A. M.; Samelson, L. E.; Klausner, R. D.: A new subunit
[68389] of the human T-cell antigen receptor complex. Nature 324: 480-482,
[68390] 1986.

[68391] 11549.

[68392] Yu, C. Y.; Milstein, C.: A physical map linking the five CD1 human
[68393] thymocyte differentiation antigen genes. EMBO J. 8: 3727-3732, 1989.
[68394] 11550.

[68395] Zeng, Z.-H.; Castano, A. R.; Segelke, B. W.; Stura, E. A.;

Peterson,
[68396] P. A.; Wilson, I. A.: Crystal structure of mouse CD1: an MHC-like
[68397] fold with a large hydrophobic binding groove. Science 277: 339-345,
[68398] 1997.

[68399] 11551.

[68400] Demczuk, S.; Aledo, R.; Zucman, J.; Delattre, O.; Desmaze, C.;
[68401] Dauphinot, L.; Jalbert, P.; Rouleau, G. A.; Thomas, G.; Aurias, A.
[68402] : Cloning of a balanced translocation breakpoint in the DiGeorge syndrome
[68403] critical region and isolation of a novel potential adhesion receptor
[68404] gene in its vicinity. Hum. Molec. Genet. 4: 551-558, 1995.

[68405] 11552.

[68406] Moretti, P. A. B.; Davidson, A. J.; Baker, E.; Lilley, B.; Zon,
[68407] L. I.; D'Andrea, R. J.: Molecular cloning of a human Vent-like homeobox
[68408] gene. Genomics 76: 21-29, 2001.

[68409] 11553.

[68410] Kim, Y. J.; Kim, K.-S.; Do, S.; Kim, C.-H.; Kim, S.-K.; Lee, Y.-C.

[68411] : Molecular cloning and expression of human alpha-2,8-sialyltransferase

[68412] (hST8Sia V). Biochem. Biophys. Res. Commun. 235: 327-330, 1997.

[68413] 11554.

[68414] Jaju, R. J.; Haas, O. A.; Neat, M.; Harbott, J.; Saha, V.; Boultwood,

[68415] J.; Brown, J. M.; Pirc-Danoewinata, H.; Krings, B. W.; Muller, U.;

[68416] Morris, S. W.; Wainscoat, J. S.; Kearney, L.: A new recurrent translocation,

[68417] t(5;11)(q35p15.5), associated with del(5q) in childhood acute myeloid

[68418] leukemia. Blood 94: 773-780, 1999.

[68419] 11555.

[68420] Kurotaki, N.; Harada, N.; Yoshiura, K.; Sugano, S.; Niikawa, N.;

[68421] Matsumoto, N.: Molecular characterization of NSD1, a human homologue

[68422] of the mouse Nsd1 gene. Gene 279: 197-204, 2001.

[68423] 11556.

[68424] Wang, X.; Yeh, S.; Wu, G.; Hsu, C.-L.; Wang, L.; Chiang, T.; Yang,

[68425] Y.; Guo, Y.; Chang, C.: Identification and characterization of a

[68426] novel androgen receptor coregulator ARA267-alpha in prostate cancer

[68427] cells. J. Biol. Chem. 276: 40417-40423, 2001.

[68428] 11557.

[68429] Lane, P. W.; Green, E. L.: Pale ear and light ear in the house

[68430] mouse: mimic mutations in linkage groups XII and XVII. J. Hered. 58:

[68431] 17-20, 1967.

[68432] 11558.

[68433] Riewald, M.; Petrovan, R. J.; Donner, A.; Mueller, B. M.; Ruf,

[68434] W.: Activation of endothelial cell protease activated receptor 1

[68435] by the protein C pathway. Science 296: 1880-1882, 2002.

[68436] 11559.

[68437] Park, B.-S.; Park, J.-S.; Lee, D. Y.; Youn, J. I.; Kim, I. G.

[68438] : Vitamin D receptor polymorphism is associated with psoriasis. J.

[68439] Invest. Derm. 112: 113-116, 1999.

[68440] 11560.

[68441] Jankowski, S. A.; De Jong, P.; Meltzer, P. S.: Genomic structure

[68442] of SAS, a member of the transmembrane 4 superfamily amplified in human

[68443] sarcomas. Genomics 25: 501-506, 1995.

[68444] 11561.

[68445] Meltzer, P. S.; Jankowski, S. A.; Dal Cin, P.; Sandberg, A. A.;

[68446] Paz, I. B.; Coccia, M. A.; Smith, S. H.: Identification and cloning

[68447] of a novel amplified DNA sequence in human malignant fibrous histiocytoma

[68448] derived from a region of chromosome 12 frequently rearranged in soft

[68449] tissue tumors. (Abstract) Cytogenet. Cell Genet. 58: 1979

only,

[68450] 1991.

[68451] 11562.

[68452] Wolfe, L. C.; John, K. M.; Falcone, J. C.; Byrne, A. M.; Lux,

[68453] S. E.: A genetic defect in the binding of protein 4.1 to spectrin

[68454] in a kindred with hereditary spherocytosis. New Eng. J. Med. 307:

[68455] 1367-1374, 1982.

[68456] 11563.

[68457] Shao, X.; Tarnasky, H. A.; Schalles, U.; Oko, R.; van der Hoorn,

[68458] F. A.: Interactional cloning of the 84-kDa major outer dense fiber

[68459] protein Odf84: leucine zippers mediate associations of Odf84 and Odf27. J.

[68460] Biol. Chem. 272: 6105-6113, 1997.

[68461] 11564.

[68462] Shao, X.; van der Hoorn, F. A.: Self-interaction of the ma-jor

[68463] 27-kilodalton outer dense fiber protein is in part medi-

ated by a leucine
[68464] zipper domain in the rat. Biol. Reprod. 55: 1343-1350, 1996.
[68465] 11565.

[68466] van der Hoorn, F. A.; Tarnasky, H. A.; Nordeen, S. K.: A new rat
[68467] gene RT7 is specifically expressed during spermatogenesis. Dev. Biol. 142:
[68468] 147-154, 1990.

[68469] 11566.

[68470] Ammer, H.; Henschen, A.; Lee, C.-H.: Isolation and amino-acid
[68471] sequence analysis of human sperm protamines P1 and P2: occurrence
[68472] of two forms of protamine P2. Biol. Chem. Hoppe-Seyler 367: 515-522,
[68473] 1986.

[68474] 11567.

[68475] Cho, C.; Willis, W. D.; Goulding, E. H.; Jung-Ha, H.; Choi, Y.-C.;
[68476] Hecht, N. B.; Eddy, E. M.: Haploinsufficiency of protamine-

1 or -2

[68477] causes infertility in mice. Nature Genet. 28: 82-86, 2001.

[68478] 11568.

[68479] Choudhary, S. K.; Wykes, S. M.; Kramer, J. A; Mohamed, A. N.; Koppitch,

[68480] F.; Nelson, J. E.; Krawetz, S. A.: A haploid expressed gene cluster

[68481] exists as a single chromatin domain in human sperm. J. Biol. Chem. 270:

[68482] 8755-8762, 1995.

[68483] 11569.

[68484] Domenjoud, L.; Nussbaum, G.; Adham, I. M.; Greeske, G.; Engel,

[68485] W.: Genomic sequences of human protamines whose genes, PRM1 and PRM2,

[68486] are clustered. Genomics 8: 127-133, 1990.

[68487] 11570.

[68488] Engel, W.: Personal Communication. Goettingen, Germany 5/17/1990.

[68489] 11571.

[68490] Hecht, N. B.; Kleene, K. C.; Yelick, P. C.; Johnson, P. A.;

Pravtcheva,

[68491] D. D.; Ruddle, F. H.: Mapping of haploid expressed genes: genes for

[68492] both mouse protamines are located on chromosome 16. Somat. Cell Molec.

[68493] Genet. 12: 203-208, 1986.

[68494] 11572.

[68495] Krawetz, S. A.; Herfort, M. H.; Hamerton, J. L.; Pon, R. T.; Dixon,

[68496] G. H.: Chromosomal localization and structure of the human P1 protamine

[68497] gene. Genomics 5: 639-645, 1989.

[68498] 11573.

[68499] Reeves, R. H.; Gallahan, D.; O'Hara, B. F.; Callahan, R.; Gearhart,

[68500] J. D.: Genetic mapping of Prm-1, Igl-1, Smst, Mtv-6, Sod-1, and Ets-2

[68501] and localization of the Down syndrome region on mouse chromosome 16. Cytogenet.

[68502] Cell Genet. 44: 76-81, 1987.

[68503] 11574.

[68504] Reeves, R. H.; Gearhart, J. D.; Hecht, N. B.; Yelick, P.; John-
son,
[68505] P.; O'Brien, S. J.: The gene encoding protamine 1 is located
on chromosome
[68506] 16 in humans and near the proximal end of mouse chro-
mosome 16, where
[68507] it is tightly linked to protamine 2. (Abstract) Cytogenet.
Cell Genet. 46:
[68508] 680 only, 1987.

[68509] 11575.

[68510] Reeves, R. H.; Gearhart, J. D.; Hecht, N. B.; Yelick, P.; John-
son,
[68511] P.; O'Brien, S. J.: The gene encoding protamine 1 is located
on human
[68512] chromosome 16, and near the proximal end of mouse
chromosome 16 where
[68513] it is tightly linked to the gene encoding protamine 2. J.
Hered. 80:
[68514] 442-446, 1989.

[68515] 11576.

[68516] Viguie, F.; Domenjoud, L.; Rousseau-Merck, M.-F.;
Dadoune, J.-P.;

[68517] Chevaillier, P.: Chromosomal localization of the human protamine
[68518] genes, PRM1 and PRM2, to 16p13.3 by in situ hybridization. Hum. Genet. 85:
[68519] 171-174, 1990.

[68520] 11577.

[68521] Lee, H. S.; Sambuughin, N.; Cervenakova, L.; Chapman, J.; Pocchiari,
[68522] M.; Litvak, S.; Qi, H. Y.; Budka, H.; del Ser, T.; Furukawa, H.; Brown,
[68523] P.; Gajdusek, D. C.; Long, J. C.; Korczyn, A. D.; Goldfarb, L. G.
[68524] : Ancestral origins and worldwide distribution of the PRNP 200K mutation
[68525] causing familial Creutzfeldt-Jakob disease. Am. J. Hum. Genet. 64:
[68526] 1063-1070, 1999.

[68527] 11578.

[68528] de The, H.; Marchio, A.; Tiollais, P.; Dejean, A.: A novel steroid
[68529] thyroid hormone receptor-related gene inappropriately expressed in

[68530] human hepatocellular carcinoma. Nature 330: 667-670, 1987.

[68531] 11579.

[68532] Clevers, H. C.; Dunlap, S.; Wileman, T. E.; Terhorst, C.: Human

[68533] CD3-epsilon gene contains three miniexons and is transcribed from

[68534] a non-TATA promoter. Proc. Nat. Acad. Sci. 85: 8156-8160, 1988.

[68535] 11580.

[68536] DeJarnette, J. B.; Sommers, C. L.; Huang, K.; Woodside, K. J.;

[68537] Emmons, R.; Katz, K.; Shores, E. W.; Love, P. E.: Specific requirement

[68538] for CD3-epsilon in T cell development. Proc. Nat. Acad. Sci. 95:

[68539] 14909-14914, 1998.

[68540] 11581.

[68541] Gold, D. P.; Puck, J. M.; Pettey, C. L.; Cho, M.; Coligan, J.;

[68542] Woody, J. N.; Terhorst, C.: Isolation of cDNA clones encoding the

[68543] 20K non-glycosylated polypeptide chain of the human T-cell receptor/T3
[68544] complex. Nature 321: 431-434, 1986.

[68545] 11582.

[68546] Le Deist, F.; Thoenes, G.; Corado, J.; Lisowska-Grospierre, B.;
[68547] Fischer, A.: Immunodeficiency with low expression of the T cell receptor/CD3
[68548] complex: effect on T lymphocyte activation. Europ. J. Immun. 21:
[68549] 1641-1647, 1991.

[68550] 11583.

[68551] Soudais, C.; de Villartay, J.-P.; Le Deist, F.; Fischer, A.; Lisowska-Grospierre,
[68552] B.: Independent mutations of the human CD3-epsilon gene resulting
[68553] in a T cell receptor/CD3 complex immunodeficiency. Nature Genet. 3:
[68554] 77-81, 1993.

[68555] 11584.

[68556] Thoenes, G.; Soudais, C.; Le Deist, F.; Griscelli, C.; Fischer,

[68557] A.; Lisowska-Grospierre, B.: Structural analysis of low TCR-CD3 complex
[68558] expression in T cells of an immunodeficient patient. J. Biol. Chem. 267:
[68559] 487-493, 1992.

[68560] 11585.

[68561] Wong, S.; Moore, S.; Orisio, S.; Millward, A.; Demaine, A. G.:
[68562] Susceptibility to type I diabetes in women is associated with the
[68563] CD3 epsilon locus on chromosome 11. Clin. Exp. Immun. 83: 69-73,
[68564] 1991.

[68565] 11586.

[68566] Andria, M. L.; Hsieh, C. L.; Oren, R.; Francke, U.; Levy, S.:

[68567] Genomic organization and chromosomal localization of the TAPA-1 gene. J.
[68568] Immun. 147: 1030-1036, 1991.

[68569] 11587.

[68570] Boismenu, R.; Rhein, M.; Fischer, W. H.; Havran, W. L.: A role

[68571] for CD81 in early T cell development. Science 271: 198-200, 1996.

[68572] 11588.

[68573] Crotta, S.; Stilla, A.; Wack, A.; D'Andrea, A.; Nuti, S.; D'Oro,

[68574] U.; Mosca, M.; Filliponi, F.; Brunetto, R. M.; Bonino, F.; Abrignani,

[68575] S.; Valiante, N. M.: Inhibition of natural killer cells through engagement

[68576] of CD81 by the major hepatitis C virus envelope protein. J. Exp.

[68577] Med. 195: 35-41, 2002.

[68578] 11589.

[68579] Oren, R.; Takahashi, S.; Doss, C.; Levy, R.; Levy, S.: TAPA-1,

[68580] the target of an antiproliferative antibody, defines a new family

[68581] of transmembrane proteins. Molec. Cell. Biol. 10: 4007-4015, 1990.

[68582] 11590.

[68583] Pileri, P.; Uematsu, Y.; Campagnoli, S.; Galli, G.; Falugi, F.;

[68584] Petracca, R.; Weiner, A. J.; Houghton, M.; Rosa, D.; Grandi,

G.; Abrignani,
[68585] S.: Binding of hepatitis C virus to CD81. Science 282:
938-941,
[68586] 1998.

[68587] 11591.

[68588] Tseng, C.-T. K.; Klimpel, G. R.: Binding of the hepatitis C
virus
[68589] envelope protein E2 to CD81 inhibits natural killer cell
functions. J.
[68590] Exp. Med. 195: 43-49, 2002.

[68591] 11592.

[68592] Virtaneva, K. I.; Emi, N.; Marken, J. S.; Aruffo, A.; Jones, C.;

[68593] Spurr, N. K.; Schroder, J. P.: Chromosomal localization of
three
[68594] human genes coding for A15, L6, and S5.7 (TAPA1): all
members of the
[68595] transmembrane 4 superfamily of proteins. Immunogenet-
ics 39: 329-334,
[68596] 1994.

[68597] 11593.

[68598] Patel, A.; Rochelle, J. M.; Jones, J. M.; Sumegi, J.; Uhl, G. R.;

[68599] Seldin, M. F.; Meisler, M. H.; Gregor, P.: Mapping of the taurine
[68600] transporter gene to mouse chromosome 6 and to the short arm of human
[68601] chromosome 3. Genomics 25: 314-317, 1995.

[68602] 11594.

[68603] Ramamoorthy, S.; Leibach, F. H.; Mahesh, V. B.; Han, H.; Yang-Feng,
[68604] T.; Blakely, R. D.; Ganapathy, V.: Functional characterization and
[68605] chromosomal localization of a cloned taurine transporter from human
[68606] placenta. Biochem. J. 300: 893-900, 1994.

[68607] 11595.

[68608] Liao, Y.-C. J.; Lebo, J.; Lebo, R. V.; Clawson, G. A.; Smuckler,
[68609] E. A.: Human prion protein cDNA: molecular cloning, chromosomal mapping,
[68610] and biological implications. Science 233: 364-367, 1986.

[68611] 11596.

[68612] Lindquist, S.: Mad cows meet Psi-chotic yeast: the expan-

sion
[68613] of the prion hypothesis. Cell 89: 495-498, 1997.

[68614] 11597.

[68615] Little, B. W.; Brown, B. W.; Rodgers-Johnson, P.; Perl, D. P.;

[68616] Gajdusek, D. C.: Familial myoclonic dementia masquerading as Creutzfeldt-Jakob
[68617] disease. Ann. Neurol. 20: 231-239, 1986.

[68618] 11598.

[68619] Lloyd, S. E.; Onwuazor, O. N.; Beck, J. A.; Mallinson, G.; Farrall,
[68620] M.; Targonski, P.; Collinge, J.; Fisher, E. M. C.: Identification
[68621] of multiple quantitative trait loci linked to prion disease incubation
[68622] period in mice. Proc. Nat. Acad. Sci. 98: 6279-6283, 2001.

[68623] 11599.

[68624] Lugaresi, E.; Medori, R.; Montagna, P.; Baruzzi, A.; Cortelli,

[68625] P.; Lugaresi, A.; Tinuper, P.; Zucconi, M.; Gambetti, P.: Fatal familial
[68626] insomnia and dysautonomia with selective degeneration

of thalamic
[68627] nuclei. New Eng. J. Med. 315: 997-1003, 1986.

[68628] 11600.

[68629] Lugaresi, E.; Montagna, P.; Baruzzi, A.; Cortelli, P.; Tinu-
per,
[68630] P.; Zucconi, M.; Gambetti, P. L.; Medori, R.: Insomnie fa-
miliale
[68631] a evolution maligne: une nouvelle maladie thalamique.
Rev. Neurol. 142:
[68632] 791-792, 1986.

[68633] 11601.

[68634] Mahal, S. P.; Asante, E. A.; Antoniou, M.; Collinge, J.: Isola-
tion
[68635] and functional characterisation of the promoter region of
the human
[68636] prion protein gene. Gene 268: 105-114, 2001.

[68637] 11602.

[68638] Mallucci, G. R.; Ratte, S.; Asante, E. A.; Linehan, J.; Gow-
land,
[68639] I.; Jefferys, J. G. R.; Collinge, J.: Post-natal knockout of
prion

[68640] protein alters hippocampal CA1 properties, but does not result in
[68641] neurodegeneration. EMBO J. 21: 202-210, 2002.

[68642] 11603.

[68643] Mallucci, G. R.; Campbell, T. A.; Dickinson, A.; Beck, J.; Holt,
[68644] M.; Plant, G.; de Pauw, K. W.; Hakin, R. N.; Clarke, C. E.; Howell,
[68645] S.; Davies-Jones, G. A. B.; Lawden, M.; Smith, C. M. L.; Ince, P.;
[68646] Ironside, J. W.; Bridges, L. R.; Dean, A.; Weeks, I.; Collinge, J.
[68647] : Inherited prion disease with an alanine to valine muta-tion at codon
[68648] 117 in the prion protein gene. Brain 122: 1823-1837, 1999.

[68649] 11604.

[68650] Manetto, V.; Medori, R.; Cortelli, P.; Montagna, P.; Tinuper,

[68651] P.; Baruzzi, A.; Rancurel, G.; Hauw, J.-J.; Vanderhaeghen, J.-J.;
[68652] Mailleux, P.; Bugiani, O.; Tagliavini, F.; Bouras, C.; Rizzuto, N.;

[68653] Lugaresi, E.; Gambetti, P.: Fatal familial insomnia: clinical and
[68654] pathologic study of 5 new cases. Neurology 42: 312-319, 1992.
[68655] 11605.

[68656] Manolakou, K.; Beaton, J.; McConnell, I.; Farquar, C.; Man-
son,
[68657] J.; Hastie, N. D.; Bruce, M.; Jackson, I. J.: Genetic and envi-
ronmental
[68658] factors modify bovine spongiform encephalopathy incu-
bation period
[68659] in mice. Proc. Nat. Acad. Sci. 98: 7402-7407, 2001.

[68660] 11606.

[68661] Manson, J. C.; Clarke, A. R.; McBride, P. A.; McConnell, I.;
Hope,
[68662] J.: PrP gene dosage determines the timing but not the final
intensity
[68663] or distribution of lesions in scrapie pathology. Neurode-
generation 3:
[68664] 331-340, 1994.

[68665] 11607.

[68666] Mastrianni, J. A.; Capellari, S.; Telling, G. C.; Han, D.; Bosque,
[68667] P.; Prusiner, S. B.; DeArmond, S. J.: Inherited prion disease caused
[68668] by the V201I mutation: transmission to transgenic mice. Neurology 57:
[68669] 2198-2205, 2001.

[68670] 11608.

[68671] Mastrianni, J. A.; Curtis, M. T.; Oberholtzser, J. C.; Da Costa,
[68672] M. M.; DeArmond, S.; Prusiner, S. B.; Garbern, J. Y.: Prion disease
[68673] (PrP-A117V) presenting with ataxia instead of dementia. Neurology 45:
[68674] 2042-2050, 1995.

[68675] 11609.

[68676] Mead, S.; Mahal, S. P.; Beck, J.; Campbell, T.; Farrall, M.; Fisher,
[68677] E.; Collinge, J.: Sporadic--but not variant--Creutzfeldt-Jakob disease
[68678] is associated with polymorphisms upstream of PRNP exon 1. Am. J.

[68679] Hum. Genet. 69: 1225-1235, 2001.

[68680] 11610.

[68681] Medori, R.: Personal Communication. New York, N. Y. 5/17/1990.

[68682]

[68683] 100. Medori, R.; Montagna, P.; Tritschler, H. J.; LeBlanc, A.; Cortelli,

[68684] P.; Tinuper, P.; Lugaresi, E.; Gambetti, P.: Fatal familial insomnia:

[68685] a second kindred with mutation of prion protein gene at codon 178. Neurology 42:

[68686] 669-670, 1992.

[68687]

[68688] 101. Medori, R.; Tritschler, H.-J.: Prion protein gene analysis in

[68689] three kindreds with fatal familial insomnia (FFI): codon 178 mutation

[68690] and codon 129 polymorphism. Am. J. Hum. Genet. 53: 822-827, 1993.

[68691]

[68692] 102. Medori, R.; Tritschler, H.-J.; LeBlanc, A.; Villare, F.;

Manetto,
[68693] V.; Chen, H. Y.; Xue, R.; Leal, S.; Montagna, P.; Cortelli, P.; Tinuper,
[68694] P.; Avoni, P.; Mochi, M.; Baruzzi, A.; Hauw, J. J.; Ott, J.; Lugaresi,
[68695] E.; Autilio-Gambetti, L.; Gambetti, P.: Fatal familial insomnia,
[68696] a prion disease with a mutation at codon 178 of the prion protein
[68697] gene. New Eng. J. Med. 326: 444-449, 1992.

[68698]

[68699] 103. Meggendorfer, F.: Klinische und genealogische Beobachtungen
[68700] bei einem Fall von spastischer Pseudosklerose Jakobs. Z. Ges. Neurol.
[68701] Psychiat. 128: 337-341, 1930.

[68702]

[68703] 104. Meiner, Z.; Gabizon, R.; Prusiner, S. B.: Familial Creutzfeldt-Jakob
[68704] disease: codon 200 prion disease in Libyan Jews. Medicine 76: 227-237,
[68705] 1997.

[68706]

[68707] 105. Mestel, R.: Putting prions to the test. Science 273: 184-189,

[68708] 1996.

[68709]

[68710] 106. Miele, G.; Jeffrey, M.; Turnbull, D.; Manson, J.; Clinton, M.

[68711] : Ablation of cellular prion protein expression affects mitochondrial

[68712] numbers and morphology. Biochem. Biophys. Res. Commun. 291: 372-377,

[68713] 2002.

[68714]

[68715] 107. Mishra, R. S.; Gu, Y.; Bose, S.; Verghese, S.; Kalepu, S.; Singh,

[68716] N.: Cell surface accumulation of a truncated transmembrane prion

[68717] protein in Gerstmann-Straussler-Scheinker disease P102L. J. Biol.

[68718] Chem. 277: 24554-24561, 2002.

[68719]

[68720] 108. Mitrova, E.; Lowenthal, A.; Appeal, B.: Familial Creutzfeldt-Jakob
[68721] disease with temporal and spatial separation of affected members. Europ.
[68722] J. Epidemiol. 6: 233-238, 1990.

[68723]

[68724] 109. Monari, L.; Chen, S. G.; Brown, P.; Parchi, P.; Pe-
tersen, R.
[68725] B.; Mikol, J.; Gray, F.; Cortelli, P.; Montagna, P.; Ghetti, B.; Goldfarb,
[68726] L. G.; Gajdusek, D. C.; Lugaresi, E.; Gambetti, P.; Autilio-Gambetti,
[68727] L.: Fatal familial insomnia and familial Creutzfeldt-Jakob disease:
[68728] different prion proteins determined by a DNA polymor-
phism. Proc.
[68729] Nat. Acad. Sci. 91: 2839-2842, 1994.

[68730]

[68731] 110. Montrasio, F.; Frigg, R.; Glatzel, M.; Klein, M. A.; Mackay,
[68732] F.; Aguzzi, A.; Weissmann, C.: Impaired prion replication in spleens

[68733] of mice lacking functional follicular dendritic cells. Science 288:
[68734] 1257-1259, 2000.

[68735]

[68736] 111. Moore, R. C.; Xiang, F.; Monaghan, J.; Han, D.; Zhang, Z.; Edstrom,
[68737] L.; Anvret, M.; Prusiner, S. B.: Huntington disease pheno-copy is
[68738] a familial prion disease. Am. J. Hum. Genet. 69: 1385-1388, 2001.

[68739]

[68740] 112. Mouillet-Richard, S.; Ermonval, M.; Chebassier, C.; Laplanche,
[68741] J. L.; Lehmann, S.; Launay, J. M.; Kellermann, O.: Signal transduction
[68742] through prion protein. Science 289: 1925-1928, 2000.

[68743]

[68744] 113. Mouillet-Richard, S.; Teil, C.; Lenne, M.; Hugon, S.; Taleb,
[68745] O.; Laplanche, J.-L.: Mutation at codon 210 (V210I) of the prion

[68746] protein gene in a North African patient with Creutzfeldt-Jakob disease. J.

[68747] Neurol. Sci. 168: 141-144, 1999.

[68748]

[68749] 114. Nieto, A.; Goldfarb, L. G.; Brown, P.; McCombie, W. R.; Trapp,

[68750] S.; Asher, D. M.; Gajdusek, D. C.: Codon 178 mutation in ethnically

[68751] diverse Creutzfeldt-Jakob disease families. (Letter) Lancet 337:

[68752] 622-623, 1991.

[68753]

[68754] 115. Nitrini, R.; Rosemberg, S.; Passos-Bueno, M. R.; da Silva, L.

[68755] S. T.; Iughetti, P.; Papadopoulos, M.; Carrilho, P. M.; Caramelli,

[68756] P.; Albrecht, S.; Zatz, M.; LeBlanc, A.: Familial spongiform encephalopathy

[68757] associated with a novel prion protein gene mutation. Ann. Neurol. 42:

[68758] 138-146, 1997.

[68759]

[68760] 116. Oesch, B.; Westaway, D.; Walchli, M.; McKinley, M. P.; Kent,

[68761] S. B. H.; Aebersold, R.; Barry, R. A.; Tempst, P.; Teplow, D. B.;

[68762] Hood, L. E.; Prusiner, S. B.; Weissmann, C.: A cellular gene encodes

[68763] scrapie PrP 27-30 protein. Cell 40: 735-746, 1985.

[68764]

[68765] 117. Owen, F.; Poulter, M.; Collinge, J.; Crow, T. J.: Codon 129

[68766] changes in the prion protein gene in Caucasians. (Letter) Am. J.

[68767] Hum. Genet. 46: 1215-1216, 1990.

[68768]

[68769] 118. Owen, F.; Poulter, M.; Collinge, J.; Crow, T. J.: A codon 129

[68770] polymorphism in the PRIP gene. Nucleic Acids Res. 18: 3103, 1990.

[68771]

[68772] 119. Owen, F.; Poulter, M.; Collinge, J.; Leach, M.; Loft-

house, R.;

[68773] Crow, T. J.; Harding, A. E.: A dementing illness associated with

[68774] a novel insertion in the prion protein gene. Molec. Brain Res. 13:

[68775] 155-157, 1992.

[68776]

[68777] 120. Owen, F.; Poulter, M.; Lofthouse, R.; Collinge, J.; Crow, T.

[68778] J.; Risby, D.; Baker, H. F.; Ridley, R. M.; Hsiao, K.; Prusiner, S.

[68779] B.: Insertion in prion protein gene in familial Creutzfeldt-Jakob

[68780] disease. (Letter) Lancet I: 51-52, 1989.

[68781]

[68782] 121. Owen, F.; Poulter, M.; Shah, T.; Collinge, J.; Loft-house, R.;

[68783] Baker, H.; Ridley, R.; McVey, J.; Crow, T. J.: An in-frame in-sertion

[68784] in the prion protein gene in familial Creutzfeldt-Jakob disease. Molec.

[68785] Brain Res. 7: 273-276, 1990.

[68786]

[68787] 122. Pablos-Mendez, A.; Netto, E. M.; Defendini, R.: Infec-
tious prions
[68788] or cytotoxic metabolites? Lancet 341: 159-161, 1993.

[68789]

[68790] 123. Palmer, M. S.; Collinge, J.: Mutations and polymor-
phisms in
[68791] the prion protein gene. Hum. Mutat. 2: 168-173, 1993.

[68792]

[68793] 124. Palmer, M. S.; Dryden, A. J.; Hughes, J. T.; Collinge, J.:
Homozygous
[68794] prion protein genotype predisposes to sporadic
Creutzfeldt-Jakob disease. Nature 352:
[68795] 340-342, 1991. Note: Erratum: Nature 352: 547 only,
1991.

[68796]

[68797] 125. Panegyres, P. K.; Toufexis, K.; Kakulas, B. A.;
Cernevakova,
[68798] L.; Brown, P.; Ghetti, B.; Piccardo, P.; Dlouhy, S. R.: A new
PRNP
[68799] mutation (G131V) associated with Gerstmann-Straus-

sler-Scheinker disease. Arch.
[68800] Neurol. 58: 1899-1902, 2001.

[68801]

[68802] 126. Peretz, D.; Williamson, R. A.; Kaneko, K.; Vergara, J.; Leclerc,
[68803] E.; Schmitt-Ulms, G.; Mehlhorn, I. R.; Legname, G.; Wormald, M. R.;
[68804] Rudd, P. M.; Dwek, R. A.; Burton, D. R.; Prusiner, S. B.: An-
tibodies
[68805] inhibit prion propagation and clear cell cultures of prion infectivity. Nature 412:
[68806] 739-743, 2001.

[68807]

[68808] 127. Perry, R. T.; Go, R. C. P.; Harrell, L. E.; Acton, R. T.: SSCP
[68809] analysis and sequencing of the human prion protein gene (PRNP) detects
[68810] two different 24 bp deletions in an atypical Alzheimer's disease family. Am.
[68811] J. Med. Genet. 60: 12-18, 1995.

[68812]

[68813] 128. Petchanikow, C.; Saborio, G. P.; Anderes, L.; Frossard, M.-J.;
[68814] Olmedo, M. I.; Soto, C.: Biochemical and structural studies of the
[68815] prion protein polymorphism. FEBS Lett. 509: 451-456, 2001.
[68816]

[68817] 129. Plaitakis, A.; Viskadouraki, A. K.; Tzagournissakis, M.; Zaganas,
[68818] I.; Verghese-Nikolakaki, S.; Karagiorgis, V.; Panagiotides, I.; Kilindireas,
[68819] C.; Patsouris, E.; Haberler, C.; Budka, H.; Sklaviadis, T.: In-creased
[68820] incidence of sporadic Creutzfeldt-Jakob disease on the is-land of Crete
[68821] associated with a high rate of PRNP 129-methionine ho-mozygosity in
[68822] the local population. Ann. Neurol. 50: 227-233, 2001.

[68823]

[68824] 130. Pocchiari, M.; Salvatore, M.; Cutruzzola, F.; Genuardi, M.; Allcatelli,
[68825] C. T.; Masullo, C.; Macchi, G.; Alema, G.; Galgani, S.; Xi, Y.

G.;

[68826] Petraroli, R.; Silvestrini, M. C.; Brunori, M.: A new point mutation

[68827] of the prion protein gene in Creutzfeldt-Jakob disease. Ann. Neurol. 34:

[68828] 802-807, 1993.

[68829]

[68830] 131. Poulter, M.; Baker, H. F.; Frith, C. D.; Leach, M.; Lofthouse,

[68831] R.; Ridley, R. M.; Shah, T.; Owen, F.; Collinge, J.; Brown, J.; Hardy,

[68832] J.; Mullan, M. J.; Harding, A. E.; Bennett, C.; Doshi, R.; Crow, T.

[68833] J.: Inherited prion disease with 144 base pair gene insertion. 1.

[68834] Genealogical and molecular studies. Brain 115: 675-685, 1992.

[68835]

[68836] 132. Prusiner, S. B.: Molecular biology and genetics of prion diseases. Cold

[68837] Spring Harbor Symp. Quant. Biol. 61: 473-493, 1996.

[68838]

[68839] 133. Prusiner, S. B.: Prions causing degenerative neurological diseases. Annu.
[68840] Rev. Med. 38: 381-398, 1987.

[68841]

[68842] 134. Prusiner, S. B.: Molecular biology of prion diseases. Science 252:
[68843] 1515-1522, 1991.

[68844]

[68845] 135. Prusiner, S. B.: Biology and genetics of prion diseases. Ann.
[68846] Rev. Microbiol. 48: 655-686, 1994.

[68847]

[68848] 136. Prusiner, S. B.: Novel proteinaceous infectious particles cause
[68849] scrapie. Science 216: 136-144, 1982.

[68850]

[68851] 137. Puckett, C.; Concannon, P.; Casey, C.; Hood, L.: Genomic structure
[68852] of the human prion protein gene. Am. J. Hum. Genet. 49: 320-329,

[68853] 1991.

[68854]

[68855] 138. Reder, A. T.; Mednick, A. S.; Brown, P.; Spire, J. P.; Cauter,

[68856] V.; Wollmann, R. L.; Cervenakova, L.; Goldfarb, L. G.; Garay, A.;

[68857] Ovsiew, F.; Gajdusek, D. C.; Roos, R. P.: Clinical and genetic studies

[68858] of fatal familial insomnia. Neurology 45: 1068-1075, 1995.

[68859]

[68860] 139. Riek, R.; Wider, G.; Billeter, M.; Hornemann, S.; Glockshuber,

[68861] R.; Wuthrich, K.: Prion protein NMR structure and familial human

[68862] spongiform encephalopathies. Proc. Nat. Acad. Sci. 95: 11667-11672,

[68863] 1998.

[68864]

[68865] 140. Rivera, H.; Zuffardi, O.; Maraschio, P.; Caiulo, A.; Anichini,

[68866] C.; Scarinci, R.; Vivarelli, R.: Alternate centromere inactivation
[68867] in a pseudodicentric (15;20)(pter;pter) associated with a progressive
[68868] neurological disorder. J. Med. Genet. 26: 626-630, 1989.

[68869]

[68870] 141. Robakis, N. K.; Devine-Gage, E. A.; Jenkins, E. C.; Kascsak,
[68871] R. J.; Brown, W. T.; Krawczun, M. S.; Silverman, W. P.: Localization
[68872] of a human gene homologous to the PrP gene on the p arm of chromosome
[68873] 20 and detection of PrP-related antigens in normal human brain. Biochem.
[68874] Biophys. Res. Commun. 140: 758-765, 1986.

[68875]

[68876] 142. Sailer, A.; Bueler, H.; Fischer, M.; Aguzzi, A.; Weissmann, C.
[68877] : No propagation of prions in mice devoid of PrP. Cell 77: 967-968,
[68878] 1994.

[68879]

[68880] 143. Sakaguchi, S.; Katamine, S.; Nishida, N.; Moriuchi, R.; Shigamatsu,

[68881] K.; Sugimoto, T.; Nakatani, A.; Kataoka, Y.; Houtani, T.; Shirabe,

[68882] S.; Okada, H.; Hasegawa, S.; Miyamoto, T.; Noda, T.: Loss of cerebellar

[68883] Purkinje cells in aged mice homozygous for a disrupted PrP gene. Nature 380:

[68884] 528-531, 1996.

[68885]

[68886] 144. Samaia, H. B.; Mari, J. J.; Vallada, H. P.; Moura, R. P.; Simpson,

[68887] A. J. G.; Brentani, R. R.: A prion-linked psychiatric disorder. Nature 390:

[68888] 241 only, 1997.

[68889]

[68890] 145. Schellenberg, G. D.; Anderson, L.; O'dahl, S.; Wisjman, E. M.;

[68891] Sadovnick, A. D.; Ball, M. J.; Larson, E. B.; Kukull, W. A.; Martin,

[68892] G. M.; Roses, A. D.; Bird, T. D.: APP-717, APP-693, and PRIP gene
[68893] mutations are rare in Alzheimer disease. Am. J. Hum. Genet. 49:
[68894] 511-517, 1991.

[68895]

[68896] 146. Schnittger, S.; Gopal Rao, V. V. N.; Deutsch, U.; Gruss, P.;
[68897] Balling, R.; Hansmann, I.: PAX1, a member of the paired box-containing
[68898] class of developmental control genes, is mapped to human chromosome
[68899] 20p11.2 by in situ hybridization (ISH and FISH). Genomics 14: 740-744,
[68900] 1992.

[68901]

[68902] 147. Scott, M.; Foster, D.; Mirenda, C.; Serban, D.; Coufal, F.; Walchli,
[68903] M.; Torchia, M.; Groth, D.; Carlson, G.; DeArmond, S. J.; Westaway,
[68904] D.; Prusiner, S. B.: Transgenic mice expressing hamster prion protein

[68905] produce species-specific scrapie infectivity and amyloid plaques. Cell 59:

[68906] 847-857, 1989.

[68907]

[68908] 148. Shibuya, S.; Higuchi, J.; Shin, R.-W.; Tateishi, J.; Kita-moto,

[68909] T.: Protective prion protein polymorphisms against spo-radic Creutzfeldt-Jakob

[68910] disease. (Letter) Lancet 351: 419 only, 1998.

[68911]

[68912] 149. Shmerling, D.; Hegyi, I.; Fischer, M.; Blattler, T.; Brandner,

[68913] S.; Gotz, J.; Rulicke, T.; Flechsig, E.; Cozzio, A.; von Mer-ing, C.;

[68914] Hangartner, C.; Aguzzi, A.; Weissmann, C.: Expression of amino-terminally

[68915] truncated PrP in the mouse leading to ataxia and specific cerebellar

[68916] lesions. Cell 93: 203-214, 1998.

[68917]

[68918] 150. Simon, E. S.; Kahana, E.; Chapman, J.; Treves, T. A.;

Gabizon,
[68919] R.; Rosenmann, H.; Zilber, N.; Korczyn, A. D.: Creutzfeldt-Jakob
[68920] disease profile in patients homozygous for the PRNP E200K mutation. Ann.
[68921] Neurol. 47: 257-260, 2000.

[68922]

[68923] 151. Sparkes, R. S.; Simon, M.; Cohn, V. H.; Fournier, R. E. K.; Lem,
[68924] J.; Klisak, I.; Heinzmann, C.; Blatt, C.; Lucero, M.; Mohan-das, T.;
[68925] DeArmond, S. J.; Westaway, D.; Prusiner, S. B.; Weiner, L. P.: Assignment
[68926] of the human and mouse prion protein genes to homolo-gous chromosomes. Proc.
[68927] Nat. Acad. Sci. 83: 7358-7362, 1986.

[68928]

[68929] 152. Speer, M. C.; Goldgaber, D.; Goldfarb, L. G.; Roses, A. D.; Pericak-Vance,
[68930] M. A.: Support of linkage of Gerstmann-Straus-sler-Scheinker syndrome
[68931] to the prion protein gene on chromosome 20p12-pter.

Genomics 9:
[68932] 366-368, 1991.

[68933]

[68934] 153. Supattapone, S.; Bosque, P.; Muramoto, T.; Wille, H.; Aagaard,
[68935] C.; Peretz, D.; Nguyen, H.-O. B.; Heinrich, C.; Torchia, M.; Safar,
[68936] J.; Cohen, F. E.; DeArmond, S. J.; Prusiner, S. B.; Scott, M.: Prion
[68937] protein of 106 residues creates an artificial transmission barrier
[68938] for prion replication in transgenic mice. Cell 96: 869-878, 1999.

[68939]

[68940] 154. Supattapone, S.; Bouzamondo, E.; Ball, H. L.; Wille, H.; Nguyen,
[68941] H.-O. B.; Cohen, F. E.; DeArmond, S. J.; Prusiner, S. B.; Scott, M.
[68942] : A protease-resistant 61-residue prion peptide causes neurodegeneration
[68943] in transgenic mice. Molec. Cell. Biol. 21: 2608-2616, 2001.

[68944]

[68945] 155. Tagliavini, F.; Lievens, P. M.-J.; Tranchant, C.; Warter, J.-M.;

[68946] Mohr, M.; Giaccone, G.; Perini, F.; Rossi, G.; Salmona, M.; Piccardo,

[68947] P.; Ghetti, B.; Beavis, R. C.; Bugiani, O.; Frangione, B.; Prelli,

[68948] F.: A 7-kDa prion protein (PrP) fragment, an integral com-ponent of

[68949] the PrP region required for infectivity, is the major amy-loid protein

[68950] in Gerstmann-Straussler-Scheinker disease A117V. J. Biol. Chem. 276:

[68951] 6009-6015, 2001.

[68952]

[68953] 156. Tagliavini, F.; Prelli, F.; Ghiso, J.; Bugiani, O.; Serban, D.;

[68954] Prusiner, S. B.; Farlow, M. R.; Ghetti, B.; Frangione, B.: Amyloid

[68955] protein of Gerstmann-Straussler-Scheinker disease (Indiana kindred)

[68956] is an 11 kd fragment of prion protein with an N-terminal

glycine at
[68957] codon 58. EMBO J. 10: 513-519, 1991.

[68958]

[68959] 157. Tagliavini, F.; Prelli, F.; Porro, M.; Rossi, G.; Giaccone, G.;
[68960] Farlow, M. R.; Dlouhy, S. R.; Ghetti, B.; Bugiani, O.; Fran-
gione,
[68961] B.: Amyloid fibrils in Gerstmann-Straussler-Scheinker dis-
ease (Indiana
[68962] and Swedish kindreds) express only PrP peptides encoded
by the mutant
[68963] allele. Cell 79: 695-703, 1994.

[68964]

[68965] 158. Tateishi, J.; Brown, P.; Kitamoto, T.; Hoque, Z. M.;
Roos, R.;
[68966] Wollman, R.; Cervenakova, L.; Gajdusek, D. C.: First ex-
perimental
[68967] transmission of fatal familial insomnia. Nature 376:
434-435, 1995.

[68968]

[68969] 159. Telling, G. C.; Parchi, P.; DeArmond, S. J.; Cortelli, P.;

Montagna,
P.; Gabizon, R.; Mastrianni, J.; Lugaresi, E.; Gambetti, P.; Prusiner,
S. B.: Evidence for the conformation of the pathologic isoform of
the prion protein enciphering and propagating prion diversity. Science 274:
2079-2082, 1996.

160. Telling, G. C.; Scott, M.; Mastrianni, J.; Gabizon, R.; Torchia,
M.; Cohen, F. E.; DeArmond, S. J.; Prusiner, S. B.: Prion propagation
in mice expressing human and chimeric PrP transgenes implicates the
interaction of cellular PrP with another protein. Cell 83:
79-90,
1995.

161. Ter-Avanesyan, M. D.; Dagkesamanskaya, A. R.; Kushnirov, V. V.;
Smirnov, V. N.: The SUP35 omnipotent suppressor gene is

involved
[68983] in the maintenance of the non-mendelian determinant [psi+] in the
[68984] yeast Saccharomyces cerevisiae. Genetics 137: 671-676, 1994.
[68985]
[68986] 162. Tobler, I.; Gaus, S. E.; Deboer, T.; Ackermann, P.; Fis-cher,
[68987] M.; Rullcke, T.; Moser, M.; Oesch, B.; McBride, P. A.; Man-son, J.
[68988] C.: Altered circadian activity rhythms and sleep in mice devoid of
[68989] prion protein. Nature 380: 639-642, 1996.
[68990]
[68991] 163. Westaway, D.; DeArmond, S. J.; Cayetano-Canlas, J.; Groth, D.;
[68992] Foster, D.; Yang, S.-L.; Torchia, M.; Carlson, G. A.; Prusiner, S.
[68993] B.: Degeneration of skeletal muscle, peripheral nerves, and the central
[68994] nervous system in transgenic mice overexpressing wild-type prion proteins. Cell 76:

[68995] 117-129, 1994.

[68996]

[68997] 164. Whittington, M. A.; Sidle, K. C. L.; Gowland, I.; Meads, J.;

[68998] Hill, A. F.; Palmer, M. S.; Jefferys, J. G. R.; Collinge, J.: Rescue

[68999] of neurophysiological phenotype seen in PrP null mice by transgene

[69000] encoding human prion protein. Nature Genet. 9: 197-201, 1995.

[69001]

[69002] 165. Wickner, R. B.: [URE3] as an altered URE2 protein: evidence

[69003] for a prior analog in Saccharomyces cerevisiae. Science 264: 566-569,

[69004] 1994.

[69005]

[69006] 166. Windl, O.; Giese, A.; Schulz-Schaeffer, W.; Zerr, I.; Skworc,

[69007] K.; Arendt, S.; Oberdieck, C.; Bodemer, M.; Poser, S.; Kretzschmar,

[69008] H. A.: Molecular genetics of human prion diseases in Germany. Hum.

[69009] Genet. 105: 244-252, 1999.

[69010]

[69011] 167. Yamada, M.; Itoh, Y.; Fujigasaki, H.; Naruse, S.; Kaneko, K.;

[69012] Kitamoto, T.; Tateishi, J.; Otomo, E.; Hayakawa, M.; Tanaka, J.; Matsushita,

[69013] M.; Miyatake, T.: A missense mutation at codon 105 with codon 129

[69014] polymorphism of the prion protein gene in a new variant of Gerstmann-Straussler-Scheinker

[69015] disease. Neurology 43: 2723-2724, 1993.

[69016] 11611.

[69017] Kreczel, W.; Ghyselinck, N.; Samad, T. A.; Dupe, V.; Kastner, P.;

[69018] Borrelli, E.; Chambon, P.: Impaired locomotion and dopamine signaling

[69019] in retinoid receptor mutant mice. Science 279: 863-867, 1998.

[69020] 11612.

[69021] Lotan, R.; Xu, X.-C.; Lippman, S. M.; Ro, J. Y.; Lee, J. S.; Lee,
[69022] J. J.; Hong, W. K.: Suppression of retinoic acid receptor-beta in
[69023] premalignant oral lesions and its up-regulation by isotretinoin. New
[69024] Eng. J. Med. 332: 1405-1410, 1995.

[69025] 11613.

[69026] Mattei, M.-G.; de The, H.; Mattei, J.-F.; Marchio, A.; Tiol-lais,
[69027] P.; Dejean, A.: Assignment of the human hap retinoic acid receptor
[69028] RAR-beta gene to the p24 band of chromosome 3. Hum. Genet. 80: 189-190,
[69029] 1988.

[69030] 11614.

[69031] Nadeau, J. H.; Compton, J. G.; Giguere, V.; Rossant, J.; Varmuza,
[69032] S.: Close linkage of retinoic acid receptor genes with homeobox-
[69033] and keratin-encoding genes on paralogous segments of mouse chromosomes

[69034] 11 and 15. Mammalian Genome 3: 202-208, 1992.

[69035] 11615.

[69036] Samad, A.; Kreczel, W.; Chambon, P.; Borrelli, E.: Regulation

[69037] of dopaminergic pathways by retinoids: activation of the D2 receptor

[69038] promoter by members of the retinoic acid receptor-retinoid X receptor

[69039] family. Proc. Nat. Acad. Sci. 94: 14349-14354, 1997.

[69040] 11616.

[69041] Flagiello, D.; Apiou, F.; Gibaud, A.; Poupon, M. F.; Dutirlaux,

[69042] B.; Malfoy, B.: Assignment of the genes for cellular retinoic acid

[69043] binding protein 1 (CRABP1) and 2 (CRABP2) to human chromosome band

[69044] 15q24 and 1q21.3, respectively, by in situ hybridization. Cytogenet.

[69045] Cell Genet. 76: 17-18, 1997.

[69046] 11617.

[69047] Geurts van Kessel, A.; de Leeuw, H.; Dekker, E. J.; Rijks, L.;

[69048] Spurr, N.; Ledbetter, D.; Kootwijk, E.; Vaessen, M. J.: Localization

[69049] of the cellular retinoic acid binding protein (CRABP) gene relative

[69050] to the acute promyelocytic leukemia-associated breakpoint on human

[69051] chromosome 15. Hum. Genet. 87: 201-204, 1991.

[69052] 11618.

[69053] MacGregor, T. M.; Copeland, N. G.; Jenkins, N. A.; Giguere, V.

[69054] : The murine gene for cellular retinoic acid-binding protein type

[69055] II: genomic organization, chromosomal localization, and post-transcriptional

[69056] regulation by retinoic acid. J. Biol. Chem. 267: 7777-7783, 1992.

[69057] 11619.

[69058] Ong, D. E.: Cellular retinoid-binding proteins. Arch. Derm. 123:

[69059] 1693-1695, 1987.

[69060] 11620.

[69061] Vaessen, M. J.; Rijks, L.; Dekker, E. J.; Kootwijk, E.; Bootsma,
[69062] D.; Westerveld, A.; Geurts van Kessel, A. H. M.: Localization of
[69063] the human cellular retinoic acid-binding protein (CRABP) on chromosome
[69064] 15. (Abstract) Cytogenet. Cell Genet. 51: 1094 only, 1989.

[69065] 11621.

[69066] Alcalay, M.; Zangrilli, D.; Pandolfi, P. P.; Longo, L.; Mencarelli,
[69067] A.; Giacomucci, A.; Rocchi, M.; Biondi, A.; Rambaldi, A.; Lo Coco,
[69068] F.; Diverio, D.; Donti, E.; Grignani, F.; Pelicci, P. G.: Translocation
[69069] breakpoint of acute promyelocytic leukemia lies within the retinoic
[69070] acid receptor alpha locus. Proc. Nat. Acad. Sci. 88: 1977-1981,
[69071] 1991.

[69072] 11622.

[69073] Arveiler, B.; Petkovich, M.; Mandel, J. L.; Chambon, P.: A PstI

[69074] RFLP for the human retinoic acid receptor in 17q21. Nucleic Acids

[69075] Res. 16: 6252 only, 1988.

[69076] 11623.

[69077] Bale, A. E.; Weinberger, C.; McBride, O. W.: A retinoic acid receptor

[69078] cDNA probe (RAR2) identifies a moderately frequent RFLP on chromosome

[69079] 17. Nucleic Acids Res. 16: 7755 only, 1988.

[69080] 11624.

[69081] Borrow, J.; Goddard, A. D.; Sheer, D.; Solomon, E.: Molecular

[69082] analysis of acute promyelocytic leukemia breakpoint cluster region

[69083] on chromosome 17. Science 249: 1577-1580, 1990.

[69084] 11625.

[69085] Brand, N.; Petkovich, M.; Krust, A.; Chambon, P.; de The, H.; Marchio,

[69086] A.; Tiollais, P.; Dejean, A.: Identification of a second human retinoic

[69087] acid receptor.(Letter) Nature 332: 850-853, 1988.

[69088] 11626.

[69089] Jeremiah, S. J.; West, L. F.; Abbott, C. M.; Murad, Z.; Povey,

[69090] S.; Thomas, H. J.; Solomon, E.; Di Scipio, R.; Fey, G. H.: Three

[69091] genes coding for late acting components of complement assigned to

[69092] chromosome 5. (Abstract) Cytogenet. Cell Genet. 51: 1019 only, 1989.

[69093] 11627.

[69094] Dodd, J.; Morton, S. B.; Karagogeos, D.; Yamamoto, M.; Jessell,

[69095] T. M.: Spatial regulation of axonal glycoprotein expression on subsets

[69096] of embryonic spinal neurons. Neuron 1: 105-116, 1988.

[69097] 11628.

[69098] Freigang, J.; Proba, K.; Leder, L.; Diederichs, K.; Sondereg-

[69099] ger, P.; Welte, W.: The crystal structure of the ligand binding module

[69100] of axonin-1/TAG-1 suggests a zipper mechanism for neural cell adhesion. Cell 101:

[69101] 425-433, 2000.

[69102] 11629.

[69103] Kenwrick, S.; Leversha, M.; Rooke, L.; Hasler, T.; Son-
deregger,
[69104] P.: Localization of the human TAX-1 gene to 1q32.1: a re-
gion implicated
[69105] in microcephaly and Van der Woude syndrome. Hum.
Molec. Genet. 2:
[69106] 1461-1462, 1993.

[69107] 11630.

[69108] Kozlov, S. V.; Giger, R. J.; Hasler, T.; Korvatska, E.;
Schorderet,
[69109] D. F.; Sonderegger, P.: The human TAX1 gene encoding
the axon-associated
[69110] cell adhesion molecule TAG-1/axonin-1: genomic struc-
ture and basic
[69111] promoter. Genomics 30: 141-148, 1995.

[69112] 11631.

[69113] Rickman, D. S.; Tyagi, R.; Zhu, X.-X.; Bobek, M. P.; Song,
S.;
[69114] Blaivas, M.; Misek, D. E.; Israel, M. A.; Kurnit, D. M.; Ross,

D.

[69115] A.; Kish, P. E.; Hanash, S. M.: The gene for the axonal cell adhesion

[69116] molecule TAX-1 is amplified and aberrantly expressed in malignant

[69117] gliomas. Cancer Res. 61: 2162-2168, 2001.

[69118] 11632.

[69119] Tsiotra, P. C.; Karagogeos, D.; Theodorakis, K.; Michaelis, T.

[69120] M.; Modi, W. S.; Furley, A. J.; Jessell, T. M.; Papamatheakis, J.

[69121] : Isolation of the cDNA and chromosomal localization of the gene (TAX1)

[69122] encoding the human axonal glycoprotein TAG-1. Genomics 18: 562-567,

[69123] 1993.

[69124] 11633.

[69125] Artavanis-Tsakonas, S.; Matsuono, K.; Fortini, M.: Notch signaling. Science 268:

[69126] 225-232, 1995.

[69127] 11634.

[69128] Bruckner, K.; Perez, L.; Clausen, H.; Cohen, S.: Glycosyl-transferase

[69129] activity of Fringe modulates Notch-Delta interactions. Nature 406:

[69130] 411-415, 2000.

[69131] 11635.

[69132] Ackerman, M. J.; Clapham, D. E.: Ion channels--basic science and

[69133] clinical disease. New Eng. J. Med. 336: 1575-1586, 1997.

[69134] 11636.

[69135] Ackerman, M. J.; Schroeder, J. J.; Berry, R.; Schaid, D. J.; Porter,

[69136] C.-B. J.; Michels, V. V.; Thibodeau, S. N.: A novel mutation in KVLQT1

[69137] is the molecular basis of inherited long QT syndrome in a near-drowning

[69138] patient's family. Pediat. Res. 44: 148-153, 1998.

[69139] 11637.

[69140] Ackerman, M. J.; Tester, D. J.; Porter, C. J.; Edwards, W. D.:

[69141] Molecular diagnosis of the inherited long-QT syndrome in a woman who

[69142] died after near-drowning. New Eng. J. Med. 341: 1121-1125, 1999.

[69143] 11638.

[69144] Barlow, D. P.: Box: KVLQT1 complexities in Beckwith-Wiedeman (sic)

[69145] syndrome. Nature Genet. 15: 114 only, 1997.

[69146] 11639.

[69147] Barlow, J. B.; Bosman, C. K.; Cochrane, J. W. C.: Congenital cardiac

[69148] arrhythmia. Lancet II: 531 only, 1964.

[69149] 11640.

[69150] Baudouy, P.; Andreassian, B.; Attuel, P.; Greze, M.; Soulie, J.;

[69151] Fruchaud, J.: Syndrome de Romano-Ward et stellectomie gauche: revue

[69152] generale a propos d'un nouveau cas. Arch. Mal. Coeur 70: 645-652,

[69153] 1977.

[69154] 11641.

[69155] Benhorin, J.; Kalman, Y. M.; Medina, A.; Towbin, J.; Rave-Harel,

[69156] N.; Dyer, T. D.; Blangero, J.; MacCluer, J. W.; Kerem, B.: Evidence

[69157] of genetic heterogeneity in the long QT syndrome. Science 260: 1960-1962,

[69158] 1993.

[69159] 11642.

[69160] Bhandari, A. K.; Scheinman, M.: The long QT syndrome. Mod. Concepts

[69161] Cardiovasc. Dis. 54: 45-50, 1985.

[69162] 11643.

[69163] Bonduelle, M.: Personal Communication. Brussels, Belgium 5/30/1993.

[69164] 11644.

[69165] Casimiro, M. C.; Knollmann, B. C.; Ebert, S. N.; Vary, J. C.,

[69166] Jr.; Greene, A. E.; Franz, M. R.; Grinberg, A.; Huang, S. P.; Pfeifer,

[69167] K.: Targeted disruption of the Kcnq1 gene produces a mouse model

[69168] of Jervell and Lange-Nielsen syndrome. Proc. Nat. Acad. Sci. 98:

[69169] 2526-2531, 2001.

[69170] 11645.

[69171] Chen, Q.; Zhang, D.; Gingell, R. L.; Moss, A. J.; Napolitano,

[69172] C.; Priori, S. G.; Schwartz, P. J.; Kehoe, E.; Robinson, J. L.;
Schulze-Bahr,

[69173] E.; Wang, Q.; Towbin, J. A.: Homozygous deletion in
KVLQT1 associated

[69174] with Jervell and Lange-Nielsen syndrome. Circulation 99:
1344-1347,

[69175] 1999.

[69176] 11646.

[69177] Cleary, M. A.; van Raamsdonk, C. D.; Levorse, J.; Zheng, B.;
Bradley,

[69178] A.; Tilghman, S. M.: Disruption of an imprinted gene clus-
ter by a

[69179] targeted chromosomal translocation in mice. Nature
Genet. 29: 78-82,

[69180] 2001.

[69181] 11647.

[69182] Curran, M.; Atkinson, D.; Timothy, K.; Vincent, G. M.;
Moss, A.

[69183] J.; Leppert, M.; Keating, M.: Locus heterogeneity of auto-

somal dominant

[69184] long QT syndrome. J. Clin. Invest. 92: 799-803, 1993.

[69185] 11648.

[69186] DeSilvey, D. L.; Moss, A. J.: Primidone in the treatment of the

[69187] long QT syndrome: QT shortening and ventricular ar-

rhythmia suppression. Ann.

[69188] Intern. Med. 93: 53-54, 1980.

[69189] 11649.

[69190] DiSegni, E.; David, D.; Katzenstein, M.; Klein, H. O.; Kaplinsky,

[69191] E.; Levy, M. J.: Permanent overdrive pacing for the sup-

pression of

[69192] recurrent ventricular tachycardia in a newborn with long QT syndrome. J.

[69193] Electrocardiol. 13: 189-192, 1980.

[69194] 11650.

[69195] Donger, C.; Denjoy, I.; Berthet, M.; Neyroud, N.; Cruaud, C.;

[69196] Bennaceur, M.; Chivoret, G.; Schwartz, K.; Coumel, P.; Guicheney,

[69197] P.: KVLQT1 C-terminal missense mutation causes a forme fruste long-QT
[69198] syndrome. Circulation 96: 2778-2781, 1997.

[69199] 11651.

[69200] Engel, J. R.; et al.; et al.: Epigenotype-phenotype correlations
[69201] in Beckwith-Wiedemann syndrome. J. Med. Genet. 37: 921-926, 2000.
[69202] 11652.

[69203] Fitzpatrick, G. V.; Soloway, P. D.; Higgins, M. J.: Regional
[69204] loss of imprinting and growth deficiency in mice with a targeted deletion
[69205] of KvDMR1. Nature Genet. 9Sept, 2002. Note: Advance Electronic Publication.
[69206] 11653.

[69207] Flugelman, M. Y.; Pollack, S.; Hammerman, H.; Riss, E.; Barzilai,
[69208] D.: Congenital prolongation of Q-T interval: a family study of three
[69209] generations. Cardiology 69: 170-174, 1982.

[69210] 11654.

[69211] Furgerg, C.; Hornell, H.: Familial QT prolongation and risk of

[69212] sudden death. Acta Paediat. Scand. 64: 777-782, 1975.

[69213] 11655.

[69214] Gale, G. E.; Bosman, C. K.; Tucker, R. B. K.; Barlow, J. B.:

[69215] Hereditary prolongation of Q-T interval: study of two families. Brit.

[69216] Heart J. 32: 505-509, 1970.

[69217] 11656.

[69218] Gamstorp, I.; Nilsen, R.; Westling, H.: Congenital cardiac arrhythmia.

[69219] (Letter) Lancet II: 965 only, 1964.

[69220] 11657.

[69221] Garza, L. A.; Vick, R. L.; Nora, J. J.; McNamara, D. G.: Heritable

[69222] Q-T prolongation without deafness. Circulation 41: 39-48, 1970.

[69223] 11658.

[69224] Giuffre, R. M.; Hejtmancik, J. F.; McCabe, E. R. B.; Towbin, J.

[69225] A.: Long QT (Romano-Ward) syndrome: molecular genetic evidence against
[69226] tight HLA linkage. (Abstract) Am. J. Hum. Genet. 47 (suppl.): A180
[69227] only, 1990.

[69228] 11659.

[69229] Gohl, K.; Feistel, H.; Weikl, A.; Bachmann, K.; Wolf, F.: Congenital
[69230] myocardial sympathetic dysinnervation (CMSD)--a structural defect
[69231] of idiopathic long QT syndrome. PACE 14: 1544-1553, 1991.
[69232] 11660.

[69233] Hashiba, K.: Hereditary QT prolongation syndrome in Japan: genetic
[69234] analysis and pathological findings of the conduction system. Jpn.
[69235] Circ. J. 42: 1133-1150, 1978.

[69236] 11661.

[69237] Horn, C. A.; Beekman, R. H.; Dick, M., II; Lacina, S. J.: The

[69238] congenital long QT syndrome: an unusual cause of child-

hood seizures. Am.
[69239] J. Dis. Child. 140: 659-661, 1986.

[69240] 11662.

[69241] Itoh, S.; Munemura, S.; Satoh, H.: A study of the inheri-
tance
[69242] pattern of Romano-Ward syndrome: prolonged Q-T inter-
val, syncope,
[69243] and sudden death. Clin. Pediat. 21: 20-24, 1982.

[69244] 11663.

[69245] Bulle, F.; Mattei, M. G.; Siegrist, S.; Pawlak, A.; Passage, E.;

[69246] Chobert, M. N.; Laperche, Y.; Guellaen, G.: Assignment of
the human
[69247] gamma-glutamyl transferase gene to the long arm of
chromosome 22. Hum.
[69248] Genet. 76: 283-286, 1987.

[69249] 11664.

[69250] Tsuchida, Y.; Yokomori, K.; Choi, S. H.: Hereditary renal
tumors:
[69251] Wilms' tumor--congenital anomalies' syndrome. Nippon
Rinsho 53:
[69252] 2742-2748, 1995.

[69253] 11665.

[69254] Turleau, C.; de Grouchy, J.; Dufier, J. L.; Phuc, L. H.; Schmelck,

[69255] P. H.; Rappaport, R.; Nihoul-Fekete, C.; Diebold, N.: Aniridia, male

[69256] pseudohermaphroditism, gonadoblastoma, mental retardation, and del

[69257] 11p13. Hum. Genet. 57: 300-306, 1981.

[69258] 11666.

[69259] Turleau, C.; de Grouchy, J.; Tournade, M.-F.; Gagnadoux, M.-F.;

[69260] Junien, C.: Del 11p/aniridia complex: report of three patients and

[69261] review of 37 observations from the literature. Clin. Genet. 26:

[69262] 356-362, 1984.

[69263] 11667.

[69264] van Heyningen, V.; Boyd, P. A.; Seawright, A.; Fletcher, J. M.;

[69265] Fantes, J. A.; Buckton, K. E.; Spowart, G.; Porteous, D. J.; Hill,

[69266] R. E.; Newton, M. S.; Hastie, N. D.: Molecular analysis of chromosome

[69267] 11 deletions in aniridia-Wilms tumor syndrome. Proc. Nat. Acad. Sci. 82:

[69268] 8592-8596, 1985.

[69269] 11668.

[69270] Varanasi, R.; Bardeesy, N.; Ghahremani, M.; Petruzzi, M.-J.; Nowak,

[69271] N.; Adam, M. A.; Grundy, P.; Shows, T. B.; Pelletier, J.: Fine structure

[69272] analysis of the WT1 gene in sporadic Wilms tumor. Proc. Nat. Acad.

[69273] Sci. 91: 3554-3558, 1994.

[69274] 11669.

[69275] Weissman, B. E.; Saxon, P. J.; Pasquale, S. R.; Jones, G. R.;

[69276] Geiser, A. G.; Stanbridge, E. J.: Introduction of normal human chromosome

[69277] into a Wilms' tumor cell line controls its tumorigenic expression. Science 236:

[69278] 175-180, 1987.

[69279] 11670.

[69280] White, G. R. M.; Kelsey, A. M.; Varley, J. M.; Birch, J. M.:

[69281] Somatic glypican 3 (GPC3) mutations in Wilms' tumour.
Brit. J. Cancer 86:

[69282] 1920-1922, 2002.

[69283] 11671.

[69284] Yunis, J. J.; Ramsay, K. C.: Familial occurrence of the
aniridia-Wilms

[69285] tumor syndrome with deletion 11p13-14.1. J. Pediat. 96:
1027-1030,

[69286] 1980.

[69287] 11672.

[69288] Brown, J. R.; Daar, I. O.; Krug, J. R.; Maquat, L. E.: Charac-
terization

[69289] of the functional gene and several processed pseudogenes
in the human

[69290] triosephosphate isomerase gene family. Molec. Cell. Biol.
5: 1694-1706,

[69291] 1985.

[69292] 11673.

[69293] Van Goethem, G.; Dermaut, B.; Lofgren, A.; Martin, J.-J.;
Van Broeckhoven,

[69294] C.: Mutation of POLG is associated with progressive external ophthalmoplegia
[69295] characterized by mtDNA deletions. Nature Genet. 28: 211-212, 2001.
[69296] 11674.

[69297] Uchida, S.; Kwon, H. M.; Yamauchi, A.; Preston, A. S.; Marumo,
[69298] F.; Handler, J. S.: Molecular cloning of the cDNA for an MDCK cell
[69299] Na(+)- and Cl(-)-dependent taurine transporter that is regulated by
[69300] hypertonicity. Proc. Nat. Acad. Sci. 89: 8230-8234, 1992.

[69301] 11675.

[69302] Neyroud, N.; Tesson, F.; Denjoy, I.; Leibovici, M.; Donger, C.;
[69303] Barhanin, J.; Faure, S.; Gary, F.; Coumel, P.; Petit, C.; Schwartz,
[69304] K.; Guicheney, P.: A novel mutation in the potassium channel gene
[69305] KVLQT1 causes the Jervell and Lange-Nielsen cardioauditory syndrome. Nature
[69306] Genet. 15: 186-189, 1997.

[69307] 11676.

[69308] Pervaiz, S.; Brew, K.: Homology of beta-lactoglobulin, serum retinol-binding
[69309] protein, and protein HC. Science 228: 335-337, 1985.

[69310] 11677.

[69311] Barber, G. N.; Edelhoff, S.; Katze, M. G.; Disteche, C. M.: Chromosomal
[69312] assignment of the interferon-inducible double-stranded RNA-dependent
[69313] protein kinase (PRKR) to human chromosome 2p21-p22 and mouse chromosome
[69314] 17 E2. Genomics 16: 765-767, 1993.

[69315] 11678.

[69316] Kuhen, K. L.; Shen, X.; Carlisle, E. R.; Richardson, A.L.; Weier,
[69317] H.-U. G.; Tanaka, H.; Samuel, C. E.: Structural organization of the
[69318] human gene (PKR) encoding an interferon-inducible RNA-dependent protein
[69319] kinase (PKR) and differences from its mouse homolog. Genomics 36:
[69320] 197-201, 1996.

[69321] 11679.

[69322] Kuhen, K. L.; Shen, X.; Samuel, C. E.: Mechanism of interferon

[69323] action: sequence of the human interferon-inducible RNA-dependent protein

[69324] kinase (PKR) deduced from genomic clones. Gene 178: 191-193, 1996.

[69325] 11680.

[69326] Hummler, E.; Barker, P.; Gatzy, J.; Beermann, F.; Verdumo, C.;

[69327] Schmidt, A.; Boucher, R.; Rossier, B. C.: Early death due to defective

[69328] neonatal lung liquid clearance in alpha-ENaC-deficient mice. Nature

[69329] Genet. 12: 325-328, 1996.

[69330] 11681.

[69331] Rogers, M. A.; Winter, H.; Langbein, L.; Krieg, T.; Schweizer,

[69332] J.: Genomic characterization of the human type I cuticular hair keratin

[69333] hHa2 and identification of an adjacent novel type I hair keratin gene

[69334] hHa5. J. Invest. Derm. 107: 633-638, 1996.

[69335] 11682.

[69336] Hansen, R. S.; Wijmenga, C.; Luo, P.; Stanek, A. M.; Can-
field,
[69337] T. K.; Weemaes, C. M. R.; Gartler, S. M.: The DNMT3B DNA
methyltransferase
[69338] gene is mutated in the ICF immunodeficiency syndrome.
Proc. Nat.
[69339] Acad. Sci. 96: 14412-14417, 1999.

[69340] 11683.

[69341] Hassan, K. M. A.; Norwood, T.; Gimelli, G.; Gartler, S. M.;
Hansen,
[69342] R. S.: Satellite 2 methylation patterns in normal and ICF
syndrome
[69343] cells and association of hypomethylation with advanced
replication. Hum.
[69344] Genet. 109: 452-462, 2001.

[69345] 11684.

[69346] Shirohzu, H.; Kubota, T.; Kumazawa, A.; Sado, T.; Chijiwa,
T.;
[69347] Inagaki, K.; Suetake, I.; Tajima, S.; Wakui, K.; Miki, Y.;

Hayashi,
[69348] M.; Fukushima, Y.; Sasaki, H.: Three novel DNMT3B mutations in Japanese
[69349] patients with ICF syndrome. Am. J. Med. Genet. 112: 31-37, 2002.
[69350] 11685.

[69351] Imataka, H.; Sogawa, K.; Yasumoto, K.; Kikuchi, Y.; Sasano, K.;
[69352] Kobayashi, A.; Hayami, M.; Fujii-Kuriyama, Y.: Two regulatory proteins
[69353] that bind to the basic transcription element (BTE), a GC box sequence
[69354] in the promoter region of the rat P-4501A1 gene. EMBO J. 11: 3663-3671,
[69355] 1992.

[69356] 11686.

[69357] Ohe, N.; Yamasaki, Y.; Sogawa, K.; Inazawa, J.; Ariyama, T.; Oshimura,
[69358] M.; Fujii-Kuriyama, Y.: Chromosomal localization and cDNA sequence
[69359] of human BTEB, a GC box binding protein. Somat. Cell Molec. Genet. 19:

[69360] 499-503, 1993.

[69361] 11687.

[69362] Salinas, M.; Duprat, F.; Heurteaux, C.; Hugnot, J.-P.; Laz-
dunski,

[69363] M.: New modulatory alpha subunits for mammalian Shab
K(+) channels. J.

[69364] Biol. Chem. 272: 24371-24379, 1997.

[69365] 11688.

[69366] Cheng, J.; Scully, P.; Shew, J. Y.; Lee, W. H.; Vila, V.; Haas,

[69367] M.: Homozygous deletion of the retinoblastoma gene in
an acute lymphoblastic

[69368] leukemia (T) cell line. Blood 75: 730-735, 1990.

[69369] 11689.

[69370] Lee, E. Y.; To, H.; Shew, J. Y.; Bookstein, R.; Scully, P.; Lee,

[69371] W. H.: Inactivation of the retinoblastoma susceptibility
gene in

[69372] human breast cancers. Science 241: 218-221, 1988.

[69373] 11690.

[69374] Lambotte, S.; Veyhl, M.; Kohler, M.; Morrison-Shetlar, A. I.;

Kinne,
[69375] R. K. H.; Schmid, M.; Koepsell, H.: The human gene of a protein that
[69376] modifies Na+-D-glucose co-transport. DNA Cell Biol. 15: 769-777,
[69377] 1996.

[69378] 11691.

[69379] Kas, K.; Roijer, E.; Voz, M.; Meyen, E.; Stenman, G.; Van de Ven,
[69380] W. J. M.: A 2-Mb YAC contig and physical map covering the chromosome
[69381] 8q12 breakpoint cluster region in pleomorphic adenomas of the salivary
[69382] glands. Genomics 43: 349-358, 1997.

[69383] 11692.

[69384] Kas, K.; Voz, M. L.; Roijer, E.; Astrom, A.-K.; Meyen, E.; Stenman,
[69385] G.; Van de Ven, W. J. M.: Promoter swapping between the genes for
[69386] a novel zinc finger protein and beta-catenin in pleiomor-phic adenomas
[69387] with t(3;8)(p21;q12) translocations. Nature Genet. 15:

170-174,
[69388] 1997.

[69389] 11693.

[69390] Tillmann, H.; Eschrich, K.: Isolation and characterization of

[69391] an allelic cDNA for human muscle fructose-
1,6-biphosphatase. Gene 212:
[69392] 295-304, 1998.

[69393] 11694.

[69394] Aderem, A.; Ulevitch, R. J.: Toll-like receptors in the in-
duction
[69395] of the innate immune response. Nature 406: 780-785,
2000.
[69396] 11695.

[69397] Szabo, G.; Dallmann, G.; Muller, G.; Patthy, L.; Soller, M.;
Varga,
[69398] L.: A deletion in the myostatin gene causes the compact
(Cmpt) hypermuscular
[69399] mutation in mice. Mammalian Genome 9: 671-672, 1998.

[69400] 11696.

[69401] Zimmers, T. A.; Davies, M. V.; Koniaris, L. G.; Haynes, P.;

Esquela,

[69402] A. F.; Tomkinson, K. N.; McPherron, A. C.; Wolfman, N. M.; Lee, S.-J.

[69403] : Induction of cachexia in mice by systemically administered myostatin. Science 296:

[69404] 1486-1488, 2002.

[69405] 11697.

[69406] Cohen, P.; Rylatt, D. B.; Nimmo, G. A.: The hormonal control of

[69407] glycogen metabolism: the amino acid sequence at the phosphorylation

[69408] site of protein phosphatase inhibitor-1. FEBS Lett. 76: 182-186,

[69409] 1977.

[69410] 11698.

[69411] Helps, N. R.; Street, A. J.; Elledge, S. J.; Cohen, P. T.: Cloning

[69412] of the complete coding region for human protein phosphatase inhibitor

[69413] 2 using the two hybrid system and expression of inhibitor 2 in E.

[69414] coli. FEBS Lett. 340: 93-98, 1994.

[69415] 11699.

[69416] Huang, F. L.; Glinsmann, W. H.: Separation and characterization

[69417] of two phosphorylase phosphatase inhibitors from rabbit skeletal muscle. Europ.

[69418] J. Biochem. 70: 419-426, 1976.

[69419] 11700.

[69420] Majer, M.; Mott, D. M.; Mochizuki, H.; Rowles, J. C.; Pedersen,

[69421] O.; Knowler, W. C.; Bogardus, C.; Prochazka, M.: Association of the

[69422] glycogen synthase locus on 19q13 with NIDDM in Pima Indians. Diabetologia 39:

[69423] 314-321, 1996.

[69424] 11701.

[69425] Permana, P. A.; Mott, D. M.: Genetic analysis of human type 1

[69426] protein phosphatase inhibitor 2 in insulin-resistant Pima Indians. Genomics 41:

[69427] 110-114, 1997.

[69428] 11702.

[69429] Prochazka, M.; Mochizuki, H.; Baier, L. J.; Cohen, P. T. W.; Bogardus,
[69430] C.: Molecular and linkage analysis of type-1 protein phos-phatase
[69431] catalytic beta-subunit gene: lack of evidence for its major role in
[69432] insulin resistance in Pima Indians. Diabetologia 38: 461-466, 1995.
[69433] 11703.

[69434] Sakagami, H.; Kondo, H.: Molecular cloning of the cDNA for rat
[69435] phosphatase inhibitor-2 and its wide gene expression in the central
[69436] nervous system. J. Chem. Neuroanat. 8: 259-266, 1995.

[69437] 11704.

[69438] Sanseau, P.; Jackson, A.; Alderton, R. P.; Beck, S.; Senger, G.;
[69439] Sheer, D.; Kelly, A.; Trowsdale, J.: Cloning and characteri-zation
[69440] of human phosphatase inhibitor-2 (IPP-2) sequences. Mammalian Genome 5:
[69441] 490-496, 1994.

[69442] 11705.

[69443] Durand, B.; Sperisen, P.; Emery, P.; Barras, E.; Zufferey, M.;

[69444] Mach, B.; Reith, W.: RFXAP, a novel subunit of the RFX DNA binding

[69445] complex is mutated in MHC class II deficiency. EMBO J. 16: 1045-1055,

[69446] 1997.

[69447] 11706.

[69448] Mach, B.: Personal Communication. Geneva, Switzerland 10/28/1998.

[69449] 11707.

[69450] Nekrep, N.; Jabrane-Ferrat, N.; Peterlin, B. M.: Mutations in

[69451] the bare lymphocyte syndrome define critical steps in the assembly

[69452] of the regulatory factor X complex. Molec. Cell Biol. 20: 4455-4461,

[69453] 2000.

[69454] 11708.

[69455] Schwartz, R. S.: The case of the bare lymphocyte syndrome: tracking

[69456] down faulty transcription factors. (Editorial) New Eng. J. Med. 337:

[69457] 781-783, 1997.

[69458] 11709.

[69459] Steimle, V.; Durand, B.; Barras, E.; Zufferey, M.; Hadam, M. R.;

[69460] Mach, B.; Reith, W.: A novel DNA binding-regulatory factor is mutated

[69461] in primary MHC class II deficiency (bare lymphocyte syndrome). Genes

[69462] Dev. 9: 1021-1032, 1995.

[69463] 11710.

[69464] Touraine, J. L.; Betuel, H.: Immunodeficiency diseases and expression

[69465] of HLA antigens. Hum. Immun. 2: 147-153, 1981.

[69466] 11711.

[69467] Villard, J.; Lisowska-Grospierre, B.; van den Elsen, P.; Fischer,

[69468] A.; Reith, W.; Mach, B.: Mutation of RFXAP, a regulator of MHC class

[69469] II genes, in primary MHC class II deficiency. New Eng. J.

Med. 337:

[69470] 748-753, 1997.

[69471] 11712.

[69472] Peijnenburg, A.; Van Eggermond, M. C. J. A.; Van den Berg, R.;

[69473] Sanal, O.; Vossen, J. M. J. J.; Van den Elsen, P. J.: Molecular analysis

[69474] of an MHC class II deficiency patient reveals a novel mutation in

[69475] the RFX5 gene. Immunogenetics 49: 338-345, 1999.

[69476] 11713.

[69477] Kim, I.; Kim, J.-H.; Ryu, Y. S.; Jung, S. H.; Nah, J. J.; Koh,

[69478] G. Y.: Characterization and expression of a novel alternatively spliced

[69479] human angiopoietin-2. J. Biol. Chem. 275: 18550-18556, 2000.

[69480] 11714.

[69481] Maisonpierre, P. C.; Suri, C.; Jones, P. F.; Bartunkova, S.; Wiegand,

[69482] S. J.; Radziejewski, C.; Compton, D.; McClain, J.; Aldrich, T. H.;

[69483] Papadopoulos, N.; Daly, T. J.; Davis, S.; Sato, T. N.; Yan-
copoulos,
[69484] G. D.: Angiopoietin-2, a natural antagonist for Tie2 that
disrupts
[69485] in vivo angiogenesis. Science 277: 55-60, 1997.

[69486] 11715.

[69487] Tanaka, S.; Mori, M.; Sakamoto, Y.; Makuuchi, M.; Sugi-
machi, K.;
[69488] Wands, J. R.: Biologic significance of angiopoietin-2 ex-
pression
[69489] in human hepatocellular carcinoma. J. Clin. Invest. 103:
341-345,
[69490] 1999.

[69491] 11716.

[69492] Chow, V. T. K.; Quek, H. H.: HEP-COP, a novel human
gene whose
[69493] product is highly homologous to the alpha-subunit of the
yeast coatomer
[69494] protein complex. Gene 169: 223-227, 1996.

[69495] 11717.

[69496] Quek, H. H.; Chow, V. T. K.: Molecular and cellular studies

of

[69497] the human homolog of the 160-kD alpha-subunit of the coatomer protein

[69498] complex. DNA Cell Biol. 16: 275-280, 1997.

[69499] 11718.

[69500] Quek, H. H.; Chow, V. T. K.: Genomic organization and mapping

[69501] of the human HEP-COP gene (COPA) to 1q. Cytogenet. Cell Genet. 76:

[69502] 139-143, 1997.

[69503] 11719.

[69504] Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore, B.; Madsen,

[69505] P.; Vandekerckhove, J.; Celis, J. E.: Identification of two human

[69506] rho GDP dissociation inhibitor proteins whose overexpres-sion leads

[69507] to disruption of the actin cytoskeleton. Exp. Cell Res. 209: 165-174,

[69508] 1993.

[69509] 11720.

[69510] Wagner, T.; Tommerup, N.; Wirth, J.; Leffers, H.; Zimmer, J.; Back,
[69511] E.; Weissenbach, J.; Scherer, G.: A somatic cell hybrid panel for
[69512] distal 17q: GDIA1 maps to 17q25.3. Cytogenet. Cell Genet. 76: 172-175,
[69513] 1997.

[69514] 11721.

[69515] Collin, T.; Lory, P.; Taviaux, S.; Courtieu, C.; Guillbault, P.;
[69516] Berta, P.; Nargeot, J.: Cloning, chromosomal location and functional
[69517] expression of the human voltage-dependent calcium-channel beta 3 subunit. Europ.
[69518] J. Biochem. 220: 257-262, 1994.

[69519] 11722.

[69520] Yamada, Y.; Masuda, K.; Li, Q.; Ihara, Y.; Kubota, A.; Miura, T.;
[69521] Nakamura, K.; Fujii, Y. Seino, S.; Seino, Y.: The structures of the
[69522] human calcium channel alpha 1 subunit (CACNL1A2) and beta subunit

[69523] (CACNLB3) genes. Genomics 27: 312-319, 1995.

[69524] 11723.

[69525] Katsanis, N.; Yaspo, M.-L.; Fisher, E. M. C.: Identification and

[69526] mapping of a novel human gene, HRMT1L1, homologous to the rat protein

[69527] arginine N-methyltransferase 1 (PRMT1) gene. Mammalian Genome 8:

[69528] 526-529, 1997.

[69529] 11724.

[69530] Lin, W. J.; Gary, J. D.; Yang, M. C.; Clarke, S.; Herschman, H.

[69531] R.: The mammalian intermediate-early TIS21 protein and the leukemia-associated

[69532] BTG1 protein interact with a protein-arginine N-methyltransferase. J.

[69533] Biol. Chem. 271: 15034-15044, 1996.

[69534] 11725.

[69535] Scott, H. S.; Antonarakis, S. E.; Lalioti, M. D.; Rossier, C.;

[69536] Silver, P. A.; Henry, M. F.: Identification and characterization

[69537] of two putative human arginine methyltransferases (HRMT1L1 and HRMT1L2). Genomics 48:
[69538] 330-340, 1998.

[69539] 11726.

[69540] Herberg, J. A.; Sgouros, J.; Jones, T.; Copeman, J.; Humphray,
[69541] S. J.; Sheer, D.; Cresswell, P.; Beck, S.; Trowsdale, J.: Ge-nomic
[69542] analysis of the Tapasin gene, located close to the TAP loci in the
[69543] MHC. Europ. J. Immun. 28: 459-467, 1998.

[69544] 11727.

[69545] Mayer, W. E.; Klein, J.: Is tapasin a modified Mhc class I molecule? Immunogenetics 53:
[69546] 719-723, 2001.

[69547] 11728.

[69548] Michalova, V.; Murray, B. W.; Sultmann, H.; Klein, J.: A contig
[69549] map of the Mhc class I genomic region in the zebrafish reveals ancient
[69550] synteny. J. Immun. 164: 5296-5305, 2000.

[69551] 11729.

[69552] Ortmann, B.; Copeman, J.; Lehner, P. J.; Sadasivan, B.; Herberg,

[69553] J. A.; Grandea, A. G.; Riddell, S. R.; Tampe, R.; Spies, T.; Trowsdale,

[69554] J.; Cresswell, P.: A critical role for tapasin in the assembly and

[69555] function of multimeric MHC class I-TAP complexes. Science 277: 1306-1309,

[69556] 1997.

[69557] 11730.

[69558] Sadasivan, B.; Lehner, P. J.; Ortmann, B.; Spies, T.; Cresswell,

[69559] P.: Roles for calreticulin and a novel glycoprotein, tapasin, in

[69560] the interaction of MHC class I molecules with TAP. Immunity 5: 103-114,

[69561] 1996.

[69562] 11731.

[69563] Teng, M. S.; Stephens, R.; Du Pasquier, L.; Freeman, T.; Lindquist,

[69564] J. A.; Trowsdale, J.: A human TAPBP (TAPASIN)-related gene, TAPBP-R. Europ.
[69565] J. Immun. 32: 1059-1068, 2002.

[69566] 11732.

[69567] Yabe, T.; Kawamura, S.; Sato, M.; Kashiwase, K.; Tanaka, H.; Ishikawa,
[69568] Y.; Asao, Y.; Oyama, J.; Tsuruta, K.; Tokunaga, K.; Tadokoro, K.;
[69569] Juji, T.: A subject with a novel type I bare lymphocyte syndrome
[69570] has tapasin deficiency due to deletion of 4 exons by Alu-mediated
[69571] recombination. Blood 100: 1496-1498, 2002.

[69572] 11733.

[69573] Blank, V.; Kim, M. J.; Andrews, N. C.: Human MAFG is a functional
[69574] partner for p45 NF-E2 in activating globin gene expression. Blood 89:
[69575] 3925-3935, 1997.

[69576] 11734.

[69577] Blank, V.; Knoll, J. H. M.; Andrews, N. C.: Molecular char-

acterization

[69578] and localization of the human MafG gene. Genomics 44: 147-149, 1997.

[69579] 11735.

[69580] Estevez, R.; Boettger, T.; Stein, V.; Birkenhager, R.; Otto, E.;

[69581] Hildebrandt, F.; Jentsch, T. J.: Barttin is a Cl- channel beta-subunit

[69582] crucial for renal Cl- reabsorption and inner ear K+ secretion. Nature 414:

[69583] 558-561, 2001.

[69584] 11736.

[69585] Ishikawa, S.; Takahashi, T.; Ogawa, M.; Nakamura, Y.: Genomic

[69586] structure of the human PLCD1 (phospholipase C delta 1) locus on 3p22-p21.3. Cytogenet.

[69587] Cell Genet. 78: 58-60, 1997.

[69588] 11737.

[69589] Branda, S. S.; Yang, Z.; Chew, A.; Isaya, G.: Mitochondrial intermediate

[69590] peptidase and the yeast frataxin homolog together main-

tain mitochondrial
[69591] iron homeostasis in Saccharomyces cerevisiae. Hum. Molec. Genet. 8:
[69592] 1099-1110, 1999.

[69593] 11738.

[69594] Walter, L.; Gunther, E.: Identification of a novel highly conserved
[69595] gene in the centromeric part of the major histocompatibility complex. Genomics 52:
[69596] 298-304, 1998.

[69597] 11739.

[69598] Howell, B. W.; Hawkes, R.; Soriano, P.; Cooper, J. A.: Neuronal
[69599] position in the developing brain is regulated by mouse disabled-1. Nature 389:
[69600] 733-737, 1997.

[69601] 11740.

[69602] Lambert de Rouvroit, C.; Goffinet, A. M.: Cloning of human DAB1
[69603] and mapping to chromosome 1p31-p32. Genomics 53: 246-247, 1998.

[69604] 11741.

[69605] Sheldon, M.; Rice, D. S.; D'Arcangelo, G.; Yoneshima, H.; Nakajima,

[69606] K.; Mikoshiba, K.; Howell, B. W.; Cooper, J. A.; Goldwitz, D.; Curran,

[69607] T.: Scrambler and yotari disrupt the disabled gene and produce a

[69608] reeler-like phenotype in mice. Nature 389: 730-733, 1997.

[69609] 11742.

[69610] Ware, M. L.; Fox, J. W.; Gonzalez, J. L.; Davis, N. M.; Lambert

[69611] de Rouvroit, C.; Russo, C. J.; Chua, S. C., Jr.; Goffinet, A. M.;

[69612] Walsh, C. A.: Aberrant splicing of a mouse disabled homolog, mdab1,

[69613] in the scrambler mouse. Neuron 19: 239-249, 1997.

[69614] 11743.

[69615] Liu, J.; Dalmau, J.; Szabo, A.; Rosenfeld, M.; Huber, J.; Furneaux,

[69616] H.: Paraneoplastic encephalomyelitis antigens bind to the AU-rich

[69617] elements of mRNA. Neurology 45: 544-550, 1995.

[69618] 11744.

[69619] Sakai, K.; Gofuku, M.; Kitagawa, Y.; Ogasawara, T.; Hirose, G.;

[69620] Yamazaki, M.; Koh, C.-S.; Yanagisawa, N.; Steinman, L.: A hippocampal

[69621] protein associated with paraneoplastic neurologic syndrome and small

[69622] cell lung carcinoma. Biochem. Biophys. Res. Commun. 199: 1200-1208,

[69623] 1994.

[69624] 11745.

[69625] Van Tine, B. A.; Knops, J. F.; Butler, A.; Deloukas, P.; Shaw,

[69626] G. M.; King, P. H.: Localization of HuC (ELAVL3) to chromosome 19p13.2

[69627] by fluorescence in situ hybridization utilizing a novel tyramide labeling

[69628] technique. Genomics 53: 296-299, 1998.

[69629] 11746.

[69630] Whitmore, T. E.; Maurer, M. F.; Sexson, S.; Raymond, F.; Conklin,

[69631] D.; Deisher, T. A.: Assignment of fibroblast growth factor 18 (FGF18)
[69632] to human chromosome 5q34 by use of radiation hybrid mapping and fluorescence
[69633] in situ hybridization. Cytogenet. Cell Genet. 90: 231-233, 2000.
[69634] 11747.

[69635] Sherrington, R.; Rogaev, E. I.; Liang, Y.; Rogaeva, E. A.; Levesque,
[69636] G.; Ikeda, M.; Chi, H.; Lin, C.; Li, G.; Holman, K.; Tsuda, T.; Mar.;
[69637] and 21 others: Cloning of a gene bearing missense mutations in
[69638] early-onset familial Alzheimer's disease. Nature 375: 754-760, 1995.
[69639] 11748.

[69640] Verdi, J. M.; Bashirullah, A.; Goldhawk, D. E.; Kubu, C. J.; Jamali,
[69641] M.; Meakin, S. O.; Lipshitz, H. D.: Distinct human NUMB isoforms
[69642] regulate differentiation vs. proliferation in the neuronal lineage. Proc.

[69643] Nat. Acad. Sci. 96: 10472-10476, 1999.

[69644] 11749.

[69645] Zhong, W.; Feder, J. N.; Jiang, M.-M.; Jan, L. Y.; Jan, Y. N.:

[69646] Asymmetric localization of a mammalian Numb homolog during mouse cortical

[69647] neurogenesis. Neuron 17: 43-53, 1996.

[69648] 11750.

[69649] Thresher, R. J.; Vitaterna, M. H.; Miyamoto, Y.; Kazantsev, A.;

[69650] Hsu, D. S.; Petit, C.; Selby, C. P.; Dawut, L.; Smithies, O.; Takahashi,

[69651] J. S.; Sancar, A.: Role of mouse cryptochrome blue-light photoreceptor

[69652] in circadian photoresponses. Science 282: 1490-1494, 1998.

[69653] 11751.

[69654] Strehl, S.; LaSalle, J. M.; Lalande, M.: High-resolution anal-ysis

[69655] of DNA replication domain organization across an R/G-band boundary. Molec.

[69656] Cell. Biol. 17: 6157-6166, 1997.

[69657] 11752.

[69658] Chow, V. T. K.; Lee, S. S.: DENN, a novel human gene differentially

[69659] expressed in normal and neoplastic cells. DNA Seq. 6: 263-273, 1996.

[69660] 11753.

[69661] Chow, V. T. K.; Lim, K. M.; Lim, D.: The human DENN gene: genomic

[69662] organization, alternative splicing, and localization to chromosome

[69663] 11p11.21-p11.22. Genome 41: 543-552, 1998.

[69664] 11754.

[69665] Peyrard, M.; Seroussi, E.; Sandberg-Nordqvist, A.-C.; Xie, Y.-G.;

[69666] Han, F.-Y.; Fransson, I.; Collins, J.; Dunham, I.; Kost-Alimova, M.;

[69667] Imreh, S.; Dumanski, J. P.: The human LARGE gene from 22q12.3-q13.1

[69668] is a new, distinct member of the glycosyltransferase gene family. Proc.

[69669] Nat. Acad. Sci. 96: 598-603, 1999.

[69670] 11755.

[69671] McKusick, V. A.: Personal Communication. Baltimore, Md. 1/1/2000.

[69672] 11756.

[69673] Mykkanen, J.; Torrents, D.; Pineda, M.; Camps, M.; Yoldi, M. E.;

[69674] Horelli-Kuitunen, N.; Huoponen, K.; Heinonen, M.; Oksa-

[69675] nen, J.; Simell, O.; Savontaus, M.-L.; Zorzano, A.; Palacin, M.; Aula, P.: Functional

[69676] analysis of novel mutations in y+LAT-1 amino acid trans-

[69677] porter gene causing lysinuric protein intolerance (LPI). Hum. Molec. Genet. 9:

[69678] 431-438, 2000.

[69679] 11757.

[69680] Noguchi, A.; Shoji, Y.; Koizumi, A.; Takahashi, T.; Shoji, Y.;

[69681] Matsumori, M.; Kayo, T.; Ohata, T.; Wada, Y.; Yoshimura, I.; Maisawa,

[69682] S.; Konishi, M.; Takasago, Y.; Takada, G.: SLC7A7 genomic structure

[69683] and novel variants in three Japanese lysinuric protein intolerance
[69684] families. Hum. Mutat. 15: 367-372, 2000.

[69685] 11758.

[69686] Rosenfeld, J.; Sloan-Brown, K.; George, A. L., Jr.: A novel muscle
[69687] sodium channel mutation causes painful congenital myotonia. Ann.
[69688] Neurol. 42: 811-814, 1997.

[69689] 11759.

[69690] Gerlai, R.; Roder, J. C.; Hampson, D. R.: Altered spatial learning
[69691] and memory in mice lacking the mGluR4 subtype of metabotropic glutamate
[69692] receptor. Behav. Neurosci. 112: 525-532, 1998.

[69693] 11760.

[69694] Chatterjee, T. K.; Eapen, A.; Kanis, A. B.; Fisher, R. A.: Genomic
[69695] organization, 5-prime-flanking region, and chromosomal localization
[69696] of the human RGS3 gene. Genomics 45: 429-433, 1997.

[69697] 11761.

[69698] Druey, K. M.; Blumer, K. J.; Kang, V. H.; Kehrl, J. H.: Inhibi-
tion
[69699] of G-protein-mediated MAP kinase activation by a new
mammalian gene
[69700] family. Nature 379: 742-746, 1996.

[69701] 11762.

[69702] Oettgen, P.; Alani, R. M.; Barcinski, M. A.; Brown, L.; Ak-
barali,
[69703] Y.; Boltax, J.; Kunsch, C.; Munger, K.; Liebermann, T. A.:
Isolation
[69704] and characterization of a novel epithelium-specific tran-
scription
[69705] factor, ESE-1, a member of the Ets family. Molec. Cell.
Biol. 17:
[69706] 4419-4433, 1997.

[69707] 11763.

[69708] Oettgen, P.; Barcinski, M.; Boltax, J.; Stolt, P.; Akbarali, Y.;

[69709] Libermann, T. A.: Genomic organization of the human
ELF3 (ESE-1/ESX)
[69710] gene, a member of the Ets transcription factor family, and

identification

[69711] of a functional promoter. Genomics 55: 358-362, 1999.

[69712] 11764.

[69713] Oettgen, P.; Carter, K. C.; Augustus, M.; Barcinski, M.; Boltax,

[69714] J.; Kunsch, C.; Libermann, T. A.: The novel epithelial-specific Ets

[69715] transcription factor gene ESX maps to human chromosome 1q32.1. Genomics 45:

[69716] 456-457, 1997.

[69717] 11765.

[69718] Lunn, C. A.; Fan, X.; Dalie, B.; Miller, K.; Zavodny, P. J.; Narula,

[69719] S. K.; Lundell, D.: Purification of ADAM 10 from bovine spleen as

[69720] a TNFalpha convertase. FEBS Lett. 400: 333-335, 1997.

[69721] 11766.

[69722] Tymms, M. J.; Ng, A. Y. N.; Thomas, R. S.; Schutte, B. C.; Zhou,

[69723] J.; Eyre, H. J.; Sutherland, G. R.; Seth, A.; Rosenberg, M.; Papas,

[69724] T.; Debouck, C.; Kola, I.: A novel epithelial-expressed ETS gene,

[69725] ELF3: human and murine cDNA sequences, murine genomic organization,

[69726] human mapping to 1q32.2 and expression in tissues and cancer. Oncogene 15:

[69727] 2449-2462, 1997.

[69728] 11767.

[69729] Wolfsberg, T. G.; Primakoff, P.; Myles, D. G.; White, J. M.: ADAM,

[69730] a novel family of membrane proteins containing a disintegrin and metalloprotease

[69731] domain: multipotential functions in cell-cell and cell-matrix interactions. J.

[69732] Cell Biol. 131: 275-278, 1995.

[69733] 11768.

[69734] Yamazaki, K.; Mizui, Y.; Tanaka, I.: Radiation hybrid mapping

[69735] of human ADAM10 gene to chromosome 15. Genomics 45: 457-459, 1997.

[69736] 11769.

[69737] Choi, D.-S.; Handa, M.; Young, H.; Gordon, A. S.; Diamond, I.;

[69738] Messing, R. O.: Genomic organization and expression of the mouse

[69739] equilibrative, nitrobenzylthioinosine-sensitive nucleoside transporter

[69740] 1 (ENT1) gene. Biochem. Biophys. Res. Commun. 277: 200-208, 2000.

[69741] 11770.

[69742] Coe, I. R.; Griffiths, M.; Young, J. D.; Baldwin, S. A.; Cass,

[69743] C. E.: Assignment of the human equilibrative nucleoside transporter

[69744] (hENT1) to 6p21.1-p21.2. Genomics 45: 459-460, 1997.

[69745] 11771.

[69746] Griffiths, M.; Beaumont, N.; Yao, S. Y. M.; Sundaram, M.; Boumah,

[69747] C. E.; Davies, A.; Kwong, F. Y. P.; Coe, I.; Cass, C. E.; Young, J.

[69748] D.; Baldwin, S. A.: Cloning of a human nucleoside transporter implicated

[69749] in the cellular uptake of adenosine and chemotherapeutic drugs. Nature

[69750] Med. 3: 89-94, 1997.

[69751] 11772.

[69752] Hu, S.-I.; Carozza, M.; Klein, M.; Nantermet, P.; Luk, D.; Crowl,

[69753] R. M.: Human HtrA, an evolutionarily conserved serine protease identified

[69754] as a differentially expressed gene product in osteoarthritic cartilage. J.

[69755] Biol. Chem. 273: 34406-34412, 1998.

[69756] 11773.

[69757] Zumbrunn, J.; Trueb, B.: Localization of the gene for a serine

[69758] protease with IGF-binding domain (PRSS11) to human chromosome 10q25.3-q26.2. Genomics 45:

[69759] 461-462, 1997.

[69760] 11774.

[69761] Zumbrunn, J.; Trueb, B.: Primary structure of a putative serine

[69762] protease specific for IGF-binding proteins. FEBS Lett. 398: 187-192,

[69763] 1996.

[69764] 11775.

[69765] Debelenko, L. V.; Emmert-Buck, M. R.; Manickam, P.; Kester, M.-B.;

[69766] Guru, S. C.; DiFranco, E. M.; Olufemi, S.-E.; Agarwal, S.; Lubensky,

[69767] I. A.; Zhuang, Z.; Burns, A. L.; Spiegel, A. M.; Liotta, L. A.; Collins,

[69768] F. S.; Marx, S. J.; Chandrasekharappa, S. C.: Haplotype analysis

[69769] defines a minimum interval for the multiple endocrine neoplasia type

[69770] 1 (MEN1) gene. Cancer Res. 57: 1039-1042, 1997.

[69771] 11776.

[69772] D'Esposito, M.; Strazzullo, M.; Cuccurese, M.; Spalluto, C.; Rocchi,

[69773] M.; D'Urso, M.; Ciccodicola, A.: Identification and assign-ment of

[69774] the human transient receptor potential channel 6 gene TRPC6 to chromosome

[69775] 11q21-q22. Cytogenet. Cell Genet. 83: 46-47, 1998.

[69776] 11777.

[69777] Hofmann, T.; Obukhov, A. G.; Schaefer, M.; Harteneck, C.; Gudermann,
[69778] T.; Schultz, G.: Direct activation of human TRPC6 and TRPC3 channels
[69779] by diacylglycerol. Nature 397: 259-263, 1999.

[69780] 11778.

[69781] Garcia, C. K.; Brown, M. S.; Pathak, R. K.; Goldstein, J. L.:
[69782] cDNA cloning of MCT2, a second monocarboxylate transporter expressed
[69783] in different cells than MCT1. J. Biol. Chem. 270: 1843-1849, 1995.

[69784] 11779.

[69785] Lin, R.-Y.; Vera, J. C.; Chaganti, R. S. K.; Golde, D. W.: Human
[69786] monocarboxylate transporter 2 (MCT2) is a high affinity pyruvate transporter. J.
[69787] Biol. Chem. 273: 28959-28965, 1998.

[69788] 11780.

[69789] Ong, O. C.; Hu, K.; Rong, H.; Lee, R. H.; Fung, B. K.-K.: Gene
[69790] structure and chromosome localization of the G-gamma-c

subunit of
[69791] human cone G-protein (GNGT2). Genomics 44: 101-109, 1997.
[69792] 11781.

[69793] Ong, O. C.; Yamane, H. K.; Phan, K. B.; Fong, H. K.; Bok, D.; Lee,
[69794] R. H.; Fung, B. K.-K.: Molecular cloning and characterization of
[69795] the G protein gamma subunit of cone photoreceptors. J. Biol. Chem. 270:
[69796] 8495-8500, 1995.

[69797] 11782.

[69798] Kohlhase, J.; Schuh, R.; Dowe, G.; Kuhnlein, R. P.; Jackle, H.;
[69799] Schroeder, B.; Schulz-Schaeffer, W.; Kretzschmar, H. A.; Kohler, A.;
[69800] Muller, U.; Raab-Vetter, M.; Burkhardt, E.; Engel, W.; Stick, R.:
[69801] Isolation, characterization, and organ-specific expression of two
[69802] novel human zinc finger genes related to the Drosophila gene spalt. Genomics 38:

[69803] 291-298, 1996.

[69804] 11783.

[69805] Park, S. H.; Ryu, S. H.; Suh, P. G.; Kim, H.: Assignment of human

[69806] PLD2 to chromosome band 17p13.1 by fluorescence in situ hybridization. Cytogenet.

[69807] Cell Genet. 82: 225 only, 1998.

[69808] 11784.

[69809] Her, C.; Kaur, G. P.; Athwal, R. S.; Weinshilboum, R. M.: Human

[69810] sulfotransferase SULT1C1: cDNA cloning, tissue-specific expression,

[69811] and chromosomal localization. Genomics 41: 467-470, 1997.

[69812] 11785.

[69813] Keon, B. H.; Schafer, S.; Kuhn, C.; Grund, C.; Franke, W. W.:

[69814] Symplekin, a novel type of tight junction plaque protein. J. Cell

[69815] Biol. 134: 1003-1018, 1996.

[69816] 11786.

[69817] Ueki, K.; Ramaswamy, S.; Billings, S. J.; Mohrenweiser, H. W.;
[69818] Louis, D. N.: Chromosomal localization to 19q13.3, partial genomic
[69819] structure and 5-prime cDNA sequence of the human symplekin gene. Somat.
[69820] Cell Molec. Genet. 23: 229-231, 1997.

[69821] 11787.

[69822] Pucharcos, C.; Fuentes, J.-J.; Casas, C.; de la Luna, S.; Al-
cantara,
[69823] S.; Arbones, M. L.; Soriano, E.; Estivill, X.; Prichard, M.: Alu-splice
[69824] cloning of human intersectin (ITSN), a putative multivalent binding
[69825] protein expressed in proliferating and differentiating neurons and
[69826] overexpressed in Down syndrome. Europ. J. Hum. Genet. 7: 704-712,
[69827] 1999.

[69828] 11788.

[69829] Lucking, C. B.; Bonifati, V.; Periquet, M.; Vanacore, N.; Brice,

[69830] A.; Meco, G.: Pseudo-dominant inheritance and exon 2 triplication
[69831] in a family with parkin gene mutations. Neurology 57: 924-927, 2001.
[69832] 11789.

[69833] Kanezaki, R.; Toki, T.; Yokoyama, M.; Yomogida, K.; Sugiyama, K.;
[69834] Yamamoto, M.; Igarashi, K.; Ito, E.: Transcription factor BACH1 is
[69835] recruited to the nucleus by its novel alternative spliced isoform. J.
[69836] Biol. Chem. 276: 7278-7284, 2001.

[69837] 11790.

[69838] Shimozawa, N.; Suzuki, Y.; Zhang, Z.; Imamura, A.; Ghaedi, K.;
[69839] Fujiki, Y.; Kondo, N.: Identification of PEX3 as the gene mutated
[69840] in a Zellweger syndrome patient lacking peroxisomal remnant structures. Hum.
[69841] Molec. Genet. 9: 1995-1999, 2000.

[69842] 11791.

[69843] Katz, P.; Whalen, G.; Kehrl, J. H.: Differential expression of

[69844] a novel protein kinase in human B lymphocytes: preferential localization

[69845] in the germinal center. J. Biol. Chem. 269: 16802-16809, 1994.

[69846] 11792.

[69847] Pombo, C. M.; Kehrl, J. H.; Sanchez, I.; Katz, P.; Avruch, J.;

[69848] Zon, L. I.; Woodgett, J. R.; Force, T.; Kyriakis, J. M.: Activation

[69849] of the SAPK pathway by the human STE20 homologue germinal centre kinase. Nature 377:

[69850] 750-754, 1995.

[69851] 11793.

[69852] Ren, M.; Zeng, J.; De Lemos-Chiarandini, C.; Rosenfeld, M.; Adesnik,

[69853] M.; Sabatini, D. D.: In its active form, the GTP-binding protein

[69854] rab8 interacts with a stress-activated protein kinase. Proc. Nat.

[69855] Acad. Sci. 93: 5151-5155, 1996.

[69856] 11794.

[69857] Bonthron, D. T.; Hayward, B. E.; Moran, V.; Strain, L.: Characterization
[69858] of TH1 and CTSZ, two non-imprinted genes downstream of GNAS1 in chromosome
[69859] 20q13. Hum. Genet. 107: 165-175, 2000.

[69860] 11795.

[69861] Santamaria, I.; Velasco, G.; Pendas, A. M.; Fueyo, A.; Lopez-Otin,
[69862] C.: Cathepsin Z, a novel human cysteine proteinase with a short propeptide
[69863] domain and a unique chromosomal location. J. Biol. Chem. 273: 16816-16823,
[69864] 1998.

[69865] 11796.

[69866] Jacquemin, P.; Chen, Z.; Martial, J. A.; Davidson, I.: Ge-nomic
[69867] structure and chromosomal mapping of the mouse tran-scription factor
[69868] TEF-5 (Tead3) gene. Mammalian Genome 10: 632-634, 1999.
[69869] 11797.

[69870] Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D. M.;

[69871] Kay, B. K.: Cloning of ligand targets: systematic isolation of SH3

[69872] domain-containing proteins. Nature Biotech. 14: 741-744, 1996.

[69873] 11798.

[69874] Blau, N.; Niederwieser, A.; Shmerling, D. H.: Peptiduria presumably

[69875] caused by aminopeptidase-P deficiency: a new inborn error of metabolism. J.

[69876] Inherit. Metab. Dis. 11 (suppl): 240-242, 1988.

[69877] 11799.

[69878] Cottrell, G. S.; Hooper, N. M.; Turner, A. J.: Cloning, expression,

[69879] and characterization of human cytosolic aminopeptidase P: a single

[69880] manganese(II)-dependent enzyme. Biochemistry 39: 15121-15128, 2000.

[69881] 11800.

[69882] Sprinkle, T. J.; Caldwell, C.; Ryan, J. W.: Cloning, chromosomal

[69883] sublocalization of the human soluble aminopeptidase P gene (XPNPEP1)
[69884] to 10q25.3 and conservation of the putative proton shuttle and metal
[69885] ligand binding sites with XPNPEP2. Arch. Biochem. Biophys. 378:
[69886] 51-56, 2000.

[69887] 11801.

[69888] Vanhoof, G.; Goossens, F.; Juliano, M. A.; Juliano, L.; Hendriks,
[69889] D.; Schatteman, K.; Lin, A. H.; Scharpe, S.: Isolation and sequence
[69890] analysis of a human cDNA clone (XPNPEPL) homologous to X-prolyl aminopeptidase
[69891] (aminopeptidase P). Cytogenet. Cell Genet. 78: 275-280, 1997.
[69892] 11802.

[69893] Vergas Romero, C.; Neudorfer, I.; Mann, K.; Schafer, W.: Purification
[69894] and amino acid sequence of aminopeptidase P from pig kidney. Europ.
[69895] J. Biochem. 229: 262-269, 1995.

[69896] 11803.

[69897] Veugelers, M.; Vermeesch, J.; Reekmans, G.; Steinfeld, R.; Marynen,

[69898] P.; David, G.: Characterization of glypican-5 and chromosomal localization

[69899] of human GPC5, a new member of the glypican gene family. Genomics 40:

[69900] 24-30, 1997.

[69901] 11804.

[69902] Hegele, R. A.; Connelly, P. W.; Scherer, S. W.; Hanley, A. J. G.;

[69903] Harris, S. B.; Tsui, L.-C.; Zinman, B.: Paraoxonase-2 gene (PON2)

[69904] G148 variant associated with elevated fasting plasma glucose in noninsulin-dependent

[69905] diabetes mellitus. J. Clin. Endocr. Metab. 82: 3373-3377, 1997.

[69906] 11805.

[69907] Heinecke, J. W.; Lusis, A. J.: Paraoxonase-gene polymorphisms

[69908] associated with coronary heart disease: support for the oxidative

[69909] damage hypothesis? (Letter) Am. J. Hum. Genet. 62: 20-24, 1998.

[69910] 11806.

[69911] Primo-Parmo, S. L.; Hsu, C.; Law, D. J.; La Du, B. N.: Loca-
tion

[69912] and arrangement of three paraoxonase genes: PON1, PON2, and PON3,

[69913] on human chromosome 7. (Abstract) Am. J. Hum. Genet. 59 (suppl.):

[69914] A406, 1996.

[69915] 11807.

[69916] Primo-Parmo, S. L.; Sorenson, R. C.; Teiber, L.; La Du, B. N.:

[69917] The human serum paraoxonase/arylesterase gene (PON1) is one member

[69918] of multigene family. Genomics 33: 498-507, 1996.

[69919] 11808.

[69920] Prochazka, M.; Thompson, D. B.; Scherer, S. W.; Tsui, L.-C.; Knowler,

[69921] W. C.; Bennett, P. H.; Bogardus, C.: Linkage and associa-
tion of insulin

[69922] resistance and NIDDM with markers at 7q21.3-q22.1 in the Pima Indians.

[69923] (Abstract) Diabetes 44 (suppl.): 42A, 1995.

[69924] 11809.

[69925] Sanghera, D. K.; Aston, C. E.; Saha, N.; Kamboh, M. I.: DNA polymorphisms

[69926] in two paraoxonase genes (PON1 and PON2) are associated with the risk

[69927] of coronary heart disease. Am. J. Hum. Genet. 62: 36-44, 1998.

[69928] 11810.

[69929] Chang, H. Y.; Nishitoh, H.; Yang, X.; Ichijo, H.; Baltimore, D.

[69930] : Activation of apoptosis signal-regulating kinase 1 (ASK1) by the

[69931] adapter protein Daxx. Science 281: 1860-1863, 1998.

[69932] 11811.

[69933] Geleziunas, R.; Xu, W.; Takeda, K.; Ichijo, H.; Greene, W. C.:

[69934] HIV-1 Nef inhibits ASK1-dependent death signalling providing a potential

[69935] mechanism for protecting the infected host cell. Nature 410: 834-838,

[69936] 2001.

[69937] 11812.

[69938] Ichijo, H.; Nishida, E.; Irie, K.; ten Dijke, P.; Saitoh, M.; Moriguchi,

[69939] T.; Takagi, M.; Matsumoto, K.; Miyazono, K.; Gotoh, Y.: In-duction

[69940] of apoptosis by ASK1, a mammalian MAPKKK that acti-vates SAPK/JNK and

[69941] p38 signaling pathways. Science 275: 90-94, 1997.

[69942] 11813.

[69943] Nishitoh, H.; Saitoh, M.; Mochida, Y.; Takeda, K.; Nakano, H.;

[69944] Rothe, M.; Miyazono, K.; Ichijo, H.: ASK1 is essential for JNK/SAPK

[69945] activation by TRAF2. Molec. Cell 2: 389-395, 1998.

[69946] 11814.

[69947] Lucking, C. B.; Durr, A.; Bonifati, V.; Vaughan, J.; De Michele,

[69948] G.; Gasser, T.; Harhangi, B. S.; Meco, G.; Denefle, P.;

Wood, N. W.;

[69949] Agid, Y.; Brice, A.: Association between early-onset Parkinson's

[69950] disease and mutations in the parkin gene. New Eng. J. Med. 342:

[69951] 1560-1567, 2000.

[69952] 11815.

[69953] Nisipeanu, P.; Inzelberg, R.; Mouch, S. A.; Carasso, R. L.; Blumen,

[69954] S. C.; Zhang, J.; Matsumine, H.; Hattori, N.; Mizuno, Y.: Parkin

[69955] gene causing benign autosomal recessive juvenile parkinsonism. Neurology 56:

[69956] 1573-1575, 2001.

[69957] 11816.

[69958] Shimura, H.; Hattori, N.; Kubo, S.; Mizuno, Y.; Asakawa, S.; Minoshima,

[69959] S.; Shimizu, N.; Iwai, K.; Chiba, T.; Tanaka, K.; Suzuki, T.: Familial

[69960] Parkinson disease gene product, parkin, is a ubiquitin-protein ligase. Nature

[69961] Genet. 25: 302-305, 2000.

[69962] 11817.

[69963] Periquet, M.; Lucking, C. B.; Vaughan, J. R.; Bonifati, V.; Durr,

[69964] A.; De Michele, G.; Horstink, M. W.; Farrer, M.; Illarioshkin, S.

[69965] N.; Pollak, P.; Borg, M.; Brefel-Courbon, C.; Denefle, P.; Meco, G.;

[69966] Gasser, T.; Breteler, M. M. B.; Wood, N. W.; Agid, Y.; Brice, A.;

[69967] French Parkinson's Disease Genetics Study Group; Euro-

[69968] pean Consortium on Genetic Susceptibility in Parkinson's Diseases: Origin of mutations

[69969] in the parkin gene in Europe: exon rearrangements are in-

[69970] dependent recurrent events, whereas point mutations may result from founder

[69971] effects. Am. J. Hum. Genet. 68: 617-626, 2001.

[69972] 11818.

[69973] Tomac, A. C; Hoffer, B. J.: Assignment of the mouse Park2 (PARKIN),

[69974] the homologue to a new human Parkinson candidate

gene, to the telomeric
[69975] region of mouse 17A3.2-3.3, by in situ hybridization. Cytogenet.
[69976] Cell Genet. 95: 120-121, 2001.

[69977] 11819.

[69978] Zhang, Y.; Gao, J.; Chung, K. K. K.; Huang, H.; Dawson, V. L.;
[69979] Dawson, T. M.: Parkin functions as an E2-dependent ubiquitin-protein
[69980] ligase and promotes the degradation of the synaptic vesicle-associated
[69981] protein, CDCrel-1. Proc. Nat. Acad. Sci. 97: 13354-13359, 2000.
[69982] 11820.

[69983] Tomasetto, C.; Regnier, C.; Moog-Lutz, C.; Mattei, M. G.; Chenard,
[69984] M. P.; Lidereau, R.; Basset, P.; Rio, M. C.: Identification of four
[69985] novel human genes amplified and overexpressed in breast carcinoma
[69986] and localized to the q11-q21.3 region of chromosome 17. Genomics 28:

[69987] 367-376, 1995.

[69988] 11821.

[69989] Chalmers, I. J.; Hofler, H.; Atkinson, M. J.: Mapping of a cadherin

[69990] gene cluster to a region of chromosome 5 subject to frequent allelic

[69991] loss in carcinoma. Genomics 57: 160-163, 1999.

[69992] 11822.

[69993] Suzuki, S.; Sano, K.; Tanihara, H.: Diversity of the cadherin

[69994] family: evidence for eight new cadherins in nervous tissue. Cell

[69995] Regul. 2: 261-270, 1991.

[69996] 11823.

[69997] Shimoyama, Y.; Gotoh, M.; Terasaki, T.; Kitajima, M.; Hirohashi,

[69998] S.: Isolation and sequence analysis of human cadherin-6 complementary

[69999] DNA for the full coding sequence and its expression in human carcinoma

[70000] cells. Cancer Res. 55: 2206-2211, 1995.

[70001] 11824.

[70002] Anderson, L. V. B.; Davison, K.; Moss, J. A.; Young, C.; Cullen,

[70003] M. J.; Walsh, J.; Johnson, M. A.; Bashir, R.; Britton, S.; Keers,

[70004] S.; Argov, Z.; Mahjneh, I.; Fougerousse, F.; Beckmann, J. S.; Bushby,

[70005] K. M. D.: Dysferlin is a plasma membrane protein and is expressed

[70006] early in human development. Hum. Molec. Genet. 8: 855-861, 1999.

[70007] 11825.

[70008] Vaz, F. M.; van Gool, S.; Ofman, R.; Ijlst, L.; Wanders, R. J.

[70009] A.: Carnitine biosynthesis: identification of the cDNA encoding human

[70010] gamma-butyrobetaine hydroxylase. Biochem. Biophys. Res. Commun. 250:

[70011] 506-510, 1998.

[70012] 11826.

[70013] Gong, B.; Almasan, A.: Genomic organization and transcriptional

[70014] regulation of human Apo2/TRAIL gene. Biochem. Biophys. Res. Commun. 278:

[70015] 747-752, 2000.

[70016] 11827.

[70017] Ohira, M.; Seki, N.; Nagase, T.; Ishikawa, K.; Nomura, N.; Ohara,

[70018] O.: Characterization of a human homolog (BACH1) of the mouse Bach1

[70019] gene encoding a BTB-basic leucine zipper transcription factor and

[70020] its mapping to chromosome 21q22.1. Genomics 47: 300-306, 1998.

[70021] 11828.

[70022] Oyake, T.; Itoh, K.; Motohashi, H.; Hayashi, N.; Hoshino, H.; Nishizawa,

[70023] M.; Yamamoto, M.; Igarashi, K.: Bach proteins belong to a novel family

[70024] of BTB-basic leucine zipper transcription factors that interact with

[70025] MafK and regulate transcription through the NF-E2 site. Molec. Cell.

[70026] Biol. 16: 6083-6095, 1996.

[70027] 11829.

[70028] Nothwang, H. G.; Rensing, C.; Kubler, M.; Denich, D.; Brandl, B;

[70029] Stubanus, M.; Haaf, T.; Kurnit, D.; Hildebrandt, F.: Identification

[70030] of a novel Ran binding protein 2 related gene (RANBP2L1) and detection

[70031] of a gene cluster on human chromosome 2q11-q12. Genomics 47: 383-392,

[70032] 1998.

[70033] 11830.

[70034] Ghanshani, S.; Coleman, M.; Gustavsson, P.; Wu, A. C.-L.; Gargus,

[70035] J. J.; Gutman, G. A.; Dahl, N.; Mohrenweiser, H.; Chandy, K. G.:

[70036] Human calcium-activated potassium channel gene KCNN4 maps to chromosome

[70037] 19q13.2 in the region deleted in Diamond-Blackfan anemia. Genomics 51:

[70038] 160-161, 1998.

[70039] 11831.

[70040] Ishii, T. M.; Silvia, C.; Hirschberg, B.; Bond, C. T.; Adelman,

[70041] J. P.; Maylie, J.: A human intermediate conductance calcium-activated

[70042] potassium channel. Proc. Nat. Acad. Sci. 94: 11651-11656, 1997.

[70043] 11832.

[70044] Joiner, W. J.; Wang, L.-Y.; Tang, M. D.; Kaczmarek, L. K.: hSK4,

[70045] a member of a novel subfamily of calcium-activated potassium channels. Proc.

[70046] Nat. Acad. Sci. 94: 11013-11018, 1997.

[70047] 11833.

[70048] Logsdon, N. J.; Kang, J.; Togo, J. A.; Christian, E. P.; Aiyar,

[70049] J.: A novel gene, hKCa4, encodes the calcium-activated potassium

[70050] channel in human T lymphocytes. J. Biol. Chem. 272: 32723-32726,

[70051] 1997.

[70052] 11834.

[70053] Matsuoka, H.; Iwata, N.; Ito, M.; Shimoyama, M.; Nagata, A.; Chihara,

[70054] K.; Takai, S.; Matsui, T.: Expression of a kinase-defective Eph-like
[70055] receptor in the normal human brain. Biochem. Biophys. Res. Commun. 235:
[70056] 487-492, 1997.

[70057] 11835.

[70058] Tang, X. X.; Zhao, H.; Robinson, M. E.; Cohen, B.; Cnaan, A.; London,
[70059] W.; Cohn, S. L.; Cheung, N.-K. V.; Brodeur, G. M.; Evans, A. E.; Ikegaki,
[70060] N.: Implications of EPHB6, EFNB2, and EFNB3 expressions in human
[70061] neuroblastoma. Proc. Nat. Acad. Sci. 97: 10936-10941, 2000.
[70062] 11836.

[70063] Meyers, R.; Cantley, L. C.: Cloning and characterization of a
[70064] wortmannin-sensitive human phosphatidylinositol 4-kinase. J. Biol.
[70065] Chem. 272: 4384-4390, 1997.

[70066] 11837.

[70067] Nakanishi, S.; Catt, K. J.; Balla, T.: A wortmannin-sensitive

[70068] phosphatidylinositol 4-kinase that regulates hormone-sensitive pools

[70069] of inositolphospholipids. Proc. Nat. Acad. Sci. 92: 5317-5321, 1995.

[70070] 11838.

[70071] Saito, T.; Seki, N.; Ishii, H.; Ohira, M.; Hayashi, A.; Kozuma,

[70072] S.; Hori, T.: Complementary DNA cloning and chromoso-mal mapping of

[70073] a novel phosphatidylinositol kinase gene. DNA Res. 4: 301-305, 1997.

[70074] 11839.

[70075] Suzuki, K.; Hirano, H.; Okutomi, K.; Suzuki, M.; Kuga, Y.; Fujiwara,

[70076] T.; Kanemoto, N.; Isono, K.; Horie, M.: Identification and characterization

[70077] of a novel human phosphatidylinositol 4-kinase. DNA Res. 4: 273-280,

[70078] 1997.

[70079] 11840.

[70080] Miura, Y.; Miyake, K.; Yamashita, Y.; Shimazu, R.; Copeland, N.

[70081] G.; Gilbert, D. J.; Jenkins, N. A.; Inazawa, J.; Abe, T.; Ki-

[70082] moto,

[70083] M.: Molecular cloning of a human RP105 homologue and chromosomal

[70084] localization of the mouse and human RP105 genes (Ly64 and LY64). Genomics 38: 299-304, 1996.

[70085] 11841.

[70086] Miura, Y.; Shimazu, R.; Miyake, K.; Akashi, S.; Ogata, H.; Yamashita,

[70087] Y.; Narisawa, Y.; Kimoto, M.: RP105 is associated with MD-1 and transmits

[70088] an activation signal in human B cells. Blood 92: 2815-2822, 1998.

[70089] 11842.

[70090] Herbarth, B.; Pingault, V.; Bondurand, N.; Kuhlbrodt, K.; Hermans-Borgmeyer,

[70091] I.; Puliti, A.; Lemort, N.; Goossens, M.; Wegner, M.: Muta-

[70092] tion of the Sry-related Sox10 gene in dominant megacolon, a

mouse model for
[70093] human Hirschsprung disease. Proc. Nat. Acad. Sci. 95:
5161-5165,
[70094] 1998.

[70095] 11843.

[70096] Khong, H. T.; Rosenberg, S. A.: The Waardenburg syn-
drome type
[70097] 4 gene, SOX10, is a novel tumor-associated antigen iden-
tified in a
[70098] patient with a dramatic response to immunotherapy. Can-
cer Res. 62:
[70099] 3020-3023, 2002.

[70100] 11844.

[70101] Kuhlbrodt, K.; Herbarth, B.; Sock, E.; Hermans-Borgmeyer,
I.; Wegner,
[70102] M.: Sox10, a novel transcriptional modulator in glial cells.
J.
[70103] Neurosci. 18: 237-250, 1998.

[70104] 11845.

[70105] Kuhlbrodt, K.; Schmidt, C.; Sock, E.; Pingault, V.; Bon-
durand,

[70106] N.; Goossens, M.; Wegner, M.: Functional analysis of Sox10 mutations
[70107] found in human Waardenburg-Hirschsprung patients. J. Biol. Chem. 273:
[70108] 23033-23038, 1998.

[70109] 11846.

[70110] Thompson, C. C.; Bottcher, M. C.: The product of a thyroid hormone-responsive
[70111] gene interacts with thyroid hormone receptors. Proc. Nat. Acad. Sci. 94:
[70112] 8527-8532, 1997.

[70113] 11847.

[70114] Zlotogorski, A.; Ahmad, W.; Christiano, A. M.: Congenital atrichia
[70115] in five Arab Palestinian families resulting from a deletion mutation
[70116] in the human hairless gene. Hum. Genet. 103: 400-404, 1998.
[70117] 11848.

[70118] Ando, A.; Kikuti, Y. Y.; Abe, K.; Shigenari, A.; Kawata, H.; Ikemura,

[70119] T.; Kimura, M.; Inoko, H.: cDNA cloning, northern hybridization,
[70120] and mapping analysis of a putative GDS-related protein gene at the
[70121] centromeric ends of the human and mouse MHC regions. Immunogenetics 49:
[70122] 354-356, 1999.

[70123] 11849.

[70124] Isomura, M.; Okui, K.; Fujiwara, T.; Shin, S.; Nakamura, Y.: Isolation
[70125] and mapping of RAB2L, a human cDNA that encodes a protein homologous
[70126] to RalGDS. Cytogenet. Cell Genet. 74: 263-265, 1996.

[70127] 11850.

[70128] Pirozzi, G.; McConnell, S. J.; Uveges, A. J.; Carter, J. M.; Sparks,
[70129] A. B.; Kay, B. K.; Fowlkes, D. M.: Identification of novel human
[70130] WW domain-containing proteins by cloning of ligand targets. J. Biol.
[70131] Chem. 272: 14611-14616, 1997.

[70132] 11851.

[70133] Chidambaram, A.; Allikmets, R.; Chandrasekarappa, S.; Guru, S.

[70134] C.; Modi, W.; Gerrard, B.; Dean, M.: Characterization of a human

[70135] homolog (OVOL1) of the Drosophila ovo gene, which maps to chromosome

[70136] 11q13. Mammalian Genome 8: 950-951, 1997.

[70137] 11852.

[70138] Motoki, K.; Megahed, M.; LaForgia, S.; Uitto, J.: Cloning and

[70139] chromosomal mapping of mouse ladinin, a novel base-ment membrane zone

[70140] component. Genomics 39: 323-330, 1997.

[70141] 11853.

[70142] Chambers, J. A.; Gardner, E.; Hauptmann, R.; Ponder, B. A.; Mulligan,

[70143] L. M.: TaqI polymorphisms at the annexin VIII locus (ANX8). Hum.

[70144] Molec. Genet. 1: 550 only, 1992.

[70145] 11854.

[70146] Chang, K.-S.; Wang, G.; Freireich, E. J.; Daly, M.; Naylor, S.

[70147] L.; Trujillo, J. M.; Stass, S. A.: Specific expression of the annexin

[70148] VIII gene in acute promyelocytic leukemia. Blood 79: 1802-1810,

[70149] 1992.

[70150] 11855.

[70151] Fernandez, M. P.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[70152] A.; Morgan, R. O.: The genetic origin of mouse annexin VIII. Mammalian

[70153] Genome 9: 8-14, 1998.

[70154] 11856.

[70155] Sarkar, A.; Yang, P.; Fan, Y.-H.; Mu, Z. M.; Hauptmann, R.; Adolf,

[70156] G. R.; Stass, S. A.; Chang, K.-S.: Regulation of the expres- sion of

[70157] annexin VIII in acute promyelocytic leukemia. Blood 84: 279-286,

[70158] 1994.

[70159] 11857.

[70160] Waterham, H. R.; Koster, J.; Romeijn, G. J.; Hennekam, R. C. M.;

[70161] Vreken, P.; Andersson, H. C.; FitzPatrick, D. R.; Kelley, R. I.; Wanders,

[70162] R. J. A.: Mutations in the 3-beta-hydroxysterol delta-24-reductase

[70163] gene cause desmosterolosis, an autosomal recessive disorder of cholesterol

[70164] biosynthesis. Am. J. Hum. Genet. 69: 685-694, 2001.

[70165] 11858.

[70166] Lechner, C.; Zahalka, M. A.; Giot, J.-F.; Moller, N. P. H.; Ullrich,

[70167] A.: ERK6, a mitogen-activated protein kinase involved in C2C12 myoblast

[70168] differentiation. Proc. Nat. Acad. Sci. 93: 4355-4359, 1996.

[70169] 11859.

[70170] Li, Z.; Jiang, Y.; Ulevitch, R. J.; Han, J.: The primary structure

[70171] of p38-gamma: a new member of p38 group of MAP kinases. Biochem.

[70172] Biophys. Res. Commun. 228: 334-340, 1996.

[70173] 11860.

[70174] Hibi, M.; Lin, A.; Smeal, T.; Minden, A.; Karin, M.: Identification

[70175] of an oncoprotein- and UV-responsive protein kinase that binds and

[70176] potentiates the c-Jun activation domain. Genes Dev. 7: 2135-2148,

[70177] 1993.

[70178] 11861.

[70179] Kallunki, T.; Su, B.; Tsigelny, I.; Sluss, H. K.; Derijard, B.;

[70180] Moore, G.; Davis, R.; Karin, M.: JNK2 contains a specificity-determining

[70181] region responsible for efficient c-Jun binding and phosphorylation. Genes

[70182] Dev. 8: 2996-3007, 1994.

[70183] 11862.

[70184] Sluss, H. K.; Barrett, T.; Derijard, B.; Davis, R. J.: Signal

[70185] transduction by tumor necrosis factor mediated by JNK protein kinases. Molec.

[70186] Cell Biol. 14: 8376-8384, 1994.

[70187] 11863.

[70188] Mohit, A. A.; Martin, J. H.; Miller, C. A.: p49(3F12) kinase:

[70189] a novel MAP kinase expressed in a subset of neurons in the human nervous

[70190] system. Neuron 14: 67-78, 1995.

[70191] 11864.

[70192] Yang, D. D.; Kuan, C. Y.; Whitmarsh, A. J.; Rincon, M.; Zheng,

[70193] T. S.; Davis, R. J.; Rakic, P.; Flavell, R. A.: Absence of exci-

totoxicity-induced

[70194] apoptosis in the hippocampus of mice lacking the Jnk3 gene. Nature 389:

[70195] 865-870, 1997.

[70196] 11865.

[70197] Yoshida, S.; Fukino, K.; Harada, H.; Nagai, H.; Imoto, I.; In-

azawa,

[70198] J.; Takahashi, H.; Teramoto, A.; Emi, M.: The c-Jun NH2-terminal

[70199] kinase 3 (JNK3) gene: genomic structure, chromosomal assignment, and

[70200] loss of expression in brain tumors. J. Hum. Genet. 46:

182-187,

[70201] 2001.

[70202] 11866.

[70203] Carpenter, N. J.; Filipovich, A.; Blaese, R. M.; Carey, T. L.;

[70204] Berkel, A. I.: Variable immunodeficiency with abnormal condensation

[70205] of the heterochromatin of chromosomes 1, 9, and 16. J. Pediat. 112:

[70206] 757-760, 1988.

[70207] 11867.

[70208] Sillen, A.; Wadelius, C.; Sundvall, M.; Ahlsten, G.; Gus-tavson,

[70209] K. H.: Hyperkalemic periodic paralysis caused by recurring mutation

[70210] in the adult muscle sodium channel alpha-subunit gene. Genet. Counsel. 7:

[70211] 267-275, 1996.

[70212] 11868.

[70213] Tahmoush, A. J.; Schaller, K. L.; Zhang, P.; Hyslop, T.; Heiman-Patterson,

[70214] T.; Caldwell, J. H.: Muscle sodium channel inactivation de-

fect in
[70215] paramyotonia congenita with the thr1313-to-met muta-
tion. Neuromusc.
[70216] Disord. 4: 447-454, 1994.

[70217] 11869.

[70218] Bunger, M. K.; Wilsbacher, L. D.; Moran, S. M.; Clendenin,
C.;
[70219] Radcliffe, L. A.; Hogenesch, J. B.; Simon, M. C.; Takahashi,
J. S.;
[70220] Bradfield, C. A.: Mop3 is an essential component of the
master circadian
[70221] pacemaker in mammals. Cell 103: 1009-1017, 2000.

[70222] 11870.

[70223] Ikeda, M.; Nomura, M.: cDNA cloning and tissue-specific
expression
[70224] of a novel basic helix-loop-helix/PAS protein (BMAL1) and
identification
[70225] of alternatively spliced variants with alternative translation
initiation
[70226] site usage. Biochem. Biophys. Res. Commun. 233:
258-264, 1997.
[70227] 11871.

[70228] Shaw, P. J.; Tononi, G.; Greenspan, R. J.; Robinson, D. F.: Stress
[70229] response genes protect against lethal effects of sleep de-
privation
[70230] in Drosophila. Nature 417: 287-291, 2002.

[70231] 11872.

[70232] Wolting, C. D.; McGlade, C. J.: Cloning and chromosomal localization
[70233] of a new member of the bHLH/PAS transcription factor family. Mammalian
[70234] Genome 9: 463-468, 1998.

[70235] 11873.

[70236] Faucheu, C.; Diu, A.; Chan, A. W. E.; Blanchet, A.-M.; Miossec,
[70237] C.; Herve, F.; Collard-Dutilleul, V.; Gu, Y.; Aldape, R. A.; Lippke,
[70238] J. A.; Rocher, C.; Su, M. S.-S.; Livingston, D. J.; Hercend, T.; Lalanne,
[70239] J.-L.: A novel human protease similar to the interleukin-
1-beta converting
[70240] enzyme induces apoptosis in transfected cells. EMBO J.
14: 1914-1922,

[70241] 1995.

[70242] 11874.

[70243] Kamens, J.; Paskind, M.; Hugunin, M.; Talanian, R. V.;
Allen, H.;
[70244] Banach, D.; Bump, N.; Hackett, M.; Johnston, C. G.; Li, P.;
Mankovich,
[70245] J. A.; Terranova, M.; Ghayur, T.: Identification and charac-
terization
[70246] of ICH-2, a novel member of the interleukin-
1-beta-converting enzyme
[70247] family of cysteine proteases. J. Biol. Chem. 270:
15250-15256, 1995.

[70248] 11875.

[70249] Munday, N. A.; Vaillancourt, J. P.; Ali, A.; Casano, F. J.;
Miller,
[70250] D. K.; Molineaux, S. M.; Yamin, T.-T.; Yu, V. L.; Nicholson,
D. W.
[70251] : Molecular cloning and pro-apoptotic activity of ICE(rel)II
and ICE(rel)III,
[70252] members of the ICE/CED-3 family of cysteine proteases. J.
Biol. Chem. 270:
[70253] 15870-15876, 1995.

[70254] 11876.

[70255] Wang, S.; Miura, M.; Jung, Y.; Zhu, H.; Li, E.; Yuan, J.: Murine

[70256] caspase-11, an ICE-interacting protease, is essential for the activation

[70257] of ICE. Cell 92: 501-509, 1998.

[70258] 11877.

[70259] Liburd, N.; Ghosh, M.; Riazuddin, S.; Naz, S.; Khan, S.; Ahmed,

[70260] Z.; Riazuddin, S.; Liang, Y.; Menon, P. S. N.; Smith, T.; Smith, A.

[70261] C. M.; Chen, K.-S.; Lupski, J. R.; Wilcox, E. R.; Potocki, L.; Friedman,

[70262] T. B.: Novel mutations of MYO15A associated with pro-found deafness

[70263] in consanguineous families and moderately severe hearing loss in a

[70264] patient with Smith-Magenis syndrome. Hum. Genet. 109: 535-541, 2001.

[70265] 11878.

[70266] Carlomagno, F.; Chang-Claude, J.; Dunning, A. M.; Pon-der, B. A.

[70267] J.: Determination of the frequency of the common 675del5 Nijmegen

[70268] breakage syndrome mutation in the German population: no association

[70269] with risk of breast cancer. Genes Chromosomes Cancer 25: 393-395,

[70270] 1999.

[70271] 11879.

[70272] Kleier, S.; Herrmann, M.; Wittwer, B.; Varon, R.; Reis, A.; Horst,

[70273] J.: Clinical presentation and mutation identification in the NBS1

[70274] gene in a boy with Nijmegen breakage syndrome. Clin. Genet. 57:

[70275] 384-387, 2000.

[70276] 11880.

[70277] Lim, D.-S.; Kim, S.-T.; Xu, B.; Maser, R. S.; Lin, J.; Petrini,

[70278] J. H. J.; Kastan, M. B.: ATM phosphorylates p95/nbs1 in an S-phase

[70279] checkpoint pathway. Nature 404: 613-617, 2000.

[70280] 11881.

[70281] Lombard, D. B.; Guarente, L.: Nijmegen breakage syndrome disease
[70282] protein and MRE11 at PML nuclear bodies and meiotic telomeres. Cancer
[70283] Res. 60: 2331-2334, 2000.

[70284] 11882.

[70285] Maser, R. S.; Zinkel, R.; Petrini, J. H. J.: An alternative mode
[70286] of translation permits production of a variant NBS1 protein from the
[70287] common Nijmegen breakage syndrome allele. Nature Genet. 27: 417-421,
[70288] 2001.

[70289] 11883.

[70290] Matsuura, S.; Tauchi, H.; Nakamura, A.; Kondo, N.; Sakamoto, S.;
[70291] Endo, S.; Smeets, D.; Solder, B.; Belohradsky, B. H.; Der Kaloustian,
[70292] V. M.; Oshimura, M.; Isomura, M.; Nakamura, Y.; Komatsu, K.: Positional
[70293] cloning of the gene for Nijmegen breakage syndrome. Nature Genet. 19:

[70294] 179-181, 1998.

[70295] 11884.

[70296] Skelly, R. H.; Korbonits, M.; Grossman, A.; Besser, G. M.; Monson,

[70297] J. P.; Geddes, J. F.; Burrin, J. M.: Expression of the pituitary

[70298] transcription factor Ptx-1, but not that of the trans-activating factor

[70299] Prop-1, is reduced in human corticotroph adenomas and is associated

[70300] with decreased alpha-subunit secretion. J. Clin. Endocr. Metab. 85:

[70301] 2537-2542, 2000.

[70302] 11885.

[70303] Morgan, R. O.; Bell, D. W.; Testa, J. R.; Fernandez, M.-P.: Human

[70304] annexin 31 genetic mapping and origin. Gene 227: 33-38, 1999.

[70305] 11886.

[70306] Morgan, R. O.; Fernandez, M.-P.: Expression profile and structural

[70307] divergence of novel human annexin 31. FEBS Lett. 434:

300-304, 1998.

[70308] 11887.

[70309] Liu, X.-Z.; Xia, X. J.; Xu, L. R.; Pandya, A.; Liang, C. Y.; Blanton,

[70310] S. H.; Brown, S. D. M.; Steel, K. P.; Nance, W. E.: Mutations in

[70311] connexin31 underlie recessive as well as dominant non-syndromic hearing

[70312] loss. Hum. Molec. Genet. 9: 63-67, 2000.

[70313] 11888.

[70314] Lopez-Bigas, N.; Olive, M.; Rabionet, R.; Ben-David, O.; Martinez-Matos,

[70315] J. A.; Bravo, O.; Banchs, I.; Volpini, V.; Gasparini, P.; Avra-ham,

[70316] K. B.; Ferrer, I.; Arbones, M. L.; Estivill, X.: Connexin 31 (GJB3)

[70317] is expressed in the peripheral and auditory nerves and causes neuropathy

[70318] and hearing impairment. Hum. Molec. Genet. 10: 947-952, 2001.

[70319] 11889.

[70320] Nadler, M. J. S.; Hermosura, M. C.; Inabe, K.; Perraud,

A.-L.;

[70321] Zhu, Q.; Stokes, A. J.; Kurosaki, T.; Kinet, J.-P.; Penner, R.; Scharenberg,

[70322] A. M.; Fleig, A.: LTRPC7 is a Mg-ATP-regulated divalent cation channel

[70323] required for cell viability. Nature 411: 590-595, 2001. Note: Erratum Nature 412:

[70324] 660 only, 2001.

[70325] 11890.

[70326] Runnels, L. W.; Yue, L.; Clapham, D. E.: TRP-PLIK, a bi-functional

[70327] protein with kinase and ion channel activities. Science 291: 1043-1047,

[70328] 2001.

[70329] 11891.

[70330] Schlingmann, K. P.; Weber, S.; Peters, M.; Nejsum, L. N.; Vitzthum,

[70331] H.; Klingel, K.; Kratz, M.; Haddad, E.; Ristoff, E.; Dinour, D.; Syrrou,

[70332] M.; Nielsen, S.; Sassen, M.; Waldegger, S.; Seyberth, H. W.; Konrad,

[70333] M.: Hypomagnesemia with secondary hypocalcemia is caused by mutations

[70334] in TRPM6, a new member of the TRPM gene family. Nature Genet. 31:

[70335] 166-170, 2002.

[70336] 11892.

[70337] Walder, R. Y.; Shalev, H.; Brennan, T. M. H.; Carmi, R.; Elbedour,

[70338] K.; Scott, D. A.; Hanauer, A.; Mark, A. L.; Patil, S.; Stone, E. M.;

[70339] Sheffield, V. C.: Familial hypomagnesemia maps to chromosome 9q,

[70340] not to the X chromosome: genetic linkage mapping and analysis of a
[70341] balanced translocation breakpoint. Hum. Molec. Genet. 6: 1491-1497,
[70342] 1997.

[70343] 11893.

[70344] Brohmann, H.; Pinnecke, S.; Hoyer-Fender, S.: Identification and
[70345] characterization of new cDNAs encoding outer dense fiber proteins
[70346] of rat sperm. J. Biol. Chem. 272: 10327-10332, 1997.

[70347] 11894.

[70348] Scott, A. F.: Personal Communication. Baltimore, Md. 9/10/1997.
[70349] 11895.

[70350] Shao, X.; Murthy, S.; Demetrick, D. J.; van der Hoorn, F. A.:
[70351] Human outer dense fiber gene, ODF2, localizes to chromosome 9q34. Cytogenet.
[70352] Cell Genet. 83: 221-223, 1998.

[70353] 11896.

[70354] Porteous, S.; Torban, E.; Cho, N.-P.; Cunliffe, H.; Chua, L.;

[70355] McNoe, L.; Ward, T.; Souza, C.; Gus, P.; Giugliani, R.; Sato, T.;

[70356] Yun, K.; Favor, J.; Sicotte, M.; Goodyer, P.; Eccles, M.: Pri-

[70357] mary renal hypoplasia in humans and mice with PAX2 muta-

[70358] tions: evidence of increased apoptosis in fetal kidneys of Pax2(1Neu) +/-

[70359] mutant mice. Hum. Molec. Genet. 9: 1-11, 2000.

[70360] 11897.

[70361] Bertazzoni, U.; Scovassi, A. I.; Brun, G. M.: Chick-embryo DNA

[70362] polymerase gamma: identity of gamma-polymerases puri-

[70363] fied from nuclei and mitochondria. Europ. J. Biochem. 81: 237-248, 1977.

[70364] 11898.

[70365] Kamath-Loeb, A. S.; Johansson, E.; Burgers, P. M. J.; Loeb, L.

[70366] A.: Functional interaction between the Werner syndrome protein and

[70367] DNA polymerase delta. Proc. Nat. Acad. Sci. 97:
4603-4608, 2000.

[70368] 11899.

[70369] Lestienne, P.: Evidence for a direct role of the DNA poly-
merase

[70370] gamma in the replication of the human mitochondrial DNA
in vitro. Biochem.

[70371] Biophys. Res. Commun. 146: 1146-1153, 1987.

[70372] 11900.

[70373] Lecrenier, N.; van der Bruggen, P.; Foury, F.: Mitochondrial
DNA

[70374] polymerases from yeast to man: a new family of poly-
merases. Gene 185:

[70375] 147-152, 1997.

[70376] 11901.

[70377] Ropp, P. A.; Copeland, W. C.: Cloning and characterization
of

[70378] the human mitochondrial DNA polymerase, DNA poly-
merase gamma. Genomics 36:

[70379] 449-458, 1996.

[70380] 11902.

[70381] Rovio, A.; Tiranti, V.; Bednarz, A. L.; Suomalainen, A.; Spelbrink,
[70382] J. N.; Lecrenier, N.; Melberg, A.; Zeviani, M.; Poulton, J.; Foury,
[70383] F.; Jacobs, H. T.: Analysis of the trinucleotide CAG repeat from
[70384] the human mitochondrial DNA polymerase gene in healthy and diseased
[70385] individuals. Europ. J. Hum. Genet. 7: 140-146, 1999.

[70386] 11903.

[70387] Rovio, A. T.; Marchington, D. R.; Donat, S.; Schuppe, H.-S.; Abel,
[70388] J.; Fritsche, E.; Elliott, D. J.; Laippala, P.; Ahola, A. L.; Mc-Nay,
[70389] D.; Harrison, R. F.; Hughes, B.; and 13 others: Mutations at the
[70390] mitochondrial DNA polymerase (POLG) locus associated with male infertility. Nature
[70391] Genet. 29: 261-262, 2001.

[70392] 11904.

[70393] Lind, B.; van Solinge, W. W.; Schwartz, M.; Thorsen, S.: Splice

[70394] site mutation in the human protein C gene associated with venous thrombosis:
[70395] demonstration of exon skipping by ectopic transcript analysis. Blood 82:
[70396] 2423-2432, 1993.

[70397] 11905.

[70398] Long, G. L.; Marshall, A.; Gardner, J. C.; Naylor, S. L.: Genes
[70399] for human vitamin K-dependent plasma proteins C and S are located
[70400] on chromosomes 2 and 3, respectively. Somat. Cell Molec. Genet. 14:
[70401] 93-98, 1988.

[70402] 11906.

[70403] Bamshad, M.; Lin, R. C.; Law, D. J.; Watkins, W. S.; Krakowiak,
[70404] P. A.; Moore, M. E.; Franceschini, P.; Lala, R.; Holmes, L. B.; Gebuhr,
[70405] T. C.; Bruneau, B. G.; Schinzel, A.; Seidman, J. G.; Seidman, C. E.;
[70406] Jorde, L. B.: Mutations in human TBX3 alter limb, apocrine and genital

[70407] development in ulnar-mammary syndrome. Nature Genet. 16: 311-315,

[70408] 1997.

[70409] 11907.

[70410] Myohanen, S.; Kauppinen, L.; Wahlfors, J.; Alhonen, L.; Janne,

[70411] J.: Human spermidine synthase gene: structure and chromosomal localization. DNA

[70412] Cell Biol. 10: 467-474, 1991.

[70413] 11908.

[70414] Wahlfors, J.; Alhonen, L.; Kauppinen, L.; Hyvonen, T.; Janne, J.;

[70415] Eloranta, T. O.: Human spermidine synthase: cloning and primary structure. DNA

[70416] Cell Biol. 9: 103-110, 1990.

[70417] 11909.

[70418] Winqvist, R.; Alanen, L.; Grzeschik, K.-H.; Janne, J.; Eloranta,

[70419] T.: Mapping of DNA sequences for spermidine synthase to human chromosome

[70420] 1p36-p22 and chromosome 3p14-q21. (Abstract) Cyto-

genet. Cell Genet. 58:
[70421] 1865 only, 1991.

[70422] 11910.

[70423] Agre, P.: Personal Communication. Baltimore, Md. 6/3/1989.

[70424] 11911.

[70425] Agre, P.; Casella, J. F.; Zinkham, W. H.; McMillan, C.; Bennett,
[70426] V.: Partial deficiency of erythrocyte spectrin in hereditary spherocytosis. Nature 314:
[70427] 380-383, 1985.

[70428] 11912.

[70429] Anderson, R.; Huestis, R. R.; Motulsky, A. G.: Hereditary spherocytosis
[70430] in the deer mouse: its similarity to the human disease. Blood 15:
[70431] 491-504, 1960.

[70432] 11913.

[70433] Barry, M.; Scheuer, P. J.; Sherlock, S.; Ross, C. F.; Williams,
[70434] R.: Hereditary spherocytosis with secondary haemochro-

matosis. Lancet II:
[70435] 481-485, 1968.

[70436] 11914.

[70437] Bass, E. B.; Smith, S. W., Jr.; Stevenson, R. E.; Rosse, W. F.

[70438] : Further evidence for location of the spherocytosis gene on chromosome 11915.
[70439] Ann. Intern. Med. 99: 192-193, 1983.

[70440]

[70441] 8. Birkenmeier, C. S.; McFarland-Starr, E. C.; Barker, J. E.: Chromosomal
[70442] location of three spectrin genes: relationship to the inher-ited hemolytic
[70443] anemias of mouse and man. Proc. Nat. Acad. Sci. 85: 8121-8125, 1988.
[70444] 11916.

[70445] Bodine, D. M., IV; Birkenmeier, C. S.; Barker, J. E.: Spectrin

[70446] deficient inherited hemolytic anemias in the mouse: char-acterization
[70447] by spectrin synthesis and mRNA activity in reticulocytes. Cell 37:
[70448] 721-729, 1984.

[70449] 11917.

[70450] Chilcote, R. R.; Le Beau, M. M.; Dampier, C.; Pergament, E.; Verlinsky,

[70451] Y.; Mohandas, N.; Frischer, H.; Rowley, J. D.: Association of red

[70452] cell spherocytosis with deletion of the short arm of chromosome 8. Blood 69:

[70453] 156-159, 1987.

[70454] 11918.

[70455] Coetzer, T. L.; Lawler, J.; Liu, S.-C.; Prchal, J. T.; Gualtieri,

[70456] R. J.; Brain, M. C.; Dacie, J. V.; Palek, J.: Partial ankyrin and

[70457] spectrin deficiency in severe, atypical hereditary spherocytosis. New

[70458] Eng. J. Med. 318: 230-234, 1988.

[70459] 11919.

[70460] Cohen, H.; Walker, H.; Delhanty, J. D. A.; Lucas, S. B.; Huehns,

[70461] E. R.: Congenital spherocytosis, B19 parvovirus infection and inherited

[70462] interstitial deletion of the short arm of chromosome 8. Brit. J.

[70463] Haemat. 78: 251-257, 1991.

[70464] 11920.

[70465] Costa, F. F.; Agre, P.; Watkins, P. C.; Winkelmann, J. C.; Tang,

[70466] T. K.; John, K. M.; Lux, S. E.; Forget, B. G.: Linkage of dominant

[70467] hereditary spherocytosis to the gene for the erythrocyte membrane-skeleton

[70468] protein ankyrin. New Eng. J. Med. 323: 1046-1050, 1990.

[70469] 11921.

[70470] de Jongh, B. M.; Blacklock, H. A.; Reekers, P.; Volkers, W. S.;

[70471] Schreuder, G. M. T.; Meera Khan, P.; Bernini, L. F.; Nijen-huis, L.

[70472] E.; van Loghem, E.; van Rood, J. J.: No evidence of linkage between

[70473] hereditary spherocytosis (SPH) and genetic markers in-cluding HLA and

[70474] IGHG.(Abstract) Cytogenet. Cell Genet. 32: 263-264, 1982.

[70475] 11922.

[70476] Duru, F.; Gurgey, A.; Ozturk, G.; Yorukan, S.; Altay, C.: Homozygosity
[70477] for dominant form of hereditary spherocytosis. Brit. J. Haemat. 82:
[70478] 596-600, 1992.

[70479] 11923.

[70480] Croce, C. M.; Isobe, M.; Palumbo, A.; Puck, J.; Ming, J.; Tweardy,
[70481] D.; Erikson, J.; Davis, M.; Rovera, G.: Gene for alpha-chain of human
[70482] T-cell receptor: location on chromosome 14 region in-volved in T-cell
[70483] neoplasms. Science 227: 1044-1047, 1985.

[70484] 11924.

[70485] Funke, B.; Epstein, J. A.; Kochilas, L. K.; Lu, M. M.; Pandita,

[70486] R. K.; Liao, J.; Bauerndistel, R.; Schuler, T.; Schorle, H.; Brown,
[70487] M. C.; Adams, J.; Morrow, B. E.: Mice overexpressing genes from the
[70488] 22q11 region deleted in velo-cardio-facial syndrome/Di-George syndrome

[70489] have middle and inner ear defects. Hum. Molec. Genet. 10: 2549-2556,

[70490] 2001.

[70491] 11925.

[70492] Guris, D. L.; Fantes, J.; Tara, D.; Druker, B. J.; Imamoto, A.

[70493] : Mice lacking the homologue of the human 22q11.2 gene CRKL phenocopy

[70494] neurocristopathies of DiGeorge syndrome. Nature Genet. 27: 293-298,

[70495] 2001.

[70496] 11926.

[70497] Halford, S.; Wilson, D. I.; Daw, S. C. M.; Roberts, C.; Wadey,

[70498] R.; Kamath, S.; Wickremasinghe, A.; Burn, J.; Goodship, J.; Mattei,

[70499] M.-G.; Moorman, A. F. M.; Scambler, P. J.: Isolation of a gene expressed

[70500] during early embryogenesis from the region of 22q11 commonly deleted

[70501] in DiGeorge syndrome.. Hum. Molec. Genet. 2: 1577-1582, 1993.

[70502] 11927.

[70503] Jerome, L. A.; Papaioannou, V. E.: DiGeorge syndrome phenotype

[70504] in mice mutant for the T-box gene, Tbx1. Nature Genet. 27: 286-291,

[70505] 2001.

[70506] 11928.

[70507] Dyson, E.; Sucov, H. M.; Kubalak, S. W.; Schmid-Schon-

[70508] bein, G. W.; DeLano, F. A.; Evans, R. M.; Ross, J., Jr.; Chien, K. R.: Atrial-like

[70509] phenotype is associated with embryonic ventricular failure in retinoid

[70510] X receptor alpha -/- mice. Proc. Nat. Acad. Sci. 92: 7386-7390,

[70511] 1995.

[70512] 11929.

[70513] Gampe, R. T., Jr.; Montana, V. G.; Lambert, M. H.; Miller, A. B.;

[70514] Bledsoe, R. K.; Milburn, M. V.; Kliewer, S. A.; Willson, T. M.; Xu,

[70515] H. E.: Asymmetry in the PPAR-gamma/RXR-alpha crystal structure reveals
[70516] the molecular basis of heterodimerization among nuclear receptors. Molec.
[70517] Cell 5: 545-555, 2000.

[70518] 11930.

[70519] Gruber, P. J.; Kubalak, S. W.; Pexieder, T.; Sucov, H. M.; Evans,
[70520] R. M.; Chien, K. R.: RXR-alpha deficiency confers genetic susceptibility
[70521] for aortic sac, conotruncal, atrioventricular cushion, and ventricular
[70522] muscle defects in mice. J. Clin. Invest. 98: 1332-1343, 1996.
[70523] 11931.

[70524] Heyman, R. A.; Mangelsdorf, D. J.; Dyck, J. A.; Stein, R. B.; Eichele,
[70525] G.; Evans, R. M.; Thaller, C.: 9-cis retinoic acid is a high affinity
[70526] ligand for the retinoid X receptor. Cell 68: 397-406, 1992.

[70527] 11932.

[70528] Hoopes, C. W.; Taketo, M.; Ozato, K.; Liu, Q.; Howard, T. A.; Linney,

[70529] E.; Seldin, M. F.: Mapping the mouse Rxr loci encoding nuclear retinoid

[70530] X receptors Rxr-alpha, Rxr-beta, and Rxr-gamma. Genomics 14: 611-617,

[70531] 1992.

[70532] 11933.

[70533] Jones, K. A.; Fitzgibbon, J.; Woodward, K. J.; Goudie, D.; Ferguson-Smith,

[70534] M. A.; Povey, S.; Wolfe, J.; Solomon, E.: Localization of the retinoid

[70535] X receptor alpha gene (RXRA) to chromosome 9q34. Ann. Hum. Genet. 57:

[70536] 195-201, 1993.

[70537] 11934.

[70538] Li, M.; Indra, A. K.; Warot, X.; Brocard, J.; Messaddeq, N.; Kato,

[70539] S.; Metzger, D.; Chambon, P.: Skin abnormalities generated by temporally

[70540] controlled RXR-alpha mutations in mouse epidermis. Nature 407: 633-636,

[70541] 2000.

[70542] 11935.

[70543] Lu, T. T.; Makishima, M.; Repa, J. J.; Schoonjans, K.; Kerr, T.

[70544] A.; Auwerx, J.; Mangelsdorf, D. J.: Molecular basis for feedback

[70545] regulation of bile acid synthesis by nuclear receptors. Molec. Cell 6:

[70546] 507-515, 2000.

[70547] 11936.

[70548] Mangelsdorf, D. J.; Ong, E. S.; Dyck, J. A.; Evans, R. M.: Nuclear

[70549] receptor that identifies a novel retinoic acid response pathway. Nature 345:

[70550] 224-229, 1990.

[70551] 11937.

[70552] Mangelsdorf, D. J.; Umesono, K.; Kliewer, S. A.; Borgmeyer, U.;

[70553] Ong, E. S.; Evans, R. M.: A direct repeat in the cellular retinol-binding

[70554] protein type II gene confers differential regulation by RXR

and RAR. Cell 66:
[70555] 555-561, 1991.

[70556] 11938.

[70557] Repa, J. J.; Turley, S. D.; Lobaccaro, J.-M. A.; Medina, J.; Li,

[70558] L.; Lustig, K.; Shan, B.; Heyman, R. A.; Dletschy, J. M.;
Mangelsdorf,
[70559] D. J.: Regulation of absorption and ABC1-mediated efflux
of cholesterol
[70560] by RXR heterodimers. Science 289: 1524-1529, 2000.

[70561] 11939.

[70562] Tontonoz, P.; Hu, E.; Spiegelman, B. M.: Stimulation of
adipogenesis
[70563] in fibroblasts by PPAR-gamma-2, a lipid-activated tran-
scription factor. Cell 79:
[70564] 1147-1156, 1994.

[70565] 11940.

[70566] Willy, P. J.; Umesono, K.; Ong, E. S.; Evans, R. M.; Heyman,
R.
[70567] A.; Mangelsdorf, D. J.: LXR, a nuclear receptor that defines
a distinct
[70568] retinoid response pathway. Genes Dev. 9: 1033-1045,

1995.

[70569] 11941.

[70570] Wan, Y.-J. Y.; An, D.; Cai, Y.; Repa, J. J.; Chen, T. H.-P.;
Flores,

[70571] M.; Postic, C.; Magnuson, M. A.; Chen, J.; Chien, K. R.;
French, S.;

[70572] Mangelsdorf, D. J.; Sucov, H. M.: Hepatocyte-specific mu-
tation establishes

[70573] retinoid X receptor alpha as a heterodimeric integrator of
multiple

[70574] physiological processes in the liver. Molec. Cell. Biol. 20:
4436-4444,

[70575] 2000.

[70576] 11942.

[70577] Zhou, M. D.; Sucov, H. M.; Evans, R. M.; Chien, K. R.:
Retinoid-dependent

[70578] pathways suppress myocardial cell hypertrophy. Proc. Nat.
Acad. Sci. 92:

[70579] 7391-7395, 1995.

[70580] 11943.

[70581] Graycar, J. L.; Miller, D. A.; Arrick, B. A.; Lyons, R. M.;

Moses,

[70582] H. L.; Derynck, R.: Human transforming growth factor-beta-3: recombinant

[70583] expression, purification, and biological activities in comparison

[70584] with transforming growth factors-beta-1 and beta-2. Molec. Endocr. 3:

[70585] 1977-1986, 1989.

[70586] 11944.

[70587] Kaartinen, V.; Voncken, J. W.; Shuler, C.; Warburton, D.; Bu, D.;

[70588] Heisterkamp, N.; Groffen, J.: Abnormal lung development and cleft

[70589] palate in mice lacking TGF-beta-3 indicates defects of epithelial-mesenchymal

[70590] interaction. Nature Genet. 11: 415-421, 1995.

[70591] 11945.

[70592] Lee, B.-S.; Nowak, R. A.: Human leiomyoma smooth muscle cells

[70593] show increased expression of transforming growth factor-beta-3 (TGF-beta-3)

[70594] and altered responses to the antiproliferative effects of

TGF-beta. J.
[70595] Clin. Endocr. Metab. 86: 913-920, 2001.

[70596] 11946.

[70597] Moren, A.; Ichijo, H.; Miyazono, K.: Molecular cloning and characterization
[70598] of the human and porcine transforming growth factor-beta type III
[70599] receptors. Biochem. Biophys. Res. Commun. 189: 356-362, 1992.
[70600] 11947.

[70601] Proetzel, G.; Pawlowski, S. A.; Wiles, M. V.; Yin, M.; Boivin,

[70602] G. P.; Howles, P. N.; Ding, J.; Ferguson, M. W. J.; Doetschman, T.
[70603] : Transforming growth factor-beta-3 is required for sec-ondary palate
[70604] fusion. Nature Genet. 11: 409-414, 1995.

[70605] 11948.

[70606] ten Dijke, P.; Geurts van Kessel, A. H. M.; Foulkes, J. G.; Le

[70607] Beau, M. M.: Transforming growth factor type beta-3 maps to human
[70608] chromosome 14, region q23-q24. Oncogene 3: 721-724,

1988.

[70609] 11949.

[70610] ten Dijke, P.; Hansen, P.; Iwata, K. K.; Pieler, C.; Foulkes, J.

[70611] G.: Identification of another member of the transforming growth factor

[70612] type beta gene family. Proc. Nat. Acad. Sci. 85: 4715-4719, 1988.

[70613] 11950.

[70614] Luerssen, H.; Mattei, M.-G.; Schroter, M.; Grzeschik, K.-H.; Adham,

[70615] I. M.; Engel, W.: Nucleotide sequence of the gene for human transition

[70616] protein 1 and its chromosomal localization on chromosome 2. Genomics 8:

[70617] 324-330, 1990.

[70618] 11951.

[70619] Yelick, P. C.; Kozak, C.; Kwon, Y. K.; Seldin, M. F.; Hecht, N.

[70620] B.: The mouse transition protein 1 gene contains a B1 repetitive

[70621] element and is located on chromosome 1. Genomics 11:

687-694, 1991.

[70622] 11952.

[70623] Yu, Y. E.; Zhang, Y.; Unni, E.; Shirley, C. R.; Deng, J. M.; Russell,

[70624] L. D.; Weil, M. M.; Behringer, R. R.; Meistrich, M. L.: Abnor-mal

[70625] spermatogenesis and reduced fertility in transition nuclear protein

[70626] 1-deficient mice. Proc. Nat. Acad. Sci. 97: 4683-4688, 2000.

[70627] 11953.

[70628] Engel, W.; Keime, S.; Kremling, H.; Hameister, H.; Schluter, G.

[70629] : The genes for protamine 1 and 2 (PRM1 and PRM2) and transition protein

[70630] 2 (TNP2) are closely linked in the mammalian genome. Cytogenet. Cell

[70631] Genet. 61: 158-159, 1992.

[70632] 11954.

[70633] Nelson, J. E.; Krawetz, S. A.: Linkage of human spermatid-specific

[70634] basic nuclear protein genes: definition and evolution of

the P1-to-P2-to-TP2
[70635] locus. J. Biol. Chem. 268: 2932-2936, 1993.

[70636] 11955.

[70637] Schluter, G.; Kremling, H.; Engel, W.: The gene for human transition
[70638] protein 2: nucleotide sequence, assignment to the pro-tamine gene cluster,
[70639] and evidence for its low expression. Genomics 14: 377-383, 1992.

[70640] 11956.

[70641] Kim, P. K. M.; Dutra, A. S.; Chandrasekharappa, S. C.; Puck, J.
[70642] M.: Genomic structure and mapping of human FADD, an intracellular
[70643] mediator of lymphocyte apoptosis. J. Immun. 157: 5461-5466, 1996.

[70644] 11957.

[70645] Lavedan, C.; Leroy, E.; Torres, R.; Dehejia, A.; Dutra, A.; Buchholtz,
[70646] S.; Nussbaum, R. L.; Polymeropoulos, M. H.: Genomic or-ganization
[70647] and expression of the human beta-synuclein gene (SNCB)

Genomics 54:
[70648] 173-175, 1998.

[70649] 11958.

[70650] Hymowitz, S. G.; Christinger, H. W.; Fuh, G.; Ultsch, M.; O'Connell,
[70651] M.; Kelley, R. F.; Ashkenazi, A.; de Vos, A. M.: Triggering cell
[70652] death: the crystal structure of Apo2L/TRAIL in a complex with death
[70653] receptor 5. Molec. Cell 4: 563-571, 1999.

[70654] 11959.

[70655] Nitsch, R.; Bechmann, I.; Deisz, R. A.; Haas, D.; Lehmann, T. N.;
[70656] Wendling, U.; Zipp, F.: Human brain-cell death induced by tumour-necrosis-factor-related
[70657] apoptosis-inducing ligand (TRAIL). Lancet 356: 827-828, 2000.
[70658] 11960.

[70659] Pitti, R. M.; Marsters, S. A.; Ruppert, S.; Donahue, C. J.; Moore,
[70660] A.; Ashkenazi, A.: Induction of apoptosis by Apo-2 ligand,

a new

[70661] member of the tumor necrosis factor cytokine family. J. Biol. Chem. 271:

[70662] 12687-12690, 1996.

[70663] 11961.

[70664] Wiley, S. R.; Schooley, K.; Smolak, P. J.; Din, W. S.; Huang, C.-P.;

[70665] Nicholl, J. K.; Sutherland, G. R.; Smith, T. D.; Rauch, C.; Smith,

[70666] C. A.; Goodwin, R. G.: Identification and characterization of a new

[70667] member of the TNF family that induces apoptosis. Immunity 3: 673-682,

[70668] 1995.

[70669] 11962.

[70670] Lyu, M. S.; Park, D. J.; Rhee, S. G.; Kozak, C. A.: Genetic mapping

[70671] of the human and mouse phospholipase C genes. Mammalian Genome 7:

[70672] 501-504, 1996.

[70673] 11963.

[70674] Shimohama, S.; Kamiya, S.; Fujii, M.; Ogawa, T.; Kanamori, M.;

[70675] Kawamata, J.; Imura, T.; Taniguchi, T.; Yagisawa, H.: Mutation in

[70676] the pleckstrin homology domain of the human phospholipase C-delta-1

[70677] gene is associated with loss of function. Biochem. Biophys. Res.

[70678] Commun. 245: 722-728, 1998.

[70679] 11964.

[70680] Iwabuchi, K.; Bartel, P. L.; Li, B.; Marraccino, R.; Fields, S.

[70681] : Two cellular proteins that bind to wild-type but not mutant p53. Proc.

[70682] Nat. Acad. Sci. 91: 6098-6102, 1994.

[70683] 11965.

[70684] Iwabuchi, K.; Li, B.; Massa, H. F.; Trask, B. J.; Date, T.; Fields,

[70685] S.: Stimulation of p53-mediated transcriptional activation by the

[70686] p53-binding proteins, 53BP1 and 53BP2. J. Biol. Chem. 273: 26061-26068,

[70687] 1998.

[70688] 11966.

[70689] Naumovski, L.; Cleary, M. L.: The p53-binding protein 53BP2 also

[70690] interacts with Bc12 and impedes cell cycle progression at G2/M. Molec.

[70691] Cell Biol. 16: 3884-3892, 1996.

[70692] 11967.

[70693] Samuels-Lev, Y.; O'Connor, D. J.; Bergamaschi, D.; Tri-giante, G.;

[70694] Hsieh, J.-K.; Zhong, S.; Campargue, I.; Naumovski, L.; Crook, T.;

[70695] Lu, X.: ASPP proteins specifically stimulate the apoptotic function

[70696] of p53. Molec. Cell 8: 781-794, 2001.

[70697] 11968.

[70698] Yang, J.-P.; Ono, T.; Sonta, S.; Kawabe, T.; Okamoto, T.: Assignment

[70699] of p53 binding protein (TP53BP2) to human chromosome band 1q42.1 by

[70700] in situ hybridization. Cytogenet. Cell Genet. 78: 61-62,

1997.

[70701] 11969.

[70702] Lamartine, J.; Seri, M.; Cinti, R.; Heitzmann, F.; Creaven, M.;

[70703] Radomski, N.; Jost, E.; Lenoir, G. M.; Romeo, G.; Sylla, B. S.: Molecular

[70704] cloning and mapping of a human cDNA (PA2G4) that encodes a protein

[70705] highly homologous to the mouse cell cycle protein p38-2G4. Cytogenet.

[70706] Cell Genet. 78: 31-35, 1997.

[70707] 11970.

[70708] Radomski, N.; Jost, E.: Molecular cloning of a murine cDNA encoding

[70709] a novel protein, p38-2G4, which varies with the cell cycle. Exp.

[70710] Cell Res. 220: 434-445, 1995.

[70711] 11971.

[70712] Crawford, M. J.; Lanctot, C.; Tremblay, J. J.; Jenkins, N.; Gilbert,

[70713] D.; Copeland, N.; Beatty, B.; Drouin, J.: Human and murine

PTX1/Ptx1

[70714] gene maps to the region for Treacher Collins syndrome. Mammalian

[70715] Genome 8: 841-845, 1997.

[70716] 11972.

[70717] Lamonerie, T.; Tremblay, J. J.; Lanctot, C.; Thierrien, M.; Gauthier,

[70718] Y.; Drouin, J.: Ptx1, a bicoid-related homeo box transcrip-tion factor

[70719] involved in transcription of the pro-opiomelanocortin gene. Genes

[70720] Dev. 10: 1284-1295, 1996.

[70721] 11973.

[70722] Shang, J.; Li, X.; Ring, H. Z.; Clayton, D. A.; Francke, U.: Backfoot,

[70723] a novel homeobox gene, maps to human chromosome 5 (BFT) and mouse

[70724] chromosome 13 (Bft). Genomics 40: 108-113, 1997.

[70725] 11974.

[70726] Szeto, D. P.; Rodriguez-Esteban, C.; Ryan, A. K.; O'Connell, S.

[70727] M.; Liu, F.; Kioussi, C.; Gleiberman, A. S.; Izpisua-Belmonte, J.

[70728] C.; Rosenfeld, M. G.: Role of the Bicoid-related homeodomain factor

[70729] Pitx1 in specifying hindlimb morphogenesis and pituitary development. Genes

[70730] Dev. 13: 484-494, 1999.

[70731] 11975.

[70732] Semina, E. V.; Altherr, M. R.; Murray, J. C.: Cloning and chromosomal

[70733] localization of two novel human genes encoding LIM-domain binding

[70734] factors CLIM1 and CLIM2/LDB1/NL1. Mammalian Genome 9: 921-924, 1998.

[70735] 11976.

[70736] Yamashita, T.; Agulnick, A. D.; Copeland, N. G.; Gilbert, D. J.;

[70737] Jenkins, N. A.; Westphal, H.: Genomic structure and chromosomal localization

[70738] of the mouse LIM domain-binding protein 1 gene, Ldb1. Genomics 48:

[70739] 87-92, 1998.

[70740] 11977.

[70741] Foley, B. T.; Moehring, J. M.; Moehring, T. J.: Mutations in the

[70742] elongation factor 2 gene which confer resistance to diphtheria toxin

[70743] and Pseudomonas exotoxin A: genetic and biochemical analyses. J.

[70744] Biol. Chem. 270: 23218-23225, 1995.

[70745] 11978.

[70746] Mattheakis, L. C.; Sor, F.; Collier, R. J.: Diphthamide synthesis

[70747] in Saccharomyces cerevisiae: structure of the DPH2 gene. Gene 132:

[70748] 149-154, 1993.

[70749] 11979.

[70750] Schultz, D. C.; Balasara, B. R.; Testa, J. R.; Godwin, A. K.:

[70751] Cloning and localization of a human diphthamide biosynthesis-like

[70752] protein-2 gene, DPH2L2. Genomics 52: 186-191, 1998.

[70753] 11980.

[70754] Abe, R.; Sakashita, E.; Yamamoto, K.; Sakamoto, H.: Two different

[70755] RNA binding activities for the AU-rich element and the poly(A) sequence

[70756] of the mouse neuronal protein mHuC. Nucleic Acids Res. 24: 4895-4901,

[70757] 1996.

[70758] 11981.

[70759] Winter, H.; Rogers, M. A.; Gebhardt, M.; Wollina, U.; Boxall, L.;

[70760] Chitayat, D.; Babul-Hirji, R.; Stevens, H. P.; Zlotogorski, A.; Schweizer,

[70761] J.: A new mutation in the type II hair cortex keratin hHb1 involved

[70762] in the inherited hair disorder monilethrix. Hum. Genet. 101: 165-169,

[70763] 1997.

[70764] 11982.

[70765] Buckanovich, R. J.; Posner, J. B.; Darnell, R. B.: Nova, the paraneoplastic

[70766] Ri antigen, is homologous to an RNA-binding protein and is specifically

[70767] expressed in the developing motor system. Neuron 11: 657-672, 1993.

[70768] 11983.

[70769] Buckanovich, R. J.; Yang, Y. Y.; Darnell, R. B.: The on-coneural

[70770] antigen Nova-1 is a neuron-specific RNA-binding protein, the activity

[70771] of which is inhibited by paraneoplastic antibodies. J. Neurosci. 16:

[70772] 1114-1122, 1996.

[70773] 11984.

[70774] Prestigiacomo, C. J.; Balmaceda, C.; Dalmau, J.: Anti-Ri-associated

[70775] paraneoplastic opsoclonus-ataxia syndrome in a man with transitional

[70776] cell carcinoma: a case report. Cancer 91: 1423-1428, 2001.

[70777] 11985.

[70778] Muzio, M.; Ni, J.; Feng, P.; Dixit, V. M.: IRAK (Pelle) family

[70779] member IRAK-2 and MyD88 as proximal mediators of IL-1 signaling. Science 278:

[70780] 1612-1615, 1997.

[70781] 11986.

[70782] Takeuchi, O.; Hoshino, K.; Akira, S.: Cutting edge:
TLR2-deficient

[70783] and MyD88-deficient mice are highly susceptible to
Staphylococcus

[70784] aureus infection. J. Immun. 165: 5392-5396, 2000.

[70785] 11987.

[70786] Eggertsen, G.; Olin, M.; Andersson, U.; Ishida, H.; Kubota,
S.;

[70787] Hellman, U.; Okuda, K.-I.; Bjorkhem, I.: Molecular cloning
and expression

[70788] of rabbit sterol 12-alpha-hydroxylase. J. Biol. Chem. 271:
32269-32275,

[70789] 1996.

[70790] 11988.

[70791] Gafvels, M.; Olin, M.; Chowdhary, B. P.; Raudsepp, T.; An-
dersson,

[70792] U.; Persson, B.; Jansson, M.; Bjorkhem, I.; Eggertsen, G.:
Structure

[70793] and chromosomal assignment of the sterol

12-alpha-hydroxylase gene

[70794] (CYP8B1) in human and mouse: eukaryotic cytochrome P-450 gene devoid

[70795] of introns. Genomics 56: 184-196, 1999.

[70796] 11989.

[70797] Zhang, M.; Chiang, J. Y. L.: Transcriptional regulation of the

[70798] human sterol 12-alpha-hydroxylase gene (CYP8B1): roles of hepatocyte

[70799] nuclear factor 4-alpha in mediating bile acid repression. J. Biol.

[70800] Chem. 276: 41690-41699, 2001.

[70801] 11990.

[70802] Brinkman, A.; van der Flier, S.; Kok, E. M.; Dorssers, L. C. J.

[70803] : BCAR1, a human homologue of the adapter protein p130Cas, and antiestrogen

[70804] resistance in breast cancer cells. J. Nat. Cancer Inst. 92: 112-120,

[70805] 2000.

[70806] 11991.

[70807] Brugge, J. S.: Casting light on focal adhesions. Nature

Genet. 19:
[70808] 309-311, 1998.

[70809] 11992.

[70810] Dorssers, L. C. J.; van Agthoven, T.; Dekker, A.; van Agth-
oven,
[70811] T. L. A.; Kok, E. M.: Induction of antiestrogen resistance in
human
[70812] breast cancer cells by random insertional mutagenesis us-
ing defective
[70813] retroviruses: identification of bcar-1, a common integra-
tion site. Molec.
[70814] Endocr. 7: 870-878, 1993.

[70815] 11993.

[70816] Honda, H.; Oda, H.; Nakamoto, T.; Honda, Z.; Sakai, R.;
Suzuki,
[70817] T.; Saito, T.; Nakamura, K.; Nakao, K.; Ishikawa, T.; Kat-
suki, M.;
[70818] Yazaki, Y.; Hirai, H.: Cardiovascular anomaly, impaired
actin bundling
[70819] and resistance to Src-induced transformation in mice
lacking p130-Cas. Nature
[70820] Genet. 19: 361-365, 1998.

[70821] 11994.

[70822] Kanner, S. B.; Reynolds, A. B.; Wang, H. C.; Vines, R. R.; Parsons,

[70823] J. T.: The SH2 and SH3 domains of pp60src direct stable association

[70824] with tyrosine phosphorylated proteins p130 and p110. EMBO J. 10:

[70825] 1689-1698, 1991.

[70826] 11995.

[70827] Kuo, C. T.; Morrisey, E. E.; Anandappa, R.; Sigrist, K.; Lu, M.

[70828] M.; Parmacek, M. S.; Soudais, C.; Leiden, J. M.: GATA4 transcription

[70829] factor is required for ventral morphogenesis and heart tube formation. Genes

[70830] Dev. 11: 1048-1060, 1997.

[70831] 11996.

[70832] Reynolds, A. B.; Kanner, S. B.; Wang, H. C.; Parsons, J. T.: Stable

[70833] association of activated pp60src with two tyrosine-phosphorylated

[70834] cellular proteins. Molec. Cell. Biol. 9: 3951-3958, 1989.

[70835] 11997.

[70836] Sakai, R.; Iwamatsu, A.; Hirano, N.; Ogawa, S.; Tanaka, T.; Mano,

[70837] H.; Yazaki, Y.; Hirai, H.: A novel signaling molecule, p130, forms

[70838] stable complexes in vivo with v-Crk and v-Src in a tyro- sine phosphorylation-dependent

[70839] manner. EMBO J. 13: 3748-3756, 1994.

[70840] 11998.

[70841] van der Flier, S.; Brinkman, A.; Look, M. P.; Kok, E. M.; Meijer-van

[70842] Gelder, M. E.; Klijn, J. G. M.; Dorssers, L. C. J.; Foekens, J. A.

[70843] : Bcar1/p130Cas protein and primary breast cancer: prog- nosis and response

[70844] to tamoxifen treatment. J. Nat. Cancer Inst. 92: 120-127, 2000.

[70845] 11999.

[70846] Yoshida, K.; Taga, T.; Saito, M.; Suematsu, S.; Kumanogoh, A.;

[70847] Tanaka, T.; Fujiwara, H.; Hirata, M.; Yamagami, T.; Nakahata, T.;

[70848] Hirabayashi, T.; Yoneda, Y.; Tanaka, K.; Wang, W. Z.; Mori, C.; Shiota,

[70849] K.; Yoshida, N.; Kishimoto, T.: Targeted disruption of gp130, a common

[70850] signal transducer for the interleukin 6 family of cytokines, leads

[70851] to myocardial and hematological disorders. Proc. Nat. Acad. Sci. 93:

[70852] 407-411, 1996.

[70853] 12000.

[70854] Soengas, M. S.; Capodieci, P.; Polsky, D.; Mora, J.; Esteller,

[70855] M.; Opitz-Araya, X.; McCombie, R.; Herman, J. G.; Gerald, W. L.; Lazebnik,

[70856] Y. A.; Cordon-Cardo, C.; Lowe, S. W.: Inactivation of the apoptosis

[70857] effector Apaf-1 in malignant melanoma. Nature 409: 207-211, 2001.

[70858] 12001.

[70859] Srinivasula, S. M.; Ahmad, M.; Fernandes-Alnemri, T.; Alnemri,

[70860] E. S.: Autoactivation of procaspase-9 by Apaf-1-mediated oligomerization. Molec.

[70861] Cell 1: 949-957, 1998.

[70862] 12002.

[70863] Yoshida, H.; Kong, Y.-Y.; Yoshida, R.; Elia, A. J.; Hakem, A.;

[70864] Hakem, R.; Penninger, J. M.; Mak, T. W.: Apaf1 is required for mitochondrial

[70865] pathways of apoptosis and brain development. Cell 94: 739-750, 1998.

[70866] 12003.

[70867] Zou, H.; Henzel, W. J.; Liu, X.; Lutschg, A.; Wang, X.: APAF-1,

[70868] a human protein homologous to C. elegans CED-4, partic-ipates in cytochrome

[70869] c-dependent activation of caspase-3. Cell 90: 405-413, 1997.

[70870] 12004.

[70871] Niederreither, K.; Abu-Abed, S.; Schuhbaur, B.; Petkovich, M.;

[70872] Chambon, P.; Dolle, P.: Genetic evidence that oxidative derivatives

[70873] of retinoic acid are not involved in retinoid signaling during mouse
[70874] development. Nature Genet. 31: 84-88, 2002.

[70875] 12005.

[70876] White, J. A.; Beckett, B.; Scherer, S. W.; Herbrick, J.-A.; Petkovich,
[70877] M.: P450RAI (CYP26A) maps to human chromosome 10q23-q24 and mouse
[70878] chromosome 19C2-3. Genomics 48: 270-272, 1998.

[70879] 12006.

[70880] White, J. A.; Beckett-Jones, B.; Guo, Y.-D.; Dilworth, F. J.; Bonasoro,
[70881] J.; Jones, G.; Petkovich, M.: cDNA cloning of human retinoic acid-metabolizing
[70882] enzyme (hP450RAI) identifies a novel family of cytochromes P450 (CYP26). J.
[70883] Biol. Chem. 272: 18538-18541, 1997.

[70884] 12007.

[70885] Beutler, E.; Gelbart, T.; West, C.; Kuhl, W.; Lee, P.: A strategy
[70886] for cloning the hereditary hemochromatosis gene. Blood

Cells Molec.
[70887] Dis. 21: 207-216, 1995.

[70888] 12008.

[70889] Chardin, P.; Paris, S.; Antonny, B.; Robineau, S.; Beraud-Dufour,
[70890] S.; Jackson, C. L.; Chabre, M.: A human exchange factor for ARF contains
[70891] Sec7- and pleckstrin-homology domains. Nature 384: 481-484, 1996.
[70892] 12009.

[70893] Perletti, L.; Talarico, D.; Trecca, D.; Ronchetti, D.; Fracchi-olla,
[70894] N. S.; Maiolo, A. T.; Neri, A.: Identification of a novel gene, PSD,
[70895] adjacent to NFKB2/lyt-10, which contains Sec7 and pleck-strin-homology
[70896] domains. Genomics 46: 251-259, 1997.

[70897] 12010.

[70898] Biunno, I.; Appierto, V.; Cattaneo, M.; Leone, B. E.; Balzano,
[70899] G.; Socci, C.; Saccone, S.; Letizia, A.; Valle, G. D.;

Sgaramella,

[70900] V.: Isolation of a pancreas-specific gene located on human chromosome

[70901] 14q31: expression analysis in human pancreatic ductal carcinomas. Genomics 46:

[70902] 284-286, 1997.

[70903] 12011.

[70904] Donoviel, D. B.; Bernstein, A.: SEL-1L maps to human chromosome

[70905] 14, near the insulin-dependent diabetes mellitus locus 11. Genomics 56:

[70906] 232-233, 1999.

[70907] 12012.

[70908] Grant, B.; Greenwald, I.: The Caenorhabditis. elegans sel-1 gene,

[70909] a negative regulator of lin-12 and glp-1, encodes a predicted extracellular

[70910] protein. Genetics 143: 237-247, 1996.

[70911] 12013.

[70912] Grant, B.; Greenwald, I.: Structure, function and expression of

[70913] SEL-1, a negative regulator of LIN-12 and GLP-1 in C. ele-
gans. Development 124:
[70914] 637-644, 1997.

[70915] 12014.

[70916] Ben Porath, I.; Kozak, C. A.; Benvenisty, N.: Chromosomal
mapping
[70917] of Tmp (Emp1), Xmp (Emp2), and Ymp (Emp3), genes en-
coding membrane
[70918] proteins related to Pmp22. Genomics 49: 443-447, 1998.

[70919] 12015.

[70920] Chen, Y.; Medvedev, A.; Ruzanov, P.; Marvin, K. W.; Jetten,
A.
[70921] M.: cDNA cloning, genomic structure, and chromosome
mapping of the
[70922] human epithelial membrane protein CL-20 gene (EMP1), a
member of the
[70923] PMP22 family. Genomics 41: 40-48, 1997.

[70924] 12016.

[70925] Liehr, T.; Kuhlenbaumer, G.; Wulf, P.; Taylor, V.; Suter, U.;
Van
[70926] Broeckhoven, C.; Lupski, J. R.; Claussen, U.; Rauten-

strauss, B.:

[70927] Regional localization of the human epithelial membrane protein genes

[70928] 1, 2, and 3 (EMP1, EMP2, EMP3) to 12p12.3, 16p13.2, and 19q13.3. Genomics 58:

[70929] 106-108, 1999.

[70930] 12017.

[70931] Marvin, K. W.; Fujimoto, W.; Jetten, A. M.: Identification and

[70932] characterization of a novel squamous cell-associated gene related

[70933] to PMP22. J. Biol. Chem. 270: 28910-28916, 1995.

[70934] 12018.

[70935] Ruegg, C. L.; Wu, H.; Fagnoni, F. F.; Engleman, E. G.; Laus, R.

[70936] : B4B, a novel growth-arrest gene, is expressed by a sub-set of progenitor/pre-B

[70937] lymphocytes negative for cytoplasmic mu-chain. J. Im-mun. 157: 72-80,

[70938] 1996.

[70939] 12019.

[70940] Taylor, V.; Suter, U.: Epithelial membrane protein-2 and epithelial
[70941] membrane protein-3: two novel members of the peripheral myelin protein
[70942] 22 gene family. Gene 175: 115-120, 1996.

[70943] 12020.

[70944] Masiakowski, P.; Carroll, R. D.: A novel family of cell surface
[70945] receptors with tyrosine kinase-like domain. J. Biol. Chem. 267:
[70946] 26181-26190, 1992.

[70947] 12021.

[70948] Adelmant, G.; Begue, A.; Stehelin, D.; Laudet, V.: A functional
[70949] Rev-erb-alpha responsive element located in the human Rev-erb-alpha
[70950] promoter mediates a repressing activity. Proc. Nat. Acad. Sci. 93:
[70951] 3553-3558, 1996.

[70952] 12022.

[70953] Chawla, A.; Lazar, M. A.: Induction of Rev-ErbA alpha, an

orphan
[70954] receptor encoded on the opposite strand of the alpha-thyroid hormone
[70955] receptor gene, during adipocyte differentiation. J. Biol. Chem. 268:
[70956] 16265-16269, 1993.

[70957] 12023.

[70958] Lazar, M. A.; Hodin, R. A.; Darling, D. S.; Chin, W. W.: A novel
[70959] member of the thyroid/steroid hormone receptor family is encoded by
[70960] the opposite strand of the rat c-erbA-alpha transcriptional unit. Molec.
[70961] Cell. Biol. 9: 1128-1136, 1989.

[70962] 12024.

[70963] Miyajima, N.; Horiuchi, R.; Shibuya, S.; Matsubara, K.; Toyoshima,
[70964] K.; Yamamoto, T.: Two erbA homologs encoding proteins with different
[70965] T(3) binding capacities are transcribed from opposite DNA strands
[70966] of the same genetic locus. Cell 57: 31-39, 1989.

[70967] 12025.

[70968] Preitner, N.; Damiola, F.; Lopez-Molina, L.; Zakany, J.; Duboule,

[70969] D.; Albrecht, U.; Schibler, U.: The orphan nuclear receptor REV-ERB-alpha

[70970] controls circadian transcription within the positive limb of the mammalian

[70971] circadian oscillator. Cell 110: 251-260, 2002.

[70972] 12026.

[70973] Zhao, Q.; Khorasanizadeh, S.; Miyoshi, Y.; Lazar, M. A.; Rastinejad,

[70974] F.: Structural elements of an orphan nuclear receptor-DNA complex. Molec.

[70975] Cell 1: 849-861, 1998.

[70976] 12027.

[70977] Saito, T.; Seki, N.; Yamauchi, M.; Tsuji, S.; Hayashi, A.; Kozuma,

[70978] S.; Hori, T.: Structure, chromosomal location, and expres-sion profile

[70979] of EXTR1 and EXTR2, new members of the multiple exos-toses gene family. Biochem.

[70980] Biophys. Res. Commun. 243: 61-66, 1998.

[70981] 12028.

[70982] Wuyts, W.; Van Hul, W.: Characterization and genomic lo-
calization
[70983] of the mouse Extl2 gene. Cytogenet. Cell Genet. 89:
185-188, 2000.
[70984] 12029.

[70985] Wuyts, W.; Van Hul, W.; Hendrickx, J.; Speleman, F.;
Wauters, J.;
[70986] De Boulle, K.; Van Roy, N.; Van Agtmael, T.; Bossuyt, P.;
Willems,
[70987] P. J.: Identification and characterization of a novel member
of the
[70988] EXT gene family, EXTL2. Europ. J. Hum. Genet. 5:
382-389, 1997.
[70989] 12030.

[70990] Yue, S.; Serra, H. G.; Zoghbi, H. Y.; Orr, H. T.: The
spinocerebellar
[70991] ataxia type 1 protein, ataxin-1, has RNA-binding activity
that is
[70992] inversely affected by the length of its polyglutamine tract.
Hum.

[70993] Molec. Genet. 10: 25-30, 2001.

[70994] 12031.

[70995] Dick, T.; Ray, K.; Salz, H. K.; Chia, W.: Cytoplasmic dynein (ddlc1)

[70996] mutations cause morphogenetic defects and apoptotic cell death in

[70997] Drosophila melanogaster. Molec. Cell. Biol. 16: 1966-1977, 1996.

[70998] 12032.

[70999] Matsushita, M.; Endo, Y.; Taira, S.; Sato, Y.; Fujita, T.; Ichikawa,

[71000] N.; Nakata, M.; Misuochi, T.: A novel human lectin with collagen-

[71001] and fibrinogen-like domains which functions as an op- sonin. J. Biol.

[71002] Chem. 271: 2448-2454, 1996.

[71003] 12033.

[71004] Azuma, N.; Hirakiyama, A.; Inoue, T.; Asaka, A.; Yamada, M.: Mutations

[71005] of a human homologue of the Drosophila eyes absent gene (EYA1) detected

[71006] in patients with congenital cataracts and ocular anterior segment
[71007] anomalies. Hum. Molec. Genet. 9: 363-366, 2000.

[71008] 12034.

[71009] Johnson, K. R.; Cook, S. A.; Erway, L. C.; Matthews, A. N.; Sanford,
[71010] L. P.; Paradies, N. E.; Friedman, R. A.: Inner ear and kidney anomalies
[71011] caused by IAP insertion in an intron of the Eya1 gene in a mouse model
[71012] of BOR syndrome. Hum. Molec. Genet. 8: 645-653, 1999.

[71013] 12035.

[71014] Kumar, S.; Deffenbacher, K.; Cremers, C. W. R. J.; Van Camp, G.;
[71015] Kimberling, W. J.: Branchio-oto-renal syndrome: identifi-cation of
[71016] novel mutations, molecular characterization, mutation distribution,
[71017] and prospects for genetic testing. Genet. Testing 1: 243-251, 1998.

[71018] 12036.

[71019] Kumar, S.; Kimberling, W. J.; Weston, M. D.; Schaefer, B. G.; Berg,
[71020] M. A.; Marres, H. A. M.; Cremers, C. W. R. J.: Identification of
[71021] three novel mutations in human EYA1 protein associated with branchio-oto-renal
[71022] syndrome. Hum. Mutat. 11: 443-449, 1998.

[71023] 12037.

[71024] Rickard, S.; Boxer, M.; Trompeter, R.; Bitner-Glindzicz, M.: Importance
[71025] of clinical evaluation and molecular testing in the branchio-oto-renal
[71026] (BOR) syndrome and overlapping phenotypes. (Letter) J. Med. Genet. 37:
[71027] 623-627, 2000.

[71028] 12038.

[71029] Stoppa-Lyonnet, D.; Carter, P. E.; Meo, T.; Tosi, M.: Clusters
[71030] of intragenic Alu repeats predispose the human C1 inhibitor locus
[71031] to deleterious rearrangements. Proc. Nat. Acad. Sci. 87: 1551-1555,

[71032] 1990.

[71033] 12039.

[71034] Vincent, C.; Kalatzis, V.; Abdelhak, S.; Chaib, H.; Compain, S.;

[71035] Helia, J.; Vaneecloo, F.-M.; Petit, C.: BOR and BO syndromes are

[71036] allelic defects of EYA1. Europ. J. Hum. Genet. 5: 242-246, 1997.

[71037] 12040.

[71038] Wallace, M. R.; Andersen, L. B.; Saulino, A. M.; Gregory, P. E.;

[71039] Glover, T. W.; Collins, F. S.: A de novo Alu insertion results in

[71040] neurofibromatosis type 1. Nature 353: 864-866, 1991.

[71041] 12041.

[71042] Xu, P.-X.; Adams, J.; Peters, H.; Brown, M. C.; Heaney, S.; Maas,

[71043] R.: Eya1-deficient mice lack ears and kidneys and show abnormal apoptosis

[71044] of organ primordia. Nature Genet. 23: 113-117, 1999.

[71045] 12042.

[71046] Xu, P.-X.; Cheng, J.; Epstein, J. A.; Maas, R. L.: Mouse Eya

[71047] genes are expressed during limb tendon development and encode a transcriptional

[71048] activation function. Proc. Nat. Acad. Sci. 94: 11974-11979, 1997.

[71049] 12043.

[71050] Xu, P.-X.; Woo, I.; Her, H.; Beier, D. R.; Maas, R. L.: Mouse

[71051] Eya homologues of the Drosophila eyes absent gene re-quire Pax6 for

[71052] expression in lens and nasal placode. Development 124: 219-231,

[71053] 1997.

[71054] 12044.

[71055] Banfi, S.; Borsani, G.; Rossi, E.; Bernard, L.; Guffanti, A.; Rubboli,

[71056] F.; Marchitiello, A.; Giglio, S.; Coluccia, E.; Zollo, M.; Zuf-fardi,

[71057] O.; Ballabio, A.: Identification and mapping of human cD-NAs homologous

[71058] to Drosophila mutant genes through EST database search-ing. Nature

[71059] Genet. 13: 167-174, 1996.

[71060] 12045.

[71061] Xu, P.-X.; Cheng, J.; Epstein, J. A.; Maas, R. L.: Mouse Eya genes

[71062] are expressed during limb tendon development and encode a transcriptional

[71063] activation function. Proc. Nat. Acad. Sci. 94: 11974-11979, 1997.

[71064] 12046.

[71065] Zimmerman, J. E.; Bui, Q. T.; Steingrimsson, E.; Nagle, D. L.;

[71066] Fu, W.; Genin, A.; Spinner, N. B.; Copeland, N. G.; Jenkins, N. A.;

[71067] Bucan, M.; Bonini, N. M.: Cloning and characterization of two vertebrate

[71068] homologs of the Drosophila eyes absent gene. Genome Res. 7: 128-141,

[71069] 1997.

[71070] 12047.

[71071] Morrisey, E. E.; Tang, Z.; Sigrist, K.; Lu, M. M.; Jiang, F.; Ip,

[71072] H. S.; Parmacek, M. S.: GATA6 regulates HNF4 and is re-

quired for

[71073] differentiation of visceral endoderm in the mouse embryo. Genes Dev. 12:

[71074] 3579-3590, 1998.

[71075] 12048.

[71076] Suzuki, E.; Evans, T.; Lowry, J.; Truong, L.; Bell, D. W.; Testa,

[71077] J. R.; Walsh, K.: The human GATA-6 gene: structure, chro-

mosomal location,

[71078] and regulation of expression by tissue-specific and mito-

gen-responsive

[71079] signals. Genomics 38: 283-290, 1996.

[71080] 12049.

[71081] Meurs, E.; Chong, K.; Galabru, J.; Shaun, N.; Thomas, S. B.; Kerr,

[71082] I. M.; Williams, B. R. G.; Hovanessian, A. G.: Molecular cloning

[71083] and characterization of the human double-stranded RNA-

activated protein

[71084] kinase induced by interferon. Cell 62: 379-390, 1990.

[71085] 12050.

[71086] Eber, S. W.; Gonzalez, J. M.; Lux, M. L.; Scarpa, A. L.; Tse,

[71087] W. T.; Dornwell, M.; Herbers, J.; Kugler, W.; Ozcan, R.;
Pekrun, A.;

[71088] Gallagher, P. G.; Schroter, W.; Forget, B. G.; Lux, S. E.:
Ankyrin-1

[71089] mutations are a major cause of dominant and recessive
hereditary spherocytosis. Nature

[71090] Genet. 13: 214-218, 1996.

[71091] 12051.

[71092] Bashir, R.; Strachan, T.; Keers, S.; Stephenson, A.;
Mahjneh, I.;

[71093] Marconi, G.; Nashef, L.; Bushby, K. M. D.: A gene for auto-
somal recessive

[71094] limb-girdle muscular dystrophy maps to chromosome 2p.
Hum. Molec.

[71095] Genet. 3: 455-457, 1994.

[71096] 12052.

[71097] Katz, M. D.; Kligman, I.; Cai, L.-Q.; Zhu, Y.-S.; Fratianni, C.

[71098] M.; Zervoudakis, I.; Rosenwaks, Z.; Imperato-McGinley, J.:
Paternity

[71099] in intrauterine insemination with sperm from a man with

5-alpha-reductase-2
[71100] deficiency. New Eng. J. Med. 336: 994-997, 1997.

[71101] 12053.

[71102] Levy, G. G.; Nichols, W. C.; Lian, E. C.; Foroud, T.; Mc-
Clintick,
[71103] J. N.; McGee, B. M.; Yang, A. Y.; Slemieniak, D. R.; Stark, K.
R.;
[71104] Gruppo, R.; Sarode, R.; Shurin, S. B.; Chandrasekaran, V.;
Stabler,
[71105] S. P.; Sabio, H.; Bouhassira, E. E.; Upshaw, J. D., Jr.; Gins-
burg,
[71106] D.; Tsai, H.-M.: Mutations in a member of the ADAMTS
gene family
[71107] cause thrombotic thrombocytopenic purpura. Nature 413:
488-494,
[71108] 2001.

[71109] 12054.

[71110] Nicolao, P.; Carella, M.; Giometto, B.; Tavolato, B.; Cattin,

[71111] R.; Giovannucci-Uzielli, M. L.; Vacca, M.; Regione, F. D.;
Piva, S.;
[71112] Bortoluzzi, S.; Gasparini, P.: DHPLC analysis of the MECP2

gene in
[71113] Italian Rett patients. Hum. Mutat. 18: 132-140, 2001.

[71114] 12055.

[71115] Sasaki, G.; Ogata, T.; Ishii, T.; Hasegawa, T.; Sato, S.; Mat-
suo,
[71116] N.: Novel mutation of TBX3 in a Japanese family with ul-
nar-mammary
[71117] syndrome: implication for impaired sex development. Am.
J. Med. Genet. 110:
[71118] 365-369, 2002.

[71119] 12056.

[71120] Fargion, S.; Cappellini, M. D.; Piperno, A.; Panajotopoulos,
N.;
[71121] Ronchi, G.; Fiorelli, G.: Association of hereditary sphero-
cytosis
[71122] and idiopathic hemochromatosis: a synergistic effect in
determining
[71123] iron overload. Am. J. Clin. Path. 86: 645-649, 1986.

[71124] 12057.

[71125] Forget, B. G.; Lux, S. E.; Agre, P.; Watkins, P. C.; John, K.;

[71126] Costa, F. F.: Dominant hereditary spherocytosis (HS) is

linked to
[71127] the gene for the erythrocyte membrane protein ankyrin.(Abstract) Cytogenet.
[71128] Cell Genet. 51: 999, 1989.

[71129] 12058.

[71130] Gallagher, P. G.; Ferreira, J. D. S.; Costa, F. F.; Saad, S. T.

[71131] O.; Forget, B. G.: A recurrent frameshift mutation of the ankyrin
[71132] gene associated with severe hereditary spherocytosis. Brit. J. Haemat. 111:
[71133] 1190-1193, 2000.

[71134] 12059.

[71135] Gallagher, P. G.; Forget, B. G.: An alternate promoter di-rects
[71136] expression of a truncated, muscle-specific isoform of the human ankyrin
[71137] 1 gene. J. Biol. Chem. 273: 1339-1348, 1998.

[71138] 12060.

[71139] Gallagher, P. G.; Sabatino, D. E.; Basseres, D. S.; Nilson, D.

[71140] M.; Wong, C.; Cline, A. P.; Garrett, L. J.; Bodine, D. M.: Ery-

throcyte

[71141] ankyrin promoter mutations associated with recessive hereditary spherocytosis

[71142] cause significant abnormalities in ankyrin expression. J. Biol. Chem. 276:

[71143] 41683-41689, 2001.

[71144] 12061.

[71145] Greenquist, A. C.; Shohet, S. B.; Bernstein, S. E.: Marked reduction

[71146] of spectrin in hereditary spherocytosis in the common house mouse. Blood 51:

[71147] 1149-1155, 1978.

[71148] 12062.

[71149] Hanspal, M.; Yoon, S.-H.; Yu, H.; Hanspal, J. S.; Lambert, S.;

[71150] Palek, J.; Prchal, J. T.: Molecular basis of spectrin and ankyrin

[71151] deficiencies in severe hereditary spherocytosis: evidence implicating

[71152] a primary defect of ankyrin. Blood 77: 165-173, 1991.

[71153] 12063.

[71154] Hayette, S.; Carre, G.; Bozon, M.; Alloisio, N.; Maillet, P.;

[71155] Wilmotte, R.; Pascal, O.; Reynaud, J.; Reman, O.; Stephan, J.-L.;

[71156] Morle, L.; Delaunay, J.: Two distinct truncated variants of ankyrin

[71157] associated with hereditary spherocytosis. Am. J. Hemat. 58: 36-41,

[71158] 1998.

[71159] 12064.

[71160] Hill, J. S.; Sawyer, W. H.; Howlett, G. J.; Wiley, J. S.: Heredi-tary

[71161] spherocytosis of man: altered binding of cytoskeletal components to

[71162] the erythrocyte membrane. Biochem. J. 201: 259-266, 1982.

[71163] 12065.

[71164] Iolascon, A.; del Giudice, E. M.; Camaschella, C.; Pinto, L.;

[71165] Nobili, B.; Perrotta, S.; Cutillo, S.: Ankyrin deficiency in dominant

[71166] hereditary spherocytosis: report of three cases. Brit. J. Haemat. 78:

[71167] 551-554, 1991.

[71168] 12066.

[71169] Jacob, H. S.: Dysfunction of the red blood cell membrane in hereditary

[71170] spherocytosis. Brit. J. Haemat. 14: 99-104, 1968.

[71171] 12067.

[71172] Jacob, H. S.: Hereditary spherocytosis: a disease of the red

[71173] cell membrane. Seminars Hemat. 2: 139-166, 1965.

[71174] 12068.

[71175] Jacob, H. S.: Abnormalities in the physiology of the erythrocyte

[71176] membrane in hereditary spherocytosis. Am. J. Med. 41: 734-741, 1966.

[71177] 12069.

[71178] Jacob, H. S.; Amsden, T.; White, J.: Experimental production

[71179] of hereditary spherocytosis (HS): role of defective membrane microfilaments

[71180] in the disorder.(Abstract) J. Clin. Invest. 50: 48A, 1971.

[71181] 12070.

[71182] Jacob, H. S.; Jandl, J. H.: Increased cell membrane perme-
ability
[71183] in the pathogenesis of hereditary spherocytosis. J. Clin.
Invest. 43:
[71184] 1704-1720, 1964.

[71185] 12071.

[71186] Jacob, H. S.; Ruby, A.; Overland, E. S.; Mazia, D.: Abnormal

[71187] membrane protein of red blood cells in hereditary sphero-
cytosis. J.
[71188] Clin. Invest. 50: 1800-1805, 1971.

[71189] 12072.

[71190] Jandl, J. H.; Cooper, R. A.: Hereditary spherocytosis.In:
Stanbury,
[71191] J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic
Basis
[71192] of Inherited Disease. New York: McGraw-Hill (pub.) (3rd
ed.)
[71193] : 1972. Pp. 1323-1337.

[71194] 12073.

[71195] Jarolim, P.; Rubin, H. L.; Brabec, V.; Palek, J.: A nonsense

[71196] mutation glu1669-to-ter within the regulatory domain of human erythroid

[71197] ankyrin leads to a selective deficiency of the major ankyrin isoform

[71198] (band 2.1) and a phenotype of autosomal dominant hereditary spherocytosis. J.

[71199] Clin. Invest. 95: 941-947, 1995.

[71200] 12074.

[71201] Jensson, O.; Jonasson, J. L.; Magnusson, S.: Studies on hereditary

[71202] spherocytosis in Iceland. Acta Med. Scand. 201: 187-195, 1977.

[71203] 12075.

[71204] Johnsson, R.; Himberg, J.-J.: Thrombocyte aggregation in hereditary

[71205] spherocytosis. Clin. Chim. Acta 119: 257-262, 1982.

[71206] 12076.

[71207] Kimberling, W. J.; Fulbeck, T.; Dixon, L.; Lubs, H. A.: Lo-calization

[71208] of spherocytosis to chromosome 8 or 12 and report of a family with

[71209] spherocytosis and a reciprocal translocation. Am. J. Hum. Genet. 27:
[71210] 586-594, 1975.

[71211] 12077.

[71212] Kirkpatrick, F. H.; Woods, G. M.; LaCelle, P. L.: Absence of
[71213] one component of spectrin adenosine triphosphatase in hereditary spherocytosis. Blood 46:
[71214] 945-954, 1975.

[71215] 12078.

[71216] Kitatani, M.; Chiyo, H.; Ozaki, M.; Shike, S.; Miwa, S.: Lo-
calization
[71217] of the spherocytosis gene to chromosome segment 8p11.22-8p21.1. Hum.
[71218] Genet. 78: 94-95, 1988.

[71219] 12079.

[71220] Lambert, S.; Yu, H.; Prchal, J. T.; Lawler, J.; Ruff, P.; Spe-
icher,
[71221] D.; Cheung, M. C.; Kan, Y. W.; Palek, J.: cDNA sequence for human
[71222] erythrocyte ankyrin. Proc. Nat. Acad. Sci. 87: 1730-1734, 1990.

[71223] 12080.

[71224] Lefrere, J. J.; Courouce, A.-M.; Girot, R.; Bertrand, Y.; Soulier,

[71225] J.-P.: Six cases of hereditary spherocytosis revealed by human parvovirus

[71226] infection. Brit. J. Haemat. 62: 653-658, 1986.

[71227] 12081.

[71228] Leite, R. C. A.; Basseres, D. S.; Ferreira, J. S.; Alberto, F.

[71229] L.; Costa, F. F.; Saad, S. T. O.: Low frequency of ankyrin mutations

[71230] in hereditary spherocytosis: identification of three novel mutations. Hum.

[71231] Mutat. 16: 529 only, 2000.

[71232] 12082.

[71233] Lux, S. E.; John, K. M.; Bennett, V.: Analysis of cDNA for human

[71234] erythrocyte ankyrin indicates a repeated structure with homology to

[71235] tissue-differentiation and cell-cycle control proteins. Na-

[71236] ture 344: 36-42, 1990.

[71237] 12083.

[71238] Lux, S. E.; Tse, W. T.; Menninger, J. C.; John, K. M.; Harris,

[71239] P.; Shalev, O.; Chilcote, R. R.; Marchesi, S. L.; Watkins, P. C.;

[71240] Bennett, V.; McIntosh, S.; Collins, F. S.; Francke, U.; Ward, D. C.;

[71241] Forget, B. G.: Hereditary spherocytosis associated with deletion

[71242] of human erythrocyte ankyrin gene on chromosome 8. Nature 345: 736-739,

[71243] 1990.

[71244] 12084.

[71245] MacKinney, A. A.: Hereditary spherocytosis: clinical family studies. Arch.

[71246] Intern. Med. 116: 257-265, 1965.

[71247] 12085.

[71248] Amrani, A.; Verdaguer, J.; Serra, P.; Tafuro, S.; Tan, R.; Santamaria,

[71249] P.: Progression of autoimmune diabetes driven by avidity maturation

[71250] of a T-cell population. Nature 406: 739-742, 2000.

[71251] 12086.

[71252] Bowcock, A. M.; Kavathas, P.; Margolskee, R. F.; Herzen-
berg, L.;

[71253] Cavalli-Sforza, L. L.: An RFLP associated with pcDLeu2-14,
a human

[71254] T-cell differentiation antigen CD8 (Leu2) cDNA mapped to
2p12. Nucleic

[71255] Acids Res. 14: 7817 only, 1986.

[71256] 12087.

[71257] Bruns, G.; Kavathas, P.; Shiloh, Y.; Sakai, K.; Schwaber, J.;
Latt,

[71258] S. A.; Herzenberg, L. A.: The human T cell antigen Leu-2
(T8) is

[71259] encoded on chromosome 2. Hum. Genet. 70: 311-314,
1985.

[71260] 12088.

[71261] Giblin, P.; Ledbetter, J. A.; Kavathas, P.: A secreted form of

[71262] the human lymphocyte cell surface molecule CD8 arises
from alternative

[71263] splicing. Proc. Nat. Acad. Sci. 86: 998-1002, 1989.

[71264] 12089.

[71265] Kavathas, P.; Sukhatme, V. P.; Herzenberg, L. A.; Parnes, J. R.

[71266] : Isolation of the gene encoding the human T-lymphocyte differentiation

[71267] antigen Leu-2 (T8) by gene transfer and cDNA subtraction. Proc. Nat.

[71268] Acad. Sci. 81: 7688-7692, 1984.

[71269] 12090.

[71270] Ledbetter, J. A.; Evans, R. L.; Lipinski, M.; Cunningham-Rundles,

[71271] C.; Good, R. A.; Herzenberg, L. A.: Evolutionary conservation of

[71272] surface molecules that distinguish T lymphocyte helper-inducer and

[71273] cytotoxic-suppressor subpopulations in mouse and man. J. Exp. Med. 153:

[71274] 310-323, 1981.

[71275] 12091.

[71276] Ledbetter, J. A.; Seaman, W. E.; Tsu, T. T.; Herzenberg, L. A.

[71277] : Lyt-2 and Lyt-3 antigens are on two different polypeptide subunits

[71278] linked by disulfide bonds: relationship of subunits to T cell cytolytic
[71279] activity. J. Exp. Med. 153: 1503-1516, 1981.

[71280] 12092.

[71281] Leishman, A. J.; Naidenko, O. V.; Attinger, A.; Koning, F.; Lena,
[71282] C. J.; Xiong, Y.; Chang, H.-C.; Reinherz, E.; Kronenberg, M.; Cheroutre,
[71283] H.: T cell responses modulated through interaction between CD8-alpha-alpha
[71284] and the nonclassical MHC class I molecule, TL. Science 294: 1936-1939,
[71285] 2001.

[71286] 12093.

[71287] Littman, D. R.: The structure of the CD4 and CD8 genes. Annu.
[71288] Rev. Immun. 5: 561-584, 1987.

[71289] 12094.

[71290] Littman, D. R.; Thomas, Y.; Maddon, P. J.; Chess, L.; Axel, R.
[71291] : The isolation and sequence of the gene encoding T8: a

molecule defining
[71292] functional classes of T lymphocytes. Cell 40: 237-246, 1985.
[71293] 12095.
[71294] Mecucci, C.; Van Den Berghe, H.: OKT8-positive T-cell lymphoma
[71295] associated with a chromosome rearrangement t(2;17) possibly involving
[71296] the T8 locus. (Letter) New Eng. J. Med. 313: 185-186, 1985.
[71297] 12096.
[71298] Nakauchi, H.; Nolan, G. P.; Hsu, C.; Huang, H. S.; Kavathas, P.;
[71299] Herzenberg, L. A.: Molecular cloning of Lyt-2, a membrane glycoprotein
[71300] marking a subset of mouse T lymphocytes: molecular homology to its
[71301] human counterpart, Leu-2/T8, and to immunoglobulin variable regions. Proc.
[71302] Nat. Acad. Sci. 82: 5126-5130, 1985.
[71303] 12097.
[71304] Lindsay, E. A.; Vitelli, F.; Su, H.; Morishima, M.; Huynh, T.;

[71305] Pramparo, T.; Jurecic, V.; Ogunrinu, G.; Sutherland, H. F.; Scambler,
[71306] P. J.; Bradley, A.; Baldini, A.: Tbx1 haploinsufficiency in the DiGeorge
[71307] syndrome region causes aortic arch defects in mice. Nature 410:
[71308] 97-101, 2001.

[71309] 12098.

[71310] Merscher, S.; Funke, B.; Epstein, J. A.; Heyer, J.; Puech, A.;

[71311] Lu, M. M.; Xavier, R. J.; Demay, M. B.; Russell, R. G.; Factor, S.;
[71312] Tokooya, K.; St. Jore, B.; and 12 others: TBX1 is responsible for
[71313] cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. Cell 104:
[71314] 619-629, 2001.

[71315] 12099.

[71316] Peukert, K.; Staller, P.; Schneider, A.; Carmichael, G.; Hanel,
[71317] F.; Eilers, M.: An alternative pathway for gene regulation by Myc. EMBO

[71318] J. 16: 5672-5686, 1997.

[71319] 12100.

[71320] Calfon, M.; Zeng, H.; Urano, F.; Till, J. H.; Hubbard, S. R.; Harding,

[71321] H. P.; Clark, S. G.; Ron, D.: IRE1 couples endoplasmic reticulum

[71322] load to secretory capacity by processing the XBP-1 mRNA. Nature 415:

[71323] 92-96, 2002.

[71324] 12101.

[71325] Liou, H.-C.; Boothby, M. R.; Finn, P. W.; Davidson, R.; Nabavi,

[71326] N.; Zeleznik-Le, N. J.; Ting, J. P.-Y.; Glimcher, L. H.: A new member

[71327] of the leucine zipper class of proteins binds to the HLA DR-alpha

[71328] promoter. Science 247: 1581-1584, 1990.

[71329] 12102.

[71330] Liou, H.-C.; Eddy, R.; Shows, T.; Lisowska-Grospierre, B.; Griscelli,

[71331] C.; Doyle, C.; Mannhalter, J.; Eibl, M.; Glimcher, L. H.: An

HLA-DR-alpha
[71332] promoter DNA-binding protein is expressed ubiquitously and maps to
[71333] human chromosomes 22 and 5. Immunogenetics 34: 286-292, 1991.
[71334] 12103.

[71335] Reimold, A. M.; Iwakoshi, N. N.; Manis, J.; Vallabhajosyula, P.;
[71336] Szomolanyi-Tsuda, E.; Gravallese, E. M.; Friend, D.; Grusby, M. J.;
[71337] Alt, F.; Glimcher, L. H.: Plasma cell differentiation requires the
[71338] transcription factor XBP-1. Nature 412: 300-307, 2001.

[71339] 12104.

[71340] Blunt, T.; Taccioli, G. E.; Priestley, A.; Hafezparast, M.; McMillan,
[71341] T.; Liu, J.; Cole, C. C.; White, J.; Alt, F. W.; Jackson, S. P.; Schurr,
[71342] E.; Lehmann, A. R.; Jeggo, P. A.: A YAC contig encompass-ing the XRCC5
[71343] (Ku80) DNA repair gene and complementation defective cells by YAC

[71344] protoplast fusion. Genomics 30: 320-328, 1995.

[71345] 12105.

[71346] Nielsen, J. B.; Henriksen, K. F.; Hansen, C.; Silahtaroglu, A.;

[71347] Schwartz, M.; Tommerup, N.: MECP2 mutations in Danish patients with

[71348] Rett syndrome: high frequency of mutations but no consistent correlations

[71349] with clinical severity or with the X chromosome inactivation pattern. Europ.

[71350] J. Hum. Genet. 9: 178-184, 2001.

[71351] 12106.

[71352] Ohki, I.; Shimotake, N.; Fujita, N.; Jee, J.-G.; Ikegami, T.;

[71353] Nakao, M.; Shirakawa, M.: Solution structure of the methyl-CpG binding

[71354] domain of human MBD1 in complex with methylated DNA. Cell 105: 487-497,

[71355] 2001.

[71356] 12107.

[71357] Oh, J.; Ho, L.; Ala-Mello, S.; Amato, D.; Armstrong, L.; Bellucci,

[71358] S.; Carakushansky, G.; Ellis, J. P.; Fong, C.-T.; Green, J. S.; Heon,

[71359] E.; Legius, E.; Levin, A. V.; Nieuwenhuis, H. K.; Pinckers,

A.; Tamura,
[71360] N.; Whiteford, M. L.; Yamasaki, H.; Spritz, R. A.: Mutation analysis
[71361] of patients with Hermansky-Pudlak syndrome: a frameshift hot spot
[71362] in the HPS gene and apparent locus heterogeneity. Am. J. Hum. Genet. 62:
[71363] 593-598, 1998.

[71364] 12108.

[71365] Orrico, A.; Lam, C.-W.; Galli, L.; Dotti, M. T.; Hayek, G.; Tong,
[71366] S.-F.; Poon, P. M. K.; Zappella, M.; Federico, A.; Sorrentino, V.
[71367] : MECP2 mutation in male patients with non-specific X-linked mental
[71368] retardation. FEBS Lett. 481: 285-288, 2000.

[71369] 12109.

[71370] Quaderi, N. A.; Meehan, R. R.; Tate, P. H.; Cross, S. H.; Bird,
[71371] A. P.; Chatterjee, A.; Herman, G. E.; Brown, S. D. M.: Ge-netic and
[71372] physical mapping of a gene encoding a methyl CpG bind-

ing protein,

[71373] Mecp2, to the mouse X chromosome. Genomics 22: 648-651, 1994.

[71374] 12110.

[71375] Shahbazian, M. D.; Antalffy, B.; Armstrong, D. L.; Zoghbi, H.

[71376] Y.: Insight into Rett syndrome: MeCP2 levels display tissue- and

[71377] cell-specific differences and correlate with neuronal maturation. Hum.

[71378] Molec. Genet. 11: 115-124, 2002.

[71379] 12111.

[71380] Tate, P.; Skarnes, W.; Bird, A.: The methyl-CpG binding protein

[71381] MeCP2 is essential for embryonic development in the mouse. Nature

[71382] Genet. 12: 205, 1996.

[71383] 12112.

[71384] Topcu, M.; Akyerli, C.; Sayi, A.; Toruner, G. A.; Kocoglu, S.

[71385] R.; Cimbis, M.; Ozcelik, T.: Somatic mosaicism for a MECP2 mutation

[71386] associated with classic Rett syndrome in a boy. Europ. J. Hum. Genet. 10:

[71387] 77-81, 2002.

[71388] 12113.

[71389] Trappe, R.; Laccone, F.; Cobilanschi, J.; Meins, M.; Huppke, P.;

[71390] Hanefeld, F.; Engel, W.: MECP2 mutations in sporadic cases of Rett

[71391] syndrome are almost exclusively of paternal origin. Am. J. Hum. Genet. 68:

[71392] 1093-1101, 2001.

[71393] 12114.

[71394] Vilain, A.; Apiou, F.; Vogt, N.; Dutrillaux, B.; Malfoy, B.:

[71395] Assignment of the gene for methyl-CpG-binding protein 2 (MECP2) to

[71396] human chromosome band Xq28 by in situ hybridization. Cytogenet. Cell

[71397] Genet. 74: 293-294, 1996.

[71398] 12115.

[71399] Wan, M.; Lee, S. S. J.; Zhang, X.; Houwink-Manville, I.; Song,

[71400] H.-R.; Amir, R. E.; Budden, S.; Naidu, S.; Pereira, J. L. P.; Lo,
[71401] I. F. M.; Zoghbi, H. Y.; Schanen, N. C.; Francke, U.: Rett syndrome
[71402] and beyond: recurrent spontaneous and familial MECP2 mutations at
[71403] CpG hotspots. Am. J. Hum. Genet. 65: 1520-1529, 1999.

[71404] 12116.

[71405] Wan, M.; Zhao, K.; Lee, S. S. J.; Francke, U.: MECP2 truncating
[71406] mutations cause histone H4 hyperacetylation in Rett syndrome. Hum.
[71407] Molec. Genet. 10: 1085-1092, 2001.

[71408] 12117.

[71409] Watson, P.; Black, G.; Ramsden, S.; Barrow, M.; Super, M.; Kerr,
[71410] B.; Clayton-Smith, J.: Angelman syndrome phenotype associated with
[71411] mutations in MECP2, a gene encoding a methyl CpG binding protein. J.
[71412] Med. Genet. 38: 224-228, 2001.

[71413] 12118.

[71414] Willard, H. F.; Hendrich, B. D.: Breaking the silence in Rett

[71415] syndrome. Nature Genet. 23: 127-128, 1999.

[71416] 12119.

[71417] Xiang, F.; Buervenich, S.; Nicolao, P.; Bailey, M. E. S.; Zhang,

[71418] Z.; Anvret, M.: Mutation screening in Rett syndrome patients. J.

[71419] Med. Genet. 37: 250-255, 2000.

[71420] 12120.

[71421] Yntema, H. G.; Oudakker, A. R.; Kleefstra, T.; Hamel, B. C. J.;

[71422] van Bokhoven, H.; Chelly, J.; Kalscheuer, V. M.; Fryns, J.-P.; Raynaud,

[71423] M.; Moizard, M.-P.; Moraine, C.: In-frame deletion in MECP2 causes

[71424] mild nonspecific mental retardation. (Letter) Am. J. Med. Genet. 107:

[71425] 81-83, 2002.

[71426] 12121.

[71427] Yusufzai, T. M.; Wolffe, A. P.: Functional consequences of Rett
[71428] syndrome mutations on human MeCP2. Nucleic Acids Res. 28: 4172-4179,
[71429] 2000.

[71430] 12122.

[71431] Fuentes, J.-J.; Pritchard, M. A.; Planas, A. M.; Bosch, A.; Ferrer,
[71432] I.; Estivill, X.: A new human gene from the Down syndrome critical
[71433] region encodes a proline-rich protein highly expressed in fetal brain
[71434] and heart. Hum. Molec. Genet. 4: 1935-1944, 1995.

[71435] 12123.

[71436] Nakamura, A.; Hattori, M.; Sakaki, Y.: Isolation of a novel human
[71437] gene from the Down syndrome critical region of chromo-some 21q22.2. J.
[71438] Biochem. 122: 872-877, 1997.

[71439] 12124.

[71440] Schallreuter, K. U.; Frenk, E.; Wolfe, L. S.; Witkop, C. J.;

Wood,

[71441] J. M.: Hermansky-Pudlak syndrome in a Swiss population. Dermatology 187:

[71442] 248-256, 1993.

[71443] 12125.

[71444] Suzuki, T.; Li, W.; Zhang, Q.; Karim, A.; Novak, E. K.; Sviderskaya,

[71445] E. V.; Hill, S. P.; Bennett, D. C.; Levin, A. V.; Nieuwenhuis, H.

[71446] K.; Fong, C.-T.; Castellan, C.; Miterski, B.; Swank, R. T.; Spritz,

[71447] R. A.: Hermansky-Pudlak syndrome is caused by mutations in HPS4,

[71448] the human homolog of the mouse light-ear gene. Nature Genet. 30:

[71449] 321-324, 2002.

[71450] 12126.

[71451] Wildenberg, S. C.; Oetting, W. S.; Almodovar, C.; Krumwiede, M.;

[71452] White, J. G.; King, R. A.: A gene causing Hermansky-Pudlak syndrome

[71453] in a Puerto Rican population maps to chromosome 10q2.

Am. J. Hum.
[71454] Genet. 57: 755-765, 1995.

[71455] 12127.

[71456] Magnussen, K.: Beitrag zur Genetik und Histologie eines isolierten
[71457] Augenalbinismus beim Kaninchen. Z. Morph. Anthrop. 44: 127-135,
[71458] 1952.

[71459] 12128.

[71460] MacKinney, A. A.; Morton, N. E.; Kosower, N. S.; Schilling, R.
[71461] F.: Ascertaining genetic carriers of hereditary spherocytosis by
[71462] statistical analysis of multiple laboratory tests. J. Clin. Invest. 41:
[71463] 554-567, 1962.

[71464] 12129.

[71465] Sukhatme, V. P.; Sizer, K. C.; Vollmer, A. C.; Hunkapiller, T.;
[71466] Parnes, J. R.: The T cell differentiation antigen leu-2/T8 is homologous

[71467] to immunoglobulin and T cell receptor variable regions. Cell 40:
[71468] 591-597, 1985.

[71469] 12130.

[71470] Chen, D. J.; Marrone, B. L.; Nguyen, T.; Stackhouse, M.; Zhao,
[71471] Y.; Siciliano, M. J.: Regional assignment of a human DNA repair gene
[71472] (XRCC5) to 2q35 by x-ray hybrid mapping. Genomics 21: 423-427, 1994.
[71473] 12131.

[71474] Kirchgessner, C. U.; Patil, C. K.; Evans, J. W.; Cuomo, C. A.;

[71475] Fried, L. M.; Carter, T.; Oettinger, M. A.; Brown, J. M.: DNA-dependent
[71476] kinase (p350) as a candidate gene for the murine SCID defect. Science 267:
[71477] 1178-1183, 1995.

[71478] 12132.

[71479] Reardon, W.; Coffey, R.; Phelps, P. D.; Luxon, L. M.; Stephens,
[71480] D.; Kendall-Taylor, P.; Britton, K. E.; Grossman, A.; Trem-

bath, R.

[71481] : Pendred syndrome--100 years of underascertainment? Quart. J. Med. 90:

[71482] 443-447, 1997.

[71483] 12133.

[71484] Royaux, I. E.; Wall, S. M.; Karniski, L. P.; Everett, L. A.; Suzuki,

[71485] K.; Knepper, M. A.; Green, E. D.: Pendrin, encoded by the Pendred

[71486] syndrome gene, resides in the apical region of renal inter-calated

[71487] cells and mediates bicarbonate secretion. Proc. Nat. Acad. Sci. 98:

[71488] 4221-4226, 2001.

[71489] 12134.

[71490] Taylor, J. P.; Metcalfe, R. A.; Watson, P. F.; Weetman, A. P.;

[71491] Trembath, R. C.: Mutations of the PDS gene, encoding pendrin, are

[71492] associated with protein mislocalization and loss of iodide efflux:

[71493] implications for thyroid dysfunction in Pendred syndrome.

J. Clin.

[71494] Endocr. Metab. 87: 1778-1784, 2002.

[71495] 12135.

[71496] Douville, P.; Seldin, M. F.; Carbonetto, S.: Genetic mapping of

[71497] the integrin alpha-1 gene (Vla1) to mouse chromosome 13. Genomics 14:

[71498] 503-505, 1992.

[71499] 12136.

[71500] Bettaieb, A.; Fromont, P.; Rodet, M.; Godeau, B.; Duedari, N.;

[71501] Bierling, P.: Br(b), a platelet alloantigen involved in neonatal

[71502] alloimmune thrombocytopenia. Vox Sang. 60: 230-234, 1991.

[71503] 12137.

[71504] Carlsson, L. E.; Santoso, S.; Spitzer, C.; Kessler, C.; Greinacher,

[71505] A.: The alpha-2 gene coding sequence T807/A873 of the platelet collagen

[71506] receptor integrin alpha-2/beta-1 might be a genetic risk

factor for
[71507] the development of stroke in younger patients. Blood 93: 3583-3586,
[71508] 1999.

[71509] 12138.

[71510] Chen, D. J.; Park, M. S.; Campbell, E.; Oshimura, M.; Liu, P.;
[71511] Zhao, Y.; White, B. F.; Siciliano, M. J.: Assignment of a human DNA
[71512] double-strand break repair gene (XRCC5) to chromosome 2. Genomics 13:
[71513] 1088-1094, 1992.

[71514] 12139.

[71515] Difilippantonio, M. J.; Zhu, J.; Chen, H. T.; Meffre, E.; Nussenzweig,
[71516] M. C.; Max, E. E.; Ried, T.; Nussenzweig, A.: DNA repair protein
[71517] Ku80 suppresses chromosomal aberrations and malignant transformation. Nature 404:
[71518] 510-514, 2000.

[71519] 12140.

[71520] Jeggo, P. A.; Hafezparast, M.; Thompson, A. F.; Broughton, B. C.;

[71521] Kaur, G. P.; Zdzienicka, M. Z.; Athwal, R. S.: Localization of a

[71522] DNA repair gene (XRCC5) involved in double-strand-break rejoining

[71523] to human chromosome 2. Proc. Nat. Acad. Sci. 89: 6423-6427, 1992.

[71524] 12141.

[71525] Taccioli, G. E.; Gottlieb, T. M.; Blunt, T.; Priestley, A.; De-mengeot,

[71526] J.; Mizuta, R.; Lehmann, A. R.; Alt, F. W.; Jackson, S. P.; Jeggo,

[71527] P. A.: Ku80: product of the XRCC5 gene and its role in DNA repair

[71528] and V(D)J recombination. Science 265: 1442-1445, 1994.

[71529] 12142.

[71530] Ohkuchi, A.; Shiraishi, H.; Minakami, H.; Eguchi, Y.; Izumi, A.;

[71531] Sato, I.: Fetus with long QT syndrome manifested by tach-yarrhythmia:

[71532] a case report. Prenat. Diag. 19: 990-992, 1999.

[71533] 12143.

[71534] Pacia, S. V.; Devinsky, O.; Luciano, D. J.; Vazquez, B.: The

[71535] prolonged QT syndrome presenting as epilepsy: a report of two cases

[71536] and literature review. Neurology 44: 1408-1410, 1994.

[71537] 12144.

[71538] Piippo, K.; Swan, H.; Pasternack, M.; Chapman, H.; Paavonen, K.;

[71539] Viitasalo, M.; Toivonen, L.; Kontula, K.: A founder mutation of the

[71540] potassium channel KCNQ1 in long QT syndrome: implications for estimation

[71541] of disease prevalence and molecular diagnostics. J. Am. Coll. Cardiol. 37:

[71542] 562-568, 2001.

[71543] 12145.

[71544] Pony, J. C.; Matheyses, M.; Daubert, J. C.; Fourdilis, M.; Gouffault,

[71545] J.: Le syndrome QT long-syncope familial: deux observations de syndrome

[71546] de Romano et Ward. Arch. Mal. Coeur 70: 1105-1114,

1977.

[71547] 12146.

[71548] Priori, S. G.; Schwartz, P. J.; Napolitano, C.; Bianchi, L.; Dennis,

[71549] A.; De Fusco, M.; Brown, A. M.; Casari, G.: A recessive variant of

[71550] the Romano-Ward Long-QT syndrome? Circulation 97: 2420-2425, 1998.

[71551] 12147.

[71552] Romano, C.: Congenital cardiac arrhythmia. (Letter) Lancet I:

[71553] 658-659, 1965.

[71554] 12148.

[71555] Romano, C.; Gemme, G.; Pongiglione, R.: Aritmie car-diache rare

[71556] dell' eta pediatrica. II. Accessi sincopali per fibrillazione ventricolare

[71557] parossistica. (Presentazione del primo caso della letter-atura pediatrica

[71558] Italiana.) Clin. Pediat. 45: 656-683, 1963.

[71559] 12149.

[71560] Roy, N.; Kahlem, P.; Dausse, E.; Bennaceur, M.; Faure, S.; Weissenbach,
[71561] J.; Komajda, M.; Denjoy, I.; Coumel, P.; Schwartz, K.; Guicheney,
[71562] P.: Exclusion of HRAS from long QT locus. (Letter) Nature Genet. 8:
[71563] 113-114, 1994.

[71564] 12150.

[71565] Roy, P. R.; Emanuel, R.; Ismail, S. A.; Tayib, M. H.: Heredi-tary
[71566] prolongation of the Q-T interval: genetic observations and management
[71567] in three families with 12 affected members. Am. J. Cardiol. 37:
[71568] 237-243, 1976.

[71569] 12151.

[71570] Russell, M. W.; Dick, M., II; Collins, F. S.; Brody, L. C.: KVLQT1
[71571] mutations in three families with familial or sporadic long QT syndrome. Hum.
[71572] Molec. Genet. 5: 1319-1324, 1996.

[71573] 12152.

[71574] Schmitt, N.; Schwarz, M.; Peretz, A.; Abitbol, I.; Attali, B.;

[71575] Pongs, O.: A recessive C-terminal Jervell and Lange-Nielsen mutation

[71576] of the KCNQ1 channel impairs subunit assembly. EMBO J. 19: 332-340,

[71577] 2000.

[71578] 12153.

[71579] Schwartz, P. J.: Cardiac sympathetic innervation and the sudden

[71580] infant death syndrome: a possible pathogenetic link. Am. J. Med. 60:

[71581] 167-172, 1976.

[71582] 12154.

[71583] Schwartz, P. J.; Priori, S. G.; Bloise, R.; Napolitano, C.; Ronchetti,

[71584] E.; Piccinini, A.; Goj, C.; Breithardt, G.; Schulze-Bahr, E.; Wedekind,

[71585] H.; Nastoli, J.: Molecular diagnosis in a child with sudden infant

[71586] death syndrome. (Letter) Lancet 358: 1342-1343, 2001.

[71587] 12155.

[71588] Schwartz, P. J.; Stramba-Badiale, M.; Segantini, A.; Austoni,

[71589] P.; Bosi, G.; Giorgetti, R.; Grancini, F.; Marni, E. D.; Perticone,

[71590] F.; Rosti, D.; Salice, P.: Prolongation of the QT interval and the

[71591] sudden infant death syndrome. New Eng. J. Med. 338: 1709-1714, 1998.

[71592] 12156.

[71593] Shalaby, F. Y.; Levesque, P. C.; Yang, W.-P.; Little, W. A.; Conder,

[71594] M. L.; Jenkins-West, T.; Blanar, M. A.: Dominant-negative KvLQT1

[71595] mutations underlie the LQT1 form of long QT syndrome. Circulation 96:

[71596] 1733-1736, 1997.

[71597] 12157.

[71598] Shimizu, W.; Kurita, T.; Matsuo, K.; Suyama, K.; Aihara, N.; Kamakura,

[71599] S.; Towbin, J. A.; Shimomura, K.: Improvement of repolarization abnormalities

[71600] by a K+ channel opener in the LQT1 form of congenital long-QT syndrome. Circulation 97:
[71601] 1581-1588, 1998.

[71602] 12158.

[71603] Singer, P. A.; Crampton, R. S.; Bass, N. H.: Familial Q-T prolongation
[71604] syndrome: convulsive seizures and paroxysmal ventricular fibrillation. Arch.
[71605] Neurol. 31: 64-66, 1974.

[71606] 12159.

[71607] Smilinich, N. J.; Day, C. D.; Fitzpatrick, G. V.; Caldwell, G.

[71608] M.; Lossie, A. C.; Cooper, P. R.; Smallwood, A. C.; Joyce, J. A.;
[71609] Schofield, P. N.; Reik, W.; Nicholls, R. D.; Weksberg, R.; Driscoll,
[71610] D. J.; Maher, E. R.; Shows, T. B.; Higgins, M. J.: A maternally methylated
[71611] CpG island in KvLQT1 is associated with an antisense paternal transcript
[71612] and loss of imprinting in Beckwith-Wiedemann syndrome. Proc. Nat.

[71613] Acad. Sci. 96: 8064-8069, 1999.

[71614] 12160.

[71615] Splawski, I.; Timothy, K. W.; Vincent, G. M.; Atkinson, D. L.;

[71616] Keating, M. T.: Molecular basis of the long-QT syndrome associated

[71617] with deafness. New Eng. J. Med. 336: 1562-1567, 1997.

[71618] 12161.

[71619] Campbell, H. D.; Webb, G. C.; Young, I. G.: A human homologue

[71620] of the Drosophila melanogaster sluggish-A (proline oxidase) gene maps

[71621] to 22q11.2, and is a candidate gene for type-I hyperprolinaemia. Hum.

[71622] Genet. 101: 69-74, 1997.

[71623] 12162.

[71624] Miura, K.; Yasuda, K.; Yanase, T.; Yamakita, N.; Sasano, H.; Nawata,

[71625] H.; Inoue, M.; Fukaya, T.; Shizuta, Y.: Mutation of cytochrome P-45017

[71626] alpha gene (CYP17) in a Japanese patient previously re-

ported as having
[71627] glucocorticoid-responsive hyperaldosteronism: with a re-
view of Japanese
[71628] patients with mutations of CYP17. J. Clin. Endocr. Metab.
81: 3797-3801,
[71629] 1996.

[71630] 12163.

[71631] New, M. I.: Male pseudohermaphroditism due to
17-alpha-hydroxylase
[71632] deficiency. J. Clin. Invest. 49: 1930-1941, 1970.

[71633] 12164.

[71634] Oshiro, C.; Takasu, N.; Wakugami, T.; Komiya, I.; Yamada,
T.;
[71635] Eguchi, Y.; Takei, H.: Seventeen alpha-hydroxylase defi-
ciency with
[71636] one base pair deletion of the cytochrome P450c17
(CYP17) gene. J.
[71637] Clin. Endocr. Metab. 80: 2526-2529, 1995.

[71638] 12165.

[71639] Picado-Leonard, J.; Miller, W. L.: Structure of the human
P450c17

[71640] gene is closely related to the P450c21 gene. (Abstract) Am. J. Hum.

[71641] Genet. 41: A234, 1987.

[71642] 12166.

[71643] Rovner, D. R.; Gordon, D. L.: Direct evidence for a functional

[71644] block with 18 oxidation in a patient with 17-alpha-hydroxylase deficiency. Trans.

[71645] Assoc. Am. Phys. 91: 416-423, 1978.

[71646] 12167.

[71647] Scaroni, C.; Biason, A.; Carpene, G.; Opocher, G.; Mantero, F.

[71648] : 17-Alpha-hydroxylase deficiency in three siblings: short- and long-term

[71649] studies. J. Endocr. Invest. 14: 99-108, 1991.

[71650] 12168.

[71651] Slominski, A.; Ermak, G.; Mihm, M.: ACTH receptor, CYP11A1, CYP17

[71652] and CYP21A2 genes are expressed in skin. J. Clin. Endocr. Metab. 81:

[71653] 2746-2749, 1996.

[71654] 12169.

[71655] Suzuki, Y.; Nagashima, T.; Nomura, Y.; Onigata, K.; Na-
gashima,
[71656] K.; Morikawa, A.: A new compound heterozygous muta-
tion (W17X, 436+5G-to-T)
[71657] in the cytochrome P450c17 gene causes
17-alpha-hydroxylase/17,20-lyase
[71658] deficiency. J. Clin. Endocr. Metab. 83: 199-202, 1998.

[71659] 12170.

[71660] Winter, J. S. D.; Couch, R. M.; Muller, J.; Perry, Y. S.; Fer-
reira,
[71661] P.; Baydala, L.; Shackleton, C. H. L.: Combined
17-hydroxylase and
[71662] 17,20-desmolase deficiencies: evidence for synthesis of a
defective
[71663] cytochrome P450(c17). J. Clin. Endocr. Metab. 68:
309-316, 1989.
[71664] 12171.

[71665] Yamaguchi, H.; Nakazato, M.; Miyazato, M.; Kangawa, K.;
Matsukura,
[71666] S.: A 5-prime-splice site mutation in the cytochrome P450
steroid

[71667] 17-alpha-hydroxylase gene in 17-alpha-hydroxylase deficiency. J.

[71668] Clin. Endocr. Metab. 82: 1934-1938, 1997.

[71669] 12172.

[71670] Yanase, T.; Kagimoto, M.; Suzuki, S.; Hashiba, K.; Simpson, E.

[71671] R.; Waterman, M. R.: Deletion of a phenylalanine in the N-terminal

[71672] region of human cytochrome P-450(17a) results in partial combined

[71673] 17-alpha-hydroxylase/17,20-lyase deficiency. J. Biol. Chem. 264:

[71674] 18076-18082, 1989.

[71675] 12173.

[71676] Yanase, T.; Sanders, D.; Shibata, A.; Matsui, N.; Simpson, E.

[71677] R.; Waterman, M. R.: Combined 17-alpha-hydroxylase/17,20-lyase deficiency

[71678] due to a 7-basepair duplication in the N-terminal region of the cytochrome

[71679] P450(17-alpha) (CYP17) gene. J. Clin. Endocr. Metab. 70: 1325-1329,

[71680] 1990.

[71681] 12174.

[71682] Yanase, T.; Simpson, E. R.; Waterman, M. R.:
17-Alpha-hydroxylase/17,20-lyase
[71683] deficiency: from clinical investigation to molecular defini-
tion. Endocr.
[71684] Rev. 12: 91-108, 1991.

[71685] 12175.

[71686] Yazaki, K.; Kuribayashi, T.; Yamamura, Y.; Kurihara, T.;
Araki,
[71687] S.: Hypokalemic myopathy associated with
17-alpha-hydroxylase deficiency:
[71688] a case report. Neurology 32: 94-97, 1982.

[71689] 12176.

[71690] Youngblood, G. L.; Sartorius, C.; Taylor, B. A.; Payne, A. H.

[71691] : Isolation, characterization, and chromosomal mapping of
mouse P450
[71692] 17-alpha-hydroxylase/C17-20 lyase. Genomics 10:
270-275, 1991.
[71693] 12177.

[71694] Zlotogora, J.; Gieselmann, V.; Bach, G.: Multiple mutations in
[71695] a specific gene in a small geographic area: a common phenomenon? (Letter) Am.
[71696] J. Hum. Genet. 58: 241-243, 1996.

[71697] 12178.

[71698] Touraine, J. L.; Betuel, H.; Souillet, G.; Jeune, M.: Com-bined
[71699] immunodeficiency disease associated with absence of cell-surface HLA-A
[71700] and -B antigens. J. Pediat. 93: 47-51, 1978.

[71701] 12179.

[71702] Wolf, H. M.; Hauber, I.; Gulle, H.; Thon, V.; Eggenbauer, H.;
[71703] Fischer, M. B.; Fiala, S.; Eibl, M. M.: Brief report: Twin boys with
[71704] major histocompatibility complex class II deficiency but inducible
[71705] immune responses. New Eng. J. Med. 332: 86-90, 1995.

[71706] 12180.

[71707] Dupuis, S.; Dargemont, C.; Fieschi, C.; Thomassin, N.;

Rosenzweig,
[71708] S.; Harris, J.; Holland, S. M.; Schreiber, R. D.; Casanova, J.-L.
[71709] : Impairment of mycobacterial but not viral immunity by a germline
[71710] human STAT1 mutation. Science 293: 300-303, 2001.

[71711] 12181.

[71712] Findlay, K. A. B.; Kaptein, E.; Visser, T. J.; Burchell, B.: Characterization
[71713] of the uridine diphosphate-glucuronosyltrans-ferase-catalyzing thyroid
[71714] hormone glucuronidation in man. J. Clin. Endocr. Metab. 85: 2879-2883,
[71715] 2000.

[71716] 12182.

[71717] Wooster, R.; Sutherland, L.; Ebner, T.; Clarke, D.; Da Cruz e Silva,
[71718] O.; Burchell, B.: Cloning and stable expression of a new member of
[71719] the human liver phenol/biliru-bin:UDP-glucuronosyltransferase cDNA
[71720] family. Biochem. J. 278: 465-469, 1991.

[71721] 12183.

[71722] Martiniuk, F.; Mehler, M.; Pellicer, A.; Tzall, S.; La Badie,

[71723] G.; Hobart, C.; Ellenbogen, A.; Hirschhorn, R.: Isolation of a cDNA

[71724] for human acid alpha-glucosidase and detection of genetic heterogeneity

[71725] for mRNA in three alpha-glucosidase-deficient patients. Proc. Nat.

[71726] Acad. Sci. 83: 9641-9644, 1986.

[71727] 12184.

[71728] Nickel, B. E.; Chudley, A. E.; Pabello, P. D.; McAlpine, P. J.

[71729] : Exclusion mapping of the GAA locus to chromosome 17q21-q25. Cytogenet.

[71730] Cell Genet. 32: 303-304, 1982.

[71731] 12185.

[71732] Nickel, B. E.; McAlpine, P. J.: Extension of human acid alpha-glucosidase

[71733] polymorphism by isoelectric focusing in polyacrylamide gel. Ann.

[71734] Hum. Genet. 46: 97-103, 1982.

[71735] 12186.

[71736] Sandison, A.; Broadhead, D. M.; Bain, A. D.: Elucidation of an

[71737] unbalanced chromosome translocation by gene dosage studies. Clin.

[71738] Genet. 22: 30-36, 1982.

[71739] 12187.

[71740] Shieh, J.-J.; Lin, C.-Y.: Frequent mutation in Chinese pa-tients

[71741] with infantile type of GSD II in Taiwan: evidence for a founder effect. Hum.

[71742] Mutat. 11: 306-312, 1998.

[71743] 12188.

[71744] Solomon, E.; Swallow, D. M.; Burgess, S.; Evans, L.: As-signment

[71745] of the human acid alpha-glucosidase gene (alpha-GLU) to chromosome

[71746] 17 using somatic cell hybrids. Ann. Hum. Genet. 42: 273-281, 1979.

[71747] 12189.

[71748] Tsunoda, H.; Ohshima, J.; Tohyama, J.; Sasaki, M.; Sakura-

gawa,
[71749] N.; Martiniuk, F.: Acid-alpha-glucosidase deficiency: identification
[71750] and expression of a missense mutation (S529V) in a Japanese adult
[71751] phenotype. Hum. Genet. 97: 496-499, 1996.

[71752] 12190.

[71753] Van der Kraan, M.; Kroos, M. A.; Joosse, M.; Bijvoet, A. G. A.;
[71754] Verbeet, M. P.; Kleijer, W. J.; Reuser, A. J. J.: Deletion of exon
[71755] 18 is a frequent mutation in glycogen storage disease type II. Biochem.
[71756] Biophys. Res. Commun. 203: 1535-1541, 1994.

[71757] 12191.

[71758] Van der Ploeg, A. T.; Hoefsloot, L. H.; Hoogeveen-Westerveld,
[71759] M.; Petersen, E. M.; Reuser, A. J. J.: Glycogenosis type II: protein
[71760] and DNA analysis in five South African families from various ethnic
[71761] origins. Am. J. Hum. Genet. 44: 787-793, 1989.

[71762] 12192.

[71763] Vorgerd, M.; Burwinkel, B.; Reichmann, H.; Malin, J.-P.; Kilimann,

[71764] M. W.: Adult-onset glycogen storage disease type II: phe-notypic and

[71765] allelic heterogeneity in German patients. Neurogenetics 1: 205-211,

[71766] 1998.

[71767] 12193.

[71768] Walvoort, H. C.; Slee, R. G.; Sluis, K. J.; Koster, J. F.; Reuser,

[71769] A. J. J.: Biochemical genetics of the Lapland dog model of glycogen

[71770] storage disease type II (acid alpha-glucosidase deficiency). Am.

[71771] J. Med. Genet. 19: 589-598, 1984.

[71772] 12194.

[71773] Yan, B.; Raben, N.; Lu, N.; Plotz, P. H.: Identification and

[71774] characterization of a tissue-specific silencer element in the first

[71775] intron of the human acid maltase gene. Hum. Genet. 109:

186-190,

[71776] 2001.

[71777] 12195.

[71778] Zhong, N.; Martiniuk, F.; Tzall, S.; Hirschhorn, R.: Identifi-
cation

[71779] of a missense mutation in one allele of a patient with
Pompe disease,

[71780] and use of endonuclease digestion of PCR-amplified RNA
to demonstrate

[71781] lack of mRNA expression from the second allele. Am. J.
Hum. Genet. 49:

[71782] 635-645, 1991.

[71783] 12196.

[71784] Tanaka, S.; Yamashita, S.; Hosaka, K.: Cloning and ex-
pression

[71785] of human cDNA encoding phosphatidylinositol transfer
protein beta. Biochim.

[71786] Biophys. Acta 1259: 199-202, 1995.

[71787] 12197.

[71788] Dijkstra, M.; Veld, G. I.; van den Berg, G. J.; Muller, M.;
Kuipers,

[71789] F.; Vonk, R. J.: Adenosine triphosphate-dependent copper transport
[71790] in isolated rat liver plasma membranes. J. Clin. Invest. 95: 412-416,
[71791] 1995.

[71792] 12198.

[71793] Figus, A.; Angius, A.; Loudianos, G.; Bertini, C.; Dessi, V.; Loi,
[71794] A.; Deiana, M.; Lovicu, M.; Olla, N.; Sole, G.; De Virgiliis, S.;
[71795] Lilliu, F.; and 21 others: Molecular pathology and haplo-type analysis
[71796] of Wilson disease in Mediterranean populations. Am. J. Hum. Genet. 57:
[71797] 1318-1324, 1995.

[71798] 12199.

[71799] Cossee, M.; Durr, A.; Schmitt, M.; Dahl, N.; Trouillas, P.; Allinson,
[71800] P.; Kostrzewa, M.; Nivelon-Chevallier, A.; Gustavson, K.-H.; Kohlschutter,
[71801] A.; Muller, U.; Mandel, J.-L.; and 11 others: Friedreich's ataxia:

[71802] point mutations and clinical presentation of compound heterozygotes. Ann.

[71803] Neurol. 45: 200-206, 1999.

[71804] 12200.

[71805] De Castro, M.; Garcia-Planells, J.; Monros, E.; Canizares, J.;

[71806] Vazquez-Manrique, R.; Vilchez, J. J.; Urtasun, M.; Lucus, M.; Navarro,

[71807] G.; Izquierdo, G.; Molto, M. D.; Palau, F.: Genotype and phenotype

[71808] analysis of Friedreich's ataxia compound heterozygous patients. Hum.

[71809] Genet. 106: 86-92, 2000.

[71810] 12201.

[71811] Delatycki, M. B.; Paris, D.; Gardner, R. J. M.; Forshaw, K.; Nicholson,

[71812] G. A.; Nassif, N.; Williamson, R.; Forrest, S. M.: Sperm DNA analysis

[71813] in a Friedreich ataxia premutation carrier suggests both meiotic and

[71814] mitotic expansion in the FRDA gene. J. Med. Genet. 35: 713-716,

[71815] 1998.

[71816] 12202.

[71817] Duclos, F.; Boschert, U.; Sirugo, G.; Mandel, J.-L.; Hen, R.;

[71818] Koenig, M.: Gene in the region of the Friedreich ataxia locus encodes

[71819] a putative transmembrane protein expressed in the nervous system. Proc.

[71820] Nat. Acad. Sci. 90: 109-113, 1993.

[71821] 12203.

[71822] Doi, A.; Shiosaka, T.; Takaoka, Y.; Yanagisawa, K.; Fujita, S.

[71823] : Molecular cloning of the cDNA encoding A+U-rich element RNA binding

[71824] factor. Biochim. Biophys. Acta 1396: 51-56, 1998.

[71825] 12204.

[71826] Kamei, D.; Tsuchiya, N.; Yamazaki, M.; Meguro, H.; Yamada, M.:

[71827] Two forms of expression and genomic structure of the human heterogeneous

[71828] nuclear ribonucleoprotein D-like JKTBP gene (HNRPDL). Gene 228:

[71829] 13-22, 1999.

[71830] 12205.

[71831] Tsuchiya, N.; Kamei, D.; Takano, A.; Matsui, T.; Yamada, M.: Cloning

[71832] and characterization of a cDNA encoding a novel heterogeneous nuclear

[71833] ribonucleoprotein-like protein and its expression in myeloid leukemia

[71834] cells. J. Biochem. 123: 499-507, 1998.

[71835] 12206.

[71836] Ma, Y. H.; Betard, C.; Roy, M.; Davignon, J.; Kessling, A. M.

[71837] : Identification of a second 'French Canadian' LDL receptor gene deletion

[71838] and development of a rapid method to detect both deletions. Clin.

[71839] Genet. 36: 219-228, 1989.

[71840] 12207.

[71841] Mandelshtam, M.; Chakir, K.; Shevtsov, S.; Golubkov, V.; Skobeleva,

[71842] N.; Lipovetsky, B.; Konstantinov, V.; Denisenko, A.; Gaitskhoki, V.;

[71843] Schwartz, E.: Prevalence of Lithuanian mutation among St. Petersburg

[71844] Jews with familial hypercholesterolemia. Hum. Mutat. 12: 255-258,

[71845] 1998.

[71846] 12208.

[71847] Meiner, V.; Landsberger, D.; Berkman, N.; Reshef, A.; Se-gal, P.;

[71848] Seftel, H. C.; van der Westhuyzen, D. R.; Jeenah, M. S.; Co-etzee,

[71849] G. A.; Leitersdorf, E.: A common Lithuanian mutation causing familial

[71850] hypercholesterolemia in Ashkenazi Jews. Am. J. Hum. Genet. 49: 443-449,

[71851] 1991.

[71852] 12209.

[71853] Miyake, Y.; Tajima, S.; Funahashi, T.; Yamamura, T.; Ya-mamoto,

[71854] A.: A point mutation of low-density-lipoprotein receptor causing

[71855] rapid degradation of the receptor. Europ. J. Biochem. 210: 1-7,

[71856] 1992.

[71857] 12210.

[71858] Moorjani, S.; Roy, M.; Torres, A.; Betard, C.; Gagne, C.; Lambert,

[71859] M.; Brun, D.; Davignon, J.; Lupien, P.: Mutations of low-density-lipoprotein-receptor

[71860] gene, variation in plasma cholesterol, and expression of coronary

[71861] heart disease in homozygous familial hypercholestero-laemia. Lancet 341:

[71862] 1303-1306, 1993.

[71863] 12211.

[71864] Oppenheim, A.; Friedlander, Y.; Dann, E. J.; Berkman, N.; Schwartz,

[71865] S. P.; Leitersdorf, E.: Hypercholesterolemia in five Israeli Christian-Arab

[71866] kindreds is caused by the 'Lebanese' allele at the low den-sity lipoprotein

[71867] receptor gene locus and by an additional independent major factor. Hum.

[71868] Genet. 88: 75-84, 1991.

[71869] 12212.

[71870] Pisciotta, L.; Cantafora, A.; De Stefano, F.; Langheim, S.; Calandra,

[71871] S.; Bertolini, S.: A 'de novo' mutation of the LDL-receptor gene

[71872] as the cause of familial hypercholesterolemia. Biochim. Biophys.

[71873] Acta 1587: 7-11, 2002.

[71874] 12213.

[71875] Rodningen, O. K.; Rosby, O.; Tonstad, S.; Ose, L.; Berg, K.; Leren,

[71876] T. P.: A 9.6 kilobase deletion in the low density lipoprotein receptor

[71877] gene in Norwegian familial hypercholesterolemia subjects. Clin. Genet. 42:

[71878] 288-295, 1992.

[71879] 12214.

[71880] Rubinsztein, D. C.; Coetzee, G. A.; Marais, A. D.; Leiters-dorf,

[71881] E.; Seftel, H. C.; van der Westhuyzen, D. R.: Identification and

[71882] properties of the proline-644-to-leucine mutant LDL receptor in South

[71883] Africans of Indian origin. J. Lipid Res. 33: 1647-1655, 1992.

[71884] 12215.

[71885] Rudiger, N. S.; Heinsvig, E. M.; Hansen, F. A.; Faergeman, O.;

[71886] Bolund, L.; Gregersen, N.: DNA deletions in the low density lipoprotein

[71887] (LDL) receptor gene in Danish families with familial hypercholesterolemia. Clin.

[71888] Genet. 39: 451-462, 1991.

[71889] 12216.

[71890] Ruffner, D. E.; Sprung, C. N.; Minghetti, P. P.; Gibbs, P. E.

[71891] M.; Dugaiczyk, A.: Invasion of the human albumin-alpha-fetoprotein

[71892] gene family by Alu, Kpn, and two novel repetitive DNA elements. Molec.

[71893] Biol. Evol. 4: 1-9, 1987.

[71894] 12217.

[71895] Russell, D. W.; Lehrman, M. A.; Sudhof, T. C.; Yamamoto,

T.; Davis,
[71896] C. G.; Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.: The LDL receptor
[71897] in familial hypercholesterolemia: use of human mutations to dissect
[71898] a membrane protein. Cold Spring Harbor Symp. Quant. Biol. 51: 811-819,
[71899] 1986.

[71900] 12218.

[71901] Russell, D. W.; Schneider, W. J.; Yamamoto, T.; Luskey, K. L.;
[71902] Brown, M. S.; Goldstein, J. L.: Domain map of the LDL receptor: sequence
[71903] homology with the epidermal growth factor precursor. Cell 37: 577-585,
[71904] 1984.

[71905] 12219.

[71906] Schuster, H.; Manke, C.; Fischer, J.; Keller, C.; Wolfram, G.;
[71907] Zollner, N.: Identification of the valine 408 to methionine mutation
[71908] in the LDL receptor in a Greek patient with homozygous

familial hypercholesterolemia. Clin.
[71909] Genet. 48: 90-92, 1995.

[71910] 12220.

[71911] Schuster, H.; Ostwald, P.; Keller, P.; Wolfram, G.; Keller, C.

[71912] : Identification of the serine-156 to leucine mutation in the low-density
[71913] lipoprotein receptor in a German family with familial hypercholesterolemia. Clin.
[71914] Investig. 71: 172-175, 1993.

[71915] 12221.

[71916] Slagel, V.; Flemington, E.; Traina-Dorge, V.; Bradshaw, H.; Deininger,
[71917] P.: Clustering and subfamily relationships of the Alu family in the
[71918] human genome. Molec. Biol. Evol. 4: 19-29, 1987.

[71919]

[71920] 100. Soutar, A. K.; Knight, B. L.; Patel, D. D.: Identification of
[71921] a point mutation in growth repeat C of the low density lipoprotein-receptor
[71922] gene in a patient with homozygous familial hypercholes-

terolemia that

affects ligand binding and intracellular movement of receptors. Proc.

Nat. Acad. Sci. 86: 4166-4170, 1989.

101. Soutar, A. K.; McCarthy, S. N.; Seed, M.; Knight, B. L.: Relationship

between apolipoprotein(a) phenotype, lipoprotein(a) concentration

in plasma, and low density lipoprotein receptor function in a large

kindred with familial hypercholesterolemia due to the pro664-to-leu

mutation in the LDL receptor gene. J. Clin. Invest. 88: 483-492,

1991.

102. Steyn, K.; Weight, M. J.; Dando, B. R.; Christopher, K. J.; Rossouw,

J. E.: The use of low density lipoprotein receptor activity of lymphocytes

to determine the prevalence of familial hypercholestero-

laemia in a
[71936] rural South African community. J. Med. Genet. 26: 32-36, 1989.
[71937]

[71938] 103. Sudhof, T. C.; Goldstein, J. L.; Brown, M. S.; Russell, D. W.
[71939] : The LDL receptor gene: a mosaic of exons shared with different proteins. Science 228:
[71940] 815-822, 1985.
[71941]

[71942] 104. Sudhof, T. C.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.;
[71943] Sanchez-Pescador, R.; Bell, G. I.: Cassette of eight exons shared
[71944] by genes for LDL receptor and EGF precursor. Science 228: 893-895,
[71945] 1985.
[71946]

[71947] 105. Sun, X.-M.; Patel, D. D.; Bhatnager, D.; Knight, B. L.; Soutar,
[71948] A. K.: Characterization of a splice-site mutation in the

gene for

[71949] the LDL receptor associated with an unpredictably severe clinical

[71950] phenotype in English patients with heterozygous FH. Arteroscler.

[71951] Thromb. Vasc. Biol. 15: 219-227, 1995.

[71952]

[71953] 106. Takahashi, M.; Ikeda, U.; Takahashi, S.; Hattori, H.; Iwasaki,

[71954] T.; Ishihara, M.; Egashira, T.; Honma, S.; Asano, Y.; Shi-mada, K.

[71955] : A novel mutation in exon 2 of the low-density lipopro-tein-receptor

[71956] gene in a patient with homozygous familial hypercholes-terolemia. (Letter) Clin.

[71957] Genet. 59: 290-292, 2001.

[71958]

[71959] 107. Taylor, R.; Bryant, J.; Gudnason, V.; Sigurdsson, G.; Humphries,

[71960] S.: A study of familial hypercholesterolaemia in Iceland using RFLPs. J.

[71961] Med. Genet. 26: 494-498, 1989.

[71962]

[71963] 108. Thiart, R.; Scholtz, C. L.; Vergotine, J.; Hoogendijk, C. F.;

[71964] de Villiers, J. N. P.; Nissen, H.; Brusgaard, K.; Gaffney, D.; Hoffs,

[71965] M. S.; Vermaak, W. J. H.; Kotze, M. J.: Predominance of a 6 bp deletion

[71966] in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia. J.

[71967] Med. Genet. 37: 514-519, 2000.

[71968]

[71969] 109. Tolleshaug, H.; Goldstein, J. L.; Schneider, W. J.; Brown, M.

[71970] S.: Posttranslational processing of the LDL receptor and its genetic

[71971] disruption in familial hypercholesterolemia. Cell 30: 715-724, 1982.

[71972]

[71973] 110. Top, B.; Koeleman, B. P. C.; Gevers Leuven, J. A.; Havekes, L.

[71974] M.; Frants, R. R.: Rearrangements in the LDL receptor gene in Dutch

[71975] familial hypercholesterolemic patients and the presence of a common

[71976] 4 kb deletion. Atherosclerosis 83: 127-136, 1990.

[71977]

[71978] 111. Top, B.; Uitterlinden, A. G.; van der Zee, A.; Kastelein, J.

[71979] J. P.; Gevers Leuven, J. A.; Havekes, L. M.; Frants, R. R.: Absence

[71980] of mutations in the promoter region of the low density lipoprotein

[71981] receptor gene in a large number of familial hypercholes-terolaemia

[71982] patients as revealed by denaturing gradient gel elec-trophoresis. Hum.

[71983] Genet. 89: 561-565, 1992.

[71984]

[71985] 112. Torrington, M.; Botha, J. L.: Familial hypercholestero-laemia

[71986] and church affiliation. (Letter) Lancet II: 1120 only, 1981.

[71987]

[71988] 113. Ullu, E.; Tschudi, C.: Alu sequences are processed

7SL RNA genes. Nature 312:
[71989] 171-172, 1984.

[71990]

[71991] 114. Varret, M.; Rabes, J.-P.; Collod-Beroud, G.; Junien, C.;
Boileau,
[71992] C.; Beroud, C.: Software and database for the analysis of
mutations
[71993] in the human LDL receptor gene. Nucleic Acids Res. 25:
172-180,
[71994] 1997.

[71995]

[71996] 115. Vergopoulos, A.; Bajari, T.; Jouma, M.; Knoblauch, H.;
Aydin,
[71997] A.; Bahring, S.; Mueller-Myhsok, B.; Dresel, A.; Joubran, R.;
Luft,
[71998] F. C.; Schuster, H.: A xanthomatosis-susceptibility gene
may exist
[71999] in a Syrian family with familial hypercholesterolemia. Eu-
rop. J.
[72000] Hum. Genet. 5: 315-323, 1997.

[72001]

[72002] 116. Vergotine, J.; Thiart, R.; Langenhoven, E.; Hillermann, R.; De

[72003] Jong, G.; Kotze, M. J.: Prenatal diagnosis of familial hypercholesterolemia:

[72004] importance of DNA analysis in the high-risk South African population. Genet.

[72005] Counsel. 12: 121-127, 2001.

[72006]

[72007] 117. Vuorio, A. F.; Turtola, H.; Piilahti, K.-M.; Repo, P.; Kanninen,

[72008] T.; Kontula, K.: Familial hypercholesterolemia in the Finnish North

[72009] Karelia: a molecular, clinical, and genealogical study. Arterioscler.

[72010] Thromb. Vasc. Biol. 17: 3127-3138, 1997.

[72011]

[72012] 118. Wilson, D. J.; Gahan, M.; Haddad, L.; Heath, K.; Whittall, R.

[72013] A.; Williams, R. R.; Humphries, S. E.; Day, I. N. M.: A World Wide

[72014] Web site for low-density lipoprotein receptor gene mutations in familial

[72015] hypercholesterolemia: sequence-based, tabular, and direct submission

[72016] data handling. Am. J. Cardiol. 81: 1509-1511, 1998.

[72017]

[72018] 119. Yamakawa, K.; Okafuji, T.; Iwamura, Y.; Yuzawa, K.; Satoh, J.;

[72019] Hattori, N.; Yamanouchi, Y.; Yanagi, H.; Kawai, K.; Tsuchiya, S.;

[72020] Russell, D. W.; Hamaguchi, H.: TaqI polymorphism in the LDL receptor

[72021] gene and a TaqI 1.5-kb band associated with familial hypercholesterolemia. Hum.

[72022] Genet. 80: 1-5, 1988.

[72023]

[72024] 120. Yamakawa, K.; Takada, K.; Yanagi, H.; Tsuchiya, S.; Kawai, K.;

[72025] Nakagawa, S.; Kajiyama, G.; Hamaguchi, H.: Three novel partial deletions

[72026] of the low-density lipoprotein (LDL) receptor gene in familial hypercholesterolemia. Hum.

[72027] Genet. 82: 317-321, 1989.

[72028]

[72029] 121. Yamamoto, T.; Davis, C. G.; Brown, M. S.; Schneider, W. J.; Casey,

[72030] M. L.; Goldstein, J. L.; Russell, D. W.: The human LDL receptor:

[72031] a cysteine-rich protein with multiple Alu sequences in its mRNA. Cell 39:

[72032] 27-38, 1984.

[72033]

[72034] 122. Zuliani, G.; Hobbs, H. H.: Personal Communication. Dallas,

[72035] Tex. 1990.

[72036] 12222.

[72037] Schomburg, L.; Turwitt, S.; Prescher, G.; Lohmann, D.; Horsthemke,

[72038] B.; Bauer, K.: Human TRH-degrading ectoenzyme: cDNA cloning, functional

[72039] expression, genomic structure and chromosomal assignment. Europ.

[72040] J. Biochem. 265: 415-422, 1999.

[72041] 12223.

[72042] Adra, C. N.; Lelias, J.-M.; Kobayashi, H.; Kaghad, M.; Morrison,
[72043] P.; Rowley, J. D.; Lim, B.: Cloning of the cDNA for a hematopoietic
[72044] cell-specific protein related to CD20 and the beta subunit of the
[72045] high-affinity IgE receptor: evidence for a family of proteins with
[72046] four membrane-spanning regions. Proc. Nat. Acad. Sci. 91: 10178-10182,
[72047] 1994.

[72048] 12224.

[72049] Nagase, T.; Kikuno, R.; Hattori, A.; Kondo, Y.; Okumura, K.; Ohara,
[72050] O.: Prediction of the coding sequences of unidentified human genes.
[72051] XIX. The complete sequences of 100 new cDNA clones from brain which
[72052] code for large proteins in vitro. DNA Res. 7: 347-355, 2000.
[72053] 12225.

[72054] O'Carroll, D.; Scherthan, H.; Peters, A. H. F. M.; Opravil, S.;

[72055] Haynes, A. R.; Laible, G.; Rea, S.; Schmid, M.; Lebersorger, A.; Jerratsch,

[72056] M.; Sattler, L.; Mattei, M. G.; Denny, P.; Brown, S. D. M.; Schweizer,

[72057] D.; Jenuwein, T.: Isolation and characterization of Suv39h2, a second

[72058] histone H3 methyltransferase gene that displays testis-specific expression. Molec.

[72059] Cell. Biol. 20: 9423-9433, 2000.

[72060] 12226.

[72061] Zhou, X. Z.; Lu, K. P.: The Pin2/TRF-1-interacting protein PinX1

[72062] is a potent telomerase inhibitor. Cell 107: 347-359, 2001.

[72063] 12227.

[72064] Davis, R. S.; Wang, Y.-H.; Kubagawa, H.; Cooper, M. D.: Identification

[72065] of a family of Fc receptor homologs with preferential B cell expression. Proc.

[72066] Nat. Acad. Sci. 98: 9772-9777, 2001.

[72067] 12228.

[72068] Xu, M.; Zhao, R.; Zhao, Z. J.: Molecular cloning and char-

acterization
[72069] of SPAP1, an inhibitory receptor. Biochem. Biophys. Res. Commun. 280:
[72070] 768-775, 2001.

[72071] 12229.

[72072] Abu-Jawdeh, G.; Comella, N.; Tomita, Y.; Brown, L. F.; Tognazzi,
[72073] K.; Sokol, S. Y.; Kocher, O.: Differential expression of fr-pHE: a
[72074] novel human stromal protein of the secreted frizzled gene family,
[72075] during the endometrial cycle and malignancy. Lab. Invest. 79: 439-447,
[72076] 1999.

[72077] 12230.

[72078] Firneisz, G.; Lakatos, P. L.; Szalay, F.; Polli, C.; Glant, T.

[72079] T.; Ferenci, P.: Common mutations of ATP7B in Wilson disease patients
[72080] from Hungary. Am. J. Med. Genet. 108: 23-28, 2002.

[72081] 12231.

[72082] Yoshikawa, T.; Sanders, A. R.; Esterling, L. E.; Detera-

Wadleigh,

[72083] S. D.: Multiple transcriptional variants and RNA editing in C18orf1,

[72084] a novel gene with LDLRA and transmembrane domains on 18p11.2. Genomics 47:

[72085] 246-257, 1998.

[72086] 12232.

[72087] Yoshikawa, T.; Sanders, A. R.; Esterling, L. E.; Overharser, J.;

[72088] Garnes, J. A.; Lennon, G.; Grewal, R.; Detera-Wadleigh, S. D.: Isolation

[72089] of chromosome 18-specific brain transcripts as positional candidates

[72090] for bipolar disorder. Am. J. Med. Genet. (Neuropsych. Genet.) 74:

[72091] 140-149, 1997.

[72092] 12233.

[72093] Anneren, C.; Reedquist, K. A.; Bos, J. L.; Welsh, M.: GTK, a Src-related

[72094] tyrosine kinase, induces nerve growth factor-independent neurite outgrowth

[72095] in PC12 cells through activation of the Rap1 pathway: re-

lationship
[72096] to Shb tyrosine phosphorylation and elevated levels of fo-
cal adhesion
[72097] kinase. J. Biol. Chem. 275: 29153-29161, 2000.

[72098] 12234.

[72099] Cance, W. G.; Craven, R. J.; Bergman, M.; Xu, L.; Alitalo, K.;

[72100] Liu, E. T.: Rak, a novel nuclear tyrosine kinase expressed
in epithelial
[72101] cells. Cell Growth Differ. 5: 1347-1355, 1994.

[72102] 12235.

[72103] Cance, W. G.; Craven, R. J.; Weiner, T. M.; Liu, E. T.: Novel

[72104] protein kinases expressed in human breast cancer. Int. J.
Cancer 54:
[72105] 571-577, 1993.

[72106] 12236.

[72107] Lee, J.; Wang, Z.; Luoh, S.-M.; Wood, W. I.; Scadden, D. T.:
Cloning
[72108] of FRK, a novel human intracellular SRC-like tyrosine ki-
nase-encoding
[72109] gene. Gene 138: 247-251, 1994.

[72110] 12237.

[72111] Scott, A. F.: Personal Communication. Baltimore, Md. 12/19/2001.

[72112] 12238.

[72113] Betz, A.; Thakur, P.; Junge, H. J.; Ashery, U.; Rhee, J.-S.; Scheuss,

[72114] V.; Rosenmund, C.; Rettig, J.; Brose, N.: Functional interaction

[72115] of the active zone proteins Munc13-1 and RIM1 in synaptic vesicle

[72116] priming. Neuron 30: 183-196, 2001.

[72117] 12239.

[72118] Castillo, P. E.; Schoch, S.; Schmitz, F.; Sudhof, T. C.; Malenka,

[72119] R. C.: RIM1-alpha is required for presynaptic long-term potentiation. Nature 415:

[72120] 327-330, 2002.

[72121] 12240.

[72122] Coppola, T.; Magnin-Luthi, S.; Perret-Menoud, V.; Gattesco, S.;

[72123] Schiavo, G.; Regazzi, R.: Direct interaction of the Rab3 ef-

fector

[72124] RIM with Ca(2+) channels, SNAP-25, and synaptotagmin. J. Biol. Chem. 276:

[72125] 32756-32762, 2001.

[72126] 12241.

[72127] Schoch, S.; Castillo, P. E.; Jo, T.; Mukherjee, K.; Geppert, M.;

[72128] Wang, Y.; Schmitz, F.; Malenka, R. C.; Sudhof, T. C.: RIM1-alpha

[72129] forms a protein scaffold for regulating neurotransmitter release at

[72130] the active zone. Nature 415: 321-326, 2002.

[72131] 12242.

[72132] Wang, Y.; Sugita, S.; Sudhof, T. C.: The RIM/NIM family of neuronal

[72133] C-2 domain proteins: interactions with Rab3 and a new class of Src

[72134] homology 3 domain proteins. J. Biol. Chem. 275: 20033-20044, 2000.

[72135] 12243.

[72136] Kitamoto, Y.; Veile, R. A.; Donis-Keller, H.; Sadler, J. E.:

cDNA

[72137] sequence and chromosomal localization of human enterokinase, the proteolytic

[72138] activator of trypsinogen. Biochemistry 34: 4562-4568, 1995.

[72139] 12244.

[72140] Kitamoto, Y.; Yuan, X.; Wu, Q.; McCourt, D. W.; Sadler, J. E.:

[72141] Enterokinase, the initiator of intestinal digestion, is a mosaic protease

[72142] composed of a distinctive assortment of domains. Proc. Nat. Acad.

[72143] Sci. 91: 7588-7592, 1994.

[72144] 12245.

[72145] Magrangeas, F.; Apiou, F.; Denis, S.; Weidle, U.; Jacques, Y.;

[72146] Minvielle, S.: Cloning and expression of CIS6, chromosome assignment

[72147] to 3p22 and 2p21 by in situ hybridization. Cytogenet. Cell Genet. 88:

[72148] 78-81, 2000.

[72149] 12246.

[72150] Vreugde, S.; Erven, A.; Kros, C. J.; Marcotti, W.; Fuches, H.;

[72151] Kurima, K.; Wilcox, E. R.; Friedman, T. B.; Griffith, A. J.;
Balling,

[72152] R.; de Angelis, M. H.; Avraham, K. B.; Steel, K. P.:
Beethoven, a

[72153] mouse model for dominant, progressive hearing loss
DFNA36. Nature

[72154] Genet. 30: 257-258, 2002.

[72155] 12247.

[72156] Grunder, S.; Geissler, H.-S.; Bassler, E.-L.; Ruppersberg, J.
P.

[72157] : A new member of acid-sensing ion channels from pitu-
itary gland. Neuroreport 11:

[72158] 1607-1611, 2000.

[72159] 12248.

[72160] Tatarelli, C.; Linnenbach, A.; Mimori, K.; Croce, C. M.:
Characterization

[72161] of the human TESTIN gene localized in the FRA7G region
at 7q31.2. Genomics 68:

[72162] 1-12, 2000.

[72163] 12249.

[72164] Tobias, E. S.; Hurlstone, A. F. L.; MacKenzie, E.; McFarlane, R.;

[72165] Black, D. M.: The TES gene at 7q31.1 is methylated in tumours and

[72166] encodes a novel growth-suppressing LIM domain protein. Oncogene 20:

[72167] 2844-2853, 2001.

[72168] 12250.

[72169] Ferdinandusse, S.; Mulders, J.; IJlst, L.; Denis, S.; Dacremont,

[72170] G.; Waterham, H. R.; Wanders, R. J. A.: Molecular cloning and expression

[72171] of human carnitine octanoyltransferase: evidence for its role in the

[72172] peroxisomal beta-oxidation of branched-chain fatty acids. Biochem.

[72173] Biophys. Res. Commun. 263: 213-218, 1999.

[72174] 12251.

[72175] Horowitz, D. S.; Kobayashi, R.; Krainer, A. R.: A new cyclophilin

[72176] and the human homologues of yeast Prp3 and Prp4 form a complex associated

[72177] with U4/U6 snRNPs. RNA 3: 1374-1387, 1997.

[72178] 12252.

[72179] Reidt, U.; Reuter, K.; Achsel, T.; Ingelfinger, D.; Luhrmann, R.;

[72180] Ficner, R.: Crystal structure of the human U4/U6 small nuclear ribonucleoprotein

[72181] particle-specific SnuCyp-20, a nuclear cyclophilin. J. Biol. Chem. 275:

[72182] 7439-7442, 2000.

[72183] 12253.

[72184] Teigelkamp, S.; Achsel, T.; Mundt, C.; Gothel, S.-F.; Cronshagen,

[72185] U.; Lane, W. S.; Marahiel, M.; Luhrmann, R.: The 20kD protein of

[72186] human [U4/U6.U5] tri-snRNPs is a novel cyclophilin that forms a complex

[72187] with the U4/U6-specific 60kD and 90kD proteins. RNA 4: 127-141,

[72188] 1998.

[72189] 12254.

[72190] Gaynor, E. C.; Mondesert, G.; Grimme, S. J.; Reed, S. I.; Or-

lean,

[72191] P.; Emr, S. D.: MCD4 encodes a conserved endoplasmic reticulum membrane

[72192] protein essential for glycosylphosphatidylinositol anchor synthesis

[72193] in yeast. Molec. Biol. Cell 10: 627-648, 1999.

[72194] 12255.

[72195] Hong, Y.; Maeda, Y.; Watanabe, R.; Ohishi, K.; Mishkind, M.; Riezman,

[72196] H.; Kinoshita, T.: Pig-n, a mammalian homologue of yeast Mcd4p, is

[72197] involved in transferring phosphoethanolamine to the first mannose

[72198] of the glycosylphosphatidylinositol. J. Biol. Chem. 274: 35099-35106,

[72199] 1999.

[72200] 12256.

[72201] Parks, W. T.; Frank, D. B.; Huff, C.; Haft, C. R.; Martin, J.;

[72202] Meng, X.; de Caestecker, M. P.; McNally, J. G.; Reddi, A.; Taylor,

[72203] S. I.; Roberts, A. B.; Wang, T.; Lechleider, R. J.: Sorting

nexin

[72204] 6, a novel SNX, interacts with the transforming growth factor-beta

[72205] family of receptor serine-threonine kinases. J. Biol. Chem. 276:

[72206] 19332-19339, 2001.

[72207] 12257.

[72208] Gao, J.; Yu, L.; Zhang, P.; Jiang, J.; Chen, J.; Peng, J.; Wei,

[72209] Y.; Zhao, S.: Cloning and characterization of human and mouse mitochondrial

[72210] elongation factor G, GFM and Gfm, and mapping of GFM to human chromosome

[72211] 3q25.1-q26.2. Genomics 74: 109-114, 2001.

[72212] 12258.

[72213] Miura, K.; Jacques, K. M.; Stauffer, S.; Kubosaki, A.; Zhu, K.;

[72214] Hirsch, D. S.; Resau, J.; Zheng, Y.; Randazzo, P. A.: ARAP1: a point

[72215] of convergence for Arf and Rho signaling. Molec. Cell 9: 109-119,

[72216] 2002.

[72217] 12259.

[72218] Hicar, M. D.; Liu, Y.; Allen, C. E.; Wu, L.-C.: Structure of the

[72219] human zinc finger protein HIVEP3: molecular cloning, expression, exon-intron

[72220] structure, and comparison with paralogous genes HIVEP1 and HIVEP2. Genomics 71:

[72221] 89-100, 2001.

[72222] 12260.

[72223] Alderuccio, F.; Chan, E. K. L.; Tan, E. M.: Molecular characterization

[72224] of an autoantigen of PM-Scl in the polymyositis/scleroderma overlap

[72225] syndrome: a unique and complete human cDNA encoding an apparent 75-kD

[72226] acidic protein of the nucleolar complex. J. Exp. Med. 173: 941-952,

[72227] 1991.

[72228] 12261.

[72229] Bates, E. E. M.; Fridman, W. H.; Mueller, C. G. F.: The ADAMDEC1

[72230] (decysin) gene structure: evolution by duplication in a metalloprotease

[72231] gene cluster on chromosome 8p12. Immunogenetics 54: 96-105, 2002.

[72232] 12262.

[72233] Bridges, L. C.; Tani, P. H.; Hanson, K. R.; Roberts, C. M.; Judkins,

[72234] M. B.; Bowditch, R. D.: The lymphocyte metalloprotease MDC-L (ADAM28)

[72235] is a ligand for the integrin alpha-4/beta-1. J. Biol. Chem. 277:

[72236] 3784-3792, 2002.

[72237] 12263.

[72238] Jury, J. A.; Perry, A. C. F.; Hall, L.: Identification, sequence

[72239] analysis and expression of transcripts encoding a putative metalloproteinase,

[72240] eMDC II, in human and macaque epididymis. Molec. Hum. Reprod. 5:

[72241] 1127-1134, 1999.

[72242] 12264.

[72243] Roberts, C. M.; Tani, P. H.; Bridges, L. C.; Laszik, Z.;

Bowditch,

[72244] R. D.: MDC-L, a novel metalloprotease disintegrin cysteine-rich protein

[72245] family member expressed by human lymphocytes. J. Biol. Chem. 274:

[72246] 29251-29259, 1999.

[72247] 12265.

[72248] Kolle, G.; Georgas, K.; Holmes, G. P.; Little, M. H.; Yamada, T.

[72249] : CRIM1, a novel gene encoding a cysteine-rich repeat protein, is

[72250] developmentally regulated and implicated in vertebrate CNS development

[72251] and organogenesis. Mech. Dev. 90: 181-193, 2000.

[72252] 12266.

[72253] Shibui, A.; Tsunoda, T.; Seki, N.; Suzuki, Y.; Sugano, S.; Sugane,

[72254] K.: Cloning, expression analysis, and chromosomal localization of

[72255] a novel butyrophilin-like receptor. J. Hum. Genet. 44: 249-252,

[72256] 1999.

[72257] 12267.

[72258] Lee, A.; Beck, L.; Markovich, D.: The human renal sodium sulfate

[72259] cotransporter (SLC13A1; hNaSi-1) cDNA and gene: organization, chromosomal

[72260] localization, and functional characterization. Genomics 70: 354-363,

[72261] 2000.

[72262] 12268.

[72263] Fitzgerald, K. A.; Palsson-McDermott, E. M.; Bowie, A. G.; Jefferies,

[72264] C. A.; Mansell, A. S.; Brady, G.; Brint, E.; Dunne, A.; Gray, P.;

[72265] Harte, M. T.; McMurray, D.; Smith, D. E.; Sims, J. E.; Bird, T. A.;

[72266] O'Neill, L. A. J.: Mal (MyD88-adapter-like) is required for Toll-like

[72267] receptor-4 signal transduction. Nature 413: 78-83, 2001.

[72268] 12269.

[72269] Horng, T.; Barton, G. M.; Medzhitov, R.: TIRAP: an adapter molecule

[72270] in the Toll signaling pathway. Nature Immun. 2: 835-841, 2001.

[72271] 12270.

[72272] Dose, A. C.; Burnside, B.: Cloning and chromosomal local-ization

[72273] of a human class III myosin. Genomics 67: 333-342, 2000.

[72274] 12271.

[72275] Walsh, T.; Walsh, V.; Vreugde, S.; Hertzano, R.; Shahin, H.; Haika,

[72276] S.; Lee, M. K.; Kanaan, M.; King, M.-C.; Avraham, K. B.: From flies'

[72277] eyes to our ears: mutations in a human class III myosin cause progressive

[72278] nonsyndromic hearing loss DFNB30. Proc. Nat. Acad. Sci. 99: 7518-7523,

[72279] 2002.

[72280] 12272.

[72281] Sohda, M.; Misumi, Y.; Yamamoto, A.; Yano, A.; Nakamura, N.; Ikehara,

[72282] Y.: Identification and characterization of a novel Golgi protein,

[72283] GCP60, that interacts with the integral membrane protein giantin. J.

[72284] Biol. Chem. 276: 45298-45306, 2001.

[72285] 12273.

[72286] Baron, M.: Genetics of schizophrenia and the new millennium: progress

[72287] and pitfalls. Am. J. Hum. Genet. 68: 299-312, 2001.

[72288] 12274.

[72289] Chakravarti, A.: A compelling genetic hypothesis for a complex

[72290] disease: PRODH2/DGCR6 variation leads to schizophrenia susceptibility. Proc.

[72291] Nat. Acad. Sci. 99: 4755-4756, 2002.

[72292] 12275.

[72293] Gogos, J. A.; Santha, M.; Takacs, Z.; Beck, K. D.; Luine, V.; Lucas,

[72294] L. R.; Nadler, J. V.; Karayiorgou, M.: The gene encoding proline

[72295] dehydrogenase modulates sensorimotor gating in mice. Nature Genet. 21:

[72296] 434-439, 1999.

[72297] 12276.

[72298] Liu, H.; Heath, S. C.; Sobin, C.; Roos, J. L.; Galke, B. L.; Blundell,

[72299] M. L.; Lenane, M.; Robertson, B.; Wijsman, E. M.; Rapoport, J. L.;

[72300] Gogos, J. A.; Karayiorgou, M.: Genetic variation at the 22q11 PRODH2/DGCR6

[72301] locus presents an unusual pattern and increases susceptibility to

[72302] schizophrenia. Proc. Nat. Acad. Sci. 99: 3717-3722, 2002.

[72303] 12277.

[72304] Scott, A.: Personal Communication. Baltimore, Md. 4/2002.

[72305] 12278.

[72306] Doege, H.; Bocianski, A.; Joost, H.-G.; Schurmann, A.: Activity

[72307] and genomic organization of human glucose transporter 9 (GLUT9), a

[72308] novel member of the family of sugar-transport facilitators predominantly

[72309] expressed in brain and leucocytes. Biochem. J. 350: 771-776, 2000.

[72310] 12279.

[72311] Scott, A. F.: Personal Communication. Baltimore, Md. 4/1/2002.

[72312] 12280.

[72313] Leo, C. P.; Hsu, S. Y.; McGee, E. A.; Salanova, M.; Hsueh, A. J.

[72314] W.: DEFT, a novel death effector domain-containing molecule predominantly

[72315] expressed in testicular germ cells. Endocrinology 139: 4839-4848,

[72316] 1998.

[72317] 12281.

[72318] Stegh, A. H.; Schickling, O.; Ehret, A.; Scaffidi, C.; Peter-hansel,

[72319] C.; Hofmann, T. G.; Grummt, I.; Krammer, P. H.; Peter, M. E.: DEDD,

[72320] a novel death effector domain-containing protein, tar-geted to the

[72321] nucleolus. EMBO J. 17: 5974-5986, 1998.

[72322] 12282.

[72323] Dixon, J.; Hovanes, K.; Shiang, R.; Dixon, M. J.: Sequence

analysis,
[72324] identification of evolutionary conserved motifs and ex-
pression analysis
[72325] of murine tcof1 provide further evidence for a potential
function
[72326] for the gene and its human homologue, TCOF1. Hum.
Molec. Genet. 6:
[72327] 727-737, 1997.

[72328] 12283.

[72329] Dixon, M. J.; Read, A. P.; Donnai, D.; Colley, A.; Dixon, J.;
Williamson,
[72330] R.: The gene for Treacher Collins syndrome maps to the
long arm of
[72331] chromosome 5. Am. J. Hum. Genet. 49: 17-22, 1991.

[72332] 12284.

[72333] Edwards, S. J.; Gladwin, A. J.; Dixon, M. J.: The mutational
spectrum
[72334] in Treacher Collins syndrome reveals a predominance of
mutations that
[72335] create a premature-termination codon. Am. J. Hum.
Genet. 60: 515-524,
[72336] 1997.

[72337] 12285.

[72338] Cabart, P.; Murphy, S.: BRFU, a TFIIB-like factor, is directly

[72339] recruited to the TATA-box of polymerase III small nuclear RNA gene

[72340] promoters through its interaction with TATA-binding protein. J. Biol.

[72341] Chem. 276: 43056-43064, 2001.

[72342] 12286.

[72343] Schramm, L.; Pendergrast, P. S.; Sun, Y.; Hernandez, N.: Different

[72344] human TFIIIB activities direct RNA polymerase III transcription from

[72345] TATA-containing and TATA-less promoters. Genes Dev. 14: 2650-2663,

[72346] 2000.

[72347] 12287.

[72348] Soccio, R. E.; Adams, R. M.; Romanowski, M. J.; Sehayek, E.; Burley,

[72349] S. K.; Breslow, J. L.: The cholesterol-regulated StarD4 gene encodes

[72350] a StAR-related lipid transfer protein with two closely re-

lated homologues,
[72351] StarD5 and StarD6. Proc. Nat. Acad. Sci. 99: 6943-6948, 2002.
[72352] 12288.

[72353] Schenck, A.; Bardoni, B.; Moro, A.; Bagni, C.; Mandel, J. L.:

[72354] A highly conserved protein family interacting with the fragile X mental
[72355] retardation protein (FMRP) and displaying selective interactions with
[72356] FMRP-related proteins FXR1P and FXR2P. Proc. Nat. Acad. Sci. 98:
[72357] 8844-8849, 2001.

[72358] 12289.

[72359] Dong, J.; Gu, T. T.; Simmons, D; MacDougall, M.: Enamelin maps
[72360] to human chromosome 4q21 within the autosomal dominant amelogenesis
[72361] imperfecta locus. Europ. J. Oral. Sci. 108: 353-358, 2000.

[72362] 12290.

[72363] Oukka, M.; Kim, S. T.; Lugo, G.; Sun, J.; Wu, L.-C.; Glimcher,

[72364] L. H.: A mammalian homolog of Drosophila schnurri, KRC, regulates

[72365] TNF receptor-driven responses and interacts with TRAF2. Molec. Cell 9:

[72366] 121-131, 2002.

[72367] 12291.

[72368] Walder, R. Y.; Landau, D.; Meyer, P.; Shalev, H.; Tsolia, M.; Borochowitz,

[72369] Z.; Boettger, M. B.; Beck, G. E.; Englehardt, R. K.; Carmi, R.; Sheffield,

[72370] V. C.: Mutation of TRPM6 causes familial hypomagnesemia with secondary

[72371] hypocalcemia. Nature Genet. 31: 171-174, 2002.

[72372] 12292.

[72373] Liang, J. C.; Chang, K. S.; Schroeder, W.; Siciliano, M.; Trujillo,

[72374] J.; Stass, S.: The human myeloperoxidase gene locates on chromosome

[72375] 17q22-24 and is translocated in acute promyelocytic leukemia. (Abstract) Am.

[72376] J. Hum. Genet. 41: A226, 1987.

[72377] 12293.

[72378] Liang, J. C.; Chang, K. S.; Schroeder, W. T.; Freireich, E. J.;

[72379] Stass, S. A.; Trujillo, J. M.: The myeloperoxidase gene is translocated
[72380] from chromosome 17 to 15 in a patient with acute promyelocytic leukemia. Cancer
[72381] Genet. Cytogenet. 30: 103-107, 1988.

[72382] 12294.

[72383] Miki, T.; Weil, S. C.; Rosner, G. L.; Reid, M. S.; Kidd, K. K.

[72384] : An MPO cDNA clone identifies an RFLP with PstI. Nucleic Acids Res. 16:
[72385] 1649, 1988.

[72386] 12295.

[72387] Morishita, K.; Kubota, N.; Asano, S.; Kaziro, Y.; Nagata, S.:

[72388] Molecular cloning and characterization of cDNA for human myeloperoxidase. J.
[72389] Biol. Chem. 262: 3844-3851, 1987.

[72390] 12296.

[72391] Murao, S.-I.; Stevens, F. J.; Ito, A.; Huberman, E.: Myeloperoxidase:
[72392] a myeloid cell nuclear antigen with DNA-binding properties. Proc.

[72393] Nat. Acad. Sci. 85: 1232-1236, 1988.

[72394] 12297.

[72395] Nauseef, W.; Cogley, M.; McCormick, S.: Effect of the R569W missense

[72396] mutation on the biosynthesis of myeloperoxidase. J. Biol. Chem. 271:

[72397] 9546-9549, 1996.

[72398] 12298.

[72399] Nauseef, W. M.; Olsson, I.; Arnljots, K.: Biosynthesis and processing

[72400] of myeloperoxidase--a marker for myeloid cell differentiation. Europ.

[72401] J. Haemat. 40: 97-110, 1988.

[72402] 12299.

[72403] Reynolds, W. F.; Hiltunen, M.; Pirskanen, M.; Mannermaa, A.; Helisalmi,

[72404] S.; Lehtovirta, M.; Alafuzoff, I.; Soininen, H.: MPO and APOE epsilon-4

[72405] polymorphisms interact to increase risk for AD in Finnish males. Neurology 55:

[72406] 1284-1290, 2000.

[72407] 12300.

[72408] Robinson, T. J.; Morris, D. J.; Ledbetter, D. H.: Chromoso-
mal
[72409] assignment and regional localization of myeloperoxidase
in the mouse. Cytogenet.
[72410] Cell Genet. 53: 83-86, 1990.

[72411] 12301.

[72412] Romano, M.; Dri, P.; Dadalt, L.; Patriarca, P.; Baralle, F. E.

[72413] : Biochemical and molecular characterization of hereditary
myeloproliferative
[72414] deficiency. Blood 90: 4126-4134, 1997.

[72415] 12302.

[72416] van Tuinen, P.; Johnson, K. R.; Ledbetter, S. A.; Nussbaum,
R.
[72417] L.; Rovera, G.; Ledbetter, D. H.: Localization of myeloper-
oxidase
[72418] to the long arm of human chromosome 17: relationship to
the 15;17
[72419] translocation of acute promyelocytic leukemia. Oncogene
1: 319-322,
[72420] 1987.

[72421] 12303.

[72422] Weil, S. C.; Rosner, G. L.; Reid, M. S.; Chisholm, R. L.; Far-
ber,
[72423] N. M.; Spitznagel, J. K.; Swanson, M. S.: cDNA cloning of
human myeloperoxidase:
[72424] decrease in myeloperoxidase mRNA upon induction of
HL-60 cells. Proc.
[72425] Nat. Acad. Sci. 84: 2057-2061, 1987.

[72426] 12304.

[72427] Weil, S. C.; Rosner, G. L.; Reid, M. S.; Chisholm, R. L.;
Lemons,
[72428] R. S.; Swanson, M. S.; Carrino, J. J.; Diaz, M. O.; Le Beau,
M. M.
[72429] : Translocation and rearrangement of myeloperoxidase
gene in acute
[72430] promyelocytic leukemia. Science 240: 790-792, 1988.

[72431] 12305.

[72432] Yamada, M.; Hur, S.-J.; Hashinaka, K.; Tsuneoka, K.; Saeki,
T.;
[72433] Nishio, C.; Sakiyama, F.; Tsunasawa, S.: Isolation and
characterization

[72434] of a cDNA coding for human myeloperoxidase. Arch. Biochem. Biophys. 255:
[72435] 147-155, 1987.

[72436] 12306.

[72437] Zaki, S. R.; Austin, G. E.; Chan, W. C.; Conaty, A. L.; Trusler,
[72438] S.; Trappier, S.; Lindsey, R. B.; Swan, D. C.: Chromosomal localization
[72439] of the human myeloperoxidase gene by in situ hybridiza-tion using oligonucleotide
[72440] probes. Genes Chromosomes Cancer 2: 266-270, 1990.

[72441] 12307.

[72442] Saiardi, A.; Erdjument-Bromage, H.; Snowman, A. M.; Tempst, P.;
[72443] Snyder, S. H.: Synthesis of diphosphoinositol pentak-isphosphate by
[72444] a newly identified family of higher inositol polyphosphate kinases. Curr.
[72445] Biol. 9: 1323-1326, 1999.

[72446] 12308.

[72447] Saiardi, A.; Nagata, E.; Luo, H. R.; Snowman, A. M.; Snyder,

S.

[72448] H.: Identification and characterization of a novel inositol hexakisphosphate

[72449] kinase. J. Biol. Chem. 276: 39179-39185, 2001.

[72450] 12309.

[72451] Morrison, B. H.; Bauer, J. A.; Hu, J.; Grane, R. W.; Ozdemir, A.

[72452] M.; Chawla-Sarkar, M.; Gong, B.; Almasan, A.; Kalvakolanu, D. V.;

[72453] Lindner, D. J.: Inositol hexakisphosphate kinase 2 sensitizes ovarian

[72454] carcinoma cells to multiple cancer therapeutics. Oncogene 21: 1882-1889,

[72455] 2002.

[72456] 12310.

[72457] Mole, S. E.; Mitchison, H. M.; Munroe, P. B.: Molecular basis

[72458] of the neuronal ceroid lipofuscinoses: mutations in CLN1, CLN2, CLN3,

[72459] and CLN5. Hum. Mutat. 14: 199-215, 1999.

[72460] 12311.

[72461] Andermann, E.; Jakob, J. S.; Andermann, F.; Carpenter, S.; Wolfe,

[72462] L.; Berkovic, S. F.: The Newfoundland aggregate of neuronal ceroid-lipofuscinosis. Am.

[72463] J. Med. Genet. Suppl. 5: 111-116, 1988.

[72464] 12312.

[72465] Andermann, E.; Scriver, C. R.; Wolfe, L. S.; Dansky, L.; Andermann,

[72466] F.: Genetic variants of Tay-Sachs disease and Sandhoff's disease

[72467] in French Canadians, juvenile Tay-Sachs disease in Lebanese Canadians,

[72468] and a Tay-Sachs screening program in the French Canadian population.In:

[72469] Kaback, M. M.; Rimoin, D. L.; O'Brien, J. S.: Tay-Sachs Disease:

[72470] Screening and Prevention. New York: Alan R. Liss (pub.) 1977.

[72471] 12313.

[72472] Berry-Kravis, E.; Sleat, D. E.; Sohar, I.; Meyer, P.; Donnelly,

[72473] R.; Lobel, P.: Prenatal testing for late infantile neuronal ceroid

[72474] lipofuscinosis. Ann. Neurol. 47: 254-257, 2000.

[72475] 12314.

[72476] Brodsky, M. C.; Baker, R. S.; Hamed, L. M.: Pediatric Neuro-Ophthalmology. New

[72477] York: Springer-Verlag , 1996. Pp. 374.

[72478] 12315.

[72479] Donahue, S.; Zeman, W.; Watanabe, I.: Electron micro-scopic observations

[72480] in Batten's disease.In: Aronson, S. M.; Volk, B. W.: Inborn Disorders

[72481] of Sphingolipid Metabolism. Oxford: Pergamon Press (pub.) 1967.

[72482] Pp. 3-22.

[72483] 12316.

[72484] Gonatas, N. K.; Gambetti, P.; Baird, H.: A second type of late

[72485] infantile amaurotic idiocy with multilamellar cytosomes. J. Neuropath.

[72486] Exp. Neurol. 27: 371-389, 1968.

[72487] 12317.

[72488] Komp, D. M.: Dr. Imperato-McGinley. (Letter) Lancet II: 262

[72489] only, 1979.

[72490] 12318.

[72491] Leshin, M.; Griffin, J. E.; Wilson, J. D.: 5-Alpha-reductase

[72492] deficiency: evidence for genetic heterogeneity. (Abstract) Clin.

[72493] Res. 26: 47A only, 1978.

[72494] 12319.

[72495] Leshin, M.; Griffin, J. E.; Wilson, J. D.: Hereditary male pseudohermaphroditism

[72496] associated with an unstable form of 5-alpha-reductase. J. Clin. Invest. 62:

[72497] 685-691, 1978.

[72498] 12320.

[72499] Makridakis, N. M.; Ross, R. K.; Pike, M. C.; Crocitto, L. E.;

[72500] Kolonel, L. N.; Pearce, C. L.; Henderson, B. E.; Reichardt, J. K.

[72501] V.: Association of mis-sense substitution in SRD5A2 gene with prostate

[72502] cancer in African-American and Hispanic men in Los Angeles, USA. Lancet 354:
[72503] 975-978, 1999.

[72504] 12321.

[72505] Moore, R. J.; Griffin, J. E.; Wilson, J. D.: Diminished 5-alpha-reductase
[72506] activity in extracts of fibroblasts cultured from patients with familial
[72507] incomplete male pseudohermaphroditism, type 2. J. Biol. Chem. 250:
[72508] 7168-7172, 1975.

[72509] 12322.

[72510] Balkan, W.; Oates, E. L.; Howard, G. A.; Roos, B. A.: Testes exhibit
[72511] elevated expression of calcitonin gene-related peptide receptor component
[72512] protein. Endocrinology 140: 1459-1469, 1999.

[72513] 12323.

[72514] Evans, B. N.; Rosenblatt, M. I.; Mnayer, L. O.; Oliver, K. R.;

[72515] Dickerson, I. M.: CGRP-RCP, a novel protein required for signal transduction

[72516] at calcitonin gene-related peptide and adrenomedullin receptors. J.
[72517] Biol. Chem. 275: 31438-31443, 2000.

[72518] 12324.

[72519] Guo, B.; Godzik, A.; Reed, J. C.: Bcl-G, a novel pro-apoptotic
[72520] member of the Bcl-2 family. J. Biol. Chem. 276: 2780-2785, 2001.
[72521] 12325.

[72522] Montpetit, A.; Boily, G.; Sinnett, D.: A detailed transcriptional
[72523] map of the chromosome 12p12 tumour suppressor locus. Europ. J. Hum.
[72524] Genet. 10: 62-71, 2002.

[72525] 12326.

[72526] Katoh, M.: Molecular cloning and characterization of RNF26 on
[72527] human chromosome 11q23 region, encoding a novel RING finger protein
[72528] with leucine zipper. Biochem. Biophys. Res. Comm. 282: 1038-1044,

[72529] 2001.

[72530] 12327.

[72531] Nishida, K.; Yoshida, Y.; Itoh, M.; Fukada, T.; Ohtani, T.; Shirogane,

[72532] T.; Atsumi, T.; Takahashi-Tezuka, M.; Ishihara, K.; Hibi, M.; Hirano,

[72533] T.: Gab-family adapter proteins act downstream of cytokine and growth

[72534] factor receptors and T- and B-cell antigen receptors. Blood 93:

[72535] 1809-1816, 1999.

[72536] 12328.

[72537] Zhao, C.; Yu, D.-H.; Shen, R.; Feng, G.-S.: Gab2, a new pleckstrin

[72538] homology domain-containing adapter protein, acts to uncouple signaling

[72539] from ERK kinase to Elk-1. J. Biol. Chem. 274: 19649-19654, 1999.

[72540] 12329.

[72541] Ge, K.; Xue, A.; Bai, J.; Wang, S.: Keshan disease-an endemic

[72542] cardiomyopathy in China. Virchows Arch. 401: 1-15, 1983.

[72543] 12330.

[72544] Lescure, A.; Gautheret, D.; Carbon, P.; Krol, A.: Novel selenoproteins

[72545] identified in silico and in vivo by using a conserved RNA structural

[72546] motif. J. Biol. Chem. 274: 38147-38154, 1999.

[72547] 12331.

[72548] Frye, R. A.: Phylogenetic classification of prokaryotic and eukaryotic

[72549] Sir2-like proteins. Biochem. Biophys. Res. Commun. 273: 793-798,

[72550] 2000.

[72551] 12332.

[72552] Berghs, S.; Aggujaro, D.; Dirkx, R., Jr.; Maksimova, E.; Stabach,

[72553] P.; Hermel, J.-M.; Zhang, J.-P.; Philbrick, W.; Slepnev, V.; Ort,

[72554] T.; Slimena, M.: Beta-IV spectrin, a new spectrin localized at axon

[72555] initial segments and nodes of Ranvier in the central and peripheral
[72556] nervous system. J. Cell Biol. 151: 985-1001, 2000.

[72557] 12333.

[72558] Bock, G. R.; Frank, M. P.; Steel, K. P.; Deol, M. S.: The quivering
[72559] mutant mouse: hereditary deafness of central origin. Acta Otolaryng. 96:
[72560] 371-377, 1983.

[72561] 12334.

[72562] Deol, M. S.; Frank, M. P.; Steel, K. P.; Bock, G. R.: Genetic

[72563] deafness of central origin. Brain Res. 258: 177-179, 1983.

[72564] 12335.

[72565] Parkinson, N. J.; Olsson, C. L.; Hallows, J. L.; McKee-Johnson,
[72566] J.; Keogh, B. P.; Noben-Trauth, K.; Kujawa, S. G.; Tempel, B. L.:
[72567] Mutant beta-spectrin 4 causes auditory and motor neuropathies in quivering
[72568] mice. Nature Genet. 29: 61-65, 2001.

[72569] 12336.

[72570] Tse, W. T.; Tang, J.; Jin, O.; Korsgren, C.; John, K. M.; Kung,
[72571] A. L.; Gwynn, B.; Peters, L. L.; Lux, S. E.: A new spectrin, beta-IV,
[72572] has a major truncated isoform that associates with promyelocytic leukemia
[72573] protein nuclear bodies and the nuclear matrix. J. Biol. Chem. 276:
[72574] 23974-23985, 2001.

[72575] 12337.

[72576] Yoon, C. H.; Les, E. P.: Quivering, a new first chromosome mutation
[72577] in mice. J. Hered. 48: 176-180, 1957.

[72578] 12338.

[72579] Saitoh, T.; Katoh, M.: Molecular cloning and characterization
[72580] of human WNT5B on chromosome 12p13.3 region. Int. J. Oncol. 19:
[72581] 347-351, 2001.

[72582] 12339.

[72583] Yousef, G. M.; Diamandis, M.; Jung, K.; Diamandis, E. P.: Molecular
[72584] cloning of a novel human acid phosphatase gene (ACPT) that is highly
[72585] expressed in the testis. Genomics 385-395, 2001.

[72586] 12340.

[72587] Miyazaki, K.; Matsuda, S.; Ichigotani, Y.; Takenouchi, Y.; Hayashi,
[72588] K.; Fukuda, Y.; Nimura, Y.; Hamaguchi, M.: Isolation and characterization
[72589] of a novel human gene (NESH) which encodes a putative signaling molecule
[72590] similar to e3B1 protein. Biochim. Biophys. Acta 1493: 237-241, 2000.

[72591] 12341.

[72592] Gomez-Fabre, P. M.; Aledo, J. C.; del Castillo-Olivares, A.; Alonso,
[72593] F. J.; Nunez de Castro, I.; Campos, J. A.; Marquez, J.: Molecular
[72594] cloning, sequencing and expression studies of the human breast cancer
[72595] cell glutaminase. Biochem. J. 345: 365-375, 2000.

[72596] 12342.

[72597] Nosaka, K.; Onozuka, M.; Kakazu, N.; Hibi, S.; Nishimura, H.; Nishino,

[72598] H.; Abe, T.: Isolation and characterization of a human thiamine pyrophosphokinase

[72599] cDNA. Biochem. Biophys. Acta 1517: 293-297, 2001.

[72600] 12343.

[72601] Nosaka, K.; Onozuka, M.; Nishino, H.; Nishimura, H.; Kawasaki,

[72602] Y.; Ueyama, H.: Molecular cloning and expression of a mouse thiamin

[72603] pyrophosphokinase cDNA. J. Biol. Chem. 274: 34129-34133, 1999.

[72604] 12344.

[72605] Zhao, R.; Gao, F.; Goldman, I. D.: Molecular cloning of human

[72606] thiamin pyrophosphokinase. Biochim. Biophys. Acta 1517: 320-322,

[72607] 2001.

[72608] 12345.

[72609] Gaire, M.; Chatton, B.; Kedinger, C.: Isolation and charac-

terization

[72610] of two novel, closely related ATF cDNA clones from HeLa cells. Nucleic

[72611] Acids Res. 18: 3467-3473, 1990.

[72612] 12346.

[72613] Lustig, L. R.; Peng, H.; Hiel, H.; Yamamoto, T.; Fuchs, P. A.:

[72614] Molecular cloning and mapping of the human nicotinic acetylcholine

[72615] receptor alpha-10 (CHRNA10). Genomics 73: 272-283, 2001.

[72616] 12347.

[72617] Leader, B.; Leder, P.: Formin-2, a novel formin homology protein

[72618] of the cappuccino subfamily, is highly expressed in the developing

[72619] and adult central nervous system. Mech. Dev. 93: 221-231, 2000.

[72620] 12348.

[72621] Mitsumoto, Y.; Oka, S.; Sakuma, H.; Inazawa, J.; Kawasaki, T.:

[72622] Cloning and chromosomal mapping of human glucuronyl-

transferase involved
[72623] in biosynthesis of the HNK-1 carbohydrate epitope. Genomics 65:
[72624] 166-173, 2000.

[72625] 12349.

[72626] Gilley, J.; Fried, M.: Extensive gene order differences within
[72627] regions of conserved synteny between the Fugu and human genomes: implications
[72628] for chromosomal evolution and the cloning of disease genes. Hum.
[72629] Molec. Genet. 8: 1313-1320, 1999.

[72630] 12350.

[72631] Harduin-Lepers, A.; Stokes, D. C.; Steelant, W. F. A.; Samyn-Petit,
[72632] B.; Krzewinski-Recchi, M.-A.; Vallejo-Ruiz, V.; Zanetta, J.-P.; Auge,
[72633] C.; Delannoy, P.: Cloning, expression and gene organization of a
[72634] human Neu5Ac-alpha-2-3Gal-beta-1-3GalNAc alpha-2,6-sialyltransferase:
[72635] hST6GalNAc IV. Biochem. J. 352: 37-48, 2000.

[72636] 12351.

[72637] Bellorini, M.; Zemzoumi, K.; Farina, A.; Berthelsen, J.; Piaggio,

[72638] G.; Mantovani, R.: Cloning and expression of human NF-YC. Gene 197:

[72639] 119-125, 1997.

[72640] 12352.

[72641] Dmitrenko, V.; Garifulin, O.; Kavsan, V.: Isolation and sequence

[72642] analysis of the cDNA encoding subunit C of human CCAAT-binding transcription

[72643] factor. Gene 197: 161-163, 1997.

[72644] 12353.

[72645] Nakshatri, H.; Bhat-Nakshatri, P.; Currie, R. A.: Subunit association

[72646] and DNA binding activity of the heterotrimeric transcription factor

[72647] NF-Y is regulated by cellular redox. J. Biol. Chem. 271: 28784-28791,

[72648] 1996.

[72649] 12354.

[72650] Scott, A. F.: Personal Communication. Baltimore, Md. 10/16/2000.

[72651] 12355.

[72652] Sinha, S.; Maity, S. N.; Seldin, M. F.; de Crombrugghe, B.: Chromosomal

[72653] assignment and tissue expression of CBF-C/NFY-C, the third subunit

[72654] of the mammalian CCAAT-binding factor. Genomics 37: 260-263, 1996.

[72655] 12356.

[72656] Reboul, J.; Gardiner, K.; Monneron, D.; Uze, G.; Lutfalla, G.:

[72657] Comparative genomic analysis of the interferon/inter- leukin-10 receptor

[72658] gene cluster. Genome Res. 9: 242-250, 1999.

[72659] 12357.

[72660] Chen, Q.; Ghilardi, N.; Wang, H.; Baker, T.; Xie, M.-H.; Gurney,

[72661] A.; Grewal, I. S.; de Sauvage, F. J.: Development of Th1-type immune

[72662] responses requires the type 1 cytokine receptor TCCR. Nature 407:

[72663] 916-920, 2000.

[72664] 12358.

[72665] Sprecher, C. A.; Grant, F. J.; Baumgartner, J. W.; Presnell, S.

[72666] R.; Schrader, S. K.; Yamagiwa, T.; Whitmore, T. E.; O'Hara, P. J.;

[72667] Foster, D. F.: Cloning and characterization of a novel class I cytokine

[72668] receptor. Biochem. Biophys. Res. Comm. 246: 82-90, 1998.

[72669] 12359.

[72670] Yoshida, H.; Hamano, S.; Senaldi, G.; Covey, T.; Faggioni, R.;

[72671] Mu, S.; Xia, M.; Wakeham, A. C.; Nishina, H.; Potter, J.; Saris, C.

[72672] J. M.; Mak, T. W.: WSX-1 is required for the initiation of Th1 responses

[72673] and resistance to L. major infection. Immunity 15: 569-578, 2001.

[72674] 12360.

[72675] Autieri, M. V.; Carbone, C. J.: 14-3-3-Gamma interacts with and

[72676] is phosphorylated by multiple protein kinase C isoforms in PDGF-stimulated

[72677] human vascular smooth muscle cells. DNA Cell Biol. 18: 555-564,

[72678] 1999.

[72679] 12361.

[72680] Yeh, W.-C.; de la Pompa, J. L.; McCurrach, M. E.; Shu, H.-B.; Elia,

[72681] A. J.; Shahinian, A.; Ng, M.; Wakeham, A.; Khoo, W.; Mitchell, K.;

[72682] El-Deiry, W. S.; Lowe, S. W.; Goeddel, D. V.; Mak, T. W.: FADD: essential

[72683] for embryo development and signaling from some, but not all, inducers

[72684] of apoptosis. Science 279: 1954-1958, 1998.

[72685] 12362.

[72686] Zhang, J.; Cado, D.; Chen, A.; Kabra, N. H.; Winoto, A.: Fas-mediated

[72687] apoptosis and activation-induced T-cell proliferation are defective

[72688] in mice lacking FADD/Mort1. Nature 392: 296-300, 1998.

[72689] 12363.

[72690] Hampe, W.; Rezgaoui, M.; Hermans-Borgmeyer, I.; Schaller, H. C.

[72691] : The genes for the human VPS10 domain-containing receptors are large

[72692] and contain many small exons. Hum. Genet. 108: 529-536, 2001.

[72693] 12364.

[72694] Torres, R.; Polymeropoulos, M. H.: Genomic organization and localization

[72695] of the human CRMP-1 gene. DNA Res. 5: 393-395, 1998.

[72696] 12365.

[72697] Koyama, K.; Sudo, K.; Nakamura, Y.: Isolation of 115 human chromosome

[72698] 8-specific expressed-sequence tags by exon amplification. Genomics 26:

[72699] 245-253, 1995.

[72700] 12366.

[72701] Woitach, J. T.; Hong, R.; Keck, C. L.; Zimonjic, D. B.; Popescu,

[72702] N. C.; Thorgeirsson, S. S.: Assignment of the Bog gene

(RBBP9) to

[72703] syntenic regions of mouse chromosome 2G1-H1 and human chromosome 20p11.2

[72704] by fluorescence in situ hybridization. Cytogenet. Cell Genet. 85:

[72705] 252-253, 1999.

[72706] 12367.

[72707] Woitach, J. T.; Zhang, M.; Niu, C.-H.; Thorgeirsson, S. S.: A

[72708] retinoblastoma-binding protein that affects cell-cycle control and

[72709] confers transforming ability. Nature Genet. 19: 371-374, 1998.

[72710] 12368.

[72711] Yokota, J.; Akiyama, T.; Fung, Y.-K. T.; Benedict, W. F.; Namba,

[72712] Y.; Hanaoka, M.; Wada, M.; Terasaki, T.; Shimosato, Y.; Sugimura,

[72713] T.; Terada, M.: Altered expression of the retinoblastoma (RB) gene

[72714] in small-cell carcinoma of the lung. Oncogene 3:

471-475, 1988.

[72715] 12369.

[72716] Katahira, J.; Sugiyama, H.; Inoue, N.; Horiguchi, Y.; Matsuda,
[72717] M.; Sugimoto, N.: Clostridium perfringens enterotoxin utilizes two
[72718] structurally related membrane proteins as functional receptors in
[72719] vivo. J. Biol. Chem. 272: 26652-26658, 1997.

[72720] 12370.

[72721] Furuse, M.; Fujita, K.; Hiiragi, T.; Fujimoto, K.; Tsukita, S.

[72722] : Claudin-1 and -2: novel integral membrane proteins localizing at
[72723] tight junctions with no sequence similarity to occludin. J. Cell
[72724] Biol. 141: 1539-1550, 1998.

[72725] 12371.

[72726] Peacock, R. E.; Keen, T. J.; Inglehearn, C. F.: Analysis of a

[72727] human gene homologous to rat ventral prostate.1 protein. Genomics 46:
[72728] 443-449, 1997.

[72729] 12372.

[72730] Biffo, S.; Sanvito, F.; Costa, S.; Preve, L.; Pignatelli, R.; Spinardi,

[72731] L.; Marchisio, P. C.: Isolation of a novel beta-4 integrin-binding

[72732] protein (p27BBP) highly expressed in epithelial cells. J. Biol. Chem. 272:

[72733] 30314-30321, 1997.

[72734] 12373.

[72735] Sanvito, F.; Arrigo, G.; Zuffardi, O.; Agnelli, M.; Marchisio,

[72736] P. C.; Biffo, S.: Localization of p27 beta-4 binding protein gene

[72737] (ITGB4BP) to human chromosome region 20q11.2. Genomics 52: 111-112,

[72738] 1998.

[72739] 12374.

[72740] Si, K.; Chaudhuri, J.; Chevesich, J.; Maitra, U.: Molecular cloning

[72741] and functional expression of a human cDNA encoding translation initiation

[72742] factor 6. Proc. Nat. Acad. Sci. 94: 14285-14290, 1997.

[72743] 12375.

[72744] Nuber, U.; Schwarz, S.; Kaiser, P.; Schneider, R.; Scheffner, M.

[72745] : Cloning of human ubiquitin-conjugating enzymes UbcH6 and UbcH7 (E2-F1)

[72746] and characterization of their interaction with E6-AP and RSP5. J.

[72747] Biol. Chem. 271: 2795-2800, 1996.

[72748] 12376.

[72749] Fuentes, J. J.; Pritchard, M. A.; Estivill, X.: Genomic organi-zation,

[72750] alternative splicing, and expression patterns of the DSCR1 (Down syndrome

[72751] candidate region 1) gene. Genomics 44: 358-361, 1997.

[72752] 12377.

[72753] Kingsbury, T. J.; Cunningham, K. W.: A conserved family of calcineurin

[72754] regulators. Genes Dev. 14: 1595-1604, 2000.

[72755] 12378.

[72756] Rothermel, B.; Vega, R. B.; Yang, J.; Wu, H.; Bassel-Duby, R.;

[72757] Williams, R. S.: A protein encoded within the Down syn-

drome critical
[72758] region is enriched in striated muscles and inhibits calcineurin signaling. J.
[72759] Biol. Chem. 275: 8719-8725, 2000.

[72760] 12379.

[72761] Pata, I.; Tensing, K.; Metspalu, A.: A human cDNA encoding the
[72762] homologue of NADH:ubiquinone oxidoreductase subunit B13. Biochim.
[72763] Biophys. Acta 1350: 115-118, 1997.

[72764] 12380.

[72765] Russell, M. W.; du Manoir, S.; Collins, F. S.; Brody, L. C.: Cloning
[72766] of the human NADH:ubiquinone oxidoreductase subunit B13: localization
[72767] to chromosome 7q32 and identification of a pseudogene on 11p15. Mammalian
[72768] Genome 8: 60-61, 1997.

[72769] 12381.

[72770] Perez Jurado, L. A.; Wang, Y.-K.; Peoples, R.; Coloma, A.; Cruces,

[72771] J.; Francke, U.: A duplicated gene in the breakpoint re-
gions of the
[72772] 7q11.23 Williams-Beuren syndrome deletion encodes the
initiator binding
[72773] protein TFII-I and BAP-135, a phosphorylation target of
BTK. Hum.
[72774] Molec. Genet. 7: 325-334, 1998.

[72775] 12382.

[72776] Yang, W.; Desiderio, S.: BAP-135, a target for Bruton's ty-
rosine
[72777] kinase in response to B cell receptor engagement. Proc.
Nat. Acad.
[72778] Sci. 94: 604-609, 1997.

[72779] 12383.

[72780] Lee, J. W.; Choi, H.-S.; Gyuris, J.; Brent, R.; Moore, D. D.:

[72781] Two classes of proteins dependent on either the presence
or absence
[72782] of thyroid hormone for interaction with the thyroid hor-
mone receptor. Molec.
[72783] Endocr. 9: 243-254, 1995.

[72784] 12384.

[72785] Venturini, L.; You, J.; Stadler, M.; Galien, R.; Lallemand, V.;

[72786] Koken, M. H. M.; Mattei, M. G.; Ganser, A.; Chambon, P.; Losson, R.;

[72787] de The, H.: TIF1-gamma, a novel member of the tran-

scriptional intermediary

[72788] factor 1 family. Oncogene 18: 1209-1217, 1999.

[72789] 12385.

[72790] Lesage, F.; Guillemare, E.; Fink, M.; Duprat, F.; Lazdunski, M.;

[72791] Romey, G.; Barhanin, J.: TWIK-1, a ubiquitous human weakly inward

[72792] rectifying K+ channel with a novel structure. EMBO J. 15: 1004-1011,

[72793] 1996.

[72794] 12386.

[72795] Lesage, F.; Mattei, M.-G.; Fink, M.; Barhanin, J.; Lazdunski, M.

[72796] : Assignment of the human weak inward rectifier K+ channel TWIK-1

[72797] gene to chromosome 1q42-q43. Genomics 34: 153-155, 1996.

[72798] 12387.

[72799] Chew, A.; Buck, E. A.; Peretz, S.; Sirugo, G.; Rinaldo, P.; Isaya,

[72800] G.: Cloning, expression, and chromosomal assignment of the human

[72801] mitochondrial intermediate peptidase gene (MIPEP). Genomics 40:

[72802] 493-496, 1997.

[72803] 12388.

[72804] Chew, A.; Sirugo, G.; Alsobrook, J. P., II; Isaya, G.: Functional

[72805] and genomic analysis of the human mitochondrial intermediate peptidase,

[72806] a putative protein partner of frataxin. Genomics 65: 104-112, 2000.

[72807] 12389.

[72808] Isaya, G.; Kalousek, F.; Rosenberg, L. E.: Sequence analysis of

[72809] rat mitochondrial intermediate peptidase: similarity to zinc metallopeptidases

[72810] and to a putative yeast homologue. Proc. Nat. Acad. Sci. 89: 8317-8321,

[72811] 1992.

[72812] 12390.

[72813] Holzmann, K.; Poltl, A.; Sauermann, G.: A novel spliced transcript

[72814] of human CLAPS2 encoding a protein alternative to clathrin adaptor

[72815] protein AP17. Gene 220: 39-44, 1998.

[72816] 12391.

[72817] Winterpacht, A.; Endele, S.; Enklaar, T.; Fuhry, M.; Zabel, B.

[72818] : Human CLAPS2 encoding AP17, a small chain of the clathrin-associated

[72819] protein complex: cDNA cloning and chromosomal assign- ment to 19q13.2-q13.3. Cytogenet.

[72820] Cell Genet. 75: 132-135, 1996.

[72821] 12392.

[72822] Alderson, M. R.; Smith, C. A.; Tough, T. W.; Davis-Smith, T.; Armitage,

[72823] R. J.; Falk, B.; Roux, E.; Baker, E.; Sutherland, G. R.; Din, W. S.;

[72824] Goodwin, R. G.: Molecular and biological characterization of human

[72825] 4-1BB and its ligand. Europ. J. Immun. 24: 2219-2227, 1994.

[72826] 12393.

[72827] Kwon, B. S.; Weissman, S. M.: cDNA sequences of two inducible

[72828] T-cell genes. Proc. Nat. Acad. Sci. 86: 1963-1967, 1989.

[72829] 12394.

[72830] Loo, D. T.; Chalupny, N. J.; Bajorath, J.; Shuford, W. W.; Mittler,

[72831] R. S.; Aruffo, A.: Analysis of 4-1BBL and laminin binding to murine

[72832] 4-1BB, a member of the tumor necrosis factor receptor superfamily,

[72833] and comparison with human 4-1BB. J. Biol. Chem. 272: 6448-6456,

[72834] 1997.

[72835] 12395.

[72836] Schwarz, H.; Arden, K.; Lotz, M.: CD137, a member of the tumor

[72837] necrosis factor receptor family, is located on chromosome 1p36, in

[72838] a cluster of related genes, and colocalizes with several malignancies. Biochem.

[72839] Biophys. Res. Commun. 235: 699-703, 1997.

[72840] 12396.

[72841] Schwarz, H.; Blanco, F. J.; von Kempis, J.; Valbracht, J.; Lotz,

[72842] M.: ILA, a member of the human nerve growth factor/tu-mor necrosis

[72843] factor receptor family, regulates T-lymphocyte prolifera-tion and survival. Blood 87:

[72844] 2839-2845, 1996.

[72845] 12397.

[72846] Schwarz, H.; Tuckwell, J.; Lotz, M.: A receptor induced by lymphocyte

[72847] activation (ILA): a new member of the human nerve-growth-factor/tumor-necrosis-factor

[72848] receptor family. Gene 134: 295-298, 1993.

[72849] 12398.

[72850] Feuchter-Murthy, A. E.; Freeman, J. D.; Mager, D. L.: Splic-ing

[72851] of a human endogenous retrovirus to a novel phospholi-

pase A2 related
[72852] gene. Nucleic Acids Res. 21: 135-143, 1993.

[72853] 12399.

[72854] Kowalski, P. E.; Freeman, J. D.; Mager, D. L.: Intergenic splicing
[72855] between a HERV-H endogenous retrovirus and two adjacent human genes. Genomics 57:
[72856] 371-379, 1999.

[72857] 12400.

[72858] Okano, M.; Xie, S.; Li, E.: Cloning and characterization of a
[72859] family of novel mammalian DNA (cytosine-5) methyltransferases. (Letter) Nature
[72860] Genet. 19: 219-220, 1998.

[72861] 12401.

[72862] Robertson, K. D.; Uzvolgyi, E.; Liang, G.; Talmadge, C.; Sumegi,
[72863] J.; Gonzales, F. A.; Jones, P. A.: The human DNA methyltransferases
[72864] (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues
[72865] and overexpression in tumors. Nucleic Acids Res. 27:

2291-2298,

[72866] 1999.

[72867] 12402.

[72868] Xie, S.; Wang, Z.; Okano, M.; Nogami, M.; Li, Y.; He, W.-W.; Okumura,

[72869] K.; Li, E.: Cloning, expression and chromosome locations of the human

[72870] DNMT3 gene family. Gene 236: 87-95, 1999.

[72871] 12403.

[72872] Moghadaszadeh, B.; Petit, N.; Jaillard, C.; Brockington, M.; Roy,

[72873] S. Q.; Merlini, L.; Romero, N.; Estournet, B.; Desguerre, I.; Chaigne,

[72874] D.; Muntoni, F.; Topaloglu, H.; Guicheney, P.: Mutations in SEPN1

[72875] cause congenital muscular dystrophy with spinal rigidity and restrictive

[72876] respiratory syndrome. Nature Genet. 29: 17-18, 2001.

[72877] 12404.

[72878] Jiang, R.; Lan, Y.; Chapman, H. D.; Shawber, C.; Norton, C. R.;

[72879] Serreze, D. V.; Weinmaster, G.; Gridley, T.: Defects in limb, craniofacial,
[72880] and thymic development in Jagged2 mutant mice. Genes Dev. 12: 1046-1057,
[72881] 1998.

[72882] 12405.

[72883] Lan, Y.; Jiang, R.; Shawber, C.; Weinmaster, G.; Gridley, T.:

[72884] The Jagged2 gene maps to chromosome 12 and is a candidate for the
[72885] lgl and sm mutations. Mammalian Genome 8: 875-876, 1997.
[72886] 12406.

[72887] Lanford, P. J.; Lan, Y.; Jiang, R.; Lindsell, C.; Weinmaster, G.;
[72888] Gridley, T.; Kelley, M. W.: Notch signalling pathway mediates hair
[72889] cell development in mammalian cochlea. Nature Genet. 21: 289-292,
[72890] 1999.

[72891] 12407.

[72892] Luo, B.; Aster, J. C.; Hasserjian, R. P.; Kuo, F.; Sklar, J.:

[72893] Isolation and functional analysis of a cDNA for human Jagged2, a gene
[72894] encoding a ligand for the Notch1 receptor. Molec. Cell. Biol. 17:
[72895] 6057-6067, 1997.

[72896] 12408.

[72897] Misaki, Y.; Pruijn, G. J. M.; van der Kemp, A. W. C. M.; van Venrooij,
[72898] W. J.: The 56K autoantigen is identical to human annexin XI. J.
[72899] Biol. Chem. 269: 4240-4246, 1994.

[72900] 12409.

[72901] Morgan, R. O.; Bell, D. W.; Testa, J. R.; Fernandez, M. P.: Genomic
[72902] locations of ANX11 and ANX13 and the evolutionary genetics of human
[72903] annexins. Genomics 48: 100-110, 1998.

[72904] 12410.

[72905] Ono, T.; Kawabe, T.; Sonta, S.; Okamoto, T.: Assignment of MARK3
[72906] alias KP78 to human chromosome band 14q32.3 by in

situ hybridization. Cytogenet.
[72907] Cell Genet. 79: 101-102, 1997.

[72908] 12411.

[72909] Parsa, I.: Loss of Mr 78,000 marker in chemically induced transplantable
[72910] carcinomas and primary carcinoma of human pancreas. Cancer Res. 48:
[72911] 2265-2272, 1988.

[72912] 12412.

[72913] Peng, C.-Y.; Graves, P. R.; Ogg, S.; Thoma, R. S.; Byrnes, M. J.,
[72914] III; Wu, Z.; Stephenson, M. T.; Piwnica-Worms, H.: C-TAK1 protein
[72915] kinase phosphorylates human Cdc25C on serine 216 and promotes 14-3-3
[72916] protein binding. Cell Growth Differ. 9: 197-208, 1998.

[72917] 12413.

[72918] Shiratsuchi, T.; Nishimori, H.; Ichise, H.; Nakamura, Y.; Tokino,
[72919] T.: Cloning and characterization of BAI2 and BAI3, novel genes homologous

[72920] to brain-specific angiogenesis inhibitor 1 (BAI1). Cyto-
genet. Cell
[72921] Genet. 79: 103-108, 1997.

[72922] 12414.

[72923] Brubaker, K.; Cowley, S. M.; Huang, K.; Loo, L.; Yochum, G.
S.;
[72924] Ayer, D. E.; Eisenman, R. N.; Radhakrishnan, I.: Solution
structure
[72925] of the interacting domains of the Mad-Sin3 complex: im-
plications for
[72926] recruitment of a chromatin-modifying complex. Cell 103:
655-665,
[72927] 2000.

[72928] 12415.

[72929] Jin, D.-Y.; Kozak, C. A.; Pangilinan, F.; Spencer, F.; Green,
E.
[72930] D.; Jeang, K.-T.: Mitotic checkpoint locus MAD1L1 maps
to human chromosome
[72931] 7p22 and mouse chromosome 5. Genomics 55: 363-364,
1999.
[72932] 12416.

[72933] Jin, D.-Y.; Spencer, F.; Jeang, K.-T.: Human T cell leukemia virus
[72934] type 1 oncoprotein Tax targets the human mitotic check-point protein
[72935] MAD1. Cell 93: 81-91, 1998.

[72936] 12417.

[72937] Tsukasaki, K.; Miller, C. W.; Greenspun, E.; Eshaghian, S.; Kawabata,
[72938] H.; Fujimoto, T.; Tomonaga, M.; Sawyers, C.; Said, J. W.; Koeffler,
[72939] H. P.: Mutations in the mitotic check point gene, MAD1L1, in human
[72940] cancers. Oncogene 20: 3301-3305, 2001.

[72941] 12418.

[72942] Kimura, M.; Kotani, S.; Hattori, T.; Sumi, N.; Yoshioka, T.; Todokoro,
[72943] T.; Okano, Y.: Cell cycle-dependent expression and spin-dle pole localization
[72944] of a novel human protein kinase Aik, related to Aurora of Drosophila
[72945] and yeast Ipl1. J. Biol. Chem. 272: 13766-13771, 1997.

[72946] 12419.

[72947] Bhatt, R. R.; Ferrell, J. E., Jr.: The protein kinase p90 Rsk

[72948] as an essential mediator of cytostatic factor activity. Sci-
ence 286:

[72949] 1362-1365, 1999.

[72950] 12420.

[72951] Bonni, A.; Brunet, A.; West, A. E.; Datta, S. R.; Takasu, M.
A.;

[72952] Greenberg, M. E.: Cell survival promoted by the Ras-MAPK
signaling

[72953] pathway by transcription-dependent and -independent
mechanisms. Science 286:

[72954] 1358-1362, 1999.

[72955] 12421.

[72956] Gross, S. D.; Schwab, M. S.; Lewellyn, A. L.; Maller, J. L.:
Induction

[72957] of metaphase arrest in cleaving Xenopus embryos by the
protein kinase

[72958] p90(Rsk). Science 286: 1365-1367, 1999.

[72959] 12422.

[72960] Pende, M.; Kozma, S. C.; Jaquet, M.; Oorschot, V.; Burcelin, R.;

[72961] Le Marchand-Brustel, Y.; Klumperman, J.; Thorens, B.; Thomas, G.:

[72962] Hypoinsulinaemia, glucose intolerance and diminished beta-cell size

[72963] in S6K1-deficient mice. Nature 408: 994-997, 2000.

[72964] 12423.

[72965] Shima, H.; Pende, M.; Chen, Y.; Fumagalli, S.; Thomas, G.; Kozma,

[72966] S. C.: Disruption of the p70(s6k)/p85(s6k) gene reveals a small mouse

[72967] phenotype and a new functional S6 kinase. EMBO J. 17: 6649-6659,

[72968] 1998.

[72969] 12424.

[72970] Goustin, A. S.: Personal Communication. Detroit, Mich. 8/17/1998.

[72971] 12425.

[72972] Chadwick, B. P.; Frischauf, A.-M.: The CD39-like gene family:

[72973] identification of three new human members (CD39L2, CD39L3, and CD39L4),

[72974] their murine homologues, and a member of the gene family from Drosophila

[72975] melanogaster. Genomics 50: 357-367, 1998.

[72976] 12426.

[72977] Kieffer, L. J.; Seng, T. W.; Li, W.; Osterman, D. G.; Handschumacher,

[72978] R. E.; Bayney, R. M.: Cyclophilin-40, a protein with homology to

[72979] the P59 component of the steroid receptor complex: cloning of the

[72980] cDNA and further characterization. J. Biol. Chem. 268: 12303-12310,

[72981] 1993.

[72982] 12427.

[72983] Kieffer, L. J.; Thalhammer, T.; Handschumacher, R. E.: Isolation

[72984] and characterization of a 40-kDa cyclophilin-related protein. J.

[72985] Biol. Chem. 267: 5503-5507, 1992.

[72986] 12428.

[72987] Ratajczak, T.; Woollatt, E.; Kumar, P.; Ward, B. K.; Minchin, R.

[72988] F.; Baker, E.: Cyclophilin 40 (PPID) gene map position 4q31.3. Chromosome

[72989] Res. 5: 151 only, 1997.

[72990] 12429.

[72991] Yokoi, H.; Shimizu, Y.; Anazawa, H.; Lefebvre, C. A.; Kor-neluk,

[72992] R. G.; Ikeda; J.-E.: The structure and complete nucleotide sequence

[72993] of the human cyclophilin 40 (PPID) gene. Genomics 35: 448-455, 1996.

[72994] 12430.

[72995] Gong, W.; Emanuel, B. S.; Collins, J.; Kim, D. H.; Wang, Z.; Chen,

[72996] F.; Zhang, G.; Roe, B.; Budarf, M. L.: A transcription map of the

[72997] DiGeorge and velo-cardio-facial syndrome minimal criti-cal region on

[72998] 22q11. Hum. Molec. Genet. 5: 789-800, 1996.

[72999] 12431.

[73000] Gong, W.; Emanuel, B. S.; Galili, N.; Kim, D. H.; Roe, B.; Driscoll,

[73001] D. A.; Budarf, M. L.: Structural and mutational analysis of a conserved

[73002] gene (DGSI) from the minimal DiGeorge syndrome critical region. Hum.

[73003] Molec. Genet. 6: 267-276, 1997.

[73004] 12432.

[73005] Lindsay, E. A.; Botta, A.; Jurecic, V.; Carattini-Rivera, S.; Cheah,

[73006] Y.-C.; Rosenblatt, H. M.; Bradley, A.; Baldini, A.: Congeni-tal heart

[73007] disease in mice deficient for the DiGeorge syndrome re-gion. Nature 401:

[73008] 379-383, 1999.

[73009] 12433.

[73010] Lindsay, E. A.; Harvey, E. L.; Scambler, P. J.; Baldini, A.: ES2,

[73011] a gene deleted in DiGeorge syndrome, encodes a nuclear protein and

[73012] is expressed during early mouse development, where it shares an expression

[73013] domain with a Goosecoid-like gene. Hum. Molec. Genet. 7: 629-635,

[73014] 1998.

[73015] 12434.

[73016] Ray, M. E.; Su, Y. A.; Meltzer, P. S.; Trent, J. M.: Isolation

[73017] and characterization of genes associated with chromosome-6 mediated

[73018] tumor suppression in human malignant melanoma. Oncogene 12: 2527-2533,

[73019] 1996.

[73020] 12435.

[73021] Ray, M. E.; Wistow, G.; Su, Y. A.; Meltzer, P. S.; Trent, J. M.

[73022] : AIM1, a novel non-lens member of the beta-gamma-crystallin superfamily,

[73023] is associated with the control of tumorigenicity in human malignant

[73024] melanoma. Proc. Nat. Acad. Sci. 94: 3229-3234, 1997.

[73025] 12436.

[73026] Teichmann, U.; Ray, M. E.; Ellison, J.; Graham, C.; Wistow, G.;

[73027] Meltzer, P. S.; Trent, J. M.; Pavan, W. J.: Cloning and tissue expression

[73028] of the mouse ortholog of AIM1, a beta-gamma-crystallin superfamily

[73029] member. Mammalian Genome 9: 715-720, 1998.

[73030] 12437.

[73031] Trent, J. M.; Stanbridge, E. J.; McBride, H. L.; Meese, E. U.;

[73032] Casey, G.; Araujo, D. E.; Witkowski, C. M.; Nagle, R. B.: Tu-morigenicity

[73033] in human melanoma cell lines controlled by introduction of human chromosome

[73034] 6. Science 247: 568-571, 1990.

[73035] 12438.

[73036] Nakamura, Y.; Miura, K.; Fujino, Y.; Iwao, H.; Ogita, S.; Ya-manaka,

[73037] S.: Evolution, structure, and expression of GNPI/oscillin orthologous

[73038] genes. Genomics 68: 179-186, 2000.

[73039] 12439.

[73040] Rogers, M. J.; Ohgi, T.; Plumbridge, J.; Soll, D.: Nucleotide

[73041] sequences of the E. coli nagE and nagB genes: the structural genes

[73042] for the N-acetylglucosamine transport protein of the bacterial phosphoenolpyruvate:

[73043] sugar phosphotransferase system and for glucosamine 6-phosphate deaminase. Gene 62:

[73044] 197-207, 1988.

[73045] 12440.

[73046] Shevchenko, V.; Hogben, M.; Ekong, R.; Parrington, J.; Lai, F.

[73047] A.: The human glucosamine-6-phosphate deaminase gene: cDNA cloning

[73048] and expression, genomic organization and chromosomal localization. Gene 216:

[73049] 31-38, 1998.

[73050] 12441.

[73051] Weidanz, J. A.; Campbell, P.; DeLucas, L. J.; Jin, J.; Moore, D.;

[73052] Roden, L.; Yu, H.; Heilmann, E.; Vezza, A. C.: Glucosamine 6-phosphate

[73053] deaminase in normal human erythrocytes. Brit. J. Haemat. 91: 72-79,

[73054] 1995.

[73055] 12442.

[73056] Bladergroen, B. A.; Strik, M. C. M.; Bovenschen, N.; van Berkum,

[73057] O.; Scheffer, G. L.; Meijer, C. J. L. M.; Hack, C. E.; Kummer, J.

[73058] A.: The granzyme B inhibitor, protease inhibitor 9, is mainly expressed

[73059] by dendritic cells and at immune-privileged sites. J. Immun. 166:

[73060] 3218-3225, 2001.

[73061] 12443.

[73062] Eyre, H. J.; Sun, J.; Sutherland, G. R.; Bird, P.: Chromosomal

[73063] mapping of the gene (PI9) encoding the intracellular serpin proteinase

[73064] inhibitor 9 to 6p25 by fluorescence in situ hybridization. Genomics 37:

[73065] 406-408, 1996.

[73066] 12444.

[73067] Krieg, S. A.; Krieg, A. J.; Shapiro, D. J.: A unique down-
stream
[73068] estrogen responsive unit mediates estrogen induction of
proteinase
[73069] inhibitor-9, a cellular inhibitor of IL-1-beta-converting
enzyme (caspase
[73070] 1). Molec. Endocr. 15: 1971-1982, 2001.

[73071] 12445.

[73072] Sun, J.; Bird, C. H.; Sutton, V.; McDonald, L.; Coughlin, P.
B.;
[73073] De Jong, T. A.; Trapani, J. A.; Bird, P. I.: A cytosolic
granzyme
[73074] B inhibitor related to the viral apoptotic regulator cytokine
response
[73075] modifier A is present in cytotoxic lymphocytes. J. Biol.
Chem. 271:
[73076] 27802-27809, 1996.

[73077] 12446.

[73078] Steimle, V.; Durand, B.; Barras, E.; Zuffrey, M.; Hadam, M.
R.;

[73079] Mach, B.; Reith, W.: A novel DNA binding regulatory factor is mutated
[73080] in primary MHC class II deficiency (bare lymphocyte syndrome). Genes
[73081] Dev. 9: 1021-1032, 1995.

[73082] 12447.

[73083] Dode, L.; Wuytack, F.; Kools, P. F. J.; Baba-Aissa, F.; Raeymaekers,
[73084] L.; Brik, F.; Van de Ven, W. J. M.; Casteels, R.: cDNA cloning, expression
[73085] and chromosomal localization of the human sarco/endoplasmic reticulum
[73086] Ca(2+)-ATPase 3 gene. Biochem. J. 318: 689-699, 1996.

[73087] 12448.

[73088] Teumer, J.; Tseng, H.; Green, H.: The human basonuclin gene. Gene 188:
[73089] 1-7, 1997.

[73090] 12449.

[73091] Tseng, H.; Green, H.: Basonuclin: a keratinocyte protein with
[73092] multiple paired zinc fingers. Proc. Nat. Acad. Sci. 89:

10311-10315,

[73093] 1992.

[73094] 12450.

[73095] Tseng, H.; Green, H.: Association of basonuclin with ability of

[73096] keratinocytes to multiply and with absence of terminal differentiation. J.

[73097] Cell Biol. 126: 495-506, 1994.

[73098] 12451.

[73099] Gibson, L.; Holmgreen, S. P.; Huang, D. C. S.; Bernard, O.; Copeland,

[73100] N. G.; Jenkins, N. A.; Sutherland, G. R.; Baker, E.; Adams, J. M.;

[73101] Cory, S.: bcl-w, a novel member of the bcl-2 family, pro- motes cell

[73102] survival. Oncogene 13: 665-675, 1996.

[73103] 12452.

[73104] Ross, A. J.; Waymire, K. G.; Moss, J. E.; Parlow, A. F.; Skin- ner,

[73105] M. K.; Russell, L. D.; MacGregor, G. R.: Testicular degener- ation

[73106] in Bclw-deficient mice. Nature Genet. 18: 251-256, 1998.

[73107] 12453.

[73108] Freedman, M. S.; Lucas, R. J.; Soni, B.; von Schantz, M.; Munoz,

[73109] M.; David-Gray, Z.; Foster, R.: Regulation of mammalian circadian

[73110] behavior by non-rod, non-cone, ocular photoreceptors. Science 284:

[73111] 502-504, 1999.

[73112] 12454.

[73113] Griffin, E. A., Jr.; Staknis, D.; Weitz, C. J.: Light-in-dependent

[73114] role of CRY1 and CRY2 in the mammalian circadian clock. Science 286:

[73115] 768-771, 1999.

[73116] 12455.

[73117] Hsu, D. S.; Zhao, X.; Zhao, S.; Kazantsev, A.; Wang, R.-P.; Todo,

[73118] T.; Wei, Y.-F.; Sancar, A.: Putative human blue-light pho-toreceptors

[73119] hCRY1 and hCRY2 are flavoproteins. Biochemistry 35:

13871-13877,

[73120] 1996.

[73121] 12456.

[73122] Kobayashi, K.; Kanno, S.; Smit, B.; van der Horst, G. T. J.; Takao,

[73123] M.; Yasui, A.: Characterization of photolyase/blue-light receptor

[73124] homologs in mouse and human cells. Nucleic Acids Res. 26: 5086-5092,

[73125] 1998.

[73126] 12457.

[73127] Kume, K.; Zylka, M. J.; Sriram, S.; Shearman, L. P.; Weaver, D.

[73128] R.; Jin, X.; Maywood, E. S.; Hastings, M. H.; Reppert, S. M.: mCRY1

[73129] and mCRY2 are essential components of the negative limb of the circadian

[73130] clock feedback loop. Cell 98: 193-205, 1999.

[73131] 12458.

[73132] Lucas, R. J.; Freedman, M. S.; Munoz, M.; Garcia-Fer-nandez, J.-M.;

[73133] Foster, R. G.: Regulation of the mammalian pineal by non-rod, non-cone,

[73134] ocular photoreceptors. Science 284: 505-507, 1999.

[73135] 12459.

[73136] Okamura, H.; Miyake, S.; Sumi, Y.; Yamaguchi, S.; Yasui, A.; Muijtjens,

[73137] M.; Hoeijmakers, J. H. J.; van der Horst, G. T. J.: Photic induction

[73138] of mPer1 and mPer2 in Cry-deficient mice lacking a biological clock. Science 286:

[73139] 2531-2534, 1999.

[73140] 12460.

[73141] Reick, M.; Garcia, J. A.; Dudley, C.; McKnight, S. L.: NPAS2:

[73142] an analog of clock operative in the mammalian forebrain. Science 293:

[73143] 506-509, 2001.

[73144] 12461.

[73145] van der Horst, G. T. J.; Muijtjens, M.; Kobayashi, K.; Takano,

[73146] R.; Kanno, S.; Takao, M.; de Wit, J.; Verkerk, A.; Eker, A. P. M.;

[73147] van Leenen, D.; Buijs, R.; Bootsma, D.; Hoeijmakers, J. H. J.; Yasui,
[73148] A.: Mammalian Cry1 and Cry2 are essential for maintenance of circadian
[73149] rhythms. Nature 398: 627-630, 1999.

[73150] 12462.

[73151] Yagita, K.; Tamanini, F.; van der Horst, G. T. J.; Okamura, H.
[73152] : Molecular mechanisms of the biological clock in cultured fibroblasts. Science 292:
[73153] 278-281, 2001.

[73154] 12463.

[73155] Villard, J.; Reith, W.; Barras, E.; Gos, A.; Morris, M. A.; Antonarakis,
[73156] S. E.; Van den Elsen, P. J.; Mach, B.: Analysis of mutations and
[73157] chromosomal localisation of the gene encoding RFX5, a novel transcription
[73158] factor affected in major histocompatibility complex class II deficiency. Hum.
[73159] Mutat. 10: 430-435, 1997.

[73160] 12464.

[73161] Sipila, L.; Szatanik, M.; Vainionpaa, H.; Ruotsalainen, H.; Myllyla,

[73162] R.; Guenet, J.-L.: The genes encoding mouse lysyl hydroxylase isoforms

[73163] map to chromosomes 4, 5, and 9. Mammalian Genome 11: 1132-1134,

[73164] 2000.

[73165] 12465.

[73166] Szpirer, C.; Szpirer, J.; Riviere, M.; Vanvooren, P.; Valtavaara,

[73167] M.; Myllyla, R.: Localization of the gene encoding a novel isoform

[73168] of lysyl hydroxylase. Mammalian Genome 8: 707-708, 1997.

[73169] 12466.

[73170] Valtavaara, M.; Papponen, H.; Pirttila, A.-M.; Hiltunen, K.; Helander,

[73171] H.; Myllyla, R.: Cloning and characterization of a novel human lysyl

[73172] hydroxylase isoform highly expressed in pancreas and muscle. J. Biol.

[73173] Chem. 272: 6831-6834, 1997.

[73174] 12467.

[73175] Hellborg, F.; Qian, W.; Mendez-Vidal, C.; Asker, C.; Kost-Alimova,

[73176] M.; Wilhelm, M.; Imreh, S.; Wiman, K. G.: Human wig-1, a p53 target

[73177] gene that encodes a growth inhibitory zinc finger protein. Oncogene 20:

[73178] 5466-5474, 2001.

[73179] 12468.

[73180] Varmeh-Ziaie, S.; Okan, I.; Wang, Y.; Magnusson, K. P.; Warthoe,

[73181] P.; Strauss, M.; Wiman, K. G.: Wig-1, a new p53-induced gene encoding

[73182] a zinc finger protein. Oncogene 15: 2699-2704, 1997.

[73183] 12469.

[73184] Modregger, J.; Ritter, B.; Witter, B.; Paulsson, M.; Plomann, M.

[73185] : All three PACSIN isoforms bind to endocytic proteins and inhibit

[73186] endocytosis. J. Cell Sci. 113: 4511-4521, 2000.

[73187] 12470.

[73188] Fardaei, M.; Rogers, M. T.; Thorpe, H. M.; Larkin, K.; Hamshere,

[73189] M. G.; Harper, P. S.; Brook, J. D.: Three proteins, MBNL, MBLL and

[73190] MBXL, co-localize in vivo with nuclear foci of expanded-re-peat transcripts

[73191] in DM1 and DM2 cells. Hum. Molec. Genet. 11: 805-814, 2002.

[73192] 12471.

[73193] Miller, J. W.; Urbinati, C. R.; Teng-umnuay, P.; Stenberg, M. G.;

[73194] Byrne, B. J.; Thornton, C. A.; Swanson, M. S.: Recruitment of human

[73195] muscleblind proteins to (CUG)n expansions associated with myotonic

[73196] dystrophy. EMBO J. 19: 4439-4448, 2000.

[73197] 12472.

[73198] Hu, C. C.; Fukae, M.; Uchida, T.; Qian, Q.; Zhang, C. H.; Ryu,

[73199] O. H.; Tanabe, T.; Yamakoshi, Y.; Murakami, C.; Dohi, N.; Shimizu,

[73200] M.; Simmer, J. P.: Cloning and characterization of porcine enamelin

[73201] mRNAs. J. Dent. Res. 76: 1720-1729, 1997.

[73202] 12473.

[73203] Hu, C. C.; Hart, T. C.; DuPont, B. R.; Chen, J. J.; Sun, X.; Qian,

[73204] Q.; Zhang, C. H.; Jiang, H.; Mattern, V. L.; Wright, J. T.; Simmer,

[73205] J. P.: Cloning human enamelin cDNA, chromosomal localization, and

[73206] analysis of expression during tooth development. J. Dent. Res. 79:

[73207] 912-919, 2000.

[73208] 12474.

[73209] Kutty, R. K.; Kutty, G.; Samuel, W.; Duncan, T.; Bridges, C. C.;

[73210] El-Sherbeeny, A.; Nagineni, C. N.; Smith, S. B.; Wiggert, B.: Molecular

[73211] characterization and developmental expression of NOR-PEG, a novel gene

[73212] induced by retinoic acid. J. Biol. Chem. 276: 2831-2840, 2001.

[73213] 12475.

[73214] Aapola, U.; Shibuya, K.; Scott, H. S.; Ollila, J.; Vihinen, M.;

[73215] Heino, M.; Shintani, A.; Kawasaki, K.; Minoshima, S.;
Krohn, K.; Antonarakis,

[73216] S. E.; Shimizu, N.; Kudoh, J.; Peterson, P.: Isolation and
initial

[73217] characterization of a novel zinc finger gene, DNMT3L, on
21q22.3,

[73218] related to the cysteine-5-methyltransferase 3 gene family.
Genomics 65:

[73219] 293-298, 2000.

[73220] 12476.

[73221] Bourc'his, D.; Xu, G.-L.; Lin, C.-S.; Bollman, B.; Bestor, T.
H.

[73222] : Dnmt3L and the establishment of maternal genomic im-
prints. Science 294:

[73223] 2536-2539, 2001.

[73224] 12477.

[73225] Chen, X.-B.; Melchionna, R.; Denis, C.-M.; Gaillard, P.-H.
L.;

[73226] Blasina, A.; Van de Weyer, I.; Boddy, M. N.; Russell, P.;

Vialard,
[73227] J.; McGowan, C. H.: Human Mus81-associated endonuclease cleaves Holliday
[73228] junctions in vitro. Molec. Cell 8: 1117-1127, 2001.

[73229] 12478.

[73230] Granjeaud, S.; Naquet, P.; Galland, F.: An ESTs description of
[73231] the new vanin gene family conserved from fly to human. Immunogenetics 49:
[73232] 964-972, 1999.

[73233] 12479.

[73234] Wang, H.; Cao, R.; Xia, L.; Erdjument-Bromage, H.; Borchers, C.;
[73235] Tempst, P.; Zhang, Y.: Purification and functional characterization
[73236] of a histone H3-lysine 4-specific methyltransferase. Molec. Cell 8:
[73237] 1207-1217, 2001.

[73238] 12480.

[73239] Brockington, M.; Blake, D. J.; Prandini, P.; Brown, S. C.; Torelli,

[73240] S.; Benson, M. A.; Ponting, C. P.; Estournet, B.; Romero, N. B.; Mercuri,
[73241] E.; Voit, T.; Sewry, C. A.; Guicheney, P.; Muntoni, F.: Mutations
[73242] in the fukutin-related protein gene (FKRP) cause a form of congenital
[73243] muscular dystrophy with secondary laminin alpha-2 deficiency and abnormal
[73244] glycosylation of alpha-dystroglycan. Am. J. Hum. Genet. 69: 1198-1209,
[73245] 2001.

[73246] 12481.

[73247] Brockington, M.; Yuva, Y.; Prandini, P.; Brown, S. C.; Torelli,
[73248] S.; Benson, M. A.; Herrmann, R.; Anderson, L. V. B.; Bashir, R.; Burgunder,
[73249] J.-M.; Fallet, S.; Romero, N.; and 10 others: Mutations in the
[73250] fukutin-related protein gene (FKRP) identify limb girdle muscular
[73251] dystrophy 2I as a milder allelic variant of congenital muscular dystrophy

[73252] MDC1C. Hum. Molec. Genet. 10: 2851-2859, 2001.

[73253] 12482.

[73254] Driss, A.; Amouri, C.; Hamida, C. B.; Souilem, N.; Gouider-
Khouja,
[73255] N.; Hamida, M. B.; Hentati, F.: A new locus for autosomal
recessive
[73256] limb girdle muscular dystrophy in a large consanguineous
Tunisian
[73257] family maps to chromosome 19q13.3. Neuromusc. Dis-
ord. 10: 240-246,
[73258] 2000.

[73259] 12483.

[73260] Mercuri, E.; Sewry, C. A.; Brown, S. C.; Brockington, M.;
Jungbluth,
[73261] H. DeVile, C.; Counsell, S.; Manzur, A.; Muntoni, F.: Con-
genital
[73262] muscular dystrophy with secondary merosin deficiency
and normal brain
[73263] MRI: a novel entity? Neuropediatrics 31: 186-189, 2000.

[73264] 12484.

[73265] Monney, L.; Sabatos, C.; Gaglia, J. L.; Ryu, A.; Waldner, H.;

Chernova,
[73266] T.; Manning, S.; Greenfield, E. A.; Coyle, A. J.; Sobel, R. A.; Freeman,
[73267] G. J.; Kuchroo, V. K.: Th1-specific cell surface protein regulates
[73268] macrophage activation and severity of an autoimmune disease. Nature 415:
[73269] 536-541, 2002.

[73270] 12485.

[73271] Hsu, S. Y.; Kudo, M.; Chen, T.; Nakabayashi, K.; Bhalla, A.; van
[73272] der Spek, P. J.; van Duin, M.; Hsueh, A. J. W.: The three subfamilies
[73273] of leucine-rich repeat-containing G protein-coupled receptors (LGR):
[73274] identification of LGR6 and LGR7 and the signaling mechanism for LGR7. Molec.
[73275] Endocr. 14: 1257-1271, 2000.

[73276] 12486.

[73277] Jourdan-Le Saux, C.; Le Saux, O.; Donlon, T.; Boyd, C. D.; Csiszar,
[73278] K.: The human lysyl oxidase-related gene (LOXL2) maps

between markers

[73279] D8S280 and D8S278 on chromosome 8p21.2-p21.3. Genomics 51: 305-307,

[73280] 1998.

[73281] 12487.

[73282] Jourdan-Le Saux, C.; Tronecker, H.; Bogic, L.; Bryant-Greenwood,

[73283] G. D.; Boyd, C. D.; Csiszar, K.: The LOXL2 gene encodes a new lysyl

[73284] oxidase-like protein and is expressed at high levels in reproductive

[73285] tissues. J. Biol. Chem. 274: 12939-12944, 1999.

[73286] 12488.

[73287] Ikeda, K.; Sato, M.; Tsutsumi, O.; Tsuchiya, F.; Tsuneizumi, M.;

[73288] Emi, M.; Imoto, I.; Inazawa, J.; Muramatsu, M.; Inoue, S.: Promoter

[73289] analysis and chromosomal mapping of human EBAG9 gene. Biochem. Biophys.

[73290] Res. Commun. 273: 654-660, 2000.

[73291] 12489.

[73292] Nakashima, M.; Sonoda, K.; Watanabe, T.: Inhibition of cell growth
[73293] and induction of apoptotic cell death by the human tumor-associated
[73294] antigen RCAS1. Nature Med. 5: 938-942, 1999.

[73295] 12490.

[73296] Kamohara, M.; Takasaki, J.; Matsumoto, M.; Saito, T.; Ohishi, T.;
[73297] Ishii, H.; Furuichi, K.: Molecular cloning and characterization of
[73298] another leukotriene B4 receptor. J. Biol. Chem. 275: 27000-27004,
[73299] 2000.

[73300] 12491.

[73301] Nilsson, N. E.; Tryselius, Y.; Owman, C.: Genomic organization
[73302] of the leukotriene B4 receptor locus of human chromosome 14. Biochem.
[73303] Biophys. Res. Commun. 274: 383-388, 2000.

[73304] 12492.

[73305] Tryselius, Y.; Nilsson, N. E.; Kotarsky, K.; Olde, B.; Owman,

C.

[73306] : Cloning and characterization of cDNA encoding a novel human leukotriene

[73307] B4 receptor. Biochem. Biophys. Res. Commun. 274: 377-382, 2000.

[73308] 12493.

[73309] Wang, S.; Gustafson, E.; Pang, L.; Qiao, X.; Behan, J.; Maguire,

[73310] M.; Bayne, M.; Laz, T.: A novel hepatointestinal leukotriene B4 receptor:

[73311] cloning and functional characterization. J. Biol. Chem. 275: 40686-40694,

[73312] 2000.

[73313] 12494.

[73314] Yokomizo, T.; Kato, K.; Terawaki, K.; Izumi, T.; Shimizu, T.:

[73315] A second leukotriene B4 receptor, BLT2: a new therapeutic target in

[73316] inflammation and immunological disorders. J. Exp. Med. 192: 421-431,

[73317] 2000.

[73318] 12495.

[73319] Lai, F.; Orelli, B. J.; Till, B. G.; Godley, L. A.; Fernald, A.

[73320] A.; Pamintuan, L.; Le Beau, M. M.: Molecular characterization of

[73321] KLHL3, a human homologue of the Drosophila kelch gene. Genomics 66:

[73322] 65-75, 2000.

[73323] 12496.

[73324] Yang, X.-H.; Huang, S.: PFM1 (PRDM4), a new member of the PR-domain

[73325] family, maps to a tumor suppressor locus on human chromosome 12q23-q24.1. Genomics 61:

[73326] 319-325, 1999.

[73327] 12497.

[73328] Jenkins, Z. A.; Haag, P. G.; Johansson, H. E.: Human EIF5A2 on

[73329] chromosome 3q25-q27 is a phylogenetically conserved vertebrate variant

[73330] of eukaryotic translation initiation factor 5A with tissue-specific

[73331] expression. Genomics 71: 101-109, 2001.

[73332] 12498.

[73333] Nakagawa, H.; Koyama, K.; Murata, Y.; Morito, M.; Akiyama, T.;

[73334] Nakamura, Y.: EB3, a novel member of the EB1 family preferentially

[73335] expressed in the central nervous system, binds to a CNS-specific APC

[73336] homologue. Oncogene 19: 210-216, 2000.

[73337] 12499.

[73338] Hocking, L. J.; Herbert, C. A.; Nicholls, R. K.; Williams, F.;

[73339] Bennett, S. T.; Cundy, T.; Nicholson, G. C.; Wuyts, W.; Van Hul, W.;

[73340] Ralston, S. H.: Genomewide search in familial Paget dis-ease of bone

[73341] shows evidence of genetic heterogeneity with candidate loci on chromosomes

[73342] 2q36, 10p13, and 5q35. Am. J. Hum. Genet. 69: 1055-1061, 2001.

[73343] 12500.

[73344] Collins, C. S.; Gould, S. J.: Identification of a common PEX1

[73345] mutation in Zellweger syndrome. Hum. Mutat. 14: 45-53, 1999.

[73346] 12501.

[73347] Gartner, J.; Preuss, N.; Brosius, U.; Biermanns, M.: Muta-
tions
[73348] in PEX1 in peroxisome biogenesis disorders: G843D and a
mild clinical
[73349] phenotype. J. Inherit. Metab. Dis. 22: 311-313, 1999.

[73350] 12502.

[73351] Imamura, A.; Tamura, S.; Shimozawa, N.; Suzuki, Y.;
Zhang, Z.;
[73352] Tsukamoto, T.; Orii, T.; Kondo, N.; Osumi, T.; Fujiki, Y.:
Temperature-sensitive
[73353] mutation in PEX1 moderates the phenotypes of peroxi-
some deficiency
[73354] disorders. Hum. Molec. Genet. 7: 2089-2094, 1998.

[73355] 12503.

[73356] Portsteffen, H.; Beyer, A.; Becker, E.; Epplen, C.; Pawlak,
A.;
[73357] Kunau, W.-H.; Dodt, G.: Human PEX1 is mutated in com-
plementation
[73358] group 1 of the peroxisome biogenesis disorders. Nature
Genet. 17:

[73359] 449-452, 1997.

[73360] 12504.

[73361] Tamura, S.; Okumoto, K.; Toyama, R.; Shimozawa, N.; Tsukamoto,

[73362] T.; Suzuki, Y.; Osumi, T.; Kondo, N.; Fujiki, Y.: Human PEX1 cloned

[73363] by functional complementation in a CHO cell mutant is responsible

[73364] for peroxisome-deficient Zellweger syndrome of complementation group

[73365] I. Proc. Nat. Acad. Sci. 95: 4350-4355, 1998.

[73366] 12505.

[73367] Walter, C.; Gootjes, J.; Mooijer, P. A.; Portsteffen, H.; Klein,

[73368] C.; Waterham, H. R.; Barth, P. G.; Epplen, J. T.; Kunau, W.-H.; Wanders,

[73369] R. J. A.; Dodt, G.: Disorders of peroxisome biogenesis due to mutations

[73370] in PEX1: phenotypes and PEX1 protein levels. Am. J. Hum. Genet. 69:

[73371] 35-48, 2001.

[73372] 12506.

[73373] Dunbar, D. R.; Shibasaki, Y.; Dobbie, L.; Andersson, B.; Brookes,
[73374] A. J.: In situ hybridisation mapping of genomic clones for five human
[73375] respiratory chain complex I genes. Cytogenet. Cell Genet. 78: 21-24,
[73376] 1997.

[73377] 12507.

[73378] Emahazion, T.; Brookes, A. J.: Mapping of the NDUFA2, NDUFA6,
[73379] NDUFA7, NDUFB8, and NDUFS8 electron transport chain genes by intron
[73380] based radiation hybrid mapping. Cytogenet. Cell Genet. 82: 114 only,
[73381] 1998.

[73382] 12508.

[73383] Ton, C.; Hwang, D. M.; Dempsey, A. A.; Liew, C.-C.: Identification

[73384] and primary structure of five human NADH-ubiquinone oxidoreductase
[73385] subunits. Biochem. Biophys. Res. Commun. 241: 589-594, 1997.
[73386] 12509.

[73387] Cheng, H. F.; Jiang, M. J.; Chen, C. L.; Liu, S. M.; Wong, L. P.;
[73388] Lomasney, J. W.; King, K.: Cloning and identification of amino acid
[73389] residues of human phospholipase C delta 1 essential for catalysis. J.
[73390] Biol. Chem. 270: 5495-5505, 1995.

[73391] 12510.

[73392] Tsukiyama-Kohara, K.; Vidal, S. M.; Gingras, A.-C.; Glover, T.
[73393] W.; Hanash, S. M.; Heng, H.; Sonenberg, N.: Tissue distribution,
[73394] genomic structure, and chromosome mapping of mouse and human eukaryotic
[73395] initiation factor 4E-binding proteins 1 and 2. Genomics 38: 353-363,
[73396] 1996.

[73397] 12511.

[73398] Lane, P. W.; Liu, H. M.: Association of megacolon with a new

[73399] dominant spotting gene (Dom) in the mouse. J. Hered. 75: 435-439,

[73400] 1984.

[73401] 12512.

[73402] Lee, M.; Goodall, J.; Verastegui, C.; Ballotti, R.; Goding, C.

[73403] R.: Direct regulation of the microphthalmia promoter by Sox10 links

[73404] Waardenburg-Shah syndrome (WS4)-associated hypopig- mentation and deafness

[73405] to WS2. J. Biol. Chem. 275: 37978-37983, 2000.

[73406] 12513.

[73407] Pingault, V.; Bondurand, N.; Kuhlbrodt, K.; Goerich, D. E.; Prehu,

[73408] M.-O.; Puliti, A.; Herbarth, B.; Hermans-Borgmeyer, I.; Legius, E.;

[73409] Matthijs, G.; Amiel, J.; Lyonnet, S.; Ceccherini, I.; Romeo, G.; Smith,

[73410] J. C.; Read, A. P.; Wegner, M.; Goossens, M.: SOX10 muta-

tions in
[73411] patients with Waardenburg-Hirschsprung disease. Nature Genet. 18:
[73412] 171-173, 1998.

[73413] 12514.

[73414] Pingault, V.; Girard, M.; Bondurand, N.; Dorkins, H.; Van Maldergem,
[73415] L.; Mowat, D.; Shimotake, T.; Verma, I.; Baumann, C.; Goossens, M.
[73416] : SOX10 mutations in chronic intestinal pseudo-obstruction suggest
[73417] a complex physiopathological mechanism. Hum. Genet. 111: 198-206,
[73418] 2002.

[73419] 12515.

[73420] Pingault, V.; Puliti, A.; Prehu, M.-O.; Samadi, A.; Bondurand,
[73421] N.; Goossens, M.: Human homology and candidate genes for the dominant
[73422] megacolon locus, a mouse model of Hirschsprung disease. Genomics 39:
[73423] 86-89, 1997.

[73424] 12516.

[73425] Pingault, V., Guiochon-Mantel, A.; Bondurand, N.; Faure, C.; Lacroix,

[73426] C.; Lyonnet, S.; Goossens, M.; Landrieu, P.: Peripheral neuropathy

[73427] with hypomyelination, chronic intestinal pseudo-obstruction and deafness:

[73428] a developmental 'neural crest syndrome' related to a SOX10 mutation. Ann.

[73429] Neurol. 48: 671-676, 2000.

[73430] 12517.

[73431] Potterf, S. B.; Furumura, M.; Dunn, K. J.; Arnheiter, H.; Pavan,

[73432] W. J.: Transcription factor hierarchy in Waardenburg syndrome: regulation

[73433] of MITF expression by SOX10 and PAX3. Hum. Genet. 107: 1-6, 2000.

[73434] 12518.

[73435] Southard-Smith, E. M.; Angrist, M.; Ellison, J. S.; Agarwala,

[73436] R.; Baxevanis, A. D.; Chakravarti, A.; Pavan, W. J.: The Sox10(Dom)

[73437] mouse: modeling the genetic variation of Waardenburg-Shah (WS4) syndrome. Genome
[73438] Res. 9: 215-225, 1999.

[73439] 12519.

[73440] Southard-Smith, E. M.; Kos, L.; Pavan, W. J.: Sox10 muta-tion
[73441] disrupts neural crest development in Dom Hirschsprung mouse model. Nature
[73442] Genet. 18: 60-64, 1998.

[73443] 12520.

[73444] Stock, D. W.; Buchanan, A. V.; Zhao, Z.; Weiss, K. M.: Nu-merous
[73445] members of the Sox family of HMG box-containing genes are expressed
[73446] in developing mouse teeth. Genomics 37: 234-237, 1996.

[73447] 12521.

[73448] Wright, E. M.; Snopek, B.; Koopman, P.: Seven new mem-bers of
[73449] the Sox gene family expressed during mouse develop-ment. Nucleic Acids
[73450] Res. 21: 744 only, 1993.

[73451] 12522.

[73452] Meluh, P. B.; Koshland, D.: Suppressors of MIF2, a putative centromere

[73453] protein gene in Saccharomyces cerevisiae. (Abstract) Molec. Biol.

[73454] Cell 6 (supp.): 360a only, 1995.

[73455] 12523.

[73456] Meluh, P. B.; Koshland, D.: Evidence that the MIF2 gene of Saccharomyces

[73457] cerevisiae encodes a centromere protein with homology to the mammalian

[73458] centromere protein CENP-C. Molec. Biol. Cell 6: 793-807, 1995.

[73459] 12524.

[73460] Cecconi, F.; Alvarez-Bolado, G.; Meyer, B. I.; Roth, K. A.; Gruss,

[73461] P.: Apaf1 (CED-4 homolog) regulates programmed cell death in mammalian

[73462] development. Cell 94: 727-737, 1998.

[73463] 12525.

[73464] Hahn, C.; Hirsch, B.; Jahnke, D.; Durkop, H.; Stein, H.:

Three
[73465] new types of Apaf-1 in mammalian cells. Biochem. Bio-
phys. Res. Commun. 261:
[73466] 746-749, 1999.

[73467] 12526.

[73468] Honarpour, N.; Du, C.; Richardson, J. A.; Hammer, R. E.;
Wang,
[73469] X.; Herz, J.: Adult Apaf-1-deficient mice exhibit male in-
fertility. Dev.
[73470] Biol. 218: 248-258, 2000.

[73471] 12527.

[73472] Honarpour, N.; Gilbert, S. L.; Lahn, B. T.; Wang, X.; Herz, J.

[73473] : Apaf-1 deficiency and neural tube closure defects are
found in fog
[73474] mice. Proc. Nat. Acad. Sci. 98: 9683-9687, 2001.

[73475] 12528.

[73476] Kim, H.; Jung, Y. K.; Kwon, Y. K.; Park, S. H.: Assignment
of
[73477] apoptotic protease activating factor-1 gene (APAF1) to
human chromosome
[73478] band 12q23 by fluorescence in situ hybridization. Cyto-

genet. Cell

[73479] Genet. 87: 252-253, 1999.

[73480] 12529.

[73481] Watanabe, T. K.; Katagiri, T.; Suzuki, M.; Shimizu, F.; Fuji-

wara,

[73482] T.; Kanemoto, N.; Nakamura, Y.; Hirai, Y.; Maekawa, H.;

Takahashi,

[73483] E.: Cloning and characterization of two novel human cD-

NAs (NELL1

[73484] and NELL2) encoding proteins with six EGF-like repeats.

Genomics 38:

[73485] 273-276, 1996.

[73486] 12530.

[73487] Imataka, H.; Olsen, H. S.; Sonenberg. N.: A new transla-

tional

[73488] regulator with homology to eukaryotic translation initia-

tion factor

[73489] 4G. EMBO J. 16: 817-825, 1997.

[73490] 12531.

[73491] Levy-Strumpf, N.; Deiss, L. P.; Berissi, H.; Kimchi, A.: DAP-

5,

[73492] a novel homolog of eukaryotic translation initiation factor 4G isolated

[73493] as a putative modulator of gamma interferon-induced programmed cell

[73494] death. Molec. Cell. Biol. 17: 1615-1625, 1997.

[73495] 12532.

[73496] Shaughnessy, J. D., Jr.; Jenkins, N. A.; Copeland, N. G.: cDNA

[73497] cloning, expression analysis, and chromosomal localization of a gene

[73498] with high homology to wheat eIF-(iso)4F and mammalian eIF-4G. Genomics 39:

[73499] 192-197, 1997.

[73500] 12533.

[73501] Yamanaka, S.; Poksay, K. S.; Arnold, K. S.; Innerarity, T. L.:

[73502] A novel translational repressor mRNA is edited extensively in livers

[73503] containing tumors caused by the transgene expression of the apoB mRNA-editing

[73504] enzyme. Genes Dev. 11: 321-333, 1997.

[73505] 12534.

[73506] Nomi, M.; Oishi, I.; Kani, S.; Suzuki, H.; Matsuda, T.; Yoda, A.;

[73507] Kitamura, M.; Itoh, K.; Takeuchi, S.; Takeda, K.; Akira, S.; Ikeya,

[73508] M.; Takada, S.; Minami, Y.: Loss of mRor1 enhances the heart and

[73509] skeletal abnormalities in mRor2-deficient mice: redundant and pleiotropic

[73510] functions of mRor1 and mRor2 receptor tyrosine kinases. Molec. Cell.

[73511] Biol. 21: 8329-8335, 2001.

[73512] 12535.

[73513] Korver, W.; Roose, J.; Heinen, K.; Weghuis, D. O.; de Bruijn, D.;

[73514] Geurts van Kessel, A.; Clevers, H.: The human TRIDENT/ HFH-11/FKHL16

[73515] gene: structure, localization, and promoter characterization. Genomics 46:

[73516] 435-442, 1997.

[73517] 12536.

[73518] Ly, D. H.; Lockhart, D. J.; Lerner, R. A.; Schultz, P. G.: Mitotic

[73519] misregulation and human aging. Science 287: 2486-2492, 2000.

[73520] 12537.

[73521] Wang, X.; Quail, E.; Hung, N.-J.; Tan, Y.; Ye, H.; Costa, R. H.

[73522] : Increased levels of forkhead box M1B transcription factor in transgenic

[73523] mouse hepatocytes prevent age-related proliferation de- fects in regenerating

[73524] liver. Proc. Nat. Acad. Sci. 98: 11468-11473, 2001.

[73525] 12538.

[73526] Westendorf, J. M.; Rao, P. N.; Gerace, L.: Cloning of cDNAs for

[73527] M-phase phosphoproteins recognized by the MPM2 mon- oclonal antibody

[73528] and determination of the phosphorylated epitope. Proc. Nat. Acad.

[73529] Sci. 91: 714-718, 1994.

[73530] 12539.

[73531] Yao, K.-M.; Sha, M.; Lu, Z.; Wong, G. G.: Molecular analysis of

[73532] a novel winged helix protein, WIN: expression pattern, DNA binding

[73533] property, and alternative splicing within the DNA binding domain. J.

[73534] Biol. Chem. 272: 19827-19836, 1997.

[73535] 12540.

[73536] Wang, X. S.; Diener, K.; Jannuzzi, D.; Trollinger, D.; Tan, T.-H.;

[73537] Lichenstein, H.; Zukowski, M.; Yao, Z.: Molecular cloning and characterization

[73538] of a novel protein kinase with a catalytic domain homologous to mitogen-activated

[73539] protein kinase kinase kinase. J. Biol. Chem. 271: 31607-31611, 1996.

[73540] 12541.

[73541] Venter, D. J.; Bevan, K. L.; Ludwig, R. L.; Riley, T. E.; Jat,

[73542] P. S.; Thomas, D. G.; Noble, M. D.: Retinoblastoma gene deletions

[73543] in human glioblastomas. Oncogene 6: 445-448, 1991.

[73544] 12542.

[73545] Richard, G.; Lin, J. P.; Smith, L.; Whyte, Y. M.; Itin, P.; Wol-

lina,
[73546] U.; Epstein, E., Jr.; Hohl, D.; Giroux, J. M.; Charnas, L.; Bale,
[73547] S. J.; DiGiovanna, J. J.: Linkage studies in erythrokeratodermias:
[73548] fine mapping, genetic heterogeneity and analysis of candidate genes. J.
[73549] Invest. Derm. 109: 666-671, 1997.

[73550] 12543.

[73551] Richard, G.; Smith, L. E.; Bailey, R. A.; Itin, P.; Hohl, D.; Epstein,
[73552] E. H., Jr.; DiGiovanna, J. J.; Compton, J. G.; Bale, S. J.: Mutations
[73553] in the human connexin gene GJB3 cause erythrokeratodermia variabilis. Nature
[73554] Genet. 20: 366-369, 1998.

[73555] 12544.

[73556] Wenzel, K.; Manthey, D.; Willecke, K.; Grzeschik, K.-H.; Traub,
[73557] O.: Human gap junction protein connexin31: molecular cloning and
[73558] expression analysis. Biochem. Biophys. Res. Commun.

248: 910-915,

[73559] 1998.

[73560] 12545.

[73561] Chuaqui, R. F.; Englert, C. R.; Strup, S. E.; Vocke, C. D.; Zhuang,

[73562] Z.; Duray, P. H.; Bostwick, D. G.; Linehan, W. M.; Liotta, L. A.;

[73563] Emmert-Buck, M. R.: Identification of a novel transcript up-regulated

[73564] in a clinically aggressive prostate carcinoma. Urology 50: 302-307,

[73565] 1997.

[73566] 12546.

[73567] Cole, K. A.; Chuaqui, R. F.; Katz, K.; Pack, S.; Zhuang, Z.; Cole,

[73568] C. E.; Lyne, J. C.; Linehan, W. M.; Liotta, L. A.; Emmert-Buck, M.

[73569] R.: cDNA sequencing and analysis of POV1 (PB39): a novel gene up-regulated

[73570] in prostate cancer. Genomics 51: 282-287, 1998.

[73571] 12547.

[73572] Bono, P.; Salmi, M.; Smith, D. J.; Leppanen, I.; Horelli-Kuitunen,
[73573] N.; Palotie, A.; Jalkanen, S.: Isolation, structural characterization,
[73574] and chromosomal mapping of the mouse vascular adhesion protein-1 gene
[73575] and promoter. J. Immun. 161: 2953-2960, 1998.

[73576] 12548.

[73577] Morris, N. J.; Ducret, A.; Aebersold, R.; Ross, S. A.; Keller,
[73578] S. R.; Lienhard, G. E.: Membrane amine oxidase cloning and identification
[73579] as a major protein in the adipocyte plasma membrane. J. Biol. Chem. 272:
[73580] 9388-9392, 1997.

[73581] 12549.

[73582] Bromme, D.; Rossi, A. B.; Smeekens, S. P.; Anderson, D. C.; Payan,
[73583] D. G.: Human bleomycin hydrolase: molecular cloning, sequencing,
[73584] functional expression, and enzymatic characterization. Biochemistry 35:

[73585] 6706-6714, 1996.

[73586] 12550.

[73587] Cloos, J.; Nieuwenhuis, E. J. C.; Boomsma, D. I.; Kuik, D. J.;

[73588] van der Sterre, M. L. T.; Arwert, F.; Snow, G. B.; Braakhuis, B. J.

[73589] M.: Inherited susceptibility to bleomycin-induced chromatid breaks

[73590] in cultured peripheral blood lymphocytes. J. Nat. Cancer Inst. 91:

[73591] 1125-1130, 1999.

[73592] 12551.

[73593] Farrer, L. A.; Abraham, C. R.; Haines, J. L.; Rogaeva, E. A.; Song,

[73594] Y.; McGraw, W. T.; Brindle, N.; Premkumar, S.; Scott, W. K.; Yamaoka,

[73595] L. H.; Saunders, A. M.; Roses, A. D.; Auerbach, S. A.; Sorbi, S.;

[73596] Duara, R.; Pericak-Vance, M. A.; St. George-Hyslop, P. H.: Association

[73597] between bleomycin hydrolase and Alzheimer's disease in Caucasians. Ann.

[73598] Neurol. 44: 808-811, 1998.

[73599] 12552.

[73600] Ferrando, A. A.; Pendas, A. M.; Llano, E.; Velasco, G.; Lid-
ereau,
[73601] R.; Lopez-Otin, C.: Gene characterization, promoter anal-
ysis, and
[73602] chromosomal localization of human bleomycin hydrolase.
J. Biol. Chem. 272:
[73603] 33298-33304, 1997.

[73604] 12553.

[73605] Ferrando, A. A.; Velasco, G.; Campo, E.; Lopez-Otin, C.:
Cloning
[73606] and expression analysis of human bleomycin hydrolase, a
cysteine proteinase
[73607] involved in chemotherapy resistance. Cancer Res. 56:
1746-1750,
[73608] 1996.

[73609] 12554.

[73610] Haston, C. K.; Amos, C. I.; King, T. M.; Travis, E. L.: Inheri-
tance
[73611] of susceptibility to bleomycin-induced pulmonary fibrosis

in the mouse. Cancer
[73612] Res. 56: 2596-2601, 1996.

[73613] 12555.

[73614] Hsu, T. C.; Johnston, D. A.; Cherry, L. M.; Ramkissoon, D.; Schantz,
[73615] S. P.; Jessup, J. M.; Winn, R. J.; Shirley, L.; Furlong, C.: Sen-
sitivity
[73616] to genotoxic effects of bleomycin in humans: possible re-
lationship
[73617] to environmental carcinogenesis. Int. J. Cancer 43:
403-409, 1989.
[73618] 12556.

[73619] Lazo, J. S.; Humphreys, C. J.: Lack of metabolism as the
biochemical
[73620] basis of bleomycin-induced pulmonary toxicity. Proc. Nat.
Acad. Sci. 80:
[73621] 3064-3068, 1983.

[73622] 12557.

[73623] Montoya, S. E.; Aston, C. E.; DeKosky, S. T.; Kamboh, M. I.; Lazo,
[73624] J. S.; Ferrell, R. E.: Bleomycin hydrolase is associated with

risk

[73625] of sporadic Alzheimer's disease. (Letter) Nature Genet. 18: 211-212,

[73626] 1998. Note: Erratum: Nature Genet. 19: 404 only, 1998.

[73627] 12558.

[73628] Montoya, S. E.; Ferrell, R. E.; Lazo, J. S.: Genomic structure

[73629] and genetic mapping of the human neutral cysteine protease bleomycin

[73630] hydrolase. Cancer Res. 57: 4191-4195, 1997.

[73631] 12559.

[73632] Zheng, W.; Johnston, S. A.; Joshua-Tor, L.: The unusual active

[73633] site of Gal6/bleomycin hydrolase can act as a carboxypeptidase, aminopeptidase,

[73634] and peptide ligase. Cell 93: 103-109, 1998.

[73635] 12560.

[73636] Xu, X.-N.; Screaton, G. R.; Gotch, F. M.; Dong, T.; Tan, R.; Almond,

[73637] N.; Walker, B.; Stebbings, R.; Kent, K.; Nagata, S.; Stott, J. E.;

[73638] McMichael, A. J.: Evasion of cytotoxic T lymphocyte (CTL)

responses

[73639] by Nef-dependent induction of Fas ligand (CD95L) expression on simian

[73640] immunodeficiency virus-infected cells. J. Exp. Med. 186: 7-16, 1997.

[73641] 12561.

[73642] Huang, L. J.; Durick, K.; Weiner, J. A.; Chun, J.; Taylor, S. S.

[73643] : Identification of a novel protein kinase A anchoring protein that

[73644] binds both type I and type II regulatory subunits. J. Biol. Chem. 272:

[73645] 8057-8064, 1997.

[73646] 12562.

[73647] Lin, R.-Y.; Moss, S. B.; Rubin, C. S.: Characterization of S-AKAP84,

[73648] a novel developmentally regulated A kinase anchor protein of male

[73649] germ cells. J. Biol. Chem. 270: 27804-27811, 1995.

[73650] 12563.

[73651] Trendelenburg, G.; Hummel, M.; Riecken, E.-O.; Hanski, C.: Molecular

[73652] characterization of AKAP149, a novel A kinase anchor protein with

[73653] a KH domain. Biochem. Biophys. Res. Commun. 225: 313-319, 1996.

[73654] 12564.

[73655] Moshous, D.; Callebaut, I.; de Chasseval, R.; Corneo, B.; Cavazzana-Calvo,

[73656] M.; Le Deist, F.; Tezcan, I.; Sanal, O.; Bertrand, Y.; Philippe, N.;

[73657] Fischer, A.; de Villartay, J.-P.: Artemis, a novel DNA dou-ble-strand

[73658] break repair/V(D)J recombination protein, is mutated in human severe

[73659] combined immune deficiency. Cell 105: 177-186, 2001.

[73660] 12565.

[73661] Chang, K.; Hanaoka, K.; Kumada, M.; Takuwa, Y.: Molecu-lar cloning

[73662] and functional analysis of a novel P2 nucleotide receptor. J. Biol.

[73663] Chem. 270: 26152-26158, 1995.

[73664] 12566.

[73665] Communi, D.; Parmentier, M.; Boeynaems, J.-M.: Cloning, functional

[73666] expression and tissue distribution of the human P2Y6 receptor. Biochem.

[73667] Biophys. Res. Commun. 222: 303-308, 1996.

[73668] 12567.

[73669] Maier, R.; Glatz, A.; Mosbacher, J.; Bilbe, G.: Cloning of P2Y6

[73670] cDNAs and identification of a pseudogene: comparison of P2Y receptor

[73671] subtype expression in bone and brain tissues. Biochem. Biophys. Res.

[73672] Commun. 237: 297-302, 1997.

[73673] 12568.

[73674] Pidlaoan, L. V.; Jin, J.; Sandhu, A. K.; Athwal, R. S.; Kunapuli,

[73675] S. P.: Colocalization of P2Y2 and P2Y6 receptor genes at human chromosome

[73676] 11q13.3-14.1. Somat. Cell Molec. Genet. 23: 291-296, 1997.

[73677] 12569.

[73678] Cahill, D. P.; Lengauer, C.; Yu, J.; Riggins, G. J.; Willson, J.

[73679] K. V.; Markowitz, S. D.; Kinzler, K. W.; Vogelstein, B.: Mu-
tations

[73680] of mitotic checkpoint genes in human cancers. Nature
392: 300-303,

[73681] 1998.

[73682] 12570.

[73683] Wang, B.; Kishihara, K.; Zhang, D.; Hara, H.; Nomoto, K.:
Molecular

[73684] cloning and characterization of a novel human receptor
protein tyrosine

[73685] phosphatase gene, hPTP-J: down-regulation of gene ex-
pression by PMA

[73686] and calcium ionophore in Jurkat T lymphoma cells.
Biochem. Biophys.

[73687] Res. Commun. 231: 77-81, 1997.

[73688] 12571.

[73689] Wang, H; Lian, Z; Lerch, M. M.; Chen, Z; Xie, W; Ullrich, A.:

[73690] Characterization of PCP-2, a novel receptor protein tyro-
sine phosphatase

[73691] of the MAM domain family. Oncogene 12: 2555-2562,

1996.

[73692] 12572.

[73693] Chinnaiyan, A. M.; O'Rourke, K.; Tewari, M.; Dixit, V. M.: FADD,

[73694] a novel death domain-containing protein, interacts with the death

[73695] domain of Fas and initiates apoptosis. Cell 81: 505-512, 1995.

[73696] 12573.

[73697] Kabra, N. H.; Kang, C.; Hsing, L. C.; Zhang, J.; Winoto, A.: T

[73698] cell-specific FADD-deficient mice: FADD is required for early T cell

[73699] development. Proc. Nat. Acad. Sci. 98: 6307-6312, 2001.

[73700] 12574.

[73701] Herrscher, R. F.; Kaplan, M. H.; Lelsz, D. L.; Das, C.; Scheuermann,

[73702] R.; Tucker, P. W.: The immunoglobulin heavy-chain ma-trix-associating

[73703] regions are bound by Bright: a B cell-specific trans-activator that

[73704] describes a new DNA-binding protein family. Genes Dev.

9: 3067-3082,
[73705] 1995.

[73706] 12575.

[73707] Kortschak, R. D.; Reimann, H.; Zimmer, M.; Eyre, H. J.; Saint,
[73708] R.; Jenne, D. E.: The human dead ringer/bright homolog, DRIL1: cDNA
[73709] cloning, gene structure, and mapping to D19S886, a marker on 19p13.3
[73710] that is strictly linked to the Peutz-Jeghers syndrome. Ge-
nomics 51:
[73711] 288-292, 1998.

[73712] 12576.

[73713] Kamei, M.; Webb, G. C.; Young, I. G.; Campbell, H. D.: SOLH, a
[73714] human homologue of the Drosophila melanogaster small optic lobes gene
[73715] is a member of the calpain and zinc-finger gene families and maps
[73716] to human chromosome 16p13.3 near CATM (cataract with microphthalmia). Genomics 51:
[73717] 197-206, 1998.

[73718] 12577.

[73719] Eriksson, I.; Sandback, D.; Ek, B.; Lindahl, U.; Kjellen, L.:

[73720] cDNA cloning and sequencing of mouse mastocytoma glucosaminyl N-deacetylase/N-sulfotransferase,

[73721] an enzyme involved in the biosynthesis of heparin. J. Biol. Chem. 269:

[73722] 10438-10443, 1994.

[73723] 12578.

[73724] Forsberg, E.; Pejler, G.; Ringvall, M.; Lunderius, C.; Tomasini-Johansson,

[73725] B.; Kusche-Gullberg, M.; Eriksson, I.; Ledin, J.; Hellman, L.; Kjellen,

[73726] L.: Abnormal mast cells in mice deficient in a heparin-synthesizing

[73727] enzyme. Nature 400: 773-776, 1999.

[73728] 12579.

[73729] Humphries, D. E.; Lanciotti, J.; Karlinsky, J. B.: cDNA cloning,

[73730] genomic organization and chromosomal localization of human heparan

[73731] glucosaminyl N-deacetylase/N-sulphotransferase-2.

Biochem. J. 332:
[73732] 303-307, 1998.

[73733] 12580.

[73734] Humphries, D. E.; Wong, G. W.; Friend, D. S.; Gurish, M. F.; Qiu,
[73735] W.-T.; Huang, C.; Sharpe, A. H.; Stevens, R. L.: Heparin is essential
[73736] for the storage of specific granule proteases in mast cells. Nature 400:
[73737] 769-772, 1999.

[73738] 12581.

[73739] Orellana, A.; Hirschberg, C. B.; Wei, Z.; Swiedler, S. J.; Ishi-hara,
[73740] M.: Molecular cloning and expression of a glycosamino-glycan N-acetylglucosaminyl
[73741] N-deacetylase/N-sulfotransferase from a heparin-pro-ducing cell line. J.
[73742] Biol. Chem. 269: 2270-2276, 1994.

[73743] 12582.

[73744] Zehnder, J. L.; Galli, S. J.: Mast-cell heparin demystified. Nature 400:

[73745] 714-715, 1999.

[73746] 12583.

[73747] Soret, J.; Gattoni, R.; Guyon, C.; Sureau, A.; Popielarz, M.; Le

[73748] Rouzic, E.; Dumon, S.; Apiou, F.; Dutrillaux, B.; Voss, H.; Ansorge,

[73749] W.; Stevenin, J.; Perbal, B.: Characterization of SRp46, a novel

[73750] human SR splicing factor encoded by a PR264/SC35 retropseudogene. Molec.

[73751] Cell. Biol. 18: 4924-4934, 1998.

[73752] 12584.

[73753] Buell, G. N.; Talabot, F.; Gos, A.; Lorenz, J.; Lai, E.; Morris,

[73754] M. A.; Antonarakis, S. E.: Gene structure and chromosomal localization

[73755] of the human P2X7 receptor. Receptors Channels 5: 347-354, 1998.

[73756] 12585.

[73757] Gu, B. J.; Zhang, W.; Worthington, R. A.; Sluyter, R.; Dao-Ung,

[73758] P.; Petrou, S.; Barden, J. A.; Wiley, J. S.: A glu-496 to ala

polymorphism

[73759] leads to loss of function of the human P2X(7) receptor. J. Biol.

[73760] Chem. 276: 11135-11142, 2001.

[73761] 12586.

[73762] Rassendren, F.; Buell, G. N.; Virginio, C.; Collo, G.; North, R.

[73763] A.; Surprenant, A.: The permeabilizing ATP receptor, P2X(7). J.

[73764] Biol. Chem. 272: 5482-5486, 1997.

[73765] 12587.

[73766] Solle, M.; Labasi, J.; Perregaux, D. G.; Stam, E.; Petrushova,

[73767] N.; Koller, B. H.; Griffiths, R. J.; Gabel, C. A.: Altered cytokine

[73768] production in mice lacking P2X7 receptors. J. Biol. Chem. 276: 125-132,

[73769] 2001.

[73770] 12588.

[73771] Wiley, J. S.; Dao-Ung, L. P.; Gu, B. J.; Sluyter, R.; Shemon, A.

[73772] N.; Li, C.; Taper, J.; Gallo, J.; Manoharan, A.: A loss-

of-function
[73773] polymorphic mutation in the cytolytic P2X7 receptor gene and chronic
[73774] lymphocyte leukaemia: a molecular study. Lancet 359: 1114-1119,
[73775] 2002.

[73776] 12589.

[73777] Worthington, R. A.; Smart, M. L.; Gu, B. J.; Williams, D. A.; Petrou,
[73778] S.; Wiley, J. S.; Barden, J. A.: Point mutations confer loss of ATP-induced
[73779] human P2X(7) receptor function. FEBS Lett. 512: 43-46, 2002.
[73780] 12590.

[73781] Hobbs, C. A.; Sherman, S. L.; Yi, P.; Hopkins, S. E.; Torfs, C.
[73782] P.; Hine, R. J.; Pogribna, M.; Rozen, R.; James, S. J.: Poly-morphisms
[73783] in genes involved in folate metabolism as maternal risk factors for
[73784] Down syndrome. Am. J. Hum. Genet. 67: 623-630, 2000.

[73785] 12591.

[73786] James, S. J.; Pogribna, M.; Pogribny, I. P.; Melnyk, S.; Hine,

[73787] R. J.; Gibson, J. B.; Yi, P.; Tafoya, D. L.; Swenson, D. H.; Wilson,

[73788] V. L.; Gaylor, D. W.: Abnormal folate metabolism and mutation in

[73789] the methylenetetrahydrofolate reductase gene may be maternal risk

[73790] factors for Down syndrome. Am. J. Clin. Nutr. 70: 495-501, 1999.

[73791] 12592.

[73792] Leclerc, D.; Wilson, A.; Dumas, R.; Gafuik, C.; Song, D.; Watkins,

[73793] D.; Heng, H. H. Q.; Rommens, J. M.; Scherer, S. W.; Rosenblatt, D.

[73794] S.; Gravel, R. A.: Cloning and mapping of a cDNA for methionine synthase

[73795] reductase, a flavoprotein defective in patients with homocystinuria. Proc.

[73796] Nat. Acad. Sci. 95: 3059-3064, 1998.

[73797] 12593.

[73798] Wilson, A.; Leclerc, D.; Rosenblatt, D. S.; Gravel, R. A.: Molecular

[73799] basis for methionine synthase reductase deficiency in patients belonging
[73800] to the cblE complementation group of disorders in folate/cobalamin
[73801] metabolism. Hum. Molec. Genet. 8: 2009-2016, 1999.

[73802] 12594.

[73803] Wilson, A.; Platt, R.; Wu, Q.; Leclerc, D.; Christensen, B.; Yang,
[73804] H.; Gravel, R. A.; Rozen, R.: A common variant in methionine synthase
[73805] reductase combined with low cobalamin (vitamin B12) increases risk
[73806] for spina bifida. Molec. Genet. Metab. 67: 317-323, 1999.

[73807] 12595.

[73808] Lavedan, C.; Buchholtz, S.; Auburger, G.; Albin, R. L.; Athanassiadou,
[73809] A.; Blancato, J.; Burguera, J. A.; Ferrell, R. E.; Kostic, V.; Leroy,
[73810] E.; Leube, B.; Mota-Vieira, L.; and 9 others: Absence of mutation
[73811] in the beta- and gamma-synuclein genes in familial autosomal dominant

[73812] Parkinson's disease. DNA Research 5: 401-402, 1998.

[73813] 12596.

[73814] Leek, J. P.; Hamlin, P. J.; Wilton, J.; Lench, N. J.: Assign-
ment
[73815] of the Rab13 gene (RAB13) to human chromosome band
12q13 by in situ
[73816] hybridization. Cytogenet. Cell Genet. 79: 210-211, 1997.

[73817] 12597.

[73818] Novick, P.; Zerial, M.: The diversity of Rab proteins in vesi-
cle
[73819] transport. Curr. Opin. Cell Biol. 9: 496-504, 1997.

[73820] 12598.

[73821] Salminen, A.; Novick, P. J.: A ras-like protein is required
for
[73822] a post-Golgi event in yeast secretion. Cell 49: 527-538,
1987.
[73823] 12599.

[73824] Zahraoui, A.; Joberty, G.; Arpin, M.; Fontaine, J. J.; Hellio,

[73825] R.; Tavitian, A.; Louvard, D.: A small rab GTPase is dis-
tributed

[73826] in cytoplasmic vesicles in nonpolarized cells but colocal-
izes with
[73827] the tight junction marker ZO-1 in polarized epithelial
cells. J.
[73828] Cell. Biol. 124: 101-115, 1994.

[73829] 12600.

[73830] Aurich-Costa, J.; Cadel, S.; Gouzy, C.; Foulon, T.; Cherif,
D.;
[73831] Cohen, P.: Assignment of the aminopeptidase B gene
(RNPEP) to human
[73832] chromosome 1 band q32 by in situ hybridization. Cyto-
genet. Cell Genet. 79:
[73833] 143-144, 1997.

[73834] 12601.

[73835] Cadel, S.; Foulon, T.; Viron, A.; Balogh, A.; Midol-Monnet,
S.;
[73836] Noel, N.; Cohen, P.: Aminopeptidase B from the rat testis
is a bifunctional
[73837] enzyme structurally related to leukotriene-A4 hydrolase.
Proc. Nat.
[73838] Acad. Sci. 94: 2963-2968, 1997.

[73839] 12602.

[73840] Cadel, S.; Pierotti, A. R.; Foulton, T.; Creminon, C.; Barre, N.;

[73841] Segretain, D.; Cohen, P.: Aminopeptidase-B in the rat testes: isolation,

[73842] functional properties and cellular localization in the seminiferous

[73843] tubules. Molec. Cell. Endocr. 110: 149-160, 1995.

[73844] 12603.

[73845] Hopsu, V. K.; Kantonen, U. M.; Glenner, G. G.: A peptidase from

[73846] rat tissues selectively hydrolyzing N-terminal arginine and lysine

[73847] residues. Life Sci. 3: 1449-1453, 1964.

[73848] 12604.

[73849] Jacquemin, P.; Martial, J. A.; Davidson, I.: Human TEF-5 is preferentially

[73850] expressed in placenta and binds to multiple functional elements of

[73851] the human chorionic somatomammotropin-B gene enhancer. J. Biol. Chem. 272:

[73852] 12928-12937, 1997.

[73853] 12605.

[73854] Jiang, S.-W.; Wu, K.; Eberhardt, N. L.: Human placental TEF-5

[73855] transactivates the human chorionic somatomammotropin gene enhancer. Molec.

[73856] Endocr. 13: 879-889, 1999.

[73857] 12606.

[73858] Bach, I.: The LIM domain: regulation by association. Mech. Dev. 91:

[73859] 5-17, 2000.

[73860] 12607.

[73861] Guy, P. M.; Kenny, D. A.; Gill, G. N.: The PDZ domain of the LIM

[73862] protein enigma binds to beta-tropomyosin. Molec. Biol. Cell 10:

[73863] 1973-1984, 1999.

[73864] 12608.

[73865] Kuroda, S.; Tokunaga, C.; Kiyohara, Y.; Higuchi, O.; Kon-ishi, H.;

[73866] Mizuno, K.; Gill, G. N.; Kikkawa, U.: Protein-protein interaction

[73867] of zinc finger LIM domains with protein kinase C. J. Biol. Chem. 271:

[73868] 31029-31032, 1996.

[73869] 12609.

[73870] Wu, R.-Y.; Gill, G. N.: LIM domain recognition of a tyrosine-containing

[73871] tight turn. J. Biol. Chem. 269: 25085-25090, 1994.

[73872] 12610.

[73873] Khoja, H.; Wang, G.; Ng, C.-T. L.; Tucker, J.; Brown, T.; Shyamala,

[73874] V.: Cloning of CCRL1, an orphan seven transmembrane receptor related

[73875] to chemokine receptors, expressed abundantly in the heart. Gene 246:

[73876] 229-238, 2000.

[73877] 12611.

[73878] Schweickart, V. L.; Epp, A.; Raport, C. J.; Gray, P. W.: CCR11

[73879] is a functional receptor for the monocyte chemoattractant

protein

[73880] family of chemokines. J. Biol. Chem. 275: 9550-9556, 2000. Note:

[73881] Erratum: J. Biol. Chem. 276: 856 only, 2001.

[73882] 12612.

[73883] Benn, P. A.; Shows, T. B.; D'Ancona, G. G.; Croce, C. M.; Orkwiszewski,

[73884] K. G.; Mellman, W. J.: Assignment of a gene for uridine diphosphate

[73885] galactose-4-epimerase to human chromosome 1 by somatic cell hybridization,

[73886] with evidence for a regional assignment to 1pter-1p21. Cytogenet.

[73887] Cell Genet. 24: 138-142, 1979.

[73888] 12613.

[73889] Lin, M. S.; Oizumi, J.; Ng, W. G.; Alfi, O. S.; Donnell, G. N.

[73890] : Regional mapping of the gene for human UDPGal 4-epimerase on chromosome

[73891] 1 in mouse-human hybrids. Cytogenet. Cell Genet. 24: 217-223, 1979.

[73892] 12614.

[73893] Lin, M. S.; Oizumi, J.; Ng, W. G.; Alfi, O. S.; Donnell, G. N.

[73894] : Assignment of UDP-gal-4-epimerase gene locus to chromosome 1 in

[73895] man. (Abstract) Am. J. Hum. Genet. 30: 132 only, 1978.

[73896] 12615.

[73897] Wang, W.; Zhou, Z.; Zhao, W.; Huang, Y.; Tang, R.; Ying, K.; Xie,

[73898] Y.; Mao, Y.: Molecular cloning, mapping and characterization of the

[73899] human neurocalcin delta gene (NCALD). Biochim. Biophys. Acta 1518:

[73900] 162-167, 2001.

[73901] 12616.

[73902] Gladyshev, V. N.; Liu, A.; Novoselov, S. V.; Krysan, K.; Sun, Q.-A.;

[73903] Kryukov, V. M.; Kryukov, G. V.; Lou, M. F.: Identification and characterization

[73904] of a new mammalian glutaredoxin (thioltransferase), Grx2. J. Biol.

[73905] Chem. 276: 30374-30380, 2001.

[73906] 12617.

[73907] Lundberg, M.; Johansson, C.; Chandra, J.; Enoksson, M.; Jacobsson,
[73908] G.; Ljung, J.; Johansson, M.; Holmgren, A.: Cloning and expression
[73909] of a novel human glutaredoxin (Grx2) with mitochondrial and nuclear
[73910] isoforms. J. Biol. Chem. 276: 26269-26275, 2001.

[73911] 12618.

[73912] Ungar, D.; Oka, T.; Brittle, E. E.; Vasile, E.; Lupashin, V. V.;

[73913] Chatterton, J. E.; Heuser, J. E.; Krieger, M.; Waters, M. G.: Characterization
[73914] of a mammalian Golgi-localized protein complex, COG, that is required
[73915] for normal Golgi morphology and function. J. Cell Biol. 157: 405-415,
[73916] 2002.

[73917] 12619.

[73918] Walter, D. M.; Paul, K. S.; Waters, M. G.: Purification and characterization
[73919] of a novel 13 S hetero oligomeric protein complex that stimulates

[73920] in vitro Golgi transport. J. Biol. Chem. 273: 29565-29576, 1998.

[73921] 12620.

[73922] Meuwissen, R. L. J.; Meerts, I.; Hoovers, J. M. N.; Leschot, N.

[73923] J.; Heyting, C.: Human synaptonemal complex protein 1 (SCP1): isolation

[73924] and characterization of the cDNA and chromosomal localization of the

[73925] gene. Genomics 39: 377-384, 1997.

[73926] 12621.

[73927] Meuwissen, R. L. J.; Offenberg, H. H.; Dietrich, A. J. J.; Riesewijk,

[73928] A.; van Iersel, M.; Heyting, C.: A coiled-coil related protein specific

[73929] for synapsed regions of meiotic prophase chromosomes. EMBO J. 11:

[73930] 5091-5100, 1992.

[73931] 12622.

[73932] Levedakou, E. N.; He, M.; Baptist, E. W.; Craven, R. J.; Cance,

[73933] W. G.; Welcsh, P. L.; Simmons, A.; Naylor, S. L.; Leach, R. L.; Lewis,

[73934] T. B.; Bowcock, A.; Liu, E. T.: Two novel human serine/threonine

[73935] kinases with homologies to the cell cycle regulating Xenopus MO15,

[73936] and NIMA kinases: cloning and characterization of their expression

[73937] pattern. Oncogene 9: 1977-1988, 1994.

[73938] 12623.

[73939] Kondoh, N.; Nishina, Y.; Tsuchida, J.; Koga, M.; Tanaka, H.; Uchida,

[73940] K.; Inazawa, J.; Taketo, M.; Nozaki, M.; Nojima, H.; Matsumiya, K.;

[73941] Namiki, M.; Okuyama, A.; Nishimune, Y.: Assignment of synaptonemal

[73942] complex protein 1 (SCP1) to human chromosome 1p13 by fluorescence

[73943] in situ hybridization and its expression in the testis. Cytogenet.

[73944] Cell Genet. 78: 103-104, 1997.

[73945] 12624.

[73946] Blondel, O.; Vandecasteele, G.; Gastineau, M.; Leclerc, S.; Dahmoune,
[73947] Y.; Langlois, M.; Fischmeister, R.: Molecular and functional characterization
[73948] of a 5-HT(4) receptor cloned from human atrium. FEBS Lett. 412:
[73949] 465-474, 1997.

[73950] 12625.

[73951] Claeysen, S.; Faye, P.; Sebben, M.; Lemaire, S.; Bockaert, J.;

[73952] Dumuis, A.; Taviaux, S.: Assignment of 5-hydroxytryptamine receptor
[73953] (HTR4) to human chromosome 5 bands q31-to-q33 by in situ hybridization. Cytogenet.
[73954] Cell Genet. 78: 133-134, 1997.

[73955] 12626.

[73956] Dumuis, A.; Bouhelal, R.; Sebben, M.; Cory, R.; Bockaert, J.:

[73957] A nonclassical 5-hydroxytryptamine receptor positively coupled with
[73958] adenylate cyclase in the central nervous system. Molec. Pharm. 34:
[73959] 880-887, 1988.

[73960] 12627.

[73961] Eglen, R. M.; Wong, E. H. F.; Dumuis, A.; Bockaert, J.: Cen-
tral
[73962] 5-HT4 receptors. Trends Pharm. Sci. 16: 391-398, 1995.

[73963] 12628.

[73964] Isomura, T.; Tamiya-Koizumi, K.; Suzuki, M.; Yoshida, S.;
Taniguchi,
[73965] M.; Matsuyama, M.; Ishigaki, T.; Sakuma, S.; Takahashi,
M.: RFP is
[73966] a DNA binding protein associated with the nuclear matrix.
Nucleic
[73967] Acids Res. 20: 5305-5310, 1992.

[73968] 12629.

[73969] Szpirer, C.; Szpirer, J.; Riviere, M.; Tazi, R.; Pontarotti, P.

[73970] : Mapping of the Olf89 and Rfp genes to the rat genome:
comparison
[73971] with the mouse and human and new insights into the evo-
lution of the
[73972] rodent genome. Cytogenet. Cell Genet. 78: 137-139,
1997.
[73973] 12630.

[73974] Adachi, O.; Kawai, T.; Takeda, K.; Matsumoto, M.; Tsutsui, H.;
[73975] Sakagami, M.; Nakanishi, K.; Akira, S.: Targeted disruption of the
[73976] MyD88 gene results in loss of IL-1- and IL-18-mediated function. Immunity 143-150,
[73977] 1998.

[73978] 12631.

[73979] Bonnert, T. P.; Garka, K. E.; Parnet, P.; Sonoda, G.; Testa, J.

[73980] R.; Sims, J. E.: The cloning and characterization of human MyD88:
[73981] a member of an IL-1 receptor related family. FEBS Lett. 402: 81-84,
[73982] 1997.

[73983] 12632.

[73984] Hardiman, G.; Jenkins, N. A.; Copeland, N. G.; Gilbert, D. J.;
[73985] Garcia, D. K.; Naylor, S. L.; Kastelein, R. A.; Bazan, J. F.: Genetic
[73986] structure and chromosomal mapping of MyD88. Genomics 45: 332-339,

[73987] 1997.

[73988] 12633.

[73989] Hayashi, F.; Smith, K. D.; Ozinsky, A.,; Hawn, T. R.; Yi, E. C.;

[73990] Goodlett, D. R.; Eng, J. K.; Akira, S.; Underhill, D. M.; Aderem,

[73991] A.: The innate immune response to bacterial flagellin is mediated

[73992] by Toll-like receptor 5. Nature 410: 1099-1103, 2001.

[73993] 12634.

[73994] Kawai, T.; Adachi, O.; Ogawa, T.; Takeda, K.; Akira, S.: Un-responsiveness

[73995] of MyD88-deficient mice to endotoxin. Immunity 11: 115-122, 1999.

[73996] 12635.

[73997] Lord, K. A.; Hoffman-Liebermann, B.; Liebermann, D. A.: Complexity

[73998] of the immediate early response of myeloid cells to termi-nal differentiation

[73999] and growth arrest includes ICAM-1, Jun-B and histone variants. Oncogene 5:

[74000] 387-396, 1990.

[74001] 12636.

[74002] Medzhitov, R.; Preston-Hurlburt, P.; Kopp, E.; Stadien, A.; Chen,

[74003] C.; Ghosh, S.; Janeway, C. A., Jr.: MyD88 is an adaptor protein in

[74004] the hToll/Il-1 receptor family signaling pathways. Molec. Cell 2:

[74005] 253-258, 1998.

[74006] 12637.

[74007] Ge, K.; Guermah, M.; Yuan, C.-X.; Ito, M.; Wallberg, A. E.; Spiegelman,

[74008] B. M.; Roeder, R. G.: Transcription coactivator TRAP220 is required

[74009] for PPAR-gamma-2-stimulated adipogenesis. Nature 417: 563-567, 2002.

[74010] 12638.

[74011] Greene, M. E.; Blumberg, B.; McBride, O. W.; Yi, H. F.; Kro-nquist,

[74012] K.; Kwan, K.; Hsieh, L.; Greene, G.; Nimer, S. D.: Isolation of the

[74013] human peroxisome proliferator activated receptor gamma cDNA: expression
[74014] in hematopoietic cells and chromosomal mapping. Gene Expr. 4: 281-299,
[74015] 1995.

[74016] 12639.

[74017] Hara, M.; Alcoser, S. Y.; Qaadir, A.; Beiswenger, K. K.; Cox,

[74018] N. J.; Ehrmann, D. A.: Insulin resistance is attenuated in women
[74019] with polycystic ovary syndrome with the Pro12Ala polymorphism in the
[74020] PPAR-gamma gene. J. Clin. Endocr. Metab. 87: 772-775, 2002.
[74021] 12640.

[74022] Harris, S. G.; Phipps, R. P.: Prostaglandin D2, its metabolite
[74023] 15-d-PGJ2, and peroxisome proliferator activated receptor-gamma agonists
[74024] induce apoptosis in transformed, but not normal, human T lineage cells. Immunology 105:
[74025] 23-34, 2002.

[74026] 12641.

[74027] Hasstedt, S. J.; Ren, Q.-F.; Teng, K.; Elbein, S. C.: Effect

[74028] of the peroxisome proliferator-activated receptor-gamma-2 Pro12Ala

[74029] variant on obesity, glucose homeostasis, and blood pressure in members

[74030] of familial type 2 diabetic kindreds. J. Clin. Endocr. Metab. 86:

[74031] 536-541, 2001.

[74032] 12642.

[74033] Dabovic, B.; Chen, Y.; Colarossi, C.; Obata, H.; Zambuto, L.; Perle,

[74034] M. A.; Rifkin, D. B.: Bone abnormalities in latent TGF-beta binding

[74035] protein (Ltbp)-3-null mice indicate a role for Ltbp-3 in modulating

[74036] TGF-beta bioavailability. J. Cell Biol. 156: 227-232, 2002.

[74037] 12643.

[74038] Li, X.; Yin, W.; Perez-Jurado, L.; Bonadio, J.; Francke, U.: Mapping

[74039] of human and murine genes for latent TGF-beta binding

protein-2 (LTBP2). Mammalian
[74040] Genome 6: 42-45, 1995.

[74041] 12644.

[74042] Sawicki, M.; Arnold, E.; Ebrahimi, S.; Duell, T.; Jin, S.; Wood,
[74043] T.; Chakrabarti, R.; Peters, J.; Wan, Y.; Samara, G.; Weier, H.-U.
[74044] G.; Udar, N.; Passaro, E., Jr.; Srivatsan, E. S.: A transcript map
[74045] encompassing the multiple endocrine neoplasia type-1 (MEN1) locus
[74046] on chromosome 11q13. Genomics 42: 405-412, 1997.

[74047] 12645.

[74048] Yin, W.; Smiley, E.; Germiller, J.; Mechan, R. P.; Florer, J. B.;
[74049] Wenstrup, R. J.; Bonadio, J.: Isolation of a novel latent transforming
[74050] growth factor-beta binding protein gene (LTBP-3). J. Biol. Chem. 270:
[74051] 10147-10160, 1995.

[74052] 12646.

[74053] Moren, A.; Olofsson, A.; Stenman, G.; Sahlin, P.; Kanzaki,

T.;

[74054] Claesson-Welsh, L.; ten Dijke, P.; Miyazono, K.; Heldin, C.-H.: Identification

[74055] and characterization of LTBP-2, a novel latent transforming growth

[74056] factor-beta-binding protein. J. Biol. Chem. 269: 32469-32478, 1994.

[74057] 12647.

[74058] Fathallah, D. M.; Cherif, D.; Dellagi, K.; Arnaout, M. A.: Molecular

[74059] cloning of a novel human hsp70 from a B cell line and its assignment

[74060] to chromosome 5. J. Immun. 151: 810-813, 1993. Note: Erratum: J.

[74061] Immun. 151: 6616 only, 1993.

[74062] 12648.

[74063] Smith, S. A.; Holik, P.; Stevens, J.; Mazoyer, S.; Melis, R.; Williams,

[74064] B.; White, R.; Albertsen, H.: Isolation of a gene (DLG3) encoding

[74065] a second member of the discs-large family on chromosome 17q12-q21. Genomics 31:

[74066] 145-150, 1996.

[74067] 12649.

[74068] Wu, S.; Wright, R. A.; Rockey, P. K.; Burgett, S. G.; Arnold, J.

[74069] S.; Rosteck, P. R., Jr.; Johnson, B. G.; Schoepp, D. D.; Bela-gaje,

[74070] R. M.: Group III human metabotropic glutamate receptors 4, 7 and

[74071] 8: molecular cloning, functional expression, and compari-son of pharmacological

[74072] properties in RGT cells. Molec. Brain Res. 53: 88-97, 1998.

[74073] 12650.

[74074] Bram, R. J.; Crabtree, G. R.: Calcium signalling in T cells stimulated

[74075] by a cyclophilin B-binding protein. Nature 371: 355-358, 1994.

[74076] 12651.

[74077] Bram, R. J.; Valentine, V.; Shapiro, D. N.; Jenkins, N. A.; Gilbert,

[74078] D. J.; Copeland, N. G.: The gene for calcium-modulating cyclophilin

[74079] ligand (CAMLG) is located on human chromosome 5q23 and a syntenic
[74080] region of mouse chromosome 13. Genomics 31: 257-260, 1996.
[74081] 12652.

[74082] Bussemakers, M. J. G.; van Bokhoven, A.; Voller, M.; Smit, F. P.;
[74083] Schalken, J. A.: The genes for the calcium-dependent cell adhesion
[74084] molecules P- and E-cadherin are tandemly arranged in the human genome. Biochem.
[74085] Biophys. Res. Commun. 203: 1291-1294, 1994.

[74086] 12653.

[74087] Carmeliet, P.; Lampugnani, M.-G.; Moons, L.; Breviario, F.; Compernolle,
[74088] V.; Bono, F.; Balconi, G.; Spagnuolo, R.; Oosthuyse, B.; Dewerchin,
[74089] M.; Zanetti, A.; Angellilo, A.; and 11 others: Targeted deficiency
[74090] of cytosolic truncation of the VE-cadherin gene in mice impairs VEGF-mediated
[74091] endothelial survival and angiogenesis. Cell 98: 147-157,

1999.

[74092] 12654.

[74093] Huber, P.; Dalmon, J.; Engiles, J.; Breviario, F.; Gory, S.; Siracusa,

[74094] L. D.; Buchberg, A. M.; Dejana, E.: Genomic structure and chromosomal

[74095] mapping of the mouse VE-cadherin gene (Cdh5). Genomics 32: 21-28,

[74096] 1996.

[74097] 12655.

[74098] Salomon, D.; Ayalon, O.; Patel-King, R.; Hynes, R. O.; Geiger,

[74099] B.: Extrajunctional distribution of N-cadherin in cultured human

[74100] endothelial cells. J. Cell Sci. 102: 7-17, 1992.

[74101] 12656.

[74102] Stock, J. L.; Brown, R. S.; Baron, J.; Coderre, J. A.; Mancilla,

[74103] E.; De Luca, F.; Ray, K.; Mericq, M. V.: Autosomal dominant hypoparathyroidism

[74104] associated with short stature and premature osteoarthritis. J. Clin.

[74105] Endocr. Metab. 84: 3036-3040, 1999.

[74106] 12657.

[74107] Yamamoto, M.; Akatsu, T.; Nagase, T.; Ogata, E.: Comparison of

[74108] hypocalcemic hypercalciuria between patients with idiopathic hypoparathyroidism

[74109] and those with gain-of-function mutations in the calcium-sensing receptor:

[74110] is it possible to differentiate the two disorders? J. Clin. Endocr.

[74111] Metab. 85: 4583-4591, 2000.

[74112] 12658.

[74113] Charrin, S.; Le Naour, F.; Oualid, M.; Billard, M.; Faure, G.,

[74114] Hanash, S. M.; Boucheix, C.; Rubinstein, E.: The major CD9 and CD81

[74115] molecular partner: identification and characterization of the complexes. J.

[74116] Biol. Chem. 276: 14329-14337, 2001.

[74117] 12659.

[74118] Nagase, T.; Kikuno, R.; Ishikawa, K.; Hirosawa, M.; Ohara, O.:

[74119] Prediction of the coding sequences of unidentified human genes. XVI.

[74120] The complete sequences of 150 new cDNA clones from brain which code

[74121] for large proteins in vitro. DNA Res. 7: 65-73, 2000.

[74122] 12660.

[74123] Orlicky, D. J.; Berry, R.; Sikela, J. M.: Human chromosome 1 localization

[74124] of the gene for a prostaglandin F-2-alpha receptor negative regulatory

[74125] protein. Hum. Genet. 97: 655-658, 1996.

[74126] 12661.

[74127] Stipp, C. S.; Orlicky, D.; Hemler, M. E.: FPRP, a major, highly

[74128] stoichiometric, highly specific CD81- and CD9-associated protein. J.

[74129] Biol. Chem. 276: 4853-4862, 2001.

[74130] 12662.

[74131] Buller, C.; Xu, X.; Marquis, V.; Schwanke, R.; Xu, P.-X.: Molecular

[74132] effects of Eya1 domain mutations causing organ defects in

BOR syndrome. Hum.
[74133] Molec. Genet. 10: 2775-2781, 2001.

[74134] 12663.

[74135] Oliver, G.; Wehr, R.; Jenkins, N. A.; Copeland, N. G.; Cheyette,
[74136] B. N. R.; Hartenstein, V.; Zipursky, S. L.; Gruss, P.: Homeobox genes
[74137] and connective tissue patterning. Development 121: 693-705, 1995.
[74138] 12664.

[74139] Casellas, R.; Jankovic, M.; Meyer, G.; Gazumyan, A.; Luo, Y.; Roeder,
[74140] R. G.; Nussenzweig, M. C.: OcaB is required for normal transcription
[74141] and V(D)J recombination of a subset of immunoglobulin kappa genes. Cell 110:
[74142] 575-585, 2002.

[74143] 12665.

[74144] Gstaiger, M.; Knoepfel, L.; Georgiev, O.; Schaffner, W.; Hovens,
[74145] C. M.: A B-cell coactivator of octamer-binding transcrip-

tion factors. Nature 373:

[74146] 360-362, 1995.

[74147] 12666.

[74148] Junker, S.; Brondum-Nielsen, K.; Newell, J. W.; Matthias, P.; Tommerup,

[74149] N.: Assignment of the human gene for Oct-binding factor-1 (OBF1),

[74150] a B-cell-specific coactivator of octamer-binding transcription factors

[74151] 1 and 2, to 11q23.1 by somatic cell hybridization and in situ hybridization. Genomics 33:

[74152] 143-145, 1996.

[74153] 12667.

[74154] Holt, K. H.; Lim, L. E.; Straub, V.; Venzke, D. P.; Duclos, F.;

[74155] Anderson, R. D.; Davidson, B. L.; Campbell, K. P.: Functional rescue

[74156] of the sarcoglycan complex in the BIO 14.6 hamster using delta-sarcoglycan

[74157] gene transfer. Molec. Cell 1: 841-848, 1998.

[74158] 12668.

[74159] Moreira, E. S.; Vainzof, M.; Marie, S. K.; Nigro, V.; Zatz, M.;

[74160] Passos-Bueno, M. R.: A first missense mutation in the delta-sarcoglycan
[74161] gene associated with a severe phenotype and frequency of limb-girdle
[74162] muscular dystrophy type 2F (LGMD2F) in Brazilian sarco-glycanopathies. J.
[74163] Med. Genet. 35: 951-953, 1998.

[74164] 12669.

[74165] Nigro, V.; Moreira, E. S.; Piluso, G.; Vainzof, M.; Belsito, A.;

[74166] Politano, L.; Puca, A. A.; Passos-Bueno, M. R.; Zatz, M.: Autosomal
[74167] recessive limb-girdle muscular dystrophy, LGMD2F, is caused by a mutation
[74168] in the delta-sarcoglycan gene. Nature Genet. 14: 195-198, 1996.
[74169] 12670.

[74170] Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.; Herschman,
[74171] H.; Witte, O.; Colicelli, J.: Protein binding and signaling properties
[74172] of RIN1 suggest a unique effector function. Proc. Nat. Acad. Sci. 94:

[74173] 4954-4959, 1997.

[74174] 12671.

[74175] Leffers, H.; Madsen, P.; Rasmussen, H. H.; Honore, B.; Andersen,

[74176] A. H.; Walbum, E.; Vandekerckhove, J.; Celis, J. E.: Molecular cloning

[74177] and expression of the transformation sensitive epithelial marker stratifin:

[74178] a member of a protein family that has been involved in the protein

[74179] kinase C signalling pathway. J. Molec. Biol. 231: 982-998, 1993.

[74180] 12672.

[74181] Jaspers, M.; Marynen, P.; Aly, M. S.; Cuppens, H.; Hilliker, C.;

[74182] Cassiman, J.-J.: Localization of the gene encoding the alpha-2 subunit

[74183] of the human VLA-2 receptor to chromosome 5q23-31. Somat. Cell Molec.

[74184] Genet. 17: 505-511, 1991.

[74185] 12673.

[74186] Yang, Y.; Hentati, A.; Deng, H.-X.; Dabbagh, O.; Sasaki, T.; Hirano,

[74187] M.; Hung, W.-Y.; Ouahchi, K.; Yan, J.; Azim, A. C.; Cole, N.; Gascon,

[74188] G.; Yagmour, A.; Ben-Hamida, M.; Pericak-Vance, M.; Hentati, F.; Siddique,

[74189] T.: The gene encoding alsin, a protein with three guanine-nucleotide

[74190] exchange factor domains, is mutated in a form of recessive amyotrophic

[74191] lateral sclerosis. Nature Genet. 29: 160-165, 2001. Note: Erratum:

[74192] Nature Genet. 29: 352 only, 2001.

[74193] 12674.

[74194] Akanuma, J.; Nishigaki, T.; Fujii, K.; Matsubara, Y.; Inui, K.;

[74195] Takahashi, K.; Kure, S.; Suzuki, Y.; Ohura, T.; Miyabayashi, S.; Ogawa,

[74196] E.; Iinuma, K.; Okada, S.; Narisawa, K.: Glycogen storage disease

[74197] type Ia: molecular diagnosis of 51 Japanese patients and characterization

[74198] of splicing mutations by analysis of ectopically transcribed

mRNA
[74199] from lymphoblastoid cells. Am. J. Med. Genet. 91:
107-112, 2000.
[74200] 12675.

[74201] Hammond, J. W.; Potter, M.; Wilcken, B.; Truscott, R.: Siblings
[74202] with gamma-glutamyltransferase deficiency. J. Inherit. Metab. Dis. 18:
[74203] 82-83, 1995.

[74204] 12676.

[74205] Heisterkamp, N.; Groffen, J.: Duplication of the bcr and gamma-glutamyl
[74206] transpeptidase genes. Nucleic Acids Res. 16: 8045-8056, 1988.
[74207] 12677.

[74208] Laperche, Y.; Bulle, F.; Aissani, T.; Chobert, M. N.; Aggerbeck,
[74209] M.; Hanoune, J.; Guellaen, G.: Molecular cloning and nucleotide sequence
[74210] of rat kidney gamma-glutamyl transpeptidase cDNA. Proc. Nat. Acad.
[74211] Sci. 83: 937-941, 1986.

[74212] 12678.

[74213] O'Daley, S.: An abnormal sulphydryl compound in urine. (Abstract) Irish

[74214] J. Med. Sci. 7: 578-579, 1968.

[74215] 12679.

[74216] Pawlak, A.; Lahuna, O.; Bulle, F.; Suzuki, A.; Ferry, N.; Siegrist,

[74217] S.; Chikhi, N.; Chobert, M. N.; Guellaen, G.; Laperche, Y.: Gamma-glutamyl

[74218] transpeptidase: a single copy gene in the rat and a multi-gene family

[74219] in the human genome. J. Biol. Chem. 263: 9913-9916, 1988.

[74220] 12680.

[74221] Rouleau, G. A.; Bazanowski, A.; Cohen, E. H.; Guellaen, G.; Gusella,

[74222] J. F.: Gamma-glutamyl transferase locus (GGT) displays a PvuII polymorphism. Nucleic

[74223] Acids Res. 16: 11848 only, 1988.

[74224] 12681.

[74225] Sakamuro, D.; Yamazoe, M.; Matsuda, Y.; Kangawa, K.;

Taniguchi,
[74226] N.; Matsuo, H.; Yoshikawa, H.; Ogasawara, N.: The primary structure
[74227] of human gamma-glutamyl transpeptidase. Gene 73: 1-9, 1988.
[74228] 12682.

[74229] Schulman, J. D.; Goodman, S. I.; Mace, J. W.; Patrick, A. D.;

[74230] Tietze, F.; Butler, E. J.: Glutathionuria: inborn error of metabolism
[74231] due to tissue deficiency of gamma-glutamyl transpeptidase. Biochem.
[74232] Biophys. Res. Commun. 65: 68-74, 1975.

[74233] 12683.

[74234] Jaeken, J.; Goemans, N.; Fryns, J.-P.; Francois, I.; de Zegher,
[74235] F.: Association of hyperprolinaemia type I and heparin cofactor II
[74236] deficiency with CATCH 22 syndrome: evidence for a contiguous gene
[74237] syndrome locating the proline oxidase gene. J. Inherit. Metab. Dis. 19:
[74238] 275-277, 1996.

[74239] 12684.

[74240] Fernie, B. A.; Wurzner, R.; Orren, A.; Morgan, B. P.; Potter,

[74241] P. C.; Platonov, A. E.; Vershinina, I. V.; Shipulin, G. A.;
Lachmann,

[74242] P. J.; Hobart, M. J.: Molecular bases of combined subtotal
deficiencies

[74243] of C6 and C7: their effects in combination with other C6
and C7 deficiencies. J.

[74244] Immun. 157: 3648-3657, 1996.

[74245] 12685.

[74246] Berard-Badier, M.; Adechy-Benkoel, L.; Chamlian, A.;
Dubois-Gambarelli,

[74247] D.; Casanova, P.; Mariani, A.: Etude ultrastructurale du
parenchyme

[74248] hepatique dans les mucopolysaccharidoses. Path. Biol.
(Paris) 18:

[74249] 117-128, 1970.

[74250] 12686.

[74251] Hoogeveen, A. T.; Verheijen, F. W.; d'Azzo, A.; Galjaard,
H.:

[74252] Genetic heterogeneity in human neuraminidase deficiency.

Nature 285:
[74253] 500-502, 1980.

[74254] 12687.

[74255] Li, Y.-T.; Muhiudeen, I. A.; DeGasperi, R.; Hirabayashi, Y.; Li,
[74256] S.-C.: Presence of activator proteins for the enzymic hydrolysis
[74257] of GM1 and GM2 gangliosides in normal human urine. Am. J. Hum. Genet. 35:
[74258] 629-634, 1983.

[74259] 12688.

[74260] Lowden, J. A.; O'Brien, J. S.: Sialidosis: a review of human
[74261] neuraminidase deficiency. Am. J. Hum. Genet. 31: 1-18, 1979.
[74262] 12689.

[74263] Delepine, M.; Nicolino, M.; Barrett, T.; Golamaully, M.; Lathrop,
[74264] G. M.; Julier, C.: EIF2AK3, encoding translation initiation factor
[74265] 2-alpha kinase 3, is mutated in patients with Wolcott-Rallison syndrome. Nature

[74266] Genet. 25: 406-409, 2000.

[74267] 12690.

[74268] Vasconcelos, O.; Sivakumar, K.; Dalakas, M. C.; Quezado, M.; Nagle,

[74269] J.; Leon-Monzon, M.; Dubnick, M.; Gajdusek, D. C.; Goldfarb, L. G.

[74270] : Nonsense mutation in the phosphofructokinase muscle subunit gene

[74271] associated with retention of intron 10 in one of the isolated transcripts

[74272] in Ashkenazi Jewish patients with Tarui disease. Proc. Nat. Acad.

[74273] Sci. 92: 10322-10326, 1995.

[74274] 12691.

[74275] Vora, S.; Corash, L.; Engel, W. K.; Durham, S.; Seaman, C.; Piomelli,

[74276] S.: The molecular mechanism of the inherited phosphofructokinase

[74277] deficiency associated with hemolysis and myopathy. Blood 55: 629-635,

[74278] 1980.

[74279] 12692.

[74280] Vora, S.; Durham, S.; de Martinville, B.; Francke, U.: As-
signment
[74281] of the human gene for muscle-type phosphofructokinase
(PFKM) to chromosome
[74282] 1 (region cen-q32) using somatic cell hybrids and mono-
clonal anti-M
[74283] antibody. Somat. Cell Genet. 8: 95-104, 1982.

[74284] 12693.

[74285] Vora, S.; Giger, U.; Turchen, S.; Harvey, J. W.: Characteri-
zation
[74286] of the enzymatic lesion in inherited phosphofructokinase
deficiency
[74287] in the dog: an animal analogue of human glycogen stor-
age disease type
[74288] VII. Proc. Nat. Acad. Sci. 82: 8109-8113, 1985.

[74289] 12694.

[74290] Vorgerd, M.; Karitzky, J.; Ristow, M.; Van Schaftingen, E.;
Tegenthoff,
[74291] M.; Jerusalem, F.; Malin, J. P.: Muscle phosphofructokinase
deficiency

[74292] in two generations. J. Neurol. Sci. 141: 95-99, 1996.

[74293] 12695.

[74294] Yamasaki,; Nakajima, H.; Kono, N.; Hotta, K.; Yamada, K.; Imai,

[74295] E.; Kuwajima, M.; Noguchi, T.; Tanaka, T.; Tarui, S.: Struc-ture of

[74296] the entire human muscle phosphofructokinase-encoding gene: a two-promoter

[74297] system. Gene 104: 277-282, 1991.

[74298] 12696.

[74299] Zanella, A.; Mariani, M.; Meola, G.; Fagnani, G.; Sirchia, G.

[74300] : Phosphofructokinase (PFK) deficiency due to a catalyti-cally inactive

[74301] mutant M-type subunit. Am. J. Hemat. 12: 215-225, 1982.

[74302] 12697.

[74303] Toda, T.; Yoshioka, M.; Nakahori, Y.; Kanazawa, I.; Naka-mura,

[74304] Y.; Nakagome, Y.: Genetic identity of Fukuyama-type con-genital muscular

[74305] dystrophy and Walker-Warburg syndrome. Ann. Neurol.

37: 99-101,

[74306] 1995.

[74307] 12698.

[74308] Aoufouchi, S.; Flatter, E.; Dahan, A.; Faili, A.; Bertocci, B.;

[74309] Storck, S.; Delbos, F.; Cocea, L.; Gupta, N.; Weill, J.-C.; Reynaud,

[74310] C.-A.: Two novel human and mouse DNA polymerases of the polX family. Nucleic

[74311] Acids Res. 28: 3684-3693, 2000.

[74312] 12699.

[74313] Murano, S.; Thweatt, R.; Reis, R. J. S.; Jones, R. A.; Moer-man,

[74314] E. J.; Goldstein, S.: Diverse gene sequences are overex-pressed in

[74315] Werner syndrome fibroblasts undergoing premature replicative senescence. Molec.

[74316] Cell. Biol. 11: 3905-3914, 1991.

[74317] 12700.

[74318] Harris, E. D.: Cellular copper transport and metabolism. Annu.

[74319] Rev. Nutr. 20: 291-310, 2000.

[74320] 12701.

[74321] Houwen, R. H. J.; Juyn, J.; Hoogenraad, T. U.; Ploos van Amstel,
[74322] J. K.; Berger, R.: H714Q mutation in Wilson disease is associated
[74323] with late, neurological presentation. J. Med. Genet. 32: 480-482,
[74324] 1995.

[74325] 12702.

[74326] Kusuda, Y.; Hamaguchi, K.; Mori, T.; Shin, R.; Seike, M.; Sakata,
[74327] T.: Novel mutations of the ATP7B gene in Japanese patients with Wilson
[74328] disease. J. Hum. Genet. 45: 86-91, 2000.

[74329] 12703.

[74330] La Fontaine, S.; Theophilos, M. B.; Firth, S. D.; Gould, R.; Parton,
[74331] R. G.; Mercer, J. F. B.: Effect of the toxic milk mutation (tx) on
[74332] the function and intracellular localization of Wnd, the murine homologue

[74333] of the Wilson copper ATPase. Hum. Molec. Genet. 10: 361-370, 2001.

[74334] 12704.

[74335] Payne, A. S.; Kelly, E. J.; Gitlin, J. D.: Functional expression

[74336] of the Wilson disease protein reveals mislocalization and impaired

[74337] copper-dependent trafficking of the common H1069Q mutation. Proc.

[74338] Nat. Acad. Sci. 95: 10854-10859, 1998.

[74339] 12705.

[74340] Petrukhin, K. E.; Lutsenko, S.; Chernov, I.; Ross, B. M.; Ka-plan,

[74341] J. H.; Gilliam, T. C.: Characterization of the Wilson disease gene

[74342] encoding a P-type copper transporting ATPase: genomic organization,

[74343] alternative splicing, and structure/function predictions. Hum. Molec.

[74344] Genet. 3: 1647-1656, 1994.

[74345] 12706.

[74346] Reed, V.; Williamson, P.; Bull, P. C.; Cox, D. W.; Boyd, Y.:

[74347] Mapping of the mouse homologue of the Wilson disease gene to mouse

[74348] chromosome 8. Genomics 28: 573-575, 1995.

[74349] 12707.

[74350] Wilson, D. C.; Phillips, M. J.; Cox, D. W.; Roberts, E. A.: Severe

[74351] hepatic Wilson's disease in preschool-aged children. J. Pediat. 137:

[74352] 719-722, 2000.

[74353] 12708.

[74354] Wu, Z.-Y.; Wang, N.; Lin, M.-T.; Fang, L.; Murong, S.-X.; Yu,

[74355] L.: Mutation analysis and the correlation between genotype and phenotype

[74356] of arg778leu mutation in Chinese patients with Wilson disease. Arch.

[74357] Neurol. 58: 971-976, 2001.

[74358] 12709.

[74359] Li, S.; Strelow, A.; Fontana, E. J.; Wesche, H.: IRAK-4: a novel

[74360] member of the IRAK family with the properties of an IRAK-

kinase. Proc.

[74361] Nat. Acad. Sci. 99: 5567-5572, 2002.

[74362] 12710.

[74363] Scanlan, M. J.; Gordon, J. D.; Williamson, B.; Stockert, E.; Bander,

[74364] N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.; Knuth, A.;

[74365] Chen, Y.-T.; Old, L. J.: Antigens recognized by autologous antibody

[74366] in patients with renal-cell carcinoma. Int. J. Cancer 83: 456-464,

[74367] 1999.

[74368] 12711.

[74369] Scott, A. F.: Personal Communication. Baltimore, Md. 4/25/2002.

[74370] 12712.

[74371] Suzuki, N.; Suzuki, S.; Duncan, G. S.; Millar, D. G.; Wada, T.;

[74372] Mirtsos, C.; Takada, H.; Wakeham, A.; Itie, A.; Li, S.; Pen-ninger,

[74373] J. M.; Wesche, H.; Ohashi, P. S.; Mak, T. W.; Yeh, W.-C.:

Severe
[74374] impairment of interleukin-1 and Toll-like receptor signalling in mice
[74375] lacking IRAK-4. Nature 416: 750-754, 2002.

[74376] 12713.

[74377] O'Brien, J. S.: Suggestions for a nomenclature for the GM2-gangliosidoses
[74378] making certain (possibly unwarranted) assumptions. (Comments). Am.
[74379] J. Hum. Genet. 30: 672-675, 1978.

[74380] 12714.

[74381] Ohno, K.; Suzuki, K.: A splicing defect due to an exon-intron
[74382] junctional mutation results in abnormal beta-hexosaminidase alpha
[74383] chain mRNAs in Ashkenazi Jewish patients with Tay-Sachs disease. Biochem.
[74384] Biophys. Res. Commun. 153: 463-469, 1988.

[74385] 12715.

[74386] Ohno, K.; Suzuki, K.: Multiple abnormal beta-hexosaminidase alpha

[74387] chain mRNAs in a compound-heterozygous Ashkenazi Jewish patient with
[74388] Tay-Sachs disease. J. Biol. Chem. 263: 18563-18567, 1988.
[74389] 12716.

[74390] Ohno, K.; Suzuki, K.: Mutation in GM2-gangliosidosis B1 variant. J.
[74391] Neurochem. 50: 316-318, 1988.

[74392] 12717.

[74393] Paw, B. H.; Kaback, M. M.; Neufeld, E. F.: Molecular basis of
[74394] adult-onset and chronic G(M2) gangliosidoses in patients of Ashkenazi
[74395] Jewish origin: substitution of serine for glycine at position 269
[74396] of the alpha-subunit of beta-hexosaminidase. Proc. Nat. Acad. Sci. 86:
[74397] 2413-2417, 1989.

[74398] 12718.

[74399] Paw, B. H.; Moskowitz, S. M.; Uhrhammer, N.; Wright, N.; Kaback,

[74400] M. M.; Neufeld, E. F.: Juvenile G(M2) gangliosidosis caused by substitution

[74401] of histidine for arginine at position 499 or 504 of the alpha-subunit

[74402] of beta-hexosaminidase. J. Biol. Chem. 265: 9452-9457, 1990.

[74403] 12719.

[74404] Paw, B. H.; Wood, L. C.; Neufeld, E. F.: A third mutation at

[74405] the CpG dinucleotide of codon 504 and a silent mutation at codon 506

[74406] of the HEX A gene. Am. J. Hum. Genet. 48: 1139-1146, 1991.

[74407] 12720.

[74408] Pennybacker, M.; Liessem, B.; Moczall, H.; Tifft, C. J.; Sandhoff,

[74409] K.; Proia, R. L.: Identification of domains in human beta-hexosaminidase

[74410] that determine substrate specificity. J. Biol. Chem. 271: 17377-17382,

[74411] 1996.

[74412] 12721.

[74413] Petersen, G. M.; Rotter, J. I.; Cantor, R. M.; Field, L. L.; Greenwald,

[74414] S.; Lim, J. S. T.; Roy, C.; Schoenfeld, V.; Lowden, J. A.; Kaback,

[74415] M. M.: The Tay-Sachs disease gene in North American Jewish populations:

[74416] geographic variations and origin. Am. J. Hum. Genet. 35: 1258-1269,

[74417] 1983.

[74418] 12722.

[74419] Petroulakis, E.; Cao, Z.; Clarke, J. T. R.; Mahuran, D. J.; Lee,

[74420] G.; Triggs-Raine, B.: W474C amino acid substitution affects early

[74421] processing of the alpha-subunit of beta-hexosaminidase A and is associated

[74422] with subacute G(M2) gangliosidosis. Hum. Mutat. 11: 432-442, 1998.

[74423] 12723.

[74424] Proia, R. L.; Kolodny, E. H.; Navon, R.: Reply to Kappler et

[74425] al. (Letter) Am. J. Hum. Genet. 47: 881-882, 1990.

[74426] 12724.

[74427] Proia, R. L.; Neufeld, E. F.: Synthesis of beta-hexosaminidase

[74428] in cell-free translation and in intact fibroblasts: an insoluble precursor

[74429] alpha chain in a rare form of Tay-Sachs disease. Proc. Nat. Acad.

[74430] Sci. 79: 6360-6364, 1982.

[74431] 12725.

[74432] Proia, R. L.; Soravia, E.: Organization of the gene encoding

[74433] the human beta-hexosaminidase alpha-chain. J. Biol. Chem. 262: 5677-5681,

[74434] 1987.

[74435] 12726.

[74436] Raghavan, S. S.; Krusell, A.; Krusell, J.; Lyerla, T. A.; Kolodny,

[74437] E. H.: G(M2)-ganglioside metabolism in hexosaminidase A deficiency

[74438] states: determination in situ using labeled G(M2) added to fibroblast

[74439] cultures. Am. J. Hum. Genet. 37: 1071-1082, 1985.

[74440] 12727.

[74441] Shore, S.; Tomczak, J.; Grebner, E. E.; Myerowitz, R.: An unusual

[74442] genotype in an Ashkenazi Jewish patient with Tay-Sachs disease. Hum.

[74443] Mutat. 1: 486-490, 1992.

[74444] 12728.

[74445] Strasberg, P.; Warren, I.; Skomorowski, M.-A.; Feigen-
baum, A.

[74446] : Homozygosity for the common Ashkenazi Jewish Tay-
Sachs +1 IVS-12

[74447] splice-junction mutation: first report. Hum. Mutat. 10:
82-83, 1997.

[74448] 12729.

[74449] Suzuki, Y.; Suzuki, K.: Partial deficiency of hex-
osaminidase

[74450] component A in juvenile Gm(2)-gangliosidosis. Neurology
20: 848-851,

[74451] 1970.

[74452] 12730.

[74453] Radisky, D. C.; Snyder, W. B.; Emr, S. C.; Kaplan, J.: Char-
acterization

[74454] of VPS41, a gene required for vacuolar trafficking and
high-affinity

[74455] iron transport in yeast. Proc. Nat. Acad. Sci. 94:
5662-5666, 1997.

[74456] 12731.

[74457] Arimura, T.; Nakamura, T.; Hiroi, S.; Satoh, M.; Takahashi, M.;

[74458] Ohbuchi, N.; Ueda, K.; Nouchi, T.; Yamaguchi, N.; Akai, J.; Matsumori,

[74459] A.; Sasayama, S.; Kimura, A.: Characterization of the human nebulette

[74460] gene: a polymorphism in an actin-binding motif is associated with

[74461] nonfamilial idiopathic dilated cardiomyopathy. Hum. Genet. 107:

[74462] 440-451, 2000.

[74463] 12732.

[74464] Millevoi, S.; Trombitas, K.; Kolmerer, B.; Kostin, S.; Schaper,

[74465] J.; Pelin, K.; Granzier, H.; Labeit, S.: Characterization of nebulette

[74466] and nebulin and emerging concepts of their roles for vertebrate Z-discs. J.

[74467] Molec. Biol. 282: 111-123, 1998.

[74468] 12733.

[74469] Popovici, C.; Mattei, M.-G.; Rattner, J. B.; Birnbaum, D.; Pebusque,

[74470] M.-J.: Assignment of the centrosomal protein 110 gene (Cep110) to

[74471] mouse chromosome bands 2B-C1 by in situ hybridization. Cytogenet.

[74472] Cell Genet. 89: 216-217, 2000.

[74473] 12734.

[74474] Shibanuma, M.; Mashimo, J.; Mita, A.; Kuroki, T.; Nose, K.: Cloning

[74475] from a mouse osteoblastic cell line of a set of transform-ing-growth-

[74476] factor-beta-1-regulated genes, one of which seems to encode a follistatin-related

[74477] polypeptide. Europ. J. Biochem. 217: 13-19, 1993.

[74478] 12735.

[74479] Tanaka, M.; Ozaki, S.; Osakada, F.; Mori, K.; Okubo, M.; Nakao,

[74480] K.: Cloning of follistatin-related protein as a novel au-toantigen

[74481] in systemic rheumatic diseases. Int. Immun. 10: 1305-1314, 1998.

[74482] 12736.

[74483] Zwijsen, A.; Blockx, H.; van Arnhem, W.; Willems, J.; Fransen,
[74484] L.; Devos, K.; Raymackers, J.; van de Voorde, A.; Slegers, H.: Characterization
[74485] of a rat C6 glioma-secreted follistatin-related protein (FRP): cloning
[74486] and sequence of the human homologue. Europ. J. Biochem. 225: 937-946,
[74487] 1994.

[74488] 12737.

[74489] Jaffrey, S. R.; Snowman, A. M.; Eliasson, M. J. L.; Cohen, N. A.;
[74490] Snyder, S. H.: CAPON: a protein associated with neuronal nitric oxide
[74491] synthase that regulates its interactions with PSD95. Neu- ron 115-124,
[74492] 1998.

[74493] 12738.

[74494] Thuresson, A.-C.; Astrom, J.; Astrom, A.; Gronvik, K.-O.; Virtanen,
[74495] A.: Multiple forms of poly(A) polymerases in human cells. Proc.

[74496] Nat. Acad. Sci. 91: 979-983, 1994.

[74497] 12739.

[74498] Yamauchi, T.; Sugimoto, J.; Hatakeyama, T.; Asakawa, S.; Shimizu,

[74499] N.; Isobe, M.: Assignment of the human poly(A) polymerase (PAP) gene

[74500] to chromosome 14q32.1-q32.2 and isolation of a polymorphic CA repeat

[74501] sequence. J. Hum. Genet. 44: 253-255, 1999.

[74502] 12740.

[74503] Fort, M. M.; Cheung, J.; Yen, D.; Li, J.; Zurawski, S. M.; Lo,

[74504] S.; Menon, S.; Clifford, T.; Hunte, B.; Lesley, R.; Muchamuel, T.;

[74505] Hurst, S. D.; Zurawski, G.; Leach, M. W.; Gorman, D. M.; Rennick,

[74506] D. M.: IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies

[74507] in vivo. Immunity 15: 985-995, 2001.

[74508] 12741.

[74509] Scott, A. F.: Personal Communication. Baltimore, Md. 2/16/2001.

[74510] 12742.

[74511] Boles, K. S.; Barten, R.; Kumaresan, P. R.; Trowsdale, J.; Mathew,

[74512] P. A.: Cloning of a new lectin-like receptor expressed on human NK

[74513] cells. Immunogenetics 50: 1-7, 1999.

[74514] 12743.

[74515] Mori, T.; Fukuda, Y.; Kuroda, H.; Matsumura, T.; Ota, S.; Sugimoto,

[74516] T.; Nakamura, Y.; Inazawa, J.: Cloning and characterization of a

[74517] novel Rab-family gene, Rab36, within the region at 22q11.2 that is

[74518] homozygously deleted in malignant rhabdoid tumors. Biochem. Biophys.

[74519] Res. Commun. 254: 594-600, 1999.

[74520] 12744.

[74521] Zhou, J.-Y.; Fogelgren, B.; Wang, Z.; Roe, B. A.; Biegel, J. A.

[74522] : Isolation of genes from the rhabdoid tumor deletion region in chromosome

[74523] band 22q11.2. Gene 241: 133-141, 2000.

[74524] 12745.

[74525] Sood, R.; Makalowska, I.; Carpten, J. D.; Robbins, C. M.; Stephan,

[74526] D. A.; Connors, T. D.; Morgenbesser, S. D.; Su, K.; Pinkett, H. W.;

[74527] Graham, C. L.; Quesenberry, M. I.; Baxevanis, A. D.; Klinger, K. W.;

[74528] Trent, J. M.; Bonner, T. I.: The human RGL (RalGDS-like) gene: cloning,

[74529] expression analysis and genomic organization. Biochim. Biophys. Acta 1491:

[74530] 285-288, 2000.

[74531] 12746.

[74532] Acquati, F.; Accarino, M.; Nucci, C.; Fumagalli, P.; Jovine, L.;

[74533] Ottolenghi, S.; Taramelli, R.: The gene encoding DRAP (BACE2), a

[74534] glycosylated transmembrane protein of the aspartic pro-tease family,

[74535] maps to the Down syndrome critical region. FEBS Lett. 468: 59-64,

[74536] 2000.

[74537] 12747.

[74538] Bennett, B. D.; Babu-Khan, S.; Loeloff, R.; Louis, J.-C.; Curran,

[74539] E.; Citron, M.; Vassar, R.: Expression analysis of BACE2 in brain

[74540] and peripheral tissues. J. Biol. Chem. 275: 20647-20651, 2000.

[74541] 12748.

[74542] Saunders, A. J.; Kim, T.-W.; Tanzi, R. E.: BACE maps to chromosome

[74543] 11 and a BACE homolog, BACE2, reside in the obligate Down syndrome

[74544] region of chromosome 21. Science 286: 1255A only, 1999.

[74545] 12749.

[74546] Solans, A.; Estivill, X.; de la Luna, S.: A new aspartyl pro-tease

[74547] on 21q22.3, BACE2, is highly similar to Alzheimer's amy-loid precursor

[74548] protein beta-secretase. Cytogenet. Cell Genet. 89: 177-184, 2000.

[74549] 12750.

[74550] Xin, H.; Stephans, J. C.; Duan, X.; Harrowe, G.; Kim, E.; Grieshammer,
[74551] U.; Kingsley, C.; Giese, K.: Identification of a novel aspartic-like
[74552] protease differentially expressed in human breast cancer cell lines. Biochim.
[74553] Biophys. Acta 1501: 125-137, 2000.

[74554] 12751.

[74555] An, G.; Ng, A. Y.; Meka, C. S. R.; Luo, G.; Bright, S. P.; Cazares,
[74556] L.; Wright, G. L., Jr.; Veltri, R. W.: Cloning and characterization
[74557] of UROC28, a novel gene overexpressed in prostate, breast, and bladder
[74558] cancers. Cancer Res. 60: 7014-7020, 2000.

[74559] 12752.

[74560] Kraemer, C.; Enklaar, T.; Zabel, B.; Schmidt, E. R.: Mapping and
[74561] structure of DMXL1, a human homologue of the DmX gene from Drosophila
[74562] melanogaster coding for a WD repeat protein. Genomics 64: 97-101,

[74563] 2000.

[74564] 12753.

[74565] Adamec, J.; Rusnak, F.; Owen, W. G.; Naylor, S.; Benson, L. M.;

[74566] Gacy, A. M.; Isaya, G.: Iron-dependent self-assembly of recombinant

[74567] yeast frataxin: implications for Friedreich ataxia. Am. J. Hum. Genet. 67:

[74568] 549-562, 2000.

[74569] 12754.

[74570] Campuzano, V.; Montermini, L.; Lutz, Y.; Cova, L.; Hinde-lang, C.;

[74571] Jiralerspong, S.; Trottier, Y.; Kish, S. J.; Faucheux, B.; Trouillas,

[74572] P.; Authier, F. J.; Durr, A.; Mandel, J.-L.; Vescovi, A.; Pan-dolfo,

[74573] M.; Koenig, M.: Frataxin is reduced in Friedreich ataxia patients

[74574] and is associated with mitochondrial membranes. Hum. Molec. Genet. 6:

[74575] 1771-1780, 1997.

[74576] 12755.

[74577] Cavadini, P.; O'Neill, H. A.; Benada, O.; Isaya, G.: Assembly

[74578] and iron-binding properties of human frataxin, the pro-
tein deficient

[74579] in Friedrich ataxia. Hum. Molec. Genet. 11: 217-227,
2002.

[74580] 12756.

[74581] Chamberlain, S.; Pook, M.; Carvajal, J.; Doudney, K.; Hiller-
mann,

[74582] R.: Frataxin fracas. (Letter) Nature Genet. 15: 337-338,
1997.

[74583] 12757.

[74584] Cossee, M.; Campuzano, V.; Koutnikova, H.; Fischbeck, K.;
Mandel,

[74585] J.-L.; Koenig, M.; Bidichandani, S. I.; Patel, P. I.; Molte, M.
D.;

[74586] Canizares, J.; De Frutos, R.; Pianese, L.; Cavalcanti, F.;
Monticelli,

[74587] A.; Cocozza, S.; Montermini, L.; Pandolfo, M.: Frataxin fra-
cas. (Letter) Nature

[74588] Genet. 15: 337-338, 1997.

[74589] 12758.

[74590] Bartles, J. R.; Wierda, A.; Zheng, L.: Identification and characterization

[74591] of espin, an actin-binding protein localized to the F-actin-rich junctional

[74592] plaques of Sertoli cell ectoplasmic specializations. J. Cell Sci. 109:

[74593] 1229-1239, 1996.

[74594] 12759.

[74595] Bartles, J. R.; Zheng, L.; Li, A.; Wierda, A.; Chen, B.: Small

[74596] espin: a third actin-bundling protein and potential forked protein

[74597] ortholog in brush border microvilli. J. Cell Biol. 143: 107-119,

[74598] 1998.

[74599] 12760.

[74600] Chen, B.; Li, A.; Wang, D.; Wang, M.; Zheng, L.; Bartles, J. R.

[74601] : Espin contains an additional actin-binding site in its N terminus

[74602] and is a major actin-bundling protein of the Sertoli cell-

spermatid
[74603] ectoplasmic specialization junctional plaque. Molec. Biol. Cell 10:
[74604] 4327-4339, 1999.

[74605] 12761.

[74606] Zheng, L.; Sekerkova, G.; Vranich, K.; Tilney, L. G.; Mug-naini,
[74607] E.; Bartles, J. R.: The deaf jerker mouse has a mutation in the gene
[74608] encoding the espin actin-bundling proteins of hair cell stereocilia
[74609] and lacks espins. Cell 102: 377-385, 2000.

[74610] 12762.

[74611] Eymard-Pierre, E.; Lesca, G.; Dollet, S.; Santorelli, F. M.; di
[74612] Capua, M.; Bertini, E.; Boespflug-Tanguy, O.: Infantile-on-set ascending
[74613] hereditary spastic paralysis is associated with mutations in the alsin
[74614] gene. Am. J. Hum. Genet. 71: 518-527, 2002.

[74615] 12763.

[74616] Hadano, S.: Personal Communication. Isehara, Japan

3/11/2002.

[74617] 12764.

[74618] Shaw, P. J.: Genetic inroads in familial ALS. Nature Genet. 29:

[74619] 103-104, 2001.

[74620] 12765.

[74621] Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Tanaka, H.; Poleshchuk,

[74622] V. V.; Markova, E. D.; Tsuji, S.: Refined genetic location of the

[74623] chromosome 2p-linked progressive muscular dystrophy gene. Genomics 42:

[74624] 345-348, 1997.

[74625] 12766.

[74626] Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Tanaka, H.; Vereshchagin,

[74627] N. V.; Markova, E. D.; Poleshchuk, V. V.; Lozhnikova, S. M.; Sukhorukov,

[74628] V. S.; Limborska, S. A.; Slominsky, P. A.; Bulayeva, K. B.; Tsuji,

[74629] S.: Clinical and molecular analysis of a large family with

three
[74630] distinct phenotypes of progressive muscular dystrophy. Brain 119:
[74631] 1895-1909, 1996.

[74632] 12767.

[74633] McNally, E. M.; Ly, C. T.; Rosenmann, H.; Rosenbaum, S. M.; Jiang,
[74634] W.; Anderson, L. V. B.; Soffer, D.; Argov, Z.: Splicing mutation
[74635] in dysferlin produces limb-girdle muscular dystrophy with inflammation. Am.
[74636] J. Med. Genet. 91: 305-312, 2000.

[74637] 12768.

[74638] Weiler, T.; Greenberg, C. R.; Nylen, E.; Halliday, W.; Morgan,
[74639] K.; Eggertson, D.; Wrogemann, K.: Limb-girdle muscular dystrophy
[74640] and Miyoshi myopathy in an aboriginal Canadian kindred map to LGMD2B
[74641] and segregate with the same haplotype. Am. J. Hum. Genet. 59: 872-878,
[74642] 1996.

[74643] 12769.

[74644] Azibi, K.: Personal Communication. Algiers, Algeria 8/1991.

[74645] 12770.

[74646] Azibi, K.; Bachner, L.; Beckmann, J. S.; Matsumura, K.; Hamouda,

[74647] E.; Chaouch, M.; Chaouch, A.; Ait-Ouarab, R.; Vignal, A.; Weissenbach,

[74648] J.; Vinet, M.-C.; Leturcq, F.; Collin, H.; Tome, F. M. S.; Reghis,

[74649] A.; Fardeau, M.; Campbell, K. P.; Kaplan, J.-C.: Severe childhood

[74650] autosomal recessive muscular dystrophy with the defi-ciency of the

[74651] 50 kDa dystrophin-associated glycoprotein maps to chro-mosome 13q12. Hum.

[74652] Molec. Genet. 2: 1423-1428, 1993.

[74653] 12771.

[74654] Azibi, K.; Chaouch, M.; Reghis, A.; Vinet, M.-C.; Vignal, A.; Becuwe,

[74655] N.; Beckman, J.; Seboun, E.; Nguyen, S.; Cometto, M.; Fardeau, M.;

[74656] Tome, R.; Leturq, F.; Chafey, P.; Bachner, L.; Kaplan, J.-C.: Linkage
[74657] analysis of 19 families with autosomal recessive (Duchenne-like) muscular
[74658] dystrophy from Algeria. (Abstract) Cytogenet. Cell Genet. 58: 1907,
[74659] 1991.

[74660] 12772.

[74661] Ben Hamida, M.; Fardeau, M.; Attia, N.: Severe childhood muscular
[74662] dystrophy affecting both sexes and frequent in Tunisia. Muscle Nerve 6:
[74663] 469-480, 1983.

[74664] 12773.

[74665] Ben Hamida, M.; Marrakchi, D.: Dystrophie musculaire progressive
[74666] de type Duchenne en Tunisie: a propos de 13 familles et 31 cas d'une
[74667] forme en apparence recessive autosomique. J. Genet. Hum. 28: 1-9,
[74668] 1980.

[74669] 12774.

[74670] Ben Jelloun-Dellagi, S.; Chaffey, P.; Hentati, F.; Ben Hamida,
[74671] C.; Tome, F.; Colin, H.; Dellagi, K.; Kaplan, J. C.; Fardeau, M.;
[74672] Ben Hamida, M.: Presence of normal dystrophin in Tunisian severe
[74673] childhood autosomal recessive muscular dystrophy. Neu-rology 40:
[74674] 1903, 1990.

[74675] 12775.

[74676] Ben Othmane, K.; Ben Hamida, M.; Pericak-Vance, M. A.; Ben Hamida,
[74677] C.; Blel, S.; Carter, S. C.; Bowcock, A. M.; Petruhkin, K.; Gilliam,
[74678] T. C.; Roses, A. D.; Hentati, F.; Vance, J. M.: Linkage of Tunisian
[74679] autosomal recessive Duchenne-like muscular dystrophy to the pericentromeric
[74680] region of chromosome 13q. Nature Genet. 2: 315-317, 1992.

[74681] 12776.

[74682] Ben Othmane, K.; Speer, M. C.; Stauffer, J.; Blel, S.; Middle-
ton,
[74683] L.; Ben Hamida, C.; Etribi, A.; Loeb, D.; Hentati, F.; Roses,
A. D.;
[74684] Ben Hamida, M.; Pericak-Vance, M. A.; Vance, J. M.: Evi-
dence for
[74685] linkage disequilibrium in chromosome 13-linked
Duchenne-like muscular
[74686] dystrophy (LGMD2C). (Letter) Am. J. Hum. Genet. 57:
732-734, 1995.
[74687] 12777.

[74688] Crosbie, R. H.; Lim, L. E.; Moore, S. A.; Hirano, M.; Hays, A.

[74689] P.; Maybaum, S. W.; Collin, H.; Dovico, S. A.; Stolle, C. A.;
Fardeau,
[74690] M.; Tome, F. M. S.; Campbell, K. P.: Molecular and genetic
characterization
[74691] of sarcospan: insights into sarcoglycan-sarcospan inter-
actions. Hum.
[74692] Molec. Genet. 9: 2019-2027, 2000.

[74693] 12778.

[74694] Dubowitz, V.: Progressive muscular dystrophy of the
Duchenne

[74695] type in females and its mode of inheritance. Brain 83: 432-439,
[74696] 1960.

[74697] 12779.

[74698] El Kerch, F.; Sefiani, A.; Azibi, K.; Boutaleb, N.; Yahyaoui,
[74699] M.; Bentahila, A.; Vinet, M.-C.; Leturcq, F.; Bachner, L.; Beckmann,
[74700] J.; Campbell, K. P.; Tome, F. M. S.; Fardeau, M.; Kaplan, J.-C.:
[74701] Linkage analysis of families with severe childhood autosomal recessive
[74702] muscular dystrophy in Morocco indicates genetic homogeneity of the
[74703] disease in North Africa. J. Med. Genet. 31: 342-343, 1994.

[74704] 12780.

[74705] Fanin, M.; Hoffman, E. P.; Angelini, C.; Pegoraro, E.: Private
[74706] beta- and gamma-sarcoglycan gene mutations: evidence of a founder
[74707] effect in northern Italy. Hum. Mutat. 16: 13-17, 2000.

[74708] 12781.

[74709] Fardeau, M.; Matsumura, K.; Tome, F. M. S.; Collin, H.; Leturcq,

[74710] F.; Kaplan, J.-C.; Campbell, K. P.: Deficiency of the 50 kDa dystrophin

[74711] associated glycoprotein (adhalin) in severe autosomal recessive muscular

[74712] dystrophies in children native from European countries. C. R. Acad.

[74713] Sci. Paris 316: 799-804, 1993.

[74714] 12782.

[74715] Francke, U.; Darras, B. T.; Hersh, J. H.; Berg, B. O.; Miller,

[74716] R. G.: Brother/sister pairs affected with early-onset, progressive

[74717] muscular dystrophy: molecular studies reveal etiologic heterogeneity. Am.

[74718] J. Hum. Genet. 45: 63-72, 1989.

[74719] 12783.

[74720] Goonewardena, P.; Gustavson, K.-H.; Gamstorp, I.; Lundstrom, N.-R.;

[74721] Pettersson, U.: A new type of muscular dystrophy in two brothers:

[74722] analysis by use of DNA probes suggests autosomal recessive inheritance. Clin.

[74723] Genet. 34: 299-305, 1988.

[74724] 12784.

[74725] Augustin, I.; Betz, A.; Herrmann, C.; Jo, T.; Brose, N.: Differential

[74726] expression of two novel Munc13 proteins in rat brain. Biochem. J. 337:

[74727] 363-371, 1999.

[74728] 12785.

[74729] Augustin, I.; Rosenmund, C.; Sudhof, T. C.; Brose, N.: Munc13-1

[74730] is essential for fusion competence of glutamatergic synaptic vesicles. Nature 400:

[74731] 457-461, 1999.

[74732] 12786.

[74733] Rhee, J.-S.; Betz, A.; Pyott, S.; Reim, K.; Varoqueaux, F.; Augustin,

[74734] I.; Hesse, D.; Sudhof, T. C.; Takahashi, M.; Rosenmund, C.; Brose,

[74735] N.: Beta phorbol ester- and diacylglycerol-induced aug-

mentation of
[74736] transmitter release is mediated by Munc13s and not by PKCs. Cell 108:
[74737] 121-133, 2002.

[74738] 12787.

[74739] Rosenmund, C.; Sigler, A.; Augustin, I.; Reim, K.; Brose, N.; Rhee,
[74740] J.-S.: Differential control of vesicle priming and short-term plasticity
[74741] by Munc13 isoforms. Neuron 33: 411-424, 2002.

[74742] 12788.

[74743] Song, Y.; Ailenberg, M.; Silverman, M.: Cloning of a novel gene
[74744] in the human kidney homologous to rat munc13s: its potential role
[74745] in diabetic nephropathy. Kidney Int. 53: 1689-1695, 1998.
[74746] 12789.

[74747] Parker, N. J.; Begley, C. G.; Smith, P. J.; Fox, R. M.: Molecular
[74748] cloning of a novel human gene (D11S4896E) at chromo-

somal region 11p15.5. Genomics 37:
[74749] 253-256, 1996.

[74750] 12790.

[74751] Sabbioni, S.; Veronese, A.; Trubia, M.; Taramelli, R.; Bar-
banti-Brodano,
[74752] G.; Croce, C. M.; Negrini, M.: Exon structure and promoter
identification
[74753] of STIM1 (alias GOK), a human gene causing growth arrest
of the human
[74754] tumor cell lines G401 and RD. Cytogenet. Cell. Genet. 86:
214-218,
[74755] 1999.

[74756] 12791.

[74757] Yoshikawa, T.; Padigaru, M.; Karkera, J. D.; Sharma, M.;
Berrettini,
[74758] W. H.; Esterling, L. E.; Detera-Wadleigh, S. D.: Genomic
structure
[74759] and novel variants of myo-inositol monophosphatase 2
(IMPA2). Molec.
[74760] Psychiat. 5: 165-171, 2000.

[74761] 12792.

[74762] Yoshikawa, T.; Turner, G.; Esterling, L. E.; Sanders, A. R.; Detera-Wadleigh,

[74763] S. D.: A novel human myo-inositol monophosphatase gene, IMP.18p,

[74764] maps to a susceptibility region for bipolar disorder. Molec. Psychiat. 2:

[74765] 393-397, 1997.

[74766] 12793.

[74767] Michaud, J.; Kudoh, J.; Berry, A.; Bonne-Tamir, B.; Lalioti, M.

[74768] D.; Rossier, C.; Shibuya, K.; Kawasaki, K.; Asakawa, S.; Minoshima,

[74769] S.; Shimizu, N.; Antonarakis, S. E.; Scott, H. S.: Isolation and

[74770] characterization of a human chromosome 21q22.3 gene (WDR4) and its

[74771] mouse homologue that code for a WD-repeat protein. Genomics 68:

[74772] 71-79, 2000.

[74773] 12794.

[74774] Boudin, H.; Doan, A.; Xia, J.; Shigemoto, R.; Huganir, R. L.; Worley,

[74775] P.; Craig, A. M.: Presynaptic clustering of mGluR7a requires the

[74776] PICK1 PDZ domain binding site. Neuron 28: 485-497, 2000.

[74777] 12795.

[74778] Dev, K. K.; Nishimune, A.; Henley, J. M.; Nakanishi, S.: The protein

[74779] kinase C-alpha binding protein PICK1 interacts with short but not

[74780] long form alternative splice variants of AMPA receptor subunits. Neuropharmacology 38:

[74781] 635-644, 1999.

[74782] 12796.

[74783] Staudinger, J.; Zhou, J.; Burgess, R.; Elledge, S. J.; Olson, E.

[74784] N.: PICK1: a perinuclear binding protein and substrate for protein

[74785] kinase C isolated by the yeast two-hybrid system. J. Cell Biol. 128:

[74786] 263-271, 1995.

[74787] 12797.

[74788] Takeya, R.; Takeshige, K.; Sumimoto, H.: Interaction of the

PDZ
[74789] domain of human PICK1 with class I ADP-ribosylation factors. Biochem.
[74790] Biophys. Res. Commun. 267: 149-155, 2000.

[74791] 12798.

[74792] Xia, J.; Zhang, X.; Staudinger, J.; Huganir, R. L.: Clustering
[74793] of AMPA receptors by the synaptic PDZ domain-containing protein PICK1. Neuron 22:
[74794] 179-187, 1999.

[74795] 12799.

[74796] Chadwick, L. H.; McCandless, S. E.; Silverman, G. L.; Schwartz,
[74797] S.; Westaway, D.; Nadeau, J. H.: Betaine-homocysteine methyltransferase-2:
[74798] cDNA cloning, gene sequence, physical mapping, and expression of the
[74799] human and mouse genes. Genomics 70: 66-73, 2000.

[74800] 12800.

[74801] Blencowe, B. J.; Bauren, G.; Eldridge, A. G.; Issner, R.; Nickerson,
[74802] J. A.; Rosonina, E.; Sharp, P. A.: The SRm160/300 splicing

coactivator
[74803] subunits. RNA 6: 111-120, 2000.

[74804] 12801.

[74805] Sawada, Y.; Miura, Y.; Umeki, K.; Tamaoki, T.; Fujinaga, K.; Ohtaki,
[74806] S.: Cloning and characterization of a novel RNA-binding protein SRL300
[74807] with RS domains. Biochim. Biophys. Acta 1492: 191-195, 2000.
[74808] 12802.

[74809] Hirose, K.; Morita, M.; Ema, M.; Mimura, J.; Hamada, H.; Fujii,
[74810] H.; Saijo, Y.; Gotoh, O.; Sogawa, K.; Fujii-Kuriyama, Y.: cDNA cloning
[74811] and tissue-specific expression of a novel basic helix-loop-helix/PAS
[74812] factor (Arnt2) with close sequence similarity to the aryl hydrocarbon
[74813] receptor nuclear translocator (Arnt). Molec. Cell. Biol. 16: 1706-1713,
[74814] 1996.

[74815] 12803.

[74816] Keith, B.; Adelman, D. M.; Simon, M. C.: Targeted mutation of
[74817] the murine arylhydrocarbon receptor nuclear translocator 2 (Arnt2)
[74818] gene reveals partial redundancy with Arnt. Proc. Nat. Acad. Sci. 98:
[74819] 6692-6697, 2001.

[74820] 12804.

[74821] Michaud, J. L.; DeRossi, C.; May, N. R.; Holdener, B. C.; Fan,
[74822] C.-M.: ARNT2 acts as the dimerization partner of SIM1 for the development
[74823] of the hypothalamus. Mech. Dev. 90: 253-261, 2000.

[74824] 12805.

[74825] Wende, H.; Volz, A.; Ziegler, A.: Extensive gene duplica-
tions
[74826] and a large inversion characterize the human leukocyte receptor cluster. Immunogenetics 51:
[74827] 703-713, 2000.

[74828] 12806.

[74829] Fukunaga-Johnson, N.; Lee, S. W.; Liebert, M.; Grossman,

H. B.

[74830] : Molecular analysis of a gene, BB1, overexpressed in bladder and

[74831] breast carcinoma. Anticancer Res. 16: 1085-1090, 1996.

[74832] 12807.

[74833] Saito, H.; Papaconstantinou, J.; Sato, H.; Goldstein, S.: Regulation

[74834] of a novel gene encoding a lysyl oxidase-related protein in cellular

[74835] adhesion and senescence. J. Biol. Chem. 272: 8157-8160, 1997.

[74836] 12808.

[74837] Hsu, S. Y.; Liang, S.-G.; Hsueh, A. J. W.: Characterization of

[74838] two LGR genes homologous to gonadotropin and thy-rotropin receptors

[74839] with extracellular leucine-rich repeats and a G protein-coupled, seven-transmembrane

[74840] region. Molec. Endocr. 12: 1830-1845, 1998.

[74841] 12809.

[74842] Loh, E. D.; Broussard, S. R.; Kolakowski, L. F.: Molecular

characterization

[74843] of a novel glycoprotein hormone G-protein-coupled receptor. Biochem.

[74844] Biophys. Res. Commun. 282: 757-764, 2001.

[74845] 12810.

[74846] Loh, E. D.; Broussard, S. R.; Liu, Q.; Copeland, N. G.; Gilbert,

[74847] D. J.; Jenkins, N. A.; Kolakowski, L. F., Jr. Chromosomal localization

[74848] of GPR48, a novel glycoprotein hormone receptor like GPCR, in human

[74849] and mouse with radiation hybrid and interspecific backcross mapping. Cytogenet.

[74850] Cell Genet. 89: 2-5, 2000.

[74851] 12811.

[74852] Giffon, T.; Lepourcelet, M.; Pichon, L.; Jezequel, P.; Bouric,

[74853] P.; Carn, G.; Pontarotti, P.; Le Gall, J.-Y.; David, V.: Cloning

[74854] of a human homologue of the mouse Tctex-5 gene within the MHC class

[74855] I region. Immunogenetics 44: 331-339, 1996.

[74856] 12812.

[74857] Lepourcelet, M.; Andrieux, N.; Giffon, T.; Pichon, L.; Hampe, A.;

[74858] Galibert, F.; Mosser, J.: Systematic sequencing of the human HLA-A/HLA-F

[74859] region: establishment of a cosmid contig and identification of a new

[74860] gene cluster within 37 kb of sequence. Genomics 37: 316-326, 1996.

[74861] 12813.

[74862] Zhang, J.; Zhang, L.; Zhao, S.; Lee, E. Y. C.: Identification

[74863] and characterization of the human HCG V gene product as a novel inhibitor

[74864] of protein phosphatase-1. Biochemistry 37: 16728-16734, 1998.

[74865] 12814.

[74866] Hayakawa, A.; Matsuda, Y.; Daibata, M.; Nakamura, H.; Sano, K.

[74867] : Genomic organization, tissue expression, and cellular localization

[74868] of AF3p21, a fusion partner of MLL in therapy-related leukemia. Genes

[74869] Chromosomes Cancer 30: 364-374, 2001.

[74870] 12815.

[74871] Sano, K.; Hayakawa, A.; Piao, J.-H.; Kosaka, Y.; Nakamura, H.:

[74872] Novel SH3 protein encoded by the AF3p21 gene is fused to the mixed

[74873] lineage leukemia protein in a therapy-related leukemia with t(3;11)(p21;q23). Blood 95:

[74874] 1066-1068, 2000.

[74875] 12816.

[74876] Coulie, P. G.; Brichard, V.; Van Pel, A.; Wolfel, T.; Schnei-der,

[74877] J.; Traversari, C.; Mattei, S.; De Plaen, E.; Lurquin, C.; Szikora,

[74878] J.-P.; Renauld, J.-C.; Boon, T.: A new gene coding for a differentiation

[74879] antigen recognized by autologous cytolytic T lymphocytes on HLA-A2

[74880] melanomas. J. Exp. Med. 180: 35-42, 1994.

[74881] 12817.

[74882] Kawakami, Y.; Eliyahu, S.; Delgado, C. H.; Robbins, P. F.; Rivoltini,

[74883] L.; Topalian, S. L.; Miki, T.; Rosenberg, S. A.: Cloning of the gene
[74884] coding for a shared human melanoma antigen recognized by autologous
[74885] T cells infiltrating into tumor. Proc. Nat. Acad. Sci. 91: 3515-3519,
[74886] 1994.

[74887] 12818.

[74888] Alagramam, K. N.; Murcia, C. L.; Kwon, H. Y.; Pawlowski, K. S.;
[74889] Wright, C. G.; Woychik, R. P.: The mouse Ames waltzer hearing-loss
[74890] mutant is caused by mutation of Pcdh15, a novel proto-cadherin gene. Nature
[74891] Genet. 27: 99-102, 2001.

[74892] 12819.

[74893] Alagramam, K. N.; Yuan, H.; Kuehn, M. H.; Murcia, C. L.; Wayne,
[74894] S.; Srisailpathy, C. R. S.; Lowry, R. B.; Knaus, R.; Van Laer, L.;
[74895] Bernier, F. P.; Schwartz, S.; Lee, C.; Morton, C. C.; Mullins, R.

[74896] F.; Ramesh, A.; Van Camp, G.; Hagemen, G. S.; Woychik, R. P.; Smith,
[74897] R. J. H.: Mutations in the novel protocadherin PCDH15 cause Usher
[74898] syndrome type 1F. Hum. Molec. Genet. 10: 1709-1718, 2001.
[74899] 12820.
[74900] Astuto, L. M.; Bork, J. M.; Weston, M. D.; Askew, J. W.; Fields,
[74901] R. R.; Orten, D. J.; Ohliger, S. J.; Riazuddin, S.; Morell, R. J.;
[74902] Khan, S.; Riazuddin, S.; Kremer, H.; and 15 others: CDH23 mutation
[74903] and phenotype heterogeneity: a profile of 107 diverse families with
[74904] Usher syndrome and nonsyndromic deafness. Am. J. Hum. Genet. 71:
[74905] 262-275, 2002.
[74906] 12821.
[74907] Di Palma, F.; Holme, R. H.; Bryda, E. C.; Belyantseva, I. A.; Pellegrino,
[74908] R.; Kachar, B.; Steel, K. P.; Noben-Trauth, K.: Mutations in Cdh23,

[74909] encoding a new type of cadherin, cause stereocilia disorganization

[74910] in waltzer, the mouse model for Usher syndrome type 1D. Nature Genet. 27:

[74911] 103-107, 2001.

[74912] 12822.

[74913] von Brederlow, B.; Bolz, H.; Janecke, A.; La O Cabrera, A.; Rudolph,

[74914] G.; Lorenz, B.; Schwinger, E.; Gal, A.: Identification and in vitro

[74915] expression of novel CDH23 mutations of patients with Usher syndrome

[74916] type 1D. Hum. Mutat. 19: 268-273, 2002.

[74917] 12823.

[74918] Cao, H.; Hegele, R. A.: Identification of single-nucleotide polymorphisms

[74919] in the human LPIN1 gene. J. Hum. Genet. 47: 370-372, 2002.

[74920] 12824.

[74921] Peterfy, M.; Phan, J.; Xu, P.; Reue, K.: Lipodystrophy in the

[74922] fld mouse results from mutation of a new gene encoding a

nuclear protein,
[74923] lipin. Nature Genet. 27: 121-124, 2001.

[74924] 12825.

[74925] Reue, K.; Xu, P.; Wang, X.-P.; Slavin, B. G.: Adipose tissue deficiency,
[74926] glucose intolerance, and increased atherosclerosis result from mutation
[74927] in the mouse fatty liver dystrophy (fld) gene. J. Lipid Res. 41:
[74928] 1067-1076, 2000.

[74929] 12826.

[74930] Shakhov, A. N.; Rubtsov, A. V.; Lyakhov, I. G.; Tumanov, A. V.;
[74931] Nedospasov, S. A.: SPLASH (PLA(2)IID), a novel member of phospholipase
[74932] A2 family, is associated with lymphotoxin-deficiency. Genes Immun. 1:
[74933] 191-199, 2000.

[74934] 12827.

[74935] Domanski, T. L.; Finta, C.; Halpert, J. R.; Zaphiropoulos, P. G.

[74936] : cDNA cloning and initial characterization of CYP3A43, a novel human

[74937] cytochrome P450. Molec. Pharm. 59: 386-392, 2001.

[74938] 12828.

[74939] Gellner, K.; Eiselt, R.; Hustert, E.; Arnold, H.; Koch, I.; Haberl,

[74940] M.; Deglmann, C. J.; Burk, O.; Buntefuss, D.; Escher, S.; Bishop,

[74941] C.; Koebe, H.-G.; Brinkmann, U.; Klenk, H.-P.; Kleine, K.; Meyer,

[74942] U. A.; Wojnowski, L.: Genomic organization of the human CYP3A locus:

[74943] identification of a new, inducible CYP3A gene. Pharmaco-genetics 11:

[74944] 111-121, 2001.

[74945] 12829.

[74946] Westlind, A.; Malmebo, S.; Johansson, I.; Otter, C.; Ander-sson,

[74947] T. B.; Ingelman-Sundberg, M.; Oscarson, M.: Cloning and tissue distribution

[74948] of a novel human cytochrome P450 of the CYP3A subfam-ily, CYP3A43. Biochem.

[74949] Biophys. Res. Commun. 281: 1349-1355, 2001.

[74950] 12830.

[74951] Furusawa, M.; Ohnishi, T.; Taira, T.; Iguchi-Ariga, S. M. M.; Ariga,

[74952] H.: AMY-1, a c-Myc-binding protein, is localized in the mitochondria

[74953] of sperm by association with S-AKAP84, an anchor protein of cAMP-dependent

[74954] protein kinase. J. Biol. Chem. 276: 36647-36651, 2001.

[74955] 12831.

[74956] Taira, T.; Maeda, J.; Onishi, T.; Kitaura, H.; Yoshida, S.; Kato,

[74957] H.; Ikeda, M.; Tamai, K.; Iguchi-Ariga, S. M. M.; Ariga, H.: AMY-1,

[74958] a novel C-MYC binding protein that stimulates transcrip-tion activity

[74959] of C-MYC. Genes Cells 3: 549-565, 1998.

[74960] 12832.

[74961] Bement, W. M.; Hasson, T.; Wirth, J. A.; Cheney, R. E.; Mooseker,

[74962] M. S.: Identification and overlapping expression of multi-

ple unconventional
[74963] myosin genes in vertebrate cell types. Proc. Nat. Acad. Sci. 91:
[74964] 6549-6553, 1994. Note: Erratum: Proc. Nat. Acad. Sci. 91: 11767 only,
[74965] 1994.

[74966] 12833.

[74967] Holt, J. R.; Gillespie, S. K. H.; Provance, D. W., Jr.; Shah, K.;

[74968] Shokat, K. M.; Corey, D. P.; Mercer, J. A.; Gillespie, P. G.: A chemical-genetic
[74969] strategy implicates myosin-1c in adaptation by hair cells. Cell 108:
[74970] 371-381, 2002.

[74971] 12834.

[74972] Kao, H. Y.; Downes, M.; Ordentlich, P.; Evans, R. M.: Isola-tion
[74973] of a novel histone deacetylase reveals that class I and class II deacetylases
[74974] promote SMRT-mediated repression. Genes Dev. 14: 55-66, 2000.
[74975] 12835.

[74976] Orimo, A.; Tominaga, N.; Yoshimura, K.; Yamauchi, Y.; Nomura, M.;

[74977] Sato, M.; Nogi, Y.; Suzuki, M.; Suzuki, H.; Ikeda, K.; Inoue, S.;

[74978] Muramatsu, M.: Molecular cloning of ring finger protein 21 (RNF21)/interferon-responsive

[74979] finger protein (ifp1), which possesses two RING-B box-coiled coil

[74980] domains in tandem. Genomics 69: 143-149, 2000.

[74981] 12836.

[74982] Gallagher, G.; Dickensheets, H.; Eskdale, J.; Izotova, L. S.; Mirochnitchenko,

[74983] O. V.; Peat, J. D.; Vazquez, N.; Pestka, S.; Donnelly, R. P.; Kotenko,

[74984] S. V.: Cloning, expression and initial characterisation of interleukin-19

[74985] (IL-19), a novel homologue of human interleukin-10 (IL-10). Genes

[74986] Immunity 1: 442-450, 2000.

[74987] 12837.

[74988] Savino, M.; d'Apolito, M.; Centra, M.; van Beerendonk, H. M.; Cleton-Jansen,

[74989] A.-M.; Whitmore, S. A.; Crawford, J.; Callen, D. F.; Zelante, L.;

[74990] Savoia, A.: Characterization of copine VII, a new member of the copine

[74991] family, and its exclusion as a candidate in sporadic breast cancers

[74992] with loss of heterozygosity at 16q24.3. Genomics 61: 219-226, 1999.

[74993] 12838.

[74994] Runnels, L. W.; Yue, L.; Clapham, D. E.: The TRPM7 chan- nel is

[74995] inactivated by PIP(2) hydrolysis. Nature Cell Biol. 4: 329-336,

[74996] 2002.

[74997] 12839.

[74998] Scott, A. F.: Personal Communication. Baltimore, Md. 3/8/2001.

[74999] 12840.

[75000] Cosman, D.; Mullberg, J.; Sutherland, C. L.; Chin, W.; Ar- mitage,

[75001] R.; Fanslow, W.; Kubin, M.; Chalupny, N. J.: ULBPs, novel MHC class

[75002] I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK

[75003] cytotoxicity through the NKG2D receptor. Immunity 14: 123-133, 2001.

[75004] 12841.

[75005] Le, L. Q.; Kabarowski, J. H. S.; Weng, Z.; Satterthwaite, A. B.;

[75006] Harvill, E. T.; Jensen, E. R.; Miller, J. F.; Witte, O. N.: Mice

[75007] lacking the orphan G protein-coupled receptor G2A develop a late-onset

[75008] autoimmune syndrome. Immunity 14: 561-571, 2001.

[75009] 12842.

[75010] Scott, A. F.: Personal Communication. Baltimore, Md. 7/20/2001.

[75011] 12843.

[75012] Koch, P.; Bohlmann, I.; Schafer, M.; Hansen-Hagge, T. E.; Kiyoi,

[75013] H.; Wilda, M.; Hameister, H.; Bartram, C. R.; Janssen, J. W. G.:

[75014] Identification of a novel putative Ran-binding protein and its close

[75015] homologue. Biochem. Biophys. Res. Commun. 278: 241-249, 2000.

[75016] 12844.

[75017] Kirikoshi, H.; Koike, J.; Sagara, N.; Saitoh, T.; Tokuhara, M.;

[75018] Tanaka, K.; Sekihara, H.; Hirai, M.; Katoh, M.: Molecular cloning

[75019] and genomic structure of human Frizzled-3 at chromosome 8p21. Biochem.

[75020] Biophys. Res. Commun. 271: 8-14, 2000.

[75021] 12845.

[75022] Sala, C. F.; Formenti, E.; Terstappen, G. C.; Caricasole, A.:

[75023] Identification, gene structure, and expression of human frizzled-3

[75024] (FZD3). Biochem. Biophys. Res. Commun. 273: 27-34, 2000.

[75025] 12846.

[75026] Eggenschwiler, J. T.; Espinoza, E.; Anderson, K. V.: Rab23 is

[75027] an essential negative regulator of the mouse Sonic hedgehog signalling

[75028] pathway. Nature 412: 194-198, 2001.

[75029] 12847.

[75030] Zhang, Q.-H.; Ye, M.; Wu, X.-Y.; Ren, S.-X.; Zhao, M.; Zhao, C.-J.;

[75031] Fu, G.; Shen, Y.; Fan, H.-Y.; Lu, G.; Zhong, M.; Xu, X.-R.; and 9

[75032] others: Cloning and functional analysis of cDNAs with open reading

[75033] frames for 300 previously undefined genes expressed in CD34+ hematopoietic

[75034] stem/progenitor cells. Genome Res. 10: 1546-1560, 2000.

[75035] 12848.

[75036] Saitoh, T.; Hirai, M.; Katoh, M.: Molecular cloning and characterization

[75037] of human Frizzled-8 gene on chromosome 10p11.2. Int. Oncol. 18:

[75038] 991-996, 2001.

[75039] 12849.

[75040] Koike, J.; Takagi, A.; Miwa, T.; Hirai, M.; Terada, M.; Katoh,

[75041] M.: Molecular cloning of Frizzled-10, a novel member of

the Frizzled
[75042] gene family. Biochem. Biophys. Res. Commun. 262:
39-43, 1999.
[75043] 12850.

[75044] Wittenberger, T.; Schaller, H. C.; Hellebrand, S.: An ex-
pressed
[75045] sequence tag (EST) data mining strategy succeeding in the
discovery
[75046] of new G-protein coupled receptors. J. Molec. Biol. 307:
799-813,
[75047] 2001.

[75048] 12851.

[75049] Communi, D.; Gonzalez, N. S.; Detheux, M.; Brezillon, S.;
Lannoy,
[75050] V.; Parmentier, M.; Boeynaems, J.-M.: Identification of a
novel human
[75051] ADP receptor coupled to G(i). J. Biol. Chem. 276:
41479-41485, 2001.
[75052] 12852.

[75053] Chen, H.; Ross, C. A.; Wang, N.; Huo, Y.; MacKinnon, D. F.;
Potash,
[75054] J. B.; Simpson, S. G.; McMahon, F. J.; DePaulo, J. R., Jr.;

McInnis,

[75055] M. G.: NEDD4L on human chromosome 18q21 has multi-
ple forms of transcripts

[75056] and is a homologue of the mouse Nedd4-2 gene. Europ. J.
Hum. Genet. 9:

[75057] 922-930, 2001.

[75058] 12853.

[75059] Erdeniz, N.; Rothstein, R.: Rsp5, a ubiquitin-protein ligase,

[75060] is involved in degradation of the single-stranded-DNA
binding protein

[75061] Rfa1 in Saccharomyces cerevisiae. Molec. Cell. Biol. 20:
224-232,

[75062] 2000.

[75063] 12854.

[75064] Riggins, G. J.; Thiagalingam, S.; Rozenblum, E.; Weinstein,
C.

[75065] L.; Kern, S. E.; Hamilton, S. R.; Willson, J. K. V.; Markowitz,
S.

[75066] D.; Kinzler, K. W.; Vogelstein, B.: Mad-related genes in the
human. Nature

[75067] Genet. 13: 347-349, 1996.

[75068] 12855.

[75069] Hanai, R.; Caron, P. R.; Wang, J. C.: Human TOP3: a single-copy

[75070] gene encoding DNA topoisomerase III. Proc. Nat. Acad. Sci. 93: 3653-3657,

[75071] 1996.

[75072] 12856.

[75073] Hu, P.; Beresten, S. F.; van Brabant, A. J.; Ye, T.-Z.; Pandolfi,

[75074] P.-P.; Johnson, F. B.; Guarente, L.; Ellis, N. A.: Evidence for BLM

[75075] and topoisomerase III-alpha interaction in genomic stability. Hum.

[75076] Molec. Genet. 10: 1287-1298, 2001.

[75077] 12857.

[75078] Li, W.; Wang, J. C.: Mammalian DNA topoisomerase III-alpha is

[75079] essential in early embryogenesis. Proc. Nat. Acad. Sci. 95: 1010-1013,

[75080] 1998.

[75081] 12858.

[75082] Harteneck, C.; Wedel, B.; Koesling, D.; Malkewitz, J.; Bohme, E.;

[75083] Schultz, G.: Molecular cloning and expression of a new alpha-subunit

[75084] of soluble guanylyl cyclase. Interchangeability of the alpha-subunits

[75085] of the enzyme. FEBS Lett. 292: 217-222, 1991.

[75086] 12859.

[75087] Yu, F.; Warburton, D.; Wellington, S.; Danziger, R. S.: Assignment

[75088] of GUCIA2, the gene coding for the alpha-2 subunit of soluble guanylyl

[75089] cyclase, to position 11q21-q22 on human chromosome 11. Genomics 33:

[75090] 334-336, 1996.

[75091] 12860.

[75092] Abe, K.; Yamamura, K.; Suzuki, M.: Molecular and embryological

[75093] characterization of a new transgene-induced null allele of mouse Brachyury

[75094] locus. Mammalian Genome 11: 238-240, 2000.

[75095] 12861.

[75096] Rozzo, C.; Fossarello, M.; Galleri, G.; Sole, G.; Serru, A.; Orzalesi,

[75097] N.; Serra, A.; Pirastu, M.: A common beta-ig-h3 gene mutation (delta-f540)

[75098] in a large cohort of Sardinian Reis Bucklers' corneal dystrophy patients.

[75099] (Abstract) Hum. Mutat. 12: 215-216, 1998.

[75100] 12862.

[75101] Skonier, J.; Bennett, K.; Rothwell, V.; Kosowski, S.; Plowman,

[75102] G.; Wallace, P.; Edelhoff, S.; Disteche, C.; Neubauer, M.; Marquardt,

[75103] H.; Rootgers, J.; Purchio, A. F.: Beta-ig-h3: a transforming growth

[75104] factor-beta-responsive gene encoding a secreted protein that inhibits

[75105] cell attachment in vitro and suppresses the growth of CHO cells in

[75106] nude mice. DNA Cell Biol. 13: 571-584, 1994.

[75107] 12863.

[75108] Skonier, J.; Neubauer, M.; Madisen, L.; Bennett, K.; Plowman,

[75109] G. D.; Purchio, A. F.: cDNA cloning and sequence analysis of beta

[75110] ig-h3, a novel gene induced in a human adenocarcinoma cell line after

[75111] treatment with transforming growth factor-beta. DNA Cell Biol. 11:

[75112] 511-522, 1992.

[75113] 12864.

[75114] Stock, E. L.; Feder, R. S.; O'Grady, R. B.; Sugar, J.; Roth, S.

[75115] I.: Lattice corneal dystrophy type IIIA: clinical and histopathologic

[75116] correlations. Arch. Ophthal. 109: 354-358, 1991.

[75117] 12865.

[75118] Arsenijevic, D.; Onuma, H.; Pecqueur, C.; Raimbault, S.; Manning,

[75119] B. S.; Miroux, B.; Couplan, E.; Alves-Guerra, M.-C.; Goubern, M.;

[75120] Surwit, R.; Bouillard, F.; Richard, D.; Collins, S.; Ricquier, D.

[75121] : Disruption of the uncoupling protein-2 gene in mice reveals a role
[75122] in immunity and reactive oxygen species production. Nature Genet. 26:
[75123] 435-439, 2000.

[75124] 12866.

[75125] Bouchard, C.; Perusse, L.; Chagnon, Y. C.; Warden, C.; Ricquier,
[75126] D.: Linkage between markers in the vicinity of the uncoupling protein
[75127] 2 gene and resting metabolic rate in humans. Hum. Molec. Genet. 6:
[75128] 1887-1889, 1997.

[75129] 12867.

[75130] Brauner, P.; Nibbelink, M.; Flachs, P.; Vitkova, I.; Kopecky, P.;
[75131] Mertelikova, I.; Janderova, L.; Penicaud, L.; Casteilla, L.; Plavka,
[75132] R.; Kopecky, J.: Fast decline of hematopoiesis and uncoupling protein
[75133] 2 content in human liver after birth: location of the protein in Kupffer

[75134] cells. Pediat. Res. 49: 440-447, 2001.

[75135] 12868.

[75136] Esterbauer, H.; Schneitler, C.; Oberkofler, H.; Ebenbichler, C.;

[75137] Paulweber, B.; Sandhofer, F.; Ladurner, G.; Hell, E.; Stros-berg, A.

[75138] D.; Patsch, J. R.; Krempler, F.; Patsch, W.: A common poly-morphism

[75139] in the promoter of UCP2 is associated with decreased risk of obesity

[75140] in middle-aged humans. Nature Genet. 28: 178-183, 2001.

[75141] 12869.

[75142] Fleury, C.; Neverova, M.; Collins, S.; Raimbault, S.; Champigny,

[75143] O.; Levi-Meyrueis, C.; Bouillaud, F.; Seldin, M. F.; Surwit, R. S.;

[75144] Ricquier, D.; Warden, C. H.: Uncoupling protein-2: a novel gene linked

[75145] to obesity and hyperinsulinemia. Nature Genet. 15: 269-272, 1997.

[75146] 12870.

[75147] Flier, J. S.; Lowell, B. B.: Obesity research springs a proton

[75148] leak. Nature Genet. 15: 223-224, 1997.

[75149] 12871.

[75150] Millet, L.; Vidal, H.; Andreelli, F.; Larrouy, D.; Riou, J.-P.;

[75151] Ricquier, D.; Laville, M.; Langin, D.: Increased uncoupling protein-2

[75152] and -3 mRNA expression during fasting in obese and lean humans. J.

[75153] Clin. Invest. 100: 2665-2670, 1997.

[75154] 12872.

[75155] Edwards, Y. H.; Putt, W.; Lekoape, K. M.; Stott, D.; Fox, M.; Hopkinson,

[75156] D. A.; Sowden, J.: The human homolog T of the mouse T (Brachyury)

[75157] gene: gene structure, cDNA sequence, and assignment to chromosome

[75158] 6q27. Genome Res. 6: 226-233, 1996.

[75159] 12873.

[75160] Morrison, K.; Papapetrou, C.; Attwood, J.; Hol, F.; Lynch, S. A.;

[75161] Sampath, A.; Hamel, B.; Burn, J.; Sowden, J.; Stott, D.; Ma-
riman,
[75162] E.; Edwards, Y. H.: Genetic mapping of the human homo-
logue (T) of
[75163] mouse T(Brachyury) and a search for allele association be-
tween human
[75164] T and spina bifida. Hum. Molec. Genet. 5: 669-674, 1996.

[75165] 12874.

[75166] Papapetrou, C.; Drummond, F.; Reardon, W.; Winter, R.;
Spitz, L.;
[75167] Edwards, Y. H.: A genetic study of the human T gene and
its exclusion
[75168] as a major candidate gene for sacral agenesis with
anorectal atresia. J.
[75169] Med. Genet. 36: 208-213, 1999.

[75170] 12875.

[75171] Shields, D. C.; Ramsbottom, D.; Donoghue, C.; Pinjon, E.;
Kirke,
[75172] P. N.; Molloy, A. M.; Edwards, Y. H.; Mills, J. L.; Mynett-
Johnson,
[75173] L.; Weir, D. G.; Scott, J. M.; Whitehead, A. S.: Association
between

[75174] historically high frequencies of neural tube defects and the human

[75175] T homologue of mouse T (Brachyury). Am. J. Med. Genet. 92: 206-211,

[75176] 2000.

[75177] 12876.

[75178] Speer, M. C.; Melvin, E. C.; Viles, K. D.; Bauer, K. A.; Ram-persaud,

[75179] E.; Drake, C.; George, T. M.; Enterline, D. S.; Mackey, J. F.; Worley,

[75180] G.; Gilbert, J. R.; Nye, J. S.; NTD Collaborative Group: T lo-cus

[75181] shows no evidence for linkage disequilibrium or mutation in American

[75182] Caucasian neural tube defect families. Am. J. Med. Genet. 110: 215-218,

[75183] 2002.

[75184] 12877.

[75185] Trembath, D.; Sherbondy, A. L.; Vandyke, D. C.; Shaw, G. M.; Todoroff,

[75186] K.; Lammer, E. J.; Finnell, R. H.; Marker, S.; Lerner, G.; Murray,

[75187] J. C.: Analysis of select folate pathway genes, PAX3, and human T

[75188] in a midwestern neural tube defect population. Teratology 59: 331-341,

[75189] 1999.

[75190] 12878.

[75191] Gunduz, M.; Ouchida, M.; Fukushima, K.; Hanafusa, H.; Etani, T.;

[75192] Nishioka, S.; Nishizaki, K.; Shimizu, K.: Genomic structure of the

[75193] human ING1 gene and tumor-specific mutations detected in head and

[75194] neck squamous cell carcinomas. Cancer Res. 60: 3143-3146, 2000.

[75195] 12879.

[75196] Kuefer, M. U.; Look, A. T.; Williams, D. C.; Valentine, V.; Naeve,

[75197] C. W.; Behm, F. G.; Mullersman, J. E.; Yoneda-Kato, N.; Montgomery,

[75198] K.; Kucherlapati, R.; Morris, S. W.: cDNA cloning, tissue distribution,

[75199] and chromosomal localization of myelodysplasia/myeloid

leukemia factor

[75200] 2 (MLF2). Genomics 35: 392-396, 1996.

[75201] 12880.

[75202] Hasegawa, H.; Kiyokawa, E.; Tanaka, S.; Nagashima, K.;
Gotoh, N.;

[75203] Shibuya, M.; Kurata, T.; Matsuda, M.: DOCK180, a major
CRK-binding

[75204] protein, alters cell morphology upon translocation to the
membrane. Molec.

[75205] Cell Biol. 16: 1770-176, 1996.

[75206] 12881.

[75207] Savill, J.: Phagocytic docking without shocking. Nature
442-443,

[75208] 1998.

[75209] 12882.

[75210] Takai, S.; Hasegawa, H.; Kiyokawa, E.; Yamada, K.; Kurata,
T.;

[75211] Matsuda, M.: Chromosomal mapping of the gene encod-
ing DOCK180, a

[75212] major Crk-binding protein, to 10q26.13-q26.3 by fluo-
rescence in situ

[75213] hybridization. Genomics 35: 403-404, 1996.

[75214] 12883.

[75215] Wu, Y.-C.; Horvitz, H. R.: C. elegans phagocytosis and cell-migration

[75216] protein CED-5 is similar to human DOCK180. Nature 392: 501-504,

[75217] 1998.

[75218] 12884.

[75219] Jamin, S. P.; Arango, N. A.; Mishina, Y.; Hanks, M. C.; Behringer,

[75220] R. R.: Requirement of Bmpr1a for Mullerian duct regression during

[75221] male sexual development. Nature Genet. 7Oct, 2002. Note: Advance

[75222] Electronic Publication.

[75223] 12885.

[75224] Zhou, X. P.; Woodford-Richens, K.; Lehtonen, R.; Kurose, K.; Aldred,

[75225] M.; Hampel, H.; Launonen, V.; Virta, S.; Pilarski, R.; Salovaara,

[75226] R.; Bodmer, W. F.; Conrad, B. A.; and 17 others: Germline

mutations
[75227] in BMPR1A/ALK3 cause a subset of cases of juvenile poly-
posis syndrome
[75228] and of Cowden and Bannayan-Riley-Ruvalcaba syn-
dromes. Am. J. Hum.
[75229] Genet. 69: 704-711, 2001.

[75230] 12886.

[75231] Alexander, J. M.; Bikkal, H. A.; Zervas, N. T.; Laws, E. R.,
Jr.;
[75232] Klibanski, A.: Tumor-specific expression and alternate
splicing of
[75233] messenger ribonucleic acid encoding activin/transforming
growth factor-beta
[75234] receptors in human pituitary adenomas. J. Clin. Endocr.
Metab. 81:
[75235] 783-790, 1996.

[75236] 12887.

[75237] Su, G. H.; Bansal, R.; Murphy, K. M.; Montgomery, E.; Yeo,
C. J.;
[75238] Hruban, R. H.; Kern, S. E.: ACVR1B (ALK4, activin receptor
type 1B)
[75239] gene mutations in pancreatic carcinoma. Proc. Nat. Acad.

Sci. 98:
[75240] 3254-3257, 2001.

[75241] 12888.

[75242] Xu, J.; Matsuzaki, K.; McKeehan, K.; Wang, F.; Kan, M.; Mc-
Keehan,
[75243] W. L.: Genomic structure and cloned cDNAs predict that
four variants
[75244] in the kinase domain of serine/threonine kinase receptors
arise by
[75245] alternative splicing and poly(A) addition. Proc. Nat. Acad.
Sci. 91:
[75246] 7957-7961, 1994.

[75247] 12889.

[75248] Zhou, Y.; Sun, H.; Danila, D. C.; Johnson, S. R.; Sigai, D. P.;

[75249] Zhang, X.; Klibanski, A.: Truncated activin type I receptor
Alk4
[75250] isoforms are dominant negative receptors inhibiting ac-
tivin signaling. Molec.
[75251] Endocr. 14: 2066-2075, 2000.

[75252] 12890.

[75253] Mann, S. S.; Pettenati, M. J.; von Kap-herr, C.; Hart, T. C.:

[75254] Reassignment of peptidyl prolyl isomerase-like 1 gene (PPIL1) to human

[75255] chromosome region 6p21.1 by radiation hybrid mapping and fluorescence

[75256] in situ hybridization. Cytogenet. Cell Genet. 83: 228-229, 1998.

[75257] 12891.

[75258] Ozaki, K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami, S.; Okuno,

[75259] S.; Takeda, S.; Shimada, Y.; Nagata, M.; Watanabe, T.; Takaichi, A.;

[75260] Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning, expression and chromosomal

[75261] mapping of a novel cyclophilin-related gene (PPIL1) from human fetal

[75262] brain. Cytogenet. Cell Genet. 72: 242-245, 1996.

[75263] 12892.

[75264] Graef, I. A.; Chen, F.; Chen, L.; Kuo, A.; Crabtree, G. R.: Signals

[75265] transduced by Ca(2+)/calcineurin and NFATc3/c4 pattern the developing

[75266] vasculature. Cell 105: 863-875, 2001.

[75267] 12893.

[75268] Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.:

[75269] Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB) and calcineurin

[75270] B (PPP3R1) are located on human chromosomes 4, 10q21-q22 and 2p16-p15,

[75271] respectively. Cytogenet. Cell Genet. 72: 236-241, 1996.

[75272] 12894.

[75273] Zeng, H.; Chattarji, S.; Barbarosie, M.; Rondi-Reig, L.; Philpot,

[75274] B. D.; Miyakawa, T.; Bear, M. F.; Tonegawa, S.: Forebrain-specific

[75275] calcineurin knockout selectively impairs bidirectional synaptic plasticity

[75276] and working/episodic-like memory. Cell 107: 617-629, 2001.

[75277] 12895.

[75278] Watanabe, T. K.; Shimizu, F.; Nagata, M.; Kawai, A.; Fuji-wara,

[75279] T.; Nakamura, Y.; Takahashi, E.; Hirai, Y.: Cloning, expres-sion,

[75280] and mapping of CKAP1, which encodes a putative cytoskeleton-associated
[75281] protein containing a CAP-GLY domain. Cytogenet. Cell Genet. 72:
[75282] 208-211, 1996.

[75283] 12896.

[75284] Kolodrubetz, D.; Burgum, A.: Sequence and genetic analysis of
[75285] NHP2: a moderately abundant high mobility group-like nuclear protein
[75286] with an essential function in Saccharomyces cerevisiae. Yeast 7:
[75287] 79-90, 1991.

[75288] 12897.

[75289] Saito, H.; Fujiwara, T.; Shin, S.; Okui, K.; Nakamura, Y.: Cloning
[75290] and mapping of a human novel cDNA (NHP2L1) that encodes a protein
[75291] highly homologous to yeast nuclear protein NHP2. Cytogenet. Cell
[75292] Genet. 72: 191-193, 1996.

[75293] 12898.

[75294] Mooseker, M. S.; Cheney, R. E.: Unconventional myosins. Annu.

[75295] Rev. Cell Dev. Biol. 11: 633-675, 1995.

[75296] 12899.

[75297] Bement, W. M.; Wirth, J. A.; Mooseker, M. S.: Cloning and mRNA

[75298] expression of human unconventional myosin-IC: a homologue of amoeboid

[75299] myosins-I with a single IQ motif and an SH3 domain. J. Molec. Biol. 243:

[75300] 356-363, 1994.

[75301] 12900.

[75302] Bement, W. M.; Hasson, T.; Wirth, J. A.; Cheney, R. E.; Mooseker,

[75303] M. S.: Identification and overlapping expression of multiple unconventional

[75304] myosin genes in vertebrate cell types. Proc. Nat. Acad. Sci. 91:

[75305] 6549-6553, 1994. Erratum: Proc. Nat. Acad. Sci. 91: 11767, 1994.

[75306] 12901.

[75307] Crozet, F.; Amraoui, A. E.; Blanchard, S.; Lenoir, M.; Ripoll,

[75308] C.; Vago, P.; Hamel, C.; Fizames, C.; Levi-Acobas, F.; De-
petris, D.;

[75309] Mattei, M.-G.; Weil, D.; Pujol, R.; Petit, C.: Cloning of the
genes

[75310] encoding two murine and human cochlear unconventional
type I myosins. Genomics 40:

[75311] 332-341, 1997.

[75312] 12902.

[75313] Dong, L. Q.; Du, H.; Porter, S. G.; Kolakowski, L. F., Jr.;
Lee,

[75314] A. V.; Mandarino, J.; Fan, J.; Yee, D.; Liu, F.: Cloning, chro-
mosome

[75315] localization, expression, and characterization of an Src
homology

[75316] 2 and pleckstrin homology domain-containing insulin re-
ceptor binding

[75317] protein hGrb10-gamma. J. Biol. Chem. 272:
29104-29112, 1997.

[75318] 12903.

[75319] Margolis, B.; Silvennoinen, O.; Comoglio, F.; Roonprapunt, C.;

[75320] Skolnik, E.; Ullrich, A.; Schlessinger, J.: High-efficiency ex-

[75321] pression/cloning of epidermal growth factor-receptor-binding proteins with Src homology

[75322] 2 domains. Proc. Nat. Acad. Sci. 89: 8894-8898, 1992.

[75323] 12904.

[75324] Skolnik, E. Y.; Margolis, B.; Mohammadi, M.; Lowenstein, E.; Fischer,

[75325] R.; Drepps, A.; Ullrich, A.; Schlessinger, J.: Cloning of PI3 kinase-

[75326] associated p85 utilizing a novel method for expression/ cloning of

[75327] target proteins for receptor tyrosine kinases. Cell 65: 83-90, 1991.

[75328] 12905.

[75329] Tanaka, S.; Mori, M.; Akiyoshi, T.; Tanaka, Y.; Mafune, K.; Wands,

[75330] J. R.; Sugimachi, K.: A novel variant of human Grb7 is as-

[75331] sociated with invasive esophageal carcinoma. J. Clin. Invest. 102:

821-827,

[75332] 1998.

[75333] 12906.

[75334] Blagitko, N.; Mergenthaler, S.; Schulz, U.; Wollmann, H. A.; Craigen,

[75335] W.; Eggermann, T.; Ropers, H.-H.; Kalscheuer, V. M.: Human GRB10

[75336] is imprinted and expressed from the paternal and maternal allele in

[75337] a highly tissue- and isoform-specific fashion. Hum. Molec. Genet. 9:

[75338] 1587-1595, 2000.

[75339] 12907.

[75340] Frantz, J. D.; Giorgetti-Peraldi, S.; Ottinger, E. A.; Shoelson,

[75341] S. E.: Human GRB-IR-beta/GRB10: splice variants of an insulin and

[75342] growth factor receptor-binding protein with PH and SH2 domains. J.

[75343] Biol. Chem. 272: 2659-2667, 1997.

[75344] 12908.

[75345] Hannula, K.; Lipsanen-Nyman, M.; Kontiokari, T.; Kere, J.: A narrow
[75346] segment of maternal uniparental disomy of chromosome 7q31-qter in
[75347] Silver-Russell syndrome delimits a candidate gene region. Am. J.
[75348] Hum. Genet. 68: 247-253, 2001.

[75349] 12909.

[75350] Jerome, C. A.; Scherer, S. W.; Tsui, L.-C.; Gietz, R. D.; Triggs-Raine,
[75351] B.: Assignment of growth factor receptor-bound protein 10 (GRB10)
[75352] to human chromosome 7p11.2-p12. Genomics 40: 215-216, 1997.
[75353] 12910.

[75354] Liu, F.; Roth, R. A.: Grb-IR: a SH2-domain-containing pro-tein
[75355] that binds to the insulin receptor and inhibits its function. Proc.
[75356] Nat. Acad. Sci. 92: 10287-10291, 1995.

[75357] 12911.

[75358] Ooi, J.; Yajnik, V.; Immanuel, D.; Gordon, M.; Moskow, J. J.;

[75359] Buchberg, A. M.; Margolis, B.: The cloning of Grb10 reveals a new

[75360] family of SH2 domain proteins. Oncogene 10: 1621-1630, 1995.

[75361] 12912.

[75362] Daly, R. J.; Sanderson, G. M.; Janes, P. W.; Sutherland, R. L.

[75363] : Cloning and characterization of GRB14, a novel member of the GRB7

[75364] gene family. J. Biol. Chem. 271: 12502-12510, 1996.

[75365] 12913.

[75366] Hakala, B. E.; White, C.; Recklies, A. D.: Human cartilage gp-39,

[75367] a major secretory product of articular chondrocytes and synovial cells,

[75368] is a mammalian member of a chitinase protein family. J. Biol. Chem. 268:

[75369] 25803-25810, 1993.

[75370] 12914.

[75371] Rehli, M.; Krause, S. W.; Andreesen, R.: Molecular characterization

[75372] of the gene for human cartilage gp-39 (CHI3L1), a member of the chitinase
[75373] protein family and marker for late stages of macrophage differentiation. Genomics 43:
[75374] 221-225, 1997.

[75375] 12915.

[75376] Dry, K.; Kenwrick, S.; Rosenthal, A.; Platzer, M.: The complete
[75377] sequence of the human locus for NgCAM-related cell adhesion molecule
[75378] reveals a novel alternative exon in chick and man and conserved genomic
[75379] organization for the L1 subfamily. Gene 273: 115-122, 2001.
[75380] 12916.

[75381] Grumet, M.; Mauro, V.; Burgoon, M. P.; Edelman, G. M.; Cunningham,
[75382] B. A.: Structure of a new nervous system glycoprotein, Nr-CAM, and
[75383] its relationship to subgroups of neural cell adhesion molecules. J.
[75384] Cell. Biol. 113: 1399-1412, 1991.

[75385] 12917.

[75386] Kayyem, J. F.; Roman, J. M.; de la Rosa, E. J.; Schwarz, U.; Dreyer,

[75387] W. J.: Bravo/Nr-CAM is closely related to the cell adhesion molecules

[75388] L1 and Ng-CAM and has a similar heterodimer structure. J. Cell. Biol. 118:

[75389] 1259-1270, 1992.

[75390] 12918.

[75391] Lane, R. P.; Chen, X.-N.; Yamakawa, K.; Vielmetter, J.; Ko-renberg,

[75392] J. R.; Dreyer, W. J.: Characterization of a highly conserved human

[75393] homolog to the chicken neural cell surface protein Bravo/ Nr-CAM that

[75394] maps to chromosome band 7q31. Genomics 35: 456-465, 1996.

[75395] 12919.

[75396] Wang, B.; Williams, H.; Du, J.-S.; Terrett, J.; Kenwrick, S.:

[75397] Alternative splicing of human NrCAM in neural and non-neural tissues. Molec.

[75398] Cell. Neurosci. 10: 287-295, 1998.

[75399] 12920.

[75400] Cates, C. A.; Michael, R. L.; Stayrook, K. R.; Harvey, K. A.; Burke,

[75401] Y. D.; Randall, S. K.; Crowell, P. L.; Crowell, D. N.: Prenylation

[75402] of oncogenic human PTP(CAAX) protein tyrosine phosphatases. Cancer

[75403] Lett. 110: 49-55, 1996.

[75404] 12921.

[75405] Montagna, M.; Serova, O.; Sylla, B. S.; Feunteun, J.; Lenoir, G.

[75406] M.: A 100-kb physical and transcriptional map around the EDH17B2

[75407] gene: identification of three novel genes and a pseudogene of a human

[75408] homologue of the rat PRL-1 tyrosine phosphatase. Hum. Genet. 96:

[75409] 532-538, 1995.

[75410] 12922.

[75411] Zeng, Q.; Hong, W.; Tan, Y. H.: Mouse PRL-2 and PRL-3,

two potentially

[75412] prenylated protein tyrosine phosphatases homologous to PRL-1. Biochem.

[75413] Biophys. Res. Commun. 244: 421-427, 1998.

[75414] 12923.

[75415] Muneer, S.; Ramalingam, V.; Wyatt, R.; Schultz, R. A.; Minna, J.

[75416] D.; Kamibayashi, C.: Genomic organization and mapping of the gene

[75417] encoding the PP2A B56-gamma regulatory subunit. Genomics 79: 344-348,

[75418] 2002.

[75419] 12924.

[75420] Deveraux, Q.; Jensen, C.; Rechsteiner, M.: Molecular cloning and

[75421] expression of a 26 S protease subunit enriched in dileucine repeats. J.

[75422] Biol. Chem. 270: 23726-23729, 1995.

[75423] 12925.

[75424] Deveraux, Q.; Ustrell, V.; Pickart, C.; Rechsteiner, M.: A 26

[75425] S protease subunit that binds ubiquitin conjugates. J. Biol. Chem. 269:

[75426] 7059-7061, 1994.

[75427] 12926.

[75428] Hu, R.-J.; Lee, M. P.; Johnson, L. A.; Feinberg, A. P.: A novel

[75429] human homologue of yeast nucleosome assembly protein, 65 kb centromeric

[75430] to the p57(KIP2) gene, is biallelically expressed in fetal and adult

[75431] tissues. Hum. Molec. Genet. 5: 1743-1748, 1996.

[75432] 12927.

[75433] Rodriguez, P.; Munroe, D.; Prawitt, D.; Chu, L. L.; Bric, E.; Kim,

[75434] J.; Reid, L. H.; Davies, C.; Nakagama, H.; Loebbert, R.; Winterpacht,

[75435] A.; Petruzzi, M.-J.; Higgins, M. J.; Nowak, N.; Evans, G.;

Shows,

[75436] T.; Weissman, B. E.; Zabel, B.; Housman, D. E.; Pelletier, J.: Functional

[75437] characterization of human nucleosome assembly protein-2 (NAP1L4) suggests

[75438] a role as a histone chaperone. Genomics 44: 253-265, 1997.

[75439] 12928.

[75440] Stegmaier, K.; Pendse, S.; Barker, G. F.; Bray-Ward, P.; Ward,

[75441] D. C.; Montgomery, K. T.; Krauter, K. S.; Reynolds, C.; Sklar, J.;

[75442] Donnelly, M.; Bohlander, S. K.; Rowley, J. D.; Sallan, S. E.; Gilliland,

[75443] D. G.; Golub, T. R.: Frequent loss of heterozygosity at the TEL gene

[75444] locus in acute lymphoblastic leukemia of childhood. Blood 86: 38-44,

[75445] 1995.

[75446] 12929.

[75447] Arakawa, H.; Nagase, H.; Hayashi, N.; Fujiwara, T.; Ogawa, M.;

[75448] Shin, S.; Nakamura, Y.: Molecular cloning and expression of a novel

[75449] human gene that is highly homologous to human FK506-binding protein

[75450] 12kDa (hFKBP-12) and characterization of two alternatively spliced

[75451] transcripts. Biochem. Biophys. Res. Commun. 200: 836-843, 1994.

[75452] 12930.

[75453] Xin, H.-B.; Senbonmatsu, T.; Cheng, D.-S.; Wang, Y.-X. Copello,

[75454] J. A.; Ji, G.-J.; Collier, M. L.; Deng, K.-Y.; Jeyakumar, L. H.; Magnuson,

[75455] M. A.; Inagami, T.; Kotlikoff, M. I.; Fleischer, S.: Oestrogen protects

[75456] FKBP12.6 null mice from cardiac hypertrophy. Nature 416: 334-337,

[75457] 2002.

[75458] 12931.

[75459] Dong, J.-T.; Isaacs, W. B.; Barrett, J. C.; Isaacs, J. T.: Genomic

[75460] organization of the human KAI1 metastasis-suppressor

gene. Genomics 41:
[75461] 25-32, 1997.

[75462] 12932.

[75463] Dong, J.-T.; Lamb, P. W.; Rinker-Schaeffer, C. W.;
Vukanovic, J.;
[75464] Ichikawa, T.; Isaacs, J. T.; Barrett, J. C.: KAI1, a metastasis
suppressor
[75465] gene for prostate cancer on human chromosome 11p11.2.
Science 268:
[75466] 884-886, 1995.

[75467] 12933.

[75468] Guo, X.-Z.; Friess, H.; Di Mola, F. F.; Heinicke, J.-M.;
Abou-Shady,
[75469] M.; Graber, H. U.; Baer, H. U.; Zimmermann, A.; Korc, M.;
Buchler,
[75470] M. W.: KAI1, a new metastasis suppressor gene, is reduced
in metastatic
[75471] hepatocellular carcinoma. Hepatology 28: 1481-1488,
1998.
[75472] 12934.

[75473] Mashimo, T.; Watabe, M.; Hirota, S.; Hosobe, S.; Miura, K.;

Tegtmeyer,

[75474] P. J.; Rinker-Shaeffer, C. W.; Watabe, K.: The expression of the

[75475] KAI1 gene, a tumor metastasis suppressor, is directly activated by

[75476] p53. Proc. Nat. Acad. Sci. 95: 11307-11311, 1998.

[75477] 12935.

[75478] Miyazaki, T.; Kato, H.; Shitara, Y.; Yoshikawa, M.; Tajima, K.;

[75479] Masuda, N.; Shouji, H.; Tsukada, K.; Nakajima, T.; Kuwano, H.: Mutation

[75480] and expression of the metastasis suppressor gene KAI1 in esophageal

[75481] squamous cell carcinoma. Cancer 89: 955-962, 2000.

[75482] 12936.

[75483] Brakebusch, C.; Jallal, B.; Fusco, O.; Iacobelli, S.; Ullrich,

[75484] A.: Expression of the 90K immunostimulator gene is controlled by

[75485] a promoter with unique features. J. Biol. Chem. 272: 3674-3682,

[75486] 1997.

[75487] 12937.

[75488] Calabrese, G.; Sures, I.; Pompetti, F.; Natoli, G.; Palka, G.;

[75489] Iacobelli, S.: The gene (LGALS3BP) encoding the serum protein 90K,

[75490] associated with cancer and infection by the human immunodeficiency

[75491] virus, maps at 17q25. Cytogenet. Cell Genet. 69: 223-225, 1995.

[75492] 12938.

[75493] Iacobelli, S.; Arno, E.; D'Orazio, A.; Coletti, G.: Detection

[75494] of antigens recognized by a novel monoclonal antibody in tissue and

[75495] serum from patients with breast cancer. Cancer Res. 46: 3005-3010,

[75496] 1986.

[75497] 12939.

[75498] Iacobelli, S.; Arno, E.; Sismondi, P.; Natoli, C.; Gentiloni, N.;

[75499] Scambia, G.; Giai, M.; Cortese, P.; Panici, P. B.; Mancuso, S.: Measurement

[75500] of a breast cancer associated antigen detected by mono-

clonal antibody

[75501] (SP-2) in sera of cancer patients. Breast Cancer Res. Treat. 11:

[75502] 19-30, 1988.

[75503] 12940.

[75504] Iacobelli, S.; Bucci, I.; D'Egidio, M.; Giuliani, C.; Natoli, C.;

[75505] Tinari, N.; Rubistein, M.; Schlessinger, J.: Purification and characterization

[75506] of a 90 kDa protein released from human tumors and tumor cell lines. FEBS

[75507] Lett. 319: 59-65, 1993.

[75508] 12941.

[75509] Gupta, P.; Soyombo, A. A.; Atashband, A.; Wisniewski, K. E.; Shelton,

[75510] J. M.; Richardson, J. A.; Hammer, R. E.; Hofmann, S. L.: Disruption

[75511] of PPT1 or PPT2 causes neuronal ceroid lipofuscinosis in knockout

[75512] mice. Proc. Nat. Acad. Sci. 98: 13566-13571, 2001.

[75513] 12942.

[75514] Mazoyer, S.; Gayther, S. A.; Nagai, M. A.; Smith, S. A.;

Dunning,

[75515] A.; van Rensburg, E. J.; Albertsen, H.; White, R.; Ponder, B. A. J.

[75516] : A gene (DLG2) located at 17q12-q21 encodes a new homologue of the

[75517] Drosophila tumor suppressor dlg-A. Genomics 28: 25-31, 1995.

[75518] 12943.

[75519] Korschen, H. G.; Beyermann, M.; Muller, F.; Heck, M.; Vantler,

[75520] M.; Koch, K.-W.; Kellner, R.; Wolfrum, U.; Bode, C.; Hofmann, K. P.;

[75521] Kaupp, U. B.: Interaction of glutamic-acid-rich proteins with the

[75522] cGMP signalling pathway in rod photoreceptors. Nature 400: 761-766,

[75523] 1999.

[75524] 12944.

[75525] Bianchi, L.: Glycogen storage disease I and hepatocellular tumours. Europ.

[75526] J. Pediat. 152 (suppl. 1): S63-S70, 1993.

[75527] 12945.

[75528] Amiel, J.; Espinosa-Parrilla, Y.; Steffann, J.; Gosset, P.; Pelet,
[75529] A.; Prieur, M.; Boute, O.; Choiset, A.; Lacombe, D.; Philip, N.; Le
[75530] Merrer, M.; Tanaka, H.; Till, M.; Touraine, R.; Toutain, A.; Vekemans,
[75531] M.; Munnich, A.; Lyonnet, S.: Large-scale deletions and SMADIP1 truncating
[75532] mutations in syndromic Hirschsprung disease with involvement of midline
[75533] structures. Am. J. Hum. Genet. 69: 1370-1377, 2001.

[75534] 12946.

[75535] Hack, A. A.; Cordier, L.; Shoturma, D. I.; Lam, M. Y.; Sweeney,
[75536] H. L.; McNally, E. M.: Muscle degeneration without mechanical injury
[75537] in sarcoglycan deficiency. Proc. Nat. Acad. Sci. 96: 10723-10728,
[75538] 1999.

[75539] 12947.

[75540] Mueller, O. T.; Henry, W. M.; Haley, L. L.; Byers, M. G.; Eddy,

[75541] R. L.; Shows, T. B.: Identification and chromosome location of genes

[75542] involved in glycoprotein neuraminidase deficiency disorders. (Abstract) Am.

[75543] J. Hum. Genet. 36: 205S only, 1984.

[75544] 12948.

[75545] Fu, G. K.; Lin, D.; Zhang, M. Y. H.; Bikle, D. D.; Shackleton,

[75546] C. H. L.; Miller, W. L.; Portale, A. A.: Cloning of human 25-hydroxyvitamin

[75547] D-1-alpha-hydroxylase and mutations causing vitamin D-dependent rickets

[75548] type 1. Molec. Endocr. 11: 1961-1970, 1997.

[75549] 12949.

[75550] Neufeld, E. F.: Personal Communication. Los Angeles, Calif.

[75551] 7/24/1987.

[75552] 12950.

[75553] Caron, H.; Peter, M.; van Sluis, P.; Speleman, F.; de Kraker, J.;

[75554] Laureys, G.; Michon, J.; Brugieres, L.; Voute, P. A.; Westerveld,

[75555] A.; Slater, R.; Delattre, O.; Versteeg, R.: Evidence for two tumour
[75556] suppressor loci on chromosomal bands 1p35-36 involved in neuroblastoma:
[75557] one probably imprinted, another associated with N-myc amplification. Hum.
[75558] Molec. Genet. 4: 535-539, 1995.

[75559] 12951.

[75560] Blair, H. J.; Ho, M.; Monaco, A. P.; Fisher, S.; Craig, I. W.;

[75561] Boyd, Y.: High-resolution comparative mapping of the proximal region
[75562] of the mouse X chromosome. Genomics 28: 305-310, 1995.
[75563] 12952.

[75564] Hayashi, Y. K.; Mizuno, Y.; Yoshida, M.; Nonaka, I.; Ozawa, E.;
[75565] Arahata, K.: The frequency of patients with 50-kd dystrophin-associated
[75566] glycoprotein (50DAG or adhalin) deficiency in a muscular dystrophy
[75567] patient population in Japan: immunocytochemical analysis of 50DAG,

[75568] 43DAG, dystrophin, and utrophin. Neurology 45: 551-554, 1995.

[75569] 12953.

[75570] Hazama, R.; Tsujihata, M.; Mori, M.; Mori, K.: Muscular dystrophy

[75571] in six young girls. Neurology 29: 1486-1491, 1979.

[75572] 12954.

[75573] Higuchi, I.; Yamada, H.; Fukunaga, H.; Iwaki, H.; Okubo, R.; Nakagawa,

[75574] M.; Osame, M.; Roberds, S. L.; Shimizu, T.; Campbell, K. P.; Matsumura,

[75575] K.: Abnormal expression of laminin suggests disturbance of sarcolemma-extracellular

[75576] matrix interaction in Japanese patients with autosomal recessive muscular

[75577] dystrophy deficient in adhalin. J. Clin. Invest. 94: 601-606, 1994.

[75578] 12955.

[75579] Jung, D.; Leturcq, F.; Sunada, Y.; Duclos, F.; Tome, F. M. S.;

[75580] Moomaw, C.; Merlini, L.; Azibi, K.; Chaouch, M.; Slaughter, C.; Fardeau,

[75581] M.; Kaplan, J.-C.; Campbell, K. P.: Absence of gamma-sarcoglycan
[75582] (35 DAG) in autosomal recessive muscular dystrophy linked to chromosome
[75583] 13q12. FEBS Lett. 381: 15-20, 1996.

[75584] 12956.

[75585] Kloepfer, H. W.; Talley, C.: Autosomal recessive inheri-tance
[75586] of Duchenne-type muscular dystrophy. Ann. Hum. Genet. 22: 138-143,
[75587] 1958.

[75588] 12957.

[75589] Lasa, A.; Piccolo, F.; de Diego, C.; Jeanpierre, M.; Colomer,

[75590] J.; Rodriguez, M. J.; Urtizberea, J. A.; Baiget, M.; Kaplan, J. C.;
[75591] Gallano, P.: Severe limb girdle muscular dystrophy in Spanish gypsies:
[75592] further evidence for a founder mutation in the gamma-sarcoglycan gene. Europ.
[75593] J. Hum. Genet. 6: 396-399, 1998.

[75594] 12958.

[75595] Leal, G. F.; Da-Silva, E. O.: Limb-girdle muscular dystro-
phy
[75596] with apparently different clinical courses within sexes in a
large
[75597] inbred kindred. J. Med. Genet. 36: 714-718, 1999.

[75598] 12959.

[75599] McNally, E. M.; Duggan, D.; Gorospe, J. R.; Bonnemann, C.
G.;
[75600] Fanin, M.; Pegoraro, E.; Lidov, H. G. W.; Noguchi, S.;
Ozawa, E.;
[75601] Finkel, R. S.; Cruse, R. P.; Angelini, C.; Kunkel, L. M.; Hoff-
man,
[75602] E. P.: Mutations that disrupt the carboxyl-terminus of
gamma-sarcoglycan
[75603] cause muscular dystrophy. Hum. Molec. Genet. 5:
1841-1847, 1996.
[75604] 12960.

[75605] Noguchi, S.; McNally, E. M.; Ben Othmane, K.; Hagiwara,
Y.; Mizuno,
[75606] Y.; Yoshida, M.; Yamamoto, H.; Bonnemann, C. G.; Gus-
soni, E.; Denton,
[75607] P. H.; Kyriakides, T.; Middleton, L.; Hentati, F.; Ben

Hamida, M.;
[75608] Nonaka, I.; Vance, J. M.; Kunkel, L. M.; Ozawa, E.: Muta-
tions in
[75609] the dystrophin-associated protein gamma-sarcoglycan in
chromosome
[75610] 13 muscular dystrophy. Science 270: 819-821, 1995.

[75611] 12961.

[75612] Passos-Bueno, M. R.; Bakker, E.; Marie, S. K.; Pavanello, R.
C.;
[75613] Vainzof, M.; Carvalho, A. A.; Cohen, D.; Beckmann, J.;
Zatz, M.:
[75614] Exclusion of the 15q locus as a candidate gene for severe
childhood
[75615] autosomal recessive Duchenne-like muscular dystrophy in
Brazilian
[75616] families. Hum. Molec. Genet. 2: 201-202, 1993.

[75617] 12962.

[75618] Passos-Bueno, M. R.; Oliveira, J. R.; Bakker, E.; Anderson,
R.
[75619] D.; Marie, S. K. N.; Vainzof, M.; Roberds, S.; Campbell, K.
P.; Zatz,
[75620] M.: Genetic heterogeneity for Duchenne-like muscular

dystrophy (DLMD)
[75621] based on linkage and 50 DAG analysis. Hum. Molec. Genet. 2: 1945-1947,
[75622] 1993.

[75623] 12963.

[75624] Piccolo, F.; Jeanpierre, M.; Leturcq, F.; Dode, C.; Azibi, K.;

[75625] Toutain, A.; Merlini, L.; Jarre, L.; Navarro, C.; Krish-
namoorthy,
[75626] R.; Tome, F. M. S.; Urtizberea, J. A.; Beckmann, J. S.; Campbell,
[75627] K. P.; Kaplan, J.-C.: A founder mutation in the gamma-sarcoglycan
[75628] gene of Gypsies possibly predating their migration out of India. Hum.
[75629] Molec. Genet. 5: 2019-2022, 1996.

[75630] 12964.

[75631] Piccolo, F.; Roberds, S. L.; Jeanpierre, M.; Leturcq, F.; Az-
ibi,
[75632] K.; Beldjord, C.; Carrie, A.; Recan, D.; Chaouch, M.; Reghis, A.;
[75633] El Kerch, F.; Sefiani, A.; Voit, T.; Merlini, L.; Collin, H.; Ey-

mard,

[75634] B.; Beckmann, J. S.; Romero, N. B.; Tome, F. M. S.; Fardeau, M.; Campbell,

[75635] K. P.; Kaplan, J.-C.: Primary adhalinopathy: a common cause of autosomal

[75636] recessive muscular dystrophy of variable severity. Nature Genet. 10:

[75637] 213-215, 1995.

[75638] 12965.

[75639] Skyring, A. P.; McKusick, V. A.: Clinical, genetic and elec-trocardiographic

[75640] studies in childhood muscular dystrophy. Am. J. Med. Sci. 242: 534-547,

[75641] 1961.

[75642] 12966.

[75643] Somer, H.; Voutilainen, A.; Knuutila, S.; Kaitila, I.; Rapola,

[75644] J.; Leinonen, H.: Duchenne-like muscular dystrophy in two sisters

[75645] with normal karyotypes: evidence for autosomal recessive inheritance. Clin.

[75646] Genet. 28: 151-156, 1985.

[75647] 12967.

[75648] Stec, I.; Kres, W.; Meng, G.; Muller, B.; Muller, C. R.; Grimm

[75649] : Estimate of severe autosomal recessive limb-girdle muscular dystrophy

[75650] (LGMD2C, LGMD2D) among sporadic muscular dystrophy males; a study

[75651] of 415 men. J. Clin. Genet. 32: 930-933, 1995.

[75652] 12968.

[75653] Todorova, A.; Ashikov, A.; Beltcheva, O.; Tournev, I.; Kremensky,

[75654] I.: C283Y mutation and other C-terminal nucleotide changes in the

[75655] gamma-sarcoglycan gene in the Bulgarian Gypsy population. Hum. Mutat. 14:

[75656] 40-44, 1999.

[75657] 12969.

[75658] Vainzof, M.; Pavanello, R. C. M.; Pavanello-Filho, I.; Rapaport,

[75659] D.; Passos-Bueno, M. R.; Zubrzycka-Gaarn, E. E.; Bulman, D. E.; Zatz,

[75660] M.: Screening of male patients with autosomal recessive

Duchenne
[75661] dystrophy through dystrophin and DNA studies. Am. J. Med. Genet. 39:
[75662] 38-41, 1991.

[75663] 12970.

[75664] Zatz, M.; Passos-Bueno, M. R.; Rapaport, D.: Estimate of the
[75665] proportion of Duchenne muscular dystrophy with autosomal recessive
[75666] inheritance. Am. J. Med. Genet. 32: 407-410, 1989.

[75667] 12971.

[75668] Aida, N.; Yagishita, A.; Takada, K.; Katsumata, Y.: Cerebellar
[75669] MR in Fukuyama congenital muscular dystrophy: polymicrogyria with
[75670] cystic lesions. Am. J. Neuroradiol. 15: 1755-1759, 1994.

[75671] 12972.

[75672] Beggs, A. H.; Neumann, P. E.; Arahata, K.; Arikawa, E.; Nonaka,
[75673] I.; Anderson, M. S.; Kunkel, L. M.: Possible influences on the expression

[75674] of X chromosome-linked dystrophin abnormalities by heterozygosity

[75675] for autosomal recessive Fukuyama congenital muscular dystrophy. Proc.

[75676] Nat. Acad. Sci. 89: 623-627, 1992.

[75677] 12973.

[75678] Colombo, R.; Bignamini, A. A.; Carobene, A.; Sasaki, J.; Tachikawa,

[75679] M.; Kobayashi, K.; Toda, T.: Age and origin of the FCMD 3-prime-untranslated-region

[75680] retrotransposal insertion mutation causing Fukuyama-type congenital

[75681] muscular dystrophy in the Japanese population. Hum. Genet. 107:

[75682] 559-567, 2000.

[75683] 12974.

[75684] Dambska, M.; Wisniewski, K.; Sher, J.; Solish, G.: Cerebro-oculo-muscular

[75685] syndrome: a variant of Fukuyama congenital cerebro-muscular dystrophy. Clin.

[75686] Neuropath. 1: 93-98, 1982.

[75687] 12975.

[75688] Fukuyama, Y.; Kawazura, M.; Haruna, H.: A peculiar form of congenital

[75689] progressive muscular dystrophy: report of fifteen cases. Paediat.

[75690] Univ. Tokyo 4: 5-8, 1960.

[75691] 12976.

[75692] Ban, Y.; Taniyama, M.; Ban, Y.: Vitamin D receptor gene polymorphism

[75693] is associated with Graves' disease in the Japanese popula- tion. J.

[75694] Clin. Endocr. Metab. 85: 4639-4643, 2000.

[75695] 12977.

[75696] Duprez, L.; Parma, J.; Van Sande, J.; Allgeier, A.; Leclere, J.;

[75697] Schvartz, C.; Delisle, M.-J.; Decoulx, M.; Orgiazzi, J.; Du- mont, J.;

[75698] Vassart, G.: Germline mutations in the thyrotropin recep- tor gene

[75699] cause non-autoimmune autosomal dominant hyperthy- roidism. Nature Genet. 7:

[75700] 396-401, 1994.

[75701] 12978.

[75702] Pearce, S. H. S.; Cheetham, T.; Imrie, H.; Vaidya, B.; Barnes,

[75703] N. D.; Bilous, R. W.; Carr, D.; Meeran, K.; Shaw, N. J.; Smith, C.

[75704] S.; Toft, A. D.; Williams, G.; Kendall-Taylor, P.: A common and recurrent

[75705] 13-bp deletion in the autoimmune regulator gene in British kindreds

[75706] with autoimmune polyendocrinopathy type 1. Am. J. Hum. Genet. 63:

[75707] 1675-1684, 1998.

[75708] 12979.

[75709] Hitzenberger, K.: Autotoxic cyanosis due to intraglobular methemoglobinemia. Wien.

[75710] Arch. Med. 23: 85-96, 1932.

[75711] 12980.

[75712] Baker, A. R.; McDonnell, D. P.; Hughes, M.; Crisp, T. M.; Mangelsdorf,

[75713] D. J.; Haussler, M. R.; Pike, J. W.; Shine, J.; O'Malley, B. W.:

[75714] Cloning and expression of full-length cDNA encoding hu-

man vitamin
[75715] D receptor. Proc. Nat. Acad. Sci. 85: 3294-3298, 1988.

[75716] 12981.

[75717] Faraco, J. H.; Morrison, N. A.; Baker, A.; Shine, J.; Frossard,

[75718] P. M.: ApaI dimorphism at the human vitamin D receptor gene locus. Nucleic
[75719] Acids Res. 17: 2150, 1989.

[75720] 12982.

[75721] Garnero, P.; Borel, O.; Sornay-Rendu, E.; Arlot, M. E.; Del-mas,
[75722] P. D.: Vitamin D receptor gene polymorphisms are not re-lated to bone
[75723] turnover, rate of bone loss, and bone mass in post-menopausal women:
[75724] the OFELY study. J. Bone Miner. Res. 11: 827-834, 1996.

[75725] 12983.

[75726] Houston, L. A.; Grant, S. F. A.; Reid, D. M.; Ralston, S. H.:

[75727] Vitamin D receptor polymorphism, bone mineral density, and osteoporotic
[75728] vertebral fracture: studies in a UK population. Bone 18:

249-252,

[75729] 1996.

[75730] 12984.

[75731] Hughes, M. R.; Malloy, P. J.; Kieback, D. G.; Kesterson, R. A.;

[75732] Pike, J. W.; Feldman, D.; O'Malley, B. W.: Point mutations in the

[75733] human vitamin D receptor gene associated with hypocalcemic rickets. Science 242:

[75734] 1702-1705, 1988.

[75735] 12985.

[75736] Hustmyer, F. G.; Peacock, M.; Hui, S.; Johnston, C. C.; Christian,

[75737] J.: Bone mineral density in relation to polymorphism at the vitamin

[75738] D receptor gene locus. J. Clin. Invest. 94: 2130-2134, 1994.

[75739] 12986.

[75740] Kelly, P. J.; Hopper, J. L.; Macaskill, G. T.; Pocock, N. A.;

[75741] Sambrook, P. N.; Eisman, J. A.: Genetic factors in bone turnover. J.

[75742] Clin. Endocr. Metab. 72: 808-813, 1991.

[75743] 12987.

[75744] Lim, S. K.; Park, Y. S.; Park, J. M.; Song, Y. D.; Lee, E. J.;

[75745] Kim, K. R.; Lee, H. C.; Huh, K. B.: Lack of association be-

tween vitamin

[75746] D receptor genotypes and osteoporosis in Koreans. J. Clin.

Endocr.

[75747] Metab. 80: 3677-3681, 1995.

[75748] 12988.

[75749] Malloy, P. J.; Eccleshall, T. R.; Gross, C.; Van Maldergem,

L.;

[75750] Bouillon, R.; Feldman, D.: Hereditary vitamin D resistant

rickets

[75751] caused by a novel mutation in the vitamin D receptor that

results

[75752] in decreased affinity for hormone and cellular hypore-

sponsiveness. J.

[75753] Clin. Invest. 99: 297-304, 1997.

[75754] 12989.

[75755] Malloy, P. J.; Hochberg, Z.; Tiosano, D.; Pike, J. W.;

Hughes,

[75756] M. R.; Feldman, D.: The molecular basis of hereditary 1,25-dihydroxyvitamin

[75757] D3 resistant rickets in seven related families. J. Clin. Invest. 86:

[75758] 2071-2079, 1990.

[75759] 12990.

[75760] Chuang, D. T.: Personal Communication. Dallas, Tex. 3/17/1995.

[75761] 12991.

[75762] Hopkinson, D. A.; Corney, G.; Cook, P. J. L.; Robson, E. B.; Harris,

[75763] H.: Genetically determined electrophoretic variants of human red

[75764] cell NADH diaphorase. Ann. Hum. Genet. 34: 1-10, 1970.

[75765] 12992.

[75766] Hors-Cayla, M. C.; Junien, C.; Heuertz, S.; Mattei, J. F.; Frezal,

[75767] J.: Regional assignment of arylsulfatase A, mitochondrial aconitase

[75768] and NADH-cytochrome b5 reductase by somatic cell hybridization. Hum.

[75769] Genet. 58: 140-143, 1981.

[75770] 12993.

[75771] Hsieh, H.-S.; Jaffe, E. R.: Electrophoretic and functional variants
[75772] of NADH-methemoglobin reductase in hereditary methemoglobinemia. J.
[75773] Clin. Invest. 50: 196-202, 1971.

[75774] 12994.

[75775] Jaffe, E. R.: The reduction of methemoglobin in erythrocytes
[75776] of a patient with congenital methemoglobinemia, subjects with erythrocyte
[75777] glucose-6-phosphate dehydrogenase deficiency, and normal individuals. Blood 21:
[75778] 561-572, 1963.

[75779] 12995.

[75780] Jaffe, E. R.; Hsieh, H. S.: DPNH-methemoglobin reductase deficiency
[75781] and hereditary methemoglobinemia. Seminars Hemat. 8: 417-437, 1971.

[75782] 12996.

[75783] Jenkins, M. M.; Prchal, J. T.: A high-frequency polymorphism
[75784] of NADH-cytochrome b5 reductase in African-Americans. Hum. Genet. 99:
[75785] 248-250, 1997.

[75786] 12997.

[75787] Johnson, C. J.; Bonrud, P. A.; Dosch, T. L.; Kilness, A. W.; Senger,
[75788] K. A.; Busch, D. C.; Meyer, M. R.: Fatal outcome of methemoglobinemia
[75789] in an infant. J.A.M.A. 257: 2796-2797, 1987.

[75790] 12998.

[75791] Junien, C.; Leroux, A.; Lostanlen, D.; Reghis, A.; Boue, J.; Nicolas,
[75792] H.; Boue, A.; Kaplan, J. C.: Prenatal diagnosis of congenital enzymopenic
[75793] methaemoglobinaemia with mental retardation due to generalized cytochrome
[75794] b5 reductase deficiency: first report of two cases. Prenatal Diag. 1:
[75795] 17-24, 1981.

[75796] 12999.

[75797] Junien, C.; Vibert, M.; Weil, D.; Van Cong, N.; Kaplan, J.-C.

[75798] : Assignment of NADH-cytochrome b5 reductase (DIA-1 locus) to human

[75799] chromosome 22. Hum. Genet. 42: 233-239, 1978.

[75800] 13000.

[75801] Kaftory, A.; Freundlich, E.; Manaster, J.; Shukri, A.; Hegesh,

[75802] E.: Prenatal diagnosis of congenital methemoglobinemia with mental

[75803] retardation. Israel J. Med. Sci. 22: 837-840, 1986.

[75804] 13001.

[75805] Kaplan, J. C.: Personal Communication. Paris, France 9/25/1981.

[75806] 13002.

[75807] Kaplan, J. C.; Beutler, E.: Electrophoresis of red cell NADH-

[75808] and NADPH-diaphorases in normal subjects and patients with congenital

[75809] methemoglobinemia. Biochem. Biophys. Res. Commun. 29: 605-610, 1967.

[75810] 13003.

[75811] Kaplan, J. C.; Leroux, A.; Beauvais, P.: Formes cliniques et

[75812] biologiques du deficit en cytochrome b5 reductase. Comp. Rend. Soc.

[75813] Biol. 173: 368-379, 1979.

[75814] 13004.

[75815] Katsube, T.; Sakamoto, N.; Kobayashi, Y.; Seki, R.; Hirano, M.;

[75816] Tanishima, K.; Tomoda, A.; Takazakura, E.; Yubisui, T.; Takeshita,

[75817] M.; Sakaki, Y.; Fukumaki, Y.: Exonic point mutations in NADH-cytochrome

[75818] B5 reductase genes of homozygotes for hereditary methe-moglobinemia,

[75819] types I and III: putative mechanisms of tissue-dependent enzyme deficiency. Am.

[75820] J. Hum. Genet. 48: 799-808, 1991.

[75821] 13005.

[75822] Kobayashi, Y.; Fukumaki, Y.; Yubisui, T.; Inoue, J.; Sakaki, Y.

[75823] : Serine-proline replacement at residue 127 of NADH-

cytochrome b5

[75824] reductase causes hereditary methemoglobinemia, generalized type. Blood 75:

[75825] 1408-1413, 1990.

[75826] 13006.

[75827] Lawson, D. L.; Miale, T. D.; Harvey, J. L.; Bucciarelli, R. L.;

[75828] Nelson, L. S.: Leukocyte diaphorase deficiency in congenital methemoglobinemia:

[75829] a valuable prognostic indicator. Biol. Neonate 32:

193-196, 1977.

[75830] 13007.

[75831] Leroux, A.; Junien, C.; Kaplan, J.-C.; Bamberger, J.: Generalised

[75832] deficiency of cytochrome b5 reductase in congenital methaemoglobinaemia

[75833] with mental retardation. Nature 258: 619-620, 1975.

[75834] 13008.

[75835] Leroux, A.; Torlinski, L.; Kaplan, J.-C.: Soluble and microsomal

[75836] forms of NADH-cytochrome b5 reductase from human placenta: similarity

[75837] with NADH-methemoglobin reductase from human erythrocytes. Biochim.

[75838] Biophys. Acta 481: 50-62, 1977.

[75839] 13009.

[75840] Lostanlen, D.; Lenoir, G.; Kaplan, J.-C.: NADH-cytochrome b5

[75841] reductase activity in lymphoid cell lines: expression of the defect

[75842] in Epstein-Barr virus transformed lymphoblastoid cell lines from patients

[75843] with recessive congenital methemoglobinemia. J. Clin. Invest. 68:

[75844] 279-285, 1981.

[75845] 13010.

[75846] Muller, J.; Murawski, K.; Szymanowska, Z.; Koziorowski, A.; Radwan,

[75847] L.: Hereditary deficiency of NADPH 2-methaemoglobin reductase. Acta

[75848] Med. Scand. 173: 243-247, 1963.

[75849] 13011.

[75850] Nagai, T.; Shirabe, K.; Yubisui, T.; Takeshita, M.: Analysis

[75851] of mutant NADH-cytochrome b5 reductase: apparent 'type III' methemoglobinemia

[75852] can be explained as type I with an unstable reductase. Blood 81:

[75853] 808-814, 1993.

[75854] 13012.

[75855] Owen, E. P.; Berens, J.; Marinaki, A. M.; Ipp, H.; Harley, E.

[75856] H.: Recessive congenital methaemoglobinaemia type II, a new mutation

[75857] which causes incorrect splicing in the NADH-cytochrome b-5 reductase

[75858] gene. J. Inherit. Metab. Dis. 20: 610 only, 1997.

[75859] 13013.

[75860] Ozsoylu, S.: Hereditary methemoglobinemic cyanosis due to diaphorase

[75861] deficiency in three successive generations. Acta Haemat. 37: 276-283,

[75862] 1967.

[75863] 13014.

[75864] Pietrini, G.; Aggujaro, D.; Carrera, P.; Malyszko, J.; Vitale,

[75865] A.; Borgese, N.: A single mRNA, transcribed from an alternative,
[75866] erythroid-specific, promoter, codes for two non-myristylated forms
[75867] of NADH-cytochrome b(5) reductase. J. Cell Biol. 117: 975-986, 1992.
[75868] 13015.

[75869] Reghis, A.; Benabadji, M.; Tchen, P.; Kaplan, J. C.: Quantitative
[75870] variations of red-cell cytochrome b5 reductase (NADH-methemoglobin-reductase)
[75871] in the Algerian population: evidence for defective alleles. Hum.
[75872] Genet. 59: 148-153, 1981.

[75873] 13016.

[75874] Reghis, A.; Troungos, C.; Lostanlen, D.; Krishnamoorthy, R.; Kaplan,
[75875] J. C.: Characterization of weak alleles at the DIA1 locus (Mustapha
[75876] 1, Mustapha 2, and Mustapha 3) in the Algerian population. Hum. Genet. 64:
[75877] 173-175, 1983.

[75878] 13017.

[75879] Rossi, E. C.; Bryan, G. T.; Schilling, R. F.; Clatanoff, D. V.

[75880] : Remission of chronic methemoglobinemia following neomycin therapy. Am.

[75881] J. Med. 40: 440-447, 1966.

[75882] 13018.

[75883] Schwartz, J. M.; Jaffe, E. R.: Hereditary methemoglobine-mia with

[75884] deficiency of NADH dehydrogenase.In: Stanbury, J. B.; Wyngaarden,

[75885] J. B.; Fredrickson, D. S.: The Metabolic Basis of Inherited Disease.

[75886] New York: McGraw-Hill (pub.) (4th ed.): 1978. Pp. 1452-1464.

[75887] 13019.

[75888] Fu, G. K.; Portale, A. A.; Miller, W. L.: Complete structure

[75889] of the human gene for the vitamin D 1-alpha-hydroxylase, P450cl-alpha. DNA

[75890] Cell Biol. 16: 1499-1507, 1997.

[75891] 13020.

[75892] Glorieux, F. H.: Personal Communication. Montreal, Quebec, Canada
[75893] 3/20/1996.

[75894] 13021.

[75895] Glorieux, F. H.; Arabian, A.; Delvin, E. E.: Pseudo-vitamin D
[75896] deficiency: absence of 25-hydroxyvitamin D 1-alpha-hydroxylase activity
[75897] in human placenta decidual cells. J. Clin. Endocr. Metab. 80: 2255-2258,
[75898] 1995.

[75899] 13022.

[75900] Hamilton, R.; Harrison, J.; Fraser, D.; Raddle, I.; Morecki, R.;
[75901] Paunier, L.: The small intestine in vitamin D dependent rickets. Pediatrics 45:
[75902] 364-373, 1970.

[75903] 13023.

[75904] Holick, M. F.; Uskokovic, M.; Henley, J. W.; MacLaughlin, J.;

[75905] Holick, S. A.; Potts, J. T., Jr.: The photoproduction of 1-alpha,

[75906] 25-dihydroxyvitamin D3 in skin: an approach to the therapy of vitamin-D-resistant

[75907] syndromes. New Eng. J. Med. 303: 350-354, 1980.

[75908] 13024.

[75909] Kitanaka, S.; Takeyama, K.; Murayama, A.; Sato, T.; Okumura, K.;

[75910] Nogami, M.; Hasegawa, Y.; Niimi, H.; Yanagisawa, J.; Tanaka, T.; Kato,

[75911] S.: Inactivating mutations in the 25-hydroxyvitamin D3-1-alpha-hydroxylase

[75912] gene in patients with pseudovitamin D-deficiency rickets. New Eng.

[75913] J. Med. 338: 653-661, 1998.

[75914] 13025.

[75915] Kong, X. F.; Zhu, X. H.; Pei, Y. L.; Jackson, D. M.; Holick, M.

[75916] F.: Molecular cloning, characterization, and promoter analysis of

[75917] the human 25-hydroxyvitamin D(3)-1-alpha-hydroxylase gene. Proc.

[75918] Nat. Acad. Sci. 96: 6988-6993, 1999.

[75919] 13026.

[75920] Kitanaka, S.; Murayama, A.; Sakaki, T.; Inouye, K.; Seino, Y.;

[75921] Fukumoto, S.; Shima, M.; Yukizane, S.; Takayanagi, M.; Niimi, H.;

[75922] Takeyama, K.-I.; Kato, S.: No enzyme activity of 25-hydroxyvitamin

[75923] D3 1-alpha-hydroxylase gene product in pseudovitamin D deficiency

[75924] rickets, including that with mild clinical manifestation. J. Clin.

[75925] Endocr. Metab. 84: 4111-4117, 1999.

[75926] 13027.

[75927] Labuda, M.; Fujiwara, T. M.; Ross, M. V.; Morgan, K.; Garcia-Heras,

[75928] J.; Ledbetter, D. H.; Hughes, M. R.; Glorieux, F. H.: Two hereditary

[75929] defects related to vitamin D metabolism map to the same region of

[75930] human chromosome 12q13-14. J. Bone Miner. Res. 7: 1447-1453, 1992.

[75931] 13028.

[75932] Labuda, M.; Morgan, K.; Glorieux, F. H.: Regional assign-

ment

[75933] of vitamin D dependent rickets type I to chromosome 12q14. (Abstract) Am.

[75934] J. Hum. Genet. 45 (suppl.): A147 only, 1989.

[75935] 13029.

[75936] Labuda, M.; Morgan, K.; Glorieux, F. H.: Autosomal recessive,

[75937] vitamin D dependency type I (VDD1) mapped to chromosome 12q by linkage

[75938] analysis. (Abstract) Cytogenet. Cell Genet. 51: 1027-1028, 1989.

[75939] 13030.

[75940] Labuda, M.; Morgan, K.; Glorieux, F. H.: Mapping autosomal recessive

[75941] vitamin D dependency type I to chromosome 12q14 by linkage analysis. Am.

[75942] J. Hum. Genet. 47: 28-36, 1990.

[75943] 13031.

[75944] Labuda, M.; Ross, M. V.; Fujiwara, T. M.; Morgan, K.; Ledbetter,

[75945] D.; Hughes, M. R.; Glorieux, F. H.: Two hereditary defects

related

[75946] to vitamin D metabolism map to the same region of human chromosome

[75947] 12q. (Abstract) Cytogenet. Cell Genet. 58: 1978 only, 1991.

[75948] 13032.

[75949] Liberman, U. A.; Eil, C.; Marx, S. J.: Resistance to 1,25-dihydroxyvitamin

[75950] D: association with heterogeneous defects in cultured skin fibroblasts. J.

[75951] Clin. Invest. 71: 192-200, 1983.

[75952] 13033.

[75953] Marx, S. J.: Vitamin D and other calciferols.In: Scriver, C.

[75954] R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: The Metabolic Basis of

[75955] Inherited Disease. New York: McGraw-Hill (pub.) (6th ed.) II:

[75956] 1989. Pp. 2029-2045.

[75957] 13034.

[75958] Monkawa, T.; Yoshida, T.; Wakino, S.; Shinki, T.; Anazawa, H.;

[75959] DeLuca, H. F.; Suda, T.; Hayashi, M.; Saruta, T.: Molecular cloning
[75960] of cDNA and genomic DNA for human 25-hydroxyvitamin D3 1 alpha-hydroxylase. Biochem.
[75961] Biophys. Res. Commun. 239: 527-533, 1997.

[75962] 13035.

[75963] Panda, D. K.; Miao, D.; Tremblay, M. L.; Sirois, J.; Farookhi,
[75964] R.; Hendy, G. N.; Goltzman, D.: Targeted ablation of the 25-hydroxyvitamin
[75965] D 1-alpha-hydroxylase enzyme: evidence for skeletal, re-productive,
[75966] and immune dysfunction. Proc. Nat. Acad. Sci. 98: 7498-7503, 2001.
[75967] 13036.

[75968] Prader, A.; Illig, R.; Heierli, E.: Eine besondere Form der primaeren
[75969] Vitamin-D-resistenten Rachitis mit Hypocalcaemie und autosomal-dominantem
[75970] Erbgang: die hereditaere Pseudo-Mangelrachitis. Helv. Paediat. Acta 16:
[75971] 452-468, 1961.

[75972] 13037.

[75973] Prader, A.; Kind, H. P.; DeLuca, H. F.: Pseudovitamin D de-
ficiency
[75974] (vitamin D dependency).In: Bickel, H.; Stern, J.: Inborn Er-
rors of
[75975] Calcium and Bone Metabolism. Baltimore: University Park
Press (pub.)
[75976] 1976. Pp. 115-123.

[75977] 13038.

[75978] Scriver, C. R.: Vitamin D dependency. (Editorial) Pediatrics
45:
[75979] 361-363, 1970.

[75980] 13039.

[75981] Scriver, C. R.; Reade, T. M.; DeLuca, H. F.; Hamstra, A. J.:

[75982] Serum 1, 25-dihydroxyvitamin D levels in normal subjects
and in patients
[75983] with hereditary rickets or bone disease. New Eng. J. Med.
299: 976-979,
[75984] 1978.

[75985] 13040.

[75986] Sinnett, D.; Deragon, J.-M.; Simard, L. R.; Labuda, D.: Alumorphs--human

[75987] DNA polymorphisms detected by polymerase chain reaction using Alu-specific

[75988] primers. Genomics 7: 331-334, 1990.

[75989] 13041.

[75990] St-Arnaud, R.; Messerlian, S.; Moir, J. M.; Omdahl, J. L.; Glorieux,

[75991] F. H.: The 25-hydroxyvitamin D1-alpha-hydroxylase gene maps to the

[75992] pseudovitamin D-deficiency rickets (PDDR) disease locus. J. Bone

[75993] Miner. Res. 12: 1552-1559, 1997.

[75994] 13042.

[75995] Takeyama, K.; Kitanaka, S.; Sato, T.; Kobori, M.; Yanagisawa,

[75996] J.; Kato, S.: 25-hydroxyvitamin D3 1-alpha-hydroxylase and vitamin

[75997] D synthesis. Science 277: 1827-1830, 1997.

[75998] 13043.

[75999] Wang, J. T.; Lin, C.-J.; Burridge, S. M.; Fu, G. K.; Labuda,

M.;

[76000] Portale, A. A.; Miller, W. L.: Genetics of vitamin D 1-alpha-hydroxylase

[76001] deficiency in 17 families. Am. J. Hum. Genet. 63: 1694-1702, 1998.

[76002] 13044.

[76003] Winkler, I.; Schreiner, F.; Harmeyer, J.: Absence of renal 25-hydroxycholecalciferol-1-hydroxylase

[76004] activity in a pig strain with vitamin D-dependent rickets. Calcif.

[76005] Tissue Int. 38: 87-94, 1986.

[76006] 13045.

[76007] Zehnder, D.; Bland, R.; Williams, M. C.; McNinch, R. W.; Howie,

[76008] A. J.; Stewart, P. M.; Hewison, M.: Extrarenal expression of 25-hydroxyvitamin

[76009] D3-1-alpha-hydroxylase. J. Clin. Endocr. Metab. 86: 888-894, 2001.

[76010] 13046.

[76011] Zietkiewicz, E.; Labuda, M.; Sinnett, D.; Glorieux, F. H.; Labuda,

[76012] D.: Linkage mapping by simultaneous screening of multi-

ple polymorphic
[76013] loci using Alu oligonucleotide-directed PCR. Proc. Nat. Acad. Sci. 89:
[76014] 8448-8451, 1992.

[76015] 13047.

[76016] Fukuyama, Y.; Osawa, M.; Suzuki, H.: Congenital progressive muscular
[76017] dystrophy of the Fukuyama type--clinical, genetic and pathological
[76018] considerations. Brain Dev. 3: 1-30, 1981.

[76019] 13048.

[76020] Cohen-Solal, L.; Bonaventure, J.; Maroteaux, P.: Dominant mutations
[76021] in familial lethal and severe osteogenesis imperfecta. Hum. Genet. 87:
[76022] 297-301, 1991.

[76023] 13049.

[76024] Pope, F. M.; Nicholls, A. C.; McPheat, J.; Talmud, P.; Owen, R.
[76025] : Collagen genes and proteins in osteogenesis imperfecta. J. Med.

[76026] Genet. 22: 466-478, 1985.

[76027] 13050.

[76028] Pruchno, C. J.; Cohn, D. H.; Wallis, G. A.; Willing, M. C.; Starman,
[76029] B. J.; Zhang, X.; Byers, P. H.: Osteogenesis imperfecta due to recurrent
[76030] point mutations at CpG dinucleotides in the COL1A1 gene of type I
[76031] collagen. Hum. Genet. 87: 33-40, 1991.

[76032] 13051.

[76033] Starman, B. J.; Eyre, D.; Charbonneau, H.; Harrylock, M.; Weis,
[76034] M. A.; Weiss, L.; Graham, J. M., Jr.; Byers, P. H.: Osteogen-
esis
[76035] imperfecta: the position of substitution for glycine by cys-
teine in
[76036] the triple helical domain of the pro-alpha-1(I) chains of type I collagen
[76037] determines the clinical phenotype. J. Clin. Invest. 84:
1206-1214,
[76038] 1989.

[76039] 13052.

[76040] Kobayashi, K.; Nakahori, Y.; Miyake, M.; Matsumura, K.; Kondo-Iida,

[76041] E.; Nomura, Y.; Segawa, M.; Yoshioka, M.; Saito, K.; Osawa, M.; Hamano,

[76042] K.; Sakakihara, Y.; Nonaka, I.; Nakagome, Y.; Kanazawa, I.; Nakamura,

[76043] Y.; Tokunaga, K.; Toda, T.: An ancient retrotransposal insertion

[76044] causes Fukuyama-type congenital muscular dystrophy. Nature 394:

[76045] 388-392, 1998.

[76046] 13053.

[76047] Kondo-Iida, E.; Kobayashi, K.; Watanabe, M.; Sasaki, J.; Kumagai,

[76048] T.; Koide, H.; Saito, K.; Osawa, M.; Nakamura, Y.; Toda, T.: Novel

[76049] mutations and genotype-phenotype relationships in 107 families with

[76050] Fukuyama-type congenital muscular dystrophy (FCMD). Hum. Molec. Genet. 8:

[76051] 2303-2309, 1999.

[76052] 13054.

[76053] Kondo-Iida, E.; Saito, K.; Tanaka, H.; Tsuji, S.; Ishihara, T.;

[76054] Osawa, M.; Fukuyama, Y.; Toda, T.: Molecular genetic evi-
dence of

[76055] clinical heterogeneity in Fukuyama-type congenital mus-
cular dystrophy. Hum.

[76056] Genet. 99: 427-432, 1997.

[76057] 13055.

[76058] Miura, K.; Shirasawa, H.: Congenital muscular dystrophy of
the

[76059] Fukuyama type (FCMD) with severe myocardial fibrosis: a
case report

[76060] with postmortem angiography. Acta Path. Jpn. 37:
1823-1835, 1987.

[76061] 13056.

[76062] Miyake, M.; Nakahori, Y.; Matsushita, I.; Kobayashi, K.;
Mizuno,

[76063] K.; Hirai, M.; Kanazawa, I.; Nakagome, Y.; Tokunaga, K.;
Toda, T.

[76064] : YAC and cosmid contigs encompassing the Fukuyama-
type congenital

[76065] muscular dystrophy (FCMD) candidate region on 9q31. Genomics 40:
[76066] 284-293, 1997.

[76067] 13057.

[76068] Murakami, T.; Konishi, Y.; Takamiya, M.; Tsukagoshi, H.: Congenital
[76069] muscular dystrophy associated with micropolygyria--report of 2 cases. Acta
[76070] Path. Jpn. 25: 599-612, 1975.

[76071] 13058.

[76072] Nonaka, I.; Sugita, H.; Takada, K.; Kumagai, K.: Muscle histochemistry
[76073] in congenital muscular dystrophy with central nervous system involvement. Muscle
[76074] Nerve 5: 102-106, 1982.

[76075] 13059.

[76076] Saito, K.; Osawa, M.; Wang, Z.-P.; Ikeya, K.; Fukuyama, Y.; Kondo-Iida,
[76077] E.; Toda, T.; Ohashi, H.; Kurosawa, K.; Wakai, S.; Kaneko, K.: Haplotype-phenotype
[76078] correlation in Fukuyama congenital muscular dystrophy.

Am. J. Med.

[76079] Genet. 92: 184-190, 2000.

[76080] 13060.

[76081] Sasaki, J.; Ishikawa, K.; Kobayashi, K.; Kondo-Iida, E.;
Fukayama,

[76082] M.; Mizusawa, H.; Takashima, S.; Sakakihara, Y.; Naka-
mura, Y.; Toda,

[76083] T.: Neuronal expression of the fukutin gene. Hum. Molec.
Genet. 9:

[76084] 3083-3090, 2000.

[76085] 13061.

[76086] Takada, K.; Nakamura, H.; Tanaka, J.: Cortical dysplasia in
congenital

[76087] muscular dystrophy with central nervous system involve-
ment (Fukuyama

[76088] type). J. Neuropath. Exp. Neurol. 43: 395-407, 1984.

[76089] 13062.

[76090] Toda, T.; Kanazawa, I.; Nakamura, Y.: Localization of a
gene

[76091] responsible for Fukuyama type congenital muscular dys-
trophy to chromosome

[76092] 9q31-33 by linkage analysis. (Abstract) Human Genome Mapping Workshop
[76093] 93 20, 1993.

[76094] 13063.

[76095] Toda, T.; Miyake, M.; Kobayashi, K.; Mizuno, K.; Saito, K.; Osawa,
[76096] M.; Nakamura, Y.; Kanazawa, I.; Nakagome, Y.; Yokunaga, K.; Nakahori,
[76097] Y.: Linkage-disequilibrium mapping narrows the Fukuyama-type congenital
[76098] muscular dystrophy (FCMD) candidate region to less than 100 kb. Am.
[76099] J. Hum. Genet. 59: 1313-1320, 1996.

[76100] 13064.

[76101] Valenzuela, D. M.; Stitt, T. N.; DiStefano, P. S.; Rojas, E.;

[76102] Mattsson, K.; Compton, D. L.; Nunez, L.; Park, J. S.; Stark, J. L.;
[76103] Gies, D. R.; Thomas, S.; Le Beau, M. M.; Fernald, A. A.; Copeland,
[76104] N. G.; Jenkins, N. A.; Burden, S. J.; Glass, D. J.; Yancopou-los, G.

[76105] D.: Receptor tyrosine kinase specific for the skeletal muscle lineage:

[76106] expression in embryonic muscle , at the neuromuscular junction, and

[76107] after injury. Neuron 15: 573-584, 1995.

[76108] 13065.

[76109] Yoshioka, M.; Kuroki, S.: Clinical spectrum and genetic studies

[76110] of Fukuyama congenital muscular dystrophy. Am. J. Med. Genet. 53:

[76111] 245-250, 1994.

[76112] 13066.

[76113] Ben-Yoseph, Y.; Momoi, T.; Baylerian, M. S.; Nadler, H. L.: K(m)

[76114] defect in neuraminidase of dysmorphic type sialidosis with and without

[76115] beta-galactosidase deficiency. Clin. Chim. Acta 123: 233-240, 1982.

[76116] 13067.

[76117] Bonten, E.; van der Spoel, A.; Fornerod, M.; Grosveld, G.; d'Azzo,

[76118] A.: Characterization of human lysosomal neuraminidase defines the
[76119] molecular basis of the metabolic storage disorder sialidosis. Genes
[76120] Dev. 10: 3156-3169, 1996.

[76121] 13068.

[76122] Bonten, E. J.; Arts, W. F.; Beck, M.; Covanis, A.; Donati, M. A.;
[76123] Parini, R.; Zammarchi., E.; d'Azzo, A.: Novel mutations in lysosomal
[76124] neuraminidase identify functional domains and determine clinical severity
[76125] in sialidosis. Hum. Molec. Genet. 9: 2715-2725, 2000.

[76126] 13069.

[76127] Cantz, M.; Gehler, J.; Spranger, J. W.: Mucolipidosis I: increased
[76128] sialic acid content and deficiency of an alpha-N-acetylneuraminidase
[76129] in cultured fibroblasts. Biochem. Biophys. Res. Commun. 74: 732-738,
[76130] 1977.

[76131] 13070.

[76132] Den Tandt, W. R.; Leroy, J. G.: Deficiency of neuraminidase in

[76133] the sialidoses and the mucolipidoses. Hum. Genet. 53: 383-388, 1980.

[76134] 13071.

[76135] Durand, P.; Gatti, R.; Cavalieri, S.; Borrone, C.; Tondeur, M.;

[76136] Michalski, J.-C.; Strecker, G.: Sialidosis (mucolipidosis I). Helv.

[76137] Paediat. Acta 32: 391-400, 1977.

[76138] 13072.

[76139] Federico, A.; Cecio, A.; Apponi Battini, G.; Michalski, J.-C.;

[76140] Strecker, G.; Guazzi, G. C.: Macular cherry-red spot and myoclonus

[76141] syndrome: juvenile form of sialidosis. J. Neurol. Sci. 48: 157-169,

[76142] 1980.

[76143] 13073.

[76144] Franceschetti, S.; Uziel, G.; Di Donato, S.; Caimi, L.; Avanzini,

[76145] G.: Cherry-red spot myoclonus syndrome and alpha-neuraminidase deficiency:
[76146] neurophysiological, pharmacological and biochemical study in an adult. J.
[76147] Neurol. Neurosurg. Psychiat. 43: 934-940, 1980.

[76148] 13074.

[76149] Gascon, G.; Wallenberg, B.; Daif, A. K.; Ozand, P.: Success-ful
[76150] treatment of cherry red spot-myoclonus syndrome with 5-hydroxytryptophan. Ann.
[76151] Neurol. 24: 453-455, 1988.

[76152] 13075.

[76153] Gillan, J. E.; Lowden, J. A.; Gaskin, K.; Cutz, E.: Congenital
[76154] ascites as a presenting sign of lysosomal storage disease. J. Pediat. 104:
[76155] 225-231, 1984.

[76156] 13076.

[76157] Gravel, R. A.; Lowden, J. A.; Callahan, J. W.; Wolfe, L. S.; Ng
[76158] Yin Kin, N. M. K.: Infantile sialidosis: a phenocopy of type 1 GM1

[76159] gangliosidosis distinguished by genetic complementation and urinary
[76160] oligosaccharides. Am. J. Hum. Genet. 31: 669-679, 1979.

[76161] 13077.

[76162] Hancock, L. W.; Thaler, M. M.; Horwitz, A. L.; Dawson, G.: Generalized
[76163] N-acetylneuraminic acid storage disease: quantitation and identification
[76164] of the monosaccharide accumulating in brain and other tissues. J.
[76165] Neurochem. 38: 803-809, 1982.

[76166] 13078.

[76167] Harada, F.; Nishimura, Y.; Suzuki, K.; Matsumoto, H.; Oohira,
[76168] T.; Matsuda, I.; Sasazuki, T.: The patient with combined deficiency
[76169] of neuraminidase and 21-hydroxylase. Hum. Genet. 75: 91-92, 1987.

[76170] 13079.

[76171] Harzer, K.; Cantz, M.; Sewell, A. C.; Dhareshwar, S. S.; Roggendorf,

[76172] W.; Heckl, R. W.; Schofer, O.; Thumler, R.; Peiffer, J.; Schlote,
[76173] W.: Normomorphic sialidosis in two female adults with severe neurologic
[76174] disease and without sialyl oligosaccharidiuria. Hum. Genet. 74: 209-214,
[76175] 1986.

[76176] 13080.

[76177] Itoh, K.; Naganawa, Y.; Matsuzawa, F.; Aikawa, S.; Doi, H.; Sasagasako,
[76178] N.; Yamada, T.; Kira, J.; Kobayashi, T.; Pshezhetsky, A. V.; Sakuraba,
[76179] H.: Novel missense mutations in the human lysosomal sialidase gene
[76180] in sialidosis patients and prediction of structural alterations of
[76181] mutant enzymes. J. Hum. Genet. 47: 29-37, 2002.

[76182] 13081.

[76183] Itoyama, Y.; Goto, I.; Kuroiwa, Y.; Takeichi, M.; Kawabuchi, M.;
[76184] Tanaka, Y.: Familial juvenile neuronal storage disease: new disease

[76185] or variant of juvenile lipidosis?. Arch. Neurol. 35: 792-800, 1978.

[76186] 13082.

[76187] Kelly, T. E.; Bartoshesky, L.; Harris, D. J.; McCauley, R. G.

[76188] K.; Feingold, M.; Schott, G.: Mucolipidosis I (acid neu-

raminidase

[76189] deficiency): three cases and delineation of the variability of the

[76190] phenotype. Am. J. Dis. Child. 135: 703-708, 1981.

[76191] 13083.

[76192] Kelly, T. E.; Graetz, G.: Isolated acid neuraminidase defi-

ciency:

[76193] a distinct lysosomal storage disease. Am. J. Med. Genet. 1: 31-46,

[76194] 1977.

[76195] 13084.

[76196] Klein, D.; Klein, J.: Polymorphism of the Apl (Neu-1) locus in

[76197] the mouse. Immunogenetics 16: 181-184, 1982.

[76198] 13085.

[76199] Klein, J.; Klein, D.; Figueroa, F.: Should the neu-
raminidase-1
[76200] locus be considered as part of the major histocompatibil-
ity complex?. Hum.
[76201] Immun. 15: 396-403, 1986.

[76202] 13086.

[76203] Lalley, P. A.; Shows, T. B.: Lysosomal acid phosphatase
deficiency:
[76204] liver specific variant in the mouse. Genetics 87: 305-317,
1977.
[76205] 13087.

[76206] Laver, J.; Fried, K.; Beer, S. I.; Iancu, T. C.; Heyman, E.;
Bach,
[76207] G.; Zeigler, M.: Infantile lethal neuraminidase deficiency
(sialidosis). Clin.
[76208] Genet. 23: 97-101, 1983.

[76209] 13088.

[76210] Loeb, H.; Teppel, M.; Cremer, N.: Clinical, biochemical and
ultrastructural
[76211] studies of an atypical form of mucopolysaccharidosis.
Acta Paediat. 58:

[76212] 220-228, 1969.

[76213] 13089.

[76214] Paschke, R.; Ludgate, M.: The thyrotropin receptor in thy-
roid
[76215] diseases. New Eng. J. Med. 337: 1675-1681, 1997.

[76216] 13090.

[76217] Thomas, J. L.; Leclere, J.; Hartemann, P.; Duheille, J.; Or-
giazzi,
[76218] J.; Petersen, M.; Janot, C.; Guedenet, J.-C.: Familial hyper-
thyroidism
[76219] without evidence of autoimmunity. Acta Endocr. 100:
512-518, 1982.

[76220] 13091.

[76221] Mitnick, M.; Reichlin, S.: Enzymatic synthesis of thy-
rotropin-releasing
[76222] hormone (TRH) by hypothalamic 'TRH synthetase.'. En-
docrinology 91:
[76223] 1145-1153, 1972.

[76224] 13092.

[76225] Cecchi, C.; Biasotto, M.; Tosi, M.; Avner, P.: The mottled
mouse

[76226] as a model for human Menkes disease: identification of mutations in
[76227] the Atp7a gene. Hum. Molec. Genet. 6: 425-433, 1997.

[76228] 13093.

[76229] Chelly, J.; Tumer, Z.; Tonnesen, T.; Petterson, A.; Ishikawa-Brush,
[76230] Y.; Tommerup, N.; Horn, N.; Monaco, A. P.: Isolation of a candidate
[76231] gene for Menkes disease that encodes a potential heavy metal binding
[76232] protein. Nature Genet. 3: 14-19, 1993.

[76233] 13094.

[76234] Dagenais, S. L.; Adam, A. N.; Innis, J. W.; Glover, T. W.: A novel
[76235] frameshift mutation in exon 23 of ATP7A (MNK) results in occipital
[76236] horn syndrome and not in Menkes disease. Am. J. Hum. Genet. 69:
[76237] 420-427, 2001.

[76238] 13095.

[76239] Das, S.; Levinson, B.; Vulpe, C.; Whitney, S.; Gitschier, J.;

Packman,
[76240] S.: Similar splicing mutations of the Menkes/mottled copper-transporting
[76241] ATPase gene in occipital horn syndrome and the blotchy mouse. Am.
[76242] J. Hum. Genet. 56: 570-576, 1995.

[76243] 13096.

[76244] Dierick, H. A.; Ambrosini, L.; Spencer, J.; Glover, T. W.; Mercer,
[76245] J. F. B.: Molecular structure of the Menkes disease gene (ATP7A). Genomics 28:
[76246] 462-469, 1995.

[76247] 13097.

[76248] Francis, M. J.; Jones, E. E.; Levy, E. R.; Ponnambalam, S.; Chelly,
[76249] J.; Monaco, A. P.: A Golgi localization signal identified in the
[76250] Menkes recombinant protein. Hum. Molec. Genet. 7: 1245-1252, 1998.
[76251] 13098.

[76252] Grimes, A.; Hearn, C. J.; Lockhart, P.; Newgreen, D. F.;

Mercer,

[76253] J. F. B.: Molecular basis of the brindled mouse mutant (Mo-br): a

[76254] murine model of Menkes disease. Hum. Molec. Genet. 6: 1037-1042,

[76255] 1997.

[76256] 13099.

[76257] Gu, Y.-H.; Kodama, H.; Murata, Y.; Mochizuki, D.; Yana-

gawa, Y.;

[76258] Ushijima, H.; Shiba, T.; Lee, C.-C.: ATP7A gene mutations in 16 patients

[76259] with Menkes disease and a patient with occipital horn syndrome. Am.

[76260] J. Med. Genet. 99: 217-222, 2001.

[76261] 13100.

[76262] Horn, N.: Personal Communication. Copenhagen, Den-

mark 2/25/1999.

[76263] 13101.

[76264] Jankov, R. P.; Boerkoel, C. F.; Hellmann, J.; Sirkin, W. L.; Tumer,

[76265] Z.; Horn, N.; Feigenbaum, A.: Lethal neonatal Menkes' dis-

ease with
[76266] severe vasculopathy and fractures. Acta Paediat. 87:
1297-1300,
[76267] 1998.

[76268] 13102.

[76269] Kaler, S. G.; Gallo, L. K.; Proud, V. K.; Percy, A. K.; Mark,

[76270] Y.; Segal, N. A.; Goldstein, D. S.; Holmes, C. S.; Gahl, W.
A.: Occipital
[76271] horn syndrome and a mild Menkes phenotype associated
with splice site
[76272] mutations at the MNK locus. Nature Genet. 8: 195-202,
1994.
[76273] 13103.

[76274] Ketterling, R. P.; Drost, J. B.; Scaringe, W. A.; Liao, D.; Liu,

[76275] J.; Kasper, C. K.; Sommer, S. S.: Reported in vivo splice-
site mutations
[76276] in the factor IX gene: severity of splicing defects and a hy-
pothesis
[76277] for predicting deleterious splice donor mutations. Hum.
Mutat. 13:
[76278] 221-231, 1999.

[76279] 13104.

[76280] Levinson, B.; Conant, R.; Schnur, R.; Das, S.; Packman, S.; Gitschier,

[76281] J.: A repeated element in the regulatory region of the MNK gene and

[76282] its deletion in a patient with occipital horn syndrome. Hum. Molec.

[76283] Genet. 5: 1737-1742, 1996.

[76284] 13105.

[76285] Mentzel, H. J.; Seidel, J.; Vogt, S.; Vogt, L.; Kaiser, W. A.

[76286] : Vascular complications (splenic and hepatic artery aneurysms) in

[76287] the occipital horn syndrome: report of a patient and re-view of the

[76288] literature. Pediat. Radiol. 29: 19-22, 1999.

[76289] 13106.

[76290] Mercer, J. F. B.; Livingston, J.; Hall, B.; Paynter, J. A.; Begy,

[76291] C.; Chandrasekharappa, S.; Lockhart, P.; Grimes, A.; Bhave, M.; Siemieniak,

[76292] D.; Glover, T. W.: Isolation of a partial candidate gene for Menkes

[76293] disease by positional cloning. Nature Genet. 3: 20-25, 1993.

[76294] 13107.

[76295] Moller, L. B.; Tumer, Z.; Lund, C.; Petersen, C.; Cole, T.; Hanusch,

[76296] R.; Seidel, J.; Jensen, L. R.; Horn, N.: Similar splice-site mutations

[76297] of the ATP7A gene lead to different phenotypes: classical Menkes disease

[76298] or occipital horn syndrome. Am. J. Hum. Genet. 66: 1211-1220, 2000.

[76299] 13108.

[76300] Packman, S.; Enns, G. M.; O'Toole, C. J.; Cox, V. A.; Golabi,

[76301] M.: Atypical severe Menkes disease presenting with neonatal cutis

[76302] laxa. (Abstract) Am. J. Hum. Genet. 61 (suppl.): A258 only, 1997.

[76303] 13109.

[76304] Petris, M. J.; Camakaris, J.; Greenough, M.; LaFontaine, S.; Mercer,

[76305] J. F. B.: A C-terminal di-leucine is required for localization of

[76306] the Menkes protein in the trans-Golgi network. Hum. Molec. Genet. 7:
[76307] 2063-2071, 1998.

[76308] 13110.

[76309] Petris, M. J.; Strausak, D.; Mercer, J. F. B.: The Menkes copper
[76310] transporter is required for the activation of tyrosinase. Hum. Molec.
[76311] Genet. 9: 2845-2851, 2000.

[76312] 13111.

[76313] Qian, Y.; Tiffany-Castiglioni, E.; Harris, E. D.: Sequence of

[76314] a Menkes-type Cu-transporting ATPase from rat C6 glioma cells: comparison
[76315] of the rat protein with other mammalian Cu-transporting ATPases. Molec.
[76316] Cell. Biochem. 181: 49-61, 1998.

[76317] 13112.

[76318] Reed, V.; Boyd, Y.: Mutation analysis provides additional proof
[76319] that mottled is the mouse homologue of Menkes' disease. Hum. Molec.

[76320] Genet. 6: 417-423, 1997.

[76321] 13113.

[76322] Katoh, M.: Molecular cloning and characterization of MFRP, a novel

[76323] gene encoding a membrane-type Frizzled-related protein. Biochem.

[76324] Biophys. Res. Commun. 282: 116-123, 2001.

[76325] 13114.

[76326] Koesters, R.; Adams, V.; Betts, D.; Moos, R.; Schmid, M.; Siermann,

[76327] A.; Hassam, S.; Weitz, S.; Lichter, P.; Heitz, P. U.; von Knebel Doeberitz,

[76328] M.; Briner, J.: Human eukaryotic initiation factor EIF2C1 gene: cDNA

[76329] sequence, genomic organization, localization to chromo-somal bands

[76330] 1q34-p35, and expression. Genomics 61: 210-218, 1999.

[76331] 13115.

[76332] Martinez, J.; Patkaniowska, A.; Urlaub, H.; Luhrmann, R.; Tuschi,

[76333] T.: Single-stranded antisense siRNAs guide target RNA

cleavage in

[76334] RNAi. Cell 110: 563-574, 2002.

[76335] 13116.

[76336] Suetsugu, S.; Miki, H.; Takenawa, T.: Identification of two human

[76337] WAVE/SCAR homologues as general actin regulatory molecules which associate

[76338] with the Arp2/3 complex. Biochem. Biophys. Res. Com-

mun. 260: 296-302,

[76339] 1999.

[76340] 13117.

[76341] Sakai, T.; Toyoda, A.; Hashimoto, K.; Maeda, H.: Isolation and

[76342] characterization of a novel zinc finger gene, ZNF219, and mapping

[76343] to the human chromosome 14q11 region. DNA Res. 7: 137-141, 2000.

[76344] 13118.

[76345] Hanaoka, E.; Ozaki, T.; Ohira, M.; Nakamura, Y.; Suzuki, M.; Takahashi,

[76346] E.; Moriya, H.; Nakagawara, A.; Sakiyama, S.: Molecular

cloning and
[76347] expression analysis of the human DA41 gene and its mapping to chromosome
[76348] 9q21.2-q21.3. J. Hum. Genet. 45: 188-191, 2000.

[76349] 13119.

[76350] Ozaki, T.; Hishiki, T.; Toyama, Y.; Yuasa, S.; Nakagawara, A.;
[76351] Sakiyama, S.: Identification of a new cellular protein that can interact
[76352] specifically with DAN. DNA Cell Biol. 16: 985-991, 1997.

[76353] 13120.

[76354] Dammann, R.; Li, C.; Yoon, J.-H.; Chin, P. L.; Bates, S.; Pfeifer,
[76355] G. P.: Epigenetic inactivation of a RAS association domain family
[76356] protein from the lung tumour suppressor locus 3p21.3. Nature Genet. 25:
[76357] 315-319, 2000.

[76358] 13121.

[76359] Dreijerink, K.; Braga, E.; Kuzmin, I.; Geil, L.; Duh, F.-M.; Angeloni,

[76360] D.; Zbar, B.; Lerman, M. I.; Stanbridge, E. J.; Minna, J. D.; Protopopov,

[76361] A.; Li, J.; Kashuba, V.; Klein, G.; Zabarovsky, E. R.: The candidate

[76362] tumor suppressor gene, RASSF1A, from human chromosome 3p21.3 is involved

[76363] in kidney tumorigenesis. Proc. Nat. Acad. Sci. 98: 7504-7509, 2001.

[76364] 13122.

[76365] Harada, K.; Toyooka, S.; Maitra, A.; Maruyama, R.; Toyooka, K.

[76366] O.; Timmons, C. F.; Tomlinson, G. E.; Mastrangelo, D.; Hay, R. J.;

[76367] Minna, J. D.; Gazdar, A. F.: Aberrant promoter methylation and silencing

[76368] of the RASSF1A gene in pediatric tumors and cell lines. Oncogene 21:

[76369] 4345-4349, 2002.

[76370] 13123.

[76371] Sekido, Y.; Ahmadian, M.; Wistuba, I. I.; Latif, F.; Bader, S.;

[76372] Wei, M.-H.; Duh, F.-M.; Gazdar, A. F.; Lerman, M. I.; Minna, J. D.

[76373] : Cloning of a breast cancer homozygous deletion junction narrows
[76374] the region of search for a 3p21.3 tumor suppressor gene. Oncogene 16:
[76375] 3151-3157, 1998.

[76376] 13124.

[76377] Dann, C. E.; Hsieh, J.-C.; Rattner, A.; Sharma, D.; Nathans, J.;
[76378] Leahy, D. J.: Insights into Wnt binding and signalling from the structures
[76379] of two Frizzled cysteine-rich domains. Nature 412: 86-90, 2001.
[76380] 13125.

[76381] Schumann, H.; Holtz, J.; Zerkowski, H.-R.; Hatzfeld, M.: Expression
[76382] of secreted frizzled related proteins 3 and 4 in human ventricular
[76383] myocardium correlates with apoptosis related gene ex- pression. Cardiovasc.
[76384] Res. 45: 720-728, 2000.

[76385] 13126.

[76386] Bouchon, A.; Dietrich, J.; Colonna, M.: Cutting edge: inflammatory
[76387] responses can be triggered by TREM-1, a novel receptor expressed on
[76388] neutrophils and monocytes. J. Immun. 164: 4991-4995, 2000.
[76389] 13127.

[76390] Bouchon, A.; Hernandez-Munain, C.; Cella, M.; Colonna, M.: A DAP12-mediated
[76391] pathway regulates expression of CC chemokine receptor 7 and maturation
[76392] of human dendritic cells. J. Exp. Med. 194: 1111-1122, 2001.
[76393] 13128.

[76394] Campbell, K. S.; Colonna, M.: DAP12: a key accessory protein for
[76395] relaying signals by natural killer cell receptors. Int. J. Biochem.
[76396] Cell Biol. 31: 631-636, 1999.

[76397] 13129.

[76398] Lanier, L. L.; Bakker, A. B. H.: The ITAM-bearing transmembrane

[76399] adaptor DAP12 in lymphoid and myeloid cell function. Immun. Today 21:

[76400] 611-614, 2000.

[76401] 13130.

[76402] Benghezal, M.; Benachour, A.; Rusconi, S.; Aebi, M.; Conzelmann,

[76403] A.: Yeast Gpi8p is essential for GPI anchor attachment onto proteins. EMBO

[76404] J. 15: 6575-6583, 1996.

[76405] 13131.

[76406] Yu, J.; Nagarajan, S.; Knez, J. J.; Udenfriend, S.; Chen, R.; Medof,

[76407] M. E.: The affected gene underlying the class K glycosylphosphatidylinositol

[76408] (GPI) surface protein defect codes for the GPI transamidase. Proc.

[76409] Nat. Acad. Sci. 94: 12580-12585, 1997.

[76410] 13132.

[76411] Walowsky, C.; Fitzhugh, D. J.; Castano, I. B.; Ju, J. Y.; Levin,

[76412] N. A.; Christman, M. F.: The topoisomerase-related function gene

[76413] TRF4 affects cellular sensitivity to the antitumor agent camptothecin. J.

[76414] Biol. Chem. 274: 7302-7308, 1999.

[76415] 13133.

[76416] Wang, Z.; Castano, I. B.; De Las Penas, A.; Adams, C.; Christman,

[76417] M. F.: Pol kappa: a DNA polymerase required for sister chromatid

[76418] cohesion. Science 289: 774-779, 2000.

[76419] 13134.

[76420] Clark, J.; Lu, Y.-J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smed-ley,

[76421] D.; Hamoudi, R.; Linehan, W. M.; Shipley, J.; Cooper, C. S.: Fusion

[76422] of splicing factor genes PSF and NonO (p54-nrb) to the TFE3 gene in

[76423] papillary renal cell carcinoma. Oncogene 15: 2233-2239, 1997.

[76424] 13135.

[76425] Patton, J. G.; Porro, E. B.; Galceran, J.; Tempst, P.; Nadal-Ginard,

[76426] B.: Cloning and characterization of PSF, a novel pre-mRNA splicing
[76427] factor. Genes Dev. 7: 393-406, 1993.

[76428] 13136.

[76429] Augood, S. J.; Penney, J. B., Jr.; Friberg, I. K.; Breakefield,

[76430] X. O.; Young, A. B.; Ozelius, L. J.; Standaert, D. G.: Expres-sion
[76431] of the early-onset torsion dystonia gene (DYT1) in human brain. Ann.
[76432] Neurol. 43: 669-673, 1998.

[76433] 13137.

[76434] Hewett, J.; Gonzalez-Agosti, C.; Slater, D.; Ziefer, P.; Li, S.;

[76435] Bergeron, D.; Jacoby, D. J.; Ozelius, L. J.; Ramesh, V.; Breakefield,
[76436] X. O.: Mutant torsinA, responsible for early-onset torsion dystonia,
[76437] forms membrane inclusions in cultured neural cells. Hum. Molec. Genet. 9:
[76438] 1403-1413, 2000.

[76439] 13138.

[76440] Piquemal, D.; Joulia, D.; Balaguer, P.; Basset, A.; Marti, J.;

[76441] Commes, T.: Differential expression of the RTP/
Drg1/Ndr1 gene product
[76442] in proliferating and growth arrested cells. Biochim. Bio-
phys. Acta 1450:
[76443] 364-373, 1999.

[76444] 13139.

[76445] LeClerc, S.; Palaniswami, R.; Xie, B.; Govdan, M. V.: Molec-
ular
[76446] cloning and characterization of a factor that binds the hu-
man glucocorticoid
[76447] receptor gene and represses its expression. J. Biol. Chem.
266:
[76448] 17333-17340, 1991.

[76449] 13140.

[76450] Tikoo, A.; Czekay, S.; Viars, C.; White, S.; Heath, J. K.; Ar-
den,
[76451] K.; Maruta, H.: p190-A, a human tumor suppressor gene,
maps to the
[76452] chromosomal region 19q13.3 that is reportedly deleted in
some gliomas. Gene 257:
[76453] 23-31, 2000.

[76454] 13141.

[76455] Wang, A. H.; Bertos, N. R.; Vezmar, M.; Pelletier, N.; Crosato,

[76456] M.; Heng, H. H.; Th'ng, J.; Han, J.; Yang, X.-J.: HDAC4, a human

[76457] histone deacetylase related to yeast HDA1, is a transcrip- tional corepressor. Molec.

[76458] Cell. Biol. 19: 7816-7827, 1999.

[76459] 13142.

[76460] Lewin, A. R.; Reid, L. E.; McMahon, M.; Stark, G. R.; Kerr, I.

[76461] M.: Molecular analysis of a human interferon-inducible gene family. Europ.

[76462] J. Biochem. 199: 417-423, 1991.

[76463] 13143.

[76464] Orimo, A.; Yamagishi, T.; Tominaga, N.; Yamauchi, Y.; Hishinuma,

[76465] T.; Okada, K.; Suzuki, M.; Sato, M.; Nogi, Y.; Suzuki, H.; Inoue,

[76466] S.; Yoshimura, K.; Shimizu, Y.; Muramatsu, M.: Molecular cloning

[76467] of testis-abundant finger protein/ring finger protein 23

(RNF23),

[76468] a novel RING-B box-coiled coil-B30.2 protein on the class I region

[76469] of the human MHC. Biochem. Biophys. Res. Commun. 276: 45-51, 2000.

[76470] 13144.

[76471] Nishimura, Y.; Hayashi, M.; Inada, H.; Tanaka, T.: Molecular cloning

[76472] and characterization of mammalian homologues of vesicle-associated

[76473] membrane protein-associated (VAMP-associated) proteins. Biochem.

[76474] Biophys. Res. Commun. 254: 21-26, 1999.

[76475] 13145.

[76476] Weir, M. L.; Klip, A.; Trimble, W. S.: Identification of a human

[76477] homologue of the vesicle-associated membrane protein (VAMP)-associated

[76478] protein of 33 kDa (VAP-33): a broadly expressed protein that binds

[76479] to VAMP. Biochem. J. 333: 247-251, 1998.

[76480] 13146.

[76481] Bahr, C.; Rohwer, A.; Stempka, L.; Rincke, G.; Marks, F.; Gschwendt,

[76482] M.: DIK, a novel protein kinase that interacts with protein kinase

[76483] C-delta: cloning, characterization, and gene analysis. J. Biol. Chem. 275:

[76484] 36350-36357, 2000.

[76485] 13147.

[76486] Chen, L.; Haider, K.; Ponda, M.; Cariappa, A.; Rowitch, D.; Pillai,

[76487] S.: Protein kinase C-associated kinase (PKK), a novel membrane-associated,

[76488] ankyrin repeat-containing protein kinase. J. Biol. Chem. 276: 21737-21744,

[76489] 2001.

[76490] 13148.

[76491] Zhou, Z.; Reed, R.: Human homologs of yeast Prp16 and Prp17 reveal

[76492] conservation of the mechanism for catalytic step II of pre-mRNA splicing. EMBO

[76493] J. 17: 2095-2106, 1998.

[76494] 13149.

[76495] Gozani, O.; Feld, R.; Reed, R.: Evidence that sequence-
independent
[76496] binding of highly conserved U2 snRNP proteins upstream
of the branch
[76497] site is required for assembly of spliceosomal complex A.
Genes Dev. 10:
[76498] 233-243, 1996.

[76499] 13150.

[76500] Perez Jurado, L. A.; Wang, Y.-K.; Francke, U.; Cruces, J.:
TBL2,
[76501] a novel transducin family member in the WBS deletion:
characterization
[76502] of the complete sequence, genomic structure, transcrip-
tional variants
[76503] and the mouse ortholog. Cytogenet. Cell Genet. 86:
277-284, 1999.

[76504] 13151.

[76505] Bergo, M. O.; Leung, G. K.; Ambroziak, P.; Otto, J. C.;
Casey,
[76506] P. J.; Gomes, A. Q.; Seabra, M. C.; Young, S. G.: Isoprenyl-
cysteine

[76507] carboxyl methyltransferase deficiency in mice. J. Biol. Chem. 276:
[76508] 841-5845, 2001.

[76509] 13152.

[76510] Dai, Q.; Choy, E.; Chiu, V.; Romano, J.; Slivka, S. R.; Steitz,
[76511] S. A.; Michaelis, S.; Philips, M. R.: Mammalian prenylcysteine carboxyl
[76512] methyltransferase is in the endoplasmic reticulum. J. Biol. Chem. 273:
[76513] 15030-15034, 1998.

[76514] 13153.

[76515] Desrosiers, R. R.; Nguyen, Q.-T.; Beliveau, R.: The carboxyl methyltransferase
[76516] modifying G proteins is a metalloenzyme. Biochem. Biophys. Res. Commun. 261:
[76517] 790-797, 1999.

[76518] 13154.

[76519] Akhmanova, A.; Hoogenraad, C. C.; Drabek, K.; Stepanova, T.; Dortland,
[76520] B.; Verkerk, T.; Vermeulen, W.; Burgering, B. M.; De Zeeuw, C. I.;

[76521] Grosveld, F.; Galjart, N.: CLASPs are CLIP-115 and -170 associating
[76522] proteins involved in the regional regulation of microtubule dynamics
[76523] in motile fibroblasts. Cell 104: 923-935, 2001.

[76524] 13155.

[76525] Cattanach, B. M.; Barr, J. A.; Beechey, C. V.; Martin, J.; Noebels,
[76526] J.; Jones, J.: A candidate model for Angelman syndrome in the mouse. Mammalian
[76527] Genome 8: 472-478, 1997.

[76528] 13156.

[76529] Dhar, M.; Webb, L. S.; Smith, L.; Hauser, L.; Johnson, D.; West,
[76530] D. B.: A novel ATPase on mouse chromosome 7 is a candidate gene for
[76531] increased body fat. Physiol. Genomics 4: 93-100, 2000.

[76532] 13157.

[76533] Halleck, M. S.; Lawler, J. F., Jr.; Blackshaw, S.; Gao, L.; Nagarajan,
[76534] P.; Hacker, C.; Pyle, S.; Newman, J. T.; Nakanishi, Y.; Ando,

H.;

[76535] Weinstock, D.; Williamson, P.; Schlegel, R. A.: Differential expression

[76536] of putative transbilayer amphipath transporters. Physiol. Genomics 1:

[76537] 139-150, 1999.

[76538] 13158.

[76539] Otsuki, T.; Kajigaya, S.; Ozawa, K.; Liu, J. M.: SNX5, a new member

[76540] of the sorting nexin family, binds to the Fanconi anemia complementation

[76541] group A protein. Biochem. Biophys. Res. Commun. 265: 630-635, 1999.

[76542] 13159.

[76543] Choi, D.-K.; Suzuki, Y.; Yoshimura, S.; Togashi, T.; Hida, M.;

[76544] Taylor, T. D.; Wang, Y.; Sugano, S.; Hattori, M.; Sakaki, Y.: Molecular

[76545] cloning and characterization of a gene expressed in mouse developing

[76546] tongue, mDscr5 gene, a homolog of human DSCR5 (Down syndrome critical

[76547] region gene 5). Mammalian Genome 12: 347-351, 2001.

[76548] 13160.

[76549] Shibuya, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.; Asakawa, S.;

[76550] Shimizu, N.: Isolation of two novel genes, DSCR5 and DSCR6, from

[76551] Down syndrome critical region on human chromosome 21q22.2. Biochem.

[76552] Biophys. Res. Commun. 271: 693-698, 2000.

[76553] 13161.

[76554] Togashi, T.; Choi, D.-K.; Taylor, T. D.; Suzuki, Y.; Sugano, S.;

[76555] Hattori, M.; Sakaki, Y.: A novel gene, DSCR5, from the dis-tal Down

[76556] syndrome critical region on chromosome 21q22.2. DNA Res. 7: 207-212,

[76557] 2000.

[76558] 13162.

[76559] Nakao, M.; Shichijo, S.; Imaizumi, T.; Inoue, Y.; Mat-sunaga, K.;

[76560] Yamada, A.; Kikuchi, M.; Tsuda, N.; Ohta, K.; Takamori, S.;

Yamana,
[76561] H.; Fujita, H.; Itoh, K.: Identification of a gene coding for a new
[76562] squamous cell carcinoma antigen recognized by the CTL. J. Immun. 164:
[76563] 2565-2574, 2000.

[76564] 13163.

[76565] Loftus, B. J.; Kim, U.-J.; Sneddon, V. P.; Kalush, F.; Bran-don,
[76566] R.; Fuhrmann, J.; Mason, T.; Crosby, M. L.; Barnstead, M.; Cronin,
[76567] L.; Mays, A. D.; Cao, Y.; Xu, R. X.; Kang, H.-L.; Mitchell, S.; Eichler,
[76568] E. E.; Harris, P. C.; Venter, J. C.; Adams, M. D.: Genome duplications
[76569] and other features in 12 Mb of DNA sequence from hu-man chromosome
[76570] 16p and 16q. Genomics 60: 295-308, 1999.

[76571] 13164.

[76572] Brauner-Osborne, H.; Krogsgaard-Larsen, P.: Sequence and expression
[76573] pattern of a novel human orphan G-protein-coupled re-

ceptor, GPRC5B,

[76574] a family C receptor with a short amino-terminal domain. Genomics 65:

[76575] 121-128, 2000.

[76576] 13165.

[76577] Robbins, M. J.; Michalovich, D.; Hill, J.; Calver, A. R.; Med-

hurst,

[76578] A. D.; Gloger, I.; Sims, M.; Middlemiss, D. N.; Pangalos, M. N.:

[76579] Molecular cloning and characterization of two novel retinoic acid-inducible

[76580] orphan G-protein-coupled receptors (GPRC5B and GPRC5C). Genomics 67:

[76581] 8-18, 2000.

[76582] 13166.

[76583] Autieri, M. V.; Feuerstein, G. Z.; Yue, T.-L.; Ohlstein, E. H.;

[76584] Douglas, S. A.: Use of differential display to identify dif-

ferentially

[76585] expressed mRNAs induced by rat carotid artery balloon angioplasty. Lab.

[76586] Invest. 72: 656-661, 1995.

[76587] 13167.

[76588] Autieri, M. V.; Haines, D. S.; Romanic, A. M.; Ohlstein, E. H.

[76589] : Expression of 14-3-3-gamma in injured arteries and growth factor-

[76590] and cytokine-stimulated human vascular smooth muscle cells. Cell

[76591] Growth Diff. 7: 1453-1460, 1996.

[76592] 13168.

[76593] Horie, M.; Suzuki, M.; Takahashi, E.; Tanigami, A.: Cloning, expression,

[76594] and chromosomal mapping of the human 14-3-3gamma gene (YWHAG) to 7q11.23. Genomics 60:

[76595] 241-243, 1999.

[76596] 13169.

[76597] Morrison, D.: 14-3-3: modulators of signaling proteins? Science 266:

[76598] 56-57, 1994.

[76599] 13170.

[76600] Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding, R.-G.; Kanbara,

[76601] N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Cleft palate and
[76602] decreased brain gamma-aminobutyric acid in mice lacking the 67-kDa
[76603] isoform of glutamic acid decarboxylase. Proc. Nat. Acad. Sci. 94:
[76604] 6496-6499, 1997.

[76605] 13171.

[76606] Brilliant, M. H.; Szabo, G.; Katarova, Z.; Kozak, C. A.; Glaser,
[76607] T. M.; Greenspan, R. J.; Housman, D. E.: Sequences ho-mologous to
[76608] glutamic acid decarboxylase cDNA are present on mouse chromosomes
[76609] 2 and 10. Genomics 6: 115-122, 1990.

[76610] 13172.

[76611] Bu, D.-F.; Tobin, A. J.: The exon-intron organization of the genes
[76612] (GAD1 and GAD2) encoding two human glutamate decar-boxylases (GAD-67
[76613] and GAD-65) suggests that they derive from a common ancestral GAD. Genomics 21:

[76614] 222-228, 1994.

[76615] 13173.

[76616] Condie, B. G.; Bain, G.; Gottlieb, D. I.; Capecchi, M. R.: Cleft

[76617] palate in mice with a targeted mutation in the gamma-aminobutyric

[76618] acid-producing enzyme glutamic acid decarboxylase 67. Proc. Nat.

[76619] Acad. Sci. 94: 11451-11455, 1997.

[76620] 13174.

[76621] Erlander, M. G.; Tillakaratne, N. J. K.; Feldblum, S.; Patel, N.;

[76622] Tobin, A. J.: Two genes encode distinct glutamate decar-boxylases. Neuron 7:

[76623] 91-100, 1991.

[76624] 13175.

[76625] Kelly, C. D.; Edwards, Y.; Johnstone, A. P.; Harfst, E.; No-gradi,

[76626] A.; Nussey, S. S.; Povey, S.; Carter, N. D.: Nucleotide se-quence

[76627] and chromosomal assignment of a cDNA encoding the

large isoform of
[76628] human glutamate decarboxylase. Ann. Hum. Genet. 56: 255-265, 1992.
[76629] 13176.

[76630] Krishnamoorthy, K. S.: Pyridoxine-dependency seizure: report
[76631] of a rare presentation. Ann. Neurol. 13: 103-104, 1983.

[76632] 13177.

[76633] Sparkes, R. S.; Kaufman, D. L.; Heinzmann, C.; Tobin, A. J.; Mohandas,
[76634] T.: Brain glutamate decarboxylase (GAD) gene assigned to human chromosome
[76635] 2 by somatic cell hybrid analysis. (Abstract) Cytogenet. Cell Genet. 46:
[76636] 696 only, 1987.

[76637] 13178.

[76638] Scott, A. F.: Personal Communication. Baltimore, Md. 1/22/2001.
[76639] 13179.

[76640] Zhu, H.; Kavsak, P.; Abdollah, S.; Wrana, J. L.; Thomsen, G. H.

[76641] : A SMAD ubiquitin ligase targets the BMP pathway and affects embryonic

[76642] pattern formation. Nature 400: 687-693, 1999.

[76643] 13180.

[76644] Drivas, G. T.; Shih, A.; Coutavas, E. E.; D'Eustachio, P.; Rush,

[76645] M. G.: Identification and characterization of a human homolog of

[76646] the Schizosaccharomyces pombe ras-like gene YPT-3. Oncogene 6: 3-9,

[76647] 1991.

[76648] 13181.

[76649] Gromov, P. S.; Celis, J. E.; Hansen, C.; Tommerup, N.; Gromova,

[76650] I.; Madsen, P.; Human rab11a: transcription, chromosome mapping and

[76651] effect on the expression levels of host GTP-binding proteins. FEBS

[76652] Lett. 429: 359-364, 1998.

[76653] 13182.

[76654] Kedra, D.; Seroussi, E.; Fransson, I.; Trifunovic, J.; Clark,

M.;

[76655] Lagercrantz, J.; Blennow, E.; Mehlin, H.; Dumanski, J.: The germinal

[76656] center kinase gene and a novel CDC25-like gene are located in the

[76657] vicinity of the PYGM gene on 11q13. Hum. Genet. 100: 611-619, 1997.

[76658] 13183.

[76659] Yao, Z.; Spriggs, M. K.; Derry, J. M. J.; Strockbine, L.; Park,

[76660] L. S.; VandenBos, T.; Zappone, J.; Painter, S. L.; Armitage, R. J.

[76661] : Molecular characterization of the human interleukin (IL)-17 receptor. Cytokine 9:

[76662] 794-800, 1997.

[76663] 13184.

[76664] Hogue, D. L.; Liu, L.; Ling, V.: Identification and characterization

[76665] of a mammalian mitochondrial ATP-binding cassette membrane protein. J.

[76666] Molec. Biol. 285: 379-389, 1999.

[76667] 13185.

[76668] Liang, H.; Guo, W.; Nagarajan, L.: Chromosomal mapping and genomic
[76669] organization of an evolutionarily conserved zinc finger gene ZNF277. Genomics 66:
[76670] 226-228, 2000.

[76671] 13186.

[76672] Ichigotani, Y.; Matsuda, S.; Machida, K.; Oshima, K.; Iwamoto,
[76673] T.; Yamaki, K.; Hayakawa, T.; Hamaguchi, M.: Molecular cloning of
[76674] a novel human gene (SIRP-B2) which encodes a new mem-ber of the SIRP/SHPS-1
[76675] protein family. J. Hum. Genet. 45: 378-382, 2000.

[76676] 13187.

[76677] Hirsch, D. S.; Pirone, D. M.; Burbelo, P. D.: A new family of

[76678] Cdc42 effector proteins, CEPs, function in fibroblast and epithelial
[76679] cell shape changes. J. Biol. Chem. 276: 875-883, 2001.

[76680] 13188.

[76681] Joberty, G.; Perlungher, R. R.; Macara, I. G.: The Borgs, a new

[76682] family of Cdc42 and TC10 GTPase-interacting proteins. Molec. Cell.

[76683] Biol. 19: 6585-6597, 1999.

[76684] 13189.

[76685] Yang, Z.-Q.; Imoto, I.; Fukuda, Y.; Pimkhaokham, A.; Shi-

[76686] mada, Y.; Imamura, M.; Sugano, S.; Nakamura, Y.; Inazawa, J.: Iden-

[76687] tification of a novel gene, GASC1, within an amplicon at 9p23-24 frequently detected

[76688] in esophageal cancer cell lines. Cancer Res. 60: 4735-4739, 2000.

[76689] 13190.

[76690] Cheung, P. C. F.; Trinkle-Mulcahy, L.; Cohen, P.; Lucocq, J. M.

[76691] : Characterization of a novel phosphatidylinositol 3-phosphate-binding

[76692] protein containing two FYVE fingers in tandem that is tar-

[76693] geted to the Golgi. Biochem. J. 355: 113-121, 2001.

[76694] 13191.

[76695] Derubeis, A. R.; Young, M. F.; Jia, L.; Robey, P. G.; Fisher, L.

[76696] W.: Double FYVE-containing protein 1 (DFCP1): isolation, cloning

[76697] and characterization of a novel FYVE finger protein from a human bone

[76698] marrow cDNA library. Gene 255: 195-203, 2000.

[76699] 13192.

[76700] Doyle, J. L.; DeSilva, U.; Miller, W.; Green, E. D.: Divergent

[76701] human and mouse orthologs of a novel gene (WBSCR15/Wbscr15) reside

[76702] within the genomic interval commonly deleted in Williams syndrome. Cytogenet.

[76703] Cell Genet. 90: 285-290, 2000.

[76704] 13193.

[76705] Lesage, F.; Maingret, F.; Lazdunski, M.: Cloning and ex-pression

[76706] of human TRAAK, a polyunsaturated fatty acids-activated and mechano-sensitive

[76707] K+ channel. FEBS Lett. 471: 137-140, 2000.

[76708] 13194.

[76709] Liu, Y.; McKenna, E.; Figueroa, D. J.; Blevins, R.; Austin, C.

[76710] P.; Bennett, P. B.; Swanson, R.: The human inward rectifier K+ channel

[76711] subunit Kir5.1 (KCNJ16) maps to chromosome 17q25 and is expressed

[76712] in kidney and pancreas. Cytogenet. Cell Genet. 90: 60-63, 2000.

[76713] 13195.

[76714] Nishimura, D. Y.; Searby, C. C.; Carmi, R.; Elbedour, K.; Van Maldergem,

[76715] L.; Fulton, A. B.; Lam, B. L.; Powell, B. R.; Swiderski, R. E.; Bugge,

[76716] K. E.; Haider, N. B.; Kwitek-Black, A. E.; Ying, L.; Duhl, D. M.;

[76717] Gorman, S. W.; Heon, E.; Iannaccone, A.; Bonneau, D.; Biesecker, L.

[76718] G.; Jacobson, S. G.; Stone, E. M.; Sheffield, V. C.: Positional cloning

[76719] of a novel gene on chromosome 16q causing Bardet-Biedl syndrome (BBS2). Hum.

[76720] Molec. Genet. 10: 865-874, 2001.

[76721] 13196.

[76722] LeCouter, J.; Kowalski, J.; Foster, J.; Hass, P.; Zhang, Z.; Dillard-Telm,

[76723] L.; Frantz, G.; Rangell, L.; DeGuzman, L.; Keller, G.-A.; Peale, F.;

[76724] Gurney, P.; Hillan, K. J.; Ferrara, N.: Identification of an angiogenic

[76725] mitogen selective for endocrine gland endothelium. Nature 412: 877-884,

[76726] 2001.

[76727] 13197.

[76728] Li, M.; Bullock, C. M.; Knauer, D. J.; Ehlert, F. J.; Zhou, Q.

[76729] Y.: Identification of two prokineticin cDNAs: recombinant proteins

[76730] potently contract gastrointestinal smooth muscle. Molec. Pharm. 59:

[76731] 692-698, 2001.

[76732] 13198.

[76733] Nagase, T.; Nakayama, M.; Nakajima, D.; Kikuno, R.; Ohara, O.:

[76734] Prediction of the coding sequences of unidentified human genes. XX.

[76735] The complete sequences of 100 new cDNA clones from brain which code

[76736] for large proteins in vitro. DNA Res. 8: 85-95, 2001.

[76737] 13199.

[76738] Telford, E. A. R.; Moynihan, L. M.; Markham, A. F.; Lench, N. J.

[76739] : Isolation and characterisation of a cDNA encoding the precursor

[76740] for a novel member of the acyl-CoA dehydrogenase gene family. Biochim.

[76741] Biophys. Acta 1446: 371-376, 1999.

[76742] 13200.

[76743] Cerretti, D. P.; DuBose, R. F.; Black, R. A.; Nelson, N.: Isolation

[76744] of two novel metalloproteinase-disintegrin (ADAM) cDNAs that show

[76745] testis-specific gene expression. Biochem. Biophys. Res. Commun. 263:

[76746] 810-815, 1999.

[76747] 13201.

[76748] Xu, R.; Cai, J.; Xu, T.; Zhou, W.; Ying, B.; Deng, K.; Zhao, S.;

[76749] Li, C.: Molecular cloning and mapping of a novel ADAM gene (ADAM29)
[76750] to human chromosome 4. Genomics 62: 537-539, 1999.

[76751] 13202.

[76752] Jacobs, S.; Schilf, C.; Fliegert, F.; Koling, S.; Weber, Y.; Schurmann,
[76753] A.; Joost, H.-G.: ADP-ribosylation factor (ARF)-like 4, 6, and 7
[76754] represent a subgroup of the ARF family characterized by rapid nucleotide
[76755] exchange and a nuclear localization signal. FEBS Lett. 456: 384-388,
[76756] 1999.

[76757] 13203.

[76758] Inoue, S.; Inoue, K.; Utsunomiya, M.; Nozaki, J.; Yamada, Y.;
[76759] Iwasa, T.; Mori, E.; Yoshinaga, T.; Koizumi, A.: Mutation analysis
[76760] in PKD1 of Japanese autosomal dominant polycystic kid-ney disease patients. Hum.
[76761] Mutat. 19: 622-628, 2002.

[76762] 13204.

[76763] Kim, K.; Drummond, I.; Ibraghimov-Beskrovnaya, O.;
Klinger, K.;
[76764] Arnaout, M. A.: Polycystin 1 is required for the structural
integrity
[76765] of blood vessels. Proc. Nat. Acad. Sci. 97: 1731-1736,
2000.

[76766] 13205.

[76767] Koptides, M.; Deltas, C. C.: Autosomal dominant polycys-
tic kidney
[76768] disease: molecular genetics and molecular pathogenesis.
Hum. Genet. 107:
[76769] 115-126, 2000.

[76770] 13206.

[76771] Lohning, C.; Nowicka, U.; Frischauf, A.-M.: The mouse ho-
molog
[76772] of PKD1: sequence analysis and alternative splicing. Mam-
malian Genome 8:
[76773] 307-311, 1997.

[76774] 13207.

[76775] Lu, W.; Peissel, B.; Babakhanlou, H.; Pavlova, A.; Geng, L.;

Fan,
[76776] X.; Larson, C.; Brent, G.; Zhou, J.: Perinatal lethality with kidney
[76777] and pancreas defects in mice with a targetted Pkd1 mutation. Nature
[76778] Genet. 17: 179-181, 1997.

[76779] 13208.

[76780] Lu, W.; Shen, X.; Pavlova, A.; Lakkis, M.; Ward, C. J.; Pritchard,
[76781] L.; Harris, P. C.; Genest, D. R.; Perez-Atayde, A. R.; Zhou, J.:
[76782] Comparison of Pkd1-targeted mutants reveals that loss of polycystin-1
[76783] causes cystogenesis and bone defects. Hum. Molec. Genet. 10: 2385-2396,
[76784] 2001.

[76785] 13209.

[76786] McCluskey, M.; Schiavello, T.; Hunter, M.; Hantke, J.; Angelicheva,
[76787] D.; Bogdanova, N.; Markoff, A.; Thomas, M.; Dworniczak, B.; Horst,
[76788] J.; Kalaydjieva, L.: Mutation detection in the duplicated re-

gion

[76789] of the polycystic kidney disease 1 (PKD1) gene in PKD1-linked Australian

[76790] families. Hum. Mutat. 19: 240-250, 2002.

[76791] 13210.

[76792] Murti, J. R.; Bumbulis, M.; Schimenti, J. C.: Gene conversion

[76793] between unlinked sequences in the germline of mice. Genetics 137:

[76794] 837-843, 1994.

[76795] 13211.

[76796] Neophytou, P.; Constantinides, R.; Lazarou, A.; Pierides, A.;

[76797] Constantinou Deltas, C.: Detection of a novel nonsense mutation and

[76798] an intragenic polymorphism in the PKD1 gene of a Cypriot family with

[76799] autosomal dominant polycystic kidney disease. Hum. Genet. 98: 437-442,

[76800] 1996.

[76801] 13212.

[76802] Olsson, P. G.; Lohning, C.; Horsley, S.; Kearney, L.; Harris,

[76803] P. C.; Frischauf, A.-M.: The mouse homologue of the polycystic kidney

[76804] disease gene (Pkd1) is a single-copy gene. Genomics 34: 233-235,

[76805] 1996.

[76806] 13213.

[76807] Peral, B.; Gamble, V.; Strong, C.; Ong, A. C. M.; Sloane-Stanley,

[76808] J.; Zerres, K.; Winearls, C. G.; Harris, P. C.: Identification of

[76809] mutations in the duplicated region of the polycystic kidney disease

[76810] 1 gene (PKD1) by a novel approach. Am. J. Hum. Genet. 60: 1399-1410,

[76811] 1997.

[76812] 13214.

[76813] Perrichot, R. A.; Mercier, B.; Simon, P. M.; Whebe, B.; Cledes,

[76814] J.; Ferec, C.: DGGE screening of PKD1 gene reveals novel mutations

[76815] in a large cohort of 146 unrelated patients. Hum. Genet.

105: 231-239,
[76816] 1999.

[76817] 13215.

[76818] Pritchard, L.; Sloane-Stanley, J. A.; Sharpe, J. A.; Aspinwall,

[76819] R.; Lu, W.; Buckle, V.; Strmecki, L.; Walker, D.; Ward, C. J.;
Alpers,
[76820] C. E.; Zhou, J.; Wood, W. G.; Harris, P. C.: A human PKD1
transgene
[76821] generates functional polycystin-1 in mice and is associ-
ated with a
[76822] cystic phenotype. Hum. Molec. Genet. 9: 2617-2627,
2000.
[76823] 13216.

[76824] Lako, M.; Lindsay, S.; Bullen, P.; Wilson, D. I.; Robson, S.
C.;
[76825] Strachan, T.: A novel mammalian Wnt gene, WNT8B, shows
brain-restricted
[76826] expression in early development, with sharply delimited
expression
[76827] boundaries in the developing forebrain. Hum. Molec.
Genet. 7: 813-822,
[76828] 1998.

[76829] 13217.

[76830] Lako, M.; Strachan, T.; Curtis, A. R. J.; Lindsay, S.: Isolation

[76831] and characterization of WNT8B, a novel human Wnt gene that maps to

[76832] 10q24. Genomics 35: 386-388, 1996.

[76833] 13218.

[76834] Tsubata, S.; Bowles, K. R.; Vatta, M.; Zintz, C.; Titus, J.; Muhonen,

[76835] L.; Bowles, N. E.; Towbin, J. A.: Mutations in the human delta-sarcoglycan

[76836] gene in familial and sporadic dilated cardiomyopathy. J. Clin. Invest. 106:

[76837] 655-662, 2000.

[76838] 13219.

[76839] Toda, T.; Iida, A.; Miwa, T.; Nakamura, Y.; Imai, T.: Isola-tion

[76840] and characterization of a novel gene encoding nuclear protein at a

[76841] locus (D11S636) tightly linked to multiple endocrine neo-plasia type

[76842] 1 (MEN1). Hum. Molec. Genet. 3: 465-470, 1994.

[76843] 13220.

[76844] Nusse, R.; Varmus, H. E.: Many tumors induced by the mouse mammary
[76845] tumor virus contain a provirus integrated in the same region of the
[76846] host genome. Cell 31: 99-109, 1982.

[76847] 13221.

[76848] Guymer, R. H.; Heon, E.; Lotery, A. J.; Munier, F. L.; Schorderet,
[76849] D. F.; Baird, P. N.; McNeil, R. J.; Haines, H.; Sheffield, V. C.;
[76850] Stone, E. M.: Variation of codons 1961 and 2177 of the Stargardt
[76851] disease gene is not associated with age-related macular degeneration. Arch.
[76852] Ophthal. 119: 745-751, 2001.

[76853] 13222.

[76854] Paoloni-Giacobino, A.; Chen, H.; Antonarakis, S. E.: Cloning of
[76855] a novel human neural cell adhesion molecule gene (NCAM2) that maps
[76856] to chromosome region 21q21 and is potentially involved

in Down syndrome. Genomics 43:
[76857] 43-51, 1997.

[76858] 13223.

[76859] Abdulkadir, S. A.; Magee, J. A.; Peters, T. J.; Kaleem, Z.; Naughton,
[76860] C. K.; Humphrey, P. A.; Milbrandt, J.: Conditional loss of Nkx3.1
[76861] in adult mice induces prostatic intraepithelial neoplasia. Molec.
[76862] Cell. Biol. 22: 1495-1503, 2002.

[76863] 13224.

[76864] Bieberich, C. J.; Fujita, K.; He, W. W.; Jay, G.: Prostate-spe-cific
[76865] and androgen-dependent expression of a novel home-obox gene. J. Biol.
[76866] Chem. 271: 31779-31782, 1996.

[76867] 13225.

[76868] Korkmaz, K. S.; Korkmaz, C. G; Ragnhildstveit, E.; Kizildag, S.;
[76869] Pretlow, T. G.; Saatcioglu, F.: Full-length cDNA sequence and genomic

[76870] organization of human NKX3A--alternative forms and regulation by both

[76871] androgens and estrogens. Gene 260: 25-36, 2000.

[76872] 13226.

[76873] He, W. W.; Sciavolino, P. J.; Wing, J.; Augustus, M.; Hudson, P.;

[76874] Meissner, P. S.; Curtis, R. T.; Shell, B. K.; Bostwick, D. G.; Tindall,

[76875] D. J.; Gelmann, E. P.; Abate-Shen, C.; Carter, K. C.: A novel human

[76876] prostate-specific, androgen-regulated homeobox gene (NKX3.1) that

[76877] maps to 8p21, a region frequently deleted in prostate cancer. Genomics 43:

[76878] 69-77, 1997.

[76879] 13227.

[76880] Gould, S.: Personal Communication. Baltimore, Md. 9/26/1997.

[76881] 13228.

[76882] Jansen, G. A.; Hogenhout, E. M.; Ferdinandusse, S.; Waterham, H.

[76883] R.; Ofman, R.; Jakobs, C.; Skjeldal, O. H.; Wanders, R. J. A.: Human

[76884] phytanoyl-CoA hydroxylase: resolution of the gene structure and the

[76885] molecular basis of Refsum's disease. Hum. Molec. Genet. 9: 1195-1200,

[76886] 2000.

[76887] 13229.

[76888] Mukherji, M.; Chien, W.; Kershaw, N. J.; Clifton, I. J.; Schofield,

[76889] C. J.; Wierzbicki, A. S.; Lloyd, M. D.: Structure-function analysis

[76890] of phytanoyl-CoA 2-hydroxylase mutations causing Refsum's disease. Hum.

[76891] Molec. Genet. 10: 1971-1982, 2001.

[76892] 13230.

[76893] Skjeldal, O. H.; Stokke, O.; Refsum, S.; Norseth, J.; Petit, H.

[76894] : Clinical and biochemical heterogeneity in conditions with phytanic

[76895] acid accumulation. J. Neurol. Sci. 77: 87-96, 1987.

[76896] 13231.

[76897] Jansen, G. A.; Mihalik, S. J.; Watkins, P. A.; Moser, H. W.; Jakobs,
[76898] C.; Denis, S.; Wanders, R. J. A.: Phytanoyl-CoA hydroxylase is present
[76899] in human liver, located in peroxisomes, and deficient in Zellweger
[76900] syndrome: direct, unequivocal evidence for the new, re-vised pathway
[76901] of phytanic acid alpha-oxidation in humans. Biochem. Biophys. Res.
[76902] Commun. 229: 205-210, 1996.

[76903] 13232.

[76904] Bilaud, T.; Brun, C.; Ancelin, K.; Koering, C. E.; Laroche, T.;
[76905] Gilson, E.: Telomeric localization of TRF2, a novel human telobox
[76906] protein. Nature Genet. 17: 236-239, 1997.

[76907] 13233.

[76908] Broccoli, D.; Smogorzewska, A.; Chong, L.; de Lange, T.: Human
[76909] telomeres contain two distinct Myb-related proteins, TRF1 and TRF2. Nature

[76910] Genet. 17: 231-235, 1997.

[76911] 13234.

[76912] Karlseder, J.; Smogorzewska, A.; de Lange, T.: Senescence induced

[76913] by altered telomere state, not telomere loss. Science 295: 2446-2449,

[76914] 2002.

[76915] 13235.

[76916] Sakaguchi, A. Y.; Padalecki, S. S.; Mattern, V.; Rodriguez, A.;

[76917] Leach, R. J.; McGill, J. R.; Chavez, M.; Giambernardi, T. A.: Chromosomal

[76918] sublocalization of the transcribed human telomere repeat binding factor

[76919] 2 gene and comparative mapping in the mouse. Somat. Cell Molec. Genet. 24:

[76920] 157-163, 1998.

[76921] 13236.

[76922] van Steensel, B.; Smogorzewska, A.; de Lange, T.: TRF2 protects

[76923] human telomeres from end-to-end fusions. Cell 92:

401-413, 1998.

[76924] 13237.

[76925] Cancel, G.; Durr, A.; Didierjean, O.; Imbert, G.; Burk, K.; Lezin,

[76926] A.; Belal, S.; Benomar, A.; Abada-Bendib, M.; Vial, C.; Guimaraes,

[76927] J.; Chneiweiss, H.; and 12 others: Molecular and clinical correlations

[76928] in spinocerebellar ataxia 2: a study of 32 families. Hum. Molec.

[76929] Genet. 6: 709-715, 1997.

[76930] 13238.

[76931] Choudhry, S.; Mukerji, M.; Srivastava, A. K.; Jain, S.; Brah-machari,

[76932] S. K.: CAG repeat instability at SCA2 locus: anchoring CAA interruptions

[76933] and linked single nucleotide polymorphisms. Hum. Molec. Genet. 10:

[76934] 2437-2446, 2001.

[76935] 13239.

[76936] Huynh, D. P.; Del Bigio, M. R.; Ho, D. H.; Pulst, S. M.: Ex-

pression
[76937] of ataxin-2 in brains from normal individuals and patients with Alzheimer's
[76938] disase and spinocerebellar ataxia 2. Ann. Neurol. 45: 232-241, 1999.
[76939] 13240.

[76940] Nechiporuk, T.; Huynh, D. P.; Figueroa, K.; Sahba, S.; Nechiporuk,
[76941] A.; Pulst, S.-M.: The mouse SCA2 gene: cDNA sequence, alternative
[76942] splicing and protein expression. Hum. Molec. Genet. 7: 1301-1309,
[76943] 1998.

[76944] 13241.

[76945] Sahba, S.; Nechiporuk, A.; Figueroa, K. P.; Nechiporuk, T.; Pulst,
[76946] S.-M.: Genomic structure of the human gene for spinocerebellar ataxia
[76947] type 2 (SCA2) on chromosome 12q24.1. Genomics 47: 359-364, 1998.
[76948] 13242.

[76949] Shibata, H.; Huynh, D. P.; Pulst, S.-M.: A novel protein with

[76950] RNA-binding motifs interacts with ataxin-2. Hum. Molec. Genet. 9:

[76951] 1303-1313, 2000.

[76952] 13243.

[76953] Parr, B. A.; McMahon, A. P.: Sexually dimorphic development of

[76954] the mammalian reproductive tract requires Wnt-7a. Nature 395: 707-710,

[76955] 1998.

[76956] 13244.

[76957] Barron-Casella, E. A.; Torres, M. A.; Scherer, S. W.; Heng, H.

[76958] H. Q.; Tsui, L.-C.; Casella, J. F.: Sequence analysis and chromosomal

[76959] localization of human Cap Z: conserved residues within the actin-binding

[76960] domain may link Cap Z to gelsolin/severin and profilin protein families. J.

[76961] Biol. Chem. 270: 21472-21479, 1995.

[76962] 13245.

[76963] Hart, M. C.; Korshunova, Y. O.; Cooper, J. A.: Mapping of

the

[76964] mouse actin capping protein alpha subunit genes and pseudogenes. Genomics 39:

[76965] 264-270, 1997.

[76966] 13246.

[76967] Cooper, J. A.: Personal Communication. St. Louis, Mo. 3/8/1999.

[76968] 13247.

[76969] Cardoso, C.; Mignon, C.; Hetet, G.; Grandchamps, B.; Fontes, M.;

[76970] Colleaux, L.: The human EZH2 gene: genomic organisation and revised

[76971] mapping in 7q35 within the critical region for malignant myeloid disorders. Europ.

[76972] J. Hum. Genet. 8: 174-180, 2000.

[76973] 13248.

[76974] Chen, H.; Rossier, C.; Antonarakis, S. E.: Cloning of a human

[76975] homolog of the Drosophila enhancer of zeste gene (EZH2) that maps

[76976] to chromosome 21q22.2. Genomics 38: 30-37, 1996.

[76977] 13249.

[76978] Dohner, K.; Brown, J.; Hehmann, U.; Hetzel, C.; Stewart, J.; Lowther,

[76979] G.; Scholl, C.; Frohling, S.; Cuneo, A.; Tsui, L. C.; Lichter, P.;

[76980] Scherer, S. W.; Dohner, H.: Molecular cytogenetic characterization

[76981] of a critical region in bands 7q35-q36 commonly deleted in malignant

[76982] myeloid disorders. Blood 92: 4031-4035, 1998.

[76983] 13250.

[76984] Laible, G.; Haynes, A. R.; Lebersorger, A.; O'Carroll, D.; Mattei,

[76985] M.-G.; Denny, P.; Brown, S. D. M.; Jenuwein, T.: The murine polycomb-group

[76986] genes Ezh1 and Ezh2 map close to Hox gene clusters on mouse chromosomes

[76987] 11 and 6. Mammalian Genome 10: 311-314, 1999.

[76988] 13251.

[76989] Varambally, S.; Dhanasekaran, S. M.; Zhou, M.; Barrette, T. R.;

[76990] Kumar-Sinha, C.; Sanda, M. G.; Ghosh, D.; Pienta, K. J.; Sewalt, R.

[76991] G. A. B.; Otte, A. P.; Rubin, M. A.; Chinnaiyan, A. M.: The polycomb

[76992] group protein EZH2 is involved in progression of prostate cancer. Nature 419:

[76993] 624-629, 2002.

[76994] 13252.

[76995] Batt, J.; Asa, S.; Fladd, C.; Rotin, D.: Pituitary, pancreatic

[76996] and gut neuroendocrine defects in protein tyrosine phosphatase-sigma-deficient

[76997] mice. Molec. Endocr. 16: 155-169, 2002.

[76998] 13253.

[76999] Elchebly, M.; Wagner, J.; Kennedy, T. E.; Lanctot, C.; Michaliszyn,

[77000] E.; Itie, A.; Drouin, J.; Tremblay, M. L.: Neuroendocrine dysplasia

[77001] in mice lacking protein tyrosine phosphatase sigma. Nature Genet. 21:

[77002] 330-333, 1999.

[77003] 13254.

[77004] Pulido, R.; Serra-Pages, C.; Tang, M.; Streuli, M.: The LAR/PTP

[77005] delta/PTP sigma subfamily of transmembrane protein-tyrosine-phosphatases:

[77006] multiple human LAR, PTP delta, and PTP sigma isoforms are expressed

[77007] in a tissue-specific manner and associate with the LAR-interacting

[77008] protein LIP.1. Proc. Nat. Acad. Sci. 92: 11686-11690, 1995.

[77009] 13255.

[77010] Wagner, J.; Gordon, L. A.; Heng, H. H. Q.; Tremblay, M. L.; Olsen,

[77011] A. S.: Physical mapping of receptor type protein tyrosine phosphatase

[77012] sigma (PTPRS) to human chromosome 19p13.3. Genomics 38: 76-78, 1996.

[77013] 13256.

[77014] Wallace, M. J.; Batt, J.; Fladd, C. A.; Henderson, J. T.; Skarnes,

[77015] W.; Rotin, D.: Neuronal defects and posterior pituitary hypoplasia

[77016] in mice lacking the receptor tyrosine phosphatase PTP-sigma. Nature

[77017] Genet. 21: 334-338, 1999.

[77018] 13257.

[77019] Bates, S.; Rowan, S.; Vousden, K. H.: Characterisation of human

[77020] cyclin G1 and G2: DNA damage inducible genes. Oncogene 13: 1103-1109,

[77021] 1996.

[77022] 13258.

[77023] Endo, Y.; Fujita, T.; Tamura, K.; Tsuruga, H.; Nojima, H.: Structure

[77024] and chromosomal assignment of the human cyclin G gene. Genomics 38:

[77025] 92-95, 1996.

[77026] 13259.

[77027] Horne, M. C.; Goolsby, G. L.; Donaldson, K. L.; Tran, D.; Neubauer,

[77028] M.; Wahl, A. F.: Cyclin G1 and cyclin G2 comprise a new family of

[77029] cyclins with contrasting tissue-specific and cell cycle-

regulated
[77030] expression. J. Biol. Chem. 271: 6050-6061, 1996.

[77031] 13260.

[77032] Jensen, M. R.; Factor, V. M.; Zimonjic, D. B.; Miller, M. J.; Keck,
[77033] C. L.; Thorgeirsson, S. S.: Chromosome localization and structure
[77034] of the murine cyclin G1 gene promoter sequence. Ge-
nomics 45: 297-303,
[77035] 1997.

[77036] 13261.

[77037] Spicer, A. P.; Augustine, M. L.; McDonald, J. A.: Molecular cloning
[77038] and characterization of a putative mouse hyaluronan syn-
thase. J.
[77039] Biol. Chem. 271: 23400-23406, 1996.

[77040] 13262.

[77041] Skalhegg, B. S.; Huang, Y.; Su, T.; Idzerda, R. L.; McKnight, G.
[77042] S.; Burton, K. A.: Mutation of the C-alpha subunit of PKA leads to

[77043] growth retardation and sperm dysfunction. Molec. Endocr. 16: 630-639,
[77044] 2002.

[77045] 13263.

[77046] Tasken, K.; Solberg, R.; Zhao, Y.; Hansson, V.; Jahnsen, T.; Siciliano,
[77047] M. J.: The gene encoding the catalytic subunit C-alpha of cAMP-dependent
[77048] protein kinase (locus PRKACA) localizes to human chromosome region
[77049] 19p13.1. Genomics 36: 535-538, 1996.

[77050] 13264.

[77051] Kim, J.; Uyemura, K.; Van Dyke, M. K.; Legaspi, A. J.; Rea, T.
[77052] H.; Shuai, K.; Modlin, R. L.: A role for IL-12 receptor expression
[77053] and signal transduction in host defense in leprosy. J. Immun. 167:
[77054] 779-786, 2001.

[77055] 13265.

[77056] Morton, S. M.; Bocaccio, I.; Depetris, D.; Mattei, M.; Des-

sein,

[77057] A.: Assignment of IL12RB2 to human chromosome 1p31.3-p31.2 between

[77058] D1S230 and D1S198. Cytogenet. Cell Genet. 79: 282-283, 1997.

[77059] 13266.

[77060] McCright, B.; Brothman, A. R.; Virshup, D. M.: Assignment of human

[77061] protein phosphatase 2A regulatory subunit genes B56-alpha, B56-beta,

[77062] B56-gamma, B56-delta, and B56-epsilon (PPP2R5A--PPP2R5E), highly expressed

[77063] in muscle and brain, to chromosome regions 1q41, 11q12, 3p21, 6p21.1,

[77064] and 7p11.2-to-p12. Genomics 36: 168-170, 1996.

[77065] 13267.

[77066] McCright, B.; Virshup, D. M.: Identification of a new family of

[77067] protein phosphatase 2A regulatory subunits. J. Biol. Chem. 270:

[77068] 26123-26128, 1995.

[77069] 13268.

[77070] Lewis, R. A.; Shroyer, N. F.; Singh, N.; Allikmets, R.; Hutchinson,

[77071] A.; Li, Y.; Lupski, J. R.; Leppert, M.; Dean, M.: Genotype/phenotype

[77072] analysis of a photoreceptor-specific ATP-binding cassette transporter

[77073] gene, ABCR, in Stargardt disease. Am. J. Hum. Genet. 64: 422-434,

[77074] 1999.

[77075] 13269.

[77076] Martinez-Mir, A.; Bayes, M.; Vilageliu, L.; Grinberg, D.; Ayuso,

[77077] C.; del Rio, T.; Garcia-Sandoval, B.; Bussaglia, E.; Baiget, M.; Gonzalez-Duarte,

[77078] R.; Balcells, S.: A new locus for autosomal recessive retinitis pigmentosa

[77079] (RP19) maps to 1p13-1p21. Genomics 40: 142-146, 1997.

[77080] 13270.

[77081] Martinez-Mir, A.; Paloma, E.; Allikmets, R.; Ayuso, C.; del Rio,

[77082] T.; Dean, M.; Vilageliu, L.; Gonzalez-Duarte, R.; Balcells,

S.: Retinitis
[77083] pigmentosa caused by a homozygous mutation in the Stargardt disease
[77084] gene ABCR. (Letter) Nature Genet. 18: 11-12, 1998.

[77085] 13271.

[77086] Maugeri, A.; Klevering, B. J.; Rohrschneider, K.; Blanke-
nagel,
[77087] A.; Brunner, H. G.; Deutman, A. F.; Hoyng, C. B.; Cremers, F. P. M.
[77088] : Mutations in the ABCA4 (ABCR) gene are the major cause of autosomal
[77089] recessive cone-rod dystrophy. Am. J. Hum. Genet. 67: 960-966, 2000.
[77090] 13272.

[77091] Maugeri, A.; van Driel, M. A.; van de Pol, D. J. R.; Klever-
ing,
[77092] B. J.; van Haren, F. J. J.; Tijmes, N.; Bergen, A. A. B.; Rohrschneider,
[77093] K.; Blankenagel, A.; Pinckers, A. J. L. G.; Dahl, N.; Brunner, H.
[77094] G.; Deutman, A. F.; Hoyng, C. B.; Cremers, F. P. M.: The 2588G-C

[77095] mutation in the ABCR gene is a mild frequent founder mutation in the
[77096] western European population and allows the classification of ABCR
[77097] mutations in patients with Stargardt disease. Am. J. Hum. Genet. 64:
[77098] 1024-1035, 1999.

[77099] 13273.

[77100] Molday, L. L.; Rabin, A. R.; Molday, R. S.: ABCR expression in
[77101] foveal cone photoreceptors and its role in Stargardt macular dystrophy. Nature
[77102] Genet. 25: 257-258, 2000.

[77103] 13274.

[77104] Nasonkin, I.; Illing, M.; Koehler, M. R.; Schmid, M.; Molday,
[77105] R. S.; Weber, B. H. F.: Mapping of the rod photoreceptor ABC transporter
[77106] (ABCR) to 1p21-p22.1 and identification of novel mutations in Stargardt's
[77107] disease. Hum. Genet. 102: 21-26, 1998.

[77108] 13275.

[77109] Paloma, E.; Martinez-Mir, A.; Vilageliu, L.; Gonzalez-Duarte,
[77110] R.; Balcells, S.: Spectrum of ABCA4 (ABCR) gene mutations in Spanish
[77111] patients with autosomal recessive macular dystrophies. Hum. Mutat. 17:
[77112] 504-510, 2001.

[77113] 13276.

[77114] Rivera, A.; White, K.; Stohr, H.; Steiner, K.; Hemmrich, N.; Grimm,
[77115] T.; Jurklies, B.; Lorenz, B.; Scholl, H. P. N.; Apfelstedt-Sylla,
[77116] E.; Weber, B. H. F.: A comprehensive survey of sequence variation
[77117] in the ABCA4 (ABCR) gene in Stargardt disease and age-related macular
[77118] degeneration. Am. J. Hum. Genet. 67: 800-813, 2000.

[77119] 13277.

[77120] Shroyer, N. F.; Lewis, R. A.; Lupski, J. R.: Complex inheri-tance
[77121] of ABCR mutations in Stargardt disease: linkage disequi-librium, complex

[77122] alleles, and pseudodominance. Hum. Genet. 106: 244-248, 2000.

[77123] 13278.

[77124] Shroyer, N. F.; Lewis, R. A.; Yatsenko, A. N.; Wensel, T. G.;

[77125] Lupski, J. R.: Cosegregation and functional analysis of mutant ABCR

[77126] (ABCA4) alleles in families that manifest both Stargardt disease and

[77127] age-related macular degeneration. Hum. Molec. Genet. 10: 2671-2678,

[77128] 2001.

[77129] 13279.

[77130] Sun, H.; Nathans, J.: Stargardt's ABCR is localized to the disc

[77131] membrane of retinal rod outer segments. (Letter) Nature Genet. 17:

[77132] 15-16, 1997.

[77133] 13280.

[77134] Sun, H.; Smallwood, P. M.; Nathans, J.: Biochemical defects in

[77135] ABCR protein variants associated with human

retinopathies. Nature
[77136] Genet. 26: 242-246, 2000.

[77137] 13281.

[77138] Weng, J.; Mata, N. L.; Azarian, S. M.; Tzekov, R. T.; Birch, D.
[77139] G.; Travis, G. H.: Insights into the function of Rim protein in photoreceptors
[77140] and etiology of Stargardt's disease from the phenotype in abcr knockout
[77141] mice. Cell 98: 13-23, 1999.

[77142] 13282.

[77143] Yatsenko, A. N.; Shroyer, N. F.; Lewis, R. A.; Lupski, J. R.:

[77144] Late-onset Stargardt disease is associated with missense mutations
[77145] that map outside known functional regions of ABCR (ABCA4). Hum. Genet. 108:
[77146] 346-355, 2001.

[77147] 13283.

[77148] Escribano, J.; Hernando, N.; Ghosh, S.; Crabb, J.; Coca-Prados,
[77149] M.: cDNA from human ocular ciliary epithelium homolo-

gous to beta
[77150] ig-h3 is preferentially expressed as an extracellular protein in the
[77151] corneal epithelium. J. Cell Physiol. 160: 511-521, 1994.

[77152] 13284.

[77153] Gupta, S. K.; Hodge, W. G.; Damji, K. F.; Guernsey, D. L.; Neumann,
[77154] P. E.: Lattice corneal dystrophy type 1 in a Canadian kindred is
[77155] associated with the arg124-to-cys mutation in the keratoepithelin
[77156] gene. Am. J. Ophthal. 125: 547-549, 1998.

[77157] 13285.

[77158] Korvatska, E.; Munier, F. L.; Djemai, A.; Wang, M. X.; Frueh, B.;
[77159] Chiou, A. G.-Y.; Uffer, S.; Ballestrazzi, E.; Braunstein, R. E.; Forster,
[77160] R. K.; Culbertson, W. W.; Boman, H.; Zografos, L.; Schorderet, D.
[77161] F.: Mutation hot spots in 5q31-linked corneal dystrophies. Am. J.
[77162] Hum. Genet. 62: 320-324, 1998.

[77163] 13286.

[77164] Mashima, Y.; Nakamura, Y.; Noda, K.; Konishi, M.; Ya-
mada, M.; Kudoh,
[77165] J.; Shimizu, N.: A novel mutation at codon 124 (R124L) in
the BIGH3
[77166] gene is associated with a superficial variant of granular
corneal
[77167] dystrophy. Arch. Ophthal. 117: 90-93, 1999.

[77168] 13287.

[77169] Munier, F. L.; Korvatska, E.; Djemai, A.; Paslier, D. L.; Zo-
grafos,
[77170] L.; Pescia, G.; Schordert, D. F.: Kerato-epithelin mutations
in four
[77171] 5q31-linked corneal dystrophies. Nat. Genet. 15:
247-251, 1997.
[77172] 13288.

[77173] Okada, M.; Yamamoto, S.; Tsujikawa, M.; Watanabe, H.;
Inoue, Y.;
[77174] Maeda, N.; Shimomura, Y.; Nishida, K.; Quantock, A. J.; Ki-
noshita,
[77175] S.; Tano, Y.: Two distinct kerato-epithelin mutations in
Reis-Bucklers

[77176] corneal dystrophy. Am. J. Ophthal. 126: 535-542, 1998.

[77177] 13289.

[77178] Okada, M.; Yamamoto, S.; Watanabe, H.; Inoue, Y.; Tsu-
jikawa, M.;

[77179] Maeda, N.; Shimomura, Y.; Nishida, K.; Kinoshita, S.; Tano,
Y.: Granular

[77180] corneal dystrophy with homozygous mutations in the ker-
ato-epithelin

[77181] gene. Am. J. Ophthal. 126: 169-176, 1998.

[77182] 13290.

[77183] Faure, S.; Meyer, L.; Costagliola, D.; Vaneensberghe, C.;
Genin,

[77184] E.; Autran, B.; French ALT and IMMUNOCO Study Groups;
Delfraisay,

[77185] J.-F.; SEROCO Study Group; McDermott, D. H.; Murphy, P.
M.; Debre,

[77186] P.; Theodorou, I.; Cambadiere, C.: Rapid progression to
AIDS in HIV+

[77187] individuals with a structural variant of the chemokine re-
ceptor CX(3)CR1. Science 287:

[77188] 2274-2277, 2000.

[77189] 13291.

[77190] Imai, T.; Hieshima, K.; Haskell, C.; Baba, M.; Nagira, M.; Nishimura,

[77191] M.; Kakizaki, M.; Takagi, S.; Nomiyama, H.; Schall, T. J.; Yoshie,

[77192] O.: Identification and molecular characterization of fractalkine

[77193] receptor CX3CR1, which mediates both leukocyte migration and adhesion. Cell 91:

[77194] 521-530, 1997.

[77195] 13292.

[77196] Moatti, D.; Faure, S.; Fumeron, F.; Amara, M. E. W.; Seknadji,

[77197] P.; McDermott, D. H.; Debre, P.; Aumont, M. C.; Murphy, P. M.; de

[77198] Prost, D.; Combadiere, C.: Polymorphism in the fractalkine receptor

[77199] CX3CR1 as a genetic risk factor for coronary artery disease. Blood 97:

[77200] 1925-1928, 2001.

[77201] 13293.

[77202] Raport, C. J.; Schweickart, V. L.; Eddy, R. L., Jr.; Shows, T.

[77203] B.; Gray, P. W.: The orphan G-protein-coupled receptor-encoding gene

[77204] V28 is closely related to genes for chemokine receptors and is expressed

[77205] in lymphoid and neural tissues. Gene 163: 295-299, 1995.

[77206] 13294.

[77207] Tripp, R. A.; Jones, L. P.; Haynes, L. M.; Zheng, H.; Murphy, P.

[77208] M.; Anderson, L. J.: CX3C chemokine mimicry by respira-tory syncytial

[77209] virus G glycoprotein. Nature Immun. 2: 732-738, 2001.

[77210] 13295.

[77211] Adra, C. N.; Zhu, S.; Ko, J.-L.; Guillemot, J.-C.; Cuervo, A. M.;

[77212] Kobayashi, H.; Horiuchi, T.; Lelias, J.-M.; Rowley, J. D.; Lim, B.

[77213] : LAPTM5: a novel lysosomal-associated multispanning membrane protein

[77214] preferentially expressed in hematopoietic cells. Genomics 35: 328-337,

[77215] 1996.

[77216] 13296.

[77217] Bondurand, N.; Kuhlbrodt, K.; Pingault, V.; Enderich, J.; Sajus,

[77218] M.; Tommerup, N.; Warburg, M.; Hennekam, R. C. M.; Read, A. P.; Wegner,

[77219] M.; Goossens, M.: A molecular analysis of the Yemenite deaf-blind

[77220] hypopigmentation syndrome: SOX10 dysfunction causes different neurocristopathies. Hum.

[77221] Molec. Genet. 8: 1785-1789, 1999.

[77222] 13297.

[77223] Hennekam, R. C. M.; Gorlin, R. J.: Confirmation of Yemenite (Warburg)

[77224] deaf-blind hypopigmentation syndrome. Am. J. Med. Genet. 65: 146-148,

[77225] 1996.

[77226] 13298.

[77227] Hanes, J.; von der Kammer, H.; Klaudiny, J.; Scheit, K. H.: Characterization

[77228] by cDNA cloning of two new human protein kinases: evi-

dence by sequence
[77229] comparison of a new family of mammalian protein ki-
nases. J. Molec.
[77230] Biol. 244: 665-672, 1994.

[77231] 13299.

[77232] Talmadge, C. B.; Finkernagel, S.; Sumegi, J.; Sciorra, L.;
Rabinow,
[77233] L.: Chromosomal mapping of three human LAMMER pro-
tein-kinase-encoding
[77234] genes. Hum. Genet. 103: 523-524, 1998.

[77235] 13300.

[77236] Fisher, R. P.; Morgan, D. O.: A novel cyclin associates with
MO15/CDK7
[77237] to form the CDK-activating kinase. Cell 78: 713-724,
1994.
[77238] 13301.

[77239] Shiekhattar, R.; Mermelstein, F.; Fisher, R. P.; Drapkin, R.;
Dynlacht,
[77240] B.; Wessling, H. C.; Morgan, D. O.; Reinberg: Cdk-
activating kinase
[77241] complex is a component of human transcription factor

TFIIH. Nature 374:
[77242] 283-287, 1995.

[77243] 13302.

[77244] Alders, M.; Hodges, M.; Hadjantonakis, A.-K.; Postmus, J.; van
[77245] Wijk, I.; Bliek, J.; de Meulemeester, M.; Westerveld, A.; Guillemot,
[77246] F.; Oudejans, C.; Little, P.; Mannens, M.: The human Achaete-Scute
[77247] homologue 2 (ASCL2,HASH2) maps to chromosome 11p15.5, close to IGF2
[77248] and is expressed in extravillus trophoblasts. Hum. Molec. Genet. 6:
[77249] 859-867, 1997.

[77250] 13303.

[77251] Guillemot, F.; Nagy, A.; Auerbach, A.; Rossant, J.; Joyner, A.
[77252] L.: Essential role of Mash-2 in extraembryonic develop-
ment. Nature 371:
[77253] 333-336, 1994.

[77254] 13304.

[77255] Banga, S. S.; Ozer, H. L.; Park, S.-K.; Lee, S.-T.: Assign-
ment
[77256] of PTK7 encoding a receptor protein tyrosine kinase-like
molecule
[77257] to human chromosome 6p21.1-p12.2 by fluorescence in
situ hybridization. Cytogenet.
[77258] Cell Genet. 76: 43-44, 1997.

[77259] 13305.

[77260] Lee, S.-T.; Strunk, K. M.; Spritz, R. A.: A survey of protein

[77261] tyrosine kinase mRNAs expressed in normal human
melanocytes. Oncogene 8:
[77262] 3403-3410, 1993.

[77263] 13306.

[77264] Mossie, K.; Jallal, B.; Alves, F.; Sures, I.; Plowman, G. D.;
Ullrich,
[77265] A.: Colon carcinoma kinase-4 defines a new subclass of
the receptor
[77266] tyrosine kinase family. Oncogene 11: 2179-2184, 1995.

[77267] 13307.

[77268] Park, S.-K.; Lee, H.-S.; Lee, S.-T.: Characterization of the
human

[77269] full-length PTK7 cDNA encoding a receptor protein tyrosine kinase-like

[77270] molecule closely related to chick KLG. J. Biochem. 119: 235-239,

[77271] 1996.

[77272] 13308.

[77273] Kohler, M.; Ansieau, S.; Prehn, S.; Leutz, A.; Haller, H.; Hartmann,

[77274] E.: Cloning of two novel human importin-alpha subunits and analysis

[77275] of the expression pattern of the importin-alpha protein family. FEBS

[77276] Lett. 417: 104-108, 1997.

[77277] 13309.

[77278] Moore, M. S.; Blobel, G.: The two steps of nuclear import, targeting

[77279] to the nuclear envelope and translocation through the nuclear pore,

[77280] require different cytosolic factors. Cell 69: 939-950, 1992.

[77281] 13310.

[77282] Takeda, S.; Fujiwara, T.; Shimizu, F.; Kawai, A.; Shinomiya, K.;

[77283] Okuno, S.; Ozaki, K.; Katagiri, T.; Shimada, Y.; Nagata, M.; Watanabe,

[77284] T.; Takaichi, A.; Kuga, Y.; Suzuki, M.; Hishigaki, H.; Takahashi,

[77285] E.; Shin, S.; Nakamura, Y.; Hirai, Y.: Isolation and mapping of karyopherin

[77286] alpha-3 (KPNA3), a human gene that is highly homologous to genes encoding

[77287] Xenopus importin, yeast SRP1 and human RCH1. Cytogenet. Cell Genet. 76:

[77288] 87-93, 1997.

[77289] 13311.

[77290] Zhang, C.-Y.; Baffy, G.; Perret, P.; Krauss, S.; Peroni, O.; Grujic,

[77291] D.; Hagen, T.; Vidal-Puig, A.; Boss, O.; Kim, Y.-B.; Zheng, X. X.;

[77292] Wheeler, M. B.; Shulman, G. I.; Chan, C. B.; Lowell, B. B.: Uncoupling

[77293] protein-2 negatively regulates insulin secretion and is a major link

[77294] between obesity, beta cell dysfunction, and type 2 diabetes. Cell 105:
[77295] 745-755, 2001.

[77296] 13312.

[77297] Edlund, A.; Johansson, T.; Leidvik, B.; Hansson, L.: Structure
[77298] of the human kappa-casein gene. Gene 174: 65-69, 1996.
[77299] 13313.

[77300] Sprecher, C. A.; Morgenstern, K. A.; Mathewes, S.; Dahlen, J. R.;
[77301] Schrader, S. K.; Foster, D. C.; Kisiel, W.: Molecular cloning, expression,
[77302] and partial characterization of two novel members of the ovalbumin
[77303] family of serine proteinase inhibitors. J. Biol. Chem. 270: 29854-29861,
[77304] 1995.

[77305] 13314.

[77306] Morahan, G.; Huang, D.; Yu, W.-P.; Cui, L.; DeAizpurua, H.; Pallen,

[77307] C. J.: Localization of the genes encoding the type I diabetes autoantigens,
[77308] protein-tyrosine phosphatases IA2 and IAR. Mammalian Genome 9: 593-594,
[77309] 1998.

[77310] 13315.

[77311] Chevalier, D.; Cauffiez, C.; Bernard, C.; Lo-Guidice, J.-M.; Allorge,
[77312] D.; Fazio, F.; Ferrari, N.; Libersa, C.; Lhermitte, M.; D'Halluin,
[77313] J.-C.; Broly, F.: Characterization of new mutations in the coding
[77314] sequence and 5-prime-untranslated region of the human prostacyclin
[77315] synthase gene (CYP8A1). Hum. Genet. 108: 148-155, 2001.
[77316] 13316.

[77317] Park, W. S.; Lee, J. H.; Shin, M. S.; Park, J. Y.; Kim, H. S.;

[77318] Lee, J. H.; Kim, Y. S.; Lee, S. N.; Xiao, W.; Park, C. H.; Lee, S.
[77319] H.; Yoo, N. J.; Lee, J. Y.: Inactivating mutations of the caspase-10

[77320] gene in gastric cancer. Oncogene 21: 2919-2925, 2002.

[77321] 13317.

[77322] Shin, M. S.; Kim, H. S.; Kang, C. S.; Park, W. S.; Kim, S. Y.;

[77323] Lee, S. N.; Lee, J. H.; Park, J. Y.; Jang, J. J.; Kim, C. W.; Kim,

[77324] S. H.; Lee, J. Y.; Yoo, N. J.; Lee, S. H.: Inactivating muta-
tions

[77325] of CASP10 gene in non-Hodgkin lymphomas. Blood 99:
4094-4099, 2002.

[77326] 13318.

[77327] Vincenz, C.; Dixit, V. M.: Fas-associated death domain
protein

[77328] interleukin-1-beta-converting enzyme 2 (FLICE2), an ICE/
Ced-3 homologue,

[77329] is proximally involved in CD95- and p55-mediated death
signaling. J.

[77330] Biol. Chem. 272: 6578-6583, 1997.

[77331] 13319.

[77332] Wang, J.; Chun, H. J.; Wong, W.; Spencer, D. M.; Lenardo,
M. J.

[77333] : Caspase-10 is an initiator caspase in death receptor sig-
naling. Proc.

[77334] Nat. Acad. Sci. 98: 13884-13888, 2001.

[77335] 13320.

[77336] Wang, J.; Zheng, L.; Lobito, A.; Chan, F. K.; Dale, J.; Sneller,

[77337] M.; Yao, X.; Puck, J. M.; Straus, S. E.; Lenardo, M. J.: Inher-
ited

[77338] human caspase 10 mutations underlie defective lympho-
cyte and dendritic

[77339] cell apoptosis in autoimmune lymphoproliferative syn-
drome type II. Cell 98:

[77340] 47-58, 1999.

[77341] 13321.

[77342] Chun, H. J.; Zheng, L.; Ahmad, M.; Wang, J.; Speirs, C. K.;
Siegel,

[77343] R. M.; Dale, J. K.; Puck, J.; Davis, J.; Hall, C. G.; Skoda-
Smith,

[77344] S.; Atkinson, T. P.; Straus, S. E.; Lenardo, M. J.: Pleiotropic
defects

[77345] in lymphocyte activation caused by caspase-8 mutations
lead to human

[77346] immunodeficiency. Nature 419: 395-399, 2002.

[77347] 13322.

[77348] Grenet, J.; Teitz, T.; Wei, T.; Valentine, V.; Kidd, V. J.: Structure

[77349] and chromosome localization of the human CASP8 gene. Gene 226: 225-232,

[77350] 1999.

[77351] 13323.

[77352] Kischkel, F. C.; Kioschis, P.; Weitz, S.; Poustka, A.; Lichter,

[77353] P.; Krammer, P. H.: Assignment of CASP8 to human chromosome band

[77354] 2q33-q34 and Casp8 to the murine syntenic region on chromosome 1B-proximal

[77355] C by in situ hybridization. Cytogenet. Cell Genet. 82: 95-96, 1998.

[77356] 13324.

[77357] Sanchez, I.; Xu, C.-J.; Juo, P.; Kakizaka, A.; Blenis, J.; Yuan,

[77358] J.: Caspase-8 is required for cell death induced by expanded polyglutamine

[77359] repeats. Neuron 22: 623-633, 1999.

[77360] 13325.

[77361] Varfolomeev, E. E.; Schuchmann, M.; Luria, V.; Chiannilkulchai,

[77362] N.; Beckmann, J. S.; Mett, I. L.; Rebrikov, D.; Brodianski, V. M.;

[77363] Kemper, O. C.; Kollet, O.; Lapidot, T.; Soffer, D.; Sobe, T.; Avraham,

[77364] K. B.; Goncharov, T.; Holtmann, H.; Lonai, P.; Wallach, D.: Targeted

[77365] disruption of the mouse Caspase 8 gene ablates cell death induction

[77366] by the TNF receptors, Fas/Apo1, and DR3 and is lethal prenatally. Immunity 9:

[77367] 267-276, 1998.

[77368] 13326.

[77369] Wang, Y.-K.; Samos, C. H.; Peoples, R.; Perez-Jurado, L. A.; Nusse,

[77370] R.; Francke, U.: A novel human homologue of the Drosophila frizzled

[77371] wnt receptor gene binds wingless protein and is in the Williams syndrome

[77372] deletion at 7q11.23. Hum. Molec. Genet. 6: 465-472, 1997.

[77373] 13327.

[77374] Kalchman, M. A.; Koide, H. B.; McCutcheon, K.; Graham, R.

K.; Nichol,
[77375] K.; Nishiyama, K.; Kazemi-Esfarjani, P.; Lynn, F. C.; Wellington,
[77376] C.; Metzler, M.; Goldberg, Y. P.; Kanazawa, I.; Gietz, R. D.; Hayden,
[77377] M. R.: HIP1, a human homologue of S. cerevisiae Slap2, interacts
[77378] with membrane-associated huntingtin in the brain. Nature Genet. 16:
[77379] 44-53, 1997.

[77380] 13328.

[77381] Wanker, E. E.; Rovira, C.; Scherzinger, E.; Hasenbank, R.; Walter,
[77382] S.; Tait, D.; Colicelli, J.; Lehrach, H.: HIP-I: a huntingtin interacting
[77383] protein isolated by the yeast two-hybrid system. Hum. Molec. Genet. 6:
[77384] 487-495, 1997.

[77385] 13329.

[77386] Cao, Y.; Traer, E.; Zimmerman, G. A.; McIntyre, T. M.; Prescott,
[77387] S. M.: Cloning, expression, and chromosomal localization

of human
[77388] long-chain fatty acid-CoA ligase 4 (FACL4). Genomics 49: 327-330,
[77389] 1998.

[77390] 13330.

[77391] Langkopf, A.; Hammarback, J. A.; Muller, R.; Vallee, R. B.; Garner,
[77392] C. C.: Microtubule-associated proteins 1A and LC2: Two proteins encoded
[77393] in one messenger RNA. J. Biol. Chem. 267: 16561-16566, 1992.
[77394] 13331.

[77395] Sun, X.-H.; Copeland, N. G.; Jenkins, N. A.; Baltimore, D.: Id
[77396] proteins Id1 and Id2 selectively inhibit DNA binding by one class
[77397] of helix-loop-helix proteins. Molec. Cell. Biol. 11: 5603-5611,
[77398] 1991.

[77399] 13332.

[77400] Kordeli, E.; Bennett, V.: Distinct ankyrin isoforms at neu-

ron
[77401] cell bodies and nodes of Ranvier resolved using erythro-
cyte ankyrin-deficient
[77402] mice. J. Cell Biol. 114: 1243-1259, 1991.

[77403] 13333.

[77404] Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu,
M.;
[77405] Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.; Spits,
H.;
[77406] de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Human
thymic
[77407] stromal lymphopoietin preferentially stimulates myeloid
cells. J.
[77408] Immun. 167: 336-343, 2001.

[77409] 13334.

[77410] Nagase, T.; Kikuno, R.; Nakayama, M.; Hirosawa, M.;
Ohara, O.:
[77411] Prediction of the coding sequences of unidentified human
genes. XVIII.
[77412] The complete sequences of 100 new cDNA clones from
brain which code
[77413] for large proteins in vitro. DNA Res. 7: 273-281, 2000.

[77414] 13335.

[77415] Kurochkin, I. V.; Yonemitsu, N.; Funahashi, S.; Nomura, H.: ALEX1,

[77416] a novel human armadillo repeat protein that is expressed differentially

[77417] in normal tissues and carcinomas. Biochem. Biophys. Res. Commun. 280:

[77418] 340-347, 2001.

[77419] 13336.

[77420] Nagase, T.; Ishikawa, K.; Miyajima, N.; Tanaka, A.; Kotani, H.;

[77421] Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified

[77422] human genes. IX. The complete sequences of 100 new cDNA clones from

[77423] brain which can code for large proteins in vitro. DNA Res. 5: 31-39,

[77424] 1998.

[77425] 13337.

[77426] Chuang, T.-H.; Ulevitch, R. J.: Cloning and characterization of

[77427] a sub-family of human Toll-like receptors: hTLR7, hTLR8 and hTLR9. Europ.

[77428] Cytokine Netw. 11: 372-378, 2000.

[77429] 13338.

[77430] Du, X.; Poltorak, A.; Wei, Y.; Beutler, B.: Three novel mam-

[77431] malian toll-like receptors: gene structure, expression, and evolu-

[77432] tion. Europ. Cytokine Netw. 11: 362-371, 2000.

[77433] 13339.

[77434] Kadowaki, N.; Ho, S.; Antonenko, S.; de Waal Malefyt, R.; Kastelein,

[77435] R. A.; Bazan, F.; Liu, Y.-J.: Subsets of human dendritic cell precursors

[77436] express different Toll-like receptors and respond to dif-

[77437] ferent microbial antigens. J. Exp. Med. 194: 863-869, 2001.

[77438] 13340.

[77439] Schwartz, J. M.; Paress, P. S.; Ross, J. M.; Dipillo, F.; Rizek,

[77440] R.: Unstable variant of NADH methemoglobin reductase in Puerto Ricans

[77441] with hereditary methemoglobinemia. J. Clin. Invest. 51: 1594-1601,
[77442] 1972.

[77443] 13341.

[77444] Roy, N.; Mahadevan, M. S.; McLean, M.; Shutler, G.; Yaraghi, Z.;
[77445] Farahani, R.; Baird, S.; Besner-Johnston, A.; Lefebvre, C.; Kang,
[77446] X.; Salih, M.; Aubry, H.; Tamai, K.; Guan, X.; Ioannou, P.; Crawford,
[77447] T. O.; de Jong, P. J.; Surh, L.; Ikeda, J.-E.; Korneluk, R. G.; MacKenzie,
[77448] A.: The gene for neuronal apoptosis inhibitory protein is partially
[77449] deleted in individuals with spinal muscular atrophy. Cell 80: 167-178,
[77450] 1995.

[77451] 13342.

[77452] Barton, D. E.; Foellmer, B. E.; Wood, W. I.; Francke, U.: Chromosome
[77453] mapping of the growth hormone receptor gene in man and mouse. Cytogenet.

[77454] Cell Genet. 50: 137-141, 1989.

[77455] 13343.

[77456] Berg, M. A.; Argente, J.; Chernausek, S.; Gracia, R.; Gue-
vara-Aguirre,

[77457] J.; Hopp, M.; Perez-Jurado, L.; Rosenbloom, A.; Toledo, S.
P. A.;

[77458] Francke, U.: Diverse growth hormone receptor gene muta-
tions in Laron

[77459] syndrome. Am. J. Hum. Genet. 52: 998-1005, 1993.

[77460] 13344.

[77461] Ezekowitz, R. A. B.; Dinauer, M. C.; Jaffe, H. S.; Orkin, S. H.;

[77462] Newburger, P. E.: Partial correction of the phagocyte de-

fect in patients

[77463] with X-linked chronic granulomatous disease by subcuta-

neous interferon

[77464] gamma. New Eng. J. Med. 319: 146-151, 1988.

[77465] 13345.

[77466] Frattini, A.; Orchard, P. J.; Sobacchi, C.; Giliani, S.; Abinun,

[77467] M.; Mattsson, J. P.; Keeling, D. J.; Andersson, A.-K.; Wall-

brandt,

[77468] P.; Zecca, L.; Notarangelo, L. D.; Vezzoni, P.; Villa, A.: De-

fects

[77469] in TCIRG1 subunit of the vacuolar proton pump are re-

sponsible for

[77470] a subset of human autosomal recessive osteopetrosis. Na-

ture Genet. 25:

[77471] 343-346, 2000.

[77472] 13346.

[77473] Afzal, A. R.; Rajab, A.; Fenske, C. D.; Oldridge, M.; Elanko,

N.;

[77474] Ternes-Pereira, E.; Tuysuz, B.; Murday, V. A.; Patton, M. A.; Wilkie,

[77475] A. O. M.; Jeffery, S.: Recessive Robinow syndrome, allelic to dominant

[77476] brachydactyly type B, is caused by mutation of ROR2. Na-

ture Genet. 25:

[77477] 419-422, 2000.

[77478] 13347.

[77479] van Bokhoven, H.; Celli, J.; Kayserili, H.; van Beusekom, E.;

[77480] Balci, S.; Brussel, W.; Skovby, F.; Kerr, B.; Percin, E. F.; Akarsu,

[77481] N.; Brunner, H. G.: Mutation of the gene encoding the ROR2 tyrosine

[77482] kinase causes autosomal recessive Robinow syndrome. Nature Genet. 25:

[77483] 423-426, 2000.

[77484] 13348.

[77485] Sorge, G.; Ruggieri, M.; Lachman, R. S.: Spondyloperiph-

eral dysplasia. Am.

[77486] J. Med. Genet. 59: 139-142, 1995.

[77487] 13349.

[77488] Sybert, V. P.; Byers, P. H.; Hall, J. G.: Variable expression

[77489] in a dominantly inherited skeletal dysplasia with similarities to

[77490] brachydactyly E and spondyloepiphyseal-spondyloperipheral dysplasia. Clin.

[77491] Genet. 15: 160-166, 1979.

[77492] 13350.

[77493] Vanek, J.: Spondyloperipheral dysplasia. J. Med. Genet. 20:

[77494] 117-121, 1983.

[77495] 13351.

[77496] Zabel, B.; Hilbert, K.; Stoss, H.; Superti-Furga, A.; Spranger,

[77497] J.; Winterpacht, A.: A specific collagen type II gene (COL2A1) mutation

[77498] presenting as spondyloperipheral dysplasia. Am. J. Med. Genet. 63:

[77499] 123-128, 1996.

[77500] 13352.

[77501] Adornato, B. T.; O'Brien, J. S.; Lampert, P. W.; Roe, T. F.;

Neustein,
[77502] H. B.: Cerebral spongy degeneration of infancy: a bio-
chemical and
[77503] ultrastructural study of affected twins. Neurology 22:
202-210,
[77504] 1972.

[77505] 13353.

[77506] Aduchi, M.; Aronson, S. M.: Studies on spongy degenera-
tion of
[77507] the central nervous system (van Bogaert-Bertrand type).In:
Aronson,
[77508] S. M.; Volk, B. W.: Inborn Disorders of Sphingolipid
Metabolism.
[77509] Oxford: Pergamon Press (pub.) 1967. Pp. 129-147.

[77510] 13354.

[77511] Banker, B. Q.; Robertson, J. T.; Victor, M.: Spongy degen-
eration
[77512] of the central nervous system in infancy. Neurology 14:
981-1001,
[77513] 1964.

[77514] 13355.

[77515] Banker, B. Q.; Victor, M.: Spongy degeneration of in-
fancy.In:
[77516] Goodman, R. E.; Motulsky, A. G.: Genetic Diseases Among
Ashkenazi
[77517] Jews. New York: Raven Press (pub.) 1979. Pp. 201-216.

[77518] 13356.

[77519] Chou, S. M.; Waisman, H. A.: Spongy degeneration of the
central
[77520] nervous system: case of homocystinuria. Arch. Path. 79:
357-363,
[77521] 1965.

[77522] 13357.

[77523] den Dunnen, J. T.; Grootscholten, P. M.; Bakker, E.; Blon-
den,
[77524] L. A. J.; Ginjaar, H. B.; Wapenaar, M. C.; van Paassen, H. M.
B.;
[77525] van Broeckhoven, C.; Pearson, P. L.; van Ommen, G. J. B.:
Topography
[77526] of the Duchenne muscular dystrophy (DMD) gene: FIGE
and cDNA analysis
[77527] of 194 cases reveals 115 deletions and 13 duplications.
Am. J. Hum.

[77528] Genet. 45: 835-847, 1989.

[77529] 13358.

[77530] Abrams, C. K.; Bennett, M. V. L.; Verselis, V. K.; Bargiello, T.

[77531] A.: Voltage opens unopposed gap junction hemichannels formed by a

[77532] connexin 32 mutant associated with X-linked Charcot-Marie-Tooth disease. Proc.

[77533] Nat. Acad. Sci. 99: 3980-3984, 2002.

[77534] 13359.

[77535] Ainsworth, P. J.; Bolton, C. F.; Murphy, B. C.; Stuart, J. A.;

[77536] Hahn, A. F.: Genotype/phenotype correlation in affected individuals

[77537] of a family with a deletion of the entire coding sequence of the connexin

[77538] 32 gene. Hum. Genet. 103: 242-244, 1998.

[77539] 13360.

[77540] Bahr, M.; Andres, F.; Timmerman, V.; Nelis, M. E.; Van Broeckhoven,

[77541] C.; Dichgans, J.: Central visual, acoustic, and motor path-way involvement

[77542] in a Charcot-Marie-Tooth family with an asn205-to-ser mutation in
[77543] the connexin 32 gene. J. Neurol. Neurosurg. Psychiat. 66: 202-206,
[77544] 1999.

[77545] 13361.

[77546] Bergoffen, J.; Scherer, S. S.; Wang, S.; Oronzi Scott, M.; Bone,
[77547] L. J.; Paul, D. L.; Chen, K.; Lensch, M. W.; Chance, P. F.; Fischbeck,
[77548] K. H.: Connexin mutations in X-linked Charcot-Marie-Tooth disease. Science 262:
[77549] 2039-2042, 1993.

[77550] 13362.

[77551] Bondurand, N.; Girard, M.; Pingault, V.; Lemort, N.; Dubourg, O.;
[77552] Goossens, M.: Human connexin 32, a gap junction protein altered in
[77553] the X-linked form of Charcot-Marie-Tooth disease, is directly regulated
[77554] by the transcription factor SOX10. Hum. Molec. Genet. 10: 2783-2795,

[77555] 2001.

[77556] 13363.

[77557] Bone, L. J.; Dahl, N.; Lensch, M. W.; Chance, P. F.; Kelly, T.;

[77558] Le Guern, E.; Magi, S.; Parry, G.; Shapiro, H.; Wang, S.; Fischbeck,

[77559] K. H.: New connexin32 mutations associated with X-linked Charcot-Marie-Tooth

[77560] disease. Neurology 45: 1863-1866, 1995.

[77561] 13364.

[77562] Corcos, I. A.; Lafreniere, R. G.; Begy, C. R.; Loch-Caruso, R.;

[77563] Willard, H. F.; Glover, T. W.: Refined localization of human connexin

[77564] 32 gene locus, GJB1, to Xq13.1. Genomics 13: 479-480, 1992.

[77565] 13365.

[77566] Fairweather, N.; Bell, C.; Cochrane, S.; Chelly, J.; Wang, S.;

[77567] Mostacciuolo, M. L.; Monaco, A. P.; Haites, N. E.: Mutations in the

[77568] connexin 32 gene in X-linked dominant Charcot-Marie-Tooth disease

[77569] (CMTX1). Hum. Molec. Genet. 3: 29-34, 1994.

[77570] 13366.

[77571] Fishman, G. I.; Eddy, R. L.; Shows, T. B.; Rosenthal, L.; Leinwand,

[77572] L. A.: The human connexin gene family of gap junction proteins: distinct

[77573] chromosomal locations but similar structure. Genomics 10: 250-256,

[77574] 1991.

[77575] 13367.

[77576] Ikegami, T.; Lin, C.; Kato, M.; Itoh, A.; Nonaka, I.; Kurimura,

[77577] M.; Hirayabashi, H.; Shinohara, Y.; Mochizuki, A.; Hayasaka, K.:

[77578] Four novel mutations of the connexin 32 gene in four Japanese families

[77579] with Charcot-Marie-Tooth disease type 1. Am. J. Med. Genet. 80:

[77580] 352-355, 1998.

[77581] 13368.

[77582] Ionasescu, V.; Ionasescu, R.; Searby, C.: Correlation be-

tween

[77583] connexin32 gene mutations and clinical phenotype in X-linked dominant

[77584] Charcot-Marie-Tooth neuropathy. Am. J. Med. Genet. 63: 486-491,

[77585] 1996.

[77586] 13369.

[77587] Ionasescu, V. V.; Searby, C.; Ionasescu, R.; Neuhaus, I. M.; Werner,

[77588] R.: Mutations of the noncoding region of the connexin 32 gene in

[77589] X-linked dominant Charcot-Marie-Tooth neuropathy. Neurology 47:

[77590] 541-544, 1996.

[77591] 13370.

[77592] Janssen, E. A. M.; Kemp, S.; Hensels, G. W.; Sie, O. G.; de Die-Smulders,

[77593] C. E. M.; Hoogendijk, J. E.; de Visser, M.; Bolhuis, P. A.: Connexin32

[77594] gene mutations in X-linked dominant Charcot-Marie-Tooth disease (CMTX1). Hum.

[77595] Genet. 99: 501-505, 1997.

[77596] 13371.

[77597] Kumar, N. M.; Gilula, N. B.: The gap junction communica-
tion channel. Cell 84:
[77598] 381-388, 1996.

[77599] 13372.

[77600] Lafreniere, R. G.; Brown, C. J.; Powers, V. E.; Carrel, L.;
Davies,
[77601] K. E.; Barker, D. F.; Willard, H. F.: Physical mapping of 60
DNA
[77602] markers in the p21.1-q21.3 region of the human X chro-
mosome. Genomics 11:
[77603] 352-363, 1991.

[77604] 13373.

[77605] Latour, P.; Levy, N.; Paret, M.; Chapon, F.; Chazot, G.;
Clavelou,
[77606] P.; Couratier, P.; Dumas, R.; Ollagnon, E.; Pouget, J.;
Setiey, A.;
[77607] Vallat, J. M.; Boucherat, M.; Fontes, M.; Vandenberghe, A.:
Mutations
[77608] in the X-linked form of Charcot-Marie-Tooth disease in
the French

[77609] population. Neurogenetics 1: 117-123, 1997.

[77610] 13374.

[77611] Nelis, E.; Simokovic, S.; Timmerman, V.; Lofgren, A.; Back-
hovens,
[77612] H.; De Jonghe, P.; Martin, J.-J.; Van Broeckhoven, C.: Mu-
tation analysis
[77613] of the connexin 32 (Cx32) gene in Charcot-Marie-Tooth
neuropathy type
[77614] 1: identification of five new mutations. Hum. Mutat. 9:
47-52, 1997.

[77615] 13375.

[77616] Raimondi, E.; Gaudi, S.; Moralli, D.; De Carli, L.; Malcovati,

[77617] M.; Simonic, T.; Tenchini, M. L.: Assignment of the human
connexin
[77618] 32 gene (GJB1) to band Xq13. Cytogenet. Cell Genet. 60:
210-211,
[77619] 1992.

[77620] 13376.

[77621] Eng, B. H.; Guerinot, M. L.; Eide, D.; Saier, M. H., Jr.: Se-
quence
[77622] analyses and phylogenetic characterization of the ZIP

family of metal
[77623] ion transport proteins. J. Membr. Biol. 166: 1-7, 1998.

[77624] 13377.

[77625] Lioumi, M.; Ferguson, C. A.; Sharpe, P. T.; Freeman, T.; Marenholz,
[77626] I.; Mischke, D.; Heizmann, C.; Ragoussis, J.: Isolation and characterization
[77627] of human and mouse ZIRTL, a member of the IRT1 family of transporters,
[77628] mapping within the epidermal differentiation complex. Genomics 62:
[77629] 272-280, 1999.

[77630] 13378.

[77631] Lioumi, M.; Olavesen, M. G.; Nizetic, D.; Ragoussis, J.: High-resolution
[77632] YAC fragmentation map of 1q21. Genomics 49: 200-208, 1998.

[77633] 13379.

[77634] Leiper, J. M,; Santa Maria, J.; Chubb, A.; MacAllister, R. J.;

[77635] Charles, I. G.; Whitley, G. S.; Vallance, P.: Identification of two

[77636] human dimethylarginine dimethylaminohydrolases with distinct tissue
[77637] distributions and homology with microbial arginine deiminases. Biochem.
[77638] J. 343: 209-214, 1999.

[77639] 13380.

[77640] Tran, C. T. L.; Fox, M. F.; Vallance, P.; Leiper, J. M.: Chromosomal
[77641] localization, gene structure, and expression pattern of DDAH1: comparison
[77642] with DDAH2 and implications for evolutionary origins. Genomics 68:
[77643] 101-105, 2000.

[77644] 13381.

[77645] Vanhalst, K.; Kools, P.; Eynde, E. V.; van Roy, F.: The human
[77646] and murine protocadherin-beta one-exon gene families show high evolutionary
[77647] conservation, despite the difference in gene number. FEBS Lett. 495:
[77648] 120-125, 2001.

[77649] 13382.

[77650] Wu, Q.; Maniatis, T.: A striking organization of a large family

[77651] of human neural cadherin like cell adhesion genes. Cell 97: 779-790,

[77652] 1999.

[77653] 13383.

[77654] Marie-Cardine, A.; Bruyns, E.; Eckerskorn, C.; Kirchgess-ner, H.;

[77655] Meuer, S. C.; Schraven, B.: Molecular cloning of SKAP55, a novel

[77656] protein that associates with the protein tyrosine kinase p59-fyn in

[77657] human T-lymphocytes. J. Biol. Chem. 272: 16077-16080, 1997.

[77658] 13384.

[77659] Katayama, H.; Ota, T.; Morita, K.; Terada, Y.; Suzuki, F.; Katoh,

[77660] O.; Tatsuka, M.: Human AIM-1: cDNA cloning and reduced expression

[77661] during endomitosis in megakaryocyte-lineage cells. Gene 224: 1-7,

[77662] 1998.

[77663] 13385.

[77664] Kimura, M.; Matsuda, Y.; Yoshioka, T.; Sumi, N.; Okano,
Y.: Identification

[77665] and characterization of STK12/Aik2: a human gene re-
lated to aurora

[77666] of Drosophila and yeast IPL1. Cytogenet. Cell Genet. 82:
147-152,

[77667] 1998.

[77668] 13386.

[77669] Tatsuka, M.; Katayama, H.; Ota, T.; Tanaka, T.; Odashima,
S.; Suzuki,

[77670] F.; Terada, Y.: Multinuclearity and increased ploidy caused
by overexpression

[77671] of the aurora- and Ipl1-like midbody-associated protein
mitotic kinase

[77672] in human cancer cells. Cancer Res. 58: 4811-4816, 1998.

[77673] 13387.

[77674] Chen, C. M.; Kraut, N.; Groudine, M.; Weintraub, H.: I-mf,
a novel

[77675] myogenic repressor, interacts with members of the MyoD

family. Cell 86:
[77676] 731-741, 1996.

[77677] 13388.

[77678] Kraut, N.: The gene encoding I-mf (Mdfi) maps to human chromosome
[77679] 6p21 and mouse chromosome 17. Mammalian Genome 8: 618-619, 1997.

[77680] 13389.

[77681] Kraut, N.; Snider, L.; Chen, C.-M. A.; Tapscott, S. J.; Grou-
dine,
[77682] M.: Requirement of the mouse I-mfa gene for placental development
[77683] and skeletal patterning. EMBO J. 17: 6276-6288, 1998.

[77684] 13390.

[77685] Inaba, S.; Okochi, K.; Yae, Y.; Niklasson, F.; de Verder, C.-H.
[77686] : Serological studies of an SLE-associated antigen-anti-
body system
[77687] discovered as a precipitation reaction in agarose gel: the Hakata
[77688] antigen-antibody system. Fukuoka Igaku Zasshi 81:

284-291, 1990.

[77689] 13391.

[77690] Sugimoto, R.; Yae, Y.; Akaiwa, M.; Kitajima, S.; Shibata, Y.; Sato,

[77691] H.; Hirata, J.; Okochi, K.; Izuhara, K.; Hamasaki, N.: Cloning and

[77692] characterization of the Hakata antigen, a member of the ficolin/opsonin

[77693] p35 lectin family. J. Biol. Chem. 273: 20721-20727, 1998.

[77694] 13392.

[77695] Llano, E.; Pendas, A. M.; Freije, J. P.; Nakano, A.; Knauper, V.;

[77696] Murphy, G.; Lopez-Otin, C.: Identification and characteri-

[77697] zation of human MT5-MMP, a new membrane-bound activator of progelatinase A overexpressed

[77698] in brain tumors. Cancer Res. 59: 2570-2576, 1999.

[77699] 13393.

[77700] Guttinger, M.; Sutti, F.; Panigada, M.; Porcellini, S.; Merati,

[77701] B.; Mariani, M.; Teesalu, T.; Consalez, G. G.; Grassi, F.: Epithelial

[77702] V-like antigen (EVA), a novel member of the immunoglob-
ulin superfamily,
[77703] expressed in embryonic epithelia with a potential role as
homotypic
[77704] adhesion molecule in thymus histogenesis. J. Cell Biol.
141: 1061-1071,
[77705] 1998.

[77706] 13394.

[77707] Bahler, M.: Myosins on the move to signal transduction.
Curr.
[77708] Opin. Cell Biol. 8: 18-22, 1996.

[77709] 13395.

[77710] Gorman, S. W.; Haider, N. B.; Grieshammer, U.; Swiderski,
R. E.;
[77711] Kim E; Welch, J. W.; Searby, C.; Leng, S.; Carmi, R.;
Sheffield, V.
[77712] C.; Duhl, D. M.: The cloning and developmental expres-
sion of unconventional
[77713] myosin IXA (MYO9A) a gene in the Bardet-Biedl syndrome
(BBS4) region
[77714] at chromosome 15q22-q23. Genomics 59: 150-160,
1999.

[77715] 13396.

[77716] Dunham, I.; Shimizu, N.; Roe, B. A.; Chissoe, S.; Hunt, A. R.;

[77717] Collins, J. E.; Bruskiewich, R.; Beare, D. M.; Clamp, M.; Smink, L.

[77718] J.; Ainscough, R.; Almeida, J. P.; and 205 others: The DNA sequence

[77719] of human chromosome 22. Nature 402: 489-495, 1999.

[77720] 13397.

[77721] Kimura, T.; Ivell, R.; Rust, W.; Mizumoto, Y.; Ogita, K.; Kusui,

[77722] C.; Matsumura, Y.; Azuma, C.; Murata, Y.: Molecular cloning of a

[77723] human MafF homologue, which specifically binds to the oxytocin receptor

[77724] gene in term myometrium. Biochem. Biophys. Res. Com-mun. 264: 86-92,

[77725] 1999.

[77726] 13398.

[77727] Hiki, K.; D'Andrea, R. J.; Furze, J.; Crawford, J.; Woollatt, E.;

[77728] Sutherland, G. R.; Vadas, M. A.; Gamble, J. R.: Cloning,

characterization,

[77729] and chromosomal location of a novel human K(+)-Cl(-) cotransporter. J.

[77730] Biol. Chem. 274: 10661-10667, 1999.

[77731] 13399.

[77732] Mount, D. B.: Personal Communication. Nashville, Tenn. 3/28/2000.

[77733] 13400.

[77734] Mount, D. B.; Mercado, A.; Song, L.; Xu, J.; George, A. L., Jr.;

[77735] Delpire, E.; Gamba, G.: Cloning and characterization of KCC3 and

[77736] KCC4, new members of the cation-chloride cotransporter gene family. J.

[77737] Biol. Chem. 274: 16355-16362, 1999.

[77738] 13401.

[77739] Herzing, L. B. K.; Kim, S.-J.; Cook, E. H., Jr.; Ledbetter, D.

[77740] H.: The human aminophospholipid-transporting ATPase gene ATP10C maps

[77741] adjacent to UBE3A and exhibits similar imprinted expres- sion. Am.

[77742] J. Hum. Genet. 68: 1501-1505, 2001.

[77743] 13402.

[77744] Dode, C.; Le Du, N.; Cuisset, L.; Letourneur, F.; Berthelot, J.-M.;

[77745] Vaudour, G.; Meyrier, A.; Watts, R. A.; Scott, D. G. I.; Nicholls,

[77746] A.; Granel, B.; Frances, C.; Garcier, F.; Edery, P.; Boulinguez, S.;

[77747] Domergues, J.-P.; Delpech, M.; Grateau, G.: New mutations of CIAS1

[77748] that are responsible for Muckle-Wells syndrome and familial cold urticaria:

[77749] a novel mutation underlies both syndromes. Am. J. Hum. Genet. 70:

[77750] 1498-1506, 2002.

[77751] 13403.

[77752] Feldmann, J.; Prieur, A.-M.; Quartier, P.; Berquin, P.; Certain,

[77753] S.; Cortis, E.; Teillac-Hamel, D.; Fischer, A.; de Saint Basile, G.

[77754] : Chronic infantile neurological cutaneous and articular syndrome

[77755] is caused by mutations in CIAS1, a gene highly expressed in polymorphonuclear

[77756] cells and chondrocytes. Am. J. Hum. Genet. 71: 198-203, 2002.

[77757] 13404.

[77758] Greeve, I.; Hermans-Borgmeyer, I.; Brellinger, C.; Kasper, D.;

[77759] Gomez-Isla, T.; Behl, C.; Levkau, B.; Nitsch, R. M.: The human DIMINUTO/DWARF1

[77760] homolog seladin-1 confers resistance to Alzheimer's disease-associated

[77761] neurodegeneration and oxidative stress. J. Neurosci. 20: 7345-7352,

[77762] 2000.

[77763] 13405.

[77764] Sarkar, D.; Imai, T.; Kambe, F.; Shibata, A.; Ohmori, S.; Siddiq,

[77765] A.; Hayasaka, S.; Funahashi, H.; Seo, H.: The human homolog of Diminuto/Dwarf1

[77766] gene (hDiminuto): a novel ACTH-responsive gene overexpressed in benign

[77767] cortisol-producing adrenocortical adenomas. J. Clin. En-

docr. Metab. 86:
[77768] 5130-5137, 2001.

[77769] 13406.

[77770] Hattori, M.; Fujiyama, A.; Taylor, T. D.; Watanabe, H.;
Yada, T.;
[77771] Park, H.-S.; Toyoda, A.; Ishii, K.; Totoki, Y.; Choi, D.-K.;
Groner,
[77772] Y.; Soeda, E.; and 52 others: The DNA sequence of human
chromosome
[77773] 21. Nature 405: 311-319, 2000. Note: Erratum: Nature:
407: 110 only,
[77774] 2000.

[77775] 13407.

[77776] Wilcox, E. R.; Burton, Q. L.; Naz, S.; Riazuddin, S.; Smith,
T.
[77777] N.; Ploplis, B.; Belyatseva, I.; Ben-Yosef, T.; Liburd, N. A.;
Morell,
[77778] R. J.; Kachar, B.; Wu, D. K.; Griffith, A. J.; Riazuddin, S.;
Friedman,
[77779] T. B.: Mutations in the gene encoding tight junction
claudin-14 cause
[77780] recessive deafness DFNB29. Cell 104: 165-172, 2001.

[77781] 13408.

[77782] Volkert, M. R.; Elliott, N. A.; Housman, D. E.: Functional genomics

[77783] reveals a family of eukaryotic oxidation protection genes. Proc.

[77784] Nat. Acad. Sci. 97: 14530-14535, 2000.

[77785] 13409.

[77786] Dunlevy, J. R.; Berryhill, B. L.; Vergnes, J.-P.; SundarRaj, N.;

[77787] Hassell, J. R.: Cloning, chromosomal localization, and characterization

[77788] of cDNA from a novel gene, SH3BP4, expressed by human corneal fibroblasts. Genomics 62:

[77789] 519-524, 1999.

[77790] 13410.

[77791] Wong, W. T.; Schumacher, C.; Salcini, A. E.; Romano, A.; Castagnino,

[77792] P.; Pelicci, P. G.; DiFiore, P. P.: A protein-binding domain, EH,

[77793] identified in the receptor tyrosine kinase substrate Eps15 and conserved

[77794] in evolution. Proc. Nat. Acad. Sci. 92: 9530-9534, 1995.

[77795] 13411.

[77796] Yamadori, T.; Baba, Y.; Matsushita, M.; Hashimoto, S.; Kurosaki,

[77797] M.; Kurosaki, T.; Kishimoto, T.; Tsukada, S.: Bruton's tyrosine kinase

[77798] activity is negatively regulated by Sab, the Btk-SH3 domain-binding

[77799] protein. Proc. Nat. Acad. Sci. 96: 6341-6346, 1999.

[77800] 13412.

[77801] Engqvist-Goldstein, A. E. Y.; Kessels, M. M.; Chopra, V. S.; Hayden,

[77802] M. R.; Drubin, D. G.: An actin-binding protein of the Sla2/Huntingtin

[77803] interacting protein 1 family is a novel component of clathrin-coated

[77804] pits and vesicles. J. Cell Biol. 147: 1503-1518, 1999.

[77805] 13413.

[77806] Baba, Y.; Matsushita, M.; Matsuda, Y.; Inazawa, J.; Yamadori, T.;

[77807] Hashimoto, S.; Kishimoto, T.; Tsukada, S.: Assignment of SH3BP5/Sh3bp5

[77808] encoding Sab, an SH3 domain-binding protein which preferentially associates

[77809] with Bruton's tyrosine kinase, to human chromosome 1q43 and mouse

[77810] chromosome 14B by in situ hybridization. Cytogenet. Cell Genet. 87:

[77811] 221-222, 1999.

[77812] 13414.

[77813] Matsushita, M.; Yamadori, T.; Kato, S.; Takemoto, Y.; In-azawa,

[77814] J.; Baba, Y.; Hashimoto, S.; Sekine, S.; Arai, S.; Kunikata, T.; Kurimoto,

[77815] M.; Kishimoto, T.; Tsukada, S.: Identification and charac-terization

[77816] of a novel SH3-domain binding protein, Sab, which pref-erentially associates

[77817] with Bruton's tyrosine kinase (Btk). Biochem. Biophys. Res. Commun. 245:

[77818] 337-343, 1998.

[77819] 13415.

[77820] Itoh, T.; Koshiba, S.; Kigawa, T.; Kikuchi, A.; Yokoyama, S.; Takenawa,

[77821] T.: Role of the ENTH domain in phosphatidylinositol-4,5-biphosphate
[77822] binding and endocytosis. Science 291: 1047-1051, 2001.

[77823] 13416.

[77824] Seki, N.; Muramatsu, M.; Sugano, S.; Suzuki, Y.; Nakagawara, A.;
[77825] Ohhira, M.; Hayashi, A.; Hori, T.; Saito, T.: Cloning, expression
[77826] analysis, and chromosomal localization of HIP1R, an isolog of huntingtin
[77827] interacting protein (HIP1). J. Hum. Genet. 43: 268-271, 1998.
[77828] 13417.

[77829] Nash, S. R.; Giros, B.; Kingsmore, S. F.; Kim, K. M.; El-Mestikawy,
[77830] S.; Dong, Q.; Fumagalli, F.; Seldin, M. F.; Caron, M. G.: Cloning,
[77831] gene structure and genomic localization of an orphan transporter from
[77832] mouse kidney with six alternatively-spliced isoforms. Receptors Channels 6:
[77833] 113-128, 1998.

[77834] 13418.

[77835] Scott, A. F.: Personal Communication. Baltimore, Md. 2/5/2001.

[77836] 13419.

[77837] Meguro, M.; Kashiwagi, A.; Mitsuya, K.; Nakao, M.; Kondo, I.; Saitoh,

[77838] S.; Oshimura, M.: A novel maternally expressed gene, ATP10C, encodes

[77839] a putative aminophospholipid translocase associated with Angelman

[77840] syndrome. Nature Genet. 28: 19-20, 2001.

[77841] 13420.

[77842] Nakakura, E. K.; Watkins, D. N.; Schuebel, K. E.; Sriuran-pong,

[77843] V.; Borges, M. W.; Nelkin, B. D.; Ball, D. W.: Mammalian Scratch:

[77844] a neural-specific Snail family transcriptional repressor. Proc. Nat.

[77845] Acad. Sci. 98: 4010-4015, 2001.

[77846] 13421.

[77847] Scott, A. F.: Personal Communication. Baltimore, Md.

6/21/2001.

[77848] 13422.

[77849] Hoatlin, M. E.; Zhi, Y.; Ball, H.; Silvery, K.; Melnick, A.; Stone,

[77850] S.; Arai, S.; Hawe, N.; Owen, G.; Zelent, A.; Licht, J. D.: A novel

[77851] BTB/POZ transcriptional repressor protein interacts with the Fanconi

[77852] anemia group C protein and PLZF. Blood 94: 3737-3747, 1999.

[77853] 13423.

[77854] Lin, W.; Lai, C.-H.; Tang, C.-J. C.; Huang, C.-J.; Tang, T. K.

[77855] : Identification and gene structure of a novel human PLZF-related

[77856] transcription factor gene, TZFP. Biochem. Biophys. Res. Commun. 264:

[77857] 789-795, 1999.

[77858] 13424.

[77859] Yokoyama-Kobayashi, M.; Yamaguchi, T.; Sekine, S.; Kato, S.: Selection

[77860] of cDNAs encoding putative type II membrane proteins on

the cell surface

[77861] from a human full-length cDNA bank. Gene 228: 161-167, 1999.

[77862] 13425.

[77863] Shiraishi, N.; Natsume, A.; Togayachi, A.; Endo, T.; Akashima,

[77864] T.; Yamada, Y.; Imai, N.; Nakagawa, S.; Koizumi, S.; Sekine, S.; Narimatsu,

[77865] H.; Sasaki, K.: Identification and characterization of three novel

[77866] beta-1,3-N-acetylglucosaminyltransferases structurally related to

[77867] the beta-1,3-galactosyltransferase family. J. Biol. Chem. 276: 3498-3507,

[77868] 2001.

[77869] 13426.

[77870] Yokoyama-Kobayashi, M.; Sugano, S.; Kato, T.; Kato, S.: A signal

[77871] sequence detection system using secreted protease activity as an indicator. Gen

[77872] e 163: 193-196, 1995.

[77873] 13427.

[77874] Brown, M. T.; Andrade, J.; Radhakrishna, H.; Donaldson, J. G.;

[77875] Cooper, J. A.; Randazzo, P. A.: ASAP1, a phospholipid-dependent Arf

[77876] GTPase-activating protein that associates with and is phosphorylated

[77877] by Src. Molec. Cell. Biol. 18: 7038-7051, 1998.

[77878] 13428.

[77879] Olivier, E.; Soury, E.; Risler, J. L.; Smith, F.; Schneider, K.;

[77880] Lochner, K.; Jouzeau, J. Y.; Fey, G. H.; Salier, J. P.: A novel set

[77881] of hepatic mRNAs preferentially expressed during an acute inflammation

[77882] in rat represents mostly intracellular proteins. Genomics 57: 352-364,

[77883] 1999.

[77884] 13429.

[77885] Olivier, E.; Soury, E.; Ruminy, P.; Husson, A.; Parmentier, F.;

[77886] Daveau, M.; Salier, J.-P.: Fetuin-B, a second member of the fetuin

[77887] family in mammals. Biochem. J. 350: 589-597, 2000.

[77888] 13430.

[77889] Hampe, J.; Grebe, J.; Nikolaus, S.; Solberg, C.; Croucher, P.
J.
[77890] P.; Mascheretti, S.; Jahnsen, J.; Moum, B.; Klump, B.;
Krawczak, M.;
[77891] Mirza, M. M.; Foelsch, U. R.; Vatn, M.; Schreiber, S.: Asso-
ciation
[77892] of NOD2 (CARD 15) genotype with clinical course of
Crohn's disease:
[77893] a cohort study. Lancet 359: 1661-1665, 2002.

[77894] 13431.

[77895] Hugot, J.-P.; Chamaillard, M.; Zouali, H.; Lesage, S.;
Cezard,
[77896] J.-P.; Belaiche, J.; Almer, S.; Tysk, C.; O'Morain, C. A.; Gas-
sull,
[77897] M.; Binder, V.; Finkel, Y.; and 8 others: Association of
NOD2 leucine-rich
[77898] repeat variants with susceptibility to Crohn's disease. Na-
ture 411:
[77899] 599-603, 2001.

[77900] 13432.

[77901] Murillo, L.; Crusius, J. B. A.; van Bodegraven, A. A.; Al-
izadeh,
[77902] B. Z.; Pena, A. S.: CARD15 gene and the classification of
Crohn's
[77903] disease. Immunogenetics 54: 59-61, 2002.

[77904] 13433.

[77905] Ogura, Y.; Bonen, D. K.; Inohara, N.; Nicolae, D. L.; Chen,
F.
[77906] F.; Ramos, R.; Britton, H.; Moran, T.; Karaliuskas, R.;
Duerr, R.
[77907] H.; Achkar, J.-P.; Brant, S. R.; Bayless, T. M.; Kirschner, B.
S.;
[77908] Hanauer, S. B.; Nunez, G.; Cho, J. H.: A frameshift muta-
tion in Nod2
[77909] associated with susceptibility to Crohn's disease. Nature
411: 603-606,
[77910] 2001.

[77911] 13434.

[77912] Ogura, Y.; Inohara, N.; Benito, A.; Chen, F. F.; Yamaoka, S.;
Nunez,

[77913] G.: Nod2, a Nod1/Apaf-1 family member that is restricted to monocytes
[77914] and activates NF-kappa-B. J. Biol. Chem. 276: 4812-4818, 2001.
[77915] 13435.

[77916] Lesage, S.; Zouali, H.; Cezard, J.-P.; EPWG-IBD Group; Colombel,
[77917] J.-F.; EPIMAD Group; Belaiche, J.; GETAID Group; Almer, S.; Tysk,
[77918] C.; O'Morain, C.; Gassull, M.; Binder, V.; Finkel, Y.; Modigliani,
[77919] R.; Gower-Rousseau, C.; Macry, J.; Merlin, F.; Chamaillard, M.; Jannot,
[77920] A.-S.; Thomas, G.; Hugot, J.-P.: CARD15/NOD2 mutational analysis
[77921] and genotype-phenotype correlation in 612 patients with inflammatory
[77922] bowel disease. Am. J. Hum. Genet. 70: 845-857, 2002.

[77923] 13436.

[77924] van Heel, D. A.; McGovern, D. P. B.; Cardon, L. R.; Dechairo, B.
[77925] M.; Lench, N. J.; Carey, A. H.; Jewell, D. P.: Fine mapping of

the
[77926] IBD1 locus did not identify Crohn disease-associated NOD2 variants:
[77927] implications for complex disease genetics. Am. J. Med. Genet. 111:
[77928] 253-259, 2002.

[77929] 13437.

[77930] Vermeire, S.; Wild, G.; Kocher, K.; Cousineau, J.; Dufresne, L.;
[77931] Bitton, A.; Langelier, D.; Pare, P.; Lapointe, G.; Cohen, A.; Daly,
[77932] M. J.; Rioux, J. D.: CARD15 genetic variation in a Quebec population:
[77933] prevalence, genotype-phenotype relationship, and haplo-type structure. Am.
[77934] J. Hum. Genet. 71: 74-83, 2002.

[77935] 13438.

[77936] Yamazaki, K.; Takazoe, M.; Tanaka, T.; Kazumori, T.; Nakamura,
[77937] Y.: Absence of mutation in the NOD2/CARD15 gene among 483 Japanese
[77938] patients with Crohn's disease. J. Hum. Genet. 47:

469-472, 2002.

[77939] 13439.

[77940] Routhier, E. L.; Burn, T. C.; Abbaszade, I.; Summers, M.; Albright,

[77941] C. F.; Prendergast, G. C.: Human BIN3 complements the F-actin localization

[77942] defects caused by loss of Hob3p, the fission yeast homolog of Rvs161p. J.

[77943] Biol. Chem. 276: 21670-21677, 2001.

[77944] 13440.

[77945] Fields, R. R.; Zhou, G.; Huang, D.; Davis, J. R.; Moller, C.; Jacobson,

[77946] S. G.; Kimberling, W. J.; Sumegi, J.: Usher syndrome type III: revised

[77947] genomic structure of the USH3 gene and identification of novel mutations. Am.

[77948] J. Hum. Genet. 71: 607-617, 2002.

[77949] 13441.

[77950] Li, B.; Oestreich, S.; de Lange, T.: Identification of human Rap1:

[77951] implications for telomere evolution. Cell 101: 471-483,

2000.

[77952] 13442.

[77953] Lieb, J. D.; Liu, X.; Botstein, D.; Brown, P. O.: Promoter-specific

[77954] binding of Rap1 revealed by genome-wide maps of protein-DNA association. Nature

[77955] Genet. 28: 327-334, 2001.

[77956] 13443.

[77957] Honore, B.; Leffers, H.; Madsen, P.; Rasmussen, H. H.; Vandekerckhove,

[77958] J.; Celis, J. E.: Molecular cloning and expression of a transformation-sensitive

[77959] human protein containing the TPR motif and sharing identity to the

[77960] stress-inducible yeast protein STI1. J. Biol. Chem. 267: 8485-8491,

[77961] 1992.

[77962] 13444.

[77963] Scheufler, C.; Brinker, A.; Bourenkov, G.; Pegoraro, S.; Moroder,

[77964] L.; Bartunik, H.; Hartl, F. U.; Moarefi, I.: Structure of TPR

domain-peptide
[77965] complexes: critical elements in the assembly of the Hsp70-Hsp90 multichaperone
[77966] machine. Cell 101: 199-210, 2000.

[77967] 13445.

[77968] Slentz-Kesler, K.; Moore, J. T.; Lombard, M.; Zhang, J.; Hollingsworth,
[77969] R.; Weiner, M. P.: Identification of the human Mnk2 gene (MKNK2)
[77970] through protein interaction with estrogen receptor beta. Genomics 69:
[77971] 63-71, 2000.

[77972] 13446.

[77973] Li, Y.; He, X.; Schembri-King, J.; Jakes, S.; Hayashi, J.: Cloning
[77974] and characterization of human Lnk, an adaptor protein with pleckstrin
[77975] homology and Src homology 2 domains that can inhibit T cell activation. J.
[77976] Immun. 164: 5199-5206, 2000.

[77977] 13447.

[77978] Takaki, S.; Sauer, K.; Iritani, B. M.; Chien, S.; Ebihara, Y.;

[77979] Tsuji, K.; Takatsu, K.; Perlmutter, R. M.: Control of B cell production

[77980] by the adaptor protein Lnk: definition of a conserved family of signal-modulating

[77981] proteins. Immunity 13: 599-609, 2000.

[77982] 13448.

[77983] Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.; Mercer,

[77984] W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.: Wip1, a

[77985] novel human protein phosphatase that is induced in response to ionizing

[77986] radiation in a p53-dependent manner. Proc. Nat. Acad. Sci. 94: 6048-6053,

[77987] 1997.

[77988] 13449.

[77989] Li, J.; Yang, Y.; Peng, Y.; Austin, R. J.; van Eyndhoven, W. G.;

[77990] Nguyen, K. C. Q.; Gabriele, T.; McCurrach, M. E.; Marks, J. R.; Hoey,

[77991] T.; Lowe, S. W.; Powers, S.: Oncogenic properties of PPM1D located

[77992] within a breast cancer amplification epicenter at 17q23. Nature Genet. 31:

[77993] 133-134, 2002.

[77994] 13450.

[77995] Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;

[77996] 1. Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.; Mercer,

[77997] W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.: Wip1, a

[77998] novel human protein phosphatase that is induced in re-sponse to ionizing

[77999] radiation in a p53-dependent manner. Proc. Nat. Acad. Sci. 94: 6048-6053,

[78000] 1997.

[78001] 13451.

[78002] Takaesu, G.; Kishida, S.; Hiyama, A.; Yamaguchi, K.; Shibuya, H.;

[78003] Irie, K.; Ninomiya-Tsuji, J.; Matsumoto, K.: TAB2, a novel adaptor

[78004] protein, mediates activation of TAK1 MAPKKK by linking TAK1 to TRAF6

[78005] in the IL-1 signal transduction pathway. Molec. Cell 5: 649-658,

[78006] 2000.

[78007] 13452.

[78008] Stover, C. M.; Schwaeble, W. J.; Lynch, N. J.; Thiel, S.; Spe-

icher,

[78009] M. R.: Assignment of the gene encoding mannan-binding lectin-associated

[78010] serine protease 2 (MASP2) to human chromosome 1p36.3-p36.2 by in situ

[78011] hybridization and somatic cell hybrid analysis. Cytogenet. Cell Genet. 84:

[78012] 148-149, 1999.

[78013] 13453.

[78014] Stover, C. M.; Thiel, S.; Thelen, M.; Lynch, N. J.; Vorup-

Jensen,

[78015] T.; Jensenius, J. C.; Schwaeble, W. J.: Two constituents of the initiation

[78016] complex of the mannan-binding lectin activation pathway of complement

[78017] are encoded by a single structural gene. J. Immun. 162: 3481-3490,

[78018] 1999.

[78019] 13454.

[78020] Takahashi, M.; Endo, Y.; Fujita, T.; Matsushita, M.: A truncated

[78021] form of mannose-binding lectin-associated serine protease (MASP)-2

[78022] expressed by alternative polyadenylation is a component of the lectin

[78023] complement pathway. Int. Immun. 11: 859-863, 1999.

[78024] 13455.

[78025] Thiel, S.; Vorup-Jensen, T.; Stover, C. M.; Schwaeble, W.; Laursen,

[78026] S. B.; Poulsen, K.; Willis, A. C.; Eggleton, P.; Hansen, S.; Holmskov,

[78027] U.; Reid, K. B. M.; Jensenius, J. C.: A second serine protease associated

[78028] with mannan-binding lectin that activates complement. Nature 386:

[78029] 506-510, 1997.

[78030] 13456.

[78031] Sundin, O. H.; Yang, J. M.; Li, Y.; Zhu, D.; Hurd, J. N.; Mitchell,

[78032] T. N.; Silva, E. D.; Maumenee, I. H.: Genetic basis of total colourblindness

[78033] among the Pingelapese islanders. Nature Genet. 25: 289-293, 2000.

[78034] 13457.

[78035] Konakova, M.; Huynh, D. P.; Yong, W.; Pulst, S. M.: Cellular distribution

[78036] of torsin A and torsin B in normal human brain. Arch. Neurol. 58:

[78037] 921-927, 2001.

[78038] 13458.

[78039] Conklin, D.; Holderman, S.; Whitmore, T. E.; Maurer, M.; Feldhaus,

[78040] A. L.: Molecular cloning, chromosome mapping and characterization

[78041] of UBQLN3 a testis-specific gene that contains an ubiquitin-like domain. Gene 249:

[78042] 91-98, 2000.

[78043] 13459.

[78044] Drewes, T.; Senkel, S.; Holewa, B.; Ryffel, G. U.: Human hepatocyte

[78045] nuclear factor 4 isoforms are encoded by distinct and differentially

[78046] expressed genes. Molec. Cell. Biol. 16: 925-931, 1996.

[78047] 13460.

[78048] Taraviras, S.; Mantamadiotis, T.; Dong-Si, T.; Mincheva, A.; Lichter,

[78049] P.; Drewes, T.; Ryffel, G. U.; Monaghan, A. P.; Schutz, G.: Primary

[78050] structure, chromosomal mapping, expression and transcriptional activity

[78051] of murine hepatocyte nuclear factor 4-gamma. Biochim. Biophys. Acta 1490:

[78052] 21-32, 2000.

[78053] 13461.

[78054] Seroussi, E.; Kedra, D.; Pan, H.-Q.; Peyrad, M.; Schwartz, C.;

[78055] Scambler, P.; Donnai, D.; Roe, B. A.; Dumanski, J. P.: Duplications

[78056] on human chromosome 22 reveal a novel Ret finger protein-like gene

[78057] family with sense and endogenous antisense transcripts. Genome Res. 9:

[78058] 803-814, 1999.

[78059] 13462.

[78060] Blencowe, B. J.; Issner, R.; Nickerson, J. A.; Sharp, P. A.: A

[78061] coactivator of pre-mRNA splicing. Genes Dev. 12: 996-1009, 1998.

[78062] 13463.

[78063] Leung, J. C.; Klein, C.; Friedman, J.; Vieregge, P.; Jacobs, H.;

[78064] Doheny, D.; Kamm, C.; DeLeon, D.; Pramstaller, P. P.; Penney, J. B.;

[78065] Eisengart, M.; Jankovic, J.; Gasser, T.; Bressman, S. B.; Corey, D.

[78066] P.; Kramer, P.; Brin, M. F.; Ozelius, L. J.; Breakefield, X. O.:

[78067] Novel mutation in the TOR1A (DYT1) gene in atypical, early onset dystonia

[78068] and polymorphisms in dystonia and early onset parkinsonism. Neurogenetics 3:

[78069] 133-143, 2001.

[78070] 13464.

[78071] Aihara, Y.; Mashima, H.; Onda, H.; Hisano, S.; Kasuya, H.; Hori,

[78072] T.; Yamada, S.; Tomura, H.; Yamada, Y.; Inoue, I.; Kojima, I.; Takeda,

[78073] J.: Molecular cloning of a novel brain-type Na(+)-dependent inorganic

[78074] phosphate cotransporter. J. Neurochem. 74: 2622-2625, 2000.

[78075] 13465.

[78076] Bellocchio, E. E.; Reimer, R. J.; Fremeau, R. T., Jr.; Edwards,

[78077] R. H.: Uptake of glutamate into synaptic vesicles by an in-organic

[78078] phosphate transporter. Science 289: 957-960, 2000.

[78079] 13466.

[78080] Ni, B.; Du, Y.; Wu, X.; DeHoff, B. S.; Rosteck, P. R., Jr.; Paul,

[78081] S. M.: Molecular cloning, expression, and chromosomal localization

[78082] of a human brain-specific Na(+)-dependent inorganic phosphate cotransporter. J.

[78083] Neurochem. 66: 2227-2238, 1996.

[78084] 13467.

[78085] Takamori, S.; Rhee, J. S.; Rosenmund, C.; Jahn, R.: Identifi-
cation
[78086] of a vesicular glutamate transporter that defines a gluta-
matergic
[78087] phenotype in neurons. Nature 407: 189-194, 2000.

[78088] 13468.

[78089] Blackwood, D.: P300, a state and a trait marker in
schizophrenia. Lancet 355:
[78090] 771-772, 2000.

[78091] 13469.

[78092] Blackwood, D. H. R.; Fordyce, A.; Walker, M. T.; St. Clair,
D.
[78093] M.; Porteous, D. J.; Muir, W. J.: Schizophrenia and affective
disorders--cosegretation
[78094] with a translocation at chromosome 1q42 that directly
disrupts brain-expressed
[78095] genes: clinical and P300 findings in a family. Am. J. Hum.
Genet. 69:
[78096] 428-433, 2001.

[78097] 13470.

[78098] Ekelund, J.; Hovatta, I.; Parker, A.; Paunio, T.; Varilo, T.; Martin,

[78099] R.; Suhonen, J.; Ellonen, P.; Chan, G.; Sinsheimer, J. S.; So-bel,

[78100] E.; Juvonen, H.; Arajarvi, R.; Partonen, T.; Suvisaari, J.; Lonnqvist,

[78101] J.; Meyer, J.; Peltonen, L.: Chromosome 1 loci in Finnish schizophrenia

[78102] families. Hum. Molec. Genet. 10: 1611-1617, 2001.

[78103] 13471.

[78104] Millar, J. K.; Wilson-Annan, J. C.; Anderson, S.; Christie, S.;

[78105] Taylor, M. S.; Semple, C. A. M.; Devon, R. S.; St. Clair, D. M.; Muir,

[78106] W. J.; Blackwood, D. H. R.; Porteous, D. J.: Disruption of two novel

[78107] genes by a translocation co-segregating with schizophre-nia. Hum.

[78108] Molec. Genet. 9: 1415-1423, 2000.

[78109] 13472.

[78110] St. Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.;

Walker,
[78111] M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association within a family
[78112] of a balanced autosomal translocation with major mental illness. Lancet 336:
[78113] 13-16, 1990.

[78114] 13473.

[78115] Bulfone, A.; Menguzzato, E.; Broccoli, V.; Marchitiello, A.; Gattuso,
[78116] C.; Mariani, M.; Consalez, G. G.; Martinez, S.; Ballabio, A.; Banfi,
[78117] S.: Barhl1, a gene belonging to a new subfamily of mammalian homeobox
[78118] genes, is expressed in migrating neurons of the CNS. Hum. Molec.
[78119] Genet. 9: 1443-1452, 2000.

[78120] 13474.

[78121] Alessi, D. R.; Deak, M.; Casamayor, A.; Caudwell, F. B.; Morrice,
[78122] N.; Norman, D. G.; Gaffney, P.; Reese, C. B.; MacDougall, C. N.; Harbison,
[78123] D.; Ashworth, A.; Bownes, M.:

3-phosphoinositide-dependent protein
[78124] kinase-1 (PDK1): structural and functional homology with the Drosophila
[78125] DSTPK61 kinase. Curr. Biol. 7: 776-789, 1997.

[78126] 13475.

[78127] Boucher, C. A.; King, S. K.; Carey, N.; Krahe, R.; Winch-
ester,
[78128] C. L.; Rahman, S.; Creavin, T.; Meghji, P.; Bailey, M. E. S.; Chartier,
[78129] F. L.; Brown, S. D.; Siciliano, M. J.; Johnson, K. J.: A novel homeodomain-encoding
[78130] gene is associated with a large CpG island interrupted by the myotonic
[78131] dystrophy unstable (CTG)n repeat. Hum. Molec. Genet. 4: 1919-1925,
[78132] 1995.

[78133] 13476.

[78134] Carango, P.; Noble, J. E.; Marks, H. G.; Funanage, V. L.: Absence
[78135] of myotonic dystrophy protein kinase (DMPK) mRNA as a result of a
[78136] triplet repeat expansion in myotonic dystrophy. Genomics

18: 304-348,

[78137] 1993.

[78138] 13477.

[78139] Fu, Y.-H.; Friedman, D. L.; Richards, S.; Pearlman, J. A.; Gibbs,

[78140] R. A.; Pizzuti, A.; Ashizawa, T.; Perryman, M. B.; Scarlato, G.; Fenwick,

[78141] R. G., Jr.; Caskey, C. T.: Decreased expression of myotonin-protein

[78142] kinase messenger RNA and protein in adult form of myotonic dystrophy. Science 260:

[78143] 235-238, 1993.

[78144] 13478.

[78145] Groenen, P. J. T. A.; Wansink, D. G.; Coerwinkel, M.; van den Broek,

[78146] W.; Jansen, G.; Wieringa, B.: Constitutive and regulated modes of

[78147] splicing produce six major myotonic dystrophy protein kinase (DMPK)

[78148] isoforms with distinct properties. Hum. Molec. Genet. 9: 605-616,

[78149] 2000.

[78150] 13479.

[78151] Jansen, G.; Bartolomei, M.; Kalscheuer, V.; Merkx, G.; Wormskamp,

[78152] N.; Mariman, E.; Smeets, D.; Ropers, H.-H.; Wieringa, B.: No imprinting

[78153] involved in the expression of DM-kinase mRNAs in mouse and human tissues. Hum.

[78154] Molec. Genet. 2: 1221-1227, 1993.

[78155] 13480.

[78156] Mahadevan, M. S.; Amemiya, C.; Jansen, G.; Sabourin, L.; Baird,

[78157] S.; Neville, C. E.; Wormskamp, N.; Segers, B.; Batzer, M.; Lamerdin,

[78158] J.; de Jong, P.; Wieringa, B.; Korneluk, R. G.: Structure and genomic

[78159] sequence of the myotonic dystrophy (DM kinase) gene. Hum. Molec.

[78160] Genet. 2: 299-304, 1993.

[78161] 13481.

[78162] Roses, A. D.; Appel, S. H.: Muscle membrane protein kinase in

[78163] myotonic muscular dystrophy. Nature 250: 245-247, 1974.

[78164] 13482.

[78165] Shaw, D. J.; McCurrach, M.; Rundle, S. A.; Harley, H. G.; Crow,

[78166] S. R.; Sohn, R.; Thirion, J.-P.; Hamshere, M. G.; Buckler, A. J.;

[78167] Harper, P. S.; Housman, D. E.; Brook, J. D.: Genomic orga-nization

[78168] and transcriptional units at the myotonic dystrophy locus. Genomics 18:

[78169] 673-679, 1993.

[78170] 13483.

[78171] Jansen, G.; Mahadevan, M.; Amemiya, C.; Wormskamp, N.; Segers,

[78172] B.; Hendriks, W.; O'Hoy, K.; Baird, S.; Sabourin, L.; Lennon, G.;

[78173] Jap, P. L.; Iles, D.; Coerwinkel, M.; Hofker, M.; Carrano, A. V.;

[78174] de Jong, P. J.; Korneluk, R. G.; Wieringa, B.: Characteriza-tion of

[78175] the myotonic dystrophy region predicts multiple protein

isoform-encoding
[78176] mRNAs. Nature Genet. 1: 261-266, 1992.

[78177] 13484.

[78178] Seznec, H.; Agbulut, O.; Sergeant, N.; Savouret, C.; Gh-
estem,
[78179] A.; Tabti, N.; Willer, J.-C.; Ourth, L.; Duros, C.; Brisson, E.;
Fouquet,
[78180] C.; Butler-Browne, G.; Delacourte, A.; Junien, C.; Gourdon,
G.: Mice
[78181] transgenic for the human myotonic dystrophy region with
expanded CTG
[78182] repeats display muscular and brain abnormalities. Hum.
Molec. Genet. 10:
[78183] 2717-2726, 2001.

[78184] 13485.

[78185] van den Broek, W. J. A. A.; Nelen, M. R.; Wansink, D. G.;
Coerwinkel,
[78186] M. M.; te Riele, H.; Groenen, P. J. T. A.; Wieringa, B.: So-
matic
[78187] expansion behaviour of the (CTG)n repeat in myotonic
dystrophy knock-in
[78188] mice is differentially affected by Msh3 and Msh6 mis-

match-repair proteins. Hum.
[78189] Molec. Genet. 11: 191-198, 2002.

[78190] 13486.

[78191] van der Ven, P. F. M.; Jansen, G.; van Kuppevelt, T. H. M. S.

[78192] M.; Perryman, M. B.; Lupa, M.; Dunne, P. W.; ter Laak, H. J.; Jap,
[78193] P. H. K.; Veerkamp, J. H.; Epstein, H. F.; Wieringa, B.: Myotonic
[78194] dystrophy kinase is a component of neuromuscular junctions. Hum.
[78195] Molec. Genet. 2: 1889-1894, 1993.

[78196] 13487.

[78197] Wang, J.; Pegoraro, E.; Menegazzo, E.; Gennarelli, M.; Hoop, R.
[78198] C.; Angelini, C.; Hoffman, E. P.: Myotonic dystrophy: evidence for
[78199] a possible dominant-negative RNA mutation. Hum. Molec. Genet. 4:
[78200] 599-606, 1995.

[78201] 13488.

[78202] Wang, Y.-H.; Amirhaeri, S.; Kang, S.; Wells, R. D.; Griffith,

[78203] J. D.: Preferential nucleosome assembly at DNA triplet repeats from
[78204] the myotonic dystrophy gene. Science 265: 669-671, 1994.
[78205] 13489.

[78206] Wang, Y.-H.; Griffith, J.: Expanded CTG triplet blocks from the
[78207] myotonic dystrophy gene create the strongest known natural nucleosome
[78208] positioning elements. Genomics 25: 570-573, 1995.

[78209] 13490.

[78210] Wochner, R. D.; Drews, G.; Strober, W.; Waldmann, T. A.: Accelerated
[78211] breakdown of immunoglobulin G (IgG) in myotonic dystrophy: a hereditary
[78212] error of immunoglobulin catabolism. J. Clin. Invest. 45: 321-329,
[78213] 1966.

[78214] 13491.

[78215] Wong, L.-J. C.; Ashizawa, T.; Monckton, D. G.; Caskey, C. T.;

[78216] Richards, C. S.: Somatic heterogeneity of the CTG repeat in myotonic
[78217] dystrophy is age and size dependent. Am. J. Hum. Genet. 56: 114-122,
[78218] 1995.

[78219] 13492.

[78220] Zhang, Y.; Monckton, D. G.; Siciliano, M. J.; Connor, T. H.; Meistrich,
[78221] M. L.: Age and insertion site dependence of repeat number instability
[78222] of a human DM1 transgene in individual mouse sperm. Hum. Molec. Genet. 11:
[78223] 791-798, 2002.

[78224] 13493.

[78225] Bertin, J.; Nir, W.-J.; Fischer, C. M.; Tayber, O. V.; Errada,
[78226] P. R.; Grant, J. R.; Keilty, J. J.; Gosselin, M. L.; Robison, K. E.;
[78227] Wong, G. H. W.; Glucksmann, M. A.; DiStefano, P. S.: Human CARD4
[78228] protein is a novel CED-4/Apaf-1 cell death family member that activates

[78229] NF-kappa-B. J. Biol. Chem. 274: 12955-12958, 1999.

[78230] 13494.

[78231] Cohen, D. E.; Green, R. M.; Wu, M. K.; Beier, D. R.: Cloning,

[78232] tissue-specific expression, gene structure and chromoso-
mal localization

[78233] of human phosphatidylcholine transfer protein. Biochim.
Biophys.

[78234] Acta 1447: 265-270, 1999.

[78235] 13495.

[78236] van Helvoort, A.; de Brouwer, A.; Ottenhoff, R.; Brouwers,
J. F.

[78237] H. M.; Wijnholds, J.; Beijnen, J. H.; Rijneveld, A.; van der
Valk,

[78238] M. A.; Majoor, D.; Voorhout, W.; Wirtz, K. W. A.; Elferink,
R. P.

[78239] J. O.; Borst, P.: Mice without phosphatidylcholine transfer
protein

[78240] have no defects in the secretion of phosphatidylcholine
into bile

[78241] or into lung airspaces. Proc. Nat. Acad. Sci. 96:
11501-11506, 1999.

[78242] 13496.

[78243] de Rooij, J.; Zwartkruis, F. J. T.; Verheijen, M. H. G.; Cool,

[78244] R. H.; Nijman, S. M. B.; Wittinghofer, A.; Bos, J. L.: Epac is a

[78245] Rap1 guanine-nucleotide-exchange factor directly activated by cyclic

[78246] AMP. Nature 396: 474-477, 1998.

[78247] 13497.

[78248] Kawasaki, H.; Springett, G. M.; Mochizuki, N.; Toki, S.; Nakaya,

[78249] M.; Matsuda, M.; Housman, D. E.; Graybiel, A. M.: A family of cAMP-binding

[78250] proteins that directly activate Rap1. Science 282: 2275-2279, 1998.

[78251] 13498.

[78252] Olsen, S. R.; Uhler, M. D.: Inhibition of protein kinase-A by

[78253] overexpression of the cloned human protein kinase inhibitor. Molec.

[78254] Endocr. 5: 1246-1256, 1991.

[78255] 13499.

[78256] Hahn, Y.; Lee, J.; Seong, C.; Yoon, J.; Chung, J. H.: Struc-

tural

[78257] analysis of phylogenetically conserved J domain protein gene. Biochim.

[78258] Biophys. Acta 1447: 325-333, 1999.

[78259] 13500.

[78260] Eberhart, C. G.; Maines, J. Z.; Wasserman, S. A.: Meiotic cell

[78261] cycle requirement for a fly homologue of human Deleted in Azoospermia. Nature 381:

[78262] 783-785, 1996.

[78263] 13501.

[78264] Rugglu, M.; Speed, R.; Taggart, M.; McKay, S. J.; Ki-lanowski, F.;

[78265] Saunders, P.; Derin, J.; Cooke, H. J.: The mouse Dazla gene encodes

[78266] a cytoplasmic protein essential for gametogenesis. Nature 389: 73-77,

[78267] 1997.

[78268] 13502.

[78269] Shan, Z.; Hirschmann, P.; Seebacher, T.; Edelmann, A.; Jauch, A.;

[78270] Morell, J.; Urbitsch, P.; Vogt, P. H.: A SPGY copy homologous to
[78271] the mouse gene Dazla and the Drosophila gene boule is autosomal and
[78272] expressed only in the human male gonad. Hum. Molec. Genet. 5: 2005-2011,
[78273] 1996.

[78274] 13503.

[78275] Slee, R.; Grimes, B.; Speed, R. M.; Taggart, M.; Maguire, S. M.;
[78276] Ross, A.; McGill, N. I.; Saunders, P. T. K.; Cooke, H. J.: A human
[78277] DAZ transgene confers partial rescue of the mouse Daz1 null phenotype. Proc.
[78278] Nat. Acad. Sci. 96: 8040-8045, 1999.

[78279] 13504.

[78280] Yen, P. H.; Chai, N. N.; Salido, E. C.: The human autosomal gene
[78281] DAZLA: testis specificity and a candidate for male infertility. Hum.
[78282] Molec. Genet. 5: 2013-2017, 1996.

[78283] 13505.

[78284] Akiyama, T. E.; Sakai, S.; Lambert, G.; Nicol, C. J.; Mat-
susue,
[78285] K.; Pimprale, S.; Lee, Y.-H.; Ricote, M.; Glass, C. K.;
Brewer, H.
[78286] B., Jr.; Gonzalez, F. J.: Conditional disruption of the perox-
isome
[78287] proliferator-activated receptor gamma gene in mice re-
sults in lowered
[78288] expression of ABCA1, ABCG1, and apoE in macrophages
and reduced cholesterol
[78289] efflux. Molec. Cell. Biol. 22: 2607-2619, 2002.

[78290] 13506.

[78291] Ardlie, K. G.; Lunetta, K. L.; Seielstad, M.: Testing for pop-
ulation
[78292] subdivision and association in four case-control studies.
Am. J.
[78293] Hum. Genet. 71: 304-311, 2002.

[78294] 13507.

[78295] Barak, Y.; Nelson, M. C.; Ong, E. S.; Jones, Y. Z.; Ruiz-
Lozano,

[78296] P.; Chien, K. R.; Koder, A.; Evans, R. M.: PPAR-gamma is required
[78297] for placental, cardiac, and adipose tissue development. Molec. Cell 4:
[78298] 585-595, 1999.

[78299] 13508.

[78300] Barroso, I.; Gurnell, M.; Crowley, V. E. F.; Agostini, M.; Schwabel,
[78301] J. W.; Soos, M. A.; Masien, G. L.; Williams, T. D. M.; Lewis, H.;
[78302] Schafer, A. J.; Chatterjee, V. K. K.; O'Rahilly, S.: Dominant negative
[78303] mutations in human PPAR-gamma associated with severe insulin resistance,
[78304] diabetes mellitus and hypertension. Nature 402: 880-883, 1999.
[78305] 13509.

[78306] Beamer, B. A.; Yen, C.-J.; Andersen, R. E.; Muller, D.; Elahi,
[78307] D.; Cheskin, L. J.; Andres, R.; Roth, J.; Shuldiner, A. R.: As-sociation
[78308] of the pro12ala variant in the peroxisome proliferator-acti-vated receptor-gamma-2

[78309] gene with obesity in two Caucasian populations. Diabetes 47: 1806-1808,

[78310] 1998.

[78311] 13510.

[78312] Beamer, B. A.; Negri, C.; Yen, C.-J.; Gavrilova, O.; Rumberger,

[78313] J. M.; Durcan, M. J.; Yarnall, D. P.; Hawkins, A. L.; Griffin, C.

[78314] A.; Burns, D. K.; Roth, J.; Reitman, M.; Shuldiner, A. R.: Chromosomal

[78315] localization and partial genomic structure of the human peroxisome

[78316] proliferator activated receptor-gamma (hPPAR-gamma) gene. Biochem.

[78317] Biophys. Res. Commun. 233: 756-759, 1997.

[78318] 13511.

[78319] Chawla, A.; Boisvert, W. A.; Lee, C.-H.; Laffitte, B. A.; Barak,

[78320] Y.; Joseph, S. B.; Liao, D.; Nagy, L.; Edwards, P. A.; Curtiss, L.

[78321] K.; Evans, R. M.; Tontonoz, P.: A PPAR-gamma-LXR-ABCA1 pathway in

[78322] macrophages is involved in cholesterol efflux and atherogenesis. Molec.

[78323] Cell 7: 161-171, 2001.

[78324] 13512.

[78325] Deeb, S. S.; Fajas, L.; Nemoto, M.; Pihlajamaki, J.; Mykkanen,

[78326] L.; Kuusisto, J.; Laakso, M.; Fujimoto, W.; Auwerx, J.: A pro12ala

[78327] substitution in PPAR-gamma-2 associated with decreased receptor activity,

[78328] lower body mass index and improved insulin sensitivity. Nature Genet. 20:

[78329] 284-287, 1998.

[78330] 13513.

[78331] Elbrecht, A.; Chen, Y.; Cullinan, C. A.; Hayes, N.; Leibowitz,

[78332] M. D.; Moller, D. E.; Berger, J.: Molecular cloning, expression and

[78333] characterization of human peroxisome proliferator activated receptors

[78334] gamma-1 and gamma-2. Biochem. Biophys. Res. Commun. 224: 431-437,

[78335] 1996.

[78336] 13514.

[78337] Fajas, L.; Auboeuf, D.; Raspe, E.; Schoonjans, K.; Lefebvre, A.

[78338] M.; Saladin, R.; Najib, J.; Laville, M.; Fruchart, J.-C.; Deeb, S.;

[78339] Vidal-Puig, A.; Flier, J.; Briggs, M. R.; Staels, B.; Vidal, H.; Auwerx,

[78340] J.: The organization, promoter analysis, and expression of the human

[78341] PPAR-gamma gene. J. Biol. Chem. 272: 18779-18789, 1997.

[78342] 13515.

[78343] Gampe, R. T., Jr.; Montana, V. G.; Lambert, M. H.; Miller, A.

[78344] B.; Bledsoe, R. K.; Milburn, M. V.; Kliewer, S. A.; Willson, T. M.;

[78345] Xu, H. E.: Asymmetry in the PPAR-gamma/RXR-alpha crystal structure

[78346] reveals the molecular basis of heterodimerization among nuclear receptors. Molec.

[78347] Cell 5: 545-555, 2000.

[78348] 13516.

[78349] Koths, K.; Taylor, E.; Halenbeck, R.; Casipit, C.; Wang, A.: Cloning

[78350] and characterization of a human Mac-2-binding protein, a new member

[78351] of the superfamily defined by the macrophage scavenger receptor cysteine-rich

[78352] domain. J. Biol. Chem. 268: 14245-14249, 1993.

[78353] 13517.

[78354] Ullrich, A.; Sures, I.; D'Egidio, M.; Jallal, B.; Powell, T. J.;

[78355] Herbst, R.; Dreps, A.; Azam, M.; Rubinstein, M.; Natoli, C.; Shawver,

[78356] L. K.; Schlessinger, J.; Iacobelli, S.: The secreted tumor-associated

[78357] antigen 90K is a potent immune stimulator. J. Biol. Chem. 269: 18401-18407,

[78358] 1994.

[78359] 13518.

[78360] Chakraborti, A.; Lippman, D. L.; Loh, H. H.; Kozak, C. A.; Lee,

[78361] N. M.: Genetic mapping of opioid binding protein gene(s)

to mouse
[78362] chromosome 9. Mammalian Genome 4: 179-182, 1993.

[78363] 13519.

[78364] Cho, T. M.; Hasegawa, J.-I.; Ge, B.-l.; Loh, H. H.: Purification
[78365] to apparent homogeneity of a mu-type opioid receptor from rat brain. Proc.
[78366] Nat. Acad. Sci. 83: 4138-4142, 1986.

[78367] 13520.

[78368] Shark, K. B.; Lee, N. M.: Cloning, sequencing and localization
[78369] to chromosome 11 of a cDNA encoding a human opioid-binding cell adhesion
[78370] molecule (OBCAM). Gene 155: 213-217, 1995.

[78371] 13521.

[78372] Burmeister, M.; Meyer, G. E.: The trefoil gene maps to mouse chromosome
[78373] 17. Mammalian Genome 8: 223-224, 1997.

[78374] 13522.

[78375] Chinery, R.; Williamson, J.; Poulsom, R.: The gene encod-

ing human
[78376] intestinal trefoil factor (TFF3) is located on chromosome 21q22.3
[78377] clustered with other members of the trefoil peptide family. Genomics 32:
[78378] 281-284, 1996.

[78379] 13523.

[78380] Mashimo , H.; Wu, D.-C.; Podolsky, D. K.; Fishman, M. C.: Impaired
[78381] defense of intestinal mucosa in mice lacking intestinal trefoil factor. Science 274:
[78382] 262-265, 1996.

[78383] 13524.

[78384] Podolsky, D. K.; Lynch-Devaney, K.; Stow, J. L.; Oates, P.; Murgue,
[78385] B.; DeBeaumont, M.; Sands, B. E.; Mahida, Y. R.: Identification of
[78386] human intestinal trefoil factor: goblet cell-specific expression of
[78387] a peptide targeted for apical secretion. J. Biol. Chem. 268: 6694-6702,
[78388] 1993.

[78389] 13525.

[78390] Probst, J. C.; Zetzsche, T.; Weber, M.; Theilemann, P.; Skutella,

[78391] T.; Landgraf, R.; Jirikowski, G. F.: Human intestinal trefoil factor

[78392] is expressed in human hypothalamus and pituitary: evidence for a novel

[78393] neuropeptide. FASEB J. 10: 1518-1523, 1996.

[78394] 13526.

[78395] Schmitt, H.; Wundrack, I.; Beck, S.; Gott, P.; Welter, C.; Shizuya,

[78396] H.; Simon, M. I.; Blin, N.: A third P-domain peptide gene (TFF3),

[78397] human intestinal trefoil factor, maps to 21q22.3. Cytogenet. Cell

[78398] Genet. 72: 299-302, 1996.

[78399] 13527.

[78400] Taupin, D.; Wu, D.-C.; Jeon, W.-K.; Devaney, K.; Wang, T. C.; Podolsky,

[78401] D. K.: The trefoil gene family are coordinately expressed immediate-early

[78402] genes: EGF receptor- and MAP kinase-dependent inter-regulation. J.

[78403] Clin. Invest. 103: R31-R38, 1999.

[78404] 13528.

[78405] Thim, L.: A new family of growth factor-like peptides: 'tre-foil'

[78406] disulphide loop structures as a common feature in breast cancer associated

[78407] peptide (pS2), pancreatic spasmolytic polypeptide (PSP), and frog

[78408] skin peptides (spasmolysins). FEBS Lett. 250: 85-90, 1989.

[78409] 13529.

[78410] Chen, Y.; Struhl, G.: Dual roles for Patched in sequestering and

[78411] transducing Hedgehog. Cell 87: 553-563, 1996.

[78412] 13530.

[78413] McIntire, J. J.; Umetsu, S. E.; Akbari, O.; Potter, M.; Kuchroo,

[78414] V. K.; Barsh, G. S.; Freeman, G. J.; Umetsu, D. T.; DeKruyff, R. H.

[78415] : Identification of Tapr (an airway hyperreactivity regulatory locus)
[78416] and the linked Tim gene family. Nature Immun. 2: 1109-1116, 2001.
[78417] 13531.

[78418] Alvarez, R. A.; Ghalayini, A. J.; Xu, P.; Hardcastle, A.; Bhattacharya,
[78419] S.; Rao, P. N.; Pettenati, M. J.; Anderson, R. E.; Baehr, W.: cDNA
[78420] sequence and gene locus of the human retinal phosphoinositide-specific
[78421] phospholipase-C-beta-4 (PLCB4). Genomics 29: 53-61, 1995.
[78422] 13532.

[78423] Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu, G.: Expression
[78424] of the p48 xeroderma pigmentosum gene is p53-dependent and is involved
[78425] in global genomic repair. Proc. Nat. Acad. Sci. 96: 424-428, 1999.
[78426] 13533.

[78427] Tahir, S. A.; Yang, G.; Ebara, S.; Timme, T. L.; Satoh, T.; Li,

[78428] L.; Goltsov, A.; Ittmann, M.; Morrisett, J. D.; Thompson, T. C.:

[78429] Secreted caveolin-1 stimulates cell survival/clonal growth and contributes

[78430] to metastasis in androgen-insensitive prostate cancer. Cancer Res. 61:

[78431] 3882-3885, 2001.

[78432] 13534.

[78433] England, S. K.; Uebele, V. N.; Kodali, J.; Bennett, P. B.; Tamkun,

[78434] M. M.: A novel K+ channel beta-subunit (hKv-beta-1.3) is produced

[78435] via alternative mRNA splicing. J. Biol. Chem. 270: 28531-28534,

[78436] 1995.

[78437] 13535.

[78438] Kratz, C. P.; Emerling, B. M.; Bonifas, J.; Wang, W.; Green, E.

[78439] D.; Le Beau, M. M.; Shannon, K. M.: Genomic structure of the PIK3CG

[78440] gene on chromosome band 7q22 and evaluation as a candidate myeloid

[78441] tumor suppressor. Blood 99: 372-374, 2002.

[78442] 13536.

[78443] Duggan, D. J.; Manchester, D.; Stears, K. P.; Mathews, D. J.; Hart,

[78444] C.; Hoffman, E. P.: Mutations in the delta-sarcoglycan gene are a

[78445] rare cause of autosomal recessive limb-girdle muscular dystrophy (LGMD2). Neurogenetics 1:

[78446] 49-58, 1997.

[78447] 13537.

[78448] Jung, D.; Duclos, F.; Apostol, B.; Straub, V.; Lee, J. C.; Allamand,

[78449] V.; Venzke, D. P.; Sunada, Y.; Moomaw, C. R.; Leveille, C. J.; Slaughter,

[78450] C. A.; Crawford, T. O.; McPherson, J. D.; Campbell, K. P.: Characterization

[78451] of delta-sarcoglycan, a novel component of the oligomeric sarcoglycan

[78452] complex involved in limb-girdle muscular dystrophy. J. Biol. Chem. 271:

[78453] 32321-32329, 1996.

[78454] 13538.

[78455] Kawada, T.; Nakazawa, M.; Nakauchi, S.; Yamazaki, K.; Shimamoto,

[78456] R.; Urabe, M.; Nakata, J.; Hemmi, C.; Masui, F.; Nakajima, T.; Suzuki,

[78457] J.-I.; Monahan, J.; Sato, H.; Masaki, T.; Ozawa, K.; Toyo-oka, T.

[78458] : Rescue of hereditary form of dilated cardiomyopathy by rAAV-mediated

[78459] somatic gene therapy: amelioration of morphological findings, sarcolemmal

[78460] permeability, cardiac performances, and the prognosis of TO-2 hamsters. Proc.

[78461] Nat. Acad. Sci. 99: 901-906, 2002.

[78462] 13539.

[78463] Nigro, V.; Okazaki, Y.; Belsito, A.; Piluso, G.; Matsuda, Y.; Politano,

[78464] L.; Nigro, G.; Ventura, C.; Abbondanza, C.; Molinari, A. M.; Acampora,

[78465] D.; Nishimura, M.; Hayashizaki, Y.; Puca, G. A.: Identifica-tion of

[78466] the Syrian hamster cardiomyopathy gene. Hum. Molec.

Genet. 6: 601-607,
[78467] 1997.

[78468] 13540.

[78469] Nigro, V.; Piluso, G.; Belsito, A.; Politano, L.; Puca, A. A.;

[78470] Papparella, S.; Rossi, E.; Vlglietto, G.; Esposito, M. G.; Ab-
bondanza,
[78471] C.; Medici, N.; Molinari, A. M.; Nigro, G.; Puca, G. A.: Iden-
tification
[78472] of a novel sarcoglycan gene at 5q33 encoding a sarcolem-
mal 35 kDa
[78473] glycoprotein. Hum. Molec. Genet. 5: 1179-1186, 1996.

[78474] 13541.

[78475] Okazaki, Y.; Okuizumi, H.; Ohsumi, T.; Nomura, O.;
Takada, S.;
[78476] Kamiya, M.; Sasaki, N.; Matsuda, Y.; Nishimura, M.;
Tagaya, O.; Muramatsu,
[78477] M.; Hayashizaki, Y.: A genetic linkage map of the Syrian
hamster
[78478] and localization of cardiomyopathy locus on chromosome
9qa2.1-b1 using

[78479] RLGS spot-mapping. Nature Genet. 13: 87-90, 1996.

[78480] 13542.

[78481] Ahmad, W.; Li, S.; Chen, H.; Tuck-Muller, C. M.; Pittler, S. J.;

[78482] Aronson, N. N., Jr.: Lysosomal chitobiase (CTB) and the G-protein

[78483] gamma-5 subunit (GNG5) genes co-localize to human chromosome 1p22. Cytogenet.

[78484] Cell Genet. 71: 44-46, 1995.

[78485] 13543.

[78486] Fisher, K. J.; Aronson, N. N.: Characterization of the cDNA and

[78487] genomic sequence of a G protein gamma subunit (gamma-5). Molec. Cell

[78488] Biol. 12: 1585-1591, 1992.

[78489] 13544.

[78490] Fiaschi, T.; Marzella, R.; Veggi, D.; Marzocchini, R.; Raugei,

[78491] G.; Rocchi, M.; Ramponi, G.: Assignment of the human erythrocyte

[78492] acylphosphatase gene (ACYP1) to chromosome band

14q24.3. Cytogenet.

[78493] Cell Genet. 81: 235-236, 1998.

[78494] 13545.

[78495] Fiaschi, T.; Raugei, G.; Marzocchini, R.; Chiarugi, P.; Cirri,

[78496] P.; Ramponi, G.: Cloning and expression of the cDNA cod-
ing for the

[78497] erythrocyte isoenzyme of human acylphosphatase. FEBS
Lett. 367:

[78498] 145-148, 1995.

[78499] 13546.

[78500] Goldowitz, D.; Smeyne, R. J.: Tune into the weaver chan-
nel. Nature

[78501] Genet. 11: 107-109, 1995.

[78502] 13547.

[78503] Hess, E. J.: Identification of the weaver mouse mutation:
the

[78504] end of the beginning. Neuron 16: 1073-1076, 1996.

[78505] 13548.

[78506] Lane, P. W.: New mutation: Weaver, wv. Mouse News Letter
32-33,

[78507] 1964.

[78508] 13549.

[78509] Lesage, F.; Duprat, F.; Fink, M.; Guillemare, E.; Coppola, T.;

[78510] Lazdunski, M.; Hugnot, J.-P.: Cloning provides evidence for a family

[78511] of inward rectifier and G-protein coupled K(+) channels in the brain. FEBS

[78512] Lett. 353: 37-42, 1994.

[78513] 13550.

[78514] Rakic, P.; Sidman, R. L.: Sequence of developmental ab-normalities

[78515] leading to granule cell deficit in cerebellar cortex of weaver mutant

[78516] mice. J. Comp. Neurol. 152: 103-132, 1973.

[78517] 13551.

[78518] Sakura, H.; Bond, C.; Warren-Perry, M.; Horsley, S.; Kear-ney, L.;

[78519] Tucker, S.; Adelman, J.; Turner, R.; Ashcroft, F. M.: Char-acterization

[78520] and variation of a human inwardly-rectifying K-channel

gene (KCNJ6):
[78521] a putative ATP-sensitive K-channel subunit. FEBS Lett.
367: 193-197,
[78522] 1995.

[78523] 13552.

[78524] Tsaur, M.-L.; Menzel, S.; Lai, F.-P.; Espinosa, R., III; Con-
cannon,
[78525] P.; Spielman, R. S.; Hanis, C. L.; Cox, N. J.; Le Beau, M. M.;
German,
[78526] M. S.; Jan, L. Y.; Bell, G. I.; Stoffel, M.: Isolation of a cDNA

[78527] clone encoding a K(ATP) channel-like protein expressed in
insulin-secreting
[78528] cells, localization of the human gene to chromosome
band 21q22.1 and
[78529] linkage studies with NIDDM. Diabetes 44: 592-596, 1995.

[78530] 13553.

[78531] Yasuda, K.; Sakura, H.; Mori, Y.; Iwamoto, K.; Shimokawa,
K.;
[78532] Kadowaki, H.; Hagura, R.; Akanuma, Y.; Adelman, J. P.;
Yazaki, Y.;
[78533] Ashcroft, F. M.; Kadowaki, T.: No evidence for mutations

in a putative
[78534] subunit of the beta-cell ATP-sensitive potassium channel (K-ATP channel)
[78535] in Japanese NIDDM patients. Biochem. Biophys. Res. Com-
mun. 211:
[78536] 1036-1040, 1995.

[78537] 13554.

[78538] Efiok, B. J. S.; Chiorini, J. A.; Safer, B.: A key transcription

[78539] factor for eukaryotic initiation factor-2-alpha is strongly homologous
[78540] to developmental transcription factors and may link metabolic genes
[78541] to cellular growth and development. J. Biol. Chem. 269: 18921-18930,
[78542] 1994.

[78543] 13555.

[78544] Gopalakrishnan, L.; Scarpulla, R. C.: Structure, expression, and
[78545] chromosomal assignment of the human gene encoding nuclear respiratory
[78546] factor 1. J. Biol. Chem. 270: 18019-18025, 1995.

[78547] 13556.

[78548] Spelbrink, J. N.; Van den Bogert, C.: The pre-mRNA of nu-
clear
[78549] respiratory factor 1, a regulator of mitochondrial biogene-
sis, is
[78550] alternatively spliced in human tissues and cell lines. Hum.
Molec.
[78551] Genet. 4: 1591-1596, 1995.

[78552] 13557.

[78553] Tiranti, V.; Rossi, E.; Rocchi, M.; DiDonato, S.; Zuffardi, O.;

[78554] Zeviani, M.: The gene (NFE2L1) for human NRF-1, an acti-
vator involved
[78555] in nuclear-mitochondrial interactions, maps to 7q32. Ge-
nomics 27:
[78556] 555-557, 1995.

[78557] 13558.

[78558] Ohshiro, T.; Yagami, T.; Zhang, C.; Matsuzaki, F.: Role of
cortical
[78559] tumour-suppressor proteins in asymmetric division of
Drosophila neuroblast. Nature 408:
[78560] 593-596, 2000.

[78561] 13559.

[78562] Peng, C.-Y.; Manning, L.; Albertson, R.; Doe, C. Q.: The tumour-suppressor

[78563] genes lgl and dlg regulate basal protein targeting in Drosophila neuroblasts. Nature 408:

[78564] 596-600, 2000.

[78565] 13560.

[78566] Strand, D.; Unger, S.; Corvi, R.; Hartenstein, K.; Schenkel, H.;

[78567] Kalmes, A.; Merdes, G.; Neumann, B.; Krieg-Schneider, F.; Coy, J.

[78568] F.; Poustka, A.; Schwab, M.; Mechler B. M.: A human homologue of

[78569] the Drosophila tumour suppressor gene l(2)gl maps to 17p11.2-12 and

[78570] codes for a cytoskeletal protein that associates with non-muscle myosin

[78571] II heavy chain. Oncogene 11: 291-301, 1995.

[78572] 13561.

[78573] Timchenko, L. T.; Miller, J. W.; Timchenko, N. A.; DeVore, D. R.;

[78574] Datar, K. V.; Lin, L.; Roberts, R.; Caskey, C. T.; Swanson, M. S.

[78575] : Identification of a (CUG)n triplet repeat RNA-binding protein and

[78576] its expression in myotonic dystrophy. Nucleic Acids Res. 24: 4407-4414,

[78577] 1996.

[78578] 13562.

[78579] Timchenko, L. T.; Timchenko, N. A.; Caskey, C. T.; Roberts, R.

[78580] : Novel proteins with binding specificity for DNA CTG re-peats and

[78581] RNA CUG repeats: implications for myotonic dystrophy. Hum. Molec.

[78582] Genet. 5: 115-121, 1996.

[78583] 13563.

[78584] Roelfsema, J. H.; Spruit, L.; Saris, J. J.; Chang, P.; Pirson,

[78585] Y.; van Ommen, G.-J. B.; Peters, D. J. M.; Breuning, M. H.: Mutation

[78586] detection in the repeated part of the PKD1 gene. Am. J. Hum. Genet. 61:

[78587] 1044-1052, 1997.

[78588] 13564.

[78589] Rossetti, S.; Strmecki, L.; Gamble, V.; Burton, S.; Sneddon, V.;

[78590] Peral, B.; Roy, S.; Bakkaloglu, A.; Komel, R.; Winearls, C. G.; Harris,

[78591] P. C.: Mutation analysis of the entire PKD1 gene: genetic and diagnostic

[78592] implications. Am. J. Hum. Genet. 68: 46-63, 2001.

[78593] 13565.

[78594] Scheffers, M. S.; van der Bent, P.; Prins, F.; Spruit, L.; Breuning,

[78595] M. H.; Litvinov, S. V.; de Heer, E.; Peters, D. J. M.: Polycystin-1,

[78596] the product of the polycystic kidney disease 1 gene, co-localizes

[78597] with desmosomes in MDCK cells. Hum. Molec. Genet. 9: 2743-2750,

[78598] 2000.

[78599] 13566.

[78600] Simon, P.; Le Goff, J.-Y.; Ang, K. S.; Charasse, C.; Le

Cacheux,
[78601] P.; Cam, G.: Donnees epidemiologiques, cliniques et pronostiques
[78602] de la maladie polykystique autosomique dominante dans une region francaise. Nephrologie 17:
[78603] 123-130, 1996.

[78604] 13567.

[78605] Terada, T.; Nakanuma, Y.: Congenital biliary dilatation in autosomal
[78606] dominant adult polycystic disease of the liver and kidneys. Arch.
[78607] Path. Lab. Med. 112: 1113-1116, 1988. 76. Trent, R. J.; Wallace,
[78608] R. C.: Alpha thalassaemia: a potential source of error in DNA linkage
[78609] studies for adult polycystic kidney disease. J. Med. Genet. 26:
[78610] 6-9, 1989.

[78611] 13568.

[78612] Thomas, R.; McConnell, R.; Whittacker, J.; Kirkpatrick, P.; Bradley,
[78613] J.; Sandford, R.: Identification of mutations in the repeated

part

[78614] of the autosomal dominant polycystic kidney disease type 1 gene, PKD1,

[78615] by long-range PCR. Am. J. Hum. Genet. 65: 39-49, 1999.

[78616] 13569.

[78617] Watnick, T.; Phakdeekitcharoen, B.; Johnson, A.; Gandolph, M.;

[78618] Wang, M.; Briefel, G.; Klinger, K. W.; Kimberling, W.; Gabow, P.;

[78619] Germino, G. G.: Mutation detection of PKD1 identifies a novel mutation

[78620] common to three families with aneurysms and/or very-early-onset disease. Am.

[78621] J. Hum. Genet. 65: 1561-1571, 1999.

[78622] 13570.

[78623] Watnick, T. J.; Gandolph, M. A.; Weber, H.; Neumann, H. P. H.;

[78624] Germino, G. G.: Gene conversion is a likely cause of mutation in

[78625] PKD1. Hum. Molec. Genet. 7: 1239-1243, 1998.

[78626] 13571.

[78627] Watnick, T. J.; Piontek, K. B.; Cordal, T. M.; Weber, H.; Gandolph,
[78628] M. A.; Qian, F.; Lens, X. M.; Neumann, H. P. H.; Germino, G. G.:
[78629] An unusual pattern of mutation in the duplicated portion of PKD1 is
[78630] revealed by use of a novel strategy for mutation detection. Hum.
[78631] Molec. Genet. 6: 1473-1481, 1997.

[78632] 13572.

[78633] Watnick, T. J.; Torres, V. E.; Gandolph, M. A.; Qian, F.; Onuchic,
[78634] L. F.; Klinger, K. W.; Landes, G.; Germino, G. G.: Somatic mutation
[78635] in individual liver cysts supports a two-hit model of cys-
togenesis
[78636] in autosomal dominant polycystic kidney disease. Molec. Cell 2:
[78637] 247-251, 1998.

[78638] 13573.

[78639] Wary, K. K.; Mariotti, A.; Zurzolo, C.; Giancotti, F. G.: A re-
quirement

[78640] for caveolin-1 and associated kinase Fyn in integrin signaling and

[78641] anchorage-dependent cell growth. Cell 94: 625-634, 1998.

[78642] 13574.

[78643] Yang, G.; Truong, L. D.; Timme, T. L.; Ren, C.; Wheeler, T. M.;

[78644] Park, S. H.; Nasu, Y.; Bangma, C. H.; Kattan, M. W.; Scardino, P.

[78645] T.; Thompson, T. C.: Elevated expression of caveolin is associated

[78646] with prostate and breast cancer. Clin. Cancer Res. 4: 1873-1880,

[78647] 1998.

[78648] 13575.

[78649] Sweadner, K. J.; Rael, E.: The FXYD gene family of small ion transport

[78650] regulators or channels: cDNA sequence, protein signature sequence,

[78651] and expression. Genomics 68: 41-56, 2000.

[78652] 13576.

[78653] Matsuoka, S.; Edwards, M. C.; Bai, C.; Parker, S.; Zhang, P.;

[78654] Baldini, A.; Harper, J. W.; Elledge, S. J.: p57(KIP2), a structurally

[78655] distinct member of the p21(CIP1) Cdk inhibitor family, is a candidate

[78656] tumor suppressor gene. Genes Dev. 9: 650-662, 1995.

[78657] 13577.

[78658] Matsuoka, S.; Thompson, J. S.; Edwards, M. C.; Barletta, J. M.;

[78659] Grundy, P.; Kalikin, L. M.; Harper, J. W.; Elledge, S. J.; Feinberg,

[78660] A. P.: Imprinting of the gene encoding a human cyclin-dependent kinase

[78661] inhibitor, p57(KIP2), on chromosome 11p15. Proc. Nat. Acad. Sci. 93:

[78662] 3026-3030, 1996.

[78663] 13578.

[78664] O'Keefe, D.; Dao, D.; Zhao, L.; Sanderson, R.; Warburton, D.;

[78665] Weiss, L.; Anyane-Yeboa, K.; Tycko, B.: Coding mutations in p57(KIP2)

[78666] are present in some cases of Beckwith-Wiedemann syndrome but are rare

[78667] or absent in Wilms tumors. Am. J. Hum. Genet. 61: 295-303, 1997.

[78668] 13579.

[78669] Tokino, T.; Urano, T.; Furuhata, T.; Matsushima, M.; Miyatsu,

[78670] T.; Sasaki, S.; Nakamura, Y.: Characterization of the human p57(KIP2)

[78671] gene: alternative splicing, insertion/deletion polymorphisms in VNTR

[78672] sequences in the coding region, and mutational analysis. Hum. Genet. 97:

[78673] 625-631, 1996.

[78674] 13580.

[78675] Zhang, P.; Leigeois, N. J.; Wong, C.; Finegold, M.; Hou, H.; Thompson,

[78676] J. C.; Silverman, A.; Harper, J. W.; DePinho, R. A.; Elledge, S. J.

[78677] : Altered cell differentiation and proliferation in mice lacking p57(KIP2)

[78678] indicates a role in Beckwith-Wiedemann syndrome. Nature

387: 151-158,
[78679] 1997.

[78680] 13581.

[78681] Hirai, H.; Roussel, M. F.; Kato, J.-Y.; Ashmun, R. A.; Sherr, C.
[78682] J.: Novel INK4 proteins, p19 and p18, are specific in-hibitors of
[78683] cyclin D-dependent kinases CDK4 and CDK6. Molec. Cell. Biol. 15:
[78684] 2672-2681, 1995.

[78685] 13582.

[78686] Okuda, T.; Hirai, H.; Valentine, V. A.; Shurtleff, S. A.; Kidd,
[78687] V. J.; Lahti, J. M.; Sherr, C. J.; Downing, J. R.: Molecular cloning,
[78688] expression pattern, and chromosomal localization of hu-man CDKN2D/INK4d,
[78689] an inhibitor of cyclin D-dependent kinases. Genomics 29: 623-630,
[78690] 1995.

[78691] 13583.

[78692] Vyas, P.; Vickers, M. A.; Picketts, D. J.; Higgs, D. R.: Con-

servation

[78693] of position and sequence of a novel, widely expressed gene containing

[78694] the major human alpha-globin regulatory element. Genomics 29: 679-689,

[78695] 1995.

[78696] 13584.

[78697] Vyas, P.; Vickers, M. A.; Simmons, D. L.; Ayyub, H.; Craddock,

[78698] C. F.; Higgs, D. R.: Cis-acting sequences regulating expression of

[78699] the human alpha-globin cluster lie within constitutively open chromatin. Cell 69:

[78700] 781-793, 1992.

[78701] 13585.

[78702] Gmachl, M.; Sagan, S.; Ketter, S.; Kreil, G.: The human sperm

[78703] protein PH-20 has hyaluronidase activity. FEBS Lett. 336: 545-548,

[78704] 1993.

[78705] 13586.

[78706] Jones, M. H.; Davey, P. M.; Aplin, H.; Affara, N. A.: Expres-
sion
[78707] analysis, genomic structure, and mapping to 7q31 of the
human sperm
[78708] adhesion molecule gene SPAM1. Genomics 29: 796-800,
1995.
[78709] 13587.

[78710] Lathrop, W. F.; Carmichael, E. P.; Myles, D. G.; Primakoff,
P.
[78711] : cDNA cloning reveals the molecular structure of a sperm
surface
[78712] protein, PH-20, involved in sperm-egg adhesion and the
wide distribution
[78713] of its gene among mammals. J. Cell Biol. 111: 2939-2949,
1990.
[78714] 13588.

[78715] Lin, Y.; Kimmel, L. H.; Myles, D. G.; Primakoff, P.: Molecu-
lar
[78716] cloning of the human and monkey sperm surface protein
PH-20. Proc.
[78717] Nat. Acad. Sci. 90: 10071-10075, 1993.

[78718] 13589.

[78719] Lin, Y.; Mahan, K.; Lathrop, W. F.; Myles, D. G.; Primakoff, P.

[78720] : A hyaluronidase activity of the sperm plasma membrane protein PH-20

[78721] enables sperm to penetrate the cumulus cell layer surrounding the

[78722] egg. J. Cell Biol. 125: 1157-1163, 1994.

[78723] 13590.

[78724] Sago, H.; Kitagawa, M.; Obata, S.; Mori, N.; Taketani, S.; Rochelle,

[78725] J. M.; Seldin, M. F.; Davidson, M.; St. John, T.; Suzuki, S. T.:

[78726] Cloning, expression, and chromosomal localization of a novel cadherin-related

[78727] protein, protocadherin-3. Genomics 29: 631-640, 1995.

[78728] 13591.

[78729] Devlin, A. M.; Ling, E.; Peerson, J. M.; Fernando, S.; Clarke,

[78730] R.; Smith, A. D.; Halsted, C. H.: Glutamate carboxypeptidase II:

[78731] a polymorphism associated with lower levels of serum folate and hyperhomocysteinemia. Hum.

[78732] Molec. Genet. 9: 2837-2844, 2000.

[78733] 13592.

[78734] Israeli, R. S.; Powell, C. T.; Fair, W. R.; Heston, W. D. W.:

[78735] Molecular cloning of a complementary DNA encoding a prostate-specific

[78736] membrane antigen. Cancer Res. 53: 227-230, 1993.

[78737] 13593.

[78738] Leek, J.; Lench, N.; Maraj, B.; Bailey, A.; Carr, I. M.; Ander-sen,

[78739] S.; Cross, J.; Whelan, P.; MacLennan, K. A.; Meredith, D. M.; Markham,

[78740] A. F.: Prostate-specific membrane antigen: evidence for the existence

[78741] of a second related human gene. Brit. J. Cancer 72: 583-588, 1995.

[78742] 13594.

[78743] Maraj, B. H.; Leek, J. P.; Karayi, M.; Ali, M.; Lench, N. J.; Markham,

[78744] A. F.: Detailed genetic mapping around a putative prostate-specific

[78745] membrane antigen locus on human chromosome 11p11.2. Cytogenet. Cell

[78746] Genet. 81: 3-9, 1998.

[78747] 13595.

[78748] O'Keefe, D. S.; Su, S. L.; Bacich, D. J.; Horiguchi, Y.; Luo, Y.;

[78749] Powell, C. T.; Zandvliet, D.; Russell, P. J.; Molloy, P. L.; Nowak,

[78750] N. J.; Shows, T. B.; Mullins, C.; Vonder Haar, R. A.; Fair, W. R.;

[78751] Heston, W. D. W.: Mapping, genomic organization and promoter analysis

[78752] of the human prostate-specific membrane antigen gene. Biochim. Biophys.

[78753] Acta 1443: 113-127, 1998.

[78754] 13596.

[78755] Rinker-Schaeffer, C. W.; Hawkins, A. L.; Su, S. L.; Israeli, R.

[78756] S.; Griffin, C. A.; Isaacs, J. T.; Heston, W. D. W.: Localization

[78757] and physical mapping of the prostate-specific membrane antigen (PSM)

[78758] gene to human chromosome 11. Genomics 30: 105-108, 1995.

[78759] 13597.

[78760] Carlson, S. G.; Eng, E.; Kim, E.-G.; Perlman, E. J.; Copeland,

[78761] T. D.; Ballermann, B. J.: Expression of SET, an inhibitor of protein

[78762] phosphatase 2A, in renal development and Wilms' tumor. J. Am. Soc.

[78763] Nephrol. 9: 1873-1880, 1998.

[78764] 13598.

[78765] Tripoulas, N.; LaJeunesse, D.; Gildea, J.; Shearn, A.: The Drosophila

[78766] ash1 gene product, which is localized at specific sites on polytene

[78767] chromosomes, contains a SET domain and a PHD finger. Genetics 143:

[78768] 913-928, 1996.

[78769] 13599.

[78770] Tschiersch, B.; Hofmann, A.; Krauss, V.; Dorn, R.; Korge, G.; Reuter,

[78771] G.: The protein encoded by the Drosophila position-effect variegation

[78772] suppressor gene Su(var)3-9 combines domains of antagonistic regulators

[78773] of homeotic gene complexes. EMBO J. 13: 3822-3831, 1994.

[78774] 13600.

[78775] von Lindern, M.; van Baal, S.; Wiegant, J.; Raap, A.; Hagemeijer,

[78776] A.; Grosveld, G.: 'can,' a putative oncogene associated with myeloid

[78777] leukemogenesis, may be activated by fusion of its 3-prime half to

[78778] different genes: characterization of the 'set' gene. Molec. Cell.

[78779] Biol. 3346-3355, 1992.

[78780] 13601.

[78781] Bartsch, J. W.; Mukai, H.; Takahashi, N.; Ronsiek, M.; Fuchs, S.;

[78782] Jockusch, H.; Ono, Y.: The protein kinase N (PKN) gene PRKCL1/Prkcl1

[78783] maps to human chromosome 19p12-p13.1 and mouse chromosome 8 with close

[78784] linkage to the myodystrophy (myd) mutation. Genomics

49: 129-132,
[78785] 1998.

[78786] 13602.

[78787] Mukai, H.; Ono, Y.: A novel protein kinase with leucine zipper-like
[78788] sequences: its catalytic domain is highly homologous to that of protein
[78789] kinase C. Biochem. Biophys. Res. Commun. 199: 897-904, 1994.
[78790] 13603.

[78791] Palmer, R. H.; Ridden, J.; Parker, P. J.: Identification of multiple,
[78792] novel, protein kinase C-related gene products. FEBS Lett. 356: 5-8,
[78793] 1994.

[78794] 13604.

[78795] Palmer, R. H.; Ridden, J.; Parker, P. J.: Cloning and expres-sions
[78796] patterns of two members of a novel protein-ki-nase-C-related kinase
[78797] family. Europ. J. Biochem. 227: 344-351, 1995.

[78798] 13605.

[78799] Yu, W.; Liu, J.; Morrice, N. A.; Wettenhall, R. E. H.: Isolation

[78800] and characterization of a structural homologue of human PRK2 from

[78801] rat liver. J. Biol. Chem. 272: 10030-10034, 1997.

[78802] 13606.

[78803] Hernandez, A.; Park, J. P.; Lyon, G. J.; Mohandas, T. K.; St. Germain,

[78804] D. L.: Localization of the type 3 iodothyronine deiodinase (DIO3)

[78805] gene to human chromosome 14q32 and mouse chromosome 12F1. Genomics 53:

[78806] 119-121, 1998.

[78807] 13607.

[78808] Honore, B.; Rasmussen, H. H.; Vorum, H.; Dejgaard, K.; Liu, X.;

[78809] Gromov, P.; Madsen, P.; Gesser, B.; Tommerup, N.; Celis, J. E.: Heterogeneous

[78810] nuclear ribonucleoproteins H, H-prime, and F are members of a ubiquitously

[78811] expressed subfamily of related but distinct proteins en-

coded by genes
[78812] mapping to different chromosomes. J. Biol. Chem. 270: 28780-28789,
[78813] 1995.

[78814] 13608.

[78815] Huang, S. A.; Tu, H. M.; Harney, J. W.; Venihaki, M.; Butte, A.
[78816] J.; Kozakewich, H. P. W.; Fishman, S. J.; Larsen, P. R.: Severe hypothyroidism
[78817] caused by type 3 iodothyronine deiodinase in infantile hemangiomas. New
[78818] Eng. J. Med. 343: 185-189, 2000.

[78819] 13609.

[78820] Salvatore, D.; Low, S. C.; Berry, M.; Maia, A. L.; Harney, J. W.;
[78821] Croteau, W.; St. German, D. L.; Larsen, P. R.: Type 3 iodothyronine
[78822] deiodinase: cloning, in vitro expression, and functional analysis
[78823] of the placental selenoenzyme. J. Clin. Invest. 96: 2421-2430, 1995.
[78824] 13610.

[78825] Ardehali, H.; Tiller, G. E.; Printz, R. L.; Mochizuki, H.; Proc-
hazka,
[78826] M.; Granner, D. K.: A novel (TA)n polymorphism in the
hexokinase
[78827] II gene: application to noninsulin-dependent diabetes
mellitus in
[78828] the Pima Indians. Hum. Genet. 97: 482-485, 1996.

[78829] 13611.

[78830] Echwald, S. M.; Bjorbaek, C.; Hansen, T.; Clausen, J. O.;
Vestergaard,
[78831] H.; Zierarth, J. R.; Printz, R. L.; Granner, D. K.; Pedersen,
O.:
[78832] Identification of four amino acid substitutions in hexoki-
nase II and
[78833] studies of relationships to NIDDM, glucose effectiveness,
and insulin
[78834] sensitivity. Diabetes 44: 347-353, 1995.

[78835] 13612.

[78836] Heikkinen, S.; Suppola, S.; Malkki, M.; Deeb, S. S.; Janne,
J.;
[78837] Laakso, M.: Mouse hexokinase II gene: structure, cDNA,
promoter analysis,

[78838] and expression pattern. Mammalian Genome 11: 91-96, 2000.

[78839] 13613.

[78840] Laakso, M.; Malkki, M.; Deeb, S. S.: Amino acid substitutions

[78841] in hexokinase II among patients with NIDDM. Diabetes 44: 330-334,

[78842] 1995.

[78843] 13614.

[78844] Lehto, M.; Xiang, K.; Stoffel, M.; Espinosa, R., III; Groop, L.

[78845] C.; Le Beau, M. M.; Bell, G. I.: Human hexokinase II: localization

[78846] of the polymorphic gene to chromosome 2. Diabetologia 36: 1299-1302,

[78847] 1993.

[78848] 13615.

[78849] Mathupala, S. P.; Heese, C.; Pedersen, P. L.: Glucose catabolism

[78850] in cancer cells: the type II hexokinase promoter contains functionally

[78851] active response elements for the tumor suppressor p53. J.

Biol. Chem. 272:
[78852] 22776-22780, 1997.

[78853] 13616.

[78854] Mathupala, S. P.; Rempel, A.; Pedersen, P. L.: Glucose catabolism
[78855] in cancer cells: isolation, sequence, and activity of the promoter
[78856] for type II hexokinase. J. Biol. Chem. 270: 16918-16925, 1995.
[78857] 13617.

[78858] Vidal-Puig, A.; Printz, R. L.; Stratton, I. M.; Granner, D. K.;

[78859] Moller, D. E.: Analysis of the hexokinase II gene in sub-jects with
[78860] insulin resistance and NIDDM and detection of a gln142-to-his substitution. Diabetes 44:
[78861] 340-346, 1995.

[78862] 13618.

[78863] Luo, Y.; Roeder, R. G.: Cloning, functional characteriza-tion,
[78864] and mechanism of action of the B-cell-specific transcrip-tional activator

[78865] OCA-B. Molec. Cell. Biol. 15: 4115-4124, 1995.

[78866] 13619.

[78867] Staudt, L. M.; Lenardo, M. J.: Immunoglobulin gene transcription. Ann.

[78868] Rev. Immun. 9: 373-398, 1991.

[78869] 13620.

[78870] Strubin, M.; Newell, J. W.; Matthias, P.: OBF-1, a novel B cell-specific

[78871] coactivator that stimulates immunoglobulin promoter activity through

[78872] association with octamer-binding proteins. Cell 80: 497-506, 1995.

[78873] 13621.

[78874] Hahm, K.; Kim, G.; Turck, C. W.; Smale, S. T.: Isolation of a

[78875] murine gene encoding a nucleic acid-binding protein with homology

[78876] to hnRNP K. Nucleic Acids Res. 21: 3894 only, 1993.

[78877] 13622.

[78878] Leffers, H.; Dejgaard, K.; Celis, J. E.: Characterisation of two

[78879] major cellular poly(rC)-binding human proteins, each containing three

[78880] K-homologous (KH) domains. Europ. J. Biochem. 230: 447-453, 1995.

[78881] 13623.

[78882] Bouchard, M. J.; Wang, L.-H.; Schneider, R. J.: Calcium signaling

[78883] by HBx protein in hepatitis B virus DNA replication. Science 294:

[78884] 2376-2378, 2001.

[78885] 13624.

[78886] Calalb, M. B.; Polte, T. R.; Hanks, S. K.: Tyrosine phosphorylation

[78887] of focal adhesion kinase at sites in the catalytic domain regulates

[78888] kinase activity: a role for Src family kinases. Molec. Cell. Biol. 15:

[78889] 954-963, 1995.

[78890] 13625.

[78891] Ganem, D.: The X files--one step closer to closure. Science 294:

[78892] 2299-2300, 2001.

[78893] 13626.

[78894] Herzog, H.; Nicholl, J.; Hort, Y. J.; Sutherland, G. R.; Shine,

[78895] J.: Molecular cloning and assignment of FAK2, a novel hu-
man focal

[78896] adhesion kinase, to 8p11.2-p22 by nonisotopic in situ
hybridization. Genomics 32:

[78897] 484-486, 1996.

[78898] 13627.

[78899] Lev, S.; Moreno, H.; Martinez, R.; Canoll, P.; Peles, E.;
Musacchio,

[78900] J. M.; Plowman, G. D.; Rudy, B.; Schlessinger, J.: Protein ty-
rosine

[78901] kinase PYK2 involved in Ca(2+)-induced regulation of ion
channel and

[78902] MAP kinase functions. Nature 376: 737-745, 1995.

[78903] 13628.

[78904] Manser, E.; Leung, T.; Salihuddin, H.; Tan, L.; Lim, L.: A
non-receptor

[78905] tyrosine kinase that inhibits the GTPase activity of
p21(cdc42). Nature 363:

[78906] 364-367, 1993.

[78907] 13629.

[78908] Matsuya, M.; Sasaki, H.; Aoto, H.; Mitaka, T.; Nagura, K.; Ohba,

[78909] T.; Ishino, M.; Takahashi, S.; Suzuki, R.; Sasaki, T.: Cell adhesion

[78910] kinase beta forms a complex with a new member, Hic-5, of proteins

[78911] localized at focal adhesions. J. Biol. Chem. 273: 1003-1014, 1998.

[78912] 13630.

[78913] Dawson, P. A.; Oelkers, P.: Bile acid transporters. Curr. Opin.

[78914] Lipid. 6: 109-114, 1995.

[78915] 13631.

[78916] Hess Thaysen, E.; Pedersen, L.: Idiopathic bile acid catharis. Gut 17:

[78917] 965-970, 1976.

[78918] 13632.

[78919] Heubi, J. E.; Balistreri, W. F.; Fondacaro, J. D.; Partin, J. C.;

[78920] Schubert, W. K.: Primary bile and malabsorption: defective in vitro
[78921] ileal active bile acid transport. Gastroenterology 83: 804-811,
[78922] 1982.

[78923] 13633.

[78924] Heubi, J. E.; Balistreri, W. F.; Partin, J. C.; Schubert, W. K.;

[78925] McGraw, C. A.: Refractory infantile diarrhea due to primary bile
[78926] acid malabsorption. J. Pediat. 94: 546-551, 1979.

[78927] 13634.

[78928] Jonas, A.; Diver-Haber, A.; Avigad, S.: Well-compensated primary
[78929] bile acid malabsorption presenting as chronic nonspecific diarrhea. J.
[78930] Pediat. Gastroent. Nutr. 5: 143-146, 1986.

[78931] 13635.

[78932] Lammert, F.; Paigen, B.; Carey, M. C.: Localization of the ileal
[78933] sodium-bile salt cotransporter gene (Slc10a2) to mouse chromosome

[78934] 8. Mammalian Genome 9: 173-174, 1998.

[78935] 13636.

[78936] Oelkers, P.; Kirby, L. C.; Heubi, J. E.; Dawson, P. A.: Pri-
mary
[78937] bile acid malabsorption caused by mutations in the ileal
sodium-dependent
[78938] bile acid transporter gene (SLC10A2). J. Clin. Invest. 99:
1880-1887,
[78939] 1997.

[78940] 13637.

[78941] Small, D. M.: Point mutations in the ileal bile salt trans-
porter
[78942] cause leaks in the enterohepatic circulation leading to se-
vere chronic
[78943] diarrhea and malabsorption. (Editorial) J. Clin. Invest. 99:
1807-1808,
[78944] 1997.

[78945] 13638.

[78946] Small, D. M.; Dowling, R. H.; Redinger, R. N.: The entero-
hepatic
[78947] circulation of bile salts. Arch. Intern. Med. 130: 552-573,

1972.

[78948] 13639.

[78949] Wong, M. H.; Oelkers, P.; Dawson, P. A.: Identification of a

[78950] mutation in the ileal sodium-dependent bile acid transporter gene

[78951] that abolishes transport activity. J. Biol. Chem. 270: 27228-27234,

[78952] 1995.

[78953] 13640.

[78954] Wong, M. H.; Rao, P. N.; Pettenati, M. J.; Dawson, P. A.: Localization

[78955] of the ileal sodium-bile acid cotransporter gene (SLC10A2) to human

[78956] chromosome 13q33. Genomics 33: 538-540, 1996.

[78957] 13641.

[78958] Luo, Z. G.; Wang, Q.; Zhou, J. Z.; Wang, J.; Lou, Z.; Liu, M.;

[78959] He, X.; Wynshaw-Boris, A.; Xiong, W. C.; Lu, B.; Mei, L.: Regulation

[78960] of AChR clustering by Dishevelled interacting with MuSK and PAK1. Neuron 35:

[78961] 489-505, 2002.

[78962] 13642.

[78963] Meyer, J.; Wirth, J.; Held, M.; Schempp, W.; Scherer, G.:
SOX20,

[78964] a new member of the SOX gene family, is located on chro-
mosome 17p13. Cytogenet.

[78965] Cell Genet. 72: 246-249, 1996.

[78966] 13643.

[78967] Vujic, M.; Rajic, T.; Goodfellow, P. N.; Stevanovic, M.: cDNA

[78968] characterization and high resolution mapping of the hu-
man SOX20 gene. Mammalian

[78969] Genome 9: 1059-1061, 1998.

[78970] 13644.

[78971] Eng, C.: Will the real Cowden syndrome please stand up:
revised

[78972] diagnostic criteria. J. Med. Genet. 37: 828-830, 2000.

[78973] 13645.

[78974] Eng, C.: To be or not to BMP. Nature Genet. 28: 105-107,
2001.

[78975] 13646.

[78976] Ide, H.; Saito-Ohara, F.; Ohnami, S.; Osada, Y.; Ikeuchi, T.;

Yoshida,

[78977] T.; Terada, M.: Assignment of the BMPR1A and BMPR1B genes to human

[78978] chromosome 10q22.3 and 4q23-q24 by in situ hybridization and radiation

[78979] hybrid mapping. Cytogenet. Cell Genet. 81: 285-286, 1998.

[78980] 13647.

[78981] Schuffenhauer, S.; Lichtner, P.; Peykar-Derakhshandeh, P.; Murken,

[78982] J.; Haas, O. A.; Back, E.; Wolff, G.; Zabel, B.; Barisic, I.; Rauch,

[78983] A.; Borochowitz, Z.; Dallapiccola, B.; Ross, M.; Meitinger, T.: Deletion

[78984] mapping on chromosome 10p and definition of a critical region for

[78985] the second DiGeorge syndrome locus (DGS2). Europ. J. Hum. Genet. 6:

[78986] 213-225, 1998.

[78987] 13648.

[78988] Carter, B. S.; Ewing, C. M.; Ward, W. S.; Treiger, B. F.; Aalders,

[78989] T. W.; Schalken, J. A.; Epstein, J. I.; Isaacs, W. B.: Allelic loss
[78990] of chromosomes 16q and 10q in human prostate cancer. Proc. Nat. Acad.
[78991] Sci. 87: 8751-8755, 1990.

[78992] 13649.

[78993] Lee, S. W.: H-cadherin, a novel cadherin with growth in-hibitory
[78994] functions and diminished expression in human breast cancer. Nature
[78995] Med. 2: 776-782, 1996.

[78996] 13650.

[78997] Sato, M.; Mori, Y.; Sakurada, A.; Fujimura, S.; Horii, A.: The
[78998] H-cadherin (CDH13) gene is inactivated in human lung cancer. Hum.
[78999] Genet. 103: 96-101, 1998.

[79000] 13651.

[79001] Tsuda, H.; Hirohashi, S.: Identification of multiple breast cancers
[79002] of multicentric origin by histological observations and distribution

[79003] of allele loss on chromosome 16q. Cancer Res. 55: 3395-3398, 1995.

[79004] 13652.

[79005] Tsuda, H.; Oda, T.; Sakamoto, M.; Hirohashi, S.: Different pattern

[79006] of chromosomal allele loss in multiple hepatocellular carcinomas as

[79007] evidence of their multifocal origin. Cancer Res. 52: 1504-1509,

[79008] 1992.

[79009] 13653.

[79010] Bedell, J. A.; Wagner-McPherson, C. B.; Bengtsson, U.; Handa, K.;

[79011] Dumars, K. W.; Marsh, J. L.; Smith, M.; McPherson, J. D.: A 1p deletion

[79012] syndrome patient is hemizygous for a human homologue of the Drosophila

[79013] dishevelled gene. (Abstract) Am. J. Hum. Genet. 59: A298 only, 1996.

[79014] 13654.

[79015] Klingensmith, J.; Nusse, R.; Perrimon, N.: The Drosophila segment

[79016] polarity gene dishevelled encodes a novel protein required for response

[79017] to the wingless signal. Genes Dev. 8: 118-130, 1994.

[79018] 13655.

[79019] Lijam, N.; Paylor, R.; McDonald, M. P.; Crawley, J. N.; Deng, C.-X.;

[79020] Herrup, K.; Stevens, K. E.; Maccaferri, G.; McBain, C. J.; Sussman,

[79021] D. J.; Wynshaw-Boris, A.: Social interaction and sensori-motor gating

[79022] abnormalities in mice lacking Dvl1. Cell 90: 895-905, 1997.

[79023] 13656.

[79024] Pizzuti, A.; Amati, F.; Calabrese, G.; Mari, A.; Colosimo, A; Silani,

[79025] V.; Giardino, L.; Ratti, A.; Penso, D.; Calza, L.; Palka, G.; Scarlato,

[79026] G.; Novelli, G.; Dallapicolla, B.: cDNA characterization and chromosomal

[79027] mapping of two human homologs of the Drosophila di-shevelled polarity

[79028] gene. Hum. Molec. Genet. 5: 953-958, 1996.

[79029] 13657.

[79030] Semenov, M. V.; Snyder, M.: Human dishevelled genes constitute

[79031] a DHR-containing multigene family. Genomics 42: 302-310, 1997.

[79032] 13658.

[79033] Wallingford, J. B.; Rowning, B. A.; Vogeli, K. M.; Rothbacher,

[79034] U.; Fraser, S. E.; Harland, R. M.: Dishevelled controls cell polarity

[79035] during Xenopus gastrulation. Nature 405: 81-85, 2000.

[79036] 13659.

[79037] Lee, J.; Hahn, Y.; Yun, J. H.; Mita, K.; Chung, J. H.: Characterization

[79038] of JDP genes, an evolutionarily conserved J domain-only protein family,

[79039] from human and moths. Biochim. Biophys. Acta 1491: 355-363, 2000.

[79040] 13660.

[79041] Carson-Walter, E. B.; Watkins, D. N.; Nanda, A.; Vogelstein, B.;

[79042] Kinzler, K. W.; St. Croix, B.: Cell surface tumor endothelial markers
[79043] are conserved in mice and humans. Cancer Res. 61: 6649-6655, 2001.
[79044] 13661.

[79045] St. Croix, B.; Rago, C.; Velculescu, V.; Traverso, G.; Romans,
[79046] K. E.; Montgomery, E.; Lal, A.; Riggins, G. J.; Lengauer, C.; Vogelstein,
[79047] B.; Kinzler, K. W.: Genes expressed in human tumor endothelium. Science 289:
[79048] 1197-1202, 2000.

[79049] 13662.

[79050] Gosling, J.; Dairaghi, D. J.; Wang, Y.; Hanley, M.; Talbot, D.;
[79051] Miao, Z.; Schall, T. J.: Cutting edge: identification of a novel
[79052] chemokine receptor that binds dendritic cell- and T cell-active chemokines
[79053] including ELC, SLC, and TECK. J. Immun. 164: 2851-2856, 2000.
[79054] 13663.

[79055] Donovan, A.; Brownlie, A.; Zhou, Y.; Shepard, J.; Pratt, S. J.;

[79056] Moynihan, J.; Paw, B. H.; Drejer, A.; Barut, B.; Zapata, Z.; Law,

[79057] T. C.; Brugnara, C.; Lux, S. E.; Pinkus, G. S.; Pinkus, J. L.; Kingsley,

[79058] P. D.; Palis, J.; Fleming, M. D.; Andrews, N. C.; Zon, L. I.: Positional

[79059] cloning of zebrafish ferroportin1 identifies a conserved vertebrate

[79060] iron exporter. Nature 403: 776-781, 2000.

[79061] 13664.

[79062] Cohn, R. D.; Campbell, K. P.: Molecular basis of muscular dystrophies. Muscle

[79063] Nerve 23: 1456-1471, 2000.

[79064] 13665.

[79065] Gillespie, C. S.; Lee, M.; Fantes, J. F.; Brophy, P. J.: The gene

[79066] encoding the Schwann cell protein periaxin localizes on mouse chromosome

[79067] 7 (Prx). Genomics 41: 297-298, 1997.

[79068] 13666.

[79069] Gillespie, C. S.; Sherman, D. L.; Fleetwood-Walker, S. M.; Cottrell,
[79070] D. F.; Tait, S.; Garry, E. M.; Wallace, V. C. J.; Ure, J.; Griffiths,
[79071] I. R.; Smith, A. Brophy, P. J.: Peripheral demyelination and neuropathic
[79072] pain behavior in periaxin-deficient mice. Neuron 26: 523-531, 2000.
[79073] 13667.

[79074] Schuler, G. D.: Sequence mapping by electronic PCR. Genome Res. 7:
[79075] 541-550, 1997.

[79076] 13668.

[79077] Takashima, H.; Boerkoel, C. F.; De Jonghe, P.; Ceuterick, C.; Martin,
[79078] J.-J.; Voit, T.; Schroder, J.-M.; Williams, A.; Brophy, P. J.; Timmerman,
[79079] V.; Lupski, J. R.: Periaxin mutations cause a broad spectrum of demyelinating
[79080] neuropathies. Ann. Neurol. 51: 709-715, 2002.

[79081] 13669.

[79082] Liu, Q.-Y.; Wang, L. F.; Miao, S. Y.; Catterall, J. F.: Expression
[79083] and characterization of a novel human sperm membrane protein. Biol.
[79084] Reprod. 54: 323-330, 1996.

[79085] 13670.

[79086] Miao, S.; Yan, Y.; Li, Y.; Bai, Y.; Wei, S.; Zong, C.; Zhao, M.;
[79087] Zong, S.; Wang, L.: cDNA encoding a human sperm membrane protein
[79088] BS-84. Prog. Natural Sci. 5: 119-122, 1995.

[79089] 13671.

[79090] Wang, H.; Miao, S.; Chen, D.; Wang, L.; Koide, S. S.: Assignment
[79091] of chromosomal locus and evidence for alternatively spliced mRNAs
[79092] of a human sperm membrane protein (hSMP-1). Biochim. Biophys. Acta 1447:
[79093] 119-124, 1999.

[79094] 13672.

[79095] Guzman-Rojas, L.; Sims, J. C.; Rangel, R.; Guret, C.; Sun, Y.;

[79096] Alcocer, J. M.; Martinez-Valdez, H.: PRELI, the human homologue of

[79097] the avian px19, is expressed by germinal center B lymphocytes. Int.

[79098] Immun. 12: 607-612, 2000.

[79099] 13673.

[79100] Blasina, A.; de Weyer, I. V.; Laus, M. C.; Luyten, W. H.; Parker,

[79101] A. E.; McGowan, C. H.: A human homologue of the checkpoint kinase

[79102] Cds1 directly inhibits Cdc25 phosphatase. Curr. Biol. 14: 1-10,

[79103] 1999.

[79104] 13674.

[79105] Brown, A. L.; Lee, C.-H.; Schwarz, J. K.; Mitiku, N.; Piwnica-Worms,

[79106] H.; Chung, J. H.: A human Cds1-related kinase that functions downstream

[79107] of ATM protein in the cellular response to DNA damage. Proc. Nat.

[79108] Acad. Sci. 96: 3745-3750, 1999.

[79109] 13675.

[79110] Chaturvedi, P.; Eng, W. K.; Zhu, Y.; Mattern, M. R.; Mishra, R.;

[79111] Hurle, M. R.; Zhang, X.; Annan, R. S.; Lu, Q.; Faucette, L. F.; Scott,

[79112] G. F.; Li, X.; Carr, S. A.; Johnson, R. K.; Winkler, J. D.; Zhou,

[79113] B. B.: Mammalian Chk2 is a downstream effector of the ATM-dependent

[79114] DNA damage checkpoint pathway. Oncogene 18: 4047-4054, 1999.

[79115] 13676.

[79116] Chehab, N. H.; Malikzay, A.; Appel, M.; Halazonetis, T. D.: Chk2/hCds1

[79117] functions as a DNA damage checkpoint in G-1 by stabiliz-ing p53. Genes

[79118] Dev. 14: 278-288, 2000.

[79119] 13677.

[79120] Ino, Y.; Wahrer, D. C. R.; Bell, D. W.; Haber, D. A.; Louis, D.

[79121] N.: Mutation analysis of the hCHK2 gene in primary hu-man malignant

[79122] gliomas. (Letter) Neurogenetics 3: 45-46, 2000.

[79123] 13678.

[79124] Lee, S. B.; Kim, S. H.; Bell, D. W.; Wahrer, D. C. R.; Schiripo,

[79125] T. A.; Jorczak, M. M.; Sgroi, D. C.; Garber, J. E.; Li, F. P.;
Nichols,

[79126] K. E.; Varley, J. M.; Godwin, A. K.; Shannon, K. M.; Harlow,
E.; Haber,

[79127] D. A.: Destabilization of CHK2 by a missense mutation as-
sociated

[79128] with Li-Fraumeni syndrome. Cancer Res. 61: 8062-8067,
2001.

[79129] 13679.

[79130] Lopes, M.; Cotta-Ramusino, C.; Pellicioli, A.; Liberi, G.; Pl-
evani,

[79131] P.; Muzi-Falconi, M. Newlon, C. S.; Foiani, M.: The DNA
replication

[79132] checkpoint response stabilizes stalled replication forks.
Nature 412:

[79133] 557-561, 2001.

[79134] 13680.

[79135] Matsuoka, S.; Huang, M.; Elledge, S. J.: Linkage of ATM to
cell

[79136] cycle regulation by the Chk2 protein kinase. Science 282: 1893-1897,

[79137] 1998.

[79138] 13681.

[79139] Meijers-Heijboer, H.; van den Ouweland, A.; Klijn, J.; Wasielewski,

[79140] M.; de Snoo, A.; Oldenburg, R.; Hollestelle, A.; Houben, M.; Crepin,

[79141] E.; van Veghel-Plandsoen, M.; Elstrodt, F.; van Duijn, C.; and 29

[79142] others: Low-penetrance susceptibility to breast cancer due to CHEK2*1100delC

[79143] in noncarriers of BRCA1 or BRCA2 mutations. Nature Genet. 31: 55-59,

[79144] 2002.

[79145] 13682.

[79146] Miller, C. W.; Ikezoe, T.; Krug, U.; Hofmann, W.-K.; Tavor, S.;

[79147] Vegesna, V.; Tsukasaki, K.; Takeuchi, S.; Koeffler, H. P.: Mutations

[79148] of the CHK2 gene are found in some osteosarcomas, but are rare in

[79149] breast, lung, and ovarian tumors. Genes Chromosomes Cancer 33: 17-21,

[79150] 2002.

[79151] 13683.

[79152] Sogo, J. M.; Lopes, M.; Foiani, M.: Fork reversal and ssDNA accumulation

[79153] at stalled replication forks owing to checkpoint defects. Science 297:

[79154] 599-602, 2002.

[79155] 13684.

[79156] Vahteristo, P.; Bartkova, J.; Eerola, H.; Syrjakoski, K.; Ojala,

[79157] S.; Kilpivaara, O.; Tamminen, A.; Kononen, J.; Aittomaki, K.; Heikkila,

[79158] P.; Holli, K.; Blomqvist, C.; Bartek, J.; Kallioniemi, O.-P.; Nevanlinna,

[79159] H.: A CHEK2 genetic variant contributing to a substantial fraction

[79160] of familial breast cancer. Am. J. Hum. Genet. 71: 432-438, 2002.

[79161] 13685.

[79162] Vahteristo, P.; Tamminen, A.; Karvinen, P.; Eerola, H.; Ek-

lund,

[79163] C.; Aaltonen, L. A.; Blomqvist, C.; Aittomaki, K.; Nevanlinna, H.

[79164] : p53, CHK2, and CHK1 genes in Finnish families with Li-Fraumeni syndrome:

[79165] further evidence of CHK2 in inherited cancer predisposition. Cancer

[79166] Res. 61: 5718-5722, 2001.

[79167] 13686.

[79168] De, S. K.; Enders, G. C.; Andrews, G. K.: High levels of metallothionein

[79169] messenger RNAs in male germ cells of the adult mouse. Molec. Endocr. 5:

[79170] 628-636, 1991.

[79171] 13687.

[79172] Salehi-Ashtiani, K.; Widrow, R. J.; Markert, C. L.; Goldberg, E.

[79173] : Testis-specific expression of a metallothionein I-driven transgene

[79174] correlates with undermethylation of the locus in testicular DNA. Proc.

[79175] Nat. Acad. Sci. 90: 8886-8890, 1993.

[79176] 13688.

[79177] Sugihara, T.; Wadhwa, R.; Kaul, S. C.; Mitsui, Y.: A novel testis-specific

[79178] metallothionein-like protein, tesmin, is an early marker of male germ

[79179] cell differentiation. Genomics 57: 130-136, 1999.

[79180] 13689.

[79181] Larrouy, D.; Vidal, H.; Andreelli, F.; Laville, M.; Langin, D.

[79182] : Cloning and mRNA tissue distribution of human PPAR-gamma coactivator-1. Int.

[79183] J. Obesity 23: 1327-1332, 1999.

[79184] 13690.

[79185] Lin, J.; Wu, H.; Tarr, P. T.; Zhang, C.-Y.; Wu, Z.; Boss, O.; Michael,

[79186] L. F.; Puigserver, P.; Isotani, E.; Olson, E. N.; Lowell, B. B.; Bassel-Duby,

[79187] R.; Spiegelman, B. M.: Transcriptional co-activator PGC-1-alpha drives

[79188] the formation of slow-twitch muscle fibres. Nature 418: 797-801,

[79189] 2002.

[79190] 13691.

[79191] Monsalve, M.; Wu, Z.; Adelmant, G.; Puigserver, P.; Fan, M.; Spiegelman,

[79192] B. M.: Direct coupling of transcription and mRNA processing through

[79193] the thermogenic coactivator PGC-1. Molec. Cell 6: 307-316, 2000.

[79194] 13692.

[79195] Puigserver, P.; Adelmant, G.; Wu, Z.; Fan, M.; Xu, J.; O'Malley,

[79196] B.; Spiegelman, B. M.: Activation of PPAR-gamma coactivator-1 through

[79197] transcription factor docking. Science 286: 1368-1371, 1999.

[79198] 13693.

[79199] Puigserver, P.; Rhee, J.; Lin, J.; Wu, Z.; Yoon, J. C.; Zhang,

[79200] C.-Y.; Krauss, S.; Mootha, V. K.; Lowell, B. B.; Spiegelman, B. M.

[79201] : Cytokine stimulation of energy expenditure through p38 MAP kinase

[79202] activation of PPAR-gamma coactivator-1. Molec. Cell 8: 971-982,

[79203] 2001.

[79204] 13694.

[79205] Puigserver, P.; Wu, Z.; Park, C. W.; Graves, R.; Wright, M.; Spiegelman,

[79206] B. M.: A cold-inducible coactivator of nuclear receptors linked to

[79207] adaptive thermogenesis. Cell 92: 829-839, 1998.

[79208] 13695.

[79209] Waite, L. L.; Person, E. C.; Zhou, Y.; Lim, K.-H.; Scanlan, T.

[79210] S.; Taylor, R. N.: Placental peroxisome proliferator-acti-vated receptor-gamma

[79211] is up-regulated by pregnancy serum. J. Clin. Endocr. Metab. 85:

[79212] 3808-3814, 2000.

[79213] 13696.

[79214] Wu, Z.; Puigserver, P.; Andersson, U.; Zhang, C.; Adelmant, G.;

[79215] Mootha, V.; Troy, A.; Cinti, S.; Lowell, B.; Scarpulla, R. C.; Spiegelman,

[79216] B. M.: Mechanisms controlling mitochondrial biogenesis and respiration

[79217] through the thermogenic coactivator PGC-1. Cell 98: 115-124, 1999.

[79218] 13697.

[79219] Yoon, J. C.; Puigserver, P.; Chen, G.; Donovan, J.; Wu, Z.; Rhee,

[79220] J.; Adelmant, G.; Stafford, J.; Kahn, C. R.; Granner, D. K.; Newgard,

[79221] C. B.; Spiegelman, B. M.: Control of hepatic gluconeogenesis through

[79222] the transcriptional coactivator PGC-1. Nature 413: 131-138, 2001.

[79223] 13698.

[79224] Ellis, J. H.; Ashman, C.; Burden, M. N.; Kilpatrick, K. E.; Morse,

[79225] M. A.; Hamblin, P. A.: GRID: a novel Grb-2-related adapter protein

[79226] that interacts with the activated T cell costimulatory receptor CD28. J.

[79227] Immun. 164: 5805-5814, 2000.

[79228] 13699.

[79229] Qiu, M.; Hua, S.; Agrawal, M.; Li, G.; Cai, J.; Chan, E.; Zhou,

[79230] H.; Luo, Y.; Liu, M.: Molecular cloning and expression of human Grap-2,
[79231] a novel leukocyte-specific SH2- and SH3-containing adaptor-like protein
[79232] that binds to Gab-1. Biochem. Biophys. Res. Commun. 253: 443-447,
[79233] 1998.

[79234] 13700.

[79235] Yoder, J.; Pham, C.; Iizuka, Y.-M.; Kanagawa, O.; Liu, S. K.; McGlade,
[79236] J.; Cheng, A. M.: Requirement for the SLP-76 adaptor GADS in T cell
[79237] development. Science 291: 1987-1991, 2001.

[79238] 13701.

[79239] Hadjantonakis, A.-K.; Formstone, C. J.; Little, P. F. R.: mCelsr1
[79240] is an evolutionarily conserved seven-pass transmembrane receptor and
[79241] is expressed during mouse embryonic development. Mech. Dev. 78:
[79242] 91-95, 1998.

[79243] 13702.

[79244] Hadjantonakis, A.-K.; Sheward, W. J.; Harmar, A. J.; de Galan,

[79245] L.; Hoovers, J. M. N.; Little, P. F. R.: Celsr1, a neural-specific

[79246] gene encoding an unusual seven-pass transmembrane receptor, maps to

[79247] mouse chromosome 15 and human chromosome 22qter. Genomics 45: 97-104,

[79248] 1997.

[79249] 13703.

[79250] Bonny, C.; Oberson, A.; Steinmann, M.; Schorderet, D. F.; Nicod,

[79251] P.; Waeber, G.: IB1 reduces cytokine-induced apoptosis of insulin-secreting

[79252] cells. J. Biol. Chem. 275: 16466-16472, 2000.

[79253] 13704.

[79254] Mooser, V.; Maillard, A.; Bonny, C.; Steinmann, M.; Shaw, P.; Yarnall,

[79255] D. P.; Burns, D. K.; Schorderet, D. F.; Nicod, P.; Waeber, G.: Genomic

[79256] organization, fine-mapping, and expression of the human islet-brain

[79257] 1 (IB1)/c-Jun-amino-terminal kinase interacting protein-1 (JIP-1)

[79258] gene. Genomics 55: 202-208, 1999.

[79259] 13705.

[79260] Thompson, N. A.; Haefliger, J.-A.; Senn, A.; Tawadros, T.; Magara,

[79261] F.; Ledermann, B.; Nicod, P.; Waeber, G.: Islet-brain1/JNK-interacting

[79262] protein-1 is required for early embryogenesis in mice. J. Biol. Chem. 276:

[79263] 27745-27748, 2001.

[79264] 13706.

[79265] Waeber, G.; Delplanque, J.; Bonny, C.; Mooser, V.; Stein-mann, M.;

[79266] Widmann, C.; Maillard, A.; Miklossy, J.; Dina, C.; Hani, E. H.; Vionnet,

[79267] N.; Nicod, P.; Boutin, P.; Froguel, P.: The gene MAPK8IP1, encoding

[79268] islet-brain-1, is a candidate for type 2 diabetes. Nature Genet. 24:

[79269] 291-295, 2000.

[79270] 13707.

[79271] Hewett-Emmett, D.; Tashian, R. E.: Functional diversity, conservation,

[79272] and convergence in the evolution of the alpha-, beta-, and gamma-carbonic

[79273] anhydrase gene families. Molec. Phylogenet. Evol. 5: 50-77, 1996.

[79274] 13708.

[79275] Hewett-Emmett, D.; Wiebauer, K.: Personal Communication. Houston,

[79276] Texas 9/1999.

[79277] 13709.

[79278] Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.-B.; Becker,

[79279] K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG repeats in

[79280] cDNAs from human brain. Hum. Genet. 103: 666-673, 1998.

[79281] 13710.

[79282] Lovejoy, D. A.; Hewett-Emmett, D.; Porter, C. A.; Cepoi, D.; Sheffield,

[79283] A.; Vale, W. W.; Tashian, R. E.: Evolutionarily conserved, 'acatalytic'

[79284] carbonic anhydrase-related protein XI contains a se-

[79285] quence motif present in the neuropeptide sauvagine: the human CA-RP XI gene (CA11) is embedded

[79286] between the secretor gene cluster and the DBP gene at 19q13.3. Genomics 54:

[79287] 484-493, 1998.

[79288] 13711.

[79289] Bellingham, J.; Greogory-Evans, K.; Gregory-Evans, C. Y.: Sequence

[79290] and tissue expression of a human novel carbonic anhy-

[79291] drase-related protein, CARP-2, mapping to chromosome 19q13.3. Biochem. Biophys.

[79292] Res. Comm. 253: 364-367, 1998.

[79293] 13712.

[79294] Fujikawa-Adachi, K.; Nishimori, I.; Taguchi, T.; Yuri, K.; Onishi,

[79295] S.: cDNA sequence, mRNA expression, and chromosomal localization

[79296] of human carbonic anhydrase-related protein, CA-RP XI. Biochim. Biophys.

[79297] Acta 1431: 518-524, 1999.

[79298] 13713.

[79299] de Veer, M. J.; Sim, H.; Whisstock, J. C.; Devenish, R. J.; Ralph,

[79300] S. J.: IFI60/ISG60/IFIT4, a new member of the human IFI54/IFIT2 family

[79301] of interferon-stimulated genes. Genomics 54: 267-277, 1998.

[79302] 13714.

[79303] Yu, M.; Tong, J.-H.; Mao, M.; Kan, L.-X.; Liu, M.-M.; Sun, Y.-W.;

[79304] Fu, G.; Jing, Y.-K.; Yu, L.; Lepaslier, D.; Lanotte, M.; Wang, Z.-Y.;

[79305] Chen, Z.; Waxman, S.; Wang, Y.-X.; Tan, J.-Z.; Chen, S.-J.: Cloning

[79306] of a gene (RIG-G) associated with retinoic acid-induced differentiation

[79307] of acute promyelocytic leukemia cells and representing a new member

[79308] of a family of interferon-stimulated genes. Proc. Nat.

Acad. Sci. 94:
[79309] 7406-7411, 1997.

[79310] 13715.

[79311] Zhu, H.; Cong, J.-P.; Shenk, T.: Use of differential display analysis
[79312] to assess the effect of human cytomegalovirus infection on the accumulation
[79313] of cellular RNAs: induction of interferon-responsive RNAs. Proc.
[79314] Nat. Acad. Sci. 94: 13985-13990, 1997.

[79315] 13716.

[79316] Devalia, V.; Carter, K.; Walker, A. P.; Perkins, S. J.; Wor-wood,
[79317] M.; May, A.; Dooley, J. S.: Autosomal dominant reticuloen-dothelial
[79318] iron overload associated with a 3-base pair deletion in the ferroportin
[79319] 1 gene (SLC11A3). Blood 100: 695-697, 2002.

[79320] 13717.

[79321] Kido, S.; Hiraoka, Y.; Ogawa, M.; Sakai, Y.; Yoshimura, Y.; Aiso,

[79322] S.: Cloning and characterization of mouse mSox13 cDNA. Gene 208:

[79323] 201-206, 1998.

[79324] 13718.

[79325] Rabin, D. U.; Pleasic, S. M.; Palmer-Crocker, R.; Shapiro, J. A.

[79326] : Cloning and expression of IDDM-specific human autoantigens. Diabetes 41:

[79327] 183-186, 1992.

[79328] 13719.

[79329] Roose, J.; Korver, W.; de Boer, R.; Kuipers, J.; Hurenkamp, J.;

[79330] Clevers, H.: The Sox-13 gene: structure, promoter characterization,

[79331] and chromosomal localization. Genomics 57: 301-305, 1999.

[79332] 13720.

[79333] Roose, J.; Korver, W.; Oving, E.; Wilson, A.; Wagenaar, G.; Markman,

[79334] M.; Lamers, W.; Clevers, H.: High expression of the HMG box factor

[79335] Sox-13 in arterial walls during embryonic development. Nucleic Acids
[79336] Res. 26: 469-476, 1998.

[79337] 13721.

[79338] Abrink, M.; Aveskogh, M.; Hellman, L.: Isolation of cDNA clones
[79339] for 42 different Kruppel-related zinc finger proteins expressed in
[79340] the human monoblast cell line U-937. DNA Cell Biol. 14: 125-136,
[79341] 1995.

[79342] 13722.

[79343] Holmes, D. I. R.; Wahab, N. A.; Mason, R. M.: Cloning and characterization
[79344] of ZNF236, a glucose-regulated Kruppel-like zinc-finger gene mapping
[79345] to human chromosome 18q22-q23. Genomics 60: 105-109, 1999.

[79346] 13723.

[79347] Hirai, H.; Tanaka, K.; Takano, S.; Ichimasa, M.; Nakamura, M.;

[79348] Nagata, K.: Cutting edge: agonistic effect of indomethacin on a prostaglandin

[79349] D2 receptor, CRTH2. J. Immun. 168: 981-985, 2002.

[79350] 13724.

[79351] Nagata, K.; Tanaka, K.; Ogawa, K.; Kemmotsu, K.; Imai, Yoshie,

[79352] O.; Abe, H.; Tada, K.; Nakamura, M.; Sugamura, K.; Takano, S.: Selective

[79353] expression of a novel surface molecule by human Th2 cells in vivo. J.

[79354] Immun. 162: 1278-1286, 1999.

[79355] 13725.

[79356] Meng, X.; Lu, X.; Morris, C. A.; Keating, M. T.: A novel hu-man

[79357] gene FKBP6 is deleted in Williams syndrome. Genomics 52: 130-137,

[79358] 1998.

[79359] 13726.

[79360] Martin, S.; Richards, A. J.; Yates, J. R. W.; Scott, J. D.; Pope,

[79361] M.; Snead, M. P.: Stickler syndrome: further mutations in COL11A1

[79362] and evidence for additional locus heterogeneity. Europ. J. Hum. Genet. 7:

[79363] 807-814, 1999.

[79364] 13727.

[79365] Sirko-Osadsa, D. A.; Zlotogora, J.; Tiller, G. E.; Knowlton, R.

[79366] G.; Warman, M. L.: A third Stickler syndrome locus is linked to COL11A1,

[79367] the gene encoding the alpha-1 subunit of collagen XI. (Abstract) Am.

[79368] J. Hum. Genet. 59 (suppl.): A17, 1996.

[79369] 13728.

[79370] Grundemann, D.; Schechinger, B.; Rappold, G. A.; Schomig, E.:

[79371] Molecular identification of the corticosterone-sensitive extraneuronal

[79372] catecholamine transporter. Nature Neurosci. 1: 349-351, 1998.

[79373] 13729.

[79374] Boettger, T.; Hubner, C. A.; Maler, H.; Rust, M. B.; Beck, F. X.;

[79375] Jentsch, T. J.: Deafness and renal tubular acidosis in mice lacking
[79376] the K-Cl co-transporter Kcc4. Nature 416: 874-878, 2002.
[79377] 13730.

[79378] Prakash, S. K.; Van den Veyver, I. B.; Franco, B.; Volta, M.; Ballabio,
[79379] A.; Zoghbi, H. Y.: Characterization of a novel chromo do-main gene
[79380] in Xp22.3 with homology to Drosophila msl-3. Genomics 59: 77-84,
[79381] 1999.

[79382] 13731.

[79383] Kotake, K.; Ozaki, N.; Mizuta, M.; Sekiya, S.; Inagaki, N.; Seino,
[79384] S.: Noc2, a putative zinc finger protein involved in exocy-tosis in
[79385] endocrine cells. J. Biol. Chem. 272: 29407-29410, 1997.

[79386] 13732.

[79387] Smith, J. S.; Tachibana, I.; Allen, C.; Chiappa, S. A.; Lee, H.

[79388] K.; McIver, B.; Jenkins, R. B.; Raffel, C.: Cloning of a human

ortholog

[79389] (RPH3AL) of (RNO)Rph3al from a candidate 17p13.3 medulloblastoma tumor

[79390] suppressor locus. Genomics 59: 97-101, 1999.

[79391] 13733.

[79392] Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayasi,

[79393] Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction

[79394] of the coding sequences of unidentified human genes. I. The coding

[79395] sequences of 40 new genes (KIAA0001-KIAA0040) de- duced by analysis

[79396] of randomly sampled cDNA clones from human immature myeloid cell line

[79397] KG-1. DNA Res. 1: 27-35, 1994.

[79398] 13734.

[79399] Kimura, M.; Okano, Y.: Identification and assignment of the human

[79400] NIMA-related protein kinase 7 gene (NEK7) to human chromosome 1q31.3. Cytogenet.

[79401] Cell Genet. 94: 33-38, 2001.

[79402] 13735.

[79403] Li, M. Z.; Yu, L.; Liu, Q.; Chu, J. Y.; Zhao, S. Y.: Assignment

[79404] of NEK6, a NIMA-related gene, to human chromosome 9q33.3-q34.11 by

[79405] radiation hybrid mapping. Cytogenet. Cell Genet. 87: 271-272, 1999.

[79406] 13736.

[79407] Suzuki, T.; Nishiyama, K.; Yamamoto, A.; Inazawa, J.; Iwaki, T.;

[79408] Yamada, T.; Kanazawa, I.; Sakaki, Y.: Molecular cloning of a novel

[79409] apoptosis-related gene, human Nap1 (NCKAP1), and its possible relation

[79410] to Alzheimer disease. Genomics 63: 246-254, 2000.

[79411] 13737.

[79412] Ruegsegger, U.; Blank, D.; Keller, W.: Human pre-mRNA cleavage

[79413] factor I-m is related to spliceosomal SR proteins and can be reconstituted

[79414] in vitro from recombinant subunits. Molec. Cell 1: 243-253, 1998.

[79415] 13738.

[79416] Hall, S. L.; Padgett, R. A.: Conserved sequences in a class of

[79417] rare eukaryotic nuclear introns with non-consensus splice sites. J.

[79418] Molec. Biol. 239: 357-365, 1994.

[79419] 13739.

[79420] Horikawa, T.; Araki, K.; Fukai, K.; Ueda, M.; Ueda, T.; Ito, S.;

[79421] Ichihashi, M.: Heterozygous HPS1 mutations in a case of Hermansky-Pudlak

[79422] syndrome with giant melanosomes. Brit. J. Derm. 143: 635-640, 2000.

[79423] 13740.

[79424] Shotelersuk, V.; Hazelwood, S.; Larson, D.; Iwata, F.; Kaiser-Kupfer,

[79425] M. I.; Kuehl, E.; Bernardini, I.; Gahl, W. A.: Three new mu-tations

[79426] in a gene causing Hermansky-Pudlak syndrome: clinical correlations. Molec.

[79427] Genet. Metab. 64: 99-107, 1998.

[79428] 13741.

[79429] Wildenberg, S. C.; Fryer, J. P.; Gardner, J. M.; Oetting, W. S.;

[79430] Brilliant, M. H.; King, R. A.: Identification of a novel tran-

script

[79431] produced by the gene responsible for the Hermansky-Pud-

lak syndrome

[79432] in Puerto Rico. J. Invest. Derm. 110: 777-781, 1998.

[79433] 13742.

[79434] Bunn, R. C.; Jensen, M. A.; Reed, B. C.: Protein interactions

[79435] with the glucose transporter binding protein GLUT1CBP that provide

[79436] a link between GLUT1 and the cytoskeleton. Molec. Biol. Cell 10:

[79437] 819-832, 1999.

[79438] 13743.

[79439] De Vries, L.; Lou, X.; Zhao, G.; Zheng, B.; Farquhar, M. G.: GIPC,

[79440] a PDZ domain containing protein, interacts specifically with the C

[79441] terminus of RGS-GAIP. Proc. Nat. Acad. Sci. 95:

12340-12345, 1998.

[79442] 13744.

[79443] Von Kap-Herr, C.; Kandala, G.; Mann, S. S.; Hart, T. C.; Pettenati,

[79444] M. J.; Setaluri, V.: Assignment of PDZ domain-containing protein

[79445] GIPC gene (C19orf3) to human chromosome band 19p13.1 by in situ hybridization

[79446] and radiation hybrid mapping. Cytogenet. Cell Genet. 89: 234-235,

[79447] 2000.

[79448] 13745.

[79449] Alessi, D. R.; James, S. R.; Downes, C. P.; Holmes, A. B.; Gaffney,

[79450] P. R. J.; Reese, C. B.; Cohen, P.: Characterization of a 3-phosphoinositide-dependent

[79451] protein kinase which phosphorylates and activates protein kinase B-alpha. Curr.

[79452] Biol. 7: 261-269, 1997.

[79453] 13746.

[79454] Adams, J. C.; Seed, B.; Lawler, J.: Muskelin, a novel intra-

cellular

[79455] mediator of cell adhesive and cytoskeletal responses to thrombospondin-1. EMBO

[79456] J. 17: 4964-4974, 1998.

[79457] 13747.

[79458] Adams, J. C.; Zhang, L.: cDNA cloning of human muskelin and localisation

[79459] of the muskelin (MKLN1) gene to human chromosome 7q32 and mouse chromosome

[79460] 6B1/B2 by physical mapping and FISH. Cytogenet. Cell Genet. 87:

[79461] 19-21, 1999.

[79462] 13748.

[79463] Moynihan, T. P.; Ardley, H. C.; Nuber, U.; Rose, S. A.; Jones,

[79464] P. F.; Markham, A. F.; Scheffner, M.; Robinson, P. A.: The ubiquitin-conjugating

[79465] enzymes UbcH7 and UbcH8 interact with RING finger/IBR motif-containing

[79466] domains of HHARI and H7-AP1. J. Biol. Chem. 274: 30963-30968, 1999.

[79467] 13749.

[79468] Tan, N. G. S.; Ardley, H. C.; Rose, S. A.; Leek, J. P.; Markham,
[79469] A. F.; Robinson, P. A.: Characterisation of the human and mouse orthologues
[79470] of the Drosophila ariadne gene. Cytogenet. Cell Genet. 90: 242-245,
[79471] 2000.

[79472] 13750.

[79473] Janssen, J. W. G.; Imoto, I.; Inoue, J.; Shimada, Y.; Ueda, M.;
[79474] Imamura, M.; Bartram, C. R.; Inazawa, J.: MYEOV, a gene at 11q13,
[79475] is coamplified with CCND1, but epigenetically inactivated in a subset
[79476] of esophageal squamous cell carcinomas. J. Hum. Genet. 47: 460-464,
[79477] 2002.

[79478] 13751.

[79479] Janssen, J. W. G.; Vaandrager, J.-W.; Heuser, T.; Jauch, A.; Kluin,
[79480] P. M.; Geelen, E.; Bergsagel, P. L.; Kuehl, W. M.; Drexler, H. G.;

[79481] Otsuki, T.; Bartram, C. R.; Schuuring, E.: Concurrent activation
[79482] of a novel putative transforming gene, myeov, and cyclin D1 in a subset
[79483] of multiple myeloma cell lines with t(11;14)(q13;q32). Blood 95:
[79484] 2691-2698, 2000.

[79485] 13752.

[79486] Di Cunto, F.; Calautti, E.; Hsiao, J.; Ong, L.; Topley, G.; Turco,
[79487] E.; Dotto, G. P.: Citron Rho-interacting kinase, a novel tissue-specific
[79488] ser/thr kinase encompassing the Rho-Rac-binding protein citron. J.
[79489] Biol. Chem. 273: 29706-29711, 1998.

[79490] 13753.

[79491] Di Cunto, F.; Imarisio, S.; Hirsch, E.; Broccoli, V.; Bulfone,
[79492] A.; Migheli, A.; Atzori, C.; Turco, E.; Triolo, R.; Dotto, G. P.;
[79493] Silengo, L.; Altruda, F.: Defective neurogenesis in citron kinase
[79494] knockout mice by altered cytokinesis and massive apop-

tosis. Neuron 28:

[79495] 115-127, 2000.

[79496] 13754.

[79497] Madaule, P.; Furuyashiki, T.; Reid, T.; Ishizaki, T.; Watanabe,

[79498] G.; Morii, N.; Narumiya, S.: A novel partner for the GTP-bound forms
[79499] of rho and rac. FEBS Lett. 377: 243-248, 1995.

[79500] 13755.

[79501] Ishizaki, J.; Suzuki, N.; Higashino, K.; Yokota, Y.; Ono, T.; Kawamoto,
[79502] K.; Fujii, N.; Arita, H.; Hanasaki, K.: Cloning and character-ization
[79503] of novel mouse and human secretory phospholipase A(2)s. J. Biol.
[79504] Chem. 274: 24973-24979, 1999.

[79505] 13756.

[79506] Baudat, F.; Manova, K.; Yuen, J. P.; Jasin, M.; Keeney, S.: Chromosome
[79507] synapsis defects and sexually dimorphic meiotic progres-sion in mice
[79508] lacking Spo11. Molec. Cell 6: 989-998, 2000.

[79509] 13757.

[79510] Romanienko, P. J.; Camerini-Otero, R. D.: The mouse Spo11 gene
[79511] is required for meiotic chromosome synapsis. Molec. Cell

6: 975-987,

[79512] 2000.

[79513] 13758.

[79514] Romanienko, P. J.; Camerini-Otero, R. D.: Cloning, charac-
terization,

[79515] and localization of mouse and human SPO11. Genomics
61: 156-169,

[79516] 1999.

[79517] 13759.

[79518] Shannon, M.; Richardson, L.; Christian, A.; Handel, M. A.;
Thelen,

[79519] M. P.: Differential gene expression of mammalian
SPO11/TOP6A homologs

[79520] during meiosis. FEBS Lett. 462: 329-334, 1999.

[79521] 13760.

[79522] Lopez-Coviella, I.; Berse, B.; Krauss, R.; Thies, R. S.; Blusz-
tajn,

[79523] J. K.: Induction and maintenance of the neuronal choliner-
gic phenotype

[79524] in the central nervous system by BMP-9. Science 289:
313-316, 2000.

[79525] 13761.

[79526] Burn, T. C.; Connors, T. D.; Van Raay, T. J.; Dackowski, W. R.;

[79527] Millholland, J. M.; Klinger, K. W.; Landes, G. M.: Generation of

[79528] a transcriptional map for a 700-kb region surrounding the polycystic

[79529] kidney disease type 1 (PKD1) and tuberous sclerosis type 2 (TSC2)

[79530] disease genes on human chromosome 16p13.3. Genome Res. 6: 525-537,

[79531] 1996.

[79532] 13762.

[79533] Pullen, N.; Dennis, P. B.; Andjelkovic, M.; Dufner, A.; Kozma,

[79534] S. C.; Hemmings, B. A.; Thomas, G.: Phosphorylation and activation

[79535] of p70(s6k) by PDK1. Science 279: 707-710, 1998.

[79536] 13763.

[79537] Stephens, L.; Anderson, K.; Stokoe, D.; Erdjument-Bro-

mage, H.;

[79538] Painter, G. F.; Holmes, A. B.; Gaffney, P. R. J.; Reese, C. B.; McCormick,
[79539] F.; Tempst, P.; Coadwell, J.; Hawkins, P. T.: Protein kinase B kinases
[79540] that mediate phosphatidylinositol 3,4,5-triphosphate-dependent activation
[79541] of protein kinase B. Science 279: 710-714, 1998.

[79542] 13764.

[79543] Uebele, V. N.; Lagrutta, A.; Wade, T.; Figueroa, D. J.; Liu, Y.;
[79544] McKenna, E.; Austin, C. P.; Bennett, P. B.; Swanson, R.: Cloning
[79545] and functional expression of two families of beta-subunits of the
[79546] large conductance calcium-activated K(+) channel. J. Biol. Chem. 275:
[79547] 23211-23218, 2000.

[79548] 13765.

[79549] Kouroku, Y.; Soyama, A.; Fujita, E.; Urase, K.; Tsukahara, T.;
[79550] Momoi, T.: RA70 is a src kinase-associated protein ex-pressed ubiquitously. Biochem.

[79551] Biophys. Res. Comm. 252: 738-742, 1998.

[79552] 13766.

[79553] Liu, J.; Kang, H.; Raab, M.; da Silva, A. J.; Kraeft, S.-K.; Rudd,

[79554] C. E.: FYB (FYN binding protein) serves as a binding part-ner for

[79555] lymphoid protein and FYN kinase substrate SKAP55 and a SKAP55-related

[79556] protein in T cells. Proc. Nat. Acad. Sci. 95: 8779-8784, 1998.

[79557] 13767.

[79558] Marie-Cardine, A.; Verhagen, A. M.; Eckerskorn, C.; Schraven, B.

[79559] : SKAP-HOM, a novel adaptor protein homologous to the FYN-associated

[79560] protein SKAP55. FEBS Lett. 435: 55-60, 1998.

[79561] 13768.

[79562] Chai, J.; Du, C.; Wu, J.-W.; Kyin, S.; Wang, X.; Shi, Y.: Structural

[79563] and biochemical basis of apoptotic activation by Smac/ DIABLO. Nature 406:

[79564] 855-862, 2000.

[79565] 13769.

[79566] Du, C.; Fang, M.; Li, Y.; Li, L.; Wang, X.: Smac, a mitochon-
drial
[79567] protein that promotes cytochrome c-dependent caspase
activation by
[79568] eliminating IAP inhibition. Cell 102: 33-42, 2000.

[79569] 13770.

[79570] Okada, H.; Suh, W.-K.; Jin, J.; Woo, M.; Du, C.; Elia, A.;
Duncan,
[79571] G. S.; Wakeham, A.; Itie, A.; Lowe, S. W.; Wang, X.; Mak, T.
W.:
[79572] Generation and characterization of Smac/DIA-
BLO-deficient mice. Molec.
[79573] Cell. Biol. 22: 3509-3517, 2002.

[79574] 13771.

[79575] Scott, A. F.: Personal Communication. Baltimore, Md.
8/18/2000.
[79576] 13772.

[79577] Verhagen, A. M.; Ekert, P. G.; Pakusch, M.; Silke, J.; Con-
nolly,

[79578] L. M.; Reid, G. E.; Moritz, R. L.; Simpson, R. J.; Vaux, D. L.: Identification
[79579] of DIABLO, a mammalian protein that promotes apoptosis by binding
[79580] to and antagonizing IAP proteins. Cell 102: 43–53, 2000.

[79581] 13773.

[79582] Brown, M. L.; Ramprasad, M. P.; Umeda, P. K.; Tanaka, A.; Kobayashi,
[79583] Y.; Watanabe, T.; Shimoyamada, H.; Kuo, W.-L.; Li, R.; Song, R.; Bradley,
[79584] W. A.; Gianturco, S. H.: A macrophage receptor for apolipoprotein
[79585] B48: cloning, expression, and atherosclerosis. Proc. Nat. Acad. Sci. 97:
[79586] 7488–7493, 2000.

[79587] 13774.

[79588] Behrens, R.; Nolting, A.; Reimann, F.; Schwarz, M.; Wald-schutz,
[79589] R.; Pongs, O.: hKCNMB3 and hKCNMB4, cloning and char-acterization
[79590] of two members of the large-conductance calcium-acti-vated potassium

[79591] channel beta subunit family. FEBS Lett. 474: 99-106, 2000.

[79592] 13775.

[79593] Brenner, R.; Jegla, T. J.; Wickenden, A.; Liu, Y.; Aldrich, R.

[79594] W.: Cloning and functional characterization of novel large conductance

[79595] calcium-activated potassium channel beta subunits, hKC-NMB3 and hKCNMB4. J.

[79596] Biol. Chem. 275: 6453-6461, 2000.

[79597] 13776.

[79598] Riazi, M. A.; Brinkman-Mills, P.; Johnson, A.; Naylor, S. L.; Minoshima,

[79599] S.; Shimizu, N.; Baldini, A.; McDermid, H. E.: Identification of

[79600] a putative regulatory subunit of a calcium-activated potassium channel

[79601] in the dup(3q) syndrome region and a related sequence on 22q11.2. Genomics 62:

[79602] 90-94, 1999.

[79603] 13777.

[79604] Hemmi, H.; Takeuchi, O.; Kawai, T.; Kaisho, T.; Sato, S.;

Sanjo,
[79605] H.; Matsumoto, M.; Hoshino, K.; Wagner, H.; Takeda, K.; Akira, S.
[79606] : A toll-like receptor recognizes bacterial DNA. Nature 408: 740-745,
[79607] 2000.

[79608] 13778.

[79609] Leadbetter, E. A.; Rifkin, I. R.; Hohlbaum, A. M.; Beaudette, B.
[79610] C.; Shlomchik, M. J.; Marshak-Rothstein, A.: Chromatin-IgG complexes
[79611] activate B cells by dual engagement of IgM and Toll-like receptors. Nature 416:
[79612] 603-607, 2002.

[79613] 13779.

[79614] Takeshita, F.; Leifer, C. A.; Gursel, I.; Ishii, K. J.; Takeshita,

[79615] S.; Gursel, M.; Klinman, D. M.: Cutting edge: role of Toll-like receptor
[79616] 9 in CpG DNA-induced activation of human cells. J. Immun. 167: 3555-3558,
[79617] 2001.

[79618] 13780.

[79619] Verthelyi, D.; Ishii, K. J.; Gursel, M.; Takeshita, F.; Klinman,

[79620] D. K.: Human peripheral blood cells differentially recog-
nize and

[79621] respond to two distinct CpG motifs. J. Immun. 166:
2372-2377, 2001.

[79622] 13781.

[79623] Bagrodia, S.; Taylor, S. J.; Jordon, K. A.; Van Aelst, L.; Ceri-
one,

[79624] R. A.: A novel regulator of p21-activated kinases. J. Biol.
Chem. 273:

[79625] 23633-23636, 1998.

[79626] 13782.

[79627] Oh, W. K.; Yoo, J. C.; Jo, D.; Song, Y. H.; Kim, M. G.; Park,
D.

[79628] : Cloning of a SH3 domain-containing proline-rich pro-
tein, p85SPR,

[79629] and its localization in focal adhesion. Biochem. Biophys.
Res. Commun. 235:

[79630] 794-798, 1997.

[79631] 13783.

[79632] Ehrmann, D. A.; Schwarz, P. E. H.; Hara, M.; Tang, X.; Horikawa,

[79633] Y.; Imperial, J.; Bell, G. I.; Cox, N. J.: Relationship of calpain-10

[79634] genotype to phenotypic features of polycystic ovary syndrome. J.

[79635] Clin. Endocr. Metab. 87: 1669-1673, 2002.

[79636] 13784.

[79637] Elbein, S. C.; Chu, W.; Ren, Q.; Hemphill, C.; Schay, J.; Cox,

[79638] N. J.; Hanis, C. L.; Hasstedt, S. J.: Role of calpain-10 gene variants

[79639] in familial type 2 diabetes in Caucasians. J. Clin. Endocr. Metab. 87:

[79640] 650-654, 2002.

[79641] 13785.

[79642] Fullerton, S. M.; Bartoszewicz, A.; Ybazeta, G.; Horikawa, Y.;

[79643] Bell, G. I.; Kidd, K. K.; Cox, N. J.; Hudson, R. R.; Di Rienzo, A.

[79644] : Geographic and haplotype structure of candidate type 2 diabetes-susceptibility

[79645] variants at the calpain-10 locus. Am. J. Hum. Genet. 70: 1096-1106,
[79646] 2002.

[79647] 13786.

[79648] Tsai, H.-J.; Sun, G.; Weeks, D. E.; Kaushal, R.; Wolujewicz, M.;
[79649] McGarvey, S. T.; Tufa, J.; Viali, S.; Deka, R.: Type 2 diabetes and
[79650] three calpain-10 gene polymorphisms in Samoans: no evidence of association. Am.
[79651] J. Hum. Genet. 69: 1236-1244, 2001.

[79652] 13787.

[79653] Weiss, K. M.; Terwilliger, J. D.: How many diseases does it take
[79654] to map a gene with SNPs? Nature Genet. 26: 151-157, 2000.
[79655] 13788.

[79656] Hanis, C. L.; Boerwinkle, E.; Chakraborty, R.; Ellsworth, D. L.;
[79657] Concannon, P.; Stirling, B.; Morrison, V. A.; Wapelhorst, B.; Spielman,

[79658] R. S.; Gogolin-Ewens, K. J.; Shephard, J. M.; Williams, S. R.; and
[79659] 21 others: A genome-wide search for human non-insulin-dependent
[79660] (type 2) diabetes genes reveals a major susceptibility locus on chromosome
[79661] 2. Nature Genet. 13: 161-166, 1996.

[79662] 13789.

[79663] Aung, T.; Ocaka, L.; Ebenezer, N. D.; Morris, A. G.; Brice, G.;
[79664] Child, A. H.; Hitchings, R. A.; Lehmann, O. J.; Bhattacharya, S. S.
[79665] : Investigating the association between OPA1 polymorphisms and glaucoma:
[79666] comparison between normal tension and high tension primary open angle
[79667] glaucoma. Hum. Genet. 110: 513-514, 2002.

[79668] 13790.

[79669] Aung, T.; Ocaka, L.; Ebenezer, N. D.; Morris, A. G.; Krawczak,
[79670] M.; Thiselton, D. L.; Alexander, C.; Votruba, M.; Brice, G.; Child,

[79671] A. H.; Francis, P. J.; Hitchings, R. A.; Lehmann, O. J.; Bhattacharya,

[79672] S. S.: A major marker for normal tension glaucoma: association with

[79673] polymorphisms in the OPA1 gene. Hum. Genet. 110: 52-56, 2002.

[79674] 13791.

[79675] Liu, Z.; Chen, Y.; Mo, R.; Hui, C.; Cheng, J.-F.; Mohandas, N.;

[79676] Huang, C.-H.: Characterization of human RhCG and mouse Rhcg as novel

[79677] nonerythroid Rh glycoprotein homologues predominantly expressed in

[79678] kidney and testis. J. Biol. Chem. 275: 25641-25651, 2000.

[79679] 13792.

[79680] Kabarowski, J. H. S.; Zhu, K.; Le, L. Q.; Witte, O. N.; Xu, Y.

[79681] : Lysophosphatidylcholine as a ligand for the immunoregulatory receptor

[79682] G2A. Science 293: 702-705, 2001.

[79683] 13793.

[79684] Wang, A.; Yang, H.-C.; Friedman, P.; Johnson, C. A.; Dennis, E.

[79685] A.: A specific human lysophospholipase: cDNA cloning, tissue distribution

[79686] and kinetic characterization. Biochim. Biophys. Acta 1437: 157-169,

[79687] 1999.

[79688] 13794.

[79689] Barker, R. L.; Gleich, G. J.; Pease, L. R.: Acidic precursor revealed

[79690] in human eosinophil granule major basic protein cDNA. J. Exp. Med. 168:

[79691] 1493-1498, 1988.

[79692] 13795.

[79693] Li, M.-S.; Sun, L.; Satoh, T.; Fisher, L. M.; Spry, C. J. F.:

[79694] Human eosinophil major basic protein, a mediator of allergic inflammation,

[79695] is expressed by alternative splicing from two promoters. Biochem.

[79696] J. 305: 921-927, 1995.

[79697] 13796.

[79698] McGrogan, M.; Simonsen, C.; Scott, R.; Griffith, J.; Ellis, N.;

[79699] Kennedy, J.; Campanelli, D.; Nathan, C.; Gabay, J.: Isolation of

[79700] a complementary DNA clone encoding a precursor to human eosinophil

[79701] major basic protein. J. Exp. Med. 168: 2295-2308, 1988.

[79702] 13797.

[79703] Plager, D. A.; Weiler, D. A.; Loegering, D. A.; Johnson, W. B.;

[79704] Haley, L.; Eddy, R. L.; Shows, T. B.; Gleich, G. J.: Comparative

[79705] structure, proximal promoter elements, and chromosome location of

[79706] the human eosinophil major basic protein genes. Genomics 71: 271-281,

[79707] 2001.

[79708] 13798.

[79709] Wasmoen, T. L.; Bell, M. P.; Loegering, D. A.; Gleich, G. J.; Prendergast,

[79710] F. G.; McKean, D. J.: Biochemical and amino acid sequence analysis

[79711] of human eosinophil granule major basic protein. J. Biol. Chem. 263:
[79712] 12559-12563, 1988.

[79713] 13799.

[79714] Weller, P. F.; Ackerman, S. J.; Smith, J. A.: Eosinophil gran-ule
[79715] proteins: major basic protein is distinct from the smaller subunit
[79716] of eosinophil peroxidase. J. Leukoc. Biol. 43: 1-4, 1988.

[79717] 13800.

[79718] Yoshimatsu, K.; Ohya, Y.; Shikata, Y.; Seto, T.; Hasegawa, Y.;
[79719] Tanaka, I.; Kawamura, T.; Kitoh, K.; Toyoshima, S.; Osawa, T.: Purification
[79720] and cDNA cloning of a novel factor produced by a human T-cell hybridoma:
[79721] sequence homology with animal lectins. Molec. Immun. 29: 537-546,
[79722] 1992.

[79723] 13801.

[79724] Khaleghpour, K.; Svitkin, Y. V.; Craig, A. W.; DeMaria, C.

T.;

[79725] Deo, R. C.; Burley, S. K.; Sonenberg, N.: Translational re-
pression

[79726] by a novel partner of human poly(A) binding protein,
Paip2. Molec.

[79727] Cell 7: 205-216, 2001.

[79728] 13802.

[79729] Scott, A. F.: Personal Communication. Baltimore, Md.
3/13/2001.

[79730] 13803.

[79731] Drane, P.; Barel, M.; Balbo, M.; Frade, R.: Identification of

[79732] RB18A, a 205 kDa new p53 regulatory protein which
shares antigenic

[79733] and functional properties with p53. Oncogene 15:
3013-3024, 1997.

[79734] 13804.

[79735] Mansharamani, M.; Hewetson, A.; Chilton, B. S.: Cloning
and characterization

[79736] of an atypical type IV P-type ATPase that binds to the
RING motif

[79737] of RUSH transcription factors. J. Biol. Chem. 276:

3641-3649, 2001.

[79738] 13805.

[79739] Sun, X. L.; Li, D.; Fang, J.; Noyes, I.; Casto, B.; Theil, K.;

[79740] Shuler, C.; Milo, G. E.: Changes in levels of normal ML-1 gene transcripts

[79741] associated with the conversion of human nontumorigenic to tumorigenic

[79742] phenotypes. Gene Expr. 8: 129-139, 1999.

[79743] 13806.

[79744] Lesage, F.; Terrenoire, C.; Romey, G.; Lazdunski, M.: Human TREK2,

[79745] a 2P domain mechano-sensitive K+ channel with multiple regulations

[79746] by polyunsaturated fatty acids, lysophospholipids, and Gs, Gi, and

[79747] Gq protein-coupled receptors. J. Biol. Chem. 275: 28398-28405, 2000.

[79748] 13807.

[79749] Nakayama, Y.; Weissman, S. M.; Bothwell, A. L. M.: BX-MAS1 identifies

[79750] a cluster of homologous genes differentially expressed in

B cells. Biochem.

[79751] Biophys. Res. Commun. 285: 830-837, 2001.

[79752] 13808.

[79753] Bashirova, A. A.; Geijtenbeek, T. B. H.; van Duijnhoven, G. C.

[79754] F.; van Vliet, S. J.; Eilering, J. B. G.; Martin, M. P.; Wu, L.; Martin,

[79755] T. D.; Viebig, N.; Knolle, P. A.; KewalRamani, V. N.; van Kooyk, Y.;

[79756] Carrington, M.: A dendritic cell-specific intercellular adhe-sion

[79757] molecule 3-grabbing nonintegrin (DC-SIGN)-related pro-tein is highly

[79758] expressed on human liver sinusoidal endothelial cells and promotes

[79759] HIV-1 infection. J. Exp. Med. 193: 671-678, 2001.

[79760] 13809.

[79761] Maguchi, M.; Nishida, W.; Kohara, K.; Kuwano, A.; Kondo, I.; Hiwada,

[79762] K.: Molecular cloning and gene mapping of human basic and acidic

[79763] calponins. Biochem. Biophys. Res. Commun. 217:

238-244, 1995.

[79764] 13810.

[79765] Miano, J. M.; Krahe, R.; Garcia, E.; Elliott, J. M.; Olson, E.

[79766] N.: Expression, genomic structure and high resolution mapping to

[79767] 19p13.2 of the human smooth muscle cell calponin gene. Gene 197:

[79768] 215-224, 1997.

[79769] 13811.

[79770] Miano, J. M.; Thomas, S.; Disteche, C. M.: Expression and chromosomal

[79771] mapping of the mouse smooth muscle calponin gene. Mammalian Genome 12:

[79772] 187-191, 2001.

[79773] 13812.

[79774] Strasser, P.; Gimona, M.; Moessler, H.; Herzog, M.; Small, J. V.

[79775] : Mammalian calponin: identification and expression of genetic variants. FEBS

[79776] Lett. 330: 13-18, 1993.

[79777] 13813.

[79778] Bogerd, H. P.; Fridell, R. A.; Madore, S.; Cullen, B. R.: Identification
[79779] of a novel cellular cofactor for the Rev/Rex class of retroviral regulatory
[79780] proteins. Cell 82: 485-494, 1995.

[79781] 13814.

[79782] Fritz, C. C.; Zapp, M. L.; Green, M. R.: A human nucleoporin-like
[79783] protein that specifically interacts with HIV Rev. Nature 376: 530-533,
[79784] 1995.

[79785] 13815.

[79786] Jones, T.; Sheer, D.; Bevec, D.; Kappel, B.; Hauber, J.; Steinkasserer,
[79787] A.: The human HIV-1 Rev binding-protein hRIP/Rab (HRB) maps to chromosome
[79788] 2q36. Genomics 40: 198-199, 1997.

[79789] 13816.

[79790] Kang-Decker, N.; Mantchev, G. T.; Juneja, S. C.; McNiven, M. A.;
[79791] van Deursen, J. M. A.: Lack of acrosome formation in Hrb-

deficient

[79792] mice. Science 294: 1531-1533, 2001.

[79793] 13817.

[79794] Salcini, A. E.; Confalonieri, S.; Doria, M.; Santolini, E.; Tassi,

[79795] E.; Minenkova, O.; Cesareni, G.; Pelicci, P. G.; Di Fiore, P. P.:

[79796] Binding specificity and in vivo targets of the EH domain, a novel

[79797] protein-protein interaction module. Genes Dev. 11: 2239-2249, 1997.

[79798] 13818.

[79799] Bullrich, F.; Druck, T.; Kunapuli, P.; Gomez, J.; Gripp, K. W.;

[79800] Schlegelberger, B.; Lasota, J.; Aronson, M.; Cannizzaro, L. A.; Huebner,

[79801] K.; Benovic, J. L.: Chromosomal mapping of the genes GPRK5 and GPRK6

[79802] encoding G protein-coupled receptor kinases GRK5 and GRK6. Cytogenet.

[79803] Cell Genet. 70: 250-254, 1995.

[79804] 13819.

[79805] Haribabu, B.; Snyderman, R.: Identification of additional members
[79806] of human G-protein-coupled receptor kinase multigene family. Proc.
[79807] Nat. Acad. Sci. 90: 9398-9402, 1993.

[79808] 13820.

[79809] Entwistle, J.; Zhang, S.; Yang, B.; Wong, C.; Li, Q.; Hall, C.

[79810] L.; Jingbo, A.; Mowat, M.; Greenberg, A. H.; Turley, E. A.: Characterization
[79811] of the murine gene encoding the hyaluronan receptor RHAMM. Gene 163:
[79812] 233-238, 1995.

[79813] 13821.

[79814] Hall, C.; Yang, B.; Yang, X.; Zhang, S.; Turley, M.; Samuel, S.;
[79815] Lange, L. A.; Wang, C.; Curpen, G. D.; Savani, R. C.; Greenberg, A.
[79816] H.; Turley, E. A.: Overexpression of the hyaluronan receptor RHAMM
[79817] is transforming and is also required for H-ras transformation. Cell 82:

[79818] 19-28, 1995.

[79819] 13822.

[79820] Hall, C. L.; Wang, C.; Lange, L. A.; Turley, E. A.: Hyaluro-
nan
[79821] and the hyaluronan receptor RHAMM promote focal adhe-
sion turnover
[79822] and transient tyrosine kinase activity. J. Cell Biol. 126:
575-588,
[79823] 1994.

[79824] 13823.

[79825] Hardwick, C.; Hoare, K.; Owens, R.; Hohn, H. P.; Moore, D.;
Cripps,
[79826] V.; Austen, L.; Nance, D. M.; Turley, E. A.: Molecular
cloning of
[79827] a novel hyaluronan receptor that mediates tumor cell
motility. J.
[79828] Cell Biol. 117: 1343-1350, 1992.

[79829] 13824.

[79830] Savani, R. C.; Wang, C.; Yang, B. H.; Zhang, S. W.; Kinsella,
M.
[79831] G.; Wight, T. N.; Stern, R.; Nance, D. M.; Turley, E. A.: Mi-

gration
[79832] of bovine aortic smooth muscle cells after wounding injury: the role
[79833] of hyaluronan and RHAMM. J. Clin. Invest. 95: 1158-1168, 1995.
[79834] 13825.

[79835] Spicer, A. P.; Roller, M. L.; Camper, S. A.; McPherson, J. D.;

[79836] Wasmuth, J. J.; Hakim, S.; Wang, C.; Turley, E. A.; McDonald, J. A.
[79837] : The human and mouse receptors for hyaluronan-mediated motility,
[79838] RHAMM, genes (HMMR) map to human chromosome 5q33.2-qter and mouse
[79839] chromosome 11. Genomics 30: 115-117, 1995.

[79840] 13826.

[79841] Cherel, M.; Sorel, M.; Apiou, F.; Lebeau, B.; Dubois, S.; Jacques,
[79842] Y.; Minvielle, S.: The human interleukin-11 receptor alpha gene (IL11RA):
[79843] genomic organization and chromosome mapping. Genomics 32: 49-53,
[79844] 1996.

[79845] 13827.

[79846] Cherel, M.; Sorel, M.; Lebeau, B.; Dubois, S.; Moreau, J.-F.; Bataille,

[79847] R.; Minvielle, S.; Jacques, Y.: Molecular cloning to two isoforms

[79848] of a receptor for the human hematopoietic cytokine interleukin-11. Blood 86:

[79849] 2534-2540, 1995.

[79850] 13828.

[79851] Neuhaus, H.; Bettenhausen, B.; Bilinski, P.; Simon-Chazottes, D.;

[79852] Guenet, J.-L.; Gossler, A.: Etl2, a novel putative type-I cytokine

[79853] receptor expressed during mouse embryogenesis at high levels in skin

[79854] and cells with skeletogenic potential. Dev. Biol. 166: 531-542,

[79855] 1994.

[79856] 13829.

[79857] Van Leuven, F.; Stas, L.; Hilliker, C.; Miyake, Y.; Bilinski, P.;

[79858] Gossler, A.: Molecular cloning and characterization of the

human

[79859] interleukin-11 receptor alpha-chain gene, IL11RA, located on chromosome

[79860] 9p13. Genomics 31: 65-70, 1996.

[79861] 13830.

[79862] Magrangeas, F.; Pitiot, G.; Dubois, S.; Bragado-Nilsson, E.; Cherel,

[79863] M.; Jobert, S.; Lebeau, B.; Boisteau, O.; Lethe, B.; Mallet, J.; Jacques,

[79864] Y.; Minvielle, S.: Cotranscription and intergenic splicing of human

[79865] galactose-1-phosphate uridylyltransferase and inter-leukin-11 receptor

[79866] alpha-chain genes generate a fusion mRNA in normal cells: implication

[79867] for the production of multidomain proteins during evolu-tion. J. Biol.

[79868] Chem. 273: 16005-16010, 1998.

[79869] 13831.

[79870] Bonnemann, C. G.; Cox, G. F.; Shapiro, F.; Wu, J.-J.; Feener, C.

[79871] A.; Thompson, T. G.; Anthony, D. C.; Eyre, D. R.; Darras, B.

T.; Kunkel,
[79872] L. M.: A mutation in the alpha 3 chain of type IX collagen causes
[79873] autosomal dominant multiple epiphyseal dysplasia with mild myopathy. Proc.
[79874] Nat. Acad. Sci. 97: 1212-1217, 2000.

[79875] 13832.

[79876] Paassilta, P.; Lohiniva, J.; Annunen, S.; Bonaventure, J.; Le Merrer,
[79877] M.; Pai, L.; Ala-Kokko, L.: COL9A3: a third locus for multi-ple epiphyseal
[79878] dysplasia. Am. J. Hum. Genet. 64: 1036-1044, 1999.

[79879] 13833.

[79880] Avraham, K. B.; Hasson, T.; Sobe, T.; Balsara, B.; Testa, J. R.;
[79881] Skvorak, A. B.; Morton, C. C.; Copeland, N. G.; Jenkins, N. A.: Characterization
[79882] of unconventional MYO6, the human homologue of the gene responsible
[79883] for deafness in Snell's waltzer mice. Hum. Molec. Genet. 6: 1225-1231,
[79884] 1997.

[79885] 13834.

[79886] Avraham, K. B.; Hasson, T.; Steel, K. P.; Kingsley, D. M.;
Russell,
[79887] L. B.; Mooseker, M. S.; Copeland, N. G.; Jenkins, N. A.: The
mouse
[79888] Snell's waltzer deafness gene encodes an unconventional
myosin required
[79889] for structural integrity of inner ear hair cells. Nature
Genet. 11:
[79890] 369-375, 1995.

[79891] 13835.

[79892] Hasson, T.; Mooseker, M. S.: Porcine myosin-VI: charac-
terization
[79893] of a new mammalian unconventional myosin. J. Cell. Biol.
127: 425-440,
[79894] 1994.

[79895] 13836.

[79896] Hasson, T.; Skowron, J. F.; Gilbert, D. J.; Avraham, K. B.;
Perry,
[79897] W. L.; Bement, W. M.; Anderson, B. L.; Sherr, E. H.; Chen,
Z.-Y.;

[79898] Greene, L. A.; Ward, D. C.; Corey, D. P.; Mooseker, M. S.; Copeland,

[79899] N. G.; Jenkins, N. A.: Mapping of unconventional myosins in mouse

[79900] and human. Genomics 36: 431-439, 1996.

[79901] 13837.

[79902] Rock, R. S.; Rice, S. E.; Wells, A. L.; Purcell, T. J.; Spudich,

[79903] J. A.; Sweeney, H. L.: Myosin VI is a processive motor with a large

[79904] step size. Proc. Nat. Acad. Sci. 98: 13655-13659, 2001.

[79905] 13838.

[79906] Wells, A. L.; Lin, A. W.; Chen, L.-Q.; Safer, D.; Cain, S. M.;

[79907] Hasson, T.; Carragher, B. O.; Milligan, R. A.; Sweeney, H. L.: Myosin

[79908] VI is an actin-based motor that moves backwards. Nature 401: 505-508,

[79909] 1999.

[79910] 13839.

[79911] Melchionda, S.; Ahituv, N.; Bisceglia, L.; Sobe, T.; Glaser, F.;

[79912] Rabionet, R.; Arbones, M. L.; Notarangelo, A.; Di Iorio, E.; Carella,
[79913] M.; Zelante, L.; Estivill, X.; Avraham, K. B.; Gasparini, P.: MYO6,
[79914] the human homologue of the gene responsible for deafness in Snell's
[79915] Waltzer mice, is mutated in autosomal dominant nonsyndromic hearing
[79916] loss. Am. J. Hum. Genet. 69: 635-640, 2001.

[79917] 13840.

[79918] Bonne, S.; van Hengel, J.; van Roy, F.: Chromosomal mapping of
[79919] human armadillo genes belonging to the p120(ctn)/plakophilin subfamily. Genomics 51:
[79920] 452-454, 1998.

[79921] 13841.

[79922] Drab, M.; Verkade, P.; Elger, M.; Kasper, M.; Lohn, M.; Lauterbach,
[79923] B.; Menne, J.; Lindschau, C.; Mende, F.; Luft, F. C.; Schedl, A.;
[79924] Haller, H.; Kurzchalia, T. V.: Loss of caveolae, vascular dysfunction,

[79925] and pulmonary defects in caveolin-1 gene-disrupted mice. Science 293:
[79926] 2449-2452, 2001.

[79927] 13842.

[79928] Engelman, J. A.; Zhang, X.; Galbiati, F.; Volonte, D.; Sotgia,

[79929] F.; Pestell, R. G.; Minetti, C.; Scherer, P. E.; Okamoto, T.; Lisanti,
[79930] M. P.: Molecular genetics of the caveolin gene family: im-plications
[79931] for human cancers, diabetes, Alzheimer disease, and muscular dystrophy. Am.
[79932] J. Hum. Genet. 63: 1578-1587, 1998.

[79933] 13843.

[79934] Engelman, J. A.; Zhang, X. L.; Galbiati, F.; Lisanti, M. P.: Chromosomal
[79935] localization, genomic organization, and developmental expression of
[79936] the murine caveolin gene family (Cav-1, -2, and -3): Cav-1 and Cav-2
[79937] genes map to a known tumor suppressor locus (6-A2/7q31). FEBS Lett. 429:

[79938] 330-336, 1998.

[79939] 13844.

[79940] Engelman, J. A.; Zhang, X. L.; Lisanti, M. P.: Genes encod-
ing

[79941] human caveolin-1 and -2 are co-localized to the D7S522
locus (7q31.1),

[79942] a known fragile site (FRA7G) that is frequently deleted in
human cancers. FEBS

[79943] Lett. 436: 403-410, 1998.

[79944] 13845.

[79945] Feron, O.; Dessy, C.; Moniotte, S.; Desager, J.-P.; Balli-
gand,

[79946] J.-L.: Hypercholesterolemia decreases nitric oxide produc-
tion by

[79947] promoting the interaction of caveolin and endothelial ni-
tric oxide

[79948] synthase. J. Clin. Invest. 103: 897-905, 1999.

[79949] 13846.

[79950] Giordano, S.; Ponzetto, C.; Di Renzo, M. F.; Cooper, C. S.;
Comoglio,

[79951] P. M.: Tyrosine kinase receptor indistinguishable from the

c-met
[79952] protein. Nature 339: 155-156, 1989.

[79953] 13847.

[79954] Glenney, J. R., Jr.: The sequence of human caveolin reveals identity
[79955] with VIP21, a component of transport vesicles. FEBS Lett. 314: 45-48,
[79956] 1992.

[79957] 13848.

[79958] Kurzchalia, T. V.; Dupree, P.; Parton, R. G.; Kellner, R.; Virta,
[79959] H.; Lehnert, M.; Simons, K.: VIP21, a 21-kD membrane protein is an
[79960] integral component of trans-Golgi-network-derived transport vesicles. J.
[79961] Cell Biol. 118 1003-1014, 1992.

[79962] 13849.

[79963] Scherer, P. E.; Okamoto, T.; Chun, M.; Nishimoto, I.; Lodish,
[79964] H. F.; Lisanti, M. P.: Identification, sequence, and expres-
sion of

[79965] caveolin-2 defines a caveolin gene family. Proc. Nat. Acad. Sci. 93:

[79966] 131-135, 1996.

[79967] 13850.

[79968] Scherer, P. E.; Tang, Z.; Chun, M.; Sargiacomo, M.; Lodish, H.

[79969] F.; Lisanti, M. P.: Caveolin isoforms differ in their N-terminal

[79970] protein sequence and subcellular distribution: identification and

[79971] epitope mapping of an isoform-specific monoclonal antibody probe. J.

[79972] Biol. Chem. 270 16395-16401, 1995.

[79973] 13851.

[79974] Razani, B.; Engelman, J. A.; Wang, X. B.; Schubert, W.; Zhang,

[79975] X. L.; Marks, C. B.; Macaluso, F.; Russell, R. G.; Li, M.; Pestell,

[79976] R. G.; Di Vizio, D.; Hou, H., Jr.; Kneitz, B.; Lagaud, G.; Christ,

[79977] G. J.; Edelmann, W.; Lisanti, M. P.: Caveolin-1 null mice are viable

[79978] but show evidence of hyperproliferative and vascular abnormalities. J.

[79979] Biol. Chem. 276: 38121-38138, 2001.

[79980] 13852.

[79981] Taipale, J.; Chen, J. K.; Cooper, M. K.; Wang, B.; Mann, R. K.;

[79982] Milenkovic, L.; Scott, M. P.; Beachy, P. A.: Effects of oncogenic

[79983] mutations in Smoothened and Patched can be reversed by cyclopamine. Nature 406:

[79984] 1005-1009, 2000.

[79985] 13853.

[79986] Taipale, J.; Cooper, M. K.; Maiti, T.; Beachy, P. A.: Patched

[79987] acts catalytically to suppress the activity of Smoothened. Nature 418:

[79988] 892-897, 2002.

[79989] 13854.

[79990] Bell, D. R.; Plant, N. J.; Rider, C. G.; Na, L.; Brown, S.; Ateitalla,

[79991] I.; Acharya, S. K.; Davies, M. H.; Elias, E.; Jenkins, N. A.; Gilbert,

[79992] D. J.; Copeland, N. G.; Elcombe, C. R.: Species-specific induction
[79993] of cytochrome P-450 4A RNAs: PCR cloning of partial guinea-pig, human
[79994] and mouse CYP4A cDNAs. Biochem. J. 294: 173-180, 1993.
[79995] 13855.

[79996] Imaoka, S.; Ogawa, H.; Kimura, S.; Gonzalez, F. J.: Complete cDNA
[79997] sequence and cDNA-directed expression of CYP4A11, a fatty acid omega-hydroxylase
[79998] expressed in human kidney. DNA Cell Biol. 12: 893-899, 1993.
[79999] 13856.

[80000] Kawashima, H.; Kusunose, E.; Kikuta, Y.; Kinoshita, H.; Tanaka,
[80001] S.; Yamamoto, S.; Kishimoto, T.; Kusunose, M.: Purification and cDNA
[80002] cloning of human liver CYP4A fatty acid omega-hydroxylase. J. Biochem. 116:
[80003] 74-80, 1994.

[80004] 13857.

[80005] Palmer, C. N. A.; Richardson, T. H.; Griffin, K. J.; Hsu, M.-H.;
[80006] Muerhoff, A. S.; Clark, J. E.; Johnson, E. F.: Characterization of
[80007] a cDNA encoding a human kidney, cytochrome P-450 4A fatty acid omega-hydroxylase
[80008] and the cognate enzyme expressed in Escherichia coli. Biochim. Biophys.
[80009] Acta 1172: 161-166, 1993.

[80010] 13858.

[80011] Aziz, N.; Maxwell, M. M.; St.-Jacques, B.; Brenner, B.: Down-regulation
[80012] of Ke6, a novel gene encoded within the major histocompatibility complex,
[80013] in murine polycystic kidney disease. Molec. Cell. Biol. 13: 1847-1853,
[80014] 1993.

[80015] 13859.

[80016] Aziz, N.; Maxwell, M. M.; St.-Jacques, B.; Brenner, B.: Co-ordinate
[80017] regulation of 11-beta-HSD and Ke6 gene in cpk mouse: implications

[80018] for steroid metabolic defect in PKD. Am. J. Physiol. 267: F791-F797,

[80019] 1994.

[80020] 13860.

[80021] Muller, J.; Ory, S.; Copeland, T.; Piwnica-Worms, H.; Morrison,

[80022] D. K.: C-TAK1 regulates Ras signaling by phosphorylating the MAPK

[80023] scaffold, KSR1. Molec. Cell 8: 983-993, 2001.

[80024] 13861.

[80025] Bione, S.; Maestrini, E.; Rivella, S.; Mancini, M.; Regis, S.;

[80026] Romeo, G.; Toniolo, D.: Identification of a novel X-linked gene responsible

[80027] for Emery-Dreifuss muscular dystrophy. Nature Genet. 8: 323-327,

[80028] 1994.

[80029] 13862.

[80030] Bione, S.; Small, K.; Aksmanovic, V. M. A.; D'Urso, M.; Ciccodicola,

[80031] A.; Merlini, L.; Morandi, L.; Kress, W.; Yates, J. R. W.; Warren,

[80032] S. T.; Toniolo, D.: Identification of new mutations in the Emery-Dreifuss
[80033] muscular dystrophy gene and evidence for genetic heterogeneity of
[80034] the disease. Hum. Mol. Genet. 4: 1859-1863, 1995.

[80035] 13863.

[80036] Bione, S.; Tamanini, F.; Maestrini, E.; Tribioli, C.; Poustka,
[80037] A.; Torri, G.; Rivella, S.; Toniolo, D.: Transcriptional organization
[80038] of a 450-kb region of the human X chromosome in Xq28. Proc. Nat.
[80039] Acad. Sci. 90: 10977-10981, 1993.

[80040] 13864.

[80041] Boswinkel, E.; Walker, A.; Hodgson, S.; Benham, F.; Bobrow, M.;
[80042] Davies, K.; Dubowitz, V.; Grenata, C.: Linkage analysis using eight
[80043] DNA polymorphisms along the length of the X chromosome locates the
[80044] gene for Emery-Dreyfuss muscular dystrophy to distal Xq. (Abstract) Cytogenet.

[80045] Cell Genet. 40: 586 only, 1985.

[80046] 13865.

[80047] Rosenthal, R. L.: Haemorrhage in PTA (factor XI) deficiency.

[80048] (Abstract) Proc. 10th Int. Cong. Soc. Hemat., Stockholm , 1964.

[80049] 13866.

[80050] Laporte, J.; Hu, L. J.; Kretz, C.; Mandel, J.-L.; Kioschis, P.;

[80051] Coy, J. F.; Klauck, S. M.; Poustka, A.; Dahl, N.: A gene mutated

[80052] in X-linked myotubular myopathy defines a new putative tyrosine phosphatase

[80053] family conserved in yeast. Nature Genet. 13: 175-182, 1996.

[80054] 13867.

[80055] Kogut, M. D.; Donnell, G. N.; Nyhan, W. L.; Sweetman, L.: Disorder

[80056] of purine metabolism due to partial deficiency of hypoxanthine-guanine

[80057] phosphoribosyltransferase. Am. J. Med. 48: 148-161, 1970.

[80058] 13868.

[80059] Koller, B. H.; Hagemann, L. J.; Doetschman, T.; Hagaman, J. R.;

[80060] Huang, S.; Williams, P. J.; First, N. L.; Maeda, N.; Smithies, O.

[80061] : Germ-line transmission of a planned alteration made in a hypoxanthine

[80062] phosphoribosyltransferase gene by homologous recombination in embryonic

[80063] stem cells. Proc. Nat. Acad. Sci. 86: 8927-8931, 1989.

[80064] 13869.

[80065] Kuehn, M. R.; Bradley, A.; Robertson, E. J.; Evans, M. J.: A

[80066] potential animal model for Lesch-Nyhan syndrome through introduction

[80067] of HPRT mutations into mice. Nature 326: 295-298, 1987.

[80068] 13870.

[80069] Lesch, M.; Nyhan, W. L.: A familial disorder of uric acid metabolism

[80070] and central nervous system function. Am. J. Med. 36: 561-570, 1964.

[80071] 13871.

[80072] Lightfoot, T.; Joshi, R.; Nuki, G.; Snyder, F. F.: The point

[80073] mutation of hypoxanthine-guanine phosphoribosyltransferase (HPRT-Edinburgh)

[80074] and detection by allele-specific polymerase chain reaction. Hum.

[80075] Genet. 88: 695-696, 1992.

[80076] 13872.

[80077] Lightfoot, T.; Lewkonia, R. M.; Snyder, F. F.: Sequence, expression

[80078] and characterization of HPRT-Moose Jaw: a point mutation resulting

[80079] in cooperativity and decreased substrate affinities. Hum. Molec.

[80080] Genet. 3: 1377-1381, 1994.

[80081] 13873.

[80082] Malleson, P. N.; Fung, M. Y.; Rosenberg, A. M.: The incidence

[80083] of pediatric rheumatic diseases: results from the Canadian Pediatric

[80084] Rheumatology Association Disease Registry. J. Rheum. 23: 1981-1987,

[80085] 1996.

[80086] 13874.

[80087] Marcus, S.; Steen, A.-M.; Andersson, B.; Lambert, B.; Kristoffersson,

[80088] U.; Francke, U.: Mutation analysis and prenatal diagnosis in a Lesch-Nyhan

[80089] family showing non-random X-inactivation interfering with carrier

[80090] detection tests. Hum. Genet. 89: 395-400, 1992.

[80091] 13875.

[80092] McDonald, J. A.; Kelley, W. N.: Lesch-Nyhan syndrome: absence

[80093] of the mutant enzyme in erythrocytes of a heterozygote for both normal

[80094] and mutant hypoxanthine-guanine phosphoribosyl transferase. Biochem.

[80095] Genet. 6: 21-26, 1972.

[80096] 13876.

[80097] McKeran, R. O.; Andrews, T. M.; Howell, A.; Gibbs, D. A.; Chinn,

[80098] S.; Watts, R. W. E.: The diagnosis of the carrier state for

the Lesch-Nyhan
[80099] syndrome. Quart. J. Med. 44: 189-206, 1975.

[80100] 13877.

[80101] Melton, D. W.; Konecki, D. S.; Brennand, J.; Caskey, C. T.: Structure,
[80102] expression, and mutation of the hypoxanthine phosphoribosyltransferase
[80103] gene. Proc. Nat. Acad. Sci. 81: 2147-2151, 1984.

[80104] 13878.

[80105] Migeon, B. R.: X-linked hypoxanthine-guanine phosphoribosyl transferase
[80106] deficiency: detection of heterozygotes by selective medium. Biochem.
[80107] Genet. 4: 377-383, 1970.

[80108] 13879.

[80109] Miller, A. D.; Jolly, D. J.; Friedmann, T.; Verma, I. M.: A transmissible
[80110] retrovirus expressing human hypoxanthine phosphoribosyltransferase
[80111] (HPRT): gene transfer into cells obtained from humans deficient in

[80112] HPRT. Proc. Nat. Acad. Sci. 80: 4709-4713, 1983.

[80113] 13880.

[80114] Mizunuma, M.; Fujimori, S.; Ogino, H.; Ueno, T.; Inoue, H.; Kamatani,

[80115] N.: A recurrent large Alu-mediated deletion in the hypox-anthine phosphoribosyltransferase

[80116] (HPRT1) gene associated with Lesch-Nyhan syndrome. Hum. Mutat. 18:

[80117] 435-443, 2001.

[80118] 13881.

[80119] Monk, M.; Handyside, A.; Hardy, K.; Whittingham, D.: Preimplantation

[80120] diagnosis of deficiency of hypoxanthine phosphoribosyl transferase

[80121] in a mouse model for Lesch-Nyhan syndrome. Lancet II: 423-425, 1987.

[80122] 13882.

[80123] Monk, M.; Handyside, A.; Muggleton-Harris, A.; Whitting-ham, D.

[80124] : Preimplantation sexing and diagnosis of hypoxanthine phosphoribosyl

[80125] transferase deficiency in mice by biochemical microassay. Am. J.

[80126] Med. Genet. 35: 201-205, 1990.

[80127] 13883.

[80128] Monnat, R. J., Jr.; Chiaverotti, T. A.; Hackmann, A. F. M.; Maresh,

[80129] G. A.: Molecular structure and genetic stability of human hypoxanthine

[80130] phosphoribosyltransferase (HPRT) gene duplications. Genomics 13:

[80131] 788-796, 1992.

[80132] 13884.

[80133] Myers, R. M.; Larin, A.; Maniatis, T.: Detection of single base

[80134] substitutions by ribonuclease cleavage at mismatches in RNA:DNA duplexes. Science 230:

[80135] 1242-1246, 1985.

[80136] 13885.

[80137] Newcombe, D. S.; Shapiro, S. L.; Sheppard, G. L., Jr.; Dreifuss,

[80138] F. E.: Treatment of X-linked primary hyperuricemia with

allopurinol. J.A.M.A. 198:
[80139] 315-317, 1966.

[80140] 13886.

[80141] Nussbaum, R. L.; Crowder, W. E.; Nyhan, W. L.; Caskey, C. T.:
[80142] A three-allele restriction-fragment-length polymorphism at the hypoxanthine
[80143] phosphoribosyltransferase locus in man. Proc. Nat. Acad. Sci. 80:
[80144] 4035-4039, 1983.

[80145] 13887.

[80146] Netchine, I.; Sobrier, M.-L.; Krude, H.; Schnabel, D.; Maghnie,
[80147] M.; Marcos, E.; Duriez, B.; Cacheux, V.; Moers, A. V.; Goossens, M.;
[80148] Gruters, A.; Amselem, S.: Mutations in LHX3 result in a new syndrome
[80149] revealed by combined pituitary hormone deficiency. Nature Genet. 25:
[80150] 182-186, 2000.

[80151] 13888.

[80152] Rosenthal, R. L.; Dreskin, O. H.; Rosenthal, N.: Plasma thromboplastin
[80153] antecedent (PTA) deficiency: clinical, coagulation, thera-peutic and
[80154] hereditary aspects of a new hemophilia-like disease. Blood 10: 120-131,
[80155] 1955.

[80156] 13889.

[80157] Seligsohn, U.: Factor XI (PTA) deficiency.In: Goodman, R. E.;
[80158] Motulsky, A. G.: Genetic Diseases Among Ashkenazi Jews. New York:
[80159] Raven Press (pub.) 1979. Pp. 141-148.

[80160] 13890.

[80161] Vinazzer, H.: Partieller familiaerer Faktor-XI-Mangel. Blut 15:
[80162] 263-267, 1967.

[80163] 13891.

[80164] Wistinghausen, B.; Reischer, A.; Oddoux, C.; Ostrer, H.; Nardi,
[80165] M.; Karpatkin, M.: Severe factor XI deficiency in an Arab

family
[80166] associated with a novel mutation in exon 11. Brit. J. Haemat. 99:
[80167] 575-577, 1997.

[80168] 13892.

[80169] Zivelin, A.; Bauduer, F.; Ducout, L.; Peretz, H.; Rosenberg, N.;
[80170] Yatuv, R.; Seligsohn, U.: Factor XI deficiency in French Basques
[80171] is caused predominantly by an ancestral cys38-to-arg mutation in the
[80172] factor XI gene. Blood 99: 2448-2454, 2002.

[80173] 13893.

[80174] Inoue, K.; Tanabe, Y.; Lupski, J. R.: Myelin deficiencies in both
[80175] the central and the peripheral nervous systems associated with a SOX10
[80176] mutation. Ann. Neurol. 46: 313-318, 1999.

[80177] 13894.

[80178] Touraine, R. L.; Attie-Bitach, T.; Manceau, E.; Korsch, E.; Sarda,

[80179] P.; Pingault, V.; Encha-Razavi, F.; Pelet, A.; Auge, J.; Nivelon-Chevallier,
[80180] A.; Holschneider, A. M.; Munnes, M.; Doerfler, W.; Goossens, M.; Munnich,
[80181] A.; Vekemans, M.; Lyonnet, S.: Neurological phenotype in Waardenburg
[80182] syndrome type 4 correlates with novel SOX10 truncating mutations and
[80183] expression in developing brain. Am. J. Hum. Genet. 66: 1496-1503,
[80184] 2000. Note: Erratum: Am. J. Hum. Genet. 66: 2020 only, 2000.
[80185] 13895.
[80186] De Smet, C.; Lurquin, C.; van der Bruggen, P.; De Plaen, E.; Brasseur,
[80187] F.; Boon, T.: Sequence and expression pattern of the human MAGE2
[80188] gene. Immunogenetics 39: 121-129, 1994.
[80189] 13896.
[80190] Chambost, H.; Van Baren, N.; Brasseur, F.; Godelaine, D.; Xerri,
[80191] L.; Landi, S. J.; Theate, I.; Plumas, J.; Spagnoli, G. C.;

Michel,
[80192] G.; Coulie, P. G.; Olive, D.: Expression of gene MAGE-A4 in Reed-Sternberg
[80193] cells. Blood 95: 3530-3533, 2000.

[80194] 13897.

[80195] Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Miyajima, N.;
[80196] Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the
[80197] coding sequences of unidentified human genes. XI. The complete sequences
[80198] of 100 new cDNA clones from brain which code for large proteins in
[80199] vitro. DNA Res. 5: 277-286, 1998.

[80200] 13898.

[80201] Hasstedt, S. J.; Atkin, C. L.: X-linked inheritance of Alport
[80202] syndrome: family P revisited. Am. J. Hum. Genet. 35: 1241-1251,
[80203] 1983.

[80204] 13899.

[80205] Scherer, S. S.; Deschenes, S. M.; Xu, Y.; Grinspan, J. B.; Fis-

chbeck,

[80206] K. H.; Paul, D. L.: Connexin32 is a myelin-related protein in the

[80207] PNS and CNS. J. Neurosci. 15: 8281-8294, 1995.

[80208] 13900.

[80209] Gospe, S. M., Jr.; Lazaro, R. P.; Lava, N. S.; Grootscholten,

[80210] P. M.; Scott, M. O.; Fischbeck, K. H.: Familial X-linked myalgia

[80211] and cramps: a nonprogressive myopathy associated with a deletion in

[80212] the dystrophin gene. Neurology 39: 1277-1280, 1989.

[80213] 13901.

[80214] Kingston, H. M.; Sarfarazi, M.; Thomas, N. S. T.; Harper, P. S.

[80215] : Localisation of the Becker muscular dystrophy gene on the short

[80216] arm of the X chromosome by linkage to cloned DNA se-

quences. Hum.

[80217] Genet. 67: 6-17, 1984.

[80218] 13902.

[80219] Kingston, H. M.; Thomas, N. S. T.; Pearson, P. L.; Sarfarazi,

[80220] M.; Harper, P. S.: Genetic linkage between Becker muscular dystrophy
[80221] and a polymorphic DNA sequence on the short arm of the X chromosome. J.
[80222] Med. Genet. 20: 255-258, 1983.

[80223] 13903.

[80224] Gilbert, F.; Kucherlapati, R. S.; Creagan, R. P.; Murnane, M.

[80225] J.; Darlington, G. J.; Ruddle, F. H.: Tay-Sachs' and Sandhoff's diseases:
[80226] the assignment of genes for hexosaminidase A and B to individual human
[80227] chromosomes. Proc. Nat. Acad. Sci. 72: 263-267, 1975.

[80228] 13904.

[80229] Derry, J. M. J.; Jess, U.; Francke, U.: Cloning and characterization
[80230] of a novel zinc finger gene in Xp11.2. Genomics 30: 361-365, 1995.
[80231] 13905.

[80232] Derry, J. M. J.; Kerns, J. A.; Francke, U.: RBM3, a novel human
[80233] gene in Xp11.23 with a putative RNA-binding domain.

Hum. Molec. Genet. 4:
[80234] 2307-2311, 1995.

[80235] 13906.

[80236] Hirst, M. C.; Barnicoat, A.; Flynn, G.; Wang, Q.; Daker, M.; Buckle,
[80237] V. J.; Davies, K. E.; Bobrow, M.: The identification of a third fragile
[80238] site, FRAXF, in Xq27-q28 distal to both FRAXA and FRAXE. Hum. Molec.
[80239] Genet. 2: 197-200, 1993.

[80240] 13907.

[80241] Ades, L. C.; Kerr, B.; Turner, G.; Wise, G.: Smith-Fine-man-Myers
[80242] syndrome in two brothers. Am. J. Med. Genet. 40: 467-470, 1991.
[80243] 13908.

[80244] Bachoo, S.; Gibbons, R. J.: Germline and gonosomal mo-saicism in
[80245] the ATR-X syndrome. Europ. J. Hum. Genet. 7: 933-936, 1999.
[80246] 13909.

[80247] Berube, N. G.; Jagla, M.; Smeenk, C.; De Repentigny, Y.; Kothary,
[80248] R.; Picketts, D. J.: Neurodevelopmental defects resulting from ATRX
[80249] overexpression in transgenic mice. Hum. Molec. Genet. 11: 253-261,
[80250] 2002.

[80251] 13910.

[80252] Berube, N. G.; Smeenk, C. A.; Picketts, D. J.: Cell cycle-dependent
[80253] phosphorylation of the ATRX protein correlates with changes in nuclear
[80254] matrix and chromatin association. Hum. Molec. Genet. 9: 539-547,
[80255] 2000.

[80256] 13911.

[80257] Cardoso, C.; Timsit, S.; Villard, L.; Khrestchatisky, M.; Fontes,
[80258] M.; Colleaux, L.: Specific interaction between the XNP/ATR-X gene
[80259] product and the SET domain of the human EZH2 protein. Hum. Molec.

[80260] Genet. 7: 679-684, 1998.

[80261] 13912.

[80262] Gibbons, R. J.; Bachoo, S.; Picketts, D. J.; Aftimos, S.; Asenbauer,

[80263] B.; Bergoffen, J.; Berry, S. A.; Dahl, N.; Fryer, A.; Keppler, K.;

[80264] Kurosawa, K.; Levin, M. L.; Masuno, M.; Neri, G.; Pierpont, M. E.;

[80265] Slaney, S. F.; Higgs, D. R.: Mutations in transcriptional regulator

[80266] ATRX establish the functional significance of a PHD-like domain. (Letter) Nature

[80267] Genet. 17: 146-148, 1998.

[80268] 13913.

[80269] Grover, S.; Fishman, G. A.; Anderson, R. J.; Lindeman, M.: A

[80270] longitudinal study of visual function in carriers of X-linked recessive

[80271] retinitis pigmentosa. Ophthalmology 107: 386-396, 2000.

[80272] 13914.

[80273] Hunter, D. G.; Fishman, G. A.; Kretzer, F. L.: Abnormal ax-

onemes
[80274] in X-linked retinitis pigmentosa. Arch. Ophthal. 106: 362-368, 1988.
[80275] 13915.

[80276] Matsumoto, T.; Kondoh, T.; Yoshimoto, M.; Fujieda, K.; Matsuura,
[80277] N.; Matsuda, I.; Miike, T.; Yano, K.; Okuno, A.; Aoki, Y.; Murano,
[80278] I.; Toyota, S.; Ohnishi, S.; Niikawa, N.: Complex glycerol kinase
[80279] deficiency: molecular-genetic, cytogenetic, and clinical studies of
[80280] five Japanese patients. Am. J. Med. Genet. 31: 603-616, 1988.
[80281] 13916.

[80282] Ament, M. E.; Ochs, H. D.: Gastrointestinal manifestations of
[80283] chronic granulomatous disease. New Eng. J. Med. 288: 382-387, 1973.
[80284] 13917.

[80285] Ariga, T.; Furuta, H.; Cho, K.; Sakiyama, Y.: Genetic analysis

[80286] of 13 families with X-linked chronic granulomatous disease reveals
[80287] a low proportion of sporadic patients and a high proportion of sporadic
[80288] carriers. Pediat. Res. 44: 85-92, 1998.

[80289] 13918.

[80290] Babior, B. M.: Personal Communication. Boston, Mass. 3/26/1974.
[80291] 13919.

[80292] Babior, B. M.; Crowley, C. A.: Chronic granulomatous disease and
[80293] other disorders of oxidative killing by phagocytes.In: Stanbury, J.
[80294] B.; Wyngaarden, J. B.; Fredrickson, D. S.; Goldstein, J. L.; Brown,
[80295] M. S.: The Metabolic Basis of Inherited Disease. New York: McGraw-Hill
[80296] (pub.) 1983. Pp. 1965-1985.

[80297] 13920.

[80298] Baehner, R. L.; Karnovsky, M. L.: Deficiency of reduced nicotinamide-adenine

[80299] dinucleotide oxidase in chronic granulomatous disease. Science 162:
[80300] 1277-1279, 1968.

[80301] 13921.

[80302] Baehner, R. L.; Kunkel, L. M.; Monaco, A. P.; Haines, J. L.; Conneally,
[80303] P. M.; Palmer, C.; Heerema, N.; Orkin, S. H.: DNA linkage analysis
[80304] of X chromosome-linked chronic granulomatous disease. Proc. Nat.
[80305] Acad. Sci. 83: 3398-3401, 1986.

[80306] 13922.

[80307] Baehner, R. L.; Nathan, D. G.: Leukocyte oxidase: defective activity
[80308] in chronic granulomatous disease. Science 155: 835-836, 1967.
[80309] 13923.

[80310] Biggar, W. D.: Phagocytosis in patients and carriers of chronic
[80311] granulomatous disease. Lancet I: 991-994, 1975.

[80312] 13924.

[80313] Nyhan, W. L.; Resek, J.; Sweetman, L.; Carpenter, D. G.; Carter,
[80314] C. H.: Genetics of an X-linked disorder of uric acid metabolism and
[80315] cerebral function. Pediat. Res. 1: 5-13, 1967.

[80316] 13925.

[80317] Ogasawara, N.; Stout, J. T.; Goto, H.; Sonta, S.; Matsumoto, A.;
[80318] Caskey, C. T.: Molecular analysis of a female Lesch-Nyhan patient. J.
[80319] Clin. Invest. 84: 1024-1027, 1989.

[80320] 13926.

[80321] Pai, G. S.; Sprenkle, J. A.; Do, T. T.; Mareni, C. E.; Migeon,
[80322] B. R.: Localization of loci for hypoxanthine phosphate de-hydrogenase
[80323] and glucose-6-phosphate dehydrogenase and biochemi-cal evidence of
[80324] nonrandom X chromosome expression from studies of a human X-autosome
[80325] translocation. Proc. Nat. Acad. Sci. 77: 2810-2813, 1980.

[80326]

[80327] 100. Palella, T. D.: Personal Communication. Ann Arbor, Mich.

[80328] 1/15/1990.

[80329]

[80330] 101. Patel, P. I.; Framson, P. E.; Caskey, C. T.; Chinault, A. C.

[80331] : Fine structure of the human hypoxanthine phosphoribo-syltransferase

[80332] gene. Molec. Cell. Biol. 6: 393-403, 1986.

[80333]

[80334] 102. Race, R. R.; Sanger, R.: Blood Groups in Man. Philadelphia:

[80335] F. A. Davis Co. (pub.) (5th ed.): 1968. Pp. 545 only.

[80336]

[80337] 103. Renwick, P. J.; Birley, A. J.; McKeown, C. M. E.; Hulten, M.

[80338] : Southern analysis reveals a large deletion at the hypox-anthine phosphoribosyltransferase

[80339] locus in a patient with Lesch-Nyhan syndrome. Clin. Genet. 48: 80-84,

[80340] 1995.

[80341]

[80342] 104. Rijksen, G.; Staal, G. E. J.; van der Vlist, M. J. M.; Beemer,

[80343] F. A.; Troost, J.; Gutensohn, W.; van Laarhoven, J. P. R. M.; de Bruyn,

[80344] C. H. M. M.: Partial hypoxanthine-guanine phosphoribosyl transferase

[80345] deficiency with full expression of the Lesch-Nyhan syndrome. Hum.

[80346] Genet. 57: 39-47, 1981.

[80347]

[80348] 105. Rosenbloom, F. M.; Kelley, W. N.; Henderson, J. F.; Seegmiller,

[80349] J. E.: Lyon hypothesis and X-linked disease. (Letter) Lancet II:

[80350] 305-306, 1967.

[80351]

[80352] 106. Rosenbloom, F. M.; Kelley, W. N.; Miller, J.; Henderson, J. F.;

[80353] Seegmiller, J. E.: Inherited disorder of purine metabolism: correlation

[80354] between central nervous system dysfunction and bio-
chemical defects. J.A.M.A. 202:
[80355] 175-177, 1967.

[80356]

[80357] 107. Rosenstraus, M.; Chasin, L. A.: Isolation of mam-
malian cell
[80358] mutants deficient in glucose-6-phosphate dehydrogenase
activity: linkage
[80359] to hypoxanthine phosphoribosyl transferase. Proc. Nat.
Acad. Sci. 72:
[80360] 493-497, 1975.

[80361]

[80362] 108. Ross, J. A.; Robison, L. L.: MLL rearrangements in in-
fant leukemia:
[80363] is there a higher frequency in females? Leukemia Res. 21:
793-795,
[80364] 1997.

[80365]

[80366] 109. Rossiter, B. J. F.; Edwards, A.; Caskey, C. T.: HPRT
mutation
[80367] and the Lesch-Nyhan syndrome.In: Brosius, J.; Freneau, R.:

Molecular
[80368] Genetic Approaches to Neuropsychiatric Disease. New York: Academic
[80369] Press (pub.) 1991.

[80370]

[80371] 110. Ruddle, F. H.: Linkage studies employing mouse-man somatic cell
[80372] hybrids. Fed. Proc. 30: 921-925, 1971.

[80373]

[80374] 111. Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.: Sheep
[80375] gene mapping by somatic cell hybridization. (Abstract) Cytogenet.
[80376] Cell Genet. 25: 200 only, 1979.

[80377]

[80378] 112. Sass, J. K.; Itabashi, H. H.; Dexter, R. A.: Juvenile gout with
[80379] brain involvement. Arch. Neurol. 13: 639-655, 1965.

[80380]

[80381] 113. Sculley, D. G.; Dawson, P. A.; Beacham, I. R.; Emmer-

son, B. T.;
[80382] Gordon, R. B.: Hypoxanthine-guanine phosphoribosyl-
transferase deficiency:
[80383] analysis of HPRT mutations by direct sequencing and al-
lele-specific
[80384] amplification. Hum. Genet. 87: 688-692, 1991.

[80385]

[80386] 114. Sculley, D. G.; Dawson, P. A.; Emmerson, B. T.; Gor-
don, R. B.
[80387] : A review of the molecular basis of hypoxanthine-guanine
phosphoribosyltransferase
[80388] (HPRT) deficiency. Hum. Genet. 90: 195-207, 1992.

[80389]

[80390] 115. Seegmiller, J. E.: Contributions of Lesch-Nyhan syn-
drome to
[80391] the understanding of purine metabolism. J. Inherit. Metab.
Dis. 12:
[80392] 184-196, 1989.

[80393]

[80394] 116. Seegmiller, J. E.; Rosenbloom, F. M.; Kelley, W. N.:
Enzyme

[80395] defect associated with a sex-linked human neurological disorder and
[80396] excessive purine synthesis. Science 155: 1682-1684, 1967.
[80397]

[80398] 117. Shapiro, S. L.; Sheppard, G. L., Jr.; Dreifuss, F. E.; Newcombe,
[80399] D. S.: X-linked recessive inheritance of a syndrome of mental retardation
[80400] with hyperuricemia. Proc. Soc. Exp. Biol. Med. 122: 609-611, 1966.
[80401]

[80402] 118. Shows, T. B.; Brown, J. A.: Human X-linked genes re-gionally
[80403] mapped utilizing X-autosome translocations and somatic cell hybrids. Proc.
[80404] Nat. Acad. Sci. 72: 2125-2129, 1975.

[80405]

[80406] 119. Shows, T. B.; Brown, J. A.; Chapman, V. M.: Compara-tive gene
[80407] mapping of HPRT, G6PD and PGK in man, mouse, and muntjac deer. Birth

[80408] Defects Orig. Art. Ser. XII(7): 436-439, 1976.

[80409]

[80410] 120. Silvers, D. N.; Cox, R. P.; Balis, M. E.; Dancis, J.: De-
tection
[80411] of the heterozygote in Lesch-Nyhan disease by hair-root
analysis. New
[80412] Eng. J. Med. 286: 390-395, 1972.

[80413]

[80414] 121. Simpson, D.; Crosby, R. M.; Skopek, T. R.: A method
for specific
[80415] cloning and sequencing of human HPRT cDNA for muta-
tion analysis. Biochem.
[80416] Biophys. Res. Commun. 151: 487-492, 1988.

[80417]

[80418] 122. Sinnett, D.; Lavergne, L.; Melancon, S. B.; Dallaire, L.;
Potier,
[80419] M.; Labuda, D.: Lesch-Nyhan syndrome: molecular inves-
tigation of
[80420] three French Canadian families using a hypoxanthine-gua-
nine phosphoribosyltransferase
[80421] cDNA probe. Hum. Genet. 81: 4-8, 1988.

[80422]

[80423] 123. Skopek, T. R.; Recio, L.; Simpson, D.; Dallaire, L.; Melancon,

[80424] S. B.; Ogier, H.; O'Neill, J. P.; Falta, M. T.; Nicklas, J. A.; Albertini,

[80425] R. J.: Molecular analyses of a Lesch-Nyhan syndrome mutation (HPRT-Montreal)

[80426] by use of T-lymphocyte cultures. Hum. Genet. 85: 111-116, 1990.

[80427]

[80428] 124. Snyder, F. F.; Chudley, A. E.; MacLeod, P. M.; Carter, R. J.;

[80429] Fung, E.; Lowe, J. K.: Partial deficiency of hypoxanthine-guanine

[80430] phosphoribosyltransferase with reduced affinity for PP-ribose-P in

[80431] four related males with gout. Hum. Genet. 67: 18-22, 1984.

[80432]

[80433] 125. Snyder, F. F.; Joyce, J. E.; Carter-Edwards, T.; Joshi, R.; Rylance,

[80434] H. L.; Wallace, R. C.; Nuki, G.: Hypoxanthine-guanine

phosphoribosyltransferase
[80435] deficiency in three brothers with gout: characterization of a variant,
[80436] HPRT(Edinburgh), having altered isoelectric point, increased thermal
[80437] lability and normal levels of messenger RNA. J. Inherit. Metab. Dis. 12:
[80438] 390-402, 1989.

[80439]

[80440] 126. Sperling, O.; Frank, M.; Ophir, R.; Liberman, U. A.; Adam, A.;
[80441] De Vries, A.: Partial deficiency of hypoxanthine-guanine phosphoribosyltransferase
[80442] associated with gout and uric acid lithiasis. Europ. J. Clin. Biol.
[80443] Res. 15: 942-947, 1970.

[80444]

[80445] 127. Srivastava, T.; O'Neill, J. P.; Dasouki, M.; Simckes, A. M.:
[80446] Childhood hyperuricemia and acute renal failure resulting from a missense
[80447] mutation in the HPRT gene. Am. J. Med. Genet. 108:

219-222, 2002.

[80448]

[80449] 128. Stout, J. T.; Caskey, C. T.: HPRT: gene structure, ex-
pression,

[80450] and mutation. Annu. Rev. Genet. 19: 127-148, 1985.

[80451]

[80452] 129. Stout, J. T.; Caskey, C. T.: Personal Communication.
Houston,

[80453] Tex. 5/5/1984.

[80454]

[80455] 130. Strauss, G. H.; Allen, E. F.; Albertini, R. J.: An enu-
merative

[80456] assay of purine analogue resistant lymphocytes in women
heterozygous

[80457] for the Lesch-Nyhan mutation. Biochem. Genet. 18:
529-547, 1980.

[80458]

[80459] 131. Strauss, M.; Lubbe, L.; Geissler, E.: HGPRT structural
gene

[80460] mutation in Lesch-Nyhan-syndrome as indicated by anti-
genic activity

[80461] and reversion of the enzyme deficiency. Hum. Genet. 57:
185-188,
[80462] 1981.

[80463]

[80464] 132. Tarle, S. A.; Davidson, B. L.; Wu, V. C.; Zidar, F. J.;
Seegmiller,
[80465] J. E.; Kelley, W. N.; Palella, T. D.: Determination of the mu-
tations
[80466] responsible for the Lesch-Nyhan syndrome in 17 subjects.
Genomics 10:
[80467] 499-501, 1991.

[80468]

[80469] 133. Toyo-Oka, T.; Hanaoka, F.; Akaoka, I.; Yamada,
M.-A.: X-linked
[80470] hypoxanthine-guanine phosphoribosyl transferase defi-
ciency without
[80471] neurological disorders: a report of a family. Clin. Genet. 7:
181-185,
[80472] 1975.

[80473]

[80474] 134. van Bogaert, P.; Ceballos, I.; Desguerre, I.; Telvi, L.;

Kamoun,
[80475] P.; Ponsot, G.: Lesch Nyhan syndrome in a girl. J. Inherit. Metab.
[80476] Dis. 15: 790-791, 1992.

[80477]

[80478] 135. Vogel, F.: A probable sex difference in some mutation rates.
[80479] (Editorial) Am. J. Hum. Genet. 29: 312-319, 1977.

[80480]

[80481] 136. Willers, I.; Held, K. R.; Singh, S.; Goedde, H. W.: Genetic
[80482] heterogeneity of hypoxanthine-phosphoribosyl transferase in human
[80483] fibroblasts of 3 families. Clin. Genet. 11: 193-200, 1977.

[80484]

[80485] 137. Wilson, J. M.; Baugher, B. W.; Landa, L.; Kelley, W. N.: Human
[80486] hypoxanthine-guanine phosphoribosyltransferase: purification and characterization
[80487] of mutant forms of the enzyme. J. Biol. Chem. 256: 10306-10312,

[80488] 1981.

[80489]

[80490] 138. Wilson, J. M.; Baugher, B. W.; Mattes, P. M.; Daddona, P. E.;

[80491] Kelley, W. N.: Human hypoxanthine-guanine phosphoribosyltransferase:

[80492] demonstration of structural variants in lymphoblastoid cells derived

[80493] from patients with a deficiency of the enzyme. J. Clin. Invest. 69:

[80494] 706-715, 1982.

[80495]

[80496] 139. Wilson, J. M.; Frossard, P.; Nussbaum, R. L.; Caskey, C. T.;

[80497] Kelley, W. N.: Human hypoxanthine-guanine phosphoribosyltransferase:

[80498] detection of a mutant allele by restriction endonuclease analysis. J.

[80499] Clin. Invest. 72: 767-772, 1983.

[80500]

[80501] 140. Wilson, J. M.; Kelley, W. N.: Molecular basis of hypox-

anthine-guanine
[80502] phosphoribosyltransferase deficiency in a patient with the Lesch-Nyhan
[80503] syndrome. J. Clin. Invest. 71: 1331-1335, 1983.

[80504]

[80505] 141. Wilson, J. M.; Kelley, W. N.: Human hypoxanthine-guanine phosphoribosyltransferase:
[80506] structural alteration in a dysfunctional enzyme variant (HPRT-Munich)
[80507] isolated from a patient with gout. J. Biol. Chem. 259: 27-30, 1984.

[80508]

[80509] 142. Wilson, J. M.; Kobayashi, R.; Fox, I. H.; Kelley, W. N.: Human
[80510] hypoxanthine-guanine phosphoribosyltransferase: molecular abnormality
[80511] in a mutant form of the enzyme (HPRT-Toronto). J. Biol. Chem. 258:
[80512] 6458-6460, 1983.

[80513]

[80514] 143. Wilson, J. M.; Stout, J. T.; Palella, T. D.; Davidson, B.

L.;

[80515] Kelley, W. N.; Caskey, C. T.: A molecular survey of hypoxanthine-guanine

[80516] phosphoribosyltransferase deficiency in man. J. Clin. Invest. 77:

[80517] 188-195, 1986.

[80518]

[80519] 144. Wilson, J. M.; Tarr, G. E.; Kelley, W. N.: Human hy-
poxanthine
[80520] (guanine) phosphoribosyltransferase: an amino acid sub-
stitution in
[80521] a mutant form of the enzyme isolated from a patient with
gout. Proc.
[80522] Nat. Acad. Sci. 80: 870-873, 1983.

[80523]

[80524] 145. Wilson, J. M.; Young, A. B.; Kelley, W. N.: Hypoxan-
thine-guanine
[80525] phosphoribosyltransferase deficiency: the molecular basis
of the clinical
[80526] syndromes. New Eng. J. Med. 309: 900-910, 1983.

[80527]

[80528] 146. Winter, R. M.: Estimation of male to female ratio of
mutation
[80529] rates from carrier-detection tests in X-linked disorders.
Am. J.
[80530] Hum. Genet. 32: 582-588, 1980.

[80531]

[80532] 147. Wong, D. F.; Harris, J. C.; Naidu, S.; Yokoi, F.;
Marenco, S.;
[80533] Dannals, R. F.; Ravert, H. T.; Yaster, M.; Evans, A.; Rousset,
O.;
[80534] Bryan, R. N.; Gjedde, A.; Kuhar, M. J.; Breese, G. R.:
Dopamine transporters
[80535] are markedly reduced in Lesch-Nyhan disease in vivo.
Proc. Nat. Acad.
[80536] Sci. 93: 5539-5543, 1996.

[80537]

[80538] 148. Wu, C.-L.; Melton, D. W.: Production of a model for
Lesch-Nyhan
[80539] syndrome in hypoxanthine phosphoribosyltransferase-defi-
cient mice. Nature
[80540] Genet. 3: 235-239, 1993.

[80541]

[80542] 149. Yang, T. P.; Patel, P. I.; Chinault, A. C.; Stout, J. T.;
Jackson,
[80543] L. G.; Hildebrand, B. M.; Caskey, C. T.: Molecular evidence
for new

[80544] mutation at the HPRT locus in Lesch-Nyhan patients. Nature 310:
[80545] 412-414, 1984.

[80546]

[80547] 150. Yang, T. P.; Stout, J. T.; Konecki, D. S.; Patel, P. I.; Alford,
[80548] R. L.; Caskey, C. T.: Spontaneous reversion of novel Lesch-Nyhan
[80549] mutation by HPRT gene rearrangement. Somat. Cell Molec. Genet. 14:
[80550] 293-303, 1988.

[80551]

[80552] 151. Yu, T.-F.; Balis, M. E.; Krenitsky, T. A.; Dancis, J.; Silvers,
[80553] D. N.; Elion, G. B.; Gutman, A. B.: Rarity of X-linked partial hypoxanthine-guanine
[80554] phosphoribosyltransferase deficiency in a large gouty population. Ann.
[80555] Intern. Med. 76: 255-264, 1972.

[80556]

[80557] 152. Zannis, V. I.; Gudas, L. J.; Martin, D. W., Jr.: Charac-

terization
[80558] of the subunit composition of HGPRTase from human erythrocytes and
[80559] cultured fibroblasts. Biochem. Genet. 18: 1-19, 1980.

[80560]

[80561] 153. Zoref, E.; Sperling, O.: Increased de novo purine synthesis
[80562] in cultured skin fibroblasts from heterozygotes for the Lesch-Nyhan
[80563] syndrome: a sensitive marker for carrier detection. Hum. Hered. 29:
[80564] 64-68, 1979.

[80565]

[80566] 154. Zoref-Shani, E.; Feinstein, S.; Frishberg, Y.; Bromberg, Y.;
[80567] Sperling, O.: Kelley-Seegmiller syndrome due to a unique variant
[80568] of hypoxanthine-guanine phosphoribosyltransferase: reduced affinity
[80569] for 5-phosphoribosyl-1-pyrophosphate manifested only at low, physiological
[80570] substrate concentrations. Biochim. Biophys. Acta 1500:

197-203,
[80571] 2000.

[80572] 13927.

[80573] Adam, A.; Ziprkowski, L.; Feinstein, A.; Sanger, R.; Tippett, P.;
[80574] Gavin, J.; Race, R. R.: Linkage relations of X-borne ichthyosis to
[80575] the Xg blood groups and to other markers of the X in Israelis. Ann.
[80576] Hum. Genet. 32: 323-332, 1969.

[80577] 13928.

[80578] Alperin, E. S.; Shapiro, L. J.: Characterization of point mutations
[80579] with X-linked ichthyosis: effects on the structure and function of
[80580] the steroid sulfatase protein. J. Biol. Chem. 272: 20756-20763,
[80581] 1997.

[80582] 13929.

[80583] Andria, G.; Ballabio, A.; Parenti, G.: X-linked ichthyosis due

[80584] to steroid sulfatase deficiency associated with hypogonadism and anosmia.

[80585] (Letter) Ann. Neurol. 22: 98, 1987.

[80586] 13930.

[80587] Andria, G.; Ballabio, A.; Parenti, G.; Di Maio, S.; Piccirillo,

[80588] A.: Steroid sulphatase deficiency is present in patients with the

[80589] syndrome 'ichthyosis and male hypogonadism' and with 'Rud syndrome.'. J.

[80590] Inherit. Metab. Dis. 7: 159-160, 1984.

[80591] 13931.

[80592] Bairoch, A.: Personal Communication. Geneva, Switzerland 5/13/1994.

[80593] 13932.

[80594] Balazs, I.; Filippi, G.; Rinaldi, A.; Grzeschik, K.-H.; Siniscalco,

[80595] M.: Studies on X-linked ichthyosis and steroid sulfatase in man,

[80596] mice and their hybrids. (Abstract) Cytogenet. Cell Genet. 25: 133,

[80597] 1979.

[80598] 13933.

[80599] Ballabio, A.: Personal Communication. Houston, Tex. 5/18/1994.

[80600] 13934.

[80601] Ballabio, A.; Parenti, G.; Carrozzo, R.; Coppa, G.; Felici, L.;

[80602] Migliori, V.; Silengo, M.; Franceschini, P.; Andria, G.: X/Y translocation

[80603] in a family with X-linked ichthyosis, chondrodysplasia punctata, and

[80604] mental retardation: DNA analysis reveals deletion of the steroid sulphatase

[80605] gene and translocation of its Y pseudogene. Clin. Genet. 34: 31-37,

[80606] 1988.

[80607] 13935.

[80608] Ballabio, A.; Parenti, G.; Carrozzo, R.; Sebastio, G.; Andria,

[80609] G.; Buckle, V.; Fraser, N.; Craig, I.; Rocchi, M.; Romeo, G.; Jobsis,

[80610] A. C.; Persico, M. G.: Isolation and characterization of a steroid

[80611] sulfatase cDNA clone: genomic deletions in patients with

X-chromosome-linked
[80612] ichthyosis. Proc. Nat. Acad. Sci. 84: 4519-4523, 1987.

[80613] 13936.

[80614] Ballabio, A.; Sebastio, G.; Carrozzo, R.; Parenti, G.; Piccir-
illo,
[80615] A.; Persico, M. G.; Andria, G.: Deletions of the steroid sul-
phatase
[80616] gene in 'classical' X-linked ichthyosis and in X-linked
ichthyosis
[80617] associated with Kallmann syndrome. Hum. Genet. 77:
338-341, 1987.
[80618] 13937.

[80619] Broughton, B. C.; Cordonnier, A.; Kleijer, W. J.; Jaspers, N.
G.
[80620] J.; Fawcett, H.; Raams, A.; Garritsen, V. H.; Stary, A.; Avril,
M.-F.;
[80621] Boudsocq, F.; Masutani, C.; Hanaoka, F.; Fuchs, R. P.;
Sarasin, A.;
[80622] Lehmann, A. R.: Molecular analysis of mutations in DNA
polymerase
[80623] eta in xeroderma pigmentosum-variant patients. Proc.
Nat. Acad. Sci. 99:

[80624] 815-820, 2002.

[80625] 13938.

[80626] Haracska, L.; Yu, S.-L.; Johnson, R. E.; Prakash, L.; Prakash,

[80627] S.: Efficient and accurate replication in the presence of 7,8-dihydro-8-oxoguanine

[80628] by DNA polymerase eta. Nature Genet. 25: 458-461, 2000.

[80629] 13939.

[80630] Johnson, R. E.; Prakash, S.; Prakash, L.: Efficient bypass of

[80631] a thymine-thymine dimer by yeast DNA polymerase, Pol-eta. Science 283:

[80632] 1001-1004, 1999.

[80633] 13940.

[80634] Limoli, C. L.; Giedzinski, E.; Morgan, W. F.; Cleaver, J. E.:

[80635] Polymerase eta deficiency in the xeroderma pigmentosum variant uncovers

[80636] an overlap between the S phase checkpoint and double-strand break

[80637] repair. Proc. Nat. Acad. Sci. 97: 7939-7946, 2000.

[80638] 13941.

[80639] Masutani, C.; Araki, M.; Yamada, A.; Kusumoto, R.; Nogi-
mori, T.;

[80640] Maekawa, T.; Iwai, S.; Hanaoka, F.: Xeroderma pigmento-
sum variant

[80641] (XP-V) correcting protein from HeLa cells has a thymine
dimer bypass

[80642] DNA polymerase activity. EMBO J. 18: 3491-3501, 1999.

[80643] 13942.

[80644] Matsuda, T.; Bebenek, K.; Masutani, C.; Hanaoka, F.;
Kunkel, T.

[80645] A.: Low fidelity DNA synthesis by human DNA poly-
merase-eta. Nature 404:

[80646] 1011-1013, 2000.

[80647] 13943.

[80648] Yamada, A.; Masutani, C. Iwai, S.; Hanaoka, F.: Comple-
mentation

[80649] of defective translesion synthesis and UV light sensitivity
in xeroderma

[80650] pigmentosum variant cells by human and mouse DNA
polymerase eta. Nucleic

[80651] Acids Res. 28: 2473-2480, 2000.

[80652] 13944.

[80653] Yuasa, M.; Masutani, C.; Eki, T.; Hanaoka, F.: Genomic structure,

[80654] chromosomal localization and identification of mutations in the xeroderma

[80655] pigmentosum variant (XPV) gene. Oncogene 19: 4721-4728, 2000.

[80656] 13945.

[80657] Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.-H.; Goldstein, L. S.

[80658] : Axonal transport of amyloid precursor protein is mediated by direct

[80659] binding to the kinesin light chain subunit of kinesin-I. Neuron 28:

[80660] 449-459, 2000.

[80661] 13946.

[80662] Luo, Y.; Bolon, B.; Kahn, S.; Bennett, B. D.; Babu-Khan, S.; Denis,

[80663] P.; Fan, W.; Kha, H.; Zhang, J.; Gong, Y.; Martin, L.; Louis, J.-C.;

[80664] Yan, Q.; Richards, W. G.; Citron, M.; Vassar, R.: Mice deficient

[80665] in BACE1, the Alzheimer's beta-secretase, have normal phenotype and

[80666] abolished beta-amyloid generation. Nature Neurosci. 4: 231-232,

[80667] 2001.

[80668] 13947.

[80669] Okamoto, N.; Hori, S.; Akazawa, C.; Hayashi, Y.; Shigemoto, R.;

[80670] Mizuno, N.; Nakanishi, S.: Molecular characterization of a new metabotropic

[80671] glutamate receptor mGluR7 coupled to inhibitory cyclic AMP signal

[80672] transduction. J. Biol. Chem. 269: 1231-1236, 1994.

[80673] 13948.

[80674] Ango, F.; Prezeau, L.; Muller, T.; Tu, J. C.; Xiao, B.; Worley,

[80675] P. F.; Pin, J. P.; Bockaert, J.; Fagni, L.: Agonist-independent activation

[80676] of metabotropic glutamate receptors by the intracellular protein Homer. Nature 411:

[80677] 962-965, 2001.

[80678] 13949.

[80679] Asai, K.; Fujita, K.; Yamamoto, M.; Hotta, T.; Morikawa, M.; Kokubo,

[80680] M.; Moriyama, A.; Kato, T.: Isolation of novel human cDNA (hGMF-gamma)

[80681] homologous to glia maturation factor-beta gene. Biochim. Biophys.

[80682] Acta 1396: 242-244, 1998.

[80683] 13950.

[80684] Peters, N.; Smith, J. S.; Tachibana, I.; Lee, H. K.; Pohl, U.;

[80685] Portier, B. P.; Louis, D. N.; Jenkins, R. B.: The human glia maturation

[80686] factor-gamma gene: genomic structure and mutation analysis in gliomas

[80687] with chromosome 19q loss. Neurogenetics 2: 163-166, 1999.

[80688] 13951.

[80689] Kowalski, P. E.; Freeman, J. D.; Nelson, D. T.; Mager, D. L.:

[80690] Genomic structure and evolution of a novel gene (PLA2L) with duplicated

[80691] phospholipase A2-like domains. Genomics 39: 38-46, 1997.

[80692] 13952.

[80693] Liu, M.; Parker, R. M. C.; Darby, K.; Eyre, H. J.; Copeland, N.

[80694] G.; Crawford, J.; Gilbert, D. J.; Sutherland, G. R.; Jenkins, N. A.;

[80695] Herzog, H.: GPR56, a novel secretin-like human G-protein-coupled

[80696] receptor gene. Genomics 55: 296-305, 1999.

[80697] 13953.

[80698] Zendman, A. J. W.; Cornelissen, I. M. H. A.; Weidle, U. H.; Ruiter,

[80699] D. J.; van Muijen, G. N. P.: TM7XN1, a novel human EGF-TM7-like cDNA,

[80700] detected with mRNA differential display using human melanoma cell

[80701] lines with different metastatic potential. FEBS Lett. 446: 292-298,

[80702] 1999.

[80703] 13954.

[80704] Champliaud, M.-F.; Burgeson, R. E.; Jin, W.; Baden, H. P.; Olson,
[80705] P. F.: cDNA cloning and characterization of sciellin, a LIM domain
[80706] protein of the keratinocyte cornified envelope. J. Biol. Chem. 273:
[80707] 31547-31554, 1998.

[80708] 13955.

[80709] Nicolaou, M.; Song, Y.-Q.; Sato, C. A.; Orlacchio, A.; Kawarai,
[80710] T.; Medeiros, H.; Liang, Y.; Sorbi, S.; Richard, E.; Rogaev, E. I.;
[80711] Moliaka, Y.; Bruni, A. C.; Jorge, R.; Percy, M.; Duara, R.; Farrer,
[80712] L. A.; St George-Hyslop, P.; Rogaeva, E. A.: Mutations in the open
[80713] reading frame of the beta-site APP cleaving enzyme (BACE) locus are
[80714] not a common cause of Alzheimer's disease. Neurogenet-ics 3: 203-206,
[80715] 2001.

[80716] 13956.

[80717] Saunders, A. J.; Kim, T.-W.; Tanzi, R. E.: BACE maps to chromosome
[80718] 11 and a BACE homolog, BACE2, reside in the obligate Down syndrome
[80719] region of chromosome 21. Science 286: 1255a, 1999. Note: Electronic
[80720] Publication.

[80721] 13957.

[80722] Sinha, S.; Anderson, J. P.; Barbour, R.; Basi, G. S.; Cac-
cavello,
[80723] R.; Davis, D.; Doan, M.; Dovey, H. F.; Frigon, N.; Hong, J.; Jacobson-Croak,
[80724] K.; Jewett, N.; and 15 others: Purification and cloning of amyloid
[80725] precursor protein beta-secretase from human brain. Na-
ture 402: 537-540,
[80726] 1999.

[80727] 13958.

[80728] Vassar, R.; Bennett, B. D.; Babu-Khan, S.; Kahn, S.; Men-
diaz,
[80729] E. A.; Dents, P.; Taplow, D. B.; Ross, S.; Amaranta, P.; Loeloff,

[80730] R.; Luo, Y.; Fisher, S.; and 12 others: Beta-secretase cleavage
[80731] of Alzheimer's amyloid precursor protein by the transmembrane aspartic
[80732] protease BACE. Science 286: 735-741, 1999.

[80733] 13959.

[80734] Yan, R.; Bienkowski, M. J.; Shuck, M. E.; Miao, H.; Tory, M. C.;
[80735] Pauley, A. M.; Brashler, J. R.; Stratman, N. C.; Mathews, W. R.; Buhl,
[80736] A. E.; Carter, D. B.; Tomasselli, A. G.; Parodi, L. A.; Heinrikson,
[80737] R. L.; Gurney, M. E.: Membrane-anchored aspartyl protease with Alzheimer's
[80738] disease beta-secretase activity. Nature 402: 533-536, 1999.

[80739] 13960.

[80740] Kusuda, J.; Hirai, M.; Toyoda, A.; Tanuma, R.; Hashimoto, K.:
[80741] Cloning and chromosome mapping of the human casein kinase I gamma-3
[80742] gene (CSNK1G3). Cytogenet. Cell Genet. 83: 101-103,

1998.

[80743] 13961.

[80744] Jagla, K.; Dolle, P.; Mattei, M.-G.; Jagla, T.; Schuhbaur, B.;

[80745] Dretzen, G.; Bellard, F.; Bellard, M.: Mouse Lbx1 and human LBX1

[80746] define a novel mammalian homeobox gene family related to the Drosophila

[80747] lady bird genes. Mech. Dev. 53: 345-356, 1995.

[80748] 13962.

[80749] Mennerich, D.; Schafer, K.; Braun, T.: Pax-3 is necessary but

[80750] not sufficient for lbx1 expression in myogenic precursor cells of

[80751] the limb. Mech. Dev. 73: 147-158, 1998.

[80752] 13963.

[80753] Schafer, K.; Braun, T.: Early specification of limb muscle precursor

[80754] cells by the homeobox gene Lbx1h. Nature Genet. 23: 213-216, 1999.

[80755] 13964.

[80756] Wilson, P. J.; McGlinn, E.; Marsh, A.; Evans, T.; Arnold, J.;

Wright,
[80757] K.; Biden, K.; Young, J.; Wainwright, B.; Wicking, C.; Chenevix-Trench,
[80758] G.: Sequence variants of DLC1 in colorectal and ovarian tumours. Hum.
[80759] Mutat. 15: 156-165, 2000.

[80760] 13965.

[80761] Yuan, B.-Z.; Miller, M. J.; Keck, C. L.; Zimonjic, D. B.; Thorgeirsson,
[80762] S. S.; Popescu, N. C.: Cloning, characterization, and chromosomal
[80763] localization of a gene frequently deleted in human liver cancer (DLC-1)
[80764] homologous to rat RhoGAP. Cancer Res. 58: 2196-2199, 1998.
[80765] 13966.

[80766] Yuan, B.-Z.; Yang, Y.; Keck-Waggoner, C. L.; Zimonjic, D. B.; Thorgeirsson,
[80767] S. S.; Popescu, N. C.: Assignment and cloning of mouse Arhgap7 to
[80768] chromosome 8A4-B2, a conserved syntenic region of human chromosome

[80769] 8p22-p21. Cytogenet. Cell Genet. 87: 189-190, 1999.

[80770] 13967.

[80771] Barbieri, A. M.; Lupo, G.; Bulfone, A.; Andreazzoli, M.; Mariani,
[80772] M.; Fougerousse, F.; Consalez, G. G.; Borsani, G.; Beck-mann, J. S.;
[80773] Barsacchi, G.; Ballabio, A.; Banfi, S.: A homeobox gene, vax2, controls
[80774] the patterning of the eye dorsoventral axis. Proc. Nat. Acad. Sci. 96:
[80775] 10729-10734, 1999.

[80776] 13968.

[80777] Mitsuuchi, Y.; Johnson, S. W.; Sonoda, G.; Tanno, S.; Golemis,
[80778] E. A.; Testa, J. R.: Identification of a chromosome 3p14.3-21.1 gene,
[80779] APPL, encoding an adaptor molecule that interacts with the oncoprotein-serine/threonine
[80780] kinase AKT2. Oncogene 18: 4891-4898, 1999.

[80781] 13969.

[80782] Fleming, R. E.; Sly, W. S.: Ferroportin mutation in autoso-

mal
[80783] dominant hemochromatosis: loss of function, gain in understanding. J.
[80784] Clin. Invest. 108: 521-522, 2001.

[80785] 13970.

[80786] Aita, V. M.; Liang, X. H.; Murty, V. V. V. S.; Pincus, D. L.; Yu,
[80787] W.; Cayanis, E.; Kalachikov, S.; Gilliam, T. C.; Levine, B.: Cloning
[80788] and genomic organization of beclin 1, a candidate tumor suppressor
[80789] gene on chromosome 17q21. Genomics 59: 59-65, 1999.

[80790] 13971.

[80791] Liang, X. H.; Jackson, S.; Seaman, M.; Brown, K.; Kempkes, B.;
[80792] Hibshoosh, H.; Levine, B.: Induction of autophagy and inhibition
[80793] of tumorigenesis by beclin 1. Nature 402: 672-676, 1999.

[80794] 13972.

[80795] Liang, X. H.; Kleeman, L. K.; Jiang, H. H.; Gordon, G.; Goldman,

[80796] J. E.; Berry, G.; Herman, B.; Levine, B.: Protection against fatal

[80797] Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. J.

[80798] Virol. 72: 8586-8596, 1998.

[80799] 13973.

[80800] Liang, X. H.; Yu, J.; Brown, K.; Levine, B.: Beclin 1 contains

[80801] a leucine-rich nuclear export signal that is required for its autophagy

[80802] and tumor suppressor function. Cancer Res. 61: 3443-3449, 2001.

[80803] 13974.

[80804] Sudbrak, R.; Brown, J.; Dobson-Stone, C.; Carter, S.; Ramser, J.;

[80805] White, J.; Healy, E.; Dissanayake, M.; Larregue, M.; Perrus-sel, M.;

[80806] Lehrach, H.; Munro, C. S.; Strachan, T.; Burge, S.; Hovna-nian, A.;

[80807] Monaco, A. P.: Hailey-Hailey disease is caused by muta-tions in ATP2C1

[80808] encoding a novel Ca2+ pump. Hum. Molec. Genet. 9: 1131-1140, 2000.

[80809] 13975.

[80810] Hatamura, I.; Kanauchi, Y.; Takahara, M.; Fujiwara, M.; Muragaki,

[80811] Y.; Ooshima, A.; Ogino, T.: A nonsense mutation in TRPS1 in a Japanese

[80812] family with tricho-rhino-phalangeal syndrome type I. (Letter) Clin.

[80813] Genet. 59: 366-367, 2001.

[80814] 13976.

[80815] Hilton, M. J.; Sawyer, J. M.; Gutierrez, L.; Hogart, A.; Kung,

[80816] T. C.; Wells, D. E.: Analysis of novel and recurrent muta-tions responsible

[80817] for the tricho-rhino-phalangeal syndromes. J. Hum. Genet. 47: 103-106,

[80818] 2002.

[80819] 13977.

[80820] Holmes, D. I. R.; Wahab, N. A.; Mason, R. M.: Identification of

[80821] glucose-regulated genes in human mesangial cells by mRNA differential

[80822] display. Biochem. Biophys. Res. Commun. 238: 179-184,

1997.

[80823] 13978.

[80824] Miller, A. F.; Harvey, S. A. K.; Thies, R. S.; Olson, M. S.: Bone

[80825] morphogenetic protein-9: an autocrine/paracrine cytokine in the liver. J.

[80826] Biol. Chem. 275: 17937-17945, 2000.

[80827] 13979.

[80828] Scott, A. F.: Personal Communication. Baltimore, Md. 7/13/2000.

[80829] 13980.

[80830] Lin, C.-Y.; Anders, J.; Johnson, M.; Sang, Q. A.; Dickson, R. B.

[80831] : Molecular cloning of cDNA for matriptase, a matrix-degrading serine

[80832] protease with trypsin-like activity. J. Biol. Chem. 274: 18231-18236,

[80833] 1999.

[80834] 13981.

[80835] Oberst, M.; Anders, J.; Xie, B.; Singh, B.; Ossandon, M.; Johnson,

[80836] M.; Dickson, R. B.; Lin, C.-Y.: Matriptase and HAI-1 are expressed
[80837] by normal and malignant epithelial cells in vitro and in vivo. Am.
[80838] J. Path. 158: 1301-1311, 2001.

[80839] 13982.

[80840] Kandil, E.; Kohda, K.; Ishibashi, T.; Tanaka, K.; Kasahara, M.
[80841] : PA28 subunits of the mouse proteasome: primary structures and chromosomal
[80842] localization of the genes. Immunogenetics 46: 337-344, 1997.
[80843] 13983.

[80844] Nikaido, T.; Shimada, K.; Shibata, M.; Hata, M.; Sakamoto, M.;
[80845] Takasaki, Y.; Sato, C.; Takahashi, T.; Nishida, Y.: Cloning and nucleotide
[80846] sequence of cDNA for Ki antigen, a highly conserved nuclear protein
[80847] detected with sera from patients with systemic lupus erythematosus. Clin.
[80848] Exp. Immun. 79: 209-214, 1990.

[80849] 13984.

[80850] Kim, Y.-O.; Park, S.-H.; Kang, Y.-J.; Koh, H.-J.; Kim, S.-H.; Park,

[80851] S.-Y.; Sohn, U.; Huh, T.-L.: Assignment of mitochondrial NAD(+)-specific

[80852] isocitrate dehydrogenase beta subunit gene (IDH3B) to human chromosome

[80853] band 20p13 by in situ hybridization and radiation hybrid mapping. Cytogenet.

[80854] Cell Genet. 86: 240-241, 1999.

[80855] 13985.

[80856] Miyake, A.; Mochizuki, S.; Yokoi, H.; Kohda, M.; Furuichi, K.:

[80857] New ether-a-go-go K(+) channel family members localized in human telencephalon. J.

[80858] Biol. Chem. 274: 25018-25025, 1999.

[80859] 13986.

[80860] Acampora, D.; Postiglione, M. P.; Avantaggiato, V.; Di Bonito,

[80861] M.; Vaccarino, F. M.; Michaud, J.; Simeone, A.: Progressive impairment

[80862] of developing neuroendocrine cell lineages in the hypothalamus of

[80863] mice lacking the Orthopedia gene. Genes Dev. 13: 2787-2800, 1999.

[80864] 13987.

[80865] Lin, X.; State, M. W.; Vaccarino, F. M.; Greally, J.; Hass, M.;

[80866] Leckman, J. F.: Identification, chromosomal assignment, and expression

[80867] analysis of the human homeodomain-containing gene Orthopedia (OTP). Genomics 60:

[80868] 96-104, 1999.

[80869] 13988.

[80870] Pessino, A.; Sivori, S.; Bottino, C.; Malaspina, A.; Morelli, L.;

[80871] Moretta, L.; Biassoni, R.; Moretta, A.: Molecular cloning of NKp46:

[80872] a novel member of the immunoglobulin superfamily involved in triggering

[80873] of natural cytotoxicity. J. Exp. Med. 188: 953-960, 1998.

[80874] 13989.

[80875] Sivori, S.; Vitale, M.; Morelli, L.; Sanseverino, L.; Au-

gugliaro,

[80876] R.; Bottino, C.; Moretta, L.; Moretta, A.: p46, a novel natural killer

[80877] cell-specific surface molecule that mediates cell activation. J.

[80878] Exp. Med. 186: 1129-1136, 1997.

[80879] 13990.

[80880] Vankayalapati, R.; Wizel, B.; Weis, S. E.; Safi, H.; Lakey, D.

[80881] L.; Mandelboim, O.; Samten, B.; Porgador, A.; Barnes, P. F.: The

[80882] NKp46 receptor contributes to NK cell lysis of mononuclear phagocytes

[80883] infected with an intracellular bacterium. J. Immun. 168: 3451-3457,

[80884] 2002.

[80885] 13991.

[80886] Chen, X.-Z.; Vassilev, P. M.; Basora, N. Peng, J.-B.; Nomura, H.;

[80887] Segal, Y.; Brown, E. M.; Reeders, S. T.; Hediger, M. A.; Zhou, J.

[80888] : Polycystin-L is a calcium-regulated cation channel per-

meable to
[80889] calcium ions. Nature 401: 383-386, 1999.

[80890] 13992.

[80891] Guo, L.; Chen, M.; Basora, N.; Zhou, J.: The human polycystic
[80892] kidney disease 2-like (PKDL) gene: exon/intron structure and evidence
[80893] for a novel splicing mechanism. Mammalian Genome 11: 46-50, 2000.
[80894] 13993.

[80895] Nomura, H.; Turco, A. E.; Pei, Y.; Kalaydjieva, L.; Schiavello,
[80896] T.; Weremowicz, S.; Ji, W.; Morton, C. C.; Meisler, M.; Reeders, S.
[80897] T.; Zhou, J.: Identification of PKDL, a novel polycystic kidney disease
[80898] 2-like gene whose murine homologue is deleted in mice with kidney
[80899] and retinal defects. J. Biol. Chem. 273: 25967-25973, 1998.
[80900] 13994.

[80901] Wu, G.; Hayashi, T.; Park, J.-H.; Dixit, M.; Reynolds, D. M.;

Li,
[80902] L.; Maeda, Y.; Cai, Y.; Coca-Prados, M.; Somlo, S.: Identification
[80903] of PKD2L, a human PKD2-related gene: tissue-specific expression and
[80904] mapping to chromosome 10q25. Genomics 54: 564-568, 1998.
[80905] 13995.

[80906] Gongora, C.; David, G.; Pintard, L.; Tissot, C.; Hua, T. D.; Dejean,
[80907] A.; Mechti, N.: Molecular cloning of a new interferon-induced PML
[80908] nuclear body-associated protein. J. Biol. Chem. 272: 19457-19463,
[80909] 1997.

[80910] 13996.

[80911] Mattei, M. G.; Tissot, C.; Gongora, C.; Mechti, N.: Assignment
[80912] of ISG20 encoding a new interferon-induced PML nuclear body-associated
[80913] protein, to chromosome 15q26 by in situ hybridization. Cytogenet.

[80914] Cell Genet. 79: 286-287, 1997.

[80915] 13997.

[80916] Pentecost, B. T.: Expression and estrogen regulation of the HEM45

[80917] mRNA in human tumor lines and in the rat uterus. J. Steroid Biochem.

[80918] Molec. Biol. 64: 25-33, 1998.

[80919] 13998.

[80920] Berchtold, S.; Muhl-Zurbes, P.; Heufler, C.; Winklehner, P.; Schuler,

[80921] G.; Steinkasserer, A.: Cloning, recombinant expression and biochemical

[80922] characterization of the murine CD83 molecule which is specifically

[80923] upregulated during dendritic cell maturation. FEBS Lett. 461: 211-216,

[80924] 1999.

[80925] 13999.

[80926] Fujimoto, Y.; Tu, L.; Miller, A. S.; Bock, C.; Fujimoto, M.; Doyle,

[80927] C.; Steeber, D. A.; Tedder, T. F.: CD83 expression influ-

ences CD4+

[80928] T cell development in the thymus. Cell 108: 755-767, 2002.

[80929] 14000.

[80930] Kozlow, E. J.; Wilson, G. L.; Fox, C. H.; Kehrl, J. H.: Subtractive

[80931] cDNA cloning of a novel member of the Ig gene superfamily expressed

[80932] at high levels in activated B lymphocytes. Blood 81: 454-461, 1993.

[80933] 14001.

[80934] Olavesen, M.G.; Bentley, E.; Mason, R. V.; Stephens, R. J.; Ragoussis,

[80935] J.: Fine mapping of 39 ESTs on human chromosome 6p23-p25. Genomics 46:

[80936] 303-306, 1997.

[80937] 14002.

[80938] Scholler, N.; Hayden-Ledbetter, M.; Hellstrom, K.-E.; Hellstrom,

[80939] I.; Ledbetter, J. A.: CD83 is a sialic acid-binding Ig-like lectin

[80940] (Siglec) adhesion receptor that binds monocytes and a

subset of activated
[80941] CD8(+) T cells. J. Immun. 166: 3865-3872, 2001.

[80942] 14003.

[80943] Stankewich, M. C.; Tse, W. T.; Peters, L. L.; Ch'ng, Y.; John,

[80944] K. M.; Stabach, P. R.; Devarajan, P.; Morrow, J. S.; Lux, S.
E.:

[80945] A widely expressed beta-III spectrin associated with Golgi
and cytoplasmic
[80946] vesicles. Proc. Nat. Acad. Sci. 95: 14158-14163, 1998.

[80947] 14004.

[80948] Bakker, A. B. H.; Baker, E.; Sutherland, G. R.; Phillips, J. H.;

[80949] Lanier, L. L.: Myeloid DAP12-associating lectin (MDL)-1 is
a cell
[80950] surface receptor involved in the activation of myeloid
cells. Proc.
[80951] Nat. Acad. Sci. 96: 9792-9796, 1999.

[80952] 14005.

[80953] St-Pierre, M. V.; Hagenbuch, B.; Ugele, B.; Meier, P. J.;
Stallmach,
[80954] T.: Characterization of an organic anion-transporting

polypeptide

[80955] (OATP-B) in human placenta. J. Clin. Endocr. Metab. 87: 1856-1863,

[80956] 2002.

[80957] 14006.

[80958] Reczek, D.; Berryman, M.; Bretscher, A.: Identification of EBP50:

[80959] a PDZ-containing phosphoprotein that associates with members of the

[80960] ezrin-radixin-moesin family. J. Cell Biol. 139: 169-179, 1997.

[80961] 14007.

[80962] Lin, B.; White, J. T.; Ferguson, C.; Bumgarner, R.; Friedman, C.;

[80963] Trask, B.; Ellis, W.; Lange, P.; Hood, L.; Nelson, P. S.: PART-1:

[80964] a novel human prostate-specific, androgen-regulated gene that maps

[80965] to chromosome 5q12. Cancer Res. 60: 858-863, 2000.

[80966] 14008.

[80967] Horowitz, D. S.; Krainer, A. R.: A human protein required

for
[80968] the second step of pre-mRNA splicing is functionally re-
lated to a
[80969] yeast splicing factor. Genes Dev. 11: 139-151, 1997.

[80970] 14009.

[80971] Boucher, C. A.; Winchester, C. L.; Hamilton, G. M.; Winter,
A.
[80972] D.; Johnson, K. J.; Bailey, M. E. S.: Structure, mapping and
expression
[80973] of the human gene encoding the homeodomain protein,
SIX2. Gene 247:
[80974] 145-151, 2000.

[80975] 14010.

[80976] Hasan, S.; Stucki, M.; Hassa, P. O.; Imhof, R.; Gehrig, P.;
Hunziker,
[80977] P.; Hubscher, U.; Hottiger, M. O.: Regulation of human flap
endonuclease-1
[80978] activity by acetylation through the transcriptional coacti-
vator p300. Molec.
[80979] Cell 7: 1221-1231, 2001.

[80980] 14011.

[80981] Hiraoka, L. R.; Harrington, J. J.; Gerhard, D. S.; Lieber, M. R.;
[80982] Hsieh, C.-L.: Sequence of human FEN-1, a structure-specific endonuclease,
[80983] and chromosomal localization of the gene (FEN1) in mouse and human. Genomics 25:
[80984] 220-225, 1995.

[80985] 14012.

[80986] Hosfield, D. J.; Mol, C. D.; Shen, B.; Tainer, J. A.: Structure

[80987] of the DNA repair and replication endonuclease and exonuclease FEN-1:
[80988] coupling DNA and PCNA binding to FEN-1 activity. Cell 95: 135-146,
[80989] 1998.

[80990] 14013.

[80991] Kucherlapati, M.; Yang, K.; Kuraguchi, M.; Zhao, J.; Lia, M.; Heyer,
[80992] J.; Kane, M. F.; Fan, K.; Russell, R.; Brown, A. M. C.; Kneitz, B.;
[80993] Edelmann, W.; Kolodner, R. D.; Lipkin, M.; Kucherlapati, R.: Haploinsufficiency

[80994] of Flap endonuclease (Fen1) leads to rapid tumor progres-
sion. Proc.
[80995] Nat. Acad. Sci. 99: 9924-9929, 2002.

[80996] 14014.

[80997] Otto, C. J.; Almqvist, E.; Hayden, M. R.; Andrew, S. E.: The
'flap'
[80998] endonuclease gene FEN1 is excluded as a candidate gene
implicated
[80999] in the CAG repeat expansion underlying Huntington dis-
ease. Clin.
[81000] Genet. 59: 122-127, 2001.

[81001] 14015.

[81002] Spiro, C.; Pelletier, R.; Rolfsmeier, M. L.; Dixon, M. J.;
Lahue,
[81003] R. S.; Gupta, G.; Park, M. S.; Chen, X.; Mariappan, S. V. S.;
McMurray,
[81004] C. T.: Inhibition of FEN-1 processing by DNA secondary
structure
[81005] at trinucleotide repeats. Molec. Cell 4: 1079-1085, 1999.

[81006] 14016.

[81007] Tishkoff, D. X.; Filosi, N.; Gaida, G. M.; Kolodner, R. D.: A

[81008] novel mutation avoidance mechanism dependent on S. cerevisiae RAD27
[81009] is distinct from DNA mismatch repair. Cell 88: 253-263, 1997.
[81010] 14017.

[81011] Vermeesch, J. R.; Mertens, G.; David, G.; Marynen, P.: Assignment
[81012] of the human glypican gene (GPC1) to 2q35-q37 by fluorescence in situ
[81013] hybridization. Genomics 25: 327-329, 1995.

[81014] 14018.

[81015] Ono, Y.; Ohno, M.; Shimura, Y.: Identification of a putative RNA
[81016] helicase (HRH1), a human homolog of yeast Prp22. Molec. Cell. Biol. 14:
[81017] 7611-7620, 1994.

[81018] 14019.

[81019] Okumura, K.; Nogami, M.; Taguchi, H.; Dean, F. B.; Chen, M.; Pan,
[81020] Z.-Q.; Hurwitz, J.; Shiratori, A.; Murakami, Y.; Ozawa, K.; Eki, T.

[81021] : Assignment of the 36.5-kDa (RFC5), 37-kDa (RFC4), 38-kDa (RFC3),
[81022] and 40-kDa (RFC2) subunit genes of human replication factor C to chromosome
[81023] bands 12q24.2-q24.3, 3q27, 13q12.3-q13, and 7q11.23. Genomics 25:
[81024] 274-278, 1995.

[81025] 14020.

[81026] Peoples, R.; Perez-Jurado, L.; Wang, Y.-K.; Kaplan, P.; Francke,
[81027] U.: The gene for replication factor C subunit 2 (RFC2) is within
[81028] the 7q11.23 Williams syndrome deletion. (Letter) Am. J. Hum. Genet. 58:
[81029] 1370-1373, 1996.

[81030] 14021.

[81031] Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin,
[81032] J.: BASC, a super complex of BRCA1-associated proteins involved in
[81033] the recognition and repair of aberrant DNA structures. Genes Dev. 14:

[81034] 927-939, 2000.

[81035] 14022.

[81036] Thim, L.; Woldike, H. F.; Nielsen, P. F.; Christensen, M.; Lynch-Devaney,

[81037] K.; Podolsky, D. K.: Characterization of human and rat intestinal

[81038] trefoil factor produced in yeast. Biochemistry 34: 4757-4764, 1995.

[81039] 14023.

[81040] Itoh, T.; Mori, T.; Ohkubo, H.; Yamaizumi, M.: A newly identified

[81041] patient with clinical xeroderma pigmentosum phenotype has a non-sense

[81042] mutation in the DDB2 gene and incomplete repair in (6-4) photoproducts. J.

[81043] Invest. Derm. 113: 251-257, 1999.

[81044] 14024.

[81045] Fernandes-Alnemri, T.; Armstrong, R. C.; Krebs, J.; Srinivasula,

[81046] S. M.; Wang, L.; Bullrich, F.; Fritz, L. C.; Trapani, J. A.; Tomaselli,

[81047] K. J.; Litwack, G.; Alnemri, E. S.: In vitro activation of CPP32
[81048] and Mch3 by Mch4, a novel human apoptotic cysteine protease containing
[81049] two FADD-like domains. Proc. Nat. Acad. Sci. 93: 7464-7469, 1996.
[81050] 14025.

[81051] Fernandes-Alnemri, T.; Litwack, G.; Alnemri, E. S.: CPP32, a novel
[81052] human apoptotic protein with homology to Caenorhabditis elegans cell
[81053] death protein Ced-3 and mammalian interleukin-1 beta-converting enzyme. J.
[81054] Biol. Chem. 269: 30761-30764, 1994.

[81055] 14026.

[81056] Fernando, P.; Kelly, J. F.; Balazsi, K.; Slack, R. S.; Megeney,

[81057] L. A.: Caspase 3 activity is required for skeletal muscle differentiation. Proc.
[81058] Nat. Acad. Sci. 99: 11025-11030, 2002.

[81059] 14027.

[81060] Huang, Y.; Shin, N.-H.; Sun, Y.; Wang, K. K. W.: Molecular

cloning

[81061] and characterization of a novel caspase-3 variant that attenuates

[81062] apoptosis induced by proteasome inhibition. Biochem. Biophys. Res.

[81063] Commun. 283: 762-769, 2001.

[81064] 14028.

[81065] Kuida, K.; Zheng, T. S.; Na, S.; Kuan, C.; Yang, D.; Karasuyama,

[81066] H.; Rakio, P.; Flavell, R. A.: Decreased apoptosis in the brain and

[81067] premature lethality in CPP32-deficient mice. Nature 384: 368-372,

[81068] 1996.

[81069] 14029.

[81070] Levkau, B.; Koyama, H.; Raines, E. W.; Clurman, B. E.; Herren,

[81071] B.; Orth, K.; Roberts, J. M.; Ross, R.: Cleavage of p21(Cip1/Waf1)

[81072] and p27(Kip1) mediates apoptosis in endothelial cells through activation

[81073] of Cdk2: role of a caspase cascade. Molec. Cell 1:

553-563, 1998.

[81074] 14030.

[81075] Nasir, J.; Theilmann, J. L.; Chopra, V.; Jones, A. M.; Walker,

[81076] D.; Rasper, D. M.; Vaillancourt, J. P.; Hewitt, J. E.; Nichol-

son,

[81077] D. W.; Hayden, M. R.: Localization of the cell death genes

CPP32

[81078] and Mch-2 to human chromosome 4q. Mammalian

Genome 8: 56-59, 1997.

[81079] 14031.

[81080] Tiso, N.; Pallavicini, A.; Muraro, T.; Zimbello, R.; Apolloni,

[81081] E.; Valle, G.; Lanfranchi, G.; Danieli, G. A.: Chromosomal

localization

[81082] of the human genes, CPP32, Mch2, Mch3, and Ich-1, in-

volved in cellular

[81083] apoptosis. Biochem. Biophys. Res. Commun. 225:

983-989, 1996.

[81084] 14032.

[81085] Woo, M.; Hakem, R.; Soengas, M. S.; Duncan, G. S.;

Shahinian,

[81086] A.; Kagi, K.; Hakem, A.; McCurrach, M.; Khoo, W.; Kauf-

man, S. A.;

[81087] Senaldi, G.; Howard, T.; Lowe, S. W.; Mak, T. W.: Essential contribution

[81088] of caspase 3/CPP32 to apoptosis and its associated nuclear changes. Genes

[81089] Dev. 12: 806-819, 1998.

[81090] 14033.

[81091] Nicholson, D. W.; Ali, A.; Thornberry, N. A.; Vaillancourt, J.

[81092] P.; Ding, C. K.; Gallant, M.; Gareau, Y.; Griffin, P. R.; La-

belle,

[81093] M.; Lazebnik, Y. A.; Munday, N. A.; Raju, S. M.; Smulson, M. E.; Yamin,

[81094] T.-T.; Yu, V. L.; Miller, D. K.: Identification and inhibition of

[81095] the ICE/CED-3 protease necessary for mammalian apoptosis. Nature 376:

[81096] 37-43, 1995. MEDLINE UID: 95319529

[81097] 14034.

[81098] Bergeron, L.; Perez, G. I.; Macdonald, G.; Shi, L.; Sun, Y.; Jurisicova,

[81099] A.; Varmuza, S.; Latham, K. E.; Flaws, J. A.; Salter, J. C. M.;

Hara,
[81100] H.; Moskowitz, M. A.; Li, E.; Greenberg, A.; Tilly, J. L.; Yuan, J.
[81101] : Defects in regulation of apoptosis in caspase-2-deficient mice. Genes
[81102] Dev. 12: 1304-1314, 1998.

[81103] 14035.

[81104] Stone, D. M.; Hynes, M.; Armanini, M.; Swanson, T. A.; Gu, Q.;
[81105] Johnson, R. L.; Scott, M. P.; Pennica, D.; Goddard, A.; Phillips,
[81106] H.; Noll, M.; Hooper, J. E.; de Sauvage, F.; Rosenthal, A.: The tumour-suppressor
[81107] gene patched encodes a candidate receptor for Sonic hedgehog. Nature 384:
[81108] 129-134, 1996.

[81109] 14036.

[81110] Zhang, X. M.; Ramalho-Santos, M.; McMahon, A. P.: Smoothened mutants
[81111] reveal redundant roles for Shh and Ihh signaling including regulation
[81112] of L/R asymmetry by the mouse node. Cell 105: 781-792,

2001.

[81113] 14037.

[81114] Geurts, J. M. W.; Schoenmakers, E. F. P. M.; Roijer, E.; Astrom,

[81115] A.-K.; Stenman, G.; van de Ven, W. J. M.: Identification of NFIB

[81116] as recurrent translocation partner gene of HMGIC in pleomorphic adenomas. Oncogene 16:

[81117] 865-872, 1998.

[81118] 14038.

[81119] Smith, S. A.; Holik, P. R.; Stevens, J.; Melis, R.; White, R.;

[81120] Albertsen, H.: Isolation and mapping of a gene encoding a novel human

[81121] ADP-ribosylation factor on chromosome 17q12-q21. Genomics 28: 113-115,

[81122] 1995.

[81123] 14039.

[81124] Fiedorek, F. T., Jr.; Kay, E. S.: Mapping of the insulin promoter

[81125] factor 1 gene (Ipf1) to distal mouse chromosome 5. Genomics 28:

[81126] 581-584, 1995.

[81127] 14040.

[81128] Hani, E. H.; Stoffers, D. A.; Chevre, J.-C.; Durand, E.;
Stanojevic,
[81129] V.; Dina, C.; Habener, J. F.; Froguel, P.: Defective muta-
tions in
[81130] the insulin promoter factor-1 (IPF-1) gene in late-onset
type 2 diabetes
[81131] mellitus. J. Clin. Invest. 104: R41-R48, 1999.

[81132] 14041.

[81133] Tang, J. Y.; Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu,
G.
[81134] : Xeroderma pigmentosum p48 gene enhances global ge-
nomic repair and
[81135] suppresses UV-induced mutagenesis. Molec. Cell 5:
737-744, 2000.
[81136] 14042.

[81137] Bermingham, J. R., Jr.; Arden, K. C.; Naumova, A. K.;
Sapienza,
[81138] C.; Viars, C. S.; Fu, X.-D.; Khotz, J.; Manley, J. L.; Rosen-
feld,

[81139] M. G.: Chromosomal localization of mouse and human genes encoding
[81140] the splicing factors ASF/SF2 (SFRS1) and SC-35 (SFRS2). Genomics 29:
[81141] 70-79, 1995.

[81142] 14043.

[81143] Wang, J.; Takagaki, Y.; Manley, J. L.: Targeted disruption of
[81144] an essential vertebrate gene: ASF/SF2 is required for cell viability. Genes
[81145] Dev. 10: 2588-2599, 1996.

[81146] 14044.

[81147] Costanzo, V.; Robertson, K.; Bibikova, M.; Kim, E.; Grieco, D.;
[81148] Gottesman, M.; Carroll, D.; Gautier, J.: Mre11 protein complex prevents
[81149] double-strand break accumulation during chromosomal DNA replication. Molec.
[81150] Cell 8: 137-147, 2001.

[81151] 14045.

[81152] Hernandez, D.; McConville, C. M.; Stacey, M.; Woods, C. G.; Brown,

[81153] M. M.; Shutt, P.; Rysiecki, G.; Taylor, A. M. R.: A family showing
[81154] no evidence of linkage between the ataxia telangiectasia gene and
[81155] chromosome 1q22-23. J. Med. Genet. 30: 135-140, 1993.

[81156] 14046.

[81157] Hopfner, K.-P.; Karcher, A.; Craig, L.; Woo, T. T.; Carney, J.

[81158] P.; Tainer, J. A.: Structural biochemistry and interaction architecture
[81159] of the DNA double-strand break repair Mre11 nuclease and Rad50-ATPase. Cell 105:
[81160] 473-485, 2001.

[81161] 14047.

[81162] Klein, C.; Wenning, G. K.; Quinn, N. P.; Marsden, C. D.: Ataxia
[81163] without telangiectasia masquerading as benign hereditary chorea. Mov.
[81164] Disord. 11: 217-220, 1996.

[81165] 14048.

[81166] Paull, T. T.; Gellert, M.: The 3-prime to 5-prime exonuclease

[81167] activity of Mre11 facilitates repair of DNA double-strand breaks. Molec.

[81168] Cell 1: 969-979, 1998.

[81169] 14049.

[81170] Petrini, J. H. J.; Walsh, M. E.; DiMare, C.; Chen, X.-N.; Korenberg,

[81171] J. R.; Weaver, D. T.: Isolation and characterization of the human

[81172] MRE11 homologue. Genomics 29: 80-86, 1995.

[81173] 14050.

[81174] Pitts, S. A.; Kullar, H. S.; Stankovic, T.; Stewart, G. S.; Last,

[81175] J. I. K.; Bedenham, T.; Armstrong, S. J.; Piane, M.; Chessa, L.; Taylor,

[81176] A. M. R.; Byrd, P. J.: hMRE11: genomic structure and a null mutation

[81177] identified in a transcript protected from nonsense-mediated mRNA decay. Hum.

[81178] Molec. Genet. 10: 1155-1162, 2001.

[81179] 14051.

[81180] Stewart, G. S.; Maser, R. S.; Stankovic, T.; Bressan, D. A.; Kaplan,

[81181] M. I.; Jaspers, N. G. J.; Raams, A.; Byrd, P. J.; Petrini, J. H. J.;

[81182] Taylor, A. M. R.: The DNA double-strand break repair gene hMRE11

[81183] is mutated in individuals with an ataxia-telangiec-tasia-like disorder. Cell 99:

[81184] 577-587, 1999.

[81185] 14052.

[81186] Stracker, T. H.; Carson, C. T.; Weitzman, M. D.: Adenovirus oncoproteins

[81187] inactivate the Mre11-Rad50-NBS1 DNA repair complex. Nature 418:

[81188] 348-352, 2002.

[81189] 14053.

[81190] Trujillo, K. M.; Yuan, S.-S. F.; Lee, E. Y.-H. P.; Sung, P.:

[81191] Nuclease activities in a complex of human recombination and DNA repair

[81192] factors Rad50, Mre11, and p95. J. Biol. Chem. 273: 21447-21450,

[81193] 1998.

[81194] 14054.

[81195] Zhong, Q.; Chen, C.-F.; Li, S.; Chen, Y.; Wang, C.-C.; Xiao, J.;

[81196] Chen, P.-L.; Sharp, Z. D.; Lee, W.-H.: Association of BRCA1 with

[81197] the hRad50-hMre11-p95 complex and the DNA damage response. Science 285:

[81198] 747-750, 1999.

[81199] 14055.

[81200] Zhu, X.-D.; Kuster, B.; Mann, M.; Petrini, J. H. J.; de Lange,

[81201] T.: Cell-cycle-regulated association of RAD50/MRE11/NBS1 with TRF2

[81202] and human telomeres. Nature Genet. 25: 347-352, 2000.

[81203] 14056.

[81204] Virbasius, C. A.; Virbasius, J. V.; Scarpulla, R. C.: NRF-1, an

[81205] activator involved in nuclear-mitochondrial interactions, utilizes

[81206] a new DNA-binding domain conserved in a family of de-velopmental regulators. Genes

[81207] Dev. 7: 2431-2445, 1993.

[81208] 14057.

[81209] Burch, G. H.; Gong, Y.; Curry, C.; Miller, W. L.; Bristow, J. D.

[81210] : Human tenascin-X deficiency causes an Ehlers-Dan-los-like phenotype.

[81211] (Abstract) Am. J. Hum. Genet. 59 (suppl.): A16, 1996.

[81212] 14058.

[81213] Albig, W.; Bramlage, B.; Gruber, K.; Klobeck, H.-G.; Kunz, J.;

[81214] Doenecke, D.: The human replacement histone H3.3B gene (H3F3B). Genomics 30:

[81215] 264-272, 1995.

[81216] 14059.

[81217] Bass, B. L.; Weintraub, H.: An unwinding activity that cova-lently

[81218] modifies its double-stranded RNA substrate. Cell 55: 1089-1098,

[81219] 1988.

[81220] 14060.

[81221] Kim, U.; Wang, Y.; Sanford, T.; Zeng, Y.; Nishikura, K.: Molecular

[81222] cloning of cDNA for double-stranded RNA adenosine

deaminase, a candidate
[81223] enzyme for nuclear RNA editing. Proc. Nat. Acad. Sci. 91:
11457-11461,
[81224] 1994.

[81225] 14061.

[81226] O'Connell, M. A.; Krause, S.; Higuchi, M.; Hsuan, J. J.;
Totty,
[81227] N. F.; Jenny, A.; Keller, W.: Cloning of cDNAs encoding
mammalian
[81228] double-stranded RNA-specific adenosine deaminase.
Molec. Cell. Biol. 15:
[81229] 1389-1397, 1995.

[81230] 14062.

[81231] Patterson, J. B.; Samuel, C. E.: Expression and regulation
by
[81232] interferon of a double-stranded-RNA-specific adenosine
deaminase from
[81233] human cells: evidence for two forms of the deaminase.
Molec. Cell.
[81234] Biol. 15: 5376-5388, 1995.

[81235] 14063.

[81236] Wang, Q.; Khillan, J.; Gadue, P.; Nishikura, K.: Requirement of
[81237] the RNA editing deaminase ADAR1 gene for embryonic erythropoiesis. Science 290:
[81238] 1765-1768, 2000.

[81239] 14064.

[81240] Wang, Y.; Zeng, Y.; Murray, J. M.; Nishikura, K.: Genomic organization
[81241] and chromosomal location of the human dsRNA adenosine deaminase gene:
[81242] the enzyme for glutamate-activated ion channel RNA editing. J. Molec.
[81243] Biol. 254: 184-195, 1995.

[81244] 14065.

[81245] Weier, H.-U. G.; George, C. X.; Greulich, K. M.; Samuel, C. E.
[81246] : The interferon-inducible, double-stranded RNA-specific adenosine
[81247] deaminase gene (DSRAD) maps to human chromosome 1q21.1-21.2. Genomics 30:
[81248] 372-375, 1995.

[81249] 14066.

[81250] Weier, H.-U. G.; George, C. X.; Lersch, R. A.; Breitweser, S.;

[81251] Cheng, J.-F.; Samuel, C. E.: Assignment of the RNA-
specific adenosine

[81252] deaminase gene (Adar) to mouse chromosome 3F2 by in
situ hybridization. Cytogenet.

[81253] Cell Genet. 89: 214-215, 2000.

[81254] 14067.

[81255] Lin, Q.; Schwarz, J.; Bucana, C.; Olson, E. N.: Control of
mouse

[81256] cardiac morphogenesis and myogenesis by transcription
factor MEF2C. Science 276:

[81257] 1404-1407, 1997.

[81258] 14068.

[81259] Burch, G. H.; Gong, Y.; Liu, W.; Dettman, R. W.; Curry, C. J.;

[81260] Smith, L.; Miller, W. L.; Bristow, J.: Tenascin-X deficiency is
associated

[81261] with Ehlers-Danlos syndrome. Nature Genet. 17:
104-108, 1997.

[81262] 14069.

[81263] Erickson, H. P.: A tenascin knockout with a phenotype. Nature
[81264] Genet. 17: 5-7, 1997.

[81265] 14070.

[81266] Gitelman, S. E.; Bristow, J.; Miller, W. L.: Mechanism and consequences
[81267] of the duplication of the human C4/P450c21/gene X locus. Molec. Cell.
[81268] Biol. 12: 2124-2134, 1992.

[81269] 14071.

[81270] Ikuta, T.; Sogawa, N.; Ariga, H.; Ikemura, T.; Matsumoto, K.:
[81271] Structural analysis of mouse tenascin-X: evolutionary aspects of reduplication
[81272] of FNIII repeats in the tenascin gene family. Gene 217: 1-13, 1998.
[81273] 14072.

[81274] Mao, J.-R.; Bristow, J.: The Ehlers-Danlos syndrome: on beyond
[81275] collagens. J. Clin. Invest. 107: 1063-1069, 2001.

[81276] 14073.

[81277] Mao, J. R.; Taylor, G.; Dean, W. B.; Wagner, D. R.; Afzal, V.;

[81278] Lotz, J. C.; Rubin, E. M.; Bristow, J.: Tenascin-X deficiency mimics

[81279] Ehlers-Danlos syndrome in mice through alteration of collagen deposition. Nature

[81280] Genet. 30: 421-425, 2002.

[81281] 14074.

[81282] Matsumoto, K.; Arai, M.; Ishihara, N.; Ando, A.; Inoko, H.; Ikemura,

[81283] T.: Cluster of fibronectin type III repeats found in the human major

[81284] histocompatibility complex class III region shows the highest homology

[81285] with the repeats in an extracellular matrix protein, tenascin. Genomics 12:

[81286] 485-491, 1992.

[81287] 14075.

[81288] Matsumoto, K.; Saga, Y.; Ikemura, T.; Sakakura, T.; Chiquet-Ehrismann,

[81289] R.: The distribution of tenascin-X is distinct and often reciprocal

[81290] to that of tenascin-C. J. Cell Biol. 125: 483-493, 1994.

[81291] 14076.

[81292] Saga, Y.; Yagi, T.; Ikawa, Y.; Sakakura, T.; Aizawa, S.: Mice

[81293] develop normally without tenascin. Genes Dev. 6:
1821-1831, 1992.

[81294] 14077.

[81295] Schalkwijk, J.; Zweers, M. C.; Steijlen, P. M.; Dean, W. B.;
Taylor,

[81296] G.; van Vlijmen, I. M.; van Haren, B.; Miller, W. L.; Bristow,
J.

[81297] : A recessive form of the Ehlers-Danlos syndrome caused
by tenascin-X

[81298] deficiency. New Eng. J. Med. 345: 1167-1175, 2001.

[81299] 14078.

[81300] Speek, M.; Barry, F.; Miller, W. L.: Alternate promoters and

[81301] alternate splicing of human tenascin-X, a gene with
5-prime and 3-prime

[81302] ends buried in other genes. Hum. Molec. Genet. 5:
1749-1758, 1996.

[81303] 14079.

[81304] Tee, M. K.; Thomson, A. A.; Bristow, J.; Miller, W. L.: Sequences
[81305] promoting the transcription of the human XA gene overlapping P450c21A
[81306] correctly predict the presence of a novel, adrenal-specific, truncated
[81307] form of tenascin-X. Genomics 28: 171-178, 1995.

[81308] 14080.

[81309] Xu, X.; Doolittle, R. F.: Presence of vertebrate fibrinogen-like
[81310] sequence in an echinoderm. Proc. Nat. Acad. Sci. 87: 2097-2101,
[81311] 1990.

[81312] 14081.

[81313] Akama, T. O.; Nakagawa, H.; Sugihara, K.; Narisawa, S.; Ohyama,
[81314] C.; Nishimura, S.-I.; O'Brien, D. A.; Moremen, K. W.; Millan, J. L.;
[81315] Fukuda, M. N.: Germ cell survival through carbohydrate-mediated interaction
[81316] with Sertoli cells. Science 295: 124-127, 2002.

[81317] 14082.

[81318] Bai, R.-Y.; Koester, C.; Ouyang, T.; Hahn, S. A.; Hammer-
schmidt,
[81319] M.; Peschel, C.; Duyster, J.: SMIF, a Smad4-interacting
protein that
[81320] functions as a co-activator in TGF-beta signalling. Nature
Cell Biol. 4:
[81321] 181-190, 2002.

[81322] 14083.

[81323] Akiyama, Y.; Tsubouchi, N.; Yuasa, Y.: Frequent somatic
mutations
[81324] of hMSH3 with reference to microsatellite instability in
hereditary
[81325] nonpolyposis colorectal cancer. Biochem. Biophys. Res.
Commun. 236:
[81326] 248-252, 1997.

[81327] 14084.

[81328] Chittenden, T.; Harrington, E. A.; O'Connor, R.; Fleming-
ton, C.;
[81329] Lutz, R. J.; Evan, G. I.; Guild, B. C.: Induction of apoptosis
by

[81330] the Bcl-2 homologue Bak. Nature 374: 733-736, 1995.

[81331] 14085.

[81332] Herberg, J. A.; Phillips, S.; Beck, S.; Jones, T.; Sheer, D.; Wu,

[81333] J. J.; Prochazka, V.; Barr, P. J.; Kiefer, M. C.; Trowsdale, J.:

[81334] Genomic structure and domain organisation of the human Bak gene. Gene 211:

[81335] 87-94, 1998.

[81336] 14086.

[81337] Kiefer, M. C.; Brauer, M. J.; Powers, V. C.; Wu, J. J.; Uman-

[81338] sky, S. R.; Tomei, L. D.; Barr, P. J.: Modulation of apoptosis by the

[81339] widely distributed Bcl-2 homologue Bak. Nature 374: 736-739, 1995.

[81340] 14087.

[81341] Ulrich, E.; Kauffmann-Zeh, A.; Hueber, A.-O.; Williamson, J.; Chittenden,

[81342] T.; Ma, A.; Evan, G.: Gene structure, cDNA sequence, and expression

[81343] of murine Bak, a proapoptotic Bcl-2 family member. Ge-

nomics 44:
[81344] 195-200, 1997.

[81345] 14088.

[81346] Saito, H.; Fujiwara, T.; Takahashi, E.; Shin, S.; Okui, K.; Nakamura,
[81347] Y.: Isolation and mapping of a novel human gene encoding a protein
[81348] containing zinc-finger structures. Genomics 31: 376-379, 1996.
[81349] 14089.

[81350] Higuchi, M.; Maas, S.; Single, F. N.; Hartner, J.; Rozov, A.; Burnashev,
[81351] N.; Feldmeyer, D.; Sprengel, R.; Seeburg, P. H.: Point mutation in
[81352] an AMPA receptor gene rescues lethality in mice deficient in the RNA-editing
[81353] enzyme ADAR2. Nature 406: 78-81, 2000.

[81354] 14090.

[81355] Lai, F.; Chen, C.-X.; Carter, K. C.; Nishikura, K.: Editing of
[81356] glutamate receptor B subunit ion channel RNAs by four alternatively

[81357] spliced DRADA2 double-stranded RNA adenosine deaminases. Molec. Cell.

[81358] Biol. 17: 2413-2424, 1997.

[81359] 14091.

[81360] Melcher, T.; Maas, S.; Herb, A.; Sprengel, R.; Seeburg, P. H.;

[81361] Higuchi, M.: A mammalian RNA editing enzyme. Nature 379: 460-463,

[81362] 1996.

[81363] 14092.

[81364] Mittaz, L.; Scott, H. S.; Rossier, C.; Seeburg, P. H.; Higuchi,

[81365] M.; Antonarakis, S. E.: Cloning of a human RNA editing deaminase

[81366] (ADARB1) of glutamate receptors that maps to chromosome 21q22.3. Genomics 41:

[81367] 210-217, 1997.

[81368] 14093.

[81369] O'Connell, M. A.; Gerber, A.; Keller, W.: Purification of human

[81370] double-stranded RNA-specific editase 1 (hRED1) involved in editing

[81371] of brain glutamate receptor B pre-mRNA. J. Biol. Chem. 272: 473-478,
[81372] 1997.

[81373] 14094.

[81374] Villard, L.; Tassone, F.; Haymowicz, M.; Welborn, R.; Gar-
diner,
[81375] K.: Map location, genomic organization and expression patterns of
[81376] the human RED1 RNA editase. Somat. Cell Molec. Genet. 23: 135-145,
[81377] 1997.

[81378] 14095.

[81379] Yang, J.-H.; Sklar, P.; Axel, R.; Maniatis, T.: Purification and
[81380] characterization of a human RNA adenosine deaminase for glutamate
[81381] receptor B pre-mRNA editing. Proc. Nat. Acad. Sci. 94: 4354-4359,
[81382] 1997.

[81383] 14096.

[81384] Zhao, Z.; Lee, C.-C.; Monckton, D. G.; Yazdani, A.; Cool-

baugh,
[81385] M. I.; Li, X.; Bailey, J.; Shen, Y.; Caskey, C. T.: Characterization
[81386] and genomic mapping of genes and pseudogenes of a new human protein
[81387] tyrosine phosphatase. Genomics 35: 172-181, 1996.

[81388] 14097.

[81389] Chopra, V. S.; Metzler, M.; Rasper, D. M.; Engqvist-Goldstein,
[81390] A. E. Y.; Singaraja, R.; Gan, L.; Fichter, K. M.; McCutcheon, K.;
[81391] Drubin, D.; Nicholson, D. W.; Hayden, M. R.: HIP12 is a non-proapoptotic
[81392] member of a gene family including HIP1, an interacting protein with
[81393] huntingtin. Mammalian Genome 11: 1006-1015, 2000.

[81394] 14098.

[81395] Bogdanova, N.; Markoff, A.; Gerke, V.; McCluskey, M.; Horst, J.;
[81396] Dworniczak, B.: Homologues to the first gene for autosomal dominant
[81397] polycystic kidney disease are pseudogenes. Genomics 74:

333-341,

[81398] 2001.

[81399] 14099.

[81400] Boletta, A.; Qian, F.; Onuchic, L. F.; Bhunia, A. K.;
Phakdeekitcharoen,

[81401] B.; Hanaoka, K.; Guggino, W.; Monaco, L.; Germino, G. G.:
Polycystin-1,

[81402] the gene product of PKD1, induces resistance to apoptosis
and spontaneous

[81403] tubulogenesis in MDCK cells. Molec. Cell 6: 1267-1273,
2000.

[81404] 14100.

[81405] Bouba, I.; Koptides, M.; Mean, R.; Costi, C.-E.; Demetriou,
K.;

[81406] Georgiou, I.; Pierides, A.; Siamopoulos, K.; Deltas, C. C.:
Novel

[81407] PKD1 deletions and missense variants in a cohort of Hel-
lenic polycystic

[81408] kidney disease families. Europ. J. Hum. Genet. 9:
677-684, 2001.

[81409] 14101.

[81410] Boulter, C.; Mulroy, S.; Webb, S.; Fleming, S.; Brindle, K.;

Sandford,
[81411] R.: Cardiovascular, skeletal, and renal defects in mice with a targeted
[81412] disruption of the Pkd1 gene. Proc. Nat. Acad. Sci. 98: 12174-12179,
[81413] 2001.

[81414] 14102.

[81415] Constantinides, R.; Xenophontos, S.; Neophytou, P.; No-mura, S.;
[81416] Pierides, A.; Constantinou Deltas, C.: New amino acid polymorphism,
[81417] Ala/Val4058, in exon 45 of the polycystic kidney disease 1 gene: evolution
[81418] of alleles. Hum. Genet. 99: 644-647, 1997.

[81419] 14103.

[81420] Delgado-Rodriguez, A.; Ortiz-Marttelo, R.; Villalobos-Pietrini,
[81421] R.; Gomez-Arroyo, S.; Graf, U.: Genotoxicity of organic extracts
[81422] of airborne particles in somatic cells of Drosophila melanogaster. Chemosphere 39:
[81423] 33-43, 1999.

[81424] 14104.

[81425] Germino, G. G.: Personal Communication. Baltimore, Md. 7/30/1998.

[81426] 14105.

[81427] Huan, Y.; van Adelsberg, J.: Polycystin-1, the PKD1 gene product,

[81428] is in a complex containing E-cadherin and the catenins. J. Clin.

[81429] Invest. 104: 1459-1468, 1999.

[81430] 14106.

[81431] Ibraghimov-Beskrovnaya, O.; Bukanov, N. O.; Donohue, L. C.; Dackowski,

[81432] W. R.; Klinger, K. W.; Landes, G. M.: Strong homophilic interactions

[81433] of the Ig-like domains of polycystin-1, the protein product of an

[81434] autosomal dominant polycystic kidney disease gene, PKD1. Hum. Molec.

[81435] Genet. 9: 1641-1649, 2000.

[81436] 14107.

[81437] Ma, T.; Yang, B.; Kuo, W.-L.; Verkman, A. S.: cDNA cloning

and
[81438] gene structure of a novel water channel expressed exclusively in human
[81439] kidney: evidence for a gene cluster of aquaporins at chromosome locus
[81440] 12q13. Genomics 35: 543-550, 1996.

[81441] 14108.

[81442] Bolino, A.; Muglia, M.; Conforti, F. L.; LeGuern, E.; Salih, M.

[81443] A. M.; Georgiou, D.-M.; Christodoulou, K.; Hausmanowa-Petrusewicz,
[81444] I.; Mandich, P.; Schenone, A.; Gambardella, A.; Bono, F.; Quattrone,
[81445] A.; Devoto, M.; Monaco, A. P.: Charcot-Marie-Tooth type 4B is caused
[81446] by mutations in the gene encoding myotubularin-related protein-2. Nature
[81447] Genet. 25: 17-19, 2000.

[81448] 14109.

[81449] Yasui, M.; Kwon, T.-H.; Knepper, M. A.; Nielsen, S.; Agre, P.:
[81450] Aquaporin-6: an intracellular vesicle water channel pro-

tein in renal

[81451] epithelia. Proc. Nat. Acad. Sci. 96: 5808-5813, 1999.

[81452] 14110.

[81453] Shan, X.; Bourdeau, A.; Rhoton, A.; Wells, D. E.; Cohen, E. H.;

[81454] Landgraf, B. E.; Palfree, R. G. E.: Characterization and mapping

[81455] to human chromosome 8q24.3 of Ly-6-related gene 9804 encoding an apparent

[81456] homologue of mouse TSA-1. J. Immun. 160: 197-208, 1998.

[81457] 14111.

[81458] Diamond, R. H.; Cressman, D. E.; Laz, T. M.; Abrams, C. S.; Taub,

[81459] R.: PRL-1, a unique nuclear protein tyrosine phosphatase, affects

[81460] cell growth. Cell. Biol. 14: 3752-3762, 1994.

[81461] 14112.

[81462] Peng, Y.; Genin, A.; Spinner, N. B.; Diamond, R. H.; Taub, R.:

[81463] The gene encoding human nuclear protein tyrosine phos-

phatase, PRL-1:
[81464] cloning, chromosomal localization, and identification of an intron
[81465] enhancer. J. Biol. Chem. 273: 17286-17295, 1998.

[81466] 14113.

[81467] Hatakeyama, S.; Osawa, M.; Omine, M.; Ishikawa, F.: JTB: a novel
[81468] membrane protein gene at 1q21 rearranged in a jumping translocation. Oncogene 18:
[81469] 2085-2090, 1999.

[81470] 14114.

[81471] Jadayel, D. M.; Osborne, L. R.; Coignet, L. J. A.; Zani, V. J.;

[81472] Tsui, L.-C.; Scherer, S. W.; Dyer, M. J. S.: The BCL7 gene family:
[81473] deletion of BCL7B in Williams syndrome. Gene 224:
35-44, 1998.
[81474] 14115.

[81475] Zani, V. J.; Asou, N.; Jadayel, D.; Heward, J. M.; Shipley, J.;

[81476] Nacheva, E.; Takasuki, K.; Catovsky, D.; Dyer, M. J. S.: Molecular
[81477] cloning of complex chromosomal translocation

t(8;14;12)(q24.1;q32.3;q24.1)

[81478] in a Burkitt lymphoma cell line defines a new gene (BCL7A) with homology

[81479] to caldesmon. Blood 87: 3124-3134, 1996.

[81480] 14116.

[81481] Joung, I.; Strominger, J. L.; Shin, J.: Molecular cloning of a

[81482] phosphotyrosine-independent ligand of the p56-lck SH2 domain. Proc.

[81483] Nat. Acad. Sci. 93: 5991-5995, 1996.

[81484] 14117.

[81485] Laurin, N.; Brown, J. P.; Morissette, J.; Raymond, V.: Recur-rent

[81486] mutation of the gene encoding sequestosome 1 (SQSTM1/p62) in Paget

[81487] disease of bone. Am. J. Hum. Genet. 70: 1582-1588, 2002.

[81488] 14118.

[81489] Park, I.; Chung, J.; Walsh, C. T.; Yun, Y.; Strominger, J. L.;

[81490] Shin, J.: Phosphotyrosine-independent binding of a 62-kDa protein

[81491] to the src homology 2 (SH2) domain of p56-lck and its

regulation by
[81492] phosphorylation of ser-59 in the lck unique N-terminal region. Proc.
[81493] Nat. Acad. Sci. 92: 12338-12342, 1995.

[81494] 14119.

[81495] Vadlamudi, R. K.; Joung, I.; Strominger, J. L.; Shin, J.: p62,

[81496] a phosphotyrosine-independent ligand of the SH2 domain of p56-lck,
[81497] belongs to a new class of ubiquitin-binding proteins. J. Biol. Chem. 271:
[81498] 20235-20237, 1996.

[81499] 14120.

[81500] Akbar, G. K. M.; Dasari, V. R.; Webb, T. E.; Ayyanathan, K.; Pillarisetti,
[81501] K.; Sandhu, A. K.; Athwal, R. S.; Daniel, J. L.; Ashby, B.; Barnard,
[81502] E. A.; Kunapuli, S. P.: Molecular cloning of a novel P2 purinoceptor
[81503] from human erythroleukemia cells. J. Biol. Chem. 271: 18363-18367,
[81504] 1996.

[81505] 14121.

[81506] Chen, X.-S.; Sheller, J. R.; Johnson, E. N.; Funk, C. D.: Role

[81507] of leukotrienes revealed by targeted disruption of 5-lipoxygenase

[81508] gene. Nature 372: 179-182, 1994.

[81509] 14122.

[81510] Devchand, P. R.; Keller, H.; Peters, J. M.; Vazquez, M.; Gonzalez,

[81511] F. J.; Wahli, W.: The PPAR alpha-leukotriene B4 pathway to inflammation

[81512] control. Nature 384: 39-43, 1996.

[81513] 14123.

[81514] Kato, K.; Yokomizo, T.; Izumi, T.; Shimizu, T.: Cell-specific

[81515] transcriptional regulation of human leukotriene B4 receptor gene. J.

[81516] Exp. Med. 192: 413-420, 2000.

[81517] 14124.

[81518] Owman, C.; Nilsson, C.; Lolait, S. J.: Cloning of cDNA encoding

[81519] a putative chemoattractant receptor. Genomics 37:
187-194, 1996.

[81520] 14125.

[81521] Samuelsson, B.; Dahlen, S. E.; Lindgren, J. A.; Rouzer, C.
A.;

[81522] Serhan, C. N.: Leukotrienes and lipoxins: structures,
biosynthesis,

[81523] and biological effects. Science 237: 1171-1176, 1987.

[81524] 14126.

[81525] Yokomizo, T.; Izumi, T.; Chang, K.; Takuwa, Y.; Shimizu,
T.: A

[81526] G-protein-coupled receptor for leukotriene B4 that medi-
ates chemotaxis. Nature 387:

[81527] 620-624, 1997.

[81528] 14127.

[81529] Fernandes-Alnemri, T.; Litwack, G.; Alnemri, E. S.: Mch2, a
new

[81530] member of the apoptotic Ced-3/Ice cysteine protease
gene family. Cancer

[81531] Res. 55: 2737-2742, 1995.

[81532] 14128.

[81533] Orth, K.; Chinnaiyan, A. M.; Garg, M.; Froelich, C. J.; Dixit,

[81534] V. M.: The CED-3/ICE-like protease Mch2 is activated during apoptosis

[81535] and cleaves the death substrate lamin A. J. Biol. Chem. 271: 16443-16446,

[81536] 1996.

[81537] 14129.

[81538] Verhaegh, G. W. C. T.; Jongmans, W.; Jaspers, N. G. J.; Natarajan,

[81539] A. T.; Oshimura, M.; Lohman, P. H. M.; Zdzienicka, M. Z.: A gene

[81540] that regulates DNA replication in response to DNA damage is located

[81541] on human chromosome 4q. Am. J. Hum. Genet. 57: 1095-1103, 1995.

[81542] 14130.

[81543] Kubo, Y.; Reuveny, E.; Slesinger, P. A.; Jan, Y. N.; Jan, L. Y.

[81544] : Primary structure and functional expression of a rat G-protein-coupled

[81545] muscarinic potassium channel. Nature 364: 802-806, 1993.

[81546] 14131.

[81547] Schoots, O.; Voskoglou, T.; Van Tol, H. H. M.: Genomic organization

[81548] and promoter analysis of the human G-protein-coupled K+ channel Kir3.1

[81549] (KCNJ3/HGIRK1). Genomics 39: 279-288, 1997.

[81550] 14132.

[81551] Schoots, O.; Yue, K.-T.; MacDonald, J. F.; Hampson, D. R.;
Nobrega,
[81552] J. N.; Dixon, L. M.; Van Tol, H. H. M.: Cloning of a G pro-
tein-activated
[81553] inwardly rectifying potassium channel from human cere-
bellum. Molec.
[81554] Brain Res. 39: 23-30, 1996.

[81555] 14133.

[81556] Stoffel, M.; Espinosa, R., III; Powell, K. L.; Philipson, L. H.;

[81557] Le Beau, M. M.; Bell, G. I.: Human G-protein-coupled in-
wardly rectifying
[81558] potassium channel (GIRK1) gene (KCNJ3): localization to
chromosome
[81559] 2 and identification of a simple tandem repeat polymor-
phism. Genomics 21:
[81560] 254-256, 1994.

[81561] 14134.

[81562] Cools, J.; Bilhou-Nabera, C.; Wlodarska, I.; Cabrol, C.; Tal-
mant,

[81563] P.; Bernard, P.; Hagemeijer, A.; Marynen, P.: Fusion of a novel gene,
[81564] BTL, to ETV6 in acute myeloid leukemias with a t(4;12)(q11-q12;p13). Blood 94:
[81565] 1820-1824, 1999.

[81566] 14135.

[81567] Phillips, H. A.; Scheffer, I. E.; Berkovic, S. F.; Hollway, G.

[81568] E.; Sutherland, G. R.; Mulley, J. C.: Localization of a gene for
[81569] autosomal dominant nocturnal frontal lobe epilepsy to chromosome 20q13.2. Nature
[81570] Genet. 10: 117-118, 1995.

[81571] 14136.

[81572] Phillips, H. A.; Scheffer, I. E.; Crossland, K. M.; Bhatia, K.

[81573] P.; Fish, D. R.; Marsden, C. D.; Howell, S. J. L.; Stephenson, J.
[81574] B. P.; Tolmie, J.; Plazzi, G.; Eeg-Olofsson, O.; Singh, R.; Lopes-Cendes,
[81575] I.; Andermann, E.; Andermann, F.; Berkovic, S. F.; Mulley, J. C.:
[81576] Autosomal dominant nocturnal frontal-lobe epilepsy: ge-

netic heterogeneity
[81577] and evidence for a second locus at 15q24. Am. J. Hum. Genet. 63:
[81578] 1108-1116, 1998.

[81579] 14137.

[81580] D'Arcangelo, G.: Personal Communication. Nutley, N. J. 6/2/1995.
[81581] 14138.

[81582] D'Arcangelo, G.; Homayouni, R.; Keshvara, L.; Rice, D. S.; Sheldon,
[81583] M.; Curran, T.: Reelin is a ligand for lipoprotein receptors. Neuron 24:
[81584] 471-479, 1999.

[81585] 14139.

[81586] D'Arcangelo, G.; Miao, G. G.; Chen, S.-C.; Soares, H. D.; Morgan,
[81587] J. I.; Curran, T.: A protein related to extracellular matrix proteins
[81588] deleted in the mouse mutant reeler. Nature 374:
719-723, 1995.
[81589] 14140.

[81590] DeSilva, U.; D'Arcangelo, G.; Braden, V. V.; Chen, J.; Miao, G.

[81591] G.; Curran, T.; Green, E. D.: The human reelin gene: isolation, sequencing,

[81592] and mapping on chromosome 7. Genome Res. 7: 157-164, 1997.

[81593] 14141.

[81594] Green, M. C.: Catalog of mutant genes and polymorphic loci.In:

[81595] Lyon, M. F.; Searle, A. G.: Genetic Variants and Strains of the Laboratory

[81596] Mouse. Oxford: Oxford Univ. Press (pub.) (2nd ed.): 1989.

[81597] 14142.

[81598] Hack, I.; Bancila, M.; Loulier, K.; Carroll, P.; Cremer, H.: Reelin

[81599] is a detachment signal in tangential chain-migration during postnatal

[81600] neurogenesis. Nature Neurosci. 5: 939-945, 2002.

[81601] 14143.

[81602] Hirotsune, S.; Takahara, T.; Sasaki, N.; Hirose, K.; Yoshiki, A.;

[81603] Ohashi, T.; Kusakabe, M.; Murakami, Y.; Muramatsu, M.; Watanabe, S.;

[81604] Nakao, K.; Katsuki, M.; Hayashizaki, Y.: The reeler gene encodes

[81605] a protein with an EGF-like motif expressed by pioneer neurons. Nature

[81606] Genet. 10: 77-83, 1995.

[81607] 14144.

[81608] Hong, S. E.; Shugart, Y. Y.; Huang, D. T.; Al Shahwan, S.; Grant,

[81609] P. E.; Hourihane, J. O.; Martin, N. D. T.; Walsh, C. A.: Autosomal

[81610] recessive lissencephaly with cerebellar hypoplasia is associated with

[81611] human RELN mutations. Nature Genet. 26: 93-96, 2000. Note: Erratum:

[81612] Nature Genet. 27: 225 only, 2001.

[81613] 14145.

[81614] Hourihane, J. O.; Bennett, C. P.; Chaudhuri, R.; Robb, S. A.;

[81615] Martin, N. D. T.: A sibship with a neuronal migration defect, cerebellar

[81616] hypoplasia and congenital lymphedema. Neuropediatrics 24: 43-46,

[81617] 1993.

[81618] 14146.

[81619] Impagnatiello, F.; Guidotti, A. R.; Pesold, C.; Dwivedi, Y.; Caruncho,

[81620] H.; Pisu, M. G.; Uzunov, D. P.; Smalheiser, N. R.; Davis, J. M.; Pandey,

[81621] G. N.; Pappas, G. D.; Tueting, P.; Sharma, R. P.; Costa, E.: A decrease

[81622] of reelin expression as a putative vulnerability factor in schizophrenia. Proc.

[81623] Nat. Acad. Sci. 95: 15718-15723, 1998.

[81624] 14147.

[81625] Magdaleno, S.; Keshvara, L.; Curran, T.: Rescue of ataxia and

[81626] preplate splitting by ectopic expression of reelin in reeler mice. Neuron 33:

[81627] 573-586, 2002.

[81628] 14148.

[81629] Royaux, I.; Lambert de Rouvroit, C.; D'Arcangelo, G.;

Demirov,
[81630] D.; Goffinet, A. M.: Genomic organization of the mouse reelin gene. Genomics 46:
[81631] 240-250, 1997.

[81632] 14149.

[81633] Yip, J. W.; Yip, Y. P. L.; Nakajima, K.; Capriotti, C.: Reelin
[81634] controls position of autonomic neurons in the spinal cord. Proc.
[81635] Nat. Acad. Sci. 97: 8612-8616, 2000.

[81636] 14150.

[81637] Bansal-Pakala, P.; Jember, A. G.-H.; Croft, M.: Signaling through
[81638] OX40 (CD134) breaks peripheral T-cell tolerance. Nature Med. 7:
[81639] 907-912, 2001.

[81640] 14151.

[81641] Birkeland, M. L.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.
[81642] A.; Barclay, A. N.: Gene structure and chromosomal localization of
[81643] the mouse homologue of rat OX40 protein. Europ. J. Im-

mun. 25: 926-930,
[81644] 1995.

[81645] 14152.

[81646] Jember, A. G.-H.; Zuberi, R.; Liu, F.-T.; Croft, M.: Development
[81647] of allergic inflammation in a murine model of asthma is dependent
[81648] on the costimulatory receptor OX40. J. Exp. Med. 193: 387-392, 2001.
[81649] 14153.

[81650] Latza, U.; Durkop, H.; Schnittger, S.; Ringeling, J.; Eitel-bach,
[81651] F.; Hummel, M.; Fonatsch, C.; Stein, H.: The human OX40 homolog:
[81652] cDNA structure, expression and chromosomal assignment of the ACT35
[81653] antigen. Europ. J. Immun. 24: 677-683, 1994.

[81654] 14154.

[81655] Anderson, D. W.; Probst, F. J.; Belyantseva, I. A.; Fridell, R.
[81656] A.; Beyer, L.; Martin, D. M.; Wu, D.; Kachar, B.; Friedman, T. B.;

[81657] Raphael, Y.; Camper, S. A.: The motor and tail regions of myosin
[81658] XV are critical for normal structure and function of auditory and
[81659] vestibular hair cells. Hum. Molec. Genet. 9: 1729-1738, 2000.
[81660] 14155.
[81661] Probst, F. J.; Fridell, R. A.; Raphael, Y.; Saunders, T. L.; Wang,
[81662] A.; Liang, Y.; Morell, R. J.; Touchman, J. W.; Lyons, R. H.; Noben-Trauth,
[81663] K.; Friedman, T. B.; Camper, S. A.: Correction of deafness in shaker-2
[81664] mice by an unconventional myosin in a BAC transgene. Science 280:
[81665] 1444-1447, 1998.
[81666] 14156.
[81667] Wang, A.; Liang, Y.; Fridell, R. A.; Probst, F. J.; Wilcox, E.
[81668] R.; Touchman, J. W.; Morton, C. C.; Morell, R. J.; Noben-Trauth, K.;
[81669] Camper, S. A.; Friedman, T. B.: Association of unconventional myosin

[81670] MYO15 mutations with human nonsyndromic deafness DFNB3. Science 280:
[81671] 1447-1451, 1998.

[81672] 14157.

[81673] Wilton, S. D.; Lim, L.; Dorosz, S. D.; Gunn, H. C.; Eyre, H. J.;
[81674] Callen, D. F.; Laing, N. G.: Assignment of the human alpha-tropomyosin
[81675] gene TPM4 to band 19p13.1 by fluorescence in situ hybridization. Cytogenet.
[81676] Cell Genet. 72: 294-296, 1996.

[81677] 14158.

[81678] Thomas, G. H.; Goldberg, M. M.; Miller, C. S.; Reynolds, L. W.
[81679] : Neuraminidase deficiency in the original patient with the Goldberg
[81680] syndrome. Clin. Genet. 16: 323-330, 1979.

[81681] 14159.

[81682] Ronce, N.; Moizard, M.-P.; Robb, L.; Toutain, A.; Villard, L.;
[81683] Moraine, C.: A C2055T transition in exon 8 of the ATP7A

gene is associated
[81684] with exon skipping in an occipital horn syndrome family. (Letter) Am.
[81685] J. Hum. Genet. 61: 233-238, 1997.

[81686] 14160.

[81687] Tsukahara, M.; Imaizumi, K.; Kawai, S.; Kajii, T.: Occipital
[81688] horn syndrome: report of a patient and review of the literature. Clin.
[81689] Genet. 45: 32-35, 1994.

[81690] 14161.

[81691] Tumer, Z.; Lund, C.; Tolshave, J.; Vural, B.; Tonnesen, T.; Horn,
[81692] N.: Identification of point mutations in 41 unrelated patients affected
[81693] with Menkes disease. Am. J. Hum. Genet. 60: 63-71, 1997.
[81694] 14162.

[81695] Tumer, Z.; Vural, B.; Tonnesen, T.; Chelly, J.; Monaco, A. P.;
[81696] Horn, N.: Characterization of the exon structure of the Menkes disease

[81697] gene using vectorette PCR. Genomics 26: 437-442, 1995.

[81698] 14163.

[81699] Vulpe, C.; Levinson, B.; Whitney, S.; Packman, S.; Gitschier,

[81700] J.: Isolation of a candidate gene for Menkes disease and evidence

[81701] that it encodes a copper-transporting ATPase. Nature Genet. 3: 7-13,

[81702] 1993.

[81703] 14164.

[81704] Lazzaro, M. A.; Picketts, D. J.: Cloning and characterization

[81705] of the murine Imitation Switch (ISWI) genes: differential expression

[81706] patterns suggest distinct developmental roles for Snf2h and Snf2l. J.

[81707] Neurochem. 77: 1145-1156, 2001.

[81708] 14165.

[81709] De Plaen, E.; Arden, K.; Traversari, C.; Gaforio, J. J.; Szikora,

[81710] J.-P.; De Smet, C.; Brasseur, F.; van der Bruggen, P.; Lethe, B.;

[81711] Lurquin, C.; Brasseur, R.; Chomez, P.; De Backer, O.; Cavenee, W.;
[81712] Boon, T.: Structure, chromosomal localization, and ex-
pression of
[81713] 12 genes of the MAGE family. Immunogenetics 40:
360-369, 1994.
[81714] 14166.

[81715] Rogner, U. C.; Wilke, K.; Steck, E.; Korn, B.; Poustka, A.: The
[81716] melanoma antigen gene (MAGE) family is clustered in the chromosomal
[81717] band Xq28. Genomics 29: 725-731, 1995.

[81718] 14167.

[81719] Chakarova, C.; Wehnert, M. S.; Uhl, K.; Sakthivel, S.; Vos-
berg,
[81720] H.-P.; van der Ven, P. F. M.; Furst, D. O.: Genomic struc-
ture and
[81721] fine mapping of the two human filamin gene paralogues FLNB and FLNC
[81722] and comparative analysis of the filamin gene family. Hum. Genet. 107:
[81723] 597-611, 2000.

[81724] 14168.

[81725] Fink, J. M.; Dobyns, W. B.; Guerrini, R.; Hirsch, B. A.: Identification

[81726] of a duplication of Xq28 associated with bilateral periventricular

[81727] nodular heterotopia. Am. J. Hum. Genet. 61: 379-387, 1997.

[81728] 14169.

[81729] Fox, J. W.; Lamperti, E. D.; Eksioglu, Y. Z.; Hong, S. E.; Feng,

[81730] Y.; Graham, D. A.; Scheffer, I. E.; Dobyns, W. B.; Hirsch, B. A.;

[81731] Radtke, R. A.; Berkovic, S. F.; Huttenlocher, P. R.; Walsh, C. A.

[81732] : Mutations in filamin 1 prevent migration of cerebral cortical neurons

[81733] in human periventricular heterotopia. Neuron 21: 1315-1325, 1998.

[81734] 14170.

[81735] Gariboldi, M.; Maestrini, E.; Canzian, F.; Manenti, G.; De Gregorio,

[81736] L.; Rivella, S.; Chatterjee, A.; Herman, G. E.; Archidiacono,

N.;

[81737] Antonacci, R.; Pierotti, M. A.; Dragani, T. A.; Toniolo, D.: Comparative

[81738] mapping of the actin-binding protein 280 genes in human and mouse. Genomics 21:

[81739] 428-430, 1994.

[81740] 14171.

[81741] Bashir, R.; Britton, S.; Strachan, T.; Keers, S.; Vafiadaki, E.;

[81742] Lako, M.; Richard, I.; Marchand, S.; Bourg, N.; Argov, Z.; Sadeh,

[81743] M.; Mahjneh, I.; Marconi, G.; Passos-Bueno, M. R.; Moreira, E. S.;

[81744] Zatz, M.; Beckmann, J. S.; Bushby, K.: A gene related to Caenorhabditis

[81745] elegans spermatogenesis factor fer-1 is mutated in limb-girdle muscular

[81746] dystrophy type 2B. Nature Genet. 20: 37-42, 1998.

[81747] 14172.

[81748] Liu, J.; Aoki, M.; Illa, I.; Wu, C.; Fardeau, M.; Angelini, C.;

[81749] Serrano, C.; Urtizberea, J. A.; Hentati, F.; Ben Hamida, M.; Bohlega,

[81750] S.; Culper, E. J.; Amato, A. A.; Bossie, K.; Oeltjen, J.; Bejaoui,
[81751] K.; McKenna-Yasek, D.; Hosler, B. A.; Schurr, E.; Arahata, K.; de
[81752] Jong, P. J.; Brown, R. H., Jr.: Dysferlin, a novel skeletal muscle
[81753] gene, is mutated in Miyoshi myopathy and limb girdle muscular dystrophy. Nature
[81754] Genet. 20: 31-36, 1998.

[81755] 14173.

[81756] Thomas, G. H.; Tipton, R. E.; Ch'ien, L. T.; Reynolds, L. W.;
[81757] Miller, C. S.: Sialidase (alpha-N-acetyl neuraminidase) deficiency:
[81758] the enzyme defect in an adult with macular cherry red spots and myoclonus
[81759] without dementia. Clin. Genet. 13: 369-379, 1978.

[81760] 14174.

[81761] Thomas, P. K.; Abrams, J. D.; Swallow, D.; Stewart, G.: Sialidosis
[81762] type I: cherry-red spot myoclonus syndrome with sialidase deficiency

[81763] and altered electrophoretic mobilities of some enzymes known to be

[81764] glycoproteins. J. Neurol. Neurosurg. Psychiat. 42: 873-880, 1979.

[81765] 14175.

[81766] Tipton, R. E.; Thomas, G. H.; Ch'ien, L. T.; Reynolds, L. W.;

[81767] Miller, C. S.: Sialidase (alpha-N-acetyl neuraminidase) de-ficiency,

[81768] macular cherry-red spots and myoclonus. (Abstract) Clin. Res. 26:

[81769] 76A only, 1978.

[81770] 14176.

[81771] Tondeur, M.; Libert, J.; Vamos, E.; Van Hoof, F.; Thomas, G. H.;

[81772] Strecker, G.: Infantile sialic acid storage disorder: clinical, ultrastructural

[81773] and biochemical studies in two siblings. Europ. J. Pediat. 139:

[81774] 142-147, 1982.

[81775] 14177.

[81776] Verheijen, F. W.; Palmieri, S.; Galjaard, H.: Purification and

[81777] partial characterization of lysosomal neuraminidase from human placenta. Europ.

[81778] J. Biochem. 162: 63-67, 1987.

[81779] 14178.

[81780] Verheijen, F. W.; Palmieri, S.; Hoogeveen, A. T.; Galjaard, H.

[81781] : Human placental neuraminidase: activation, stabilization and association

[81782] with beta-galactosidase and its 'protective' protein. Europ. J. Biochem. 149:

[81783] 315-321, 1985.

[81784] 14179.

[81785] Winter, R. M.; Swallow, D. M.; Baraitser, M.; Purkiss, P.: Sialidosis

[81786] type 2 (acid neuraminidase deficiency): clinical and bio-chemical features

[81787] of a further case. Clin. Genet. 18: 203-210, 1980.

[81788] 14180.

[81789] Womack, J. E.; David, C. S.: Mouse gene for neuraminidase activity

[81790] (Neu-1) maps to the D end of H-2. Immunogenetics 16:

177-180, 1982.

[81791] 14181.

[81792] Womack, J. E.; Eicher, E.: Liver-specific lysosomal acid phosphatase

[81793] deficiency (Apl) on mouse chromosome 17. Molec. Genet. 155: 315-317,

[81794] 1977.

[81795] 14182.

[81796] Womack, J. E.; Yan, D. L. S.; Potier, M.: Gene for neu-raminidase

[81797] activity on mouse chromosome 17 near H-2: pleiotropic effects on multiple

[81798] hydrolases. Science 212: 63-65, 1981.

[81799] 14183.

[81800] Womack, J. E.; Yan, D. L. S.; Potier, M.: Liver neu-raminidase

[81801] deficiency inherited as a single gene on mouse chromo-some 17. (Abstract) Am.

[81802] J. Hum. Genet. 32: 59A only, 1980.

[81803] 14184.

[81804] Yamamoto, A.; Adachi, S.; Kawamura, S.; Takahashi, M.;

Kitano,
[81805] T.; Ohtori, T.; Shinji, Y.; Nishikawa, M.: Localized beta-galactosidase
[81806] deficiency. Arch. Intern. Med. 134: 627-634, 1974.

[81807] 14185.

[81808] Young, I. D.; Young, E. P.; Mossman, J.; Fielder, A. R.; Moore,
[81809] J. R.: Neuraminidase deficiency: case report and review of the phenotype. J.
[81810] Med. Genet. 24: 283-290, 1987.

[81811] 14186.

[81812] Rosenbloom, A. L.; Guevara-Aguirre, J.; Rosenfeld, R. G.; Fiedler,
[81813] P. J.: The little women of Loja-- growth hormone-receptor deficiency
[81814] in an inbred population of southern Ecuador. New Eng. J. Med. 323:
[81815] 1367-1374, 1990.

[81816] 14187.

[81817] Woods, K. A.; Fraser, N. C.; Postel-Vinay, M.-C.; Savage, M. O.;

[81818] Clark, A. J. L.: A homozygous splice site mutation affecting the

[81819] intracellular domain of the growth hormone (GH) receptor resulting

[81820] in Laron syndrome with elevated GH-binding protein. J. Clin. Endocr.

[81821] Metab. 81: 1686-1690, 1996.

[81822] 14188.

[81823] Aghai, E.; Yaniv, I.; David, M.: Factor XI deficiency in an Arab

[81824] Moslem family in Israel. Scand. J. Haemat. 32: 327-331, 1984.

[81825] 14189.

[81826] Asakai, R.; Chung, D. W.; Davie, E. W.; Seligsohn, U.: Factor

[81827] XI deficiency in Ashkenazi Jews in Israel. New Eng. J. Med. 325:

[81828] 153-158, 1991.

[81829] 14190.

[81830] Asakai, R.; Chung, D. W.; Ratnoff, O. D.; Davie, E. W.: Factor

[81831] XI (plasma thromboplastin antecedent) deficiency in Ashkenazi Jews
[81832] is a bleeding disorder that can result from three types of point mutations. Proc.
[81833] Nat. Acad. Sci. 86: 7667-7671, 1989.

[81834] 14191.

[81835] Asakai, R.; Davie, E. W.; Chung, D. W.: Organization of the gene
[81836] for human factor XI. Biochemistry 26: 7221-7228, 1987.

[81837] 14192.

[81838] Bauduer, F.; Dupreuilh, F.; Ducout, L.; Marti, B.: Factor XI deficiency
[81839] in the French Basque country. Haemophilia 5: 187-190, 1999.
[81840] 14193.

[81841] Bertina, R. M.; Veltkamp, J. J.: A genetic variant of factor XI
[81842] with decreased capacity for Ca ion binding. Brit. J. Haemat. 42:
[81843] 623-635, 1979.

[81844] 14194.

[81845] Biggs, R.; MacFarlane, R. G.: Human Blood Coagulation and Its
[81846] Disorders. Oxford: Blackwell (pub.) (3rd ed.): 1962.
[81847] 14195.
[81848] Bolton-Maggs, P. H. B.: Factor XI deficiency. Bailliere's Clin.
[81849] Haemat. 9: 355-368, 1996.
[81850] 14196.
[81851] Braunstein, K. M.; Noyes, C. M.; Griffith, M. J.; Lundblad, R.
[81852] L.; Roberts, H. R.: Characterization of the defect in activation
[81853] of factor IX Chapel Hill by human factor Xla. J. Clin. Invest. 68:
[81854] 1420-1426, 1981.
[81855] 14197.
[81856] Buetow, K. H.; Shiang, R.; Yang, P.; Nakamura, Y.; Lathrop, G.
[81857] M.; White, R.; Wasmuth, J. J.; Wood, S.; Berdahl, L. D.; Leysens,
[81858] N. J.: A detailed multipoint map of human chromosome 4 provides evidence

[81859] for linkage heterogeneity and position-specific recombination rates. Am.

[81860] J. Hum. Genet. 48: 911-925, 1991.

[81861] 14198.

[81862] Clarkson, K.; Rosenfeld, B.; Fair, J.; Klein, A.; Bell, W.: Factor

[81863] XI deficiency acquired by liver transplantation. Ann. Intern. Med. 115:

[81864] 877-879, 1991.

[81865] 14199.

[81866] Dodds, W. J.; Kull, J. E.: Canine factor XI (plasma thromboplastin

[81867] antecedent) deficiency. J. Lab. Clin. Med. 78: 746-752, 1971.

[81868] 14200.

[81869] Dzik, W. H.; Arkin, C. F.; Jenkins, R. L.: Transfer of congenital

[81870] factor XI deficiency from a donor to a recipient by liver transplantation.

[81871] (Letter) New Eng. J. Med. 316: 1217-1218, 1987.

[81872] 14201.

[81873] Fujikawa, K.; Chung, D. W.; Hendrickson, L. E.; Davie, E. W.:

[81874] Amino acid sequence of human factor XI, a blood coagulation factor

[81875] with four tandem repeats that are highly homologous with plasma prekallikrein. Biochemistry 25:

[81876] 2417-2424, 1986.

[81877] 14202.

[81878] Imanaka, Y.; Lal, K.; Nishimura, T.; Bolton-Maggs, P. H. B.; Tuddenham,

[81879] E. G. D.; McVey, J. H.: Identification of two novel mutations in

[81880] non-Jewish factor XI deficiency. Brit. J. Haemat. 90: 916-920, 1995.

[81881] 14203.

[81882] Kato, A.; Asakai, R.; Davie, E. W.; Aoki, N.: Factor XI gene

[81883] (F11) is located on the distal end of the long arm of human chromosome

[81884] 4. Cytogenet. Cell Genet. 52: 77-78, 1989.

[81885] 14204.

[81886] Litz, C. E.; Swaim, W. R.; Dalmasso, A. P.: Factor XI defi-

ciency:
[81887] genetic and clinical studies of a single kindred. Am. J. Hemat. 28:
[81888] 8-12, 1988.

[81889] 14205.

[81890] Mannhalter, C.; Hellstern, P.; Deutsch, E.: Identification of

[81891] a defective factor XI cross-reacting material in a factor XI-deficient
[81892] patient. Blood 70: 31-37, 1987.

[81893] 14206.

[81894] Meijers, J. C. M.; Davie, E. W.; Chung, D. W.: Expression of

[81895] human blood coagulation factor XI: characterization of the defect
[81896] in factor XI type III deficiency. Blood 79: 1435-1440, 1992.
[81897] 14207.

[81898] Mitchell, M.; Cutler, J.; Thompson, S.; Moore, G.; Rees, E. J.;
[81899] Smith, M.; Savidge, G.; Alhaq, A.: Heterozygous factor XI deficiency
[81900] associated with three novel mutations. Brit. J. Haemat.

107: 763-765,
[81901] 1999.

[81902] 14208.

[81903] Muir, W. A.; Ratnoff, O. D.: The prevalence of plasma thromboplastin
[81904] antecedent (PTA factor XI) deficiency. Blood 44: 569-570, 1974.
[81905] 14209.

[81906] Niskanen, E. O.; Saito, H.; Cline, M. J.: Plasma thrombo-plastin
[81907] antecedent (factor XI) deficiency in a black family. Arch. Intern.
[81908] Med. 141: 936-941, 1981.

[81909] 14210.

[81910] Peretz, U.; Zivelin, A.; Usher, S.; Eichel, R.; Seligsohn, U.

[81911] : Identification of a new mutation in the factor XI gene of an Ashkenazi-Jew
[81912] with severe factor XI deficiency. (Abstract) Blood 82 (suppl. 1):
[81913] 66a only, 1993.

[81914] 14211.

[81915] Pugh, R. E.; McVey, J. H.; Tuddenham, E. G. D.; Hancock, J. F.

[81916] : Six point mutations that cause factor XI deficiency. Blood 85:

[81917] 1509-1516, 1995.

[81918] 14212.

[81919] Ragni, M. V.; Sinha, D.; Seaman, F.; Lewis, J. H.; Spero, J. A.;

[81920] Walsh, P. N.: Comparison of bleeding tendency, factor XI coagulant

[81921] activity, and factor XI antigen in 25 factor XI-deficient kindreds. Blood 65:

[81922] 719-724, 1985.

[81923] 14213.

[81924] Rapaport, S. I.; Proctor, R. R.; Patch, M. J.; Yettra, M.: The

[81925] mode of inheritance of PTA deficiency: evidence for the existence

[81926] of major PTA deficiency and minor PTA deficiency. Blood 18: 149-165,

[81927] 1961.

[81928] 14214.

[81929] Rimon, A.; Schiffman, S.; Feinstein, D. I.; Rapaport, S. I.:

[81930] Factor-XI activity and factor-XI antigen in homozygous and heterozygous

[81931] factor-XI deficiency. Blood 48: 165-174, 1976.

[81932] 14215.

[81933] Foresti, V.; Ferrari, C.: Central hypothyroidism: isolated thyrotropin-releasing

[81934] hormone deficiency or resistance of pituitary thyrotropes?. (Letter) J.

[81935] Endocr. Invest. 8: 577 only, 1985.

[81936] 14216.

[81937] Katakami, H.; Kato, Y.; Inada, M.; Imura, H.: Hypothalamic hypothyroidism

[81938] due to isolated thyrotropin-releasing hormone (TRH) deficiency. J.

[81939] Endocr. Invest. 7: 231-233, 1984.

[81940] 14217.

[81941] Niimi, H.; Inomata, H.; Sasaki, N.; Nakajima, H.: Congenital isolated

[81942] thyrotrophin releasing hormone deficiency. Arch. Dis. Child. 57:

[81943] 877-878, 1982.

[81944] 14218.

[81945] Roller, M. L.; Camper, S. A.: Localization of the thy-
rotropin-releasing
[81946] hormone gene, Trh, on mouse chromosome 6. Mam-
malian Genome 6: 443-444,
[81947] 1995.

[81948] 14219.

[81949] Yamada, M.; Radovick, S.; Wondisford, F. E.; Nakayama, Y.;
Weintraub,
[81950] B. D.; Wilber, J. F.: Cloning and structure of human ge-
nomic DNA
[81951] and hypothalamic cDNA encoding human preprothy-
rotropin-releasing hormone. Molec.
[81952] Endocr. 4: 551-556, 1990.

[81953] 14220.

[81954] Yamada, M.; Satoh, T.; Monden, T.; Mori, M.: Assignment
of the
[81955] thyrotropin-releasing hormone gene (TRH) to human
chromosome 3q13.3-q21
[81956] by in situ hybridization. Cytogenet. Cell Genet. 87: 275

only, 1999.

[81957] 14221.

[81958] Yamada, M.; Wondisford, F. E.; Radovick, S.; Nakayama, Y.; Weintraub,

[81959] B. D.; Wilber, J. F.: Assignment of human preprothy-

rotropin-releasing

[81960] hormone (TRH) gene to chromosome 3. Somat. Cell Molec. Genet. 17:

[81961] 97-100, 1991.

[81962] 14222.

[81963] Abramowicz, M. J.; Duprez, L.; Parma, J.; Vassart, G.; Heinrichs,

[81964] C.: Familial congenital hypothyroidism due to inactivating mutation

[81965] of the thyrotropin receptor causing profound hypoplasia of the thyroid

[81966] gland. J. Clin. Invest. 99: 3018-3024, 1997.

[81967] 14223.

[81968] Biebermann, H.; Schoneberg, T.; Krude, H.; Schultz, G.; Gudermann,

[81969] T.; Gruters, A.: Mutations of the human thyrotropin recep-

tor gene

[81970] causing thyroid hypoplasia and persistent congenital hypothyroidism. J.

[81971] Clin. Endocr. Metab. 82: 3471-3480, 1997.

[81972] 14224.

[81973] Clifton-Bligh, R. J.; Gregory, J. W.; Ludgate, M.; John, R.; Persani,

[81974] L.; Asteria, C.; Beck-Peccoz, P.; Chatterjee, V. K. K.: Two novel

[81975] mutations in the thyrotropin (TSH) receptor gene in a child with resistance

[81976] to TSH. J. Clin. Endocr. Metab. 82: 1094-1100, 1997.

[81977] 14225.

[81978] de Roux, N.; Misrahi, M.; Brauner, R.; Houang, M.; Carel, J. C.;

[81979] Granier, M.; le Bouc, Y.; Ghinea, N.; Boumedienne, A.; Toublanc, J.

[81980] E.; Milgrom, E.: Four families with loss of function mutations of

[81981] the thyrotropin receptor. J. Clin. Endocr. Metab. 81: 4229-4235,

[81982] 1996.

[81983] 14226.

[81984] Sunthornthepvarakul, T.; Gottschalk, M. E.; Hayashi, Y.; Refetoff,

[81985] S.: Resistance to thyrotropin caused by mutations in the thyrotropin-receptor

[81986] gene. New Eng. J. Med. 332: 155-160, 1995.

[81987] 14227.

[81988] Morrison, N. A.; Qi, J. C.; Tokita, A.; Kelly, P. J.; Crofts,

[81989] L.; Nguyen, T. V.; Sambrook, P. N.; Elsman, J. A.: Predic-tion of

[81990] bone density from vitamin D receptor alleles. Nature 367: 284-287,

[81991] 1994.

[81992] 14228.

[81993] Szpirer, J.; Szpirer, C.; Riviere, M.; Levan, G.; Marynen, P.;

[81994] Cassiman, J.-J.; Wiese, R.; DeLuca, H. F.: The Sp1 tran-scription

[81995] factor gene (SP1) and the 1,25-dihydroxyvitamin D(3) re-ceptor gene

[81996] (VDR) are colocalized on human chromosome arm 12q and rat chromosome

[81997] 7. Genomics 11: 168-173, 1991.

[81998] 14229.

[81999] Van Maldergem, L.; Bachy, A.; Feldman, D.; Bouillon, R.; Maassen,

[82000] J.; Dreyer, M.; Rey, R.; Holm, C.; Gillerot, Y.: Syndrome of lipoatrophic

[82001] diabetes, vitamin D resistant rickets, and persistent Mulle-rian ducts

[82002] in a Turkish boy born to consanguineous parents. Am. J. Med. Genet. 64:

[82003] 506-513, 1996.

[82004] 14230.

[82005] McDermott, J. C.; Cardoso, M. C.; Yu, Y.-T.; Andres, V.; Leifer,

[82006] D.; Krainc, D.; Lipton, S. A.; Nadal-Ginard, B.: hMEF2C gene encodes

[82007] skeletal muscle- and brain-specific transcription factors. Molec.

[82008] Cell. Biol. 13: 2564-2577, 1993.

[82009] 14231.

[82010] Anderson, G. J.; Murphy, T. L.; Cowley, L.; Evans, B. A.;

Halliday,
[82011] J. W.; McLaren, G. D.: Mapping the gene for sex-linked anemia: an
[82012] inherited defect of intestinal iron absorption in the mouse. Genomics 48:
[82013] 34-39, 1998.

[82014] 14232.

[82015] Bannerman, R. M.: Genetic defects of iron transport. Fed. Proc. 35:
[82016] 2281-2285, 1976.

[82017] 14233.

[82018] Edwards, J. A.; Hoke, J. E.; Mattioli, M.; Reichlin, M.: Fer-
ritin
[82019] distribution and synthesis in sex-linked anemia. J. Lab. Clin. Med. 90:
[82020] 68-76, 1977.

[82021] 14234.

[82022] Falconer, D. S.; Isaacson, J. H.: The genetics of sex-linked anaemia
[82023] in the mouse. Genet. Res. 3: 248-250, 1962.

[82024] 14235.

[82025] Grewal, M. S.: A sex-linked anaemia in the mouse. Genet. Res. 3:

[82026] 238-247, 1962.

[82027] 14236.

[82028] Lee, G. R.; Nacht, S.; Lukens, J. N.; Cartwright, G. E.: Iron

[82029] metabolism in copper-deficient swine. J. Clin. Invest. 47: 2058-2069,

[82030] 1968.

[82031] 14237.

[82032] Vulpe, C. D.; Kuo, Y.-M.; Murphy, T. L.; Cowley, L.; Askwith, C.;

[82033] Libina, N.; Gitschier, J.; Anderson, G. J.: Hephaestin, a ceruloplasmin

[82034] homologue implicated in intestinal iron transport, is de-fective in

[82035] the sla mouse. Nature Genet. 21: 195-199, 1999.

[82036] 14238.

[82037] Karumanchi, S. A.; Jha, V.; Ramchandran, R.; Karihaloo, A.; Tsiokas,

[82038] L.; Chan, B.; Dhanabal, M.; Hanai, J.; Venkataraman, G.; Shriver,

[82039] Z.; Keiser, N.; Kalluri, R.; and 9 others: Cell surface glypi-
cans
[82040] are low-affinity endostatin receptors. Molec. Cell 7:
811-822, 2001.
[82041] 14239.

[82042] Veugelers, M.; Vermeesch, J.; Watanabe, K.; Yamaguchi,
Y.; Marynen,
[82043] P.; David, G.: GPC4, the gene for human K-glypican, flanks
GPC3 on
[82044] Xq26: deletion of the GPC3-GPC4 gene cluster in one
family with Simpson-Golabi-Behmel
[82045] syndrome. Genomics 53: 1-11, 1998.

[82046] 14240.

[82047] Alitalo, T.; Francis, F.; Kere, J.; Lehrach, H.; Schlessinger,

[82048] D.; Willard, H. F.: A 6-Mb YAC contig in Xp22.1-p22.2
spanning the
[82049] DXS69E, XE59, GLRA2, PIGA, GRPR, CALB3, and PHKA2
genes. Genomics 25:
[82050] 691-700, 1995.

[82051] 14241.

[82052] de Conciliis, L.; Marchitiello, A.; Wapenaar, M. C.; Borsani,

G.;

[82053] Giglio, S.; Mariani, M.; Consalez, G. G.; Zuffardi, O.; Franco, B.;

[82054] Ballabio, A.; Banfi, S.: Characterization of Cxorf5 (71-7A), a novel

[82055] human cDNA mapping to Xp22 and encoding a protein containing coiled-coil

[82056] alpha-helical domains. Genomics 51: 243-250, 1998.

[82057] 14242.

[82058] de Martinville, B.; Kunkel, L. M.; Bruns, G.; Morle, F.; Koenig,

[82059] M.; Mandel, J. L.; Horwich, A.; Latt, S. A.; Gusella, J. F.; Housman,

[82060] D.; Francke, U.: Localization of DNA sequences in region Xp21 of

[82061] the human X chromosome: search for molecular markers close to the

[82062] Duchenne muscular dystrophy locus. Am. J. Hum. Genet. 37: 235-249,

[82063] 1985.

[82064] 14243.

[82065] Emes, R. D.; Ponting, C. P.: A new sequence motif linking

lissencephaly,

[82066] Treacher Collins and oral-facial-digital type 1 syndromes, microtubule

[82067] dynamics and cell migration. Hum. Molec. Genet. 10: 2813-2820, 2001.

[82068] 14244.

[82069] Ferrante, M. I.; Giorgio, G.; Feather, S. A.; Bulfone, A.; Wright,

[82070] V.; Ghiani, M.; Selicorni, A.; Gammaro, L.; Scolari, F.; Woolf, A.

[82071] S.; Sylvie, O.; Le Marec, B.; Malcolm, S.; Winter, R.; Ballabio, A.;

[82072] Franco, B.: Identification of the gene for oral-facial-digital type

[82073] I syndrome. Am. J. Hum. Genet. 68: 569-576, 2001.

[82074] 14245.

[82075] Kunkel, L. M.; Tantravahi, U.; Kurnit, D. M.; Eisenhard, M.; Bruns,

[82076] G. P.; Latt, S. A.: Identification and isolation of transcribed human

[82077] X chromosome DNA sequences. Nucleic Acids Res. 11: 7961-7979, 1983.

[82078] 14246.

[82079] Odent, S.; Le Marec, B.; Toutain, A.; David, A.; Vigneron, J.;

[82080] Treguier, C.; Jouan, H.; Milon, J.; Fryns, J.-P.; Verloes, A.: Central

[82081] nervous system malformations and early end-stage renal disease in

[82082] oro-facial-digital syndrome type I: a review. Am. J. Med. Genet. 75:

[82083] 389-394, 1998.

[82084] 14247.

[82085] Scolari, F.; Valzorio, B.; Carli, O.; Vizzardi, V.; Costantino,

[82086] E.; Grazioli, L.; Bondioni, M. P.; Savoldi, S.; Maiorca, R.: Oral-facial-digital

[82087] syndrome type I: an unusual cause of hereditary cystic kidney disease. Nephrol.

[82088] Dial Transplant. 12: 1247-1250, 1997.

[82089] 14248.

[82090] Copley, L. M.; Zhao, W. D.; Kopacz, K.; Herman, G. E.; Kioschis,

[82091] P.; Poustka, A.; Taudien, S.; Platzer, M.: Exclusion of mutations

[82092] in the MTMR1 gene as a frequent cause of X-linked myotubular myopathy.

[82093] (Letter) Am. J. Med. Genet. 107: 256-258, 2002.

[82094] 14249.

[82095] Kioschis, P.; Wiemann, S.; Heiss, N. S.; Francis, F.; Gotz, C.;

[82096] Poustka, A.; Taudien, S.; Platzer, M.; Wiehe, T.; Beckmann, G.; Weber,

[82097] J.; Nordsiek, G.; Rosenthal, A.: Genomic organization of a 225-kb

[82098] region in Xq28 containing the gene for X-linked myotubular myopathy

[82099] (MTM1) and a related gene (MTMR1). Genomics 54: 256-266, 1998.

[82100] 14250.

[82101] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Tentler, D.; Kretz, C.;

[82102] Dahl, N.; Mandel, J.-L.: Characterization of the myotubularin dual

[82103] specificity phosphatase gene family from yeast to human. Hum. Molec.

[82104] Genet. 7: 1703-1712, 1998.

[82105] 14251.

[82106] Divry, P.; Vianey-Liaud, C.; Gay, C.; Macabeo, V.; Rapin, F.; Echenne,

[82107] B.: N-acetylaspartic aciduria: report of three new cases in children

[82108] with a neurological syndrome associating macrocephaly and leucodystrophy. J.

[82109] Inherit. Metab. Dis. 11: 307-308, 1988.

[82110] 14252.

[82111] Gorlin, J. B.; Henske, E.; Warren, S. T.; Kunst, C. B.; D'Urso,

[82112] M.; Palmieri, G.; Hartwig, J. H.; Bruns, G.; Kwiatkowski, D. J.:

[82113] Actin-binding protein (ABP-280) filamin gene (FLN) maps telomeric

[82114] to the color vision locus (R/GCP) and centromeric to G6PD in Xq28. Genomics 17:

[82115] 496-498, 1993.

[82116] 14253.

[82117] Rouger, H.; LeGuern, E.; Birouk, N.; Gouider, R.; Tardieu, S.;

[82118] Plassart, E.; Gugenheim, M.; Vallat, J.-M.; Louboutin, J.-P.;

Bouche,
[82119] P.; Agid, Y.; Brice, A.: Charcot-Marie-Tooth disease with intermediate
[82120] motor nerve conduction velocities: characterization of 14 CX32 mutations
[82121] in 35 families. Hum. Mutat. 10: 443-452, 1997.

[82122] 14254.

[82123] Jager, R. J.; Anvret, M.; Hall, K.; Scherer, G.: A human XY female
[82124] with a frame shift mutation in the candidate testis-determining gene
[82125] SRY. Nature 348: 452-454, 1990.

[82126] 14255.

[82127] Koopman, P.; Gubbay, J.; Vivian, N.; Goodfellow, P.; Lovell-Badge,
[82128] R.: Male development of chromosomally female mice transgenic for
[82129] Sry. Nature 351: 117-121, 1991.

[82130] 14256.

[82131] Koopman, P.; Munsterberg, A.; Capel, B.; Vivian, N.; Lovell-Badge,

[82132] R.: Expression of a candidate sex-determining gene dur-
ing mouse testis
[82133] differentiation. Nature 348: 450-452, 1990.

[82134] 14257.

[82135] McElreavey, K. D.; Vilain, E.; Boucekkine, C.; Vidaud, M.;
Jaubert,
[82136] F.; Richaud, F.; Fellous, M.: XY sex reversal associated with
a nonsense
[82137] mutation in SRY. Genomics 13: 838-840, 1992.

[82138] 14258.

[82139] Page, D. C.; Mosher, R.; Simpson, E. M.; Fisher, E. M. C.;
Mardon,
[82140] G.; Pollack, J.; McGillivray, B.; de la Chapelle, A.; Brown, L.
G.
[82141] : The sex-determining region of the human Y chromo-
some encodes a finger
[82142] protein. Cell 51: 1091-1104, 1987.

[82143] 14259.

[82144] Taylor, H.; Barter, R. H.; Jacobson, C. B.: Neoplasms of
dysgenetic
[82145] gonads. Am. J. Obstet. Gynec. 96: 816-823, 1966.

[82146] 14260.

[82147] Vilain, E.; McElreavey, K.; Jaubert, F.; Raymond, J.-P.; Richaud,

[82148] F.; Fellous, M.: Familial case with sequence variant in the testis-determining

[82149] region associated with two sex phenotypes. Am. J. Hum. Genet. 50:

[82150] 1008-1011, 1992.

[82151] 14261.

[82152] Elpeleg, O. N.; Anikster, Y.; Barash, V.; Branski, D.; Shaag, A.

[82153] : The frequency of the C854 mutation in the aspartoacy-lase gene in

[82154] Ashkenazi Jews in Israel. Am. J. Hum. Genet. 55: 287-288, 1994.

[82155] 14262.

[82156] Feigelman, T.; Shih, V. E.; Buyse, M. L.: Prolonged survival in

[82157] Canavan disease. Dysmorph. Clin. Genet. 5: 107-110, 1991.

[82158] 14263.

[82159] Hagenfeldt, L.; Bollgren, I.; Venizelos, N.: N-acetylaspartic
[82160] aciduria due to aspartoacylase deficiency--a new etiology of childhood
[82161] leukodystrophy. J. Inherit. Metab. Dis. 10: 135-141, 1967.
[82162] 14264.
[82163] Hogan, G. R.; Richardson, E. P., Jr.: Spongy degeneration of
[82164] the nervous system (Canavan's disease): report of a case in an Irish-American
[82165] family. Pediatrics 35: 284-294, 1965.
[82166] 14265.
[82167] Kaul, R.; Balamurugan, K.; Gao, G. P.; Matalon, R.: Canavan disease:
[82168] genomic organization and localization of human ASPA to 17p13-ter and
[82169] conservation of the ASPA gene during evolution. Genomics 21: 364-370,
[82170] 1994.
[82171] 14266.
[82172] Kaul, R.; Gao, G. P.; Aloya, M.; Balamurugan, K.; Petrosky, A.;

[82173] Michals, K.; Matalon, R.: Canavan disease: mutations among Jewish
[82174] and non-Jewish patients. Am. J. Hum. Genet. 55: 34-41, 1994.
[82175] 14267.

[82176] Kaul, R.; Gao, G. P.; Balamurugan, K.; Matalon, R.: Cloning of
[82177] the human aspartoacylase cDNA and a common missense mutation in Canavan
[82178] disease. Nature Genet. 5: 118-123, 1993.

[82179] 14268.

[82180] Kaul, R.; Gao, G. P.; Matalon, R.; Aloya, M.; Su, Q.; Jin, M.;

[82181] Johnson, A. B.; Schutgens, R. B. H.; Clarke, J. T. R.: Identi-
fication
[82182] and expression of eight novel mutations among non-Jewish patients
[82183] with Canavan disease. Am. J. Hum. Genet. 59: 95-102, 1996.
[82184] 14269.

[82185] Kaul, R.; Gao, G. P.; Michals, K.; Whelan, D. T.; Levin, S.; Matalon,

[82186] R.: Novel (cys152-to-arg) missense mutation in an Arab patient with
[82187] Canavan disease. Hum. Mutat. 5: 269-271, 1995.

[82188] 14270.

[82189] Kvittingen, E. A.; Guldal, G.; Borsting, S.; Skalpe, I. O.; Stokke,
[82190] O.; Jellum, E.: N-acetylaspartic aciduria in a child with a progressive
[82191] cerebral atrophy. Clin. Chim. Acta 158: 217-227, 1986.

[82192] 14271.

[82193] Mahloudji, M.; Daneshbod, K.; Karjoo, M.: Familial spongy degeneration
[82194] of the brain. Arch. Neurol. 22: 294-298, 1970.

[82195] 14272.

[82196] Matalon, R.: Personal Communication. Miami, Fla. 11/3/1990.
[82197] 14273.

[82198] Matalon, R.; Kaul, R.; Casanova, J.; Michals, K.; Johnson, A.;
[82199] Rapin, I.; Gashkoff, P.; Deanching, M.: Aspartoacylase de-ficiency:

[82200] the enzyme defect in Canavan disease. J. Inherit. Metab. Dis. 12
[82201] (suppl. 2): 329-331, 1989.

[82202] 14274.

[82203] Matalon, R.; Michals, K.; Sebesta, D.; Deanching, M.; Gashkoff,
[82204] P.; Casanova, J.: Aspartoacylase deficiency and N-acetylaspartic
[82205] aciduria in patients with Canavan disease. Am. J. Med. Genet. 29:
[82206] 463-471, 1988.

[82207] 14275.

[82208] Morcaldi, L.; Salvati, G.; Giordano, G. G.; Guazzi, G. C.: Congenital
[82209] van Bogaert-Bertrand disease in a non-Jewish family. Acta Genet.
[82210] Med. Gemellol. 18: 142-157, 1969.

[82211] 14276.

[82212] Ozand, P. T.; Gascon, G. G.; Dhalla, M.: Aspartoacylase deficiency
[82213] and Canavan disease in Saudi Arabia. Am. J. Hum. Genet.

35: 266-268,
[82214] 1990.

[82215] 14277.

[82216] Propheta, O.; Magal, N.; Shohat, M.; Eyal, N.; Navot, N.; Horowitz,
[82217] M.: A benign polymorphism in the aspartoacylase gene may cause misinterpretation
[82218] of Canavan gene testing. Europ. J. Hum. Genet. 6: 635-637, 1998.
[82219] 14278.

[82220] Rady, P. L.; Vargas, T.; Tyring, S. K.; Matalon, R.; Langen-beck,
[82221] U.: Novel missense mutation (Y231C) in a Turkish patient with Canavan
[82222] disease. (Letter) Am. J. Med. Genet. 87: 273-275, 1999.

[82223] 14279.

[82224] Schmidt, H.; Rott, H.-D.; Neuhauser, G.; Neumann, W.: Spongiose
[82225] Hirndystrophie im fruhen Kindesalter (Typ Canavan-van Bogaert-Bertrand):
[82226] Erkrankung von 3 Geschwistern einer nichtjudischen Fam-

ilie aus Oberfranken. Klin.
[82227] Paediat. 190: 580-585, 1978.

[82228] 14280.

[82229] Shaag, A.; Anikster, Y.; Christensen, E.; Glustein, J. Z.; Fois,
[82230] A.; Michelakakis, H.; Nigro, F.; Pronicka, E.; Ribes, A.; Zabot, M.
[82231] T.; Elpeleg, O. N.: The molecular basis of Canavan (aspartoacylase
[82232] deficiency) disease in European non-Jewish patients. Am. J. Hum.
[82233] Genet. 57: 572-580, 1995.

[82234] 14281.

[82235] Sistermans, E. A.; de Coo, R. F. M.; van Beerendonk, H. M.; Poll-The,
[82236] B. T.; Kleijer, W. J.; van Oost, B. A.: Mutation detection in the
[82237] aspartoacylase gene in 17 patients with Canavan disease: four new
[82238] mutations in the non-Jewish population. Europ. J. Hum. Genet. 8:
[82239] 557-560, 2000.

[82240] 14282.

[82241] Ungar, M.; Goodman, R. M.: Spongy degeneration of the brain in

[82242] Israel: a retrospective study. Clin. Genet. 23: 23-29, 1983.

[82243] 14283.

[82244] van Bogaert, L.: Familial spongy degeneration of the brain. (Complementary

[82245] study of the family R). Acta Psychiat. Neurol. Scand. 39: 107-113,

[82246] 1963.

[82247] 14284.

[82248] ZuRhein, G. M.; Eichman, P. L.; Puletti, F.: Familial idiocy

[82249] with spongy degeneration of the central nervous system of van Bogaert-Bertrand

[82250] type. Neurology 10: 998-1006, 1960.

[82251] 14285.

[82252] Gorlin, J. B.; Yamin, R.; Egan, S.; Stewart, M.; Stossel, T. P.;

[82253] Kwiatkowski, D. J.; Hartwig, J. H.: Human endothelial actin-binding

[82254] protein (ABP-280, nonmuscle filamin): a molecular leaf

spring. J.
[82255] Cell Biol. 111: 1089-1105, 1990.

[82256] 14286.

[82257] Huttenlocher, P. R.; Taravath, S.; Mojtahedi, S.: Periven-
tricular
[82258] heterotopia and epilepsy. Neurology 44: 51-55, 1994.

[82259] 14287.

[82260] Kunst, C. B.; Henske, E.; Hartwig, J. H.; Kwiatkowski, D. J.;
D'Urso,
[82261] M.; Bruns, G.; Warren, S. T.; Gorlin, J. B.: The dystrophin-
like
[82262] actin binding protein 280 gene maps between DXS52 and
G6PD overlapping
[82263] the Emery-Dreifuss muscular dystrophy locus. (Abstract)
Am. J. Hum.
[82264] Genet. 51: A21, 1992.

[82265] 14288.

[82266] Maestrini, E.; Patrosso, C.; Mancini, M.; Rivella, S.; Rocchi,

[82267] M.; Repetto, M.; Villa, A.; Frattini, A.; Zoppe, M.; Vezzoni,
P.;
[82268] Toniolo, D.: Mapping of two genes encoding isoforms of

the actin
[82269] binding protein ABP-280, a dystrophin like protein, to Xq28 and to
[82270] chromosome 7. Hum. Molec. Genet. 2: 761-766, 1993.

[82271] 14289.

[82272] Maestrini, E.; Rivella, S.; Tribioli, C.; Purtilo, D.; Rocchi,

[82273] M.; Archidiacono, N.; Toniolo, D.: Probes for CpG islands on the
[82274] distal long arm of the human X chromosome are clustered in Xq24 and
[82275] Xq28. Genomics 8: 664-670, 1990.

[82276] 14290.

[82277] Patrosso, M. C.; Repetto, M.; Villa, A.; Milanesi, L.; Frattini,

[82278] A.; Faranda, S.; Mancini, M.; Maestrini, E.; Toniolo, D.; Vezzoni,
[82279] P.: The exon-intron organization of the human X-linked gene (FLN1)
[82280] encoding actin-binding protein 280. Genomics 21: 71-76, 1994.
[82281] 14291.

[82282] Sheen, V. L.; Dixon, P. H.; Fox, J. W.; Hong, S. E.; Kinton,

L.;

[82283] Sisodiya, S. M.; Duncan, J. S.; Dubeau, F.; Scheffer, I. E.; Schachter,

[82284] S. C.; Wilner, A.; Henchy, R.; and 18 others: Mutations in the

[82285] X-linked filamin 1 gene cause periventricular nodular het-erotopia

[82286] in males as well as in females. Hum. Molec. Genet. 10: 1775-1783,

[82287] 2001.

[82288] 14292.

[82289] Small, K.; Wagener, M.; Warren, S. T.: Isolation and char-acterization

[82290] of the complete mouse emerin gene. Mammalian Genome 8: 337-341,

[82291] 1997.

[82292] 14293.

[82293] Vadlamudi, R. K.; Li, F.; Adam, L.; Nguyen, D.; Ohta, Y.; Stossel,

[82294] T. P.; Kumar, R.: Filamin is essential in actin cytoskeletal assembly

[82295] mediated by p21-activated kinase 1. Nature Cell Biol. 4:

681-690,
[82296] 2002.

[82297] 14294.

[82298] Faranda, S.; Frattini, A.; Vezzoni, P.: The human genes en-
coding
[82299] renin-binding protein and host cell factor are closely
linked in Xq28
[82300] and transcribed in the same direction. Gene 155:
237-239, 1995.
[82301] 14295.

[82302] Frattini, A.; Chatterjee, A.; Faranda, S.; Sacco, M. G.; Villa,

[82303] A.; Herman, G. E.; Vezzoni, P.: The chromosome localiza-
tion and the
[82304] HCF repeats of the human host cell factor gene (HCFC1)
are conserved
[82305] in the mouse homologue. Genomics 32: 277-280, 1996.

[82306] 14296.

[82307] Frattini, A.; Faranda, S.; Redolfi, E.; Zucchi, I.; Villa, A.;

[82308] Patrosso, M. C.; Strina, D.; Susani, L.; Vezzoni, P.: Genomic
organization
[82309] of the human VP16 accessory protein, a housekeeping

gene (HCFC1) mapping
[82310] to Xq28. Genomics 23: 30-35, 1994.

[82311] 14297.

[82312] Wilson, A. C.; LaMarco, K.; Peterson, M. G.; Herr, W.: The VP16
[82313] accessory protein HCF is a family of polypeptides processed from a
[82314] large precursor protein. Cell 74: 115-125, 1993.

[82315] 14298.

[82316] Wilson, A. C.; Parrish, J. E.; Massa, H. F.; Nelson, D. L.; Trask,
[82317] B. J.; Herr, W.: The gene encoding the VP16-accessory protein HCF
[82318] (HCFC1) resides in human Xq28 and is highly expressed in fetal tissues
[82319] and the adult kidney. Genomics 25: 462-468, 1995.

[82320] 14299.

[82321] Zoppe, M.; Frattini, A.; Faranda, S.; Vezzoni, P.: The complete
[82322] sequence of the host cell factor 1 (HCFC1) gene and its promoter:

[82323] a role for YY1 transcription factor in the regulation of its expression. Genomics 34:

[82324] 85-91, 1996.

[82325] 14300.

[82326] Maestrini, E.; Tamagnone, L.; Longati, P.; Cremona, O.; Gulisano,

[82327] M.; Bione, S.; Tamanini, F.; Neel, B. G.; Toniolo, D.; Comoglio, P.

[82328] M.: A family of transmembrane proteins with homology to the MET-hepatocyte

[82329] growth factor receptor. Proc. Nat. Acad. Sci. 93: 674-678, 1996.

[82330] 14301.

[82331] Tamagnone, L.; Artigiani, S.; Chen, H.; He, Z.; Ming, G.; Song,

[82332] H.; Chedotal, A.; Winberg, M. L.; Goodman, C. S.; Poo, M.; Tessier-Lavigne,

[82333] M.; Comoglio, P. M.: Plexins are a large family of receptors for

[82334] transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. Cell 99:

[82335] 71-80, 1999.

[82336] 14302.

[82337] Nolte, D.; Ramser, J.; Niemann, S.; Lehrach, H.; Sudbrak, R.; Muller,

[82338] U.: ACRC codes for a novel nuclear protein with unusual acidic repeat

[82339] tract and maps to DYT3 (dystonia parkinsonism) critical interval in

[82340] Xq13.1. Neurogenetics 3: 207-213, 2001.

[82341] 14303.

[82342] Kitagawa, H.; Fujita, M.; Ito, N.; Sugahara, K.: Molecular cloning

[82343] and expression of a novel chondroitin 6-O-sulfotransferase. J. Biol.

[82344] Chem. 275: 21075-21080, 2000.

[82345] 14304.

[82346] Becker, P. E.: Two new families of benign sex-linked re-cessive

[82347] muscular dystrophy. Rev. Canad. Biol. 21: 551-566, 1962.

[82348] 14305.

[82349] Becker, P. E.: Eine neue X-chromosomale Muskeldystro-phie. Acta

[82350] Psychiat. Neurol. Scand. 193: 427, 1955.

[82351] 14306.

[82352] Becker, P. E.: Neue Ergebnisse der Genetik der Muskeldys-
trophien. Acta
[82353] Genet. Statist. Med. 7: 303-310, 1957.

[82354] 14307.

[82355] Bushby, K. M. D.; Cleghorn, N. J.; Curtis, A.; Haggerty, I.
D.;
[82356] Nicholson, L. V. B.; Johnson, M. A.; Harris, J. B.; Bhat-
tacharya,
[82357] S. S.: Identification of a mutation in the promoter region
of the
[82358] dystrophin gene in a patient with atypical Becker muscular
dystrophy. Hum.
[82359] Genet. 88: 195-199, 1991.

[82360] 14308.

[82361] Doriguzzi, C.; Palmucci, L.; Mongini, T.; Chiado-Piat, L.;
Restagno,
[82362] G.; Ferrone, M.: Exercise intolerance and recurrent myo-
globinuria
[82363] as the only expression of Xp21 Becker type muscular dys-

trophy. J.

[82364] Neurol. 240: 269-271, 1993.

[82365] 14309.

[82366] England, S. B.; Nicholson, L. V. B.; Johnson, M. A.; Forrest,

[82367] S. M.; Love, D. R.; Zubrzycka-Gaarn, E. E.; Bulman, D. E.;
Harris,

[82368] J. B.; Davies, K. E.: Very mild muscular dystrophy associ-
ated with

[82369] the deletion of 46% dystrophin. Nature 343: 180-182,
1990.

[82370] 14310.

[82371] Derry, J. M. J.; Barnard, P. J.: Physical linkage of the A-
raf-1,

[82372] properdin, synapsin I, and TIMP genes on the human and
mouse X chromosomes. Genomics 12:

[82373] 632-638, 1992.

[82374] 14311.

[82375] Beales, P. L.; Katsanis, N.; Lewis, R. A.; Ansley, S. J.; El-
cioglu,

[82376] N.; Raza, J.; Woods, M. O.; Green, J. S.; Parfrey, P. S.;
Davidson,

[82377] W. S.; Lupski, J. R.: Genetic and mutational analyses of a large
[82378] multiethnic Bardet-Biedl cohort reveal a minor involvement of BBS6
[82379] and delineate the critical intervals of other loci. Am. J. Hum. Genet. 68:
[82380] 606-616, 2001.

[82381] 14312.

[82382] Slavotinek, A. M.; Searby, C.; Al-Gazali, L.; Hennekam, R. C. M.;
[82383] Schrander-Stumpel, C.; Orcana-Losa, M.; Pardo-Reoyo, S.; Cantani,
[82384] A.; Kumar, D.; Capellini, Q.; Neri, G.; Zackai, E.; Biesecker, L.
[82385] G.: Mutation analysis of the MKKS gene in McKusick-Kaufman syndrome
[82386] and selected Bardet-Biedl syndrome patients. Hum. Genet. 110: 561-567,
[82387] 2002.

[82388] 14313.

[82389] Stone, D. L.; Slavotinek, A.; Bouffard, G. G.; Banerjee-Basu, S.;

[82390] Baxevanis, A. D.; Barr, M.; Biesecker, L. G.: Mutation of a gene
[82391] encoding a putative chaperonin causes McKusick-Kaufman syndrome. Nature
[82392] Genet. 25: 79-82, 2000.

[82393] 14314.

[82394] Buhman, K. K.; Smith, S. J.; Stone, S. J.; Repa, J. J.; Wong, J.
[82395] S.; Knapp, F. F., Jr.; Burri, B. J.; Hamilton, R. L.; Abumrad, N.
[82396] A.; Farese, R. V., Jr.: DGAT1 is not essential for intestinal triacylglycerol
[82397] absorption or chylomicron synthesis. J. Biol. Chem. 277: 25474-25479,
[82398] 2002.

[82399] 14315.

[82400] Cases, S.; Smith, S. J.; Zheng, Y.-W.; Myers, H. M.; Lear, S. R.;
[82401] Sande, E.; Novak, S.; Collins, C.; Welch, C. B.; Lusis, A. J.; Erickson,
[82402] S. K.; Farese, R. V., Jr.: Identification of a gene encoding an acyl

[82403] CoA:diacylglycerol acyltransferase, a key enzyme in tria-
cylglycerol
[82404] synthesis. Proc. Nat. Acad. Sci. 95: 13018-13023, 1998.

[82405] 14316.

[82406] Cheng, D.; Meegalla, R. L.; He, B.; Cromley, D. A.; Bill-
heimer,
[82407] J. T.; Young, P. R.: Human acyl-CoA:diacylglycerol acyl-
transferase
[82408] is a tetrameric protein. Biochem. J. 359: 707-714, 2001.

[82409] 14317.

[82410] Grisart, B.; Coppieters, W.; Farnir, F.; Karim, L.; Ford, C.;
Berzi,
[82411] P.; Cambisano, N.; Mni, M.; Reid, S.; Simon, P.; Spelman,
R.; Georges,
[82412] M.; Snell, R.: Positional candidate cloning of a QTL in dairy
cattle:
[82413] identification of a missense mutation in the bovine DGAT1
gene with
[82414] major effect on milk yield and composition. Genome Res.
12: 222-231,
[82415] 2002.

[82416] 14318.

[82417] Haile, D. J.: Assignment of Slc11a3 to mouse chromosome 1 band

[82418] 1B and SLC11A3 to human chromosome 2q21 by in situ hybridization. Cytogenet.

[82419] Cell Genet. 88: 328-329, 2000.

[82420] 14319.

[82421] McKie, A. T.; Marciani, P.; Rolfs, A.; Brennan, K.; Wehr, K.; Barrow,

[82422] D.; Miret, S.; Bomford, A.; Peters, T. J.; Farzaneh, F.; Hediger,

[82423] M. A.; Hentze, M. W.; Simpson, R. J.: A novel duodenal iron-regulated

[82424] transporter, IREG1, implicated in the basolateral transfer of iron

[82425] to the circulation. Molec. Cell 5: 299-309, 2000.

[82426] 14320.

[82427] Montosi, G.; Donovan, A.; Totaro, A.; Garuti, C.; Pignatti, E.;

[82428] Cassanelli, S.; Trenor, C. C.; Gasparini, P.; Andrews, N. C.; Pietrangelo,

[82429] A.: Autosomal-dominant hemochromatosis is associated with a mutation
[82430] in the ferroportin (SLC11A3) gene. J. Clin. Invest. 108: 619-623,
[82431] 2001.

[82432] 14321.

[82433] Njajou, O. T.; Vaessen, N.; Joosse, M.; Berghuis, B.; van Dongen,
[82434] J. W. F.; Breuning, M. H.; Snijders, P. J. L. M.; Rutten, W. P. F.;
[82435] Sandkuijl, L. A.; Oostra, B. A.; van Duijn, C. M.; Heutink, P.: A
[82436] mutation in SLC11A3 is associated with autosomal dominant hemochromatosis. Nature
[82437] Genet. 28: 213-214, 2001.

[82438] 14322.

[82439] Roetto, A.; Merryweather-Clarke, A. T.; Daraio, F.; Livesey, K.;
[82440] Pointon, J. J.; Barbabietola, G.; Piga, A.; Mackie, P. H.; Robson,
[82441] K. J. H.; Camaschella, C.: A valine deletion of ferroportin 1: a

[82442] common mutation in hemochromatosis type 4? (Letter) Blood 100: 733-734,
[82443] 2002.

[82444] 14323.

[82445] Wallace, D. F.; Pedersen, P.; Dixon, J. L.; Stephenson, P.; Searle,
[82446] J. W.; Powell, L. W.; Subramaniam, V. N.: Novel mutation in ferroportin1
[82447] is associated with autosomal dominant hemochromatosis. Blood 100:
[82448] 692-694, 2002.

[82449] 14324.

[82450] Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S. L.;
[82451] Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J: The NIK
[82452] protein kinase and C17orf1 genes: chromosomal mapping, gene structures
[82453] and mutational screening in frontotemporal dementia and parkinsonism
[82454] linked to chromosome 17. Hum. Genet. 103: 340-345, 1998.

[82455] 14325.

[82456] Malinin, N. L.; Boldin, M. P.; Kovalenko, A. V.; Wallach, D.:

[82457] MAP3K-related kinase involved in NF-kappaB induction by TNF, CD95

[82458] and IL-1. Nature 385: 540-544, 1997.

[82459] 14326.

[82460] Shinkura, R.; Kitada, K.; Matsuda, F.; Tashiro, K.; Ikuta, K.;

[82461] Suzuki, M.; Kogishi, K.; Serikawa, T.; Honjo, T.: Alympho-plasia is

[82462] caused by a point mutation in the mouse gene encoding Nf-kappa-b-inducing

[82463] kinase. Nature Genet. 22: 74-77, 1999.

[82464] 14327.

[82465] Smith, C.; Andreakos, E.; Crawley, J. B.; Brennan, F. M.; Feldmann,

[82466] M.; Foxwell, B. M. J.: NF-kappa-B-inducing kinase is dis-pensable

[82467] for activation of NF-kappa-B in inflammatory settings but essential

[82468] for lymphotoxin beta receptor activation of NF-kappa-B in primary

[82469] human fibroblasts. J. Immun. 167: 5895-5903, 2001.

[82470] 14328.

[82471] Yin, L.; Wu, L.; Wesche, H.; Arthur, C. D.; White, J. M.;
Goeddel,
[82472] D. V.; Schreiber, R. D.: Defective lymphotoxin-beta recep-
tor-induced
[82473] NF-kappa-B transcriptional activity in NIK-deficient mice.
Science 291:
[82474] 2162-2165, 2001.

[82475] 14329.

[82476] Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S.
L.;
[82477] Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J.:
The NIK
[82478] protein kinase and C17orf1 genes: chromosomal map-
ping, gene structures
[82479] and mutational screening in frontotemporal dementia and
parkinsonism
[82480] linked to chromosome 17. Hum. Genet. 103: 340-345,
1998.
[82481] 14330.

[82482] Yayoshi-Yamamoto, S.; Taniuchi, I.; Watanabe, T.: FRL, a novel
[82483] formin-related protein, binds to Rac and regulates cell motility and
[82484] survival of macrophages. Molec. Cell. Biol. 20: 6872-6881, 2000.
[82485] 14331.

[82486] Borggrefe, T.; Masat, L.; Wabl, M.; Riwar, B.; Cattoretti, G.;
[82487] Jessberger, R.: Cellular, intracellular, and developmental expression
[82488] patterns of murine SWAP-70. Europ. J. Immun. 29: 1812-1822, 1999.
[82489] 14332.

[82490] Borggrefe, T.; Wabl, M.; Akhmedov, A. T.; Jessberger, R.: A B-cell-specific
[82491] DNA recombination complex. J. Biol. Chem. 273: 17025-17035, 1998.
[82492] 14333.

[82493] Masat, L.; Caldwell, J.; Armstrong, R.; Khoshnevisan, H.; Jessberger,
[82494] R.; Herndier, B.; Wabl, M.; Ferrick, D.: Association of SWAP-70 with

[82495] the B cell antigen receptor complex. Proc. Nat. Acad. Sci. 97: 2180-2184,
[82496] 2000.

[82497] 14334.

[82498] Masat, L.; Liddell, R. A.; Mock, B. A.; Kuo, W.-L.; Jessberger,
[82499] R.; Wabl, M.; Morse, H. C., III: Mapping of the SWAP70 gene to mouse
[82500] chromosome 7 and human chromosome 11p15. Immunogenetics 51: 16-19,
[82501] 2000.

[82502] 14335.

[82503] Shinohara, M.; Terada, Y.; Iwamatsu, A.; Shinohara, A.; Mochizuki,
[82504] N.; Higuchi, M.; Gotoh, Y.; Ihara, S.; Nagata, S.; Itoh, H.; Fukui,
[82505] Y.; Jessberger, R.: SWAP-70 is a guanine-nucleotide-exchange factor
[82506] that mediates signalling of membrane ruffling. Nature 416: 759-763,
[82507] 2002.

[82508] 14336.

[82509] Kourlas, P. J.; Strout, M. P.; Becknell, B.; Veronese, M. L.; Croce,

[82510] C. M.; Theil, K. S.; Krahe, R.; Ruutu, T.; Knuutila, S.; Bloomfield,

[82511] C. D.; Caligiuri, M. A.: Identification of a gene at 11q23 encoding

[82512] a guanine nucleotide exchange factor: evidence for its fu- sion with

[82513] MLL in acute myeloid leukemia. Proc. Nat. Acad. Sci. 97: 2145-2150,

[82514] 2000.

[82515] 14337.

[82516] Miura, K.; Suzuki, K.; Tokino, T.; Isomura, M.; Inazawa, J.; Matsuno,

[82517] S.; Nakamura, Y.: Detailed deletion mapping in squamous cell carcinomas

[82518] of the esophagus narrows a region containing a putative tumor suppressor

[82519] gene to about 200 kilobases on distal chromosome 9q. Cancer Res. 56:

[82520] 1629-1634, 1996.

[82521] 14338.

[82522] Nishiwaki, T.; Daigo, Y.; Kawasoe, T.; Nakamura, Y.: Isola-
tion
[82523] and mutational analysis of a novel human cDNA, DEC1
(deleted in esophageal
[82524] cancer 1), derived from the tumor suppressor locus in
9q32. Genes
[82525] Chromosomes Cancer 27: 169-176, 2000.

[82526] 14339.

[82527] Delettre, C.; Griffoin, J.-M.; Kaplan, J.; Dollfus, H.; Lorenz,
[82528] B.; Faivre, L.; Lenaers, G.; Belenguer, P.; Hamel, C. P.: Mu-
tation
[82529] spectrum and splicing variants in the OPA1 gene. Hum.
Genet. 109:
[82530] 584-591, 2001.

[82531] 14340.

[82532] Thompson, M. A.; Moon, E.; Kim, U.-J.; Xu, J.; Siciliano, M.
J.;
[82533] Weinshilboum, R. M.: Human indolethylamine N-
methyltransferase: cDNA
[82534] cloning and expression, gene cloning, and chromosomal

localization. Genomics 61:
[82535] 285-297, 1999.

[82536] 14341.

[82537] Bespalova, I. N.; Burmeister, M.: Identification of a novel LIM
[82538] domain gene, LMCD1, and chromosomal localization in human and mouse. Genomics 63:
[82539] 69-74, 2000.

[82540] 14342.

[82541] Valladeau, J.; Clair-Moninot, V.; Dezutter-Dambuyant, C.; Pin,
[82542] J.-J.; Kissenpfennig, A.; Mattei, M.-G.; Ait-Yahia, S.; Bates, E.
[82543] E. M.; Malissen, B.; Koch, F.; Fossiez, F.; Romani, N.; Lebecque,
[82544] S.; Saeland, S.: Identification of mouse langerin/CD207 in Langerhans
[82545] cells and some dendritic cells of lymphoid tissues. J. Im-
mun. 168:
[82546] 782-792, 2002.

[82547] 14343.

[82548] Valladeau, J.; Ravel, O.; Dezutter-Dambuyant, C.; Moore, K.; Kleijmeer,

[82549] M.; Liu, Y.; Duvert-Frances, V.; Vincent, C.; Schmitt, D.; Davoust,

[82550] J.; Caux, C.; Lebecque, S.; Saeland, S.: Langerin, a novel C-type

[82551] lectin specific to Langerhans cells, is an endocytic recep-tor that

[82552] induces the formation of Birbeck granules. Immunity 12: 71-81, 2000.

[82553] 14344.

[82554] Ruiz, A.; Kuehn, M. H.; Andorf, J. L.; Stone, E.; Hageman, G. S.;

[82555] Bok, D.: Genomic organization and mutation analysis of the gene encoding
[82556] lecithin retinol acyltransferase in human retinal pigment epithelium. Invest.
[82557] Ophthal. Vis. Sci. 42: 31-37, 2001.

[82558] 14345.

[82559] Ruiz, A.; Winston, A.; Lim, Y.-H.; Gilbert, B. A.; Rando, R. R.;
[82560] Bok, D.: Molecular and biochemical characterization of lecithin retinol
[82561] acyltransferase. J. Biol. Chem. 274: 3834-3841, 1999.

[82562] 14346.

[82563] Thompson, D. A.; Li, Y.; McHenry, C. L.; Carlson, T. J.; Ding,
[82564] X.; Sieving, P. A.; Apfelstedt-Sylla, E.; Gal, A.: Mutations in the
[82565] gene encoding lecithin retinol acyltransferase are associated with
[82566] early-onset severe retinal dystrophy. Nature Genet. 28: 123-124,
[82567] 2001.

[82568] 14347.

[82569] Reichenberger, E.; Tiziani, V.; Watanabe, S.; Park, L.; Ueki, Y.;

[82570] Santanna, C.; Baur, S. T.; Shiang, R.; Grange, D. K.; Beighton, P.;

[82571] Gardner, J.; Hamersma, H.; Sellars, S.; Ramesar, R.; Lidral, A. C.;

[82572] Sommer, A.; Raposo do Amaral, C. M.; Gorlin, R. J.; Mul-

liken, J. B.;

[82573] Olsen, B. R.: Autosomal dominant craniometaphyseal dys-

plasia is caused

[82574] by mutations in the transmembrane protein ANK. Am. J. Hum. Genet. 68:

[82575] 1321-1326, 2001.

[82576] 14348.

[82577] Im, D.-S.; Heise, C. E.; Ancellin, N.; O'Dowd, B. F.; Shei, G.;

[82578] Heavens, R. P.; Rigby, M. R.; Hla, T.; Mandala, S.; McAllis-

ter, G.;

[82579] George, S. R.; Lynch, K. R.: Characterization of a novel sphingosine

[82580] 1-phosphate receptor, Edg-8. J. Biol. Chem. 275: 14281-14286, 2000.

[82581] 14349.

[82582] Li, Z.; Yao, K.; Cao, Y.: Molecular cloning of a novel tissue-
specific
[82583] gene from human nasopharyngeal epithelium. Gene 237:
235-240, 1999.
[82584] 14350.

[82585] Derst, C.; Engel, H.; Grzeschik, K.-H.; Daut, J.: Genomic
structure
[82586] and chromosome mapping of human and mouse RAMP
genes. Cytogenet.
[82587] Cell Genet. 90: 115-118, 2000.

[82588] 14351.

[82589] Amler, L. C.; Bauer, A.; Corvi, R.; Dihlmann, S.; Praml, C.;
Cavenee,
[82590] W. K.; Schwab, M.; Hampton, G. M.: Identification and
characterization
[82591] of novel genes located at the t(1;15)(p36.2;q24) translo-
cation breakpoint
[82592] in the neuroblastoma cell line NGP. Genomics 64:
195-202, 2000.
[82593] 14352.

[82594] Yanagisawa, H.; Bundo, M.; Miyashita, T.; Okamura-Oho, Y.; Tadokoro,

[82595] K.; Tokunaga, K.; Yamada, M.: Protein binding of a DRPLA family through

[82596] arginine-glutamic acid dipeptide repeats is enhanced by extended polyglutamine. Hum.

[82597] Molec. Genet. 9: 1433-1442, 2000.

[82598] 14353.

[82599] Oh, J.; Takahashi, R.; Kondo, S.; Mizoguchi, A.; Adachi, E.; Sasahara,

[82600] R. M.; Nishimura, S.; Imamura, Y.; Kitayama, H.; Alexander, D. B.;

[82601] Ide, C.; Horan, T. P.; Arakawa, T.; Yoshida, H.; Nishikawa, S.; Itoh,

[82602] Y.; Seiki, M.; Itohara, S.; Takahashi, C.; Noda, M.: The membrane-anchored

[82603] MMP inhibitor RECK is a key regulator of extracellular matrix integrity

[82604] and angiogenesis. Cell 107: 789-800, 2001.

[82605] 14354.

[82606] Takahashi, C.; Sheng, Z.; Horan, T. P.; Kitayama, H.; Maki, M.;

[82607] Hitomi, K.; Kitaura, Y.; Takai, S.; Sasahara, R. M.; Horimoto, A.;
[82608] Ikawa, Y.; Ratzkin, B. J.; Arakawa, T.; Noda, M.: Regulation of matrix
[82609] metalloproteinase-9 and inhibition of tumor invasion by the membrane-anchored
[82610] glycoprotein RECK. Proc. Nat. Acad. Sci. 95: 13221-13226, 1998.
[82611] 14355.

[82612] Nagase, T.; Ishikawa, I.; Nakajima, D.; Ohira, M.; Seki, N.; Miyajima,
[82613] N.; Tanaka, A.; Kotani, H.; Nomura, N.; O'Hara, O.: Prediction of
[82614] the coding sequences of unidentified human genes. VII. The complete
[82615] sequences of 100 new cDNA clones from brain which can code for large
[82616] proteins in vitro. DNA Res. 4: 141-150, 1997.

[82617] 14356.

[82618] Johnston, R. L.; Seller, M. J.; Behnam, J. T.; Burdon, M. A.; Spalton,
[82619] D. J.: Dominant optic atrophy: refining the clinical diag-

nostic criteria
[82620] in light of genetic linkage studies. Ophthalmology 106: 123-128,
[82621] 1999.

[82622] 14357.

[82623] Jones, B. A.; Fangman, W. L.: Mitochondrial DNA maintenance in
[82624] yeast requires a protein containing a region related to the GTP-binding
[82625] domain of dynamin. Genes Dev. 6: 380-389, 1992.

[82626] 14358.

[82627] Pelloquin, L.; Belenguer, P.; Menon, Y.; Ducommun, B.: Identification
[82628] of a fission yeast dynamin-related protein involved in mitochondrial
[82629] DNA maintenance. Biochem. Biophys. Res. Commun. 251: 720-726, 1998.
[82630] 14359.

[82631] Pelloquin, L.; Belenguer, P.; Menon, Y.; Gas, N.; Ducommun, B.
[82632] : Fission yeast Msp1 is a mitochondrial dynamin-related

protein. J.

[82633] Cell Sci. 112: 4151-4161, 1999.

[82634] 14360.

[82635] Pesch, U. E. A.; Leo-Kottler, B.; Mayer, S.; Jurklies, B.; Kellner,

[82636] U.; Apfelstedt-Sylla, E.; Zrenner, E.; Alexander, C.; Wissinger, B.

[82637] : OPA1 mutations in patients with autosomal dominant optic atrophy

[82638] and evidence for semi-dominant inheritance. Hum. Molec. Genet. 10:

[82639] 1359-1368, 2001.

[82640] 14361.

[82641] Thiselton, D. L.; Alexander, C.; Morris, A.; Brooks, S.; Rosenberg,

[82642] T.; Eiberg, H.; Kjer, B.; Kjer, P.; Bhattacharya, S. S.; Votruba,

[82643] M.: A frameshift mutation in exon 28 of the OPA1 gene explains the

[82644] high prevalence of dominant optic atrophy in the Danish population:

[82645] evidence for a founder effect. Hum. Genet. 109: 498-502,

2001.

[82646] 14362.

[82647] Toomes, C.; Marchbank, N. J.; Mackey, D. A.; Craig, J. E.; Newbury-Ecob,

[82648] R. A.; Bennett, C. P.; Vize, C. J.; Desai, S. P.; Black, G. C. M.;

[82649] Patel, N.; Teimory, M.; Markham, A. F.; Inglehearn, C. F.; Churchill,

[82650] A. J.: Spectrum, frequency and penetrance of OPA1 muta-tions in dominant

[82651] optic atrophy. Hum. Molec. Genet. 10: 1369-1378, 2001.

[82652] 14363.

[82653] Muto, A.; Hoshino, H.; Madisen, L.; Yanai, N.; Obinata, M.; Karasuyama,

[82654] H.; Hayashi, N.; Nakauchi, H.; Yamamoto, M.; Groudine, M.; Igarashi,

[82655] K.: Identification of Bach2 as a B-cell-specific partner for small

[82656] Maf proteins that negatively regulate the immunoglobulin heavy chain

[82657] gene 3-prime enhancer. EMBO J. 17: 5734-5743, 1998.

[82658] 14364.

[82659] Sasaki, S.; Ito, E.; Toki, T.; Maekawa, T.; Kanezaki, R.; Umenai,
[82660] T.; Muto, A.; Nagai, H.; Kinoshita, T.; Yamamoto, M.; In-azawa, J.;
[82661] Taketo, M. M.; Nakahata, T.; Igarashi, K.; Yokoyama, M.: Cloning
[82662] and expression of human B cell-specific transcription fac-tor BACH2
[82663] mapped to chromosome 6q15. Oncogene 19: 3739-3749, 2000.
[82664] 14365.

[82665] Shimaoka, T.; Kume, N.; Minami, M.; Hayashida, K.; Kataoka, H.;
[82666] Kita, T.; Yonehara, S.: Molecular cloning of a novel scav-enger receptor
[82667] for oxidized low density lipoprotein, SR-PSOX, on macrophages. J.
[82668] Biol. Chem. 275: 40663-40666, 2000.

[82669] 14366.

[82670] Wilbanks, A.; Zondlo, S. C.; Murphy, K.; Mak, S.; Soler, D.; Langdon,
[82671] P.; Andrew, D. P.; Wu, L.; Briskin, M.: Expression cloning of

the
[82672] STRL33/BONZO/TYMSTR ligand reveals elements of CC, CXC, CX3C chemokines. J.
[82673] Immun. 166: 5145-5154, 2001.

[82674] 14367.

[82675] Matsumoto, T.; Shimamoto, A.; Goto, M.; Furuichi, Y.: Im-
paired
[82676] nuclear localization of defective DNA helicases in Werner's syndrome.
[82677] (Letter) Nature Genet. 16: 335-336, 1997.

[82678] 14368.

[82679] Meisslitzer, C.; Ruppitsch, W.; Weirich-Schwaiger, H.; Weirich,
[82680] H. G.; Jabkowsky, J.; Klein, G.; Schweiger, M.; Hirsch-Kauffmann,
[82681] M.: Werner syndrome: characterization of mutations in the WRN gene
[82682] in an affected family. Europ. J. Hum. Genet. 5: 364-370, 1997.
[82683] 14369.

[82684] Monnat, R. J., Jr.: Unusual features of thyroid carcinomas

in
[82685] Japanese patients with Werner syndrome and possible genotype-phenotype
[82686] relations to cell type and race. (Letter) Cancer 86: 728-729, 1999.
[82687] 14370.
[82688] Moser, M. J.; Oshima, J.; Monnat, R. J., Jr.: WRN mutations in
[82689] Werner syndrome. Hum. Mutat. 13: 271-279, 1999.
[82690] 14371.
[82691] Oshima, J.; Yu, C.-E.; Piussan, C.; Klein, G.; Jabkowski, J.;
[82692] Balci, S.; Miki, T.; Nakura, J.; Ogihara, T.; Ells, J.; Smith, M.
[82693] A. C.; Melaragno, M. I.; Fraccaro, M.; Scappaticci, S.; Matthews,
[82694] J.; Ouais, S.; Jarzebowicz, A.; Schellenberg, G. D.; Martin, G. M.
[82695] : Homozygous and compound heterozygous mutations at the Werner syndrome
[82696] locus. Hum. Molec. Genet. 5: 1909-1913, 1996.
[82697] 14372.
[82698] Sinclair, D. A.; Guarente, L.: Extrachromosomal rDNA cir-

cles--a
[82699] cause of aging in yeast. Cell 91: 1033-1042, 1997.

[82700] 14373.

[82701] Wyllie, F. S.; Jones, C. J.; Skinner, J. W.; Haughton, M. F.;

[82702] Wallis, C.; Wynford-Thomas, D.; Faragher, R. G. A.;
Kipling, D.:
[82703] Telomerase prevents the accelerated cell ageing of Werner
syndrome
[82704] fibroblasts. (Letter) Nature Genet. 24: 16-17, 2000.

[82705] 14374.

[82706] Yamagata, K.; Kato, J.; Shimamoto, A.; Goto, M.; Furuichi,
Y.;
[82707] Ikeda, H.: Bloom's and Werner's syndrome genes suppress
hyperrecombination
[82708] in yeast sgs1 mutant: implication for genomic instability
in human
[82709] diseases. Proc. Nat. Acad. Sci. 95: 8733-8738, 1998.

[82710] 14375.

[82711] Yan, H.; Chen, C.-Y.; Kobayashi, R.; Newport, J.: Replica-
tion
[82712] focus-forming activity 1 and the Werner syndrome gene

product. Nature
[82713] Genet. 19: 375-378, 1998.

[82714] 14376.

[82715] Yan, H.; Newport, J.: FFA-1, a protein that promotes the formation
[82716] of replication centers within nuclei. Science 269: 1883-1885, 1995.
[82717] 14377.

[82718] Hughes-Benzie, R. M.; Pilia, G.; Xuan, J. Y.; Hunter, A. G. W.;
[82719] Chen, E.; Golabi, M.; Hurst, J. A.; Kobori, J.; Marymee, K.; Pagon,
[82720] R. A.: Simpson-Golabi-Behmel syndrome: genotype/phenotype analysis
[82721] of 18 affected males from 7 unrelated families. Am. J. Med. Genet. 66:
[82722] 227-234, 1996.

[82723] 14378.

[82724] Lindsay, S.; Ireland, M.; O'Brien, O.; Clayton-Smith, J.; Hurst,
[82725] J. A.; Mann, J.; Cole, T.; Sampson, J.; Slaney, S.; Sch-

lessinger,

[82726] D.; Burn, J.; Pilia, G.: Large scale deletions in the GPC3 gene may

[82727] account for a minority of cases of Simpson-Golabi-Behmel syndrome. J.

[82728] Med. Genet. 34: 480-483, 1997.

[82729] 14379.

[82730] Pilia, G.; Hughes-Benzie, R. M.; MacKenzie, A.; Baybayan, P.; Chen,

[82731] E. Y.; Huber, R.; Neri, G.; Cao, A.; Forabosco, A.; Sch-lessinger,

[82732] D.: Mutations in GPC3, a glypican gene, cause the Simp-son-Golabi-Behmel

[82733] overgrowth syndrome. Nature Genet. 12: 241-247, 1996.

[82734] 14380.

[82735] Shen, T.; Sonoda, G.; Hamid, J.; Li, M.; Filmus, J.; Buick, R.

[82736] N.; Testa, J. R.: Mapping of the Simpson-Golabi-Behmel overgrowth

[82737] syndrome gene (GPC3) to chromosome X in human and rat by fluorescence

[82738] in situ hybridization. Mammalian Genome 8: 72 only,

1997.

[82739] 14381.

[82740] Veugelers, M.; De Cat, B.; Muyldermans, S. Y.; Reekmans, G.; Delande,

[82741] N.; Frints, S.; Legius, E.; Fryns, J.-P.; Schrander-Stumpel, C.; Weidle,

[82742] B.; Magdalena, N.; David, G.: Mutational analysis of the GPC3/GPC4

[82743] glypican gene cluster on Xq26 in patients with Simpson-Go-labi-Behmel

[82744] syndrome: identification of loss-of-function mutations in the GPC3

[82745] gene. Hum. Molec. Genet. 9: 1321-1328, 2000.

[82746] 14382.

[82747] Weksberg, R.; Squire, J. A.; Templeton, D. M.: Glypicans: a growing

[82748] trend. Nature Genet. 12: 225-227, 1996.

[82749] 14383.

[82750] Xuan, J. Y.; Hughes-Benzie, R. M.; MacKenzie, A. E.: A small

[82751] interstitial deletion in the GPC3 gene causes Simpson-Go-

labi-Behmel
[82752] syndrome in a Dutch-Canadian family. J. Med. Genet. 36: 57-58, 1999.
[82753] 14384.
[82754] Brown, C. J.; Miller, A. P.; Carrel, L.; Rupert, J. L.; Davies,
[82755] K. E.; Willard, H. F.: The DXS423E gene in Xp11.21 escapes X chromosome
[82756] inactivation. Hum. Molec. Genet. 4: 251-255, 1995.
[82757] 14385.
[82758] Sumara, I.; Vorlaufer, E.; Gieffers, C.; Peters, B. H.; Peters,
[82759] J.-M.: Characterization of vertebrate cohesin complexes and their
[82760] regulation in prophase. J. Cell Biol. 151: 749-761, 2000.
[82761] 14386.
[82762] Teague, P. W.; Aldred, M. A.; Jay, M.; Dempster, M.; Harri-son,
[82763] C.; Carothers, A. D.; Hardwick, L. J.; Evans, H. J.; Strain, L.; Brock,
[82764] D. J. H.; Bundey, S.; Jay, B.; Bird, A. C.; Bhattacharya, S. S.; Wright,
[82765] A. F.: Heterogeneity analysis in 40 X-linked retinitis pig-

mentosa
[82766] families. Am. J. Hum. Genet. 55: 105-111, 1994.

[82767] 14387.

[82768] Bodrug, S. E.; Ray, P. N.; Gonzalez, I. L.; Schmickel, R. D.;

[82769] Sylvester, J. E.; Worton, R. G.: Molecular analysis of a constitutional
[82770] X-autosome translocation in a female with muscular dystrophy. Science 237:
[82771] 1620-1624, 1987.

[82772] 14388.

[82773] Boyce, F. M.; Beggs, A. H.; Feener, C.; Kunkel, L. M.: Dystrophin
[82774] is transcribed in brain from a distant upstream promoter. Proc. Nat.
[82775] Acad. Sci. 88: 1276-1280, 1991.

[82776] 14389.

[82777] Boyd, Y.; Buckle, V. J.: Cytogenetic heterogeneity of translocations
[82778] associated with Duchenne muscular dystrophy. Clin. Genet. 29: 108-115,
[82779] 1986.

[82780] 14390.

[82781] Bulman, D. E.; Gangopadhyay, S. B.; Bebchuck, K. G.; Wor-
ton, R.
[82782] G.; Ray, P. N.: Point mutation in the human dystrophin
gene: identification
[82783] through Western blot analysis. Genomics 10: 457-460,
1991.
[82784] 14391.

[82785] Burke, J. F.; Mogg, A. E.: Suppression of a nonsense muta-
tion
[82786] in mammalian cells in vivo by the aminoglycoside antibi-
otics G-418
[82787] and paromomycin. Nucleic Acids Res. 13: 6265-6272,
1985.
[82788] 14392.

[82789] Burnette, W. N.: 'Western blotting': electrophoretic transfer
[82790] of proteins from sodium dodecyl sulfate-polyacrylamide
gels to unmodified
[82791] nitrocellulose and radiographic detection with antibody
and radioiodinated
[82792] protein A. Anal. Biochem. 112: 195-203, 1981.

[82793] 14393.

[82794] Chamberlain, J. S.; Pearlman, J. A.; Muzny, D. M.; Gibbs, R. A.;

[82795] Ranier, J. E.; Reeves, A. A.; Caskey, C. T.: Expression of the murine

[82796] Duchenne muscular dystrophy gene in muscle and brain. Science 239:

[82797] 1416-1418, 1988.

[82798] 14394.

[82799] Chelly, J.; Concordet, J.-P.; Kaplan, J.-C.; Kahn, A.: Illegiti-mate

[82800] transcription: transcription of any gene in any cell type. Proc.

[82801] Nat. Acad. Sci. 86: 2617-2621, 1989.

[82802] 14395.

[82803] Chelly, J.; Gilgenkrantz, H.; Hugnot, J. P.; Hamard, G.; Lambert,

[82804] M.; Recan, D.; Akli, S.; Cometto, M.; Kahn, A.; Kaplan, J. C.: Illegitimate

[82805] transcription: application to the analysis of truncated transcripts

[82806] of the dystrophin gene in nonmuscle cultured cells from Duchenne and
[82807] Becker patients. J. Clin. Invest. 88: 1161-1166, 1991.

[82808] 14396.

[82809] Chelly, J.; Hamard, G.; Koulakoff, A.; Kaplan, J.-C.; Kahn, A.;
[82810] Berwald-Netter, Y.: Dystrophin gene transcribed from different promoters
[82811] in neuronal and glial cells. Nature 344: 64-65, 1990.

[82812] 14397.

[82813] Chelly, J.; Kaplan, J.-C.; Maire, P.; Gautron, S.; Kahn, A.:

[82814] Transcription of the dystrophin gene in human muscle and non-muscle
[82815] tissues. Nature 333: 858-860, 1988.

[82816] 14398.

[82817] Clemens, P. R.; Ward, P. A.; Caskey, C. T.; Bulman, D. E.; Fenwick,
[82818] R. G.: Premature chain termination mutation causing Duchenne muscular
[82819] dystrophy. Neurology 42: 1775-1782, 1992.

[82820] 14399.

[82821] Cooper, B. J.; Valentine, B. A.; Wilson, S.; Patterson, D. F.;

[82822] Concannon, P. W.: Canine muscular dystrophy: confirma-
tion of X-linked

[82823] inheritance. J. Hered. 79: 405-408, 1988.

[82824] 14400.

[82825] Covone, A. E.; Lerone, M.; Romeo, G.: Genotype-pheno-
type correlation

[82826] and germline mosaicism in DMD/BMD patients with dele-
tions of the dystrophin

[82827] gene. Hum. Genet. 87: 353-360, 1991.

[82828] 14401.

[82829] Cox, G. A.; Cole, N. M.; Matsumura, K.; Phelps, S. F.;
Hauschka,

[82830] S. D.; Campbell, K. P.; Faulkner, J. A.; Chamberlain, J. S.:
Overexpression

[82831] of dystrophin in transgenic mdx mice eliminates dys-
trophic symptoms

[82832] without toxicity. Nature 364: 725-729, 1993.

[82833] 14402.

[82834] Cox, G. A.; Sunada, Y.; Campbell, K. P.; Chamberlain, J. S.:

[82835] Dp71 can restore the dystrophin-associated glycoprotein complex in

[82836] muscle but fails to prevent dystrophy. Nature Genet. 8: 333-339,

[82837] 1994.

[82838] 14403.

[82839] Comi, G. P.; Ciafaloni, E.; de Silva, H. A. R.; Prelle, A.; Bar-doni,

[82840] A.; Rigoletto, C.; Robotti, M.; Bresolin, N.; Moggio, M.; Fortunato,

[82841] F.; Ciscato, P.; Turconi, A.; Rose, A. D.; Scarlato, G.: A G(+1)-to-A

[82842] transversion at the 5-prime splice site of intron 69 of the dystrophin

[82843] gene causing the absence of peripheral nerve Dp116 and severe clinical

[82844] involvement in a DMD patient. Hum. Molec. Genet. 4: 2171-2174, 1995.

[82845] 14404.

[82846] Crawford, G. E.; Lu, Q. L.; Partridge, T. A.; Chamberlain, J.

[82847] S.: Suppression of revertant fibers in mdx mice by expres-
sion of
[82848] a functional dystrophin. Hum. Molec. Genet. 10:
2745-2750, 2001.
[82849] 14405.

[82850] Darras, B. T.; Blattner, P.; Harper, J. F.; Spiro, A. J.; Alter,

[82851] S.; Francke, U.: Intragenic deletions in 21 Duchenne mus-
cular dystrophy
[82852] (DMD)/Becker muscular dystrophy (BMD) families studied
with the dystrophin
[82853] cDNA: location of breakpoints on HindIII and BglII exon-
containing
[82854] fragment maps, meiotic and mitotic origin of the muta-
tions. Am. J.
[82855] Hum. Genet. 43: 620-629, 1988.

[82856] 14406.

[82857] Darras, B. T.; Francke, U.: Normal human genomic restric-
tion-fragment
[82858] patterns and polymorphisms revealed by hybridization
with the entire
[82859] dystrophin cDNA. Am. J. Hum. Genet. 43: 612-619, 1988.

[82860] 14407.

[82861] Darras, B. T.; Francke, U.: A partial deletion of the muscu-
lar
[82862] dystrophy gene transmitted twice by an unaffected male.
Nature 329:
[82863] 556-558, 1987.

[82864] 14408.

[82865] Davies, K. E.; Smith, T. J.; Bundey, S.; Read, A. P.; Flint, T.;

[82866] Bell, M.; Speer, A.: Mild and severe muscular dystrophy
associated
[82867] with deletions in Xp21 of the human X chromosome. J.
Med. Genet. 25:
[82868] 9-13, 1988.

[82869] 14409.

[82870] De Angelis, F. G.; Sthandier, O.; Berarducci, B.; Toso, S.;
Galluzzi,
[82871] G.; Ricci, E.; Cossu, G.; Bozzoni, I.: Chimeric snRNA
molecules carrying
[82872] antisense sequences against the splice junctions of exon
51 of the
[82873] dystrophin pre-mRNA induce exon skipping and restora-

tion of a dystrophin
[82874] synthesis in delta-48-50 DMD cells. Proc Nat. Acad. Sci. 99: 9456-9461,
[82875] 2002.

[82876] 14410.

[82877] den Dunnen, J. T.; Bakker, E.; Klein Breteler, E. G.; Pearson,

[82878] P. L.; van Ommen, G. J. B.: Direct mutation of more than 50% of the
[82879] Duchenne muscular dystrophy mutations by field inversion gels. Nature 329:
[82880] 640-642, 1987.

[82881] 14411.

[82882] Dickson, G.; Pizzey, J. A.; Elsom, V. E.; Love, D.; Davies, K.

[82883] E.; Walsh, F. S.: Distinct dystrophin mRNA species are expressed
[82884] in embryonic and adult mouse skeletal muscle. FEBS Lett. 242: 47-52,
[82885] 1988.

[82886] 14412.

[82887] Dominguez-Steglich, M.; Meng, G.; Bettecken, T.; Muller,

C. R.;

[82888] Schmid, M.: The dystrophin gene is autosomally located on a microchromosome

[82889] in chicken. Genomics 8: 536-540, 1990.

[82890] 14413.

[82891] Doolittle, R. F.: Similar amino acid sequences: chance or common

[82892] ancestry? Science 214: 149-159, 1981.

[82893] 14414.

[82894] Zhou, J.; Mochizuki, T.; Smeets, H.; Antignac, C.; Laurila, P.;

[82895] de Paepe, A.; Tryggvason, K.; Reeders, S. T.: Deletion of the paired

[82896] alpha-5(IV) and alpha-6(IV) collagen genes in inherited smooth muscle

[82897] tumors. Science 261: 1167-1169, 1993.

[82898] 14415.

[82899] Bolscher, B. G. J. M.; de Boer, M.; de Klein, A.; Weening, R.

[82900] S.; Roos, D.: Point mutations in the beta-subunit of cytochrome b(558)

[82901] leading to X-linked chronic granulomatous disease. Blood

77: 2482-2487,

[82902] 1991.

[82903] 14416.

[82904] Borregaard, N.; Cross, A. R.; Herlin, T.; Jones, O. T. G.; Se-
gal,

[82905] A. W.; Valerius, N. H.: A variant form of X-linked chronic
granulomatous

[82906] disease with normal nitroblue tetrazolium slide test and
cytochrome

[82907] b. Europ. J. Clin. Invest. 13: 243-248, 1983.

[82908] 14417.

[82909] Branch, D. R.; Gaidulis, L.; Lazar, G. S.: Human granulo-
cytes

[82910] lack red cell Kx antigen. Brit. J. Haemat. 62: 747-755,
1986.

[82911] 14418.

[82912] Brandrup, F.; Koch, C.; Petri, M.; Schiodt, M.; Johansen, K.
S.

[82913] : Discoid lupus erythematosus-like lesions and stomatitis
in female

[82914] carriers of X-linked chronic granulomatous disease. Brit. J.

Derm. 104:
[82915] 495-505, 1981.

[82916] 14419.

[82917] Brockdorff, N.; Fisher, E. M. C.; Orkin, S. H.; Lyon, M. F.; Brown,
[82918] S. D. M.: Localization of the human X-linked gene for chronic granulomatous
[82919] disease to the mouse X chromosome: implications for X-chromosome evolution. Cytogenet.
[82920] Cell Genet. 48: 124-125, 1988.

[82921] 14420.

[82922] Brzica, S. M., Jr.; Rhodes, K. H.; Pineda, A. A.; Taswell, H.
[82923] F.: Chronic granulomatous disease and the McLeod phenotype: successful
[82924] treatment of infection with granulocyte transfusions resulting in
[82925] subsequent hemolytic transfusion reaction. Mayo Clin. Proc. 52:
[82926] 153-156, 1977.

[82927] 14421.

[82928] Buescher, E. S.; Alling, D. W.; Gallin, J. I.: Use of an X-

linked
[82929] human neutrophil marker to estimate timing of lyonization and size
[82930] of the dividing stem cell pool. J. Clin. Invest. 76:
1581-1584,
[82931] 1985.

[82932] 14422.

[82933] Carson, M. J.; Chadwick, D. L.; Brubaker, C. A.; Cleland, R. S.;
[82934] Landing, B. H.: Thirteen boys with progressive septic granulomatosis. Pediatrics 35:
[82935] 405-412, 1965.

[82936] 14423.

[82937] Clark, R. A.; Malech, H. L.; Gallin, J. I.; Nunoi, H.; Volpp,
[82938] B. D.; Pearson, D. W.; Nauseef, W. M.; Curnutte, J. T.: Genetic variants
[82939] of chronic granulomatous disease: prevalence of deficiencies of two
[82940] cytosolic components of the NADPH oxidase system. New Eng. J. Med. 321:
[82941] 647-652, 1989.

[82942] 14424.

[82943] Condino-Neto, A.; Newburger, P. E.: Interferon-gamma improves

[82944] splicing efficiency of CYBB gene transcripts in an inter-feron-responsive

[82945] variant of chronic granulomatous disease due to a splice site consensus

[82946] region mutation. Blood 95: 3548-3554, 2000.

[82947] 14425.

[82948] Cross, A. R.; Curnutte, J. T.; Rae, J.; Heyworth, P. G.: Hematologically

[82949] important mutations: X-linked chronic granulomatous disease. Blood

[82950] Cells Mol. Dis. 22: 90-95, 1996.

[82951] 14426.

[82952] Curnutte, J. T.: Classification of chronic granulomatous disease. Hemat.

[82953] Oncol. Clin. North Am. 2: 241-252, 1988.

[82954] 14427.

[82955] Curnutte, J. T.; Hopkins, P. J.; Kuhl, W.; Beutler, E.: Study-ing

[82956] X inactivation. (Letter) Lancet 339: 749 only, 1992.

[82957] 14428.

[82958] Curnutte, J. T.; Kipnes, R. S.; Babior, B. M.: Defect in pyri-
dine
[82959] nucleotide dependent superoxide production by a partic-
ulate fraction
[82960] from the granulocytes of patients with chronic granulo-
matous disease. New
[82961] Eng. J. Med. 293: 628-632, 1975.

[82962] 14429.

[82963] Curnutte, J. T.; Whitten, D. M.; Babior, B. M.: Defective su-
peroxide
[82964] production by granulocytes from patients with chronic
granulomatous
[82965] disease. New Eng. J. Med. 290: 593-597, 1974.

[82966] 14430.

[82967] D'Amelio, R.; Bellavite, P.; Bianco, P.; De Sole, P.; Le Moli,

[82968] S.; Lippa, S.; Seminara, R.; Vercelli, B.; Rossi, F.; Rocchi, G.;

[82969] Aiuti, F.: Chronic granulomatous disease in two sisters. J.
Clin.

[82970] Immun. 4: 220-227, 1984.

[82971] 14431.

[82972] Danks, D. M.: Personal Communication. Melbourne, Australia

[82973] 1966.

[82974] 14432.

[82975] de Boer, M.; Bakker, E.; Van Lierde, S.; Roos, D.: Somatic triple

[82976] mosaicism in a carrier of X-linked chronic granulomatous disease. Blood 91:

[82977] 252-257, 1998.

[82978] 14433.

[82979] de Boer, M.; Bolscher, B. G. J. M.; Sijmons, R. H.; Scheffer,

[82980] H.; Weening, R. S.; Roos, D.: Prenatal diagnosis in a family with

[82981] X-linked chronic granulomatous disease with the use of the polymerase

[82982] chain reaction. Prenatal Diag. 12: 773-777, 1992.

[82983] 14434.

[82984] Densen, P.; Wilkinson-Kroovand, S.; Mandell, G. L.; Sulli-

van,

[82985] G.; Oyen, R.; Marsh, W. L.: Kx: its relationship to chronic granulomatous

[82986] disease and genetic linkage with Xg. Blood 58: 34-37, 1981.

[82987] 14435.

[82988] Dilworth, J. A.; Mandell, G. L.: Adults with chronic granu-lomatous

[82989] disease of 'childhood.'. Am. J. Med. 63: 233-243, 1977.

[82990] 14436.

[82991] Dinauer, M.; Parkos, C. A.; Jesaitis, A. J.; Orkin, S. H.: Iden-tification

[82992] of the in vivo protein encoded by the gene mutated in X-linked chronic

[82993] granulomatous disease (X-CGD). (Abstract) Clin. Res. 35: 598A only,

[82994] 1987.

[82995] 14437.

[82996] Dinauer, M. C.; Curnutte, J. T.; Rosen, H. R.; Orkin, S. H.:

[82997] A missense mutation in the neutrophil cytochrome b heavy chain in

[82998] cytochrome-positive X-linked chronic granulomatous disease. J. Clin.

[82999] Invest. 84: 2012-2016, 1989.

[83000] 14438.

[83001] Dinauer, M. C.; Orkin, S. H.: Chronic granulomatous disease:

[83002] molecular genetics. Hemat. Oncol. Clin. North Am. 2: 225-240, 1988.

[83003] 14439.

[83004] Dinauer, M. C.; Orkin, S. H.; Brown, R.; Jesaitis, A. J.; Parkos,

[83005] C. A.: The glycoprotein encoded by the X-linked chronic granulomatous

[83006] disease locus is a component of the neutrophil cytochrome b complex. Nature 327:

[83007] 717-720, 1987.

[83008] 14440.

[83009] Dinauer, M. C.; Orkin, S. H.; Hurst, J. K.; Parkos, C. A.; Jesaitis,

[83010] A. J.; Rosen, H.; Curnutte, J. T.: A missense mutation in the neutrophil

[83011] cytochrome b heavy chain leading to X-linked chronic granulomatous

[83012] disease (X-CGD). (Abstract) Clin. Res. 37: 544A only, 1989.

[83013] 14441.

[83014] Edwards, J. H.: Inheritance of chronic granulomatous disease.

[83015] (Letter) Lancet II: 850-851, 1969.

[83016] 14442.

[83017] Eissa, N. T.; Strauss, A. J.; Haggerty, C. M.; Choo, E. K.; Chu,

[83018] S. C.; Moss, J.: Alternative splicing of human inducible nitric-oxide

[83019] synthase mRNA. Tissue-specific regulation and induction by cytokines. J.

[83020] Biol. Chem. 271: 27184-27187, 1996.

[83021] 14443.

[83022] Ezekowitz, R. A. B.: What is the best way to treat inherited

[83023] disorders? (Editorial) New Eng. J. Med. 344: 926-927, 2001.

[83024] 14444.

[83025] Fikrig, S. M.; Phillipp, J. C. D.; Smithwick, E. M.; Oyen, R.;

[83026] Marsh, W. L.: Chronic granulomatous disease and McLeod syndrome in

[83027] a black child. Pediatrics 66: 403-404, 1980.

[83028] 14445.

[83029] Bassi, M. T.; Bergen, A. A. B.; Bitoun, P.; Charles, S. J.; Clementi,

[83030] M.; Gosselin, R.; Hurst, J.; Lewis, R. A.; Lorenz, B.; Meitinger,

[83031] T.; Messiaen, L.; Ramesar, R. S.; Ballabio, A.; Schiaffino, M. V.

[83032] : Diverse prevalence of large deletions within the OA1 gene in ocular

[83033] albinism type 1 patients from Europe and North America. Hum. Genet. 108:

[83034] 51-54, 2001.

[83035] 14446.

[83036] Oldfors, A.; Kyllerman, M.; Wahlstrom, J.; Darnfors, C.; Henriksson,

[83037] K. G.: X-linked myotubular myopathy: clinical and patho-logical findings

[83038] in a family. Clin. Genet. 36: 5-14, 1989.

[83039] 14447.

[83040] Samson, F.; Mesnard, L.; Heimburger, M.; Hanauer, A.; Chevallay,
[83041] M.; Mercadier, J. J.; Pelissier, J. F.; Feingold, N.; Junien, C.;
[83042] Mandel, J.-L.; Fardeau, M.: Genetic linkage heterogeneity in myotubular
[83043] myopathy. Am. J. Hum. Genet. 57: 120-126, 1995.

[83044] 14448.

[83045] Sarnat, H. B.: Myotubular myopathy: arrest of morphogen-esis of
[83046] myofibres associated with persistence of fetal vimentin and desmin:
[83047] four cases compared with fetal and neonatal muscle. Can. J. Neurol.
[83048] Sci. 17: 109-123, 1990.

[83049] 14449.

[83050] Sarnat, H. B.: Vimentin and desmin in maturing skeletal muscle
[83051] and developmental myopathies. Neurology 42: 1616-1624, 1992.

[83052] 14450.

[83053] Sarnat, H. B.; Roth, S. I.; Jimenez, J. F.: Neonatal myotubu-
lar
[83054] myopathy: neuropathy and failure of postnatal maturation
of fetal
[83055] muscle. Canad. J. Neurol. Sci. 8: 313-320, 1981.

[83056] 14451.

[83057] Sawchak, J. A.; Sher, J. H.; Norman, M. G.; Kula, R. W.;
Shafiq,
[83058] S. A.: Centronuclear myopathy heterogeneity: distinction
of clinical
[83059] types by myosin isoform patterns. Neurology 41:
135-140, 1991.
[83060] 14452.

[83061] Spiro, A. J.; Shy, G. M.; Gonatas, N. K.: Myotubular myopa-
thy. Arch.
[83062] Neurol. 14: 1-14, 1966.

[83063] 14453.

[83064] Starr, J.; Lamont, M.; Iselius, L.; Harvey, J.; Heckmatt, J.:

[83065] A linkage study of a large pedigree with X linked cen-
tronuclear myopathy. J.

[83066] Med. Genet. 27: 281-283, 1990.

[83067] 14454.

[83068] Sutton, I. J.; Winer, J. B.; Norman, A. N.; Liechti-Gallati, S.;

[83069] MacDonald, F.: Limb girdle and facial weakness in female carriers

[83070] of X-linked myotubular myopathy mutations. Neurology 57: 900-902,

[83071] 2001.

[83072] 14455.

[83073] Tanner, S. M.; Laporte, J.; Guiraud-Chaumeil, C.; Liechti-Gallati,

[83074] S.: Confirmation of prenatal diagnosis results of X-linked recessive

[83075] myotubular myopathy by mutational screening, and de-scription of three

[83076] new mutations in the MTM1 gene. Hum. Mutat. 11: 62-68, 1998.

[83077] 14456.

[83078] Tanner, S. M.; Orstavik, K. H.; Kristiansen, M.; Lev, D.; Ler-man-Sagie,

[83079] T.; Sadeh, M.; Liechti-Gallati, S.: Skewed X-inactivation in

a manifesting

[83080] carrier of X-linked myotubular myopathy and in her non-manifesting

[83081] carrier mother. Hum. Genet. 104: 249-253, 1999.

[83082] 14457.

[83083] Taylor, G. S.; Maehama, T.; Dixon, J. E.: Myotubularin, a protein

[83084] tyrosine phosphatase mutated in myotubular myopathy, dephosphorylates

[83085] the lipid second messenger, phosphatidylinositol 3-phosphate. Proc.

[83086] Nat. Acad. Sci. 97: 8910-8915, 2000.

[83087] 14458.

[83088] Thomas, N. S. T.; Sarfarazi, M.; Roberts, K.; Williams, H.; Cole,

[83089] G.; Liechti-Gallati, S.; Harper, P. S.: X-linked myotubular myopathy

[83090] (MTM1): evidence for linkage to Xq28 DNA mark-ers.(Abstract) Cytogenet.

[83091] Cell Genet. 46: 704, 1987.

[83092] 14459.

[83093] Thomas, N. S. T.; Williams, H.; Cole, G.; Roberts, K.; Clarke,
[83094] A.; Liechti-Gallati, S.; Braga, S.; Gerber, A.; Meier, C.; Moser,
[83095] H.; Harper, P. S.: X linked neonatal centronuclear/myotubu-lar myopathy:
[83096] evidence for linkage to Xq28 DNA marker loci. J. Med. Genet. 27:
[83097] 284-287, 1990.

[83098] 14460.

[83099] Torres, C. F.; Griggs, R. C.; Goetz, J. P.: Severe neonatal centronuclear
[83100] myopathy with autosomal dominant inheritance. Arch. Neurol. 42:
[83101] 1011-1014, 1985.

[83102] 14461.

[83103] Van Wijngaarden, G. K.; Fleury, P.; Bethlem, J.; Meijer, A. E.

[83104] F. H.: Familial 'myotubular' myopathy. Neurology 19: 901-908, 1969.
[83105] 14462.

[83106] Vincent, M. C.; Guiraud-Chaumeil, C.; Laporte, J.; Manou-

vrier-Hanu,

[83107] S.; Mandel, J. L.: Extensive germinal mosaicism in a family with

[83108] X linked myotubular myopathy simulates genetic hetero-geneity. J.

[83109] Med. Genet. 35: 241-243, 1998.

[83110] 14463.

[83111] Wallgren-Pettersson, C.; Clarke, A.; Samson, F.; Fardeau, M.;

[83112] Dubowitz, V.; Moser, H.; Grimm, T.; Barohn, R. J.; Barth, P. G.:

[83113] The myotubular myopathies: differential diagnosis of the X linked

[83114] recessive, autosomal dominant, and autosomal recessive forms and present

[83115] state of DNA studies. J. Med. Genet. 32: 673-679, 1995.

[83116] 14464.

[83117] Williams, H.; Cole, G.; Thomas, N.; Brown, C.; Sarfarazi, M.:

[83118] Lethal X-linked myotubular myopathy.(Abstract) J. Med. Genet. 22:

[83119] 138, 1985.

[83120] 14465.

[83121] Williams, H.; Sarfarazi, M.; Roberts, K.; Thomas, N. S. T.; Cole,

[83122] G.; Leichti-Gallati, S.; Harper, P. S.: X-linked myotubular myopathy

[83123] (XLMTM): another muscle disease gene localized to Xq28.(Abstract) Am.

[83124] J. Hum. Genet. 41: A110, 1987.

[83125] 14466.

[83126] Bassi, M. T.; Schiaffino, M. V.; Renieri, A.; De Nigris, F.; Galli,

[83127] L.; Bruttini, M.; Gebbia, M.; Bergen, A. A. B.; Lewis, R. A.; Ballabio,

[83128] A.: Cloning of the gene for ocular albinism type 1 from the distal

[83129] short arm of the X chromosome. Nature Genet. 10: 13-19, 1995.

[83130] 14467.

[83131] Bergen, A. A. B.; Samanns, C.; Schuurman, E. J. M.; van Osch, L.;

[83132] van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker, E.; Gal, A.; van Ommen,

[83133] G. J. B.; Bleeker-Wagemakers, E. M.: Multipoint linkage analysis

[83134] in X-linked ocular albinism of the Nettleship-Falls type. Hum. Genet. 88:

[83135] 162-166, 1991.

[83136] 14468.

[83137] Bergen, A. A. B.; Schuurman, E. J. M.; van den Born, L. I.; Samanns,

[83138] C.; van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker, E.; van Ommen,

[83139] G. J. B.; Gal, A.; Bleeker-Wagemakers, E. M.: Carrier detection in

[83140] X-linked ocular albinism of the Nettleship-Falls type by DNA analysis. Clin.

[83141] Genet. 41: 135-138, 1992.

[83142] 14469.

[83143] Bergen, A. A. B.; Zijp, P.; Schuurman, E. J. M.; Bleeker-Wagemakers,

[83144] E. M.; Apkarian, P.; van Ommen, G.-J. B.: Refinement of the localization

[83145] of the X-linked ocular albinism gene. Genomics 16: 272-273, 1993.

[83146] 14470.

[83147] Bouloux, P.-M. G.; Kirk, J.; Munroe, P.; Duke, V.; Meindl, A.;

[83148] Hilson, A.; Grant, D.; Carter, N.; Betts, D.; Meitinger, T.; Besser,

[83149] G. M.: Deletion analysis maps ocular albinism proximal to the steroid

[83150] sulphatase locus. Clin. Genet. 43: 169-173, 1993.

[83151] 14471.

[83152] Charles, S. J.; Green, J. S.; Moore, A. T.; Barton, D. E.; Yates,

[83153] J. R. W.: Genetic mapping of X-linked ocular albinism: linkage analysis

[83154] in a large Newfoundland kindred. Genomics 16: 259-261, 1993.

[83155] 14472.

[83156] Charles, S. J.; Moore, A. T.; Yates, J. R. W.: Genetic map-ping

[83157] of X linked ocular albinism: linkage analysis in British families. J.

[83158] Med. Genet. 29: 552-554, 1992.

[83159] 14473.

[83160] Cooper, D. N.; Krawczak, M.: Mechanisms of insertional mutagenesis

[83161] in human genes causing genetic disease. Hum. Genet. 87: 409-415,

[83162] 1991.

[83163] 14474.

[83164] Creel, D.; O'Donnell, F. E., Jr.; Witkop, C. J., Jr.: Visual

[83165] system anomalies in human ocular albinos. Science 201: 931-933,

[83166] 1978.

[83167] 14475.

[83168] d'Addio, M.; Pizzigoni, A.; Bassi, M. T.; Baschirotto, C.; Valetti,

[83169] C.; Incerti, B.; Clementi, M.; De Luca, M.; Ballabio, A.; Schi-affino,

[83170] M. V.: Defective intracellular transport and processing of OA1 is

[83171] a major cause of ocular albinism type 1. Hum. Molec. Genet. 9: 3011-3018,

[83172] 2000.

[83173] 14476.

[83174] Engelhard, C. F.: Eine Familie mit hereditaerem Nystag-
mus. Z.
[83175] Ges. Neurol. Psychiat. 28: 319-338, 1915.

[83176] 14477.

[83177] Fialkow, P. J.; Giblett, E. R.; Motulsky, A. G.: Measurable
linkage
[83178] between ocular albinism and Xg. Am. J. Hum. Genet. 19:
63-69, 1967.
[83179] 14478.

[83180] Francois, J.; Deweer, J. P.: Albinisme oculaire lie au sexe et

[83181] alterations caracteristiques du fond d'oeil chez les
femmes heterozygotes. Ophthalmologica 126:
[83182] 209-221, 1953.

[83183] 14479.

[83184] Gillespie, F. D.: Ocular albinism with report of a family
with
[83185] female carriers. Arch. Ophthal. 66: 774-777, 1961.

[83186] 14480.

[83187] Incerti, B.; Cortese, K.; Pizzigoni, A.; Surace, E. M.; Varani,

[83188] S.; Coppola, M.; Jeffery, G.; Seeliger, M.; Jaissle, G.; Bennett,

[83189] D. C.; Marigo, V.; Schiaffino, M. V.; Tacchetti, C.; Ballabio, A.

[83190] : Oa1 knock-out: new insights on the pathogenesis of ocular albinism

[83191] type 1. Hum. Molec. Genet. 9: 2781-2788, 2000.

[83192] 14481.

[83193] Jaeger, C.; Jay, B.: X-linked ocular albinism: a family containing

[83194] a manifesting heterozygote, and an affected male married to a female

[83195] with autosomal recessive ocular albinism. Hum. Genet. 56: 299-304,

[83196] 1981.

[83197] 14482.

[83198] Johnson, G. J.; Gillan, J. G.; Pearce, W. G.: Ocular albinism

[83199] in Newfoundland. Can. J. Ophthal. 6: 237-248, 1971.

[83200] 14483.

[83201] Kidd, J. R.; Castiglione, C. M.; Davies, K. E.; Pakstis, A. J.;

[83202] Gusella, J.; Sparkes, R. S.; Pearson, P.; Willard, H.; Kidd, K. K.

[83203] : Mapping the locus for X-linked ocular albinism (OA). (Abstract) Am.

[83204] J. Hum. Genet. 37: A161 only, 1985.

[83205] 14484.

[83206] Krawczak, M.; Reiss, J.; Cooper, D. N.: The mutational spectrum

[83207] of single base-pair substitutions in mRNA splice junctions of human

[83208] genes: causes and consequences. Hum. Genet. 90: 41-54, 1992.

[83209] 14485.

[83210] Lein, J. N.; Stewart, C. T.; Moll, F. C.: Sex-linked hereditary

[83211] nystagmus. Pediatrics 18: 214-217, 1956.

[83212] 14486.

[83213] Lyon, M. F.: Sex chromatin and gene action in the mammalian X-chromosome. Am.

[83214] J. Hum. Genet. 14: 135-148, 1962.

[83215] 14487.

[83216] Meindl, A.; Hosenfeld, D.; Bruckl, W.; Schuffenhauer, S.; Jenderny,
[83217] J.; Bacskulin, A.; Oppermann, H.-C.; Swensson, O.; Bouloux, P.; Meitinger,
[83218] T.: Analysis of a terminal Xp22.3 deletion in a patient with six
[83219] monogenic disorders: implications for the mapping of X linked ocular
[83220] albinism. J. Med. Genet. 30: 838-842, 1993.

[83221] 14488.

[83222] Negrelli, B. C.: L'albinisme oculaire lie au sexe dans le cadre
[83223] du depistage des heterozygotes en ophtalmologie. J. Genet. Hum. 8:
[83224] 108 only, 1959.

[83225] 14489.

[83226] Newton, J. M.; Orlow, S. J.; Barsh, G. S.: Isolation and char-acterization
[83227] of a mouse homolog of the X-linked ocular albinism (OA1) gene. Genomics 37:
[83228] 219-225, 1996.

[83229] 14490.

[83230] O'Donnell, F. E., Jr.; Green, W. R.; Fleishman, J. A.; Ham-
brick,
[83231] G. W.: X-linked ocular albinism in Blacks: ocular albinism
cum pigmento. Arch.
[83232] Ophthal. 96: 1189-1192, 1978.

[83233] 14491.

[83234] O'Donnell, F. E., Jr.; Hambrick, G. W., Jr.; Green, W. R.; Iliff,
[83235] W. J.; Stone, D. L.: X-linked ocular albinism: an oculocuta-
neous
[83236] macromelanosomal disorder. Arch. Ophthal. 94:
1883-1892, 1976.
[83237] 14492.

[83238] Oetting, W. S.: New insights into ocular albinism type 1
(OA1):
[83239] mutations and polymorphisms of the OA1 gene. Hum.
Mutat. 19: 85-92,
[83240] 2002.

[83241] 14493.

[83242] Pearce, W. G.; Johnson, G. J.; Gillan, J. G.: Nystagmus in a

[83243] female carrier of ocular albinism. J. Med. Genet. 9:
126-128, 1972.
[83244] 14494.
[83245] Pearce, W. G.; Johnson, G. J.; Sanger, R.: Ocular albinism
and
[83246] Xg. (Letter) Lancet I: 1072 only, 1971.
[83247] 14495.
[83248] Pearce, W. G.; Sanger, R.: X mapping in man: evidence
against
[83249] direct measurable linkage between ocular albinism and
deutan colour
[83250] blindness. J. Med. Genet. 13: 319 only, 1976.
[83251] 14496.
[83252] McCabe, E. R. B.: Human glycerol kinase deficiency: an in-
born
[83253] error of compartment metabolism. Biochem. Med. 30:
215-230, 1983.
[83254] 14497.
[83255] Basler, E.; Grompe, M.; Parenti, G.; Yates, J.; Ballabio, A.:
[83256] Identification of point mutations in the steroid sulfatase
gene of

[83257] three patients with X-linked ichthyosis. Am. J. Hum. Genet. 50:
[83258] 483-491, 1992.

[83259] 14498.

[83260] Bernatowicz, L. F.; Li, X.-M.; Carrozzo, R.; Ballabio, A.; Mohandas,
[83261] T.; Yen, P. H.; Shapiro, L. J.: Sequence analysis of a partial deletion
[83262] of the human steroid sulfatase gene reveals 3 bp of homology at deletion
[83263] breakpoints. Genomics 13: 892-893, 1992.

[83264] 14499.

[83265] Bonifas, J. M.; Morley, B. J.; Oakey, R. E.; Kan, Y. W.; Epstein,
[83266] E. H., Jr.: Cloning of a cDNA for steroid sulfatase: frequent occurrence
[83267] of gene deletions in patients with recessive X chromosome-linked ichthyosis. Proc.
[83268] Nat. Acad. Sci. 84: 9248-9251, 1987.

[83269] 14500.

[83270] Buckle, V. J.; Edwards, J. H.; Evans, E. P.; Jonasson, J. A.;

[83271] Lyon, M. F.; Peters, J.; Searle, A. G.: Comparative maps of human
[83272] and mouse X chromosomes. (Abstract) Cytogenet. Cell Genet. 40: 594-595,
[83273] 1985.

[83274] 14501.

[83275] Burns, G.: On the identity of arylsulphatase C and steroid sulphatase. Hum.
[83276] Genet. 65: 189, 1983.

[83277] 14502.

[83278] Chance, P. F.; Gartler, S. M.: Evidence for a dosage effect at
[83279] the X-linked steroid sulfatase locus in human tissues. Am. J. Hum.
[83280] Genet. 35: 234-240, 1983.

[83281] 14503.

[83282] Chang, P. L.; Lafferty, K. I.; Rosa, N. E.; Davidson, R. G.:

[83283] Arylsulfatase-C isozymes in human tissues. (Abstract) Am. J. Hum.
[83284] Genet. 33: 38A, 1981.

[83285] 14504.

[83286] Cockayne, E. A.: Inherited Abnormalities of the Skin and Its

[83287] Appendages. London: Oxford Univ. Press (pub.) 1933. Pp. 213 only.

[83288] 14505.

[83289] Conary, J. T.; Lorkowski, G.; Schmidt, B.; Pohlmann, R.; Nagel,

[83290] G.; Meyer, H. E.; Krentler, C.; Cully, J.; Hasilik, A.; von Figura,

[83291] K.: Genetic heterogeneity of steroid sulfatase deficiency revealed

[83292] with cDNA for human steroid sulfatase. Biochem. Biophys. Res. Commun. 144:

[83293] 1010-1017, 1987.

[83294] 14506.

[83295] Cooper, D. W.; McAllan, B. M.; Donald, J. A.; Dawson, G.; Dobrovic,

[83296] A.; Marshall Graves, J. A.: Steroid sulphatase is not de-tected on

[83297] the X chromosome of Australian marsupials. (Abstract) Cytogenet.

[83298] Cell Genet. 37: 439, 1984.

[83299] 14507.

[83300] Craig, I. W.; Tolley, E.: Steroid sulphatase and the conser-
vation
[83301] of mammalian X chromosomes. Trends Genet. 2:
201-204, 1986.

[83302] 14508.

[83303] Csorsz, K.: Ichthyosis (X-linked). Mschr. Unfallheilk. Med.
2:
[83304] 180, 1928. Note: Alternate: Haut. Geschlechtskr. 26: 463,
1928...

[83305] 14509.

[83306] Cuevas-Covarrubias, S. A.; Kofman-Alfaro, S.; Orozco
Orozco, E.;
[83307] Diaz-Zagoya, J. C.: The biochemical identification of car-
rier state
[83308] in mothers of sporadic cases of X-linked recessive
ichthyosis. Genet.
[83309] Counsel. 6: 103-107, 1995.

[83310] 14510.

[83311] Czeizel, A.: A historical evaluation of the doctrine of here-

dodegeneration. Orv.

[83312] Hetil. 120: 722, and 840, and 963, 1979.

[83313] 14511.

[83314] DeUnamuno, P.; Martin-Pascual, A.; Garcia-Perez, A.: X-linked

[83315] ichthyosis. Brit. J. Derm. 97: 53-58, 1977.

[83316] 14512.

[83317] Eicher, E. M.: Personal Communication. Bar Harbor, Me. 1974.

[83318] 14513.

[83319] Elias, P. M.; Williams, M. L.; Maloney, M. E.; Bonifas, J. A.;

[83320] Brown, B. E.; Grayson, S.; Epstein, E. H., Jr.: Stratum corneum lipids

[83321] in disorders of cornification: steroid sulfatase and choles-terol sulfate

[83322] in normal desquamation and the pathogenesis of reces-sive X-linked

[83323] ichthyosis. J. Clin. Invest. 74: 1414-1421, 1984.

[83324] 14514.

[83325] Epstein, E. H., Jr.; Bonifas, J. M.: Recessive X-linked

ichthyosis:
[83326] lack of immunologically detectable steroid sulfatase enzyme protein. Hum.
[83327] Genet. 71: 201-205, 1985.

[83328] 14515.

[83329] Epstein, E. H., Jr.; Krauss, R. M.; Shackleton, C. H. L.: X-linked
[83330] ichthyosis: increased blood cholesterol sulfate and electrophoretic
[83331] mobility of low-density lipoprotein. Science 214: 659-660, 1981.
[83332] 14516.

[83333] Epstein, E. H., Jr.; Williams, M. L.; Elias, P. M.: Steroid sulfatase,
[83334] X-linked ichthyosis, and stratum corneum cell cohesion. Arch. Derm. 117:
[83335] 761-763, 1981.

[83336] 14517.

[83337] Filippi, G.; Meera Khan, P.: Linkage studies on X-linked ichthyosis
[83338] in Sardinia. Am. J. Hum. Genet. 20: 564-569, 1968.

[83339] 14518.

[83340] France, J. T.; Downey, J. A.: A study of arylsulfatase activity

[83341] in children born of pregnancies affected with placental sulfatase

[83342] deficiency. Biochem. Med. 10: 167-174, 1974.

[83343] 14519.

[83344] France, J. T.; Liggins, G. C.: Placental sulfatase deficiency. J.

[83345] Clin. Endocr. 29: 138-141, 1969.

[83346] 14520.

[83347] France, J. T.; Seddons, R. J.; Liggins, G. C.: A study of a pregnancy

[83348] with low estrogen production due to placental sulfatase deficiency. J.

[83349] Clin. Endocr. 36: 1-9, 1973.

[83350] 14521.

[83351] Freiberg, R. A.; Choate, K. A.; Deng, H.; Alperin, E. S.; Shapiro,

[83352] L. J.; Khavari, P. A.: A model of corrective gene transfer in X-linked

[83353] ichthyosis. Hum. Molec. Genet. 6: 927-933, 1997.

[83354] 14522.

[83355] Gant, N. F.; Milewich, L.; Calvert, M. E.; MacDonald, P. C.:

[83356] Steroid sulfatase activity in human fetal membranes. J. Clin. Endocr. 45:

[83357] 965-972, 1977.

[83358] 14523.

[83359] Garcia Perez, A.; Crespo, M.: X-linked ichthyosis associated

[83360] with hypertrophic pyloric stenosis in three brothers. Clin. Exp.

[83361] Derm. 6: 159-161, 1981.

[83362] 14524.

[83363] Gartler, S. M.; Andina, R. J.: Mammalian X-chromosome inactivation. Adv.

[83364] Hum. Genet. 7: 99-140, 1976.

[83365] 14525.

[83366] Gartler, S. M.; Rivest, M.: Evidence for X-linkage of steroid

[83367] sulfatase in the mouse: steroid sulfatase levels in oocytes

of XX
[83368] and XO mice. Genetics 103: 137-141, 1983.

[83369] 14526.

[83370] Gillard, E. F.; Affara, N. A.; Yates, J. R. W.; Goudie, D. R.;

[83371] Cooke, A.; Lambert, J.; Aitken, D. A.; Ferguson-Smith, M. A.: Deletion
[83372] of a DNA probe DXS237 (GMGX9) in twelve of fifteen males with X-linked
[83373] ichthyosis (steroid sulphatase deficiency). (Abstract) Cyto-genet.
[83374] Cell Genet. 46: 620, 1987.

[83375] 14527.

[83376] Gillard, E. F.; Affara, N. A.; Yates, J. R. W.; Goudie, D. R.;

[83377] Lambert, J.; Aitken, D. A.; Ferguson-Smith, M. A.: Deletion of a
[83378] DNA sequence in eight of nine families with X-linked ichthyosis (steroid
[83379] sulphatase deficiency). Nucleic Acids Res. 15: 3977-3985, 1987.

[83380] 14528.

[83381] Gladstein, K.; Shapiro, L. J.; Spence, M. A.: Estimating sex

[83382] ratio biases in X-linked disorders: is there an excess of males in
[83383] families with X-linked ichthyosis?. Am. J. Hum. Genet. 31: 741-746,
[83384] 1979.

[83385] 14529.

[83386] Gohlke, B. C.; Haug, K.; Fukami, M.; Friedl, W.; Noeker, M.; Rappold,
[83387] G. A.; Haverkamp, F.: Interstitial deletion in Xp22.3 is as-sociated
[83388] with X linked ichthyosis, mental retardation, and epilepsy. J. Med.
[83389] Genet. 37: 600-602, 2000.

[83390] 14530.

[83391] Beresford, R. G.; Tatlidil, C.; Riddell, D. C.; Welch, J. P.;

[83392] Ludman, M. D.; Neumann, P. E.; Greer, W. L.: Absence of fragile X
[83393] syndrome in Nova Scotia. J. Med. Genet. 37: 77-79, 2000.

[83394] 14531.

[83395] Blomquist, H. K.; Bohman, M.; Edvinsson, S. O.; Gillberg, C.;

[83396] Gustavson, K.-H.; Holmgren, G.; Wahlstrom, J.: Frequency of the fragile
[83397] X syndrome in infantile autism: a Swedish multicenter study. Clin.
[83398] Genet. 27: 113-117, 1985.

[83399] 14532.

[83400] Blomquist, H. K.; Gustavson, K.-H.; Holmgren, G.; Nor-
denson, I.;
[83401] Sweins, A.: Fragile site X chromosomes and X-linked mental retardation
[83402] in severely retarded boys in a northern Swedish county: a prevalence
[83403] study. Clin. Genet. 21: 209-214, 1982.

[83404] 14533.

[83405] Bontekoe, C. J. M.; Bakker, C. E.; Nieuwenhuizen, I. M.; van der
[83406] Linde, H.; Lans, H.; de Lange, D.; Hirst, M. C.; Oostra, B. A.: Instability
[83407] of a (CGG)98 repeat in the Fmr1 promoter. Hum. Molec. Genet. 10:
[83408] 1693-1699, 2001.

[83409] 14534.

[83410] Bowen, P.; Biederman, B.; Swallow, K. A.: The X-linked syndrome

[83411] of macro-orchidism and mental retardation: further observations. Am.

[83412] J. Med. Genet. 2: 409-414, 1978.

[83413] 14535.

[83414] Brookwell, R.; Turner, G.: High resolution banding and the locus

[83415] of the Xq fragile site. Hum. Genet. 63: 77, 1983.

[83416] 14536.

[83417] Brown, V.; Jin, P.; Ceman, S.; Darnell, J. C.; O'Donnell, W. T.;

[83418] Tenenbaum, S. A.; Jin, X.; Feng, Y.; Wilkinson, K. D.; Keene, J. D.;

[83419] Darnell, R. B.; Warren, S. T.: Microarray identification of FMRP-associated

[83420] brain mRNAs and altered mRNA translational profiles in fragile X syndrome. Cell 107:

[83421] 477-487, 2001.

[83422] 14537.

[83423] Brown, W. T.; Gross, A.; Chan, C.; Jenkins, E. C.; Mandel, J.

[83424] L.; Oberle, I.; Arveiler, B.; Novelli, G.; Thibodeau, S.; Hagerman,

[83425] R.; and 18 others: Multilocus analysis of the fragile X syndrome. Hum.

[83426] Genet. 78: 201-205, 1988.

[83427] 14538.

[83428] Brown, W. T.; Gross, A. C.; Chan, C. B.; Jenkins, E. C.: Genetic

[83429] linkage heterogeneity in the fragile X syndrome. Hum. Genet. 71:

[83430] 11-18, 1985.

[83431] 14539.

[83432] Bundey, S.: Julia Bell, MRCS LRCP FRCP (1879-1979): Steamboat

[83433] lady, statistician and geneticist. J. Med. Bio. 4: 8-13, 1996.

[83434] 14540.

[83435] Camerino, G.; Mattei, M. G.; Mattei, J. F.; Jaye, M.; Mandel,

[83436] J. L.: Close linkage of fragile X-linked mental retardation syndrome

[83437] to haemophilia B and transmission through a normal male. Nature 306:

[83438] 701-707, 1983.

[83439] 14541.

[83440] Cantu, J. M.; Scaglia, H. E.; Gonzalez-Diddi, M.; Hernan-dez-Jauregui,

[83441] P.; Morato, T.; Moreno, M. E.; Giner, J.; Alcantar, A.; Her-rera, D.;

[83442] Perez-Palacios, G.: Inherited congenital normofunctional testicular

[83443] hyperplasia and mental deficiency. Hum. Genet. 41: 331-339, 1978.

[83444] 14542.

[83445] Cantu, J. M.; Scaglia, H. E.; Medina, M.; Gonzalez-Diddi, M.;

[83446] Morato, T.; Moreno, M. E.; Perez-Palacios, G.: Inherited congenital

[83447] normofunctional testicular hyperplasia and mental defi-ciency. Hum.

[83448] Genet. 33: 23-33, 1976.

[83449] 14543.

[83450] Carmi, R.; Meryash, D. L.; Wood, J.; Gerald, P. S.: Fragile-X
[83451] syndrome ascertained by the presence of macro-or-
chidism in a 5-month-old
[83452] infant. Pediatrics 74: 883-886, 1984.
[83453] 14544.
[83454] Cavalli-Sforza, L.; Menozzi, P.; Piazza, A.: The History and
[83455] Geography of Human Genes. Princeton: Princeton Univ.
Press , 1994.
[83456] 14545.
[83457] Chakravarti, A.: Fragile X founder effect? Nature Genet. 1:
[83458] 237-238, 1992.
[83459] 14546.
[83460] Chiurazzi, P.; Pomponi, M. G.; Pietrobono, R.; Bakker, C.
E.;
[83461] Neri, G.; Oostra, B. A.: Synergistic effect of histone hyper-
acetylation
[83462] and DNA demethylation in the reactivation of the FMR1
gene. Hum.
[83463] Molec. Genet. 8: 2317-2323, 1999.
[83464] 14547.

[83465] Chiurazzi, P.; Pomponi, M. G.; Willemsen, R.; Oostra, B. A.; Neri,
[83466] G.: In vitro reactivation of the FMR1 gene involved in fragile X
[83467] syndrome. Hum. Molec. Genet. 7: 109-113, 1998.

[83468] 14548.

[83469] Choo, K. H.; George, D.; Fillby, G.; Halliday, J. L.; Leversha,
[83470] M.; Webb, G.; Danks, D. M.: Linkage analysis of X-linked mental retardation
[83471] with and without fragile-X using factor IX gene probe. (Letter) Lancet II:
[83472] 349, 1984.

[83473] 14549.

[83474] Crawford, D. C.; Meadows, K. L.; Newman, J. L.; Taft, L. F.; Scott,
[83475] E.; Leslie, M.; Shubek, L.; Holmgreen, P.; Yeargin-Allsopp, M.; Boyle,
[83476] C.; Sherman, S. L.: Prevalence of the fragile X syndrome in African-Americans. Am.
[83477] J. Med. Genet. 110: 226-233, 2002.

[83478] 14550.

[83479] Crawford, D. C.; Wilson, B.; Sherman, S. L.: Factors involved
[83480] in the initial mutation of the fragile X CGG repeat as determined
[83481] by sperm small pool PCR. Hum. Molec. Genet. 9: 2909-2918, 2000.
[83482] 14551.

[83483] Daker, M. G.; Chidiac, P.; Fear, C. N.; Berry, A. L.: Fragile
[83484] X in a normal male: a cautionary tale. Lancet I: 780, 1981.

[83485] 14552.

[83486] Dar, H.; Schaap, T.; Bait-Or, H.; Borochowitz, Z.; Gelman-Kohan,
[83487] Z.; Chemke, T.; Chaki, R.; Cohen, H.; Falik-Borenstein, Z.; Chemke,
[83488] J.: Ethnic distribution of the fragile X syndrome in Israel: evidence
[83489] of founder chromosomes(?). Israel J. Med. Sci. 31: 323-325, 1995.
[83490] 14553.

[83491] Darnell, J. C.; Jensen, K. B.; Jin, P.; Brown, V.; Warren, S.
[83492] T.; Darnell, R. B.: Fragile X mental retardation protein tar-

gets
[83493] G quartet mRNAs important for neuronal function. Cell 107: 489-499,
[83494] 2001.

[83495] 14554.

[83496] Davids, J. R.; Hagerman, R. J.; Eilert, R. E.: Orthopaedic as-
pects
[83497] of fragile-X syndrome. J. Bone Joint Surg. 72A: 889-896,
1990.
[83498] 14555.

[83499] Crawford, D. C.; Meadows, K. L.; Newman, J. L.; Taft, L. F.;
Pettay,
[83500] D. L.; Gold, L. B.; Hersey, S. J.; Hinkle, E. F.; Stanfield, M.
L.;
[83501] Holmgreen, P.; Yeargin-Allsopp, M.; Boyle, C.; Sherman, S.
L.: Prevalence
[83502] and phenotype consequence of FRAXA and FRAXE alleles
in a large, ethnically
[83503] diverse, special education-needs population. Am. J. Hum.
Genet. 64:
[83504] 495-507, 1999.

[83505] 14556.

[83506] Gaugitsch, H. W.; Prieschl, E. E.; Kalthoff, F.; Huber, N. E.;

[83507] Baumruker, T.: A novel transiently expressed, integral membrane protein

[83508] linked to cell activation: molecular cloning via the rapid degradation

[83509] signal AUUUA. J. Biol. Chem 267: 11267-11273, 1992.

[83510] 14557.

[83511] Kanai, Y.; Segawa, H.; Miyamoto, K.; Uchino, H.; Takeda, H.; Endou,

[83512] H.: Expression cloning and characterization of a trans- porter for

[83513] large neutral amino acids activated by the heavy chain of 4F2 antigen

[83514] (CD98). J. Biol. Chem. 273: 23629-23632, 1998.

[83515] 14558.

[83516] Amagai, M.; Wang, Y.; Minoshima, S.; Kawamura, K.; Green, K. J.;

[83517] Nishikawa, T.; Shimizu, N.: Assignment of the human genes for desmocollin

[83518] 3 (DSC3) and desmocollin 4 (DSC4) to chromosome 18q12. Genomics 25:

[83519] 330-332, 1995.

[83520] 14559.

[83521] Kawamura, K.; Watanabe, K.; Suzuki, T.; Yamakawa, T.; Kamiyama,
[83522] T.; Nakagawa, H.; Tsurufuji, S.: cDNA cloning and expression of a
[83523] novel human desmocollin. J. Biol. Chem. 269: 26295-26302, 1994.

[83524] 14560.

[83525] King, I. A.; Sullivan, K. H.; Bennett, R., Jr.; Buxton, R. S.:
[83526] The desmocollins of human foreskin epidermis: identification and chromosomal
[83527] assignment of a third gene and expression patterns of the three isoforms. J.
[83528] Invest. Derm. 105: 314-321, 1995.

[83529] 14561.

[83530] Chan, E. K. L.; Takano, S.; Andrade, L. E. C.; Hamel, J. C.; Matera,
[83531] A. G.: Structure, expression and chromosomal localization of human
[83532] p80-coilin gene. Nucleic Acids Res. 22: 4462-4469, 1994.

[83533] 14562.

[83534] Warburg, O.: On the origin of cancer cells. Science 123: 309-314,

[83535] 1956.

[83536] 14563.

[83537] Dubrovsky, A. L.; Taratuto, A. L.; Sevlever, G.; Schultz, M.;

[83538] Pegoraro, E.; Hoop, R. C.; Hoffman, E. P.: Duchenne muscular dystrophy

[83539] and myotonic dystrophy in the same patient. Am. J. Med. Genet. 55:

[83540] 342-348, 1995.

[83541] 14564.

[83542] Finlay, A. Y.; Kingston, H. M.; Holt, P. J. A.: Chronic granulomatous

[83543] disease carrier geno-dermatosis (CGDCGD). Clin. Genet. 23: 276-280,

[83544] 1983.

[83545] 14565.

[83546] Goodfellow, P. N.; Tippett, P.: A human quantitative polymorphism

[83547] related to Xg blood groups. Nature 289: 404-409, 1981.

[83548] 14566.

[83549] Davies, K. E.; Mattei, M. G.; Mattei, J. F.; Veenema, H.; McGlade,

[83550] S.; Harper, K.; Tommerup, N.; Nielsen, K. B.; Mikkelsen, M.; Beighton,

[83551] P.; Drayna, D.; White, R.; Pembrey, M. E.: Linkage studies of X-linked

[83552] mental retardation: high frequency of recombination in the telomeric

[83553] region of the human X chromosome (fragile site/linkage/recombination/X

[83554] chromosome). Hum. Genet. 70: 249-255, 1985.

[83555] 14567.

[83556] Cazzola, M.; Bergamaschi, G.: X-linked Wiskott-Aldrich syndrome

[83557] in a girl. (Letter) New Eng. J. Med. 338: 1850 only, 1998.

[83558] 14568.

[83559] Emery, A. E. H.: Duchenne Muscular Dystrophy. Oxford, UK: Oxford

[83560] University Press (pub.) (2nd ed.): 1993.

[83561] 14569.

[83562] Fabb, S. A.; Wells, D. J.; Serpente, P.; Dickson, G.: Adeno-associated

[83563] virus vector gene transfer and sarcolemmal expression of a 144 kDa

[83564] micro-dystrophin effectively restores the dystrophin-associated protein

[83565] complex and inhibits myofibre degeneration in nude/mdx mice. Hum.

[83566] Molec. Genet. 11: 733-741, 2002.

[83567] 14570.

[83568] Feener, C. A.; Boyce, F. M.; Kunkel, L. M.: Rapid detection of

[83569] CA polymorphisms in cloned DNA: application to the 5-prime region

[83570] of the dystrophin gene. Am. J. Hum. Genet. 48: 621-627, 1991.

[83571] 14571.

[83572] Ferlini, A.; Galie, N.; Merlini, L.; Sewry, C.; Branzi, A.; Muntoni,

[83573] F.: A novel Alu-like element rearranged in the dystrophin gene causes

[83574] a splicing mutation in a family with X-linked dilated car-
diomyopathy. Am.
[83575] J. Hum. Genet. 63: 436-446, 1998.

[83576] 14572.

[83577] Finnegan, D. J.: Eukaryotic transposable elements and
genome
[83578] evolution. Trends Genet. 5: 103-107, 1989.

[83579] 14573.

[83580] Forrest, S. M.; Cross, G. S.; Speer, A.; Gardner-Medwin, D.;
Burn,
[83581] J.; Davies, K. E.: Preferential deletion of exons in
Duchenne and
[83582] Becker muscular dystrophies. Nature 329: 638-640, 1987.

[83583] 14574.

[83584] Francke, U.; Ochs, H. D.; de Martinville, B.; Giacalone, J.;
Lindgren,
[83585] V.; Disteche, C.; Pagon, R. A.; Hofker, M. H.; van Ommen,
G.-J. B.;
[83586] Pearson, P. L.; Wedgwood, R. J.: Minor Xp21 chromosome
deletion in
[83587] a male associated with expression of Duchenne muscular

dystrophy,
[83588] chronic granulomatous disease, retinitis pigmentosa, and McLeod syndrome. Am.
[83589] J. Hum. Genet. 37: 250-267, 1985.

[83590] 14575.

[83591] Furst, D.; Nave, R.; Osborn, M.; Weber, K.; Bardosi, A.; Archidiacono,
[83592] N.; Ferro, M.; Romano, V.; Romeo, G.: Nebulin and titin expression
[83593] in Duchenne muscular dystrophy appears normal. FEBS Lett. 224: 49-53,
[83594] 1987.

[83595] 14576.

[83596] Giacalone, J. P.; Francke, U.: Common sequence motifs at the
[83597] rearrangement sites of a constitutional X/autosome translocation and
[83598] associated deletion. Am. J. Hum. Genet. 50: 725-741, 1992.
[83599] 14577.

[83600] Gillard, E. F.; Chamberlain, J. S.; Murphy, E. G.; Duff, C. L.;

[83601] Smith, B.; Burghes, A. H. M.; Thompson, M. W.; Sutherland, J.; Oss,
[83602] I.; Bodrug, S. E.; Klamut, H. J.; Ray, P. N.; Worton, R. G.: Molecular
[83603] and phenotypic analysis of patients with deletions within the deletion-rich
[83604] region of the Duchenne muscular dystrophy (DMD) gene. Am. J. Hum.
[83605] Genet. 45: 507-520, 1989.

[83606] 14578.

[83607] Ginjaar, I. B.; Kneppers, A. L. J.; Meulen, J.-D. M.; Anderson,
[83608] L. V. B.; Bremmer-Bout, M.; van Deutekom, J. C. T.; Weegenaar, J.;
[83609] den Dunnen, J. T.; Bakker, E.: Dystrophin nonsense mutation induces
[83610] different levels of exon 29 skipping and leads to variable phenotypes
[83611] within one BMD family. Europ. J. Hum. Genet. 8: 793-796, 2000.

[83612] 14579.

[83613] Greenberg, D. S.; Sunada, Y.; Campbell, K. P.; Yaffe, D.;

Nudel,
[83614] U.: Exogenous Dp71 restores the levels of dystrophin associated proteins
[83615] but does not alleviate muscle damage in mdx mice. Nature Genet. 8:
[83616] 340-344, 1994.

[83617] 14580.

[83618] Gussoni, E.; Soneoka, Y.; Strickland, C. D.; Buzney, E. A.; Khan,
[83619] M. K.; Flint, A. F.; Kunkel, L. M.; Mulligan, R. C.: Dystrophin expression
[83620] in the mdx mouse restored by stem cell transplantation. Nature 401:
[83621] 390-394, 1999.

[83622] 14581.

[83623] Hagiwara, Y.; Mizuno, Y.; Takemitsu, M.; Matsuzaki, T.; Nonaka,
[83624] I.; Ozawa, E.: Dystrophin-positive muscle fibers following C2 myoblast
[83625] transplantation into mdx nude mice. Acta Neuropath. 90:
592-600,
[83626] 1995.

[83627] 14582.

[83628] Hagiwara, Y.; Nishio, H.; Kitoh, Y.; Takeshima, Y.; Narita, N.;

[83629] Wada, H.; Yokoyama, M.; Nakamura, H.; Matsuo, M.: A novel point mutation

[83630] (G(-1) to T) in a 5-prime splice donor site of intron 13 of the dystrophin

[83631] gene results in exon skipping and is responsible for Becker muscular

[83632] dystrophy. Am. J. Hum. Genet. 54: 53-61, 1994.

[83633] 14583.

[83634] Hammonds, R. G., Jr.: Protein sequence of DMD gene is related

[83635] to actin-binding domain of alpha-actinin. (Letter) Cell 51: 1, 1987.

[83636] 14584.

[83637] Harper, S. Q.; Hauser, M. A.; DelloRusso, C.; Duan, D.; Crawford,

[83638] R. W.; Phelps, S. F.; Harper, H. A.; Robinson, A. S.; Engel-hardt,

[83639] J. F.; Brooks, S. V.; Chamberlain, J. S.: Modular flexibility of

[83640] dystrophin: implications for gene therapy of Duchenne muscular dystrophy. Nature

[83641] Med. 8: 253-261, 2002.

[83642] 14585.

[83643] Hart, K. A.; Hodgson, S.; Walker, A.; Cole, C. G.; Johnson, L.;

[83644] Dubowitz, V.; Bobrow, M.: DNA deletions in mild and se-vere Becker

[83645] muscular dystrophy. Hum. Genet. 75: 281-285, 1987.

[83646] 14586.

[83647] Hodgson, S. V.; Abbs, S.; Clark, S.; Manzur, A.; Heckmatt, J.

[83648] Z. H.; Dubowitz, V.; Bobrow, M.: Correlation of clinical and deletion

[83649] data in Duchenne and Becker muscular dystrophy, with special reference

[83650] to mental ability. Neuromusc. Disord. 2: 269-276, 1992.

[83651] 14587.

[83652] Hoffman, E. P.; Brown, R. H., Jr.; Kunkel, L. M.: The protein

[83653] product of the Duchenne muscular dystrophy locus. Cell 51: 919-928,

[83654] 1987.

[83655] 14588.

[83656] Hoffman, E. P.; Knudson, C. M.; Campbell, K. P.; Kunkel, L. M.

[83657] : Subcellular fractionation of dystrophin to the triads of skeletal

[83658] muscle. Nature 330: 754-758, 1987.

[83659] 14589.

[83660] Hoffman, E. P.; Monaco, A. P.; Feener, C. C.; Kunkel, L. M.:

[83661] Conservation of the Duchenne muscular dystrophy gene in mice and humans. Science 238:

[83662] 347-350, 1987.

[83663] 14590.

[83664] Hoop, R. C.; Russo, L. S.; Riconda, D. L.; Schwartz, L. S.; Hoffman,

[83665] E. P.: Restoration of half the normal dystrophin sequence in a double-deletion

[83666] Duchenne muscular dystrophy family. Am. J. Med. Genet. 49: 323-327,

[83667] 1994.

[83668] 14591.

[83669] Hoffman, E. P.; Fischbeck, K. H.; Brown, R. H.; Johnson, M.; Medori,

[83670] R.; Loike, J. D.; Harris, J. B.; Waterston, R.; Brooke, M.; Specht,

[83671] L.; Kupsky, W.; Chamberlain, J.; Caskey, C. T.; Shapiro, F.; Kunkel,

[83672] L. M.: Characterization of dystrophin in muscle-biopsy specimens

[83673] from patients with Duchenne's or Becker's muscular dystrophy. New

[83674] Eng. J. Med. 318: 1363-1368, 1988.

[83675] 14592.

[83676] Howard, P. L.; Dally, G. Y.; Wong, M. H.; Ho, A.; Weleber, R.

[83677] G.; Pillers, D.-A. M.; Ray, P. N.: Localization of dystrophin isoform

[83678] Dp71 to the inner limiting membrane of the retina suggests a unique

[83679] functional contribution of Dp71 in the retina. Hum. Molec. Genet. 7:

[83680] 1385-1391, 1998.

[83681] 14593.

[83682] Hu, X.; Burghes, A. H. M.; Bulman, D. E.; Ray, P. N.; Wor-
ton,
[83683] R. G.: Evidence for mutation by unequal sister chromatid
exchange
[83684] in the Duchenne muscular dystrophy gene. Am. J. Hum.
Genet. 44:
[83685] 855-863, 1989.

[83686] 14594.

[83687] Alon, U.; Chan, J. C. M.: Hydrochlorothiazide-amiloride in
the
[83688] treatment of congenital nephrogenic diabetes insipidus.
Am. J. Nephrol. 5:
[83689] 9-13, 1985.

[83690] 14595.

[83691] Andreoli, T. E.; Schafer, J. A.: Nephrogenic diabetes in-
sipidus.In:
[83692] Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.:
Metabolic
[83693] Basis of Inherited Disease. New York: McGraw-Hill (pub.)
(4th

[83694] ed.): 1978. Pp. 1634-1659.

[83695] 14596.

[83696] Barak, L. S.; Oakley, R. H.; Laporte, S. A.; Caron, M. G.: Constitutive

[83697] arrestin-mediated desensitization of a human vasopressin receptor

[83698] mutant associated with nephrogenic diabetes insipidus. Proc. Nat.

[83699] Acad. Sci. 98: 93-98, 2001.

[83700] 14597.

[83701] Bichet, D. G.: Personal Communication. Montreal, Quebec, Canada

[83702] 5/1/1994.

[83703] 14598.

[83704] Bichet, D. G.; Arthus, M.-F.; Lonergan, M.; Hendy, G. N.; Paradis,

[83705] A. J.; Fujiwara, T. M.; Morgan, K.; Gregory, M. C.; Rosen-thal, W.;

[83706] Didwania, A.; Antaramian, A.; Birnbaumer, M.: X-linked nephrogenic

[83707] diabetes insipidus mutations in North America and the

Hopewell hypothesis. J.
[83708] Clin. Invest. 92: 1262-1268, 1993.

[83709] 14599.

[83710] Bichet, D. G.; Razi, M.; Arthus, M.-F.; Lonergan, M.; Tittley,

[83711] P.; Smiley, R. K.; Rock, G.; Hirsch, D. J.: Epinephrine and dDAVP
[83712] administration in patients with congenital nephrogenic diabetes insipidus:
[83713] evidence for a pre-cyclic AMP V(2) receptor defective mechanism. Kidney
[83714] Int. 36: 859-866, 1989.

[83715] 14600.

[83716] Bichet, D. G.; Razi, M.; Lonergan, M.; Arthus, M.-F.; Papukna,
[83717] V.; Kortas, C.; Barjon, J.-N.: Hemodynamic and coagulation responses
[83718] to 1-desamino(8-D-arginine) vasopressin in patients with congenital
[83719] nephrogenic diabetes insipidus. New Eng. J. Med. 318: 881-887, 1988.

[83720] 14601.

[83721] Birnbaumer, M.: Vasopressin receptors. TEM 11: 406-410, 2000.

[83722] 14602.

[83723] Birnbaumer, M.; Seibold, A.; Gilbert, S.; Ishido, M.; Barberis,

[83724] C.; Antaramian, A.; Brabet, P.; Rosenthal, W.: Molecular cloning

[83725] of the receptor for human antidiuretic hormone. Nature 357: 333-335,

[83726] 1992.

[83727] 14603.

[83728] Gabig, T. G.; Lefker, B. A.: Deficient flavoprotein component

[83729] of the NADPH-dependent superoxide-generating oxidase in the neutrophils

[83730] from three male patients with chronic granulomatous disease. J. Clin.

[83731] Invest. 73: 701-705, 1984.

[83732] 14604.

[83733] Heyworth, P. G.; Curnutte, J. T.; Rae, J.; Noack, D.; Cross, A.

[83734] R.: Hematologically important mutations: X-linked chronic granulomatous
[83735] disease--an update. Blood Cells Molec. Dis. 23: 443-450, 1997.
[83736] 14605.

[83737] Ho, C. M. L.; Vowels, M. R.; Lockwood, L.; Ziegler, J. B.: Successful
[83738] bone marrow transplantation in a child with X-linked chronic granulomatous
[83739] disease. Bone Marrow Transplant. 18: 213-215, 1996.

[83740] 14606.

[83741] Hohn, D. C.; Lehrer, R. I.: Identification of the defect in X-linked
[83742] chronic granulomatous disease. (Abstract) Clin. Res. 22: 394A only,
[83743] 1974.

[83744] 14607.

[83745] Holmes, B.; Page, A. R.; Good, R. A.: Studies of the metabolic
[83746] activity of leukocytes from patients with a genetic abnormality of

[83747] phagocytic function. J. Clin. Invest. 46: 1422-1432, 1967.

[83748] 14608.

[83749] Horn, D. C.; Lehrer, R. I.: NADPH oxidase deficiency in X-
linked
[83750] chronic granulomatous disease. J. Clin. Invest. 55:
707-713, 1975.
[83751] 14609.

[83752] Horwitz, M. E.; Barrett, A. J.; Brown, M. R.; Carter, C. S.;
Childs,
[83753] R.; Gallin, J. I.; Holland, S. M.; Linton, G. F.; Miller, J. A.;
Leitman,
[83754] S. F.; Read, E. J.; Malech, H. L.: Treatment of chronic gran-
ulomatous
[83755] disease with nonmyeloablative conditioning and a T-
cell-depleted hematopoietic
[83756] allograft. New Eng. J. Med. 344: 881-888, 2001.

[83757] 14610.

[83758] Hui, Y. F.; Chan, S. Y.; Lau, Y. L.: Identification of muta-
tions
[83759] in seven Chinese patients with X-linked chronic granulo-
matous disease. Blood 88:

[83760] 4021-4028, 1996.

[83761] 14611.

[83762] Ishibashi, F.; Mizukami, T.; Kanegasaki, S.; Motoda, L.; Kakinuma,

[83763] R.; Endo, F.; Nunoi, H.: Improved superoxide-generating ability by

[83764] interferon-gamma due to splicing pattern change of transcripts in

[83765] neutrophils from patients with a splice site mutation in CYBB gene. Blood 98:

[83766] 436-441, 2001.

[83767] 14612.

[83768] Ishibashi, F.; Nunoi, H.; Endo, F.; Matsuda, I.; Kanegasaki, S.

[83769] : Statistical and mutational analysis of chronic granulomatous disease

[83770] in Japan with special reference to gp91-phox and p22-phox deficiency. Hum.

[83771] Genet. 106: 473-481, 2000.

[83772] 14613.

[83773] Ito, K.; Mukumoto, Y.; Konishi, H.; Sakura, N.; Usui, T.: Kell

[83774] phenotypes in 15 Japanese patients with chronic granulomatous disease. Vox
[83775] Sang. 37: 39-40, 1979.

[83776] 14614.

[83777] Johnston, R. B., Jr.: Defects of neutrophil function. (Editorial) New
[83778] Eng. J. Med. 307: 434-436, 1982.

[83779] 14615.

[83780] Johnston, R. B., Jr.; Keele, B. B., Jr.; Misra, H. P.; Lehmeyer,
[83781] J. E.; Webb, L. S.; Baehner, R. L.; Rajagopalan, K. V.: The role
[83782] of superoxide anion generation in phagocytic bactericidal activity:
[83783] studies with normal and chronic granulomatous disease leukocytes. J.
[83784] Clin. Invest. 55: 1357-1372, 1975.

[83785] 14616.

[83786] Klebanoff, S. J.: Personal Communication. Seattle, Wash. 1/29/1982.
[83787] 14617.

[83788] Klebanoff, S. J.; Clark, R. A.: The Neutrophil: Function and

[83789] Clinical Disorders. Amsterdam: North Holland (pub.) 1978.

[83790] 14618.

[83791] Kontras, S. B.; Bodenbender, J. G.; McClave, C. R.; Smith, J.

[83792] P.: Interstitial cystitis in chronic granulomatous disease. J. Urol. 105:

[83793] 575-578, 1971.

[83794] 14619.

[83795] Kragballe, K.; Borregaard, N.; Brandrup, F.; Koch, C.; Jo-hansen,

[83796] K. S.: Relation of monocyte and neutrophil oxidative metabolism to

[83797] skin and oral lesions in carriers of chronic granulomatous disease. Clin.

[83798] Exp. Immun. 43: 390-398, 1981.

[83799] 14620.

[83800] Kunkel, L. M.; Monaco, A. P.; Middlesworth, W.; Ochs, H. D.; Latt,

[83801] S. A.: Specific cloning of DNA fragments absent from the DNA of a

[83802] male patient with an X chromosome deletion. Proc. Nat. Acad. Sci. 82:
[83803] 4778-4782, 1985.

[83804] 14621.

[83805] Leusen, J. H. W.; de Boer, M.; Bolscher, B. G. J. M.; Hilarius,

[83806] P. M.; Weening, R. S.; Ochs, H. D.; Roos, D.; Verhoeven, A. J.: A
[83807] point mutation in gp91-phox of cytochrome b(558) of the human NADPH
[83808] oxidase leading to defective translocation of the cytosolic proteins
[83809] p47-phox and p67-phox. J. Clin. Invest. 93: 2120-2126, 1994.
[83810] 14622.

[83811] Lew, P. D.; Southwick, F. S.; Stossel, T. P.; Whitin, J. C.; Si-mons,
[83812] E.; Cohen, H. J.: A variant of chronic granulomatous dis-ease: deficient
[83813] oxidative metabolism due to a low-affinity NADPH oxi-dase. New Eng.
[83814] J. Med. 305: 1329-1333, 1981.

[83815] 14623.

[83816] Lischner, H. W.; Martyn, L. J.: Chorioretinal lesions, sea-
blue
[83817] histiocytes and other manifestations in familial chronic
granulomatous
[83818] disease. Birth Defects Orig. Art. Ser. XI(1): 73-76, 1975.

[83819] 14624.

[83820] MacFarlane, P. S.; Speirs, A. L.; Sommerville, R. G.: Fatal
granulomatous
[83821] disease of childhood and benign lymphocytic infiltration
of the skin
[83822] (congenital dysphagocytosis). Lancet I: 408-410, 1967.

[83823] 14625.

[83824] Macher, A. M.; Casale, T. B.; Fauci, A. S.: Chronic granulo-
matous
[83825] disease of childhood and Chromobacterium violaceum in-
fections in the
[83826] southeastern United States. Ann. Intern. Med. 97: 51-55,
1982.
[83827] 14626.

[83828] Marsh, W. L.; Oyen, R.; Nichols, M. E.; Allen, F. H., Jr.:

Chronic
granulomatous disease and Kell blood groups. Brit. J. Haemat. 29:
247-262, 1975.

14627.

Marsh, W. L.; Uretsky, S. C.; Douglas, S. D.: Antigens of the
Kell blood group system on neutrophils and monocytes: their relation
to chronic granulomatous disease. J. Pediat. 87:
1117-1120, 1975.

14628.

Matthay, K. K.; Golbus, M. S.; Wara, D. W.; Mentzer, W. C.: Prenatal
diagnosis of chronic granulomatous disease. Am. J. Med. Genet. 17:
731-739, 1984.

14629.

McPhail, L. C.; DeChatelet, L. R.; Shirley, P. S.; Wilfert, C.;

Johnston, R. B., Jr.; McCall, C. E.: Deficiency of NADPH oxi-
dase

[83842] activity in chronic granulomatous disease. J. Pediat. 90: 213-217,
[83843] 1977.

[83844] 14630.

[83845] Meischl, C.; de Boer, M.; Ahlin, A.; Roos, D.: A new exon created
[83846] by intronic insertion of a rearranged LINE-1 element as the cause
[83847] of chronic granulomatous disease. Europ. J. Hum. Genet. 8: 697-703,
[83848] 2000.

[83849] 14631.

[83850] Hameister, H.; Wolff, G.; Laurietzen, C. H.; Lehmann, W. O.; Hauser,
[83851] A.; Ropers, H. H.: Clinical and biochemical investigations on patients
[83852] with partial deficiency of placental steroid sulfatase. Hum. Genet. 46:
[83853] 199-207, 1979.

[83854] 14632.

[83855] Happle, R.: X-linked dominant ichthyosis. Clin. Genet. 15:

[83856] 239-240, 1979.

[83857] 14633.

[83858] Harris, H.: A pedigree of sex-linked ichthyosis vulgaris. Ann.

[83859] Eugen. 14: 9, 1947.

[83860] 14634.

[83861] Hernandez-Martin, A.; Gonzalez-Sarmiento, R.; De Unamuno, P.:

[83862] X-linked ichthyosis: an update. Brit. J. Derm. 141: 617-627, 1999.

[83863] 14635.

[83864] Jay, B.; Blach, R. K.; Wells, R. S.: Ocular manifestations of

[83865] ichthyosis. Brit. J. Ophthal. 52: 217-226, 1968.

[83866] 14636.

[83867] Jobsis, A. C.; De Groot, W. P.; Meijer, A. E. F. H.; van der Loos,

[83868] C. M.: A new method for the determination of steroid sulphatase activity

[83869] in leukocytes in X-linked recessive ichthyosis. Brit. J. Derm. 108:

[83870] 567-572, 1983.

[83871] 14637.

[83872] Jobsis, A. C.; De Groot, W. P.; Tigges, A. J.; De Bruijn, H. W.;

[83873] Rijken, Y.; Meijer, A. E. F. H.; Marinkovic-Ilsen, A.: X-linked ichthyosis

[83874] and X-linked placental sulfatase deficiency: a disease entity. Histochemical

[83875] observations. Am. J. Path. 99: 279-290, 1980.

[83876] 14638.

[83877] Jobsis, A. C.; van Duuren, C. Y.; de Vries, G. P.; Koppe, J. G.;

[83878] Rijken, Y.; van Kempen, G. M. J.; de Groot, W. P.: Trophoblast sulphatase

[83879] deficiency associated with X-chromosomal ichthyosis. (Abstract) Nederl.

[83880] T. Geneesk. 120: 1980, 1976.

[83881] 14639.

[83882] Kawano, J.; Kotani, T.; Ohtaki, S.; Mimamino, N.; Matsuo, H.;

[83883] Oinuma, T.; Aikawa, E.: Characterization of rat and human

steroid
[83884] sulfatases. Biochim. Biophys. Acta 997: 199-205, 1989.

[83885] 14640.

[83886] Keitges, E.; Gartler, S. M.: Dosage of the Sts gene in the mouse. Am.
[83887] J. Hum. Genet. 39: 470-476, 1986.

[83888] 14641.

[83889] Keitges, E.; Rivest, M.; Siniscalco, M.; Gartler, S. M.: X-linkage
[83890] of steroid sulphatase in the mouse is evidence for a functional Y-linked
[83891] allele. Nature 315: 226-227, 1985.

[83892] 14642.

[83893] Kerr, C. B.; Wells, R. S.: Sex-linked ichthyosis. Ann. Hum.
[83894] Genet. 29: 33-50, 1965.

[83895] 14643.

[83896] Kerr, C. B.; Wells, R. S.; Sanger, R.: X-linked ichthyosis and
[83897] the Xg groups. Lancet II: 1369-1370, 1964.

[83898] 14644.

[83899] Kipling, D.; Salido, E. C.; Shapiro, L. J.; Cooke, H. J.: High

[83900] frequency de novo alterations in the long-range genomic structure

[83901] of the mouse pseudoautosomal region. Nature Genet. 13: 78-82, 1996.

[83902] 14645.

[83903] Koppe, J. G.; Marinkovic-Ilsen, A.; Rijken, Y.; De Groot, W. P.;

[83904] Jobsis, A. C.: X-linked ichthyosis: a sulphatase deficiency. Arch.

[83905] Dis. Child. 53: 803-806, 1978.

[83906] 14646.

[83907] Koppe, J. G.; Rijken, Y.; Jobsis, A. C.; Marinkovic-Ilsen, A.

[83908] : X-linked ichthyosis, a sulfatase deficiency. (Abstract) Vth Int.

[83909] Conf. on Birth Defects, Montreal , 8/1977.

[83910] 14647.

[83911] Kubilus, J.; Tarascio, A. J.; Baden, H. P.: Steroid-sulfatase

[83912] deficiency in sex-linked ichthyosis. Am. J. Hum. Genet. 31: 50-53,

[83913] 1979.

[83914] 14648.

[83915] Laval, S. H.; Boyd, Y.: Partial inversion of gene order within

[83916] a homologous segment on the X chromosome. Mam-
malian Genome 4: 119-123,

[83917] 1993.

[83918] 14649.

[83919] Li, X. M.; Salido, E. C.; Gong, Y.; Kitada, K.; Serikawa, T.;

[83920] Yen, P. H.; Shapiro, L. J.: Cloning of the rat steroid sulfa-
tase

[83921] gene (Sts), a non-pseudoautosomal X-linked gene that
undergoes X inactivation. Mammalian

[83922] Genome 7: 420-424, 1996.

[83923] 14650.

[83924] Lykkesfeldt, G.; Bennett, P.; Lykkesfeldt, A. E.; Micic, S.;
Moller,

[83925] S.; Svenstrup, B.: Abnormal androgen and oestrogen
metabolism in

[83926] men with steroid sulphatase deficiency and recessive X-
linked ichthyosis. Clin.

[83927] Endocr. 23: 385-393, 1985.

[83928] 14651.

[83929] Lykkesfeldt, G.; Hoyer, H.: Topical cholesterol treatment of

[83930] recessive X-linked ichthyosis. Lancet II: 1337-1338, 1983.

[83931] 14652.

[83932] Lykkesfeldt, G.; Hoyer, H.; Ibsen, H. H.; Brandrup, F.: Steroid

[83933] sulphatase deficiency disease. Clin. Genet. 28: 231-237, 1985.

[83934] 14653.

[83935] Lykkesfeldt, G.; Hoyer, H.; Lykkesfeldt, A. E.; Skakkebaek, N.

[83936] E.: Steroid sulphatase deficiency associated with testis cancer. Lancet II:

[83937] 1456, 1983.

[83938] 14654.

[83939] Lykkesfeldt, G.; Lykkesfeldt, A. E.; Skakkebaek, N. E.: Steroid

[83940] sulphatase in man: a non inactivated X-locus with partial gene dosage

[83941] compensation. Hum. Genet. 65: 355-357, 1984.

[83942] 14655.

[83943] Macsai, M. S.; Doshi, H.: Clinical pathologic correlation of

[83944] superficial corneal opacities in X-linked ichthyosis. Am. J. Ophthal. 118:

[83945] 477-484, 1994.

[83946] 14656.

[83947] Marinkovic-Ilsen, A.; Koppe, J. G.; Jobsis, A. C.; De Groot, W.

[83948] P.: Enzymatic basis of typical X-linked ichthyosis. (Letter) Lancet II:

[83949] 1097, 1978.

[83950] 14657.

[83951] Metaxotou, C.; Ikkos, D.; Panagiotopoulou, P.; Alevizaki, M.;

[83952] Mavrou, A.; Tsenghi, C.; Matsaniotis, N.: A familial X/Y translocation

[83953] in a boy with ichthyosis, hypogonadism and mental retar- dation. Clin.

[83954] Genet. 24: 380-383, 1983.

[83955] 14658.

[83956] Meyer, J. C.; Weiss, H.; Grundemann, H. P.; Wurseh, T. G.; Schnyder,

[83957] U. W.: Deficiency of arylsulfatase C in cultured skin fibroblasts

[83958] of X-linked ichthyosis. Hum. Genet. 53: 115-116, 1979.

[83959] 14659.

[83960] Migeon, B. R.; Shapiro, L. J.; Norum, R. A.; Mohandas, T.; Axelman,

[83961] J.; Dabora, R. L.: Differential expression of steroid sulphatase

[83962] locus on active and inactive human X chromosome. Nature 299: 838-840,

[83963] 1982.

[83964] 14660.

[83965] Mohandas, T.; Shapiro, L. J.; Sparkes, R. S.; Sparkes, M. C.:

[83966] Regional assignment of the steroid sulfatase--X-linked ichthyosis

[83967] locus: implications for a non-inactivated region on the short arm

[83968] of the human X-chromosome. Proc. Nat. Acad. Sci. 76:

5779-5783,

[83969] 1979.

[83970] 14661.

[83971] Mohandas, T.; Sparkes, R. S.; Hellkuhl, B.; Grzeschik, K. H.;

[83972] Shapiro, L. J.: Expression of an X-linked gene from an in-
active human

[83973] X chromosome in mouse-human hybrid cells: further evi-
dence for the

[83974] noninactivation of the steroid sulfatase locus in man. Proc.
Nat.

[83975] Acad. Sci. 77: 6759-6763, 1980.

[83976] 14662.

[83977] Mohandas, T. K.; Stern, H. J.; Meeker, C. A.; Passage, M. B.;

[83978] Muller, U.; Page, D. C.; Yen, P. H.; Shapiro, L. J.: Steroid
sulfatase

[83979] gene in XX males. Am. J. Hum. Genet. 46: 369-376, 1990.

[83980] 14663.

[83981] Monroe, D. G.; Chang, P. L.: Tissue-specific expression of
human

[83982] arylsulfatase-C isozymes and steroid sulfatase. Am. J.

Hum. Genet. 40:
[83983] 102-114, 1987.

[83984] 14664.

[83985] Allingham-Hawkins, D. J.; Ray, P. N.: FRAXE expansion is not a
[83986] common etiological factor among developmentally de-layed males. Am.
[83987] J. Hum. Genet. 57: 72-76, 1995.

[83988] 14665.

[83989] Barnicoat, A. J.; Wang, Q.; Turk, J.; Green, E.; Mathew, C. G.;
[83990] Flynn, G.; Buckle, V.; Hirst, M.; Davies, K.; Bobrow, M.: Clinical,
[83991] cytogenetic, and molecular analysis of three families with FRAXE. J.
[83992] Med. Genet. 34: 13-17, 1997.

[83993] 14666.

[83994] Brown, W. T.: The FRAXE syndrome: is it time for routine screening?
[83995] (Editorial) Am. J. Hum. Genet. 58: 903-905, 1996.

[83996] 14667.

[83997] Chakrabarti, L.; Knight, S. J. L.; Flannery, A. V.; Davies, K.

[83998] E.: A candidate gene for mild mental handicap at the FRAXE fragile

[83999] site. Hum. Molec. Genet. 5: 275-282, 1996.

[84000] 14668.

[84001] Ennis, S.; Murray, A.; Morton, N. E.: Haplotypic determi-nants

[84002] of instability in the FRAX region: concatenated mutation or founder

[84003] effect? Hum. Mutat. 18: 61-69, 2001.

[84004] 14669.

[84005] Gecz, J.; Bielby, S.; Sutherland, G. R.; Mulley, J. C.: Gene structure

[84006] and subcellular localization of FMR2, a member of a new family of

[84007] putative transcription activators. Genomics 44: 201-213, 1997.

[84008] 14670.

[84009] Gecz, J.; Gedeon, A. K.; Sutherland, G. R.; Mulley, J. C.: Identification

[84010] of the gene FMR2, associated with FRAXE mental retarda-

tion. Nature
[84011] Genet. 13: 105-108, 1996.

[84012] 14671.

[84013] Gecz, J.; Oostra, B. A.; Hockey, A.; Carbonell, P.; Turner, G.;
[84014] Haan, E. A.; Sutherland, G. R.; Mulley, J. C.: FMR2 expression in
[84015] families with FRAXE mental retardation. Hum. Molec. Genet. 6: 435-441,
[84016] 1997.

[84017] 14672.

[84018] Gu, Y.; Shen, Y.; Gibbs, R. A.; Nelson, D. L.: Identification
[84019] of FMR2, a novel gene associated with the FRAXE CCG repeat and CpG
[84020] island. Nature Genet. 13: 109-113, 1996.

[84021] 14673.

[84022] Yoshima, T.; Yura, T.; Yanagi, H.: Novel testis-specific protein
[84023] that interacts with heat shock factor 2. Gene 214: 139-146, 1998.
[84024] 14674.

[84025] Kools, P.; Vanhalst, K.; Wan den Eynde, E.; van Roy, F.: The human

[84026] cadherin-10 gene: complete coding sequence, predominant expression

[84027] in the brain and mapping on chromosome 5p13-13. FEBS Lett. 452:

[84028] 328-334, 1999.

[84029] 14675.

[84030] Leder, S.; Weber, Y.; Altafaj, X.; Estivill, X.; Joost, H.-G.;

[84031] Becker, W.: Cloning and characterization of DYRK1B, a novel member

[84032] of the DYRK family of protein kinases. Biochem. Biophys. Res. Commun. 254:

[84033] 474-479, 1999.

[84034] 14676.

[84035] Lee, K.; Deng, X.; Friedman, E.: Mirk protein kinase is a mitogen-activated

[84036] protein kinase substrate that mediates survival of colon cancer cells. Cancer

[84037] Res. 60: 3631-3637, 2000.

[84038] 14677.

[84039] Coyle, A. J.; Lehar, S.; Lloyd, C.; Tian, J.; Delaney, T.; Manning,

[84040] S.; Nguyen, T.; Burwell, T.; Schneider, H.; Gonzalo, J. A.; Gosselin,

[84041] M.; Owen, L. R.; Rudd, C. E.; Gutierriez-Ramos, J. C.: The CD28-related

[84042] molecule ICOS is required for effective T cell-dependent immune responses. Immunity 13:

[84043] 95-105, 2000.

[84044] 14678.

[84045] Dong, C.; Juedes, A. E.; Temann, U. A.; Shresta, S.; Allison, J.

[84046] P.; Ruddle, N. H.; Flavell, R. A.: ICOS co-stimulatory receptor is

[84047] essential for T-cell activation and function. Nature 409: 97-101,

[84048] 2001.

[84049] 14679.

[84050] Haimila, K. E.; Partanen, J. A.; Holopainen, P. M.: Genetic polymorphism

[84051] of the human ICOS gene. Immunogenetics 53: 1028-1032, 2002.

[84052] 14680.

[84053] Hutloff, A.; Dittrich, A. M.; Beier, K. C.; Eljaschewitsch, B.;

[84054] Kraft, R.; Anagnostopoulos, I.; Kroczek, R. A.: ICOS is an inducible

[84055] T-cell co-stimulator structurally and functionally related to CD28. Nature 397:

[84056] 263-266, 1999.

[84057] 14681.

[84058] McAdam, A. J.; Greenwald, R. J.; Levin, M. A.; Chernova, T.; Malenkovich,

[84059] N.; Ling, V.; Freeman, G. J.; Sharpe, A. H.: ICOS is critical for

[84060] CD40-mediated antibody class switching. Nature 409: 102-105, 2001.

[84061] 14682.

[84062] Schillace, R. V.; Andrews, S. F.; Liberty, G. A.; Davey, M. P.;

[84063] Carr, D. W.: Identification and characterization of myeloid translocation

[84064] gene 16b as a novel A kinase anchoring protein in T lym-phocytes. J.

[84065] Immun. 168: 1590-1599, 2002.

[84066] 14683.

[84067] Pack, S. D.; Pak, E.; Tanigami, A.; Ledbetter, D. H.; Fukuda, M.

[84068] N.: Assignment of the bystin gene BYSL to human chromosome band 6p21.1

[84069] by in situ hybridization. Cytogenet. Cell Genet. 83: 76-77, 1998.

[84070] 14684.

[84071] Leegwater, P. A. J.; Vermeulen, G.; Konst, A. A. M.; Naidu, S.;

[84072] Mulders, J.; Visser, A.; Kersbergen, P.; Mobach, D.; Fonds, D.; van

[84073] Berkel, C. G. M.; Lemmers, R. J. L. F.; Frants, R. R.; Oudejans, C.

[84074] B. M.; Schutgens, R. B. H.; Pronk, J. C.; van der Knaap, M. S.: Subunits

[84075] of the translation initiation factor eiF2B are mutant in leukoencephalopathy

[84076] with vanishing white matter. Nature Genet. 29: 383-388, 2001.

[84077] 14685.

[84078] Medina, M.; Marinescu, R. C.; Overhauser, J.; Kosik, K. S.:

Hemizygosity
[84079] of delta-catenin (CTNND2) is associated with severe mental retardation
[84080] in cri-du-chat syndrome. Genomics 63: 157-164, 2000.

[84081] 14686.

[84082] Paffenholz, R.; Franke, W. W.: Identification and localization
[84083] of a neurally expressed member of the plakoglobin/armadillo multigene
[84084] family. Differentiation 61: 293-304, 1997.

[84085] 14687.

[84086] Zhou, J..; Liyanage, U.; Medina, M.; Ho, C.; Simmons, A. D.; Lovett.
[84087] M.; Kosik, K. S.: Presenilin 1 interaction in the brain with a novel
[84088] member of the armadillo family. Neuroreport 8: 2085-2090, 1997.
[84089] 14688.

[84090] Burger, J.; Fonknechten, N.; Hoeltzenbein, M.; Neumann, L.; Bratanoff,
[84091] E.; Hazan, J.; Reis, A.: Hereditary spastic paraplegia caused

by
[84092] mutations in the SPG4 gene. Europ. J. Hum. Genet. 8:
771-776, 2000.
[84093] 14689.
[84094] Burger, J.; Metzke, H.; Paternotte, C.; Schilling, F.; Hazan,
J.;
[84095] Reis, A.: Autosomal dominant spastic paraplegia with an-
ticipation
[84096] maps to a 4-cM interval on chromosome 2p21-p24 in a
large German family. Hum.
[84097] Genet. 98: 371-375, 1996.
[84098] 14690.
[84099] Errico, A.; Ballabio, A.; Rugarli, E. I.: Spastin, the protein
[84100] mutated in autosomal dominant hereditary spastic para-
plegia, is involved
[84101] in microtubule dynamics. Hum. Molec. Genet. 11:
153-163, 2002.
[84102] 14691.
[84103] Lindsey, J. C.; Lusher, M. E.; McDermott, C. J.; White, K. D.;
[84104] Reid, E.; Rubinsztein, D. C.; Bashir, R.; Hazan, J.; Shaw, P.
J.;

[84105] Bushby, K. M. D.: Mutation analysis of the spastin gene (SPG4) in
[84106] patients with hereditary spastic paraplegia. J. Med. Genet. 37:
[84107] 759-765, 2000.

[84108] 14692.

[84109] Takahashi, N.; Tuiki, H.; Saya, H.; Kaibuchi, K.: Localization
[84110] of the gene coding for ROCK II/Rho kinase on human chromosome 2p24. Genomics 55:
[84111] 235-237, 1999.

[84112] 14693.

[84113] Gruber, A. D.; Pauli, B. U.: Clustering of the human CLCA gene
[84114] family on the short arm of chromosome 1 (1p22-31). Genome 42: 1030-1032,
[84115] 1999.

[84116] 14694.

[84117] Leegwater, P. A. J.; Yuan, B. Q.; van der Steen, J.; Mulders, J.;
[84118] Konst, A. A. M.; Ilja Boor, P. K.; Mejaski-Bosnjak, V.; van

der Maarel,
[84119] S. M.; Frants, R. R.; Oudejans, C. B. M.; Schutgens, R. B. H.; Pronk,
[84120] J. C.; van der Knapp, M. S.: Mutations of MLC1 (KIAA0027), encoding
[84121] a putative membrane protein, cause megalencephalic leukoencephalopathy
[84122] with subcortical cysts. Am. J. Hum. Genet. 68: 831-838, 2001.
[84123] 14695.

[84124] Kedra, D.; Pan, H.-Q.; Seroussi, E.; Fransson, I.; Guilbaud, C.;
[84125] Collins, J. E.; Dunham, I.; Blennow, E.; Roe, B. A.; Piehl, F.; Dumanski,
[84126] J. P.: Characterization of the human synaptogyrin gene family. Hum.
[84127] Genet. 103: 131-141, 1998.

[84128] 14696.

[84129] Hazan, J.; Davoine, C. S.; Mavel, D.; Fonknechten, N.; Pa-
ternotte,
[84130] C.; Fizames, C.; Cruaud, C.; Samson, D.; Muselet, D.; Vega-Czarny,

[84131] N.; Brice, A.; Gyapay, G.; Heilig, R.; Fontaine, B.; Weis-
senbach,
[84132] J.: A fine integrated map of the SPG4 locus excludes an
expanded
[84133] CAG repeat in chromosome 2p-linked autosomal domi-
nant spastic paraplegia. Genomics 60:
[84134] 309-319, 1999.

[84135] 14697.

[84136] Hazan, J.; Fonknechten, N.; Mavel, D.; Paternotte, C.; Sam-
son,
[84137] D.; Artiguenave, F.; Davoine, C.-S.; Cruaud, C.; Durr, A.;
Wincker,
[84138] P.; Brottier, P.; Cattolico, L.; Barbe, V.; Burgunder, J.-M.;
Prud'homme,
[84139] J.-F.; Brice, A.; Fontaine, B.; Heilig, R.; Weissenbach, J.:
Spastin,
[84140] a new AAA protein, is altered in the most frequent form of
autosomal
[84141] dominant spastic paraplegia. Nature Genet. 23: 296-303,
1999.
[84142] 14698.

[84143] Bedford, F. K.; Ashworth, A.; Enver, T.; Wiedemann, L. M.:

HEX:
[84144] a novel homeobox gene expressed during haematopoiesis and conserved
[84145] between mouse and human. Nucleic Acids Res. 21: 1245-1249, 1993.
[84146] 14699.

[84147] Crompton, M. R.; Bartlett, T. J.; MacGregor, A. D.; Manfio-letti,
[84148] G.; Buratti, E.; Giancotti, V.; Goodwin, G. H.: Identification of
[84149] a novel vertebrate homeobox gene expressed in haematopoietic cells. Nucleic
[84150] Acids Res. 20: 5661-5667, 1992.

[84151] 14700.

[84152] D'Elia, A. V.; Tell, G.; Russo, D.; Arturi, F.; Puglisi, F.; Man-fioletti,
[84153] G.; Gattei, V.; Mack, D. L.; Cataldi, P.; Filetti, S.; Di Loreto,
[84154] C.; Damante, G.: Expression and localization of the home-odomain-containing
[84155] protein HEX in human thyroid tumors. J. Clin. Endocr. Metab. 87:
[84156] 1376-1383, 2002.

[84157] 14701.

[84158] Hromas, R.; Radich, J.; Collins, S.: PCR cloning of an or-
phan
[84159] homeobox gene (PRH) preferentially expressed in myeloid
and liver
[84160] cells. Biochem. Biophys. Res. Commun. 195: 976-983,
1993.
[84161] 14702.

[84162] Morgutti, M.; Demori, E.; Pecile, V.; Amoroso, A.; Rustighi,
A.;
[84163] Manfioletti, G.: Genomic organization and chromosome
mapping of the
[84164] human homeobox gene HHEX. Cytogenet. Cell Genet. 94:
30-32, 2001.
[84165] 14703.

[84166] Tanaka, T.; Inazu, T.; Yamada, K.; Myint, Z.; Keng, V. W;
Inoue,
[84167] Y.; Taniguchi, N.; Noguchi, T.: cDNA cloning and expres-
sion of rat
[84168] homeobox gene, Hex, and functional characterization of
the protein. Biochem.
[84169] J. 339: 111-117, 1999.

[84170] 14704.

[84171] Miles, M. F.; Barhite, S.; Sganga, M.; Elliott, M.: Phosducin-like

[84172] protein: an ethanol-responsive potential modulator of guanine nucleotide-binding

[84173] protein function. Proc. Nat. Acad. Sci. 90: 10831-10835, 1993.

[84174] 14705.

[84175] Thibault, C.; Wang, J. F.; Charnas, R.; Mirel, D.; Barhite, S.;

[84176] Miles, M. F.: Cloning and characterization of the rat and human phosducin-like

[84177] protein genes: structure, expression and chromosomal localization. Biochim.

[84178] Biophys. Acta 1444: 346-354, 1999.

[84179] 14706.

[84180] Her, C.; Wood, T. C.; Eichler, E. E.; Mohrenweiser, H. W.; Ramagli,

[84181] L. S.; Siciliano, M. J.; Weinshilboum, R. M.: Human hydroxysteroid

[84182] sulfotransferase SULT2B1: two enzymes encoded by a single chromosome

[84183] 19 gene. Genomics 53: 284-295, 1998.

[84184] 14707.

[84185] Pope, R. K.; Pestonjamasp, K. N.; Smith, K. P.; Wulfkuhle, J. D.;

[84186] Strassel, C. P.; Lawrence, J. B.; Luna, E. J.: Cloning, characterization,

[84187] and chromosomal localization of human supervillin (SVIL). Genomics 52:

[84188] 342-351, 1998.

[84189] 14708.

[84190] Ting, H.-J.; Yeh, S.; Nishimura, K.; Chang, C.: Supervillin associates

[84191] with androgen receptor and modulates its transcriptional activity. Proc.

[84192] Nat. Acad. Sci. 99: 661-666, 2002.

[84193] 14709.

[84194] Bianchi, V.; Robles, R.; Alberio, L.; Furlan, M.; Lammle, B.:

[84195] Von Willebrand factor-cleaving protease (ADAMTS13) in thrombocytopenic

[84196] disorders: a severely deficient activity is specific for thrombotic

[84197] thrombocytopenic purpura. Blood 100: 710-713, 2002.

[84198] 14710.

[84199] Fujikawa, K.; Suzuki, H.; McMullen, B.; Chung, D.: Purification

[84200] of human von Willebrand factor-cleaving protease and its identification

[84201] as a new member of the metalloproteinase family. Blood 98: 1662-1666,

[84202] 2001.

[84203] 14711.

[84204] Furlan, M.; Robles, R.; Solenthaler, M.; Wassmer, M.; Sandoz, Pl;

[84205] Lammle, B.: Deficient activity of von Willebrand factor-cleaving

[84206] protease in chronic relapsing thrombotic thrombocytopenic purpura. Blood 89:

[84207] 3097-3103, 1997.

[84208] 14712.

[84209] Gerritsen, H. E.; Robles, R.; Lammle, B.; Furlan, M.: Partial

[84210] amino acid sequence of purified von Willebrand factor-cleaving protease. Blood 98:

[84211] 1654-1661, 2001.

[84212] 14713.

[84213] Kokame, K.; Matsumoto, M.; Soejima, K.; Yagi, H.; Ishiza-
shi, H.;

[84214] Funato, M.; Tamai, H.; Konno, M.; Kamide, K.; Kawano, Y.;
Miyata,

[84215] T.; Fujimura, Y.: Mutations and common polymorphisms
in ADAMTS13

[84216] gene responsible for von Willebrand factor-cleaving pro-
tease activity. Proc.

[84217] Nat. Acad. Sci. 99: 11902-11907, 2002.

[84218] 14714.

[84219] Re, F.; Strominger, J. L.: Toll-like receptor 2 (TLR2) and
TLR4

[84220] differentially activate human dendritic cells. J. Biol. Chem.
276:

[84221] 37692-37699, 2001.

[84222] 14715.

[84223] Wolfs, T. G. A. M.; Buurman, W. A.; van Schadewijk, A.; de
Vries,

[84224] B.; Daemen, M. A. R. C.; Hiemstra, P. S.; van't Veer, C.: In

vivo

[84225] expression of Toll-like receptor 2 and 4 by renal epithelial cells:

[84226] IFN-gamma and TNF-alpha mediated up-regulation during inflammation. J.

[84227] Immun. 168: 1286-1293, 2002.

[84228] 14716.

[84229] Arbour, N. C.; Lorenz, E.; Schutte, B. C.; Zabner, J.; Klein, J.

[84230] N.; Jones, M.; Frees, K.; Watt, J. L.; Schwartz, D. A.: TLR4 mutations

[84231] are associated with endotoxin hyporesponsiveness in humans. Nature

[84232] Genet. 25: 187-191, 2000.

[84233] 14717.

[84234] Frantz, S.; Kobzik, L.; Kim, Y.-D.; Fukazawa, R.; Medzhitov, R.;

[84235] Lee, R. T.; Kelly, R. A.: Toll4 (TLR4) expression in cardiac myocytes

[84236] in normal and failing myocardium. J. Clin. Invest. 104: 271-280,

[84237] 1999.

[84238] 14718.

[84239] Kiechl, S.; Lorenz, E.; Reindl, M.; Wiedermann, C. J.; Oberhollenzer,

[84240] F.; Bonora, E.; Willeit, J.; Schwartz, D. A.: Toll-like receptor

[84241] 4 polymorphisms and atherogenesis. New Eng. J. Med. 347: 185-192,

[84242] 2002.

[84243] 14719.

[84244] Medzhitov, R.; Preston-Hurlburt, P.; Janeway, C. A., Jr.: A human

[84245] homologue of the Drosophila Toll protein signals activation of adaptive

[84246] immunity. Nature 388: 394-397, 1997.

[84247] 14720.

[84248] Michalek, S. M.; Moore, R. N.; McGhee, J. R.; Rosenstreich, D.

[84249] L.; Mergenhagen, S. E.: The primary role of lymphoreticular cells

[84250] in the mediation of host responses to bacterial endotoxin. J. Infect.

[84251] Dis. 141: 55-63, 1980.

[84252] 14721.

[84253] Nagai, Y.; Akashi, S.; Nagafuku, M.; Ogata, M.; Iwakura, Y.; Akira,

[84254] S.; Kitamura, T.; Kosugi, A.; Kimoto, M.; Miyake, K.: Essen-tial role

[84255] of MD-2 in LPS responsiveness and TLR4 distribution. Na-ture Immun. 3:

[84256] 667-672, 2002.

[84257] 14722.

[84258] Poltorak, A.; He, X.; Smirnova, I.; Liu, M.-Y.; Van Huffel, C.;

[84259] Du, X.; Birdwell, D.; Alejos, E.; Silva, M.; Galanos, C.; Freudenberg,

[84260] M.; Ricciardi-Castagnoli, P.; Layton, B.; Beutler, B.: Defec-tive

[84261] LPS signaling in C3H/HeJ and C57BL/10ScCr mice: muta-tions in Tlr4

[84262] gene. Science 282: 2085-2088, 1998.

[84263] 14723.

[84264] Rehli, M.; Poltorak, A.; Schwarzfischer, L.; Krause, S. W.; Andreesen,

[84265] R.; Beutler, B.: PU.1 and interferon consensus sequence-

binding protein

[84266] regulate the myeloid expression of the human Toll-like receptor 4

[84267] gene. J. Biol. Chem. 275: 9773-9781, 2000.

[84268] 14724.

[84269] Advani, R. J.; Bae, H.-R.; Bock, J. B.; Chao, D. S.; Doung, Y.-C.;

[84270] Prekeris, R.; Yoo, J.-S.; Scheller, R. H.: Seven novel mam-

malian

[84271] SNARE proteins localize to distinct membrane compart-

ments. J. Biol.

[84272] Chem. 273: 10317-10324, 1998.

[84273] 14725.

[84274] Bock, J. B.; Scheller, R. H.: A fusion of new ideas. Nature 387:

[84275] 133-135, 1997.

[84276] 14726.

[84277] Bui, T. D.; Wong, S. H.; Lu, L.; Hong, W.: Endobrevin maps to

[84278] chromosome 2 in human and chromosome 6 in mouse. Genomics 54: 579-580,

[84279] 1998.

[84280] 14727.

[84281] Wong, S. H.; Zhang, T.; Xu, Y.; Subramaniam, V. N.; Griffiths,

[84282] G.; Hong, W.: Endobrevin, a novel synaptobrevin/ VAMP-like protein

[84283] preferentially associated with the early endosome. Molec. Biol. Cell 9:

[84284] 1549-1563, 1998.

[84285] 14728.

[84286] Soilleux, E. J.; Barten, R.; Trowsdale, J.: Cutting edge: DC-SIGN;

[84287] a related gene, DC-SIGNR; and CD23 form a cluster on 19p13. J. Immun. 165:

[84288] 2937-2942, 2000.

[84289] 14729.

[84290] Matoskova, B.; Wong, W. T.; Nomura, N.; Robbins, K. C.; Di Fiore,

[84291] P. P.: RN-tre specifically binds to the SH3 domain of eps8 with high

[84292] affinity and confers growth advantage to NIH3T3 upon

carboxy-terminal
[84293] truncation. Oncogene 12: 2679-2688, 1996.

[84294] 14730.

[84295] Matoskova, B.; Wong, W. T.; Seki, N.; Nagase, T.; Nomura, N.; Robbins,
[84296] K. C.; Di Fiore, P. P.: RN-tre identifies a family of tre-related
[84297] proteins displaying a novel potential protein binding domain. Oncogene 12:
[84298] 2563-2571, 1996.

[84299] 14731.

[84300] Lee, M. P.; Brandenburg, S.; Landes, G. M.; Adams, M.; Miller,
[84301] G.; Feinberg, A. P.: Two novel genes in the center of the 11p15 imprinted
[84302] domain escape genomic imprinting. Hum. Molec. Genet. 8: 683-690,
[84303] 1999.

[84304] 14732.

[84305] Nakamura, M.; Masuda, H.; Horii, J.; Kuma, K.; Yokoyama, N.; Ohba,

[84306] T.; Nishitani, H.; Miyata, T.; Tanaka, M.; Nishimoto, T.: When overexpressed,

[84307] a novel centrosomal protein, RanBPM, causes ectopic microtubule nucleation

[84308] similar to gamma-tubulin. J. Cell Biol. 143: 1041-1052, 1998.

[84309] 14733.

[84310] Nishitani, H.; Hirose, E.; Uchimura, Y.; Nakamura,; M.; Umeda,

[84311] M.; Nishii, K.; Mori, N.; Nishimoto, T.: Full-sized RanBPM cDNA encodes

[84312] a protein possessing a long stretch of proline and glutamine within

[84313] the N-terminal region, comprising a large protein complex. Gene 272:

[84314] 25-33, 2001.

[84315] 14734.

[84316] Ohura, T.; Kobayashi, K.; Tazawa, Y.; Nishi, I.; Abukawa, D.; Sakamoto,

[84317] O.; Iinuma, K.; Saheki, T.: Neonatal presentation of adult-onset

[84318] type II citrullinemia. Hum. Genet. 108: 87-90, 2001.

[84319] 14735.

[84320] Tazawa, Y.; Kobayashi, K.; Ohura, T.; Abukawa, D.; Nishi-
nomiya,
[84321] F.; Hosoda, Y.; Yamashita, M.; Nagata, I.; Kono, Y.; Ya-
suda, T.; Yamaguchi,
[84322] N.; Saheki, T.: Infantile cholestatic jaundice associated
with adult-onset
[84323] type II citrullinemia. J. Pediat. 138: 735-740, 2001.

[84324] 14736.

[84325] Tomomasa, T.; Kobayashi, K.; Kaneko, H.; Shimura, H.;
Fukusato,
[84326] T.; Tabata, M.; Inoue, Y.; Ohwada, S.; Kasahara, M.; Mor-
ishita, Y.;
[84327] Kimura, M.; Saheki, T.; Morikawa, A.: Possible clinical and
histologic
[84328] manifestations of adult-onset type II citrullinemia in early
infancy. J.
[84329] Pediat. 138: 741-743, 2001.

[84330] 14737.

[84331] Bellefroid, E. J.; Marine, J. C.; Ried, T.; Lecocq, P. J.; Riviere,

[84332] M.; Amemiya, C.; Poncelet, D. A.; Coulie, P. G.; de Jong, P.;

Szpirer,

[84333] C.; Ward, D. C.; Martial, J. A.: Clustered organization of homologous

[84334] KRAB zinc-finger genes with enhanced expression in human T lymphoid

[84335] cells. EMBO J. 12: 1363-1374, 1993.

[84336] 14738.

[84337] Bellefroid, E. J.; Poncelet, D. A.; Lecocq, P. J.; Revelant, O.;

[84338] Martial, J. A.: The evolutionarily conserved Kruppel-associated box

[84339] domain defines a subfamily of eukaryotic multifingered proteins. Proc.

[84340] Nat. Acad. Sci. 88: 3608-3612, 1991.

[84341] 14739.

[84342] Williams, A. J.; Khachigian, L. M.; Shows, T.; Collins, T.: Isolation

[84343] and characterization of a novel zinc-finger protein with transcriptional

[84344] repressor activity. J. Biol. Chem. 270: 22143-22152, 1995.

[84345] 14740.

[84346] Grand, R. J. A.; Milner, A. E.; Mustoe, T.; Johnson, G. D.; Owen,
[84347] D.; Grant, M. L.; Gregory, C. D.: A novel protein expressed in mammalian
[84348] cells undergoing apoptosis. Exp. Cell Res. 218: 439-451, 1995.
[84349] 14741.
[84350] Twist, C. J.; Beier, D. R.; Disteche, C. M.; Edelhoff, S.; Ted-der,
[84351] T. F.: The mouse Cd83 gene: structure, domain organiza-tion, and chromosome
[84352] localization. Immunogenetics 48: 383-393, 1998.
[84353] 14742.
[84354] Lee, S. B.; Rao, A. K.; Lee, K.-H.; Yang, X.; Bae, Y. S.; Rhee,
[84355] S. G.: Decreased expression of phospholipase C-b2 isozyme in human
[84356] platelets with impaired function. Blood 88: 1684-1691, 1996.
[84357] 14743.
[84358] Mao, G. F.; Vaidyula, V. R.; Kunapuli, S. P.; Rao, A. K.: Lin-eage-specific

[84359] defect in gene expression in human platelet phospholipase C-beta-2

[84360] deficiency. Blood 99: 905-911, 2002.

[84361] 14744.

[84362] Park, D.; Jhon, D.-Y.; Kriz, R.; Knopf, J.; Rhee, S. G.: Cloning,

[84363] sequencing, expression, and Gq-independent activation of phospholipase

[84364] C-beta-2. J. Biol. Chem. 267: 16048-16055, 1992.

[84365] 14745.

[84366] Park, S. H.; Ryu, S. H.; Suh, P. G.; Kim, H.: Assignment of human

[84367] PLCB2 encoding PLC-beta-2 to human chromosome 15q15 by fluorescence

[84368] in situ hybridization. Cytogenet. Cell Genet. 83: 48-49, 1998.

[84369] 14746.

[84370] Rao, A. K.; Kowalska, M. A.; Disa, J.: Impaired cytoplasmic ionized

[84371] calcium mobilization in inherited platelet secretion defects. Blood 74:

[84372] 664-672, 1989.

[84373] 14747.

[84374] Yang, X.; Sun, L.; Ghosh, S.; Rao, A. K.: Human platelet signaling

[84375] defect characterized by impaired production of 1,4,5 inositol triphosphate

[84376] and phosphatic acid, and diminished pleckstrin phosphorylation. Evidence

[84377] for detective phospholipase C activation. Blood 88: 1676-1683, 1996.

[84378] 14748.

[84379] Schweitzer, B.; Taylor, V.; Welcher, A. A.; McClelland, M.; Suter,

[84380] U.: Neural membrane protein 35 (NMP35): a novel member of a gene

[84381] family which is highly expressed in the adult nervous system. Molec.

[84382] Cell. Neurosci. 11: 260-273, 1998.

[84383] 14749.

[84384] Somia, N. V.; Schmitt, M. J.; Vetter, D. E.; Van Antwerp, D.; Heinemann,

[84385] S. F.; Verma, I. M.: LFG: an anti-apoptotic gene that provides protection
[84386] from Fas-mediated cell death. Proc. Nat. Acad. Sci. 96: 12667-12672,
[84387] 1999.

[84388] 14750.

[84389] Allen, M.; Chu, S.; Brill, S.; Stotler, C.; Buckler, A.: Restricted
[84390] tissue expression pattern of a novel human rasGAP-related gene and
[84391] its murine ortholog. Gene 218: 17-25, 1998.

[84392] 14751.

[84393] Gillen, C. M.; Brill, S.; Payne, J. A.; Forbush, B., III: Molecular
[84394] cloning and functional expression of the K-Cl cotransporter from rabbit,
[84395] rat, and human: a new member of the cation-chloride cotransporter
[84396] family. J. Biol. Chem. 271: 16237-16244, 1996.

[84397] 14752.

[84398] Larsen, F.; Solheim, J.; Kristensen, T.; Kolsto, A. B.; Prydz,

[84399] H.: A tight cluster of five unrelated human genes on chromosome 16q22.1. Hum.
[84400] Molec. Genet. 2: 1589-1595, 1993.

[84401] 14753.

[84402] Pellegrino, C. M.; Rybicki, A. C.; Musto, S.; Nagel, R. L.; Schwartz,
[84403] R. S.: Molecular identification and expression of erythroid K:Cl
[84404] cotransporter in human and mouse erythroleukemic cells. Blood Cells
[84405] Molec. Dis. 24: 31-40, 1998.

[84406] 14754.

[84407] Hammond, E. M.; Brunet, C. L.; Johnson, G. D.; Parkhill, J.; Milner,
[84408] A. E.; Brady, G.; Gregory, C. D.; Grand, R. J. A.: Homology between
[84409] a human apoptosis specific protein and the product of APG5, a gene
[84410] involved in autophagy in yeast. FEBS Lett. 425: 391-395, 1998.
[84411] 14755.

[84412] Schmeiser, K.; Armstrong, S.; Hammond, E. M.; Grand, R. J. A.:

[84413] Assignment of the yeast APG5 human homologue APG5L to chromosome band

[84414] 6q21 by fluorescence in situ hybridisation. Cytogenet. Cell Genet. 87:

[84415] 213-214, 1999.

[84416] 14756.

[84417] Nakayama, M.; Nakajima, D.; Nagase, T.; Nomura, N.; Seki, N.; Ohara,

[84418] O.: Identification of high-molecular-weight proteins with multiple

[84419] EGF-like motifs by motif-trap screening. Genomics 51: 27-34, 1998.

[84420] 14757.

[84421] Wu, Q.; Maniatis, T.: Large exons encoding multiple ectodomains

[84422] are a characteristic feature of protocadherin genes. Proc. Nat. Acad.

[84423] Sci. 97: 3124-3129, 2000.

[84424] 14758.

[84425] Girard, J.-P.; Baekkevold, E. S.; Feliu, J.; Brandtzaeg, P.; Amalric,
[84426] F.: Molecular cloning and functional analysis of SUT-1, a sulfate
[84427] transporter from human high endothelial venules. Proc. Nat. Acad.
[84428] Sci. 96: 12772-12777, 1999.

[84429] 14759.

[84430] Rubio, J. P.; Levy, E. R.; Dobson-Stone, C.; Monaco, A. P.: Genomic
[84431] organization of the human G-alpha-14 and G-alpha-Q genes and mutation
[84432] analysis in chorea-acanthocytosis (CHAC). Genomics 57: 84-93, 1999.
[84433] 14760.

[84434] Paine-Saunders, S.; Viviano, B. L.; Saunders, S.: GPC6, a novel
[84435] member of the glypican gene family, encodes a product structurally
[84436] related to GPC4 and is colocalized with GPC5 on human chromosome 13. Genomics 57:
[84437] 455-458, 1999.

[84438] 14761.

[84439] Veugelers, M.; De Cat, B.; Ceulemans, H.; Bruystens, A. M.; Coomans,

[84440] C.; Durr, J.; Vermeesch, J.; Marynen, P.; David, G.: Glypican-6,

[84441] a new member of the glypican family of cell surface heparan sulfate

[84442] proteoglycans. J. Biol. Chem. 274: 26968-26977, 1999.

[84443] 14762.

[84444] Jacobs, S.; Schurmann, A.; Becker, W.; Bockers, T. M.; Copeland,

[84445] N. G.; Jenkins, N. A.; Joost, H.-G.: The mouse ADP-ribosylation factor-like

[84446] 4 gene: two separate promoters direct specific transcription in tissues

[84447] and testicular germ cell. Biochem. J. 335: 259-265, 1998.

[84448] 14763.

[84449] Oelkers, P.; Behari, A.; Cromley, D.; Billheimer, J. T.; Sturley,

[84450] S. L.: Characterization of two human genes encoding acyl coenzyme

[84451] A: cholesterol acyltransferase-related enzymes. J. Biol. Chem. 273:

[84452] 26765-26771, 1998.

[84453] 14764.

[84454] Bootcov, M. R.; Bauskin, A. R.; Valenzuela, S. M.; Moore, A. G.;

[84455] Bansal, M.; He, X. Y.; Zhang, H. P.; Donnellan, M.; Mahler, S.; Pryor,

[84456] K.; Walsh, B. J.; Nicholson, R. C.; Fairlie, W. D.; Por, S. B.; Robbins,

[84457] J. M.; Breit, S. N.: MIC-1, a novel macrophage inhibitory cytokine,

[84458] is a divergent member of the TGF-beta superfamily. Proc. Nat. Acad.

[84459] Sci. 94: 11514-11519, 1997.

[84460] 14765.

[84461] Bottner, M.; Laaff, M.; Schechinger, B.; Rappold, G.; Unsicker,

[84462] K.; Suter-Crazzolara, C.: Characterization of the rat, mouse, and

[84463] human genes of growth/differentiation factor-15/macrophage inhibiting

[84464] cytokine-1 (GDF-15/MIC-1). Gene 237: 105-111, 1999.

[84465] 14766.

[84466] Brown, D. A.; Breit, S. N.; Buring, J.; Fairlie, W. D.; Bauskin,

[84467] A. R.; Liu, T.; Ridker, P. M.: Concentration in plasma of macrophage

[84468] inhibitory cytokine-1 and risk of cardiovascular events in women:

[84469] a nested case-control study. Lancet 359: 2159-2163, 2002.

[84470] 14767.

[84471] Fairlie, W. D.; Russell, P. K.; Wu, W. M.; Moore, A. G.; Zhang,

[84472] H.-P.; Brown, P. K.; Bauskin, A. R.; Breit, S. N.: Epitope mapping

[84473] of the transforming growth factor-beta superfamily pro-tein, macrophage

[84474] inhibitory cytokine-1 (MIC-1): identification of at least five distinct

[84475] epitope specificities. Biochemistry 40: 65-73, 2001.

[84476] 14768.

[84477] Hromas, R.; Hufford, M.; Sutton, J.; Xu, D.; Li, Y.; Lu, L.:

PLAB,

[84478] a novel placental bone morphogenetic protein. Biochim. Biophys. Acta 1354:

[84479] 40-44, 1997.

[84480] 14769.

[84481] Moore, A. G.; Brown, D. A.; Fairlie, W. D.; Bauskin, A. R.; Brown,

[84482] P. K.; Munier, M. L. C.; Russell, P. K.; Salamonsen, L. A.; Wallace,

[84483] E. M.; Breit, S. N.: The transforming growth factor-beta superfamily

[84484] cytokine macrophage inhibitory cytokine-1 is present in high concentrations

[84485] in the serum of pregnant women. J. Clin. Endocr. Metab. 85: 4781-4788,

[84486] 2000.

[84487] 14770.

[84488] Paralkar, V. M.; Vail, A. L.; Grasser, W. A.; Brown, T. A.; Xu,

[84489] H.; Vukicevic, S.; Ke, H. Z.; Qi, H.; Owen, T. A.; Thompson, D. D.

[84490] : Cloning and characterization of a novel member of the

transforming
[84491] growth factor-beta/bone morphogenetic protein family. J. Biol. Chem. 273:
[84492] 13760-13767, 1998.

[84493] 14771.

[84494] Yokoyama-Kobayashi, M.; Saeki, M.; Sekine, S.; Kato, S.: Human
[84495] cDNA encoding a novel TGF-beta superfamily protein highly expressed
[84496] in placenta. J. Biochem. 122: 622-626, 1997.

[84497] 14772.

[84498] Conklin, D. C.; Rixon, M. W.; Kuestner, R. E.; Maurer, M. F.; Whitmore,
[84499] T. E.; Millar, R. P.: Cloning and gene expression of a novel human
[84500] ribonucleoprotein. Biochim. Biophys. Acta 1492: 465-469, 2000.
[84501] 14773.

[84502] Kataoka, N.; Yong, J.; Kim, V. N.; Velazquez, F.; Perkinson, R.
[84503] A.; Wang, F.; Dreyfuss, G.: Pre-mRNA splicing imprints

mRNA in the
[84504] nucleus with a novel RNA-binding protein that persists in the cytoplasm. Molec.
[84505] Cell 6: 673-682, 2000.

[84506] 14774.

[84507] Salicioni, A. M.; Xi. M.; Vanderveer, L. A.; Balsara, B.; Testa,
[84508] J. R.; Dunbrack, R. L., Jr.; Godwin, A. K.: Identification and structural
[84509] analysis of human RBM8A and RBM8B: two highly con-served RNA-binding
[84510] motif proteins that interact with OVCA1, a candidate tu-mor suppressor. Genomics 69:
[84511] 54-62, 2000.

[84512] 14775.

[84513] Zhao, X.-F.; Nowak, N. J.; Shows, T. B.; Aplan, P. D.: MAGOH interacts
[84514] with a novel RNA-binding protein. Genomics 63: 145-148, 2000.
[84515] 14776.

[84516] Fischle, W.; Emiliani, S.; Hendzel, M. J.; Nagase, T.; No-

mura,
[84517] N.; Voelter, W.; Verdin, E.: A new family of human histone deacetylases
[84518] related to Saccharomyces cerevisiae HDA1p. J. Biol. Chem. 274: 11713-11720,
[84519] 1999.

[84520] 14777.

[84521] Pazin, M. J.; Kadonaga, J. T.: What's up and down with histone
[84522] deacetylation and transcription? Cell 89: 325-328, 1997.

[84523] 14778.

[84524] Zhou, L.-J.; Tedder, T. F.: Human blood dendritic cells selectively
[84525] express CD83, a member of the immunoglobulin superfamily. J. Immun. 154:
[84526] 3821-3835, 1995.

[84527] 14779.

[84528] Zhou, L. J.; Schwarting, R.; Smith, H. M.; Tedder, T. F.: A novel
[84529] cell-surface molecule expressed by human interdigitating reticulum

[84530] cells, Langerhans cells, and activated lymphocytes is a new member

[84531] of the Ig superfamily. J. Immun. 149: 735-742, 1992.

[84532] 14780.

[84533] Bost-Usinger, L.; Chen, R. J.; Hillman, D.; Park, P.; Burn-side,

[84534] B.: Multiple kinesin family members expressed in teleost retina and

[84535] RPE include a novel C-terminal kinesin. Exp. Eye Res. 64: 781-794,

[84536] 1997.

[84537] 14781.

[84538] Hoang, E. H.; Whitehead, J. L.; Dose, A. C.; Burnside, B.: Cloning

[84539] of a novel C-terminal kinesin (KIFC3) that maps to human chromosome

[84540] 16q13-q21 and thus is a candidate gene for Bardet-Biedl syndrome. Genomics 52:

[84541] 219-222, 1998.

[84542] 14782.

[84543] Kiss, H.; Kedra, D.; Yang, Y.; Kost-Alimova, M.; Kiss, C.;

O'Brien,
[84544] K. P.; Fransson, I.; Klein, G.; Imreh, S.; Dumanski, J. P.: A novel
[84545] gene containing LIM domains (LIMD1) is located within the common eliminated
[84546] region 1 (C3CER1) in 3p21.3. Hum. Genet. 105: 552-559, 1999.
[84547] 14783.

[84548] Szeles, A.; Yang, Y.; Sandlund, A. M.; Kholodnyuck, I.; Kiss, H.;
[84549] Kost-Alimova, M.; Zabarovsky, E. R.; Stanbridge, E.; Klein, G.; Imreh,
[84550] S.: Human/mouse microcell hybrid based on elimination test reduces
[84551] the putative tumor suppressor region at 3p21.3 to 1.6 cM. Genes Chromosomes
[84552] Cancer 20: 329-336, 1997.

[84553] 14784.

[84554] Norman, D. A. M.; Barton, P. J. R.: Isolation, sequence, and chromosomal
[84555] localisation of the human I(kappa)BR gene. Ann. Hum. Genet. 64:

[84556] 15-23, 2000.

[84557] 14785.

[84558] Ray, P.; Zhang, D.-H.; Elias, J. A.; Ray, A.: Cloning of a dif-
ferentially

[84559] expressed I-kappa-B-related protein. J. Biol. Chem. 270:
10680-10685,

[84560] 1995.

[84561] 14786.

[84562] Lovering, R.; Trowsdale, J.: A gene encoding 22 highly re-
lated

[84563] zinc fingers is expressed in lymphoid cell lines. Nucleic
Acids Res. 19:

[84564] 2921-2928, 1991.

[84565] 14787.

[84566] Uchiumi, T.; Hinoshita, E.; Haga, S.; Nakamura, T.; Tanaka,
T.;

[84567] Toh, S.; Furukawa, M.; Kawabe, T.; Wada, M.; Kagotani, K.;
Okumura,

[84568] K.; Kohno, K.; Akiyama, S.; Kuwano, M.: Isolation of a
novel human

[84569] canalicular multispecific organic anion transporter,

cMOAT2/MRP3,

[84570] and its expression in cisplatin-resistant cancer cells with decreased

[84571] ATP-dependent drug transport. Biochem. Biophys. Res. Commun. 252:

[84572] 103-110, 1998.

[84573] 14788.

[84574] Deng, X.; Ito, T.; Carr, B.; Mumby, M.; May, W. S., Jr.: Re-versible

[84575] phosphorylation of Bcl2 following interleukin 3 or bryo-statin 1 is

[84576] mediated by direct interaction with protein phosphatase 2A. J. Biol.

[84577] Chem. 273: 34157-34163, 1998.

[84578] 14789.

[84579] Fujigasaki, H.; Verma, I. C.; Camuzat, A.; Margolis, R. L.; Zander,

[84580] C.; Lebre, A.-S.; Jamot, L.; Saxena, R.; Anand, I.; Holmes, S. E.;

[84581] Ross, C. A.; Durr, A.; Brice, A.: SCA12 is a rare locus for autosomal

[84582] dominant cerebellar ataxia: a study of an Indian family.

Ann. Neurol. 49:

[84583] 117-121, 2001.

[84584] 14790.

[84585] Holmes, S. E.; O'Hearn, E. E.; McInnis, M. G.; Gorelick-Feldman,

[84586] D. A.; Kleiderlein, J. J.; Callahan, C.; Kwak, N. G.; Ingersoll-Ashworth,

[84587] R. G.; Sherr, M.; Sumner, A. J.; Sharp, A. H.; Ananth, U.; Seltzer,

[84588] W. K.; Boss, M. A.; Vieria-Saecker, A.-M.; Epplen, J. T.; Riess, O.;

[84589] Ross, C. A.; Margolis, R. L.: Expansion of a novel CAG trin-ucleotide

[84590] repeat in the 5-prime region of PPP2R2B is associated with SCA12.

[84591] (Letter) Nature Genet. 23: 391-392, 1999.

[84592] 14791.

[84593] Mayer, R. E.; Hendrix, P.; Cron, P.; Matthies, R.; Stone, S. R.;
[84594] Goris, J.; Merlevede, W.; Hofsteenge, J.; Hemmings, B. A.: Structure
[84595] of the 55-kDa regulatory subunit of protein phosphatase 2A: evidence
[84596] for a neuronal-specific isoform. Biochemistry 30: 3589-3597, 1991.
[84597] 14792.

[84598] Mayer-Jaekel, R. E.; Ohkura, H.; Gomes, R.; Sunkel, C. E.; Baumgartner,
[84599] S.; Hemmings, B. A.; Glover, D. M.: The 55 kd regulatory subunit
[84600] of Drosophila protein phosphatase 2A is required for anaphase. Cell 72:
[84601] 621-633, 1993.

[84602] 14793.

[84603] Millward, T. A.; Zolnierowicz, S.; Hemmings, B. A.: Regula-
[84604] tion of protein kinase cascades by protein phosphatase 2A. Trends Biochem.
[84605] Sci. 24: 186-191, 1999.

[84606] 14794.

[84607] Santoro, M. F.; Annand, R. R.; Robertson, M. M.; Peng, Y.-W.; Brady,

[84608] M. J.; Mankovich, J. A.; Hackett, M. C.; Ghayur, T.; Walter, G.; Wong,

[84609] W. W.; Giegel, D. A.: Regulation of protein phosphatase 2A activity

[84610] by caspase-3 during apoptosis. J. Biol. Chem. 273: 13119-13128,

[84611] 1998.

[84612] 14795.

[84613] Schols, L.; Amoiridis, G.; Buttner, T.; Przuntek, H.; Epplen, J.

[84614] T.; Riess, O.: Autosomal dominant cerebellar ataxia: phe-notypic differences

[84615] in genetically defined subtypes? Ann. Neurol. 42: 924-932, 1997.

[84616] 14796.

[84617] Sontag, E.; Nunbhakdi-Craig, V.; Lee, G.; Brandt, R.; Kamibayashi,

[84618] C.; Kuret, J.; White, C. L., III; Mumby, M. C.; Bloom, G. S.: Molecular

[84619] interactions among protein phosphatase 2A, tau, and microtubules. J.

[84620] Biol. Chem. 274: 25490-25498, 1999.

[84621] 14797.

[84622] Almeida, R.; Levery, S. B.; Mandel, U.; Kresse, H.; Schwientek,

[84623] T.; Bennett, E. P.; Clausen, H.: Cloning and expression of a proteoglycan

[84624] UDP-galactose:beta-xylose beta-1,4-galactosyltransferase I: a seventh

[84625] member of the human beta4-galactosyltransferase gene family. J. Biol.

[84626] Chem. 274: 26165-26171, 1999.

[84627] 14798.

[84628] Kuroiwa, A.; Matsuda, Y.; Okajima, T.; Furukawa, K.: Assignment

[84629] of human xylosylprotein beta-1,4-galactosyltransferase gene (B4GALT7)

[84630] to human chromosome 5q35.2-q35.3 by in situ hybridization. Cytogenet.

[84631] Cell Genet. 89: 8-9, 2000.

[84632] 14799.

[84633] Bruhn, S. L.; Pil, P. M.; Essigmann, J. M.; Housman, D. E.; Lippard,

[84634] S. J.: Isolation and characterization of human cDNA clones encoding

[84635] a high mobility group box protein that recognizes structural distortions

[84636] to DNA caused by binding of the anticancer agent cisplatin. Proc.

[84637] Nat. Acad. Sci. 89: 2307-2311, 1992.

[84638] 14800.

[84639] Orphanides, G.; LeRoy, G.; Chang, C.-H.; Luse, D. S.; Reinberg,

[84640] D.: FACT, a factor that facilitates transcript elongation through

[84641] nucleosomes. Cell 92: 105-116, 1998.

[84642] 14801.

[84643] Endele, S.; Fuhry, M.; Pak, S. J.; Zabel, B. U.; Winterpacht, A.

[84644] : LETM1, a novel gene encoding a putative EF-hand Ca(2+)-binding protein,

[84645] flanks the Wolf-Hirschhorn syndrome (WHS) critical region and is deleted

[84646] in most WHS patients. Genomics 60: 218-225, 1999.

[84647] 14802.

[84648] Taki, T.; Kano, H.; Taniwaki, M.; Sako, M.; Yanagisawa, M.; Hayashi,

[84649] Y.: AF5q31, a newly identified AF4-related gene, is fused to MLL

[84650] in infant acute lymphoblastic leukemia with ins(5;11)(q31;q31q23). Proc.

[84651] Nat. Acad. Sci. 96: 14535-14540, 1999.

[84652] 14803.

[84653] Caira, F.; Antonson, P.; Pelto-Huikko, M.; Treuter, E.; Gustafsson,

[84654] J.-A.: Cloning and characterization of RAP250, a novel nuclear receptor

[84655] coactivator. J. Biol. Chem. 275: 5308-5317, 2000.

[84656] 14804.

[84657] Guan, X. Y.; Xu, J.; Anzick, S. L.; Zhang, H.; Trent, J. M.; Meltzer,

[84658] P. S.: Hybrid selection of transcribed sequences from mi-

crodissected

[84659] DNA: isolation of genes within amplified region at 20q11-q13.2 in

[84660] breast cancer. Cancer Res. 56: 3446-3450, 1996.

[84661] 14805.

[84662] Ko, L.; Cardona, G. R.; Chin, W. W.: Thyroid hormone receptor-binding

[84663] protein, an LXXLL motif-containing protein, functions as a general

[84664] coactivator. Proc. Nat. Acad. Sci. 97: 6212-6217, 2000.

[84665] 14806.

[84666] Lee, S.-K.; Anzick, S. L.; Choi, J.-E.; Bubendorf, L.; Guan, X.-Y.;

[84667] Jung, Y.-K.; Kallioniemi, O. P.; Kononen, J.; Trent, J. M.; Azorsa,

[84668] D.; Jhun, B.-H.; Cheong, J. H.; Lee, Y. C.; Meltzer, P. S.; Lee, J.

[84669] W.: A nuclear factor, ASC-2, as a cancer-amplified transcriptional

[84670] coactivator essential for ligand-dependent transactivation by nuclear

[84671] receptors in vivo. J. Biol. Chem. 274: 34283-34293, 1999.

[84672] 14807.

[84673] Mahajan, M. A.; Samuels, H. H: A new family of nuclear receptor

[84674] coregulators that integrate nuclear receptor signaling through CREB-binding

[84675] protein. Molec. Cell. Biol. 20: 5048-5063, 2000.

[84676] 14808.

[84677] Zhu, Y.; Kan, L.; Qi, C.; Kanwar, Y. S.; Yeldandi, A. V.; Rao,

[84678] M. S.; Reddy, J. K.: Isolation and characterization of peroxisome

[84679] proliferator-activated receptor (PPAR) interacting protein (PRIP)

[84680] as a coactivator for PPAR. J. Biol. Chem. 275: 13510-13516, 2000.

[84681] 14809.

[84682] Glockner, G.; Scherer, S.; Schattevoy, R.; Boright, A.; Weber,

[84683] J.; Tsui, L.-C.; Rosenthal, A.: Large-scale sequencing of two regions

[84684] in human chromosome 7q22: analysis of 650 kb of genomic sequence around

[84685] the EPO and CUTL1 loci reveals 17 genes. Genome Res. 8: 1060-1073,

[84686] 1998.

[84687] 14810.

[84688] Yokouchi, M.; Suzuki, R.; Masuhara, M.; Komiya, S.; Inoue, A.;

[84689] Yoshimura, A.: Cloning and characterization of APS, an adaptor molecule

[84690] containing PH and SH2 domains that is tyrosine phosphorylated upon

[84691] B-cell receptor stimulation. Oncogene 15: 7-15, 1997.

[84692] 14811.

[84693] Still, I. H.; Hamilton, M.; Vince, P.; Wolfman, A.; Cowell, J.

[84694] K.: Cloning of TACC1, an embryonically expressed, potentially transforming

[84695] coiled coil containing gene, from the 8p11 breast cancer amplicon. Oncogene 18:

[84696] 4032-4038, 1999.

[84697] 14812.

[84698] Burmester, T.; Weich, B.; Reinhardt, S.; Hankeln, T.: A vertebrate

[84699] globin expressed in the brain. Nature 407: 520-523, 2000.

[84700] 14813.

[84701] Moens, L.; Dewilde, S.: Globins in the brain. Nature 407: 461-462,

[84702] 2000.

[84703] 14814.

[84704] Cadigan, K. M.; Nusse, R.: Wnt signaling: a common theme in animal

[84705] development. Genes Dev. 11: 3286-3305, 1997.

[84706] 14815.

[84707] Parr, B. A.; Shea, M. J.; Vassileva, G.; McMahon, A. P.: Mouse

[84708] Wnt genes exhibit discrete domains of expression in the early embryonic

[84709] CNS and limb buds. Development 119: 247-261, 1993.

[84710] 14816.

[84711] Rankin, J.; Strachan, T.; Lako, M.; Lindsay, S.: Partial cloning

[84712] and assignment of WNT6 to human chromosome band 2q35 by in situ hybridization. Cytogenet.

[84713] Cell Genet. 84: 50-52, 1999.

[84714] 14817.

[84715] Vainio, S.; Parr, B.; McMahon, A.: Sexually dimorphic de-
velopment
[84716] of the mammalian reproductive system is regulated by
WNT-signaling.
[84717] (Abstract) Molec. Biol. Cell (suppl.) 8: 349a only, 12/1997.

[84718] 14818.

[84719] Ann, K.; Kowalchyk, J. A.; Loyet, K. M.; Martin, T. F. J.:
Novel
[84720] Ca(2+)-binding protein (CAPS) related to UNC-31 required
for Ca(2+)-activated
[84721] exocytosis. J. Biol. Chem. 272: 19637-19640, 1997.

[84722] 14819.

[84723] Berwin, B.; Floor, E.; Martin, T. F. J.: CAPS (mammalian
UNC-31)
[84724] protein localizes to membranes involved in dense-core
vesicle exocytosis. Neuron 21:
[84725] 137-145, 1998.

[84726] 14820.

[84727] Hirosawa, M.; Nagase, T.; Ishikawa, K.; Kikuno, R.; No-
mura, N.;
[84728] Ohara, O.: Characterization of cDNA clones selected by
the GeneMark
[84729] analysis from size-fractionated cDNA libraries from hu-
man brain. DNA
[84730] Res. 6: 329-336, 1999.

[84731] 14821.

[84732] Walent, J. H.; Porter, B. W.; Martin, T. F. J.: A novel 145 kd

[84733] brain cytosolic protein reconstitutes Ca(2+)-regulated se-
cretion in
[84734] permeable neuroendocrine cells. Cell 70: 765-775, 1992.

[84735] 14822.

[84736] Veldhuisen, B.; Spruit, L.; Dauwerse, H. G.; Breuning, M.
H.; Peters,
[84737] D. J.: Genes homologous to the autosomal dominant poly-
cystic kidney
[84738] disease genes (PKD1 and PKD2). Europ. J. Hum. Genet. 7:
860-872,
[84739] 1999.

[84740] 14823.

[84741] Hughes, J.; Ward, C. J.; Aspinwall, R.; Butler, R.; Harris, P.

[84742] C.: Identification of a human homologue of the sea urchin receptor

[84743] for egg jelly: a polycystic kidney disease-like protein. Hum. Molec.

[84744] Genet. 8: 543-549, 1999.

[84745] 14824.

[84746] Feinberg, H.; Mitchell, D. A.; Drickamer, K.; Weis, W. I.: Structural

[84747] basis for selective recognition of oligosaccharides by DC-SIGN and

[84748] DC-SIGNR. Science 294: 2163-2166, 2001.

[84749] 14825.

[84750] Xia, J.; Liu, C.; Tang, B.; Pan, Q.; Huang, L.; Dai, H.; Zhang,

[84751] B.; Xie, W.; Hu, D.; Zheng, D.; Shi, X.; Wang, D.; Xia, K.; Yu, K.;

[84752] Liao, X.; Feng, Y.; Yang, Y.; Xiao, J.; Xie, D.; Huang, J.: Mu-tations

[84753] in the gene encoding gap junction protein beta-3 associ-ated with autosomal

[84754] dominant hearing impairment. Nature Genet. 20:

370-373, 1998. Note:

[84755] Erratum: Nature Genet.:21: 241 only, 1999.

[84756] 14826.

[84757] Mueller, L.; Cordes, V. C.; Bischoff, F. R.; Ponstingl, H.: Human

[84758] RanBP3, a group of nuclear RanGTP binding proteins. FEBS Lett. 427:

[84759] 330-336, 1998.

[84760] 14827.

[84761] Rasooly, R. S.: Personal Communication. Baltimore, Md. 11/30/1998.

[84762] 14828.

[84763] Good, P.; Yoda, A.; Sakakibara, S.; Yamamoto, A.; Imai, T.; Sawa,

[84764] H.; Ikeuchi, T.; Tsuji, S.; Satoh, H.; Okano, H.: The human Musashi

[84765] homolog 1 (MSI1) gene encoding the homologue of Musashi/Nrp-1, a neural

[84766] RNA-binding protein putatively expressed in CNS stem cells and neural

[84767] progenitor cells. Genomics 52: 382-384, 1998.

[84768] 14829.

[84769] Crackower, M. A.; Sinasac, D. S.; Xia, J.; Motoyama, J.; Prochazka,

[84770] M.; Rommens, J. M.; Scherer, S. W.; Tsui, L.-C.: Cloning and characterization

[84771] of two cytoplasmic dynein intermediate chain genes in mouse and human. Genomics 55:

[84772] 257-267, 1999.

[84773] 14830.

[84774] Rodrigues Pereira, R.; Scholte, H. R.; Luyt-Houwen, I. E. M.; Vaandrager-Verduin,

[84775] M. H. M.: Cardiomyopathy associated with carnitine loss in kidneys

[84776] and small intestine. Europ. J. Pediat. 148: 193-197, 1988.

[84777] 14831.

[84778] Scholte, H. R.; Rodrigues Pereira, R.; de Jonge, P. C.; Luyt-Houwen,

[84779] I. E. M.; Hedwig, M.; Verduin, M.; Ross, J. D.: Primary carnitine

[84780] deficiency. J. Clin. Chem. Clin. Biochem. 28: 351-357, 1990.

[84781] 14832.

[84782] Tang, N. L.; Hui, J.; Law, L. K.; To, K. F.; Ruiter, J. P.; IJlst,

[84783] L.; Wanders, R. J.; Ho, C. S.; Fok, T. F.; Yuen, P. M.; Hjelm, N.

[84784] M.: Primary plasmalemmal carnitine transporter defect manifested

[84785] with dicarboxylic aciduria and impaired fatty acid oxidation. J.

[84786] Inherit. Metab. Dis. 21: 423-425, 1998.

[84787] 14833.

[84788] Tang, N. L. S.; Ganapathy, V.; Wu, X.; Hui, J.; Seth, P.; Yuen,

[84789] P. M. P.; Fok, T. F.; Hjelm, N. M.: Mutations of OCTN2, an organic

[84790] cation/carnitine transporter, lead to deficient cellular carnitine

[84791] uptake in primary carnitine deficiency. Hum. Molec. Genet. 8: 655-660,

[84792] 1999.

[84793] 14834.

[84794] Vaz, F. M.; Scholte, H. R.; Ruiter, J.; Hussaarts-Odijk, L. M.;

[84795] Rodrigues Pereira, R.; Schweitzer, S.; de Klerk, J. B. C.; Waterham,
[84796] H. R.; Wanders, R. J. A.: Identification of two novel mutations in
[84797] OCTN2 of three patients with systemic carnitine deficiency. Hum.
[84798] Genet. 105: 157-161, 1999.

[84799] 14835.

[84800] Wang, Y.; Taroni, F.; Garavaglia, B.; Longo, N.: Functional analysis
[84801] of mutations in the OCTN2 transporter causing primary carnitine deficiency:
[84802] lack of genotype-phenotype correlation. Hum. Mutat. 16: 401-407,
[84803] 2000.

[84804] 14836.

[84805] Wang, Y.; Ye, J.; Ganapathy, V.; Longo, N.: Mutations in the
[84806] organic cation/carnitine transporter OCTN2 in primary carnitine deficiency. Proc.
[84807] Nat. Acad. Sci. 96: 2356-2360, 1999.

[84808] 14837.

[84809] Wu, X.; Prasad, P. D.; Leibach, F. H.; Ganapathy, V.: cDNA sequence,

[84810] transport function, and genomic organization of human OCTN2, a new

[84811] member of the organic cation transporter family. Biochem. Biophys.

[84812] Res. Commun. 246: 589-595, 1998.

[84813] 14838.

[84814] Hart, M. J.; Callow, M. G.; Souza, B.; Polakis, P.: IQGAP1, a

[84815] calmodulin-binding protein with a rasGAP-related do- main, is a potential

[84816] effector for cdc42Hs. EMBO J. 15: 2997-3005, 1996.

[84817] 14839.

[84818] Sugimoto, N.; Imoto, I.; Fukuda, Y.; Kurihara, N.; Kuroda, S.;

[84819] Tanigami, A.; Kaibuchi, K.; Kamiyama, R.; Inazawa, J.: IQ- GAP1, a

[84820] negative regulator of cell-cell adhesion, is upregulated by gene amplification

[84821] at 15q26 in gastric cancer cell lines HSC39 and 40A. J.

Hum. Genet. 46:
[84822] 21-25, 2001.

[84823] 14840.

[84824] Weissbach, L.; Settleman, J.; Kalady, M. F.; Snijders, A. J.; Murthy,
[84825] A. E.; Yan, Y.-X.; Bernards, A.: Identification of a human RasGAP-related
[84826] protein containing calmodulin-binding motifs. J. Biol. Chem. 269:
[84827] 20517-20521, 1994.

[84828] 14841.

[84829] Brocker, F.; Bardenheuer, W.; Vieten, L.; Julicher, K.; Werner,
[84830] N.; Marquitan, G.; Michael, D.; Opalka, B.; Schutte, J.: As-signment
[84831] of human filamin gene FLNB to human chromosome band 3p14.3 and identification
[84832] of YACs containing the complete FLNB transcribed region. Cytogenet.
[84833] Cell Genet. 85: 267-268, 1999.

[84834] 14842.

[84835] Leedman, P. J.; Faulkner-Jones, B.; Cram, D. S.; Harrison, P. J.;

[84836] West, J.; O'Brien, E.; Simpson, R.; Coppel, R. L.; Harrison, L. C.

[84837] : Cloning from the thyroid of a protein related to actin binding protein

[84838] that is recognized by Graves disease immunoglobulins. Proc. Nat.

[84839] Acad. Sci. 90: 5994-5998, 1993.

[84840] 14843.

[84841] Harrington, J. J.; Lieber, M. R.: Functional domains within FEN-1

[84842] and RAD2 define a family of structure-specific endonucleases: implications

[84843] for nucleotide excision repair. Genes Dev. 8: 1344-1355, 1994.

[84844] 14844.

[84845] Dahl, M. R.; Thiel, S.; Matsushita, M.; Fujita, T.; Willis, A.

[84846] C.; Christensen, T.; Vorup-Jensen, T.; Jensenius, J. C.: MASP-3 and

[84847] its association with distinct complexes of the mannan-binding lectin

[84848] complement activation pathway. Immunity 15: 127-135, 2001.

[84849] 14845.

[84850] Endo, Y.; Sato, T.; Matsushita, M.; Fujita, T.: Exon structure

[84851] of the gene encoding the human mannose-binding protein-associated

[84852] serine protease light chain: comparison with complement C1r and C1s

[84853] genes. Int. Immun. 8: 1355-1358, 1996.

[84854] 14846.

[84855] Matsushita, M.; Thiel, S.; Jensenius, J. C.; Terai, I.; Fujita,

[84856] T.: Proteolytic activities of two types of mannose-binding lectin-associated

[84857] serine protease. J. Immun. 165: 2637-2642, 2000.

[84858] 14847.

[84859] Sato, T.; Endo, Y.; Matsushita, M.; Fujita, T.: Molecular characterization

[84860] of a novel serine protease involved in activation of the complement

[84861] system by mannose-binding protein. Int. Immun. 6:

665-669, 1994.

[84862] 14848.

[84863] Takada, F.; Seki, N.; Matsuda, Y.; Takayama, Y.; Kawakami, M.:

[84864] Localization of the genes for the 100-kDa complement-acti-vating components

[84865] of Ra-reactive factor (CRARF and Crarf) to human 3q27-q28 and mouse

[84866] 16B2-B3. Genomics 25: 757-759, 1995.

[84867] 14849.

[84868] Takada, F.; Takayama, Y.; Hatsuse, H.; Kawakami, M.: A new member

[84869] of the C1s family of complement proteins found in a bac-tericidal factor,

[84870] Ra-reactive factor, in human serum. Biochem. Biophys. Res. Commun. 196:

[84871] 1003-1009, 1993.

[84872] 14850.

[84873] Takayama, Y.; Takada, F.; Nowatari, M.; Kawakami, M.; Matsu-ura,

[84874] N.: Gene structure of the P100 serine-protease compo-

nent of the human
[84875] Ra-reactive factor. Molec. Immun. 36: 505-514, 1999.

[84876] 14851.

[84877] Hansen, L.; Urioste, S.; Petersen, H. V.; Jensen, J. N.; Eiberg,
[84878] H.; Barbetti, F.; Serup, P.; Hansen, T.; Pedersen, O.: Mis-sense mutations
[84879] in the human insulin promoter factor-1 gene and their re-lation to
[84880] maturity-onset diabetes of the young and late-onset type 2 diabetes
[84881] mellitus in Caucasians. J. Clin. Endocr. Metab. 85: 1323-1326, 2000.
[84882] 14852.

[84883] Johansson, K. A.; Grapin-Botton, A.: Development and diseases
[84884] of the pancreas. Clin. Genet. 62: 14-23, 2002.

[84885] 14853.

[84886] Jonnson, J.; Carlsson, L.; Edlund, T.; Edlund, H.: Insulin-promoter-factor
[84887] 1 is required for pancreas development in mice. Nature

371: 606-609,

[84888] 1994.

[84889] 14854.

[84890] Leonard, J.; Peers, B.; Johnson, T.; Ferreri, I.; Lee, K.;
Montminy,

[84891] M. R.: Characterization of somatostatin transactivating
factor-1,

[84892] a novel homeobox factor that stimulates somatostatin ex-
pression in

[84893] pancreatic islet cells. Molec. Endocr. 7: 1275-1283, 1993.

[84894] 14855.

[84895] Macfarlane, W. M.; Frayling, T. M.; Ellard, S.; Evans, J. C.;
Allen,

[84896] L. I. S.; Bulman, M. P.; Ayres, S.; Shepherd, M.; Clark, P.;
Millward,

[84897] A.; Demaine, A.; Wilkin, T.; Docherty, K.; Hattersley, A. T.:
Missense

[84898] mutations in the insulin promoter factor-1 gene predis-
pose to type

[84899] 2 diabetes. J. Clin. Invest. 104: R33-R39, 1999.

[84900] 14856.

[84901] Miller, C. P.; McGehee, R. E., Jr.; Habener, J. F.: IDX-1: a

[84902] new homeodomain transcription factor expressed in rat pancreatic islets

[84903] and duodenum that transactivates the somatostatin gene. EMBO J. 13:

[84904] 1145-1156, 1994.

[84905] 14857.

[84906] Ohlsson, H.; Karlsson, K.; Edlund, T.: IPF1, a home-odomain-containing

[84907] transactivator of the insulin gene. EMBO J. 12: 4251-4259, 1993.

[84908] 14858.

[84909] Sharma, S.; Jhala, U. S.; Johnson, T.; Ferreri, K.; Leonard, J.;

[84910] Montminy, M.: Hormonal regulation of an islet-specific enhancer in

[84911] the pancreatic homeobox gene STF-1. Molec. Cell. Biol. 17: 2598-2604,

[84912] 1997.

[84913] 14859.

[84914] Stoffel, M.; Stein, R.; Wright, C. V. E.; Espinosa, R., III; Le

[84915] Beau, M. M.; Bell, G. I.: Localization of human homeodomain transcription
[84916] factor insulin promoter factor 1 (IPF1) to chromosome band 13q12.1. Genomics 28:
[84917] 125-126, 1995.

[84918] 14860.

[84919] Stoffers, D. A.; Ferrer, J.; Clarke, W. L.; Habener, J. F.: Early-onset
[84920] type-II diabetes mellitus (MODY4) linked to IPF1. (Letter) Nature
[84921] Genet. 17: 138-141, 1997.

[84922] 14861.

[84923] Stoffers, D. A.; Stanojevic, V.; Habener, J. F.: Insulin promoter
[84924] factor-1 gene mutation linked to early-onset type 2 diabetes mellitus
[84925] directs expression of a dominant negative isoprotein. J. Clin. Invest. 102:
[84926] 232-241, 1998.

[84927] 14862.

[84928] Watada, H.; Kajimoto, Y.; Kaneto, H.; Matsuoka, T.; Fuji-

tani,

[84929] Y.; Miyazaki, J.; Yamasaki, Y.: Involvement of the homeodomain-containing

[84930] transcription factor PDX-1 in islet amyloid polypeptide gene transcription. Biochem.

[84931] Biophys. Res. Commun. 229: 746-751, 1996.

[84932] 14863.

[84933] Wright, N. M.; Metzger, D. L.; Borowitz, S. M.; Clarke, W. L.

[84934] : Permanent neonatal diabetes mellitus and pancreatic exocrine insufficiency

[84935] resulting from congenital pancreatic agenesis. Am. J. Dis. Child. 147:

[84936] 607-609, 1993.

[84937] 14864.

[84938] Ashford, M. L. J.; Bond, C. T.; Blair, T. A.; Adelman, J. P.:

[84939] Cloning and functional expression of a rat heart KATP channel. Nature 370:

[84940] 456-459, 1994.

[84941] 14865.

[84942] Bond, C. T.; Pessia, M.; Xia, X.-M.; Lagrutta, A.; Ka-

vanaugh, M.

[84943] P.; Adelman, J. P.: Cloning and expression of a family of inward

[84944] rectifier potassium channels. Receptors Channels 2: 183-191, 1994.

[84945] 14866.

[84946] Tucker, S. J.; James, M. R.; Adelman, J. P.: Assignment of K(ATP)-1,

[84947] the cardiac ATP-sensitive potassium channel gene (KCNJ5), to human

[84948] chromosome 11q24. Genomics 28: 127-128, 1995.

[84949] 14867.

[84950] Wickman, K.; Seldin, M. F.; Gendler, S. J.; Clapham, D. E.: Partial

[84951] structure, chromosome localization, and expression of the mouse Girk4

[84952] gene. Genomics 40: 395-401, 1997.

[84953] 14868.

[84954] Bange, F.-C.; Vogel, U.; Flohr, T.; Kiekenbeck, M.; De-

necke, B.;

[84955] Boettger, E. C.: IFP 35 is an interferon-induced leucine

zipper protein
[84956] that undergoes interferon-regulated cellular redistribution. J. Biol.
[84957] Chem. 269: 1091-1098, 1994.

[84958] 14869.

[84959] Brown, M. A.; Jones, K. A.; Nicolai, H.; Bonjardim, M.; Black,
[84960] D.; McFarlane, R.; De Jong, P.; Quirk, J. P.; Lehrach, H.; Solomon,
[84961] E.: Physical mapping, cloning, and identification of genes within
[84962] a 500-kb region containing BRCA1. Proc. Nat. Acad. Sci. 92: 4362-4366,
[84963] 1995.

[84964] 14870.

[84965] Smith, T. M.; Lee, M. K.; Szabo, C. I.; Jerome, N.; McEuen, M.;
[84966] Taylor, M.; Hood, L.; King, M.-C.: Complete genomic sequence and
[84967] analysis of 117 kb of human DNA containing the gene BRCA1. Genomic
[84968] Res. 6: 1029-1049, 1996.

[84969] 14871.

[84970] Khurana, T. S.; Engle, E. C.; Bennett, R. R.; Silverman, G. A.;

[84971] Selig, S.; Bruns, G. A. P.; Kunkel, L. M.: (CA) repeat poly-morphism

[84972] in the chromosome 18 encoded dystrophin-like protein. Hum. Molec.

[84973] Genet. 3: 841 only, 1994.

[84974] 14872.

[84975] Wunderle, V.; Ramkissoon, Y. D.; Kwok, C.; Korn, R. M.; King,

[84976] V. E.; Goodfellow, P. N.: Breakpoint break for consortium studying

[84977] adult polycystic kidney disease. Cell 77: 785-786, 1994.

[84978] 14873.

[84979] Eisenberg, I.; Avidan, N.; Potikha, T.; Hochner, H.; Chen, M.;

[84980] Olender, T.; Barash, M.; Shemesh, M.; Sadeh, M.; Grabov-Nardini, G.;

[84981] Shmilevich, I.; Friedmann, A.; Karpati, G.; Bradley, W. G.; Baumbach,

[84982] L.; Lancet, D.; Ben Asher, E.; Beckmann, J. S.; Argov, Z.; Mitrani-Rosenbaum,
[84983] S.: The UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase
[84984] gene is mutated in recessive hereditary inclusion body myopathy. Nature
[84985] Genet. 29: 83-87, 2001.

[84986] 14874.

[84987] Metzinger, L.; Blake, D. J.; Squier, M. V.; Anderson, L. V. B.;

[84988] Deconinck, A. E.; Nawrotzki, R.; Hilton-Jones, D.; Davies, K. E.:
[84989] Dystrobrevin deficiency at the sarcolemma of patients with muscular
[84990] dystrophy. Hum. Molec. Genet. 6: 1185-1191, 1997.

[84991] 14875.

[84992] Sadoulet-Puccio, H. M.; Feener, C. A.; Schaid, D. J.; Thi-bodeau,
[84993] S. N.; Michels, V. V.; Kunkel, L. M.: The genomic organiza-tion of
[84994] human dystrobrevin. Neurogenetics 1: 37-42, 1997.

[84995] 14876.

[84996] Sadoulet-Puccio, H. M.; Khurana, T. S.; Cohen, J. B.; Kunkel, L.
[84997] M.: Cloning and characterization of the human homologue of a dystrophin
[84998] related phosphoprotein found at the Torpedo electric organ post-synaptic
[84999] membrane. Hum. Molec. Genet. 5: 489-496, 1996.

[85000] 14877.

[85001] Yoshida, M.; Hama, H.; Ishikawa-Sakurai, M.; Imamura, M.; Mizuno,
[85002] Y.; Araishi, K.; Wakabayashi-Takai, E.; Noguchi, S.; Sasaoka, T.;
[85003] Ozawa, E.: Biochemical evidence for association of dystrobrevin with
[85004] the sarcoglycan-sarcospan complex as a basis for understanding sarcoglycanopathy. Hum.
[85005] Molec. Genet. 9: 1033-1040, 2000.

[85006] 14878.

[85007] Chae, Y.-J.; Chung, C.-E.; Kim, B.-J.; Lee, M.-H.; Lee, H.: The
[85008] gene encoding guanidinoacetate methyltransferase (GAMT) maps to human

[85009] chromosome 19 at band p13.3 and to mouse chromosome 10. Genomics 49:
[85010] 162-164, 1998.

[85011] 14879.

[85012] Isbrandt, D.; von Figura, K.: Cloning and sequence analysis of
[85013] human guanidinoacetate N-methyltransferase cDNA. Biochim. Biophys.
[85014] Acta 1264: 265-267, 1995.

[85015] 14880.

[85016] Schulze, A.; Hess, T.; Wevers, R.; Mayatepek, E.; Bachert, P.;
[85017] Marescau, B.; Knopp, M. V.; De Deyn, P. P.; Bremer, H. J.; Rating,
[85018] D.: Creatine deficiency syndrome caused by guanidinoacetate methyltransferase
[85019] deficiency: diagnostic tools for a new inborn error of metabolism. J.
[85020] Pediat. 131: 626-631, 1997.

[85021] 14881.

[85022] Stockler, S.; Holzbach, U.; Hanefeld, F.; Marquardt, I.;

Helms,
[85023] G.; Requart, M.; Hanicke, W.; Frahm, J.: Creatine deficiency in the
[85024] brain: a new, treatable inborn error of metabolism. Pediat. Res. 36:
[85025] 409-413, 1994.

[85026] 14882.

[85027] Stockler, S.; Isbrandt, D.; Hanefeld, F.; Schmidt, B.; von Figura,
[85028] K.: Guanidinoacetate methyltransferase deficiency: the first inborn
[85029] error of creatine metabolism in man. Am. J. Hum. Genet. 58: 914-922,
[85030] 1996.

[85031] 14883.

[85032] Verhoeven, N. M.; Guerand, W. S.; Struys, E. A.; Bouman, A. A.;
[85033] van der Knaap, M. S.; Jakobs, C.: Plasma creatinine assess-ment in
[85034] creatine deficiency: a diagnostic pitfall. J. Inherit. Metab. Dis. 23:
[85035] 835-840, 2000.

[85036] 14884.

[85037] Elsea, S. H.; Fritz, E.; Schoener-Scott, R.; Meyn, M. S.; Pa-
tel,
[85038] P. I.: Gene for topoisomerase III maps within the Smith-
Magenis syndrome
[85039] critical region: analysis of cell-cycle distribution and radi-
ation
[85040] sensitivity. Am. J. Med. Genet. 75: 104-108, 1998.

[85041] 14885.

[85042] Fritz, E.; Elsea, S. H.; Patel, P. I.; Meyn, M. S.: Overexpres-
sion
[85043] of a truncated human topoisomerase III partially corrects
multiple
[85044] aspects of the ataxia-telangiectasia phenotype. Proc. Nat.
Acad.
[85045] Sci. 94: 4538-4542, 1997.

[85046] 14886.

[85047] Chappell, T. G.; Welch, W. J.; Schlossman, D. M.; Palter, K.
B.;
[85048] Schlesinger, M. J.; Rothman, J. E.: Uncoating ATPase is a
member

[85049] of the 70 kilodalton family of stress proteins. Cell 45:
3-13, 1986.

[85050] 14887.

[85051] Dworniczak, B.; Mirault, M.-E.: Structure and expression
of a

[85052] human gene coding for a 71 kd heat shock 'cognate' pro-
tein. Nucleic

[85053] Acids Res. 15: 5181-5197, 1987.

[85054] 14888.

[85055] Tavaria, M.; Gabriele, T.; Anderson, R. L.; Mirault, M.-E.;
Baker,

[85056] E.; Sutherland, G.; Kola, I.: Localization of the gene encod-
ing the

[85057] human heat shock cognate protein, HSP73, to chromo-
some 11. Genomics 29:

[85058] 266-268, 1995.

[85059] 14889.

[85060] Ungewickell, E.: The 70-kd mammalian heat shock pro-
teins are structurally

[85061] and functionally related to the uncoating protein that re-
leases clathrin

[85062] triskelia from coated vesicles. EMBO J. 4: 3385-3391, 1985.

[85063] 14890.

[85064] Polymeropoulos, M. H.; Ide, S.; Soares, M. B.; Lennon, G. G.:

[85065] Sequence characterization and genetic mapping of the human VSNL1 gene,

[85066] a homologue of the rat visinin-like peptide RNVP1. Genomics 29:

[85067] 273-275, 1995.

[85068] 14891.

[85069] Camoretti-Mercado, B.; Forsythe, S. M.; LeBeau, M. M.; Espinosa,

[85070] R., III; Vieira, J. E.; Halayko, A. J.; Willadsen, S.; Kurtz, B.;

[85071] Ober, C.; Evans, G. A.; Thweatt, R.; Shapiro, S.; Niu, Q.; Qin, Y.;

[85072] Padrid, P. A.; Solway, J.: Expression and cytogenetic localization

[85073] of the human SM22 gene (TAGLN). Genomics 49: 452-457, 1998.

[85074] 14892.

[85075] Lawson, D.; Harrison, M.; Shapland, C.: Fibroblast transgelin
[85076] and smooth muscle SM22-alpha are the same protein, the expression
[85077] of which is down-regulated in many cell lines. Cell Motil. Cytoskeleton 38:
[85078] 250-257, 1997.

[85079] 14893.

[85080] Lees-Miller, J. P.; Heeley, D. H.; Smillie, L. B.; Kay, C. M.:

[85081] Isolation and characterization of an abundant and novel 22-kDa protein
[85082] (SM22) from chicken gizzard smooth muscle. J. Biol. Chem. 262: 2988-2993,
[85083] 1987.

[85084] 14894.

[85085] Nishida, W.; Kitami, Y.; Abe, M.; Hiwada, K.: Gene cloning and
[85086] nucleotide sequence of SM22-alpha from the chicken gizzard smooth
[85087] muscle. Biochem. Int. 23: 663-668, 1991.

[85088] 14895.

[85089] Shapland, C.; Hsuan, J. J.; Totty, N. F.; Lawson, D.: Purification

[85090] and properties of transgelin: a transformation and shape change sensitive

[85091] actin-gelling protein. J. Cell Biol. 121: 1065-1073, 1993.

[85092] 14896.

[85093] Shapland, C.; Lowings, P.; Lawson, D.: Identification of new actin-associated

[85094] polypeptides that are modified by viral transformation and changes

[85095] in cell shape. J. Cell Biol. 107: 153-161, 1988.

[85096] 14897.

[85097] Solway, J.; Seltzer, J.; Samaha, F. F.; Kim, S.; Alger, L. E.;

[85098] Niu, Q.; Morrisey, E. E.; Ip, H. S.; Parmacek, M. S.: Structure and

[85099] expression of a smooth muscle cell-specific gene, SM22-alpha. J.

[85100] Biol. Chem. 270: 13460-13469, 1995.

[85101] 14898.

[85102] Stanier, P.; Abu-Hayyeh, S.; Murdoch, J. N.; Eddleston, J.; Copp,

[85103] A. J.: Paralogous Sm22-alpha (Tagln) genes map to mouse chromosomes
[85104] 1 and 9: further evidence for a paralogous relationship. Genomics 51:
[85105] 144-147, 1998.

[85106] 14899.

[85107] Thweatt, R.; Lumpkin, C. K., Jr.; Goldstein, S.: A novel gene
[85108] encoding a smooth muscle protein is overexpressed in senescent human
[85109] fibroblasts. Biochem. Biophys. Res. Commun. 187: 1-7, 1992.
[85110] 14900.

[85111] Tamanini, F.; Bontekoe, C.; Bakker, C. E.; van Unen, L.; Anar,
[85112] B.; Willemsen, R.; Yoshida, M.; Galjaard, H.; Oostra, B. A.; Hoogeveen,
[85113] A. T.: Different targets for the fragile X-related proteins revealed
[85114] by their distinct nuclear localizations. Hum. Molec. Genet. 8: 863-869,
[85115] 1999.

[85116] 14901.

[85117] Tamanini, F.; Kirkpatrick, L. L.; Schonkeren, J.; van Unen, L.;

[85118] Bontekoe, C.; Bakker, C.; Nelson, D. L.; Galjaard, H.; Oostra, B.

[85119] A.; Hoogeveen, A. T.: The fragile X-related proteins FXR1P and FXR2P

[85120] contain a functional nucleolar-targeting signal equivalent to the

[85121] HIV-1 regulatory proteins. Hum. Molec. Genet. 9: 1487-1493, 2000.

[85122] 14902.

[85123] Radice, P.; Pensotti, V.; Jones, C.; Perry, H.; Pierotti, M. A.;

[85124] Tunnacliffe, A.: The human archain gene, ARCN1, has highly conserved

[85125] homologs in rice and Drosophila. Genomics 26: 101-106, 1995.

[85126] 14903.

[85127] Tunnacliffe, A.; van de Vrugt, H.; Pensotti, V.; Radice, P.: The

[85128] coatomer protein delta-COP, encoded by the archain gene, is conserved

[85129] across diverse eukaryotes. Mammalian Genome 7:
784-786, 1996.
[85130] 14904.
[85131] Hawtin, S. R.; Wesley, V. J.; Parslow, R. A.; Simms, J.; Miles,
[85132] A.; McEwan, K.; Wheatley, M.: A single residue (arg46) lo-
cated within
[85133] the N-terminus of the V(1a) vasopressin receptor is criti-
cal for binding
[85134] vasopressin but not peptide or nonpeptide antagonists.
Molec. Endocr. 16:
[85135] 600-609, 2002.
[85136] 14905.
[85137] Morel, A.; O'Carroll, A.-M.; Brownstein, M. J.; Lolait, S. J.:
[85138] Molecular cloning and expression of a rat V1a arginine
vasopressin
[85139] receptor. Nature 356: 523-526, 1992.
[85140] 14906.
[85141] Thibonnier, M.; Auzan, C.; Madhun, Z.; Wilkins, P.; Berti-
Mattera,
[85142] L.; Clauser, E.: Molecular cloning, sequencing, and func-
tional expression

[85143] of a cDNA encoding the human V1a vasopressin receptor. J. Biol. Chem. 269:
[85144] 3304-3310, 1994.

[85145] 14907.

[85146] Thibonnier, M.; Graves, M. K.; Wagner, M. S.; Auzan, C.; Clauser,
[85147] E.; Willard, H. F.: Structure, sequence, expression, and chromosomal
[85148] localization of the human V(1a) vasopressin receptor gene. Genomics 31:
[85149] 327-334, 1996.

[85150] 14908.

[85151] Young, L. J.; Nilsen, R.; Waymire, K. G.; MacGregor, G. R.; Insel,
[85152] T. R.: Increased affiliative response to vasopressin in mice expressing
[85153] the V(1A) receptor from a monogamous vole. Nature 400:
766-768,
[85154] 1999.

[85155] 14909.

[85156] Weber, B. H. F.; Brohm, M.; Stec, I.; Backe, J.; Caffier, H.:

[85157] A somatic truncating mutation in BRCA2 in a sporadic breast tumor. Am.

[85158] J. Hum. Genet. 59: 962-964, 1996.

[85159] 14910.

[85160] Huh, T.-L.; Kim, Y.-O.; Oh, I.-U.; Song, B. J.; Inazawa, J.: Assignment

[85161] of the human mitochondrial NAD(+)-specific isocitrate dehydrogenase

[85162] alpha subunit (IDH3A) gene to 15q25.1-q25.2 by in situ hybridization. Genomics 31:

[85163] 295-296, 1996.

[85164] 14911.

[85165] Kim, Y.-O.; Oh, I.-U.; Park, H.-S.; Jeng, J.; Song, B. J.; Huh,

[85166] T.-L.: Characterization of a cDNA clone for human NAD(+)-specific

[85167] isocitrate dehydrogenase alpha-subunit and structural comparison with

[85168] its isoenzymes from different species. Biochem. J. 308: 63-68, 1995.

[85169] 14912.

[85170] Amann, J.; Kidd, V. J.; Lahti, J. M.: Characterization of pu-

tative

[85171] human homologues of the yeast chromosome transmis-
sion fidelity gene,

[85172] CHL1. J. Biol. Chem. 272: 3823-3832, 1997.

[85173] 14913.

[85174] Amann, J.; Valentine, M.; Kidd, V. J.; Lahti, J. M.: Localiza-
tion

[85175] of Chl1-related helicase genes to human chromosome re-
gions 12p11 and

[85176] 12p13: similarity between parts of these genes and con-
served human

[85177] telomeric-associated DNA. Genomics 32: 260-265, 1996.

[85178] 14914.

[85179] Frank, S.; Werner, S.: The human homologue of the yeast
CHL1 gene

[85180] is a novel keratinocyte growth factor-regulated gene. J.
Biol. Chem. 271:

[85181] 24337-24340, 1996.

[85182] 14915.

[85183] Barnes, L. D.; Garrison, P. N.; Siprashvili, Z.; Guranowski,
A.;

[85184] Robinson, A. K.; Ingram, S. W.; Croce, C. M.; Ohta, M.; Huebner, K.

[85185] : Fhit, a putative tumor suppressor in humans, is a dinu-

cleoside 5-prime,5-triple

[85186] prime-P(1),P(3)-triphosphate hydrolase. Biochemistry 35: 11529-11535,

[85187] 1996.

[85188] 14916.

[85189] Bernar, J.; Funderburk, S. J.; Sparkes, R. S.: The inducible fragile

[85190] site on chromosome 3. (Letter) Hum. Genet. 66: 373 only, 1984.

[85191] 14917.

[85192] Rajan, J. V.; Wang, M.; Marquis, S. T.; Chodosh, L. A.: Brca2

[85193] is coordinately regulated with Brca1 during proliferation and differentiation

[85194] in mammary epithelial cells. Proc. Nat. Acad. Sci. 93: 13078-13083,

[85195] 1996.

[85196] 14918.

[85197] Rebbeck, T. R.; Lynch, H. T.; Neuhausen, S. L.; Narod, S. A.;

[85198] van't Veer, L.; Garber, J. E.; Evans, G.; Isaacs, C.; Daly, M. B.;

[85199] Matloff, E.; Olopade, O. I.; Weber, B. L. Prophylactic oophorectomy

[85200] in carriers of BRCA1 or BRCA2 mutations. New Eng. J. Med. 346: 1616-1622,

[85201] 2002.

[85202] 14919.

[85203] Risch, H. A.; McLaughlin, J. R.; Cole, D. E. C.; Rosen, B.; Bradley,

[85204] L.; Kwan, E.; Jack, E.; Vesprini, D. J.; Kuperstein, G.; Abra-

[85205] hamson, J. L. A.; Fan, I.; Wong, B.; Narod, S. A.: Prevalence and pen-

[85206] etrance of germline BRCA1 and BRCA2 mutations in a population series of 649

[85207] women with ovarian cancer. Am. J. Hum. Genet. 68: 700-710, 2001.

[85208] 14920.

[85209] Roa, B. B.; Boyd, A. A.; Volcik, K.; Richards, C. S.: Ashke-

nazi

[85210] Jewish population frequencies for common mutations in BRCA1 and BRCA2. Nature

[85211] Genet. 14: 185-187, 1996.

[85212] 14921.

[85213] Sarantaus, L.; Huusko, P.; Eerola, H.; Launonen, V.; Vehmanen,

[85214] P.; Rapakko, K.; Gillanders, E.; Syrjakoski, K.; Kainu, T.; Vahteristo,

[85215] P.; Krahe, R.; Paakkonen, K.; and 14 others: Multiple founder effects

[85216] and geographical clustering of BRCA1 and BRCA2 families in Finland. Europ.

[85217] J. Hum. Genet. 8: 757-763, 2000.

[85218] 14922.

[85219] Sarantaus, L.; Vahteristo, P.; Bloom, E.; Tamminen, A.; Unkila-Kallio,

[85220] L.; Butzow, R.; Nevanlinna, H. BRCA1 and BRCA2 mutations among

[85221] 233 unselected Finnish ovarian carcinoma patients. Europ. J. Hum.

[85222] Genet. 9: 424-430, 2001.

[85223] 14923.

[85224] Sharan, S. K.; Morimatsu, M.; Albrecht, U.; Lim, D.-S.; Regel,

[85225] E.; Dinh, C.; Sands, A.; Eichele, G.; Hasty, P.; Bradley, A.: Embryonic

[85226] lethality and radiation hypersensitivity mediated by Rad51 in mice

[85227] lacking Brca2. Nature 386: 804-810, 1997.

[85228] 14924.

[85229] Sigbjornsdottir, B. I.; Ragnarsson, G.; Agnarsson, B. A.; Huiping,

[85230] C.; Barkardottir, R. B.; Egilsson, V.; Ingvarsson, S.: Chro-mosome

[85231] 8p alterations in sporadic and BRCA2 999del5 linked breast cancer. J.

[85232] Med. Genet. 37: 342-347, 2000.

[85233] 14925.

[85234] Spain, B. H.; Larson, C. J.; Shihabuddin, L. S.; Gage, F. H.;

[85235] Verma, I. M.: Truncated BRCA2 is cytoplasmic: implica-tions for cancer-linked

[85236] mutations. Proc. Nat. Acad. Sci. 96: 13920-13925, 1999.

[85237] 14926.

[85238] Stratton, M. R.; Ford, D.; Neuhasen, S.; Seal, S.; Wooster, R.;

[85239] Friedman, L. S.; King, M.-C.; Egilsson, V.; Devilee, P.; Mc-Manus,

[85240] R.; Daly, P. A.; Smyth, E.; Ponder, B. A. J.; Peto, J.; Cannon-Albright,

[85241] L.; Easton, D. F.; Goldgar, D. E.: Familial male breast can-cer is

[85242] not linked to the BRCA1 locus on chromosome 17q. Na-ture Genet. 7:

[85243] 103-107, 1994.

[85244] 14927.

[85245] Struewing, J. P.; Hartge, P.; Wacholder, S.; Baker, S. M.; Berlin,

[85246] M.; McAdams, M.; Timmerman, M. M.; Brody, L. C.; Tucker, M. A.: The

[85247] risk of cancer associated with specific mutations of BRCA1 and BRCA2

[85248] among Ashkenazi Jews. New Eng. J. Med. 336: 1401-1408, 1997.

[85249] 14928.

[85250] Suzuki, A.; de la Pompa, J. L.; Hakem, R.; Elia, A.; Yoshida,

[85251] R.; Mo, R.; Nishina, H.; Chuang, T.; Wakeham, A.; Itie, A.; Koo, W.;

[85252] Billia, P.; Ho, A.; Fukumoto, M.; Hui, C. C.; Mak, T. W.: Brca2 is

[85253] required for embryonic cellular proliferation in the mouse. Genes

[85254] Dev. 11: 1242-1252, 1997.

[85255] 14929.

[85256] Szabo, C. I.; King, M.-C.: Population genetics of BRCA1 and BRCA2.

[85257] (Editorial) Am. J. Hum. Genet. 60: 1013-1020, 1997.

[85258] 14930.

[85259] Taillon-Miller, P.; Bauer-Sardina, I.; Zakeri, H.; Hillier, L.;

[85260] Mutch, D. G.; Kwok, P.-Y.: The homozygous complete hydatidiform mole:

[85261] a unique resource for genome studies. Genomics 46: 307-310, 1997.

[85262] 14931.

[85263] Tavtigian, S. V.; Simard, J.; Rommens, J.; Couch, F.; Shattuck-Eidens,

[85264] D.; Neuhausen, S.; Merajver, S.; Thorlacius, S.; Offit, K.; Stoppa-Lyonnet,

[85265] D.; Belanger, C.; Bell, R.; and 37 others: The complete BRCA2 gene

[85266] and mutations in chromosome 13q-linked kindreds. Nature Genet. 12:

[85267] 333-337, 1996.

[85268] 14932.

[85269] Teng, D. H.-F.; Bogden, R.; Mitchell, J.; Baumgard, M.; Bell,

[85270] R.; Berry, S.; David, T.; Ha, P. C.; Kehrer, R.; Jammulapati, S.;

[85271] Chen, Q.; Offit, K.; Skolnick, M. H.; Tavtigian, S. V.; Jhanwar, S.;

[85272] Swedlund, B.; Wong, A. K. C.; Kamb, A.: Low incidence of BRCA2 mutations

[85273] in breast carcinoma and other cancers. Nature Genet. 13: 241-244,

[85274] 1996.

[85275] 14933.

[85276] Tesoriero, A.; Andersen, C.; Southey, M.; Somers, G.; McKay, M.;

[85277] Armes, J.; McCredie, M.; Giles, G.; Hopper, J. L.; Venter, D.: De
[85278] novo BRCA1 mutation in a patient with breast cancer and an inherited
[85279] BRCA2 mutation. (Letter) Am. J. Hum. Genet. 65: 567-569, 1999.
[85280] 14934.

[85281] Thompson, D.; Easton, D.: Variation in cancer risks, by mutation
[85282] position, in BRCA2 mutation carriers. Am. J. Hum. Genet. 68: 410-419,
[85283] 2001.

[85284] 14935.

[85285] Thorlacius, S.; Olafsdottir, G.; Tryggvadottir, L.; Neuhausen,
[85286] S.; Jonasson, J. G.; Tavtigian, S. V.; Tulinius, H.; Ogmunds-dottir,
[85287] H. M.; Eyfjord, J. E.: A single BRCA2 mutation in male and female
[85288] breast cancer families from Iceland with varied cancer phenotypes. Nature
[85289] Genet. 13: 117-122, 1996.

[85290] 14936.

[85291] Thorlacius, S.; Sigurdsson, S.; Bjarnadottir, H.; Olafsdottir,

[85292] G.; Jonasson, J. G.; Tryggvadottir, L.; Tulinius, H.; Eyfjord, J.

[85293] E.: Study of a single BRCA2 mutation with high carrier frequency

[85294] in a small population. Am. J. Hum. Genet. 60: 1079-1084, 1997.

[85295] 14937.

[85296] Thorlacius, S.; Tryggvadottir, L.; Olafsdottir, G. H.; Jonasson,

[85297] J. G.; Ogmundsdottir, H. M.; Tulinius, H.; Eyfjord, J. E.: Linkage

[85298] to BRCA2 region in hereditary male breast cancer. Lancet 346: 544-545,

[85299] 1995.

[85300] 14938.

[85301] Tonin, P. N.; Mes-Masson, A.-M.; Futreal, P. A.; Morgan, K.; Mahon,

[85302] M.; Foulkes, W. D.; Cole, D. E. C.; Provencher, D.; Ghadirian, P.;

[85303] Narod, S. A.: Founder BRCA1 and BRCA2 mutations in French Canadian
[85304] breast and ovarian cancer families. Am. J. Hum. Genet. 63: 1341-1351,
[85305] 1998.

[85306] 14939.

[85307] van der Luijt, R. B.; van Zon, P. H. A.; Jansen, R. P. M.; van
[85308] der Sijs-Bos, C. J. M.; Warlam-Rodenhuis, C. C.; Ausems, M. G. E.
[85309] M.: De novo recurrent germline mutation of the BRCA2 gene in a patient
[85310] with early onset breast cancer. J. Med. Genet. 38: 102-105, 2001.
[85311] 14940.

[85312] Wagner, T. M. U.; Hirtenlehner, K.; Shen, P.; Moeslinger, R.;
[85313] Muhr, D.; Fleischmann, E.; Concin, H.; Doeller, W.; Haid, A.; Lang,
[85314] A. H.; Mayer, P.; Petru, E.; and 9 others: Global sequence diversity
[85315] of BRCA2: analysis of 71 breast cancer families and 95 control individuals

[85316] of worldwide populations. Hum. Molec. Genet. 8:
413-423, 1999.

[85317] 14941.

[85318] Warren, M.; Smith, A.; Partridge, N.; Masabanda, J.; Griffin,

[85319] D.; Ashworth, A.: Structural analysis of the chicken BRCA2
gene facilitates

[85320] identification of functional domains and disease causing
mutations. Hum.

[85321] Molec. Genet. 11: 841-851, 2002.

[85322] 14942.

[85323] Auroy, S.; Avril, M.-F.; Chompret, A.; Pham, D.; Goldstein,
A.

[85324] M.; Bianchi-Scarra, G.; Frebourg, T.; Joly, P.; Spatz, A.; Ru-
bino,

[85325] C.; Demenais, F.; French Hereditary Melanoma Study
Group; Bressac-de

[85326] Paillerets, B.: Sporadic multiple primary melanoma cases:
CDKN2A

[85327] germline mutations with a founder effect. Genes Chromo-
somes Cancer 32:

[85328] 195-202, 2001.

[85329] 14943.

[85330] Bahuau, M.; Vidaud, D.; Jenkins, R. B.; Bieche, I.; Kimmel, D.

[85331] W.; Assouline, B.; Smith, J. S.; Alderete, B.; Cayuela, J.-M.; Harpey,

[85332] J.-P.; Caille, B.; Vidaud, M.: Germ-line deletion involving the INK4

[85333] locus in familial proneness to melanoma and nervous system tumors. Cancer

[85334] Res. 58: 2298-2303, 1998.

[85335] 14944.

[85336] Bartsch, D.; Shevlin, D. W.; Tung, W. S.; Kisker, O.; Wells, S.

[85337] A., Jr.; Goodfellow, P. J.: Frequent mutations of CDKN2 in primary

[85338] pancreatic adenocarcinomas. Genes Chromosomes Cancer 14: 189-195,

[85339] 1995.

[85340] 14945.

[85341] Borg, A.; Johansson, U.; Johannsson, O.; Hakansson, S.; Westerdahl,

[85342] J.; Masback, A.; Olsson, H.; Ingvar, C.: Novel germline p16 mutation
[85343] in familial malignant melanoma in southern Sweden. Cancer Res. 56:
[85344] 2497-2500, 1996.

[85345] 14946.

[85346] Cairns, J.; Mao, L.; Merlo, A.; Lee, D. J.; Schwab, D.; Eby, Y.;
[85347] Tokino, K.; van der Riet, P.; Blaugrund, J. E.; Sidransky, D.: Rates
[85348] of p16 (MTS1) mutations in primary tumors with 9p loss. Science (Letter) 265:
[85349] 415-417, 1994.

[85350] 14947.

[85351] Hey, P. J.; Twells, R. C. J.; Phillips, M. S.; Nakagawa, Y.; Brown,
[85352] S. D.; Kawaguchi, Y.; Cox, R.; Xie, G.; Dugan, V.; Hammond, H.; Metzker,
[85353] M. L.; Todd, J. A.; Hess, J. F.: Cloning of a novel member of the
[85354] low-density lipoprotein receptor family. Gene 216:
103-111, 1998.

[85355] 14948.

[85356] Weber, Y.; Steinberger, D.; Deuschl, G.; Benecke, R.; Muller,

[85357] U.: Two previously unrecognized splicing mutations of GCH1 in dopa-responsive

[85358] dystonia: exon skipping and one base insertion. Neuroge-netics 1:

[85359] 125-127, 1997.

[85360] 14949.

[85361] Witter, K.; Werner, T.; Blusch, J. H.; Schneider, E.-M.; Riess,

[85362] O.; Ziegler, I.; Rodl, W.; Bacher, A.; Gutlich, M.: Cloning, sequencing

[85363] and functional studies of the gene encoding human GTP cyclohydrolase

[85364] I. Gene 171: 285-290, 1996.

[85365] 14950.

[85366] Bai, C.; Sen, P.; Hofmann, K.; Ma, L.; Goebl, M.; Harper, J. W.;

[85367] Elledge, S. J.: SKP1 connects cell cycle regulators to the ubiquitin

[85368] proteolysis machinery through a novel motif, the F-box.

Cell 86:
[85369] 263-274, 1996.

[85370] 14951.

[85371] Kraus, B.; Pohlschmidt, M.; Leung, A. L. S.; Germino, G. G.; Snarey,
[85372] A.; Schneider, M. C.; Reeders, S. T.; Frischauf, A.-M.: A novel cyclin
[85373] gene (CCNF) in the region of the polycystic kidney disease gene (PKD1). Genomics 24:
[85374] 27-33, 1994.

[85375] 14952.

[85376] Obermayr, F.; Sutherland, H. F.; Kraus, B.; Frischauf, A.-M.:
[85377] Mouse cyclin F maps to a conserved linkage group on mouse chromosome
[85378] 17. Mammalian Genome 6: 149-150, 1995.

[85379] 14953.

[85380] Barker, P. M.; Gowen, C. W.; Lawson, E. E.; Knowles, M. R.: Decreased
[85381] sodium ion absorption across nasal epithelium of very premature infants

[85382] with respiratory distress syndrome. J. Pediat. 130: 373-377, 1997.

[85383] 14954.

[85384] Bonny, O.; Chraibi, A.; Loffing, J.; Jaeger, N. F.; Grunder, S.;

[85385] Horisberger, J.-D.; Rossier, B. C.: Functional expression of a pseudohypoaldosteronism

[85386] type I mutated epithelial Na(+) channel lacking the pore-forming region

[85387] of its alpha subunit. J. Clin. Invest. 104: 967-974, 1999.

[85388] 14955.

[85389] Canessa, C. M.; Schild, L.; Buell, G.; Thorens, B.; Gautschi, I.;

[85390] Horisberger, J.-D.; Rossier, B. C.: Amiloride-sensitive epithelial

[85391] Na(+)-channel is made of three homologous subunits. Nature 367:

[85392] 463-467, 1994.

[85393] 14956.

[85394] Donaldson, S. H.; Hirsh, A.; Li, D. C.; Holloway, G.; Chao, J.;

[85395] Boucher, R. C.; Gabriel, S. E.: Regulation of the epithelial sodium
[85396] channel by serine proteases in human airways. J. Biol. Chem. 277:
[85397] 8338-8345, 2002.

[85398] 14957.

[85399] Harvey, K. F.; Dinudom, A.; Cook, D. I.; Kumar, S.: The Nedd4-like
[85400] protein KIAA0439 is a potential regulator of the epithelial sodium
[85401] channel. J. Biol. Chem. 276: 8597-8601, 2001.

[85402] 14958.

[85403] Iwai, N.; Baba, S.; Mannami, T.; Ogihara, T.; Ogata, J.: Association
[85404] of a sodium channel alpha subunit promoter variant with blood pressure. J.
[85405] Am. Soc. Nephrol. 13: 80-85, 2002.

[85406] 14959.

[85407] Ludwig, M.; Bolkenius, U.; Wickert, L.; Marynen, P.; Bidlingmaier,
[85408] F.: Structural organisation of the gene encoding the al-

pha-subunit
[85409] of the human amiloride-sensitive epithelial sodium chan-
nel. Hum.
[85410] Genet. 102: 576-581, 1998.

[85411] 14960.

[85412] Meisler, M. H.; Barrow, L. L.; Canessa, C. M.; Rossier, B. C.

[85413] : SCNN1, an epithelial cell sodium channel gene in the
conserved linkage
[85414] group on mouse chromosome 6 and human chromosome
12. Genomics 24:
[85415] 185-186, 1994.

[85416] 14961.

[85417] O'Brodovich, H. M.: Immature epithelial Na+ channel ex-
pression
[85418] is one of the pathogenetic mechanisms leading to human
neonatal respiratory
[85419] distress syndrome. Proc. Assoc. Am. Phys. 108: 345-355,
1996.
[85420] 14962.

[85421] Oh, Y. S.; Lee, S.-I.; Won, C.-K.; Warnock, D. G.: An Alu
cassette

[85422] in the human epithelial sodium channel. Biochim. Biophys. Acta 1520:
[85423] 94-98, 2001.

[85424] 14963.

[85425] Pathak, B. G.; Shaughnessy, J. D., Jr.; Meneton, P.; Greeb, J.;
[85426] Shull, G. E.; Jenkins, N. A.; Copeland, N. G.: Mouse chromosomal
[85427] location of three epithelial sodium channel subunit genes and an apical
[85428] sodium chloride cotransporter gene. Genomics 33: 124-127, 1996.
[85429] 14964.

[85430] Reddy, M. M.; Light, M. J.; Quinton, P. M.: Activation of the
[85431] epithelial Na(+) channel (ENaC) requires CFTR CI(-) channel function. Nature 402:
[85432] 301-304, 1999.

[85433] 14965.

[85434] Rossier, B. C.: 1996 Homer Smith award lecture: Cum grano salis:
[85435] the epithelial sodium channel and the control of blood

pressure. J.

[85436] Am. Soc. Nephrol. 8: 980-992, 1997.

[85437] 14966.

[85438] Schaedel, C.; Marthinsen, L.; Kristoffersson, A.-C.; Korn-
falt,

[85439] R.; Nilsson, K. O.; Orlenius, B.; Holmberg, L.: Lung symp-
toms in

[85440] pseudohypoaldosteronism type 1 are associated with de-
ficiency of the

[85441] alpha-subunit of the epithelial sodium channel. J. Pediat.
135:

[85442] 739-745, 1999.

[85443] 14967.

[85444] Voilley, N.; Lingueglia, E.; Champigny, G.; Mattei, M.-G.;
Waldmann,

[85445] R.; Lazdunski, M.; Barbry, P.: The lung amiloride-sensitive
Na(+)

[85446] channel: biophysical properties, pharmacology, ontogene-
sis, and molecular

[85447] cloning. Proc. Nat. Acad. Sci. 91: 247-251, 1994.

[85448] 14968.

[85449] Arriza, J. L.; Kavanaugh, M. P.; Fairman, W. A.; Wu, Y.-N.; Murdoch,
[85450] G. H.; North, R. A.; Amara, S. G.: Cloning and expression of a human
[85451] neutral amino acid transporter with structural similarity to the glutamate
[85452] transporter gene family. J. Biol. Chem. 268: 15329-15332, 1993.
[85453] 14969.

[85454] Hofmann, K.; Duker, M.; Fink, T.; Lichter, P.; Stoffel, W.: Human
[85455] neutral amino acid transporter ASCT1: structure of the gene (SLC1A4)
[85456] and localization to chromosome 2p13-p15. Genomics 24: 20-26, 1994.
[85457] 14970.

[85458] Shafqat, S.; Tamarappoo, B. K.; Kilberg, M. S.; Puranam, R. S.;
[85459] McNamara, J. O.; Guadano-Ferraz, A.; Fremeau, R. T., Jr.: Cloning
[85460] and expression of a novel Na(+)-dependent neutral amino acid transporter

[85461] structurally related to mammalian Na(+)/glutamate co-transporters. J.

[85462] Biol. Chem. 268: 15351-15355, 1993.

[85463] 14971.

[85464] Zerangue, N.; Kavanaugh, M. P.: ASCT-1 is a neutral amino acid

[85465] exchanger with chloride channel activity. J. Biol. Chem. 271: 27991-27994,

[85466] 1996.

[85467] 14972.

[85468] Fukuda, R.; McNew, J. A.; Weber, T.; Parlati, F.; Engel, T.; Nickel,

[85469] W.; Rothman, J. E.; Sollner, T. H.: Functional architecture of an

[85470] intracellular membrane t-SNARE. Nature 407: 198-202, 2000.

[85471] 14973.

[85472] Grabs, D.; Bergmann, M.; Urban, M.; Post, A.; Gratzl, M.: Rab3

[85473] proteins and SNAP-25, essential components of the exo-cytosis machinery

[85474] in conventional synapses, are absent from ribbon synapses of the mouse
[85475] retina. Europ. J. Neurosci. 8: 162-168, 1996.

[85476] 14974.

[85477] Zhao, N.; Hashida, H.; Takahashi, N.; Sakaki, Y.: Cloning and
[85478] sequence analysis of the human SNAP25 cDNA. Gene 145: 313-314, 1994.
[85479] 14975.

[85480] Newton, J. S.; Deed, R. W.; Mitchell, E. L. D.; Murphy, J. J.;

[85481] Norton, J. D.: A B cell specific immediate early human gene is located
[85482] on chromosome band 1q31 and encodes an alpha helical basic phosphoprotein. Biochim.
[85483] Biophys. Acta 1216: 314-316, 1993.

[85484] 14976.

[85485] Sierra, D. A.; Gilbert, D. J.; Householder, D.; Grishin, N. V.;

[85486] Yu, K.; Ukidwe, P.; Barker, S. A.; He, W.; Wensel, T. G.; Otero, G.;
[85487] Brown, G.; Copeland, N. G.; Jenkins, N. A.; Wilkie, T. M.: Evolution

[85488] of the regulators of G-protein signaling multigene family in mouse

[85489] and human. Genomics 79: 177-185, 2002.

[85490] 14977.

[85491] Chiao, J. H.; Roy, K.; Tolner, B.; Yang, C.-H.; Sirotnak, F. M.

[85492] : RFC-1 gene expression regulates folate absorption in mouse small

[85493] intestine. J. Biol. Chem. 272: 11165-11170, 1997.

[85494] 14978.

[85495] Nguyen, T. T.; Dyer, D. L.; Dunning, D. D.; Rubin, S. A.; Grant,

[85496] K. E.; Said, H. M.: Human intestinal folate transport: cloning, expression,

[85497] and distribution of complementary RNA. Gastroenterology 112: 783-791,

[85498] 1997.

[85499] 14979.

[85500] Prasad, P. D.; Ramamoorthy, S.; Leibach, F. H.; Ganapathy, V.:

[85501] Molecular cloning of the human placental folate trans-

porter. Biochem.

[85502] Biophys. Res. Commun. 206: 681-687, 1995.

[85503] 14980.

[85504] Shaw, G. M.; Lammer, E. J.; Zhu, H.; Baker, M. W.; Neri, E.; Finnell,

[85505] R. H.: Maternal periconceptional vitamin use, genetic vari-ation of

[85506] infant reduced folate carrier (A80G), and risk of spina bi-fida. Am.

[85507] J. Med. Genet. 108: 1-6, 2002.

[85508] 14981.

[85509] Tolner, B.; Roy, K.; Sirotnak, F. M.: Structural analysis of the

[85510] human RFC-1 gene encoding a folate transporter reveals multiple promoters

[85511] and alternatively spliced transcripts with 5-prime end heterogeneity. Gene 211:

[85512] 331-341, 1998.

[85513] 14982.

[85514] Tse, A.; Brigle, K.; Taylor, S. M.; Moran, R. G.: Mutations in

[85515] the reduced folate carrier gene which confer dominant re-

sistance to

[85516] 5,10-dideazatetrahydrofolate. J. Biol. Chem. 273: 25953-25960, 1998.

[85517] 14983.

[85518] Williams, F. M. R.; Flintoff, W. F.: Structural organization of

[85519] the human reduced folate carrier gene: evidence for 5-prime heterogeneity

[85520] in lymphoblast mRNA. Somat. Cell Molec. Genet. 24: 143-156, 1998.

[85521] 14984.

[85522] Williams, F. M. R.; Flintoff, W. F.: Isolation of a human cDNA

[85523] that complements a mutant hamster cell defective in methotrexate uptake. J.

[85524] Biol. Chem. 270: 2987-2992, 1995.

[85525] 14985.

[85526] Wong, S. C.; Proefke, S. A.; Bhushan, A.; Matherly, L. H.: Isolation

[85527] of human cDNAs that restore methotrexate sensitivity and reduced folate

[85528] carrier activity in methotrexate transport-defective Chi-

nese hamster

[85529] ovary cells. J. Biol. Chem. 270: 17468-17475, 1995.

[85530] 14986.

[85531] Yang-Feng, T. L.; Ma, Y.-Y.; Liang, R.; Prasad, P. D.; Leibach,

[85532] F. H.; Ganapathy, V.: Assignment of the human folate transporter

[85533] gene to chromosome 21q22.3 by somatic cell hybrid analysis and in

[85534] situ hybridization. Biochem. Biophys. Res. Commun. 210: 874-879,

[85535] 1995.

[85536] 14987.

[85537] Lees, J. A.; Saito, M.; Vidal, M.; Valentine, M.; Look, T.; Harlow,

[85538] E.; Dyson, N.; Helin, K.: The retinoblastoma protein binds to a family

[85539] of E2F transcription factors. Molec. Cell. Biol. 13: 7813-7825,

[85540] 1993.

[85541] 14988.

[85542] Cloud, J. E.; Rogers, C.; Reza, T. L.; Ziebold, U.; Stone, J. R.;
[85543] Picard, M. H.; Caron, A. M.; Bronson, R. T.; Lees, J. A.: Mutant
[85544] mouse models reveal the relative roles of E2F1 and E2F3 in vivo. Molec.
[85545] Cell. Biol. 22: 2663-2672, 2002.

[85546] 14989.

[85547] He, Y.; Armanious, M. K.; Thomas, M. J.; Cress, W. D.: Identification
[85548] of E2F-3B, an alternative form of E2F-3 lacking a conserved N-terminal
[85549] region. Oncogene 19: 3422-3433, 2000.

[85550] 14990.

[85551] Halford, M. M.; Armes, J.; Buchert, M.; Meskenaite, V.; Grail,
[85552] D.; Hibbs, M. L.; Wilks, A. F.; Farlie, P. G.; Newgreen, D. F.; Hovens,
[85553] C. M.; Stacker, S. A.: Ryk-deficient mice exhibit craniofacial defects
[85554] associated with perturbed Eph receptor crosstalk. Nature Genet. 25:

[85555] 414-418, 2000.

[85556] 14991.

[85557] Nakamura, S.; Stock, D. W.; Wydner, K. L.; Bollekens, J. A.; Takeshita,

[85558] K.; Nagai, B. M.; Chiba, S.; Kitamura, T.; Freeland, T. M.; Zhao,

[85559] Z.; Minowada, J.; Lawrence, J. B.; Weiss, K. M.; Ruddle, F. H.: Genomic

[85560] analysis of a new mammalian distal-less gene: Dlx7. Genomics 38:

[85561] 314-324, 1996.

[85562] 14992.

[85563] Adams, R. H.; Diella, F.; Hennig, S.; Helmbacher, F.; Deutsch,

[85564] U.; Klein, R.: The cytoplasmic domain of the ligand ephrinB2 is required

[85565] for vascular morphogenesis but not cranial neural crest migration. Cell 104:

[85566] 57-69, 2001.

[85567] 14993.

[85568] Bennett, B. D.; Zeigler, F. C.; Gu, Q.; Fendly, B.; Goddard,

A.

[85569] D.; Gillett, N.; Matthews, W.: Molecular cloning of a ligand for

[85570] the EPH-related receptor protein-tyrosine kinase Htk. Proc. Nat.

[85571] Acad. Sci. 92: 1866-1870, 1995.

[85572] 14994.

[85573] Bonaldo, M. F.; Yu, M. T.; Jelenc, P.; Brown, S.; Su, L.; Law-ton,

[85574] L.; Deaven, L.; Efstratiadis, A.; Warburton, D.; Soares, M. B.: Selection

[85575] of cDNAs using chromosome-specific genomic clones: application to

[85576] human chromosome 13. Hum. Molec. Genet. 3: 1663-1673, 1994.

[85577] 14995.

[85578] Cerretti, D. P.; Vanden Bos, T.; Nelson, N.; Kozlosky, C. J.; Reddy,

[85579] P.; Maraskovsky, E.; Park, L. S.; Lyman, S. D.; Copeland, N. G.; Gilbert,

[85580] D. J.; Jenkins, N. A.; Fletcher, F. A.: Isolation of LERK-5: a ligand

[85581] of the EPH-related receptor tyrosine kinases. Molec. Immun. 32:
[85582] 1197-1205, 1995.

[85583] 14996.

[85584] Singh, J.; Itahana, Y.; Parrinello, S.; Murata, K.; Desprez, P.-Y.
[85585] : Molecular cloning and characterization of a zinc finger protein
[85586] involved in Id-1-stimulated mammary epithelial cell growth. J. Biol.
[85587] Chem. 276: 11852-11858, 2001.

[85588] 14997.

[85589] Kumar, S.; Kinoshita, M.; Noda, M.; Copeland, N. G.; Jenkins, N.
[85590] A.: Induction of apoptosis by the mouse Nedd2 gene, which encodes
[85591] a protein similar to the product of the Caenorhabditis elegans cell
[85592] death gene ced-3 and the mammalian IL-1-beta-converting enzyme. Genes
[85593] Dev. 8: 1613-1626, 1994.

[85594] 14998.

[85595] Bollag, R. J.; Siegfried, Z.; Cebra-Thomas, J. A.; Garvey, N.;

[85596] Davison, E. M.; Silver, L. M.: An ancient family of embry-
onically

[85597] expressed mouse genes sharing a conserved protein motif
with the T

[85598] locus. Nature Genet. 7: 383-389, 1994.

[85599] 14999.

[85600] Campbell, C.; Goodrich, K.; Casey, G.; Beatty, B.: Cloning
and

[85601] mapping of a human gene (TBX2) sharing a highly con-
served protein

[85602] motif with the Drosophila omb gene. Genomics 28:
255-260, 1995.

[85603] 15000.

[85604] Campbell, C. E.; Casey, G.; Goodrich, K.: Genomic struc-
ture of

[85605] TBX2 indicates conservation with distantly related T-box
genes. Mammalian

[85606] Genome 9: 70-73, 1998.

[85607] 15001.

[85608] Jacobs, J. J. L.; Keblusek, P.; Robanus-Maandag, E.; Kristel, P.;

[85609] Lingbeek, M.; Nederlof, P. M.; van Welsem, T.; van de Vijver, M. J.;

[85610] Koh, E. Y.; Daley, G. Q.; van Lohuizen, M.: Senescence bypass screen

[85611] identifies TBX2, which represses Cdkn2a(p19ARF) and is amplified in

[85612] a subset of human breast cancers. Nature Genet. 26: 291-299, 2000.

[85613] 15002.

[85614] Law, D. J.; Gebhur, T.; Garvey, N.; Agulnik, S. I.; Silver, L.

[85615] M.: Identification, characterization, and localization to chromosome

[85616] 17q21-22 of the human TBX2 homolog, member of a conserved developmental

[85617] gene family. Mammalian Genome 6: 793-797, 1995.

[85618] 15003.

[85619] Pflugfelder, G. O.; Roth, H.; Poeck, B.: A homology domain shared

[85620] between Drosophila optomotor-blind and mouse Brachyury is involved

[85621] in DNA binding. Biochem. Biophys. Res. Commun. 186: 918-925, 1992.

[85622] 15004.

[85623] Sowden, J. C.; Holt, J. K. L.; Meins, M.; Smith, H. K.; Bhat-tacharya,

[85624] S. S.: Expression of Drosophila omb-related T-box genes in the developing

[85625] human and mouse neural retina. Invest. Ophthal. Vis. Sci. 42: 3095-3102,
[85626] 2001.

[85627] 15005.

[85628] Walter, L.; Dirks, B.; Rothermel, E.; Heyens, M.; Szpirer, C.;

[85629] Levan, G.; Gunther, E.: A novel, conserved gene of the rat that is
[85630] developmentally regulated in the testis. Mammalian Genome 5: 216-221,
[85631] 1994.

[85632] 15006.

[85633] Walter, L.; Marynen, P.; Szpirer, J.; Levan, G.; Gunther, E.:

[85634] Identification of a novel conserved human gene, TEGT. Genomics 28:
[85635] 301-304, 1995.

[85636] 15007.

[85637] Xu, Q.; Reed, J. C.: Bax inhibitor-1, a mammalian apoptosis suppressor
[85638] identified by functional screening in yeast. Molec. Cell 1: 337-346,

[85639] 1998.

[85640] 15008.

[85641] Hsu, Y.-C.; Perin, M. S.: Human neuronal pentraxin II (NPTX2):

[85642] conservation, genomic structure, and chromosomal localization. Genomics 28:

[85643] 220-227, 1995.

[85644] 15009.

[85645] Schlimgen, A. K.; Helms, J. A.; Vogel, H.; Perin, M. S.: Neuronal

[85646] pentraxin, a secreted protein with homology to acute phase proteins

[85647] of the immune system. Neuron 14: 519-526, 1995.

[85648] 15010.

[85649] Kumar, S.; White, D. L.; Takai, S.; Turczynowicz, S.; Juttner,

[85650] C. A.; Hughes, T. P.: Apoptosis regulatory gene NEDD2 maps to human

[85651] chromosome segment 7q34-35, a region frequently affected in haematological

[85652] neoplasms. Hum. Genet. 95: 641-644, 1995.

[85653] 15011.

[85654] Lassus, P.; Opitz-Araya, X.; Lazebnik, Y.: Requirement for caspase-2

[85655] in stress-induced apoptosis before mitochondrial permeabilization. Science 297:

[85656] 1352-1354, 2002.

[85657] 15012.

[85658] Fuhlbrigge, R. C.; Kieffer, J. D.; Armerding, D.; Kupper, T. S.

[85659] : Cutaneous lymphocyte antigen is a specialized form of PSGL-1 expressed

[85660] on skin-homing T cells. Nature 389: 978-981, 1997.

[85661] 15013.

[85662] Herron, M. J.; Nelson, C. M.; Larson, J.; Snapp, K. R.; Kansas,

[85663] G. S.; Goodman, J. L.: Intracellular parasitism by the human granulocytic

[85664] ehrlichiosis bacterium through the P-selectin ligand, PSGL-1. Science 288:

[85665] 1653-1656, 2000.

[85666] 15014.

[85667] Veldman, G. M.; Bean, K. M.; Cumming, D. A.; Eddy, R. L.; Sait,
[85668] S. N. J.; Shows, T. B.: Genomic organization and chromosomal localization
[85669] of the gene encoding human P-selectin glycoprotein ligand. J. Biol.
[85670] Chem. 270: 16470-16475, 1995.

[85671] 15015.

[85672] Yang, J.; Galipeau, J.; Kozak, C. A.; Furie, B. C.; Furie, B.:

[85673] Mouse P-selectin glycoprotein ligand-1: molecular cloning, chromosomal
[85674] localization, and expression of a functional P-selectin receptor. Blood 87:
[85675] 4176-4186, 1996.

[85676] 15016.

[85677] Brown, C. B.; Boyer, A. S.; Runyan, R. B.; Barnett, J. V.: Requirement
[85678] of type III TGF-beta receptor for endocardial cell transformation
[85679] in the heart. Science 283: 2080-2082, 1999.

[85680] 15017.

[85681] Lewis, K. A.; Gray, P. C.; Blount, A. L.; MacConell, L. A.; Wiater,
[85682] E.; Bilezikjian, L. M.; Vale, W.: Betaglycan binds inhibin and can
[85683] mediate functional antagonism of activin signalling. Nature 404:
[85684] 411-414, 2000.

[85685] 15018.

[85686] Hu, Y.-F.; Luscher, B.; Admon, A.; Mermod, N.; Tjian, R.: Transcription
[85687] factor AP-4 contains multiple dimerization domains that regulate dimer
[85688] specificity. Genes Dev. 4: 1741-1752, 1990.

[85689] 15019.

[85690] Mermod, N.; Williams, T. J.; Tjian, R.: Enhancer binding factors
[85691] AP-4 and AP-1 act in concert to activate SV40 late transcription in
[85692] vitro. Nature 332: 557-561, 1988.

[85693] 15020.

[85694] Carr, C. S.; Sharp, P. A.: A helix-loop-helix protein related

[85695] to the immunoglobulin E box-binding proteins. Molec. Cell. Biol. 10:

[85696] 4384-4388, 1990.

[85697] 15021.

[85698] Bonner, C. A.; Loftus, S. K.; Wasmuth, J. J.: Isolation, characterization,

[85699] and precise physical localization of human CDX1, a caudal-type homeobox

[85700] gene. Genomics 28: 206-211, 1995.

[85701] 15022.

[85702] Subramanian, V.; Meyer, B. I.; Gruss, P.: Disruption of the murine

[85703] homeobox gene Cdx1 affects axial skeletal identities by altering the

[85704] mesodermal expression domains of Hox genes. Cell 83: 641-653, 1995.

[85705] 15023.

[85706] Schwientek, T.; Nomoto, M.; Levery, S. B.; et al: Control of O-glycan

[85707] branch formation. J. Biol. Chem. 274: 4504-4512, 1999.

[85708] 15024.

[85709] Walczak, H.; Degli-Esposti, M. A.; Johnson, R. S.; Smolak, P. J.;
[85710] Waugh, J. Y.; Boiani, N.; Timour, M. S.; Gerhart, M. J.; Schooley,
[85711] K. A.; Smith, C. A.; Goodwin, R. G.; Rauch, C. T.: TRAIL-R2: a novel
[85712] apoptosis-mediating receptor for TRAIL. EMBO J. 16: 5386-5397, 1997.
[85713] 15025.

[85714] Wu, G. S.; Burns, T. F.; McDonald, E. R., III; Jiang, W.; Meng,
[85715] R.; Krantz, I. D.; Kao, G.; Gan, D.-D.; Zhou, J.-Y.; Muschel, R.;
[85716] Hamilton, S. R.; Spinner, N. B.; Markowitz, S.; Wu, G.; El-Deiry,
[85717] W. S.: KILLER/DR5 is a DNA damage-inducible p53-regulated death receptor
[85718] gene. Nature Genet. 17: 141-143, 1997.

[85719] 15026.

[85720] Degli-Esposti, M. A.; Smolak, P. J.; Walczak, H.; Waugh, J.; Huang,
[85721] C.-P.; DuBose, R. F.; Goodwin, R. G.; Smith, C. A.: Cloning

and characterization
[85722] of TRAIL-R3, a novel member of the emerging TRAIL receptor family. J.
[85723] Exp. Med. 186: 1165-1170, 1997.

[85724] 15027.

[85725] Kanaar, R.; Troelstra, C.; Swagemakers, S. M. A.; Essers, J.; Smit,
[85726] B.; Franssen, J.-H.; Pastink, A.; Bezzubova, O. Y.; Buerstedde, J.-M.;
[85727] Clever, B.; Heyer, W.-D.; Hoeijmakers, J. H. J.: Human and mouse
[85728] homologs of the Saccharomyces cerevisiae RAD54 DNA repair gene: evidence
[85729] for functional conservation. Curr. Biol. 6: 828-838, 1996.

[85730] 15028.

[85731] Matsuda, M.; Miyagawa, K.; Takahashi, M.; Fukuda, T.; Kataoka,
[85732] T.; Asahara, T.; Inui, H.; Watatani, M.; Yasutomi, M.; Kamada, N.;
[85733] Dohi, K.; Kamiya, K.: Mutations in the RAD54 recombination gene in
[85734] primary cancers. Oncogene 18: 3427-3430, 1999.

[85735] 15029.

[85736] Tucker, J. E.; Winkfein, R. J.; Cooper, C. B.; Schnetkamp, P.
P.

[85737] : cDNA cloning of the human retinal rod Na-Ca + K ex-
changer: comparison

[85738] with a revised bovine sequence. Invest. Ophthal. Vis. Sci.
39: 435-440,

[85739] 1998.

[85740] 15030.

[85741] Tucker, J. E.; Winkfein, R. J.; Murthy, S. K.; Friedman, J. S.;

[85742] Walter, M. A.; Demetrick, D. J.; Schnetkamp, P. P. M.:
Chromosomal

[85743] localization and genomic organization of the human reti-
nal rod Na-Ca+K

[85744] exchanger. Hum. Genet. 103: 411-414, 1998.

[85745] 15031.

[85746] Dennis, J. A.; Healy, P. J.; Beaudet, A. L.; O'Brien, W. E.:

[85747] Molecular definition of bovine argininosuccinate syn-
thetase deficiency. Proc.

[85748] Nat. Acad. Sci. 86: 7947-7951, 1989.

[85749] 15032.

[85750] McCarrey, J. R.; Riggs, A. D.: Determinator-inhibitor pairs as

[85751] a mechanism for threshold setting in development: a possible function

[85752] for pseudogenes. Proc. Nat. Acad. Sci. 83: 679-683, 1986.

[85753] 15033.

[85754] Su, T.-S.; Nussbaum, R. L.; Airhart, S.; Ledbetter, D. H.; Mohandas,

[85755] T.; O'Brien, W. E.; Beaudet, A. L.: Human chromosomal assignments

[85756] for 14 argininosuccinate synthetase pseudogenes: cloned DNAs as reagents

[85757] for cytogenetic analysis. Am. J. Hum. Genet. 36: 954-964, 1984.

[85758] 15034.

[85759] Kobayashi, K.; Sinasac, D. S.; Iijima, M.; Boright, A. P.; Begum,

[85760] L.; Lee, J. R.; Yasuda, T.; Ikeda, S.; Hirano, R.; Terazono, H.; Crackower,

[85761] M. A.; Kondo, I.; Tsui, L.-C.; Scherer, S. W.; Saheki, T.: The gene